

ANTIBODY MOLECULES TO APRIL AND USES THEREOF

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 62/259,897, filed November 25, 2015, U.S. Provisional Application No. 62/313,684, filed March 25, 2016, U.S. Provisional Application No. 62/399,087, filed September 23, 2016, U.S. and U.S. Provisional Application No. 62/422,848, filed November 16, 2016. The contents of the aforementioned applications are hereby incorporated by reference in their entirety.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on November 21, 2016, is named P2029-7004WO_SL.txt and is 259,298 bytes in size.

BACKGROUND

IgA nephropathy is one of the most prevalent, chronic glomerular diseases worldwide. Conservative epidemiological estimates cite a global incidence of approximately 5-50 cases/million (children) and 10-40 cases /million (adults). This incidence of disease presents a regional bias with a higher prevalence in Asia and the Americas, with a particularly higher disease burden in Japan and regions of China. Biopsy confirmed cases of IgA nephropathy in Japan are projected at approximately 350,000. In the US, this projection is approximately 100,000—as such, it is the most frequently diagnosed 1° glomerular disease in adults. While a relatively indolent disease, IgA nephropathy leads to end stage renal disease (ESRD), *i.e.*, renal failure in 20-50% of patients within a 20-30 year span. These numbers are likely grossly underreported given the need to confirm the disease by kidney biopsy, a protocol that is variably practiced in various clinical settings. The disease has a complex pathogenesis with genetic, epidemiological, and potentially environmental components to disease etiology, pathology, and progression. It likewise has a variable clinical presentation ranging from asymptomatic to end-stage renal failure (ESRD). IgA nephropathy is caused by the deposition of IgA, typically in the form of immune complexes in the mesangium of the kidney. There are currently no disease-specific treatments to address primary disease or progression.

There is a need for developing new approaches for treating, preventing and diagnosing IgA nephropathy and other disorders that share similar disease mechanisms.

SUMMARY

This disclosure provides, at least in part, antibody molecules that bind to APRIL, *e.g.*, human and/or mouse APRIL, and that comprise one or more functional and structural properties disclosed herein. In an embodiment, the antibody molecule binds to and/or reduces (*e.g.*, inhibits, blocks or

neutralizes) one or more activities of APRIL. In an embodiment, the antibody molecule binds to a region in APRIL that interacts with TACI (*e.g.*, the CRD2 domain of TACI). In an embodiment, the antibody molecule binds to one or more residues within a region of human APRIL as defined in any of **Tables 3-4 or 7-8**. While not wishing to be bound by theory, it is believed that in an embodiment, improved or optimal inhibition of APRIL activities can be achieved, by targeting certain region(s) on APRIL (*e.g.*, the region(s) associated with the interactions between APRIL and the CDR2 domain of TACI). In an embodiment, the antibody molecule is selected from **Table 1 or 5**, or competes for binding to APRIL with an antibody molecule selected from **Table 1 or 5**. In an embodiment, the antibody molecule binds to the same or overlapping epitope as the epitope recognized by an antibody molecule selected from **Table 1 or 5**. In an embodiment, the antibody molecule comprises one or more heavy chain variable regions and/or one or more light chain variable regions described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one or more heavy chain CDRs and/or one or more light chain CDRs described in **Table 1 or 5**. In an embodiment, nucleic acid molecules encoding the antibody molecules, expression vectors, host cells, compositions (*e.g.*, pharmaceutical compositions), kits, containers, and methods for making the antibody molecules, are also provided. The antibody molecules disclosed herein can be used (alone or in combination with other agents or therapeutic modalities) to treat, prevent and/or diagnose disorders associated with APRIL, such as IgA nephropathy.

Accordingly, this disclosure provides an antibody molecule, *e.g.*, an antibody molecule described herein, having one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23) of the following properties:

- a) Binds to human APRIL with high affinity, *e.g.*, with a dissociation constant (K_D) of less than about 100 nM, typically about 10 nM, and more typically, about 10-0.001 nM, about 10-0.01 nM, about 5-0.01 nM, about 3-0.05 nM, about 1-0.1 nM, or stronger, *e.g.*, less than about 80, 70, 60, 50, 40, 30, 20, 10, 8, 6, 4, 3, 2, 1, 0.5, 0.2, 0.1, 0.05, 0.01, 0.005, or 0.001 nM,
- b) Binds to mouse APRIL with high affinity, *e.g.*, with a dissociation constant (K_D) of less than about 100 nM, typically about 10 nM, and more typically, about 10-0.001 nM, about 10-0.01 nM, about 5-0.01 nM, about 3-0.05 nM, about 1-0.1 nM, or stronger, *e.g.*, less than about 80, 70, 60, 50, 40, 30, 20, 10, 8, 6, 4, 3, 2, 1, 0.5, 0.2, 0.1, 0.05, 0.01, 0.005, or 0.001 nM,
- c) Does not bind to mouse APRIL, or binds mouse APRIL with low affinity, *e.g.*, with a dissociation constant (K_D) of greater than about 500 nM, *e.g.*, greater than about 1000 nM,
- d) Does not bind, or binds with low affinity, *e.g.*, with a dissociation constant (K_D) of greater than about 500 nM, *e.g.*, greater than about 1000 nM, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7,

8, or more) cytokines from the TNF superfamily (TNFSF) other than APRIL (*e.g.*, TNF α , CD40 (TNFSF4), FasL (TNFSF6), TRAIL (TNFSF10), RANKL (TNFSF11), Tweak (TNFSF12), BAFF (TNFSF13B), or LIGHT (TNFSF14)),

- 5 e) Binds to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) residues within a region of APRIL as defined in **Table 3**, or binds specifically to an epitope on APRIL, *e.g.*, an epitope comprising one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) residues described in **Table 3**,
- 10 f) Binds to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all) residues within a region of APRIL as defined in **Table 4**, or binds specifically to an epitope on APRIL, *e.g.*, an epitope comprising one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all) residues described in **Table 4**,
- 15 g) Binds to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all) residues within a region of APRIL as defined in **Table 7**, or binds specifically to an epitope on APRIL, *e.g.*, an epitope comprising one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all) residues described in **Table 7**,
- 20 h) Binds to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) residues within a region of APRIL as defined in **Table 8**, or binds specifically to an epitope on APRIL, *e.g.*, an epitope comprising one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) residues described in **Table 8**,
- 25 i) Binds specifically to an epitope on APRIL, *e.g.*, the same, similar, or overlapping epitope as the epitope recognized by a monoclonal antibody described in **Table 1 or 5**, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237,
- 30 j) Reduces (*e.g.*, inhibits, blocks, or neutralizes) one or more biological activities of APRIL (*e.g.*, human APRIL, mouse APRIL, or both), *in vitro*, *ex vivo*, or *in vivo*,
- k) Reduces (*e.g.*, inhibits, blocks, or neutralizes) binding of human APRIL to TACI, *e.g.*, at an IC₅₀ of about 50 nM or less, typically about 0.01-50 nM, 0.1-25 nM, 0.1-10 nM, 0.5-5 nM, or 1-5 nM, *e.g.*, less than about 40, 30, 20, 10, 5, 1, 0.5, 0.2, 0.1, 0.05, or 0.01 nM, *e.g.*, as determined by a method described herein,
- 35 l) Reduces (*e.g.*, inhibits, blocks, or neutralizes) binding of mouse APRIL to TACI, *e.g.*, at an IC₅₀ of about 100 nM or less, typically about 0.01-75 nM, 0.1-50 nM, 0.1-25 nM, 0.1-10 nM, 0.5-5 nM, or 1-5 nM, *e.g.*, less than about 80, 60, 40, 20, 10, 5, 1, 0.5, 0.2, 0.1, 0.05, or 0.01 nM, *e.g.*, as determined by a method described herein,

- m) Reduces (*e.g.*, inhibits, blocks, or neutralizes) binding of human APRIL to BMCA, *e.g.*, at an IC₅₀ of about 50 nM or less, typically about 0.01-50 nM, 0.1-25 nM, 0.1-10 nM, 0.5-5 nM, or 1-5 nM, *e.g.*, less than about 40, 30, 20, 10, 5, 1, 0.5, 0.2, 0.1, 0.05, or 0.01 nM, *e.g.*, as determined by a method described herein,
- 5 n) Reduces (*e.g.*, inhibits, blocks, or neutralizes) binding of mouse APRIL to BMCA, *e.g.*, at an IC₅₀ of about 200 nM or less, typically about 0.01-200 nM, 0.1-150 nM, 0.1-100 nM, 0.1-50 nM, 0.1-25 nM, 0.1-10 nM, 0.5-5 nM, or 1-5 nM, *e.g.*, less than about 150, 100, 50, 40, 30, 20, 10, 5, 1, 0.5, 0.2, 0.1, 0.05, or 0.01 nM, *e.g.*, as determined by a method described herein,
- 10 o) Shows the same or similar binding affinity or specificity, or both, as a monoclonal antibody described in **Table 1 or 5**, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063,
- 15 4540-033, 4439, or 4237,
- p) Shows the same or similar binding affinity or specificity, or both, as an antibody molecule comprising a heavy chain variable region and/or light chain variable region described in **Table 1 or 5**, *e.g.*, a heavy chain variable region and/or light chain variable region of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406,
- 20 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237,
- q) Shows the same or similar binding affinity or specificity, or both, as an antibody molecule comprising one or more (*e.g.*, two or three) heavy chain CDRs and/or one or more (*e.g.*,
- 25 two or three) light chain CDRs described in **Table 1 or 5**, *e.g.*, one or more (*e.g.*, two or three) heavy chain CDRs and/or one or more (two or three) light chain CDRs of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934,
- 30 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237,
- r) Shows the same or similar binding affinity or specificity, or both, as an antibody molecule comprising an amino acid sequence shown in **Table 1 or 5**,
- s) Shows the same or similar binding affinity or specificity, or both, as an antibody molecule comprising an amino acid sequence encoded by a nucleotide sequence shown in **Table 2**,
- 35 t) Inhibits, *e.g.*, competitively inhibits, the binding of a second antibody molecule to human APRIL, mouse APRIL, or both, wherein the second antibody molecule is an antibody molecule chosen from **Table 1 or 5**, *e.g.*, any of monoclonal antibodies 2218, 2419,

2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806,
2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922,
3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540,
4540-063, 4540-033, 4439, or 4237,

- 5 u) Competes for binding with a second antibody molecule to human APRIL, mouse APRIL,
or both, wherein the second antibody molecule is a monoclonal antibody chosen from
Table 1 or 5, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205,
2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205,
2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525,
10 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033,
4439, or 4237,
- v) Has one or more biological properties of a monoclonal antibody chosen from **Table 1 or**
5, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206,
2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210,
15 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621,
4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or
4237,
- w) Has one or more structural properties of a monoclonal antibody chosen from **Table 1 or**
5, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206,
20 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210,
2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621,
4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or
4237, or
- x) Has one or more pharmacokinetic properties of a monoclonal antibody chosen from
25 **Table 1 or 5**, *e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205,
2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205,
2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525,
3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033,
4439, or 4237.

30 In an aspect, the disclosure features an anti-APRIL antibody molecule, which:

- (i) binds, or substantially binds, to human APRIL;
(ii) binds, or substantially binds, to mouse APRIL;
(iii) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL,
35 or both) to TACI (*e.g.*, human TACI, mouse TACI, or both); and
(iv) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL,
or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both).

In an embodiment, the antibody molecule is a synthetic antibody molecule. In an embodiment, the antibody molecule is an isolated antibody molecule.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), *e.g.*, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, or between 0.1 nM and 1 nM, *e.g.*, as determined by a method described herein.

nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule is an IgG antibody molecule, *e.g.*, comprising a heavy chain constant region of IgG, *e.g.*, chosen from IgG1, IgG2 (*e.g.*, IgG2a), IgG3, or IgG4, *e.g.*, IgG2 or IgG4. In an embodiment, the antibody molecule is an IgG1 antibody molecule, *e.g.*, having an IgG1 constant region described herein. In another embodiment, the antibody molecule is an IgG2 antibody molecule *e.g.*, having an IgG2 constant region described herein. In an embodiment, the antibody molecule comprises a light chain constant region of kappa or lambda light chain.

In an embodiment, the antibody molecule comprises an Fc region. In an embodiment, the Fc region comprises one or more mutations located at the interface between the CH2 and CH3 domains (*e.g.*, to increase the binding affinity to neonatal receptor FcRn and/or the half-life of the antibody molecule). In an embodiment, the Fc region comprises one or more mutations, *e.g.*, one or more (*e.g.*, 2, 3, 4, 5, 6 or all) mutations chosen from T250Q, M252Y, S254T, T256E, M428L, H433K, N434F, or any combination thereof, of IgG1. In an embodiment, the Fc region comprises one or more mutations at positions 233-236 or 322 of human IgG1 or IgG2, or one or more substitutions at positions 327, 330 or 331 of human IgG4 (*e.g.*, to reduce complement-dependent cytotoxicity (CDC)). In an embodiment, the Fc region comprises one or more (*e.g.*, 2, 3, 4, 5, 6, 7 or all) mutations chosen from E233P, L234V, L235A, G236, K322A, A327G, A330S, P331S, or any combination thereof.

In an embodiment, the antibody molecule is a humanized antibody molecule, *e.g.*, comprising one or more framework regions derived from human framework germline sequence. In an embodiment, the antibody molecule comprises a heavy chain variable region (VH) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises a light chain variable region (VL) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises a heavy chain variable region (VH) and a light chain variable region (VL) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one, two, or three CDRs of a heavy chain variable region (VH) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one, two, or three CDRs of a light chain variable region (VL) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one, two, or three CDRs of a heavy chain variable region (VH) described in **Table 1 or 5**, and one, two, or three CDRs of a light chain variable region (VL) described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises two heavy chain variable regions and two light chain variable regions. In an embodiment, the antibody molecule is a Fab, F(ab')₂, Fv, Fd, or a single chain Fv fragment (scFv).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the

amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); or (ii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,

or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70).

In an embodiment the antibody molecule is monoclonal antibody 3530. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3530.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid

residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprises the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50).

In an embodiment the antibody molecule is monoclonal antibody 3525. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3525.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid

residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprises the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122).

In an embodiment the antibody molecule is monoclonal antibody 3833. In an embodiment, monoclonal antibody 3833 is a humanized monoclonal antibody 3833. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 246-250, a VL comprising the amino acid sequence of any of SEQ ID NO: 251-253, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an

HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprises the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132).

In an embodiment the antibody molecule is monoclonal antibody 3631. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3631.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid

residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 138); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 138); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprises the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 138); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141).

In an embodiment the antibody molecule is monoclonal antibody 3732. In an embodiment, monoclonal antibody 3732 is a humanized monoclonal antibody 3732.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology

with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises: (i) a VH comprising one, two, or all of the following: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and (ii) a VL comprising one, two, or all of the following: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or
 5 has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256); and (ii) a VL comprising the amino
 10 acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261).

In an embodiment, the antibody molecule is monoclonal antibody 4540, 4540-063, or 4540-033. In an embodiment, monoclonal antibody 4540 is a humanized monoclonal antibody 4540 (*e.g.*, antibodies 4540-063 or 4540-033). In an embodiment, the antibody molecule comprises a VH
 15 comprising the amino acid sequence of any of SEQ ID NO: 254-258, a VL comprising the amino acid sequence of any of SEQ ID NO: 259-261, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within
 20 a region of human APRIL as defined in any of **Tables 3-4 or 7-8**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9,
 25 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as
 30 shown in **Table 3**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, residues within a region of human APRIL as defined in **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all, of the human APRIL residues from **Table**
 35 **4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises one or

more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, residues within a region of human APRIL as defined in **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 7**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, residues within a region of human APRIL as defined in **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 8**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL from positions 96-105. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one, two or all of Asp129, Arg233, or His203 of human APRIL. In an embodiment, the epitope is a conformational epitope.

In an embodiment, binding of the antibody molecule to APRIL (*e.g.*, human APRIL) inhibits, or substantially inhibits, the binding of the CRD2 domain of TACI (*e.g.*, human TACI) to APRIL (*e.g.*, human APRIL). In another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the APRIL residues from **Table 3**. In yet another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all, of the human APRIL residues from **Table 4**. In still another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In still another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**. In another embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of human BCMA, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11,

12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**.

In an aspect, the disclosure features an antibody molecule that binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in any of **Tables 3-4 or 7-8**.

In an embodiment, the anti-APRIL antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, of the human APRIL residues from any of **Tables 3-4 or 7-8**. In an embodiment, the antibody molecule binds, or substantially binds, to a conformational epitope.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in **Table 3**. In an embodiment, the anti-APRIL antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 3**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all, residues within a region of human APRIL as defined in **Table 4**. In an embodiment, the epitope comprises consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the APRIL residues from **Table 4**. In an embodiment, the epitope comprises or consists of one or more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, residues within a region of human APRIL as defined in **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 7**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, residues within a region of human APRIL as defined in **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from

Table 8. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 8**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL from positions 96-105. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one, two or all of Asp129, Arg233, or His203 of human APRIL.

In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, or all) of human APRIL residues from **Table 6**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, or all) of the amino acid residues of human APRIL chosen from V174, F176, Q190, R195, R206, or Y208. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from V181, S226, I228, or N237. In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or I228. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one or both of the amino acid residues of human APRIL chosen from Y208 or N237. In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, or all) of the amino acid residues of human APRIL chosen from V174, R206, or Y208. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or N237.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL and mouse APRIL. In an embodiment, the antibody molecule binds, or substantially binds to, human APRIL, but does not bind to mouse APRIL, or binds to mouse APRIL with low affinity.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, *e.g.*, 10 nM

or less, *e.g.*, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule does not bind to mouse APRIL, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both). In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, binding of the antibody molecule to APRIL (*e.g.*, human APRIL) inhibits, or substantially inhibits, the binding of the CRD2 domain of TACI (*e.g.*, human TACI) to APRIL (*e.g.*, human APRIL). In an embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 3**. In an embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all, of the human APRIL residues from **Table 4**. In an embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In an embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**. In another embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of

human BCMA, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both).

5 In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), *e.g.*, at an IC₅₀ of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule is a synthetic antibody molecule. In an embodiment, the antibody molecule is an isolated antibody molecule. In an embodiment, the antibody molecule is an IgG antibody molecule, *e.g.*, comprising a heavy chain constant region of IgG, *e.g.*, chosen from IgG1, IgG2 (*e.g.*, IgG2a), IgG3, or IgG4, *e.g.*, IgG2 or IgG4. In an embodiment, the antibody molecule is an IgG1 antibody molecule. In an embodiment, the antibody molecule is an IgG2 antibody molecule. In an embodiment, the antibody molecule comprises a light chain constant region of kappa or lambda light chain.

In an embodiment, the antibody molecule comprises an Fc region. In an embodiment, the Fc region comprises one or more mutations located at the interface between the CH2 and CH3 domains (*e.g.*, to increase the binding affinity to neonatal receptor FcRn and/or the half-life of the antibody molecule). In an embodiment, the Fc region comprises one or more mutations, *e.g.*, one or more (*e.g.*, 2, 3, 4, 6 or all) mutations chosen from T250Q, M252Y, S254T, T256E, M428L, H433K, N434F, or any combination thereof, of IgG1. In an embodiment, the Fc region comprises one or more mutations at positions 233-236 or 322 of human IgG1 or IgG2, or one or more substitutions at positions 327, 330 or 331 of human IgG4 (*e.g.*, to reduce complement-dependent cytotoxicity (CDC)). In an embodiment, the Fc region comprises one or more (*e.g.*, 2, 3, 4, 6 7 or all) mutations chosen from E233P, L234V, L235A, G236, K322A, A327G, A330S, P331S, or any combination thereof.

In an embodiment, the antibody molecule is a humanized antibody molecule, *e.g.*, as described in **Table 1 or 5**, *e.g.*, comprising one or more framework regions derived from human framework germline sequence.

In an embodiment, the antibody molecule comprises two heavy chain variable regions and two light chain variable regions. In an embodiment, the antibody molecule is a Fab, F(ab')₂, Fv, Fd, or a single chain Fv fragment (scFv).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 1); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 2); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 4); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 1); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 2); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*,

SEQ ID NO: 4); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 1); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 2); and an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3), and (ii) a VL comprising: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 4); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); and an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises a VH comprising one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 7); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 8); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3).

In an embodiment, the antibody molecule comprises a VL comprising one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 4); (ii) an LCDR2

comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 7); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 8); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 4); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 7); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 8); and an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3), and (ii) a VL comprising: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 4); an LCDR2 comprising

an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); and an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 9). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 10).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 9); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 10). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 9); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 10).

In an embodiment the antibody molecule is monoclonal antibody 2218. In an embodiment, monoclonal antibody 2218 is a humanized monoclonal antibody 2218. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 190-201, a VL comprising the amino acid sequence of any of SEQ ID NO: 202-208, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 11) or a 2419-related antibody; (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 12) or a 2419-related

antibody; or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody.

5 In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino
10 acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15) or a 2419-related antibody; or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3
15 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid
20 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 11) or a 2419-related antibody; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID
25 NO: 12) or a 2419-related antibody; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody, and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid
30 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID
35 NO: 15) or a 2419-related antibody; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology

with, the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 11) or a 2419-related antibody; an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 12) or a 2419-related antibody; or an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody, and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15) or a 2419-related antibody; and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 17) or a 2419-related antibody; (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 18) or a 2419-related antibody; or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15) or a 2419-related antibody; or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid

sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 17) or a 2419-related antibody; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 18) or a 2419-related antibody; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody, and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15) or a 2419-related antibody; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 17) or a 2419-related antibody; an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 18) or a 2419-related antibody; or an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13) or a 2419-related antibody, and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14) or a 2419-related antibody; an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15) or a 2419-related antibody; and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid

sequence of the VH of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 19) or a 2419-related antibody. In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 20) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 19) or a 2419-related antibody; and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 20) or a 2419-related antibody.

In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 19) or a 2419-related antibody; and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 20) or a 2419-related antibody.

In an embodiment the antibody molecule is monoclonal antibody 2419. In an embodiment, monoclonal antibody 2419 is a humanized monoclonal antibody 2419. In an embodiment, the antibody molecule is a 2419-related antibody molecule, *e.g.*, any of antibodies 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, or 2419-1406, *e.g.*, as disclosed in **Table 1 or 5**. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NOS: 209-214, 283, 288, 289, 291, 292, 294, 296, or 317, a VL comprising the amino acid sequence of any of SEQ ID NOS: 215-219, 284, 286, 295, or 316, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 21); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 32); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least

85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 21); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 32); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 21); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 32); and an HCDR3 comprising the amino acid sequence of the HCDR3 of

monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 37); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 38); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 37); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 38); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,

or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 37); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 38); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 39). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 40).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 39); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 40). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the

VH of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 39); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 40).

In an embodiment the antibody molecule is monoclonal antibody 2922. In an embodiment, the antibody molecule is a humanized monoclonal antibody 2922.

5

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 51); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 52); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 51); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 52); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,

or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 51); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 52); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 57); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 58); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino

acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); or (iii) an LCDR3 comprising an amino acid
5 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid
10 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 57); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 58); or an HCDR3
15 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99
20 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,
25 or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 57); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3327
30 (*e.g.*, SEQ ID NO: 58); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal
35 antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 59). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 60).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 59); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 60). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 59); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 60).

In an embodiment the antibody molecule is monoclonal antibody 3327. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3327.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); (ii) an LCDR2

comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one,

two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or an LCDR3

comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 65); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70).

In an embodiment the antibody molecule is monoclonal antibody 3530. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3530.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more

than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100%

homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 64); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 65); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50).

In an embodiment the antibody molecule is monoclonal antibody 3525. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3525.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 21); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 22); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 21); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 22); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 21); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 22); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 27); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 28); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least

85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 27); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 28); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 27); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 28); and an HCDR3 comprising the amino acid sequence of the HCDR3 of

monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 29). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 30).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 29); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 30). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 29); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 30).

In an embodiment the antibody molecule is monoclonal antibody 2621. In an embodiment, the antibody molecule is a humanized monoclonal antibody 2621.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 11); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 42); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least

85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 11); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 42); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 11); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 42); and an HCDR3 comprising the amino acid sequence of the HCDR3 of

monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 47); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 48); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 47); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 48); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,

or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 47); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 48); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 49). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 50).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 49); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 50). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the

VH of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 49); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 50).

In an embodiment the antibody molecule is monoclonal antibody 3125. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3125.

5

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 93); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 94); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 93); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 94); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino

acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 93); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 94); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 99); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 100) or 4035-062 (*e.g.*, SEQ ID NO: 273); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than

1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 99); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 100) or 4035-062 (*e.g.*, SEQ ID NO: 273); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 99); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 100) or 4035-062 (*e.g.*, SEQ ID NO: 273); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 95), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 97); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

97); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4035 or 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 101) or 4035-062 (*e.g.*, SEQ ID NO: 225). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 102) or 4035-062 (*e.g.*, SEQ ID NO: 229).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 101) or 4035-062 (*e.g.*, SEQ ID NO: 225); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 102) or 4035-062 (*e.g.*, SEQ ID NO: 229). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 101) or 4035-062 (*e.g.*, SEQ ID NO: 225); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 102) or 4035-062 (*e.g.*, SEQ ID NO: 229).

In an embodiment, the antibody molecule is monoclonal antibody 4035. In an embodiment, monoclonal antibody 4035 is a humanized monoclonal antibody 4035 (*e.g.*, antibody 4035-062). In another embodiment, the antibody molecule is antibody 4035-062. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NOS: 220-227 or 262-265, a VL comprising the amino acid sequence of any of SEQ ID NOS: 228-234, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 103); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the

HCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 104); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 105).

5 In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino
10 acid sequence of the LCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 106); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 107); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99
15 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99
20 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 103); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 104); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 105), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3934 (*e.g.*,
30 SEQ ID NO: 106); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 107); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of
35 monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO:

103); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3934 (e.g., SEQ ID NO: 104); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (e.g., SEQ ID NO: 105), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3934 (e.g., SEQ ID NO: 106); an
 5 LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (e.g., SEQ ID NO: 107); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (e.g., SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining
 10 regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (e.g., SEQ ID NO: 109); (ii) an
 HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid
 15 residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3934 (e.g., SEQ ID NO: 110); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (e.g., SEQ ID NO: 105).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining
 20 regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino
 25 acid sequence of the LCDR1 of monoclonal antibody 3934 (e.g., SEQ ID NO: 106); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (e.g., SEQ ID NO: 107); or (iii) an LCDR3 comprising an amino acid
 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99
 30 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (e.g., SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid
 35 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (e.g., SEQ ID NO: 109); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid

sequence of the HCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 110); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 105), and

5 (ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 106); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid
10 sequence of the LCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 107); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1
15 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 109); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 110); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 105), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 106); an
20 LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 107); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid
25 residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 111). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3934
30 (*e.g.*, SEQ ID NO: 112).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 111); and (ii) a VL comprising
35 an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 112). In an

embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 111); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 112).

In an embodiment the antibody molecule is monoclonal antibody 3934. In an embodiment,
5 the antibody molecule is a humanized monoclonal antibody 3934.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one,
10 two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 112); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid
15 HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114).

In an embodiment, the antibody molecule comprises a light chain variable region (VL),
20 wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115); (ii) an LCDR2
25 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*,
30 SEQ ID NO: 117).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*,
35 SEQ ID NO: 113); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114); or an HCDR3

comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than

1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 119); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 120); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122).

In an embodiment the antibody molecule is monoclonal antibody 3833. In an embodiment, monoclonal antibody 3833 is a humanized monoclonal antibody 3833. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 246-250, a VL comprising the amino acid sequence of any of SEQ ID NO: 251-253, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining

regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); (ii) an LCDR2
 5 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*,
 10 SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*,
 15 SEQ ID NO: 123); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of
 20 monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2,
 25 or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and (ii) a VL comprising: an LCDR1 comprising
 30 the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ

ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO:45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 129); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 130); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132).

In an embodiment the antibody molecule is monoclonal antibody 3631. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3631.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH),
5 wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); (ii) an
10 HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal
15 antibody 3732 (*e.g.*, SEQ ID NO: 135).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than
20 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid
25 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid
30 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); or an HCDR3
35 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 133); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 138); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,

or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*,
 5 SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*,
 10 SEQ ID NO: 138); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of
 15 monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2,
 20 or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 138); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 139); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and (ii) a VL comprising: an LCDR1 comprising
 30 the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid
 35 sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140). In an embodiment, the

antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141).

5 In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15
10 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141).

15 In an embodiment the antibody molecule is monoclonal antibody 3732. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3732.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining
20 regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 11); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid
25 residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 142); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143).

30 In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino
35 acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the

LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

5 In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 11); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or
10 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 142); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); an LCDR2 comprising an amino acid sequence that differs by no more than
15 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1
25 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 11); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 142); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146);
30 an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining
35 regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the

amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 149); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 150); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 149); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 150); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid

residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 149); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 150); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 151). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 152 or 153).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 151); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 152 or 153). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 150); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 152 or 153).

In an embodiment the antibody molecule is monoclonal antibody 4338. In an embodiment, the antibody molecule is a humanized monoclonal antibody 4338.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the

amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid

residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises: (i) a VH comprising three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 154); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 155); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two,

or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); (ii) an LCDR2 comprising an amino acid
 5 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the
 10 LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*,
 15 SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or an HCDR3 comprising an amino acid sequence that differs by
 20 no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99
 25 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or
 30 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises: (i) a VH comprising one, two, or all of
 35 the following: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 159), 4540-063 (*e.g.*, SEQ ID NO: 276), or 4540-033 (*e.g.*, SEQ ID NO: 159); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody

4540 (*e.g.*, SEQ ID NO: 160), 4540-063 (*e.g.*, SEQ ID NO: 277), or 4540-033 (*e.g.*, SEQ ID NO: 278); or an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 156), and (ii) a VL comprising one, two, or all of the following: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody
 5 4540 (*e.g.*, SEQ ID NO: 116), 4540-063 (*e.g.*, SEQ ID NO: 274), or 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157), 4540-063 (*e.g.*, SEQ ID NO: 275), or 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4540, 4540-063, or 4540-033 (*e.g.*, SEQ ID NO: 158).

10 In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256). In an embodiment, the antibody molecule comprises
 15 a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid
 20 sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or
 25 has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161), 4540-063 (*e.g.*, SEQ ID NO: 258), or 4540-033 (*e.g.*, SEQ ID NO: 256); and (ii) a VL comprising the amino
 30 acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162), 4540-063 (*e.g.*, SEQ ID NO: 261), or 4540-033 (*e.g.*, SEQ ID NO: 261).

In an embodiment the antibody molecule is monoclonal antibody 4540, 4540-063, or 4540-033. In an embodiment, monoclonal antibody 4540 is a humanized monoclonal antibody 4540 (*e.g.*, antibodies 4540-063 or 4540-033). In an embodiment, the antibody molecule comprises a VH
 35 comprising the amino acid sequence of any of SEQ ID NOS: 254-258, a VL comprising the amino acid sequence of any of SEQ ID NOS: 259-261, or both.

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 163); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 164); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 163); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 164); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*,

SEQ ID NO: 166); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 163); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 164); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: (i) an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 169); (ii) an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 170); or (iii) an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165).

In an embodiment, the antibody molecule comprises a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: (i) an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); (ii) an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); or (iii) an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99

or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises:

(i) a VH comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 169); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 170); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165), and

(ii) a VL comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises: (i) a VH comprising: an HCDR1 comprising the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 169); an HCDR2 comprising the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 170); and an HCDR3 comprising the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165), and (ii) a VL comprising: an LCDR1 comprising the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); an LCDR2 comprising the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); and an LCDR3 comprising the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 171). In an embodiment, the antibody molecule comprises a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96,

97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 172).

In an embodiment, the antibody molecule comprises: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 171); and (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 172). In an embodiment, the antibody molecule comprises: (i) a VH comprising the amino acid sequence of the VH of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 171); and (ii) a VL comprising the amino acid sequence of the VL of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 172).

In an embodiment the antibody molecule is monoclonal antibody 4237. In an embodiment, monoclonal antibody 4237 is a humanized monoclonal antibody 4237. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NOS: 235-240, a VL comprising the amino acid sequence of any of SEQ ID NOS: 241-245, or both.

In another aspect, the disclosure features an anti-APRIL antibody molecule, which:

(i) binds, or substantially binds, to human APRIL;

(ii) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both);

(iii) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both); and

(iv) binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in any of **Tables 3-4 or 7-8**.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC₅₀ of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule does not bind to mouse APRIL, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), *e.g.*, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to one or more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both. In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL from positions 96-105. In an embodiment, the antibody molecule does not bind, or binds with low affinity, to one, two or all of Asp129, Arg233, or His203 of human APRIL.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH) comprising one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of a monoclonal antibody chosen from antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3125, 2621, 4035, 4035-062, 3934, 4338, 4439, or

4237; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of the (same) monoclonal antibody; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of the (same) monoclonal antibody, or

(ii) a light chain variable region (VL) comprising one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of the (same) monoclonal antibody; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of the (same) monoclonal antibody; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of the (same) monoclonal antibody.

In an embodiment, the antibody molecule comprises one or both of:

(i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of a monoclonal antibody chosen from antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3125, 2621, 4035, 4035-062, 3934, 4338, 4439, or 4237; or

(ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of the (same) monoclonal antibody.

In an embodiment, the antibody molecule is a synthetic antibody molecule. In an embodiment, the antibody molecule is an isolated antibody molecule. In an embodiment, the antibody molecule is a humanized antibody molecule, *e.g.*, comprising one or more framework regions derived from human framework germline sequence.

In an embodiment, the antibody molecule is an IgG antibody molecule, *e.g.*, comprising a heavy chain constant region of IgG, *e.g.*, chosen from IgG1, IgG2 (*e.g.*, IgG2a), IgG3, or IgG4, *e.g.*, IgG2 or IgG4. In an embodiment, the antibody molecule is an IgG1 antibody molecule. In an embodiment, the antibody molecule is an IgG2 antibody molecule. In an embodiment, the antibody molecule comprises a light chain constant region of kappa or lambda light chain.

In an embodiment, the antibody molecule comprises an Fc region. In an embodiment, the Fc region comprises one or more mutations located at the interface between the CH2 and CH3 domains (*e.g.*, to increase the binding affinity to neonatal receptor FcRn and/or the half-life of the antibody molecule). In an embodiment, the Fc region comprises one or more mutations, *e.g.*, one or more (*e.g.*,

2, 3, 4, 6 or all) mutations chosen from T250Q, M252Y, S254T, T256E, M428L, H433K, N434F, or any combination thereof, of IgG1. In an embodiment, the Fc region comprises one or more mutations at positions 233-236 or 322 of human IgG1 or IgG2, or one or more substitutions at positions 327, 330 or 331 of human IgG4 (*e.g.*, to reduce complement-dependent cytotoxicity (CDC)). In an
 5 embodiment, the Fc region comprises one or more (*e.g.*, 2, 3, 4, 6 7 or all) mutations chosen from E233P, L234V, L235A, G236, K322A, A327G, A330S, P331S, or any combination thereof.

In an embodiment, the antibody molecule comprises two heavy chain variable regions and two light chain variable regions. In an embodiment, the antibody molecule is a Fab, F(ab')₂, Fv, Fd, or a single chain Fv fragment (scFv).

10 In an aspect, the disclosure features an anti-APRIL antibody, which:

a) competes for binding to APRIL with an antibody molecule comprising the heavy chain complementary determining regions (HCDR1, HCDR2 and HCDR3) and the light chain complementary determining regions (LCDR1, LCDR2 and LCDR3) of any of monoclonal antibodies
 15 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237, *e.g.*, as described in **Table 1 or 5**; or

b) binds, or substantially binds, to an epitope that completely or partially overlaps with the
 20 epitope of an antibody molecule comprising the heavy chain complementary determining regions (HCDR1, HCDR2 and HCDR3) and the light chain complementary determining regions (LCDR1, LCDR2 and LCDR3) of any of monoclonal antibodies 2218 (*e.g.*, SEQ ID NOS: 1-6 according to Chothia numbering or SEQ ID NOS: 3-8 according to Kabat numbering), 2419 (*e.g.*, SEQ ID NOS: 11-16 according to Chothia numbering or SEQ ID NOS: 13-18 according to Kabat numbering), 2419-
 25 0105 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17 and 280-282 according to Kabat numbering), 2419-0205 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17 and 280-282 according to Kabat numbering), 2419-0206 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 285 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 282 and 285 according to Kabat numbering), 2419-0406
 30 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 285 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 285 and 290 according to Kabat numbering), 2419-0605 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17 and 280-282 according to Kabat numbering), 2419-0805 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 281 and 287 according to Kabat numbering), 2419-0806 (*e.g.*, SEQ
 35 ID NOS: 11-13, 16, 280 and 285 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 285 and 287 according to Kabat numbering), 2419-1204 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 293 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 282 and 293 according to Kabat

numbering), 2419-1205 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17 and 280-282 according to Kabat numbering), 2419-1210 (*e.g.*, SEQ ID NOS: 11-13, 16, 314 and 315 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 282, 314 and 315 according to Kabat numbering), 2419-1305 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 281 according to Chothia numbering or SEQ ID NOS: 13, 16, 17 and 280-282 according to Kabat numbering), 2419-1306 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 285 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 282 and 285 according to Kabat numbering), 2419-1310 (*e.g.*, SEQ ID NOS: 11-13, 16, 314 and 315 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 282, 314 and 315 according to Kabat numbering), 2419-1406 (*e.g.*, SEQ ID NOS: 11-13, 16, 280 and 285 according to Chothia numbering or SEQ ID NOS: 13, 16, 17, 280, 282 and 285 according to Kabat numbering), 2922 (*e.g.*, SEQ ID NOS: 21 and 32-36 according to Chothia numbering or SEQ ID NOS: 33-38 according to Kabat numbering), 3327 (*e.g.*, SEQ ID NOS: 51-56 according to Chothia numbering or SEQ ID NOS: 53-58 according to Kabat numbering), 3530 (*e.g.*, SEQ ID NOS: 61-63, 67, 45 and 46 according to Chothia numbering or SEQ ID NOS: 63-65, 67, 45 and 46 according to Kabat numbering), 3525 (*e.g.*, SEQ ID NOS: 44-46 and 61-63 according to Chothia numbering or SEQ ID NOS: 44-46 and 63-65 according to Kabat numbering), 3125 (*e.g.*, SEQ ID NOS: 11 and 42-46 according to Chothia numbering or SEQ ID NOS: 43-48 according to Kabat numbering), 2621 (*e.g.*, SEQ ID NOS: 21-26 according to Chothia numbering or SEQ ID NOS: 23-28 according to Kabat numbering), 4035 (*e.g.*, SEQ ID NOS: 93-98 according to Chothia numbering or SEQ ID NOS: 95-100 according to Kabat numbering), 4035-062 (*e.g.*, SEQ ID NOS: 93-98 according to Chothia numbering or SEQ ID NOS: 95-99 and 273 according to Kabat numbering), 3934 (*e.g.*, SEQ ID NOS: 103-108 according to Chothia numbering or SEQ ID NOS: 105-110 according to Kabat numbering), 3833 (*e.g.*, SEQ ID NOS: 113-118 according to Chothia numbering or SEQ ID NOS: 115-120 according to Kabat numbering), 3631 (*e.g.*, SEQ ID NOS: 123-128 according to Chothia numbering or SEQ ID NOS: 125-130 according to Kabat numbering), 3732 (*e.g.*, SEQ ID NOS: 127 and 133-137 according to Chothia numbering or SEQ ID NOS: 127 and 135-139 according to Kabat numbering), 4338 (*e.g.*, SEQ ID NOS: 11, 107 and 142-145, or SEQ ID NOS: 11, 142, 143 and 146-148 according to Chothia numbering; or SEQ ID NOS: 107, 143-145 and 149-150, or SEQ ID NOS: 143 and 146-150 according to Kabat numbering), 4540 (*e.g.*, SEQ ID NOS: 116 and 154-158 according to Chothia numbering or SEQ ID NOS: 116 and 156-160 according to Kabat numbering), 4540-063 (*e.g.*, SEQ ID NOS: 154-156, 158, 274 and 275 according to Chothia numbering or SEQ ID NOS: 156, 158 and 274-277 according to Kabat numbering), 4540-033 (*e.g.*, SEQ ID NOS: 154-156, 158, 274 and 275 according to Chothia numbering or SEQ ID NOS: 156, 158, 159, 274, 275 and 278 according to Kabat numbering), 4439 (*e.g.*, SEQ ID NOS: 146-148 and 266-268 according to Chothia numbering or SEQ ID NOS: 146-148 and 269-270 according to Kabat numbering), or 4237 (*e.g.*, SEQ ID NOS: 163-168 according to Chothia numbering or SEQ ID NOS: 165-170 according to Kabat numbering), *e.g.*, as described in **Table 1 or 5**.

In an embodiment, the antibody molecule is a synthetic antibody molecule. In an embodiment, the antibody molecule is an isolated antibody molecule.

In an embodiment, the antibody molecule competes for binding with two, three, four, five, six, seven, eight, nine, ten, or more of the antibody molecules that comprise the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3 of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237.

In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that completely or partially overlaps with the epitopes of two, three, four, five, six, seven, eight, nine, ten, or more of the antibody molecules that comprise the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3 of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237.

In an embodiment, the antibody molecule that comprises the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3 of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 3934, 3833, 3631, 3732, 4338, 4540, 4439, or 4237 comprises a heavy chain variable region and a light chain variable region of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 3934, 3833, 3631, 3732, 4338, 4540, 4439, or 4237.

In an embodiment, the antibody molecule that comprises the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3 of any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 3934, 3833, 3631, 3732, 4338, 4540, 4439, or 4237 is monoclonal antibody 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 3934, 3833, 3631, 3732, 4338, 4540, 4439, or 4237.

In an embodiment, the antibody molecule is a humanized monoclonal antibody 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237. In an embodiment, the antibody molecule comprises a heavy chain variable region (VH) having an

amino acid sequence described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises a light chain variable region (VL) having an amino acid sequence described in **Table 1 or 5**. In an antibody molecule comprises a heavy chain variable region (VH) having an amino acid sequence described in **Table 1 or 5** and a light chain variable region (VL) having an amino acid sequence described in **Table 1 or 5**.

In an embodiment, the antibody molecule competes for binding to human APRIL, mouse APRIL, or both. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or

less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

5 In an embodiment, binding of the antibody molecule to APRIL (*e.g.*, human APRIL) inhibits, or substantially inhibits, the binding of the CRD2 domain of TACI (*e.g.*, human TACI) to APRIL (*e.g.*, human APRIL).

In an embodiment, binding of the antibody molecule to human APRIL inhibits, or substantially inhibits, the binding of human TACI to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 10 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 3**. In an embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the human APRIL residues from **Table 4**. In an embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 15 10, 11, 12, 13, 14, 15, 16, 17, or all of the human APRIL residues from **Table 7**. In an embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 8**.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL 20 (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both).

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), *e.g.*, at an IC₅₀ of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM 25 or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

30 In an embodiment, the antibody molecule does not inhibit, or does not substantially inhibit, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both).

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within 35 a region of human APRIL as defined in any of **Tables 3-4 or 7-8**. In an embodiment, the antibody molecule binds, or substantially binds, to a conformational epitope.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in **Table 3**. In an embodiment, the antibody molecule binds to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 3**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, residues within a region of human APRIL as defined in **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises one or more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, residues within a region of human APRIL as defined in **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 7**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, residues within a region of human APRIL as defined in **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 8**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL from positions 96-105. In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one, two or all of Asp129, Arg233, or His203 of human APRIL.

In an embodiment, the antibody molecule is an IgG antibody molecule, *e.g.*, comprising a heavy chain constant region of IgG, *e.g.*, chosen from IgG1, IgG2 (*e.g.*, IgG2a), IgG3, or IgG4, *e.g.*, IgG2 or IgG4. In an embodiment, the antibody molecule is an IgG1 antibody molecule. In another

embodiment, the antibody molecule is an IgG2 antibody molecule. In an embodiment, the antibody molecule comprises a light chain constant region of kappa or lambda light chain.

In an embodiment, the antibody molecule comprises an Fc region. In an embodiment, the Fc region comprises one or more mutations located at the interface between the CH2 and CH3 domains (e.g., to increase the binding affinity to neonatal receptor FcRn and/or the half-life of the antibody molecule). In an embodiment, the Fc region comprises one or more mutations, e.g., one or more (e.g., 2, 3, 4, 6 or all) mutations chosen from T250Q, M252Y, S254T, T256E, M428L, H433K, N434F, or any combination thereof, of IgG1. In an embodiment, the Fc region comprises one or more mutations at positions 233-236 or 322 of human IgG1 or IgG2, or one or more substitutions at positions 327, 330 or 331 of human IgG4 (e.g., to reduce complement-dependent cytotoxicity (CDC)). In an embodiment, the Fc region comprises one or more (e.g., 2, 3, 4, 6 or all) mutations chosen from E233P, L234V, L235A, G236, K322A, A327G, A330S, P331S, or any combination thereof.

In an embodiment, the antibody molecule is a humanized antibody molecule, e.g., comprising one or more framework regions derived from human framework germline sequence. In an embodiment, the antibody molecule comprises two heavy chain variable regions and two light chain variable regions. In an embodiment, the antibody molecule is a Fab, F(ab')2, Fv, Fd, or a single chain Fv fragment (scFv).

The disclosure features an anti-APRIL antibody molecule described herein, e.g., a synthetic or isolated anti-APRIL antibody molecule described herein.

In another aspect, the present disclosure features an anti-APRIL antibody molecule, comprising one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following:

an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of a monoclonal antibody chosen from 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237;

an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of the monoclonal antibody; or

an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of the monoclonal antibody, or

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following:

an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of the monoclonal antibody;

an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of the monoclonal antibody; or

an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of the monoclonal antibody; and/or

which comprises one or both of:

(i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of the monoclonal antibody; or

(ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of the monoclonal antibody.

In another aspect, the present disclosure features an anti-APRIL antibody molecule, comprising one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following:

an HCDR1 comprising an amino acid sequence of G-Y-T-F-T-D-Y (SEQ ID NO: 11);

an HCDR2 comprising an amino acid sequence of Y-P-L-R-G-S (SEQ ID NO: 12); or

an HCDR3 comprising an amino acid sequence of H-G-A-Y-Y-S-N-A-F-D-Y (SEQ ID NO: 13), or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following:

an LCDR1 comprising an amino acid sequence of X1-X2-S-X4-S-V-D-N-D-G-I-R-F-X14-H (SEQ ID NO: 327), wherein X1 is R or K; X2 is A or S; X4 is E or Q; and X14 is M or L;

an LCDR2 comprising an amino acid sequence of R-A-S-X4-X5-X6-X7 (SEQ ID NO: 328), wherein X4 is N or T; X5 is L or R; X6 is E or A; and X7 is S or T; or

an LCDR3 comprising an amino acid sequence of Q-Q-S-N-K-D-P-Y-T (SEQ ID NO: 16); or comprising one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following:

an HCDR1 comprising an amino acid sequence of D-Y-T-I-H (SEQ ID NO: 17);

5 an HCDR2 comprising an amino acid sequence of W-I-Y-P-L-R-G-S-I-N-Y-X12-X13-X14-F-X16-X17 (SEQ ID NO: 329), wherein X12 is N, S, or A, X13 is E, P, or Q; X14 is K or S; X16 is K or Q; and X17 is D or G; or

an HCDR3 comprising an amino acid sequence of H-G-A-Y-Y-S-N-A-F-D-Y (SEQ ID NO: 13), or

10 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following:

an LCDR1 comprising an amino acid sequence of X1-X2-S-X4-S-V-D-N-D-G-I-R-F-X14-H (SEQ ID NO: 327), wherein X1 is R or K; X2 is A or S; X4 is E or Q; and X14 is M or L;

15 an LCDR2 comprising an amino acid sequence of R-A-S-X4-X5-X6-X7 (SEQ ID NO: 328), wherein X4 is N or T; X5 is L or R; X6 is E or A; and X7 is S or T; or

an LCDR3 comprising an amino acid sequence of Q-Q-S-N-K-D-P-Y-T (SEQ ID NO: 16).

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises
20 three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 1 or 7); an HCDR2 comprising an amino acid sequence that differs
25 by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 2 or 8); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 3), and

30 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2218

(*e.g.*, SEQ ID NO: 4); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 5); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 6).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 9); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2218 (*e.g.*, SEQ ID NO: 10).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 71 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 72 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 2218. In an embodiment, monoclonal antibody 2218 is humanized monoclonal antibody 2218. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 190-201, a VL comprising the amino acid sequence of any of SEQ ID NO: 202-208, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 1 nM or less, *e.g.*, about 0.6 nM. In an embodiment, the antibody molecule does not bind to mouse APRIL, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 1 nM or less, *e.g.*, about 0.74 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 0.5 nM or less, *e.g.*, about 0.22 nM.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence of G-Y-T-F-T-D-Y (SEQ ID NO: 11); an HCDR2 comprising an amino acid sequence of Y-P-L-R-G-S (SEQ ID NO: 12); or an HCCDR3 comprising an amino acid sequence of H-G-A-Y-Y-S-N-A-F-D-Y (SEQ ID NO: 13), or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence of X1-X2-S-X4-S-V-D-N-D-G-I-R-F-X14-H (SEQ ID NO: 327), wherein X1 is R or K; X2 is A or S; X4 is E or Q; and X14 is M or L; an LCDR2 comprising an amino acid sequence of R-A-S-X4-X5-X6-X7 (SEQ ID NO: 328), wherein X4 is N or T; X5 is L or R; X6 is E or A; and X7 is S or T; or an LCDR3 comprising an amino acid sequence of Q-Q-S-N-K-D-P-Y-T (SEQ ID NO: 16).

In another embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence of D-Y-T-I-H (SEQ ID NO: 17); an HCDR2 comprising an amino acid sequence of W-I-Y-P-L-R-G-S-I-N-Y-X12-X13-X14-F-X16-X17 (SEQ ID NO: 329), wherein X12 is N, S, or A, X13 is E, P, or Q; X14 is K or S; X16 is K or Q; and X17 is D or G; or an HCCDR3 comprising an amino acid sequence of H-G-A-Y-Y-S-N-A-F-D-Y (SEQ ID NO: 13), or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence of X1-X2-S-X4-S-V-D-N-D-G-I-R-F-X14-H (SEQ ID NO: 327), wherein X1 is R or K; X2 is A or S; X4 is E or Q; and X14 is M or L; an LCDR2 comprising an amino acid sequence of R-A-S-X4-X5-X6-X7 (SEQ ID NO: 328), wherein X4 is N or T; X5 is L or R; X6 is E or A; and X7 is S or T; or an LCDR3 comprising an amino acid sequence of Q-Q-S-N-K-D-P-Y-T (SEQ ID NO: 16).

In an embodiment, the antibody molecule is any of antibodies 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, or 2419-1406.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 0.01 nM or less, *e.g.*, about 0.001-0.005 nM or 0.002-0.004 nM, *e.g.*, about 0.001, 0.002, 0.003, 0.004, or 0.005 nM. In an embodiment, the antibody molecule does not bind to mouse APRIL, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1

nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 0.5 nM or less, *e.g.*, about 0.1-0.5 nM or 0.2-0.4 nM, *e.g.*, about 0.1, 0.2, 0.3, 0.4, or 0.5 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 0.5 nM or less, *e.g.*, about 0.1-0.5 nM or 0.2-0.4 nM, *e.g.*, about 0.1, 0.2, 0.3, 0.4, or 0.5 nM.

In another embodiment, the antibody molecule comprises (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 11 or 17); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 12 or 18); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 13), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 14); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino

acid sequence of the LCDR2 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 15); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 16).

5 In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 19); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,
10 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2419 (*e.g.*, SEQ ID NO: 20).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 73 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 74 (or a nucleotide sequence substantially
15 identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 2419. In an embodiment, monoclonal antibody 2419 is humanized monoclonal antibody 2419. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NOS: 209-214, a VL comprising the amino acid sequence of any of SEQ ID NOS: 215-219, or both.

20 In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between
25 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 1 nM or less, *e.g.*, about 0.8 nM, about 0.003 nM, or about 0.002 nM. In
30 an embodiment, the antibody molecule does not bind, or bind to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 500 nM or more.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human
35 APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1

nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC₅₀ of 1 nM or less, *e.g.*,
 5 about 0.74 nM, about 0.4 nM, 0.3 nM, or 0.2 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC₅₀ of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6
 10 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method
 15 described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC₅₀ of 5 nM or less, *e.g.*, about 4 nM, about 2 nM, or about 1 nM, or 0.5 nM or less, *e.g.*, about 0.22 nM, about 1 nM, about 0.7 nM, about 0.3 nM, about 0.2 nM, or about 0.1 nM.

In another embodiment, the antibody molecule comprises:

20 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of a 2419-related
 25 antibody (*e.g.*, SEQ ID NO: 11 or 17); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of the 2419-related antibody (*e.g.*, SEQ ID NOS: 12, 282, 287, or 290); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid
 30 sequence of the HCDR3 of the 2419-related antibody (*e.g.*, SEQ ID NO: 13), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90,
 35 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of the 2419-related antibody (*e.g.*, SEQ ID NOS: 280 or 314); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with,

the amino acid sequence of the LCDR2 of the 2419-related antibody (*e.g.*, SEQ ID NOS: 281, 285, 293, or 315); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of the 2419-related antibody (*e.g.*, SEQ ID NO: 16).

5 In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of the 2419-related antibody (*e.g.*, SEQ ID NOS: 283, 288, 289, 291, 292, 294, 296, or 317); or (ii) a VL comprising an amino acid sequence that differs by no more than 1,
10 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of the 2419-related antibody (*e.g.*, SEQ ID NOS: 284, 286, 295, or 316).

In an embodiment, the antibody molecule comprises a VH encoded by the VH nucleotide sequence of the 2419-related antibody (*e.g.*, SEQ ID NOS: 304, 307, 308, 309, 310, 311, 313, or 319)
15 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the VL nucleotide sequence of the 2419-related antibody (*e.g.*, SEQ ID NOS: 305, 306, 312, or 318) (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the 2419-related antibody molecule is chosen from antibodies 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-
20 1210, 2419-1305, 2419-1306, 2419-1310, or 2419-1406. In an embodiment, the 2419-related antibody is humanized antibody molecule. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NOS: 209-214, 283, 288, 289, 291, 292, 294, 296, or 317, a VL comprising the amino acid sequence of any of SEQ ID NOS: 215-219, 284, 286, 295, or 316, or both.

25 In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as
30 determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 1 nM or less, *e.g.*, about 0.8 nM, about 0.003 nM, or about 0.002 nM.

In an embodiment, the antibody molecule does not bind, or bind to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 500 nM or more.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 1 nM or less, *e.g.*, about 0.74 nM, about 0.4 nM, 0.3 nM, or 0.2 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 5 nM or less, *e.g.*, about 4 nM, about 2 nM, or about 1 nM, or 0.5 nM or less, *e.g.*, about 0.22 nM, about 1 nM, about 0.7 nM, about 0.3 nM, about 0.2 nM, or about 0.1 nM.

In another embodiment, the antibody molecule comprises (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 21 or 37); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 32 or 38); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 33), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 34); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 35); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 36).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 39); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2922 (*e.g.*, SEQ ID NO: 40).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 77 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 78 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 2922. In an embodiment, the antibody molecule is humanized monoclonal antibody 2922.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 5 nM or less, *e.g.*, about 3.3 nM. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 50 nM or less, *e.g.*, about 31.64 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the IC_{50} is 50 nM or less. In an embodiment, the antibody molecule inhibits binding of human TACI to human BCMA at an IC_{50} of 25 nM or less, *e.g.*, about 21.96 nM.

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 51 or 57); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 52 or 58); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 53), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 54); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 55); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 56).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 59); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3327 (*e.g.*, SEQ ID NO: 60).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 81 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 82 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3327. In an embodiment, the antibody molecule is humanized antibody 3327.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 3.16 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the IC_{50} is 50 nM or less. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 5 nM or less, *e.g.*, about 2.35 nM.

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 93 or 99); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 94 or 100); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 95), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 97); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 101); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4035 (*e.g.*, SEQ ID NO: 102).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 173 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 174 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4035. In an embodiment, monoclonal antibody 4035 is humanized monoclonal antibody 4035. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 220-227 or 262-265, a VL comprising the amino acid sequence of any of SEQ ID NO: 228-234, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC₅₀ of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC₅₀ of 0.01 nM or less, *e.g.*, about 0.001-0.002 nM. In an embodiment, the antibody molecule does

not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 3.16 nM, or about 0.1-0.5 nM or 0.2-0.4 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 5 nM or less, *e.g.*, about 2.35 nM, or about 0.1-0.5 nM or 0.1-0.2 nM.

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 93 or 99); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 94 or 273); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 95), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 96); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 97); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 225); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4035-062 (*e.g.*, SEQ ID NO: 229).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 299 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 300 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4035-062.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 0.01 nM or less, *e.g.*, about 0.001-0.002 nM. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 1 nM or less, *e.g.*, about 0.1-0.5 nM or 0.2-0.4 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.1-0.5 nM or 0.1-0.2 nM.

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising the amino acid sequence of I-Y-D-V-H (SEQ ID NO: 99); an HCDR2 comprising the amino acid sequence of V-I-W-S-D-G-S-T-D-Y-N-X12-X13-X14-X15-S (SEQ ID NO: 342), X12 is A or P, X13 is A or S, X14 is F or L, and X15 is I or K; or an HCDR3 comprising the amino acid sequence of N-W-V-D-Q-A-W-F-A-Y (SEQ ID NO: 95), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising the amino acid sequence of R-A-S-K-N-I-Y-S-Y-L-A (SEQ ID NO: 96); an LCDR2 comprising the amino acid sequence of N-A-K-T-L-P-E (SEQ ID NO: 97); or an LCDR3 comprising the amino acid sequence of Q-H-H-Y-G-T-P-L-T (SEQ ID NO: 98).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of SEQ ID NO: 101 or 225; or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of SEQ ID NO: 102 or 229.

In an embodiment, the antibody molecule is monoclonal antibody 4035. In an embodiment, the antibody molecule is monoclonal antibody 4035-062.

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 103 or 109); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 104 or 110); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 105), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 106); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 107); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 108).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 111); or (ii) a VL

comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3934 (*e.g.*, SEQ ID NO: 112).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 175 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 176 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3934. In an embodiment, the antibody molecule is humanized monoclonal antibody 3934.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 3.16 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6

nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC₅₀ of 5 nM or less, *e.g.*, about 2.35 nM.

In an embodiment, the antibody molecule comprises:

- (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 11 or 149); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 142 or 150); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 143), and
- (ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 144 or 146); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 107 or 147); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 145 or 148).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 151); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4338 (*e.g.*, SEQ ID NO: 152 or 153).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 183 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 184 or 185 (or a nucleotide sequence substantially identical thereto), or both.

5 In an embodiment, the antibody molecule is monoclonal antibody 4338. In an embodiment, the antibody molecule is humanized monoclonal antibody 4338.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less,
 10 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM
 15 and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL
 20 (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or
 25 less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 3.16
 30 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6
 35 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM

and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC₅₀ of 5 nM or less, *e.g.*, about 2.35 nM.

5 In another embodiment, the antibody molecule comprises (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100%
10 homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 163 or 169); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 164 or 170); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,
15 or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 165), and

(ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino
20 acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 166); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 167); or an LCDR3
25 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 168).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15
30 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 171); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4237 (*e.g.*, SEQ ID NO: 172).

35 In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 188 (or a nucleotide sequence substantially identical thereto) or a VL

encoded by the nucleotide sequence of SEQ ID NO: 189 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4237. In an embodiment, monoclonal antibody 4237 is humanized monoclonal antibody 4237. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 235-240, a VL comprising the amino acid sequence of any of SEQ ID NO: 241-245, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 3.16 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM

and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC₅₀ of 5 nM or less, *e.g.*, about 2.35 nM.

5 In another embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence of G-Y-X3-X4-T-X6-X7-Y (SEQ ID NO: 330), wherein X3 is S or T; X4 is I or
10 F; X6 is S or absent; and X7 is G, D or S; an HCDR2 comprising an amino acid sequence of X3-X4-X5-X6-X7-X8 (SEQ ID NO: 331), wherein X3 is absent, N or Y; X4 is S or P, X5 is Y, L or R; X6 is D, N or R; X7 is G or S; and X8 is Y, D or S; or an HCDR3 comprising an amino acid sequence of X1-X2-X3-X4-Y-X6-X7-X8-X9-F-X11-X12 (SEQ ID NO: 332), wherein X1 is Y, E or H; X2 is absent or G; X3 is Y, D or A; X4 is D, G or Y; X6 is E, absent or D; X7 is D, Y, S or K; X8 is W, N
15 or R; X9 is Y, A or G; X11 is G or D; and X12 is V or Y, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence of X1-A-S-X4-S-V-X7-X8-X9-G-X11-X12-X13-X14-X15 (SEQ ID NO: 333),
20 wherein X1 is R or K; X4 is E or Q; X7 is D or S; X8 is N, F, I or N; X9 is Y, A, I or D; X11 is I or T; X12 is S, N or R; X13 is F, L or S; X14 is M or I; and X15 is N or H; an LCDR2 comprising an amino acid sequence of X1-A-S-N-X5-X6-X7 (SEQ ID NO: 334), wherein X1 is A, R or H; X5 is Q or L; X6 is G or E; and X7 is S, P or T; or an LCDR3 comprising an amino acid sequence of X1-Q-S-X4-X5-X6-P-X8-T (SEQ ID NO: 335), wherein X1 is Q or L; X4 is K, R or N; X5 is E or K; X6 is V,
25 Y, I or D; and X8 is R, W or Y.

In an embodiment, the antibody molecule is any of monoclonal antibodies 2218, 2419, 2922, or 3327.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC₅₀ of
30 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between
35 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to

mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, or 0.1 nM or less, *e.g.*, between 0.1 and 50 nM, *e.g.*, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In another embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence of X6-X7-Y-X9-X10-X11 (SEQ ID NO: 336), wherein X6 is S or absent; X7 is G, D or S; X9 is Y, F, T or D; X10 is W, M, I or V; and X11 is N, H or F; an HCDR2 comprising an amino acid sequence of X1-I-X3-X4-X5-X6-X7-X8-X9-X10-Y-N-X13-X14-X15-K-X17 (SEQ ID NO: 337), wherein X1 is Y, R or W; X3 is absent, N or Y; X4 is S or P; X5 is Y, L or R; X6 is D, N or R; X7 is G or S; X8 is Y, D or S; X9 is N, T or I; X10 is N, F or K; X13 is P, Q or E; X14 is S or K; X15 is L or F; and X17 is N, G or D; or an HCDR3 comprising an amino acid sequence of X1-X2-X3-X4-Y-X6-X7-X8-X9-F-X11-X12 (SEQ ID NO: 332), wherein X1 is Y, E or H; X2 is absent or G; X3 is Y, D or A; X4 is D, G or Y; X6 is E, absent or D; X7 is D, Y, S or K; X8 is W, N or R; X9 is Y, A or G; X11 is G or D; and X12 is V or Y, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino

acid sequence of X1-A-S-X4-S-V-X7-X8-X9-G-X11-X12-X13-X14-X15 (SEQ ID NO: 333), wherein X1 is R or K; X4 is E or Q; X7 is D or S; X8 is N, F, I or N; X9 is Y, A, I or D; X11 is I or T; X12 is S, N or R; X13 is F, L or S; X14 is M or I; and X15 is N or H; an LCDR2 comprising an amino acid sequence of X1-A-S-N-X5-X6-X7 (SEQ ID NO: 334), wherein X1 is A, R or H; X5 is Q or L; X6 is G or E; and X7 is S, P or T; or an LCDR3 comprising an amino acid sequence of X1-Q-S-X4-X5-X6-P-X8-T (SEQ ID NO: 335), wherein X1 is Q or L; X4 is K, R or N; X5 is E or K; X6 is V, Y, I or D; and X8 is R, W or Y.

In an embodiment, the antibody molecule is any of monoclonal antibodies 2218, 2419, 2922, or 3327. In an embodiment, the antibody molecule is a humanized monoclonal antibody 2218, 2419, 2922, or 3327.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human TACI, at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of human APRIL to human BCMA, at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM

or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

5

In another embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 61 or 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 62 or 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 63), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 67); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 66); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3530 (*e.g.*, SEQ ID NO: 70).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 83 (or a nucleotide sequence substantially identical thereto) or a VL

encoded by the nucleotide sequence of SEQ ID NO: 84 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3530. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3530.

5 In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL, mouse APRIL, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 5 nM or less, *e.g.*, about 2.7 nM.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*,

as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 4.95 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both),
 5 at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10
 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.68 nM.

15 In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least
 20 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 61 or 64); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 62 or 65); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or
 25 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 63), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino
 30 acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 45); or an LCDR3
 35 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 66); or (ii) a VL
 5 comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3525 (*e.g.*, SEQ ID NO: 50).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 83 (or a nucleotide sequence substantially identical thereto) or a VL
 10 encoded by the nucleotide sequence of SEQ ID NO: 80 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3525. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3525.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL,
 15 mouse APRIL, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to
 20 human APRIL at an EC_{50} of 5 nM or less, *e.g.*, about 2.5 nM.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined
 30 by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an

embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 4.05 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.85 nM.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 113 or 119); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 114 or 120); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 115), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino

acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 117); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 118).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 121); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3833 (*e.g.*, SEQ ID NO: 122).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 177 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 178 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3833. In an embodiment, monoclonal antibody 3833 is a humanized monoclonal antibody 3833. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 246-250, a VL comprising the amino acid sequence of any of SEQ ID NO: 251-253, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL, mouse APRIL, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 5 nM or less, *e.g.*, about 2.5 nM.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or

less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 4.05 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.85 nM.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least

85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 123 or 129); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 124 or 130); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 125), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 126); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 128).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 131); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3631 (*e.g.*, SEQ ID NO: 132).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 179 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 180 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3631. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3631.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL, mouse APRIL, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC₅₀ of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005

nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC₅₀ of 5 nM or less, *e.g.*, about 2.5 nM.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC₅₀ of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both. In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC₅₀ of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC₅₀ of 5 nM or less, *e.g.*, about 4.05 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), at an IC₅₀ of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an

embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.85 nM.

In an embodiment, the antibody molecule comprises:

- 5 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal
 - 10 antibody 3732 (*e.g.*, SEQ ID NO: 133 or 138); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 134 or 139); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid
 - 15 sequence of the HCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 135), and
 - (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90,
 - 20 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 136); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 127); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,
 - 25 or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 137).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the

- 30 amino acid sequence of the VH of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 140); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3732 (*e.g.*, SEQ ID NO: 141).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide

- 35 sequence of SEQ ID NO: 181 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 182 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3732. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3732.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL, mouse APRIL, or both.

5 In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between
10 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 5 nM or less, *e.g.*, about 2.5 nM.

15 In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC_{50} of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less,
20 or 0.001 nM or less, *e.g.*, between 0.001 nM and 100 nM, *e.g.*, between 0.001 nM and 50 nM, between 0.01 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM or between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein.

25 In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM,
30 *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 4.05 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both), at an IC₅₀ of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC₅₀ of 1 nM or less, *e.g.*, about 0.85 nM.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 154 or 159); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 155 or 160); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 156), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 116); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 157); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15

amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 161); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540 (*e.g.*, SEQ ID NO: 162).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 186 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 187 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4540. In an embodiment, monoclonal antibody 4540 is a humanized monoclonal antibody 4540. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of any of SEQ ID NO: 254-258, a VL comprising the amino acid sequence of any of SEQ ID NO: 259-261, or both.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 154 or 276); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 155 or 277); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 156), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 258); or (ii) a
 5 VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540-063 (*e.g.*, SEQ ID NO: 261).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 301 (or a nucleotide sequence substantially identical thereto) or a VL
 10 encoded by the nucleotide sequence of SEQ ID NO: 302 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4540-063.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises
 15 three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 154 or 159); an HCDR2 comprising an amino acid sequence
 20 that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 155 or 278); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 156), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three
 25 light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody
 30 4540-033 (*e.g.*, SEQ ID NO: 274); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 275); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the
 35 LCDR3 of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15

amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 256); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4540-033 (*e.g.*, SEQ ID NO: 261).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 303 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 302 (or a nucleotide sequence substantially identical thereto), or both.

10 In an embodiment, the antibody molecule is monoclonal antibody 4540-033.

In an embodiment, the antibody molecule comprises:

- (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the VH comprises one, two, or all of the following: an HCDR1 comprising the amino acid sequence of D-Y-Y-X4-N (SEQ ID NO: 343), where X4 is I or M; an HCDR2 comprising the amino acid sequence of W-I-F-P-G-S-G-S-T-Y-Y-X12-X13-K-X15-X16-G, where X12 is N or A, X13 is E or Q, X15 is F or L, and X16 is K or Q (SEQ ID NO: 344); or an HCDR3 comprising the amino acid sequence of G-D-S-G-R-A-M-D-Y (SEQ ID NO: 156), and
- 20 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the VL comprises one, two, or all of the following: an LCDR1 comprising the amino acid sequence of X1-A-S-Q-D-I-N-K-Y-I-A, wherein X1 is K or Q (SEQ ID NO: 345); an LCDR2 comprising the amino acid sequence of Y-T-S-T-L-X6-X7, wherein X6 is Q or E, and X7 is S or T (SEQ ID NO: 346); or an
- 25 LCDR3 comprising the amino acid sequence of L-Q-Y-D-N-L-L-T (SEQ ID NO: 158).

In another embodiment, the antibody molecule comprises:

- (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the VH comprises one, two, or all of the following: an HCDR1 comprising the amino acid sequence of G-Y-T-F-A-D-Y (SEQ ID NO: 154); an HCDR2 comprising the amino acid sequence of F-P-G-S-G-S (SEQ ID NO: 155); or an HCDR3 comprising the amino acid sequence of G-D-S-G-R-A-M-D-Y (SEQ ID NO: 156), and
- 30 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the VL comprises one, two, or all of the following: an LCDR1 comprising the amino acid sequence of X1-A-S-Q-D-I-N-K-Y-I-A, wherein X1 is K or Q (SEQ ID NO: 345); an LCDR2 comprising the amino acid
- 35

sequence of Y-T-S-T-L-X6-X7 (SEQ ID NO: 346), wherein X6 is Q or E, and X7 is S or T; or an LCDR3 comprising the amino acid sequence of L-Q-Y-D-N-L-L-T (SEQ ID NO: 158).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of SEQ ID NOS: 161, 256 or 258; or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of SEQ ID NO: 162 or 261.

In an embodiment, the antibody molecule is monoclonal antibody 4540. In another embodiment, the antibody molecule is monoclonal antibody 4540-063. In yet another embodiment, the antibody molecule is monoclonal antibody 4540-033.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL, mouse APRIL, or both.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC₅₀ of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, or 0.1 nM or less, *e.g.*, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC₅₀ of 5 nM or less, *e.g.*, about 2.5 nM.

In an embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL at an EC₅₀ of 100 nM or less, *e.g.*, 80 nM or less, 60 nM or less, 40 nM or less, 20 nM or less, 10 nM or less, 9 nM or less or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, or 0.1 nM or less, *e.g.*, between 0.1 nM and 20 nM, *e.g.*, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), BCMA (*e.g.*, human BCMA), or both.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both), at an IC₅₀ of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM,

e.g., as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 5 nM or less, *e.g.*, about 4.05 nM.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both),
 5 at an IC_{50} of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10
 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA at an IC_{50} of 1 nM or less, *e.g.*, about 0.85 nM.

15 In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least
 20 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 21 or 27); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 22 or 28); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or
 25 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 23), and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino
 30 acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 24); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 25); or an LCDR3
 35 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 26).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 29); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 2621 (*e.g.*, SEQ ID NO: 30).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 75 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 76 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 2621. In an embodiment, the antibody molecule is a humanized monoclonal antibody 2621.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 1 nM or less, *e.g.*, about 0.7 nM. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more, *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI). In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of about 1 nM or less.

In an embodiment, the antibody molecule does not inhibit, or does not substantially inhibit, binding of APRIL (*e.g.*, human APRIL) to BCMA (*e.g.*, human BCMA).

In an embodiment, the antibody molecule comprises:

- 5 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal
 - 10 antibody 3125 (*e.g.*, SEQ ID NO: 11 or 47); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 42 or 48); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid
 - 15 sequence of the HCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 43), and
- (ii) a light chain variable region (VH), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90,
 - 20 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 44); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 45); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from,
 - 25 or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 46).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 49); or (ii) a VL comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 3125 (*e.g.*, SEQ ID NO: 50).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 79 (or a nucleotide sequence substantially identical thereto) or a VL encoded by the nucleotide sequence of SEQ ID NO: 80 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 3125. In an embodiment, the antibody molecule is a humanized monoclonal antibody 3125.

In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL. In an embodiment, the antibody molecule binds, or substantially binds, to human APRIL at an EC_{50} of
 5 20 nM or less, *e.g.*, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, 0.01 nM or less, 0.005 nM or less, 0.002 nM or less, or 0.001 nM or less, *e.g.*, between 0.001 nM and 20 nM, *e.g.*, between 0.01 nM and 20 nM, between 0.1 nM and 20 nM, between 0.1 nM and 10 nM, between 0.5 nM and 5 nM, between
 10 1 nM and 5 nM, between 0.001 nM and 0.1 nM, between 0.001 nM and 0.01 nM, between 0.001 nM and 0.005 nM, between 0.01 nM and 0.05 nM, or between 0.01 nM and 0.1 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule binds to human APRIL at an EC_{50} of 20 nM or less, *e.g.*, about 13 nM. In an embodiment, the antibody molecule does not bind, or binds to mouse APRIL with low affinity, *e.g.*, at an EC_{50} of 1000 nM or more, *e.g.*, 2000 nM or more,
 15 *e.g.*, as determined by a method described herein.

In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI). In an embodiment, the antibody molecule inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL) to TACI (*e.g.*, human TACI), at an IC_{50} of 50 nM or less, *e.g.*, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9
 20 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, 1 nM or less, 0.8 nM or less, 0.6 nM or less, 0.4 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, 0.02 nM or less, or 0.01 nM or less, *e.g.*, between 0.01 nM and 50 nM, between 0.1 nM and 50 nM, between 0.1 nM and 25 nM, between 0.1 nM and 10 nM, between 0.1 nM and 5 nM, between 0.1 nM and 1 nM, between 0.1 nM and 0.5 nM, between 0.5 nM and 5 nM, or between 1 nM
 25 and 5 nM, *e.g.*, as determined by a method described herein. In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI at an IC_{50} of 150 nM or less, *e.g.*, about 112.97 nM. In an embodiment, the antibody molecule does not inhibit, or does not substantially inhibit, binding of APRIL (*e.g.*, human APRIL) to BCMA (*e.g.*, human BCMA).

30 In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least
 35 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR1 of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 266 or 269); an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100%

homology with, the amino acid sequence of the HCDR2 of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 267 or 270); or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the HCDR3 of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 268), and

5 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of monoclonal antibody 4439
10 (*e.g.*, SEQ ID NO: 146); an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR2 of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 147); or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR3 of
15 monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 148).

In an embodiment, the antibody molecule comprises one or both of: (i) a VH comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VH of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 271); or (ii) a VL
20 comprising an amino acid sequence that differs by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid residues from, or has at least 85, 90, 95, 96, 97, 98, 99, or 100% homology with, the amino acid sequence of the VL of monoclonal antibody 4439 (*e.g.*, SEQ ID NO: 272).

In an embodiment, the antibody molecule comprises a VH encoded by the nucleotide sequence of SEQ ID NO: 297 (or a nucleotide sequence substantially identical thereto) or a VL
25 encoded by the nucleotide sequence of SEQ ID NO: 298 (or a nucleotide sequence substantially identical thereto), or both.

In an embodiment, the antibody molecule is monoclonal antibody 4439. In an embodiment, monoclonal antibody 4439 is humanized monoclonal antibody 4439.

30 In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in any of **Tables 3-4 or 7-8**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within
35 a region of human APRIL as defined in **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from

Table 3. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 3**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, residues within a region of human APRIL as defined in **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all, of the human APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises one or more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, residues within a region of human APRIL as defined in **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all, of the human APRIL residues from **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 7**.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of human APRIL as defined in **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all, of the human APRIL residues from **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 8**.

In an embodiment, the antibody molecule is an IgG antibody molecule, *e.g.*, comprising a heavy chain constant region of IgG, *e.g.*, chosen from IgG1, IgG2 (*e.g.*, IgG2a), IgG3, or IgG4, *e.g.*, IgG2 or IgG4. In an embodiment, the antibody molecule is an IgG1 antibody molecule. In another embodiment, the antibody molecule is an IgG2 antibody molecule. In an embodiment, the antibody molecule comprises a light chain constant region of kappa or lambda light chain.

In an embodiment, the antibody molecule comprises an Fc region. In an embodiment, the Fc region comprises one or more mutations located at the interface between the CH2 and CH3 domains (*e.g.*, to increase the binding affinity to neonatal receptor FcRn and/or the half-life of the antibody molecule). In an embodiment, the Fc region comprises one or more mutations, *e.g.*, one or more (*e.g.*, 2, 3, 4, 6 or all) mutations chosen from T250Q, M252Y, S254T, T256E, M428L, H433K, N434F, or any combination thereof, of IgG1. In an embodiment, the Fc region comprises one or more mutations at positions 233-236 or 322 of human IgG1 or IgG2, or one or more substitutions at positions 327, 330 or 331 of human IgG4 (*e.g.*, to reduce complement-dependent cytotoxicity (CDC)). In an embodiment, the Fc region comprises one or more (*e.g.*, 2, 3, 4, 6 7 or all) mutations chosen from E233P, L234V, L235A, G236, K322A, A327G, A330S, P331S, or any combination thereof.

In an embodiment, the antibody molecule is a humanized antibody molecule, *e.g.*, as described in **Table 5**, *e.g.*, comprising one or more framework regions derived from human framework germline sequence.

In an embodiment, the antibody molecule comprises two heavy chain variable regions and two light chain variable regions. In an embodiment, the antibody molecule is a Fab, F(ab')₂, Fv, Fd, or a single chain Fv fragment (scFv).

The disclosure features a composition, *e.g.*, pharmaceutical composition, comprising an antibody molecule described herein. In an embodiment, the composition further comprises a pharmaceutical acceptable carrier. In another aspect, the present disclosure features a composition comprising an antibody molecule as disclosed herein, optionally, wherein the composition is a pharmaceutical composition, optionally further comprising a pharmaceutically acceptable carrier.

In an aspect, the disclosure features a nucleic acid molecule encoding a heavy chain variable region (VH), a light chain variable region (VL), or both, of an antibody molecule described herein.

In an aspect, the disclosure features a vector comprising a nucleic acid molecule described herein.

In an aspect, the disclosure features a cell, *e.g.*, an isolated cell, comprising a nucleic acid molecule described herein or a vector described herein.

In an aspect, the disclosure features a kit comprising an antibody molecule described herein and instructions to use of the antibody molecule.

In an aspect, the disclosure features a container comprising an antibody molecule described herein or a composition described herein.

In an aspect, the disclosure features a method of producing an anti-APRIL antibody molecule, the method comprising culturing a cell described herein under conditions that allow production of an antibody molecule, thereby producing the antibody molecule.

In an embodiment, the method further comprises isolating the antibody molecule.

The disclosure also features a method of treating IgA nephropathy, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating IgA nephropathy. In another aspect, the present disclosure features the use of an antibody molecule described herein, or a composition
5 described herein, in the manufacture of a medicament for treating IgA nephropathy in a subject.

In an embodiment, the antibody molecule is administered to the subject intravenously.

In an embodiment, the antibody molecule is administered to the subject at a dose between 0.1 mg/kg and 50 mg/kg, *e.g.*, between 0.2 mg/kg and 25 mg/kg, between 0.5 mg/kg and 10 mg/kg, between 0.5 mg/kg and 5 mg/kg, between 0.5 mg/kg and 3 mg/kg, between 0.5 mg/kg and 2.5 mg/kg,
10 between 0.5 mg/kg and 2 mg/kg, between 0.5 mg/kg and 1.5 mg/kg, between 0.5 mg/kg and 1 mg/kg, between 1 mg/kg and 1.5 mg/kg, between 1 mg/kg and 2 mg/kg, between 1 mg/kg and 2.5 mg/kg, between 1 mg/kg and 3 mg/kg, between 1 mg/kg and 2.5 mg/kg, or between 1 mg/kg and 5 mg/kg.

In an embodiment, the antibody molecule is administered to the subject at a fixed dose between 10 mg and 1000 mg, *e.g.*, between 10 mg and 500 mg, between 10 mg and 250 mg, between
15 10 mg and 150 mg, between 10 mg and 100 mg, between 10 mg and 50 mg, between 250 mg and 500 mg, between 150 mg and 500 mg, between 100 mg and 500 mg, between 50 mg and 500 mg, between 25 mg and 250 mg, between 50 mg and 150 mg, between 50 mg and 100 mg, between 100 mg and 150 mg, between 100 mg and 200 mg, or between 150 mg and 250 mg.

In an embodiment, the antibody molecule is administered once a week, twice a week, once
20 every two weeks, once every three weeks, once every four weeks, once every eight weeks, once a month, once every two months, or once every three months.

In an embodiment, administration of the antibody molecule reduces the level of IgA in a peripheral tissue, *e.g.*, in serum, mucosal tissue, bone marrow, or any combination thereof.

In an embodiment, administration of the antibody molecule reduces the level of IgA1. In an
25 embodiment, administration of the antibody molecule reduces the level of IgA1 in polymeric form (pIgA1). In an embodiment, administration of the antibody molecule reduces the level of IgA1 with O-linked glycosylation variants (*e.g.*, aberrant or reduced composition of galactose in CH1 hinge region).

In an embodiment, the method further comprises determining the level of IgA in a peripheral
30 tissue sample from the subject, *e.g.*, chosen from serum, mucosal tissue, or bone marrow.

In an embodiment, the method further comprises administering to the subject a second therapy for IgA nephropathy. In an embodiment, the second therapy is chosen from an angiotensin-converting enzyme (ACE) inhibitor, an angiotensin receptor blocker (ARB), omega-3 fatty acids, an immunosuppressant (*e.g.*, a corticosteroid, *e.g.*, prednisone), a statin, mycophenolate mofetil, or any
35 combination thereof.

The disclosure also features a method of treating diabetic nephropathy, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating diabetic nephropathy. In another aspect, the present disclosure features the use of an antibody molecule described herein, or a
5 composition described herein, in the manufacture of a medicament for treating diabetic nephropathy in a subject.

The disclosure further features a method of treating cancer, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating cancer. In another aspect, the present disclosure features the use of an antibody molecule described herein, or a composition described
5 herein, in the manufacture of a medicament for treating cancer in a subject.

In an embodiment, the cancer is a hematological cancer. In an embodiment, the hematological cancer is chosen from B-cell non-Hodgkin's lymphoma, chronic lymphocytic leukemia (CLL), Hodgkin's lymphoma, multiple myeloma, Waldenström macroglobulinemia, or lymphoplasmacytic lymphoma. In an embodiment, the cancer is a multiple myeloma.

10 The disclosure also features a method of treating an immunoproliferative disorder, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating the immunoproliferative disorder. In another aspect, the present disclosure features the use of an antibody molecule as described herein, or a composition of described herein, in the manufacture of a medicament for treating an
15 immunoproliferative disorder in a subject.

In an embodiment, the immunoproliferative disorder is monoclonal IgA hypergammaglobulinemia.

The disclosure also features a method of treating vasculitis, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described
20 herein or a composition described herein, thereby treating vasculitis. In another aspect, the present disclosure features the use of an antibody molecule as described herein, or a composition of as described herein, in the manufacture of a medicament for treating vasculitis in a subject.

In an embodiment, the vasculitis is kidney vasculitis. In an embodiment, the vasculitis is an IgA associated vasculitis (*e.g.*, Henoch-Schonlein purpura) or post-streptococcal glomerulonephritis.

25 The disclosure further features a method of treating an autoimmune disorder, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating the autoimmune disorder. In yet another aspect, the present disclosure features the use of an antibody molecule described herein, or a composition described herein, in the manufacture of a medicament for treating an autoimmune
30 disorder in a subject.

In an embodiment, the autoimmune disorder is chosen from rheumatoid arthritis, systemic lupus erythematosus, a linear IgA bullous disease (*e.g.*, linear immunoglobulin A (IgA) dermatosis), or IgA-mediated epidermolysis bullosa acquisita (EBA).

The disclosure features a method of treating IgA pemphigus, the method comprising
35 administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating IgA pemphigus. In another aspect, the present disclosure features the use of an antibody molecule described herein, or a

composition described herein, in the manufacture of a medicament for treating IgA pemphigus in a subject.

5 The disclosure also features a method of treating celiac disease, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating celiac disease. In a further aspect, the present disclosure features the use of an antibody molecule described herein, or a composition described herein, in the manufacture of a medicament for treating celiac disease in a subject.

10 The disclosure further features a method of treating alcoholic cirrhosis, the method comprising administering to a subject in need thereof an effective amount of an antibody molecule described herein or a composition described herein, thereby treating alcoholic cirrhosis. In another aspect, the present disclosure features the use of an antibody molecule described herein, or a composition described herein, in the manufacture of a medicament for treating alcoholic cirrhosis in a subject.

15 In an aspect, the disclosure features a method of reducing the level of IgA in a cell or subject, the method comprising contacting the cell or subject, or administering to a subject in need thereof an

effective amount of, an antibody molecule described herein or a composition described herein, thereby reducing the level of IgA.

In an embodiment, the antibody molecule is administered to the subject intravenously.

5 In an embodiment, the antibody molecule is administered to the subject at a dose between 0.1 mg/kg and 50 mg/kg, *e.g.*, between 0.2 mg/kg and 25 mg/kg, between 0.5 mg/kg and 10 mg/kg, between 0.5 mg/kg and 5 mg/kg, between 0.5 mg/kg and 3 mg/kg, between 0.5 mg/kg and 2.5 mg/kg, between 0.5 mg/kg and 2 mg/kg, between 0.5 mg/kg and 1.5 mg/kg, between 0.5 mg/kg and 1 mg/kg, between 1 mg/kg and 1.5 mg/kg, between 1 mg/kg and 2 mg/kg, between 1 mg/kg and 2.5 mg/kg, between 1 mg/kg and 3 mg/kg, between 1 mg/kg and 2.5 mg/kg, or between 1 mg/kg and 5 mg/kg.

10 In an embodiment, the antibody molecule is administered to the subject at a fixed dose between 10 mg and 1000 mg, *e.g.*, between 10 mg and 500 mg, between 10 mg and 250 mg, between 10 mg and 150 mg, between 10 mg and 100 mg, between 10 mg and 50 mg, between 250 mg and 500 mg, between 150 mg and 500 mg, between 100 mg and 500 mg, between 50 mg and 500 mg, between 25 mg and 250 mg, between 50 mg and 150 mg, between 50 mg and 100 mg, between 100 mg and 150 mg, between 100 mg and 200 mg, or between 150 mg and 250 mg.

In an embodiment, the antibody molecule is administered once a week, twice a week, once every two weeks, once every three weeks, once every four weeks, once every eight weeks, once a month, once every two months, once every three months.

20 In an embodiment, administration of the antibody molecule reduces the level of IgA in a peripheral tissue, *e.g.*, in serum, mucosal tissue, bone marrow, or any combination thereof.

In an embodiment, administration of the antibody molecule reduces the level of IgA1. In an embodiment, administration of the antibody molecule reduces the level of IgA1 in polymeric form (pIgA1). In an embodiment, administration of the antibody molecule reduces the level of IgA1 with O-linked glycosylation variants (*e.g.*, aberrant or reduced composition of galactose in CH1 hinge region).

In an aspect, the disclosure features use of an antibody molecule described herein or a composition described herein in the treatment, or in the manufacture of a medicament for the treatment, of a disorder described herein.

30 The disclosure also features an antibody molecule described herein or a composition described herein for use in the treatment of a disorder described herein.

In an aspect, the disclosure features a method of detecting an APRIL molecule, the method comprising contacting a cell or a sample from a subject with an antibody molecule described herein, thereby detecting the APRIL molecule.

In an embodiment, the antibody molecule is coupled with a detectable label. In an embodiment, the APRIL molecule is detected *in vitro* or *ex vivo*. In another embodiment, the APRIL molecule is detected *in vivo*.

The disclosure contemplates all combinations of any one or more of the foregoing aspects and/or embodiments, as well as combinations with any one or more of the embodiments set forth in the detailed description and examples.

Other features, objects, and advantages of the compositions and methods herein will be apparent from the description and drawings, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B depict the anti-APRIL antibody profiling from mouse-derived hybridomas. CD-1 mice were immunized with recombinant multimeric APRIL (human or mouse) as described. The splenocytes from anti-APRIL seropositive mice were immortalized through myeloma fusion to generate hybridomas. ELISA-based screening methods were used for evaluating anti-APRIL antibodies present in conditioned media of immunoglobulin producing hybridomas. Cell culture supernatants were initially screened both for binding to target (APRIL) by indirect ELISA and for functional activity with respect to ability to block interaction of APRIL with the recombinant, soluble human TNF receptor TACI-Fc (blocking ELISA). Screening against both human and mouse APRIL was carried out to identify potentially cross-reactive antibodies. Relative activities are based on single point measurements following normalization of immunoglobulin concentration to 10 µg/mL when possible. APRIL immunogen for final tail vein boost is noted in bold. Hybridoma-derived antibodies lacking blocking activity to either human or mouse APRIL (as initially defined by less than 10% receptor blocking in this first-pass screening assay) are not included in this summary table. Ig isotypes for select hybridomas are noted.

FIGS. 2A-2B depict the APRIL receptor blocking activity of exemplary anti-APRIL antibodies. Functional activities of select anti-APRIL antibodies purified from mouse-derived hybridoma clones were assessed by ELISA using recombinant TACI-Fc as soluble receptor. Assay involves two steps: 1) preincubation of APRIL with varying concentrations of purified mouse antibody; 2) subsequent measurement of APRIL binding to immobilized TACI-Fc as quantified by ELISA using the appropriate secondary antibody for detection of epitope tagged APRIL. Interference of APRIL-receptor binding was measured as a loss of A₄₅₀. Hybridoma antibodies are depicted by solid symbols. Control and comparator antibodies are depicted by open symbols. Apry-1-1 represents a mouse-specific blocking antibody used as a control. Human neutralizing antibodies 0201 and 1313 were likewise used as controls and for comparative purposes to antibodies described in the scientific literature. Data presented is representative of assay and method for assessing antibody inhibition

(receptor blocking). **FIG. 2A** (human APRIL); **FIG. 2B** (mouse APRIL). IC_{50} values are reported in **FIG. 3**.

FIG. 3 depicts the activity profiles of select, purified mouse derived anti-APRIL antibodies. Binding and blocking activities are extrapolated from data summarized in **FIGS. 2A-2B** and are based on a non-linear regression of antibody titrations with 3 parameter curve fitting using Graphpad Prism. Dashes indicate no activity.

FIG. 4 depicts the inhibition of APRIL-mediated TACI signaling in HEK 293 cells. Functional inhibition of human APRIL-mediated receptor activation was assessed using a 293-derived cell line with stably transfected NF- κ B reporter and transiently transfected, human APRIL receptor TACI.

HA-tagged GCN4 human APRIL (R&D Systems) was used as source of APRIL. Pre-existing mAb 1313 (human specific, with pre-established blocking activity) was used as a positive control; mouse specific anti-APRIL antibody Apry-1-1 was used as a negative control. Data is illustrative of an orthogonal, biologically relevant assay to assess anti-APRIL activity of mouse derived antibodies.

FIG. 5 depicts the binding of anti-APRIL antibodies to human APRIL. Relative binding of select anti-APRIL antibodies was measured by indirect ELISA. Select antibodies based on prior hybridoma screening were recombinantly produced (based on immunoglobulin VH and VL gene sequencing) as described. HA-tagged GCN4 human APRIL R&D Systems was used as source of APRIL. Binding data was analyzed by non-linear regression using a three parameter fit. MAbs 1313, 0201, and Aprily-5 were included for comparative purposes; Apry-1-1 was used as a negative control.

Extrapolated EC_{50} values are summarized in **FIG. 6**.

FIG. 6 depicts the relative binding affinities of select anti-APRIL antibodies. Data was derived from indirect ELISA (summarized in **FIG. 5**). Dashes indicate no binding activity.

FIG. 7 depicts the antibody inhibition of APRIL binding to TACI. Assay is based on blocking ELISA using recombinant human APRIL (R&D Systems) and Human TACI-Fc. Inhibition was analyzed by non-linear regression using a four parameter fit. MAbs 1313 and 0201 were used as controls and for comparative purposes. Non-neutralizing anti-APRIL mAb Aprily-5 was used as a negative control.

FIG. 8 depicts the antibody inhibition of APRIL binding to BCMA. Assay is based on blocking ELISA using recombinant human APRIL and Human BCMA-Fc. *See* the description of **FIG. 7** for details.

FIG. 9 depicts antibody inhibition of APRIL binding to both human TACI-Fc and human BCMA-Fc. Relative inhibitory activities are summarized. Data derived from non-linear regression analyses of antibody inhibition curves depicted in **FIGS. 7-8** using a 4-parameter fit. % inhibition was normalized to the no antibody control (100% inhibition) following subtraction of background (0% inhibition). Dashes represent lack of calculated IC_{50} values due to poor or zero blocking activity measured.

FIGS. 10A-10B depict APRIL species cross reactivity of anti-APRIL antibodies. Anti-APRIL antibodies with blocking activity (summarized in **FIG. 9**) were assessed for ability to block both human and mouse APRIL binding to human TACI-Fc and BCMA-Fc. Recombinant mouse and human APRIL with otherwise identical HA-tagged GCN4 N-terminal fusions (R&D Systems) were used as receptor ligands. Select data is included for illustrative purposes. Antibodies 3530 and 3525 demonstrated any cross-species activity with respect to TACI-Fc receptor blocking (**FIG. 10A**). Analogous cross-neutralization with respect to APRIL binding to BCMA-Fc was only partially achieved (**FIG. 10B**). Neutralizing Antibody 1313 (human specific) and antibody Apyr-1-1 (mouse specific) were used as controls. Closed symbols, human APRIL; open symbols, mouse APRIL.

FIGS. 11A-11B depict antibody Inhibition of APRIL-mediated receptor signaling. Inhibition of APRIL-receptor mediated NF- κ B intracellular signaling was evaluated using the HEK 293 NF- κ B reporter cell line following transient transfection of either full-length human TNF family receptors TACI or BCMA full-length cDNA expression vectors. Data are normalized to activity vs. no antibody treatment. Scale of Y axis displayed ranges from 0 to 1. Inhibition of APRIL-mediated receptor signaling was measured at three different antibody concentrations (0.08 μ g/mL, 0.4 μ g/mL, and 10 μ g/mL). **FIG. 11A** shows the inhibition of TACI-mediated NFB signaling; **FIG. 11B** shows the inhibition of BCMA mediated NF- κ B signaling.

FIGS. 12A-12B depict *in vivo* potency of an anti-APRIL antibody in reducing serum IgA levels. The suitability of use of anti-APRIL antibody for modulating IgA production *in vivo* was evaluated based directly on a reduction of serum IgA levels following the administration of a neutralizing anti-APRIL antibody in a laboratory rodent model. For this purpose mouse-APRIL specific, blocking antibody Apyr-1-1 (Adipogen) hitherto used as a control for assessing anti-APRIL antibody activity *in vitro* was also used as a test antibody to demonstrate proof-of-concept. Age-matched male B6C3F1 mice (6-10 weeks old) were dosed with 20 mg/kg antibody two times a week via i.p. injection for a total of 8 weeks. Saline for injection was used as the negative (vehicle) control (VC). Serum isotype specific immunoglobulin levels (IgG, IgM, and IgA) were monitored individually by ELISA approximately every 12 days. **FIG. 12A** shows the effect of treatment on total serum IgA levels; **FIG. 12B** shows the effect of treatment vs. control on total immunoglobulin levels in sera. Data represent summation of IgG + IgA + IgM levels measured separately.

FIG. 13 depicts sequence alignment of human and mouse APRIL (SEQ ID NOS: 85 and 91, respectively). Sequence alignments of soluble APRIL were determined using CLUSTALW. Amino acid sequences correspond to SwissProt accession numbers O75888 (human) and Q9D777 (mouse). **FIG. 14** depicts the structural definition of exemplary APRIL epitope for antibody targeting. A space filling model of trimeric APRIL is depicted. Exemplary (spatially defined) region to be targeted by an antibody molecule is depicted in darker gray and indicated by arrow. This epitope includes positions in APRIL that bridge monomers. Differences between mouse and human APRIL sequences are

highlighted with corresponding amino acid differences at these positions noted. The putative N-glycosylation site (N124) is noted in black and indicated by arrow.

FIG. 15 depicts the structural definition of exemplary APRIL epitope for antibody targeting. Ribbon model of trimeric APRIL is depicted and core epitope for anti-APRIL targeting is highlighted in dark gray.

FIG. 16 depicts the binding of exemplary anti-APRIL antibodies to human APRIL. Relative binding of exemplary anti-APRIL antibodies was measured by indirect ELISA. Extrapolated EC₅₀ values are summarized in **FIG 17**.

FIG. 17 depicts the relative binding affinities of exemplary anti-APRIL antibodies. Data was derived from indirect ELISA (summarized in **FIG.16**).

FIGS. 18A-18B depict the antibody inhibition of APRIL-mediated receptor signaling. Inhibition of APRIL-receptor mediated NFκB intracellular signaling was evaluated using the HEK 293 NFκB reporter cell line following transient transfection of either full-length human TNF family receptors TACI (**FIG. 18A**) or BCMA (**FIG. 18B**) full-length cDNA expression vectors. Exemplary antibodies were shown for illustrative purposes.

FIGS. 19A-19B depict the antibody inhibition of APRIL binding to TNFSF receptors TACI (**FIG. 19A**) and BCMA (**FIG. 19B**). Assay is based on blocking ELISA using recombinant human APRIL (R&D Systems) and Human TACI-Fc. Inhibition was analyzed by non-linear regression using a four parameter fit. IC₅₀ values are summarized in **FIG. 20**.

FIG. 20 depicts the antibody inhibition of APRIL binding to both human TACI-Fc and human BCMA-Fc (summary of relative inhibitory activities). Data derived from non-linear regression analyses of antibody inhibition curves depicted in **FIG. 19** using a 4-parameter fit. % inhibition was normalized to the no antibody control (100% inhibition) following subtraction of background (0% inhibition). Dashes represent lack of calculated IC₅₀ values due to poor or partial blocking activity measured.

FIGS. 21A-21B depict APRIL species cross blocking activities of anti-APRIL antibodies. Ability of anti-APRIL antibodies to block binding of both human (**FIG. 21A**) and mouse (**FIG. 21B**) APRIL to BCMA was evaluated by ELISA. Respective blocking activities of exemplary antibodies (3833, 4540, 3530, 3631, and 3732) with previously determined cross-species (mouse and human) APRIL binding are shown.

FIG. 22 depicts human APRIL site-directed variants used for epitope mapping. APRIL is depicted as a trimer. Typical epitope containing CRD2 high affinity receptor binding site is depicted in dark gray. Positions of amino acid changes are noted with wildtype amino acid preceding number and mutation following (*e.g.*, R233G represents mutation of arginine at position 233 to glycine).

FIGS. 23A-23B depict epitope mapping of antibody 4035 (**FIG. 23A**) (comparison is made to reference antibody 1313; *see* **FIG. 23B**). Antibody binding to human and mouse variants was assessed by ELISA. FLAG-tagged APRIL was captured from cell culture media using anti-FLAG

antibody. Human APRIL variants are depicted as open bars; mouse APRIL variants are depicted as solid bars.

FIGS. 24A-24B depict epitope mapping of antibody 2419 (**FIG. 24A**) (comparison is made to reference antibody 1313; *see* **FIG. 24B**). Antibody binding to human and mouse variants was assessed by ELISA. Human APRIL variants are depicted as open bars; mouse APRIL variants are depicted as solid bars.

FIGS. 25A-25B depict epitope mapping of antibody 3833 (**FIG. 25A**) (comparison is made to reference antibody 1313; *see* **FIG. 25B**). Antibody binding to human and mouse variants was assessed by ELISA. FLAG-tagged APRIL was captured from cell culture media using anti-FLAG antibody. Human APRIL variants are depicted as open bars; mouse APRIL variants are depicted as solid bars.

FIG. 26 depicts differentiated epitope mapping of anti-APRIL antibodies by site directed mutagenesis. Primary characterization of human APRIL binding site was carried out by site-directed mutagenesis of select amino acid positions within APRIL followed by evaluation of antibody binding to these variants by ELISA. Exemplary data for three anti-APRIL antibodies 4035 (solid circles), 2419 (solid squares), and 1313 (open triangles) are shown for illustrative purposes.

FIGS. 27A-27B depict the binding of exemplary anti-APRIL antibodies to human APRIL. Antibodies include humanized variants of mouse antibodies 4035 (**FIG. 27A**) and 2419 (**FIG. 27B**). Relative binding of exemplary anti-APRIL antibodies was measured by indirect ELISA. Comparison of humanized anti-APRIL antibodies to parental (non-humanized) mouse antibodies is made for comparative purposes. Extrapolated EC_{50} values are summarized in **FIG 29A**.

FIGS. 28A-28B depict the binding of humanized variant of human-mouse cross reactive antibody 4540-063 to both human APRIL (**FIG. 28A**) and mouse APRIL (**FIG. 28B**). Relative binding of exemplary anti-APRIL antibody was measured by indirect ELISA. Comparison of parental (non-humanized) antibody is made for comparative purposes. Extrapolated EC_{50} values are summarized in **FIG 29B**.

FIGS. 29A-29B depict the relative binding affinities of exemplary anti-APRIL antibodies. Data (EC_{50} values) were derived from indirect ELISA (summarized in **FIGS. 27A-27B** and **FIGS 28A-28B**). **FIG. 29A** shows relative binding affinities of exemplary, humanized anti-APRIL antibodies based on 2419 and 4035. **FIG. 29B** shows relative binding affinities of cross-reactive antibody 4540 and its humanized variant 4540-063. Binding data for both human and mouse APRIL are included.

FIGS. 29C-29D depict the binding affinity of antibody 2419-1406 and 4035-062 to trimeric human APRIL measured by ELISA. Trimer was stabilized by N-terminal fusion of APRIL with isoleucine zipper (GCN4) domain. Binding of APRIL to human TACI-Fc is shown for comparative purposes.

The EC_{50} values derived from the binding curves depicted in **FIG. 29C** are summarized in **FIG. 29D**.

FIGS. 30A-30B depict antibody inhibition of APRIL binding to TNFSF receptors TACI and BCMA by humanized IgG2 κ variants of parental, murine derived antibody 2419. Assay is based on receptor

blocking ELISA using recombinant human APRIL (R&D Systems) and either human TACI-Fc (**FIG. 30A**) or BCMA-Fc (**FIG. 30B**). Parental, chimeric 2419 (mouse VH-VL grafted on to human IgG1 κ constant regions) was included for comparative purposes as was chimeric anti-human APRIL antibody 4035. Inhibition was analyzed by non-linear regression using a four parameter curve fit following normalization to 100% activity (no antibody control). IC₅₀ values are summarized in **FIG. 33**.

FIGS. 31A-31B depict the inhibition of APRIL binding to TNFSF receptors TACI (**FIG. 31A**) and BCMA (**FIG. 31B**) by additional humanized variants of 2419 (IgG2 κ) 2419-0205 and 2419-1406 and humanized variant (4035-062) of parental antibody 4035. Humanized 4035-062 is of the IgG1 κ subtype. Chimeric, non-humanized versions of mouse derived antibodies 4035 and 2419, and 1313 are included for comparative purposes. mAb1313 is a control anti-APRIL antibody. TACI-Fc and BCMA-Fc were used. IC₅₀ values are summarized in **FIG. 33**.

FIGS. 32A-32B depict the antibody inhibition of APRIL binding to TNFSF receptors TACI (**FIG. 32A**) and BCMA (**FIG. 32B**) by humanized variants of mouse/human APRIL cross neutralizing antibody 4540. Humanized antibody 4540 is of the IgG1 κ subtype. Parental 4540 (non-humanized chimera) and humanized 4035-062 (**FIGS. 30A-30B**) are included for comparative purposes. Inhibition was analyzed by non-linear regression using a four parameter curve fit as described for **FIGS. 29A-29B** and **FIGS. 30A-30B**. IC₅₀ values are also summarized in **FIG. 33**.

FIG. 33 depicts IC₅₀ values of antibody inhibition of APRIL-receptor binding. IC₅₀ values are based on a non-linear regression analysis of data from **FIGS. 30A-32B**. Data are also normalized (relative activity) to parental, mouse derived antibody 2419.

FIGS. 34A-34B depict the antibody inhibition of APRIL-mediated receptor signaling. Inhibition of APRIL-receptor mediated NF κ B intracellular signaling was evaluated using the HEK 293 NF κ B reporter cell line following transient transfection of either full-length human TNF family receptors TACI or BCMA cDNA expression vectors. Data are normalized to no antibody control (100%).

FIG. 34A shows inhibition of TACI-mediated NF κ B signaling. **FIG. 34B** shows inhibition of BCMA-mediated-NF κ B signaling.

FIG. 35 depicts the approximate IC₅₀ values of antibody inhibition of APRIL-mediated receptor signaling. Data are extrapolated from **FIGS. 34A-34B** based on a non-linear regression analysis using a variable slope, three parameter fit of antibody concentration vs. response. Negative antibody control (no APRIL binding) demonstrated no activity in this assay (data not shown).

FIG 36 depicts analysis of APRIL antibody binding reactivity to other members of the TNFSF13 family of cytokines. A select panel of recombinant TNFSF-related cytokines (Adipogen) was used to test for antibody cross-reactivity as measured by ELISA. Panel included human APRIL (specific target) and structurally related BAFF. Exemplary antibodies are included for illustrative purposes. As predicted, strong binding of anti-APRIL antibodies to human APRIL was detected. Cross-reactivity to BAFF and other members other than the target (human APRIL), however, was generally

not substantially detected above assay background (measured here using BSA only and assay diluent, 1x PBS controls). Antibody 4338, a previously identified BAFF cross-reactive antibody, was used as a positive (BAFF-reactive) control. A control antibody, mAb1313, was also included.

FIGS. 37A-37C depict the minimal protein cross-reactivity of mAb 2419-1406 and mAb 4035-062 in an extensive array of over 4500 heterologously expressed human membrane proteins in an HEK293-based assay and the confirmatory/secondary screen. An example of protein expression array design is shown in **FIG. 37A**. **FIGS. 37B-37C** depict the confirmatory/secondary screen of 12 proteins.

Expression levels of secondary screening array of 12 human membrane proteins in 293 cells based on GFP are shown in left panels of **FIGS. 37B-37C**. Antibody binding to this same array is shown in right panels of **FIGS. 37B-37C**.

FIG. 38A depicts the molecular engagement of the anti-APRIL mAb 2419 binding to APRIL based on low resolution X-ray co-crystallographic data of human APRIL (amino acids 115-250) and Fab region of mouse 2419. Structure was solved by molecular replacement using mouse APRIL described in the protein structure database. 2419 epitope within APRIL is indicated by arrow.

Variable heavy and light chains of mAb 2419 are also indicated. Structure of APRIL is depicted as a trimer.

FIG. 38B depicts residues within APRIL that comprise a subset of the 2419 epitope. This structural depiction of the 2419 epitope also highlights the fact that 2419 binds to a non-linear, quaternary epitope spanning two different monomers of APRIL within a larger trimeric complex depicted here as monomer A and monomer B. Epitope on monomer A and epitope on monomer B are indicated. 2419 epitope substantially overlaps with the high affinity receptor binding site (CRD2) and lower affinity receptor binding site (CRD1) critical for APRIL-mediated receptor signaling. CRD2 receptor binding site and CRD1 receptor binding site are outlined and indicated.

FIGS. 39A-39B depict the thermal stability of antibodies mAb 2419-1406 and 4035-062 as measured using Sypro-Orange® fluorescence scanning assay. The thermal melting temperatures (T_m values) for both 2419-1406 and 4035-062 are listed in **FIG. 39B**.

FIGS. 40A-40B depict the PK profiles of mAb 2419-1406 and 4035-062 in humanized FcRn transgenic mouse strain tg32. Tg32 mice were administered a single 5 mg/kg dose of mAb 2419-1406 or mAb 4035-062 by intravenous injection. Antibody levels in sera were determined by ELISA following timepoints taken out to 20 days. PK values are listed in **FIG. 40B**.

FIG. 41 depicts the reduction of basal serum IgA levels in mice treated with mAb 4540 and mAb 3833. 11 week old C57/BL6 mice were sub-chronically dosed (1x weekly by i.p injection) with 20 mg/kg of mAb 4540 or isotype control antibody for seven weeks. Basal serum levels of total IgA were quantified by ELISA. Median IgA levels graphed; N=5 mice per group.

DETAILED DESCRIPTION

Disclosed herein are antibody molecules that bind to APRIL, *e.g.*, human APRIL, mouse APRIL, or both, with high affinity and specificity. Advantageously, several of the antibody molecules describe herein have improved ability to reduce (*e.g.*, inhibit, block, or neutralize) one or more biological activities of APRIL. Nucleic acid molecules encoding the antibody molecules, expression vectors, host cells, compositions (*e.g.*, pharmaceutical compositions), kits, and methods for making the antibody molecules, are also provided. The antibody molecules and pharmaceutical compositions disclosed herein can be used (alone or in combination with other agents or therapeutic modalities) to treat, prevent and/or diagnose disorders and conditions, *e.g.*, disorders and conditions associated with APRIL, *e.g.*, IgA nephropathy.

Definitions

As used herein, the articles “a” and “an” refer to one or to more than one (*e.g.*, to at least one) of the grammatical object of the article.

The term “or” is used herein to mean, and is used interchangeably with, the term “and/or”, unless context clearly indicates otherwise.

“About” and “approximately” shall generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Exemplary degrees of error are within 20 percent (%), typically, within 10%, and more typically, within 5% of a given value or range of values.

The compositions and methods disclosed herein encompass polypeptides and nucleic acids having the sequences specified, or sequences substantially identical or similar thereto, *e.g.*, sequences at least 85%, 90%, 95% identical or higher to the sequence specified.

In the context of an amino acid sequence, the term “substantially identical” is used herein to refer to a first amino acid that contains a sufficient or minimum number of amino acid residues that are i) identical to, or ii) conservative substitutions of aligned amino acid residues in a second amino acid sequence such that the first and second amino acid sequences can have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain having at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence, *e.g.*, a sequence provided herein.

In the context of nucleotide sequence, the term “substantially identical” is used herein to refer to a first nucleic acid sequence that contains a sufficient or minimum number of nucleotides that are identical to aligned nucleotides in a second nucleic acid sequence such that the first and second nucleotide sequences encode a polypeptide having common functional activity, or encode a common structural polypeptide domain or a common functional polypeptide activity. For example, nucleotide sequences having at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence, *e.g.*, a sequence provided herein.

The term “functional variant” refers polypeptides that have a substantially identical amino acid sequence to the naturally-occurring sequence, or are encoded by a substantially identical nucleotide sequence, and are capable of having one or more activities of the naturally-occurring sequence.

5 Calculations of homology or sequence identity between sequences (the terms are used interchangeably herein) are performed as follows.

To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and
10 non-homologous sequences can be disregarded for comparison purposes). In a typical embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, *e.g.*, at least 40%, 50%, 60%, *e.g.*, at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or
15 nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position.

The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

20 The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In some embodiments, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch ((1970) *J. Mol. Biol.* 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package (available at www.gcg.com), using either a Blossum 62 matrix or a PAM250 matrix, and a
25 gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In certain embodiments, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at www.gcg.com), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. One suitable set of
30 parameters (and the one that should be used unless otherwise specified) are a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller ((1989) CABIOS, 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

35 The nucleic acid and protein sequences described herein can be used as a “query sequence” to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0)

of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid as described herein. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules described herein. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See www.ncbi.nlm.nih.gov.

As used herein, the term “hybridizes under low stringency, medium stringency, high stringency, or very high stringency conditions” describes conditions for hybridization and washing. Guidance for performing hybridization reactions can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6, which is incorporated by reference. Aqueous and nonaqueous methods are described in that reference and either can be used. Specific hybridization conditions referred to herein are as follows: 1) low stringency hybridization conditions in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by two washes in 0.2X SSC, 0.1% SDS at least at 50°C (the temperature of the washes can be increased to 55°C for low stringency conditions); 2) medium stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C; 3) high stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C; and preferably 4) very high stringency hybridization conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Very high stringency conditions 4) are suitable conditions and the ones that should be used unless otherwise specified.

It is understood that the molecules described herein may have additional conservative or non-essential amino acid substitutions, which do not have a substantial effect on their functions.

The term “amino acid” is intended to embrace all molecules, whether natural or synthetic, which include both an amino functionality and an acid functionality and capable of being included in a polymer of naturally-occurring amino acids. Exemplary amino acids include naturally-occurring amino acids; analogs, derivatives and congeners thereof; amino acid analogs having variant side chains; and all stereoisomers of any of any of the foregoing. As used herein the term “amino acid” includes both the D- or L- optical isomers and peptidomimetics.

A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine,

methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

The terms “polypeptide,” “peptide” and “protein” (if single chain) are used interchangeably herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling component. The polypeptide can be isolated from natural sources, can be produced by recombinant techniques from a eukaryotic or prokaryotic host, or can be a product of synthetic procedures.

The terms “nucleic acid,” “nucleic acid sequence,” “nucleotide sequence,” or “polynucleotide sequence,” and “polynucleotide” are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. The polynucleotide may be either single-stranded or double-stranded, and if single-stranded may be the coding strand or non-coding (antisense) strand. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. The nucleic acid may be a recombinant polynucleotide, or a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which either does not occur in nature or is linked to another polynucleotide in a non-natural arrangement.

The term “isolated,” as used herein, refers to material that is removed from its original or native environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated by human intervention from some or all of the co-existing materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of the environment in which it is found in nature.

As used herein, the term “treat,” *e.g.*, IgA nephropathy, means that a subject (*e.g.*, a human) who has a disorder, *e.g.*, IgA nephropathy, and/or experiences a symptom of a disorder, *e.g.*, IgA nephropathy, will, in an embodiment, suffer less a severe symptom and/or recover faster when an antibody molecule is administered than if the antibody molecule were never administered. In an embodiment, when IgA nephropathy is treated, a kidney biopsy will show less or no IgA deposits, *e.g.*, in the form of immune complexes in the mesangium of the kidney, after effective treatment for IgA nephropathy. For example, a diagnostic assay using immunofluorescence or electron microscopy will detect less or no IgA deposits in a biological sample of a subject after administration of an antibody molecule described herein for the effective treatment of IgA nephropathy. Other assays, urine tests,

blood tests, iothalamate clearance tests, or kidney imaging (*e.g.*, ultrasound, X-rays, or cystoscopy), can also be used to monitor treatment in a patient, or to detect the presence, *e.g.*, decreased presence (or absence), of a symptom of IgA nephropathy, after treatment of IgA nephropathy in the subject. Treatment can, *e.g.*, partially or completely, alleviate, ameliorate, relieve, inhibit, or reduce the severity of, and/or reduce incidence, and optionally, delay onset of, one or more manifestations of the effects or symptoms, features, and/or causes of a disorder, *e.g.*, IgA nephropathy. In an embodiment, treatment is of a subject who does not exhibit certain signs of a disorder, *e.g.*, IgA nephropathy, and/or of a subject who exhibits only early signs of a disorder, *e.g.*, nephropathy. In an embodiment, treatment is of a subject who exhibits one or more established signs of a disorder, *e.g.*, IgA nephropathy. In an embodiment, treatment is of a subject diagnosed as suffering from a disorder, *e.g.*, IgA nephropathy.

As used herein, the term “prevent,” a disorder, *e.g.*, IgA nephropathy, means that a subject (*e.g.*, a human) is less likely to have the disorder, *e.g.*, IgA nephropathy, if the subject receives the antibody molecule.

Various aspects of the compositions and methods herein are described in further detail below. Additional definitions are set out throughout the specification.

APRIL

APRIL (A Proliferation Inducing Ligand), also known as CD256, TNF- and APOL-related Leukocyte Expressed Ligand 2 (TALL-2), or TNF-related Death Ligand 1 (TRDL-1), is a TNF family cytokine encoded by the Tumor Necrosis Factor Ligand Superfamily Member 13 (*TNFSF13*) gene (also known as *APRIL*, *TALL2*, or *ZTNF2*). APRIL plays a role in a number of biological processes such as signal transduction, regulation of cell proliferation, and IgA class switching (Hahne *et al.* (1998) *J. Exp. Med.* 188:1185-1190 (1998); Castigli *et al. Proc. Natl. Acad. Sci. U.S.A.* 101:3903-3908 (2004)).

APRIL is both functionally and structurally related to BAFF (B Cell Activating Factor F13B) also known as BLyS (B lymphocyte stimulator). Both cytokines are involved in regulating keys aspects of innate and adaptive immune functions. Both APRIL and BAFF bind the lymphocyte receptors TACI (transmembrane activator and CAML interactor) and BCMA (B cell maturation antigen). APRIL and BAFF appear to heterologously interact with each other through protein-protein interactions. While both APRIL and BAFF share biochemical (receptor binding), immunological and even some structural overlap (*e.g.*, as it relates to the three-dimensional topology of their respective receptor binding domains), the two cytokines, nevertheless, are both structurally and functionally distinct. APRIL binds to biologically relevant heparan sulfate (present in the extracellular matrices of cells as heparan sulfate proteoglycans); BAFF does not. This interaction plays a critical biological function with respect to promoting the oligomerization state of APRIL in concert with its localized interaction with TACI, which likewise requires HSPGS for full activity. Unlike BAFF which acts as

a potent activator of B cells inclusive of both proliferation and differentiation, APRIL would appear to function more particularly with respect to the modulation of B cell phenotype, *e.g.*, as it relates to IgA production and the differentiation/survival of IgA positive plasma cells. As such, a targeted disruption in APRIL-receptor signaling is expected to have less perturbative effects on B cell homeostasis and overall immune function in comparison to other immune related therapeutics that target BAFF (*e.g.*, belimumab) or anti CD20 therapies (*e.g.*, rituximab) that largely target pre and early B cells. APRIL has also been shown to be expressed at high levels on other myeloid related cells and lymphoid tissues, as well as hematological cancers (*e.g.*, myeloma, chronic lymphocytic leukemia (CLL)) and solid tumors (*e.g.*, colon, thyroid, and breast).

Exemplary amino acid and nucleotide sequences of human APRIL are described, *e.g.*, in Hahne *et al. J. Exp. Med.* 188:1185-1190 (1998); Shu *et al. J. Leukoc. Biol.* 65:680-683 (1999); Kelly *et al. Cancer Res.* 60:1021-1027(2000); and Pradet-Balade *et al. EMBO J.* 21:5711-5720 (2002).

The amino acid sequence of human APRIL (isoform alpha, also referred to as the “canonical” sequence (SEQ ID NO: 85)) is provided as follows.

```
>huAPRIL
MPASSPFLAPKGPPGNMGGPVREPALSVALWLSWGAALGAVACAMALLTQQTELQSLRREVSRLQGT
GGPSQNGEGYPWQSLPEQSSDALEAWENGERSRKRRVLTQKQKKQHSVLHLVPINATSKDDSDVTEV
MWQPALRRGRGLQAQGYGVRIQDAGVYLLYSQVLFQDVTFTMGQVVSREGQGRQETLFRICIRSMPSHP
DRAYNSCYSAGVFHLHQGDILSVIIPRARA KLNLSPHGTFLGFVKL
```

There are several isoforms of human APRIL produced by alternative splicing.

Isoform beta has the following amino acid sequence (SEQ ID NO: 86):

```
>sp|O75888-2|TNF13_HUMAN Isoform Beta of Tumor necrosis factor
ligand superfamily member 13 OS=Homo sapiens GN=TNFSF13
MPASSPFLAPKGPPGNMGGPVREPALSVALWLSWGAALGAVACAMALLTQQTELQSLRREVSRLQGT
GGPSQNGEGYPWQSLPEQSSDALEAWENGERSRKRRVLTQKQKNSDVTEVMWQPALRRGRGLQAQG
YGVRIQDAGVYLLYSQVLFQDVTFTMGQVVSREGQGRQETLFRICIRSMPSHPDRAYNSCYSAGVFHLH
QGDILSVIIPRARA KLNLSPHGTFLGFVKL
```

The sequence of isoform beta differs from the canonical sequence as follows: amino acids 113-129 of SEQ ID NO: 85: KQHSVLHLVPINATSKD → N

Isoform gamma has the following amino acid sequence (SEQ ID NO: 87):

```
>sp|O75888-3|TNF13_HUMAN Isoform Gamma of Tumor necrosis factor
ligand superfamily member 13 OS=Homo sapiens GN=TNFSF13
MPASSPFLAPKGPPGNMGGPVREPALSVALWLSWGAALGAVACAMALLTQQTELQSLRREVSRLQGT
GGPSQNGEGYPWQSLPEQSSDALEAWENGERSRKRRVLTQKQKKQHSVLHLVPINATSKDDSDVTEV
MWQPALRRGRGLQAQGYGVRIQDAGVYLLYSQVLFQDVTFTMGQVVSREGQGRQETLFRICIRSMPSHP
DRAYNSCYSAGVFHLHQGDILSVIIPRARA KLNLSPHGTFLGL
```

The sequence of isoform gamma differs from the canonical sequence as follows: amino acids 247-249: Missing.

Isoform 4 has the following amino acid sequence (SEQ ID NO: 88):

```
>sp|O75888-4|TNF13_HUMAN Isoform 4 of Tumor necrosis factor ligand
superfamily member 13 OS=Homo sapiens GN=TNFSF13
```

MPASSPFLAPKGPFGNMGFPVREFALSVALWLSWGAALGAVACAMALLTQQTELQSLRREVSRLQGT
GGPSQNGEGYPWQSLPEQHSVLHLVLPINATSKDDSDVTEVMWQPALRRGRGLQAQGYGVRIQDAGVYL
LYSQVLFQDVTFTMGQVVSREGQGRQETLFRICIRSMPSHPDRAYNSCYSAGVFHLHQGDILSVIIPRA
RAKLNLSPHGTFLGFVKL

5

The sequence of isoform 4 differs from the canonical sequence as follows: amino acids 86-113: Missing.

Isoform TWE-PRIL has the following amino acid sequence (SEQ ID NO: 89):

>sp|O43508-2|TNF12_HUMAN Isoform TWE-PRIL of Tumor necrosis factor
ligand superfamily member 12 OS=Homo sapiens GN=TNFSF12
MAARRSQRRRGRRGEPGTALLVPLALGLGLALACLGLLLAVVSLGSRASLSAQEPAQEELVAEEDQDP
SELNPQTEESQDPAPFLNRLVRPRRSAPKGRKTRARRAIAAHYEVHPRPGQDGAQAGVDGTVSGWEEA
RINSSPLRYNRQIGEFIVTRAGLYLYLYCQSSDALEAWENGERSRKRRAVLTQKQKKQHSVLHLVLPIN
ATSKDDSDVTEVMWQPALRRGRGLQAQGYGVRIQDAGVYLLYSQVLFQDVTFTMGQVVSREGQGRQET
LFRICIRSMPSHPDRAYNSCYSAGVFHLHQGDILSVIIPRAKLNLSPHGTFLGFVKL

15

Isoform 5 has the following amino acid sequence (SEQ ID NO: 90):

>sp|O75888-5|TNF13_HUMAN Isoform 5 of Tumor necrosis factor ligand
superfamily member 13 OS=Homo sapiens GN=TNFSF13
MGGFVREFALSVALWLSWGAALGAVACAMALLTQQTELQSLRREVSRLQGTGGPSQNGEGYPWQSLPE
QHSVLHLVLPINATSKDDSDVTEVMWQPALRRGRGLQAQGYGVRIQDAGVYLLYSQVLFQDVTFTMGQV
VSREGQGRQETLFRICIRSMPSHPDRAYNSCYSAGVFHLHQGDILSVIIPRAKLNLSPHGTFLGFVK
L

25

The sequence of isoform 5 differs from the canonical sequence as follows: amino acids 1-17: Missing; amino acids 87-114: Missing.

Other variant and alternative sequences of human APRIL are described, *e.g.*, in The MGC Project Team, *Genome Res.* 14:2121-2127 (2004); Ota *et al. Nat. Genet.* 36:40-45 (2004); and Kelly *et al. Cancer Res.* 60:1021-1027 (2000).

30

As used herein, when an anti-APRIL antibody molecule binds, or substantially binds, to human APRIL, it binds, or substantially binds, to one or more isoforms of human APRIL, *e.g.*, one or more isoforms of human APRIL described herein. In an embodiment, the antibody molecule binds or substantially binds to human APRIL having the amino acid sequence of SEQ ID NO: 85.

35

Exemplary amino acid and nucleotide sequences of mouse APRIL are described, *e.g.*, in Yu *et al. Nat. Immunol.* 1:252-256 (2000); Carninci *et al. Science* 309:1559-1563 (2005); The MGC Project Team, *Genome Res.* 14:2121-2127 (2004); and Bossen *et al. J. Biol Chem.* 281: 13964-13971 (2006).

The amino acid sequence of mouse APRIL isoform 1 (SEQ ID NO: 91) is provided as follows.

40

>muAPRIL
MPASSPGHMGGSVREFALSVALWLSWGAALGAVTCAVALLIQQTELQSLRREVSRLQRSGGPSQKQGE
RPWQSLWEQSPDVLEAWKDGAKSRRRRRAVLTQKHKKKHSVLHLVFPVNITSKADSDVTEVMWQPVLRG
RGLEAQGDIVRVWDTGIYLLYSQVLFHDVTFTMGQVVSREGQGRRETFRICIRSMPSDPDRAYNSCYS
AGVFHLHQGDIITVKIPRANAKLSLSPHGTFLGFVKL

45

The amino acid sequence of mouse APRIL isoform 2 (SEQ ID NO: 92) is provided as follows.

5 MPASSPGHMGGSVREPALSVALWLSWGAVLGAVTCAVALLIQQTELOSLRREVSRLQRSGGPSQKQGE
 RPWQSLWEQSPDVLEAWKDGAKSRRRRRAVLTKHKKKHSVLHLPVNVNITSKDSVDTEVMWQPVLRGR
 GLEAQGDIVRVWDTGIYLLYSQVLFHDVTFMTGQVVSREGQGRRETLFRCIRSMPSDPDRAYNSCYSA
 GVFHLHQGDIITVKIPRANAKLSLSPHGTFLGFVKL

As used herein, when an anti-APRIL antibody molecule binds, or substantially binds, to
 10 mouse APRIL, it binds, or substantially binds, to one or more isoforms of mouse APRIL, *e.g.*, one or
 more isoforms of mouse APRIL described herein. In an embodiment, the antibody molecule binds or
 substantially binds to mouse APRIL having the amino acid sequence of SEQ ID NO: 91, SEQ ID NO:
 92, or both.

As used herein, when an anti-APRIL antibody molecule does not bind, or does not
 15 substantially bind, to mouse APRIL, it does not bind, or does not substantially bind, to one or more
 isoforms of mouse APRIL, *e.g.*, one or more isoforms of mouse APRIL described herein. In an
 embodiment, the antibody molecule does not bind, or does not substantially bind, to mouse APRIL
 having the amino acid sequence of SEQ ID NO: 91 or 92. In a typical embodiment, the antibody
 molecule does not bind, or does not substantially bind, to mouse APRIL having the amino acid
 20 sequence of SEQ ID NO: 91 and mouse APRIL having the amino acid sequence of SEQ ID NO: 92.

Sequence alignment of exemplary human and mouse APRIL proteins (SEQ ID NOS: 85 and
 91, respectively) is shown in **FIG. 13**.

Epitope

25 The antibody molecule described herein can bind to an epitope on APRIL (*e.g.*, human
 APRIL, mouse APRIL, or both). For example, an epitope bound by an antibody molecule described
 herein can include one or more epitope contact points described herein.

In an embodiment, the antibody molecule contacts (*e.g.*, binds, or substantially binds, to) one
 or more residues, or one or more regions, as described in any of **Tables 3-4 or 6-8**, or any of **FIGS.**
 30 **14, 22, 23A-23B, 24A-24B, 25A-25B, or 38A-38B**.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one
 or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all)
 of the amino acid residues shown in **Table 3**. In an embodiment, the antibody molecule contacts
 (*e.g.*, binds or substantially binds to) all of the amino acid residues shown in **Table 3**. For example,
 35 the antibody molecules described herein can contact the amino acid residues shown in **Table 3** in a
 manner that includes binding across two APRIL monomers (*e.g.*, as depicted positionally in **Table 3**
 as A vs. B). While not wishing to be bound by theory, it is believed that in an embodiment, at least
 some of the amino acid residues shown in **Table 3** contribute to high affinity interactions between
 APRIL and the CDR2 domain of TACI. In an embodiment, contacting one or more of the amino acid

residues in **Table 3** with an antibody molecule described herein inhibits, or substantially inhibits, binding of APRIL to TACI.

Exemplary human APRIL amino acid residues that can bind to the anti-APRIL antibody molecules described herein are shown in **Table 3**. A structural representation of this epitope (*e.g.*, defined both spatially and conformationally) is depicted in **FIG. 14**.

Table 3. Exemplary Human APRIL Amino Acid Residues that Bind to Anti-APRIL Antibodies (amino acid numbering based on SEQ ID NO: 85)

Monomer	Amino Acid Position	Amino Acid
A	130	Asp
A	131	Ser
A	132	Asp
A	174	Val
A	175	Thr
A	176	Phe
A	177	Thr
A	178	Met
A	179	Gly
A	180	Gln
A	181	Val
A	192	Thr
A	195	Arg
A	196	Cys
A	197	Ile
A	200	Met
A	201	Pro
A	202	Ser
A	208	Tyr
A	230	Pro
A	231	Arg
A	232	Ala
A	241	His
B	170	Leu
B	205	Asp
B	206	Arg

10 In another embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all) of the amino acid residues shown in **Table 4**. In another embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) all of the amino acid residues shown in **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds to, the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*,
15 the loop connecting β -sheets G and H), or both, on APRIL.

A structural (spatial) representation of this epitope (sometimes referred herein as “core region”) is depicted in **FIG. 15**. As shown in **FIG. 15**, each APRIL protein molecule contains two packed antiparallel eight-stranded β -sheets (A to G), one inner and one outer, in a β -jelly roll topology. These B sheets are connected by loops that also define (based on secondary structure definitions) a desired epitope. While not wishing to be bound by theory, it is believed that as these positions/structures define a subset of key interactions with APRIL and the CRD2 domain of TACI, optimal inhibition of APRIL binding to TACI by such an antibody would be achieved.

Table 4. Exemplary Human APRIL Amino Acid Residues that Bind to Anti-APRIL Antibodies
(amino acid numbering based on SEQ ID NO: 85)

Amino Acid Position	Amino Acid
174	Val
175	Thr
176	Phe
177	Thr
178	Met
179	Gly
180	Gln
181	Val
230	Pro
231	Arg
232	Ala

In another embodiment, the antibody molecule does not bind to one, two, or all of Asp129, Arg233, or His203, on human APRIL (*e.g.*, SEQ ID NO: 85). For example, one or more mutations at these positions, *e.g.*, Asp129Ala, Arg233Asn, His203Asp, or any combination thereof, would not reduce, or substantially reduce, the binding affinity of the antibody molecule to human APRIL, or the inhibitory effect of the antibody molecule on a human APRIL activity (*e.g.*, neutralization of APRIL binding to TACI).

In yet another embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL (*e.g.*, SEQ ID NO: 85) from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL (*e.g.*, SEQ ID NO: 91) from positions 96-105.

In another embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all) of the amino acid residues shown in **Table 7**. In another embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) all of the amino acid residues shown in **Table 7**.

Table 7. Exemplary Human APRIL Amino Acid Residues that Bind to Anti-APRIL Antibodies
(amino acid numbering based on SEQ ID NO: 85)

Amino Acid Position	Amino Acid
132	Asp
170	Leu
175	Thr
176	Phe
177	Thr
178	Met
181	Val
192	Thr
195	Arg
197	Ile
205	Asp
206	Arg
208	Tyr
228	Iso
230	Pro
231	Arg
232	Ala
241	His

In an embodiment, the antibody molecule, *e.g.*, an anti-APRIL antibody molecule having one,
5 two, three, four, five or six CDRs of any of monoclonal antibodies 2419, 2419-0105, 2419-0205,
2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1210, 2419-1305, 2419-
1306, 2419-1310, or 2419-1406, binds to one or more amino acids described in **Table 7**. In another
embodiment, the antibody molecule, *e.g.*, a human-specific, anti-APRIL antibody molecule, *e.g.*,
having one, two, three, four, five or six CDRs of any of monoclonal antibodies 2419, 2419-0105,
10 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1210, 2419-
1305, 2419-1306, 2419-1310, 2419-1406, binds to mouse APRIL when one or more (*e.g.*, 2, 3, 4 or
all) following positions within mouse APRIL (mouse APRIL numbering applies) are mutated, *e.g.*, to
the following: A120D, N224R, H163Q, K219I, or R181Q. In yet another embodiment, the antibody
molecule, *e.g.*, a human-specific, anti-APRIL antibody molecule, *e.g.*, having one, two, three, four,
15 five or six CDRs of any of monoclonal antibodies 2419, 2419-0105, 2419-0205, 2419-0206, 2419-
0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1210, 2419-1305, 2419-1306, 2419-1310,
2419-1406, binds to mouse APRIL when the lysine at position 219 (mouse APRIL numbering
applies) is mutated, *e.g.*, to an isoleucine (*i.e.*, K219I).

20 In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one
or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27,
28, 29, 30, 31, 32, or all) of the amino acid residues of human APRIL shown in **Table 6**. In an

embodiment, the antibody molecule is an antibody molecule described herein, *e.g.*, monoclonal antibody 2218, 2419, 2621, 2622, 3125, 3327, 3525, 3530, 4035, 3934, 3833, 3631, 3732, 4338, 4540, or 4237.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) of the amino acid residues of human APRIL chosen from D132, V174, F176, V181, Q190, R195, R206, Y208, I228, or N237. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, or all) of the amino acid residues of human APRIL chosen from V174, F176, Q190, R195, R206, or Y208. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or I228. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, or all) of the amino acid residues of human APRIL chosen from V174, R206, or Y208.

In an embodiment, the antibody molecule does not contact (*e.g.*, does not bind or does not substantially bind to) at least one (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) of the amino acid residues of human APRIL shown in **Table 6**. In an embodiment, the antibody molecule is an antibody molecule described herein, *e.g.*, monoclonal antibody 2218, 2419, 2621, 2622, 3125, 3327, 3525, 3530, 4035, 3934, 3833, 3631, 3732, 4338, 4540, or 4237.

In an embodiment, the antibody molecule does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, 4, 5, 6, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, S226, I228, Y208, or N237. In an embodiment, the antibody molecule does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from V181, S226, I228, or N237. In an embodiment, the antibody molecule does not contact (*e.g.*, does not bind or does not substantially bind to) one or both of the amino acid residues of human APRIL chosen from Y208 or N237. In an embodiment, the antibody molecule does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or N237.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, or all) of the amino acid residues of human APRIL chosen from V174, F176, Q190, R195, R206, or Y208; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from V181, S226, I228, or N237. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or both of the amino acid residues of human APRIL chosen from V174 or R206; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or both of the amino acid residues of human APRIL chosen from V181 or N237 (and optionally S226). In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) heavy chain CDRs, one or more (*e.g.*, two or three) light chain CDRs, or both of monoclonal antibody 4035. In an embodiment, the

antibody molecule comprises a heavy chain region, a light chain variable region, or both, of monoclonal antibody 4035. In an embodiment, monoclonal antibody 4035 is a humanized antibody molecule.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or I228; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or both of the amino acid residues of human APRIL chosen from Y208 or N237. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) amino acid residue I228 of human APRIL; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or both of the amino acid residues of human APRIL chosen from Y208 or N237. In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) heavy chain CDRs, one or more (*e.g.*, two or three) light chain CDRs, or both of monoclonal antibody 2419. In an embodiment, the antibody molecule comprises a heavy chain region, a light chain variable region, or both, of monoclonal antibody 2419. In an embodiment, monoclonal antibody 2419 is a humanized antibody molecule.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, or all) of the amino acid residues of human APRIL chosen from V174, R206, or Y208; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or N237. In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or both of the amino acid residues of human APRIL chosen from V174 or R206; and does not contact (*e.g.*, does not bind or does not substantially bind to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from F176, V181, Q190, or N237. In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) heavy chain CDRs, one or more (*e.g.*, two or three) light chain CDRs, or both of monoclonal antibody 3833. In an embodiment, the antibody molecule comprises a heavy chain region, a light chain variable region, or both, of monoclonal antibody 3833. In an embodiment, monoclonal antibody 3833 is a humanized antibody molecule.

In an embodiment, the epitope overlaps with a CRD2 receptor binding site. In an embodiment, the epitope is non-linear epitope, *e.g.*, that spans across a monomer interface. In an embodiment, the epitope is in a region associated with both TACI and BCMA receptor blocking.

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or all) of the amino acid residues of human APRIL chosen from V133, V181, E185, Q187, G188, R189, Q190, E191, T192, R195, H218, L219, H220, S226, I228, P230 (located in monomer A). In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) of the amino acid residues of human APRIL chosen from V121, I123, Q139, P140, A141, L142, N237, S239, P240, or H241 (located in monomer B). In an embodiment, the antibody molecule contacts (*e.g.*, binds or

substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) of the amino acid residues of human APRIL chosen from V133, V181, E185, Q187, G188, R189, Q190, E191, T192, R195, H218, L219, H220, S226, I228, P230 (located in monomer A); V121, I123, Q139, P140, A141, L142, N237, S239, P240, or H241 (located in monomer B).

In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, or all) of the amino acid residues of human APRIL chosen from V181, Q190, T192, and I228 (located in monomer A). In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or both of the amino acid residues of human APRIL chosen from A141 or H241 (located in monomer B). In an embodiment, the antibody molecule contacts (*e.g.*, binds or substantially binds to) one or more (*e.g.*, 2, 3, 4, 5, or all) of the amino acid residues of human APRIL chosen from V181, Q190, T192, and I228 (located in monomer A); A141 or H241 (located in monomer B).

In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) heavy chain CDRs, one or more (*e.g.*, two or three) light chain CDRs, or both of monoclonal antibody 2419. In an embodiment, the antibody molecule comprises a heavy chain region, a light chain variable region, or both, of monoclonal antibody 2419. In an embodiment, monoclonal antibody 2419 is a humanized antibody molecule.

In an embodiment, the epitope comprise one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) of the amino acid residues of human APRIL chosen from V133, V181, E185, Q187, G188, R189, Q190, E191, T192, R195, H218, L219, H220, S226, I228, P230 (located in monomer A); V121, I123, Q139, P140, A141, L142, N237, S239, P240, or H241 (located in monomer B). In an embodiment, the epitope comprises one or more (*e.g.*, 2, 3, 4, 5, or all) of the amino acid residues of human APRIL chosen from V181, Q190, T192, and I228 (located in monomer A); A141 or H241 (located in monomer B).

In an embodiment, a structural representation of this epitope is depicted in **FIG. 38B**. In an embodiment, the epitope comprises one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all) of the amino acid residues shown in **Table 8**.

In an embodiment, the antibody molecule contacts (*e.g.*, binds, or substantially binds, to) all of the amino acid residues shown in any of **Tables 3-4 or 7-8**. In an embodiment, the epitope comprises, or consists of, all of the amino acid residues shown in any of **Tables 3-4 or 7-8**.

In an embodiment, the antibody molecule has one or more of the following properties described herein, *e.g.*, one or more (*e.g.*, two, three or all) of: (i) binds, or substantially binds, to human APRIL; (ii) binds, or substantially binds, to mouse APRIL; (iii) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to TACI (*e.g.*, human TACI, mouse TACI, or both); or (iv) inhibits, or substantially inhibits, binding of APRIL (*e.g.*, human APRIL, mouse APRIL, or both) to BCMA (*e.g.*, human BCMA, mouse BCMA, or both). In an

embodiment, the antibody molecule binds, or substantially binds, to mouse APRIL. In another embodiment, the antibody molecule does not bind, or binds with low affinity, to mouse APRIL.

Antibody Molecules

5 Disclosed herein are antibody molecules that bind to APRIL, *e.g.*, an APRIL molecule described herein.

As used herein, the term “antibody molecule” refers to a protein, *e.g.*, an immunoglobulin chain or a fragment thereof, comprising at least one immunoglobulin variable domain sequence. The term “antibody molecule” includes, for example, full-length, mature antibodies and antigen-binding
 10 fragments of an antibody. For example, an antibody molecule can include a heavy (H) chain variable domain sequence (abbreviated herein as VH), and a light (L) chain variable domain sequence (abbreviated herein as VL). In another example, an antibody molecule includes two heavy (H) chain variable domain sequences and two light (L) chain variable domain sequence, thereby forming two antigen binding sites, such as Fab, Fab', F(ab')₂, Fc, Fd, Fd', Fv, single chain antibodies (scFv for
 15 example), single variable domain antibodies, diabodies (Dab) (bivalent and bispecific), and chimeric (*e.g.*, humanized) antibodies, which may be produced by the modification of whole antibodies or those synthesized *de novo* using recombinant DNA technologies. These functional antibody fragments retain the ability to selectively bind with their respective antigen or receptor. Antibodies and antibody fragments can be from any class of antibodies including, but not limited to, IgG, IgA, IgM, IgD, and IgE, and from any subclass (*e.g.*, IgG1, IgG2, IgG3, and IgG4) of antibodies. The
 20 antibody molecules can be monoclonal or polyclonal. The antibody molecule can also be a human, humanized, CDR-grafted, or *in vitro* generated antibody. The antibody molecule can have a heavy chain constant region chosen from, *e.g.*, IgG1, IgG2, IgG3, or IgG4. The antibody molecule can also have a light chain chosen from, *e.g.*, kappa or lambda. The term “immunoglobulin” (Ig) is used
 25 interchangeably with the term “antibody” herein.

Examples of antigen-binding fragments include: (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a
 30 single arm of an antibody, (v) a diabody (dAb) fragment, which consists of a VH domain; (vi) a camelid or camelized variable domain; (vii) a single chain Fv (scFv), see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; (viii) a single domain antibody. These antibody fragments may be obtained using any suitable method, including several conventional techniques known to those with skill in the art, and the fragments can
 35 be screened for utility in the same manner as are intact antibodies.

The term “antibody” includes intact molecules as well as functional fragments thereof. Constant regions of the antibodies can be altered, *e.g.*, mutated, to modify the properties of the

antibody (*e.g.*, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, or complement function).

The antibody molecule can be a single chain antibody. A single-chain antibody (scFv) may be engineered (*see*, for example, Colcher, D. *et al.* (1999) *Ann N Y Acad Sci* 880:263-80; and Reiter, Y. (1996) *Clin Cancer Res* 2:245-52). The single chain antibody can be dimerized or multimerized to generate multivalent antibodies having specificities for different epitopes of the same target protein.

The antibody molecules disclosed herein can also be single domain antibodies. Single domain antibodies can include antibodies whose complementary determining regions are part of a single domain polypeptide. Examples include, but are not limited to, heavy chain antibodies, antibodies naturally devoid of light chains, single domain antibodies derived from conventional 4-chain antibodies, engineered antibodies and single domain scaffolds other than those derived from antibodies. Single domain antibodies may be any of the art, or any future single domain antibodies. Single domain antibodies may be derived from any species including, but not limited to mouse, human, camel, llama, fish, shark, goat, rabbit, and bovine. According to some aspects, a single domain antibody is a naturally occurring single domain antibody known as heavy chain antibody devoid of light chains. Such single domain antibodies are disclosed in WO 94/04678, for example. For clarity reasons, this variable domain derived from a heavy chain antibody naturally devoid of light chain is known herein as a VHH or nanobody to distinguish it from the conventional VH of four chain immunoglobulins. Such a VHH molecule can be derived from antibodies raised in *Camelidae* species, for example in camel, llama, dromedary, alpaca and guanaco. Other species besides *Camelidae* may produce heavy chain antibodies naturally devoid of light chain; such VHHs are also contemplated.

The VH and VL regions can be subdivided into regions of hypervariability, termed “complementarity determining regions” (CDR), interspersed with regions that are more conserved, termed “framework regions” (FR or FW). The terms “complementarity determining region,” and “CDR,” as used herein refer to the sequences of amino acids within antibody variable regions which confer antigen specificity and binding affinity. As used herein, the terms “framework,” “FW” and “FR” are used interchangeably.

The extent of the framework region and CDRs has been precisely defined by a number of methods (*see*, Kabat, E. A., *et al.* (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Chothia, C. *et al.* (1987) *J. Mol. Biol.* 196:901-917; and the AbM definition used by Oxford Molecular’s AbM antibody modeling software. *See*, generally, *e.g.*, Protein Sequence and Structure Analysis of Antibody Variable Domains. In: Antibody Engineering Lab Manual (Ed.: Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg). In an embodiment, the following definitions are used: AbM definition of CDR1 of the heavy chain variable domain and Kabat definitions for the other CDRs. In an embodiment, Kabat definitions are used for all CDRs. In addition, embodiments

described with respect to Kabat or AbM CDRs may also be implemented using Chothia hypervariable loops. Each VH and VL typically includes three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4.

As used herein, an “immunoglobulin variable domain sequence” refers to an amino acid sequence which can form the structure of an immunoglobulin variable domain. For example, the sequence may include all or part of the amino acid sequence of a naturally-occurring variable domain. For example, the sequence may or may not include one, two, or more N- or C-terminal amino acids, or may include other alterations that are compatible with formation of the protein structure.

The term “antigen-binding region” refers to the part of an antibody molecule that comprises determinants that form an interface that binds to an antigen, *e.g.*, APRIL, or an epitope thereof. With respect to proteins (or protein mimetics), the antigen-binding region typically includes one or more loops (of at least, *e.g.*, four amino acids or amino acid mimics) that form an interface that binds to the antigen, *e.g.*, APRIL. Typically, the antigen-binding region of an antibody molecule includes at least one or two CDRs and/or hypervariable loops, or more typically at least three, four, five or six CDRs and/or hypervariable loops.

The terms “compete” or “cross-compete” are used interchangeably herein to refer to the ability of an antibody molecule to interfere with binding of an anti-APRIL antibody molecule, *e.g.*, an anti-APRIL antibody molecule provided herein, to a target, *e.g.*, APRIL. The interference with binding can be direct or indirect (*e.g.*, through an allosteric modulation of the antibody molecule or the target). The extent to which an antibody molecule is able to interfere with the binding of another antibody molecule to the target, and therefore whether it can be said to compete, can be determined using a competition binding assay, for example, a FACS assay, an ELISA or BIACORE assay. In an embodiment, a competition binding assay is a quantitative competition assay. In an embodiment, a first anti-APRIL antibody molecule is said to compete for binding to the target with a second anti-APRIL antibody molecule when the binding of the first antibody molecule to the target is reduced by 10% or more, *e.g.*, 20% or more, 30% or more, 40% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, 99% or more in a competition binding assay (*e.g.*, a competition assay described herein).

The terms “monoclonal antibody” or “monoclonal antibody composition” as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope. A monoclonal antibody can be made by hybridoma technology or by methods that do not use hybridoma technology (*e.g.*, recombinant methods).

An “effectively human” protein is a protein that does not evoke a neutralizing antibody response, *e.g.*, the human anti-murine antibody (HAMA) response. HAMA can be problematic in a number of circumstances, *e.g.*, if the antibody molecule is administered repeatedly, *e.g.*, in treatment

of a chronic or recurrent disease condition. A HAMA response can make repeated antibody administration potentially ineffective because of an increased antibody clearance from the serum (*see, e.g., Saleh et al., Cancer Immunol. Immunother.*, 32:180-190 (1990)) and also because of potential allergic reactions (*see, e.g., LoBuglio et al., Hybridoma*, 5:5117-5123 (1986)).

5 The antibody molecule can be a polyclonal or a monoclonal antibody. In some embodiments, the antibody can be recombinantly produced, *e.g.,* produced by any suitable phage display or combinatorial methods.

 Various phage display and combinatorial methods for generating antibodies are known in the art (as described in, *e.g.,* Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* International Publication
10 No. WO 92/18619; Dower *et al.* International Publication No. WO 91/17271; Winter *et al.* International Publication WO 92/20791; Markland *et al.* International Publication No. WO 92/15679; Breitling *et al.* International Publication WO 93/01288; McCafferty *et al.* International Publication No. WO 92/01047; Garrard *et al.* International Publication No. WO 92/09690; Ladner *et al.* International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982, the contents of all of which are incorporated by reference herein).

20 In an embodiment, the antibody molecule is a fully human antibody (*e.g.,* an antibody made in a mouse which has been genetically engineered to produce an antibody from a human immunoglobulin sequence), or a non-human antibody, *e.g.,* a rodent (mouse or rat), goat, primate (*e.g.,* monkey), camel antibody. In an embodiment, the non-human antibody is a rodent (mouse or rat antibody). Methods of producing rodent antibodies are known in the art.

25 Human monoclonal antibodies can be generated using transgenic mice carrying the human immunoglobulin genes rather than the mouse system. Splenocytes from these transgenic mice immunized with the antigen of interest are used to produce hybridomas that secrete human mAbs with specific affinities for epitopes from a human protein (*see e.g.,* Wood *et al.* International Application WO 91/00906, Kucherlapati *et al.* PCT publication WO 91/10741; Lonberg *et al.* International
30 Application WO 92/03918; Kay *et al.* International Application 92/03917; Lonberg, N. *et al.* 1994 *Nature* 368:856-859; Green, L.L. *et al.* 1994 *Nature Genet.* 7:13-21; Morrison, S.L. *et al.* 1994 *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Bruggeman *et al.* 1993 *Year Immunol* 7:33-40; Tuailon *et al.* 1993 *PNAS* 90:3720-3724; Bruggeman *et al.* 1991 *Eur J Immunol* 21:1323-1326).

35 An antibody can be one in which the variable region, or a portion thereof, *e.g.,* the CDRs, are generated in a non-human organism, *e.g.,* a rat or mouse. Chimeric, CDR-grafted, and humanized antibodies are within the invention. Antibodies generated in a non-human organism, *e.g.,* a rat or

mouse, and then modified, *e.g.*, in the variable framework or constant region, to decrease antigenicity in a human are within the invention.

Chimeric antibodies can be produced by any suitable recombinant DNA technique. Several are known in the art (*see* Robinson *et al.*, International Patent Application Publication No.

- 5 WO1987/002671; Akira, *et al.*, European Patent Application Publication No. 184,187; Taniguchi, M., European Patent Application Publication No. 171,496; Morrison *et al.*, European Patent Application Publication No. 173,494; Neuberger *et al.*, International Patent Application Publication No. WO 86/01533; Cabilly *et al.* U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent Application Publication No. 125,023; Better *et al.* (1988 *Science* 240:1041-1043); Liu *et al.* (1987) *PNAS*
 10 84:3439-3443; Liu *et al.*, 1987, *J. Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.*, 1987, *Canc. Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; and Shaw *et al.*, 1988, *J. Natl Cancer Inst.* 80:1553-1559).

- A humanized or CDR-grafted antibody will have at least one or two but generally all three recipient CDRs (of heavy and or light immunoglobulin chains) replaced with a donor CDR. The
 15 antibody may be replaced with at least a portion of a non-human CDR or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to lipopolysaccharide. In an embodiment, the donor will be a rodent antibody, *e.g.*, a rat or mouse antibody, and the recipient will be a human framework or a human consensus framework. Typically, the immunoglobulin providing the CDRs is called the
 20 “donor” and the immunoglobulin providing the framework is called the “acceptor.” In some embodiments, the donor immunoglobulin is a non-human (*e.g.*, rodent). The acceptor framework is typically a naturally-occurring (*e.g.*, a human) framework or a consensus framework, or a sequence about 85% or higher, *e.g.*, 90%, 95%, 99% or higher identical thereto.

- As used herein, the term “consensus sequence” refers to the sequence formed from the most
 25 frequently occurring amino acids (or nucleotides) in a family of related sequences (See *e.g.*, Winnaker, From Genes to Clones (Verlagsgesellschaft, Weinheim, Germany 1987). In a family of proteins, each position in the consensus sequence is occupied by the amino acid occurring most frequently at that position in the family. If two amino acids occur equally frequently, either can be included in the consensus sequence. A “consensus framework” refers to the framework region in the consensus
 30 immunoglobulin sequence.

An antibody can be humanized by any suitable method, and several such methods known in the art (*see e.g.*, Morrison, S. L., 1985, *Science* 229:1202-1207, by Oi *et al.*, 1986, *BioTechniques* 4:214, and by Queen *et al.* US 5,585,089, US 5,693,761 and US 5,693,762, the contents of all of which are hereby incorporated by reference).

- 35 Humanized or CDR-grafted antibodies can be produced by CDR-grafting or CDR substitution, wherein one, two, or all CDRs of an immunoglobulin chain can be replaced. *See e.g.*, U.S. Patent 5,225,539; Jones *et al.* 1986 *Nature* 321:552-525; Verhoeyan *et al.* 1988 *Science*

239:1534; Beidler *et al.* 1988 *J. Immunol.* 141:4053-4060; Winter US 5,225,539, the contents of all of which are hereby expressly incorporated by reference. Winter describes a CDR-grafting method which may be used to prepare humanized antibodies (UK Patent Application GB 2188638A, filed on March 26, 1987; Winter US 5,225,539), the contents of which is expressly incorporated by reference.

Also provided are humanized antibodies in which specific amino acids have been substituted, deleted or added. Criteria for selecting amino acids from the donor are described in, *e.g.*, US 5,585,089, *e.g.*, columns 12-16 of US 5,585,089, the contents of which are hereby incorporated by reference. Other techniques for humanizing antibodies are described in Padlan *et al.* EP 519596 A1, published on December 23, 1992.

In an embodiment, the antibody molecule has a heavy chain constant region chosen from, *e.g.*, the heavy chain constant regions of IgG1, IgG2 (*e.g.*, IgG2a), IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE; particularly, chosen from, *e.g.*, the (*e.g.*, human) heavy chain constant regions of IgG1, IgG2, IgG3, and IgG4. In another embodiment, the antibody molecule has a light chain constant region chosen from, *e.g.*, the (*e.g.*, human) light chain constant regions of kappa or lambda. The constant region can be altered, *e.g.*, mutated, to modify the properties of the antibody molecule (*e.g.*, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, and/or complement function). In an embodiment, the antibody molecule has effector function and can fix complement. In another embodiment, the antibody molecule does not recruit effector cells or fix complement. In certain embodiments, the antibody molecule has reduced or no ability to bind an Fc receptor. For example, it may be an isotype or subtype, fragment or other mutant, which does not support binding to an Fc receptor, *e.g.*, it has a mutagenized or deleted Fc receptor binding region.

In an embodiment, a constant region of the antibody molecule is altered. Methods for altering an antibody constant region are known in the art. Antibody molecules with altered function, *e.g.*, altered affinity for an effector ligand, such as FcR on a cell, or the C1 component of complement can be produced by replacing at least one amino acid residue in the constant portion of the antibody with a different residue (see *e.g.*, EP 388,151 A1, U.S. Pat. No. 5,624,821 and U.S. Pat. No. 5,648,260, the contents of all of which are hereby incorporated by reference). Amino acid mutations which stabilize antibody structure, such as S228P (EU nomenclature, S241P in Kabat nomenclature) in human IgG4 are also contemplated. Similar type of alterations could be described which if applied to the murine, or other species immunoglobulin would reduce or eliminate these functions.

In an embodiment, the only amino acids in the antibody molecule are canonical amino acids. In an embodiment, the antibody molecule comprises naturally-occurring amino acids; analogs, derivatives and congeners thereof; amino acid analogs having variant side chains; and/or all stereoisomers of any of any of the foregoing. The antibody molecule may comprise the D- or L-optical isomers of amino acids and peptidomimetics.

A polypeptide of an antibody molecule described herein may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The antibody molecule may also be modified; for example, by disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling
5 component. The polypeptide can be isolated from natural sources, can be a produced by recombinant techniques from a eukaryotic or prokaryotic host, or can be a product of synthetic procedures.

The antibody molecule described herein can be used alone in unconjugated form, or can be bound to a substance, *e.g.*, a toxin or moiety (*e.g.*, a therapeutic drug; a compound emitting radiation; molecules of plant, fungal, or bacterial origin; or a biological protein (*e.g.*, a protein toxin) or particle
10 (*e.g.*, a recombinant viral particle, *e.g.*, via a viral coat protein). For example, the anti-APRIL antibody can be coupled to a radioactive isotope such as an α -, β -, or γ -emitter, or a β - and γ -emitter.

An antibody molecule can be derivatized or linked to another functional molecule (*e.g.*, another peptide or protein). As used herein, a “derivatized” antibody molecule is one that has been modified. Methods of derivatization include but are not limited to the addition of a fluorescent
15 moiety, a radionucleotide, a toxin, an enzyme or an affinity ligand such as biotin. Accordingly, the antibody molecules are intended to include derivatized and otherwise modified forms of the antibodies described herein, including immunoadhesion molecules. For example, an antibody molecule can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (*e.g.*, a bispecific
20 antibody or a diabody), a detectable agent, a toxin, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

Some types of derivatized antibody molecule are produced by crosslinking two or more antibodies (of the same type or of different types, *e.g.*, to create bispecific antibodies). Suitable
25 crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (*e.g.*, m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (*e.g.*, disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

Useful detectable agents with which an anti-dengue antibody molecule may be derivatized (or
30 labeled) to include fluorescent compounds, various enzymes, prosthetic groups, luminescent materials, bioluminescent materials, fluorescent emitting metal atoms, *e.g.*, europium (Eu), and other anthanides, and radioactive materials (described below). Exemplary fluorescent detectable agents include fluorescein, fluorescein isothiocyanate, rhodamine, 5dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. An antibody may also be derivatized with detectable enzymes,
35 such as alkaline phosphatase, horseradish peroxidase, β -galactosidase, acetylcholinesterase, glucose oxidase and the like. When an antibody is derivatized with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a detectable reaction product. For

example, when the detectable agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody molecule may also be derivatized with a prosthetic group (*e.g.*, streptavidin/biotin and avidin/biotin). For example, an antibody may be derivatized with biotin, and detected through indirect measurement of avidin or streptavidin binding. Examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of bioluminescent materials include luciferase, luciferin, and aequorin.

Labeled antibody molecules can be used, for example, diagnostically and/or experimentally in a number of contexts, including (i) to isolate a predetermined antigen by standard techniques, such as affinity chromatography or immunoprecipitation; (ii) to detect a predetermined antigen (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the protein; (iii) to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to determine the efficacy of a given treatment regimen.

An antibody molecule may be conjugated to another molecular entity, typically a label or a therapeutic (*e.g.*, antimicrobial (*e.g.*, antibacterial or bactericidal), immunomodulatory, immunostimulatory, cytotoxic, or cytostatic) agent or moiety. Radioactive isotopes can be used in diagnostic or therapeutic applications. Radioactive isotopes that can be coupled to the antibody molecules include, but are not limited to α -, β -, or γ -emitters, or β - and γ -emitters. Such radioactive isotopes include, but are not limited to iodine (^{131}I or ^{125}I), yttrium (^{90}Y), lutetium (^{177}Lu), actinium (^{225}Ac), praseodymium, astatine (^{211}At), rhenium (^{186}Re), bismuth (^{212}Bi or ^{213}Bi), indium (^{111}In), technetium ($^{99\text{m}}\text{Tc}$), phosphorus (^{32}P), rhodium (^{188}Rh), sulfur (^{35}S), carbon (^{14}C), tritium (^3H), chromium (^{51}Cr), chlorine (^{36}Cl), cobalt (^{57}Co or ^{58}Co), iron (^{59}Fe), selenium (^{75}Se), or gallium (^{67}Ga). Radioisotopes useful as therapeutic agents include yttrium (^{90}Y), lutetium (^{177}Lu), actinium (^{225}Ac), praseodymium, astatine (^{211}At), rhenium (^{186}Re), bismuth (^{212}Bi or ^{213}Bi), and rhodium (^{188}Rh). Radioisotopes useful as labels, *e.g.*, for use in diagnostics, include iodine (^{131}I or ^{125}I), indium (^{111}In), technetium ($^{99\text{m}}\text{Tc}$), phosphorus (^{32}P), carbon (^{14}C), and tritium (^3H), or one or more of the therapeutic isotopes listed above.

The present disclosure provides radiolabeled antibody molecules and methods of labeling the same. In an embodiment, a method of labeling an antibody molecule is disclosed. The method includes contacting an antibody molecule, with a chelating agent, to thereby produce a conjugated antibody. The conjugated antibody is radiolabeled with a radioisotope, *e.g.*, ^{111}In , ^{90}Y and ^{177}Lu , to thereby produce a labeled antibody molecule.

In some aspects, this disclosure provides a method of making an antibody molecule disclosed herein. The method includes: providing an antigen, *e.g.*, APRIL or a fragment thereof; obtaining an antibody molecule that specifically binds to the antigen; evaluating efficacy of the antibody molecule in modulating activity of the antigen and/or organism expressing the antigen, *e.g.*, APRIL. The

method can further include administering the antibody molecule, including a derivative thereof (*e.g.*, a humanized antibody molecule) to a subject, *e.g.*, a human.

This disclosure provides an isolated nucleic acid molecule encoding the above antibody molecule, vectors and host cells thereof. The nucleic acid molecule includes, but is not limited to,
5 RNA, genomic DNA and cDNA.

Amino acid and nucleotide sequences of exemplary antibody molecules are described in **Tables 1 and 2**, respectively. Amino acid sequences of additional exemplary humanized antibody molecules are described in **Table 5**.

10

Table 1. The amino acid sequences of the heavy chain variable region (VH) and light chain variable region (VL) of the exemplary anti-APRIL antibodies are provided as follows. CDRs, defined according to the Kabat system, are underlined and bold, while CDRs defined according to the Chothia system are italicized.

Antibody	Chain	Amino Acid Sequence	SEQ ID NO	Chothia CDR		SEQ ID NO	Kabat CDR		SEQ ID NO
2218	VH	DVQLQESGPGLVKPSQSLSLTCSVTGYSIT <u>SGYYW</u> <u>NWIRQFPGNKLEWMGYI</u> <u>SYDGYNNYNPSLKNRISI</u> TRDTSKNQFFLKLNSVTTEDTATYYCAN <u>YYDYEDW</u> <u>YFGVWGTGTTVTVSS</u>	9	HCDR1	GYSITSGY	1	HCDR1	SGYYWN	7
				HCDR2	SYDGY	2	HCDR2	YISYDGYNNYNPSLKN	8
				HCDR3	YYDYEDWYFGV	3	HCDR3	YYDYEDWYFGV	3
	VL	DIVLTQSPASLAMS LGKRATIS <u>RASESVSIIGTN</u> <u>SIHWYQQKPGQP</u> <u>PKLLIY</u> <u>HASNLET</u> GVPAREFSGSG SRTDFTLTIDPVEEDDAFYVC <u>LQSRKIPYT</u> FGGG TKLEIK	10	LCDR1	RASESVSIIGTNSIH	4	LCDR1	RASESVSIIGTNSIH	4
				LCDR2	HASNLET	5	LCDR2	HASNLET	5
				LCDR3	LQSRKIPYT	6	LCDR3	LQSRKIPYT	6
2419	VH	QVQLQQSGAELVKPGASVRLSCEASGYTFT <u>DYTIH</u> WVKQRSGQGLEWIG <u>WI</u> <u>YPLRGS</u> SINYNEKFKDKATL TADKSSSTVYLELGRITSKDSAVYFCAR <u>HGAYYSN</u> <u>AFDYWGQGTTLTVSS</u>	19	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINYNEKFKD	18
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	NIVMTQSPASLAVSLGQRATIS <u>RASESVDNDGIR</u> <u>FMHWYQQKPGQP</u> <u>PKLLIY</u> <u>RASNLES</u> GIPAREFSGSG SRTDFTLTINPVETDDVATYYC <u>QOSNKDPYT</u> FGGG TKLELK	20	LCDR1	RASESVDNDGIRFMH	14	LCDR1	RASESVDNDGIRFMH	14
				LCDR2	RASNLES	15	LCDR2	RASNLES	15
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1305	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>DYTIH</u> WVRQATGQGLEWM <u>WI</u> <u>YPLRGS</u> SINYAQKFQGRVTM TANKSISTVYMEISSLRSEDAVYFCAR <u>HGAYYSN</u> <u>AFDYWGQGTTLTVSS</u>	283	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINYAQKFQG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLS <u>RASESVDNDGIR</u> <u>FLHWYQQKPGQAPRLIY</u> <u>RASNRET</u> GIPAREFSGSG SGTEFTLTISLQSEDAVYVC <u>QOSNKDPYT</u> FGGG TKVEIK	284	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1306	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>DYTIH</u> WVRQATGQGLEWM <u>WI</u> <u>YPLRGS</u> SINYAQKFQGRVTM TANKSISTVYMEISSLRSEDAVYFCAR <u>HGAYYSN</u> <u>AFDYWGQGTTLTVSS</u>	283	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINYAQKFQG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLS <u>RASESVDNDGIR</u> <u>FLHWYQQKPGQAPRLIY</u> <u>RASTRAT</u> GIPAREFSGSG SRTEFTLTISLQSEDAVYVC <u>QOSNKDPYT</u> FGGG TKVEIK	286	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASTRAT	285	LCDR2	RASTRAT	285
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1310	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>DYTIH</u> WVRQATGQGLEWM <u>WI</u> <u>YPLRGS</u> SINYAQKFQGRVTM TANKSISTVYMEISSLRSEDAVYFCAR <u>HGAYYSN</u>	283	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINYAQKFQG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13

		<u>AFDYWGQGLTVTVSS</u>							
	VL	DIVMTQSPDSLAVSLGERATINC <u>KSSQSVNDNGIR</u> <u>FLHWYQQKPGQPPKLLIYRASTR</u> ESGVDPDRFSGSG SGIDFTLTIISSLQAEDVAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	316	LCDR1	KSSQSVNDNGIRFLH	314	LCDR1	KSSQSVNDNGIRFLH	314
				LCDR2	RASTR	315	LCDR2	RASTR	315
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0806	VH	EVQLVQSGAEVKKPGESLKISCKASGYTFTDYTIH WVRQMPGKGLEWMGWIYPLRGSSINYS PSFQ QVTTI SADKSISTVYLQWSSLKASDIAM YFCAR HGAYYSN <u>AFDYWGQGLTVTVSS</u>	288	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSSINYS PSFQ	287
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSC <u>RASESVNDNGIR</u> <u>FLHWYQQKPGQAPRLLIYRASTR</u> ATGIPARFSGSG SRTEFTLTIISSLQSEDFAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	286	LCDR1	RASESVNDNGIRFLH	280	LCDR1	RASESVNDNGIRFLH	280
				LCDR2	RASTRAT	285	LCDR2	RASTRAT	285
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0205	VH	QVQLVQSGAEVKKPGSSVKVSKASGYTFTDYTIH WVRQAPGQGLEWMGWIYPLRGSSIN YAEK FQGRVTI TADKSTSTAYMELSSLRSEDTAVYFCARHGAYYSN <u>AFDYWGQGLTVTVSS</u>	289	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSSIN YAEK FQGRVTI	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSC <u>RASESVNDNGIR</u> <u>FLHWYQQKPGQAPRLLIYRASNRET</u> GIPARFSGSG SGTEFTLTIISSLQSEDFAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	284	LCDR1	RASESVNDNGIRFLH	280	LCDR1	RASESVNDNGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0406	VH	QVQLVQSGAEVKKPGSSVKVSKASGYTFTDYTIH WVRQAPGQGLEWMGWIYPLRGSSIN YAEK FQGRVTL TADKSTSTVYMELSSLRSEDTAVYFCARHGAYYSN <u>AFDYWGQGLTVTVSS</u>	291	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSSIN YAEK FQGRVTL	290
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSC <u>RASESVNDNGIR</u> <u>FLHWYQQKPGQAPRLLIYRASTR</u> ATGIPARFSGSG SRTEFTLTIISSLQSEDFAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	286	LCDR1	RASESVNDNGIRFLH	280	LCDR1	RASESVNDNGIRFLH	280
				LCDR2	RASTRAT	285	LCDR2	RASTRAT	285
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0605	VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTDYTIH WVRQAPGQGLEWMGWIYPLRGSSIN YAEK FQGRVTL TADKSTSTVYMELSSLRSEDTAVYFCARHGAYYSN <u>AFDYWGQGLTVTVSS</u>	317	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSSIN YAEK FQGRVTL	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSC <u>RASESVNDNGIR</u> <u>FLHWYQQKPGQAPRLLIYRASNRET</u> GIPARFSGSG SGTEFTLTIISSLQSEDFAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	284	LCDR1	RASESVNDNGIRFLH	280	LCDR1	RASESVNDNGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0805	VH	EVQLVQSGAEVKKPGESLKISCKASGYTFTDYTIH WVRQMPGKGLEWMGWIYPLRGSSINYS PSFQ QVTTI SADKSISTVYLQWSSLKASDIAM YFCAR HGAYYSN <u>AFDYWGQGLTVTVSS</u>	288	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSSINYS PSFQ	287
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSC <u>RASESVNDNGIR</u> <u>FLHWYQQKPGQAPRLLIYRASNRET</u> GIPARFSGSG SGTEFTLTIISSLQSEDFAVYYC <u>QOSNKDPYTFGGG</u> TKVEIK	284	LCDR1	RASESVNDNGIRFLH	280	LCDR1	RASESVNDNGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16

2419-0105	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYTIH WVRQAPGQGLEWMGWIYPLRGSINIAQKFQGRVTM TADKSISTVYMELSLRLSDDIAVYYCARHGAYYSN AFDYWGQGTILVTVSS	292	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSCRAESVDNDGIR FLHWYQQKPGQAPRLLIYRASNRETGIPARFSGSG SGTEFTLTITISLQSEDAVYYCQOSNKDPYTFGGG TKVEIK	284	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1204	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYTIH WVRQATGQGLEWMGWIYPLRGSINIAQKFQGRVTM TANKSSSTVYMELSSLRSEDAVYFCARHGAYYSN AFDYWGQGTILVTVSS	294	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSCRAESVDNDGIR FLHWYQQKPGQAPRLLIYRASTLETGIPARFSGSG SGTEFTLTITISLQSEDAVYYCQOSNKDPYTFGGG TKVEIK	295	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASTLET	293	LCDR2	RASTLET	293
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1205	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYTIH WVRQATGQGLEWMGWIYPLRGSINIAQKFQGRVTM TANKSSSTVYMELSSLRSEDAVYFCARHGAYYSN AFDYWGQGTILVTVSS	294	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSCRAESVDNDGIR FLHWYQQKPGQAPRLLIYRASNRETGIPARFSGSG SGTEFTLTITISLQSEDAVYYCQOSNKDPYTFGGG TKVEIK	284	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASNRET	281	LCDR2	RASNRET	281
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1210	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYTIH WVRQATGQGLEWMGWIYPLRGSINIAQKFQGRVTM TANKSSSTVYMELSSLRSEDAVYFCARHGAYYSN AFDYWGQGTILVTVSS	294	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	DIVMTQSPDSLAVSLGERATINCKSSQSVNDGIR FLHWYQQKPGQPPKLLIYRASTRESGVPDRFSGSG SGTDFTLTITISLQAEDVAVYYCQOSNKDPYTFGGG TKVEIK	316	LCDR1	KSSQSVNDGIRFLH	314	LCDR1	KSSQSVNDGIRFLH	314
				LCDR2	RASTRES	315	LCDR2	RASTRES	315
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-1406	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYTIH WVRQATGQGLEWMGWIYPLRGSINIAQKFQGRVTM TADKSISTVYMELSSLRSEDAVYFCARHGAYYSN AFDYWGQGTILVTVSS	296	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSCRAESVDNDGIR FLHWYQQKPGQAPRLLIYRASTRATGIPARFSGSG SRTEFTLTITISLQSEDAVYYCQOSNKDPYTFGGG TKVEIK	286	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280
				LCDR2	RASTRAT	285	LCDR2	RASTRAT	285
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2419-0206	VH	QVQLVQSGAEVKKPGSSVKVSCKASGYTFTDYTIH WVRQAPGQGLEWMGWIYPLRGSINIAQKFQGRVTI TADKSTSTAYMELSSLRSEDAVYFCARHGAYYSN AFDYWGQGTILVTVSS	289	HCDR1	GYTFTDY	11	HCDR1	DYTIH	17
				HCDR2	YPLRGS	12	HCDR2	WIYPLRGSINIAQKFQGG	282
				HCDR3	HGAYYSNAFDY	13	HCDR3	HGAYYSNAFDY	13
	VL	EIVMTQSPATLSVSPGERATLSCRAESVDNDGIR	286	LCDR1	RASESVDNDGIRFLH	280	LCDR1	RASESVDNDGIRFLH	280

		<u>FLH</u> WYQQKPGQAPRLLIY <u>RASTRAT</u> GIPARFSGSG SRTEFTLTISLQSEDFAVYYC <u>QOSNKDPYT</u> FGGG TKVEIK		LCDR2	RASTRAT	285	LCDR2	RASTRAT	285
				LCDR3	QOSNKDPYT	16	LCDR3	QOSNKDPYT	16
2621	VH	EVQLQQSGAELVRPGSSVKMSCKTSGYFTT <u>SYGIN</u> WVKQRPQGGLIEWIGYIYIGNGYAEYNERFKGKATL TSDTSSSTAYMQLSSLTSEDSAIYFCALYYPWFTY WGQGTILVTVSA	29	HCDR1	GYFTTSY	21	HCDR1	SYGIN	27
				HCDR2	YIGNGY	22	HCDR2	YIYIGNGYAEYNERFKG	28
				HCDR3	YYPWFTY	23	HCDR3	YYPWFTY	23
	VL	DIQMTQSPASISASVGDSTITC <u>RASENIYSYLA</u> YQQKQGKSPQLLVYNAKTLAEGVPSRFSGSGSGTQ FSLKINSLQPEDFGNYCQHHDTPFTFGGGTKLE IK	30	LCDR1	RASENIYSYLA	24	LCDR1	RASENIYSYLA	24
				LCDR2	NAKTLAE	25	LCDR2	NAKTLAE	25
				LCDR3	QHHDTPFT	26	LCDR3	QHHDTPFT	26
2922	VH	QVQLHQSGPELVKPGASVKLSCKTSGYFTT <u>SYDVF</u> WVKQRPQGGLIEWIGWIYPRDSSTKYNEKFKGKATL TVDTSSSTAYMELHSLTSEDSAVYFCAKEGYDYDK RGFDYWGQGTTLTVSS	39	HCDR1	GYFTTSY	21	HCDR1	SYDVF	37
				HCDR2	YPRDSS	32	HCDR2	WIYPRDSSTKYNEKFKG	38
				HCDR3	EGYDYDKRGFDY	33	HCDR3	EGYDYDKRGFDY	33
	VL	DIVLTQSPASISAVSLGQRATISCKASQSVSFAGTN <u>LMH</u> WYQQRPQGQPKLLIY <u>RASNLEP</u> GVPTRFSGSG SRTDFTLNHPVEEDDAATYYCQOSREYPWTFGGG TKLEIK	40	LCDR1	KASQSVSFAGTNLMH	34	LCDR1	KASQSVSFAGTNLMH	34
				LCDR2	RASNLEP	35	LCDR2	RASNLEP	35
				LCDR3	QOSREYPWT	36	LCDR3	QOSREYPWT	36
3125	VH	QVQLQQSGAELVRPGASVTLSCASGYFTT <u>DYEMH</u> WVKQTPVHGLEWIGAIIDPETGGTAYNQRFKGKATL TTDKSSITAYMELRSLTSEDSAVYYCTR <u>WNDGDY</u> WGQGTTLTVSS	49	HCDR1	GYFTTDY	11	HCDR1	DYEMH	47
				HCDR2	DPETGG	42	HCDR2	AIDPETGGTAYNQRFKG	48
				HCDR3	WNDGDY	43	HCDR3	WNDGDY	43
	VL	DVVMVTQTPLSLSVTIGQPASISCKSSQSLLYSNGK <u>TYLN</u> WFQQRPQGSPKRLMYQVSKLDPGIPDRFSGS GSETDFTLKISRVEAEDLGLYYC <u>LQGTYYPYT</u> FGG GTKLEIK	50	LCDR1	KSSQSLLYSNGKTYLN	44	LCDR1	KSSQSLLYSNGKTYLN	44
				LCDR2	QVSKLDP	45	LCDR2	QVSKLDP	45
				LCDR3	LQGTYYPYT	46	LCDR3	LQGTYYPYT	46
3327	VH	EVQLQQSGPELVKPGASVKMSCKASGYFTT <u>GYFMN</u> WVKQSHGKSLEWIGRINPYNGDTFYNQKFKGKATL TVDKSSSTAHMELRSLTSEDSALYYCASEGDGYW YFDVWGAGTIVTVSS	59	HCDR1	GYSFTGY	51	HCDR1	GYFMN	57
				HCDR2	NPYNGD	52	HCDR2	RINPYNGDTFYNQKFKG	58
				HCDR3	EGDGYWYFDV	53	HCDR3	EGDGYWYFDV	53
	VL	DIVLTQSPASISAVSLGQRATISCKASEVDNYGIS <u>FMN</u> WFQQKPGQPPKLLIYAASNQGSVPARFSGSG SGTDFSLNIHPMEEDDTAMYFCQOSKEVPRTFGGG TKLEIK	60	LCDR1	RASEVDNYGISFMN	54	LCDR1	RASEVDNYGISFMN	54
				LCDR2	AASNQGS	55	LCDR2	AASNQGS	55
				LCDR3	QOSKEVPRT	56	LCDR3	QOSKEVPRT	56
3525	VH	QVQLQQSGAELVRPGASVKLSCKASGYFTT <u>DHEMH</u> WVRQTPVHGLEWIGVIDPDTGDTTYNQKFKGKATL TADKSSSTAYMDLRSLTSEDSAVFYCTR <u>WTGGDY</u> GHGTTTLTVSS	66	HCDR1	GYFTTDH	61	HCDR1	DHEMH	64
				HCDR2	DPDTGD	62	HCDR2	VIDPDTGDTTYNQKFKG	65
				HCDR3	WTGGDY	63	HCDR3	WTGGDY	63
	VL	DVVMVTQTPLSLSVTIGQPASISCKSSQSLLYSNGK <u>TYLN</u> WFQQRPQGSPKRLMYQVSKLDPGIPDRFSGS GSETDFTLKISRVEAEDLGLYYC <u>LQGTYYPYT</u> FGG GTKLEIK	50	LCDR1	KSSQSLLYSNGKTYLN	44	LCDR1	KSSQSLLYSNGKTYLN	44
				LCDR2	QVSKLDP	45	LCDR2	QVSKLDP	45
				LCDR3	LQGTYYPYT	46	LCDR3	LQGTYYPYT	46
3530	VH	QVQLQQSGAELVRPGASVKLSCKASGYFTT <u>DHEMH</u>	66	HCDR1	GYFTTDH	61	HCDR1	DHEMH	64

		WVRQTPVHGLEWIGVIDPDTGDTTYNQKFKGKATL TADKSSSTAYMDLRSLTSEDSAVFYCTRWTGGDYW GHGTTLTVSS		HCDR2	DPDTGD	62	HCDR2	VIDPDTGDTTYNQKFKG	65
				HCDR3	WTGGDY	63	HCDR3	WTGGDY	63
	VL	DAVMTQTPLSLSVTIGQPASISCKSSQSLLYSDGK TYLNWFQQRPGQSPKRLMYQVSKLDPGIPDRFSGS GSETDFTLKISRVEAEDLGVYYCLOGTYYPYTFGS GTKLEIK		LCDR1	KSSQSLLYSDGKTYLN	67	LCDR1	KSSQSLLYSDGKTYLN	67
	4035	VH	70	LCDR2	QVSKLDP	45	LCDR2	QVSKLDP	45
				LCDR3	LQGTYYPYT	46	LCDR3	LQGTYYPYT	46
		VH	101	HCDR1	GFSLTIY	93	HCDR1	IYDVH	99
				HCDR2	WSDGS	94	HCDR2	VIWSDGSTDYNAAFIS	100
				HCDR3	NWVDQAWFAY	95	HCDR3	NWVDQAWFAY	95
		VL	102	LCDR1	RASKNIYSYLA	96	LCDR1	RASKNIYSYLA	96
				LCDR2	NAKTLPE	97	LCDR2	NAKTLPE	97
				LCDR3	QHHYGTPLT	98	LCDR3	QHHYGTPLT	98
	4035-062	VH	225	HCDR1	GFSLTIY	93	HCDR1	IYDVH	99
				HCDR2	WSDGS	94	HCDR2	VIWSDGSTDYNPSLKS	273
				HCDR3	NWVDQAWFAY	95	HCDR3	NWVDQAWFAY	95
		VL	229	LCDR1	RASKNIYSYLA	96	LCDR1	RASKNIYSYLA	96
				LCDR2	NAKTLPE	97	LCDR2	NAKTLPE	97
				LCDR3	QHHYGTPLT	98	LCDR3	QHHYGTPLT	98
	3934	VH	111	HCDR1	GYIFTDY	103	HCDR1	DY TIN	109
				HCDR2	YPGSGN	104	HCDR2	WIYPGSGNRKYNDKFKG	110
				HCDR3	ESNYVGYAMDY	105	HCDR3	ESNYVGYAMDY	105
		VL	112	LCDR1	RSSQSVVNSNGN	106	LCDR1	RSSQSVVNSNGNTYLE	106
				LCDR2	KVSNRFS	107	LCDR2	KVSNRFS	107
				LCDR3	FQGSHPWT	108	LCDR3	FQGSHPWT	108
	3833	VH	121	HCDR1	GYFTNY	113	HCDR1	NYWIG	119
				HCDR2	YPGGIGGGY	114	HCDR2	DIYPGGIGGGYTRYNEK FKG	120
				HCDR3	SETGRAMDY	115	HCDR3	SETGRAMDY	115
		VL	122	LCDR1	KASQDINKYIA	116	LCDR1	KASQDINKYIA	116
				LCDR2	YTSTLKP	117	LCDR2	YTSTLKP	117
				LCDR3	LQYDNLNT	118	LCDR3	LQYDNLNT	118
	3631	VH	131	HCDR1	GYSFTDY	123	HCDR1	DYNIY	129
				HCDR2	DPSNGG	124	HCDR2	YIDPSNGGPGYNQKFRG	130
				HCDR3	RDNYGSGTMDY	125	HCDR3	RDNYGSGTMDY	125
		VL	132	LCDR1	KASQNVGTDVS	126	LCDR1	KASQNVGTDVS	126

		YQQKPGKSPKPLIY <u>WASNRFT</u> GVPDRFIGSGSGTD FTLTISNVQSEDLADYFC <u>EQYSIYPLT</u> FGAGTKLE LK		LCDR2	WASNRFT	127	LCDR2	WASNRFT	127
				LCDR3	EQYSIYPLT	128	LCDR3	EQYSIYPLT	128
3732	VH	EIQLQQSGPELVKPGASVKVSCASGYST <u>DDNMY</u> WVKQSHGKSLEWIG <u>YIDPLNGGT</u> GYNOQKFKGKATL TVDKSSSTAFLHLSLTSEDSAVYYCAR <u>RDNYATG</u> <u>TMDYWGQTSVTVSS</u>	140	HCDR1	GYSTDD	133	HCDR1	DDNMY	138
				HCDR2	DPLNGG	134	HCDR2	YIDPLNGGTGYNOQKFKG	139
				HCDR3	RDNYATGTMDY	135	HCDR3	RDNYATGTMDY	135
	VL	DIVMTQSQKFMSTSVGDRVSITC <u>KASKNVGTDVSW</u> YQQKPGKSPKPLIY <u>WASNRFT</u> GVPDRFTGSGSGTD FTLTINNVSQSEDLADYFC <u>EQYSSYPLT</u> FGAGTKLE LK	141	LCDR1	KASKNVGTDVS	136	LCDR1	KASKNVGTDVS	136
				LCDR2	WASNRFT	127	LCDR2	WASNRFT	127
				LCDR3	EQYSSYPLT	137	LCDR3	EQYSSYPLT	137
4338	VH	EVQLQQSGPELVKPGASVKISCKASGYTFT <u>DYNMD</u> WVKQSHGKSLEWIG <u>NIYPINGYT</u> GYNQRFKNKATL TVDKSSSTAYMELHLSLTSEDSAVYYCAR <u>DSNYVGW</u> <u>YFDVWGAGTIVTVSS</u>	151	HCDR1	GYTFTDY	11	HCDR1	DYNMD	149
				HCDR2	YPINGY	142	HCDR2	NIYPINGYTGYNQRFKN	150
				HCDR3	DSNYVGWYFDV	143	HCDR3	DSNYVGWYFDV	143
	VL	DVVMTQTPLSLPVSLGDAQSISCR <u>RSSQSLVHSNGN</u> <u>TYLHWYLQKPGQSPKLLIY</u> <u>KVSNRFS</u> GVPDRFSGS SGSDFTFKISRVEAEDLGVIYFC <u>SQSTHVPRT</u> FGG GTKLEIK	152	LCDR1	RSSQSLVHSNGNTYLH	144	LCDR1	RSSQSLVHSNGNTYLH	144
				LCDR2	KVSNRFS	107	LCDR2	KVSNRFS	107
				LCDR3	SQSTHVPRT	145	LCDR3	SQSTHVPRT	145
	VL	DIVLTQSPASLTVSLGQRATFSC <u>RASKSVSTSGYS</u> <u>YMHYQQKPGQPPKLLIY</u> <u>FTSDLEP</u> GVPARFTGSG SGDFTLNHPVEEEDAATYYC <u>QHSRELPYP</u> FGGG TRLEIK	153	LCDR1	RASKSVSTSGYSYM	146	LCDR1	RASKSVSTSGYSYM	146
				LCDR2	FTSDLEP	147	LCDR2	FTSDLEP	147
				LCDR3	QHSRELPYP	148	LCDR3	QHSRELPYP	148
4540	VH	QVQLVQSGPELVKPGASVKISCKASGYTF <u>DYYIN</u> WVKQRPQGQLEWIG <u>WIFPGSGSTYYNEK</u> FKGKATL TVDKSSSTAYMLLSLTSEDSAVYFCAR <u>GDSGRAM</u> <u>DYWGQGTIVTVSS</u>	161	HCDR1	GYTFADY	154	HCDR1	DYYIN	159
				HCDR2	FPGSGS	155	HCDR2	WIFPGSGSTYYNEKFKG	160
				HCDR3	GDSGRAMDY	156	HCDR3	GDSGRAMDY	156
	VL	DIQMTQSPSSLSASLGKVTITC <u>KASQDINKYIAW</u> YQHKGPKGPRLLIH <u>YTSTLQS</u> GIPSRFSGSGSGRD YSFISISNLEPEDNATYYC <u>LQYDNLLT</u> FGAGTKLEL K	162	LCDR1	KASQDINKYIA	116	LCDR1	KASQDINKYIA	116
				LCDR2	YTSTLQS	157	LCDR2	YTSTLQS	157
				LCDR3	LQYDNLLT	158	LCDR3	LQYDNLLT	158
4540-063	VH	QVQLVQSGAELKKPGASVKVSCASGYTF <u>DYYMN</u> WVRQAPGQGLEWMG <u>WIFPGSGSTYYNQK</u> FQGRVTM TVDKSSSTAYMELSRRLSDDTAVYYCAR <u>GDSGRAM</u> <u>DYWGQGTIVTVSS</u>	258	HCDR1	GYTFADY	154	HCDR1	DYYMN	276
				HCDR2	FPGSGS	155	HCDR2	WIFPGSGSTYYNQKFG	277
				HCDR3	GDSGRAMDY	156	HCDR3	GDSGRAMDY	156
	VL	DIQMTQSPSSLSASVGRVTITC <u>QASQDINKYLAW</u> YQHKGPKAPKLLIH <u>YTSTLET</u> GVPSRFSGSGSGTD FTFTISSLPEDIATYYC <u>LQYDNLLT</u> FGGGTKVEI K	261	LCDR1	QASQDINKYLA	274	LCDR1	QASQDINKYLA	274
				LCDR2	YTSTLET	275	LCDR2	YTSTLET	275
				LCDR3	LQYDNLLT	158	LCDR3	LQYDNLLT	158
4540-033	VH	QVQLVQSGAEVKKPGASVKVSCASGYTF <u>DYYIN</u> WVRQAPGQGLEWMG <u>WIFPGSGSTYYAQK</u> LQGRVTM TTDTSTAYMELRSLRSDDTAVYYCAR <u>GDSGRAM</u> <u>DYWGQGTIVTVSS</u>	256	HCDR1	GYTFADY	154	HCDR1	DYYIN	159
				HCDR2	FPGSGS	155	HCDR2	WIFPGSGSTYYAQKLQ	278
				HCDR3	GDSGRAMDY	156	HCDR3	GDSGRAMDY	156
	VL	DIQMTQSPSSLSASVGRVTITC <u>QASQDINKYLAW</u>	261	LCDR1	QASQDINKYLA	274	LCDR1	QASQDINKYLA	274

		YQHKPGKAPKLLIH YTSTLET GVPSRFSGSGSGTD FTFTISSLQPEDIAITYYCLQYDNLLTFGGGKVEI K		LCDR2	YTSTLET	275	LCDR2	YTSTLET	275
				LCDR3	LQYDNLLT	158	LCDR3	LQYDNLLT	158
4237	VH	QAHLKESGPGLVAPSQSL SITCTVSGFSLTDYDVH WVRQSPGKGLEWLGVIWWDGSTDYNTAFISRLTIS KDNSKRSQVFFKMNSLQADDTAIYYCARNWYGGYWF AYWGQGTILVTVA	171	HCDR1	GFSLTDY	163	HCDR1	DYDVH	169
				HCDR2	WWDGS	164	HCDR2	VIWWDGSTDYNTAFIS	170
				HCDR3	NWYGGYWFAY	165	HCDR3	NWYGGYWFAY	165
	VL	DIQMTQSPASLSASAGETVTITCRSSENIYSYLA YQQKQKSPQLLVYANALAEGVPSRFSGSGSVTQ FSLKINSLQPEDFGSYYCQHGYGTPFTFGSGTKLE IK	172	LCDR1	RSSENIYSYLA	166	LCDR1	RSSENIYSYLA	166
				LCDR2	NANALAE	167	LCDR2	NANALAE	167
				LCDR3	QHRYGTPFT	168	LCDR3	QHRYGTPFT	168
4439	VH	EIQLQQSGAELVKPGASVKISCKASDYSFTGYNMN WVMQSHGKSLEWIGNIHPYYGGTSFNQKFMGKATL TADKSSSTAYMQLNSLTSEDSAVYYCARERSNFHA LDYWGQGTISVTSS	271	HCDR1	DYSFTGY	266	HCDR1	GYNMN	269
				HCDR2	HPYYGG	267	HCDR2	NIHPYYGGTSFNQKFMG	270
				HCDR3	ERSNFHALDY	268	HCDR3	ERSNFHALDY	268
	VL	DIVLTQSPASLTVSLGQRATFSCRASKSVSTSGYS YMHYQQKPGQPPKLLIYFTSDLEPGVPAREFTGSG SGTDFTLNIHPVEEEDAATYYCQHSRELPYPFGGG TKLEIK	272	LCDR1	RASKSVSTSGYSYM	146	LCDR1	RASKSVSTSGYSYM	146
				LCDR2	FTSDLEP	147	LCDR2	FTSDLEP	147
				LCDR3	QHSRELPYP	148	LCDR3	QHSRELPYP	148

Table 2. Nucleotide sequences of heavy chain variable regions (VHs) and light chain variable regions (VLs) of exemplary antibody molecules

Antibody	Chain	Nucleotide Sequence	SEQ ID NO
2218	VH	GATGTACAGCCTTCAGGAGTCAGGACCTGGCCTCGTGAAACCTTCTCAGTCTCTGTCTCTCACCTGCTCTGTCACTGGC TACTCCATCACCAGTGGTTATTACTGGAACCTGGATCCGGCAGTTTTCCAGGAAACAAACTGGAATGGATGGGCTACATA AGCTACGATGGTTACAATAACTACAACCCATCTCTCAAAAATCGAATCTCCATCACTCGTGACACATCTAAGAACCAG TTTTTCCTGAAGTTGAATTCGTGTGACTACTGAGGACACAGCCACATATTACTGTGCAAACCTACTATGATTACGAAGAC TGGTACTTCGGTGTCTGGGGCACAGGGACCAACGGTCACCGTCTCCTCA	71
	VL	GACATTGTGCTGACCCAAATCTCCAGCTTCTTTGGCTATGTCTCTAGGGAAGAGGGGCCACCATCTCCTGCAGAGCCAGC GAAAGTGTCACTATTATTGGTACTAATTCAAATACACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTCCTCATC TATCATGCATCCAAACCTAGAAACTGGAGTCCCTGCCAGGTTCACTGGCAGTGGGTCTAGAACAGACTTCACCCCTCACC ATTGATCCTGTGGAGGAAGATGATGTTGCAATCTATTACTGTCTGCAAAGTAGGAAGATTCCGTACACGTTCCGGAGGG GGGACCAAGCTGGAAATAAAA	72
2419	VH	CAGGTCCAGCTGCAGCAGTCTGGAGCTGAGCTGGTGAAACCCGGGGCATCAGTGAGGCTGTCTCTGCGAGGCTTCTGGC TACACCTTCACGGACTATACTATACTGGGTAAAGCAGAGGTCTGGACAGGGTCTTGAGTGGATTGGATGGATTAC CCTCTAAGAGGTAGTATAAACTACAATGAGAAATTCAGGACAAGGCCACATTGACTGCGGACAAATCCTCCAGCACA GTCTATTTGGAGCTTGGTAGATTGACATCTAAGGACTCTGCGGTCTATTTCTGTGCAAGACACGGAGCCTACTATAGT AACGCCCTTGACTACTGGGGCCAAGGCACCACTCTCAGTCTCCTCA	73
	VL	AACATTGTAATGACCCAATCTCCAGCTTCATTGGCTGTGTCTCTAGGTCAGAGGGCCACCATCTCCTGCAGAGCCAGC GAGAGTGTGATAATGATGGCATTAGATTTATGCACTGCTACCAGCAGAAACCAGGACAGCCACCCAAACTCCTCATC TATCGTGCATCCAAACCTAGAATCTGGGATCCCTGCCAGGTTCACTGGCAGTGGGTCTAGGACAGACTTCACCCCTCACT ATTAATCCTGTGGAGACTGATGATGTTGCAACCTATTACTGTGCAAGTAATAAGGATCCGTACACGTTCCGGAGGG GGGACCAAGCTGGAGCTGAAA	74
2419-1305	VH	CAAGTTCAGTTGGTGCAAAAGCGGGGCAGAAGTGAAGAAAACCTGGTGCTTCTGTGAAAAGTTTCTCTGCAAGGCCAGCGGC TACACCTTTACTGATTACACAATACACTGGGTACGGCAGGCAACTGGGCAAGGATTGGAATGGATGGGGTGGATATAC CCATTGCGAGGGTCTATAAACTACGCACAGAAATTTCAAGGTGCGAGTAACAATGACAGCCAACAAATCAATAAGCACC GTTTATATGGAACCTCTCATCTCTCAGGAGTGAGGATACCGCCGTGTATTTCTGCGCACGACACGGTGCATATTACTCA AACGCTTTTCGACTATTGGGGCCAGGGCACCCCTTGTGACTGTTAGTAGC	304
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAAGCCAGGGGAGCGAGCAACCCCTCAGTTGCAGAGCATCT GAGAGTGTGATAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCCCTGCACGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTCGCTGTGTATTACTGTGCAATCCAACAAGGACCCATACACTTTCCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-1306	VH	CAAGTTCAGTTGGTGCAAAAGCGGGGCAGAAGTGAAGAAAACCTGGTGCTTCTGTGAAAAGTTTCTCTGCAAGGCCAGCGGC TACACCTTTACTGATTACACAATACACTGGGTACGGCAGGCAACTGGGCAAGGATTGGAATGGATGGGGTGGATATAC CCATTGCGAGGGTCTATAAACTACGCACAGAAATTTCAAGGTGCGAGTAACAATGACAGCCAACAAATCAATAAGCACC GTTTATATGGAACCTCTCATCTCTCAGGAGTGAGGATACCGCCGTGTATTTCTGCGCACGACACGGTGCATATTACTCA AACGCTTTTCGACTATTGGGGCCAGGGCACCCCTTGTGACTGTTAGTAGC	304
	VL	GAGATAGTTATGACTCAGTCTCCCGCCACACTTTCAGTAAGTCCCGGTGAACGCGCCACCCCTGTCTGCGGTGCTTCC GAATCAGTGGATAATGACGGCATTAGGTTTTTGTCACTGGTACCAACAAAAAGCCCGGACAGGCCCCCGCCTGCTGATA	306

		TATCGTGCATCAACACGAGCAACAGGGATCCCCGCTCGATTTAGTGGATCCGGAAGCAGGACCGAATTTACACTTACC ATTTCCCTCACTTCAGTCAGAAGATTTTCGCCGTTTACTACTGTGTCAGCAGTCAAATAAGGATCCTTACACATTTGGGGGC GGTACAAAAGTCGAGATCAAA	
2419-1310	VH	CAAGTTCAGTTGGTGC AAAAGCGGGGCAGAAGTGAAGAAAACCTGGTGCTTCTGTGAAAAGTTTCCGTGCAAGGCCAGCGGC TACACCTTTACTGATTACACAATAACACTGGGTACGGCAGGCAACTGGGCAAGGATTGGAATGGATGGGGTGGATATAC CCATTGCGAGGGTCTATAAACTACGCACAGAAAATTTCAAGGTGCGAGTAACAATGACAGCCAACAAAATCAATAAGCACC GTTTATATGGAACCTCATCTCTCAGGAGTGAGGATACCGCCGTGTATTTCTGCGCACGACACGGTGCATATTACTCA AACGCTTTTCGACTATTGGGGCCAGGGCACCCCTTGTGACTGTTAGTAGC	304
	VL	GACATTGTAATGACCCAGTCTCCCGATAGCCTCGCTGTCTCACTCGGAGAACGCGCAACCATCAAACTGCAAGTCCCTCC CAAAGCGTTGACAATGACGGCATTAGGTTTTTGCAGTGGTACCAGCAGAAAACCCGGTCAACCTCCTAAGTTGCTCATT TACCGAGCATCTACCCGCGAGTCAGGAGTACCTGATCGCTTTTCCGGTAGCGGTAGTGAACAGATTTTACTCTGACC ATTAGTTCAGTCCAGGCAGAAGATGTGGCTGTCTACTACTGCCAACAGTCAAATAAAGACCCTTATACCTTCGGTGGG GGTACCAAAGTAGAGATCAAA	318
2419-0806	VH	GAGGTCCAGTTGGTCCAGTCAGGAGCCGAAGTCAAGAAGCCTGGGGAAAAGCCTGAAAATAAGTTGCAAAGCTAGTGGA TATACATTTACAGATTATACCATTCATTGGGTCCGGCAAATGCCAGGAAAAGGCTTGGAGTGGATGGGGTGGATTTAT CCCCCTCCGAGGCTCAATAAATTATAGTCCTAGTTTTTCAGGGGCAGGTAACCTATTAGCGCTGATAAAAGTATTTCTACA GTTTATTTGCAGTGGAGTTCATTGAAGGCTAGTGACACCGCTATGTATTTCTGCGCTAGACATGGTGCATATTATTCA AATGCCCTTCGACTATTGGGGCCAGGGCACCCCTCGTCACTGTGAGTTCC	307
	VL	GAGATAGTTATGACTCAGTCTCCCGCCACACTTTTCAGTAAGTCCCGGTGAACGCGCCACCCTGTCTGCGGTGCTTCC GAATCAGTGGATAATGACGGCATTAGGTTTTTGCAGTGGTACCAACAAAAAGCCCGGACAGGCCCCCCCGCCTGCTGATA TATCGTGCATCAACACGAGCAACAGGGATCCCCGCTCGATTTAGTGGATCCGGAAGCAGGACCGAATTTACACTTACC ATTTCCCTCACTTCAGTCAGAAGATTTTCGCCGTTTACTACTGTGTCAGCAGTCAAATAAGGATCCTTACACATTTGGGGGC GGTACAAAAGTCGAGATCAAA	306
2419-0205	VH	CAGGTGCAACTTGTTCAGTCAGGGGCTGAAGTAAAGAAGCCAGGCTCATCAGTCAAGGTATCATGCAAAGCATCTGGC TATACATTTACAGATTACACCATTCAGTGGGTGAGGCAAGCTCCCGGTCAAGGTCTCGAGTGGATGGGGTGGATATAC CCTCTCAGAGGCTCTATAAATTACGCTCAGAAAATTTCAAGGGAGAGTTACAATTACTGCTGATAAAAGTACCAGCACT GCTTATATGGAGCTTTCCCTCACTTCGTTTCAAGGACACCGCCGTTTACTTTTGTGCCCGGCATGGTGCCTATTATTCA AATGCCCTTCGATTATTGGGGGCAGGGAACCTTGGTCAAGTTTCATCT	308
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAAGCCCAGGGGAGCGAGCAACCCCTAGTTGCAGAGCATCT GAGAGTGTTGATAAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCTGTCACGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTCGCTGTGTATTACTGTGTCAGCAATCCAACAAGGACCCATACACTTTTCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-0406	VH	CAAGTTCAACTTGTCCAAAAGTGGGGCTGAAGTTAAAAAACCTGGATCATCAGTCAAGGTTTTCATGCAAAAGCCAGCGGT TACACATTTACAGACTATACAATACATTGGGTTTCGACAGGCTCCCGGGCAAGGGCTCGAATGGATGGGATGGATTTAT CCCCCTCAGGGGCTCAATTAACCTATGCTGAGAAAATTTAAGGGTCGTGTAACACTCACCGCCGATAAATCCACCTCAACC GTATATATGGAGCTTTCTTCTCTTCGCTCTGAAGATACCGCCGTCTATTTCTGCGCACGACACGGGGCATACTATTCT AATGCTTTTGACTIONTGGGGACAAGGGACACTTGTGACCGTTAGTAGC	309
	VL	GAGATAGTTATGACTCAGTCTCCCGCCACACTTTTCAGTAAGTCCCGGTGAACGCGCCACCCTGTCTGCGGTGCTTCC GAATCAGTGGATAATGACGGCATTAGGTTTTTGCAGTGGTACCAACAAAAAGCCCGGACAGGCCCCCCCGCCTGCTGATA	306

		TATCGTGCATCAACACGAGCAACAGGGATCCCCGCTCGATTTAGTGGATCCGGAAGCAGGACCGAATTTACACTTACC ATTTCCCTCACTTCAGTCAGAAGATTTTCGCCGTTTACTACTGTCCAGCAGTCAAATAAGGATCCTTACACATTTGGGGGC GGTACAAAAGTCGAGATCAAA	
2419-0605	VH	CAGGTGCAGTTGGTCCAGAGCGGGGCAGAGGTTAAGAAGCCTGGGGCCTCAGTAAAGGTATCCTGCAAGGCTTCTGGG TACACCTTCACAGATTACACTATTTCATTGGGTGCGCCAAGCACCTGGTCAAGGCCTTGAAATGGATGGGATGGATTTAC CCCTTGCGAGGGAGTATTAATTATGCACAGAAGTTCCAGGGAAGGGTTACTCTTACCGCCGACAAGTCCACATCAACC GTTTACATGGAGCTTTCCCTCTCTCAGGTCCGAAGACACTGCTGTATATTTCTGCGCTCGGCATGGGGCTTATTACAGC AACGCCTTCGATTACTGGGGTCAGGGTACATTGGTTCACAGTGTCCAGT	319
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAGCCCAGGGGAGCGAGCAACCCCTCAGTTGCAGAGCATCT GAGAGTGTGATAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCTGACAGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTCGCTGTGTATTACTGTTCAGCAATCCAACAAGGACCCATACACTTTTCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-0805	VH	GAGGTCCAGTTGGTCCAGTCAGGAGCCGAAGTCAAGAAGCCTGGGGAAAGCCTGAAAATAAGTTGCAAAGCTAGTGGA TATACATTTACAGATTATACCAATTCATTGGGTCCGGCAAATGCCAGGAAAAGGCTTGGAGTGGATGGGGTGGATTTAT CCCCCTCCGAGGCTCAATAAATTATAGTCTAGTTTTTCAGGGGCAGGTAAGTATTAGCGCTGATAAAAGTATTTCTACA GTTTATTTGCAGTGGAGTTCATTGAAGGCTAGTGACACCGCTATGTATTTCTGCGCTAGACATGGTGCATATTATTCA AATGCCCTTCGACTATTGGGGCCAGGGCACCCCTCGTCACTGTGAGTTCC	307
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAGCCCAGGGGAGCGAGCAACCCCTCAGTTGCAGAGCATCT GAGAGTGTGATAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCTGACAGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTCGCTGTGTATTACTGTTCAGCAATCCAACAAGGACCCATACACTTTTCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-0105	VH	CAAGTGCAGTTGGTCCAGAGTGGAGCAGAGGTGAAGAAGCCTGGTGCTTCCGTCAAGGTGAGTTGCAAGGCATCTGGT TATACTTTCACTGACTACACAATTCATTGGGTCCAGGCAGGCCCTTGGACAGGGACTGGAATGGATGGGATGGATCTAT CCACTTAGAGGATCAATCAACTATGCTCAAAAAGTTCCAGGGTCGTGTAACAATGACCGCAGACAAAAGTATCTCAACT GTATACATGGAATTTGTCGGATTGAGGAGCGACGACACAGCCGTATATTATTGTGCCAGGCACGGAGCCTACTACAGT AATGCCCTTCGACTACTGGGGGCAGGGCACCCCTTGTTACCGTGTCCAGC	310
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAGCCCAGGGGAGCGAGCAACCCCTCAGTTGCAGAGCATCT GAGAGTGTGATAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCTGACAGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTCGCTGTGTATTACTGTTCAGCAATCCAACAAGGACCCATACACTTTTCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-1204	VH	CAAGTGCAGCTCGTTCACTCTGGCGCAGAAGTGAAGAAGCCAGGAGCTTCCGTTAAAGTGTCTGTAAAGCCTCTGGA TATACATTCACAGATTATACAATTCAGTGGGTGAGACAAGCAACCGGTCAAGGTCTCGAATGGATGGGCTGGATATAC CCCCCTCCGAGGTTCCATCAACTACGCTCAAAAATTCGAAGGACGAGTCACTATGACAGCAAAACAAGAGTTCTTCCACT GTATATATGGAACCTCTCTAGTTTGGCGCTCTGAAGACACCGCCGTGTAATTTCTGTGCCAGGCACGGCGCATACTATTCT AATGCATTTGACTATTGGGGGCAGGGCACATTGGTAACAGTTAGTTCC	311
	VL	GAAATTGTAATGACCCAGAGCCCCGCCACCCCTTAGTGTFCCCCAGGCGAGAGGGCCACTCTTTCTTGCCGCGCAAGC GAATCCGTAGACAACGATGGTATAAGATTTTTGCATTGGTATCAGCAAAAGCCAGGCCAGGCACCCCGGCTTCTCATC	312

		TACAGAGCTAGCACCCCTCGAAACTGGAATCCCCGCTCGTTTTTCAGGATCTGGTAGCGGAACAGAATTTACTTTGACA ATTAGTAGTTTTGCAGTCAGAGGACTTTGCTGTCTATTATTGCCAGCAGTCTAATAAAGATCCATACACCTTCGGCGGA GGGACCAAAGTAGAGATTTAAA	
2419-1210	VH	CAAGTGCAGCTCGTTTCAGTCTGGCGCAGAAGTGAAGAAGCCAGGAGCTTCCGTTAAAGTGTCTGTAAAGCCTCTGGA TATACATTACAGATTATACAATTCACTGGGTGAGACAAGCAACCGGTCAAGGTCTCGAATGGATGGGCTGGATATAC CCCCCTCCGAGGTTCCATCAACTACGCTCAAAAAATTCCAAGGACGAGTCACTATGACAGCAAACAAGAGTTCCCTCCACT GTATATATGGAACCTCTAGTTTGGCGCTCTGAAGACACCGCCGTGTACTTCTGTGCCAGGCACGGCGCATACTATTCT AATGCATTTGACTATTGGGGGCAGGGCACATTGGTAACAGTTAGTTCC	311
	VL	GACATTGTAATGACCCAGTCTCCCGATAGCCTCGCTGTCTCACTCGGAGAACCGCGCAACCATCAACTGCAAGTCTCTCC CAAAGCGTTGACAATGACGGCATTAGGTTTTTTGCACTGGTACCAGCAGAAAACCCGGTCAACCTCCTAAGTTGCTCATT TACCGAGCATCTACCCGCGAGTCAAGGAGTACCTGATCGCTTTTTCCGGTAGCGGTAGTGAACAGATTTTACTCTGACC ATTAGTTCACTCCAGGCAGAAGATGTGGCTGTCTACTACTGCCAACAGTCAAATAAAGACCCTTATACCTTCGGTGGG GGTACCAAAGTAGAGATCAAA	318
2419-1406	VH	CAAGTTCAGTTGGTGC AAAAGCGGGGCAGAAGTGAAGAAACCTGGTGTCTCTGTGAAAGTTTCTGCAAGGCCAGCGGC TACACCTTTACTGATTACACAATACACTGGGTACGGCAGGCAACTGGGCAAGGATTGGAATGGATGGGGTGGATATAC CCATTGCCAGGGTCTATAAACTACGCACAGAAAATTTCAAGGTCCGACTACAATGACAGCCGACAAAATCAATAAGCACC GTTTATATGGAACCTCTCATCTCTCAGGAGTGAGGATACCGCCGTGTATTTCTGCGCACGACACGGTGCATATTACTCA AACGCTTTTCGACTATTGGGGCCAGGGCACCCCTTGTGACTGTTAGTAGC	313
	VL	GAGATAGTTATGACTCAGTCTCCCGCCACACTTTTCAGTAAGTCCCGGTGAACGCGCCACCCTGTCTGCGGTGCTTCC GAATCAGTGGATAATGACGGCATTAGGTTTTTTGCACTGGTACCAACAAAAAGCCCGGACAGGCCCCCCCGCCTGCTGATA TATCGTGCATCAACACGAGCAACAGGGATCCCCGCTCGATTTAGTGGATCCGGAAGCAGGACCGAATTTTACACTTACC ATTTCCCTCACTTCAGTCAAGAAGATTTTCGCCGTTTACTACTGTCAAGCAGTCAAATAAGGATCCTTACACATTTGGGGGC GGTACAAAAGTTCGAGATCAAA	306
2419-1205	VH	CAAGTGCAGCTCGTTTCAGTCTGGCGCAGAAGTGAAGAAGCCAGGAGCTTCCGTTAAAGTGTCTGTAAAGCCTCTGGA TATACATTACAGATTATACAATTCACTGGGTGAGACAAGCAACCGGTCAAGGTCTCGAATGGATGGGCTGGATATAC CCCCCTCCGAGGTTCCATCAACTACGCTCAAAAAATTCCAAGGACGAGTCACTATGACAGCAAACAAGAGTTCCCTCCACT GTATATATGGAACCTCTAGTTTGGCGCTCTGAAGACACCGCCGTGTACTTCTGTGCCAGGCACGGCGCATACTATTCT AATGCATTTGACTATTGGGGGCAGGGCACATTGGTAACAGTTAGTTCC	311
	VL	GAGATAGTAATGACTCAGTCTCCCGCTACACTTAGTGTAAAGCCCAGGGGAGCGAGCAACCCCTCAGTTGCAGAGCATCT GAGAGTGTTGATAATGATGGAATACGTTTTCTCCATTGGTATCAACAAAAACCAGGGCAGGCCCCAGATTGCTGATC TACCGTGCTTCCAATCGCGAGACTGGCATTCTCTGCACGTTTCAGCGGCAGCGGCTCCGGAACCGAGTTTACACTTACT ATTAGCTCACTCCAGTCTGAAGACTTTCGCTGTGTATTACTGTCAAGCAATCCAACAAGGACCCATACACTTTTCGGAGGC GGCACTAAGGTTGAGATCAAA	305
2419-0206	VH	CAGGTGCAACTTGTTCAGTCAGGGGCTGAAGTAAAGAAGCCAGGCTCATCAGTCAAGGTATCATGCAAAAGCATCTGGC TATACATTTACAGATTACACCATTCAGTGGGTGAGGCAAGCTCCCGGTCAAGGTCTCGAGTGGATGGGGTGGATATAC CCTCTCAGAGGCTCTATAAAATTACGCTCAGAAAATTTCAAGGGAGAGTTACAATTACTGCTGATAAAAGTACCAGCACT GCTTATATGGAGCTTTCTCTCACTTCGTTTCAGAGGACACCGCCGTTTACTTTTTGTGCCCGGCATGGTGCCTATTATTCA AATGCCCTTCGATTATTGGGGGCAGGGAACTTTGGTTCACAGTTTCATCT	308
	VL	GAGATAGTTATGACTCAGTCTCCCGCCACACTTTTCAGTAAGTCCCGGTGAACGCGCCACCCTGTCTGCGGTGCTTCC GAATCAGTGGATAATGACGGCATTAGGTTTTTTGCACTGGTACCAACAAAAAGCCCGGACAGGCCCCCCCGCCTGCTGATA	306

		TATCGTGCATCAACACGAGCAACAGGGATCCCCGCTCGATTTAGTGGATCCGGAAGCAGGACCGAATTTTACACTTACC ATTTCCCTCAGTTTCAGTCAGAAGATTTTCGCCGTTTACTACTGTTCAGCAGTCAAATAAGGATTCCTTACACATTTGGGGGC GGTACAAAAGTCGAGATCAAA	
2621	VH	GAGGTCCAGCTTCAGCAGTCTGGAGCTGAGCTGGTGAGGCCCTGGGTCCCTCAGTGAAGATGTCTCTGCAAGACTTCTGGA TATACTTTCACAAGCTACGGTATAAACTGGGTGAAGCAGAGGCCCTGGACAGGGCCCTGGAATGGATTGGATATATTTAT ATTGGAAATGGTTATGCTGAGTACAATGAGAGGTTCAAGGGCAAGGCCACACTGACTTCAGACACATCCTCCAGCACA GCCTACATGCAGCTCAGCAGCCTGACATCTGAGGACTCTGCAATCTATTTCTGTGCACTATACTATCCCTGGTTTACT TACTGGGGCCAGGGGACTCTGGTCACTGTCTCTGCA	75
	VL	GACATCCAGATGACTCAGTCTCCAGCCTCCCTTTTCTGCATCTGTGGGAGATTCTGTCAACCATCACATGTCTGAGCAAGT GAGAATATTTACAGTTATTTAGCATGGTATCAGCAGAAAACAGGGAAAAATCTCCTCAGCTCCTGGTCTATAATGCAAAA ACCTTAGCTGAAGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGCACACAGTTTTCTCTGAAGATCAACAGCCTG CAGCCTGAAGATTTTGGGAATTATTACTGTCAACATCATTATGATACTCCGTTACGTTCCGAGGGGGGACCAAGCTG GAAATAAAA	76
2922	VH	CAGGTTTCAGCTGCACCAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTTCAGTGAAGTTGTCTCTGCAAGACTTCTGGC TACACCTTCACAAGCTACGATGTCTTCTGGGTGAAGCAGAGGCCCTGGACAGGGACTTGAGTGGATTGGATGGATTTAT CCTAGAGATAGTAGTACTAAATACAATGAGAAGTTCAAGGGCAAGGCCACATTGACTGTAGACACATCCTCCAGCACA GCATACATGGAGCTCCACAGCCTGACATCTGAGGACTCTGCCGTCTATTTCTGTGCAAAAGAGGGGTATGATTATGAC AAGAGGGGCTTTGACTACTGGGGCCAAGGCCACCACTCTCACAGTCTCCTCA	77
	VL	GACATTGTGCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGGCAGAGGGCCATCATCTCCTGCAAGGGCCAGC CAAAGTGTCAGTTTTTGTGGTACTAATTTAATGCACTGGTACCAACAGAGACCAGGGCAGCAACCCAAAACCTCCTCATC TATCGTGCATCCAACCTAGAACCTGGGGTTCCCTACCAGGTTTAGTGGCAGTGGGTCTAGGACAGACTTCACCCCTCAAT ATCCATCCTGTGGAGGAAGATGATGCTGCAACCTATTACTGTTCAGCAAAAGTAGGGAATATCCGTGGACGTTCCGGTGG GGCACCAAGCTGGAAATCAAA	78
3125	VH	CAGGTTCAACTGCAGCAGTCTGGGGCTGAGCTGGTGAGGCCCTGGGGCTTCAGTGACGCTGTCTCTGCAAGGCTTCGGGC TACACTTTTTACTGACTATGAAATGCACTGGGTGAAGCAGACACCTGTGCATGGCCTGGAATGGATTGGAGCTATTGAT CCTGAAACTGGTGGTACTGCCTACAATCAGAGGTTCAAGGGCAAGGCCATACTGACTACAGACAAATCCTCCATCACA GCCTACATGGAGCTCCGCAGCCTGACATCTGAGGACTCTGCCGTCTATTACTGTACAAGATGGAATGATGGCGACTAC TGGGGCCAAGGCCACCACTCTCACAGTCTCCTCA	79
	VL	GATGTTGTGATGACCCAGACTCCACTGTCTTTGTGGTTACCAATTGGACAACCAGCCTCCATTTCTTGCAAGTCAAGT CAGAGCCTCTTATACAGTAATGGAAAGACATATTTGAATTTGGTTTCAACAGAGGCCCTGGCCAGTCTCCAAGCGCCTA ATGTATCAGGTGTCCAAACTGGACCCTGGCATCCCTGACAGGTTTCAGTGGCAGTGGATCAGAAACAGATTTTACACTT AAAATCAGCAGAGTGGAGGCTGAAGATTTGGGACTTTATTACTGCTTGCAAGGTACATATTATCCGTACACGTTCCGGA GGGGGGACCAAGCTGGAAATAAAA	80
3327	VH	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTTCAGTGAAGATGTCTCTGCAAGGCTTCTGGT TACTCCTTTTACTGGCTACTTTTGAAGTGGGTGAAGCAGAGCCATGAAAAGAGCCTTGAGTGGATTGGACGTATTAAT CCTTACAATGGTGATACTTTCTACAACCAGAAGTTCAAGGGCAAGGCCACATTGACTGTAGACAAATCCTCTAGCACA GCCCACATGGAGCTCCGGAGCCTGACATCTGAGGACTCTGCACTCTATTATTGTGCAAGCGAAGGTGATGGTTACTAC TGGTACTTCGATGTCTGGGGCGCAGGGACCACGGTCACCGTCTCCTCA	81
	VL	GACATTGTGCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGGCAGAGGGCCACCATCTCCTGCAGAGCCAGC GAAAGTGTGATAATTATGGCATTAGTTTTATGAACTGGTTCCAACAGAAACCAGGACAGCCACCCAAAACCTCCTCATC	82

		TATGCTGCATCCAACCAAGGATCCGGGGTCCCTGCCAGGTTTAGTGGCAGTGGGTCTGGGACAGACTTCAGCCTCAAC ATCCATCCTATGGAGGAGATGATACTGCAATGTATTTCTGTGCAGCAAAGTAAGGAGGTTTCCTCGGACGTTTCGGTGGA GGCACCAAGCTGGAAATCAAA	
3525	VH	CAGGTTCAACTGCAGCAGTCTGGGGCTGAGCTGGTGAGGCCCTGGGGCTTCAGTGAAGCTGTCTCTGCAAGGCTTCGGGC TACACATTTACTGACCATGAAATGCACTGGGTGAGACAGACACCTGTGCATGGCCTGGAATGGATTGGAGTTATTGAT CCTGACACTGGTGATACTACCTACAATCAGAAATTCAGGGGCAAGGCCACACTGACTGCAGACAAATCCTCCAGCACA GCCTACATGGACCTCCGCAGCCTGACATCTGAGGACTCTGCCGTCTTTTACTGTACACGGTGGACTGGGGGGGACTAC TGGGGCCATGGCACCACTCTCACAGTCTCCTCA	83
	VL	GATGTTGTGATGACCCAGACTCCACTGTCTTTGTGCGTTTACCATTGGACAACCAGCCTCCATTTCTTGCAAGTCAAGT CAGAGCCTCTTATACAGTAATGGAAAAGACATATTTGAATTGGTTTCAACAGAGGCCCTGGCCAGTCTCCAAAGCGCCTA ATGTATCAGGTGTCCAAACTGGACCTGGCATCCCTGACAGGTTTCAGTGGCAGTGGATCAGAAACAGATTTTACACTT AAAAATCAGCAGAGTGGAGGCTGAAGATTTGGGACTTTATTACTGCTTGCAAGGTACATATTATCCGTACACGTTTCGGA GGGGGGACCAAGCTGGAAATAAAA	80
3530	VH	CAGGTTCAACTGCAGCAGTCTGGGGCTGAGCTGGTGAGGCCCTGGGGCTTCAGTGAAGCTGTCTCTGCAAGGCTTCGGGC TACACATTTACTGACCATGAAATGCACTGGGTGAGACAGACACCTGTGCATGGCCTGGAATGGATTGGAGTTATTGAT CCTGACACTGGTGATACTACCTACAATCAGAAATTCAGGGGCAAGGCCACACTGACTGCAGACAAATCCTCCAGCACA GCCTACATGGACCTCCGCAGCCTGACATCTGAGGACTCTGCCGTCTTTTACTGTACACGGTGGACTGGGGGGGACTAC TGGGGCCATGGCACCACTCTCACAGTCTCCTCA	83
	VL	GATGCTGTGATGACCCAGACTCCACTGTCTTTGTGCGTTTACCATTGGACAACCAGCCTCTATCTCTTGCAAGTTCGAGT CAGAGCCTCTTATATAGTGATGGAAAAGACATATTTGAATTGGTTTCCAAACAGAGGCCAGGCCAGTCTCCAAAGCGCCTA ATGTATCAGGTGTCCAAACTGGACCTGGCATCCCTGACAGGTTTCAGTGGCAGTGGATCAGAGACAGATTTTACACTT AAAAATCAGCAGAGTGGAGGCTGAGGATTTGGGAGTTTATTACTGCTTGCAAGGTACATATTATCCGTATACGTTTCGGA TCGGGGACCAAGCTGGAAATAAAA	84
4035	VH	CAGGTGCAGCTGAAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCCTGTCCATCACCTGCACAGTCTCTGGT TTCTCATTAACCATCTATGATGTACACTGGGTTTCGCCAGTCTCCAGGAAAGGGTCTGGAGTGGCTGGGAGTGATATGG AGTGATGGAAGCACAGACTATAATGCAGCTTTCATATCTAGACTGAGCATCAGCAAGGACAACCTCCAAGAGCCAAGTT TTCTTTAAAAATGAACAGTCTGCAAGCTGATGACACAGCCATATACTACTGTGCCAGAAATTGGGTGACACAGGCCTGG TTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTCTCTGCA	173
	VL	GACATCCAGATGACTCAGTCTCCAGCCTCCCTATCTGCATCTGTGGGAGAAACTATCACCATCACATGTGCGAGCAAGT AAGAATATTTACAGTTATTTAGCATGGTATCAGCAGAAACAGGGAAAAATCTCCTCAGCTCCTGGTCTATAATGCAAAA ACCTTACCAGAAGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGCACACAGTTTTCTCTGAAGATCAACAGCCTG CAGCCTGAAGATTTTGGGAGTTATTACTGTCAACATCATTATGGTACTCCGCTCACGTTTCGGTGCTGGGACCAAGCTG GAGCTGAAA	174
4035-062	VH	CAGGTACAACCTCCAGGAATCCGGGCTGGGCTCGTCAAAACCAAGCGAAAACACTCTCTCTCACTGCACCGTTTCTGGG TTTTCTCTTACTATCTATGACGTACATTGGGTAAGGCAACCACCCGGGAAGGGGCTGGAGTGGATCGGTGTAATCTGG TCAGATGGATCTACAGACTACAACCCATCCCTTAAAAAGCAGGGTGACCATTTCTAAGGACACTTCCAAGAACCAAGTA TCCCTTAAATTTGCTCTCTGTAAACCGCAGCAGACACCGCAGTTTACTACTGCGCACGAAATTGGGTTGACCAAGCATGG TTTGCAATTTGGGGACAGGGAACCTTTGTCAGTGTGTCTTCA	299
	VL	GATATTCAAATGACCCAATCCCCCTCATCACTTTTCAGCATCTGTGCGTGATCGGGTCACCATTACTTGCAGAGCCAGT AAGAATATCTACAGTACCTGGCTTGGTATCAGCAAAAACCTGGTAAGGCCCTAAACTTCTCGTTTACAATGCTAAG	300

		ACCCTTCCCGAGGGAGTTCCCTTCCAGGTTTTCCGGTAGCGGGAGTGGAACAGATTTACCTTGACTATTTCTAGCTTG CAGCCCGAGGATTTCCGCTACATACTACTGCCAGCATCACTATGGAACCCCCCTGACCTTCGGTCAGGGAACCAAGCTC GAGATCAAA	
3934	VH	CAGGTCCAACATGCAGCAGTCTGGACCTGAGCTGGTGAAGCCTGGAGCTTCAGTGAAGCTGTCTCTGCAAGGCTGCTGGC TACATCTTCACTGACTATACTATAAACTGGGTGAAGCAGAGTCTCTGGACAGGGACTTGAGTGGATTGGATGGATTTAT CCTGGAAGTGGTAATCGTAAATACAATGACAAGTTCAAGGGCAAGGCCACAATGACTGCAGACAAAATCCTCCAGCACA GCCTACATGCAGCTCAGCAGCCTGACCTCTGAGGATTCTGCGGTCTATTTCTGTGCAAGAGAGAGTAACACGTGGGG TACTATGCTATGGACTATTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCA	175
	VL	GATGTTTTGATGACCCAAACTCCACTCTCCCTGCCTGTCTAGTCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGT CAGAGCGTTGTAAATAGTAATGGAAACACCTATTTAGAATGGTACCTGCAGAAAACCAGGCCAGTCTCCAAATCTCCTG ATCTACAAAAGTTTCCAATCGATTTTCTGGGGTCCCAGACAGGTTTCAGTGGCAGTGGATCGGGGACAGATTTACACTC AAGATCAGCAGAGTGGAGGCTGAGGATCTGGGAGTTTATTACTGTTTTCAAGGTTACATGTTCCGTGGACGTTCCGT GGAGGCACCAAGCTGGAATCAAA	176
3833	VH	CAGGTCCAGCTGCAGCAGTCTGGAGCTGAGCTGGTGAAGCCTGGGACTTCAGTGAAGATGTCTCTGCAAGGCTGCTGGA TACACCTTCACAAACTACTGGATAGGTTGGGTAAAGCAGAGGCCTGGACATGGCCTTGAGTGGATTGGAGATATTTAC CCTGGAGGTATAGGAGGTGGTTATACTAAGTACAATGAGAAGTTCAAGGGCAAGGCCACACTGACTGCAGACACATCC TCCAGCACAGCCTACATGCAGCTCGGCAGCCTGACATCTGAGGACTCTGCCATCTATTTCTGTTCAAGATCGGAAACT GGACGGGCTATGGACTACTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCA	177
	VL	GACATCCAGATGACACAGTCTCCATCCTCACTGTCTGCATCTCTGGGAGGCCAAAGTCACCATCACTTGCAAGGCAAGC CAAGACATTAATAAGTATATAGCTTGGTACCAACACAAGCCTGGAAAAAGGTCTAGGCTGCTCATAACATTACACATCT ACATTAAAGCCAGGCATCCCATCAAGGTTTCAGTGGAGTGGGTCTGGGAGAGATTATTCCTTCAGCATCAGTGACCTG GAGCCTGAAGATATTGCAACTTATTATTTGTCTACAGTATGATAATCTGAACACGTTTCGGAGGGGGGACCAAGCTGGAA ATAAAA	178
3631	VH	GAGATCCAGCTGCAGCAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTTCAGTGAAGGTGTCTCTGCAAGGCTTCTGGT TATTCATTCACTGACTACAACATCTACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGATATATTGAT CCTTCCAATGGTGGTCTCTGGCTACAACCAGAAAGTTCAAGGGCAAGGCCACATTGACTGTTGACAAGTCTCTCCAGCACA GCCTTCTGTCATCTCAACAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGAAGGGACAACCTACGGCTCG GGGACTATGGACTACTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCA	179
	VL	GACATTGTGATGACCCAGTCTCAAAAAATTCATGTCCACATCAGTAGGAGACAGGGTCAGCATCACCTGTAAGGCCAGT CAGAATGTGGGTACTGATGTATCCTGGTATCAACAGAAACCAGGGAAATCTCCTAAACCACTGATTTACTGGGCATCA AACC GGTTCACTGGAGTCCCTGATCGCTTCATAGGTAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAATGTG CAGTCTGAAGACTTGGCAGATTATTTCTGTGAGCAATATAGCATCTATCCGCTCACGTTCCGGTGTCTGGGACCAAGCTG GAGCTGAAA	180
3732	VH	GAGATCCAGCTGCAGCAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCGTCAGTGAAGGTATCTCTGCAAGGCTTCTGGT TACTCATTCAGTGACGACAACATGTACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGATATATTGAT CCTCTCAATGGTGGTACTGGCTACAACCAGAAATTCAGGGCAAGGCCACACTGACTGTTGACAAGTCTCTCCAGCACA GCCTTCTGTCATCTCAACAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGAAGGGACAACCTACGCCACG GGGACTATGGACTACTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCA	181
	VL	GACATTGTGATGACCCAGTCTCAAAAAATTCATGTCCACATCAGTAGGAGACAGGGTCAGCATCACCTGCAAGGCCAGT AAGAATGTGGGTACTGATGTATCCTGGTATCAACAGAAACCAGGGAAATCTCCTAAACCACTGATTTACTGGGCATCA	182

		AACCGGTTCACTGGAGTCCCTGATCGCTTCACAGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAACAATGTG CAGTCTGAAGACTTGGCAGATTATTTCTGTGAGCAATATAGCAGCTATCCGCTCACGTTCCGGTGCTGGGACCAAGCTG GAGCTGAAA	
4338	VH	GAGGTCCAGCTGCAGCAGTCTGGCCCTGAGCTGGTGAAGCCTGGGGCTTCAGTGAAGATATCCTGCAAGGCTTCTGGA TACACATTCACTGACTACAACATGGACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAAATATTTAT CCTATCAATGGTTATACTGGCTACAACCAGAGGTTCAAGAACAAGGCCACATTGACTGTAGACAAGTCCCTCCAGCACA GCCTACATGGAACTCCACAGCCTGACATCTGAGGACTCTGCGGTCTATTACTGCGCAAGAGATAGTAACTACGTTGGC TGGTACTTCGATGTCTGGGGCGCAGGGACACGGTCACCGTCTCCTCA	183
	VL	GATGTTGTGATGACCCAAACTCCACTCTCCCTGCCTGTGAGTCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGT CAGAGCCTTGATACACAGTAATGGAAACACCTATTTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTCCTG ATCTACAAAAGTTTCCAACCGATTTTCTGGGGTCCCAGACAGGTTTCAGTGGCAGTGGATCAGGGACAGATTTACATTC AAGATCAGCAGAGTGGAGGCTGAGGATCTGGGAGTTTATTTCTGCTCTCAAAGTACACATGTTCTCGGACGTTCCGT GGAGGCACCAAGCTGGAAATCAAA	184
	VL	GACATTGTGCTGACACAGTCTCCTGCTTCCTTAAGTGTATCTCTGGGGCAGAGGGCCACCTTCTCATGCAGGGCCAGC AAAAGTGTGAGTACATCTGGCTATAGTTATATGCACTGGTACCAACAGAAAACCAGGACAGCCACCCAAAAGTCTCATC TATTTTACATCCGACCTAGAACCCTGGGGTCCCTGCCAGGTTCACTGCCAGTGGGTCTGGGACAGACTTCACCCCTAAC ATCCATCCTGTGGAGGAGGAGGATGCTGCAACCTATTACTGTGACACAGTAGGGAGCTTCCGTACCCCTTCGGAGGG GGGACCAAGTTGGAAATAAAA	185
4540	VH	caggtccagctacagcagtcctggacctgagctggggaagcctggggcttcagtggaagatccctgcaaggcttctggc tacaccttcgctgactactatataaaactgggtgaagcagaggcctggacagggacttgagtggaattggatggatTTTT cctggaagtggtagtacttactacaatgagaagttcaagggcaaggccacacttaactgtagacaaatcctccagcaca gctacatgttgctcagcagcctgacctctgaggactctgcggtctatTTTctgtgcaagaggggactccggtagggtct atggactactggggtcaaggaaacctcagtcacccgtctcctca	186
	VL	gacatccagatgacacagtcctccatcctcactgtctgcatctctgggaggcaagtcacccatcacttgcaaggcaagc caagacattaacaaatatatagcttggtaccaacacaagcctggaaaaggctcctaggctgctcatacattacacatct acattacagtcaggcatcccatcaagggttcagtggaagtgggtctgggagagattattccttcagcatcagcaacctg gagcctgaagataatgcaacttattattgtctacagtatgataatcttctcacgttcggtgctgggaccaagctggag ctgaaa	187
4540-063	VH	CAAGTCCAGCTCGTACAGAGCGGGGCAGAGCTGAAGAAGCCTGGGGCCTCCGTCAAGGTCTCCTGTAAGGCTTCTGGT TACACATTTGCCGACTACTACATGAACCTGGGTACGGCAAGCCCCAGGTCAAGGGCTGGAATGGATGGGATGGATTTTT CCAGGGAGCGGCAGCACTTACTACAACCAGAAAATTTCAAGGTCTGTGTGACAATGACCGTGGATAAAAAGCAGCTCTACA GCTTACATGGAGCTTTTCCCGCTTGAGGTCCGATGATACTGCCGTATATTATTGTGCCCGTGGTGACTCAGGTAGGGCC ATGGACTATTGGGGACAGGGCACCCCTCGTGACCGTGTCCAGC	301
	VL	GATATCCAGATGACACAATCCCTTCATCCTTGAGCGCATCAGTTGGCGACAGGGTCACCATAACTTGTGAGGCTAGT CAGGATATTAACAAGTACCTGGCTTGGTATCAACACAAGCCTGGAAAGGCCCCCAAAATTGCTGATTCACTACACCTCT ACATTGGAAACTGGCGTACCCAGTTCGCTTTTTCTGGGAGTGAAGCGGAACTGATTTCACTTTCACTATATCCAGTCTT CAGCCAGAAGATATCGCAACTTACTATTGTCTTCAGTATGATAACTTGCTTACTTTTCGGAGGAGGGACCAAGTTGAA ATCAAG	302
4540-033	VH	CAGGTGCAGTTGGTCCAATCCGGGGCTGAGGTGAAGAAGCCTGGGGCCTCTGTTAAAGTTAGTTGCAAGGCATCAGGC TACACCTTCGCTGACTACTACATCAACTGGGTTAGACAGGCCCCCGGCAGGGGTGGAGTGGATGGGTTGGATTTTT	303

		CCAGGATCAGGTTCAACATATTACGCACAAAAACTGCAAGGTAGAGTAACCATGACAACTGATACTAGCACCTCCACA GCCATATATGGAACCTCCGCTCTCTCAGGAGTGACGATACAGCCGTTTATTACTGCGCCCGTGGGGATTTCAGGCCCGTGCA ATGGATTACTGGGGTCAAGGGACCCCTCGTGACCCGTAAGTTCA	
	VL	GATATCCAGATGACACAATCCCTTCATCCTTGAGCGCATCAGTTGGCGACAGGGTCACCATAACTTGTGAGGCTAGT CAGGATATTAACAAGTACCTGGCTTGGTATCAACACAAGCCTGGAAAGGCCCCCAAATTGCTGATTCACTACACCTCT ACATTGGAAACTGGCGTACCCAGTCGCTTTTCTGGGAGTGGAAGCGGAACTGATTTCACTTTCACTATATCCAGTCTT CAGCCAGAAGATATCGCAACTTACTATTGTCTTCAGTATGATAACTTGCTTACTTTTCGGAGGAGGGACCAAAGTTGAA ATCAAG	302
4237	VH	CAGGCGCACCTGAAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCCTGTCCATCACCTGCACAGTCTCTGGT TTCTCATTAAACCGACTATGATGTACACTGGGTTTCGCCAGTCTCCAGGAAAAGGGTCTGGAGTGGCTGGGAGTGATATGG AATGATGGAAGCACAGACTATAATACAGCTTTTCATATCTAGACTGACCATCAGCAAGGACAACTCCAAGAGCCAAGTT TTCTTTAAAAATGAACAGTCTGCAAGCTGATGACACAGCCATATACTACTGTGCCAGAAATTGGTATGGTGGCTACTGG TTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTCTCTGCA	188
	VL	GACATCCAGATGACTCAGTCTCCAGCCTCCCTATCTGCATCTGCGGGAGAACTGTCACCATCACATGTGATCAAGT GAGAATATTTACAGTTATTTAGCATGGTATCAGCAGAAAACAGGGAAAAATCTCCTCAGCTCCTAGTCTATAATGCAAAAT GCCTTAGCAGAAGGTGTGCCATCGAGGTTCACTGGCAGTGGATCAGTCACACAGTTTTCTCTGAAGATCAACAGCCTG CAGCCTGAAGATTTTGGGAGTTATTACTGTCAACATCATTATGGTACTCCATTACGTTTCGGCTCGGGGACAAAGTTG GAAATAAAA	189
4439	VH	GAGATCCAGCTGCAGCAGTCTGGAGCTGAACTGGTGAAGCCTGGGGCTTCAGTGAAGATATCCTGCAAGGCTTCTGAT TACTCATTCACTGGCTACAACATGAACTGGGTGATGCAGAGCCATGGAAAAGAGCCTTGAGTGGATTGGAAATATTCAT CCTTACTATGGTGGTACTAGCTTCAATCAGAAGTTCATGGGCAAGGCCACATTGACTGCAGACAAATCTTCCAGCACA GCCTACATGCAGCTCAACAGCCTGACATCTGAAGACTCTGCAGTCTATTACTGTGCAAGAGAGAGAAGTAACCTTCCAT GCTCTGGACTACTGGGGTCAGGGAACTTCAGTCAACCGTCTCCTCA	297
	VL	GACATTGTGCTGACACAGTCTCCTGCTTCCTTAACTGTATCTCTGGGGCAGAGGGCCACCTTCTCATGCAGGGCCAGC AAAAAGTGTCAGTACATCTGGCTATAGTTATATGCACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTCCTCATC TATTTTACATCCGACCTAGAACCTGGGGTCCCTGCCAGGTTCACTGGCAGTGGGTCTGGGACAGACTTCACCCCTCAAC ATCCATCCTGTGGAGGAGGAGGATGCTGCAACCTATTACTGTGACACAGTAGGGAGCTTCCGTACCCCTTCGGAGGG GGGACCAAGTTGGAAATAAAA	298

Table 5. The Amino Acid Sequences of the Heavy Chain Variable Region (VH) and Light Chain Variable Region (VL) of the Exemplary Humanized Anti-APRIL Antibodies Are Provided As Follows.

Antibody Chain	Amino Acid Sequence	SEQ ID NO
>Hu_2218_VH01	QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYYWNWIRQHPGKGLEWIGYISYDGYNNYNPSLKNRVTISVDTSKNQF SLKLSSVTAADTAVYYCARYDYEDWYFGVWGQGTMTVSS	190
>Hu_2218_VH02	QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYYWSWIRQHPGKGLEWIGYISYDGYTYYNPSLKSRTISVDTSKNQF SLKLSSVTAADTAVYYCARYDYEDWYFGVWGQGTMTVSS	191
>Hu_2218_VH03	QVQLQESGPGLVKPSQTLSTCTVSGYSITSGYYWNWIRQHPGKGLEWIGYISYDGYNNYNPSLKSRTISRDTSKNQF SLKLSSVTAADTAVYYCARYDYEDWYFGVWGQGTMTVSS	192
>Hu_2218_VH04	QVQLQQWGAGLLKPSETLSLTCAVYGYISITSGYYWNWIRQPPGKGLEWIGYISYDGYNNYNPSLKNRVTISVDTSKNQF SLKLSSVTAADTAVYYCANYDYEDWYFGVWGQGTMTVSS	193
>Hu_2218_VH05	QVQLQQWGAGLVKPSSETLSLTCAVYGYISITSGYYWNWIRQPPGKGLEWIGYISYDGYNNYNPSLKNRVTISRDTSKNQF SLKLSSVTAADTAVYYCANYDYEDWYFGVWGQGTMTVSS	194
>Hu_2218_VH06	QVQLQESGPGLVKPSSETLSLTCTVSGYSITSGYYWNWIRQPPGKGLEWIGYISYDGYNNYNPSLKNRVTISVDTSKNQF SLKLSSVTAADTAVYYCANYDYEDWYFGVWGQGTMTVSS	195
>Hu_2218_VH07	QVQLQESGPGLVKPSSETLSLTCTVSGYSITSGYYWNWIRQPPGKGLEWIGYISYDGYNNYNPSLKNRVTISRDTSKNQF SLKLSSVTAADTAVYYCANYDYEDWYFGVWGQGTMTVSS	196
>Hu_2218_VH08	QVQLQESGPGLMKPSETLSLTCSVSGYSITSGYYWSWIRKPPGKLEYIGYVSYDGYSTYYNPSLKSRTISVDTSKNRF SLKLNSVTAADTAVYYCANYDYEDWYFGYWGQGLLVTVSS	197
>Hu_2218_VH09	QVQLQESGPGLMKPSETLSLTCSVSGYSITSGYYWNWIRKPPGKLEYIGYVSYDGYSTYYNPSLKSRTISRDTSKNRF SLKLNSVTAADTAVYYCANYDYEDWYFGVWGQGLLVTVSS	198
>Hu_2218_VH10	EVQLVESGGGLVQPGGSLRLSCAVSGYSITSGYYWNWIRQAPGKGLEWVASISYDGYNNYNPSVKGRITISRDDSKNTF YLQMNSLRAEDTAVYYCANYDYEDWYFGVWGQGLLVTVSS	199
>Hu_2218_VH11	EVQLVESGGGLVQPGGSLRLSCAVSGYSITSGYYWNWIRQAPGKGLEWVAYISYDGYNNYNPSVKGRITISRDTSKNTF YLQMNSLRAEDTAVYYCANYDYEDWYFGVWGQGLLVTVSS	200
>Hu_2218_VH12	QVQLVESGGGVVQPGGRSLRLSCAASGYISITSGYYWNWVRQAPGKGLEWVAYISYDGYNNYNPSVKGRFTISRDNKNTL YLQMNSLRAEDTAVYYCANYDYEDWYFGVWGQGTMTVSS	201
>Hu_2218_VL01	EIVLTQSPATLSLSPGERATLSCRASESVSIIGTNSIHWHYQQKPGQAPRLLIYHASNLETGIPARFSGSGPGTDFTLTI SSLEPEDFAVYYCLQSRKIPYTFGGGTKLEIK	202
>Hu_2218_VL02	DIVLTQSPASLAVSPGQRATITCRASESVSIIGTNSIHWHYQQKPGQPPKLLIYHASNLETGVPARFSGSGSGTDFTLTI NPVEANDTANYCYCLQSRKIPYTFGGGTKLEIK	203
>Hu_2218_VL03	EIVMTQSPATLSVSPGERATLSCRASESVSIIGTNSLHWYQQKPGQAPRLLIYHASQSISGIPARFSGSGSGTEFTLTI SSLQSEDFAVYYCQQSRKIPYTFGGGTKVEIK	204
>Hu_2218_VL04	EIVMTQSPATLSVSPGERATLSCRASESVSIIGTNSIHWHYQQKPGQAPRLLIYHASNLETGIPARFSGSGSRTEFTLTI SSLQSEDFAVYYCLQSRKIPYTFGGGTKVEIK	205
>Hu_2218_VL05	DIQLTQSPSSLSASVGRVTITCRASESVSIIGTNSMNWYQQKPGKAPKLLIYHASYLESGVPSRFSGSGSGTDFTLTI SSLQPEDFATYYCLQSRKIPYTFGGGTKVEIK	206
>Hu_2218_VL06	DIQLTQSPSSLSASVGRVTITCRASESVSIIGTNSMHWYQQKPGKAPKLLIYHASNLESGVPSRFSGSGSRTDFTLTI	207

	SSLQPEDFATYYCLQSRKIPYTFGQGTKVEIK	
>Hu_2218_VL07	DIQMTQSPSSLSASVGDRTITCRASESVSIIGTNSMHWYQQKPGKAPKLLIYHASNLESGVPSRFSGSGSGTDFTFTI SSLQPEDFATYYCLQSRKIPYTFGQGTKVEIK	208
>Hu_2419_VH01	QVQLVQSGAEVKKPGASVKVSCKASGYTFDTYTIHWVRQAPGQGLEWMGWIYPLRGSINYNEKFKDRVTSTRDTSISTA YMELSLRLSDDTVVYYCARHGAYYSNAFDYWGQGTLLVTVSS	209
>Hu_2419_VH02	QVQLVQSGAEVKKPGASVKVSCKASGYTFDTYIMHWVRQAPGQGLEWMGRIYPLRGSTNYAQKFQGRVTSTRDTSISTA YMELSLRLSDDTVVYYCARHGAYYSNAFDYWGQGTLLVTVSS	210
>Hu_2419_VH03	QVQLVQSGAEVKKPGASVKVSCKASGYTFDTYTIHWVRQAPGQGLEWMGWIYPLRGSINYNEKFKDRVTMTADTSSSTA YMELSLRLSDDTVVYYCARHGAYYSNAFDYWGQGTLLVTVSS	211
>Hu_2419_VH04	QVQLVQSGAEVKKPGASVKVSCEASGYTFDTYTIHWVRQAPGKGLEWMGWIYPLRGSINYNEKFKDRVTMTADTSTDTA YMELSSRLSKDTAVYYCARHGAYYSNAFDYWGQGTLLVTVSS	212
>Hu_2419_VH05	QVQLVQSGAEVVKPGASVKLSCKASGYTFDTYTMVWKQAPGQGLEWIGEYIYPLRGSINFNEKFKSKATLTVDKSASTA YMELSSLRSEDTAVYYCARHGAYYSNAFDYWGQGTLLVTVSS	213
>Hu_2419_VH06	QVQLVQSGAEVVKPGASVKLSCKASGYTFDTYTMHWVKQAPGQGLEWIGEYIYPLRGSINFNEKFKSKATLTVDKSASTA YMELSSLRSEDTAVYYCARHGAYYSNAFDYWGQGTLLVTVSS	214
>Hu_2419_VL01	EIVLTQSPATLSLSPGERATLSCRASESDNDGIRFMHWYQQKPGQAPRLLIYRASNLESGIPARFSGSGPGTDFTLTI SSLEPEDFAVYYCQQSNKDPYTFGQGTKLEIK	215
>Hu_2419_VL02	EIVLTQSPATLSLSPGERATLSCRASESDNDGIRFMHWYQQKPGQAPRLLIYRASNLESGIPARFSGSGPGTDFTLTI SSLEPEDFAVYYCQQSNKDPYTFGQGTKLEIK	216
>Hu_2419_VL03	DIVMTQSPASLAVSLGERATINCRASESDNDGIRFMHWYQQKPGQPPKLLIYRASNLESGVPDRFSGSGSRTDFTLTI SSLQAEDFAVYYCQQSNKDPYTFGGGTKVEIK	217
>Hu_2419_VL04	DIVMTQSPASLAVSLGERATINCRASESDNDGIRFMHWYQQKPGQPPKLLIYRASNLESGVPDRFSGSGSRTDFTLTI NSLQAEDFAVYYCQQSNKDPYTFGGGTKVELK	218
>Hu_2419_VL05	DIVLTQSPATLSVSPGERATISCRASESDNDGIRFMHWYQQKPGQPPKLLIYRASNLESGVPARFSGSGSRTDFTLTI SSVEPEDFATYYCQHSWEIPPTFGGGTKLEIK	219
>hu_4035_VH01	QVQLVESGGGVVQPGRSLRLSCAASGFSLTIIYDVHWVRQAPGKGLEWVAVIWSDGSTDYNAAFISRFTISRDNKNTLY LQMNSLRAEDTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	220
>hu_4035_VH02	QVQLVESGGGVVQPGRSLRLSCAASGFSLTIIYDVHWVRQAPGKGLEWVGVVWSDGSTDYNAAFISRFTISKDNKNTLY LQMNSLRAEDTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	221
>hu_4035_VH03	QVQLVESGGGVVQPGRSLRLSCAASGFSLTIIYDVHWVRQAPGKGLEWVAVIWSDGSTDYADSVKGRFTISKDNKNTLY LQMNSLRAEDTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	222
>hu_4035_VH04	QVQLVESGGGVVQPGRSLRLSCAVSGFSLTIIYDVHWVRQAPGKGLEWVGVVWSDGSTDYADSVKGRFTISKDNKNTVY LQMNSLRAEDTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	223
>hu_4035_VH05	QVQLQESGPGLVKPSSETLSLTCTVSGFSLTIIYDVHWIRQPPGKGLEWIGVIWSDGSTDYNAAFISRVTISVDTSKNQFS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	224
>hu_4035_VH06	QVQLQESGPGLVKPSSETLSLTCTVSGFSLTIIYDVHWVRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSKDTSKNQVS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	225
>hu_4035_VH07	QVQLQESGPGLMKPSSETLSLTCSVSGDSITIIYDWHWIRQPPGKGLEWIGVVWSDGSTDYNPSLKSRTVTSVDTSKNRFS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGTLLVTVSS	226
>hu_4035_VH07	QVQLQESGPGLVKPSSETLSLTCTVSGFSLTIIYDVHWIRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSKDNKSNQFS	227

	LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGLTVTVSS	
>hu_4035_VH09	QVQLQESGPGLVKPSSETLSLTCTVSGFSLTIYDVHWIRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSKDTSKNQVS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGLTVTVSS	262
>hu_4035_VH10	QVQLQESGPGLVKPSSETLSLTCTVSGGSITIIDWHWVRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSKDTSKNQFS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGLTVTVSS	263
>hu_4035_VH11	QVQLQESGPGLVKPSSETLSLTCTVSGGSITIIDWHWVRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSVDTSKNQFS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGLTVTVSS	264
>hu_4035_VH12	QVQLQESGPGLVKPSSETLSLTCTVSGGSITIIDWHWIRQPPGKGLEWIGVIWSDGSTDYNPSLKSRTVTSVDTSKNQFS LKLSSVTAADTAVYYCARNWVDQAWFAYWGQGLTVTVSS	265
>hu_4035_VL01	DIQMTQSPSSLSASVGDRTVITCRASKNIYSYLAWYQQKPGKAPKLLIYNAKTLPEGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQHHYGTPLTFGGQGTKLEIK	228
>hu_4035_VL02	DIQMTQSPSSLSASVGDRTVITCRASKNIYSYLAWYQQKPGKAPKLLVYNAKTLPEGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQHHYGTPLTFGGQGTKLEIK	229
>hu_4035_VL03	EIVLTQSPATLSLSPGERATLSCRASKNIYSYLAWYQQKPGQAPRLLIYNAKTRATGIPARFSGSGSGTDFTLTISSE PEDFAVYYCQHHYGTPLTFGGQGTKLEIK	230
>hu_4035_VL04	EIVLTQSPATLSLSPGERATLSCRASKNIYSYLAWYQQKPGQAPRLLVYNAKTLPEGIPARFSGSGSGTDFTLTISSE PEDFAVYYCQHHYGTPLTFGGQGTKLEIK	231
>hu_4035_VL05	EIVMTQSPATLSVSPGERATLSCRASKNIYSYLAWYQQKPGQAPRLLIYNAKTRATGIPARFSGSGSGTEFTLTISSE SEDFAVYYCQHHYGTPLTFGGGTKVEIK	232
>hu_4035_VL06	DIQLTQSPSFLSASVGDRTVITCRASKNIYSYLAWYQQKPGKAPKLLIYNAKSLQSGVPSRFSGSGSGTEFTLTISSE PEDFATYYCQHHYGTPLTFGGGTKLEIK	233
>hu_4035_VL07	DIQLTQSPSFLSASVGDRTVITCRASKNIYSYLAWYQQKPGKAPKLLIYNAKSLQSGVPSRFSGSGSGTEFTLTISSE PEDFATYYCQHHYGTPLTFGGGTKLEIK	234
>hu_4237_VH01	QLQLQESGSGLVKPSQTLSTCAVSGFSLTDYDVHWVRQPPGKGLEWIGVIWNDGSTDYNPSLISRTVTSKDNSKNQVS LKLSSVTAADTAVYYCARNWYGGYWFAYWGQGLTVTVSS	235
>hu_4237_VH02	QLQLQESGSGLVKPSQTLSTCAVSGGSITDWDHWVRQPPGKGLEWIGVIWNDGSTDYNPSLISRTVTSVDNSKNQFS LKLSSVTAADTAVYYCARNWYGGYWFAYWGQGLTVTVSS	236
>hu_4237_VH03	QVQLVESGGGVVQPGRSRLRLSCAASGFSTFDYDMHWVRQAPGKGLEWVAVIWNNDGSTDYATSVIGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCARNWYGGYWFAYWGQGLTVTVSS	237
>hu_4237_VH04	QVQLVESGGGVVQPGRSRLRLSCAASGFSTFDYDMHWVRQAPGKGLEWVGVWVWVNDGSTDYATSVIGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCARNWYGGYWFAYWGQGLTVTVSS	238
>hu_4237_VH05	QVQLQESGPGLMKPSSETLSLTCSVSGGSITDWDHWIRQPPGKGLEWIGVWVWVNDGSTDYNPSLKSRTVTSVDTSKNRF LKLNSVTAADTAVYYCARNWYGGYWFAYWGQGLTVTVSS	239
>hu_4237_VH06	QVTLKESGPALVKPTQTTLTCTFSGFSLTDYDVHWIRQPPGKALEWLAVIWNNDGSTDYSPSLKSRLTITKDTSKNQVV LTMTNMDPVDATATYYCARNWYGGYWFAYWGQGLTVTVSS	240
>hu_4237_VL01	DIQMTQSPSSLSASVGDRTVITCRSSENIYSYLAWYQQKPGKAPKLLVYNANALAEVPSRFSGSGSVTDFTLTISSE PEDFATYYCQHHYGTPTFTFGGQGTKLEIK	241
>hu_4237_VL02	DIQMTQSPSTLSASVGDRTVITCRSSENIYSYLAWYQQKPGKAPKLLVYNANALAEVPSRFSGSGSVTEFTLTISSE PDDEFATYYCQHHYGTPTFTFGGQGTKLEIK	242
>hu_4237_VL03	EIVMTQSPATLSVSPGERATLSCRASENIYSYLAWYQQKPGQAPRLLIYNANASAEVPSRFSGSGSGTEFTLTISSE LQMNSLRAEDTAVYYCARNWYGGYWFAYWGQGLTVTVSS	243

	SEDFAVYYCQQHYGTPFTFGGGTKVEIK	
>hu_4237_VL04	DIQMTQSPSSLSASVGDRTTITCRASENIYSYLAWYQQKPGKAPKLLLYNANRLESGVPSRFSGSGSGTDFTLTISSLQ PEDFASYCQHHYGTPFTFGSGTKLEIK	244
>hu_4237_VL05	DIQMTQSPSSLSASVGDRTTITCRASENIYSYLAWYQQKPGKAPKLLLYNANRLESGVPSRFSGSGSGTDYTLTISSLQ PEDFASYCQHHYGTPFTFGSGTKLEIK	245
>hu_3833_VH01	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIGWVRQAPGQGLEWMGDIYPGGIGGGYTKYNEKFKGRVTMTADTST STAYMELRSLRSDDTAVYYCSRSETGRAMDYWGQGTLLVTVSS	246
>hu_3833_VH02	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIGWVRQAPGQGLEWMGDIYPGGIGGGYTKYAQKLQGRVTMTADTST STAYMELRSLRSDDTAVYYCSRSETGRAMDYWGQGTLLVTVSS	247
>hu_3833_VH03	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIGWVRQAPGQGLEWMGDIYPGGIGGGYTKYAQKFQGRVTMTADTST STAYMELSSLRSEDTAVYYCSRSETGRAMDYWGQGTLLVTVSS	248
>hu_3833_VH04	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIGWVRQAPGQGLEWMGDIYPGGIGGGYTKYNEKFKGRVTMTADTST STAYMELSSLRSEDTAVYFCRSRSETGRAMDYWGQGTLLVTVSS	249
>hu_3833_VH05	QVQLVQSGAELKRPASVKVSCKASGYTFITNYWVGWVKQAPGQGLEWMGDIYPGGIGGGYTNYAQKFQKATMTADTSS STAYMQLSRLRSEDTAVYYCSRSETGRAMDYWGQGTLLVTVSS	250
>hu_3833_VL01	DIQMTQSPSSLSASVGDRTTITCQASQDINKYLAWYQQKPGKAPKLLIHYTSTLKPQVPSRFSGSGSGTDFTFTISSLQ PEDIATYYCLQYDNLNTFGGGTKLEIK	251
>hu_3833_VL02	DIQMTQSPSSLSASVGDRTTITCKASQDINKYIAWYQQKPGKAPKLLIHYTSTLKPQVPSRFSGSGSGRDYFTFTISSLQ PEDIATYYCLQYDNLNTFGGGTKLEIK	252
>hu_3833_VL03	DIQMTQSPSSLSASVGDRTTITCQASQDINKYLAWYQQKPGKAPKLLIYYTSTLETGVPSPRFSGSGSGTDFTFSISSLQ PEDIATYYCLQYDNLNTFGGGTKLEIK	253
>hu_4540_VH01	QVQLVQSGAEVKKPGASVKVSCKASGYTFADYYINWVRQAPGKGLEWMGWIFPGSGSTYYNEKFKGRVTMTVDKSTSTA YMELSSLRSEDTAVYFCARGDSGRAMDYWGQGTLLVTVSS	254
>hu_4540_VH02	QVQLVQSGAEVKKPGASVKVSCKASGYTFADYYMNWVRQAPGQGLEWMGWIFPGSGSTYYAEKFKGRVTSTRDTSISTA YMELSRSLRSDDTVVYYCARGDSGRAMDYWGQGTLLVTVSS	255
>hu_4540_VH03	QVQLVQSGAEVKKPGASVKVSCKASGYTFADYYINWVRQAPGQGLEWMGWIFPGSGSTYYAQKLQGRVTMTTDTSTSTA YMELRSLRSDDTAVYYCARGDSGRAMDYWGQGTLLVTVSS	256
>hu_4540_VH04	QVQLVQSGAEVKKPGASVKVSCKASGYTFADYYINWVRQAPGQGLEWMGWIFPGSGSTYYAQKLQGRVTMTVDKSSSTA YMELRSLRSDDTAVYYCARGDSGRAMDYWGQGTLLVTVSS	257
>hu_4540_VH05	QVQLVQSGAELKKPGASVKVSCKASGYTFADYYMNWVRQAPGQGLEWMGWIFPGSGSTYYNQKFQGRVTMTVDKSSSTA YMELSRSLRSDDTAVYYCARGDSGRAMDYWGQGTLLVTVSS	258
>hu_4540_VL01	DIQMTQSPSSLSASVGDRTTITCKASQDINKYIAWYQQKPGKAPKLLIHYTSTLQSGVPSRFSGSGSGTDFTFTISSLQ PEDIATYYCLQYDNLNTFGQGQTKLEIK	259
>hu_4540_VL02	DIQMTQSPSSLSASVGDRTTITCKASQDINKYIAWYQHKPGKAPKLLIHYTSTLQSGVPSRFSGSGSGRDYFTFTISSLQ PEDIATYYCLQYDNLNTFGQGQTKLEIK	260
>hu_4540_VL03	DIQMTQSPSSLSASVGDRTTITCQASQDINKYLAWYQHKPGKAPKLLIHYTSTLETGVPSPRFSGSGSGTDFTFTISSLQ PEDIATYYCLQYDNLNTFGGGTKVEIK	261

In an embodiment, the antibody molecule comprises one, two, or three CDRs of the VH region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, 4439, or 4237), using the Kabat or Chothia definitions of CDRs. In an embodiment, the antibody molecule comprises one, two, or three CDRs of the VL region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, 4439, or 4237), using the Kabat or Chothia definitions of CDRs. In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) CDRs of the VH region and/or one or more (*e.g.*, two or three) CDRs of the VL region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, 4439, or 4237), using the Kabat or Chothia definitions of CDRs.

In an embodiment, the antibody molecule comprises one, two, or three VH CDRs described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one, two, or three VL CDRs described in **Table 1 or 5**. In an embodiment, the antibody molecule comprises one or more (*e.g.*, two or three) VH CDRs and/or one or more (*e.g.*, two or three) VL CDRs described in **Table 1 or 5**.

In an embodiment, the antibody molecule comprises one, two, three, or four frameworks of the VH region of an antibody molecule described in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, 4439, or 4237). In an embodiment, the antibody molecule comprises one, two, three, or four frameworks of the VL region of an antibody molecule described in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, 4439, or 4237). In an embodiment, the antibody molecule comprises one or more (*e.g.*, two, three, or four) frameworks of the

VH region and/or one or more (*e.g.*, two, three, or four) frameworks of the VL region of an antibody molecule described in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237).

In an embodiment, the antibody molecule comprises a heavy chain variable region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237). In an embodiment, the antibody molecule comprises a light chain variable region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237). In an embodiment, the antibody molecule comprises a heavy chain variable region and a light chain variable region of an antibody molecule described herein, *e.g.*, in **Table 1 or 5** (*e.g.*, any of monoclonal antibodies 2218, 2419, 2419-0105, 2419-0205, 2419-0206, 2419-0406, 2419-0605, 2419-0805, 2419-0806, 2419-1204, 2419-1205, 2419-1210, 2419-1305, 2419-1306, 2419-1310, 2419-1406, 2922, 3327, 3530, 3525, 3125, 2621, 4035, 4035-062, 3934, 3833, 3631, 3732, 4338, 4540, 4540-063, 4540-033, 4439, or 4237).

In an embodiment, the antibody molecule comprises a heavy chain variable region having an amino acid sequence described in **Table 1 or 5**, or an amino acid sequence substantially identical thereof. In an embodiment, the antibody molecule comprises a light chain variable region having an amino acid sequence described in **Table 1 or 5**, or an amino acid sequence substantially identical thereof. In an embodiment, the antibody molecule comprises a heavy chain variable region having an amino acid sequence described in **Table 1 or 5** (or an amino acid sequence substantially identical thereof) and a light chain variable region having an amino acid sequences described in **Table 1 or 5** (or an amino acid sequence substantially identical thereof).

In an embodiment, the antibody molecule comprises a heavy chain variable region encoded by a nucleotide sequence described in **Table 2**, or a nucleotide sequence substantially identical thereof. In an embodiment, the antibody molecule comprises a light chain variable region encoded by a nucleotide sequence described in **Table 2**, or a nucleotide sequence substantially identical thereof. In an embodiment, the antibody molecule comprises a heavy chain variable region encoded by a nucleotide sequence described in **Table 2** (or a nucleotide sequence substantially identical thereof) and a light chain

variable region encoded by a nucleotide sequence described in **Table 2** (or a nucleotide sequence substantially identical thereof).

In an embodiment, the antibody molecule further comprises a heavy chain constant region. In an embodiment, the heavy chain constant region is an IgG1 constant region, *e.g.*, any of SEQ ID NOS: 320-322, or a functional portion thereof. In another embodiment, the heavy chain constant region is an IgG2 constant region, *e.g.*, any of SEQ ID NOS: 323-326, or a functional portion thereof. In an embodiment, the antibody molecule further comprises a light chain constant region. In an embodiment, the antibody molecule further comprises a heavy chain constant region and a light chain constant region. In an embodiment, the antibody molecule comprises a heavy chain constant region, a light chain constant region, and heavy and light chain variable regions of an antibody molecule described in **Table 1 or 5**. In certain embodiments, the antibody molecule comprises a heavy chain constant region, a light chain constant region, and variable regions that comprise one, two, three, four, five, or six CDRs of an antibody molecule described in **Table 1 or 5**.

Exemplary heavy chain constant regions are described below.

Exemplary IgG1 constant regions

>IGHG1*01

STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
PSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL
PAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVL
DSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSP (SEQ ID NO: 320)

>IGHG1*03

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
VPSSSLGTQTYICNVNHKPSNTKVDKRVKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA
LPAPIEKTISKAKGQPREPQVYTLPPSRDEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPV
LDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSP (SEQ ID NO: 321)

>IGHG1*04

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA
LPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPV
LDSDGSFFLYSKLTVDKSRWQQGNIFSCFSVMHEALHNHYTQKSLSLSP (SEQ ID NO: 322)

Exemplary IgG2 constant regions

>IGHG2*01

STKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
PSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV
VVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPI

EKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDG
SFFLYSKLTVDKSRWQQGNVFSVSMHEALHNHYTQKSLSLSP (SEQ ID NO: 323)

>IGHG2*02

5 STKGPSVFPFLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV
TSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLFPPKPKDITLMISRTPEVTCV
VVDVSHEDPEVQFNWYVDGMEVHNAKTKPREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPI
EKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDG
SFFLYSKLTVDKSRWQQGNVFSVSMHEALHNHYTQKSLSLSP (SEQ ID NO: 324)

10

>IGHG2*04

STKGPSVFPFLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV
PSSSLGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLFPPKPKDITLMISRTPEVTCV
VVDVSHEDPEVQFNWYVDGMEVHNAKTKPREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPI
15 EKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDG
SFFLYSKLTVDKSRWQQGNVFSVSMHEALHNHYTQKSLSLSP (SEQ ID NO: 325)

>IGHG2*06

STKGPSVFPFLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV
20 PSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLFPPKPKDITLMISRTPEVTCV
VVDVSHEDPEVQFNWYVDGMEVHNAKTKPREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPI
EKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDG
SFFLYSKLTVDKSRWQQGNVFSVSMHEALHNHYTQKSLSLSP (SEQ ID NO: 326)

25

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or
30 100% homology with, the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 12; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

35 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2
40 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 285; or

an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

5 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 12; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

10 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 285; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises one or both of:

15 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 17; an HCDR2 comprising an amino acid
20 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 282; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

25 (ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 280; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95,
30 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 285; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 17; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 282; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of SEQ ID NO: 280; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 285; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 296. In an embodiment, the antibody molecule comprises a VL comprising the amino acid sequence of SEQ ID NO: 286. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 296 and a VL comprising the amino acid sequence of SEQ ID NO: 286.

In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 313. In an embodiment, the antibody molecule comprises a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 306. In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 313 and a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 306.

In an embodiment, the antibody molecule further comprises a heavy constant region of IgG2, *e.g.*, any of SEQ ID NOS: 323-326.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 12; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 285; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 12; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 285; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 17; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 282; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or

100% homology with, the amino acid sequence of SEQ ID NO: 280; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 285; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least
5 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 17; an
10 HCDR2 comprising the amino acid sequence of SEQ ID NO: 282; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of SEQ ID NO: 280; an
15 LCDR2 comprising the amino acid sequence of SEQ ID NO: 285; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 289. In an embodiment, the antibody molecule comprises a VL comprising the amino acid sequence of SEQ ID NO: 286. In an embodiment, the antibody molecule comprises a VH comprising the
20 amino acid sequence of SEQ ID NO: 289 and a VL comprising the amino acid sequence of SEQ ID NO: 286.

In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 308. In an embodiment, the antibody molecule comprises a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 305. In an embodiment,
25 the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 308 and a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 306.

In an embodiment, the antibody molecule further comprises a heavy constant region of IgG2, *e.g.*, any of SEQ ID NOS: 323-326.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid

sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 12; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 281; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 11; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 12; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of SEQ ID NO: 280; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 281; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 17; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 282; or an HCDR3 comprising an amino

acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 13, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 280; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 281; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 17; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 282; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 13, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of SEQ ID NO: 280; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 281; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 16.

In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 289. In an embodiment, the antibody molecule comprises a VL comprising the amino acid sequence of SEQ ID NO: 284. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 289 and a VL comprising the amino acid sequence of SEQ ID NO: 284.

In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 308. In an embodiment, the antibody molecule comprises a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 305. In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 308 and a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 305.

In an embodiment, the antibody molecule further comprises a heavy constant region of IgG2, *e.g.*, any of SEQ ID NOS: 323-326.

In an embodiment, the antibody molecule comprises one or both of:

5 (i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 93; an HCDR2 comprising an amino acid
10 sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 94; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 95, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three
15 light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the LCDR1 of SEQ ID NO: 96; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least
20 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 97; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 98.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three
25 heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 93; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 94; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 95, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three
30 light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of the LCDR1 of SEQ ID NO: 96; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 97; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 98.

In an embodiment, the antibody molecule comprises one or both of:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises one, two, or all of the following: an HCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 99; an HCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 273; or an HCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 95, or

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises one, two, or all of the following: an LCDR1 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 96; an LCDR2 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of the SEQ ID NO: 97; or an LCDR3 comprising an amino acid sequence that differs by no more than 1, 2, or 3 amino acid residues from, or has at least 85, 90, 95, 99 or 100% homology with, the amino acid sequence of SEQ ID NO: 98.

In an embodiment, the antibody molecule comprises:

(i) a heavy chain variable region (VH), wherein the heavy chain variable region comprises three heavy chain complementarity determining regions (HCDR1, HCDR2, and HCDR3), wherein the heavy chain variable region comprises: an HCDR1 comprising the amino acid sequence of SEQ ID NO: 99; an HCDR2 comprising the amino acid sequence of SEQ ID NO: 273; and an HCDR3 comprising the amino acid sequence of SEQ ID NO: 95, and

(ii) a light chain variable region (VL), wherein the light chain variable region comprises three light chain complementarity determining regions (LCDR1, LCDR2, and LCDR3), wherein the light chain variable region comprises: an LCDR1 comprising the amino acid sequence of SEQ ID NO: 96; an LCDR2 comprising the amino acid sequence of SEQ ID NO: 97; or an LCDR3 comprising the amino acid sequence of SEQ ID NO: 98.

In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 225. In an embodiment, the antibody molecule comprises a VL comprising the amino acid sequence of SEQ ID NO: 229. In an embodiment, the antibody molecule comprises a VH comprising the amino acid sequence of SEQ ID NO: 225 and a VL comprising the amino acid sequence of SEQ ID NO: 229.

In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 299. In an embodiment, the antibody molecule comprises a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 300. In an embodiment, the antibody molecule comprises a VH encoded by a nucleic acid comprising the nucleotide sequence of
 5 SEQ ID NO: 299 and a VL encoded by a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 300.

In an embodiment, the antibody molecule further comprises a heavy chain constant region of IgG1, *e.g.*, any of SEQ ID NOS: 320-322.

10 In an embodiment, the antibody molecule described herein has one or more (*e.g.*, 2, 3, 4, 5, or all) of the following properties: (a) is a humanized antibody molecule; (b) binds to human APRIL at an EC_{50} of 60 pM or less, as determined by ELISA; (c) inhibits binding of human APRIL to TACI, *e.g.*, *in vitro*, at an IC_{50} of 0.5 nM or less; (d) inhibits binding of human APRIL to BCMI, *e.g.*, *in vitro*, at an IC_{50} of 0.6 nM or less; (e) is an IgG2 κ ; or (f) has an Fc region engineered to reduce complement activation. In an
 15 embodiment, the antibody molecule comprises one or more (*e.g.*, 2, 3, 4, 5, or all) CDRs, one or both of heavy chain variable region or light chain variable regions, or one or both of heavy chain or light chain, of any of antibody molecules 2419-1406, 2419-0205, or 2419-0206. In an embodiment, the antibody molecule is suitable for use in treating a disorder in kidney, *e.g.*, IgA nephropathy. In another embodiment, the antibody molecule is suitable for use in treating a cancer, *e.g.*, a multiple myeloma.

20 In an embodiment, the antibody molecule described herein has one or more (*e.g.*, 2, 3, 4, 5, or all) of the following properties: (a) is a humanized antibody molecule; (b) binds to human APRIL at an EC_{50} of 50 pM or less, as determined by ELISA; (c) inhibits binding of human APRIL to TACI, *e.g.*, *in vitro*, at an IC_{50} of 0.3 nM or less; (d) inhibits binding of human APRIL to BCMA, *e.g.*, *in vitro*, at an IC_{50} of 0.2 nM or less; (e) is an IgG1 κ ; or (f) has higher BCMA neutralization activity, *e.g.*, has an IC_{50} of 0.1
 25 nM or less. In an embodiment, the antibody molecule comprises one or more (*e.g.*, 2, 3, 4, 5, or all) CDRs, one or both of heavy chain variable region or light chain variable regions, or one or both of heavy chain or light chain, of antibody molecule 4035-062. In an embodiment, the antibody molecule is suitable for use in treating a cancer or an autoimmune disorder.

30 The antibody molecules described herein can have several advantageous properties. For example, the antibody molecules can be used to effectively treat, prevent or diagnose a disorder associated with APRIL, *e.g.*, a disorder described herein, *e.g.*, IgA nephropathy.

In an embodiment, the antibody molecule is capable of binding, or substantially binding, to human APRIL and mouse APRIL. In an embodiment, the antibody molecule is capable of binding, or

substantially binding, to human APRIL, but is not capable of binding, or substantially binding to mouse APRIL. In an embodiment, the antibody molecule binds to APRIL with high affinity, *e.g.*, with a dissociation constant (K_D) of less than about 100 nM, typically about 10 nM, and more typically, about 10-0.001 nM, about 10-0.01 nM, about 10-0.01 nM, about 5-0.01 nM, about 3-0.05 nM, about 1-0.1 nM, or stronger, *e.g.*, less than about 80, 70, 60, 50, 40, 30, 20, 10, 8, 6, 4, 3, 2, 1, 0.5, 0.2, 0.1, 0.05, 0.01, 0.005, or 0.001 nM. In an embodiment, the antibody molecule binds to APRIL with a K_{off} slower than 1×10^{-4} , 5×10^{-5} , or $1 \times 10^{-5} \text{ s}^{-1}$. In an embodiment, the antibody molecule binds to APRIL with a K_{on} faster than 1×10^4 , 5×10^4 , 1×10^5 , or $5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$.

In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to TACI. In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to TACI. In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to BCMA. In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to TACI and BCMA. In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to TACI, but is not capable of inhibiting, or substantially inhibiting, binding of human APRIL to BCMA. In an embodiment, the antibody molecule is capable of inhibiting, or substantially inhibiting, binding of human APRIL to BCMA, but is not capable of inhibiting, or substantially inhibiting, binding of human APRIL to TACI.

In an embodiment, the antibody molecule inhibits binding of human APRIL to human TACI by 50% or more, *e.g.*, 60% or more, 70% or more, 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more, 99% or more, or 100%, as determined by a method described herein (*e.g.*, normalized to the no antibody control).

In an embodiment, the antibody molecule inhibits binding of human APRIL to human BCMA by 30% or more, *e.g.*, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more, 99% or more, or 100%, as determined by a method described herein (*e.g.*, normalized to the no antibody control).

In an embodiment, the antibody molecule does not substantially inhibit binding of human APRIL to human BCMA, *e.g.*, inhibits binding of human APRIL to human BCMA by less than 10%, as determined by a method described herein (*e.g.*, normalized to the no antibody control).

In an embodiment, the antibody molecule binds to a linear or conformational epitope on APRIL. In an embodiment, the antibody molecule binds to an epitope conserved between human APRIL and mouse APRIL. In an embodiment, the antibody molecule binds to an epitope described herein. In an embodiment, the antibody molecule binds, or substantially binds, to the same, similar, or overlapping epitope on APRIL, as a second antibody molecule (*e.g.*, a monoclonal antibody described in **Table 1** or

5). In an embodiment, the antibody molecule competes with a second antibody molecule (*e.g.*, a monoclonal antibody described in **Table 1 or 5**) for binding to APRIL.

In an embodiment, the antibody molecule binds, or substantially binds, one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of APRIL as defined in **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 3**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises APRIL residues from two monomers, *e.g.*, one or more residues from monomer A and monomer B as shown in **Table 3**.

In an embodiment, the antibody molecule binds, or substantially binds, one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, residues within a region of APRIL as defined in **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the human APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 4**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises one or more APRIL residues from the C-D loop (*e.g.*, the loop connecting β -sheets C and D), the G-H loop (*e.g.*, the loop connecting β -sheets G and H), or both.

In an embodiment, the antibody molecule binds, or substantially binds, to one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of human APRIL from positions 105-114 and/or one or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or all) residues of mouse APRIL from positions 96-105.

In an embodiment, the antibody molecule binds, or substantially binds, one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or more, residues within a region of APRIL as defined in **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all of the human APRIL residues from **Table 7**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 7**.

In an embodiment, the antibody molecule binds, or substantially binds, one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, residues within a region of APRIL as defined in **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that comprises or consists of one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15,

16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 8**. In an embodiment, the antibody molecule binds, or substantially binds, to an epitope that overlaps an epitope that comprises or consists of all of the human APRIL residues from **Table 8**.

In an embodiment, the epitope is a conformational epitope.

5 In an embodiment, the antibody molecule does not bind, or does not substantially bind, to one, two or all of Asp129, Arg233, or His203 of human APRIL.

In an embodiment, binding of the antibody molecule to APRIL (*e.g.*, human APRIL) inhibits, or substantially inhibits, the binding of the CRD2 domain of TACI (*e.g.*, human TACI) to APRIL (*e.g.*, human APRIL). In another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the APRIL residues from **Table 3**. In yet another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, or all of the human APRIL residues from **Table 4**. In still another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or all of the human APRIL residues from **Table 7**. In still another embodiment, binding of the antibody molecule to human APRIL, inhibits, or substantially inhibits, the binding of human TACI, to one or more, *e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or all of the human APRIL residues from **Table 8**.

20

Animal Models

The antibody molecules described herein can be evaluated *in vivo*, *e.g.*, using various animal models. For example, an animal model can be used to test the efficacy of an antibody molecule described herein in inhibiting APRIL and/or in treating or preventing a disorder described herein, *e.g.*, IgA nephropathy. Animal models can also be used, *e.g.*, to investigate for side effects, measure concentrations of antibody molecules *in situ*, demonstrate correlations between an APRIL function and a disorder described herein (*e.g.*, IgA nephropathy).

Exemplary animal models for IgA nephropathy that can be used for evaluating an antibody molecule described herein include, but are not limited to, a ddY mouse model for spontaneous IgA nephritis (Imai *et al. Kidney Int.* 1985; 27(5):756-761); a mouse model utilizing inert proteins or a common viral pathogen as the inciting antigen (Emancipator *et al. Curr. Protoc. Immunol.* 2001 May; Chapter 15: Unit 15.11), a rat model by noninfectious protein antigens (Emancipator *et al. Curr. Protoc. Immunol.* 2001 May; Chapter 15: Unit 15.11); a chronic mouse model of IgA immune-complex-associated nephropathy (Montinaro *et al. Nephrol. Dial. Transplant.* 1995; 10(11): 2035-2042); the Gne

30

M712T mouse as a model for human glomerulopathy (Kakani *et al. Am. J. Pathol.* 2012; 180(4):1431-1440); a mouse IgA nephropathy model with the MBP-20-peptide fusion protein (Zhang *et al. Anat. Rec. (Hoboken).* 2010; 293(10): 1729-1737); and a mouse model for IgA immune complex nephritis (Rifai *et al. J Exp Med.* 1979; 150(5):1161-1173). Other animal models for IgA nephropathy are described, *e.g.*,
5 in Tomino *et al. J. Nephrol.* 2008; 21(4):463-467; Endo *Ren. Fail.* 1997; 19(3):347-371; and Rifai *Kidney Int.* 1987; 31(1):1-7.

Exemplary animal models for other disorders described herein are also known in the art. Exemplary types of animals that can be used to evaluate the antibody molecules described herein include, but are not limited to, mice, rats, rabbits, guinea pigs, and monkeys.

Pharmaceutical Compositions and Kits

In some aspects, this disclosure provides compositions, *e.g.*, pharmaceutically acceptable compositions, which include an antibody molecule described herein (*e.g.*, a humanized antibody molecule described herein), formulated together with a pharmaceutically acceptable carrier.

As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, isotonic and absorption delaying agents, and the like that are physiologically compatible. The carrier can be suitable for intravenous, intramuscular, subcutaneous, parenteral, rectal, spinal or epidermal administration (*e.g.*, by injection or infusion). In certain embodiments, less than about 5%, *e.g.*, less than about 4%, 3%, 2%, or 1% of the antibody molecules in the pharmaceutical composition are present as
20 aggregates. In other embodiments, at least about 95%, *e.g.*, at least about 96%, 97%, 98%, 98.5%, 99%, 99.5%, 99.8%, or more of the antibody molecules in the pharmaceutical composition are present as monomers. In some embodiments, the level of aggregates or monomers is determined by chromatography, *e.g.*, high performance size exclusion chromatography (HP-SEC).

The compositions set out herein may be in a variety of forms. These include, for example, liquid,
25 semi-solid and solid dosage forms, such as liquid solutions (*e.g.*, injectable and infusible solutions), dispersions or suspensions, liposomes, and suppositories. A suitable form depends on the intended mode of administration and therapeutic application. Typical suitable compositions are in the form of injectable or infusible solutions. One suitable mode of administration is parenteral (*e.g.*, intravenous, subcutaneous, intraperitoneal, intramuscular). In some embodiments, the antibody molecule is administered by
30 intravenous infusion or injection. In certain embodiments, the antibody is administered by intramuscular or subcutaneous injection.

The phrases “parenteral administration” and “administered parenterally” as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital,

intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

Therapeutic compositions typically should be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high antibody concentration. Sterile injectable solutions can be prepared by incorporating the active compound (*i.e.*, antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

The antibody molecules described herein can be administered by a variety of methods. Several are known in the art, and for many therapeutic, prophylactic, or diagnostic applications, an appropriate route/mode of administration is intravenous injection or infusion. For example, the antibody molecules can be administered by intravenous infusion at a rate of less than 10mg/min; preferably less than or equal to 5 mg/min to reach a dose of about 1 to 100 mg/m², preferably about 5 to 50 mg/m², about 7 to 25 mg/m² and more preferably, about 10 mg/m². As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, *e.g.*, *Sustained and Controlled Release Drug Delivery Systems*, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

In certain embodiments, an antibody molecule can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The antibody molecule (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the antibody molecule may be

incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer an antibody molecule by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. Therapeutic, prophylactic, or diagnostic compositions can also be administered with medical devices, and several are known in the art.

Dosage regimens are adjusted to provide the desired response (*e.g.*, a therapeutic, prophylactic, or diagnostic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms are dictated by and directly dependent on (a) the unique characteristics of the antibody molecule and the particular therapeutic, prophylactic, or diagnostic effect to be achieved, and (b) the limitations inherent in the art of compounding such an antibody molecule for the treatment of sensitivity in individuals.

An exemplary, non-limiting range for a therapeutically, prophylactically, or diagnostically effective amount of an antibody molecule is about 0.1-50 mg/kg body weight of a subject, *e.g.*, about 0.1-30 mg/kg, *e.g.*, about 1-30, 1-15, 1-10, 1-5, 5-10, or 1-3 mg/kg, *e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, or 50 mg/kg. The antibody molecule can be administered by intravenous infusion at a rate of less than 10 mg/min, *e.g.*, less than or equal to 5 mg/min to reach a dose of about 1 to 100 mg/m², *e.g.*, about 5 to 50 mg/m², about 7 to 25 mg/m², *e.g.*, about 10 mg/m². It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

The pharmaceutical compositions herein may include a “therapeutically effective amount,” “prophylactically effective amount,” or “diagnostically effective amount” of an antibody molecule described herein.

A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody molecule may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual.

A therapeutically effective amount is also one in which any toxic or detrimental effect of the antibody molecule is outweighed by the therapeutically beneficial effects. A “therapeutically effective dosage” typically inhibits a measurable parameter by at least about 20%, *e.g.*, by at least about 40%, by at least about 60%, or by at least about 80% relative to untreated subjects. The measurable parameter may be, *e.g.*, hematuria, colored urine, foamy urine, pain, swelling (edema) in the hands and feet, or high blood pressure. The ability of an antibody molecule to inhibit a measurable parameter can be evaluated in an animal model system predictive of efficacy in treating or preventing IgA nephropathy. Alternatively, this property of a composition can be evaluated by examining the ability of the antibody molecule to inhibit APRIL, *e.g.*, by an *in vitro* assay.

A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

A “diagnostically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired diagnostic result. Typically, a diagnostically effective amount is one in which a disorder, *e.g.*, a disorder described herein, *e.g.*, IgA nephropathy, can be diagnosed *in vitro*, *ex vivo*, or *in vivo*.

Also within this disclosure is a kit that comprises an antibody molecule, described herein. The kit can include one or more other elements including: instructions for use; other reagents, *e.g.*, a label, a therapeutic agent, or an agent useful for chelating, or otherwise coupling, an antibody molecule to a label or therapeutic agent, or a radioprotective composition; devices or other materials for preparing the antibody molecule for administration; pharmaceutically acceptable carriers; and devices or other materials for administration to a subject.

Nucleic Acids

The present disclosure also features nucleic acids comprising nucleotide sequences that encode the antibody molecules (*e.g.*, heavy and light chain variable regions and CDRs of the antibody molecules), as described herein.

For example, the present disclosure features a first and second nucleic acid encoding heavy and light chain variable regions, respectively, of an antibody molecule chosen from one or more of the antibody molecules disclosed herein, *e.g.*, an antibody molecule of **Table 1 or 5**, or a portion of an antibody molecule, *e.g.*, the variable regions of **Table 2**. The nucleic acid can comprise a nucleotide sequence encoding any one of the amino acid sequences in the tables herein, or a sequence substantially

identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in the tables herein).

In certain embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs from a heavy chain variable region having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions). In some embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs from a light chain variable region having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions). In some embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, three, four, five, or six CDRs from heavy and light chain variable regions having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions).

In certain embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs from a heavy chain variable region having the nucleotide sequence as set forth in **Table 2**, a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein). In some embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs from a light chain variable region having the nucleotide sequence as set forth in **Table 2**, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein). In certain embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, three, four, five, or six CDRs from heavy and light chain variable regions having the nucleotide sequence as set forth in **Table 2**, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein).

In certain embodiments, the nucleic acid comprises a nucleotide sequence as set forth in **Table 2** or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein). In some embodiments, the nucleic acid comprises a portion of a nucleotide sequence as set forth in **Table 2** or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein).

The portion may encode, for example, a variable region (*e.g.*, VH or VL); one, two, or three or more CDRs; or one, two, three, or four or more framework regions.

The nucleic acids disclosed herein include deoxyribonucleotides or ribonucleotides, or analogs thereof. The polynucleotide may be either single-stranded or double-stranded, and if single-stranded may be the coding strand or non-coding (antisense) strand. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. The nucleic acid may be a recombinant polynucleotide, or a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which either does not occur in nature or is linked to another polynucleotide in a non-natural arrangement.

In some aspects, the application features host cells and vectors containing the nucleic acids described herein. The nucleic acids may be present in a single vector or separate vectors present in the same host cell or separate host cell, as described in more detail below.

Vectors

Further provided herein are vectors that comprise nucleotide sequences encoding an antibody molecule described herein.

In an embodiment, the vector comprises a nucleotide encoding an antibody molecule described herein, *e.g.*, as described in **Table 1 or 5**. In another embodiment, the vector comprises a nucleotide sequence described herein, *e.g.*, in **Table 2**. The vectors include, but are not limited to, a virus, plasmid, cosmid, lambda phage or a yeast artificial chromosome (YAC).

Numerous vector systems can be employed. For example, one class of vectors utilizes DNA elements which are derived from animal viruses such as, for example, bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (Rous Sarcoma Virus, MMTV or MOMLV) or SV40 virus. Another class of vectors utilizes RNA elements derived from RNA viruses such as Semliki Forest virus, Eastern Equine Encephalitis virus and Flaviviruses.

Additionally, cells which have stably integrated the DNA into their chromosomes may be selected by introducing one or more markers which allow for the selection of transfected host cells. The marker may provide, for example, prototrophy to an auxotrophic host, biocide resistance (*e.g.*, antibiotics), or resistance to heavy metals such as copper, or the like. The selectable marker gene can be either directly linked to the DNA sequences to be expressed, or introduced into the same cell by cotransformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcriptional promoters, enhancers, and termination signals.

Once the expression vector or DNA sequence containing the constructs has been prepared for expression, the expression vectors may be transfected or introduced into an appropriate host cell. Various techniques may be employed to achieve this, such as, for example, protoplast fusion, calcium phosphate precipitation, electroporation, retroviral transduction, viral transfection, gene gun, lipid based transfection
5 or other conventional techniques. In the case of protoplast fusion, the cells are grown in media and screened for the appropriate activity.

Methods and conditions for culturing the resulting transfected cells and for recovering the antibody molecule produced are known to those skilled in the art, and may be varied or optimized depending upon the specific expression vector and mammalian host cell employed, based upon the
10 present description.

Cells

The present disclosure also provides cells (*e.g.*, host cells) comprising a nucleic acid encoding an antibody molecule as described herein. For example, the host cells may comprise a nucleic acid molecule
15 having a nucleotide sequence described in **Table 2**, a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein), or a portion of one of said nucleic acids. Additionally, the host cells may comprise a nucleic acid molecule encoding an amino acid sequence of **Table 1 or 5**, a sequence substantially homologous thereto (*e.g.*, a sequence at least about 80%, 85%, 90%, 95%, 99% or
20 more identical thereto), or a portion of one of said sequences.

In some embodiments, the host cells are genetically engineered to comprise nucleic acids encoding the antibody molecule described herein.

In certain embodiments, the host cells are genetically engineered by using an expression cassette. The phrase “expression cassette,” refers to nucleotide sequences, which are capable of affecting
25 expression of a gene in hosts compatible with such sequences. Such cassettes may include a promoter, an open reading frame with or without introns, and a termination signal. Additional factors necessary or helpful in effecting expression may also be used, such as, for example, an inducible promoter.

The disclosure also provides host cells comprising the vectors described herein.

The cell can be, but is not limited to, a eukaryotic cell, a bacterial cell, an insect cell, or a human
30 cell. Suitable eukaryotic cells include, but are not limited to, Vero cells, HeLa cells, COS cells, CHO cells, HEK293 cells, BHK cells and MDCKII cells. Suitable insect cells include, but are not limited to, Sf9 cells. In an embodiment, the cell (*e.g.*, host cell) is an isolated cell.

Uses of Antibody Molecules

The antibody molecules disclosed herein, as well as the pharmaceutical compositions disclosed herein, have *in vitro*, *ex vivo*, and *in vivo* therapeutic, prophylactic, and/or diagnostic utilities.

In an embodiment, the antibody molecule reduces (*e.g.*, inhibits, blocks, or neutralizes) one or more biological activities of APRIL. For example, these antibodies molecules can be administered to cells in culture, *in vitro* or *ex vivo*, or to a subject, *e.g.*, a human subject, *e.g.*, *in vivo*, to reduce (*e.g.*, inhibits, blocks, or neutralizes) one or more biological activities of APRIL. In an embodiment, the antibody molecule inhibits, or substantially inhibit, binding of APRIL, *e.g.*, human APRIL, to TACI, BCMA, or both. Accordingly, in an aspect, the disclosure provides a method of treating, preventing, or diagnosing a disorder, *e.g.*, a disorder described herein (*e.g.*, IgA nephropathy), in a subject, comprising administering to the subject an antibody molecule described herein, such that the disorder is treated, prevented, or diagnosed. For example, the disclosure provides a method comprising contacting the antibody molecule described herein with cells in culture, *e.g.* *in vitro* or *ex vivo*, or administering the antibody molecule described herein to a subject, *e.g.*, *in vivo*, to treat, prevent, or diagnose a disorder, *e.g.*, a disorder associated with APRIL (*e.g.*, IgA nephropathy).

As used herein, the term “subject” is intended to include human and non-human animals. In some embodiments, the subject is a human subject, *e.g.*, a human patient having a disorder described herein (*e.g.*, IgA nephropathy), or at risk of having a disorder described herein (*e.g.*, IgA nephropathy). The term “non-human animals” includes mammals and non-mammals, such as non-human primates. In some embodiments, the subject is a human. The methods and compositions described herein are suitable for treating human patients a disorder described herein (*e.g.*, IgA nephropathy). Patients having a disorder described herein (*e.g.*, IgA nephropathy) include those who have developed a disorder described herein (*e.g.*, IgA nephropathy) but are (at least temporarily) asymptomatic, patients who have exhibited a symptom of a disorder described herein (*e.g.*, IgA nephropathy), or patients having a disorder related to or associated with a disorder described herein (*e.g.*, IgA nephropathy).

Methods of Treating or Preventing Disorders

The antibody molecules described herein can be used to treat or prevent disorders associated with APRIL or symptoms thereof.

Exemplary disorders or conditions that can be associated with APRIL include, but are not limited to IgA nephropathy, diabetic nephropathy, cancer (*e.g.*, hematological cancer (*e.g.*, B-cell non-Hodgkin’s lymphoma, chronic lymphocytic leukemia, Hodgkin’s lymphoma, multiple myeloma, Waldenström macroglobulinemia, and lymphoplasmacytic lymphoma) or solid tumors (*e.g.*, colorectal cancer, breast cancer (*e.g.*, breast carcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), brain cancer (*e.g.*,

glioblastoma), and kidney cancer (*e.g.*, renal cell carcinoma)), immunoproliferative disorders (*e.g.*, monoclonal IgA hypergammaglobulinemia), vasculitis (*e.g.*, kidney vasculitis, Henoch-Schonlein purpura (IgA associated vasculitis), and post-streptococcal glomerulonephritis), autoimmune disorders (*e.g.*, rheumatoid arthritis, systemic lupus erythematosus, linear IgA bullous disease/linear immunoglobulin A (IgA) dermatosis, and IgA-mediated epidermolysis bullosa acquisita), IgA pemphigus, celiac disease, and alcoholic cirrhosis. In an embodiment, the disorder is associated with aberrant expression of IgA. In an embodiment, the antibody molecule is used to treat a subject having a disorder described herein, or is at risk of developing a disorder described herein.

The antibody molecules described herein are typically administered at a frequency that keeps a therapeutically effective level of antibody molecules in the patient's system until the patient recovers. For example, the antibody molecules may be administered at a frequency that achieves a serum concentration sufficient for at least about 1, 2, 5, 10, 20, 30, or 40 antibody molecules to bind each APRIL molecule. In an embodiment, the antibody molecules are administered every 1, 2, 3, 4, 5, 6, or 7 days, every 1, 2, 3, 4, 5, or 6 weeks, or every 1, 2, 3, 4, 5, or 6 months.

Methods of administering various antibody molecules are known in the art and are described below. Suitable dosages of the antibody molecules used will depend on the age and weight of the subject and the particular drug used.

In an embodiment, the antibody molecule is administered to the subject (*e.g.*, a human subject) intravenously. In an embodiment, the antibody molecule is administered to the subject at a dose between 0.1 mg/kg and 50 mg/kg, *e.g.*, between 0.2 mg/kg and 25 mg/kg, between 0.5 mg/kg and 10 mg/kg, between 0.5 mg/kg and 5 mg/kg, between 0.5 mg/kg and 3 mg/kg, between 0.5 mg/kg and 2.5 mg/kg, between 0.5 mg/kg and 2 mg/kg, between 0.5 mg/kg and 1.5 mg/kg, between 0.5 mg/kg and 1 mg/kg, between 1 mg/kg and 1.5 mg/kg, between 1 mg/kg and 2 mg/kg, between 1 mg/kg and 2.5 mg/kg, between 1 mg/kg and 3 mg/kg, between 1 mg/kg and 2.5 mg/kg, or between 1 mg/kg and 5 mg/kg. In an embodiment, the antibody molecule is administered to the subject at a fixed dose between 10 mg and 1000 mg, *e.g.*, between 10 mg and 500 mg, between 10 mg and 250 mg, between 10 mg and 150 mg, between 10 mg and 100 mg, between 10 mg and 50 mg, between 250 mg and 500 mg, between 150 mg and 500 mg, between 100 mg and 500 mg, between 50 mg and 500 mg, between 25 mg and 250 mg, between 50 mg and 150 mg, between 50 mg and 100 mg, between 100 mg and 150 mg, between 100 mg and 200 mg, or between 150 mg and 250 mg. In an embodiment, the antibody molecule is administered once a week, twice a week, once every two weeks, once every three weeks, once every four weeks, once every eight weeks, once a month, once every two months, or once every three months. In an embodiment, the antibody molecule is administered between 0.5 mg/kg and 3 mg/kg or between 50 mg and 150 mg, once a week, twice a week, once every two weeks, or once every four weeks.

The antibody molecules can be used by themselves or conjugated to a second agent, *e.g.*, a bacterial agent, toxin, or protein, *e.g.*, a second anti-APRIL antibody molecule. This method includes: administering the antibody molecule, alone or conjugated to a second agent, to a subject requiring such treatment. The antibody molecules can be used to deliver a variety of therapeutic agents, *e.g.*, a toxin, or mixtures thereof.

IgA Nephropathy

IgA nephropathy (also known as Berger's disease, Berger disease, Berger's syndrome, Berger syndrome, IgA nephritis, IgAN, or synpharyngitic glomerulonephritis) is the most prevalent, chronic glomerular disease worldwide. Conservative epidemiological estimates cite a global incidence of approximately 5-50 cases/million (children) and 10-40 cases /million (adults). This incidence of disease presents a regional bias with a higher prevalence in Asia and the Americas, with a particularly higher disease burden in Japan and regions of China. Biopsy confirmed cases of IgA nephropathy in Japan are projected at approximately 350,000. In the US, this projection is approximately 100,000—as such, it is the most frequently diagnosed 1° glomerular disease in adults. While a relatively indolent disease, IgA nephropathy leads to end stage renal disease (ESRD), *i.e.*, renal failure in 20-50% of patients within a 20-30 year span. These numbers are likely grossly underreported given the need to confirm the disease by kidney biopsy, a protocol that is variably practiced in various clinical settings. The disease has a complex pathogenesis with genetic, epidemiological, and potentially environmental components to disease etiology, pathology, and progression. It likewise has a variable clinical presentation ranging from asymptomatic to end-stage renal failure (ESRD). There are currently no disease-specific treatments to address primary disease or progression.

The etiology of this disease, as its name implies, has been established. In brief, the disease is caused by the deposition of IgA, typically in the form of immune complexes in the mesangium of the kidney. A molecular characterization of these particular immunoglobulins has been carried out. These IgAs are of the A1 subclass (IgA1 vs. IgA2), predominantly polymeric (with J chain-mediated linkages), and apparently differentially o-glycosylated in the hinge region that is intervening between CH1 and CH2 domains. In particular, these o-glycans are heterogeneously lacking β 1,3 galactose linkages and, as such, are commonly referred to as galactose-deficient IgA1 (or gdlIgA1). As the pathogenesis of this disease can involve a polygenic, multi-hit mechanism for inducing renal pathology and aberrant physiology, IgA1 may be viewed as the so-called auto-antigen representing this first critical “hit” in a multi-hit model for IgA nephropathy. A set of autoantibodies for this disease has likewise been defined and it relates to immunoglobulins (predominantly IgG) that specifically recognize this differentially glycosylated epitope and promote the formation of immune complexes (representing so-called “hit 2”). It should also be noted

that IgA itself is subject to aggregation due to misfolding, conformational changes, and potential changes in the N-glycosylation state of the CH2/CH3 glycans.

Without wishing to be bound by theory, it is believed that in an embodiment, aberrantly glycosylated IgA1 levels correlate with disease and clinical outcomes in IgA nephropathy. Aberrantly glycosylated IgA1 has been characterized directly from kidney biopsies and increased production of aberrantly glycosylated IgA1 was observed in B cells (tonsillar, PBMC) in IgA nephropathy patients. The level of galactose-deficient IgA1 in the sera of patients with IgA nephropathy is associated with disease progression (Zhao *et al. Kidney Int.* 2012; 82(7):790-6). Differential lectin staining demonstrated elevated levels of aberrantly glycosylated IgA1 in serum and glomeruli of IgA nephropathy patients relative to healthy controls (Allen *et al. Kidney Int.* 2001; 60(3):969-73).

Based on this evolving disease model, IgA nephropathy may be appropriately viewed as an autoimmune disease with strong and critical extra-renal involvement. The identification and validation of select immune-based targets proposed to play a critical role in disease pathogenesis, namely the production of IgA and subsequent production of autoreactive antibodies to this target, represent a logical therapeutic strategy for treatment. APRIL (TNFSF13) represents particular area of focus for this reason. Additional rationale for targeting APRIL include emerging genetic data based on multiple, comprehensive genome wide association (GWAS) studies along with IgA related genetic disorders *e.g.*, IgA hypogammaglobulinemia related common variable immunoglobulin deficiency (CVID) whose locus maps to defects in TNFRSF13B (TACI) with direct implications of the role of APRIL-TACI interactions in regulating IgA synthesis.

IgA nephropathy often does not cause symptoms in the early stages. The disease can go unnoticed for years and is sometimes first diagnosed when routine tests reveal protein and red blood cells in urine that cannot be seen without a microscope (microscopic hematuria). Signs and symptoms of IgA nephropathy when kidney function is impaired include, *e.g.*, cola- or tea-colored urine (caused by red blood cells in the urine); repeated episodes of cola- or tea-colored urine, sometimes even visible blood in the urine, usually during or after an upper respiratory or other type of infection; pain in the side(s) of the back below the ribs (flank); foam in the toilet water from protein in the urine; swelling (edema) in the hands and feet; and high blood pressure. In an embodiment, the sign or symptom includes, *e.g.*, one or more of hematuria, proteinuria, albuminuria, hypertension, or an early stage kidney disease (*e.g.*, requiring dialysis or transplantation). In an embodiment, the sign or symptom is associated with, *e.g.*, one or more of aberrantly glycosylated IgA1, auto-antibody formation, deposition of nephritogenic immune complexes in the kidney, or inflammation and loss of kidney function.

The classic presentation (in about 40-50% of the cases, more common in younger adults) of IgA nephropathy is episodic hematuria which usually starts within a day or two of a non-specific upper

respiratory tract infection (hence synpharyngitic). Less commonly gastrointestinal or urinary infection can be the inciting agent. All of these infections have in common the activation of mucosal defenses and hence IgA antibody production. These episodes can occur on an irregular basis every few months and in most patients eventually subsides. Renal function usually remains normal, though rarely, acute kidney failure may occur.

A smaller proportion (in about 20-30% of the cases, usually the older population) of IgA nephropathy patients have microscopic hematuria and proteinuria (less than 2 gram/day). These patients may not have any symptoms and are only clinically found if a doctor decides to take a urine sample. Hence, the disease is more commonly diagnosed in situations where screening of urine is compulsory, *e.g.*, school children in Japan.

Some (about 5% each) IgA nephropathy patients have the following disease presentation: nephrotic syndrome (*e.g.*, 3-3.5 grams of protein loss in the urine, associated with a poorer prognosis); acute kidney failure (*e.g.*, either as a complication of the frank hematuria, when it usually recovers, or due to rapidly progressive glomerulonephritis which often leads to chronic kidney failure); chronic kidney failure (*e.g.*, no previous symptoms, presents with anemia, hypertension and other symptoms of kidney failure, in people who probably had longstanding undetected microscopic hematuria and/or proteinuria).

A variety of systemic diseases can be associated with IgA nephropathy such as liver failure, celiac disease, rheumatoid arthritis, reactive arthritis, ankylosing spondylitis and HIV. Diagnosis of IgA nephropathy and a search for any associated disease occasionally reveals such an underlying serious systemic disease. Occasionally, there are simultaneous symptoms of Henoch-Schönlein purpura. Some HLA alleles have been suspected along with complement phenotypes as being genetic factors.

IgA nephropathy can be diagnosed by various tests, *e.g.*, urine test, blood tests (*e.g.*, to show increased blood levels of the waste product creatinine), iothalamate clearance test, kidney imaging (*e.g.*, ultrasound, X-rays, or cystoscopy), kidney biopsy, or a combination thereof.

For an adult patient with isolated hematuria, tests such as ultrasound of the kidney and cystoscopy are usually done first to pinpoint the source of the bleeding. These tests would rule out kidney stones and bladder cancer, two other common urological causes of hematuria. In children and younger adults, the history and association with respiratory infection can raise the suspicion of IgA nephropathy. A kidney biopsy is often necessary to confirm the diagnosis. The biopsy specimen shows proliferation of the mesangium, with IgA deposits on immunofluorescence and electron microscopy. However, patients with isolated microscopic hematuria (*i.e.*, without associated proteinuria and with normal kidney function) are not usually biopsied since this is associated with an excellent prognosis. A urinalysis will show red blood cells, usually as red cell urinary casts. Proteinuria, usually less than 2 grams per day, also may be present. Other renal causes of isolated hematuria include, *e.g.*, thin basement membrane disease and Alport

syndrome, the latter being a hereditary disease associated with hearing impairment and eye problems. Other blood tests done to aid in the diagnosis include CRP or ESR, complement levels, ANA, and LDH. Protein electrophoresis and immunoglobulin levels can show increased IgA in 50% of all patients.

Treatment with a number of medications can slow the progress of the disease and help manage symptoms such as high blood pressure, protein in the urine (proteinuria), and swelling (edema) in the hands and feet. Exemplary therapies for IgA nephropathy include, *e.g.*, high blood pressure medications (*e.g.*, angiotensin-converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs)), omega-3 fatty acids, immunosuppressants (*e.g.*, corticosteroid medications, such as prednisone), statin therapy, mycophenolate mofetil, ciclosporin, mizoribine, cyclophosphamide (*e.g.*, in combination with anti-platelet/anticoagulants, or in combination with steroids and azathioprine), kidney dialysis, or kidney transplantation. Exemplary therapies for IgA nephropathy are also described in Floege and Eitner *J. Am. Soc. Nephrol.* 22: 1785-1794, 2011. Other exemplary therapies for IgA nephropathy are described in the section of "Combination Therapies" herein.

Without wishing to be bound by theory, it is believed that in an embodiment, targeting APRIL selectively reduces IgA. APRIL^{-/-} mice have normal T and B lymphocyte development, normal T and B cell proliferation *in vitro*, but decreased serum IgA levels (Castigli *et al. Proc Natl Acad Sci U S A.* 2004; 101(11):3903-8). Discovery of new risk loci for IgA nephropathy implicates genes involved in immunity against intestinal pathogens (Kiryuk *et al. Nat Genet.* 2014; 46(11):1187-96). Serum levels and B cell production of APRIL are elevated in patients with IgA nephropathy and correlate with aberrantly glycosylated IgA levels (Zhai *et al. Medicine* (Baltimore). 2016; 95(11):e3099). Plasma levels of APRIL (TNFSF13) correlate with progression of chronic kidney disease in IgA nephropathy (Han *et al. J Am Soc Nephrol.* 2016; 27(2):439-53). Treatment with anti-APRIL antibody results in reduction of serum IgA, clearing of kidney mesangium, and reduction of inflammatory cell infiltration and glomerular injury, in mice (Kim *et al. PLoS One.* 2015; 10(9):e0137044). Anti-APRIL antibody preserves immune cell homeostasis in bone marrow and spleen (Kim *et al. PLoS One.* 2015; 10(9):e0137044).

APRIL (TNFSF13) represents a logical biological and therapeutic target for the treatment of IgA nephropathy. Without wishing to be bound by theory, it is believed that in an embodiment, the efficacy of the antibody molecules described herein with respect to the targeted modulation of APRIL-mediated immunobiological mechanisms is directly relevant to treatment of IgA nephropathy. The anti-APRIL antibody molecules described herein (*e.g.*, humanized anti-APRIL antibody molecules), *e.g.*, with high biological potency and/or low complement activation, can be used to treat IgA nephropathy. In an embodiment, the antibody molecule has picomolar APRIL binding affinity and sub-nanomolar receptor blocking activity to both TACI and BCMA, *e.g.*, *in vitro*. In another embodiment, the antibody molecule functionally interfere with APRIL mediated downstream cellular signaling, *e.g.*, through the canonical

NFκB activation pathway. In an embodiment, the antibody molecule is engineered, *e.g.*, as an IgG2 subtype, for purposes of clinically mitigating against antibody-dependent exacerbation of complement recruitment, *e.g.*, in the kidneys of IgA nephropathy patients. In an embodiment, an antibody molecule described herein can have an improved safety profile in comparison to more depletive B cell-based therapeutic approaches, *e.g.*, due to a lesser perturbation of B and T cell homeostasis as shown in a murine model (Kim *et al. PLoS One.* 2015;10(9):e0137044).

The antibody molecules described herein can be used to treat or prevent different stages of IgA nephropathy. In an embodiment, the antibody molecule is used to treat a symptom associated with IgA nephropathy, *e.g.*, hematuria, proteinuria, albuminuria, hypertension, an early stage kidney disease (*e.g.*, requiring dialysis or transplantation), or a combination thereof. In an embodiment, the antibody molecule reduces aberrantly glycosylated IgA1, auto-antibody formation, deposition of nephritogenic immune complexes in the kidney, inflammation and loss of kidney function, or a combination thereof. In an embodiment, the subject is at low risk, *e.g.*, having minor urinary abnormalities (*e.g.*, micro-hematuria), normal glomerular filtration rate (GFR), and/or no hypertension. In another embodiment, the subject is at moderate to high risk, *e.g.*, having proteinuria greater than 0.5-1 g/d and/or GFR reduced (*e.g.*, below 30-50 ml/min) and/or hypertension. In yet another embodiment, the subject has acute or rapid GFR loss, *e.g.*, having nephrotic syndrome or rapidly progressive glomerulonephritis (RPGN), or acute kidney injury (AKI) due to macro-hematuria or other common cause. In an embodiment, the subject has proteinuria greater than 0.5 g/day, *e.g.*, between 0.5-1 g/day or greater than 1 g/day. In an embodiment, the subject treated for IgA nephropathy has glomerular filtration rate (GFR) less than 50 ml/min, *e.g.*, less than 30 ml/min. In an embodiment, the antibody molecule does not significantly change (*e.g.*, capable of preserving) immune cell homeostasis. In another embodiment, the antibody molecule results in a reduction of IgA not total ablation of IgA.

Diabetic Nephropathy

The antibody molecule described herein can be used to treat or prevent diabetic nephropathy. Diabetic nephropathy (or known as diabetic kidney disease) is a progressive kidney disease caused, *e.g.*, by damage to the capillaries in the kidneys' glomeruli. It is typically characterized by nephrotic syndrome and diffuse scarring of the glomeruli. It is often due to longstanding diabetes mellitus, and is a prime reason for dialysis. It is classified as a small blood vessel complication of diabetes.

Exemplary symptoms of diabetic nephropathy include, but are not limited to, severe tiredness, headaches, a general feeling of illness, nausea, vomiting, frequent voiding, lack of appetite, itchy skin, or leg swelling. The cause of diabetic nephropathy can include, *e.g.*, high blood sugar, advanced glycation end product formation. Cytokines may be involved in the development of diabetic nephropathy.

Diabetes can cause a number of changes to the body's metabolism and blood circulation, which likely combine to produce excess reactive oxygen species. These changes damage the kidney's glomeruli, which leads to the hallmark feature of albuminuria (Cao and Cooper *J Diabetes Investig.* 2011; 2(4): 243-247). As diabetic nephropathy progresses, the glomerular filtration barrier (GFB), which is composed of the fenestrated endothelium, glomerular basement membrane, and epithelial podocytes, is increasingly damaged (Mora-Fernández *et al. J. Physiol. (Lond.)* 2014; 592 (Pt 18): 3997-4012). Damage to the glomerular basement membrane allows proteins in the blood to leak through, leading to accumulation in Bowman's space as distinct periodic-acid schiff positive nodules (Kimmelstiel–Wilson nodules).

Diagnosis of diabetic nephropathy can be based on the measurement of high levels of albumin in the urine or evidence of reduced kidney function (Lewis and Maxwell *Practitioner.* 2014; 258(1768):13-7, 2). Albumin measurements can be defined as follows: normal albuminuria: urinary albumin excretion <30 mg/24h; microalbuminuria: urinary albumin excretion in the range of 30–299 mg/24h; clinical (overt) albuminuria: urinary albumin excretion ≥300 mg/24h. To test kidney function, the person's estimated glomerular filtration rate (eGFR) is measured from a blood sample. Normal eGFR ranges from 90 to 120 ml/min/1.73 m².

Other treatments that can be used in combination with the antibody molecule described herein to treat diabetic nephropathy include, *e.g.*, an angiotensin-converting enzyme (ACE) inhibitor (*e.g.*, captopril, enalapril, lisinopril, or ramipril), an angiotensin II receptor blocker (ARB) (*e.g.*, candesartan cilexetil, irbesartan, losartan, or telmisartan), a calcium channel blocker (*e.g.*, amlodipine, diltiazem, or verapamil), a diuretic (*e.g.*, chlorthalidone, hydrochlorothiazide, or spironolactone), a beta-blocker (*e.g.*, atenolol, carvedilol, or metoprolol), and diabetes management (*e.g.*, control of high blood pressure or blood sugar levels, or reduction of dietary salt intake).

Cancer

The antibody molecule described herein can be used to treat or prevent a cancer. Exemplary cancers that can be treated or prevented by the antibody molecules described herein include, but are not limited to, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical carcinoma, Kaposi sarcoma, an AIDS-related lymphoma, primary central nervous system (CNS) lymphoma, anal cancer, appendix cancer, astrocytoma, atypical teratoid/rhabdoid tumor, basal cell carcinoma, bile duct cancer, bladder cancer, bone cancer (*e.g.*, Ewing sarcoma or osteosarcoma and malignant fibrous histiocytoma), brain tumor (*e.g.*, astrocytomas, brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumor, central nervous system germ cell tumor, craniopharyngioma, or ependymoma), breast cancer, bronchial tumor, Burkitt lymphoma, carcinoid tumor (*e.g.*, gastrointestinal carcinoid tumor), cardiac (heart) tumor, embryonal

tumor, germ cell tumor, lymphoma, cervical cancer, cholangiocarcinoma, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative neoplasm, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, ductal carcinoma *in situ* (DCIS), endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, Ewing sarcoma, 5 extracranial germ cell tumor, extragonadal germ cell tumor, eye cancer (*e.g.*, intraocular melanoma or retinoblastoma), fallopian tube cancer, fibrous histiocytoma of bone, osteosarcoma, gallbladder cancer, gastric (stomach) cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumor (*e.g.*, central nervous system tumor, extracranial tumor, extragonadal tumor, ovarian cancer, or testicular cancer), gestational trophoblastic disease, glioma, hairy cell leukemia, head and neck cancer, 10 hepatocellular (liver) cancer, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumor, pancreatic neuroendocrine tumor, Kaposi sarcoma, kidney cancer (*e.g.*, renal cell cancer or Wilms tumor), Langerhans cell histiocytosis (LCH), laryngeal cancer, leukemia (*e.g.*, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), or hairy cell leukemia), lip and oral cavity cancer, liver cancer, lung 15 cancer (*e.g.*, non-small cell lung cancer (NSCLC) or small cell lung cancer), lymphoma (*e.g.*, aids-related, Burkitt lymphoma, cutaneous T-cell lymphoma, Hodgkin lymphoma, non-Hodgkin lymphoma, or primary central nervous system (CNS) lymphoma), Waldenström macroglobulinemia, male breast cancer, malignant fibrous histiocytoma of bone and osteosarcoma, melanoma (*e.g.*, intraocular (eye) melanoma), Merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer, midline tract carcinoma, mouth 20 cancer, multiple endocrine neoplasia syndrome, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndrome, myelodysplastic/myeloproliferative neoplasm, chronic myeloproliferative neoplasm, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, oral cancer, lip and oral cavity cancer, oropharyngeal cancer, osteosarcoma and malignant fibrous histiocytoma of bone, ovarian cancer (*e.g.*, epithelial ovarian cancer or germ cell ovarian tumor), 25 pancreatic cancer, pancreatic neuroendocrine tumors (islet cell tumors), papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytoma, pituitary tumor, pleuropulmonary blastoma, peritoneal cancer, prostate cancer, rectal cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, sarcoma (*e.g.*, Ewing sarcoma, Kaposi sarcoma, osteosarcoma, rhabdomyosarcoma, soft tissue sarcoma, or uterine sarcoma), Sézary syndrome, 30 skin cancer (*e.g.*, melanoma, Merkel cell carcinoma, or nonmelanoma skin cancer), small intestine cancer, squamous cell carcinoma, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, transitional cell cancer of the renal pelvis and ureter, urethral cancer, endometrial uterine cancer, vaginal cancer, vulvar cancer, or a metastatic lesion thereof.

In an embodiment, the cancer is a hematological cancer, *e.g.*, a lymphoma or leukemia, *e.g.*, chosen from B-cell non-Hodgkin's lymphoma, chronic lymphocytic leukemia (CLL), Hodgkin's lymphoma, multiple myeloma, Waldenström macroglobulinemia, or lymphoplasmacytic lymphoma. In an embodiment, the cancer is a multiple myeloma. In another embodiment, the cancer is a solid tumor,
5 *e.g.*, chosen from colorectal cancer, breast cancer (*e.g.*, breast carcinoma), esophageal cancer (*e.g.*, esophageal adenocarcinoma), brain cancer (*e.g.*, glioblastoma), or kidney cancer (*e.g.*, renal cell carcinoma).

In an embodiment, the antibody molecule is used to treat a lymphoma. Other treatments that can be used in combination with the antibody molecule described herein to treat lymphoma include, *e.g.*,
10 chemotherapy, immunotherapy, targeted drug therapy, radiation therapy, and stem cell transplant. Exemplary targeted drug therapy includes a CD20 inhibitor (*e.g.*, rituximab (RITUXAN® or ibritumomab tiuxetan (ZEVALIN®)).

In an embodiment, the antibody molecule is used to treat a leukemia. Other treatments that can be used in combination with the antibody molecule described herein to treat leukemia include, *e.g.*,
15 chemotherapy, immunotherapy, targeted drug therapy, radiation therapy, and stem cell transplant. Exemplary targeted drug therapy includes a tyrosine kinase inhibitor (*e.g.*, imatinib (GLEEVEC®)).

In an embodiment, the antibody molecule is used to treat a multiple myeloma. Other treatments that can be used in combination with the antibody molecule described herein to treat multiple myeloma include, *e.g.*, chemotherapy, corticosteroids, immunotherapy, targeted drug therapy, radiation therapy,
20 and stem cell transplant. Exemplary targeted drug therapy includes, *e.g.*, a thalidomide analog (*e.g.*, thalidomide (THALOMID®), lenalidomide (REVLIMID®), or pomalidomide (POMALYST®)).

In an embodiment, the antibody molecule is used to treat Waldenström macroglobulinemia. Other treatments that can be used in combination with the antibody molecule described herein to treat Waldenström macroglobulinemia include, *e.g.*, plasma exchange, chemotherapy, immunotherapy,
25 targeted drug therapy, and stem cell transplant.

In an embodiment, the antibody molecule is used to treat a colorectal cancer. Other treatments that can be used in combination with the antibody molecule described herein to treat colorectal cancer include, *e.g.*, surgery, chemotherapy, radiation therapy, immunotherapy, and targeted drug therapy. Exemplary targeted drug therapy includes, *e.g.*, a VEGF inhibitor (*e.g.*, bevacizumab (AVASTIN®)), an
30 EGFR inhibitor (*e.g.*, cetuximab (ERBITUX®), panitumumab (VECTIBIX®)), and dual VEGFR2-TIE2 tyrosine kinase inhibitor (*e.g.*, regorafenib (STIVARGA®)).

In an embodiment, the antibody molecule is used to treat a breast cancer, *e.g.*, a breast carcinoma. Other treatments that can be used in combination with the antibody molecule described herein to treat breast cancer include, *e.g.*, surgery, chemotherapy, radiation therapy, hormone therapy, immunotherapy,

and targeted drug therapy. Exemplary target drug therapy includes, *e.g.*, an HER2 inhibitor (*e.g.*, trastuzumab (HERCEPTIN®), pertuzumab (PERJETA®), ado-trastuzumab (KADCYLA®), or lapatinib (TYKERB®)) or a VEGF inhibitor (*e.g.*, bevacizumab (AVASTIN®)).

In an embodiment, the antibody molecule is used to treat an esophageal cancer, *e.g.*, an esophageal adenocarcinoma. Other treatments that can be used in combination with the antibody molecule described herein to treat esophageal cancer include, *e.g.*, surgery, chemotherapy, radiation therapy, and immunotherapy.

In an embodiment, the antibody molecule is used to treat a brain cancer, *e.g.*, a glioblastoma. Other treatments that can be used in combination with the antibody molecule described herein to treat brain cancer include, *e.g.*, surgery, chemotherapy, radiation therapy, radiosurgery, immunotherapy, and targeted drug therapy. Exemplary targeted drug therapy includes, *e.g.*, a VEGF inhibitor (*e.g.*, bevacizumab (AVASTIN®)).

In an embodiment, the antibody molecule is used to treat a kidney cancer, *e.g.*, a renal cell carcinoma. Other treatments that can be used in combination with the antibody molecule described herein to treat kidney cancer include, *e.g.*, surgery, cryoablation, radiofrequency ablation, radiation therapy, immunotherapy, and targeted drug therapy. Exemplary targeted drug therapy includes, *e.g.*, a VEGF inhibitor (*e.g.*, bevacizumab (AVASTIN®)), a tyrosine kinase inhibitor (*e.g.*, axitinib (INLYTA®), pazopanib (VOTRIENT®), sorafenib (NEXAVAR®), or sunitinib (SUTENT®), or an mTOR inhibitor (*e.g.*, temsirolimus (TORISEL®) or everolimus (AFINITOR®)).

Immunoproliferative Disorders

The antibody molecule described herein can be used to treat or prevent an immunoproliferative disorder. Immunoproliferative disorders (also known as immunoproliferative diseases or immunoproliferative neoplasms) are disorders of the immune system that are characterized by the abnormal proliferation of the primary cells of the immune system (*e.g.*, B cells, T cells and Natural killer (NK) cells) or by the excessive production of immunoglobulins (*e.g.*, antibodies).

Exemplary immunoproliferative disorders include, but are not limited to, lymphoproliferative disorders (LPDs), hypergammaglobulinemia, and paraproteinemia. Lymphoproliferative disorders include several conditions in which lymphocytes are produced in excessive quantities. They typically occur in patients who have compromised immune systems. Hypergammaglobulinemia is often characterized by increased levels of immunoglobulins in the blood serum. Paraproteinemia or monoclonal gammopathy is the presence of excessive amounts of a single monoclonal gammaglobulin (*e.g.*, a paraprotein) in the blood. In an embodiment, the antibody molecule is used to treat monoclonal IgA hypergammaglobulinemia.

Vasculitis

The antibody molecule described herein can be used to treat or prevent vasculitis. Vasculitis is a group of disorders that destroy blood vessels by inflammation. Vasculitis is primarily caused by leukocyte migration and resultant damage. Exemplary types of vasculitis include, but are not limited to, microscopic polyarteritis (poly-angiitis), Wegener's granulomatosis, Henoch Schonlein purpura and polyarteritis nodosa.

In an embodiment, the antibody molecule is used to treat Henoch-Schonlein purpura (IgA associated vasculitis).

Henoch-Schönlein purpura (HSP, also known as anaphylactoid purpura, purpura rheumatica, or Schönlein-Henoch purpura) is a disease of the skin and other organs that most commonly affects children. HSP is a systemic vasculitis (inflammation of blood vessels) and is characterized by deposition of immune complexes of IgA and complement component 3 (C3) on arterioles, capillaries, and venules. In the skin, the disease causes palpable purpura (small hemorrhages); often with joint and abdominal pain. With kidney involvement, there may be a loss of small amounts of blood and protein in the urine; in a small proportion of cases, the kidney involvement proceeds to chronic kidney disease even irreversible kidney damage. HSP is often preceded by an infection, such as a throat infection.

Symptoms of Henoch-Schönlein purpura include, *e.g.*, rash (purpura), swollen or sore joints (arthritis), gastrointestinal symptoms (*e.g.*, abdominal pain, nausea, vomiting or bloody stools), and kidney involvement (*e.g.*, protein or blood in the urine). Serum levels of IgA are high in HSP patients.

Standards for defining Henoch-Schönlein purpura include, *e.g.*, the 1990 American College of Rheumatology (ACR) classification (Mills *et al.* (1990). *Arthritis and Rheumatism* 33 (8): 1114-21), the 1994 Chapel Hill Consensus Conference (CHCC) (Jennette *et al.* (1994) *Arthritis and Rheumatism* 37 (2): 187-92), and the 2006 European League Against Rheumatism (EULAR) and Pediatric Rheumatology Society (PRoS) classification, which includes palpable purpura as a mandatory criterion, together with at least one of the following findings: diffuse abdominal pain, predominant IgA deposition (confirmed on skin biopsy), acute arthritis in any joint, and renal involvement (as evidenced by the presence of blood and/or protein in the urine) (Ozen *et al.* (2006) *Annals of Rheumatic Diseases* 65 (7): 936-41).

Other treatments that can be used in combination with the antibody molecule described herein to treat Henoch-Schönlein purpura include, *e.g.*, analgesics for the abdominal and joint pains, steroids (*e.g.*, oral steroids or a combination of intravenous methylprednisolone (steroid), cyclophosphamide and dipyridamole followed by prednisone). Other regimens also include, *e.g.*, steroids/azathioprine, and steroids/cyclophosphamide (with or without heparin and warfarin), or intravenous immunoglobulin (IVIG).

In another embodiment, the antibody molecule is used to treat acute proliferative glomerulonephritis, *e.g.*, post-streptococcal glomerulonephritis.

Acute proliferative glomerulonephritis is a disorder of the glomeruli (glomerulonephritis), or small blood vessels in the kidneys. It is a common complication of bacterial infections, typically skin infection by *Streptococcus* bacteria types 12, 4 and 1 (impetigo) but also after streptococcal pharyngitis, for which it is also known as postinfectious or poststreptococcal glomerulonephritis. The infection causes blood vessels in the kidneys to develop inflammation, which hampers the renal organs ability to filter urine.

The pathophysiology of this disorder is consistent with an immune complex mediated mechanism. This disorder produces proteins that have different antigenic determinants, which in turn have an affinity for sites in the glomerulus. As soon as binding occurs to the glomerulus, via interaction with properdin, complement is activated. Complement fixation causes the generation of additional inflammatory mediators.

Symptoms of acute proliferative glomerulonephritis include, *e.g.*, hematuria, oliguria, edema, hypertension, fever, headache, malaise, anorexia, and nausea.

Other treatments that can be used in combination with the antibody molecule described herein to treat acute proliferative glomerulonephritis includes, *e.g.*, blood pressure (BP) control and control of the amount of potassium in individuals with oliguric acute kidney injury.

Autoimmune Disorders

The antibody molecule described herein can be used to treat or prevent an autoimmune disorder. Exemplary autoimmune disorders that can be treated or prevented by the antibody molecule described herein include, but are not limited to, acute Disseminated Encephalomyelitis (ADEM), acute necrotizing hemorrhagic leukoencephalitis, Addison's disease, agammaglobulinemia, alopecia areata, amyloidosis, ankylosing spondylitis, anti-GBM/anti-TBM nephritis, antiphospholipid syndrome (APS), autoimmune angioedema, autoimmune aplastic anemia, autoimmune dysautonomia, autoimmune hepatitis, autoimmune hyperlipidemia, autoimmune immunodeficiency, autoimmune inner ear disease (AIED), autoimmune myocarditis, autoimmune oophoritis, autoimmune pancreatitis, autoimmune retinopathy, autoimmune thrombocytopenic purpura (ATP), autoimmune thyroid disease, autoimmune urticaria, axonal & neuronal neuropathies, Balo disease, Behcet's disease, bullous pemphigoid, cardiomyopathy, Castleman disease, celiac disease, Chagas disease, chronic fatigue syndrome, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss syndrome, cicatricial pemphigoid/benign mucosal pemphigoid, Crohn's disease, Cogan's syndrome, cold agglutinin disease, congenital heart block, coxsackie myocarditis, CREST disease, essential mixed

cryoglobulinemia, demyelinating neuropathies, dermatitis herpetiformis, dermatomyositis, Devic's
 disease (neuromyelitis optica), discoid lupus, Dressler's syndrome, endometriosis, eosinophilic
 esophagitis, eosinophilic fasciitis, Erythema nodosum, experimental allergic encephalomyelitis, Evans
 syndrome, fibromyalgia, fibrosing alveolitis, giant cell arteritis (temporal arteritis), giant cell myocarditis,
 5 glomerulonephritis, Goodpasture's syndrome, granulomatosis with polyangiitis (GPA) (formerly called
 Wegener's Granulomatosis), Graves' disease, Guillain-Barre syndrome, Hashimoto's encephalitis,
 Hashimoto's thyroiditis, hemolytic anemia, Henoch-Schonlein purpura, Herpes gestationis,
 hypogammaglobulinemia, idiopathic thrombocytopenic purpura (ITP), IgA nephropathy, IgG4-related
 sclerosing disease, immunoregulatory lipoproteins, inclusion body myositis, interstitial cystitis, juvenile
 10 arthritis, juvenile diabetes (type 1 diabetes), juvenile myositis, Kawasaki syndrome, Lambert-Eaton
 syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, linear
 IgA disease (LAD), lupus (SLE), Lyme disease, chronic, Meniere's disease, microscopic polyangiitis,
 mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, multiple sclerosis,
 myasthenia gravis, myositis, narcolepsy, neuromyelitis optica (Devic's), neutropenia, ocular cicatricial
 15 pemphigoid, optic neuritis, palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric
 Disorders Associated with Streptococcus), paraneoplastic cerebellar degeneration, paroxysmal nocturnal
 hemoglobinuria (PNH), Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis (peripheral
 uveitis), pemphigus, peripheral neuropathy, perivenous encephalomyelitis, pernicious anemia, POEMS
 syndrome, polyarteritis nodosa, Type I, II, & III autoimmune polyglandular syndromes, polymyalgia
 20 rheumatica, polymyositis, postmyocardial infarction syndrome, postpericardiotomy syndrome,
 progesterone dermatitis, primary biliary cirrhosis, primary sclerosing cholangitis, psoriasis, psoriatic
 arthritis, idiopathic pulmonary fibrosis, pyoderma gangrenosum, pure red cell aplasia, raynauds
 phenomenon, reactive arthritis, reflex sympathetic dystrophy, reiter's syndrome, relapsing polychondritis,
 restless legs syndrome, retroperitoneal fibrosis, rheumatic fever, rheumatoid arthritis, sarcoidosis,
 25 Schmidt syndrome, scleritis, scleroderma, Sjogren's syndrome, sperm & testicular autoimmunity, stiff
 person syndrome, subacute bacterial endocarditis (SBE), Susac's syndrome, sympathetic ophthalmia,
 Takayasu's arteritis, temporal arteritis/Giant cell arteritis, thrombocytopenic purpura (TTP), Tolosa-Hunt
 syndrome, transverse myelitis, Type 1 diabetes, ulcerative colitis, undifferentiated connective tissue
 disease (UCTD), uveitis, vasculitis, vesiculobullous dermatosis, vitiligo, Wegener's granulomatosis (also
 30 known as Granulomatosis with Polyangiitis (GPA)).

In an embodiment, the autoimmune disorder is rheumatoid arthritis, systemic lupus
 erythematosus, a linear IgA bullous disease (*e.g.*, linear immunoglobulin A (IgA) dermatosis), or IgA-
 mediated epidermolysis bullosa acquisita.

In an embodiment, the antibody molecule is used to treat rheumatoid arthritis. Other treatments that can be used in combination with the antibody molecule described herein to treat rheumatoid arthritis includes, *e.g.*, an NSAID, a steroid (*e.g.*, corticosteroid), a disease-modifying antirheumatic drug (DMARD) (*e.g.*, methotrexate (TREXALL®), leflunomide (ARAVA®), hydroxychloroquine (PLAQUENIL®), or sulfasalazine (AZULFIDINE®), a biologic response modifier (*e.g.*, abatacept (ORENCIA®), adalimumab (HUMIRA®), anakinra (KINERET®), certolizumab (CIMZIA®), etanercept (ENBREL®), golimumab (SIMPONI®), infliximab (REMICADE®), rituximab (RITUXAN®) and tocilizumab (ACTEMRA®), or Tofacitinib (XELJANZ®)), or surgery.

In an embodiment, the antibody molecule is used to treat systemic lupus erythematosus. Other treatments that can be used in combination with the antibody molecule described herein to treat rheumatoid arthritis includes, *e.g.*, an NSAID, an antimalarial drug (*e.g.*, hydroxychloroquine (PLAQUENIL®), corticosteroid (*e.g.*, prednisone), an immunosuppressant (*e.g.*, azathioprine (IMURAN®, AZASAN®), mycophenolate (CELLCEPT®), leflunomide (ARAVA®), or methotrexate (TREXALL®)), or a BAFF inhibitor (*e.g.*, belimumab (BENLYSTA®).

In an embodiment, the antibody molecule is used to treat a linear IgA bullous disease (*e.g.*, linear immunoglobulin A (IgA) dermatosis). Other treatments that can be used in combination with the antibody molecule described herein to treat a linear IgA bullous disease (*e.g.*, linear immunoglobulin A (IgA) dermatosis) include, *e.g.*, corticosteroids (*e.g.*, prednisone or prednisolone), an antibiotic (*e.g.*, tetracycline, erythromycin, sulfapyridine), colchicine, or mycophenolate mofetil.

In an embodiment, the antibody molecule is used to treat IgA-mediated epidermolysis bullosa acquisita. Other treatments that can be used in combination with the antibody molecule described herein to treat IgA-mediated epidermolysis bullosa acquisita includes, *e.g.*, an antibiotic, an anti-inflammatory drug (*e.g.*, corticosteroid), or surgery.

Other Disorders

The antibody molecule described herein can be used to treat or prevent other disorders, *e.g.*, IgA pemphigus, celiac disease, or alcoholic cirrhosis.

In an embodiment, the antibody molecule is used to treat or prevent IgA pemphigus. Other treatments that can be used in combination with the antibody molecule described herein to treat IgA pemphigus include, *e.g.*, corticosteroid, an immunosuppressant (*e.g.*, azathioprine (IMURAN®), methotrexate (TREXALL®), or mycophenolate mofetil (CELLCEPT®)), an CD-20 inhibitor (*e.g.*, rituximab (RITUXAN®), an antibiotic, an antiviral agent, or an antifungal agent.

In an embodiment, the antibody molecule is used to treat or prevent celiac disease. Other treatments that can be used in combination with the antibody molecule described herein to treat celiac disease include, *e.g.*, a gluten-free diet, a vitamin or mineral supplement, or a steroid.

In an embodiment, the antibody molecule is used to treat or prevent alcoholic cirrhosis. Other treatments that can be used in combination with the antibody molecule described herein to treat alcoholic cirrhosis include, *e.g.*, an immunosuppressant (*e.g.*, azathioprine, prednisone, azathioprine, cyclosporine, or methotrexate) or liver transplant.

Combination Therapies

The antibody molecules can be used in combination with other therapies. For example, the combination therapy can include an antibody molecule co-formulated with, and/or co-administered with, one or more additional therapeutic agents, *e.g.*, one or more additional therapeutic agents described herein. In other embodiments, the antibody molecules are administered in combination with other therapeutic treatment modalities, *e.g.*, other therapeutic treatment modalities described herein. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

Administered "in combination", as used herein, means that two (or more) different treatments are delivered to the subject before, or during the course of the subject's affliction with a disorder. In an embodiment, two or more treatments are delivered prophylactically, *e.g.*, before the subject has the disorder or is diagnosed with the disorder. In another embodiment, the two or more treatments are delivered after the subject has developed or diagnosed with the disorder. In some embodiments, the delivery of one treatment is still occurring when the delivery of the second begins, so that there is overlap. This is sometimes referred to herein as "simultaneous" or "concurrent delivery." In other embodiments, the delivery of one treatment ends before the delivery of the other treatment begins. In some embodiments of either case, the treatment is more effective because of combined administration. For example, the second treatment is more effective, *e.g.*, an equivalent effect is seen with less of the second treatment, or the second treatment reduces symptoms to a greater extent, than would be seen if the second treatment were administered in the absence of the first treatment, or the analogous situation is seen with the first treatment. In some embodiments, delivery is such that the reduction in a symptom, or other parameter related to the disorder is greater than what would be observed with one treatment delivered in the absence of the other. The effect of the two treatments can be partially additive, wholly additive, or greater than additive. The delivery can be such that an effect of the first treatment delivered is still detectable when the second is delivered.

In certain embodiments, the additional agent is a second antibody molecule, *e.g.*, an antibody molecule different from a first antibody molecule. Exemplary antibody molecules that can be used in combination include, but are not limited to, any combination of the antibody molecules listed in **Table 1 or 5**.

5 In an embodiment, the antibody molecule is administered in combination with a second therapy to treat or prevent IgA nephropathy.

In an embodiment, the antibody molecule is administered in combination with an angiotensin-converting-enzyme (ACE) inhibitor or an angiotensin receptor blocker (ARB).

10 In an embodiment, the antibody molecule is administered in combination with an Fc decoy receptor, *e.g.*, a soluble Fc receptor. In an embodiment, the soluble Fc receptor is a soluble Fc-gamma receptor IIB. In an embodiment, the soluble Fc receptor is SM101/BAX 1810 (Baxalta). In an embodiment, the soluble Fc receptor is administered at a dose between 1 mg/kg and 50 mg/kg, *e.g.*, between 5 mg/kg and 15 mg/kg, between 12 mg/kg and 24 mg/kg, or between 20 mg/kg and 30 mg/kg.

15 In an embodiment, the antibody molecule is administered in combination with repository corticotropin (ACTHAR®). Repository corticotropin is an adrenocorticotrophic hormone (ACTH) analogue. In an embodiment, repository corticotropin is administered at a dose between 50 U and 150 U, *e.g.*, between 80 U and 120 U, by subcutaneous injection, twice or three times a week. In an embodiment, repository corticotropin is administered at a dose of 120 U, by subcutaneous injection, *e.g.*, once, twice, or three times a week.

20 In an embodiment, the antibody molecule is administered in combination with mycophenolate mofetil (MMF). Mycophenolate mofetil is the 2-morpholinoethyl ester of mycophenolic acid (MPA), an immunosuppressive agent and inosine monophosphate dehydrogenase (IMPDH) inhibitor. In an embodiment, mycophenolate mofetil is administered at a dose of between 0.5 g and 2 g, *e.g.*, between 1 g and 1.5 g or between 1.5 g and 2 g, orally or intravenously, *e.g.*, once, twice, or three times a day.

25 In an embodiment, the antibody molecule is administered in combination with bortezomib (VELCADE®). Bortezomib, also known as [(1R)-3-methyl-1-((2S)-3-phenyl-2-[(pyrazin-2-ylcarbonyl)amino]propanoyl)amino)butyl]boronic acid, is a proteasome inhibitor. In an embodiment, bortezomib is administered at a dose at between 0.5 mg/m² and 2.5 mg/m², *e.g.*, between 1 mg/m² and 1.5 mg/m², *e.g.* every three days or every week.

30 In an embodiment, the antibody molecule is administered in combination with allopurinol (ZYLOPRIM®). Allopurinol, also known as 1*H*-pyrazolo[3,4-*d*]pyrimidin-4(2*H*)-one, is a purine analog. In an embodiment, allopurinol is administered at a dose between about 50 mg and 1000 mg, *e.g.*, between 100 mg and 600 mg or between 200 and 300 mg, orally, *e.g.*, once a day or once two days.

In an embodiment, the antibody molecule is administered in combination with prednisone and/or cyclophosphamide. In an embodiment, prednisone is administered at a dose between 0.2 mg/kg and 2 mg/kg, *e.g.*, between 0.5 mg/kg and 1 mg/kg, *e.g.*, once a day. In an embodiment, cyclophosphamide is administered at a dose between 0.2 g and 2 g, *e.g.*, between 0.5 g and 1 g, *e.g.*, once a day.

5 In an embodiment, the antibody molecule is administered in combination with rituximab (RITUXAN®). Rituximab is a chimeric anti-CD20 monoclonal antibody. In an embodiment, rituximab is administered at a dose between 100 mg/m² and 500 mg/m², *e.g.*, between 200 mg/m² and 450 mg/m² or between 300 mg/m² and 400 mg/m², intravenously, *e.g.*, once weekly, once every two weeks, once every four weeks, or once every eight weeks.

10 In an embodiment, the antibody molecule is administered in combination with blisibimod. Blisibimod, also known as A-623 or AMG 623, is a selective antagonist of B-cell activating factor (BAFF, also known as B-lymphocyte stimulator or BLyS).

In an embodiment, the antibody molecule is administered with budesonide. In an embodiment, the budesonide is NEFECON®, an oral formulation that releases budesonide.

15 In an embodiment, the antibody molecule is administered with valsartan and/or probucol. In an embodiment, valsartan is administered at a dose between 50 mg/day and 200 mg/day, *e.g.*, between 80 mg/day and 160 mg/day. In an embodiment, probucol is administered at a dose between 500 mg/day and 1000 mg/day, *e.g.*, between 700 mg/day and 800 mg/day.

In an embodiment, the antibody molecule is administered in combination with OPL-CCL2-LPM.
20 OPL-CCL2-LPM is a recombinant fusion protein comprised of the human CCL2 (monocyte chemoattractant protein-1) chemokine fused to a truncated form of the enzymatically active A1 domain of *Shigella dysenteriae* holotoxin (SA1). In an embodiment, OPL-CCL2-LPM is administered at a dose between 0.001 mg/kg and 1 mg/kg, *e.g.*, between 0.01 mg/kg and 0.5 mg/kg or 0.05 mg/kg and 0.1 mg/kg, *e.g.*, intravenously.

25 In an embodiment, the antibody molecule is administered in combination with methylprednisolone. In an embodiment, methylprednisolone is administered at a dose between 0.1 mg/kg and 2 mg/kg/day, *e.g.*, between 0.2 mg/kg/day and 1.5 mg/kg/day or 0.5 mg/kg/day and 1 mg/kg/day, *e.g.*, orally.

In an embodiment, the antibody molecule is administered in combination with sirolimus.
30 Sirolimus, also known as rapamycin, can inhibit IL-2 and other cytokines receptor-dependent signal transduction mechanisms, via action on mTOR, and thereby block activation of T and B cells. In an embodiment, sirolimus is administered at dose between 0.2 mg/day and 2 mg/day, *e.g.*, between 0.5 mg/day and 1 mg/day.

In an embodiment, the antibody molecule is administered in combination with a renin-angiotensin system (RAS) blocker. For example, the RAS blocker can be an angiotensin-converting enzyme (ACE) inhibitor or an AT1 receptor blocker (ARB). Exemplary ACE inhibitors that can be used in combination with the antibody molecule described herein include, *e.g.*, benazepril (LOTENSIN®), captopril, enalapril (VASOTEC®), fosinopril, lisinopril (ZESTRIL®), moexipril (UNIVASC®), perindopril (ACEON®), quinapril (ACCUPRIL®), ramipril (ALTACE®), or trandolapril (MAVIK®). Exemplary AT1 receptor blockers that can be used in combination with the antibody molecule described herein include, *e.g.*, candesartan (ATACAND®), eprosartan (TEVETEN®), irbesartan (AVAPRO®), losartan (COZAAR®), olmesartan (BENICAR®), telmisartan (MICARDIS®), or valsartan (DIOVAN®).

In an embodiment, the antibody molecule is administered in combination with fostamatinib. Fostamatinib is a prodrug of the active compound taminib (R-406), which is an inhibitor of the enzyme spleen tyrosine kinase (Syk). In an embodiment, fostamatinib is administered at a dose between about 50 mg and 200 mg, *e.g.*, between 100 mg and 150 mg, *e.g.*, orally, *e.g.*, every day.

In an embodiment, the antibody molecule is administered in combination with paricalcitol. In an embodiment, paricalcitol is administered at a dose between about 0.2 mg and 2 mg, *e.g.*, between 0.5 mg and 1 mg, *e.g.*, every day.

In an embodiment, the antibody molecule is administered in combination with ramipril. In an embodiment, ramipril is administered at a dose between about 0.5 mg and 5 mg, *e.g.*, between 1 mg and 4 mg or between 2 mg and 3 mg, *e.g.*, every day.

In an embodiment, the antibody molecule is administered in combination with an angiotensin-converting-enzyme (ACE) inhibitor. In an embodiment, the ACE inhibitor is enalapril (VASOTEC®).

In an embodiment, the antibody molecule is administered in combination with an immunosuppressant. In an embodiment, the immunosuppressant is tacrolimus. Tacrolimus, also known as FK-506 or fujimycin, is a macrolide calcineurin inhibitor.

In an embodiment, the antibody molecule is administered in combination with omega-3 fatty acids.

In an embodiment, the antibody molecule is administered in combination with CCX168. CCX168 is an orally administered C5aR inhibitor.

Exemplary therapies that can be used in combination with an antibody molecule or composition described herein to treat or prevent other disorders are also described in the section of "Methods of Treating or Preventing Disorders" herein.

Methods of Diagnosis

In some aspects, the present disclosure provides a diagnostic method for detecting the presence of APRIL *in vitro* (*e.g.*, in a biological sample, such as a biopsy or blood sample) or *in vivo* (*e.g.*, *in vivo* imaging in a subject). The method includes: (i) contacting the sample with an antibody molecule described herein, or administering to the subject, the antibody molecule; (optionally) (ii) contacting a reference sample, *e.g.*, a control sample (*e.g.*, a control biological sample, such as a biopsy or blood sample) or a control subject with an antibody molecule described herein; and (iii) detecting formation of a complex between the antibody molecule and APRIL in the sample or subject, or the control sample or subject, wherein a change, *e.g.*, a statistically significant change, in the formation of the complex in the sample or subject relative to the control sample or subject is indicative of the presence of APRIL in the sample. The antibody molecule can be directly or indirectly labeled with a detectable substance to facilitate detection of the bound or unbound antibody. Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials, as described above and described in more detail below.

The term “sample,” as it refers to samples used for detecting a polypeptide (*e.g.*, APRIL) or a nucleic acid encoding the polypeptide includes, but is not limited to, cells, cell lysates, proteins or membrane extracts of cells, body fluids such as blood, or tissue samples such as biopsies.

Complex formation between the antibody molecule, and APRIL, can be detected by measuring or visualizing either the antibody molecule bound to APRIL or unbound antibody molecule. Any suitable detection assays can be used, and conventional detection assays include an enzyme-linked immunosorbent assays (ELISA), a radioimmunoassay (RIA) or tissue immunohistochemistry. Alternative to labeling the antibody molecule, the presence of APRIL can be assayed in a sample by a competition immunoassay utilizing standards labeled with a detectable substance and an unlabeled antibody molecule. In this assay, the biological sample, the labeled standards and the antibody molecule are combined and the amount of labeled standard bound to the unlabeled binding molecule is determined. The amount of APRIL in the sample is inversely proportional to the amount of labeled standard bound to the antibody molecule.

The antibody molecules described herein can be used to diagnose disorders that can be treated or prevented by the antibody molecules described herein. The detection or diagnostic methods described herein can be used in combination with other methods described herein to treat or prevent a disorder described herein.

EXAMPLES

Example 1: Immunization and Selection of Anti-APRIL Antibodies

CD-1 IGS (outbred stock) mice (Charles River Laboratories), female (20-25 g weight), 7-8 weeks old were immunized intraperitoneally (i.p.) with 10 µg of recombinant, oligomeric FLAG-

ACRP30^{headless} APRIL herein referred to as FLAG-ACR-APRIL. For purposes of potentially generating a species cross reactive (mouse and human) anti-APRIL antibody response, both autologous and heterologous immunizations were carried out and comprised the use of human and/or mouse APRIL as immunogens. FLAG-tagged APRIL immunogens were formulated in a 200 µl volume consisting of 100 µl sterile PBS and 100 µl emulsified RIBI adjuvant system (Sigma Aldrich) comprised of a defined mixture of Monophosphoryl Lipid A (MPL, isolated from *Salmonella minnesota*) and synthetic trehalose dicorynomycolate (TDM, an analogue of trehalose dimycolate from the cord factor of the tubercle bacillus). 3 mice per arm were immunized twice weekly for up to four weeks. Serum titers of anti-APRIL antibodies were detected subsequently by indirect ELISA using FLAG-GCN4 APRIL, R&D Systems. In brief, 50 ng FLAG-GCN4 hAPRIL (105-250) or FLAG-GCN4 mAPRIL (96-241) in PBS were coated on Maxisorp 96-well flat bottom plates(NUNC # 439454), overnight at 4°C. Coated plates were blocked in 1 x blocking buffer containing 5% BLOTTO™ in PBS and 0.05% Tween-20 (PBST) for 1 hour at room temperature. All subsequent incubation steps were followed out with an intervening 3X wash step in PBST. Anti- APRIL antibody titers were determined from a fold-dilution of mouse sera (in PBS) initially starting at 1:1000 and followed by incubation of a 1:5000 HRP conjugated rabbit anti-mouse IgG secondary antibody(Jackson Immunoresearch Laboratories) for 1 hour at room temperature. Anti-APRIL immunoglobulin reactivity was visualized using 100 µl/well of freshly prepared TMB substrate (KPL). Colorimetric development was carried out for up to 10 minutes at room temperature before quenching enzymatic reaction by the addition of 100 µl of 1N sulfuric acid and quantification by absorbance at 450 nm. Mice with strong seropositive titers against primary immunogen (human or mouse APRIL) were boosted by tail vein injection three days prior to sacrifice, removal of spleen and isolation of splenocyte fusions. Mice with preferable species cross-reactivity from serum profiling were noted.

P3X63Ag8.653 plasmacytomas (ATCC #CRL-1580), herein referred to as P3X cells were used as source of fusion partner myelomas. Splenically-derived B cell clones were immortalized using published methods with modification. In brief, P3X cells were cultured at least 1 week prior to use and maintained in log phase to achieve a target cell density of between 6×10^5 and 1.2×10^6 cells/mL and 95% viability the day prior to subsequently performing the splenic fusion. Spleen cells were isolated from 2-3 mice per immunization arm following euthanization and cardiac puncture and collected into DMEM + 1% antibiotic (penicillin/streptomycin), followed by gently washing centrifugation (2X) to pellet tissue debris and clarify suspended splenocytes. Splenocytes were then pelleted by centrifugation for 10min at 400xg

at 4°C, and red blood cells lysed at room temperature for 5 minutes following gentle resuspension of cell pellet in 1X red blood cell lysis buffer. Splenocytes were collected by centrifugation (2X) following dilution with ice cold DMEM. P3X cells were also washed 3X in DMEM prior to fusion.

Mouse splenocytes were fused with P3X cells in fusion medium (50% PEG 1450, Sigma Aldrich)) at a 3:1 ratio in accordance with established methods. In brief, pre-warmed PEG was added gradually to pelleted mixture of splenocytes and P3X cells (37°C, with gentle resuspension) followed by gradual addition of pre-warmed DMEM. Fused cells were collected by low speed centrifugation and resuspended in hybridoma selective media (hypoxanthine-aminopterin-thymidine, Sigma Aldrich) followed by incubation at 37°C for 30 minutes. Fused cells were plated in a 96 well plate at a density of approximately 2.0×10^6 spleen cells per plate (20,000 cells per well).

Hybridoma supernatants were screened by ELISA on day 14 post-fusion as described (Example 2). In brief, supernatants from conditioned media were quantified for total IgG by bioinference using AMC anti mouse IgG quantification kit (Pall Biosciences). Supernatants from hybridoma conditioned media were normalized to 10 µg/mL when possible and assayed for APRIL reactivity to both FLAG-GCN4-hAPRIL and FLAG GCN4-mAPRIL as described. A counter screen using non-APRIL FLAG-tagged protein (FLAG-ACR30-myc-his) was also included to exclude clones with a strong immunoreactivity to either FLAG or ACRP30 specific epitopes (non-relevant epitopes present in original immunogen). APRIL positive hybridomas (human APRIL or mouse) were screened for receptor blocking activity by ELISA as described in Example 3. In brief, recombinant TACI-Fc was coated on to Maxisorp 96-well flat bottom plates at 100 ng/well in 0.1 M Carbonate-Bicarbonate Buffer (pH 9.6), overnight at 4°C. Plates were blocked with 1% BSA in 1xPBS for 1 hour at 37°C followed by 3X washing in 1X PBST (with 0.025% Tween). Recombinant FLAG-tagged APRIL (mouse or human) at a concentration of 50 ng/mL was premixed in binding buffer with supernatant from hybridoma media normalized when possible to 10 µg/mL IgG concentration. Antibody-APRIL preincubation was carried out for 1 hour at 30°C with mixing prior to adding to TACI-Fc coated plates, followed by a 1-hour incubation, likewise at 30°C. Detection of FLAG-APRIL bound to TACI-Fc was quantified using anti-FLAG M2 antibody conjugated with HRP (Sigma Aldrich) used at 1:10000 dilution as described in Example 3. APRIL immunoreactive Hybridomas which also exhibited at least 10% inhibition to either human APRIL or mouse APRIL were isolated, subcloned by limited dilution, and reassessed for APRIL binding and blocking activity and IgG titer by ELISA as described. Hybridomas with positive activity but low IgG titers were further isotyped for determination of potential IgM-producing clones. Positive hybridomas were selected for culture scale up, antibody purification and further characterization as described in Example 2.

Example 2: Purification and Characterization of Anti-APRIL Antibodies Derived from Mouse Hybridomas

Thirteen hybridoma clones from Example 1 were cultured at sequentially higher scale from 96 well plates to 24 well plates and subsequently to T150 flasks (20 mL culture volume). Prior to
 5 purification, cells were transferred out of HAT selective media into pre-defined, low Ig media. Supernatants were harvested 3-5 days after media transfer and clarified by centrifugation, followed by sterile filtration through a 0.22 μ m PES membranes (Corning). IgG titers were confirmed by Bioinframetry as described. Supernatants were diluted 1:1 with 2x Protein G binding buffer (1M glycine, 2M NaCl, pH 9.0.). Antibodies were purified by Protein G affinity chromatography using 1 mL
 10 Protein G HiTrap columns (GE Health Care) at a flow rate of 1 ml/min and as per the manufacturer's recommendations. IgG was eluted from the protein G column by lowering pH using 0.1M glycine buffer, pH 2.8 followed by immediate neutralization using 2M TRIS, pH 8.5. Purified antibodies were reformulated by dialysis in 1x PBS, pH 7.4 followed by concentration by ultrafiltration using an Ultra-30 AMICON 30kD MWCO filtration unit. Final antibody concentration was determined
 15 spectrophotometrically by NanoDrop using a generalized extinction coefficient for murine antibodies (IgG1). Antibody purity and integrity was confirmed by SDS-PAGE under both reducing and non-reducing conditions. Antibody isotype was determined using the Rapid ELISA Mouse mAb isotyping kit (Pierce/ThermoFisher Scientific) in conjunction with preliminary sequence analysis (*see* Example 3). All purified antibodies were determined to be predominantly a distribution of IgG1 and G2a; light chains
 20 were all determined to be of kappa class. A relatively smaller subset of APRIL immunoreactive antibodies derived from mice B-002/B-003 (*e.g.*, 02-009, 02-016, 046, **FIGS. 1A-1B**) were determined to be IgMs and not carried forward for purification.

Purified antibodies were further characterized for APRIL binding, APRIL species cross reactivity (human APRIL vs. mouse APRIL), and receptor blocking activity using TACI-Fc. For binding, an
 25 APRIL-based indirect ELISA was used to determine by first approximation the relative affinity of purified anti-APRIL antibodies to either human APRIL or mouse APRIL. ELISA method was as generally described above. In brief, HA-GCN4 hAPRIL (amino acid residues 105-250) and HA-GCN4 mAPRIL (amino acid residues 96-241) were coated at a density of 50ng/well. Blocking and wash steps were completed as described. Binding of anti-APRIL antibodies to APRIL was quantified using an 8
 30 point dilution of test antibody that spanned over a four log scale. Antibody binding to APRIL was detected using a rabbit anti-mouse IgG (H+L)-HRP conjugate (Jackson ImmunoResearch Laboratories). Antibody binding data was analyzed by non-linear regression analysis using a 3 parameter fit to determine max binding and apparent EC₅₀ values. The results are shown in **FIG. 3**. Human specific, anti-APRIL monoclonal antibodies h01A (described, *e.g.*, in Guadagnoli, M. *et al. Blood* 117, 6856-6865

(2011)), herein described as mAb 1313, and A019C11 (described, *e.g.*, in Jagessar, S. *et al. J Neuroimmune Pharmacol* 7:557–570 (2012)), herein referred to as mAb 0201, were used as positive controls for binding to human APRIL and for comparative purposes. Mouse specific antibody Aprry-1-1 (Adipogen) was likewise used for quantification of antibody binding to mouse APRIL.

5 Receptor blocking activity of anti-APRIL antibodies was likewise measured by ELISA using recombinant TACI or BCMA (extracellular domain(s) expressed and purified as Fc fusion proteins in a binding competition-based experiment. For this experiment, recombinant TACI-Fc or BCMA-Fc were produced in HEK 293 cells following transient transfection of these cells using the Fc expression vector pc-tPA-Fc. In brief, this vector was constructed from parental mammalian expression vector pcDNA3.1
10 (Life Technologies) using standard molecular cloning techniques. Vector was designed to include a 5' Kozak translation initiation consensus sequence followed by an N-terminal tPA signal sequence for processing and optimized secretion of recombinant protein into the media as a soluble protein. The c-termini of the extracellular APRIL binding domains of human TACI or human BCMA were fused in frame to the Fc portion of human IgG1 beginning at position E98 in CH1. DNA sequences were
15 synthesized following codon optimization for mammalian expression and initially cloned into pcDNA3.1 typically as a Bam HI-Xba cassette. Receptor variants were cloned into resultant vector as AscI-BbsI or AscI-Not I DNA fragments depending on design and cloning strategy. TACI-Fc comprises human TACI sequences 29-110 that includes both CRD1 and CRD2 domains and generally corresponds to the TACI-Fc based receptor decoy atacept. This sequence is herein described as HuTACI-Fc-001 (or more
20 commonly as TACI-Fc unless otherwise noted). A variant of TACI, likewise expressed as an Fc fusion protein only includes CRD2. TACI may include the C-terminal region (or so-called “stalk”) from amino acids 110 to approximately 166 that immediately precede the transmembrane domain. BCMA-Fc comprises the extracellular cytokine binding domain (amino acid residues 1-54) and herein described as HuBCMA-Fc-001 (or more commonly as BCMA-Fc unless otherwise noted).

25 Recombinant TACI-Fc or BCMA-Fc was coated on 96 well plates in 0.1 M carbonate-bicarbonate buffer (pH 9.6), overnight at 4°C. Plates were washed three times with 1x PBST (PBS + 0.025% Tween 20) followed by blocking with 1% BSA in 1x PBS, 37°C for 1 hour. All subsequent wash steps were carried out with 1x PBST. To assess antibody blocking, either 15 ng/mL or 50 ng/mL HA-GCN4-huAPRIL was preincubated with varying antibody concentrations ranging from 0.03–30 µg/mL in
30 binding buffer comprised of 1% BSA and 0.025% Tween-20 in 1X PBS. Preincubation was carried out at 30°C for 1 hour with mixing. Antibody-APRIL premix with 50 ng/mL APRIL or 15 ng/mL APRIL was then added to TACI-Fc coated or BCMA-Fc coated plates, respectively, and incubated for an additional hour at 30°C. HA-tagged APRIL binding to either receptor was quantified using a goat polyclonal anti-HA tag-HRP antibody (Abcam) added at a 1:10000 dilution followed by colorimetric development using

100 µl/well of freshly prepared TMB substrate (KPL) carried out for up to 30 minutes at room temperature before quenching enzymatic reaction by the addition of 1N sulfuric acid. ELISA signal was quantified by absorbance at 450 nm. The results are shown in **FIGS. 2A-2B** and **FIG. 3**. ELISA data was analyzed by non-linear regression. IC₅₀ values were calculated based on a 4 parameter fit of antibody titration curves. Where appropriate, % inhibition is calculated based on normalization of data to no antibody control (0% inhibition) vs. background (no APRIL), set as 100% inhibition. Anti-human mAb 1313 and mAb 0201 were used as positive controls and for comparative purposes. Non-blocking antibodies Aprly-1 or Aprly-5 (Enzo Biosciences) were used as negative controls. Mouse specific, blocking antibody aprily-1-1 (Adipogen) was generally used as a positive control in experiments using mouse APRIL (HA-GCN4-mAPRIL).

The functional activity of anti-APRIL antibodies was also evaluated using a cell-based receptor signaling transduction assay. In this assay, binding of APRIL to either TACI or BCMA results in receptor activation leading, in turn, to downstream activation of NF-κB, a transcription factor that ultimately mediates programmed changes in B cell gene expression and phenotype. The use of established NF-κB reporter cell lines for this purpose has been described in the literature. The use of heterologous (non-lymphoid) cell lines lacking TACI or BCMA expression but wherein TACI or BCMA can be introduced exogenously through transfection allows for controlled, receptor defined analysis of APRIL signaling and inhibition of this signal by anti-APRIL antibodies. For this purpose, the commercial 293TN-derived NF-κB reporter cell line NF-κB /293/GFP-LucTM (System Biosciences) was chosen. These cells are stably transfected with the genes for both GFP and firefly luciferase placed in tandem under the transcriptional control of a minimal CMV promoter (mCMV) and multiple copies of the NF-κB recognition element.

NF-κB /293/GFP-LucTM cells were maintained and grown in 293TN Cell growth medium (DMEM base medium supplemented with GlutaMAX and FBS) as per manufacturer's instruction. cDNA Expression plasmids pcMV6-XL4/TACI and pcMV6-XL4/BCMA (Origene) were used for transfections. These plasmids encode full-length human TACI (TNFRSF13B, accession number NM_012452) or BCMA (TNFRSF17, accession number NM_001192). 293TN reporter cells were transfected at a density of approximately 6x10⁵ cells/mL (>90% viability) using PEI-MAX as follows: For pcMV6-XL4/TACI, 10.4 ng/mL (~1 ng/well); for pcMV6-XL4/BCMA 83 ng/mL cell culture (~8 ng/well). Total plasmid concentration was held constant at 1.67 µg/mL culture volume (167 ng/well) using empty vector pcMV6-XL4 as needed to maintain constant plasmid amounts for each transfection. Transfections were scaled appropriately based on number of plates needed. 100 µl of transfection mix was transferred to each well. Plates were transferred to 37°C incubator with 5% CO₂ for 20-24 hours.

On day 2, recombinant APRIL (HA-GCN4-APRIL) was preincubated with serially diluted

antibody in complete 293TN culture media prior to addition to cells. In brief, for APRIL-mediated activation of TACI signal transduction, 40 ng/mL APRIL (2X target concentration) was mixed 1:1 with serially diluted antibody (likewise diluted in cell culture media) in a 96 well plate; for APRIL-mediated activation of BCMA signal transduction, 200 ng/mL APRIL (2X target concentration) was likewise mixed 1:1. Antibody-APRIL mix was incubated for 1 hour at 37°C with shaking. 40 µl of preincubation was added to transfected cell culture following removal of spent cell culture media. Cells were incubated at 37°C as above for an additional 20-24 hours. NF-κB-driven luciferase activity was subsequently quantified using ONE-Glo™ Luciferase kit (Promega) generally in accordance with manufacturer protocol, but with minor modifications. Relative fluorescent units (RLUs) were measured in opaque white 96 -well plates using a luminometer plate reader. The results are shown in **FIG. 4**. Data was normalized to no Ab control after correcting for assay signal background (no APRIL). IC₅₀ values were derived from a non-linear regression analysis of antibody titration curves fit to a four fit parameter. Antibodies 1313 and 0201 typically were used as positive controls. In the case of using recombinant mouse APRIL, apry1-1, was used as a positive control. Non-neutralizing antibodies anti human APRIL antibodies Aprily-1 or Aprily-5 were typically used as negative controls.

Example 3: Determination and Molecular Cloning of Anti-APRIL Immunoglobulin Sequences

VH and VL gene sequences of mouse antibodies derived from hybridoma screening were initially determined by reverse transcriptase PCR of B cell RNA using a pool of pre-defined set of mouse Ig sequence-specific primers of varying degeneracy. 5' Primer design for VH sequencing was based on a comprehensive analysis of the mouse immunoglobulin database with corresponding alignment to variable leader sequences. From this analysis, VH leader sequences were clustered (or binned based on sequence relatedness and representation of germline "families"); a unique set of primers, each predicted to anneal more specifically to these binned VH sequence families were designed and used as a cocktail in the RT-PCR reaction. 3' primers were designed to anneal in the constant region of the heavy chain. This primer set was naturally less complex and corresponded to unique sequences in CH1 that define the four known mouse IgG constant regions (IgG1, IgG2a, IgG2b and IgG3). IgM related VH sequences were amplified as above but with substitution of an IgM isotype 3' primer. Similarly, a so-called "pooled primer" RT-PCR approach was used to amplify the corresponding VL sequences from mouse hybridoma RNA. A systematic query of all known mouse VL leader sequences was likewise performed. As kappa and lambda light chains share neither the constant region nor variable region sequences, separate primer sets (kappa vs. lambda specific) were designed. 3' primers were designed based on isotype specific light chain constant region sequence (kappa vs. lambda) in a manner analogous to the one described above for heavy chain sequences. RT-PCR amplification of hybridoma gene sequences from B cell RNA was

completed using otherwise established methods. In brief, RNA was extracted from $0.5\text{-}2 \times 10^6$ cells using the RNeasy kit (Life Technologies) as per manufacturer's instructions. Cell lysis was facilitated using Qiashredder or related method for initial nucleic acid extraction. Purified RNA was quantified by UV absorbance. cDNA synthesis and subsequent PCR amplification (using Platinum Taq polymerase and primer mixes described above) were completed in tandem using Superscript III One Step RT-PCR kit (Life Technologies). PCR amplicons were purified using Qiaquick PCR clean up kit (Life Technologies) and quantified by UV absorbance at 260 and 280 nm using a Nanodrop spectrophotometer. PCR products were also analyzed by agarose gel electrophoresis to confirm predicted size and gel purified as needed. VH and VL gene sequences were determined by directly sequencing of PCR products using nested primers. Ambiguous sequence data was followed by re-amplification of cell RNA by RT PCR as described above but with modification to protocol and using a subset of smaller pooled primer sets; if necessary PCR products were cloned by TA cloning into an intermediate vector) and transformed into chemically competent TOP10 (Life Technologies) or DH5 α (New England Biolabs) as per the manufacturers protocols. DNA sequence data was analyzed using publically available databases (*e.g.*, International Immunogenetics Information system (IMGT), VBase, or NCBI Ig-Blast) to evaluate germline usage, identify CDR sequences and assign putative isotype when possible. In general, this sequencing strategy led to the identification of unique VH sequences for each hybridoma of interest; several clones resulted in the identification of multiple light chains.

Productive VH and VL Ig sequences were amplified by PCR and cloned separately into mammalian Ig expression vectors o-pcMG2 and o-pcMK2, respectively, for recombinant production in HEK293 cells as paired mouse IgG2 (HC) and kappa (LC) isotyped antibodies. Gene-specific primers were designed based on VH and VL sequences identified as described above. Primer design included 18-23 overlapping nucleotides complementary to the corresponding framework regions of the variable gene sequences in addition to vector complementary sequences designed to enable recombination based cloning by modified Gibson assembly fused in frame to variable region sequences through DNA ligation as described below. Primer design was assisted through the use of NEB Builder (New England Biolabs) or Primer3 software. Additional Primer design included the incorporation of restriction endonuclease recognition sequences on respective 5' ends for subcloning as needed. In the case of ambiguous variable sequences, primer design incorporated the use of modestly degenerate nucleotide sequences (at 1-2 positions) or surrogate ("best guess") codons guided by the knowledge of the predicted germline framework identified in the original VH and VL sequence analysis.

For molecular cloning by RT PCR, RNA was extracted from hybridomas generally as described. cDNA first strand was synthesized using Superscript III First Strand Synthesis Supermix (Life Technologies) as per the manufacturer's protocol. 2.5 μ l of cDNA template DNA was amplified by PCR

using Q5 high fidelity DNA polymerase (New England BioLabs). Amplification included a total of 35 cycles by “touch-up” PCR” using initially a three step amplification for 10 cycles followed by 25 cycles involving 2 step amplification (annealing and extension both at 72°C). PCR amplicons were evaluated by agarose gel electrophoresis to confirm purity, correct size, and approximate amounts. Gel purification of PCR products was carried when necessary using Qiaquick gel purification kit (Qiagen). HC and LC Ig expression vectors were linearized and prepared for cloning by restriction endonuclease double digestion. 5' ends of digested vectors were subsequently dephosphorylated with shrimp alkaline phosphatase followed by heat inactivation of enzymes. PCR products were ligated into 50 ng linearized vector (gel purified) using NEB Builder (New England Biolabs) at a 2:1 mole ratio of insert: vector. 2 µl of ligation reaction was transformed into chemically competent *E. coli* (DH5α, New England Biolabs) and plated on LB with antibiotic. Recombinant clones were selected by colony DNA sequencing followed by plasmid purification of positive clones at 200 mL *E. coli* scale using low endotoxin Purelink Maxiprep kits (Life Technologies) as per the manufacturer’s protocols.

For recombinant antibody production, approximately 225 µg each of purified Ig expression vectors (HC and LC) were used to transiently transfect HEK 293F cells. About 2×10^6 cells/mL cells were transfected using PEI-Max-based transfection reagent and cultured in Freestyle 293 cell media for a total of 5-7 days. Antibody titer was quantified by bioinferometry using Protein A-immobilized biosensors (Pall Biosensors).

Recombinant antibodies were purified from culture supernatant following clarification by low speed centrifugation and sterile filtration through 0.22 µm PES membranes followed by addition of a protease inhibitor cocktail (Cocktail III, Thermofisher) to mitigate against proteolysis. Antibodies were purified by Protein A affinity chromatography using a 1 mL Protein A Hitrap column (GE Healthcare) with low pH elution as described followed by neutralization by TRIS, pH 8.5.. Antibodies were reformulated and concentrated in 1x PBS, pH 7.4 by tandem dialysis and ultrafiltration using Amicon-Ultra 30 centrifugation concentrating units

Example 4: *In Vitro* Binding and Receptor Blocking Activities of Anti-APRIL Antibodies

Recombinant anti-APRIL antibodies were characterized with respect to both APRIL binding and receptor blocking activities using both ELISA-based and cell-signaling methods as described. As shown in FIGS. 5-6, first-pass antibody binding to human APRIL by indirect ELISA indicated several antibodies with low and subnanomolar binding affinities to the cytokine target based on EC₅₀ values extrapolated from nonlinear regression analyses of antibody titration curves. Antibodies 2218, 2419, and 2621 in particular bound human APRIL with apparent target binding affinities of less than 0.2 nM; antibodies 3530 and 2922 bound human APRIL with apparent target binding affinities between 0.2 nM

and 1 nM. 3125 bound human APRIL with an apparently lower affinity (>1 nM). Similar analysis of monoclonal antibody binding to mouse APRIL homologue indicated only mAb 3530 having appreciable cross-species binding to target (data not shown). Functional analysis of antibody blocking activity was evaluated by competition ELISA using receptor-Fc as the APRIL capture ligand in a 96 well based format as described. This analysis included the use of both biologically relevant APRIL TNFR-related receptors (human TACI and human BCMA) for purposes of evaluating any selectivity with respect to the antibody-mediated antagonism of APRIL-receptor interactions. As shown in **FIG. 9**, antibody IC₅₀ values were calculated from a non-linear analysis of antibody titration curves. Anti-human blocking antibodies 1313 and 0201 were used as positive controls and for comparative purposes. Non-neutralizing antibody April-5 (Enzo Biosciences) was used as a negative control. Based on this *in vitro* data, all of the recombinant antibodies demonstrated at least partial blocking of human APRIL to human TACI-Fc. As shown in **FIG. 7**, monoclonal antibodies 2218, 2419, 2621, 3327, and 3530 blocked APRIL-TACI binding with corresponding IC₅₀ values in the low or sub-nanomolar range. MAbs 3125 and 2922 would appear to block with somewhat lower potency, with IC₅₀ values greater than 10 nM. As shown in **FIG. 8**, a similar evaluation of receptor blocking using BCMA-Fc indicates antibodies 2218, 2419, and 3327 are likewise able to block BCMA binding with low or subnanomolar potency. MAb 3530 would appear to be relatively selective with respect to blocking APRIL binding to TACI-Fc in comparison to BCMA-Fc; as such this antibody may be viewed as being "TACI selective." MAb 2621 did not appear to block APRIL binding to BCMA-Fc as evaluated in this assay; as such, mAb 2621 may be viewed as being a potentially "TACI-specific" antibody.

Functional (receptor antagonism) activity of anti-APRIL antibodies was further evaluated using an orthogonal, cell-based NF- κ B transcriptional reporter assay to assess inhibition of APRIL-mediated receptor signal activation. This assay involved the heterologous expression of full-length (transmembrane) TACI or BCMA by transient plasmid transfection in an engineered HEK293 cell line possessing a stably transfected NF- κ B –transcriptionally activated luciferase reporter gene. Assay was generally carried out as described using recombinant Hu APRIL as the exogenous source of receptor activating cytokine. Data was normalized to the minus antibody control after subtraction of signal background (no APRIL or Ab). The data are summarized in **FIGS. 11A-11B**. Based on these data, the potent receptor blocking activities of monoclonal antibodies 2218 and 2419 were further confirmed in an antibody-dose dependent manner. These activities included blocking of both BCMA and TACI receptors. Monoclonal antibodies 2922 and 3125 qualitatively exhibited lesser activity, consistent with their relatively lower activities in the Receptor-Fc blocking ELISA (*i.e.*, in comparison to mAbs 2218 and 2419). Apparent discrepancies between these two assays may be attributed to differential receptor expression levels, protein turnover, or other biological factors not present in the less complex ELISA

based binding assays. Nevertheless, these data, taken collectively, demonstrate clear functional activity of the recombinant anti-APRIL antibodies with respect to antagonism of APRIL activity described herein.

Additional experimental data include, for example, the following. Binding of exemplary anti-APRIL antibodies to human APRIL is shown in **FIG. 16**. Relative binding affinities of exemplary anti-APRIL antibodies are shown in **FIG. 17**. Antibody inhibition of APRIL-mediated receptor signaling is shown in **FIGS. 18A-18B**. Antibody inhibition of APRIL binding to TNFSF receptors TACI and BCMA is shown in **FIGS. 19A-19B**. Antibody inhibition of APRIL binding to both human TACI-Fc and human BCMA-Fc is summarized in the table in **FIG. 20**.

Example 5: Species Cross-Reactivity of Anti-APRIL Antibodies

In addition to demonstrating the functional activity of several anti-APRIL antibodies, the cross-reactivity of these antibodies with respect to binding and blocking both mouse and human APRIL was also evaluated. This characterization employed the same set of *in vitro* assays as described but with the inclusion of analogously HA-tagged mouse APRIL (R&D Systems). For illustrative purposes, a subset of data from the receptor-Fc based blocking ELISA (both TACI-Fc and BCMA-Fc) is included. In this analysis, mAb 3530 was compared to antibodies 1313 (human specific anti-APRIL blocking antibody), Apry-1-1 (mouse specific anti-APRIL blocking antibody) and non-neutralizing antibody Aprly-5 (negative control). Antibody-titered receptor blocking activity data are summarized in **FIGS. 10A-10B** for blocking APRIL binding to human TACI-Fc and human BCMA-Fc, respectively. Consistent with other data, mAb 3530 would appear to be a "TACI-selective" antibody based on its relative neutralization profile (TACI vs. BCMA). Moreover, this antibody was able to apparently block both mouse and APRIL binding to TACI-Fc with comparable potency (apparent IC_{50} values). To our knowledge, this the first example of a species cross reactive anti-APRIL antibody with both implications with respect to epitope specificity as well as use in preclinical development of this antibody using disease relevant (syngeneic) rodent models.

APRIL species cross blocking activities of anti-APRIL antibodies are also shown in **FIGS. 21A-21B**.

Example 6: Functional Activity of Anti-APRIL Antibody for Reduction of Serum IgA *In Vivo*

In addition to its role in promotion of tumor growth, the cytokine APRIL plays several critical roles in the regulation of adaptive and mucosal immunity vis-à-vis the modulation of B and T cell function. This immunological activity includes induction of IgA production in B cells through receptor-mediated induction of Ig class switching, B cell proliferation, and survival of IgA+ related plasma cells. This central role of APRIL in IgA production leads to its potential as a therapeutic target for diseases

involving the dysregulated production of IgA and/or formation of IgA-containing immune complexes. Such diseases would include but not limited to IgA nephropathy, IgA-related vasculitis (*e.g.*, Henoch-Schodein purpura), SLE, IgA-related monoclonal gammopathies, alcoholic liver disease, etc. The biological potency of anti-APRIL therapeutic antibody can therefore be evaluated *in vivo* based directly on a reduction of serum IgA levels following the administration of such an antibody in a laboratory rodent model. Toward this end, the mouse-APRIL specific, blocking antibody Apry-1-1 (Adipogen) used as described herein as a control for assessing anti-APRIL antibody activity *in vitro* was also used as a test article. Age-matched male B6C3F1 mice (6-10 weeks old) were dosed with 20 mg/kg antibody two times a week via IP injection.12 for 8 weeks. Saline for injection was used as the negative (vehicle) control. Serum Isotype specific immunoglobulin levels (IgG, IgM, and IgA) were monitored individually by ELISA approximately every 12 days. Body weights were monitored 3x weekly, hematology and serum chemistries, and general health of the animals was monitored on a regular basis. Blood was drawn prior to dosing (day 0, pre-bleeds). Survival bleeds occurred at the end of 2 weeks (end of phase I) at 4 weeks, and 6 weeks. Terminal bleeds occurred at the end of 8 weeks. The results are shown in **FIGS. 12A-12B**.

In this study, chronic administration of a functional anti-APRIL antibody with *in vitro* validated blocking activity led to a reduction in serum IgA levels below 50%. Reduction was observed by day 24 and was sustained over the course of antibody treatment. Treatment with the mouse anti-APRIL antibody had no statistically differential effects on total Ig serum levels relative to the control; hematology and blood chemistry likewise were not affected.

Example 7: Epitope Mapping

Human APRIL site-directed variants were used for epitope mapping. As shown in **FIG. 22**, APRIL is depicted as a trimer (cyan, green, and magenta). Typical epitope containing CRD2 high affinity receptor binding site is depicted in dark blue. Positions of amino acid changes are noted with wildtype amino acid preceding number and mutation following (*e.g.*, R233G represents mutation of arginine at position 233 to glycine).

Epitope mapping of exemplary antibodies 4035, 2419, and 3833 was performed. Antibody binding to human and mouse variants was assessed by ELISA. Comparison is made to reference antibody 1313. FLAG-tagged APRIL was captured from cell culture media using anti-FLAG antibody. The results are shown in **FIGS. 23A-23B, 24A-24B, and 25A-25B**. Differentiated epitope mapping of exemplary anti-APRIL antibodies was performed by site directed mutagenesis. Primary characterization of human APRIL binding site was carried out by site-directed mutagenesis of select amino acid positions within APRIL followed by evaluation of antibody binding to these variants by ELISA. Exemplary data for three anti-APRIL antibodies 4035 (solid circles), 2419 (solid squares), and 1313 (open triangles) are

shown in **FIG. 26** for illustrative purposes.

Differential binding profiles of exemplary anti-APRIL antibodies are shown in **Table 6**.

Exemplary amino acid residues that bind to the anti-APRIL antibody molecules described herein are

listed. One or more of these residues form at least part of a binding region on APRIL. Exemplary

5 residues within the binding region predicted to have an impact on binding (*e.g.*, based on the site directed mutagenesis studies described herein) are noted by an “X”. Exemplary residues predicted to have lesser or no impact on binding (*e.g.*, based on the site directed mutagenesis studies described herein) are noted by an “N” for comparative purposes.

10 **Table 6.** Differential Binding Profiles of Exemplary Anti-APRIL Antibodies (Table discloses SEQ ID NOS: 338-341, respectively, in order of appearance)

		1313		4035		3833		2419	
		Impact	No impact	Impact	No impact	Impact	No impact	Impact	No impact
AA position	AA								
129	Asp								
130	Asp								
131	Ser								
132	Asp	X							
170	Leu								
174	Val		N	X		X			
175	Thr								
176	Phe	X		X			N	X	
177	Thr								
178	Met								
179	Gly								
180	Gln								
181	Val	X			N		N	X	
190	Gln	X		X			N	X	
192	Thr								
195	Arg	X		X					
196	Cys								
197	Ile								
200	Met								
201	Pro								
202	Ser								
203	His								

205	Asp								
206	Arg		N	X		X			
208	Tyr	X		X		X			N
226	Ser				N				
228	Ile		N		N			X	
230	Pro								
231	Arg								
232	Ala								
233	Arg								
237	Asn	X			N		N		N
241	His								

Example 8: Humanization of Mouse Derived Anti-APRIL Antibodies

Select anti-APRIL antibodies were derived from mouse immunization as described herein. The variable regions of select antibodies, namely 2419, 4035, and 4540 were subsequently humanized for purposes of potential therapeutic use and mitigation of immunogenicity. In brief, humanization was performed by identifying human germlines proximal to the mouse variable heavy (VH) and variable light (VL). Once identified, the complementarity determining regions (CDRs) from mAb2419 VH and VL were grafted on the human VH and VL germline templates respectively using structure-guided design. Additional mutations (including back mutations to the parental residue in the mouse mAb) were selectively introduced based on visual inspection of the structural model. The humanized antibody constructs were recombinantly produced in HEK293 cells following transient transfection of separate vectors for heavy and light chain expression. Recombinant antibodies were purified by Protein A affinity chromatography using standard methods. Humanized antibodies were subsequently tested for binding to APRIL, functional receptor blocking (TACI-APRIL and BCMA-APRIL interactions) and thermal stability using a differential scanning fluorescence protein unfolding assay. Relative activity and stability profiles were compared to parental, mouse antibodies upon which humanization was based.

Example 9: *In Vitro* Binding and Receptor Blocking Activities of Humanized Anti-APRIL Antibodies

Relative binding of exemplary anti-APRIL antibodies was measured by indirect ELISA. The binding of humanized variants of mouse antibodies 4035 and 2419 to human APRIL is shown in **FIG. 27A-27B**, respectively. The binding of a humanized variant of human-mouse cross-reactive antibody 4540-063 to both human APRIL and mouse APRIL is shown in **FIGS. 28A-28B**, respectively. Comparison of humanized anti-APRIL antibodies to parental (non-humanized) mouse antibodies is made

for comparative purposes. Extrapolated EC₅₀ values are summarized in **FIGS. 29A-29B**. The exemplary antibody molecules bind to APRIL with picomolar affinity.

The apparent binding affinities of antibodies 2419-1406 and 4035-062 to trimeric human APRIL were also measured by ELISA. APRIL trimer was stabilized by the N-terminal fusion of APRIL with an isoleucine zipper (GCN4) oligomerization domain. Binding of APRIL to human TACI-Fc is shown for comparative purposes. The ELISA binding results are shown in **FIG. 29C**. Picomolar EC₅₀ values derived from the binding curves depicted in **FIG. 29C** are summarized in **FIG. 29D**. The antibody showed picomolar (pM) binding to trimeric human APRIL as measured by ELISA. Higher affinity binding of 2419-1406 and 4035-062 to APRIL relative to APRIL binding to its own receptor (TACI-Fc) was observed.

The inhibition of APRIL binding to TNFSF receptors TACI and BCMA by humanized IgG2κ variants of parental, murine derived antibody 2419 was measured. Assay is based on receptor blocking ELISA using recombinant human APRIL (R&D Systems) and either human TACI-Fc (**FIG. 30A**) or BCMA-Fc (**FIG. 30B**). Parental, chimeric 2419 (mouse VH-VL grafted on to human IgG1κ constant regions) is included for comparative purposes as is chimeric anti-human APRIL antibody 4035. Inhibition was analyzed by non-linear regression using a four parameter curve fit following normalization to 100% activity (no antibody control). IC₅₀ values are summarized in **FIG. 33**.

The inhibition of APRIL binding to TNFSF receptors TACI (**FIG. 31A**) and BCMA (**FIG. 31B**) by additional humanized variants of 2419 (IgG2κ) 2419-0205 and 2419-1406 and humanized variant (4035-062) of parental antibody 4035 was measured. Humanized 4035-062 is of the IgG1κ subtype. Chimeric, non-humanized versions of mouse derived antibodies 4035 and 2419, and 1313 are included for comparative purposes. mAb1313 is a control anti-APRIL antibody. TACI-Fc and BCMA-Fc were used. IC₅₀ values are summarized in **FIG. 33**.

The inhibition of APRIL binding to TNFSF receptors TACI (**FIG. 32A**) and BCMA (**FIG. 32B**) by humanized variants of mouse/human APRIL cross-neutralizing antibody 4540 was measured. Humanized antibody 4540 is of the IgG1κ subtype. Parental 4540 (non-humanized chimera) and humanized 4035-062 (**FIGS. 30A-30B**) are included for comparative purposes. Inhibition was analyzed by non-linear regression using a four parameter curve fit as described for **FIGS. 29A-29B** and **FIGS. 30A-30B**. IC₅₀ values are also summarized in **FIG. 33**. Sub-nanomolar blocking of APRIL-receptor binding was observed for most of the tested antibodies.

These results indicate that humanization and reformatting (*e.g.*, as IgG2) generally, if not always, leads to comparable retention of receptor blocking activities.

The antibody inhibition of APRIL-mediated receptor signaling was evaluated. Inhibition of APRIL-receptor mediated NFκB intracellular signaling was evaluated using the HEK 293 NFκB reporter

cell line following transient transfection of either full-length human TNF family receptors TACI or BCMA cDNA expression vectors. Data are normalized to no antibody control (100%). The inhibition of TACI- and BCMA-mediated NF κ B signaling is shown in **FIGS. 34A-34B**, respectively.

FIG. 35 depicts the approximate IC₅₀ values of antibody inhibition of APRIL-mediated receptor signaling. Data are extrapolated from **FIGS. 34A-34B** based on a non-linear regression analysis using a variable slope, three parameter fit of antibody concentration vs. response. Negative antibody control (no APRIL binding) demonstrated no activity in this assay (data not shown). These results indicate potent inhibition of APRIL-mediated NF κ B downstream signaling pathway (with low or sub-nM IC₅₀) by exemplary anti-APRIL antibody molecule. Blocking occurs with both TACI and BCMA receptors.

Example 10: Evaluation of Anti-APRIL Antibody Binding Specificity to APRIL

Selected anti-APRIL antibodies were evaluated for potential cross-reactivity with other members of the TNF α superfamily (TNFSF) family of cytokines, including: human TNF α (TNFSF2; Adipogen), human CD40 (TNFSF4, Adipogen), human FasL (TNFSF6, Adipogen), human TRAIL (TNFSF10, Adipogen), human RANKL (TNFSF11; Adipogen), human Tweak (TNFSF12; Abcam), human and mouse BAFF (TNFSF13B, AB Biosciences and Adipogen, respectively), and human LIGHT (TNFSF14, Adipogen). These cytokines share variable degree of sequence as well as higher order structural homologies. Among them, BAFF appears to be the most closely related to APRIL, with 29% identity and 53% amino acid sequence similarity.

Binding was evaluated by an indirect ELISA protocol using similar methods described for evaluation of APRIL binding; cytokines were immobilized to ELISA plate at 50 ng per well and then exposed to a solution of each of the test antibodies at a fixed concentration of 10 μ g/ml Anti-human or anti-mouse Ig- HRP polyclonal antibody conjugates (Jackson Laboratories) were used for detection of antibody binding. The results presented in **FIG. 36** indicated eleven of the twelve antibodies tested in this assay to specifically bind APRIL with no measurable cross-reactivity with the other TNFSF members substantially above assay background. Antibody 4338 demonstrated detectable binding to both the human and mouse versions of BAFF and was used as an assay control.

mAbs 2419-1406 and 4035-062 demonstrate minimal or no extraneous protein cross-reactivity in extensive array of over 4500 human membrane proteins (RETROGENIX™). An example of protein expression array design is shown in **FIG. 37A**. Confirmatory/secondary screen was performed on 12 proteins (**FIGS. 37B-37C**). Binding to Fc receptors was observed for both 2419-1406 and 4035-062 as predicted. Weaker binding to Fc γ R1 was observed for antibody 2419-1406 consistent with it having an IgG2 isotype. Negligible binding to PDGFR was also detected for 2419-1406. Binding to any other membrane targets except those described was not observed Antibody 4035-062

Example 11: Identification of Target Epitope for Maximal Anti-APRIL Antibody Potency

The epitope of an exemplary anti-APRIL antibody molecule, mAb 2419, was mapped using a combination of empirical and computational tools and data. These methods and resultant data included 1) low resolution crystallography of mAb 2419 Fab in complex with human APRIL (amino acids 115-250) in tandem with 2) structural modeling (**FIG. 38A**), and 3) APRIL saturation mutagenesis at select positions within the surface of APRIL carried out in combination with APRIL surface display in yeast (**FIG 38B**). As shown in both **FIGS. 38A-38B**, the antibody molecule targets a non-linear, quaternary epitope spanning two different monomers of APRIL within a larger trimeric complex. The epitope of 2419 also substantially overlaps with a region of APRIL corresponding to the high affinity receptor binding site (CRD2 site) critical for both TACI and BCMA receptor blocking (**FIG. 38B**). A secondary receptor binding site (CRD1 site) also overlaps with the 2419 epitope with implications for blocking TACI-APRIL interactions. Based on this analysis, positions within APRIL that define the epitope of 2419 include V133, V181, E185, Q187, G188, R189, Q190, E191, T192, R195, H218, L219, H220, S226, I228, P230 (located in monomer A); V121, I123, Q139, P140, A141, L142, N237, S239, P240, and H241 (located in monomer B).

A more focused analysis of this epitope by mutagenesis of select surface-accessible positions of APRIL point to a subset of positions within the larger epitope of 2419 (structurally depicted in **FIG. 38A**) that appear to particularly critical for antibody binding. These so-called “hotspot residues”, *i.e.*, those residues that are critical to the interaction between APRIL and mAb 2419, are empirically defined) using a combination of the methods described above) as those positions where mutation from the wildtype sequence to nearly any other amino residue resulted in a substantial loss of binding of 2419 to APRIL. Examples of such positions include V181, Q190, T192, and I228 on one monomer, and A141 and H241 on an adjacent monomer.

Table 8. Exemplary Human APRIL Amino Acid Residues that Bind to mAb 2419 (amino acid numbering based on SEQ ID NO: 85)

Position	Monomer	Hotspot
Human		
V133	A	
V181	A	Y
E185	A	
Q187	A	
G188	A	
R189	A	

Q190	A	Y
E191	A	
T192	A	Y
R195	A	
H218	A	
L219	A	
H220	A	
S226	A	
I228	A	Y
P230	A	
V121	B	
I123	B	
Q139	B	
P140	B	
A141	B	Y
L142	B	
N237	B	
S239	B	
P240	B	
H241	B	Y

In summary, the epitope overlaps predicted regions for maximal receptor blocking, and targets a non-linear, quaternary epitope spanning two different monomers of APRIL, implications for neutralizing biologically most active form of APRIL (trimer).

5

Example 12: Pharmaceutical Properties of Anti-APRIL Antibodies

Thermal stability of exemplary anti-APRIL molecules, 2419-1406 and 4035-062, were measured using Sypro-Orange® fluorescence scanning assay (**FIG. 39A**). The thermal melting temperatures (T_m values) for both 2419-1406 and 4035-062 are listed in **FIG. 39B**. mAb 2419-1406 and mAb 4035-062 exhibit high thermal stability.

10

The tg32 mouse model was used as a surrogate for predicting the PK of antibody pharmacokinetics (PK) in humans. PK of control antibody (IgG1) with pre-established PK of approximately 25 days (in humans) was also evaluated in this study for comparative purposes. As shown in **FIGS. 40A-40B**, favorable PK profile of an exemplary anti-APRIL antibody molecule, 2419-1406 and 4035-062, in humanized FcRn transgenic mouse strain tg32 was observed.

15

Example 13: Species Cross-Reactive Anti-APRIL Antibodies Effectively Reduce Serum IgA Levels in Mice

Mouse mAb 4540 is a cross-species reactive (mouse and human) anti-APRIL with receptor neutralizing activity. mAb 4540 targets an overlapping APRIL epitope to mAb 2419-1406. mAb4540 was used as a surrogate to evaluate the effect of anti-APRIL mAb treatment on a reduction of serum IgA levels in C57/BL6 mice. 11 week old mice were sub-chronically dosed (1x weekly by i.p injection) with 20 mg/kg of mAb 4540, mAb 3833, or isotype control antibody for seven weeks. Basal serum levels of total IgA were quantified by ELISA. As shown in **FIG. 41**, treatment of mAb 4540 or mAb 3833 reduced serum IgA levels in C57/BL6 mice.

INCORPORATION BY REFERENCE

All publications, patents, and Accession numbers mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

EQUIVALENTS

While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification and the claims below. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

CLAIMS

1. An anti-APRIL antibody molecule, comprising a heavy chain variable region (VH) comprising the amino acid sequence of SEQ ID NO: 296, and a light chain variable region (VL) comprising the amino acid sequence of SEQ ID NO: 286.

5

2. The antibody molecule of claim 1, which is a synthetic or isolated antibody molecule, and/or is a humanized antibody molecule.

3. The antibody molecule of any of claims 1-2, which comprises a heavy chain constant region of IgG1, IgG2, IgG3, or IgG4, and/or comprises a light chain constant region of kappa or lambda light chain.

10

4. The antibody molecule of any of claims 1-3, which comprises a heavy chain constant region of IgG2.

15

5. The antibody molecule of any of claims 1-4, which comprises an Fc region.

6. The antibody molecule of any of claims 1-5, which comprises two heavy chain variable regions and two light chain variable regions, or is a Fab, F(ab')₂, Fv, Fd, or a single chain Fv fragment (scFv).

20

7. A pharmaceutical composition comprising an antibody molecule of any of claims 1-6 and a pharmaceutically acceptable carrier.

25

8. A nucleic acid molecule encoding a heavy chain variable region (VH) and a light chain variable region (VL) of an antibody molecule of any of claims 1-6.

9. A vector comprising a nucleic acid molecule of claim 8.

30

10. An isolated cell comprising (i) a nucleic acid molecule of claim 8, (ii) a nucleic acid molecule encoding a heavy chain variable region (VH) and a nucleic acid molecule encoding a

light chain variable region (VL), of an antibody molecule of any of claims 1-6, or (iii) a vector of claim 9.

11. A kit comprising an antibody molecule of any of claims 1-6 and instructions to use of the antibody molecule.

12. A container comprising an antibody molecule of any of claims 1-6, or a composition of claim 7.

13. A method of producing an anti-APRIL antibody molecule, the method comprising culturing a cell of claim 10 under conditions that allow production of an antibody molecule, thereby producing the antibody molecule, optionally, further comprising isolating or purifying the antibody molecule.

14. Use of an antibody molecule of any of claims 1-6, or a composition of claim 7, for the manufacture of a medicament for treating IgA nephropathy in a subject.

15. The use of claim 14, wherein the antibody molecule is to be administered to the subject intravenously or subcutaneously, and/or is to be administered to the subject at a dose between 0.1 mg/kg and 50 mg/kg, between 0.2 mg/kg and 25 mg/kg, between 0.5 mg/kg and 10 mg/kg, between 0.5 mg/kg and 5 mg/kg, between 0.5 mg/kg and 3 mg/kg, between 0.5 mg/kg and 2.5 mg/kg, between 0.5 mg/kg and 2 mg/kg, between 0.5 mg/kg and 1.5 mg/kg, between 0.5 mg/kg and 1 mg/kg, between 1 mg/kg and 1.5 mg/kg, between 1 mg/kg and 2 mg/kg, between 1 mg/kg and 2.5 mg/kg, between 1 mg/kg and 3 mg/kg, between 1 mg/kg and 10 mg/kg, or between 1 mg/kg and 5 mg/kg, or 1 mg/kg, 2 mg/kg, 4 mg/kg, or 8 mg/kg, or

at a fixed dose between 10 mg and 1000 mg, between 10 mg and 500 mg, between 10 mg and 250 mg, between 10 mg and 150 mg, between 10 mg and 100 mg, between 10 mg and 50 mg, between 250 mg and 500 mg, between 150 mg and 500 mg, between 100 mg and 500 mg, between 50 mg and 500 mg, between 25 mg and 250 mg, between 50 mg and 150 mg, between 50 mg and 100 mg, between 100 mg and 150 mg, between 100 mg and 200 mg, or between 150 mg and 250 mg,

optionally, wherein the antibody molecule is to be administered once a week, twice a week, once every two weeks, or once every four weeks, or once a month, once every two months, or once every three months.

- 5 16. The use of claim 14 or 15, wherein administration of the antibody molecule:
- (i) reduces the level of IgA in a peripheral tissue, optionally, wherein the peripheral tissue is serum, mucosal tissue, bone marrow, or any combination thereof, and/or
- (ii) reduces the level of a variant of IgA, optionally, wherein the variant of IgA is IgA1, IgA1 in polymeric form (pIgA1), IgA1 with an O-linked glycosylation variant, or any
- 10 combination thereof.

17. The use of any of claims 14-16, wherein the level of IgA is determined in a peripheral tissue sample from the subject, optionally, wherein the peripheral tissue is serum, mucosal tissue, or bone marrow.

- 15
18. A method of detecting an APRIL molecule, the method comprising contacting a cell or a sample from a subject with an antibody molecule of any of claims 1-6, thereby detecting the APRIL molecule, wherein the APRIL molecule is detected *in vitro* or *ex vivo*, optionally, wherein the antibody molecule is coupled with a detectable label.