PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :
A61K 9/26, 31/4402, A61P 1/08

A1

(11) International Publication Number: WO 00/53162

(43) International Publication Date: 14 September 2000 (14.09.00)

(21) International Application Number: PCT/EP00/01642

(22) International Filing Date: 28 February 2000 (28.02.00)

(30) Priority Data:
MI99A000454 5 March 1999 (05.03.99) IT

(71) Applicant *(for all designated States except US)*: FARMACEUTICI FORMENTI S.P.A. [IT/IT]; Via Correggio, 45, I–20149 Milano (IT).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: FABIANI, Flavio [IT/IT]; Via di Vittorio, 2, I–21040 Origgio (IT). FRIMONTI, Enrico [IT/IT]; Via di Vittorio, 2, I–21040 Origgio (IT). VALENTI, Mauro [IT/IT]; Via di Vittorio, 2, I–21040 Origgio (IT).

(74) Agents: MINOJA, Fabrizio et al.; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I–20122 Milano (IT).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: CONTROLLED–RELEASE COMPOSITIONS OF BETAHISTINE

(57) Abstract

Oral solid controlled–release formulations of betahistine, obtained subjecting the active ingredient to a melt–granulation process with a fatty compound and then mixing the obtained granulate with a hydrophilic polymer and with conventional excipients.