




Description

The present invention relates to the use of a class of 1-aryl-3-(4-piperidyl)-indole derivatives for the manufacture of medicaments for the treatment of anxiety in mammals, including humans, and certain of the compounds used which are novel compounds.

US patent No 4,710,500 corresponding to European patent No 0200322 discloses in general optionally 5-substituted 1-aryl-3-(4-piperidyl)- (I'), 1-aryl-3-(1-piperazinyl)- (II), or 1-aryl-3-(1,2,3,6-tetrahydro-4-pyridyl)-indole (III) derivatives having the formulas:




The compounds are claimed to be potent and long-lasting dopamine antagonists, and accordingly to be useful in the treatment of psychoses and additionally to be strong 5-HT₂ antagonists indicating effects in the treatment of negative symptoms of schizophrenia and depression and cardiovascular diseases. One of the tests used to show blockade of dopaminergic activity was a catalepsy test, catalepsy being at that time regarded as a test for antidopaminergic activity. However, at present catalepsy is considered to be a measure of the propensity of an antipsychotic compound to induce extrapyramidal side effects.

Today therapeutical treatment with benzodiazepines almost reign supreme for the treatment of anxiety. However, though they show activity against anxiety, the benzodiazepines have severe well known side effects. The most important of these side effects are sedation, dizziness, balance disturbances, interaction with alcohol, amnesia and a rapid addiction or habituation.

As a consequence the use of the benzodiazepines in the treatment of anxiety has become controversial, and accordingly there is a strong need for new means for the treatment of anxiety.

Surprisingly, it has now been found that certain 1-aryl-3-(4-piperidyl)-indole derivatives having the above formula I' have very potent anxiolytic activity substantially without the side effects mentioned and that they are without or only have weak cataleptogenic activity, thereby being useful in the treatment of anxiety.

Accordingly, the present invention provides the use of an 1-aryl-3-(4-piperidyl)-indole derivative having the general formula:




wherein

R¹ is hydrogen, halogen, lower alkyl, lower alkoxy, hydroxy, cyano, nitro, lower alkylthio, trifluoromethyl, trifluoromethylthio, lower alkylsulfonyl, amino, lower alkylamino or lower dialkylamino;

R is phenyl optionally substituted with one or more substituents independently selected from the following: halogen, lower alkyl, lower alkoxy, hydroxy, trifluoromethyl, and cyano, or R is a hetero aromatic group, preferably 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-thiazolyl, 2-oxazolyl, 2-imidazolyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl; and

R² is hydrogen, cycloalkyl, cycloalkylmethyl lower alkyl or lower alkenyl, optionally substituted with one or two hydroxy groups, any hydroxy group present being optionally esterified with an aliphatic carboxylic acid having from two to twentyfour carbon atoms inclusive, or R² is a group of the formula IV :



wherein n is an integer from 2 - 6;

W is oxygen, or sulphur;

U is nitrogen or CH;

Z is $-(CH_2)_m-$, m being 2 or 3, or Z is $-CH=CH-$ or 1,2-phenylene optionally substituted with halogen or trifluoromethyl, or Z is $-COCH_2-$ or $-CSC_2-$;

V is oxygen, sulphur, CH₂, or NR³, wherein R³ is hydrogen, lower alkyl or lower alkenyl optionally substituted with one or two hydroxy groups, or a cycloalkyl or cycloalkylmethyl group; or a pharmaceutically acceptable acid addition salt thereof for the manufacture of a medicament for therapeutical treatment of anxiety in mammals, including humans.

The term "lower alkyl" is intended to mean a straight or branched alkyl group having from one to four carbon atoms, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, etc. Lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkylamino and lower dialkylamino similarly designate such groups wherein the alkyl moiety is a lower alkyl group as defined above.

Lower alkenyl is intended to mean an alkenyl group containing from 2 to 4 carbon atoms, for example ethenyl, 1-propenyl, 2-butenyl, etc., and cycloalkyl is a cyclic alkyl group having from three to six carbon atoms inclusive

The pharmaceutically acceptable acid addition salts of the compounds used in the invention are salts formed with non-toxic organic or inorganic acids. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, embonic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanesulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene sulfonic and theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromo-theophylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids.

The compounds of the formula I and the pharmaceutically acceptable acid addition salts thereof may be administered in any suitable way, e.g. orally or parenterally, and the compounds may be presented in any suitable form for such administration, eg. in the form of tablets, capsules, powders, syrups or solutions or dispersions for injection.

An effective daily dose of a compound of the formula I or a pharmaceutically acceptable salt thereof is from 1.0 ng/Kg to 1.0 mg/Kg body weight.

The compounds used in accordance with the present invention have been shown to be highly active in animal models predictive of anxiolytic effects in man and they have been found not to induce catalepsy or only induce weak catalepsy which today is regarded as an indication of extrapyramidal side effects. It is indeed very surprising that the present compounds are non-cataleptogenic whereas the compounds of the formulas II and III of the above US patent have proved to be cataleptogenic (c.f. the pharmacological data in the following). The mechanisms behind this are not fully understood.

In addition to the benzodiazepines other compounds which have proven to be or are potentially effective in the treatment of anxiety in humans and are active in some animal models indicative of anxiety are buspirone (a partial 5-HT_{1A} agonist), ritanserin (a 5-HT₂ antagonist) and ondansetron (a selective 5-HT₃ antagonist). However said compounds are completely different from the present compounds in structure

and their mechanisms of action seems to be different from those of the present compounds (c.f. the description of the pharmacology in the following).

In a preferred embodiment of the invention the compound used is a compound of the formula I as defined in the foregoing wherein

- 5 R is phenyl substituted with halogen, or R is 2- or 3- thienyl;
 R¹ is hydrogen; halogen, lower alkyl, lower alkoxy, cyano, trifluoromethyl, or lower alkylsulphonyl;
 R² is a group having the formula IV as defined in the foregoing wherein n = 2 - 6;
 W is oxygen or sulphur;
 U is nitrogen;
- 10 Z is -(CH₂)₂-, -(CH₂)₃-, or -CH=CH-; and
 V is oxygen, CH₂ or NR³, R³ being hydrogen or lower alkyl;
 or a pharmaceutically acceptable acid addition salt thereof.

Most preferably the compound used is a compound having the general formula I as defined above, wherein:

- 15 R is phenyl substituted in 4 position with fluoro; R¹ is hydrogen, halogen, or lower alkyl; R² is a group having the formula IV as defined in the foregoing wherein n = 2; W is oxygen; U is nitrogen; Z is -(CH₂)₂;
 and V is NR³, R³ being hydrogen or lower alkyl; or a pharmaceutically acceptable acid addition salt thereof.

Though US patent No 4,710,500 generally comprises the 3-(4-piperidyl)-compounds of the formula I mentioned above, only five compounds have been specifically mentioned.

- 20 Accordingly the present invention also provides the novel 1-aryl-3-(4-piperidyl)-indole derivatives having the general formula I as defined above, provided that in case R is 4-fluorophenyl: R² may not be CH₃ when R¹ is CH₃ or CF₃, R¹ may not be hydrogen or chloro when R² is an 2-(2-imidazolidinon-1-yl)ethyl group and R¹ may not be CF₃ when R² is an 2-(1-pyrrolidin-2-on-yl)ethyl group; and pharmaceutically acceptable acid addition salts thereof.

- 25 The compounds of the formula I used in the invention may be prepared according to methods (b), (c), or (d) described in US patent No. 4,710,500. 2-pyrrolidinthiones are prepared from the corresponding lactame derivatives according to literature methods (Bull.Soc.Chim.Belg. 87, 223, 229, 299, 525 (1978)) by using Lawesson's reagent or phosphorous pentasulphide at appropriate temperatures. Imidazolidin-2-thion derivatives are prepared by ringclosure reactions from properly substituted ethylenediamines with carbon-
- 30 disulphide, thiophosgen or corresponding thiocarbonyl precursor compounds.

When R¹ is hydroxy the compound may be prepared by conventional methods of demethylation of the corresponding methyl ether. Pyridine hydrochloride, hydrobromic acid or methionine in methanesulphonic acid may be used to split off the methyl group.

- 35 The compounds wherein R¹ is cyano may be prepared by substitution of 5-bromo or 5-iodo in the appropriate substituted compounds using CuCN in an aprotic polar solvent such as N,N-dimethylformamide, N-methyl-2-pyrrolidone (NMP) or HMPA at elevated temperatures.

In compounds wherein R³ is hydroxy substituted alkyl the hydroxy group may be introduced by deprotection of a corresponding labile ether derivative, eg. by treating the benzyl ether in strong mineral acid, such as hydrobromic acid, or by catalytic debenzylation using Pd or Pt as catalyst.

- 40 The acid addition salts of the compounds used in the invention are easily prepared by methods well known in the art. The base is reacted with either the calculated amount of organic or inorganic acid in an aqueous miscible solvent, such as acetone or ethanol, with isolation of the salt by concentration and cooling, or an excess of the acid in an aqueous immiscible solvent such as ethyl ether or chloroform with the desired salt separating directly. Of course, these salts may also be prepared by the classical method of
- 45 double decomposition of appropriate salts.

The following compounds were prepared according to methods (b), (c), or (d) described in US patent No. 4,710,500, or from the corresponding lactame derivatives according to literature methods (Bull.Soc.Chim.Belg. 87, 223, 229, 299, 525 (1978)) by using Lawesson's reagent or phosphorous pentasulphide at appropriate temperatures:

- 50 1-(4-fluorophenyl)-5-methyl-3-(1-methyl-4-piperidyl)-1H-indole, hydrobromide, Lu 21-037, MP :254-256 °C
 1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, Lu 23-086, MP : 174-175 °C
 1-(4-fluorophenyl)-3-[1-[2-(2-pyrrolidinon-1-yl)ethyl]-4-piperidyl]-5-trifluoromethyl-1H-indole, fumarate, Lu 23-158, MP : 240-241 °C
 1-(4-fluorophenyl)-3-(1-methyl-4-piperidyl)-5-trifluoromethyl-1H-indole, oxalate, Lu 21-131, MP : 251-252 °C
- 55 5-chloro-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, Lu 23-174, MP : 162 °C
 5-chloro-1-(4-fluorophenyl)-3-[1-(2-hydroxyethyl)-4-piperidyl]-1H-indole, hydrochloride, 1 MP : 266-269 °C
 5-chloro-1-(4-fluorophenyl)-3-[1-[2-(2-oxazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, fumarate, 2, MP : 203-

205 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-(3-methyl-2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, fumarate, 3, MP : 198-199 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-(2-pyrrolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, fumarate, 4, MP : 209-211 °C

1-(4-fluorophenyl)-3-[1-[2-(3-methyl-2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-trifluoromethyl-1H-indole, 5, MP : 144-145 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-(2-pyrrolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, fumarate, 7, MP : 195-199 °C

1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-methylsulfonyl-1H-indole, fumarate, 8, MP : 188-192 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[6-(2-pyrrolidinon-1-yl)-1-hexyl]-4-piperidyl]-1H-indole, hydrochloride, 9, MP : 123-128 °C

5-chloro-1-(4-fluorophenyl)-3-(4-piperidyl)-1H-indole, fumarate, 10, MP : 196-201 °C

1-(4-fluorophenyl)-3-(4-piperidyl)-5-trifluoromethyl-1H-indole, hydrochloride, 11 MP : 281-284 °C

1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-trifluoromethyl-1H-indole, 12, MP : 169-171 °C

1-(4-fluorophenyl)-3-[1-[6-(2-pyrrolidinon-1-yl)-1-hexyl]-4-piperidyl]-5-trifluoromethyl-1H-indole, oxalate, 13, MP : 85-87 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1H-indole, oxalate, 14, MP : 92-96 °C

5-fluoro-1-(4-fluorophenyl)-3-(4-piperidyl)-1H-indole, fumarate, 15, MP : 198-200 °C

5-fluoro-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, oxalate, 16, MP : 188-190 °C

5-fluoro-1-(4-fluorophenyl)-3-[1-[2-(2-pyrrolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, fumarate, 17, MP : 178-180 °C

5-fluoro-1-(4-fluorophenyl)-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1H-indole, fumarate, 18, MP : 115-120 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[5-(2-imidazolidinon-1-yl)-1-pentyl]-4-piperidyl]-1H-indole, oxalate, 19, MP : 145-147 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[4-(2-imidazolidinon-1-yl)-1-butyl]-4-piperidyl]-1H-indole, oxalate, 20, MP : 178-179 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[6-(2-imidazolidinon-1-yl)-1-hexyl]-4-piperidyl]-1H-indole, oxalate, 21, MP : 156-158 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-(hydantoin-2-yl)ethyl]-4-piperidyl]-1H-indole, 22, MP : 174-176 °C

5-fluoro-1-(4-fluorophenyl)-3-[1-[6-(2-pyrrolidinon-1-yl)-1-hexyl]-4-piperidyl]-1H-indole, 23, oil

1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-methyl-1H-indole, 24, MP : 187-189 °C

1-(4-fluorophenyl)-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-5-methyl-1H-indole, hydrochloride, hydrate, 25, MP : 214-215 °C

1-(4-fluorophenyl)-3-[1-[2-(2-pyrrolidinon-1-yl)ethyl]-4-piperidyl]-5-methyl-1H-indole, hydrochloride, hemihydrate, 26, 265-266 °C

1-(4-fluorophenyl)-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-5-trifluoromethyl-1H-indole, 27, MP : 99-100 °C

3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1-(3-thienyl)-1H-indole, oxalate 28, MP : 139-140 °C

1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-methoxy-1H-indole, 29, MP : 167 °C

5-fluoro-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1-(3-thienyl)-1H-indole, oxalate, hemihydrate, 30, MP : 95-97 °C

5-fluoro-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1-(2-thienyl)-1H-indole, dioxalate, 31, MP : 173-174 °C

5-bromo-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, 32, MP : 171-172 °C

1-(4-fluorophenyl)-3-[1-[2-[3-(2-propyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1H-indole, hydrochloride, 33, MP : 226-227 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[3-(2-imidazolidinon-1-yl)-1-propyl]-4-piperidyl]-1H-indole, fumarate, 34, MP : 203-205 °C

5-chloro-1-(4-fluorophenyl)-3-[1-[2-[3-(2-hydroxyethyl)-2-imidazolidinon-1-yl]ethyl]-4-piperidyl]-1H-indole, fumarate, 35, MP : 180-182 °C

In the following example the preparation of compounds used in accordance with the invention is further illustrated by way of examples:

EXAMPLE 1

5-chloro-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinethion-1-yl)ethyl]-4-piperidyl]-1H-indole, oxalate, 35, MP : 150 °C

To a solution of 5-chloro-1-(4-fluorophenyl)-3-(4-piperidyl)-1H-indole (25 g) in N-methyl-2-pyrrolidone (150 ml) were added chloroacetonitrile (6 g) and triethylamine (10 ml). The reaction mixture was heated at 60 °C for one hour and subsequently poured onto crushed ice. The precipitated 5-chloro-3-(1-cyanomethyl-4-piperidyl)-1-(4-fluorophenyl)-1H-indole was filtered off and washed with water. Yield 20 g. MP : 170-172 °C.

A solution of the thus isolated cyanomethyl derivative (24 g) in dry THF (150 ml) was added dropwise to a previously prepared solution of AlH_3 (from 8 g of LiAlH_4 and 8 g of AlCl_3) in dry diethyl ether (250 ml). The mixture was heated at reflux for one hour and finally hydrolyzed by carefully adding a conc. aqueous solution of NaOH (10 ml) under simultaneous cooling. Inorganic salts were filtered off and were subsequently carefully washed with hot dichloromethane (2 x 100 ml). The combined organic phases were dried (anh. MgSO_4) and finally evaporated leaving 3-[1-(2-aminoethyl)-4-piperidyl]-5-chloro-1-(4-fluorophenyl)-1H-indole (25 g) as an oil.

Without further purification this product (12 g) and triethylamine (4.2 g) were heated in 1,1,1-trichloroethane (100 ml) at 50-55 °C. A solution of chloroacetonitrile (3.6 g) in 1,1,1-trichloroethane (10 ml) was added dropwise during 10 minutes. The mixture was heated for another 4 hours at 50 °C. Ethyl acetate (200 ml) was added and the mixture was poured into ice cooled dil. aqueous NaOH solution (400 ml). The organic phase was separated, washed with brine, dried (anh. MgSO_4) and the solvents evaporated leaving 5-chloro-3-[1-[2-(N-cyanomethyl)aminoethyl]-4-piperidyl]-1-(4-fluorophenyl)-1H-indole (14 g) as an oil.

The oil thus isolated was dissolved in dry THF (100 ml) and added dropwise to a previously prepared solution of AlH_3 (from 6 g of LiAlH_4 and 6 g of AlCl_3) in dry diethyl ether (200 ml). The mixture was refluxed for one hour and finally hydrolyzed by cautiously adding a conc. aqueous solution of NaOH (8 ml) under simultaneous cooling. Inorganic salts were filtered off and were subsequently washed with hot dichloromethane (2 x 100 ml). The combined organic phases were dried (anh. MgSO_4) and finally evaporated leaving 3-[1-[N-(2-aminoethyl)-2-aminoethyl]-4-piperidyl]-5-chloro-1-(4-fluorophenyl)-1H-indole (8.5 g) as an oil. This oil (4.5 g) was dissolved in 1-pentanol (50 ml) and carbondisulphide (5 ml) was added. After stirring for 2 hours at room temperature the resulting suspension was heated to 140 °C for 1.5 hours. Excess CS_2 was flushed away by a gentle stream of N_2 gas. Finally most of the 1-pentanol was evaporated at reduced pressure. The remaining oil was purified by column chromatography on silica gel (eluted with ethyl acetate/ethanol/triethylamine - 80/20/4). The oxalate salt of the title compound 35 crystallized from acetone.

Yield 250 mg. MP : 150 °C.

EXAMPLE 2

1-(4-Fluorophenyl)-5-hydroxy-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, 36, MP: 220 °C

Pyridinhydrochloride (60 g) and 1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-5-methoxy-1H-indole, compound 29, (6 g) were mixed and heated to 180 °C under N_2 for 1 1/2 hours. After cooling, water (300 ml) and ethyl acetate (100 ml) were added. By addition of NH_4OH solution the pH was adjusted to >9. The organic phase was separated, washed with water (50 ml), dried (anh. MgSO_4), and the solvent evaporated leaving the phenolic crude title compound as an oil. Purification by column chromatography on silica gel (eluted with ethyl acetate/dichloromethane/ethanol/triethylamine 60:20:20:5) afforded the title compound 36 as a crystalline material. Yield: 1.9 g. MP: 220 °C.

EXAMPLE 3

5-Cyano-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-4-piperidyl]-1H-indole, 37, MP: 209 °C.

To a solution of 5-bromo-1-(4-fluorophenyl)-3-(4-piperidyl)-1H-indole (17 g) in dichloromethane (170 ml) was added a solution of di-tert.-butyloxycarbonate (12 g) in dichloromethane (30 ml). After stirring for 30 minutes at room temperature the dichloromethane was evaporated in vacuo. 5-Bromo-3-(1-tert-butyloxycarbonyl-4-piperidyl)-1-(4-fluorophenyl)-1H-indole crystallized from n-heptane. Yield: 14 g. MP: 155 °C. All the crystalline material was dissolved in N-methyl-2-pyrrolidone (75 ml) and CaCN (5 g) was added. The mixture was heated at 160 °C for 6 hours. The mixture was then poured into a solution of NaCN (10 g)

in water (200 ml) and stirred for 20 minutes. Diethyl ether (200 ml) was added. The ether phase was separated, washed with brine (50 ml), dried (anh. MgSO_4), and the ether evaporated leaving a mixture of 5-bromo and 5-cyano compounds which were separated by column chromatography on silica gel (eluted with diethyl ether). The 3-(1-tert.butyloxycarbonyl-4-piperidyl)-5-cyano-1-(4-fluorophenyl)-1H-indole was isolated
 5 an an oil. Yield: 4.5 g. The protecting group - tert.butyloxycarbonyl - was splitted off by standard acidic (CF_3COOH) decomposition. The thus obtained 5-cyano-1-(4-fluorophenyl)-3-(4-piperidyl)-1H-indole (3.2 g) was dissolved in methyl isobutyl ketone (90 ml). Potassium carbonate (4.5 g), potassium iodide (0.5 g) and 1-(2-chloroethyl)-2-imidazolidinone (2.3 g) were added. The mixture was refluxed for 16 hours. After cooling inorganic salts were filtered off, and the organic solvent evaporated. Water (100 ml) and ethyl acetate (50
 10 ml) were added. The organic phase was separated, dried (anh. MgSO_4), and finally ethyl acetate evaporated leaving the crude title compound as an oil. Purification by column chromatography on silica gel (eluted with ethyl acetate / ethanol / triethylamine 80:20:4) afforded 2.1 g of pure crystalline title compound, 37. MP: 209 °C.

15 PHARMACOLOGY

The compounds used in the invention were tested in accordance with well recognized and reliable test methods. The tests were as follows:

20 CATALEPSY TEST

Evaluation of catalepsy was made according to Arnt (Eur. J. Pharmacol. 90, 47 - 55 (1983)). Catalepsy is an indicator of dopamine receptor blockade *in vivo* which correlates to the clinical potency of the classical neuroleptics. However, catalepsy is also considered to indicate extrapyramidal side effects (Arnt, J.
 25 et al.; *Neuropharmacology* 1981, 20, 1331-1334.

Procedure

In this test test compound is given s.c. in different doses. The rat (170 - 240 g) is placed on a vertical
 30 wire mesh (mesh diameter 12 mm). The rat is considered cataleptogenic if it remains immobile for more than 15 sec. The maximum number of rats showing catalepsy within the first 6 hours is recorded for each dose group. The results are recorded in fractions and an ED_{50} value is calculated by means of log-probit analysis. The results are shown in Table 1.

The following corresponding 1-aryl-3-(1,2,3,6-tetrahydropyridyl)- or 1-aryl-3-(piperazinyl)indole deriva-
 35 tives which are analogues of Lu 23-174 and Compound No 12, respectively, were included in the test as comparing compounds:

- A) 1-(4-Fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-1,2,3,6-tetrahydropyridin-4-yl]-5-trifluoromethyl-1H-indole
- B) 5-Chloro-1-(4-fluorophenyl)-3-[1-[2-(2-imidazolidinon-1-yl)ethyl]-1,2,3,6-tetrahydro-4-pyridyl]-1H-indole
- 40 C) 5-Chloro-1-(4-fluorophenyl)-3-[4-[2-(2-imidazolidinon-1-yl)ethyl]-1-piperazinyl]-1H-indole

TABLE 1

Compound	$\text{ED}_{50}(\text{s.c.}) (\mu\text{mol/kg})$
Lu 23-174	>98
comp. No 12	38
comp. No 2	>18
comp. No 3	31
45 comp. No 4	23
comp. No 14	>69
comp. No 16	>78
comp. No 24	>95
comp. A	0.49
50 comp. B	2.2
55 comp. C	4.5

Further ED₅₀ values of corresponding 1-aryl-3-(1,2,3,6-tetrahydro-4-pyridyl)- or 1-aryl-3-(1-piperazinyl)-indole derivatives are given in US patent No 4,710,500.

LIGHT/DARK DISCRIMINATION TEST IN MICE

This test was carried out in accordance with the method described in Costall et al. Br. J. Pharmacol. 90 275P (1987).

The test was conducted using a two compartment activity box in which the actions of anxiolytic compounds to reduce aversion against a brightly-lit environment may be readily detected. The box was designed as an open-top experimental box (45*27*27cm) one third of which was partitioned from the rest, painted black and illuminated with red light. The remainder of the box was painted white and brightly illuminated (1000 W). The floor of each area was lined into squares. Behavioural changes were determined for each area from video recordings for periods of 40 min. Data obtained from dose groups of 5 animals (male albino B6W mice, 25 - 30 g) were analysed using single factor analysis of variance, and Dunnett's t-test. Test compounds were given intraperitoneally 45 min before testing.

The results are given in the following Table 2 as minimum doses (mg/kg i.p.) tested showing effect (p<0.001). The behavioural changes used were exploratory rearings and line crossings.

The known substances diazepam, and ondansetron were included in the test for comparison purposes.

TABLE 2

Compound	Rearings Minimum effective dose (μg/kg)	Line crossings Minimum effective dose (μg/kg)
No 16	0.001<ME<0.1	0.001<ME<0.1
No 3	0.001	0.001
Lu 23-174	0.000001	0.000001
Diazepam	0.13	0.13
Ondansetron	0.05	0.05

The well known 5-HT₂ antagonist ritanserin did not show effect in this test.

LIGHT/DARK DISCRIMINATION TEST IN RATS.

The test was carried out similarly to the test in mice described above, however modified in accordance with F.C.Colpaert et al., Psychopharmacology (1985) 86: 45-54. The test used Wistar WU rats.

In this test the compound Lu 23-174 according to the invention showed effect in nano molar doses/kg (s.c.; observation 2h).

The well known 5-HT₂ antagonist ritanserin was only active in μmolar doses/kg.

INHIBITION OF AGGRESSION IN MICE.

This test is used to evaluate the ability of a test substance to inhibit isolation induced aggressive behaviour in mice and is considered to be a test model for anxiolytic effect. The method is a modified version of the method described by McMillen et al. in Drug Develop. Research 12, 53-62, 1988.

Procedure.

Male mice (NMRI/BOM,SPF) weighing 16-18 g were used. The mice were housed individually in macrolon cages for 3 weeks before training to attack an intruder mouse. The mice were trained daily for a period of 1-2 weeks until a satisfactory response was obtained. During the first days the mice were trained against each other. When a satisfactory response was obtained group-housed mice were used as intruders. The time elapsed until attack was recorded. The intruder mouse was removed as soon as an attack had occurred or after 180 sec. After the training period 60-70 percent of the mice had latencies of less than 10 sec.

The mice were pretested before drug treatment. Only mice attacking within 25 sec. were used. The animals were treated with test substance or saline. The mice were tested 30 or 120 minutes later, respectively. Nonspecific motoric effects were also recorded. 8 mice were used per group.

The results were stated as fractions of mice with attack latencies greater than or equal to a threshold value. As threshold values were used 180 and 90 sec. ED₅₀ values were calculated by log-probit analysis.

The results are given in the following table as ED₅₀ values in $\mu\text{mol/kg}$. The known compounds diazepam, buspirone and ritanserin were included in the test for the purpose of comparison.

TABLE 3

Compound	Threshold Value ED ₅₀ ($\mu\text{mol/kg}$)			
	30 min Pretreatment		120 min Pretreatment	
	90 sec.	180 sec.	90 sec.	180 sec.
Lu 23-174	4.4	3.2		
Lu-23-096			1.9	1.7
Compound No 16			2.1	1.6
Compound No 24			3.2	2.5
Diazepam	26	15		
Buspirone	9.6	5.4		
Ritanserin	>21	>21	>10	>10

It appears from the above results that the compounds used in the present invention have very weak or no cataleptic effect and have potent anxiolytic effects. So it is seen from table 1 that the cataleptic effects are much weaker than those of the corresponding 1-aryl-3-(1,2,3,6-tetrahydropyridyl)- or 1-aryl-3-(piperaziny)indole derivatives (a factor of about 100).

Furthermore the results show that the present compounds have much more potent anxiolytic effects than diazepam (until 5 orders of magnitude in the mouse dark/ light discrimination test and a factor 6-10 in the aggression test).

In the light/dark discrimination test in mice the present compounds are very potent (effect in nano molar doses) whereas the well known 5HT₂ antagonist ritanserin is inactive. Additionally ritanserin showed no effect in the aggression test. This shows that the anxiolytic activity cannot be a consequence of the 5HT₂ action alone.

Furthermore, the compounds used in the present invention appear to be more potent than the substances proven to be effective in humans, such as benzodiazepines and buspirone, and the recent potential anxiolytics, such as ritanserin and ondansetron.

FORMULATION EXAMPLES

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example: Tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive colourings, aroma, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by solving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilization of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Typical examples of recipes for the formulations of the invention are as follows:

- 1) Tablets containing 0.5 milligrams of Lu 23-174 calculated as the free base:

Lu 23-174	0.5 mg
Lactose	18 mg
Potato starch	27 mg
Saccharose	58 mg
Sorbitol	3 mg
Talcum	5 mg
Gelatine	2 mg
Povidone	1 mg
Magnesium stearate	0.5 mg

2) Tablets containing 1 milligrams of compound No 3 calculated as the free base:

Comp. 3	1.0 mg
Lactose	16 mg
Potato starch	45 mg
Saccharose	106 mg
Sorbitol	6 mg
Talcum	9 mg
Gelatine	4 mg
Povidone	3 mg
Magnesium stearate	0.6 mg

3) Syrup containing per milliliter:

Comp. 16	5.0 mg
Sorbitol	500 mg
Tragacanth	7 mg
Glycerol	50 mg
Methyl-paraben	1 mg
Propyl-paraben	0.1 mg
Ethanol	0.005 ml
Water	ad 1 ml

4) Solution for injection containing per milliliter:

Lu 23-174	0.2 mg
Acetic acid	17.9 mg
Sterile water	ad 1 ml


5) Solution for injection containing per milliliter:

Comp. 3	0.5 mg
Sorbitol	42.9 mg
Acetic acid	0.63 mg
Sodium hydroxide	22 mg
Sterile water	ad 1 ml



Claims


1. Use of an 1-aryl-3-(4-piperidyl)-indole derivative having the general formula:



wherein

R¹ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, cyano, nitro, C₁-C₄ alkylthio, trifluoromethyl, trifluoromethylthio, C₁-C₄ alkylsulfonyl, amino, C₁-C₄ alkylamino or di(C₁-C₄)alkylamino; R is phenyl optionally substituted with one or more substituents independently selected from the following: halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, trifluoromethyl, and cyano, or R is a hetero aromatic group, preferably 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-thiazolyl, 2-oxazolyl, 2-imidazolyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl; and

R² is hydrogen, cycloalkyl, cycloalkylmethyl, C₁-C₄ alkyl or C₂-C₄ alkenyl, optionally substituted with one or two hydroxy groups, any hydroxy group present being optionally esterified with an aliphatic carboxylic acid having from two to twentyfour carbon atoms inclusive, or R² is a group of the formula IV :



wherein n is an integer from 2 - 6;

W is oxygen, or sulphur;

U is nitrogen or CH;

Z is -(CH₂)_m-, m being 2 or 3, or Z is -CH=CH- or 1,2-phenylene optionally substituted with halogen or trifluoromethyl, or Z is -COCH₂- or -CSCH₂-;

V is oxygen, sulphur, CH₂, or NR³, wherein R³ is hydrogen, C₁-C₄ alkyl or C₂-C₄ alkenyl optionally substituted with one or two hydroxy groups, or a cycloalkyl or cycloalkylmethyl group;

or a pharmaceutically acceptable acid addition salt thereof for the manufacture of a medicament for therapeutic treatment of anxiety in mammals, including humans.

2. Use according to claim 1, characterized in that the compound used is a compound of the formula I as defined in claim 1 wherein R is phenyl substituted with halogen, or R is 2- or 3- thienyl;

R¹ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, cyano, trifluoromethyl, or C₁-C₄ alkylsulphonyl;

R² is a group having the formula IV as defined in the foregoing wherein n = 2 - 6 ;

W is oxygen or sulphur;

U is nitrogen;

Z is -(CH₂)₂-, -(CH₂)₃-, or -CH=CH-; and

V is oxygen, CH₂ or NR³, R³ being hydrogen or C₁-C₄ alkyl;

or a pharmaceutically acceptable acid addition salt thereof.


3. Use according to claim 1 or 2, characterized in that the compound used is a compound of the formula I as defined in claim 1 wherein

R is phenyl substituted in the 4-position with fluoro; R¹ is hydrogen, halogen, or C₁-C₄ alkyl; R² is a group having the formula IV as defined in claim 1 or 2 wherein n = 2; W is oxygen; U is nitrogen; Z is



-(CH₂)₂-; and V is NR³, R³ being hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable acid addition salt thereof.

4. An 1-aryl-3-(4-piperidyl)-indole derivative having the general formula:




wherein

R¹ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, cyano, nitro, C₁-C₄ alkylthio, trifluoromethyl, trifluoromethylthio, C₁-C₄ alkylsulfonyl, amino, C₁-C₄ alkylamino or lower di(C₁-C₄)-alkylamino;

R is phenyl optionally substituted with one or more substituents independently selected from the following: halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, trifluoromethyl, and cyano, or R is a hetero aromatic group, preferably 2-thienyl, 3-thienyl, 2-furanyl, 3-furanyl, 2-thiazolyl, 2-oxazolyl, 2-imidazolyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl; and

R² is hydrogen, cycloalkyl, cycloalkylmethyl, C₁-C₄ alkyl or C₂-C₄ alkenyl, optionally substituted with one or two hydroxy groups, any hydroxy group present being optionally esterified with an aliphatic carboxylic acid having from two to twentyfour carbon atoms inclusive, or R² is a group of the formula IV



wherein n is an integer from 2 - 6;

W is oxygen, or sulphur;

U is nitrogen or CH;

Z is -(CH₂)_m-, m being 2 or 3, or Z is -CH=CH- or 1,2-phenylene optionally substituted with halogen or trifluoromethyl, or Z is -COCH₂- or -CSCH₂-;

V is oxygen, sulphur, CH₂, or NR³, wherein R³ is hydrogen, C₁-C₄ alkyl or C₂-C₄ alkenyl optionally substituted with one or two hydroxy groups, or a cycloalkyl or cycloalkylmethyl group;

provided that in case R is 4-fluorophenyl: R² may not be CH₃ when R¹ is CH₃ or CF₃; R¹ may not be hydrogen or chloro when R² is an 2-(2-imidazolidinon-1-yl)-ethyl group and R¹ may not be CF₃ when R² is an 2-(1-pyrrolidin-2-on-yl)ethyl group; and pharmaceutically acceptable acid addition salts thereof.

5. A compound according to claim 4, characterized in that

R is phenyl substituted with halogen, or R is 2- or 3- thienyl;

R¹ is hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, cyano, trifluoromethyl, or C₁-C₄ alkylsulphonyl;

R² is a group having the formula IV as defined in the foregoing wherein n = 2 - 6 ;

W is oxygen or sulphur;

U is nitrogen;

Z is -(CH₂)₂-, -(CH₂)₃-, or -CH=CH-; and


V is oxygen, CH₂ or NR³, R³ being hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable acid addition salt thereof.



6. A compound according to claim 5, characterized in that R is phenyl substituted in the 4-position with fluoro; R¹ is hydrogen, halogen, or C₁-C₄ alkyl; R² is a group having the formula IV as defined in claim 1 or 2 wherein n = 2; W is oxygen; U is nitrogen; Z is -(CH₂)₂-; and V is NR³, R³ being hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable acid addition salt thereof.

Patentansprüche


1. Verwendung eines 1-Aryl-3-(4-piperidyl)-indolderivates der allgemeinen Formel:



worin R¹ Wasserstoff, Halogen, C₁-C₄ Alkyl, C₁-C₄ Alkoxy, Hydroxy, Cyan, Nitro, C₁-C₄ Alkylthio, Trifluormethyl, Trifluormethylthio, C₁-C₄ Alkylsulphonyl, Amino, C₁-C₄ Alkylamino oder di(C₁-C₄) Alkylamino ist;

R ist ein Phenylrest, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig voneinander aus den Folgenden ausgewählt: Halogen, C₁-C₄ Alkyl, C₁-C₄ Alkoxy, Hydroxy, Trifluormethyl und Cyan, oder R ist eine heteroaromatische Gruppe, vorzugsweise 2-Thienyl, 3-Thienyl, 2-Furanyl, 3-Furanyl, 2-Thiazolyl, 2-Oxazolyl, 2-Imidazolyl, 2-Pyridyl, 3-Pyridyl, oder 4-Pyridyl; und

R² ist Wasserstoff, Cycloalkyl, Cycloalkylmethyl, C₁-C₄ Alkyl, oder C₂-C₄ Alkenyl, gegebenenfalls substituiert mit einer oder zwei Hydroxygruppen, wobei jede anwesende Hydroxygruppe gegebenenfalls mit einer aliphatischen Carbonsäure, mit zwischen einschließlich zwei und einschließlich vierundzwanzig Kohlenstoffatomen verestert ist, oder R² ist eine Gruppe der Formel IV:



worin n eine ganze Zahl von 2-6 ist;

W ist Sauerstoff, oder Schwefel;

U ist Stickstoff oder Kohlenstoff;

Z ist -(CH₂)_m-, wobei m gleich 2 oder 3 ist, oder Z ist -CH=CH- oder 1,2-Phenylen, gegebenenfalls substituiert mit Halogen oder Trifluormethyl, oder Z ist -COCH₂- oder CSCH₂-;

V ist Sauerstoff, Schwefel, CH₂, oder NR³, worin R³ Wasserstoff ist, C₁-C₄ Alkyl, oder C₂-C₄ Alkenyl, gegebenenfalls substituiert mit einer oder zwei Hydroxygruppen, oder Cycloalkyl, oder Cycloalkylmethylgruppe,

oder ein pharmazeutisch annehmbares Säureadditionssalz derselben zur Herstellung eines Medikaments für die therapeutische Behandlung von Angstzuständen bei Säugetieren, einschließlich Menschen.

2. Verwendung nach Anspruch 1, dadurch gekennzeichnet, daß die verwendete Verbindung eine Verbindung der Formel 1, wie in Anspruch 1 definiert, ist, worin

R ein mit Halogen substituierter Phenylrest ist, oder R ist 2- oder 3-Thienyl;

R¹ ist Wasserstoff, Halogen, C₁-C₄ Alkyl, C₁-C₄ Alkoxy, Cyan, Trifluormethyl, oder C₁-C₄ Alkylsulphonyl,

R² ist eine Gruppe der Formel IV, wie in der vorhergehenden Formel definiert, worin n = 2-6;



W ist Sauerstoff oder Schwefel;


U ist Stickstoff;

Z ist $-(CH_2)_2-$, $-(CH_2)_3-$, oder $-CH=CH-$, und

V ist Sauerstoff, CH_2 oder NR^3 , wobei R^3 Wasserstoff oder C_1 - C_4 Alkyl ist; oder ein pharmazeutisch annehmbares Säureadditionssalz derselben.

3. Verwendung nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß die verwendete Verbindung eine Verbindung der Formel I, wie in Anspruch 1 definiert, ist, worin R ein in der Position 4 mit Fluor substituierter Phenylrest ist; R^1 ist Wasserstoff, Halogen oder C_1 - C_4 Alkyl; R^2 ist eine Gruppe der Formel IV, wie in Anspruch 1 oder 2 definiert, worin n gleich 2 ist; W ist Sauerstoff; U ist Stickstoff; Z ist $-(CH_2)_2-$; und V ist NR^3 , wobei R^3 Wasserstoff oder C_1 - C_4 Alkyl ist; oder ein pharmazeutisch annehmbares Säureadditionssalz derselben.


4. Ein 1-Aryl-3-(4-piperidyl)-indolderivat, der allgemeinen Formel:



worin R^1 Wasserstoff, Halogen, C_1 - C_4 Alkyl, C_1 - C_4 Alkoxy, Hydroxy, Cyan, Nitro, C_1 - C_4 Alkylthio, Trifluormethyl, Trifluormethylthio, C_1 - C_4 Alkylsulphonyl, Amino, C_1 - C_4 Alkylamino oder ein niedriger di- (C_1-C_4) Alkylaminorest ist;

R ist eine Phenylgruppe, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig voneinander aus den Folgenden ausgewählt: Halogen, C_1 - C_4 Alkyl, C_1 - C_4 Alkoxy, Hydroxy, Trifluormethyl und Cyan, oder R ist eine heteroaromatische Gruppe, vorzugsweise eine 2-Thienyl, 3-Thienyl, 2-Furanyl, 3-Furanyl, 2-Thiazolyl, 2-Oxazolyl, 2-Imidazolyl, 2-Pyridyl, 3-Pyridyl, oder 4-Pyridyl; und

R^2 ist Wasserstoff, Cycloalkyl, Cycloalkylmethyl, C_1 - C_4 Alkyl, oder C_2 - C_4 Alkenyl, gegebenenfalls substituiert mit einer oder zwei Hydroxygruppen, wobei jede anwesende Hydroxygruppe gegebenenfalls mit einer aliphatischen Carbonsäure, mit zwischen einschließlich zwei und einschließlich vierundzwanzig Kohlenstoffatomen verestert ist, oder R^2 ist eine Gruppe der Formel IV:



worin n eine ganze Zahl von 2-6 ist;

W ist Sauerstoff, oder Schwefel;

U ist Stickstoff oder Kohlenstoff;

Z ist $-(CH_2)_m-$, wobei m gleich 2 oder 3 ist, oder Z ist $-CH=CH-$ oder ein 1,2-Phenylen, gegebenenfalls substituiert mit Halogen oder Trifluormethyl, oder Z ist $-COCH_2-$ oder $CSCH_2-$;

V ist Sauerstoff, Schwefel, CH_2 , oder NR^3 , worin R^3 Wasserstoff ist, C_1 - C_4 Alkyl, oder C_2 - C_4 Alkenyl, gegebenenfalls substituiert mit einer oder zwei Hydroxygruppen, oder einer Cycloalkyl, oder einer Cycloalkylmethylgruppe;

vorausgesetzt, daß im Fall R gleich 4-Fluorphenyl: R^2 kann nicht CH_3 sein, wenn R^1 gleich CH_3 oder CF_3 ist; R^1 kann nicht Wasserstoff oder Chlor sein, wenn R^2 ein 2-(2-Imidazolidinon-1-yl)ethylrest ist, und R^1 kann nicht CF_3 sein, wenn R^2 ein 2-(1-Pyrrolidin-2-on-yl)ethylrest ist;



und pharmazeutisch annehmbare Säureadditionssalze derselben.

5. Eine Verbindung nach Anspruch 4, dadurch gekennzeichnet, daß R eine mit Halogen substituierte Phenylgruppe ist, oder R ist 2- oder 3-Thienyl;

R¹ ist Wasserstoff, Halogen, C₁-C₄ Alkyl, C₁-C₄ Alkoxy, Cyan, Trifluormethyl, oder C₁-C₄ Alkylsulphonyl,

R² ist ein Rest der Formel IV, wie in der vorhergehenden Formel definiert, worin n = 2-6;

W ist Sauerstoff oder Schwefel;

U ist Stickstoff;


Z ist -(CH₂)₂-, -(CH₂)₃-, oder -CH=CH-, und

V ist Sauerstoff, CH₂ oder NR³, wobei R³ Wasserstoff oder C₁-C₄ Alkyl ist; oder ein pharmazeutisch annehmbares Säureadditionssalz derselben.

6. Eine Verbindung nach Anspruch 5, dadurch gekennzeichnet, daß R ein in der Position 4 mit Fluor substituierter Phenylrest ist; R¹ ist Wasserstoff, Halogen oder C₁-C₄ Alkyl; R² ist ein Rest der Formel IV, wie in Anspruch 1 oder 2 definiert, worin n = 2 ist; W ist Sauerstoff; U ist Stickstoff; Z ist -(CH₂)₂-; und V ist NR³, wobei R³ Wasserstoff oder C₁-C₄ Alkyl ist; oder ein annehmbares Säureadditionssalz derselben.

Revendications

1. Utilisation d'un dérivé de 1-aryl-3-(4-pipéridyl)-indole ayant la formule générale :




dans laquelle

R¹ est hydrogène, halogène, alkyle en C₁-C₄; alkoxy en C₁-C₄, hydroxy, cyano, nitro, alkylthio en C₁-C₄, trifluorométhyle, trifluorométhylthio, alkylsulfonyl en C₁-C₄, amino, alkylamino en C₁-C₄ ou di(C₁-C₄) alkylamino ;

R est phényle, facultativement substitué par un ou plusieurs substituants choisis indépendamment parmi : halogène, alkyle en C₁-C₄, alkoxy en C₁-C₄, hydroxy, trifluorométhyle, et cyano, ou R est un groupe hétéro-aromatique, de préférence 2-thiényle, 3-thiényle, 2-furanyl, 3-furanyl, 2-thiazolyle, 2-oxazolyle, 2-imidazolyle, 2-pyridyle, 3-pyridyle, ou 4-pyridyle ; et

R² est hydrogène, cycloalkyle, cycloalkylméthyle, alkyle en C₁-C₄ ou alcényle en C₂-C₄, facultativement substitués par un ou deux groupes hydroxy, tout groupe hydroxy présent étant facultativement estérifié avec un acide carboxylique aliphatique ayant de 2 à 24 atomes de carbone inclus, ou R² est un groupe de la formule IV :



dans laquelle n est un nombre entier de 2 à 6 ;

W est oxygène ou soufre ;

U est azote ou CH

Z est -(CH₂)_m-, m étant 2 ou 3, ou Z est CH=CH, ou 1,2-phénylène, facultativement substitué par



halogène ou trifluorométhyle, ou Z est $-\text{COCH}_2-$ ou $-\text{CSCH}_2-$;

V est oxygène, soufre, CH_2 , ou NR^3 , dans lequel R^3 est hydrogène, en C_1 - C_4 alkyle ou en C_2 - C_4 alcényle, facultativement substitué par un ou deux groupes hydroxy, ou un groupe cycloalkyle ou cycloalkylméthyle ;

ou son sel d'addition d'acide acceptable pharmaceutiquement pour la fabrication d'un médicament destiné au traitement thérapeutique de l'anxiété chez les mammifères y compris chez l'homme.

2. Utilisation selon la revendication 1, caractérisée en ce que le composé utilisé est un composé de la formule I, définie dans la revendication 1, dans laquelle R est phényle, substitué par halogène ou R est 2 ou 3-thiényle ;

R^1 est hydrogène, halogène, alkyle C_1 - C_4 -, alcoxy C_1 - C_4 -, cyano, trifluorométhyle, ou alkylsulfonyl C_1 - C_4 ;

R^2 est un groupe ayant la formule IV définie ci-dessus, dans laquelle

$\text{N} = 2 - 6$;

W est oxygène ou soufre ;


U est azote ;

Z est $-(\text{CH}_2)_2-$, $-(\text{CH}_2)_3-$, ou $\text{CH}=\text{CH}$; et

V est oxygène- CH_2 ou NR^3 , R^3 étant hydrogène ou alkyle C_1 - C_4 ou son sel d'addition d'acide acceptable pharmaceutiquement.

3. Utilisation selon la revendication 1 ou 2, caractérisée en ce que le composé utilisé est un composé de la formule I définie dans la revendication 1, dans laquelle R est phényle, substitué dans la position 4 par fluoro ; R^1 est hydrogène, halogène ou alkyle en C_1 - C_4 ; R^2 est un groupe ayant la formule IV définie dans la revendication 1 ou 2 dans laquelle $n = 2$; W est oxygène ; U est azote, Z est $-(\text{CH}_2)_2-$; et V est NR^3 , R^3 étant hydrogène ou alkyle en C_1 - C_4 ; ou son sel d'addition d'acide acceptable pharmaceutiquement.

4. Dérivé de 1-aryle-3-(4-pipéridyle)-indole ayant la formule générale




dans laquelle

R^1 est hydrogène, halogène, alkyle en C_1 - C_4 , alcoxy en C_1 - C_4 , hydroxy, cyano, nitro, alkylthio en C_1 - C_4 , trifluorométhyle, trifluorométhylthio, alkylsulfonyl en C_1 - C_4 , amino, alkylamino en C_1 - C_4 ou un di- $(\text{C}_1$ - C_4) alkylamino inférieur ;

R est phényle, facultativement substitué par un ou plusieurs substituants, choisis indépendamment parmi : halogène, alkyle en C_1 - C_4 , alcoxy en C_1 - C_4 , hydroxy, trifluorométhyle, et cyano ou R est un groupe hétéro-aromatique, de préférence 2-thiényle, 3-thiényle, 2 furanyle, 3-furanyl, 2-thiazolyne, 2-oxazolyne, 2-imidazolyne, 2-pyridyle, 3-pyridyle ou 4-pyridyle ; et

R^2 est hydrogène, cycloalkyle, cycloalkylméthyle, alkyle en C_1 - C_4 ou alcényle en C_2 - C_4 , facultativement substitués par un ou deux groupes hydroxy, tout groupe hydroxy présent étant facultativement estérifié par un acide carboxylique aliphatique ayant de 2 à 24 atomes de carbone inclus, ou R^2 est un groupe de la formule IV :



dans laquelle n est un nombre entier de 2 à 6 ;

W est oxygène, ou soufre ;

U est azote ou CH

Z est $-(CH_2)_m-$, m étant 2 ou 5 ou Z est $CH=CH$ ou 1,2 phénylène, facultativement substitué par halogène ou trifluorométhyle ou Z est $-COCH_2-$ ou $-CSCCH_2-$;

V est oxygène, soufre, CH_2 , ou NR^3 , dans lequel R^3 est hydrogène, alkyle en C_1-C_4 ou alcényle, en C_2-C_4 , facultativement substitués par un ou deux groupe hydroxy, ou un groupe cycloalkyle ou cycloalkylméthyle ;

dans le cas où R est 4-fluorophényle : R^2 peut ne pas être CH_3 lorsque R^1 est CH_3 ou CF_3 ; R^1 peut ne pas être hydrogène ou chloro lorsque R^2 est un groupe 2-(2-imidazolidinon-1-yl)-éthyle et R^1 peut ne pas être CF_3 lorsque R^2 est un groupe 2-(1-pyrrolidin-2-on-yl)-éthyle ;

et leurs sels d'addition d'acide pharmaceutiquement acceptables.

5. Composé selon la revendication 4, caractérisé en ce que

R est phényle substitué par halogène, ou R est 2 ou 3-thiényle,

R^1 est hydrogène, halogène, alkyle en C_1-C_4 , alcoxy en C_1-C_4 , cyano, trifluorométhyle, ou alkylsulphonyle en C_1-C_4 ;

R^2 est un groupe ayant la formule IV, définie ci dessus, dans laquelle $n = 2 - 6$;

W est oxygène ou soufre ;

U est azote ;

Z est $-(CH_2)_2-$, $-(CH_2)_3-$, ou $CH=CH$; et

V est oxygène, CH_2 ou NR^3 , R^3 étant hydrogène ou alkyle en C_1-C_4 ; ou son sel d'addition d'acide pharmaceutiquement acceptable.

6. Composé selon la revendication 5, caractérisé en ce que

R est phényle, substitué dans la position 4 par fluoro ; R^1 est hydrogène, halogène, ou alkyle en C_1-C_4 ; R^2 est un groupe ayant la formule IV, définie dans la revendication 1 ou 2, dans laquelle $n = 2$; W

est oxygène ; U est azote ; Z est $-(CH_2)_2-$; et V est NR^3 , R^3 étant hydrogène ou alkyle en C_1-C_4 ; ou son sel d'addition d'acide pharmaceutiquement acceptable.