


Claims

1. Use of a compound of the formula



or a diastereomer, an enantiomer, a salt, a solvate, a solvate of the salt thereof, in the manufacture of a medicament for the treatment, and/or prophylaxis of allergic dermatitis in a dog, wherein the compound or the diastereomer, the enantiomer, the salt, the solvate, the solvate of the salt thereof, is to be administered to the dog orally,

wherein the allergic dermatitis is Atopic Dermatitis or Flea Allergy Dermatitis.

2. Use of a compound selected from the group consisting of:
- 1) N-[6-(2-Hydroxypropan-2-yl)-2-(2-methoxyethyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 2) N-[6-(Hydroxymethyl)-2-(2-methoxyethyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 3) N-[6-(2-Hydroxypropan-2-yl)-2-(3-methoxypropyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 4) N-[6-(Hydroxymethyl)-2-(3-methoxypropyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 5) N-[2-(2-Hydroxyethyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 6) N-[6-(2-Hydroxypropan-2-yl)-2-(3-hydroxypropyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 7) N-[2-(2-Hydroxyethyl)-6-(hydroxymethyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
 - 8) N-[6-(2-Hydroxypropan-2-yl)-2-(oxetan-3-ylmethyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide

- 9) N-[6-(Hydroxymethyl)-2-(oxetan-3-ylmethyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
- 10) N-{6-(2-Hydroxypropan-2-yl)-2-[3-(methylsulphonyl)propyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide
- 11) N-[2-(3-Hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
- 12) N-{6-(2-Hydroxypropan-2-yl)-2-[2-(methylsulphonyl)ethyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide
- 13) 6-(Difluoromethyl)-N-[2-(3-hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]pyridine-2-carboxamide
- 14) 6-(Difluoromethyl)-N-{6-(2-hydroxypropan-2-yl)-2-[2-(methylsulphonyl)ethyl]-2H-indazol-5-yl}pyridine-2-carboxamide
- 15) 6-(Difluoromethyl)-N-[6-(2-hydroxypropan-2-yl)-2-(3-hydroxypropyl)-2H-indazol-5-yl]pyridine-2-carboxamide
- 16) N-[6-(2-Hydroxypropan-2-yl)-2-(4,4,4-trifluorobutyl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide
- 17) N-{6-(2-Hydroxypropan-2-yl)-2-[3-(trifluoromethoxy)propyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide
- 18) N-{6-(2-Hydroxypropan-2-yl)-2-[3-(2,2,2-trifluoroethoxy)propyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide
- 19) 5-Fluoro-N-[2-(3-hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-methylpyridine-2-carboxamide
- 20) N-[2-(3-Hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-methylpyridine-2-carboxamide
- 21) 6-(2-Hydroxypropan-2-yl)-N-[6-(2-hydroxypropan-2-yl)-2-(4,4,4-trifluorobutyl)-2H-indazol-5-yl]pyridine-2-carboxamide; and
- 22) N-{2-[2-(1-Hydroxycyclopropyl)ethyl]-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide; or

or a diastereomer, an enantiomer, a salt, a solvate, a solvate of the salt thereof, in the manufacture of a medicament for the treatment, and/or prophylaxis of allergic dermatitis in a dog, wherein the compound or the diastereomer, the enantiomer, the salt, the solvate, the solvate of the salt thereof, is to be administered to the dog orally,

wherein the allergic dermatitis is Atopic Dermatitis or Flea Allergy Dermatitis.

3. The use of claim 1, wherein the allergic dermatitis is Atopic Dermatitis.

4. The use of claim 1, wherein the allergic dermatitis is Flea Allergy Dermatitis.
5. The use of claim 1, wherein the compound or a pharmaceutically acceptable salt thereof is to be administered to the dog at a daily dose of 10 to 20 mg per kg body weight.
6. The use of claim 4, wherein the compound or a pharmaceutically acceptable salt thereof is to be orally administered to the dog at a dose of about 10 mg per kg body weight.
7. The use of claim 4, wherein the compound is to be orally administered to the dog at a dose of about 20 mg per kg body weight.
8. The use of claim 3, wherein the compound or a pharmaceutically acceptable salt thereof is to be orally administered to the dog at a dose of about 10 mg per kg body weight.
9. The use of claim 3, wherein the compound is to be orally administered to the dog at a dose of about 20 mg per kg body weight.
10. The use of claim 5, wherein the compound or a pharmaceutically acceptable salt thereof is to be orally administered to the dog in the form of a tablet or capsule comprising inert, nontoxic, pharmaceutically suitable excipients.
11. The use of claim 1, wherein the compound is N-{6-(2-Hydroxypropan-2-yl)-2-[2-(methylsulphonyl)ethyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide.
12. The use of claim 2, wherein the compound is selected from the group consisting of
 - 11) N-[2-(3-Hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-(trifluoromethyl)pyridine-2-carboxamide;
 - 12) N-{6-(2-Hydroxypropan-2-yl)-2-[2-(methylsulphonyl)ethyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide;

- 13) 6-(Difluoromethyl)-N-[2-(3-hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]pyridine-2-carboxamide; and
 - 19) 5-Fluoro-N-[2-(3-hydroxy-3-methylbutyl)-6-(2-hydroxypropan-2-yl)-2H-indazol-5-yl]-6-methylpyridine-2-carboxamide.
13. The use of claim 2, wherein the compound is N-{6-(2-Hydroxypropan-2-yl)-2-[2-(methylsulphonyl)ethyl]-2H-indazol-5-yl}-6-(trifluoromethyl)pyridine-2-carboxamide.