

IgA MONOCLONAL ANTIBODIES FOR TREATING FLAVIVIRUS INFECTION

CROSS REFERENCE TO RELATED APPLICATIONS

- 5 [0001] This application claims the benefit of the filing date of U.S. Provisional Application No. 63/235,325 filed August 20, 2021. The content of this earlier filed application is hereby incorporated by reference herein in its entirety.

REFERENCE TO A SEQUENCE LISTING

- [0002] This application contains a Sequence Listing in computer readable form, which
10 is incorporated herein by reference.

FIELD OF THE INVENTION

- [0003] The present disclosure relates to immunoglobulin class switching or class-switch recombination, immunoglobulins that neutralize flavivirus infection and/or mitigate antibody-dependent enhancement (ADE) of flavivirus virus infection such as
15 ADE associated with secondary heterologous dengue infections.

BACKGROUND

- [0004] Dengue virus (DENV) is one of the most widespread vector-borne viral pathogens in the world. Including four immunologically and genetically distinct serotypes (DENV-1, -2, -3, and -4), DENV is borne primarily by the tropical and
20 subtropical mosquitoes *Aedes aegypti* and *A. albopictus* [1, 2]. DENV and its mosquito vectors can currently be found across Central and South America, South and South-East Asia, the Western Pacific, and sub-Saharan Africa, meaning 40% of the world's population is currently at risk of exposure and infection [1-3]. Consequently, an estimated 400 million DENV infections are thought to occur every year, resulting in
25 100 million clinically apparent infection [2]. Approximately 500,000 cases per year progress to severe dengue—characterized by thrombocytopenia, vascular leakage and hemorrhage—resulting in nearly 20,000 deaths [4-7].

[0005] A distinct epidemiological feature of dengue as compared to other flaviviral diseases is the increased risk for severe disease upon heterologous secondary

infection [8]. While the risk factors associated with developing severe dengue upon secondary DENV exposure are complex and incompletely understood, the leading mechanistic explanation for this phenomenon is a process known as antibody-dependent enhancement (ADE) [9, 10]. ADE is thought to occur when poorly-neutralizing or sub-neutralizing concentrations of DENV-reactive IgG opsonizes DENV and facilitates its entry into permissive FcγR-bearing cells [11]. Various lines of evidence support the association of ADE with severe dengue, including increased incidence of severe dengue in infants born to dengue-immune mothers [12-14]; increased viremia in interferon (IFN) receptor-deficient mice or non-primates passively immunized with anti-DENV antibodies [15, 16]; and increased incidence of severe dengue during the second of sequential/heterologous DENV outbreaks and in patients with a narrow range of preexisting anti-DENV antibody titers [17, 18]. Furthermore, *in-vitro* assessments of serum ADE activity in dengue-primed non-human primates have been shown to correlate with viral titers following heterologous attenuated DENV infection [19].

[0006] The increased risk of severe dengue upon secondary heterologous infection also presents a challenge to vaccine development as incomplete or waning vaccine-elicited immunity may place recipients at an increased risk of developing severe dengue should they be exposed following vaccination [20]. This is most significantly highlighted by the revelation that the only currently available DENV vaccine (DENVAXIA®) fails to protect previously DENV naïve individuals from infection and can increase the risk of hospitalization with virologically confirmed dengue [21-23]. Accordingly, understanding the subtleties of both natural and vaccine-elicited DENV humoral immunity is critical for further the understanding of disease risk and infection-associated immunopathogenesis.

[0007] To date, the literature on dengue serology has overwhelmingly focused on the contribution of immunoglobulin isotypes IgM and IgG to functional dengue immunity and infection-associated immunopathogenesis. During both primary and secondary dengue infection, these isotype antibodies follow a highly predictable pattern of induction, with an IgM response preceding the rise of DENV-reactive IgG, and DENV-reactive IgG reaching significantly higher titers during secondary infection [24]. These characteristics, as well as the assumed importance of IgG-mediated ADE, have left the role of other serum antibody isotypes relatively unexamined. Notably, this includes IgA1, the second most prevalent antibody isotype in serum and one that has been

suggested to play a unique and non-redundant role in many viral infections. Most work on DENV-reactive serum IgA has focused on its potential as a diagnostic tool [25], with a small literature examining DENV-reactive serum IgA as a possible correlate of severe disease [26-29].

5 [0008] The role for IgA1 during primary dengue was recently described, where IgA1 appears to be the dominant isotype-switched antibody expressed by DENV-elicited plasmablasts [30, 31]. While IgA expressing plasmablasts were also observed following secondary dengue, they constituted a significantly smaller fraction of the total infection-elicited immune response [30, 31]. Importantly, the IgA1 antibodies
10 expressed by these DENV-elicited plasmablasts exhibited both DENV-binding and DENV-neutralization activity, comparable to what was observed for IgG isotype antibodies derived from contemporaneous samples [30].

[0009] Prior-art-of-interest includes U.S. Patent No. 9,073,981 entitled Dengue virus neutralizing antibodies and use thereof (herein incorporated by reference). However,
15 the reference is deficient in that it fails to identify the immunoglobulins of the present disclosure and uses thereof.

[0010] Accordingly, there is a continuing need for materials and methods for preventing flavivirus infection such as dengue virus without increasing the risk of antibody-dependent enhancement of infection.

20 SUMMARY

[0011] The present disclosure is based, in part, on the observation of milder symptoms and lower viral burden typically associated with primary dengue relative to secondary dengue, and that DENV-reactive IgA1 plays a role in limiting DENV propagation and potentially the immune-mediated enhancement of disease. Below, isotype-switched
25 antibodies show conversion of IgG1 to IgA1 does not impact the ability of a monoclonal antibody to either bind whole DENV virions or to neutralize DC-SIGN-dependent DENV infection of a susceptible cell line. Moreover, the inventors have found that IgG1 antibodies subjected to immunoglobulin class switching, wherein an Fc region include an IgA Fc domain or segment, provide DENV-reactive antibodies suitable for
30 prophylaxis for flavivirus infection such as dengue virus and are capable of treating flavivirus disease such as dengue disease, and/or mediating ADE associated therewith.

[0012] In embodiments, immunoglobulins of the present disclosure include antibodies including the heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain or segment, and cocktails of immunoglobulins including antibodies including the heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain or segment, which neutralize dengue virus infection without contributing to antibody-dependent enhancement of dengue virus infection. Accordingly, in one aspect of the invention, the present disclosure includes a human antibody, an antibody variant, or an antigen binding fragment thereof, that neutralize a flavivirus virus infection such as dengue virus, wherein the antibody, antibody variant, or antigen binding fragment does not contribute to antibody-dependent enhancement of dengue virus infection.

[0013] In embodiments, the present disclosure includes treatments wherein adding antibodies of the present disclosure such as DENV-reactive monoclonal IgA1 to either an enhancing concentration of monoclonal IgG1 or to an enhancing dilution of dengue-immune plasma, antagonizes ADE in a dose-dependent fashion.

[0014] In embodiments, the present disclosure relates to an isolated monoclonal antibody, including: a heavy chain having an amino acid sequence of SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain or segment. In embodiments, the IgA Fc domain is an IgA1 Fc domain.

[0015] In embodiments, the present disclosure relates to an isolated monoclonal antibody for targeting a fusion loop epitope of dengue virus, including: a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain; and a light chain having an amino acid sequence having at least 90% sequence identity to SEQ ID. NO. 2.

[0016] In some embodiments, the present disclosure relates to an isolated monoclonal antibody, including: a heavy chain including or consisting of an amino acid sequence of SEQ. ID NO. 1; and a light chain including or consisting of an amino acid sequence of SEQ ID NO: 2.

[0017] In some embodiments, the present disclosure relates to an isolated monoclonal antibody, including: a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain.

[0018] In some embodiments the present disclosure relates to a nucleic acid polymer encoding a monoclonal antibody, wherein the polymer includes or consists of SEQ. ID NO. 1 and/or a second nucleic acid polymer encoding a monoclonal antibody, wherein the second polymer includes or consists of SEQ. ID NO. 2.

- 5 [0019] In some embodiments, the present disclosure relates to a complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 1.

[0020] In some embodiments, the present disclosure relates to a complementary deoxynucleotide (cDNA) sequence including or consisting of a nucleic acid sequence
10 of SEQ ID NO: 3.

[0021] In some embodiments, the present disclosure relates to a complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 2.

- [0022] In some embodiments, the present disclosure relates to a complementary
15 deoxynucleotide (cDNA) sequence including or consisting of a nucleic acid sequence of SEQ ID NO: 4.

[0023] In some embodiments, the present disclosure relates to a method for preventing or treating a flavivirus virus infection such as dengue viral infection, the method including administering a therapeutically effective amount of the monoclonal
20 antibody of the present disclosure to a subject in need thereof under conditions effective to treat the viral infection.

[0024] In some embodiments, the present disclosure relates to a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including: administering a therapeutically effective amount of a monoclonal
25 antibody including a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is characterized as an isotypic commutation.

[0025] In some embodiments, the present disclosure relates to a method for
30 preventing or treating antibody-dependent enhancement of a viral infection, the method including: administering a therapeutically effective amount of a monoclonal antibody including a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is formed by class-

switch recombination.

[0026] In some embodiments, the present disclosure relates to a method for preventing or treating antibody-dependent enhancement of a flavivirus infection, the method including: administering a therapeutically effective amount of an immunoglobulin such as: a) a monoclonal antibody encoded by a DNA polymer of the present disclosure; b) an antibody encoded by one or more cDNAs of the present disclosure; or c) a monoclonal antibody of the present disclosure. In embodiments, the monoclonal antibody is disposed with a serum including one or more additional antibodies or fragments thereof directed against dengue virus or bind one or more epitopes of dengue virus.

[0027] In embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including administering a therapeutically effective amount of an antibody including a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection.

[0028] In embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including administering a therapeutically effective amount of a monoclonal antibody to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection, wherein the monoclonal antibody is characterized as an IgA isoform.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE LISTING

[0029] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[0030] Embodiments of the present disclosure, briefly summarized above and discussed in greater detail below, can be understood by reference to the illustrative embodiments of the disclosure depicted in the appended drawings. However, the appended drawings illustrate only typical embodiments of the disclosure and are therefore not to be considered limiting of scope, for the disclosure may admit to other equally effective embodiments.

[0031] FIGS. 1A, 1B and 1C depict an isotype conversion scheme of the present disclosure, DENV binding, and DENV neutralization capacity of VDB33 and VDB50 mAbs.

5 [0032] FIGS. 2A, 2B, 2C and 2D depict data indicating IgG1, but not IgA1, mediates ADE of DENV infection.

[0033] FIGS. 3A, 3B, 3C and 3D depict data and the fractional addition of DENV-reactive IgA1 significantly reduced the ADE activity observed in cultures containing either VDB33-IgG1 or VDB50-IgG1.

10 [0034] FIGS. 4A-4C depict Monoclonal IgA1 antagonizes ADE mediated by polyclonal DENV-immune plasma.

[0035] FIG. 5 depicts a gating scheme and representative plots from FlowNT assays.

[0036] FIG. 6 shows a gating scheme and representative plots from ADE assays.

[0037] FIGS. 7A, 7B and 7C show DENV-3 IgM, IgG, and IgA titers in DENV-immune plasma samples.

15 [0038] FIGS. 8A and 8B depict DENV-3 neutralization activity of DENV-immune plasma as assessed by FlowNT, and DENV-3 ADE activity of DENV-immune plasma as assessed by K562 infection.

[0039] FIG. 9 depicts Isotype distribution of plasmablasts captured during acute primary or secondary DENV infection by single cell RNA sequencing.

20 [0040] FIG. 10 shows isotype distribution of plasmablasts captured during acute secondary DENV infection in individuals that progressed to develop mild and severe dengue.

[0041] FIG. 11 depicts a proposed model for IgA antagonism of IgG-mediated DENV ADE.

25 [0042] FIG. 12 shows an analysis on samples collected around the day of fever abatement, prior to the critical phase of dengue where there is an increased probability of developing clinical manifestations of severe dengue.

[0043] FIGS. 13A, 13B, 13C, and 13D Example of IgG mediated enhancement of DENV infection in K562 cells expressing FcγR. Fold enhancement of infection calculated relative to infection level achieved in the absence of recombinant antibody

30 [0044] FIG. 14 depicts an annotated sequence of a synthesized antibody in accordance with the present disclosure.

[0045] FIG. 15 depicts an annotated sequence in accordance with the present disclosure.

[0046] FIG. 16 depicts DENV1_E_protein (Wp74, UniProt P17763.2) and target epitopes suitable for binding in accordance with the present disclosure.

[0047] FIG. 17 depicts VDB50_IgA_heavy_chain_AA and VDB50_IgA_light_chain_AA suitable for combination into an antibody suitable for use in accordance with the present disclosure.

[0048] FIG. 18 depicts DENV1_E_protein (Wp74, UniProt P17763.2) and target epitopes suitable for binding in accordance with the present disclosure.

[0049] SEQ ID NO:1 depicts a peptide sequence for VDB33_IgA1_HC.

[0050] SEQ ID NO:2 depicts a peptide sequence for VDB33_IgA1_LC.

[0051] SEQ ID NO:3 depicts a nucleotide sequence for VDB33-IgA1_heavy_chain.

[0052] SEQ ID NO:4 depicts a nucleotide sequence for VDB33-IgA1_light_chain.

[0053] SEQ ID NO:5 depicts DENV1_E_protein including VDB33 target epitopes: W101, G106, L107, F108.

[0054] SEQ ID NO: 6 depicts a peptide sequence for VDB50-IgG1_heavy_chain.

[0055] SEQ ID NO:7 depicts a peptide sequence for VDB50-IgG1_light_chain.

[0056] SEQ ID NO:8 depicts a peptide sequence for VDB50_IgA_heavy_chain_nt.

[0057] SEQ ID NO:9 depicts a peptide sequence for VDB50_IgA_light_chain_nt.

[0058] SEQ ID NO:10 depicts a peptide sequence for VDB50_IgA_heavy_chain_AA.

[0059] SEQ ID NO:11 depicts a peptide sequence for VDB50_IgA_light_chain_AA.

[0060] It is noted that the drawings of the disclosure are not necessarily to scale. The drawings are intended to depict only typical aspects of the disclosure, and therefore should not be considered as limiting the scope of the disclosure. In the drawings, like numbering represents like elements between the drawings.

DETAILED DESCRIPTION

[0061] The present disclosure relates to one or more immunoglobulins or functional fragments thereof, such as an antibody or functional fragment thereof having a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain or an IgA Fc segment. In embodiments, one or more non-IgA antibodies for targeting a flavivirus virus such as dengue virus are subjected to isotype-switching to include a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain. In embodiments, one or more immunoglobulins that neutralize flavivirus infection such as dengue and/or mitigate antibody-dependent

enhancement (ADE) of flavivirus virus infection such as ADE associated with secondary heterologous dengue infections formed by class switching or class-switch recombination are provided. In embodiments, wild-type IgA antibodies are suitable for use in accordance with the present disclosure.

- 5 [0062] Advantages of the immunoglobulins, antibodies, or functional fragments thereof include excellent prophylaxis and treatment of flavivirus disease such as dengue disease associated with one or more dengue virus strains.

DEFINITIONS

- 10 [0063] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

- [0064] As used herein, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus, for example, references to "a compound" include the use of one or more compound(s). "A step" of a method means
15 at least one step, and it could be one, two, three, four, five or even more method steps.

- [0065] As used herein the terms "about," "approximately," and the like, when used in connection with a numerical variable, generally refers to the value of the variable and to all values of the variable that are within the experimental error (e.g., within the 95% confidence interval [CI 95%] for the mean) or within $\pm 10\%$ of the indicated value,
20 whichever is greater.

- [0066] The term "antibody" as used herein refers to an immunoglobulin molecule capable of specific binding to a target antigen or biomarker, such as a carbohydrate, polynucleotide, lipid, polypeptide, peptide etc., via at least one antigen recognition site (also referred to as a binding site), located in the variable region of the immunoglobulin
25 molecule. In embodiments, the term "antibody" refers to a selective binding compound. In embodiments, antibodies, or functional fragments thereof may selectively bind or target any portion of one or more E proteins or fragments thereof of dengue virus, or a variant thereof. In embodiments, antibodies or functional fragments thereof refers to a compound that selectively binds to a fusion loop (FL) domain in a dengue viral
30 envelope (E) protein associated with fusion of the viral membrane to a cellular membrane.

[0067] As used herein, the terms "bind" and "binding" generally refer to the non-covalent interaction between a pair of partner molecules or portions thereof (e.g.,

antigenic protein- binding partner complexes) that exhibit mutual affinity or binding capacity. In embodiments, binding can occur such that the partners are able to interact with each other to a substantially higher degree than with other, similar substances. This specificity can result in stable complexes (e.g., antigenic protein-binding partner
5 complexes or bound biomarkers-of-interest) that remain bound during handling steps such as chromatography, centrifugation, filtration, and other techniques typically used for separations and other processes. In embodiments, the interaction between a target region of an antigenic protein and a binding partner that binds specifically thereto is a non-covalent interaction. In some instances, the interaction between a binding partner
10 and a non-target region of an antigenic protein is a non-covalent interaction. However, in other instances, the interaction between a binding partner and a non-target region of an antigenic protein may be a covalent interaction. In embodiments, a protein complex comprising the antigenic protein and the binding partner may be contacted with a chemical crosslinking reagent that causes covalent bonds between the
15 antigenic protein and the binding partner to be formed. In another example, the antigenic protein may contain a first reactive chemical moiety (handle) and the one or more binding partners may each contain a second reactive chemical moiety (handle), wherein the first and second chemical reactive moieties can react with each other to form a covalent bond. Exemplary reactive chemical moieties include those useable in
20 "click" chemistry, which is a class of biocompatible small molecule reactions commonly used in bioconjugation, allowing the joining of substrates of choice with specific biomolecules. Click chemistry is not a single specific reaction, but refers to a way of generating products that follow examples in nature, which also generates substances by joining small modular units. In one example, the antigenic protein may have a first
25 reactive chemical moiety such as a clickable handle like an azide, and the binding partner(s) could have a complementary reactive handle such as, for example a strained cyclooctyne, or vice versa. When these reactive chemical moieties come into proximity when the antigenic protein and the one or binding partners interact to form a protein complex, they can react with each other to form a covalently bond between
30 the proteins.

[0068] As used herein the term "cDNA" refers to a DNA molecule that can be prepared by reverse transcription from an RNA molecule obtained from a eukaryotic or prokaryotic cell, a virus, or from a sample solution. In embodiments, cDNA lacks introns or intron sequences that may be present in corresponding genomic DNA. In

embodiments, cDNA may refer to a nucleotide sequence that corresponds to the nucleotide sequence of an RNA from which it is derived. In embodiments, cDNA refers to a double-stranded DNA that is complementary to and derived from mRNA.

[0069] As used herein the term "competitive inhibitor" refers to one or more substances that binds to or blocks another substance from participating in a reaction. Non-limiting examples of competitive inhibitors of the present disclosure include one or more antibodies of the present disclosure or fragments thereof that bind an envelope protein epitope or Dengue virus viral envelope glycoprotein, E, or one or more fusion loops disposed therein. In embodiments, antibodies of the present disclosure target the E dimer epitope (EDE), readily exposed at an E dimer interface over a region of a conserved fusion loop. In embodiments, antibodies of the present disclosure target a conserved fusion loop (FL) domain among dengue virus strains in a viral envelope (E) protein associated with fusion of the viral membrane to a cellular membrane.

[0070] The terms "deoxyribonucleotide" and "DNA" refer to a nucleotide or polynucleotide including at least one ribosyl moiety that has an H at the 2' position of a ribosyl moiety. In embodiments, a deoxyribonucleotide is a nucleotide having an H at its 2' position.

[0071] As used herein the terms "drug," "drug substance," "active pharmaceutical ingredient," and the like, refer to a compound (e.g., immunoglobulin or antibody) that may be used for treating a subject in need of treatment.

[0072] As used herein the term "excipient" or "adjuvant" refers to any inert substance.

[0073] As used herein the terms "drug product," "pharmaceutical dosage form," "dosage form," "final dosage form" and the like, refer to a pharmaceutical composition that is administered to a subject in need of treatment and generally may be in the form of inhalers, tablets, capsules, sachets containing powder or granules, liquid solutions or suspensions, patches, and the like.

[0074] As used herein "dengue virus" refers to one of any of four related viruses: Dengue virus 1, 2, 3, and 4. In embodiments, dengue virus includes a single strand of RNA, or positive-sense *RNA* that can be directly translated into proteins. In embodiments, the viral genome encodes ten genes. In embodiments, the genome is translated as a single, long polypeptide and then cut into ten proteins. In embodiments, the dengue virus genome encodes three structural (capsid [C], membrane [M], and envelope [E]) and seven nonstructural (NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5) proteins. In embodiments, the dengue virus includes a highly conserved fusion

loop (FL) domain in the viral envelope (E) protein associated with fusion of the viral membrane to a cellular membrane.

[0075] As used herein the term "flavivirus" refers to any of a group of RNA viruses, mostly having arthropod vectors, that cause a number of serious human diseases including yellow fever, dengue, various types of encephalitis, and hepatitis C.

[0076] As used herein the term "fragment" means a polypeptide having one or more (e.g., several) amino acids absent from the amino and/or carboxyl terminus of a mature polypeptide or domain. In embodiments, a fragment is able to bind to Dengue virus viral envelope glycoprotein, E, or one or more fusion loops disposed therein. In 10 embodiments, a fragment is able to target fusion loop (FL) domain in the viral envelope (E) protein associated with fusion of the viral membrane to a cellular membrane. In embodiments, a fragment contains at least 70% to 99%, at least 90% to 99% or about 95 to 99% of the number of amino acids of the mature polypeptide of SEQ ID NO: 1 and/or SEQ ID NO:2.

15 [0077] By "hybridizable" or "complementary" or "substantially complementary" a nucleic acid (e.g. RNA, DNA) includes a sequence of nucleotides that enables it to non-covalently bind, i.e. form Watson-Crick base pairs and/or G/U base pairs, "anneal", or "hybridize," to another nucleic acid in a sequence-specific, antiparallel, manner (i.e., a nucleic acid specifically binds to a complementary nucleic acid) under 20 the appropriate in vitro and/or in vivo conditions of temperature and solution ionic strength. Standard Watson-Crick base-pairing includes adenine/adenosine (A) pairing with thymidine/thymine (T), A pairing with uracil/uridine (U), and guanine/guanosine (G) pairing with cytosine/cytidine (C). In addition, for hybridization between two RNA molecules (e.g., dsRNA), and for hybridization of a DNA molecule 25 with an RNA molecule (e.g., when a DNA target nucleic acid base pairs with a guide RNA, etc.): G can also base pair with U. For example, G/U base-pairing is partially responsible for the degeneracy (i.e., redundancy) of the genetic code in the context of tRNA anti-codon base-pairing with codons in mRNA. In embodiments, hybridization requires that the two nucleic acids contain complementary sequences, although 30 mismatches between bases are possible. The conditions appropriate for hybridization between two nucleic acids depend on the length of the nucleic acids and the degree of complementarity, variables well known in the art. The greater the degree of complementarity between two nucleotide sequences, the greater the value of the melting temperature (T_m) for hybrids of nucleic acids having those sequences.

Typically, the length for a hybridizable nucleic acid is 8 nucleotides or more (e.g., 10 nucleotides or more, 12 nucleotides or more, 15 nucleotides or more, 20 nucleotides or more, 22 nucleotides or more, 25 nucleotides or more, or 30 nucleotides or more). It is understood that the sequence of a polynucleotide need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, a polynucleotide may hybridize over one or more segments such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure or hairpin structure, a `bulge`, and the like). In embodiments, a polynucleotide can include 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, 99% or more, 99.5% or more, or 100% sequence complementarity to a target region within the target nucleic acid sequence to which it will hybridize. For example, an antisense nucleic acid in which 18 of 20 nucleotides of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. The remaining noncomplementary nucleotides may be clustered or interspersed with complementary nucleotides and need not be contiguous to each other or to complementary nucleotides. Percent complementarity between particular stretches of nucleic acid sequences within nucleic acids can be determined using any convenient method. Example methods include BLAST programs (basic local alignment search tools) and PowerBLAST programs (Altschul et al., J. Mol. Biol., 1990, 215, 403-410; Zhang and Madden, Genome Res., 1997, 7, 649-656) or by using the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), e.g., using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489).

[0078] The term "isolated" means a substance in a form or environment that does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally occurring substance, (2) any substance such as a variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated.

[0079] The term "mature polypeptide" means a polypeptide in its final form following

translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, etc.

[0080] The term "nucleotide" refers to a ribonucleotide or a deoxyribonucleotide or modified form thereof, as well as an analog thereof.

- 5 [0081] The terms "peptide," "polypeptide," and "protein" are used interchangeably herein, and refer to a polymeric form of amino acids of any length, which can include coded and non-coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified peptide backbones.

- [0082] The terms "polynucleotide" and "nucleic acid," used interchangeably herein, 10 refer to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. Thus, terms "polynucleotide" and "nucleic acid" encompass single-stranded DNA; double-stranded DNA; multi-stranded DNA; single-stranded RNA; double-stranded RNA; multi-stranded RNA; genomic DNA; cDNA; DNA-RNA hybrids; and a polymer including purine and pyrimidine bases or other natural, 15 chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The terms "polynucleotide" and "nucleic acid" should be understood to include, as applicable to the embodiments being described, single-stranded (such as sense or antisense) and double-stranded polynucleotides.

- [0083] The terms "sequence identity", "identity" and the like as used herein with 20 respect to polynucleotide or polypeptide sequences refer to the nucleic acid residues or amino acid residues in two sequences that are the same when aligned for maximum correspondence over a specified comparison window. Thus, "percentage of sequence identity", "percent identity" and the like refer to the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the 25 polynucleotide or polypeptide sequence in the comparison window may include additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage may be calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the 30 number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the results by 100 to yield the percentage of sequence identity.

[0084] In embodiments, when calculating sequence identity between a DNA sequence and an RNA sequence, T residues of the DNA sequence align with, and can be

considered "identical" with, U residues of the RNA sequence. For purposes of determining "percent complementarity" of first and second polynucleotides, one can obtain this by determining (i) the percent identity between the first polynucleotide and the complement sequence of the second polynucleotide (or vice versa), for example, and/or (ii) the percentage of bases between the first and second polynucleotides that would create canonical Watson and Crick base pairs.

[0085] In embodiments, the degree of sequence identity between a query sequence and a reference sequence is determined by: 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty; 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment; and 3) dividing the number of exact matches with the length of the reference sequence. In one embodiment, the degree of sequence identity between a query sequence and a reference sequence is determined by: 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty; 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid; or nucleotide in the two aligned sequences on a given position in the alignment; and 3) dividing the number of exact matches with the length of the longest of the two sequences. In some embodiments, the degree of sequence identity refers to and may be calculated as described under "Degree of Identity" in U.S. Patent No. 10,531,672 starting at Column 11, line 56. U.S. Patent No. 10,531,672 is incorporated by reference in its entirety. In embodiments, an alignment program suitable for calculating percent identity performs a global alignment program, which optimizes the alignment over the full-length of the sequences. In embodiments, the global alignment program is based on the Needleman-Wunsch algorithm (Needleman, Saul B.; and Wunsch, Christian D. (1970), "*A general method applicable to the search for similarities in the amino acid sequence of two proteins*", Journal of Molecular Biology 48 (3): 443-53). Examples of current programs performing global alignments using the Needleman-Wunsch algorithm are EMBOSS Needle and EMBOSS Stretcher programs, which are both available on the world wide web at www.ebi.ac.uk/Tools/psa/. In some embodiments a global alignment program uses the Needleman-Wunsch algorithm, and the sequence identity is calculated by identifying the number of exact matches identified by the program divided by the "alignment

length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences. In embodiments, the mafft alignment program is suitable for use herein.

[0086] The term "substantially purified," as used herein, refers to a component of interest that may be substantially or essentially free of other components which normally accompany or interact with the component of interest prior to purification. In embodiments, a component of interest may be "substantially purified" when the preparation of the component of interest contains less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, less than about 5%, less than about 4%, less than about 3%, less than about 2%, or less than about 1% (by dry weight) of contaminating components. Thus, a "substantially purified" component of interest may have a purity level of about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or greater. In embodiments, a component of interest includes a virus-of-interest, such as dengue virus or a variant thereof.

[0087] "Substantially similar" refers to nucleic acid molecules wherein changes in one or more nucleotide bases result in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. "Substantially similar" also refers to nucleic acid molecules wherein changes in one or more nucleotide bases do not affect the ability of the nucleic acid molecule to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid molecules of the instant disclosure (such as deletion or insertion of one or more nucleotide bases) that do not substantially affect the functional properties of the resulting transcript vis-a-vis the ability to mediate alteration of gene expression by antisense or co-suppression technology or alteration of the functional properties of the resulting protein molecule. The disclosure encompasses more than the specific exemplary sequences.

[0088] As used herein the term "pharmaceutically acceptable" substances refers to those substances, such as e.g., antibodies of the present disclosure and functional fragments thereof, which are within the scope of sound medical judgment suitable for use in contact with the tissues of subjects without undue toxicity, irritation, allergic response, and the like, and effective for their intended use.

[0089] As used herein the term "pharmaceutical composition" refers to the combination of one or more drug substances such as e.g., antibodies of the present disclosure or

functional fragments thereof and one or more excipients and one or more pharmaceutically acceptable vehicles with which the one or more antibodies or functional fragments thereof is administered to a subject.

[0090] As used herein the term "pharmaceutically acceptable vehicle" refers to a diluent, adjuvant, excipient or carrier with which a compound, such as e.g., an antibody of the present disclosure or functional fragment thereof, is administered.

[0091] As used herein the term "prevent", "preventing" and "prevention" of dengue means (1) reducing the risk of a patient who is not experiencing symptoms of dengue virus infection from developing dengue, or (2) reducing the frequency of, the severity of, or a complete elimination of dengue symptoms already being experienced by a subject.

[0092] The term "prophylactically effective amount," as used herein, refers to that amount of a composition, such as e.g., antibody of the present disclosure or a functional fragment thereof, administered to a subject which will relieve to some extent one or more of the symptoms of a disease, likelihood of becoming diseased, or condition or disorder being treated. In such prophylactic applications, such amounts may depend on the subject's state of health, weight, and the like. It is considered well within the skill of the art for one to determine such prophylactically effective amounts by routine experimentation, including, but not limited to, a dose escalation clinical trial.

[0093] The term "recombinant" when used herein to characterize a DNA sequence such as a plasmid, vector, or construct refers to an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis and/or by manipulation of isolated segments of nucleic acids by genetic engineering techniques.

[0094] As used herein the term "subject" includes humans, animals or mammals. The terms "subject" and "patient" may be used interchangeably herein.

[0095] As used herein, the term "selective binding compound" refers to a compound that selectively binds to any portion of one or more target proteins.

[0096] As used herein, the term "target activity" refers to a biological activity capable of being modulated by a selective modulator. Certain exemplary target activities include, but are not limited to, binding affinity, signal transduction, enzymatic activity, tumor growth, inflammation or inflammation-related processes, and amelioration of one or more symptoms associated with a disease or condition.

[0097] As used herein, the term "target protein" refers to a molecule or a portion of a protein capable of being bound by a selective binding compound. In certain

embodiments, a target protein is one or more E proteins or fragments thereof of dengue virus (including variants thereof). In embodiments, a target protein is a fusion loop (FL) domain in a dengue viral envelope (E) protein associated with fusion of the viral membrane to a cellular membrane.

5 [0098] As used herein the term "therapeutically effective amount" means the amount of a compound that, when administered to a subject for treating or preventing flavivirus infection such as a dengue virus infection, is sufficient to effect such treatment or prevention of flavivirus disease such as dengue and related symptoms. A "therapeutically effective amount" can vary depending, for example, on the compound,
10 the severity of the dengue infection, the etiology of the dengue infection, one or more prior dengue infections, the age of the subject to be treated, comorbidities of the subject to be treated, existing health conditions of the subject, and/or the weight of the subject to be treated. A "therapeutically effective amount" is an amount sufficient to alter the subjects' natural state.

15 [0099] As used herein the term "treat", "treating" and "treatment" of flavivirus disease such as dengue means an intervention for reducing the frequency of symptoms of flavivirus disease such as dengue, eliminating the symptoms of flavivirus disease such as dengue, avoiding or arresting the development of symptoms of flavivirus disease such as dengue, ameliorating or curing an existing or undesirable symptom caused by
20 flavivirus disease such as dengue, and/or reducing the severity of symptoms of flavivirus disease such as dengue.

[00100] In embodiment the term "variant" means a polypeptide including an alteration, i.e., a substitution, insertion, and/or deletion, at one or more (e.g., several) positions. A substitution means replacement of the amino acid occupying a position
25 with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding one or more (e.g., several) amino acids, e.g., 1-10 amino acids, adjacent to the amino acid occupying a position.

[00101] General methods in molecular and cellular biochemistry can be found in such standard textbooks as Molecular Cloning: A Laboratory Manual, 3rd Ed. (Sambrook et al., HaRBor Laboratory Press 2001); Short Protocols in Molecular
30 Biology, 4th Ed. (Ausubel et al. eds., John Wiley & Sons 1999); Protein Methods (Bollag et al., John Wiley & Sons 1996); Nonviral Vectors for Gene Therapy (Wagner et al. eds., Academic Press 1999); Viral Vectors (Kapliff & Loewy eds., Academic Press 1995); Immunology Methods Manual (I. Lefkovits ed., Academic Press 1997);

and Cell and Tissue Culture: Laboratory Procedures in Biotechnology (Doyle & Griffiths, John Wiley & Sons 1998), the disclosures of which are incorporated herein by reference.

[00102] Before embodiments are further described, it is to be understood that
5 this disclosure is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

[00103] Where a range of values is provided, it is understood that each
10 intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or
15 both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[00104] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or
20 equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

25

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

[00105] In embodiments, the present disclosure includes one or more immunoglobulins or functional fragments thereof, such as an antibody or functional fragment thereof having a heavy chain or a segment of the heavy chain including an
30 Fc region characterized as IgA Fc domain or an IgA Fc segment suitable for prophylaxis and treatment of flavivirus disease such as dengue disease associated with one or more dengue virus strains.

[00106] In embodiments, the present disclosure includes an antibody, including: a heavy chain having an amino acid sequence of SEQ. ID NO. 1, wherein the heavy

chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain. In embodiments, the IgA Fc domain is an IgA1 Fc domain, or IgA2 Fc domain. In embodiments, the antibody is characterized as isolated and/or monoclonal. In embodiments, the antibody is chimeric or humanized. In some embodiments, an isolated monoclonal antibody includes: a light chain having an amino acid sequence of SEQ ID NO. 2. In embodiments, an antibody, such as an isolated monoclonal antibody binds an epitope of a Deng virus, wherein the epitope is characterized as a loop.

[00107] In embodiments, the Fc region characterized as IgA Fc domain includes or consists of the amino acids of SEQ ID NO: 1. In embodiments, the Fc region characterized as IgA Fc domain includes or consists of the amino acids between S31 and Y475 of SEQ ID NO. 1, an amino acid segment between G51 and Y475 of SEQ ID NO: 1, an amino acid segment between L101, or an amino acid segment between A181 and Y475, wherein SEQ ID NO: 1 is used for numbering.

[00108] In embodiments, an antibody, such as an isolated monoclonal antibody or functional fragment thereof is suitable for targeting a fusion loop epitope of dengue virus. In embodiments, such antibodies include a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain; and a light chain having an amino acid sequence having at least 90% sequence identity to SEQ ID. NO. 2. In embodiments, the heavy chain includes an amino acid sequence having at least 95%, 97%, 98%, 99% sequence identity to SEQ. ID NO. 1. In embodiments, the light chain includes an amino acid sequence having at least 95%, 97%, 98%, 99% sequence identity to SEQ. ID NO. 2.

[00109] In embodiments, the immunoglobulins of the present disclosure include an isolated monoclonal antibody, including: a heavy chain consisting of an amino acid sequence of SEQ. ID NO. 1; and a light chain consisting of an amino acid sequence of SEQ ID NO: 2. In embodiments, the isolated monoclonal antibody binds a fusion loop epitope (FLE) of a Dengue virus.

[00110] In embodiments, the immunoglobulins of the present disclosure include an isolated monoclonal antibody, including: a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain includes an Fc region characterized as IgA Fc domain. In embodiments, the isolated monoclonal antibody binds an epitope of a

Dengue virus, wherein the epitope is characterized as a loop.

[00111] In embodiments, the present disclosure includes a nucleic acid or nucleic acid polymer encoding a monoclonal antibody, wherein the polymer includes or consists of SEQ. ID NO. 1. In embodiments, the present disclosure includes a nucleic acid polymer encoding a monoclonal antibody, wherein the polymer includes or consists of SEQ. ID NO. 2.

[00112] In embodiments, the present disclosure includes a complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence or antibody of the present disclosure. In embodiments, the present disclosure includes a complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 1. In embodiments, the present disclosure includes a complementary deoxynucleotide (cDNA) sequence including or consisting of a nucleic acid sequence of SEQ ID NO: 3.

[00113] In embodiments, the present disclosure includes a complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 2. In embodiments, the present disclosure includes a complementary deoxynucleotide (cDNA) sequence including or consisting of a nucleic acid sequence of SEQ ID NO: 4.

[00114] In some embodiments, the present disclosure includes a method for preventing or treating a dengue viral infection, the method including administering a therapeutically effective amount of the immunoglobulins described herein, including a monoclonal antibody, to a subject in need thereof under conditions effective to treat the viral infection. In embodiments, the method is used for preventing or treating antibody-dependent enhancement of a viral infection.

[00115] In some embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including: administering a therapeutically effective amount of a monoclonal antibody including a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is characterized as an isotypic commutation. In embodiments, the viral infection is any one of respiratory syncytial virus (RSV), influenza, and coronavirus infection, or a combination thereof.

In embodiments, the viral infection is a flavivirus infection. In embodiments, the flavivirus infection is a Zika virus, or a dengue virus.

[00116] In some embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including: administering a therapeutically effective amount of a monoclonal antibody including a heavy chain or a segment of the heavy chain including an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is formed by class-switch recombination.

10 [00117] In some embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a flavivirus infection, the method including administering a therapeutically effective amount of: a) a monoclonal antibody encoded by a DNA polymer of the present disclosure; b) a monoclonal antibody encoded by a cDNA of the present disclosure; or c) a monoclonal antibody of the present disclosure. In embodiments, the monoclonal antibody is disposed with a serum including one or more additional antibodies or fragments thereof directed against dengue virus or bind one or more epitopes of dengue virus.

[00118] In some embodiments, the present disclosure to IgA1 class-switched monoclonal antibody therapy, which may be an effective therapy for preventing dengue virus infection without risking immune mediated enhancement of disease due to waning antibody titers. Further, the present invention relates to the fields of immunology and virology, including methods of assessing monoclonal antibody and plasma DENV-reactivity using DENV-capture ELISA protocol, assessing neutralizing titers of monoclonal antibodies and heat-inactivated plasma using a neutralization assay, and quantification of DENV-3 infection using ADE assay.

25 [00119] In some embodiments, the present disclosure includes an Immunoglobulin A1 (IgA1) monoclonal antibody; an Immunoglobulin G1 (IgG1) monoclonal antibody; a method of administering a therapeutic treatment for DENV; a method of making a therapeutic treatment for DENV; isolated isotype-switched monoclonal antibodies comprising vdb33 IgA1 and vdb50 IgG1; a method of quantification of DENV-3 infection using *in vitro* ADE assay; and an IgA1 class-switched monoclonal antibody therapy, and combinations of these.

[00120] In some embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the

method including: administering a therapeutically effective amount of an antibody including a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection. In
5 embodiments, the viral infection is any one of respiratory syncytial virus (RSV), influenza, and coronavirus infection, or a combination thereof. In embodiments, the viral infection is a flavivirus infection. In embodiments, the flavivirus infection is a Zika virus. In embodiments, the flavivirus infection is a dengue virus. In embodiments, the method further includes increasing the concentration of antibodies characterized as
10 IgA in a serum, such as a pharmaceutically acceptable serum, including one or more additional antibodies. In embodiments, the antibody is a wild-type IgA isoform that binds dengue protein E. In embodiments the antibody is monoclonal and or characterized as pharmaceutically acceptable. In embodiments, the antibody is synthetic, formed by recombinant technology, or characterized as isolated and/or
15 pharmaceutically acceptable. In embodiments, antibodies of the present disclosure are disposed within a pharmaceutically acceptable compositions and may include additional excipients. In embodiments, antibodies of the present disclosure are disposed within a pharmaceutically acceptable vehicle suitable for administration to a subject in need thereof.

20 [00121] In embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the method including administering a therapeutically effective amount of a monoclonal antibody to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection, wherein the monoclonal
25 antibody is characterized as an IgA isoform. In embodiments, the monoclonal antibody includes a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain. In embodiments, the Fc region characterized as IgA Fc domain includes or consists of the amino acids of SEQ ID NO: 1. In embodiments, the Fc region characterized as IgA Fc domain includes or consists of the amino acids
30 between S31 and Y475 of SEQ ID NO. 1, an amino acid segment between G51 and Y475 of SEQ ID NO: 1, an amino acid segment between L101, or an amino acid segment between A181 and Y475, wherein SEQ ID NO: 1 is used for numbering.

[00122] In embodiments, the present disclosure includes a method for preventing or treating antibody-dependent enhancement of a viral infection, the

method including administering a therapeutically effective amount of a polypeptide to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection, wherein the polypeptide is characterized as an IgA antibody isoform. In embodiments, the polypeptide is an immunoglobulin and includes a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain.

EXAMPLE I

Materials and Methods

[00123] **Viruses:** DENV-3 (strain CH53489) propagated in Vero cells were utilized for ELISA, FlowNT50, and ADE assays. Virus for ELISA was purified by ultracentrifugation through a 30% sucrose solution and the virus pellet was resuspended in PBS.

[00124] **Cell lines:** Human K562 cells were maintained in IMDM supplemented with 10% FBS, penicillin, and streptomycin. U937-DC-SIGN cells were maintained in RPMI supplemented with 10% FBS, L-glutamine, penicillin, and streptomycin.

[00125] **Monoclonal antibodies and serum:** The variable regions from the heavy and light chains were codon optimized, synthesized *in vitro* and subcloned into a pcDNA3.4 vector containing the human IgG1 or IgA1 Fc region by a commercial partner (Genscript). Transfection grade plasmids were purified by maxiprep and transfected into a 293-6E expression system. Cells were grown in serum-free FreeStyle 293 Expression Medium (Thermo Fisher), and the cell supernatants collected on day 6 for antibody purification. Following centrifugation and filtration, the cell culture supernatant was loaded onto an affinity purification column, washed, eluted, and buffer exchanged to the final formulation buffer (PBS). Antibody lot purity was assessed by SDS-PAGE, and the final concentration determined by 280 nm absorption. The clonotype information for all monoclonal antibodies generated as part of this study is listed in **Table 1**.

[00126] **Table 1. Sequence information of DENV-reactive monoclonal antibodies**

30

Clone name	VDB33	VDB50
Parental Isotype	IgG1	IgA1
Infecting Serotype	DENV-3	DENV-1
Primary/Secondary	Secondary	Primary
H_c CDR3aa	CARLLQYKWNWLFDPW	CAKASQMATVFIDYW
H_c V	IGHV4-39*01	IGHV3-23*03
H_c D	IGHD1-7*01	IGHD5-24*01
H_c J	IGHJ5*02	IGHJ4*02
H_c Total SHM	26	13
L_c CDR3aa	CQVWDSDDHPVF	CQSYDSSLGGVF
L_c V	IGLV3-21*03	IGLV1-40*01
L_c J	IGLJ3*02	IGLJ3*02
L_c Total SHM	14	8
Target residues	W101, G106, L107, F108	G100, W101, F108
E protein epitope	Fusion loop	Fusion loop

[00127] Dengue IgG antibody positive plasma was purchased from SeraCare. Donor ID and batch numbers are shown in Supplemental Table 1.

[00128] **Supplemental Table 1. Dengue immune plasma used in this study**

Supplier	Product #	Donor ID	Batch number
SeraCare	0325-0014	BD250524	10127363
SeraCare	0325-0014	BD250525	10127364
SeraCare	0325-0014	BD250535	10127374
SeraCare	0325-0014	BD250543	10127383

5

[00129] **DENV- capture ELISA:** Monoclonal antibody and plasma DENV-reactivity was assessed using a 4G2 DENV capture ELISA protocol. In short, 96 well NUNC MaxSorb flat-bottom plates were coated with 2 µg/ml flavivirus group-reactive mouse monoclonal antibody 4G2 (Envigo Bioproducts, Inc.) diluted in borate saline

buffer. Plates were washed and blocked with 0.25% BSA + 1% Normal Goat Serum in PBS after overnight incubation. DENV-3 (strain CH53489) diluted in blocking buffer was captured for 2 hr, followed by extensive washing with PBS + 0.1% Tween 20. Serially diluted monoclonal antibody samples were incubated for 1 hr at RT on the captured virus, and DENV-specific antibody binding quantified using anti-human IgG HRP (Sigma-Aldrich, SAB3701362). Secondary antibody binding was quantified using the TMB Microwell Peroxidase Substrate System (KPL, cat. #50-76-00) and Synergy HT plate reader (BioTek, Winooski, VT). Antibody data were analyzed by nonlinear regression (One site total binding) to determine EC50 titers in GraphPad Prism 8 (GraphPad Software, La Jolla, CA).

[00130] **Neutralization Assay:** Neutralizing titers of monoclonal antibodies and heat-inactivated plasma were assessed using a flow cytometry-based neutralization assay in U937 cells expressing DC-SIGN as previously described [32, 33]. Four-fold dilutions of antibody or sera were mixed with an equal volume of virus diluted to a concentration to achieve 10%–15% infection of U937-DC-SIGN cells in the absence of antibody. The antibody/virus mixture was incubated for 1 h at 37 °C, after which an equal volume of medium (RPMI-1640 supplemented with 10% FBS, 1% penicillin/streptomycin, 1% l-glutamine (200 mM) containing 5×10^4 U937-DC-SIGN cells was added to each well and incubated 18–20 hr overnight in a 37 °C, 5% CO₂, humidified incubator. Following overnight incubation, the cells were fixed with IC Fixation Buffer (Invitrogen, 00-82222-49), permeabilized using IC Permeabilization Buffer (Invitrogen, 00-8333-56) and immunostained with flavivirus group-reactive mouse monoclonal antibody 4G2 (Envigo Bioproducts, Inc.), and secondary polyclonal goat anti-mouse IgG PE-conjugated antibody (#550589, BD Biosciences). The percentage of infected cells were quantified on a BD Accuri C6 Plus flow cytometer (BD Biosciences). Data were analyzed by nonlinear regression to determine 50% neutralization titers in GraphPad Prism 8 (GraphPad Software, La Jolla, CA).

[00131] **ADE Assay:** *In vitro* antibody-dependent enhancement (ADE) of DENV-3 infection was quantified as previously described [30, 34]. Four-fold serial dilutions of antibody or heat-inactivated sera were incubated with virus (in sufficient amounts to infect 10%–15% of U937-DC-SIGN cells) at a 1:1 ratio for 1 h at 37 °C. This mixture was then added to a 96-well plate containing 5×10^4 K562 cells per well in duplicate. Cells were cultured for 18–20 hr overnight in a 37 °C, 5% CO₂, humidified incubator. Processing and quantification continued as outlined in the FlowNT50 methods.

[00132] **Statistical Analysis:** All statistical analysis was performed using GraphPad Prism 8 Software (GraphPad Software, La Jolla, CA). A P-value < 0.05 was considered significant.

Results

- 5 [00133] **DENV binding and neutralizing is unaffected by antibody Fc isotype.** To assess the potential contribution of DENV-reactive IgA1 to a functional anti-DENV humoral immune response, two pairs of DENV-reactive monoclonal antibodies with either an IgG1 or an IgA1 Fc domain (**Figure 1A**). Both mAbs selected for this analysis were previously determined to bind the fusion loop of the DENV E protein and to react with all 4 DENV serotypes [30]. However, VDB33 was initially identified as an IgG1 clone, while VDB50 was discovered as an IgA1 clone (**Table 1**). This cross-conversion strategy was chosen so as to determine if the native Fc configuration of a given antibody influenced its functionality as either an IgG1 or IgA1 protein product.
- 10 [00134] The DENV-binding capacity of the IgG1 and IgA1 versions of VDB33 and VDB50 was initially assessed with a DENV virion-capture ELISA. For this analysis, DENV-3 was chosen as the prototypic DENV serotype as previous work demonstrated that the IgG1 versions of both VDB33 and VDB50 exhibited significant DENV-3 reactivity [30]. It was found that both VDB33 and VDB50 exhibited potent DENV-3 binding activity with VDB33 demonstrating ~200 fold higher affinity for DENV-3 than VDB50 (**FIG. 1A, Table 2**).
- 15 [00134] The DENV-binding capacity of the IgG1 and IgA1 versions of VDB33 and VDB50 was initially assessed with a DENV virion-capture ELISA. For this analysis, DENV-3 was chosen as the prototypic DENV serotype as previous work demonstrated that the IgG1 versions of both VDB33 and VDB50 exhibited significant DENV-3 reactivity [30]. It was found that both VDB33 and VDB50 exhibited potent DENV-3 binding activity with VDB33 demonstrating ~200 fold higher affinity for DENV-3 than VDB50 (**FIG. 1A, Table 2**).
- 20 [00134] The DENV-binding capacity of the IgG1 and IgA1 versions of VDB33 and VDB50 was initially assessed with a DENV virion-capture ELISA. For this analysis, DENV-3 was chosen as the prototypic DENV serotype as previous work demonstrated that the IgG1 versions of both VDB33 and VDB50 exhibited significant DENV-3 reactivity [30]. It was found that both VDB33 and VDB50 exhibited potent DENV-3 binding activity with VDB33 demonstrating ~200 fold higher affinity for DENV-3 than VDB50 (**FIG. 1A, Table 2**).

Table 2. Functional characteristics isotype-switched monoclonal antibodies

Clone name	VDB33 IgG	VDB33 IgA	VDB50 IgG	VDB50 IgA
Kd (ng/mL)	0.3962	0.8296	19.92	31.12
IC50 (ug/mL)	0.5339	0.3139	0.2391	0.2439

- [00135] However, the DENV-binding capacity of the two mAbs was not impacted by their conversion to either an IgG1 or IgA1 format (**FIG. 1A, Table 2**). Furthermore, this cross-conversion of VDB33 and VDB50 to either an IgG1 or IgA1 format minimally
- 25 [00135] However, the DENV-binding capacity of the two mAbs was not impacted by their conversion to either an IgG1 or IgA1 format (**FIG. 1A, Table 2**). Furthermore, this cross-conversion of VDB33 and VDB50 to either an IgG1 or IgA1 format minimally

impacted the DENV-3 neutralization activity of the clones when assessed using a flow cytometry-based neutralization assay (FIG. 1B, Table 2). These results indicate that both IgG1 and IgA1 isotype antibodies are equally capable of binding and neutralizing DENV, reaffirming that antibody epitope/paratope interactions occur independently of an antibody's F_c domain.

[00136] **DENV-reactive IgA1 is incapable of mediating ADE.** Having demonstrated that the antigen binding and neutralization capacity of DENV-reactive monoclonal antibodies is negligibly impacted by the isotype of the construct, determination of whether the infection-enhancing capability of these antibodies was impacted by their isotype conversion was investigated. To this end, a K562-based ADE assay was used, wherein antibody/DENV immune complexes were pre-formed and added to the F_c-receptor expressing K652 cell line to assess the ability of defined antibody complexes to enhance DENV infection.

[00137] More specifically, FIGS. 1A and FIG. 1B depict an isotype conversion scheme of the present disclosure, DENV binding, and DENV neutralization capacity of VDB33 and VDB50 mAbs. FIG. 1A depicts a schematic of isotype conversion of VDB33 and VDB50 from respective parental isotypes, indicating conservation of antigen-binding domains and alteration of F_c domains. FIG. 1B depicts DENV-3 binding capability of VDB33-IgG1, VDB33-IgA1, VDB50-IgG, and VDB50-IgA measured by DENV virus-capture ELISA. FIG. 1C depicts DENV-3 neutralization capability of VDB33-IgG, VDB33-IgA, VDB50-IgG, and VDB50-IgA as assessed by FlowNT. Neutralization data are presented as a percent of the positive (no neutralizing mAb) control for each replicate. Error bars +/- SEM.

[00138] The IgG1 versions of both VDB33 and VDB50 exhibited potent infection-enhancing activity in the K562 ADE assay, with both antibodies capable of facilitating DENV infection/enhancement in a dose-dependent fashion (See FIG. 2A and FIG. 2B). Consistent with their relative EC₅₀/IC₅₀ values, VDB33-IgG1 exhibited notably higher ADE activity than VDB50-IgG1, but with the peak of ADE activity occurring at a similar antibody concentration. However, no infection enhancement was observed when the same assay was performed with either VDB33-IgA1 or VDB50-IgA1 (See FIG. 2A and FIG. 2B). This was despite the fact that these IgA1 isotype antibodies exhibit nearly identical virus binding and neutralization activity as their IgG1 counterparts, underlining the obligate role of an antibody's F_c domain in determining the ADE potential of an antibody.

[00139] More specifically, FIGS. 2A-2D depict data indicating IgG1, but not IgA1, mediates ADE of DENV infection. FIG. 2A depicts ADE activity of VDB33-IgG and VDB33-IgA against DENV-3 in K562 cells. FIG. 2B depicts AUC values of 7 independent replicates of DENV-3 ADE assay with VDB33-IgG and VDB33-IgA. FIG. 2C depicts ADE activity of VDB50-IgG and VDB50-IgA against DENV-3 in K562 cells. FIG. 2D depicts AUC values of 7 independent replicates of DENV-3 ADE assay with VDB50-IgG and VDB50-IgA. Error bars +/- SEM. ** $p < 0.01$, **** $p < 0.0001$, unpaired t test.

[00140] **DENV-reactive IgA1 antagonizes IgG1 mediated enhancement of DENV infection.** In light of the inability of VDB33-IgA1 and VDB50-IgA1 to facilitate ADE of DENV-3, how DENV-reactive IgG1 and IgA1 behave in a polyclonal/competitive setting was determined. IgG1 and IgA1 antibodies are never found in isolation in a dengue immune individual, so determining how these antibodies function in a complex/poly-immune setting is critical for understanding their potential contribution to function anti-DENV immunity.

[00141] The same K562 ADE assay as previously described but used a fractional IgG1/IgA1 replacement strategy wherein the total amount of antibody remained the same across the different titration schemes but the ratio of IgG1 to IgA1 was varied from 100:0 to 0:100. The fractional addition of DENV-reactive IgA1 significantly reduced the ADE activity observed in cultures containing either VDB33-IgG1 or VDB50-IgG1 (See e.g., FIGS. 3A-3D). While both VDB33-IgA1 and VDB50-IgA1 were capable of antagonizing IgG1-mediated ADE of DENV-3, the highly avid yet non-enhancing VDB33-IgA1 antibody was capable of dramatically blunting IgG1-mediated ADE even when used at low fractional concentrations. Of note, the addition of DENV-reactive IgA1 to these ADE assays does not appear to shift the antibody dilution at which maximal ADE activity is observed for any of the cultures. Rather, the addition of DENV-reactive IgA1 reduces the magnitude of infection achieved at any given antibody dilution. These results are consistent with IgA1 actively antagonizing IgG1 mediated ADE by competing with DENV-reactive IgG1 for the same viral epitopes.

[00142] More specifically, it is noted that FIGS. 3A-3D depict homotypic and heterotypic monoclonal IgA1 antagonizes IgG-mediated antibody-dependent enhancement. FIG. 3A depicts DENV-3 ADE activity of VDB33-IgG when antagonized with VDB33-IgA. Total antibody concentration for each dilution point was held constant, with varying ratios of VDB33-IgG and VDB33-IgA as indicated. AUC of each

ADE titration was calculated and normalized to that of the 100% IgG condition. FIG. 3B depicts DENV-3 ADE activity of VDB33-IgG when antagonized with VDB50-IgA. The AUC of each ADE titration was calculated and normalized to that of the 100% VDB33-IgG condition. FIG. 3C depicts DENV-3 ADE activity of VDB50-IgG when antagonized with VDB33-IgA. AUC of each ADE titration was calculated and normalized to that of the 100% VDB50-IgG condition FIG. 3D depicts DENV-3 ADE activity of VDB50-IgG when antagonized with VDB50-IgA. AUC of each ADE titration was calculated and normalized to that of the 100% VDB33-IgG condition. Blue = 100% IgG / 0% IgA. Green = 90% IgG1 / 10% IgA1. Orange = 50% IgG1 / 50% IgA1. Red = 0% IgG1 / 100% IgA1. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ 1-way ANOVA with Dunnett correction for multiple comparisons

[00143] **DENV-reactive IgA1 antagonizes DENV-immune serum mediated enhancement of DENV infection.** A limitation of the analysis presented thus far is that all the monoclonal antibodies used in this analysis have the same antigen specificity; namely the fusion loop of the DENV E protein. Therefore, it is unclear what impact—if any—DENV-reactive IgA1 would have in the presence of a polyclonal IgG1 repertoire of divergent DENV antigen specificity. Therefore, we endeavored to determine how the presence of either VDB33-IgA1 or VDB50-IgA1 impacts the infection-enhancing potential of polyclonal/DENV-immune serum.

[00144] Plasma from DENV-immune donors were screened to identify samples with both high DENV-3 reactive IgG titers by ELISA as well as DENV-3 enhancing activity in the K562 ADE assay. Samples from four subjects were selected for additional analysis based on these criteria (See FIGS. 4A, 4B, and 4C, FIGS. 7A, 7B, and 7C, and FIGS. 8A and 8B).

[00145] VDB33-IgA1 or VDB50-IgA1 were then titrated into cultures containing this enhancing DENV-immune plasma to determine if IgA1 isotype monoclonal antibodies could antagonize polyclonal enhancement of DENV-3 infection.

[00146] Consistent with what was observed with IgG1 monoclonal antibodies, the addition of VDB33-IgA1 or VDB50-IgA1 significantly suppressed ADE-mediated K562 infection with DENV-3 (FIG. 4B, FIG. 4C). The additional of DENV-reactive IgA1 in these assays suppressed ADE-mediated infection by 75%-90% in a dose-dependent fashion, a result consistent with the concept that IgG antibodies targeting the fusion loop of the DENV E protein are particularly amenable to facilitating ADE activity [35]. These data also indicate that even modest concentrations of DENV-

reactive IgA1 can significantly antagonize polyclonal IgG1-mediated enhancement of DENV infection, signifying that the presence of DENV E reactive IgA1 (especially fusion loop reactive IgA1) has the potential to significantly modulate DENV infection and associated immunopathogenesis.

- 5 [00147] **FIGS. 4A-4C depicts Monoclonal IgA1 antagonizes ADE mediated by polyclonal DENV-immune plasma.** FIG. 4A depicts DENV immune plasma enhances DENV-3 infection of K562 cells. Each datapoint represents a unique plasma donor (n = 4). FIG. 4B depicts VDB33-IgA antagonizes *in vitro* enhancement of DENV-3 infection mediated by polyclonal DENV-immune serum. Serum used at a 1:50
- 10 dilution for ADE assay, n = 4 unique plasma donors. The percentage of DENV-positive cells was normalized to that observed in the plasma-only condition. FIG. 4C depicts VDB50-IgA antagonizes *in vitro* enhancement of DENV-3 infection mediated by polyclonal DENV-immune serum. Serum used at a 1:50 dilution for ADE assay, n = 4 unique plasma donors. The percentage of DENV-positive cells was normalized to that
- 15 observed in the plasma-only condition. *** p < 0.001, **** p < 0.0001 1-way ANOVA with Dunnett correction for multiple comparisons

Discussion

- [00148] In this study it was demonstrate that DENV-reactive IgA1 monoclonal antibodies can bind and neutralize DENV but are incapable of facilitating ADE.
- 20 Furthermore, the addition of DENV-reactive IgA1 can significant blunt the DENV-infection enhancing activity of monoclonal and polyclonal DENV-reactive antibodies in a complete fashion. These results suggest an unappreciated role for DENV-reactive IgA during the humoral response to DENV infection and raise the potential that IgA could act as regulator of dengue severity and infection-attendant inflammation.
- 25 [00149] While the invention has been shown and described with reference to certain embodiments of the present invention thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present disclosure.
- [00150] The entire disclosure of all applications, patents, and publications cited
- 30 herein are herein incorporated by reference in their entirety. While the foregoing is directed to embodiments of the present disclosure, other and further embodiments of the disclosure may be devised without departing from the basic scope thereof.

SEQUENCE LISTING

<110> The Research Foundation for the State University of New York

<120> IgA MONOCLONAL ANTIBODIES FOR TREATING FLAVIVIRUS INFECTION

<130> 110-2073P01

<160> 11

<170> PatentIn version 3.5

<210> 1

<211> 475

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 1

Gln Leu Gln Leu Gln Ala Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ile Ser Ser
20 25 30

Ser Tyr Phe Trp Gly Trp Ile Arg Gln Pro Pro Glu Lys Glu Leu Gln
35 40 45

Trp Leu Gly Ser Ile Phe Ser Arg Gly Asn Ala Tyr Tyr Asn Pro Ser
50 55 60

Leu Lys Ser Arg Val Thr Val Ser Val Asp Thr Ser Lys Asn Gln Phe
65 70 75 80

Ser Leu Lys Leu Thr Ser Val Thr Ala Thr Asp Thr Ala Val Tyr Tyr
85 90 95

Cys Ala Arg Leu Leu Gln Tyr Lys Trp Asn Trp Leu Phe Asp Pro Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Pro Thr Ser Pro
 115 120 125

Lys Val Phe Pro Leu Ser Leu Cys Ser Thr Gln Pro Asp Gly Asn Val
 130 135 140

Val Ile Ala Cys Leu Val Gln Gly Phe Phe Pro Gln Glu Pro Leu Ser
 145 150 155 160

Val Thr Trp Ser Glu Ser Gly Gln Gly Val Thr Ala Arg Asn Phe Pro
 165 170 175

Pro Ser Gln Asp Ala Ser Gly Asp Leu Tyr Thr Thr Ser Ser Gln Leu
 180 185 190

Thr Leu Pro Ala Thr Gln Cys Leu Ala Gly Lys Ser Val Thr Cys His
 195 200 205

Val Lys His Tyr Thr Asn Pro Ser Gln Asp Val Thr Val Pro Cys Pro
 210 215 220

Val Pro Ser Thr Pro Pro Thr Pro Ser Pro Ser Thr Pro Pro Thr Pro
 225 230 235 240

Ser Pro Ser Cys Cys His Pro Arg Leu Ser Leu His Arg Pro Ala Leu
 245 250 255

Glu Asp Leu Leu Leu Gly Ser Glu Ala Asn Leu Thr Cys Thr Leu Thr
 260 265 270

Gly Leu Arg Asp Ala Ser Gly Val Thr Phe Thr Trp Thr Pro Ser Ser
 275 280 285

Gly Lys Ser Ala Val Gln Gly Pro Pro Glu Arg Asp Leu Cys Gly Cys
 290 295 300

Tyr Ser Val Ser Ser Val Leu Pro Gly Cys Ala Glu Pro Trp Asn His

<210>	2
<211>	214
<212>	PRT
<213>	Artificial Sequence
<220>	
<223>	Synthetic sequence

<400> 2

Ser Tyr Val Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Lys
 1 5 10 15

Thr Ala Arg Ile Thr Cys Gly Gly Asn Asn Ile Glu Ser Lys Ser Val
 20 25 30

His Trp Tyr Gln Gln Lys Ser Arg Gln Ala Pro Val Leu Val Phe Tyr
 35 40 45

Asp His Ser Asp Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Ala Ser
 50 55 60

Asn Ser Gly His Thr Ala Thr Leu Ile Ile Ser Gly Val Glu Ala Gly
 65 70 75 80

Asp Glu Ala Asp Tyr His Cys Gln Val Trp Asp Ser Asp Ser Asp His
 85 90 95

Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro Lys
 100 105 110

Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln
 115 120 125

Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly
 130 135 140

Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys Ala Gly
 145 150 155 160

Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala
 165 170 175

Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser
 180 185 190

Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr Val
 195 200 205

Ala Pro Thr Glu Cys Ser
 210

<210> 3
 <211> 1485
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 3
 atgggctggg cctgcatcat tctgtttctg gtggccacag ccaccggcgt gcactctcaa 60
 ctgcaactgc aggcttctgg ccctggcctg gtcaagcctt ctgagacact gagcctgacc 120
 tgtaccgtgt ctggcggcag cattatcagc agcagctact tctggggctg gatcagacag 180
 cctcctgaga aagaactgca gtggctgggc agcatcttct ccagaggcaa cgcctactac 240
 aaccccagcc tgaagtccag agtgaccgtg tccgtggaca ccagcaagaa ccagttctcc 300
 ctgaagctga ccagcgtgac cgccacagat accgccgtgt actactgtgc cagactgctg 360
 cagtacaagt ggaactggct gttcgaccct tggggccagg gaacactggg cacagtgtct 420
 agcgcctctc caacaagccc caaggtgttc cctctgagcc tgtgtagcac acagcccagc 480
 ggcaatgtcg tgatcgcttg tctggtgcag ggattcttcc cacaagagcc cctgtccgtg 540
 acttggagcg aatctggaca gggcgtgaca gccagaaact tcccacctag ccaggatgcc 600
 agcggcgatc tgtacacaac aagcagccag ctgaccctgc ctgccacaca atgtctggcc 660
 ggcaagtctg tgacctgcca cgtgaagcac tacaccaatc caagccagga cgtgaccgtg 720
 ccttgtcctg tgcctagcac acctcctaca ctttctccaa gcacaccacc aactccatct 780
 ccatcctgct gtcaccccag gctgtctctg catagaccgg ctctggaaga tctgctgctg 840
 ggctctgagg ccaacctgac atgtacactg accggcctga gagatgcctc cggcgtgacc 900
 ttacatgga cacctagctc tggcaagagc gccgttcagg gacctcctga aagggatctg 960

tgccgctgtt acagcgtgtc ctctgtgctg cctggatgtg ccgagccttg gaatcacggc	1020
aagaccttta cctgcaccgc cgcctatcct gagagcaaga cacctctgac agccacactg	1080
agcaagagcg gcaacacctt cagacccgaa gtgcatctgc tgcctccacc atctgaagaa	1140
ctggccctga acgagctggg cactctgaca tgtctggcta gaggcttcag ccctaaggac	1200
gtgctcgtca gatggctgca gggctctcaa gagctgccta gagagaagta cctgacctgg	1260
gccagcagac aagagccttc tcagggcacc accacctttg ccgtgaccag cattctgaga	1320
gtggccgccc aggattggaa gaagggcgat accttcagct gcatggctcg acacgaagcc	1380
ctgcctctgg cttcacaca gaaaaccatc gatcggctgg ccggaaagcc cacacatgtg	1440
aatgtgtccg tcgtgatggc cgaggtggac ggcacatgtt attga	1485

<210> 4
 <211> 702
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 4	
atgggatggg catgtattat tctgtttctg gtcgcaactg caaccggcgt gcatagcagc	60
tacgtgctga cacagcctcc atccgtgtct gtggcccctg gaaagaccgc cagaatcaca	120
tgccggcggca acaacatcga gagcaagagc gtgcactggg atcagcagaa gtccagacag	180
gcccctgtgc tgggtgttcta cgaccacagc gatagacca gcggcatccc cgagagattc	240
agcgcctcta atagcggcca caccgccaca ctgatcatct ctggtgttga ggccggcgac	300
gaggccgatt accattgcca agtgtgggac agcgacagcg atcacctgt ttttggcgga	360
ggcaccaagc tgacagtgtc ggggcagccc aaggccgctc ctagctgac actgtttccc	420
ccttcctccg aggagctgca ggccaacaag gccaccctgg tgtgcctgat ctccgacttc	480
tatcctggcg ccgtgacagt ggcctggaag gctgattcta gccagtgaa ggctggcgtg	540
gagaccacaa cccctccaa gcagtctaac aataagtatg ccgcttcctc ttacctgagc	600
ctgacaccag agcagtggaa gtcccaccgg tcttacagct gccaggtcac tcacgaaggc	660

tctaccgtgg aaaagacagt cgcacccacc gaatgctcat ga

702

<210> 5
 <211> 495
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> synthetic sequence

<400> 5

Met Arg Cys Val Gly Ile Gly Asn Arg Asp Phe Val Glu Gly Leu Ser
 1 5 10 15

Gly Ala Thr Trp Val Asp Val Val Leu Glu His Gly Ser Cys Val Thr
 20 25 30

Thr Met Ala Lys Asp Lys Pro Thr Leu Asp Ile Glu Leu Leu Lys Thr
 35 40 45

Glu Val Thr Asn Pro Ala Val Leu Arg Lys Leu Cys Ile Glu Ala Lys
 50 55 60

Ile Ser Asn Thr Thr Thr Asp Ser Arg Cys Pro Thr Gln Gly Glu Ala
 65 70 75 80

Thr Leu Val Glu Glu Gln Asp Thr Asn Phe Val Cys Arg Arg Thr Phe
 85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly Ser
 100 105 110

Leu Ile Thr Cys Ala Lys Phe Lys Cys Val Thr Lys Leu Glu Gly Lys
 115 120 125

Ile Val Gln Tyr Glu Asn Leu Lys Tyr Ser Val Ile Val Thr Val His
 130 135 140

Thr Gly Asp Gln His Gln Val Gly Asn Glu Thr Thr Glu His Gly Thr

39

Thr Ala Asn Pro Ile Val Thr Asp Lys Glu Lys Pro Val Asn Ile Glu
 355 360 365

Ala Glu Pro Pro Phe Gly Glu Ser Tyr Ile Val Val Gly Ala Gly Glu
 370 375 380

Lys Ala Leu Lys Leu Ser Trp Phe Lys Lys Gly Ser Ser Ile Gly Lys
 385 390 395 400

Met Phe Glu Ala Thr Ala Arg Gly Ala Arg Arg Met Ala Ile Leu Gly
 405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Ile Gly Gly Val Phe Thr Ser Val
 420 425 430

Gly Lys Leu Ile His Gln Ile Phe Gly Thr Ala Tyr Gly Val Leu Phe
 435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Gly Ile Gly Ile Leu Leu Thr
 450 455 460

Trp Leu Gly Leu Asn Ser Arg Ser Thr Ser Leu Ser Met Thr Cys Ile
 465 470 475 480

Ala Val Gly Met Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
 485 490 495

<210> 6
 <211> 1417
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 6
 atgggctggg catgcattat tctgtttctg gtcgcaactg ctacaggcgt gcatagtgaa 60
 gtgcagctgc tggaatctgg cggaggactg gttcaacctg gcggctctct gagactgtct 120

tgtgccgcca gcggcttcac cttcagcagc tttgtgatgg cctgggtccg acaggccct	180
ggcaaaggac ttgaatgggt gtccgtgatc tacgacggcg gcagcagcac ctactacgcc	240
gattctgtga agggcagatt caccatcagc cgggacaaca gcaagaacac cctgtacctg	300
cagatgaaca gcctgagagc cgaggacacc gccgtgtact attgtgcca ggccagccag	360
atggccaccg tgttcacga ttattggggc cagggcaccc tggtcaccgt ttcttctgcc	420
agcaccaagg gcccttcctg gtttccactg gccccctcct ctaaattcac atctggcggc	480
accgccgccc tgggctgtct ggtgaaggac tacttcccag agcctgtgac agtgtcctgg	540
aactctggcg ccctgacatc cggcgtgcac acatttccag ccgtgctgca gagctccggc	600
ctgtacagcc tgtctagcgt ggtgacagt ccctcctcta gcctgggcac acagacctat	660
atctgcaacg tgaatcacaa gccaagcaat accaagggtg acaagaagg ggagcccaag	720
tcctgtgata agacacacac ctgccccct tgtcctgtc ccgagctgt gggcggccct	780
agcgtgttcc tgtttccacc caagcctaag gacaccctga tgatctccc gacaccgag	840
gtgacctgcg tgggtgtgga cgtgtctcac gaggatcctg aggtgaagtt caactggtat	900
gtggatggcg tggaggtgca caatgccaa accaagccca gagaggagca gtacaactct	960
acatataggg tggtagcgt gctgaccgt ctgcaccagg actggctgaa cggcaaggag	1020
tataagtgca aggtgtccaa taaggccctg cccgccccca tcgagaagac aatcagcaag	1080
gccaagggcc agcctcggga gccacaggtg tacaccctgc ctccatccag agacgagctg	1140
acaaagaacc aggtgtctct gacatgtctg gtgaagggt tctatcctag cgatatcgcc	1200
gtggagtggg agtccaatgg ccagccagag aacaattaca agaccacacc ccctgtgctg	1260
gactccgatg gctccttctt tctgtattcc aagctgaccg tggataagtc tcggtggcag	1320
cagggaacg tgttcagctg ttccgtgatg cacgaagccc tgcataatca ctatactcag	1380
aaatccctgt ccctgtcacc tggaaagtga taagctt	1417

<210> 7

<211> 711

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic sequence

<400> 7

atgggatggt catgtattat tctgtttctg gtcgcaactg caaccggcgt gcatagccag	60
tctgtgctga cacagcctcc atctgtgtct ggcgctccag gccagagagt gatcatcagc	120
tgtacaggca gcagcagcaa catcggagcc ggctttgacg tgcactggta tcagcagctg	180
cctggcacag cccctaaact gctgatctac ggcaacaaca acagaccag cgccgtgcct	240
gatagattca gcggctctaa gagcggcaca tctgccagcc tggccattac tggactgcag	300
gccgaagatg aggccgacta ctactgccag agctacgaca gctctctgtc tggcggagtt	360
tttggcggag gcaccaagct gacagtgtg gggcagccca aggccgctcc tagcgtgaca	420
ctgtttcccc cttcctccga ggagctgcag gccacaagg ccaccctgggt gtgcctgac	480
tccgacttct atcctggcgc cgtgacagtg gcctggaagg ctgattctag ccagtgaa	540
gctggcgtgg agaccacaac cccctccaag cagtctaaca ataagtatgc cgcttcctct	600
tacctgagcc tgacaccaga gcagtggaag tcccaccggt cttacagctg ccaggctact	660
cacgaaggct ctaccgtgga aaagacagtc gcaccaccg aatgctcatg a	711

<210> 8

<211> 1422

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic sequence

<400> 8

gaagttcagc tgcttgagtc tggcggcgga ctggttcaac ctggcggatc tctgagactg	60
agctgtgccg ccagcggctt caccttcagc agctttgtga tggcctgggt ccgacaggcc	120
cctggcaaag gacttgaatg ggtgtccgtg atctacgacg gcggcagcag cacctactac	180
gccgattctg tgaagggcag attcaccatc agccgggaca acagcaagaa caccctgtac	240
ctgcagatga acagcctgag agccgaggac accgccgtgt actattgtgc caaggccagc	300
cagatggcca ccgtgttcat cgattattgg ggccagggca ccctggtcac cgtgtcatct	360

gctagcccta caagcccaaa ggtgttccct ctgagcctgt gtagcacaca gcccgcggc 420
 aatgtcgtga tcgcttgtct ggtgcaggga ttcttccac aagagcccct gtccgtgact 480
 tggagcgaat ctggacaggg cgtgaccgcc agaaacttcc caccttctca ggatgccagc 540
 ggcgacctgt acacaacaag cagccaactg accctgcctg ccacacagtg tctggccgga 600
 aagtctgtga cctgccacgt gaagcactac aaaaaccca gccaggacgt gaccgtgcct 660
 tgtcctgttc ctagcacacc tcctacacct tctccaagca caccaccaac tccatctcca 720
 tcctgtctgc accccagact gagcctgcat agaccgctc tggaagatct gctgtgggc 780
 tctgaggcca acctgacatg tacactgacc ggcctgagag atgcctccgg cgtgacctt 840
 acatggacac ctagcagcgg caagagcgc gttcaaggac ctctgagag ggatctgtgc 900
 ggctgttaca gcgtgtctc tgtgctgcct ggatgtgccg agccttggaa tcacggcaag 960
 accttcacct gtaccgccgc ctatctgag agcaagacc ctctgacagc cacttgagc 1020
 aagagcggca acaccttctg gcccgaagt catcttctgc ctccacctag cgaagaactg 1080
 gccctgaatg agctggtcac cctgacatgc ctggccagag gcttcagccc taaggatgtg 1140
 ctctgcagat ggctgcaggg cagccaagag ctgccagag agaagtatct gacctgggcc 1200
 agcagacaag agcctagcca gggaaccacc acctttgccg tgaccagcat tctgagagt 1260
 gccgccgagg attggaagaa gggcgatacc ttcagctgca tggctggaca cgaagccctg 1320
 ccaactggcct tcacacagaa aaccatcgac agactggccg gcaagcccac acatgtgaat 1380
 gtgtctgtgg tcatggccga ggtggacggc acatgttatt ga 1422

<210> 9
 <211> 654
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 9
 cagtctgtgc tgacacagcc tccatctgtg tctggcgctc caggccagag agtgatcatc 60
 agctgtacag gcagcagcag caacatcgga gccggctttg acgtgcactg gtatcagcag 120

ctgcctggca cagcccctaa actgctgac tacggcaaca acaacagacc cagcgccgtg 180
 cctgatagat tcagcggctc taagagcggc acatctgcca gcctggccat tactggactg 240
 caggccgaag atgaggccga ctactactgc cagagctacg acagctctct gtctggcgga 300
 gtttttggcg gaggcaccaa gctgacagtg ctggggcagc ccaaggccgc tcctagcgtg 360
 aactgtttc ccccttctc cgaggagctg caggccaaca aggccaccct ggtgtgcctg 420
 atctccgact tctatcctgg cgccgtgaca gtggcctgga aggctgattc tagcccagtg 480
 aaggctggcg tggagaccac aacccccctc aagcagtcta acaataagta tgccgcttcc 540
 tcttacctga gcctgacacc agagcagtgg aagtcaccac ggtcttacag ctgccaggtc 600
 actcacgaag gctctaccgt ggaaaagaca gtcgcacca ccgaatgctc atga 654

<210> 10
 <211> 473
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 10

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe
 20 25 30

Val Met Ala Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Val Ile Tyr Asp Gly Gly Ser Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

45

Ser Ala Val Gln Gly Pro Pro Glu Arg Asp Leu Cys Gly Cys Tyr Ser
 290 295 300

Val Ser Ser Val Leu Pro Gly Cys Ala Glu Pro Trp Asn His Gly Lys
 305 310 315 320

Thr Phe Thr Cys Thr Ala Ala Tyr Pro Glu Ser Lys Thr Pro Leu Thr
 325 330 335

Ala Thr Leu Ser Lys Ser Gly Asn Thr Phe Arg Pro Glu Val His Leu
 340 345 350

Leu Pro Pro Pro Ser Glu Glu Leu Ala Leu Asn Glu Leu Val Thr Leu
 355 360 365

Thr Cys Leu Ala Arg Gly Phe Ser Pro Lys Asp Val Leu Val Arg Trp
 370 375 380

Leu Gln Gly Ser Gln Glu Leu Pro Arg Glu Lys Tyr Leu Thr Trp Ala
 385 390 395 400

Ser Arg Gln Glu Pro Ser Gln Gly Thr Thr Thr Phe Ala Val Thr Ser
 405 410 415

Ile Leu Arg Val Ala Ala Glu Asp Trp Lys Lys Gly Asp Thr Phe Ser
 420 425 430

Cys Met Val Gly His Glu Ala Leu Pro Leu Ala Phe Thr Gln Lys Thr
 435 440 445

Ile Asp Arg Leu Ala Gly Lys Pro Thr His Val Asn Val Ser Val Val
 450 455 460

Met Ala Glu Val Asp Gly Thr Cys Tyr
 465 470

<210> 11

<211> 217
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic sequence

<400> 11

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

Arg Val Ile Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
 20 25 30

Phe Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
 35 40 45

Leu Ile Tyr Gly Asn Asn Asn Arg Pro Ser Ala Val Pro Asp Arg Phe
 50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu
 65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser
 85 90 95

Leu Ser Gly Gly Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 115 120 125

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 130 135 140

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
 145 150 155 160

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys

165 170 175

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
180 185 190

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
195 200 205

Lys Thr Val Ala Pro Thr Glu Cys Ser
210 215

WHAT IS CLAIMED IS:

1. An isolated monoclonal antibody, comprising: a heavy chain having an amino acid sequence of SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain comprises an Fc region characterized as IgA Fc domain.
5
2. The isolated monoclonal antibody of claim 1, wherein the IgA Fc domain is an IgA1 Fc domain.
3. The isolated monoclonal antibody of claim 1, wherein the IgA Fc domain is an IgA2 Fc domain.
- 10 4. The isolated monoclonal antibody of claim 1, wherein the antibody is chimeric or humanized.
5. The isolated monoclonal antibody of claim 1, further comprising: a light chain having an amino acid sequence of SEQ ID. NO. 2.
- 15 6. The isolated monoclonal antibody of claim 1, wherein the isolated monoclonal antibody binds an epitope of a Deng virus, wherein the epitope is characterized as a loop.
7. An isolated monoclonal antibody for targeting a fusion loop epitope of dengue virus, comprising:

a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain comprises an Fc region characterized as IgA Fc domain; and
20

a light chain having an amino acid sequence having at least 90% sequence identity to SEQ ID. NO. 2.
8. The isolated monoclonal antibody of claim 7, wherein the heavy chain comprises an amino acid sequence having at least 95%, 97%, 98%, 99% sequence identity to SEQ. ID NO. 1.
25

9. The isolated monoclonal antibody of claim 7, wherein the light chain comprises an amino acid sequence having at least 95%, 97%, 98%, 99% sequence identity to SEQ. ID NO. 2.
10. An isolated monoclonal antibody, comprising: a heavy chain consisting of an amino acid sequence of SEQ. ID NO. 1; and
5 a light chain consisting of an amino acid sequence of SEQ ID NO: 2.
11. The isolated monoclonal antibody of claim 10, wherein the isolated monoclonal antibody binds a fusion loop epitope (FLE) of a Dengue virus.
12. An isolated monoclonal antibody, comprising: a heavy chain having an amino acid sequence having at least 90% sequence identity to SEQ. ID NO. 1, wherein the heavy chain or a segment of the heavy chain comprises an Fc region characterized as IgA Fc domain.
10
13. The isolated monoclonal antibody of claim 11, wherein the isolated monoclonal antibody binds an epitope of a Dengue virus, wherein the epitope is characterized as a loop.
15
14. A nucleic acid polymer encoding a monoclonal antibody, wherein the polymer comprises SEQ. ID NO. 1.
15. A nucleic acid polymer encoding a monoclonal antibody, wherein the polymer comprises SEQ. ID NO. 2.
- 20 16. A complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 1.
17. A complementary deoxynucleotide (cDNA) sequence comprising a nucleic acid sequence of SEQ ID NO: 3.
- 25 18. A complementary deoxynucleotide (cDNA) sequence encoding an amino acid sequence having at least at least 90%, at least 95%, or at least 99% sequence identity to SEQ ID NO: 2.

19. A complementary deoxynucleotide (cDNA) sequence comprising a nucleic acid sequence of SEQ ID NO: 4.

20. A method for preventing or treating a dengue viral infection, the method comprising: administering a therapeutically effective amount of the monoclonal antibody of any of claims 1-13 to a subject in need thereof under conditions effective to treat the viral infection.

21. The method according to claim 20, wherein the method is used for preventing or treating antibody-dependent enhancement of a viral infection.

22. A method for preventing or treating antibody-dependent enhancement of a viral infection, the method comprising:

administering a therapeutically effective amount of a monoclonal antibody comprising a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is characterized as an isotypic commutation.

23. The method according to claim 22, wherein the viral infection is any one of respiratory syncytial virus (RSV), influenza, and coronavirus infection, or a combination thereof.

24. The method according to claim 20, wherein the viral infection is a flavivirus infection.

25. The method according to claim 24, wherein the flavivirus infection is a Zika virus.

26. The method according to claim 25, wherein the flavivirus infection is a dengue virus.

27. A method for preventing or treating antibody-dependent enhancement of a viral infection, the method comprising:

administering a therapeutically effective amount of a monoclonal antibody

comprising a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to treat the viral infection, wherein the IgA Fc domain is formed by class-switch recombination.

- 5 28. A method for preventing or treating antibody-dependent enhancement of a flavivirus infection, the method comprising:

administering a therapeutically effective amount of a monoclonal antibody of:

a) a monoclonal antibody encoded by a DNA polymer of claims 14 and 15;

b) a cDNA of claims 16 and 17; or

- 10 c) a monoclonal antibody of claims 1-13.

29. The method of claim 28, where in the monoclonal antibody is disposed with a serum comprising one or more additional antibodies or fragments thereof directed against dengue virus or bind one or more epitopes of dengue virus.

- 15 30. A method for preventing or treating antibody-dependent enhancement of a viral infection, the method comprising:

administering a therapeutically effective amount of an antibody comprising a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain to a subject in need thereof under conditions effective to prevent or treat antibody-dependent enhancement of a viral infection.

- 20 31. The method according to claim 30, wherein the viral infection is any one of respiratory syncytial virus (RSV), influenza, and coronavirus infection, or a combination thereof.

32. The method according to claim 30, wherein the viral infection is a flavivirus infection.

- 25 33. The method according to claim 32, wherein the flavivirus infection is a Zika virus.

34. The method according to claim 33, wherein the flavivirus infection is a dengue virus.

35. The method of claim 30, further comprising, increasing a concentration of antibodies characterized as IgA in a serum comprising one or more additional
5 antibodies.

36. The method of claim 30, wherein the antibody is a wild-type IgA isoform that binds dengue protein E.

37. The method of claim 30, wherein the antibody is monoclonal.

38. The method of claim 30, wherein the antibody is synthetic, formed by
10 recombinant technology, or characterized as isolated.

39. A method for preventing or treating antibody-dependent enhancement of a viral infection, the method comprising:

administering a therapeutically effective amount of a monoclonal antibody to a subject in need thereof under conditions effective to prevent or treat antibody-
15 dependent enhancement of a viral infection, wherein the monoclonal antibody is characterized as an IgA isoform.

40. The method of claim 39, wherein the monoclonal antibody comprises a heavy chain or a segment of the heavy chain comprising an Fc region characterized as IgA Fc domain.

20 41. The method of claim 40, wherein the Fc region characterized as IgA Fc domain comprises an amino acid sequence of SEQ ID NO: 1.

42. The method of claim 40, wherein the Fc region characterized as IgA Fc domain comprises an amino acid segment between S31 and Y475 of SEQ ID NO: 1, an amino acid segment between G51 and Y475 of SEQ ID NO: 1, an amino acid segment
25 between L101, or an amino acid segment between A181 and Y475, wherein SEQ ID NO: 1 is used for numbering.