


or a pharmaceutically acceptable salt, enantiomer, stereoisomer, or tautomer thereof, wherein

A is taken from the group consisting of $-N(R_2)R_3$ and G;

G is selected from the group consisting of



and wherein the symbol (**) is the point of attachment to the pyrimidine ring;

each G moiety may be further substituted with one, two, or three R_4 moieties;

W is C5-C6heteroaryl, phenyl, $-NHC(O)R_6$, $-NHC(O)R_7$, $-NHC(O)N(R_8)R_9$ or $-C(O)N(R_8)R_9$, and wherein each C5-C6heteroaryl or phenyl is optionally substituted by one, two, or three R_5 moieties;

X_1 and X_2 are individually and independently hydrogen or C1-C6 alkyl;

R_1 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, or branched C3-C8 alkyl;

R_2 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, branched C3-C8 alkyl, C3-C8 cycloalkyl, fluoroC1-C6alkyl wherein the alkyl is fully or partially fluorinated, $-(CH_2)_m-OR_8$, or a 3-8 membered non-aromatic heterocycle, wherein each alkylene is optionally substituted with C1-C4 alkyl;

R_3 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, branched C3-C8 alkyl, C3-C8 cycloalkyl, fluoroC1-C6alkyl wherein the alkyl is fully or partially fluorinated or a 3-8 membered non-aromatic heterocycle;


each R_4 is individually and independently hydrogen, halogen, C1-C6 alkyl, fluoro-C1-C6 alkyl wherein the alkyl chain is partially or completely fluorinated, branched C3-C8 alkyl, C3-C8 cycloalkyl, $-(CH_2)_m-OR_8$, $-(CH_2)_m-NR_8(R_9)$, $-(CH_2)_m-R_7$, or cyano, wherein each alkylene is optionally substituted with C1-C4 alkyl;

each R_5 is individually and independently hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, branched C3-C8 alkyl, halogen, cyano, fluoro-C1-C6 alkyl wherein the alkyl chain is partially or completely fluorinated, $-(CH_2)_m-C(O)NR_8(R_9)$, $-(CH_2)_m-C(O)R_7$, $-(CH_2)_m-OR_8$, $-(CH_2)_m-NR_8(R_9)$, or $-(CH_2)_m-R_7$, wherein each alkylene is optionally substituted with C1-C4 alkyl;

CLAIMS

What is claimed is:

1. A compound of **Formula I**,




Formula I

or a pharmaceutically acceptable salt, enantiomer, stereoisomer, or tautomer thereof, wherein

A is taken from the group consisting of $-N(R_2)R_3$ and G;

G is selected from the group consisting of



and wherein the symbol (**) is the point of attachment to the pyrimidine ring;

each G moiety may be further substituted with one, two, or three R_4 moieties;

W is C5-C6heteroaryl, phenyl, $-NHC(O)R_6$, $-NHC(O)R_7$, $-NHC(O)N(R_8)R_9$ or $-C(O)N(R_8)R_9$, and wherein each C5-C6heteroaryl or phenyl is optionally substituted by one, two, or three R_5 moieties;

X_1 and X_2 are individually and independently hydrogen or C1-C6 alkyl;

R_1 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, or branched C3-C8 alkyl;

R_2 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, branched C3-C8 alkyl, C3-C8 cycloalkyl, fluoroC1-C6alkyl wherein the alkyl is fully or partially fluorinated, $-(CH_2)_m-OR_8$, or a 3-8 membered non-aromatic heterocycle, wherein each alkylene is optionally substituted with C1-C4 alkyl;

R_3 is hydrogen, C1-C6 alkyl, deuterio-C1-C6 alkyl wherein the alkyl chain is partially or completely deuterated, branched C3-C8 alkyl, C3-C8 cycloalkyl, fluoroC1-C6alkyl wherein the alkyl is fully or partially fluorinated or a 3-8 membered non-aromatic heterocycle;