CLAIMS

We claim:

1. A method for preparing remnant tumor infiltrating lymphocytes (rTILs) for adoptive T cell therapy, the method comprising:

   (a) obtaining tumor tissue from a patient, wherein the tumor tissue comprises tumor infiltrating lymphocytes (TILs);

   (b) fragmenting the tumor tissue;

   (c) treating the tumor tissue in a gas permeable container with a first cell culture medium and interleukin 2 (IL-2) to provide tumor remnants and emergent TILs (eTILs);

   (d) removing at least a plurality of the eTILs;

   (e) enzymatically digesting the tumor remnants into tumor remnant cells using a digest mixture; and

   (f) expanding the tumor remnant cells with a second cell culture medium comprising cell culture media, irradiated feeder cells, OKT-3 antibody, and IL-2 in a gas permeable container to provide an expanded number of remnant tumor infiltrating lymphocytes (rTILs);

   wherein the rTILs express reduced levels of a T cell exhaustion marker relative to the eTILs, and wherein the T cell exhaustion marker is selected from the group consisting of TIM3, LAG3, TIGIT, PD-1, CTLA-4, and combinations thereof.

2. The method of Claim 1, wherein the tumor tissue is selected from the group consisting of melanoma tumor tissue, head and neck tumor tissue, breast tumor tissue, renal tumor tissue, pancreatic tumor tissue, glioblastoma tumor tissue, lung tumor tissue, colorectal tumor tissue, sarcoma tumor tissue, triple negative breast tumor tissue, cervical tumor tissue, ovarian tumor tissue, and acute myeloid leukemia bone marrow or tumor tissue.

3. The method of Claim 1 or 2, wherein the irradiated feeder cells comprise irradiated allogeneic peripheral blood mononuclear cells.

4. The method of any one of Claims 1 to 3, wherein IL-2 is present in the second cell culture medium at an initial concentration of about 3000 IU/mL and OKT-3 antibody is present in the second cell culture medium at an initial concentration of about 30 ng/mL.

5. The method of any one of Claims 1 to 4, wherein at least one T cell exhaustion marker in $CD8^+$ and $CD4^+$ T cells in the rTILs is reduced by at least 10% relative to the eTILs.

6. The method of any one of Claims 1 to 5, wherein the T cell exhaustion marker is a LAG3 marker in $CD8^+$ T cells, and wherein the LAG3 marker in the rTILs is reduced by at least 2-fold relative to the eTILs.

7. The method of any one of Claims 1 to 6, wherein the T cell exhaustion marker is a TIM3 marker in $CD8^+$ T cells, and wherein the LAG3 marker in the rTILs is reduced by at least 3-fold relative to the eTILs.

8. The method of any one of Claims 1 to 7, wherein the T cell exhaustion marker is a TIM3 marker in $CD4^+$ T cells, and wherein the LAG3 marker in the rTILs is reduced by at least 2-fold relative to the eTILs.

9. The method of any one of Claims 1 to 8, wherein the TIM3 marker and the LAG3 marker in the rTILs are undetectable by flow cytometry.

10. The method of any one of Claims 1 to 9, wherein $CD56^+$ expression in the rTILs is reduced by at least 3-fold relative to $CD56^+$ expression in the eTILs.

11. The method of any one of Claims 1 to 10, wherein $CD69^+$ expression in the rTILs is increased by at least 2-fold relative to $CD69^+$ expression in the eTILs.

12. The method of any one of Claims 1 to 11, wherein the digest mixture comprises deoxyribonuclease, collagenase, and hyaluronidase.

13. The method of any one of Claims 1 to 12, wherein the first cell culture medium further comprises a cytokine selected from the group consisting of IL-4, IL-7, IL-15, IL-21, and combinations thereof.

14. The method of any one of Claims 1 to 13, wherein the second cell culture medium further comprises a cytokine selected from the group consisting of IL-4, IL-7, IL-15, IL-21, and combinations thereof.

15. The method of any one of Claims 1 to 14, further comprising the steps of cryopreserving the expanded number of rTILs to form a cryopreserved rTIL population.

16. The method of Claim 15, further comprising the step of thawing the cryopreserved rTIL population.

17. The method of any one of Claims 1 to 16, wherein the rTILs are substantially free of eTILs.

18. The method of any one of Claims 1 to 17, wherein the step of expanding the tumor remnant cells with the second cell culture medium comprises adding eTILs to the second cell culture medium to provide a selected rTIL to eTIL ratio.

19. The method of any one of Claims 1 to 17, further comprising the step of adding eTIL to the rTIL to provide a selected rTIL to eTIL ratio.

20. The method of Claim 18 or 19, wherein the selected rTIL to eTIL ratio is at least 1% rTIL to eTIL.

21. The method of Claim 20, wherein the selected rTIL to eTIL ratio is at most 99.9% rTIL to eTIL.

22. The method of Claim 18 or 19, wherein the selected rTIL to eTIL ratio is selected from the group consisting of about 1:99, about 5:95, about 10:90, about 15:85, about 20:80, about 25:75, about 30:70, about 35:65, about 40:60, about 45:55, about 50:50, about 55:45, about 60:40, about 65:35, about 70:30, about 75:25, about 80:20, about 85:15, about 90:10, about 95:5, and about 99:1 rTIL to eTIL.

23. A method of treating a cancer in a patient in need of such treatment, wherein the treatment comprises delivering a therapeutically effective amount of rTILs to a patient, wherein the rTILs are prepared according to the method of any one of Claims 1 to 22.

24. A method of treating a cancer in a patient in need of such treatment, comprising the steps of:

(a) obtaining rTILs from a tumor resected from a patient according to the method of any one of Claims 1 to 22;

(b) treating the patient with a non-myeloablative lymphodepletion regimen prior to administering the rTILs to the patient;

(c) administering a therapeutically effective amount of rTILs to the patient, wherein the

therapeutically effective amount of rTILs may be optionally cryopreserved and thawed prior to the administering to the patient; and

(d) treating the patient with a high-dose IL-2 regimen starting on the day after administration of the rTILs to the patient.

25. The method of any one of Claims 23 to 24, wherein a therapeutically effective amount of eTILs are simultaneously administered to the patient in a mixture with the rTILs.

26. The method of any one of Claims 23 to 25, wherein the non-myeloablative lymphodepletion regimen comprises the steps of administration of cyclophosphamide at a dose of 60 mg/m^2/day for two days followed by administration of fludarabine at a dose of 25 mg/m^2/day for five days.

27. The method of any one of Claims 23 to 26, wherein the high-dose IL-2 regimen comprises 600,000 or 720,000 IU/kg of aldesleukin, or a biosimilar or variant thereof, administered as a 15-minute bolus intravenous infusion every eight hours until tolerance.

28. The method of any one of Claims 23 to 27, wherein the cancer is selected from the group consisting of melanoma, double-refractory melanoma, uveal melanoma, ovarian cancer, cervical cancer, lung cancer, bladder cancer, breast cancer, head and neck cancer, renal cell carcinoma, acute myeloid leukemia, colorectal cancer, and sarcoma.

29. The method of Claim 28, wherein the cancer is breast cancer, and wherein the breast cancer is selected from the group consisting of triple negative breast cancer, estrogen receptor-positive breast cancer, progesterone receptor-positive breast cancer, and estrogen receptor-positive/progesterone receptor-positive breast cancer.

30. The method of Claim 28, wherein the cancer is lung cancer, and the lung cancer is selected from the group consisting of non-small cell lung cancer and small cell lung cancer.