

A MEDICINAL COMPOSITION OF AMARANTH EXTRACT ORIGIN HAVING ENRICHED NITRATE CONTENT AND A METHOD OF PREPARING THE SAME

FIELD OF THE INVENTION

5 The disclosure relates to a medicinal composition of Amaranth extract origin having enriched nitrate content, L-arginine, flavonoids, carbohydrate, protein, saponins, alkaloids, potassium and having negligible amount of oxalic acid or oxalate content for lowering the blood pressure (hypertension) and increasing the endurance of mammals especially human beings. The said enriched extract of
10 Amaranth is useful for enhancing the level of nitric oxide in the blood plasma and saliva. The enriched extract of Amaranth is also useful for enhancing the level of nitrite and nitrate in the blood plasma and saliva and for lowering the blood pressure (hypertension) and increasing the endurance.

The disclosure also relates to a process for manufacturing the said enriched
15 extract of Amaranth having negligible amount of oxalic acid or oxalate content. More particularly, the disclosure relates to a process of extraction of Amaranth and enrichment of the extract mainly with nitrates to get percentage of nitrates in a range of 3-20%. Further the disclosure also discloses a method of treatment for enhancing the level of nitric oxide in the blood and saliva and also useful for
20 enhancing the level of nitrite and nitrate in the blood plasma and saliva. Disclosure discloses a method of treatment for lowering the blood pressure (hypertension) and increasing the endurance by delivering the nitrate enriched formulations orally as fast melt tablets, lozenge, candy, chewing gum, beverage etc. to make it convenient for absorption from the mouth and also in the form of
25 tablets, capsules, pills, powders etc to be absorbed from the stomach.

BACKGROUND OF THE INVENTION

Nitric oxide is an important cellular signaling molecule involved in many physiological and pathological processes. It is a powerful vasodilator with a short half-life of a few seconds in the blood. Nitric Oxide is a free form of gas

produced by the cells in our body. It is produced when enzymes in the body break down the amino acid Arginine, and is mainly used for intra-cellular communication. Nitric Oxide is required for key physiological functions within our body. It travels freely from one cell to another assisting in a variety of biological functions. Depending on the requirement, Nitric Oxide can act as a hormone, a neurotransmitter and an intracellular messenger.

Nitric Oxide is crucial for a variety of body functions, and it is essential that the cells produce adequate amount of this gas within our bodies. Body builders or those who perform rigorous physical activities need a higher amount of NO, as it allows for an increase in blood flow while building muscles. Signs of NO deficiency include extreme fatigue and physical weakness. It is useful in treating a variety of conditions such as insomnia, obesity, diabetes and sexual problems. In the cardiovascular system, NO is an important determinant of basal vascular tone, prevents platelet activation, limits leukocyte adhesion to the endothelium, and regulates myocardial contractility. NO may also play a role in the pathogenesis of common cardiovascular disorders, including hypotension accompanying shock states, essential hypertension, and atherosclerosis. Nitric oxide regulates blood circulation throughout the body, increases the diameter of blood vessels and prevents formation of clots. It assists the endothelial cells in controlling and relaxing blood vessels. Nitric oxide is a cytotoxic agent of macrophages, a messenger molecule of neurons, and a vasodilator produced by endothelial cells. The Immune cells within our body release Nitric Oxide to destroy bacteria, virus and other harmful foreign organisms that can cause infections. NO is also known to prevent tumour and cancerous growths within the body cells.

Nitric Oxide has been shown to be mediator of erectile function. Nitric Oxide stimulates, invigorates and amplifies the sexual response mechanism within the body. Sensory and mental stimulation causes the nerve cells to release nitric oxide. This causes the penis muscles to relax, allowing blood to flow into the penis and create an erection. The process remains the same for women too, as

blood flow increases in their vaginal tissues. Nitric Oxide increases the endurance level of the muscle cells, enabling you to lift heavier load and perform strenuous activities with ease. Nitric Oxide acts as an intracellular messenger between various cells in the body, including the nerve cells. With adequate amount of NO present in the body, the communication between nerve cells is faster, leading to quick responses and an increase in focus and vigilance. Common disorders that promote atherosclerosis, such as hypertension, hyperlipidemia, smoking, and diabetes, are all associated with abnormal endothelial function, one manifestation of which is a comparative deficiency of bioactive NO. A deficiency of NO producing neurons in the gastrointestinal tract is believed to be responsible for certain abnormalities in gastrointestinal motility, such as Hirschsprung's disease, achalasia, and chronic intestinal pseudo-obstruction. NO is also believed to play an important role in gastric cytoprotection, possibly by way of increased mucosal blood flow and the modulation of gastric epithelial function.

15 *Production of Nitric oxide in human body (Biosynthesis)*

NO is produced from the amino acid L-arginine by the enzymatic action of nitric oxide synthase (NOS). There are two endothelial forms of NOS: constitutive NOS (cNOS; type III) and inducible NOS (iNOS; type II). Co-factors for NOS include oxygen, NADPH, tetrahydrobiopterin and flavin adenine nucleotides. In addition to endothelial NOS, there is a neural NOS (nNOS; type I) that serves as a transmitter in the brain and in different nerves of the peripheral nervous system, such as non-adrenergic, non-cholinergic autonomic nerves that innervate penile erectile tissues and other specialized tissues in the body to produce vasodilation.

Under normal, basal condition in blood vessels, NO is continually being produced by cNOS. The activity of cNOS is calcium and calmodulin-dependent. There are two basic pathways for the stimulation of cNOS, both of which involve release of calcium ions from subsarcolemmal storage sites. First, shearing forces acting on the vascular endothelium generated by blood flow causes a release of calcium and subsequent cNOS activation. Therefore, increases in blood flow stimulate NO formation (flow-dependent NO formation). Second, endothelial

receptors for a variety of ligands stimulate calcium release and subsequent NO production (receptor-stimulated NO formation). Included are receptors for acetylcholine, bradykinin, substance-P, adenosine, and many others vasoactive substances.

- 5 The other isoform of endothelial NOS is iNOS. It differs, in part, from cNOS in that its activation is calcium independent. Under normal, basal conditions, the activity of iNOS is very low. The activity of iNOS is stimulated during inflammation by bacterial endotoxins (e.g., lipopolysaccharide) and cytokines such as tumor necrosis factor (TNF) and interleukins. During inflammation, the
10 amount of NO produced by iNOS may be a 1,000-fold greater than that produced by cNOS.

Production of NO from dietary nitrates

- Dietary nitrate is an important source of nitric oxide in mammals. Green, leafy vegetables, and some root vegetables (such as beetroot) have high concentrations
15 of nitrate. When eaten and absorbed into the bloodstream nitrate is concentrated in saliva (about 10 fold) and is reduced to nitrite on the surface of the tongue by a biofilm of commensal facultative anaerobic bacteria. This nitrite is swallowed and reacts with acid and reducing substances in the stomach (such as ascorbate) to produce high concentrations of nitric oxide.

- 20 An Acceptable Daily Intake (ADI) for nitrate of 3.7 mg/kg b.w./day, equivalent to 222 mg nitrate per day for a 60 kg adult was established by the former Scientific Committee on Food (SCF) and was reconfirmed by the Joint FAO/WHO Expert Committee on Food Additives (JECFA).

- The bioactivation of nitrate from dietary or endogenous sources requires its initial
25 reduction to nitrite, and because mammals lack specific and effective nitrate reductase enzymes, this conversion is mainly carried out by commensal bacteria in the gastrointestinal tract and on body surfaces. Nitrite is unique to the nitrogen oxides in its redox position between oxidative (NO_2 radical) and reductive (NO radical) signalling and its relative stability in blood and tissue. Once nitrite is

formed, there are numerous pathways in the body for its further reduction to NO, involving haemoglobin, myoglobin, xanthine oxidoreductase, ascorbate, polyphenols and protons. The generation of NO by these pathways is greatly enhanced during hypoxia and acidosis, thereby ensuring NO production in situations for which the oxygen-dependent NOS enzyme activities are compromised. Dietary nitrate is rapidly absorbed in the upper gastrointestinal tract. In the blood, it mixes with the nitrate formed from the oxidation of endogenous NO produced from the NOS enzymes.

Once in the oral cavity, commensal facultative anaerobic bacteria use nitrate as an alternative electron acceptor to oxygen during respiration, effectively reducing salivary nitrate to nitrite by the action of nitrate reductases. Human nitrate reduction requires the presence of these bacteria — suggesting a functional symbiosis relationship — as mammalian cells cannot effectively metabolize this anion. The salivary nitrate levels can approach 10 mM and nitrite levels 1–2 mM after a dietary nitrate load. When saliva enters the acidic stomach (1–1.5 l per day), much of the nitrite is rapidly protonated to form nitrous acid (HNO_2 ; $\text{pK}_a \sim 3.3$), which decomposes further to form NO and other nitrogen oxides. Nitrite reduction to NO is greatly enhanced by reducing compounds such as vitamin C and polyphenols, both of which are abundant in the diet.

The percentage of nitrate content in leafy vegetables is very less to significantly increase the nitric oxide level in the blood when taken orally. A larger quantity is required to be ingested at regular interval to get significant level of nitric oxide in the blood. Consumption of large quantity of leafy vegetables on regular basis is not convenient in such a busy life. Thus, providing a supplement developed from nitrate containing plant in the form of compact delivery system will be a convenient alternate to generate higher nitric oxide content in the blood. Moreover, as nitrates are required to be converted into nitrites in the mouth itself (by commensal facultative anaerobic bacteria), it will be more advantageous if the nitrate rich formulation can be delivered to oral cavity in the form of chewing

gum, lozenge, candy etc. Delivery via oral cavity offers certain advantages over conventional delivery of drugs.

The sites of drug administration in the oral cavity include the floor of the mouth (sublingual), the inside of the cheeks (buccal) and the gums (gingival). Drug

5 administration via oral cavity offers several distinct advantages:

- The buccal mucosa is relatively permeable with a rich blood supply, robust in comparison to the other mucosal tissues.
- Bypass the first-pass effect and non-exposure of the drugs to the gastrointestinal fluids.
- 10 • Improve the performance of many drugs, as they are having prolonged contact time with the mucosa.
- High patient acceptance compared to other non-oral routes of drug administration.
- As a result of adhesion and intimate contact, the formulation stays longer at the delivery site improving API bioavailability using lower API concentrations for disease treatment.
- 15 • Harsh environmental factors that exist in oral delivery of a drug are circumvented by buccal drug delivery.

A lot of work has been done for enhancing the level of nitric oxide in the blood for human beings and method of administration of medicaments / formulations, including those originated from plants (botanical source) available in nature in plenty. A short review of some of the prior art will elucidate how we have addressed the issue compared to other prior art and how effectively we could address enhancing the level of nitric oxide in the blood for human beings and the method of treating diseases as well as source for overcoming certain deficiencies of human beings.

20

25

US patent No.8303995, Bryan et al, discloses a nitrite formulations and their use as nitric oxide prodrugs. The Compositions comprising from about 40 weight

parts to about 1000 weight parts of a botanical nitrate source; from about 20 weight parts to about 500 weight parts of a botanical source of nitrite reduction activity; and from about 4 weight parts to about 100 weight parts of a nitrite salt. The botanical source of nitrite reduction activity is selected from the group
5 consisting of hawthorn berry, Schisandra, green tea, beet root, pine bark, holy basil, gymnema sylvestre, ashwagandha root, salvia, St. John wort, broccoli, stevia, spinach, ginkgo, kelp, tribulus, eleuthero, epimedium, eucommia, rhodiola, green tea, codonopsys, panax ginseng, astragalus, dodder seed, cordyceps, berries, tea, beer, grapes, wine, olive oil, chocolate, cocoa, coffee,
10 walnuts, peanuts, borojo, pomegranates, popcorn, yerba mate, and mixtures thereof. Methods of reducing triglycerides or reducing C-reactive protein levels are also provided.

Patent US patent No.8298589, Bryan et al, is related to 'Nitrite formulations and their use as nitric oxide prodrugs having a composition comprising nitrite salt
15 (10mg to 100mg), nitrate salt (50mg to 500 mg), ascorbic acid (100mg to 2000mg) of sodium and potassium.

Patent publication No US 20130071494, Bryan et al, also discloses a nitrite formulations and their use as nitric oxide prodrugs. The compositions comprise of from about 40 weight parts to about 1000 weight parts of a botanical nitrate
20 source; from about 20 weight parts to about 500 weight parts of a botanical source of nitrite reduction activity; and from about 4 weight parts to about 100 weight parts of a nitrite salt Use of said composition in methods of reducing triglycerides or reducing C-reactive protein levels are also disclosed here.

Patent publication No US 20120321724, and WO 201323217 Bryan et al,
25 discloses a method of measuring in vivo nitric oxide and nitrite levels in individuals by providing a salivary nitrite test substrate, testing salivary nitrite levels with the test substrate, measuring nitrite levels detected in the testing; and correlating the measured nitrite levels with in vivo nitric oxide bio-availability.

In patent No. US 8435570, Bryan et al, disclose a composition comprising a
30 nitrite salt, a nitrate salt, and ascorbic acid. A method of enhancing

cardiovascular performance or treating adverse cardiovascular event in a mammal is also provided.

Patent publication No 20130071371, Bryan et al, is related to a composition and method of providing nitric oxide and nitrite therapy to patients whereby a
5 therapeutic amount including a dispersable medium is bioavailable within approximately 30 minutes of administration including. In embodiments of the disclosure, nitric oxide is produced in the oral cavity.

In the above patents, the nitrate source is either synthetic one or from a botanical nitrate source which includes spinach, beet root, artichoke, holybasil, green tea
10 etc.

Supplementation of Amaranth extract enriched with nitrate enhances the nitrate and nitrite levels in the body. Presence of vitamin C promotes the conversion of nitrite to nitric oxide in body thereby increasing the antioxidant potential in body. The formulation of nitrate enriched Amaranth extract with Vitamin C in the
15 present formulation further improves the generation of nitric oxide by promoting the conversion of nitrite delivered from Amaranth extract to nitric oxide in the body. Vitamin C used may be of synthetic or natural origin.

Amaranthus are dietary leafy vegetables belonging to the family Amaranthaceae. Amaranth belongs to the genus Amaranthus and various species include
20 *Amaranthus dubius*, *Amaranthus spinosus*, *Amaranthus fimbriatus*, *Amaranthus floridanus*, *Amaranthus graecizans* etc. Other dietary sources of nitrate are Spinach (*Spinacia oleracea*, family: Amaranthaceae), beet root (*Beta vulgaris*, family: Amaranthaceae), lettuce (*Lactuca sativa*, family: Asteraceae) etc.

The genus amaranthus contains approximately 70 species of worldwide
25 distribution including pigweeds, water hemsps, and grain amaranthus. For human consumption there are cultivated grain amaranths –*A.caudatus*, *A. cruentus* and *A. hypochondriacus* and vegetable amaranths – mainly *A.dubius*, *A.tricolor* and *A.cruentus*. Nowadays, the grain amaranthus are cultivated from the temperate to

tropical zone and the vegetable amaranths mainly in the South Africa and South Asia.

These plants, which grow 90 centimeters to 150 centimeters tall, are abundant weeds in many parts of the world. All amaranth have alternate simple leaves.

5 They may have some red color present on the stems. They bear minute, greenish flowers in dense clusters at the top of the plants. Their seeds may be brown or black in weedy species and light-colored in domestic species. Some amaranth species have been grown as a grain crop and a garden vegetable in various parts of the world, especially in South America.

10 Amaranthus are very promising crops. The main reasons could be content of protein, fat and active substances. The content of seed protein is in the range 13 – 18% with very good balanced amino acids. The lysine content is relatively high in the comparison with common cereals. The content of crude proteins in leaves is from 27 to 49% in d.m. what is more than in the leaves in the spinach.

15 Amaranthus have comparable or higher amounts of essential amino acids as whole egg protein. The fat content is in the range 0.8-8.0%. The linoleic acid is the predominant fatty acid, with lesser amount of oleic and palmitic acids.

SUMMARY OF THE INVENTION:

Disclosure provides a medicinal composition obtained from the extract of
20 Amaranth. Disclosure provides an extract of Amaranth having low oxalic acid content. The Amaranth extract includes

about 0.1% to about 20% nitrates;

about 0.1% to about 10% of L-arginine;

about 0.001% to about 10% of flavonoids;

25 about 0.01% to about 15% of saponins;

about 0.01% to about 10% of alkaloids;

about 10% to about 30% of carbohydrates;

about 5% to about 25% of proteins;

about 5 % to about 25% potassium; and,

about 0.01% to about 50% of total oxalic acid. Determination of total oxalic acid content included measurement of free oxalic acid and oxalates in the extract.

5 The disclosure also provides oral dosage forms of the composition of Amaranth such as fast melt tablets, lozenge, chewing gum etc to make it convenient for buccal absorption. The disclosure also provides dosage forms such as tablets, capsules, pills, powders etc for absorption from the stomach.

10 The disclosure provides methods for producing the Amaranth extract. The extract of Amaranth can be prepared from fresh or from dried leaves and stem of Amaranth.

Some embodiments of the composition are found to be useful for the enhancing the amount of nitric oxide in blood plasma. Some embodiments enhance the amount of nitric oxide in saliva. Some embodiments enhance the level of nitrate and nitrite in blood plasma. Some embodiments of the extract enhance the level
15 of nitrate and nitrite in saliva. Some embodiments enhance the antioxidant activity in mammals. In some embodiments, the composition lowers blood pressure (hypertension). Some embodiments of the composition increase endurance.

20 In some embodiments, the Amaranth extract can be fortified by adding amla extract.

BRIEF DESCRIPTION OF THE DRAWINGS

Other features and advantages of the disclosure should become apparent from the following description of the preferred process and read in conjunction with the figures of the accompanying drawings, which demonstrate the process steps
25 involved in various process trials carried under certain preferred embodiment of the invention as well as the findings.

Fig. 1: Flow chart- Extraction of Fresh Amaranth with methanol.

Fig. 2: Flow chart- Extraction of Fresh Amaranth with water.

- Fig. 3: Flow chart- Extraction of Dried Amaranth with methanol.
- Fig. 4: Flow chart- Extraction of dried Amaranth with water.
- Fig. 5: Flow chart- Extraction of Fresh Amaranth by treating with pectinase.
- Fig. 6: Flow chart- Extraction of Fresh Amaranth by treating with pectinase and
5 removal of Oxalic acid.
- Fig. 7: Flow chart- Hexane Extract of Fresh Amaranth after treating with
pectinase and removal of Oxalic acid.
- Fig. 8: Flow chart-Chloroform Extract of Fresh Amaranth after treating with
pectinase and removal of Oxalic acid.
- 10 Fig. 9: Flow chart- EDC Extract of Fresh Amaranth after treating with pectinase
and removal of Oxalic acid.
- Fig. 10: Flow chart- Extraction of Amla.
- Fig. 11: Flow chart- Hexane Extract of Fresh Amaranth after treating with
pectinase and removal of Oxalic acid in combination with Amla.
- 15 Fig. 12: Flow chart- Preparation of dried Amaranth leaf powder.
- Fig. 13: Flow chart - Preparation of Amaranth Juice.
- Fig. 14 (Part A): Table showing the percentage of actives in different extract of
Amaranth and fast dissolving tablets
- Fig. 14 (Part B): Table showing the percentage of actives in different extract of
20 Amaranth and fast dissolving tablets

DETAILED DESCRIPTION THE INVENTION

The disclosure provides an entirely different medicinal composition prepared
from Amaranth as nitrate source by a unique method of extraction. The disclosed
25 processes provide an extract of Amaranth enriched with nitrate, L-arginine,
flavonoids, carbohydrate, protein, saponins, alkaloids, potassium and having

negligible oxalic acid or oxalate content. Many vegetables contain high amount of total oxalic acid in the form of free oxalic acid or oxalate salts and the oxalic or oxalate content is toxic to human health. The extracts from vegetables contain oxalic acid along with other active constituents. The extract of Amaranth also
5 contains oxalic acid and oxalates. But in the disclosed methods, we removed the oxalic acid and oxalates from the Amaranth extract to provide an extract of Amaranth having negligible oxalic acid or oxalate content. The total oxalic acid content was measured by HPLC method and provides the total of both free oxalic acid and oxalate salts in the extract.

10 The composition obtained from the enriched extract of amaranth is useful for lowering the blood pressure (hypertension), enhancing the level of nitric oxide in the blood plasma and saliva, enhancing the level of nitrate and nitrite in blood plasma and saliva, enhance the antioxidant activity and increase the endurance. The disclosure relates to a medicinal composition of Amaranth extract. The
15 composition has an enriched nitrate content, L-arginine, flavonoids, carbohydrate, protein, saponins, alkaloids, potassium and a negligible amount of oxalic acid or oxalate.

The Amaranth extract can be obtained from both fresh and dry leaves and stem of Amaranth. The extract when administered to mammals especially to human
20 beings can lower the blood pressure (hypertension) and increase the endurance. If the same extract is further enriched by enzyme treatment it can enhance the capability of lowering the blood pressure (hypertension) and increasing the endurance.

It is further noted that the composition of enriched Amaranth extract origin has
25 the property to enhance the amount of nitric oxide in blood plasma and saliva as well as the property to enhance the level of nitrate and nitrite in blood plasma and saliva. The composition is also found to be useful for enhancing the antioxidant activity.

Some embodiments provide a composition obtained from the extract of Amaranth
30 from fresh leaves and stem of Amaranth comprising enriched nitrate content, L-

Arginine, flavonoids, saponins, alkaloids, potassium and having negligible amount of oxalic acid or oxalate content wherein nitrates are present in the extract in a range of 3% to 20%, L-Arginine are present in the extract from 1% to 10%, flavonoids are present in the extract in a range of 1% and 10%, saponins are present in the range of 5% to 15%, alkaloids are present in the range of 3% to 10%. The extract also contains carbohydrate from 10% to 30%, potassium content from 5% to 25% and proteins are present in a range of 5% to 25%.

Some embodiments provide an extract of Amaranth having low total oxalic acid content. The extract includes:

- 10 about 0.1% to about 20% nitrates;
about 0.1% to about 10% of L-arginine; (0.1-2%; 0.1-6%; 0.1-10%)
about 0.001% to about 10% of flavonoids;
about 0.01% to about 15% of saponins;
about 0.01% to about 10% of alkaloids; (0.01-5%)
- 15 about 10% to about 30% of carbohydrates;
about 5% to about 25% of proteins;
about 5 % to about 25% potassium; and,
about 0.01% to about 50% of total oxalic acid. The total oxalic acid includes free oxalic acid and oxalate salts in the extract.
- 20 In some embodiments of the extract, nitrates ranges from about 0.1% to about 3%. In some embodiments of the extract, nitrates ranges from about 1% to about 10%. In some embodiments of the extract, nitrates ranges from about 10% to about 20%. In some embodiments of the extract, nitrates ranges from about 3% to about 20%. In some embodiments of the extract, L-arginine ranges from
- 25 about 0.1% to about 2%. In some embodiments of the extract, L-arginine ranges from about 0.1% to about 6%. In some embodiments of the extract, L-arginine ranges from about 0.1% to about 10%. In some embodiments of the extract, alkaloids ranges from about 0.01% to about 5%. In some embodiments of the

extract, total oxalic acid ranges from about 0.01% to about 0.1%. Total oxalic acid content in the extracts was determined by HPLC method to measure both free oxalic acid and oxalates content in the extracts. In some embodiments of the extract, total oxalic acid ranges from about 0.01% to about 1%. In some
5 embodiments of the extract, total oxalic acid ranges from about 0.1% to about 1%. In some embodiments of the extract, total oxalic acid ranges from about 1% to about 10%. In some embodiments of the extract, total oxalic acid ranges from about 1% to about 20%. In some embodiments of the extract, total oxalic acid ranges from about 1% to about 30%. In some embodiments of the extract, total
10 oxalic acid ranges from about 1% to about 40%. In some embodiments of the extract, total oxalic acid ranges from about 1% to about 50%. In some embodiments of the extract, total oxalic acid ranges from about 10% to about 20%.

Some embodiments provide an extract of Amaranth having low total oxalic acid
15 content. The extract includes

about 3% to about 20% nitrates;
about 1% to about 10% of L-arginine;
about 1% to about 10% of flavonoids;
about 5% to about 15% of saponins;
20 about 3% to about 10% of alkaloids;
about 10% to about 30% of carbohydrates;
about 5% to about 25% of proteins;
about 1% to about 5% of potassium; and,
about 0.1% total oxalic acid. Total oxalic acid was measured by HPLC method
25 to determine free oxalic acid and oxalates in the extract.

The disclosure provides extracts having enriched nitrate, L-arginine, flavonoids, carbohydrate, protein, saponins, alkaloids, potassium and having negligible oxalic acid or oxalate content wherein the extracts may be obtained from red

Amaranth or Spleen amaranth or red Spinach (*Amaranthus dubius*) or any related species of amaranth like *A. caudatus*, *A. cruentus*, *A. hypochondriacus*, *A. tricolor*, *A. blitum*, *A. viridis* etc.

The disclosure also describes delivery of the nitrate enriched formulations orally
5 as fast melt tablets, lozenge, candy, chewing gum, beverage etc to make it convenient for absorption from the mouth and also in the form of tablets, capsules, pills, powders etc to be absorbed from the stomach. Some embodiments provide a dosage form having the Amaranth extract. The dosage form is selected from the group consisting of fast melt tablet, lozenge, candy, chewing gum,
10 beverage, tablet, capsule, pill, and powder.

Some embodiments provide a method of lowering blood pressure by administering the extract of Amaranth. Some embodiments provide a method of enhancing nitric oxide concentration in blood following administration of the Amaranth extract. Some embodiments provide, a method of increasing physical
15 endurance by administering an extract of Amaranth. Some embodiments provide a method for increasing swimming endurance by administering the Amaranth extract. Some embodiments provide a method for increasing running endurance by administering an Amaranth extract. Some embodiments provide a method for improving sexual performance by administering an Amaranth extract. Some
20 embodiments provide a method of enhancing nitrate level in blood by administering an extract of Amaranth. Some embodiments provide a method of enhancing nitrite level in blood by administering an extract of Amaranth. Some embodiments provide a method of enhancing nitrate level in saliva by administering an extract of Amaranth. Some embodiments provide a method of
25 enhancing nitrite level in saliva by administering an extract of Amaranth. Some embodiments provide a method of enhancing nitric oxide level in saliva by administering an extract of Amaranth.

In some embodiments as shown in Example 26, administering fast dissolving tablet containing Amaranth extract enriched with nitrate, led to increased levels
30 of NO₃, NO₂ and NO content in plasma compared to plasma levels prior to

administration of tablet. In some embodiments, NO₃ level increases about 6 fold the baseline value after 1 hour. In some embodiments, NO₂ level increases more than 6 fold. In some embodiments, NO level increases by about 6 fold.

5 In some embodiments as shown in Example 27, administering fast dissolving tablet containing Amaranth extract enriched with nitrate, led to increased NO₃, NO₂ and NO levels in saliva when compared to baseline prior to administration of the tablet. In some embodiments, NO₃ level increases about 6.9 fold of the baseline value after 1 hour. In some embodiments, NO₂ level increases more than 9.2 fold. In some embodiments, NO level increases 7.5 fold.

10 In some embodiments as shown in Example 28, administering Amaranth extract enriched with nitrate mixed with water, led to increased levels of NO₃, NO₂ and NO level in saliva when compared to levels in saliva prior to administration of extract. In some embodiments, NO₃ level increases 6.6 fold the baseline value after 1 hour. In some embodiments, NO₂ level increases 9.1 fold. In some
15 embodiments, NO level increases 7.3 fold.

In some embodiments, administration of Amaranth extract enhanced plasma level of nitrate by 2 fold to 10 fold. In some embodiment, administration of Amaranth extract enhanced plasma level of nitrite by about 3 fold to about 15 fold. In some
20 embodiments, administration of Amaranth extract enhanced plasma level of nitric oxide by about 2 fold to about 10 fold.

In some embodiments, administration of Amaranth extract enhanced saliva level of nitrate by 2 fold to 10 fold. In some embodiment, administration of Amaranth extract enhanced saliva level of nitrite by about 3 fold to about 15 fold. In some
25 embodiments, administration of Amaranth extract enhanced saliva level of nitric oxide by about 2 fold to about 10 fold.

In some embodiments, administering Amaranth extract enriched with nitrate enhanced the nitrate and nitrite levels in the body. It is likely that presence of vitamin C promotes the conversion of nitrite to nitric oxide in body thereby increasing the antioxidant potential in body. Some embodiments provide a

formulation of nitrate enriched Amaranth extract with Vitamin C, wherein administering the formulation further improves the generation of nitric oxide by promoting the conversion of nitrite delivered from Amaranth extract to nitric oxide in the body. Vitamin C added to the Amaranth extract may be of synthetic or natural origin.

The disclosure provides different extracts of Amaranth extracted by using solvents like water, methanol, ethanol, isopropanol, n-butanol, methyl acetate, ethyl acetate, propyl acetate, n-butyl acetate and combinations thereof.

Low molecular weight alcohols that can be used in preparation of the extract include methanol, ethanol, isopropanol, n-butanol and combinations thereof. Esters that can be used for extract preparation include methyl acetate, ethyl acetate, propyl acetate, n-butyl acetate and combinations thereof. Alkanes that can be used for preparation of the extract include pentane, hexane, heptane, isooctane, and combinations thereof.

Various species of Amaranth from which extract can be prepared include, *Amaranthus acanthochiton*, *Amaranthus acutilobus*, *Amaranthus viridis*, *Amaranthus albus*, *Amaranthus arenicola*, *Amaranthus australis*, *Amaranthus bigelovii*, *Amaranthus blitoides*, *Amaranthus blitum*, *Amaranthus brownie*, *Amaranthus californicus*, *Amaranthus cannabinus*, *Amaranthus caudatus*, *Amaranthus chihuahuensis*, *Amaranthus chlorostachys*, *Amaranthus crassipes*, *Amaranthus crispus*, *Amaranthus cruentus*, *Amaranthus deflexus*, *Amaranthus dubius*, *Amaranthus fimbriatus*, *Amaranthus floridanus*, *Amaranthus graecizans*, *Amaranthus greggii*, *Amaranthus hybridus*, *Amaranthus hypochondriacus*, *Amaranthus leucocarpus*, *Amaranthus lineatus*, *Amaranthus lividus*, *Amaranthus mantegazzianus*, *Amaranthus minimus*, *Amaranthus muricatus*, *Amaranthus obcordatus*, *Amaranthus oleraceus*, *Amaranthus palmeri*, *Amaranthus paniculus*, *Amaranthus polygonoides*, *Amaranthus powellii*, *Amaranthus pringlei*, *Amaranthus pumilus*, *Amaranthus quitensis*, *Amaranthus retroflexus* etc.

In some embodiments, the extract of Amaranth includes a second extract. The second extract is selected from the group consisting of amla extract, turmeric

extract, grape seed extract, green tea extract, pomegranate extract, cocoa extract, coconut root extract, rosemary extract, mint leaf extract, star anise, sweet basil extract, cinnamon extract/clove extract, ginger extract, cumin seed extract, black pepper extract, fenugreek extract, or combinations thereof.

- 5 In some embodiments of the extract of Amaranth, the Amaranth is selected from the group consisting of *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus tricolor*, *Amaranthus blitum*, *Amaranthus viridis*, *Amaranthus dubis* or combinations thereof.

Various methods for the preparation of the extract enriched with nitrate content,
10 L-arginine, flavonoids, carbohydrate, protein, saponins, alkaloids, potassium and having negligible oxalic acid or oxalate content prepared by the extraction of fresh leaves of Amaranth are as under:

In some embodiments (refer Fig. 1), the fresh leaves and stem of Amaranth is cleaned, crushed and extracted for 5 hrs using methanol at 65°C in a Soxhlet
15 extractor and then filtered. The resultant extract is concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C to form concentrated extract. Later the concentrated extract is dried under vacuum at above 500 mm of mercury to form powder of "methanol extract" of fresh Amaranth.

In some embodiments, a method of preparing a powdered extract of fresh
20 Amaranth is provided. The method includes crushing fresh leaves and stem of Amaranth to obtain a crushed material. Then extracting the crushed material with methanol to obtain a supernatant and residue. Then filtering the supernatant to obtain a filtrate. Then concentrating the filtrate to obtained a concentrated filtrate. Then drying the concentrated filtrate to obtain the powdered extract of
25 fresh Amaranth.

According to another embodiment (refer Fig. 2), the fresh leaves and stem of Amaranth is cleaned, crushed and extracted for about 1 hr using water in an extractor with reflux condenser to obtain residue and supernatant. The residue and supernatant is separated by draining out the supernatant from the extractor

bottom through the filter cloth. The resultant supernatant is concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated extract. Later the concentrated extract is dried under vacuum at above 500 mm of mercury to form powder of "water extract" of fresh Amaranth.

- 5 Some embodiments provide a method of preparing a powdered extract of fresh Amaranth. The method includes crushing fresh leaves and stem of Amaranth to obtain a crushed material. Then extracting the crushed material with water to obtain a supernatant and a residue. Next, concentrating the supernatant to obtain a concentrated water extract. Then drying the concentrated water extract to
10 obtain a powdered extract of fresh Amaranth.

In some embodiments (refer to fig-3), the fresh leaves and stem of Amaranth is cleaned, dried and powdered. Then the powder is extracted for 5 hrs using methanol at 65°C in a Soxhlet extractor and then filtered. The resultant extract is concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C
15 to form concentrated extract. Later the concentrated extract is dried under vacuum at above 500 mm of mercury to form powder of "methanol extract" of dried Amaranth.

Some embodiments provide a method of preparing a powdered extract of dried Amaranth. The method includes powdering dried leaves and stem of Amaranth
20 to obtain a powdered material. Next, the powdered material is extracted with methanol to obtain a supernatant. Then the supernatant is filtered to obtain a filtrate. Next, concentrating the filtrate to obtain a concentrated methanol filtrate. Then drying the concentrated methanol filtrate to obtain the powdered extract of dried Amaranth.

- 25 According to another embodiment (refer to Fig. 4), the fresh leaves and stem of amaranth is cleaned, dried and powdered. The powder is extracted for about 1hr using water in an extractor with reflux condenser to obtain residue and supernatant. The residue and supernatant is separated by draining out the supernatant from the extractor bottom through the filter cloth. The resultant
30 supernatant is concentrated in an Agitated thin film evaporator (ATFE) at a

temperature of 85°C to form concentrated extract. Later the concentrated extract is dried under vacuum at above 500 mm of mercury to form powder of “water extract” of dried Amaranth.

Some embodiments provide a method of preparing a powdered extract of dried Amaranth. The method includes powdering dried leaves and stem of Amaranth to obtain a powdered material. Then, the powdered material is extracted with water to obtain a supernatant and a residue. Next, the supernatant is concentrated to obtain a concentrated water extract. The concentrated water extract is dried to obtain a powdered extract of dried Amaranth.

10 In another embodiment (refer to Fig. 5), the fresh leaves and stem of amaranth is cleaned and crushed to form slurry. The slurry is treated with pectinase (1%) for 2 hrs. Then the enzyme is deactivated by heating the slurry at 90°C for 10 minutes. The pectinase treated amaranth slurry is extracted for about 1 hr using water in an extractor with reflux condenser to obtain residue and supernatant. The residue and supernatant is separated by draining out the supernatant from the extractor bottom through the filter cloth. The resultant supernatant is concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated extract. Later the concentrated extract is dried under vacuum at above 500 mm of mercury to form powder of pectinase treated “water extract” of Amaranth.

In some embodiments of the method of preparing the Amaranth extract, the method includes crushing fresh leaves and stem of Amaranth to obtain a slurry. Then treating the slurry with pectinase to obtain a pectinase treated material. Then heating the pectinase treated material to obtain a pectinase-deactivated material. Next, extracting the pectinase-deactivated material with water to obtain a supernatant and a residue. Then, concentrating the supernatant to obtain a concentrated supernatant. And drying the concentrated supernatant to obtain a powdered extract of fresh Amaranth.

30 In yet another embodiment, (refer to Fig. 6), the fresh leaves and stem of Amaranth is cleaned and crushed to form slurry. The slurry is treated with

pectinase (1%) for 2 hrs. Then the enzyme is deactivated by heating the slurry at 90°C for 10 minutes. The pectinase treated Amaranth slurry is extracted for about 1 hr using water in an extractor with reflux condenser to obtain residue and supernatant. The residue and supernatant is separated by draining out the supernatant from the extractor bottom through the filter cloth. The resultant supernatant is concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated extract. Keep the concentrated water extract at 10°C for 48 hr to crystallize oxalic acid or oxalates. Decant the supernatant and again keep the supernatant at 10°C for another 24 hr to crystallize remaining oxalic acid or oxalates. Decant and filter the supernatant and dry under vacuum at above 500 mm of mercury to get oxalic acid or oxalate free extract of Amaranth.

In some embodiments a method of preparing an extract of Amaranth is provided. The method includes crushing fresh leaves and stem of Amaranth to obtain a slurry, then treating the slurry with pectinase to obtain a pectinase treated material, followed by heating the pectinase treated material to obtain a pectinase-deactivated material. Next the pectinase-deactivated material is extracted with water to obtain a supernatant and a residue. The supernatant is concentrated to obtain a concentrated water extract. The concentrated water extract is cooled at 10°C for 48 hours then filtered to obtain crystals of oxalic acid or oxalate and a second supernatant. The second supernatant is cooled at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid or oxalate crystals. Then the third supernatant is filtered to obtain a filtrate. Then the filtrate is concentrated to obtain a concentrated filtrate. And the concentrated filtrate is dried to obtain a powdered extract of Amaranth. In some embodiments, the powdered extract has less than 0.1% oxalic acid.

In accordance with another embodiment (refer to Fig. 7) an Amaranth extract having enriched nitrate content is provided. A dried powder of oxalic acid or oxalate free extract of Amaranth is obtained after treating with pectinase and removal of oxalic acid is further extracted with hexane for 6 hours in a soxhlet

extractor. Then filter and concentrate the hexane extract in an Agitated thin film evaporator (ATFE). The concentrated hexane extract is dried under vacuum to get powder of hexane extract of fresh Amaranth after treating with pectinase and removal of oxalic acid and having enriched nitrate content.

5 Some embodiments provide a method of preparing an extract of Amaranth having enriched nitrate content. The method includes crushing fresh leaves and stem of Amaranth to obtain a slurry. The slurry is treated with pectinase to obtain a pectinase treated material. Next, the pectinase treated material is heated to deactivate the pectinase and obtain a pectinase-deactivated material. Next, the
10 pectinase-deactivated material is extracted with water to obtain a supernatant and a residue. Next, the supernatant is concentrated to obtain a concentrated water extract. The concentrated water extract is cooled at 10°C for 48 hours and then filtered to obtain crystals of oxalic acid or oxalate and a second supernatant. The second supernatant is cooled at 10°C for 24 hours to obtain a third supernatant
15 and a second oxalic acid or oxalate crystals. The third supernatant is filtered to obtain a filtrate. The filtrate is concentrated to obtain a concentrated filtrate. The concentrated filtrate is dried to obtain a powdered extract. Next the powdered extract is treated with hexane to obtain a mixture. Then the mixture is filtered to obtain a filtrate. Then the filtrate is concentrated to obtain a concentrated hexane
20 extract. Then the concentrated hexane extract is dried to obtain a powdered hexane extract.

In another embodiment (refer Fig. 8), the dried powder of oxalic acid or oxalate free extract of Amaranth obtained after treating with pectinase and removal of oxalic acid is further extracted with chloroform by cold extraction instead of
25 hexane. Filter and concentrate the chloroform extract in an Agitated thin film evaporator (ATFE). Then it is dried under vacuum to get powder of chloroform extract of fresh Amaranth after treating with pectinase and removal of oxalic acid.

Some embodiments provide a method of preparing an extract of Amaranth. The
30 method includes crushing fresh leaves and stem of Amaranth to obtain a slurry.

Then slurry is treated with pectinase to obtain a pectinase treated material. Then the pectinase treated material is heated to obtain a pectinase-deactivated material. Then the pectinase-deactivated material is extracted with water to obtain a supernatant and a residue. Then the supernatant is concentrated to obtain a concentrated water extract. Then the concentrated water extract is cooled at 10 C for 48 hours followed by filtering to obtain crystals of oxalic acid or oxalate and a second supernatant. Then the second supernatant is cooled at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid or oxalate crystals. Then the third supernatant is filtered to obtain a filtrate. Then the filtrate is concentrated to obtain a concentrated filtrate. Then the concentrated filtrate is dried to obtain a powdered extract. Then the powdered extract is treated with chloroform to obtain a mixture. The mixture is filtered to obtain a filtrate. The filtrate is concentrated to obtain a concentrated chloroform extract. The concentrated chloroform extract is dried to obtain a powdered chloroform extract.

15 According to yet another embodiment (refer to Fig. 9), the dried powder of oxalic acid or oxalate free extract of Amaranth obtained after treating with pectinase and removal of oxalic acid is further extracted with ethylene dichloride (EDC) for 6 hours in a soxhlet extractor instead of Hexane/ Chloroform. Filter and concentrate the chloroform extract in an Agitated thin film evaporator (ATFE).

20 Then it is dried under vacuum to get powder of chloroform extract of fresh Amaranth after treating with pectinase and removal of oxalic acid.

Some embodiments provide a method of preparing an extract of Amaranth. The method includes crushing fresh leaves and stem of Amaranth to obtain a slurry. The slurry is treated with pectinase to obtain a pectinase treated material. The pectinase treated material is heated to obtain a pectinase-deactivated material. The pectinase-deactivated material is extracted with water to obtain a supernatant and a residue. The supernatant is concentrated to obtain a concentrated water extract. The concentrated water extract is cooled at 10°C for 48 hours followed by filtering to obtain crystals of oxalic acid or oxalate and a second supernatant.

30 The second supernatant is cooled at 10°C for 24 hours to obtain a third

supernatant and a second set of crystals of oxalic acid or oxalate. The third supernatant is filtered to obtain a filtrate. The filtrate is concentrated to obtain a concentrated filtrate. The concentrated filtrate is dried to obtain a powdered extract. The powdered extract is dried with ethylene dichloride to obtain a mixture. The mixture is filtered to obtain a second filtrate. The second filtrate is concentrated to obtain a concentrated ethylene dichloride extract. The concentrated ethylene dichloride extract is dried to obtain a powdered ethylene dichloride extract.

In another embodiment (refer to Fig. 11), the dried powder of oxalic acid or oxalate free extract of Amaranth obtained after treating with pectinase and removal of oxalic acid is further extracted with hexane for 6 hours in a soxhlet extractor. Then filter and concentrate the hexane extract in an Agitated thin film evaporator (ATFE). The concentrated hexane extract is dried under vacuum to get powder of hexane extract. It is dissolved in methanol and Amla extract already prepared is added and stirred for 1 hr at 100 rpm using mechanical stirrer. Evaporate and dry under vacuum to form powder. Thus a hexane extract of fresh Amaranth after treating with pectinase and removal of oxalic acid in combination with Amla is obtained.

The Amla extract (refer to fig.10) used here can be prepared according to the following method.

Pulp the fruits of amla and extract the pulp using methanol at 65°C in a Soxhlet extractor for 5 hrs. The resultant methanol extract is filtered and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C to form concentrated extract. Dry the concentrated extract under vacuum at above 500 mm of mercury to form powder of methanol extract of Amla.

Some embodiments provide a method of preparing an extract of Amaranth. The method includes crushing fresh leaves and stem of Amaranth to obtain a slurry. Then the slurry is treated with pectinase to obtain a pectinase treated material. The pectinase treated material is heated to obtain a pectinase-deactivated material. The pectinase-deactivated material is extracted with water to obtain a

supernatant and a residue. The supernatant is concentrated to obtain a concentrated water extract. The concentrated water extract is cooled at 10°C for 48 hours followed by filtering to obtain crystals of oxalic acid or oxalate and a second supernatant. The second supernatant is cooled at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid or oxalate crystals. The third supernatant is filtered to obtain a filtrate. The filtrate is concentrated to obtain a concentrated filtrate. The concentrated filtrate is dried to obtain a powdered extract. The powdered extract is treated with hexane to obtain a mixture. The mixture is filtered to obtain a second filtrate. The second filtrate is concentrated to obtain a concentrated hexane extract. The concentrated hexane extract is dried to obtain a powdered hexane extract. The powdered hexane extract is dissolved in methanol to obtain a solution of hexane extract of Amaranth. A powder of methanol extract of Amla is added to the solution of hexane extract of Amaranth to obtain a second mixture. The second mixture is evaporated to obtain an evaporate. The evaporate is dried to obtain a powder of Amaranth and Amla extract. The powdered methanol extract of Amla extract is prepared by a process including pulping fresh Amla (*Emblica officinalis*) fruit to obtain a pulp. The pulp is extracted with methanol to obtain a filtrate. The filtrate is concentrated to obtain a concentrated methanol extract of Amla. The concentrated methanol extract of Amla is dried to obtain the powdered methanol extract of Amla.

In some embodiments, amla extract / turmeric extract/ grape seed extract/ green tea extract/ pomegranate extract/cocoa extract/ coconut root extract/ rosemary extract/ mint leaf extract/ star anise /sweet basil extract/cinnamon extract/clove extract/ginger extract/cumin seed extract/black pepper extract/ fenugreek extract are added to the pectinase treated fresh leaves of Amaranth having low total oxalic acid or oxalate content to fortify the Amaranth extract. The combination extracts can be administered to a mammal for enhancing the amount of nitric oxide in blood.

In some embodiments (refer to Fig. 13), the fresh leaves of amaranth is cleaned and crushed in the screw expeller. Initial Juice obtained on the tray of the screw expeller is collected in the feed tank under room temperature of 27°C. Initial juice

is centrifuged to obtain filtrate and residue is removed. Filtrate is clarified in a rotating disc type clarifier (RPM- 18000) to obtain final filtrate and residue. Residue is removed and final filtrate is collected (clarified liquid juice of leaves of Amaranth).

- 5 Details of some of the trials/ experiments carried out and findings are explained below by way of examples.

Example 1

Preparation of Amaranth fresh leaves extract with methanol.

- 10 Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed. Crushed materials were filled in the Soxhlet extractor and extracted with methanol (300 L). The extraction was carried out for 5 hrs at a temperature of about 65° C. After the completion of extraction, the supernatant was filtered and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C to form concentrated methanol extract. Concentrated
15 methanol extract was dried under vacuum at above 500 mm of mercury to get powder of methanol extract of fresh Amaranth (Yield 2Kg). See also extract preparation in Fig. 1.

The nitrate content in methanol extract of fresh Amaranth by ion chromatography was found to be 1.5%.

20 Example 2

Preparation of Amaranth fresh leaves extract with water.

- 25 Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed. Water in an amount ten times the quantity of crushed material of Amaranth was added to form a mixture. The extraction was performed using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor

bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant.

- 5 All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract. Concentrated water extract was dried under vacuum at above 500 mm of mercury to get powder of water extract of fresh Amaranth. (Yield 3%). See also extract preparation in Fig. 2.
- 10 The nitrate content in water extract of fresh Amaranth by ion chromatography was found to be 1.6%.

Example 3

Preparation of Amaranth dried leaves extract with methanol.

- Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and dried. Dried leaves and stem were powdered. Powdered Amaranth were filled in the Soxhlet apparatus and extracted with methanol (500 L). The extraction was carried out for 5 hrs at a temperature of about 65°C. After the completion of extraction, the supernatant was filtered and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C to form concentrated methanol extract. Concentrated methanol extract was dried under vacuum at above 500 mm of mercury to get powder of methanol extract of dried amaranth. When starting with 100 kg of fresh amaranth, a yield of 24 Kg of powder of methanol extract of dried amaranth was obtained. See also extract preparation in Fig. 3.

- 25 The nitrate content in methanol extract of dried Amaranth by ion chromatography was found to be 1.65%.

Example 4

Preparation of Amaranth dried leaves extract with water .

Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and dried. Dried Amaranth was powdered. Water in an amount ten times the quantity of powder of dried Amaranth was added to the powder of Amaranth to form a mixture. The extraction was performed using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract. Concentrated water extract was dried under vacuum at above 500 mm of mercury to get powder of water extract of dried Amaranth. (Yield 25%). See also extract preparation in Fig. 4.

The nitrate content in water extract of dried Amaranth by ion chromatography was found to be 1.7%.

Example 5

Enhanced yield of Amaranth extract by treatment with pectinase.

Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed to make slurry. The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was deactivated by heating the slurry at 90°C for 10 minutes. After the enzyme treatment, the pectinase treated fresh leaves were extracted with water. The extraction was performed with water using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the

- polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract. Concentrated water extract was dried under vacuum at above 500 mm of mercury to get powder of water extract of fresh Amaranth (Yield 3.5Kg). See also extract preparation in Fig. 5.
- 10 Nitrate content of the extract was found to be 2% as determined by ion chromatography.

Example 6

Oxalic acid or oxalates free Amaranth extract.

- Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed to make slurry. The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was deactivated by heating the slurry at 90°C for 10 minutes. After the enzyme treatment, the pectinase treated fresh leaves were extracted with water. The extraction was performed with water using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract.

Concentrated water extract was kept at 10°C for 48 hr to crystallize oxalic acid or oxalates. The supernatant was decanted off and again kept at 10°C for 24 hr to crystallize remaining oxalic acid or oxalates. The supernatant from this solution was decanted, filtered and evaporated to dryness under vacuum to get oxalic acid or oxalate free Amaranth extract (2.2 Kg). See also extract preparation in Fig. 6

Nitrate content of the extract was found to be 5% as determined by ion chromatography.

Example 7

Enrichment of nitrates in Amaranth extract by extraction with hexane.

10 Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed to make slurry. The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was deactivated by heating the slurry at 90°C for 10 minutes. The extraction was performed with water using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene
15 (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and
20 supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract.

Concentrated water extract was kept at 10°C for 48 hr to crystallize oxalic acid or
25 oxalates. The supernatant was decanted off and again kept at 10°C for 24 hr to crystallize remaining oxalic acid or oxalates. The supernatant from this solution was decanted, filtered and evaporated to dryness under vacuum to get oxalic acid or oxalate free Amaranth extract (2.2 Kg).

This oxalic acid or oxalate free Amaranth extract was further extracted with hexane for 6 hours in a soxhlet extractor. The extract was filtered and evaporated to dryness under vacuum to get nitrate enriched Amaranth extract (1.6 Kg). Nitrate content of the extract was found to be 18% as determined by ion chromatography. See also extract preparation in Figure 7.

Example 8

Enrichment of nitrates in Amaranth extract by extraction with chloroform.

Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed to make slurry. The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was deactivated by heating the slurry at 90°C for 10 minutes. The extraction was performed with water using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract.

Concentrated water extract was kept at 10°C for 48 hr to crystallize oxalic acid or oxalates. The supernatant was decanted off and again kept at 10°C for 24 hr to crystallize remaining oxalic acid or oxalates. The supernatant from this solution was decanted, filtered and evaporated to dryness under vacuum to get oxalic acid or oxalate free Amaranth extract (2.2 Kg).

This oxalic acid or oxalate free Amaranth extract was further extracted with chloroform by cold extraction. The extract was filtered and evaporated to dryness

under vacuum to get nitrate enriched Amaranth extract (1.6 Kg). See also extract preparation in Fig. 8.

Nitrate content of the extract was found to be 16% as determined by ion chromatography.

5 **Example 9**

Enrichment of nitrates in Amaranth extract by extraction with ethylene dichloride (EDC).

Fresh Amaranth were collected (100 Kg). Leaves and stem of fresh Amaranth were cleaned and crushed to make slurry. The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was deactivated by heating the slurry at 90°C for 10 minutes. The extraction was performed with water using an extractor with a reflux condenser. The bottom of the extractor was fitted with a polypropylene (100 microns) filter cloth. The mixture was refluxed for one hour to obtain a first residue and supernatant. The residue and supernatants were separated by draining out the supernatant from the extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract.

Concentrated water extract was kept at 10°C for 48 hr to crystallize oxalic acid or oxalates. The supernatant was decanted off and again kept at 10°C for 24 hr to crystallize remaining oxalic acid or oxalates. The supernatant from this solution was decanted, filtered and evaporated to dryness under vacuum to get oxalic acid or oxalate free Amaranth extract (2.2 Kg).

This oxalic acid or oxalate free Amaranth extract was further extracted with EDC for 8 hours in a soxhlet extractor. The extract was filtered and evaporated to

dryness under vacuum to get nitrate enriched Amaranth extract (1.6 Kg). See also extract preparation in Fig. 9.

Nitrate content of the extract was found to be 17% as determined by ion chromatography.

5 **Example 10**

Preparation of Amla extract.

Fresh Amla fruits weighing 10 Kg were pulped and the pulp was extracted with methanol at 65°C for 8 hours in a soxhlet apparatus. After the completion of extraction, the supernatant was filtered and concentrated at 60°C to form
10 concentrated methanol extract of Amla. Concentrated methanol extract was dried under vacuum at above 500 mm of mercury to get methanol extract of fresh Amla fruits (Yield: 500 gm). See also extract preparation in Fig. 10.

Example 11

Amaranth extract enriched with Amla extract.

15 1.5 Kg of fresh Amaranth leaves extract (from Example 7) was taken in a beaker and dissolved in methanol (5 L). 150 gm of Amla extract (from Example 10) was added to this solution and stirred for 1 hr at 100 rpm using mechanical stirrer. The obtained mixture was evaporated to dryness under vacuum to get combined extract. See also extract preparation in Fig. 11

20 Nitrate content of this combined extract was 17%.

Example 12

Method of preparation of Amaranth leaf powder .

Fresh Amaranth were collected (100 Kg). Leaves of fresh Amaranth were cleaned and dried. Dried leaves were powdered to get Amaranth leaf powder. See
25 also extract preparation in Fig. 12

Example 13

Method of preparation of Juice from Amaranth leaves.

Fresh Amaranth leaves were collected (2 Kg). Leaves of fresh Amaranth were cleaned. The cleaned leaves were crushed in the screw expeller. The initial juice obtained on the tray of the screw expeller was collected in the feed tank under room temperature of 27°C. The initial juice was centrifuged to obtain filtrate which was collected and the residue which was removed. The collected filtrate was clarified in a rotating disc type clarifier (RPM – 18000) to obtain final filtrate and some residue. After removing the residue, the final filtrate (clarified liquid juice of leaves of Amaranth) was collected with a yield 750 ml juice from 2Kg Amaranth leaves. See also extract preparation in Fig. 13.

10 **Example 14**

Preparation of methanol extract of Amaranth fresh leaves using different varieties of Amaranth.

Fresh leaves and stem of *Amaranthus tricolor* were collected (100 Kg). Leaves and stem of fresh *Amaranthus tricolor* were cleaned and crushed. Crushed materials were filled in the Soxhlet extractor and extracted with methanol (300 L). The extraction was carried out for 5 hrs at a temperature of about 65°C. After the completion of extraction, the supernatant was filtered and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 60°C to form concentrated methanol extract. Concentrated methanol extract was dried under vacuum at above 500 mm of mercury to get powder of methanol extract of fresh *Amaranthus tricolor* (Product 1) (Yield 2.1Kg).

The nitrate content in methanol extract of fresh *Amaranthus tricolor* by HPLC was found to be 1.48%.

Methanol extract of different species of Amaranth were prepared followed by the same procedure as mentioned above.

Different species of *Amaranthus* that were used for the extraction are *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus blitum* and *Amaranthus viridis*. After methanol extraction of different species yields

1. Powder of methanol extract of fresh *Amaranthus caudatus*(Product II)
(Yield 1.95Kg).

The nitrate content in methanol extract of fresh *Amaranthus caudatus* by HPLC was found to be 1.55%.

- 5 2. Powder of methanol extract of fresh *Amaranthus cruentus* (Product III)
(Yield 2 Kg).

The nitrate content in methanol extract of fresh *Amaranthus cruentus* by HPLC was found to be 1.53%.

- 10 3. Powder of methanol extract of fresh *Amaranthus blitum* (Product IV)
(Yield 2.05Kg).

The nitrate content in methanol extract of fresh *Amaranthus blitum* by HPLC was found to be 1.5%.

4. Powder of methanol extract of fresh *Amaranthus viridis* (Product V)
(Yield 2.15Kg).

- 15 The nitrate content in methanol extract of fresh *Amaranthus viridis* by
HPLC was found to be 1.49%.

Example 15

Enrichment of nitrates in Amaranth (different varieties) extract by extraction with hexane.

- 20 Fresh leaves and stem of *Amaranthus tricolor* were collected (100 Kg). Leaves
and stem of fresh *Amaranthus tricolor* were cleaned and crushed to make slurry.
The slurry was treated with pectinase (1%) for 2 Hrs. The enzyme was
deactivated by heating the slurry at 90°C for 10 minutes. The extraction was
performed with water using an extractor with a reflux condenser. The bottom of
25 the extractor was fitted with a polypropylene (100 microns) filter cloth. The
mixture was refluxed for one hour to obtain a first residue and supernatant. The
residue and supernatants were separated by draining out the supernatant from the

extractor bottom through the polypropylene filter cloth using a centrifugal pump. After the first extraction, the first residue was further extracted with ten times the quantity of water to get second residue and supernatant. The second residue was further extracted with ten times the quantity of water to get third residue and supernatant. All the supernatants were pooled and concentrated in an Agitated thin film evaporator (ATFE) at a temperature of 85°C to form concentrated water extract.

Concentrated water extract was kept at 10°C for 48 hr to crystallize oxalic acid or oxalates. The supernatant was decanted off and again kept at 10°C for 24 hr to crystallize remaining oxalic acid or oxalates. The supernatant from this solution was decanted, filtered and evaporated to dryness under vacuum to get oxalic acid or oxalate free extract of *Amaranthus tricolor* (2.22 Kg).

This oxalic acid or oxalate free extract of *Amaranthus tricolor* was further extracted with hexane for 6 hours in a soxhlet extractor. The extract was filtered and evaporated to dryness under vacuum to get nitrate enriched extract of *Amaranthus tricolor* (Product I) (1.63 Kg).

Nitrate content of the extract was found to be 17.7% as determined by HPLC.

Nitrate enriched extract of different species of *Amaranthus* were prepared followed by the same procedure as mentioned above.

Different species of *Amaranthus* that were used for the extraction are *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus blitum* and *Amaranthus viridis*. After extraction of different species yields

1. Powder of nitrate enriched extract of fresh *Amaranthus caudatus* (Product II) (Yield 1.62Kg)

The nitrate content in nitrate enriched extract of fresh *Amaranthus caudatus* by HPLC was found to be 17.9%.

2. Powder of nitrate enriched extract of fresh *Amaranthus cruentus* (Product III) (Yield 1.6 Kg)

The nitrate content in nitrate enriched extract of fresh *Amaranthus cruentus* by HPLC was found to be 18%.

3. Powder of nitrate enriched extract of fresh *Amaranthus blitum* (Product IV) (Yield 1.58Kg)

5 The nitrate content in nitrate enriched extract of fresh *Amaranthus blitum* by HPLC was found to be 18.3%.

4. Powder of nitrate enriched extract of fresh *Amaranthus viridis* (Product V) (Yield 1.61Kg)

10 The nitrate content in nitrate enriched extract of fresh *Amaranthus viridis* by HPLC was found to be 18.1%.

Example 16

Determination of nitrate and nitrite content in Amaranth by ion chromatography.

15 A 50 microliter aliquot of sample was pipette into a 300 microliter glass tube (Sci-Vi, Chromacol, Welwyn, UK). Acetonitrile (50 microliter) was added, and the tube with an autosampler limited-volume insert spring was put in a 4-ml WISP vial, capped, mixed and centrifuged at 2000 g for 2 min. The supernatant was injected for analysis.

20 882 Compact IC plus chromatograph (Metrohm, USA) fitted with a WISP autosampler (Waters, Watford, UK). The column was a Metrosep A Supp 5 – 150/4.0 ion-exchange column, 150 mm x 4 mm (Metrohm), and the eluent was 1.7 mM Sodium bicarbonate (NaHCO_3), 1.8 mM Sodium carbonate (Na_2CO_3), pumped at 1.5 ml/ min. Detection was by conductivity (DX100) and absorbance at 214 nm using a 441 detector fitted with a zinc lamp (Waters). Data was acquired and processed using an 840 data system (Waters).

25 Example 17

Determination of nitrate and nitrite by HPLC method.

Standard for nitrate (NO₃) was prepared by weighing 81.53 mg potassium nitrate (99% pure) in to 100 ml standard flask and made up to mark with water. 1 ml from the 100 ml solution was pipetted out into 50 ml standard flask and made up to mark with water.

- 5 Standard for nitrite (NO₂) was prepared by weighing 92.49 mg potassium nitrite (96% pure) in to 100 ml standard flask and made up to mark with water. 1 ml from the 100 ml solution was pipetted out into 50 ml standard flask and made up to mark with water. Again 1ml from the 50 ml solution was pipetted out into 50 ml standard flask and again made up to mark with water.
- 10 Added 1ml standard of nitrite and 1ml mobile phase A (Weighed 1.3 g of the Tetra Butyl ammonium Hydroxide Solution (10%aqu) in to 1000ml standard flask. Made up to mark with Water .Then PH adjusted to 2.50 with Con: H₂SO₄), 200 µ Griess A (Weighed 50.00mg sulphanilic acid and 1.5 ml acetic acid and 3 ml purified water was added and dissolved) and after 2 min added 200 µ Griess B
- 15 (Weighed 4.00mg 1-naphthylamine and 4.0 ml acetic acid was added and dissolved) in to a plastic tube.Filtered through 0.45µm membrane filter before injection.

Sample (Amaranth extract) was prepared by weighing accurately 20-25 mg of the extract powder to 25 ml std flask and made up the with demineralized water.

- 20 Filtered through 0.45 Micron membrane filter and divided into 2 portions. One portion was used for Nitrate and other portion was used for Nitrite determinations.

- NO₃ was analyzed by HPLC on a BETASIL C-18 column (250 x 4.6mm) using Tetra butyl ammonium hydroxide in De-ionized water, acetonitrile and methanol
- 25 as mobile phase and Photo diode array detection at 222 nm. NO₂ was analysed at 520 nm using the same conditions.

Nitrite% = $\frac{\text{Area of Nitrite peak in sample} \times \text{Amount of Std} \times \text{Purity of Std}}{\text{Area of Nitrite peak in Standard} \times \text{Weight of the Sample}}$

Area of Nitrite peak in Standard X Weight of the Sample

$$\text{Nitrate \%} = \frac{\text{Area of Nitrate peak in sample} \times \text{Amount of Std} \times \text{Purity of Std}}{\text{Area of Nitrate peak in Standard} \times \text{Weight of the Sample}}$$

Example 18

5 Determination of nitric oxide(NO).

Total NO content in plasma can be interpreted as the sum of total nitrate(NO₃) and nitrite(NO₂) in plasma.

(Journal of Medical sciences (2010):3(3): 153-159, Plasma Endothelin -1, Homocysteine, and Oxide Levels in a Multiethnic Hypertensive Cohort from the
10 United Arab Emirates.)

NO levels was recorded as the sum of the two major metabolites of NO, nitrite and nitrate.

Example 19

Determination of total oxalic acid content.

15 Oxalic acid was estimated by high performance liquid chromatography (HPLC-DAD) on a C18 column (250x 4.6 mm, Gemini 5µm, USA.). The mobile phase was 5mM H₂SO₄ and used under isocrating condition with an eluent flow rate of 1 ml/min. Oxalic acid was detected at 210 nm.

Standard was prepared by weighing 5 mg of standard oxalic acid (95% purity)
20 and was made up to 50 ml with 6mM H₂SO₄. Sample was prepared by weighing 50 mg of the dry extract of Amaranth and was made up to 50 ml with 6mM H₂SO₄. Both the sample and standard were filtered separately through a 0.2 µm membrane filter before injection into the HPLC column. The injection volume was 20µl. Oxalic acid was detected at 210 nm. By comparing the area of
25 standard and sample, the percentage of oxalic acid present in the sample was quantified.

Oxalic acid % = $\frac{\text{Area of sample} \times \text{amount of standard}}{\text{Amount of sample} \times \text{Area of standard}}$ x purity of standard

Amount of sample x Area of standard

Example 20

Determination of potassium content.

- 5 Potassium was determined by Atomic absorption spectroscopy (AAS). A 100 ppm solution of potassium Standard (KCl) was prepared. Seven different standards of potassium in different concentrations were prepared by simple dilution of the above 100 ppm solution. The sample solution and standard were measured in the AAS machine at 766.5nm to create a calibration curve for
10 determination of potassium content in the sample solutions.

Example 21

Preparation of fast dissolving tablets.

- Fast dissolving tablets were made by mixing Fast-Melt (119.2 gm) with amaranth nitrate enriched (18% nitrate) extract (80 gm) (as per Example 7). Magnesium
15 stearate (0.8 gm) was added to the mixture before tablet punching. Tablets weighing 200 mg were punched on automatic tablet punch machine using 8 mm die to get 5 Kg hardness. The final composition of each tablet was 119.2 mg F-Melt, 80 mg extract (approx. 14.4mg nitrate) and 0.8 mg of magnesium stearate. Disintegration time of tablets was approximately 30 seconds.

20 Example 22

Preparation of fast dissolving tablets.

- Fast dissolving tablets were made by mixing Fast-Melt (745 gm) with amaranth nitrate enriched (18% nitrate) extract (500 gm) prepared as per example 7. Magnesium stearate (5 gm) was added to the mixture before tablet punching.
25 Tablets weighing 1250 mg were punched on automatic tablet punch machine using 12 mm die to get 5 Kg hardness. The final composition of each tablet was 745 mg F-Melt, 500 mg extract (approx.90 mg nitrate) and 5 mg of magnesium stearate. Disintegration time of tablets was approximately 30 seconds.

Similarly fast melt tablets were made by substitute the above extract with

1. Methanol extract of fresh leaves of Amaranth (1.5% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 7.5 mg nitrate) and 5 mg of magnesium stearate)
- 5 2. Water extract of fresh leaves of Amaranth (1.6% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 8 mg nitrate) and 5 mg of magnesium stearate)
3. Methanol extract of dried leaves of Amaranth (1.65% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 8.25 mg nitrate) and 5 mg of magnesium stearate)
- 10 4. Water extract of dried leaves of Amaranth (1.7% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 8.5 mg nitrate) and 5 mg of magnesium stearate)
5. Pectinase treated water extract of fresh leaves of Amaranth (2% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 10 mg nitrate) and 5 mg of magnesium stearate)
- 15 6. Extract of *Amaranthus blitum* with enriched nitrate (18.3% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 91.5mg nitrate) and 5 mg of magnesium stearate)
- 20 7. Extract of *Amaranthus viridis* with enriched nitrate (18.1% nitrate) (Final composition of Fast melt tablet contain 750 mg F-Melt, 500 mg extract (approx 90.5mg nitrate) and 5 mg of magnesium stearate)

Example 23

Preparation of chewing gum.

- 25 Chewing gums were prepared by adding amaranth nitrate enriched extract (80 mg for one chewing gum) to the base material (920 mg for one chewing gum). The base material consists of polyvinyl acetate in an amount by weight of about 38%, natural rubber in an amount by weight of about 10%, partially hydrogenated

soybean oil in an amount by weight of about 11%, polyisobutylene in an amount by weight of about 12%, dicalcium phosphate in an amount by weight of about 13.92%, triacetin in an amount by weight of about 3%, mono-di-glycerides in an amount by weight of about 7%, liquid glucose in an amount by weight of about 5%, and BHT (butylated hydroxyl toluene) in an amount by weight of about 0.08%.

The chewing gum components (except liquid glucose and BHT) were mixed in a mixer with strong horizontally placed Z-blades, which processes the components into a homogeneous gum mass. The mixer was heated to a temperature of approx. 50-60°C. The mixing process starts with mixing and melting the gum base for 3 minutes.

The liquid glucose was then added and mixed for 4 minutes. Subsequent, the amaranth nitrate enriched extract were added and the mixture was mixed for further 6 minutes. The BHT was then added and mixed for further 5 minutes.

When the mixing was completed, the gum mass was taken out onto trays and rolled out to a 1-2 cm thick sheet. The chewing gum mass was subsequent cooled for approx. 15-20 minutes until the right texture of the gum mass was achieved.

The formed and cooled chewing gum cores were then coated and polished with a coating suspension in a round stainless steel coating kettle that rotate during the coating and polishing process. The coating suspension consists of Xylitol (58.2%), Mannitol (10.6%), Gelatin (1.2%), Titanium dioxide (0.9%), Acesulfame K (0.1%) and water (29%).

The polishing was done in rotating kettles in which carnauba wax (0.1%) was added to the coated chewing gums in one portion. Polishing was done until a shining surface is achieved, typically for 10-20 minutes. Each chewing gum contained 80 mg of extract (approx. 14.4 mg nitrate).

Example 24

Nitric oxide level in rats.

18 albino rats (Sprague-Dawley (SD) strain) weighing 200-250 gm were taken for the study. The animals were acclimatized to the animal house condition ($24\pm 2^{\circ}\text{C}$ temperature, 65% relative humidity, 12 hr light/dark cycle) for one week. The animals were divided into three groups comprising of six animals in each group. The animals were fasted overnight and blood was collected to determine the baseline nitric oxide level. The animals were treated as follows:

Group 1: Vehicle control

Group 2: Amaranth dried leaves powder – 1gm/kg

Group 3: Amaranth nitrate enriched extract (from example 7) – 100 mg/kg

After 30 minutes of feeding, blood was collected from all the animals in EDTA (ethylenediaminetetraacetic acid) coated tubes and plasma was separated immediately by centrifuging the tubes at 4000 rpm (rotations per minute) for 10 minutes. The plasma was stored below -80°C until analysis.

The nitric oxide (NO) content of the plasma was reported as $\mu\text{mol/L}$.

15

S. No.	Groups	Baseline Nitric oxide level ($\mu\text{mol/L}$)	Nitric oxide level after 30 min ($\mu\text{mol/L}$)
1.	Vehicle control	18.2	18.4
2.	Amaranth dried leaves powder	17.9	18.2
3.	Amaranth nitrate enriched extract (from example 7)	18.6	42.3

Example 25

Nitric oxide level in mice.

18 swiss albino mice weighing 25-30 gm were taken for the study. The animals were acclimatized to the animal house condition ($24\pm 2^{\circ}\text{C}$ temperature, 65% relative humidity, 12 hr light/dark cycle) for one week. The animals were divided into three groups comprising of six animals in each group. The animals were

fasted overnight and blood was collected to determine the baseline nitric oxide level. The animals were treated as follows:

Group 1: Vehicle control

Group 2: Amaranth dried leaves powder (from example 12) – 1gm/kg

- 5 Group 3: Amaranth extract enriched with Amla extract (from example 11) – 200 mg/kg

After 30 minutes of feeding, blood was collected from all the animals in EDTA (ethylenediaminetetraaceticacid) coated tubes and plasma was separated immediately by centrifuging the tubes at 4000 rpm for 10 minutes. The plasma
10 was stored below -80°C until analysis.

The nitric oxide (NO) content of the plasma is reported as $\mu\text{mol/L}$.

S. No.	Groups	Baseline Nitric oxide level ($\mu\text{mol/L}$)	Nitric oxide level after 30 min ($\mu\text{mol/L}$)
1.	Vehicle control	21.9	21.7
2.	Amaranth dried leaves powder (from example 12)	20.8	22.6
3.	Amaranth extract enriched with Amla extract (from example 11)	21.6	76.4

Example 26

Oral toxicity study in rats.

- 15 This study was conducted as per OECD Guidelines (Organisation for economic co-operation and development) for Testing of chemicals (No.-408). 100 Wistar albino rats (50 males/50 females) were divided into four groups of 20 animals (10 males and 10 Females) in each and two groups (Satellite groups) of 10 animals (5 males and 5 Females) in each group. The animals were acclimatized for seven
20 days before the commencement of dosing. Three groups of 20 rats each (10 male and 10 female) were administered with amaranth nitrate enriched extract (from

example 8) at the dosage levels of 100 mg/kg body weight (low dose), 500 mg/kg b.wt (body weight) (intermediate dose) and 1000 mg/kg body weight (high dose) respectively for seven days a week for 90 days with the help of cannula attached with the syringe. Similarly, fourth group of 20 rats (10 male and 10 female rats) were orally administered with corn oil only (vehicle) for 90 days and was designated as control group.

Two additional satellite groups of 10 rats (5 male and 5 female) each were also kept and designated as 'Satellite control' and 'Satellite high dose' and were administered with corn oil (vehicle) and amaranth nitrate enriched extract (1000 mg/kg) respectively for 90 days. After terminal sacrifice of the test and control group animals, both Satellite group animals (Satellite control and Satellite high dose) were kept under observation for an additional 28 days, to check the reversibility, persistence or delayed toxic effect, if any. The animals were observed daily for behavior, appearance and toxicological signs and symptoms. Blood was collected from all the animals before terminal sacrifice for detailed hematological and biochemical evaluation. Urine samples were also collected from all animals at the termination of the experiment. Criteria used to evaluate compound related effects included; appearance, behavior, morbidity, mortality, body weights, Feed consumption, hematological and biochemical analysis, urine analysis, organ weights, necropsy and histopathology.

No treatment related toxic sign and symptoms were observed in low dose (100 mg/kg body weight), intermediate dose (500 mg/kg body weight), high dose (1000 mg/kg body weight.) and satellite high dose (1000 mg/kg body weight) group animals when compared to their respective control counter parts. Body weights of all the test and control group animals were recorded weekly. Body weight gain of all the treatment groups and satellite group animals (LD, ID, HD & satellite HD) was comparable to their control counterparts. Feed consumption of the animals of low dose, intermediate dose, high dose and satellite high dose were comparable to the control group and satellite control group animals.

There were no variations in the haematological parameters of animals of low dose (100 mg/kg Body weight), intermediate dose (500 mg/kg body weight), high dose (1000 mg/kg body weight) and satellite high dose (1000 mg/kg body weight) groups when compared to the control group animals. Similarly, the
5 biochemical parameters of animals of all the treatment groups i.e. low dose, intermediate dose and high dose were comparable to the biochemical parameters of the control group animals at the terminal sacrifice. The biochemical parameters of satellite high dose group were also comparable to its satellite control counterparts. The haematological as well as biochemical parameters of
10 test groups were not significantly different from respective control groups.

Urine samples were collected from all animals at the termination of the experiment. No significant changes were noted in the urine parameters of any of the dose group animals when compared to the control group. Animals of satellite groups were sacrificed after 28 days post treatment. None of the animals died
15 during the study in any of the treated groups as well as the satellite groups. After completion of dosing period of 90 days, all the groups (treatment and control) except both the satellite groups were sacrificed and were examined for gross pathological findings. Organs of all the animals (treatment and control) were trimmed of any adherent fat tissue and their weights taken. Organ weights of
20 animals of all treatment groups were comparable to their respective control counterparts. There were no significant histopathological changes in the animals of low dose group, intermediate dose group, high dose group and satellite high dose group when compared to their control counterparts.

In this study, the repeated administration of amaranth nitrate enriched extract for
25 90 days, by oral route, to wistar rats at the dosage level of 1000 mg/kg body weight did not induce any observable toxic effects, when compared to its corresponding control group of animals.

Example 27

Nitric oxide level in human subjects.

18 healthy human volunteers (age 35-55 years) were taken for the study. The subjects were put on standard diet (devoid of nitrate/nitrite containing food) one week prior to the study. The subjects were divided into three groups comprising of six subjects in each group. The subjects were fasted overnight and blood was collected to determine the baseline nitric oxide level. After blood collection, the subjects were treated as follows:

Group 1: Vehicle control

Group 2: Amaranth fresh leaves as such – 100 gm

Group 3: Amaranth fast dissolving tablet containing amaranth nitrate enriched extract (14.4 mg nitrate per tablet prepared as per example 21 and extract used for making the tablet prepared as per example 7)

After 30 minutes of feeding, blood was collected from all the subjects in EDTA (ethylene di amine tetra acetic acid) coated tubes and plasma was separated immediately by centrifuging the tubes at 4000 rpm for 10 minutes. The plasma was stored below -80°C until analysis for nitric oxide content. The nitric oxide (NO) content of the plasma is reported as $\mu\text{mol/L}$.

S. No.	Groups	Baseline Nitric oxide level ($\mu\text{mol/L}$)	Nitric oxide level after 30 min ($\mu\text{mol/L}$)
1.	Vehicle control	28.5	28.2
2.	Amaranth fresh leaves as such	29.2	30.1
3.	Amaranth fast dissolving tablet(Fast melt prepared as per Example 21 and extract used for making the tablet prepared as per example 7)	29.1	78.9

Example 28

Anti-hypertensive activity in human subjects.

10 Subjects (6 males and 4 females) aged 40 to 60 years with pre-hypertension [SBP (systolic blood pressure), 130 to 140 mmHg and DBP (diastolic blood pressure), 85 to 90 mmHg] or hypertension (SBP (systolic blood pressure), 140 mmHg or higher and DBP (diastolic blood pressure), 90 mmHg or higher) were enrolled in the study. Exclusion criteria included pregnant and lactating women, people who had received trace element supplements in the previous three months, people receiving gastric or diuretic treatments, patients with acute renal failure, or people with recent history of surgery or acute infections. All subjects were informed of the purposes of the study, were free to ask questions throughout the study, and signed an informed consent form witnessed by one of the investigators.

S. No.	Subjects	Baseline blood pressure (SBP/DBP in mmHg)	Blood pressure after 15 days (SBP/DBP in mmHg)	Blood pressure after 30 days (SBP/DBP in mmHg)
1.	Subject 1	138/87	135/84	130/82
2.	Subject 2	142/90	138/88	133/84
3.	Subject 3	140/89	137/87	131/83
4.	Subject 4	139/88	136/86	132/83
5.	Subject 5	143/91	138/88	134/85
6.	Subject 6	141/90	136/87	132/84
7.	Subject 7	139/90	135/83	129/81
8.	Subject 8	138/86	134/83	128/80
9.	Subject 9	142/90	137/87	131/84
10.	Subject 10	142/90	136/86	132/83

The patients were provided with a bottle containing 60 fast melt mouth dissolving tablets (from Example 21) and instructed to take one tablet twice daily at 12 hour interval for 30 days. Eating habits were not restricted during the treatment.

5 However, to avoid possible interference with the effects of PG, the consumption of vitamin supplements and functional foods was prohibited during the trial. Each subject's BP (blood pressure) was measured at the beginning, 15 days and end of the trial. Their BP (blood pressure) was measured using a digital BP analyzer and recorded as the average of two measurements. Measurements were always made

10 at the same place and time after 15 min of rest.

Example 29

Swimming endurance test in mice.

For swimming endurance study, prior to experiment, each mouse was tested for feasibility testing. All mice in swimming groups could swim. The mice were

15 divided into 3 groups comprising of 6 animals in each group. Following treatment was given to the respective group of mice:

Group 1: Vehicle control

Group 2: Amaranth dried leaves powder (from example 12)– 1gm/kg

Group 3: Amaranth nitrate enriched extract (from example 8) – 100 mg/kg

20 Treatment was given for four weeks.

All mice were subjected to swim in a water pool (48x28x24 inch), filled 19 inch deep of water maintained at 25°C. During swimming testing, each mouse was fixed four paper clips with a plastic string on the tail. The paper clips and plastic string weighted about 1.81 grams. Bearing this weight, all mice were forced to

25 swim very hard. All mice were subjected to swim until exhaustion (cease movement of limbs and float). The maximum time swimming until mice exhausted was used as the index of swimming endurance capacity. The test was conducted in the beginning and then weekly till four weeks. Blood was also

collected in the beginning and after each week to determine the nitric oxide level in plasma.

S. No.	Groups	Swimming endurance (seconds)				
		Baseline	1 Week	2 Weeks	3 Weeks	4 Weeks
1.	Vehicle control	130	126	128	130	127
2.	Amaranth dried leaves powder (from example 12)	132	146	168	172	179
3.	Amaranth nitrate enriched extract (from example 8)	128	155	171	185	193

Nitric oxide level

S. No.	Groups	Nitric oxide level ($\mu\text{mol/L}$)				
		Baseline	1 Week	2 Weeks	3 Weeks	4 Weeks
1.	Vehicle control	17.6	17.3	18.2	18.0	17.9
2.	Amaranth dried leaves powder (from example 12)	17.9	18.1	18.3	17.7	17.9
3.	Amaranth nitrate enriched extract (from example 8)	17.7	32.1	46.5	51.3	56.9

5

Example 30

Running endurance test in mice.

For running endurance study, individual mouse was trained for running on a Rota-Rod apparatus (treadmill) for three days prior to running study. During training, the speed of the Rota-rod was increased incrementally to make animals exercise more intensely and running for at least one minute. The mice were divided into 3 groups comprising of 6 animals in each group. Following treatment was given to the respective group of mice:

Group 1: Vehicle control

Group 2: Amaranth dried leaves powder (from example 12) – 1gm/kg

Group 3: Amaranth nitrate enriched extract (from example 9) – 100 mg/kg

Treatment was given for four weeks.

The Rota-Rod treadmills had five testing zones, each zone had a photo bean that sensed when the animal dropped off the treadmills which was connected with computer to calculate running time on the treadmills automatically. The speed of the Rota-rod was set to 35 RPM (rotation per minute). The maximum time running until exhaustion (drops from the Rota-rod) was used as the index of running endurance capacity. The maximum time of running until exhaustion in each mouse was recorded. The test was conducted in the beginning and then weekly till four weeks.

S. No	Groups	Running endurance (seconds)				
		Baseline	1 Week	2 Weeks	3 Weeks	4 Weeks
1.	Vehicle control	452	445	454	450	448
2.	Amaranth dried leaves powder (from example 12)	436	451	472	481	486
3.	Amaranth nitrate enriched extract (from example 9)	441	468	489	497	510

Example 31

Sexual performance study in rats.

Twelve male wistar rats (200-250 gm) were randomized into 2 groups comprising of 6 animals each. The animals were treated for 30 days with vehicle/extract and
5 various test parameters were evaluated.

Group I (vehicle only) served as control.

Group II Amaranth extract enriched with amla extract (from example 11) - 100 mg/Kg (p.o.) daily

Orientation behavior analysis

10 The analysis of orientation activity was carried out and analyzed in three segments. Orientation behavior of male rats was determined using following method of scoring: Orientation towards female – (1 for every sniffing and 2 for every licking)

Orientation towards self – (1 for non-genital grooming and 2 for genital
15 grooming)

Orientation towards environment – (1 for exploration, 2 for *rearing* and 3 for climbing)

Rats were observed daily for their orientation activity. The cumulative score after 0, 15, 30 days of the treatment were recorded.

20 Sexual behavior analysis

Male rat was placed in the observation glass chambers in order to acclimatize it with the cage environment. Sexually receptive female rat was then allowed to enter the test cage silently from a side door inside the cage. The behavioral observations were carried out taking into account the following parameters.

25 Mounting Behavior – It was determined and characterized by following parameters.

(A) Mount frequency – average number of mount during 30 min observation.

(B) Mount latency- The leg time from the introduction of female in the cage to first mount.

Intromission Behavior – It was evaluated as follows.

(A) Intromission frequency – average number of Intromission during 30 min observation.

(B) Intromission latency- Intromission latency (IL) was considered as the time for first intromission after introduction of female in the cage.

Penile Erection Index

Penile Erection (PE) was determined when the rats bent down to lick their erect penis during the observation period. *Penile erection index (PI)* was determined by multiplying the percentage of rats exhibiting at least one episode of penile erection during 30-min observation period with the mean number of penile erections.

PI = % of rats exhibiting penile erection X Mean number of erections.

15

Table: Effect of enriched amaranth extract on orientation activities in male rats

Parameters for sexual behavior analysis	Groups	0 days	15 days of treatment	30 days of treatment
Mean activity Score Towards Female (Licking & Anogenital smelling)	Vehicle control	7.5	7.6	7.5
	Amaranth extract enriched with Amla extract	7.4	11.2	16.3
Mean activity Score Towards Environment (Exploration, Rearing and Climbing)	Vehicle control	13.2	13.3	13.6
	Amaranth extract enriched with Amla extract	13.3	16.5	22.2

Mean activity Score Towards Self (Nongenital grooming and Genital grooming)	Vehicle control	8.1	8.0	8.1
	Amaranth extract enriched with Amla extract	7.9	12.6	15.3

Table: Effect of enriched amaranth extract on sexual performance parameters in male rats

Parameters for sexual behavior analysis	Groups	0 days	15 days of treatment	30 days of treatment
Mount latency (time in seconds)	Vehicle control	194.3	195.6	195.2
	Amaranth extract enriched with Amla extract	193.5	160.2	124.3
Mount frequency	Vehicle control	3.3	3.5	3.8
	Amaranth extract enriched with Amla extract	3.4	10.5	12.4
Intromission latency (time in seconds)	Vehicle control	305.2	304.3	302.6
	Amaranth extract enriched with Amla extract	309.5	275.3	254.4
Intromission frequency	Vehicle control	1.6	1.7	1.7
	Amaranth extract enriched with Amla extract	1.5	4.2	6.3
Penile Erection Index (PI)	Vehicle control	23.3	22.7	24.5
	Amaranth extract enriched with Amla extract	21.5	45.3	62.6

Example 32

Nitrate (NO_3) Nitrite (NO_2) and nitric oxide (NO) content in plasma of human subjects.

32 Healthy human volunteers (age 35-55 years) were taken for the study. The subjects were put on standard diet one week prior to the study. The subjects were divided into 8 groups comprising of four subjects in each group. The subjects were fasted overnight and the blood was collected to determine the baseline NO_3 and NO_2 levels. After baseline blood collection, the subjects were treated as follows:

10 Group 1: Vehicle Control.

Group 2: Amaranth leaves fresh juice as such -100 ml (100 mg Nitrate /100ml of juice) (Prepared as per example No-13)

Group 3: Amaranth fast dissolving tablet containing Methanol extract of Amaranth fresh leaves (7.5mg nitrate per tablet) (Tablet prepared as per example No.22 and extract used for preparing the tablet is as per example 1)

Group 4: Amaranth fast dissolving tablet containing Methanol extract of Amaranth dried leaves (8.25mg nitrate per tablet) (Tablet prepared as per example No.22 and extract used for preparing the tablet is as per example 3)

Group 5: Amaranth fast dissolving tablet containing water extract of Amaranth fresh leaves (8 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 2)

Group 6: Amaranth fast dissolving tablet containing water extract of Amaranth dried leaves (8.5mg nitrate per tablet) (Tablet prepared as per example No 22 and extract used for preparing the tablet is as per example 4)

25 Group 7: Amaranth fast dissolving tablet containing Pectinase treated water extract of Amaranth fresh leaves (10 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 5)

Group 8: Amaranth fast dissolving tablet containing amaranth nitrate enriched extract (90 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 7)

5 All subjects were allowed to consume the water/Juice orally(Group No-1 to 2).Amaranth Tablet were kept in oral cavity and allow to dissolve (Group No-3 to 8).

10 After 1 hour, 2 hour and 3 hour from feeding of tablet/juice/vehicle, blood was drawn from all the subjects in EDTA coated tubes. Plasma was separated immediately by centrifuging the tubes at 5000 rpm for 15 min. The plasma samples were stored below -80 degree Celsius until analysis. The plasma thus obtained was deproteinized using Acetonitrile and the supernatant liquid was used for nitrate/nitrite determination using HPLC. NO₃ was analyzed by HPLC on a BETASIL C-18 column (250 x 4.6mm) using Tetra butyl ammonium hydroxide in De-ionized water, acetonitrile and methanol as mobile phase and 15 Photo diode array detection at 222 nm. NO₂ was analysed at 520 nm using the same conditions. The NO₃ and NO₂ levels in plasma were reported as µmol/L .

20

25

Table: Nitrate (NO₃) content in plasma

Sl NO	Groups	Baseline NO ₃ levels (μmol/l)	NO ₃ levels after 1 hour (μmol/l)	NO ₃ levels after 2 hour (μmol/l)	NO ₃ levels after 3 hour (μmol/l)
1.	Vehicle Control	28.92	28.95	27.88	28.81
2.	Amaranth fresh juice as such	28.55	30.11	29.45	28.56
3.	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	27.88	30.12	29.88	28.65
4	Amaranth fast dissolving tablet containing water extract of fresh leaves	26.55	30.19	30.02	28.89
5	Amaranth fast dissolving tablet containing methanol extract of dried leaves .	29.52	33.20	30.25	29.56
6	Amaranth fast dissolving tablet containing water extract of dried leaves	29.10	32.8	31.02	29.77
7	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	29.11	32.92	31.95	30.10
8	Amaranth fast dissolving tablet containing enriched extract of nitrate	28.88	174.40	110.22	70.89

Table: Nitrite (NO₂) content in plasma

Sl No	Groups	Base line NO ₂ levels (μmol/l)	NO ₂ levels after 1 hour (μmol/l)	NO ₂ levels after 2 hour (μmol/l)	NO ₂ levels after 3 hour (μmol/l)
1.	Vehicle Control	0.19	0.20	0.23	0.18
2.	Amaranth fresh juice as such	0.21	0.25	0.24	0.20
3.	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	0.20	0.27	0.26	0.24
4	Amaranth fast dissolving tablet containing water extract of fresh leaves	0.21	0.29	0.27	0.25
5	Amaranth fast dissolving tablet containing methanol extract of dried leaves .	0.19	0.28	0.27	0.24
6	Amaranth fast dissolving tablet containing water extract of dried leaves	0.21	0.31	0.35	0.30
7	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	0.19	0.30	0.28	0.25
8	Amaranth fast dissolving tablet containing enriched extract of nitrate	0.20	1.20	0.98	0.60

Table: Total nitric oxide (NO) content in plasma

Sl NO	Groups	Base line NO levels ($\mu\text{mol/l}$)	NO levels after 1 hour ($\mu\text{mol/l}$)	NO levels after 2 hour ($\mu\text{mol/l}$)	NO levels after 3 hour ($\mu\text{mol/l}$)
1.	Vehicle Control	29.11	29.15	28.11	28.99
2.	Amaranth fresh juice as such	28.76	30.36	29.69	28.76
3.	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	28.08	30.39	30.14	28.89
4.	Amaranth fast dissolving tablet containing water extract of fresh leaves	26.76	30.48	30.29	29.14
5.	Amaranth fast dissolving tablet containing methanol extract of dried leaves .	29.71	33.48	30.52	29.80
6.	Amaranth fast dissolving tablet containing water extract of dried leaves	29.31	33.11	31.37	30.07
7.	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	29.30	33.22	32.23	30.35
8.	Amaranth fast dissolving tablet containing enriched extract of nitrate	29.08	175.60	111.2	71.49

Amaranth fresh juice consumption as such slightly increased the NO_3 content after one hour. Methanol extract from amaranth fresh leaves and dried leaves also

showed a slight increase in NO_3 content. Tablet made with pectinase treated water extract of fresh leaves of Amaranth shown a little more increase than the other tablets consumptions. But after consumption of fast dissolving tablet containing Amaranth extract enriched with nitrate, NO_3 level increased to 6 times the baseline value after 1 hour and the increase from base line is maintained up to 3 hour. There was no significant change in NO_3 level after administering the vehicle.

In case of NO_2 also, the fast dissolving tablet containing Amaranth extract enriched with nitrate, increased the NO_2 level more than 6 times and the higher levels maintained for 3 hours. Amaranth fresh juice consumption as such slightly increased the NO_2 content after one hour. Methanol extract from amaranth fresh leaves and dried leaves also showed a slight increase in NO_2 content. Here also vehicle did not influence NO_2 level in the subjects .

Overall conclusion is that the increased levels of NO_3 and NO_2 increase observed in Amaranth fast dissolving tablet was more when compared with levels of nitrate and nitrite in Amaranth fresh juice, and methanol extract from fresh of dried leaves. The increase in total nitric oxide(NO) content of "Amaranth fast dissolving tablet containing enriched extract of nitrate" was 6 times from base line and it remained increased for 3 hours. Total NO is determined from the sum of total NO_3 and NO_2 . (ie. Level of NO_3 and NO_2 can be interpreted as a measure of NO level in human body) So an increase in NO_3 and NO_2 level is interpreted as an increase in NO level. NO levels was recorded as the sum of the two major metabolites of NO , nitrite and nitrate.

Example 33

Nitrate (NO_3), Nitrite (NO_2) and nitric oxide(NO) content in saliva of human subjects.

32 Healthy human volunteers (age 35-55 years) were taken for the study. The subjects were put on controlled diet, one week prior to the study. The subjects

were divided into eight groups comprising of four subjects in each group. The subjects were fasted overnight and saliva was collected to determine the base line NO_3 and NO_2 levels. After saliva collection, the subjects were treated as follows:

Group 1: Vehicle Control.

- 5 Group 2: Amaranth leaves fresh juice as such -100 ml (100 mg Nitrate /100ml of juice) (Prepared as per example No.13).

Group 3: Amaranth fast dissolving tablet containing Methanol extract of Amaranth fresh leaves(7.5mg nitrate per tablet) (Tablet prepared as per example No 22 and extract used for preparing the tablet is as per example 1).

- 10 Group 4: Amaranth fast dissolving tablet containing Methanol extract of Amaranth dried leaves (8.25mg nitrate per tablet) (Tablet prepared as per example No.22 and extract used for preparing the tablet is as per example 3).

- 15 Group 5: Amaranth fast dissolving tablet containing water extract of Amaranth fresh leaves (8mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 2).

Group 6: Amaranth fast dissolving tablet containing water extract of Amaranth dried leaves (8.5mg nitrate per tablet) (Tablet prepared as per example No 22 and extract used for preparing the tablet is as per example 4)

- 20 Group 7: Amaranth fast dissolving tablet containing Pectinase treated water extract of Amaranth fresh leaves (10 mg nitrate per tablet) (Tablet prepared as per example No 22 and extract used for preparing the tablet is as per example 5)

Group 8: Amaranth fast dissolving tablet containing amaranth nitrate enriched extract (90 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 7)

All subjects were allowed to consume the water/Juice orally (Group No-1 to 2). Amaranth Tablet were kept in oral cavity and allow to dissolve (Group No-3 to 8)

After each of the time point 1 hour, 2 hour and 3 hour from feeding, saliva was collected from all subjects in 5.0ml PTFE tubes. This saliva was stored below -80 degree Celsius until analysis for NO_3 and NO_2 content. The saliva thus obtained was deprotenized using Acetonitrile and the supernatant liquid was used for NO_3 and NO_2 determination using HPLC. NO_3 was analyzed by HPLC on a BETASIL C-18 column (250 x 4.6mm) using Tetra butyl ammonium hydroxide in De-ionized water, acetonitrile and methanol as mobile phase and Photo diode array detection at 222 nm. NO_2 was analyzed at 520 nm using the same conditions. The Nitrate and Nitrite of the saliva are reported as $\mu\text{mol/L}$.

15

20

Table: Nitrate (NO₃) content in saliva

SI NO	Groups	Base line NO ₃ levels (μmol/l)	NO ₃ levels after 1 hour (μmol/l)	NO ₃ levels after 2 hour (μmol/l)	NO ₃ levels after 3 hour (μmol/l)
1.	Vehicle Control	292.54	295.66	295.95	290.57
2.	Amaranth fresh juice as such	295.56	298.25	297.20	295.57
3.	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	296.55	299.35	299.10	297.01
4.	Amaranth fast dissolving tablet containing water extract of fresh leaves	298.82	302.72	300.2	299.84
5.	Amaranth fast dissolving tablet containing methanol extract of dried leaves	295.02	299.06	298.35	296.77
6.	Amaranth fast dissolving tablet containing water extract of dried leaves	298.25	302.35	301.5	299.56
7.	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	296.50	300.78	298.8	297.23
8.	Amaranth fast dissolving tablet containing enriched extract of nitrate	297.80	2062.25	2120.2	1895.5 1

Table: Nitrite (NO₂) content in saliva

SI NO	Groups	Base line NO ₂ levels (μmol/l)	NO ₂ levels after 1 hour (μmol/l)	NO ₂ levels after 2 hour (μmol/l)	NO ₂ levels after 3 hour (μmol/l)
1.	Vehicle Control	112.54	113.55	112.95	115.39
2.	Amaranth fresh juice as such	115.26	122.57	120.68	116.56
3	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	113.50	121.00	120.65	119.82
4	Amaranth fast dissolving tablet containing water extract of fresh leaves	118.82	126.42	124.66	118.56
5	Amaranth fast dissolving tablet containing methanol extract of dried leaves .	114.93	122.63	120.44	119.64
6	Amaranth fast dissolving tablet containing water extract of dried leaves	115.65	123.45	120.23	117.38
7	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	118.56	126.46	125.56	120.25
8	Amaranth fast dissolving tablet containing enriched extract of nitrate	115.78	1075.20	1005.20	782.50

Table: Total nitric oxide (NO)content in saliva

SI NO	Groups	Base line NO levels ($\mu\text{mol/l}$)	NO levels after 1 hour ($\mu\text{mol/l}$)	NO levels after 2 hour ($\mu\text{mol/l}$)	NO levels after 3 hour ($\mu\text{mol/l}$)
1.	Vehicle Control	405.08	409.21	408.9	405.96
2.	Amaranth fresh juice as such	410.82	420.82	417.88	412.13
3.	Amaranth fast dissolving tablet containing methanol extract of fresh leaves	410.05	420.35	419.75	416.83
4.	Amaranth fast dissolving tablet containing water extract of fresh leaves	417.64	429.14	424.86	418.80
5.	Amaranth fast dissolving tablet containing methanol extract of dried leaves .	409.95	421.69	418.79	416.41
6.	Amaranth fast dissolving tablet containing water extract of dried leaves	413.90	425.80	421.73	416.94
7.	Amaranth fast dissolving tablet containing pectinase treated water extract of fresh leaves	415.06	427.24	424.36	417.48
8.	Amaranth fast dissolving tablet containing enriched extract of nitrate	413.58	3137.45	3125.40	2678.01

Amaranth fresh juice consumption as such slightly increased the NO_3 content after one hour. Methanol extract from amaranth fresh leaves and dried leaves also showed a slight increase in NO_3 content. Tablet made with pectinase treated water extract of fresh leaves of Amaranth shown a little more increase than the other tablets consumptions. But after consumption of fast dissolving tablet

containing Amaranth extract enriched with nitrate, NO_3 level increased to 6.9 fold the baseline value after 1 hour and the increase in NO_3 was maintained for 3 hours. There was no significant change in NO_3 level after administering the vehicle.

- 5 In case of NO_2 also, the fast dissolving tablet containing Amaranth extract enriched with nitrate, increased the NO_2 level more than 9.2 fold and the increase in NO_2 was maintained for 3 hours. Amaranth fresh juice consumption as such slightly increased the NO_2 content after one hour. Methanol extract from amaranth fresh leaves and dried leaves also showed a slight increase in NO_2
10 content. Here also vehicle did not influenced NO_2 level in the subjects .

Overall conclusion is that the total NO_3 and NO_2 level increase was observed in Amaranth fast dissolving tablet was more compared with - Amaranth fresh juice, Methanol extract from fresh and dried leaves . The increase in total NO content of "Amaranth fast dissolving tablet containing enriched extract of nitrate" is 7.5
15 fold from base line and it remained increased for 3 hours. Total NO was interpreted as the sum of total NO_3 and NO_2 . Level of NO_3 and NO_2 was used to interpret the level of NO level in human body. So an increase in NO_3 and NO_2 level was interpreted as an increase in NO level. NO levels was recorded as the sum of the two major metabolites of NO, namely, nitrite and nitrate. (Journal of
20 Medical sciences (2010):3(3): 153-159, Plasma Endothelin -1, Homocysteine, and Oxide Levels in a Multiethnic Hypertensive Cohort from the United Arab Emirates.)

Example 34.

Nitrate (NO_3) Nitrite (NO_2) and nitric oxide(NO) content in saliva of human
25 subjects.

18 Healthy human volunteers (age 35-55 years) were taken for the study. The subjects were put on controlled diet, one week prior to the study. The subjects were divided into three groups comprising of six subjects in each group. The

subjects were fasted overnight and saliva was collected to determine the base line NO_3 and NO_2 levels. After saliva collection, the subjects were treated as follows:

Group 1: Vehicle control.(Water)

Group 2: Amaranth leaves fresh juice as such -100 ml (100 mg Nitrate /100ml of
5 juice) (Prepared as per example No-13)

Group 3: Amaranth extract with enriched nitrate (18% nitrate prepared as per
example 7) is mixed with water.(560 mg of extract in 300 ml water)

All subjects were allowed to consume the water/Juice orally(Group No-1 to 2).
Amaranth extract enriched with nitrate mixed with water was consumed directly
10 in the oral cavity (Group No-3)

After each of the time point 1 hour, 2 hour and 3 hour from feeding, saliva was
collected from all subjects in 5.0ml PTFE tubes. This saliva was stored below -80
degree Celsius until analysis for NO_3 and NO_2 content. The saliva thus obtained
was deprotenized using Acetonitrile and the supernatant liquid was used for NO_3
15 and NO_2 determination using HPLC. NO_3 was analyzed by HPLC on a BETASIL
C-18 column (250 x 4.6mm) using Tetra butyl ammonium hydroxide in De-
ionized water, acetonitrile and methanol as mobile phase and Photo diode array
detection at 222 nm. NO_2 was analyzed at 520 nm using the same conditions. The
Nitrate and Nitrite of the saliva were reported as $\mu\text{mol/L}$.

20

25

Table: Nitrate (NO₃) content in saliva

SI NO	Groups	Base line NO ₃ levels (μmol/l)	NO ₃ levels after 1 hour (μmol/l)	NO ₃ levels after 2 hour (μmol/l)	NO ₃ levels after 3 hour (μmol/l)
1.	Vehicle Control	295.65	296.55	293.78	296.58
2.	Amaranth fresh juice as such	296.23	300.02	297.02	296.33
3	Amaranth extract with enriched nitrate mixed with water	295.68	1961.51	1901.27	789.56

Table: Nitrite (NO₂) content in saliva

SI NO	Groups	Base line NO ₂ levels (μmol/l)	NO ₂ levels after 1 hour (μmol/l)	NO ₂ levels after 2 hour (μmol/l)	NO ₂ levels after 3 hour (μmol/l)
1.	Vehicle Control	110.56	112.56	110.94	113.24
2.	Amaranth fresh juice as such	113.54	120.56	119.85	113.55
3	Amaranth extract with enriched nitrate mixed with water	115.60	1060.45	1055.33	814.34

Table: Total nitric oxide (NO) content in Saliva

SI NO	Groups	Base line NO levels ($\mu\text{mol/l}$)	NO levels after 1 hour ($\mu\text{mol/l}$)	NO levels after 2 hour ($\mu\text{mol/l}$)	NO levels after 3 hour ($\mu\text{mol/l}$)
1.	Vehicle Control	406.21	409.11	404.72	409.82
2.	Amaranth fresh juice as such	409.77	420.58	416.87	409.88
3	Amaranth extract with enriched nitrate mixed with water	411.28	3021.96	2956.60	1603.90

Amaranth fresh juice consumption as such slightly increased the NO_3 content after one hour. But after consumption of Amaranth extract with enriched nitrate mixed with water, NO_3 level increased to 6.6 fold the baseline value after 1 hour and remained increased for 3 hours. There was no significant change in NO_3 level after administering the vehicle.

In case of NO_2 also, Amaranth extract with enriched nitrate mixed with water increased the NO_2 level 9.1 fold and remained increased for 3 hours. Amaranth fresh juice consumption as such slightly increased the NO_2 content after one hour.

Overall conclusion is that the total NO_3 and NO_2 level increase observed by administering Amaranth enriched extract mixed with water was more when compared with administering Amaranth fresh juice. The increase in total NO content of "Amaranth enriched extract mixed with water" was 7.3 times from baseline and it remained increased for 3 hours.

Example 35

Nitrate (NO_3), Nitrite (NO_2) and nitric oxide (NO) content in plasma and saliva of human subjects after administering extract of *Amaranthus blitum* and *Amaranthus viridis* with enriched nitrate.

- 5 Nine healthy human volunteers (age 35-55 years) were taken for the study. The subjects were put on standard diet one week prior to the study. The subjects were divided into three groups comprising of three subjects in each group. The subjects were fasted overnight and the blood and saliva was collected to determine the baseline NO_3 and NO_2 levels. After baseline blood and saliva collection, the
- 10 subjects were treated as follows:

Group 1: Vehicle Control.

Group 2: Amaranth fast dissolving tablet containing extract of *Amaranthus blitum* with enriched nitrate (91.5 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 15).

- 15 Group 3: Amaranth fast dissolving tablet containing extract of *Amaranthus viridis* with enriched nitrate (90.5 mg nitrate per tablet) (Tablet prepared as per example No22 and extract used for preparing the tablet is as per example 15).

All subjects were allowed to consume the water orally (Group No.1). Amaranth Tablets were kept in oral cavity and allow to dissolve (Group No. 2 to 3).

- 20 After 1 hour, 2 hour and 3 hour from feeding of tablet/ vehicle, blood was drawn from all the subjects in EDTA coated tubes and saliva was collected from the same subjects in 5.0 ml PTFE tubes. Plasma was separated immediately by centrifuging the tubes at 5000 rpm for 15 min. The plasma samples and saliva were stored below -80 degree Celsius until analysis. The plasma and saliva were
- 25 deproteinized using Acetonitrile and the supernatant liquid was used for nitrate/nitrite determination using HPLC. NO_3 was analyzed by HPLC on a BETASIL C-18 column (250 x 4.6mm) using Tetra butyl ammonium hydroxide in De-ionized water, acetonitrile and methanol as mobile phase and photo diode

array detection at 222 nm. NO₂ was analysed at 520 nm using the same conditions. The NO₃ and NO₂ levels in plasma were reported as µmol/L.

Nitrate (NO₃), nitrite (NO₂) and nitric oxide (NO) level in plasma

Table: Nitrate (NO₃) content in plasma

SI NO	Groups	Baseline NO ₃ levels (µmol/l)	NO ₃ levels after 1 hour (µmol/l)	NO ₃ levels after 2 hour (µmol/l)	NO ₃ levels after 3 hour (µmol/l)
1.	Vehicle Control	28.92	28.95	27.88	28.81
2	Fast dissolving tablet containing extract of <i>Amaranthus blitum</i> with enriched nitrate.	28.41	172.50	109.43	70.5
3	Fast dissolving tablet containing extract of <i>Amaranthus viridis</i> with enriched nitrate.	27.65	168.92	110.22	70.89

5

Table: Nitrite (NO₂) content in plasma

SI No	Groups	Base line NO ₂ levels (µmol/l)	NO ₂ levels after 1 hour (µmol/l)	NO ₂ levels after 2 hour (µmol/l)	NO ₂ levels after 3 hour (µmol/l)
1.	Vehicle Control	0.19	0.20	0.23	0.18
2	Fast dissolving tablet containing extract of <i>Amaranthus blitum</i> with enriched nitrate.	0.19	1.15	0.94	0.59

3	Fast dissolving tablet containing extract of <i>Amaranthus viridus</i> with enriched nitrate.	0.21	1.27	0.97	0.65
---	---	------	------	------	------

Table: Total nitric oxide (NO) content in plasma

Sl NO	Groups	Base line NO levels ($\mu\text{mol/l}$)	NO levels after 1 hour ($\mu\text{mol/l}$)	NO levels after 2 hour ($\mu\text{mol/l}$)	NO levels after 3 hour ($\mu\text{mol/l}$)
1.	Vehicle Control	29.11	29.15	28.11	28.99
2.	Fast dissolving tablet containing extract of <i>Amaranthus blitum</i> with enriched nitrate	28.6	173.65	110.37	71.09
3.	Fast dissolving tablet containing extract of <i>Amaranthus viridus</i> with enriched nitrate.	27.86	170.19	111.19	71.54

After consumption of fast dissolving tablet containing extract of *Amaranthus blitum* and *Amaranthus viridus* with enriched nitrate, NO_3 level increased to 6 fold the baseline value after 1 hour and the increase from base line was maintained up to 3 hour. There was no significant change in NO_3 level after administering the vehicle.

In case of NO_2 also, the fast dissolving tablet containing extract of *Amaranthus blitum* and *Amaranthus viridus* with enriched nitrate, increased the NO_2 level more than 6 times and the higher levels maintained for 3 hours. Here also vehicle did not influence NO_2 level in the subjects.

The increase in total nitric oxide (NO) content of “Amaranth fast dissolving tablet containing extract of Amaranthus blitum and Amaranthus viridus with enriched nitrate” was 6 times from base line and it remained increased for 3 hours. Total NO was determined from the sum of total NO₃ and NO₂. (ie. Level of NO₃ and NO₂ can be interpreted as a measure of NO level in human body). So an increase in NO₃ and NO₂ level was interpreted as an increase in NO level.

Nitrate (NO₃), nitrite (NO₂) and nitric oxide (NO) level in saliva

Table: Nitrate (NO₃) content in saliva

Sl NO	Groups	Base line NO ₃ levels (μmol/l)	NO ₃ levels after 1 hour (μmol/l)	NO ₃ levels after 2 hour (μmol/l)	NO ₃ levels after 3 hour (μmol/l)
1.	Vehicle Control	292.54	295.66	295.95	290.57
2.	Fast dissolving tablet containing extract of Amaranthus blitum with enriched nitrate	297.10	2058.9	2184.2	1796.51
3.	Fast dissolving tablet containing extract of Amaranthus viridus with enriched nitrate	296.7	2047.52	2147.2	1801.12

Table: Nitrite (NO₂) content in saliva

SI NO	Groups	Base line NO ₂ levels (μmol/l)	NO ₂ levels after 1 hour (μmol/l)	NO ₂ levels after 2 hour (μmol/l)	NO ₂ levels after 3 hour (μmol/l)
1	Vehicle Control	112.54	113.55	112.95	115.39
2	Fast dissolving tablet containing extract of <i>Amaranthus blitum</i> with enriched nitrate	115.75	1064.9	1021.1	796.70
3	Fast dissolving tablet containing extract of <i>Amaranthus viridus</i> with enriched nitrate	113.58	1033.57	1001.4	701.3

Table: Total nitric oxide (NO) content in saliva

SI NO	Groups	Base line NO levels (μmol/l)	NO levels after 1 hour (μmol/l)	NO levels after 2 hour (μmol/l)	NO levels after 3 hour (μmol/l)
1.	Vehicle Control	405.08	409.21	408.9	405.96
2.	Fast dissolving tablet containing extract of <i>Amaranthus blitum</i> with enriched nitrate	412.85	3123.8	3205.3	2593.21
3.	Fast dissolving tablet containing extract of <i>Amaranthus viridus</i> with enriched nitrate	410.28	3081.09	3148.6	2502.42

After consumption of fast dissolving tablet containing extract of *Amaranthus blitum* and *Amaranthus viridus* with enriched nitrate, NO_3 level increased to 6.9 fold the baseline value after 1 hour and the increase in NO_3 was maintained for 3 hours. There was no significant change in NO_3 level after administering the vehicle.

In case of NO_2 also, the fast dissolving tablet containing extract of *Amaranthus blitum* and *Amaranthus viridus* with enriched nitrate, increased the NO_2 level more than 9.2 fold and the increase in NO_2 was maintained for 3 hours. Here also vehicle did not influenced NO_2 level in the subjects.

The increase in total NO content of "Amaranth fast dissolving tablet containing extract of *Amaranthus blitum* and *Amaranthus viridus* with enriched nitrate" was 7.5 fold from base line and it remained increased for 3 hours. Total NO was interpreted as the sum of total NO_3 and NO_2 . (Level of NO_3 and NO_2 is used to interpret the level of NO level in human body). So an increase in NO_3 and NO_2 level was interpreted as an increase in NO level.

I have brought out the novel features of the invention by explaining some of the preferred embodiments under the invention, enabling those in the art to understand and visualize the disclosure. It is also to be understood that the invention is not limited in its application to the details set forth in the above description or as illustrated in the examples. Although the invention has been described in considerable detail with reference to certain preferred embodiments thereof, various modifications can be made without departing from the spirit and scope of the invention as described herein above.

I claim:

1. An extract of Amaranth, the extract comprising
 - about 0.1% to about 20% nitrates;
 - about 0.1% to about 10% of L-arginine;
 - 5 about 0.001% to about 10% of flavonoids;
 - about 0.01% to about 15% of saponins;
 - about 0.01% to about 10% of alkaloids;
 - about 10% to about 30% of carbohydrates;
 - about 5% to about 25% of proteins;
 - 10 about 5 % to about 25% potassium; and,
 - about 0.01% to about 50% of the total oxalic acid, wherein the total oxalic acid comprises free oxalic acid and oxalates.
2. The extract of Amaranth of claim 1 further comprising a second extract selected from the group consisting of amla extract, turmeric extract, grape seed extract, green tea extract, pomegranate extract, cocoa extract, coconut root extract, rosemary extract, mint leaf extract, star anise, sweet basil extract, cinnamon extract/clove extract, ginger extract, cumin seed extract, black pepper extract, fenugreek extract, and combinations thereof.
3. The extract of Amaranth of claim 1 further comprising Vitamin C.
- 20 4. The extract of Amaranth of claim 1, wherein the Amaranth is selected from the group consisting of *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus tricolor*, *Amaranthus blitum*, *Amaranthus viridis*, *Amaranthus dubius* and combinations thereof.

5. An extract of Amaranth having low total oxalic acid content, the extract comprising
- about 3% to about 20% nitrates;
 - about 1% to about 10% of L-arginine;
 - 5 about 1% to about 10% of flavonoids;
 - about 5% to about 15% of saponins;
 - about 3% to about 10% of alkaloids;
 - about 10% to about 30% of carbohydrates;
 - about 5% to about 25% of proteins;
 - 10 about 1% to about 5% of potassium; and,
 - about 0.1% total oxalic acid, wherein the total oxalic acid comprises free oxalic acid and oxalates.
6. The extract of Amaranth of claim 5 further comprising a second extract selected from the group consisting of amla extract, turmeric extract, grape
- 15 seed extract, green tea extract, pomegranate extract, cocoa extract, coconut root extract, rosemary extract, mint leaf extract, star anise, sweet basil extract, cinnamon extract/clove extract, ginger extract, cumin seed extract, black pepper extract, fenugreek extract, and combinations thereof.
7. The extract of claim 5 further comprising Vitamin C.
- 20 8. The extract of Amaranth of claim 5, wherein the Amaranth is selected from the group consisting of *Amaranthus caudatus*, *Amaranthus cruentus*, *Amaranthus tricolor*, *Amaranthus blitum*, *Amaranthus viridis*, *Amaranthus dubius* and combinations thereof.

9. A dosage form comprising the extract of any one of claims 1 to 8, the dosage form is selected from the group consisting of fast melt tablet, lozenge, candy, chewing gum, beverage, tablet, capsule, pill, and powder.
10. A method of lowering blood pressure comprising administering the extract of claims 1 to 8.
11. A method of increasing physical endurance comprising administering the extract of claims 1 to 8.
12. A method of increasing swimming endurance comprising administering the extract of claims 1 to 8.
13. A method of increasing running endurance comprising administering the extract of claims 1 to 8.
14. A method of improving sexual performance comprising administering the extract of claims 1 to 8.
15. A method of enhancing nitric oxide concentration in blood comprising administering the extract of claims 1 to 8.
16. A method of enhancing nitrate concentration in blood comprising administering the extract of claims 1 to 8.
17. A method of enhancing nitrite concentration in blood comprising administering the extract of claims 1 to 8.
18. A method of enhancing nitric oxide concentration in saliva comprising administering the extract of claims 1 to 8.
19. A method of enhancing nitrate concentration in saliva comprising administering the extract of claims 1 to 8.
20. A method of enhancing nitrite concentration in saliva comprising administering the extract of claims 1 to 8.

21. A method of enhancing nitrate level in blood by about 2 fold to about 10 fold comprising administering the extract of any one of claims 1 to 8.
22. A method of enhancing nitrite level in blood by about 3 fold to about 15 fold comprising administering the extract of claims 1 to 8.
- 5 23. A method of enhancing nitric oxide level in blood by about 2 fold to about 10 fold comprising administering the extract of claims 1 to 8.
24. A method of enhancing nitrate level in saliva by about 2 fold to about 10 fold comprising administering the extract of any of claims 1 to 8.
25. A method of enhancing nitrite in saliva level by about 3 fold to about 15 fold
10 comprising administering the extract of claims 1 to 8.
26. A method of enhancing nitric oxide in saliva level by about 2 fold to about 10 fold comprising administering the extract of claims 1 to 8.
27. A method of preparing a powdered extract of fresh Amaranth, the method comprising:
15 crushing fresh leaves and stem of Amaranth to obtain a crushed material,
extracting the crushed material with methanol to obtain a supernatant and residue,
filtering the supernatant to obtain a filtrate;
concentrating the filtrate to obtained a concentrated filtrate, and,
20 drying the concentrated filtrate to obtain a powdered extract of fresh Amaranth.
28. A method of preparing a powdered extract of fresh Amaranth, the method comprising:
crushing fresh leaves and stem of Amaranth to obtain a crushed material,

extracting the crushed material with water to obtain a supernatant and a residue, and,

concentrating the supernatant to obtain a concentrated water extract,

drying the concentrated water extract to obtain a powdered extract of fresh
5 Amaranth.

29. A method of preparing a powdered extract of dried Amaranth, the method comprising:

powdering dried leaves and stem of Amaranth to obtain a powdered material,

10 extracting the powdered material with methanol to obtain a supernatant,

filtering the supernatant to obtain a filtrate,

concentrating the filtrate to obtain a concentrated methanol filtrate, and,

drying the concentrated methanol filtrate to obtain a powdered extract of dried Amaranth.

15 30. A method of preparing a powdered extract of dried Amaranth, the method comprising:

powdering dried leaves and stem of Amaranth to obtain a powdered material,

20 extracting the powdered material with water to obtain a supernatant and a residue,

concentrating the supernatant to obtain a concentrated water extract, and,

drying the concentrated water extract to obtain a powdered extract of dried Amaranth.

31. A method preparing an extract of Amaranth, the method comprising:

- crushing fresh leaves and stem of Amaranth to obtain a slurry,
- treating the slurry with pectinase to obtain a pectinase treated material,
- heating the pectinase treated material to obtain a pectinase-deactivated material,
- 5 extracting the pectinase-deactivated material with water to obtain a supernatant and a residue,
- concentrating the supernatant to obtain a concentrated supernatant,
- drying the concentrated supernatant to obtain a powdered extract of fresh Amaranth.
- 10 32. A method of preparing an extract of Amaranth, the method comprising:
- crushing fresh leaves and stem of Amaranth to obtain a slurry,
- treating the slurry with pectinase to obtain a pectinase treated material,
- heating the pectinase treated material to obtain a pectinase-deactivated material,
- 15 extracting the pectinase-deactivated material with water to obtain a supernatant and a residue,
- concentrating the supernatant to obtain a concentrated water extract,
- cooling the concentrated water extract at 10°C for 48 hours then filtering to obtain crystals of oxalic acid and/or oxalate and a second supernatant;
- 20 cooling the second supernatant at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid and/or oxalate crystals;
- filtering the third supernatant to obtain a filtrate,
- concentrating the filtrate to obtain a concentrated filtrate, and,

drying the concentrated filtrate to obtain a powdered extract, wherein the powdered extract comprises less than 0.1% total oxalic acid, wherein total oxalic acid comprises free oxalic acid and oxalates.

33. A method of preparing an extract of Amaranth, the method comprising:

- 5 crushing fresh leaves and stem of Amaranth to obtain a slurry,
treating the slurry with pectinase to obtain a pectinase treated material,
heating the pectinase treated material to obtain a pectinase-deactivated material,
extracting the pectinase-deactivated material with water to obtain a
10 supernatant and a residue,
concentrating the supernatant to obtain a concentrated water extract,
cooling the concentrated water extract at 10° C for 48 hours then filtering to obtain crystals of oxalic acid and/or oxalate and a second supernatant;
cooling the second supernatant at 10° C for 24 hours to obtain a third
15 supernatant and a second oxalic acid and/or oxalate crystals;
filtering the third supernatant to obtain a filtrate,
concentrating the filtrate to obtain a concentrated filtrate, and,
drying the concentrated filtrate to obtain a powdered extract,
treating the powdered extract with hexane to obtain a mixture,
20 filtering the mixture to obtain a filtrate,
concentrating the filtrate to obtain a concentrated hexane extract,
drying the concentrated hexane extract to obtain a powdered hexane extract.

34. A method of preparing an extract of Amaranth, the method comprising:

- crushing fresh leaves and stem of Amaranth to obtain a slurry,
treating the slurry with pectinase to obtain a pectinase treated material,
heating the pectinase treated material to obtain a pectinase-deactivated material,
- 5 extracting the pectinase-deactivated material with water to obtain a supernatant and a residue,
concentrating the supernatant to obtain a concentrated water extract,
cooling the concentrated water extract at 10° C for 48 hours then filtering to obtain crystals of oxalic acid and/or oxalates and a second supernatant;
- 10 cooling the second supernatant at 10° C for 24 hours to obtain a third supernatant and a second oxalic acid and/or oxalate crystals;
filtering the third supernatant to obtain a filtrate,
concentrating the filtrate to obtain a concentrated filtrate, and,
drying the concentrated filtrate to obtain a powdered extract,
- 15 treating the powdered extract with chloroform to obtain a mixture,
filtering the mixture to obtain a filtrate,
concentrating the filtrate to obtain a concentrated chloroform extract,
drying the concentrated chloroform extract to obtain a powdered chloroform extract.
- 20 35. A method of preparing an extract of Amaranth, the method comprising:
crushing fresh leaves and stem of Amaranth to obtain a slurry,
treating the slurry with pectinase to obtain a pectinase treated material,

- heating the pectinase treated material to obtain a pectinase-deactivated material,
- extracting the pectinase-deactivated material with water to obtain a supernatant and a residue,
- 5 concentrating the supernatant to obtain a concentrated water extract,
- cooling the concentrated water extract at 10°C for 48 hours then filtering to obtain crystals of oxalic acid and/or oxalate and a second supernatant;
- cooling the second supernatant at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid and/or oxalate crystals;
- 10 filtering the third supernatant to obtain a filtrate,
- concentrating the filtrate to obtain a concentrated filtrate, and,
- drying the concentrated filtrate to obtain a powdered extract,
- treating the powdered extract with ethylene dichloride to obtain a mixture,
- filtering the mixture to obtain a second filtrate,
- 15 concentrating the second filtrate to obtain a concentrated ethylene dichloride extract,
- drying the concentrated ethylene dichloride extract to obtain a powdered ethylene dichloride extract.
36. A method of preparing an extract of Amaranth additionally with Amla
20 extract, the method comprising:
- crushing fresh leaves and stem of Amaranth to obtain a slurry,
- treating the slurry with pectinase to obtain a pectinase treated material,
- heating the pectinase treated material to obtain a pectinase-deactivated material,

- extracting the pectinase-deactivated material with water to obtain a supernatant and a residue,
- concentrating the supernatant to obtain a concentrated water extract,
- cooling the concentrated water extract at 10°C for 48 hours then filtering to obtain crystals of oxalic acid and/or oxalate and a second supernatant;
- 5 cooling the second supernatant at 10°C for 24 hours to obtain a third supernatant and a second oxalic acid and/or oxalate crystals;
- filtering the third supernatant to obtain a filtrate,
- concentrating the filtrate to obtain a concentrated filtrate, and,
- 10 drying the concentrated filtrate to obtain a powdered extract,
- treating the powdered extract with hexane to obtain a mixture,
- filtering the mixture to obtain a second filtrate,
- concentrating the second filtrate to obtain a concentrated hexane extract,
- drying the concentrated hexane extract to obtain a powdered hexane extract,
- 15 dissolving the powdered hexane extract in methanol to obtain a solution of hexane extract of Amaranth,
- adding a powder of methanol extract of Amla to the solution of hexane extract of Amaranth to obtain a second mixture,
- evaporating the second mixture to obtain an concentrate,
- 20 drying the concentrate to obtain a powder of Amaranth and Amla extract,
- wherein the powdered methanol extract of Amla extract is prepared by a process comprising:
- pulping fresh Amla fruit to obtain a pulp,

extracting the pulp with methanol to obtain a filtrate,

concentrating the filtrate to obtain a concentrated methanol extract of Amla,

drying the concentrated methanol extract of Amla to obtain the powdered methanol extract of Amla.