CLAIMS

1. A fungicidal composition comprising:
   (a) oxathiapiprolin,
   (b) metalaxyl-M, and
   (c) a lactamide compound of formula I

$$CH_3CH(OH)C(=O)NR_1R_2 \quad (I)$$

wherein R_1 and R_2 are each independently hydrogen; or C_{1-6} alkyl, C_{2-6} alkenyl or C_{3-6} cycloalkyl, each of which is optionally substituted by up to three substituents independently selected from phenyl, hydroxy, C_{1-5} alkoxy, morpholinyl and NR_3R_4 where R_3 and R_4 are each independently C_{1-3} alkyl; or phenyl optionally substituted by up to three substituents independently selected from C_{1-3} alkyl; or R_1 and R_2 together with the nitrogen atom to which they are attached form a morpholinyl, pyrrolidinyl, piperidinyl or azepanyl ring, each of which is optionally substituted by up to three substituents independently selected from C_{1-3} alkyl,

the composition comprising at least 40% by weight of said lactamide compound, over the total weight of the composition, and

wherein said composition comprises at least 5% by weight of metalaxyl-M, over the total weight of the composition.

2. A composition according to claim 1, characterized in that the composition comprising at least 50% by weight of said lactamide compound, and preferably at least 60% by weight of said lactamide compound, over the total weight of the composition.

3. A composition according to any one of the preceding claims, characterized in that the composition comprises at least 10% by weight of metalaxyl-M, over the total weight of the composition.

4. A composition according to any one of the preceding claims, characterized in that the composition comprises:
   (a) from 1% to 10% by weight of oxathiapiprolin, over the total weight of the composition,
   (b) from 5% to 30% by weight of metalaxyl-M, over the total weight of the composition, and
   (c) from 40% to 85% by weight of said lactamide compound, over the total weight of the composition.

5. A composition according to any one of the preceding claims, characterized in that R_1 and R_2 are each independently hydrogen or C_{1-6} alkyl.

6. A composition according to any one of the preceding claims, characterized in that the composition further comprises one or several emulsifiers.

7. A composition according to claim 6, characterized in that the composition comprises from 0.01% to 40% by weight of emulsifier, preferably from 0.5% to 30% by weight of emulsifier, and more preferably from 0.5% to 20% by weight of emulsifier, over the total weight of the composition.

8. A composition according to any one of the preceding claims, characterized in that the composition further comprises one or several dispersing agents.

9. A composition according to claim 8, characterized in that the composition comprises from 0.01% to 20% by weight of dispersing agent, and preferably from 0.5% to 10% by weight of dispersing agent, over the total weight of the composition.

10. A composition according to any one of the preceding claims, characterized in that the composition is a dispersible concentrate or an emulsifiable concentrate.

11. A method of controlling or preventing an infestation of plants by fungal organisms, which comprises the foliar and/or the soil application(s) of the composition according to any one of the preceding claims.

12. A method according to claim 11, characterized in that the plant is selected among grapes, vegetables and permanent crops.

13. Use of a composition according of any one of claims 1 to 10 to control or prevent an infestation of plants by fungal organisms.