PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

IPOS COPY

20000196 4-6

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :
A61K 31/00, 31/44, 31/425

A1

(11) International Publication Number: WO 99/18943

(43) International Publication Date: 22 April 1999 (22.04.99)

(21) International Application Number: PCT/GB98/03064

(22) International Filing Date: 12 October 1998 (12.10.98)

(30) Priority Data:
9721693.1 13 October 1997 (13.10.97) GB

(71) Applicant *(for all designated States except US)*: SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: BUCKINGHAM, Robin, Edwin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). SMITH, Stephen, Alistair [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).

(74) Agent: GIDDINGS, Peter, John; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: USE OF THIAZOLIDINEDIONES FOR THE TREATMENT OF HYPERGLYCAEMIA

(57) Abstract

A method for the treatment of hyperglycaemia wherein plasma glucose levels are in the range of elevated normal to ≤126mg/dl, which method comprises administering an effective non–toxic and pharmaceutically acceptable amount of an insulin sensitiser, to a mammal in need thereof.