

1/15




FIG. 1

**Table 1.** Clinical characteristics of study participants (Guangzhou sample)

| Group                   | HCC<br>(n=75) | Hepatitis<br>(n=128) |
|-------------------------|---------------|----------------------|
| Gender                  |               |                      |
| Male                    | 70 (93%)      | 99 (77%)             |
| Female                  | 5 (7%)        | 29 (23%)             |
| Age (years)             | 52 (29-76)    | 49 (18-69)           |
| >50                     | 38 (51%)      | 26 (20%)             |
| AFP ng/ml <sup>a</sup>  |               |                      |
| ≤ 20                    | 25 (34%)      | 66 (50%)             |
| >20                     | 48 (64%)      | 36 (28%)             |
| ALB g/l <sup>b</sup>    |               |                      |
| ≥35                     | 59 (79%)      | 74 (58%)             |
| ALT U/L <sup>c</sup>    |               |                      |
| ≤ 40                    | 34 (45%)      | 22 (17%)             |
| Cirrhosis <sup>d</sup>  | 54 (72%)      | 29 (27%)             |
| Tumor size <sup>e</sup> |               |                      |
| ≤ 3cm                   | 11 (15%)      | NA                   |
| 3-5 cm                  | 22 (31%)      | NA                   |
| >5 cm                   | 39 (54%)      | NA                   |

a. 73 out of 75 HCC patients have AFP values;  
100 out of 128 hepatitis patients have AFP values.  
b. 118 out of 128 hepatitis patients have ALB values.  
c. 123 out of 128 hepatitis patients have ALT values.  
d. 29 hepatitis patients were confirmed with cirrhosis.  
e. 72 out of 75 HCC patients have tumor size information

*FIG. 2A*

Table 2. Clinical characteristics of study participants (NCC/SGH sample)

| Group                   | HCC<br>(n=35) | Hepatitis<br>(n=12) |
|-------------------------|---------------|---------------------|
| Gender                  |               |                     |
| Male                    | 17 (49%)      | 9 (75%)             |
| Female                  | 4 (11%)       | 2 (17%)             |
| Age (years)             |               |                     |
| >50                     | 20 (57%)      | 0 (0%)              |
| AFP ng/ml <sup>a</sup>  |               |                     |
| ≤ 20                    | 19 (54%)      | 7 (58%)             |
| >20                     | 11 (31%)      | 0 (0%)              |
| ALB g/l <sup>b</sup>    |               |                     |
| ≥35                     | 17 (49%)      | 4 (33%)             |
| ALT U/L <sup>c</sup>    |               |                     |
| ≤ 40                    | 11 (31%)      | 5 (42%)             |
| Cirrhosis <sup>d</sup>  | 13 (37%)      | 6 (0.5%)            |
| Tumor size <sup>e</sup> |               |                     |
| ≤ 3cm                   | 5 (14%)       | NA                  |
| 3-5 cm                  | 5 (14%)       | NA                  |
| >5 cm                   | 15 (43%)      | NA                  |

a. 31 out of 35 HCC patients have AFP values;  
7 out of 12 hepatitis patients have AFP values.

b. 5 out of 12 hepatitis patients have ALB values.

c. 6 out of 12 hepatitis patients have ALT values.

d. 6 out of 12 hepatitis patients were confirmed with cirrhosis.

e. 26 out of 35 HCC patients have tumor size information

FIG. 2B

4/15

**Table 3** Differential expression and diagnostic performance of the 9 significant genes in the training group (HCC n=50, CHB n=50).

| Gene Name | Fold Change | AUC  | Std. Error | P-value | 95% Confidence Interval |
|-----------|-------------|------|------------|---------|-------------------------|
| GIMAP6    | -1.94801    | .841 | .039       | .000    | .764 - .919             |
| GIMAP4    | -2.11316    | .808 | .043       | .000    | .723 - .892             |
| GIMAP5    | -2.70232    | .906 | .030       | .000    | .848 - .965             |
| GIMAP8    | -2.45173    | .801 | .043       | .000    | .716 - .885             |
| TNFAIP    | 4.67122     | .943 | .024       | .000    | .896 - .989             |
| AREG      | 8.28786     | .926 | .027       | .000    | .873 - .978             |
| NFKBIA    | 4.64282     | .869 | .036       | .000    | .799 - .940             |
| NKFBIZ    | 3.55833     | .885 | .032       | .000    | .823 - .947             |
| CD83      | 2.82843     | .776 | .049       | .000    | .679 - .872             |

*FIG. 3A*






FIG. 3B

6/15

**FIG. 3C**

**Training Group**  
50 HCC and 50 Hepatitis



Diagonal segments are produced by ties.

FIG. 4A

| Area Under the Curve    |      |            |                 | Area Under the Curve               |             |      |            |
|-------------------------|------|------------|-----------------|------------------------------------|-------------|------|------------|
| Test Result Variable(s) | Area | Std. Error | Asymptotic Sig. | Asymptotic 95% Confidence Interval |             | Area | Std. Error |
|                         |      |            |                 | Lower Bound                        | Upper Bound |      |            |
| HCC_p_TNFAIP3_AREG_G5   | .977 | .014       | .000            | .940                               | 1.000       | .909 | .028       |
| TNFAIP3                 | .943 | .024       | .000            | .896                               | .989        | .891 | .029       |


| Test Result Variable(s) | Area | Std. Error | Asymptotic Sig. | Asymptotic 95% Confidence Interval |             |
|-------------------------|------|------------|-----------------|------------------------------------|-------------|
|                         |      |            |                 | Lower Bound                        | Upper Bound |
| HCC_p_TNFAIP3_AREG_G5   | .909 | .028       | .000            | .853                               | .964        |
| TNFAIP3                 | .891 | .029       | .000            | .834                               | .947        |

ROC curves analysis of different marker models in the training group (A) and testing group (B). Area Under Curve (AUC) is shown with 95% confidence interval. Models were developed by stepwise logistic regression (forward method). The probability of being HCC was calculated from the odds ratio and was given as a score ranging from 0 to 100.

8/15

**Testing Group**

69 HCC and 93 Hepatitis

**FIG. 4B**

Diagonal segments are produced by ties.


| Area Under the Curve    |      |            |                 | Area Under the Curve |            |                 |                                    |
|-------------------------|------|------------|-----------------|----------------------|------------|-----------------|------------------------------------|
| Test Result Variable(s) | Area | Std. Error | Asymptotic Sig. | Area                 | Std. Error | Asymptotic Sig. | Asymptotic 95% Confidence Interval |
| HCC_p_TNFAIP3_AREG_G5   | .977 | .014       | .000            | .909                 | .028       | .000            | Lower Bound .853 Upper Bound .964  |
| TNFAIP3                 | .943 | .024       | .000            | .891                 | .029       | .000            | Lower Bound .834 Upper Bound .947  |

ROC curves analysis of different marker models in the training group (A) and testing group (B). Area Under Curve (AUC) is shown with 95% confidence interval. Models were developed by stepwise logistic regression (forward method). The probability of being HCC was calculated from the odds ratio and was given as a score ranging from 0 to 100.

**Table 3.** Sensitivity (TPR) and specificity (1-FPR) from ROC analysis of training and testing groups at different cutoff points.

| Model                              | Cut-off | Training |       | Testing |       |
|------------------------------------|---------|----------|-------|---------|-------|
|                                    |         | TPR      | 1-FPR | TPR     | 1-FPR |
| TNFAIP3 alone                      | > 0.58  | 80%      | 92%   | 80%     | 88%   |
|                                    | > 0.73  | 76%      | 98%   | 75%     | 94%   |
|                                    | > 0.90  | 58%      | 100%  | 55%     | 96%   |
| TNFAIP3 Set (TNFAIP, AREG, GIMAP5) | > 0.58  | 92%      | 96%   | 85%     | 87%   |
|                                    | > 0.73  | 88%      | 96%   | 82%     | 90%   |
|                                    | > 0.90  | 72%      | 100%  | 77%     | 97%   |

*FIG. 5*



Diagonal segments are produced by ties.


Sensitivity (TPR) and specificity (1-FPR) of AFP and WBC gene markers (HCC n=104; CHB n=108).

| Classifier                      | Cut-off | TPR | 1-FPR |
|---------------------------------|---------|-----|-------|
| TNFAIP3 (probability score)     | > 55    | 81% | 90%   |
|                                 | > 69    | 79% | 95%   |
|                                 | > 90    | 57% | 98%   |
| TNFAIP3 Set (probability score) | > 57    | 90% | 92%   |
|                                 | > 70    | 87% | 93%   |
|                                 | > 90    | 74% | 99%   |
| AFP (ng/ml)                     | > 20    | 58% | 67%   |
|                                 | > 107   | 47% | 89%   |
|                                 | > 209   | 43% | 95%   |

| Test Result Variable(s) | Area Under the Curve |                         |                              | Asymptotic 95% Confidence Interval |             |
|-------------------------|----------------------|-------------------------|------------------------------|------------------------------------|-------------|
|                         | Area                 | Std. Error <sup>a</sup> | Asymptotic Sig. <sup>b</sup> | Lower Bound                        | Upper Bound |
| HCC_p_TNFAIP_AREG_G5    | .951                 | .015                    | .000                         | .921                               | .981        |
| TNFAIP                  | .925                 | .019                    | .000                         | .889                               | .961        |
| AFP                     | .697                 | .036                    | .000                         | .626                               | .769        |

FIG. 6

ROC curves analysis of WBC gene markers and serum AFP in 104 HCC and 108 CHB patients.




Diagonal segments are produced by ties.

| Test Result Variable(s) | Area Under the Curve |                         |                              |                                    |             |
|-------------------------|----------------------|-------------------------|------------------------------|------------------------------------|-------------|
|                         | Area                 | Std. Error <sup>a</sup> | Asymptotic Sig. <sup>b</sup> | Asymptotic 95% Confidence Interval |             |
|                         |                      |                         |                              | Lower Bound                        | Upper Bound |
| HCC_p_TNFAIP_AREG_G5    | .963                 | .031                    | .000                         | .000                               | 1.000       |
| TNFAIP3                 | .985                 | .009                    | .000                         | .959                               | 1.000       |
| AFP                     | .618                 | .073                    | .153                         | .475                               | .761        |

FIG. 7

ROC curves analysis of WBC gene markers and serum AFP in 14 BCLC stage A HCC patients (single nodule 3 cm, no vascular invasion and 140 CHB patients).

12/15

**FIG. 8**HCC vs Hepatitis ( $p < 0.0003$ ,  $q < 0.03$ ,  $FC > 1.5$ )HCC vs Normal ( $p < 0.0003$ ,  $q < 0.001$ ,  $FC > 1.5$ )Normal vs Hepatitis ( $p < 0.0003$ ,  $q < 0.001$ ,  $FC > 1.5$ )

Venn Diagram for three pairwise comparisons. Candidate gene markers were selected from those genes that are differentially expressed in HCC compared with CHB and healthy subjects.

13/15

**Table 4** Differential gene expression of the 9 significant genes in HCC and CHB from gene microarray analysis.

| Gene Symbol | Gene Title                                                                          | p-value   | q-value | HCC/CHB |
|-------------|-------------------------------------------------------------------------------------|-----------|---------|---------|
| TNFAIP3     | Tumor necrosis factor, alpha-induced protein 3                                      | 3.68E-005 | 0.015   | 3.72    |
| NFKBIA      | Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha | 0.00013   | 0.027   | 3.06    |
| NFKBIZ      | Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta  | 5.05E-005 | 0.017   | 3.8     |
| AREG        | Amphiregulin                                                                        | 5.04E-007 | 0.0004  | 8.6     |
| CD83        | CD83 molecule                                                                       | 0.0014    | 0.080   | 4.83    |
| GIMAP4      | GTPase, IMAP family member 4                                                        | 2.05E-005 | 0.013   | -2.04   |
| GIMAP5      | GTPase, IMAP family member 5                                                        | 1.4E-05   | 0.008   | -2.05   |
| GIMAP6      | GTPase, IMAP family member 6                                                        | 5.11E-005 | 0.017   | -1.70   |
| GIMAP8      | GTPase, IMAP family member 8                                                        | 0.0001    | 0.024   | -2.34   |

E/C 0

14/15




**FIG. 10** Differential expression of WBC gene markers in the training group validated by q-PCR (HCC n=50; CHB n=50; Healthy n=6). Gene expression levels were normalized to that of CD45 and presented as percentage of CD45 expression level. Box refers to the 25<sup>th</sup> and 75<sup>th</sup> percentile, with the line indicating the median. Whiskers represent the minimum and maximum values. Mann-Whitney test was performed to determine the significance.

15/15

|           | <u>Forward 5'-3'</u>     | <u>Reverse 5'-3'</u>      |
|-----------|--------------------------|---------------------------|
| 1 NFKBIZ  | TCCTGTTGCACATCCGAAGTC    | TCCATCAGACAACGAATCGGG     |
| 2 GIMAP8  | GGGTCGCTCTCCGGCCATTTC    | CAGGCTCCCGCTTGTCTGGG      |
| 3 GIMAP5  | GTGCAGCTGAGTCATGGAGCTT   | TTCTCTCCAGAAACGGTTGTGTGC  |
| 4 AREG    | GTGGTGCTGTCGCTCTTGATACTC | TCAATCCATCAGCACTGTGGTC    |
| 5 CD83    | TGCACTCTGCAGGACCCGGA     | TGTAGCCGTGCAAACTTACAAGTGA |
| 6 NFKBIA  | CTCCGAGACTTTCGAGGAAATAC  | GCCATTGTAGTTGGTAGCCTTCA   |
| 7 GIMAP6  | GTCTTCGAGTCTAAACTCAGCAC  | TGGGTGTGTCAATCACCTCAA     |
| 8 TNFAIP3 | TTGTCCTCAGTTTCGGGAGAT    | TTCTCGACACCCAGTTGAGTTTC   |
| 9 GIMAP4  | GCCCAATACGGCAGTATGAG     | CCTGCTCCGGTTTTACCCAC      |

FIG. 11