

HBV POLYMERASE MUTANTS

FIELD OF THE INVENTION

5 The present invention relates to polymerase HBV mutant polypeptides comprising a mutated polymerase domain which is functionally disrupted for polymerase activity and fusion proteins comprising such polymerase mutant polypeptide. The present invention also relates to a nucleic acid molecule and an expression vector for expressing said polymerase mutant polypeptide as well as a
10 composition which can be used for eliciting an immune response to HBV with the goal of providing a protective or therapeutic effect against HBV infection. The invention is of very special interest in the field of immunotherapy, and more particular for treating patients infected with HBV, especially those chronically infected.

15 BACKGROUND OF THE INVENTION

Hepatitis B is a major public health problem with more than 350 million persons chronically infected worldwide, 20 to 40% of them being at risk of developing chronic liver disease, cirrhosis and hepatocellular carcinoma. Despite the existence of effective preventive vaccines, the hepatitis B virus (HBV) infection is still rampant in many
20 countries, even developed ones, with an estimation of 4.5 millions of new cases of infection per year worldwide. Unlike the WHO recommendation which is to implement universal vaccination, the coverage of full course preventive vaccination varies from 25% in Asia to 75-90% in Europe. Currently hepatitis B is the 10th cause of mortality (around 1 million of deaths/year) and HBV related liver carcinoma, the 5th most
25 frequent cancer. Geographic repartition of HBV infection is uneven with prevalence lower than 1% in Western countries to more than 10% in South Eastern countries, most part of Africa and Equatorial South America.

Hepatitis B virus is a member of the hepadnaviridae and primarily infects the liver, replicating in hepatocytes. The infectious particles are the so called 42-45nm
30 "Dane particles" which consist of an outer lipoprotein envelope with three different surface proteins (HBs) and an inner nucleocapsid, the major structural protein of which is the core protein (HBcAg). Within the nucleocapsid is a single copy of the HBV

genome linked to the viral polymerase protein (P). In addition to 42–45nm virions, the blood of HBV-infected patients contains 20-nm spheres made of HBsAg and host-derived lipids which are released from infected cells. These spheres outnumber the virions by a factor of 10^4 – 10^6 .

5 After virions enter hepatocytes, by an as-yet-unknown receptor, nucleocapsids transport the genomic HBV DNA to the nucleus, where the relaxed circular DNA is converted to covalently closed circular DNA (cccDNA). The cccDNA functions as the template for the transcription of four viral RNAs, which are exported to the cytoplasm and used as mRNAs for translation of the HBV proteins. The longest (pre-genomic)
10 RNA also functions as the template for HBV replication, which occurs in nucleocapsids in the cytoplasm. Some of the HBV DNA and polymerase-containing capsids are then transported back to the nucleus, where they release the newly generated relaxed circular DNA to form additional cccDNA. With a half-life longer than the one of hepatocytes, the cccDNA is responsible for the persistence of HBV. Other capsids are enveloped by
15 budding into the endoplasmic reticulum and secreted after passing through the Golgi complex.

Structural and functional organization of the HBV genome has been investigated for more than 30 years. The HBV genome is a relaxed circular partially double-stranded DNA of approximately 3,200 nucleotides consisting of a full-length negative strand and
20 a shorter positive strand. It contains 4 overlapping open reading frames (ORFs), C, S, P and X. The C ORF encodes the core protein (or HBcAg), a 183 amino acid-long protein constitutive of the nucleocapsid and a second protein found in the serum of patients during virus replication known as HBeAg which contains a precore N-terminal extension and a part of HBcAg. The C-terminus of the core protein is very basic and
25 contains 4 Arg-rich domains which are predicted to bind nucleic acids as well as numerous phosphorylation sites. The S ORF encodes three surface proteins all of which have the same C terminus but differ at their N-termini due to the presence of three in-frame ATG start codons that divide the S ORF into three regions, S (226 amino acids), pre-S2 (55 amino acids) and pre-S1 (108 amino acids), respectively. The large-surface
30 antigen protein (L) is produced following translation initiation at the first ATG start codon and comprises 389 amino acid residues (preS1-preS2-S). The middle surface antigen protein (M) results from translation of the S region and the pre-S2 region

starting at the second start ATG whereas the small surface antigen protein of 226 amino acids (S, also designated HBsAg) results from translation of the S region initiated at the third start ATG codon. The HBV surface proteins are glycoproteins with carbohydrate side chains (glycans) attached by N-glycosidic linkages. The P ORF encodes the viral
5 polymerase and the X ORF a protein known as the X protein which is thought to be a transcriptional activator.

The viral polymerase is about 832-845 amino acid residues long according to the HBV genotype and it is encoded in a long open reading frame ("P") that overlaps the
10 3'end of the core gene and all the surface protein genes. The viral polymerase is a multifunctional protein composed of four domains, including three functional domains, respectively the terminal protein, polymerase and RNase H domains that catalyse the major steps in HBV replication (priming, DNA synthesis and removal of RNA templates) as well as a non-essential spacer domain present between the terminal protein
15 and polymerase domains (see for example Radziwill et al., 1990, J. Virol. 64:613; Bartenschlager et al., 1990, J. Virol. 64, 5324). The catalytic sites responsible for enzymatic activities have been characterized. In this regard, four residues forming the conserved YMDD motif (residues 538 to 541 numbered with respect to the 832 residue long polymerase) have been shown essential to the DNA- and RNA-dependent DNA
20 polymerase activity whereas RNase H activity is based on a DEDD motif involving four non-consecutive amino acid residues, respectively Asp (D) in position 689, Glu (E) in position 718, Asp (D) in position 737 and Asp (D) in position 777 as well as few other amino acid residues including Val (V) in position 769 and Thr (T) in position 776. Different mutations have been described in the art that abolish the RT polymerase and
25 RNase H activities (Chang et al., 1990, J. Virol. 64: 5553; Bartenschlager et al., 1990, J. Virol. 64, 5324, Radziwill et al., 1990, J. Virol. 64:613 and Chen et al., 1996, J. Virol. 70:6151). Several groups have succeeded in expressing HBV polymerase protein in various host system, but its expression has been reported toxic for the expressing cells, requiring the use of inducible promoters (Choi et al., 2002, Antiviral Res. 55:279;
30 Karimi et al., 2002, J. Virol. 76:8609).

A number of preclinical and clinical studies have emphasized the importance of CD4⁺ and CD8⁺ T cell immune responses for effective anti-viral response. It was indeed observed that patients naturally having recovered from hepatitis B mounted multi-specific and sustained responses mediated by T helper (T_H) and cytotoxic T (CTL) lymphocytes which are readily detectable in peripheral blood. Appearance of anti-HBe and anti-HBs antibodies indicates a favorable outcome of infection. HBsAg-specific antibodies are neutralizing, mediate protective immunity and persist for life after clinical recovery.

Chronic HBV infection is, however, only rarely resolved by the immune system. The vast majority of chronically infected patients show weak and temporary CD4 and CD8 T cell immune responses that are antigenically restricted and ineffective to clear viral infection. The reason for this alteration of the effector functions of the cellular immune response in chronic hepatitis B is currently not well-understood even if the involvement of different inhibitory molecules that are up-regulated in HBV chronically infected patients, such PD-1, CTLA4...etc, has been observed. Therefore, there is a need for immunomodulatory strategies capable of inducing an effective T-cell response.

Conventional treatment of chronic hepatitis B includes pegylated interferon-alpha (IFNa) and nucleoside/nucleotide analogues (NUCs) such as lamivudine, and more recently entecavir, telbivudine, adefovir and tenofovir. IFNa is a potent antiviral molecule, whereby inhibiting viral replication, which however, causes serious side effects in merely 25 - 30% of patients. NUCs act as competitive inhibitors of HBV polymerase aimed to inhibit the reverse transcription of the pre-genomic RNA into the negative DNA strand and then the double stranded viral DNA. They limit the formation of new virions, but are ineffective to eliminate the supercoiled cccDNA hidden in the nucleus of infected hepatocytes which constitutes a source of new progeny viruses. This can explain why NUC efficacy is temporary and viral rebound occurs immediately after cessation of treatment, requiring patients to stay lifelong under treatment. In addition, long-term efficacy is also limited due to emergence of resistant HBV mutants (more than 24% after one year and approximately 66% after four years of lamivudine treatment as reported in some studies although newer NUCs showed much fewer occurrences of drug-resistant HBV mutants). A number of HBV strains exhibiting a decreased sensitivity to anti-viral agents have now been isolated and genome

sequencing revealed high spot of substitution mutations in the polymerase domain, including in the YMDD motif (US2008-0233557; Zoulim and Locarnini, 2009, *Gastroenterology*, 137:1593).

Besides antiviral therapies, efforts are currently made to develop supplemental
5 therapies aiming at improved host's immune responses, specifically those mediated by cytotoxic T and helper T lymphocytes. Several encouraging vaccine strategies have focused on HBV surface proteins S, preS1 and/or preS2 (Zanetti et al., 2008, *Vaccine* 26: 6266; Mancini-Bourguine et al., 2006, *Vaccine* 24:4482) as well as on multivalent immunotherapy approaches aimed to simultaneously target multiple HBV antigens. For
10 example, immunization with a polyepitope DNA vaccine encoding multiple envelope, core and polymerase epitopes was shown to elicit CTL and T_H responses in preclinical mouse models (Depla et al., 2008, *J. Virol.* 82: 435). An approach based on a mixture of DNA plasmids encoding HBsAg, HBcAg and HBV polymerase (WO2005/056051; WO2008/020656) demonstrated specific anti-HBV cellular and humoral responses in
15 transgenic mouse model of chronic hepatitis B (Chae Young Kim et al., 2008, *Exp. Mol. Medicine* 40: 669). Phase I clinical trials were initiated in South Korea in HBV carriers in combination with lamivudine treatment (Yang et al., 2006, *Gene Ther.* 13: 1110). Another approach recently investigated involves the use of a vectored therapeutic vaccine encoding a combination of HBc and HBV polymerase together with Hbs
20 immunogenic domains (WO2011/01565). Mice immunized with Ad-vectorized vaccine showed T cell response against all expressed HBV antigens, especially against polymerase.

One may expect that HBV will continue to be a serious global health threat for
25 many years due to the chronic and persistent nature of the infection, its high prevalence, the continuing transmission of HBV and the significant morbidity of the associated diseases. Thus, there is an important need to develop more effective approaches for improving prevention and treatment of HBV infections or HBV-associated diseases or disorders. In particular, there still exists a need for approaches that conciliate T cell-
30 mediated immunity against the targeted HBV antigen(s), especially against core, and low potential toxicity. Such approaches are especially useful for treating subjects chronically infected with HBV.

This technical problem is solved by the provision of the embodiments as defined in the claims.

Other and further aspects, features and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention. These embodiments are given for the purpose of disclosure.

SUMMARY OF THE INVENTION

In one aspect, the present invention concerns a mutant polymerase polypeptide which comprises at least 500 amino acid residues of a native HBV polymerase wherein said mutant polymerase polypeptide comprises a polymerase domain with an internal deletion that functionally disrupts the polymerase activity and wherein said internal deletion includes at least the YMDD motif naturally present in the polymerase domain of a native polymerase.

The present invention also concerns a nucleic acid molecule encoding said mutant polymerase polypeptide, a vector comprising said nucleic acid molecule, or a composition comprising or encoding said mutant polymerase polypeptide.

The present invention also relates to the use of this mutant polymerase polypeptide, nucleic acid molecule, vector or composition, preferably in combination with additional polypeptides (e.g. with one or more HBV polypeptide(s)) for the purpose of treating, preventing or inhibiting an HBV infection or ameliorating a condition associated with an HBV infection.

Still a further aspect of the present invention includes a method of treating, preventing or inhibiting HBV infection or ameliorating a condition associated with HBV infection in a subject in need thereof, comprising providing or administering this mutant polymerase polypeptide, nucleic acid molecule, vector or composition, eventually in combination with additional polypeptides (e.g. with one or more HBV polypeptide(s)) and/or with the standard of care.

Still yet a further aspect of the present invention concerns a method of eliciting an immune response in a subject in need thereof, comprising providing or administering this mutant polymerase polypeptide, nucleic acid molecule, vector or composition, eventually in combination with additional polypeptides (e.g. with one or more HBV

polypeptide(s)) and/or with the standard of care, for the purpose of inducing or stimulating an immune response in this subject or for treating an HBV infection or ameliorating a condition or symptom associated with HBV infection.

Still more aspect of the present invention provides a kit of parts comprising a plurality of containers and instructions for providing or administering to a subject this mutant polymerase polypeptide, nucleic acid molecule, vector or composition, eventually in combination with additional polypeptides (e.g. with one or more HBV polypeptide(s)), in accordance with the compositions and methods described herein.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a mutant polymerase polypeptide which comprises a mutated polymerase domain with an internal deletion that functionally disrupts the polymerase activity, wherein said internal deletion comprises at least the YMDD motif naturally present in the polymerase domain of a native HBV polymerase. Such a mutant polymerase polypeptide or vector encoding it can be used in compositions and methods for the treatment or prevention of an HBV infection or a condition associated with an HBV infection, eventually in combination with other HBV polypeptides and/or standard of care. This invention permits to envisage expression and production of the mutant polymerase polypeptide in various vector systems due to the disruption of the associated enzymatic activities. The invention is also particularly adapted for human use and may be used to reinforce standard therapies (e.g. SOC). Immunization of animal models with a vector encoding this mutant polymerase polypeptide in fusion with HBc and Hbs immunogenic domains, elicited HBV specific T cell responses, and surprisingly a strong immunity against both HBc and Polymerase was observed

The following section provides a greater explanation of the meaning of some of the terms used herein.

Definitions

30 As used herein throughout the entire application, the terms "a" and "an" are used in the sense that they mean "at least one", "at least a first", "one or more" or "a plurality" of the referenced compounds or steps, unless the context dictates otherwise.

The term "and/or" wherever used herein includes the meaning of "and", "or" and "all or any other combination of the elements connected by said term".

The term "about" or "approximately" as used herein means within 10%, preferably within 8%, and more preferably within 5% of a given value or range.

- 5 The terms "amino acids", "residues" and "amino acid residues" are synonyms and encompass natural amino acids as well as amino acid analogs (e.g. non-natural, synthetic and modified amino acids, including D or L optical isomers).

 The terms "polypeptide", "peptide" and "protein" refer to polymers of amino acid residues which comprise at least nine or more amino acids bonded via peptide
10 bonds. The polymer can be linear, branched or cyclic and may comprise naturally occurring and/or amino acid analogs and it may be interrupted by non-amino acids. As a general indication, if the amino acid polymer is more than 50 amino acid residues, it is preferably referred to as a polypeptide or a protein whereas if it is 50 amino acids long or less, it is referred to as a "peptide".

- 15 As used herein, when used to define products, compositions and methods, the term "comprising" (and any form of comprising, such as "comprise" and "comprises"), "having" (and any form of having, such as "have" and "has"), "including" (and any form of including, such as "includes" and "include") or "containing" (and any form of containing, such as "contains" and "contain") are open-ended and do not exclude
20 additional, unrecited elements or method steps. Thus, a polypeptide "comprises" an amino acid sequence when the amino acid sequence might be part of the final amino acid sequence of the polypeptide. Such a polypeptide can have up to several hundred additional amino acids residues. "Consisting essentially of" means excluding other components or steps of any essential significance. Thus, a composition consisting
25 essentially of the recited components would not exclude trace contaminants and pharmaceutically acceptable carriers. A polypeptide "consists essentially of" an amino acid sequence when such an amino acid sequence is present with eventually only a few additional amino acid residues. "Consisting of" means excluding more than trace elements of other components or steps. For example, a polypeptide "consists of" an
30 amino acid sequence when the polypeptide does not contain any amino acids but the recited amino acid sequence.

As used herein, "HBV" and "hepatitis B virus" are used interchangeably and refer to any member of the hepadnaviridae (see e.g. Ganem and Schneider in Hepadnaviridae (2001) "The viruses and their replication", pp2923-2969, Knipe DM et al., eds. Fields Virology, 4th ed. Philadelphia, Lippincott Williams & Wilkins or
5 subsequent edition). Amino acid sequences of the various HBV polypeptides and the encoding nucleotide sequences can be found in specialized data banks (e.g. those mentioned above) and in the literature (see e.g. Valenzuela et al., 1980, The nucleotide sequence of the hepatitis B viral genome and the identification of the major viral genes (pp57-70) in "Animal Virus Genetics"; eds B. Fields, et al.; Academic Press Inc., New
10 York and Vaudin et al., 1988, J. Gen. Virol. 69: 1383).

As used herein, the term "HBV polymerase" refers to a polypeptide that retains at least 500 amino acid residues comprised in a native HBV polymerase protein. Desirably, such at least 500 amino acid residues are spread over the three functional domains and preferably over the four domains normally present in a native HBV
15 polymerase. This term encompasses native (i.e. naturally-occurring) polymerase polypeptides of any HBV strain, isolate or genotype that can be found, isolated, obtained from a source of HBV in nature such as those cited above in connection with the term "HBV" as well as modified polymerase (i.e. mutant polymerase polypeptide) and fragments thereof. For purpose of illustration, the amino acid residues for HBV
20 polymerase described herein are numbered by reference to a 832 amino acids long polymerase with the residue Tyr in the motif Tyr Met Asp Asp (YMDD) being residue number 538. It is within the reach of the skilled person to adapt the numeration of the amino acid residues to other polymerases (e.g. 843 or 845 amino acid long).

As used herein, the term "native" or "naturally-occurring" when used in
25 connection with any amino acid sequence (e.g. peptide, polypeptide, protein, etc) or nucleotide sequence (e.g. gene, nucleic acid molecule, polynucleotide, etc) refers to an amino acid sequence or to a nucleotide sequence that can be found, isolated, obtained from a source in nature as distinct from one being artificially modified or mutated by man in the laboratory (i.e. mutant). Such sources in nature include biological samples
30 (e.g. blood, plasma, sera, semen, saliva, tissue sections, biopsy specimen etc.) collected from an organism infected or that has been exposed to HBV, cultured cells (such as HepG2.2.15, HuH6-C15 (Sureau et al., 1986, Cell 47:37; Sells et al., 1987, Proc. Natl.

Acad. Sci. 84(4):1005); HuH7.TA61 or HuH7.TA62 (Sun et al., 2006, J Hepatol. 45(5):636), tissue cultures as well as recombinant materials. Recombinant materials include without limitation HBV isolates (e.g. available in depositary institutions), HBV genome, genomic RNA or cDNA libraries, vectors containing HBV genome or
5 fragment(s) thereof or any prior art vector known to include such elements).

For purpose of illustration, a “native HBV polymerase” means a HBV polymerase encoded by the ORF P of any naturally-occurring HBV genotype, strain or isolate described in the art (e.g. a polypeptide of 832 to 845 amino acids depending of the genotype) or fragment thereof. The term “native” also encompasses HBV
10 polymerase polypeptide/peptides that are representative of a specific genotype, and thus comprise an amino acid sequence corresponding to a consensus or near consensus sequence which is typically determined after sequence alignment of various HBV polymerases of a particular genotype.

The term "mutant" as used herein refers to a polypeptide exhibiting one or more
15 mutation(s) with respect to the native counterpart. For illustrative purposes, a “mutant polymerase polypeptide” refers to a polymerase polypeptide that originates from a native polymerase after being artificially mutated or altered by man in the laboratory as described herein. Any mutation(s) can be envisaged, including substitution, insertion and/or deletion of one or more nucleotide/amino acid residue(s), non-natural
20 arrangements (e.g. fusion with foreign polypeptides/peptides) as well as any combination of these possibilities. When several mutations are contemplated, they can concern consecutive residues and/or non-consecutive residues. Mutation(s) can be generated by a number of ways known to those skilled in the art, such as site-directed mutagenesis (e.g. using the SculptorTM *in vitro* mutagenesis system of Amersham, Les
25 Ullis, France), PCR mutagenesis, DNA shuffling and by chemical synthetic techniques (e.g. resulting in a synthetic nucleic acid molecule). According to preferred embodiments, the mutation(s) contemplated by the present invention encompass deletion(s) and/or substitution(s) of one or more amino acid residue(s) (consecutive or not) involved directly or indirectly in at least one enzymatic activity exhibited by a
30 native HBV polymerase, with the aim of disrupting said at least one enzymatic activity such as the polymerase activity and/or the RNaseH activity. In the context of the invention, the resulting mutant polymerase polypeptide globally retains a high degree of

identity (e.g. at least 80%) with the corresponding native HBV polymerase in the non-mutated portions.

The term “disrupt” as used herein in connection with a given enzymatic activity or any derivative such as “disrupting” means “abolish” (no residual activity at all) or
5 “significantly reduce” (residual activity of less than 20% of the activity exhibited by the native polymerase).

The term “identity” refers to an exact amino acid to amino acid or nucleotide to nucleotide correspondence between two polypeptide or nucleotide sequences. The percentage of identity between two sequences is a function of the number of identical
10 positions shared by the sequences, taking into account the number of gaps which need to be introduced for optimal alignment and the length of each gap. Various computer programs and mathematical algorithms are available in the art to determine the percentage of identity between amino acid sequences, such as for example the Blast program available at NCBI or ALIGN in Atlas of Protein Sequence and Structure
15 (Dayhoffed., 1981, Suppl., 3 482-489). Programs for determining homology between nucleotide sequences are also available in specialized data base (e.g. Genbank, the Wisconsin Sequence Analysis Package, BESTFIT, FASTA and GAP programs). For illustrative purposes, “at least 80% sequence identity” as used herein means 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,
20 97%, 98%, 99% or 100%.

As used herein, the term “isolated” refers to a protein, polypeptide, peptide, polynucleotide, plasmid vector, viral vector, or host cell that is removed from its natural environment (i.e. separated from at least one other component(s) with which it is naturally associated).

25

HBV sequence

A number of HBV sequences are suitable for use in the embodiments described herein including such sequences that are readily available to investigators in the field, including, but not limited to, HBV sequences described in Genbank and PubMed. For
30 illustrative purposes, extensive phylogenetic analyses have led to the classification of hepatitis B viruses into 8 major genotypes (A to H) which show distinct geographic distribution and clinical outcome although exhibiting high degree of sequence

conservation. The various HBV were also classified in nine different subtypes (*ayw1*, *ayw2*, *ayw3*, *ayw4*, *ayr*, *adw2*, *adw4*, *adrq+* and *adqr-*) in connection with HBsAg-associated serology (see review by Mamum-Al Mahtab et al., 2008, Hepatobiliary Pancrease Dis Int 5: 457; Schaeffer, 2007, World Gastroenterol. 7: 14). Each genotype and serotype encompasses different HBV strains and isolates. An isolate corresponds to a specific virus isolated from a particular source of HBV (e.g. a patient sample or other biological HBV reservoir) whereas a strain encompasses various isolates which are very close each other in terms of genomic sequences.

Exemplary HBV of genotype A include without limitation isolate HB-JI444AF and strain HB-JI444A (accession number AP007263). Exemplary HBV of genotype B include without limitation clone pJDW233 (accession number D00329), isolate HBV/14611 (accession number AF121243), HBV-B1 (GenBank accession number AF282917.1), HBV strain Whutj-37 (GenBank accession number AY2933309.1), the Chinese HBV strain GDH1 (GenBank accession number AY766463.1) and HBV isolate 57-1 subtype adw (GenBank accession number AY518556.1). Exemplary HBV of genotype C include without limitation isolate AH-1-ON980424 (accession number AB113879), strain HCC-3-TT (accession number AB113877), HBV isolate SWT3.3 (GenBank accession number EU916241.1), HBV isolate H85 (GenBank accession number AY306136.1), HBV strain C1248 (GenBank accession number DQ975272.1), HBV isolate CHN-H155 (GenBank accession number DQ478901.1) and HBV isolate GZ28-1 (GenBank accession number EF688062). Exemplary HBV of genotype D include without limitation isolates KAMCHATKA27 (accession number AB188243), ALTAY136 (accession number AB188245) and Y07587 (Genbank accession number Y07587 and Stoll-Becker et al., 1997, J. Virol. 71: 5399) as well as the HBV isolate described under accession number AB267090. Exemplary HBV of genotype E include without limitation isolate HB-JI411F and strain HB-JI411 (accession number AP007262). Exemplary HBV of genotype F include without limitation isolates HBV-BL597 (accession number AB214516) and HBV-BL592 (accession number AB166850). Exemplary HBV of genotype G include without limitation isolate HB-JI444GF and strain HB-JI444G (accession number AP007264). Exemplary HBV of genotype H include without limitation isolate HBV ST0404

(accession number AB298362) and isolate HB-JI260F and strain HB-JI260 (accession number AP007261).

It is intended that the present invention is not limited to these exemplary HBV sequences. Indeed the nucleotide and amino acid sequences of any or all of the HBV polypeptides/peptides used in accordance with the present invention can vary between different HBV isolates and genotypes and this natural genetic variation is included within the scope of the invention. Moreover, the HBV polypeptides/peptides in use in the invention can be representative of a specific genotype, and thus comprise an amino acid sequence corresponding to a consensus or near consensus sequence.

In addition, each of HBV polypeptides/peptides may originate independently from any HBV genotype, strain or isolate identified at present time, such as any of those described above in connection with the term "HBV". Such a configuration may permit to provide protection against a broader range of HBV genotypes or adaptation to a specific geographic region by using HBV genotype(s) that is/are endemic in this region or to a specific population of patients. In this regard, genotypes A and C are the most prevalent in the United States, genotypes A and D in Western European countries and genotype D in the Mediterranean basin whereas genotypes B and C are the most common in China. Limited data from India suggest that genotypes A and D are most prevalent in India. It is within the reach of the skilled person to choose appropriate HBV genotypes, serotypes, strains and/or isolates according to the population and/or geographic region to be treated by.

According to a preferred embodiment, the HBV polypeptides/peptides in use in the invention originate from a genotype D virus, with a specific preference for HBV isolate Y07587.

25

Mutant HBV polymerase

The mutant polymerase polypeptide of the invention comprises a mutated polymerase domain with an internal deletion that functionally disrupts the polymerase activity and includes at least the YMDD motif naturally present in the polymerase domain of a native polymerase. The disruption of the polymerase activity exhibited by the resulting mutant polymerase polypeptide can be evaluated using assays well known

in the art (e.g. the endogenous polymerase assays described in Radziwill et al., 1990, J Virol. 64:613).

A generic amino acid sequence encompassing the polymerase domain of native HBV polymerases of genotypes B, C and D is provided in SEQ ID NO: 1, with the
5 residue Xaa in position 7 being Thr (T) or Ala (A); the residue Xaa in position 13 being Asn (N), Arg (R) or His (H); the residue Xaa in position 16 being Ile (I) or Thr (T); the residue Xaa in position 38 being Thr (T) or Ala (A); the residue Xaa in position 53 being Ser (S) or Asn (N); the residue Xaa in position 54 being Thr (T) or Tyr (Y); the residue Xaa in position 55 being His (H) or Arg (R); the residue Xaa in position 91
10 being Ile (I) or Leu (L); the residue Xaa in position 109 being Pro (P) or Ser (S); the residue Xaa in position 118 being Thr (T) or Asn (N); the residue Xaa in position 121 being Asn (N) or Ile (I); the residue Xaa in position 122 being Ile (I) or Phe (F); the residue Xaa in position 124 being Tyr (Y) or Asn (N); the residue Xaa in position 127 being Gly (G) or Arg (R); the residue Xaa in position 131 being Asp (D) or Asn (N); the
15 residue Xaa in position 134 being Asp (D) or Asn (N); the residue Xaa in position 145 being Leu (L) or Met (M); the residue Xaa in position 149 being Lys (K) or Gln (Q); the residue Xaa in position 151 being Phe (F) or Tyr (Y); the residue Xaa in position 221 being Phe (F) or Tyr (Y); the residue Xaa in position 222 being Thr (T) or Ala (A); the residue Xaa in position 223 being Ser (S) or Ala (A); the residue Xaa in position 224
20 being Ile (I) or Val (V); the residue Xaa in position 238 being Asn (N) or His (H); the residue Xaa in position 248 being Asn (N) or His (H); the residue Xaa in position 256 being Ser (S) or Cys (C); the residue Xaa in position 257 being Trp (W) or Tyr (Y); the residue Xaa in position 259 being Thr (T) or Ser (S); the residue Xaa in position 263 being Glu (E) or Asp (D); the residue Xaa in position 266 being Val (V) or Ile (I); the
25 residue Xaa in position 267 being Leu (L) or Gln (Q); the residue Xaa in position 271 being Gln (Q), Met (M) or Glu (E); the residue Xaa in position 317 being Ser (S) or Ala (A); and the residue Xaa in position 332 being Cys (T) or Ser (S).

In accordance with the present invention, the mutated polymerase domain comprised in the mutant polymerase polypeptide of the invention lacks at least the
30 YMDD motif present from position 203 to position 206 of such generic polymerase domain of SEQ ID NO: 1.

The present invention also encompasses any other internal deletion of at least 4 amino acid residues and at most 30 amino acid residues which comprises at least this YMDD motif.

A representative mutant polymerase polypeptide according to the invention
5 comprises a mutated polymerase domain comprising the amino acid sequence shown in SEQ ID NO:1 but lacking at least the Tyr residue in position 203, the Met residue in position 204, the Asp residue in position 205 and the Asp residue in position 206.

Further to the YMDD motif, it is preferred that the internal deletion also encompasses all or parts of the neighboring VVL motif present at the C terminus of the
10 YMDD motif in a native HBV polymerase domain (corresponding to residues in positions 207-209 of SEQ ID NO: 1 and to residues in position 542-544 of a native polymerase of 832 amino acids). Such VVL motif can indeed contribute to the formation of “junctional” epitopes (e.g. colinearly synthesized new epitopes) which are at risk of reducing or silencing the host’s immune response directed against one or more
15 of the HBV polymerase-associated epitopes.

Preferably, the mutant polymerase polypeptide of the invention comprises a mutated polymerase domain having the amino acid sequence shown in SEQ ID NO: 1 but lacking at least the Tyr residue in position 203, the Met residue in position 204, the Asp residue in position 205, the Asp residue in position 206, the Val residue in position
20 207, the Val residue in position 208 and the Leu residue in position 209.

More preferably, the mutant polymerase polypeptide of the invention comprises a polymerase domain comprising, alternatively essentially consisting of, or alternatively consisting of an amino acid sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, preferably at least 90% of identity, more
25 preferably at least 95% of identity, and even more preferably 100% identity with the amino acid sequence shown in SEQ ID NO: 2. Even more preferably, the mutated polymerase domain comprises the amino acid sequence shown in SEQ ID NO: 2.

Alternatively or in combination, the mutant polymerase polypeptide of the
30 invention also comprises a mutated RNaseH domain comprising mutation(s) of one or more amino acid residue(s) that functionally disrupt the RNaseH activity normally exhibited by a native HBV polymerase.

As discussed above, the functional domain involved in RNase H activity has been mapped within the C-terminal portion, more particularly from position 680 to the C-terminus of a native 832 amino acid long HBV polymerase (or from position 693 to the C-terminus of a native 845 amino acid long HBV polymerase) and the present invention encompasses any mutation(s) in this domain that correlate with disruption of the RNase H activity (i.e. eventually leading to a weak residual activity less than 20% of the native RNaseH activity). Disruption of the RNase H activity exhibited by the mutant polymerase polypeptide can be evaluated using assays well known in the art (e.g. in vitro RNaseH activity assays or DNA-RNA tandem molecule analysis described in Radziwill et al., 1990, J Virol. 64:613 or in Lee et al., 1997, Biochem. Bioph. Res. Commun. 233(2):401).

A generic amino acid sequence encompassing the RNaseH domain of native HBV polymerases of genotypes B, C and D is provided in SEQ ID NO: 3, with the residue Xaa in position 2 being Ser (S) or Pro (P); the residue Xaa in position 19 being Ala (A) or Val (V); the residue Xaa in position 20 being Ile (I) or Met (M); the residue Xaa in position 30 being Val (V) or Leu (L); the residue Xaa in position 31 being Ala (A) or Ser (S); the residue Xaa in position 53 being Lys (K) or Asn (N); the residue Xaa in position 54 being Leu (L) or Ile (I); the residue Xaa in position 55 being Leu (L) or Ile (I); the residue Xaa in position 97 being Ala (A) or Thr (T); the residue Xaa in position 108 being Tyr (Y) or Ser (S); the residue Xaa in position 115 being Pro (P) or Leu (L); the residue Xaa in position 116 being Phe (F) or Tyr (Y); the residue Xaa in position 128 being Val (V) or Asp (D).

Advantageously, the one or more mutation(s) comprised in the RNaseH domain of the mutant polymerase polypeptide of the invention are selected from the group consisting of:

- a deletion of at least 8 amino acids and at most 60 amino acids including at least the portion of SEQ ID NO: 3 extending from approximately the Glu residue (E) in position 39 to approximately the Ala (A) residue in position 46 (del ELLAACFA);
- the substitution of the Asp (D) residue in position 10 of SEQ ID NO: 3 with an amino acid residue other than D;

- the substitution of the Val (V) residue in position 90 of SEQ ID NO: 3 with an amino acid residue other than V;
- the substitution of the Thr (T) or Ala (A) residue in position 97 of SEQ ID NO: 3 with an amino acid residue other than T or A;
- 5 - the substitution of the Asp (D) residue in position 98 of SEQ ID NO: 3 with an amino acid residue other than D; and
- any combination thereof.

Representative examples of appropriate combinations include without limitation (a) the substitutions of amino acid residues in position 10, 90, 97 and 98; (b) the deletion of 8 to 60 amino acid residues including the GLLAACFA motif and the substitutions of amino acid residues at any of the cited positions; or (c) the combination of all listed mutations.

Suitably, the substituted residue(s) in position 10, 90, 97 or 98 of SEQ ID NO: 3 are individually replaced with a His (H) residue or with a Tyr (Y) residue, with a specific preference for the residue in position 10 of SEQ ID NO: 3 substituted with a His (H) residue (D689H), the residue in position 90 of SEQ ID NO: 3 substituted with a Tyr (Y) residue (V769Y), the residue in position 97 of SEQ ID NO: 3 substituted with a Tyr (Y) residue (T776Y or A776Y) and/or the residue in position 98 of SEQ ID NO: 3 substituted with a His (H) residue (D777H).

Suitably, the deletion comprised in the mutated RNase H domain includes a portion of at least 19 amino acid residues extending from approximately the Glu residue (E) in position 39 to approximately the Thr (T) residue in position 57 of SEQ ID NO: 3, preferably a portion of at least 25 amino acids extending from approximately the Glu residue (E) in position 39 to approximately the Leu (L) residue in position 63 of SEQ ID NO: 3, and more preferably a portion of at least 33 amino acids extending from approximately the residue Xaa (A or S) in position 31 to approximately the Leu (L) residue in position 63 of SEQ ID NO: 3.

Preferably, the mutant polymerase polypeptide of the invention comprises a mutated RNaseH domain comprising the amino acid sequence shown in SEQ ID NO: 3 but (a) lacking the portion of 33 amino acid residues extending from the residue Xaa (X) in position 31 to approximately the Leu (L) residue in position 63 and comprising (b) the substitution of the Asp (D) residue in position 10 with a His (H) residue

(D689H); (c) the substitution of the Val (V) residue in position 90 with a Tyr (Y) residue (V769Y); (d) the substitution of the residue in position 97 with a Tyr (Y) residue (T/A776Y) and (e) the substitution of the Asp (D) residue in position 98 with a His (H) residue (D777H).

5 More preferably, the mutant polymerase polypeptide of the invention comprises a mutated RNaseH domain comprising, alternatively essentially consisting of, or alternatively consisting of an amino acid sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, preferably at least 90% of identity, more preferably at least 95% of identity, and even more preferably 100% identity with
10 the amino acid sequence shown in SEQ ID NO: 4.

In an even more preferred embodiment, the mutant polymerase polypeptide of the invention comprises, alternatively essentially consists of, or alternatively consists of an amino acid sequence which exhibits at least 80% of identity, advantageously at least
15 85% of identity, preferably at least 90% of identity, more preferably at least 95% of identity, and even more preferably 100% identity with the amino acid sequence shown in SEQ ID NO: 5. Still even more preferred is a mutant polymerase polypeptide comprising the amino acid sequence shown in SEQ ID NO: 5.

20 In the context of the invention, the mutant polymerase polypeptide of the invention can comprise additional mutation(s). However, it is preferred to avoid modification(s) that can be detrimental to the immunogenic activity, especially in portions rich in B, CTL and/or T_H epitopes.

Exemplary additional modifications include N-terminal truncation. Particularly
25 appropriate is truncation of at least 20 amino acid residues and at most 100 amino acid residues normally present at the N-terminus of a native HBV polymerase or of SEQ ID NO: 5, with a specific preference for a truncation extending from position 1 (Met initiator) or 2 to approximately position 47 of SEQ ID NO: 5. This modification is particularly relevant for mutant polymerase polypeptide used in combination with a
30 native HBV core polypeptide due to the fact that such a N-terminal truncation contributes to reduce or delete the overlapping portions between these two polypeptides. However, the same can be achieved by using a non-truncated mutant polymerase

polypeptide in combination with a C-terminal truncated HBV core polypeptide as described below.

Desirably, the resulting mutant polymerase polypeptide retains immunogenic properties, in particular a capacity to stimulate a cell-mediated immune response, within
5 the same range as or alternatively higher than the native polymerase.

Combination/fusion with other polypeptide(s)

In another embodiment the mutant polymerase polypeptide of the invention can be used in combination with one or more additional polypeptide(s) or peptide(s).

10 The term “combination” and variation such as “combined use” refers to the action of administering in the same host organism two or more entities, one of which being an object of the invention. Typically, the at least two entities can be administered by different routes and according to different time schedule. A suitable combination includes without limitation the combination of the mutant polymerase polypeptide
15 described herein (or a vector encoding it) with antiviral(s) (SOC) and/or with additional polypeptide(s) or vector(s) encoding such additional polypeptide(s). Such a combination can be in the form of (a) a mixture (e.g. mixture of two or more polypeptides or vectors), (b) a fusion between the two or more entities or (c) through specific expression design (e.g. bicistronic or independent expression). For example, the two or more
20 entities may be expressed independently in the same vector using distinct regulatory elements (e.g. distinct promoter and termination sequences). The independent expression design is particularly adapted for expression from plasmid or measles vectors. Alternatively, the two or more entities may be expressed in a bicistronic manner under the control of the same promoter and termination sequences but requiring the use of
25 additional regulatory elements such as IRES (for internal ribosome entry site) permitting translation of two or more cistrons from the same mRNA. A large choice of IRES is available in the art such as those originating from the poliovirus, hepatitis C virus and encephalomyocarditis (EMCV) viruses (e.g. see WO95/24485). The bicistronic design is particularly adapted for expression from vectors with more limited
30 cloning capacity such as adenovirus vector.

The term "fusion" or "fusion protein" as used herein refers to the combination with two or more polypeptides/peptides in a single polypeptide chain. Preferably, the fusion is performed by genetic means, i.e. by fusing in frame the nucleotide sequences encoding each of said polypeptides/peptides. By "fused in frame", it is meant that the expression of the fused coding sequences results in a single protein without any translational terminator between each of the fused polypeptides/peptides. The fusion can be direct (i.e. without any additional amino acid residues in between) or through a linker (e.g. 3 to 30 amino acids long peptide composed of repeats of amino acid residues such as glycine, serine, threonine, asparagine, alanine and/or proline).

Additional polypeptide(s) or peptide(s) for use in combination with the mutant polymerase polypeptide of the invention is/are preferably a polypeptide or peptide encoded by an HBV genome, such as any native HBV polypeptide(s), modified derivative(s) and/or fragment(s) thereof. Representative examples of such HBV polypeptides or peptides include without limitation HBc (core), HBs, X protein and any immunogenic fragment thereof.

In accordance with the invention, as mentioned above, any of the additional HBV polypeptide(s) or peptide(s) in use in the invention can originate from different or from the same HBV genotype, strain or isolate as the HBV genotype, strain or isolate from which originates the mutant polymerase polypeptide of the invention. Preferably, such additional HBV polypeptide(s) originate(s) from a genotype D HBV, and especially from the Y07587 isolate.

A preferred combination is in the form of a fusion protein between the mutant polymerase polypeptide described herein and the additional polypeptide(s). Such a fusion is preferably direct without any linker between the fused entities.

25

Combination with core

As used herein, the term "core polypeptide" refers to a polypeptide that retains at least 100 amino acid residues comprised in a native HBV core (HBc) protein. This term encompasses native (i.e. naturally-occurring) core polypeptides of any HBV strain, isolate or genotype that can be found, isolated, obtained from a source of HBV in nature such as those cited above in connection with the term "HBV" as well as modified core and fragments thereof.

The HBV core polypeptide in use in the invention can originate from an HBV virus having the same as or a different genotype than the one from which originates the mutant polymerase polypeptide. Preferably, they both originate from a genotype D virus and more particularly from the Y07587 isolate. Core polypeptides and their encoding
5 sequences can be generated by a number of ways known to those skilled in the art, such as by chemical synthesis of the encoding sequence (e.g. resulting in a synthetic nucleic acid molecule) or by recombinant means (e.g. site-directed mutagenesis of the corresponding nucleotide sequence, PCR mutagenesis, DNA shuffling).

Any modification(s) can be envisaged, provided that the resulting core retains a
10 significant immunogenic activity when combined or fused with the mutant polymerase polypeptide described herein, preferably in the same range as or higher than the native core counterpart.

Suitable modifications include truncation of at least 10 amino acid residues and at most 41 amino acid residues normally present at the C-terminus of a native core
15 polypeptide or within the C-terminal part thereof, with a special preference for a truncation extending from residue 143, 144, 145, 146, 147, 148 or residue 149 to the C-terminus (residue 183) of the native core polypeptide. Other suitable modifications include internal deletion of one or more amino acid residues, especially in the surface-exposed region(s) such as the region located in the vicinity of residue 80 that is
20 predicted to form an outer loop (Argos et al. 1988, EMBO J. 7: 819; Borisova et al., 1993, J. Virol. 67: 3696; Schodel et al., 1992, J. Virol. 66: 106 ; Yon et al., 1992, J. Gen Virol. 73: 2569; and Pumpens et al., 1995, Intervirology 38: 63).

In a preferred embodiment, the combination is in the form of a fusion.
25 Accordingly, the invention relates to a fusion protein comprising the mutant polymerase polypeptide described herein and a fusion partner. Preferably, the fusion partner is an HBV core polypeptide, with a specific preference for a core polypeptide which is C-terminally truncated and especially truncated at residue 148.

Preferably, the HBV core polypeptide is fused in frame to the N-terminus of the
30 mutant polymerase polypeptide described herein, resulting in a fusion protein starting with an initiator Met, the core polypeptide (modified or native) without any stop codon, the mutant polymerase polypeptide (without any Met initiator) and a stop codon.

A preferred fusion protein comprises, alternatively essentially consists of, or alternatively consists of an amino acid sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, preferably at least 90% of identity, more preferably at least 95% of identity, and even more preferably 100% identity with the amino acid sequence shown in SEQ ID NO: 6. More preferably, the fusion protein of the invention comprises the amino acid sequence shown in SEQ ID NO: 6.

Combination with immunogenic HbsAg domains

Alternatively or in combination with the previous embodiment (combination with a core polypeptide), the mutant polymerase polypeptide of the invention can be used in combination with an HbsAg or immunogenic fragment(s)/domain(s) thereof.

As used herein, the term “immunogenic domain” refers to a polypeptide having from approximately 15 to approximately 100 amino acid residues, and preferably at least 20 and at most 60 consecutive amino acids comprising at least one B and/or T cell epitope specific for T helper (T_H) cells and/or for cytotoxic T (CTL) cells normally present in a native HBsAg protein. Moreover such epitope(s) can be restricted to various MHC class I and/or class II antigens (e.g. A2, A24, DR, DP, etc). Preferably, the one or more HBsAg immunogenic domains used in the invention do not include any portions of HBV preS1 and preS2 polypeptides.

Each of the one or more immunogenic domain(s) can independently originate from the same or different HBV virus(es) which can be the same or different with respect to HBV virus from which originates the mutant polymerase polypeptide described herein (and eventually the core polypeptide). Preferably, the one or more immunogenic domains originate from a genotype D HBV, and especially from the Y07587 isolate.

Exemplary immunogenic domains that can be used in the invention are described in the art (e.g. WO93/03764; WO94/19011; Desombere et al., 2000, Clin. Exp. Immunol 122: 390; Loirat et al., 2000, J. Immunol. 165: 4748; Schirmbeck et al., 2002, J. Immunol 168: 6253; Depla et al., 2008, J. Virol. 82: 435 and WO2011/015656). Particularly preferred immunogenic domains include the env1 and env2 domains described in WO2011/015656. “Env1” corresponds to the portion of a

native HBsAg from approximately position 14 to approximately position 51 and env2 to the HBsAg portion from approximately position 165 to approximately position 194.

In a preferred embodiment, the combination is in the form of a fusion and the invention relates to a fusion protein comprising the mutant polymerase polypeptide 5 described herein and one or more HBsAg immunogenic domain(s) or to the fusion protein defined above (comprising at least the mutant polymerase polypeptide described herein and an Hbc polypeptide) further comprising one or more HBsAg immunogenic domain(s). The one or more immunogenic domains can be positioned in the fusion protein at the N-terminus, at the C-terminus and/or internally, e.g. within the mutant 10 polymerase polypeptide (for example in place of the portion lacking in the mutated polymerase and/or RNaseH domains) or in between the core and the mutant polymerase polypeptide. It is within the reach of the skilled person to define accordingly the need and location of the translation-mediating regulatory elements (e.g. the initiator Met and codon STOP at the N- and C-termini of the fusion protein).

15 Fusion proteins of particular interest comprise the mutant polymerase polypeptide described herein, the core polypeptide and two HBsAg immunogenic domains, with a specific preference for a fusion comprising at its N-terminus a core polypeptide (e.g. native 183 residues or truncated with 148 residues with an initiator Met) fused to the mutant polymerase polypeptide (without initiator Met) comprising one 20 or two HbsAg immunogenic domains fused in place of the internal deletion in the mutated polymerase domain (e.g. env1) and/or in place of the deletion in the mutated RNaseH domain (e.g. env2).

In a preferred aspect of this embodiment, the fusion protein of the invention comprises, alternatively essentially consists of, or alternatively consists of an amino 25 acid sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, preferably at least 90% of identity, more preferably at least 95% of identity, and even more preferably 100% identity with any of the amino acid sequence shown in SEQ ID NO: 7-9. A particularly preferred embodiment is directed to a fusion protein comprising the amino acid sequence shown in SEQ ID NO: 8.

In the context of the invention, the mutant polymerase polypeptide of the invention or the fusion protein of the invention may further comprise additional structural features.

In one embodiment, it can comprise additional compound(s) (e.g. peptide or
5 polypeptide) aimed to improve its immunogenic activity in a host organism. Such compounds capable of enhancing immunogenicity have been described in the literature and include, without limitation, calreticulin (Cheng et al., 2001, J. Clin. Invest. 108: 669), Mycobacterium tuberculosis heat shock protein 70 (HSP70) (Chen et al., 2000, Cancer Res. 60: 1035), ubiquitin (Rodriguez et al., 1997, J. Virol. 71: 8497), bacterial
10 toxin such as the translocation domain of *Pseudomonas aeruginosa* exotoxin A (ETA(dIII)) (Hung et al., 2001 Cancer Res. 61: 3698) as well as T helper epitope(s) such as Pan-Dr peptide (Sidney et al., 1994, Immunity 1: 751), pstS1 GCG epitope (Vordermeier et al., 1992, Eur. J. Immunol. 22: 2631), tetanus toxoid peptides P2TT (Panina-Bordignon et al., 1989, Eur. J. Immunol. 19: 2237) and P30TT (Demotz et al.,
15 1993, Eur. J. Immunol. 23: 425), influenza epitope (Lamb et al., 1982, Nature 300: 66) and hemagglutinin epitope (Rothbard et al., 1989, Int. Immunol. 1: 479).

Other suitable structural features are those which are beneficial to the synthesis, processing, stability and solubility of the mutant polymerase polypeptide or fusion protein of the invention (e.g. those aimed to modify potential cleavage sites, potential
20 glycosylation sites and/or membrane anchorage so as to improve presentation to the cell membrane).

It could be beneficial for immune response to direct the synthesis of the mutant polymerase polypeptide or fusion protein described herein at the cell surface by using appropriate sequences well known in the art such as signal and/or trans-membrane
25 peptides. Briefly, signal peptides are generally present at the N-terminus of membrane-presented or secreted polypeptides and initiate their passage into the endoplasmic reticulum (ER). They usually comprise 15 to 35 essentially hydrophobic amino acids which are then removed by a specific ER-located endopeptidase to give the mature polypeptide. Trans-membrane peptides are also highly hydrophobic in nature and serve
30 to anchor the polypeptides within cell membrane. The choice of the trans-membrane and/or signal peptides which can be used in the context of the present invention is vast. They may be obtained from any membrane-anchored and/or secreted polypeptide (e.g.

cellular or viral polypeptides) such as those of immunoglobulins, tissue plasminogen activator, insulin, rabies glycoprotein, the HIV virus envelope glycoprotein or the measles virus F protein or may be synthetic.

In one embodiment, the mutant polymerase polypeptide or fusion protein of the invention is fused in frame to a signal peptide which is inserted at the N-terminus downstream of the codon for initiation of translation. In another embodiment, the mutant polymerase polypeptide or fusion protein of the invention is fused in frame to a signal peptide (e.g. inserted at its N-terminus) and to a trans-membrane peptide (e.g. inserted at the C-terminus, for example immediately upstream of the stop codon). Preferably, the signal and trans-membrane peptides employed in the context of the invention originate from the rabies glycoprotein (see e.g; WO99/03885 or WO2008/138649). Preferred embodiments are directed to a HBV polymerase mutant polypeptide and a fusion protein comprising, alternatively essentially consist of, or alternatively consists of an amino acid sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, particularly at least 90% of identity, preferably at least 95% of identity and more preferably 100% identity with the amino acid sequence shown in SEQ ID NO: 10, SEQ ID NO: 11 or SEQ ID NO: 12.

Nucleic acid molecule

In another aspect, the present invention provides isolated nucleic acid molecules encoding mutant polymerase polypeptides and fusion proteins described herein.

Within the context of the present invention, the terms “nucleic acid”, “nucleic acid molecule”, “polynucleotide” and “nucleotide sequence” are used interchangeably and define a polymer of any length of either polydeoxyribonucleotides (DNA) (e.g., cDNA, genomic DNA, plasmids, vectors, viral genomes, isolated DNA, probes, primers and any mixture thereof) or polyribonucleotides (RNA) (e.g., mRNA, antisense RNA) or mixed polyribo-polydeoxyribonucleotides. They encompass single or double-stranded, linear or circular, natural or synthetic polynucleotides. Moreover, a polynucleotide may comprise non-naturally occurring nucleotides and may be interrupted by non-nucleotide components.

The nucleic acid molecules of the present invention can be generated from any source using sequence data accessible in the art and the sequence information provided herein. For example, the DNA sequence coding for the HBV polymerase and if needed core polypeptide and HBsAg immunogenic domains can be isolated independently from
5 HBV-containing cells, cDNA and genomic libraries, viral genomes or any prior art vector known to include it, and then suitably linked together by conventional molecular biology or PCR techniques. Alternatively, the nucleic acid molecules of the invention can also be generated by chemical synthesis in automatized process (e.g. assembled from overlapping synthetic oligonucleotides or synthetic gene). Modification(s) can be
10 generated by a number of ways known to those skilled in the art, such as chemical synthesis, site-directed mutagenesis, PCR mutagenesis, DNA shuffling, etc.

Of particular interest is any of a nucleic acid molecule selected from the group consisting of:

- A nucleic acid molecule which encodes a mutant polymerase polypeptide
15 comprising a polymerase domain having the amino acid sequence shown in SEQ ID NO: 1 or 2;
- A nucleic acid molecule which encodes a mutant polymerase polypeptide comprising a RNaseH domain having the amino acid sequence shown in SEQ ID NO: 3 or 4;
- 20 - A nucleic acid molecule which encodes a mutant polymerase polypeptide comprising an amino acid sequence which exhibits at least 80% of identity (e.g. 80%, 85%, 90%, 95%, 97%, 100%) with the amino acid sequence shown in SEQ ID NO: 5; or
- A nucleic acid molecule which encodes a fusion protein comprising an amino
25 acid sequence which exhibits at least 80% of identity (e.g. 80%, 85%, 90%, 95%, 97%, 100%) with the amino acid sequence shown in any of SEQ ID NO: 6-12.

The present invention is not limited to these exemplary nucleotide sequences and encompasses any modifications aimed to improve cloning, expression, stability of the
30 nucleic acid molecules in use in the invention (e.g. introduction of appropriate restriction sites degeneration and/or optimisation of nucleotide sequence to optimize translation in a given host cell and/or suppression of potentially negative elements that

may destabilize the nucleic acid molecule or its transcript). When several modifications are contemplated, they can concern consecutive residues and/or non-consecutive residues. The modification(s) contemplated by the present invention encompass silent modifications that do not change the amino acid sequence of the encoded polypeptides
5 and fusion proteins, as well as modifications that are translated into the encoded polypeptides and fusion proteins.

In one embodiment, the nucleic acid molecule of the invention can be degenerated over the full length nucleotide sequence or portion(s) thereof so as to reduce sequence homology between nucleic acid molecule(s) used in the context of the
10 invention or in the host cell. It is indeed advisable to degenerate the portions of nucleic acid sequences that show a high degree of nucleotide sequence identity and the skilled person is capable of identifying such portions by sequence alignment. For example if a vector carries a nucleic acid molecule encoding a mutant polymerase polypeptide as described herein and a nucleic acid molecule encoding another HBV polypeptide
15 encoded by overlapping sequences in the HBV genome, it may be advantageous to degenerate one or both nucleic acid molecule(s) in the overlapping portions so as to avoid homologous recombination problems during production process.

Alternatively or in combination, the nucleic acid molecule of the invention can be optimized for providing high level expression in a particular host cell or organism. It
20 has been indeed observed that, when more than one codon is available to code for a given amino acid, the codon usage patterns of organisms are highly non-random and the utilisation of codons may be markedly different between different hosts. As the nucleotide sequences encompassed by the invention are mostly of viral origin (HBV), they may have an inappropriate codon usage pattern for efficient expression in host cells
25 such as bacterial, lower or higher eukaryotic cells. Typically, codon optimisation can be performed by replacing one or more "native" (e.g. HBV) codon corresponding to a codon infrequently used in the host cell/organism of interest by one or more codon encoding the same amino acid which is more frequently used in the host cell/organism of interest. It is not necessary to replace all native codons corresponding to infrequently
30 used codons since increased expression can be achieved even with partial replacement. Moreover, some deviations from strict adherence to optimised codon usage may be

made to accommodate the introduction of restriction site(s) into the resulting nucleic acid molecule.

Further, expression in the host cell or organism can be improved through additional modifications of the nucleotide sequence aimed to prevent clustering of rare, non-optimal codons and/or to suppress or modify at least partially negative sequence elements which are expected to negatively influence expression levels (e.g. AT-rich or GC-rich sequence stretches; unstable direct or inverted repeat sequences; RNA secondary structures; and/or internal cryptic regulatory elements such as internal TATA-boxes, chi-sites, ribosome entry sites, and/or splicing donor/acceptor sites).

10 A particularly preferred embodiment of the present invention is directed to a nucleic acid molecule comprising, alternatively essentially consisting of or alternatively consisting of a nucleotide sequence which exhibits at least 80% of identity, advantageously at least 85% of identity, preferably at least 90% of identity, more preferably at least 95% of identity, and even more preferably 100% identity with the
15 nucleotide sequence shown in any of SEQ ID NO: 13 to 17.

Another embodiment of the invention pertains to fragments of the nucleic acid molecules of the invention, e.g. restriction endonuclease and PCR-generated fragments. Such fragments can be used as probes, primers or fragments encoding an immunogenic portion of the encoded immunogenic polypeptide.

20

Vectors

In another aspect, the present invention provides vectors comprising a nucleic acid molecule of the present invention.

The term "vector" as used herein refers to a vehicle, preferably a nucleic acid molecule or a viral particle that contains the elements necessary to allow delivery, propagation and/or expression of one or more nucleic acid molecule(s) within a host cell or organism. This term encompasses vectors for maintenance (cloning vectors) or vectors for expression in various host cells or organisms (expression vectors), extrachromosomal vectors (e.g. multicopy plasmids) or integration vectors (e.g. designed to integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates) as well as shuttle vectors (e.g. functioning in both prokaryotic and/or eukaryotic hosts) and transfer vectors (e.g. for
25
30

transferring nucleic acid molecule(s) in a viral genome). For the purpose of the invention, the vectors may be of naturally occurring genetic sources, synthetic or artificial, or some combination of natural and artificial genetic elements.

In the context of the invention, the term “vector” has to be understood broadly as including plasmid and viral vectors. A "plasmid vector" as used herein refers to a replicable DNA construct. Usually plasmid vectors contain selectable marker genes that allow host cells carrying the plasmid vector to be selected for or against in the presence of a corresponding selective drug. A variety of positive and negative selectable marker genes are known in the art. By way of illustration, an antibiotic resistance gene can be used as a positive selectable marker gene that allows a host cell to be selected in the presence of the corresponding antibiotic.

The term "viral vector" as used herein refers to a nucleic acid vector that includes at least one element of a virus genome and may be packaged into a viral particle or to a viral particle. The terms “virus”, “virions”, viral particles” and “viral vector particle” are used interchangeably to refer to viral particles that are formed when the nucleic acid vector is transduced into an appropriate cell or cell line according to suitable conditions allowing the generation of infectious viral particles. In the context of the present invention, the term “viral vector” has to be understood broadly as including nucleic acid vector (e.g. DNA viral vector) as well as viral particles generated thereof. The term “infectious” refers to the ability of a viral vector to infect and enter into a host cell or organism. Viral vectors can be replication-competent or -selective (e.g. engineered to replicate better or selectively in specific host cells), or can be genetically disabled so as to be replication-defective or replication-impaired.

Vectors which are appropriate in the context of the present invention, include, without limitation, bacteriophage, plasmid or cosmid vectors for expression in prokaryotic host cells such as bacteria (e.g. *E. coli*, *Bacillus subtilis* or *Listeria*); vectors for expression in yeast (e.g. *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris*); baculovirus vectors for expression in insect cell systems (e.g. Sf 9 cells); viral and plasmid vectors for expression in plant cell systems (e.g. Ti plasmid, cauliflower mosaic virus CaMV; tobacco mosaic virus TMV); as well as viral and plasmid vectors for expression in higher eukaryotic cells or organisms.

Typically, such vectors are commercially available (e.g. in Invitrogen, Stratagene, Amersham Biosciences, Promega, etc.) or available from depositary institutions such as the American Type Culture Collection (ATCC, Rockville, Md.) or have been the subject of numerous publications describing their sequence, organization
5 and methods of producing, allowing the artisan to apply them.

Representative examples of suitable plasmid vectors include, without limitation, pREP4, pCEP4 (Invitrogen), pCI (Promega), pVAX (Invitrogen) and pgWiz (Gene Therapy System Inc).

Representative examples of suitable viral vectors are generated from a variety of
10 different viruses (e.g. retrovirus, adenovirus, adenovirus-associated virus (AAV), poxvirus, herpes virus, measles virus, foamy virus, alphavirus, vesicular stomatitis virus, etc). As described above, the term "viral vector" encompasses vector DNA, genomic DNA as well as viral particles generated thereof.

The present invention also encompasses vectors (e.g. plasmid DNA) complexed
15 to lipids or polymers to form particulate structures such as liposomes, lipoplexes or nanoparticles.

In one embodiment, the vector of the invention is an adenoviral vector. It can be derived from a variety of human or animal adenoviruses (e.g. canine, ovine, simian, etc). Any serotype can be employed. Desirably, the adenoviral vector is replication-
20 defective and originates from a human Ad, and more particularly from a human Ad of a rare serotype, or from a chimpanzee Ad. Representative examples of human adenoviruses include subgenus C Ad2 Ad5 and Ad6, subgenus B Ad11, Ad34 and Ad35 and subgenus D Ad19, Ad24, Ad48 and Ad49. Representative examples of chimp Ad include without limitation AdCh3 (Peruzzi et al., 2009, Vaccine 27: 1293), AdCh63
25 (Dudareva et al., 2009, vaccine 27: 3501) and any of those described in the art (see for example WO03/000283; WO03/046124; WO2005/071093; WO2009/073103; WO2009/073104; WO2009/105084; WO2009/136977 and WO2010/086189).

Replication-defective adenoviral vectors can be obtained as described in the art, e.g by deletion of at least a region of the adenoviral genome or portion thereof essential
30 to the viral replication, with a specific preference for deletion of E1 region comprising E1 coding sequences (e.g. extending from approximately positions 459 to 3510 by reference to the sequence of the human adenovirus type 5 disclosed in the GeneBank

under the accession number M 73260 and in Chroboczek et al., 1992, Virol. 186:280). The present invention also encompasses vectors having additional deletion(s)/modification(s) within the adenoviral genome (all or part of the non-essential E3 region or of other essential E2, E4 regions as described in WO94/28152; 5 Lusky et al., 1998, J. Virol 72: 2022).

The nucleic acid molecule of the present invention can be inserted in any location of the adenoviral genome, and may be positioned in sense or antisense orientation relative to the natural transcriptional direction of the region in question. Preferably, the nucleic acid molecule of the invention is inserted in replacement of the 10 adenoviral E1 region and placed under the control the CMV promoter.

Other viral vectors suitable in the context of the invention are poxviral vectors which can be obtained from any member of the poxviridae with a specific preference for a poxviral vector originating from a canarypox, fowlpox or vaccinia virus, the latter being preferred. Suitable vaccinia viruses include without limitation the Copenhagen 15 strain (Goebel et al., 1990, Virol. 179: 247; Johnson et al., 1993, Virol. 196: 381), the Wyeth strain and particularly the modified Ankara (MVA) strain (Antoine et al., 1998, Virol. 244: 365). The general conditions for constructing recombinant poxvirus are well known in the art. The nucleic acid molecule of the present invention is preferably inserted within the poxviral genome in a non-essential locus. Thymidine kinase gene is 20 particularly appropriate for insertion in Copenhagen vaccinia vectors and deletion II or III for insertion in MVA vector. Preferably, the nucleic acid molecule of the invention is inserted in deletion III of the MVA vector and placed under the control the vaccinia 7.5K or pH5R promoter.

Other viral vectors suitable in the context of the invention are morbillivirus 25 which can be obtained from the paramyxoviridae family, with a specific preference for measles virus. Various attenuated strains are available in the art (Brandler et al, 2008, CIMID, 31: 271; Singh et al., 1999, J. virol. 73(6): 4823), such as and without limitation, the Edmonston A and B strains (Griffin et al., 2001, Field's in Virology, 1401-1441), the Schwartz strain (Schwarz A, 1962, Am J Dis Child, 103: 216), the S- 30 191 or C-47 strains (Zhang et al., 2009, J Med Virol. 81 (8): 1477). Insertion between P and M genes is particularly appropriate.

In accordance with the present invention, the nucleic acid molecule(s) comprised in the vector of the invention is in a form suitable for expression in a host cell or organism, which means that the nucleic acid molecule is placed under the control of appropriate regulatory sequences. As used herein, the term “regulatory elements” refers
5 to any element that allows, contributes or modulates the expression of a nucleic acid molecule in a given host cell or organism, including replication, duplication, transcription, splicing, translation, stability and/or transport of the nucleic acid or its derivative (i.e. mRNA).

It will be appreciated by those skilled in the art that the choice of the regulatory
10 sequences can depend on such factors as the vector itself, the host cell, the level of expression desired, etc. The promoter is of special importance. In the context of the invention, it can be constitutive directing expression of the nucleic acid molecule in many types of host cells or specific to certain host cells (e.g. liver-specific regulatory sequences) or regulated in response to specific events or exogenous factors (e.g. by
15 temperature, nutrient additive, hormone, etc) or according to the phase of a viral cycle (e.g. late or early). One may also use promoters that are repressed during the production step in response to specific events or exogenous factors, in order to optimize vector production and circumvent potential toxicity of the expressed polypeptide(s).

Promoters suitable for constitutive expression in mammalian cells include but
20 are not limited to the cytomegalovirus (CMV) immediate early promoter (Boshart et al., 1985, Cell 41: 521), the RSV promoter, the adenovirus major late promoter, the phosphoglycerate kinase (PGK) promoter (Adra et al., 1987, Gene 60: 65), the thymidine kinase (TK) promoter of herpes simplex virus (HSV)-1 and the T7 polymerase promoter. Vaccinia virus promoters are particularly adapted for expression in poxviral
25 vectors. Representative examples include without limitation the vaccinia 7.5K, H5R, 11K7.5 (Erbs et al., 2008, Cancer Gene Ther. 15: 18), TK, p28, p11 and K1L promoter, as well as synthetic promoters such as those described in Chakrabarti et al. (1997, Biotechniques 23: 1094), Hammond et al. (1997, J. Virological Methods 66: 135) and Kumar and Boyle (1990, Virology 179: 151) as well as early/late chimeric promoters.
30 Promoters suitable for measles-mediated expression include without limitation any promoter directing expression of measles transcription units (Brandler and Tangy, 2008, CID 31: 271). Liver-specific promoters include without limitation those of HMG-

CoA reductase (Luskey, 1987, Mol. Cell. Biol. 7: 1881); sterol regulatory element 1 (SRE-1; Smith et al., 1990, J. Biol. Chem. 265: 2306); albumin (Pinkert et al., 1987, Genes Dev. 1: 268); phosphoenol pyruvate carboxy kinase (PEPCK) (Eisenberger et al., 1992, Mol. Cell Biol. 12: 1396); alpha-1 antitrypsin (Ciliberto et al., 1985, Cell 41: 531); human transferrin (Mendelzon et al., 1990, Nucleic Acids Res. 18: 5717); and FIX (US 5,814,716) genes.

Those skilled in the art will appreciate that the regulatory elements controlling the expression of the nucleic acid molecule of the invention may further comprise additional elements for proper initiation, regulation and/or termination of transcription (e.g. polyA transcription termination sequences), mRNA transport (e.g. nuclear localization signal sequences), processing (e.g. splicing signals), and stability (e.g. introns and non-coding 5' and 3' sequences), translation (e.g. an initiator Met, tripartite leader sequences, IRES ribosome binding sites, Shine-Dalgarno sequences, etc.) into the host cell or organism and purification steps (e.g. a tag).

Particularly preferred embodiments of the invention are directed to vectors (or viral particles) selected from the group consisting of:

- A defective Ad vector comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter, and encoding a mutant polymerase polypeptide comprising an amino acid sequence as shown in SEQ ID NO: 5 or a fusion protein comprising an amino acid sequence as shown in SEQ ID NO: 6 or SEQ ID NO: 8;
- A defective Ad vector comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter, and comprising the nucleotide sequence shown in SEQ ID NO: 13, SEQ ID NO: 14 or SEQ ID NO: 15;
- A replication-defective Ad vector especially a defective AdCh3 comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter and comprising the nucleotide sequence shown in SEQ ID NO: 16 or SEQ ID NO: 17;
- A MVA vector comprising a nucleic acid molecule placed under the control of a vaccinia promoter such as the 7.5K or pH5R promoter, and encoding a mutant polymerase polypeptide comprising the amino acid sequence shown in SEQ ID

- NO: 5 or SEQ ID NO: 10 or a fusion protein comprising the amino acid sequence shown in SEQ ID NO: 6, SEQ ID NO: 8 or SEQ ID NO: 12; and
- A MVA vector comprising a nucleic acid molecule placed under the control of a vaccinia promoter such as the 7.5K or pH5R promoter, and comprising the nucleotide sequence shown in SEQ ID NO: 13, SEQ ID NO: 14 or SEQ ID NO: 15. Preferably, said nucleic acid molecule is inserted in deletion III of the MVA genome.

If needed, the vector of the invention can further comprise one or more transgene(s), e.g. a gene of interest to be expressed together with the nucleic acid molecule of the invention in a host cell or organism aimed to improve therapeutic or protective activity to an HBV infection or any disease or condition caused by or associated with an HBV infection. Suitable transgenes include without limitation immunomodulators such as cytokines and any other antigen originating from a potentially co-infecting organism (e.g. HIV, tuberculosis mycobacterium, etc). If a transgene is used, it can be expressed from the vector of the invention or from an independent vector for use in combination which can be the same or different with respect to the vector of the invention. For example, one may envisage using in combination an adenovirus expressing the mutant polymerase polypeptide or the fusion protein of the invention and an adenovirus expressing an immunomodulator.

According to a preferred embodiment, the vector of the invention is in the form of infectious viral particles. Typically, such viral particles are produced by a process comprising the steps of (a) introducing the viral vector of the invention into a suitable cell line, (b) culturing said cell line under suitable conditions so as to allow the production of said infectious viral particle, (c) recovering the produced viral particle from the culture of said cell line, and (d) optionally purifying said recovered viral particle.

When the viral vector is defective, the particles are usually produced in a complementation cell line or via the use of a helper virus, which supplies *in trans* the non-functional viral genes. For example, suitable cell lines for complementing E1-deleted adenoviral vectors include the 293 cells (Graham et al., 1997, J. Gen. Virol. 36:

59-72) as well as the HER-96 and PER-C6 cells (e.g. Fallaux et al., 1998, Human Gene Ther. 9: 1909-1917; WO97/00326) or any derivative of these cell lines. But any other cell line described in the art can also be used in the context of the present invention, especially any cell line used for producing product for human use such as Vero cells, 5 HeLa cells and avian cells particularly suitable for propagating poxvirus vectors. Suitable avian cells include without limitation primary chicken embryo fibroblasts (CEF) prepared from chicken embryos obtained from fertilized eggs, and duck cell lines (e.g. as described in WO03/076601, WO2009/004016, WO2010/130756 and US2011-008872).

10 The infectious viral particles may be recovered from the culture supernatant and/or from the cells after lysis. They can be further purified according to standard techniques (chromatography, ultracentrifugation in a cesium chloride gradient as described for example in WO96/27677, WO98/00524, WO98/22588, WO98/26048, WO00/40702, EP1016700 and WO00/50573).

15 The present invention also encompasses vectors or viral particles that have been modified to allow preferential targeting to a specific host cell. A characteristic feature of targeted vectors is the presence at their surface of a ligand capable of recognizing and binding to a cellular and surface-exposed component such as a cell-specific marker (e.g. an HBV-infected cell), a tissue-specific marker (e.g. a liver-specific marker), as well as 20 a viral (e.g. HBV) antigen. Examples of suitable ligands include antibodies or fragments thereof directed to an HBV antigenic domain. Targeting can be carried out by genetically inserting the ligand into a polypeptide present on the surface of the virus (e.g. adenoviral fiber, penton, pIX or vaccinia p14 gene product).

25 Host cells

In another aspect, the invention also relates to host cells which comprise the nucleic acid molecules or vectors (or viral particles) of the invention.

As used herein, the term "host cell" should be understood broadly without any limitation concerning particular organization in tissue, organ, or isolated cells. Such 30 cells may be of a unique type of cells or a group of different types of cells such as cultured cell lines, primary cells and proliferative cells. In the context of the invention, the term "host cells" include prokaryotic cells, lower eukaryotic cells such as yeast, and

other eukaryotic cells such as insect cells, plant and mammalian (e.g. human or non-human) cells as well as cells capable of producing the vector of the invention (e.g. 293, HER96, PERC.6 cells, Vero, HeLa, CEF, duck cell lines, etc). This term includes cells which can be or has been the recipient of the vector described herein as well as progeny
5 of such cells. Host cells can be cultured in conventional fermentation bioreactors, flasks, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a given host cell. No attempts will be made here to describe in detail the various prokaryote and eukaryotic host cells and methods known for the production of the polypeptides and vectors in use in the invention.

10 According to a specific embodiment of the invention, the host cell can be further encapsulated. Cell encapsulation technology is known in the art.

Still a further aspect of the present invention is a method for recombinant production of the mutant polymerase polypeptide or the fusion protein of the invention,
15 employing the vectors (or infectious viral particles) and/or host cells of the invention. Typically, the method comprises (a) introducing a vector into a suitable host cell to produce a transfected or infected host cell, (b) culturing *in-vitro* said transfected or infected host cell under conditions suitable for growth of the host cell, (c) recovering the cell culture, and (d) optionally, purifying the mutant polymerase polypeptide or the
20 fusion protein from the recovered cell and/or culture supernatant.

It is expected that those skilled in the art are knowledgeable in the numerous expression systems available in the art for expressing the mutant polymerase polypeptide or the fusion protein in appropriate host cells (such as those described above) and of the methods for introducing a vector into a host cell. Such methods
25 include, but are not limited to microinjection, CaPO_4 - mediated transfection, DEAE-dextran-mediated transfection, electroporation, lipofection/liposome fusion, gene guns, transduction, viral infection as well as direct administration into a host organism via various means. The vector of the invention can be used in association with transfection reagents in order to facilitate introduction in the host cell, such as polycationic polymers
30 (e.g. chitosan, polymethacrylate, PEI, etc) and cationic lipids (e.g. DC-Chol/DOPE, transfectam lipofectin now available from Promega).

Host cells can be cultured in conventional fermentation bioreactors, flasks, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a given host cell. The mutant polymerase polypeptide or the fusion protein can then be purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, gel electrophoresis; filtration and chromatographic methods (e.g. reverse phase, size exclusion, ion exchange, affinity, hydrophobic-interaction, hydroxylapatite, high performance liquid chromatography, etc). The conditions and technology to be used depend on factors such as net charge, molecular weight, hydrophobicity, hydrophilicity and will be apparent to those having skill in the art. Moreover, the level of purification will depend on the intended use.

Compositions

In another aspect, this invention provides a composition comprising at least the mutant polymerase polypeptide or the fusion protein, the nucleic acid molecule, the vector (e.g. the infectious viral particle), or the host cell described herein (also referred herein to "active agent") or any combination thereof (e.g. combination of different polypeptides or vectors/viral particles as described herein or combination of different genotypes). Preferably, the composition is a pharmaceutical composition which comprises a pharmaceutically acceptable vehicle further to a therapeutically effective amount of the active agent(s).

As used herein, a "pharmaceutically acceptable vehicle" is intended to include any and all carriers, solvents, diluents, excipients, adjuvants, dispersion media, coatings, antibacterial and antifungal agents, and absorption delaying agents, and the like, compatible with administration in a host organism and in particular in human.

As used herein a "therapeutically effective amount" is a dose sufficient for the alleviation of one or more symptoms normally associated with an HBV infection or any disease or condition caused by or associated with an HBV infection. When prophylactic use is concerned, this term means a dose sufficient to prevent or to delay the establishment of an HBV infection. "Therapeutic" compositions are designed and administered to a host organism already infected by an HBV with the goal of reducing or ameliorating at least one disease or condition caused by or associated with said HBV infection, eventually in combination with one or more conventional therapeutic

modalities as described herein (e.g. treatment with nucleoside, nucleotide analogs and/or IFN-based therapy). For example, a therapeutically effective amount for inducing an immune response could be that amount necessary to cause activation of the immune system (e.g. resulting in the development of an anti-HBV response).

5 The term “host organism” generally refers to a vertebrate, particularly a member of the mammalian species and especially domestic animals, farm animals, sport animals, and primates including humans for whom any product and method of the invention is needed or may be beneficial such as organisms who have been diagnosed as being or at risk of being infected with an HBV and thus are susceptible of having or at
10 risk of having a disease or condition caused by or associated with an HBV infection. In preferred embodiments, the host organism is a human patient chronically infected with an HBV virus or alternatively co-infected with an HBV virus and another virus (e.g. the human immunodeficiency virus HIV). The infecting HBV can be from the same genotype, strain or isolate as any HBV from which originates the mutant polymerase
15 polypeptide or any other HBV polypeptide/peptide in use in the present invention (e.g. genotype D) or it can be from a different genotype (e.g. genotype B or C). The cross reactive potential of a genotype D-based vaccine composition was recently investigated by the inventors (see US application 13/423,193). A vast *in silico* study highlighted that the amino acid sequences of HBV polymerase, core and Env proteins are highly
20 conserved among genotypes B, C and D at the global protein level but also at the T cell epitope level. *In vivo* immunization in a suitable animal model supported the ability to induce cross reactive T cell responses recognizing genotype B and C epitopes. Even if this study is limited to HLA-A2 epitopes, it provides a proof of concept that a vaccine composition based on genotype D antigens is able to induce T cell responses that are
25 cross reactive with other HBV genotypes.

The composition of the invention is suitably buffered in order to be appropriate for human use at a physiological or slightly basic pH (e.g. from approximately pH 7 to approximately pH 9). Suitable buffers include without limitation phosphate buffer (e.g.
30 PBS), bicarbonate buffer and/or Tris buffer.

The composition of the invention can further comprise a diluent appropriate for human or animal use. It is preferably isotonic, hypotonic or weakly hypertonic and has a relatively low ionic strength. Representative examples include sterile water, physiological saline (e.g. sodium chloride), Ringer's solution, glucose, trehalose or
5 saccharose solutions, Hank's solution, and other aqueous physiologically balanced salt solutions (see for example the most current edition of Remington : The Science and Practice of Pharmacy, A. Gennaro, Lippincott, Williams&Wilkins).

The pharmaceutically acceptable vehicles included in the composition of the invention must also permit to preserve its stability under the conditions of manufacture
10 and long-term storage (i.e. at least one month with a preference for at least one year) at freezing (e.g. -70°C, -20°C), refrigerated (e.g. 4°C) or ambient temperatures. In this respect, formulations which are particularly adapted to the composition of the invention include (a) 1M saccharose, 150 mM NaCl, 1mM MgCl₂, 54 mg/l Tween 80, 10 mM Tris pH 8.5 (especially when the active agent is an adenoviral vector), (b) 10 mg/ml
15 mannitol, 1 mg/ml HSA, 20 mM Tris, pH 7.2, and 150 mM NaCl and (c) physiological saline.

Additional pharmaceutically acceptable excipients may be used for providing desirable pharmaceutical or pharmacodynamic properties, including for example modifying or maintaining the pH, osmolarity, viscosity, clarity, colour, sterility,
20 stability, rate of dissolution of the formulation, modifying or maintaining release or absorption into an the human or animal organism, promoting transport across the blood barrier or penetration in a particular organ (e.g. liver).

In addition, the composition of the invention may comprise one or more adjuvant(s) suitable for systemic or mucosal application in humans. Preferably, the
25 adjuvant is capable of stimulating immunity to the composition of the invention, especially a T cell-mediated immunity e.g. through the toll-like receptors (TLR), such as TLR-7, TLR-8 and TLR-9. Representative examples of useful adjuvants include without limitation alum, mineral oil emulsion such as Freund's complete and incomplete (IFA), lipopolysaccharide or a derivative thereof (Ribi et al., 1986, Immunology and
30 Immunopharmacology of Bacterial Endotoxins, Plenum Publ. Corp., NY, p407-419), saponins such as QS21 (Sumino et al., 1998, J.Virol. 72: 4931; WO 98/56415), imidazo-quinoline compounds such as Imiquimod (Suader, 2000, J. Am Acad

Dermatol. 43: S6), S-27609 (Smorlesi, 2005, Gene Ther. 12: 1324) and related compounds such as those described in WO2007/147529, cytosine phosphate guanosine oligodeoxynucleotides such as CpG (Chu et al., 1997, J. Exp. Med. 186: 1623; Tritel et al., 2003, J. Immunol. 171: 2358) and cationic peptides such as IC-31 (Kritsch et al.,
5 2005, J. Chromatogr Anal. Technol Biomed Life Sci 822: 263).

The composition of the present invention is suitable for a variety of modes of administration.

The term “administration” (and any form of administration such as
10 “administered”) as used herein refers to the delivery of a therapeutic agent such as the mutant polymerase polypeptide, the fusion protein, the nucleic acid molecule, the vector described herein into a host cell or organism. A number of methods and means are available in the art such as direct administration to a host organism.

Direct administration can be performed by systemic, topical or mucosal routes
15 Systemic administration includes for example subcutaneous, intradermal, intramuscular, intravenous (e.g. injection into a vein feeding liver such as the portal vein), intraperitoneal, intratumoral, intravascular, intraarterial injection (e.g. by hepatic artery infusion) as well as scarification. Injections can be made with conventional syringes and needles, or any other appropriate devices available in the art (e.g. electroporation).
20 Alternatively the composition of the present invention may be administered via a mucosal route, such as the oral/alimentary, intranasal, intratracheal, intrapulmonary, intravaginal or intra-rectal route. Administration in the respiratory tract can be performed through nebulisation or aerosolization of droplet, spray, or dry powdered compositions using appropriate dispenser. Topical administration can also be performed
25 using transdermal means (e.g. patch and the like).

In the context of the invention, the composition is preferably formulated for intramuscular, subcutaneous, intradermal administration or scarification.

The composition of the invention can be in various forms, e.g. solid, liquid or frozen. Solid (e.g. dry powdered or lyophilized) compositions can be obtained by a
30 process involving vacuum drying and freeze-drying. For mucosal administration, the compositions can be formulated as gastroresistant capsules and granules for oral administration, suppositories for rectal or vaginal administration, eventually in

combination with absorption enhancers useful to increase the pore size of the mucosal membranes. Such absorption enhancers are typically substances having structural similarities to the phospholipid domains of the mucosal membranes such as sodium deoxycholate, sodium glycocholate, dimethyl-beta-cyclodextrin, lauryl-1-lysophosphatidylcholine).

The appropriate dosage can be adapted as a function of various parameters, in particular the mode of administration; the composition employed; the age, health, and weight of the host organism; the nature and extent of symptoms; kind of concurrent treatment; the frequency of treatment; and/or the need for prevention or therapy. Further refinement of the calculations necessary to determine the appropriate dosage for treatment is routinely made by a practitioner, in the light of the relevant circumstances.

For general guidance, suitable dosage for a vector-comprising composition varies from about 10^5 to about 10^{13} vp (viral particles), iu (infectious unit) or pfu (plaque-forming units) depending on the vector and the quantitative technique used. Techniques available to evaluate the quantity of vp, iu and pfu present in a sample are conventional in the art. For example, the number of adenoviral particles (vp) is usually determined by measuring the A260 absorbance or HPLC, iu titers by quantitative DBP immunofluorescence and pfu by counting the number of plaques following infection of permissive cells. Preferably the vp/iu ratio is below 100 in accordance with FDA guidelines. Preferred doses contain from about 10^5 to about 10^{12} vp, with a specific preference for doses of about 5×10^8 , about 10^9 , about 5×10^9 , about 10^{10} , about 5×10^{10} vp or about 10^{11} vp of an adenoviral vector of the invention. Doses from about 5×10^5 to about 10^9 pfu are preferred for vaccinia (MVA)-based composition with a specific preference for doses of about 5×10^6 , about 10^7 , about 5×10^7 , about 10^8 or about 5×10^8 pfu. A composition based on vector plasmids may be administered in doses of between 10 μ g and 20 mg, advantageously between 100 μ g and 2 mg. A protein composition may be administered in one or more doses of between 10 ng and 20 mg, with a special preference for a dosage from about 0.1 μ g to about 2 mg of the immunogenic polypeptide per kg body weight. The administration may take place in a single dose or a dose repeated one or several times after a certain time interval.

In another specific embodiment, the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell or composition of the invention can be used in combination with additional polypeptides or peptides or vector encoding such additional polypeptides or peptides. Preferably, said additional polypeptide or peptide is
5 a HBV antigen with a specific preference for an Hbc polypeptide and/or one or more HBs immunogenic domains as described herein. The HBV polypeptide or peptide can be expressed from a vector, in particular a vector selected from the group consisting of plasmid DNA, adenoviral (e.g. Ad5, AdCh3, AdCh63, etc), poxviral (e.g. vaccinia such as MVA) and measles vectors. Accordingly the invention also relates to a composition
10 comprising a mixture of a vector encoding a mutant polymerase polypeptide or a fusion protein of the invention and a vector encoding at least one additional polypeptide/peptide such as an HBV core and/or HBs immunogenic domain(s) as described herein.

15 The mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell or composition of the invention may be employed in methods for treating a variety of diseases and pathologic conditions, especially those caused by or associated with an HBV infection. Accordingly, the present invention also encompasses the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host
20 cell or composition of the invention for use for treating or preventing HBV infections, HBV-associated diseases and pathologic conditions, according to the modalities described herein, and particularly chronic HBV infection. The present invention also relates to a method of treatment in an organism in need thereof comprising at least one administration to said organism of at least one of such active agents in an amount
25 sufficient to treat or prevent HBV infections (e.g. particularly chronic HBV infection) or alleviate one or more symptoms related to HBV-associated diseases and pathologic conditions, according to the modalities described herein. In a particular embodiment, the active agent(s) and method(s) of the invention may be employed according to the modalities described herein to break HBV-specific immune tolerance usually
30 encountered in HBV chronic subjects.

The term “treating” (and any form of treating such as “treatment”) as used herein refers to prophylaxis (e.g. prevention of a subject at risk of being infected with HBV) and/or therapy (e.g. in a subject diagnosed as being infected with an HBV). It is especially useful for treating HBV chronic infection and/or liver lesions in HBV-
5 infected patients including cirrhosis and liver cancer. Treatment requires administer externally or internally to a host cell or organism a therapeutic agent such as the mutant polymerase polypeptide described herein, eventually in combination with other HBV polypeptide(s) or with the standard of care (SOC) (e.g. treatment with nucleoside or nucleotide analogs).

10 Typically, upon administration into a host organism according to the modalities described herein, the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention provides a therapeutic benefit to the treated host organism over the baseline status or over the expected status if not treated. The therapeutic benefit can be evidenced by any relevant clinical
15 measurement typically used by physicians or other skilled healthcare staff, including, for instance, a decrease of the HBV viral load quantified in blood, plasma, or sera of a treated organism, and/or a decrease of the level of liver enzyme activity (e.g. alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST)), and/or a stabilized (not worsening) state of disease (e.g. stabilization or decrease of conditions typically
20 associated with HBV infection such as liver inflammation/steatosis/fibrosis), and/or the reduction of the level of sero markers such as HBeAg or HBsAg (e.g. HBe or HBs seroconversion) and/or an improved response of the treated organism to conventional therapies and/or a survival extension as compared to expected survival if not receiving treatment.

25 In the context of the invention, the therapeutic benefit can be transient (for one or a couple of months after cessation of administration) or sustained (for several months or years). As the natural course of clinical status which may vary considerably from a subject to another, it is not required that the therapeutic benefit be observed in each organism treated but in a significant number of organisms (e.g. statistically significant
30 differences between two groups can be determined by any statistical test known in the art, such as a Tukey parametric test, the Kruskal-Wallis test the U test according to Mann and Whitney, the Student’s t-test, the Wilcoxon test, etc).

Such measurements can be performed before the administration of the described herein (baseline) and at various time points during treatment and at least for 12 weeks after cessation of the treatment. For general guidance, the viral load can be determined using a quantitative PCR assay or any other methodology accepted in the art (e.g. Roche
5 Ampli Prep/Cobas taqman assay v2.0, Abbott real-time hepatitis B virus performance assay). In preferred embodiments, the administration of the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention results in a reduction of the viral load whether transient or sustained of at least one $\log_{10}$, preferably at least 1.5 $\log_{10}$ and more preferably at least 2 $\log_{10}$ as
10 compared to the viral load measured at baseline or as compared to the control group (non-treated subjects). The administration of the active agent(s) described herein can result in a at least transient return to normal ALT and/or AST values as compared to baseline or to the control group. The levels of liver enzyme activity can be evaluated routinely in medical laboratories and hospitals. Alternatively, the administration of the
15 active agent(s) described herein results in a at least transient reduction of seromarker HBe and/or HBs of at least one $\log_{10}$, preferably at least 1.5 $\log_{10}$ and more preferably at least 2 $\log_{10}$ as compared to the seromarker level measured at baseline or as compared to the control group (non-treated subjects). The levels of HBV seromarker can be evaluated routinely in medical laboratories and hospitals and a large number of kits are
20 available commercially (e.g. immunoassays developed by Abbott Laboratories, Organon Technika).

Preferably, the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention are/is used or
25 administered for eliciting or stimulating an immune response in the treated organism. Accordingly, the present invention also encompasses a method for eliciting or stimulating an immune response against HBV upon administration in a host organism of the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention.

The elicited or stimulated immune response can be specific (i.e. directed to HBV epitopes/antigen) and/or non-specific (innate), humoral and/or cellular. In the context of the invention, the immune response is preferably a T cell response CD4+ or CD8+-mediated or both, directed to an HBV polypeptide/epitope.

5 The ability of the active agents(s) described herein to elicit an immune response can be evaluated either *in vitro* or *in vivo* using a variety of direct or indirect assays which are standard in the art. Testing and validation are also illustrated in the appended Example section.

For a general description of techniques available to evaluate the onset and
10 activation of an immune response, see for example Coligan et al. (1992 and 1994, Current Protocols in Immunology ; ed J Wiley & Sons Inc, National Institute of Health or subsequent editions). The ability to stimulate a humoral response may be determined by antibody binding and/or competition in binding (see for example Harlow, 1989, Antibodies, Cold Spring Harbor Press).

15 Evaluation of non-specific immunity can be performed by for example measurement of the NK/NKT-cells (e.g. representativity and level of activation), as well as, IFN-related cytokine and/or chemokine producing cascades, activation of TLRs and other markers of innate immunity (Scott-Algara et al., 2010 PLOS One 5(1), e8761; Zhou et al., 2006, Blood 107, 2461-2469; Chan, 2008, Eur. J. Immunol. 38, 2964-2968).

20 Evaluation of cellular immunity can be performed for example by quantification of cytokine(s) produced by activated T cells including those derived from CD4+ and CD8+ T-cells using routine bioassays (e.g. characterization and/or quantification of T cells by ELISpot, by multiparameters flow cytometry or ICS, by cytokine profile analysis using multiplex technologies or ELISA), by determination of the proliferative
25 capacity of T cells (e.g. T cell proliferation assays by [³H] thymidine incorporation assay), by assaying cytotoxic capacity for antigen-specific T lymphocytes in a sensitized subject or by immunization of appropriate animal models.

The immunogenic capacity of the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention can
30 also be further validated in animal models, which can be challenged with an appropriate infectious or tumor-inducing agent (e.g. a vaccinia virus or a *Listeria Monocytogenes* bacteria expressing HBV gene products) or injected by a DNA encoding the full-length

HBV genome (such as described in Huan et al., 2010, Proc. Natl. Acad. Sci. 107: 9340 to determine neutralization of the infectious or tumor-inducing agent and eventually partial resistance to the associated symptoms, reflecting an induction or an enhancement of an anti-HBV immune response. Exemplary animal models include without limitation 5 the HLA-A2.1 transgenic mice described in Examples, and HBV transgenic mice such as those described in Chisari et al. (1996, Curr. Top. Microbiol. Immunol., 206: 149) and Halverscheid al. (2008, J. Med. Virol. 80: 583).

Said use or method comprises one or more administration(s) (1, 2, 3, 4, 5, 6, 7, 10 8, 9, 10, etc) of a therapeutically effective amount of said active agent(s), said administrations being separated from each other by an appropriate period of time and being carried out by the same route of administration or by different routes of administrations at the same site or at different sites. Intramuscular and subcutaneous routes are particularly preferred in the context of the invention. Three subcutaneous 15 administrations separated from each other by approximately one week are particularly suitable for MVA-based compositions and vectors whereas one or two intramuscular or subcutaneous administration(s) are particularly suitable for Ad-based compositions and vectors which can be separated from each other by approximately one month or more. The first series of administrations can be followed by one or more subsequent 20 administration(s) using the same active agent(s) two or several months after so as to recall the anti-HBV immune response.

If desired, the method or use of the invention can be carried out in combination with one or more conventional therapeutic modalities (e.g. radiation, chemotherapy and/or surgery). Preferably, the method or use of the invention is associated with the 25 one or more drugs which are available for treating or preventing HBV infections, HBV-associated diseases and pathologic conditions. Their administration may precede, be concomitant, or subsequent to the administration of the active agent of the invention. Representative examples of suitable drugs include without limitation polymerase inhibitors, RNase H inhibitors, nucleoside analogs, nucleotide analogs, TLR agonists, 30 IFN, N-glycosylation inhibitors, siRNA, antisense oligonucleotides, anti-HBV antibodies, immune modulators, therapeutic vaccines and antitumor agents usually used in the treatment of HBV-associated liver cancers (e.g. adriamycin, adriamycin with

lipiodol or sorafenib). Moreover, the active agent(s) may also be used in combination with other therapeutic vaccines such as synthetic peptides, recombinant antigens, VLPs, vectors encoding HBV proteins (Core, preS1, PreS2, S and/or polymerase) which are particularly suited to trigger an anti-HBV humoral response. A particularly suitable method or use according to the invention is in combination with standard of care and especially treatment with cytokines (e.g. IFN α , pegylated IFN α 2a or 2b such as Pegasys (Roche), Peginteron (Schering Plough) or IntronA (Schering Plough)) and/or with nucleotide or nucleoside analogs (NUCs) such as lamivudine, entecavir, telbivudine, adefovir, dipivoxil or tenofovir. The treatment with NUCs is only partially effective (infection resolution is observed in only 3-5% of subjects after 1 year of treatment) and needs long term therapy (may be life-long). It is expected that the active agents and methods of the invention bring an immune dimension that permits to complement NUC's action on viral replication, thus resulting in an improvement of such treatments (e.g. by decreasing doses of NUCs or length of NUC treatment required to achieve a therapeutic benefit) or an increase of the percentage of infection resolution (greater than 5%).

In a specific embodiment, the method or use of the invention can be carried out according to prime boost modality which comprises sequential administrations of one or more priming composition(s) and one or more boosting composition(s). Typically, the priming and the boosting compositions use different vectors which comprise or encode at least an antigenic domain in common. Moreover, the priming and boosting compositions can be administered at the same site or at alternative sites by the same route or by different routes of administration. For example, compositions based on polypeptide can be administered by a mucosal route whereas compositions based on vectors are preferably injected, e.g. subcutaneous injection for a MVA vector, intramuscular injection for a DNA plasmid and subcutaneous or intramuscular injection for an adenoviral vector.

In one embodiment, the priming is carried out with a MVA vector and the boosting with an Ad vector, with a specific preference for the MVA and/or the Ad vector encoding a mutant polymerase protein or a fusion protein described herein, e.g. the fusion protein shown in SEQ ID NO: 8. The MVA vector is administered to the organism one or more times followed by the administration of the adenoviral vector one

or more times with a specific preference for at least 3 subcutaneous administrations of the MVA vector separated by a period of time varying from 3 days to 3 months followed by a intramuscular or subcutaneous boost of the adenovirus vector (e.g. from approximately 1 month to 1 year after the MVA prime).

5 In another embodiment, the priming is carried out with a plasmid DNA vector and the boosting with a MVA vector, with a specific preference for the plasmid and/or the MVA vector encoding a mutant polymerase protein or a fusion protein described herein, e.g. the fusion protein shown in SEQ ID NO: 8). The DNA vector is administered to the organism one or more times followed by the administration of the
10 MVA vector one or more times with a specific preference for at least 3 intramuscular administrations of the DNA vector separated by a period of time varying from 2 weeks to 3 months and at least one subcutaneous boost of the MVA vector (e.g. from approximately 1 month to 1 year after the DNA prime). Preferably, the DNA vector is administered through electroporation.

15 The present invention also relates to a kit of parts for use in the treatment of an HBV infection or an HBV-associated disease or pathologic condition, wherein said kit comprises a plurality of active agents selected from the group consisting of the mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector, host cell and/or composition of the invention and instructions for administering said plurality of active
20 agents to an organism in need thereof. More preferably, the organism is a patient chronically infected with HBV.

All of the above cited disclosures of patents, publications and database entries
25 are specifically incorporated herein by reference in their entirety to the same extent as if each such individual patent, publication or entry were specifically and individually indicated to be incorporated by reference.

DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the mutant polymerase polypeptides, fusion proteins and antigenic combinations described in the invention.

5

Figure 2 illustrates Elispot IFN γ assays carried out following immunization of HLA-A2 transgenic mice with plasmids pTG18188 (Core-Pol-Env1-Pol-Env2-Pol), pTG18194 (Core-Pol) or pTG13135 (Empty). Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each HBV HLA-A2 peptides evaluated in the experiment for 10^6 spleen cells of immunized mice. Each bar represent an individual mouse vaccinated by one or the other plasmid and indicated by its reference number (1.1 or 2.3 ...etc) and the mean of all mice immunized with the same plasmid is also represented for each group (bar indicated as “mean” on the graph). For individual mouse and the means, frequency of IFN γ producing cells specific of the different tested peptides are piled. Bars are filled with different symbols, each symbol representing the response against one specific HBV peptide as indicated by the legend on the graph.

Figures 3A-F illustrates ICS assays carried out following immunization of HLA-A2 transgenic mice with plasmids pTG18188 (Core-P-E1-P-E2-P or Core-Pol-Env1-Pol-Env2-Pol), pTG18194 (Core-Pol) or pTG13135 (empty). Results are presented as the percentage of CD8 (Figures 3A, 3B, 3D and 3F), or CD4 (Figures 3C and 3E) T cells producing IFN γ (alone or combined with TNF α) specific of each HBV HLA-A2 peptides (Figure 3A) or peptide pools covering HBV core (Figures 3B and 3C), polymerase (Figured 3D and 3E) and env (Figure 3F) antigens. Each bar represent an individual mouse vaccinated by one or the other plasmid and indicated by its reference number (1.1 or 2.3 ...etc) and the mean of all mice immunized with the same plasmid is also represented for each group (bar indicated as “mean” on the graph). For individual mouse and the means, frequency of CD8 or CD4 T cells specific of the different tested peptides are piled. Bars are filled with different symbols, each symbol representing the response against one specific HBV peptide as indicated by the legend on the graph.

Figure 4 illustrates Elispot IFN γ assays carried out following immunization of HLA-A2 transgenic mice with Adenovirus AdTG18201 (Core-Pol-Env Ad), AdTG18202 (Core-Pol Ad), AdTG18203 (Pol Ad) and AdTG15149 (Empty Ad). Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each peptide pools covering the antigen of interest, evaluated in the experiment for 10^6 spleen cells of immunized mice. Each bar represent an individual mouse vaccinated by one or the other plasmid and indicated by its reference number (1.1 or 2.3 ...etc) and the mean of all mice immunized with the same plasmid is also represented for each group (bar indicated as “mean” on the graph).

For individual mouse and the means, frequency of IFN γ producing cells specific of the different tested peptides are piled. Bars are filled with different symbols, each symbol representing the response against one specific HBV peptide as indicated by the legend on the graph.

Figure 5 illustrates Elispots IFN γ assays carried out following immunization of HLA-A2 transgenic mice with AdTG18201 (Core-Pol-Env Ad). Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each HBV HLA-A2 epitope, or of an irrelevant peptide, evaluated in the experiment for 10^6 spleen cells of immunized mice. Each bar represents an individual mouse vaccinated by the AdTG18201 (indicated with its reference number, 3.1 to 3.8) and the median of all these mice immunized with AdTG18201 is also represented (bar indicated as « Median » on the graph). For individual mouse and median, frequency of IFN γ producing cells specific of the different HBV HLA-A2 epitopes are piled whereas the frequency of IFN γ producing cells observed in presence of an irrelevant peptide is represented on a separate bar (indicated « Irrel » on the graph for each mouse). Bars are filled with different symbols, each symbol representing the response against one specific HBV peptide as indicated by the legend on the graph.

Figure 6 illustrates ICS assays carried out following immunization of HLA-A2 mice with AdTG18201 (Core-Pol-Env Ad) or AdTG15149 (Empty Ad). Results are presented as the percentage of CD8 T cells producing IFN γ alone or in combination with TNF α , specific of two selected peptide pools, called PP8 and PC1, and covering

respectively a part of the HBV polymerase (amino acids 725 to 835) and a part of the HBV Core protein (amino acids 1 to 100). Each bar represents an individual mouse vaccinated by one or the other adenovirus and indicated by its reference number (1.1 or 3.2...etc) and the median of all mice immunized with the same adenovirus is also represented for each group (bar indicated as “median” on the graph). Bars are filled with different symbols, each symbol representing the cytokine(s) produced by the detected CD8 T cells as indicated by the legend on the graph.

Figure 7 illustrates *in vivo* CTL assays carried out following immunization of HLA-A2 mice with AdTG18201 (Core-Pol-Env Ad) or AdTG15149 (Empty Ad). Results are presented as the percentage of *in vivo* specific lysis, ie lysis specific of the HBV HLA-A2 epitopes that were tested. Each square or triangle symbols represents an individual mouse and the mean of all mice immunized with the same adenovirus is also represented for each group (represented by a thick bar symbol on the graph).

15

Figure 8 illustrates ICS assays carried out following immunization of HBV transgenic mice with AdTG18201 (Core-Pol-Env Ad) or AdTG15149 (Empty Ad). Results are presented as the percentage of CD8 T cells producing both IFN γ and TNF α specific of epitopes from the HBV polymerase (mix of 2 peptides, called VSA and N13F) or from the HBV envelope (1 peptide called F13L) and found both in spleens and in livers of vaccinated mice. Each bar represents an individual mouse vaccinated by one or the other adenovirus and indicated by its reference number (1.1 or 3.2...etc) and the median of all mice immunized with the same adenovirus is also represented for each group (bar indicated as “median” on the graph). The dotted line represented on graphs corresponds to the cut-off of the experiment, ie the threshold above which observed percentage of CD8 T cells is considered as a positive immune response.

Figure 9 illustrates Elispots IFN γ assays carried out following immunization of HLA-A2 transgenic mice with AdTG18201 (Core-Pol-Env Ad) with different doses, from 10^5 iu to 10^9 iu. Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each HBV HLA-A2 epitope, or of an irrelevant peptide, or in presence of medium only, evaluated in the

30

experiment for 10^6 spleen cells of immunized mice. Each solid or dotted bar represents an individual mouse vaccinated by the AdTG18201 and the mean of all mice immunized with AdTG18201 at one specific dose is also represented (hatched bars). The different doses are represented by different colors or symbols as indicated by the legend on the graph. The dotted-line represents the cut-off value, defined as described in Material and Methods, above which observed T cell responses are considered as positive.

Figure 10 illustrated Elispots IFN γ assays carried out following immunization of HLA-A2 transgenic mice with AdTG18202 (Core-Pol Ad) according to different schedules of injections. Mice were immunized either once (squares) or received 3 injections at 1 week interval (triangles) or 6 injections at 1 week interval (circles). Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each HBV HLA-A2 epitope, or of the adenovirus vector or an irrelevant peptide or in presence of medium only evaluated in the experiment for 10^6 spleen cells of immunized mice. Each symbol (square, triangle or circle) represents an individual mouse vaccinated by the AdTG18202 and the mean of all mice immunized with AdTG18202 with one of the tested schedules is represented by a solid thick line. The dotted-line represents the cut-off value, defined as described in Material and Methods, above which observed T cell responses are considered as positive.

Figure 11 illustrated Elispots IFN γ assays carried out following immunization of HLA-A2 transgenic mice with AdTG18202 (Core-Pol Ad) according to different schedules of injections. Mice were immunized either once 2 weeks before the monitoring of T cell responses (group 1 represented by squares) or once 20 weeks before the monitoring of T cell responses (group 2 represented by triangles), or twice at 2 months interval (group 3 represented by circles), or twice at 4 months interval (group 4 represented by crosses) or three times at 2 month interval (group 5 represented by rhombuses). For all groups except group 2, T cell responses were monitored 2 weeks after the last injection. Results are presented as the number of spots for 10^6 cells corresponding to the frequency of IFN γ producing cells specific of each HBV HLA-A2

epitope or an irrelevant peptide or in presence of medium only, evaluated in the experiment for 10^6 spleen cells of immunized mice. Each symbol (square, triangle, circle, cross, rhombus) represents an individual mouse vaccinated by the AdTG18202 and the mean of all mice immunized with AdTG18202 with one of the tested schedules is represented by a solid thick line. The dotted-line represents the cut-off value, defined as described in Material and Methods, above which observed T cell responses are considered as positive.

10 **EXAMPLES**

1. Material and Methods

The constructions described below (see Figure 1) are carried out according to the general genetic engineered and molecular cloning techniques detailed in Maniatis et al. (1989, Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor NY or subsequent editions) or according to the manufacturer's recommendations when a commercial kit is used. PCR amplification techniques are known to the person skilled in the art (see for example PCR protocols –A guide to methods and applications, 1990, published by Innis, Gelfand, Sninsky and White, Academic Press). The recombinant plasmids carrying the ampicillin resistance gene are replicated in the *E. coli* C600 (Stratagene) on agar or liquid medium supplemented with 100µg/ml of antibiotic.

MVA vector construction are generated by homologous recombination between a shuttle plasmid and the MVA genome as previously described in Erbs et al. (2008, Cancer gene Ther. 15: 18). The “basic” shuttle plasmid contains a multiple cloning site, a vaccinia virus (VV) promoter surrounded by the flanking sequences of deletion III as well as the *E. Coli* xanthine-guanine phosphoribosyl-transferase (GPT) selection gene under the control of p11K7.5 vaccinia promoter (Falkner and Moss, 1988). Briefly, CEF cells were infected with a genomic MVA without any inserted transgene (MVA-null) and then transfected by CaCl_2 precipitation with the shuttle plasmid carrying the gene of interest cloned downstream the VV promoter. Homologous recombination occurred between MVA-null and the shuttle plasmid and recombinant viruses were isolated by

multiple steps of mycophenolic acid selection. Recombinant MVA viruses were controlled by PCR, amplified in CEF and virus stocks were titrated on CEF by plaque assay.

For adenoviral vector construction, an adenoviral shuttle plasmid is first constructed by inserting the nucleic acid molecule of interest into the “basic” shuttle plasmid pTG13135. Typically, the nucleic acid molecule is inserted into the *NheI* and *NotI* restriction sites of pTG13135 containing a CMV-driven expression cassette surrounded by adenoviral sequences (adenoviral nucleotides 1-454 and nucleotides 3513-5781 respectively) to allow further generation of the vector genome by homologous recombination (Chartier et al., 1996, J. Virol. 70:4805). The adenoviral vector is then obtained by homologous recombination between the recombinant shuttle vector digested by *BstI* and *PacI* and pTG15375 (encoding the complete adenoviral genome) linearized by *Clal* digestion. The resulting adenoviral vector is E3 (nucleotides 27867-30743) and E1 (nucleotides 455-3512) deleted, with the E1 region replaced by the expression cassette containing, from 5' to 3', the CMV immediate-early enhancer/promoter, a chimeric human β -globin/IgG intron (as found in pCI vector available in Promega), the nucleic acid molecule of interest and the SV40 late polyadenylation signal. Adenoviral particles are then obtained by transfecting the *PacI* linearized viral genome into an E1 complementation cell line. Virus propagation, purification and titration is made as described previously (Erbs et al., 2000, Cancer Res. 60:3813)

1.1. Vectors constructions and production

The vectors illustrated hereinafter have been engineered to express the mutant polymerase polypeptide eventually fused to the Core polypeptide and/or immunogenic domains of the envelope protein. All HBV sequences originate from HBV strain Y07587 which sequence is described in international databases (Genbank Y07587) and in different publications. It is a genotype D virus of serotype ayw.

The following examples illustrate the fusion of a truncated Core polypeptide (aa 1-148) with a mutated polymerase polypeptide (designated Pol*) comprising two internal deletions (from positions 538 to 544 and from positions 710 to 742) and 4 amino acid substitutions (D689H, V769Y, V776Y and D777H respectively) as

represented in SEQ ID NO: 6 as well as a longer fusion further comprising two immunogenic Env domains (Env1 and Env2 respectively extending from amino acids 14 to 51 and from amino acids 165 to 194 of the HBs protein) inserted in place of the deleted pol regions as represented in SEQ ID NO: 8.

5

1.1.1. Construction and production of plasmid and adenovirus vectors expressing truncated Core-Pol*-Env1-Env2 (or Core-Pol-Env1-Pol-Env2-Pol) fusion

A synthetic gene (3024 nucleotides described in SEQ ID NO: 15) encoding the truncated Core-Pol*-Env1-Env2 fusion protein (amino acid sequence is shown in SEQ ID NO: 8) was synthesized by GENEART (Regensburg, Germany). The synthetic fragment was inserted into the *NheI* and *NotI* restriction sites of pTG13135 shuttle plasmid, providing pTG18188. An adenoviral vector was then obtained by homologous recombination between pTG18188 digested by *BstI107I* and *PacI* and pTG15375 linearized by *ClaI* digestion. The resulting adenoviral vector pTG18201 is E3 and E1 deleted, with the E1 region replaced by the expression cassette containing the synthetic sequence encoding the truncated Core-Pol*-Env1-Env2 driven by the CMV promoter. Adenoviral particles (AdTG18201) were obtained by transfecting the *PacI* linearized viral genome into an E1 complementation cell line.

20 1.1.2. Construction and production of plasmid and adenovirus vectors expressing truncated Core-Pol*

A synthetic gene (2820 nucleotides described in SEQ ID NO: 14) encoding a truncated Core-Pol* fusion protein was synthesized by GENEART (Regensburg, Germany). The synthetic fragment was inserted into the *NheI* and *NotI* restriction sites of pTG13135 shuttle plasmid, providing pTG18194. An adenoviral vector was then obtained by homologous recombination between pTG18194 digested by *BstI107I* and *PacI* and pTG15375 linearized by *ClaI* digestion. The resulting adenoviral vector pTG18202 is E3 and E1 deleted, with the E1 region replaced by the expression cassette containing the synthetic sequence encoding the truncated Core-Pol* driven by the CMV promoter. Adenoviral particles (AdTG18202) were obtained by transfecting the *PacI* linearized viral genome into an E1 complementation cell line.

30

1.1.3. Construction and production of plasmid and adenovirus vectors expressing Pol*

A synthetic gene (2379 nucleotides described in SEQ ID NO: 13) encoding the Pol mutant polypeptide was synthesized by GENEART (Regensburg, Germany). The synthetic fragment was inserted into the *NheI* and *NotI* restriction sites of pTG13135 shuttle plasmid, providing pTG18195. An adenoviral vector was then obtained by homologous recombination between pTG18195 digested by *Bst1107I* and *PacI* and pTG15375 linearized by *ClaI* digestion. The resulting adenoviral vector pTG18203 is E3 and E1 deleted, with the E1 region replaced by the expression cassette containing the synthetic sequence encoding Pol* driven by the CMV promoter. Adenoviral particles (AdTG18203) were obtained by transfecting the *PacI* linearized viral genome into an E1 complementation cell line.

1.2. Immunogenicity evaluation in a mouse model

Antigen immunogenicity was evaluated *in vivo* by Elispot IFN γ assays and Intracellular cytokine staining (ICS) following immunization of HLA transgenic mice.

1.2.1 Mouse model

The HLA-A2.1 transgenic mice used in the study were described by Pascolo et al. (1997, J. Exp. Med. 185:2043). These mice have the H-2D^b and murine β_2 -microglobulin genes knocked-out and express a transgenic monochain histocompatibility class I molecule (HHD molecule) in which the C-terminus of the human β_2m is covalently linked to the N-terminus of a chimeric heavy chain (HLA-A*0201 $\alpha 1$ - $\alpha 2$, H-2D^b $\alpha 3$ transmembrane and intracytoplasmic domains). Seven to 10 weeks-old mice (male and female) were immunized. Average weight of the mice is around 25-30 g.

The HBV transgenic mice used in the study were described by Halverscheid et al (2008, J. Med. Virol. 80: 583-590) and kindly provided by Reinhold Schirmbeck. These mice are on a C57Bl/6J genetic background and transgenic for the HBV genome (1.4 copy of the HBV genome with a mutation at position 1438 (T to C) which avoid the expression of the small form of the HBsAg protein and inhibit the formation of HBV

infectious particles). Ten to 16 weeks-old mice (male and female) were immunized. Average weight of the mice is around 25-30g.

1.2.2. Immunization protocols

5 *1.2.2.1 DNA immunization protocols*

DNA immunization protocols were run in order to evaluate the immunogenicity of the different fusion proteins encoded by the plasmids illustrated in Example 1.1. The DNA used for immunization was produced in endotoxin-free conditions. Mice were immunized twice at 15-day interval with 100µg/injection of each tested plasmid via
10 intramuscular route in the tibialis anterior muscle. A cardiotoxin injection was done prior to the 1st DNA injection in order to favor DNA immunogenicity. Cellular immune response was evaluated 15 days following the last DNA injection.

1.2.2.2 Adenovirus immunization protocols

Adenovirus immunization protocols were run in order to compare the
15 immunogenicity of the different fusion proteins encoded by the Ad vectors which were produced as described in Example 1.1. Mice were immunized once with the adenovirus encoding the different fusion proteins (10^8 iu/mouse/injection) via sub-cutaneous route at the base of the tail. Cellular immune response was evaluated 15 days following the last adenovirus injection.

20 Different doses of adenoviruses were also evaluated with the AdTG18201. Mice were immunized once with 10^5 , 10^6 , 10^7 , 10^8 or 10^9 iu of AdTG18201 by sub-cutaneous route at the base of the tail.

Different schedules of immunization were also tested with the AdTG18202 and mice were injected one, two, three or 6 times at different time interval. Each injection
25 was performed with 10^8 iu/mouse via sub-cutaneous route at the base of the tail. One, 3 or 6 injections at 1 week interval were compared side by side. One injection 2 weeks or 20 weeks before the time of monitoring of induced T cell responses, 2 injections at 2 or 4 month interval and 3 injections at 2 month interval were also compared side by side.

1.2.3 Peptides

Peptides used for cells stimulation *in vitro* are either short peptides of 9 to 10 amino acids which are described or predicted as HLA-A2 restricted epitopes or long peptides of 15 amino acids included in peptide libraries covering all the antigens of interest.

Short peptides corresponding to described or predicted HLA-A2 restricted epitopes of Core protein, Pol protein or Env domains were synthesized by Eurogentec (Belgium) and were dissolved in 100% DMSO (sigma, D2650) at a concentration of 10mM.

Peptides libraries covering the whole Core, Pol and Envelope domains were synthesized by ProImmune (Oxford, United Kingdom). The Core, Pol and Env libraries were composed of 15 mer peptides overlapping by 11 amino acids. Each crude peptide was dissolved in 100% DMSO (sigma, D2650) at a concentration of 50mg/ml. For each library, peptides were pooled to a concentration of 2mg/ml per peptide:

-HBV Core protein was covered by 2 pools of 21 and 22 peptides (Pool 1 (PC1): 22 peptides covering Core residues 1 to 100; Pool 2 (PC2): 21 peptides covering Core residues 89 to 183);

-HBV Pol protein was covered by 8 pools of 24 peptides (Pool 1 (PP1): 24 peptides covering aa 45 to 151; Pool 2 (PP2): 24 peptides covering aa 141 to 251 (peptide from aa 205 to 219 was excluded because of insolubility in 100 % DMSO or DMSO + Tris 100 mM pH9; peptide from aa 221 to 235 was dissolved in DMSO + Tris 100 mM pH9 because of insolubility in 100 % DMSO); Pool 3 (PP3): 24 peptides covering aa 241 to 347; Pool 4 (PP4): 24 peptides covering aa 337 to 447 (peptide from aa 373 to 387 was excluded because of insolubility in 100 % DMSO or DMSO + Tris 100 mM pH9); Pool 5 (PP5): 24 peptides covering aa 437 to 543; Pool 6 (PP6): 24 peptides covering aa 533 to 639; Pool 7 (PP7): 24 peptides covering aa 629 to 735; Pool 8 (PP8): 24 peptides covering aa 725 to 835);

-Env domains were covered by 2 pools of 9 and 10 peptides (Pool 1 (PE1): 10 peptides covering HBs residues 9 to 59; Pool 2 (PE2): 9 peptides covering HBs residues 157 to 194).

For experiments performed in HBV transgenic mice, with a C57BL/6J genetic background, HBV peptides described in the literature or identified in previous experiments as being reactive in mice with a C57BL/6J genetic background were used for cell stimulation *in vitro*. They are either short peptide (VSAAFYHLPL for polymerase; SEQ ID NO: 24) or long peptides (NLNVSIPWTHKVGNF called N13F for polymerase (SEQ ID NO: 25) and FLWEWASARFSWLSL called F13L for envelope protein (SEQ ID NO: 26)). They were synthesized by Eurogentec (Belgium) or by ProImmune (Oxford, United Kingdom). Each peptide was dissolved in 100% DMSO (sigma D2650) at a concentration of 10mM. They were used at a concentration of 10µM during the ICS assays (even when tested as a mix of 2 peptides).

1.2. 4. IFNγ Elispot assays

Splenocytes from immunized mice were collected and red blood cells were lysed (Sigma, R7757). $2 \cdot 10^5$ cells per well were cultured in triplicate for 40 h in Multiscreen 15 plates (Millipore, MSHA S4510) coated with an anti-mouse IFNγ monoclonal antibody (BD Biosciences; 10 µg/ml, 551216) in MEM culture medium (Gibco, 22571) supplemented with 10 % FCS (JRH, 12003-100M), 80 U/mL penicillin / 80 µg/mL streptomycin (PAN, P06-07-100), 2 mM L-glutamine (Gibco, 25030), 1x non-essential amino acids (Gibco, 11140), 10 mM Hepes (Gibco, 15630), 1 mM sodium pyruvate 20 (Gibco, 31350) and 50 µM β-mercaptoethanol (Gibco, 31350) and in presence of 10 units/ml of recombinant murine IL2 (Peprotech, 212-12), alone as negative control, or with:

- 10 µM of a selected HLA-A2 restricted peptide present in HBV antigens encoded by plasmids (FLP, ILC for Core, VLQ, FLG and GLS for Env and SLY for Pol) described in SEQ ID NO: 18-23) or an irrelevant one;
- a pool of peptides at a final concentration of 5µg/ml per peptide
- 5 µg/ml of Concanavalin A (Sigma, C5275) for positive control.

IFNγ-producing T cells were quantified by Elispot (cytokine-specific enzyme linked immunospot) assay as previously described (Himoudi et al., 2002, J. Virol. 30 76:12735). The number of spots (corresponding to the IFNγ-producing T cells) in negative control wells were subtracted from the number of spots detected in experimental wells containing HBV peptides. Results are shown as the mean value

obtained for triplicate wells. An experimental threshold of positivity for observed responses (or cut-off) was determined by calculating a threshold value which corresponds to the mean value of spots observed with medium alone + 2 standard deviations, reported to 10^6 cells. A technical cut-off linked to the CTL Elispot reader was also defined as being 50 spots/ 10^6 cells (which is the value above which the CV (coefficient of variation) of the reader was systematically less than 20%). The highest value between the technical cut-off and the experimental threshold calculated for each experiment was taken into account to define the cut-off value of each experiment. Statistical analyses of Elispot responses were conducted by using a Mann-Whitney test. P value equal or inferior to 0.05 will be considered as significant.

1.2. 5. Intracellular cytokine staining (ICS) assays

ICS were performed on splenocytes from each animal of each group. Following red blood cells lysis with lysis buffer (Sigma, R7757), 2×10^6 cells per well in flat-bottom 96-well plate were incubated in complete alpha MEM culture medium (Gibco BRL, 22571) in the presence of 10 units/ml of murine recombinant IL-2 (Peprotech, 212-12) alone as negative control or with 10 μ M of specific HBV peptide or with a pool of peptides at a final concentration of 5 μ g/ml per peptide or with 10 μ M of an irrelevant peptide. The GolgiPlug (BD Biosciences, 555029) was immediately added at a 1 μ l/ml final concentration for 5 h. Then, cells were harvested in V-bottom 96-well plates and washed with 1 % FCS-PBS. Staining was performed using monoclonal antibodies against CD3 (hamster MAb anti-CD3e-PE, dilution 1/200), CD8 (rat MAb anti CD8a-APC, dilution 1/600) and CD4 (rat MAb anti-CD4-PerCP, dilution 1/600) (all from BD Biosciences, 553063, 553035 and 553052 respectively) in 50 μ l of 1 % FCS-PBS for 15 min at room temperature. After washing, cells were fixed and permeabilized with Cytofix/Cytoperm and washed with Perm/Wash solution (BD Biosciences, 554714). Then, the anti-mouse IFN γ -PE antibodies (BD Biosciences, 554412557724) and anti-mouse TNF α -Alexa488 antibodies (BD Biosciences, 557719) or the anti-mouse IFN γ -PE antibodies (BD Biosciences, 554412557724) were added for 15 min at room temperature and after washing with Perm/Wash, cells were resuspended in 1% FCS-PBS and analysed by flow cytometry using a FacsCalibur (Becton Dickinson). CD3e $^{+}$, CD8a $^{+}$ cells or CD3e $^{+}$, CD4 $^{+}$ cells were gated to determine percentages of IFN γ $^{+}$

CD8⁺ or IFN γ ⁺ CD4⁺ T or TNF α ⁺ CD8⁺ or TNF α ⁺ CD4⁺ T or IFN γ ⁺ TNF α ⁺ CD8⁺ or IFN γ ⁺ TNF α ⁺ CD4⁺ T cell population. The percentage obtained in medium only was considered as background.

For experiments performed in HBV transgenic mice, ICS were also performed
5 on liver cells of each animal of each group. After euthanasia of the mouse, the liver was perfused *in situ* by the hepatic portal vein with cold PBS until the organ becomes pale. The liver was harvested, placed in PBS + FCS 2 % solution, cut into small pieces, pressed gently through a 70 μ m cell-strainer and then suspended in cold PBS + 2% FCS solution. After centrifugation, cells were washed again with cold PBS + 2% FCS
10 solution. After a new centrifugation, the pellet containing cells was resuspended in 10mL of Percoll solution, centrifuged for 12 minutes at 700g at room temperature and washed again with PBS + 2% FCS solution. Then red blood cells were lysed as described before for splenocytes and all subsequent steps were performed as described in the previous paragraph for splenocytes. Of note, for the liver, as the quantity of T
15 lymphocytes collected is limited, number of cells per well is variable: all obtained cells were cultured in a way that an equivalent quantity of cells was dispatched in all wells.

1.2.6 *In vivo* CTL assays

In vivo CTL assay was performed as described by Fournillier et al. (2007,
20 Vaccine, 25: 7339-53) in HLA-A2 transgenic mice. Splenocytes suspensions were obtained from syngenic mice spleens and adjusted to 20×10^6 cells/mL after lysis of red blood cells. Half of the cells were incubated with the HBV peptide of interest (SLY, FLP or ILC) at 10 μ M final concentration for 1 h at 37 °C and half of the cells was left unpulsed. 5(6)-carboxyfluorescein diacetate succinimidyl ester (CFSE) (Molecular
25 probes, C1157) was then added at 10 μ M (CFSE-high) to unpulsed cells and 1 μ M (CFSE-low) to HBV-peptide pulsed cells for 10 min. After washing with PBS, all populations were mixed and 20×10^6 total cells were injected to anaesthetized mice via the retro-orbital sinus, mice being previously immunized (2 weeks earlier) by AdTG18201 or AdTG15149. Thus, CFSE-low population represented specific targets
30 supposed to be lysed by cytotoxic T cells induced by the vaccination and CFSE-high population was an internal reference allowing assay normalization. Splenocytes from recipient mice were analyzed 24 h later by flow cytometer to detect the CFSE-labeled

cells. For each animal, ratio between peptide-pulsed targets and unpulsed targets was calculated ($R = \text{Number CFSE-low cells} / \text{Number CFSE-high cells}$). The percentage of specific lysis for each animal was determined by the following formula: $\% \text{ lysis} = (1 - R_{\text{mouse}} / R_{\text{reference}}) \times 100$ where $R_{\text{reference}}$ is the mean R obtained for 2 naïve HLA-A2 mice which were injected with the same suspension of CFSE-labeled targets. A response was considered positive if the percentage of specific lysis was higher than 10 %.

1.3 *In vitro* analysis of AdTG18201 by electron microscopy

A549 cells (Human lung adenocarcinoma epithelial cell line) were infected in suspension and under reduced medium volume conditions with AdTG18201 at different MOI (25 to 100) and then cultured for 16H, 24H or 48H before being collected for analysis. Cells were collected at these different timepoints and then fixed using glutaraldehyde 2% diluted in sodium cacodylate buffer 0.2M. Cells were then dried, included in blocks of resin and then cut in ultra-thin sections. Obtained grids were then stained using uranyl acetate and lead citrate and observed by electron microscopy.

2. Results

2.1. *Immunogenicity of HBV fusion proteins expressed by DNA plasmids pTG18188 and pTG18194*

The immunogenicity of the HBV fusion proteins expressed by DNA plasmids was assessed in HLA-A2 transgenic mice. Following two intramuscular injections of either pTG18188 (tCore-Pol*-Env1-Env2) or pTG18194 (tCore-Pol*) or pTG13135 as negative control (empty plasmid), specific T cell responses were evaluated by Elispot IFN γ and ICS (IFN γ /TNF α) using known HLA-A2 epitopes present in Polymerase, Core or the envelope domains and/or pools of overlapping peptides covering the HBV antigens of interest.

2.1.1. HBV specific IFN γ producing cell evaluation by Elispot assays

As illustrated in Figure 2, immunization with the plasmid pTG18194 encoding the HBV fusion protein “tCore-Pol*” induced IFN γ producing cells specific of the HLA-A2 restricted SLY epitope (SEQ ID NO: 23) located within the HBV polymerase

(positions 816-824). Immunization with the plasmid pTG18194 also resulted in the induction of high frequency of IFN γ producing cells specific for 2 Core HLA-A2 restricted epitopes FLP (SEQ ID NO: 18, located within the HBV Core protein at position 18-27) and ILC (SEQ ID NO: 19 located within the HBV Core protein at position 99-108). Positive responses were observed in 4 out of the 8 vaccinated mice.

As illustrated in Figure 2, the plasmid pTG18188 encoding the HBV fusion protein “Core-Pol*-Env1-Env2” also induced IFN γ producing cells specific of the pol HLA-A2 epitope SLY and of the 2 Core HLA-A2 restricted epitopes FLP and ILC. Positive responses were observed in 4 out of the 8 vaccinated mice. In addition, IFN γ -producing cells specific of HLA-A2 GLS epitope (SEQ ID NO: 22 located within Env2 at positions 185-194 of HBsAg) were also detected although at a weak frequency and in 1 vaccinated mouse.

2.1.2. Evaluation of induced HBV specific IFN γ producing T CD8 $^{+}$ and CD4 $^{+}$ cells by intracellular staining assays

2.1.2.1. CD8 T cell response specific of HLA-A2 restricted epitopes

The percentage of CD8 T cells producing either IFN γ alone or combined with TNF α targeting HLA-A2 restricted epitopes included into polymerase (SLY), Core (FLP and ILC) and envelope domains (VLQ, FLG and GLS) was evaluated by ICS assay. The results are shown in Figure 3A as percentages of CD8 $^{+}$ T cell specific of these epitopes and producing IFN γ (sum of single IFN γ producing cells or double IFN γ and TNF α producing cells). Four out of 8 animals immunized with pTG18194 (expressing tCore-Pol*) mounted IFN γ producing CD8 $^{+}$ T cells specific of FLP, ILC and SLY HLA-A2 restricted epitopes located respectively in Core and Pol antigens. Similarly, 4 out of 8 animals immunized with pTG18188 expressing tCore-Pol*-Env1-Env2) also mounted IFN γ producing T CD8 $^{+}$ T cells specific of FLP, ILC and SLY epitopes. In addition as already observed in ELISPOT assay, 1 out of 8 mice immunized with the plasmid pTG18188 displayed a response specific of the GLS HLA-A2 restricted epitope located within the Env2 domain mediated by IFN γ producing CD8 $^{+}$ T cells. Immunization with pTG13135 did not induce any specific response as expected.

2.1.2.2 CD8 and CD4 T cell response specific of pools of peptides covering the Core protein, Polymerase protein and Env domains.

Responses specific of pools of peptides covering the Core protein

The percentage of CD8 and CD4 T cells able to produce either IFN γ alone or
5 combined with TNF α in response to pools of peptides covering the Core protein (PC)
was evaluated by ICS assay. The results are expressed as percentages of CD8+ or CD4
+ T cell specific of these pools of peptides and producing IFN γ (sum of single IFN γ
producing cells or double IFN γ and TNF α producing cells).

As shown in Figure 3B, a positive percentage of CD8+ T cells producing IFN γ
10 was detected against the 2 pools of peptides covering the Core protein (PC1 and PC2),
with a CD8+ T cell response mainly focused on peptides of Pool Core 1. The
percentage of reactive CD8+ T cells observed in mice vaccinated by either the
pTG18194 or the pTG18188 was significantly different from the percentage that was
observed for mice vaccinated with the negative control (pTG13135) ($p < 0.05$, Mann
15 Withney test) for both peptide pools (1 and 2).

As shown in Figure 3C, positive percentage of CD4+ T cells producing IFN γ
was also detected against one pool of peptides covering the Core protein, the pool Core
2 in the two groups of mice vaccinated with pTG18194 or pTG18188. The percentage
of reactive CD4+ T cells observed in mice vaccinated by the pTG18188 was
20 significantly different from the percentage that was observed for mice vaccinated with
the negative control (pTG13135) ($p < 0.05$, Mann Withney test) for pool Core 2.

Responses specific of pools of peptides covering the polymerase protein

The percentage of CD8 and CD4 T cells able to produce either IFN γ alone or
combined with TNF α in response to pools of peptides covering the polymerase protein
25 was evaluated by ICS assay. The results are expressed as percentages of CD8+ or CD4
+ T cell specific of these pools of peptides and producing IFN γ (sum of single IFN γ
producing cells or double IFN γ and TNF α producing cells).

As shown in Figure 3D, a positive percentage of CD8+ T cells producing IFN γ
was mainly detected against one pool of peptides, PP8. Specifically for PP8, the
30 percentage of reactive CD8+ T cells observed in mice vaccinated by either the
pTG18194 or the pTG18188 was significantly different from the percentage that was
observed for mice vaccinated with the negative control (pTG13135) ($p < 0.05$, Mann

Withney test). Of note, one mouse in the group of mice vaccinated with pTG18188 also displayed positive percentage of IFN γ producing CD8 $^{+}$ T cells against pool 4, pool 5 and pool 6.

As shown in Figure 3E, a weak but positive percentage of CD4 $^{+}$ T cells producing IFN γ was detected against 4 pools of peptides covering the Pol protein, the pool Pol 1, pool Pol 4, pool Pol 5 and pool Pol 6 in the two groups of mice vaccinated with pTG18194 or pTG18188, with at least 3 out of the 8 tested mice in each group displaying responses.

Responses specific of pools of peptides covering the Envelope domains

The percentage of CD8 and CD4 T cells able to produce either IFN γ alone or combined with TNF α in response to pools of peptides covering the Envelope domains, Env1 and Env2, was evaluated by ICS assay. No specific CD4 $^{+}$ T cell response was detected during this experiment. The results for CD8 $^{+}$ T cell response are shown in Figure 3F as percentages of CD8 $^{+}$ T cells specific of these pools of peptides and producing IFN γ (sum of single IFN γ producing cells or double IFN γ and TNF α producing cells). Specifically, a weak but positive percentage of CD8 $^{+}$ T cells producing IFN γ was detected for 1 mouse vaccinated with the pTG18188 against one pool of peptides, pool Env2.

2.2. Immunogenicity of HBV fusion proteins expressed by Adenovirus AdTG18201, AdTG18202 and AdTG18203

2.2.1. Evaluation of HBV-specific IFN γ producing T cells by Elispots IFN γ using pools of overlapping peptides

The immunogenicity of the HBV Pol mutant and fusion proteins expressed by human adenovirus 5 was assessed in HLA-A2 transgenic mice immunized with either AdTG18201 or AdTG18202 or AdTG18203 or AdTG15149 (empty adenovirus used as negative control). Specific T cell responses induced following one subcutaneous injection of adenovirus were evaluated by Elispot IFN γ using pools of overlapping peptides covering the HBV antigens of interest, Core (PC1-2), Polymerase (PP1-8) and Env (PE1-2) domains.

As illustrated in Figure 4, AdTG18203 encoding the HBV mutant polymerase polypeptide alone is able to induce IFN γ producing cells specific of polymerase peptide pools 4, 5, 6 and 8. All immunized mice displayed specific T cell responses with a high frequency of IFN γ producing cells mainly against the polymerase peptide pools 4 and 8.

5 As illustrated in Figure 4, AdTG18202 encoding the HBV fusion protein “tCore-Pol*” induced IFN γ producing cells specific of peptide pools PP2, PP3, PP4, PP5 and PP8, the polymerase-specific response being mainly focused on against PP2, PP3 et PP8. Immunization with AdTG18202 also resulted in the induction of high frequency of IFN γ producing cells specific for the 2 Core peptide pools PC1 and PC2 with a higher
10 frequency of T cells targeting PC1. Positive responses targeting both the polymerase and core antigens were observed in 5 out of the 5 vaccinated mice.

AdTG18201 encoding the HBV fusion protein “Core-Pol*-Env1-Env2” was also found immunogenic as illustrated in Figure 4. More specifically, IFN γ producing cells specific of polymerase peptide pools PP2, PP3, and PP8 were induced in all vaccinated
15 mice, as well as against PP4 and PP5 although with weaker spots and lower responding mice frequencies. Immunization with AdTG18201 also resulted in the induction of high frequency of IFN γ producing cells specific for the 2 Core peptide pools PC1 and PC2, in all vaccinated mice (5/5). Immunisation with AdTG18201 also induced specific T cell responses against the Env domains, even if those responses are weak and sporadic
20 with 1 out of 5 mice displaying responses targeting PE1 and 2 out of 5 mice displaying responses targeting PE2.

2.2.2. Evaluation of HBV specific IFN γ producing T cells by Elispots IFN γ using HLA-A2 peptides following immunization with AdTG18201.

25 The immunogenicity of one of the HBV fusion protein expressed by AdTG18201 was assessed in HLA-A2 transgenic mice. The animals were immunized by one subcutaneous injection of either AdTG18201 or AdTG15149 (empty adenovirus used as negative control). Specific T cell responses were evaluated by Elispot IFN γ using HLA-A2 restricted epitopes contained in Polymerase (SLY), Core (FLP and ILC)
30 and Envelope (VLQ and GLS).

As illustrated in Figure 5, AdTG18201 encoding the HBV fusion protein « Core-Pol*-Env1-Env2 » was found immunogenic. More specifically, IFN γ producing cells

specific of the polymerase HLA-A2 epitope, SLY, were induced in all AdTG18201-vaccinated mice. At the same time, AdTG18201 also induced IFN γ -producing cells specific of the 2 HLA-A2 epitopes of the Core protein, FLP and ILC, with high frequencies. Immunization with AdTG18201 also induced specific T cell responses
5 against the Env domains, although the frequencies and number of responding mice are lower, with 2 out of 8 tested mice displaying positive T cell response against the VLQ peptide and 5 out of 8 tested mice displaying positive T cell response against the GLSP peptide.

10 2.2.3. Evaluation of HBV specific IFN γ and/or TNF α producing CD8 $^{+}$ T cells by intracellular staining assays following immunization of HLA-A2 mice with the AdTG18201 and using selected pools of peptides.

The percentage of CD8 $^{+}$ T cells able to produce either IFN γ alone or combined with TNF α in response to selected pools of peptides, covering a part of the polymerase
15 protein (PP8, amino acids 725 to 835) and a part of the HBV Core protein (PC1, amino acids 1 to 100), was evaluated by ICS assay. The result is expressed as percentage of CD8 $^{+}$ T cells specific of these pools of peptides and producing IFN γ alone and IFN γ combined with TNF α .

As shown in Figure 6, AdTG18201 is specifically capable of inducing high
20 percentages of CD8 $^{+}$ T cells producing both IFN γ alone as well as IFN γ combined with TNF α recognizing peptides of PP8 and PC1 pools. All vaccinated mice displayed a high percentage of both single producing (IFN γ alone) and double producing (IFN γ and TNF α) specific CD8 $^{+}$ T cells.

Of note, similar experiments performed in another mouse model, C57Bl6 mice,
25 displayed similar results of immunogenicity of the AdTG18201 (not shown)

2.2.4. Evaluation of the induction of *in vivo* functional CD8 $^{+}$ T cells using an *in vivo* CTL assays following immunization of HLA-A2 mice with the AdTG18201.

The capacity of AdTG18201 to induce *in vivo* functional CD8 T cells displaying
30 cytolytic activity was evaluated by *in vivo* cytolytic (or CTL) assay in HLA-A2 mice following immunization with AdTG18201 or AdTG15149 (as negative control) and

using 3 of the HLA-A2 epitopes already shown as being targeted by induced CD8+ T cells producing IFN γ (SLY, FLP and ILC).

As illustrated by Figure 7, the AdTG18201 is able to induce high percentage of *in vivo* specific lysis against the polymerase epitope, SLY, with a specific lysis detected for all immunized mice and a percentage ranging from 42% to 75% (Figure 7a). It was also shown that the AdTG18201 is able to induce high percentage of *in vivo* specific lysis against the 2 tested HLA-A2 epitopes of the Core protein, FLP (Figure 7a) and ILC (Figure 7b), with percentages ranging from 32% to 69% and 3% to 64% respectively. Response against env epitope was detectable but at low levels.

These data clearly demonstrate the ability of the AdTG18201 to induce *in vivo* functional CD8+ T cells displaying cytolytic activity and targeting both the HBV polymerase and the HBV core proteins.

2.2.5. Evaluation of the induction of functional CD8+ T cells in HBV transgenic mice following immunization with AdTG18201 and using an ICS assay.

The capacity of AdTG18201 to induce functional T cells in a tolerant mouse model was evaluated in HBV transgenic mice. In fact, these mice are transgenic for the HBV genome and, thus, tolerant to HBV antigens mimicking, to some extent, the tolerance encountered in HBV chronic patients. The HBV transgenic mice were immunized by one subcutaneous injection of AdTG18201 (10^8 iu) or AdTG15149 as negative control. Induced T cells were monitored both in spleens and livers of vaccinated mice by ICS (detection of CD8+ T cells producing both IFN γ and TNF α). In this specific model, peptides identified to be reactive in C57Bl/6J mice were used to screen the induced T cell response: a pool of the VSA and the N13F peptides for the polymerase and the F13L peptide for the envelope.

As illustrated in Figure 8, functional CD8+ T cells producing both IFN γ and TNF α were detected in spleens and livers of AdTG18201-vaccinated mice, with 4 out of 5 tested mice displaying functional IFN γ /TNF α producing CD8 + T cells specific of polymerase and with 2 out of 5 tested mice displaying functional IFN γ /TNF α producing CD8+ T cells specific of Envelope in both organs. As expected, no responses were detected in mice immunized with the empty AdTG15149 or when stimulation is using an irrelevant peptide.

All together, these data demonstrate the ability of the viral vector AdTG18201 expressing a fusion protein containing a RNaseH-defective and YMDD-deleted pol mutant, env domains and core to induce functional CD8⁺ T cells, producing both IFN γ and TNF α , in a HBV tolerant model.

5

2.3 Evaluation of different doses and schedules of immunization with the AdTG18201 or the AdTG18202.

2.3.1. Adenovirus dose evaluation.

The immunogenicity of the HBV fusion protein expressed by AdTG18201 was
10 assessed in HLA-A2 transgenic mice at different doses. The animals were immunized by one subcutaneous injection of either AdTG18201 at a dose of 10⁵iu or 10⁶iu or 10⁷iu or 10⁸iu or 10⁹iu or AdTG15149 at 10⁹iu (empty adenovirus used as negative control). Specific T cell responses were evaluated by Elispot IFN γ using HLA-A2 restricted epitopes contained in Polymerase (SLY), Core (FLP and ILC) and Envelope (VLQ and
15 GLS).

As illustrated in Figure 9, AdTG18201 encoding the HBV fusion protein « Core-Pol*-Env1-Env2 » was found immunogenic when injected at doses of 10⁷iu, 10⁸iu and 10⁹iu. More specifically, no IFN γ producing cells specific of the tested HLA-A2 epitopes of Core, Polymerase or Env domains were detected with the doses of 10⁵ and
20 10⁶iu. Specific IFN γ producing cells targeting the 2 tested core epitopes and the tested epitope of Pol were detected for doses of 10⁷, 10⁸ and 10⁹ iu. A dose effect is observed for the 3 epitopes (SLY, FLP and ILC). For the 2 epitopes of the Env domains (VLQ and GLS), frequencies of IFN γ producing cells are low for doses of 10⁷ and 10⁸ whereas frequencies are clearly increased with a dose of 10⁹ iu.

25

2.3.2. Evaluation of multiple immunization schedule at short term interval.

The immunogenicity of one of the HBV fusion protein expressed by AdTG18202 was assessed in HLA-A2 transgenic mice according to different schedules of immunization. AdTG18202 was either administred once or 3 times (1 injection/week
30 during 3 weeks) or 6 times (1 injection/week during 6 weeks) and the induced immune T cell responses was assessed 2 weeks after the last injection by an Elispots IFN γ assay

and using HLA-A2 restricted epitopes, SLY (Pol) and FLP and ILC (Core). Some mice were immunized 6 times with an empty adenovirus as a negative control (not shown)

As illustrated in Figure 10, AdTG18202 encoding for the fusion protein "Core-Pol*" was found immunogenic whatever the tested schedules. More particularly, no specific T cell response was detected with medium alone and irrelevant peptide whereas high and similar frequencies of IFN γ producing cells were detected in presence of the 3 tested HBV epitopes, SLY, FLP and ILC. Frequencies of detected IFN γ producing T cells appeared comparable between groups of mice injected once, 3 times or 6 times at 1 week interval, without the appearance, at the IFN γ production level, of a T cell exhaustion due to a too high number of immunizations in a short time-interval. The adenovirus specific T cell responses appeared higher when mice were injected 6 times than when they were injected once or 3 times. As expected, no HBV-specific T cell responses were observed following immunization with an empty adenovirus.

2.3.3. Evaluation of multiple immunization schedule at long term interval.

The immunogenicity of one of the HBV fusion protein expressed by AdTG18202 was assessed in HLA-A2 transgenic mice according to different schedules of immunization. AdTG18202 was either administered once (2 (group 1) or 20 (group 2) weeks before the monitoring of T cell responses) or twice (2 injections at 2 (group 3) or 4 (group 4) month interval, monitoring of T cell responses 2 weeks after the last immunization) or three times (at 2 month interval (group 5), monitoring of T cell responses 2 weeks after the last injection). Induced T cells were monitored by an Elispots IFN γ assay and using HLA-A2 restricted epitopes, SLY (Pol) and FLP and ILC (Core). Some mice were immunized either once or three times at 2 month interval with an empty adenovirus as a negative control (not shown).

As illustrated by Figure 11, AdTG18202 encoding for the fusion protein "Core-Pol*" was found immunogenic whatever the tested schedule whereas no specific T cell response was detected following immunizations with AdTG18202 in presence of medium alone or of an irrelevant peptide. More particularly, observed specific T cell responses in group 2 showed that even if lower than those observed in group 1, 2 weeks after 1 immunization, induced T cell responses after one injection of AdTG18202 still exist 20 weeks after the injection. Observed T cell responses in group 3 and 4 showed that a 2nd immunization 2 or 4 months after the first one was able to recall T cell

responses specific of HBV epitopes at least at the level of the primary immune response observed in group 1, even slightly higher for the SLY epitope. A similar observation was done with mice immunized three times at 2 month interval (group 5) with a recall of induced T cell responses through the 2nd and 3rd injections to a level similar to the one observed in group 1. As expected, no HBV-specific T cell responses were observed following immunization with an empty adenovirus

2.4 Electron microscopy observation

A549 cells were infected *in vitro* by AdTG18201 at MOI 25, 50 or 100 and cells were collected at either 16h, 24h or 48 post-infection. Collected cells were then treated to be observed by electron microscopy.

Some virus-like particles (VLP) were observed in the nucleus and cytoplasm of AdTG18201 infected cells whereas none of these structures were observed in cells infected by an empty adenovirus. These VLP were mainly located within the nucleus. In some cells both protein aggregates and VLP were observed.

CLAIMS

1. A mutant polymerase polypeptide which comprises a mutated polymerase domain with an internal deletion that functionally disrupts the polymerase activity, wherein said internal deletion comprises at least the YMDD motif naturally present in the polymerase domain of a native HBV polymerase.
2. The mutant polymerase polypeptide according to claim 1, wherein said internal deletion is of at least 4 amino acid residues and at most 30 amino acid residues and comprises at least the YMDD motif.
3. The mutant polymerase polypeptide according to claim 1 or 2, wherein said mutated polymerase domain comprises the amino acid sequence shown in SEQ ID NO: 1 but lacks at least the Tyr residue in position 203, the Met residue in position 204, the Asp residue in position 205 and the Asp residue in position 206.
4. The mutant polymerase polypeptide according to claim 3, wherein said mutated polymerase domain comprises the amino acid sequence shown in SEQ ID NO: 1 but lacks at least the Tyr residue in position 203, the Met residue in position 204, the Asp residue in position 205, the Asp residue in position 206, the Val residue in position 207, the Val residue in position 208 and the Leu residue in position 209.
5. The mutant polymerase polypeptide according to claim 4, wherein said mutated polymerase domain comprises an amino acid sequence which exhibits at least 80% identity with the amino acid sequence shown in SEQ ID NO: 2.
6. The mutant polymerase polypeptide according to claim 5, wherein said mutated polymerase domain comprises the amino acid sequence shown in SEQ ID NO: 2.

7. The mutant polymerase polypeptide according to any one of claims 1 to 5, which further comprises a mutated RNaseH domain comprising mutation(s) of one or more amino acid residue(s) that functionally disrupt(s) the RNaseH activity normally exhibited by a native HBV polymerase.
- 5
8. The mutant polymerase polypeptide according to claim 7, wherein said one or more mutation(s) comprised in the mutated RNaseH domain are selected from the group consisting of:
- a deletion of at least 8 amino acids and at most 60 amino acids including at least the portion of SEQ ID NO: 3 extending from approximately the Glu residue (E) in position 39 to approximately the Ala (A) residue in position 46;
 - the substitution of the Asp (D) residue in position 10 of SEQ ID NO: 3 with an amino acid residue other than Asp (D);
 - 15 - the substitution of the Val (V) residue in position 90 of SEQ ID NO: 3 with an amino acid residue other than Val (V);
 - the substitution of the Thr (T) or Ala (A) residue in position 97 of SEQ ID NO: 3 with an amino acid residue other than Thr (T) or Ala (A);
 - the substitution of the Asp (D) residue in position 98 of SEQ ID NO: 3 with an amino acid residue other than Asp (D); and
 - 20 - any combination thereof.
9. The mutant polymerase polypeptide according to claim 8, wherein said substituted residue(s) in position 10, 90, 97 or 98 of SEQ ID NO: 3 are replaced with a His (H) residue or with a Tyr (Y) residue.
- 25
10. The mutant polymerase polypeptide according to claim 9, wherein said residue in position 10 of SEQ ID NO: 3 is substituted with a His (H) residue, said residue in position 90 of SEQ ID NO: 3 is substituted with a Tyr (Y) residue, said residue in position 97 of SEQ ID NO: 3 is substituted with a Tyr (Y) residue and/or said residue in position 98 of SEQ ID NO: 3 is substituted with a His (H) residue.
- 30

11. The mutant polymerase polypeptide according to any one of claims 8 to 10, wherein said deletion comprised in the mutated RNase H domain comprises a portion of at least 25 amino acids extending from approximately the Glu residue (E) in position 39 to approximately the Leu (L) residue in position 63 of SEQ ID NO: 3, and more preferably a portion of at least 33 amino acids extending from approximately the Xaa residue in position 31 to approximately the Leu (L) residue in position 63 of SEQ ID NO: 3.
12. The mutant polymerase polypeptide according to claim 11, wherein the mutated RNaseH domain comprises the amino acid sequence shown in SEQ ID NO: 3 but (a) lacking the portion of 33 amino acid residues extending from the residue Xaa in position 31 to approximately the Leu (L) residue in position 63 and comprises (b) the substitution of the Asp (D) residue in position 10 with a His (H) residue (D689H); (c) the substitution of the Val (V) residue in position 90 with a Tyr (Y) residue (V769Y); (d) the substitution of the residue in position 97 with a Tyr (Y) residue (T/A776Y) and (e) the substitution of the Asp (D) residue in position 98 with a His (H) residue (D777H).
13. The mutant polymerase polypeptide according to claim 12, which comprises a mutated RNaseH domain comprising an amino acid sequence exhibiting at least 80% identity with the amino acid sequence shown in SEQ ID NO: 4.
14. The mutant polymerase polypeptide according to any one of claims 4 to 12, which comprises an amino acid sequence exhibiting at least 80% identity with the amino acid sequence shown in SEQ ID NO: 5.
15. The mutant polymerase polypeptide according to claim 14, which comprises the amino acid sequence shown in SEQ ID NO: 5.
16. The mutant polymerase polypeptide according to any one of claims 1 to 15, used in combination with one or more additional polypeptide(s) or peptide(s).

17. The mutant polymerase polypeptide according to claim 16, wherein said one or more additional polypeptide(s) or peptide(s) is an HBV polypeptide or peptide selected from the group consisting of HBc, HBs, X protein and immunogenic fragments thereof.
18. The mutant polymerase polypeptide according to claim 17, wherein said additional HBV polypeptide(s) or peptide(s) originate from a genotype D HBV such as the Y07587 isolate.
19. A fusion protein comprising the mutant polymerase polypeptide according to any one of claims 1 to 18 and a HBV core polypeptide.
20. The fusion protein according to claim 19, wherein said HBV core polypeptide is C-terminally truncated and especially truncated at residue 148 or 149.
21. The fusion protein according to claim 19 or 20, wherein said HBV core polypeptide is fused in frame to the N-terminus of the mutant polymerase polypeptide.
22. The fusion protein according to claim 21, wherein said fusion protein comprises an amino acid sequence exhibiting at least 80% identity with the amino acid sequence shown in SEQ ID NO: 6.
23. The fusion protein according to claim 22, wherein said fusion protein comprises the amino acid sequence shown in SEQ ID NO: 6.
24. A fusion protein comprising the mutant polymerase polypeptide according to any one of claims 1 to 18 and one or more HBsAg immunogenic domain(s) or the fusion protein according to any one of claims 19 to 23 further comprising one or more HBsAg immunogenic domain(s).

25. The fusion protein according to any one of claims 19 to 24, comprising at its N-terminus a core polypeptide fused to the mutant polymerase polypeptide and one or two HbsAg immunogenic domains fused in place of the internal deletion in the mutated polymerase domain and/or in place of the deletion in the mutated RNaseH domain.
26. The fusion protein according to claim 25, comprising an amino acid sequence exhibiting at least 80% identity with the amino acid sequence shown in any of SEQ ID NO: 7 to 9.
27. The fusion protein according to claim 26, comprising the amino acid sequence shown in SEQ ID NO: 8.
28. The mutant polymerase polypeptide according to any one of claims 1 to 18 and the fusion protein according any one of claims 19 to 27, which is fused in frame to a signal peptide and to a trans-membrane peptide.
29. The mutant polymerase polypeptide and the fusion protein according to claim 28, comprising an amino acid sequence exhibiting at least 80% identity with the amino acid sequence shown in any of SEQ ID NO: 10 to 12.
30. A nucleic acid molecule coding for the polymerase mutant polypeptide according to any one of claims 1 to 18 and 28 to 29 or the fusion protein according to any one of claims 19 to 29.
31. The nucleic acid molecule according to claim 30 which is selected from the group consisting of:
- A nucleic acid molecule which encodes a mutant polymerase polypeptide comprising a polymerase domain having the amino acid sequence shown in SEQ ID NO: 1 or 2;
 - A nucleic acid molecule which encodes a mutant polymerase polypeptide comprising a RNaseH domain having the amino acid sequence shown in SEQ ID NO: 3 or 4;

- A nucleic acid molecule which encodes a mutant polymerase polypeptide comprising an amino acid sequence which exhibits at least 80% of identity with the amino acid sequence shown in SEQ ID NO: 5; or
- A nucleic acid molecule which encodes a fusion protein comprising an amino acid sequence which exhibits at least 80% of identity with the amino acid sequence shown in any of SEQ ID NO: 6-12.

32. The nucleic acid molecule according to claim 30 which comprises a nucleotide sequence exhibiting at least 80% of identity with the nucleotide sequence shown in any of SEQ ID NO: 13-17.

33. A vector comprising a nucleic acid molecule according to any one of claims 30 to 32.

34. The vector according to claim 33, wherein said vector is a plasmid or a viral vector for expression in higher eukaryotic cells or organisms.

35. The vector according to claim 34, wherein said vector is a viral vector originating from a retrovirus, adenovirus, adenovirus-associated virus (AAV), poxvirus, herpes virus, measles virus, foamy virus, alphavirus, or vesicular stomatis virus.

36. The vector according to claim 35, wherein said vector is a replication-defective adenoviral vector originating from a human or from a chimpanzee adenovirus.

37. The vector according to claim 36, wherein the nucleic acid molecule is inserted in the adenoviral E1 region and placed under the control of a CMV promoter.

38. The vector according to claim 35, wherein said vector is a poxviral vector originating from a canarypox, a fowlpox or a vaccinia virus.

39. The vector according to claim 38, wherein said vaccinia virus is the Copenhagen strain, the Wyeth strain and the modified Ankara (MVA) strain.
40. The vector according to claim 39, wherein the nucleic acid molecule is inserted
5 in deletion III of the MVA vector and placed under the control the vaccinia 7.5K or pH5R promoter.
41. The vector according to any one of claims 35 to 40, wherein said vector is selected from the group consisting of:
- 10 - A defective Ad vector comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter, and encoding a mutant polymerase polypeptide comprising an amino acid sequence as shown in SEQ ID NO: 5 or a fusion protein comprising an amino acid sequence as shown in SEQ ID
15 NO: 6 or SEQ ID NO: 8;
- A replication-defective Ad vector comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter, and comprising the nucleotide sequence shown in SEQ ID NO: 13, SEQ ID NO: 14 or SEQ ID NO: 15;
- 20 - A replication-defective Ad vector, especially a defective AdCh3 comprising inserted in place of the E1 region a nucleic acid molecule placed under the control of a promoter such as the CMV promoter and comprising the nucleotide sequence shown in SEQ ID NO: 16 or SEQ ID NO: 17;
- 25 - A MVA vector comprising a nucleic acid molecule placed under the control of a vaccinia promoter such as the 7.5K or pH5R promoter, and encoding a mutant polymerase polypeptide comprising an amino acid sequence as shown in SEQ ID NO: 5 or SEQ ID NO: 10 or a fusion protein comprising an amino acid sequence as shown in SEQ ID NO: 6,
30 SEQ ID NO: 8 or SEQ ID NO: 12; and
- A MVA vector comprising a nucleic acid molecule placed under the control of a vaccinia promoter such as the 7.5K or pH5R promoter, and

comprising the nucleotide sequence shown in SEQ ID NO: 13, SEQ ID NO: 14 or SEQ ID NO: 15.

5 42. The vector according to any one of claims 35 to 41, wherein said vector is in the form of infectious viral particles.

43. A process of producing the vector of claim 42 comprising the steps of introducing the viral vector into a suitable cell line, culturing said cell line under suitable conditions so as to allow the production of said infectious viral particle,
10 recovering the produced infectious viral particle from the culture of said cell line and optionally purifying said viral particle.

44. A host cell comprising the nucleic acid molecule according to any one of claims 30 to 32 or the vector according to any one of claims 33 to 42.

15

45. A method for recombinant production of the mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29 or the fusion protein according to any one of claims 19 to 29, comprising the steps of introducing the vector according to any one of claims 33 to 42 into a suitable host cell to produce
20 a transfected or infected host cell, culturing in vitro said transfected or infected host cell under conditions suitable for growth of the host cell, recovering the cell culture and optionally purifying the produced mutant polymerase polypeptide or fusion protein.

25 46. A composition comprising at least the mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29, the fusion protein according to any one of claims 19 to 29, the nucleic acid molecule according to any one of claims 30 to 32, the vector according to any one of claims 33 to 42, the host cell according to claim 44 or any combination thereof.

30

47. The composition according to claim 46, further comprising a pharmaceutically acceptable vehicle.

48. The composition according to claim 46 or 47, wherein said composition is formulated for intramuscular, subcutaneous, intradermal administration or scarification.
- 5 49. The composition according to any one of claims 46 to 48, wherein said composition comprises doses of about 5×10^8 , about 10^9 , about 5×10^9 , about 10^{10} , about 5×10^{10} vp or about 10^{11} vp of an adenoviral vector.
- 10 50. The composition according to any one of claims 46 to 48, wherein said composition comprises doses of about 5×10^6 , about 10^7 , about 5×10^7 , about 10^8 , or about 5×10^8 pfu of a MVA vector.
- 15 51. The mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29, the fusion protein according to any one of claims 19 to 29, the nucleic acid molecule according to any one of claims 30 to 32, the vector according to any one of claims 33 to 42, the host cell according to claim 44 or the composition according to any one of claims 46 to 50 for use for treating or preventing an HBV infection or HBV-associated diseases and pathological conditions.
- 20 52. The mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29, the fusion protein according to any one of claims 19 to 29, the nucleic acid molecule according to any one of claims 30 to 32, the vector according to any one of claims 33 to 42, the host cell according to claim 44 or the composition according to any one of claims 46 to 50 for use for treating a chronic HBV infection.
- 25 53. The mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29, the fusion protein according to any one of claims 19 to 29, the nucleic acid molecule according to any one of claims 30 to 32, the vector according to any one of claims 33 to 42, the host cell according to claim 44 or the composition according to any one of claims 46 to 50 for use for eliciting or stimulating an immune response in the treated organism.
- 30

54. The use according to claim 53, wherein said elicited or stimulated immune response is specific and/or non-specific, humoral and/or cellular.
55. The use according to claim 54, wherein said immune response is a T cell response CD4+ or CD8+-mediated or both, directed to an HBV polypeptide/epitope.
56. The use according to any one of claims 51 to 55, wherein said use comprises one or more administration(s) of a therapeutically effective amount of said mutant polymerase polypeptide, fusion protein, nucleic acid molecule, vector or composition.
57. The use according to any one of claims 51 to 55, wherein said vector is a MVA vector and said use comprises three subcutaneous administrations separated from each other by approximately one week.
58. The use according to any one of claims 51 to 55, wherein said vector is an adenoviral vector and said use comprises one or two intramuscular or subcutaneous administrations.
59. The use according to any one of claims 51 to 58, wherein said use is carried out in combination with the standard of care.
60. The use according to any one of claims 51 to 59, wherein said use is carried out according to prime boost modality.
61. The use according to claim 60, wherein the priming is carried out with a MVA vector and the boosting with an Ad vector.
62. The use according to claim 61, and wherein the MVA and/or the Ad vector encodes the fusion protein shown in SEQ ID NO: 8.

63. The use according to claim 61 or 62 comprising at least 3 subcutaneous administrations of the MVA vector separated by a period of time varying from 3 days to 3 months followed by a intramuscular or subcutaneous boost of the adenovirus vector.
- 5
64. The use according to claim 60, wherein the priming is carried out with a plasmid DNA vector and the boosting with a MVA vector.
65. The use according to claim 64, wherein the plasmid and/or the Ad vector
10 encodes the fusion protein shown in SEQ ID NO: 8.
66. The use according to claim 64 or 65, comprising at least 3 intramuscular administrations of the DNA vector separated by a period of time varying from 2 weeks to 3 months and at least one subcutaneous boost of the MVA vector.
- 15
67. A kit of parts for use in the treatment of an HBV infection or for eliciting an immune response in a subject, wherein said kit comprises a plurality of active agents selected from the group consisting of the mutant polymerase polypeptide according to any one of claims 1 to 18 and 28 to 29, the fusion protein according
20 to any one of claims 19 to 29, the nucleic acid molecule according to any one of claims 30 to 32, the vector according to any one of claims 33 to 42, the host cell according to claim 44 or the composition according to any one of claims 46 to 50.