

HERBICIDAL EMULSIFIABLE CONCENTRATE COMPOSITIONS
OF IMIDAZOLINONE HERBICIDES

BACKGROUND OF THE INVENTION

United States Patent 4,749,404 describes herbicidal liquid concentrate compositions of salts of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid which may be combined with an organic dinitroaniline composition and United States Patent 4,822,405 describes concentrated oil in water emulsion combination compositions of water soluble salts of imidazolinone herbicides and dinitroaniline herbicides. However, the ratio of herbicides which may be employed is limited because of problems with crystallization which may occur with more concentrated formulations especially when storing the compositions at low temperatures. In addition, these compositions employ salts of imidazolinones and require both aqueous and organic solvents.

REPLACED SHEETS		SG
☒	Description : paged ² to paged ⁴	
☒	Claim(s) : paged ¹⁴ to paged ¹⁷	
☐	Drawing(s) : paged to paged	
☐	Abstract : paged to paged	
DATE OF RECEIPT		

SUMMARY OF THE INVENTION

The present invention relates to a herbicidal emulsifiable concentrate composition which comprises 0.5% to 5% by weight of an imidazolinyl acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.


It is therefore an object of the present invention to advantageously provide an emulsifiable concentrate combination composition in which all of the active herbicidal agents are dissolved in a single solvent which may be stored at lower temperatures and for a longer period of time than suspension concentrate compositions or concentrated oil in water emulsion compositions.

It is also an object of the present invention to provide herbicidal emulsifiable concentrate compositions of free acids of imidazolinone herbicides and 2,6-dinitroaniline herbicides which are physically and chemically stable.

DETAILED DESCRIPTION OF THE INVENTION

A preferred embodiment of the invention is herbicidal emulsifiable concentrate compositions comprising on a weight basis about 0.5% to 5% of an imidazolinyl acid represented by structural formula I





(I)

wherein


W is oxygen or sulfur;

R₁ is C₁-C₄ alkyl;

5 R₂ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R₁ and R₂ are taken together with the carbon to which they are attached they may represent C₃-C₆ cycloalkyl optionally substituted with methyl;


10 X is hydrogen, halogen, hydroxyl or methyl;

Y and Z are independently hydrogen, halogen, C₁-C₆ alkyl, C₁-C₄ alkoxyalkyl, hydroxyloweralkyl, C₁-C₆ alkoxy, C₁-C₄ alkylthio, phenoxy, C₁-C₄ haloalkyl, nitro, cyano, C₁-C₄ alkylamino, 15 diloweralkylamino, or C₁-C₄ alkylsulfonyl group or phenyl optionally substituted with one C₁-C₄ alkyl, C₁-C₄ alkoxy or halogen; and when taken together, Y and Z may form a ring in which YZ are represented by the structure: 20 -(CH₂)_n-, where n is an integer of 3 or 4, provided that X is hydrogen; or



where L, M, Q and R are independently 25 hydrogen, halogen, C₁-C₄ alkyl or C₁-C₄ alkoxy;

0% to about 40% of a herbicidal 2,6-dinitroaniline derivative represented by structural formula II



(II)

wherein

- 5 R₃ is hydrogen, C₂-C₄ alkyl or chloroethyl;
 R₄ is C₂-C₅ alkyl, chloroethyl, 2-methallyl or
 cyclopropylmethyl;
 R₅ is hydrogen, methyl or amino;
 R₆ is trifluoromethyl, C₁-C₃ alkyl, -SO₂NH₂ or
10 SO₂CH₃;
2% to 6% of an alkylarylsulfonic acid; 2% to 6% of an
ethylene oxide/propylene oxide block copolymer; 1% to
6% of an alkylphenol polyethylene oxide condensate;
0.5% to 10% of an antigelation agent and a sufficient
15 amount of an aromatic solvent to total 100%.

A most preferred embodiment of the present invention comprises on a weight basis 0.5% to 5% of a
formula I imidazolinyl acid, 20% to 40% of a formula II
20 2,6-dinitroaniline derivative, 2% to 6% of an
alkylarylsulfonic acid, 2% to 6% of an ethylene
oxide/propylene oxide block copolymer, 1% to 6% of an
alkylphenol polyethylene oxide condensate, 0.5% to 10%
of an antigelation agent and a sufficient amount of an
25 aromatic solvent to total 100%.



Surprisingly, it has been found that the compositions of the present invention provide physically and chemically stable herbicidal emulsifiable concentrate compositions of formula (I) imidazolinyl acids and formula (II) 2,6-dinitroaniline derivatives which remain free-flowing and homogeneous for extended periods of time, and also remain physically and chemically stable after repeated freezing and thawing cycles without precipitating insoluble solids.

The alkylarylsulfonic acid of the present invention is an especially important element of the present compositions and should be present in at least an amount equal to about 1.2 times the percent by weight of the imidazolinyl acid to ensure that the acid completely dissolves in the aromatic solvent. Alkylarylsulfonic acids suitable for use in the compositions of the invention include C_8 - C_{18} alkylbenzenesulfonic acids, with dodecylbenzenesulfonic acid being most preferred.

Antigelation agents are used in the compositions of the present invention to ensure that the compositions remain homogeneous and free-flowing. Antigelaⁿ agents which are suitable for use in the compositions of the present invention include N-methylpyrrolidone, cyclohexanone, alcohols such as ethanol and methanol, glycols such as propylene glycol and ethylene glycol and the like with N-methylpyrrolidone being most preferred.

A preferred group of ethylene oxide/propylene oxide block copolymers suitable for use in the compositions of the present invention are

butyl-omega-hydroxypoly(oxypropylene)block polymer with poly(oxyethylene) having an average molecular weight in a range of 2,400 to 3,500, with alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymers having an HLB of 12 and a viscosity at 25°C of 2,000 CPS, (TOXIMUL® 8320, Stepan Chemical Co.) being a most preferred member of this class of emulsifiers.

Preferred alkylphenol polyethylene oxide condensates suitable for use in the compositions of the present invention are the nonylphenol ethoxylates, with nonylphenol ethoxylate (9 to 10 mols of ethylene oxide) (FLO MO®9N, DeSoto, Inc., Sellers Chemical Div.) being a most preferred member of this class of emulsifiers.

Aromatic solvents suitable for use in the compositions of this invention include aromatic hydrocarbon and chlorinated aromatic hydrocarbon solvents and mixtures thereof; such as toluene, xylenes, polynuclear aromatic hydrocarbons such as naphthalenes and alkyl-naphthalenes and mixtures thereof, many of which are available from the fractionation of crude oil and in general have distillation ranges in the temperature range of from about 140°C to 305°C, with those having a distillation range of from about 183°C to 280°C being most preferred, and are commercially available under a variety of tradenames; and mono or polychlorobenzene and toluenes.

Imidazolinyl acids particularly useful in the compositions of the present invention are 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinoline-carboxylic acid, 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 2-(4-isopropyl-

4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and
2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-
(methoxymethyl)nicotinic acid.

5 2,6-Dinitroaniline derivatives which are especial-
ly suitable for use in the compositions of the present
invention are pendimethalin and trifluralin.

10 The emulsifiable concentrate compositions of the
present invention may conveniently be prepared by
admixing the formula (I) imidazolinyl acid with a
solution of the desired antigelation agent, alkyl-
arylsulfonic acid, ethylene oxide/propylene oxide block
15 copolymer, alkylphenol polyethylene oxide condensate
and aromatic solvent until all of the solids are
dissolved, then the formula (II) 2,6-dinitroaniline
derivative is added to the solution and mixing is
continued until a homogeneous solution is obtained.

20 The following examples are provided to further
illustrate the compositions of the present invention
but are not limitative of the invention described
herein.

EXAMPLE 1

Preparation of herbicidal emulsifiable concentrate
compositions

5 Solid 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid (2.62g, 93.4% pure, 2.62% on a weight basis) is added to a stirred solution of N-methylpyrrolidone (4g, 4% on a weight basis) and dodecylbenzenesulfonic acid (4g, 4% on a weight basis). The mixture is stirred until all of the solids dissolve. An aromatic hydrocarbon mixture (C_{10} to C_{13}) aromatics having a distillation range of from about 226^o-279^oC (AROMATIC® 200, Exxon) (45.68g, 45.68% on a weight basis) is added to the mixture and stirring is continued until a homogeneous solution of obtained. Pendimethalin (35.7g, 92.3% pure, 35.7% on a weight basis) is added to the homogeneous solution and stirring is continued until the solids dissolve. Next, nonylphenol ethoxylate (9 to 10 mols of ethylene oxide) (4g, 4% on a weight basis) and alpha-butyl-omega-hydroxy-ethylene oxide propylene oxide block copolymer having a HLB of 12 and a viscosity at 25^oC of 2,000 CPS (4g, 4% on a weight basis) are added to the mixture and stirring is continued until a homogeneous solution is obtained which is then filtered through 1.5 micron filter paper to obtain a clear dark brown solution having a density at 20^oC of 1.0683 g/mL.

25 Following the procedure of Example 1, but using the compounds listed in Table I below gives the emulsifiable concentrate compositions listed in Table II below.

TABLE I

Herbicidal Imidazolinyl Acid

- a. 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid

5

Herbicidal 2,6-Dinitroaniline Derivative

- b. pendimethalin
- c. trifluralin

Alkylarylsulfonic Acid

10

- d. dodecylbenzene sulfonic acid

Antigelation Agent

15

- e. N-methylpyrrolidone
- f. cyclohexanone
- g. dipropylene glycol

Ethylene Oxide/propylene Oxide Block Copolymer

20

- h. alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymer having an HLB of 12 and a viscosity at 25°C of 2,000 CPS

Alkylphenol Polyethylene Oxide Condensate

25

- i. nonylphenol ethoxylate (6 mols of ethylene oxide)
- j. nonylphenol ethoxylate (9-10 mols of ethylene oxide)
- k. nonylphenol ethoxylate (11 mols of ethylene oxide)

- 1. nonylphenol ethoxylate (13 mols of ethylene oxide)
- m. nonylphenol ethoxylate (15 mols of ethylene oxide)
- 5 n. nonylphenol ethoxylate (30 mols of ethylene oxide)

Aromatic Solvent

- 10 o. aromatic hydrocarbon mixture (C_{10} to C_{13} aromatics, distillation range 226° - 279° C)
(AROMATIC®200, Exxon)
- 15 p. aromatic hydrocarbon mixture (C_9 to C_{12} aromatics, distillation range 183° - 210° C)
(AROMATIC®150, Exxon)

Emulsifiable Concentrate Compositions

Compo- sition	Imidazo- linyl acid/ % W/W	Ethylene oxide/ propylene oxide block copolymer/ oxide Condense- ment					
		2,6-Dinitro- aniline/ % W/W	Alkylaryl- sulfonic acid/ % W/W	Anti- gelation agent/ % W/W	Alkylphenol		Aromatic Solvent/ % W/W
					propylene oxide block copolymer/ oxide Condense- ment	Alkylphenol polyethylene oxide	
1	a/2.62	b/35.70	d/4.0	e/4.0	h/4.0	j/4.0	o/45.68
2	a/2.40	b/33.40	d/4.0	e/8.0	h/4.0	j/4.0	o/44.20
3	a/2.60	b/35.80	d/4.0	e/8.0	h/4.0	j/2.0	o/43.60
4	a/2.40	b/33.40	d/4.0	e/4.0	h/4.0	j/4.0	o/48.20
5	a/2.40	b/33.40	d/4.0	e/4.0	h/3.0	j/4.0	o/49.20
6	a/2.40	b/33.40	d/4.0	e/4.0	h/2.0	j/4.0	o/50.20
7	a/2.40	b/33.40	d/4.0	e/4.0	h/3.0	j/3.0	o/50.20
8	a/2.40	b/33.40	d/4.0	e/4.0	h/2.0	j/3.0	o/51.20
9	a/2.60	b/35.80	d/4.0	e/0.5	h/4.0	j/4.0	o/50.90
10	a/2.60	b/35.80	d/4.0	f/1.0	h/4.0	j/4.0	o/48.60
11	a/2.60	b/35.80	d/4.0	g/1.0	h/4.0	j/4.0	o/48.60
12	a/2.60	b/35.80	d/4.0	e/8.0	h/4.0	n/1.0	o/44.60
13	a/2.50	b/34.60	d/4.0	e/1.0	h/4.0	n/1.0	o/51.90
14	a/2.65	c/29.66	d/3.0	e/2.0	h/3.0	j/3.0	p/56.69

TABLE II (continued)

Compo- sition	Imidazo- lanyl	2,6-Dinitro- aniline/ acid/ % W/W	Alkylaryl- sulfonic acid/ % W/W	Anti- gelation agent/ % W/W	Ethylene oxide/ propylene oxide block copolymer/ % W/W		Alkylphenol polyethylene oxide Conden- sate/% W/W	Aromatic Solvent/ % W/W
15	a/2.65	c/29.66	d/3.0	e/4.0	h/3.0	j/3.0		p/54.69
16	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	i/3.0		p/54.89
17	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	k/3.0		p/54.89
18	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	l/3.0		p/54.89
19	a/2.45	c/29.66	d/3.0	e/4.0	h/3.0	m/3.0		p/54.89
20	a/2.62	c/29.66	d/4.0	e/8.0	h/4.0	j/4.0		p/47.72
21	a/2.45	c/29.44	d/3.0	e/4.0	h/3.0	j/3.0		o/55.11

EXAMPLE 2

Low temperature stability of emulsifiable concentrate compositions of the invention

Two 10mL samples of compositions 2 and 4 prepared in the above example are placed in a constant temperature bath maintained at 5°C and the temperature of the samples is allowed to equilibrate for two hours. Seeds of pendimethalin crystals and 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid powder are added to the samples. After 48 hours, the samples are visually inspected for evidence of crystallization. The temperature of the bath is then lowered in 1°C increments holding the sample at each temperature for 48 hours until crystallization is observed. A comparative control composition containing on a weight basis 31.71% pendimethalin, 2.26% 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 0.5% ammonium hydroxide, 27.42% aromatic mixture (AROMATIC® 200, Exxon), 6.04% surfactants and 32.07% water is also evaluated similarly except the initial testing temperature is set at 15°C instead of 5°C. The results of these experiments are summarized in Table III below.

TABLE III

Low Temperature Experiments


<u>Composition</u>	<u>Crystallization Temperature °C</u>
Control	13°C
2	-1°C
4	-3°C
9	0°C
11	0°C



WHAT IS CLAIMED IS:

1 1. A herbicidal emulsifiable concentrate
2 composition comprising about 0.5% to 5% by weight of an
3 imidazoliny acid; 0% to about 40% by weight of a 2,6-
4 dinitroaniline derivative; 2% to 6% by weight of an
5 alkylarylsulfonic acid; ethylene oxide/propylene oxide
6 block copolymer and alkylphenol polyethylene oxide
7 condensate emulsifying agents; an antigelation agent;
8 and an aromatic solvent.

1 2. The composition according to Claim 1
2 wherein the imidazoliny acid has the structural
3 formula (I) below



(I)

5 wherein

6 W is oxygen or sulfur;

7 R₁ is C₁-C₄ alkyl;

8 R₂ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R₁
9 and R₂ are taken together with the carbon to
10 which they are attached they may represent
11 C₃-C₆ cycloalkyl optionally substituted with
12 methyl;

13 X is hydrogen, halogen, hydroxyl or methyl;





14 Y and Z are independently hydrogen, halogen, C₁-C₆
 15 alkyl, C₁-C₄ alkoxyalkyl, hydroxyloweralkyl,
 16 C₁-C₆ alkoxy, C₁-C₄ alkylthio, phenoxy, C₁-C₄
 17 haloalkyl, nitro, cyano, C₁-C₄ alkylamino,
 18 diloweralkylamino, or C₁-C₄ alkylsulfonyl
 19 group or phenyl optionally substituted with
 20 one C₁-C₄ alkyl, C₁-C₄ alkoxy or halogen; and
 21 when taken together, Y and Z may form a ring
 22 in which YZ are represented by the structure:
 23 $-(CH_2)_n-$, where n is an integer of 3 or 4,
 24 provided that X is hydrogen; or



26 where L, M, Q and R are independently
 27 hydrogen, halogen, C₁-C₄ alkyl or C₁-C₄
 28 alkoxy;

29 and the 2,6-dinitroaniline derivative has the structur-
 30 al formula (II) below



(II)

32 wherein

33 R₃ is hydrogen, C₂-C₄ alkyl or chloroethyl;
 34 R₄ is C₂-C₅ alkyl, chloroethyl, 2-methyl or
 35 cyclopropylmethyl;
 36 R₅ is hydrogen, methyl or amino;
 37 R₆ is trifluoromethyl, C₁-C₃ alkyl, -SO₂NH₂ or
 38 SO₂CH₃.





1 3. The composition according to Claim 2
2 comprising on a weight basis 2% to 6% of the ethylene
3 oxide/propylene oxide block copolymer, 1% to 6% of the
4 alkylphenol polyethylene oxide condensate, 0.5% to 10%
5 of the antigelation agent and a sufficient amount of
6 the aromatic solvent to total 100%.

1 4. The composition according to Claim 3
2 comprising on a weight basis 20% to 40% of the
3 2,6-dinitroaniline derivative.

1 5. The composition according to Claim 4
2 wherein the alkylarylsulfonic acid is a C_8 - C_{18} alkyl-
3 benzenesulfonic acid, the ethylene oxide/propylene
4 oxide block copolymer is a butyl-omega-hydroxy-poly-
5 (oxypropylene) block polymer with poly(oxyethylene),
6 the alkylphenol polyethylene oxide condensate is a
7 nonylphenol ethoxylate, the antigelation agent is
8 N-methylpyrrolidone, cyclohexanone, an alcohol or a
9 glycol and the aromatic solvent is an aromatic hydro-
10 carbon mixture having a distillation range of from
11 about 140°C to 305°C.

1 6. The composition according to Claim 5
2 wherein the C_8 - C_{18} alkylbenzenesulfonic acid is
3 dodecylbenzenesulfonic acid, the butyl-omega-hydroxy-
4 poly(oxypropylene) block polymer with poly(oxyethylene)
5 is alpha-butyl-omega-hydroxy-ethylene oxide-propylene
6 oxide block copolymer having a HLB of 12 and a vis-
7 cosity at 25°C of 2,000 CPS, the nonylphenol ethoxylate
8 is nonylphenol ethoxylate (9 to 10 mols of ethylene
9 oxide), the antigelation agent is N-methylpyrrolidone,
10 and the aromatic hydrocarbon mixture has a distillation
11 range of from about 183°C to 280°C.





7. The composition according to Claim 5 wherein the imidazolinyl acid is selected from the group consisting of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid, 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-(methoxymethyl) nicotinic acid; and the 2,6-dinitroaniline derivative is selected from the group consisting of pendimethalin and trifluralin.

8. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is pendimethalin.

9. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is trifluralin.

10. A method of preparing a herbicidal emulsifiable concentrate composition comprising admixing 0.5% to 5% by weight of an imidazolinyl acid; 0% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% by weight of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.



HERBICIDAL EMULSIFIABLE CONCENTRATE COMPOSITIONS
OF IMIDAZOLINONE HERBICIDES

ABSTRACT OF THE INVENTION

The present invention provides herbicidal emulsifiable concentrate compositions of imidazolinone herbicides and dinitroaniline herbicides.

SUMMARY OF THE INVENTION

The present invention relates to a herbicidal emulsifiable concentrate composition which comprises 0.5% to 5% by weight of an imidazolinyl acid; 20% to about 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.

It is therefore an object of the present invention to advantageously provide an emulsifiable concentrate combination composition in which all of the active herbicidal agents are dissolved in a single solvent which may be stored at lower temperatures and for a longer period of time than suspension concentrate compositions or concentrated oil in water emulsion compositions.

It is also an object of the present invention to provide herbicidal emulsifiable concentrate compositions of free acids of imidazolinone herbicides and 2,6-dinitroaniline herbicides which are physically and chemically stable.

DETAILED DESCRIPTION OF THE INVENTION


A preferred embodiment of the invention is herbicidal emulsifiable concentrate compositions comprising on a weight basis about 0.5% to 5% of an imidazolinyl acid represented by structural formula I

REPLACEMENT SHEETS		83
☒	Description :	paged 24 to paged 27
☒	Claim(s) :	paged 14 to paged 17
☐	Drawing(s) :	paged to paged
☐	Abstract :	paged to paged
DATE OF RECEIPT		10/12/97

This is the replacement sheet

Date: 11 0 DEC 1997

20% to about 40% of a herbicidal 2,6-dinitroaniline derivative represented by structural formula II



wherein

R₃ is hydrogen, C₂-C₄ alkyl or chloroethyl;

R₄ is C₂-C₅ alkyl, chloroethyl, 2-methallyl or cyclopropylmethyl;

R₅ is hydrogen, methyl or amino;

R₆ is trifluoromethyl, C₁-C₃ alkyl, -SO₂NH₂ or SO₂CH₃;


2% to 6% of an alkylarylsulfonic acid; 2% to 6% of an ethylene oxide/propylene oxide block copolymer; 1% to 6% of an alkylphenol polyethylene oxide condensate; 0.5% to 10% of an antigelation agent and a sufficient amount of an aromatic solvent to total 100%.

A most preferred embodiment of the present invention comprises on a weight basis 0.5% to 5% of a formula I imidazolinyl acid, 20% to 40% of a formula II 2,6-dinitroaniline derivative, 2% to 6% of an alkylarylsulfonic acid, 2% to 6% of an ethylene oxide/propylene oxide block copolymer, 1% to 6% of an alkylphenol polyethylene oxide condensate, 0.5% to 10% of an antigelation agent and a sufficient amount of an aromatic solvent to total 100%.

Claims

1. A herbicidal emulsifiable concentrate composition comprising 0.5% to 5% by weight of an imidazolinyl acid; 20% to 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% by weight of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.

2. The composition according to Claim 1 wherein the imidazolinyl acid has the structural formula (I) below



(I)

wherein

W is oxygen or sulfur;

R₁ is C₁-C₄ alkyl;

R₂ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R₁ and R₂ are taken together with the carbon to which they are attached they may represent C₃-C₆ cycloalkyl optionally substituted with methyl;


X is hydrogen, halogen, hydroxyl or methyl;



This is the replacement sheet


Date: 10 DEC 1997

Y and Z are independently hydrogen, halogen, C_1-C_6 alkyl, C_1-C_4 alkoxyalkyl, hydroxyloweralkyl, C_1-C_6 alkoxy, C_1-C_4 alkylthio, phenoxy, C_1-C_4 haloalkyl, nitro, cyano, C_1-C_4 alkylamino, diloweralkylamino, or C_1-C_4 alkylsulfonyl group or phenyl optionally substituted with one C_1-C_4 alkyl, C_1-C_4 alkoxy or halogen; and when taken together, Y and Z may form a ring in which YZ are represented by the structure: $-(CH_2)_n-$, where n is an integer of 3 or 4, provided that X is hydrogen; or



where L, M, Q and R are independently hydrogen, halogen, C_1-C_4 alkyl or C_1-C_4 alkoxy;

and the 2,6-dinitroaniline derivative has the structural formula (II) below



(II)

wherein

- R_3 is hydrogen, C_2-C_4 alkyl or chloroethyl;
- R_4 is C_2-C_5 alkyl, chloroethyl, 2-methylallyl or cyclopropylmethyl;
- R_5 is hydrogen, methyl or amino;
- R_6 is trifluoromethyl, C_1-C_3 alkyl, $-SO_2NH_2$ or SO_2CH_3 .



This is the replacement sheet

Date: 10 DEC 1997

3. The composition according to Claim 2 comprising on a weight basis 2% to 6% of the ethylene oxide/propylene oxide block copolymer, 1% to 6% of the alkylphenol polyethylene oxide condensate, 0.5% to 10% of the antigelation agent and a sufficient amount of the aromatic solvent to total 100%.

4. The composition according to Claim 4 wherein the alkylarylsulfonic acid is a C_8 - C_{18} alkylbenzenesulfonic acid, the ethylene oxide/propylene oxide block copolymer is a butyl-omega-hydroxy-poly(oxypropylene) block polymer with poly(oxyethylene), the alkylphenol polyethylene oxide condensate is a nonylphenol ethoxylate, the antigelation agent is N-methylpyrrolidone, cyclohexanone, an alcohol or a glycol and the aromatic solvent is an aromatic hydrocarbon mixture having a distillation range of from 140°C to 305°C .

5. The composition according to Claim 5 wherein the C_8 - C_{18} alkylbenzenesulfonic acid is dodecylbenzenesulfonic acid, the butyl-omega-hydroxy-poly(oxypropylene) block polymer with poly(oxyethylene) is alpha-butyl-omega-hydroxy-ethylene oxide-propylene oxide block copolymer having a HLB of 12 and a viscosity at 25°C of 2,000 CPS, the nonylphenol ethoxylate is nonylphenol ethoxylate (9 to 10 mols of ethylene oxide), the antigelation agent is N-methylpyrrolidone, and the aromatic hydrocarbon mixture has a distillation range of from 183°C to 280°C .



* C L M S *

This is the replacement sheet

Date: 10 DEC 1997

6. The composition according to Claim 5 wherein the imidazolinyl acid is selected from the group consisting of 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-3-quinolinecarboxylic acid, 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid, 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and 2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-(methoxymethyl) nicotinic acid; and the 2,6-dinitroaniline derivative is selected from the group consisting of pendimethalin and trifluralin.

7. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is pendimethalin.

8. The composition according to Claim 7 wherein the imidazolinyl acid is 5-ethyl-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)nicotinic acid and the 2,6-dinitroaniline derivative is trifluralin.

9. A method of preparing a herbicidal emulsifiable concentrate composition comprising admixing 0.5% to 5% by weight of an imidazolinyl acid; 20% to 40% by weight of a 2,6-dinitroaniline derivative; 2% to 6% by weight of an alkylarylsulfonic acid; ethylene oxide/propylene oxide block copolymer and alkylphenol polyethylene oxide condensate emulsifying agents; an antigelation agent; and an aromatic solvent.



This is the replacement sheet

Date: 10 DEC 1997