



[DRAWINGS]


[FIG. 1]




[FIG. 2]




[FIG. 3]




[FIG. 4]




[FIG. 5]




[FIG. 6]



[FIG. 7]




[FIG. 8]



- Allogeneic transplantation – excipient
- △ Allogeneic transplantation – Prednisolone 2 mg/kg
- ◇ Allogeneic transplantation – Ruxolitinib 30 mg/kg
- Allogeneic transplantation – GLP-2 derivative long-acting conjugate 4613 µg/kg
- ◆ Allogeneic transplantation – Prednisolone 2 mg/kg +
Ruxolitinib 30 mg/kg (from D+1)
- Allogeneic transplantation – Prednisolone 2 mg/kg +
GLP-2 derivative long-acting conjugate 4613 µg/kg (from D+1)

[FIG. 9]



- Allogeneic transplantation – excipient
- Allogeneic transplantation – Prednisolone 2 mg/kg
- Allogeneic transplantation – Ruxolitinib 30 mg/kg
- Allogeneic transplantation – GLP-2 derivative long-acting conjugate 4613 µg/kg
- Allogeneic transplantation – Prednisolone 2 mg/kg + Ruxolitinib 30 mg/kg (from D+1)
- Allogeneic transplantation – Prednisolone 2 mg/kg + GLP-2 derivative long-acting conjugate 4613 µg/kg (from D+1)