

COMPOUNDS USEFUL AS INHIBITORS OF ISOPRENYLCYSTEINE CARBOXYL METHYLTRANSFERASE

Technical Field

5 The present invention generally relates to compounds which act as inhibitors of isoprenylcysteine carboxyl methyltransferase (ICMT). The present invention also relates to the compounds for use in therapy, methods of treating ICMT-related disorders, or uses of the compounds in the manufacture of a medicament for the treatment of ICMT-related disorders. The present invention further relates to processes for synthesizing said compounds.

10 Background Art

The members of the Ras family of signalling molecules and other proteins that terminate with a -CaaX motif undergo a three step post translational modification to carry out their cellular function. In the first step of the modification, a 15-carbon farnesyl or a 20-carbon geranylgeranyl lipid is covalently attached to the Cysteine
15 residue of the CaaX motif. This is carried out by farnesyltransferase (FTase) or geranylgeranyltransferase (GGTase) type I enzymes. The second step involves the cleaving of the three amino acids (i.e., the -aaX) by the membrane associated RceI (Ras converting enzyme 1) protease. In the last step of the modification the newly exposed isoprenylated cysteine residue is methylated by ICMT. The post translational
20 modification increases the hydrophobicity of the signalling molecules and directs it to the membrane.

The Ras family of GTPases are implicated in oncogenesis and tumour progression. K-Ras is the most commonly mutated form of Ras found in human malignancies, particularly in solid malignancies. Activating mutations in Ras have been
25 found in almost 30% of all cancers, including 50% of colon cancers and up to 90% of pancreatic cancers. The transforming abilities of Ras require processing through the prenylation pathway. Hence, the protein prenyltransferases are important targets for many drug discovery programs. Inhibitors of FTase (FTIs) have been explored in the past but have not proven effective in clinical trials. One of the grounds of lack of
30 efficacy might be due to the ability of GGTase to be effective in the post translational modification in the absence of FTase.

Genetic and pharmacological inactivation of ICMT have defined the specific requirement of CAAX motif methylation for Ras function. ICMT is required for the proper membrane targeting of H-, N-, and K-RAS. It has been shown that, inactivating
35 ICMT in mouse fibroblasts inhibited cell growth and K-RAS-induced oncogenic transformation. Further, ICMT inactivation ameliorates K-Ras induced myeloproliferative disease. Hence, inhibitors of ICMT may be a better strategy to overcome the RAS proteins from associating with the plasma membrane.

With rising evidence for the importance of ICMT-catalyzed CaaX protein methylation in oncogenesis, there is a clear need for specific pharmacological agents to target this process. However, the only such agents available to date have been analogs of the substrate, like isoprenoid-modified analogs of the minimal ICMT substrate *N*-acetyl- *S*-farnesyl cysteine prenylcysteine or indole-based small-molecule inhibitor of Icmt. These compounds either had low potency or suffered from poor physicochemical and PK properties.

The other rationale for targeting ICMT includes treating progeroid disorders. Hutchinson-Gilford progeria syndrome (HGPS) is a rare dominant genetic disorder that mimics premature, rapid aging. This disease is caused by the accumulation of a mutant form of prelamin A that cannot be processed to mature lamin A. HGPS is most commonly caused by a *de novo* point mutation in exon 11 of *LMNA*. This mutation, which occurs in codon 608, activates a cryptic splice site and leads to the in-frame deletion of 50 aa within prelamin A. This deletion leaves the CAAX motif intact; hence, the mutant prelamin A (progerin) is predicted to undergo farnesylation, release of the AAX, and carboxyl methylation. However, the site for the second endoproteolytic cleavage step, ZMPSTE24-recognition site is eliminated by the deletion.

A more severe form of progeria is caused by the loss of ZMPSTE24 protease activity through genetic mutation. In all forms of progeria, the lack of proteolysis of farnesylated and carboxymethylated prelamin A results in the accumulation of an incompletely processed molecule, termed progerin, on the nuclear envelope, where it triggers a range of molecular disturbances that lead to premature ageing. The presence of progerin adversely affects the integrity of the nuclear lamina, resulting in misshapen nuclei and nuclear blebs.

A possible treatment strategy for HGPS was tested using FTIs. A gene-targeted mouse model of HGPS was generated with genetically identical primary mouse embryonic fibroblasts and then FTIs effect on nuclear blebbing was examined. Using immunofluorescence microscopy the mislocalized progerin was observed to be away from the nuclear envelope to the nucleoplasm. So far, only modest effect has been observed in clinical trials.


Another strategy is to target ICMT in Progeria. The introduction of a hypomorphic allele of ICMT into the *Zmpste24* $-/-$ mouse model of progeria largely rescued both the premature ageing phenotype and premature death. Thus, reducing isoprenylcysteine methylation pharmacologically may be helpful for children affected by Progeria.

ICMT has emerged as a promising target for drug discovery due to its role in protein prenylation pathway, a pathway that can be targeted for cancer and premature ageing disorders.

There is therefore a need to provide compounds that inhibit ICMT, and overcome, or at least ameliorate, one or more of the disadvantages described above.

Summary

In one aspect of the present disclosure, there is provided a compound having the following Formula (I):



and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof;

wherein:

10  represents an aromatic ring system;

Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

B₁, B₂ and B₃ are independently selected from C, CH, or N;

15 Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

20 R₁, R₂, and R₃ are independently absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;


R₄ and R₅ are independently selected from H or aliphatic;

n is an integer selected from 0 to 5, wherein when n is more than 1, each R₁ substituted on Ring A may be the same or different;

m is an integer selected from 0 to 4, wherein when m is more than 1, each R₂ substituted on Ring B may be the same or different; and

5 p is an integer selected from 0 to 5, wherein when p is more than 1, each R₃
substituted on Ring C may be the same or different.

In another aspect of the present disclosure, there is provided a compound having the following Formula (I):



10 and/or tautomers, enantiomers, solvates, hydrates, prodrugs and
pharmaceutically acceptable salts thereof;

wherein:

 represents an aromatic ring system;

15 Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

B₁, B₂ and B₃ are independently selected from C, CH, or N, wherein when one of B₁, B₂ or B₃ is N, the remaining B₁, B₂ or B₃ is C or CH;

Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

20 R₁ is absent or selected from H, OH, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted

cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

5 R_2 is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF_3 , CHF_2 , optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

10 R_3 is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF_3 , CHF_2 , optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, or optionally substituted aryl;

15 R_4 and R_5 are independently selected from H or aliphatic;

n is an integer selected from 0 to 5, wherein when n is more than 1, each R_1 substituted on Ring A may be the same or different;

20 m is an integer selected from 0 to 4, wherein when m is more than 1, each R_2 substituted on Ring B may be the same or different, and m is 1 when R_2 is halogen; and

p is an integer selected from 0 to 5, wherein when p is more than 1, each R_3 substituted on Ring C may be the same or different.

25 In yet another aspect of the present disclosure, there is provided a pharmaceutical composition comprising a compound as defined herein, or a pharmaceutically acceptable form or prodrug thereof, and a pharmaceutically acceptable excipient.

30 In a further aspect of the present disclosure, there is provided a method of inhibiting methylation of isoprenylated cysteine or isoprenylcysteine caused by isoprenylcysteine carboxyl methyltransferase (ICMT) comprising contacting the compound as defined herein, or a pharmaceutical composition as defined herein, with ICMT.

35 In another aspect of the present disclosure, there is provided a method of inhibiting ICMT in a cell comprising contacting said cell with a compound as defined herein, or a pharmaceutically acceptable form or prodrug thereof, or a pharmaceutical composition as defined herein.

In yet another aspect of the present disclosure, there is provided a method of treating a ICMT-related disorder comprising administering to a subject in need of treatment a compound as defined herein, or pharmaceutical, or a composition as defined herein.

- 5 In a further aspect of the present disclosure, there is provided a compound as defined herein, or a pharmaceutically form or prodrug thereof, or a composition as defined herein for use in therapy.

- 10 In another aspect of the present disclosure, there is provided a compound as defined herein, or a pharmaceutically form or prodrug thereof, or a composition as defined herein for use in the treatment and/or prevention of cancer and/or progeroid diseases.

In yet another aspect of the present disclosure, there is provided a compound as defined herein, or a pharmaceutically form or prodrug thereof, or a composition as defined herein, for use in the treatment of a ICMT-related disorder.


- 15 In a further aspect of the present disclosure, there is provided a use of a compound as defined herein, or a pharmaceutically form or prodrug thereof, or a composition as defined herein, in the manufacture of a medicament for the treatment and/or prevention of cancer and/or progeroid diseases.

- 20 In another aspect of the present disclosure, there is provided a use of a compound as defined herein, or a pharmaceutically form or prodrug thereof, or a composition as defined herein, in the manufacture of a medicament for the treatment of a ICMT-related disorder.

In yet another aspect of the present disclosure, there is provided a process for synthesizing a compound as defined herein, comprising:

- 25 (a) reacting a compound of formula (III):


7



wherein R_1 , R_2 , R_4 , R_5 , B_1 , B_2 , B_3 , n and m are as defined herein, and R_6 is a leaving group,

5

in an organic solvent in the presence of a base with a compound of formula (IV):




wherein R_3 and p are as defined herein, and R_7 is $-OH$;

or


(b) reacting a compound of formula (III):

10



wherein R_1 , R_2 , R_4 , R_5 , B_1 , B_2 , B_3 , n and m are as defined herein, and R_6 is OH or an alkoxide,

in an organic solvent in the presence of a base with a compound of formula (IV):



(IV)

wherein R_3 and p are as defined herein, and R_7 is halogen.

Advantageously, the compounds of the present disclosure and pharmaceutical compositions comprising them may inhibit ICMT effectively at low concentrations. Said compounds and pharmaceutical compositions comprising them may be useful for the prophylaxis and/or treatment of diseases linked to ICMT. Advantageously, the compounds or pharmaceutical compositions comprising them may offer a treatment to cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS), including without limitation a cancer linked to mutant Ras overactivity such as hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukaemia, and other diseases linked to high ICMT expression.

Advantageously, compared to known compounds of the prior art, the compounds of the present disclosure may show high effectiveness in ICMT inhibition combined with a desirable excellent anti-proliferative and anti-ageing activity. The compounds of the present disclosure can further be processed well into formulation for further drug testing.

Definitions

The following words and terms used herein shall have the meaning indicated:

"Acyl" means an $R-C(=O)-$ group in which the R group may be an optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl group as defined herein. Examples of acyl include acetyl, benzoyl and amino acid derived aminoacyl. The group may be a terminal group or a bridging group. If the group is a

terminal group it is bonded to the remainder of the molecule through the carbonyl carbon.

“Aliphatic” refers to an organic moiety wherein the carbon and hydrogen atoms are arranged in saturated or unsaturated straight or branched chains, including alkanes, alkenes and alkynes.

“Alkenyl” as a group or part of a group denotes an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched preferably having 2-12 carbon atoms, more preferably 2-10 carbon atoms, most preferably 2-6 carbon atoms, in the normal chain. The group may contain a plurality of double bonds in the normal chain and the orientation about each is independently E or Z. Exemplary alkenyl groups include, but are not limited to, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl and nonenyl. The group may be a terminal group or a bridging group.

“Alkoxy” refers to an alkyl-O- group in which alkyl is as defined herein. Preferably the alkyloxy is a C₁-C₆ alkyloxy. Examples include, but are not limited to methoxy and ethoxy. The group may be a terminal group or a bridging group. The term alkoxy may be used interchangeably with the term “alkyloxy”.

“Alkyl” or “alkylene” as a group or part of a group refers to a straight or branched aliphatic hydrocarbon group, preferably a C₁-C₁₂ alkyl, more preferably a C₁-C₁₀ alkyl, most preferably C₁-C₆ unless otherwise noted. Examples of suitable straight and branched C₁-C₆ alkyl substituents include methyl, ethyl, n-propyl, 2-propyl, n-butyl, sec-butyl, t-butyl, hexyl, and the like. The group may be a terminal group or a bridging group.

“Alkynyl” as a group or part of a group means an aliphatic hydrocarbon group containing a carbon-carbon triple bond and which may be straight or branched preferably having from 2-12 carbon atoms, more preferably 2-10 carbon atoms, more preferably 2-6 carbon atoms in the normal chain. Exemplary structures include, but are not limited to, ethynyl and propynyl. The group may be a terminal group or a bridging group.

“Amino” refers to groups of the form -NR_aR_b wherein R_a and R_b are individually selected from the group including but not limited to hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, and optionally substituted aryl groups.

“Aryl” as a group or part of a group denotes (i) an optionally substituted monocyclic, or fused polycyclic, aromatic carbocycle (ring structure having ring atoms that are all carbon) preferably having from 6 to 12 atoms per ring. Examples of aryl groups include phenyl, naphthyl, and the like; (ii) an optionally substituted partially saturated bicyclic aromatic carbocyclic moiety in which a phenyl and a C₅₋₇ cycloalkyl

or C₅₋₇ cycloalkenyl group are fused together to form a cyclic structure, such as tetrahydronaphthyl, indenyl or indanyl. The group may be a terminal group or a bridging group. Typically an aryl group is a C₆-C₁₈ aryl group.

5 "Azidoalkyl" refers to an alkyl group in which alkyl is as defined herein and which is substituted by an -N₃ group. Preferably the azidoalkyl is a C₁-C₆ azidoalkyl. Examples include, but are not limited to azidomethyl and azidoethyl.

10 "Carbocyclic" refers to any stable 3, 4, 5, 6, or 7-membered monocyclic or bicyclic or 7, 8, 9, 10, 11, 12, or 13-membered bicyclic or tricyclic ring system, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Preferred carbocycles, unless otherwise specified, are cyclopropyl, cyclobutyl, cyclopentyl, 15 cyclohexyl, phenyl, naphthyl, and indanyl. When the term "carbocycle" is used, it is intended to include "aryl". Carbocycles may be optionally substituted.

20 "Cycloalkenyl" means a non-aromatic monocyclic or multicyclic ring system containing at least one carbon-carbon double bond and preferably having from 5-10 carbon atoms per ring. Exemplary monocyclic cycloalkenyl rings include cyclopentenyl, cyclohexenyl or cycloheptenyl. The cycloalkenyl group may be substituted by one or more substituent groups. A cycloalkenyl group typically is a C₃-C₁₂ alkenyl group. The group may be a terminal group or a bridging group.

25 "Cycloalkyl" refers to a saturated monocyclic or fused or bridged or spiro polycyclic, carbocycle preferably containing from 3 to 9 carbon atoms per ring, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like, unless otherwise specified. It includes monocyclic systems such as cyclopropyl and cyclohexyl, bicyclic systems such as decalin, and polycyclic systems such as adamantane. A cycloalkyl group typically is a C₃-C₁₂ alkyl group. The group may be a terminal group or a bridging group.

30 "Haloalkyl" refers to an alkyl group in which alkyl is as defined herein and the alkyl is substituted by at least one halogen. Preferably the alkyl group is substituted by 1 to 5 halogen atoms. Preferably the alkyloxy is a C₁-C₆ haloalkyl. Examples include, but are not limited to chloromethyl, fluoromethyl and trifluoromethyl. The group may be a terminal group or a bridging group.

35 "Halogen" or "halo" represents chlorine, fluorine, bromine or iodine.

"Heteroaryl" either alone or part of a group refers to groups containing an aromatic ring (preferably a 5 or 6 membered aromatic ring) having one or more heteroatoms as ring atoms in the aromatic ring with the remainder of the ring atoms

being carbon atoms. Suitable heteroatoms include nitrogen, oxygen and sulphur. Examples of heteroaryl include thiophene, benzothiophene, benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole, naphtho[2,3-b]thiophene, furan, isoindolizine, xantholene, phenoxatine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, tetrazole, indole, isoindole, 1H-indazole, purine, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, cinnoline, carbazole, phenanthridine, acridine, phenazine, thiazole, isothiazole, phenothiazine, oxazole, isooxazole, furazane, phenoxazine, 2-, 3- or 4- pyridinyl, 2-, 3-, 4-, 5-, or 8- quinolinyl, 1-, 3-, 4-, or 5- isoquinolinyl 1-, 2-, or 3- indolyl, and 2-, or 3-thiophenyl. A heteroaryl group is typically a C₁-C₁₈ heteroaryl group. A heteroaryl group may comprise 3 to 8 ring atoms. A heteroaryl group may comprise 1 to 3 heteroatoms independently selected from the group consisting of N, O and S. The group may be a terminal group or a bridging group.

“Heterocyclyl” refers to saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system containing at least one heteroatom selected from the group consisting of nitrogen, sulfur and oxygen as a ring atom. Examples of heterocyclic moieties include heterocycloalkyl and heterocycloalkenyl.

“Heterocycloalkenyl” refers to a heterocycloalkyl as defined herein but containing at least one double bond. A heterocycloalkenyl group typically is a C₂-C₁₂ heterocycloalkenyl group. The group may be a terminal group or a bridging group.

“Heterocycloalkyl” refers to a saturated monocyclic, fused or bridged or spiro polycyclic ring containing at least one heteroatom selected from nitrogen, sulfur, oxygen, preferably from 1 to 3 heteroatoms in at least one ring. Each ring is preferably from 3 to 10 membered, more preferably 4 to 7 membered. Examples of suitable heterocycloalkyl substituents include pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiofuranyl, piperidinyl, piperazinyl, tetrahydropyranyl, morpholino, 1,3-diazapane, 1,4-diazapane, 1,4-oxazepane, and 1,4-oxathiapane. A heterocycloalkyl group typically is a C₂-C₁₂ heterocycloalkyl group. A heterocycloalkyl group may comprise 3 to 9 ring atoms. A heterocycloalkyl group may comprise 1 to 3 heteroatoms independently selected from the group consisting of N, O and S. The group may be a terminal group or a bridging group.

It is understood that included in the family of compounds of Formula (I) are isomeric forms including diastereomers, enantiomers, tautomers, and geometrical isomers in “E” or “Z” configurational isomer or a mixture of E and Z isomers. It is also understood that some isomeric forms such as diastereomers, enantiomers, and geometrical isomers can be separated by physical and/or chemical methods and by those skilled in the art.

Some of the compounds of the disclosed embodiments may exist as single stereoisomers, racemates, and/or mixtures of enantiomers and/or diastereomers. All

such single stereoisomers, racemates and mixtures thereof, are intended to be within the scope of the subject matter described and claimed.

The term “optionally substituted” as used herein means the group to which this term refers may be unsubstituted, or may be substituted with one or more groups independently selected from optionally substituted aryl, optionally substituted heteroaryl, alkyl, including an alkylene bridge representing two substituents, alkenyl, alkynyl, thioalkyl, cycloalkyl, aminocycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkylalkenyl, heterocycloalkyl, cycloalkylheteroalkyl, cycloalkoxy, cycloalkylaminocarbonyl, cycloalkenyloxy, cycloamino, halo, carboxyl, haloalkyl, haloalkenyl, haloalkynyl, alkynyloxy, heteroalkyl, heteroalkyloxy, hydroxyl, hydroxyalkyl, alkoxy, alkoxyalkyl, alkyloxyalkyloxyalkyl, cycloalkylalkyloxyalkyl, thioalkoxy, alkenyloxy, haloalkoxy, haloalkenyloxy, nitro, halogen, amino, aminocarbonyl, aminocarbonylalkyl, azidoalkyl, nitroalkyl, nitroalkenyl, nitroalkynyl, nitroheterocycl, alkylamino, alkylaminocarbonyl, dialkylamino, dialkylaminoalkyl, alkenylamine, alkylcarbonylamino, aminoalkyl, alkynylamino, acyl, oxo, alkyloxy, alkyloxyalkyl, alkyloxyaryl, alkyloxycarbonyl, alkyloxycycloalkyl, alkyloxyheteroaryl, alkyloxyheterocycloalkyl, alkenoyl, alkynoyl, acylamino, diacylamino, acyloxy, alkylsulfonyloxy, heterocyclic, heterocycloalkenyl, heterocycloalkyl, heterocycloalkylalkyl, heterocycloalkylalkenyl, heterocycloalkylheteroalkyl, heterocycloalkyloxy, heterocycloalkenyloxy, heterocycloxy, heterocycloamino, haloheterocycloalkyl, alkylsulfinyl, alkylsulfonyl, alkylsulfenyl, alkylcarbonyloxy, alkylthio, acylthio, aminosulfonyl, phosphorus-containing groups such as phosphono and phosphinyl, sulfinyl, sulfinylamino, sulfonyl, sulfonylamino, alkylsulfamoyl, aryl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroalkyl, heteroarylamino, heteroaryloxy, arylalkenyl, arylalkyl, alkylaryl, alkylheteroaryl, aryloxy, arylsulfonyl, cyano, cyanate, isocyanate, -C(O)NH(alkyl), and -C(O)N(alkyl)₂. The number of carbon and hetero atoms in the groups of the optional substituents is as defined for the groups mentioned above e.g. every alkyl or alkylene moiety can be a C₁-C₁₂ alkyl, methyl, ethyl, n-propyl, 2-propyl, etc. Accordingly a haloalkoxy group is for instance defined as an alkoxy group as referred to above which is substituted by at least one halogen atom. Where the optional substituents are themselves optionally substituted there substituents can be chosen from the list of optional substituents and such substituents are not further substituted.

Preferred optional substituents are defined as one to three groups selected from halogen, C₁-C₃-alkyl, or C₃-C₇-cycloalkyl, nitro, halogen, C₁-C₄ haloalkyl having 1 to 5 halogen atoms, C₁-C₄ hydroxyalkyl, optionally halogen substituted C₃-C₇-cycloalkyl, aminocarbonyl-C₁-C₃-alkyl, optionally C₁-C₃-alkyl substituted and/or halogen substituted heteroaryl having 5 to 6 ring members and 1 to two hetero atoms selected from N or O, or optionally halogen, C₁-C₃-alkyl, C₁-C₄ hydroxyalkyl and/or azido C₁-C₃-alkyl substituted phenyl. Preferred optional substituents are also two substituents forming an optionally C₁-C₃-alkyl substituted C₁-C₄-alkylene bridge.

The term "pharmaceutically acceptable salts" refers to salts that retain the desired biological activity of the above-identified compounds, and include pharmaceutically acceptable acid addition salts and base addition salts. Suitable pharmaceutically acceptable acid addition salts of compounds of Formula (I) may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, sulfuric, and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, heterocyclic carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, fumaric, maleic, alkyl sulfonic, arylsulfonic. Additional information on pharmaceutically acceptable salts can be found in Remington's Pharmaceutical Sciences, 19th Edition, Mack Publishing Co., Easton, PA 1995. In the case of agents that are solids, it is understood by those skilled in the art that the inventive compounds, agents and salts may exist in different crystalline or polymorphic forms, all of which are intended to be within the scope of the present disclosure and specified formulae.

"Prodrug" means a compound that undergoes conversion to a compound of formula (I) within a biological system, usually by metabolic means (e.g. by hydrolysis, reduction or oxidation). For example an ester prodrug of a compound of formula (I) containing a hydroxyl group may be converted by hydrolysis *in vivo* to the parent molecule. Suitable esters of compounds of formula (I) containing a hydroxyl group, are for example formates, acetates, citrates, lactates, tartrates, malonates, oxalates, salicylates, propionates, succinates, fumarates, maleates, methylene-bis- β -hydroxynaphthoates, gentisates, isethionates, di-*p*-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, *p*-toluenesulfonates, cyclohexylsulfamates, and quinates. As another example an ester prodrug of a compound of formula (I) containing a carboxy group may be convertible by hydrolysis *in vivo* to the parent molecule. (Examples of ester prodrugs are those described by F.J. Leinweber, Drug Metab. Res., 18:379, 1987). Similarly, an acyl prodrug of a compound of formula (I) containing an amino group may be converted by hydrolysis *in vivo* to the parent molecule (Many examples of prodrugs for these and other functional groups, including amines, are described in Prodrugs: Challenges and Rewards (Parts 1 and 2); Ed V. Stella, R. Borchardt, M. Hageman, R. Oliyai, H. Maag and J Tilley; Springer, 2007).

In the context of this invention the term "administering" and variations of that term including "administer" and "administration", includes contacting, applying, delivering or providing a compound or composition of the invention to an organism, or a surface by any appropriate means.

The term "treatment", refers to any and all uses which remedy a disease state or symptoms, prevent the establishment of disease, or otherwise prevent, hinder, retard, or reverse the progression of disease or other undesirable symptoms in any way whatsoever.

The word “substantially” does not exclude “completely” e.g. a composition which is “substantially free” from Y may be completely free from Y. Where necessary, the word “substantially” may be omitted from the definition of the invention.

- 5 Unless specified otherwise, the terms "comprising" and "comprise", and grammatical variants thereof, are intended to represent "open" or "inclusive" language such that they include recited elements but also permit inclusion of additional, unrecited elements.

- 10 As used herein, the term "about", in the context of concentrations of components of the formulations, typically means +/- 5% of the stated value, more typically +/- 4% of the stated value, more typically +/- 3% of the stated value, more typically, +/- 2% of the stated value, even more typically +/- 1% of the stated value, and even more typically +/- 0.5% of the stated value.

- 15 Throughout this disclosure, certain embodiments may be disclosed in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the disclosed ranges. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, description of a range
20 such as from 1 to 6 should be considered to have specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

- 25 Certain embodiments may also be described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the disclosure. This includes the generic description of the embodiments with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

30

Brief Description of Drawings

- 35 The accompanying drawings illustrate a disclosed embodiment and serves to explain the principles of the disclosed embodiment. It is to be understood, however, that the drawings are designed for purposes of illustration only, and not as a definition of the limits of the invention.

Fig.1

[Fig. 1] shows dose response curves showing the effects of compounds on the enzymatic activity of isoprenylcysteine carboxyl methyltransferase; for compound **7** (Fig. 1A); for compound **71** (Fig. 1B); for compound **53** (Fig. 1C); for compound **70** (Fig. 1D); for compound **59** (Fig. 1E); for compound **69** (Fig. 1F); and for compound **78** (Fig. 1G).

Fig. 2


[Fig. 2] shows a Western blot image showing Prelamin A accumulation in MIAPaCa-2 cells following treatment with various concentrations of compound **59**, **69**, **70** and **71**.

Fig. 3

[Fig. 3] is the dose response curves showing the effects of compounds on the growth of MIAPaCa-2 cells; for compound **71** (Fig. 3A); for compound **53** (Fig. 3B); for compound **70** (Fig. 3C); for compound **59** (Fig. 3D); for compound **69** (Fig. 3E); and for compound **78** (Fig. 3F).

Detailed Description of Embodiments

The present invention relates to compounds having the following Formula (I):



and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof;

wherein:

 represents an aromatic ring system;

Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

B₁, B₂ and B₃ are independently selected from C, CH, or N;

5 Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

R₁, R₂, and R₃ are independently absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, 10 optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

R₄ and R₅ are independently selected from H or aliphatic;


n is an integer selected from 0 to 5, wherein when n is more than 1, each R₁ substituted on Ring A may be the same or different;

15 m is an integer selected from 0 to 4, wherein when m is more than 1, each R₂ substituted on Ring B may be the same or different; and

p is an integer selected from 0 to 5, wherein when p is more than 1, each R₃ substituted on Ring C may be the same or different.

The present invention also relates to compounds having the following Formula

20 (I):



and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof;

wherein:



represents an aromatic ring system;

Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

5 B₁, B₂ and B₃ are independently selected from C, CH, or N, wherein when one of B₁, B₂ or B₃ is N, the remaining B₁, B₂ or B₃ is C or CH;

Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

10 R₁ is absent or selected from H, OH, halogen, optionally substituted alkyl, optionally substituted alkenyl, haloalkyl, CF₃, CHF₂, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

15 R₂ is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

20 R₃ is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, or optionally substituted aryl;


25 R₄ and R₅ are independently selected from H or aliphatic;

n is an integer selected from 0 to 5, wherein when n is more than 1, each R₁ substituted on Ring A may be the same or different;

30 m is an integer selected from 0 to 4, wherein when m is more than 1, each R₂ substituted on Ring B may be the same or different, and m is 1 when R₂ is halogen; and

p is an integer selected from 0 to 5, wherein when p is more than 1, each R₃ substituted on Ring C may be the same or different.

35 The present invention further relates to compounds having the following Formula (IA):




(IA)

and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof, wherein:

- 5  represents an aromatic ring system; and


A_1 , A_2 , A_3 , A_4 and A_5 , R_1 , n , B_1 , B_2 , B_3 , R_2 , m , R_4 , R_5 , C_1 , C_2 , C_3 , C_4 , C_5 , C_6 , R_3 and p are as defined herein.

The present invention also relates to compounds of the following Formula (IE) or Formula (IF):



(IE)

19




(IF)

and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof, wherein:

- 5  represents an aromatic ring system; and

A₁, A₂, A₃, A₄ and A₅, R₁, n, B₁, B₂, B₃, R₂, m, R₄, R₅, C₁, C₂, C₃, C₄, C₅, C₆, R₃ and p are as defined herein.

In the formulas disclosed herein, Ring A may be of the formula (aa):




(aa)

wherein:

* represents the point of attachment to Ring B; and

A₁, A₂, A₃, A₄ and A₅ are independently selected from C, CH, N, O or S.

In the formulas disclosed herein, Ring A may be of the formula (ab):



(ab)

wherein:

* represents the point of attachment to Ring B; and

A₁ is N or C;


A₂ is N, S, CH, or C;

A₃ is C, CH, N, or O;

A₄ is N; and

5 A₅ is C, CH, O, or N.

In the formulas disclosed herein, Ring A may be of formula (ac):



(ac)

wherein:

10 * represents the point of attachment to Ring B; and

A₁ is N or C;

A₂ is N, S, CH, or C;

A₃ is C, CH, N, or O;

A₄ is N; and

15 A₅ is C, O, or N.

In the formulas disclosed herein, Ring A may be selected from the group consisting of formulas (a1) to (a8):



(a1)



(a2)



(a3)



(a4)



(a5)



(a6)




(a7)




(a8)


20 In the formulas disclosed herein, Ring A may be selected from the group consisting of formulas (a1') to (a8'):




(a1')




(a2')



(a3')




(a4')




(a5')



(a6')




(a7')




(a8')

In the formulas disclosed herein, Ring A may be selected from the group consisting of formulas (a1'') to (a10''):




(a1'')




(a2'')




(a3'')




(a4'')




(a5'')




(a6'')




(a7'')



(a8'')



(a9'')



(a10'')

wherein * represents the point of attachment to Ring B.

In the formulas disclosed herein, R₁ may be absent or H, OH, cyano, halogen (such as F, Cl, Br, I), optionally substituted alkyl, haloalkyl, CF₃, CHF₂, CH₂F,

optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl.

- 5 The optionally substituted alkyl of R₁ may be an optionally substituted C₁-C₁₂ alkyl, or optionally substituted C₁ alkyl, optionally substituted C₂ alkyl, optionally substituted C₃ alkyl, optionally substituted C₄ alkyl, optionally substituted C₅ alkyl, optionally substituted C₆ alkyl, optionally substituted C₇ alkyl, optionally substituted C₈ alkyl, optionally substituted C₉ alkyl, optionally substituted C₁₀ alkyl, optionally substituted C₁₁ alkyl, or optionally substituted C₁₂ alkyl. The optionally substituted alkyl may of R₁ be methyl, ethyl, propyl, butyl, pentyl, hexyl, or a haloalkyl, such as CF₃, CHF₂, or CH₂F. The optionally substituted alkoxy of R₁ may be an optionally substituted C₁-C₁₂ alkoxy, or optionally substituted C₁ alkoxy, optionally substituted C₂ alkoxy, optionally substituted C₃ alkoxy, optionally substituted C₄ alkoxy, optionally substituted C₅ alkoxy, optionally substituted C₆ alkoxy, optionally substituted C₇ alkoxy, optionally substituted C₈ alkoxy, optionally substituted C₉ alkoxy, optionally substituted C₁₀ alkoxy, optionally substituted C₁₁ alkoxy, or optionally substituted C₁₂ alkoxy. The optionally substituted cycloalkyl of R₁ may be an optionally substituted C₃-C₉ cycloalkyl, or optionally substituted C₃ cycloalkyl, optionally substituted C₄ cycloalkyl, optionally substituted C₅ cycloalkyl, optionally substituted C₆ cycloalkyl, optionally substituted C₇ cycloalkyl, optionally substituted C₈ cycloalkyl, or optionally substituted C₉ cycloalkyl. The optionally substituted cycloalkyl may be cyclopropyl. The optionally substituted cycloalkenyl of R₁ may be an optionally substituted C₃-C₉ cycloalkenyl, or optionally substituted C₃ cycloalkenyl, optionally substituted C₄ cycloalkenyl, optionally substituted C₅ cycloalkenyl, optionally substituted C₆ cycloalkenyl, optionally substituted C₇ cycloalkenyl, optionally substituted C₈ cycloalkenyl, or optionally substituted C₉ cycloalkenyl. The optionally substituted heterocycloalkyl of R₁ may be an optionally substituted heterocycloalkyl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group consisting of N, O and S. The optionally substituted aryl of R₁ may be an optionally substituted C₆-C₁₈ aryl, or optionally substituted C₆ aryl, optionally substituted C₇ aryl, optionally substituted C₈ aryl, optionally substituted C₉ aryl, optionally substituted C₁₀ aryl, optionally substituted C₁₁ aryl, or optionally substituted C₁₂ aryl, optionally substituted C₁₃ aryl, optionally substituted C₁₄ aryl, or optionally substituted C₁₅ aryl, optionally substituted C₁₆ aryl, optionally substituted C₁₇ aryl, or optionally substituted C₁₈ aryl. The optionally substituted heteroaryl of R₁ may be an optionally substituted heteroaryl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group

consisting of N, O and S. The optionally substituted alkenyl of R₁ may be an optionally substituted C₂-C₁₂ alkenyl, or optionally substituted C₂ alkenyl, optionally substituted C₃ alkenyl, optionally substituted C₄ alkenyl, optionally substituted C₅ alkenyl, optionally substituted C₆ alkenyl, optionally substituted C₇ alkenyl, optionally substituted C₈ alkenyl, optionally substituted C₉ alkenyl, optionally substituted C₁₀ alkenyl, optionally substituted C₁₁ alkenyl, or optionally substituted C₁₂ alkenyl. The optionally substituted alkynyl of R₁ may be an optionally substituted C₂-C₁₂ alkynyl, or optionally substituted C₂ alkynyl, optionally substituted C₃ alkynyl, optionally substituted C₄ alkynyl, optionally substituted C₅ alkynyl, optionally substituted C₆ alkynyl, optionally substituted C₇ alkynyl, optionally substituted C₈ alkynyl, optionally substituted C₉ alkynyl, optionally substituted C₁₀ alkynyl, optionally substituted C₁₁ alkynyl, or optionally substituted C₁₂ alkynyl.

In the formulas disclosed herein, R₂ may be absent or H, OH, cyano, halogen (such as F, Cl, Br, I), optionally substituted alkyl, haloalkyl, CF₃, CHF₂, CH₂F, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl.

The optionally substituted alkyl of R₂ may be an optionally substituted C₁-C₁₂ alkyl, or optionally substituted C₁ alkyl, optionally substituted C₂ alkyl, optionally substituted C₃ alkyl, optionally substituted C₄ alkyl, optionally substituted C₅ alkyl, optionally substituted C₆ alkyl, optionally substituted C₇ alkyl, optionally substituted C₈ alkyl, optionally substituted C₉ alkyl, optionally substituted C₁₀ alkyl, optionally substituted C₁₁ alkyl, or optionally substituted C₁₂ alkyl. The optionally substituted alkoxy of R₂ may be an optionally substituted C₁-C₁₂ alkoxy, or optionally substituted C₁ alkoxy, optionally substituted C₂ alkoxy, optionally substituted C₃ alkoxy, optionally substituted C₄ alkoxy, optionally substituted C₅ alkoxy, optionally substituted C₆ alkoxy, optionally substituted C₇ alkoxy, optionally substituted C₈ alkoxy, optionally substituted C₉ alkoxy, optionally substituted C₁₀ alkoxy, optionally substituted C₁₁ alkoxy, or optionally substituted C₁₂ alkoxy. The optionally substituted cycloalkyl of R₂ may be an optionally substituted C₃-C₉ cycloalkyl, or optionally substituted C₃ cycloalkyl, optionally substituted C₄ cycloalkyl, optionally substituted C₅ cycloalkyl, optionally substituted C₆ cycloalkyl, optionally substituted C₇ cycloalkyl, optionally substituted C₈ cycloalkyl, or optionally substituted C₉ cycloalkyl. The optionally substituted cycloalkenyl of R₂ may be an optionally substituted C₃-C₉ cycloalkenyl, or optionally substituted C₃ cycloalkenyl, optionally substituted C₄ cycloalkenyl, optionally substituted C₅ cycloalkenyl, optionally substituted C₆ cycloalkenyl, optionally substituted C₇ cycloalkenyl, optionally substituted C₈ cycloalkenyl, or optionally substituted C₉ cycloalkenyl. The optionally substituted heterocycloalkyl of R₂ may be

an optionally substituted heterocycloalkyl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group consisting of N, O and S. The optionally substituted aryl of R₂ may be an optionally substituted C₆-C₁₈ aryl, or optionally substituted C₆ aryl, optionally substituted C₇ aryl, optionally substituted C₈ aryl, optionally substituted C₉ aryl, optionally substituted C₁₀ aryl, optionally substituted C₁₁ aryl, or optionally substituted C₁₂ aryl, optionally substituted C₁₃ aryl, optionally substituted C₁₄ aryl, or optionally substituted C₁₅ aryl, optionally substituted C₁₆ aryl, optionally substituted C₁₇ aryl, or optionally substituted C₁₈ aryl. The optionally substituted heteroaryl of R₂ may be an optionally substituted heteroaryl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group consisting of N, O and S. The optionally substituted alkenyl of R₂ may be an optionally substituted C₂-C₁₂ alkenyl, or optionally substituted C₂ alkenyl, optionally substituted C₃ alkenyl, optionally substituted C₄ alkenyl, optionally substituted C₅ alkenyl, optionally substituted C₆ alkenyl, optionally substituted C₇ alkenyl, optionally substituted C₈ alkenyl, optionally substituted C₉ alkenyl, optionally substituted C₁₀ alkenyl, optionally substituted C₁₁ alkenyl, or optionally substituted C₁₂ alkenyl. The optionally substituted alkynyl of R₂ may be an optionally substituted C₂-C₁₂ alkynyl, or optionally substituted C₂ alkynyl, optionally substituted C₃ alkynyl, optionally substituted C₄ alkynyl, optionally substituted C₅ alkynyl, optionally substituted C₆ alkynyl, optionally substituted C₇ alkynyl, optionally substituted C₈ alkynyl, optionally substituted C₉ alkynyl, optionally substituted C₁₀ alkynyl, optionally substituted C₁₁ alkynyl, or optionally substituted C₁₂ alkynyl. R₂ may be H or F.

In the formulas disclosed herein, R₃ may be absent or H, OH, cyano, halogen (such as F, Cl, Br, I), optionally substituted alkyl, haloalkyl, CF₃, CHF₂, CH₂F, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl.

The optionally substituted alkyl of R₃ may be an optionally substituted C₁-C₁₂ alkyl, or optionally substituted C₁ alkyl, optionally substituted C₂ alkyl, optionally substituted C₃ alkyl, optionally substituted C₄ alkyl, optionally substituted C₅ alkyl, optionally substituted C₆ alkyl, optionally substituted C₇ alkyl, optionally substituted C₈ alkyl, optionally substituted C₉ alkyl, optionally substituted C₁₀ alkyl, optionally substituted C₁₁ alkyl, or optionally substituted C₁₂ alkyl. The optionally substituted alkoxy of R₃ may be an optionally substituted C₁-C₁₂ alkoxy, or optionally substituted C₁ alkoxy, optionally substituted C₂ alkoxy, optionally substituted C₃ alkoxy, optionally substituted C₄ alkoxy, optionally substituted C₅ alkoxy, optionally substituted C₆ alkoxy,

optionally substituted C₇ alkoxy, optionally substituted C₈ alkoxy, optionally substituted C₉ alkoxy, optionally substituted C₁₀ alkoxy, optionally substituted C₁₁ alkoxy, or optionally substituted C₁₂ alkoxy. The optionally substituted cycloalkyl of R₃ may be an optionally substituted C₃-C₉ cycloalkyl, or optionally substituted C₃ cycloalkyl, optionally substituted C₄ cycloalkyl, optionally substituted C₅ cycloalkyl, optionally substituted C₆ cycloalkyl, optionally substituted C₇ cycloalkyl, optionally substituted C₈ cycloalkyl, or optionally substituted C₉ cycloalkyl. The optionally substituted cycloalkenyl of R₃ may be an optionally substituted C₃-C₉ cycloalkenyl, or optionally substituted C₃ cycloalkenyl, optionally substituted C₄ cycloalkenyl, optionally substituted C₅ cycloalkenyl, optionally substituted C₆ cycloalkenyl, optionally substituted C₇ cycloalkenyl, optionally substituted C₈ cycloalkenyl, or optionally substituted C₉ cycloalkenyl. The optionally substituted heterocycloalkyl of R₃ may be an optionally substituted heterocycloalkyl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group consisting of N, O and S. The optionally substituted aryl of R₃ may be an optionally substituted C₆-C₁₈ aryl, or optionally substituted C₆ aryl, optionally substituted C₇ aryl, optionally substituted C₈ aryl, optionally substituted C₉ aryl, optionally substituted C₁₀ aryl, optionally substituted C₁₁ aryl, or optionally substituted C₁₂ aryl, optionally substituted C₁₃ aryl, optionally substituted C₁₄ aryl, or optionally substituted C₁₅ aryl, optionally substituted C₁₆ aryl, optionally substituted C₁₇ aryl, or optionally substituted C₁₈ aryl. The optionally substituted heteroaryl of R₃ may be an optionally substituted heteroaryl having a ring atom number of 3 to 8 (such as a ring atom number of 3, 4, 5, 6, 7, 8) and having 1 to 3 heteroatoms (such as 1, 2, 3 heteroatoms) independently selected from the group consisting of N, O and S. The optionally substituted alkenyl of R₃ may be an optionally substituted C₂-C₁₂ alkenyl, or optionally substituted C₂ alkenyl, optionally substituted C₃ alkenyl, optionally substituted C₄ alkenyl, optionally substituted C₅ alkenyl, optionally substituted C₆ alkenyl, optionally substituted C₇ alkenyl, optionally substituted C₈ alkenyl, optionally substituted C₉ alkenyl, optionally substituted C₁₀ alkenyl, optionally substituted C₁₁ alkenyl, or optionally substituted C₁₂ alkenyl. The optionally substituted alkynyl of R₃ may be an optionally substituted C₂-C₁₂ alkynyl, or optionally substituted C₂ alkynyl, optionally substituted C₃ alkynyl, optionally substituted C₄ alkynyl, optionally substituted C₅ alkynyl, optionally substituted C₆ alkynyl, optionally substituted C₇ alkynyl, optionally substituted C₈ alkynyl, optionally substituted C₉ alkynyl, optionally substituted C₁₀ alkynyl, optionally substituted C₁₁ alkynyl, or optionally substituted C₁₂ alkynyl. R₃ may be halogen (such as F, Cl, Br, I), -CN, alkoxy (such as methoxy, ethoxy, propoxy, butoxy, pentoxy), amino, haloalkyl (such as -CF₃, -CHF₂, -CH₂F), haloalkyloxy (such as -OCF₃, -OCHF₂, -OCH₂F), or alkyl (such as methyl, ethyl, propyl, butyl, pentyl, hexyl).


R₃ may be fluoro, chloro, bromo, -CN, methoxy, methyl, -NH₂, -OCF₃ or -CF₃.

- R₄ may be H or aliphatic. R₄ may be H or optionally substituted alkyl. R₄ may be H or optionally substituted C₁-C₁₂ alkyl, optionally substituted C₁-C₁₂ alkenyl, or optionally substituted C₁-C₁₂ alkynyl. R₄ may be optionally substituted C₁ alkyl, optionally substituted C₂ alkyl, optionally substituted C₃ alkyl, optionally substituted C₄ alkyl, optionally substituted C₅ alkyl, optionally substituted C₆ alkyl, optionally substituted C₇ alkyl, optionally substituted C₈ alkyl, optionally substituted C₉ alkyl, optionally substituted C₁₀ alkyl, optionally substituted C₁₁ alkyl, or optionally substituted C₁₂ alkyl. R₄ may be optionally substituted C₂ alkenyl, optionally substituted C₃ alkenyl, optionally substituted C₄ alkenyl, optionally substituted C₅ alkenyl, optionally substituted C₆ alkenyl, optionally substituted C₇ alkenyl, optionally substituted C₈ alkenyl, optionally substituted C₉ alkenyl, optionally substituted C₁₀ alkenyl, optionally substituted C₁₁ alkenyl, or optionally substituted C₁₂ alkenyl. R₄ may be optionally substituted C₂ alkynyl, optionally substituted C₃ alkynyl, optionally substituted C₄ alkynyl, optionally substituted C₅ alkynyl, optionally substituted C₆ alkynyl, optionally substituted C₇ alkynyl, optionally substituted C₈ alkynyl, optionally substituted C₉ alkynyl, optionally substituted C₁₀ alkynyl, optionally substituted C₁₁ alkynyl, or optionally substituted C₁₂ alkynyl. R₄ may be H or methyl, ethyl, propyl, butyl, pentyl, or hexyl.
- R₅ may be H or aliphatic. R₅ may be H or optionally substituted alkyl. R₅ may be H or optionally substituted C₁-C₁₂ alkyl, optionally substituted C₁-C₁₂ alkenyl, or optionally substituted C₁-C₁₂ alkynyl. R₅ may be optionally substituted C₁ alkyl, optionally substituted C₂ alkyl, optionally substituted C₃ alkyl, optionally substituted C₄ alkyl, optionally substituted C₅ alkyl, optionally substituted C₆ alkyl, optionally substituted C₇ alkyl, optionally substituted C₈ alkyl, optionally substituted C₉ alkyl, optionally substituted C₁₀ alkyl, optionally substituted C₁₁ alkyl, or optionally substituted C₁₂ alkyl. R₅ may be optionally substituted C₂ alkenyl, optionally substituted C₃ alkenyl, optionally substituted C₄ alkenyl, optionally substituted C₅ alkenyl, optionally substituted C₆ alkenyl, optionally substituted C₇ alkenyl, optionally substituted C₈ alkenyl, optionally substituted C₉ alkenyl, optionally substituted C₁₀ alkenyl, optionally substituted C₁₁ alkenyl, or optionally substituted C₁₂ alkenyl. R₅ may be optionally substituted C₂ alkynyl, optionally substituted C₃ alkynyl, optionally substituted C₄ alkynyl, optionally substituted C₅ alkynyl, optionally substituted C₆ alkynyl, optionally substituted C₇ alkynyl, optionally substituted C₈ alkynyl, optionally substituted C₉ alkynyl, optionally substituted C₁₀ alkynyl, optionally substituted C₁₁ alkynyl, or optionally substituted C₁₂ alkynyl. R₅ may be H or methyl, ethyl, propyl, butyl, pentyl, or hexyl.


R₄ and R₅ may be H; or R₄ is H and R₅ is methyl; or R₄ is methyl and R₅ is H; or R₄ is methyl and R₅ is methyl.

In the formulas disclosed herein, Ring B may be of formula (bb1) or formula (bb2):

5



(bb1)



(bb2)


wherein:

* represents the point of attachment to Ring A; and


** represents the point of attachment to the -O(CHR₄)- moiety of formula (I).

10


In the formulas disclosed herein, Ring B may be selected from the group consisting of formulas (b1) to (b5):




(b1)




(b2)



(b3)




(b4)




(b5)

15


In the formulas disclosed herein, Ring B may be selected from the group consisting of formulas (b1') to (b4'):




(b1')



(b2')




(b3')



(b4')

28




wherein:

* represents the point of attachment to Ring A; and

5 ** represents the point of attachment to the $-\text{O}(\text{CHR}_4)-$ moiety of formula (I).

In the formulas disclosed herein, Ring C may be of formula (ca):



wherein:


10  represents an aromatic ring system;

*** represents the point of attachment to the $-\text{O}(\text{CHR}_5)-$ moiety of formula (I);
and

C_1 , C_2 , C_3 , C_4 , C_5 and C_6 are independently selected from C, CH, or N.

In the formulas disclosed herein, Ring C may be of the formula (cb):

15



wherein:

○ represents an aromatic ring system;


*** represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I);

C_1 and C_2 are independently C, CH or N;

5 C_4 , C_5 , and C_6 are independently C or CH; and

C_3 is C.

In the formulas disclosed herein, Ring C may be of formula (cc):



(cc)

wherein:

○ represents an aromatic ring system;


*** represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I);

C_1 and C_2 are independently CH or N;


15 C_5 is CH; and

C_3 , C_4 , and C_6 are C.


In the formulas disclosed herein, Ring C may be selected from the group consisting of formulas (c1) to (c4):




(c1)



(c2)



(c3)




(c4)

20

In the formulas disclosed herein, Ring C may be selected from the group consisting of formulas (c1') to (c6'):


30




wherein:

*** represents the point of attachment to the $-\text{O}(\text{CHR}_5)-$ moiety of formula (I).

- 5 In the formulas disclosed herein, Ring C may be selected from the group consisting of (c1'') to (c12''):



31















wherein:









- 5 *** represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I).

















Ring A may be a 5-membered carbocyclic ring system, wherein 1 to 4 (such as 1, 2, 3, 4) carbon atoms is replaced with a heteroatom; and Ring B is of formula (bb1) or formula (bb2).


- 10 Ring A may be a 5-membered heteroaryl, and Ring B is of formula (bb1) or formula (bb2).

















The compounds of the formulas disclosed herein may be selected from the group consisting of:





















































There is provided a pharmaceutical composition comprising a compound
 5 disclosed herein, or a pharmaceutically acceptable form or prodrug thereof, and a pharmaceutically acceptable excipient.

The amount of compound in the compositions may be such that it is effective to measurably inhibit ICMT in a biological sample or in a patient. The composition may be formulated for administration to a patient in need of such composition.

10 In using the compounds, they may be administered in any form or mode which may make the compound bioavailable. One skilled in the art of preparing formulations can readily select the proper form and mode of administration depending upon the particular characteristics of the compound selected, the condition to be treated, the stage of the condition to be treated and other relevant circumstances.

15 The term "pharmaceutically acceptable excipient" may refer to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this disclosure may include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins,
 20 such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol,
 25 sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol or wool fat.

Compositions as defined above may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this disclosure may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

Pharmaceutically acceptable compositions as defined above may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Pharmaceutical compositions for parenteral injection may comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such

as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of
5 dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of micro-organisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may
10 also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminium monostearate and gelatin.

If desired, and for more effective distribution, the compounds may be
15 incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres.

The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved or dispersed in sterile water or other
20 sterile injectable medium just prior to use.

Alternatively, pharmaceutically acceptable compositions as defined above may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum
25 to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

Pharmaceutically acceptable compositions as defined above may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the
30 lower intestinal tract. Suitable topical formulations may be readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract may be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

35 For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of compounds as

defined above may include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions may be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers may include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetaryl alcohol, 2 octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

Pharmaceutically acceptable compositions as defined above may also be administered by nasal aerosol or inhalation. Such compositions may be prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

Most preferably, pharmaceutically acceptable compositions as defined above may be formulated for oral administration. Such formulations may be administered with or without food. In some embodiments, pharmaceutically acceptable compositions as defined above may be administered without food. In other embodiments, pharmaceutically acceptable compositions as defined above may be administered with food.

The amount of compound that may be combined with the carrier materials to produce a composition in a single dosage form may vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present disclosure in the composition will also depend upon the particular compound in the composition.

There is provided a method of inhibiting methylation of isoprenylated cysteine or isoprenylcysteine caused by isoprenylcysteine carboxyl methyltransferase (ICMT) comprising contacting the compound disclosed herein, or a pharmaceutical composition disclosed herein, with ICMT.

- 5 There is also provided a method of inhibiting ICMT in a cell comprising contacting said cell with a compound disclosed herein, or a pharmaceutically acceptable form or prodrug thereof, or a pharmaceutical composition disclosed herein.

The inhibition of ICMT may further comprise the inhibition of cell proliferation.

- 10 There is further provided a method of treating a ICMT-related disorder comprising administering to a subject in need of treatment a compound disclosed herein, or pharmaceutical, or a composition disclosed herein. The method may further comprise the step of administering an additional therapeutic agent to the subject.

- 15 The disorder may be cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS). The cancer may be linked to mutant Ras overactivity. The cancer may be selected from the group consisting of hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.

There is also provided a compound disclosed herein, or a pharmaceutically form or prodrug thereof, or a composition disclosed herein for use in therapy. The compound may be administered in combination with an additional therapeutic agent.

- 20 There is also provided a compound disclosed herein, or a pharmaceutically form or prodrug thereof, or a composition disclosed herein for use in the treatment and/or prevention of cancer and/or progeroid diseases.

- 25 There is also provided a compound disclosed herein, or a pharmaceutically form or prodrug thereof, or a composition disclosed herein for use in the treatment of a ICMT-related disorder. The disorder may be cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS). The cancer may be one linked to mutant Ras overactivity. The cancer may be hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.

- 30 There is also provided a use of a compound disclosed herein, or a pharmaceutically form or prodrug thereof, or a composition disclosed herein, in the manufacture of a medicament for the treatment and/or prevention of cancer and/or progeroid diseases. The medicament may be administered in combination with an additional therapeutic agent.


- 35 There is also provided a use of a compound disclosed herein, or a pharmaceutically form or prodrug thereof, or a composition disclosed herein, in the manufacture of a medicament for the treatment of a ICMT-related disorder. The

disorder may be cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS). The cancer may be one linked to mutant Ras overactivity. The cancer may be hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.

- 5 The compounds of the present invention may have an ICMT IC₅₀ of less than about 5 μ M. The compounds of the present invention may have an ICMT IC₅₀ of about 0.001 μ M to about 5 μ M.


There is also provided a process for synthesizing a compound disclosed herein, comprising:

- 10 (a) reacting a compound of formula (III):



wherein R₁, R₂, R₄, R₅, B₁, B₂, B₃, n and m are as defined herein, and R₆ is a leaving group,

- 15 in an organic solvent in the presence of a base with a compound of formula (IV):



wherein R₃ and p are as defined herein, and R₇ is -OH.

- 20 R₆ being a leaving group may be any suitable leaving group and may be a mesylate (OMs), tosylate (OTs), perfluoroalkylsulfonates (trifluoromethanesulfonate), bromide or iodide.

The organic solvent for the above process (a) may be N-methylpyrrolidone, dimethylformamide, tetrahydrofuran, ethyl acetate, acetone, acetonitrile or dimethylsulfoxide.


5 The base for the above process (a) may be selected from a variety of bases including inorganic bases or nitrogen bases such as triethylamine, pyridine, sodium hydroxide, potassium hydroxide, lithium hydroxide, ammonium hydroxide, calcium hydroxide, lithium carbonate, sodium carbonate, potassium carbonate caesium carbonate, sodium hydride or sodium bicarbonate.

10 The above process (a) may be performed at a temperature that depends on the type of solvent and base used. As a guideline, the temperature may generally be in the range of about 40 °C to about 80 °C. Process (a) may be performed at a temperature range of about 40 °C to about 80 °C, about 45 °C to about 80 °C, about 50 °C to about 80 °C, about 55 °C to about 80 °C, about 60 °C to about 80 °C, about 65 °C to about 80 °C, about 70 °C to about 80 °C, about 75 °C to about 80 °C, about 40 °C to about 75 °C, about 40 °C to about 70 °C, about 40 °C to about 65 °C, about 40 °C to about 60 °C, about 40 °C to about 55 °C, about 40 °C to about 50 °C, about 40 °C to about 45 °C, or about 40 °C, about 45 °C, about 50 °C, about 55 °C, about 60 °C, about 65 °C, about 70 °C, about 75 °C, about 80 °C, or any value or range therein,

20 The above process (a) may be performed for a time duration that depends on the type of solvent and base used. As a guideline, the time duration may generally be in the range of about 12 hours to about 20 hours. Process (a) may be performed for about 12 hours to about 20 hours, about 13 hours to about 20 hours, about 14 hours to about 20 hours, about 15 hours to about 20 hours, about 16 hours to about 20 hours, about 17 hours to about 20 hours, about 18 hours to about 20 hours, about 19 hours to about 20 hours, about 12 hours to about 19 hours, about 12 hours to about 18 hours, about 12 hours to about 17 hours, about 12 hours to about 16 hours, about 12 hours to about 15 hours, about 12 hours to about 14 hours, about 12 hours to about 13 hours, or about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, or any value or range therein.


30 There is also provided another process for synthesizing a compound disclosed herein, comprising:

(b) reacting a compound of formula (III):



wherein R_1 , R_2 , R_4 , R_5 , B_1 , B_2 , B_3 , n and m are as defined herein, and R_6 is OH or an alkoxide,

in an organic solvent in the presence of a base with a compound of formula (IV):



wherein R_3 and p are as defined herein, and R_7 is halogen.

The organic solvent for the above process (b) may be N-methylpyrrolidone, dimethylformamide, tetrahydrofuran, ethyl acetate, acetone, acetonitrile, dimethylsulfoxide, propylene carbonate or nitromethane.

The base for the above process (b) may be selected from a variety of bases including inorganic bases or nitrogen bases such as triethylamine, pyridine, sodium hydroxide, potassium hydroxide, lithium hydroxide, lithium carbonate, sodium carbonate, potassium carbonate or caesium carbonate.

The above process (b) may be performed at a temperature that depends on the type of solvent and base used. As a guideline, the temperature may generally be in the range of about 80 °C to about 120 °C. Process (b) may be performed at a temperature range of about 80 °C to about 120 °C, about 85 °C to about 120 °C, about 90 °C to about 120 °C, about 95 °C to about 120 °C, about 100 °C to about 120 °C, about 105 °C to about 120 °C, about 110 °C to about 120 °C, about 115 °C to about 120 °C, about 80 °C to about 115 °C, about 80 °C to about 110 °C, about 80 °C to about 105 °C, about 80 °C to about 100 °C, about 80 °C to about 95 °C, about 80 °C to about 90 °C, about

80 °C to about 85 °C, or about 80 °C, about 85 °C, about 90 °C, about 95 °C, about 100 °C, about 105 °C, about 110 °C, about 115 °C, about 120 °C, or any value or range therein.

5 The above process (b) may be performed for a time duration that depends on the type of solvent and base used. As a guideline, the time duration may generally be in the range of about 12 hours to about 20 hours. Process (b) may be performed for about 12 hours to about 20 hours, about 13 hours to about 20 hours, about 14 hours to about 20 hours, about 15 hours to about 20 hours, about 16 hours to about 20 hours, about 17 hours to about 20 hours, about 18 hours to about 20 hours, about 19 hours to about 20
10 hours, about 12 hours to about 19 hours, about 12 hours to about 18 hours, about 12 hours to about 17 hours, about 12 hours to about 16 hours, about 12 hours to about 15 hours, about 12 hours to about 14 hours, about 12 hours to about 13 hours, or about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, or any value or range therein.

15

Examples

Non-limiting examples of the invention will be further described in greater detail by reference to specific Examples, which should not be construed as in any way limiting the scope of the invention.

20 **Example 1: Expression and Purification of ICMT enzyme**

Recombinant human ICMT was prepared by infection of Sf9 cells with a recombinant baculovirus containing the entire open reading frame of the human ICMT cDNA. Briefly, Sf9 cells in log phase growth were diluted to 1×10^6 cells/ml and infected with recombinant baculovirus at a multiplicity of infection of 2. Cells
25 producing ICMT were harvested 66 hours post-infection. The cells are homogenized and resuspended in 5 mM NaHPO₄ (pH 7.0), and a cocktail of protease inhibitors. The membrane was pelleted, after removal of nuclei and debris, at 100,000 xg for 1 hour. Membranes were resuspended in 5 mM NaHPO₄ (pH 7.0) with 5 mM EDTA at 10 - 12 mg/ml. The suspension was stored at -80°C in multiple aliquots.

30 **Example 2: ICMT Biochemical Assay**

An ICMT enzymatic assay was developed using Promega's Methyltransferase-GloTM reagents. In the assay, ICMT catalyzes the methylation of the N-Acetyl-S-farnesyl-L-cysteine by transferring a methyl group from SAM to the farnesylated cysteine and further converts the SAM to SAH. The methyltransferase activity is
35 measured based on the amount of SAH produced from the reaction through the use of coupling enzymes that convert the SAH to ATP. The MTase-Glo detection solution then catalyzes the formation of light from ATP.

For the IC₅₀ determination, the compounds were incubated for 30 minutes with 0.04 µg/well ICMT membrane extract from Sf-9 cells. A final concentration of 2.0 µM and 20 µM of SAM and peptide were added and further incubated for 90 minutes at room temperature before adding the MTase Glo and detection reagent. Reaction signals were detected using microplate readers on luminescent mode (Safire Tecan). The IC₅₀ was determined by nonlinear regression, using GraphPad Prism version, 5.03.

Example 3: Prelamin A accumulation assay

The MIAPaCa-2 cells were seeded in 6-well plates. After seeding for 24 hours, the cells were treated with either DMSO or varying concentration of the compound and incubated for 48 hours. The cells were trypsinized and the lysate was extracted with RIPA buffer (Santa Cruz). The total protein concentration of lysate is quantified using the standard Bradford assay (Biorad protein assay, microplate standard assay).

Western blot analysis was performed using antibody against Prelamin A in MIAPaCa cells treated with compound at various concentration for 48 hours. 15.0 µg of cell lysate was run on a SDS-PAGE and then transferred to nitrocellulose membrane. The membrane was then incubated in blocking buffer [PBS (phosphate buffered saline) with 0.1% Tween 20 and 5% dry milk] at room temperature for 1 hour, followed by incubation with Prelamin A antibody (rat monoclonal antibody) at 1:2500 dilution in PBS, 0.1% Tween 20 and 5% dry milk overnight at 4 °C, followed by three washes (15 minutes each wash) in PBS, 0.1% Tween 20 on the next day.

The nitrocellulose membrane was then incubated with secondary antibody solution at room temperature for 1 hour, washed three times before developed with enhanced Chemiluminescence (ECL) mixture (Amersham, Aylesbury, United Kingdom), incubated for 5 minutes and exposed using FluorChem E System instrument (Protein Simple).

Example 4: Soft Agar Assay

MIAPaCa-2 cells were purchased from ATCC. For the soft agar assay, 600 µL of 0.6% agar was added to 24-well plate to form the base layer. This is followed by the addition of 500 µL of 0.36% agar middle layer (containing MIAPaCa-2 cells). Lastly, 500 µL of fresh growth medium (containing the corresponding serially diluted compound) was added above the middle layer. The plates were incubated at 37 °C with 5% carbon dioxide in a humidified incubator for 1 to 2 weeks, with media changes every 3 days. 70 µL of thiazolyl blue tetrazolium bromide (5 mg/mL, Sigma Catalogue No: M5655) was added to each well and the plates were incubated at room temperature overnight. The plates with colonies imaged with dissecting microscope (4 X magnification). The images were then analysed with ImageJ software for colony counts, which were plotted against compound concentrations using Graphpad Prism software.

In addition, the software was used to perform non-linear curve fitting and the calculation of IC₅₀.

Example 5: Cell Proliferation Assay


Cell proliferation assay was performed using CellTiter-Glo Luminescent Cell Viability Assay (Promega) following manufacturer's instructions. The MIAPaCa-2 cells (ATCC) were treated with compounds that were serially diluted in the media (DMEM). Plates were incubated for 3 days at 37 °C in 5% CO₂. After 3 days, an equal volume of Cell Titer Glo reagent was added. Plates were rocked on a rotator for 2 hours. 100 µL of each well was transferred to a 96-well opaque plate, and luminescence emitted was measured with the Tecan Safire II. The data for selected compounds is summarized in Table 1.

Table 1. Summary of MIAPaCa-2 cells proliferation assay data for selected compounds.

Compound	GI ₅₀ (µM)
71	12.66
53	2.2
70	5.99
59	5.52
69	5.05
78	6.02

Example 6: Preparation of phenol intermediates

General procedure 1




To a well stirred solution of the respective halide, **1** (1 molar equiv.) in 4:1 ratio of 1,4-dioxane-water (0.6 M) were added boronic acid, **2** (1.2 molar equiv.), K₃PO₄ (2 molar equiv.) and Pd(dppf)Cl₂.DCM (0.05 molar equiv.) sequentially at room temperature and degassed with argon for 5 minutes. The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled to room temperature, poured into

ice-cold water and extracted with ethyl acetate. The combined organics was washed with water, brine, dried over sodium sulphate, filtered and concentrated under reduced pressure to give the crude product. This crude product was purified by column chromatography to afford the desired product, **3**.


- 5 To a well stirred solution of **3** in excess HBr in acetic acid was heated at 80 °C for 16 hours. The reaction mixture was concentrated under reduced pressure. The crude material was basified with saturated sodium bicarbonate and pH was adjusted to 8 and extracted with ethyl acetate. The combined organics was washed with water and brine solution, dried over anhydrous sodium sulphate, filtered, concentrated under reduced
10 pressure to afford the desired phenol, **4**.

Preparation of 3-(thiazol-5-yl)phenol



- ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.75 (s, 1H), 8.07 (s, 1H), 7.33 (t, J = 8.0 Hz, 1H),
15 7.20-7.15 (m, 1H), 7.11 (t, J = 2.2 Hz, 1H), 6.91-6.88 (m, 1H), 3.86 (s, 3H). MS (ESI) m/z 178.2 [C₉H₇NOS+H]⁺.


Preparation of 3-(1-methyl-1H-pyrazol-5-yl) phenol



- 20 ¹H NMR (400 MHz, CDCl₃) δ (ppm): 7.52 (d, J = 2.0 Hz, 1H), 7.32-7.25 (m, 1H), 6.96-6.86 (m, 3H), 6.38 (t, J = 2.4 Hz, 1H), 6.30 (dd, J = 6.8, 2.0 Hz, 1H), 3.93 (s, 3H). MS (ESI) m/z 175.2 [C₁₀H₁₀N₂O+H]⁺.


Preparation of 2-(1-methyl-1H-pyrazol-5-yl) pyridin-4-ol

25




- ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.8 (s, 1H), 8.49 (d, J = 6.0 Hz, 1H), 7.49 (s, 1H), 7.07 (s, 1H), 6.79 (t, J = 2.0 Hz, 1H), 6.54 (s, 1H), 4.2 (s, 3H). MS (ESI) m/z 176.2 [C₉H₉N₃O+H]⁺.
30


35

Preparation of 3-(isoxazol-4-yl)phenol


¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.66 (d, J = 6.8 Hz, 1H), 8.52 (d, J = 3.6 Hz, 1H), 7.27 (t, J = 7.6 Hz, 1H), 7.05 (d, J = 8.0 Hz, 1H), 6.81 (dd, J = 8.4, 2.0 Hz, 1H), 4.98 (s, 1H). MS (ESI) m/z 162.2 [C₉H₇NO₂+H]⁺.

Preparation of 3-(isoxazol-4-yl)phenol


¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.66 (d, J = 6.8 Hz, 1H), 8.52 (d, J = 3.6 Hz, 1H), 7.27 (t, J = 7.6 Hz, 1H), 7.05 (d, J = 8.0 Hz, 1H), 6.81 (dd, J = 8.4, 2.0 Hz, 1H), 4.98 (s, 1H). MS (ESI) m/z 162.2 [C₉H₇NO₂+H]⁺.

Preparation of 3-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenol


¹H NMR (DMSO-d₆, 400 MHz): δ 10.527 (brs, 1H), 7.434 (d, 1H, J = 1.6 Hz), 6.765 (d, 1H, J = 9.6 Hz), 6.715 (s, 1H), 6.623 (dd, 1H, J = 10.8 & 2.0 Hz), 6.379 (d, 1H, J = 1.6 Hz), 3.815 (s, 3H). MS (ESI) m/z 193.1 [C₁₀H₉FN₂O+H]⁺.

Preparation of 5-(1-methyl-1H-pyrazol-5-yl)pyridin-3-ol


MS (ESI) m/z 176.1 [C₉H₉N₃O]⁺.

Preparation of 6-(1-methyl-1H-pyrazol-5-yl)pyridin-2-ol

¹H NMR (CDCl₃, 400 MHz): δ 7.542 (t, 0.5H, J = 8.0 Hz), 7.444 (t, 0.5H, J = 8.0 Hz), 7.082 (d, 0.5H, J = 7.2 Hz), 6.924 (d, 0.5H, J = 7.2 Hz), 6.735 (d, 0.5H, J = 8.0 Hz), 6.703 (d, 0.5H, J = 8.0 Hz), 4.462-4.440 (m, 2H), 3.964-3.942 (m, 2H). MS (ESI) m/z 176.1 [C₉H₉N₃O+H]⁺.


Preparation of 4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenol

^1H NMR (CDCl_3 , 400 MHz): δ 7.54-7.53 (d, J = 1.6 Hz, 1H), 7.10 (t, J = 9.0 Hz, 1H), 6.97-6.93 (m, 1H), 6.87-6.85 (m, 1H), 6.32-6.31 (d, J = 2.0 Hz, 1H), 4.09-4.07 (m, 2H), 3.98-3.96 (m, 2H), 3.82 (s, 3H). MS (ESI) m/z 193.1 $[\text{C}_{10}\text{H}_9\text{FN}_2\text{O}+\text{H}]^+$.


General procedure 2

To a well stirred solution of 1-bromo-3-methoxybenzene, **1** (1 molar equiv.) in DMF (0.85 M) were added the respective amine, **1a** (1.5 molar equiv.) and caesium carbonate (2 molar equiv.) and copper(I) iodide (0.1 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured into water and extracted with ethyl acetate. The combined organics was washed with water and brine, dried over sodium sulphate and concentrated under reduced pressure. The crude product was purified by column chromatography to afford the desired intermediate, **2**.

To a well stirred solution of **2** in excess HBr in acetic acid was heated at 80 °C for 16 hours. The reaction mixture was concentrated under reduced pressure. The crude material was basified with saturated sodium bicarbonate and pH was adjusted to 8 and extracted with ethyl acetate. The combined organics was washed with water and brine solution, dried over anhydrous sodium sulphate, filtered, concentrated under reduced pressure to afford the desired phenol, **3**.


Preparation of 3-(1H-1,2,4-triazol-1-yl)phenol

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 9.97 (s, 1H), 9.22 (s, 1H), 8.19 (s, 1H), 7.34-7.24 (m, 3H), 6.80-6.78 (m, 1H). MS (m/z): 162.2 $[\text{C}_8\text{H}_7\text{N}_3\text{O}+\text{H}]^+$.


Preparation of 3-(4H-1,2,4-triazol-4-yl)phenol

^1H NMR (400 MHz, CDCl_3) δ (ppm): 9.94 (s, 1H), 9.23 (s, 1H), 8.19 (s, 1H), 7.35-7.24 (m, 3H), 6.93 (dd, $J = 1.6$ Hz, 8.0 Hz, 1H). MS (ESI) m/z 162.2 $[\text{C}_8\text{H}_7\text{N}_3\text{O}+\text{H}]^+$.


5

Preparation of 2-(3-methyl-4H-1,2,4-triazol-4-yl)pyridin-4-ol

MS (ESI) m/z 177.1 $[\text{C}_8\text{H}_8\text{N}_4\text{O}]^+$.


10 **Preparation of 2-(1H-tetrazol-1-yl)pyridin-4-ol**

MS (ESI) m/z 177.1 $[\text{C}_6\text{H}_5\text{N}_5\text{O}]^+$.

Preparation of 3-(2-methyl-1H-imidazol-1-yl)phenol


15

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 9.91 (s, 1H), 7.32-7.28 (m, 1H), 7.23 (m, 1H), 6.68 (m, 1H), 6.84-6.81 (m, 2H), 6.75 (m, 1H), 2.26 (s, 3H); MS (ESI) m/z 175.1 $[\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}+\text{H}]^+$.


20 **Preparation of 4-methoxy-2-(2-methyl-1H-imidazol-1-yl)pyridine**

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 11.08 (s, 1H), 8.23 (d, $J = 6.0$ Hz, 1H), 7.45 (d, $J = 1.6$ Hz, 1H), 6.87-6.80 (m, 3H), 2.23 (s, 3H). MS (ESI) m/z 176.2 $[\text{C}_9\text{H}_9\text{N}_3\text{O}+\text{H}]^+$.


25

Preparation of 6-(2-methyl-1H-imidazol-1-yl)pyridin-2-ol


MS (ESI) m/z 176.0 $[\text{C}_9\text{H}_9\text{N}_3\text{O}]^+$.

Preparation of 3-(2-cyclopropyl-1H-imidazol-1-yl)phenol


¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 9.89 (s, 1H), 7.32 (t, J = 8.4 Hz, 1H), 7.19 (d, J = 1.1 Hz, 1H), 6.89 (dd, J = 1.4 and 2.5 Hz, 1H), 6.84-6.82 (m, 3H), 1.80-1.74 (m, 5
1H), 0.88-0.82 (m, 4H). MS (ESI) m/z 201.2 [C₁₂H₁₂N₂O+H]⁺.

Preparation of 4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenol

¹H NMR (DMSO, 400 MHz,) δ 9.88 (s, 1H), 7.27 (t, J = 9.4 Hz, 1H), 7.22 (s, 1H), 6.92
10 (s, 1H), 6.91-6.87 (m, 1H), 6.80-6.78 (m, 1H), 2.17 (s, 3H). MS (ESI) m/z 192.8 [C₁₀H₉FN₂O+H]⁺.

Preparation of Ethyl 2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propanoate

15 To a well stirred solution of 3-(2-methyl-1H-imidazol-1-yl)phenol (1 molar equiv.) in DMF (0.3 M) were added ethyl 2-bromopropanoate (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured in to water (200 mL) and extracted with EtOAc (3 x 200 mL). The combined organic layer was washed with water and brine, dried over Na₂SO₄
20 and concentrated under reduced pressure. The crude product was purified by column chromatography to afford ethyl 2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propanoate as a colorless gummy liquid.
1H NMR (400 MHz, CDCl₃) δ (ppm): 7.37 (t, J = 8.0 Hz, 1H), 7.01-6.89 (m, 4H), 6.81 (d, J = 1.6 Hz, 1H), 4.78-4.73 (m, 1H), 4.25-4.20 (s, 2H), 2.37 (d, J = 6.0 Hz, 3H), 1.28
25 (t, J = 7.6 Hz, 3H).


Preparation of 2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-1-ol

To a well stirred solution of ethyl 2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propanoate (0.07 M) in THF (20 mL) were added LAH (2 M in THF) (2 molar equiv.). The reaction mixture was cooled at 0 °C to room temperature for 16 hours. The reaction mixture was cooled and quenched with saturated NaSO₄ and extracted with EtOAc. The combined organic layer was washed with water and brine,
30

dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford 2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-1-ol.

1H NMR (400 MHz, CDCl₃) δ (ppm): 7.38 (t, 1H), 7.05-6.90 (m, 4H), 6.85 (d, 1H), 4.58 (m, 1H), 3.78 (s, 2H), 2.37 (s, 3H), 1.28 (s, 3H).


Preparation of 1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-ol



To a well stirred solution of 3-(1-methyl-1H-pyrazol-5-yl) phenol (500 mg, 2.8 mmol) in DMSO (0.5 M) were added 1-chloropropan-2-ol (1.5 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured in to water and extracted with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford 1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-ol.


1H NMR (400 MHz, CDCl₃) δ (ppm): 7.50 (d, *J* = 1.6 Hz, 1H), 7.39-7.24 (m, 1H), 7.02-6.95 (m, 3H), 6.29 (d, *J* = 2.0 Hz, 1H), 4.25-4.20 (m, 1H), 4.04-3.96 (m, 1H), 3.96-3.82 (m, 4H), 2.31 (d, *J* = 3.6 Hz, 1H), 1.31-1.24 (m, 3H).

Preparation of 1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-ol




To a well stirred solution of 3-(2-methyl-1H-imidazol-1-yl) phenol (1 equiv.) in DMSO (0.2 M) were added 1-chloropropan-2-ol (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured in to water and extracted with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford 1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-ol.


1H NMR (400 MHz, CDCl₃) δ (ppm): 7.37 (t, *J* = 8.0 Hz, 1H), 7.27 (d, *J* = 1.2 Hz, 1H), 7.03-6.97 (m, 3H), 6.89 (d, *J* = 1.2 Hz, 1H), 4.89 (d, *J* = 4.4 Hz, 1H), 3.97-3.84 (m, 3H), 2.29 (s, 3H), 1.15 (d, *J* = 6.4 Hz, 3H).

Preparation of (S)-1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-ol

- To a well stirred solution of 3-(2-cyclopropyl-1H-imidazol-1-yl)phenol (1 molar equiv.) in DMSO (0.3 M) were added (S)-1-chloropropan-2-ol (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured in to water and extracted with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford (S)-1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-ol.
- 1H NMR (400 MHz, CDCl₃) δ (ppm): 7.41-7.36 (m, 1H), 7.02 (t, *J* = 8.0 Hz, 4H), 4.25-4.20 (m, 1H), 3.97 (dd, *J* = 4.0, 12.0 Hz, 1H), 3.85 (t, *J* = 8.0 Hz, 1H), 2.34 (s, 1H), 1.82-1.75 (m, 1H), 1.30 (d, *J* = 4.0 Hz, 3H), 1.12-1.09 (m, 2H), 0.92-0.87 (m, 2H); MS (ESI) *m/z* 259.21 [C₁₅H₁₈N₂O₂+H]⁺.

15 Preparation of (R)-1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-ol

- To a well stirred solution of 3-(2-cyclopropyl-1H-imidazol-1-yl)phenol (1 molar equiv.) in DMSO (0.3 M) were added (R)-1-chloropropan-2-ol (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured in to water and extracted with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford (R)-1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-ol.
- 1H NMR (400 MHz, CDCl₃) δ (ppm): 7.41-7.36 (m, 1H), 7.02 (t, *J* = 8.0 Hz, 4H), 4.25-4.20 (m, 1H), 3.97 (dd, *J* = 4.0, 12.0 Hz, 1H), 3.85 (t, *J* = 8.0 Hz, 1H), 2.34 (s, 1H), 1.82-1.75 (m, 1H), 1.30 (d, *J* = 4.0 Hz, 3H), 1.12-1.09 (m, 2H), 0.92-0.87 (m, 2H); MS (ESI) *m/z* 259.21 [C₁₅H₁₈N₂O₂+H]⁺.


Preparation of 3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-one

- To a well stirred solution of 3-(2-methyl-1H-imidazol-1-yl)phenol (1 molar equiv.) in DMSO (0.14 M) were added 3-bromobutan-2-one (1.5 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction

mixture was cooled and poured in to water and extracted with EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to afford 3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-one.

- 5 MS (ESI) m/z 245.2 [C₁₄H₁₆N₂O₂+H]⁺.

Preparation of 3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-ol




- 10 To a well stirred solution of 3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-one (1 molar equiv.) in methanol (0.12 M) was added NaBH₄ (1.5 molar equiv.) at 0-5 °C and stirred the reaction mixture at room temperature for 4 hours. The reaction mixture was concentrated and diluted with water and extracted into EtOAc. The combined organic layer was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure to give crude product. This crude product was purified by column chromatography to afford 3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-ol.
- 15 MS (ESI) m/z 247.2 [C₁₄H₁₈N₂O₂+H]⁺.

Example 7: Preparation of final compounds

General Procedure A

20




- 25 To a well stirred solution of the respective phenol, **1** (1 molar equiv.) in DMSO (0.25 M) were added 2-chloroethanol (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. Upon completion of reaction, the reaction mixture was cooled and poured into water and extracted with ethyl acetate. The combined organics was washed with water and brine, dried over sodium sulphate and concentrated under reduced pressure. The crude product was purified by column chromatography to afford the desired product, **2**.

To a solution of the respective compound 2 (1 molar equiv.) in dichloromethane (0.4 M) was added trimethylamine (3 molar equiv.) and tosyl chloride (1.5 molar equiv.) at 0 °C. The reaction mixture was stirred at room temperature for 16 hours. Upon completion of the reaction, the crude product was extracted with dichloromethane and washed with water. The combined organics was dried over sodium sulphate and concentrated under reduced pressure to obtain the crude compound which was purified by column chromatography to afford the respective tosylate, **3**.


To a well stirred solution of the respective phenol, **4** (1 molar equiv.) in DMF (0.15 M) was added the respective tosylate, **3** (1.1 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 60 °C for 16 hours. Upon completion of reaction, the reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude product was diluted with ethyl acetate and washed with water. The combined organics was dried over sodium sulphate and washed concentrated under reduced pressure to obtain the crude product which was then purified further by column chromatography.

Preparation of 4-(2-(3-(1H-1,2,4-triazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 1




¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.55 (s, 1H), 8.11 (s, 1H), 7.63-7.60 (m, 2H), 7.43 (t, *J* = 8.0 Hz, 1H), 7.34 (t, *J* = 2.4 Hz, 1H), 7.29-7.28 (m, 1H), 7.03-6.99 (m, 2H), 6.97 (dd, *J* = 0.8 and 5.6 Hz, 1H), 4.42-4.41 (m, 4H). MS (*m/z*): 307.1 [C₁₇H₁₄N₄O₂+H]⁺.


Preparation of 4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 2




¹H NMR (400 MHz, CDCl₃) δ (ppm): 7.62 (d, *J* = 8.8 Hz, 2H), 7.40 (t, *J* = 8.0 Hz, 1H), 7.02-6.98 (m, 3H), 7.01 (d, *J* = 8.8 Hz, 2H), 6.94-6.91 (m, 1H), 6.87 (t, *J* = 2.0 Hz, 1H), 4.41-4.37 (m, 4H), 2.37 (s, 3H). MS (*m/z*): 320.2 [C₁₉H₁₇N₃O₂+H]⁺.

Preparation of 4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 3


1H NMR (400 MHz, CDCl₃) δ (ppm): 8.76 (s, 1H), 8.07 (s, 1H), 7.62 (d, J = 8.8 Hz, 2H), 7.35 (t, J = 8.0 Hz, 1H), 7.22 (d, J = 7.6 Hz, 1H), 7.16 (s, 1H), 7.04 (d, J = 8.8 Hz, 2H), 6.92 (dd, J = 2.4 and 8.2 Hz, 1H), 4.40 (s, 4H). MS (m/z): 322.9 [C₁₈H₁₄N₂O₂S+H]⁺.

Preparation of 5-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)picolinonitrile, Compound 4

1H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.50 (d, J = 2.0 Hz, 1H), 8.03 (d, J = 8.8 Hz, 1H), 7.66 (dd, J = 5.2, 2.4 Hz, 3H), 7.45 (t, J = 8.4 Hz, 1H), 7.28 (s, 1H), 7.04 (dd, J = 16.4, 7.2 Hz, 3H), 6.90 (s, 1H), 4.54-4.43 (m, 4H), 2.29 (s, 3H); MS (ESI) m/z 321.23 [C₁₈H₁₆N₄O₂+H]⁺.


Preparation of 3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 5

1H NMR (400 MHz, CDCl₃) δ (ppm): 7.67 (d, J = 1.9 Hz, 1H), 7.57 (dd, J = 2.0 and 8.5 Hz, 1H), 7.40 (t, J = 8.1 Hz, 1H), 7.06 (d, J = 8.5 Hz, 1H), 7.04-7.00 (m, 3H), 6.96-6.90 (m, 1H), 6.89 (t, J = 2.1 Hz, 1H), 4.48-4.43 (m, 4H), 2.37 (s, 3H). MS (m/z): 354.3 [C₁₉H₁₆ClN₃O₂+H]⁺.

Preparation of 3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 6


1H NMR (400 MHz, DMSO-d₆) δ (ppm): 7.85 (d, J = 11.2 Hz, 1H), 7.70 (d, J = 8.4 Hz, 1H), 7.45 (t, J = 8.8 Hz, 2H), 7.29 (s, 1H), 7.05 (dd, J = 17.6, 6.8 Hz, 3H), 6.90 (s, 1H), 4.53 (d, J = 4.0 Hz, 2H), 4.44 (d, J = 3.2 Hz, 2H), 2.30 (s, 3H). MS (ESI) m/z 338.20 [C₁₉H₁₆FN₃O₂+H]⁺.

Preparation of 3-chloro-4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 9




1H NMR (400 MHz, CDCl₃) δ (ppm): 7.68 (d, J = 1.9 Hz, 1H), 7.56 (dd, J = 1.9 and 8.5 Hz, 1H), 7.41 (t, J = 7.8 Hz, 1H), 7.07-6.98 (m, 6H), 4.48-4.44 (m, 4H), 1.82-1.75 (m, 1H), 1.13-1.09 (m, 2H), 0.92-0.86 (m, 2H); MS (m/z): 380.3 [C₂₁H₁₈ClN₃O₂+H]⁺.

Preparation of 3-chloro-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 11




1H NMR (400 MHz, CDCl₃) δ (ppm): 8.76 (s, 1H), 8.07 (s, 1H), 7.68 (d, J = 2.0 Hz, 1H), 7.57 (dd, J = 2.0 and 8.5 Hz, 1H), 7.36 (t, J = 7.9 Hz, 1H), 7.21 (dd, J = 0.8 and 7.7 Hz, 1H), 7.15 (t, J = 2.2 Hz, 1H), 7.07 (d, J = 8.5 Hz, 1H), 6.96-6.93 (m, 1H), 4.47 (s, 4H); MS (m/z): 357.3 [C₁₈H₁₃ClN₂O₂S+H]⁺.

Preparation of 3-fluoro-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 12



1H NMR (400 MHz, CDCl₃) δ (ppm): 8.76 (s, 1H), 8.07 (s, 1H), 7.47-7.43 (m, 1H), 7.40 (dd, J = 1.9 and 10.4 Hz, 1H), 7.35 (t, J = 7.9 Hz, 1H), 7.23-7.20 (m, 1H), 7.15 (t, J = 2.24 Hz, 1H), 7.11 (t, J = 8.3 Hz, 1H), 6.95-6.91 (m, 1H), 4.50-4.47 (m, 2H), 4.44-4.42 (m, 2H); MS (m/z): 339.4 [C₁₈H₁₃FN₂O₂S+H]⁺.


Preparation of 5-(3-(2-(4-fluoro-2-methylphenoxy)ethoxy)phenyl)-1-methyl-1H-pyrazole, Compound 27



1H NMR (400MHz, DMSO-d₆) δ (ppm): 7.45 (d, J = 2 Hz, 1H), 7.41 (t, J = 8.2 Hz, 1H), 7.11-7.09 (m, 2H), 7.08-7.05 (m, 1H), 7.03-6.93 (m, 3H), 6.40 (d, J = 2 Hz, 1H),

4.40-4.38 (m, 2H), 4.32-4.30 (m, 2H), 3.85 (s, 3H), 2.12 (s, 3H); MS (ESI) m/z 327.1 $[C_{19}H_{19}FN_2O_2+H]^+$


5 **Preparation of 1-(3-(2-(4-chloro-2-fluorophenoxy)ethoxy)phenyl)-2-cyclopropyl-1H-imidazole, Compound 50**



1H NMR (400 MHz, DMSO- d_6) δ 7.48-7.42 (m, 2H), 7.31-7.22 (m, 3H), 7.10-7.07 (m, 3H), 6.85 (d, $J=1.2$ Hz, 1H), 4.42 (s, 4H), 1.85-1.79 (m, 1H), 0.90-0.85 (m, 4H); MS (ESI) m/z 373.1 $[C_{20}H_{18}ClFN_2O_2+H]^+$.

10


Preparation of 4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 57



1H NMR (400 MHz, DMSO- d_6) δ 7.80-7.77 (m, 2H), 7.45 (d, $J = 2.0$ Hz, 1H), 7.42 (t, $J = 8.0$ Hz, 1H), 7.19-7.16 (m, 2H), 7.12-7.10 (m, 2H), 7.07-7.04 (m, 1H), 6.41 (d, $J = 1.6$ Hz, 1H), 4.45-4.41 (m, 4H), 3.85 (s, 3H); MS (ESI) m/z 320.1 $[C_{19}H_{17}N_3O_2+H]^+$.

15


Preparation of 1-(3-(2-(2-fluoro-4-methylphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 58



20


1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.38 (t, $J = 8.0$ Hz, 1H), 7.0 (d, $J = 8.4$ Hz, 3H), 6.94-6.84 (m, 5H), 4.37 (d, $J = 5.2$ Hz, 4H), 2.37 (s, 3H), 2.28 (s, 3H); MS (ESI) m/z 327.14 $[C_{19}H_{19}FN_2O_2+H]^+$.

25 **Preparation of 4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 62**




30


1H NMR (400 MHz, DMSO- d_6) δ 7.80-7.78 (m, 2H), 7.47 (t, $J = 7.6$ Hz, 1H), 7.26 (d, $J = 0.8$ Hz, 1H), 7.18-7.16 (m, 2H), 7.10-7.07 (m, 3H), 6.85 (s, 1H), 4.44-4.43 (m, 4H), 1.84-1.80 (m, 1H), 0.90-0.85 (m, 4H); MS (ESI) m/z 346.1 $[C_{21}H_{19}N_3O_2+H]^+$.

Preparation of 4-(2-(3-(isoxazol-4-yl)phenoxy)ethoxy)benzonitrile, Compound 67


- 1H NMR (400 MHz, CDCl₃) δ (ppm): 8.66 (s, 1H), 8.53 (s, 1H), 7.61 (d, J = 8.8 Hz, 2H), 7.35 (t, J = 8.4 Hz, 1H), 7.10 (d, J = 7.6 Hz, 1H), 7.05 (dd, J = 19.2, 2.0 Hz, 3H), 6.90 (dd, J = 8.4, 1.6 Hz, 1H), 4.39 (m, 4H); MS (ESI) m/z 307.22 [C₁₈H₁₄N₂O₃+H]⁺.

Preparation of 4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 85

- 10 1H NMR (400 MHz, CDCl₃) δ 8.41-8.40 (m, 1H), 7.64-7.61 (m, 2H), 7.25 (s, 1H), 7.03-6.99 (m, 3H), 6.89-6.86 (m, 2H), 4.46-4.43 (m, 4H), 2.60 (s, 3H); MS (ESI) m/z 321.1 [C₁₈H₁₆N₄O₂+H]⁺.


Preparation of 3-fluoro-4-(2-((6-(2-methyl-1H-imidazol-1-yl)pyridin-2-yl)oxy)ethoxy)benzonitrile, Compound 86

- 20 1H NMR (400 MHz, DMSO-d₆) δ 7.95-7.87 (m, 1H), 7.83 (dd, J = 11.2, 1.6 Hz, 1H), 7.72-7.65 (m, 1H), 7.57 (s, 1H), 7.48-7.40 (m, 1H), 7.19 (d, J = 7.6 Hz, 1H), 6.93-6.85 (m, 2H), 4.75-4.65 (m, 2H), 4.60-4.50 (m, 2H), 2.54 (s, 3H); MS (ESI) m/z 339.1 [C₁₈H₁₅N₄O₂+H]⁺.

Preparation of 5-fluoro-6-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile, Compound 87


- 25 1H NMR (CDCl₃, 400 MHz,) δ 8.22-8.21 (d, J = 2.8 Hz, 1H), 7.68-7.65 (dd, J = 6.8, 2.8 Hz, 1H), 7.24 (t, J = 6.0 Hz, 1H), 7.19-7.17 (m, 1H), 7.04-6.99 (m, 2H), 6.88-6.86 (m, 1H), 4.76 (t, J = 4.8 Hz, 1H), 4.35 (t, J = 4.6 Hz, 1H), 2.35 (s, 3H); MS (ESI) m/z 357.2 [C₁₈H₁₄F₂N₄O₂+H]⁺.

Preparation of 4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)-4-fluorophenoxy)ethoxy)-3-fluorobenzonitrile, Compound 88




1H NMR (400 MHz, DMSO-d₆) δ 7.88-7.84 (dd, J =11.2, 1.6Hz, 1H), 7.71-7.69 (m, 1H), 7.48-7.41 (m, 2H), 7.33-7.28 (m, 1H), 7.14-7.09 (m, 2H), 6.4 (s, 1H), 4.52-4.51 (m, 2H), 4.42 (m, 2H), 3.60-3.55 (m, 1H), 0.92-0.91 (m, 2H), 0.9-0.87 (m, 2H); MS (ESI) m/z 353.1 [C₁₉H₁₇FN₄O₂+H]⁺.

Preparation of 4-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 89



1H NMR (400 MHz, DMSO-d₆) δ 8.39 (d, J = 5.6 Hz, 1H), 7.81-7.78 (m, 2H), 7.49 (s, 1H), 7.25 (d, J = 2.0 Hz, 1H), 7.19-7.16 (m, 2H), 7.11-7.09 (m, 1H), 6.85 (s, 1H), 4.56-4.55 (m, 2H), 4.49-4.48 (m, 2H), 2.40-2.34 (m, 1H), 0.90-0.88 (m, 4H); MS (ESI) m/z 347.1 [C₂₀H₁₈N₄O₂+H]⁺.

Preparation of 3-fluoro-4-(2-((5-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)oxy)ethoxy)benzonitrile, Compound 92




1H NMR (400 MHz, DMSO-d₆) δ 8.39-8.36 (m, 1H), 7.88-7.85 (m, 1H), 7.73-7.70 (m, 1H), 7.63-7.62 (m, 1H), 7.51 (d, J = 2.0 Hz, 1H), 7.45 (t, J = 8.4 Hz, 1H), 6.54 (d, J = 1.6 Hz, 1H), 4.56 (s, 4H), 3.89 (s, 3H); MS (ESI) m/z 339.1 [C₁₈H₁₅FN₄O₂]⁺.

25

30


65

Preparation of 4-(2-(3-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 94




5 ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.80-7.78 (m, 2H), 7.47 (m, 1H), 7.18-7.16 (m, 2H), 7.02-6.98 (m, 3H), 6.48 (s, 1H), 4.44 (s, 4H), 3.88 (m, 3H); MS (ESI) *m/z* 338.1 [C₁₉H₁₆FN₃O₂]⁺.

Preparation of 3-fluoro-4-(2-((2-(3-methyl-4H-1,2,4-triazol-4-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 95




10 ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.40 (d, *J* = 6.0 Hz, 1H), 8.06 (s, 1H), 7.88-7.85 (m, 1H), 7.72-7.69 (m, 1H), 7.45 (t, *J* = 8.4 Hz, 1H), 7.39 (d, *J* = 2.4 Hz, 1H), 7.13-7.11 (m, 1H), 4.58 (s, 4H), 2.74 (s, 3H); MS (ESI) *m/z* 340.1 [C₁₇H₁₄FN₅O₂]⁺.

Preparation of 4-(2-(2-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 96




20 ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.80 (s, 1H), 7.78 (s, 1H), 7.46 (s, 1H), 7.40-7.30 (m, 2H), 7.18 (s, 1H), 7.16 (s, 1H), 7.14-7.08 (m, 1H), 6.42 (s, 1H), 4.55-4.50 (m, 2H), 4.50-4.45 (m, 2H), 3.85 (s, 3H); MS (ESI) *m/z* 338.1 [C₁₉H₁₆FN₃O₂+H]⁺.

Preparation of 5-fluoro-6-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile, Compound 97




25 ¹H NMR (CDCl₃, 400 MHz): δ 8.54-8.53 (d, *J* = 4.8 Hz, 1H), 8.49-8.43 (m, 2H), 7.46(d, *J* = 2.0 Hz, 1H), 7.37 (d, *J* = 2.0 Hz, 1H), 7.04-7.02(m, 1H), 6.82(d, *J* = 1.6 Hz, 1H), 4.77-4.75 (m, 2H), 4.56-4.54(m, 2H), 4.12 (s, 3H); MS (ESI) *m/z* 340.16 [C₁₇H₁₄FN₅O₂+H]⁺.

Preparation of 4-(2-(4-chloro-2-fluorophenoxy)ethoxy)-2-(1-methyl-1H-pyrazol-5-yl)pyridine, Compound 101




1H NMR (400 MHz, DMSO- d_6) δ 8.49 (d, J = 6.0 Hz, 1H), 7.47-7.43 (m, 2H), 7.38 (d, J = 2.4 Hz, 1H), 7.31-7.22 (m, 2H), 7.04-7.01 (m, 1H), 6.82 (d, J = 2.0 Hz, 1H), 4.53-4.51 (m, 2H), 4.46-4.43 (m, 1H), 4.13 (s, 3H); MS (ESI) m/z 348.0 [$C_{17}H_{15}ClFN_3O_2+H$] $^+$.

Preparation of 1-(3-(2-(4-chloro-2-fluorophenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 102




1H NMR (400 MHz, DMSO- d_6) δ 7.42 - 7.45 (m, 2H), 7.21 - 7.30 (m, 3H), 7.00 - 7.07 (m, 3H), 6.89 (s, 1H), 4.41 (s, 4H), 2.29 (s, 3H); MS (ESI) m/z 347.0 [$C_{18}H_{16}ClFN_2O_2+H$] $^+$.

Preparation of 4-(2-(4-chloro-2-fluorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine, Compound 103




1H NMR (400 MHz, DMSO- d_6) δ 8.37 (d, J = 5.6 Hz, 1H), 7.56 (d, J = 1.2 Hz, 1H), 7.45 (dd, J = 11.2, 2.4 Hz, 1H), 7.32-7.18 (m, 3H), 7.10-7.03 (m, 1H), 6.90 (d, J = 1.2 Hz, 1H), 4.60-4.50 (m, 2H), 4.50-4.40 (m, 2H), 2.49 (s, 3H); MS (ESI) m/z 348.1 [$C_{17}H_{15}ClFN_3O_2+H$] $^+$.

Preparation of 4-(2-(4-bromo-2-fluorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine, Compound 104




1H NMR (400 MHz, DMSO- d_6) δ 8.37 (d, J = 6.0 Hz, 1H), 7.60-7.50 (m, 2H), 7.40-7.32 (m, 1H), 7.28-7.20 (m, 2H), 7.10-7.03 (m, 1H), 6.90 (d, J = 1.6 Hz, 1H), 4.60-4.50 (m, 2H), 4.50-4.40 (m, 2H), 2.49 (s, 3H); MS (ESI) m/z 394.0 [$C_{17}H_{15}BrFN_3O_2+H$] $^+$.

Preparation of 5-fluoro-6-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile, Compound 107




1H NMR (CDCl₃, 400 MHz) δ 8.21 (d, J = 3.2 Hz, 1H), 7.67-7.64 (dd, J = 3.2, 7.2 Hz, 1H), 7.53 (d, J = 1.6 Hz, 1H), 7.11 (t, J = 9.2 Hz, 1H), 7.01-6.97 (m, 1H), 6.89-6.87 (dd, J = 3.2, 5.6 Hz, 1H), 6.32-6.31 (d, J = 2.0 Hz, 1H), 4.75 (t, J = 4.8 Hz, 2H), 4.34 (t, J = 5.0 Hz, 2H), 3.82-3.81 (d, J = 1.2 Hz, 1H); MS (ESI) m/z 357.1 [C₁₈H₁₄F₂N₄O₂+H]⁺.

Preparation of 6-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile, Compound 109




1H NMR (CDCl₃, 400 MHz) δ 8.48 (d, J = 2.4 Hz, 1H), 7.82-7.80 (dd, J = 8.8, 2.4 Hz, 1H), 7.18 (t, J = 9.2 Hz, 1H), 7.05 (s, 1H), 7.01-6.96 (m, 1H), 6.95 (s, 1H), 6.89-6.87 (d, J = 8.4 Hz, 1H), 6.86-6.85 (m, 1H), 4.74 (t, J = 4.6 Hz, 1H), 4.31 (t, J = 4.8 Hz, 1H), 2.30 (s, 3H); MS (ESI) m/z 339.1 [C₁₈H₁₅FN₄O₂+H]⁺

Preparation of 5-(3-(2-(4-chlorophenoxy)ethoxy)phenyl)-1-methyl-1H-pyrazole, Compound 110




1H NMR (400 MHz, DMSO-d₆) δ 7.51 (d, J = 2.0 Hz, 1H), 7.38-7.36 (m, 1H), 7.26-7.24 (m, 2H), 7.04-7.02 (m, 1H), 7.01-6.98 (m, 2H), 6.90-6.88 (m, 2H), 6.3 (d, J = 1.6 Hz, 1H), 4.35-4.32 (m, 4H), 3.89 (s, 3H); MS (ESI) m/z 329.1 [C₁₈H₁₇ClN₂O₂+H]⁺.

Preparation of 1-(3-(2-(4-chlorophenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 112




1H NMR (CDCl₃, 400 MHz) δ 7.41 (t, J = 8.1 Hz, 1H), 7.28-7.26 (m, 2H), 7.04-7.03 (m, 3H), 6.94-6.89 (m, 4H), 4.37-4.33 (m, 4H), 2.39 (s, 3H). MS (ESI) m/z 329.2 [C₁₈H₁₇ClN₂O₂+H]⁺.

Preparation of 3-fluoro-4-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 113




- 1H NMR (CDCl₃, 400 MHz) δ 7.56 (d, 1H, J = 2.0 Hz), 7.44 (dt, 1H, J = 8.8 & 0.8 Hz), 7.39 (dd, 1H, J = 10.4 & 2.0 Hz), 7.15-7.07 (m, 2H), 6.984 (dt, 1H, J = 8.8 & 3.6 Hz), 6.89 (dd, 1H, J = 6.0 & 3.0 Hz), 4.47-4.45 (m, 2H), 4.38-4.36 (m, 2H), 3.83 (d, 3H, J = 1.2 Hz); MS (ESI) m/z 356.1 [C₁₉H₁₅F₂N₃O₂+H]⁺.

Preparation of 4-(2-(4-chlorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine, Compound 115




- 1H NMR (400 MHz, DMSO-d₆) δ 8.38-8.36 (d, J = 6.0 Hz, 1H), 7.57 (d, J = 1.2 Hz, 1H), 7.35-7.32 (m, 2H), 7.21-7.20 (d, J = 1.2 Hz, 1H), 7.09-7.07 (dd, J = 6.0, 1.2 Hz, 1H), 7.03-7.00 (m, 2H), 6.92 (d, J = 1.2 Hz, 1H), 4.53-4.51 (m, 2H), 4.37-4.34 (m, 2H); MS (ESI) m/z 330.1 [C₁₇H₁₆ClN₃O₂+H]⁺.

Preparation of 5-fluoro-2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 119




- 1H NMR (400 MHz, DMSO-d₆) δ 8.38-8.37 (d, J = 5.6 Hz, 1H), 7.77-7.74 (dd, J = 8.4, 2.8 Hz, 1H), 7.62-7.57 (m, 2H), 7.39-7.35 (dd, J = 9.2, 4.0 Hz, 1H), 7.23-7.22 (d, J = 2.4 Hz, 1H), 7.10-7.08 (dd, J = 6.0, 2.4 Hz, 1H), 6.96-6.95 (d, J = 1.6 Hz, 1H), 4.58-4.56 (m, 2H), 4.54-4.53 (m, 2H); MS (ESI) m/z 339.2 [C₁₈H₁₅FN₄O₂+H]⁺.

Preparation of 4-(2-((2-(1H-tetrazol-1-yl)pyridin-4-yl)oxy)ethoxy)-3-fluorobenzonitrile, Compound 120




- 1H NMR (400 MHz, DMSO-d₆) δ 10.15 (s, 1H), 8.47 (t, J = 6.0 Hz, 1H), 7.88-7.85 (m, 1H), 7.73-7.70 (m, 1H), 7.65 (d, J = 2.4 Hz, 1H), 7.45 (t, J = 8.8 Hz, 1H), 7.28-7.26 (m, 1H), 4.67-4.65 (m, 2H), 4.60-4.58 (m, 2H); MS (ESI) m/z 327.1 [C₁₅H₁₁FN₆O₂+H]⁺.

Preparation of 4-(2-(4-bromo-2-fluorophenoxy)ethoxy)-2-(1-methyl-1H-pyrazol-5-yl)pyridine, Compound 121



1H NMR (CDCl₃, 400 MHz,) δ 8.51 (d, J = 6Hz, 1H), 7.50 (d, J = 2Hz, 1H), 7.28-7.25 (m, 2H), 7.23-7.20 (m, 1H), 7.12 (d, J = 2.4Hz, 1H), 6.91 (t, J = 8.4Hz, 1H), 6.83 (dd, J = 6 Hz, 2.4 Hz, 1H), 6.55 (d, J = 2Hz, 1H), 4.45-4.40 (m, 4H), 4.20 (s, 3H); MS (ESI) m/z 392.1 [C₁₇H₁₅BrFN₃O₂+H]⁺.

General Procedure B




To a well stirred solution of the respective phenol, **1** (1 molar equiv.) in DMSO (0.25M) were added 2-chloroethanol (2 molar equiv.) and K₂CO₃ (3 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. Upon completion of reaction, the reaction mixture was cooled and poured into water and extracted with ethyl acetate. The combined organics was washed with water and brine, dried over sodium sulphate and concentrated under reduced pressure. The crude product was purified by column chromatography to afford the desired product, **2**.

To a well stirred solution of **2** (1 molar equiv.) in DMF (0.1 M) were added the respective fluoro compound, **3** (1.3 molar equiv.) and Cs₂CO₃ (3.2 molar equiv.). The reaction mixture was heated at 100 °C for 16 hours. The reaction mixture was cooled and poured into water and extracted with ethyl acetate. The combined organics was washed with water and brine, dried over sodium sulphate and concentrated under reduced pressure to give the crude product. This crude product was purified by column chromatography to afford the desired product.


70

Preparation of 3-chloro-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 7



1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.02 (d, J = 2.0 Hz, 1H), 7.84 (dd, J = 8.4, 2.0 Hz, 1H), 7.45-7.39 (m, 3H), 7.12-7.05 (m, 3H), 6.40 (d, J = 1.2 Hz, 1H), 4.55 (dd, J = 4.8, 3.6 Hz, 2H), 4.45 (dd, J = 6.4, 2.0 Hz, 2H), 3.85 (s, 3H); MS (ESI) m/z 354.21 [$C_{19}H_{16}ClN_3O_2+H$] $^+$.


Preparation of 3-methoxy-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 8



1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.46-7.41 (m, 3H), 7.29 (d, J = 0.8 Hz, 1H), 7.20 (d, J = 8.4 Hz, 1H), 7.07-7.01 (m, 3H), 6.89 (d, J = 0.8 Hz, 1H), 4.54-4.31 (m, 4H), 3.78 (s, 3H), 2.30 (s, 3H); MS (ESI) m/z 350.28 [$C_{20}H_{19}N_3O_3+H$] $^+$.

15


Preparation of 6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile, Compound 10



1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.70 (d, J = 1.6 Hz, 1H), 8.17 (d, J = 8.8, 2.0 Hz, 1H), 7.44 (t, J = 16 Hz, 1H), 7.28 (s, 1H), 7.07-7.01 (m, 4H), 6.90 (s, 1H), 4.68 (d, J = 4.4 Hz, 2H), 4.41 (d, J = 4.4 Hz, 2H), 2.29 (s, 3H); MS (ESI) m/z 321.23 [$C_{18}H_{16}N_4O_2+H$] $^+$.

20

Preparation of 5-methyl-2-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 13




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.53-7.42 (m, 3H), 7.31 (s, 1H), 7.23 (d, J = 8.8, Hz, 1H), 7.10-7.02 (m, 3H), 6.93 (s, 1H), 4.46 (d, J = 4.8 Hz, 2H), 4.43 (d, J = 5.6 Hz, 2H), 2.32 (s, 6H); MS (ESI) m/z 334.26 [$C_{20}H_{19}N_3O_2+H$] $^+$.

30


71

Preparation of 3-fluoro-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 14




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.92-7.84 (m, 1H), 7.75-7.69 (t, $J = 16.4$ Hz, 1H), 7.57-7.39 (m, 3H), 7.12-7.05 (m, 3H), 6.40 (d, $J = 2.0$ Hz, 1H), 4.53 (dd, $J = 4.4, 3.6$ Hz, 2H), 4.43 (dd, $J = 6.0, 2.0$ Hz, 2H), 3.85 (s, 3H); MS (ESI) m/z 338.27 [$C_{19}H_{16}FN_3O_2 + H$] $^+$.

Preparation of 3-methoxy-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 15




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.46-7.39 (m, 4H), 7.20 (d, $J = 7.6$ Hz, 1H), 7.12-7.04 (m, 3H), 6.41 (d, $J = 1.2$ Hz, 1H), 4.41 (m, 4H), 3.80 (s, 6H); MS (ESI) m/z 350.28 [$C_{20}H_{19}N_3O_3 + H$] $^+$.

Preparation of 3-methoxy-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 16




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 9.08 (s, 1H), 8.35 (s, 1H), 7.45-7.43 (m, 3H), 7.31-7.20 (m, 3H), 7.01-6.99 (dd, $J = 2.0, 2.0$ Hz, 1H), 4.42 (m, 4H), 3.79 (s, 3H); MS (ESI) m/z 353.16 [$C_{19}H_{16}N_2O_3S + H$] $^+$.

Preparation of 4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-fluorobenzonitrile, Compound 17




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.88-7.85 (m, 1H), 7.70 (d, $J = 8.4$ Hz, 1H), 7.48-7.42 (m, 2H), 7.26 (d, $J = 1.2$ Hz, 1H), 7.10-7.06 (m, 3H), 6.85 (d, $J = 1.2$ Hz, 1H), 4.55-4.53 (m, 2H), 4.46-4.44 (m, 2H), 1.84-1.80 (m, 1H), 0.90-0.84 (m, 4H); MS (ESI) m/z 364.1 [$C_{21}H_{18}FN_3O_2$] $^+$.

Preparation of 5-methyl-2-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 18




1H NMR (400MHz, DMSO -d₆) δ 7.54 (s, 1H), 7.49 (d, J = 8.8 Hz, 1H), 7.46 (d, J = 1.6 Hz, 1H), 7.42 (t, J = 8 Hz, 1H), 7.24 (d, J = 8.8 Hz, 1H), 7.12-7.10 (m, 2H), 7.09-7.06 (m, 1H), 6.41 (d, J = 1.6 Hz, 1H), 4.49-4.47 (m, 2H), 4.43-4.41 (m, 2H), 3.95 (s, 3H), 2.27 (s, 3H) ; MS (ESI) m/z 334.4 [C₂₀H₁₉N₃O₂+H]⁺

Preparation of 4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-methoxybenzonitrile, Compound 19




1H NMR (400MHz, DMSO -d₆) δ 7.50-7.38 (m, 3H), 7.27 (d, J = 0.8 Hz, 1H), 7.23-7.17 (m, 1H), 7.15-7.05 (m, 3H), 6.85 (d, J = 0.8 Hz, 1H), 4.43 (s, 4H), 3.78 (s, 3H), 1.90-1.80 (m, 1H), 0.95-0.80 (m, 4H) ; MS (ESI) m/z 376.1 [C₂₂H₂₁N₃O₃+H]⁺

Preparation of 4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 20




1H NMR (400MHz, DMSO-d₆) δ 7.79 (d, J = 8.8 Hz, 2H), 7.46 (t, J = 8.0 Hz, 1H), 7.27 (d, J = 1.6 Hz, 1H), 7.20-7.13 (m, 2H), 7.13-7.05 (m, 3H), 6.85 (d, J = 1.2 Hz, 1H), 4.50-4.40 (m, 4H), 1.85-1.75 (m, 1H), 0.95-0.80 (m, 4H) ; MS (ESI) m/z 346.1 [C₂₁H₁₉N₃O₂+H]⁺

Preparation of -(4H-1,2,4-triazol-4-yl)phenoxy)ethoxy)benzonitrile, Compound 21




1H NMR (400 MHz, CDCl₃) δ (ppm): 8.54 (s, 1H), 8.09 (s, 1H), 7.60 (d, J = 8.8 Hz, 2H), 7.42 (t, J = 8.0 Hz, 1H), 7.33-7.25 (m, 2H), 7.02-6.96 (m, 3H), 4.43-4.39 (m, 4H); MS (ESI) m/z 307.29 [C₁₇H₁₄N₄O₂ +H]⁺.

Preparation of 3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 22




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.99 (d, *J* = 1.2 Hz, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.42 (dd, *J* = 16.8, 16.4 Hz, 2H), 7.25 (s, 1H), 7.05 (d, *J* = 6.4 Hz, 2H), 7.00 (d, *J* = 8.0 Hz, 1H), 6.88 (s, 1H), 5.01-4.98 (m, 1H), 4.38 (dd, *J* = 16.8, 16 Hz, 2H), 2.27 (s, 3H), 1.39 (d, *J* = 5.6 Hz, 3H); MS (ESI) *m/z* 368.33 [C₂₀H₁₈ClN₃O₂+H]⁺.

Preparation of 3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 23




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.81 (d, *J* = 11.2 Hz, 1H), 7.66 (d, *J* = 8.8 Hz, 1H), 7.41 (d, *J* = 8.4 Hz, 2H), 7.25 (s, 1H), 7.05 (d, *J* = 8.4 Hz, 2H), 6.98 (d, *J* = 8.4 Hz, 1H), 6.88 (s, 1H), 5.01-4.64 (m, 1H), 4.38 (t, *J* = 2.4 Hz, 2H), 2.27 (s, 3H), 1.38 (d, *J* = 6.4 Hz, 3H); MS (ESI) *m/z* 352.12 [C₂₀H₁₈FN₃O₂+H]⁺.

Preparation of 3-chloro-4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 24




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.41 (d, *J* = 5.6 Hz, 1H), 8.03 (d, *J* = 2.0 Hz, 1H), 7.86-7.84 (dd, *J* = 2.0 Hz, 1H), 7.75 (s, 1H), 7.41 (d, *J* = 8.8 Hz, 1H), 7.32 (d, *J* = 2.0 Hz, 1H), 7.21-7.26 (m, 2H), 4.62-4.58 (dd, *J* = 4.8, 5.2 Hz, 4H), 2.58 (s, 3H); MS (ESI) *m/z* 355.27 [C₁₈H₁₅ClN₄O₂+H]⁺.

Preparation of 3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 25




1H NMR (400 MHz, CDCl₃) δ (ppm): 7.42-7.35 (m, 3H), 7.14 (t, J = 8.4 Hz, 1H), 7.02-6.89 (m, 4H), 6.81 (t, J = 2.0 Hz, 1H), 2.36 (s, 3H), 1.50 (d, J = 6.4 Hz, 3H); MS (ESI) m/z 352.14 [C₂₀H₁₈FN₃O₂+H]⁺.

Preparation of 1-(3-(2-(4-ethynylphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 26




1H NMR (400 MHz, DMSO-d₆) δ (ppm): 7.47-7.39 (m, 3H), 7.28 (d, J = 1.6 Hz, 1H), 7.10-6.96 (m, 5H), 6.89 (d, J = 1.2 Hz, 1H), 4.45-4.30 (m, 4H), 4.01 (s, 1H), 2.30 (s, 3H); MS (ESI) m/z 319.1 [C₂₀H₁₈N₂O₂+H]⁺.

Preparation of 4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-fluorobenzonitrile, Compound 28



1H NMR (400 MHz, DMSO-d₆) δ (ppm): 7.80 (dd, J = 11.2 Hz, 2.0 Hz, 1H), 7.64 (d, J = 8.4 Hz, 1H), 7.49-7.39 (m, 2H), 7.22 (s, 1H), 7.06-6.98 (m, 3H), 6.82 (s, 1H), 5.90-5.05 (m, 1H), 4.29-4.21 (m, 2H), 1.80-1.73 (m, 1H), 1.36 (t, J = 6.4 Hz, 3H), 0.91-0.79 (m, 4H); MS (ESI) m/z 378.21 [C₂₂H₂₀FN₃O₂+H]⁺.


Preparation of 3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 29



1H NMR (400 MHz, CDCl₃) δ (ppm): 7.36 (t, J = 8.0 Hz, 1H), 7.26-7.24 (m, 1H), 7.10 (d, J = 8.8 Hz, 1H), 7.06-6.88 (m, 5H), 6.83 (s, 1H), 4.86-4.82 (m, 1H), 4.26-4.22 (m, 1H), 4.12-4.09 (m, 1H), 3.84 (s, 3H), 2.36 (s, 3H), 1.49 (d, J = 6.4 Hz, 3H); MS (ESI) m/z 364.16 [C₂₁H₂₁N₃O₃+H]⁺.


30

Preparation of 3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 30




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.50-7.42 (m, 3H), 7.38-7.34 (m, 1H), 7.03-6.92 (m, 4H), 6.28 (d, *J* = 2.0 Hz, 1H), 4.87-4.82 (dd, *J* = 6.4, 5.6 Hz, 1H), 4.23-4.19 (dd, *J* = 6.4, 5.6 Hz, 1H), 4.11-3.88 (d, *J* = 4.4 Hz, 1H), 3.88 (s, 3H), 2.19 (s, 3H), 1.48 (d, *J* = 6.4 Hz, 3H); MS (ESI) *m/z* 348.29 [C₂₁H₂₁N₃O₂+H]⁺.

Preparation of 3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 31




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.50 (d, *J* = 1.6 Hz, 1H), 7.37-7.31 (m, 1H), 7.26 (s, 1H), 7.09 (d, *J* = 1.2 Hz, 1H), 7.04-6.99 (m, 2H), 6.93 (d, *J* = 7.6 Hz, 2H), 6.28 (d, *J* = 1.2 Hz, 1H), 4.84 (dd, *J* = 6.0, 2.0 Hz, 1H), 4.26 (dd, *J* = 9.6, 6.4 Hz, 1H), 4.13-4.09 (m, 1H), 3.88 (s, 3H), 3.84 (s, 3H), 1.50 (d, *J* = 6.4 Hz, 3H); MS (ESI) *m/z* 364.26 [C₂₁H₂₁N₃O₃+H]⁺.

Preparation of (R)-3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 32




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.99 (d, *J* = 1.2 Hz, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.42 (dd, *J* = 16.8, 16.4 Hz, 2H), 7.25 (s, 1H), 7.05 (d, *J* = 6.4 Hz, 2H), 7.00 (d, *J* = 8.0 Hz, 1H), 6.88 (s, 1H), 5.01-4.98 (m, 1H), 4.38 (dd, *J* = 16.8, 16 Hz, 2H), 2.27 (s, 3H), 1.39 (d, *J* = 5.6 Hz, 3H); MS (ESI) *m/z* 368.33 [C₂₀H₁₈ClN₃O₂+H]⁺.

Preparation of (S)-3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 33




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.99 (d, *J* = 1.2 Hz, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.42 (dd, *J* = 16.8, 16.4 Hz, 2H), 7.25 (s, 1H), 7.05 (d, *J* = 6.4 Hz, 2H), 7.00 (d, *J* = 8.0 Hz, 1H), 6.88 (s, 1H), 5.01-4.98 (m, 1H), 4.38 (dd, *J* = 16.8, 16 Hz, 2H), 2.27 (s, 3H), 1.39 (d, *J* = 5.6 Hz, 3H); MS (ESI) *m/z* 368.33 [C₂₀H₁₈ClN₃O₂+H]⁺.

Preparation of (R)-3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 34




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.84 (dd, *J* = 11.6, 1.6 Hz, 1H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.44-7.39 (m, 2H), 7.26 (d, *J* = 1.2 Hz, 1H), 7.05-6.98 (m, 3H), 6.89 (d, *J* = 1.2 Hz, 1H), 5.01-4.97 (m, 1H), 4.38-4.31 (m, 2H), 2.28 (s, 3H), 1.37 (d, *J* = 6.0 Hz, 3H); MS (ESI) *m/z* 352.25 [C₂₀H₁₈FN₃O₂+H]⁺.

Preparation of (S)-3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile, Compound 35




¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.84 (dd, *J* = 11.6, 1.6 Hz, 1H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.44-7.39 (m, 2H), 7.26 (d, *J* = 1.2 Hz, 1H), 7.05-6.98 (m, 3H), 6.89 (d, *J* = 1.2 Hz, 1H), 5.01-4.97 (m, 1H), 4.38-4.31 (m, 2H), 2.28 (s, 3H), 1.37 (d, *J* = 6.0 Hz, 3H); MS (ESI) *m/z* 352.25 [C₂₀H₁₈FN₃O₂+H]⁺.

Preparation of 4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-(trifluoromethoxy)benzonitrile, Compound 36




5 ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.78-7.69 (m, 2H), 7.49-7.39 (m, 2H), 7.18 (d, *J* = 1.2 Hz, 1H), 7.13-7.08 (m, 1H), 7.02-6.95 (m, 3H), 4.60-4.50 (m, 2H), 4.50-4.40 (m, 2H), 2.33 (m, 3H); MS (ESI) *m/z* 404.1 [C₂₀H₁₆F₃N₃O₃+H]⁺.

Preparation of 4-(2-(3-(4H-1,2,4-triazol-4-yl)phenoxy)ethoxy)-3-fluorobenzonitrile, Compound 38




10 ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.54 (s, 1H), 8.09 (s, 1H), 7.45-7.26 (m, 5H), 7.10 (t, *J* = 8.4 Hz, 1H), 6.97 (dd, *J* = 2.0, 8.0 Hz, 1H), 4.49-4.45 (m, 4H); MS (ESI) *m/z* 325.10 [C₁₇H₁₃FN₄O₂+H]⁺.

Preparation of 3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 37




20 ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.50 (d, *J* = 2.0 Hz, 1H), 7.42-7.34 (m, 3H), 7.17 (t, 3H), 7.03-6.99 (m, 1H), 6.92-6.91 (m, 2H), 6.28 (d, *J* = 1.6 Hz, 1H), 4.91-4.87 (dd, *J* = 10.8, 6.4 Hz, 1H), 4.25-4.21 (m, 1H), 4.14-4.10 (m, 1H), 3.88 (d, *J* = 4.4 Hz, 1H), 1.55-1.47 (m, 3H); MS (ESI) *m/z* 352.25 [C₂₀H₁₈FN₃O₂+H]⁺.

Preparation of (R)-3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 39




25 ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.50-7.42 (m, 3H), 7.38-7.34 (m, 1H), 7.03-6.92 (m, 4H), 6.28 (d, *J* = 2.0 Hz, 1H), 4.87-4.82 (dd, *J* = 6.4, 5.6 Hz, 1H), 4.23-4.19 (dd, *J* = 6.4, 5.6 Hz, 1H), 4.11-3.88 (d, *J* = 4.4 Hz, 1H), 3.88 (s, 3H), 2.19 (s, 3H), 1.48 (d, *J* = 6.4 Hz, 3H); MS (ESI) *m/z* 348.29 [C₂₁H₂₁N₃O₂+H]⁺.

Preparation of (S)-3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 40




- 1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.50-7.42 (m, 3H), 7.38-7.34 (m, 1H), 7.03-6.92 (m, 4H), 6.28 (d, $J = 2.0$ Hz, 1H), 4.87-4.82 (dd, $J = 6.4, 5.6$ Hz, 1H), 4.23-4.19 (dd, $J = 6.4, 5.6$ Hz, 1H), 4.11-3.88 (d, $J = 4.4$ Hz, 1H), 3.88 (s, 3H), 2.19 (s, 3H), 1.48 (d, $J = 6.4$ Hz, 3H); MS (ESI) m/z 348.29 [$C_{21}H_{21}N_3O_2 + H$] $^+$.

Preparation of (R)-4-(1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy-3-fluorobenzonitrile, Compound 41




- 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.41-7.36 (m, 1H), 7.02 (t, $J = 8.0$ Hz, 4H), 4.25-4.20 (m, 1H), 3.97 (dd, $J = 4.0, 12.0$ Hz, 1H), 3.85 (t, $J = 8.0$ Hz, 1H), 2.34 (s, 1H), 1.82-1.75 (m, 1H), 1.30 (d, $J = 4.0$ Hz, 3H), 1.12-1.09 (m, 2H), 0.92-0.87 (m, 2H); MS (ESI) m/z 259.21 [$C_{15}H_{18}N_2O_2 + H$] $^+$.

Preparation of (S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-fluorobenzonitrile, Compound 42



- 20 1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.84 (dd, $J = 2.0$ Hz, 11.2 Hz, 1H), 7.66 (d, $J = 8.0$ Hz, 1H), 7.52-7.42 (m, 2H), 7.26 (d, $J = 0.8$ Hz, 1H), 7.09-7.01 (m, 3H), 6.86 (s, 1H), 5.11-5.07 (m, 1H), 4.31-4.23 (m, 2H), 1.81-1.77 (m, 1H), 1.39 (d, $J = 6.4$ Hz, 3H), 0.91-0.82 (m, 4H); MS (ESI) m/z 378.26 [$C_{22}H_{20}FN_3O_2 + H$] $^+$.

Preparation of (S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile, Compound 43




- 1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.43-7.35 (m, 3H), 7.23 (dd, $J = 2.0$ Hz, 3.2 Hz, 2H), 7.06-6.99 (m, 3H), 6.82 (s, 1H), 4.95 (dd, $J = 13.2$ Hz, 11.2 Hz, 1H), 4.22 (d, J

= 4.4 Hz, 2H), 3.73 (s, 3H), 1.79-1.75 (m, 1H), 1.34 (dd, $J = 8.0, 11.2$ Hz, 3H), 0.88-0.79 (m, 4H); MS (ESI) m/z 390.27 [$C_{23}H_{23}N_3O_3+H$] $^+$.

Preparation of 3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 44


5



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.66 (d, $J = 2.0$ Hz, 1H), 7.53 (dd, $J = 2.0$ Hz, 8.8 Hz, 1H), 7.37 (t, $J = 8.4$ Hz, 1H), 7.15 (d, $J = 8.8$ Hz, 1H), 7.01-6.90 (m, 4H), 6.82 (s, 1H), 4.91-4.87 (m, 1H), 4.27-4.23 (m, 1H), 4.15-4.12 (m, 1H), 2.36 (s, 3H), 1.54 (d, $J = 13.2$ Hz, 3H); MS (ESI) m/z 368.11 [$C_{20}H_{18}ClN_3O_2+H$] $^+$.

10

Preparation of 1-(3-(2-(4-chloro-2-methoxyphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 45




1H NMR (400 MHz, $DMSO-d_6$) δ (ppm): 7.92-7.84 (m, 1H), 7.75-7.69 (t, $J = 16.4$ Hz, 1H), 7.57-7.39 (m, 3H), 7.12-7.05 (m, 3H), 6.40 (d, $J = 2.0$ Hz, 1H), 4.53 (dd, $J = 4.4, 3.6$ Hz, 2H), 4.43 (dd, $J = 6.0, 2.0$ Hz, 2H), 3.85 (s, 3H); MS (ESI) m/z 338.27 [$C_{19}H_{16}FN_3O_2+H$] $^+$.

15

Preparation of (R)-3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 46


20



1H NMR (400 MHz, $DMSO-d_6$) δ (ppm): 7.51 (d, $J = 2.0$ Hz, 1H), 7.43-7.34 (m, 3H), 7.17-7.13 (m, 3H), 7.03-7.01 (m, 1H), 6.92-6.91 (m, 2H), 6.29 (d, $J = 1.6$ Hz, 1H), 4.91-4.87 (dd, $J = 10.8, 6.4$ Hz, 1H), 4.26-4.21 (m, 1H), 4.14-4.11 (m, 1H), 3.89 (d, $J = 4.4$ Hz, 1H), 1.52-1.47 (m, 3H); MS (ESI) m/z 352.25 [$C_{20}H_{18}FN_3O_2+H$] $^+$.


25

Preparation of (S)-3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 47




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.51 (d, J = 2.0 Hz, 1H), 7.43-7.34 (m, 3H), 7.17-7.13 (m, 3H), 7.03-7.01 (m, 1H), 6.92-6.91 (m, 2H), 6.29 (d, J = 1.6 Hz, 1H), 4.91-4.87 (dd, J = 10.8, 6.4 Hz, 1H), 4.26-4.21 (m, 1H), 4.14-4.11 (m, 1H), 3.89 (d, J = 4.4 Hz, 1H), 1.52-1.47 (m, 3H); MS (ESI) m/z 352.25 [$C_{20}H_{18}FN_3O_2+H$] $^+$.

Preparation of (R)-3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 48




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.51 (s, 1H), 7.36 (t, 1H), 7.26 (s, 1H), 7.10 (s, 1H), 7.04-7.01 (m, 2H), 6.96-6.92 (m, 2H), 6.29 (s, 2H), 4.90-4.83 (m, 1H), 4.26 (dd, J = 10.0, 6.4 Hz, 1H), 4.11 (dd, J = 10.0, 4.4 Hz, 1H), 3.89 (s, 1H), 3.84 (s, 3H), 1.51 (d, J = 6.4 Hz, 3H); MS (ESI) m/z 364.26 [$C_{21}H_{21}N_3O_3+H$] $^+$.

Preparation of (S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile, Compound 49




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.62 (t, J = 8.4 Hz, 2H), 7.43 (t, J = 8.0 Hz, 1H), 7.24 (d, J = 7.6 Hz, 2H), 7.05 (t, J = 10.8 Hz, 3H), 6.84 (s, 1H), 4.99 (d, J = 4.4 Hz, 1H), 4.27 (d, J = 13.2 Hz, 2H), 2.09 (s, 3H), 1.76 (dd, J = 4.8, 10 Hz, 1H), 1.37 (d, J = 6.4 Hz, 3H), 0.88-0.80 (m, 4H); MS (ESI) m/z 374.30 [$C_{23}H_{23}N_3O_2+H$] $^+$.

Preparation of 1-(3-(2-(4-ethynyl-2-fluorophenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole, Compound 51




1H NMR (400MHz, DMSO - d_6) δ 7.44 (t, J = 8.2 Hz, 1H), 7.37 (dd, J =1.6, 2 Hz, 1H), 7.31-7.22 (m, 3H), 7.08-7.06 (m, 2H), 7.02 (d, J = 8.4 Hz, 1H), 6.89 (d, J = 0.8 Hz, 1H), 4.45-4.41 (m, 4H), 4.13 (s, 1H), 2.30 (s, 3H) ; MS (ESI) m/z 337.1 [$C_{20}H_{17}FN_2O_2+H$] $^+$

Preparation of 3-amino-4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 52




1H NMR (400 MHz, DMSO- d_6) δ 7.47 (t, J = 8.0 Hz, 1H), 7.17-7.05 (m, 4H), 7.03-6.95 (m, 3H), 6.88-6.92 (m, 1H), 4.50-4.40 (m, 4H), 1.90-1.75 (m, 1H), 1.00-0.85 (m, 4H); MS (ESI) m/z 361.1 [$C_{21}H_{20}N_4O_2+H$] $^+$.

Preparation of 3-fluoro-4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 53




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.37 (d, J = 6.0 Hz, 1H), 7.84-7.85 (dd, J = 1.6, 11.2 Hz, 1H), 7.71 (d, J = 8.0 Hz, 1H), 7.56 (s, 1H), 7.44 (t, 1H), 7.21 (d, J = 2.0 Hz, 1H), 7.09-7.07 (dd, J = 2.0, 6.0 Hz, 1H), 6.91 (s, 1H), 4.58-4.57 (m, 4H), 3.31 (s, 3H); MS (ESI) m/z 339.23 [$C_{18}H_{15}FN_4O_2+H$] $^+$.

Preparation of 5-fluoro-2-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 54




1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.50 (d, J = 1.6 Hz, 1H), 7.38 (t, J = 7.6 Hz, 1H), 7.29 (s, 1H), 7.26 (d, J = 7.2 Hz, 1H), 7.05-6.97 (m, 4H), 6.30 (d, J = 2.0 Hz, 1H), 4.43 (q, J = 2.8 Hz, 4H), 3.89 (s, 3H); MS (ESI) m/z 338.26 [$C_{19}H_{16}FN_3O_2+H$] $^+$.

Preparation of (R)-3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 55




1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.66 (d, J = 2.0 Hz, 1H), 7.53 (dd, J = 2.0 Hz, 8.8 Hz, 1H), 7.37 (t, J = 8.4 Hz, 1H), 7.15 (d, J = 8.8 Hz, 1H), 7.01-6.90 (m, 4H), 6.82 (s, 1H), 4.91-4.87 (m, 1H), 4.27-4.23 (m, 1H), 4.15-4.12 (m, 1H), 2.36 (s, 3H), 1.54 (d, J = 13.2 Hz, 3H); MS (ESI) m/z 368.11 [$C_{20}H_{18}ClN_3O_2+H$] $^+$.

Preparation of (S)-3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 56




1H NMR (400 MHz, CDCl₃) δ (ppm): 7.66 (d, J = 2.0 Hz, 1H), 7.53 (dd, J = 2.0 Hz, 8.8 Hz, 1H), 7.37 (t, J = 8.4 Hz, 1H), 7.15 (d, J = 8.8 Hz, 1H), 7.01-6.90 (m, 4H), 6.82 (s, 1H), 4.91-4.87 (m, 1H), 4.27-4.23 (m, 1H), 4.15-4.12 (m, 1H), 2.36 (s, 3H), 1.54 (d, J = 13.2 Hz, 3H); MS (ESI) m/z 368.11 [C₂₀H₁₈ClN₃O₂+H]⁺.

Preparation of 4-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 59




1H NMR (400 MHz, CDCl₃) δ (ppm): 8.52 (d, J = 9.6 Hz, 1H), 7.61 (d, J = 8.8 Hz, 2H), 7.49 (s, 1H), 7.11 (s, 1H), 7.03 (d, J = 8.8 Hz, 2H), 6.82 (d, J = 4.0 Hz, 1H), 6.53 (s, 1H), 4.42 (d, J = 6.4 Hz, 4H), 4.20 (s, 3H); MS (ESI) m/z 321.30 [C₁₈H₁₆N₄O₂+H]⁺.

Preparation of (S)-3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 60



1H NMR (400 MHz, DMSO-d₆) δ (ppm): 7.50 (d, J = 1.6 Hz, 1H), 7.37-7.31 (m, 1H), 7.26 (s, 1H), 7.09 (d, J = 1.2 Hz, 1H), 7.04-6.99 (m, 2H), 6.93 (d, J = 7.6 Hz, 2H), 6.28 (d, J = 1.2 Hz, 1H), 4.84 (dd, J = 6.0, 2.0 Hz, 1H), 4.26 (dd, J = 9.6, 6.4 Hz, 1H), 4.13-4.09 (m, 1H), 3.88 (s, 3H), 3.84 (s, 3H), 1.50 (d, J = 6.4 Hz, 3H); MS (ESI) m/z 364.26 [C₂₁H₂₁N₃O₃+H]⁺.


Preparation of 4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile, Compound 61



1H NMR (400 MHz, DMSO-d₆) δ (ppm): 7.62 (t, J = 8.4 Hz, 2H), 7.43 (t, J = 8.0 Hz, 1H), 7.24 (d, J = 7.6 Hz, 2H), 7.05 (t, J = 10.8 Hz, 3H), 6.84 (s, 1H), 4.99 (d, J = 4.4

Hz, 1H), 4.27 (d, $J = 13.2$ Hz, 2H), 2.09 (s, 3H), 1.76 (dd, $J = 4.8, 10$ Hz, 1H), 1.37 (d, $J = 6.4$ Hz, 3H), 0.88-0.80 (m, 4H); MS (ESI) m/z 374.30 [$C_{23}H_{23}N_3O_2+H$] $^+$.


Preparation of 3-fluoro-4-(3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-yloxy)benzonitrile, Compound 63



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.40-7.33 (m, 3H), 7.11-7.03 (m, 3H), 6.98-6.87 (m, 2H), 6.81 (t, $J = 2.4$ Hz, 1H), 4.65-4.56 (m, 2H), 2.37 (d, $J = 5.2$ Hz, 3H), 1.57-1.40 (m, 6H); MS (ESI) m/z 366.15 [$C_{21}H_{20}FN_3O_2+H$] $^+$.

10


Preparation of 3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 64



1H NMR (400 MHz, $DMSO-d_6$) δ (ppm): 8.62 (d, $J = 4.0$ Hz, 1H), 7.82 (dd, $J = 2.0, 11.6$ Hz, 1H), 7.65 (d, $J = 8.8$ Hz, 1H), 7.51-7.39 (m, 4H), 7.04 (dd, $J = 1.6, 5.2$ Hz, 2H), 5.10-5.06 (m, 1H), 4.30-4.23 (m, 2H), 1.35 (d, $J = 5.6$ Hz, 3H); MS (ESI) m/z 339.19 [$C_{19}H_{15}FN_3O_2+H$] $^+$.

15


Preparation of (R)-3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 65



1H NMR (400 MHz, $DMSO-d_6$) δ (ppm): 8.62 (d, $J = 4.0$ Hz, 1H), 7.82 (dd, $J = 2.0, 11.6$ Hz, 1H), 7.65 (d, $J = 8.8$ Hz, 1H), 7.51-7.39 (m, 4H), 7.04 (dd, $J = 1.6, 5.2$ Hz, 2H), 5.10-5.06 (m, 1H), 4.30-4.23 (m, 2H), 1.35 (d, $J = 5.6$ Hz, 3H); MS (ESI) m/z 339.19 [$C_{19}H_{15}FN_3O_2+H$] $^+$.


25

Preparation of (S)-3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 66




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.62 (d, J = 4.0 Hz, 1H), 7.82 (dd, J = 2.0, 11.6 Hz, 1H), 7.65 (d, J = 8.8 Hz, 1H), 7.51-7.39 (m, 4H), 7.04 (dd, J = 1.6, 5.2 Hz, 2H), 5.10-5.06 (m, 1H), 4.30-4.23 (m, 2H), 1.35 (d, J = 5.6 Hz, 3H); MS (ESI) m/z 339.19 [$C_{19}H_{15}FN_3O_2+H$] $^+$.

Preparation of (R)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile, Compound 68




1H NMR (400 MHz, DMSO- d_6) δ (ppm): 7.62 (t, J = 8.4 Hz, 2H), 7.43 (t, J = 8.0 Hz, 1H), 7.24 (d, J = 7.6 Hz, 2H), 7.05 (t, J = 10.8 Hz, 3H), 6.84 (s, 1H), 4.99 (d, J = 4.4 Hz, 1H), 4.27 (d, J = 13.2 Hz, 2H), 2.09 (s, 3H), 1.76 (dd, J = 4.8, 10 Hz, 1H), 1.37 (d, J = 6.4 Hz, 3H), 0.88-0.80 (m, 4H); MS (ESI) m/z 374.30 [$C_{23}H_{23}N_3O_2+H$] $^+$.

Preparation of 3-fluoro-4-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 69



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.52 (d, J = 6.0 Hz, 1H), 7.5 (d, J = 1.6 Hz, 1H), 7.41 (dd, J = 8.8 Hz, 19.6 Hz, 2H), 7.12-7.07 (m, 2H), 6.83-6.81 (m, 1H), 6.54 (d, J = 2 Hz, 1H), 4.48 (s, 4H), 4.20 (s, 3H); MS (ESI) m/z 339.19 [$C_{18}H_{15}FN_4O_2+H$] $^+$.


Preparation of 6-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile, Compound 70



1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.72 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 5.6 Hz, 1H), 8.19-8.17 (dd, J = 2.4, 8.8 Hz, 1H), 7.57 (s, 1H), 7.21 (d, J = 2.0 Hz, 1H), 7.08-


7.06 (m, 2H), 6.93 (s, 1H), 4.73-4.71 (m, 2H), 4.56-4.54 (m, 2H), 2.50 (s, 3H); MS (ESI) m/z 322.29 [$C_{17}H_{15}N_5O_2+H$]⁺

Preparation of 6-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile, Compound 71




1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.60 (d, J = 5.6 Hz, 2H), 7.8 (dd, J = 1.6, 8 Hz, 1H), 7.49 (d, J = 1.6 Hz, 1H), 7.12 (d, J = 2.0 Hz, 1H), 6.88 (d, J = 4.8 Hz, 1H), 6.81 (t, J = 6.0 Hz, 1H), 6.53 (d, J = 2.0 Hz, 1H), 4.78 (t, J = 4.4 Hz, 2H), 4.42 (t, J = 4.8 Hz, 2H), 4.20 (s, 3H); MS (ESI) m/z 322.12 [$C_{17}H_{15}N_5O_2+H$]⁺.

Preparation of (S)-3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 72




1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.36 (t, J = 8.0 Hz, 1H), 7.26-7.24 (m, 1H), 7.10 (d, J = 8.8 Hz, 1H), 7.06-6.88 (m, 5H), 6.83 (s, 1H), 4.86-4.82 (m, 1H), 4.26-4.22 (m, 1H), 4.12-4.09 (m, 1H), 3.84 (s, 3H), 2.36 (s, 3H), 1.49 (d, J = 6.4 Hz, 3H); MS (ESI) m/z 364.16 [$C_{21}H_{21}N_3O_3+H$]⁺.

Preparation of 6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)pyridazine-3-carbonitrile, Compound 73




1H NMR (400 MHz, $DMSO-d_6$) δ 8.25 (d, J = 9.2 Hz, 1H), 7.56 (d, J = 9.2 Hz, 1H), 7.44 (t, J = 8.8 Hz, 1H), 7.28 (d, J = 1.2 Hz, 1H), 7.08-7.07 (m, 2H), 7.02 (d, J = 8.4 Hz, 1H), 6.89 (d, J = 1.2 Hz, 1H), 4.89 (d, J = 4.0 Hz, 2H), 4.48 (d, J = 4.4 Hz, 2H), 2.30 (s, 3H); MS (ESI) m/z 321.9 [$C_{17}H_{15}N_5O_2+H$]⁺.

Preparation of 5-fluoro-6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile, Compound 74




1H NMR (400 MHz, DMSO- d_6) δ 8.53 (d, J = 3.2 Hz, 1H), 8.45-8.43 (m, 1H), 7.44 (t, J = 8.0 Hz, 1H), 7.28 (d, J = 1.2 Hz, 1H), 7.08-7.01 (m, 3H), 6.89 (d, J = 1.2 Hz, 1H), 4.74 (t, J = 4.0 Hz, 2H), 4.44 (t, J = 5.2 Hz, 2H), 2.29 (s, 3H); MS (ESI) m/z 339.9 [$C_{18}H_{15}FN_4O_2+H$] $^+$.

Preparation of 6-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile, Compound 75




1H NMR (400 MHz, DMSO- d_6) δ 8.71 (d, J = 2.0 Hz, 1H), 8.19-8.16 (m, 1H), 7.46-7.39 (m, 2H), 7.11-7.04 (m, 4H), 6.40 (d, J = 2.0 Hz, 1H), 4.70 (d, J = 4.4 Hz, 2H), 4.41 (d, J = 4.4 Hz, 2H), 3.85 (s, 3H); MS (ESI) m/z 320.9 [$C_{18}H_{16}N_4O_2+H$] $^+$.

Preparation of 4-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)-3-fluorobenzonitrile, Compound 76



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.42 (d, J = 5.6 Hz, 1H), 7.46-7.38 (m, 2H), 7.29 (d, J = 1.2 Hz, 1H), 7.08 (t, J = 8.4 Hz, 2H), 6.98-6.97 (m, 1H), 6.88 (dd, J = 2.4, 6.0 Hz, 1H), 4.50 (s, 4H), 2.19-2.14 (m, 1H), 1.17-1.13 (m, 2H), 0.98-0.94 (m, 2H); MS (ESI) m/z 365.23 [$C_{20}H_{17}FN_4O_2+H$] $^+$.


Preparation of (R)-3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 77



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.36 (t, J = 8.0 Hz, 1H), 7.26-7.24 (m, 1H), 7.10 (d, J = 8.8 Hz, 1H), 7.06-6.88 (m, 5H), 6.83 (s, 1H), 4.86-4.82 (m, 1H), 4.26-4.22

(m, 1H), 4.12-4.09 (m, 1H), 3.84 (s, 3H), 2.36 (s, 3H) 1.49 (d, $J = 6.4$ Hz, 3H); MS (ESI) m/z 364.16 [$C_{21}H_{21}N_3O_3+H$] $^+$.


Preparation of (R)-3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 78



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.42-7.35 (m, 3H), 7.14 (t, $J = 8.4$ Hz, 1H), 7.02-6.89 (m, 4H), 6.81 (t, $J = 2.0$ Hz, 1H), 2.36 (s, 3H), 1.50 (d, $J = 6.4$ Hz, 3H); MS (ESI) m/z 352.14 [$C_{20}H_{18}FN_3O_2+H$] $^+$.

10


Preparation of (S)-3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile, Compound 79



1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.42-7.35 (m, 3H), 7.14 (t, $J = 8.4$ Hz, 1H), 7.02-6.89 (m, 4H), 6.81 (t, $J = 2.0$ Hz, 1H), 2.36 (s, 3H), 1.50 (d, $J = 6.4$ Hz, 3H); MS (ESI) m/z 352.14 [$C_{20}H_{18}FN_3O_2+H$] $^+$.

15

Preparation of 4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile, Compound 80




20

1H NMR (400 MHz, $CDCl_3$) δ (ppm): 7.41-7.36 (m, 1H), 7.04-6.96 (m, 4H), 4.25-4.20 (m, 1H), 3.95 (dd, $J = 3.2, 9.2$ Hz, 1H), 3.85 (t, $J = 8.0$ Hz, 1H), 2.37 (s, 1H), 1.82-1.75 (m, 1H), 1.30 (d, $J = 6.4$ Hz, 3H), 1.12-1.08 (m, 2H), 0.92-0.87 (m, 2H); MS (ESI) m/z 259.21 [$C_{15}H_{18}N_2O_2+H$] $^+$.

25


30

Preparation of (S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile, Compound 81




1H NMR (400 MHz, CDCl₃) δ (ppm): 7.41-7.36 (m, 1H), 7.04-6.96 (m, 4H), 4.25-4.20 (m, 1H), 3.95 (dd, J = 3.2, 9.2 Hz, 1H), 3.85 (t, J = 8.0 Hz, 1H), 2.37 (s, 1H), 1.82-1.75 (m, 1H), 1.30 (d, J = 6.4 Hz, 3H), 1.12-1.08 (m, 2H), 0.92-0.87 (m, 2H); MS (ESI) m/z 259.21 [C₁₅H₁₈N₂O₂+H]⁺.

Preparation of 5-fluoro-6-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile, Compound 82




1H NMR (400 MHz, DMSO-d₆) δ 8.53 (d, J = 3.2 Hz, 1H), 8.45-8.42 (m, 1H), 7.46-7.39 (m, 2H), 7.12-7.05 (m, 3H), 6.41 (d, J = 1.6 Hz, 1H), 4.74 (d, J = 4.0 Hz, 2H), 4.43 (d, J = 4.8 Hz, 2H), 3.85 (s, 3H); MS (ESI) m/z 339.1 [C₁₈H₁₅FN₄O₂+H]⁺.

Preparation of 6-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-5-fluoronicotinonitrile, Compound 83




1H NMR (400 MHz, DMSO-d₆) δ 8.53 (d, J = 3.2 Hz, 1H), 8.45-8.42 (m, 1H), 7.46 (t, J = 7.6 Hz, 1H), 7.26 (d, J = 1.2 Hz, 1H), 7.10-7.08 (m, 3H), 6.84 (d, J = 0.8 Hz, 1H), 4.74 (t, J = 4.0 Hz, 1H), 4.45 (t, J = 4.4 Hz, 1H); MS (ESI) m/z 365.1 [C₂₀H₁₇FN₄O₂+H]⁺.

Preparation of 3-fluoro-4-(2-(3-(isoxazol-4-yl)phenoxy)ethoxy)benzonitrile, Compound 84




1H NMR (400 MHz, CDCl₃) δ (ppm): 8.66 (s, 1H), 8.54 (s, 1H), 7.45-7.33 (m, 3H), 7.08 (dd, J = 17.6, 8.0 Hz, 3H), 6.90 (dd, J = 8.4, 1.2 Hz, 1H), 4.49 (t, J = 3.6 Hz, 2H), 4.42 (t, J = 4.8 Hz, 2H); MS (ESI) m/z 325.24 [C₁₈H₁₃FN₂O₃+H]⁺.

Preparation of 6-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)-5-fluoronicotinonitrile, Compound 90




- 1H NMR (400 MHz, DMSO- d_6) δ 8.54 (d, J = 2.8 Hz, 1H), 8.46-8.43 (m, 1H), 8.39 (d, J = 6.0 Hz, 1H), 7.48 (s, 1H), 7.24 (d, J = 2.0 Hz, 1H), 7.11-7.10 (m, 1H), 6.85 (s, 1H), 4.77 (t, J = 4.0 Hz, 2H), 4.59 (t, J = 4.0 Hz, 2H), 2.37-2.33 (m, 1H), 0.90-0.87 (m, 4H); MS (ESI) m/z 366.1 [$C_{19}H_{16}FN_5O_2+H$] $^+$.

Preparation of 5-fluoro-6-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile, Compound 91




- 1H NMR (400 MHz, $CDCl_3$) δ 8.39 (d, J = 6.0 Hz, 1H), 8.23 (d, J = 2.8 Hz, 1H), 7.69-7.67 (m, 1H), 7.26-7.25 (m, 1H), 7.01 (d, J = 1.6 Hz, 1H), 6.89-6.87 (m, 1H), 6.84 (d, J = 0.8 Hz, 1H), 4.80 (t, J = 4.4 Hz, 2H), 4.47 (t, J = 4.8 Hz, 2H), 2.59 (s, 3H); MS (ESI) m/z 340.1 [$C_{17}H_{14}FN_5O_2$] $^+$.

Preparation of 4-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 93



- 1H NMR ($CDCl_3$, 400 MHz): δ 7.61 (d, J = 8.8 Hz, 2H), 7.54 (s, 1H), 7.12 (t, J = 9.2 Hz, 1H), 7.02-6.88 (m, 4H), 6.31 (s, 1H), 4.35 (d, J = 7.2 Hz, 4H), 3.81 (s, 3H); MS (ESI) m/z 338.12 [$C_{19}H_{16}FN_3O_2+H$] $^+$


Preparation of 5-chloro-2-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 98



- 1H NMR (400 MHz, DMSO- d_6) δ 8.50 (d, J = 6.0 Hz, 1H), 7.92 (d, J = 2.4 Hz, 1H), 7.77-7.75 (m, 1H), 7.47 (d, J = 2.0 Hz, 1H), 7.40-7.37 (m, 2H), 7.05-7.03 (m, 1H), 6.82

(d, $J = 2.0$ Hz, 1H), 4.55 (s, 4H), 4.12 (s, 3H) ; MS (ESI) m/z 355.1 $[C_{18}H_{15}ClN_4O_2+H]^+$.


Preparation of 5-chloro-2-((2-((2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 99



1H NMR (400 MHz, $CDCl_3$) δ 8.41 (d, $J = 5.6$ Hz, 1H), 7.56-7.52 (m, 2H), 7.27 (s, 1H), 7.05 (s, 1H), 6.99 (d, $J = 8.8$ Hz, 1H), 6.92 (d, $J = 6.0$ Hz, 1H), 6.86 (s, 1H), 4.52-4.51 (m, 2H), 4.48-4.47 (m, 2H), 2.63 (s, 3H); MS (ESI) m/z 355.0 $[C_{18}H_{15}ClN_4O_2+H]^+$.

10


Preparation of 3-chloro-4-((2-((1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile, Compound 100



1H NMR (400 MHz, $DMSO-d_6$) δ 8.50 (d, $J = 5.6$, 1H), 8.03 (d, $J = 1.6$ Hz, 1H), 7.88-7.80 (m, 1H), 7.50-7.39 (m, 3H), 7.06-7.00 (s, 1H), 6.83 (d, $J = 2.0$ Hz, 1H), 4.58 (s, 4H), 4.12 (s, 3H); MS (ESI) m/z 355.1 $[C_{18}H_{15}ClN_4O_2+H]^+$.

15


Preparation of 3-fluoro-4-((2-((3-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 105



1H NMR ($CDCl_3$, 400 MHz): δ 7.508 (d, 1H, $J = 2.0$ Hz), 7.460-7.376 (m, 2H), 7.092 (t, 1H, $J = 8.4$ Hz), 6.783-6.700 (m, 3 H), 6.310 (d, 1H, $J = 1.6$ Hz), 4.485-4.463 (m, 2H), 4.413-4.391 (m, 2H), 3.902 (s, 3H); MS (ESI) m/z 356.1 $[C_{19}H_{15}F_2N_3O_2+H]^+$.

20

Preparation of 3-fluoro-4-((2-((6-(1-methyl-1H-pyrazol-5-yl)pyridin-2-yl)oxy)ethoxy)benzonitrile, Compound 106




1H NMR ($CDCl_3$, 400 MHz) δ 7.662 (dd, 1H, $J = 7.6$ & 7.2 Hz), 7.497 (d, 1H, $J = 2.0$ Hz), 7.418 (dt, 1H, $J = 8.4$ & 2.0 Hz), 7.375 (dd, 1H, $J = 10.4$ & 2.0 Hz), 7.225 (dd, 1H, $J = 7.6$ & 0.8 Hz), 7.086 (t, 1H, $J = 8.0$ Hz), 6.747 (dd, 1H, $J = 8.0$ & 0.8 Hz), 6.578 (d,

30


¹H, *J* = 2.0 Hz), 4.772 (t, 2H, *J* = 4.8 Hz), 4.481 (t, 2H, *J* = 4.8 Hz), 4.238 (s, 3H); MS (ESI) *m/z* 339.1 [C₁₈H₁₅FN₄O₂+H]⁺.

Preparation of 4-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 108




¹H NMR (CDCl₃, 400 MHz): δ 7.62 (m, 1H), 7.60 (m, 1H), 7.19 (t, *J* = 9.2 Hz, 1H), 7.06-7.05 (d, *J* = 1.2 Hz, 1H), 7.02-6.97 (m, 3H), 6.95 (s, 1H), 6.87-6.85 (dd, *J* = 6.4, 3.0 Hz, 1H), 4.36 (m, 2H), 4.34 (m, 2H), 2.31 (s, 3H); MS (ESI) *m/z* 338.2 [C₁₉H₁₆FN₃O₂+H]⁺.

Preparation of 4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-(trifluoromethyl)benzonitrile, Compound 111




¹H NMR (CDCl₃, 400 MHz): δ 7.89-7.88 (d, *J* = 2.0 Hz, 1H), 7.84-7.81 (dd, *J* = 8.4, 2.0 Hz, 1H), 7.40 (t, *J* = 8.2 Hz, 1H), 7.18-7.16 (d, *J* = 8.8 Hz, 1H), 7.03-6.99 (m, 3H), 6.92-6.91 (m, 1H), 6.84 (t, *J* = 2.2 Hz, 1H), 4.55-4.50 (m, 2H), 4.49-4.40 (m, 2H), 2.37 (s, 3H). MS (ESI) *m/z* 388.1 [C₂₀H₁₆F₃N₃O₂+H]⁺.

Preparation of 5-fluoro-2-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile, Compound 114




¹H NMR (DMSO-d₆, 400 MHz): δ 7.77-7.74 (dd, *J* = 8.0, 3.2 Hz, 1H), 7.61-7.56 (dt, *J* = 8.8, 3.2 Hz, 1H), 7.44 (t, *J* = 8.0 Hz, 1H), 7.39-7.35 (dd, *J* = 9.2, 4.0 Hz, 1H), 7.29 (d, *J* = 1.2 Hz, 1H), 7.09-7.01 (m, 3H), 6.90 (s, 1H), 4.51-4.49 (m, 2H), 4.43-4.41 (m, 2H), 2.29 (s, 3H); MS (ESI) *m/z* 338.2 [C₁₉H₁₆FN₃O₂+H]⁺.

Preparation of 4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)-3-fluorobenzonitrile, Compound 116




1H NMR (DMSO-d₆, 400 MHz): δ 7.88-7.84 (dd, J = 11.4, 2.0, 1H), (td, J = 8.8, 1.4, 1H), 4.47-7.30 (m, 3H), 7.24-7.22 (m, 2H), 7.06-7.03 (m, 1H), 6.43-6.42 (d, J = 1.6, 1H), 4.55-4.52 (m, 2H), 4.44-4.42 (m, 2H), 3.77-3.72 (m, 1H), 0.97-0.91 (m, 4H); MS (ESI) m/z 364.2 [C₂₁H₁₈FN₃O₂+H]⁺.

Preparation of 4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 117



1H NMR (DMSO-d₆, 400 MHz) : δ 7.63-7.61 (d, J = 8.8, 2H), 7.47 (s, 1H), 7.40 (t, J = 8.0, 1H), 7.22-7.20 (d, J = 7.6, 1H), 7.14 (s, 1H), 7.04-7.02 (d, J = 8.8, 2H), 7.01-6.98 (d, J = 8.8, 1H), 6.32 (s, 1H), 4.41 (s, 4H), 3.62 (m, 1H), 1.17 (m, 2H), 0.98-0.97 (m, 2H); MS (ESI) m/z 346.2 [C₂₁H₁₉N₃O₂+H]⁺.

Preparation of 3-chloro-4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile, Compound 118









1H NMR (DMSO-d₆, 400 MHz): δ 7.69-7.68 (d, J = 1.6, 1H), 7.59-7.56 (dd, J = 8.4, 1.6, 1H), 7.47 (s, 1H), 7.40 (t, J = 8.0, 1H), 7.23-7.21 (d, J = 7.2, 1H), 7.16 (s, 1H), 7.09-7.07 (d, J = 8.8, 1H), 7.02-7.00 (d, J = 7.6, 1H), 6.32 (s, 1H), 4.48 (s, 4H), 3.62 (m, 1H), 1.17 (m, 2H), 0.98-0.97 (m, 2H); MS (ESI) m/z 380.2 [C₂₁H₁₈ClN₃O₂+H]⁺.




Example 8: Compound List




The following Table 2 shows a list representing the exemplified compounds of this disclosure, together with the biological activity data.




Table 2. Table listing the structure and IC50 of exemplified compounds




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
1		4-(2-(3-(1H-1,2,4-triazol-1-yl)phenoxy)ethoxy)benzonitrile	0.56
2		4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.027
3		4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile	0.63; 0.49




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
4		5-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)picolinonitrile	0.26; 0.22; 0.39
5		3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.0003; 0.0002; 0.0004
6		3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.002; 0.001; 0.0024




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
7		3-chloro-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.003; 0.003
8		3-methoxy-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.0002; 0.00036
9		3-chloro-4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.0001; 0.00018




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
10		6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile	0.08
11		3-chloro-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile	0.02
12		3-fluoro-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile	0.21

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
13		5-methyl-2-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.011
14		3-fluoro-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.015
15		3-methoxy-4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.0011; 0.0009

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
16		3-methoxy-4-(2-(3-(thiazol-5-yl)phenoxy)ethoxy)benzonitrile	0.008, 0.011
17		4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-fluorobenzonitrile	0.016
18		5-methyl-2-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.003




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
19		4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-methoxybenzonitrile	0.0045
20		4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.00017; 0.00016
21		4-(2-(3-(4H-1,2,4-triazol-4-yl)phenoxy)ethoxy)benzonitrile	0.298




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
22		3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	0.003
23		3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	0.061
24		3-chloro-4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.001




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
25		3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.023
26		1-(3-(2-(4-ethynylphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole	0.069
27		5-(3-(2-(4-fluoro-2-methylphenoxy)ethoxy)phenyl)-1-methyl-1H-pyrazole	0.29




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
28		4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-fluorobenzonitrile	0.0016
29		3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.00008
30		3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0016




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
31		3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0012
32		(R)-3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	0.2
33		(S)-3-chloro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	0.0015




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
34		(R)-3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	1.47
35		(S)-3-fluoro-4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propoxy)benzonitrile	0.0036
36		4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-(trifluoromethoxy)benzonitrile	0.14




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
37		3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0007
38		4-(2-(3-(4H-1,2,4-triazol-4-yl)phenoxy)ethoxy)-3-fluorobenzonitrile	0.31
39		(R)-3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0047




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
40		(S)-3-methyl-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0095
41		(R)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-fluorobenzonitrile	0.000002
42		(S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-fluorobenzonitrile	0.012




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
43		(S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile	0.00068
44		3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.78
45		1-(3-(2-(4-chloro-2-methoxyphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole	0.076




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
46		(R)-3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.001
47		(S)-3-fluoro-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0068, 0.00038
48		(R)-3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.00064




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
49		(S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile	0.0003
50		1-(3-(2-(4-chloro-2-fluorophenoxy)ethoxy)phenyl)-2-cyclopropyl-1H-imidazole	0.012
51		1-(3-(2-(4-ethynyl-2-fluorophenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole	0.02




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
52		3-amino-4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.0036
53		3-fluoro-4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.00044
54		5-fluoro-2-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.032

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
55		(R)-3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.011
56		(S)-3-chloro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.00015
57		4-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.009




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
58		1-(3-(2-(2-fluoro-4-methylphenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole	0.06
59		4-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.02
60		(S)-3-methoxy-4-((1-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.81nM




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
61		4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile	0.88nM
62		4-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.0085
63		3-fluoro-4-((3-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)butan-2-yl)oxy)benzonitrile	0.018; 0.006




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
64		3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.27; 0.47
65		(R)-3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	4.67; 4.39
66		(S)-3-fluoro-4-((1-(3-(isoxazol-5-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.18; 0.19




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
67		4-(2-(3-(isoxazol-4-yl)phenoxy)ethoxy)benzonitrile	0.14; 0.10
68		(R)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methylbenzonitrile	0.0015; 0.013
69		3-fluoro-4-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.0028; 0.0014; 0.0015




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
70		6-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile	0.027
71		6-(2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile	0.044; 0.041
72		(S)-3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.0045




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
73		6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)pyridazine-3-carbonitrile	0.95
74		5-fluoro-6-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile	0.065
75		6-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile	0.06; 0.062




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
76		4-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)-3-fluorobenzonitrile	0.39nM
77		(R)-3-methoxy-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.7nM
78		(R)-3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	75pM




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
79		(S)-3-fluoro-4-((1-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)benzonitrile	0.031
80		4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile	3pM
81		(S)-4-((1-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)propan-2-yl)oxy)-3-methoxybenzonitrile	0.001




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
82		5-fluoro-6-(2-(3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile	0.21; 0.19
83		6-(2-(3-(2-cyclopropyl-1H-imidazol-1-yl)phenoxy)ethoxy)-5-fluoronicotinonitrile	0.0037
84		3-fluoro-4-(2-(3-(isoxazol-4-yl)phenoxy)ethoxy)benzonitrile	0.028

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
85		4-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.015
86		3-fluoro-4-(2-(((6-(2-methyl-1H-imidazol-1-yl)pyridin-2-yl)oxy)ethoxy)benzonitrile	0.017
87		5-fluoro-6-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile	0.033




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
88		4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)-4-fluorophenoxy)ethoxy)-3-fluorobenzonitrile	0.018
89		4-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.0028
90		6-(2-((2-(2-cyclopropyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)-5-fluoronicotinonitrile	0.037




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
91		5-fluoro-6-(2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile	0.12
92		3-fluoro-4-(2-((5-(1-methyl-1H-pyrazol-5-yl)pyridin-3-yl)oxy)ethoxy)benzonitrile	0.3-1
93		4-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.0066




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
94		4-(2-(3-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.0016
95		3-fluoro-4-(2-((2-(3-methyl-4H-1,2,4-triazol-4-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.01
96		4-(2-(2-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.4-3




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
97		5-fluoro-6-((2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)nicotinonitrile	0.47-1
98		5-chloro-2-((2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.01
99		5-chloro-2-((2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.0026




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
100		3-chloro-4-((2-((2-(1-methyl-1H-pyrazol-5-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.0018
101		4-(2-(4-chloro-2-fluorophenoxy)ethoxy)-2-(1-methyl-1H-pyrazol-5-yl)pyridine	0.029
102		1-(3-(2-(4-chloro-2-fluorophenoxy)phenyl)-2-methyl-1H-imidazole	0.068




Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
103		4-(2-(4-chloro-2-fluorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine	0.0031
104		4-(2-(4-bromo-2-fluorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine	0.0017
105		3-fluoro-4-(2-(3-fluoro-5-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.006


Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
106		3-fluoro-4-((6-(1-methyl-1H-pyrazol-5-yl)pyridin-2-yl)oxy)ethoxybenzonitrile	0.3-1
107		5-fluoro-6-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)nicotinonitrile	0.15
108		4-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.051

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
109		6-(2-(4-fluoro-3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)nicotinonitrile	0.17
110		5-(3-(2-(4-chlorophenoxy)ethoxy)phenyl)-1-methyl-1H-pyrazole	0.27
111		4-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)-3-(trifluoromethyl)benzonitrile	0.18

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
112		1-(3-(2-(4-chlorophenoxy)ethoxy)phenyl)-2-methyl-1H-imidazole	0.25
113		3-fluoro-4-(2-(4-fluoro-3-(1-methyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.022
114		5-fluoro-2-(2-(3-(2-methyl-1H-imidazol-1-yl)phenoxy)ethoxy)benzonitrile	0.082

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
115		4-(2-(4-chlorophenoxy)ethoxy)-2-(2-methyl-1H-imidazol-1-yl)pyridine	0.16
116		4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)-3-fluorobenzonitrile	0.0095
117		4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.055

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
118		3-chloro-4-(2-(3-(1-cyclopropyl-1H-pyrazol-5-yl)phenoxy)ethoxy)benzonitrile	0.0022
119		5-fluoro-2-((2-(2-methyl-1H-imidazol-1-yl)pyridin-4-yl)oxy)ethoxy)benzonitrile	0.08
120		4-(2-((2-(1H-tetrazol-1-yl)pyridin-4-yl)oxy)ethoxy)-3-fluorobenzonitrile	0.012

Compound #	Structure	IUPAC Name	ICMT IC50 (μM)
121		4-(2-(4-bromo-2-fluorophenoxy)ethoxy)-2-(1-methyl-1H-pyrazol-5-yl)pyridine	0.07


Industrial Applicability

The compounds as defined above may find a multiple number of applications in which their ability to inhibit methyltransferases such as ICMT is useful. The compounds may also be used in treating or preventing a condition or disorder in which inhibition of a protein methyl transferase and/or co-factor thereof and/or via an unspecified mechanism prevents, inhibits or ameliorates a pathology or a symptomology of the condition. The condition or disorder may be cancer or progeroid disease. The compounds may be particularly useful in treating cancer such as breast cancer, ovarian cancer, pancreatic cancer, leukemia, colorectal carcinoma, lung cancer, hepatocellular carcinoma cancer and other hypervascular tumors as well as angiogenic diseases.

It will be apparent that various other modifications and adaptations of the invention will be apparent to the person skilled in the art after reading the foregoing disclosure without departing from the spirit and scope of the invention and it is intended that all such modifications and adaptations come within the scope of the appended claims.

CLAIMS

1. A compound having the following Formula (I):



and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof;

wherein:

 represents an aromatic ring system;

Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

B₁, B₂ and B₃ are independently selected from C, CH, or N;

Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

R₁, R₂, and R₃ are independently absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;


R₄ and R₅ are independently selected from H or aliphatic;

n is an integer selected from 0 to 5, wherein when n is more than 1, each R₁ substituted on Ring A may be the same or different;

m is an integer selected from 0 to 4, wherein when m is more than 1, each R₂ substituted on Ring B may be the same or different; and

p is an integer selected from 0 to 5, wherein when p is more than 1, each R_3 substituted on Ring C may be the same or different.

2. A compound having the following Formula (I):



and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof;

wherein:

○ represents an aromatic ring system;

Ring A is a 5- or 6- membered carbocyclic ring system, wherein 1 to 4 carbon atoms may be optionally replaced with a heteroatom;

B_1 , B_2 and B_3 are independently selected from C, CH, or N, wherein when one of B_1 , B_2 or B_3 is N, the remaining B_1 , B_2 or B_3 is C or CH;

Ring C is a 5- or 6- membered carbocyclic ring system, wherein 1 to 3 carbon atoms may be optionally replaced with a heteroatom;

R_1 is absent or selected from H, OH, halogen, optionally substituted alkyl, haloalkyl, CF_3 , CHF_2 , optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

R_2 is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF_3 , CHF_2 , optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted

cycloalkenyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl;

R₃ is absent or selected from H, OH, cyano, halogen, optionally substituted alkyl, haloalkyl, CF₃, CHF₂, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted amino, optionally substituted acyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocycloalkyl, or optionally substituted aryl;


R₄ and R₅ are independently selected from H or aliphatic;

n is an integer selected from 0 to 5, wherein when n is more than 1, each R₁ substituted on Ring A may be the same or different;

m is an integer selected from 0 to 4, wherein when m is more than 1, each R₂ substituted on Ring B may be the same or different, and m is 1 when R₂ is halogen; and

p is an integer selected from 0 to 5, wherein when p is more than 1, each R₃ substituted on Ring C may be the same or different.

3. The compound of claim 1 or 2, wherein Ring A is of the formula (aa):



(aa)


wherein:

* represents the point of attachment to Ring B; and

A₁, A₂, A₃, A₄ and A₅ are independently selected from C, CH, N, O or S.

4. The compound of any one of claims 1 to 3, wherein Ring A is of the formula (ab):

137



(ab)

wherein:

* represents the point of attachment to Ring B; and

A₁ is N or C;


A₂ is N, S, CH or C;

A₃ is C, CH, N, or O;

A₄ is N; and


A₅ is C, CH, O, or N.

5. The compound of any one of claims 1 to 4, wherein Ring A is of formula (ac):




(ac)

6. The compound of any one of claims 1 to 4, wherein Ring A is selected from the group consisting of formulas (a1) to (a8):



wherein * represents the point of attachment to Ring B.

7. The compound of any one of claims 1 to 6, wherein Ring A is selected from the group consisting of formulas (a1') to (a8'):




wherein * represents the point of attachment to Ring B.

8. The compound of any one of claims 1 to 7, wherein R_1 is alkyl or cycloalkyl.


9. The compound of any one of claims 1 to 8, wherein R_1 is methyl or cyclopropyl.

10. The compound of any one of claims 1 to 9, wherein Ring A is selected from the group consisting of formulas (a1'') to (a10''):



wherein * represents the point of attachment to Ring B.

11. The compound of any one of claims 1 to 10, wherein Ring B is of formula (bb1) or formula (bb2):




wherein:

* represents the point of attachment to Ring A; and

** represents the point of attachment to the $-O(CHR_4)-$ moiety of formula (I).


12. The compound of any one of claims 1 to 11, wherein Ring A is a 5-membered carbocyclic ring system, wherein 1 to 4 carbon atoms is replaced with a heteroatom; and Ring B is of formula (bb1) or formula (bb2).

13. The compound of any one of claims 1 to 12, wherein Ring B is selected from the group consisting of formulas (b1) to (b5):




14. The compound of any one of claims 1 to 13, wherein R_2 is H or F.

15. The compound of any one of claims 1 to 14, wherein Ring B is selected from the group consisting of formulas (b1') to (b9'):



141




wherein:

* represents the point of attachment to Ring A; and

** represents the point of attachment to the $-O(CHR_4)-$ moiety of formula (I).

16. The compound of any one of claims 1 to 15, wherein Ring C is of formula (ca):



wherein:


○ represents an aromatic ring system;

*** represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I);

and

C₁, C₂, C₃, C₄, C₅ and C₆ are independently selected from C, CH, or N.

17. The compound of any one of claims 1 to 16, wherein Ring C is of the formula (cb):



wherein:

○ represents an aromatic ring system;


*** represents the point of attachment to the $-\text{O}(\text{CHR}_5)-$ moiety of formula (I);

C_1 and C_2 are independently C, CH or N;

C_4 , C_5 , and C_6 are C or CH; and

C_3 is C.

18. The compound of any one of claims 1 to 17, wherein Ring C is of formula (cc):



wherein:

○ represents an aromatic ring system;

*** represents the point of attachment to the $-\text{O}(\text{CHR}_5)-$ moiety of formula (I);


C_1 and C_2 are independently CH or N;

C_5 is CH; and


C_3 , C_4 , and C_6 are C.

19. The compound of any one of claims 1 to 18, wherein Ring C is selected from the group consisting of formulas (c1) to (c4):

143



20. The compound of any one of claims 1 to 19, wherein Ring C is selected from the group consisting of formulas (c1') to (c6'):




wherein:


$***$ represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I).

21. The compound of any one of claims 1 to 20, wherein R_3 is halogen, $-CN$, alkoxy, amino, haloalkyl, haloalkyloxy, or alkyl.

22. The compound of any one of claims 1 to 21, wherein R_3 is fluoro, chloro, bromo, $-CN$, methoxy, methyl, $-NH_2$, $-OCF_3$, CF_3 , or CHF_2 .

23. The compound of any one of claims 1 to 22, wherein Ring C is selected from the group consisting of (c1'') to (c22''):





wherein:


*** represents the point of attachment to the $-O(CHR_5)-$ moiety of formula (I).

24. A compound of any one of claims 1 to 23, wherein R_4 and R_5 are independently H or alkyl.

25. A compound of any one of claims 1 to 24, wherein R_4 and R_5 are independently H or methyl, ethyl, propyl, or butyl.

26. A compound of any one of claims 1 to 25, wherein R_4 and R_5 are H; or R_4 is H and R_5 is methyl; or R_4 is methyl and R_5 is H; or R_4 is methyl and R_5 is methyl.

27. A compound of claim 1 or claim 2, having the following Formula (IA):




and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof, wherein:


○ represents an aromatic ring system; and

A₁, A₂, A₃, A₄ and A₅, R₁, n, B₁, B₂, B₃, R₂, m, R₄, R₅, C₁, C₂, C₃, C₄, C₅, C₆, R₃ and p are as defined in any one of claims 1 to 27.

28. A compound of claim 1 or claim 2, having the following Formula (IE) or Formula (IF):



(IE)















(IF)









and/or tautomers, enantiomers, solvates, hydrates, prodrugs and pharmaceutically acceptable salts thereof, wherein:

















○ represents an aromatic ring system; and

















A₁, A₂, A₃, A₄ and A₅, R₁, n, B₁, B₂, B₃, R₂, m, R₄, R₅, C₁, C₂, C₃, C₄, C₅, C₆, R₃ and p are as defined in any one of claims 1 to 27.

















29. A compound of any one of claims 1 to 28, selected from the group consisting of:




































































30. A pharmaceutical composition comprising a compound of any one of claims 1 to 29, or a pharmaceutically acceptable form or prodrug thereof, and a pharmaceutically acceptable excipient.

31. A method of inhibiting methylation of isoprenylated cysteine or isoprenylcysteine caused by isoprenylcysteine carboxyl methyltransferase (ICMT) comprising contacting the compound of any one of claims 1 to 29, or a pharmaceutical composition according to claim 30, with ICMT.

32. A method of inhibiting ICMT in a cell comprising contacting said cell with a compound of any one of claims 1 to 29, or a pharmaceutically acceptable form or prodrug thereof, or a pharmaceutical composition according to claim 30.

33. The method according to claim 32, wherein the inhibition of ICMT further comprises the inhibition of cell proliferation.

34. A method of treating a ICMT-related disorder comprising administering to a subject in need of treatment a compound of any one of claims 1 to 29, or pharmaceutical, or a composition according to claim 30.

35. The method according to claim 34, wherein the disorder is cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS).
36. The method according to claim 35, wherein the cancer is one linked to mutant Ras overactivity.
37. The method according to claim 35 wherein the cancer is selected from the group consisting of hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.
38. The method according to any one of claims 33 to 37, further comprising the step of administering an additional therapeutic agent to the subject.
39. The compound according to any one of claims 1 to 29, or a pharmaceutically form or prodrug thereof, or a composition according to claim 30 for use in therapy.
40. A compound of any one of claims 1 to 29, or a pharmaceutically form or prodrug thereof, or a composition according to claim 30 for use in the treatment and/or prevention of cancer and/or progeroid diseases.
41. The compound according to any one of claims 1 to 29, or a pharmaceutically form or prodrug thereof, or a composition according to claim 30, for use in the treatment of a ICMT-related disorder.
42. The compound or pharmaceutically form or prodrug thereof according to claim 41, wherein the disorder is cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS).
43. The compound or pharmaceutically form or prodrug thereof according to claim 42, wherein the cancer is one linked to mutant Ras overactivity.
44. The compound or pharmaceutically form or prodrug thereof of claim 42 or claim 43, wherein the cancer is hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.

45. The compound or pharmaceutically form or prodrug thereof of any one of claims 39 to 44, wherein the compound is to be administered in combination with an additional therapeutic agent.

46. Use of a compound according to any one of claims 1 to 29, or a pharmaceutically form or prodrug thereof, or a composition according to claim 30, in the manufacture of a medicament for the treatment and/or prevention of cancer and/or progeroid diseases.

47. Use of a compound according to any one of claims 1 to 29, or a pharmaceutically form or prodrug thereof, or a composition according to claim 30, in the manufacture of a medicament for the treatment of a ICMT-related disorder.

48. The use of claim 47, wherein the disorder is cancer, premature ageing, or Hutchinson-Gilford progeria syndrome (HGPS).

49. The use of claim 48, wherein the cancer is one linked to mutant Ras overactivity.

50. The use of claim 48 or 49, wherein the cancer is hepatocellular carcinoma cancer, breast cancer, ovarian cancer, colorectal carcinoma, lung cancer, pancreatic cancer or leukemia.

51. The use of any one of claims 46 to 50, wherein the medicament is to be administered in combination with an additional therapeutic agent.


52. The compound of any one of claims 1 to 29, having an ICMT IC₅₀ of less than about 5 μ M.

53. The compound of any one of claims 1 to 29, having an ICMT IC₅₀ of about 0.001 μ M to about 5 μ M.

54. A process for synthesizing a compound according to any one of claims 1 to 29, comprising:


(a) reacting a compound of formula (III):

158



wherein R_1 , R_2 , R_4 , R_5 , B_1 , B_2 , B_3 , n and m are as defined in claim 1 or 2, and R_6 is a leaving group,


in an organic solvent in the presence of a base with a compound of formula (IV):



wherein R_3 and p are as defined in claim 1 or 2, and R_7 is $-OH$;


or

(b) reacting a compound of formula (III):



wherein R_1 , R_2 , R_4 , R_5 , B_1 , B_2 , B_3 , n and m are as defined in claim 1 or 2, and R_6 is OH or an alkoxide,

in an organic solvent in the presence of a base with a compound of formula (IV):



(IV)

wherein R_3 and p are as defined in claim 1 or 2, and R_7 is halogen.