

The present invention relates to storage-stable detergents which, in addition to at least diperoxydodecanedioic acid or salts thereof, contain at least one specific fluorescent whitening agent of the stilbene type, and to their use for washing textiles.


Because of the increased content of synthetic fibres and fibre blends in the textiles manufactured today and the desire no longer to have to wash coloured laundry separately and for energy saving measures, in many countries laundry is no longer washed at 90°C-95°C or at the boil but at lower temperatures. This means that it has been necessary for the perborates which have usually so far been contained in detergents and act as bleaching agents to be activated by auxiliaries, such as tetraacetythylenediamine (TAED) in order to achieve acceptable bleaching effects even at washing temperatures of 60-80°C. At still lower washing temperatures, even the perborate/activator systems no longer give satisfactory results.

Detergents which contain stronger bleaching agents, for example peracids, have therefore been described for some time (German Offenlegungsschrift 2,756,583, EP-A-145,438, GB 2,141,754, GB 2,141,755, US 4,028,263 and GB 59,272). Although these novel bleaching agents on the one hand already exhibit outstanding bleaching effects at temperatures from 20°C, on the other hand they destroy the customary fluorescent whitening agents contained in detergents.

It has now been found that specific stilbene compounds surprisingly have an excellent stability in detergents which contain such strong bleaching agents. Under average storage conditions and even under intensified conditions (temperatures above 30°C and atmospheric humidity above 60%), these specific stilbene fluorescent whitening agents are completely stable, or are at most degraded to a degree which causes no trouble in practice, in the detergent for several months.

The present invention thus relates to storage-stable dry detergents comprising 0.5 to 30% of an organic peracid or salts thereof or mixtures thereof with other peracids or salts thereof, and a fluorescent whitening agent or a mixture of fluorescent whitening agents, which contain


- a) diperoxydodecanedioic acid as the organic peracid and
b) 0.03% to 0.5% of a fluorescent whitening agent or a mixture of fluorescent whitening agents of the formula (I)




in which A is hydrogen, $-\text{SO}_3\text{M}$, halogen, phenyl or phenylpyrazol-1-yl; B is benzofuran-2-yl if A is hydrogen, 2-(stilben-4-yl)-oxadiazol-5-yl if A is halogen, phenylpyrazol-1-yl if A is phenylpyrazol-1-yl, benzoxazol-2-yl if A is $-\text{SO}_3\text{M}$ or phenyl, or naphthoxazol-2-yl if A is hydrogen; R_1 and R_2 independently of one another are hydrogen, halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; M is hydrogen or one equivalent of a non-chromophoric cation and n is zero to 2, but where the compound of the formula (I) contains at least one $-\text{SO}_3\text{M}$ group.

M in formula (I) is, for example, an alkaline earth metal, such as magnesium or calcium, but preferably an alkali metal, such as lithium, sodium or potassium, or substituted or unsubstituted ammonium, such as ammonium, mono-, di- or triethanol-ammonium, mono-, di- or tripropanol-ammonium or tri- or tetramethyl-ammonium.



In particular, the detergents contain as whitening agents compounds of the formulae




in which R is H or halogen and M is as defined above,



in which R_1 and R_2 are H or C_1 - C_4 alkyl, R_3 is $-\text{SO}_3\text{M}$ or



in which M is as defined above, and/or



in which M is as defined above.

All these fluorescent whitening agents of the formula I to V are known and can be prepared by known methods.

The peracids and salts thereof are inorganic or organic compounds which are described in the literature or are commercially available and which preferably already bleach textiles at temperatures from 20°C. The organic peracid is diperoxydodecanedioic acid (DPDDA) in question [sic]. The amount of organic peracid to be used is preferably 0.5% to 10%, in particular 1% to 5%. However, it is also possible for highly active inorganic peracids, such as persulfate and/or percarbonate, to be used. The amount of these inorganic peracids is 1% to 20%, in particular 10% to 20%, based on the total weight of detergent. Optionally and advantageously they are used in combination with small amounts of catalytically acting divalent metal salts, such as are described in US 4,655,782 and US 4,655,953. Metal salts of copper and/or manganese are preferably used.

Mixtures of the organic and inorganic peracids or peracid salts or only mixtures of the inorganic peracids can of course also be used.

The peracids are added to the detergent by dry mixing, for example with the aid of automatic screw metering systems and/or fluidized bed mixers.

The detergents are dry detergents of customary compositions. As a rule, in addition to the combination according to the invention of peracid and fluorescent whitening agent, they contain, for example, anionic, nonionic, amphoteric and/or cationic surfactants, builders, for example pentasodium tripolyphosphate or substitute products, such as phosphonates, polycarboxylates, acrylic/maleic copolymers, zeolites, nitrilotriacetate and ethylenediaminetetraacetic acid, soil-suspending agents, for example sodium carboxymethylcellulose, salts for adjusting the pH, for example alkali metal or alkaline earth metal silicates, foam regulators, for example soaps, salts for adjusting the spray-drying and granulation properties, for example sodium sulfate, perfumes and if appropriate antistatic and softening agents, enzymes, photobleaching agents, pigments and/or toning agents. These constituents should of course be stable towards the bleaching system used.

As a result of the combination according to the invention, it is possible to provide detergents which meet the customary standard, for example as regards to washing power, spot removal and refreshing the appearance of the washed articles, even if washing is carried out at temperatures of 20°C-60°C. Coloured laundry and white laundry can thus advantageously be washed together regardless of the fibre.

The following examples illustrate the invention without limiting it thereto. Percentage data are percentages by weight.

Detergent formulation

The detergents of the following examples are prepared in two stages by customary processes:

Granules A are first prepared by drying and granulating a slurry consisting of about 1 part of water and 1 part of detergent of the following composition:

- 84 g of linear dodecylbenzenesulfonate
- 31 g of tallow alcohol tetradecane-ethylene glycol ether (14 mol of ethylene oxide)
- 37 g of Na soap (chiefly of behenic acid and C₁₄-C₂₀)
- 458 g of Na tripolyphosphate

79 g of Na silicate
20 g of Mg silicate
12 g of carboxymethylcellulose
2 g of ethylenediaminetetraacetate
222 g of Na sulfate
1 g of fluorescent whitening agent (FWA) according to the formula
(II) - (V).

The peracid B (15 g of DPDDA) is homogeneously admixed in the dry state to 500 g of these granules A, which have a residual moisture content of 5% after drying.

Storage test

Samples of the detergent D (granules A + peracid B) thus obtained are:







- on the one hand, for control of the starting value, immediately analyzed for the content of FWA by extraction and measurement of the spectrophotometric absorbance (theoretical: 0.1% of FWA, based on the weight of the granules A),
- on the other hand kept in cardboard packets suitable for commercial washing powder, that is to say with a coating, under selected and controlled conditions of temperature and atmospheric humidity for certain periods of time. After storage, the FWA content of each detergent is immediately determined. The difference from the starting value is expressed as a percentage and is a measure of the stability of the FWA towards the corresponding bleaching agent in the washing powder.


The abovementioned FWA determination is carried out as follows:

The washing powder is homogenized thoroughly by grinding and 1 g thereof is admixed with 200 ml of solvent consisting of 9 parts of dimethyl sulfoxide and 1 part of water, and the mixture is stirred at room temperature for 30 minutes. It is then centrifuged for 30 minutes. A sample of the clear solution thus obtained is transferred with a pipette into a 1 cm quartz cell and its absorbance is measured in the UV range at the absorption maximum against a standard solution of the particular FWA. The absorbance is proportional to the FWA concentration. The


reproducibility of the results is about $\pm 1\%$ if the test conditions are kept exactly the same.

Example : The percentage FWA loss is determined under the conditions described above. Storage is in a closed packet at 20 to 25°C. After 6 months, the FWA loss is as follows:





2 %




0 %



Patent claims

1. A storage-stable dry detergent comprising 0.5 to 30% of an organic peracid or salts thereof or mixtures thereof with other peracids or salts thereof, and a fluorescent whitening agent or a mixture of fluorescent whitening agents, which contains

- a) diperoxidodecanedioic acid as the organic peracid and
- b) 0.03% to 0.5% of a fluorescent whitening agent or a mixture of fluorescent whitening agents of the formula (I)




in which A is hydrogen, $-\text{SO}_3\text{M}$, halogen, phenyl or phenylpyrazol-1-yl; B is benzofuran-2-yl if A is hydrogen, 2-(stilben-4-yl)-oxadiazol-5-yl if A is halogen, phenylpyrazol-1-yl if A is phenylpyrazol-1-yl, benzoxazol-2-yl if A is $-\text{SO}_3\text{M}$ or phenyl, or naphthoxazol-2-yl if A is hydrogen; R_1 and R_2 independently of one another are hydrogen, halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; M is hydrogen or one equivalent of a non-chromophoric cation and n is zero to 2, but where the compound of the formula (I) contains at least one $-\text{SO}_3\text{M}$ group.

2. A detergent according to claim 1, which contains peracids or salts thereof which bleach textile material at a temperature from 20°C .

3. A detergent according to claim 2, which contains, as a further peracid or peracid salt, a persulfate and/or a percarbonate with the addition of a catalyst.


4. A detergent according to claim 1, which contains, as the fluorescent whitening agent, a compound of the formula (II)




in which R is H or halogen and M is as defined in claim 1.




5. A detergent as claimed in claim 1, which contains, as the fluorescent whitening agent, a compound of the formula (III)



in which R_1 and R_2 are H or C_1 - C_4 alkyl, R_3 is $-SO_3M$ or




6. A detergent as claimed in claim 1, which contains, as the fluorescent whitening agent, a compound of the formula (IV)



in which M is as defined in claim 1.

7. A detergent according to claim 1, which contains, as the fluorescent whitening agent, a compound of the formula (V)



in which M is as defined in claim 1.

8. The use of a detergent according to any one of claims 1 to 7 for washing textiles at temperatures of 20-60°C.

9. The preparation of a detergent according to any one of claims 1 to 7 by dry mixing with the aid of automatic screw metering systems and/or fluidized bed mixers.