5-MEMBERED HETEROCYCLE-BASED P38 KINASE INHIBITORS

FIELD

Provided herein are 5-membered heterocycle-, including pyrazole- and imidazole-, based compounds which have cytokine inhibitory activity. Also provided are uses of the compounds for treating conditions associated with p38α and β kinases and for treating p38 kinase-associated conditions.

BACKGROUND

A large number of cytokines participate in the inflammatory response, including IL-1, IL6, IL-8 and TNF-α. Overproduction of cytokines such as IL-1 and TNF-α are implicated in a wide variety of diseases, including inflammatory bowel disease, rheumatoid arthritis, psoriasis, multiple sclerosis, endotoxin shock, osteoporosis, Alzheimer's disease, and congestive heart failure, among others (Henry *et al.*, Drugs Fut., 24:1345-1354 (1999); Salituro *et al.*, Curr. Med. Chem., 6:807-823 (1999)). Evidence in human patients indicates that protein antagonists of cytokines are effective in treating chronic inflammatory diseases, such as, for example, a monoclonal antibody to TNF-α (Remicade) (Rankin *et al.*, Br. J. Rheumatol., 34:334-342 (1995)), and a soluble TNF-α receptor-Fc fusion protein (Etanercept) (Moreland *et al.*, 25 Ann. Intern. Med., 130:478-486 (1999)).

The biosynthesis of TNF-α occurs in many cell types in response to an external stimulus, such as, for example, a mitogen, an infectious organism, or trauma. Important mediators of TNF-α production are the mitogen-activated protein (MAP) kinases, and in particular, p38 kinases. These kinases are activated in response to various stress stimuli, including but not limited to proinflammatory cytokines, endotoxin, ultraviolet light, and osmotic shock. Activation of p38 requires dual phosphorylation by upstream MAP kinase kinases (MKK3 and MKK6) on threonine and tyrosine within a Thr-Gly-Tyr motif characteristic of p38 isozymes.

There are four known isoforms of p38, *i.e.*, p38α, p38β, p38γ, and p38δ. The α and β isoforms are expressed in inflammatory cells and are key modulators of TNF-α production. Inhibiting the p38α and β enzymes in cells results in reduced levels of TNF-α expression. Also, administering inhibitors of p38α and β in animal models of inflammatory disease has proven that such inhibitors are effective in treating


those diseases. Accordingly, the p38 enzymes serve an important role in inflammatory processes mediated by IL-1 and TNF-α. See, *e.g.*, U.S. Patent Nos. 6,277,989, 6,130,235, 6,147,080, 5,945,418, 6,251,914, 5,977,103, 5,658,903, 5,932,576, and 6,087,496; and in International Patent Application Publication Nos. WO 00/56738, WO 01/27089, WO 01/34605, WO 00/12497, WO 00/56738, WO 00/12497 and WO 00/12074. See also, U.S. Patent Nos. 6,376,527; 6,316,466 and 6,444,696; and International Patent Application Publication Nos. WO 99/57101, WO 02/40486, WO 03/032970, WO 03/033482, WO 03/032971, WO 03/032986, WO 03/032980, WO 03/032987, WO 03/033483, WO 03/033457 and WO 03/032972.

Thus, there is a need for inhibitors of p38 kinases, including p38a and p38b kinase, for treatment, prevention, or amelioration of one or more symptoms of diseases and disorders associated with p38 kinase activity.

SUMMARY

Provided herein are compounds, compositions and methods of treating, preventing, or ameliorating one or more symptoms of conditions associated with p38 kinase activity. In one embodiment, the compounds for use in the compositions and methods are pyrazole- or imidazole-based compounds. In another embodiment, the pyrazole- or imidazole-based compounds are useful as kinase inhibitors, including p38α and p38β kinases.

In one embodiment, the compounds provided herein have the formula:



or a pharmaceutically acceptable derivative thereof, where:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, optioally substituted alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted alkoxy, optionally substituted heterocyclyloxy or alkylamino;

G is an aryl, aralkyl, cycloalkyl, heteroaryl, heteroaralkyl or a heterocyclyl ring optionally fused to a phenyl ring, and is substituted with R^3 and R^4, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom, or G is OR^{83} or $NR^{80}R^{81}$;

B is an aryl or heteroaryl ring;

C is a 5-membered heteroaryl ring containing one or two heteroatoms in the ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$;

each R^{80} and R^{81} is independently hydrogen, alkyl, cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl;

R^{83} is hydrogen, alkyl, cycloalkyl, heteroaryl or optionally substituted heteroaryl;

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n-$ (where n is an integer from 0 to 2); and R^9 is halo, cyano, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(t) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) -OC(O)-alkylene-CO_2H or -OC(O)-NR'R" (where R' and R" are independently hydrogen or alkyl);

(bb) heteroarylalkenylene or heteroarylalkynylene;

(cc) hydrogen;

(dd) halo;

(ee) pseudohalo;

(ff) hydroxy;

(gg) optionally substituted alkoxy;

(hh) $C(L)R^{40}$, where L is O, S or NR^{55}; R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{56}$, halo pseudohalo, OR^{55}, SR^{55}, $NR^{57}R^{58}$ or $SiR^{52}R^{53}R^{54}$; where R^{52}, R^{53} and R^{54} are selected as in (i) or (ii) as follows (i) R^{52}, R^{53} and R^{54} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or (ii) any two of R^{52}, R^{53} and R^{54} together form alkylene, alkenylene, alkynylene, heteroalkylene; and the other is selected as in (i); R^{55} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{56} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{64}R^{65}$; where R^{64} and R^{65} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{66} or $NR^{62}R^{63}$, or R^{64} and R^{65} together form alkylene, alkenylene, alkynylene, heteroalkylene, where R^{66} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{57} and R^{58} are selected as in (i) or (ii) as follows (i) R^{57} and R^{58} are each independently hydrogen, optionally substituted alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55}, $NR^{67}R^{68}$ or $C(L)R^{69}$, where R^{67} and R^{68} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl, or together form alkylene, alkenylene, alkynylene, heteroalkylene; and R^{69} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$, where R^{70} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl; or (ii) R^{57} and R^{58} together form alkylene, alkenylene, alkynylene, heteroalkylenem or alkylenoxyalkylene; R^{62} and R^{63} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, or R^{62} and R^{63} together form alkylene, alkenylene, alkynylene, heteroalkylene; and

(ii) optionally substituted alkyl;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(1) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy.

Also provided herein are pharmaceutical compositions containing a compound provided herein in combination with a pharmaceutically acceptable carrier.

Methods of treating, preventing or ameliorating one or more symptoms of cytokine mediated disease in a mammal, by administering to a mammalian patient, in need of such treatment, a compound of formula I are provided. Diseases and disorders treated, prevented, or whose symptoms are ameliorated, include, but are not limited to, chronic inflammatory diseases, inflammatory bowel disease, rheumatoid arthritis, psoriasis, multiple sclerosis, endotoxin shock, osteoporosis, Alzheimer's disease, and congestive heart failure.

Methods of preventing or inhibiting inflammatory responses using the compounds and compositions provided herein are also provided.

Further provided are methods of inhibiting p38 kinases, including p38α and p38β kinases, using the compounds and compositions provided herein.

Articles of manufacture are provided containing packaging material, a compound or composition provided herein which is useful for treating, preventing, or ameliorating one or more symptoms of p38 kinase-mediated diseases or disorders, and a label that indicates that the compound or composition is useful for treating, preventing, or ameliorating one or more symptoms of p38 kinase-mediated diseases or disorders.

DETAILED DESCRIPTION

A. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which the invention(s) belong. All patents, patent applications, published applications and publications, Genbank sequences, databases, websites and other published materials referred to throughout the entire disclosure herein, unless noted otherwise, are incorporated by reference in their entirety. In the event that there are a plurality of definitions for terms herein, those in this section prevail. Where reference is made to a URL or other such identifier or address, it understood that such identifiers can change and particular information on the internet can come and go, but equivalent information can be found by searching the internet. Reference thereto evidences the availability and public dissemination of such information.

As used herein, p38α refers to the enzyme disclosed in Han *et al.* (1995) *Biochim.*

Biophys. Acta 1265(2-3):224-7. As used herein, p38β refers to the enzyme disclosed in Jiang *et al.* (1996) *J. Biol. Chem. 271(30)*:17920-6. As used herein, p38γ refers to the enzyme disclosed in Li *et al.* (1996) *Biochem. Biophys. Res. Commun. 228*: 334-340. As used herein, p38δ refers to the enzyme disclosed in Wang *et al.* (1997) *J. Biol. Chem. 272(38)*:23668-74.

As used herein, pharmaceutically acceptable derivatives of a compound include salts, esters, enol ethers, enol esters, acetals, ketals, orthoesters, hemiacetals, hemiketals, acids, bases, solvates, hydrates or prodrugs thereof. Such derivatives may be readily prepared by those of skill in this art using known methods for such derivatization. The compounds produced may be administered to animals or humans without substantial toxic effects and either are pharmaceutically active or are prodrugs. Pharmaceutically acceptable salts include, but are not limited to, amine salts, such as but not limited to N,N'-dibenzylethylenediamine, chloroprocaine, choline, ammonia, diethanolamine and other hydroxyalkylamines, ethylenediamine, N-methylglucamine, procaine, N-benzylphenethylamine, 1-para-chlorobenzyl-2-pyrrolidin-1'-ylmethyl-benzimidazole, diethylamine and other alkylamines, piperazine and tris(hydroxymethyl)aminomethane; alkali metal salts, such as but not limited to lithium, potassium and sodium; alkali earth metal salts, such as but not limited to barium, calcium and magnesium; transition metal salts, such as but not limited to zinc; and other metal salts, such as but not limited to sodium hydrogen phosphate and disodium phosphate; and also including, but not limited to, nitrates, borates, methanesulfonates, benzenesulfonates, toluenesulfonates, salts of mineral acids, such as but not limited to hydrochlorides, hydrobromides, hydroiodides and sulfates; and salts of organic acids, such as but not limited to acetates, trifluoroacetates, oxalates, benzoates, salicylates, maleates, lactates, malates, tartrates, citrates, ascorbates, succinates, butyrates, valerates and fumarates. In addition, zwitterions ("inner salts") may be formed. In certain embodiments, salt forms of the compounds improve the compounds' dissolution rate and oral bioavailability. Pharmaceutically acceptable esters include, but are not limited to, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cycloalkyl and heterocyclyl esters of acidic groups, including, but not limited to, carboxylic acids, phosphoric acids, phosphinic acids, sulfonic acids, sulfinic acids and boronic acids. Pharmaceutically acceptable enol ethers include, but

are not limited to, derivatives of formula C=C(OR) where R is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cycloalkyl or heterocyclyl. Pharmaceutically acceptable enol esters include, but are not limited to, derivatives of formula C=C(OC(O)R) where R is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cycloalkyl or heterocyclyl. Pharmaceutically acceptable solvates and hydrates are complexes of a compound with one or more solvent or water molecules, or 1 to about 100, or 1 to about 10, or one to about 2, 3 or 4, solvent or water molecules.

"Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl, pentyl, and the like.

The term "cycloalkyl" refers to a saturated or partially unsaturated nonaromatic cyclic hydrocarbon ring system, preferably containing 1 to 3 rings and 3 to 7 carbons per ring which may be further fused with an unsaturated C_3-C_7 carbocylic ring. Exemplary groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cycloctyl, cyclodecyl, cyclododecyl, and adamantyl. A "substituted cycloalkyl" is substituted with one or more alkyl or substituted alkyl groups as described above, or one or more groups described above as alkyl substituents. The expression "lower cycloalkyl" refers to an unsubstituted saturated or unsaturated nonaromatic cyclic hydrocarbon ring system containing 3 to 5 carbon atoms.

"Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, propylene, 2-methylpropylene, pentylene, and the like.

"Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenyl, propenyl, and the like.

"Alkenylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenylene, propenylene, and the like.

"Alkynyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, e.g., ethynyl, propynyl, and the like.

"Alkynylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, e.g., ethynylene, propynylene, and the like.

"Alkoxy" means a radical -OR where R is alkyl as defined above, e.g., methoxy, ethoxy, propoxy, 2-propoxy, the like.

"Acyl" means a radical -C(O)R where R is alkyl or haloalkyl e.g., acetyl, trifluoroacetyl, and the like.

"Acylamino" means a radical -NRC(O)R' where R is hydrogen or alkyl, and R' is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, e.g., acetylamino, 2-amino-2-methylpropionamide, and the like.

"Halo" means fluoro, chloro, bromo, or iodo, generally fluoro and chloro.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g., $-CH_2Cl$, $-CF_3$, $-CH_2CF_3$, $-CH_2CCl_3$, and the like.

"Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms e.g., phenyl, 1-naphthyl, 2-naphthyl, and the like. The aryl ring may optionally be fused to a 5-, 6- or 7- membered monocyclic saturated ring optionally containing 1 or 2 heteroatoms independently selected from oxygen, nitrogen or sulfur, the remaining ring atoms being C where one or two C atoms are optionally replaced by a carbonyl group. Representative aryl radicals with fused rings include, but are not limited to, 2,3-dihydrobenzo[1,4]dioxan, chroman, isochroman, 2,3-dihydrobenzofuran, 1,3-dihydroisobenzofuran, benzo[1,3]dioxole, 1,2,3,4-tetrahydroisoquinoline, 1,2,3,4tetrahydroquinoline, 2,3-dihydro-lH-indole, 2,3-dihydro-lH-isoindole, benzimidazol-2-one, 3H-benzoxazol-2-one, and the like.

"Heteroaryl" means a monovalent monocyclic or bicyclic aromatic radical of 5 to 10 ring atoms containing one, two, or three ring heteroatoms selected from N, 0, or S, the remaining ring atoms being C. The term also includes those radicals where a heteroatom within the ring has been oxidized or quaternized, such as, for example, to form an N-oxide or a quaternary salt. Representative examples include,

but are not limited to, thienyl, benzothienyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinolinyl, quinoxalinyl, imidazolyl, furanyl, benzofuranyl, thiazolyl, isoxazolyl, benzisoxazolyl, benzimidazolyl, triazolyl, pyrazolyl, pyrrolyl, indolyl, 2-pyridonyl, 4-pyridonyl, N-alkyl-2-pyridonyl, pyrazinonyl, pyridazinonyl, pyrimidinonyl, oxazolonyl, and their corresponding N-oxides, (e.g. pyridyl N-oxide, quinolinyl N-oxide), their quaternary salts and the like.

"Heterocycle" or "heterocyclyl" means a cyclic nonaromatic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from N, 0, or $S(O)_n$, (where n is an integer from 0 to 2), the remaining ring atoms being C where one or two C atoms are optionally replaced by a carbonyl group. The term also includes those radicals where a ring nitrogen atom has been oxidized or quaternized, such as, for example, to form an N-oxide or a quaternary salt. Representative examples include, but are not limited to, tetrahydropyranyl, tetrahydrofuranyl, tetrahydrothiophenyl, piperidino, morpholino, piperazino, pyrrolidino, oxiranyl, dioxane, 1,3-dioxolanyl, 2,2-dimethyl-1,3-dioxalanyl, sulfolanyl, 2-oxazolidonyl, 2-imidazolidonyl, S,S-dioxo-thiomorpholino, and the like.

"Heterocycloamino" means a saturated monovalent cyclic group of 4 to 8 ring atoms, wherein at least one ring atom is N and optionally contains one additional ring atom selected from N or O, the remaining ring atoms being C. The term includes groups such as pyrrolidino, piperidino, morpholino, piperazino and the like.

"Optionally substituted alkyl, alkenyl, alkynyl, alkoxy or cycloalkyl" means an alkyl, alkenyl, alkynyl, alkoxy or cycloalkyl group, as defined herein, which is optionally substituted independently with one or two substituents selected from alkyl, phenyl, benzyl, haloalkyl, heteroalkyl, halo, cyano, heterocyclyl, acyl, -OR (where R is hydrogen or alkyl), -NRR' (where R and R' are independently selected from hydrogen, acyl, or alkyl which is optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), -NHCOR (where R is alkyl which is optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), $-NRS(O)_nR'$ (where R is hydrogen or alkyl, n is an integer from 0 to 2; and R' is hydrogen, alkyl or heteroalkyl, and is optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), $-NRS(O)_nNR'R''$ (where R is hydrogen or alkyl, n is an integer from 0 to 2; and R' and

R" are independently hydrogen, alkyl or heteroalkyl and are optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), -$S(O)_nR$ (where n is an integer from 0 to 2; and R is hydrogen, alkyl or heteroalkyl and is optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), -$S(O)_nNRR'$ (where n is an integer from 0 to 2; and R and R' are independently hydrogen, alkyl or heteroalkyl and are optionally substituted with hydroxy, alkoxy, cyano, halo or heterocyclyl), -COOR, -(alkylene)COOR (where R is hydrogen or alkyl), -CONR'R" or -(alkylene)CONR'R" (where R' and R" are independently hydrogen or alkyl, or together form a heterocyclyl ring with the nitrogen atom to which they are attached).

"Optionally substituted aryl, heteroaryl or heterocyclyl" means an aryl, heteroaryl or heterocyclyl ring as defined above, which is optionally substituted independently with one or two substituients selected from alkyl, phenyl, benzyl, haloalkyl, heteroalkyl, halo, cyano, acyl, -OR (where R is hydrogen or alkyl), -NRR' (where R and R' are independently selected from hydrogen, alkyl or acyl), -NHCOR (where R is alkyl), -$NRS(O)_nR'$ (where R is hydrogen or alkyl, n is an integer from 0 to 2 and R' is hydrogen, alkyl or heteroalkyl), -$NRS(O)_nNR'R"$ (where R is hydrogen or alkyl, n is an integer from 0 to 2 and R' and R" are independently hydrogen, alkyl or heteroalkyl), -$S(O)_nR$ (where n is an integer from 0 to 2 and R is hydrogen, alkyl or heteroalkyl), -$S(O)_nNRR'$ (where n is an integer from 0 to 2 and R and R' are independently hydrogen, alkyl or heteroalkyl), -COOR, - (alkylene)COOR (where R is hydrogen or alkyl), -CONR'R" or - (alkylene)CONR'R" (where R' and R" are independently hydrogen or alkyl).

"Heteroalkyl" means an alkyl radical as defined above, carrying one, two or three substituents selected from -NR^aR^b, -OR^c wherein R^a, R^b and R^c are independently of each other hydrogen, alkyl or acyl, or R^a and R^b together form heterocycloamino group. Representative examples include, but are not limited to, hydroxymethyl, acetoxymethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 2-methoxyethyl, 2-aminoethyl, 2-dimethylaminoethyl, 2-acetylaminoethyl, 3-[pyrrolidin-1-yl]ethyl and the like.

"Heteroalkenyl" means an alkenyl radical as defined above, carrying one or two substituents selected from -NR^aR^b, -OR^c or -$S(O)_nR^d$ wherein R^a, R^b and R^c are independently of each other hydrogen or alkyl, and R^d is alkyl or –NRR' (where R

and R' are independently of each other hydrogen or alkyl. Representative examples include, but are not limited to, 3-hydroxy-1-propenyl, 3-aminoprop-l-enyl, 2-aminosulfonylethenyl, 2methylsulfonylethenyl, and the like.

"Heteroalkynyl" means an alkynyl radical as defined above, carrying one or two substituents selected $-NR^aR^b$, $-OR^c$, $-S(O)_nR^d$ or $-S(O)_nNRR'$ (where R and R' are independently of each other hydrogen or alkyl) wherein R^a, R^b and R^c are independently of each other hydrogen or alkyl, and R^d is alkyl and n is an integer from zero to two. Representative examples include, but are not limited to, 3-hydroxy-l-propynyl, 3dimethylaminoprop-1-ynyl and the like.

"Heteroalkoxy" means a radical -OR where R is heteroalkyl group as defined above, e.g., 2-hydroxyethoxy, 3-hydroxypropoxy, 2,3-dihydroxypropoxy, 2-aminoethoxy, and the like.

"Heteroalkylamino" means a radical $-NR^aR^b$ where R^a is hydrogen or alkyl, and R^b is a heteroalkyl group as defined above, e.g., 2-hydroxyethylamino, 3-dimethylaminopropylamino, and the like.

"Optionally substituted heterocyclylalkyl" means a radical $-R^aR^b$ where R^a is an alkylene group, and R^b is an optionally substituted heterocyclyl group as defined above e.g., 2-(morpholin-4-yl)ethyl, 3(piperidin-1-yl)-2-methylpropyl, and the like.

"Optionally substituted heterocyclylalkenyl" means a radical $-R^aR^b$ where R^a is an alkenylene group and R^b is an optionally substituted heterocyclyl group as defined above e.g., 3-(morpholin-4-yl)prop-1-enyll, 3-(piperidin-1-yl)prop-1-enyl, 3-(4-methylpiperazin-1-yl)prop-1-enyl, and the like.

"Optionally substituted heterocyclylalkynyl" means a radical $-R^aR^b$ where R^a is an alkynyl group and R^b is an optionally substituted heterocyclyl group as defined above e.g., 3-(morpholin-4-yl)prop-1-ynyl, 3-(piperidin-1-yl)prop-1-ynyl, and the like.

"Optionally substituted heterocyclylalkoxy" means a radical -OR where R is an optionally substituted heterocyclylalkyl group as defined above, e.g., 2-(morpholin-4-yl)-ethoxy, 3-(piperazin-1-yl)propoxy, 2-[2-oxopyrrolidin-1-yl]ethoxy, and the like.

"Optionally substituted heterocyclylalkylamino" means a radical $-NR^aR^b$ where R^a is hydrogen or alkyl and R^b is an optionally substituted heterocyclylalkyl

group as defined above, e.g., 2-(pyrrolidin-2-yl)ethylamino, 3-(piperidin-1-yl)propylamino, and the like.

"Optionally substituted heteroaralkyloxy means a radical -O-R^a where R^a is a heteroaralkyl radical e.g. 2-(pyridin-3-yl)ethoxy, 2-[3(2H)-pyridazon-1-yl]ethoxy and the like.

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "aryl group optionally mono- or di-substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the aryl group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

"Amino protecting group" refers to those organic groups intended to protect nitrogen atoms against undesirable reactions during synthetic procedures e.g., benzyl, benzyloxycarbonyl (CBZ), tert-butoxycarbonyl (Boc), trifluoroacetyl, and the like.

Throughout the specification, groups and substituents thereof may be chosen by one skilled in the field to provide stable moieties and compounds. It is also understood that the chemical groups, as described herein, can be substituted or unsubstituted, branched or unbranched, as appropriate and desired.

All stereoisomers of the compounds provided herein are contemplated, either in admixture or in pure or substantially pure form. The definition of compounds provided herein embraces all the possible stereoisomers and their mixtures. It embraces the racemic forms and the isolated optical isomers having the specified activity. The racemic forms can be resolved by physical methods, such as, for example, fractional crystallization, separation or crystallization of diastereomeric derivatives or separation by chiral column chromatography. The individual optical isomers can be obtained from the racemates from the conventional methods, such as, for example, salt formation with an optically active acid followed by crystallization.

The compounds provided herein may also have prodrug forms. Any compound that will be converted *in vivo* to provide the bioactive agent is a prodrug. Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see, *e.g.*:

a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol.42, p. 309-396, edited by K. Widder, et al. (Acamedic Press, 1985);

b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, p. 113-191 (1991); and

c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992), each of which is incorporated herein by reference.


As used herein, treatment means any manner in which one or more of the symptoms of a disease or disorder are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compounds and compositions herein, such as use for treating p38 kinase mediated diseases or disorders, or diseases or disorders in which p38 kinase activity, including p38α and p38β kinase activity, is implicated.

As used herein, amelioration of the symptoms of a particular disorder by administration of a particular compound or pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

As used herein, IC_{50} refers to an amount, concentration or dosage of a particular test compound that achieves a 50% inhibition of a maximal response, such as modulation of p38a kinase activity, in an assay that measures such response.

B. Compounds

The compounds provided herein for use in the compositions and methods are active in assays that measure p38 kinase activity, including, but not limited to, p38α and p38β kinase activity. In one embodiment, the compounds provided herein have formulae I:



I

or a pharmaceutically acceptable derivative thereof, where:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

C is a 5-membered heteroaryl ring containing one or two heteroatoms in the ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$, where R^{80} and R^{81} are independently hydrogen, alkyl, cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl;

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n-$ (where n is an integer from 0 to 2); and R^9 is halo, cyano, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(u) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) -OC(O)-alkylene-CO_2H or -OC(O)-NR'R" (where R' and R" are independently hydrogen or alkyl);

(bb) heteroarylalkenylene or heteroarylalkynylene;

(cc) hydrogen;

(dd) halo;

(ee) pseudohalo;

(ff) hydroxy;

(gg) optionally substituted alkoxy;

(hh) $C(L)R^{40}$, where L is O, S or NR^{55}; R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{56}$, halo pseudohalo, OR^{55}, SR^{55}, $NR^{57}R^{58}$ or $SiR^{52}R^{53}R^{54}$; where R^{52}, R^{53} and R^{54} are selected as in (i) or (ii) as follows (i) R^{52}, R^{53} and R^{54} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or (ii) any two of R^{52}, R^{53} and R^{54} together form alkylene, alkenylene, alkynylene, heteroalkylene; and the other is selected as in (i); R^{55} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{56} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{64}R^{65}$; where R^{64} and R^{65} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{66} or $NR^{62}R^{63}$, or R^{64} and R^{65} together form alkylene, alkenylene, alkynylene, heteroalkylene, where R^{66} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{57} and R^{58} are selected as in (i) or (ii) as follows (i) R^{57} and R^{58} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55}, $NR^{67}R^{68}$ or $C(L)R^{69}$, where R^{67} and R^{68} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl, or together form alkylene, alkenylene,

alkynylene, heteroalkylene; and R^{69} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$, where R^{70} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl; or (ii) R^{57} and R^{58} together form alkylene, alkenylene, alkynylene, heteroalkylene; R^{62} and R^{63} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, or R^{62} and R^{63} together form alkylene, alkenylene, alkynylene, heteroalkylene; and

(ii) optionally substituted alkyl;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(1) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;


(c) alkyl; and

(d) alkoxy.

In one embodiment, C is a 5-membered heteroaryl ring containing one or two heteroatoms in the ring. In another embodiment, C is selected from pyrazole, imidazole, pyrrole, thiazole, isothiazole, oxazole, isoxazole, furan and thiophene rings. In another embodiment, C is a pyrazole or imidazole ring. In another embodiment, C is an imidazole ring. In another embodiment, C is a pyrazole ring.

1. Pyrazole-based compounds


In one embodiment, C is a pyrazole ring and the compounds provided herein have formulae II:



II

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein. In this embodiment, the hydrogen of the ring NH group may be replaced by one of the substituents shown in the structure (*i.e.*, -C(O)-A(R^3)(R^4), -R^2, or -B(D)(R^6)(R^5)).


In another embodiment, the compounds provided herein have formulae III:



III

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein.


In another embodiment, the compounds provided herein have formulae IV:



IV

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein.


In another embodiment, the compounds provided herein have formulae V:



V

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein.

In another embodiment, the compounds provided herein have formulae Va:




Va

or a pharmaceutically acceptable derivative thereof, wherein A, D, R^3, R^4, and R^6 are as defined elsewhere herein.

2. Imidazole-based compounds


In another embodiment, the compounds are imidazole-based compounds of formulae VI:



VI

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein. In this embodiment, the hydrogen of the ring NH group may be replaced by one of the substituents shown in the structure structure (*i.e.*, -C(O)-A(R^3)(R^4), -R^2, or -B(D)(R^6)(R^5).


In another embodiment, the compounds provided herein have formulae VII:



VII

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein.

In another embodiment, the compounds provided herein have formulae VIII:



VIII

or a pharmaceutically acceptable derivative thereof, where the variables are as defined elsewhere herein.

3. Other embodiments

In other embodiments, the compounds for use in the compositions and methods provided herein have the above formulae, or pharmaceutically acceptable derivatives thereof, where R^1 is hydrogen. In another embodiment, R^2 is hydrogen or lower alkyl. In another embodiment, R^2 is hydrogen.

In another embodiment, G is OR^{83} or $NR^{80}R^{81}$. In another embodiment, R^{83} is alkyl or cycloalkyl. In another embodiment, R^{83} is alkyl. In another embodiment, R^{83} is ethyl. In another embodiment, R^{80} and R^{81} are each independently hydrogen, alkyl or cycloalkyl. In another embodiment, R^{80} and R^{81} are each independently hydrogen or cycloalkyl. In another embodiment, R^{80} and R^{81} are each independently hydrogen or cyclohexyl. In another embodiment, G is NH_2 or NH(cyclohexyl).

In another embodiment, G is aryl, heteroaryl, cycloalkyl, heterocyclyl or heterocyclyl optionally fused to phenyl, and is substituted with R^3 and R^4, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom. In another embodiment, G is phenyl, cyclohexyl, cyclopentyl or benzyl, and is substituted with R^3 and R^4. In another embodiment, G is phenyl and is substituted with R^3 and R^4.

In another embodiment, A is an aryl ring. In another embodiment, A is a phenyl ring.

In another embodiment, B is an aryl ring. In another embodiment, B is a phenyl ring.

In another embodiment, D is $-C(O)NR^{80}R^{81}$. In another embodiment, R^{80} and R^{81} are each independently hydrogen, cycloalkyl or alkoxy. In another embodiment, R^{80} is hydrogen. In another embodiment, R^{81} is cycloalkyl or alkoxy. In another embodiment, R^{81} is C_{3-6}cycloalkyl or C_{1-6}alkoxy. In another embodiment, R^{81} is cyclopropyl or methoxy.

In another embodiment, D is optionally substituted heteroaryl. In another embodiment, D is optionally substituted triazolyl. In another embodiment, D is 1,2,4-triazol-3-yl.

In another embodiment, R^3 is hydrogen, optionally substituted heterocyclyl, optionally substituted alkyl, $C(L)R^{40}$, halo, pseudohalo or OR^{41}; where L is O, S or NR^{55}; R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{56}$, halo pseudohalo, OR^{55}, SR^{55}, $NR^{57}R^{58}$ or $SiR^{52}R^{53}R^{54}$; R^{41} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{59}$, $NR^{60}R^{61}$ or $SiR^{52}R^{53}R^{54}$; where R^{52}, R^{53} and R^{54} are selected as in (i) or (ii) as follows (i) R^{52}, R^{53} and R^{54} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or (ii) any two of R^{52}, R^{53} and R^{54} together form alkylene, alkenylene, alkynylene, heteroalkylene; and the other is selected as in (i); R^{55} is

hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{56} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{64}R^{65}$; where R^{64} and R^{65} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{66} or $NR^{62}R^{63}$, or R^{64} and R^{65} together form alkylene, alkenylene, alkynylene, heteroalkylene, where R^{66} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{57} and R^{58} are selected as in (i) or (ii) as follows (i) R^{57} and R^{58} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55}, $NR^{67}R^{68}$ or $C(L)R^{69}$, where R^{67} and R^{68} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl, or together form alkylene, alkenylene, alkynylene, heteroalkylene; and R^{69} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$, where R^{70} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl; or (ii) R^{57} and R^{58} together form alkylene, alkenylene, alkynylene, heteroalkylene; R^{59} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$; R^{60} and R^{61} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl or $C(L)R^{71}$, where R^{71} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or R^{60} and R^{61} together form alkylene, alkenylene, alkynylene, heteroalkylene; R^{62} and R^{63} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, or R^{62} and R^{63} together form alkylene, alkenylene, alkynylene, heteroalkylene.

In another embodiment, R^3 is hydrogen, optionally substituted heterocyclyl, optionally substituted alkyl, $C(L)R^{40}$, halo or OR^{41}. In another embodiment, R^3 is hydrogen, optionally substituted heterocyclyl, optionally substituted alkyl, $C(L)R^{40}$, iodo, chloro or OR^{41}. In another embodiment, R^3 is hydrogen, optionally substituted dioxolanyl, optionally substituted methyl, $C(L)R^{40}$, iodo, chloro or OR^{41}. In another embodiment, R^3 is hydrogen, 2-dioxolanyl, optionally substituted methyl, $C(O)R^{40}$, iodo, chloro or OR^{41}. In another embodiment, R^3 is

hydrogen, 2-dioxolanyl, optionally substituted methyl, CHO, iodo, chloro or OR^{41}. In another embodiment, R^3 is hydrogen.

In another embodiment, R^3 is optionally substituted methyl. In another embodiment, R^3 is methyl which is optionally substituted with heterocyclyl, hydroxy, aralkylamino or heterocyclylalkylamino. In another embodiment, R^3 is *N*-morpholinylmethyl, hydroxymethyl, *N*-(2-(3-chlorophenyl)-1-ethyl)aminomethyl, *N*-(2-morpholinyl-1-ethyl)aminomethyl or 4-piperizinylmethyl.

In another embodiment, A is O. In another embodiment, R^{40} is hydrogen, optionally substituted alkyl or cycloalkyl. In another embodiment, R^{40} is hydrogen or alkyl. In another embodiment, R^{40} is hydrogen.

In another embodiment, R^{41} is hydrogen or optionally substituted alkyl. In another embodiment, R^{41} is hydrogen, or alkyl optionally substituted with heterocyclyl, aryl, dialkylamino, halo or hydroxy. In another embodiment, R^{41} is hydrogen, or C_{1-3}alkyl optionally substituted with heterocyclyl, phenyl, dialkylamino, halo or hydroxy. In another embodiment, R^{41} is hydrogen, 2-(N-morpholinyl)eth-1-yl, benzyl, 2-(*N,N*-di-(2-hydroxy-1-ethyl)amino)-1-ethyl, 2-bromo-1-ethyl, 2,2-dioxolan-4-ylmethyl, 2-(4-methylpiperazin-1-yl)-1-ethyl or 2,3-dihydroxy-1-propyl. In another embodiment, R^{41} is (*S*)-2,3-dihydroxy-1-propyl.

In another embodiment, R^4 is hydrogen. In another embodiment, R^5 is alkyl. In another embodiment R^5 is methyl. In another embodiment, R^6 is hydrogen.

In another embodiment, the compounds for use in the compositions and methods provide herein have the above formulae, including formulae I-VIII, or a pharmaceutically acceptable derivative thereof, where:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$ (where R^{80} and R^{81} are independently hydrogen, alkyl, cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl);

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n$ (where n is an integer from 0 to 2); and R^9 is cyano, optionally substituted heteroaryl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16},

R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$- where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) $-OC(O)$-alkylene-CO_2H or -OC(O)-NR'R" (where R' and R" are independently hydrogen or alkyl); and

(bb) heteroarylalkenylene or heteroarylalkynylene;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(l) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy;

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

In another embodiment, the compounds are those wherein:

R^1 is hydrogen or acyl;

R^2 is hydrogen or alkyl;

A is an aryl or heteroaryl ring.

In another embodiment, the compounds are those wherein:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

R^3 is selected from the group consisting of:

(a) amino;

(b) acylamino;

(c) optionally substituted heterocycle;

(d) heteroaryl optionally substituted with a substituent selected from halo, alkyl or alkoxy;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n-$ (where n is an integer from 0 to 2); and R^9 is cyano, heteroaryl, -COOH, $-COR^{10}$, $-COOR^{11}$ $-CONR^{12}R^{13}$, $-SO_2R^{14}$,$-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$ where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(t) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$ where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -0- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26,}$ where R^{25} and R25 independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy;

R^5 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(l) optionally substituted heterocyclylalkyl; and

(m) optionally substituted heterocyclylalkoxy;

R^6 is selected from a group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy.

In another embodiment the compounds are those where R^3 is:

(a) optionally substituted heterocyclyl;

(b) aryl or heteroaryl both optionally substituted with a substituent selected from halo, alkyl, amino, alkoxy, carboxy, lower alkoxy carbonyl, SO_2R' (where R' is alkyl) or $-O_2NHR'R''$ (where R' and R" are independently hydrogen or alkyl);

(c) heteroalkyl;

(d) heteroalkenyl;

(e) heteroalkylamino;

(f) heteraloxy

(g) optionally substituted heterocyclylalkyl or heterocyclyloxy;

(h) optionally substituted heterocyclylalkenyl;

(i) optionally substituted heterocyclylalkynyl;

(j) optionally substituted heterocyclylalkoxy;

(k) optionally substituted heterocyclylalkylamino;

(l) optionally substituted heterocyclylalkylcarbonyl:

(s) -Y-(alkylene)-R^9 where Y is a single bond, -O or -NH- and R^9 is optionally substituted heteroaryl, -$CONR^{12}R^{13}$, SO_2R^{14}, -$SO_2NR^{15}R^{16}$ -$NHSO_2R^{17}$ or -$NHSO_2NR^{18}R^{19}$ where R^{12}, R^{13}, R^{14}, R^{15}, R^{16} R^{17}, R^{18} and R^{19} are independently of each other heteroalkyl; hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(m) arylaminoalkylene or heteroarylaminoalkylene;

or

(n) Z-alkylene-$NR^{30}R^{31}$ where Z is -NH-, -N(alkyl)- or -0-, and R^{30} and R^{31} are independently of each other, hydrogen, alkyl or heteroalkyl.

In still another embodiment, the compounds are those where R^1 and R^2 are hydrogen and B is phenyl. In an additional embodiment, the compounds are those wherein R^4 is hydrogen and R^5 is halo or alkyl. In another embodiment, the compounds are those wherein R^5 is chloro, fluoro or methyl and R^6 is hydrogen, chloro, fluoro, methyl or methoxy. In another embodiment, the compounds are those wherein R^3 is optionally substituted heteroaryl.

In yet another embodiment, the compounds are those wherein R^3 is pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxidopyridin-2-yl, N-oxidopyridin-3-yl, Noxidopyridin-4-yl or pyridon-2-yl, all optionally substituted. In a further embodiment, the compounds are those wherein R^3 is at the 3-position. In still another embodiment, the compounds are those wherein R^5 is 4-F or 2-Me, and R^6 is hydrogen. In another embodiment, the compounds are those wherein R^3 is optionally substituted phenyl.

In a further embodiment, the compounds are those wherein R^3 is 3-sulfamoylphenyl, 3-methylsulfonylphenyl, 3-carboxyphenyl or 3-ethoxycarbonylphenyl. In yet another embodiment, the compounds are those wherein R^5 is 4-F and R^6 is hydrogen.

In another embodiment, the compounds are those wherein R^3 is:

(a) heteroalkyl;

(b) heteroalkoxy;

(c) heteroalkylamino;

(d) optionally substituted heterocyclylalkyl;

(e) optionally substituted heterocyclylalkoxy;

(f) optionally substituted heterocyclylalkylamino;

(g) Y-(alkylene)-R^9 where Y is a single bond, -O- or –NH- and R^9 is optionally substituted heteroaryl, -$CONR^{12}R^{13}$, -SO_2R^{14}, -$SO_2NR^{15}R^{16}$ -$NHSO_2R^{17}$ or -$NHSO_2NR^{18}R^{19}$ where R_{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are independently of each other hydrogen, alkyl or heteroalkyl; or

(h) Z-alkylene-$NR^{30}R^{31}$ where Z is -NH-, -N(alkyl)- or -O-, and R^{30} and R^{31} are independently of each other, hydrogen, alkyl or heteroalkyl.

In a further embodiment, the compounds are those wherein R^3 is heteroalkyl. In another embodiment, the compounds are those wherein R^3 is at the 3-position and is selected from the group consisting of 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino, hydroxymethyl, 1,2-dihydroxyethyl, 3-hydroxy-3-methyl-l-butyl or 3-hydroxybutyl. In yet another embodiment, the compounds are those wherein R^5 is 2-F and R^6 is 4-F.

In still another embodiment, the compounds are those wherein R^5 is 2-Me and R^6 is hydrogen. In an additional embodiment, the compounds are those wherein R^9 is heteroalkoxy or heteroalkylamino. In yet another embodiment, the compounds are those wherein R^3 is at the 3-position and is selected from the group consisting of 3-dimethylaminopropoxy, 2dimethylaminoethoxy, 2-hydroxyethoxy, 2,3-dihydroxypropoxy, 2-dimethylaminoethylamino and 3-dimethylaminopropylamino.

In yet another embodiment, the compounds are those wherein R^3 is optionally substituted heterocyclylalkyl, optionally substituted heterocyclylalkoxy

or optionally substituted heterocyclylalkylamino. In still another embodiment, the compounds are those wherein R^3 is at the 3-position and is selected from the group consisting of 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 2-(2-oxo-pyrrolidin-1-yl)ethoxy, 3(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)ethylamino, 4-hydroxypiperidinylmethyl, 2-(S,S-dioxothiamorpholin-4-yl)ethyl, 3-(S,S-dioxo-thiamorpholin-4-yl)propyl and N-methylpiperazinylmethyl.

In an additional embodiment, the compounds are those wherein R^3 is -Y-(alkylene)-R^9 where Y is a single bond, -O- or -NH- and R^9 is optionally substituted heteroaryl, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$ where R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are independently of each other hydrogen, alkyl or heteroalkyl. In a further embodiment, the compounds are those wherein Y is a single bond and R^9 is $-SO_2R^{14}$ or $-SO_2NR^{15}R^{16}$.

In an additional embodiment, the compounds are those wherein R^3 is 5-methylsulfonylethyl or sulfamoylethyl.

Also provided herein is a compound selected from the group consisting of 5-amino-l-(4-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole, 5-amino-l-(2,4-difluorophenyl)-4-[3-(3-morpholin-4ylpropyl)benzoyl]pyrazole, 5-amino-4-(3-aminobenzoyl)-1-(4-fluorophenyl)pyrazole, 5-amino-l-(4-fluorophenyl)-4- [3-(3-morpholin-4-ylpropyl)benzoyl] pyrazole, 5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl] -1-(4fluorophenyl)pyrazole, 5-amino-4-(3-acetylaminobenzoyl)-1-phenylpyrazole, 5-amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)pyrazole, 5-amino-l-(4-fluorophenyl)-4- [3-(3-morpholin-4-ylpropylamino) benzoyl] pyrazole, 5-amino-4-[3-(2-aminosulfonylethyl)benzoyl]-1-(4-fluorophenyl)pyrazole and 5-amino-l-(4-fluorophenyl)-4-(3-pyridin-3-ylbenzoyl)pyrazole.

Also provided herein is a compound selected from the group consisting of: 5-amino-l-(2-methylphenyl)-4-[3-pyridin-3-yl)benzoyl]pyrazole, 5-amino-l-(2-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]pyrazole, 5-amino-4-[3-(2,3-dihydroxypropoxy)benzoyl]-1-(4-fluorophenyl) pyrazole, 5-amino-4-[3-(1,2-dihydroxyethyl)benzoyl]-1-(4-fluorophenyl) pyrazole, 5-amino-l-(4-fluorophenyl)-4-[3-(sulfamoylbenzoyl] pyrazole, 5-amino-l-(4-fluorophenyl)-4-[3-(3-hydroxy-3-

methylbutyl) benzoyl]pyrazole, 5-amino-1-(4-fluorophenyl)-4-[3-(2-(1-hydroxycyclopentyl)ethyl) benzoyl]pyrazole, 5-amino-4-[3-(2-methylsulfonylethyl)benzoyl]-1-(4-fluorophenyl) pyrazole, and 5-amino-l-(2,4-difluorophenyl)-4-[3-(2-hydroxyethylsulfonyl) benzoyl]pyrazole.

Further provided herein is a compound selected from 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone; 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; and 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

Also provided herein is a compound selected from 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone; 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-{5-

amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; or 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

Also provided is a compound selected from:

3-{5-Amino-4-[3-(2-dimethylamino-ethylcarbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(5-chloro-thiophene-2-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-hydrazinocarbonyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclohexanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclopentanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-phenylacetyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; and

3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide.

Also provided is a compound selected from:

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester;

3-(5-Amino-4-cyclopentanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-hydrazinocarbonyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid benzylamide;

3-(5-Amino-4-cyclohexanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; and

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylamide.

Also provided is a compound selected from:

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester;

5-Amino-3-[(3-chloro-benzylcarbamoyl)-methoxy]-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester;

3-[5-Amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-(2-hydroxy-ethoxy)-1H-pyrazole-4-carboxylic acid ethyl ester;

4-[5-Amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

3-(5-Amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; and

3-[5-Amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide.

Also provided is a compound selected from:

3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide;

3-(5-Amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide;

3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid;

3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid;

3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

{5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone;

3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclohexanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclopentanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-phenylacetyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(2-dimethylamino-ethylcarbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-methylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-cyclopentylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(morpholine-4-carbonyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(cyclopropylmethyl-carbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-ethoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-benzoyl-3-(2-benzyloxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

4-[5-amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

3-[5-amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide, trifluoroacetate salt;

3-(5-amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide;


5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide; and

5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester.

C. Preparation of the Compounds


Also provided herein is a process for preparing a compound of formula (I), which process involves the steps of:

(i) reacting a 2-keto-3-phenylaminoacrylonitrile of formula 1:



1


with a hydrazine of formula 2:



2


where R^3, R^4 R^5 and R^6 are as defined herein to provide a compound of formula (I) where R^1 is hydrogen; or

(ii) reacting a 2-keto-3-phenylaminoacrylonitrile of formula 3:



3

where Z is either hydroxy, nitro or halo group and R^4 is as defined herein with a hydrazine of formula 2 to provide a compound of formula 4:



4

followed by conversion of the Z group to the desired R^3 group to provide a compound of formula (I) where R^1 is hydrogen;


(iii) optionally modifying any of the R^1, R^3, R', R^5 or R^6 groups;

(iv) optionally converting the compound of formula (I) prepared in steps (i), (ii) or (iii) above to the corresponding acid addition salt by treatment with an acid;

(v) optionally converting the compound of formula (I) prepared in steps (i), (ii) or (iii) above, to the corresponding free base by treatment with a base; and


(vi) optionally separating a mixture of stereoisomers of a compound of formula (I) prepared in steps (i) - (v) above, to give a single stereoisomer.

Also provided herein is a process for preparing a compound of formula (I), which process involves reacting a compound of formula 5:



5

where R^5 and R^6 are as as defined herein and L is a leaving group under organometallic displacement reaction conditions, including, but not limited to, halo, pseudohalo, aryloxy, perfluoroaryloxy, N-alkoxyamino, including N-methoxyamino, with an organometallic reagent of formula:



where R^3 and R^4 are as defined herein and M is a metallic moiety, including, but not limited to, an alkali metal, an alkaline earth metal, and a transitional metal, such as Li, K and Mg, to provide a compound of formula (I) where R^1 is hydrogen;

(ii) optionally modifying any of the R^1, R^3, R^4, R^5 or R^6 groups;

(iii) optionally converting the compound of formula (I) prepared in steps (i) or (ii) above, to the corresponding acid addition salt by treatment with an acid;

(iv) optionally converting the compound of formula (I) prepared in steps (i) or (ii) above, to the corresponding free base by treatment with a base; and optionally separating a mixture of stereoisomers of a compound of formula (I) prepared in steps (i) or (iv) above, to give a single stereoisomer.

The compounds disclosed herein are merely exemplary, and one skilled in the art can be readily prepare compounds in the same manner as that disclosed herein using well known methods of chemical synthesis, including methods similar to those exemplified herein.

Compounds provided herein may generally be prepared according to the following schemes and the knowledge of one skilled in the art. Additional


synthetic methods are described, for example, in U.S. Patent Nos. 6,376,527; 6,316,466 and 6,444,696, and International Patent Application Publication No. WO 99/57101, each of which is incorporated herein by reference.

In addition to the documents incorporated by reference we disclose the following non-limiting examples of methods useful for the production of compounds provided herein (see, Schemes 1-8, *infra*).

Amines attached to aryl or heteroaryl ring systems are useful as intermediates in the preparation of the compounds provided herein. There are many methods of preparing such intermediates known to one skilled in the art of organic chemistry. Several methods of preparing amines useful in the preparation of the compounds provided herein are illustrated in schemes 1-7.

Substituted aniline of type (II) useful herein can be prepared from commercially available 3-amino-4-methylbenzoic acid as depicted in Scheme 1, using methods similar to those disclosed in International Patent Application Publication No. WO 02/40486. Aniline is protected by a Boc group. This is followed by condensation with methylamine using the coupling agent EDC and HOBt. The Boc group is then removed by HCl in dioxane to give the desired substituted aniline of type (II) as a hydrochloride salt.


Scheme 1



(II)

Substituted anilines of type (III) useful herein can be prepared from commercially available 3-amino-4-methylbenzoic acid as depicted in Scheme 2. Condensation with cyclopropylamine using the coupling agent EDC affords an aniline of type (III).


Scheme 2



(III)

Substituted anilines of type (IV) useful herein can be prepared from commercially available 4-methyl-3-nitrobenzoic acid as depicted in Scheme 3, using methods similar to those disclosed in International Patent Application Publication No. WO 03/033482. Condensation of the acid with t-butoxy-carbonylhydrazide using the coupling agent HBTU and HOBt affords the protected acyl hydrazide. Deprotection with TFA, followed by condensation with triethyl orthoformate yields the oxadiazolyl-substituted nitrotoluene. Hydrogenation of the nitro group gives the desired aniline of type (IV).


Scheme 3



(IV)

Substituted anilines of type (V) useful herein can be prepared from commercially available 4-methyl-3-nitrobenzamide as depicted in Scheme 4, using methods described by Han *et al.* [*J. Med. Chem.*, 41, 2019-2028 (1998)]. The indicated aryl carboxamide was condensed with N,N-dimethylformamide diethyl acetal. This was followed by reaction with hydrazine in acetic acid to form a triazolyl-substituted nitrotoluene. Hydrogenation of the nitro group gives the desired aniline of type (V).


Scheme 4



(V)

Substituted anilines of type (VI) useful herein can be prepared from commercially available methyl 4-iodobenzoate as depicted in Scheme 5. Nitration of an aromatic precursor followed by reduction of the nitro group yields the aniline. Palladium-catalyzed coupling with ethynyltrimethylsilane, followed by desilylation and saponification gives the desired ethynyl-substituted aminobenzoic acid. Coupling with methoxyamine using the coupling agent EDC affords the desired aniline (VI). See, *e.g.*, *Eur. J. Org. Chem.*, 4607 (2001).

Scheme 5




(VI)

Alternatively, a substituted aniline of type (VI) useful herein can be prepared from 4-amino-3-nitrobenzoic acid as depicted in Scheme 6. Iodide substitution of the aryldiazonium salt, followed by esterification with methanol gives methyl 4-iodo-3-nitrobenzoate. The nitro group can be reduced by $SnCl_2$ to give the desired aniline. Palladium catalyzed coupling with ethynyltrimethyl-silane, followed by desilylation and saponification yields the ethynyl-substituted

aminobenzoic acid. Coupling with methoxyamine using the coupling agent EDC affords the aniline (VI). See, *e.g.*, *Eur. J. Org. Chem.*, 4607 (2001).


Scheme 6



(VI)

As depicted in Scheme 7, substituted anilines of type (VII) useful herein can be prepared from intermediate methyl 4-iodo-3-nitrobenzoate, which can be synthesized as shown in Scheme 6. Palladium catalyzed coupling with vinyltributyltin followed by carbene addition to the resulting styrene double bond gives the cyclopropyl substituted methyl nitrobenzoate. Reduction of the nitro group followed by Boc protection and saponification gives the protected 3-amino-4-cyclopropylbenzoic acid. Coupling with an alkoxyamine using the coupling agent EDC affords the desired aniline (VII). See, *e.g.*, International Patent Application Publication Nos. WO 02/092087 and WO 02/40486.

Scheme 7



Hydrazines attached to aryl or heteroaryl ring systems are useful as intermediates herein. There are many methods of preparing such intermediates


known to one skilled in the art of organic chemistry. One method of preparing some of the hydrazines useful herein is illustrated in Scheme 8.

Aryl hydrazines of type (VIII) useful herein can be prepared from 3-amino-*N*-methoxy-4-methylbenzamide hydrochloride, which itself can be prepared according the methods disclosed in International Patent Application Publication No. WO 02/40486. Through the formation of the aryldiazonium salt and its subsequent reduction by $SnCl_2$, the desired hydrazine of type (VIII) is obtained.


Also as depicted in Scheme 8, aryl hydrazines of type (IX) useful herein can be prepared from 3-amino-*N*-cyclopropyl-4-methylbenzamide, which itself can be prepared according to the methods depicted in Scheme 2. Through the formation of the aryldiazonium salt and its subsequent reduction by $SnCl_2$, the desired hydrazine of type (IX) is obtained.

Similarly, other hydrazines can also be prepared from amines such as those described above in Schemes 1-7.

Scheme 8




(VIII)



(IX)


Also, as depicted in Scheme 9, acrylonitriles of type (X) useful herein can be prepared from an aryl ester and acetonitrile. Acetonitrile can be treated with lithium diisopropylamide in THF at -78 °C, followed by the addition of the aryl ester to give the corresponding aryloylacetonitrile. This intermediate is then reacted with N,N'-diphenylformamidine in a solvent such as toluene at relfux to give the desired corresponding acrylonitrile of type (X).

Scheme 9




Also, as depicted in Scheme 10, aminopyrazoles of type (XI) useful herein can be prepared from an acrylonitrile of type (X), which itself can be prepared according to the methods depicted in Scheme 9, and hydrazines such as those of type (VIII) and (IX), which themselves can be prepared according to the methods depicted in Scheme 8. The acrylonitrile and the hydrazine are heated to 60 to 100 °C in a solvent such as DMF or ethanol to give the desired aminopyrazole of type (XI).

Scheme 10




Also, as depicted in Scheme 11, aminopyrazoles of type (XIII) useful herein can be prepared from an acrylonitrile of type (XII), which itself can be prepared by treatment of an acrylonitrile of type (X), prepared according to the methods depicted in Scheme 9, with sodium hydride and carbon disulfide followed by treatment with iodomethane, and hydrazines such as those of type (VIII) and (IX), which themselves can be prepared according to the methods depicted in Scheme 8. The acrylonitrile and the hydrazine are heated to 60 to 100 °C in a solvent such as DMF or ethanol to give the desired aminopyrazole of type (XIII).

Scheme 11




Also, as depicted in Scheme 12, aminopyrazoles of type (XIV) useful herein can be prepared from an acrylonitrile of type (XII), which itself can be prepared according to the methods depicted in Scheme 11, and hydrazines such as those of type (VIII) and (IX), which themselves can be prepared according to the methods depicted in Scheme 8. Treatement of this intermediate (XII) with an alcohol alkoxide prior to with heating at 60 to 100 °C with the hydrazine in a solvent such as DMF or ethanol gives the desired aminopyrazole of type (XIII).

Scheme 12



Also, as depicted in Scheme 13, aminoimidazoles of type (XV) useful herein can be prepared from substituted anilines of type (III), which themselves can be prepared according to the methods depicted in Scheme 2. The aniline is heated in triethyl orthoformate. After removal of the solvent *in vacuo*, the product was reacted with aminomalononitrile *p*-toluenesulfonate and sodium acetate in acetic acid to give the aminocyanoimidazole intermediate. Reaction of this intermediate with a Grignard reagent gives the desired aminoimidazole of type (XV).

Scheme 13



Compounds provided herein can be prepared from hydrazines attached to aryl or heteroaryl ring systems using methods disclosed in U.S. Patents Nos. 6,316,466, 6,376,527 and 6,444,696.

Additional synthetic methods useful in the synthesis of the compounds provided herein are disclosed in the following, the disclosures of which are incorporated herein by reference in their entirety:

1) J. Heterocyclic Chem. 17, 631 (1980)
2) Tetrahedron 55(48), 13703 (1999)
3) European Patent No. EP 0 713 876
4) Chemische Berichte 126(10), 2317 (1993)
5) Journal of Organic Chem. 58(24), 6620 (1993)
6) Tetrahedron Letters 35, 3239 (1973)
7) Journal of Chemical Research, Synopses 1, 2 (1997)
8) Boletin de la Sociedad Quimica del Peru 53(3), 150 (1987)
9) Journal of the Chemical Society, Chemical Communications 2, 35 (1973)
10) Comptes Rendus des Seances de l'Academie des Sciences, Series C: Sciences Chimiques 274(20), 1703 (1972)

Also provided herein are compounds prepared according to a process disclosed herein.

D. Formulation of pharmaceutical compositions

Also provided herein is a pharmaceutical composition comprising a compound provided herein. The composition can be used, for example, as a medicament. The composition can contain, for example, a pharmaceutically acceptable excipient or carrier. A composition or medicament provided herein can be used for the treatment, prevention or amelioration of one or more symptoms of p38 kinase mediated diseases or disorders, including inflammatory diseases.

Thus, provided herein are pharmaceutical compositions capable of treating p38-kinase-associated conditions, including TNF-α, IL-1, and/or IL-8 mediated conditions, as described above. The compositions may contain other therapeutic agents, as described herein, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (*e.g.*, excipients, binders, preservatives, stabilizers, flavors, *etc.*) according to techniques such as those well known in the art of pharmaceutical formulation.

The compounds provided herein may be administered by any means suitable for the condition to be treated, which may depend on the need for site-specific treatment or quantity of drug to be delivered. Topical administration is generally useful for skin-related diseases, and systemic treatment is generally used for cancerous or pre-cancerous conditions, although other modes of delivery are contemplated. For example, the compounds may be delivered orally, such as in the form of tablets, capsules, granules, powders, or liquid formulations including syrups; topically, such as in the form of solutions, suspensions, gels or ointments; sublingually; bucally; parenterally, such as by subcutaneous, intravenous, intramuscular or intrasternal injection or infusion techniques (*e.g.*, as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; rectally such as in the form of suppositories; or liposomally. Dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents may be administered. The compounds may be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved with suitable pharmaceutical compositions or, particularly in the case of extended release, with

devices such as subcutaneous implants or osmotic pumps. Exemplary compositions for topical administration include a topical carrier such as PLASTIBASE® (mineral oil gelled with polyethylene).

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The inventive compounds may also be orally delivered by, sublingual and/or buccal administration, *e.g.*, with molded, compressed, or freeze-dried tablets. Exemplary compositions may include fast-dissolving diluents such as mannitol, lactose, sucrose, and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (AVICEL®) or polyethylene glycols (PEG); an excipient to aid mucosal adhesion such as hydroxypropyl cellulose (HPC), hydroxypropyl methyl cellulose (HPMC), sodium carboxymethyl cellulose (SCMC), and/or maleic anhydride copolymer (*e.g.*, GANTREZ®); and agents to control release such as polyacrylic copolymer (*e.g.*, CARBOPOL 934®). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal aerosol or inhalation administration include solutions which may contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance absorption and/or bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or suspensions which may contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Exemplary compositions for rectal administration include suppositories which may contain, for example, suitable non-irritating excipients, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures but liquefy and/or dissolve in the rectal cavity to release the drug.

The effective amount of a compound provided herein may be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for a mammal of from about 0.05 to 100 mg/kg of body weight of active compound per day, which may be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors, including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition. Subjects for treatment include animals, generally mammalian species such as humans, and domestic animals such as dogs, cats, horses, and the like. Thus, when the term "patient" is used herein, this term is intended to include all subjects, in particular mammalian species,, including humans, that are affected by mediation of p38 enzyme levels.

Also provided in one embodiment is a process for the manufacture of medicaments which process involves bringing a compound provided herein together with a pharmaceutically acceptable excipient and bringing the mixture into a galenical administration form.

E. Methods of use of the compounds and compositions

In a further embodiment, the compounds provided herein can be used in the treatment, prevention, or amelioration of one or more symptoms of inflammatory diseases. A compound provided herein can be used, in another embodiment, for the manufacture of a medicament for the treatment orprophylaxis of inflammatory diseases.

The compounds provided herein are selective inhibitors of p38 kinase activity, and in particular, isoforms p38α and p38β. Accordingly, compounds provided herein are useful for treating conditions associated with p38 kinase activity. Such conditions

include diseases in which cytokine levels are modulated as a consequence of intracellular signaling *via* p38, and in particular, diseases that are associated with an overproduction of cytokines IL-l, IL-4, IL-8, and TNF-α. Provided herein are methods of treating a disease by administering a compound provided herein that inhibits p38 kinase activity. Also provided herein are methods for inhibiting or delaying the onset of a disease or disorder by administering a compound provided herein. Methods provided herein can be used to achieve a full or partial reduction of the symptoms of a disease or disease state, and/or to alleviate, ameliorate, or lessen, the disease or disorder and/or its symptoms. When reference is made herein to inhibition of "p-38α/βkinase," this means that either p38α and/or p38β kinase are inhibited. Thus, reference to an IC_{50} value for inhibiting p-38α/β kinase means that the compound has such effectiveness for inhibiting at least one of, or both of, p38α and p38β kinases.

In view of their activity as inhibitors of p38α/β kinase, compounds provided herein are useful in treating p-38 associated conditions including, but not limited to, inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, angiogenic disorders, infectious diseases, neurodegenerative diseases, and viral diseases.

More particularly, the specific conditions or diseases that may be treated with the inventive compounds include, without limitation, pancreatitis (acute or chronic), asthma, allergies, adult respiratory distress syndrome, chronic obstructive pulmonary disease, glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosis, scleroderma, chronic thyroiditis, Grave's disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, graft vs. host disease, inflammatory reaction induced by endotoxin, tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, pancreatic β-cell disease; diseases characterized by massive neutrophil infiltration; rheumatoid spondylitis, gouty arthritis and other arthritic conditions, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption disease, allograft rejections, fever and myalgias

due to infection, cachexia secondary to infection, meloid formation, scar tissue formation, ulcerative colitis, pyresis, influenza, osteoporosis, osteoarthritis and multiple myeloma-related bone disorder, acute myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, multiple myeloma, sepsis, septic shock, and Shigellosis; Alzheimer's disease, Parkinson's disease, cerebral ischemias or neurodegenerative disease caused by traumatic injury; angiogenic disorders including solid tumors, ocular neovasculization, and infantile haemangiomas; viral diseases including acute hepatitis infection (including hepatitis A, hepatitis B and hepatitis C), HIV infection and CMV retinitis, AIDS, SARS, ARC or malignancy, and herpes; stroke, myocardial ischemia, ischemia in stroke heart attacks, organ hyposia, vascular hyperplasia, cardiac and renal reperfusion injury, thrombosis, cardiac hypertrophy, thrombin induced platelet aggregation, endotoxemia and/or toxic shock syndrome, and conditions associated with prostaglandin endoperoxidase synthase-2.

In addition, p38 inhibitors provided herein inhibit the expression of inducible pro-inflammatory proteins such as prostaglandin endoperoxide synthase-2 (PGHS-2), also referred to as cyclooxygenase-2 (COX-2). Accordingly, additional p38-associated conditions include edema, analgesia, fever and pain, such as neuromuscular pain, headache, pain caused by cancer, dental pain and arthritis pain. The inventive compounds also may be used to treat veterinary viral infections, such as lentivirus infections, including, but not limited to equine infectious anemia virus; or retro virus infections, including feline immunodeficiency virus, bovine immunodeficiency virus, and canine immunodeficiency virus. When the terms "p38-associated condition" or "p38-associated disease or disorder" are used herein, each is intended to encompass all of the conditions identified above as if repeated at length, as well as any other condition that is affected by p38 kinase activity.

Thus, provided herein are methods for treating such conditions, involving administering to a subject in need thereof an effective amount of at least one compound provided herein or a pharmaceutically acceptable derivative thereof. The methods of treating p38 kinase-associated conditions may involve administering compounds provided herein alone or in combination with each other and/or other suitable therapeutic agents useful in treating such conditions. Exemplary of such other therapeutic agents include corticosteroids, rolipram, calphostin, CSAIDs, 4-

substituted imidazo[1,2-A]quinoxalines as disclosed in U.S. Patent No. 4,200,750 and in S. Ceccarelli *et al.* (1998) *European Journal of Medicinal Chemistry 33*:943-955; interleukin-10, glucocorticoids, salicylates, nitric oxide, and other immunosuppressants; nuclear translocation inhibitors, such as deoxyspergualin (DSG); non-steroidal antiinflammatory drugs (NSAIDs) such as ibuprofen, celecoxib and rofecoxib; steroids such as prednisone or dexamethasone; antiviral agents such as abacavir; antiproliferative agents such as methotrexate, leflunomide, FK506 (tacrolimus, Prograf); cytotoxic drugs such as azathioprine and cyclophosphamide; TNF-α inhibitors such as tenidap, anti-TNF antibodies or soluble TNF receptor, and rapamycin (sirolimus or Rapamune) or derivatives thereof.

The above other therapeutic agents, when employed in combination with the compounds provided herein, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art. In the methods provided herein, such other therapeutic agent(s) may be administered prior to, simultaneously with, or following the administration of the inventive compounds.

The following Examples illustrate embodiments herein, and are not intended to limit the scope of the claims. Abbreviations employed in the Examples are defined below. Compounds of the Examples are identified by the example and step in which they are prepared (for example, "1A" denotes the title compound of step A of Example 1), or by the example only where the compound is the title compound of the example (for example, "2" denotes the title compound of Example 2).

Abbreviations

Ph = phenyl

Bz = benzyl

t-Bu = tertiary butyl

Me = methyl

Et = ethyl

Pr = propyl

Iso-P or i-Pr = isopropyl

MeOH = methanol

EtOH = ethanol

EtOAc = ethyl acetate

Boc = tert-butyloxycarbonyl

CBZ = carbobenzyloxy or carbobenzoxy or benzyloxycarbonyl

DCM or CH_2Cl_2 = dichloromethane

DCE = 1,2-dichloroethane

DMF = dimethyl formamide

DMSO = dimethyl sulfoxide

TFA = trifluoroacetic acid

THF = tetrahydrofuran

HATU = *O*-(7-Azabenzotriazol-1-yl-*N,N,N',N'*-tetramethyluronim hexafluorophosphate

KOH = potassium hydroxide

K_2CO_3 = potassium carbonate

$POCl_3$ = phosphorous oxychloride

KOtBu = potassium t-butoxide

EDC or EDCI = 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

DIPEA = diisopropylethylamine

HOBt = 1-hydroxybenzotriazole hydrate

m-CPBA = m-chloroperbenzoic acid

NaH = sodium hydride

NaOH = sodium hydroxide

$Na_2S_2O_3$ = sodium thiosulfate

Na_2SO_4 = sodium sulfate

Pd = palladium

Pd/C = palladium on carbon

min = minute(s)

L = liter

mL = milliliter

μL= microliter

g = gram(s)

mg = milligram(s)

mol = moles

mmol = millimole(s)

meq = milliequivalent

RT or rt = room temperature


ret. t. or t_R = HPLC retention time (minutes)

sat or sat'd = saturated

General Methods. Mass spectral data were obtained on a Thermo Finnigan LCQ Duo Ion Trap mass spectrometer. In the Examples: "HPLC (6 minute gradient)" refers to Keystone C18 Beta Basic column, 0.4 mL/min flow rate, 6 minute linear gradient elution (start solvent %B = 0; final solvent %B = 100), solvent A: acetonitrile + 0.025% TFA; solvent B = H_2O + 0.025% TFA ."HPLC (4 minute gradient)" refers to Keystone C18 Beta Basic column, 0.5 mL/min flow rate, 4 minute linear gradient elution (start solvent %B = 0; final solvent %B = 100), solvent A: acetonitrile + 0.025% TFA; solvent B = H_2O + 0.025% TFA .

EXAMPLE 1

Preparation of 3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide



A. 3-Hydrazino-N-methoxy-4-methyl-benzamide

To a stirred solution of 3-amino-N-methoxy-4-methyl-benzamide **1** (102 mg, 0.56 mmol, preparation: International Patent Application Publication No. WO 02/40486 A2, pg. 66) in water (5 ml) at 0 °C was added conc. HCl (5 mL) followed by the addition of sodium nitrite (43 mg, 0.62 mmol). The reaction mixture was stirred at


0 °C for 40 min then a solution of tin(II)chloride (241 mg, 1.27 mmol) in conc. HCl (1 mL) was added and the mixture was stirred for 1 hr then allowed to stand at –20 °C for 20 hr before it was warmed to RT and concentrated to a white solid. The solid was triturated with ethanol, the solids were filtered, and the filtrate concentrated to provide 3-hydrazino-N-methoxy-4-methyl-benzamide **2** as white solid (486 mg) as a mixture with tin salts and ethanol which was used without further purification. HPLC (6 minute gradient) t_R 0.78 min; MS m/z 195.9 $[M+H]^+$.

B. 3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide

To a stirred solution 3-hydrazino-N-methoxy-4-methyl-benzamide **2** (116 mg, estimated 0.14 mmol) in EtOH (10 ml) was added 2-(3-iodo-benzoyl)-3-phenylamino-acrylonitrile **3** (59 mg, 0.14 mmol, preparation: International Patent Application Publication No. WO 02/57101 A1, pg. 84) and mixture was heated (bath T = 65-70 °C) for 4 hr, when additional 3-hydrazino-N-methoxy-4-methyl-benzamide **2** (80 mg, 0.11 mmol) was added and the mixture was heated at the same temperature for 27 hr. It was cooled to room temperature, concentrated and redissolved in EtOAc before it was washed with water and brine, dried over Na_2SO_4, concentrated to give a crude semisolid. The mixture was then purified by flash chromatography, eluting with 1:1 EtOAc/hexanes to remove impurities then 100% EtOAc give 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide **4** as an off-white solid (38 mg, 0.08 mmol, 55%). HPLC (6 minute gradient) t_R 3.49 min; MS m/z 476.96 $[M+H]^+$; ^1H NMR (CD_3OD, 300 MHz) δ 8.10 (s, 1 H), 7.95 (d, J = 8.0, 1 H), 7.88 (d, J = 8.0, 1 H), 7.82 (m, 2 H), 7.78 (s, 1 H), 7.58 (d, J = 8.0, 1 H), 7.33 (t, J = 7.8, 1 H), 5.02 (s. 3 H), 2.33 (s, 3 H) ppm; ^{13}C NMR (DMSO-d_6, 125MHz) δ 189.3, 154.3, 143.5, 143.3, 142.0, 138.3, 137.4, 133.5, 132.9, 132.0, 130.4, 128.8, 128.4, 104.8, 95.4, 64.8, 18.2 ppm.

EXAMPLE 2


Preparation of 3-(5-Amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide



To a stirred solution of 3-amino-N-methoxy-4-methyl-benzamide **1** (104 mg, 0.58 mmol) in water (2 ml) at 0 °C was added conc. HCl (2 mL) followed by the addition of sodium nitrite (44 mg, 0.63 mmol). The reaction mixture was stirred at 0 °C for 40 minutes then a solution of tin(II)chloride (245 mg, 1.30 mmol) in conc. HCl (1 mL) was added and the mixture was stirred for 40 minutes then allowed to stand at –20 °C for 20 hours before it was warmed to RT and concentrated to a white solid. The solid was triturated with ethanol, the solids were removed, and 2-benzoyl-3-phenylamino-acrylonitrile **5** (144 mg, 0.58 mmol, preparation: Grothaus, J. Am. Chem. Soc. 58, 1334 (1936)) was added and the mixture heated (bath T = 65-70 °C) for 16 hr. The mixture was cooled to RT, concentrated and purified by flash chromatography, eluting with 1:1 EtOAc/hexanes to remove impurities then 100% EtOAc to give 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide **6** as an off-white solid (41 mg, 0.12 mmol, 28%). HPLC (4 minute gradient) t_R 1.93 min; MS m/z 351.1 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 11.88 (s 1 H, NH), 7.803 (m, 4 H), 7.56 (m, 4 H), 7.01 (s, 2 H, NH_2), 3.32 (s, 3 H), 2.162 (s, 3 H) ppm; ^{13}C NMR (DMSO-d_6, 125 MHz) δ 187.6, 151.9, 141.2, 139.7, 139.6, 135.7, 131.4, 131.2, 130.9, 128.5, 128.1, 127.8, 126.4, 102.6, 63.2, 17.2 ppm.

EXAMPLE 3

Preparation of 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid



A. 3-Hydrazino-4-methyl-benzoic acid hydrochloride

To a stirred solution of 3-amino-4-methylbenzoic acid **1** (5.64 g, 31.2 mmol, 1.0 eq) in 100 mL of dioxane and 100 mL of water at 0 °C was added 100 mL of conc. HCl followed by the portionwise addition of sodium nitrite (2.82 g, 40.9 mmol, 1.1 eq) as a solid at a rate to control gas evolution and foaming over 45 minutes. A clear brown solution resulted. Anhydrous tin(II)chloride (15.62 g, 83.7 mmol, 2.25 eq) was dissolved in conc. HCl (25 mL) and added dropwise over 25 mL at 0 °C. After 1 hour, the precipitate was filtertered and washed with dioxanes then dried under vacuum to provide 3-hydrazino-4-methyl-benzoic acid hydrochloride **2** as a tan solid (4.98 g, 66%): HPLC (4 minute gradient) t_R 0.97 min; MS m/z 167 $[M+H]^+$; ^1H (300 MHz, DMSO-d_6) δ 10.03 (s, 1 H, COOH), 7.89 (s, 1 H), 7.51 (s, 1 H), 7.27 (d, J = 8.0, 1 H), 3.38 (s, 3 H, NHNH$_2$), 2.23 (s, 3 H) ppm.


B. 3-(5-Amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid

To a stirred solution of 3-hydrazino-4-methyl-benzoic acid hydrochloride **2** (242 mg, 1.19 mmol, 1.0 eq) in 25 mL of ethanol was added 2-benzoyl-3-phenylamino-acrylonitrile **4** (296 mg, 1.19 mmol, 1.0 eq, preparation: Grothasu, Davis, *J. Am. Chem. Soc.*, 58, 1334 (1936)) and triethylamine (161 µL, 1.19 mmol, 1.0 eq) and the mixture was heated to 65 °C. All solids dissolved when temperature reached 65 °C. After three hours, LC-MS indicates consumption of the hydrazine. The solids were filtered to provide 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid **5** (95 mg, 25%) as a beige solid: HPLC (4 minute gradient) t_R 2.10 min;

MS m/z 322 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 7.99 (d, J = 7.6, 1 H), 7.81 (s, 2 H), 7.78 (s, 1 H), 7.57 (m, 4 H), 7.02 (s, 2 H, $N\underline{H}_2$),.2.18 (s, 3 H) ppm; ^{13}C NMR (DMSO-d_6, 75 MHz) δ187.6, 166.3, 152.0, 141.3, 141.2, 139.6, 135.9, 131.6, 131.2, 130.1, 129.8, 128.6, 127.8, 102.6, 17.4 ppm.

EXAMPLE 4

Preparation of 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide




5 6

To a stirred solution of 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid **5** (Example 3, 700 mg, 2.18 mmol, 1.0 eq) in 30 mL of DMF was added EDCI (855 mg 4.35 mmol, 2.0 eq), HOBt (589 mg, 4.35 mmol, 2.0 eq), and diisopropylethylamine (1.59 mL, 8.71 mmol, 4.0 eq) and the solution was stirred for 15 minutes at room temperature when cyclopropylamine (302 μL, 4.35 mmol, 2.0 eq) was added and the reation stirred for 1 hour. The mixture was diluted with EtOAc (300 mL) and washed with water (2 x 25 mL) and brine (25 mL), dried (Na_2SO_4) and concentrated. The product was purified by flash chromatography on silica gel eluted with 8/2 EtOAc/MeOH to provide the product as a brown oil. The product was further puriried by trituration with 1/1/1 EtOAc/hexanes/CH_2Cl_2 and dried under vacuum to provide 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **6** (387 mg, 50%) as a white powder: HPLC (4 minute gradient) t_R 2.11 min; MS m/z 361 $[M+H]^+$; ^1H NMR (CD_3OD, 300 MHz) δ 7.92 (d, J = 7.6, 1 H), 7.81 (m, 4 H), 7.54 (m, 4 H), 2.85 (m, 1 H), 2.22 (s, 3 H), 0.80 (d, J =5.7, 2 H), 0.63

(s, 2 H) ppm; ^{13}C NMR (CD_3OD, 75MHz) δ 191.2, 170.1, 153.8, 143.3, 142.0, 141.1, 136.9, 134.8, 132.9, 132.7, 130.1, 129.7, 129.2, 128.1, 104.8, 24.1, 17.7, 6.6 ppm.

EXAMPLE 5


Preparation of 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid




To a stirred solution of 3-hydrazino-4-methyl-benzoic acid hydrochloride **1** (Example 3A, 314 mg, 1.54 mmol, 1.0 eq) in 50 mL of ethanol and 5 mL of methanol was added 2-(3-iodo-benzoyl)-3-phenylamino-acrylonitrile **2** (579 mg, 1.54 mmol, preparation: International Patent Application Publication No. WO 02/57101 A1, pg. 84). The mixture was heated to 75 °C for 18 h. The precipitated solids were collected on a fritted filter and were washed with ethanol to provide 3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid **3** (153 mg, 22%) as a white solid. MS m/z 448 $[M+H]^+$; ^1H NMR (300 MHz, DMSO-d_6) δ2.18 (s, 3H), 7.05 (bs, 2H), 7.34 (dd, $J_1 = J_2 = 7.7$ Hz), 7.58 (d, $J = 8.1$ Hz, 1H), 7.78 (s, 1H), 7.81 (m, 2H), 7.94 (m, 1H), 7.98 (m, 1H), 8.03 (m, 1H) ppm.

EXAMPLE 6

Preparation of 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide





A. 3-(N'-tert-Butoxycarbonyl-hydrazino)-4-methyl-benzoic acid

3-Hydrazino-4-methyl-benzoic acid hydrochloride **1** (Example 3A, 13 g, 64.5 mmol) was dissolved into dioxane (200 mL) and H_2O (100 mL). Aqueous NaOH (5.16 g NaOH in 100 mL H_2O, 2×64.5 mmol) was added followed by the addition of $(Boc)_2O$ (15.5 g, 1.1×64.5 mmol) immediately. The reaction mixture was stirred at room temperature for 2 hrs. Concentrated on rotavapor. Then H_2O and CH_2Cl_2 (some MeOH) were added. With stirring strong H^+ resin was added to neutralize the mixture to pH < 2. Filtered and the resin was washed with CH_2Cl_2 and MeOH. The aqueous layer was washed with CH_2Cl_2 (added some MeOH) for two times. The combined organic layer was dried over Na_2SO_4 (some EtOAc was added). Filtration and concentration gave 3-(N'-tert-butoxycarbonyl-hydrazino)-4-methyl-benzoic acid **4** (16 g, 94%) as a white solid. ^1H NMR (300 MHz, $CDCl_3$) δ 1.48 (s, 9H), 2.27 (s, 3H), 5.76 (s, 1H), 7.14 (d, J = 7.7 Hz, 1 H), 7.56 (d, J = 7.7 Hz, 1H), 7.61 (s, 1H) ppm.

B. N'-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-hydrazinecarboxylic acid tert-butyl ester

3-(N'-tert-Butoxycarbonyl-hydrazino)-4-methyl-benzoic acid **4** (14 g, 52.6 mmol) was dissolved in DMF (250 mL). EDCI (20 g, 105.2 mmol) and HOBt (16 g, 105.2 mmol) were added. The mixture was stirred at room temperature for 30 min. Cyclopropylamine (7.4 mL, 105.2 mmol) was added, followed by DIPEA (37 mL, 4×52.6 mmol). The reaction mixture was stirred at room temperature for 18 h. After concentration of the reaction mixture *in vacuo*, H_2O was added. The mixture was then extracted with CH_2Cl_2 three times. The organic layer was washed with aqueous NaCl solution. Dried over Na_2SO_4, filtration and concentration gave a white solid.

The crude product was dissolved in CH_2Cl_2/MeOH, and then purified by silica gel column chromatography (CH_2Cl_2/EtOAc, gradient 2/1 to1/1) to give the desire product. The product was further purified by recrystallization from EtOAc/CH_2Cl_2; washing of the collected solids with EtOAc gave N'-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-hydrazinecarboxylic acid tert-butyl ester **5** (11 g, 69%). ^{1}H NMR (300 MHz, $CDCl_3$) δ0.58 (m, 2H), 0.84 (m, 2H), 1.48 (s, 9H), 2.23 (s, 3H), 2.87 (m, 1H), 5.69 (bs, 1H), 6.17 (bs, 1H), 6.39 (brs, 1H), 7.70 (m, 2H), 7.32 (s, 1H) ppm.

C. N-Cyclopropyl-3-hydrazino-4-methyl-benzamide, trifluoroacetic acid salt


N'-(5-Cyclopropylcarbamoyl-2-methyl-phenyl)-hydrazinecarboxylic acid tert-butyl ester **5** was dissolved in CH_2Cl_2/TFA (2/1) with 2% H_2O and the mixture was stirred at room temperature for 3 hrs. Concentration *in vacuo* gave a syrup. CH_2Cl_2 and toluene was added, and concentration *in vacuo* again gave N-cyclopropyl-3-hydrazino-4-methyl-benzamide, trifluoroacetic acid salt **6** as an off-white solid. (Yield: 100%). MS m/z 206 $[M+H]^+$; ^{1}H NMR (300 MHz, D_2O) δ0.68 (m, 2H), 0.88 (m, 2H), 2.31 (s, 3H), 2.79 (m, 1H), 7.31 (s, 1H), 7.36 (bs, 2H) ppm.

D. 3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

To a stirred solution of N-cyclopropyl-3-hydrazino-4-methyl-benzamide, trifluoroacetic acid salt **6** (648 mg, 1.74 mmol, 1.0 eq) in 2 mL of ethanol was added 2-(3-iodo-benzoyl)-3-phenylamino-acrylonitrile **2** (550 mg, 1.74 mmol, preparation: International Patent Application Publication No. WO 02/57101 A1, pg. 84) and DIEA (0.50 mL, 2.9 mmol). The mixture was heated to 160 °C for 40 min using microwave. The mixture was concentrated *in vacuo*. The crude product was purified by silica gel column chromatography (EtOAc/hex, gradient 1/1 to 2/1) to give 3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **7** (589 mg, 70%) as a white solid. MS m/z 497 $[M+H]^+$; ^{1}H NMR (300 MHz, $CDCl_3$) δ0.60 (m, 2H), 0.86 (m, 2H), 2.24 (s, 3H), 2.87 (m, 1H), 5.81 (bs, 2H), 6.35 (bs, 1H), 7.25 (dd, $J_1 = J_2 = 7.8$ Hz, 1H), 7.45 (d, $J = 8.1$ Hz, 1H), 7.69 (d, $J = 1.7$ Hz, 1H), 7.80 (m, 3H), 7.89 (m, 1H), 8.00 (s, 1H) ppm.

EXAMPLE 7

Preparation of {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone



A. N-Dimethylaminomethylene-4-methyl-3-nitro-benzamide

4-Methyl-3-nitro-benzamide **1** (10 g, 56 mmol) was suspended in 80 mL of N,N-dimethylformamide dimethylacetal and was then heated to 95 °C for 2 h. The red solution was allowed to cool to room temperature with stirring. After another 2 h, the resulting red precipitate was collected on a fritted filter, and was washed three times with Et_2O to give N-dimethylaminomethylene-4-methyl-3-nitro-benzamide **2** as a red solid (8.7 g, 66%). HPLC (4 minute gradient) t_R 1.76 min; MS m/z 236.0 $[M+H]^+$.

B. 3-(4-Methyl-3-nitro-phenyl)-4H-[1,2,4]triazole

To a solution of N-dimethylaminomethylene-4-methyl-3-nitro-benzamide **2** (8.6 g, 37 mmol) in 250 mL of acetic acid was added dropwise anhydrous hydrazine (4.7 mL, 180 mL). The now light orange solution was heated to 95 °C for 1.5 h before allowing to cool and to stir at room temperature for 18 h. Acetic acid was removed *in vacuo*, and the residue was partitioned between H_2O and EtOAc. The organic layer was washed twice with saturated $NaHCO_3$ solution, then dried over

$MgSO_4$. After filtration and concentration *in vacuo*, the residue was triturated with warm EtOAc, and the resulting off-white solid was collected on a fritted filter to give 3-(4-methyl-3-nitro-phenyl)-4H-[1,2,4]triazole **3** (5.8 g, 77%). HPLC (4 minute gradient) t_R 1.82 min; MS m/z 205.1 $[M+H]^+$; ^1H NMR (300 MHz, DMSO-d_6) δ 8.57 (bs, 1 H), 8.46 (s, 1 H), 8.12 (d, J = 7.9, 1 H), 7.56 (d, J = 7.9, 1 H), 2.5 (s, 3 H) ppm; ^{13}C NMR (DMSO-d_6, 125 MHz) δ 149.1, 133.5, 130.1, 121.3, 19.5 ppm.

C. 2-Methyl-5-(4H-[1,2,4]triazol-3-yl)-phenylamine

3-(4-Methyl-3-nitro-phenyl)-4H-[1,2,4]triazole **3** (5.75 g, 28.2 mmol) was suspended in 220 mL of ethanol with 2.35 mL (*ca.* 28.2 mmol) of concentrated aqueous HCl. Under nitrogen, 900 mg of 10% palladium on activated carbon (dry) was added carefully. Hydrogen gas was bubbled through the reaction mixture *via* a balloon attached to a syringe needle for 5 minutes. The reaction mixture was then stirred under an atmosphere of hydrogen gas maintained by a balloon at room temperature for 5 h. The catalyst was removed by filtration through a short pad a Celite. The filtrate was concentrated *in vacuo*, and the residue was neutralized with saturated $NaHCO_3$ solution. The mixture was extracted with EtOAc six times and the combined organic layers were dried over $MgSO_4$. After filtration and concentration *in vacuo*, the residue was recrystallized in EtOAc to give 2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenylamine **4** (4.5 g, 92%) as an off-white solid. HPLC (4 minute gradient) t_R 0.79 min; MS m/z 175.2 $[M+H]^+$; ^1H NMR (300 MHz, MeOH-d_3) δ 8.10 (bs, 1 H), 7.54 (s, 1 H), 7.29 (m, 2 H), 7.15 (d, J = 7.6, 2 H), 2.23 (s, 3 H) ppm.

D. [2-Methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-hydrazine


2-Methyl-5-(4H-[1,2,4]triazol-3-yl)-phenylamine **4** (200 mg, 1.15 mmol, preparation) in dioxane (5 ml) and water (5 ml) at 0 °C was added conc. HCl (10 mL) followed by the addition of sodium nitrite (87 mg, 1.26 mmol). The reaction mixture was stirred at 0°C for 40 min then a solution of tin(II)chloride (481 mg, 2.59 mmol) in conc. HCl (1 mL) was added dropwise. The mixture was stirred for 2 hr at 0 °C which resulted in a white solid precipitate. The solid was collected by filtration and identified as [2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-hydrazine **5** (261mg) which was used without further purification. HPLC (4 minute gradient) t_R 0.71 min; MS m/z 190.1 $[M+H]^+$.

E. {5-Amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl-phenyl-methanone

To a stirred solution of [2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-hydrazine **5** (261 mg, estimated 1.15 mmol) in EtOH (25 ml) was added 2-benzoyl-3-phenylamino-acrylonitrile **6** (285 mg, 1.15 mmol, preparation: Grothaus, J. Am. Chem. Soc. 58, 1334 (1936)) and the mixture was heated to 65-70°C for 12-16 hours. It was cooled to room temperature, concentrated to give crude product. The mixture was then purified by flash chromatography, eluting with 1:1 EtOAc/Hexanes to give {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone **7** as a white solid (21 mg, 0.09 mmol, 8%). HPLC (4 minute gradient) t_R 1.89 min; MS m/z 345.2 $[M+H]^+$; ^1H NMR (DMSO-d_6, 500 MHz) δ 8.40 (d, 1 H), 8.24 (bs, 1 H), 8.14 (s, 1 H), 8.12 (d, 2 H), 7.34 (bs, 2 H), 7.92-7.85 (m, 4 H), 2.49 (s, 3 H) ppm; ^{13}C NMR (DMSO-d_6, 500 MHz) δ152.9, 142.2, 140.5, 137.1, 132.7 132.2, 129.4, 128.8, 127.6, 126.0, 103.6, 18.1 ppm.

EXAMPLE 8

Preparation of 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



A. 3-[1,3]Dioxolan-2-yl-benzoic acid methyl ester

A mixture of 3-formyl-benzoic acid methyl ester **1** (6.09 g, 37.2 mmol), ethylene glycol (2.28 mL, 40.9 mmol) and *p*-toluenesulfonic acid monohydrate (0.78 g, 4.09 mmol) was refluxed overnight with Dean-Stark apparatus. TLC plate showed that all the starting material has disappeared. Poured the reaction mixture into a mixture of cooled aqueous $NaHCO_3$ and EtOAc. The organic layer was separated and dried over Na_2SO_4. Filtered and concentrated to give the crude product that was purified by silica gel chromatography (eluent: 8/1 hexane/ethyl acetate). The desired 3-[1,3]dioxolan-2-yl-benzoic acid methyl ester **2** was obtained as a colorless oil (6.41 g, 83%). ^{1}H NMR (300 MHz, $CDCl_3$) δ 3.92 (s, 3H), 4.01 (m, 2 H), 4.13 (m, 2 H), 5.85 (s, 1 H), 7.46 (dd, $J_1 = J_2 = 7.7$ Hz, 1H), 7.68 (dt, $J_1 = 7.7$ Hz, $J_2 = 1.2$ Hz, 1 H), 8.05 (dt, $J_1 = 7.7$ Hz, $J_2 = 1.5$ Hz, 1 H), 8.16 (dd, $J_1 = J_2 = 1.5$ Hz, 1H) ppm.

B. 3-(3-[1,3]Dioxolan-2-yl-phenyl)-3-oxo-propionitrile

To a mixture of acetonitrile (1.90 mL, 36.4 mmol) in THF (60 mL) was added LDA (1.8 M in THF, 33.9 mL) at –78 °C. After stirring the mixture for 20 min at –78 °C, 3-[1,3]dioxolan-2-yl-benzoic acid methyl ester **2** (6.06 g, 29.1 mmol) in THF (20 ml) was added all at once. The mixture was stirred at –78 °C for 1.5 h and then warmed to 0 °C and stirred for 1 h at this temperature. Saturated NH_4Cl was added to quench the reaction. The mixture was extracted with EtOAc three times. Organic layers were combined and dried over Na_2SO_4. Filtration and concentration *in vacuo* gave a residue that was purified by silica gel chromatography (CH_2Cl_2, then 20/1 CH_2Cl_2/ethyl acetate). The desired 3-(3-[1,3]dioxolan-2-yl-phenyl)-3-oxo-propionitrile **3** was obtained as a white solid (5.71 g, 96%). ^{1}H NMR (300 MHz, $CDCl_3$) δ 4.12 (m, 6H), 5.86 (s, 1H), 7.55 (dd, $J_1 = J_2 = 7.7$ Hz, 1H), 7.77 (d, $J = 7.7$ Hz, 1 H), 7.92 (d, $J = 7.7$ Hz, 1 H), 8.01 (s, 1H) ppm.

C. 2-(3-[1,3]Dioxolan-2-yl-benzoyl)-3-phenylamino-acrylonitrile

The mixture of 3-(3-[1,3]dioxolan-2-yl-phenyl)-3-oxo-propionitrile **3** (3.07 g, 15.0 mmol) and *N, N'*-diphenylformamidine (4.10 g, 21 mmol) in toluene was heated to reflux for 18 h. Concentration *in vacuo* gave a residue that was purified by silica gel chromatography (hexanes/EtOAc, gradient from 3/1 to 2/1 then 1/1). The desired 2-(3-[1,3]dioxolan-2-yl-benzoyl)-3-phenylamino-acrylonitrile **4** was obtained as a yellow solid (3.88 g, 81%). ^{1}H NMR (300 MHz, $CDCl_3$) δ 4.12 (m, 4H), 5.92 (s, 1H),


7.26 (m, 4H), 7.47 (m, 3H), 7.65 (d, J = 7.7 Hz, 1H), 7.94 (dt, J_1 = 7.7 Hz, J_2 = 1.3 Hz, 1H), 8.05 (m, 2H) ppm.

D. 3-[5-Amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

A mixture of 2-(3-[1,3]dioxolan-2-yl-benzoyl)-3-phenylamino-acrylonitrile **4** (0.20 g, 0.62 mmol), N-cyclopropyl-3-hydrazino-4-methyl-benzamide, trifluoroacetic acid salt **8** (Example 6C, 0.20 g, 0.62 mmol) and DIEA (0.5 mL) in DMF (3 mL) was heated to 160 °C for 40 min using microwave. The mixture was then cooled down to room temperature and concentrated. The obtained residue was purified by silica gel chromatography (gradient from 1/4 Hexanes/EtOAc to 100% EtOAc). The desired 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **5** was obtained as an orange solid (0.26 g, 100%). MS m/z 433.2 $[M+H]^+$; ^{1}H NMR (300 MHz, $CDCl_3$) δ0.60 (m, 2H), 0.86 (m, 2H), 2.24 (s, 3H), 2.87 (m, 1H), 4.11 (m, 4H), 5.85 (m, 3H), 6.51 (bs, 1H), 7.43 (d, J = 8.0 Hz, 1 H), 7.45 (dd, $J_1 = J_2$ = 7.6 Hz, 1H), 7.70 (m, 2H), 7.82 (m, 3H), 7.95 (s, 1H), 8.00 (s, 1H) ppm.

EXAMPLE 9

Preparation of 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide




A mixture of 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **5** (Example 8, 1.14 g, 2.6 mmol) was suspended in aqueous AcOH (10 mL, 1.5 M in H_2O). Glacial AcOH was added dropwise until a clear solution was obtained. The reaction mixture was stirred at room temperature overnight. Evaporation of the solvent under reduced pressure to give a residue. Toluene and EtOAc were added and the mixture was concentrated again to give the

crude product that could be purified by silica gel chromatography (hexanes/EtOAc: 1/4). The desired 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** was obtained as a yellow foamy solid (0.91 g, 89%). MS m/z 389.1 $[M+H]^+$; ^1H NMR (300 MHz, $CDCl_3$) δ0.61 (m, 2 H), 0.85 (dd, J_1 = 6.9 Hz, J_2 = 12.4 Hz, 2 H), 2.25 (s, 3 H), 2.87 (m, 1 H), 5.95 (brs, 2 H), 6.57 (brs, 1 H), 7.44 (d, J = 8.0 Hz, 1 H), 7.67-7.78 (m, 4 H), 8.07 (s, 1 H), 8.09 (d, J = 1.3 Hz, 1 H), 8.31 (s, 1 H), 10.12 (s, 1 H) ppm.

EXAMPLE 10


Preparation of 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



To a mixture of 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** (38 mg, 0.098 mmol) in methanol (3 mL) was added $NaBH_4$ (11 mg, 0.29 mmol). The reaction mixture was stirred at room temperature for 1 h. Aqueous NaOH was added to quench the reaction and the mixture was extracted with ethyl acetate. The organic layer was separated and washed with water and saturated NaCl solution. The organic was then dried over Na_2SO_4, filtered and concentrated to give a residue that could be purified by silica gel chromatography (eluent: 1/4 hexanes/EtOAc). The desired 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **7** was obtained as a white solid (22 mg, 58%). MS m/z 391.2 $[M+H]^+$; ^1H NMR (300 MHz, $CDCl_3$) δ0.56 (m, 2H), 0.82 (m, 2H), 2.22 (s, 3H), 2.82 (m, 1H), 4.75 (s, 2H), 5.91 (brs, 2H), 6.75 (brs, 1H), 7.49 (m, 3H), 7.76 (m, 4H), 7.99 (s, 1H) ppm.

EXAMPLE 11


Preparation of 3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



To a mixture of 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** (Example 9, 1.0 eq) and 1-methyl-piperazine (1.09 eq) in equal volume of 1,2-dichloroethane and dichloromethane was added AcOH (0.96 eq) Sodium triacetoxyborohydride (1.5 eq) was then added. The reaction mixture was stirred at room temperature for 2 h. Aqueous NaOH was added to quench the reaction and the mixture was extracted with ethyl acetate. The organic layer was separated and washed with water and saturated NaCl solution. The organic layer was then dried over Na_2SO_4, filtered and concentrated to give a residue that could be purified by silica gel chromatography (eluent: 9/1 CH_2Cl_2/methanol then 9/1/0.05 CH_2Cl_2/MeOH/NH_3H_2O). Yield 67%. MS m/z 473.3 $[M+H]^+$; ^1H NMR (300 MHz, $CDCl_3$) δ0.61 (m, 2H), 0.86 (m, 2H), 2.26 (s, 3H), 2.30 (s, 3H), 2.52 (brs, 8 H), 2.90 (m, 1H), 3.60 (s, 2H), 5.87 (brs, 1H), 6.30 (brs, 2H), 7.45 (m, 3H), 7.70 (m, 2H), 7.83 (m, 3H) ppm.

EXAMPLE 12


Preparation of 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 11 except morpholine was used in place of 1-methyl-piperidine. Yield 45%. MS m/z 460.2 $[M+H]^+$; ^1H NMR (300 MHz, $CDCl_3$) δ0.61 (m, 2H), 0.87 (m, 2H), 2.26 (s, 3H), 2.25 (s, 3H), 2.49 (m, 4H), 2.90 (m, 1H), 3.59 (s, 2H), 3.72 (t, J = 4.4 Hz, 2H), 5.85 (brs, 2H), 6.35 (brs, 1H), 7.45 (m, 3H), 7.71 (m, 2H), 7.83 (m, 3H) ppm.

EXAMPLE 13

Preparation of 3-{5-Amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



A. 3-(2-Morpholin-4-yl-ethoxy)-benzoic acid ethyl ester

Potassium carbonate (6.2 g, 45 mmol) was added to a DMF solution (100 mL) of ethyl 3-hydroxybenzoate **1** (3.32 g, 20 mmol), 4-(2-chloroethyl)-morpholine hydrochloride (5.58 g, 30 mmol) and 18-crown-6 (20 mg). The mixture was stirred at 100 °C for 20 hr. Solvent was removed *in vacuo* and the residue was suspended in ethyl acetate. The organic layer was washed by saturated $NaHCO_3$ solution and then brine, and was dried over sodium sulfate. Solvent was evaporated to yield the product 3-(2-morpholin-4-yl-ethoxy)-benzoic acid ethyl ester **2** as a light yellow oil (5.3 g, 95%). HPLC (4 minute gradient) t_R 1.47 min; MS m/z 280.2 $[M+H]^+$.

B. 3-[3-(2-Morpholin-4-yl-ethoxy)-phenyl]-3-oxo-propionitrile

Lithium diisopropylamide (16.4 mL, 29.6 mmol, 1.8M solution in heptane/tetrahydrofuran/ethylbenzene) was added dropwise to the solution of acetonitrile (1.2 g, 29.6 mmol) in dry tetrahydrofuran (20 mL) at – 78 °C in nitrogen atmosphere. After stirring the reaction mixture for 30 min, a solution of 3-(2-morpholin-4-yl-ethoxy)-benzoic acid ethyl ester **2** (5.5 g, 19.7 mmol) in dry tetrahydrofuran (20 mL) was added dropwise and stirred at – 78 °C for 1 hr. Water was added and the aqueous layer was separated and acidified with dilute hydrochloric acid to pH 7. The product was extracted into ethyl acetate. The organic layer was washed with brine and then dried over sodium sulfate. The solvent was removed *in vacuo* to give 3-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-3-oxo-propionitrile **3** as a light yellow oil (4.8 g). HPLC (4 minute gradient) t_R 1.11 min; MS m/z 275.2 $[M+H]^+$.

C. 2-[3-(2-Morpholin-4-yl-ethoxy)-benzoyl]-3-phenylamino-acrylonitrile

A mixture of 3-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-3-oxo-propionitrile **3** (4.8 g, 17.5 mmol) and N,N-diphenylformamidine (1.2 g, 24.5 mmol) in dry toluene (100 mL) was heated at 110 °C for 3 hr under a nitrogen atmosphere. Solvent was removed and the oily residue was subjected to silica gel column chromatography (gradient 100% EtOAc to 100/10/1 EtOAc/MeOH/Et_3N) to yield 2-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-3-phenylamino-acrylonitrile **4** light yellow solid (3.1g, 47 %). HPLC (4 minute gradient) t_R 2.04 min; MS m/z 378.2 $[M+H]^+$.

D. 3-{5-Amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-4-methyl-benzoic acid


2-[3-(2-Morpholin-4-yl-ethoxy)-benzoyl]-3-phenylamino-acrylonitrile **4** (189 mg, 0.5 mmol) and 3-hydrazino-4-methyl-benzoic acid hydrochloride **7** (Example 3A 152 mg, 0.75 mmol) were suspended in N,N-dimethylformamide (5 mL) and heated at 160 °C using microwave for 30 min. Solvent was evaporated and the residue was subjected to column chromatography (EtOAc ~ MeOH). Product 3-{5-amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-4-methyl-benzoic acid **5** was obtained as a light yellow solid (200 mg). HPLC (4 minute gradient) t_R 1.60 min; MS m/z 451.2 $[M+H]^+$.

E. 3-{5-Amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

A mixture of 3-{5-amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-4-methyl-benzoic acid **5** (400 mg, 0.89 mmol), cyclopropylamine (0.89 mmol), EDCI (340 mg, 1.78 mmol), HOBt (272 mg, 1.78 mmol) and diisopropylethylamine (459 mg, 3.56 mmol) in dry N,N-dimethylformamide (10 mL) was stirred at room temperature for 18 h. Solvent was evaporated and the residue was suspended in EtOAc and washed by water, saturated $NaHCO_3$ solution and brine. The organic layer was dried over sodium sulfate. Product 3-{5-amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide **6** was obtained as a light yellow solid (45 mg, 10%) after purification by column chromatography (100/10/1 EtOAc/MeOH/Et_3N). HPLC (4 minute gradient) t_R 1.69 min; MS m/z 490.24 $[M+H]^+$.

EXAMPLE 14

Preparation of 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



A. 3-Benzyloxy-benzoic acid ethyl ester

K_2CO_3 (6.9 g, 50 mmol) and 18-crown-6 were added to a solution of 3-hydroxy-benzoic acid ethyl ester **1** (8.3 g, 50 mmol) in acetone (100 mL) and the mixture was stirred at room temperature for 18 h. Solid was removed by filtration. Filtrate was concentrated *in vacuo* to give 3-benzyloxy-benzoic acid ethyl ester **2** as a colorless liquid.

B. 3-(3-Benzyloxy-phenyl)-3-oxo-propionitrile

LDA (1.8 M, 100 mmol, 56 mL) was added to a solution of acetonitrile (4.1 g, 100 mmol) in THF (100 mL, dry) at -78 °C under nitrogen. The mixture was stirred at – 78 °C for 30 min. A solution of 3-benzyloxy-benzoic acid ethyl ester **2** in 50 mL anhydrous THF was then added dropwise to the reaction mixture. The mixture was stirred at – 78 °C for 1 h before water was added. The organic phase was separated. The aqueous phase was acidified by hydrochloride acid until pH ~ 2, and was extracted by EtOAc. THF and EtOAc layers were combined and washed by water, brine, dried over Na_2SO_4. Solvent was removed *in vacuo* and the solid residue was triturate with Et_2O and dried in vacuo. The desired product 3-(3-benzyloxy-phenyl)-3-oxo-propionitrile **3** was obtained as a light brown solid (11.0 g, 87%).

C. 2-(3-Benzyloxy-benzoyl)-3-dimethylamino-acrylonitrile

N,N-Dimethylformamide dimethyl acetal (10 mL) was added to the solution of 3-(3-benzyloxy-phenyl)-3-oxo-propionitrile **3** (2.5 g, 10 mmoL) in DMF (20 mL, dry) and the mixture was stirred at 100 °C for 3 hr. Solvent was removed and the residue was purified by silica gel column chromatography (EtOAc as eluent). Product 2-(3-benzyloxy-benzoyl)-3-dimethylamino-acrylonitrile **4** was obtained as a light yellow solid (2.6 g, 90 %).


D. 3-[5-Amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

2-(3-Benzyloxy-benzoyl)-3-dimethylamino-acrylonitrile **4** (147 mg, 0.5 mmol) and N-cyclopropyl-3-hydrazino-4-methyl-benzamide, trifluoroacetic acid salt **8** (Example 6C, 240 mg, 0.75 mmol) were dissolved in DMF (5 mL) and was heated at 160 °C in microwave oven for 30 min. Solvent was removed and the residue was purified by column (3:1 EtOAc/hexanes). Product 3-[5-amino-4-(3-benzyloxy-

benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** was obtained as a light yellow solid (120 mg, 52%).

EXAMPLE 15

Preparation of 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide





6 7

3-[5-Amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** (200 mg, 0.43 mmol) was dissolved in MeOH (10 mL). Catalyst 10% palladium on activated carbon (dry) was added and was stirred in an atmosphere of hydrogen for 2 h. The catalyst was removed by filtration and solvent was removed *in vacuo*. Product 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **7** was obtained as a light yellow solid (140 mg, 87%).

EXAMPLE 16

Preparation of 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide





A. 2-(4-Methyl-benzoyl)-3-phenylamino-acrylonitrile


A mixture of 4-toluoylacetonitrile **1** (4.0 g, 25 mmol) and N,N-diphenylformamidine (4.9 g, 25 mmol) in dry toluene (50 mL) was heated at 85 °C for 16 h under nitrogen. The mixture was cooled to room temperature and 170 mL of hexanes was added. A yellow precipitate was formed after the mixture was stirred at room temperature for 5 minutes. The solid was collected of a fritted flask, and was washed with hexanes to give pure 2-(4-methyl-benzoyl)-3-phenylamino-acrylonitrile **2** (4.5 g, 68%). ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.04 (d, J = 13.0, 1 H), 7.86 (d, J = 7.9, 2 H), 7.42 (t, J = 7.4, 1 H), 7.28-7.25 (m, 3 H), 7.19 (d, J = 7.6, 1 H), 2.41 (s, 3H) ppm.

B. 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

A mixture of 2-(4-methyl-benzoyl)-3-phenylamino-acrylonitrile **2** (205 mg, 0.78 mmol), N-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt **7** (Example 6C, 250 mg, 0.78 mmol) and diisopropylethylamine (0.14 mL, 0.78 mmol) in 8 mL of ethanol was heated at 65 °C in for 18 h. Solvent was removed and the residue was purified by silica gel column chromatography (EtOAc/hexanes, gradient from 1/3 to 3/1). The product can be further purified by trituration with Et_2O to give 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **3** as a white solid (64 mg, 22%). HPLC (4 minute gradient) t_R 2.26 min; MS m/z 375.2 $[M+H]^+$; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.51 (d, J = 4.0, 1 H), 7.93 (d, J = 8.0, 1 H), 8.51 (d, J = 4.0, 1 H), 7.83 (bs, 1 H), 7.82 (s, 1 H), 7.70 (d, J = 7.9, 2 H), 7.53 (d, J = 8.0, 1 H), 7.35 (d, J = 7.9, 2 H), 6.95 (bs, 2 H), 2.86 (m, 1 H), 2.40 (s, 3 H), 2.14 (s, 3 H), 0.68 (m, 2 H), 0.56 (m, 2 H); ^{13}C NMR (300 MHz, DMSO-d_6) δ 187.4, 166.0, 151.9, 141.3, 141.1, 139.2, 136.9, 135.6, 133.0, 131.2, 129.1, 128.3, 128.0, 126.5, 102.6, 23.1, 21.0, 17.2, 5.6 ppm.

EXAMPLE 17

Preparation of 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



A. 3-(5-Amino-4-cyano-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide


A mixture of 3-amino-N-cyclopropyl-4-methyl-benzamide **1** (380 mg, 2.0 mmol) in 2.0 mL of triethyl orthoformate was stirred at 120°C in microwave for 20 minutes. The solvent was removed under reduced pressure. The residue was dissolved in 5 mL of acetic acid and then was added aminomalononitrile *p*-toluenesulfonate (506 mg, 2.0 mmol) and sodium acetate (164 mg, 2.0 mmol). The reaction mixture was stirred at room temperature overnight. The mixture was diluted with 20 mL of water and then its pH was adjusted to 8.0 by aqueous NaOH. The resulting mixture was extracted with EtOAc (3 × 50 mL). The combined organic layers were washed with water (10 mL) and brine (10 mL), dried over $MgSO_4$, filtered and concentrated *in vacuo* evaporated. The residue was purified by silica gel column chromatography (10/1, methylene chloride : methanol) to give 3-(5-amino-4-cyano-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **3** as a colorless solid (170 mg, 30%). HPLC (4 minute gradient) t_R = 1.39 min; MS m/z 282 [M+H].

B. 3-(5-Amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

To a solution of 3-(5-amino-4-cyano-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **3** (56.4 mg, 0.2 mmol) in dry THF (10 ml) under nitrogen was added phenylmagnesium bromide (1M, 1 mL) at room temperature. After 1 h, HCl solution (3N, 10 ml) was added and the mixture was stirred overnight. The solution was neutralized with dilute aqueous NaOH. The mixture was extracted with ethyl acetate (100 mL x2), washed with water and dried over Na_2SO_4. Evaporation of the solvent gave a residue which was purifiedd by HPLC to give 3-(5-Amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide as a while solid (56 mg, 78%). LCMS (4 minute gradient) t_R = 2.07 min; MS m/z 361.17 $[M + H]^+$

EXAMPLE 18


Preparation of 3-(5-Amino-4-cyclohexanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 17 except cyclohexylmagnesium bromide was used in place of phenyl magnesium bromide. HPLC (4 minute 10-90 gradient) t_R 2.01 min; MS m/z 367.29 $[M+H]^+$.

EXAMPLE 19


Preparation of 3-(5-Amino-4-cyclopentanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 17 except cyclopentylmagnesium bromide was used in place of phenyl magnesium bromide. HPLC (4 minute 10-90 gradient) t_R 1.92 min; MS m/z 353.22 $[M+H]^+$.

EXAMPLE 20


Preparation of 3-(5-Amino-4-phenylacetyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 17 except benzylmagnesium bromide was used in place of phenyl magnesium bromide. HPLC (4 minute 10-90 gradient) t_R 2.14 min; MS m/z 375.20 $[M+H]^+$.

EXAMPLE 21

Preparation of 3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



A. 3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-benzoic acid

To a solution of 3-iodobenzoic acid tert-butyl ester (4.6g) in THF (20 mL) at -40 °C under N_2 was added cyclohexylmagnesium chloride (2M in THF, 8.5 mL). The solution was kept at -40 °C to 0 °C for 20 min., when 3-(5-amino-4-cyano-imidazol-1-yl)-*N*-cyclopropyl-4-methyl-benzamide was added and the reaction was kept at rt


for 1h. Then HCl (4 M, 10 mL) was added and the mixuture was heated at 40 to 45 °C overnight. The mixture was neutralized with K_2CO_3 solution and extracted with EtOAc (2 X 100 mL) and the combined organics dried over Na_2SO_4, and concentrated. Purification of the crude product by column chromatography (EtOAc: MeOH = 6:1) gave the desired product (0.46g).

B. 3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

A solution of acid **4** (160 mg), EDCI (90 mg), and N-hydroxysucinimide (53 mg) in DMF (2 mL) was reacted at rt overnight. Water (12 ml) was added and the solution was extracted with EtOAc (15 mL X 2), dried over Na_2SO_4. Evaporation of solvent gave a residue, into which EtOAc (4 mL) and 2-propylamine (1.2 eq) was added. The reaction was kept at rt for 1 h., then concentrated and the crude product purified by column chromatography to give the desired product (yield: 80%). HPLC (4 minute gradient) t_R = 2.00 min; MS m/z 446.19 $[M + H]^+$.

EXAMPLE 22


Preparation of 3-{5-Amino-4-[3-(2-dimethylamino-ethylcarbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except 2-dimethylaminoethylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 2.18 min; MS m/z 475.15 $[M+H]^+$.

EXAMPLE 23


Preparation of 3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except ethylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.70 min; MS m/z 432.18 $[M+H]^+$.

EXAMPLE 24


Preparation of 3-[5-Amino-4-(3-methylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except methylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.61 min; MS m/z 418.15 $[M+H]^+$.

EXAMPLE 25


Preparation of 3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except cyclopropylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.74 min; MS m/z 444.14 $[M+H]^+$.

EXAMPLE 26


Preparation of 3-[5-Amino-4-(3-cyclopentylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except cyclopentylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.95 min; MS m/z 472.24 $[M+H]^+$.

EXAMPLE 27


Preparation of 3-{5-Amino-4-[3-(morpholine-4-carbonyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except morpholine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.67 min; MS m/z 474.17 $[M+H]^+$.

EXAMPLE 28


Preparation of 3-{5-Amino-4-[3-(cyclopropylmethyl-carbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except cyclopropylmethylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.86 min; MS m/z 458.23 $[M+H]^+$.

EXAMPLE 29

Preparation of 3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



A. 4-bromo-tetrahydro-pyran.

Tetrahydro-4*H*-pyran-4-ol (1.0g, 10mmol), carbon tetrabromide (3.6g, 11mmol) and triphenylphosphine (3.1g, 12mmol) were dissolved in CH_2Cl_2 (25 mL) and stirred at room temperature overnight. The crude reaction mixture was concentrated then purified by flash chromatography on silica gel (EtOAc:Hexanes = 1:20), and the product was obtained as a colorless oil (1.4g, 87%).


B. 3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

A solution of 4-bromo-tetrahydro-pyran (0.82g, 5mmol) in dry THF (10mL) was added dropwise to the suspension of magnesium (132mg, 5.5 mmol) and iodine (25mg) in dry THF (20 mL) at 50 °C under N_2. The mixture was stirred for 30 min after addition at 50 °C, then cooled to room temperature. Then a THF (10mL) solution of 3-(5-amino-4-cyano-imidazol-1-yl)-*N*-cyclopropyl-4-methyl-benzamide (90 mg, 0.32 mmol) was added to the reaction mixture and it was stirred at room temperature for 3h then quenched with HCl (2N) and stirred at room temperature overnight. The pH of the solution was adjusted pH ~ 8 with saturated aqueous K_2CO_3 was and it was extracted with EtOAc. The organic layer was washed by water and brine, dried over Na_2SO_4, and concentrated. The crude product was purified by column chromatography on silical gel (EtOAc ~ EtOAc:MeOH:Et_3N = 100:10:1),

and the product was obtained as a beige solid (35 mg, 30 %). HPLC (4 minute gradient) $t_R = 3.05$ min; MS m/z 369.18 $[M + H]^+$.

EXAMPLE 30

Preparation of 3-(5-amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



A. 2-Benzoyl-3,3-bis-methylsulfanyl-acrylonitrile


To a stirred solution of benzoylacetonitrile 1 (7.50 g, 51.7 mmol) in THF (100 ml) at 0 °C was added dry sodium hydride (2.61 g, 103 mmol). The resulting suspension was stirred at 0 °C for 45 min before carbon disulfide (2.39 ml, 54.8 mmol) was added. The reaction was then stirred at room temperature for 2 h. The resulting red solution was cooled to 0 °C, and iodomethane (6.75 ml, 109 mmol) was added. The mixture was stirred at room temperature for 18 h. Solvent was removed *in vacuo*. The residue was diluted in ether and was washed with brine. The aqueous layer was extracted twice with ether. The combined organic layers were washed twice with 5% sodium thiosulfate, and then brine. The organic layer was dried over $MgSO_4$, filtered and concentrated to give 2-benzoyl-3,3-bis-methylsulfanyl-acrylonitrile 2 as a yellow powder (9.5 g, 74%). The product was used in the next step without further purification.


B. 3-(5-Amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

Sodium (317 mg, 13.8 mmol) was added to methanol (10 ml) at 0 °C. After all of the sodium was consumed, this solution of sodium methoxide was added to a stirred solution of 2-benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (3.12 g, 12.5 mmol) in dioxane (30 ml) at 0 °C. The reaction was warmed to room temperature and then heated to 80 °C for 3 h. The dark red solution was cooled to room temperature and was added to a solution of N-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (4.00 g, 12.5 mmol) and diisopropylethylamine (2.18 ml, 12.5 mmol) in dioxane (15 ml). The mixture was heated to 85 °C for another 6 h. The solvent was removed *in vacuo*. The residue was diluted in saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 75 to 90% EtOAc/hexanes) and recrystallization from EtOAc to give the desired 3-(5-amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **3** as a white solid (950 mg, 19%). HPLC (4 minute 10-90 gradient) t_R 2.33 min; MS m/z 391.2 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.50 (d, J = 3.4, 1 H), 7.91 (d, J = 7.9, 1 H), 7.85 (s, 1 H), 7.61 (d, J = 6.9, 2 H), 7.41-7.62 (m, 4 H), 7.00 (bs, 2 H), 3.66 (s, 3 H), 2.87 (m. 1 H), 2.20 (s, 3 H), 0.70 (m, 2 H), 0.85 (m, 2 H) ppm; ^{13}C NMR (DMSO-d_6, 125MHz) δ 188.1, 166.0, 159.5, 152.8, 140.1, 139.5, 135.5, 132.9, 131.1, 130.6, 128.1, 128.0, 127.4, 126.8, 91.0, 55.2, 23.0, 15.1, 5.6 ppm.

EXAMPLE 31

Preparation of 3-(5-amino-4-benzoyl-3-ethoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide






Ethanol (0.47 ml, 8.0 mmol) was added to a suspension of dry sodium hydride (41 mg, 1.6 mmol) in dioxane (2 ml). The mixture was stirred at room temperature for 10 min. 2-Benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (0.20 g, 0.80 mmol) was added, and the mixture was stirred at 85 °C for 2.5 h. The mixture was cooled to room temperature. *N*-Cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.26 g, 0.80 mmol) was added and the reaction was heated to 85 °C for another 3 h. Solvents were removed *in vacuo*. The residue was diluted in saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 75 to 85% EtOAc/hexanes). The product was further purified by washing with a warm mixture of EtOAc and hexanes to give the desired 3-(5-amino-4-benzoyl-3-ethoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide 4 as a white solid (27 mg, 8.3%) . HPLC (4 minute 10-90 gradient) t_R 2.37 min; MS m/z 405.2 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.49 (d, J = 4.0, 1 H), 7.90 (d, J = 8.0, 1 H), 7.85 (s, 1 H), 7.62 (d, J = 7.1, 2 H), 7.50 (d, J = 7.5, 2 H), 7.41-7.45 (m, 2 H), 6.99 (bs, 2 H), 4.05 (q, J = 7.0, 2 H), 2.87 (m. 1 H), 2.19 (s, 3 H), 1.09 (t, J = 7.0, 3 H), 0.69 (m, 2 H), 0.58 (m, 2 H) ppm; ^{13}C NMR (DMSO-d_6, 125MHz) δ 188.2, 166.0, 158.9, 152.6, 140.0, 139.5, 135.5, 132.9, 131.1, 130.6, 128.1, 127.2, 126.7, 91.2, 63.4, 23.0, 17.3, 14.2, 5.6 ppm.

EXAMPLE 32


Preparation of 3-[5-amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide




Dry sodium hydride (21 mg, 0.84 mmol) was added to a solution of 2-methoxyethanol (0.63 ml, 8.0 mmol) in dioxane (2 ml) at 0 °C. The mixture was stirred at room temperature for 30 min. 2-Benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (0.20 g, 0.80 mmol) was added, and the mixture was stirred at 85 °C for 4 h. The mixture was cooled to room temperature. *N*-Cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.26 g, 0.80 mmol), followed by diisopropylethylamine (0.14 ml, 0.80 mmol), was added and the reaction was heated to 85 °C for another 11 h. Solvents were removed *in vacuo*. The residue was diluted in saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 70 to 90% EtOAc/hexanes). The product was further purified by washing with a warm mixture of EtOAc and hexanes to give the desired 3-[5-amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **5** as a white solid (60 mg, 17%) . HPLC (4 minute 10-90 gradient) t_R 2.17 min; MS m/z 435.2 $[M+H]^+$; ^1H NMR (DMSO-d_6, 500 MHz) δ 8.47 (s, 1 H), 7.89 (d, J = 7.9, 1 H), 7.83 (s, 1 H), 7.64 (d, J = 7.4, 2 H), 7.49 (d, J = 7.7, 2 H), 7.40-7.43 (m, 2 H), 6.99 (bs, 2 H), 4.12 (m, 2 H), 3.42 (m, 2 H), 3.12 (s, 3 H), 2.86 (m, 1 H), 2.07 (s, 3 H), 0.69 (m, 2 H), 0.56 (m, 2 H) ppm.

EXAMPLE 33

Preparation of 3-[5-amino-4-benzoyl-3-(2-benzyloxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide






Dry sodium hydride (21 mg, 0.84 mmol) was added to a solution of 2-benzyloxyethanol (1.1 ml, 8.0 mmol) in dioxane (2 ml) at 0 °C. The mixture was stirred at room temperature for 35 min. 2-Benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (0.20 g, 0.80 mmol) was added, and the mixture was stirred at 80 °C for 2.5 h. The mixture was cooled to room temperature. *N*-Cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.26 g, 0.80 mmol) was added and the reaction was heated to 80 °C for another 8.5 h. After the reaction mixture was cooled to room temperature, solvents were removed *in vacuo*. The residue was diluted in saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 60 to 85% EtOAc/hexanes). The product was further purified by washing with warm EtOAc to give the desired 3-[5-amino-4-benzoyl-3-(2-benzyloxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide **6** as an off-white solid (74 mg, 18%) . HPLC (4 minute 10-90 gradient) t_R

2.57 min; MS m/z 511.2 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.49 (d, J = 4.0, 1 H), 7.90 (d, J = 8.0, 1 H), 7.85 (s, 1 H), 7.66 (d, J = 7.1, 2 H), 7.49 (m, 2 H), 7.27-7.40 (m, 5 H), 7.22 (d, J = 6.7, 2 H), 7.02 (bs, 2 H), 4.34 (s, 2 H), 4.19 (m, 2 H), 3.57 (m, 2 H), 2.87 (m, 1 H), 2.17 (s, 3 H), 0.69 (m, 2 H), 0.59 (m, 2 H) ppm.

EXAMPLE 34

Preparation of 4-[5-amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester




Dry sodium hydride (41.0 mg, 1.60 mmol) was added to a solution of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester **7** (0.565 g, 2.81 mmol) in dioxane (2 ml) at 0 °C. The mixture was stirred at room temperature for 45 min. 2-Benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (0.20 g, 0.80 mmol) was added, and the mixture was stirred at 65 °C for 4 h. The mixture was cooled to room temperature. *N*-Cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.26 g, 0.80 mmol) was added and the reaction was heated to 80 °C for another 3 h. After the reaction mixture was cooled to room temperature, solvents were removed *in vacuo*.

The residue was diluted in saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 65 to 85% EtOAc/hexanes). The product was further purified by washing the with warm EtOAc to give the desired 4-[5-amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester **8** as an off-white solid (70 mg, 16%) . HPLC (4 minute 10-90 gradient) t_R 2.63 min; MS m/z 559.9 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.49 (d, J = 3.6, 1 H), 7.90 (d, J = 8.1, 1 H), 7.84 (s, 1 H), 7.59 (d, J = 7.7, 2 H), 7.40-7.51 (m, 4 H), 6.99 (bs, 2 H), 4.75 (m, 1 H), 3.18 (m, 2 H), 2.99 (m, 2 H), 2.87 (m, 1 H), 2.19 (s, 3 H), 1.68 (m, 2 H), 1.43 (m, 2 H), 1.37 (s, 9 H), 0.71 (m, 2 H), 0.58 (m, 2 H) ppm.

EXAMPLE 35


Preparation of 3-[5-amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide, trifluoroacetate salt



To a solution of 4-[5-amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester **8** (5.0 mg, 0.0089 mmol) in dichloromethane (2.0 ml) was added trifluoroacetic acid (0.5 ml). The mixture was stirred at room temperature for 3 h. Volatiles were removed *in vacuo* and the residue was washed with ether and a small amount of EtOAc to give the desired 3-[5-amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide, trifluoroacetate salt **9** as a white solid (3.0 mg, 59%). HPLC (4 minute 10-90 gradient) t_R 1.75 min; MS m/z 460.1 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.50 (d, J = 4.0, 1 H), 8.33 (bs, 2 H), 7.90 (d, J = 8.0, 1 H), 7.83 (s, 1 H), 7.62 (d, J = 7.0, 2 H), 7.44-7.52 (m, 4 H), 7.03 (bs, 2 H), 4.82 (m, 1 H), 2.97 (m, 2 H), 2.85 (m, 1 H), 2.73 (m, 2 H), 2.27 (s, 3 H), 1.91 (m, 2 H), 1.73 (m, 2 H), 0.71 (m, 2 H), 0.57 (m, 2 H) ppm.

EXAMPLE 36

Preparation of 3-(5-amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide





To a solution of benzoyl-3,3-bis-methylsulfanyl-acrylonitrile **2** (0.218 g, 0.874 mmol) in ethanol (5 ml) was added *N*-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.243 g, 0.874 mmol) and diisopropylethylamine (0.152 ml, 0.874 mmol). The mixture was heated to 65 °C for 18 h. The mixture was cooled to room temperature. Solvents were removed *in vacuo*. The residue was diluted in

saturated $NaHCO_3$ solution and was extracted three times with EtOAc. Combined organic layers were dried over $MgSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (SiO_2, gradient of 65 to 100% EtOAc/hexanes). The product was further purified by washing with a warm mixture of EtOAc and hexanes to give the desired 3-(5-amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **10** as an off-white solid (57 mg, 16%). HPLC (4 minute 10-90 gradient) t_R 2.34 min; MS m/z 407.1 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.50 (d, J = 4.0, 1 H), 7.93 (dd, J = 1.2, 7.9, 1 H), 7.86 (s, 1 H), 7.45-7.57 (m, 6 H), 6.85 (bs, 2 H), 2.88 (m, 1 H), 2.23 (s, 3 H), 2.18 (s, 3 H), 0.69 (m, 2 H), 0.58 (m, 2 H) ppm; ^{13}C NMR (DMSO-d_6, 125MHz) δ 189.3, 165.9, 152.8, 148.0, 140.4, 139.3, 135.4, 132.9, 131.2, 130.6, 128.4, 128.1, 127.2, 126.6, 102.0, 23.0, 17.3, 13.4, 5.6 ppm.


EXAMPLE 37

Preparation of 3-(5-amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide





10




11

To a suspension of 3-(5-amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **10** (40 mg, 0.098 mmol) in dichloromethane (1 ml) was added 3-chloroperoxybenzoic acid (70-75%, 53 mg, 0.22 mmol). The resulting solution was stirred at room temperature for 2 h, and then stored at 4 °C overnight. Upon warming to room temperature, the product began to precipitate. The white solid

was collected on a fritted funnel and was washed with dichloromethane and ether to give the desired 3-(5-amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide **11** (27 mg, 63%). HPLC (4 minute 10-90 gradient) t_R 1.98 min; MS m/z 439.08 $[M+H]^+$; ^1H NMR (DMSO-d_6, 300 MHz) δ 8.53 (d, J = 3.7, 1 H), 7.97 (d, J = 8.1, 1 H), 7.92 (s, 1 H), 7.76 (d, J = 7.2, 1 H), 7.48-7.64 (m, 5 H), 6.30 (bs, 2 H), 3.29 (s, 3 H), 2.88 (m, 1 H), 2.17 (s, 3 H), 0.69 (m, 2 H), 0.58 (m, 2 H) ppm.

EXAMPLE 38

Preparation of 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide





To a solution of 2-cyano-3,3-bis-methylsulfanyl-acrylamide **12** (100 mg, 0.574 mmol) in ethanol (5 ml) was added *N*-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.183 g, 0.574 mmol) and diisopropylethylamine (0.100 ml, 0.574 mmol). The mixture was heated to 65 °C for 18 h. The mixture was cooled to room temperature. Solvents were removed *in vacuo*. EtOAc was added to the residue and a solid precipitated. The solid was collected on a fritted funnel and was washed with EtOAc and ether to give the desired 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide **13**. HPLC (4 minute 10-90 gradient) t_R 1.98 min; MS m/z 439.08

$[M+H]^+$; ^1H NMR (DMSO-d_6, 500 MHz) δ 8.47 (d, J = 3.7, 1 H), 7.90 (d, J = 7.9, 1 H), 7.78 (s, 1 H), 7.49 (d, J = 8.0, 1 H), 6.81 (bs, 2 H), 6.30 (s, 2 H), 2.86 (m, 1 H), 2.45 (s, 3 H), 2.11 (s, 3 H), 0.68 (m, 2 H), 0.57 (m, 2 H) ppm.

EXAMPLE 39

Preparation of 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide






13 **14**

To a suspension of 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide **13** (100 mg, 0.289 mmol) in dichloromethane (3 ml) was added 3-chloroperoxybenzoic acid (70-75%, 157 mg, 0.637 mmol). The resulting clear solution was stirred at room temperature for 16 h. Saturated $NaHCO_3$ solution was added and the mixture was vigorously stirred for f min. The resulting suspension was filtered on a fritted funnel, and the collected solid was washed three times with H_2O, and three times with ether to give the desired 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide **14** as a white solid (87 mg, 80%). HPLC (4 minute 10-90 gradient) t_R 1.66 min; MS m/z 378.1 $[M+H]^+$; ^1H NMR (DMSO-d_6, 500 MHz) δ 8.49 (s, 1 H), 7.95 (d, J = 7.6, 1 H), 7.83 (s, 1 H), 7.53 (d, J = 7.8, 1 H), 7.46 (bs, 2 H), 6.74 (bs, 2 H), 3.40 (s, 3 H), 2.84 (m, 1 H), 2.09 (s, 3 H), 0.68 (m, 2 H), 0.55 (m, 2 H) ppm.

EXAMPLE 40


Preparation of 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester



To a solution of 2-cyano-3,3-bis-methylsulfanyl-acrylic acid ethyl ester **15** (78.0 mg, 0.359 mmol) in ethanol (3 ml) was added *N*-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetic acid salt (0.100 g, 0.313 mmol) and diisopropylethylamine (0.0626 ml, 0.359 mmol). The mixture was heated to 65 °C for 2 h. The mixture was cooled to room temperature. Solvents were removed *in vacuo*. EtOAc and ether was added to the residue and a solid precipitated. The solid was collected on a fritted funnel and was washed with EtOAc and ether to give the desired 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester **16** as a white solid (80 mg, 59%). HPLC (4 minute 10-90 gradient) t_R 2.18 min; MS m/z 375.1 $[M+H]^+$; ^{1}H NMR (DMSO-d_6, 300 MHz) δ 8.47 (d, J = 3.7, 1 H), 7.90 (d, J = 7.9, 1 H), 7.78 (s, 1 H), 7.49 (d, J = 8.0, 1 H), 6.81 (bs, 2 H), 6.30 (s, 2 H), 2.86 (m, 1 H), 2.45 (s, 3 H), 2.11 (s, 3 H), 0.68 (m, 2 H), 0.57 (m, 2 H) ppm.HPLC (4 minute 10-90 gradient) t_R 1.66 min; MS m/z 378.1 $[M+H]^+$; ^{1}H NMR (DMSO-d_6, 300 MHz) δ 8.50 (d, J = 2.6, 1 H), 7.92 (d, J = 7.9, 1 H), 7.81 (s, 1 H), 7.51 (d, J = 7.9, 1 H), 6.24 (bs, 2 H), 4.22 (q, J = 6.6, 2 H), 2.88 (m, 1 H), 2.35 (s, 3 H), 2.14 (s, 3 H), 1.29 (t, J = 6.7, 3 H), 0.72 (m, 2 H), 0.58 (m, 2 H) ppm; ^{13}C NMR (DMSO-d_6, 125MHz) δ 166.0, 163.1, 151.8, 148.5, 139.4, 135.8, 132.9, 131.1, 128.2, 126.7, 91.3, 58.9, 23.0, 17.2, 14.4, 12.3, 5.6 ppm..

EXAMPLE 41

Preparation of 3-[5-Amino-4-(3-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide



A. 2-(3-Chlorobenzoyl)-3-phenylaminoacrylonitrile


A solution of 3-chlorobenzoylacetonitrile (476 mg, 2.66 mmol, 1.0 eq) and diphenylformamidine (522 mg, 2.66 mmol, 1.0 eq) in 25 mL of toluene was stirred at room temperature for 2h then heated to 100 °C overnight. The solution was cooled and diluted with hexanes. The resulting solid was filtered and dried to provide the desired product (566 mg, 75%). HPLC (4 minute 10-95 gradient) t_R 2.97 min; MS m/z 283.2 $[M+H]^+$; ^1H NMR ($CDCl_3$), δ 8.06 (d, J = 13.2 Hz, 1 H), 7.85 (m, 2 H), 7.46 (m, 4 H), 7.27 (m, 4 H) ppm.

B. 3-[5-Amino-4-(3-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

A solution of 2-(3-chlorobenzoyl)-3-phenylaminoacrylonitrile (63 mg, 0.22 mmol, 1.0 eq), *N*-cyclopropyl-3-hydrazino-4-methylbenzamide trifluoroacetate (72 mg, 0.22 mmol, 1 eq), and triethylamine (31 μL, 0.22 mmol, 1.0 eq) in 10 mL of ethanol was heated to 65 °C for 20h. After cooling, the mixture was concentrated and the residue purified by flash chromatography on silica gel packed and eluted with 7/3 hexanes/ethyl acetate to remove byproducts followed by 3/2 ethyl acetate/hexanes to elute the title compound (33 mg, 38%) as a brown solid. HPLC (4 minute 10-95 gradient) t_R 2.35 min; MS m/z 395.1 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.92 (d, J = 7.2 Hz, 1 H), 7.77 (m, 4 H), 7.55 (m, 3 H), 2.85 (m, 1 H), 2.23 (s, 3 H), 0.80 (d, J = 5.5 Hz, 2 H), 0.63 (d, J = 2.0 Hz, 2 H) ppm; ^{13}C NMR (CD_3OD), δ 187.1, 168.1, 151.9, 141.0, 140.9, 140.0, 134.9, 133.8, 132.8, 130.9, 130.5, 129.4128.1, 127.0, 126.1, 125.6, 102.6, 22.1, 15.7, 4.6 ppm.

EXAMPLE 42


Preparation of 3-[5-Amino-4-(3-methyl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



Similar procedure as Example 41 except that 3-methylbenzoylacetonitrile was used in place of 3-chlorobenzoylacetonitrile. HPLC (4 minute 10-95 gradient) t_R 2.27 min; MS m/z 375.16 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.92 (d, J = 7.0 Hz, 1 H), 7.83 (s, 1 H), 7.80 (s, 1 H), 7.57 (m, 3 H), 7.42 (m, 2 H), 2.85 (heptet, J = 3.6 Hz, 1 H), 2.45 (s, 3 H), 2.23 (s, 3 H), 0.80 (d, J = 5.4 Hz, 2 H), 0.64 (s, 2 H) ppm.

EXAMPLE 43


Preparation of 3-[5-Amino-4-(2-methylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide



Similar procedure as Example 41 except that 2-methylbenzoylacetonitrile was used in place of 3-chlorobenzoylacetonitrile. HPLC (4 minute 10-90 gradient) t_R 2.21 min; MS m/z 375.15 $[M+H]^+$.

EXAMPLE 44


Preparation of 3-[5-Amino-4-(2-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as Example 41 except that 2-methoxybenzoylacetonitrile was used in place of 3-chlorobenzoylacetonitrile. HPLC (4 minute 10-90 gradient) t_R 2.03 min; MS m/z 391.16 $[M+H]^+$.

EXAMPLE 45


Preparation of 3-[5-Amino-4-(4-chlorobenzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as Example 41 except that 3-methylbenzoylacetonitrile was used in place of 3-chlorobenzoylacetonitrile. HPLC (4 minute 10-90 gradient) t_R 1.65 min; MS m/z 394.2 $[M+H]^+$.

EXAMPLE 46

Preparation of 3-[5-Amino-4-(2-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide



A. 2-(2-Chlorobenzoyl)-3-phenylaminoacrylonitrile

A solution of 2-chlorobenzoylacetonitrile (1.0 g, 5.6 mmol, 1.0 eq) and diphenylformamidine (1.10g, 5.6 mmol, 1.0 eq) in 50 mL of toluene was heated to 85 °C overnight. The heat source was removed and desired product slowly began to precipitate from solution. The resulting solid was filtered and dried to provide the desired product (826 mg, 52%). HPLC (4 minute 10-90 gradient) t_R 3.13 min; MS m/z 283.2 $[M+H]^+$.


B. 3-[5-Amino-4-(2-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

A solution of 2-(2-chlorobenzoyl)-3-phenylaminoacrylonitrile (93 mg, 0.33 mmol, 1.0 eq), *N*-cyclopropyl-3-hydrazino-4-methylbenzamide trifluoroacetate (104 mg, 0.33 mmol, 1 eq), and triethylamine (31 µL, 0.22 mmol, 1.0 eq) in 20 mL of ethanol was heated to 60 °C for 48h. After cooling, the mixture was concentrated and

the residue was dissolved in minimal ethyl acetate. 100 ml diethyl ether was added and the precipitate was filtered and dried to provide desired product (50 mg, 39%). HPLC (4 minute 10-90 gradient) t_R 2.51 min; MS m/z 395.1 $[M+H]^+$; ^1H NMR (DMSO), δ 8.50 (d, *J* = 3.8 Hz, 1 H), 7.93 (d, *J* = 8.0 Hz, 1 H), 7.84 (s, 1 H), 7.57 (m, 5 H) 7.32 (s, 1 H) , 7.01 (s, 2 H) 3.37 (m, 2 H), 2.86 (m, 1 H), 2.14 (s, 3 H), 1.09 (t, 2 H), 0.68 (m, 2 H), 0.58 (m, 2 H) ppm

EXAMPLE 47

Preparation of 3-[5-Amino-4-(3-methoxy-benzoyl)-pyrazol-1-yl]-4,N-dimethyl-benzamide



A. 3-methxoybenzoylacetonitrile

To a stirred solution of ethyl 3-methoxybenozoate (3.05 mL, 18.6 mmol, 1. eq) and acetonitrile (1.19 mL, 22.9 mmol, 1.23 eq) in 5 mL of THF at – 50 °C under N_2 was added via cannula a freshly prepared solution of LDA (diisopropylamine, 5.3 mL, 38.0 mmol, 2.04 eq and 2.5 *M* n-butyllithium in hexanes, 15.25 mL, 38.0 mmol, 2.04 eq). The reaction was stirred at this temperature for 3h then warmed to 0 °C for 1h. The reaction was quenched with 10 mL of sat. NH_4Cl and allowed to warm to room temperature. The mixture was extracted with EtOAc and the organic layer washed with water and brine, dried (Na_2SO_4), and concentrated. The residue was purified by flash chromatography on silica gel to provide the product as an off-white solid.

B. 2-(3-Methoxybenzoyl)-3-phenylamino-acrylonitrile

A solution of 3-methoxybenzoylacetonitrile (1.20 g, 68.5 mmol, 1.0 eq) and

diphenylformamidine (1.34 g, 68.5 mmol, 1.0 eq) in 25 mL of toluene was stirred at room temperature for 2h then heated to 100 °C overnight. The solution was cooled and diluted with hexanes. The resulting solid was filtered and dried to provide the desired product. HPLC (4 minute 10-90 gradient) t_R 3.05 min; MS m/z 279.2 $[M+H]^+$.

C. 3-[5-Amino-4-(3-methoxybenzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid


A solution of 2-(3-chlorobenzoyl)-3-phenylaminoacrylonitrile (63 mg, 0.22 mmol, 1.0 eq), 3-hydrazino-4-methylbenzoic acid hydrochloride (72 mg, 0.22 mmol, 1 eq), and triethylamine (31 μL, 0.22 mmol, 1.0 eq) in 10 mL of ethanol was heated to 65 °C for 20h. After cooling, the mixture was concentrated and the residue purified by flash chromatography on silica gel packed and eluted with 7/3 hexanes/ethyl acetate to remove byproducts followed by 3/2 ethyl acetate/hexanes to elute the title compound (15 mg, 32%) as a brown solid. HPLC (4 minute 10-90 gradient) t_R 2.13.min; MS m/z 352.2 $[M+H]^+$.

D. 3-[5-Amino-4-(3-methoxybenzoyl)-pyrazol-1-yl]-4,N-dimethyl-benzamide

To a stirred solution of 3-[5-Amino-4-(3-methoxybenzoyl)-pyrazol-1-yl]-4-methylbenzoic acid C (50 mg, 0.14 mmol, 1.0 eq) in 10 mL of DMF was added EDCI (41 mg 0.21 mmol, 1.5 eq), HOBt (29 mg,1.5 mmol, 2.0 eq), and diisopropylethylamine (55 mg, 0.43 mmol, 3.0 eq) and the solution was stirred for 15 minutes at room temperature when methylamine hydrochloride (13 mg, 0.19 mmol, 1.5 eq) was added and the reation stirred for 1 hour. The mixture was diluted with EtOAc (300 mL) and washed with water (2 x 25 mL) and brine (25 mL), dried (Na_2SO_4) and concentrated. The product was purified by flash chromatography on silica gel to provide the product to provide the product (15 mg, 32%) as a brown solid: HPLC (4 minute 10-90 gradient) t_R 1.97 min; MS m/z 365.2 $[M+H]^+$.

EXAMPLE 48

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester



A. 5-Amino-1-(5-carboxy-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

To a stirred solution of 3-hydrazino-4-methylbenzoic acid hydrochloride (Example 3A, 478 mg, 2.36 mmol, 1.0 eq) in 20 mL of ethanol were added ethyl(ethoxymethylene)cyanoacrylate (399 mg, 2.36 mmol, 1.0 eq) and triethylamine (329 μL, 2.36 mmol, 1.0 eq) and the mixture was heated at 65 °C for 5 hrs. After standing at room temperature overnight, additional 3-hydrazino-4-methylbenzoic acid hydrochloride (159 mg, 0.78 mmol, 0.3 eq) and triethylamine (110 μL, 0.78 mmol, 0.3 eq) and heated for 2.5. The mixture was cooled to room temperature and concentrated. The crude residue was purified by flash chromatography on silica gel (gradient elution from 7/3 hexanes/EtOAc to 1/1 to elute byproducts followed by EtOAc and 9/1 EtOAc/MeOH to elute product) to provide the product as a brown solid (464 mg, 68%). HPLC (4 minute 10-95 gradient) t_R 1.87 min; MS m/z 290.1 $[M+H]^+$; ^1H NMR (CD_3OD), δ 8.08 (d, J = 7.0 Hz, 1 H), 7.93 (s, 1 H), 7.76 (s, 1 H), 7.54 (d, J = 8.0 Hz, 1 H), 4.29 (q, J = 7.1 Hz, 2 H), 2.19 (s, 3 H), 1.35 (d, J = 7.1 Hz, 3 H), ppm; ^{13}C NMR (CD_3OD), δ 166.5, 163.8, 150.4, 141.7, 139.9, 135.3, 130.8, 129.5, 128.6, 93.8, 58.8, 15.8, 12.9 ppm.


B. 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

To a solution of 5-amino-1-(5-carboxy-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester (47 mg, 0.16 mmol, 1.0 eq), EDCI (62 mg, 0.32 mmol, 2.0 eq), HOBt (44 mg, 0.32 mmol, 2.0 eq), and diisopropylethyl amine (119 μL, 0.32 mmol, 2.0 eq) in DMF (5 mL) which had been stirred at RT for 15 min was added cyclopropylamine (23 μL, 0.32 mmol, 2.0 eq). After stirring overnight, the solution was diluted with EtOAc and water and the organic layer was washed with water and brine, dried (Na_2SO_4) and concentrated. The crude residue was purified by flash

chromatography on silica gel eluted with 8/2 EtOAc/hexanes to provide the product as a colorless oil (42 mg, 79%). HPLC (4 minute 10-95 gradient) t_R 1.84 min; MS m/z 329.09 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.96 (s, 1 H, NH), 7.88 (d, J = 7.9 Hz, 1 H), 7.78 (s, 1 H), 7.75 (s, 1 H), 7.50 (d, J = 8.0 Hz, 1 H), 4.28 (q, J = 7.1 Hz, 2 H), 2.83 (m, 1 H), 2.16 (s, 3 H), 1.35 (d, J = 7.0 Hz, 3 H), 0.78 (dd, J = 12.3, 7.0 Hz, 2 H), 0.63 (dd, J = 7.0, 4.5 Hz, 2 H) ppm; ^{13}C NMR (CD_3OD), δ 170.1, 165.8, 152.3, 142.1, 141.8, 137.2, 134.6, 132.8, 130.0, 128.2, 128.1, 95.8, 60.8, 24.1, 17.6, 14.9, 6.6 ppm.

EXAMPLE 49

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester



A. 5-Amino-1-(5-carboxy-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester

To a stirred solution of 3-hydrazino-4-methylbenzoic acid hydrochloride (Example 3A, 353 mg, 1.74 mmol, 1.0 eq) in 15 mL of ethanol were added 2-cyano-3-ethoxy-but-2-enoic acid ethyl ester (prepared as described by Xia et al., *J. Med. Chem.*, **1997**, *40*, 4372) (319 mg, 1.746 mmol, 1.0 eq) and triethylamine (242 μL, 1.74 mmol, 1.0 eq) and the mixture was heated at 65 °C overnight. The mixture was cooled to room temperature and concentrated. The crude residue was purified by flash chromatography on silica gel (loaded with CH_2Cl_2, and packed and eluted with gradient from 6/4 hexanes/EtOAc to elute byproducts followed by 8/2 EtOAc/hexanes and 8/2 EtOAc/MeOH to elute product) to provide the product as a brown solid (464 mg, 68%). HPLC (4 minute 10-95 gradient) t_R 1.97 min; MS m/z 304.1 $[M+H]^+$


B. 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester

To a solution of 5-amino-1-(5-carboxy-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester (150 mg, 0.49 mmol, 1.0 eq), EDCI (190 mg,

0.98 mmol, 2.0 eq), HOBt (134 mg, 0.98 mmol, 2.0 eq), and diisopropylethyl amine (362 μL, 0.98 mmol, 2.0 eq) in DMF (5 mL) which had been stirred at RT for 15 min was added cyclopropylamine (68 μL, 0.98 mmol, 2.0 eq). After stirring overnight, the solution was diluted with EtOAc and water and the organic layer was washed with water (2x) and brine, dried (Na_2SO_4) and concentrated. The crude residue was purified by flash chromatography on silica gel (gradient elution, 3/2 EtOAc/hexanes then 100% EtOAc) to provide the product as a white solid (29 mg, 17%). HPLC (4 minute 10-95 gradient) t_R 1.97 min; MS m/z 343 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.87 (d, J = 7.1 Hz, 1 H), 7.76 (s, 1 H), 7.50 (d, J = 8.0 Hz, 1 H), 4.29 (q, J = 7.0 Hz, 2 H), 2.84 (m, 1 H), 2.34 (s, 3 H), 2.18 (s, 3 H), 1.36 (t, J = 7.0 Hz, 3 H), 0.79 (d, J = 5.5 Hz, 2 H), 0.62 (s, 2 H) ppm; ^{13}C NMR (CD_3OD), δ 168.1, 164.4, 151.3, 149.8, 140.1, 135.2, 132.6, 130.7, 127.9, 126.3, 92.0, 58.6, 22.1, 15.7, 12.9, 12.6, 4.6 ppm.

EXAMPLE 50

Preparation of 5-Amino-3-[(3-chloro-benzylcarbamoyl)-methoxy]-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester



A. 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-(2-hydroxy-ethoxy)-1H-pyrazole-4-carboxylic acid ethyl ester

A stirred solution of cyano-[1,3]dioxolan-2-ylidene-acetic acid ethyl ester (prepared as described by Neidlein and Kikelj, *Synthesis*, **1988**, 981, 266 mg, 1.45

mmol, 1.0 eq), *N*-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetate (Example 6C, 463 mg, 1.45 mmol, 1.0 eq), and triethylamine (405 µL, 2.9 mmol, 2.0 eq) in 20 mL of ethanol was heated to 65 °C overnight. After cooling to room temperature, the mixture was concentrated and the residue purified by flash chromatography on silica gel (eluted with 1/1 hexanes/EtOAc followed by 100% EtOAc) to provide the desired compound as a tan solid (350 mg, 62%). HPLC (4 minute 10-95 gradient) t_R 1.59 min; MS m/z 389.06 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.87 (d, J = 7.9 Hz, 1 H), 7.78 (s, 1 H), 7.49 (d, J = 7.8 Hz, 1 H), 4.29 (dd, J = 14.9, 6.9 Hz, 2 H), 4.19 (d, J = 4.3 Hz, 2 H), 3.84 (d, J = 4.4 Hz, 2 H), 2.84 (m, 1 H), 2.22 (s, 3 H), 1.35 (t, J = 7.3 Hz, 3 H), 0.81 (d, J = 5.3 Hz, 2 H), 0.63 (s, 2 H) ppm.

B. 5-Amino-3-carboxymethoxy-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

To a stirred solution of alcohol 50A (48 mg, 0.12 mmol, 1.0 eq) in 5 mL of acetonitrile was added 2,2',6,6'-tetramethylpiperidinyloxyl (TEMPO) (catalytic) and the solution heated to 35 °C. Then a solution of sodium chlorite (17 mg, 0.24 mmol, 2.0 eq) in 2 mL of water (2mL) and an aqueous solution of sodium hypochlorite diluted to 2% (1 mL) were added simultaneously dropwise and the heating was continued for 24 hours. A bright orange color developed. The reaction was cooled to RT and diluted with water then quenched with 1*M* Na_2SO_3 and stirred for 30 minutes. The mixture was washed with EtOAc, then the pH of the aqueous layer was adjusted from pH = 8 to pH = 2 with 3 *M* HCl, and extracted with CH_2Cl_2. The CH_2Cl_2 extracts were dried and concentrated to provide the product as a yellow solid. HPLC (4 minute 10-95 gradient) t_R 1.70 min; MS m/z 403.02 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.83 (d, J = 7.9 Hz, 1 H), 7.52 (s, 1 H), 7.46 (d, J = 7.9 Hz, 1 H), 4.73, (s, 2 H), 4.19 (q, J = 7.1 Hz, 2 H), 2.84 (m, 1 H), 2.18 (s, 3 H), 1.31 (t, J = 7.0 Hz, 3 H), 0.78 (d, J = 6.1 Hz, 2 H), 0.62 (s, 2 H) ppm; ^{13}C NMR (CD_3OD), δ 171.1, 170.2, 168.4, 163.6, 159.7, 151.3, 140.5, 135.2, 132.5, 130.7, 127.6, 126.2, 80.9, 59.6, 58.7, 22.1, 15.8, 12.5, 4.6 ppm.


C. 5-Amino-3-[(3-chloro-benzylcarbamoyl)-methoxy]-1-(5-

cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

To a stirred solution of acid 50B (28 mg, 0.7 mmol, 1.0 eq), EDCI (32 mg, 0.17 mmol, 2.4 eq), HOBt (22 mg, 0.16 mmol, 2.4 mmol) in 3.0 mL of DMF at room temperature was added 3-chlorobenzylamine (18 μL, 0.07 mmol, 1.0 eq) and the mixture was stirred overnight. The mixture was diluted with EtOAc, washed with water (x2) and brine, dried (Na_2SO_4), and concentrated. The residue was purified by flash chromatography on silica gel eluted with 9/1 EtOAc/hexanes to provide the product as a clear oil (16 mg, 44%). HPLC (4 minute 10-95 gradient) t_R 2.34 min; MS m/z 525.99 $[M+H]^+$; ^1H NMR (CD_3OD), δ 7.85 (d, J = 7.9 Hz, 1 H), 7.56 (s, 1 H), 7.47 (d, J = 8.0 Hz, 1 H), 7.25 (m, 3 H), 4.73, (s, 2 H), 4.45 (s, 2 H), 4.25 (q, J = 7.0 Hz, 2 H), 2.84 (m, 1 H), 2.17 (s, 3 H), 1.27 (t, J = 7.0 Hz, 3 H), 0.80 (dd, J = 12.3, 6.7 Hz, 2 H), 0.62 (d, J = 2.2 Hz, 2 H) ppm; ^{13}C NMR (CD_3OD), δ 168.9, 168.3, 163.4, 159.6, 151.1, 140.3, 140.1, 135.2, 133.6, 132.6, 130.8, 129.1, 127.7, 126.6, 126.4, 125.0, 66.3, 58.8, 41.1, 22.1, 15.9, 12.9, 4.6 ppm.

EXAMPLE 51

Preparation of 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



A. 3-Hydrazino-4-methylbenzoic acid hydrochloride

A solution of 3-amino-4-methyl benzoic acid **1** (100 g, 0.66 mol, 1.0 equiv) in water (1.78 L) was cooled to 0 – 5 °C using ice-water. Conc. HCl (1.78 L) and

sodium nitrite (68.5 g, 0.99 mol, 1.5 equiv) were added in sequence at 0 – 5 °C. The reaction mixture was stirred at the 0 – 5 °C for 1 hour. Stannous chloride dihydrate (336 g, 1.488 mol, 2.25 equiv) in conc. HCl (540 ml) was added at 0 – 5 °C. The mixture was stirred at the same temperature for 2 hours. Solid formed during the course of the reaction was filtered and washed with water (3 x 500 ml). Dried under vacuum at 25 – 30 °C for 15 hours to provide then the crude material was(110 g) dissolved in ethanol (1 L) and stirred at 70° C for 1 hour. The material is filtered at hot and washed with ethanol (50 ml) and air dried to get the pure hydrazine **2** (60 g, 45 %) as an off white solid.

B. 3-(5-Amino-4-benzoyl-pyrazol-1-yl)-4-methylbenzoic acid

To a stirred solution of hydrazine **2** (59 g, 0.29 mol, 1.0 equiv) in ethanol (4.5 L) was added **3** (65g, 0.262 mol, 0.9 equiv, preparation: Grothasu, Davis, *J. Am. Chem. Soc.*, 58, 1334 (1936)) and triethylamine (29 g, 0.29 mol, 1.0 equiv). The mixture was heated to 65 °C. At 65 °C the reaction mixture became homogenous and was stirred at 65 °C for 4 hours. Product was precipitated out during the reaction. The solids were filtered in hot condition and dried to provide acid **4** (45 g, 53 %) as an off-white crystalline solid. HPLC (Waters X-Terra 5 micron C18 column 4.6mm x 250 mm, 1.0 mL/min, mobile phase: 0.1% TEA in H_2O/acetonitrile 40/60, 30 min elution) t_R 2.12 min, 96.6% purity; ^{1}H NMR (DMSO-d_6, 400 MHz) is consistent with Example 3.


C. 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methylbenzamide

To a stirred solution of acid **4** (46 g, 0.143 mol, 1.0 equiv) in DMF (1.9 L) was added EDCI (57.5 g, 0.299 mol, 2.09 equiv), HOBt (41.4 g, 0.306, 2.14 equiv) and diisopropylethylamine (76.6 g, 0.59 mol, 4.15 equiv) and the solution was stirred for 20 minutes at room temperature. Then it was cooled to 15-20 °C and cyclopropylamine (20.6 g, 0.36 mol, 2.51 equiv) was added and stirred at room temperature. The reaction was monitored by TLC. After 14 hours, since the reaction was not complete and additional cyclopropylamine (9.36 g, 0.16 mol, 1.14 equiv) was added and stirring was continued for two hours. DMF was removed under reduced pressure at 50 – 55 °C. To the residue EtOAc (1 L) and water (500 ml) were added

and the mixture was stirred for 10 minutes. The mixture was extracted and the organic layer was collected. The aqueous layer was extracted with EtOAc (2 x 250 ml). The combined organic layer was washed with sodium bicarbonate (2 x 500 ml) and brine (2 x 500 ml), dried over anhydrous sodium sulfate and concentrated. To the residue EtOAc/dichloromethane /hexane (50 ml/50 ml/50 ml) was added, the mixtue was stirred for 10 minutes, and filtered to provide the product (34.1 g, 65.7 %) as an off white crystalline solid. HPLC (Waters X-Terra 5 micron C18 column 4.6mm x 250 mm, 1.0 mL/min, mobile phase: 0.1% TEA in H_2O/acetonitrile 50/50, 30 min elution) t_R 5.53 min, 99.3% purity; MS m/z 360 $[M]^+$; ^1H NMR (DMSO-d_6, 400 MHz) is consistent with Example 4.

EXAMPLE 52


Preparation of 3-[5-Amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



To a solution of 3-[5-amino-4-(3-iodobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (110mg, 0.23 mmol) in DMF (5mL) was added CuCN (40mg, 0.45 mmol) and tetrakis(triphenylphosphine)palladium (catalytic) and the mixture was heated at 100 °C over night under N_2. The solvent was removed and the residue was suspended in EtOAc, and solids were filtered off. The filtrate was washed by water, brine, dried over Na_2SO_4 and concentrated. The crude product was purified by column chromatography on silica gel (EtOAc). Product was obtained as a beige solid (30mg, 34%). HPLC (4 minute 10-90 gradient) t_R 2.02 min; MS m/z 386.13 $[M+H]^+$.

EXAMPLE 53

Preparation of 3-[5-Amino-4-(3-[1,3,4]oxadiazol-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



A. 3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carbonyl]-benzoic acid

To a stirred solution of 3-[5-amino-4-(3-formylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (900mg) in CH_3CN (25mL) were added NaH_2PO_4 (55mg in 2 mL water) and H_2O_2 (1.3g, 30% solution in water) followed by the dropwise addition of an aqueous solution of $NaClO_2$ (365mg) at 10 °C. The mixture was stirred at this temperature for 4h before Na_2SO_3 was added. Solvent was removed and residue was dissolved in EtOAc, the organic layer was washed with water and brine and concentrated. The crude product was purified by column chromatography on silica gel eluted with EtOAc followed by EtOAC : AcOH = 100 :1), to provide the desired intermediate as a beige foam (345 mg, 37%).

B. 3-[5-Amino-4-(3-hydrazinocarbonylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

Compound 53A (50 mg, 0.12 mmol), *t*-butyl carbazate (33 mg, 0.24 mmol), EDCI (46mg, 0.24mmol) and HOBt (37mg, 0.24mmol) were dissolved in DMF (5 mL, dry) and stirred at room temperature over night. Solvent was removed, the residue was dissolved in EtOAc and the organics were washed by water, K_2CO_3 aqueous solution, brine, and dried over Na_2SO_4. Then TFA/DCE (5 mL, 1 : 1) was


added and stirred at room temperature for 30 min. Solvent was removed, the residue was dissolved in EtOAc, washed with K_2CO_3 aqueous solution and dried over Na_2SO_4. Solvent was removed to provide compound B as a beige solid (45 mg, 88%).

C. 3-[5-Amino-4-(3-[1,3,4]oxadiazol-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

Trimethyl orthoformate (2mL) was added to the solution of compound 53B in MeOH (2 mL) and stirred at room temperature over night. Solvent was removed, solid residue was dissolved in 1,4-dioxane, and five drops of a 4M solution of HCl in dioxane was added and the mixture was heated in the microwave at 120 °C for 30 min. The solvent was removed, the residue was dissolved in EtOAc, and the organics were washed with water, brine and the crude product was purified by preparative TLC (EtOAc : MeOH = 95 : 5) to provide product as a beige solid (25 mg, 63%). HPLC (4 minute 10-90 gradient) t_R 1.81 min; MS m/z 429.13 $[M+H]^+$.

EXAMPLE 54


Preparation of 3-{5-Amino-4-[3-(5-methyl-[1,3,4]oxadiazol-2-yl)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 53 except trimethylorthoacetate was used in place of trimethyl orthoformate. HPLC (4 minute 10-90 gradient) t_R 1.84 min; MS m/z 443.15 $[M+H]^+$.

EXAMPLE 55


Preparation of 3-{5-Amino-4-[3-(pyrrolidine-1-carbonyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except pyrrolidine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.93 min; MS m/z 458.2 $[M+H]^+$.

EXAMPLE 56


Preparation of 3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except cyclopropylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.74 min; MS m/z 444.14 $[M+H]^+$.

EXAMPLE 57


Preparation of 3-[5-Amino-4-(3-carbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 21 except ammonia was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.51 min; MS m/z 404.2 $[M+H]^+$.

EXAMPLE 58

Preparation of 3-[5-Amino-4-(3-isopropylcarbamoylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide









Similar procedure as in Example 21B except 3-[5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-1H-pyrazole-4-carbonyl]-benzoic acid was






used in place of 3-[5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-benzoic acid. HPLC (4 minute 10-90 gradient) t_R 1.89 min; MS m/z 446.2 $[M+H]^+$.

EXAMPLES 59 - 69

The following examples were prepared with a procedure similar to Example 58 except the appropriate amine was used in place of isoprpylamine.

Table 1.



Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
59		3-[5-Amino-4-(4-methylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.03	418.2
60		3-[5-Amino-4-(4-cyclopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.77	444.26
61		3-[5-Amino-4-(3-carbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.64	404.16
62		3-{5-Amino-4-[3-(piperazine-1-carbonyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide	2.27	473.21
63		3-[5-Amino-4-(3-dimethylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.71	432.21
64		3-{5-Amino-4-[3-(cyclopropylmethyl-carbamoyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide	1.9	458.25





65		3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.77	432.18
66		3-[5-Amino-4-(3-methylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.69	418.17
67		3-[5-Amino-4-(3-cyclopentylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.0	472.23
68		3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.89	446.2
69		3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.82	444.15

EXAMPLES 70 - 75

The following examples were prepared with a procedure similar to Example 11 except the appropriate amine was used in place of 1-methylpiperazine.

Table 2.



Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
70		3-(5-Amino-4-{3-[(3-chloro-benzylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.74	514.17
71		3-[5-Amino-4-(3-{[2-(3-chloro-phenyl)-ethylamino]-methyl}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.74	528.2









72		3-(5-Amino-4-{3-[4-(3-chloro-phenyl)-piperazin-1-ylmethyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.82	569.23
73		3-[5-Amino-4-(3-{[2-(2-benzyloxy-5-chloro-phenyl)-ethylamino]-methyl}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.11	634.2
74		3-(5-Amino-4-{3-[(2-morpholin-4-yl-ethylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.03	503.22
75		3-(5-Amino-4-{3-[(3-morpholin-4-yl-propylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.02	517.25









EXAMPLES 76 - 93

The following examples were prepared with a procedure similar to Example 17 except the appropriate grignard reagent was used in place of phenylmagnesium bromide.

Table 3.


Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
76		3-[5-Amino-4-(pyridine-2-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.56	362.23
77		3-[5-Amino-4-(2-methyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.95	375.1

78		3-[5-Amino-4-(3,4-difluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.53	397.17
79		3-[5-Amino-4-(3-fluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.23	379.2
80		3-[5-Amino-4-(3,4-dichloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.95	429.11
81		3-[5-Amino-4-(3-methoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.22	391.19
82		3-[5-Amino-4-(4-fluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.29	379.1
83		3-[5-Amino-4-(3,5-dichloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	3.07	429.13
84		3-[5-Amino-4-(4-methoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.12	391.11
85		3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-benzoic acid tert-butyl ester	2.53	461.05

86		3-[5-Amino-4-(3,5-difluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.58	397.18
87		3-[5-Amino-4-(benzo[1,3]dioxole-5-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.12	405.11
88		3-[5-Amino-4-(4-chloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.54	395.15
89		3-[5-Amino-4-(3,4-dimethoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.95	420.1
90		3-[5-Amino-4-(3-benzyloxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.51	467
91		3-[5-Amino-4-(4-fluoro-3-methyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methylbenzamide	2.25	393
92		{3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-phenoxy}-acetic acid tert-butyl ester	2.23	491
93		3-[5-Amino-4-(3-chloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.54	395.17

EXAMPLE 94


Preparatation of 3-(5-Amino-4-benzoyl-2-methyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in except 17 except triethylorthoacetate was used in place of triethylorthoformate. HPLC (4 minute 10-90 gradient) t_R 1.56 min; MS m/z 375 $[M+H]^+$.

EXAMPLE 95


Preparatation of 3-(5-Amino-4-benzoyl-2-propyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in except 17 except triethylorthbutyrate was used in place of triethylorthoformate. HPLC (4 minute 10-90 gradient) t_R 2.14 min; MS m/z 403 $[M+H]^+$.

EXAMPLE 96

Preparation of 3-[5-Amino-4-(3-carbamoylmethoxy-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



A. {3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carbonyl]-phenoxy}-acetic acid

To a stirred solution of 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (400mg, 1.06mmol) and *t*-butyl chloroacetate

(319mg, 2.12mmol) in DMF (20 mL) was added K_2CO_3 (292mg, 2.12 mmol) and the mixture was heated at 100 °C overnight. The solvent was removed, and the residue was suspended in EtOAc, washed by water, brine, dried over Na_2SO_4, and concentrated. The crude product was purified by column chromatography on silica gel (EtOAc : Hexane = 3 :1) to provide the product as a light yellow oil (140 mg, 27%).

B. {3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carbonyl]-phenoxy}-acetic acid

The oil (180mg, 0.37 mmol) from last step was dissolved in DCM (5 mL), TFA (5 mL) was added and stirred at room temperature over night. Volatile organics were removed, toluene was added then removed *in vacuo* to provide the product as a white solid (140 mg, 88%).


C. 3-[5-Amino-4-(3-carbamoylmethoxy-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide








To a solution of the intermediate obtained in the previous step in THF (5 mL) was added $SOCl_2$ (1mL) and the mixture was stirred at room temperature for 2 h. The volatiles were removed, then NH_3 (0.5 *M* dioxane solution) was added and the mixture was stirred at room temperature for 30 min. The solvent was removed, and the residue was purified by preparative TLC (EtOAc : MeOH : Et_3N = 100: 10: 1) and then by preparatory HPLC to provide the product as a beige solid (4.2mg, 10%). HPLC (4 minute 10-90gradient) t_R 1.78 min; MS m/z 434.14 $[M+H]^+$.


EXAMPLEs 97 - 105

The following examples were prepared with a procedure similar to Example 96 except the appropriate amine was used in place of ammonia.

Table 4

Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
97		3-(5-Amino-4-{3-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.56	517.22


98		3-{5-Amino-4-[3-(2-oxo-2-piperazin-1-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide	1.48	503.21
99		3-(5-Amino-4-{3-[2-(3-amino-pyrrolidin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.63	503.2
100		3-(5-Amino-4-{3-[2-(3-methylamino-pyrrolidin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.51	517.22
101		3-(5-Amino-4-{3-[2-(3,5-dimethyl-piperazin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	1.62	531.26
102		3-{5-Amino-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide	1.92	504.2
103		3-[5-Amino-4-(3-{[(1H-benzoimidazol-2-ylmethyl)-carbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	1.54	564.21
104		3-[5-Amino-4-(3-{[2-(2-benzyloxy-5-chloro-phenyl)-ethylcarbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	3.2	678.26

105		3-[5-Amino-4-(3-{[2-(5-chloro-2-hydroxy-phenyl)-ethylcarbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.49	588.19

.

EXAMPLE 106


Preparation of 3-[5-Amino-4-(3-pyrazin-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



To a stirred solution of 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (130mg, 0.27 mmol) and 2-tributylstannylpyrazine (119 mg, 0.32 mmol) in DMF (2 mL) was added tetrakis(triphenylphosphine) palladium (catalytic) and the mixture was heated in microwave at 160 °C for 15 min. The solvent was removed and the residue was dissolved in EtOAc. The organics were washed by water, brine and concentrated. The crude material was purified by preparatory HPLC to provide the desired product as beige solid (6.2 mg, 5%). HPLC (4 minute10–90 gradient) t_R 2.12 min; MS m/z 439.19 $[M+H]^+$.

EXAMPLE 107


Preparation of 3-[5-Amino-4-(3-pyridin-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



Similar procedure as in Example 106 except 2-tributylstannylpyridine was used in place of 2-tributylstannylpyrazine. HPLC (4 minute 10-90 gradient) t_R 1.94 min; MS m/z 438.26 $[M+H]^+$.

EXAMPLE 108

Preparation of 3-[5-Amino-4-(pyridine-2-carbonyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide



A. 3-(5-Amino-4-cyano-pyrazol-1-yl)-*N*-cyclopropyl-4-methylbenzamide


DIPEA (3.4g, 26.5 mmol) was added to the solution of *N*-cyclopropyl-3-hydrazino-4-methyl-benzamide trifluoroacetate (8.45g, 26.5 mmol) and 2-ethoxymethylene malononitrile (3.2g, 26.5 mmol) in EtOH (100mL) and stirred at 65 °C for 3h. The solvent was removed the residue was suspended in EtOAc (~100 mL), and water was added to this suspension. The solid product was filtered. The filtrate was washed by water, brine, dried over Na_2SO_4, concentrated, and purified by column chromatography on silica gel eluted with EtOAc. The product, combined from filtration and chromatography, was obtained as a beige solid (7.1g, 96%).

B. 3-[5-Amino-4-(pyridine-2-carbonyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

Cyclohexylmagnesium chloride (5 mL, 2.0 *M* in Et_2O) was added dropwise to the solution of 2-iodopyridine (1.03g, 5 mmol) in THF (15mL) at -20 °C. The mixture was stirred at this temperature for 20 min before 3-(5-Amino-4-cyano-pyrazol-1-yl)-*N*-cyclopropyl-4-methyl-benzamide (140 mg, 0.5 mmol) was added, then stirred at room temperature over night. The reaction was quenched by the addition of K_2CO_3 aqueous solution, then the mixture was extracted with EtOAc and the organic layers washed with water and brine. Purification by column chromatography on silica gel eluted with EtOAc provided the product as a white solid (60mg, 33%). HPLC (4 minute 10-90 gradient) t_R 2.09 min; MS m/z 362.20 $[M+H]^+$.

EXAMPLE 108

Preparation of 3-(5-Amino-4-cyclopentanecarbonyl-pyrazol-1-yl)-*N*-cyclopropyl-4-methylbenzamide















Similar procedure as in Example 17B except 3-[5-amino-4-(3-cyano-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide was used in place of 3-(5-amino-4-cyano-imidazol-1-yl)-*N*-cyclopropyl-4-methyl-benzamide and cyclopentylmagnesium bromide was used in place of phenyl magnesium bromide. HPLC (4 minute 10-90 gradient) t_R 2.54 min; MS m/z 353.19 $[M+H]^+$.






EXAMPLES 110 - 126

The following examples were prepared with a procedure similar to Example 109 except the appropriate grignard reagent was used in place of cyclopentylmagnesium bromide.

Table 5.


Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
110		3-[5-Amino-4-(benzo[1,3]dioxole-5-carbonyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide	2.67	405.16
111		3-[5-Amino-4-(2-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.05	379.13
112		3-[5-Amino-4-(3-ethoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.58	405.17

113		3-[5-Amino-4-(3-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.37	391.17
114		3-[5-Amino-4-(3-cyclopropoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.68	417.2
115		3-[5-Amino-4-(4-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.24	391.19
116		3-[5-Amino-4-(3,4-dimethoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.11	421.17
117		3-[5-Amino-4-(4-methylsulfanyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.59	407.14
118		3-[5-Amino-4-(3-methylsulfanyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.61	407.15
119		3-[5-Amino-4-(4-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.18	379.23
120		3-[5-Amino-4-(3-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.18	379.15
121		3-[5-Amino-4-(3,4-difluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.25	3.97.16

122		3-[5-Amino-4-(3,5-difluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.28	397.17
123		3-[5-Amino-4-(4-fluoro-3-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.32	393.23
124		3-[5-Amino-4-(4-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.15	379.17
125		3-(5-Amino-4-cyclohexanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide	2.75	367.22
126		3-[5-Amino-4-(3-vinyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.64	403.15

EXAMPLE 127

Preparation of 3-{5-Amino-4-[3-([1,3,4]oxadiazol-2-ylmethoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide



A. 3-[5-Amino-4-(3-hydrazinocarbonylmethoxy-benzoyl)-pyrazol-1-yl]-*N*-

cyclopropyl-4-methylbenzamide


Compound 97A (300 mg, 0.69 mmol), *t*-butyl carbazate (182 mg, 1.38 mmol), EDCI (263 mg, 1.38 mmol) and HOBt (211mg, 1.38 mmol) were dissolved in DMF (10 mL, dry) and stirred at room temperature for 2h. Solvent was removed, residue was dissolved in EtOAc, washed by water, K_2CO_3 aqueous solution, brine, and dried over Na_2SO_4. Then TFA/DCE (5 mL, 1 : 1) was added and stirred at room temperature for 30 min. Solvent was removed, residue was dissolved in EtOAc, washed with K_2CO_3 aqueous solution and dried over Na_2SO_4. The crude resiude was purified by column chromatography on silica gel (EtOAc : MeOH = 10 : 1), and the desired compound was obtained as a white solid (88 mg, 28%).


B. 3-{5-Amino-4-[3-([1,3,4]oxadiazol-2-ylmethoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide

Trimethyl orthoformate (3mL) was added to the solution of compound 52B (48mg, 0.11mmol) in MeOH (3 mL) and stirred at room temperature over night. Solvent was removed, solid residue was dissolved in 1,4-dioxane, five drops of HCl in dioxane (4M) was added, and the mixture heated in microwave at 120 °C for 30 min. Solvent was removed, residue was dissolved in EtOAc, washed with water and brine. The crude product was purified by preparatory HPLC, and the product was obtained as a white solid (3.3 mg, 6.7%). HPLC (4 minute 10–90 gradient) t_R 2.07 min; MS m/z 459.13 $[M+H]^+$.

EXAMPLE 128

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-methyl-cyclohexyl)-amide





A. 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid








A solution of 5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester (1.0g, see Example 44), LiOH (1.2 g) in water/ethanol (15 ml /20 ml) was heated at 50 °C overnight. The solution was neutralized with dilute HCl (2M), and extracted with ethyl acetate (200 ml x 2), dried over Na_2SO_4. The solvent was evaporated under reduced pressure and ethyl acetate and diethyl ether was added. The resulting solid was filtered to give the desired 5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-1H-pyrazole-4-carboxylic acid. (0.8g; yield: 88%.).









B. 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-methyl-cyclohexyl)-amide









A solution of 5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-1H-pyrazole-4-carboxylic acid (21 mg), 2-methylcyclohexylamine (10 mg), EDCI (28 mg) and HOBt (12 mg) in DMF (0.75 ml) was reacted at room temperature for 24 h. Water (4 ml) was added and the solution was extracted with ethyl acetate (3 mL x 2). The organic phase was then washed with water (3 mL), dried over Na_2SO_4 and evaporated. The residue was purified by preparative TLC plate to give the desired product 5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-1H-pyrazole-4-carboxylic acid (2-methylcyclohexyl)-amide (19 mg, 70%). HPLC (4 minute gradient) t_R 2.34 min; MS m/z 396.31 $[M+H]^+$.






EXAMPLES 129 – 156

The following examples were prepared with a procedure similar to Example 128 except the appropriate amine was used in place of 2-methylcyclohexylamine. Table 6.

Ex	Structure	Name	HPLC t_R(min)	MS m/z $[M+H]^+$
129		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid phenylamide	2.4	376.2
130		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethyl-propyl)-amide	2.16	370.3
131		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid bicyclo[2.2.1]hept-2-ylamide	2.41	394.3
132		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethy	2.57	406.2
133		3-[5-Amino-4-(2,5-dimethyl-pyrrolidine-1-carbonyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.17	383.29
134		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 4-methoxy-benzylamide	2.15	420.3
135		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid indan-1-ylamide	2.35	416.75


136		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid benzylamide	2.04	390.23
137		3-[5-Amino-4-(piperidine-1-carbonyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide	2.05	368.16
138		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylamide	2.19	382.25
139		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclopropylamide	1.72	340.23
140		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-chlorobenzylamide	2.31	424.31
141		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclopentylamide	2.03	368.29
142		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 2,4-dichloro-benzylamide	2.54	458.24
143		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylmethyl-amide	2.38	396.4

144		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3,4-dichloro-benzylamide	2.5	458.23
145		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (4-methyl-cyclohexyl)-amide	2.45	396.32
146		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-trifluoromethyl-benzylamide	2.44	458.25
147		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (4-tert-butyl-cyclohexyl)-amide	2.86	438.4
148		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2,2-dimethyl-propyl)-amide	2.19	370.28
149		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-methoxy-benzylamide	2.13	420.29
150		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 4-fluoro-benzylamide	2.19	408.26
151		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethyl-pyrrolidin-2-ylmethyl)-amide	1.41	411.28

152		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 2-methyl-benzylamide	2.26	404.33
153		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide	1.39	391.14
154		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-morpholin-4-yl-ethyl)-amide	1.12	413.25
155		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-cyclohexyl-ethyl)-amide	2.53	410.45
156		5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-phenyl-ethyl)-amide	2.22	404.28

EXAMPLE 157

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2,2-dimethyl-propyl)-amide




Similar procedure as in Example 128 except neopentylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 2.19 min; MS m/z 370.32 $[M+H]^+$.

EXAMPLE 158

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-


pyrazole-4-carboxylic acid (1-ethyl-pyrrolidin-2-ylmethyl)-amide



Similar procedure as in Example 128 except 2-(aminomethyl)-1-ethylpyrrolidine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.41 min; MS m/z 411.25 $[M+H]^+$.

EXAMPLE 159


Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide



Similar procedure as in Example 128 except 1-(2-aminoethyl)pyrrolidine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 1.34 min; MS m/z 397.22 $[M+H]^+$.

EXAMPLE 160


Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexyl-methyl-amide



Similar procedure as in Example 57 except *N*-methylcyclohexylamine was used in place of isopropylamine. HPLC (4 minute 10-90 gradient) t_R 2.37 min; MS m/z 396.3 $[M+H]^+$.

EXAMPLE 161

Preparation of 3-[5-Amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide



A. 3-[5-amino-4-(3-iodobenzoyl)-pyrazol-1-yl]-4-methylbenzoic acid

To a stirred solution of hydrazine **2** (32.5 g, 0.160 mol, 1.0 equiv) in ethanol (3.6 L) was added **1** (60g, 0.160 mol, 1.0 equiv, preparation: International Patent Application Publication No. WO 02/57101 A1, pg. 84)) and triethylamine (16.56 g, 0.16 mol, 1.0 equiv) and the mixture was heated to 65 °C. All solids dissolved when temperature reached 65°C. After cooling, the solids were filtered to provide acid **3** (22 g, 30 %) as a light brown solid. HPLC (Waters X-Terra 5 micron C18 column 4.6mm x 250 mm, 1.0 mL/min, mobile phase: 0.1% TEA in H_2O/acetonitrile 40/60, 30 min elution) t_R 6.74 min, 95.2% purity;

B. 3-[5-amino-4-(3-iodobenzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

To a stirred solution of acid **3** (21 g, 0.0469 mol, 1.0 equiv) in DMF (30 ml) was added EDCI (17 g, 0.0886 mol, 2.0 equiv), HOBt (12.6 g, 0.0939, 2.0 equiv) and diisopropylethylamine (24.2 g, 0.18 mol, 4.0 equiv) and the solution was stirred for 15 minutes at room temperature when cyclopropylamine (10.7 g, 0.0939, 2.0 equiv) was added and the reaction stirred for 1 hour. The mixture was added to water and extracted with EtOAc (1 L) and washed with water (2 x 200 ml) and brine (200 ml), dried over anhydrous sodium sulfate and concentrated. The product was purified by flash chromatography on silica gel eluted with 8/2 EtOAc/hexanes to provide the product as brown oil. The product was further purified by trituration with Isopropyl ether (1 L) and dried under vacuum to provide the amide **4** (12 g, 50 %) as an off


white solid.


C. 3-[5-Amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

To a stirred solution of iodide **4** (7.4 g, 0.015 mol, 1.0 equiv) in N,N-dimethyl formamide (25 ml) under N_2 was added copper cyanide. The resultant suspension was refluxed for 1 hour. The reaction was monitored by TLC and checked for the completion. Heating was discontinued and the reaction mixture was cooled down to 80°C. Ice water (15 ml), 25 % aqueous ammonia (15 ml), and ethyl acetate (50 ml) were added to quench the reaction. The mixture was filtered to remove the solids. From the filtrate organic layer was separated. The organic layer was washed with water, saturated brine and dried over anhydrous sodium sulfate. Solvent was removed under reduced pressure and the crude material dried at 60°C under high vacuum then purified by column with ethyl acetate as eluent to provide the product (3 g, 51 %) as an off white solid. HPLC (Waters X-Terra 5 micron C18 column 4.6mm x 250 mm, 1.0 mL/min, mobile phase: 0.1% TEA in H_2O/acetonitrile 70/30, 30 min elution) t_R 23.70 min, 99.4% purity; ^{1}H NMR (400 MHz, $CDCl_3$) δ8.10 (t, J = 2.9 Hz, 1 H), 8.06 (t, J = 2.7 Hz, 1 H), 8.04 (m, 1 H), 7.85 (t, J = 2.5 Hz, 1 H), 7.83 (m, 1 H), 7.81 (d, J = 1.8 Hz, 1 H), 7.95 (d, J = 1.6 Hz, 1 H), 7.37 (s, 1 H), 7.65 (t, J = 7.8 Hz, 1 H), 7.45 (d, J = 8.0 Hz, 1 H), 6.41 (s, 1 H), 5.94 (s, 2 H), 2.88 (m, 1 H), 2.24 (s, 3 H), 0.86 (dd, J = 6.9, 6.1 Hz, 2 H) 0.61 (m, 2 H) ppm; ^{13}C NMR (400 MHz, $CDCl_3$) δ186.6, 167.2, 151.5, 141.3, 140.7, 140.3, 135.1, 134.5, 134.0, 132.2, 131.8, 129.5, 128.7, 126.2, 118.1, 113.0, 103.7, 30.9, 23.2, 17.7, 6.8 ppm.

EXAMPLE 162

Preparation of 3-[5-Amino-4-(3-pyrazin-2-yl-benzoyl)-imidazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide





A. Preparation of 3-[5-Amino-4-(3-bromobenzoyl)-imidazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide


To a solution of 3-bromo-1-iodobenzene (2.82g) in THF (20 mL) at -40 °C under N_2 was added cyclohexylmagnesium chloride (2M in THF, 6 mL). The solution was kept at -40 °C to 0 °C for 20 min. then 3-(5-amino-4-cyano-imidazol-1-yl)-*N*-cyclopropyl-4-methyl-benzamide (0.18g) was added and the reaction was kept at rt for 1h. Then HCl (4 M, 20 mL) was added and the mixture was kept at rt for two days. The mixture was neutralized with K_2CO_3 solution and extracted with EtOAc (2 X 100 mL) and the combined organics dried over Na_2SO_4, and concentrated. Purification of the crude product by column chromatography (EtOAc) gave the desired product (0.15g, 54%). HPLC (4 minute 10-90 gradient) t_R 1.85 min; MS m/z 438.10, 441.07 $[M+H]^+$.

B. Preparation of 3-[5-Amino-4-(3-pyrazin-2-yl-benzoyl)-imidazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

A solution of 3-[5-amino-4-(3-bromobenzoyl)-imidazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (60 mg), 2-tributyltinpyrazine (120 uL) and tetrakis(triphenylphosphine)palladium(0) (30 mg) in 1,4-dioxane (0.8 mL) was heated under microwave radiation at 160 °C for 25 min. Water (3 ml) was added and the mixture was extracted with ethyl acetate (4 mL x 2). Evaporation of the solvent and purification by preparatory TLC (EtOAc: MeOH = 10:1) gave the desired product (38 mg, 63%). HPLC (4 minute 10-90 gradient) t_R 2.18 min; MS m/z 439.27 $[M+H]^+$.

EXAMPLE 163


Preparation of 3-[5-Amino-4-(3-pyrimidin-5-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 162 except 4-tributylstannylpyrimidine was used in place of 2-tributylstannylpyrazine. HPLC (4 minute 10-90 gradient) t_R 1.93 min; MS m/z 439.19 $[M+H]^+$.

EXAMPLE 164


Preparation of 3-[5-Amino-4-(3-pyrimidin-5-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 162 except 2-tributylstannylpyridine was used in place of 2-tributylstannylpyrazine. HPLC (4 minute 10-90 gradient) t_R 2.04 min; MS m/z 438.25 $[M+H]^+$.

EXAMPLE 165


Preparation of 3-[5-Amino-4-(3-pyrimidin-2-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 162 except 2-tributylstannylpyrimidine was used in place of 2-tributylstannylpyrazine. HPLC (4 minute 10-90 gradient) t_R 2.49 min; MS m/z 439.24 $[M+H]^+$.

EXAMPLE 166


Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-methylsulfonylbenzoyl-1H-imidazole



A solution of 5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-methylsulfonylbenzoyl-1H-imidazole (64 mg) and Oxone® (242 mg) in water/methanol (2ml/2ml) was reacted at room temperature for 2 hours.. The solution was evaporated and THF/MeOH (2ml/2ml) was added. The mixture was heated to dissolve the organic materials and filtered. The filtrate was concentrated under reduced pressure and the residue was purified by preparative HPLC to provide the product (1.6 mg). HPLC (4 minute 10-90 gradient) t_R 1.90 min; MS m/z 439.15 $[M+H]^+$.

EXAMPLE 167

Preparation of 5-Amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-[3-(5-methyl)oxadiazol-3-ylbenzoly]-1H-imidazole





A mixture of *N*-hydroxy-3-iodobenzimidine (1.31g), acetic anhydride (1 mL) and catalytic amount of *p*-toluenesulfonic acid was heated in acetic acid (10 mL) at 90 °C for 6 hour. The solvent was evaporated and water and methanol (1:1; 100 mL)) was added. The resulting precipitate was filtered to give the desired product (1.1 g).

To a solution 3-(3-iodophenyl)-5-methyloxadiazole (0.286g) in THF (15 ml) under N_2 was added cyclopentylmagnesiumbromide (2 M, 1.1 mL) at -20 °C. The temperature was gradually raised to 5-10 °C. It took about 20 minutes. Then 5-amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-cyanoimidazole (96 mg) was added and the reaction was kept at room temperature for 0.5 hour. Then diluted HCl (4M, 12 mL) was added and the reaction was heated at 60 °C for 3 hour. After petition in water and ethyl acetate, the final product was purified by column chromatography (Ethyl acetate/Hexane = 1:1). HPLC (4 minute 10-90 gradient) t_R 2.03 min; MS m/z 443.20 $[M+H]^+$.

EXAMPLE 168

Preparation of 3-(5-Amino-4-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide





A. 3-{5-Amino-4-[3-(2-bromo-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methylbenzamide


To a solution of 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide (472 mg, 1.26 mmol) in THF (dry, 20 mL) at 0°C was added of 2-bromoethanol (785 mg, 6.3 mmol), PPh_3 (1.3g, 5.04 mmol) and diethyl azodicarboxylate (877mg, 5.04 mmol) and the mixture was stirred at room temperature for 72 h. An aqueous solution of NH_4Cl was added then the THF layer was isolated. The aqueous layer was extracted by EtOAc. The combined organic phase was washed by brine, purified by preparatory HPLC, to provide the product as a white solid (360 mg, 59%). HPLC (4 minute 10-90 gradient) t_R 2.72 min; MS m/z 483.14/485.09 $[M+H]^+$.

B. 3-(5-Amino-4-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methylbenzamide

To the suspension of 3-{5-amino-4-[3-(2-bromoethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide (24 mg, 0.05mmol) in EtOH (1ml) was added 1-methylpiperazine (100mg, 1 mmol) and the mixture was stirred at 80 °C overnight. The solvent was removed, residue was dissolved in EtOAc, washed by water, dried over Na_2SO_4. Solvent was removed, desired product was obtained as a colorless oil (18 mg, 72%). HPLC (4 minute gradient) t_R 1.33 min; MS m/z 503.29 $[M+H]^+$.

EXAMPLE 169


Preparation of 3-[5-Amino-4-(3-{2-[bis-(2-hydroxy-ethyl)-amino]-ethoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 168 except 2-[bis-(2-hydroxyethyl)-amino]-ethanol was used in place of 1-methylpiperazine. HPLC (4 minute 10-90 gradient) t_R 1.39 min; MS m/z 508.24 $[M+H]^+$.

EXAMPLE 170


Preparation of 3-{5-Amino-4-[3-(2-dimethylamino-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 168 except 2-dimethylaminoethanol was used in place of 1-methylpiperazine. HPLC (4 minute 10-90 gradient) t_R 1.59 min; MS m/z 448.22 $[M+H]^+$.

EXAMPLE 171

Preparation of 3-{5-Amino-4-[3-(2,3-dihydroxy-propoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide



A. 3-{5-Amino-4-[3-(2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide


To a stirred solution of 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methylbenzamide (75mg, 0.2 mmol) and 4-chloromethyl-2,2-dimethyl-[1,3]dioxolane (36 mg, 0.24mmol) were dissolved in DMF (4ml) was added K_2CO_3 and the mixture stirred at 150 °C for 2 h in microwave oven. The solvent was removed, residue was dissolved in EtOAc, washed by water, brine, dried over Na_2SO_4, and concentrated. The residue was purified by column chromatography (EtOAc : Hex = 3 :1) and the product was obtained as a colorless oil (18mg, 18%). HPLC (4 minute gradient) t_R 2.57 min; MS m/z 491.11 $[M+H]^+$.

B. 3-{5-Amino-4-[3-(2,3-dihydroxy-propoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide

Compound 171A (10 mg) was dissolved in HOAc (1ml), 0.5 ml water was added, and the mixture was stirred at 50 °C overnight. The solvent was removed and the residue was purified by preparative TLC (EtOAc), to provide the product as a light yellow oil (4.1 mg, 45%). HPLC (4 minute 10-90 gradient) t_R 1.73 min; MS m/z 451.21 $[M+H]^+$.

EXAMPLE 172

Preparation of 3-(5-Amino-4-{3-[2-(4-chloro-phenoxy)-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide




To a stirred solution of 3-{5-amino-4-[3-(2-bromoethoxy)-benzoyl]-pyrazol-1-

yl}-N-cyclopropyl-4-methyl-benzamide (24 mg, 0.05 mmol) and 4-chlorophenol (128mg, 1mmol) in DMF (1mL), was added K_2CO_3 (138 mg, 1mmol) and the mixture was stirred at 100 °C for 2h. The mixture was cooled and the solids were filtered off. The filtrated was concentrated and purified by preparatory HPLC, to provide the product as a beige solid (10mg, 38%). HPLC (4 minute 10-90 gradient) t_R 2.77 min; MS m/z 531.21 $[M+H]^+$.

EXAMPLE 173


Preparation of 3-{5-Amino-4-[3-(4H-[1,2,4]triazol-3-yl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



To a stirred solution of 3-[5-amino-4-(3-carbamoyl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide (120 mg) in DMF (2 mL), was added N,N-dimethylformamide dimethyl acetal (5 ml) and the solution was stirred at 80 °C overnight. The solvent was removed, residue was dissolved in AcOH (5ml), hydrazine monohydrate (3ml) was added and the mixture stirred at 90 °C overnight. The solution was acidified with hydrochloric acid to pH ~ 1and stirred at 80 °C overnight. The solvent was removed and the residue was resuspended in EtOAc, then washed by aqueous K_2CO_3 solution, water, and brine, and concentrated. The crude product was purified by column chromatography on silica gel eluted with EtOAc to provide product as a white solid (65 mg, 51%). HPLC (4 minute 10-90 gradient) t_R 1.58 min; MS m/z 428.18 $[M+H]^+$.

EXAMPLE 174

Preparation of 3-{5-Amino-4-[3-(4H-[1,2,4]triazol-3-yl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide



Similar procedure as in Example 173 except 3-[5-amino-4-(3-carbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide was used in place of 3-[5-amino-4-(3-carbamoyl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide. HPLC (4 minute 10-90 gradient) t_R 1.58 min; MS m/z 428.24 $[M+H]^+$.

EXAMPLE 175

The ability of the compounds provided herein to inhibit the synthesis or the activity of cytokines can be demonstrated using the following *in vitro* assays.

Generation of p38 kinases

cDNAs of human p38α and β were cloned by PCR. The α and β cDNAs were subcloned into DEST2 plasmid (Gateway, InVitrogen). His_6-p38 fusion protein was expressed in *E. coli* and purified from bacterial lysates by affinity chromatography using Ni^{+2}-NTA-agarose. His_6-p38 protein was activated by incubating with constitutively active MKK6. Active p38 was separated from MKK6 by affinity chromatography. Constitutively active MKK6 was generated in a manner similar to Raingeaud *et al.* [*Mol. Cell. Biol.*, 1247-1255 (1996)].

TNF-α Production by LPS-Stimulated PBMCs

Heparinized human whole blood was obtained from healthy volunteers. Peripheral blood mononuclear cells (PBMCs) were purified from human whole blood by Accu-paque density gradient centrifugation and resuspended at a concentration of 5 x 10^6/ml in assay medium (RPMI medium containing 10% fetal bovine serum). 175 μL of cell suspension was incubated with 10 μL of test compound (in 4% DMSO) in 96-well tissue culture plates for 30 minutes at RT. 15 μL of LPS (13.33 ug/ml stock) was then added to the cell suspension and the plate was incubated for 18 hours at 37°C in a humidified atmosphere containing 5% CO_2. Following incubation, the culture medium was collected and stored at -20°C.

THP-1 cells (TIB-202, ATCC) were washed and resuspended at a concentration of 1 x 10^5/ml in assay medium (RPMI medium containing 3% fetal bovine serum). 175 μL of cell suspension was incubated with 10 μL of test compound (in 4% DMSO) in 96-well tissue culture plates for 30 minutes at RT. 15 μL of LPS (13.33 ug/ml stock) was then added to the cell suspension and the plate was incubated for 18 hours at 37°C in a humidified atmosphere containing 5% CO_2. Following incubation, the culture medium was collected and stored at -20°C.

TNF-α concentration in the medium was quantified using a standard ELISA kit (BioSource International, Camarillo, CA). Concentrations of TNF-α and IC_{50} values for test compounds (concentration of compound that inhibited LPS-stimulated TNF-α production by 50%) were calculated by four parameter logistic curve (SigmaPlot, SPSS, Inc.).

p38α Assay

The p38α assay employed is based on measurement of ADP released in the reaction of interest through NADH oxidation obtained by coupling with pyruvate kinase and lactate dehydrogenase reactions. The assays were performed in 384-well UV-plates. The final volume was 25 μL prepared from the addition of 2.5 μL compound dissolved in 10% DMSO, 17.5 μL of assay buffer and 5 μL of ATP. Assay buffer contains the following reagents to give final concentration in the assay: 25 mM HEPES, 20 mM 2-glycerophosphate, pH 7.6, 10 mM $MgCl_2$, 0.1 mM sodium orthovanadate, 0.5 mM phosphoenolpyruvate, 0.12 mM NADH, 3.1 mg/ml LDH, 6.67 mg/ml pyruvate kinase, 0.25 mM peptide substrate, 2 mM DTT, 0.005% Tween 80 and 20 nM p38α kinase from Upstate. Test compounds are preincubated with p38α kinase for 60 min and the reaction started by addition of ATP to 0.15 mM final concentration. Reaction rates were measured at 340 nm using SpectraMax plate-reading spectrophotometer for 10 min at 37°C. Inhibition data were analyzed by non-linear least-squares regression using SigmaPlot.

TNF-α Production by LPS-Stimulated Mice

Mice (Balb/c female, 6-8 weeks of age, Taconic Labs; n=8/treatment group) were injected intraperitoneally with lipopolysaccharide (LPS) (50 ug/kg of *E. coli* strain 0111:B4, Sigma) suspended in sterile saline. Ninety minutes later, mice were

sedated by CO_2:O_2 inhalation and a blood sample was obtained. Serum was separated and analyzed for TNF-α concentrations by commercial ELISA assay per the manufacturer's instructions (BioSource International). Test compounds were administered orally at various times before LPS injection. The compounds were dosed either as suspensions or as solutions in various vehicles or solubilizing agents.

Results

The compounds exemplified herein all showed activity in the above assays as inhibitors of p38 kinase. The p38 inhibitory activity of certain compounds provided herein are shown in the Table below. For p38 kinase IC_{50} values, "+++" represents < 1 μM, "++" represents between 1.0 and 10 μM and "+" represents > 10 μM.


Example	**p38α IC_{50}**
1	+++
2	+++
3	++
4	+++
5	++
6	+++
7	+++
8	+++
9	+++
10	+++
18	+++
29	+++
30	+++
33	+++
34	+++
48	+++
49	+++
50	+++
52	+++

Example	p38α IC_{50}
53	+++
76	+++
87	+++
128	+++
129	+++
135	+++
148	+++
171	+++

Since modifications will be apparent to those of skill in the art, the claimed subject matter is intended to be limited only by the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound of the formula:



or a pharmaceutically acceptable derivative thereof, wherein:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, optionally substituted alkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted alkoxy, optionally substituted heterocyclyloxy, or alkylamino;

G is an aryl, aralkyl, cycloalkyl, heteroaryl, heteroaralkyl or a heterocyclyl ring optionally fused to a phenyl ring, and is substituted with R^3 and R^4, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom, or G is OR^{83} or $NR^{80}R^{81}$;

B is an aryl or heteroaryl ring;

C is a 5-membered heteroaryl ring containing one or two heteroatoms in the ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$;

each R^{80} and R^{81} is independently hydrogen, alkyl, optionally substituted cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl, or together from optionally substituted alkylene or heteroalkylene;

R^{83} is hydrogen, alkyl, cycloalkyl, heteroaryl or optionally substituted heteroaryl;

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n-$ (where n is an integer from 0 to 2); and R^9 is halo, cyano, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(w) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) $-OC(O)$-alkylene-CO_2H or -OC(O)-NR'R" where R' and R" are independently hydrogen or alkyl;

(bb) heteroarylalkenylene or heteroarylalkynylene;

(cc) hydrogen;

(dd) halo;

(ee) pseudohalo;

(ff) hydroxy;

(gg) optionally substituted alkoxy;

(hh) $C(L)R^{40}$, where L is O, S or NR^{55}; R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{56}$, halo pseudohalo, OR^{55}, SR^{55}, $NR^{57}R^{58}$ or $SiR^{52}R^{53}R^{54}$; where R^{52}, R^{53} and R^{54} are selected as in (i) or (ii) as follows (i) R^{52}, R^{53} and R^{54} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or (ii) any two of R^{52}, R^{53} and R^{54} together form alkylene, alkenylene, alkynylene, heteroalkylene; and the other is selected as in (i); R^{55} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{56} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{64}R^{65}$; where R^{64} and R^{65} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{66} or $NR^{62}R^{63}$, or R^{64} and R^{65} together form alkylene, alkenylene, alkynylene, heteroalkylene, where R^{66} is

hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{57} and R^{58} are selected as in (i) or (ii) as follows (i) R^{57} and R^{58} are each independently hydrogen, optionally substituted alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55}, $NR^{67}R^{68}$ or $C(L)R^{69}$, where R^{67} and R^{68} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl, or together form alkylene, alkenylene, alkynylene, heteroalkylene; and R^{69} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$, where R^{70} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl; or (ii) R^{57} and R^{58} together form alkylene, alkenylene, alkynylene, heteroalkylene, or alkylenoxyalkylene; R^{62} and R^{63} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, or R^{62} and R^{63} together form alkylene, alkenylene, alkynylene, heteroalkylene; and

(ii) optionally substituted alkyl;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

or R^3 and R^4, which substitute adjacent atoms on a ring, together form alkylenedioxy, thioalkylenoxy or alkylenedithioxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(l) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy.

2. The compound of claim 1, wherein G is OR^{83} or $NR^{80}R^{81}$.

3. The compound of claim 1 or claim 2, wherein, R^{83} is alkyl or cycloalkyl.

4. The compound of any of claims 1-3, wherein R^{83} is alkyl.

5. The compound of any of claims 1-4, wherein R^{83} is ethyl.

6. The compound of claim 1 or claim 2, wherein R^{80} and R^{81} are each independently hydrogen, optionally substituted alkyl or optionally substituted cycloalkyl, or together from optionally substituted alkylene or optionally substituted heteroalkylene.

7. The compound of any of claims 1, 2 and 6, wherein R^{80} and R^{81} are each independently hydrogen or optionally substituted cycloalkyl.

8. The compound of any of claims 1, 2, 6 and 7, wherein R^{80} and R^{81} are each independently hydrogen, cyclohexyl, 2-methylcyclohexyl, 3-pentyl, bicyclo[2.2.1]heptyl, 1-ethynylcyclohexyl, methoxybenzyl, benzyl, indanyl, cyclopropyl, 3-chlorobenzyl, cyclopentyl, 2,4-dichlorobenzyl, cyclohexylmethyl, 3,4-dichlorobenzyl, 4-methylcyclohexyl, 3-trifluoromethylbenzyl, 4-tert-butylcyclohexyl, neopentyl, fluorobenzyl, N-ethyl-2-pyrrolydinylmethyl, methylbenzyl, pyridylmethyl, morpholinylethyl, 1-cyclohexylethyl, 1-phenylethyl or 2-pyrrolidinylethyl, or R^{80} and R^{81} together form pentylene or 2,5-hexylene.


9. The compound of any of claims 1, 2 and 6-8, wherein G is NH_2 or NH(cyclohexyl).

10. The compound of claim 1, wherein G is aryl, heteroaryl, cycloalkyl, heterocyclyl or heterocyclyl optionally fused to phenyl, and is substituted with R^3 and R^4, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom.

11. The compound of claim 1 or claim 10, wherein G is phenyl, pyridyl, cyclohexyl, cyclopentyl or benzyl, and is substituted with R^3 and R^4.

12. The compound of any of claims 1, 10 or 11, wherein G is phenyl and is substituted with R^3 and R^4.

13. The compound of any of claims 1 or 10-12 that has formula (I):



I

wherein:

A is an aryl, heteroaryl, cycloalkyl or a heterocyclyl ring optionally fused to a phenyl ring, provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom.


14. The compound of any of claims 1-13, wherein C is selected from pyrazole, imidazole, pyrrole, thiazole, isothiazole, oxazole, isoxazole, furan and thiophene rings.

15. The compound of any of claims 1-14, wherein C is a pyrazole or imidazole ring.

16. The compound of any one of claims 1-15, wherein C is an imidazole ring.

17. The compound of any one of claims 1-16, wherein C is a pyrazole ring.


18. The compound of any one of claims 1-15 and 17, wherein compound has formula II:



II

or a pharmaceutically acceptable derivative thereof, wherein the ring NH group may be replaced by one of the -C(O)-A(R^3)(R^4), -R^2, or -B(D)(R^6)(R^5) substituents.


19. The compound of any one of claims 1-15, 17 and 18, wherein the compound has formula III:



III

or a pharmaceutically acceptable derivative thereof.


20. The compound of any one of claims 1-15, 17 and 18, wherein the compound has formula IV:



IV

or a pharmaceutically acceptable derivative thereof.


21. The compound of any one of claims 1-15, 17 and 18, wherein the compound has formula V:



V

or a pharmaceutically acceptable derivative thereof.


22. The compound of any one of claims 1-15, 17 and 18, wherein the compound has formula Va:



Va

or a pharmaceutically acceptable derivative thereof.


23. The compound of any one of claims 1-16, wherein the compound has formula VI:



VI

or a pharmaceutically acceptable derivative thereof, wherein the ring NH group may be replaced by one of the -C(O)-A(R^3)(R^4), -R^2, or -B(D)(R^6)(R^5) substituents.


24. The compound of any one of claims 1-16 and 23, wherein the compound has formula VII:



VII

or a pharmaceutically acceptable derivative thereof.

25. The compound of any one of claims 1-16 and 23, wherein the compound has formula VIII:



VIII

or a pharmaceutically acceptable derivative thereof.

26. The compound of any one of claims 1-25, wherein R^1 is hydrogen.

27. The compound of any one of claims 1-26, wherein R^2 is hydrogen or lower alkyl.

28. The compound of any one of claims 1-27, wherein R^2 is hydrogen, methyl, n-propyl or 3-chlorobenzylaminocarbonylmethoxy.

29. The compound of any one of claims 1-28, wherein A is an aryl ring.

30. The compound of any one of claims 1-29, wherein A is a phenyl ring.

31. The compound of any one of claims 1-30, wherein B is an aryl ring.

32. The compound of any one of claims 1-31, wherein B is a phenyl ring.

33. The compound of any one of claims 1-32, wherein D is $-C(O)NR^{80}R^{81}$.

34. The compound of any one of claims 1-33, wherein R^{80} and R^{81} are each independently hydrogen, cycloalkyl or alkoxy.

35. The compound of any one of claims 1-34, wherein R^{80} is hydrogen.

36. The compound of any one of claims 1-35, wherein R^{81} is cycloalkyl or alkoxy.

37. The compound of any one of claims 1-36, wherein R^{81} is C_{3-6}cycloalkyl or C_{1-6}alkoxy.

38. The compound of any one of claims 1-37, wherein R^{81} is cyclopropyl or methoxy.

39. The compound of any one of claims 1-32, wherein D is optionally substituted heteroaryl.

40. The compound of any one of claims 1-32 and 39, wherein D is optionally substituted triazolyl.

41. The compound of any one of claims 1-32, 39 and 40, wherein D is 1,2,4-triazol-3-yl.

42. The compound of any one of claims 1-41, wherein R^3 is hydrogen, optionally substituted heterocyclyl, optionally substituted alkyl, $C(L)R^{40}$, halo, pseudohalo or OR^{41}; where L is O, S or NR^{55}; R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{56}$, halo pseudohalo, OR^{55}, SR^{55}, $NR^{57}R^{58}$ or $SiR^{52}R^{53}R^{54}$; R^{41} is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heteroarylium, optionally substituted cycloalkyl, optionally substituted heterocyclyl, $C(L)R^{59}$, $NR^{60}R^{61}$ or $SiR^{52}R^{53}R^{54}$; where R^{52}, R^{53} and R^{54} are selected as in (i) or (ii) as follows (i) R^{52}, R^{53} and R^{54} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or (ii) any two of R^{52}, R^{53} and R^{54} together form alkylene, alkenylene, alkynylene, heteroalkylene; and the other is selected as in (i); R^{55} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{56} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{64}R^{65}$; where R^{64} and R^{65} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{66} or

$NR^{62}R^{63}$, or R^{64} and R^{65} together form alkylene, alkenylene, alkynylene, heteroalkylene, where R^{66} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl; R^{57} and R^{58} are selected as in (i) or (ii) as follows (i) R^{57} and R^{58} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55}, $NR^{67}R^{68}$ or $C(L)R^{69}$, where R^{67} and R^{68} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl or heterocyclyl, or together form alkylene, alkenylene, alkynylene, heteroalkylene; and R^{69} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$, where R^{70} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl; or (ii) R^{57} and R^{58} together form alkylene, alkenylene, alkynylene, heteroalkylene; R^{59} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{70} or $NR^{62}R^{63}$; R^{60} and R^{61} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl or $C(L)R^{71}$, where R^{71} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, OR^{55} or $NR^{62}R^{63}$; or R^{60} and R^{61} together form alkylene, alkenylene, alkynylene, heteroalkylene; R^{62} and R^{63} are each independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heteroarylium, cycloalkyl, heterocyclyl, or R^{62} and R^{63} together form alkylene, alkenylene, alkynylene, heteroalkylene.

43. The compound of any one of claims 1-42, wherein R^3 is hydrogen, optionally substituted heterocyclyl, optionally substituted alkyl, $C(L)R^{40}$, halo, cyano, optionally substituted heteroaryl, $S(O)_nR^{27}$ or OR^{41}.

44. The compound of any one of claims 1-43, wherein R^3 is hydrogen, optionally substituted heterocyclyl, cyano, optionally substituted heteroaryl, optionally substituted alkyl, $C(L)R^{40}$, iodo, chloro, fluoro, $S(O)_nR^{27}$ or OR^{41}.

45. The compound of any one of claims 1-44, wherein R^3 is hydrogen, optionally substituted dioxolanyl, pyrimidinyl, pyridyl, pyrazinyl, cyano, oxadiazolyl, triazolyl, methylthio, methanesulfonyl, optionally substituted methyl, $C(L)R^{40}$, iodo, chloro or OR^{41}.

46. The compound of any one of claims 1-45, wherein R^3 is hydrogen, 2-dioxolanyl, pyrimidinyl, pyridyl, pyrazinyl, cyano, oxadiazolyl, triazolyl, methylthio,

methanesulfonyl, optionally substituted methyl, $C(O)R^{40}$, iodo, chloro or optionally substituted methoxy.

47. The compound of any one of claims 1-46, wherein R^3 is hydrogen, 2-dioxolanyl, pyrimidinyl, pyridyl, pyrazinyl, cyano, oxadiazolyl, triazolyl, methylthio, methanesulfonyl, 5-methyloxadiazolyl, methyl, CHO, iodo, chloro, methoxy, benzyloxy, ethoxy, cyclopropoxy, t-butoxycarbonylbutoxy or carbamoylmethoxy.

48. The compound of any one of claims 1-47, wherein R^3 is optionally substituted methyl.

49. The compound of any one of claims 1-48, wherein R^3 is methyl which is optionally substituted with heterocyclyl, hydroxy, aralkylamino or heterocyclylalkylamino.

50. The compound of any one of claims 1-49, wherein R^3 is *N*-morpholinylmethyl, hydroxymethyl, *N*-(2-(3-chlorophenyl)-1-ethyl)aminomethyl, *N*-(2-morpholinyl-1-ethyl)aminomethyl, (3-chloro-benzylamino)-methyl, 4-(3-chloro-phenyl)-piperazin-1-ylmethyl, [2-(2-benzyloxy-5-chloro-phenyl)-ethylamino]-methyl, (3-morpholin-4-yl-propylamino)-methyl, carbamoylmethoxy, 2-(4-methyl-piperazin-1-yl)-2-oxo-ethoxy, 2-oxo-2-piperazin-1-yl-ethoxy, 2-(3-amino-pyrrolidin-1-yl)-2-oxo-ethoxy, 2-(3-methylamino-pyrrolidin-1-yl)-2-oxo-ethoxy, 2-(3,5-dimethyl-piperazin-1-yl)-2-oxo-ethoxy, 2-morpholin-4-yl-2-oxo-ethoxy, [(1H-benzoimidazol-2-ylmethyl)-carbamoyl]-methoxy, [2-(2-benzyloxy-5-chloro-phenyl)-ethylcarbamoyl]-methoxy, [2-(5-chloro-2-hydroxy-phenyl)-ethylcarbamoyl]-methoxy or 4-piperizinylmethyl.

51. The compound of any one of claims 1-50, wherein L is O.

52. The compound of any one of claims 1-51, wherein R^{40} is hydrogen, optionally substituted alkyl, optionally substituted alkoxy, $NR^{57}R^{58}$ or cycloalkyl.

53. The compound of any one of claims 1-52, wherein R^{40} is hydrogen, alkyl, alkoxy or $NR^{57}R^{58}$.

54. The compound of any one of claims 1-53, wherein R^{40} is hydrogen, cyclopropylamino, NH2, isopropylamino, methylamino, piperazinyl, dimethylamino, cyclopropylmethylamino, ethylamino, cyclopentylamino or tert-butoxy.

55. The compound of any one of claims 1-54, wherein R^{41} is hydrogen, optionally substituted alkenyl, or optionally substituted alkyl.

56. The compound of any one of claims 1-55, wherein R^{41} is hydrogen,

or alkyl or alkenyl each optionally substituted with heterocyclyl, aryl, heteroaryl, dialkylamino, halo or hydroxy.

57. The compound of any one of claims 1-56, wherein R^{41} is hydrogen, or C_{1-3}alkyl or C_{1-3}alkenyl each optionally substituted with heterocyclyl, phenyl, heteroaryl, dialkylamino, halo or hydroxy.

58. The compound of any one of claims 1-57, wherein R^{41} is hydrogen, 2-(N-morpholinyl)eth-1-yl, benzyl, 2-(*N,N*-di-(2-hydroxy-1-ethyl)amino)-1-ethyl, 2-bromo-1-ethyl, 2,2-dioxolan-4-ylmethyl, 2-(4-methylpiperazin-1-yl)-1-ethyl, oxadiazolylmethyl, ethenyl, 2-dimethylaminoethyl, 4-chlorophenoxyethyl, or 2,3-dihydroxy-1-propyl.

59. The compound of any one of claims 1-58, wherein R^{41} is (*S*)-2,3-dihydroxy-1-propyl.

60. The compound of any one of claims 1-59, wherein R^4 is hydrogen.

61. The compound of any one of claims 1-60, wherein R^5 is alkyl.

62. The compound of any one of claims 1-61, wherein R^5 is methyl.

63. The compound of any one of claims 1-62, wherein R^6 is hydrogen.

64. The compound of any one of claims 1-25, wherein:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$ (where R^{80} and R^{81} are independently hydrogen, alkyl, cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl);

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n$ (where n is an integer from 0 to 2); and R^9 is cyano, optionally substituted heteroaryl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$- where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) $-OC(O)$-alkylene-CO_2H or -OC(O)-NR'R" (where R' and R" are independently hydrogen or alkyl); and

(bb) heteroarylalkenylene or heteroarylalkynylene;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(l) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy;

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

65. The compound of any one of claims 1-25, wherein:

R^1 is hydrogen or acyl;

R^2 is hydrogen or alkyl; and

A is an aryl or heteroaryl ring.

66. The compound of any one of claims 1-25, wherein:

R^1 is hydrogen, acyl or $-P(O)(OH)_2$;

R^2 is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

R^3 is selected from the group consisting of:

(a) amino;

(b) acylamino;

(c) optionally substituted heterocycle;

(d) heteroaryl optionally substituted with a substituent selected from halo, alkyl or alkoxy;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted heterocyclylalkoxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n-$ (where n is an integer from 0 to 2); and R^9 is cyano, heteroaryl, -COOH, $-COR^{10}$, $-COOR^{11}$ $-CONR^{12}R^{13}$, $-SO_2R^{14}$,$-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$ where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl,

and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(t) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$ where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -0- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26,}$ where R^{25} and R25 independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy;

R^5 is selected from the group consisting of:

(n)hydrogen;

(o) halo;

(p) alkyl;

(q) haloalkyl;

(r) thioalkyl;

(s) hydroxy;

(t) amino;

(u) alkylamino;

(v) dialkylamino;

(w) heteroalkyl;

(x) optionally substituted heterocycle;

(y) optionally substituted heterocyclylalkyl; and

(z) optionally substituted heterocyclylalkoxy;

R^6 is selected from a group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy.

67. The compound of any one of claims 1-25, wherein R^3 is:

(a) optionally substituted heterocyclyl;

(b) aryl or heteroaryl both optionally substituted with a substituent selected from halo, alkyl, amino, alkoxy, carboxy, lower alkoxy carbonyl, SO_2R' (where R' is alkyl) or $-O_2NHR'R''$ (where R' and R'' are independently hydrogen or alkyl);

(c) heteroalkyl;

(d) heteroalkenyl;

(e) heteroalkylamino;

(f) heteraloxy

(g) optionally substituted heterocyclylalkyl or heterocyclyloxy;

(h) optionally substituted heterocyclylalkenyl;

(i) optionally substituted heterocyclylalkynyl;

(j) optionally substituted heterocyclylalkoxy;

(k) optionally substituted heterocyclylalkylamino;

(l) optionally substituted heterocyclylalkylcarbonyl:

(s) -Y-(alkylene)-R^9 where Y is a single bond, -O or -NH- and R^9 is optionally substituted heteroaryl, $-CONR^{12}R^{13}$, SO_2R^{14}, $-SO_2NR^{15}R^{16}$ $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$ where R^{12}, R^{13}, R^{14}, R^{15}, R^{16} R^{17}, R^{18} and R^{19} are independently of each other heteroalkyl; hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(m) arylaminoalkylene or heteroarylaminoalkylene;

or

(n) Z-alkylene-$NR^{30}R^{31}$ where Z is -NH-, -N(alkyl)- or -0-, and R^{30} and R^{31} are independently of each other, hydrogen, alkyl or heteroalkyl.

68. The compound of any one of claims 1-25, wherein R^1 and R^2 are hydrogen and B is phenyl.

69. The compound of any one of claims 1-25, wherein R^4 is hydrogen and R^5 is halo or alkyl.

70. The compound of any one of claims 1-25, wherein R^5 is chloro, fluoro or methyl and R^6 is hydrogen, chloro, fluoro, methyl or methoxy.

71. The compound of any one of claims 1-25, wherein R^3 is optionally substituted heteroaryl.

72. The compound of any one of claims 1-25, wherein R^3 is pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxidopyridin-2-yl, N-oxidopyridin-3-yl, Noxidopyridin-4-yl or pyridon-2-yl, all optionally substituted.

73. The compound of any one of claims 1-72, wherein R^3 is at the 3-position.

74. The compound of any one of claims 1-73, wherein R^5 is 4-F or 2-Me, and R^6 is hydrogen.

75. The compound of any one of claims 1-25, wherein R^3 is optionally substituted phenyl.

76. The compound of any one of claims 1-25, wherein R^3 is 3-sulfamoylphenyl, 3-methylsulfonylphenyl, 3-carboxyphenyl or 3-ethoxycarbonylphenyl.

77. The compound of any one of claims 1-25, wherein R^5 is 4-F and R^6 is hydrogen.

78. The compound of any one of claims 1-25, wherein R^3 is:

(a) heteroalkyl;

(b) heteroalkoxy;

(c) heteroalkylamino;

(d) optionally substituted heterocyclylalkyl;

(e) optionally substituted heterocyclylalkoxy;

(f) optionally substituted heterocyclylalkylamino;

(g) Y-(alkylene)-R^9 where Y is a single bond, -O- or –NH- and R^9 is optionally substituted heteroaryl, -$CONR^{12}R^{13}$, -SO_2R^{14}, -$SO_2NR^{15}R^{16}$ -$NHSO_2R^{17}$ or -$NHSO_2NR^{18}R^{19}$ where R_{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are independently of each other hydrogen, alkyl or heteroalkyl; or

(h) Z-alkylene-$NR^{30}R^{31}$ where Z is -NH-, -N(alkyl)- or -O-, and R^{30} and R^{31} are independently of each other, hydrogen, alkyl or heteroalkyl.

79. The compound of any one of claims 1-25, wherein R^3 is heteroalkyl.

80. The compound of any one of claims 1-25, wherein R^3 is at the 3-position and is selected from the group consisting of 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino, hydroxymethyl, 1,2-dihydroxyethyl, 3-hydroxy-3-methyl-l-butyl or 3-hydroxybutyl.

81. The compound of any one of claims 1-25, wherein R^5 is 2-F and R^6 is 4-F.

82. The compound of any one of claims 1-25, wherein R^5 is 2-Me and R^6 is hydrogen.

83. The compound of any one of claims 1-25, wherein R^9 is heteroalkoxy or heteroalkylamino.

84. The compound of any one of claims 1-25, wherein R^3 is at the 3-position and is selected from the group consisting of 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, 2-hydroxyethoxy, 2,3-dihydroxypropoxy, 2-dimethylaminoethylamino and 3-dimethylaminopropylamino.

85. The compound of any one of claims 1-25, wherein R^3 is optionally substituted heterocyclylalkyl, optionally substituted heterocyclylalkoxy or optionally substituted heterocyclylalkylamino.

86. The compound of any one of claims 1-25, wherein R^3 is at the 3-position and is selected from the group consisting of 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 2-(2-oxo-pyrrolidin-1-yl)ethoxy, 3(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4yl)ethylamino, 4-hydroxypiperidinylmethyl, 2-(S,S-dioxo-thiamorpholin-4-yl)ethyl, 3-(S,S-dioxo-thiamorpholin-4-yl)propyl and N-methylpiperazinylmethyl.

87. The compound of any one of claims 1-25, wherein R^3 is -Y-(alkylene)-R^9 where Y is a single bond, -O- or -NH- and R^9 is optionally substituted heteroaryl, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$ where R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are independently of each other hydrogen, alkyl or heteroalkyl.

88. The compound of any one of claims 1-25, wherein Y is a single bond and R^9 is $-SO_2R^{14}$ or $-SO_2NR^{15}R^{16}$.

89. The compound of any one of claims 1-25, wherein R^3 is 5-methylsulfonylethyl or sulfamoylethyl.

90. The compound of any one of claims 1-89 selected from 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone; 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-

hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; and 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

91. The compound of any one of claims 1-90 selected from 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone; 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; or 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

92. The compound of any one of claims 1-91 that is 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

93. The compound of any one of claims 1-89 selected from 5-amino-l-(4-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole, 5-amino-l-(2,4-difluorophenyl)-4-[3-(3-morpholin-4ylpropyl)benzoyl]pyrazole, 5-amino-4-(3-aminobenzoyl)-1-(4-fluorophenyl)pyrazole, 5-amino-l-(4-fluorophenyl)-4- [3-(3-morpholin-4-ylpropyl)benzoyl] pyrazole, 5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl] -1-(4fluorophenyl)pyrazole, 5-amino-4-(3-acetylaminobenzoyl)-1-phenylpyrazole, 5-amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)pyrazole, 5-amino-l-(4-fluorophenyl)-4- [3-(3-morpholin-4-ylpropylamino) benzoyl] pyrazole, 5-amino-4-[3-(2-aminosulfonylethyl)benzoyl]-1-

(4-fluorophenyl)pyrazole and 5-amino-l-(4-fluorophenyl)-4-(3-pyridin-3-ylbenzoyl)pyrazole.

94. The compound of any one of claims 1-89 selected from 5-amino-l-(2-methylphenyl)-4-[3-pyridin-3-yl)benzoyl]pyrazole, 5-amino-l-(2-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]pyrazole, 5-amino-4-[3-(2,3-dihydroxypropoxy)benzoyl]-1-(4-fluorophenyl) pyrazole, 5-amino-4-[3-(1,2-dihydroxyethyl)benzoyl]-1-(4-fluorophenyl) pyrazole, 5-amino-l-(4-fluorophenyl)-4-[3-(sulfamoylbenzoyl] pyrazole, 5-amino-l-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methylbutyl) benzoyl]pyrazole, 5-amino-1-(4-fluorophenyl)-4-[3-(2-(1-hydroxycyclopentyl)ethyl) benzoyl]pyrazole, 5-amino-4-[3-(2-methylsulfonylethyl)benzoyl]-1-(4-fluorophenyl) pyrazole, and 5-amino-l-(2,4-difluorophenyl)-4-[3-(2-hydroxyethylsulfonyl) benzoyl]pyrazole.

95. The compound of any one of claims 1-89 selected from 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid; 3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid; 3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; {5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone; 3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; 3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; and 3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

96. The compound of any one of claims 1-89 selected from:

3-{5-Amino-4-[3-(2-dimethylamino-ethylcarbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(5-chloro-thiophene-2-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-hydrazinocarbonyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclohexanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclopentanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-phenylacetyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide; and

3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide.

97. The compound of any one of claims 1-89 selected from:

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester;

3-(5-Amino-4-cyclopentanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-hydrazinocarbonyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid benzylamide;

3-(5-Amino-4-cyclohexanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; and

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylamide.

98. The compound of any one of claims 1-89 selected from:
5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide;
5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide;
5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester;
5-Amino-3-[(3-chloro-benzylcarbamoyl)-methoxy]-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester;
3-[5-Amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;
3-(5-Amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;
3-(5-Amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;
5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-(2-hydroxy-ethoxy)-1H-pyrazole-4-carboxylic acid ethyl ester;
4-[5-Amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;
3-(5-Amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide; and
3-[5-Amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide.

99. The compound of any one of claims 1-89 selected from:
3-[5-Amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-methoxy-4-methyl-benzamide;
3-(5-Amino-4-benzoyl-pyrazol-1-yl)-N-methoxy-4-methyl-benzamide;
3-(5-amino-4-benzoyl-pyrazol-1-yl)-4-methyl-benzoic acid;
3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;
3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-4-methyl-benzoic acid;
3-[5-amino-4-(3-iodo-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;
{5-amino-1-[2-methyl-5-(4H-[1,2,4]triazol-3-yl)-phenyl]-1H-pyrazol-4-yl}-phenyl-methanone;

3-[5-amino-4-(3-[1,3]dioxolan-2-yl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-formyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-hydroxymethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-amino-4-[3-(4-methyl-piperazin-1-ylmethyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-morpholin-4-ylmethyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(2-morpholin-4-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-benzyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(3-hydroxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-(4-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclohexanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-cyclopentanecarbonyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-Amino-4-phenylacetyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(2-dimethylamino-ethylcarbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-methylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(3-cyclopentylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(morpholine-4-carbonyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-{5-Amino-4-[3-(cyclopropylmethyl-carbamoyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide;

3-[5-Amino-4-(tetrahydro-pyran-4-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-ethoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-benzoyl-3-(2-methoxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

3-[5-amino-4-benzoyl-3-(2-benzyloxy-ethoxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide;

4-[5-amino-4-benzoyl-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazol-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

3-[5-amino-4-benzoyl-3-(piperidin-4-yloxy)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide, trifluoroacetate salt;

3-(5-amino-4-benzoyl-3-methylsulfanyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

3-(5-amino-4-benzoyl-3-methanesulfonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide;

5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid amide;

5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methanesulfonyl-1H-pyrazole-4-carboxylic acid amide; and

5-amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methylsulfanyl-1H-pyrazole-4-carboxylic acid ethyl ester.

100. The compound of any one of claims 1-89 selected from:

3-[5-Amino-4-(3-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methyl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(2-methylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(2-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-chlorobenzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(2-chlorobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methoxy-benzoyl)-pyrazol-1-yl]-4,N-dimethyl-benzamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester

5-Amino-3-[(3-chloro-benzylcarbamoyl)-methoxy]-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid ethyl ester

3-(5-amino-4-benzoyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(3-[1,3,4]oxadiazol-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-{5-Amino-4-[3-(5-methyl-[1,3,4]oxadiazol-2-yl)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(pyrrolidine-1-carbonyl)-benzoyl]-imidazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-carbamoyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-isopropylcarbamoylbenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(4-methylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-cyclopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-carbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(piperazine-1-carbonyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-dimethylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(cyclopropylmethyl-carbamoyl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-ethylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-cyclopentylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-isopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-cyclopropylcarbamoyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[(3-chloro-benzylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{[2-(3-chloro-phenyl)-ethylamino]-methyl}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[4-(3-chloro-phenyl)-piperazin-1-ylmethyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{[2-(2-benzyloxy-5-chloro-phenyl)-ethylamino]-methyl}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[(2-morpholin-4-yl-ethylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[(3-morpholin-4-yl-propylamino)-methyl]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(pyridine-2-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(2-methyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,4-difluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-fluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,4-dichloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-fluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,5-dichloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-methoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-benzoic acid tert-butyl ester

3-[5-Amino-4-(3,5-difluoro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(benzo[1,3]dioxole-5-carbonyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-chloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,4-dimethoxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-benzyloxy-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-fluoro-3-methyl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methylbenzamide

{3-[5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-imidazole-4-carbonyl]-phenoxy}-acetic acid tert-butyl ester

3-[5-Amino-4-(3-chloro-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-benzoyl-2-methyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-benzoyl-2-propyl-imidazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-carbamoylmethoxy-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-(5-Amino-4-{3-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(2-oxo-2-piperazin-1-yl-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[2-(3-amino-pyrrolidin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[2-(3-methylamino-pyrrolidin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-{3-[2-(3,5-dimethyl-piperazin-1-yl)-2-oxo-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(2-morpholin-4-yl-2-oxo-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{[(1H-benzoimidazol-2-ylmethyl)-carbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{[2-(2-benzyloxy-5-chloro-phenyl)-ethylcarbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{[2-(5-chloro-2-hydroxy-phenyl)-ethylcarbamoyl]-methoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-pyrazin-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(3-pyridin-2-yl-benzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(pyridine-2-carbonyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-cyclopentanecarbonyl-pyrazol-1-yl)-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(benzo[1,3]dioxole-5-carbonyl)-pyrazol-1-yl]-N-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(2-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-ethoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-cyclopropoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-methoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,4-dimethoxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-methylsulfanyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-methylsulfanyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,4-difluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3,5-difluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-fluoro-3-methyl-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(4-fluoro-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-(5-Amino-4-cyclohexanecarbonyl-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-vinyloxy-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-([1,3,4]oxadiazol-2-ylmethoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-methyl-cyclohexyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid phenylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethyl-propyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid bicyclo[2.2.1]hept-2-ylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethy

3-[5-Amino-4-(2,5-dimethyl-pyrrolidine-1-carbonyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 4-methoxy-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid indan-1-ylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid benzylamide

3-[5-Amino-4-(piperidine-1-carbonyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclopropylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-chlorobenzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclopentylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 2,4-dichloro-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexylmethyl-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3,4-dichloro-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (4-methyl-cyclohexyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-trifluoromethyl-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (4-tert-butyl-cyclohexyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2,2-dimethyl-propyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 3-methoxy-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 4-fluoro-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethyl-pyrrolidin-2-ylmethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid 2-methyl-benzylamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-morpholin-4-yl-ethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-cyclohexyl-ethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-phenyl-ethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2,2-dimethyl-propyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (1-ethyl-pyrrolidin-2-ylmethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methyl-phenyl)-1H-pyrazole-4-carboxylic acid cyclohexyl-methyl-amide

3-[5-Amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(3-pyrazin-2-yl-benzoyl)-imidazol-1-yl]-*N*-cyclopropyl-4-methylbenzamide

3-[5-Amino-4-(3-pyrimidin-5-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-pyrimidin-5-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-pyrimidin-2-yl-benzoyl)-imidazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

5-Amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-methylsulfonylbenzoyl-1H-imidazole

5-Amino-1-(5-cyclopropylcarbamoyl-2-methylphenyl)-4-[3-(5-methyl)oxadiazol-3-ylbenzoly]-1H-imidazole

3-(5-Amino-4-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-[5-Amino-4-(3-{2-[bis-(2-hydroxy-ethyl)-amino]-ethoxy}-benzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(2-dimethylamino-ethoxy)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(2,3-dihydroxy-propoxy)-benzoyl]-pyrazol-1-yl}-*N*-cyclopropyl-4-methylbenzamide

3-(5-Amino-4-{3-[2-(4-chloro-phenoxy)-ethoxy]-benzoyl}-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(4H-[1,2,4]triazol-3-yl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide

3-{5-Amino-4-[3-(4H-[1,2,4]triazol-3-yl)-benzoyl]-pyrazol-1-yl}-N-cyclopropyl-4-methyl-benzamide.

101. The compound of any one of claims 1-25, wherein:

R^1 is hydrogen or acyl;

R^2 is hydrogen or alkyl;

A is an aryl or heteroaryl ring;

B is an aryl or heteroaryl ring;

D is heteroaryl, optionally substituted heteroaryl or $-C(O)NR^{80}R^{81}$ (where R^{80} and R^{81} are independently hydrogen, alkyl, cycloalkyl, alkoxy, hydroxy, heteroaryl or optionally substituted heteroaryl);

R^3 is selected from the group consisting of:

(a) amino, alkylamino or dialkylamino;

(b) acylamino;

(c) optionally substituted heterocyclyl;

(d) optionally substituted aryl or heteroaryl;

(e) heteroalkyl;

(f) heteroalkenyl;

(g) heteroalkynyl;

(h) heteroalkoxy;

(i) heteroalkylamino;

(j) optionally substituted heterocyclylalkyl;

(k) optionally substituted heterocyclylalkenyl;

(l) optionally substituted heterocyclylalkynyl;

(m) optionally substituted cycloalkoxy, cycloalkylalkyloxy, heterocyclylalkoxy or heterocyclyloxy;

(n) optionally substituted heterocyclylalkylamino;

(o) optionally substituted heterocyclylalkylcarbonyl;

(p) heteroalkylcarbonyl;

(q) $-NHSO_2R^6$ where R^6 is alkyl, heteroalkyl or optionally substituted heterocyclylalkyl;

(r) $-NHSO_2NR^7R^8$ where R^7 and R^8 are, independently of each other, hydrogen, alkyl or heteroalkyl;

(s) -Y-(alkylene)-R^9 where: Y is a single bond, -O-, -NH- or $-S(O)_n$ (where n is an integer from 0 to 2); and R^9 is cyano, optionally substituted heteroaryl, -COOH, $-COR^{10}$, $-COOR^{11}$, $-CONR^{12}R^{13}$, $-SO_2R^{14}$, $-SO_2NR^{15}R^{16}$, $-NHSO_2R^{17}$ or $-NHSO_2NR^{18}R^{19}$, where R^{10} is alkyl or optionally substituted heterocycle, R^{11} is alkyl, and R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18} and R^{19} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(y) $-C(=NR^{20})(NR^{21}R^{22})$ where R^{20}, R^{21} and R^{22} independently represent hydrogen, alkyl or hydroxy, or R^{20} and R^{21} together are $-(CH_2)_n-$- where n is 2 or 3 and R^{22} is hydrogen or alkyl;

(u) $-NHC(X)NR^{23}R^{24}$ where X is -O- or -S-, and R^{23} and R^{24} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(v) $-CONR^{25}R^{26}$ where R^{25} and R^{26} independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclylalkyl, or R^{25} and R^{26} together with the nitrogen to which they are attached form an optionally substituted heterocyclyl ring;

(w) $-S(O)_nR^{27}$ where n is an integer from 0 to 2, and R^{27} is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or $-NR^{28}R^{29}$ where R^{28} and R^{29} are, independently of each other, hydrogen, alkyl or heteroalkyl;

(x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;

(y) arylaminoalkylene or heteroarylaminoalkylene;

(z) Z-alkylene-$NR^{30}R^{31}$ or Z-alkylene-OR^{32} where Z is-NH-, -N(lower alkyl)- or -O-, and R^{30}, $R^{31}$and R^{32} are independently of each other, hydrogen, alkyl or heteroalkyl;

(aa) $-OC(O)$-alkylene-CO_2H or -OC(O)-NR'R" (where R' and R" are independently hydrogen or alkyl); and

(bb) heteroarylalkenylene or heteroarylalkynylene;

R^4 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) alkoxy; and

(e) hydroxy;

R^5 is selected from the group consisting of

(a) hydrogen;

(b) halo;

(c) alkyl;

(d) haloalkyl;

(e) thioalkyl;

(f) hydroxy;

(g) amino;

(h) alkylamino;

(i) dialkylamino;

(j) heteroalkyl;

(k) optionally substituted heterocycle;

(l) optionally substituted heterocyclylalkyl;

(m) optionally substituted heterocyclylalkoxy;

(n) alkylsulfonyl;

(o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;

(p) heteroalkoxy; and

(q) carboxy;

R^6 is selected from the group consisting of:

(a) hydrogen;

(b) halo;

(c) alkyl; and

(d) alkoxy;

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

102. The compound of any one of claims 1-28, 31-63 and 67-89 wherein A or G is a cycloalkyl group.

103. The compound of any one of claims 1-28, 31-63, 67-89 and 102, wherein A or G is a cyclopentyl or cyclohexyl group.

104. The compound of any of claims 1-89 that is 3-(5-amino-4-benzoyl-3-methoxy-pyrazol-1-yl)-N-cyclopropyl-4-methyl-benzamide.

105. The compound of any of claims 1-89 that is 3-[5-amino-4-(3-cyanobenzoyl)-pyrazol-1-yl]-N-cyclopropyl-4-methylbenzamide.

106. A pharmaceutical composition, comprising a compound of any one of claims 1-105 and a pharmaceutically acceptable carrier.

107. The pharmaceutical composition of claim 106 that is formulated for single dosage administration.

108. An article of manufacture, comprising packaging material, a compound of any one of claims 1-105 which is useful for treating, preventing, or ameliorating one or more symptoms of p38 kinase-mediated diseases or disorders, and a label that indicates that the compound is useful for treating, preventing, or ameliorating one or more symptoms of p38 kinase-mediated diseases or disorders.

109. A method of treating, preventing, or ameliorating one or more symptoms of p38 kinase-mediated diseases or disorders, comprising administering to a subject in need thereof a compound of any one of claims 1-105.

110. The method of claim 109, wherein the disease or disorder is selected from inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, angiogenic disorders, infectious diseases, neurodegenerative diseases, and viral diseases.

111. The method of claim 109 or claim 110, wherein the disease or disorder is selected from pancreatitis (acute or chronic), asthma, allergies, adult respiratory distress syndrome, chronic obstructive pulmonary disease, glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosis, scleroderma, chronic thyroiditis, Grave's disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, graft vs. host disease, inflammatory reaction induced by endotoxin, tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, gout, traumatic arthritis, rubella arthritis, acute synovitis, pancreatic

β-cell disease; diseases characterized by massive neutrophil infiltration; rheumatoid spondylitis, gouty arthritis and other arthritic conditions, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption disease, allograft rejections, fever and myalgias due to infection, cachexia secondary to infection, meloid formation, scar tissue formation, ulcerative colitis, pyresis, influenza, osteoporosis, osteoarthritis and multiple myeloma-related bone disorder, acute myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, multiple myeloma, sepsis, septic shock, and Shigellosis; Alzheimer's disease, Parkinson's disease, cerebral ischemias or neurodegenerative disease caused by traumatic injury; angiogenic disorders including solid tumors, ocular neovasculization, and infantile haemangiomas; viral diseases including acute hepatitis infection (including hepatitis A, hepatitis B and hepatitis C), HIV infection and CMV retinitis, AIDS, SARS, ARC or malignancy, and herpes; stroke, myocardial ischemia, ischemia in stroke heart attacks, organ hyposia, vascular hyperplasia, cardiac and renal reperfusion injury, thrombosis, cardiac hypertrophy, thrombin induced platelet aggregation, endotoxemia and/or toxic shock syndrome, and conditions associated with prostaglandin endoperoxidase synthase-2.

112. A method of inhibiting the expression of inducible pro-inflammatory proteins, comprising administering to a subject in need thereof a compound of any one of claims 1-105.

113. The method of claim 112, wherein the inducible pro-inflammatory protein is prostaglandin endoperoxide synthase-2 (PGHS-2), also referred to as cyclooxygenase-2 (COX-2).

114. A method of treating, preventing, or ameliorating one or more symptoms of diseases or disorders associated with inducible pro-inflammatory proteins, comprising administering to a subject in need thereof a compound of any one of claims 1-105.

115. The method of claim 114, wherein the disease or disorder is selected from edema, analgesia, fever, pain, neuromuscular pain, headache, pain caused by cancer, dental pain and arthritis pain.

116. The method of claim 110, wherein the viral infection is a veterinary viral infection.

117. The method of claim 116, wherein the veterinary viral infection is lentivirus infection, equine infectious anemia virus; retro virus infection, feline immunodeficiency virus, bovine immunodeficiency virus, and canine immunodeficiency virus.

118. A method of treating, preventing, or ameliorating one or more symptoms of a cytokine mediated disease or disorder, comprising administering to a subject in need thereof a compound of any one of claims 1-105.

119. The method of any one of claims 109-118, further comprising administering a corticosteroid, rolipram, calphostin, a CSAID, a 4-substituted imidazo[1,2-A]quinoxaline, interleukin-10, a glucocorticoid, a salicylate, nitric oxide, an immunosuppressant, a nuclear translocation inhibitor, deoxyspergualin (DSG); a non-steroidal antiinflammatory drug (NSAID), ibuprofen, celecoxib, rofecoxib; a steroid, prednisone, dexamethasone; an antiviral agent, abacavir; an antiproliferative agent, methotrexate, leflunomide, FK506; a cytotoxic drug, azathioprine, cyclophosphamide, a TNF-α inhibitor, tenidap, an anti-TNF antibody, a soluble TNF receptor, and rapamycin, or derivatives thereof.


120. The pharmaceutical composition of claim 106 or claim 107, further comprising one or more of the following: corticosteroid, rolipram, calphostin, a CSAID, a 4-substituted imidazo[1,2-A]quinoxaline, interleukin-10, a glucocorticoid, a salicylate, nitric oxide, an immunosuppressant, a nuclear translocation inhibitor, deoxyspergualin (DSG); a non-steroidal antiinflammatory drug (NSAID), ibuprofen, celecoxib, rofecoxib; a steroid, prednisone, dexamethasone; an antiviral agent, abacavir; an antiproliferative agent, methotrexate, leflunomide, FK506; a cytotoxic drug, azathioprine, cyclophosphamide, a TNF-α inhibitor, tenidap, an anti-TNF antibody, a soluble TNF receptor, and rapamycin, or derivatives thereof.

121. A method of inhibiting a p38 kinase, comprising contacting the kinase with a compound of any one of claims 1-105.

122. The method of claim 121, wherein the kinase is p38α kinase or p38β kinase.


123. A process for preparing a compound of claim 18, comprising the steps of (i) and (iii)-(vi), or the steps of (ii)-(vi), as follows:

(i) reacting a 2-keto-3-phenylaminoacrylonitrile of formula 1:



1


with a hydrazine of formula 2:



2


to provide a compound of formula (I) where R^1 is hydrogen; or

(ii) reacting a 2-keto-3-phenylaminoacrylonitrile of formula 3:



3

wherein Z is hydroxy, nitro or halo, with a hydrazine of formula 2 to provide a compound of formula 4:



4

followed by conversion of the Z group to the desired R^3 group to provide a compound of claim 1 where R^1 is hydrogen;


(iii) optionally modifying any of the R^1, R^3, R', R^5 or R^6 groups;

(iv) optionally converting the compound of formula (I) prepared in steps (i), (ii) or (iii) above to the corresponding acid addition salt by treatment with an acid;

(v) optionally converting the compound of formula (I) prepared in steps (i), (ii) or (iii) above, to the corresponding free base by treatment with a base; and


(vi) optionally separating a mixture of stereoisomers of a compound of formula (I) prepared in steps (i) - (v) above, to give a single stereoisomer.

124. A process for preparing a compound of claim 18, which process involves reacting a compound of formula 5:



5

wherein L is a leaving group under organometallic displacement reaction conditions, with an organometallic reagent of formula:



wherein M is a metallic moiety, to provide a compound of formula (I) where R^1 is hydrogen;

(ii) optionally modifying any of the R^1, R^3, R^4, R^5 or R^6 groups;

(iii) optionally converting the compound of formula (I) prepared in steps (i) or (ii) above, to the corresponding acid addition salt by treatment with an acid;

(iv) optionally converting the compound of formula (I) prepared in steps (i) or (ii) above, to the corresponding free base by treatment with a base; and

(v) optionally separating a mixture of stereoisomers of the compound prepared in steps (i) or (iv) above, to give a single stereoisomer.

125. A compound prepared by the process of claim 123 or claim 124.

126. A compound of any one of claims 1-105 when used in the treatment, prevention, or amelioration of one or more symptoms of a p38 kinase mediated disease or disorder.

127. Use of a compound of any one of claims 1-105 for the manufacture of a medicament for the treatment, prevention, or amelioration of one or more symptoms of a p38 kinase mediated disease or disorder.

128. A method of mediating cytokine response, comprising administering to a subject in need thereof a compound of any one of claims 1-105.

129. The method of claim 116, wherein the cytokine response is induced by p38 kinase activity.