

NOVEL PEPTIDES FOR TREATING AND PREVENTING IMMUNE-RELATED DISORDERS, INCLUDING TREATING AND PREVENTING INFECTION BY MODULATING INNATE IMMUNITY

PRIOR RELATED APPLICATIONS

[0001] This application claims priority from United States Provisional Patent Applications, 60/722,962; 60/722,958; and 60/722,959, filed October 4, 2005, all entitled, "Novel Peptides For Treating And Preventing Infection By Modulating Innate Immunity", and which are all incorporated by reference herein.

FIELD OF THE INVENTION

[0002] This invention relates to peptides for use in treating and preventing immune-related disorders, including treating and preventing infection by modulating innate immunity. In one aspect, the invention relates to compositions and uses thereof for modulating innate immunity. In another aspect, the invention provides novel peptides and uses thereof effective in reducing DPPIV activity.

BACKGROUND OF THE INVENTION

[0003] A variety of microorganisms, including viruses, bacteria, fungi, and parasites, can cause disease. Microbial cells are distinct from the cells of animals and plants – which are unable to live alone in nature, existing only as parts of multicellular organisms. Microbial cells can be pathogenic or non-pathogenic, depending, in part, on the microorganism and the status of the host. For example, in an immunocompromised host, a normally harmless bacterium can become a pathogen. Entry into host cells is critical for the survival of bacterial pathogens that replicate in an intracellular milieu. For organisms that replicate at extracellular sites, the significance of bacterial entry into host cells is less well defined.

[0004] Drug resistance remains an obstacle in the ongoing effort to fight infection. For example, penicillin was effective in treating *Staphylococcus aureus*, until the bacterium became resistant. Throughout the second half of the 20th century, new antibiotics, such as vancomycin and methicillin, were developed; these successfully cured *S. aureus* infections. However, methicillin-resistant strains of *S. aureus* evolved in the 1970s, and have been

plaguing hospitals worldwide ever since. More recently, vancomycin-resistant strains of *S. aureus* have surfaced.

[0005] With the increasing threat of resistance to antimicrobial drugs and the emergence of new infectious diseases, there exists a continuing need for novel therapeutic compounds. Therapeutics that act on the host, not the pathogen, are desirable, because they do not encourage pathogenic resistance. In particular, drugs that act on the host *via* the innate immune system provide a promising source of therapeutics.

[0006] Host defense against microorganisms begins with the epithelial barriers of the body and the innate immune system, and culminates in the induction of the adaptive immune response. The host innate immune response encompasses a set of highly-conserved mechanisms that recognize and counter microbial infections. Elements of innate immunity are continuously maintained at low levels, and are activated very rapidly when stimulated. The innate immune response begins with events that occur immediately after exposure to a microbial pathogen. Events associated with adaptive immunity, such as rearrangement of immunoglobulin receptor genes, are not considered part of the innate response.

[0007] There is evidence to indicate that innate responses are instrumental in controlling most infections, and also contribute to inflammatory responses. Inflammatory responses triggered by infection are known to be central components of disease pathogenesis. The importance of Toll-like receptors (TLRs) in the innate immune response has also been well characterized. The mammalian family of TLRs recognizes conserved molecules, many of which are found on the surfaces of, or are released by, microbial pathogens. There are numerous other mechanisms, less well characterized, that initiate and/or contribute to the host innate defense.

[0008] The innate immune system provides a range of protective mechanisms, including epithelial-barrier function and secretion of cytokines and chemokines. To date, four families of chemokines have been categorized, according to the number of conserved N-terminal cysteine motifs: C, CC, CXC, and CX3C, where X is a non-conserved amino acid residue. The CXC chemokines are known to be chemotactic for cells bearing the CXCR3 receptor, including monocytes, activated T cells (Th1), and NK cells. Primary human airway epithelial cells, and the cell line 16-HBE, constitutively express the CXCR3 receptor and its

ligands, IP-10, I-TAC, and MIG (Kelsen *et al.*, The chemokine receptor CXCR3 and its splice variant are expressed in human airway epithelial cells, *Am. J. Physiol. Lung Cell Mol. Physiol.*, 287:L584, 2004). Furthermore, CXCR3 ligands induce chemotactic responses and actin reorganization in 16-HBE cells (Kelsen *et al.*, The chemokine receptor CXCR3 and its splice variant are expressed in human airway epithelial cells, *Am. J. Physiol. Lung Cell Mol. Physiol.*, 287:L584, 2004).

[0009] Further, the type II transmembrane serine protease dipeptidyl peptidase IV (DPPIV), also known as CD26 or adenosine deaminase binding protein, is a major regulator of various physiological processes including immune functions. CD26/DPPIV is a 110-kD cell surface glycoprotein that is mainly expressed on mature thymocytes, activated T-cells, B-cells, NK-cells, macrophages, and epithelial cells. It has at least two functions, a signal transduction function and a proteolytic function (Morimoto C, Schlossman SF. The structure and function of CD26 in the T-cell immune response. *Immunol. Review.* 1998, 161: 55-70.). One of its cellular roles involves modulation of chemokine activity by cleaving dipeptides from the chemokine N-terminus. The modulation of the NH₂ termini of chemokines is of great importance not only for binding to their receptors and the following reactions but also for altering the receptor specificity of the processed chemokine. DPPIV activity has been associated with a number of immune-related conditions.

SUMMARY OF THE INVENTION

[0010] The inventors have discovered that peptides having the amino acid sequence of one of the peptides listed and described in TABLE 1 or an analogue, derivative, or variant thereof can enhance a host's innate immunity. In one aspect, the immunomodulatory peptides of the invention were found to lack antimicrobial activity while demonstrating an ability to improve survival in infected hosts. In another aspect, the invention provides peptides that modulate DPPIV activity. In one aspect the invention provides peptides that reduce DPPIV activity. In yet another aspect, the invention provides peptides which can be used in the diagnosis, treatment or prevention of an immunological disorder, such as one associated with DPPIV activity and/or innate immunity..

[0011] Accordingly, in one aspect, the present invention provides an isolated peptide that includes the amino acid sequence of any one of TABLE 1 or an analogue, derivative, or

variant thereof or obvious chemical equivalent thereof or a peptide comprising said peptide. In one embodiment the peptide is up to 10 amino acids comprising said peptide. By way of example, the isolated peptide may have a modified C-terminus (e.g., an amidated C-terminus) and/or a modified N-terminus. The isolated peptide of the invention may further include the amino acid sequence of TABLE 1 as modified by at least one substitution of a D amino acid. The isolated peptide may further include a modified backbone, by way of example, wherein the N-terminus is modified from an amide to an N-methyl. In one aspect, those modified peptides which retain the immunological activity of the parent peptide and obvious chemical equivalent thereto which retain said activity are encompassed within the scope of the present invention.

[0012] In another aspect, the present invention further provides an agent reactive with an isolated peptide that includes the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof. In one embodiment, the agent is a non-naturally occurring antibody (e.g., a polyclonal or monoclonal antibody). In one embodiment, the antibody is made using a MAPS Antigen attached to the peptide of the present invention via 2 glycine residues inserted at the C-terminus of the peptide. The construct can then be administered to an animal, such as a rabbit and the antibody harvested using procedures well known in the art. In one aspect, such agents can be labeled or used to label peptides of the invention. In another aspect such agents can be used in diagnostic and screening methods to monitor agents that may modulate peptide activity or to quantitate the amount of the peptide.

[0013] In yet another aspect, the present invention provides an isolated nucleic acid molecule encoding an isolated peptide having or comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof. Also provided is a recombinant nucleic acid construct that includes the nucleic acid molecule operably linked to an expression vector.

[0014] In a further aspect, the present invention provides at least one host cell comprising the recombinant nucleic acid construct of the invention. Also provided is a method for producing a peptide having or comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof, by: (a) culturing the at least one host cell,

under conditions allowing expression of the peptide; and (b) recovering the peptide from the at least one host cell or culture medium thereof.

[0015] In still another aspect, the present invention provides a pharmaceutical composition that includes an isolated peptide having or comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof (including a pharmaceutically-acceptable salt, addition salt, or ester of any of the foregoing or polymorph), in combination with a pharmaceutically-acceptable carrier, diluent, or excipient.

[0016] In another aspect, the present invention provides a method for treating and/or preventing infection (*e.g.*, a microbial infection) in a subject, by administering to the subject a peptide having or comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof or obvious chemical equivalent thereof. By way of example, the subject may have, or be at risk of having, infection. In one embodiment, the peptide modulates innate immunity in the subject, thereby treating and/or preventing the infection in the subject. The present invention further provides a method for identifying a microbial infection that can be treated with a peptide of the invention. In another aspect, the invention provides a method for treating or preventing a DPPIV-related condition or disorder.

[0017] Exemplary infections which may be treated and/or prevented by the method of the present invention include an infection by a bacterium (*e.g.*, a Gram-positive or Gram-negative bacterium), an infection by a fungus, an infection by a parasite, and an infection by a virus. In one embodiment of the present invention, the infection is a bacterial infection (*e.g.*, infection by *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella spp.*, *Staphylococcus aureus*, *Streptococcus spp.*, or vancomycin-resistant enterococcus). In another embodiment, the infection is a fungal infection (*e.g.*, infection by a mould, a yeast, or a higher fungus). In still another embodiment, the infection is a parasitic infection (*e.g.*, infection by a single-celled or multicellular parasite, including *Giardia duodenalis*, *Cryptosporidium parvum*, *Cyclospora cayetanensis*, and *Toxoplasma gondii*). In yet another embodiment, the infection is a viral infection (*e.g.*, infection by a virus associated with AIDS, avian flu, chickenpox, cold sores, common cold, gastroenteritis, glandular fever, influenza, measles, mumps, pharyngitis, pneumonia, rubella, SARS, and lower or upper respiratory tract infection (*e.g.*, respiratory syncytial virus)).

[0018] In accordance with the method of the present invention, a peptide having or comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof may be administered to the subject directly (*i.e.*, by administering the peptide itself) or indirectly (*e.g.*, by administering to the subject a nucleic acid sequence encoding the peptide, in a manner permitting expression of the peptide in the subject). The peptide of the invention (or nucleic acid encoding same) may be administered to the subject orally, parenterally (*e.g.*, intradermally, intramuscularly, intraperitoneally, intravenously, or subcutaneously), transdermally, intranasally, by pulmonary administration (*e.g.*, by intratracheal administration), and/or by osmotic pump.

[0019] In yet another aspect, the present invention provides a method for predicting whether a subject would be responsive to treatment with a peptide comprising the amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof, by assaying a diagnostic sample of the subject for DPPIV activity, wherein modulation, such as reduction of DPPIV activity is indicative that the subject would be responsive to treatment by the peptide. In one aspect, the subject has or is suspected of having a DPPIV-related condition or disorder.

[0020] Additional aspects and advantages of the present invention will be apparent in view of the description which follows. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE FIGURES

[0021] The invention will now be described in relation to the drawings, in which:

[0022] FIG. 1 A and B depicts the results of the experiment described in Example 2. %Viability = the amount of bacterial growth relative to the vehicle control (Tris), which is set to 100% bacterial survival with respective peptides SEQ. ID. NOs. 5, and 47 ; Erythr. = erythromycin.

[0023] FIG. 2 A – G depicts the results of the experiment described in Example 3. The graph shows colony-forming units per ml (CFU/ml) on the Y-axis, and treatment group

(control = no peptide; SEQ. ID. NOs. 1, 4, 5, 6, 45 and 47 = treatment with a peptide having the respective amino acid sequence) on the X-axis. The bacterial count of individual mice is shown.

[0024] FIG. 3 A and B depicts the results of the experiment described in Example 4. The graph shows colony-forming units per ml (CFU/ml) on the Y-axis, and treatment group (control = no peptide; SEQ. ID. NOs. 1 and 5 = treatment with a peptide having the respective amino acid sequence) on the X-axis. The bacterial count of individual mice is shown.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0025] “DPPIV-related disorder” or “DPPIV-related condition” or “DPPIV associated condition” as used herein means any medical condition that has been correlated with DPPIV activity and wherein modulation of said activity can be used to treat and/or prevent or diagnose said condition. Examples of such conditions include, but are not limited to: HIV/AIDS, autoimmune conditions, such as Rheumatoid Arthritis, multiple sclerosis, cancer (e.g. colon and lung), diabetes, and Graves disease.

[0026] “Immune-related disorder” is a condition that is associated with the immune system of a subject, either through activation or inhibition of the immune system, or that can be treated, prevented or diagnosed by targeting a certain component of the immune response in a subject, such as the innate immune response.

[0027] “Immunologically active” as used herein refers to innate immune activity (e.g. the ability to modulate the innate immune response or component thereof in a subject) or the ability to modulate DPPIV activity.

[0028] “Modulate” or “Modulating” as used herein, for instance such as modulating DPPIV activity or a particular response, encompasses the increase or decrease of activity or response in relation to a control or the normal or baseline level of activity or response under certain conditions. It can also encompass the maintaining of a level of activity or response

under conditions that would normally increase or decrease the level of activity of the peptide or response.

[0029] "Pharmaceutically acceptable salts" refer to the non-toxic alkali metal, alkaline earth metal, and ammonium salts commonly used in the pharmaceutical industry including the sodium, potassium, lithium, calcium, magnesium, barium, ammonium, and protamine zinc salts, which are prepared by methods well known in the art. The term also includes non-toxic acid addition salts, which are generally prepared by reacting the compounds of this invention with a suitable organic or inorganic acid. Representative salts include the hydrochloride, hydrobromide, sulfate, bisulfate, acetate, oxalate, valerate, oleate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, napsylate, trifluoroacetate and the like.

[0030] "Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as trifluoroacetic acid, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, menthanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. For a description of pharmaceutically acceptable acid addition salts as prodrugs, see Bundgaard, H., ed., (1985) *Design of Prodrugs*, Elsevier Science Publishers, Amsterdam.

[0031] "Pharmaceutically acceptable ester" refers to those esters which retain, upon hydrolysis of the ester bond, the biological effectiveness and properties of the carboxylic acid or alcohol and are not biologically or otherwise undesirable. For a description of pharmaceutically acceptable esters as prodrugs, see Bundgaard, H., *supra*. These esters are typically formed from the corresponding carboxylic acid and an alcohol. Generally, ester formation can be accomplished via conventional synthetic techniques. (See, e.g., March, *Advanced Organic Chemistry*, 3rd Ed., John Wiley & Sons, New York (1985) p. 1157 and references cited therein, and Mark et al., *Encyclopedia of Chemical Technology*, John Wiley & Sons, New York (1980).) The alcohol component of the ester will generally comprise (i) a

C.₂-C.₁₂. aliphatic alcohol that can or can not contain one or more double bonds and can or can not contain branched carbon chains or (ii) a C.₇- C.₁₂ aromatic or heteroaromatic alcohols. This invention also contemplates the use of those compositions which are both esters as described herein and at the same time are the pharmaceutically acceptable acid addition salts thereof.

[0032] "Pharmaceutically acceptable amide" refers to those amides which retain, upon hydrolysis of the amide bond, the biological effectiveness and properties of the carboxylic acid or amine and are not biologically or otherwise undesirable. For a description of pharmaceutically acceptable amides as prodrugs, see Bundgaard, H., ed., *supra*. These amides are typically formed from the corresponding carboxylic acid and an amine. Generally, amide formation can be accomplished via conventional synthetic techniques. (See, e.g., March, *Advanced Organic Chemistry*, 3rd Ed., John Wiley & Sons, New York (1985) p. 1152 and Mark et al., *Encyclopedia of Chemical Technology*, John Wiley & Sons, New York (1980).) This invention also contemplates the use of those compositions which are both amides as described herein and at the same time are the pharmaceutically acceptable acid addition salts thereof.

[0033] "Pharmaceutically or therapeutically acceptable carrier" refers to a carrier medium which does not interfere with the effectiveness of the biological activity of the active ingredients and which is not toxic to the host or patient.

[0034] "Stereoisomer" refers to a chemical compound having the same molecular weight, chemical composition, and constitution as another, but with the atoms grouped differently. That is, certain identical chemical moieties are at different orientations in space and, therefore, when pure, has the ability to rotate the plane of polarized light. However, some pure stereoisomers may have an optical rotation that is so slight that it is undetectable with present instrumentation. The compounds of the instant invention may have one or more asymmetrical carbon atoms and therefore include various stereoisomers. All immunologically active stereoisomers are included within the scope of the invention.

[0035] "Therapeutically or pharmaceutically effective amount" as applied to the compositions of the instant invention refers to the amount of composition sufficient to induce a desired biological result. That result can be alleviation of the signs, symptoms, or causes of

a disease, or any other desired alteration of a biological system. For instance, in the present invention, the result will typically involve enhancement of the innate immune response, reduction of DPPIV activity and/or modulation (such as inhibition or reduction or non-stimulation) of the inflammatory responses to infection or tissue injury.

[0036] Amino acid residues in peptides are abbreviated as follows: Phenylalanine is Phe or F; Leucine is Leu or L; Isoleucine is Ile or I; Methionine is Met or M; Valine is Val or V; Serine is Ser or S; Proline is Pro or P; Threonine is Thr or T; Alanine is Ala or A; Tyrosine is Tyr or Y; Histidine is His or H; Glutamine is Gln or Q; Asparagine is Asn or N; Lysine is Lys or K; Aspartic Acid is Asp or D; Glutamic Acid is Glu or E; Cysteine is Cys or C; Tryptophan is Trp or W; Arginine is Arg or R; and Glycine is Gly or G. In addition, the abbreviation Nal is used to denote 1-naphthylalanine; Ornithine is Orn or O, Cit is citrulline Hci is citrulline with one more methylene groups, and Vx or Valine x, wherein the "x" refers to a variation in the backbone of the amino acid, wherein the amino acid linkage is no longer an amide bond, but a methylated amine, this similarly applies to other amino acids with the "x" designation. Also, 2,4-diaminobutyric acid is Dab; 2,3-diaminopropionic acid is Dpr or Dapa; N-(4-aminobutyl)-glycine is Nlys; hSer is homoserine; Hyp is hydroxyproline; Val(betaOH) is hydroxyvaline; D-Pro is 3,4-dehydropyrrolidine; Pyr is pyroglutamine (proline with C=O in ring); Proline with fluorine substitutions on the ring; 1,3-thiazolidine-4-carboxylic acid (proline with S in ring); Thi is beta-(2thienyl)-alanine; Abu is 2-aminobutyric acid; Nva is norvaline; Nle is norleucine; Hol is homoleucine; and Aib is alpha-aminoisobutyric acid.

[0037] In addition to peptides consisting only of naturally-occurring amino acids, peptidomimetics or peptide analogs are also provided. Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics" (Fauchere, J., *Adv. Drug Res.* 15:29 (1986); Veber and Freidinger, *TINS* p. 392 (1985); and Evans et al., *J Med. Chem.* 30:1229 (1987), which are incorporated herein by reference). Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent or enhanced therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm peptide (i.e., a peptide that has a biological or pharmacological activity), such as naturally-occurring receptor-binding

peptide, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: $--CH_2NH--$, $--CH_2S--$, $--CH_2=CH_2--$, $--CH=CH--$ (cis and trans), $--COCH_2--$, $--CH(OH)CH_2--$, and $--CH_2SO--$, by methods known in the art and further described in the following references: Spatola, A. F. in *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983); Spatola, A. F., *Vega Data* (March 1983), Vol. 1, Issue 3, PEPTIDE BACKBONE MODIFICATIONS (general review); Morley, *Trends Pharm Sci* (1980) pp. 463 468 (general review); Hudson, D. et al., *Int J Pept Prot Res* 14:177 185 (1979): ($--CH_2NH--$, CH_2CH_2--); Spatola et al., *Life Sci* 38:1243 1249 (1986): ($--CH_2--S--$); Hann J. Chem. Soc. *Perkin Trans. I* 307 314 (1982): ($--CH--CH--$, cis and trans); Almquist et al., *J Med Chem* 23:1392 1398 (1980): ($--COCH_2--$); Jennings-White et al., *Tetrahedron Lett* 23:2533 (1982): ($--COCH_2--$); Szelke et al., *European Application. EP 45665 CA: 97:39405* (1982) ($--CH(OH)CH_2--$); Holladay et al., *Tetrahedron Lett* 24:4401 4404 (1983): ($--C(OH)CH_2--$); and Hruby *Life Sci* 31:189 199 (1982): ($--CH_2--S--$); each of which is incorporated herein by reference. In one aspect, the non-peptide linkage is $--CH_2NH--$. Such peptide mimetics may have significant advantages over polypeptide embodiments, including, for example: more economical production, greater chemical stability, enhanced pharmacological properties (half-life, absorption, potency, efficacy, etc.), altered specificity (e.g., a broad-spectrum of biological activities), reduced antigenicity, and others. Labeling of peptidomimetics usually involves covalent attachment of one or more labels, directly or through a spacer (e.g., an amide group), to non-interfering position(s) on the peptidomimetic that are predicted by quantitative structure-activity data and/or molecular modeling. Such non-interfering positions generally are positions that do not form direct contacts with the macromolecules(s) (e.g., immunoglobulin superfamily molecules) to which the peptidomimetic binds to produce the therapeutic effect. Derivatization (e.g., labeling) of peptidomimetics should not substantially interfere with the desired biological or pharmacological activity of the peptidomimetic. Generally, peptidomimetics of receptor-binding peptides bind to the receptor with high affinity and possess detectable biological activity (i.e., are agonistic or antagonistic to one or more receptor-mediated phenotypic changes).

[0038] Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used to generate more stable peptides. It is appreciated that those D-amino acid substitutions wherein immunological activity of the peptide is retained are desired.

Description

[0039] As described herein, the inventors have identified novel peptides having and/or comprising the amino acid sequence as shown in TABLE 1 or an analogue, derivative, or variant of the amino acid sequences disclosed therein. The inventors have also demonstrated that a peptide having or comprising one of the amino acid sequences of TABLE 1 and an amidated C-terminus has therapeutic utility in the enhancement of innate immunity. In particular, the inventors have shown that a peptide comprising an amino acid sequence of TABLE 1 lacked antimicrobial efficacy against *S. aureus*, yet provided *in vivo* protection in mice infected with *S. aureus*. The peptide enhanced the host response to infection, resulting in improved bacterial clearance and host survival. Thus, the novel peptides described can be used as a therapeutic for the treatment of infectious disease. In another embodiment, the peptides of the invention have been shown to reduce DPPIV activity, which has been shown to be related to a number of immune-related disorders, such as , AIDS and HIV disease progression (Blazquez et al. 1992; Vanham et al. 1993; Schols et al. 1998 Oravec et al. 1995), Graves' disease (Eguchi et al. 1989; Nishikawa et al. 1995), and cancer (Stecca et al. 1997), such as lung and colon cancer, and diabetes (Hinke et al. 2000; Marguet et al. 2000). Further, DPPIV as an indicator of T-cell activation has been shown to fluctuate in parallel with several autoimmune diseases such as rheumatoid arthritis (Nakao et al., 1989) and autoimmune thyroiditis (Eguchi et al., 1989). DPPIV has been described as a marker that correlates well with the level of activity of these diseases. It has furthermore been studied as an indicator of disease progression in chronic progressive multiple sclerosis (Constantinescu et al., 1995). The peptides of the invention can be used in the treatment of such conditions.

Peptides of the Invention

[0040] Accordingly, the present invention provides isolated peptides having or comprising the amino acid sequence of TABLE 1 or an immunologically active analogue, derivative, or variant thereof. Also provided are pharmaceutically-acceptable salts, acid addition salts, and esters of the peptides, analogues, derivatives, and variants of the invention, including those described herein, such as conservative substitution, and, N and C terminus modifications and backbone modifications, as described herein. As used herein, an "isolated" peptide of the invention is a peptide which either has no naturally-occurring counterpart or has been separated or purified from components which naturally accompany it. An isolated peptide of the invention can be obtained, for example, by expression of a recombinant nucleic acid encoding the peptide or by chemical synthesis. Because a peptide that is chemically synthesized is, by its nature, separated from the components that naturally accompany it, the synthetic peptide is "isolated".

[0041] In one aspect, the isolated peptide of the invention comprises the amino acid sequence having the formula: "X₁X₂P" (SEQ. ID. NO. 55), wherein: X₁ is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH), or in another embodiment is selected from the group consisting of K, R, S, O, and Cit, or in another embodiment, selected from the group consisting of K, R, and S, or is R; and wherein X₂ is selected from the group consisting of V, I, K, P, and H. In one embodiment, the isolated peptide of the invention is SEQ. ID. NO. 55. In another aspect, it is a peptide of up to 10 amino acids comprising an amino acid sequence of SEQ. ID. NO. 55. In one embodiment, the isolated peptide of SEQ. ID. NO. 55 is SEQ. ID. NOs. 8, 9, 26, 39, 40, 41, and 45-53, or an isolated peptide of up to 10 amino acids comprising said sequences. In another embodiment, the isolated peptide comprising SEQ. ID. NO. 55 is SEQ. ID. NO. 44, which is up to 13 amino acids.

[0042] In another embodiment, the invention provides an isolated peptide comprising the formula, "X₁X₂X₃P" (SEQ. ID. NO. 56) wherein X₁ is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH), or in another embodiment selected from the group consisting of K, H, R, S, T, and O, or in another embodiment, K, H, R, S, and T, or in another embodiment, K, H, R, S and O, or in another

embodiment, R, H, K and S; and wherein X_2 is selected from the group consisting of A, I, L, V, K, P, G, H, R, S, O, Dab, Dpr, Cit, Hci, Abu, Nva, Nle and where X_2 can be N-methylated, or in another embodiment, selected from the group consisting of A, I, L, V, K, P, G, H, and R, where it can be N-methylated; and wherein X_3 is selected from the group consisting of I, V, P, wherein in one embodiment, X_3 is not N-methylated. In one embodiment, the isolated peptide can be an amino acid sequence of up to 10 amino acids, comprising SEQ. ID. NO. 56, including SEQ. ID. NOs. 1, 3-7, 10-16, 18, 21 - 25, 27, 28, 31-39, 42, 43, or 47, or an isolated peptide of up to 11 amino acids comprising SEQ. ID. NO. 54. However, in one embodiment when SEQ. ID. NO. 56 is a hexamer, it is not SEQ. ID. NO. 2, or when in one embodiment, when it is a pentamer, it is not SEQ. ID. NO. 17. In one embodiment, the isolated peptide of the invention does not comprise a peptide comprising SEQ. ID. NOs. 2 or 17.

[0043] In another embodiment, the invention provides an isolated peptide comprising the peptide comprising the formula of SEQ. ID. NO. 56 in a pentamer or hexamer. In one embodiment, said peptide is immunologically active.

[0044] In one embodiment, the isolated peptide of the invention comprises a peptide of formula, " $aX_1X_2X_3P$ " (SEQ. ID. NO. 57) wherein X_1 , X_2 and X_3 are defined as for SEQ. ID. NO. 56, and wherein " a " is selected from the group consisting of S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, M, and F, or in another embodiment is selected from the group consisting of, S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, and M, or in another embodiment is selected from the group consisting of, S, P, I, R, and C, or in another embodiment is S. In one embodiment, the isolated peptide comprises SEQ. ID. NO. 57, or is a peptide of up to 10 amino acids comprising said sequence. In another embodiment, the isolated peptide is SEQ. ID. NOs. 4, 47, or when it is a hexamer, SEQ. ID. NO. 39, or an isolated peptide up to 10 amino acids comprising said sequences.

[0045] In another embodiment, the isolated peptide of the invention comprises a peptide of formula, " $X_1X_2X_3Pb$ " (SEQ. ID. NO. 58) wherein $X_1X_2X_3$ are as defined in SEQ. ID. NO. 56 and " b " is selected from the group consisting of A, A*, G, S, L, F, K, C, I, V, T, Y, R, H, O, and M, but in one embodiment not P, or in another embodiment selected from the group consisting of A, A*, G, S, L, F, and K, or in another embodiment selected from the

group consisting of A, A*,G, S, L, K and C, or in one embodiment, selected from the group consisting of A, A*,G, S, L, and K. Wherein A* denotes a D amino acid of Alanine. In one embodiment, the isolated peptide is an amino acid of up to 10 amino acids comprising SEQ. ID. NO. 58. In one embodiment, the isolated peptide is or comprises SEQ. ID. NOs. 5 – 8, 10, 11, 13-16, 21-25, 27, 28, 31, 33-38 and 42-43. In another embodiment, the peptide is of SEQ. ID. NO. 58, wherein "b" is not P or Y. or not RIVPP (SEQ. ID. NO. 17); or where X₃ is not G or not RIGPA, or X₃ is not Vx or not RIVxPA.

[0046] In one embodiment, the isolated peptide of the invention is or comprises a peptide similar to SEQ. ID. NO. 58, but wherein X₁ is instead selected from the group consisting of G, GG, or Cit, or wherein "b" is A, X₂ is I, X₃ is V, X₁ is G, GG, or Cit, or the peptide is SEQ. ID. NOs. 19, 20 and 36. In one embodiment, the isolated peptide comprises SEQ. ID. NO. 31. In another embodiment, the isolated peptide comprises a reverse sequence of SEQ. ID. NO. 58, or comprises SEQ. ID. NO. 30.

[0047] In one embodiment, the isolated peptide of the invention is or comprises a peptide having the amino acid sequence of SEQ. ID. NO. 29.

[0048] The peptide of the invention also provides an isolated peptide comprising the formula, "a₁a₂ X₁X₂X₃P" (SEQ. ID. NO. 59), where X₁, X₂ and X₃ are as defined in SEQ. ID. NO. 56 and a₁ is selected from the group consisting of K, I R, H, O, L, V, A, and G, or in one embodiment, K and I, or in one embodiment K and a₂ is selected from the group consisting of S, P, R T, H, K, O, L, V, A, G, S, and I or in one embodiment, S, P, and R, or in another embodiment, S and P, or in another embodiment P. In one embodiment, a₁ is not acetylated, or where a₁ is K, K is not acetylated or not SEQ. ID. NO. 2. In one embodiment, the isolated peptide is or comprises SEQ. ID. NOs, 1, and 47 or a peptide of up to 10 amino acids comprising SEQ. ID. NO. 59.

[0049] In another embodiment, the isolated peptide of the invention is or comprises a peptide of the formula , "a X₁X₂X₃Pb" (SEQ. ID. NO. 60) where X₁, X₂ and X₃ are as defined in SEQ. ID. NO. 56 and where "a" is selected from the group consisting of S, R, K, H, O, T, I, L, V, A, G or in another embodiment, S, R and I, or in another embodiment S and R, and wherein "b" is selected from the group consisting of A, V, I, L, G, K, H, R, O, S, T, F or in another embodiment, A. In another embodiment, the peptide of SEQ. ID. NO. 60 is

SEQ. ID. NOs. 3, 12 and 39, or a peptide of up to 10 amino acids comprising SEQ. ID. NO. 60 or SEQ. ID. NOs. 3, 12 or 39.

[0050] As used herein, a "peptide comprising an amino acid sequence of a sequence of TABLE 1" or a "peptide comprising an amino acid sequence of a sequence of TABLE 1" includes the peptide itself, obvious chemical equivalents thereto, isomers thereof (*e.g.*, isomers, stereoisomers, retro isomers, retro-inverso isomers, all-[D] isomers, all-[L] isomers, or mixed [L] and [D] isomers thereof), conservative substitutions therein, precursor forms thereof, endoproteolytically-processed forms thereof, such as cleavage of single amino acids from N or C terminals or immunologically active metabolites of the peptides of the invention, pharmaceutically-acceptable salts and esters thereof, and other forms resulting from post-translational modification. Also included is any parent sequence, up to and including 10, 9, 8, 7, 6, 5 and 4 amino acids in length (cyclized, or linear, or branched from the core parent sequence), for which the specified sequence is a subsequence. A person skilled in the art would appreciate that where the peptide in the table is a trimer, it could be a subsequence of a 10, 9, 8, 7, 6, 5, and 4 mer, whereas if the peptide listed in TABLE 1 is a hexamer, it could be a subsequence of a 10, 9, 8, and 7 mer, but not a 5 or 4 mer. In addition, the invention comprises sequences that are greater than 10 mer, SEQ. ID. NOs. 44 and 54. Those modified peptides which retain the immunological activity of the peptides of the invention are encompassed within the scope of the present invention.

[0051] As further used herein, an "obvious chemical equivalent" of a peptide of the invention is a molecule which possesses the same desired activity, *e.g.* immunological activity, as peptides described herein, and exhibits a trivial chemical different, or a molecule which is converted, under mild conditions, into a peptide of the invention (*e.g.*, esters, ethers, reduction products, and complexes of the peptides of the invention).

[0052] Additionally, as used herein, "conservative substitutions" are those amino acid substitutions which are functionally equivalent to the substituted amino acid residue, either because they have similar polarity or steric arrangement, or because they belong to the same class as the substituted residue (*e.g.*, hydrophobic, acidic, or basic). The term "conservative substitutions", as defined herein, includes substitutions having an inconsequential effect on the ability of the peptide of the invention to enhance innate

immunity. Examples of conservative substitutions include the substitution of a polar (hydrophilic) residue for another (*e.g.*, arginine/lysine, glutamine/asparagine, or threonine/serine); the substitution of a non-polar (hydrophobic) residue (*e.g.*, isoleucine, leucine, methionine, phenylalanine, tyrosine, or valine) for another; the substitution of an acidic residue (*e.g.*, aspartic acid or glutamic acid) for another; or the substitution of a basic residue (*e.g.*, arginine, histidine, lysine or ornithine) for another.

[0053] The term "analogue", as used herein, includes any peptide having an amino acid sequence substantially identical to a sequence described herein, in which at least one residue has been conservatively substituted with a functionally-similar residue. An "analogue" has 60% or greater (preferably, 70% or greater, 80% or greater, 85% or greater, 90% or greater, 95% or greater, or 99%) amino-acid-sequence homology with an amino acid sequence of TABLE 1, and is a functional variant thereof. As further used herein, the term "functional variant" refers to the activity of a peptide that demonstrates an ability to enhance innate immunity or reduce DPPIV activity, as described herein. An "analogue" includes a variant of an amino acid of TABLE 1 that has an homologous three-dimensional conformation. An "analogue" further includes any pharmaceutically-acceptable salt of an analogue as described herein. A "variant" further includes any pharmaceutically-acceptable salt of a variant as described herein.

[0054] A "derivative", as used herein, refers to a peptide of the invention having one or more amino acids chemically derivatized by reaction of a functional side group. Exemplary derivatized molecules include, without limitation, peptide molecules in which free amino groups have been derivatized to form salts or amides, by adding acetyl groups, amine hydrochlorides, carbobenzoxy groups, chloroacetyl groups, formyl groups, p-toluene sulfonyl groups, or t-butyloxycarbonyl groups. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. Furthermore, free carboxyl groups may be derivatized to form salts, esters (*e.g.*, methyl and ethyl esters), or hydrazides. Thus, a "derivative" further includes any pharmaceutically-acceptable salt of a derivative as described herein.

[0055] In one embodiment of the present invention, the isolated peptide of the invention has a modified C-terminus and/or a modified N-terminus. For example, the isolated peptide may have an amidated C-terminus. For example, the amino terminus can be

acetylated (Ac) or the carboxy terminus can be amidated (NH₂). However, in one embodiment of the invention, the peptides of the invention are not preferably acetylated if such a modification would result in loss of desired immunological activity. Amino terminus modifications include methylating (i.e., --NHCH₃ or --NH(CH₃)₂), acetylating, adding a carbobenzoyl group, or blocking the amino terminus with any blocking group containing a carboxylate functionality defined by RCOO--, where R is selected from the group consisting of naphthyl, acridinyl, steroidyl, and similar groups. Carboxy terminus modifications include replacing the free acid with a carboxamide group or forming a cyclic lactam at the carboxy terminus to introduce structural constraints.

[0056] In one embodiment backbone substitutions can be made, such as NH to NCH₃. The isolated peptide may also be a modification (e.g., a point mutation, such as an insertion or a deletion, or a truncation) of or comprising an amino acid sequence of TABLE 1. By way of example, the peptide may comprise an amino acid sequence of TABLE 1 as modified by at least one point insertion of a D amino acid as long as desired immunological activity is retained. In particular, proline analogs in which the ring size of the proline residue is changed from 5 members to 4, 6, or 7 members can be employed. Cyclic groups can be saturated or unsaturated, and if unsaturated, can be aromatic or non-aromatic.

[0057] One can replace the naturally occurring side chains of the 20 genetically encoded amino acids (or D amino acids) with other side chains with similar properties, for instance with groups such as alkyl, lower alkyl, cyclic 4-, 5-, 6-, to 7-membered alkyl, amide, amide lower alkyl, amide di(lower alkyl), lower alkoxy, hydroxy, carboxy and the lower ester derivatives thereof, and with 4-, 5-, 6-, to 7-membered heterocyclic.

[0058] Such substitutions can include but are not necessarily limited to: (1) non-standard positively charged amino acids, like: ornithine, Dab; 2,4-diaminobutyric acid, which is like ornithine minus one methylene group (or lysine minus two methylene groups), Dpr or Dapa; 2,3-diaminopropionic acid, which is like ornithine minus two methylene group (or lysine minus three methylene groups, or serine with an amino group instead of hydroxyl), Nlys; N-(4-aminobutyl)-glycine which has the lysine side chain attached to the "N-terminus". and compounds with aminopropyl or aminoethyl groups attached to the amino group of glycine. (2), Non-naturally occurring amino acids like arganine, no charge, such as,

Cit; citrulline and Hci; citrulline with one more methylene group; (3) non-standard non-naturally occurring amino acids with OH (e.g., like serine), such as, hSer; homoserine (one more methylene group, Hyp; hydroxyproline, Val(betaOH); hydroxyvaline, Pen; penicillamin, (Val(betaSH); (4) proline derivatives, such as, D-Pro, such as, 3,4-dehydropoline, Pyr; pyroglutamine (proline with C=O in ring), Proline with fluorine substitutions on the ring, 1,3-thiazolidine-4-carboxylic acid (proline with S in ring); (5) Histidine derivative, such as, Thi; beta-(2thienyl)-alanine; or (6) alkyl derivatives, such as, Abu; 2-aminobutyric acid (ethyl group on Calpha),, Nva; norvaline (propyl group on Calpha), Nle; norleucine (butyl group on Calpha), Hol; homoleucine (propyl group on Calpha), Aib, alpha-aminoisobutyric acid (valine without methylene group). A person skilled in the art would appreciate that those substitutions that retain the immunological activity of the parent peptide/sequence.

[0059] In another alternative embodiment, the C-terminal carboxyl group or a C-terminal ester can be induced to cyclize by internal displacement of the --OH or the ester (--OR) of the carboxyl group or ester respectively with the N-terminal amino group to form a cyclic peptide. For example, after synthesis and cleavage to give the peptide acid, the free acid is converted to an activated ester by an appropriate carboxyl group activator such as dicyclohexylcarbodiimide (DCC) in solution, for example, in methylene chloride (CH₂Cl₂), dimethyl formamide (DMF) mixtures. The cyclic peptide is then formed by internal displacement of the activated ester with the N-terminal amine. Internal cyclization as opposed to polymerization can be enhanced by use of very dilute solutions. Such methods are well known in the art.

[0060] One can also cyclize the peptides of the invention, or incorporate a desamino or descarboxy residue at the termini of the peptide, so that there is no terminal amino or carboxyl group, to decrease susceptibility to proteases or to restrict the conformation of the peptide. C-terminal functional groups of the compounds of the present invention include amide, amide lower alkyl, amide di(lower alkyl), lower alkoxy, hydroxy, and carboxy, and the lower ester derivatives thereof, and the pharmaceutically acceptable salts thereof.

[0061] One can also cyclize the peptide by adding an N and/or C terminal cysteine and cyclizing the peptide through disulfide linkages or other side chain interactions.

[0062] One can also incorporate a desamino or descarboxy residue at the termini of the peptide, so that there is no terminal amino or carboxyl group, to decrease susceptibility to proteases or to restrict the conformation of the peptide.

Method of Making Peptides

[0063] The present invention contemplates peptides, including peptide analogues, derivatives, and variants, that are produced synthetically, generated recombinantly, or isolated from native cells. A peptide of the invention may be synthesized by methods commonly known to one skilled in the art (e.g., as described in *Modern Techniques of Peptide and Amino Acid Analysis* (New York: John Wiley & Sons, 1981; and Bodansky, M., *Principles of Peptide Synthesis* (New York: Springer-Verlag N.Y., Inc., 1984). Examples of methods that may be employed in the synthesis of the peptides of the invention include, but are not limited to, solid-phase peptide synthesis, solution or liquid-method peptide synthesis, and synthesis using any of the commercially-available peptide synthesizers. In one embodiment, a peptide of the invention is synthesized *in vitro*, e.g., by chemical means or *in vitro* translation of mRNA. In another embodiment, a peptide of the invention is produced recombinantly, using conventional techniques and cDNA encoding the peptide. The amino acid sequences of the present invention may further comprise coupling agents and protecting groups which are used in the synthesis of peptide sequences, and which are well known to one of skill in the art.

[0064] Peptide analogues, derivatives, and variants of the invention can be made by a wide variety of different mutagenesis techniques well known to those skilled in the art. These techniques can be found in any molecular biology laboratory manual, including, for example, Sambrook *et al.*, *Molecular Cloning – A Laboratory Manual*, 2nd ed. (Plainview, NY: Cold Spring Harbor Press, 1989); or Ausubel *et al.*, *Current Protocols in Molecular Biology* (John Wiley & Sons). Mutagenesis kits are also available from many commercial molecular biology suppliers. Methods are available to make site-directed, regio-specific, or random mutagenesis in the initial amino acid sequence. After the analogues, derivatives, and variants are produced, they can be screened for the desired ability to enhance innate immunity, as described herein.

Agents Reactive With Peptide

[0065] The present invention further provides an agent reactive with a peptide comprising an amino acid sequence of TABLE 1 or an analogue, derivative, or variant of thereof. As used herein, "reactive" means the agent has affinity for, binds to, or is directed against the peptide of the invention. As further used herein, an "agent" shall include a protein, polypeptide, peptide, nucleic acid (including DNA or RNA), a non-naturally occurring antibody, Fab fragment, F(ab')₂ fragment, molecule, compound, antibiotic, drug, and any combination(s) thereof. A Fab fragment is a univalent antigen-binding fragment of an antibody, which is produced by papain digestion. A F(ab')₂ fragment is a divalent antigen-binding fragment of an antibody, which is produced by pepsin digestion. Preferably, the agent of the present invention is labeled with a detectable marker or label. A non-naturally occurring antibody means, an antibody that is generated with the peptide associated with another compound, such as two C-terminal glycine residues and MAPS. MAPS Antigen is attached to the peptide of the present invention via 2 glycine residues inserted at the C-terminus of the peptide. The construct can then be administered to an animal, such as a rabbit and the antibody harvested using procedures well known in the art.

[0066] In one embodiment of the present invention, the agent reactive with the peptide of the invention is an antibody. As used herein, the antibody of the present invention may be polyclonal or monoclonal. In addition, the antibody of the present invention may be produced by techniques well known to those skilled in the art. Polyclonal antibody, for example, may be produced by immunizing a mouse, rabbit, or rat with a purified peptide of the invention. Monoclonal antibody then may be produced by removing the spleen from the immunized animal, and fusing the spleen cells with myeloma cells to form a hybridoma which, when grown in culture, will produce a monoclonal antibody. See, e.g., J.G.R. Hurrel, *Monoclonal Hybridoma Antibodies: Techniques and Applications* (Boca Raton, FL: CRC Press Inc., 1982).

[0067] The antibody of the invention may be labeled with a detectable marker or label. Labeling of an antibody may be accomplished using one of a variety of labeling techniques, including peroxidase, chemiluminescent labels known in the art, and radioactive labels known in the art. The detectable marker or label of the present invention may be, for example, a nonradioactive or fluorescent marker, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine, which can be detected using fluorescence and other

imaging techniques readily known in the art. Alternatively, the detectable marker or label may be a radioactive marker, including, for example, a radioisotope. The radioisotope may be any isotope that emits detectable radiation, such as ^{35}S , ^{32}P , ^{125}I , ^3H , or ^{14}C . Radioactivity emitted by the radioisotope can be detected by techniques well known in the art. For example, gamma emission from the radioisotope may be detected using gamma imaging techniques, particularly scintigraphic imaging. Preferably, the agent of the present invention is a high-affinity antibody labeled with a detectable marker or label.

Isolated Nucleic Acid Molecules

[0068] In addition, the present invention provides an isolated nucleic acid molecule encoding a peptide comprising an amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof, including a conjugated peptide (e.g. a carrier-peptide construct) or other peptide, or a pro-peptide that metabolizes or cleaves to an immunologically active peptide of TABLE 1. Due to the degeneracy of the genetic code, the nucleic acid molecule of the invention includes a multitude of nucleic acid substitutions that will also encode a peptide of the invention. The present invention further provides a nucleic acid which hybridizes to the isolated nucleic acid molecule encoding an amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof.

[0069] The nucleic acid molecules of the present invention may be DNA or RNA. They may be prepared by a variety of techniques known to those skilled in the art, including, without limitation, automated synthesis of oligonucleotides using commercially-available oligonucleotide synthesizers, such as the Applied Biosystems Model 392 DNA/RNA synthesizer. In addition, the nucleic acid molecules of the present invention may be labeled with one or more detectable markers or labels. Labeling of the nucleic acid molecules may be accomplished using one of a number of methods known in the art – e.g., nick translation, end labeling, fill-in end labeling, polynucleotide kinase exchange reaction, random priming, or SP6 polymerase (for riboprobe preparation) – along with one of a variety of labels – e.g., radioactive labels, such as ^{35}S , ^{32}P , or ^3H , or nonradioactive labels, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine (ROX).

[0070] The present invention also provides a recombinant nucleic acid construct comprising a nucleic acid molecule of the invention operably linked to an expression vector.

As used herein, an "expression vector" is a DNA construct containing a DNA sequence which is operably linked to a suitable control sequence capable of effecting the expression of the DNA in a suitable host. The vector may be, for example, a plasmid, a phage particle, or a potential genomic insert. As further used herein, the term "operably linked" describes a functional relationship between two DNA regions. Expression vectors suitable for use in the present invention comprise at least one expression control element (*e.g.*, operator, promoter, lac system, leader sequence, termination codon, and/or polyadenylation signal) operably linked to the nucleic acid molecule encoding a peptide of the invention. In one embodiment, the expression vector is a eukaryotic expression vector that functions in eukaryotic cells (*e.g.*, a retroviral vector, a vaccinia virus vector, an adenovirus vector, a herpes virus vector, or a fowl pox virus vector).

[0071] Once operably linked to a nucleic acid molecule of the invention, the expression vector may be introduced into a recipient cell by any *in vivo* or *ex vivo* means suitable for transfer of nucleic acid, including, without limitation, electroporation, DEAE Dextran transfection, calcium phosphate transfection, lipofection, monocationic liposome fusion, polycationic liposome fusion, protoplast fusion, creation of an *in vivo* electrical field, DNA-coated microprojectile bombardment, injection with recombinant replication-defective viruses, homologous recombination, viral vectors, naked DNA transfer, or any combination thereof. Recombinant viral vectors suitable for transfer of nucleic acid include, but are not limited to, vectors derived from the genomes of viruses such as retrovirus, HSV, adenovirus, adeno-associated virus, Semiliki Forest virus, cytomegalovirus, and vaccinia virus.

[0072] The present invention further provides at least one host cell comprising the recombinant nucleic acid construct of the invention. The host cell of the invention is transformed with the nucleic acid construct described herein. The host cell may be eukaryotic (*e.g.*, an animal, plant, insect, or yeast cell) or prokaryotic (*e.g.*, *E. coli*).

[0073] In addition, the present invention provides a method for producing a peptide comprising an amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof. The method comprises the steps of: (a) culturing at least one host cell comprising a recombinant nucleic acid construct, as described herein, under conditions allowing expression of the peptide; and (b) recovering the peptide from the at least one host cell or

from the culture medium thereof. The recombinant peptide can be recovered as a crude lysate; it can also be purified by standard protein purification procedures known in the art, including, without limitation, affinity and immunoaffinity chromatography, differential precipitation, gel electrophoresis, ion-exchange chromatography, isoelectric focusing, size-exclusion chromatography, and the like.

Pharmaceutical Composition

[0074] The present invention further provides a pharmaceutical composition comprising a peptide comprising an amino acid sequence of TABLE 1 or SEQ. ID. NOs. 1, 3-16, or 18-60, or an analogue, derivative, or variant thereof (which includes a pharmaceutically-acceptable salt, acid addition salt or ester of any of the foregoing), in combination with at least one pharmaceutically-acceptable carrier, diluent, or excipient. The pharmaceutically-acceptable carrier, diluent, or excipient must be "acceptable" in the sense of being compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. Examples of acceptable pharmaceutical carriers, diluents, and excipients include, without limitation, carboxymethyl cellulose, crystalline cellulose, glycerin, gum arabic, lactose, magnesium stearate, methyl cellulose, powders, saline, sodium alginate, sucrose, starch, talc, and water, among others. Formulations of the pharmaceutical composition of the invention, as described herein, may be conveniently presented in unit dosage.

Uses

[0075] The peptides of the invention have been shown to have therapeutic utility in enhancing innate immunity. The enhancement of innate immunity is demonstrated by the lack of antimicrobial activity (Example 2) and the protection against infection in *in vivo* models (Examples 3 and 4) and also by the DPPIV assays of Example 5. Accordingly, the present invention also provides a method for treating and/or preventing infection in a subject. As used herein, the "subject" is a bird (e.g., a chicken, turkey, etc.) or a mammal (e.g., a cow, dog, human, monkey, mouse, pig, rat, etc.). In one embodiment, the subject is a human. The subject may have, or be at risk of having, an infection. By way of example, the infection may be a microbial infection. Microbial infections which may be treated by the method of

the present invention include, without limitation, infection by a bacterium, infection by a fungus, infection by a parasite, and infection by a virus.

[0076] Most bacterial pathogens are present in the general environment, or in the host's normal bacterial flora. Bacteria have evolved the ability to cause severe disease by acquiring different mechanisms (called virulence factors) which enable them to colonize, disseminate within, and invade host tissues. When these pathogenicity factors are suppressed, bacteria are no longer able to maintain themselves in host tissues, and, therefore, cannot cause disease. Exemplary bacteria which may be treated by the method of the present invention include, without limitation, *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella spp.* (e.g., *Salmonella typhimurium*), *Staphylococcus aureus*, *Streptococcus spp.*, and vancomycin-resistant enterococcus.

[0077] *Pseudomonas aeruginosa* is a ubiquitous Gram-negative bacterium that is noted for its environmental versatility, its ability to cause disease in susceptible individuals, and its resistance to antibiotics. It is a versatile organism that grows in soil, marshes, and coastal marine habitats, and on plant and animal tissues. The most serious complication of cystic fibrosis is respiratory tract infection by *P. aeruginosa*. Cancer and burn patients also commonly suffer serious infections by this organism, as do certain other individuals with immune system deficiencies. Unlike many environmental bacteria, *P. aeruginosa* has a remarkable capacity to cause disease in susceptible hosts.

[0078] *Staphylococcus aureus* is a Gram-positive spherical bacterium, about 1 micrometer in diameter, that thrives in microscopic clusters. It is one of the most important human pathogens, causing both community-acquired and nosocomial infections that range from endocarditis to pneumonia. Although *S. aureus* is generally classified as an extracellular pathogen, recent data have revealed its ability to infect various types of host cells, e.g., both professional phagocytes and non-phagocytes, including endothelial cells, fibroblasts, and others. This invasion is initiated by the adherence of *S. aureus* to the cell surface, a process in which staphylococcal fibronectin-binding proteins play a prominent role. Phagocytosed *S. aureus* can either induce apoptosis of the host cell or survive for several days in the cytoplasm – which is thought to be devoid of anti-staphylococcal effector mechanisms.

[0079] *S. aureus* colonizes nasal passages, skin surfaces, mucous membranes, and areas around the mouth, genitals, and rectum. *S. aureus* may cause superficial skin lesions, such as boils, styes, and furuncles. More serious infections include pneumonia, mastitis, phlebitis, meningitis, and urinary tract infections; deep-seated infections include osteomyelitis and endocarditis.

[0080] Exemplary fungi which may be treated by the method of the present invention include, without limitation, moulds, yeasts, and higher fungi. All fungi are eukaryotic, and have sterols, but not peptidoglycan, in their cell membranes. Fungal infections, or mycoses, are classified according to the degree of tissue involvement and the mode of entry into the host. In the immunocompromised host, a variety of non-pathogenic fungi, or fungi that are normally mild, can cause potentially fatal infections.

[0081] Parasites are organisms that derive nourishment and protection from other living organisms (known as hosts). They may be transmitted from animals to humans, from humans to humans, or from humans to animals. Several parasites have emerged as significant causes of food-borne and water-borne disease. They may be transmitted from host to host through consumption of contaminated food and water, or through ingestion of a substance that has come into contact with the stool (feces) of an infected person or animal. Parasites live and reproduce within the tissues and organs of infected human and animal hosts, and are often excreted in feces. There are different types of parasites, ranging in size from tiny, single-celled, microscopic organisms (protozoa), to larger, multi-cellular worms (helminths) that may be seen without a microscope. Examples of common parasites which may be treated by the method of the present invention include, without limitation, *Giardia duodenalis*, *Cryptosporidium parvum*, *Cyclospora cayetanensis*, and *Toxoplasma gondii*.

[0082] Viruses are unlike fungi and bacteria, lacking many of the attributes of free-living cells. A single virus particle is a static structure, quite stable and unable to change or replace its parts. Only when associated with a cell does a virus become capable of replicating and acquiring some of the attributes of a living system. Viruses cause numerous diseases, including such upper respiratory tract infections (URTIs) as the common cold and pharyngitis (sore throat). Other examples of viruses which may be treated by the method of the present invention include, without limitation, viruses associated with AIDS, avian flu,

chickenpox, cold sores, common cold, gastroenteritis (especially in children), glandular fever, influenza, measles, mumps, pharyngitis, pneumonia, rubella, SARS, and lower respiratory tract infection (*e.g.*, respiratory syncytial virus, or RSV)).

[0083] The inventors have demonstrated herein that peptides comprising the amino acid sequence of TABLE 1 or SEQ. ID. NO. 1, 3-16, 18-60, or analogue, derivative, or variant thereof or obvious chemical equivalent thereof have efficacy in the prevention and/or treatment of infection. Accordingly, the present method of treating and/or preventing infection in a subject comprises administering to the subject a peptide comprising the amino acid sequence of TABLE 1 or SEQ. ID. NO. 1, 3-16, 18-60, or analogue, derivative, or variant thereof or obvious chemical equivalent thereof. It is within the confines of the present invention that the peptide of the invention may be linked to another agent or administered in combination with another agent, such as an antibiotic (*e.g.*, penicillin, methicillin, or vancomycin), in order to increase the effectiveness of the treatment and/or prevention of infection, and/or increase the efficacy of targeting.

[0084] In one embodiment of the present invention, the peptide of the invention comprises the amino acid sequence of TABLE 1 or SEQ. ID. NO. 1, 3-16, 18-60, or analogue, derivative, or variant thereof or obvious chemical equivalent thereof. In another embodiment, the peptide of the invention modulates innate immunity in the subject, thereby treating and/or preventing the infection in the subject. The innate immune response is the front line response to a pathogen encounter. It comprises a multiplicity of mechanisms to prevent development of infectious disease. One such mechanism involves the priming and recruitment of immune effector cells.

[0085] In one embodiment, the peptides of the invention can enhance innate immunity or the innate immune response, while limiting inflammation.

[0086] In another embodiment, the peptides of the invention have been shown to be modulators of DPPIV activity. They have been shown to reduce DPPIV activity. As such, they would be useful in the screening of subjects who may benefit from administration of the peptides to treat a particular immunological condition, comprising taking a sample from a subject suspected or known to have a DPPIV-related condition, incubating it together with a peptide of the invention and a DPPIV substrate and then monitoring the effect of the peptide

on DPPIV activity in comparison to a control wherein a reduction in activity would indicate the potential benefit of administration of the peptide to the subject to treat a DPPIV-related condition. In another embodiment modulation of DPPIV activity in the presence of the peptide as compared to the control can be indicative of a DPPIV-related condition. As such, the peptides of the invention can be used in the diagnosis of DPPIV-related conditions. In another aspect the peptides of the invention would be useful in the treatment of a number of immunological disorders, such as DPPIV-related disorder, such as : HIV/AIDS, Grave's disease, cancer (such as lung and colon cancer), diabetes, and autoimmune disorders such as rheumatoid arthritis and multiple sclerosis.

Administration

[0087] In accordance with the method of the present invention, a peptide of the present invention as described herein may be administered to the subject directly, in an amount effective to treat and/or prevent infection in the subject and or to treat or prevent a DPPIV-related condition, e.g. a therapeutic effective amount. Similarly, a peptide as described herein may be administered to the subject indirectly, by administering to the subject a nucleic acid sequence encoding the peptide, in a manner permitting expression of the peptide in the subject, and in an amount effective to treat and/or prevent infection.

[0088] Furthermore, a peptide of the invention, or a nucleic acid molecule encoding same, may be administered to a subject in an amount effective to treat the infection in the subject. As used herein, the phrase "effective to treat the infection" means effective to ameliorate or minimize the clinical impairment or symptoms resulting from infection (by a bacterium, fungus, parasite, virus, etc.). For example, where the subject is infected with a microbe, the amount of peptide (or nucleic acid encoding same) which is effective to treat the microbial infection is that which can ameliorate or minimize the symptoms of the microbial infection, including, without limitation, headache, stiff neck, anorexia, nausea, vomiting, diarrhea, abdominal discomfort, acute renal failure, changing manifestations of ischemic damage to multiple organs, fever, and thrombocytopenia. The amount of peptide (or nucleic acid encoding same) effective to treat an infection in a subject will vary depending on the particular factors of each case, including the subject's weight and the severity of the subject's condition. The appropriate amount of peptide (or nucleic acid

encoding same) can be readily determined by the skilled artisan. Similarly the amount effective to treat a DPPIV-related condition can vary depending on a number of similar factors known to a person skilled in the art.

[0089] Similarly, in the method of the present invention, a peptide of the invention, or a nucleic acid molecule encoding same, may also be administered to a subject at risk of developing an infection, in an amount effective to prevent the infection in the subject. As used herein, the phrase "effective to prevent the infection" includes effective to hinder or prevent the development or manifestation of clinical impairment or symptoms resulting from infection (by a bacterium, fungus, parasite, virus, etc.). The amount of peptide (or nucleic acid encoding same) effective to prevent an infection in a subject will vary depending on the particular factors of each case, including the subject's sex, weight and the severity of the subject's condition, nature of condition, site of infection, and mode of administration. The appropriate amount of peptide (or nucleic acid encoding same) can be readily determined by the skilled artisan.

[0090] The peptide of the invention, or the nucleic acid sequence encoding same, as disclosed herein, may be administered to a human or animal subject by known procedures, including, without limitation, oral administration, parenteral administration (*e.g.*, epifascial, intracapsular, intracutaneous, intradermal, intramuscular, intraorbital, intraperitoneal, intraspinal, intrasternal, intravascular, intravenous, parenchymatous, or subcutaneous administration), transdermal administration, intranasal administration, pulmonary administration (*e.g.*, intratracheal administration), and administration by osmotic pump. In one embodiment, the method of administration is parenteral administration, by intravenous or subcutaneous injection.

[0091] For oral administration, the formulation of the peptide (or nucleic acid encoding same) may be presented as capsules, tablets, powders, granules, or as a suspension or liquid. The formulation may have conventional additives, such as lactose, mannitol, corn starch, or potato starch. The formulation also may be presented with binders, such as crystalline cellulose, cellulose derivatives, acacia, corn starch, or gelatins. Additionally, the formulation may be presented with disintegrators, such as corn starch, potato starch, or sodium carboxymethylcellulose. The formulation may be further presented with dibasic

calcium phosphate anhydrous or sodium starch glycolate. Finally, the formulation may be presented with lubricants, such as talc or magnesium stearate.

[0092] For parenteral administration, the peptide (or nucleic acid encoding same) may be combined with a sterile aqueous solution, which is preferably isotonic with the blood of the subject. Such a formulation may be prepared by dissolving a solid active ingredient in water containing physiologically-compatible substances, such as sodium chloride, glycine, and the like, and having a buffered pH compatible with physiological conditions, so as to produce an aqueous solution, then rendering said solution sterile. The formulation may be presented in unit or multi-dose containers, such as sealed ampoules or vials. The formulation also may be delivered by any mode of injection, including any of those described herein.

[0093] For transdermal administration, the peptide (or nucleic acid encoding same) may be combined with skin penetration enhancers, such as propylene glycol, polyethylene glycol, isopropanol, ethanol, oleic acid, N-methylpyrrolidone, and the like, which increase the permeability of the skin to the peptide or nucleic acid, and permit the peptide or nucleic acid to penetrate through the skin and into the bloodstream. The composition of enhancer and peptide or nucleic acid also may be further combined with a polymeric substance, such as ethylcellulose, hydroxypropyl cellulose, ethylene/vinylacetate, polyvinyl pyrrolidone, and the like, to provide the composition in gel form, which may be dissolved in solvent, such as methylene chloride, evaporated to the desired viscosity, and then applied to backing material to provide a patch. The peptide or nucleic acid may be administered transdermally, at or near the site on the subject where the infection may be localized. Alternatively, the peptide or nucleic acid may be administered transdermally at a site other than the affected area, in order to achieve systemic administration.

[0094] For intranasal administration (e.g., nasal sprays) and/or pulmonary administration (administration by inhalation), formulations of the peptide or nucleic acid, including aerosol formulations, may be prepared in accordance with procedures well known to persons of skill in the art. Aerosol formulations may comprise either solid particles or solutions (aqueous or non-aqueous). Nebulizers (e.g., jet nebulizers, ultrasonic nebulizers, etc.) and atomizers may be used to produce aerosols from solutions (e.g., using a solvent such as ethanol); metered-dose inhalers and dry-powder inhalers may be used to generate

small-particle aerosols. The desired aerosol particle size can be obtained by employing any one of a number of methods known in the art, including, without limitation, jet-milling, spray drying, and critical-point condensation.

[0095] Pharmaceutical compositions for intranasal administration may be solid formulations (*e.g.*, a coarse powder), and may contain excipients (*e.g.*, lactose). Solid formulations may be administered from a container of powder held up to the nose, using rapid inhalation through the nasal passages. Compositions for intranasal administration may also comprise aqueous or oily solutions of nasal spray or nasal drops. For use with a sprayer, the formulation of peptide or nucleic acid may comprise an aqueous solution and additional agents, including, for example, an excipient, a buffer, an isotonicity agent, a preservative, or a surfactant. A nasal spray may be produced, for example, by forcing a suspension or solution of the peptide or nucleic acid through a nozzle under pressure.

[0096] Formulations of the peptide or nucleic acid for pulmonary administration may be presented in a form suitable for delivery by an inhalation device, and may have a particle size effective for reaching the lower airways of the lungs or sinuses. For absorption through mucosal surfaces, including the pulmonary mucosa, the formulation of the present invention may comprise an emulsion that includes, for example, a bioactive peptide, a plurality of submicron particles, a mucoadhesive macromolecule, and/or an aqueous continuous phase. Absorption through mucosal surfaces may be achieved through mucoadhesion of the emulsion particles.

[0097] Pharmaceutical compositions for use with a metered-dose inhaler device may include a finely-divided powder containing the peptide or nucleic acid as a suspension in a non-aqueous medium. For example, the peptide or nucleic acid may be suspended in a propellant with the aid of a surfactant (*e.g.*, sorbitan trioleate, soya lecithin, or oleic acid). Metered-dose inhalers typically use a propellant gas (*e.g.*, a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon) stored in a container (*e.g.*, a cannister) as a mixture (*e.g.*, as a liquefied, compressed gas). Inhalers require actuation during inspiration. For example, actuation of a metering valve may release the mixture as an aerosol. Dry-powder inhalers use breath-actuation of a mixed powder.

[0098] The peptide or nucleic acid of the present invention also may be released or delivered from an osmotic mini-pump or other timed-release device. The release rate from an elementary osmotic mini-pump may be modulated with a microporous, fast-response gel disposed in the release orifice. An osmotic mini-pump would be useful for controlling release, or targeting delivery, of the peptide or nucleic acid.

[0099] In accordance with methods described herein, the peptide of the invention may be administered to a subject by introducing to the subject the peptide itself, or by introducing to the subject a nucleic acid encoding the peptide in a manner permitting expression of the peptide. Accordingly, in one embodiment of the present invention, infection in a subject may be treated or prevented by administering to the subject an amount of a peptide of the invention. In a further embodiment of the present invention, infection in the subject may be treated or prevented by administering to the subject a nucleic acid sequence encoding a peptide of the invention, in a manner permitting expression of the peptide in the subject.

[00100] The peptides of the present invention may be administered or introduced to a subject by known techniques used for the introduction of proteins and other drugs, including, for example, injection and transfusion. Where an infection is localized to a particular portion of the body of the subject, it may be desirable to introduce the therapeutic peptide directly to that area by injection or by some other means (*e.g.*, by introducing the peptide into the blood or another body fluid). The amount of peptide to be used is an amount effective to treat and/or prevent the infection in the subject, as defined above, and may be readily determined by the skilled artisan.

[00101] In the method of the present invention, the peptide also may be administered or introduced to the subject by introducing into a sufficient number of cells of the subject a nucleic acid encoding the peptide, in a manner permitting expression of the peptide. The amount of nucleic acid encoding the therapeutic peptide is an amount that will produce the peptide in an amount effective to treat and/or prevent infection, as defined above, in the subject. This amount may be readily determined by the skilled artisan.

[00102] Nucleic acid encoding the peptide of the present invention may be introduced to the subject using conventional procedures known in the art, including, without limitation,

electroporation, DEAE Dextran transfection, calcium phosphate transfection, lipofection, monocationic liposome fusion, polycationic liposome fusion, protoplast fusion, creation of an *in vivo* electrical field, DNA-coated microprojectile bombardment, injection with recombinant replication-defective viruses, homologous recombination, *in vivo* gene therapy, *ex vivo* gene therapy, viral vectors, naked DNA transfer, or any combination thereof. Recombinant viral vectors suitable for gene therapy include, but are not limited to, vectors derived from the genomes of viruses such as retrovirus, HSV, adenovirus, adeno-associated virus, Semiliki Forest virus, cytomegalovirus, and vaccinia virus.

[00103] It is also within the confines of the present invention that a nucleic acid encoding a peptide of the invention may be introduced into suitable cells *in vitro*, using conventional procedures, to achieve expression of the therapeutic peptide in the cells. Cells expressing the peptide then may be introduced into a subject to treat and/or prevent infection *in vivo*. In such an *ex vivo* gene therapy approach, the cells are preferably removed from the subject, subjected to DNA techniques to incorporate nucleic acid encoding the therapeutic peptide, and then reintroduced into the subject.

[00104] It is also within the confines of the present invention that a formulation containing a peptide of the invention, or a nucleic acid encoding same, may be further associated with a pharmaceutically-acceptable carrier, diluent, or excipient, thereby comprising a pharmaceutical composition. Pharmaceutical compositions of the invention, and exemplary carriers, diluents, and excipients, are described above.

[00105] The formulations of the present invention may be prepared by methods well-known in the pharmaceutical arts. For example, the peptide of the invention, or a nucleic acid encoding same, may be brought into association with a carrier, diluent, or excipient, as a suspension or solution. Optionally, one or more accessory ingredients (*e.g.*, buffers, flavoring agents, surface active agents, and the like) also may be added. The choice of carrier will depend upon the route of administration. The pharmaceutical composition would be useful for administering the peptide of the present invention, or a nucleic acid molecule encoding same, to a subject, in order to treat and/or prevent infection. The peptide or nucleic acid is provided in an amount that is effective to treat and/or prevent infection in a subject to

whom the pharmaceutical composition is administered. This amount may be readily determined by the skilled artisan, as described above.

Diagnostics and Screening Assays

[00106] The present invention provides a method for diagnosing a subject who is suspected of having an innate immune condition, or DPPIV-related condition, for predicting whether a subject would be responsive to treatment with a peptide of the invention, such as those listed in TABLE 1, or an analogue, derivative or variant thereof, and to screening for agents that would modulate (e.g., enhance, inhibit or mimic) the immunological effect of the peptides of the invention. In another embodiment, the invention provides methods for screening for immunologically active analogues, derivatives, and variants of the peptides of the invention or those listed in TABLE 1 or to immunologically active modifications thereof.

[00107] In one embodiment, a method for predicting whether a patient with a immunological disorder, such as an innate immune-related condition would be responsive to treatment with a peptide of the invention comprises obtaining a biological sample from the subject, administering a peptide of the invention to said sample, and monitoring levels of a predetermined marker that is indicative of the condition, such as DPPIV for a DPPIV-related condition, an inflammatory biomarker for an infection, cell viability or bacterial load, in comparison to a positive and/or negative control. The positive control can be a sample from a subject with a known immunological condition. A negative control can be a sample from the same subject that is not administered the peptide. If the peptide modulates the activity, level of marker, or cell viability in relation to the control, the subject may have such immunological disorder and may benefit from treatment with the peptide.

[00108] More particularly, in one aspect of the invention, if the subject has or is suspected of having a DPPIV-related condition, then monitoring DPPIV activity as the marker for the condition would be appropriate. In one aspect, reduction of DPPIV activity in comparison to the control would be indicative that the subject would be responsive to treatment with the peptide. Alternatively, if the subject has or was suspected of having an infection, then obtaining a sample from the patient, monitoring it for pathogen load or cell viability in comparison to a sample from the patient after administration of the peptide, wherein pathogen load is less or cell viability is higher in the patient after administration of

the peptide, is indicative that the subject would benefit from peptide treatment or has an immunological disorder.

[00109] In another embodiment, if one wishes to see whether a peptide or modification of a peptide of TABLE 1 or other agent would have the same immunological activity as the peptide of the invention, one can monitor the effect of the peptide on DPPIV activity in comparison to the reference peptide with known modulatory affects, on a sample (either from a mouse infected with an agent or a known DPPIV-related condition), or to monitor prevention, administration of the peptide or agent to a sample, inducing said infection or DPPIV-related condition in said sample and then monitoring whether the peptide modulated or inhibited development of said infection or DPPIV-related condition, or immune response. Said sample can be an animal model, wherein induction of the condition or infection is done in an accepted animal model in accordance with ethical guidelines and then the animal or appropriate biological sample of the animal is screened for effect of the peptide.

[00110] The present invention further provides a method for predicting whether a subject would be responsive to treatment for a microbial infection wherein the treatment comprises administering to the subject a peptide comprising an amino acid sequence of TABLE 1 or an analogue, derivative, or variant thereof. The method includes assaying a diagnostic sample of the subject for one or more biomarkers (such as an inflammatory biomarker), wherein the presence of at least one biomarker (such as an inflammatory biomarker) is indicative that the subject would be responsive to the treatment.

[00111] As used herein a "biomarker" or "marker is any suitable biomarker known to be, or recognized as being, related to the condition (e.g. immune condition, infection, inflammatory condition, DPPIV-related condition, innate immune condition), and includes any molecule derived from a gene (e.g., a transcript of the gene), a sense (coding) or antisense (non-coding) probe sequence derived from a gene, or a partial-length or full-length translation product of a gene, or an antibody thereto, which can be used to monitor a condition, disorder, or disease associated with the immune response, innate immune response, inflammation, and/or a DPPIV-related condition.

[00112] According to the method of the present invention, the diagnostic sample of a subject may be assayed *in vitro* or *in vivo*. Where the assay is performed *in vitro*, a diagnostic sample from the subject may be removed using standard procedures. The diagnostic sample may be tissue, including any muscle tissue, skin tissue, or soft tissue, which may be removed by standard biopsy. In addition, the diagnostic sample may be a bodily fluid, including blood, saliva, serum, or urine. The subject or patient may be known to have a microbial infection or other immunological disorder such as a DPPIV-related condition, suspected of having a microbial infection or other immunological condition, such as an innate immune condition or DPPIV-related condition, or believed not to have a microbial infection or other immunological condition, such as an innate-immune condition, or DPPIV-related condition..

[00113] In accordance with the method of the present invention, a diagnostic sample of the subject may be assayed for expression of one or more desired markers. As used herein, "expression" means the transcription of an inflammatory-marker gene into at least one mRNA transcript, or the translation of at least one mRNA into a marker protein. Accordingly, a diagnostic sample may be assayed for marker expression by assaying for a marker protein, marker cDNA, or marker mRNA. The appropriate form of the marker will be apparent based on the particular techniques discussed herein.

[00114] Protein to be assayed may be isolated and purified from the diagnostic sample of the subject or patient using standard methods known in the art, including, without limitation, extraction from a tissue (*e.g.*, with a detergent that solubilizes the protein) where necessary, followed by affinity purification on a column, chromatography (*e.g.*, FPLC and HPLC), immunoprecipitation (with an antibody to an inflammatory marker of interest), and precipitation (*e.g.*, with isopropanol and a reagent such as Trizol). Isolation and purification of the protein may be followed by electrophoresis (*e.g.*, on an SDS-polyacrylamide gel). It is contemplated that the diagnostic sample may be assayed for expression of any or all forms of marker protein (including precursor, endoproteolytically-processed forms, and other forms resulting from post-translational modification). Nucleic acid may be isolated from a diagnostic sample using standard techniques known to one of skill in the art.

[00115] In accordance with the method of the present invention, a diagnostic sample of a subject may be assayed for marker expression, and marker expression may be detected in a diagnostic sample, using assays and detection methods readily determined from the known art (*e.g.*, immunological techniques, hybridization analysis, fluorescence imaging techniques, and/or radiation detection), as well as any assays and detection methods disclosed herein (*e.g.*, immunoprecipitation, Western blot analysis, etc.). For example, a diagnostic sample of a subject may be assayed for marker expression using an agent reactive with an inflammatory marker. As used herein, "reactive" means the agent has affinity for, binds to, or is directed against the marker. As further used herein, an "agent" shall include a protein, polypeptide, peptide, nucleic acid (including DNA or RNA), antibody, Fab fragment, F(ab')₂ fragment, molecule, compound, antibiotic, drug, and any combination(s) thereof. Preferably, the agent of the present invention is labeled with a detectable marker or label, in accordance with techniques described herein. In one embodiment of the present invention, the agent reactive with a marker is an antibody.

[00116] Where the agent of the present invention is an antibody reactive with the desired marker, a diagnostic sample taken from the subject may be purified by passage through an affinity column which contains antibody to the marker, attached as a ligand to a solid support (*e.g.*, an insoluble organic polymer in the form of a bead, gel, or plate). The antibody attached to the solid support may be used in the form of a column. Examples of suitable solid supports include, without limitation, agarose, cellulose, dextran, polyacrylamide, polystyrene, sepharose, and other insoluble organic polymers. The antibody to the marker may be further attached to the solid support through a spacer molecule, if desired. Appropriate binding conditions (*e.g.*, temperature, pH, and salt concentration) for ensuring binding of the agent and the antibody may be readily determined by the skilled artisan. In a preferred embodiment, the antibody to the marker is attached to a sepharose column, such as Sepharose 4B.

[00117] Additionally, where the agent is an antibody, a diagnostic sample of the subject may be assayed for expression of the immunological marker using binding studies that utilize one or more antibodies immunoreactive with the marker, along with standard immunological detection techniques. For example, the marker protein eluted from the affinity column may be subjected to an ELISA assay, Western blot analysis, flow cytometry,

or any other immunostaining method employing an antigen-antibody interaction. Preferably, the diagnostic sample is assayed for marker expression using Western blotting.

[00118] Alternatively, a diagnostic sample of a subject may be assayed for -marker expression using hybridization analysis of nucleic acid extracted from the diagnostic sample taken from the subject. According to this method of the present invention, the hybridization analysis may be conducted using Northern blot analysis of mRNA. This method also may be conducted by performing a Southern blot analysis of DNA using one or more nucleic acid probes which hybridize to nucleic acid encoding the marker. The nucleic acid probes may be prepared by a variety of techniques known to those skilled in the art, including, without limitation, the following: restriction enzyme digestion of -marker nucleic acid; and automated synthesis of oligonucleotides having sequences which correspond to selected portions of the nucleotide sequence of the marker nucleic acid, using commercially-available oligonucleotide synthesizers, such as the Applied Biosystems Model 392 DNA/RNA synthesizer.

[00119] Nucleic acid probes used in the present invention may be DNA or RNA, and may vary in length from about 8 nucleotides to the entire length of the inflammatory-marker nucleic acid. In addition, the nucleic acid probes of the present invention may be labeled with one or more detectable markers or labels. Labeling of the nucleic acid probes may be accomplished using one of a number of methods known in the art, including any of those described herein. Combinations of two or more nucleic acid probes (or primers), corresponding to different or overlapping regions of the marker nucleic acid, also may be used to assay a diagnostic sample for marker expression, using, for example, PCR or RT-PCR.

[00120] The detection of marker expression in the method of the present invention may be followed by an assay to measure or quantify the extent of marker expression in a diagnostic sample of a subject. Such assays are well known to one of skill in the art, and may include immunohistochemistry/immunocytochemistry, flow cytometry, mass spectroscopy, Western blot analysis, or an ELISA for measuring amounts of marker protein. For example, to use an immunohistochemistry assay, histological (paraffin-embedded) sections of tissue may be placed on slides, and then incubated with an antibody against a

marker. The slides then may be incubated with a second antibody (against the primary antibody), which is tagged to a dye or other calorimetric system (e.g., a fluorochrome, a radioactive agent, or an agent having high electron-scanning capacity), to permit visualization of the marker present in the sections.

[00121] The present invention is described in the following Examples, which are set forth to aid in the understanding of the invention, and should not be construed to limit in any way the scope of the invention as defined in the claims which follow thereafter.

EXAMPLES

EXAMPLE 1 -PEPTIDE SYNTHESIS

[00122] The peptides in TABLE 1 were synthesized using a solid phase peptide synthesis technique.

[00123] All the required Fmoc-protected amino acids were weighed in three-fold molar excess relative to the 1 mmole of peptide desired. The amino acids were then dissolved in Dimethylformamide (DMF) (7.5 ml) to make a 3mMol solution. The appropriate amount of Rink amide MBHA resin was weighed taking in to account the resin's substitution. The resin was then transferred into the automated synthesizer reaction vessel and was pre-soaked with Dichloromethane (DCM) for 15 minutes.

[00124] The resin was de-protected by adding 25% piperidine in DMF (30 ml) to the resin and mixing for 20 minutes. After de-protection of the resin the first coupling was made by mixing the 3mMol amino acid solution with 4mMol 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and 8mMol N,N-diisopropylethylamine (DIEPA). The solution was allowed to pre-activate for 5 minutes before being added to the resin. The amino acid was allowed to couple for 45 minutes.

[00125] After coupling the resin was thoroughly rinsed with DMF and Dimethylacetamide (DMA). The attached Fmoc protected amino acid was deprotected in the same manner described above and the next amino acid was attached using the same coupling scheme AA:HBTU:DIEPA.

[00126] After the completion of the synthesis the peptide was cleaved from the resin with the use of a cleavage cocktail containing 97.5 % Trifluoroacetic acid (TFA) and 2.5% water. The resin was allowed to swim in the cleavage cocktail for 1 ½ hours. The solution was then filtered by gravity using a Buchner funnel and the filtrate was collected in a 50 ml centrifugation tube. The peptide was isolated by precipitating with chilled diethyl ether. After centrifuging and decanting diethyl ether the crude peptide was washed with diethyl ether once more before being dried in a vacuum desiccator for 2 hours. The peptide was then dissolved in de-ionized water (10 ml), frozen at -80°C and lyophilized. The dry peptide was then ready for HPLC purification.

[00127] Due to the hydrophilic nature of these peptides the diethyl ether peptide isolation did not work. Therefore a chloroform extraction was required. The TFA was evaporated and the resulting peptide residue was dissolved in 10% acetic acid (15 ml). The impurities and scavengers were removed from the acetic acid peptide solution by washing the solution twice with chloroform (30ml). The aqueous peptide solution was then frozen at -80°C and lyophilized resulting in a powdered peptide ready for HPLC purification.

[00128] Peptides SEQ. ID. NOs. 33 and 34 each contained one N-methyl amino acid. This coupling was carried out by combining the N-methyl amino acid, PyBroP and N-hydroxybenzotriazole*H₂O (HOBt) and DIEPA solutions together in the RV containing the resin. After allowing to couple for 45 minutes the N-methyl amino acid was then doubled coupled to ensure complete coupling. It was observed that the coupling following the N-methyl amino acid was not fully complete. Therefore this coupling was performed using N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (HATU) instead of HBTU. This still resulted in a crude peptide that typically contained two impurities totaling 30-40% of the total purity. The peptide was purified under modified HPLC conditions to isolate the pure peptide peak away from the closely eluting impurities.

EXAMPLE 2 – NON-ANTIMICROBIAL ACTIVITY

[00129] Bacteria (*S. aureus* 25923) were seeded into wells containing peptide (200 µM), vehicle (Tris), or antibiotic (erythromycin; 120 µg/ml). The bacteria were allowed to grow for 2 hours. Thereafter, bacterial viability was determined utilizing a WST-1

colorimetric viability assay (catalogue number 1 644 807; Roche Diagnostics). DMEM and DMEM+WST-1 were included as background controls. As shown in FIG. 1 A and B, the peptide of SEQ ID NOs: 5 and 47 clearly show a lack of activity, as compared with an antibiotic control.

EXAMPLE 3 – *IN VIVO* PROTECTION

[00130] Mice were infected with *S. aureus* 25923 *via* intraperitoneal (IP) injection. Four hours later, the peptide of SEQ ID NOs:1, 4, 5, 6, 45, and 47 were administered at 12 mg/kg and 24 mg/kg for SEQ. ID. NO. 1 (FIG. 2A and 2B), 9.6 mg/kg for SEQ. ID. NO. 5 (FIG. 2C), 13 mg/kg for SEQ. ID. NO. 47 (FIG. 2D), 12 mg/kg for SEQ. ID. NO. 4 (FIG. 2E), 9 mg/kg for SEQ. ID. NO. 6 (FIG. 2F), and 13 mg/kg for SEQ. ID. NO. 45 (FIG. 2G), *via* IP injection. Twenty-four hours post-infection, surviving animals were sacrificed, and intraperitoneal lavage fluid was plated to determine residual bacterial counts (# colony forming units per ml (CFU/ml)) in the presence and absence of peptide treatment.

[00131] Dead animals were assigned the highest bacterial count of any animal in the study. The peptide of SEQ ID NOs:1, 4, 5, 6, 45, and 47 clearly demonstrated protection, as compared with the control. as shown in FIG. 2 A – G.

EXAMPLE 4 – PROPHYLACTIC *IN VIVO* PROTECTION

[00132] Twenty-four hours prior to infection, peptide was administered at 12 mg/kg (SEQ. ID. NO. 1, FIG. 3A) and 11.5 mg/kg (SEQ. ID. No. 5, FIG. 3B), *via* IP injection. Mice were then infected with *S. aureus* 25923 *via* IP injection. Twenty-four hours post-infection, surviving animals were sacrificed and intraperitoneal lavage fluid was plated to determine residual bacterial counts (# colony forming units per ml (CFU/ml)) in the presence and absence of peptide treatment.

[00133] Dead animals were assigned the highest bacterial count of any animal in the study. The peptides of SEQ. ID. NO:1 and 5 clearly demonstrated protection ((0 mouse dead (peptide treatment) vs. 2 mice dead (control)). Please see Figures 3 A and B.

[00134] Discussed below are results obtained by the inventors in connection with the experiments of Examples 1-4:

[00135] The inventors have shown that a peptide having the amino acid sequence of those shown in TABLE 1 or as described herein as part of the invention can enhance innate immunity. Specifically, the peptides of SEQ ID NOs:1, 4, 5, 6, 45, and 47 had the ability to prevent and protect against infection, as demonstrated in *in vivo* models (FIG. 2 and Example 3; FIG. 3 and Example 4) However, the peptide of SEQ ID NOs:5 and 47 lacked antimicrobial activity, as shown in Example 1 and FIG. 1. Accordingly, modulation of innate immunity, *via* the peptide of SEQ ID NOs:5 and/or 47, indicate that these peptides can be used as a therapeutic for the treatment of infectious disease.

EXAMPLE 5 -PLASMA DPPIV ACTIVITY ASSAY WITH MOUSE BLOOD

[00136] Mouse blood was obtained by cardiac puncture from ICR mice and collected in heparinized blood collection tubes. Blood from several mice was pooled and aliquoted into 300 μ L aliquots. The peptide was dissolved in acetate buffered saline, pH 5.5, to a concentration of 9 mM. Of this stock solution 30 μ L were added to 300 μ L of blood and mixed by resuspension (concentration in blood 0.82 mM). For the control, 30 μ L of blank acetate buffered saline was added to 300 μ L of blood. Each peptide group was prepared in triplicate, whereas the control was prepared in six replicates. The samples were incubated at 37°C in closed microtubes for two hours. After incubation the plasma was isolated from the samples by centrifugation at 4000 rcf. The plasma was transferred to a 96-well assay plate for the DPPIV assay. The assay was started by adding 5 μ L of the DPPIV substrate gly-pro-p-nitroanilide (16 mM in de-ionized water) to 95 μ L of plasma (concentration in plasma 0.8 mM) and the increase in UV absorbance (405 nm) was monitored over a 20 min time period. The rate of the production of p-nitroaniline by enzymatic cleavage of gly-pro-p-nitroanilide was taken as the activity of DPPIV (Durinx C et al., (2001) "Reference values for plasma dipeptidyl-peptidase IV activity and their association with other laboratory parameters". Clin Chem Lab Med. 39(2):155-9.)

[00137] The results can be seen in TABLE 1. The effect of the peptides on the activity of DPPIV was observed. Results are presented as normalized, averaged %activity relative to saline control (set to 100%). Anything less than 100% activity represents a reduction in DPPIV activity.

[00138] In one aspect of the invention, a reduction by of DPPIV activity by about, or in one embodiment, at least, 25% (i.e. to about 75%) was deemed to be active. A person skilled in the art would appreciate that the desired level of activity may vary depending on the use of the peptides.

Discussion

[00139] The type II transmembrane serine protease dipeptidyl peptidase IV (DPPIV), also known as CD26 or adenosine deaminase binding protein, is a major regulator of various physiological processes including immune functions. CD26/DPPIV is a 110-kD cell surface glycoprotein that is mainly expressed on mature thymocytes, activated T-cells, B-cells, NK-cells, macrophages, and epithelial cells. It has at least two functions, a signal transduction function and a proteolytic function (Morimoto C, Schlossman SF. The structure and function of CD26 in -. The T-cell immune response. Immunol. Review. 1998, 161: 55-70.). One of its cellular roles involves modulation of chemokine activity by cleaving dipeptides from the chemokine N-terminus. The modulation of the NH₂ termini of chemokines is of great importance not only for binding to their receptors and the following reactions but also for altering the receptor specificity of the processed chemokine. Furthermore, it was demonstrated that soluble rCD26 enhances transendothelial migration of T cells whereas it reduces the migratory response of monocytes [Oravecz, T. et. al., (1997) Regulation of the receptor specificity and formation of the chemokine RANTES (regulated on activation, normal T cell expressed and secreted) by dipeptidyl peptidase IV (CD26)-mediated cleavage. J. Exp. Med. 186:1865-1872; Iwata, S., et. al., (1999) CD26/dipeptidyl peptidase IV differentially regulates the chemotaxis of T cells and monocytes toward RANTES: possible mechanism for the switch from innate to acquired immune response. Int. Immunol. 11:417-426). These results indicate that CD26/DPPIV differentially regulate the chemotactic response of T cells and monocytes and is involved in the switch from innate to acquired immune response. As such, a reduction in activity of DPPIV would then have the opposite effect, promoting an innate immune response and macrophage migratory responses. It has also been reported that pharmacological inhibition of DPPIV enzyme activity could reduce the progression of arthritis in an experimental rat model of RA (Tanaka S et al., Anti-arthritis effects of the novel dipeptidyl peptidase IV inhibitors TMC-2A and TSL-225.

Immunopharmacology 1998, 40:21-26; Tanaka S, et al.,: Suppression of arthritis by the inhibitors of dipeptidyl peptidase IV. Int J Immunopharmacol 1997, 19:15-24), suggesting that decreases in DPPIV-activity may alleviate inflammation under some circumstances. Together the anti-inflammatory role and its modulation of chemokine activity make DPPIV a good molecule for screening novel compounds for these activities.

[00140] CD26/DPPIV is involved in the pathology of a variety of diseases, such as AIDS and HIV disease progression (Blazquez et al. 1992; Vanham et al. 1993; Schols et al. 1998 Oravec et al. 1995), 'Graves' disease (Eguchi et al. 1989; Nishikawa et al. 1995), and cancer (Stecca et al. 1997) and diabetes (Hinke et al. 2000; Marguet et al. 2000).

[00141] Further, CD26 as an indicator of T-cell activation has been shown to fluctuate in parallel with several autoimmune diseases such as rheumatoid arthritis (Nakao et al., 1989) and autoimmune thyroiditis (Eguchi et al., 1989). CD26 has been described as a marker that correlates well with the level of activity of these diseases. It has furthermore been studied as an indicator of disease progression in chronic progressive multiple sclerosis (Constantinescu et al., 1995).

[00142] The peptides of the present invention have demonstrated that they can reduce the activity of DPPIV. As such, they can be used in the treatment of certain immunological conditions, such as DPPIV- related or associated conditions, and may, in one aspect modulate innate immunity and inflammation, such as inflammation leading to sepsis.

[00143] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art, from a reading of the disclosure, that various changes in form and detail can be made without departing from the true scope of the invention in the appended claims.

TABLE 1

all C-terminal amidated unless otherwise indicated****

SEQ ID	Description	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Length	Net charge	%DPPV Activity (Saline)**
1																	6	3	74
2																	6	2	
3																	6	2	54
4																	5	2	62
5																	5	2	68
6																	5	2	62
7																	5	2	50
8																	4	2	58
9																	4	2	54
10																	5	1	51
11																	5	2	41
12																	6	3	58
13																	5	3	49
14																	5	3	67
15																	5	2	20
16																	5	2	71
17																	5	2	
18																	7	2	67
19																	6	1	67
20																	5	1	72
21																	5	2	64
22																	5	2	88
23																	5	2	69
24																	5	2	72
25																	5	27	40
26																	5	2	-11
27																	5	2	68
28																	5	2	19
29																	5	2	-11
30																	5	2	33
31																	5	1	
32																	7	1	27
33																	5	2	70
34																	5	2	75
35																	5	2	74
36																	5	1	

TABLE 1

all C-terminal amidated unless otherwise indicated****

SEQ ID	Description	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	Length	Net charge	%DPPIV Activity (saline)**
37																	5	2	39
38																	5	17	51
39																	6	3	46
40																	5	2	66
41																	5	2	60
42																	5	2	48
43																	5	1	73
44																	13	3	73
45																	7	3	32
46																	6	2	69
47																	6	3	50
48																	5	2	68
49																	4	2	64
50																	8	2	65
51																	3	2	73
52																	6	1	71
53																	6	27	69
54																	11	2	-2
55	See Note 1																	3	
56	See Note 2																	4	
57	See Note 3																	5	
58	See Note 4																	5	
59	See Note 5																	6	
60	See Note 6																	6	

** % DPPIV Activity (Saline), where control is 100% activity (saline or vehicle alone without the peptide). About 75% or less activity relative to saline control is desirable.
 **** OH indicates the free acid form of the peptide. Ac indicates acetylated. O indicates Ornithine, Cit indicates Citrulline, x indicates NMe backbone (versus amide backbone).

TABLE 1 CONTINUED

Note 1 of Table 1:

X₁ is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH)

X₂ is selected from the group consisting of V, I, K, P, and H including an isolated peptide of up to 10 amino acids comprising an amino acid sequence of SEQ. ID. NO. 55.

Note 2 of Table 1:

wherein X₁ is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH) and wherein X₂ is selected from the group consisting of A, I, L, V, K, P, G, H, R, S, O, Dab, Dpr, Cit, Hci, Abu, Nva, Nle and where X₂ can be N-methylated, and wherein X₃ is selected from the group consisting of I, V, P, wherein in one embodiment, X₃ is not N-methylated. In one embodiment, the isolated peptide can be an amino acid sequence of up to 10 amino acids, but is not SEQ. ID. NOs. 2 or 17.

Note 3 of Table 1:

wherein X₁, X₂ and X₃ are defined as in Seq. ID. NO. 56, and wherein "a" is selected from the group consisting of S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, M, and F or an isolated peptide up to 10 amino acids comprising said sequences.

Note 4 of Table 1:

wherein X₁X₂X₃P are as defined in SEQ. ID. NO. 56 and "b" is selected from the group consisting of A, A*, G, S, L, F, K, C, I, V, T, Y, R, H, O, and M, but in one embodiment not P. In one embodiment, the isolated peptide is a peptide of up to 10 amino acids comprising SEQ. ID. NO. 58 but not SEQ. ID. NO. 17.

Note 5 of Table 1:

where X_1 , X_2 and X_3 are as defined in SEQ. ID. NO. 56 and " a_1 " is selected from the group consisting of K, I, R, H, O, L, V, A, and G and " a_2 " is selected from the group consisting of S, P, R, T, H, K, O, L, V, A, G, S, and I. In one embodiment, " a_1 " is not acetylated, or where a_1 is K, K is not acetylated or not SEQ. ID. NO. 2. In one embodiment, the isolated peptide comprises up to 10 amino acids comprising SEQ. ID. NO. 59.

Note 6 of Table 1:

where X_1 , X_2 and X_3 are as defined in SEQ. ID. NO. 56 and where " a " is selected from the group consisting of S, R, K, H, O, T, I, L, V, A, and G and wherein " b " is selected from the group consisting of A, V, I, L, G, K, H, R, O, S, T, and F or a peptide of up to 10 amino acids comprising SEQ. ID. NO. 60.

REFERENCES CITED

Blazquez MV, Madueno JA, Gonzalez R, Jurado R, Bachovchin WW, Pena J, Munoz E. Selective decrease of CD26 expression in T cells from HIV-1-infected individuals. *J Immunol.* 1992 Nov 1;149(9):3073-7.

Vanham G, Kestens L, De Meester I, Vingerhoets J, Penne G, Vanhoof G, Scharpe S, Heyligen H, Bosmans E, Ceuppens JL, et al. Decreased expression of the memory marker CD26 on both CD4+ and CD8+ T lymphocytes of HIV-infected subjects. *J Acquir Immune Defic Syndr.* 1993 Jul;6(7):749-57.

Schols D, Proost P, Struyf S, Wuyts A, De Meester I, Scharpe S, Van Damme J, De Clercq E. CD26-processed RANTES(3-68), but not intact RANTES, has potent anti-HIV-1 activity. *Antiviral Res.* 1998 Oct;39(3):175-87. *Erratum in: Antiviral Res* 1999 Jan;40(3):189-90.

Oravec T, Roderiquez G, Koffi J, Wang J, Ditto M, Bou-Habib DC, Lusso P, Norcross MA. CD26 expression correlates with entry, replication and cytopathicity of monocytotropic HIV-1 strains in a T-cell line. *Nat Med.* 1995 Sep;1(9):919-26. *Comment in: Nat Med.* 1995 Sep;1(9):881-2.

Nishikawa Y, Nakamura M, Fukumoto K, Matsumoto M, Matsuda T, Tanaka Y, Yoshihara H. [Adenosine deaminase isoenzymes in patients with Graves' disease] *Rinsho Byori.* 1995 Oct;43(10):1057-60. [Article in Japanese]

Eguchi K, Ueki Y, Shimomura C, Otsubo T, Nakao H, Migita K, Kawakami A, Matsunaga M, Tezuka H, Ishikawa N, et al. Increment in the Ta1+ cells in the peripheral blood and thyroid tissue of patients with Graves' disease. *J Immunol.* 1989 Jun 15;142(12):4233-40.

Stecca BA, Nardo B, Chieco P, Mazziotti A, Bolondi L, Cavallari A. Aberrant dipeptidyl peptidase IV (DPP IV/CD26) expression in human hepatocellular carcinoma. *J Hepatol.* 1997 Aug;27(2):337-45.

Hinke SA, Pospisilik JA, Demuth HU, Mannhart S, Kuhn-Wache K, Hoffmann T, Nishimura E, Pederson RA, McIntosh CH. Dipeptidyl peptidase IV (DPIV/CD26) degradation of glucagon. Characterization of glucagon degradation products and DPIV-resistant analogs. *J Biol Chem.* 2000 Feb 11;275(6):3827-34.

Marguet D, Baggio L, Kobayashi T, Bernard AM, Pierres M, Nielsen PF, Ribet U, Watanabe T, Drucker DJ, Wagtmann N. Enhanced insulin secretion and improved glucose tolerance in mice lacking CD26. *Proc Natl Acad Sci U S A.* 2000 Jun 6;97(12):6874-9.

Nakao H, Eguchi K, Kawakami A, Migita K, Otsubo T, Ueki Y, Shimomura C, Tezuka H, Matsunaga M, Maeda K, et al. Increment of Tal positive cells in peripheral blood from patients with rheumatoid arthritis. *J Rheumatol.* 1989 Jul;16(7):904-10.

Constantinescu CS, Kamoun M, Dotti M, Farber RE, Galetta SL, Rostami A. A longitudinal study of the T cell activation marker CD26 in chronic progressive multiple sclerosis. *J Neurol Sci.* 1995 Jun;130(2):178-82.

CLAIMS

What is claimed is:

1. An isolated peptide consisting of up to 10 amino acids, said peptide comprising the amino acid sequence of

$X_1X_2X_3P$ (SEQ. ID. NO. 56)

wherein

X_1 is R

X_2 is I or V, where it can be N-methylated;

X_3 is I, or V and can be N-methylated,

P is a proline or a proline analogue,

or an analogue, derivative, or functional variant thereof having the same activity thereof, or a pharmaceutical salt thereof, but wherein SEQ. ID. NO. 56 is not SEQ. ID. NO. 2 or 17.

2. The isolated peptide of claim 1, wherein each of X_2 and X_3 are I or V.
3. The isolated peptide of claim 1 or 2 wherein X_2 is I.
4. The isolated peptide of claim 1 or 3 wherein X_3 is V.
5. The isolated peptide of any one of claims 1 to 4 consisting of up to 7 amino acids.
6. The isolated peptide of any one of claim 1 to 5 comprising the formula:
$$aX_1X_2X_3P$$
 (SEQ. ID. NO. 57)
wherein X_1 , X_2 , and X_3 are as defined in any one of claims 1 to 5 and "a" is S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, M, or F.
7. The isolated peptide of any one of claims 1 to 5 wherein SEQ. ID. NO. 56 is the first four amino acids at the N-terminus of the peptide.

8. An isolated peptide of any one of claims 1 to 7, comprising the formula:



wherein X_1 , X_2 , and X_3 are as defined in any one of claims 1 to 6 and "b" is A, A* G, S, L, F, K, C, I, V, T, Y, R, H, O, or M, wherein A* denotes a D amino acid of Alanine.

9. An isolated peptide of claim 8 wherein "b" is A.

10. An isolated peptide of any one of claims 1 to 6, having the formula:



wherein X_1 , X_2 , and X_3 are as defined in any one of claims 1 to 6 and " a_1 " is K, I, R, H, O, L, V, A, or G, and a_2 is S, P, R, T, H, K, O, L, V, A, G, S, or I.

11. An isolated peptide of any one of claims 1 to 6, having the formula:



wherein X_1 , X_2 , and X_3 are as defined in any one of claims 1 to 6 and "a" is S, R, K, H, O, T, I, L, V, A or G, and "b" is A, V, I, L, G, K, H, R, O, S, T or F.

12. An isolated peptide of claim 10, wherein "b" is A.

13. An isolated peptide of claim 1 comprising any one of SEQ. ID. NOs. 1, 3, 4, 5, 7, 10, 12, 14, 18, 22, 23, 24, 31, 32-35.

14. An isolated peptide of claim 1 consisting of any one of SEQ. ID. NOs. 1, 3, 4, 5, 7, 10, 12, 14, 18, 22, 23, 24, 31, 32-35 or an analogue, derivative, or functional variant thereof having the same activity thereof, or a pharmaceutical salt thereof.

15. An isolated peptide of any one of claims 1 to 12 comprising SEQ. ID. NO. 5.

16. An isolated peptide consisting of SEQ. ID. NO. 5, or an analogue, derivative, or functional variant thereof having the same activity thereof, or a pharmaceutical salt thereof.
17. An isolated peptide of any one of claims 1 to 16 that has DPPIV activity of about 75% or less as compared to a saline control.
18. An isolated peptide of any one of claims 1 to 16 having immunological activity.
19. The isolated peptide of any one of claims 1 to 18, having a modified C-terminus and/or a modified N-terminus.
20. The isolated peptide of claim 19, having a modified C-terminus.
21. The isolated peptide of claim 20, having an amidated C-terminus.
22. The isolated peptide of any one of claims 1 to 21, modified by at least one substitution of a D amino acid.
23. A pharmaceutical composition comprising the peptide of any one of claims 1 to 22 and a pharmaceutically-acceptable carrier, diluent, or excipient.
24. A use of a peptide comprising an amino acid sequence listed in any one of claims 1 to 22 or a pharmaceutical composition of claim 23 for treating and/or preventing infection in a subject.
25. The use of claim 24, wherein the peptide modulates innate immunity in the subject, thereby treating and/or preventing the infection in the subject.
26. The use of claim 24, wherein the infection is a microbial infection.

27. The use of claim 26, wherein the infection is selected from the group consisting of an infection by a bacterium, an infection by a fungus, an infection by a parasite, and an infection by a virus.

28. The use of claim 27, wherein the bacterium is a Gram-positive or Gram-negative bacterium.

29. The use of claim 27, wherein the bacterium is selected from the group consisting of *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella spp.*, *Staphylococcus aureus*, *Streptococcus spp.*, and vancomycin-resistant enterococcus.

30. The use of claim 27, wherein the fungus is selected from the group consisting of a mould, a yeast, and a higher fungus.

31. The use of claim 27 wherein the parasite is single-celled or multicellular.

32. The use of claim 31 wherein the parasite is selected from the group consisting of *Giardia duodenalis*, *Cryptosporidium parvum*, *Cyclospora cayetanensis*, and *Toxoplasma gondii*.

33. The use of claim 27, wherein the virus is associated with a condition selected from the group consisting of AIDS, avian flu, chickenpox, cold sores, common cold, gastroenteritis, glandular fever, influenza, lower respiratory tract infection, measles, mumps, pharyngitis, pneumonia, rubella, SARS, and upper respiratory tract infection.

34. The use of claim 33, wherein the virus is respiratory syncytial virus (RSV).

35. The use of claim 24, wherein the subject has, or is at risk of having, infection.

36. The use of claim 24, wherein the peptide is administered orally, parenterally, transdermally, intranasally, by pulmonary administration, or by osmotic pump.

37. A use of a peptide of any one of claims 1 to 22 by assaying a diagnostic sample of a subject for dipeptidyl peptidase-4 (DPPIV) activity and administering to the sample the peptide of the invention in the presence of a DPPIV substrate under conditions that would permit DPPIV to react with the substrate, wherein the reduction of DPPIV activity as compared to a control without the peptide is indicative that the subject would be responsive to the treatment.

38. An isolated peptide consisting essentially of an amino acid sequence of TABLE 1, or an analogue, derivative, or variant thereof or pharmaceutical acceptable salt thereof.

39. An isolated peptide of claim 38 but not comprising an amino acid sequence of SEQ. ID. NOs. 2, 17, or 36.

40. An isolated peptide of 4 to 10 amino acids comprising the amino acid sequence of TABLE 1, or an analogue, derivative, or variant thereof or pharmaceutical acceptable salt thereof, but not comprising an amino acid sequence of SEQ. ID. NOs. 2, 17, 36, 44 and 54.

41. An isolated peptide consisting essentially of a 4 to 10 amino acid peptide, said peptide comprising the formula:



wherein

X₁ is K, H, R, S, O, glycine based compound with a basic functional group substituted on the N-terminal, hSer, or Val (beta OH), Cit, hci, Dab, or Dpr;

X₂ is A, I, L, V, K, P, G, H, R, S, O, Dab, Dpr, Cit, Hci, Abu, Nva, Nle, where it can be N-methylated;

X₃ is I, V, P, and can be N-methylated,

P is a proline or a proline analogue,

or an analog, derivative or functional variant thereof, or pharmaceutical salt thereof.

but wherein SEQ. ID. NO. 56 is not SEQ. ID. NO. 2, 17 or 36.

42. The isolated peptide of claim 41, wherein X₁ is K, R, S, O or a glycine based compound with a basic functional group substituted on the N-terminal.

43. The isolated peptide of claim 42, wherein X₁ is R.


44. The isolated peptide of any one of claims 41 to 43 wherein X₂ is I.

45. The isolated peptide of any one of claims 41 to 44 wherein X₃ is V.

46. The isolated peptide of 4 amino acids of any one of claims 41 to 45.


47. The isolated peptide of 5 amino acids of any one of claims 41 to 45.

48. The isolated peptide of any one of claims 41 to 45 or 47 comprising SEQ. ID. NO. 57:



wherein X₁, X₂, X₃, and P are defined as in any one of claims 41 to 45 and "a" is selected from the group consisting of S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, M and F;

49. The isolated peptide of any one of claims 41 to 45 or 47 comprising SEQ. ID. NO. 58:




wherein X_1 , X_2 , X_3 , and P are defined as in any one of claims 41 to 45 and "b" is selected from the group consisting of A, A*, G, S, L, F, , C, I, V, T, Y, R, H, O and M.

50. The isolated peptide of claim 49 wherein "b" is A.


51. An isolated peptide of 6 amino acids comprising any one of claims 41 to 45.

52. An isolated peptide of any one of claims 41 to 45 or 51 comprising SEQ. ID. NO. 59:



wherein X_1 , X_2 , X_3 , and P are defined as in any one of claims 41 to 45, " a_1 " is K, I, R, H, O, L, V, A, or G, and " a_2 " is S, P, R, T, H, K, O, L, V, A, G, S, or I.

53. An isolated peptide of any one of claims 41 to 45 or 51 comprising SEQ. ID. NO. 60:



X_1 , X_2 , X_3 , and P are defined as in any one of claims 41 to 45, "a" is S, R, K, H, O, T, I, L, V, A, or G and "b" is A, V, I, L, G, K, H, R, O, S, T, and F.

54. The isolated peptide of any one of claims 41 to 53 wherein the Proline analogue is selected from the group consisting of Pip, Thz, Fpro, or Dhp.

55. The isolated peptide of any one of claims 38 to 54, having a modified C-terminus and/or a modified N-terminus.

56. The isolated peptide of claim 55, having a modified C-terminus.

57. The isolated peptide of claim 56, having an amidated C-terminus.

58. The isolated peptide of any one of claims 38 to 51 modified by at least one substitution of a D amino acid.

59. The isolated peptide of any one of claims 41 to 45 and 47 to 53, wherein the isolated peptide consists essentially of 5 to 10 amino acids and SEQ. ID. NO. 56, 57, 58, 59 or 60 is at the N-terminal of said peptide.

60. The isolated peptide of claim 41 comprising an amino acid sequence selected from the group consisting of SEQ. ID. NOs. 1, 3-7, 8, 9, 10-16, 18, 19, 20, 21-25, 26, 27, 28, 31-35, 37-39, 42, 43, 46, 47, and 50.

61. The isolated peptide of claim 41 comprising an amino acid sequence selected from the group consisting of SEQ. ID. NOs. 1, 3-7, 10-16, 18, 19, 20, 21-25, 27, 28, 31-35, 37-39, 42, 43, 46, or 47.

62. The isolated peptide of claim 41 comprising an amino acid sequence selected from the group consisting of SEQ. ID. NOs. 1, 3-7, 10-16, 18, 21-25, 27, 28, 31-35, 37-39, 42, 43, 46, or 47.

63. An isolated peptide of any one of claims 41 to 62 having DPPIV activity of about 75% or less as compared to a saline control.

64. An isolated peptide of any one of claims 41 to 62 having innate immunity modulating activity.

65. A pharmaceutical composition comprising the peptide of any one of claims 38 to 64 and a pharmaceutically-acceptable carrier, diluent, or excipient.

66. A pharmaceutical composition of claim 65 further comprising an antibiotic.

67. A use of a peptide of any one of claims 41 to 64 or a pharmaceutical composition of claim 65 for treating and/or preventing infection in a subject.

68. A use of a peptide of any one of claims 41 to 64 or a pharmaceutical composition of claim 65 in the treatment of a DPPIV-related condition.

69. A use a peptide of any one of claims 41 to 64 or a pharmaceutical composition of claim 65 for the prevention and/or treatment of an innate immunity-related condition.

70. The use of claim 67, wherein the peptide or pharmaceutical composition modulates innate immunity in the subject, thereby treating and/or preventing the infection in the subject.

71. The use of claim 67, wherein the infection is a microbial infection.

72. The use of claim 71, wherein the infection is selected from the group consisting of an infection by a bacterium, an infection by a fungus, an infection by a parasite, and an infection by a virus.

73. The use of claim 72, wherein the bacterium is a Gram-positive or Gram-negative bacterium.

74. The use of claim 73, wherein the bacterium is selected from the group consisting of *E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella* spp., *Staphylococcus aureus*, *Streptococcus* spp., and vancomycin-resistant enterococcus.

75. The use of claim 72, wherein the fungus is selected from the group consisting of a mould, a yeast, and a higher fungus.

76. The use of claim 72, wherein the parasite is single-celled or multicellular.

77. The use of claim 76, wherein the parasite is selected from the group consisting of *Giardia duodenalis*, *Cryptosporidium parvum*, *Cyclospora cayetanensis*, and *Toxoplasma gondii*.

78. The use of claim 72, wherein the virus is associated with a condition selected from the group consisting of AIDS, avian flu, chickenpox, cold sores, common cold, gastroenteritis, glandular fever, influenza, lower respiratory tract infection, measles, mumps, pharyngitis, pneumonia, rubella, SARS, and upper respiratory tract infection.

79. The use of claim 78, wherein the virus is respiratory syncytial virus (RSV).

80. The use of claim 67, wherein the subject has, or is at risk of having, infection.

81. The use of claim 80, wherein the peptide is administered orally, parenterally, transdermally, intranasally, by pulmonary administration, or by osmotic pump.

82. The use of an antibiotic in the use of any one of claims 67 to 81.

83. The use of claim 82, wherein the antibiotic is used with the peptide.

84. The use of claim 83 wherein the antibiotic and peptide are used in the form of a pharmaceutical composition of claim 65.

85. The use of claim 82, wherein the antibiotic is used after use of the peptide.

86. The use of any one of claims 67 to 85 wherein the peptide is used in the form of a pro-peptide or substance that metabolizes once administered to a subject into the peptide.

87. The use of claim 86 wherein the pro-peptide is a fusion protein comprising the peptide and a delivery vehicle.

88. A method for predicting whether a subject would be responsive to treatment for a microbial infection or an immunity-related disorder, wherein the treatment comprises administering to the subject a peptide comprising an amino acid sequence of claim 39 or an analogue, derivative, or variant thereof, the method comprising assaying a diagnostic sample of the subject for DPPIV activity by administering to the sample the peptide of the invention in the presence of a DPPIV substrate under conditions that would permit DPPIV to react with the substrate, wherein the reduction of DPPIV activity as compared to a control without the peptide is indicative that the subject would be responsive to the treatment.

89. An agent reactive with the peptide of any one of claims 38 to 64.

90. The agent of claim 89, which is an antibody.

91. The agent of claim 90, which is a monoclonal antibody.

92. An isolated nucleic acid molecule encoding the peptide of any one of claims 38 to 64.

93. A recombinant nucleic acid construct comprising the nucleic acid molecule of claim 92 operably linked to an expression vector.

94. At least one host cell comprising the recombinant nucleic acid construct of claim 93.

95. A method for producing a peptide comprising an amino acid sequence of any one of claims 38 to 64 or an analogue, derivative, or variant thereof, said method comprising the steps of:

(a) culturing the at least one host cell of claim 88, under conditions allowing expression of the peptide; and

(b) recovering the peptide from the at least one host cell or culture medium thereof.

SEQUENCE LISTING

<110> Inimex Pharmaceuticals, Inc.
 <120> NOVEL PEPTIDES FOR TREATING AND PREVENTING IMMUNE-RELATED
 DISORDERS, INCLUDING TREATING AND PREVENTING INFECTION BY
 MODULATING INNATE IMMUNITY
 <130> 190476-382091
 <140> N/A
 <141> 2006-10-04
 <150> US 60/722,962
 <151> 2005-10-04
 <150> US 60/722,958
 <151> 2005-10-04
 <150> US 60/722,959
 <151> 2005-10-04
 <160> 60
 <170> PatentIn version 3.3
 <210> 1
 <211> 6
 <212> PRT
 <213> Artificial
 <220>
 <223> IMMUNOLOGICAL PEPTIDE
 <400> 1
 Lys Ser Arg Ile Val Pro
 1 5
 <210> 2
 <211> 6
 <212> PRT
 <213> Artificial
 <220>
 <223> IMMUNOLOGICAL PEPTIDE
 <220>
 <221> MISC_FEATURE
 <223> X is equal to Acetylated K
 <220>
 <221> misc_feature
 <222> (1)..(1)

<223> xaa can be any naturally occurring amino acid

<400> 2

Xaa Ser Arg Ile Val Pro
1 5

<210> 3

<211> 6

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 3

Ser Arg Ile Val Pro Ala
1 5

<210> 4

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 4

Ser Arg Ile Val Pro
1 5

<210> 5

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 5

Arg Ile Val Pro Ala
1 5

<210> 6

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 6

Lys Ile Val Pro Ala
1 5

<210> 7

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<220>

<221> MISC_FEATURE

<223> Xaa is equal to D- Amino Acid of Ala

<220>

<221> misc_feature

<222> (5)..(5)

<223> Xaa can be any naturally occurring amino acid

<400> 7

Arg Ile Val Pro Xaa
1 5

<210> 8

<211> 4

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 8

Arg Val Pro Ala
1

<210> 9

<211> 4

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 9

Arg Ile Pro Ala

1

<210> 10

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<220>

<221> MISC_FEATURE

<223> Xaa is equal to Ala -OH

<220>

<221> misc_feature

<222> (5)..(5)

<223> Xaa can be any naturally occurring amino acid

<400> 10

Arg Ile Val Pro Xaa

1 5

<210> 11

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 11

Arg Ala Val Pro Ala

1 5

<210> 12

<211> 6

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 12

Arg Arg Ile Val Pro Ala

1 5

<210> 13
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 13

Arg Lys Val Pro Ala
1 5

<210> 14
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 14

Arg Ile Val Pro Lys
1 5

<210> 15
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 15

Arg Pro Val Pro Ala
1 5

<210> 16
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 16

Arg Ile Pro Pro Ala
1 5

<210> 17
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 17

Arg Ile Val Pro Pro
1 5

<210> 18
<211> 7
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 18

Arg Ile Val Pro Gly Gly Ala
1 5

<210> 19
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 19

Gly Gly Ile Val Pro Ala
1 5

<210> 20
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 20

Gly Ile Val Pro Ala
1 5

<210> 21
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 21

Arg Gly Val Pro Ala
1 5

<210> 22
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 22

Arg Ile Val Pro Gly
1 5

<210> 23
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 23

Arg Ile Val Pro Ser
1 5

<210> 24
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 24

Arg Ile Val Pro Leu
1 5

<210> 25
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 25

Arg His Val Pro Ala
1 5

<210> 26
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 26

Arg Ile Pro Val Ala
1 5

<210> 27
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 27

Arg Val Ile Pro Ala
1 5

<210> 28
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 28

Arg Ile Ile Pro Ala
1 5

<210> 29
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 29

Ala Val Pro Ile Arg
1 5

<210> 30
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 30

Ala Pro Val Ile Arg
1 5

<210> 31
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 31

Arg Ile Val Pro Ala
1 5

<210> 32
<211> 7
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 32

Cys Arg Ile Val Pro Ala Cys
1 5

<210> 33
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> MISC_FEATURE
<223> Xaa is equal to I with an NMe backbone

<220>
<221> misc_feature
<222> (2)..(2)
<223> Xaa can be any naturally occurring amino acid

<400> 33

Arg Xaa Val Pro Ala
1 5

<210> 34
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> MISC_FEATURE
<223> Xaa is equal to A with an N-methylated backbone

<220>
<221> misc_feature
<222> (5)..(5)
<223> Xaa can be any naturally occurring amino acid

<400> 34

Arg Ile Val Pro Xaa
1 5

<210> 35
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 35

Arg Ile Val Pro Phe
1 5

<210> 36

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<220>

<221> MISC_FEATURE

<223> Xaa is equal to Citrulline

<220>

<221> misc_feature

<222> (1)..(1)

<223> Xaa can be any naturally occurring amino acid

<400> 36

Xaa Ile Val Pro Ala
1 5

<210> 37

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 37

Arg Leu Val Pro Ala
1 5

<210> 38

<211> 5

<212> PRT

<213> Artificial

<220>

<223> IMMUNOLOGICAL PEPTIDE

<400> 38

His Ile Val Pro Ala
1 5

<210> 39
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 39

Ile Arg Arg Val Pro Ala
1 5

<210> 40
<211> 5
<212> PRT
<213> Artificial

<220>
<223>

<400> 40

Ala Arg Val Pro Ala
1 5

<210> 41
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 41

Ile Arg Val Pro Ala
1 5

<210> 42
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> MISC_FEATURE
<223> xaa is equal to Ornithine

<220>
<221> misc_feature
<222> (1)..(1)
<223> Xaa can be any naturally occurring amino acid

<400> 42

Xaa Ile Val Pro Ala
1 5

<210> 43
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 43

Ser Ile Val Pro Ala
1 5

<210> 44
<211> 13
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 44

Val Ser Ile Ile Lys Pro Ala Arg Val Pro Ser Leu Leu
1 5 10

<210> 45
<211> 7
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 45

Lys Pro Ala Arg Val Pro Ser
1 5

<210> 46
<211> 6

<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 46

Arg Val Pro Ser Leu Leu
1 5

<210> 47
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 47

Lys Pro Arg Ala Val Pro
1 5

<210> 48
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 48

Pro Ala Arg Val Pro
1 5

<210> 49
<211> 4
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 49

Ile Arg Val Pro
1

<210> 50
<211> 4

<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 50

Arg Val Pro Ser
1

<210> 51
<211> 3
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 51

Arg Val Pro
1

<210> 52
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 52

Pro Ser Val Pro Gly Ser
1 5

<210> 53
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 53

Gly Leu Lys His Pro Ser
1 5

<210> 54
<211> 11

<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<400> 54

Arg Ile Val Pro Ala Ile Pro Val Ser Leu Leu
1 5 10

<210> 55
<211> 3
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (1)..(2)
<223> X1 is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH).
X2 is selected from the group consisting of V, I, K, P, and H [including an isolated peptide of up to 10 amino acids comprising an amino acid sequence of SEQ. ID. NO. 55.]

<400> 55

X1 X2 Pro
1

<210> 56
<211> 4
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (1)..(3)
<223> Wherein X1 is selected from the group consisting of K, H, R, S, T, O, Cit, Hci, Dab, Dpr, or glycine based compounds with basic functional groups substituted on the N-terminal (e.g., Nlys), hSer, Val(betaOH) and wherein X2 is selected from the group consisting of A, I, L, V, K, P, G, H, R, S, O, Dab, Dpr, Cit, Hci, Abu, Nva, Nle and where X2 can be N-methylated, and wherein X3 is selected from the group consisting of I, V, P, wherein in one embodiment,

X3 is not N-methylated. In one embodiment, the peptide can be an amino acid sequence of up to 10 amino acids comprising SEQ. ID. NO. 56, but not SEQ. ID. NOS. 2 or 17.

<400> 56

X1 X2 X3 Pro
1

<210> 57
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (2)..(4)
<223> wherein X1, X2 and X3 are defined as in Seq. ID. NO. 56, and wherein "a" is selected from the group consisting of S, P, I, R, C, T, L, V, A, G, K, H, R, O, C, M, and F or an isolated peptide up to 10 amino acids comprising SEQ. ID. NO. 57.

<400> 57

a X1 X2 X3 Pro
1 5

<210> 58
<211> 5
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (1)..(3)
<223> wherein X1X2X3P are as defined in SEQ. ID. NO. 56 and "b" is selected from the group consisting of A, A*,G, S, L, F, K, C, I, V, T, Y, R, H, O, and M, but in one embodiment not P. In one embodiment, the isolated peptide is a peptide of up to 10 amino acids comprising SEQ. ID. NO. 58. In one embodiment, the peptide is not SEQ. ID. NO. 17.

<400> 58

X1 X2 X3 Pro b

1

5

<210> 59
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (3)..(5)
<223> Where X1, X2 and X3 are as defined in SEQ. ID.NO. 56 and a1 is selected from the group consisting of K, I, R, H, O, L, V, A, and G and a2 is selected from the group consisting of S, P, R, T, H, K, O, L, V, A, G, S, I. In one embodiment, a1 is not acetylated, or where a1 is K, K is not acetylated or not SEQ. ID. NO. 2. In one embodiment, the isolated peptide comprises up to 10 amino acids comprising SEQ. ID. NO. 59.

<400> 59

a1	a2	X1	X2	X3	Pro
1					5

<210> 60
<211> 6
<212> PRT
<213> Artificial

<220>
<223> IMMUNOLOGICAL PEPTIDE

<220>
<221> misc_feature
<222> (2)..(4)
<223> Where X1, X2 and X3 are as defined in SEQ. ID.NO. 56 and where "a" is selected from the group consisting of S, R, K, H, O, T, I, L, V, A, and G and wherein "b" is selected from the group consisting of A, V, I, L, G, K, H, R, O, S, T, and F or a peptide of up to 10 amino acids comprising SEQ. ID. NO. 60.

<400> 60

a	X1	X2	X3	Pro	b
1					5