

1-(ARYLMETHYL)QUINAZOLINE-2,4(1H,3H)-DIONES AS PARP INHIBITORS AND THE USE THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention is in the field of medicinal chemistry. In particular, the invention relates to 1-(arylmethyl)quinazoline-2,4(1H,3H)-diones, and the use of these compounds as PARP inhibitors and anti-cancer drugs.

Related Art


[0002] Poly (ADP-ribose) polymerase (PARP) catalyzes the addition of poly (ADP-ribose) to the target protein using NAD⁺ that is an important process in DNA repair. This is an essential process for maintaining DNA and chromosome integrity and stability, and for ensuring the survival of mammalian cells. PARP-1 catalyzes the majority of the intracellular ADP-ribose polymerization reactions, although PARP-2 and other subtypes also have this function. The PARP-1 knockout mice do not have the repair function for single-stranded DNA damages (Krishnakumar and Kraus, 2010, Molecular Cell 39:8). Cancer cells with DNA repair defects, such as BRCA1 (breast cancer 1) or BRCA2 (breast cancer 2) deficiency, are particularly sensitive to DNA damaging anticancer agents, including platinum chemotherapy drugs, DNA methylation anticancer drugs and DNA topoisomerase inhibitors, or radiation therapy. Phase II clinical trial data have shown that PARP-1 inhibitor olaparib (AZD2281) was effective for the treatment of advanced breast cancer (Andrew Tutt et al., 2009, J. Clin. Oncol 27:18 s ; Andrew Tutt et al., 2010 Lancet 376:235; RA Dent et al., 2010 J. Clin. Oncol. 28:15 s). These scientific and clinical results demonstrated that PARP-1 inhibitors may be used as effective anti-cancer drugs to treat a variety of cancers. The applications of PARP-1 inhibitors for the treatment of cancer are mainly based on two mechanisms. First, because of the rapid growth, DNA replication is much higher in cancer cells than in normal cells. Drugs that cause DNA damage will induce cancer cell death selectively. However, due to the presence of DNA repair enzymes such as PARP-1, the therapeutic effects of these drugs can not be fully materialized. By inhibiting the DNA repair mechanism, PARP-1 inhibitors in combination with commonly used DNA damaging anti-cancer drugs, such as temozolomide, can achieve synergy effects and greatly enhance the anticancer effects of currently used anticancer drugs. Second, for cancer cells with DNA repair deficiency, such as BRCA1 or BRCA2 deficient triple-negative breast

cancer, PARP-1 inhibitors can directly kill the cancer cells and function as anticancer drugs independently. According to statistics, about 10-15% of breast cancer patients have family history of genetic factors, in which the BRCA1 or BRCA2 gene mutations account for 15-20% of all hereditary breast cancers. Since PARP-1 is involved in DNA repair, utilization of PARP-1 inhibitors to inhibit DNA repair may be an effective and selective treatment for cancers with DNA repair genetic defect, including triple-negative breast cancers. Furthermore, PARP-1 inhibitors may also be used to treat diseases due to excessive cell death, including central nervous system diseases such as stroke and neurodegenerative diseases (Akinori Iwashita et al., 2004, *J. Pharmacol. Exp. Thera.* 310: 425).


[0003] The inhibitory activity of PARP-1 inhibitors can be measured by directly using PARP-1 enzymes. In addition, since PARP-1 inhibitors can increase the cytotoxicity of DNA damaging anti-cancer drugs such as methyl methanesulfonate (MMS) on cancer cells, the activity of PARP-1 inhibitors can also be determined by measuring cell viability, such as using a MTT assay, in the presence of MMS and PARP-1 inhibitors. Furthermore, Cancer cells with deficiency in DNA repair, such as in the case of BRCA1 or BRCA2 deficient triple-negative breast cancer, can be killed by PARP-1 inhibitors alone. Therefore the anticancer activity of PARP-1 inhibitors can be determined by measuring the inhibitory effect of these compounds on cell growth of BRCA-2 deficient CAPAN-1 human pancreatic cancer cells.

[0004] It has been known that many cancer chemotherapeutic drugs trigger cancer cells to undergo apoptosis. The mechanism of apoptosis involves a cascade of initiator and effector caspases that are activated sequentially. Caspases are a family of cysteine proteases that require aspartic acid residues at the P₁ position of substrates for efficient cleavage. Among these caspases, caspase-3, 6, and 7 are key effector caspases that cleave multiple protein substrates in cells, leading to cell death. Cellular caspase activity can be determined using caspase substrates and used as a measurement of cell apoptosis. PARP-1 inhibitors can increase the apoptosis-inducing activity of many DNA damaging anticancer drugs such as MMS. Therefore, the activity of PARP-1 inhibitors can be determined via measuring the intracellular caspase activity of cancer cells treated with DNA damaging anticancer drugs in combination with PARP-1 inhibitors.

[0005] JP2007137818 disclosed the preparation of 8-hydroxyquinazoline-2,4(1H,3H)-dione derivatives as poly(ADP-ribose) polymerase (PARP) inhibitors, wherein X = (CH₂)_n; n = an integer of 1-4; Y = H, NR₁R₂, 1,2,3,4-tetrahydroisoquinolyl, decahydroisoquinolyl, 1,3-dioxo-1,3-dihydro-2H-isoindolyl, 3-oxo-3,4-dihydrobenz[1,4]oxazinyl, pyridyl, benzyl, (un)substituted optionally fused 5-membered N-heterocyclyl, and (un)substituted aryloxy, etc.



[0006] WO2006003148 disclosed the preparation of quinazolinone derivatives as PARP inhibitors for the treatment of PARP mediated diseases, wherein X and Y are each independently N or CH; L_1 = a bond or alkylene; L_2 = a bond, CO, alkylene, CO-alkylene, etc.; R_1 = H or OH; Z = H or (un)substituted (hetero)aryl; etc.



SUMMARY OF THE INVENTION

[0007] The invention provides novel 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones, as represented in Formulae I, II and III. These compounds have PARP inhibitory activities.

[0008] The present invention also provides pharmaceutical compositions comprising a compound of Formula I, II or III in an effective amount for the treatment of cancer.

[0009] The invention also provides a pharmaceutical composition useful for the treatment of cancer, containing an effective amount of a compound of one of the Formula I, II or III in admixture with one or more pharmaceutically acceptable carriers or diluents.


[0010] The invention also provides a pharmaceutical composition useful for the treatment of cancer, containing an effective amount of a compound of one of the Formula I, II or III, in combination with one known anticancer drugs or its pharmaceutically acceptable salts.

[0011] The invention also is directed to methods for the preparation of novel compounds of Formulae I, II and III.

DETAILED DESCRIPTION OF THE INVENTION

[0012] The novel and potent PARP inhibitors of the present invention include 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones, as represented in Formulae I, II and III.

[0013] Specifically, compounds of the present invention are represented by Formula I:




or pharmaceutically acceptable salts or prodrugs thereof, wherein:

Ar is an optionally substituted aryl or an optionally substituted heteroaryl;

R₁-R₆ independently are hydrogen, halo, optionally substituted amino, optionally substituted alkoxy, optionally substituted C₁₋₁₀ alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, cyano, acylamido, hydroxy, thiol, acyloxy, azido, carboxy, ethylenedioxy, carbonylamido or optionally substituted alkylthiol.

[0014] Preferred compounds of Formula I include compounds wherein Ar is an optionally substituted phenyl, pyridyl or furanyl. More preferably, Ar is phenyl, pyridyl or furanyl, optionally substituted with a substituted carbonyl or methyl, preferably carbonyl, at the meta-position. Another group of preferred compounds of Formula I include compounds wherein R₅ and R₆ are hydrogen.

[0015] One group of preferred compounds of the present invention are represented by Formula II:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

R₁-R₄ independently are hydrogen, halo, optionally substituted amino, optionally substituted alkoxy, optionally substituted C₁₋₁₀ alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, cyano, acylamido, hydroxy, thiol, acyloxy, azido, carboxy, ethylenedioxy, carbonylamido or optionally substituted alkylthiol;


R₇-R₁₀ independently are hydrogen, halo, optionally substituted amino, alkoxy, C₁₋₁₀ alkyl, haloalkyl, aryl, heteroaryl, a carbocyclic group, a heterocyclic group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, hydroxyalkoxy, aminoalkyl, aminoalkoxy, carboxyalkyl, carboxyalkoxy, nitro, cyano, acylamido, aminocarbonyl, hydroxy, thiol, acyloxy,

azido, carboxy, carbonylamido, alkylsulfonyl, aminosulfonyl, dialkylaminosulfonyl, alkylsulfinyl, alkylthiol, or substituted carbonyl;

R₁₁ is an optionally substituted amino, hydrazine, alkoxy, C₁₋₁₀ alkyl, haloalkyl, aryl, heteroaryl, carbocyclic group, heterocyclic group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, hydroxyalkoxy, aminoalkyl, aminoalkoxy, carboxyalkyl, carboxyalkoxy, acylamido, hydroxy, thiol, acyloxy, carbonylamido, or alkylthiol.

[0016] One group of preferred compounds of Formula II includes compounds wherein R₇, R₈, R₉ or R₁₀ is hydrogen or halo, preferably fluoro. Another group of preferred compounds of Formula II includes compounds wherein R₁ or R₂ is hydrogen, fluoro, chloro, bromo or methyl. Another group of preferred compounds of Formula II includes compounds wherein R₄ is hydrogen, fluoro, methyl, methoxy or hydroxy. Another group of preferred compounds of Formula II includes compounds wherein R₁₁ is an optionally substituted amino, more preferably substituted piperazine or piperidine.

[0017] One group of preferred compounds of the present invention are represented by Formula III:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

R₁-R₄ independently are hydrogen, halo, optionally substituted amino, optionally substituted alkoxy, optionally substituted C₁₋₁₀ alkyl, haloalkyl, alkenyl, alkynyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, cyano, acylamido, hydroxy, thiol, acyloxy, azido, carboxy, ethylenedioxy, carbonylamido or optionally substituted alkylthiol;

R₇-R₁₀ independently are hydrogen, halo, optionally substituted amino, alkoxy, C₁₋₁₀ alkyl, haloalkyl, aryl, heteroaryl, a carbocyclic group, a heterocyclic group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, hydroxyalkoxy, aminoalkyl, aminoalkoxy, carboxyalkyl, carboxyalkoxy, nitro, cyano, acylamido, aminocarbonyl, hydroxy, thiol, acyloxy, azido, carboxy, carbonylamido, alkylsulfonyl, aminosulfonyl, dialkylaminosulfonyl, alkylsulfinyl, alkylthiol, or substituted carbonyl;

R₁₂ is an optionally substituted C₁₋₁₀ alkyl, haloalkyl, cycloalkyl, aryl, heteroaryl, carbocyclic group, heterocyclic group, alkenyl, alkynyl, acyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, heterocyclocarbonyl, aminocarbonyl, alkylsulfonyl, cycloalkylsulfonyl or aminosulfonyl.

[0018] In one preferred embodiment, one group of preferred compounds of Formula III includes compounds wherein R₁₂ is selected from the group consisting of cycloalkyl, aryl, heteroaryl, carbocyclic group, heterocyclic group, arylalkyl, heteroarylalkyl, carbocycloalkyl, heterocycloalkyl, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, heteroarylcarbonyl or heterocyclocarbonyl. Another group of preferred compounds of Formula III includes compounds wherein R₁ or R₂ is hydrogen, fluoro, chloro, bromo or methyl; R₄ is hydrogen, fluoro, methoxy or hydroxy; R₇, R₈, R₉ or R₁₀ is hydrogen or fluoro.

[0019] Exemplary preferred compounds of Formulae I, II and III include, without limitation:

- 1-(3-Methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-Carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-Cyclohexylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-([1,2,4]Triazolo[4,3-*b*]pyridazin-6-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-Ethylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-Benzoylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(4-Fluorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(4-Chlorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(4-Bromobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(4-Methoxybenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Tetrahydro-2*H*-pyran-4-yl)carbonylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Ethylsulfonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-Acetylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Phenylpiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-Phenylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Pyrazin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(4-Fluoro-3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-Carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(4-Fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Piperidin-1-yl)piperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Pyridin-4-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(6-Fluorobenzo[*d*]isoxazol-3-yl)piperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Furan-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Thiophene-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Pyridine-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Pyridine-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Pyridine-4-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-Phenoxypiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(pyrazin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Chloro-3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Chloro-3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Chloro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(2-Chloro-3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(2-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(2-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-2-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(5-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-5-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Chloro-3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-((2-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-((2-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-((2-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-((2-(4-(Pyridin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-((2-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylmethyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylmethyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylsulfonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

7-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-methylquinazoline-2,4(1*H*,3*H*)-dione;

7-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;

6-Bromo-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Bromo-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Bromo-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione;

6-Nitro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-Benzoylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(Pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(4-Methoxybenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(4-Fluorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(4-Bromobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-8-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-methylquinazoline-2,4(1*H*,3*H*)-dione;

7-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

7-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-(1-(3-(4-benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)quinazoline)-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Chloro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Cyclopentylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(6-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(6-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylcarbamoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride;

5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride;

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride;

1-(3-((4-(Pyridin-2-yl)piperazin-1-yl)methyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(Naphthalen-2-yl)acetamidobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(3,4-Dimethoxyphenyl)acetamidobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-([1,2,4]Triazolo[4,3-*a*]pyridine-6-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Fluoro-1-((2-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-methoxybenzylcarbamoyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(3-Chlorobenzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(Benzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(5-Bromopyrimidin-2-aminocarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-7-(trifluoromethyl)quinazoline-2,4(1*H*,3*H*)-dione;

6,7-Ethylenedio-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(6-methoxy-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

7-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-(tetrahydrofuran-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-nitro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-Cyclohexylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-phenylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5-Fluoro-1-(4-fluoro-3-(4-phenylpiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(4-Bromo-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

6,7-Methylenedio-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(3-(4-(Cyclohexylmethyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

8-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

6-Amino-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

1-(2-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Hydroxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

and pharmaceutically acceptable salts or prodrugs thereof.

- [0020] The term “alkyl” as employed herein by itself or as part of another group refers to both straight and branched chain radicals of up to ten carbons. Useful alkyl groups include straight-chained and branched C₁₋₁₀ alkyl groups, more preferably C₁₋₆ alkyl groups. Typical C₁₋₁₀ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, 3-pentyl, hexyl and octyl groups, which may be optionally substituted.
- [0021] The term “alkenyl” as employed herein by itself or as part of another group means a straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, including at least one double bond between two of the carbon atoms in the chain. Typical alkenyl groups include ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl and 2-butenyl.
- [0022] The term “alkynyl” is used herein to mean a straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, wherein there is at least one triple bond between two of the carbon atoms in the chain. Typical alkynyl groups include ethynyl, 1-propynyl, 1-methyl-2-propynyl, 2-propynyl, 1-butyne and 2-butyne.
- [0023] Useful alkoxy groups include oxygen substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted. Alkoxy substituents include, without limitation, halo, morpholino, amino including alkylamino and dialkylamino, and carboxy including esters thereof.
- [0024] Useful alkylthio groups include sulfur substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted. Also included are the sulfoxides and sulfones of such alkylthio groups.
- [0025] Useful amino and optionally substituted amino groups include -NH₂, -NHR₁₅ and -NR₁₅R₁₆, wherein R₁₅ and R₁₆ are optionally substituted C₁₋₁₀ alkyl, cycloalkyl, aryl, heteroaryl, or amino; or R₁₅ and R₁₆ are combined with the N to form a 5-8 membered heterocyclic ring structure, such as a piperidine; or R₁₅ and R₁₆ are combined with the N and an additional N or O atom to form a 5-8 membered heterocyclic ring, such as a piperazine, which are optionally substituted.
- [0026] The groups as described herein, such as alkyl, alkoxy, alkylthio, alkenyl, alkynyl, cycloalkyl, carbonyl, carbocyclic and heterocyclic groups, aryl, arylalkyl, arylalkenyl, arylalkynyl and heteroaryl and heteroarylalkyl groups, may be optionally substituted. Generally, the term “optionally substituted” used herein indicates that the group that is “optionally substituted” may be optionally substituted by one or more (such as 1, 2, 3, or 4) substituents

selected from the group consisting of halo, hydroxy, carboxyl, amino, nitro, cyano, C₁-C₆ acylamino, C₁-C₆ acyloxy, C₁-C₆ alkoxy, aryloxy, alkylthio, C₁-C₆ alkyl, C₆-C₁₀ aryl, C₃-C₈ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, saturated and unsaturated heterocyclic and heteroaryl, methylenedioxy, C₁-C₆ haloalkyl, C₆-C₁₀ aryl(C₁-C₆)alkyl, C₁-C₆ hydroxyalkyl, ureido, thiol, azido, C₁-C₆ alkoxy, carbonyl, di(C₁₋₁₀ alkyl)amino, alkylsulfonyl, aminosulfonyl, dialkylaminosulfonyl, and alkylsulfinyl, and the like. The substituent itself may also be optionally substituted.

[0027] Optional substituents on the alkyl, alkoxy, alkylthio, alkenyl, alkynyl, cycloalkyl, carbonyl, carbocyclic and heterocyclic groups may be one or more (such as 1, 2, 3, or 4) groups selected from the group consisting of halo, hydroxy, carboxyl, amino, nitro, cyano, C₁-C₆ acylamino, C₁-C₆ acyloxy, C₁-C₆ alkoxy, aryloxy, alkylthio, C₆-C₁₀ aryl, C₃-C₈ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, saturated and unsaturated heterocyclic and heteroaryl.

[0028] Optional substituents on the aryl, arylalkyl, arylalkenyl, arylalkynyl and heteroaryl and heteroarylalkyl groups may be one or more (such as 1, 2, 3, or 4) groups selected from the group consisting of halo, methylenedioxy, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, C₃-C₈ cycloalkyl, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₁-C₆)alkyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, C₁-C₆ hydroxyalkyl, nitro, amino, ureido, cyano, C₁-C₆ acylamino, hydroxy, thiol, C₁-C₆ acyloxy, azido, C₁-C₆ alkoxy, carbonyl, carboxy, di(C₁₋₁₀ alkyl)amino, alkylsulfonyl, aminosulfonyl, dialkylaminosulfonyl, and alkylsulfinyl.

[0029] The term "aryl" as employed herein by itself or as part of another group refers to monocyclic, bicyclic or tricyclic aromatic groups containing from 6 to 14 carbons in the ring portion.

[0030] Useful aryl groups include C₆₋₁₄ aryl, preferably C₆₋₁₀ aryl. Typical C₆₋₁₄ aryl groups include phenyl, naphthyl, phenanthrenyl, anthracenyl, indenyl, azulenyl, biphenyl, biphenylenyl and fluorenyl groups.

[0031] The term "carbocycle" as employed herein include cycloalkyl and partially saturated carbocyclic groups. Useful cycloalkyl groups are C₃₋₈ cycloalkyl. Typical cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

[0032] Useful saturated or partially saturated carbocyclic groups are cycloalkyl groups as described above, as well as cycloalkenyl groups, such as cyclopentenyl, cycloheptenyl and cyclooctenyl.

[0033] Useful halo or halogen groups include fluorine, chlorine, bromine and iodine.

- [0034] The term "arylalkyl" is used herein to mean any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Preferably the arylalkyl group is benzyl, phenethyl or naphthylmethyl.
- [0035] The term "arylalkenyl" is used herein to mean any of the above-mentioned C₂₋₁₀ alkenyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups.
- [0036] The term "arylalkynyl" is used herein to mean any of the above-mentioned C₂₋₁₀ alkynyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups.
- [0037] The term "aryloxy" is used herein to mean oxygen substituted by one of the above-mentioned C₆₋₁₄ aryl groups, which may be optionally substituted. Useful aryloxy groups include phenoxy and 4-methylphenoxy.
- [0038] The term "arylalkoxy" is used herein to mean any of the above mentioned C₁₋₁₀ alkoxy groups substituted by any of the above-mentioned aryl groups, which may be optionally substituted. Useful arylalkoxy groups include benzyloxy and phenethyloxy.
- [0039] Useful haloalkyl groups include alkyl groups, such as C₁₋₁₀ alkyl, or preferably C₁₋₆ alkyl substituted by one or more fluorine, chlorine, bromine or iodine atoms, e.g., fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl, chloromethyl, chlorofluoromethyl and trichloromethyl groups.
- [0040] Useful acylamino (acylamido) groups are any C₁₋₆ acyl (alkanoyl) attached to an amino nitrogen, e.g., acetamido, chloroacetamido, propionamido, butanoylamido, pentanoylamido and hexanoylamido, as well as aryl-substituted C₁₋₆ acylamino groups, e.g., benzoylamido, and pentafluorobenzoylamido. Useful acyl includes C₁₋₆ acyl, such as acetyl.
- [0041] Useful acyloxy groups are any C₁₋₆ acyl (alkanoyl) attached to an oxy (–O–) group, e.g., formyloxy, acetoxy, propionyloxy, butanoyloxy, pentanoyloxy and hexanoyloxy.
- [0042] The term heterocycle (heterocyclic group) is used herein to mean a saturated or partially saturated 3-7 membered monocyclic, or 7-10 membered bicyclic ring system, which consists of carbon atoms and one to four heteroatoms independently selected from the group consisting of O, N, and S, wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, the nitrogen can be optionally quaternized. The term also includes any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring of heterocycle can be substituted on carbon or on a nitrogen atom if the resulting compound is stable.
- [0043] Useful saturated or partially saturated heterocyclic groups include tetrahydrofuranyl, pyranal, piperidinyl, piperazinyl, pyrrolidinyl, imidazolidinyl, imidazoliny, indoliny, isoindoliny, quinuclidiny, morpholiny, isochromanyl, chromanyl, pyrazolidinyl pyrazoliny, tetronoyl and tetramoyl groups, which are optionally substituted.

- [0044] The term “heteroaryl” as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 π electrons shared in a cyclic array; and containing, as ring atom, carbon atoms and 1-3 heteroatoms selected from oxygen, nitrogen and sulfur.
- [0045] Useful heteroaryl groups include thienyl (thiophenyl), benzo[*d*]isothiazol-3-yl, benzo[*b*]thienyl, naphtho[2,3-*b*]thienyl, thianthrenyl, furyl (furan), pyran, isobenzofuran, chromenyl, xanthenyl, phenoxanthiyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl (pyridinyl), including without limitation 2-pyridyl, 3-pyridyl, and 4-pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizyl, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinolizyl, isoquinolyl, quinolyl, phthalzyl, naphthyridinyl, quinoxalyl, cinnolyl, pteridinyl, carbazolyl, β -carbolyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, phenoxazinyl, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-*a*]pyrimidin-4-one, tetrahydrocyclopenta[*c*]pyrazol-3-yl, pyrazolo[1,5-*a*]pyrimidinyl, benzoisoxazolyl such as 1,2-benzoisoxazol-3-yl, benzimidazolyl, 2-oxindolyl, thiadiazolyl, and 2-oxobenzimidazolyl. Where the heteroaryl group contains a nitrogen atom in a ring, such nitrogen atom may be in the form of an *N*-oxide, e.g., a pyridyl *N*-oxide, pyrazinyl *N*-oxide and pyrimidinyl *N*-oxide.
- [0046] The term “heteroaryloxy” is used herein to mean oxygen substituted by one of the above-mentioned heteroaryl groups, which may be optionally substituted. Useful heteroaryloxy groups include pyridyloxy, pyrazinyloxy, pyrrolyloxy, pyrazolyloxy, imidazolyloxy and thiophenyloxy.
- [0047] The term “heteroarylalkoxy” is used herein to mean any of the above-mentioned C₁₋₁₀ alkoxy groups substituted by any of the above-mentioned heteroaryl groups, which may be optionally substituted.
- [0048] Some of the compounds of the present invention may exist as stereoisomers including optical isomers. The invention includes all stereoisomers and both the racemic mixtures of such stereoisomers as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art.
- [0049] Examples of pharmaceutically acceptable addition salts include inorganic and organic acid addition salts, such as hydrochloride, hydrobromide, phosphate, sulphate, citrate, lactate, tartrate, maleate, fumarate, mandelate and oxalate; and inorganic and organic base addition salts with bases, such as sodium hydroxy, Tris(hydroxymethyl)aminomethane (TRIS, tromethane) and *N*-methyl-glucamine.
- [0050] Examples of prodrugs of the compounds of the invention include the simple esters of carboxylic acid containing compounds (e.g., those obtained by condensation with a C₁₋₄ alcohol according to methods known in the art); esters of hydroxy containing compounds (e.g., those

obtained by condensation with a C₁₋₄ carboxylic acid, C₃₋₆ dioic acid or anhydride thereof, such as succinic and fumaric anhydrides according to methods known in the art); imines of amino containing compounds (e.g., those obtained by condensation with a C₁₋₄ aldehyde or ketone according to methods known in the art); carbamate of amino containing compounds, such as those described by Leu, *et al.*, (*J. Med. Chem.* 42:3623-3628 (1999)) and Greenwald, *et al.*, (*J. Med. Chem.* 42:3657-3667 (1999)); and acetals and ketals of alcohol containing compounds (e.g., those obtained by condensation with chloromethyl methyl ether or chloromethyl ethyl ether according to methods known in the art).


[0051] The compounds of this invention may be prepared using methods known to those skilled in the art, or the novel methods of this invention. Specifically, the compounds of this invention with Formula I, II or III can be prepared as illustrated by the exemplary reaction in Scheme 1 (Shunsuke Goto et al. Organic Process Research & Development, 2003, 7, 700-706). Reaction of quinazoline-2,4(1*H*,3*H*)-dione with hexamethyldisilazane (HMDS) in toluene in the presence of sulfuric acid produced the intermediate 2,4-di(trimethylsiloxy)quinazoline. Reaction of 2,4-di(trimethylsiloxy)quinazoline with a substituted alkyl 3-(halomethyl)benzoate, such as methyl 3-(bromomethyl)benzoate in DMF, followed by treatment with 1,4-dioxane and methanol, produced 1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Treatment of the ester with NaOH in water-methanol produced 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Coupling of the acid with a substituted amine, such as 1-(pyridin-2-yl)piperazine, in the presence of coupling agents, such as 2-(7-aza-1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) and *N,N*-diisopropylethylamine (DIPEA) in DMF, produced the targeted compound 1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione.

[0052] Other related compounds can be prepared similarly. For example, replacement of methyl 3-(bromomethyl)benzoate with methyl 6-(bromomethyl)pyridine-2-carboxylate produced the targeted compound 1-((2-(4-(pyridin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione. Replacement of methyl 3-(bromomethyl)benzoate with ethyl 5-(chloromethyl)furan-2-carboxylate produced the targeted compound 1-((2-(4-cyclopentylcarbonylpiperazine-1-carbonyl)furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione. Replacement of methyl 3-(bromomethyl)benzoate with methyl 3-(2-bromoethyl)benzoate produced the targeted compound 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)phenethyl)quinazoline-2,4(1*H*,3*H*)-dione.

[0053] For quinazoline-2,4(1*H*,3*H*)-dione with a substituent in the 8-position, such as 8-chloro quinazoline-2,4(1*H*,3*H*)-dione, reaction with hexamethyldisilazane (HMDS) followed by treatment of the intermediate 8-chloro-2,4-di(trimethylsiloxy)quinazoline with methyl 3-

(bromomethyl)benzoate produced a mixture of 1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione and 3-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione. The mixture can be separated, hydrolyzed, and coupled with a substituted amine to produce the 1-substituted compound and 3-substituted compound, respectively. Alternatively, the 3-substituted compound also can be prepared from reaction of quinazoline-2,4(1*H*,3*H*)-dione with methyl 3-(bromomethyl)benzoate and K_2CO_3 in DMF, followed by hydrolysis and coupling of the acid with a substituted amine.

Scheme 1




[0054] Compounds of this invention can be prepared as illustrated by the exemplary reaction in Scheme 2. Reaction of the intermediate 2,4-di(trimethylsiloxy)quinazoline with a substituted alkyl 3-(halomethyl)benzoate, such as methyl 5-(bromomethyl)-2-fluorobenzoate in DMF, followed by treatment with 1,4-dioxane and methanol, produced 1-(4-fluoro-3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Treatment of the ester with NaOH in water-methanol produced 1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Coupling of the acid with a substituted amine, such as 1-cyclopentylcarbonylpiperazine, in the presence of coupling agents, such as HATU and DIPEA in DMF, produced the targeted compound 1-(3-(4-cyclopentylcarbonylpiperazine-1-carboxyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione.

[0055] Other related compounds can be prepared similarly. For example, replacement of quinazoline-2,4(1*H*,3*H*)-dione with a substituted quinazoline-2,4(1*H*,3*H*)-dione, such as 5-fluoroquinazoline-2,4(1*H*,3*H*)-dione, which can be prepared from reaction of 2-amino-6-fluorobenzoic acid with potassium cyanate, produced the targeted compound 1-(3-(4-cyclopentylcarbonylpiperazine-1-carboxyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione. Replacement of methyl 5-(bromomethyl)-2-fluorobenzoate with another substituted methyl 3-(bromomethyl)benzoate, such as methyl 3-(bromomethyl)-4-fluorobenzoate, produced

the targeted compound 1-(3-(4-cyclopentylcarbonylpiperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Replacement of 1-cyclopentylcarbonylpiperazine with another substituted amine, such as 1-(pyrimidin-2-yl)piperazine, produced the targeted compound 1-(4-fluoro-3-(4-(pyrimidin-2-yl)carbonylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione.


Scheme 2



The compounds of this invention can be prepared as illustrated by the exemplary reaction in Scheme 3. Reaction of 2-amino-3-methylbenzoic acid with triphosgene in THF produced 8-methyl-1*H*-benzo[*d*][1,3]oxazine-2,4-dione, which was treated with tert-butylamine and DMAP in DMF to produce 2-amino-*N*-tert-butyl-3-methylbenzamide. Reaction of the benzamide with CDI in THF produced 3-tert-butyl-8-methylquinazoline-2,4(1*H*,3*H*)-dione, which was reacted with methyl 3-(bromomethyl)benzoate and MeONa in DMF, followed by treatment with hydrochloric acid aqueous solution, to produce 8-methyl-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione. Coupling of the acid with 2-(piperazin-1-yl)pyrimidine in the presence of coupling agents, such as HATU and DIPEA in DMF, produced the targeted compound 8-methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione.


Other related compounds can be prepared similarly. For example, replacement of 2-amino-3-methylbenzoic acid with other substituted 2-aminobenzoic acid, such as 2-amino-3-methoxybenzoic acid, produced the targeted compound 8-methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione.

Scheme 3



The 1-phenyl substituted compound can be prepared as illustrated by the exemplary reaction in Scheme 4. Reaction of 2-bromobenzamide with oxalyl chloride in DCM, followed by addition of methyl 3-aminobenzoate produced methyl 3-(3-(2-bromobenzoyl)ureido)benzoate. Reaction of the benzoate with potassium *tert*-butoxide in DMF produced 1-(3-methoxycarbonylphenyl)quinazoline-2,4(1*H*,3*H*)-dione. Hydrolysis of the ester with NaOH in water-methanol produced 1-(3-carboxyphenyl)quinazoline-2,4(1*H*,3*H*)-dione. Coupling of the acid with 2-(piperazin-1-yl)pyrimidine in the presence of coupling agents, such as HATU and DIPEA in DMF, produced the targeted compound 1-(3-(1-(pyrimidin-2-yl)piperazine-4-carbonyl)phenyl)quinazoline-2,4(1*H*,3*H*)-dione.

Scheme 4



[0056] An important aspect of the present invention is the discovery that compounds having Formula I, II or III are PARP inhibitors. Therefore, these compounds are useful for the

treatment of a variety of clinical conditions responsive to the inhibition of PARP activity, such as cancer.

[0057] The present invention includes a therapeutic method comprising administering to a mammal an effective amount of a compound of Formula I, II or III, or a pharmaceutically acceptable salt or prodrug thereof, wherein said therapeutic method is useful for the treatment of diseases due to abnormal PARP activity, such as cancer. Such diseases that can be treated or prevented by the method or pharmaceutical composition of the present invention include, but are not limited to, liver cancer, melanoma, Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic leukemia, chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, choriocarcinoma, mycosis fungoides, head or neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer, and prostatic carcinoma.

[0058] Compounds of the present invention also are useful for the treatment or prevention of other clinical conditions due to abnormal PARP activity, such as excessive cell death, including central nervous system diseases such as stroke and neurodegenerative diseases.

[0059] In practicing the therapeutic methods, effective amounts of compositions containing therapeutically effective concentrations of the compounds of Formula I, II or III formulated for oral, intravenous, local or topical application, for the treatment of neoplastic diseases and other diseases, are administered to an individual exhibiting the symptoms of one or more of these disorders. The amounts are effective to ameliorate or eliminate one or more symptoms of the disorders. An effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce, the symptoms associated with the disease. Such amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Typically, repeated administration is required to achieve the desired amelioration of symptom.

[0060] In another embodiment, a pharmaceutical composition comprising a compound of Formula I, II or III or a pharmaceutically acceptable salt thereof, which functions as PARP inhibitor, in combination with a pharmaceutically acceptable vehicle, is provided.

[0061] Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound of Formula I, II or III, or a pharmaceutically acceptable salt or prodrug thereof, which functions as a PARP inhibitor, in combination with at least one known anticancer agent or a pharmaceutically acceptable salt thereof. Examples of known anticancer agents which may be used for combination therapy include, but not are limited to alkylating agents, such as busulfan, melphalan, chlorambucil, cyclophosphamide, ifosfamide, temozolomide, bendamustine, cis-platin, mitomycin C, bleomycin, and carboplatin; topoisomerase I inhibitors, such as camptothecin, irinotecan, and topotecan; topoisomerase II inhibitors, such as doxorubicin, epirubicin, aclarubicin, mitoxantrone, elliptinium and etoposide; RNA/DNA antimetabolites, such as 5-azacytidine, gemcitabine, 5-fluorouracil and methotrexate; DNA antimetabolites, such as 5-fluoro-2'-deoxy-uridine, fludarabine, nelarabine, ara-C, alanosine, pralatrexate, pemetrexed, hydroxyurea and thioguanine; antimitotic agents, such as colchicine, vinblastine, vincristine, vinorelbine, paclitaxel, ixabepilone, cabazitaxel, and docetaxel; antibodies, such as campath, Panitumumab, Ofatumumab, Avastin, Herceptin®, Rituxan®; kinase inhibitors such as imatinib, gefitinib, erlotinib, lapatinib, sorafenib, sunitinib, nilotinib, dasatinib, pazopanib, temsirolimus and everolimus; HDAC inhibitors such as vorinostat and romidepsin. Other known anticancer agents which may be used for combination therapy include tamoxifen, letrozole, fulvestrant, mitoguanzone, octreotide, retinoic acid, arsenic trioxide, zoledronic acid, bortezomib, thalidomide and lenalidomide.

[0062] In practicing the methods of the present invention, the compound of the invention may be administered together with at least one known anticancer agent as part of a unitary pharmaceutical composition. Alternatively, the compound of the invention may be administered apart from at least one known anticancer agent. In one embodiment, the compound of the invention and at least one known anticancer agent are administered substantially simultaneously, i.e. the compounds are administered at the same time or one after the other, so long as the compounds reach therapeutic levels in the blood at the same time. In another embodiment, the compound of the invention and at least one known anticancer agent are administered according to their individual dose schedule, so long as the compounds reach therapeutic levels in the blood.

[0063] Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a bioconjugate of a compound described herein, which functions as a PARP inhibitor, in bioconjugation with at least one known therapeutically useful antibody,

such as Herceptin[®] or Rituxan[®], growth factors, such as DGF, NGF; cytokines, such as IL-2, IL-4, or any molecule that binds to the cell surface. The antibodies and other molecules will deliver a compound described herein to its targets and make it an effective anticancer agent. The bioconjugates could also enhance the anticancer effect of the therapeutically useful antibodies, such as Herceptin[®] or Rituxan[®].

[0064] Similarly, another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound of Formula I, II or III, or its pharmaceutically acceptable salt or prodrug, which functions as a PARP inhibitor, in combination with radiation therapy. In this embodiment, the compound of the invention may be administered at the same time as the radiation therapy is administered or at a different time.

[0065] Yet another embodiment of the present invention is directed to a composition effective for post-surgical treatment of cancer, comprising a compound of Formula I, II or III, or its pharmaceutically acceptable salt or prodrug, which functions as a PARP inhibitor. The invention also relates to a method of treating cancer by surgically removing the cancer and then treating the mammal with one of the pharmaceutical compositions described herein.

[0066] Pharmaceutical compositions within the scope of this invention include all compositions wherein the compounds of the present invention are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the compounds may be administered to mammals, orally at a dose of 0.0025 to 50 mg/kg of body weight, per day, or an equivalent amount of the pharmaceutically acceptable salt thereof, to a mammal being treated. Preferably, approximately 0.01 to approximately 10 mg/kg of body weight is orally administered. If a known anticancer agent is also administered, it is administered in an amount that is effective to achieve its intended purpose. The amounts of such known anticancer agents effective for cancer are well known to those skilled in the art.

[0067] The unit oral dose may comprise from approximately 0.01 to approximately 50 mg, preferably approximately 0.1 to approximately 10 mg of the compound of the invention. The unit dose may be administered one or more times daily, as one or more tablets, each containing from approximately 0.1 to approximately 50 mg, conveniently approximately 0.25 to 10 mg of the compound or its solvates.

[0068] In a topical formulation, the compound may be present at a concentration of approximately 0.01 to 100 mg per gram of carrier.

[0069] In addition to administering the compound as a raw chemical, the compounds of the invention may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate

processing of the compounds into preparations that may be used pharmaceutically. Preferably, the preparations, particularly those preparations which may be administered orally and that may be used for the preferred type of administration, such as tablets, dragees, and capsules, as well as suitable solutions for administration by injection or orally, contain from approximately 0.01 to 99 percent, preferably from approximately 0.25 to 75 percent of active compound(s), together with the excipient.

[0070] Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the compounds of the present invention with a solution of a pharmaceutically acceptable non-toxic acid, such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like. Base addition salts are formed by mixing a solution of the compounds of the present invention with a solution of a pharmaceutically acceptable non-toxic base, such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, Tris, *N*-methylglucamine and the like.

[0071] The pharmaceutical compositions of the invention may be administered to any mammal, which may experience the beneficial effects of the compounds of the invention. Foremost among such mammals are humans and veterinary animals, although the invention is not intended to be so limited.

[0072] The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

[0073] The pharmaceutical preparations of the present invention are manufactured in a manner, which is itself known, e.g., by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use may be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

[0074] Suitable excipients are, in particular: fillers, such as saccharides, e.g. lactose or sucrose, mannitol or sorbitol; cellulose preparations and/or calcium phosphates, e.g. tricalcium phosphate or calcium hydrogen phosphate; as well as binders, such as starch paste, using, e.g.,

maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added, such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, e.g., silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, e.g., for identification or in order to characterize combinations of active compound doses.

[0075] Other pharmaceutical preparations, which may be used orally, include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules may contain the active compounds in the form of granules, which may be mixed with fillers, such as lactose; binders, such as starches; and/or lubricants, such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

[0076] Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, e.g., water-soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, e.g., sesame oil, or synthetic fatty acid esters, e.g., ethyl oleate or triglycerides or polyethylene glycol-400, or cremophor, or cyclodextrins. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, e.g., sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

[0077] In accordance with one aspect of the present invention, compounds of the invention are employed in topical and parenteral formulations and are used for the treatment of skin cancer.

[0078] The topical compositions of this invention are formulated preferably as oils, creams, lotions, ointments and the like by choice of appropriate carriers. Suitable carriers include vegetable or mineral oils, white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high molecular weight alcohol (greater than C₁₂). The preferred carriers are

those in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included, as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers may be employed in these topical formulations. Examples of such enhancers are found in U.S. Patent Nos. 3,989,816 and 4,444,762.

- [0079] Creams are preferably formulated from a mixture of mineral oil, self-emulsifying beeswax and water in which mixture of the active ingredient, dissolved in a small amount of an oil, such as almond oil, is admixed. A typical example of such a cream is one which includes approximately 40 parts water, approximately 20 parts beeswax, approximately 40 parts mineral oil and approximately 1 part almond oil.
- [0080] Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil, such as almond oil, with warm soft paraffin and allowing the mixture to cool. A typical example of such an ointment is one which includes approximately 30 % almond oil and approximately 70 % white soft paraffin by weight.
- [0081] The present invention also includes the use of the compounds of the subject invention in the manufacture of a medicament for treating or preventing a disorder responsive to the inhibition of PARP activity in a mammal suffering therefrom. The medicament may include, such as the pharmaceutical compositions as described above.
- [0082] The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

EXAMPLES

General remarks

All reagents were of commercial quality. Solvents were dried and purified by standard methods. Mass spectrum analyses were recorded on a Platform II (Agilent 6110) quadrupole mass spectrometer fitted with an electrospray rinterface. ¹H NMR spectra was recorded at 300 MHz and at 300 K, on a Brücker AMX 300 apparatus. Chemical shifts were recorded as parts per million (ppm) downfield from TMS (0.00 ppm), and *J* coupling constants were reported in hertz (Hz).

Example 1

1-(3-Methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione

Quinazoline-2,4(1*H*,3*H*)-dione (1.0 g, 6.2 mmol), hexamethyldisilazane (HMDS, 2.5 g, 15.4 mmol), and toluene (50 mL) were added into 100 mL two-mouth flask. Concentrated sulfuric acid (0.06 g) was added into the flask when stirring. The mixture was refluxed and reacted for 8h. Toluene and excessive HMDS were removed via evaporation under reduced pressure, thus obtaining an intermediate product 2,4-di(trimethylsilyloxy)quinazoline.

The intermediate product 2,4-di(trimethylsilyloxy)quinazoline, methyl 3-(bromomethyl)benzoate (2.1 g, 9.2 mmol) and DMF (1 mL) were added in turn into a 50 mL single-mouth flask, and then the temperature were raised to 115-130°C and reacted for 3 hours. 1,4-dioxane (6 mL) and methanol (10 mL) were added after the reaction mixture was cooled to 100°C. Then the mixture was refluxed for 30 minutes. The mixture was filtered after it was cooled to room temperature. The precipitate was washed with water (20 mL) and methanol (10mL), respectively, and dried to give the title compound (1.6 g, 83.7% yield) as white powder. ¹H NMR (DMSO-*d*₆): 11.76 (s, 1H), 8.00 (dd, *J* = 7.8 and 1.2 Hz, 1H), 7.89 (s, 1H), 7.83 (d, *J* = 7.8 Hz, 1H), 7.65-7.43 (m, 3H), 7.25-7.20 (m, 2H), 5.36 (s, 2H), 3.80 (s, 3H). MS: *m/z* 311.1 [M+H]⁺.

Example 2

1-(3-Carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione

1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione (1.6 g, 5.2 mmol), NaOH (0.31g, 7.7 mmol), water (40 mL) and methanol (40 mL) were added into a 250 mL three-mouth flask. The mixture was refluxed and reacted for 3 hours. The methanol was removed via evaporation under reduced pressure after the reaction was finished. The mixture was adjusted to pH=2-3 by 3N HCl, forming many precipitates. The mixture was filtered and the solid was washed by water and methanol. The solids were dried to give the title compound as white solid (1.2 g, 78% yield). ¹H NMR (300 M, DMSO-*d*₆): 13.09 (brs, 1H), 11.77 (s, 1H), 8.01 (dd, *J* = 7.7 and 1.4 Hz, 1H), 7.84-7.80 (m, 2H), 7.66-7.61 (m, 1H), 7.53 (d, *J* = 7.8 Hz, 1H), 7.44 (t, *J* = 7.7 Hz, 1H), 7.27-7.21 (m, 2H), 5.36 (s, 2H). MS: *m/z* 297.1 [M+H]⁺.

Example 3

1-(3-(4-(Pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (0.2 mmol), 1-(pyridin-2-yl)piperazine (0.2 mmol), 2-(7-aza-1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU, 0.26 mmol), *N,N*-diisopropylethylamine (DIPEA, 0.4 mmol) and DMF (5 mL) were added into a 50mL two-mouth flask. The obtained mixture was stirred at room temperature for 3 hours. To the mixture was added 50 mL of water and it was extracted by dichloromethane

(DCM, 50 mL x 2). The organic layer was washed by 1N HCl (50 mL x 2) and saturated NaCl aqueous solution (50 mL x 2), dried with anhydrous sodium sulphate, and evaporated under reduced pressure to give the crude product. The crude product was purified by column chromatography (ethyl acetate) to give the title compound as white solid (7.7 mg, 8.6% yield). ¹H NMR (DMSO-d₆): 11.69 (s, 1H), 8.09 (dd, *J* = 5.0 and 1.7 Hz, 1H), 7.99 (dd, *J* = 8.0 and 1.7 Hz, 1H), 7.64 (t, *J* = 7.1 Hz, 1H), 7.53 (t, *J* = 7.8 Hz, 1H), 7.40-7.21 (m, 6H), 6.77 (d, *J* = 8.7 Hz, 1H), 6.64 (dd, *J* = 7.1 and 5.0 Hz, 1H), 5.34 (s, 2H), 3.40-3.20 (m, 8H). MS: *m/z* 442.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine or piperidine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 4

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ¹H NMR (DMSO-d₆): 11.71 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 8.01 (d, *J* = 6.9 Hz, 1H), 7.67-7.61 (m, 1H), 7.42-7.24 (m, 6H), 6.65 (t, *J* = 4.8 Hz, 1H), 5.35 (s, 2H), 3.40-3.20 (m, 8H). MS: *m/z* 443.3 [M+H]⁺.

Example 5

1-(3-(4-Cyclohexylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ¹H NMR (DMSO-d₆): 11.74 (s, 1H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.62 (t, *J* = 7.5 Hz, 1H), 7.43-7.38 (m, 2H), 7.26-7.20 (m, 4H), 5.33 (s, 2H), 3.40-3.20 (m, 4H), 2.21-2.15 (m, 4H), 1.96-1.70 (m, 5H), 1.21-1.08 (m, 6H). MS: *m/z* 447.3 [M+H]⁺.

Example 6

1-(3-(4-([1,2,4]Triazolo[4,3-*b*]pyridazin-6-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ¹H NMR (DMSO-d₆): 11.73 (s, 1H), 9.24 (s, 1H), 8.12 (d, *J* = 10.2 Hz, 1H), 8.02 (d, *J* = 7.8 Hz, 1H), 7.66 (t, *J* = 6.9 Hz, 1H), 7.42-7.24 (m, 7H), 5.36 (s, 2H), 3.40-3.20 (m, 8H). MS: *m/z* 483.3 [M+H]⁺.

Example 7

1-(3-(4-Ethylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.72 (s, 1H), 8.01 (dd, $J = 7.8$ and 1.2 Hz, 1H), 7.63 (t, $J = 8.0$ Hz, 1H), 7.42-7.38 (m, 2H), 7.26-7.23 (m, 4H), 5.34 (s, 2H), 3.40-3.20 (m, 4H), 2.35-2.29 (m, 6H), 0.97 (t, $J = 7.1$ Hz, 3H). MS: m/z 393.2 $[\text{M}+\text{H}]^+$.

Example 8

1-(3-(4-Benzoylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.71 (s, 1H), 7.97 (d, $J = 8.1$ Hz, 1H), 7.60-7.56 (m, 1H), 7.45-7.18 (m, 11H), 5.34 (s, 2H), 3.40-3.20 (m, 8H). MS: m/z 469.3 $[\text{M}+\text{H}]^+$.

Example 9

1-(3-(4-(4-Fluorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.71 (s, 1H), 7.99 (d, $J = 7.8$ Hz, 1H), 7.60-7.19 (m, 11H), 5.34 (s, 2H), 3.60-3.20 (m, 8H). MS: m/z 487.3 $[\text{M}+\text{H}]^+$.

Example 10

1-(3-(4-(4-Chlorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.73 (s, 1H), 7.98-7.97 (m, 1H), 7.63-7.57 (m, 1H), 7.48-7.19 (m, 10H), 5.33 (s, 2H), 3.75-3.15 (m, 8H). MS: m/z 503.2, 505.2 $[\text{M}+\text{H}]^+$.

Example 11

1-(3-(4-(4-Bromobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.74 (s, 1H), 8.02-7.95 (m, 2H), 7.68-7.59 (m, 2H), 7.40-7.15 (m, 8H), 5.35 (s, 2H), 3.40-3.20 (m, 8H). MS: m/z 547.2 $[\text{M}+\text{H}]^+$.

Example 12

1-(3-(4-(4-Methoxybenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.73 (s, 1H), 7.99 (d, $J = 7.2$ Hz, 1H), 7.70-7.50 (m, 1H), 7.40-7.29 (m, 6H), 7.21-7.18 (m, 2H), 6.99-6.96 (m, 2H), 5.33 (s, 2H), 3.77 (s, 3H), 3.56-3.15 (m, 8H). MS: m/z 499.3 $[\text{M}+\text{H}]^+$.

Example 13

1-(3-(4-(Tetrahydro-2H-pyran-4-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (DMSO- d_6): 11.74 (s, 1H), 8.02 (d, $J = 7.5$ Hz, 1H), 7.65 (t, $J = 7.8$ Hz, 1H), 7.45-7.38 (m, 2H), 7.32-7.23 (m, 4H), 5.36 (s, 2H), 3.86-3.82 (m, 2H), 3.75-3.10 (m, 10H), 2.90-2.80 (m, 1H), 1.64-1.48 (m, 4H). MS: m/z 477.3 $[\text{M}+\text{H}]^+$.

Example 14

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

White solid. ^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.01 (d, $J = 6.9$ Hz, 1H), 7.70-7.60 (m, 1H), 7.41-7.39 (m, 2H), 7.30-7.21 (m, 4H), 5.34 (s, 2H), 3.70-3.10 (m, 8H), 2.95-2.89 (m, 1H), 1.80-1.40 (m, 8H). MS: m/z 461.3 $[\text{M}+\text{H}]^+$.

Example 15

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

White solid. ^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.01 (d, $J = 6.6$ Hz, 1H), 7.63 (t, $J = 7.2$ Hz, 1H), 7.41-7.21 (m, 6H), 5.34 (s, 2H), 3.80-3.15 (m, 8H), 1.94-1.91 (m, 1H), 0.72-0.69 (m, 4H). MS: m/z 433.2 $[\text{M}+\text{H}]^+$.

Example 16

1-(3-(4-(Ethylsulfonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

White solid. ^1H NMR (DMSO- d_6): 11.74 (s, 1H), 8.01 (dd, $J = 7.8$ and 1.2 Hz, 1H), 7.70-7.55 (m, 1H), 7.45-7.35 (m, 2H), 7.31-7.19 (m, 4H), 5.35 (s, 2H), 3.75-3.11 (m, 6H), 3.10-2.90 (m, 4H), 1.20 (t, $J = 7.4$ Hz, 3H). MS: m/z 457.2 $[\text{M}+\text{H}]^+$.

Example 17

1-(3-(4-Acetylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

White solid. ^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.01 (d, $J = 6.6$ Hz, 1H), 7.63 (t, $J = 7.2$ Hz, 1H), 7.43-7.37 (m, 2H), 7.30-7.21 (m, 4H), 5.34 (s, 2H), 3.70-3.10 (m, 8H), 1.93 (s, 3H). MS: m/z 407.2 $[\text{M}+\text{H}]^+$.

Example 18

1-(3-(4-Phenylpiperidine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

White solid. ^1H NMR (300 M, DMSO- d_6): 11.75 (s, 1H), 7.99 (dd, $J = 7.8$ and 1.2 Hz, 1H), 7.61 (t, $J = 7.2$ Hz, 1H), 7.41-7.17 (m, 11H), 5.37 (s, 2H), 3.55-3.42 (m, 2H), 3.06-2.99 (m, 1H), 2.78-2.71 (m, 2H), 1.97-1.51 (m, 4H). MS: m/z 407.2 $[\text{M}+\text{H}]^+$.

Example 19

1-(3-(4-Phenylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ¹H NMR (300 M, DMSO-*d*₆): 11.75 (s, 1H), 8.03 (d, *J* = 6.6 Hz, 1H), 7.67 (t, *J* = 7.2 Hz, 1H), 7.46-7.39 (m, 2H), 7.34-7.21 (m, 6H), 6.93-6.91 (m, 2H), 6.81 (t, *J* = 7.2 Hz, 1H), 5.37 (s, 2H), 3.85-2.85 (m, 8H). MS: *m/z* 441.3 [M+H]⁺.

Example 20

1-(3-(4-(Pyrazin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ¹H NMR (300 M, DMSO-*d*₆): 11.75 (s, 1H), 8.31 (s, 1H), 8.11 (s, 1H), 8.02 (d, *J* = 7.8 Hz, 1H), 7.87 (d, *J* = 2.7 Hz, 1H), 7.67 (t, *J* = 7.2 Hz, 1H), 7.44-7.24 (m, 6H), 5.37 (s, 2H), 3.80-3.25 (m, 8H). MS: *m/z* 443.3 [M+H]⁺.

Example 21

1-(4-Fluoro-3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione

The title compound was prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using a procedure similar to those described for the synthesis of compound of Example 1. ¹H NMR (DMSO-*d*₆): 11.75 (s, 1H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.84 (d, *J* = 6.9 Hz, 1H), 7.67-7.55 (m, 2H), 7.32-7.21 (m, 3H), 5.32 (s, 2H), 3.81 (s, 3H). MS: *m/z* 329.1 [M+H]⁺.

Example 22

1-(3-Carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

The title compound was prepared from 1-(4-fluoro-3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione using a procedure similar to those described for the synthesis of compound of Example 2. ¹H NMR (DMSO-*d*₆): 13.23 (brs, 1H), 11.75 (s, 1H), 8.00 (d, *J* = 7.5 Hz, 1H), 7.80 (dd, *J* = 6.9 and 2.1 Hz, 1H), 7.67-7.54 (m, 2H), 7.30-7.21 (m, 3H), 5.31 (s, 2H). MS: *m/z* 315.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 23

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (300 M, DMSO- d_6): 11.73 (s, 1H), 8.00 (d, $J = 6.9$ Hz, 1H), 7.64 (t, $J = 7.8$ Hz, 1H), 7.43-7.21 (m, 5H), 5.30 (s, 2H), 3.56-2.97 (m, 8H), 2.89-2.84 (m, 1H), 1.80-1.40 (m, 8H). MS: m/z 479.3 $[\text{M}+\text{H}]^+$.

Example 24

1-(4-Fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (300 M, DMSO- d_6): 11.74 (s, 1H), 8.12 (d, $J = 3.3$ Hz, 1H), 8.02 (d, $J = 6.6$ Hz, 1H), 7.67 (t, $J = 8.4$ Hz, 1H), 7.58-7.53 (m, 1H), 7.47-7.39 (m, 2H), 7.32-7.24 (m, 3H), 6.82 (d, $J = 8.4$ Hz, 1H), 6.67 (t, $J = 6.9$ Hz, 1H), 5.32 (s, 2H), 3.75-3.67 (m, 2H), 3.61-3.53 (m, 2H), 3.42-3.37 (m, 2H), 3.28-3.20 (m, 2H). MS: m/z 460.3 $[\text{M}+\text{H}]^+$.

Example 25

1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

White solid. ^1H NMR (300 M, DMSO- d_6): 11.72 (s, 1H), 8.37 (d, $J = 4.8$ Hz, 2H), 8.00 (d, $J = 7.8$ Hz, 1H), 7.65 (t, $J = 6.8$ Hz, 1H), 7.48-7.38 (m, 2H), 7.30-7.22 (m, 3H), 6.66 (t, $J = 4.8$ Hz, 1H), 5.31 (s, 2H), 3.70-3.10 (m, 8H). MS: m/z 461.3 $[\text{M}+\text{H}]^+$.

The following compounds were prepared from 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine or piperidine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 26

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.75 (s, 1H), 8.02 (d, $J = 7.8$ Hz, 1H), 7.64 (t, $J = 7.2$ Hz, 1H), 7.50-7.45 (m, 2H), 7.35-7.20 (m, 4H), 5.35 (s, 2H), 3.50-3.10 (m, 9H), 1.80-1.25 (m, 10H). MS: m/z 475.3 $[\text{M}+\text{H}]^+$.

Example 27

1-(3-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.75 (s, 1H), 8.08-8.06 (m, 2H), 8.01 (d, $J = 7.8$ Hz, 1H), 7.87 (s, 1H), 7.67-7.54 (m, 2H), 7.48-7.30 (m, 5H), 7.27-7.20 (m, 2H), 5.37 (s, 2H), 3.80-3.31 (m, 8H). MS: m/z 498.3 $[\text{M}+\text{H}]^+$.

Example 28

1-(3-(4-(Piperidin-1-yl)piperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.76 (s, 1H), 8.03 (d, *J* = 7.2 Hz, 1H), 7.65 (t, *J* = 7.8 Hz, 1H), 7.42-7.21 (m, 6H), 5.41-5.32 (m, 2H), 3.10-2.60 (m, 9H), 1.90-1.30 (m, 10H). MS: *m/z* 447.3 [M+H]⁺.

Example 29

1-(3-(4-(Pyridin-4-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.19 (d, *J* = 6.6 Hz, 2H), 8.02 (d, *J* = 6.3 Hz, 1H), 7.68-7.63 (m, 1H), 7.46-7.39 (m, 2H), 7.34-7.23 (m, 4H), 6.89 (d, *J* = 6.6 Hz, 2H), 5.35 (s, 2H), 3.84-3.31 (m, 8H). MS: *m/z* 442.3 [M+H]⁺.

Example 30

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.03 (d, *J* = 6.4 Hz, 1H), 7.66 (t, *J* = 7.8 Hz, 1H), 7.46-7.39 (m, 2H), 7.32-7.23 (m, 4H), 5.36 (s, 2H), 3.52-3.31 (m, 9H), 2.19-1.75 (m, 6H). MS: *m/z* 447.3 [M+H]⁺.

Example 31

1-(3-(4-(6-Fluorobenzo[*d*]isoxazol-3-yl)piperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.06 (dd, *J* = 8.7 and 6.4 Hz, 1H), 7.95 (dd, *J* = 7.8 and 1.5 Hz, 1H), 7.70 (dd, *J* = 9.3 and 2.1 Hz, 1H), 7.64-7.59 (m, 1H), 7.46-7.14 (m, 7H), 5.36 (s, 2H), 3.45-3.28 (m, 5H), 2.10-2.04 (m, 1H), 1.99-1.94 (m, 3H). MS: *m/z* 499.3 [M+H]⁺.

Example 32

1-(3-(4-(Thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 8.00 (dd, *J* = 8.1 and 1.2 Hz, 1H), 7.76 (d, *J* = 5.1 Hz, 1H), 7.64-7.59 (m, 1H), 7.44-7.11 (m, 8H), 5.34 (s, 2H), 3.81-3.40 (m, 8H). MS: *m/z* 475.2 [M+H]⁺.

Example 33

1-(3-(4-(Furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.02 (d, *J* = 6.9 Hz, 1H), 7.86 (s, 1H), 7.65 (t, *J* = 7.5 Hz, 1H), 7.50-7.15 (m, 6H), 7.00 (d, *J* = 3.3 Hz, 1H), 6.70-6.60 (m, 1H), 5.37 (s, 2H), 3.85-3.20 (m, 8H). MS: *m/z* 459.2 [M+H]⁺.

Example 34

1-(3-(4-(Furan-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.15-7.90 (m, 2H), 7.73 (s, 1H), 7.62 (t, *J* = 7.5 Hz, 1H), 7.50-7.10 (m, 6H), 6.64 (s, 1H), 5.34 (s, 2H), 3.80-3.10 (m, 8H). MS: *m/z* 459.3 [M+H]⁺.

Example 35

1-(3-(4-(Thiophene-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 7.99 (d, *J* = 7.5 Hz, 1H), 7.77 (s, 1H), 7.63-7.58 (m, 2H), 7.41-7.39 (m, 2H), 7.30-7.28 (m, 2H), 7.22-7.17 (m, 3H), 5.34 (s, 2H), 3.81-3.40 (m, 8H). MS: *m/z* 475.3 [M+H]⁺.

Example 36

1-(3-(4-(Pyridine-3-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.67 (dd, *J* = 4.8 and 1.5 Hz, 1H), 8.61 (s, 1H), 8.01 (d, *J* = 7.2 Hz, 1H), 7.82 (d, *J* = 6.8 Hz, 1H), 7.52-7.20 (m, 8H), 5.36 (s, 2H), 3.52-3.31 (m, 8H). MS: *m/z* 470.3 [M+H]⁺.

Example 37

1-(3-(4-(Pyridine-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.78 (s, 1H), 8.68 (d, *J* = 4.2 Hz, 1H), 8.20-7.95 (m, 2H), 7.68-7.27 (m, 9H), 5.43 (s, 2H), 3.90-3.50 (m, 8H). MS: *m/z* 470.3 [M+H]⁺.

Example 38

1-(3-(4-(Pyridine-4-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.62 (s, 1H), 8.58 (d, *J* = 4.2 Hz, 2H), 7.91-7.89 (m, 1H), 7.57-7.44 (m, 1H), 7.29-7.10 (m, 8H), 5.25 (s, 2H), 3.80-2.75 (m, 8H). MS: *m/z* 470.3 [M+H]⁺.

Example 39

1-(3-(4-Phenoxypiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.95 (d, *J* = 8.1 Hz, 1H), 7.57 (t, *J* = 7.8 Hz, 1H), 7.38-7.30 (m, 2H), 7.27-7.14 (m, 6H), 6.92-6.85 (m, 3H), 5.30 (s, 2H), 4.58-4.52 (m, 1H), 3.90-3.10 (m, 4H), 1.95-1.35 (m, 4H). MS: *m/z* 456.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 40

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.01 (d, *J* = 6.6 Hz, 1H), 7.65 (t, *J* = 7.8 Hz, 1H), 7.39-7.22 (m, 5H), 5.31 (s, 2H), 3.52-3.31 (m, 8H), 2.01-1.86 (m, 1H), 0.73-0.70 (m, 4H). MS: *m/z* 451.3 [M+H]⁺.

Example 41

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.00 (dd, *J* = 7.8 and 1.2 Hz, 1H), 7.64 (t, *J* = 7.4 Hz, 1H), 7.43-7.21 (m, 5H), 5.30 (s, 2H), 3.58-3.10 (m, 9H), 1.68-1.00 (m, 10H). MS: *m/z* 493.3 [M+H]⁺.

Example 42

1-(4-Fluoro-3-(4-(pyrazin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.29 (s, 1H), 8.08 (s, 1H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.86 (d, *J* = 2.1 Hz, 1H), 7.65 (t, *J* = 7.5 Hz, 1H), 7.46-7.36 (m, 2H), 7.30-7.22 (m, 3H), 5.31 (s, 2H), 3.81-3.40 (m, 8H). MS: *m/z* 461.3 [M+H]⁺.

Example 43

1-(3-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.08-8.05 (m, 2H), 8.00 (dd, *J* = 7.8 and 1.2 Hz, 1H), 7.64 (t, *J* = 7.2 Hz, 1H), 7.56 (t, *J* = 7.7 Hz, 1H), 7.46-7.41 (m, 3H), 7.31-7.19 (m, 3H), 5.31 (s, 2H), 3.87-3.81 (m, 2H), 3.49-3.31 (m, 6H). MS: *m/z* 516.2 [M+H]⁺.

Example 44

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.64 (t, *J* = 8.0 Hz, 1H), 7.44-7.41 (m, 1H), 7.35 (d, *J* = 5.7 Hz, 1H), 7.28-7.21 (m, 3H), 5.30 (s, 2H), 3.70-3.00 (m, 9H), 2.25-1.60 (m, 6H). MS: *m/z* 465.3 [M+H]⁺.

Example 45

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 7.99 (d, *J* = 7.8 Hz, 1H), 7.70-7.10 (m, 11H), 5.30 (s, 2H), 3.80-3.10 (m, 8H). MS: *m/z* 487.3 [M+H]⁺.

Example 46

1-(4-Fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 8.00 (d, *J* = 7.5 Hz, 1H), 7.75 (d, *J* = 5.1 Hz, 1H), 7.62 (t, *J* = 7.7 Hz, 1H), 7.50-7.15 (m, 6H), 7.12 (t, *J* = 4.4 Hz, 1H), 5.30 (s, 2H), 3.80-3.10 (m, 8H). MS: *m/z* 493.2 [M+H]⁺.

Example 47

1-(4-Fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 8.00 (d, *J* = 6.6 Hz, 1H), 7.83 (s, 1H), 7.64 (t, *J* = 7.1 Hz, 1H), 7.50-7.35 (m, 2H), 7.30-7.15 (m, 3H), 6.99 (d, *J* = 3.3 Hz, 1H), 6.70-6.55 (m, 1H), 5.31 (s, 2H), 3.85-3.10 (m, 8H). MS: *m/z* 477.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-4-fluorobenzoate using a procedure similar to those used for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 48

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.75 (s, 1H), 8.03 (dd, *J* = 7.8 and 1.2 Hz, 1H), 7.66 (t, *J* = 7.8 Hz, 1H), 7.38-7.35 (m, 2H), 7.26 (t, *J* = 7.7 Hz, 1H), 7.20-7.15 (m, 2H), 5.35 (s, 2H), 3.38-3.15 (m, 9H), 1.70-1.55 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 49

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.75 (s, 1H), 8.04 (d, *J* = 7.8 Hz, 1H), 7.67 (d, *J* = 7.4 Hz, 1H), 7.50-7.10 (m, 5H), 5.37 (s, 2H), 3.70-2.90 (m, 9H), 1.80-1.00 (m, 10H). MS: *m/z* 493.3 [M+H]⁺.

Example 50

1-(6-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.35 (d, *J* = 4.5 Hz, 2H), 8.01 (d, *J* = 7.6 Hz, 1H), 7.67 (t, *J* = 8.4 Hz, 1H), 7.40-7.17 (m, 5H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.35 (s, 2H), 3.81-3.14 (m, 8H). MS: *m/z* 461.2 [M+H]⁺.

Example 51

1-(6-Fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.12 (d, *J* = 3.3 Hz, 1H), 8.03 (d, *J* = 6.6 Hz, 1H), 7.70 (t, *J* = 8.0 Hz, 1H), 7.56 (t, *J* = 8.0 Hz, 1H), 7.42-7.18 (m, 5H), 6.76 (d, *J* = 8.4 Hz, 1H), 6.67 (dd, *J* = 6.9 and 5.1 Hz, 1H), 5.38 (s, 2H), 3.70-3.00 (m, 8H). MS: *m/z* 460.3 [M+H]⁺.

Example 52

1-(6-Fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.01 (d, *J* = 8.1 Hz, 1H), 7.76 (d, *J* = 4.8 Hz, 1H), 7.62 (t, *J* = 7.8 Hz, 1H), 7.43-7.32 (m, 3H), 7.22-7.12 (m, 4H), 5.35 (s, 2H), 3.80-2.90 (m, 8H). MS: *m/z* 493.2 [M+H]⁺.

Example 53

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.02 (d, *J* = 7.8 Hz, 1H), 7.65 (t, *J* = 8.0 Hz, 1H), 7.36-7.13 (m, 5H), 5.34 (s, 2H), 3.60-2.80 (m, 9H), 2.20-1.70 (m, 6H). MS: *m/z* 465.3 [M+H]⁺.

Example 54

1-(6-Fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.01 (d, $J = 6.6$ Hz, 1H), 7.84 (s, 1H), 7.64 (t, $J = 7.5$ Hz, 1H), 7.43-7.32 (m, 2H), 7.25-7.16 (m, 3H), 6.96 (d, $J = 3.0$ Hz, 1H), 7.63-6.62 (m, 1H), 5.35 (s, 2H), 3.80-2.90 (m, 8H). MS: m/z 477.3 $[\text{M}+\text{H}]^+$.

The following compounds were prepared from 1-(3-carboxy-6-chlorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-4-chlorobenzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 55

1-(6-Chloro-3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.75 (s, 1H), 8.05 (d, $J = 7.8$ Hz, 1H), 7.69-7.64 (m, 2H), 7.37 (d, $J = 8.1$ Hz, 1H), 7.28 (t, $J = 7.3$ Hz, 1H), 7.08-7.02 (m, 2H), 5.33 (s, 2H), 3.70-2.80 (m, 9H), 1.80-1.55 (m, 8H). MS: m/z 495.3 $[\text{M}+\text{H}]^+$.

Example 56

1-(6-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.36 (d, $J = 4.8$ Hz, 2H), 8.03 (d, $J = 6.6$ Hz, 1H), 7.69-7.64 (m, 2H), 7.38 (d, $J = 6.9$ Hz, 1H), 7.28 (t, $J = 7.5$ Hz, 1H), 7.09-7.04 (m, 2H), 6.65 (t, $J = 4.8$ Hz, 1H), 5.33 (s, 2H), 3.80-3.00 (m, 8H). MS: m/z 477.2 $[\text{M}+\text{H}]^+$.

Example 57

1-(6-Chloro-3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.02 (d, $J = 7.5$ Hz, 1H), 7.75-7.55 (m, 2H), 7.34 (d, $J = 8.1$ Hz, 1H), 7.25 (t, $J = 7.5$ Hz, 1H), 7.10-6.90 (m, 2H), 5.30 (s, 2H), 3.60-2.80 (m, 9H), 1.80-1.00 (m, 10H). MS: m/z 509.3 $[\text{M}+\text{H}]^+$.

Example 58

1-(6-Chloro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.73 (s, 1H), 8.11 (d, $J = 3.6$ Hz, 1H), 8.03 (d, $J = 7.8$ Hz, 1H), 7.80-7.60 (m, 2H), 7.55 (t, $J = 6.9$ Hz, 1H), 7.39 (d, $J = 6.6$ Hz, 1H), 7.30 (t, $J = 7.5$ Hz, 1H),

7.15-7.00 (m, 2H), 6.72 (d, $J = 8.4$ Hz, 1H), 6.66 (dd, $J = 6.8$ and 5.0 Hz, 1H), 5.34 (s, 2H), 3.70-3.00 (m, 8H). MS: m/z 476.2 $[M+H]^+$.

The following compounds were prepared from 1-(3-carboxyl-2-chlorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-2-chlorobenzoate using a procedure similar to those used for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 59

1-(2-Chloro-3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.78 (s, 1H), 8.07 (d, $J = 7.8$ Hz, 1H), 7.67 (t, $J = 7.5$ Hz, 1H), 7.45-7.20 (m, 3H), 7.15-6.95 (m, 2H), 5.50-5.15 (m, 2H), 3.80-3.10 (m, 8H), 3.04-2.94 (m, 1H), 1.90-1.35 (m, 8H). MS: m/z 495.3 $[M+H]^+$.

Example 60

1-(2-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.88 (s, 1H), 8.49 (d, $J = 4.8$ Hz, 2H), 8.16 (d, $J = 7.8$ Hz, 1H), 7.77 (t, $J = 7.8$ Hz, 1H), 7.50-7.30 (m, 3H), 7.25-7.05 (m, 2H), 6.78 (t, $J = 7.7$ Hz, 1H), 5.60-5.30 (m, 2H), 4.10-3.70 (m, 8H). MS: m/z 477.2 $[M+H]^+$.

The following compounds were prepared from 1-(3-carboxy-2-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-2-fluorobenzoate using a procedure similar to those used for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis compound of Example 3.

Example 61

1-(2-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.71 (s, 1H), 8.37 (d, $J = 4.8$ Hz, 2H), 8.03 (d, $J = 6.6$ Hz, 1H), 7.66 (t, $J = 7.1$ Hz, 1H), 7.45-7.05 (m, 5H), 6.66 (t, $J = 4.7$ Hz, 1H), 5.35 (s, 2H), 3.90-3.60 (m, 8H). MS: m/z 461.3 $[M+H]^+$.

Example 62

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-2-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.74 (s, 1H), 8.03 (d, *J* = 7.5 Hz, 1H), 7.66 (t, *J* = 7.7 Hz, 1H), 7.40-7.10 (m, 5H), 5.34 (s, 2H), 3.80-3.20 (m, 8H), 3.10-2.80 (m, 1H), 1.90-1.40 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-5-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-5-fluorobenzoate using a procedure similar to those used for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis compound of Example 3.

Example 63

1-(5-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 8.00 (d, *J* = 7.8 Hz, 1H), 7.64 (t, *J* = 8.0 Hz, 1H), 7.30-7.15 (m, 5H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.34 (s, 2H), 3.80-3.50 (m, 8H). MS: *m/z* 461.3 [M+H]⁺.

Example 64

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-5-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 8.00 (d, *J* = 9.0 Hz, 1H), 7.64 (t, *J* = 7.1 Hz, 1H), 7.31-7.15 (m, 5H), 5.34 (s, 2H), 3.70-3.10 (m, 8H), 2.94-2.86 (m, 1H), 1.80-1.40 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-4-chlorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-chlorobenzoate using a procedure similar to those used for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis compound of of Example 3.

Example 65

1-(4-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.38 (d, *J* = 4.6 Hz, 2H), 8.16 (d, *J* = 9.0 Hz, 1H), 7.69 (t, *J* = 8.7 Hz, 1H), 7.54-7.29 (m, 5H), 6.67 (t, *J* = 4.6 Hz, 1H), 5.51-5.39 (m, 2H), 3.98-3.57 (m, 8H). MS: *m/z* 477.2 [M+H]⁺.

Example 66

1-(4-Chloro-3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.60 (s, 1H), 7.91 (d, *J* = 7.8 Hz, 1H), 7.53 (t, *J* = 7.5 Hz, 1H), 7.39 (d, *J* = 9.0 Hz, 1H), 7.28-7.26 (m, 2H), 7.16-7.07 (m, 2H), 5.31-5.13 (m, 2H), 3.48-2.90 (m, 9H), 1.59-1.09 (m, 10H). MS: *m/z* 509.3 [M+H]⁺.

The following compounds were prepared from 1-((2-carboxy-furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and ethyl 5-(chloromethyl)furan-2-carboxylate using a procedure similar to those described for the synthesis of compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 67

1-((2-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.75 (s, 1H), 8.01 (d, *J* = 7.5 Hz, 1H), 7.74 (t, *J* = 7.5 Hz, 1H), 7.57 (d, *J* = 8.1 Hz, 1H), 7.28 (t, *J* = 7.5 Hz, 1H), 6.96 (d, *J* = 3.3 Hz, 1H), 6.59 (d, *J* = 3.3 Hz, 1H), 5.36 (s, 2H), 3.61-3.14 (m, 9H), 1.73-1.54 (m, 8H). MS: *m/z* 451.3 [M+1]⁺.

Example 68

1-((2-(4-(Benzo[*d*]isothiazol-3-yl)piperazine-1-carbonyl)furan-5-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.74 (s, 1H), 8.10-8.08 (m, 2H), 8.02 (d, *J* = 7.8 Hz, 1H), 7.76 (t, *J* = 4.5 Hz, 1H), 7.61-7.57 (m, 2H), 7.48 (t, *J* = 7.5 Hz, 1H), 7.30-7.25 (m, 1H), 7.00 (d, *J* = 3.3 Hz, 1H), 6.60 (d, *J* = 3.3 Hz, 1H), 5.39 (s, 2H), 3.85-3.78 (m, 4H), 3.49-3.40 (m, 4H). MS: *m/z* 488.2 [M+1]⁺.

The following compounds were prepared from 1-((2-carboxypyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 6-(bromomethyl)pyridine-2-carboxylate using a procedure similar to those described for

the synthesis of compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 69

1-((2-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.16 (d, *J* = 6.3 Hz, 1H), 7.98-7.96 (m, 1H), 7.64-7.59 (m, 3H), 7.32-7.26 (m, 2H), 5.57 (s, 2H), 3.70-3.58 (m, 4H), 3.31-2.92 (m, 5H), 1.84-1.70 (m, 8H). MS: *m/z* 462.3 [M+H]⁺.

Example 70

1-((2-(4-(Pyridin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.15-8.12 (m, 2H), 7.96 (t, *J* = 7.5 Hz, 1H), 7.67-7.58 (m, 4H), 7.30-7.25 (m, 2H), 6.80-6.71 (m, 2H), 5.57 (s, 2H), 3.81-3.78 (m, 2H), 3.59-3.55 (m, 2H), 3.44-3.41 (m, 2H), 3.23-3.19 (m, 2H). MS: *m/z* 443.3 [M+H]⁺.

Example 71

1-((2-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.36 (d, *J* = 4.8 Hz, 2H), 8.14 (d, *J* = 6.9 Hz, 1H), 7.96 (t, *J* = 7.5 Hz, 1H), 7.65-7.58 (m, 3H), 7.31-7.28 (m, 2H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.57 (s, 2H), 3.85-3.82 (m, 2H), 3.78-3.73 (m, 2H), 3.57-3.53 (m, 2H), 3.39-3.35 (m, 2H). MS: *m/z* 444.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 72

1-(3-(4-(Thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.03 (d, *J* = 7.3 Hz, 1H), 7.66 (t, *J* = 7.7 Hz, 1H), 7.46-7.23 (m, 6H), 7.18 (d, *J* = 3.6 Hz, 1H), 6.88 (d, *J* = 3.6 Hz, 1H), 5.37 (s, 2H), 3.80-3.40 (m, 8H). MS: *m/z* 448.3 [M+H]⁺.

Example 73

1-(3-(4-(Cyclohexylmethyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.75 (s, 1H), 8.02 (dd, *J* = 8.0 and 1.4 Hz, 1H), 7.67-7.62 (m, 1H), 7.41-7.39 (m, 2H), 7.27-7.23 (m, 4H), 5.35 (s, 2H), 3.54-3.15 (m, 4H), 2.32-2.03 (m, 6H), 1.73-1.44 (m, 5H), 1.23-1.06 (m, 6H). MS: *m/z* 461.3 [M+1]⁺.

Example 74

1-(3-(4-(Cyclopentylmethyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.73 (s, 1H), 8.03 (d, *J* = 7.5 Hz, 1H), 7.65 (d, *J* = 7.5 Hz, 1H), 7.42-7.23 (m, 6H), 5.36 (s, 2H), 3.54-2.87 (m, 6H), 2.25-2.20 (m, 4H), 1.80-1.24 (m, 9H). MS: *m/z* 447.3 [M+1]⁺.

Example 75

1-(3-(4-(Cyclohexylsulfonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 8.01 (dd, *J* = 8.0 and 1.4 Hz, 1H), 7.64-7.60 (m, 1H), 7.42-7.38 (m, 2H), 7.29-7.16 (m, 4H), 5.35 (s, 2H), 3.56-3.52 (m, 2H), 3.20-3.05 (m, 6H), 1.96-1.93 (m, 1H), 1.80-1.76 (m, 2H), 1.62-1.58 (m, 2H), 1.27-1.22 (m, 6H). MS: *m/z* 511.3 [M+1]⁺.

Example 76

7-Fluoroquinazoline-2,4(1*H*,3*H*)-dione

To a mixture of 2-amino-4-fluorobenzoic acid (4.0 g, 26 mmol) and potassium cyanate (3.5 g, 43 mmol) in water (200 mL) was added acetic acid (3 mL, 45 mmol) and the mixture was stirred at room temperature for about 5 h. To the mixture was added NaOH (15 g, 375 mmol) and it was stirred for another 1 h. The mixture was filtered and the solid was mixed in hot water, and it was adjusted to pH=5~6 by addition of acetic acid. The mixture was filtered and washed with water, dried to give the title compound (3.0 g, 65% yield) as gray solid. MS: *m/z* 181.1 [M+H]⁺

The following compounds were prepared from 1-(3-carboxybenzyl)-7-fluoroquinazoline-2,4(1*H*,3*H*)-dione (prepared from 7-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 77

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.81 (s, 1H), 8.08 (t, *J* = 7.6 Hz, 1H), 7.44-7.31 (m, 4H), 7.18-7.12 (m, 2H), 5.34 (s, 2H), 3.70-2.80 (m, 9H), 1.80-1.55 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 78

7-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.79 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 8.05 (t, *J* = 7.6 Hz, 1H), 7.45-7.31 (m, 4H), 7.17-7.07 (m, 2H), 6.65 (t, *J* = 4.8 Hz, 1H), 5.33 (s, 2H), 3.78-3.55 (m, 8H). MS: *m/z* 461.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)-7-methylquinazoline-2,4(1*H*,3*H*)-dione (prepared from 7-methylquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 79

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.04 (d, *J* = 8.1 Hz, 1H), 7.51-7.49 (m, 2H), 7.40-7.38 (m, 2H), 7.15-7.11 (m, 2H), 5.45 (s, 2H), 3.75-3.50 (m, 8H), 3.12-3.09 (m, 1H), 2.39 (s, 3H), 1.88-1.64 (m, 8H). MS: *m/z* 475.3 [M+H]⁺.

Example 80

7-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.38-8.36 (m, 2H), 8.04 (d, *J* = 7.8 Hz, 1H), 7.52-7.50 (m, 2H), 7.42-7.40 (m, 2H), 7.16-7.13 (m, 2H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.46 (s, 2H), 3.92-3.56 (m, 8H), 2.40 (s, 3H). MS: *m/z* 457.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-methylquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 81

6-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.34 (d, *J* = 4.8 Hz, 2H), 7.94 (s, 1H), 7.49-7.47 (m, 3H), 7.39-7.34 (m, 2H), 7.18 (d, *J* = 8.4 Hz, 1H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.44 (s, 2H), 3.89-3.39 (m, 8H), 2.39 (s, 3H). MS: *m/z* 457.3 [M+H]⁺.

Example 82

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 7.95 (s, 1H), 7.49-7.47 (m, 3H), 7.38-7.35 (m, 2H), 7.18 (d, *J* = 8.7 Hz, 1H), 5.43 (s, 2H), 3.71-3.69 (m, 4H), 3.49-3.47 (m, 4H), 3.09-3.08 (m, 1H), 2.39 (s, 3H), 1.88-1.63 (m, 8H). MS: *m/z* 475.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-methylquinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 83

1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.37 (d, *J* = 4.8 Hz, 2H), 7.96 (s, 1H), 7.54-7.51 (m, 2H), 7.40 (dd, *J* = 6.3 and 2.1 Hz, 1H), 7.28-7.21 (m, 2H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.41 (s, 2H), 3.93-3.56 (m, 8H), 2.41 (s, 3H). MS: *m/z* 475.3 [M+H]⁺.

Example 84

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.54 (s, 1H), 7.71 (s, 1H), 7.38-7.24 (m, 3H), 7.16 (t, *J* = 9.0 Hz, 1H), 7.04 (d, *J* = 8.7 Hz, 1H), 5.19 (s, 2H), 3.45-2.99 (m, 9H), 2.22 (s, 3H), 1.59-1.49 (m, 5H), 1.22-1.13 (m, 5H). MS: *m/z* 507.3 [M+H]⁺.

Example 85

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.63 (s, 1H), 7.80 (s, 1H), 7.46-7.22 (m, 4H), 7.12 (d, *J*=8.4 Hz, 1H), 5.27 (s, 2H), 3.55-2.97 (m, 8H), 2.99-2.84 (m, 1H), 2.30 (s, 3H), 1.73-1.49 (m, 8H). MS: *m/z* 493.3 [M+H]⁺.

The following compounds were prepared from 6-bromo-1-(3-carboxybenzyl) quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-bromoquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 86

6-Bromo-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 8.05 (d, *J*=2.4 Hz, 1H), 7.79 (dd, *J*=9.0 and 2.4 Hz, 1H), 7.40-7.38 (m, 2H), 7.30-7.27 (m, 2H), 7.16 (d, *J*=9.0 Hz, 1H), 5.32 (s, 2H), 3.80-3.11 (m, 9H), 1.68-1.59 (m, 5H), 1.32-1.19 (m, 5H). MS: *m/z* 553.3 [M+H]⁺.

Example 87

6-Bromo-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 8.04 (d, *J*=2.4 Hz, 1H), 7.78 (dd, *J*=9.0 and 2.4 Hz, 1H), 7.41-7.38 (m, 2H), 7.29-7.27 (m, 2H), 7.15 (d, *J*=8.7 Hz, 1H), 5.32 (s, 2H), 3.57-2.90 (m, 9H), 1.71-1.51 (m, 8H). MS: *m/z* 539.2 [M+H]⁺.

Example 88

6-Bromo-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.38 (d, *J*=4.8 Hz, 2H), 8.25-8.24 (m, 1H), 7.78 (dd, *J*=8.9 and 2.3 Hz, 1H), 7.52-7.50 (m, 2H), 7.42-7.41 (m, 2H), 7.25 (d, *J*=9.3 Hz, 1H), 6.66 (t, *J*=4.6 Hz, 1H), 5.45 (s, 2H), 3.93-3.40 (m, 8H). MS: *m/z* 521.2 [M+H]⁺.

Example 89

6-Nitroquinazoline-2,4(1*H*,3*H*)-dione

A mixture of 2-amino-5-nitrobenzoic acid (0.588 g, 3.23 mmol) and urea (1.164 g, 19.38 mmol) was heated at 200°C under N₂ for 1 h. The mixture was cooled to room temperature and 4 M NaOH was added until pH = 14. It was acidified to pH = 5.0 via addition of AcOH. The mixture was filtered and the yellow solid was dried to give the title compound (0.49 g, 72.8%) as a yellow solid. MS: m/z 208.1 [M+1]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-nitroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those for compounds of Example 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 90

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 12.22 (s, 1H), 8.77 (d, *J* = 2.4 Hz, 1H), 8.51 (dd, *J* = 9.3 and 2.4 Hz, 1H), 7.52-7.46 (m, 3H), 7.43-7.39 (m, 2H), 5.50 (s, 2H), 3.58-2.96 (m, 9H), 1.68-1.31 (m, 8H). MS: m/z 506.3 [M+H]⁺.

Example 91

6-Nitro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 12.10 (s, 1H), 8.66 (d, *J* = 2.7 Hz, 1H), 8.44 (d, *J* = 9.3 Hz, 1H), 8.33 (d, *J* = 4.5 Hz, 2H), 7.45-7.39 (m, 3H), 7.34-7.32 (m, 2H), 6.65 (t, *J* = 4.6 Hz, 1H), 5.42 (s, 2H), 3.77-3.46 (m, 8H). MS: m/z 488.3 [M+H]⁺.

Example 92

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 12.21 (s, 1H), 8.77 (d, *J* = 2.7 Hz, 1H), 8.51 (dd, *J* = 9.2 and 2.6 Hz, 1H), 7.52-7.50 (m, 3H), 7.44-7.39 (m, 2H), 5.50 (s, 2H), 3.68-3.24 (m, 9H), 1.78-1.68 (m, 5H), 1.45-1.23 (m, 5H). MS: m/z 520.3 [M+H]⁺.

The following compounds were prepared from 1-(4-carboxybenzyl)quinazoline

-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 4-(bromomethyl)benzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 93

1-(4-(4-Benzoylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.17 (d, *J* = 7.5 Hz, 1H), 7.68-7.63 (m, 1H), 7.53-7.41 (m, 9H), 7.30-7.27 (m, 2H), 5.46 (s, 2H), 3.76-3.40 (m, 8H). MS: *m/z* 469.2 [M+H]⁺.

Example 94

1-(4-(4-(Pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.15 (d, *J* = 8.1 Hz, 1H), 8.11 (d, *J* = 5.1 Hz, 1H), 7.67-7.59 (m, 2H), 7.49-7.43 (m, 4H), 7.32-7.27 (m, 2H), 6.86 (d, *J* = 8.4 Hz, 1H), 6.72 (t, *J* = 6.0 Hz, 1H), 5.46 (s, 2H), 3.85-3.42 (m, 8H). MS: *m/z* 442.2 [M+H]⁺.

Example 95

1-(4-(4-(4-Methoxybenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.15 (d, *J* = 7.5 Hz, 1H), 7.64 (t, *J* = 7.5 Hz, 1H), 7.43-7.41 (m, 6H), 7.31-7.26 (m, 2H), 7.02-6.99 (m, 2H), 5.45 (s, 2H), 3.84 (s, 3H), 3.80-3.40 (m, 8H). MS: *m/z* 499.2 [M+H]⁺.

Example 96

1-(4-(4-(4-Fluorobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.15 (d, *J* = 6.9 Hz, 1H), 7.66-7.61 (m, 1H), 7.59-7.44 (m, 6H), 7.28-7.21 (m, 4H), 5.45 (s, 2H), 3.65-3.40 (m, 8H). MS: *m/z* 487.2 [M+H]⁺.

Example 97

1-(4-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.35 (d, *J* = 4.8 Hz, 2H), 8.16 (dd, *J* = 8.0 and 1.1 Hz, 1H), 7.65 (t, *J* = 6.9 Hz, 1H), 7.49-7.41 (m, 4H), 7.32-7.28 (m, 2H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.46 (s, 2H), 3.80-3.40 (m, 8H). MS: *m/z* 443.3 [M+H]⁺.

Example 98

1-(4-(4-(4-Bromobenzoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

-dione

¹H NMR (CD₃OD): 8.16 (d, *J* = 7.8 Hz, 1H), 7.88-7.63 (m, 3H), 7.49-7.39 (m, 6H), 7.32-7.27 (m, 2H), 5.45 (s, 2H), 3.90-3.42 (m, 8H). MS: 547.2 [M+H]⁺.

Example 99

1-(4-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.15 (dd, *J* = 8.1 and 1.5 Hz, 1H), 7.64 (t, *J* = 8.0 Hz, 1H), 7.49-7.42 (m, 4H), 7.31-7.26 (m, 2H), 5.45 (s, 2H), 3.83-3.39 (m, 8H), 3.09-3.00 (m, 1H), 2.02-1.51 (m, 8H). MS: 461.3 [M+H]⁺.

The following compounds were prepared from 5-chloro-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 5-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 100

5-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.38 (d, *J* = 4.8 Hz, 2H), 7.60-7.50 (m, 3H), 7.45-7.39 (m, 2H), 7.35-7.20 (m, 2H), 7.26 (d, *J* = 8.4 Hz, 1H), 6.66 (t, *J* = 4.7 Hz, 1H), 5.48 (s, 2H), 4.00-3.40 (m, 8H). MS: *m/z* 477.2 [M+H]⁺.

Example 101

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 7.60-7.49 (m, 3H), 7.40-7.35 (m, 2H), 7.31 (d, *J* = 9.0 Hz, 1H), 7.23 (d, *J* = 9.0 Hz, 1H), 5.46 (s, 2H), 3.85-2.90 (m, 9H), 2.00-1.55 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

Example 102

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 7.53 (t, *J* = 8.3 Hz, 1H), 7.45-7.39 (m, 2H), 7.30-7.20 (m, 3H), 7.15 (d, *J* = 8.7 Hz, 1H), 5.34 (s, 2H), 3.70-3.00 (m, 9H), 1.80-1.00 (m, 10H). MS: *m/z* 509.3 [M+H]⁺.

The following compounds were prepared from 5-chloro-1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 5-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 103

5-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 8.39 (d, *J* = 4.5 Hz, 2H), 7.50 (t, *J* = 7.8 Hz, 1H), 7.47-7.39 (m, 2H), 7.32-7.25 (m, 2H), 7.20 (d, *J* = 9.0 Hz, 1H), 6.67 (t, *J* = 4.5 Hz, 1H), 5.32 (s, 2H), 3.81-3.33 (m, 8H). MS: *m/z* 495.2 [M+H]⁺.

Example 104

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 7.58 (t, *J* = 7.8 Hz, 1H), 7.47-7.42 (m, 1H), 7.38 (d, *J* = 6.0 Hz, 1H), 7.31-7.25 (m, 2H), 7.18-7.14 (m, 1H), 5.31 (s, 2H), 3.70-3.00 (m, 9H), 1.85-1.40 (m, 8H). MS: *m/z* 513.3 [M+H]⁺.

Example 105

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.53 (t, *J* = 8.3 Hz, 1H), 7.45-7.41 (m, 1H), 7.35 (dd, *J* = 6.2 and 2.0 Hz, 1H), 7.31-7.20 (m, 2H), 7.15 (s, 1H), 5.29 (s, 2H), 3.70-3.00 (m, 9H), 1.80-1.00 (m, 10H). MS: *m/z* 527.3 [M+H]⁺.

Example 106

5-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.66 (s, 1H), 8.11 (d, *J* = 1.8 Hz, 1H), 7.57-7.52 (m, 2H), 7.45-7.37 (m, 2H), 7.31-7.20 (m, 2H), 7.17 (d, *J* = 8.7 Hz, 1H), 6.81 (d, *J* = 8.7 Hz, 1H), 6.56 (t, *J* = 6.0 Hz, 1H), 5.30 (s, 2H), 3.80-3.10 (m, 8H). MS: *m/z* 494.2 [M+H]⁺.

Example 107

5-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 7.76 (dd, *J* = 5.0 and 1.1 Hz, 1H), 7.46 (t, *J* = 8.4 Hz, 1H), 7.49-7.36 (m, 3H), 7.30-7.19 (m, 2H), 7.15-7.05 (m, 2H), 5.30 (s, 2H), 3.80-3.15 (m, 8H). MS: *m/z* 527.2 [M+H]⁺.

The following compounds were prepared from 5-chloro-1-(6-fluoro-3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 5-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-4-fluorobenzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 108

5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 7.55 (t, *J* = 8.7 Hz, 1H), 7.45-7.34 (m, 2H), 7.27 (d, *J* = 7.5 Hz, 1H), 7.17 (d, *J* = 6.9 Hz, 1H), 7.13 (d, *J* = 7.8 Hz, 1H), 5.33 (s, 2H), 3.80-3.00 (m, 9H), 1.80-1.10 (m, 10H). MS: *m/z* 527.3 [M+H]⁺.

Example 109

5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 7.57 (t, *J* = 8.3 Hz, 1H), 7.45-7.25 (m, 3H), 7.20 (d, *J* = 7.5 Hz, 1H), 7.14 (d, *J* = 8.7 Hz, 1H), 5.35 (s, 2H), 3.70-2.80 (m, 9H), 1.85-1.40 (m, 8H). MS: *m/z* 513.3 [M+H]⁺.

Example 110

5-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 8.35 (d, *J* = 4.5 Hz, 2H), 7.56 (t, *J* = 8.4 Hz, 1H), 7.45-7.27 (m, 3H), 7.22 (d, *J* = 6.6 Hz, 1H), 7.15 (d, *J* = 8.7 Hz, 1H), 6.64 (t, *J* = 4.5 Hz, 1H), 5.34 (s, 2H), 3.80-3.10 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

The following compounds were prepared from 6-chloro-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-

(bromomethyl)benzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 111

6-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.37 (d, *J* = 4.8 Hz, 2H), 8.09 (d, *J* = 2.7 Hz, 1H), 7.64 (dd, *J* = 9.2 and 2.6 Hz, 1H), 7.55-7.45 (m, 2H), 7.45-7.35 (m, 2H), 7.32 (d, *J* = 9.0 Hz, 1H), 6.66 (t, *J* = 4.7 Hz, 1H), 5.45 (s, 2H), 4.00-3.33 (m, 8H). MS: *m/z* 477.2 [M+H]⁺.

Example 112

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.87 (s, 1H), 7.94 (d, *J* = 2.7 Hz, 1H), 7.70 (dd, *J* = 9.0 and 2.7 Hz, 1H), 7.46-7.38 (m, 2H), 7.31-7.23 (m, 3H), 5.35 (s, 2H), 3.70-2.80 (m, 9H), 1.90-1.40 (m, 8H). MS: *m/z* 495.2 [M+H]⁺.

Example 113

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.88 (s, 1H), 7.94 (d, *J* = 2.4 Hz, 1H), 7.74 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.46-7.37 (m, 2H), 7.32-7.23 (m, 3H), 5.35 (s, 2H), 3.80-3.10 (m, 9H), 1.80-1.10 (m, 10H). MS: *m/z* 509.3 [M+H]⁺.

Example 114

6-Chloro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.95 (s, 1H), 8.21-8.20 (m, 1H), 8.02 (s, 1H), 7.82-7.78 (m, 1H), 7.66-7.60 (m, 1H), 7.49-7.42 (m, 5H), 6.89 (d, *J* = 3.0 Hz, 1H), 6.75 (t, *J* = 6.0 Hz, 1H), 5.44 (s, 2H), 3.85-3.29 (m, 8H). MS: *m/z* 476.3 [M+H]⁺.

Example 115

6-Chloro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.87 (s, 1H), 7.92 (d, *J* = 2.7 Hz, 1H), 7.76 (d, *J* = 4.8 Hz, 1H), 7.67 (dd, *J* = 8.9 and 2.6 Hz, 1H), 7.50-7.35 (m, 3H), 7.32-7.29 (m, 2H), 7.23 (d, *J* = 9.0 Hz, 1H), 7.12 (dd, *J* = 4.8 and 3.6 Hz, 1H), 5.33 (s, 2H), 3.79-3.45 (m, 8H). MS: *m/z* 509.2 [M+H]⁺.

Example 116

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.87 (s, 1H), 7.93 (d, *J* = 2.4 Hz, 1H), 7.68 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.43-7.35 (m, 2H), 7.31-7.22 (m, 3H), 5.33 (s, 2H), 3.60-3.00 (m, 9H), 2.20-1.60 (m, 6H). MS: *m/z* 481.3 [M+H]⁺.

The following compounds were prepared from 6-chloro-1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 117

6-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.84 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 7.92 (s, 1H), 7.69 (dd, *J* = 8.9 and 2.6 Hz, 1H), 7.47-7.36 (m, 2H), 7.30-7.24 (m, 2H), 6.65 (t, *J* = 4.8 Hz, 1H), 5.30 (s, 2H), 3.78-3.21 (m, 8H). MS: *m/z* 495.2 [M+H]⁺.

Example 118

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.85 (s, 1H), 7.92 (d, *J* = 2.4 Hz, 1H), 7.68 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.44-7.41 (m, 1H), 7.36-7.34 (m, 1H), 7.30-7.15 (m, 2H), 5.29 (s, 2H), 3.70-2.80 (m, 9H), 1.80-1.40 (m, 8H). MS: *m/z* 513.3 [M+H]⁺.

Example 119

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.85 (s, 1H), 7.92 (d, *J* = 2.1 Hz, 1H), 7.68 (dd, *J* = 8.7 and 2.4 Hz, 1H), 7.45-7.39 (m, 1H), 7.35 (d, *J* = 5.4 Hz, 1H), 7.31-7.15 (m, 2H), 5.29 (s, 2H), 3.70-3.00 (m, 9H), 1.80-1.10 (m, 10H). MS: *m/z* 527.3 [M+H]⁺.

Example 120

6-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 8.10 (d, *J* = 3.6 Hz, 1H), 7.92 (s, 1H), 7.69 (d, *J* = 9.0 Hz, 1H), 7.54 (t, *J* = 9.0 Hz, 1H), 7.45-7.36 (m, 2H), 7.30-7.23 (m, 2H), 6.81 (d, *J* = 7.8 Hz, 1H), 6.66 (t, *J* = 6.3 Hz, 1H), 5.30 (s, 2H), 3.68-3.21 (m, 8H). MS: *m/z* 494.2 [M+H]⁺.

Example 121

6-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.86 (s, 1H), 7.92 (s, *J* = 2.4 Hz, 1H), 7.76 (d, *J* = 4.2 Hz, 1H), 7.68 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.46-7.36 (m, 3H), 7.30-7.22 (m, 2H), 7.12 (dd, *J* = 5.0 and 3.8 Hz, 1H), 5.30 (s, 2H), 3.68-3.21 (m, 8H). MS: *m/z* 527.2 [M+H]⁺.

Example 122

6-Chloro-1-(3-(4-(cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.85 (s, 1H), 7.94 (s, 1H), 7.70 (dd, *J* = 8.4 and 2.1 Hz, 1H), 7.47-7.23 (m, 4H), 5.31 (s, 2H), 3.59-3.09 (m, 8H), 2.01-1.96 (m, 1H), 0.77-0.69 (m, 4H). MS: *m/z* 485.3 [M+H]⁺.

The following compounds were prepared from 6-chloro-1-(6-fluoro-3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-chloroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-4-fluorobenzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 123

6-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.88 (s, 1H), 8.36 (d, *J* = 4.5 Hz, 2H), 7.93 (d, *J* = 2.4 Hz, 1H), 7.73 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.36-7.20 (m, 4H), 6.65 (t, *J* = 4.8 Hz, 1H), 5.37 (s, 2H), 3.78-3.21 (m, 8H). MS: *m/z* 495.2 [M+H]⁺.

Example 124

6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.87 (s, 1H), 7.94 (d, *J* = 2.4 Hz, 1H), 7.71 (dd, *J* = 9.2 and 2.3 Hz, 1H), 7.42-7.32 (m, 2H), 7.23-7.17 (m, 2H), 5.34 (s, 2H), 3.59-3.09 (m, 9H), 1.71-1.56 (m, 8H). MS: *m/z* 513.3 [M+H]⁺.

Example 125

6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.87 (s, 1H), 7.93 (d, *J* = 2.4 Hz, 1H), 7.70 (dd, *J* = 8.9 and 2.6 Hz, 1H), 7.41-7.31 (m, 2H), 7.22-7.16 (m, 2H), 5.33 (s, 2H), 3.54-3.10 (m, 9H), 1.70-1.15 (m, 10H). MS: *m/z* 527.3 [M+H]⁺.

Example 126

6-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.86 (s, 1H), 7.93 (d, *J* = 2.4 Hz, 1H), 7.76 (d, *J* = 4.8 Hz, 1H), 7.67 (dd, *J* = 9.0 and 1.8 Hz, 1H), 7.43-7.32 (m, 3H), 7.22-7.19 (m, 2H), 7.12 (t, *J* = 4.2 Hz, 1H), 5.33 (s, 2H), 3.54-3.19 (m, 8H). MS: *m/z* 527.2 [M+H]⁺.

The following compounds were prepared from 5-fluoro-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 5-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 127

5-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.35 (d, *J* = 4.8 Hz, 2H), 7.61 (td, *J* = 8.4 and 5.7 Hz, 1H), 7.50-7.32 (m, 4H), 7.06 (d, *J* = 8.4 Hz, 1H), 6.99 (dd, *J* = 10.8 and 8.4 Hz, 1H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.48 (s, 2H), 3.89-3.38 (m, 8H). MS: *m/z* 461.2 [M+H]⁺.

Example 128

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.65-7.57 (m, 1H), 7.41-7.39 (m, 2H), 7.31-7.27 (m, 2H), 7.05-6.97 (m, 2H), 5.32 (s, 2H), 3.69-2.90 (m, 9H), 1.81-1.51 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 129

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 7.62 (td, *J* = 8.4 and 6.0 Hz, 1H), 7.46-7.41 (m, 2H), 7.32-7.29 (m, 2H), 7.05-6.99 (m, 2H), 5.34 (s, 2H), 3.79-3.11 (m, 9H), 1.71-1.61 (m, 5H), 1.33-1.17 (m, 5H). MS: *m/z* 493.3 [M+H]⁺.

Example 130

5-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 8.10 (d, *J* = 3.6 Hz, 1H), 7.66-7.51 (m, 2H), 7.44-7.29 (m, 4H), 7.04-6.98 (m, 2H), 6.80 (d, *J* = 8.7 Hz, 1H), 6.64 (t, *J* = 6.6 Hz, 1H), 5.33 (s, 2H), 3.79-3.41 (m, 8H). MS: *m/z* 460.3 [M+H]⁺.

Example 131

5-Fluoro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.77 (d, *J* = 5.1 Hz, 1H), 7.59 (td, *J* = 6.0 and 8.4 Hz, 1H), 7.44-7.29 (m, 5H), 7.14 (d, *J* = 3.9 Hz, 1H), 7.02-6.95 (m, 2H), 5.33 (s, 2H), 3.64-3.41 (m, 8H). MS: *m/z* 493.2 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxy-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione (prepared from 5-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using a procedure similar to those used for

Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 132

5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.65 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 7.95-7.66 (m, 1H), 7.47-7.38 (m, 2H), 7.28 (t, *J* = 9.0 Hz, 1H), 7.05-6.99 (m, 2H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.30 (s, 2H), 3.82-3.79 (m, 2H), 3.69-3.61 (m, 4H), 3.23-3.21 (m, 2H). MS: *m/z* 479.2 [M+H]⁺.

Example 133

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.66 (s, 1H), 7.65-7.58 (m, 1H), 7.47-7.41 (m, 1H), 7.38 (d, *J* = 5.7 Hz, 1H), 7.27 (t, *J* = 9.0 Hz, 1H), 7.05-6.98 (m, 2H), 5.29 (s, 2H), 3.59-2.89 (m, 9H), 1.79-1.45 (m, 8H). MS: *m/z* 497.3 [M+H]⁺.

Example 134

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.66-7.59 (m, 1H), 7.47-7.43 (m, 1H), 7.38 (d, *J* = 6.0 Hz, 1H), 7.28 (t, *J* = 8.9 Hz, 1H), 7.05-6.99 (m, 2H), 5.30 (s, 2H), 3.55-3.12 (m, 9H), 1.62-1.06 (m, 10H). MS: *m/z* 511.3 [M+H]⁺.

Example 135

5-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.65 (s, 1H), 8.11 (dd, *J* = 5.0 and 1.7 Hz, 1H), 7.70-7.50 (m, 2H), 7.48-7.37 (m, 2H), 7.27 (t, *J* = 9.0 Hz, 1H), 7.04-6.98 (m, 2H), 6.81 (d, *J* = 8.7 Hz, 1H), 6.66 (t, *J* = 5.7 Hz, 1H), 5.29 (s, 2H), 3.70-3.68 (m, 2H), 3.57-3.54 (m, 2H), 3.38-3.34 (m, 2H), 3.24-3.22 (m, 2H). MS: *m/z* 478.2 [M+H]⁺.

Example 136

5-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.64 (s, 1H), 7.77 (d, *J* = 4.8 Hz, 1H), 7.64-7.56 (m, 1H), 7.46-7.37 (m, 3H), 7.27 (t, *J* = 9.0 Hz, 1H), 7.13 (t, *J* = 4.8 Hz, 1H), 7.02-6.97 (m, 2H), 5.30 (s, 2H), 3.70-3.23 (m, 8H). MS: *m/z* 511.2 [M+H]⁺.

Example 137

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.67 (s, 1H), 7.60-7.27 (m, 9H), 7.05-6.95 (m, 2H), 5.30 (s, 2H), 3.66-3.29 (m, 8H). MS: *m/z* 505.3 [M+H]⁺.

Compound of Example 138 was prepared from 1-(3-carboxybenzyl)-8-fluoroquinazoline-2,4(1*H*,3*H*)-dione (prepared from 8-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those used for Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 138

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-8-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.90 (s, 1H), 7.91 (d, *J* = 7.2 Hz, 1H), 7.59-7.51 (m, 1H), 7.44-7.33 (m, 2H), 7.29-7.28 (m, 3H), 5.40 (s, 2H), 3.59-3.19 (m, 9H), 1.61-1.24 (m, 10H). MS: *m/z* 493.3 [M+H]⁺.

The following compounds were prepared from 6-fluoro-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using procedures similar to those of compounds of Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 139

6-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.86 (s, 1H), 8.38 (d, *J* = 4.5 Hz, 2H), 7.73 (dd, *J* = 8.1 and 3.0 Hz, 1H), 7.57 (td, *J* = 9.0 and 3.3 Hz, 1H), 7.36-7.27 (m, 5H), 6.67 (t, *J* = 4.7 Hz, 1H), 5.36 (s, 2H), 3.65-3.30 (m, 8H). MS: *m/z* 461.2 [M+H]⁺.

Example 140

6-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione
¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 8.13 (d, *J* = 3.3 Hz, 1H), 7.73 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.61-7.53 (m, 2H), 7.46-7.38 (m, 2H), 7.35-7.28 (m, 3H), 6.82 (d, *J* = 8.4 Hz, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.36 (s, 2H), 3.76-3.33 (m, 8H). MS: *m/z* 460.3 [M+H]⁺.

Example 141

6-Fluoro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 7.77-7.76 (m, 2H), 7.73 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.54-7.14 (m, 6H), 7.13 (t, *J* = 4.4 Hz, 1H), 5.35 (s, 2H), 3.66-3.30 (m, 8H). MS: *m/z* 493.2 [M+H]⁺.

Example 142

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 7.74 (dd, *J* = 8.1 and 3.0 Hz, 1H), 7.56 (m, 1H), 7.43-7.41 (m, 2H), 7.32-7.29 (m, 3H), 5.36 (s, 2H), 3.60-3.10 (m, 9H), 1.61-1.29 (m, 10H). MS: *m/z* 493.3 [M+H]⁺.

Example 143

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 7.84 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.52-7.43 (m, 5H), 7.34 (dd, *J* = 9.2 and 4.1 Hz, 1H), 5.46 (s, 2H), 3.72-3.11 (m, 9H), 1.88-1.66 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

The following compounds were prepared from 6-fluoro-1-(3-carboxy-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 5-(bromomethyl)-2-fluorobenzoate using procedures similar to those of compounds of Examples 1 and 2), and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 144

6-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.83 (s, 1H), 8.39 (d, *J* = 4.8 Hz, 2H), 7.74 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.60-7.54 (m, 1H), 7.41-7.29 (m, 4H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.32 (s, 2H), 3.81-3.23 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 145

6-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.85 (s, 1H), 8.12 (dd, *J* = 4.8 and 1.5 Hz, 1H), 7.72 (dd, *J* = 8.3 and 3.2 Hz, 1H), 7.60-7.53 (m, 2H), 7.47-7.40 (m, 2H), 7.32-7.26 (m, 2H), 6.83 (d, *J* = 8.7 Hz, 1H), 6.68 (dd, *J* = 6.9 and 5.1 Hz, 1H), 5.33 (s, 2H), 3.72-3.25 (m, 8H). MS: *m/z* 478.3 [M+H]⁺.

Example 146

6-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.83 (s, 1H), 7.77-7.70 (m, 2H), 7.57-7.51 (m, 1H), 7.41-7.37 (m, 3H), 7.31-7.25 (m, 2H), 7.12 (t, *J* = 4.2 Hz, 1H), 5.31 (s, 2H), 3.70-3.16 (m, 8H). MS: *m/z* 511.2 [M+H]⁺.

Example 147

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.83 (s, 1H), 7.72 (dd, *J* = 8.3 and 3.2 Hz, 1H), 7.57-7.50 (m, 1H), 7.45-7.41 (m, 1H), 7.36 (d, *J* = 6.3 Hz, 1H), 7.29-7.23 (m, 2H), 5.30 (s, 2H), 3.53-3.10 (m, 9H), 1.60-1.21 (m, 10H). MS: *m/z* 511.3 [M+H]⁺.

Example 148

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.83 (s, 1H), 7.73 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.55 (t, *J* = 8.4 Hz, 1H), 7.46-7.42 (m, 1H), 7.36 (d, *J* = 6.0 Hz, 1H), 7.30-7.24 (m, 2H), 5.31 (s, 2H), 3.56-3.12 (m, 9H), 1.60-1.56 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

The following compounds were prepared from 6-fluoro-1-(3-carboxy-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 6-fluoroquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)-4-fluorobenzoate using procedures similar to those of

compounds of Examples 1 and 2), and the corresponding substituted piperazine or piperidine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 149

6-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 8.38 (d, *J* = 4.5 Hz, 2H), 7.74 (dd, *J* = 8.3 and 3.2 Hz, 1H), 7.62-7.55 (m, 1H), 7.39-7.33 (m, 2H), 7.30-7.20 (m, 2H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.37 (s, 2H), 3.65-3.30 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 150

6-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 8.12 (d, *J* = 3.3 Hz, 1H), 7.74 (d, *J* = 8.4 Hz, 1H), 7.63-7.52 (m, 2H), 7.40-7.26 (m, 3H), 7.21 (d, *J* = 7.2 Hz, 1H), 6.77 (d, *J* = 8.4 Hz, 1H), 6.66 (t, *J* = 5.9 Hz, 1H), 5.37 (s, 2H), 3.51-3.32 (m, 8H). MS: *m/z* 478.3 [M+H]⁺.

Example 151

6-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.83 (s, 1H), 7.75-7.69 (m, 2H), 7.52-7.49 (m, 1H), 7.39-7.30 (m, 3H), 7.23-7.17 (m, 2H), 7.11 (t, *J* = 4.2 Hz, 1H), 5.33 (s, 2H), 3.70-3.13 (m, 8H). MS: *m/z* 511.2 [M+H]⁺.

Example 152

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 7.76 (dd, *J* = 8.1 and 3.0 Hz, 1H), 7.60-7.54 (m, 1H), 7.39-7.35 (m, 2H), 7.24 (dd, *J* = 9.2 and 4.1 Hz, 1H), 7.18 (d, *J* = 6.9 Hz, 1H), 5.36 (s, 2H), 3.53-3.10 (m, 9H), 1.70-1.25 (m, 10H). MS: *m/z* 511.3 [M+H]⁺.

Example 153

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 7.72 (dd, $J = 8.6$ and 3.9 Hz, 1H), 7.58-7.51 (m, 1H), 7.36-7.33 (m, 2H), 7.23 (dd, $J = 9.5$ and 3.8 Hz, 1H), 7.16 (d, $J = 7.2$ Hz, 1H), 5.33 (s, 2H), 3.31-2.63 (m, 9H), 1.73-1.43 (m, 8H). MS: m/z 479.3 $[\text{M}+\text{H}]^+$.

The following compounds were prepared from 5-fluoro-1-(3- carboxy-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 154

5-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline -
2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.68 (s, 1H), 8.38 (d, $J = 4.5$ Hz, 2H), 7.67-7.65 (m, 1H), 7.40-7.33 (m, 2H), 7.21 (d, $J = 7.2$ Hz, 1H), 7.08-7.00 (m, 2H), 6.66 (t, $J = 4.8$ Hz, 1H), 5.35 (s, 2H), 3.81-3.23 (m, 8H). MS: m/z 479.3 $[\text{M}+\text{H}]^+$.

Example 155

5-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline -2,4(1*H*,3*H*)-
dione

^1H NMR (DMSO- d_6): 11.66 (s, 1H), 8.10 (d, $J = 3.3$ Hz, 1H), 7.69-7.61 (m, 1H), 7.53 (t, $J = 6.9$ Hz, 1H), 7.40-7.31 (m, 2H), 7.21 (d, $J = 6.6$ Hz, 1H), 7.08-6.99 (m, 2H), 6.76 (d, $J = 8.7$ Hz, 1H), 6.65 (t, $J = 6.9$ Hz, 1H), 5.34 (s, 2H), 3.51-3.30 (m, 8H). MS: m/z 478.3 $[\text{M}+\text{H}]^+$.

Example 156

5-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-
2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.69 (s, 1H), 7.78 (d, $J = 4.5$ Hz, 1H), 7.64-7.60 (m, 1H), 7.38-7.35 (m, 3H), 7.22 (d, $J = 6.0$ Hz, 1H), 7.16-7.13 (m, 1H), 7.00-6.94 (m, 2H), 5.33 (s, 2H), 3.32-3.15 (m, 8H). MS: m/z 511.2 $[\text{M}+\text{H}]^+$.

Example 157

1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-
2,4(1*H*,3*H*)-dione

^1H NMR (DMSO- d_6): 11.70 (s, 1H), 7.68-7.61 (m, 1H), 7.39-7.33 (m, 2H), 7.21 (d, $J = 7.2$ Hz, 1H), 7.06-6.97 (m, 2H), 5.34 (s, 2H), 3.53-3.10 (m, 9H), 1.70-1.24 (m, 10H). MS: m/z 511.3 $[\text{M}+\text{H}]^+$.

Example 158

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 7.67-7.60 (m, 1H), 7.41-7.32 (m, 2H), 7.20 (d, *J* = 7.5 Hz, 1H), 7.06-6.97 (m, 2H), 5.33 (s, 2H), 3.60-3.15 (m, 9H), 1.60-1.55 (m, 8H). MS: *m/z* 497.3 [M+H]⁺.

The following compounds were prepared from 5-methyl-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 159

5-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 8.37 (d, *J* = 4.8 Hz, 2H), 7.49-7.47 (m, 3H), 7.42-7.36 (m, 2H), 7.14-7.08 (m, 2H), 6.66 (t, *J* = 5.0 Hz, 1H), 5.46 (s, 2H), 3.91-3.62 (m, 8H), 2.79 (s, 3H). MS: *m/z* 457.3 [M+H]⁺.

Example 160

1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-methylquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (CD₃OD): 7.49-7.36 (m, 5H), 7.14-7.08 (m, 2H), 5.46 (s, 2H), 3.82-3.10 (m, 9H), 2.80 (s, 3H), 1.88-1.56 (m, 8H). MS: *m/z* 475.3 [M+H]⁺.

The following compounds were prepared from 1-(3-carboxybenzyl)-7-chloroquinazoline-2,4(1*H*,3*H*)-dione and the corresponding substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 161

7-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.81 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 8.00 (d, *J* = 8.4 Hz, 1H), 7.40-7.27 (m, 6H), 6.65 (t, *J* = 4.7 Hz, 1H), 5.36 (s, 2H), 3.77-3.63 (m, 8H). MS: *m/z* 477.2 [M+H]⁺.

Example 162

7-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 8.02 (d, *J* = 9.0 Hz, 1H), 7.44-7.29 (m, 6H), 5.36 (s, 2H), 3.40-3.10 (m, 9H), 1.81-1.51 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

The following compounds were prepared from the corresponding substituted 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and substituted piperazine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 163

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.65 (s, 1H), 7.65-7.57 (m, 1H), 7.46-7.41 (m, 1H), 7.38-7.35 (m, 1H), 7.30-7.23 (m, 1H), 7.04-6.98 (m, 2H), 5.29 (s, 2H), 3.75-3.15 (m, 8H), 1.99-1.96 (m, 1H), 0.72-0.69 (m, 4H). MS: *m/z* 469.5 [M+H]⁺.

Example 164

6-Chloro-1-(3-(4-benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.85 (s, 1H), 7.91 (s, 1H), 7.68-7.62 (m, 1H), 7.44-7.35 (m, 7H), 7.28-7.22 (m, 2H), 5.29 (s, 2H), 3.64-3.21 (m, 8H). MS: *m/z* 521.5 [M+H]⁺.

Example 165

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.66 (s, 1H), 7.64-7.57 (m, 1H), 7.43-7.36 (m, 2H), 7.29-7.26 (m, 2H), 7.04-6.98 (m, 2H), 5.32 (s, 2H), 3.56-3.00 (m, 9H), 2.17-1.66 (m, 6H). MS: *m/z* 465.6 [M+H]⁺.

Example 166

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.65 (s, 1H), 7.65-7.57 (m, 1H), 7.46-7.40 (m, 1H), 7.36 (d, *J* = 5.4 Hz, 1H), 7.26 (t, *J* = 9.0 Hz, 1H), 7.04-6.99 (m, 2H), 5.28 (s, 2H), 3.55-3.09 (m, 9H), 2.18-1.67 (m, 6H). MS: *m/z* 483.3 [M+H]⁺.

Example 167

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.87 (s, 1H), 7.95 (d, *J* = 2.4 Hz, 1H), 7.69 (dd, *J* = 8.9 and 2.6 Hz, 1H), 7.42-7.30 (m, 2H), 7.22-7.15 (m, 2H), 5.33 (s, 2H), 3.55-3.09 (m, 9H), 2.14-1.72 (m, 6H). MS: *m/z* 499.3 [M+H]⁺.

Example 168

5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 7.55 (t, *J* = 8.3 Hz, 1H), 7.46-7.42 (m, 1H), 7.37 (d, *J* = 6.3 Hz, 1H), 7.30-7.24 (m, 2H), 7.18-7.13 (m, 1H), 5.31 (s, 2H), 3.56-3.11 (m, 9H), 2.10-1.73 (m, 6H). MS: *m/z* 499.3 [M+H]⁺.

Example 169

6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.88 (s, 1H), 7.94 (s, 1H), 7.70 (dd, *J* = 9.0 and 2.7 Hz, 1H), 7.46-7.42 (m, 1H), 7.36 (dd, *J* = 6.3 and 1.8 Hz, 1H), 7.31-7.21 (m, 2H), 5.30 (s, 2H), 3.56-3.11 (m, 9H), 2.50-1.73 (m, 6H). MS: *m/z* 499.3 [M+H]⁺.

Example 170

5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.69 (s, 1H), 7.57 (t, *J* = 8.3 Hz, 1H), 7.38-7.29 (m, 3H), 7.20 (d, *J* = 6.9 Hz, 1H), 7.15 (d, *J* = 8.7 Hz, 1H), 5.35 (s, 2H), 3.56-3.00 (m, 9H), 2.12-1.74 (m, 6H). MS: *m/z* 499.4 [M+H]⁺.

Example 171

5-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 7.79 (d, *J* = 5.1 Hz, 1H), 7.54 (t, *J* = 8.3 Hz, 1H), 7.39-7.36 (m, 3H), 7.23 (m, 2H), 7.14-7.11 (m, 2H), 5.35 (s, 2H), 3.66-3.10 (m, 8H). MS: *m/z* 527.2 [M+H]⁺.

Example 172

5-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 7.85 (s, 1H), 7.66-7.58 (m, 1H), 7.46-7.39 (m, 2H), 7.29 (t, *J* = 9.0 Hz, 1H), 7.04-6.98 (m, 3H), 6.64-6.63 (m, 1H), 5.31 (s, 2H), 3.73-3.70 (m, 4H), 3.55-3.54 (m, 2H), 3.23-3.22 (m, 2H). MS: *m/z* 495.3 [M+H]⁺.

Example 173

5-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 8.12 (dd, *J* = 4.8 and 1.2 Hz, 1H), 7.62-7.52 (m, 2H), 7.45-7.31 (m, 3H), 7.22 (d, *J* = 7.5 Hz, 1H), 7.15 (d, *J* = 8.4 Hz, 1H), 6.75 (d, *J* = 8.7 Hz, 1H), 6.68-6.64 (m, 1H), 5.36 (s, 2H), 3.69-3.21 (m, 8H). MS: *m/z* 494.3 [M+H]⁺.

Example 174

6-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 8.08 (d, *J* = 3.9 Hz, 1H), 7.92 (d, *J* = 2.4 Hz, 1H), 7.72 (dd, *J* = 9.0 and 2.4 Hz, 1H), 7.66-7.59 (m, 1H), 7.44-7.32 (m, 2H), 7.24 (d, *J* = 9.0 Hz, 1H), 7.20 (d, *J* = 7.8 Hz, 1H), 6.84 (d, *J* = 7.5 Hz, 1H), 6.70 (t, *J* = 5.4 Hz, 1H), 5.34 (s, 2H), 3.69-3.21 (m, 8H). MS: *m/z* 494.3 [M+H]⁺.

Example 175

6-Chloro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.83 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 7.92 (d, *J* = 2.4 Hz, 1H), 7.70 (d, *J* = 8.7 Hz, 1H), 7.24-7.18 (m, 4H), 6.64 (t, *J* = 4.8 Hz, 1H), 5.34 (s, 2H), 3.69-3.21 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

Example 176

5-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.63 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 7.66-7.61 (m, 1H), 7.29 (d, *J* = 9.3 Hz, 1H), 7.19-7.16 (m, 2H), 7.05-6.98 (m, 2H), 6.65 (t, *J* = 4.8 Hz, 1H), 5.33 (s, 2H), 3.80-3.20 (m, 8H). MS: *m/z* 479.3 [M+H]⁺.

Example 177

1-(3-(4-Cyclopentylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.02 (dd, *J* = 8.0 and 1.4 Hz, 1H), 7.63 (t, *J* = 7.1 Hz, 1H), 7.43-7.36 (m, 2H), 7.26-7.21 (m, 4H), 5.34 (s, 2H), 3.65-3.30 (m, 9H), 1.48-1.22 (m, 8H). MS: *m/z* 433.3 [M+H]⁺.

Example 178

1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 7.74 (dd, *J* = 8.1 and 3.0 Hz, 1H), 7.56 (t, *J* = 8.4 Hz, 1H), 7.45-7.39 (m, 2H), 7.31-7.25 (m, 2H), 5.32 (s, 2H), 3.54-3.16 (m, 8H), 1.92 (m, 1H), 0.73-0.70 (m, 4H). MS: *m/z* 469.3 [M+H]⁺.

Example 179

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 7.75 (dd, *J* = 8.3 and 2.9 Hz, 1H), 7.56 (t, *J* = 8.7 Hz, 1H), 7.43-7.40 (m, 2H), 7.31-7.26 (m, 3H), 5.35 (s, 2H), 3.52-3.31 (m, 9H), 2.18-1.75 (m, 6H). MS: *m/z* 465.3 [M+H]⁺.

Example 180

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 7.74 (d, *J* = 8.1 Hz, 1H), 7.59-7.53 (m, 1H), 7.46-7.44 (m, 1H), 7.37 (d, *J* = 5.7 Hz, 1H), 7.31-7.25 (m, 2H), 5.31 (s, 2H), 3.57-3.11 (m, 9H), 2.14-1.74 (m, 6H). MS: *m/z* 483.3 [M+H]⁺.

Example 181

1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 7.74 (dd, *J* = 8.1 and 2.7 Hz, 1H), 7.55-7.53 (m, 1H), 7.46-7.42 (m, 7H), 7.30-7.24 (m, 2H), 5.31 (s, 2H), 3.67-3.20 (m, 8H). MS: *m/z* 505.3 [M+H]⁺.

Example 182

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-
2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.68 (s, 1H), 7.66-7.59 (m, 1H), 7.36-7.33 (m, 2H), 7.18 (d, *J* = 7.2 Hz, 1H), 7.07-6.96 (m, 2H), 5.32 (s, 2H), 3.60-3.01 (m, 9H), 2.11-1.73 (m, 6H). MS: *m/z* 483.4 [M+H]⁺.

Example 183

1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-
2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 7.73 (dd, *J* = 8.1 and 2.7 Hz, 1H), 7.56 (t, *J* = 8.4 Hz, 1H), 7.36-7.30 (m, 2H), 7.24-7.14 (m, 2H), 5.34 (s, 2H), 3.60-3.01 (m, 9H), 2.14-1.72 (m, 6H). MS: *m/z* 483.3 [M+H]⁺.

Example 184

5-Fluoro-1-(6-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-
2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 7.86 (s, 1H), 7.67-7.59 (m, 1H), 7.39-7.36 (m, 2H), 7.23 (d, *J* = 7.2 Hz, 1H), 7.05-6.97 (m, 3H), 6.65-6.63 (m, 1H), 5.34 (s, 2H), 3.70-3.10 (m, 8H). MS: *m/z* 495.4 [M+H]⁺.

Example 185

6-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-
2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 7.84 (s, 1H), 7.74 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.59-7.53 (m, 1H), 7.45-7.39 (m, 2H), 7.32-7.26 (m, 2H), 7.02 (d, *J* = 3.3 Hz, 1H), 6.63 (dd, *J* = 3.3 Hz and 1.8 Hz, 1H), 5.32 (s, 2H), 3.67-3.23 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

Example 186

6-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.83 (s, 1H), 8.39 (d, *J* = 4.5 Hz, 2H), 7.74 (dd, *J* = 8.3 and 3.2 Hz, 1H), 7.60-7.54 (m, 1H), 7.33-7.29 (m, 2H), 7.27-7.18 (m, 2H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.36 (s, 2H), 3.80-3.35 (m, 8H). MS: *m/z* 479.4 [M+H]⁺.

Example 187

1-(6-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.86 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 7.76-7.69 (m, 1H), 7.65 (d, *J* = 8.1 Hz, 1H), 7.60-7.59 (m, 1H), 7.38 (dd, *J* = 8.3 and 1.7 Hz, 1H), 7.13-7.09 (m, 2H), 6.65 (t, *J* = 4.7 Hz, 1H), 5.32 (s, 2H), 3.73-3.03 (m, 8H). MS: *m/z* 495.3 [M+H]⁺.

Example 188

6-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.82 (s, 1H), 7.71 (dd, *J* = 8.4 and 3.0 Hz, 1H), 7.58-7.51 (m, 1H), 7.43-7.38 (m, 2H), 7.30-7.23 (m, 2H), 7.17 (d, *J* = 3.6 Hz, 1H), 6.88 (d, *J* = 3.6 Hz, 1H), 5.30 (s, 2H), 3.73-3.26 (m, 8H). MS: *m/z* 484.3 [M+H]⁺.

Example 189

5-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.65 (s, 1H), 7.65-7.57 (m, 1H), 7.43-7.38 (m, 2H), 7.27 (t, *J* = 9.0 Hz, 1H), 7.17 (d, *J* = 3.6 Hz, 1H), 7.04-6.98 (m, 2H), 6.87 (d, *J* = 3.6 Hz, 1H), 5.29 (s, 2H), 3.73-3.26 (m, 8H). MS: *m/z* 484.3 [M+H]⁺.

Example 190

1-(4-Fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 8.00 (d, *J* = 8.1 Hz, 1H), 7.64 (t, *J* = 7.9 Hz, 1H), 7.46-7.38 (m, 2H), 7.30-7.22 (m, 3H), 7.17 (d, *J* = 3.6 Hz, 1H), 6.87 (d, *J* = 3.6 Hz, 1H), 5.31 (s, 2H), 3.79-3.23 (m, 8H). MS: *m/z* 466.3 [M+H]⁺.

Example 191

1-(3-(4-(Cyclohexylcarbamoyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.67 (s, 1H), 8.01 (d, *J* = 7.8 Hz, 1H), 7.63 (t, *J* = 7.8 Hz, 1H), 7.40-7.37 (m, 2H), 7.30-7.22 (m, 4H), 6.20 (d, *J* = 4.0 Hz, 1H), 5.33 (s, 2H), 3.80-3.00 (m, 9H), 1.80-1.00 (m, 10H). MS: *m/z* 490.4 [M+H]⁺.

The following salts were prepared from the corresponding free base via stirring the free base in HCl/ethyl acetate for 2.5 h at room temperature, followed by evaporation of the solvents.

Example 192

1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 8.00 (d, *J* = 6.9 Hz, 1H), 7.64 (t, *J* = 7.8 Hz, 1H), 7.46-7.39 (m, 2H), 7.30-7.21 (m, 3H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.31 (s, 2H), 3.81-3.78 (m, 2H), 3.67-3.62 (m, 4H), 3.24-3.20 (m, 2H). MS: *m/z* 461.2 [M+H]⁺.

Example 193

5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride

¹H NMR (DMSO-*d*₆): 11.66 (s, 1H), 8.41 (d, *J* = 4.5 Hz, 2H), 7.67-7.59 (m, 1H), 7.47-7.39 (m, 2H), 7.28 (t, *J* = 4.5 Hz, 1H), 7.06-6.99 (m, 2H), 6.70 (t, *J* = 4.8 Hz, 1H), 5.31 (s, 2H), 3.82-3.81 (m, 2H), 3.69-3.64 (m, 4H), 3.25-3.22 (m, 2H). MS: *m/z* 479.3 [M+H]⁺.

Example 194

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride

¹H NMR (DMSO-*d*₆): 11.72 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 8.01 (d, *J* = 6.9 Hz, 1H), 7.64 (t, *J* = 6.9 Hz, 1H), 7.50-7.20 (m, 6H), 6.66 (t, *J* = 4.7 Hz, 1H), 5.35 (s, 2H), 4.10-3.20 (m, 8H). MS: *m/z* 443.3 [M+H]⁺.

Example 195

1-(3-((4-(Pyridin-2-yl)piperazin-1-yl)methyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

- a) 1-(3-(Hydroxymethyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: To the solution of 1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione (0.5 g, 1.6 mmol) in THF (30 mL) and ethanol (10 mL) was added NaBH₄ (1.2 g, 32 mmol) at room temperature under N₂. It was stirred at room temperature for 4 h, then refluxed overnight. The reaction mixture was quenched with HCl (2N), then concentrated. The residue was diluted with H₂O (10 mL), and it was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, and concentrated to afford the crude product, which was used for the next reaction without purification.
- b) 1-(3-(Methanesulfonylmethyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: To a solution of 1-(3-(hydroxymethyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (0.1 g, 0.35 mmol) and triethylamine (0.1 g, 1.1 mmol) in dry DCM (2 mL) was added MsCl dropwise at 0°C under N₂. The mixture was stirred for 2 h, and was used for the next reaction directly.

c) 1-(3-((4-(Pyridin-2-yl)piperazin-1-yl)methyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: A solution of 1-(pyridin-2-yl)piperazine (0.057 g, 0.35 mmol) in dry DCM (0.2 mL) was added to the above solution at room temperature under N₂, and the mixture was refluxed overnight. The reaction mixture was concentrated, and the residue was diluted with H₂O (2 mL). It was extracted with ethyl acetate (3 x 10 mL). The combined organic layers was washed with brine, dried with anhydrous Na₂SO₄, and concentrated. The crude product was purified by preparative TLC to give the title compound (5 mg) as white solid. ¹H NMR (DMSO-*d*₆): 8.09 (d, *J* = 3.3 Hz, 1H), 8.02 (d, *J* = 6.6 Hz, 1H), 7.66-7.61 (m, 1H), 7.54-7.48 (m, 1H), 7.29-7.15 (m, 6H), 6.78 (d, *J* = 8.7 Hz, 1H), 6.64-6.59 (m, 1H), 5.32 (s, 2H), 3.42-3.39 (m, 4H), 2.40-2.37 (m, 4H). MS: *m/z* 428.3 [M+H]⁺.

The following compounds were prepared from 1-(3-aminobenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and 1-(bromomethyl)-3-nitrobenzene followed by reduction) and the corresponding substituted acetic acid using a procedure similar to those described for the synthesis of compound of Example 3.

Example 196

1-(3-(Naphthalen-2-yl)acetamidobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.72 (brs, 1H), 10.23 (s, 1H), 8.10-7.99 (m, 1H), 7.90-7.76 (m, 4H), 7.62-7.56 (m, 2H), 7.46-7.38 (m, 4H), 7.26-7.16 (m, 3H), 7.05-6.90 (m, 1H), 5.25 (s, 2H), 3.75 (s, 2H). MS: *m/z* 436.4 [M+H]⁺.

Example 197

1-(3-(3,4-Dimethoxyphenyl)acetamidobenzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (brs, 1H), 10.03 (s, 1H), 8.01 (dd, *J* = 7.8 and 1.2 Hz, 1H), 7.65-7.55 (m, 2H), 7.34 (s, 1H), 7.27-7.17 (m, 3H), 6.98 (d, *J* = 7.8 Hz, 1H), 6.83-6.77 (m, 3H), 5.26 (s, 2H), 3.69 (s, 6H), 3.47 (s, 2H). MS: *m/z* 446.3 [M+H]⁺.

Example 198

1-(3-([1,2,4]Triazolo[4,3-*a*]pyridine-6-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

a) 1-(3-(*N*-Methoxy-*N*-methylcarbamoyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: To a solution of 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (0.5 g, 1.68 mmol), HATU (0.96 g, 2.53 mmol) and DIEA (0.9 g, 7 mmol) in dry DCM (20 mL) was added *N*,*O*-dimethylhydroxylamine hydrochloride (0.2 g, 2.0 mmol) at room temperature under N₂, then the mixture was heated to 4°C and stirred overnight. The reaction solution was

concentrated, the residue was diluted with H₂O and it was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with water and brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified via chromatography to give the title compound (0.4 g, 70% yield).

- b) 1-(3-([1,2,4]triazolo[4,3-*a*]pyridine-6-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: To a solution of 6-bromo-[1,2,4]triazolo[4,3-*a*]pyridine (35 mg, 0.18 mmol) in anhydrous THF (1 mL) was added *n*-BuLi (0.15 mL, 0.38 mmol) dropwise at -65°C under N₂, then it was maintained at this temperature for 3 h. To this solution was added dropwise a solution of 1-(3-(*N*-methoxy-*N*-methylcarbamoyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (50 mg, 0.15 mmol) in THF (1 mL), then it was stirred for 4 h at -65 °C. It was allowed to warm to 0 °C for over 4 h. The reaction solution was poured into saturated aqueous NH₄Cl, extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, and concentrated. The crude product was purified by preparative TLC to give the title compound (6.5 mg, 11% yield) as white solid. ¹H NMR (CD₃OD+CDCl₃): 9.42 (d, *J* = 7.2 Hz, 1H), 8.41-8.39 (m, 2H), 8.10 (d, *J* = 7.2 Hz, 1H), 7.95 (d, *J* = 9.0 Hz, 1H), 7.58-7.45 (m, 4H), 7.18-7.11 (m, 3H), 5.38 (s, 2H). MS: *m/z* 398.2 [M+H]⁺.

The following compounds were prepared from the corresponding substituted 1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione or 1-((2-carboxylpyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione and substituted amine using a procedure similar to those described for the synthesis of compound of Example 3.

Example 199

6-Fluoro-1-((2-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.84 (s, 1H), 8.39 (d, *J* = 4.8 Hz, 2H), 7.94 (t, *J* = 7.8 Hz, 1H), 7.72-7.70 (m, 1H), 7.57-7.53 (m, 3H), 7.34-7.30 (m, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.46 (s, 2H), 3.73-3.63 (m, 4H), 3.42-3.32 (m, 4H). MS: *m/z* 462.5 [M+H]⁺.

Example 200

5-Fluoro-1-(4-fluoro-3-(4-methoxybenzylcarbamoyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 8.79 (t, *J* = 5.3 Hz, 1H), 7.66-7.57 (m, 2H), 7.46-7.41 (m, 1H), 7.27-7.20 (m, 3H), 7.08-6.99 (m, 2H), 6.90-6.87 (m, 2H), 5.30 (s, 2H), 4.37 (d, *J* = 6.0 Hz, 2H), 3.73 (s, 3H). MS: *m/z* 452.2 [M+H]⁺.

Example 201

1-(3-(3-Chlorobenzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 9.01 (t, *J* = 5.1 Hz, 1H), 7.75-7.68 (m, 2H), 7.57-7.52 (m, 1H), 7.48-7.32 (m, 5H), 7.18-7.08 (m, 2H), 5.31 (s, 2H), 4.45 (d, *J* = 5.7 Hz, 2H). MS: *m/z* 456.1 [M+H]⁺.

Example 202

1-(3-(Benzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.70 (s, 1H), 8.87 (t, *J* = 5.1 Hz, 1H), 7.67-7.59 (m, 2H), 7.47-7.42 (m, 1H), 7.36-7.22 (m, 6H), 7.09-6.99 (m, 2H), 5.31 (s, 2H), 4.45 (d, *J* = 6.0 Hz, 1H). MS: *m/z* 422.3 [M+H]⁺.

Example 203

5-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.32 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 7.53 (t, *J* = 8.4 Hz, 1H), 7.44-7.29 (m, 4H), 6.85 (d, *J* = 8.4 Hz, 1H), 6.75 (d, *J* = 8.4 Hz, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.33 (s, 2H), 3.84 (s, 3H), 3.79-3.42 (m, 8H). MS: *m/z* 473.3 [M+H]⁺.

Example 204

6-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.71 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 7.47-7.40 (m, 3H), 7.34-7.26 (m, 3H), 7.20 (d, *J* = 9.3 Hz, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.35 (s, 2H), 3.80 (s, 3H), 3.63-3.46 (m, 8H). MS: *m/z* 473.3 [M+H]⁺.

Example 205

1-(3-(5-Bromopyrimidin-2-aminocarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-*d*₆): 11.73 (s, 1H), 10.27 (s, 1H), 9.43 (s, 1H), 8.50 (s, 2H), 7.68-7.52 (m, 3H), 7.28 (t, *J* = 9.3 Hz, 1H), 7.11-7.00 (m, 2H), 5.33 (s, 2H). MS: *m/z* 503.0 [M+H]⁺.

Example 206

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-7-(trifluoromethyl)quinazoline-2,4(1*H*,3*H*)-dione

^1H NMR (300 M, DMSO- d_6): δ 11.96 (s, 1H), 8.36 (d, $J = 4.8$ Hz, 2H), 8.19 (d, $J = 8.1$ Hz, 1H), 7.55 (d, $J = 8.4$ Hz, 1H), 7.47 (s, 1H), 7.43-7.39 (m, 3H), 7.33-7.30 (m, 1H), 6.65 (t, $J = 4.7$ Hz, 1H), 5.44 (s, 2H), 3.59-3.46 (m, 8H). MS: m/z 511.2 $[\text{M}+\text{H}]^+$.

Example 207

6,7-Ethylenedio-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

^1H NMR (DMSO- d_6): 11.55 (s, 1H), 8.39 (d, $J = 4.8$ Hz, 2H), 7.44-7.34 (m, 5H), 6.75 (s, 1H), 6.67 (t, $J = 4.8$ Hz, 1H), 5.29 (s, 2H), 4.32-4.27 (m, 4H), 3.88-3.56 (m, 8H). MS: m/z 501.2 $[\text{M}+\text{H}]^+$.

Example 208

5-Fluoro-1-(6-methoxy-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

^1H NMR (DMSO- d_6): 11.65 (s, 1H), 8.37 (d, $J = 4.8$ Hz, 2H), 7.67-7.60 (m, 1H), 7.39 (dd, $J = 8.4$ and 1.8 Hz, 1H), 7.16 (d, $J = 8.4$ Hz, 1H), 7.07-6.98 (m, 2H), 6.87 (d, $J = 8.4$ Hz, 1H), 6.65 (t, $J = 4.7$ Hz, 1H), 5.20 (s, 2H), 3.95 (s, 3H), 3.82-3.16 (m, 8H). MS: m/z 491.2 $[\text{M}+\text{H}]^+$.

Example 209

7-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

^1H NMR (DMSO- d_6): 11.54 (s, 1H), 8.37 (d, $J = 4.8$ Hz, 2H), 7.92 (d, $J = 8.7$ Hz, 1H), 7.42-7.30 (m, 4H), 6.84 (dd, $J = 8.9$ and 2.3 Hz, 1H), 6.68-6.63 (m, 2H), 5.34 (s, 2H), 3.76 (s, 3H), 3.71-3.51 (m, 8H). MS: m/z 473.3 $[\text{M}+\text{H}]^+$.

Example 210

5-Fluoro-1-(4-fluoro-3-(4-(tetrahydrofuran-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

^1H NMR (DMSO- d_6): 11.67 (s, 1H), 7.66-7.59 (m, 1H), 7.47-7.42 (m, 1H), 7.38 (d, $J = 6.0$ Hz, 1H), 7.28 (t, $J = 9.0$ Hz, 1H), 7.10-6.95 (m, 2H), 5.30 (s, 2H), 4.75-4.50 (m, 1H), 3.85-3.45 (m, 8H), 3.25-3.00 (m, 2H), 2.15-1.70 (m, 4H). MS: m/z 499.3 $[\text{M}+\text{H}]^+$.

Example 211

5-Fluoro-1-(4-nitro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1H,3H)-dione

¹H NMR (DMSO-d₆): 11.71 (s, 1H), 8.40 (d, *J* = 3.9 Hz, 2H), 8.17 (d, *J* = 8.7 Hz, 1H), 7.63-7.57 (m, 3H), 7.07-6.94 (m, 2H), 6.69 (t, *J* = 4.4 Hz, 1H), 5.45 (s, 2H), 3.85-3.84 (m, 2H), 3.70-3.69 (m, 2H), 3.61-3.59 (m, 2H), 3.21-3.18 (m, 2H). MS: *m/z* 506.2 [M+H]⁺.

Example 212

1-(3-(4-Cyclohexylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.68 (s, 1H), 7.66-7.58 (m, 1H), 7.45-7.40 (m, 1H), 7.30-7.23 (m, 2H), 7.06-6.99 (m, 2H), 5.30 (s, 2H), 3.56-3.07 (m, 8H), 2.33-2.25 (m, 1H), 1.72-1.04 (m, 10H). MS: *m/z* 483.5 [M+H]⁺.

Example 213

5-Fluoro-1-(4-fluoro-3-(4-phenylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.68 (s, 1H), 7.67-7.60 (m, 1H), 7.48-7.38 (m, 2H), 7.32-7.20 (m, 3H), 7.07-7.00 (m, 2H), 6.94-6.91 (m, 2H), 6.82 (t, *J* = 7.2 Hz, 1H), 5.31 (s, 2H), 3.85-3.65 (m, 2H), 3.45-3.10 (m, 4H), 3.10-2.90 (m, 2H). MS: *m/z* 477.4 [M+H]⁺.

Example 214

5-Fluoro-1-(4-fluoro-3-(4-phenylpiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.68 (s, 1H), 7.62-7.56 (m, 1H), 7.45-7.37 (m, 2H), 7.33-7.18 (m, 6H), 7.04-6.93 (m, 2H), 5.45-5.15 (m, 2H), 4.70-4.55 (m, 1H), 3.50-3.05 (m, 2H), 2.95-2.65 (m, 2H), 1.80-1.20 (m, 4H). MS: *m/z* 476.4 [M+H]⁺.

Example 215

1-(4-Bromo-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.67 (s, 1H), 8.39 (d, *J* = 4.8 Hz, 2H), 7.66-7.58 (m, 2H), 7.37 (d, *J* = 1.8 Hz, 1H), 7.30 (dd, *J* = 8.4 and 2.1 Hz, 1H), 7.06-6.97 (m, 2H), 6.68 (t, *J* = 4.8 Hz, 1H), 5.37-5.22 (m, 2H), 3.82-3.11 (m, 8H). MS: *m/z* 539.3 [M+H]⁺.

Example 216

6,7-Methylenedioxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.64 (s, 1H), 8.39 (d, *J* = 4.5 Hz, 2H), 7.44-7.32 (m, 5H), 6.95 (s, 1H), 6.67 (t, *J* = 4.4 Hz, 1H), 6.14 (s, 2H), 5.33 (s, 2H), 3.81-3.65 (m, 8H). MS: *m/z* 487.2 [M+H]⁺.

Example 217

1-(3-(4-(Cyclohexylmethyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.70 (s, 1H), 7.66-7.58 (m, 1H), 7.44-7.40 (m, 1H), 7.32-7.23 (m, 2H), 7.05-6.99 (m, 2H), 5.30 (s, 2H), 3.70-3.50 (m, 2H), 3.20-3.00 (m, 2H), 2.38-2.28 (m, 2H), 2.20-2.10 (m, 2H), 2.05 (d, *J* = 6.6 Hz, 2H), 1.73-1.63 (m, 5H), 1.45-1.39 (m, 1H), 1.30-1.00 (m, 3H), 0.90-0.70 (m, 2H). MS: *m/z* 497.3 [M+H]⁺.

Example 218

8-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

¹H NMR (DMSO-d₆): 11.93 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 7.89 (d, *J* = 7.8 Hz, 1H), 7.60-7.52 (m, 1H), 7.45-7.22 (m, 5H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.40 (s, 2H), 3.90-3.20 (m, 8H). MS: *m/z* 461.2 [M+H]⁺.

Example 219

6-Amino-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

A mixture of 6-nitro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-quinazoline-2,4(1*H*,3*H*)-dione (50 mg, 0.10 mmol) in EtOH (10 mL) with Fe powder (33.5 mg, 0.60 mmol) and sat. NH₄Cl solution (0.2 mL) were refluxed under N₂ overnight. The mixture was filtered to remove the solid. The solution was concentrated and purified by flash chromatography (DCM:MeOH = 50:3) to give the title compound (10.79 mg, 23.6% yield) as yellow solid. ¹H NMR (DMSO-d₆): 11.46 (s, 1H), 8.39 (d, *J* = 4.8 Hz, 2H), 7.42-7.31 (m, 4H), 7.20 (d, *J* = 2.7 Hz, 1H), 7.00 (d, *J* = 8.7 Hz, 1H), 6.89 (dd, *J* = 8.7 and 2.7 Hz, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.26 (s, 4H), 3.81-3.68 (m, 8H). MS: *m/z* 458.3 [M+H]⁺.

Example 220

1-(2-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

The title compound was prepared from 1-(2-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 2-(bromomethyl)benzoate using a procedure similar to those used for compounds of Example 1 and 2), and 2-(piperazin-1-yl)pyrimidine using a procedure similar to that described for the synthesis of compound of

Example 3. ^1H NMR (DMSO- d_6): 11.76 (s, 1H), 8.40 (d, $J = 4.8$ Hz, 2H), 8.03-8.00 (m, 1H), 7.70-7.50 (m, 1H), 7.35-7.00 (m, 6H), 6.68 (t, $J = 4.8$ Hz, 1H), 5.25 (s, 2H), 4.10-3.60 (m, 8H). MS: m/z 443.2 $[\text{M}+\text{H}]^+$.

Example 221

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)phenyl)quinazoline-2,4(1*H*,3*H*)-dione

- a) Methyl 3-(3-(2-bromobenzoyl)ureido)benzoate: To a solution of 2-bromobenzamide (2.0 g, 10.0 mmol) in DCM (30 mL) was added oxalyl chloride (1.65 g, 13.0 mmol) and it was heated under reflux for 23 h. To the reaction mixture was added methyl 3-aminobenzoate (1.66 g, 11.0 mmol) and it was stirred for 20 min. The mixture was poured into water, and the precipitate was filtered. It was recrystallized from ethanol to give the crude product (1.80g, 48%), which was used for the next step without further purification.
- b) 1-(3-Methoxycarbonylphenyl)quinazoline-2,4(1*H*,3*H*)-dione: A mixture of methyl 3-(3-(2-bromobenzoyl)ureido)benzoate (1.8 g, 4.8 mmol), potassium *tert*-butoxide (2.69 g, 24 mmol) and DMF (20 mL) was stirred at 70-80°C under nitrogen for 2.25 h. It was cooled, diluted with water, and the mixture was filtered. The crude product was purified by preparative TLC (EtOAc) to give the title compound (0.34 g, 24% yield) as white solid. MS: m/z 297 $[\text{M}+1]^+$.
- c) 1-(3-Carboxyphenyl)quinazoline-2,4(1*H*,3*H*)-dione: The title compound was prepared from 1-(3-methoxycarbonylphenyl)quinazoline-2,4(1*H*,3*H*)-dione and NaOH using a procedure similar to those of compound of Example 2, and was isolated as white solid. MS: m/z 283 $[\text{M}+\text{H}]^+$.
- d) 1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)phenyl)quinazoline-2,4(1*H*,3*H*)-dione: The title compound was prepared from 1-(3-carboxyphenyl)quinazoline-2,4(1*H*,3*H*)-dione and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those of compound of Example 3, and was isolated as white solid. ^1H NMR (DMSO- d_6): 11.72 (s, 1H), 8.36 (d, $J = 4.8$ Hz, 2H), 8.02 (d, $J = 6.6$ Hz, 1H), 7.69-7.55 (m, 5H), 7.25 (t, $J = 7.5$ Hz, 1H), 6.65 (t, $J = 4.7$ Hz, 1H), 6.48 (d, $J = 8.4$ Hz, 1H), 3.80-3.20 (m, 8H). MS: m/z 429.1 $[\text{M}+\text{H}]^+$.

Example 222

8-Chloro-1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione and 8-chloro-3-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione

To a suspension of 8-chloroquinazoline-2,4(1*H*,3*H*)-dione (5.1 g, 25.9 mmol) in toluene (50 mL) and hexamethyldisilazane (HMDS, 10.4 g, 64.9 mmol) was added sulfuric acid (0.25 g, 2.6 mmol). The mixture was heated to reflux and refluxed for 8 h until a clear solution was

obtained. The solvent and excessive HMDS was removed via vacuum evaporation, and methyl 3-(bromomethyl)benzoate (5.9 g, 25.9 mmol) and DMF (1 mL) was added to the residue. The reaction mixture was heated to 160°C and was stirred at this temperature for 3 h. It was diluted with 1,4-dioxane (6 mL) at 100°C, and then methanol (10 mL) was added at 70°C and it was stirred for 30 min. The mixture was cooled to 5°C and the precipitates were collected. The solids was washed with methanol (10 mL) and water (20 mL), dried under vacuum for 12 h to give the title compound 8-chloro-3-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione (2.6 g, 29.1% yield) as gray solid. MS: *m/z* 477.1 [M+H]⁺. The filtrate was concentrated, and the residue was purified by preparative TLC (PE:EA = 1:1) to give the other title compound 8-chloro-1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione (0.2 g, 1.8%) as yellow solid. MS: *m/z* 477.1 [M+H]⁺.

Example 223

8-Chloro-3-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

The compound was prepared from 8-chloro-3-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 8-chloro-3-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione and NaOH using a procedur similar to those of compound of Example 2), and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis of compound of Example 3. ¹H NMR (DMSO-*d*₆): 11.15 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 7.95 (d, *J* = 7.8 Hz, 1H), 7.82 (d, *J* = 7.5 Hz, 1H), 7.50-7.30 (m, 4H), 7.23 (t, *J* = 7.8 Hz, 1H), 6.67 (t, *J* = 4.8 Hz, 1H), 5.14 (s, 2H), 3.90-3.55 (m, 8H). MS: *m/z* 477.1 [M+H]⁺.

Example 224

8-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

The compound was prepared from 8-chloro-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 8-chloro-1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione and NaOH using a procedure similar to those of compound of Examples 2), and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis of compound of Example 3. ¹H NMR (DMSO-*d*₆): 11.94 (s, 1H), 8.38 (d, *J* = 4.8 Hz, 2H), 8.05 (d, *J* = 6.6 Hz, 1H), 7.77 (d, *J* = 6.6 Hz, 1H), 7.50-7.20 (m, 5H), 6.66 (t, *J* = 4.8 Hz, 1H), 5.53 (s, 2H), 3.90-3.50 (m, 8H). MS: *m/z* 477.1 [M+H]⁺.

Example 225

8-Methyl-3-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

The compound was prepared from 8-methyl-3-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from 8-methylquinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(bromomethyl)benzoate using a procedure similar to those used for Examples 222 and 223), and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis of compound of Example 3. ¹H NMR (DMSO-*d*₆): 10.81 (s, 1H), 8.36 (d, *J* = 4.5 Hz, 2H), 7.83 (d, *J* = 7.5 Hz, 1H), 7.52 (d, *J* = 7.2 Hz, 1H), 7.45-7.30 (m, 4H), 7.13 (t, *J* = 7.7 Hz, 1H), 6.67 (t, *J* = 4.7 Hz, 1H), 5.15 (s, 2H), 3.92-3.46 (m, 8H), 2.36 (s, 3H). MS: *m/z* 457.3 [M+H]⁺.

Example 226

8-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

- a) 8-Methyl-1*H*-benzo[*d*][1,3]oxazine-2,4-dione: To a solution of 2-amino-3-methylbenzoic acid (5.03 g, 33.3 mmol) in THF (50 mL) was added triphosgene (9.92 g, 33.4 mmol). The mixture was stirred at room temperature for 15 min and then filtered. The precipitate was washed by THF and water, dried to give the title compound (5.10 g, 86.4% yield) as white solid. MS: *m/z* 178.1 [M+H]⁺.
- b) 2-Amino-*N*-tert-butyl-3-methylbenzamide: To a solution of 8-methyl-1*H*-benzo[*d*][1,3]oxazine-2,4-dione (3.89 g, 22.0 mmol) in DMF (30 mL) were added DMAP (0.27 g, 2.2 mmol) and tert-butylamine (2.55 mL, 24.1 mmol). It was stirred at room temperature overnight, then the mixture was poured into water and the precipitate was filtered. The solid was washed with water and dried to give the product (1.68g, 37.1% yield). MS: *m/z* 207.2 [M+H]⁺.
- c) 3-Tert-butyl-8-methylquinazoline-2,4(1*H*,3*H*)-dione: To a solution of 2-amino-*N*-tert-butyl-3-methylbenzamide (0.52 g, 2.5 mmol) in THF (20 mL) were added CDI (0.85 g, 5.2 mmol). It was refluxed overnight, cooled to room temperature, then the mixture was poured into water and the precipitate was filtered. The solid was washed with water and dried to give the title compound (0.25 g, 42.6% yield). ¹H NMR (DMSO-*d*₆): 10.33 (s, 1H), 7.68 (d, *J* = 7.8 Hz, 1H), 7.41 (d, *J* = 7.2 Hz, 1H), 7.04 (t, *J* = 7.6 Hz, 1H), 2.31 (s, 3H), 1.66 (s, 9H).
- d) 3-Tert-butyl-8-methyl-1-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione: A mixture of 3-tert-butyl-8-methylquinazoline-2,4(1*H*,3*H*)-dione (1.00 g, 4.3 mmol), methyl 3-(bromomethyl)benzoate (0.99 g, 4.3 mmol) and MeONa (0.28 g, 5.2 mmol) in DMF (15 mL) was stirred at 50°C overnight. The resulting mixture was used directly for next step reaction. MS: *m/z* 403.2 [M+Na]⁺.
- e) 8-Methyl-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione: To the reaction mixture from above was added 6N HCl (a.q., 40 mL), and the mixture was heated at 80°C for 6 h. It was cooled to room temperature, poured into water and the precipitate was filtered. The solid

was washed with water and dried to give the title compound (0.27 g, 20.1% yield). MS: m/z 311.1 $[M+H]^+$.

- f) 8-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: The title compound was prepared from 8-methyl-1-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis of compound of Example 3. 1H NMR (DMSO- d_6): 11.70 (s, 1H), 8.38 (d, $J = 4.8$ Hz, 2H), 7.92 (d, $J = 6.9$ Hz, 1H), 7.50-7.26 (m, 5H), 7.17 (t, $J = 7.5$ Hz, 1H), 6.65 (t, $J = 4.8$ Hz, 1H), 5.35 (s, 2H), 3.85-3.15 (m, 8H), 2.35 (s, 3H). MS: m/z 457.2 $[M+H]^+$.

Example 227

8-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

The compound was prepared from 2-amino-3-methoxybenzoic acid using procedures similar to those described for the synthesis of compound of Example 226. 1H NMR (DMSO- d_6): 11.76 (s, 1H), 8.38 (d, $J = 4.8$ Hz, 2H), 7.65 (d, $J = 6.6$ Hz, 1H), 7.45-7.17 (m, 6H), 6.67 (t, $J = 4.7$ Hz, 1H), 5.51 (s, 2H), 3.80-3.64 (m, 8H), 3.56 (s, 3H). MS: m/z 473.2 $[M+H]^+$.

Example 228

3-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

- a) 3-(3-Methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione: A suspension of quinazoline-2,4(1*H*,3*H*)-dione (2.5 g, 15.4 mmol), methyl 3-(bromomethyl)benzoate (3.56 g, 15.4 mmol) and K_2CO_3 (4.26 g, 30.8 mmol) in DMF (30 mL) was stirred at room temperature overnight. To the mixture was added 150 mL water and the precipitates were collected. The crude product was purified by chromatography on silica (CH_2Cl_2 :MeOH=200:1) to give the title compound (0.87 g, 18.2% yield) as white solid. 1H NMR (DMSO- d_6): 11.51 (s, 1H), 8.00-7.90 (m, 2H), 7.84 (d, $J = 7.5$ Hz, 1H), 7.75-7.55 (m, 2H), 7.47 (t, $J = 7.8$ Hz, 1H), 7.30-7.15 (m, 2H), 5.14 (s, 2H), 3.83 (s, 3H). MS: m/z 311.2 $[M+H]^+$.
- b) 3-(3-Carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione: The title compound was prepared from 3-(3-methoxycarbonylbenzyl)quinazoline-2,4(1*H*,3*H*)-dione using a procedure similar to those described for the synthesis of compound of Example 2. MS: m/z 297.2 $[M+H]^+$.
- c) 3-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione: The title compound was prepared from 3-(3-carboxybenzyl)quinazoline-2,4(1*H*,3*H*)-dione and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis

of compound of Example 3. ¹H NMR (DMSO-*d*₆): 11.55 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 7.93 (d, *J* = 6.9 Hz, 1H), 7.66 (t, *J* = 7.1 Hz, 1H), 7.45-7.26 (m, 4H), 7.25-7.12 (m, 2H), 6.65 (t, *J* = 4.7 Hz, 1H), 5.11 (s, 2H), 3.90-3.20 (m, 8H). MS: *m/z* 443.3 [M+H]⁺.

Example 229

1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)phenethyl)quinazoline-2,4(1*H*,3*H*)-dione

The compound was prepared from 1-(3-carboxyphenethyl)quinazoline-2,4(1*H*,3*H*)-dione (prepared from quinazoline-2,4(1*H*,3*H*)-dione and methyl 3-(2-bromoethyl)benzoate using a procedure similar to those used for Examples 1 and 2) and 2-(piperazin-1-yl)pyrimidine using a procedure similar to those described for the synthesis of compound of Example 3. ¹H NMR (DMSO-*d*₆): 11.54 (s, 1H), 8.37 (d, *J* = 4.8 Hz, 2H), 7.78 (dd, *J* = 7.8 and 1.5 Hz, 1H), 7.73 (t, *J* = 6.9 Hz, 1H), 7.52 (d, *J* = 7.2 Hz, 1H), 7.35-7.22 (m, 5H), 6.66 (t, *J* = 4.7 Hz, 1H), 4.28 (t, *J* = 7.5 Hz, 2H), 3.80-3.20 (m, 8H), 2.95 (t, *J* = 7.4 Hz, 1H). MS: *m/z* 457.4 [M+H]⁺.

Example 230

8-Hydroxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione

A solution of BBr₃ (4N in DCM, 10 mL, 40 mmol) was added dropwise to a mixture of 8-methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (60 mg, 0.13 mmol) in DCM (25 mL) at -70°C. The mixture was warmed to room temperature and stirred at r.t overnight. After cooled at ice-bath, 10 mL MeOH was added dropwise and then NH₄Cl (a.q.) was added. The mixture was extracted by DCM (50 mL x 2), the combined organic layer was dried to give crude product which was purified by TLC (DCM/MeOH = 20:1) to give the target product (7.64 mg, 13.1% yield). ¹H NMR (DMSO-*d*₆): 11.64 (s, 1H), 10.34 (s, 1H), 8.36 (d, *J* = 4.8 Hz, 2H), 7.51 (dd, *J* = 6.8 and 2.6 Hz, 1H), 7.39-7.34 (m, 1H), 7.27-7.23 (m, 3H), 7.10-7.03 (m, 2H), 6.65 (t, *J* = 4.7 Hz, 1H), 5.63 (s, 2H), 3.85-3.53 (m, 8H). MS: *m/z* 459.4 [M+H]⁺.

Example 231

Evaluation of the synergistic potentiation effects of 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione and analogs on the growth inhibiting activity of MMS using a MTT based cell viability assay

The synergistic potentiation effect of PARP inhibitors on the growth inhibiting activity of DNA damaging anticancer drugs such as methyl methanesulphonate (MMS) were measured with SW620 human colorectal cancer cells using a MTT based cell viability assay.

Briefly, SW620 human colorectal cancer cells were used to measure the cell growth inhibiting activity of DNA alkylating anticancer drug MMS. The MMS concentration used in the assay was carefully titrated to a level at which MMS only had a minimal inhibitory effect on the cell growth, and under this conditions it was sensitive to detect the potentiation effect of 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones on MMS's inhibiting activity on cell growth. SW620 cells were grown and maintained in RPMI1640 (Gibco) medium supplemented with 10% FBS (Hyclone). In the first day of experiment, 4000 cells were seeded to each well of a 96-well cell culture plate and incubated at 37°C and 5% CO₂ in a cell culture incubator overnight. In the next day, the cell culture medium was removed. One hundred and eighty µl fresh medium containing 1.5 µg/ml MMS and 20 µl of 10-fold concentration of testing compounds or reference compounds (AZD2281 and ABT-888) were added to each well sequentially. The serial dilutions of the reference compounds and the compounds to be tested were made with a 1:3 and 1:10 fashion in DMSO. The 10-fold concentrated solutions of these compounds were made by mixing 10µl of the serial dilutions in DMSO with 90 µl fresh medium. The final concentration of DMSO in the medium was 1%. The cells were incubated at 37°C in 5% CO₂ cell culture incubator for additional 5 days (120 hours). Afterwards, the 96-well cell culture plate was taken out and 20 µl of MTT solution was added to each well and incubated at 37°C for 4 h. The medium was removed and 100 µl DMSO was added to each well. The 96-well cell culture plates were shaken adequately for 10 min and readed in a Varioskan Flash plate reader (Thermo Fisher Scientific) at 520 and 690 nm. The data were analyzed by Prism 5 software (GraphPad). The obtained 520 nm absorbance readings subtracting corresponding 690 nm reading were analyzed and plotted against the Log scale of the compound concentrations. The curves were fitted using the following equation to calculate the IC₅₀ value of each compound, $Y (\text{absorbance}) = \text{minimal absorbance value} + (\text{maximal absorbance value} - \text{minimal absorbance value}) / (1 + 10^{-(\text{LogC} - \text{LogIC}_{50})})$, C was the compound concentration.

The calculated IC₅₀ values are listed in Table 1. These values reflect the potentiation effect of 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones on the growth inhibiting activity of a fixed concentration of MMS in a MTT based cell viability assays.

Table 1. Substituted 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones potentiate the growth inhibiting activity of MMS (IC₅₀)

Example #	1	2	3	4	5	6	7	8
IC ₅₀ (nM)	>10000	>10000	90	12	5356	7026	5269	112

Example #	9	10	11	12	13	14	15	16
IC ₅₀ (nM)	500	439	618	533	727	102	666	7259
Example #	17	18	19	20	21	22	23	24
IC ₅₀ (nM)	>10000	>10000	>10000	391	>10000	>10000	8.2	26
Example #	25	26	27	28	29	30	31	32
IC ₅₀ (nM)	9.8	149	83	>10000	>10000	119	2595	106
Example #	33	34	35	36	37	38	39	40
IC ₅₀ (nM)	130	294	265	>10000	>10000	>10000	604	182
Example #	41	42	43	44	45	46	47	48
IC ₅₀ (nM)	11	78	51	6.8	34	21	54	56
Example #	49	50	51	52	53	54	55	56
IC ₅₀ (nM)	32	37	33	55	59	217	89	30
Example #	57	58	59	60	61	62	63	64
IC ₅₀ (nM)	46	171	>10000	>10000	61	816	50	177
Example #	65	66	67	68	69	70	71	72
IC ₅₀ (nM)	417	991	2054	673	352	136	70	80
Example #	73	74	75	76	77	78	79	80
IC ₅₀ (nM)	3379	>10000	2661	ND	2964	269	5543	>10000
Example #	81	82	83	84	85	86	87	88
IC ₅₀ (nM)	99	459	72	52	25	86	35	137
Example #	89	90	91	92	93	94	95	96
IC ₅₀ (nM)	ND	>10000	>10000	>10000	>10000	>10000	>10000	>10000
Example #	97	98	99	100	101	102	103	104
IC ₅₀ (nM)	>10000	>10000	>10000	202	117	87	45	8.3
Example #	105	106	107	108	109	110	111	112
IC ₅₀ (nM)	7.9	105	6.3	78	48	72	27	14
Example #	113	114	115	116	117	118	119	120
IC ₅₀ (nM)	9.6	51	19	48	5.1	0.82	0.72	42
Example #	121	122	123	124	125	126	127	128
IC ₅₀ (nM)	4.5	Td	15	9.9	6.7	18	13	37
Example #	129	130	131	132	133	134	135	136
IC ₅₀ (nM)	18	15	17	4.1	4.1	5.8	6.8	7.1
Example #	137	138	139	140	141	142	143	144
IC ₅₀ (nM)	7.0	351	16	37	12	22	15	2.8

Example #	145	146	147	148	149	150	151	152
IC ₅₀ (nM)	12	3.4	2.7	2.1	14	25	18	13
Example #	153	154	155	156	157	158	159	160
IC ₅₀ (nM)	9.2	7.3	23	28	18	7.8	648	394
Example #	161	162	163	164	165	166	167	168
IC ₅₀ (nM)	506	1193	28	6.2	49	4.0	5.2	15
Example #	169	170	171	172	173	174	175	176
IC ₅₀ (nM)	3.4	120	70	9.5	228	94	16	15
Example #	177	178	179	180	181	182	183	184
IC ₅₀ (nM)	>10000	7.9	12	3.1	4.9	16	13	62
Example #	185	186	187	188	189	190	191	192
IC ₅₀ (nM)	6.7	24	27	7.8	10	21	192	7.2
Example #	193	194	195	196	197	198	199	200
IC ₅₀ (nM)	4.5	18	1440	985	>10000	>10000	19	>10000
Example #	201	202	203	204	205	206	207	208
IC ₅₀ (nM)	>10000	>10000	>10000	>10000	>10000	>10000	>10000	37
Example #	209	210	211	212	213	214	215	216
IC ₅₀ (nM)	103	10	145	19	58	517	1388	>10000
Example #	217	218	219	220	221	223	224	225
IC ₅₀ (nM)	2.5	103	481	>10000	>10000	>10000	1113	>10000
Example #	226	227	228	229	230	AZD2281	ABT-888	
IC ₅₀ (nM)	84	13	>10000	>10000	2.6	28	2642	

ND, not determined.

In summary, 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (Example 4) and analogs have excellent potentiation effects on the growth inhibiting activity of DNA damaging anticancer agent MMS in a MTT based cell viability assay.

Example 232

Determination of the synergistic potentiation effect of 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione and analogs on MMS induced apoptosis using a cell based caspase-3 activity assay

Human breast cancer cell line T47D was used to determine the potentiation effect of PARP inhibitors on the apoptosis inducing activity of DNA damaging anticancer drugs such as MMS. Intracellular caspase-3 activity was used to measure cell apoptosis. The dose response

effect of MMS to induce intracellular caspase activity was carefully titrated and the concentration that only induces a minimal level of caspase activity was used to test the potentiation effect of PARP inhibitors. Briefly, T47D cells were grown in DMEM/F12 cell culture medium (Hyclone) supplemented with 0.2 U/ml insulin (Genview) and 10% FBS (Hyclone). A day before experiment, 20000 of T47D cells were seeded to each well of a 96-well cell culture plate and maintained at 37°C and 5% CO₂ in a cell culture incubator overnight. On the day of experiment, cell culture medium was removed. One hundred and eighty µl of fresh medium containing 100 nM MMS was added to each well and followed by 20 µl medium containing 10 folds of the concentrations of experimental drugs or reference compounds (ADZ2281 and ABT-888). The serial dilutions of the compounds to be tested and the reference compounds were made in a 1:3 and 1:10 fashion in DMSO. The 10-fold compound solutions were made by mixing 10 µl DMSO serial dilution solutions with 90 µl fresh growth medium. Twenty four hours later, the cells in the 96-well plates were centrifuged at 1000 g for 5 min and the supernatants were removed. Fifty µl lysis buffer (10mM Tris, pH7.5, 0.1M NaCl, 1 mM EDTA, 0.01% Triton X-100) was added to each well and the plates were shaken horizontally for 30 min at 4°C. After centrifugation at 1000 g at 4°C for 10 min, 20 µl of supernatant was transferred from each well to a corresponding well in a 384-well black plate. Twenty µl of buffer (20mM PIPES, pH7.4, 4 mM EDTA and 0.2% CHAPS) containing 20 µM fluorescent caspase-3 substrate ((Ac-DEVD)₂-R110, AnaSpec Cat #60304-5) was added to each well afterwards. The plates were shaken to uniformly mix the wells and incubated at 37°C for 3 h. The fluorescence intensity was measured using the following wavelength: ex: 496nm, em: 520nm using a fluorescence plate reader (Varioskan Flash, Thermo Fisher Scientific). The caspase-3 activity induced by the compounds was expressed as a relative fluorescence unit (RFU). The obtained fluorescence readings were analyzed using a commercial graphic software (GraphPad Prism 5) and plotted against the Log value of the compound concentrations. The EC₅₀ values were obtained by fitting the data points with the equation of $Y (\text{fluorescence reading}) = \text{minimal fluorescence reading} + (\text{maximal fluorescence reading} - \text{minimal fluorescence reading}) / (1 + 10^{-(\text{LogEC}_{50} - \text{LogC})})$, where C is the concentration of the testing compound.

The synergistic potentiation effect of compounds on MMS induced apoptosis in T47D cells is expressed as EC₅₀ values and listed in Table 2.

Table 2. Substituted 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones potentiate the apoptosis inducing activity of MMS (EC₅₀)

Example #	1	2	3	4	5	6	7	8
EC ₅₀ (nM)	>10000	>10000	56	12	450	2518	2655	120
Example #	9	10	11	12	13	14	15	16
EC ₅₀ (nM)	426	400	214	318	6601	39	803	9206
Example #	17	18	19	20	21	22	23	24
EC ₅₀ (nM)	5018	>10000	>10000	139	5797	>10000	4.0	20
Example #	25	26	27	28	29	30	31	32
EC ₅₀ (nM)	3.9	20	51	5894	>10000	39	112	96
Example #	33	34	35	36	37	38	39	40
EC ₅₀ (nM)	96	104	238	>10000	1914	267	614	29
Example #	41	42	43	44	45	46	47	48
EC ₅₀ (nM)	2.2	11	18	5.7	11	6.6	2.6	8.2
Example #	49	50	51	52	53	54	55	56
EC ₅₀ (nM)	10	13	20	22	18	124	20	16
Example #	57	58	59	60	61	62	63	64
EC ₅₀ (nM)	22	108	>10000	>10000	17	379	15	51
Example #	65	66	67	68	69	70	71	72
EC ₅₀ (nM)	373	604	4820	635	76	34	67	41
Example #	73	74	75	76	77	78	79	80
EC ₅₀ (nM)	811	2111	2869	ND	5206	112	3219	1192
Example #	81	82	83	84	85	86	87	88
EC ₅₀ (nM)	63	154	21	18	7.6	27	27	67
Example #	89	90	91	92	93	94	95	96
EC ₅₀ (nM)	ND	4168	1987	2926	>10000	>10000	>10000	>10000
Example #	97	98	99	100	101	102	103	104
EC ₅₀ (nM)	>10000	>10000	>10000	70	20	26	29	2.6
Example #	105	106	107	108	109	110	111	112
EC ₅₀ (nM)	14	217	9.5	68	23	32	1.7	2.9
Example #	113	114	115	116	117	118	119	120
EC ₅₀ (nM)	17	19	23	13	5.8	3.1	3.0	8.8
Example #	121	122	123	124	125	126	127	128
EC ₅₀ (nM)	4.9	8.4	11	3.8	22	6.5	4.1	12
Example #	129	130	131	132	133	134	135	136
EC ₅₀ (nM)	49	87	20	0.93	2.1	4.7	9.8	3.2

Example #	137	138	139	140	141	142	143	144
EC ₅₀ (nM)	2.1	281	4.0	8.9	13	3.8	6.0	3.4
Example #	145	146	147	148	149	150	151	152
EC ₅₀ (nM)	6.5	4.1	3.4	2.3	5.8	16	5.2	5.1
Example #	153	154	155	156	157	158	159	160
EC ₅₀ (nM)	2.0	6.7	12	14	26	8.6	220	261
Example #	161	162	163	164	165	166	167	168
EC ₅₀ (nM)	405	495	42	2.1	18	1.6	5.0	3.8
Example #	169	170	171	172	173	174	175	176
EC ₅₀ (nM)	1.0	43	23	3.2	122	37	14	10
Example #	177	178	179	180	181	182	183	184
EC ₅₀ (nM)	743	11	7.4	1.4	2.4	14	6.1	18
Example #	185	186	187	188	189	190	191	192
EC ₅₀ (nM)	2.1	6.8	30	ND	ND	ND	ND	3.1
Example #	193	194	195	196	197	198	199	200
EC ₅₀ (nM)	1.6	11	471	>10000	>10000	>10000	4.9	>10000
Example #	201	202	203	204	205	206	207	208
EC ₅₀ (nM)	>10000	>10000	>10000	>10000	>10000	>10000	>10000	24
Example #	209	210	211	212	213	214	215	216
EC ₅₀ (nM)	294	15	166	45	590	654	743	>10000
Example #	217	218	219	220	221	223	224	225
EC ₅₀ (nM)	19	37	445	>10000	>10000	>10000	274	>10000
Example #	226	227	228	229	230	AZD2281	ABT-888	
EC ₅₀ (nM)	24	40	>10000	>10000	3.2	24	3494	

ND, not determined.

In summary, as measured by the intracellular caspase activity, 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (Example 4) and its analogs have shown excellent potentiation effect on DNA damaging anticancer drug MMS induced cancer cell apoptosis.

Exmaple 233

Determination of the cell growth inhibiting activity of 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione and analogs on BRCA-2 deficient CAPAN-1 pancreatic duck cancer cells using a MTT based cell viability assay

BRCA-deficient cancer cells are sensitive to PARP inhibition. In the presence of PARP inhibitors, BRCA-deficient cancer cells could not grow and eventually will lead to death of cells. BRCA-2 deficient CAPAN-1 human pancreatic cancer cells were used to test the growth inhibiting activity of substituted 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones. Briefly, CAPAN-1 cells are grown and maintained in IMEM medium supplemented with 20% FBS. The assay takes 10 days to finish. On day 1, 6000 CAPAN-1 cells are seeded to each well of a 96-well plate. The cells are incubated at 37°C in a cell culture incubator overnight. In the next day, serial dilutions of compounds to be tested and reference compound (AZD2281) were prepared with a 1:3 and 1:10 fashion in DMSO. Ten-fold compound solutions were made by mixing 10 µl DMSO serial dilution solutions with 90 µl fresh growth medium. The cell culture medium was removed and 180 µl fresh media with FBS and 20 µl of compound solutions were added to each well. The final concentration of DMSO was 1%. Cells were then returned to the CO₂ cell culture incubator and incubated at 37°C for two days (day 3 and day 4). On day 5, the same procedure as what was done on day 2 was repeated. Cell culture medium was removed and fresh medium containing serial dilutions of compounds or reference compounds were added. Cells were returned to the CO₂ cell culture incubator and incubated for two more days. On day 8, same procedure was repeated again and fresh medium containing the testing compounds or reference compounds were added and incubated for an additional 2 days. On day 10, 20 µl of MTT reagent (5mg/ml) was added to each well of the cell culture plate and incubated for 4 h. The cell culture media was removed and 100 µl DMSO was added to each well. The 96-well cell culture plates were shaken horizontally for 10 min and readed in a Varioskan Flash plate reader (Thermo Fisher Scientific) at 520 and 690 nm. The obtained 520 nm absorbance readings subtracting corresponding 690 nm reading were analyzed using a commercial graphic software (GraphPad Prism 5) and plotted against the Log scale of the compound concentrations. The curves were fitted using the following equation to calculate the IC₅₀ value of each compound, $Y(520\text{ nm} - 690\text{ nm}) = \text{minimal absorbance value} + (\text{maximal absorbance value} - \text{minimal absorbance value}) / (1 + 10^{(\text{LogC} - \text{LogIC}_{50})})$, where C was the concentration of testing compounds.

The inhibitory effect of compounds on the CAPAN-1 cell growth is expressed as IC₅₀ values and listed in Table 3.

Table 3. Growth inhibition of CAPAN-1 cells by selected substituted 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones (IC₅₀)

Example #	3	4	8	11	15	20	23	24
-----------	---	---	---	----	----	----	----	----

IC ₅₀ (nM)	445	55	2360	1841	>10000	4942	160	735
Example #	25	26	27	30	40	41	42	43
IC ₅₀ (nM)	98	400	281	1785	1739	208	2539	974
Example #	44	45	46	48	49	50	51	56
IC ₅₀ (nM)	318	668	446	168	421	173	402	1181
Example #	65	69	70	81	83	85	100	101
IC ₅₀ (nM)	3189	1982	339	421	746	291	312	1049
Example #	103	104	105	107	111	112	113	117
IC ₅₀ (nM)	665	452	97	161	77	767	170	39
Example #	118	119	123	124	127	128	132	133
IC ₅₀ (nM)	54	56	111	137	44	442	17	22
Example #	134	136	139	140	142	143	144	145
IC ₅₀ (nM)	45	40	36	155	223	73	78	146
Example #	147	149	150	152	154	155	157	161
IC ₅₀ (nM)	47	52	176	121	56	142	202	3547
Example #	168	172	176	185	186	188	189	190
IC ₅₀ (nM)	1465	389	187	358	167	472	736	1383
Example #	199	210	212	217	AZD2281			
IC ₅₀ (nM)	25	948	3445	>10000	460			

In summary, 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (Example 4) and analogs are excellent inhibitors of CAPAN-1 cell growth, suggesting that they are good PARP inhibitors.

Exmample 234

Measurement of the inhibitory effect of 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione and analogs on PARP-1 enzymtic activity

The measurement of PARP-1 enzymtic activity was achieved by using a commercial 96-well colorimetric assay kit (4676-096-K, Trevigen, Inc. Gaithersburg, MD 20877 USA). Briefly, PARP-1 catalyzes the NAD-dependent addition of poly(ADP-ribose) to its nuclear protein substrates such as histones. The assay kit measures the incorporation of biotinylated poly(ADP-ribose) onto histone proteins in a 96-well format.

Reference compound (AZD2281) and compounds to be tested are serially diluted 1:10 with a 1X buffer. To each well of histone pre-coated plate 10 μ l of 5-fold concentrations of testing compounds or reference compound, 15 μ l of PARP-1 enzyme (0.5 unit) and 25 μ l

reaction buffer were added and the plates were incubated at room temperature for 60 min. The plates were washed with 200 μ l PBS with 0.1% Triton X-100 twice and then with 200 μ l PBS twice. The residual liquid was removed by carefully tapping the plates on paper towels. Equal volumes of PeroxyGlowTM solution A and B were mixed and 100 μ l of the solution was added to each well. The luminescence readings were read immediately in a Varioskan Flash plate reader (Thermo Fisher Scientific). The obtained luminescence readings were analyzed using a commercial graphic software (GraphPad Prism 5) and plotted against the Log scale of the compound concentrations. The IC₅₀ values were obtained by fitting the data points with the equation of Y (Luminescence reading) = minimal luminescence reading + (maximal luminescence reading - minimal luminescence reading)/(1+10^{^(LogC - LogEC₅₀)}), where C is the concentration of the testing compound.

The inhibitory effect of compounds on the PARP-1 enzymatic activity is expressed as IC₅₀ values and listed in Table 4.

Table 4. The inhibitory effect of substituted 1-(arylmethyl)quinazoline-2,4(1*H*,3*H*)-diones on PARP-1 enzymatic activity (IC₅₀)


Example #	3	4	8	14	25	44	127	132	135	139
IC ₅₀ (nM)	4.2	0.96	5.3	2.3	0.21	0.88	0.37	1.19	1.74	1.3
Example #	144	147	189	AZD2281						
IC ₅₀ (nM)	1.15	1.2	2.20	2.2						

In summary, as measured in the PARP-1 enzymatic assay, 1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione (Example 4) and analogs are potent PARP-1 inhibitors.

[0083] Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

CLAIMS:

1. A compound having Formula III:



or pharmaceutically acceptable salts thereof, wherein:

R₁-R₄ are selected from:

- 5 (a) R₁ is halo or C₁₋₆ alkyl; R₂ is H, halo, -NH₂ or C₁₋₆ alkyl; R₃ is H, halo, C₁₋₆ alkoxy or C₁₋₆ alkyl; and R₄ is H, halo, hydroxyl, C₁₋₆ alkoxy or C₁₋₆ alkyl;
- (b) R₁ is H, halo or C₁₋₆ alkyl; R₂ is halo, -NH₂ or C₁₋₆ alkyl; R₃ is H, halo, C₁₋₆ alkoxy or C₁₋₆ alkyl; and R₄ is H, halo, hydroxyl, C₁₋₆ alkoxy or C₁₋₆ alkyl;
- 10 (c) R₁ is H, halo or C₁₋₆ alkyl; R₂ is H, halo, -NH₂ or C₁₋₆ alkyl; R₃ is halo, C₁₋₆ alkoxy or C₁₋₆ alkyl; and R₄ is H, halo, hydroxyl, C₁₋₆ alkoxy or C₁₋₆ alkyl; or
- (d) R₁ is H, halo or C₁₋₆ alkyl; R₂ is H, halo, -NH₂ or C₁₋₆ alkyl; R₃ is H, halo, C₁₋₆ alkoxy or C₁₋₆ alkyl; and R₄ is halo, hydroxyl, C₁₋₆ alkoxy or C₁₋₆ alkyl;

R₇-R₁₀ independently are hydrogen, halo or C₁₋₆ alkyl; and

- 15 R₁₂ is C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylcarbonyl, pyridyl, pyrimidinyl, benzoyl, phenyl, piperidinyl, thienylcarbonyl, furyl or pyrazinyl, optionally substituted by 1, 2, 3 or 4 substituents selected from the group consisting of halo and C₁-C₆ alkyl.

2. The compound of claim 1, wherein

- 20 R₁ is F, Cl, Br or methyl;
- R₂ is H, F, Cl, Br or methyl;
- R₃ is H, F, Cl or methyl;
- R₄ is H, F, Cl, methyl, methoxy or hydroxy; and
- each of R₇, R₈, R₉, or R₁₀ is H or F.

3. The compound of claim 1, wherein
 R₂ is F, Cl, Br or methyl;
 R₁ is H, F, Cl, Br or methyl;
 R₃ is H, F, Cl or methyl;
 5 R₄ is H, F, methoxy or hydroxy; and
 each of R₇, R₈, R₉, or R₁₀ is H or F.
4. The compound of claim 1, wherein
 R₄ is F, methoxy or hydroxy;
 10 R₁ is H, F, Cl, Br or methyl;
 R₂ is H, F, Cl, Br or methyl;
 R₃ is H, F, Cl or methyl; and
 each of R₇, R₈, R₉, or R₁₀ is H or F
- 15 5. The compound of claim 1, wherein
 R₁ is halo;
 R₂ is H, halo or C₁-C₆ alkyl;
 R₃ is H, halo or C₁-C₆ alkyl;
 R₄ is H, halo or C₁-C₆ alkyl;
 20 R₇-R₁₀ independently are selected from the group consisting of H and halo; and
 R₁₂ is pyrimidinyl or pyridyl.
6. A compound selected from the group consisting of:
 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-fluoroquinazoline-
 25 2,4(1*H*,3*H*)-dione;
 7-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;
 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-7-methylquinazoline-
 2,4(1*H*,3*H*)-dione;
 30 7-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;
 6-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;

- 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(4-Fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;
- 5 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-methylquinazoline-2,4(1*H*,3*H*)-dione;
- 10 6-Bromo-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Bromo-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Bromo-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 15 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione;
- 6-Nitro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-nitroquinazoline-2,4(1*H*,3*H*)-dione;
- 20 5-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 25 5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 30 5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

- 5-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5 5-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 10 6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 15 6-Chloro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 20 6-Chloro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 25 6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 30 6-Chloro-1-(3-(4-(cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

- 6-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5 6-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 10 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 15 5-Fluoro-(1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline)-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 20 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 25 5-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 1 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-8-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 30 6-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

- 6-Fluoro-1-(3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(4-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 10 2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(4-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 15 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 20 2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 25 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(6-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 30 2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclohexylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;

- 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5 1-(3-(4-(Cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)-5-methylquinazoline-2,4(1*H*,3*H*)-dione;
- 7-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 7-Chloro-1-(3-(4-(cyclopentylcarbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 10 1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-(1-(3-(4-benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)quinazoline)-2,4(1*H*,3*H*)-dione;
- 15 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 20 5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 25 5-Chloro-1-(3-(4-(cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Chloro-1-(6-fluoro-3-(4-(thiophene-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 30 5-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Chloro-1-(6-fluoro-3-(4-(pyridin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

- 6-Chloro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5 1-(3-(4-(Cyclopropylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 10 1-(3-(4-Benzoylpiperazine-1-carbonyl)-4-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 15 1-(3-(4-(Cyclobutylcarbonyl)piperazine-1-carbonyl)-6-fluorobenzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(6-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 6-Fluoro-1-(4-fluoro-3-(4-(furan-2-carbonyl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 20 6-Fluoro-1-(5-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 1-(6-Chloro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-6-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
- 25 6-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-(thiazol-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione hydrochloride;
- 30 6-Fluoro-1-((2-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)pyridin-6-yl)methyl)quinazoline-2,4(1*H*,3*H*)-dione;
- 5-Fluoro-1-(4-fluoro-3-(4-methoxybenzylcarbamoyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

- 1-(3-(3-Chlorobenzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
 1-(3-(Benzylcarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-2,4(1*H*,3*H*)-dione;
 5-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 5 6-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 1-(3-(5-Bromopyrimidin-2-aminocarbamoyl)-4-fluorobenzyl)-5-fluoroquinazoline-
 2,4(1*H*,3*H*)-dione;
 1-(3-(4-(Pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-7-(trifluoromethyl)quinazoline-
 10 2,4(1*H*,3*H*)-dione;
 5-Fluoro-1-(6-methoxy-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 7-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 15 5-Fluoro-1-(4-fluoro-3-(4-(tetrahydrofuran-2-carbonyl)piperazine-1-
 carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;
 5-Fluoro-1-(4-nitro-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 1-(3-(4-Cyclohexylpiperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-
 20 2,4(1*H*,3*H*)-dione;
 5-Fluoro-1-(4-fluoro-3-(4-phenylpiperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;
 5-Fluoro-1-(4-fluoro-3-(4-phenylpiperidine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;
 25 1-(4-Bromo-3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)-5-fluoroquinazoline-
 2,4(1*H*,3*H*)-dione;
 6,7-Methylenedioxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-
 2,4(1*H*,3*H*)-dione;
 1-(3-(4-(Cyclohexylmethyl)piperazine-1-carbonyl)-4-fluorobenzyl)-5-fluoroquinazoline-
 30 2,4(1*H*,3*H*)-dione;
 8-Fluoro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;
 6-Amino-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-
 dione;

8-Chloro-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Methyl-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

5 8-Methoxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

8-Hydroxy-1-(3-(4-(pyrimidin-2-yl)piperazine-1-carbonyl)benzyl)quinazoline-2,4(1*H*,3*H*)-dione;

or a pharmaceutically acceptable salt or prodrug thereof.

10 7. Compound of any of claims 1-6, or a pharmaceutically acceptable salt thereof, for use as a medicament.

8. Compound for use according to claim 6, for use in treating or preventing a disorder responsive to the inhibition of PARP activity in a mammal suffering therefrom; preferably wherein said disorder is cancer.

15 9. Compound for use according to claim 8, wherein said cancer is liver cancer, melanoma, Hodgkin's disease, non-Hodgkin's lymphomas, acute or chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain
20 carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, malignant melanoma, choriocarcinoma, mycosis fungoide, head or neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal
25 carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer, or prostatic carcinoma.

10. Use of the compound of any of claims 1-6 in the manufacture of a medicament for treating or preventing a disorder responsive to the inhibition of PARP activity in a mammal
30 suffering therefrom.

11. The use according to claim 10, wherein said disorder is cancer.

12. The use according to claim 11, wherein said cancer is liver cancer, melanoma, Hodgkin's disease, non-Hodgkin's lymphomas, acute or chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor,

cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, malignant melanoma, choriocarcinoma, mycosis fungoide, head or neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer, or prostatic carcinoma.

13. The use according to claim 12, further comprising administering at least one known anticancer agent, or a pharmaceutically acceptable salt of said agent.

14. The use according to claim 12, wherein said compound is administered together with at least one compound selected from the group consisting of busulfan, melphalan, chlorambucil, cyclophosphamide, ifosfamide, temozolomide, bendamustine, cis-platin, mitomycin C, bleomycin, carboplatin, camptothecin, irinotecan, topotecan, doxorubicin, epirubicin, aclarubicin, mitoxantrone, elliptinium, etoposide, 5-azacytidine, gemcitabine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, fludarabine, nelarabine, ara-C, alanosine, pralatrexate, pemetrexed, hydroxyurea, thioguanine, colchicine, vinblastine, vincristine, vinorelbine, paclitaxel, ixabepilone, cabazitaxel, docetaxel, campath, Panitumumab, Ofatumumab, Avastin, Herceptin®, Rituxan®, imatinib, gefitinib, erlotinib, lapatinib, sorafenib, sunitinib, nilotinib, dasatinib, pazopanib, temsirolimus, everolimus, vorinostat, romidepsin, tamoxifen, letrozole, fulvestrant, mitoguazone, octreotide, retinoic acid, arsenic trioxide, zoledronic acid, bortezomib, thalidomide or lenalidomide.

15. The use according to claim 12, further comprising treating said mammal with radiation-therapy.

16. A pharmaceutical composition comprising the compound of any of claims 1 to 6 and a pharmaceutically acceptable carrier.

17. The pharmaceutical composition of claim 16, further comprising at least one known anticancer agent, or a pharmaceutically acceptable salt of said agent.

18. The pharmaceutical composition of claim 16, further comprising at least one compound selected from the group consisting of busulfan, melphalan, chlorambucil, cyclophosphamide, ifosfamide, temozolomide, bendamustine, cis-platin, mitomycin C, bleomycin, carboplatin, camptothecin, irinotecan, topotecan, doxorubicin, epirubicin,

- aclarubicin, mitoxantrone, elliptinium, etoposide, 5-azacytidine, gemcitabine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, fludarabine, nelarabine, ara-C, alanosine, pralatrexate, pemetrexed, hydroxyurea, thioguanine, colchicine, vinblastine, vincristine, vinorelbine, paclitaxel, ixabepilone, cabazitaxel, docetaxel, campath, Panitumumab, Ofatumumab, Avastin, 5 Herceptin®, Rituxan®, imatinib, gefitinib, erlotinib, lapatinib, sorafenib, sunitinib, nilotinib, dasatinib, pazopanib, temsirolimus, everolimus, vorinostat, romidepsin, tamoxifen, letrozole, fulvestrant, mitoguazone, octreotide, retinoic acid, arsenic trioxide, zoledronic acid, bortezomib, thalidomide or lenalidomide.