

DESCRIPTION

TITLE OF THE INVENTION

PRIMER SET FOR PCR FOR BACTERIAL DNA AMPLIFICATION, KIT FOR
5 DETECTION AND/OR IDENTIFICATION OF BACTERIAL SPECIES, AND,
METHOD OF DETECTION AND/OR IDENTIFICATION OF BACTERIAL
SPECIES

Technical Field

10 [0001]

The present invention relates to a primer set useful for detection
and/or identification of a bacterial species by an examination method using
PCR, particularly real time PCR, a kit containing this primer set for detection
or identification of a bacterial species, and a method of detection and/or
15 identification of bacterial species using them.

Background Art

[0002]

Microorganism examinations are technologies required in research
20 facilities, medical fronts, food manufacturing floors, etc. In medical fronts and
food manufacturing floors, smearing, microscopy, cultivation, etc. are
conventionally carried out as routine examination methods. In medical fronts,
a drug susceptibility test is often carried out concurrently since some
microorganisms may have antibiotic-resistant genes. However, the above-
25 described general examination methods such as cultivation, etc. need fairly
long times (days), thus, speeding up and higher sensitivity of examinations

are recently required and studied eagerly. There are various examination methods developed owing to analysis of pathogenic genes, analysis of toxin genes, analysis of antibiotic-resistant genes, analysis of DNA sequences (rDNA) coding rRNA under phylogenetic viewpoint, etc., particularly due to accumulation of the published database based on gene analysis.

[0003]

Now, gene examination methods are classified roughly into DNA probe methods and gene amplification methods (PCR). Especially, the PCR method is capable of amplifying a trace amount of DNA present in a sample in a short period of time and has high detection sensitivity as compared with the DNA probe method, and accordingly, its application range is wide.

[0004]

Under such a background, there is an investigation on detection of unidentified bacteria using PCR, and there is a trial of amplifying a trace amount of DNA of a sepsis-derived microorganism by PCR, hybridizing the amplified causal microorganism DNA with a nucleotide probe specific to bacterial species targeting an empirically supposed microorganism, and detecting the causal microorganism and identifying the bacterial species (JP-A No. 6-90799).

[0005]

Further, there is an investigation on development of sepsis examining technologies based on real time PCR using a hybridization probe, intending quickness of detection and identification of bacterial species (Journal of Analytical Bio-Science, Vol. 28, No. 5, (2005), 400 - 40). Furthermore, there is an investigation on quick detection of a causal bacterium and identification of the bacterial species, by conducting gene amplification by PCR utilizing

microbial DNA as a template and using several specific primer sets and then analyzing a combination of melting temperatures (T_m value) according to the resultant several DNA fragments or a difference between the T_m values (International Publication WO 2007/097323).

5 [0006]

However, it can be said that concomitant satisfaction of high sensitivity and high specificity is important in PCR, since precision should be ensured even in the result obtained in a short period of time. There prior technologies are a method limited to a supposed target microorganism though a gene
10 amplification technology by PCR is applied.

[0007]

In contrast, a method of producing a DNA polymerase preparation by a gene recombination technology is investigated, for providing a DNA polymerase used in PCR (JP-A No. 2006-180886). There is a suggestion on
15 a method of suppressing amplification of a non-specific nucleic acid in conducting amplification of an extremely trace amount of sample bacterial DNA by a PCR method using a heat-resistant DNA preparation produced by using eukaryote as a host (International Publication WO 2010/082640). By using, in PCR, this heat-resistant DNA polymerase preparation produced by
20 using eukaryote as a host, an analysis method which is suitable for novel examination of an extremely trace amount of a sample bacterium can be provided, and this technology is an extremely epoch-making method as a method of analyzing an extremely trace amount of a sample bacterium.

[0008]

25 Further, a method is supposed in which curves are obtained by high resolution melting (HRM) curve analysis for three DNA fragments amplified

from a bacterium, using a latest real time PCR apparatus, and the bacterial species is judged according to a combination of their shapes (Journal of Clinical Microbiology 2009, p.2252-2255: Journal of Clinical Microbiology 2010, p.3410-3413). However, these documents do not disclose clear

5 criteria for judging conformity or inconformity of wave shapes. In the above-described documents, judgment of 60 bacteria is performed, however, it is guessed to be difficult to identify a wide range of bacterium species over 100 kinds by way of three curves by HRM from three DNA fragments.

Additionally, a sample after hemoculture is used as a sample for extraction of

10 DNA in this method, needing a time required for culture.

[0009]

Further additionally, there is disclosed a method in which the T_m values of a amplified 16S rDNA fragment and several probes are measured using a real time PCR apparatus, and the bacterial species is specified

15 based on the T_m values (Journal of Clinical Microbiology 2010, p.258-267). Also in this method, a sample after hemoculture is used as a sample for extracting DNA, needing a time necessary for culture.

Prior Art Document

20

Patent Document

[0010]

Patent document 1: Japanese Patent Laid-Open No. 90799/1994 (JP H06-90799 A)

25 Patent document 2: International Publication WO 2007/097323

Patent document 3: Japanese Patent Laid-Open No. 2006-180886 (JP2006-180886 A)

Patent document 4: International Publication WO 2010/082640

5 Non-patent Document

[0011]

Non-patent document 1: Journal of Analytical Bio-Science, Vol. 28, No. 5, (2005), 400-40

Non-patent document 2: Journal of Clinical Microbiology 2009, p.2252-2255

10 Non-patent document 3: Journal of Clinical Microbiology 2010, p.3410-3413

Non-patent document 4: Journal of Clinical Microbiology 2010, p.258-267

Summary of the Invention

Problems to be solved by the Invention

15 [0012]

Based on the technology disclosed in Patent document 4, it is believed that if DNA can be amplified simply from an extremely trace amount of a sample bacterium and if analysis, particularly quantification or identification analysis can be effected in a short period of time, then, even a gene at
20 extremely trace amount level unanalyzable to date can be analyzed, and additionally, quick and correct judgment is attained in various sample analysis fields such as the medical field, the veterinary field, the fields of daily water and food, etc. However, further improvement in detection sensitivity and quickness are needed from the clinical sites, food production floors, etc.

25 [0013]

As described above, a detection method using a PCR method gives higher quickness as compared with a DNA probe method, and is suitable also for detection with high sensitivity, thus, is useful as a quick and highly sensitive detection method in the clinical sites, food production floors, etc..

- 5 However, conventional detection methods using a PCR method generate a case in which detection is impossible when the bacterial species contained in a sample is unclear and a case in which furthermore improvement in detection sensitivity and quickness cannot be sufficiently satisfied due to a longer time necessary for bacterial proliferation in a sample for ensuring the amount of nucleic acids necessary for detection, etc.
- 10

[0014]

- For example, in Non-patent document 4, a fair amount of bacterial bodies necessary for analysis after culture, etc. is understood also from a use amount of bacterial DNA subjected to PCR of 2-10 ng. When *E. coli* is exemplified as the bacterium, the genome size of the *E. coli* MG1655 strain is
- 15 4639675 bp, and if the average molecular weight of one base pair is calculated from this and further if the presence of one copy (1 molecule) of a genome per one bacterium cell is taken into consideration, then, it is understood that the amount of the genome DNA per 1 cell of *E. coli* is about
- 20 5 fg and the bacterial amount corresponding to the amount of DNA subjected to PCR is the fair amount, leading to a supposition that the detection sensitivity of this method is not sufficient.

[0015]

- In a case of detection and identification of a bacterium using a PCR
- 25 method, selection of a primer set used in a PCR method is one important factor for enhancing sensitivity of detection and identification of an

examination. For example, the amplification efficiency in PCR, the threshold cycle, the production ratio of the unintended amplification product, etc., vary depending on selection of a primer set, and for enhancing sensitivity of detection and identification of an examination, a primer set optimizing these factors is necessary. Depending on the kind of a sample, there is also a case where large amount of DNA derived from human cells other than the DNA as an analyte to be detected is present in the sample, such as a sample derived from blood, etc. For example, large amount of DNA derived from human cells such as leucocytes, desquamated epithelial cells, etc. is present in a blood specimen. According to investigations of the present inventors, when the content of the bacterial DNA as a target to be tested is small in such samples, selectivity by a primer set for PCR to bacterial DNA to be detected or productivity of the amplification product become low. As a result, there are cases where an amount of amplification product necessary for identification of species as a target to be tested cannot be obtained, or where rising of the background in identification occurs due to increase of undesired DNA amplification products. In Patent document 4, remarkable improvement in the identification sensitivity is attained by using a heat-resistant DNA polymerase preparation containing a suppressed amount of bacteria-derived DNA produced by using eukaryote as a host for identification of a bacterium as an analyte using a PCR method, in an identification method by a conventional technology which Patent document 4 utilizes, however, it is required to enable identification at further higher sensitivity also for a sample in which the amount of DNA of a bacterium as an analyte is further smaller for a bacterium other than the bacterium as an analyte present in large amount. Also from such a standpoint, design of a primer giving higher

selectivity, not generating unintended amplification products, is important for a primer set to be used in a PCR method. However, a primer set designed noticing the technical problems in using samples such as blood, etc. as an analyte as described above has not been found in conventional technologies,
5 under present circumstances.

[0016]

Further, in analyzing a intended bacterium to be detected and identifying an unidentified bacterium using the T_m value of a amplification product, it is preferable to select a primer set giving a amplification product
10 having a T_m value suitable for enhancing quickness of an examination and examination sensitivity.

[0017]

In contrast, also the property of a heat-resistant DNA polymerase preparation used for amplification by a PCR method and the use condition
15 thereof are one of important factors for enhancing the detection sensitivity of an examination using a PCR method.

[0018]

Commercially available heat-resistant DNA polymerase preparations generally used in a PCR reaction include also highly purified preparations
20 supplied in the market, and in a PCR reaction using these highly purified preparations, amplification products of unknown reasons and derivation are detected when a gene amplification reaction is required to be repeated in a frequency over usual 30 cycles, for attaining detection at high sensitivity, thus, use thereof is limited.

25 [0019]

Patent document 4 describes mixing of host-derived bacterial DNA of a DNA polymerase, however, does not describe mixing of bacteria-derived DNA as the other contamination source from reagents and instruments. In contrast, it has been found by investigation of the present inventors that
5 mixing of bacteria-derived DNA as a contamination source from reagents and instruments exerts an extremely critical influence on detection of an extremely trace amount of bacterial DNA in a sample and identification of the bacterial species.

[0020]

10 Means for controlling sensitivity and specificity together at higher levels in PCR analysis for amplifying an extremely trace amount of bacteria-derived DNA in a sample have not been provided yet, and also identification of bacterial species at high sensitivity targeting bacteria-derived DNA in a sample has not been sufficiently attained.

15 [0021]

By a detection method using a heat-resistant DNA polymerase preparation produced by using eukaryote as a host disclosed in Patent document 2, the detection limit of bacteria-derived DNA can be lowered to 10 fg/ μ L, and analysis of sample-derived DNA at higher sensitivity becomes
20 possible as compared with a conventional detection method using PCR.

[0022]

According to the method disclosed in Patent document 2, bacterial DNA of a concentration of 10 fg/ μ L or more can be detected when a PCR method is carried out using SEQ ID No. 51 and SEQ ID No. 52, however,
25 DNA of a concentration of less than 10 fg/ μ L cannot be detected. It can be confirmed from this that if amplification of an intended product cannot be

observed in performing PCR using SEQ ID No. 51 and SEQ ID No. 52 by the method disclosed in Patent document 2 using a certain solution as a template, then, the concentration of bacteria-derived DNA mixed in its solution is less than 10 fg/ μ L.

5 [0023]

For conducting not only detection of bacterial species in a sample but also identification of bacterial species in a sample at high sensitivity with high precision, it is necessary that a sample-derived DNA amplification product is obtained in an amount sufficient for identification of bacterial species, in
10 addition to lowering of the detection limit of the amount of bacterial-derived DNA. For example, for obtaining the highly reliable measurement result of the T_m value in performing identification of bacterial species in a sample by measuring the T_m value of the amplification product of sample-derived DNA, it is important that the amplification product in an amount sufficient for this is
15 obtained by PCR. In other words, when sufficient amplification of DNA is not conducted, the T_m value cannot be measured correctly, and identification with high precision becomes difficult.

[0024]

For example, in the case of a detection method in which the amount of
20 genome DNA per one cell of *E. coli* is about 5 fg and the detection limit is 10 fg/ μ L, these amount of DNA per 1 μ L of a sample provided to PCR corresponds theoretically to two cells of *E. coli* and, thus, detection of *E. coli* at high sensitivity in a sample becomes possible. In contrast, if the amount of *E. coli*-derived DNA in a sample which is necessary at minimum for
25 measuring the reliable T_m value of a DNA amplification product, namely, the detection limit is assumed to 20 pg/ μ L, it is supposed that the bacterial

amount corresponding to the amount of DNA provided to PCR is a fair amount.

[0025]

Investigation of the present inventors obtained novel findings that it is
5 effective to use a primer set giving further improved amplification efficiency
of a amplification product and selectivity, for further lowering the detection
limit and conducting identification of bacterial species at high sensitivity with
good precision in a measurement system capable of detecting bacterial
species at high sensitivity. For example, primers made of sequence regions
10 common to 16SrRNA gene of all bacteria used in Patent document 2 and
Patent document 4 have the same sequence, and if a primer set giving
further improved amplification efficiency and selectivity of the amplification
product than primer sets disclosed in these patent documents can be
provided, it becomes possible to further improve not only the detection
15 sensitivity of bacterial species but also the identification sensitivity of bacterial
species.

[0026]

Therefore, the present invention has an object of providing a primer
set which is useful for further improvement of sensitivity in detection or
20 identification of bacterial species in a sample by a PCR method, a set for
PCR using this, and a method of detection or identification of bacterial
species in a sample.

[0027]

The present invention has another object of attaining further
25 improvement of sensitivity in detection or identification of bacterial species in

a sample, by conducting a PCR method using the above-described primer set while minimizing the mixing amount of bacteria-derived nucleic acids.

Means for Solving the Problem

5 [0028]

The present inventors have intensively studied in view of the above-described problems and resultantly found that bacterial species identification kit capable of carrying out detection of an extremely trace amount of bacterial DNA in a sample and identification of the bacterial species at high sensitivity
10 with high precision can be obtained by using a specific primer set and further minimizing mixing of bacteria-derived DNA as a contamination source of reagents and instruments to be used.

[0029]

A primer set for polymerase chain reaction (PCR) for amplification of
15 bacterial DNA according to the present invention is characterized in comprising at least one primer pair of seven primer pairs obtained by selecting one primer pair from each group of Groups S1 to S7 as described below.

[0030]

20 A kit for detection of a bacterial species according to the present invention is characterized in comprising at least one primer pair of seven primer pairs obtained by selecting one primer pair from each of Groups K1 to K7 as described below.

[0031]

25 The above kit may further comprise at least one of the reagents and the parts selected from an enzyme for PCR, a pH buffer, fluorochrome, dNTP,

MgCl₂ and a vessel for measurement, which may have, independently or as a whole, a contamination amount of bacterial nucleic acid of 10 fg or less than 10 fg.

[0032]

5 A method of detection and/or identification of an bacterial species as a target in a sample according to the present invention is characterized in comprising:

 a step of conducting PCR using bacterial genome DNA prepared from the sample, primers to obtain amplification product comprising a target gene
10 specific to the bacterial species as a target, a thermostable DNA polymerase, and

 a step of detecting and/or identifying the bacterial species as a target in the sample by detecting the target gene in the amplification product by PCR, or analyzing the amplification product by PCR,

15 wherein, as the primers, the above described primer set is used.

Effect of the Invention

[0033]

 The present invention can provide a primer set for bacterial DNA
20 amplification PCR useful for detection and/or identification of bacterial species at high sensitivity and a kit for detection and/or identification of bacterial species. Further, sensitivity in detection and/or identification of bacterial species can be further improved by minimizing mixing of bacteria-derived DNA as a contamination source of reagents and instruments used in
25 this kit. Therefore, according to the present invention, detection of an extremely trace amount of bacteria-derived DNA in a sample and detection

and/or identification of bacterial species in the medical field, food field, etc.
can be carried out at higher sensitivity and with higher precision than
conventional examination kits.

5 Brief Explanation of Drawings

[0034]

[Figure 1A] The figure is a view showing a part of the results of
electrophoresis in Example 2 and Comparative Example 2.

10 [Figure 1B] The figure is a graph showing a part (amplification curve)
of the results of real time PCR in Example 3 and Comparative Example 3.

[Figure 1C] The figure is a graph showing a part (melting curve) of
the results of real time PCR of Example 3 and Comparative Example 3.

[Fig. 2] The figure is a view showing the result in Example 12.

[Fig. 3] The figure is a view showing the result in Example 13.

15 [Fig. 4] The figure is a view showing the result in Example 14.

Lines represent the amplification results of the following primer pairs. (1) a
combination of SEQ ID No. 15 and 16, (2) a combination of SEQ ID No. 43
and 50, (3) a combination of SEQ ID No. 51 and 52, (4) a combination of
SEQ ID No. 53 and 57, (5) a combination of SEQ ID No. 62 and 70, (6) a
20 combination of SEQ ID No. 81 and 103 and (7) a combination of SEQ ID No.
115 and 5.

[Fig. 5] The figure is a view showing the result in Example 16.

Modes for Carrying Out the Invention

25 [0035]

<Primer>

The first point of the present invention resides in a specific primer set capable of obtaining an effect of improvement of detection sensitivity and an effect of improvement of identification precision based thereon.

[0036]

5 Primers in primer sets described in Patent document 4 have DNA sequences hybridizing to bacterial-common conservation regions in DNA (16S rDNA) coding 16S rRNA of the bacteria (SEQ ID No. 1, SEQ ID No. 16, SEQ ID No. 28, SEQ ID No. 44, SEQ ID No. 51, SEQ ID No. 52, SEQ ID No. 53, SEQ ID No. 57, SEQ ID No. 62, SEQ ID No. 70, SEQ ID No. 76, SEQ ID
10 No. 91, SEQ ID No. 104, SEQ ID No. 2). As shown in lane 2 in Fig. 1A, however, bands of other components other than the analyte to be detected appeared and the intended amplification product could not be detected in electrophoresis analysis by the primer sets described in Patent document 4. In addition, cases with low amplification efficiencies were observed
15 occasionally in real time PCR (see dotted line in Fig. 1B). Furthermore, a peak (or shoulder) other than the main peak emerged in melting curve analysis in real time PCR (see dotted line in Fig. 1C). These may constitute an obstacle to further improvement of detection sensitivity in detection of an extremely trace amount of sample-derived DNA and identification of bacterial
20 species.

[0037]

 The seven regions recognized by the primer sets of Patent document 2 were investigated, while there are commonly conserved bacterial regions on 16S rDNA. Specifically, regions were selected base on the primer sets of
25 Patent Document 2, which shifted from several bases to several tens of bases in the directions to the 3' terminals and the 5' terminals of the

commonly conserved regions, respectively, and, then, each of forward and reverse primers was designed in a length of a base number of ten to several tens. The effects were confirmed for the primer pairs corresponding to the seven designed conserved regions, and it was found that the primer pairs
5 classified to the following groups S1 through S7 and the primer pairs classified to the following groups K1 through K7 were suitable.

[0038]

Thus, a primer set for PCR for amplification of bacterial DNA, which is applicable for detection and/or identification of a bacterial species, is
10 prepared by selecting one primer pair from every one of Groups S1 to S7 to obtain seven primer pairs, i. e., a first to a seventh primer pair, and using at least one primer pairs, preferably at least four primer pairs from the primer pair group thus selected.

A primer set for identification of a bacterial species may be prepared
15 by using at least 4 primer pairs of the first to the seventh primer pairs thus selected.

(Primer Pairs for Primer Set)

Group S1:

20 1-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 1 and a reverse primer consisting of any one sequence of SEQ ID: No. 22 and 27;

1-2A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 10 and a reverse primer
25 consisting of any one sequence of SEQ ID: No. 20 and 21;

1-3A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 11 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 18 and 26;

5 1-4; a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 12 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 13 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

10 1-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 15 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-8A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 6 and a reverse primer
15 consisting of any one sequence of SEQ ID: No. 16, 19 to 21, 23, 24, 26 and 27;

1-9A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 7 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21, 23, 24, 26 and 27;
20 or

1-10A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 8 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21, 23, 24, 26 and 27;
Group S2:

2-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 28 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, and 47 to 50;

5 2-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 29 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45, 47, 48 and 50;

2-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 30 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, and 47 to 50;

10 2-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 31 and a reverse primer consisting of any one sequence of SEQ ID: No. 46 and 47;

2-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 32 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, and 47 to 49;

2-6A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 33 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45 and 47 to 50;

20 2-7A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 34 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, 47 and 48;

2-8A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 35 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45, 48 and 49;

2-9A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 36 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45, 48 and 49;

2-10A: a primer pair, which consists of a combination of a forward
5 primer consisting of a sequence of SEQ ID: No. 37 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, and 47 to 50;

2-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 38 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

10 2-12A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 39 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 48 and 50;

2-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 40 and a reverse primer
15 consisting of any one sequence of SEQ ID: No. 44 to 50;

2-14A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 41 and a reverse primer consisting of any one sequence of SEQ ID: No. 45 and 46;

2-15A: a primer pair, which consists of a combination of a forward
20 primer consisting of a sequence of SEQ ID: No. 42 and a reverse primer consisting of any one sequence of SEQ ID: No. 48 and 49; or

2-16A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 46 to 48 and 50;

25 Group S3:

a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

Group S4:

5 4-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 61;

 4-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 54 and a reverse primer consisting
10 of any one sequence of SEQ ID: No. 59 and 61;

 4-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 55 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61; or

 4-4: a primer pair, which consists of a combination of a forward primer
15 consisting of a sequence of SEQ ID: No. 56 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61;

Group S5:

 5-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer
20 consisting of any one sequence of SEQ ID: No. 70, 73 to 75;

 5-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 63 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

 5-3: a primer pair, which consists of a combination of a forward primer
25 consisting of a sequence of SEQ ID: No. 64 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 65 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5 5-5A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 66 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 72, 73 and 75;

5-6A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 67 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 71 and 73 to 75;

10 5-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 68 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75; or

5-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 69 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

Group S6:

6-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 76 and a reverse primer consisting of any one sequence of SEQ ID: No. 93, 96 and 101 to 103;

20 6-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 77 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, and 101 to 103;

6-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 78 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

25

6-4A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 79 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

5 6-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 80 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, and 100 to 103;

6-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, and 101 to 103;

10 6-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 82 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, 97, and 101 to 103;

15 6-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 83 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 101 to 103;

6-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 84 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

20 6-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 85 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

6-11A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 86 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, and 101 to 103;

6-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 87 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, and 101 to 103;

5 6-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 88 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

6-14: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 89 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103; or

10 6-15: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 90 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 92, 96, and 101 to 103;

Group S7:

15 7-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 104 and a reverse primer consisting of any one sequence of SEQ ID: No. 5, and 118 to 125, 127, 128, 130 and 131;

20 7-2A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 105 and a reverse primer consisting of any one sequence of SEQ ID: No. 5, and 118 to 125;

7-3A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 106 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, 116, and 118 to 131;

25 7-4A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 107 and a reverse primer consisting of any one sequence of SEQ ID: No. 120 and 121;

7-5A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 108 and a reverse primer consisting of any one sequence of SEQ ID: No. 120 and 123;

5 7-6A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 109 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 118 to 122, 124 and 126;

7-7A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 110 and a reverse primer consisting of any one sequence of SEQ ID: No. 118, 120, 121, 124, and 126
10 to 130;

7-8A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 111 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 118 to 131;

7-9A: a primer pair, which consists of a combination of a forward
15 primer consisting of a sequence of SEQ ID: No. 112 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116, and 118 to 125;

7-10A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 113 and a reverse primer
20 consisting of any one sequence of SEQ ID: No. 2, 5, 118 to 126, 130 and 131;

7-11A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 114 and a reverse primer consisting of a sequence of SEQ ID: No. 5; or

7-12A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of any one sequence of SEQ ID: No. 2 and 5.

Furthermore, it has been revealed that the primer pairs divided into the following Groups K1 to K7 is suitable to compose a kit for detection of a bacterial species.

[0039]

Thus, a kit for PCR for detection and/or identification of a bacterial species, is prepared by selecting one primer pair from every one of Groups K1 to K7 to obtain seven primer pairs, i. e., a first to a seventh primer pair, and using at least one primer pairs, preferably at least 4 primer pairs from the primer pair group thus selected.

A kit for identification of a bacterial species may be prepared by using at least 4 primer pairs of the first to the seventh primer pairs thus selected.

[0040]

(Primer Pairs for Kit)

Group K1:

1-1: a primer pair , which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 1 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 10 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 18, 20 and 21;

1-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 11 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21, and 23 to 26;

1-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 12 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

5 1-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 13 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 14 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

10 1-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 15 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

15 1-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 6 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 7 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

20 1-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 8 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27; or

1-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 9 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 17, 19 and 26;

25 Group K2:

2-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 28 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45, and 47 to 50;

5 2-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 29 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 45, 47, 48 and 50;

2-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 30 and a reverse primer consisting of any one sequence of SEQ ID: No. 45, and 47 to 50;

10 2-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 31 and a reverse primer consisting of any one sequence of SEQ ID: No. 46 and 47;

2-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 32 and a reverse primer consisting
15 of any one sequence of SEQ ID: No. 45, and 47 to 49;

2-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 33 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

2-7: a primer pair, which consists of a combination of a forward primer
20 consisting of a sequence of SEQ ID: No. 34 and a reverse primer consisting of any one sequence of SEQ ID: No. 45 to 50;

2-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 35 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

2-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 36 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 46, and 48 to 50;

5 2-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 37 and a reverse primer consisting of any one sequence of SEQ ID: No. 45 to 50;

2-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 38 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

10 2-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 39 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

2-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 40 and a reverse primer
15 consisting of any one sequence of SEQ ID: No. 44 to 50;

2-14: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 41 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50;

2-15: a primer pair, which consists of a combination of a forward
20 primer consisting of a sequence of SEQ ID: No. 42 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 50; or

2-16: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 48, and 50;

25 Group K3:

a primer pair , which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

Group K4:

5 4-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 61;

 4-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 54 and a reverse primer consisting
10 of any one sequence of SEQ ID: No. 59 and 61;

 4-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 55 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59, and 61; or

 4-4: a primer pair, which consists of a combination of a forward primer
15 consisting of a sequence of SEQ ID: No. 56 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59, and 61;

Group K5:

 5-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting
20 of any one sequence of SEQ ID: No. 70, 71, and 73 to 75;

 5-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 63 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

 5-3: a primer pair, which consists of a combination of a forward primer
25 consisting of a sequence of SEQ ID: No. 64 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 65 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5 5-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 66 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 67 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

10 5-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 68 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75; or

5-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 69 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

Group K6:

6-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 76 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, 96, and 101 to 103;

20 6-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 77 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, and 101 to 103;

6-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 78 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

25

6-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 79 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 97, and 101 to 103;

5 6-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 80 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, and 100 to 103;

6-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, and 101 to 103;

10 6-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 82 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, 97, and 101 to 103;

15 6-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 83 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 101 to 103;

6-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 84 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

20 6-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 85 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

6-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 86 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, 97, and 101 to 103;

6-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 87 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, and 101 to 103;

5 6-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 88 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103;

6-14: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 89 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, and 101 to 103; or

10 6-15: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 90 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 92, 96, and 101 to 103;

Group K7:

15 7-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 104 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

7-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 105 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

20 7-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 106 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

25 7-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 107 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

7-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 108 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

5 7-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 109 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

7-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 110 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

10 7-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 111 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

7-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 112 and a reverse primer consisting
15 of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

7-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 113 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, and 116 to 131;

20 7-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 114 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 125; or

7-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, 116, 118, 123, and 125
25 to 128.

[0041]

The above primer set, which comprises a primer pair selected from Groups S1 to S7, preferably comprises at least one primer pair from the following seven primer pairs of S-1) to S-7).

5 S-1) as the primer set of Group S1, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 15 and a reverse primer consisting of a sequence of SEQ ID: No. 16;

S-2) as the primer set of Group S2, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of a sequence of SEQ ID: No. 50;

10 S-3) as the primer set of Group S3, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

S-4) as the primer set of Group S4, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of a sequence of SEQ ID: No. 57;

S-5) as the primer set of Group S5, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting of a sequence of SEQ ID: No. 70;

20 S-6) as the primer set of Group S6, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of a sequence of SEQ ID: No. 103; and

S-7) as the primer set of Group S7, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of a sequence of SEQ ID: No. 5.

25 [0042]

In addition, the above primer set comprises preferably any one of the primer pairs of the above S-1) to S-7) as an essential component for detection of bacterial species, and comprises preferably any one combination of the following A to G for detection and/or identification of bacterial species.

5 (A) when the primer set essentially comprises primer set S-1), the primer set further comprises at least 3 primer pairs selected from S-2) to S-7);

(B) when the primer set essentially comprises primer set S-2), the primer set further comprises at least 3 primer pairs selected from S-1) and S-10 3) to S-7);

(C) when the primer set essentially comprises primer pair S-3), the primer set further comprises at least 3 primer pairs selected from S-1), S-2) and S-4) to S-7);

(D) when the primer set essentially comprises primer pair S-4), the 15 primer set further comprises at least 3 primer pairs selected from S-1) to S-3) and S-5) to S-7);

(E) when the primer set essentially comprises primer pair S-5), the primer set further comprises at least 3 primer pairs selected from S-1) to S-4) and S-6) to S-7);

20 (F) when the primer set essentially comprises primer pair S-6), the primer set further comprises at least 3 primer pairs selected from S-1) to S-5) and S-7);

(G) when the primer set essentially comprises primer pair S-7), the primer set further comprises at least 3 primer pairs selected from S-1) to S-25 6).

[0043]

The above primer set, which comprises a primer pair selected from the above Groups K1 to K7, preferably comprises at least one primer pair from the seven primer pairs of the following K-1) to K-7).

5 K-1) as the primer set of Group K1, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 15 and a reverse primer consisting of a sequence of SEQ ID: No. 16;

K-2) as the primer set of Group K2, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of a sequence of SEQ ID: No. 50;

10 K-3) as the primer set of Group K3, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

K-4) as the primer set of Group K4, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of a sequence of SEQ ID: No. 57;

K-5) as the primer set of Group K5, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting of a sequence of SEQ ID: No. 70;

20 K-6) as the primer set of Group K6, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of a sequence of SEQ ID: No. 103; and

K-7) as the primer set of Group K7, a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of a sequence of SEQ ID: No. 5.

25 [0044]

In addition, the above primer set for a kit comprises preferably any one of the primer pairs in the above K-1) to K-7) as an essential component for detection of bacterial species, and comprises preferably any one combination of the following A to G for detection and/or identification of bacterial species.

5 (A) when the primer set essentially comprises primer set K-1), the primer set further comprises at least 3 primer pairs selected from K-2) to K-7);

(B) when the primer set essentially comprises primer set K-2), the primer set further comprises at least 3 primer pairs selected from K-1) and K-
10 3) to K-7);

(C) when the primer set essentially comprises primer pair K-3), the primer set further comprises at least 3 primer pairs selected from K-1), K-2) and K-4) to K-7);

(D) when the primer set essentially comprises primer pair K-4), the
15 primer set further comprises at least 3 primer pairs selected from K-1) to K-3) and K-5) to K-7);

(E) when the primer set essentially comprises primer pair K-5), the primer set further comprises at least 3 primer pairs selected from K-1) to K-4) and K-6) to K-7);

20 (F) when the primer set essentially comprises primer pair K-6), the primer set further comprises at least 3 primer pairs selected from K-1) to K-5) and K-7);

(G) when the primer set essentially comprises primer pair K-7), the primer set further comprises at least 3 primer pairs selected from K-1) to K-
25 6).

[0045]

By combining four from the primer pairs S-1 to S-7 in the above-described primer sets, identification of bacterial species can be conducted, and it is preferable that the number of pairs selected from these 7 primer pairs is increased to 5 or more, and it is more preferable that all of the seven primer pairs S-1 to S-7 are combined. Identification precision can be further improved by thus increasing the number of pairs to be selected.

[0046]

Likewise, by combining four out of primer pairs K-1 to K-7 in the above-described primer sets, identification of bacterial species can be conducted, and it is preferable that the number of pairs selected from these 7 primer pairs is increased to 5 or more, and it is more preferable that all of the seven primer pairs K-1 to K-7 are combined. Identification precision can be further improved by thus increasing the number of pairs to be selected.

[0047]

Those, in which some bases in a primer sequence are converted, can also be applied, provided that the effect of the present invention can be obtained. Basically, those hybridizable with complementary strands of DNA fragments having SEQ ID Numbers described in the claims of the present application are regarded as equivalent to the primer of the present invention. The primer sequences obtained by addition, deletion or substitution of 1 to 2 bases of primer sequences are regarded also as equivalent to the primer of the present invention.

[0048]

It is noted that the primer sets of the present invention have been found for the first time after confirmation of their effects based on a large amount of combinations and could not be attained by simple confirmation of

the effects, and that the primer sets have been confirmed as the novel combinations as the primer sets for detection and/or identification of bacterial species, as a result of sequence search.

[0049]

5 The present invention does not prevent concomitant use of several known primers for fungi, specific microorganisms or viruses, in addition to the primer set of the present invention.

[0050]

10 The primer set of the present invention is suitably used for the purpose of amplifying the intended DNA fragment by PCR from bacterial DNA extracted from a sample, measuring T_m of the amplified DNA fragment, measuring the molecular weight thereof, and decoding its sequence. Specifically, it is suitably used for identifying bacterial species by the T_m value of a DNA fragment composed of a part of 16S rDNA sequence of a
15 bacterium to be amplified by the primer set of the present invention.

[0051]

20 In amplification of DNA by PCR and in measurement of T_m , DNA containing the intended sequence may be previously amplified by a pair of forward and reverse primers having a base number of 15 to 30 b containing a DNA sequence hybridizing to bacterial-common conservation region on 16S rDNA before using at least one primer pair out of the above-described first to seventh primer pairs for a primer set or for a kit, and PCR amplification may be further performed on the amplified sample by the above-described primer set (so-called, Nested PCR method and Semi-nested PCR method).

25 [0052]

The pair of forward and reverse primers having a base number of 15 to 30 b used for first amplification of DNA by PCR is not particularly restricted providing it contains a DNA sequence hybridizing to a bacterial-common conservation region on 16S rDNA sequence, and it is preferable that the
5 DNA chain length of the PCR product using this primer pair is 500 bp to 1500 bp. In particular, primers including those which generate amplification product, to which a primer set for the second amplification hybridizes, concretely such the primer pair described in Groups K 1 to K7, and, thus, from which at least four DNA fragments are obtained, are preferable.

10 [0053]

The primer set used in the present invention is suitably used for detection of an extremely trace amount of sample bacterial DNA and/or identification of bacterial species. Therefore, it is necessary that contamination of bacteria-derived DNA as a contamination source into each
15 primer pair is suppressed to minimum. The amount of contamination of bacteria-derived DNA as a contamination source is less than 10 fg, preferably less than 5 fg, further preferably less than 1 fg with respect to a primer pair preparation solution.

[0054]

20 The primer concentration in PCR is not particularly restricted and in general, it is appropriately set depending on investigation in a range of 0.05 μM to 1.0 μM .

[0055]

<Reagent>

25 As the reagents other than the primer set used for PCR analysis in the present invention, any known reagents can be used in any known

combinations. The reagents necessary for measurement include an enzyme for PCR, a pH buffer solution, dNTP, a Mg^{2+} source, and sterile water purified by a filtration treatment, etc. In real time PCR analysis, a fluorochrome is required in addition to the above-described materials.

5 [0056]

<Enzyme for PCR>

As the enzyme for PCR used in the present invention, a heat-resistant DNA polymerase in which the amount of contamination of bacteria-derived DNA is suppressed to minimum is preferably used. Specific examples
10 thereof include, but not limited to, those produced by eukaryotes, those highly purified, or those treated with EMA (ethidium monoazide) or PMA (propidium monoazide) as a selectively membrane-permeable dye.

[0057]

In the present invention, the amount of mixing of bacteria-derived DNA
15 from a host or contamination source is less than 10 fg, preferably less than 5 fg, further preferably less than 1 fg with respect to an enzyme preparation solution. In the case of use of a heat-resistant DNA polymerase preparation (preparation solution) produced by an eukaryote utilizing a bacteria-derived heat-resistant DNA polymerase gene in which DNA from a bacteria-derived
20 heat-resistant DNA polymerase gene is mixed, the amount of mixing of bacteria-derived DNA other than a gene coding the heat-resistant DNA polymerase is not higher than the above-described upper limit. Taq DNA Polymerase having a bacteria-derived DNA content less than 10 fg/ μ L is preferable as the enzyme for PCR.

25 [0058]

The gene coding the enzyme for PCR used in the present invention may be any gene such as cDNA coding a heat-resistant DNA polymerase, genome DNA, synthetic DNA, etc., and may be a single-stranded chain or a double-stranded chain having its complementary strand, and may contain a natural or artificial nucleotide derivative. Further, when the heat-resistant DNA polymerase is derived from an organism, the derivation of the heat-resistant DNA polymerase is not particularly restricted.

[0059]

Specific examples of the heat-resistant, i.e., thermostable DNA polymerase used in the present invention include heat-resistant DNA polymerases derived from *Thermus aquaticus*, *Thermus thermophilus*, *Bacillus stearothermophilus*, *Thermococcus gorgonarius*, *Thermococcus kodakaraensis* KOD1, *Pyrococcus woesei*, *Pyrococcus furiosus*, *Aeropyrum pernix*, *Aquifex aeolicus*, *Sulfolobus tokodaii*, *Pyrolobus fumarii* or *Methanopyrus kandleri*.

[0060]

Further, the heat-resistant DNA polymerase may be a biogenetically artificially synthesized heat-resistant DNA polymerase.

[0061]

The enzyme for PCR used in the present invention is preferably an organism-derived heat-resistant DNA polymerase having heat resistance, more preferably one derived from a prokaryote such as methane bacteria, thermoacidophiles, thermophiles, hyperthermophiles, etc.

[0062]

(Production by eukaryote)

The enzyme for PCR used in the present invention is preferably an enzyme produced by using a eukaryote as a host.

[0063]

5 The eukaryotic cell includes fungi, animal cells, plant cells, insect cells, etc., and the host cell is not particularly restricted and may be a cell derived from a eukaryote.

[0064]

10 The fungi include ascomycetes such as yeasts, fungi, etc., filamentous bacteria, basidiomycetes, zygomycetes, etc., and of them, yeasts or filamentous bacteria are preferable. Specific examples thereof include Saccharomyces, Schizosaccharomyces, Candida, Pichia, Hansenula, Kluyveromyces, Zygosaccharomyces, Yarrowia, Trichosporon, Rhodosporidi, Aspergillus, Fusarium, Trichoderma, etc.

[0065]

15 Specifically, Saccharomyces cerevisiae, Schizosaccharomyces pombe, Candida utilis, Candida boidini, Pichia metanolica, Pichia angusta, Pichia pastoris, Pichia anomala, Hansenula polymorpha, Kluyveromyces lactis, Zygosaccharomyces rouxii, Yarrowia lipolytica, Trichosporon pullulans, Rhodosporidium toruloides, Aspergillus niger, Aspergillus nidulans, 20 Aspergillus awamori, Aspergillus oryzae, Trichoderma reesei, etc. are mentioned.

[0066]

25 The animal cells include human-derived cultured cells, mouse-derived cultured cells, etc., and specific examples thereof include CHO cells, Hela cells, etc. The plant cells may be cells derived from plants, are desirably established cultured cells and include Nicotiana cells, Arabidopsis cells,

Ipomoea cells, Daucus cells, Oryza cells, etc. Specific examples thereof include *Nicotiana tabacum* BY-2 cultured cells, *Arabidopsis thaliana* cultured cells, *Ipomoea batatas* cultured cells, *Daucus carota* cultured cells, *Oryza sativa* cultured cells, etc. The insect cells may be cells derived from insects, are desirably established cultured cells and include *Spodoptera frugiperda* ovary cell-derived cultured cell sf9 strain and sf21 strain, *Bombix mori* cultured cell Bm-N strain, etc. The host cells are preferably microorganisms or eukaryotes of quick multiplication such as yeasts, etc., and examples thereof include yeasts typically including *Saccharomyces* such as *Saccharomyces cerevisiae*, etc., plant cells typically including cultured cells of *Nicotiana* plants such as *Nicotiana tabacum*, etc., and filamentous bacteria typically including *Aspergillus* such as *Aspergillus oryzae*, etc.

[0067]

The production method is not particularly restricted, and production can be carried out appropriately according to, for example, a method described in Patent document 4.

[0068]

(High purification treatment)

Even general enzymes, which was produced by using bacteria such as *E. coli*, etc. as a host, can be used in the present invention provided that they were treated by a high purification treatment to physically remove host bacterium-derived DNA using a DNA degrading enzyme, an ion exchange resin, etc. and in which the amount of contamination of bacteria-derived DNA as a contamination source is less than 10 fg with respect to the enzyme preparation.

[0069]

(Treatment with selectively membrane-permeable dye)

By mixing the enzyme for PCR with EMA (ethidium monoazide) or PMA (propidium monoazide) as a selectively membrane-permeable dye and irradiating it with light, bacteria-derived DNA as a contamination source, particularly, killed bacteria-derived DNA can be inactivated. Since the inactivated DNA does not present amplification by PCR, only DNA derived from an intended bacterium to be detected present in a sample can be detected and the bacterial species can be identified. The treatment conditions comply with conditions recommended by a company of manufacturing and distributing the reagent and the reagent treating agent.

[0070]

<Other reagent, instrument, etc.>

The pH buffer solution is used for adjusting pH of a sample provided to PCR analysis to 7.0 to 10.0 (more preferably, pH 8.0 to 9.0). Specific examples thereof include a Tris-HCl buffer solution, etc. For example, the Tris-HCl buffer solution is used at a concentration of 5 mM to 100 mM.

[0071]

dNTP is a nucleoside source for amplification of DNA by PCR, and four kinds: dATP, dGTP, dCTP and dTTP are necessary. As the dNTP, one chemically modified for application in a hot start method, for example, CleanAmp™ dNTP manufactured by TriLink BioTechnologies, Inc. may be used. Regarding the use amount, dATP, dGTP, dCTP and dTTP are used each at a concentration of around 0.2 mM.

[0072]

In amplification of DNA by PCR, Mg^{2+} is necessary. The Mg^{2+} source includes $MgCl_2$, $MgSO_4$, etc. $MgCl_2$ is preferable and used appropriately in a use amount of 0.5 mM to 5.0 mM depending on investigation.

[0073]

5 The fluorochrome is used for the purpose of detecting a DNA amplification product by real time PCR, and further, for the purpose of measuring T_m of an amplified DNA fragment, and various known fluorochromes can be used. For example, a method of using an intercalator having a labeling function, a method of using a probe obtained by connecting
10 a fluorescent substance to a nucleotide hybridizing specifically to a DNA sequence to be amplified, etc., are mentioned. The intercalator includes ethidium bromide and SYBR Green I as unsaturated fluorochromes; and Resolight (manufactured by Roche), EvaGreen (manufactured by Biotim), etc., as saturated fluorochromes, etc. Preferable intercalators are SYBR
15 Green I as an unsaturated fluorochrome and EvaGreen and Resolight as a saturated fluorochrome, more preferable intercalators are EvaGreen and Resolight as a saturated fluorochrome. The use amount complies with recommendation of a company of producing and distributing the fluorochrome to be used.

20 [0074]

Additionally, genome DNA of a bacterium which is used as a positive control of PCR and sterile water which is used as a negative control may be contained in the kit.

As the vessel for PCR analysis and measurement used in the present
25 invention (for example, tube for analysis and measurement), those having a shape and a structure appropriate to a measuring machine can be selected

and used. Those recommended by a measuring machine distributing source can be used for measurement without particular problems.

[0075]

Other necessary instruments in carrying out the present invention
5 include instruments widely used in molecular biological experiments such as pipettes, pipette tips, 1.5 ml microtubes, etc., and apparatuses thereof include means widely used in molecular biological experiments such as PCR, a clean bench, a centrifugal machine for a tube, etc.

[0076]

10 <Contamination level>

The second point of the present invention is reduction of the level of contamination of bacteria-derived DNA as a contamination source in the whole sample solution prepared for PCR measurement including a primer used for detection and/or identification having the reduced level thereof,
15 which does not disturb detection of an extremely trace amount of a sample bacterium and identification of bacterial species.

[0077]

Patent document 4 describes that the amount of contamination of bacteria-derived DNA in a heat-resistant polymerase preparation solution is preferably 10 fg or less. As a result of investigation by the present inventors,
20 however, it was found that also bacterial contamination in reagents and instruments is an extremely important factor for detection of an extremely trace amount of a sample bacterium and identification of bacterial species. When commercially available reagents and instruments were used directly to
25 analysis, as they are, PCR amplifications was occasionally observed (Comparative Example 11).

[0078]

In the present invention, suppression of contamination of bacteria-derived DNA as a contamination source is necessary, not only for reagents to be used, but also for instruments and filtered water used for preparation of solutions for measurement. Regarding the instruments, the instruments are washed with water for washing capable of dissolving DNA, which has no DNA contamination in an operation space closing off DNA contamination from the environment, such as a clean room, etc. DNA contamination can be checked by analyzing the washing water after washing by a PCR method.

10 When appropriate, contamination of DNA may be checked by using this method.

[0079]

In the present invention, the amount of contamination of bacteria-derived DNA as a contamination source is 10 fg or less, preferably 5 fg or less, further preferably 1 fg or less with respect to the whole kit for identification. For respective reagents and instruments, the amount is lower than 10 fg, preferably lower than 5 fg, further preferably lower than 1 fg with respect to an individual preparation solution, depending on the embodiment of a kit described later. By suppressing the amount of mixing of bacteria-derived DNA as a contamination source to such a level, there are no emergence of unnecessary peaks and no deviation of the T_m value even in execution of amplification of 30 cycles or more in PCR analysis necessary for detection of an extremely trace amount of a sample bacterium and identification of bacterial species, thus, detection of bacterial species and

20 identification of bacterial species can be effected at high sensitivity with high precision.

25

[0080]

< Method of removing contaminant >

The method of suppressing contamination of bacteria-derived DNA as a contamination source in reagents and instruments used in the present invention includes an UV irradiation treatment, a gamma ray sterilization treatment, an ultrafiltration treatment, an EMA treatment, etc. When it is evident that the amount of mixing of bacteria-derived DNA as a contamination source is 10 fg or less by a receiving inspection, etc., the following treatments are not necessarily required.

10 [0081]

(Gamma ray sterilization treatment)

Reagents and instruments used in the present invention can be directly irradiated with a gamma ray to break bacteria-derived DNA as a contamination source. Gamma ray irradiation conditions vary depending on the kind of the material to be irradiated. In general, 5 kGy to 30 kGy, preferably 10 kGy to 25 kGy are applied.

[0082]

For specific fluorochromes such as Resolight, EvaGreen, SYBR Green I, etc., primers and instruments made of specific materials such as polypropylene, etc., however, a gamma ray sterilization treatment cannot be applied depending on conditions since they deteriorate by gamma ray irradiation. Example 15 described later shows the effect of a gamma ray sterilization treatment.

[0083]

25 (Ultrafiltration treatment)

Bacteria-derived DNA as a contamination source can be separated and removed, by treating reagents used in the present invention with ultrafiltration. Concretely, when bacteria-derived DNA is bacterial cells per se or genome, their molecular weights are so large. Thus, ultrafiltration of the reagents can be carried out by selecting an ultrafiltration membrane through which the reagents, etc., pass, but genomic DNA does not pass. As the ultrafiltration membrane, a membrane having a molecular weight cut off of 1 kDa to 1000 kDa, preferably 10 kDa to 100 kDa, more preferably 30 kDa to 50 kDa is preferably used. An ultrafiltration membrane having a molecular weight cut off of 10 kDa to 1000 kDa, preferably 30 kDa to 100 kDa, more preferably 30 kDa to 50 kDa is used for filtration of a primer(s). The material of the ultrafiltration membrane is not particularly restricted and includes regenerated cellulose, cellulose acetate, polyether sulfone, etc. As the pressure source for ultrafiltration, any of a vapor pressurization method, a centrifugal method, etc. may be used, and a preferable method is selected depending on the treating amount.

[0084]

When the treating amount is small, a centrifugal membrane filtration treatment is preferable. In the case of the centrifugal membrane filtration treatment, the amount of a treating solution and the centrifugal force (G) in a centrifugal operation comply with recommendation by a company of producing and distributing a filtration apparatus to be used.

[0085]

Example 13, Example 14, Comparative Example 10 and Comparative Example 11 described later show the effect of a centrifugal membrane filtration treatment.

[0086]

(Treatment with selectively membrane-permeable dye)

Reagents and instruments can be treated with EMA (ethidium monoazide) or PMA (propidium monoazide) as a selectively membrane-permeable dye and irradiated with light to inactivate bacteria-derived DNA as a contamination source, particularly, killed bacteria-derived DNA. Since inactivated DNA does not manifest multiplication by PCR, only DNA derived from an intended bacterium to be detected present in a sample can be detected and identification of bacterial species can be conducted.

10 [0087]

Treatment conditions comply with conditions recommended by a company of producing and distributing the reagents and the reagent treating agents. For example, it is disclosed for "Viable Bacteria Selection Kit for PCR" manufactured by TAKARA BIO Inc. that appended Solution A-gn (10 μ l) and Solution B-gn (5 μ l) are added to 40 μ l of a sample solution to be treated, the solution is allowed to stand still and reacted under light shielding for 5 to 15 minutes, then, irradiated with light for about 5 minutes. Example 16 described later shows the effect of an EMA treatment.

<Embodiment of kit>

20 The bacterial species identification kit of the present invention is a kit for identifying the bacterial species by analyzing a DNA fragment composed of a part of 16S rDNA sequence of a bacterium to be amplified by a primer set of the present invention. The analysis of a DNA fragment includes analysis of the molecular weight of a DNA fragment, measurement of the T_m value, etc. Measurement of the T_m value is preferable, and real time PCR is

25

suitably used for this purpose. The molecular weight can be analyzed by electrophoresis, a mass spectrometer, etc.

[0088]

As the primer pair composing a kit for detection of bacterial species,
5 the primer pairs classified into the following groups K1 to K7 were found to be suitable. Then, each one primer pair was selected from respective groups K1 to K7, and at least one, preferably at least four of thus obtained seven primer pairs were used to constitute bacterial species detection kit.

[0089]

10 (Primer pair for kit)

Group K1:

1-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 1 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

15 1-2: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 10 and a reverse primer consisting of any one of SEQ ID No. 16, 18, 20 and 21,

1-3: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 11 and a reverse primer consisting
20 of any one of SEQ ID No. 16 to 21, and 23 to 26,

1-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 12 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

1-5: a primer pair consisting of a combination of a forward primer
25 consisting of a sequence of SEQ ID No. 13 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

1-6: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 14 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

5 1-7: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 15 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

1-8: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 6 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

10 1-9: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 7 and a reverse primer consisting of any one of SEQ ID No. 16 to 27,

1-10: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 8 and a reverse primer consisting of
15 any one of SEQ ID No. 16 to 27, or

1-11: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 9 and a reverse primer consisting of any one of SEQ ID No. 16, 17, 19 and 26,

Group K2:

20 2-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 28 and a reverse primer consisting of any one of SEQ ID No. 44, 45, and 47 to 50,

2-2: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 29 and a reverse primer consisting
25 of any one of SEQ ID No. 44, 45, 47, 48 and 50,

2-3: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 30 and a reverse primer consisting of any one of SEQ ID No. 45, and 47 to 50,

5 2-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 31 and a reverse primer consisting of any one of SEQ ID No. 46 and 47,

2-5: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 32 and a reverse primer consisting of any one of SEQ ID No. 45, and 47 to 49,

10 2-6: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 33 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

15 2-7: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 34 and a reverse primer consisting of any one of SEQ ID No. 45 to 50,

2-8: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 35 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

20 2-9: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 36 and a reverse primer consisting of any one of SEQ ID No. 44 to 46, and 48 to 50,

2-10: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 37 and a reverse primer consisting of any one of SEQ ID No. 45 to 50,

2-11: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 38 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

2-12: a primer pair consisting of a combination of a forward primer
5 consisting of a sequence of SEQ ID No. 39 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

2-13: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 40 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

10 2-14: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 41 and a reverse primer consisting of any one of SEQ ID No. 44 to 50,

2-15: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 42 and a reverse primer consisting
15 of any one of SEQ ID No. 44 to 50, or

2-16: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 43 and a reverse primer consisting of any one of SEQ ID No. 44 to 48, and 50,

Group K3:

20 a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 51 and a reverse primer consisting of a sequence of SEQ ID No. 52,

Group K4:

25 4-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 53 and a reverse primer consisting of any one of SEQ ID No. 57 to 61,

4-2: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 54 and a reverse primer consisting of any one of SEQ ID No. 59 and 61,

4-3: a primer pair consisting of a combination of a forward primer
5 consisting of a sequence of SEQ ID No. 55 and a reverse primer consisting of any one of SEQ ID No. 57 to 59 and 61, or

4-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 56 and a reverse primer consisting of any one of SEQ ID No. 57 to 59, and 61,

10 Group K5:

5-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 62 and a reverse primer consisting of any one of SEQ ID No. 70, 71, and 73 to 75,

5-2: a primer pair consisting of a combination of a forward primer
15 consisting of a sequence of SEQ ID No. 63 and a reverse primer consisting of any one of SEQ ID No. 70 to 75,

5-3: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 64 and a reverse primer consisting of any one of SEQ ID No. 70 to 75,

20 5-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 65 and a reverse primer consisting of any one of SEQ ID No. 70 to 75,

5-5: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 66 and a reverse primer consisting
25 of any one of SEQ ID No. 70 to 75,

5-6: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 67 and a reverse primer consisting of any one of SEQ ID No. 70 to 75,

5 5-7: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 68 and a reverse primer consisting of any one of SEQ ID No. 70 to 75, or

5-8: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 69 and a reverse primer consisting of any one of SEQ ID No. 70 to 75,

10 Group K6:

6-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 76 and a reverse primer consisting of any one of SEQ ID No. 91, 93, 96, and 101 to 103,

15 6-2: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 77 and a reverse primer consisting of any one of SEQ ID No. 91, 96, and 101 to 103,

6-3: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 78 and a reverse primer consisting of any one of SEQ ID No. 91, and 101 to 103,

20 6-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 79 and a reverse primer consisting of any one of SEQ ID No. 91, 97, and 101 to 103,

25 6-5: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 80 and a reverse primer consisting of any one of SEQ ID No. 91, 93, and 100 to 103,

6-6: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 81 and a reverse primer consisting of any one of SEQ ID No. 91, 93, and 101 to 103,

5 6-7: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 82 and a reverse primer consisting of any one of SEQ ID No. 91, 96, 97, and 101 to 103,

6-8: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 83 and a reverse primer consisting of any one of SEQ ID No. 91, and 101 to 103,

10 6-9: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 84 and a reverse primer consisting of any one of SEQ ID No. 91, and 101 to 103,

6-10: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 85 and a reverse primer consisting
15 of any one of SEQ ID No. 91, and 101 to 103,

6-11: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 86 and a reverse primer consisting of any one of SEQ ID No. 91, 93, 97, and 101 to 103,

20 6-12: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 87 and a reverse primer consisting of any one of SEQ ID No. 91, 96, and 101 to 103,

6-13: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 88 and a reverse primer consisting of any one of SEQ ID No. 91, and 101 to 103,

6-14: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 89 and a reverse primer consisting of any one of SEQ ID No. 91, and 101 to 103, or

6-15: a primer pair consisting of a combination of a forward primer
5 consisting of a sequence of SEQ ID No. 90 and a reverse primer consisting of any one of SEQ ID No. 91, 92, 96, and 101 to 103,

Group K7:

7-1: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 104 and a reverse primer consisting
10 of any one of SEQ ID No. 2, 5, and 116 to 131,

7-2: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 105 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

7-3: a primer pair consisting of a combination of a forward primer
15 consisting of a sequence of SEQ ID No. 106 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

7-4: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 107 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

20 7-5: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 108 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

7-6: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 109 and a reverse primer consisting
25 of any one of SEQ ID No. 2, 5, and 116 to 131,

7-7: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 110 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

5 7-8: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 111 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

7-9: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 112 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

10 7-10: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 113 and a reverse primer consisting of any one of SEQ ID No. 2, 5, and 116 to 131,

7-11: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 114 and a reverse primer consisting
15 of any one of SEQ ID No. 2, 5 and 125, or

7-12: a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 115 and a reverse primer consisting of any one of SEQ ID No. 2, 5, 116, 118, 123, and 125 to 128.

[0090]

20 It is preferable that the above-described primer sets and kits contain at least one primer pair from the following seven primer pairs 1) to 7).

1) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 15 and a reverse primer consisting of a sequence of SEQ ID No. 16,

2) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 43 and a reverse primer consisting of a sequence of SEQ ID No. 50,

3) a primer pair consisting of a combination of a forward primer
5 consisting of a sequence of SEQ ID No. 51 and a reverse primer consisting of a sequence of SEQ ID No. 52,

4) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 53 and a reverse primer consisting of a sequence of SEQ ID No. 57,

10 5) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 62 and a reverse primer consisting of a sequence of SEQ ID No. 70,

6) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 81 and a reverse primer consisting
15 of a sequence of SEQ ID No. 103, and

7) a primer pair consisting of a combination of a forward primer consisting of a sequence of SEQ ID No. 115 and a reverse primer consisting of a sequence of SEQ ID No. 5.

[0091]

20 Further, the above-described kit containing as an essential component at least one of the above-described primer pairs 1) to 7) are preferable, and those containing at least one of the following combinations A to G are preferable for identification of bacterial species.

(A) A combination essentially containing the above-described primer
25 set 1) and further containing at least three primer pairs selected from the above-described 2) to 7).

(B) A combination essentially containing the above-described primer set 2) and further containing at least three primer pairs selected from the above-described 1) and 3) to 7).

5 (C) A combination essentially containing the above-described primer pair 3) and further containing at least three primer pairs selected from the above-described 1), 2) and 4) to 7).

(D) A combination essentially containing the above-described primer pair 4) and further containing at least three primer pairs selected from the above-described 1) to 3) and 5) to 7).

10 (E) A combination essentially containing the above-described primer pair 5) and further containing at least three primer pairs selected from the above-described 1) to 4) and 6) to 7).

(F) A combination essentially containing the above-described primer pair 6) and further containing at least three primer pairs selected from the
15 above-described 1) to 5) and 7).

(G) A combination essentially containing the above-described primer pair 7) and further containing at least three primer pairs selected from the above-described 1) to 6).

[0092]

20 The kit for detection and/or identification of bacterial species of the present invention has a constitution containing one or more out of the above-described primer pairs. In addition to this primer, if necessary, at least one reagent and/or part selected from reagents such as an enzyme for PCR, a pH buffer solution, $MgCl_2$, dNTP (or CleanAmp-dNTP), sterile water, etc. and
25 parts such as vessels for PCR analysis and measurement (for example, tube for analysis and measurement), other necessary instruments, etc. can be

contained in the kit. Further, in analysis in real time PCR, a fluorochrome can be added as a component. Depending on various embodiments of a PCR method, components necessary for a PCR method, in addition to these components, may be added to a kit. These reagents or parts may be packed
5 in any form such as an individual form, a partial form, an integrated form, etc. Specifically, the following forms are exemplified.

1) Reagents to be used are divided individually.

2) Reagents are divided into three components: a mixture obtained by previously mixing a pH buffer solution, $MgCl_2$, dNTP (or CleanAmp-dNTP),
10 and a fluorochrome in analysis by real time PCR; a primer pair; and an enzyme.

3) Reagents are divided into two components: a mixture obtained by previously mixing a pH buffer solution, $MgCl_2$, dNTP (or CleanAmp-dNTP), a fluorochrome in analysis by real time PCR and a primer pair; and an enzyme.

15 4) A pH buffer solution, $MgCl_2$, dNTP (or CleanAmp-dNTP), a fluorochrome in analysis by real time PCR, a primer and an enzyme are all previously mixed.

[0093]

In the above-described kit, the amount of mixing of bacteria-derived
20 DNA as a contamination source is 10 fg or less on the whole, the enzyme to be used is one having undergone production by an eukaryote, a high purification treatment or a treatment with a selectively membrane-permeable dye such as EMA or the like, and the reagents and instruments to be used (including tube for PCR analysis) are those individually or wholly having
25 undergone a gamma ray sterilization treatment, a centrifugal membrane filtration treatment or a treatment with a selectively membrane-permeable

dye, if needed. In packing reagents and parts, various treatments are so conducted that the amount of mixing of bacteria-derived DNA in a package after packing and sealing is 10 fg or less.

[0094]

5 <Application field, Sample>

The kit for detection and/or identification of bacterial species of the present invention can be used for detection of a bacterium and identification of bacterial species in various fields such as the medical field, food field, environmental analysis, etc. Particularly, in the medical field, the kit is useful
10 since it can detect an infecting bacterium in an infectious disease and/or identify bacterium species.

[0095]

The sample analyzed by the bacteria test is not particularly restricted, and a wide variety of samples can be applied. Specifically, in the medical
15 field, the sample includes blood, spinal fluid, lacrimal fluid, amnion fluid, other body fluids, materials adhered to medical devices such as catheters, etc. Blood, amnion fluid and spinal fluid are particularly useful for detection of a pathogenic bacterium of a serious infectious disease. In the food field, the sample includes foods themselves, liquid or solid samples in the course of
20 production, materials adhered to a machine in the production process, etc.

[0096]

<DNA extraction treatment>

In execution of detection of a bacterium in a sample and/or identification of bacterium species using the identification kit of the present
25 invention, it is necessary that DNA is previously extracted from a bacterium possibly present in a sample. As the DNA extraction method, an alkali

melting method, a boiling method, a phenol extraction method, etc. are currently known, and dedicated DNA extraction kits are marketed from various makers.

[0097]

- 5 The DNA extraction method in the present invention is not particularly restricted, and it is desirable to select a method proper for a subject sample since the suitable method varies depending on a sample. The suitably applicable means for DNA extraction includes QIAamp DNA Mini Kit manufactured by Qiagen and High pure PCR template preparation kit
- 10 manufactured by Roche used in the examples of the present application. As described in the manual of QIAamp DNA Mini Kit manufactured by Qiagen, the elution amount can be changed in a range of 50 μ l to 200 μ l according to necessity.

[0098]

- 15 <PCR analysis>

- The primer set or kit of the present invention is used in a PCR method. As the PCR method, various PCR methods for amplification of a target gene for detecting intended bacteria can be used. Real time PCR is a preferable analysis method, and a combination of real time PCR and analysis of a
- 20 melting curve for T_m measurement is more preferable. In this case, it is preferable that the performance of controlling temperature is $\pm 0.1^\circ\text{C}$ as the specification of a machine of real time PCR to be used. Specific examples of PCR and real time PCR and measurement of T_m using them are listed below.

- 25 [0099]

In PCR and real time PCR, apparatuses, methods, etc. which are usually known can be used.

[0100]

Conditions of thermal cycles in PCR (temperature, time, temperature elevating and dropping rate, cycle number) are not particularly restricted and may be appropriately set depending on the properties of a primer, an enzyme, a template, etc. to be used and the sensitivity of a DNA detection method after PCR. For setting such conditions, a lot of documents are already known.

10 [0101]

In PCR, a template double-stranded DNA thermal denaturation step, a primer annealing step and a DNA elongation step with an enzyme are repeated, in general. The thermal denaturation step may include temperature and time at which a template double-stranded DNA dissociates into single strands, and for example, a step of heating at 90°C to 98°C for several seconds to several minutes is set. In initiation of PCR, a process of thermal denaturation of several minutes to 10 minutes is often added only in the first cycle. The primer annealing step is set according to the base sequence of a primer and the number of bases, and a step of heating at 40°C to 72°C for several seconds to dozens of seconds is often set. In the DNA elongation step, the temperature complies with the property of the enzyme such as its optimum temperature, etc., and, for example, the range of 58°C to 76°C is general. Regarding the time of the DNA elongation step, the necessary time is roughly estimated and set from the chain length of DNA to be amplified and the DNA synthesis speed of an enzyme. The intended DNA is amplified by repeating steps of thermal denaturation, annealing and

elongation, and the number of this repetition may be appropriately altered depending on the amount of template DNA, the amount of an enzyme and the sensitivity of a DNA detection method after PCR, and in general, 10 to 50 times are exemplified. When the annealing temperature and the DNA
5 elongation temperature are approximately the same, it is also possible to carry out the both steps simultaneously.

[0102]

Also for real time PCR, conditions for thermal denaturation, annealing and DNA elongation necessary for DNA amplification, and further, the
10 numbers of repetitions of these steps, are the same as those for the above-described PCR. In real time PCR, the amount of amplified DNA can be quantified and estimated by measuring the fluorescent intensity derived from an intercalator and a probe, before and after the step of DNA elongation. The temperature for measuring fluorescent intensity can be appropriately
15 altered depending on the kind of a probe, etc. The temperature for measuring fluorescent intensity may be, for example, the temperature in DNA elongation as it stands in the case of execution using an intercalator, and when the chain length of intended DNA to be amplified is relatively long and its T_m value is relatively high, the temperature for measuring fluorescent
20 intensity may be set at temperatures between the T_m value of intended DNA and the T_m value of non-specifically amplified DNA typically including the intermediate value therebetween, by utilizing a fact that non-specifically amplified unintended DNA having relatively short chain length such as a primer dimer, etc. (also referred to as non-specifically amplified DNA) has
25 relatively low T_m value. By this, it is possible that, particularly in execution using an intercalator, only non-specifically amplified DNA such as a primer

dimer, etc. dissociates into single strands from a double-stranded structure, and the amount of DNA quantified or estimated from fluorescent intensity is a value for intended DNA.

[0103]

5 Further, in real time PCR, T_m of amplified DNA can be measured by analysis of a melting curve after completion of the DNA amplification step. In a melting curve analysis, dissociation of DNA from a double-stranded structure into single strands according to temperature change is observed, however, its temperature and detection conditions are not particularly
10 restricted. In general, stages of thermal denaturation (Denaturation) (90°C to 98°C), double strand formation (Annealing) (40°C to 80°C) and melting (Melting) (gradually rising from temperature of double strand formation up to around 98°C) are implemented and the change of fluorescent intensity in the melting step is monitored to obtain a melting curve, and the T_m value can be
15 obtained from this. Such measurement is possible in a lot of models of the real time PCR apparatus, and can be carried out according to device usage instructions.

[0104]

<Method of detection and/or identification>

20 As a method of detection or identification of bacterial species in a sample according to the present invention, the method comprising the following steps can be used.

(1) A step of conducting PCR using genome DNA of bacterium prepared from the above-described sample, a primer for obtaining an
25 amplification product containing an intended gene specific to the intended

bacterial species to be detected as a target, and a heat-resistant DNA polymerase.

- (2) A step of conducting detection or identification of the intended bacterial species to be detected in the above-described sample, by detection
5 of the above-described intended gene in the amplification product in the above-described PCR or analysis of the amplification product.

[0105]

- By using a primer set according to the present invention as the primer, identification of bacterial species at high sensitivity can be performed.
10 Further, by using reagents and instruments having controlled contamination level described above, further higher sensitivity and higher treating speed can be attained.

[0106]

- In this detection and/or identification method, the amplification step is
15 preferably conducted under suppression of amplification of genes other than the intended gene (unintended gene). For this suppression of amplification of unintended genes, PCR by a hot start method can be used. As an example, a hot start method using an anti-DNA polymerase antibody is mentioned. In this case, it is preferable to use an anti-DNA polymerase
20 antibody in an amount surplus to heat-resistant DNA polymerase 1U. Also, a hot start method in which reversible chemical modification is performed on a DNA polymerase as disclosed in Japanese Patent Laid-Open No. 2000-4847 (JP 2000 4847 A) and Japanese Patent Laid-Open No. 276776/1998 (JP H10 276776 A) can be suitably used. Further, a hot start method in which
25 chemically modified dNTP is used or a chemically modified primer is used as described in US Patent Publication No. 20070281308 can also be suitably

used. Furthermore, a hot start method in which a DNA polymerase and constituent components (for example, primer, dNTP, Mg^{2+} salt) indispensable for DNA amplification by a DNA polymerase are physically isolated using a wax melting by heating can also be suitably used.

5 [0107]

The step of detection and/or identification of a amplification product can be carried out by measuring T_m by real time PCR using an intercalator or a probe having a fluorescent label for detection. Preferable is real time PCR using an intercalator, and the intercalator includes, preferably,
10 unsaturated fluorochromes: SYBR Green I, saturated fluorochromes: EvaGreen and Resolight; more preferably, saturated fluorochromes: EvaGreen and Resolight.

[0108]

In the step of detection and/or identification by measurement of T_m
15 using real time PCR, the amplification product of the above-described intended gene can be made detectable and the amplification products of unintended genes other than this can be made undetectable. For this purpose, a method is preferable, which detects the amplification product of the intended gene by setting the conditions that detection of the amplification
20 product of the intended gene can be detected, while the amplification products of unintended genes other than the intended gene can be not detected, by:

(1) designing the above-described primer so that the melting temperature (T_m^A) of the amplification product of an intended gene is higher
25 than the melting temperature (T_m^B) of the amplification products of unintended genes, and

(2) performing detection of the amplification product at temperatures between T_m^A and T_m^B .

[0109]

Further, a method can be used in which the amplification step and the
5 step of detection and/or identification are conducted by real time PCR using a display indicating the amount of the amplification product, and the amplification products of unintended genes are hidden on the above-described display.

[0110]

10 The step of detection and/or identification can also be conducted by analysis of the amplification product in which the amplification product is developed by electrophoresis on a gel, etc. and visualized. The step of detection and/or identification can also be conducted by analysis of the amplification product in which the base sequence of the amplification product
15 is read. Further, this step can also be conducted by a method in which the molecular weight of the amplification product is measured and analyzed by a mass spectrometer.

[0111]

<Identification method>

20 For identifying the detected bacterium, the T_m value of a DNA fragment obtained from the detected bacterium using a primer set of the present invention can be utilized. In the algorithm for identification, a process can be added in which an influence of measurement errors such as, for example, a measuring error in every trial run of an apparatus is minimized by
25 effecting identification utilizing not only a combination of the T_m values

themselves described above but also a combination of differences between the T_m values.

[0112]

As the method for correcting the measuring error in every trial run of an apparatus described above, "the average value of a combination of T_m values is calculated, a combination of the relative values of respective T_m values from its average value" can be utilized. Namely, it is a method in which the configuration of a combination of T_m values is identified as "shape". "Shape" shown in two dimensions of the configuration of a combination of T_m values is not influenced by a measuring error. For example, a combination of T_m values (n (n is an integer of 4 or more and 7 or less)) specific to a detected bacterium are represented as T_{1db} to T_{ndb} (db is database), and the relative values from its average value are represented as d_{1db} to d_{ndb}, respectively. In the same way, a combination of T_m values (n (n is an integer of 4 or more and 7 or less)) of unknown intended organisms to be detected obtained from a sample are represented as T_{1ref} to T_{nref} (ref is reference), and the relative values from its average value are represented as d_{1ref} to d_{nref}, respectively. After comparison with the database, "one having a near combination of relative values = one in which "shape" of the configuration of a combination of T_m values is close" is utilized as the identification algorithm.

[0113]

Specific calculation methods include, but not limited to, for example, a method of calculating the distance between two points on the Euclidean space (formula 1).

[0114]

[Formula: 1]

$$\text{Dist.} = \sqrt{[(D1_{db} - D1_{ref})^2 + (D2_{db} - D2_{ref})^2 + \dots + (Dn_{db} - Dn_{ref})^2]}$$

In the calculation method by Formula 1, one in which the D value obtained by this calculation formula is nearest to 0 is identified as a required bacterium species to be detected. However, in view of the measuring error of a PCR apparatus to be used, the allowable range of the D value is 0 to 0.37, preferably 0 to 0.30 depending on the temperature control specification of the apparatus and the number of primers.

[0115]

10 The above-described algorithm can be used as a data base type identification software on a computer.

[0116]

<Identifiable bacterial species>

Regarding the identifiable microorganisms, any microorganisms corresponding to bacteria according to the taxonomic designation can be detected and identified according to the system.

Concrete bacterial species include: *Achromobacter denitrificans*, *Achromobacter xylosoxidans*, *Acinetobacter baumannii*, *Acinetobacter calcoaceticus*, *Actinomyces israelii*, *Aerococcus christensenii*, *Aeromonas hydrophila*, *Aeromonas sobria*, *Aggregatibacter actinomycetemcomitans*, *Alcaligenes faecalis*, *Alistipes onderdonkii*, *Anaerococcus vaginalis*, *Anaeroglobus geminatus*, *Arcanobacterium haemolyticum*, *Arcanobacterium pyogenes*, *Arthrobacter cummingsii*, *Atopobium vaginae*, *Bacillus anthracis*, *Bacillus cereus*, *Bacillus coagulans*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus sphaericus*, *Bacillus subtilis*, *Bacteroides dorei*, *Bacteroides finegoldii*,

- Bacteroides fragilis*, *Bacteroides nordii*, *Bacteroides salyersiae*,
Bacteroides thetaiotaomicron, *Bacteroides uniformis*, *Bacteroides*
vulgatus, *Bartonella henselae*, *Bartonella quintana*, *Bifidobacterium*
bifidum, *Bifidobacterium breve*, *Bilophila wadsworthia*, *Bordetella*
5 *pertussis*, *Borrelia burgdorferi*, *Borrelia recurrentis*, *Brevibacillus*
laterosporus, *Brucella abortus*, *Brucella melitensis*, *Brucella suis*,
Burkholderia cepacia, *Burkholderia mallei*, *Burkholderia pseudomallei*,
Campylobacter coli, *Campylobacter curvus*, *Campylobacter jejuni*,
Campylobacter rectus, *Capnocytophaga gingivalis*, *Capnocytophaga*
10 *granulosa*, *Capnocytophaga haemolytica*, *Capnocytophaga sputigena*,
Cardiobacterium hominis, *Chryseobacterium meningosepticum*, *Citrobacter*
amalonaticus, *Citrobacter freundii*, *Citrobacter koseri*, *Clostridium*
butyricum, *Clostridium difficile*, *Clostridium histolyticum*, *Clostridium*
hylemonae, *Clostridium paraputrificum*, *Clostridium perfringens*,
15 *Clostridium septicum*, *Clostridium sporogenes*, *Clostridium subterminale*,
Clostridium tertium, *Clostridium tetani*, *Corynebacterium amycolatum*,
Corynebacterium confusum, *Corynebacterium diphtheriae*,
Corynebacterium glucuronolyticum, *Corynebacterium jeikeium*,
Corynebacterium kroppenstedtii, *Corynebacterium macginleyi*,
20 *Corynebacterium minutissimum*, *Corynebacterium pseudodiphtheriticum*,
Corynebacterium pseudotuberculosis, *Corynebacterium riegelii*,
Corynebacterium tuberculostearicum, *Corynebacterium ulcerans*,
Corynebacterium xerosis, *Edwardsiella tarda*, *Eggerthella lenta*, *Eikenella*
corrodens, *Elizabethkingia meningoseptica*, *Empedobacter brevis*,
25 *Enterobacter aerogenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*,
Enterobacter sakazakii, *Enterococcus avium*, *Enterococcus bovis*,

- Enterococcus casseliflavus, Enterococcus cecorum, Enterococcus dispar,
 Enterococcus durans, Enterococcus faecium, Enterococcus flavescens,
 Enterococcus gallinarum, Enterococcus gilvus, Enterococcus hirae,
 Enterococcus italicus, Enterococcus malodoratus, Enterococcus mundtii,
 5 Enterococcus pallens, Enterococcus pseudoavium, Enterococcus
 raffinosus, Enterococcus sanguinicola, Erysipelothrix rhusiopathiae,
 Escherichia albertii, Escherichia coli, Eubacterium lentum, Eubacterium
 limosum, Finegoldia magna, Francisella tularensis, Fusobacterium
 necrophorum, Fusobacterium nucleatum, Fusobacterium periodonticum,
 10 Fusobacterium varium, Gardnerella vaginalis, Gemella morbillorum,
 Geobacillus stearothermophilus, Granulicatella adiacens, Haemophilus
 ducreyi, Haemophilus influenzae, Haemophilus parainfluenzae, Hafnia
 alvei, Halomonas venusta, Helicobacter cinaedi, Helicobacter pylori,
 Kingella kingae, Klebsiella granulomatis, Klebsiella oxytoca, Klebsiella
 15 pneumoniae, Lactobacillus acidophilus, Lactobacillus crispatus,
 Lactobacillus delbrueckii, Lactobacillus jensenii, Lactococcus garvieae,
 Legionella pneumophila, Leptospira interrogans, Listeria monocytogenes,
 Micrococcus luteus, Moraxella catarrhalis, Morganella morganii,
 Mycoplasma genitalium, Mycoplasma hominis, Neisseria gonorrhoeae,
 20 Neisseria meningitidis, Nocardia cyriacigeorgica, Odoribacter
 splanchnicus, Pantoea agglomerans, Parabacteroides distasonis,
 Parvimonas micra, Pasteurella multocida, Pediococcus damnosus,
 Peptoniphilus asaccharolyticus, Peptoniphilus gorbachii,
 Peptostreptococcus anaerobius, Plesiomonas shigelloides,
 25 Porphyromonas asaccharolytica, Porphyromonas gingivalis, Prevotella
 bivia, Prevotella bivia, Prevotella corporis, Prevotella intermedia,

- Prevotella melaninogenica, Prevotella nigrescens, Prevotella timonensis, Prevotella veroralis, Propionibacterium acnes, Propionibacterium avidum, Propionibacterium granulosum, Proteus mirabilis, Proteus vulgaris, Providencia rettgeri, Providencia stuartii, Pseudomonas aeruginosa,
- 5 Pseudomonas fluorescens, Pseudomonas putida, Rothia dentocariosa, Rothia mucilaginosa, Salmonella enterica, Serratia marcescens, Serratia plymuthica, Shigella boydii, Shigella dysenteriae, Shigella flexneri, Shigella sonnei, Spirillum minus, Staphylococcus aureus, Staphylococcus auricularis, Staphylococcus capitis, Staphylococcus caprae,
- 10 Staphylococcus carnosus, Staphylococcus cohnii, Staphylococcus epidermidis, Staphylococcus haemolyticus, Staphylococcus hominis, Staphylococcus lugdunensis, Staphylococcus pasteurii, Staphylococcus pettenkoferi, Staphylococcus pulvereri, Staphylococcus saccharolyticus, Staphylococcus saprophyticus, Staphylococcus schleiferi, Staphylococcus
- 15 simulans, Staphylococcus warneri, Staphylococcus xylosus, Stenotrophomonas maltophilia, Streptobacillus moniliformis, Streptococcus agalactiae, Streptococcus anginosus, Streptococcus bovis, Streptococcus canis, Streptococcus constellatus, Streptococcus dysgalactiae, Streptococcus equi, Streptococcus gallolyticus, Streptococcus gordonii,
- 20 Streptococcus infantarius, Streptococcus iniae, Streptococcus intermedius, Streptococcus lutetiensis, Streptococcus mitis, Streptococcus mutans, Streptococcus oralis, Streptococcus pasteurianus, Streptococcus pneumoniae, Streptococcus porcinus, Streptococcus pyogenes, Streptococcus salivarius, Streptococcus salivarius, Streptococcus
- 25 sanguinis, Streptococcus sobrinus, Streptococcus suis, Streptococcus vestibularis, Sutterella wadsworthensis, Treponema pallidum, Ureaplasma

parvum, *Vagococcus fluvialis*, *Veillonella atypica*, *Veillonella parvula*,
Vibrio alginolyticus, *Vibrio cholerae*, *Vibrio fluvialis*, *Vibrio*
parahaemolyticus, *Vibrio vulnificus*, *Yersinia enterocolitica*, *Yersinia*
pestis, *Yersinia pseudotuberculosis*. The identifiable microorganisms are
5 not limited to the above bacteria.

EXAMPLES

[0117]

The present invention will be illustrated specifically by examples and
10 test examples below, but the present invention is not limited to them.

[0118]

(Production Example 1) DNAP preparation method

e-DNAP was prepared by the following method according to
paragraphs 0195 to 0201 of Patent document 4, and used in examples
15 below.

(1) Synthesis of DNA

The entire DNA sequence of a heat-resistant DNA polymerase derived
from *T.aquaticus* was synthesized by GenScript. In this procedure, the
codon sequence was optimized to a yeast host, *S.cerevisiae*. The
20 synthesized DNA was incorporated into a plasmid pUC57 and supplied from
GenScript, and a vector pUC-TA01 was obtained. The gene coding the heat-
resistant DNA polymerase was so designed that Hind III added into the 5'
end sequence and an EcoRI restriction enzyme site added into the 3' end
sequence.

25 (2) Construction of vector for expression of *T. aquaticus*-derived heat-
resistant DNA polymerase

The gene coding the synthesized *T. aquaticus*-derived heat-resistant DNA polymerase was inserted into a plasmid pYES2 (Invitrogen), to construct a vector pYES-TA01. For the gene coding the heat-resistant DNA polymerase, pUC-TA01 was digested by restriction enzymes HindIII, EcoRI (TaKaRa Bio), electrophoresed on 1% agarose gel (Wako), and the gene coding the heat-resistant DNA polymerase was recovered by QIAquick gel extraction kit (Qiagen). The plasmid pYES2 was digested by EcoRI, NotI (TaKaRa Bio), and the gene coding the heat-resistant DNA polymerase and pYES2 were connected by DNA Ligation Kit Ver. 2.1 (TaKaRa Bio).

10 (3) Transformation of *S. cerevisiae*

PCR was conducted using the genome of *Saccharomyces cerevisiae* X2180 as a template and using a forward primer of SEQ ID No. 134 and a reverse primer of SEQ ID No. 135, to obtain a fragment A of about 550 bp. Likewise, PCR was conducted using the genome of *Saccharomyces*
 15 *cerevisiae* X2180 as a template and using a forward primer of SEQ ID No. 136 and a reverse primer of SEQ ID No. 137, to obtain a fragment B of about 550 bp. Next, PCR was conducted using the fragment A and the fragment B as a template and using a forward primer of SEQ ID No. 134 and a reverse primer of SEQ ID No. 137, to obtain a fragment AB. Next, *Saccharomyces*
 20 *cerevisiae* X2180 was cultured, then, a competent cell was made, and the fragment AB was transformed using FastTrack™-Yeast Transformation Kit (Geno Technology). The transformant was applied to the minimal agar medium in which 5-FOA was contained so as to give a final concentration of 1 mg/ml. This agar medium was cultured at 28°C for 3 days, then, a
 25 growable strain was obtained in the minimal agar medium containing 5-FOA, and named *Saccharomyces cerevisiae* X2180-S strain. The yeast

Saccharomyces cerevisiae X2180 strain is available from American Type Culture Collection as a bank for cells, bacteria and genes.

[0119]

This obtained vector pYES-TA01 was introduced into the yeast
5 (*Saccharomyces cerevisiae* X2180-S strain). As the host, the other yeast
can also be used providing it is a uracil auxotroph. For transformation,
FastTrack™-Yeast Transformation Kit (Geno Technology) was used.
(4) Production of *T. aquaticus*-derived heat-resistant DNA polymerase by
S.cerevisiae

10 In the following experiment, the instruments and ultrapure water used
were free of DNA. The resultant transformant was cultured at 28°C for 72
hours under vibration in 100 ml of an SD medium (0.67% Bacto yeast
nitrogen base, 2% Galactose). These were centrifugally separated at 5000
rpm for 10 minutes and harvested, and suspended in a disrupting buffer
15 solution (50 mM Tris-HCl pH7.5, 50 mM KCl), and bacterial bodies were
disrupted using 0.5 mm glass beads, then, centrifugally separated at 12000
rpm for 30 minutes, to obtain the yeast disrupting solution supernatant and a
precipitate as a cell extract. Next, the above-described cell extract was
thermally treated at 70°C for 60 minutes, and after the thermal treatment,
20 centrifugal separation was carried out at 5000 rpm for 30 minutes at 4°C and
the supernatant was recovered. To this supernatant, polyethyleneimine
(Sigma Aldrich) was added so as to give a final concentration of 0.1%, and
the mixture was stirred vigorously for 1 minute, then, allowed to stand at
room temperature for 5 minutes. Thereafter, the mixture was centrifugally
25 separated at 8000 rpm at 4°C for 30 minutes and the supernatant was

recovered, and this supernatant was allowed to pass through a filter of 0.45 mm.

[0120]

(5) Purification of *T. aquaticus*-derived heat-resistant DNA polymerase

5 In the following experiment, the instruments and ultrapure water used were free of DNA, and the experiment was conducted in a low temperature cabinet. The recovered coarse extracted solution was subjected to HiTrap Q FF (GE) equilibrated with a buffer solution A (50 mM Tris-HCl pH7.5), and thereafter, a 5% buffer solution B (50 mM Tris-HCl pH7.5, 1M NaCl) was
10 allowed to flow through a column to elute foreign substances. Next, a 15% buffer solution B was allowed to flow through a column to elute a heat-resistant DNA polymerase. The eluted enzyme solution was subjected to HiTrap Heparin equilibrated with a 15% buffer solution B, the content of the buffer solution B was increased gradually up to 50%, and an enzyme eluted
15 fraction was recovered. The buffer solution of this enzyme fraction was replaced by a preservation buffer solution (40 mM Tris-HCl pH8.0, 1M NaCl, 2 mM DTT, 0.2 mM EDTA), and further, a preservation solution (98% glycerol, 1% Tween20, 1% NP-40) was added in an equal amount and the mixture was stirred, then, preserved as an enzyme solution at -20°C.

20 (6) Measurement of activity of *T. aquaticus*-derived heat-resistant DNA polymerase

 First, a solution shown in Table 1 was prepared, and thermally insulated at 70°C for 5 minutes. Using this solution, a reaction solution was prepared as shown in Table 2, and the solution was reacted at 74°C for 5
25 minutes, and 50 µL of 10 mM EDTA was added to stop the reaction. After stopping of the reaction, 10 µL of SYBER Green I and 15 µL of sterile water

were added and the mixture was allowed to stand at room temperature for 5 minutes, then, fluorescence thereof was measured (Em: 497 nm, Ex: 528 nm). Activity of the heat-resistant DNA polymerase was defined based on the unit number of Thunder Taq (Nippon Gene Co., Ltd.) as a standard.

5 [0121]

[Table 1]

Table 1 Composition of reaction solution 1 for measurement of activity

1μg/μL M13mp18 DNA	28μL
10μM M13 primer	2.4μL
10mM Tris-HCl (pH8.8)	
50mM KCl	
0.08% Nonidet P-40	
2mM MgCl ₂	
DW	Appropriate amount
Total	140μL

[0122]

[Table 2]

Table 2 Composition of reaction solution 2 for measurement of activity

Solution of Table 14	2.4μL
2mM dNTPs	5μL
10mM Tris-HCl (pH8.8)	
50mM KCl	
0.08% Nonidet P-40	
2mM MgCl ₂	
Enzyme solution	1μL
DW	Appropriate amount
Total	25μL

10 [0123]

(Example 1) Bacterial identification by nested-PCR

In the present example, E. coli DNA was extracted from the culture solution of E. coli MG1655 using QIAamp DNA purification kit (QIAGEN), and

after extraction, the DNA amount was measured by an absorption spectrometer and diluted with ultrapure water, and EC1 and EC3 shown in Table 3, out of the diluted solutions, were used as a template of the reaction. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus. The instruments and ultrapure water used were free of DNA. The *E. coli* MG1655 strain is available from American Type Culture Collection as a bank for cells, bacteria and genes.

[0124]

[Table 3]

10

Table 3 *E. coli* genome solution used in experiment

Name	<i>E. coli</i> genome concentration (ng/mL)
EC1	10.0
EC2	1.0
EC3	0.1

[0125]

In conducting nested-PCR, a reaction solution having a formulation shown in Table 4 was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 40 times.

15

[0126]

[Table 4]

Table 4 Reaction solution composition 1

Template	2.0	μL
10X Thnuder Taq Buffer	2.0	μL
25mM MgCl ₂	1.6	μL
e-DNAP	1.0	Unit
2mM CleanAmp-dNTP	2.0	μL
EvaGreen (Biotium)	1.0	μL
10μM forward primer	0.6	μL
10μM reverse primer	0.6	μL
DW	Appropriate amount	
Total	20.0	μL

[0127]

- 5 As the primer, a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 2 were used. After completion of PCR, the reaction solution was recovered, and diluted 500-fold with DNA-free ultrapure water. The reaction was carried out using this diluted solution as a template, with a formulation shown in Table 5.

10 [0128]

[Table 5]

Table 5 Reaction solution composition 2

Template	2.0	μL
10X Thnuder Taq Buffer	2.0	μL
25mM MgCl ₂	2.0	μL
e-DNAP	1.0	Unit
2mM CleanAmp-dNTP	2.0	μL
EvaGreen (Biotium)	1.0	μL
10μM forward primer	0.6	μL
10μM reverse primer	0.6	μL
DW	Appropriate amount	
Total	20.0	μL

[0129]

Regarding the method of PCR, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times. For the primer, 7 sets including (1) a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16, (2) a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44, (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52, (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57, (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70, (6) a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and (7) a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2 were adopted, amplification of DNA thereof was performed, then, the DNA dissociation curve thereof was made and the T_m values were measured. In creation of the DNA dissociation curve, the reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise. At each stage, the reaction solution was incubated for 2 seconds, and the data were obtained. The resultant T_m value was compared with the T_m value of E. coli previously obtained and the D value was determined, as described in Patent document 4. When the D value is 0.3 or less, they were judged to be homogeneous. This result is shown in Table 6 together with the result of Comparative Example 1.

[0130]

(Comparative Example 1) (Direct-PCR) compared with Example 1 (nested-PCR)

The following comparative example is a replication experiment of Patent document 4. *E. coli* DNA was extracted from the culture solution of *E. coli* MG1655 using QIAamp DNA purification kit (QIAGEN), and after extraction, the DNA amount was measured by an absorption spectrometer and diluted with DNA free ultrapure water, and EC1 and EC3 shown in Table 3, out of the diluted solutions, were used as a template of the reaction. The reaction was conducted with a formulation shown in Table 5. The instruments and ultrapure water used were free of DNA. Regarding the method of PCR, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 20 seconds was repeated 35 times. For the primer, 7 sets including (1) a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16, (2) a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44, (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52, (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57, (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70, (6) a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and (7) a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2 were adopted, amplification of DNA thereof was performed, then, the DNA dissociation curve thereof was made and the T_m values were measured. The resultant T_m value was compared with the T_m value of *E. coli* previously obtained and the D value was determined, according to Example 7 of Patent document 4. This result is shown in Table 6 together with the result of Example 1. When the D value is 0.3 or less, they were judged to be homogeneous.

[0131]

[Table 6]

Table 6 Comparison between direct-PCR and nested-PCR

Template	Comparative example 1	Example 1
EC1	o	o
EC3	x	o

o: Identification was successful

x: Identification was a failure

[0132]

5 (Example 2) Verification of primer (Electrophoresis with purified coli genome)

- In the present example, EC3 shown in Table 3 was used as a template of the reaction. The instruments and ultrapure water used were free of DNA. For the primer combination, the reaction solution was so added as to give (1) respective combinations of a forward primer consisting of SEQ ID
- 10 No. 1 and reverse primers consisting of SEQ ID No. 2 to 5, (2) respective combinations of forward primers consisting of SEQ ID No. 1 and 6 to 15 and reverse primers consisting of SEQ ID No. 16 to 27, (3) respective combinations of forward primers consisting of SEQ ID No. 28 to 43 and reverse primers consisting of SEQ ID No. 44 to 50, (4) respective
- 15 combinations of forward primers consisting of SEQ ID No. 53 to 56 and reverse primers consisting of SEQ ID No. 57 to 61, (5) respective combinations of forward primers consisting of SEQ ID No. 62 to 69 and reverse primers consisting of SEQ ID No. 70 to 75, (6) respective combinations of forward primers consisting of SEQ ID No. 76 to 90 and
- 20 reverse primers consisting of SEQ ID No. 91 to 103, (7) respective combinations of forward primers consisting of SEQ ID No. 104 to 113 and

reverse primers consisting of SEQ ID No. 2 and 116 to 131, respective combinations of a forward primer consisting of SEQ ID No. 114 and reverse primers consisting of SEQ ID No. 2 and 5 and 125, respective combinations of a forward primer consisting of SEQ ID No. 115 and reverse primers
5 consisting of SEQ ID No. 2 and 5 and 125, and PCR thereof was performed. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0133]

10 For (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For (2) to (7) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a
15 process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times. The solution after the reaction was electrophoresed on 2% agarose gel and stained with ethidium bromide, then, the amplification amount of the intended product and amplification of the non-specific product were verified. Here, the combination
20 of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, the combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, the combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, the combination of a forward primer
25 consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, the combination of a forward primer consisting of SEQ ID No. 62 and a

reverse primer consisting of SEQ ID No. 70, the combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and the combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2 correspond to
 5 replication experiments of Patent document 4. The results are shown in Tables 7-1 to 7-7 together with the result of Comparative Example 2.
 [0134]

Evaluation in Table 7 was conducted according to the following criteria.

- 10 Amount of intended product:
 ◎ : amplification is particularly good
 ○: amplification is good
 Δ: amplification is poor
 ×: amplification is particularly poor
- 15 Non-specific amplification product
 ○: non-specific product is not found
 Δ: non-specific product is found
 ×: non-specific product is abundant

(Comparative Example 2) Comparative example for Example 2

- 20 The following comparative example is a replication experiment of Patent document 4. In the present comparative example, EC3 shown in Table 3 was used as a template of the reaction. The instruments and ultrapure water used were free of DNA. The reaction solution was so added as to give (1) a combination of a forward primer consisting of SEQ ID No. 1
 25 and a reverse primer consisting of SEQ ID No. 2, (2) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of

SEQ ID No. 16, (3) a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, (4) a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, (5) a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, (6) a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and (7) a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2, and PCR thereof was performed. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0135]

For (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For (2) to (7) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times. The solution after the reaction was electrophoresed on 2% agarose gel and stained with ethidium bromide, then, the amplification amount of the intended product and amplification of the non-specific product were verified. The results are shown in Tables 7-1 to 7-7 together with the result of Example 2.

[0136]

[Table 7-1]

Table 7-1

EC3	Electrophoresis : Amount of intended product				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	◎	○	○
	Electrophoresis : Non specificity				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	○	○	○

5

Table 7-2

Electrophoresis: Amount of intended product												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	○	×	×	△	×	△	⊗	×	△	×	×	×
SEQ ID No.6	⊗	×	×	⊗	⊗	△	×	△	△	△	⊗	⊗
SEQ ID No.7	⊗	⊗	⊗	⊗	⊗	⊗	×	○	○	×	⊗	⊗
SEQ ID No.8	⊗	⊗	⊗	⊗	⊗	⊗	△	○	△	△	△	△
SEQ ID No.9	△	△	×	△	×	×	×	×	×	×	△	×
SEQ ID No.10	△	×	△	×	△	△	×	×	×	×	×	×
SEQ ID No.11	△	△	△	△	△	△	×	×	×	×	△	×
SEQ ID No.12	×	○	○	○	○	○	×	△	△	×	⊗	○
SEQ ID No.13	⊗	⊗	⊗	⊗	⊗	△	△	⊗	○	×	⊗	⊗
SEQ ID No.14	△	△	△	△	△	△	×	×	△	△	○	○
SEQ ID No.15	⊗	⊗	⊗	⊗	⊗	⊗	×	⊗	⊗	⊗	⊗	⊗
Electrophoresis : Non specificity												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	○	×	×	○	×	○	○	×	○	×	×	×
SEQ ID No.6	○	×	×	○	○	○	×	○	○	○	○	○
SEQ ID No.7	○	○	○	○	○	○	×	○	○	×	○	○
SEQ ID No.8	○	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.9	△	○	×	○	×	×	×	×	×	×	○	×
SEQ ID No.10	○	×	○	×	○	○	×	×	×	×	×	×
SEQ ID No.11	○	○	○	○	○	○	×	×	×	×	○	×
SEQ ID No.12	×	○	○	○	○	○	×	○	○	×	○	○
SEQ ID No.13	○	○	○	○	○	○	○	○	○	×	○	○
SEQ ID No.14	○	○	○	○	○	○	×	×	○	○	○	○
SEQ ID No.15	○	○	○	○	○	○	×	○	○	○	○	○

EC3

[01371]
[Table 7-2]

[0138]

[Table 7-3]

Table 7-3

	Electrophoresis : Amount of intended product							
		SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
EC3	SEQ ID No.28	x	△	x	△	○	x	x
	SEQ ID No.29	△	△	x	△	○	x	x
	SEQ ID No.30	x	x	x	○	x	x	x
	SEQ ID No.31	x	x	△	○	x	x	x
	SEQ ID No.32	x	○	x	○	○	○	x
	SEQ ID No.33	○	⊙	x	○	○	x	x
	SEQ ID No.34	x	○	x	○	x	x	x
	SEQ ID No.35	△	x	x	x	○	○	x
	SEQ ID No.36	x	○	x	x	○	○	x
	SEQ ID No.37	x	⊙	x	○	○	x	△
	SEQ ID No.38	x	x	○	x	○	○	x
	SEQ ID No.39	⊙	x	⊙	x	○	x	x
	SEQ ID No.40	○	x	△	x	○	△	△
	SEQ ID No.41	x	x	○	x	x	x	x
	SEQ ID No.42	x	x	x	x	x	△	x
	SEQ ID No.43	x	x	x	x	x	x	x
	Electrophoresis : Non specificity							
		SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
	SEQ ID No.28	x	○	x	○	○	x	x
	SEQ ID No.29	○	○	x	○	○	x	x
	SEQ ID No.30	x	x	x	○	x	x	x
	SEQ ID No.31	x	x	○	○	x	x	x
	SEQ ID No.32	x	○	x	○	○	○	x
	SEQ ID No.33	○	△	x	○	○	x	x
	SEQ ID No.34	x	○	x	○	x	x	x
	SEQ ID No.35	○	x	x	x	○	○	x
	SEQ ID No.36	x	○	x	x	○	○	x
	SEQ ID No.37	x	○	x	○	○	x	○
	SEQ ID No.38	x	x	○	x	○	○	x
	SEQ ID No.39	○	x	○	x	○	x	x
	SEQ ID No.40	○	x	○	x	○	○	○
	SEQ ID No.41	x	x	○	x	x	x	x
	SEQ ID No.42	x	x	x	x	x	○	x
	SEQ ID No.43	x	x	x	x	x	x	x

[0139]

[Table 7-4]

Table 7-4

EC3	Electrophoresis : Amount of intended product					
		SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	×	△	○	×	◎
	SEQ ID No.54	×	×	△	×	△
	SEQ ID No.55	○	○	○	×	○
	SEQ ID No.56	○	○	○	×	◎
	Electrophoresis : Non specificity					
		SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	×	○	○	×	○
	SEQ ID No.54	×	×	○	×	○
	SEQ ID No.55	○	○	○	×	○
	SEQ ID No.56	○	○	○	×	○

EC3

Electrophoresis : Amount of intended product						
	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
SEQ ID No.62	○	×	-	○	○	○
SEQ ID No.63	◎	×	△	△	○	×
SEQ ID No.64	◎	×	×	×	○	×
SEQ ID No.65	◎	○	△	○	×	×
SEQ ID No.66	△	×	×	×	×	△
SEQ ID No.67	△	×	×	○	×	×
SEQ ID No.68	○	×	×	×	×	×
SEQ ID No.69	○	×	×	×	×	×

Electrophoresis : Non specificity						
	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
SEQ ID No.62	△	×	-	△	○	○
SEQ ID No.63	○	×	○	○	○	×
SEQ ID No.64	○	×	×	×	○	×
SEQ ID No.65	○	×	○	○	×	×
SEQ ID No.66	○	×	×	×	×	○
SEQ ID No.67	○	×	×	○	×	×
SEQ ID No.68	○	×	×	×	×	×
SEQ ID No.69	○	×	×	×	×	×

Table 7-5

Electrophoresis : Amount of intended product													
	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	△	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.77	○	x	x	x	x	x	x	x	x	x	△	△	○
SEQ ID No.78	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.79	○	x	x	x	x	x	x	x	x	x	○	○	x
SEQ ID No.80	○	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.81	○	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.82	⊗	x	x	x	x	x	x	x	x	x	⊗	⊗	○
SEQ ID No.83	○	x	x	x	x	x	x	x	x	x	⊗	⊗	x
SEQ ID No.84	○	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.85	○	x	x	x	x	x	x	x	x	x	⊗	⊗	○
SEQ ID No.86	○	x	x	x	x	x	x	x	x	x	⊗	⊗	○
SEQ ID No.87	○	x	x	x	x	x	x	x	x	x	⊗	⊗	○
SEQ ID No.88	○	x	x	x	x	x	x	x	x	x	⊗	⊗	x
SEQ ID No.89	⊗	x	x	x	x	x	x	x	x	x	x	⊗	x
SEQ ID No.90	○	x	x	x	x	x	x	x	x	x	○	○	○

Electrophoresis : Non specificity													
	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.77	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.78	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.79	○	x	x	x	x	x	x	x	x	x	△	○	x
SEQ ID No.80	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.81	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.82	○	x	x	x	x	x	x	x	x	x	○	△	○
SEQ ID No.83	○	x	x	x	x	x	x	x	x	x	○	○	x
SEQ ID No.84	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.85	○	x	x	x	x	x	x	x	x	x	○	△	△
SEQ ID No.86	○	x	x	x	x	x	x	x	x	x	△	○	△
SEQ ID No.87	○	x	x	x	x	x	x	x	x	x	○	○	△
SEQ ID No.88	○	x	x	x	x	x	x	x	x	x	○	x	x
SEQ ID No.89	○	x	x	x	x	x	x	x	x	x	x	○	x
SEQ ID No.90	○	x	x	x	x	x	x	x	x	x	○	○	○

EC3

[Table 7-5]
[0140]

5

Table 7-6

	Electrophoresis : Amount of intended product									
	SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123	
EC3	SEQ ID No.104	○	△	△	◎	◎	◎	△	◎	◎
	SEQ ID No.105	○	○	○	◎	◎	◎	◎	◎	◎
	SEQ ID No.106	×	△	×	○	○	○	○	△	○
	SEQ ID No.107	△	×	×	×	△	△	△	×	△
	SEQ ID No.108	○	×	×	○	○	○	○	×	○
	SEQ ID No.109	○	△	△	○	○	○	○	○	○
	SEQ ID No.110	×	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	△	△	○	○	○	○	○	○
	SEQ ID No.112	△	△	△	○	○	○	○	○	○
	SEQ ID No.113	△	△	△	×	○	○	○	○	○
	SEQ ID No.114	○								
	SEQ ID No.115	○								

[0141]
[Table 7-6]

EC3	Electrophoresis : Amount of intended product									
		SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
	SEQ ID No.104	◎	◎	◎	×	△	◎	△	△	◎
	SEQ ID No.105	◎	◎	◎	×	×	×	×	×	×
	SEQ ID No.106	△	○	○	×	△	△	○	◎	△
	SEQ ID No.107	△	△	△	×	△	△	△	△	×
	SEQ ID No.108	×	○	○	×	×	△	×	×	×
	SEQ ID No.109	○	○	○	◎	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○	○
	SEQ ID No.113	○	○	○	○	○	○	○	○	○
	SEQ ID No.114		○	○						
	SEQ ID No.115		×	○						

5

Table 7-7

	Electrophoresis	Non specific product								
		SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
EC3	SEQ ID No.104	○	○	○	○	○	○	△	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○	○
	SEQ ID No.106	x	○	x	○	○	○	○	○	○
	SEQ ID No.107	○	x	x	x	○	○	○	x	○
	SEQ ID No.108	○	x	x	○	○	○	○	x	○
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	x	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○	○
	SEQ ID No.113	○	○	○	x	x	○	○	○	○
	SEQ ID No.114	○								
	SEQ ID No.115	○								

[0142]
[Table 7-7]

	Electrophoresis	Non specific product								
		SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
EC3	SEQ ID No.104	○	○	○	x	○	○	○	○	○
	SEQ ID No.105	○	○	○	x	x	x	x	x	x
	SEQ ID No.106	○	○	○	x	○	○	○	○	○
	SEQ ID No.107	○	○	○	x	○	○	○	○	x
	SEQ ID No.108	x	○	○	x	x	○	x	x	x
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○	○
	SEQ ID No.113	○	x	○	○	○	○	○	○	○
	SEQ ID No.114		○	x						
	SEQ ID No.115		x	○						

[0143]

(Example 3) Verification of primer (PCR with purified coli genome and melting thereof)

In the present example, EC3 in Table 3 was used as a template. The
5 instruments and ultrapure water used were free of DNA. The reaction
solution was so added as to give (1) respective combinations of a forward
primer consisting of SEQ ID No. 1 and reverse primers consisting of SEQ ID
No. 2 to 5, (2) respective combinations of forward primers consisting of SEQ
ID No. 1 and 6 to 15 and reverse primers consisting of SEQ ID No. 16 to 27,
10 (3) respective combinations of forward primers consisting of SEQ ID No. 28
to 43 and reverse primers consisting of SEQ ID No. 44 to 50, (4) respective
combinations of forward primers consisting of SEQ ID No. 53 to 56 and
reverse primers consisting of SEQ ID No. 57 to 61, (5) respective
combinations of forward primers consisting of SEQ ID No. 62 to 69 and
15 reverse primers consisting of SEQ ID No. 70 to 75, (6) respective
combinations of forward primers consisting of SEQ ID No. 76 to 90 and
reverse primers consisting of SEQ ID No. 91 to 103, (7) respective
combinations of forward primers consisting of SEQ ID No. 104 to 113 and
reverse primers consisting of SEQ ID No. 2 and 116 to 131, respective
20 combinations of a forward primer consisting of SEQ ID No. 114 and reverse
primers consisting of SEQ ID No. 2 and 5 and 125 and respective
combinations of a forward primer consisting of SEQ ID No. 115 and reverse
primers consisting of SEQ ID No. 2 and 5 and 125, and PCR thereof was
performed. The reaction was conducted with a formulation shown in Table 4
25 for the above-described combination (1), and with a formulation shown in
Table 5 for the above-described combinations (2) to (7).

[0144]

Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the program thereof, for (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times, and for (2) to (7), the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. The reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds, and the data were obtained. The threshold cycle of the amplification curve and the dissociation curve were verified. Here, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2 correspond to

replication experiments of Patent document 4. The results are shown in Tables 8-1 to 8-7 together with the result of Comparative Example 3.

[0145]

Evaluation in Table 8 was carried out according to the following

5 criteria.

Threshold cycle:

⊙ : threshold cycle is fast (decreased by 2 cycles or more)

○: threshold cycle is somewhat fast (decreased by 0-1 cycle)

△: threshold cycle is somewhat slow (increased by 1-2 cycles)

10 ×: threshold cycle is slow (increased by 3 cycles or more)

Melting analysis

○: only one peak

△: small peaks

×: several peaks

15 -: no peak

(Comparative Example 3) Comparative example for Example 3

The following comparative example is a replication experiment of Patent document 4. In the present comparative example, EC3 shown in Table 3 was used as a template of the reaction. The instruments and

20 ultrapure water used were free of DNA, and the operation was conducted in a clean bench. The reaction solution was so added as to give (1) a

combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, (2) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No.

25 16, (3) a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, (4) a combination of a forward

primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, (5) a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, (6) a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and (7) a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2, and PCR thereof was conducted. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

10 [0146]

Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the program thereof, for (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times, and for (2) to (7), the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. In creation of the DNA dissociation curve, the reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise. At each stage, the reaction solution was incubated for 2 seconds, and the data were obtained. The threshold cycle of the amplification curve and the dissociation curve were verified. The results are shown in Tables 8-1 to 8-7 together with the result in Example 3.

25 [0147]

[Table 8-1]

Table 8-1

EC3	PCR: Threshold cycle				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	○	○	○
	PCR: Melting analysis				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	×	○	○	○

Table 8-2

PCR: Threshold cycle												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	○	-	-	-	-	-	△	-	○	-	-	-
SEQ ID No.6	○	-	-	△	△	⊗	-	△	△	-	⊗	⊗
SEQ ID No.7	○	○	○	⊗	⊗	○	-	△	△	-	○	○
SEQ ID No.8	⊗	○	△	○	○	○	-	○	○	-	⊗	⊗
SEQ ID No.9	○	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.10	-	-	-	-	○	×	-	-	-	-	-	-
SEQ ID No.11	⊗	○	×	-	-	-	-	-	-	-	△	-
SEQ ID No.12	-	⊗	⊗	⊗	⊗	○	△	△	△	×	⊗	⊗
SEQ ID No.13	⊗	⊗	⊗	⊗	⊗	⊗	-	⊗	△	-	△	△
SEQ ID No.14	-	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.15	⊗	⊗	⊗	⊗	⊗	⊗	-	⊗	⊗	⊗	⊗	⊗
PCR: Melting analysis												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	△	-	-	-	-	-	△	-	△	-	-	-
SEQ ID No.6	○	-	-	○	○	×	-	○	○	-	△	△
SEQ ID No.7	○	○	○	○	○	○	-	○	○	-	○	○
SEQ ID No.8	○	○	○	○	○	○	-	○	○	-	△	△
SEQ ID No.9	△	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.10	-	-	-	-	○	○	-	-	-	-	-	-
SEQ ID No.11	△	△	○	-	-	-	-	-	-	-	○	-
SEQ ID No.12	-	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.13	○	○	○	○	○	△	-	○	○	-	○	○
SEQ ID No.14	-	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.15	○	○	○	○	○	△	-	○	○	○	○	○

EC3

[Table 8-2]

[0148]

[0149]

[Table 8-3]

Table 8-3

EC3	PCR: Threshold cycle							
		SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
	SEQ ID No.28	-	○	-	○	○	○	○
	SEQ ID No.29	○	○	-	○	○	-	○
	SEQ ID No.30	-	⊙	-	○	○	⊙	⊙
	SEQ ID No.31	-	-	-	⊙	-	-	-
	SEQ ID No.32	-	○	-	○	○	○	-
	SEQ ID No.33	○	○	-	○	○	○	⊙
	SEQ ID No.34	-	⊙	-	○	⊙	-	-
	SEQ ID No.35	⊙	-	-	-	○	○	-
	SEQ ID No.36	-	○	-	-	○	⊙	-
	SEQ ID No.37	-	○	-	⊙	○	○	⊙
	SEQ ID No.38	-	○	○	○	○	-	○
	SEQ ID No.39	-	○	-	○	○	-	○
	SEQ ID No.40	-	○	○	○	○	-	○
	SEQ ID No.41	-	○	-	-	-	-	-
	SEQ ID No.42	-	-	-	-	○	-	-
	SEQ ID No.43	-	-	-	-	-	-	-
	PCR: Melting analysis							
		SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
	SEQ ID No.28	-	△	-	△	△	△	△
	SEQ ID No.29	△	○	-	○	△	-	△
	SEQ ID No.30	-	×	-	○	×	×	×
	SEQ ID No.31	-	-	-	-	-	-	-
	SEQ ID No.32	-	△	-	△	○	△	-
	SEQ ID No.33	○	×	-	×	×	×	×
	SEQ ID No.34	-	△	-	△	△	-	-
	SEQ ID No.35	△	-	-	-	○	△	-
	SEQ ID No.36	-	○	-	-	○	○	-
	SEQ ID No.37	-	○	-	○	○	△	△
	SEQ ID No.38	-	△	△	△	×	-	△
	SEQ ID No.39	-	○	-	○	△	-	×
	SEQ ID No.40	-	△	○	○	×	-	△
	SEQ ID No.41	-	○	-	-	-	-	-
	SEQ ID No.42	-	-	-	-	○	-	-
	SEQ ID No.43	-	-	-	-	-	-	-

[0150]

[Table 8-4]

Table 8-4

EC3	PCR: Threshold cycle					
	—	SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	-	○	○	○	○
	SEQ ID No.54	-	-	○	-	○
	SEQ ID No.55	○	○	○	-	○
	SEQ ID No.56	○	○	○	-	○
	PCR: Melting analysis					
	—	SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	-	○	○	△	○
	SEQ ID No.54	-	-	○	-	○
	SEQ ID No.55	○	○	○	-	○
	SEQ ID No.56	○	○	○	-	○

EC3	PCR: Threshold cycle						
	—	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
	SEQ ID No.62	-	-		○	-	-
	SEQ ID No.63	○	○	-	⊙	⊙	○
	SEQ ID No.64	-	-	-	-	-	○
	SEQ ID No.65	○	○	-	○	○	-
	SEQ ID No.66	○	-	-	○	-	-
	SEQ ID No.67	○	○	-	⊙	○	○
	SEQ ID No.68	○	○	○	○	○	○
	SEQ ID No.69	○	○	○	○	○	○
	PCR: Melting analysis						
	—	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
	SEQ ID No.62	-	-		○	-	-
	SEQ ID No.63	○	○	-	△	×	○
	SEQ ID No.64	-	-	-	-	-	○
	SEQ ID No.65	○	△	-	○	○	-
	SEQ ID No.66	○	-	-	×	-	-
	SEQ ID No.67	△	○	-	△	△	○
	SEQ ID No.68	×	○	○	△	○	○
	SEQ ID No.69	△	○	○	×	○	△

Table 8-5

PCR: Threshold cycle													
○	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	x	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.77	-	-	-	-	-	-	-	-	-	-	○	△	△
SEQ ID No.78	-	-	-	-	-	-	-	-	-	-	△	x	x
SEQ ID No.79	x	-	-	-	-	-	x	-	-	-	△	△	○
SEQ ID No.80	x	-	-	-	-	x	x	x	x	x	○	○	○
SEQ ID No.81	x	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.82	△	-	-	-	-	-	x	-	x	-	○	○	○
SEQ ID No.83	△	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.84	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.85	△	-	-	-	-	-	x	-	x	-	○	○	○
SEQ ID No.86	△	-	-	-	-	-	x	-	-	-	○	○	○
SEQ ID No.87	△	-	-	-	-	-	-	-	-	-	△	○	○
SEQ ID No.88	⊗	-	-	-	-	-	-	-	-	-	△	-	-
SEQ ID No.89	○	-	-	-	-	-	-	-	-	-	⊗	○	○
SEQ ID No.90	x	-	-	-	-	-	-	-	-	-	○	○	△

PCR: Melting analysis													
○	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.77	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.78	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.79	○	-	-	-	-	-	△	-	-	-	○	○	△
SEQ ID No.80	○	-	-	-	-	x	x	x	x	x	○	△	○
SEQ ID No.81	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.82	○	-	-	-	-	-	△	-	x	-	○	○	○
SEQ ID No.83	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.84	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.85	△	-	-	-	-	-	x	-	x	-	○	○	○
SEQ ID No.86	○	-	-	-	-	-	△	-	-	-	○	△	○
SEQ ID No.87	○	-	-	-	-	-	-	-	-	-	○	△	○
SEQ ID No.88	△	-	-	-	-	-	-	-	-	-	○	-	-
SEQ ID No.89	○	-	-	-	-	-	-	-	-	-	△	△	○
SEQ ID No.90	○	-	-	-	-	-	-	-	-	-	△	△	○

EC3

[0151]
[Table 8-5]

5

Table 8-6

EC3	PCR;Threshold cycle									
		SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
	SEQ ID No.104	○	○	○	○	○	○	○	○	⊗
	SEQ ID No.105	-	○	○	○	○	○	○	○	○
	SEQ ID No.106	⊗	○	○	○	○	○	○	○	○
	SEQ ID No.107	○	-	-	○	○	○	○	○	○
	SEQ ID No.108	○	○	-	○	○	○	○	○	○
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	-	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○	○
	SEQ ID No.113	○	-	-	-	○	○	○	○	○
SEQ ID No.114										
SEQ ID No.115										

PCR;Threshold cycle										
	SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131	
EC3	SEQ ID No.104	○	○	○	-	-	○	○	⊗	○
	SEQ ID No.105	○	○	○	-	-	-	-	-	-
	SEQ ID No.106	○	○	○	○	-	○	○	○	○
	SEQ ID No.107	○	○	○	-	-	-	-	-	-
	SEQ ID No.108	○	○	○	-	-	-	-	-	-
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	-	-	-	-	-	-
	SEQ ID No.112	○	○	○	-	-	-	-	-	-
	SEQ ID No.113	○	-	○	-	-	-	-	-	-
	SEQ ID No.114									
	SEQ ID No.115									

[0152]
[Table 8-6]

Table 8-7

Table 6-7

PCR: Melting analysis									
	SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
EC3	SEQ ID No.104	○	△	○	○	○	×	○	○
	SEQ ID No.105	-	○	△	○	○	○	△	○
	SEQ ID No.106	○	○	△	○	○	○	○	○
	SEQ ID No.107	○	-	-	×	○	○	○	○
	SEQ ID No.108	○	△	-	○	○	○	○	○
	SEQ ID No.109	○	×	×	○	○	△	○	○
	SEQ ID No.110	-	○	×	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○
	SEQ ID No.113	△	-	-	-	△	○	○	○
	SEQ ID No.114								
	SEQ ID No.115								

[0153]
[Table 8-7]

PCR: Melting analysis										
	SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131	
EC3	SEQ ID No.104	○	○	○	-	-	○	○	×	○
	SEQ ID No.105	○	○	○	-	-	-	-	-	-
	SEQ ID No.106	○	○	○	○	-	-	○	○	○
	SEQ ID No.107	○	○	○	-	-	-	-	-	-
	SEQ ID No.108	○	○	○	-	-	-	-	-	-
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	-	-	-	-	-	-
	SEQ ID No.112	○	○	○	-	-	-	-	-	-
	SEQ ID No.113	○	-	○	-	-	-	-	-	-
	SEQ ID No.114									
	SEQ ID No.115									

[0154]

(Example 4) Verification of primer (Electrophoresis with simulated DNA sample)

In the present example, a solution HE prepared by adding E. coli genomic DNA to 100 µg/ml human genomic DNA (clonotech) to reach a final concentration of 10 pg/ml was used as a template. The instruments and ultrapure water used were free of DNA. With respect to the primer combination, the reaction solution was so added as to give (1) respective combinations of a forward primer consisting of SEQ ID No. 1 and reverse primers consisting of SEQ ID No. 2 to 5, (2) respective combinations of forward primers consisting of SEQ ID No. 1 and 6 to 15 and reverse primers consisting of SEQ ID No. 16 to 27, (3) respective combinations of a forward primer consisting of SEQ ID No. 28 and reverse primers consisting of SEQ ID No. 44 and 45 and 47, a combination of a forward primer consisting of SEQ ID No. 30 and a reverse primer consisting of SEQ ID No. 48, a combination of a forward primer consisting of SEQ ID No. 31 and a reverse primer consisting of SEQ ID No. 47, a combination of a forward primer consisting of SEQ ID No. 32 and a reverse primer consisting of SEQ ID No. 48, respective combinations of forward primers consisting of SEQ ID No. 33 to 43 and reverse primers consisting of SEQ ID No. 44 to 50, (4) respective combinations of a forward primer consisting of SEQ ID No. 53 and reverse primers consisting of SEQ ID No. 57 and 61, respective combinations of a forward primer consisting of SEQ ID No. 54 and reverse primers consisting of SEQ ID No. 59 and 61, respective combinations of a forward primer consisting of SEQ ID No. 55 reverse primers consisting of SEQ ID No. 57 to 59 and 61, a combination of a forward primer consisting of SEQ ID No. 56

and a reverse primer consisting of SEQ ID No. 61, (5) respective combinations of a forward primer consisting of SEQ ID No. 62 and reverse primers consisting of SEQ ID No. 70 and 71 and 73 to 75, respective combinations of forward primers consisting of SEQ ID No. 63 to 69 and
5 reverse primers consisting of SEQ ID No. 70 to 75, (6) respective combinations of forward primers consisting of SEQ ID No. 76 to 90 and reverse primers consisting of SEQ ID No. 91 to 103 and (7) respective combinations of forward primers consisting of SEQ ID No. 104 to 113 and reverse primers consisting of SEQ ID No. 2 and 116 to 131, and PCR thereof
10 was conducted. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0155]

For (1) out of the above-described primer combinations, the reaction
15 solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For (2) to (7) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds
20 and at 72°C for 10 seconds was repeated 35 times. The solution after the reaction was electrophoresed on 2% agarose gel and stained with ethidium bromide, then, the amplification amount of the intended product and amplification of the non-specific product were verified. Here, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer
25 consisting of SEQ ID No. 2, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, a

combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2 correspond to replication experiments of Patent document 4. The results are shown in Tables 9-1 to 9-7 together with the result of Comparative Example 4.

[0156]

Evaluation in Table 9 was carried out according to the following criteria.

Amount of intended product:

15 ⊙ : amplification is particularly good

○: amplification is good

△: amplification is poor

×: amplification is particularly poor

Non-specific product amplification

20 ○: non-specific product is not found

△: non-specific product is found

×: non-specific product is abundant

(Comparative Example 4) Comparative example for Example 4

The following comparative example is a replication experiment of Patent document 4. In the present example, a solution HE prepared by adding E. coli genomic DNA to 100 µg /ml human genomic DNA (clonotech)

to reach a final concentration of 10 pg/ml was used as a template. The instruments and ultrapure water used were free of DNA, and the operation was conducted in a clean bench. The reaction solution was so added as to give (1) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, (2) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, (3) a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, (4) a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, (5) a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, (6) a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and (7) a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2, and PCR thereof was conducted. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0157]

For (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For (2) to (7) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times. The solution after the

reaction was electrophoresed on 2% agarose gel and stained with ethidium bromide, then, the amplification amount of the intended product and amplification of the non-specific product were verified. The results are shown in Tables 9-1 to 9-7 together with the result of Example 4.

5 [0158]

[Table 9-1]

Table 9-1

HE	Electrophoresis : Amount of intended product				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	○	○	○
	Electrophoresis : Non specificity				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	○	○	○

5

Table 9-2

Electrophoresis : Amount of Intended product												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	O	O	O	O	O	O	Δ	O	O	O	O	Δ
SEQ ID No.6	O	O	Δ	O	O	O	O	O	O	O	O	O
SEQ ID No.7	O	O	O	O	O	O	Δ	O	O	O	O	O
SEQ ID No.8	O	O	O	O	O	O	O	O	O	O	O	Δ
SEQ ID No.9	Δ	Δ	Δ	Δ	Δ	Δ	x	Δ	x	x	Δ	x
SEQ ID No.10	x	x	Δ	Δ	Δ	Δ	x	Δ	Δ	Δ	Δ	Δ
SEQ ID No.11	O	O	O	O	O	O	Δ	O	O	O	O	Δ
SEQ ID No.12	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗
SEQ ID No.13	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗
SEQ ID No.14	O	O	O	O	O	O	O	O	O	O	O	O
SEQ ID No.15	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗	⊗
Electrophoresis : Non specificity												
	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	O	Δ	Δ	O	O	O	Δ	O	O	O	O	Δ
SEQ ID No.6	O	O	O	O	O	O	Δ	O	O	O	O	Δ
SEQ ID No.7	Δ	Δ	Δ	Δ	Δ	O	O	O	O	O	O	Δ
SEQ ID No.8	Δ	Δ	Δ	Δ	Δ	Δ	Δ	Δ	Δ	Δ	O	O
SEQ ID No.9	x	x	x	x	x	x	x	x	x	x	x	x
SEQ ID No.10	x	x	x	x	x	x	x	x	x	x	x	x
SEQ ID No.11	Δ	Δ	Δ	Δ	Δ	Δ	x	Δ	Δ	Δ	Δ	x
SEQ ID No.12	O	O	O	O	Δ	O	O	O	O	O	O	x
SEQ ID No.13	O	O	O	O	O	O	O	O	O	O	O	x
SEQ ID No.14	O	O	O	O	O	O	O	Δ	O	O	O	Δ
SEQ ID No.15	O	O	O	Δ	O	O	O	O	O	O	Δ	Δ

HE

[0159]
[Table 9-2]

[0160]

[Table 9-3]

Table 9-3

	Electrophoresis : Amount of intended product						
	SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
SEQ ID No.28	○	△		○			
SEQ ID No.29							
SEQ ID No.30					○		
SEQ ID No.31				○			
SEQ ID No.32		×			○		
SEQ ID No.33	×	○	○	×	△	×	◎
SEQ ID No.34	×	×	○	△	○	○	○
SEQ ID No.35	△	◎	○	○	○	○	○
SEQ ID No.36	×	×	△	×	○	△	○
SEQ ID No.37	×	△	○	○	○	△	○
SEQ ID No.38	○	×	○	○	○	×	○
SEQ ID No.39	×	×	×	○	△	△	×
SEQ ID No.40	×	○	○	×	○	△	×
SEQ ID No.41	○	○	○	×	○	△	×
SEQ ID No.42	×	×	○	×	○	○	△
SEQ ID No.43	○	○	○	◎	◎	—	◎
HE	Electrophoresis : Non specificity						
	SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
SEQ ID No.28	○	×		△			
SEQ ID No.29							
SEQ ID No.30					△		
SEQ ID No.31				×			
SEQ ID No.32		×			△		
SEQ ID No.33	×	○	△	×	○	×	△
SEQ ID No.34	×	×	△	×	○	○	○
SEQ ID No.35	×	○	×	×	○	○	○
SEQ ID No.36	×	×	×	×	×	×	○
SEQ ID No.37	×	×	○	○	○	×	○
SEQ ID No.38	△	×	○	○	○	×	○
SEQ ID No.39	×	×	×	○	×	×	×
SEQ ID No.40	×	△	○	×	△	×	×
SEQ ID No.41	○	○	○	×	○	×	×
SEQ ID No.42	×	×	×	×	△	△	×
SEQ ID No.43	○	×	○	○	○	—	○

[0161]

[Table 9-4]

Table 9-4

Table 3

HE	Electrophoresis : Amount of intended product						
		SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61	
	SEQ ID No.53	x				⊙	
	SEQ ID No.54			x		⊙	
	SEQ ID No.55	⊙	⊙	⊙		⊙	
	SEQ ID No.56					x	
	Electrophoresis : Non specificity						
		SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61	
	SEQ ID No.53	x				○	
	SEQ ID No.54			x		○	
	SEQ ID No.55	○	○	○		○	
SEQ ID No.56					x		

HE

Electrophoresis : Amount of intended product						
	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
SEQ ID No.62	○	△	-	○	○	○
SEQ ID No.63	○	○	○	○	○	○
SEQ ID No.64	△	×	△	×	△	△
SEQ ID No.65	○	○	○	○	○	○
SEQ ID No.66	○	○	○	○	○	○
SEQ ID No.67	○	○	○	○	○	○
SEQ ID No.68	○	○	○	○	○	○
SEQ ID No.69	○	△	○	○	○	○

Electrophoresis : Non specificity						
	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
SEQ ID No.62	○	○	-	○	○	○
SEQ ID No.63	△	○	○	△	○	△
SEQ ID No.64	△	×	×	×	○	×
SEQ ID No.65	△	○	○	△	○	○
SEQ ID No.66	○	○	○	○	○	○
SEQ ID No.67	○	○	○	○	○	○
SEQ ID No.68	○	○	○	○	○	○
SEQ ID No.69	○	△	○	○	○	○

Table 9-5

Electrophoresis : Amount of intended product													
	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	x	x	x	x	x	○	x	x	x	x	x	x	x
SEQ ID No.77	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.78	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.79	○	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.80	○	x	x	x	x	x	x	x	x	○	○	○	○
SEQ ID No.81	⊗	x	△	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.82	○	x	x	x	x	⊗	○	x	x	x	⊗	⊗	⊗
SEQ ID No.83	○	x	x	x	x	x	x	x	x	x	⊗	⊗	○
SEQ ID No.84	△	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.85	x	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.86	⊗	x	△	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.87	⊗	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.88	○	x	x	x	x	x	x	x	x	x	⊗	△	△
SEQ ID No.89	○	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗
SEQ ID No.90	○	x	x	x	x	x	x	x	x	x	⊗	⊗	⊗

Electrophoresis : Non specificity													
	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	x	x	x	x	x	○	x	x	x	x	x	x	x
SEQ ID No.77	△	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.78	△	x	x	x	x	x	x	x	x	x	○	△	○
SEQ ID No.79	△	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.80	△	x	x	x	x	x	x	x	x	△	△	△	△
SEQ ID No.81	△	x	x	x	x	x	x	x	x	x	△	△	△
SEQ ID No.82	x	x	x	x	x	△	△	x	x	x	△	△	△
SEQ ID No.83	△	x	x	x	x	x	x	x	x	x	△	△	○
SEQ ID No.84	x	x	x	x	x	x	x	x	x	x	△	○	△
SEQ ID No.85	x	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.86	△	x	x	x	x	x	x	x	x	x	△	△	△
SEQ ID No.87	x	x	x	x	x	x	x	x	x	x	○	△	○
SEQ ID No.88	△	x	x	x	x	x	x	x	x	x	○	○	x
SEQ ID No.89	△	x	x	x	x	x	x	x	x	x	○	○	○
SEQ ID No.90	x	x	x	x	x	x	x	x	x	x	○	○	○

[Table 9-5]
[0162]

5

Table 9-6

Electrophoresis : Amount of intended product									
	SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
HE	SEQ ID No.104	○	×	○	○	○	○	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○
	SEQ ID No.106	△	△	△	△	△	△	△	○
	SEQ ID No.107	○	○	○	○	○	○	○	○
	SEQ ID No.108	○	○	△	○	○	△	△	○
	SEQ ID No.109	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○
	SEQ ID No.113	○	○	△	○	△	△	△	△
	SEQ ID No.114								
	SEQ ID No.115	○							

[Table 9-6]
(0163)

Electrophoresis : Amount of intended product									
	SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
HE	SEQ ID No.104	○	○	○	○	○	○	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○
	SEQ ID No.106	○	○	○	○	○	○	○	○
	SEQ ID No.107	○	○	○	○	○	○	○	○
	SEQ ID No.108	△	○	○	○	○	○	○	○
	SEQ ID No.109	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	×
	SEQ ID No.111	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○
	SEQ ID No.113	△	○	○	○	○	○	○	○
	SEQ ID No.114		○						
	SEQ ID No.115			○					

5

Table 9-7

	Electrophoresis	Non specific product								
		SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
HE	SEQ ID No.104	○	×	○	○	○	△	○	○	○
	SEQ ID No.105	○	△	○	○	○	△	△	△	△
	SEQ ID No.106	○	×	×	○	○	○	○	○	△
	SEQ ID No.107	○	○	○	○	○	△	△	△	○
	SEQ ID No.108	○	○	○	○	○	△	○	○	○
	SEQ ID No.109	○	○	×	×	×	×	○	○	○
	SEQ ID No.110	○	×	×	×	×	×	○	×	○
	SEQ ID No.111	○	×	×	×	×	×	×	×	○
	SEQ ID No.112	○	×	×	×	×	×	×	○	○
	SEQ ID No.113	○	×	×	○	×	×	×	×	×
	SEQ ID No.114									
	SEQ ID No.115	△								

[Table 9-7]
[0164]

	Electrophoresis	Non specific product								
		SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
HE	SEQ ID No.104	○	○	○	○	○	○	○	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○	○
	SEQ ID No.106	△	○	○	○	○	○	○	○	△
	SEQ ID No.107	△	○	○	○	○	○	○	○	○
	SEQ ID No.108	○	○	○	○	○	○	△	△	○
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	×
	SEQ ID No.111	×	○	○	△	△	△	△	△	△
	SEQ ID No.112	○	○	○	△	△	△	△	△	△
	SEQ ID No.113	×	○	○	△	△	△	△	△	△
	SEQ ID No.114		○							
	SEQ ID No.115			○						

[0165]

(Example 5) Verification of primer (PCR with simulated DNA sample and melting thereof)

In the present example, a solution HE prepared by adding E. coli genomic DNA to 100 µg /ml human genomic DNA (clonotech) to reach a final concentration of 10 pg/ml was used as a template. The instruments and ultrapure water used were free of DNA. With respect to the primer combination, the reaction solution was so added as to give (1) respective combinations of a forward primer consisting of SEQ ID No. 1 and reverse primers consisting of SEQ ID No. 2 to 5, (2) respective combinations of forward primers consisting of SEQ ID No. 1 and 6 to 15 and reverse primers consisting of SEQ ID No. 16 to 27, (3) respective combinations of a forward primer consisting of SEQ ID No. 28 and reverse primers consisting of SEQ ID No. 44 and 45 and 47, a combination of a forward primer consisting of SEQ ID No. 30 and a reverse primer consisting of SEQ ID No. 48, a combination of a forward primer consisting of SEQ ID No. 31 and a reverse primer consisting of SEQ ID No. 47, a combination of a forward primer consisting of SEQ ID No. 32 and a reverse primer consisting of SEQ ID No. 48, respective combinations of forward primers consisting of SEQ ID No. 33 to 43 and reverse primers consisting of SEQ ID No. 44 to 50, (4) respective combinations of a forward primer consisting of SEQ ID No. 53 and reverse primers consisting of SEQ ID No. 57 and 61, respective combinations of a forward primer consisting of SEQ ID No. 54 and reverse primers consisting of SEQ ID No. 59 and 61, respective combinations of a forward primer consisting of SEQ ID No. 55 and reverse primers consisting of SEQ ID No. 57 to 59 and 61, a combination of a forward primer consisting of SEQ ID No.

56 and a reverse primer consisting of SEQ ID No. 61, (5) respective combinations of a forward primer consisting of SEQ ID No. 62 and reverse primers consisting of SEQ ID No. 70 and 71 and 73 to 75, respective combinations of forward primers consisting of SEQ ID No. 63 to 69 and reverse primers consisting of SEQ ID No. 70 to 75, (6) respective combinations of forward primers consisting of SEQ ID No. 76 to 90 and reverse primers consisting of SEQ ID No. 91 to 103, (7) respective combinations of forward primers consisting of SEQ ID No. 104 to 113 and reverse primers consisting of SEQ ID No. 2 and 116 to 131, a combination of a forward primer consisting of SEQ ID No. 114 and a reverse primer consisting of SEQ ID No. 5 and respective combinations of a forward primer consisting of SEQ ID No. 115 and reverse primers consisting of SEQ ID No. 2 and 5, and PCR thereof was conducted. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0166]

Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the PCR program thereof, for (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times, and for (2) to (7), the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. The reaction

solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds, and the data were obtained. The threshold cycle of the amplification curve and the dissociation curve were verified. Here, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2 correspond to replication experiments of Patent document 4. The results are shown in Tables 10-1 to 10-7 together with the result of Comparative Example 5. [0167]

Evaluation in Table 10 was carried out according to the following criteria.

Threshold cycle:

- ⊙ : threshold cycle is fast (decreased by 2 cycles or more)
- : threshold cycle is somewhat fast (decreased by 0-1 cycle)
- △ : threshold cycle is somewhat slow (increased by 1-2 cycles)
- × : threshold cycle is slow (increased by 3 cycles or more)

Melting analysis:

o: only one peak

Δ: small peaks

×: several peaks

5 -: no peak

(Comparative Example 5) Comparative example for Example 5

The following comparative example is a replication experiment of Patent document 4. In the present example, a solution HE prepared by adding E. coli genomic DNA to 100 µg /ml human genomic DNA (clonetech) to reach a final concentration of 10 pg/ml was used as a template. The instruments and ultrapure water used were free of DNA. The reaction solution was so added as to give (1) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 2, (2) a combination of a forward primer consisting of SEQ ID No. 1 and a reverse primer consisting of SEQ ID No. 16, (3) a combination of a forward primer consisting of SEQ ID No. 28 and a reverse primer consisting of SEQ ID No. 44, (4) a combination of a forward primer consisting of SEQ ID No. 53 and a reverse primer consisting of SEQ ID No. 57, (5) a combination of a forward primer consisting of SEQ ID No. 62 and a reverse primer consisting of SEQ ID No. 70, (6) a combination of a forward primer consisting of SEQ ID No. 76 and a reverse primer consisting of SEQ ID No. 91 and (7) a combination of a forward primer consisting of SEQ ID No. 104 and a reverse primer consisting of SEQ ID No. 2, and PCR thereof was conducted. The reaction was conducted with a formulation shown in Table 4 for the above-described combination (1), and with a formulation shown in Table 5 for the above-described combinations (2) to (7).

[0168]

Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the PCR program thereof, for (1) out of the above-described primer combinations, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times, and for (2) to (7), the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. The reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds, and the data were obtained. The threshold cycle of the amplification curve and the dissociation curve were verified. The results are shown in Tables 10-1 to 10-7 together with the result of Example 5.

[0169]

[Table 10-1]

Table 10-1

HE	PCR: Melting analysis				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	⊙	⊙	○
	Electrophoresis : Non specificity				
		SEQ ID No. 2	SEQ ID No. 3	SEQ ID No. 4	SEQ ID No. 5
	SEQ ID No. 1	○	○	○	○

5

Table 10-2

HE

PCR: Threshold cycle												
○	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	○	○	○	○	○	○	*	△	△	△	○	⊗
SEQ ID No.6	○	○	○	○	○	○	-	△	△	△	⊗	△
SEQ ID No.7	○	○	○	○	○	○	*	○	○	△	○	⊗
SEQ ID No.8	○	○	○	○	○	○	-	○	△	△	○	⊗
SEQ ID No.9	-	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.10	-	-	-	-	-	-	-	-	-	*	-	-
SEQ ID No.11	x	-	-	x	x	x	-	x	x	-	x	-
SEQ ID No.12	△	△	△	△	△	△	*	△	△	*	△	○
SEQ ID No.13	○	○	○	△	○	○	*	○	△	△	○	○
SEQ ID No.14	○	○	○	○	○	○	*	△	△	△	△	△
SEQ ID No.15	⊗	⊗	○	○	○	○	*	○	○	○	○	⊗

PCR: Melting analysis

○	SEQ ID No.16	SEQ ID No.17	SEQ ID No.18	SEQ ID No.19	SEQ ID No.20	SEQ ID No.21	SEQ ID No.22	SEQ ID No.23	SEQ ID No.24	SEQ ID No.25	SEQ ID No.26	SEQ ID No.27
SEQ ID No.1	○	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.6	○	○	○	○	○	○	-	○	○	○	○	○
SEQ ID No.7	○	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.8	○	△	○	○	○	○	-	○	○	○	○	△
SEQ ID No.9	-	-	-	-	-	-	-	-	-	-	-	-
SEQ ID No.10	-	-	-	-	-	-	-	-	-	△	-	-
SEQ ID No.11	x	-	-	○	○	△	-	○	○	-	△	-
SEQ ID No.12	○	○	○	○	○	○	○	○	○	○	○	△
SEQ ID No.13	○	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.14	○	○	○	○	○	○	○	○	○	○	○	○
SEQ ID No.15	○	○	○	○	○	○	○	○	○	○	○	○

[Table 10-2]
[0170]

[0171]
[Table 10-3]

Table 10-3

HE	PCR: Threshold cycle							
	○	SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
	SEQ ID No.28	○	△		○			
	SEQ ID No.29							
	SEQ ID No.30					○		
	SEQ ID No.31				⊙			
	SEQ ID No.32		⊙			△		
	SEQ ID No.33	⊙	-	-	-	-	-	⊙
	SEQ ID No.34	-	-	○	-	○	-	○
	SEQ ID No.35	⊙	-	-	-	○	-	○
	SEQ ID No.36	⊙	○	○	-	-	-	○
	SEQ ID No.37	-	-	△	○	△	-	△
	SEQ ID No.38	⊙	-	△	△	○	-	○
	SEQ ID No.39	-	-	x	x	x	-	△
	SEQ ID No.40	-	-	-	-	○	-	△
	SEQ ID No.41	○	○	○	○	○	-	△
	SEQ ID No.42	○	○	△	△	○	-	△
	SEQ ID No.43	⊙	○	⊙	△	△		△
	PCR: Melting analysis							
	○	SEQ ID No.44	SEQ ID No.45	SEQ ID No.46	SEQ ID No.47	SEQ ID No.48	SEQ ID No.49	SEQ ID No.50
	SEQ ID No.28	○	△		○			
	SEQ ID No.29							
	SEQ ID No.30					○		
	SEQ ID No.31				x			
	SEQ ID No.32		x			○		
	SEQ ID No.33	x	-	-	-	-	-	○
	SEQ ID No.34	-	-	○	-	△	-	○
	SEQ ID No.35	x	-	-	-	x	-	○
	SEQ ID No.36	x	x	○	-	-	-	○
	SEQ ID No.37	-	-	○	x	△	-	○
	SEQ ID No.38	○	-	△	△	○	-	○
	SEQ ID No.39	-	-	△	x	△	-	○
	SEQ ID No.40	-	-	-	-	○	-	△
	SEQ ID No.41	○	x	○	x	○	-	○
	SEQ ID No.42	○	x	○	△	x	-	○
	SEQ ID No.43	△	○	△	○	○		○

[0172]

[Table 10-4]

Table 10-4

HE	PCR: Threshold cycle					
	—	SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	-				⊙
	SEQ ID No.54			-		⊙
	SEQ ID No.55	⊙	⊙	⊙		⊙
	SEQ ID No.56					-
	PCR: Melting analysis					
	—	SEQ ID No.57	SEQ ID No.58	SEQ ID No.59	SEQ ID No.60	SEQ ID No.61
	SEQ ID No.53	-				○
	SEQ ID No.54			-		○
	SEQ ID No.55	○	○	○		○
	SEQ ID No.56					-

HE	PCR: Threshold cycle						
	○	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
	SEQ ID No.62	○	△		○	⊙	○
	SEQ ID No.63	⊙	⊙	⊙	⊙	⊙	○
	SEQ ID No.64	⊙	⊙	⊙	⊙	⊙	○
	SEQ ID No.65	⊙	○	⊙	⊙	⊙	⊙
	SEQ ID No.66	⊙	○	⊙	⊙	○	○
	SEQ ID No.67	⊙	○	○	⊙	⊙	⊙
	SEQ ID No.68	○	○	○	○	○	⊙
	SEQ ID No.69	⊙	⊙	○	⊙	○	○
	PCR: Melting analysis						
	○	SEQ ID No.70	SEQ ID No.71	SEQ ID No.72	SEQ ID No.73	SEQ ID No.74	SEQ ID No.75
	SEQ ID No.62	○	○		○	○	○
	SEQ ID No.63	△	○	○	△	○	○
	SEQ ID No.64	×	○	×	△	△	×
	SEQ ID No.65	○	○	○	○	○	○
	SEQ ID No.66	○	○	○	○	○	○
	SEQ ID No.67	○	○	○	○	○	○
	SEQ ID No.68	○	○	○	○	○	○
	SEQ ID No.69	○	○	○	○	○	○

Table 10-5

PCR: Threshold cycle

○	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	-	-	Δ	-	-	x	x	-	-	-	-	-	-
SEQ ID No.77	Δ	x	x	-	-	x	x	x	x	x	○	Δ	Δ
SEQ ID No.78	x	-	x	-	-	-	-	x	x	-	Δ	Δ	Δ
SEQ ID No.79	-	-	-	-	-	-	-	-	-	-	Δ	Δ	Δ
SEQ ID No.80	Δ	-	○	-	-	-	-	-	-	-	○	○	○
SEQ ID No.81	x	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.82	x	-	-	-	-	-	-	-	-	-	x	Δ	x
SEQ ID No.83	-	-	-	-	-	-	-	-	-	-	Δ	Δ	Δ
SEQ ID No.84	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.85	-	-	-	-	-	-	-	-	-	-	Δ	Δ	Δ
SEQ ID No.86	Δ	-	-	-	-	-	-	-	-	-	Δ	Δ	Δ
SEQ ID No.87	Δ	-	-	-	-	x	-	-	-	-	○	○	Δ
SEQ ID No.88	Δ	-	-	-	-	-	-	-	-	-	Δ	○	-
SEQ ID No.89	Δ	-	-	-	-	-	-	-	-	-	Δ	Δ	Δ
SEQ ID No.90	x	x	-	-	-	x	-	x	x	x	x	x	x

HE

PCR: Melting analysis

○	SEQ ID No.91	SEQ ID No.92	SEQ ID No.93	SEQ ID No.94	SEQ ID No.95	SEQ ID No.96	SEQ ID No.97	SEQ ID No.98	SEQ ID No.99	SEQ ID No.100	SEQ ID No.101	SEQ ID No.102	SEQ ID No.103
SEQ ID No.76	-	-	x	-	-	○	x	-	-	-	-	-	-
SEQ ID No.77	○	x	x	-	-	Δ	x	x	x	x	○	○	○
SEQ ID No.78	Δ	-	x	-	-	-	-	x	x	-	○	○	Δ
SEQ ID No.79	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.80	x	-	x	-	-	-	-	-	-	-	○	○	○
SEQ ID No.81	x	-	-	-	-	-	-	-	-	-	○	Δ	○
SEQ ID No.82	x	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.83	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.84	-	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.85	-	-	-	-	-	-	-	-	-	-	○	○	Δ
SEQ ID No.86	x	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.87	x	-	-	-	-	○	-	-	-	-	○	Δ	○
SEQ ID No.88	x	-	-	-	-	-	-	-	-	-	○	Δ	-
SEQ ID No.89	○	-	-	-	-	-	-	-	-	-	○	○	○
SEQ ID No.90	Δ	○	-	-	-	○	-	x	x	x	○	Δ	○

HE

[Table 10-5]

[0173]

5

Table 10-6

PCR;Threshold cycle										
HE	○	SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
	SEQ ID No.104	△	-	x	○	○	○	○	○	○
	SEQ ID No.105	△	△	x	○	○	○	○	○	△
	SEQ ID No.106	○	○	△	○	○	⊗	⊗	○	○
	SEQ ID No.107	x	x	x	△	△	○	○	△	x
	SEQ ID No.108	x	x	x	△	△	○	△	△	○
	SEQ ID No.109	○	△	x	○	○	○	○	○	△
	SEQ ID No.110	△	△	x	○	△	○	○	△	△
	SEQ ID No.111	○	△	x	○	○	⊗	⊗	⊗	○
	SEQ ID No.112	○	○	△	○	⊗	⊗	⊗	○	○
	SEQ ID No.113	○	△	△	○	○	⊗	⊗	⊗	⊗
	SEQ ID No.114									
	SEQ ID No.115	○								

HE	PCR;Threshold cycle									
	○	SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
	SEQ ID No.104	○	○	○	△	○	△	△	△	△
	SEQ ID No.105	○	○	○	△	△	△	△	△	△
	SEQ ID No.106	○	○	○	○	○	○	○	○	○
	SEQ ID No.107	△	x	x	x	x	x	x	x	x
	SEQ ID No.108	x	△	x	x	x	x	x	x	x
	SEQ ID No.109	○	△	△	x	x	x	x	x	x
	SEQ ID No.110	○	△	△	○	○	○	○	○	-
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	△	△	△	△	△	△
	SEQ ID No.113	⊗	○	○	○	△	△	△	○	○
	SEQ ID No.114			○						
	SEQ ID No.115			○						

[0174]
[Table 10-6]

5

Table 10-7

PCR: Melting analysis										
HE	○	SEQ ID No.2	SEQ ID No.116	SEQ ID No.117	SEQ ID No.118	SEQ ID No.119	SEQ ID No.120	SEQ ID No.121	SEQ ID No.122	SEQ ID No.123
	SEQ ID No.104	○	-	○	○	○	○	○	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○	○
	SEQ ID No.106	○	○	○	○	○	○	○	○	○
	SEQ ID No.107	○	○	○	○	○	△	△	○	○
	SEQ ID No.108	○	○	○	○	○	△	○	○	△
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	○
	SEQ ID No.111	○	○	○	○	△	×	△	×	○
	SEQ ID No.112	○	○	×	○	○	○	○	○	○
	SEQ ID No.113	○	○	×	○	△	△	△	△	○
	SEQ ID No.114									
SEQ ID No.115	○	○	×	○	△	△	△	△	○	

PCR: Melting analysis										
HE	○	SEQ ID No.124	SEQ ID No.125	SEQ ID No.5	SEQ ID No.126	SEQ ID No.127	SEQ ID No.128	SEQ ID No.129	SEQ ID No.130	SEQ ID No.131
	SEQ ID No.104	○	○	○	○	○	○	○	○	○
	SEQ ID No.105	○	○	○	○	○	○	○	○	○
	SEQ ID No.106	○	○	○	○	○	○	○	○	○
	SEQ ID No.107	○	○	○	○	○	○	○	○	○
	SEQ ID No.108	○	○	○	○	○	○	○	○	○
	SEQ ID No.109	○	○	○	○	○	○	○	○	○
	SEQ ID No.110	○	○	○	○	○	○	○	○	-
	SEQ ID No.111	○	○	○	○	○	○	○	○	○
	SEQ ID No.112	○	○	○	○	○	○	○	○	○
	SEQ ID No.113	△	○	○	○	○	○	△	△	△
	SEQ ID No.114			○						
SEQ ID No.115	△	○	○	○	○	○	△	△	△	

[0175]
[Table 10-7]

[0176]

(Example 6) Comparison with Patent document 4 in direct-PCR

In the present example, E. coli DNA was extracted from the culture solution of E. coli MG1655 using QIAamp DNA purification kit (QIAGEN), and
5 after extraction, the DNA amount was measured by an absorption spectrometer, and EC1 and EC2 solutions shown in Table 3 diluted with ultrapure water were used as a template of the reaction. The instruments and ultrapure water used were free of DNA. The reaction was carried out with a formulation shown in Table 5. Rotor-Gene Q MDx 5plex HRM
10 (QIAGEN) was used as a real time PCR apparatus, and in the program thereof, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. The reaction
15 solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds and the data were obtained, and the T_m value of the amplification product was measured. Regarding the primer, PCR using the
20 following all seven sets was carried out:

(1) a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,

(2) a combination of a forward primer SEQ ID No. 43 and a reverse primer SEQ ID No. 50,

25 (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,

(4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,

(5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

5 (6) a combination of a forward primer SEQ ID No. 81 and a reverse primer SEQ ID No. 103 and

(7) a combination of a forward primer SEQ ID No. 115 and a reverse primer SEQ ID No. 5.

10 Then, the T_m value of the each resultant fragment was compared with the T_m value of *E. coli* previously obtained and the D value was determined, as described in Patent document 4. When the D value is 0.3 or less, they were judged to be homogeneous. This result is shown in Table 11 together with the result of Comparative Example 6.

[0177]

15 (Comparative Example 6) Primer part in Patent document 4 in Example 6

The following comparative example is a replication experiment of Patent document 4. In the present comparative example, *E. coli* DNA was extracted from the culture solution of *E. coli* MG1655 using QIAamp DNA purification kit (QIAGEN), and after extraction, the DNA amount was
20 measured by an absorption spectrometer, and EC1 and EC2 solutions shown in Table 3 diluted with ultrapure water were used as a template of the reaction. The reaction was carried out with a formulation shown in Table 5. The instruments and ultrapure water used were free of DNA. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in
25 the program thereof, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10

seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each amplification product was measured. Regarding the primer, PCR was carried out with the following all seven sets:

- 5 (1) a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16,
- (2) a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44,
- (3) a combination of a forward primer SEQ ID No. 51 and a reverse
10 primer SEQ ID No. 52,
- (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,
- 15 (6) a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and
- (7) a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2.

20 Then, the T_m value of the each resultant fragment was compared with the T_m value of E. coli previously obtained and the D value was determined, as described in Patent document 4. This result is shown in Table 11 together with the result of Example 6. When the D value is 0.3 or less, they were judged to be homogeneous.

[0178]

[Table 11]

Table 11 Comparison of Primers in direct-PCR

Template	Comparative Example 6	Example 6
EC1	○	○
EC2	X	×

○: Identification was successful.

×: Identification was a failure.

5 [0179]

(Example 7) Detection sensitivity with simulated sample and comparison of nested PCR with primer in Patent document 4

In the present example, an *E. coli* suspension was added to 2 ml of whole blood of a healthy individual as shown in Table 12 to prepare a

10 solution. Experiments were all conducted in clinical laboratory center of Toyama University Hospital (2630 Sugitani, Toyama city, Toyama prefecture, 930-0194, Japan).

[0180]

[Table 12]

Table 12 Concentration in model blood sample

Name	<i>E. coli</i> concentration (CFU/ml)
B1	2560
B2	50

15 [0181]

The solution prepared by adding an *E. coli* suspension to whole blood was centrifugally separated at 100 g for 5 minutes and the plasma fraction was recovered, then, the supernatant thereof was further centrifugally separated at 13000 g for 5 minutes, and the precipitate was recovered. DNA

solutions B1 and B2 extracted from this precipitate using QIAamp DNA purification kit (QIAGEN) were used as a template and PCR was carried out. The instruments and ultrapure water used were free of DNA. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in

5 the PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 40 times. For the primer, a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 5 was adopted and the reaction was conducted with a

10 formulation shown in Table 4. Next, this reaction product was diluted 500-fold with DNA-free ultrapure water, and further, this was used as a template and heated at 95°C for 5 minutes with a formulation shown in Table 5, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the DNA

15 dissociation curve of the amplification product was made. As the primer, the following all seven sets were used:

- (1) a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,
- (2) a combination of a forward primer SEQ ID No. 43 and a reverse
- 20 primer SEQ ID No. 50,
- (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,
- (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- 25 (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

(6) a combination of a forward primer SEQ ID No. 81 and SEQ ID No. 103 and

(7) a combination of a forward primer SEQ ID No. 115 and a reverse primer SEQ ID No. 5.

5 The resultant T_m value was compared with the T_m value of *E. coli* previously obtained and verified, as described in Patent document 4. The reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was
10 incubated for 2 seconds and the data were obtained, and the T_m value of the amplification product was measured. The results are shown in Table 13 together with the result of Comparative Example 7. When the D value is 0.3 or less, they were judged to be homogeneous.

[0182]

15 (Comparative Example 7) Primer part in Patent document 4 in Example 7

 The following comparative example is a replication experiment of Patent document 4. In the present comparative example, an *E. coli* suspension was added to 2 ml of whole blood of a healthy individual as shown in Table 12 to prepare a solution. Experiments were all conducted in
20 clinical laboratory center of Toyama University Hospital. The solution prepared by adding an *E. coli* suspension to whole blood was centrifugally separated at 100 g for 5 minutes and the plasma fraction was recovered, then, the supernatant thereof was further centrifugally separated at 13000 g for 5 minutes, and the precipitate was recovered. A DNA solution extracted
25 from this precipitate using QIAamp DNA purification kit (QIAGEN) was used as a template and PCR was carried out. The instruments and ultrapure

water used were free of DNA. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at

5 72°C for 30 seconds was repeated 40 times. For the primer, a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 2 was adopted and the reaction was conducted with a formulation shown in Table 1. Next, this reaction product was diluted 500-fold with DNA-free ultrapure water, and this was used as a template and heated at 95°C for 5 minutes

10 with a formulation shown in Table 2, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each amplification product was measured. Regarding the primer, the following all seven sets were used:

15 (1) a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16,

(2) a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44,

(3) a combination of a forward primer SEQ ID No. 51 and a reverse

20 primer SEQ ID No. 52,

(4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,

(5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

25 (6) a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and

(7) a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2.

The resultant T_m value was compared with the T_m value of *E. coli* previously obtained and verified, as described in Patent document 4. The results are shown in Table 13 together with the result of Example 7. When the D value is 0.3 or less, they were judged to be homogeneous.

[0183]

[Table 13]

Table 13 Investigation of identification sensitivity using model blood sample

Template	Comparative example 7	Example 7
B1	○	○
B2	X	○

○ : Identification was successful.

x: Identification was a failure.

[0184]

(Example 8) Verification of primer with multiple bacterial species (electrophoresis)

In the present example, genomes extracted from cultured bacteria of *Bacteroides flagilis*, *Campylobacter jejuni*, *Clostridium perfringens*, *Acinetobacter calcoaceticus*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Corynebacterium xerosis* and *Enterobacter cloacae* preserved in Toyama University (clinical laboratory center of Toyama University Hospital) using QIAamp DNA purification kit (QIAGEN) were used as a template of the reaction. The reaction was carried out with a formulation shown in Table 5.

The instruments and ultrapure water used were free of DNA. In the PCR program, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times. PCR was carried out for

5 the following primer combination:

- (1) a combination of SEQ ID No. 13 and SEQ ID No. 17,
- (2) a combination of SEQ ID No. 34 and SEQ ID No. 50,
- (3) a combination of SEQ ID No. 43 and SEQ ID No. 50,
- (4) a combination of SEQ ID No. 54 and SEQ ID No. 61,
- 10 (5) a combination of SEQ ID No. 55 and SEQ ID No. 58,
- (6) a combination of SEQ ID No. 55 and SEQ ID No. 61,
- (7) a combination of SEQ ID No. 68 and SEQ ID No. 72,
- (8) a combination of SEQ ID No. 77 and SEQ ID No. 103,
- (9) a combination of SEQ ID No. 81 and SEQ ID No. 103,
- 15 (10) a combination of SEQ ID No. 114 and SEQ ID No. 125 and
- (11) a combination of SEQ ID No. 115 and SEQ ID No. 5.

Then, the presence or absence of amplification of the intended fragment was verified. The results are shown in Tables 14-1 to 14-5 together with the result of Comparative Example 8.

20 [0185]

Bacterial strains of *Bacteroides flagilis*, *Campylobacter jejuni*, *Clostridium perfringes*, *Acinetobacter calcoaceticus*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Corynebacterium xerosis* and *Enterobacter*
 25 *cloacae* are available from RIKEN BioResource Center and Japan Culture Collection of Microorganisms.

[0186]

Evaluation in Table 14 was carried out according to the following criteria.

Amount of intended product:

5 ⊙ : amplification is particularly good

 ○: amplification is good

 Δ: amplification is poor

 ×: amplification is particularly poor

Amplification of non-specific product

10 ○: non-specific product is not found

 Δ: non-specific product is found

 ×: non-specific product is abundant

Threshold cycle:

 ⊙ : threshold cycle is fast (decreased by 2 cycles or more)

15 ○: threshold cycle is somewhat fast (decreased by 0-1 cycle)

 Δ: threshold cycle is somewhat slow (increased by 1-2 cycles)

 ×: threshold cycle is slow (increased by 3 cycles or more)

Melting analysis:

 ○: only one peak

20 Δ: small peaks

 ×: several peaks

 -: no peak

(Comparative Example 8) Verification of primer with multiple bacterial species (electrophoresis)

25 The following comparative example is a replication experiment of Patent document 4. In the present comparative example, genomes extracted

from cultured bacteria of *Bacteroides flagilis*, *Campylobacter jejuni*,
Clostridium perfringens, *Acinetobacter calcoaceticus*, *Klebsiella pneumoniae*,
Staphylococcus aureus, *Enterococcus faecalis*, *Bacillus cereus*,
Pseudomonas aeruginosa, *Corynebacterium xerosis* and *Enterobacter*
5 *cloacae* preserved in Toyama University (clinical laboratory center of Toyama
University Hospital) using QIAamp DNA purification kit (QIAGEN) were used
as a template of the reaction. The reaction was carried out with a formulation
shown in Table 5. The instruments and ultrapure water used were free of
DNA. In the PCR program, the reaction solution was heated at 95°C for 5
10 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C
for 10 seconds and at 72°C for 10 seconds was repeated 35 times.

[0187]

PCR was carried out for the following primers:

- (1) a combination of SEQ ID No. 1 and SEQ ID No. 16,
- 15 (2) a combination of SEQ ID No. 28 and SEQ ID No. 44,
- (3) a combination of SEQ ID No. 51 and SEQ ID No. 52,
- (4) a combination of SEQ ID No. 53 and SEQ ID No. 57,
- (5) a combination of SEQ ID No. 62 and SEQ ID No. 70,
- (6) a combination of SEQ ID No. 76 and SEQ ID No. 91 and
- 20 (7) a combination of SEQ ID No. 104 and SEQ ID No. 2.

Then, the presence or absence of amplification of the intended
fragment was verified. The results are shown in Table 14 together with the
result of Example 8.

[0188]

- 25 (Example 9) Verification of primer with multiple bacterial species (PCR)

In the present example, genomes extracted from cultured bacteria of *Bacteroides flagilis*, *Campylobacter jejuni*, *Clostridium perfringens*, *Acinetobacter calcoaceticus*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus cereus*, *Pseudomonas aeruginosa*,
5 *Corynebacterium xerosis* and *Enterobacter cloacae* preserved in Toyama University (clinical laboratory center of Toyama University Hospital) using QIAamp DNA purification kit (QIAGEN) were used as a template of the reaction. The reaction was carried out with a formulation shown in Table 5. The instruments and ultrapure water used were free of DNA. Rotor-Gene Q
10 MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the program thereof, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times, then, the DNA dissociation curve of the amplification product was made. The reaction
15 solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve.

PCR was carried out for the following primers:

- (1) a combination of SEQ ID No. 13 and SEQ ID No. 17,
- 20 (2) a combination of SEQ ID No. 34 and SEQ ID No. 50,
- (3) a combination of SEQ ID No. 43 and SEQ ID No. 50,
- (4) a combination of SEQ ID No. 54 and SEQ ID No. 61,
- (5) a combination of SEQ ID No. 55 and SEQ ID No. 58,
- (6) a combination of SEQ ID No. 55 and SEQ ID No. 61,
- 25 (7) a combination of SEQ ID No. 68 and SEQ ID No. 72,
- (8) a combination of SEQ ID No. 77 and SEQ ID No. 103,

(9) a combination of SEQ ID No. 81 and SEQ ID No. 103,

(10) a combination of SEQ ID No. 114 and SEQ ID No. 125 and

(11) a combination of SEQ ID No. 115 and SEQ ID No. 5.

Then, the DNA dissociation curve of the amplification product was
5 made. The reaction solution was heated at 95°C for 10 seconds, then,
incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise,
in order to make the DNA dissociation curve. At each stage, the reaction
solution was incubated for 2 seconds and the data were obtained, and the
T_m value of the amplification product was measured. Based on the results,
10 the threshold cycle of the amplification curve and the dissociation curve were
verified. The results are shown in Table 14 together with the result of
Comparative Example 5.

[0189]

(Comparative Example 9) Verification of primer with multiple bacterial
15 species. (PCR)

The following comparative example is a replication experiment of
Patent document 4. In the present comparative example, genomes extracted
from cultured bacteria of *Bacteroides flagilis*, *Campylobacter jejuni*,
Clostridium perfringens, *Acinetobacter calcoaceticus*, *Klebsiella pneumoniae*,
20 *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus cereus*,
Pseudomonas aeruginosa, *Corynebacterium xerosis* and *Enterobacter*
cloacae preserved in Toyama University (clinical laboratory center of Toyama
University Hospital) using QIAamp DNA purification kit (QIAGEN) were used
as a template of the reaction. The reaction was carried out with a formulation
25 shown in Table 5. The instruments and ultrapure water used were free of
DNA. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time

PCR apparatus, and in the program, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 35 times.

[0190]

- 5 For the primer, a combination of SEQ ID No. 1 and SEQ ID No. 16, a combination of SEQ ID No. 28 and SEQ ID No. 44, a combination of SEQ ID No. 51 and SEQ ID No. 52, a combination of SEQ ID No. 53 and SEQ ID No. 57, a combination of SEQ ID No. 62 and SEQ ID No. 70, a combination of SEQ ID No. 76 and SEQ ID No. 91 and a combination of SEQ ID No. 104
- 10 and SEQ ID No. 2 were used for PCR, and thereafter, the DNA dissociation curve of the amplification product was made. The reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds and
- 15 the data were obtained, and the T_m value of the amplification product was measured. Based on the results, the threshold cycle of the amplification curve and the dissociation curve were verified. The results are shown in Table 14 together with the result of Example 9.

Table 14-1

		<i>Bacteroides flagilis</i>				<i>Campylobacter jejuni</i>				<i>Clostridium perfringens</i>			
Forward	Reverse	Electrophoresis		PCR		Electrophoresis		PCR		Electrophoresis		PCR	
		Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting
		-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis
SEQ ID No.1	SEQ ID No.16	Δ	Δ	○	○	Δ	Δ	○	Δ	-	-	-	-
SEQ ID No.13	SEQ ID No.17	Δ	Δ	×	○	Δ	Δ	○	○	○	○	⊗	○
SEQ ID No.28	SEQ ID No.44	Δ	Δ	○	×	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.34	SEQ ID No.50	○	○	○	×	○	○	○	○	-	-	-	-
SEQ ID No.43	SEQ ID No.50	○	○	○	×	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.51	SEQ ID No.52	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.53	SEQ ID No.57	×	-	-	-	Δ	Δ	○	○	-	-	-	-
SEQ ID No.54	SEQ ID No.61	○	Δ	⊗	○	Δ	×	×	Δ	-	-	-	-
SEQ ID No.55	SEQ ID No.58	○	○	⊗	○	○	Δ	○	○	○	○	○	○
SEQ ID No.55	SEQ ID No.61	○	Δ	⊗	○	-	-	-	-	○	○	○	○
SEQ ID No.62	SEQ ID No.70	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.68	SEQ ID No.72	×	×	×	Δ	Δ	Δ	⊗	○	Δ	Δ	○	○
SEQ ID No.76	SEQ ID No.91	×	×	○	Δ	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.77	SEQ ID No.103	○	○	⊗	○	○	○	⊗	○	○	○	○	○
SEQ ID No.81	SEQ ID No.103	○	Δ	⊗	○	○	○	⊗	○	○	○	○	○
SEQ ID No.104	SEQ ID No.2	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.114	SEQ ID No.125	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.115	SEQ ID No.5	○	○	○	○	Δ	Δ	○	○	Δ	Δ	○	○

[0191]
[Table 14-1]

Table 14-2

		<i>Acinetobacter calcoaceticus</i>				<i>Klebsiella pneumoniae</i>				<i>Staphylococcus aureus</i>			
Forward	Reverse	Electrophoresis		PCR		Electrophoresis		PCR		Electrophoresis		PCR	
		Amplifi -cation	Non- specificity	Threshold cycle	Melting Analysis	Amplifi -cation	Non- specificity	Threshold cycle	Melting Analysis	Amplifi -cation	Non- specificity	Threshold cycle	Melting Analysis
SEQ ID No.1	SEQ ID No.16	Δ	Δ	0	0	-	-	-	-	-	-	-	-
SEQ ID No.13	SEQ ID No.17	Δ	Δ	0	0	0	0	0	0	0	Δ	0	0
SEQ ID No.28	SEQ ID No.44	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0
SEQ ID No.34	SEQ ID No.50	0	Δ	x	0	Δ	Δ	0	0	-	-	-	-
SEQ ID No.43	SEQ ID No.50	Δ	Δ	0	0	-	-	-	-	Δ	Δ	⊗	0
SEQ ID No.51	SEQ ID No.52	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0
SEQ ID No.53	SEQ ID No.57	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0
SEQ ID No.54	SEQ ID No.61	Δ	Δ	Δ	0	0	0	⊗	0	0	0	Δ	0
SEQ ID No.55	SEQ ID No.58	Δ	Δ	x	0	Δ	Δ	⊗	0	Δ	Δ	⊗	0
SEQ ID No.55	SEQ ID No.61	Δ	Δ	Δ	0	0	0	⊗	0	0	0	⊗	0
SEQ ID No.62	SEQ ID No.70	Δ	Δ	0	0	-	-	-	-	Δ	Δ	0	0
SEQ ID No.68	SEQ ID No.72	Δ	Δ	0	0	0	0	⊗	0	Δ	0	0	0
SEQ ID No.76	SEQ ID No.91	-	-	-	-	-	-	-	-	0	0	0	0
SEQ ID No.77	SEQ ID No.103	Δ	Δ	⊗	0	0	0	⊗	0	Δ	Δ	0	0
SEQ ID No.81	SEQ ID No.103	0	0	⊗	0	0	0	⊗	0	0	Δ	0	0
SEQ ID No.104	SEQ ID No.2	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0
SEQ ID No.114	SEQ ID No.125	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0
SEQ ID No.115	SEQ ID No.5	Δ	Δ	0	0	Δ	Δ	0	0	Δ	Δ	0	0

[0192]
[Table 14-2]

[0193]

[Table 14-3]

Table 14-3

		<i>Enterococcus faecalis</i>			
		Electrophoresis		PCR	
Forward	Reverse	Amplifi -cation	Non- specificity	Threshold cycle	Melting Analysis
SEQ ID No.1	SEQ ID No.16	△	△	○	○
SEQ ID No.13	SEQ ID No.17	○	○	△	○
SEQ ID No.28	SEQ ID No.44	△	△	○	○
SEQ ID No.34	SEQ ID No.50	△	△	△	○
SEQ ID No.43	SEQ ID No.50	△	△	○	○
SEQ ID No.51	SEQ ID No.52	△	△	○	○
SEQ ID No.53	SEQ ID No.57	△	△	○	○
SEQ ID No.54	SEQ ID No.61	△	△	○	○
SEQ ID No.55	SEQ ID No.58	△	△	○	○
SEQ ID No.55	SEQ ID No.61	△	△	○	○
SEQ ID No.62	SEQ ID No.70	△	△	○	○
SEQ ID No.68	SEQ ID No.72	△	△	○	○
SEQ ID No.76	SEQ ID No.91	△	△	○	○
SEQ ID No.77	SEQ ID No.103	△	△	○	○
SEQ ID No.81	SEQ ID No.103	△	△	○	○
SEQ ID No.104	SEQ ID No.2	△	×	○	○
SEQ ID No.114	SEQ ID No.125	○	○	○	○
SEQ ID No.115	SEQ ID No.5	○	○	○	○

5

10

Table 14-4

		<i>Bacillus cereus</i>				<i>Streptococcus agalactiae</i>				<i>Pseudomonas aeruginosa</i>			
Forward	Reverse	Electrophoresis		PCR		Electrophoresis		PCR		Electrophoresis		PCR	
		Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting
		-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis
SEQ ID No.1	SEQ ID No.16	Δ	Δ	○	○	Δ	Δ	○	○	×	×	○	○
SEQ ID No.13	SEQ ID No.17	Δ	Δ	Δ	○	Δ	Δ	×	○	○	○	Δ	○
SEQ ID No.28	SEQ ID No.44	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.34	SEQ ID No.50	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.43	SEQ ID No.50	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.51	SEQ ID No.52	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.53	SEQ ID No.57	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.54	SEQ ID No.61	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.55	SEQ ID No.58	Δ	Δ	○	○	Δ	Δ	○	○	-	-	-	-
SEQ ID No.55	SEQ ID No.61	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.62	SEQ ID No.70	-	-	-	-	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.68	SEQ ID No.72	○	○	⊗	○	Δ	Δ	Δ	○	Δ	Δ	Δ	○
SEQ ID No.76	SEQ ID No.91	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.77	SEQ ID No.103	Δ	Δ	⊗	○	Δ	Δ	○	○	○	○	⊗	○
SEQ ID No.81	SEQ ID No.103	Δ	Δ	⊗	○	Δ	Δ	Δ	○	○	○	⊗	○
SEQ ID No.104	SEQ ID No.2	Δ	Δ	○	○	Δ	Δ	○	○	×	×	○	○
SEQ ID No.114	SEQ ID No.125	○	○	Δ	○	○	○	○	○	○	○	○	○
SEQ ID No.115	SEQ ID No.5	○	○	Δ	○	○	○	○	○	Δ	Δ	○	○

[0194]
[Table 14-4]

Table 14-5

Forward	Reverse	<i>Corynebacterium xerosis</i>				<i>Enterobacter cloacae</i>				<i>Escherichia coli</i>			
		Electrophoresis		PCR		Electrophoresis		PCR		Electrophoresis		PCR	
		Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting	Amplifi	Non-	Threshold	Melting
		-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis	-cation	specificity	cycle	Analysis
SEQ ID No.1	SEQ ID No.16	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.13	SEQ ID No.17	Δ	Δ	Δ	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.28	SEQ ID No.44	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.34	SEQ ID No.50	○	○	×	○	Δ	Δ	⊗	○	Δ	Δ	Δ	○
SEQ ID No.43	SEQ ID No.50	○	○	○	○	Δ	Δ	⊗	○	Δ	Δ	○	○
SEQ ID No.51	SEQ ID No.52	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.53	SEQ ID No.57	Δ	Δ	○	○	Δ	×	○	○	Δ	Δ	○	○
SEQ ID No.54	SEQ ID No.61	○	○	⊗	○	○	○	Δ	○	○	○	⊗	○
SEQ ID No.55	SEQ ID No.58	Δ	Δ	⊗	○	○	○	Δ	○	○	○	○	○
SEQ ID No.55	SEQ ID No.61	○	○	⊗	○	○	○	Δ	○	-	-	-	-
SEQ ID No.62	SEQ ID No.70	Δ	Δ	○	○	Δ	Δ	○	Δ	Δ	Δ	○	○
SEQ ID No.68	SEQ ID No.72	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.76	SEQ ID No.91	Δ	Δ	○	○	Δ	Δ	○	Δ	Δ	Δ	○	○
SEQ ID No.77	SEQ ID No.103	○	○	×	○	Δ	Δ	○	Δ	Δ	Δ	○	○
SEQ ID No.81	SEQ ID No.103	○	○	×	○	Δ	Δ	○	Δ	Δ	Δ	○	○
SEQ ID No.104	SEQ ID No.2	Δ	Δ	○	○	Δ	Δ	○	○	Δ	Δ	○	○
SEQ ID No.114	SEQ ID No.125	Δ	Δ	○	○	Δ	Δ	○	○	○	○	○	○
SEQ ID No.115	SEQ ID No.5	Δ	Δ	○	○	Δ	Δ	○	○	○	○	○	○

[0195]
[Table 14-5]

[0196]

(Example 10) Verification with simulated blood sample of other bacterial species than *E. coli*

In the present example, cultured bacteria of *Bacillus cereus*,

- 5 *Pseudomonas aeruginosa*, *Enterobacter aerogenes* and *Streptococcus agalctiae* preserved in Toyama University (clinical laboratory center of Toyama University Hospital) were added to 2 ml of whole blood from a healthy individual so as to give concentrations shown in Table 15 to prepare solutions, being prepared along with the efforts of Toyama University
- 10 Hospital. Experiments were all conducted in clinical laboratory center of Toyama University Hospital. The solution prepared by adding a bacterium suspension to whole blood was centrifugally separated at 100 g for 5 minutes and the plasma fraction was recovered, then, the supernatant thereof was further centrifugally separated at 13000 g for 5 minutes, and the precipitate
- 15 was recovered. A DNA solution extracted from this precipitate using QIAamp DNA purification kit (QIAGEN) was used as a template and PCR was carried out in clinical laboratory center of Toyama University Hospital. The instruments and ultrapure water used were free of DNA. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a real time PCR apparatus, and in the
- 20 PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 40 times. For the primer, a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 5 was adopted. The reaction was conducted with a formulation
- 25 shown in Table 4. This reaction product was diluted 500-fold with DNA-free ultrapure water, and further, this was used as a template and heated at 95°C

for 5 minutes with a formulation shown in Table 5, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the DNA dissociation curve of the amplification product was made. The reaction solution was heated at 95°C
5 for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds and the data were obtained, and the T_m value of the amplification product was measured.

As the primer, the following seven sets were used:

- 10 a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,
- a combination of a forward primer SEQ ID No. 43 and a reverse primer SEQ ID No. 50,
- a combination of a forward primer SEQ ID No. 51 and a reverse
15 primer SEQ ID No. 52,
- a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,
- 20 a combination of a forward primer SEQ ID No. 81 and SEQ ID No. 103 and
- a combination of a forward primer SEQ ID No. 115 and a reverse primer SEQ ID No. 5.

Then, the resultant T_m value was compared with the T_m values of
25 various bacterial species previously obtained and verified, as described in Patent document 4.

The results are shown in Table 15.

[0197]

[Table 15]

Table 15 Concentrations of model blood samples prepared by using bacteria other than *E. coli* and results

Name	Species	CFU/ml	Identification
Ak19Ba	<i>Bacillus cereus</i>	5040	○
AK24St	<i>Streptococcus agalactiae</i>	548	○
AK24Ps	<i>Pseudomonas aeruginosa</i>	530	○
AEa24	<i>Enterobacter aerogenes</i>	262	○

Name	Species	CFU/ml	Identification
PA-Bc	<i>Bacillus cereus</i>	5040	○
PA-Sa	<i>Streptococcus agalactiae</i>	548	○
PA-Ps	<i>Pseudomonas aeruginosa</i>	530	○
PA-Ea	<i>Enterobacter aerogenes</i>	262	○

5 [0198]

(Example 11) Verification with clinical sample of Hospital of Toyama University

10 In the present example, a DNA solution extracted from a blood sample of a patient suspected of sepsis in Toyama University Hospital was used as a template, and experiments were all conducted in a laboratory of the examination department of Toyama University Hospital. The instruments and ultrapure water used were free of DNA. Rotor-Gene Q MDx 5plex HRM was used as a real time PCR apparatus, and in the PCR program thereof, first,

the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 40 times. For the primer, a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 5 was adopted. The reaction was conducted with a formulation shown in Table 4. This reaction product was diluted 500-fold with DNA-free ultrapure water, and further, this was used as a template and heated at 95°C for 5 minutes with a formulation shown in Table 5, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the DNA dissociation curve of the amplification product was made. The reaction solution was heated at 95°C for 10 seconds, then, incubated at 72°C for 90 seconds, and heated up to 95°C by 0.5°C stepwise, in order to make the DNA dissociation curve. At each stage, the reaction solution was incubated for 2 seconds and the data were obtained, and the T_m value of the amplification product was measured. As the primer, the following seven sets were used:

- a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,
- a combination of a forward primer SEQ ID No. 43 and a reverse primer SEQ ID No. 50,
- a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,
- a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

a combination of a forward primer SEQ ID No. 81 and SEQ ID No. 103
and

a combination of a forward primer SEQ ID No. 115 and a reverse
primer SEQ ID No. 5

- 5 Then, the resultant T_m value was compared with data in the data base
containing input T_m values of various bacterial species previously obtained
and verified, as described in Patent document 4. The results are shown in
Table 16.

Table 16 Identification results of clinical samples in Toyama University

	Results of Example 11	Blood culture method	Consistency of results
CS-1	<i>Bacteroides vulgatus</i>	<i>Bacteroides vulgatus</i>	○
CS-2	<i>Klebsiella pneumoniae</i>	<i>Klebsiella pneumoniae</i> , <i>Corynebacterium sp.</i>	○
CS-3	<i>Staphylococcus haemolyticus</i> or <i>aureus</i>	<i>Staphylococcus haemolyticus</i>	○
CS-4	<i>Citrobacter freundii</i>	<i>Citrobacter freundii</i>	○
CS-5	<i>Bacillus cereus</i>	<i>Bacillus cereus</i>	○
CS-6	<i>Staphylococcus capitis</i> or <i>epidermidis</i>	<i>Staphylococcus epidermidis</i>	○
CS-7	<i>Staphylococcus aureus</i>	<i>Staphylococcus aureus</i>	○
CS-8	<i>Klebsiella pneumoniae</i>	<i>Klebsiella pneumoniae</i>	○
CS-9	<i>Escherichia coli</i>	<i>Escherichia coli</i>	○
CS-10	<i>Enterobacter cloacae</i>	<i>Enterobacter cloacae</i>	○
CS-11	<i>Escherichia coli</i>	<i>Escherichia coli</i>	○

[0200]

(Example 12) Regarding hot start PCR

In the present example, a DNA solution (HE) was used as a template, and an amplification reaction was conducted using CleanAmp dNTPs (Sigma-Aldrich) as dNTP for hot start and using dNTP conventionally used, and the results were compared. The reaction was conducted with a formulation shown in Tables 4 and 17.

[0201]

[Table 17]

10

Table 17 Composition of Reaction solution 3

Template	2.0 μ L
10X Thnuder Taq Buffer	2.0 μ L
25mM MgCl ₂	1.6 μ L
e-DNAP	1.0 Unit
2mM dNTP	2.0 μ L
EvaGreen (Biotium)	1.0 μ L
10 μ M forward primer	0.6 μ L
10 μ M reverse primer	0.6 μ L
DW	Appropriate amount
Total	20.0 μ L

[0202]

The instruments and ultrapure water used were free of DNA. As the template, EC3 shown in Table 3 was used. For the primer, a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 2 was adopted, and in the PCR program, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. The solution after the reaction was electrophoresed on 2% agarose gel and

stained with ethidium bromide, then, the amplification of the intended product and amplification of the non-specific product were verified.

[0203]

The results are shown in Fig. 2.

5 [0204]

(Example 13) DNA removal treatment (ultrafiltration, E. coli genome addition experiment)

The instruments and ultrapure water used were free of DNA. Five hundred microliter of sterile water was added into Amicon UFC50508 (Millipore) and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Next, 500 μ L of 1 N hydrochloric acid was supplied and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Finally, 500 μ L of sterile water was supplied and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Such an operation was repeated twice. Next, 100 μ L of a mixed solution composed of components obtained after removal of the template and EvaGreen from a formulation of the reaction solution shown in Table 4 was prepared, and further, 4 pg of E. coli DNA was added. The resultant solution was put into Amicon UFC50508 after the above operations and centrifugal separation was performed at 5000 g for 5 minutes. Then, the filtrate was recovered. EvaGreen and sterile water were added to the resultant filtrate and PCR was conducted. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and in the PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating

at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For the primer, a combination of SEQ ID No. 1 and 2 was adopted. The product of this reaction solution was diluted 500-fold, and further, PCR was conducted with a formulation shown in Table 5 using this

5 diluted product as a template. The reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each amplification product was measured. PCR was carried out with

10 the following seven sets:

- a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16,
- a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44,
- 15 a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,
- a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- a combination of a forward primer SEQ ID No. 62 and a reverse
- 20 primer SEQ ID No. 70,
- a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and
- a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2.

Then, the presence or absence of amplification of each DNA sample was verified. The results are shown in Fig. 3 together with the result of Comparative Example 10.

[0205]

- 5 (Comparative Example 10) DNA removal treatment (ultrafiltration, negative control of *E. coli* genome addition experiment)

Five hundred microliter of sterile water was added into Amicon UFC50508 (Millipore) and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were
10 removed. Next, 500 μ L of 1 N hydrochloric acid was supplied and centrifugal separation was performed at 5000 g for 5 minutes, then remove the filtrate and the solution remaining in the filter. Finally, 500 μ L of sterile water was supplied and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed.
15 Such an operation was repeated twice. Next, 100 μ L of a mixed solution composed of components obtained after removal of the template and EvaGreen from a formulation of the reaction solution shown in Table 4 was prepared. The resultant solution was put into Amicon UFC50508 after the above operations and centrifugal separation was performed at 5000 g for 5
20 minutes. Then, the filtrate was recovered. EvaGreen and sterile water were added to the resultant filtrate and PCR was conducted. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and in the PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10
25 seconds and at 72°C for 30 seconds was repeated 35 times. For the primer, a combination of SEQ ID No. 1 and 2 was adopted. The product of this

reaction solution was diluted 500-fold, and further, PCR was conducted with a formulation shown in Table 5 using this diluted product as a template. The reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10
5 seconds was repeated 30 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each amplification product was measured. PCR was carried out with the following seven sets:

a combination of a forward primer SEQ ID No. 1 and a reverse primer
SEQ ID No. 16,

10 a combination of a forward primer SEQ ID No. 28 and a reverse
primer SEQ ID No. 44,

a combination of a forward primer SEQ ID No. 51 and a reverse
primer SEQ ID No. 52,

15 a combination of a forward primer SEQ ID No. 53 and a reverse
primer SEQ ID No. 57,

a combination of a forward primer SEQ ID No. 62 and a reverse
primer SEQ ID No. 70,

a combination of a forward primer SEQ ID No. 76 and a reverse
primer SEQ ID No. 91 and

20 a combination of a forward primer SEQ ID No. 104 and a reverse
primer SEQ ID No. 2.

Then, the presence or absence of amplification of each DNA sample was verified. The results are shown in Fig. 3 together with the result of
Example 13.

25 [0206]

(Example 14) DNA removal treatment (Purification of contaminated reagent, direct-PCR)

The instruments and ultrapure water used were free of DNA. Five hundred microliter of sterile water was added into Amicon UFC50508 (Millipore) and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Next, 500 μ L of 1 N hydrochloric acid was supplied and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Finally, 500 μ L of sterile water was supplied and centrifugal separation was performed at 5000 g for 5 minutes, and, then, the filtrate and the solution remaining in the filter were removed. Such an operation was repeated twice. Next, 100 μ L of a mixed solution composed of components obtained after removal of the template and EvaGreen from a formulation of the reaction solution shown in Table 4 was prepared. The resultant solution was put into Amicon UFC50508 after the above operations and centrifugal separation was performed at 5000 g for 5 minutes. Then, the filtrate was recovered. EvaGreen and sterile water were added to the resultant filtrate and PCR was conducted. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and in the program of PCR, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For the primer, a combination of SEQ ID No. 1 and 2 was adopted. The product of this reaction solution was diluted 500-fold, and further, PCR was conducted with a formulation shown in Table 5 using this diluted product as a template. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus,

and the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 26 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each
5 amplification product was measured. PCR was carried out with the following seven sets:

- (1) a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,
- (2) a combination of a forward primer SEQ ID No. 43 and a reverse
10 primer SEQ ID No. 50,
- (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,
- (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- 15 (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,
- (6) a combination of a forward primer SEQ ID No. 81 and a reverse primer SEQ ID No. 103 and
- (7) a combination of a forward primer SEQ ID No. 115 and a reverse
20 primer SEQ ID No. 5.

Then, the presence or absence of amplification of the intended fragment was verified. The results are shown in Fig. 4 together with the result of Comparative Example 11.

[0207]

- 25 (Comparative Example 11) DNA removal treatment (Purification of contaminated reagent, direct-PCR)

The instruments and ultrapure water used were free of DNA. A solution having a reaction formulation shown in Table 4 was prepared, and PCR was conducted. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and in the PCR program thereof, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 35 times. For the primer, a combination of SEQ ID No. 1 and 2 was adopted. The product of this reaction solution was diluted 500-fold, and further, PCR was conducted with a formulation shown in Table 5 using this diluted product as a template. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 26 times, then, the DNA dissociation curve of the amplification product was made, and the T_m value of each amplification product was measured. PCR was carried out with the following seven sets:

- (1) a combination of a forward primer SEQ ID No. 15 and a reverse primer SEQ ID No. 16,
- (2) a combination of a forward primer SEQ ID No. 43 and a reverse primer SEQ ID No. 50,
- (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,
- (4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,
- (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

(6) a combination of a forward primer SEQ ID No. 81 and a reverse primer SEQ ID No. 103 and

(7) a combination of a forward primer SEQ ID No. 115 and a reverse primer SEQ ID No. 5.

5 Then, the presence or absence of amplification of the intended fragment was verified. The results are shown in Fig. 4 together with the result of Example 14.

[0208]

(Example 15) DNA removal treatment (gamma ray sterilization, E. coli DNA is
10 treated with gamma ray)

 The instruments and ultrapure water used were free of DNA. A solution was prepared by adding 500 fg of E. coli DNA to 100 μ L of sterile water, and this was irradiated with a gamma ray of 10 kGy and 25 kGy. This solution was used as a template and PCR was conducted under the following
15 conditions. The reaction was carried out with a formulation shown in Table 1. Rotor-Gene Q MDx 5plex HRM (QIAGEN) was used as a PCR apparatus, and in the program of PCR, first, the reaction solution was heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 65°C for 10 seconds and at 72°C for 30 seconds was repeated 40 times, then, the
20 DNA dissociation curve was made. For the primer, a combination of SEQ ID No. 1 and 2 was adopted. The product of this reaction solution was diluted 500-fold, and further, this was used as a template and heated at 95°C for 5 minutes, then, a process including heating at 94°C for 10 seconds, at 60°C for 10 seconds and at 72°C for 10 seconds was repeated 30 times, then, the
25 DNA dissociation curve of the amplification product was made. PCR was carried out with the following seven sets:

(1) a combination of a forward primer SEQ ID No. 1 and a reverse primer SEQ ID No. 16,

(2) a combination of a forward primer SEQ ID No. 28 and a reverse primer SEQ ID No. 44,

5 (3) a combination of a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52,

(4) a combination of a forward primer SEQ ID No. 53 and a reverse primer SEQ ID No. 57,

10 (5) a combination of a forward primer SEQ ID No. 62 and a reverse primer SEQ ID No. 70,

(6) a combination of a forward primer SEQ ID No. 76 and a reverse primer SEQ ID No. 91 and

(7) a combination of a forward primer SEQ ID No. 104 and a reverse primer SEQ ID No. 2..

15 Then, the presence or absence of amplification of each DNA sample was verified. The results are shown in Table 18.

[0209]

[Table 18]

Table 18 gamma ray sterilization

Amplification of target product	
No treatment	○
10kGy	△
25kGy	x

○ : Amplification was observed.

△ : Amplification was slightly observed.

x: No amplification was observed.

[0210]

(Example 16) DNA removal treatment (EMA)

The instruments and ultrapure water used were free of DNA, and the operation was conducted in a clean bench. A solution was prepared by omitting the template, the primers and EvaGreen from the composition of the reaction liquid formulation of Table 5, but adding 25 ng of E. coli genomic DNA. Next, Ethidium Monoazide Bromid (Molecular Probe) was added to the resultant solution so that the final concentration of Ethidium Monoazide Bromid was 20 μ M to provide a solution containing Ethidium Monoazide Bromid, while the above resultant solution without addition of Ethidium Monoazide Bromid was provided as the negative control. These solutions were placed under an LED bulb of 570 lm and irradiated with light for 15 minutes. Thereafter, primers and EvaGreen were added, and each mixture was heated at 95°C for 10 minutes, then, a process including heating at 95°C for 10 seconds, at 60°C for 30 seconds and at 72°C was repeated 60 times, then, the DNA dissociation curve of the amplification product was made. The presence or absence of amplification of the intended product was verified. For the primer, a forward primer SEQ ID No. 51 and a reverse primer SEQ ID No. 52 were adopted. The results are shown in Fig. 5.

20 [0211]

(Example 17) Identification of bacterial species by four T_m values

A blind test was conducted by preparing the unidentified solution having no bacterial names of 4 bacterial species. *Escherichia coli*, *Clostridium difficile*, *Klebsiella pneumoniae* and *Enterobacter cloacae* were named as Bacterium No. 1 to No. 4, respectively and DNA was extracted from each bacterium to try to identify them. PCR reactions were carried out

in the same manner as Example 11. The identification was made only on the basis of a combination of four T_m values among them. In view of the measurement error of Rotor-Gene Q, those in which $\text{Dist} < 0.2$ or less were regarded as homogeneous.

5 [0212]

[Table 19]

Table 19 Discrimination of bacterial species by four T_m values

Sample name	Name of bacterial species	Dist.
Bacterium No. 1	<i>E. coli</i>	≤ 0.18
Bacterium No. 2	<i>C. difficile</i>	≤ 0.15
Bacterium No. 3	<i>K. pneumoniae</i>	≤ 0.19
Bacterium No. 4	<i>E. cloacae</i>	≤ 0.19

Claim 1

A primer set for polymerase chain reaction (PCR) for amplification of bacterial DNA, which comprises one primer pair obtained by selecting from

group S2:

2-16A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of any one sequence of SEQ ID: No. 44, 46 to 48 and 50.

Claim 2

A primer set according to claim 1, which further comprises at least one primer pair of seven primer pairs obtained by selecting one primer pair from each group of the S1 and S3 to S7 groups:

Group S1:

1-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 1 and a reverse primer consisting of any one sequence of SEQ ID: No. 22 and 27;

1-2A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 10 and a reverse primer consisting of any one sequence of SEQ ID: No. 20 and 21;

1-3A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 11 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 18 and 26;

1-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 12 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 13 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-8A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 6 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 19 to 21, 23, 24, 26 and 27;

1-9A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 7 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21, 23, 24, 26 and 27; or

1-10A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 8 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21, 23, 24, 26 and 27;

Group S3:

a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

Group S4:

4-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 61;

4-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 54 and a reverse primer consisting of any one sequence of SEQ ID: No. 59 and 61;

4-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 55 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61; or

4-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 56 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61;

Group S5:

5-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 73 to 75;

5-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 63 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 64 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 65 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-5A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 66 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 72, 73 and 75;

5-6A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 67 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 71 and 73 to 75;

5-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 68 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75; or

5-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 69 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

Group S6:

6-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93 and 101 to 103;

6-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 76 and a reverse primer consisting of any one sequence of SEQ ID: No. 93, 96 and 101 to 103;

6-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 77 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96 and 101 to 103;

6-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 78 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-4A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 79 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 80 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93 and 100 to 103;

6-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 82 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, 97 and 101 to 103;

6-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 83 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 101 to 103;

6-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 84 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 85 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-11A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 86 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93 and 101 to 103;

6-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 87 and a reverse primer consisting of any one sequence of SEQ ID: Nos. 91, 96 and 101 to 103;

6-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 88 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-14: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 89 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103; or

6-15: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 90 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 92, 96 and 101 to 103;

Group S7:

7-12A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of any one sequence of SEQ ID: No. 2 and 5;

7-1A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 104 and a reverse primer consisting of any one sequence of SEQ ID: No. 5 and 118 to 125, 127, 128, 130 and 131;

7-2A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 105 and a reverse primer consisting of any one sequence of SEQ ID: No. 5 and 118 to 125;

7-3A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 106 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 and 118 to 131;

7-4A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 107 and a reverse primer consisting of any one sequence of SEQ ID: No. 120 and 121;

7-5A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 108 and a reverse primer consisting of any one sequence of SEQ ID: No. 120 and 123;

7-6A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 109 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 118 to 122, 124 and 126;

7-7A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 110 and a reverse primer consisting of any one sequence of SEQ ID: No. 118, 120, 121, 124 and 126 to 130;

7-8A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 111 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 118 to 131;

7-9A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 112 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116, and 118 to 125;

7-10A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 113 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, 118 to 126, 130 and 131; or

7-11A: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 114 and a reverse primer consisting of a sequence of SEQ ID: No. 5.

Claim 3

A kit for detection of a bacterial species, which comprises one primer pair obtained by selecting from the following group K2:

2-16: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of any one sequence of SEQ ID: No. 44 to 48 and 50.

Claim 4

A kit according to claim 3, which further comprises at least one primer pair of seven primer pairs obtained by selecting one primer pair from each group of the K1 and K3 to K7 groups:

Group K1:

1-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 1 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 10 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 18, 20 and 21;

1-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 11 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 21 and 23 to 26;

1-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 12 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 13 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 14 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 6 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 7 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27;

1-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 8 and a reverse primer consisting of any one sequence of SEQ ID: No. 16 to 27; or

1-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 9 and a reverse primer consisting of any one sequence of SEQ ID: No. 16, 17, 19 and 26;

Group K3:

a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

Group K4:

4-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 61;

4-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 54 and a reverse primer consisting of any one sequence of SEQ ID: No. 59 and 61;

4-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 55 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61; or

4-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 56 and a reverse primer consisting of any one sequence of SEQ ID: No. 57 to 59 and 61;

Group K5:

5-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting of any one sequence of SEQ ID: No. 70, 71 and 73 to 75;

5-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 63 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 64 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 65 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 66 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 67 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

5-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 68 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75; or

5-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 69 and a reverse primer consisting of any one sequence of SEQ ID: No. 70 to 75;

Group K6:

6-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93 and 101 to 103;

6-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 76 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, 96 and 101 to 103;

6-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 77 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96 and 101 to 103;

6-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 78 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 79 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 97 and 101 to 103;

6-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 80 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93 and 100 to 103;

6-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 82 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96, 97 and 101 to 103;

6-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 83 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 101 to 103;

6-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 84 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 85 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 86 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 93, 97 and 101 to 103;

6-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 87 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 96 and 101 to 103;

6-13: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 88 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103;

6-14: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 89 and a reverse primer consisting of any one sequence of SEQ ID: No. 91 and 101 to 103; or

6-15: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 90 and a reverse primer consisting of any one sequence of SEQ ID: No. 91, 92, 96 and 101 to 103;

Group K7:

7-12: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5, 116, 118, 123 and 125 to 128;

7-1: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 104 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-2: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 105 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-3: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 106 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-4: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 107 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-5: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 108 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-6: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 109 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-7: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 110 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-8: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 111 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-9: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 112 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131;

7-10: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 113 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 116 to 131; or

7-11: a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 114 and a reverse primer consisting of any one sequence of SEQ ID: No. 2, 5 and 125.

Claim 5

A primer set according to claim 2 or a kit according to claim 4, which comprises at least one primer pair of the following seven primer pairs 1) to 7):

1) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 15 and a reverse primer consisting of a sequence of SEQ ID: No. 16;

2) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 43 and a reverse primer consisting of a sequence of SEQ ID: No. 50;

3) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 81 and a reverse primer consisting of a sequence of SEQ ID: No. 103;

4) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 115 and a reverse primer consisting of a sequence of SEQ ID: No. 5;

5) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 51 and a reverse primer consisting of a sequence of SEQ ID: No. 52;

6) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 53 and a reverse primer consisting of a sequence of SEQ ID: No. 57; and

7) a primer pair, which consists of a combination of a forward primer consisting of a sequence of SEQ ID: No. 62 and a reverse primer consisting of a sequence of SEQ ID: No. 70.

Claim 6

A primer set or a kit according to claim 5, comprising any one of the primer pairs 1) to 7).

Claim 7

A primer set or a kit according to claim 6, comprising any one of the following combinations A to G:

(A) when the primer set comprises primer pair 1), the primer set further comprises at least 3 primer pairs selected from 2) to 7);

(B) when the primer set comprises primer pair 2), the primer set further comprises at least 3 primer pairs selected from 1) and 3) to 7);

(C) when the primer set comprises primer pair 3), the primer set further comprises at least 3 primer pairs selected from 1), 2) and 4) to 7);

(D) when the primer set comprises primer pair 4), the primer set further comprises at least 3 primer pairs selected from 1) to 3) and 5) to 7);

(E) when the primer set comprises primer pair 5), the primer set further comprises at least 3 primer pairs selected from 1) to 4) and 6) to 7);

(F) when the primer set comprises primer pair 6), the primer set further comprises at least 3 primer pairs selected from 1) to 5) and 7);

(G) when the primer set comprises primer pair 7), the primer set further comprises at least 3 primer pairs selected from 1) to 6).

Claim 8

A primer set according to any one of claims 1, 2 or 5 to 7 for detection and/or identification of a bacterial species.

Claim 9

A kit according to any one of claims 3 to 7 for identification of a bacterial species.

Claim 10

A primer set according to any one of claims 2, or 5 to 8 or a kit according to any one of claims 4, 5 to 7, or 9, wherein the primer set or kit comprises at least four primer pairs of the seven primer pairs.

Claim 11

A kit according to any one of claims 3 to 7, or 9, which further comprises an additional primer pair of a forward primer consisting of a sequence of SEQ ID: No. 1 and a reverse primer consisting of any one sequence of SEQ ID: No. 2 to 5, such as wherein the additional primer pair consists of a forward primer consisting of a sequence of SEQ ID No. 1 and a reverse primer consisting of a sequence of SEQ ID No. 5.

Claim 12

A kit according to any one of claims 3 to 7, or 9 which further comprises an enzyme for PCR.

Claim 13

A kit according to claim 12, wherein

(i) the primers and the enzyme for PCR have, independently or as a whole, a contamination amount of bacterial nucleic acid of 10 fg or less than 10 fg; Or

(ii) which further comprises at least one of a pH buffer, dNTP, MgCl₂ and a vessel for measurement, and preferably the pH buffer, dNTP, MgCl₂ and the vessel for measurement have, independently or as a whole, a contamination amount of bacterial nucleic acid of 10 fg or less than 10 fg, and more preferably the primers, the enzyme for PCR, the pH buffer, dNTP, MgCl₂ and the vessel for measurement are individually packaged and the contamination amount of bacterial nucleic acid in each of the packages is 10 fg or less than 10 fg.

Claim 14

A kit according to claim 12 or 13, part (i), which further comprises fluorochrome.

Claim 15

A kit according to claim 14, wherein the primer, the enzyme for PCR, the pH buffer, dNTP, the fluorochrome, MgCl₂ and the vessel for measurement have, independently or as a whole, a contamination amount of bacterial nucleic acid of 10 fg or less than 10 fg.

Claim 16

A kit according to claim 15, wherein the primer, the enzyme for PCR, the pH buffer, dNTP, the fluorochrome, MgCl₂ and the vessel for measurement are individually packaged and the contamination amount of bacterial nucleic acid in each of the packages is 10 fg or less than 10 fg.

Claim 17

A kit according to claim 13 or claim 14, wherein dNTP is chemically modified for hot start PCR.

Claim 18

A kit according to any one of claims 12 to 17, wherein the enzyme for PCR is a Taq DNA Polymerase having an amount of bacterial DNA of less than 10 fg/μl.

Claim 19

A method of detection and/or identification of a bacterial species as a target in a sample, comprising:

a step of conducting PCR using bacterial genome DNA prepared from the sample, primers to obtain amplification product comprising a target gene specific to the bacterial species as a target, a thermostable DNA polymerase, and

a step of detecting and/or identifying the bacterial species as a target in the sample by detecting the target gene in the amplification product by PCR, or analyzing the amplification product by PCR,

wherein, as the primers, the primer set according to claims 1, 2, and 5 to 8, or the kit according to any one of claims 3 to 7, and 9 to 16 is used.

Claim 20

A method of identification of a bacterial species according to claim 17, wherein a kit according to any one of claims 3 to 7, 9 to 16 is used.