



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 59/347, 51/567	A1	(11) International Publication Number: WO 96/35658 (43) International Publication Date: 14 November 1996 (14.11.96)
(21) International Application Number: PCT/EP96/01934 (22) International Filing Date: 6 May 1996 (06.05.96) (30) Priority Data: 95201233.4 / 12 May 1995 (12.05.95) HP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): QUEST INTERNATIONAL B.V. [NL/NL], Huizerstraatweg 28, NL-1411 GP Naarden (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): COENDERS, Peter, Joannes [NL/NL]; Pearl Buckerf 98, NL-3315 BB Dordrecht (NL). GALEMA, Saskia, Alexandra [NL/NL]; Bernweg 173, NL-3137 NC Vlaardingen (NL). GOMBERT, Johannes, Durk [NL/NL]; Kerkpad 10, NL-6883 AA Velp (NL). MEER, Anna, Sophie, E., ter [NL/NL]; Aalscholverlaan 266, NL-3136 KL Vlaardingen (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: COMPOSITIONS OF ENANTIOMERIC DIACYL TARTARIC ANHYDRIDES AND PREPARATION THEREOF (57) Abstract <p>The present invention relates to compositions comprising mixtures of two enantiomeric forms of diacyl tartaric anhydride and mixtures obtainable by reacting an organic compound with a mixture comprising two enantiomeric forms of diacyl tartaric anhydride. The invention also relates to a method for the preparation of such enantiomeric mixtures, starting from a mixture comprising D-tartaric acid and L-tartaric acid. The method involves reacting tartaric acid with a fatty acid anhydride under dielectric heating conditions.</p>		