

DESCRIPTION

Title of Invention: METHOD FOR PRODUCING RETINAL TISSUE
[Technical Field]
[0001]

5 The present invention relates to a method for producing
retinal cells or a retinal tissue from pluripotent stem cells.
[Background Art]
[0002]

As a method for producing a neural tissue such as a
10 retinal tissue from pluripotent stem cells, a method for
producing a neural tissue which comprises forming uniformed
aggregates of pluripotent stem cells in a serum-free medium,
culturing them in suspension, culturing them in suspension in a
medium for differentiation induction in the presence of a
15 differentiation-inducing factor and the like as appropriate to
induce differentiation of pluripotent stem cells into the
intended neural cells has been reported (patent document 1 and
non-patent document 1). For example, a method for obtaining a
multi-layered retinal tissue from pluripotent stem cells (non-
20 patent document 2 and patent document 2), and a method for
obtaining a multi-layered retinal tissue which comprises
forming uniformed aggregates of pluripotent stem cells in a
serum-free medium containing a Wnt signal transduction pathway
inhibiting substance, followed by culturing them in suspension
25 in the presence of a basement membrane preparation, and then
culturing them in suspension in a serum-containing medium (non-
patent document 3 and patent document 3), and a method for
obtaining a retinal tissue by culturing an aggregate of
pluripotent stem cells in suspension in a medium containing a
30 BMP signal transduction pathway activating substance (patent
document 5 and non-patent document 10) are known. In addition,
a method for inducing differentiation of pluripotent stem cells
into a hypothalamic tissue (patent document 4 and non-patent
document 4), and a method inducing differentiation of
35 pluripotent stem cells into neural progenitor cells (non-patent

document 5 and 6) have also been reported.

The pluripotent stem cells as a starting material of these production methods, particularly in the case of primate pluripotent stem cells, could be cultured while maintaining an undifferentiated state in the presence of feeder cells and with the addition of a factor for maintaining undifferentiated state. In recent years, improvement has been made in the culture to maintain undifferentiated state, and a method of culturing primate pluripotent stem cells in the absence of feeder cells (feeder-free) with the addition of a factor for maintaining undifferentiated state has been reported (non-patent documents 7, 8 and 9). A method for stably producing neural cells or a neural tissue such as retina and the like, which uses pluripotent stem cells subjected to feeder-free culture by this method as a starting material has been desired.

[Document List]

[Patent documents]

[0003]

patent document 1: WO 2009/148170

patent document 2: WO 2011/055855

patent document 3: WO 2013/077425

patent document 4: WO 2013/065763

patent document 5: WO 2015/025967

[non-patent documents]

[0004]

non-patent document 1: Cell Stem Cell, 3, 519-32 (2008)

non-patent document 2: Nature, 472, 51-56 (2011)

non-patent document 3: Cell Stem Cell, 10(6), 771-775 (2012)

non-patent document 4: Nature, 480, 57-62 (2011)

non-patent document 5: Nature Biotechnology, 27(3), 275-80 (2009)

non-patent document 6: Proc Natl Acad Sci USA, 110(50), 20284-9 (2013)

non-patent document 7: Nature Methods, 8, 424-429 (2011)

non-patent document 8: Scientific Reports, 4, 3594 (2014)

non-patent document 9: In Vitro Cell Dev Biol Anim., 46, 247-58
(2010)

non-patent document 10: Nature Communications, 6, 6286 (2015)

[Summary of Invention]

5 [Technical Problem]

[0005]

The problem to be solved by the present invention is
provision of a method for producing retinal cells or a retinal
tissue from pluripotent stem cells cultured while maintaining
10 an undifferentiated state in the absence of feeder cells.

[Solution to Problem]

[0006]

The present inventors have conducted intensive studies in
an attempt to solve the aforementioned problems and found that
15 a cell aggregate maintaining an undifferentiated state can be
formed with high efficiency by culturing pluripotent stem cells
in the absence of feeder cells (under feeder-free conditions)
in a medium containing a TGF β family signal transduction
pathway inhibiting substance and/or a Sonic hedgehog signal
20 transduction pathway activating substance, and then culturing
them in suspension in a medium containing a Wnt signal
transduction pathway inhibiting substance. In addition, the
present inventors have found that neural tissues such as a
retinal tissue and the like and neural cells can be induced
25 with high efficiency by using this high quality cell aggregate,
which resulted in the completion of the present invention.

That is, the present invention relates to the following.

[0007]

[1] A method for producing a retinal cell or a retinal tissue,
30 comprising the following steps (1) - (3):

(1) a first step of culturing pluripotent stem cells in the
absence of feeder cells in a medium containing 1) a TGF β family
signal transduction pathway inhibiting substance and/or a Sonic
hedgehog signal transduction pathway activating substance, and
35 2) a factor for maintaining undifferentiated state,

- (2) a second step of culturing the cells obtained in the first step in suspension in a medium containing a Wnt signal transduction pathway inhibiting substance to form a cell aggregate, and
- 5 (3) a third step of culturing the aggregate obtained in the second step in suspension in the presence or absence of a Wnt signal transduction pathway inhibiting substance in a medium containing a BMP signal transduction pathway activating substance to obtain an aggregate containing a retinal cell or a
- 10 retinal tissue.
- [2] The production method of [1], wherein the pluripotent stem cells are cultured for 0.5 hr - 144 hr in the first step.
- [3] The production method of [1] or [2], wherein the culturing in the first step is performed by adhesion culture.
- 15 [4] The production method of any of [1] to [3], wherein, in the second step, the cells obtained in the first step are dispersed, and the dispersed cells are cultured in suspension.
- [5] The production method of any of [1] to [4], wherein the factor for maintaining undifferentiated state comprises at
- 20 least an FGF signal transduction pathway activating substance.
- [6] The production method of [5], wherein the FGF signal transduction pathway activating substance is bFGF.
- [7] The production method of any of [1] to [6], wherein the medium used for suspension culture in the second step further
- 25 comprises a Sonic hedgehog signal transduction pathway activating substance.
- [8] The production method of [7], wherein the pluripotent stem cells are human pluripotent stem cells and, in the second step, the concentration of the Sonic hedgehog signal transduction
- 30 pathway activating substance in the medium is a concentration corresponding to Sonic hedgehog signal transduction activity of SAG at 10 nM to 700 nM.
- [9] The production method of any of [1] to [8], wherein the TGF β family signal transduction pathway inhibiting substance is
- 35 a TGF β signal transduction pathway inhibiting substance or a

BMP signal transduction pathway inhibiting substance.

[10] The production method of any of [1] to [9], wherein the TGF β family signal transduction pathway inhibiting substance is one or more substances selected from the group consisting of
5 Lefty, SB431542, A-83-01 and LDN193189.

[11] The production method of any of [1] to [10], wherein the Sonic hedgehog signal transduction pathway activating substance is one or more substances selected from the group consisting of Shh, SAG and Purmorphamine.

10 [12] The production method of any of [1] to [11], wherein, in the third step, the BMP signal transduction pathway activating substance is added to the medium between day 1 and day 9 from the start of the second step.

[13] The production method of [12], wherein, in the third step,
15 the BMP signal transduction pathway activating substance is added to the medium between day 1 and day 6 from the start of the second step.

[14] The production method of [13], wherein, in the third step, the BMP signal transduction pathway activating substance is
20 added to the medium between day 1 and day 3 from the start of the second step.

[15] The production method of any of [1] to [14], wherein the BMP signal transduction pathway activating substance is one or more proteins selected from the group consisting of BMP2, BMP4,
25 BMP7 and GDF7.

[16] The production method of any of [1] to [15], wherein the BMP signal transduction pathway activating substance is BMP4.

[17] The production method of any of [1] to [16], wherein, in the third step, the aggregate is cultured in a medium
30 containing a Sonic hedgehog signal transduction pathway activating substance at a concentration not more than a concentration corresponding to Sonic hedgehog signal transduction activity of SAG at 700 nM.

[18] The production method of any of [1] to [16], wherein, in
35 the third step, the concentration of the Sonic hedgehog signal

transduction pathway activating substance in the medium is not more than a concentration corresponding to Sonic hedgehog signal transduction activity of SAG at 700 nM.

[19] The production method of any of [1] to [18], wherein the
5 culturing is performed in a medium containing a Wnt signal transduction pathway inhibiting substance for 3 days to for 18 days from the start of the second step.

[20] The production method of any of [1] to [19], wherein the
10 culturing is performed in a medium containing a Wnt signal transduction pathway inhibiting substance for 10 days from the start of the second step.

[21] The production method of any of [1] to [20], wherein the Wnt signal transduction pathway inhibiting substance is IWR-1-endo.

15 [22] The production method of any of [1] to [21], wherein the third step comprises the following steps:

- (i) a step of culturing the aggregate obtained in the second step in suspension in the presence or absence of a Wnt signal transduction pathway inhibiting substance in a medium
20 containing a BMP signal transduction pathway activating substance to obtain a cell aggregate containing a retinal cell or a retinal tissue and having an existence ratio of Chx10 positive cells of not less than 20% and not more than 100%;
- (ii) a step of culturing the cell aggregate obtained in step
25 (i) in a serum-free medium or serum-containing medium containing a Wnt signal transduction pathway activating substance and/or an FGF signal transduction pathway inhibiting substance for only a period before the appearance of a cell expressing RPE65 gene;
- 30 (iii) a step of culturing the cell aggregate obtained in step (ii), in which the RPE65 gene-expressing cell does not appear, in a serum-free medium or a serum-containing medium without a Wnt signal transduction pathway activating substance.

[23] The production method of any of [1] to [22], wherein the
35 aggregate obtained in the third step comprises one or more

cells selected from the group consisting of retinal progenitor cell, neural retinal progenitor cell, photoreceptor precursor cell, photoreceptor cell, rod photoreceptor cell, cone photoreceptor cell, horizontal cell, amacrine cell, interneuron, 5 ganglion cell, bipolar cell, retinal pigment epithelial cell, and ciliary marginal zone cell.

[24] The production method of any of [1] to [23], wherein the pluripotent stem cells are human pluripotent stem cells.

[25] The production method of any of [1] to [24], wherein the 10 pluripotent stem cells are induced pluripotent stem cells.

[26] The production method of any of [1] to [25], wherein a uniformed aggregate is formed in the second step.

[27] The production method of any of [1] to [26], wherein the suspension culture is performed in the absence of a basement 15 membrane preparation.

[28] A reagent for evaluating toxicity or efficacy of a test substance, comprising a retinal cell or a retinal tissue produced by the method of any of [1] to [27].

[29] A method for evaluating toxicity or efficacy of a test 20 substance, comprising bringing the substance into contact with a retinal cell or a retinal tissue produced by the method of any of [1] to [27], and detecting an influence of the substance on the cell or tissue.

[30] A medicament for treating a disease due to a disorder of a 25 retinal cell or a retinal tissue, comprising a retinal cell or a retinal tissue produced by the method of any of [1] to [27].

[31] The medicament of [30], wherein the retinal cell or retinal tissue is a retinal progenitor cell, a retinal layer-specific neural cell or a retinal tissue.

30 [32] A method for treating a disease due to a disorder of a retinal cell or a retinal tissue, comprising transplanting an effective amount of a retinal cell or a retinal tissue produced by the method of any of [1] to [27] to a subject in need of the transplantation.

35 [33] A retinal cell or a retinal tissue produced by the method

of any of [1] to [27] for use in the treatment of a disease due to a disorder of a retinal cell or a retinal tissue.

[34] A pharmaceutical composition comprising a retinal cell or a retinal tissue produced by the method of any of [1] to [27]
5 as an active ingredient.

[Advantageous Effects of Invention]

[0008]

According to the present invention, a high quality cell aggregate, as well as retinal cells or retinal tissues can be
10 produced with high efficiency from pluripotent stem cells cultured in the absence of feeder cells.

[Brief Description of Drawings]

[0009]

Fig. 1 shows the comparison results of morphology of cell
15 aggregates by bright field observation between the conditions of with (C, D) or without (A, B) Precondition.

Fig. 2 shows the comparison results of expression of Rx (A-D) and Chx10 (E-H) in cell aggregates by immunohistostaining between the conditions of with (C, D, G, H) or without (A, B, E,
20 F) Precondition.

Fig. 3 shows the comparison results of morphology of cell aggregates formed from human iPS cells under various culture conditions by bright field observation.

Fig. 4 shows the comparison results of expression of Rx (A-D) and Chx10 (E-H) in cell aggregates formed from human iPS
25 cells under various culture conditions by immunohistostaining.

Fig. 5 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

30 Fig. 6 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 7 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under
35 various culture conditions by immunohistostaining.

Fig. 8 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining. Counterstained with DAPI.

5 Fig. 9 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining. Counterstained with DAPI.

Fig. 10 shows the comparison results of expression of
10 Chx10 (A, C) and Rx (B, D) in cell aggregates formed from human iPS cells (A, B: TFH-R1-10-2 strain, C, D: TFH-R2-10-F8 strain) under various culture conditions by immunohistostaining.

Fig. 11 shows bright field observation results of a photoreceptor cell-containing retinal tissue induced from human
15 iPS cells (A). Observation results (B-E) of expression of Rx, Chx10 and Crx in the photoreceptor cell-containing retinal tissue by immunohistostaining are shown. Counterstained with DAPI (E).

Fig. 12 shows the comparison results of expression of
20 Chx10 (A, D, G), Rx (B, E, H) and Pax6 (C, F, I) in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 13 shows the comparison results of expression of Chx10 (A, D, G), Rx (B, E, H) and Pax6 (C, F, I) in cell
25 aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 14 shows bright field observation results of a photoreceptor cell-containing retinal tissue induced from human iPS cells (A). Observation results (B-E) of expression of Crx,
30 Chx10, Rx, Recoverin, NRL, RXRG, N-cadherin and Aqp1 in the photoreceptor cell-containing retinal tissue by immunohistostaining are shown. Counterstained with DAPI (B-E).

Fig. 15 shows bright field observation results of a photoreceptor cell-containing retinal tissue induced from human
35 iPS cells (A). Observation results (B-E) of expression of Crx,

Chx10, Rx, Recoverin, NRL, RXRG, N-cadherin and Aqp1 in the photoreceptor cell-containing retinal tissue by immunohistostaining are shown. Counterstained with DAPI (B-E).

Fig. 16 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 17 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 18 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 19 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 20 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 21 shows the comparison results of expression of Chx10 in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 22 shows the comparison results of expression of Chx10 and Rx in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 23 shows the comparison results of expression of Chx10, Rx, Crx, and Brn3b in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 24 shows the comparison results of expression of Chx10 and Rx in cell aggregates formed from human iPS cells under various culture conditions by immunohistostaining.

Fig. 25 shows the comparison results of expression of Chx10 and Rx in cell aggregates formed from human iPS cells by immunohistostaining.

[Description of Embodiments]

[0010]

1. Definition

In the present invention, "stem cell" means an undifferentiated cell having differentiation potency and proliferative capacity (particularly self-renewal competence) maintaining differentiation potency. The stem cell includes subpopulations such as pluripotent stem cell, multipotent stem cell, unipotent stem cell and the like according to the differentiation potency. Pluripotent stem cell refers to a stem cell capable of being cultured in vitro and having a potency to differentiate into any cell lineage belonging to three germ layers (ectoderm, mesoderm, endoderm) and/or extraembryonic tissues (pluripotency). The multipotent stem cell means a stem cell having a potency to differentiate into plural types of tissues or cells, though not all kinds. The unipotent stem cell means a stem cell having a potency to differentiate into a particular tissue or cell.

[0011]

Pluripotent stem cell can be induced from fertilized egg, clone embryo, germ stem cell, stem cell in a tissue, somatic cell and the like. Examples of the pluripotent stem cell include embryonic stem cell (ES cell), EG cell (embryonic germ cell), induced pluripotent stem cell (iPS cell) and the like. Muse cell (Multi-lineage differentiating stress enduring cell) obtained from mesenchymal stem cell (MSC), and GS cell produced from reproductive cell (e.g., testis) are also encompassed in the pluripotent stem cell. Embryonic stem cell was first established in 1981, and has also been applied to the generation of knockout mouse since 1989. In 1998, human embryonic stem cell was established, which is also being utilized for regenerative medicine. ES cell can be produced by culturing an inner cell mass on a feeder cell or in a medium containing LIF. The production methods of ES cell are described in, for example, WO 96/22362, WO 02/101057, US 5,843,780, US 6,200,806, US 6,280,718 and the like. Embryonic

stem cells are available from given organizations, or a commercially available product can be purchased. For example, human embryonic stem cells, KhES-1, KhES-2 and KhES-3, are available from Kyoto University's Institute for Frontier
5 Medical Sciences. Rx::GFP strain (derived from KhES-1), which is a human embryonic stem cell, is available from Incorporated Administrative Agency RIKEN. EB5 cell, which is a mouse embryonic stem cell, is available from Incorporated Administrative Agency RIKEN, and D3 cell line, which is a mouse
10 embryonic stem cell, is available from ATCC.

[0012]

Nuclear transfer ES cell (ntES cell), which is one of the ES cells, can be established from a clone embryo produced by transplanting the nucleus of a somatic cell into an enucleated
15 egg.

[0013]

EG cell can be produced by culturing a primordial germ cell in a medium containing mSCF, LIF and bFGF (Cell, 70: 841-847, 1992).

20 [0014]

The "induced pluripotent stem cell" in the present invention is a cell induced to have pluripotency by reprogramming a somatic cell by a known method and the like. Specifically, a cell induced to have pluripotency by
25 reprogramming differentiated somatic cells such as fibroblast, peripheral blood mononuclear cell and the like by the expression of a combination of a plurality of genes selected from the group consisting of reprogramming genes including Oct3/4, Sox2, Klf4, Myc (c-Myc, N-Myc, L-Myc), Glis1, Nanog,
30 Sall4, lin28, Esrrb and the like can be mentioned. Examples of preferable combination of reprogramming factors include (1) Oct3/4, Sox2, Klf4, and Myc (c-Myc or L-Myc), and (2) Oct3/4, Sox2, Klf4, Lin28 and L-Myc (Stem Cells, 2013; 31:458-466).

Induced pluripotent stem cell was established by Yamanaka
35 et al. in mouse cell in 2006 (Cell, 2006, 126(4), pp.663-676).

In 2007, Induced pluripotent stem cell was also established from human fibroblast, and has pluripotency and self-renewal competence similar to those of embryonic stem cells (Cell, 2007, 131(5), pp.861-872; Science, 2007, 318(5858), pp.1917-1920; Nat. Biotechnol., 2008, 26(1), pp.101-106).

Besides the production method based on direct reprogramming by gene expression, induced pluripotent stem cell can also be obtained from somatic cell by the addition of a compound and the like (Science, 2013, 341, pp. 651-654).

It is also possible to obtain established induced pluripotent stem cell and, for example, human induced pluripotent cell lines established by Kyoto University such as 201B7 cell, 201B7-Ff cell, 253G1 cell, 253G4 cell, 1201C1 cell, 1205D1 cell, 1210B2 cell or, 1231A3 cell and the like are available from Kyoto University and iPS Academia Japan, Inc. As the established induced pluripotent stem cell, for example, Ff-I01 cell, Ff-I14 cell and QHJI01s04 cell established by Kyoto University are available from Kyoto University.

[0015]

While the somatic cell used for producing induced pluripotent stem cell is not particularly limited, tissue-derived fibroblast, blood-lineage cells (e.g., peripheral blood mononuclear cell (PBMC), T cell), hepatocyte, pancreatic cell, intestinal epithelial cell, smooth muscle cell and the like can be mentioned.

[0016]

When an induced pluripotent stem cell is produced by reprogramming by the expression of several kinds of genes, the means for gene expression is not particularly limited. Examples of the aforementioned means include an infection method using a virus vector (e.g., retrovirus vector, lentivirus vector, adenovirus vector, adeno-associated virus vector, Sendaivirus vector which is cytosolic RNA vector), a gene transfer method using a plasmid vector which is a non-viral vector (e.g., plasmid vector, episomal vector) (e.g.,

calcium phosphate method, lipofection method, RetroNectin method, electroporation method), a gene transfer method using an RNA vector (e.g., calcium phosphate method, lipofection method, electroporation method), a method with direct injection
5 of protein (e.g., method using a needle, lipofection method, electroporation method) and the like.

[0017]

An induced pluripotent stem cell can be produced in the presence of a feeder cell or in the absence of feeder cells
10 (feeder-free). When an induced pluripotent stem cell is produced in the presence of a feeder cell, the induced pluripotent stem cell can be produced by a known method in the presence of a factor for maintaining undifferentiated state. While a medium to be used for producing an induced pluripotent
15 stem cell in the absence of feeder cells, i.e., a medium containing a factor for maintaining undifferentiated state (undifferentiation maintenance medium) is not particularly limited, a known maintenance medium for embryonic stem cells and/or induced pluripotent stem cells, and a medium well known
20 to those of ordinary skill in the art as a medium for establishing an induced pluripotent stem cell under feeder-free conditions can be used. As the undifferentiation maintenance medium for establishing an induced pluripotent stem cell under feeder-free conditions, many synthetic media have been
25 developed and are commercially available and, for example, Essential 8 medium (manufactured by Life Technologies) can be mentioned. Essential 8 medium is DMEM/F12 medium containing L-ascorbic acid-2-phosphate magnesium (64 mg/l), sodium selenite (14 µg/l), insulin (19.4 mg/l), NaHCO₃ (543 mg/l), transferrin
30 (10.7 mg/l), bFGF (100 ng/mL), and a TGFβ family signal transduction pathway activating substance (TGFβ 1 (2 ng/mL) or Nodal (100 ng/mL)) as additives (Nature Methods, 8, 424-429 (2011)). Examples of other commercially available feeder-free medium (undifferentiation maintenance medium) include Essential
35 6 medium (manufactured by Life Technologies), Stabilized

Essential 8 medium (manufactured by Life Technologies), S-medium (manufactured by DS Pharma Biomedical), StemPro (manufactured by Life Technologies), hESF9 (Proc Natl Acad Sci U S A. 2008 Sep 9; 105(36):13409-14), TeSR medium (manufactured
5 by STEMCELL Technologies), mTeSR1 (manufactured by STEMCELL Technologies), mTeSR2 (manufactured by STEMCELL Technologies), and TeSR-E8 (manufactured by STEMCELL Technologies). In addition to these, examples of other feeder-free medium (undifferentiation maintenance medium) include StemFit
10 (registered trade mark) (manufactured by Ajinomoto Co., Inc.). When an induced pluripotent stem cell is to be produced, for example, an induced pluripotent stem cell can be produced by gene transfer of 4 factors of Oct3/4, Sox2, Klf4, and Myc into somatic cell by using a Sendaivirus vector in the absence of
15 feeder cells.

[0018]

The pluripotent stem cell to be used in the present invention is preferably ES cell or induced pluripotent stem cell, more preferably induced pluripotent stem cell.

20 [0019]

As the multipotent stem cell, tissue stem cells (also called stem cell in a tissue, tissue-specific stem cell or somatic stem cell) such as hematopoietic stem cell, neural stem cell, retinal stem cell, mesenchymal stem cell and the like can
25 be mentioned.

[0020]

Genetically-modified pluripotent stem cells can be produced by using, for example, a homologous recombination technique. Examples of the gene on the chromosome to be
30 modified include a cell marker gene, a histocompatibility antigen gene, a gene related to a disease due to a disorder of neural cell and so on. A target gene on the chromosome can be modified using the methods described in Manipulating the Mouse Embryo, A Laboratory Manual, Second Edition, Cold Spring Harbor
35 Laboratory Press (1994); Gene Targeting, A Practical Approach,

IRL Press at Oxford University Press (1993); Biomanual Series 8, Gene Targeting, Making of Mutant Mouse using ES cell, YODOSHA CO., LTD. (1995); and so on.

[0021]

5 To be specific, for example, the genomic DNA comprising the target gene to be modified (e.g., cell marker gene, histocompatibility antigen gene, disease-related gene and so on) is isolated, and a targetting vector used for homologous recombination of the target gene is produced using the isolated
10 genomic DNA. The produced targetting vector is introduced into stem cells and the cells that showed homologous recombination between the target gene and the targetting vector are selected, whereby stem cells having the modified gene on the chromosome can be produced.

15 [0022]

Examples of the method for isolating genomic DNA comprising the target gene include known methods described in Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press (1989), Current Protocols in
20 Molecular Biology, John Wiley & Sons (1987-1997) and so on. The genomic gene comprising the target gene can also be isolated using genomic DNA library screening system (manufactured by Genome Systems), Universal GenomeWalker Kits (manufactured by CLONTECH) and so on. A polynucleotide
25 encoding the target protein can also be used instead of genome DNA. The polynucleotide can be obtained by amplifying the corresponding polynucleotide by the PCR method.

[0023]

Production of targetting vector used for homologous
30 recombination of the target gene, and efficient selection of a homologous recombinant can be performed according to the methods described in Gene Targeting, A Practical Approach, IRL Press at Oxford University Press (1993); Biomanual Series 8, Gene Targeting, Making of Mutant Mouse using ES cell, YODOSHA
35 CO., LTD. (1995); and so on. As the targetting vector, any of

replacement type or insertion type can be used. As the selection method, methods such as positive selection, promoter selection, negative selection, polyA selection and so on can be used.

5 Examples of a method for selecting the desired homologous recombinant from the selected cell lines include Southern hybridization method, PCR method and so on for the genomic DNA. [0024]

10 In addition, a genetically modified pluripotent stem cell can also be produced by genome editing. The genome editing is a genetic modification technique to perform gene-specific destruction, knock-in of reporter gene and the like by techniques such as Zinc Finger system, CRISPR/Cas9 system, Transcription Activator-Like Effector Nucleases (TALEN) and the
15 like. [0025]

20 In the CRISPR/Cas9 genome editing system, in general, an expression vector or mRNA of Cas9 (DNA cleaving enzyme) and an expression vector or guide RNA itself expressing the guide RNA under the control of polymerase III promoter and the like are introduced into cells. The guide RNA may be a fusion RNA of RNA (crRNA) complementary to the target genome sequence and tracrRNA. When protospacer adjacent motif (PAM - sequence NGG) is present at the 3' terminal of the target genomic sequence,
25 Cas9 dissociates the DNA double strand, recognizes the target sequence by the guide RNA, and cleaves both strands, and a mutation is introduced in the process of repairing the cleavage site. [0026]

30 The transcription activator-like effector nuclease (TALEN) is a system using a TAL effector produced by plant pathogenic bacterium *Xanthomonas* spp. TALEN is, in general, an artificial nuclease in which the DNA binding domain of TAL effector and the DNA cleavage domain of FokI nuclease are fused.
35 The DNA binding domain consists of a repeat sequence of 34

amino acids, and one repeat recognizes one base of the target DNA. The 12th and the 13th amino acids in the repeat sequence are called repeat variable di-residues (RVD) and the target base is determined by this sequence. The FokI domain dimerizes
5 on the target sequence and shows nuclease activity by designing two pairs of TALEN molecules to face each other on a specific sequence on the genome. The cleaved DNA double strand is repaired by the mechanism in the cell and a mutation is introduced during the process. When genome editing by TALEN is
10 performed, an expression vector or mRNA of TALEN is introduced into the cell.

[0027]

The "mammal" in the present invention encompasses rodents, ungulata, carnivora, primates and the like. The rodents
15 encompass mouse, rat, hamster, guinea pig and the like. Ungulata encompass swine, bovine, goat, horse, sheep and the like. Carnivora encompasses dog, cat and the like. The "primates" in the present invention refers to mammals belonging to the primate, and the primates include prosimian such as
20 lemur, loris, tupai and the like, and anthropoidea such as monkey, ape, human and the like.

[0028]

The pluripotent stem cells to be used in the present invention are mammalian pluripotent stem cells, preferably
25 pluripotent stem cells of rodents (e.g., mouse, rat) or primates (e.g., human, monkey), more preferably a human pluripotent stem cell, more preferably a human induced pluripotent stem cell (iPS cell) or human embryonic stem cell (ES cell).

30 [0029]

The "suspension culture" or "suspension culture method" in the present invention refers to culture while maintaining a state in which cells or cell aggregates are suspended in a medium and a method of performing the culture. That is, the
35 suspension culture is performed under conditions in which cells

or cell aggregates are not adhered to a culture vessel and the like, and culture performed under conditions permitting adhesion to a culture vessel and the like (adhesion culture or adhesion culture method) is not included in the category of
5 suspension culture. In this case, adhesion of cell means that a strong cell-substratum junction is formed between a cell or cell aggregate and a culture vessel. More particularly, suspension culture refers to culture under conditions in which a strong cell-substratum junction is not formed between a cell
10 or cell aggregate and a culture vessel, and adhesion culture refers to culture under conditions in which a strong cell-substratum junction is formed between a cell or cell aggregate and a culture vessel and the like.

[0030]

15 In a cell aggregate in suspension culture, a planar cell-cell adhesion is formed. In cell aggregates in suspension culture, a cell-substratum junction is hardly formed with a culture vessel and the like and, even if it is formed, its contribution is small. In some embodiments, an endogenous
20 cell-substratum junction is present inside the aggregate, but a cell-substratum junction is hardly formed with a culture vessel and the like and, even if it is formed, its contribution is small.

[0031]

25 The planar cell-cell adhesion (plane attachment) means that a cell attaches to another cell via planes. More particularly, the planar cell-cell adhesion means that, for example, not less than 1%, preferably not less than 3%, more preferably not less than 5%, of the surface area of a cell
30 adheres to the surface of another cell. A surface of a cell can be observed by staining with a reagent (e.g., DiI) that stains membranes, immunostaining of cell adhesion factors (e.g., E-cadherin and N-cadherin).

[0032]

35 The culture vessel to be used when performing suspension

culture is not particularly limited as long as it enables "culture in suspension" and those of ordinary skill in the art can appropriately determine same. Examples of such culture vessel include flask, tissue culture flask, culture dish (dish),
5 petri dish, tissue culture dish, multidish, microplate, microwell plate, micropore, multiplate, multiwell plate, chamber slide, schale, tube, tray, culture bag, spinner flask, Erlenmeyer flask, and roller bottle. To enable suspension culture, these culture vessels are preferably non-cell-adhesive.
10 Useful non-cell-adhesive culture vessels include culture vessels whose surfaces have not undergone an artificial treatment for improving adhesiveness to cells (e.g., surface treatment with extracellular matrix such as basement membrane preparation, laminin, entactin, collagen, gelatin etc., and the
15 like, or coating treatment with polymer such as polylysine, polyornithine and the like or positive electric charge treatment and the like), and the like. As a non-cell-adhesive culture vessel, culture vessels whose surfaces have been artificially treated to decrease adhesiveness to the cells
20 (e.g., superhydrophilic treatment with MPC polymer and the like, protein low adsorption treatment etc.) and the like can be used. Roller culture using spinner flask, roller bottle and the like may be performed. The culture surface of the culture vessel may be a flat bottom or may have concaves and convexes.

25 [0033]

A culture vessel used for adhesion culture is not particularly limited as long as "adhesion culture" can be performed, and those of ordinary skill in the art can appropriately select a culture vessel suitable according to the
30 culture scale, culture conditions and period for the culture. Examples of such culture vessel include flasks, tissue culture flasks, culture dishes (dishes), tissue culture dishes, multi-dishes, microplates, micro-well plates, multi-plates, multi-well plates, chamber slides, schale, tubes, trays, culture bags,
35 microcarrier, bead, stack plate, spinner flask and roller

bottles. To enable adhesion culture, these culture vessels are preferably cell-adhesive. Cell-adhesive culture vessels include culture vessels whose surfaces have been artificially treated to improve adhesiveness to cells, and specifically, a surface-processed culture vessel, or, a culture vessel whose inside is coated with a coating agent can be mentioned. Examples of the coating agent include extracellular matrix such as laminin [including laminin $\alpha 5 \beta 1 \gamma 1$ (hereinafter laminin 511), laminin $\alpha 1 \beta 1 \gamma 1$ (hereinafter laminin 111) and the like and laminin fragment (laminin 511E8 etc.)], entactin, collagen, gelatin, vitronectin, Synthemax (Corning Incorporated), Matrigel and the like, or polymers such as polylysine, polyornithine and the like. Examples of the surface-processed culture vessel include culture vessels surface-processed by a positive electric charge treatment and the like.

[0034]

The medium to be used for culturing cells in the present invention can be prepared from a medium generally used for culturing animal cells as a basal medium. Examples of the basal medium include media that can be used for culturing animal cells such as BME medium, BGJb medium, CMRL1066 medium, Glasgow MEM (GMEM) medium, Improved MEM Zinc Option medium, IMDM medium, Medium199 medium, Eagle MEM medium, α MEM medium, DMEM medium, F-12 medium, DMEM/F-12 medium, IMDM/F12 medium, Ham's medium, RPMI1640 medium, Fischer's medium, and mixed medium thereof etc.

[0035]

The "serum-free medium" in the present invention means a medium free of unadjusted or unpurified serum. In the present invention, a medium containing purified blood-derived components and animal tissue-derived components (e.g., growth factor) is also included in a serum-free medium unless unadjusted or unpurified serum is contained therein.

[0036]

The serum-free medium may contain a serum alternative.

Examples of the serum alternative include one appropriately containing albumin, transferrin, fatty acid, collagen precursor, trace element, 2-mercaptoethanol or 3' thioglycerol, or equivalents of these etc., and so on. Such serum alternative
5 may be prepared by, for example, the method described in WO98/30679. The serum alternative may be a commercially available product. Examples of such commercially available serum alternative include KnockoutTM Serum Replacement (Life Technologies, now ThermoFisher: hereinafter sometimes to be
10 indicated as KSR), Chemically Defined Lipid Concentrated (manufactured by Life Technologies) and GlutamaxTM (manufactured by Life Technologies), B27 (manufactured by Life Technologies), N2 supplement (manufactured by Life Technologies) and ITS supplement (manufactured by Life
15 Technologies).

[0037]

The serum-free medium to be used for suspension culture may appropriately contain a fatty acid or lipid, amino acid (e.g., non-essential amino acids), vitamin, growth factor,
20 cytokine, antioxidant, 2-mercaptoethanol, pyruvic acid, buffering agent, inorganic salts and so on.

[0038]

To avoid complicated preparation, a serum-free medium supplemented with an appropriate amount (e.g., about 0.5% to
25 about 30%, preferably about 1% to about 20%) of commercially available KSR (manufactured by Life Technologies) (e.g., medium of 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, Chemically-defined Lipid concentrated, and 450 μ M 1-monothioglycerol) may be used as such serum-free medium. In
30 addition, as a product equivalent to KSR, the medium disclosed in JP-A-2001-508302 can be mentioned.

[0039]

The "serum-containing medium" in the present invention means a medium containing unadjusted or unpurified serum. The
35 medium may contain a fatty acid or lipid, amino acid (e.g.,

non-essential amino acids), vitamin, growth factor, cytokine, antioxidant, 2-mercaptoethanol, 1-monothioglycerol, pyruvic acid, buffering agent, inorganic salts and so on.

[0040]

5 In one embodiment, the culture in the present invention may be preferably performed under xeno-free conditions. The "xeno-free" means conditions eliminating components derived from species different from that of the cell to be cultured.

[0041]

10 In the present invention, the "medium containing a substance X" and "in the presence of a substance X" refer to a medium supplemented with an exogenous substance X or a medium containing an exogenous substance X, or in the presence of an exogenous substance X. That is, when the cells or tissues
15 present in the medium endogenously express, secrete or produce a substance X, the endogenous substance X is distinguished from the exogenous substance X, and a medium free of exogenous substance X is understood to fall outside the category of the "medium containing a substance X", even when it contains the
20 endogenous substance X.

[0042]

For example, a "medium containing a TGF β family signal transduction pathway inhibiting substance" is a medium supplemented with an exogenous TGF β family signal transduction
25 pathway inhibiting substance or a medium containing an exogenous TGF β family signal transduction pathway inhibiting substance.

[0043]

In the present invention, a "feeder cell" refers to a
30 cell other than a stem cell that co-exists when culturing the stem cell. Examples of the feeder cells used for culturing pluripotent stem cells while maintaining undifferentiated state include mouse fibroblasts (MEF etc.), human fibroblasts, SNL cells and the like. As the feeder cells, feeder cells that
35 underwent a growth suppression treatment is preferable.

Examples of the growth suppression treatment include treatment with a growth inhibitor (e.g., mitomycin C), gamma irradiation, UV irradiation and the like. Feeder cells used for culturing pluripotent stem cells while maintaining undifferentiated state
5 contributes to the maintenance of undifferentiated state of pluripotent stem cell by secretion of a humoral factor (preferably factor for maintaining undifferentiated state), or production of a scaffold for cell adhesion (extracellular matrix).

10 [0044]

In the present invention, "the absence of feeder cells (feeder-free)" means culture in the absence of feeder cells. The absence of feeder cells means, for example, conditions free of addition of feeder cells, or conditions substantially free
15 of feeder cells (e.g., the ratio of the number of feeder cells relative to the total number of cells is not more than 3%, preferably not more than 0.5%).

[0045]

In the present invention, an "aggregate" of cells refers
20 to a clump formed by assembly of cells dispersed in a medium, wherein the cells are adhered to each other. Cell clumps, embryoid bodies, spheres, spheroids are also encompassed in the cell aggregates. Preferably, a planar cell-cell adhesion is formed in the aggregate of cells. In some embodiments, cells
25 sometimes form a cell-cell junction and/or a cell adhesion, for example, adherence junction, in some or all of the aggregates. The "aggregate" in the present invention specifically includes an aggregate produced in the second step of the above-mentioned present invention [1], which is formed by cells dispersed at
30 the time of the start of the suspension culture, and an aggregate produced in the third step of the above-mentioned present invention [1], which contains induced retinal cells differentiated from pluripotent stem cell, and the "aggregate" also includes an aggregate already formed at the time of the
35 start of suspension culture of the second step in the above-

mentioned present invention [1]. The cell aggregate formed in the second step encompasses "embryoid body (EB)".

[0046]

In the present invention, "uniformed aggregates" means
5 that the size of each aggregate is constant when a plurality of aggregates are cultured, and that the variance in the length of the maximum diameter is small when the size of the aggregates are evaluated by the length of the maximum diameter. More specifically, it means that not less than 75% of aggregates in
10 the whole aggregate population are within mean $\pm$ 100%, preferably mean $\pm$ 50%, more preferably mean $\pm$ 20%, of the maximum diameter in the population of the aggregates.

[0047]

In the present invention, to "form uniformed cell
15 aggregates" means to "rapidly aggregate a given number of dispersed cells" to form cell aggregates uniform in size, when gathering the cells to form cell aggregates and culturing the aggregates in suspension.

[0048]

"Dispersion" refers to dividing cells or a tissue into
20 small cell clusters (not less than 2 cells and not more than 100 cells, preferably not more than 50 cells) or single cells by a dispersion treatment such as enzymatic treatment, physical treatment and the like. A given number of dispersed cells is a
25 collection of a certain number of cell clusters or single cells.

[0049]

Examples of the method of dispersing pluripotent stem cells include a mechanical dispersion treatment, a cell dispersion solution treatment, and a cell protecting agent
30 addition treatment. These treatments may be performed in combination. Preferably, a cell dispersion solution treatment is performed and then a mechanical dispersion treatment is performed.

[0050]

35 As a method of mechanical dispersion treatment, a

pipetting treatment or scraping operation by a scraper can be mentioned.

[0051]

As a cell dispersion solution to be used for the cell
5 dispersion solution treatment, a solution containing any of
enzymes such as trypsin, collagenase, hyaluronidase, elastase,
pronase, DNase, papain and so on, and a chelating agent such as
ethylenediaminetetraacetic acid and so on can be mentioned. A
commercially available cell dispersion solution such as TrypLE
10 Select (manufactured by Life Technologies) and TrypLE Express
(manufactured by Life Technologies) can also be used.

[0052]

When pluripotent stem cells are dispersed, cell death of
the pluripotent stem cells may be suppressed by treating with a
15 cell protecting agent. As a cell protecting agent to be used
for the cell protecting agent treatment, an FGF signal
transduction pathway activating substance, heparin, an IGF
signal transduction pathway activating substance, serum, and
serum alternative can be mentioned. To suppress cell death
20 induced by dispersion (particularly, cell death of human
pluripotent stem cells), a Rho-associated coiled-coil kinase
(ROCK) inhibiting substance or a Myosin inhibiting substance
may be added at the time of dispersion. As a ROCK inhibiting
substance, Y-27632, Fasudil (HA1077), H-1152 and the like can
25 be mentioned. As a Myosin inhibiting substance, Blebbistatin
can be mentioned. As a preferable cell protecting agent, a
ROCK inhibiting substance can be mentioned.

[0053]

For example, a method for dispersing pluripotent stem
30 cells includes a method involving treating a colony of
pluripotent stem cells with a cell dispersion solution (TrypLE
Select) in the presence of a ROCK inhibiting substance as a
cell protecting agent, and further dispersing them by pipetting.

[0054]

35 In the production method of the present invention, it is

preferable to form an aggregate of pluripotent stem cells by rapidly gathering the pluripotent stem cells. When an aggregate of pluripotent stem cells is formed in such a manner, an epithelium-like structure can be formed with good
5 reproducibility in the cells induced and differentiated from the formed aggregate. Examples of the experimental operation to form an aggregate include a method involving keeping cells in a small space by using a plate with small wells (e.g., plate with wells having a base area of about 0.1 - 2.0 cm² when
10 calculated in terms of flat bottom), micropore and so on, a method involving aggregating cells by centrifugation for a short time using a small centrifugation tube. As a plate with small wells, for example, 24 well plate (area of about 1.88 cm² when calculated in terms of flat bottom), 48 well plate (area
15 of about 1.0 cm² when calculated in terms of flat bottom), 96 well plate (area of about 0.35 cm² when calculated in terms of flat bottom, inner diameter about 6 - 8 mm), and 384 well plate can be mentioned. Preferred is 96 well plate. As a shape of the plate with small wells, the shape of the bottom surface
20 when the well is seen from above is, for example, polygon, rectangle, ellipse, true circle, preferably true circle. As a shape of the plate with small wells when the well is seen from the side well, the shape of the bottom surface may be a flat bottom structure or a structure having high outer circumference
25 and low inner concave. The shape of the bottom surface includes, for example, U-bottom, V-bottom, M-bottom, preferably M-bottom or V-bottom. As a plate with small wells, a cell culture dish (e.g., 60 mm - 150 mm dish, culture flask) with a concave convex, or dent on the bottom surface may also be used.
30 The bottom surface of a plate with small wells is preferably a non-cell-adhesive bottom surface, preferably the aforementioned non-cell-adhesive-coated bottom surface.

[0055]

Formation of aggregates of pluripotent stem cells or a
35 cell population containing pluripotent stem cells, and

uniformity thereof can be determined based on the size of the aggregate mass and the number of cells therein, macroscopic morphology, microscopic morphology by tissue staining analysis and homogeneity thereof, and the like. In addition, formation
5 of an epithelial-like structure in the aggregate, and uniformity thereof can be determined based on the macroscopic morphology of the aggregate, microscopic morphology by tissue staining analysis and uniformity thereof, expression of differentiation and undifferentiation markers and uniformity
10 thereof, control of expression of differentiation marker and synchronism thereof, reproducibility of differentiation efficiency between aggregates, and so on.

[0056]

The "tissue" in the present invention refers to a
15 structure of a cell population having a structure in which one kind of cells having a uniformed morphology or property, or plural types of cells having different morphologies and properties are sterically arranged in a given pattern.

[0057]

20 In the present invention, the "neural tissue" refers to a tissue constituted of neural cells including cerebrum, midbrain, cerebellum, spinal cord, retina, peripheral nerve, forebrain, hindbrain, telencephalon, diencephalon and the like in the developing stage or adult stage. A neural tissue in a cell
25 aggregate can be confirmed by microscopic observation with optical microscopes (e.g., bright field microscope, phase contrast microscope, differential interference contrast microscope, stereo microscope etc.) or using expression of neural tissue markers such as PSA-NCAM, N-cadherin or the like
30 as an index.

[0058]

A neural tissue can form a "neuroepithelial tissue" (epithelial structure with layer structure). The characteristics of the neuroepithelial tissue can be identified
35 by indices such as the morphology of the cells contained in the

tissue, the orientation of the cell body in the tissue, the transparency of the whole tissue, the macroscopic form and the like. The amount of neuroepithelial tissues in cell aggregates can be evaluated by bright field observation using an optical
5 microscope.

[0059]

In the present invention, the "neural cell" refers to a cell other than epidermal lineage cell in a tissue derived from ectoderm. That is, it includes cells such as neural precursor
10 cell, neuron (neuronal cell), glia, neural stem cell, neuron precursor cell, glial precursor cell and the like. The neural cell can be identified by using Nestin, TuJ1, PSA-NCAM, N-cadherin and the like as a marker. Neuron (or neuronal cell) is a functional cell that forms a neural circuit and
15 contributes to signal transmission, and can be identified by using the expression of immature neuronal markers such as TuJ1, Dcx, HuC/D and the like and/or mature neuronal cell markers such as Map2, NeuN and the like as an index.

[0060]

20 Neural cell encompasses retinal cell described below.

[0061]

In the present invention, the "retinal cell" means a cell constituting each retinal layer or a progenitor/precursor cell thereof in retina in vivo, and nonlimitatively include retinal
25 progenitor cell, neural retinal progenitor cell, photoreceptor precursor cell, photoreceptor cell, rod photoreceptor cell, cone photoreceptor cell, horizontal cell, amacrine cell, interneuron, retinal ganglion cell (ganglion cell), bipolar cell, retinal pigment epithelial cell (RPE), ciliary marginal
30 zone cell, progenitor/precursor cells of these and the like. In the present invention, the "retinal tissue" means a tissue in which one type or at least two or more types of the above-mentioned cells that constitute respective retinal layers in retina in vivo are sterically arranged in layers. The retinal
35 layer which is constituted by each cell can be confirmed by a

known method, for example, presence or absence of the expression of a cell marker or the level thereof, etc.

[0062]

The "retinal layer" in the present invention means each
5 layer constituting the retina. Specific examples thereof
include retinal pigment epithelial layer, photoreceptor cell
layer, external limiting membrane, outer nuclear layer, outer
plexiform layer, inner nuclear layer, inner plexiform layer,
ganglion cell layer, nerve fiber layer and inner limiting
10 membrane.

[0063]

The "retinal progenitor cell" in the present invention
refers to a progenitor cell capable of differentiating into any
mature retinal layer-specific neural cells including
15 photoreceptor cell, rod photoreceptor cell, cone photoreceptor
cell, horizontal cell, bipolar cell, amacrine cell, retinal
ganglion cell, retinal pigment epithelial cell and the like.

[0064]

In the present invention, the "neural retinal progenitor
20 cell" refers to a progenitor cell capable of differentiating
into any one of or plural mature retinal layer-specific neural
cells including photoreceptor cell, horizontal cell, bipolar
cell, amacrine cell, retinal ganglion cell, and the like. In
general, a neural retinal progenitor cell does not
25 differentiate into a retinal pigment epithelial cell.

[0065]

The precursor cell of photoreceptor, precursor cell of
horizontal cell, precursor cell of bipolar cell, precursor cell
of amacrine cell, precursor cell of retinal ganglion cell, and
30 precursor cell of retinal pigment epithelial progenitor cell
respectively refers to precursor cell committed to
differentiate into photoreceptor cell, horizontal cell, bipolar
cell, amacrine cell, retinal ganglion cell, and retinal pigment
epithelial cell.

35 [0066]

In the present invention, the "retinal layer-specific neural cell" is a cell constituting a retina layer and is a neuronal cell specific to the retinal layer. Examples of the retinal layer-specific neural cell include bipolar cell,
5 retinal ganglion cells, amacrine cell, horizontal cell, photoreceptor cell, retinal pigment epithelial cell, rod photoreceptor cell and cone photoreceptor cell.

[0067]

Examples of the retinal cell marker include Rx (also
10 referred to as Rax), PAX6 and Chx10 expressed in retinal progenitor cell, Nkx2.1 expressed in progenitor cell of hypothalamus neuron but not expressed in retinal progenitor cell, Sox1 expressed in hypothalamus neuroepithelium but not expressed in retina, Crx and Blimp1 expressed in progenitor
15 cell of photoreceptor cell, and the like. Examples of the marker of the retinal layer-specific neural cell include Chx10, PKC α and L7 expressed in bipolar cell, TUJ1 and Brn3 expressed in retinal ganglion cells, Calretinin expressed in amacrine cell, Calbindin expressed in horizontal cell, Rhodopsin and
20 Recoverin expressed in mature photoreceptor cell, Nrl and Rhodopsin expressed in rod photoreceptor cell, Rxr-gamma, S-Opsin and M/L-Opsin expressed in cone photoreceptor cell, RPE65 and Mitf expressed in retinal pigment epithelial cell, Rdh10 and SSEA1 expressed in ciliary marginal zone cell and the like.

25 [0068]

2. Method for producing retinal cells or retinal tissue

One embodiment of the present invention is a method for producing retinal cells or a retinal tissue, comprising the following steps (1) - (3):

- 30 (1) a first step of culturing pluripotent stem cells in the absence of feeder cells in a medium containing 1) a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and 2) a factor for maintaining undifferentiated state,
35 (2) a second step of culturing the cells obtained in the first

step in suspension in a medium containing a Wnt signal transduction pathway inhibiting substance to form a cell aggregate, and

(3) a third step of culturing the aggregate obtained in the second step in suspension in the presence or absence of a Wnt signal transduction pathway inhibiting substance in a medium containing a BMP signal transduction pathway activating substance to obtain an aggregate containing a retinal cell or a retinal tissue.

10 [0069]

2-1. Step (1)

In step (1), pluripotent stem cells are cultured in the absence of feeder cells in a medium containing 1) a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and 2) a factor for maintaining undifferentiated state.

[0070]

A preferable pluripotent stem cell in step (1) is an induced pluripotent stem cell or an embryonic stem cell (ES cell), more preferably a human induced pluripotent stem cell or a human embryonic stem cell (ES cell).

[0071]

The production method of induced pluripotent stem cells is not particularly limited, and it can be produced by a method well known to those of ordinary skill in the art as mentioned above. It is also desirable to perform a step for preparing induced pluripotent stem cells (that is, a step of reprogramming somatic cells to establish pluripotent stem cells) under feeder-free condition.

30 [0072]

While the production method of embryonic stem cells (ES cells) is not particularly limited, and can be produced by a method well known to those of ordinary skill in the art as mentioned above, it is also desirable to perform a step for preparing embryonic stem cells (ES cells) under feeder-free

condition.

[0073]

The maintenance culture or expansion culture of pluripotent stem cells to be used in step (1) can be performed
5 by a method well known to those of ordinary skill in the art. While the maintenance culture and expansion culture of pluripotent stem cells can be performed by adhesion culture or suspension culture, it is preferably performed by adhesion culture. While the maintenance culture and expansion culture
10 of pluripotent stem cells may be performed in the presence of feeders or under feeder-free condition, it is preferably performed under feeder-free condition. The absence of feeder cells (feeder-free) in maintenance culture and expansion culture of pluripotent stem cells means a condition
15 substantially free of feeder cells (e.g., the ratio of number of feeder cells relative to the total number of cells is not more than 3%). Preferably, the maintenance culture and expansion culture of pluripotent stem cells is performed under conditions free of feeder cells.

20 [0074]

The absence of feeder cells (feeder-free) in step (1) means a condition substantially free of feeder cells (e.g., the ratio of the number of feeder cells relative to the total number of cells is not more than 3%). Preferably, step (1) is
25 performed under a condition free of feeder cells. The medium to be used in step (1) is not particularly limited as long as it is a medium enabling culture of pluripotent stem cells to maintain undifferentiated state under feeder-free conditions (feeder-free medium). To enable culture to maintain
30 undifferentiated state, it contains a factor for maintaining undifferentiated state.

[0075]

The factor for maintaining undifferentiated state is not particularly limited as long as it is a substance having an
35 action to suppress differentiation of pluripotent stem cells.

Examples of the factor for maintaining undifferentiated state widely used by those of ordinary skill in the art include an FGF signal transduction pathway activating substance, a TGF β family signal transduction pathway activating substance, 5 insulin and the like in the case of primed pluripotent stem cells (e.g., human ES cells, human iPS cells). As the FGF signal transduction pathway activating substance, fibroblast growth factors (e.g., bFGF, FGF4, FGF8) can be specifically mentioned. As the TGF β family signal transduction pathway 10 activating substance, a TGF β signal transduction pathway activating substance, a Nodal/Activin signal transduction pathway activating substance can be mentioned. As the TGF β signal transduction pathway activating substance, for example, TGF β 1, TGF β 2 can be mentioned. As the Nodal/Activin signal 15 transduction pathway activating substance, for example, Nodal, Activin A, Activin B can be mentioned. The factor for maintaining undifferentiated state may contain one or more kinds of these. When human pluripotent stem cells (human ES cells, human iPS cells etc.) are cultured, the medium in step 20 (1) preferably contains bFGF as a factor for maintaining undifferentiated state.

[0076]

The factor for maintaining undifferentiated state to be used in the present invention is generally a factor for 25 maintaining undifferentiated state of mammals. The mammals are, for example, those mentioned above. Since the factor for maintaining undifferentiated state may have cross-reactivity among mammal species, a factor for maintaining undifferentiated state of any mammal may also be used as long as the 30 undifferentiated state of the pluripotent stem cells to be cultured can be maintained. Preferably, a factor for maintaining undifferentiated state of a mammal of the same species as the cells to be cultured is used. For example, for the culture of human pluripotent stem cells, human factor for 35 maintaining undifferentiated states (e.g., bFGF, FGF4, FGF8,

EGF, Nodal, Activin A, Activin B, TGF β 1, TGF β 2 etc.) are used. Here, the "human protein X" means that protein X has the amino acid sequence of protein X naturally expressed in human in vivo. [0077]

5 The factor for maintaining undifferentiated state to be used in the present invention is preferably isolated. Being "isolated" means that an operation to remove factors other than the intended component or cell has been performed, and the component or cell is no longer in a naturally occurring state. 10 Therefore, "isolated protein X" does not include an endogenous protein X produced from the cells or tissues to be cultured, and contained in a cell or tissue or in the medium. The purity of the "isolated protein X" (percentage of the weight of protein X to the total protein weight) is generally not less 15 than 70%, preferably not less than 80%, more preferably not less than 90%, further preferably not less than 99%, further preferably 100%. Therefore, in one embodiment, the present invention comprises a step of providing an isolated factor for maintaining undifferentiated state. In one embodiment, it 20 includes a step of exogenously adding an isolated factor for maintaining undifferentiated state to a medium used in step (1). Alternatively, a factor for maintaining undifferentiated state may be added in advance to a medium to be used in step (1). [0078]

25 The concentration of the factor for maintaining undifferentiated state in the medium to be used in step (1) is a concentration capable of maintaining the undifferentiated state of the pluripotent stem cells to be cultured, and can be appropriately determined by those of ordinary skill in the art. 30 For example, specifically, when bFGF is used as a factor for maintaining undifferentiated state in the absence of feeder cells, the concentration thereof is generally about 4 ng - 500 ng/mL, preferably about 10 ng - 200 ng/mL, more preferably about 30 ng - 150 ng/mL.

35 [0079]

As the feeder free medium (i.e., undifferentiation maintenance medium) to be used in step (1), many synthetic media have been developed and are commercially available. For example, Essential 8 (manufactured by Life Technologies) medium
5 can be mentioned. Essential 8 medium is DMEM/F12 medium containing L-ascorbic acid-2-phosphate magnesium (64 mg/l), sodium selenite (14 µg/l), insulin (19.4 mg/l), NaHCO₃ (543 mg/l), transferrin (10.7 mg/l), bFGF (100 ng/mL), and a TGFβ family signal transduction pathway activating substance (TGFβ 1
10 (2 ng/mL) or Nodal (100 ng/mL)) as additives (Nature Methods, 8, 424-429 (2011)). Examples of other commercially available feeder-free medium (undifferentiation maintenance medium) include Essential 6 medium (manufactured by Life Technologies), Stabilized Essential 8 medium (manufactured by Life
15 Technologies), S-medium (manufactured by DS Pharma Biomedical Co., Ltd.), StemPro (manufactured by Life Technologies), hESF9 (Proc Natl Acad Sci U S A. 2008 Sep 9; 105(36):13409-14), TeSR medium (manufactured by STEMCELL Technologies), mTeSR1 (manufactured by STEMCELL Technologies),
20 mTeSR2 (manufactured by STEMCELL Technologies), TeSR-E8 (manufactured by STEMCELL Technologies). In addition to these, StemFit (registered trade mark) (manufactured by Ajinomoto Co., Inc.) can be mentioned as the feeder-free medium. The present invention can be performed conveniently by using these in the
25 above-mentioned step (1).

[0080]

In step (1), the pluripotent stem cells may be cultured under any conditions of suspension culture and adhesion culture, preferably adhesion culture.

30 [0081]

While a culture vessel used for adhesion culture is not particularly limited as long as "adhesion culture" can be performed, a cell-adhesive culture vessel is preferable. Cell-adhesive culture vessels include culture vessels whose surfaces
35 have been artificially treated to improve adhesiveness to cells,

and specifically, the above-mentioned culture vessel whose inside is coated with a coating agent can be mentioned. Examples of the coating agent include extracellular matrix such as laminin [including laminin $\alpha 5 \beta 1 \gamma 1$ (hereinafter laminin 511),
5 laminin $\alpha 1 \beta 1 \gamma 1$ (hereinafter laminin 111) and the like and laminin fragment (laminin 511E8 etc.)], entactin, collagen, gelatin, vitronectin, Synthemax (Corning Incorporated), Matrigel and the like, or polymer such as polylysine, polyornithine and the like, and the like. It is also possible
10 to use a culture container whose surface is processed by a positive electric charge treatment and the like. Preferred is laminin and more preferred is laminin 511E-8. Laminin 511E-8 can be a commercially available product (e.g., iMatrix-511, Nippi).

15 [0082]

The medium used in step (1) contains a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance. Specifically, for example, a TGF β family signal transduction
20 pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance can be added to the aforementioned undifferentiation maintenance medium. In the first step, pluripotent stem cells are treated with a TGF β family signal transduction pathway inhibiting substance and/or
25 a Sonic hedgehog signal transduction pathway activating substance and subjected to suspension culture in the second step, whereby the state of the cells changes, the quality of the aggregates is improved, and cell aggregates maintaining an undifferentiated state can be produced with high efficiency.
30 The thus-obtained cell aggregates are expected to show characteristics of, for example, spherical cell aggregates with a smooth surface and dense inside.

[0083]

The TGF β family signal transduction pathway (i.e., TGF β
35 superfamily signal transduction pathway) is a signal

transduction pathway intracellularly transduced by Smad family with TGF β , Nodal/Activin or BMP as a ligand.

[0084]

The TGF β family signal transduction pathway inhibiting substance is a substance that inhibits TGF β family signal transduction pathway, that is, a signal transduction pathway transduced by the Smad family. Specifically, a TGF β signal transduction pathway inhibiting substance, a Nodal/Activin signal transduction pathway inhibiting substance and a BMP
10 signal transduction pathway inhibiting substance can be mentioned.

[0085]

The TGF β signal transduction pathway inhibiting substance is not particularly limited as long as it is a substance
15 inhibiting a signal transduction pathway caused by TGF β , and may be any of nucleic acid, protein and low-molecular organic compound. As the substance, for example, a substance directly acting on TGF β (e.g., protein, antibody, aptamer etc.), a substance suppressing expression of gene encoding TGF β (e.g.,
20 antisense oligonucleotide, siRNA etc.), a substance that inhibits the binding of TGF β receptor and TGF β , and a substance that inhibits physiological activity caused by signal transduction by the TGF β receptor (e.g., TGF β receptor inhibitor, Smad inhibitor etc.) can be mentioned. As a protein
25 known as a TGF β signal transduction pathway inhibiting substance, Lefty and the like can be mentioned. As a TGF β signal transduction pathway inhibiting substance, compounds well known to those of ordinary skill in the art can be used and, specifically, SB431542, LY-364947, SB-505124, A-83-01 and
30 the like can be mentioned. SB431542 (4-(5-benzol[1,3]dioxol-5-yl-4-pyridin-2-yl-1H-imidazol-2-yl)-benzamide) and A-83-01 (3-(6-methyl-2-pyridinyl)-N-phenyl-4-(4-quinolinyl)-1H-pyrazole-1-carbothioamide) is a compound known as an inhibitor of TGF β receptor (ALK5) and an Activin receptor (ALK4/7) (i.e., TGF β R
35 inhibitor). One or more kinds of these may be contained as a

TGF β signal transduction pathway inhibiting substances. The TGF β signal transduction pathway inhibiting substance is preferably SB431542 or A-83-01.

[0086]

5 The Nodal/Activin signal transduction pathway inhibiting substance is not particularly limited as long as it is a substance inhibiting a signal transduction pathway caused by Nodal or Activin, and may be any of nucleic acid, protein and a low-molecule organic compound. Examples of the substance
10 include a substance that directly acts on Nodal or Activin (e.g., antibody, aptamer etc.), a substance that suppresses expression of gene encoding Nodal or Activin (e.g., antisense oligonucleotide, siRNA etc.), a substance that suppresses binding of Nodal/Activin receptor and Nodal/Activin, and a
15 substance that suppresses physiological activity caused by signal transduction by Nodal/Activin receptor. As the Nodal/Activin signal transduction pathway inhibiting substance, a compound well known to those of ordinary skill in the art can be used and, specifically, SB431542, A-83-01 and the like can
20 be mentioned. In addition, a protein (Lefty, Cerberus etc.) known as a Nodal/Activin signal transduction pathway inhibiting substance may be used. One or more kinds of these may be contained as a Nodal/Activin signal transduction pathway inhibiting substance. The Nodal/Activin signal transduction
25 pathway inhibiting substance is preferably SB431542, A-83-01 or Lefty.

[0087]

 The BMP signal transduction pathway inhibiting substance is not particularly limited as long as it a substance
30 inhibiting a signal transduction pathway caused by BMP, and may be any of nucleic acid, protein and a low-molecule organic compound. As BMP here, BMP2, BMP4, BMP7 and GDF7 can be mentioned. Examples of the substance include a substance that directly acts on BMP (e.g., antibody, aptamer etc.), a
35 substance that suppresses expression of gene encoding BMP (e.g.,

antisense oligonucleotide, siRNA etc.), a substance that suppresses binding of BMP receptor (BMPR) and BMP, and a substance that suppresses physiological activity caused by signal transduction by BMP receptor. As BMPR, ALK2 or ALK3 can
5 be mentioned. As the BMP signal transduction pathway inhibiting substance, a compound well known to those of ordinary skill in the art can be used and, specifically, LDN193189, Dorsomorphin and the like can be mentioned. Here, LDN193189 (4-[6-(4-piperazin-1-ylphenyl)pyrazolo[1,5-
10 alpyrimidin-3-yl]quinoline) is a known BMPR (ALK2/3) inhibitor (hereinafter BMPR inhibitor) and is generally commercially available in the form of hydrochloride. In addition, a protein known as a BMP signal transduction pathway inhibiting substance (Chordin, Noggin etc.) may also be used. One or more kinds of
15 these may be contained as a BMP signal transduction pathway inhibiting substance. The BMP signal transduction pathway inhibiting substance is preferably LDN193189.

[0088]

The TGF β family signal transduction pathway inhibiting
20 substance is preferably Lefty, SB431542, A-83-01 or LDN193189.

[0089]

Multiple kinds of TGF β family signal transduction pathway inhibiting substances with different points of action may be used in combination. By combining them, it is expected that
25 the improving effect on the quality of the aggregate will be enhanced. For example, a combination of a TGF β signal transduction pathway inhibiting substance and a BMP signal transduction pathway inhibiting substance, a combination of a TGF β signal transduction pathway inhibiting substance and
30 Nodal/Activin signal transduction pathway inhibiting substance, and a combination of a BMP signal transduction pathway inhibiting substance and a Nodal/Activin signal transduction pathway inhibiting substance can be mentioned. Preferably, a TGF β signal transduction pathway inhibiting substance and a BMP
35 signal transduction pathway inhibiting substance are used in

combination. A specific preferable combination is a combination of SB431542 and LDN193189.

[0090]

The Sonic hedgehog (hereinafter sometimes to be indicated
5 as Shh) signal transduction pathway activating substance is a substance capable of enhancing signal transduction mediated by Shh. Examples of the Shh signal transduction pathway activating substance include proteins belonging to the Hedgehog family (e.g., Shh and Ihh), Shh receptor, Shh receptor agonist,
10 PMA (Purmorphamine; 9-cyclohexyl-N-[4-(4-morpholinyl)phenyl]-2-(1-naphthalenyloxy)-9H-purin-6-amine), SAG (Smoothed Agonist; N-methyl-N'-(3-pyridinylbenzyl)-N'-(3-chlorobenzo[b]thiophene-2-carbonyl)-1,4-diaminocyclohexane) and the like. The Shh signal transduction pathway activating substance may contain
15 one or more kinds of these. A preferred Shh signal transduction pathway activating substances are Shh protein (Genbank accession numbers: NM_000193, NP_000184), SAG and PMA.
[0091]

A TGF β family signal transduction pathway inhibiting
20 substance and a Shh signal transduction pathway activating substance may be used in combination. By combining them, it is expected that the improving effect on the quality of the aggregate will be enhanced. Examples of the specific combination include a combination of any TGF β family signal
25 transduction pathway inhibiting substance selected from the group consisting of Lefty, SB431542, A-83-01 and LDN193189, and any Shh signal transduction pathway activating substance selected from the group consisting of Shh protein, SAG and PMA. When a TGF β family signal transduction pathway inhibiting
30 substance and a Shh signal transduction pathway activating substance are used in combination, the cells may be cultured in a medium containing both a TGF β family signal transduction pathway inhibiting substance and a Shh signal transduction pathway activating substance, or the cells may be treated with
35 any one of a TGF β family signal transduction pathway inhibiting

substance and a Shh signal transduction pathway activating substance and then the cells may be continuously treated with either or both of them. Alternatively, the cells may be treated with both a TGF β family signal transduction pathway
5 inhibiting substance and a Shh signal transduction pathway activating substance, and then continuously treated with either of them.

[0092]

The concentration of the TGF β family signal transduction
10 pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance can be appropriately determined to fall within a range capable of achieving the aforementioned effects. For example, SB431542 is generally used at a concentration of 0.1 - 200 μ M, preferably 2 - 50 μ M.
15 A-83-01 is generally used at a concentration of 0.05 - 50 μ M, preferably 0.5 - 5 μ M. LDN193189 is generally used at a concentration of 1 - 2000 nM, preferably 10 - 300 nM. Lefty is generally used at a concentration of 5 - 200 ng/ml, preferably 10 - 50 ng/ml. Shh protein is generally used at a
20 concentration of 20 - 1000 ng/ml, preferably 50 - 300 ng/ml. SAG is generally used at a concentration of 1 - 2000 nM, preferably 10 - 700 nM, more preferably 30 - 600 nM. PMA is generally used at a concentration of 0.002 - 20 μ M, preferably 0.02 - 2 μ M. In one embodiment, a TGF β family signal
25 transduction pathway inhibiting substance can be appropriately used in an amount showing a TGF β family signal transduction pathway inhibiting activity equivalent to that of SB431542 at the aforementioned concentration. In one embodiment, a Sonic hedgehog signal transduction pathway activating substance can
30 be appropriately used in an amount showing a Shh signal transduction promoting activity equivalent to that of SAG at the aforementioned concentration.

[0093]

The TGF β family signal transduction pathway inhibiting
35 activity of SB431542, LDN193189 and the like can be determined

by detecting, for example, phosphorylation of Smad by a method well known to those of ordinary skill in the art such as a Western blotting method (Mol Cancer Ther. (2004) 3, 737-45.). Sonic hedgehog signal transduction promoting activity of SAG
5 and the like can be determined by a method well known to those of ordinary skill in the art, for example, a reporter gene assay taking note of the expression of the Gli1 gene (Oncogene (2007) 26, 5163-5168).

[0094]

10 While the medium used for step (1) may be a serum-containing medium or a serum-free medium, it is preferably a serum-free medium, to avoid contamination with chemically-undefined components.

[0095]

15 To avoid contamination with a chemically-undefined component, a medium to be used for step (1) may be a medium whose components are chemically-defined.

[0096]

For culturing pluripotent stem cells under feeder-free
20 conditions in step (1), the aforementioned feeder-free medium (That is, undifferentiation maintenance medium) can be used as a medium. As a feeder-free medium (undifferentiation maintenance medium), Essential 8 medium, Essential 6 medium, Stabilized Essential 8 medium, S-medium, StemPro medium, hESF9
25 medium, TeSR medium, mTeSR medium (mTeSR1, mTeSR2 and the like), TeSR-E8, StemFit (registered trade mark) medium and the like can be mentioned, and Essential 8 medium or StemFit (registered trade mark) medium is preferably used.

[0097]

30 For culturing pluripotent stem cells under feeder-free conditions in step (1), an appropriate matrix may be used as a scaffold to provide a scaffold in stead of the feeder cells to the pluripotent stem cell. The pluripotent stem cells are subjected to adhesion culture in a cell container whose surface
35 is coated with a matrix as a scaffold.

[0098]

As a matrix available as a scaffold, laminin (Nat Biotechnol 28, 611-615 (2010)), laminin fragment (Nat Commun 3, 1236 (2012)), basement membrane preparation (Nat Biotechnol 19, 5 971-974 (2001)), gelatin, collagen, heparan sulfate proteoglycan, entactin, vitronectin and the like can be mentioned.

[0099]

"Laminin" is a heterotrimer molecule consisting of α , β , 10 γ chains and an extracellular matrix protein containing isoforms having different subunit chain compositions. Specifically, laminin has about 15 species of isoforms based on the combinations of heterotrimers with 5 species of α chains, 4 species of β chains and 3 species of γ chains. The name of 15 laminin is determined by combining respective numbers of α chain ($\alpha 1 - \alpha 5$), β chain ($\beta 1 - \beta 4$) and γ chain ($\gamma 1 - \gamma 3$). For example, a laminin having a combination of $\alpha 5$ chain, $\beta 1$ chain, $\gamma 1$ chain is named laminin 511. In the present invention, laminin 511 is preferably used (Nat Biotechnol 28, 611-615 20 (2010)).

[0100]

Laminin to be used in the present invention is generally a mammalian laminin. As the mammal, those mentioned above can be recited. To achieve xeno-free conditions, laminin of a 25 mammal of the same species as the cell to be cultured is preferably used. For example, human laminin (preferably, human laminin 511) is used for culturing human pluripotent stem cells.

[0101]

A laminin fragment to be used in the present invention is 30 not particularly limited as long as it has adhesiveness to pluripotent stem cells and enables maintenance culture of pluripotent stem cell under feeder-free conditions, and is preferably E8 fragment. Laminin E8 fragment was identified as a fragment with strong cell adhesion activity among the 35 fragments obtained by digestion of laminin 511 with elastase

(EMBO J., 3:1463-1468, 1984, J. Cell Biol., 105:589-598, 1987).
In the present invention, E8 fragment of laminin 511 is
preferably used (Nat Commun 3, 1236 (2012), Scientific Reports
4, 3549 (2014)). The laminin E8 fragment to be used in the
5 present invention is not required to be an elastase-digestion
product of laminin and may be a recombinant. To avoid
contamination of unidentified components, a recombinant laminin
fragment is preferably used in the present invention. An E8
fragment of laminin 511 is commercially available and can be
10 purchased from, for example, Nippi, Inc. and the like.

[0102]

The laminin or laminin fragment to be used in the present
invention is preferably isolated.

[0103]

15 The "basement membrane preparation" in the present
invention refers to one containing basement membrane-
constituting components having a function to control cell
morphology, differentiation, growth, motility, expression of
function and so on which are similar to those of epithelial
20 cell, when intended cells capable of forming a basement
membrane are plated thereon and cultured. For example, retinal
cells and retinal tissues produced by the present invention may
be dispersed, and cultured in the presence of a basement
membrane preparation when further adhesion culture is performed.
25 Here, the "basement membrane constituting components" refers to
extracellular matrix molecules in the form of a thin membrane
present between epithelial cell layer and interstitial cell
layer and so on in animal tissues. A basement membrane
preparation can be produced by, for example, removing cells
30 capable of forming a basement membrane, which adhere onto a
support via a basement membrane, from a support with a solution
capable of dissolving the lipid of the cells, an alkali
solution and so on. Examples of the basement membrane
preparation include products commercially available as basement
35 membrane preparation (e.g., Matrigel™ (manufactured by Corning

Incorporated: hereinafter sometimes referred to as Matrigel)),
Geltrex™ (manufactured by Life Technologies), and
extracellular matrix molecules known as basement membrane
components (e.g., laminin, type IV collagen, heparan sulfate
5 proteoglycan, entactin and so on).

[0104]

Matrigel™ is a basement membrane preparation extracted
from Engelbreth Holm Swarn (EHS) mouse sarcoma. The main
component of Matrigel™ is type IV collagen, laminin, heparan
10 sulfate proteoglycan, and entactin. In addition to these, TGF-
β, FGF, tissue plasminogen activator, and a growth factor
naturally produced by EHS tumor are contained. The "growth
factor reduced product" of Matrigel™ has a lower growth factor
concentration than common Matrigel™, and the standard
15 concentration thereof is <0.5 ng/ml for EGF, <0.2 ng/ml for NGF,
<5 pg/ml for PDGF, 5 ng/ml for IGF1, and 1.7 ng/ml for TGFβ.

[0105]

To avoid contamination of unidentified components, an
isolated laminin or laminin fragment is preferably used in the
20 present invention.

[0106]

Preferably, in the culture of pluripotent stem cells
under feeder-free conditions in step (1), the human pluripotent
stem cells are cultured in an adhered state in a cell container
25 with surface coated with isolated laminin 511 or E8 fragment of
laminin 511 (more preferably, E8 fragment of laminin 511).

[0107]

While the period for the culture of pluripotent stem
cells in step (1) is not particularly limited as long as the
30 effect of improving the quality of the aggregate formed in step
(2) can be achieved, it is generally 0.5 - 144 hr. The culture
period of pluripotent stem cells in step (1) is preferably not
less than 1 hr, not less than 2 hr, not less than 6 hr, not
less than 12 hr, not less than 18 hr, not less than 20 hr or
35 not less than 24 hr. The culture period of pluripotent stem

cells in step (1) is preferably within 96 hr, within 72 hr, within 60 hr, within 48 hr or within 28 hr. In one embodiment, the range of the culture period of pluripotent stem cells in step (1) is preferably 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60
5 hr, 18 - 60 hr, 18 - 48 hr or 18 - 28 hr (e.g., 24 hr). That is, the first step is started 0.5 - 144 hr (preferably, 12 - 60 hr, 18 - 48 hr or 18 - 28 hr) before the start of the suspension culture in step (2), and step (2) is continuously performed after completion of step (1). When the cells are
10 treated with either of a TGF β family signal transduction pathway inhibiting substance and a Shh signal transduction pathway activating substance and then continuously treated with the other, the treatment time of each can be set to independently fall within the range of the aforementioned
15 culture period.

[0108]

When the cells are treated with both of the TGF β family signal transduction pathway inhibiting substance and the Shh signal transduction pathway activating substance and then
20 continuously treated with one of them, and when the cells are treated with either of the TGF β family signal transduction pathway inhibiting substance and the Shh signal transduction pathway activating substance and then continuously treated with the both, the treatment time of each can also be set to
25 independently fall within the range of the aforementioned culture period.

[0109]

In one embodiment, the cells can be cultured with either of the TGF β family signal transduction pathway inhibiting
30 substance (e.g., SB431542 or LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and then further cultured with either of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542 or LDN193189) and the Shh signal
35 transduction pathway activating substance (e.g., SAG) for 18 -

28 hr (e.g., 24 hr). In this case, the concentration of the TGF β family signal transduction pathway inhibiting substance is, for example, a concentration showing a TGF β signal transduction pathway inhibiting activity equivalent to that of SB431542 at 3 μ M - 10 μ M, or a concentration showing a BMP signal transduction pathway inhibiting activity equivalent to that of LDN193189 at 50 nM - 200 nM, and a concentration of the Shh signal transduction pathway activating substance is, for example, a concentration showing a signal transduction promoting activity equivalent to that of SAG at 100 nM - 500 nM. [0110]

In another embodiment, the cells can be cultured with both of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542 or LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and then further cultured with either of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542 or LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr). In this case, the concentration of the TGF β family signal transduction pathway inhibiting substance is, for example, a concentration showing a TGF β signal transduction pathway inhibiting activity equivalent to that of SB431542 at 3 μ M - 10 μ M, or a concentration showing a BMP signal transduction pathway inhibiting activity equivalent to that of LDN193189 at 50 nM - 200 nM, and a concentration of the Shh signal transduction pathway activating substance is, for example, a concentration showing a signal transduction promoting activity equivalent to that of SAG at 100 nM - 500 nM. [0111]

In another embodiment, the cells can be cultured with either of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542 or LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and then further cultured with

both of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542 or LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr). In this case, the concentration of the
5 TGF β family signal transduction pathway inhibiting substance is, for example, a concentration showing a TGF β signal transduction pathway inhibiting activity equivalent to that of SB431542 at 3 μ M - 10 μ M, or a concentration showing a BMP signal transduction pathway inhibiting activity equivalent to that of
10 LDN193189 at 50 nM - 200 nM, and a concentration of the Shh signal transduction pathway activating substance is, for example, a concentration showing a signal transduction promoting activity equivalent to that of SAG at 100 nM - 500 nM.
[0112]

15 The culture conditions such as culture temperature, and CO₂ concentration in step (1) can be appropriately determined. While the culture temperature is, for example, about 30°C to about 40°C, preferably about 37°C. The CO₂ concentration is, for example, about 1% to about 10%, preferably about 5%.

20 [0113]

In one preferable embodiment, human pluripotent stem cells (e.g., human iPS cells) are cultured in an adhered state in the absence of feeder cells and in a serum-free medium (undifferentiation maintenance medium) containing bFGF. The
25 adhesion culture is preferably performed in a cell container with surface coated with laminin 511, E8 fragment of laminin 511 or vitronectin. The adhesion culture is preferably performed using Essential 8, TeSR medium, mTeSR medium, mTeSR-E8 medium, or StemFit (registered trade mark) medium, more
30 preferably Essential 8 or StemFit (registered trade mark) medium, as a feeder-free medium (undifferentiation maintenance medium).

[0114]

In one embodiment, human pluripotent stem cells (e.g.,
35 human iPS cells) are cultured in suspension in the absence of

feeder cells and in a serum-free medium containing bFGF. In the suspension culture, human pluripotent stem cells may form an aggregate of human pluripotent stem cells.

[0115]

5 In a preferable embodiment, the cell obtained in step (1) is a cell maintaining pluripotent-like properties (pluripotent-like state) and the pluripotent-like properties are maintained throughout step (1). The pluripotent-like properties mean that at least a part of the characteristics unique to and common to
10 the pluripotent stem cells including pluripotency is maintained. The pluripotent-like properties do not require strict pluripotency. Specifically, a state expressing all or some of the markers to be the index of pluripotent properties (pluripotent state) is included in the "pluripotent-like
15 properties". As a marker of the pluripotent-like properties, Oct3/4 positive, alkaline phosphatase positive and the like can be mentioned. In one embodiment, a cell maintaining the pluripotent-like properties is Oct3/4 positive. Even if the expression level of Nanog is lower than that of ES cell or iPS
20 cell, it is included in "cell showing pluripotent-like properties".

[0116]

In one embodiment, the cells obtained in step (1) are stem cells having an ability to differentiate into at least a
25 retinal cell or retinal tissue (preferably, retinal tissue, retinal progenitor cell, or retinal layer-specific neural cell). In one embodiment, the cells obtained in step (1) are Oct3/4 positive stem cells having an ability to differentiate into at least a retinal cell or retinal tissue (preferably, retinal
30 tissue, retinal progenitor cell or retinal layer-specific neural cell). In one embodiment, the cells obtained in step (1) contain not less than 60%, for example, not less than 90%, of Oct3/4 positive stem cells.

[0117]

35 In a preferable embodiment, in step (1), human

pluripotent stem cells (e.g., iPS cell, ES cell) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium containing a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and bFGF.

[0118]

The above-mentioned adhesion culture in step (1) is preferably performed in a cell container with a surface coated with laminin 511 or E8 fragment of laminin 511. The TGF β family signal transduction pathway inhibiting substance is preferably a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty), a Nodal/Activin signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189, Chordin, Noggin), or a combination of these (e.g., SB431542 and LDN193189). The TGF β family signal transduction pathway inhibiting substance is further preferably Lefty, SB431542, A-83-01, or LDN193189, or a combination of these (e.g., SB431542 and LDN193189). The Sonic hedgehog signal transduction pathway activating substance is preferably Shh protein, SAG or Purmorphamine (PMA), more preferably SAG. The TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and the Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) may be used in combination. The culture period is 0.5 - 144 hr (preferably 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60 hr, 18 - 60 hr, 18 - 48 hr or 18 - 28 hr (e.g., 24 hr)). In the course of the culture, the TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189), the Shh signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) or the combination thereof may be changed. In one embodiment, the cells can be cultured with both of a TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and a Shh signal transduction

pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and then further cultured with either of a TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and a Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr). In another embodiment, the cells can be cultured with either of a TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and a Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and then further cultured with both or either of a TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and a Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr). [0119]

For example, pluripotent stem cells are subjected to maintenance culture in the absence of feeder cells in a medium containing a factor for maintaining undifferentiated state before the start of step (1), a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance is/are added to the culture and culturing is continued to perform step (1). Specifically, for example, when step (1) is started after completion of maintenance culture, an undifferentiation maintenance medium added with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance may be added or a part or all of the undifferentiation maintenance medium may be exchanged and culturing is continued. [0120]

For example, human pluripotent stem cells (e.g., human iPS cells) are subjected to a maintenance culture in the absence of feeder cells and in a serum-free medium containing bFGF. The maintenance culture is preferably performed in adhesion culture. The adhesion culture is preferably performed in a cell container with a surface coated with vitronectin,

laminin 511 or E8 fragment of laminin 511. The period of maintenance culture is not particularly limited. A pluripotent stem cell is a cell capable of proliferation (preferably autonomously replicating) maintaining differentiation potency,
5 and maintenance culture is a culture method capable of maintaining pluripotency. Thus, infinite maintenance culture is possible if properly operated. In some embodiments, during maintenance culture, pluripotent stem cells are frozen to produce a frozen cell stock, and the frozen cell stock is
10 cultivated and maintenance culture can be continued. While the culture period after the cultivation is not particularly limited, it is, for example, about 1 day - 1000 days, preferably about 7 days - 150 days, more preferably about 10 days - 90 days. In some embodiments, passage operation of the
15 cells can be performed during maintenance culture.

[0121]

In step (1), a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance are/is added to a
20 medium containing human pluripotent stem cells subjected to a maintenance culture as mentioned above and culturing is continued. The TGF β family signal transduction pathway inhibiting substance is preferably a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty),
25 a Nodal/Activin signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination thereof (e.g., SB431542 and LDN193189). The TGF β family signal transduction pathway inhibiting substance is
30 further preferably Lefty, SB431542, A-83-01, LDN193189, or a combination thereof (e.g., SB431542 and LDN193189). The Sonic hedgehog signal transduction pathway activating substance is preferably Shh protein, SAG or PMA. The TGF β family signal transduction pathway inhibiting substance (e.g., Lefty,
35 SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal

transduction pathway activating substance (e.g., Shh protein, SAG, PMA) may be used in combination. After addition, culturing is continued for 0.5 - 144 hr (preferably, 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60 hr, 18 - 60 hr, 18 - 48 hr, or 18
5 - 28 hr (e.g., 24 hr)). The TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189), the Shh signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) or a combination thereof may be changed during the culture. In one embodiment,
10 cells can be cultured using both the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and further cultured using either of the TGF β family signal transduction
15 pathway inhibiting substance (e.g., SB431542, LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr). In another embodiment, cells can be cultured using either of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542,
20 LDN193189) and the Shh signal transduction pathway activating substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr) and further cultured using either or both of the TGF β family signal transduction pathway inhibiting substance (e.g., SB431542, LDN193189) and the Shh signal transduction pathway activating
25 substance (e.g., SAG) for 18 - 28 hr (e.g., 24 hr).

[0122]

2-2. Step (2)

The step (2) for culturing the cells obtained in step (1) in suspension in a medium containing a Wnt signal transduction
30 pathway inhibiting substance to form a cell aggregate is explained.

[0123]

The time point when the suspension culture is started in step (2) of the present invention is the time point when the
35 following operation is performed.

.operation: operation including medium exchange of undifferentiation maintenance medium (that is, medium containing 1) a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal
5 transduction pathway activating substance, and 2) a factor for maintaining undifferentiated state) used in step (1) and proceeding to suspension culture.

The medium exchange in the above-mentioned operation is exchange with a medium other than the undifferentiation
10 maintenance medium and is not particularly limited as long as the medium is capable of inducing differentiation into the retinal cell and/or the retinal tissue of the present invention, or an intermediate thereof. For example, a medium not containing an "undifferentiation maintenance factor" such as a
15 basal medium to be described later and the like can be mentioned. It may or may not contain a Wnt signal transduction pathway inhibiting substance at the time point of the start of step (2), that is, at the time point of medium exchange, and a Wnt signal transduction pathway inhibiting substance may be
20 added after medium exchange as described later.

[0124]

The medium (basal medium) to be used in step (2) is not particularly limited and the basal medium described in the above-mentioned section of definition can be appropriately
25 selected. The medium to be used in step (2) may be a serum-containing medium or serum-free medium. To avoid contamination of chemically-undefined components, a serum-free medium is preferably used in the present invention. To avoid complicated preparation, for example, a serum-free medium supplemented with
30 an appropriate amount of a commercially available serum alternative such as KSR and so on (e.g., medium of 1:1 mixture of IMDM and F-12, which is supplemented with 10% KSR, 450 μ M 1-monothioglycerol and 1x Chemically Defined Lipid Concentrate, or GMEM medium supplemented with 5% - 20% KSR, NEAA, pyruvic
35 acid, 2-mercaptoethanol) is preferably used. The amount of KSR

to be added to a serum-free medium in the case of human pluripotent stem cell is generally about 1% to about 30%, preferably about 2% to about 20% (e.g., about 5%, about 10%).
[0125]

5 The medium used in step (2) contains a Wnt signal transduction pathway inhibiting substance. In step (1), pluripotent stem cells are treated with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and,
10 in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium (preferably serum-free medium) containing a Wnt signal transduction pathway inhibiting substance to form a cell aggregate, whereby the quality of the aggregates is further improved, and cell aggregates maintaining
15 an undifferentiated state and aggregates suitable for differentiation into retinal cells can be formed with high efficiency. The thus-obtained cell aggregates are expected to show characteristics of, for example, being spherical and having a smooth surface, an uncollapsed shape and dense inside.
20 [0126]

 The Wnt signal transduction pathway inhibiting substance to be used in step (2) is not particularly limited as long as it can suppress signal transduction mediated by Wnt and may be any of protein, nucleic acid, low-molecular-weight compound and
25 the like. The signal mediated by Wnt is transduced via Wnt receptors that exist as heterodimers of Frizzled (Fz) and LRP5/6 (low-density lipoprotein receptor-related protein 5/6). Examples of the Wnt signal transduction pathway inhibiting substance include, but are not limited to, a substance that
30 directly acts on Wnt or Wnt receptor (anti-Wnt neutralizing antibody, anti-Wnt receptor neutralizing antibody etc.), a substance that suppresses expression of gene encoding Wnt or Wnt receptor (e.g., antisense oligonucleotide, siRNA etc.), a substance that suppresses binding of Wnt receptor and Wnt
35 (soluble Wnt receptor, dominant negative Wnt receptor etc., Wnt

antagonist, Dkk1, Cerberus protein etc.), and a substance that suppresses physiological activity caused by signal transduction by Wnt receptor [low-molecular-weight compounds such as CKI-7 (N-(2-aminoethyl)-5-chloroisoquinoline-8-sulfonamide), D4476
5 (4-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)-5-(2-pyridinyl)-1H-imidazol-2-yl]benzamide), IWR-1-endo (IWR1e) (4-[(3aR,4S,7R,7aS)-1,3,3a,4,7,7a-hexahydro-1,3-dioxo-4,7-methano-2H-isoindol-2-yl]-N-8-quinolinyl-benzamide), and IWP-2 (N-(6-methyl-2-benzothiazolyl)-2-[(3,4,6,7-tetrahydro-4-oxo-3-phenylthieno[3,2-d]pyrimidin-2-yl)thio]acetamide) and the like,
10 and the like] and the like. One or more kinds of these may be contained as a Wnt signal transduction pathway inhibiting substance. CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2 and the like are known Wnt signal transduction pathway inhibiting
15 substances, and commercially available products and the like are appropriately available. IWR1e is preferably used as a Wnt signal transduction pathway inhibiting substance.
[0127]

The concentration of the Wnt signal transduction pathway
20 inhibiting substance only needs to be a concentration capable of inducing formation of a good cell aggregate. For example, IWR-1-endo is added to a medium such that the concentration is about 0.1 μM to about 100 μM , preferably about 0.3 μM to about 30 μM , more preferably about 1 μM to about 10 μM , further
25 preferably about 3 μM . When a Wnt signal transduction pathway inhibiting substance other than IWR-1-endo is used, it is desirably used at a concentration exhibiting a Wnt signal transduction pathway inhibiting activity equivalent to that of IWR-1-endo at the above-mentioned concentration.

30 [0128]

The timing of addition of a Wnt signal transduction pathway inhibiting substance to the medium is not particularly limited as long as the above-mentioned effects can be afforded, but a higher effect can be obtained when it is added earlier.
35 A Wnt signal transduction pathway inhibiting substance is added

to the medium generally within 6 days, preferably within 3 days, more preferably within 1 day, more preferably within 12 hours, from the start of the suspension culture in step (2), and further preferably at the time of the start of the suspension
5 culture in step (2). Specifically, for example, it is possible to add a basal medium to which a Wnt signal transduction pathway inhibiting substance is added, or exchange a part or the whole of the medium into the basic medium. While the period during which the cells obtained in step (1) are treated
10 with the Wnt signal transduction pathway inhibiting substance in step (2) is not particularly limited as long as the above-mentioned effect can be achieved, preferably, the substance is added to the medium when the suspension culture is started in step (2) and acted until the end of step (2) (immediately
15 before addition of the BMP signal transduction pathway activating substance). Further preferably, as mentioned later, the cells are continuously exposed to the Wnt signal transduction pathway inhibiting substance even after completion of step (2) (that is, during the period of step (3)). In one
20 embodiment, as described later, the Wnt signal transduction pathway inhibiting substance may be allowed to continuously act even after completion of step (2) (that is, during the period of step (3)) until a neuroepithelial tissue and/or a neural tissue are/is formed. Formation of a neuroepithelial tissue
25 and/or a neural tissue can be confirmed by the above-described method.

[0129]

For formation of an aggregate, a dispersing operation of the cells obtained in step (1) is first performed to give
30 dispersed cells. The "dispersed cells" obtained by the dispersing operation refers to a state where, for example, not less than 70% of cells are single cells and not more than 30% of cells are clumps of 2 - 50 cells. Preferably, as the dispersed cells, a state where not less than 80% of cells are
35 single cells, and not more than 20% of cells are clumps of 2 -

50 cells can be mentioned. The dispersed cells refer to a state almost free of mutual adhesion of cells (e.g., plane attachment). In some embodiments, dispersed cells refer to a state almost free of cell-cell junction (e.g., adhesive bond).

5 [0130]

A dispersion operation of the cells obtained in step (1) may contain the above-mentioned mechanical dispersion treatment, cell dispersion solution treatment, and cell protecting agent treatment. These treatments may be performed in combination.

10 Preferably, a cell dispersion solution treatment is performed simultaneously with a cell protecting agent treatment and then a mechanical dispersion treatment is performed.

[0131]

As a cell protecting agent to be used for the cell
15 protecting agent treatment, an FGF signal transduction pathway activating substance, serum, and serum alternative can be mentioned. Also, as a cell protecting agent to suppress cell death of pluripotent stem cells (particularly, cell death of human pluripotent stem cells) induced by dispersion, a Rho-
20 associated coiled-coil kinase (ROCK) inhibitor or a Myosin inhibitor may be added. To suppress cell death of pluripotent stem cells (particularly, human pluripotent stem cells) induced by dispersion, and protect the cells, a Rho-associated coiled-coil kinase (ROCK) inhibitor or a Myosin inhibitor may be added
25 from the start of the second step culture. As a ROCK inhibitor, Y-27632, Fasudil (HA1077), H-1152 and the like can be mentioned. As a Myosin inhibitor, Blebbistatin can be mentioned.

[0132]

As a cell dispersion solution to be used for the cell
30 dispersion solution treatment, a solution containing any of enzymes such as trypsin, collagenase, hyaluronidase, elastase, pronase, DNase, papain and so on, and a chelating agent such as ethylenediaminetetraacetic acid and so on can be mentioned. A commercially available cell dispersion solution such as TrypLE
35 Select (manufactured by Life Technologies) and TrypLE Express

(manufactured by Life Technologies) can also be used.

[0133]

As a method of mechanical dispersion treatment, a pipetting treatment or scraping by a scraper can be mentioned.

5 [0134]

The dispersed cells are suspended in the above-mentioned medium. For example, the dispersed cells are suspended in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (and a cell protecting agent such as a
10 ROCK inhibitor and the like as necessary), whereby a treatment with a Wnt signal transduction pathway inhibiting substance is possible from the start of the suspension culture in step (2).

[0135]

Then, a suspension of the dispersed cells is seeded in
15 the above-mentioned culture vessel, and the dispersed cells are cultured under a condition non-adhesive to the culture vessel, whereby plural cells are gathered to form an aggregate.

[0136]

In this case, plural cell aggregates may be
20 simultaneously formed in one culture vessel by seeding the dispersed cells in a comparatively large culture vessel such as a 10 cm dish. However, the size of the aggregates varies in this case. Thus, for example, a given number of dispersed cells are placed in each well of a multiwell plate (U-bottom,
25 V-bottom, M-bottom) such as a 96-well microplate, and static culture is performed, whereby the cells are rapidly aggregated to form one aggregate in each well. The aggregates are recovered from plural wells, whereby a population of uniformed aggregates can be obtained.

30 [0137]

The concentration of the cells in step (2) can be appropriately set so that cell aggregates can be more uniformly and efficiently formed. For example, when human cells (e.g., cells obtained from human iPS in step (1)) are cultured in
35 suspension using a 96-well microwell plate, a liquid prepared

to achieve about 1×10^3 cells to about 1×10^5 cells,
preferably about 3×10^3 cells to about 5×10^4 cells, more
preferably about 4×10^3 cells to about 2×10^4 cells, further
preferably about 4×10^3 cells to about 1.6×10^4 cells,
5 further more preferably about 8×10^3 cells to about 1.2×10^4
cells, per well is added to the wells, and the plate is stood
to form aggregates.

[0138]

The culture conditions such as culture temperature, CO_2
10 concentration and so on in step (2) can be appropriately
determined. The culture temperature is, for example, about
 30°C to about 40°C , preferably about 37°C . The CO_2
concentration is, for example, about 1% to about 10%,
preferably about 5%.

15 [0139]

In step (2), when a medium exchange operation is
performed, for example, an operation to add a fresh medium
without discarding the existing medium (medium addition
operation), an operation to discard about a half amount of the
20 existing medium (about 30 - 90%, for example, 40 - 60% of the
volume of the existing medium) and add about a half amount of a
fresh medium (30 - 90%, for example, about 40 - 60% of the
volume of the existing medium) (half-medium exchange operation),
and an operation to discard about the whole amount of the
25 existing medium (not less than 90% of the amount of the
existing medium) and add about the whole amount of a fresh
medium (not less than 90% of the amount of the existing medium)
(full-medium exchange operation) can be mentioned.

[0140]

30 When a particular component (e.g., differentiation-
inducing factor) is added at a certain time point, for example,
an operation to calculate the final concentration, to discard
about a half amount of the existing medium, and to add about a
half amount of a fresh medium containing a particular component
35 at a concentration higher than the final concentration,

specifically 1.5 times - 3.0 times the final concentration, for example, about 2 times the final concentration (half-medium exchange operation, half-medium exchange) may be performed.
[0141]

5 When the concentration of a particular component contained in the existing medium is maintained at a certain time point, for example, an operation to discard about a half amount of the existing medium and to add about a half amount of a fresh medium containing the particular component at a
10 concentration same as that in the existing medium may be performed.
[0142]

 When the concentration of a component contained in the existing medium is to be decreased by dilution at a certain
15 time pointed, for example, the medium exchange operation may be performed plural times per day, preferably plural times (e.g., 2 - 3 times) within 1 hr. Also, when the concentration of a component contained in the existing medium is to be decreased by dilution at a certain time point, the cells or aggregates
20 may be transferred to another culture container.
[0143]

 While the tool used for the medium exchange operation is not particularly limited, for example, pipetter, micropipette, multichannel micropipette, continuous dispenser, and the like
25 can be mentioned. For example, when a 96 well plate is used as a culture vessel, a multichannel micropipette may be used.
[0144]

 The period for suspension culture necessary for forming a cell aggregate can be determined as appropriate according to
30 the cell to be used, so that the cells can be aggregated uniformly. To form uniformed cell aggregates, it is desirably as short as possible. The steps for the dispersed cells to form cell aggregates can be divided into a step for gathering cells, and a step for forming cell aggregates from the gathered
35 cells. In a step of seeding the dispersed cells (i.e., at the

time of the start of suspension culture) to gather the cells in case of human pluripotent stem cells (e.g., stem cells obtained from human iPS in step (1)), for example, the gathered cells are formed preferably within about 24 hr, more preferably
5 within about 12 hr. The period from the time point of seeding the dispersed cells (i.e., at the time of the start of suspension culture) to that of forming an aggregate in the case of human pluripotent stem cells (e.g., human iPS cells), the aggregate is formed, for example, preferably within about 72 hr,
10 more preferably within about 48 hr. The period for cell aggregate formation can be appropriately adjusted by controlling the tools for aggregating the cells, centrifugation conditions and so on.

[0145]

15 Formation of cell aggregates and uniformity thereof can be determined based on the size and cell number of the aggregates, macroscopic morphology, microscopic morphology and uniformity thereof by tissue staining analysis, expression of markers for differentiation and undifferentiated state and
20 uniformity thereof, control of expression of differentiation marker and synchronism thereof, reproducibility of differentiation efficiency between the aggregates, and so on.

[0146]

After aggregate formation, the aggregate may be
25 continuously cultured as it is. The period for suspension culture in step (2) is generally 24 hr - 6 days, preferably 24 hr - 3 days, further preferably about 24 hr - 48 hr.

[0147]

In one embodiment, the medium used in step (2) may
30 further contain a Sonic hedgehog signal transduction pathway activating substance. In step (1), pluripotent stem cells are treated with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance and, in step (2), the
35 cells obtained in step (1) are subjected to suspension culture

in a medium (preferably serum-free medium) containing a Wnt signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance to form a cell aggregate, whereby it is expected that the quality
5 of the aggregate is further improved, and cell aggregates maintaining an undifferentiated state (e.g., a spherical, smooth surfaced, uncollapsed, and dense inside aggregate of cells maintaining undifferentiated properties) can be formed with high efficiency.

10 [0148]

As the Sonic hedgehog signal transduction pathway activating substance, those mentioned above can be used. Preferably, the Sonic hedgehog signal transduction pathway activating substance is SAG, Purmorphamine (PMA) or Shh protein.
15 In one embodiment, a Sonic hedgehog signal transduction pathway activating substance of the same kind as that used in step (1) can be used in step (2). The concentration of the Sonic hedgehog signal transduction pathway activating substance in the medium can be appropriately determined to fall within a
20 range capable of achieving the aforementioned effects. SAG is used at a concentration of generally 1 - 2000 nM, preferably 10 nM - 1000 nM, more preferably 10 nM - 700 nM, further preferably 50 nM - 700 nM, further preferably 100 nM - 600 nM, further preferably 100 nM - 500 nM, in step (2). In another
25 embodiment, SAG is used at a concentration of generally 1 - 2000 nM, preferably 3 nM - 100 nM, more preferably 5 nM - 50 nM, further preferably 10 nM - 30 nM, in step (2). PMA is used at a concentration of generally 0.002 - 20 μ M, preferably 0.02 - 2 μ M, in step (2). Shh protein is used at a concentration of
30 generally 20 - 1000 ng/ml, preferably 50 - 300 ng/ml, in step (2). When a Sonic hedgehog signal transduction pathway activating substance other than SAG, PMA, Shh protein is used, it is desirably used at a concentration showing a Sonic hedgehog signal transduction promoting activity equivalent to
35 that of SAG at the above-mentioned concentration (e.g., 5 nM -

50 nM).

[0149]

The timing of addition of a Sonic hedgehog signal transduction pathway activating substance to the medium is not particularly limited as long as the above-mentioned effects can be afforded, but a higher effect can be obtained when it is added earlier. A Sonic hedgehog signal transduction pathway activating substance is added to the medium generally within 6 days, preferably within 3 days, more preferably within 1 day, more preferably within 12 hr, from the start of suspension culture in step (2), and further preferably at the time of the start of suspension culture in step (2). The Sonic hedgehog signal transduction pathway activating substance may be contained in the medium for any period (e.g., about 24 hr, about 48 hr, until the end of step (2)) up to the completion of step (2) (immediately before addition of a BMP signal transduction pathway activating substance). In one embodiment, a Sonic hedgehog signal transduction pathway activating substance is added simultaneously with a Wnt signal transduction pathway inhibiting substance and contained in the medium for the same period in step (2) (e.g., until the end of step (2)).

[0150]

In one embodiment, the medium used in step (2) may further contain a TGF β signal transduction pathway inhibiting substance. In step (1), pluripotent stem cells are treated with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance and, in the second step, the cells obtained in the first step are subjected to suspension culture in a medium (preferably serum-free medium) containing a Wnt signal transduction pathway inhibiting substance and a TGF β family signal transduction pathway inhibiting substance or a medium (preferably serum-free medium) containing a Wnt signal transduction pathway inhibiting substance, a Sonic hedgehog

signal transduction pathway activating substance and a TGF β family signal transduction pathway inhibiting substance to form a cell aggregate, whereby it is expected that the quality of the aggregate is further improved, and cell aggregates
5 maintaining an undifferentiated state (e.g., a spherical, smooth surfaced, uncollapsed, and dense inside aggregate of cells maintaining undifferentiated properties) can be formed with high efficiency.

[0151]

10 As the TGF β signal transduction pathway inhibiting substance, those mentioned above can be used. Preferably, the TGF β signal transduction pathway inhibiting substance is SB431542 or A83-01. In one embodiment, a TGF β signal transduction pathway inhibiting substance of the same kind as
15 that used in step (1) can be used in step (2). The concentration of the TGF β signal transduction pathway inhibiting substance in the medium can be appropriately determined to fall within the range capable of achieving the aforementioned effect. SB431542 is used at a concentration of
20 generally 0.1 - 200 μ M, preferably 2 - 50 μ M, more preferably 3 - 10 μ M. A-83-01 is used at a concentration of generally 0.05 - 50 μ M, preferably 0.5 - 5 μ M. When a TGF β signal transduction pathway inhibiting substance other than SB431542 or A-83-01 is used, it is desirably used at a concentration showing a TGF β
25 signal transduction pathway inhibiting activity equivalent to that of SB431542 at the above-mentioned concentration.

[0152]

The timing of addition of a TGF β signal transduction pathway inhibiting substance to the medium is not particularly
30 limited as long as the above-mentioned effects can be afforded, but a higher effect can be obtained when it is added earlier. A TGF β signal transduction pathway inhibiting substance is added to the medium generally within 6 days, preferably within 3 days, more preferably within 1 day, more preferably within 12
35 hours, from the start of the suspension culture in step (2),

and further preferably at the time of the start of the suspension culture in step (2). In step (2), the TGF β signal transduction pathway inhibiting substance is contained in the medium for any period (e.g., about 24 hr, about 48 hr, until
5 the end of step (2)) up to the completion of step (2) (immediately before addition of the BMP signal transduction pathway activating substance). In one embodiment, a TGF β signal transduction pathway inhibiting substance is added simultaneously with a Wnt signal transduction pathway
10 inhibiting substance and contained in the medium for the same period (e.g., until the end of step (2)).
[0153]

The medium used in step (2) may be substantially free of a TGF β signal transduction pathway inhibiting substance. In
15 the present specification, when the medium used in step (1) contains a TGF β signal transduction pathway inhibiting substance, and when the concentration of the TGF β signal transduction pathway inhibiting substance carried over from step (1) to step (2) during medium exchange (e.g., medium
20 exchange of about total amount) is suppressed to a level at which the biological activity (namely, TGF β signal transduction pathway inhibiting activity) thereof is not expressed (e.g., not more than 5 nM for SB431542), it can be regarded as a medium substantially free of a TGF β signal transduction pathway
25 inhibiting substance. In the present specification, substances other than the TGF β signal transduction pathway inhibiting substance are also interpreted similarly. For example, by a medium exchange operation of a medium free of a TGF β signal transduction pathway inhibiting substance, carrying over of the
30 TGF β signal transduction pathway inhibiting substance (e.g., SB431542) from step (1) is suppressed to not more than 3%, preferably not more than 1%, not more than 0.3%, not more than 0.1%, not more than 0.03%, or not more than 0.01% of the concentration of the TGF β signal transduction pathway
35 inhibiting substance (e.g., SB431542) in the medium used in

step (1) and the concentration of a TGF β signal transduction pathway inhibiting substance in the medium is suppressed to a level at which the biological activity (i.e., TGF β signal transduction pathway inhibiting activity) thereof is not
5 expressed (e.g., not more than 5 nM for SB431542), whereby a medium substantially free of a TGF β signal transduction pathway inhibiting substance can be obtained. Carrying over of the component in the medium to a fresh medium can be suppressed to the minimum by, during medium exchange (e.g., medium exchange
10 of about total amount), removing the old medium and pre-washing the cells and the culture container with a fresh medium one or plural times.

[0154]

The medium used in step (2) may be substantially free of
15 a Sonic hedgehog signal transduction pathway activating substance. In the present specification, when the medium used in step (1) contains a Sonic hedgehog signal transduction pathway activating substance, and when the concentration of the Sonic hedgehog signal transduction pathway activating substance
20 carried over from step (1) to step (2) during medium exchange (e.g., medium exchange of about total amount) is suppressed to a level at which the biological activity (namely, Sonic hedgehog signal transduction pathway activation) thereof is not expressed (e.g., not more than 0.03 nM for SAG), it can be
25 regarded as a medium substantially free of a Sonic hedgehog signal transduction pathway activating substance. For example, by a medium exchange operation of a medium free of a Sonic hedgehog signal transduction pathway activating substance, carrying over of the Sonic hedgehog signal transduction pathway
30 activating substance (e.g., SAG) from step (1) is suppressed to not more than 3%, preferably not more than 1%, not more than 0.3%, not more than 0.1%, not more than 0.03%, or not more than 0.01% of the concentration of the Sonic hedgehog signal transduction pathway activating substance (e.g., SAG) in the
35 medium used in step (1) and the concentration of a Sonic

hedgehog signal transduction pathway activating substance in the medium is suppressed to a level at which the biological activity (i.e., Sonic hedgehog signal transduction pathway activation) thereof is not expressed (e.g., not more than 0.03 nM for SAG), whereby a medium substantially free of a Sonic hedgehog signal transduction pathway activating substance can be obtained. Carrying over of the component in the medium to a fresh medium can be suppressed to the minimum by, during medium exchange (e.g., medium exchange of about total amount), removing the old medium and pre-washing the cells and the culture container with a fresh medium one or plural times. [0155]

In a preferable embodiment, in step (2), the human cells obtained in step (1) (e.g., cells obtained from human iPS cells in step (1)) are subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) to form an aggregate. The medium may further contain a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01). The time of addition of the Wnt signal transduction pathway inhibiting substance and that of the Sonic hedgehog signal transduction pathway activating substance and/or the TGF β signal transduction pathway inhibiting substance may be the same or different. Preferably, a Wnt signal transduction pathway inhibiting substance, a Sonic hedgehog signal transduction pathway activating substance and/or a TGF β signal transduction pathway inhibiting substance are both preferably contained in the medium from the time of the start of suspension culture. A ROCK inhibitor (e.g., Y-27632) may also be added to the medium. The period for the culture in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

[0156]

For example, the human cells obtained in step (1) (e.g., cells obtained from human iPS cells in step (1)) are recovered, dispersed to single cells or a state close thereto in a serum-free medium containing a Wnt signal transduction pathway
5 inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2), and a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01) and subjected to suspension culture.
10 The serum-free medium may contain a ROCK inhibitor (e.g., Y-27632). A suspension of human pluripotent stem cells (e.g., human iPS cells) is seeded in the above-mentioned culture vessel and the dispersed pluripotent stem cells are cultured under conditions where they are non-adhesive to the culture
15 vessel, whereby plural pluripotent stem cells are assembled to form an aggregate. The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.
[0157]

20 In one preferable embodiment, in step (1), pluripotent stem cells are cultured (preferably, adhesion culture) in a medium containing a TGF β signal transduction pathway inhibiting substance and containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG,
25 PMA, Shh protein) and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and containing or not containing a Sonic hedgehog signal transduction pathway
30 activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01). Preferably, SB431542 can be used as a TGF β signal transduction pathway inhibiting substance in step (1). The period for the culturing in step (2) is 24 hr - 6
35 days, preferably 24 hr - 48 hr. The aggregates formed are

preferably uniformed aggregates.

[0158]

In one preferable embodiment, in step (1), pluripotent stem cells are cultured (preferably, adhesion culture) in a medium containing a BMP signal transduction pathway inhibiting substance and containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01). Preferably, LDN193189 or Dorsomorphin, more preferably LDN193189, can be used as a BMP signal transduction pathway inhibiting substance in step (1). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

[0159]

In one preferable embodiment, in step (1), pluripotent stem cells are cultured (preferably, adhesion culture) in a medium containing a Sonic hedgehog signal transduction pathway activating substance and containing or not containing a TGF β signal transduction pathway inhibiting substance and/or a BMP signal transduction pathway inhibiting substance and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01). Preferably, SAG can be used as a Sonic hedgehog signal

transduction pathway activating substance in step (1). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

5 [0160]

In one preferable embodiment, in step (1), pluripotent stem cells (e.g., human pluripotent stem cell) are cultured (preferably, adhesion culture) in a medium containing a TGF β family signal transduction pathway inhibiting substance (e.g.,
10 a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a Nodal/Activin signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination thereof (e.g., SB431542 and
15 LDN193189) etc.); a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA); or a combination of a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway
20 activating substance (e.g., Shh protein, SAG, PMA) and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a Sonic hedgehog signal transduction
25 pathway activating substance (e.g., SAG, PMA, Shh protein) (in one embodiment, not containing a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty)), or a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2), a Sonic hedgehog
30 signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably
35 uniformed aggregates.

[0161]

In another embodiment, in step (1), pluripotent stem cells (e.g., human pluripotent stem cell) are cultured (preferably, adhesion culture) in a medium containing a TGF β family signal transduction pathway inhibiting substance (e.g., a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a Nodal/Activin signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination thereof (e.g., SB431542 and LDN193189) etc.); a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA); or a combination of a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2); a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) or a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2): or a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

[0162]

In another embodiment, in step (1), pluripotent stem cells (e.g., human pluripotent stem cell) are cultured (preferably, adhesion culture) in a medium containing a combination of a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) for 18 hr - 30 hr (e.g., about 24 hr), after which further cultured (preferably, adhesion culture) in a medium containing a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) for 18 hr - 30 hr (e.g., about 24 hr) and, in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2); a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) or a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2): or a medium not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) but containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

[0163]

In another embodiment, in step (1), pluripotent stem cells (e.g., human pluripotent stem cell) are cultured (preferably, adhesion culture) in a medium containing a

combination of a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) for 18 hr - 5 30 hr (e.g., about 24 hr), after which further cultured (preferably, adhesion culture) in a medium containing a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) for 18 hr - 30 hr (e.g., about 24 hr) and, in step (2), the cells obtained in step (1) are subjected 10 to suspension culture in a medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction 15 pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01). The period for the culturing in step (2) is 24 hr - 6 days, preferably 24 hr - 48 hr. The aggregates formed are preferably uniformed aggregates.

[0164]

20 In any embodiment, the medium in step (2) preferably contains a ROCK inhibiting substance (e.g., Y-27632).

[0165]

By performing step (2) in this manner, an aggregate of the cells obtained in step (1), or the cells derived therefrom 25 are formed. The present invention also provides a method for producing such aggregate. The aggregate obtained in step (2) has higher quality than the one formed when a treatment with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating 30 substance is not performed in step (1). To be specific, for example, a population of aggregates having a high ratio of spherical cell aggregates having a smooth surface, a dense inside, and uncollapsed shape can be obtained. In one embodiment, when aggregates (e.g., not less than 100 35 aggregates) are randomly selected on day 6 from the start of

the second step, the sum of the ratios of uncollapsed aggregates and/or non-cystic aggregates is, for example, not less than 70%, preferably not less than 80%.

[0166]

5 The aggregate obtained in step (2) has a potency to differentiate into a retinal cell or retinal tissue (e.g., retinal tissue, retinal progenitor cell, or retinal layer-specific neural cell).

[0167]

10 In one embodiment, by using the stem cells obtained in step (1) and having a potency to differentiate into at least a retinal cell or a retinal tissue (e.g., retinal tissue, retinal progenitor cell, or retinal layer-specific neural cell) (e.g., Oct3/4 positive stem cells having a potency to differentiate
15 into at least a retinal cell or a retinal tissue (e.g., retinal tissue, retinal progenitor cell, or retinal layer-specific neural cell)) in step (2), an aggregate containing stem cells (preferably Oct3/4 positive stem cells) having a potency to differentiate into at least a retinal cell or a retinal tissue
20 (e.g., retinal tissue, retinal progenitor cell, or retinal layer-specific neural cell) can be obtained. Retinal cells and retinal tissues can be induced with high efficiency by culturing the aggregate obtained in step (2) under appropriate differentiation conditions.

25 [0168]

 In one embodiment, the aggregate obtained in step (2) contains cells corresponding to the cells in an intermediate stage between cells maintaining pluripotent-like properties (specifically, expressing Oct3/4) obtained on completion of
30 step (1), and retinal cells or retinal tissues. These cells express any of (i) pluripotent property marker Oct3/4, and (ii) the aforementioned retinal cell marker (Rx, PAX6, Chx10, Crx, Blimp1) and/or the aforementioned retina layer-specific neural cell marker. That is, in one embodiment, the aggregate
35 obtained in step (2) contains a mixture of cells expressing any

of (i) pluripotent property marker Oct3/4, and (ii) the
aforementioned retinal cell marker (Rx, PAX6, Chx10, Crx,
Blimp1) and/or the aforementioned retinal layer-specific neural
cell marker. That is, the aggregate obtained in step (2)
5 contains stem cells having a potency to differentiate into at
least a retinal cell or retinal tissue, and/or progenitor cells
of retinal cells or retinal tissues. The progenitor cells are
characterized in that they show an ability (competence) to
express the aforementioned retinal cell or retina layer-
10 specific neural cell markers when they are cultured under known
appropriate culture conditions. Therefore, in one embodiment,
the aggregate obtained in step (2) contains Oct3/4 positive
stem cells having a potency to differentiate into at least a
retinal cell or retinal tissue, and/or progenitor cells of a
15 retinal cell or retinal tissue. A part of the cells contained
in the aggregate obtained in step (2) may express the
aforementioned retina layer-specific cell marker. In one
embodiment, the aggregate obtained in step (2) may contain
Oct3/4 positive cells at a proportion of not less than 50%, for
20 example, not less than 70%, of the total cells.

[0169]

2-3. step (3)

Step (3) where an aggregate containing retinal cells or a
retinal tissue are induced from the aggregate obtained in step
25 (2) is explained.

[0170]

The medium (basal medium) to be used in step (3) is not
particularly limited, and the basal medium described in the
above-mentioned definitions can be appropriately selected. The
30 medium to be used in step (3) is, for example, a serum-free
medium or a serum-containing medium (preferably serum-free
medium) supplemented with a BMP signal transduction pathway
activating substance.

[0171]

35 A serum-free medium or serum-containing medium to be used

for such medium is not particularly limited as long as it is as mentioned above. To avoid complicated preparation, for example, a serum-free medium supplemented with an appropriate amount of a commercially available serum alternative such as KSR and so
5 on (e.g., medium of 1:1 mixture of IMDM and F-12 supplemented with 10% KSR, 450 μ M 1-monothioglycerol and 1x Chemically Defined Lipid Concentrate) is preferably used. The amount of KSR to be added to a serum-free medium in the case of human pluripotent stem cell (e.g., iPS cell) is generally about 1% to
10 about 20%, preferably about 2% to about 20%.

[0172]

As the medium (preferably serum-free medium) to be used in step (3), the medium (preferably serum-free medium) used in step (2) may be directly used, or may be replaced with a fresh
15 medium (preferably serum-free medium). When the serum-free medium free of a BMP signal transduction pathway substance used in step (2) is directly used for step (3), a BMP signal transduction pathway activating substance may be added to the medium.

20 [0173]

Examples of the BMP signal transduction pathway activating substance to be used in step (3) include BMP proteins such as BMP2, BMP4, BMP7 etc., GDF proteins such as GDF7 etc., anti-BMP receptor antibody, BMP partial peptide and
25 so on. BMP2 protein, BMP4 protein and BMP7 protein are available from, for example, R&D Systems, and GDF7 protein is available from, for example, Wako Pure Chemical Industries, Ltd. The BMP signal transduction pathway activating substance is preferably BMP4.

30 [0174]

The concentration of the BMP signal transduction pathway activating substance may be a concentration at which differentiation of the cells forming an aggregate of pluripotent stem cells into retinal cells can be induced. For
35 example, in the case of human BMP4, it is added to the medium

to a concentration of about 0.01 nM to about 1 μ M, preferably about 0.1 nM to about 100 nM, more preferably about 1 nM to about 10 nM, further preferably about 1.5 nM (55 ng/mL). When a BMP signal transduction pathway activating substance other
5 than BMP4 is used, it is desirably used at a concentration at which a BMP signal transduction promoting activity equivalent to that of BMP4 at the above-mentioned concentration is exerted.
[0175]

The concentration of the BMP signal transduction pathway
10 activating substance in the medium may be varied during the period of step (3). For example, the BMP signal transduction pathway activating substance is provided to fall within the above-mentioned range at the time of the start of step (3), and the concentration may be gradually or stepwisely decreased at a
15 ratio of 40 - 60% per 2 - 4 days. In one embodiment, the cells are cultured with a given concentration (e.g., about 1.5 nM) of a BMP signal transduction pathway activating substance for a given period (e.g., 4 - 10 days) from the start of step (3), and the concentration may be gradually or stepwisely decreased
20 at a ratio of 40 - 60% per 2 - 4 days.
[0176]

A BMP signal transduction pathway activating substance only needs to be added after about 24 hr or later from the start of the suspension culture in step (2), and may also be
25 added to the medium within several days (e.g., within 15 days) from the start of the suspension culture. Preferably, a BMP signal transduction pathway activating substance is added to the medium at any time point between day 1 and day 15, more preferably between day 1 and day 9, day 1 and day 8, day 1 and
30 day 7, day 1 and day 6, day 1 and day 5 or day 1 and day 4, more preferably day 1 and day 3, from the start of the suspension culture in step (2). In the present specification, "N days from the start of suspension culture in step (2)" means a period of from N days after the start of the suspension
35 culture in step (2) (N x 24 hr later) to immediately before

progress of N+1 days ((N+1) x 24 hr). For example, 2 days from the start of suspension culture in step (2) means 48 hr from the start of suspension culture in step (2) as the standard to immediately before progress of 72 hr.

5 [0177]

In a specific embodiment, the medium may be partly or entirely exchanged with a medium containing BMP4 on, for example, day 1 - 9, preferably day 1 - 3, after the start of suspension culture in step (2) to adjust the final
10 concentration of BMP4 to about 1 - 10 nM, and the cells may be cultured in the presence of BMP4 for, for example, about 1 - 12 days, preferably 2 - 9 days, more preferably 2 - 5 days. To maintain the concentration of BMP4 at the same level, the medium may be partly or entirely exchanged one or two times
15 with a medium containing BMP4. Alternatively, as mentioned above, the concentration of BMP4 may also be stepwisely reduced.
[0178]

After the addition of a BMP signal transduction pathway activating substance to the medium and the start of the
20 differentiation induction of cells forming an aggregate into retinal cells, addition of the BMP signal transduction pathway activating substance to the medium is not necessary, and the medium may be exchanged with a serum-free medium or serum-containing medium each free of a BMP signal transduction
25 pathway activating substance. In one embodiment, after the start of the differentiation induction into retinal cells, the concentration of the BMP signal transduction pathway activating substance in the medium is gradually or stepwisely decreased at a ratio of 40 - 60% per 2 - 4 days by exchanging the medium
30 with a serum-free medium or a serum-containing medium, each free of a BMP signal transduction pathway activating substance. The cells in which induction of differentiation into retinal cells has been started can be confirmed by, for example, detecting the expression of retinal progenitor cell marker gene
35 (e.g., Rx gene (alias Rax), Pax6 gene, Chx10 gene) in the cells.

The aggregate formed in step (2) by using pluripotent stem cells in which a fluorescent reporter protein gene such as GFP and so on is knocked-in into the Rx gene locus is cultured in suspension in the presence of a BMP signal transduction pathway activating substance at a concentration necessary for differentiation induction into retinal cell, and fluorescence emitted from the expressed fluorescence reporter protein is detected, whereby the time point when differentiation induction into retinal cell was started can be also confirmed. As one embodiment of step (3), a step of culturing the aggregate formed in step (2) in suspension in a serum-free medium or serum-containing medium containing a BMP signal transduction pathway activating substance at a concentration necessary for differentiation induction into retinal cell, until a cell expressing retinal progenitor cell marker gene (e.g., Rx gene, Pax6 gene, Chx10 gene) begins appearing, thereby obtaining an aggregate comprising retinal progenitor cells can be mentioned.

[0179]

In step (3), when a medium exchange operation is performed, for example, an operation to add a fresh medium without discarding the existing medium (medium addition operation), an operation to discard about a half amount of the existing medium (about 40 - 80% of the volume of the existing medium) and add about a half amount of a fresh medium (40 - 80% of the volume of the existing medium) (half-medium exchange operation), and an operation to discard about the whole amount of the existing medium (not less than 90% of the amount of the existing medium) and add about the whole amount of a fresh medium (not less than 90% of the amount of the existing medium) (full-medium exchange operation) can be mentioned.

[0180]

When a particular component (e.g., BMP4) is added at a certain time point, for example, an operation to calculate the final concentration, to discard about a half amount of the existing medium, and to add about a half amount of a fresh

medium containing a particular component at a concentration higher than the final concentration (specifically 1.5 - 3.0 times the final concentration, for example, about 2 times the final concentration) (half-medium exchange operation, half-medium exchange) may be performed.

[0181]

When the concentration of a particular component contained in the existing medium is to be maintained at a certain time point, for example, an operation to discard about a half amount of the existing medium and to add about a half amount of a fresh medium containing the particular component at a concentration same as that in the existing medium may be performed.

[0182]

When the concentration of a component contained in the existing medium is to be decreased by dilution at a certain time pointed, for example, the medium exchange operation may be performed plural times per day, preferably plural times (e.g., 2 - 3 times) within 1 hr. Also, when the concentration of a component contained in the existing medium is to be decreased by dilution at a certain time point, the cell or aggregate may be transferred to another culture container.

[0183]

While the tool used for the medium exchange operation is not particularly limited, for example, pipetter, micropipette, multichannel micropipette, continuous dispenser, and the like can be mentioned. For example, when a 96 well plate is used as a culture vessel, a multichannel micropipette may be used.

[0184]

In one embodiment, when the concentration of the Shh signal transduction pathway activating substance added to the medium in step (2) is comparatively low (e.g., not more than 300 nM, preferably not more than 30 nM, further preferably not more than 3 nM, for SAG, and a concentration conferring Shh signal transduction promoting activity equivalent to or lower

than that of SAG at the above-mentioned concentration, for other Shh signal transduction pathway activating substances), medium exchange is not necessary, and a BMP signal transduction action substance (e.g., BMP4) may be added to the medium used
5 in step (2). On the other hand, when the concentration of the Shh signal transduction pathway activating substance is comparatively high (e.g., exceeding 700 nM or not less than 1000 nM for SAG, and a concentration conferring a Shh signal transduction promoting activity equivalent to that of SAG at
10 the above-mentioned concentration, for other Shh signal transduction pathway activating substances), it is desirable to exchange the medium to a fresh medium containing a BMP signal transduction action substance (e.g., BMP4) and free of a Shh signal transduction pathway activating substance to suppress an
15 influence of the Shh signal transduction pathway activating substance remaining when a BMP signal transduction action substance is added.

[0185]

In a preferable embodiment, the concentration of a Shh
20 signal transduction pathway activating substance in the medium to be used in step (3) is a concentration not imparting an adverse influence on selective differentiation into a retinal progenitor cell or a retinal tissue. Concretely, it is, when calculated in terms of Shh signal transduction promoting
25 activity of SAG, not more than 700 nM, preferably not more than 300 nM, more preferably not more than 10 nM, further preferably not more than 0.1 nM, further preferably substantially free of a Shh signal transduction pathway activating substance. The medium "substantially free of a Shh signal transduction pathway
30 activating substance" means a medium in which the concentration of the Shh signal transduction pathway activating substance in the medium is suppressed to a level at which the biological activity thereof (i.e., Sonic hedgehog signal transduction pathway activation) is not expressed (e.g., not more than 0.03
35 nM for SAG). The medium "free of a Shh signal transduction

pathway activating substance" also includes a medium substantially not supplemented with a Shh signal transduction pathway activating substance, for example, a medium not supplemented with a Shh signal transduction pathway activating substance at a concentration at which the biological activity thereof (i.e., Sonic hedgehog signal transduction pathway activation) is expressed, that is, a concentration imparting an adverse influence on selective differentiation into a retinal progenitor cell or a retinal tissue.

10 [0186]

In step (3), the aggregates obtained in step (2) are cultured in the presence or absence of a Wnt signal transduction pathway inhibiting substance in a medium (preferably serum-free medium) containing a BMP signal transduction pathway activating substance (e.g., BMP4). That is, the medium used in step (3) may or may not contain, in addition to the BMP signal transduction pathway activating substance (e.g., BMP4), a Wnt signal transduction pathway inhibiting substance. In a preferable embodiment, the aggregates obtained in step (2) are cultured in the presence of a Wnt signal transduction pathway inhibiting substance in a medium (preferably serum-free medium) containing a BMP signal transduction pathway activating substance (e.g., BMP4). That is, in this embodiment, the medium (preferably serum-free medium) used in step (3) contains a BMP signal transduction pathway activating substance (e.g., BMP4), and a Wnt signal transduction pathway inhibiting substance.

[0187]

As the Wnt signal transduction pathway inhibiting substance, any of the aforementioned Wnt signal transduction pathway inhibiting substances can be used. Preferably, the same kind of Wnt signal transduction pathway inhibiting substance as in step (2) is used in step (3).

[0188]

35 The concentration of the Wnt signal transduction pathway

inhibiting substance only needs to be a concentration capable of inducing a retinal progenitor cell and retinal tissue. For example, IWR-1-endo is added to a medium such that the concentration is about 0.1 μ M to about 100 μ M, preferably about 5 0.3 μ M to about 30 μ M, more preferably about 1 μ M to about 10 μ M, further preferably about 3 μ M. When a Wnt signal transduction pathway inhibiting substance other than IWR-1-endo is used, it is desirably used at a concentration exhibiting a Wnt signal transduction pathway inhibiting activity equivalent 10 to that of IWR-1-endo at the above-mentioned concentration. The concentration of the Wnt signal transduction pathway inhibiting substance in the medium of step (3) is preferably 50 - 150, more preferably 80 - 120, further preferably 90 - 110, when the concentration of the Wnt signal transduction pathway 15 inhibiting substance in the medium of step (2) is 100, and more preferably equivalent to the concentration of the Wnt signal transduction pathway inhibiting substance in the medium of step (2).

[0189]

20 The timing of addition of a Wnt signal transduction pathway inhibiting substance to the medium is not particularly limited as long as an aggregate containing a retinal cell or retinal tissue can be formed, but earlier addition is more preferable. Preferably, a Wnt signal transduction pathway 25 inhibiting substance is added to the medium at the time of the start of step (3). More preferably, a Wnt signal transduction pathway inhibiting substance is added in step (2) and continuously contained in the medium also in step (3) (i.e., from the time of the start of step (3)). Further preferably, a 30 Wnt signal transduction pathway inhibiting substance is added at the time of the start of suspension culture in step (2) and continuously contained in the medium also in step (3). For example, a BMP signal transduction action substance (e.g., BMP4) is added to the culture obtained in step (2) (suspension 35 of aggregate in the medium containing a Wnt signal transduction

pathway inhibiting substance).

[0190]

The period of action of the Wnt signal transduction pathway inhibiting substance is not particularly limited as long as the above-mentioned effects can be afforded.

Preferably, when a Wnt signal transduction pathway inhibiting substance is added at the time of the start of suspension culture in step (2), the period is 2 days to 30 days, more preferably 6 days to 20 days, 8 days to 18 days, 10 days to 18 days, or 10 days to 17 days (e.g., 10 days), with the time of the start of suspension culture in step (2) as the starting point. In another embodiment, when the Wnt signal transduction pathway inhibiting substance is added at the time of the start of suspension culture in step (2), the period of action of the Wnt signal transduction pathway inhibiting substance is preferably 3 days to 15 days (e.g., 5 days, 6 days, 7 days), more preferably 6 days to 10 days (e.g., 6 days), with the time of the start of suspension culture in step (2) as the starting point.

[0191]

In one embodiment, medium used in step (3) may further contain a TGF β signal transduction pathway inhibiting substance. Particularly, when a medium containing a TGF β signal transduction pathway inhibiting substance is used in the suspension culture of step (2), the medium containing a TGF β signal transduction pathway inhibiting substance is preferably used continuously in step (3) as well.

[0192]

As the TGF β signal transduction pathway inhibiting substance, any of the aforementioned TGF β signal transduction pathway inhibiting substances can be used. Preferably, those of the same kind as the TGF β signal transduction pathway inhibiting substance used in step (2) are used in step (3).

[0193]

The concentration of the TGF β signal transduction pathway

inhibiting substance only needs to be a concentration capable of inducing a retinal progenitor cell and a retinal tissue. For example, SB431542 is generally used at a concentration of 0.1 - 200 μ M, preferably 2 - 50 μ M, more preferably 3 - 10 μ M. 5 A-83-01 is generally used at a concentration of 0.05 - 50 μ M, preferably 0.5 - 5 μ M. When a TGF β signal transduction pathway inhibiting substance other than SB431542 or A-83-01 is used, it is desirably used at a concentration exhibiting a TGF β signal transduction pathway inhibiting substance equivalent to that of 10 SB431542 at the above-mentioned concentration. The concentration of the TGF β signal transduction pathway inhibiting substance in the medium of step (3) is preferably 50 - 150, more preferably 80 - 120, further preferably 90 - 110, when the concentration of the TGF β signal transduction pathway 15 inhibiting substance in the medium of step (2) is 100. It is more preferably equivalent to the concentration of the TGF β signal transduction pathway inhibiting substance in the medium of step (2).

[0194]

20 The addition time and action period of the TGF β signal transduction pathway inhibiting substance in the medium is not particularly limited as long as formation of an aggregate containing a retinal cell or a retinal tissue can be achieved. When a TGF β signal transduction pathway inhibiting substance is 25 added at the time of the start of suspension culture in step (2), it is preferably 2 days to 30 days, more preferably 6 days to 20 days, 8 days to 18 days, 10 days to 18 days, or 10 days to 17 days (e.g., 10 days), with the time of the start of suspension culture in step (2) as the starting point. In a 30 preferable embodiment, the addition time and exposure period of the TGF β signal transduction pathway inhibiting substance in the medium is the same as the aforementioned addition time and action period of the Wnt signal transduction pathway inhibiting substance. That is, in the embodiment, a TGF β signal 35 transduction pathway inhibiting substance is added to the

medium simultaneously with a Wnt signal transduction pathway inhibiting substance, and contained in the medium for the same period as the Wnt signal transduction pathway inhibiting substance.

5 [0195]

For example, in step (1), pluripotent stem cells are treated with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance; in step (2), the
10 cells obtained in step (1) are subjected to suspension culture in a medium (preferably serum-free medium) containing a Wnt signal transduction pathway inhibiting substance and a TGF β family signal transduction pathway inhibiting substance or a medium (preferably serum-free medium) containing a Wnt signal
15 transduction pathway inhibiting substance, a Sonic hedgehog signal transduction pathway activating substance and a TGF β family signal transduction pathway inhibiting substance to form a cell aggregate; and in step (3), the aggregates obtained in step (2) are cultured in the presence of a Wnt signal
20 transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance in a medium (preferably serum-free medium) containing a BMP signal transduction pathway activating substance (e.g., BMP4). That is, in this embodiment, the medium (preferably serum-free
25 medium) used in step (3) contains a BMP signal transduction pathway activating substance (e.g., BMP4), a TGF β signal transduction pathway inhibiting substance and a Wnt signal transduction pathway inhibiting substance.

[0196]

30 In one embodiment, after suspension culture of step (3), the medium of step (3) is exchanged with a medium substantially free of an exogenous BMP signal transduction action substance and containing a Sonic hedgehog signal transduction pathway activating substance and a serum and suspension culture can be
35 continued. In step (1), pluripotent stem cells are treated

with a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and in step (2), the cells obtained in step (1) are subjected to suspension culture in a medium
5 (preferably serum-free medium and optionally further containing a TGF β signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance) containing or not containing a Wnt signal transduction pathway inhibiting substance to form an aggregate,
10 the aggregates obtained in step (2) are cultured in a medium (preferably serum-free medium and optionally further containing a Wnt signal transduction pathway inhibiting substance and/or a TGF β signal transduction pathway inhibiting substance) containing a BMP signal transduction pathway activating
15 substance (e.g., BMP4) for a given period, the medium is exchanged to a medium substantially free of an exogenous BMP signal transduction action substance (e.g., BMP4) containing a Sonic hedgehog signal transduction pathway activating substance and a serum and culturing is continued. That is, in this
20 embodiment, after culturing in step (3), the medium is exchanged from a medium (preferably serum-free medium and optionally further containing a Wnt signal transduction pathway inhibiting substance and/or a TGF β signal transduction pathway inhibiting substance) containing a BMP signal transduction
25 pathway activating substance (e.g., BMP4) to a medium substantially free of an exogenous BMP signal transduction action substance and containing a Sonic hedgehog signal transduction pathway activating substance and a serum.

[0197]

30 As the serum, for example, serum of mammals such as bovine serum, calf serum, fetal calf serum, horse serum, foal serum, foal fetal serum, rabbit serum, leveret rabbit serum, rabbit fetal serum, human serum and the like can be used. The serum is added at a concentration of about 1 - 30%, preferably
35 about 3 - 20%, more preferably about 10%. In this embodiment,

an appropriate amount of a serum alternative such as commercially available KSR and the like may be used instead of the serum.

[0198]

5 As the Sonic hedgehog signal transduction pathway activating substance, any of the aforementioned Sonic hedgehog signal transduction pathway activating substance can be used, preferably SAG. For example, in the case of SAG, it is generally added at a concentration of about 0.1 nM - 10 μ M, 10 preferably about 10 nM - 1 μ M, more preferably about 100 nM.

[0199]

When the concentration of a BMP signal transduction action substance (e.g., BMP4) to be carried over from step (3) is suppressed to a level at which the biological activity 15 (namely, BMP signal transduction action) thereof is not expressed (e.g., concentration of less than 0.01 nM for BMP4), it can be regarded as a medium substantially free of a BMP signal transduction action substance. For example, by a medium exchange operation with a medium free of a BMP signal 20 transduction action substance, carrying over of the BMP signal transduction action substance (e.g., BMP4) from step (3) is suppressed to not more than 3%, preferably not more than 1%, not more than 0.3%, not more than 0.1%, not more than 0.03% or not more than 0.01% of the concentration of the BMP signal 25 transduction action substance (e.g., BMP4) in the medium used in step (3), and the concentration of a BMP signal transduction action substance in the medium is suppressed to a level at which the biological activity (i.e., BMP signal transduction action) thereof is not expressed (e.g., concentration of BMP4 30 is less than 0.01 nM), whereby a medium substantially free of a BMP signal transduction action substance can be obtained.

[0200]

In one embodiment, exchange to a medium substantially free of an exogenous BMP signal transduction action substance 35 and containing a Sonic hedgehog signal transduction pathway

activating substance and a serum is performed after 7 days,
more preferably after 9 days (for example, 10 days), from the
start of suspension culture (step (2)). Here, exchange to a
medium containing a Sonic hedgehog signal transduction pathway
5 activating substance and a serum may also be performed when the
formation of the neuroepithelium is started. Formation of the
neuroepithelium can be confirmed by observing the sequence .
morphology of the cells by bright field image observation using
a microscope. In one embodiment, the cells are cultured for 3
10 days to 20 days, preferably 5 days to 15 days, 7 days to 10
days, after exchange to a medium containing a Sonic hedgehog
signal transduction pathway activating substance and a serum.
[0201]

In the method of the present invention, suspension
15 culture is preferably performed through step (2) and step (3)
in the absence of a basement membrane preparation.
[0202]

The culture conditions such as culture temperature, CO₂
concentration and so on in step (3) can be appropriately
20 determined. The culture temperature is, for example, about
30°C to about 40°C, preferably about 37°C. The CO₂
concentration is, for example, about 1% to about 10%,
preferably about 5%.
[0203]

25 By such culture, differentiation of the cells forming the
aggregate obtained in step (2) into retinal progenitor cells is
induced, whereby an aggregate containing the retinal progenitor
cells can be obtained. The present invention also provides a
method for producing such aggregate containing retinal
30 progenitor cell. That an aggregate comprising retinal
progenitor cells was obtained can be confirmed by, for example,
detecting the presence of cells expressing Rax, PAX6 or Chx10,
which is a retinal progenitor cell marker, in the aggregate.
One embodiment of step (3) is a step of culturing the aggregate
35 formed in step (2) in suspension in a serum-free medium or

serum-containing medium containing a BMP signal transduction pathway activating substance at a concentration necessary for differentiation induction into retinal cell, until a cell expressing Rx gene begins appearing, whereby obtaining an
5 aggregate comprising retinal progenitor cells. In one embodiment, the culturing of step (3) is performed until not less than 20% (preferably, not less than 30%, not less than 40%, not less than 50%, not less than 60%, not less than 70%, not less than 80%, not less than 90%) of the cells contained in the
10 aggregate express Rx. In this embodiment, the Rx expressing cell preferably coexpresses Chx10.

[0204]

In one embodiment, the aggregate obtained in step (3) contains a retinal tissue and is substantially free of non-
15 neural head ectoderm. In an aggregate containing a retinal tissue and substantially free of non-neural head ectoderm, for example, an Rx-positive tissue is observed and an Rx-negative tissue is not observed on the outside thereof in the immunostaining images of the aforementioned aggregate
20 cryosection.

[0205]

In a preferable embodiment of producing a retinal cell and/or a retinal tissue, in step (1), human pluripotent stem cells (e.g., human iPS cell) are subjected to adhesion culture
25 in the absence of feeder cells in a serum-free medium containing a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01) and bFGF for preferably 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60 hr, 18 - 60 hr, 18 - 48 hr or 18 - 28 hr (e.g., 24 hr), and in step (2), the cells
30 obtained in step (1) are subjected to suspension culture in the presence of a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) in a serum-free medium not containing or containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein), and in step (3), the aggregates are
35

subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a BMP signal transduction pathway activating substance (e.g.,
5 BMP4).

[0206]

In a preferable embodiment of producing a retinal cell and/or a retinal tissue, in step (1), human pluripotent stem cells (e.g., human iPS cell) are subjected to adhesion culture
10 in the absence of feeder cells in a serum-free medium containing a BMP signal transduction pathway inhibiting substance (e.g., LDN193189) and bFGF for preferably 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60 hr, 18 - 60 hr, 18 - 48 hr or 18 - 28 hr (e.g., 24 hr), and in step (2), the cells obtained in
15 step (1) are subjected to suspension culture in the presence of a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) in a serum-free medium not containing or containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA), and
20 in step (3), the aggregates are subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a BMP signal transduction pathway activating substance (e.g., BMP4).

25 [0207]

In a preferable embodiment in the production of retinal progenitor cells and/or retinal tissues, human pluripotent stem cells (e.g., human iPS cells) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium
30 containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA) and bFGF for preferably 2 - 96 hr, 6 - 72 hr, 6 - 60 hr, 12 - 60 hr, 18 - 60 hr, 18 - 48 hr, or 18 - 28 hr (e.g., for 24 hr) in step (1) and in step (2), the cells obtained in step (1) are subjected to suspension
35 culture in the presence of a Wnt signal transduction pathway

inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) in a serum-free medium not containing or containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA), and in step (3), the aggregates are subjected
5 to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) and a BMP signal transduction pathway activating substance (e.g., BMP4).

[0208]

10 In a preferable embodiment in the production of retinal cells and/or retinal tissues, human pluripotent stem cells (e.g., human iPS cells) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium containing a TGF β family signal transduction pathway inhibiting substance
15 (e.g., a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a Nodal/Activin signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination of
20 these (e.g., SB431542 and LDN193189) etc.); a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA); or a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway
25 activating substance (e.g., Shh protein, SAG, PMA) in combination; and bFGF in step (1), in step (2), the cells obtained in step (1) are subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e),
30 IWP-2) (containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01)) to form a cell aggregate, and in step (3), the aggregates are subjected to
35 suspension culture in a serum-free medium containing a Wnt

signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo(IWR1e), IWP-2) and a BMP signal transduction pathway activating substance (e.g., BMP4) (containing or containing a TGF β signal transduction pathway inhibiting
5 substance (e.g., Lefty, SB431542, A-83-01)) to give an aggregate containing a retinal cell (e.g., retinal progenitor cell, retinal layer-specific neural cell) or retinal tissue.

The medium in step (2) preferably contains a ROCK inhibiting substance (e.g., Y-27632).

10 [0209]

In a preferable embodiment in the production of retinal cells and/or retinal tissues, human pluripotent stem cells (e.g., human iPS cells) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium containing a
15 TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) in combination; and bFGF for 18 hr - 30 hr (e.g., about 24 hr) and thereafter adhesion culture in a
20 serum-free medium containing a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) and bFGF for 18 hr - 30 hr (e.g., about 24 hr), in step (2), the cells obtained in step (1) are subjected to suspension culture in a serum-free medium containing a Wnt
25 signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) (containing or not containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty,
30 SB431542, A-83-01)) to form a cell aggregate, and in step (3), the aggregates are subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo(IWR1e), IWP-2) and a BMP signal transduction pathway activating
35 substance (e.g., BMP4) (containing or containing a TGF β signal

transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01)) to give an aggregate containing a retinal cell (e.g., retinal progenitor cell, retinal layer-specific neural cell) or retinal tissue.

- 5 The medium in step (2) preferably contains a ROCK inhibiting substance (e.g., Y-27632).
[0210]

In a preferable embodiment in the production of retinal cells and/or retinal tissues, human pluripotent stem cells
10 (e.g., human iPS cells) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium containing a TGF β family signal transduction pathway inhibiting substance (e.g., a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a Nodal/Activin signal
15 transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination of these (e.g., SB431542 and LDN193189) etc.); a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh
20 protein, SAG, PMA); or a TGF β family signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) in
25 obtained in step (1) are subjected to suspension culture in a serum-free medium or a serum-free medium containing a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) (containing or not containing a Sonic hedgehog signal transduction pathway activating
30 substance (e.g., SAG, PMA, Shh protein) and/or a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01)) to form a cell aggregate, and in step (3), the aggregates are subjected to suspension culture in a serum-free medium containing a Wnt signal transduction pathway
35 inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo(IWR1e),

IWP-2) and a BMP signal transduction pathway activating substance (e.g., BMP4) for 10 days, and the medium was exchanged to a medium not containing an exogenous BMP signal transduction action substance (e.g., BMP4) and containing a
5 Sonic hedgehog signal transduction pathway activating substance (e.g., SAG) and a serum and the suspension culture was continued to give an aggregate containing a retinal cell (e.g., retinal progenitor cell, retinal layer-specific neural cell) or retinal tissue.

10 The medium in step (2) preferably contains a ROCK inhibiting substance (e.g., Y-27632).
[0211]

In a preferable embodiment in the production of retinal cells and/or retinal tissues, human pluripotent stem cells
15 (e.g., human iPS cells) are subjected to adhesion culture in the absence of feeder cells in a serum-free medium containing a TGF β family signal transduction pathway inhibiting substance (e.g., a TGF β signal transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a Nodal/Activin signal
20 transduction pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01), a BMP signal transduction pathway inhibiting substance (e.g., LDN193189), or a combination of these (e.g., SB431542 and LDN193189) etc.); a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA); or a TGF β family signal transduction
25 pathway inhibiting substance (e.g., Lefty, SB431542, A-83-01, LDN193189) and a Sonic hedgehog signal transduction pathway activating substance (e.g., Shh protein, SAG, PMA) in combination; and bFGF in step (1), in step (2), the cells
30 obtained in step (1) are subjected to suspension culture in the presence of a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-endo (IWR1e), IWP-2) in a serum-free medium containing a Sonic hedgehog signal transduction pathway activating substance (e.g., SAG, PMA, Shh
35 protein) and/or a TGF β signal transduction pathway inhibiting

substance (e.g., SB431542, A-83-01, Lefty) to form a cell aggregate, and in step (3), the aggregates are subjected to suspension culture in the presence of a Wnt signal transduction pathway inhibiting substance (e.g., CKI-7, D4476, IWR-1-
5 endo(IWR1e), IWP-2) and a BMP signal transduction pathway activating substance (e.g., BMP4) in a serum-free medium containing a TGF β signal transduction pathway inhibiting substance (e.g., SB431542, A-83-01, Lefty) to give an aggregate containing a retinal cell (e.g., retinal progenitor cell,
10 retinal layer-specific neural cell) or retinal tissue.

The medium in step (2) preferably contains a ROCK inhibiting substance (e.g., Y-27632).

[0212]

The obtained aggregate containing retinal progenitor
15 cells may be used as it is as a reagent for evaluating toxicity or efficacy. An aggregate containing retinal progenitor cells is subjected to a dispersion treatment (e.g., trypsin/EDTA treatment or papain treatment), and the obtained cells are subjected to a selection using FACS or MACS, whereby highly
20 pure retinal progenitor cells can also be obtained.

[0213]

Furthermore, the aggregate containing the retinal progenitor cell obtained by the aforementioned step (1) - step (3) may be continuously cultured in a serum-free medium or
25 serum-containing medium to make the retinal progenitor cell further differentiate, whereby a neuroepithelial structure-like retinal tissue may be produced.

[0214]

Such medium (basal medium) is not particularly limited,
30 and the basal medium described in the above-mentioned definition can be appropriately selected. A serum-free medium or serum-containing medium to be used for such medium is not particularly limited as long as it is as mentioned above. For example, a serum-containing medium which is a DMEM-F12 medium
35 supplemented with 10% fetal bovine serum, N2 supplement, 100 μ M

taurine, and 500 nM retinoic acid, or a serum-free medium supplemented with an appropriate amount of a commercially available serum alternative such as KSR and so on (e.g., medium of 1:1 mixture of IMDM and F-12 supplemented with 10% KSR, 450
5 μ M 1-monothioglycerol and 1x Chemically Defined Lipid Concentrate) and the like can be mentioned.

[0215]

While the period of the culture for inducing a retinal tissue from retinal progenitor cells varies depending on the
10 intended retinal layer-specific neural cell, it is, for example, about 7 days to about 200 days.

[0216]

The retinal tissue exists covering the surface of the aggregate. After completion of the suspension culture, the
15 aggregate may be fixed with a fixative such as para-formaldehyde solution and so on, and a cryosection is prepared, then formation of a retinal tissue having a layer structure may be confirmed by immunostaining and the like. Since respective layers of a retinal tissue are composed of different retinal
20 progenitor cell (photoreceptor cell, horizontal cell, bipolar cell, amacrine cell, retinal ganglion cell), formation of a layer structure can be confirmed using antibodies against the aforementioned markers expressed in these cells by the immunostaining. In one embodiment, the retinal tissue is an
25 Rx- or Chx10-positive neuroepithelial structure.

[0217]

The retinal tissue existing on the surface of the aggregate can be physically excised from the aggregate by using tweezers and the like. In this case, since a neural tissue
30 other than the retinal tissue may be formed on the surface of each aggregate, a part of the neural tissue excised from the aggregate may be cut out and subjected to confirmation by the below-mentioned immunostaining and the like, whereby the tissue can be confirmed to be a retinal tissue.

35 [0218]

In one embodiment, the aggregate obtained in step (3) contains a retinal tissue and is substantially free of non-neural head ectoderm. In an aggregate containing a retinal tissue and substantially free of non-neural head ectoderm, for example, an Rx-positive tissue is observed and an Rx-negative tissue is not observed on the outside thereof in the immunostaining images of the aforementioned aggregate cryosection.

[0219]

One embodiment of step (3) is a step of culturing the aggregate formed in step (2) in suspension in a serum-free medium or serum-containing medium (as mentioned above, optionally containing a Wnt signal transduction pathway inhibiting substance and/or a TGF β signal transduction pathway inhibiting substance) containing a BMP signal transduction pathway activating substance (e.g., BMP4) at a concentration necessary for differentiation induction into retinal cells, until a cell expressing Rx gene and/or Chx10 gene begins appearing to give an aggregate comprising retinal progenitor cells, and subsequently culturing the aggregate containing the retinal progenitor cells in suspension in a serum-free medium or serum-containing medium until a retinal tissue is formed, whereby obtaining an aggregate comprising a retinal tissue. When the aggregate containing the retinal progenitor cells is subsequently cultured in suspension in a serum-free medium or serum-containing medium until a retinal tissue is formed, the concentration of the BMP signal transduction pathway activating substance in the medium in order to induce retinal progenitor cells may be gradually or stepwisely decreased at a ratio of 40 - 60% per 2 - 4 days by exchanging the medium with a serum-free medium or a serum-containing medium, each free of a BMP signal transduction pathway activating substance (e.g., BMP4). In this case, the concentration of the Wnt signal transduction pathway inhibiting substance and/or the TGF β signal transduction pathway inhibiting substance may be decreased at

the same proportion as in the BMP signal transduction pathway activating substance, and preferably maintained at a constant level. In one embodiment, suspension culture of an aggregate containing retinal progenitor cells is performed until not less
5 than 20% (preferably, not less than 30%, not less than 40%, not less than 50%, not less than 60%, not less than 70%, not less than 80%, not less than 90%) of the cells contained in the aggregate expresses Chx10.

[0220]

10 In one embodiment of step (3), the aggregate obtained in step (2), or an aggregate obtained by culturing the aggregate obtained in step (2) in suspension by the above-mentioned method may be subjected to adhesion culture to form an adhered aggregate. The adhered aggregate is cultured in an adhered
15 state in a serum-free medium or serum-containing medium containing a BMP signal transduction pathway activating substance at a concentration necessary for differentiation induction into a retinal cell (the medium optionally contains a Wnt signal transduction pathway inhibiting substance and/or a
20 TGF β signal transduction pathway inhibiting substance), until a cell expressing Rx gene and/or Chx10 gene begins appearing to give an adhered aggregate containing retinal progenitor cells. The aggregate containing the retinal progenitor cells is continuously cultured in an adhered state in a serum-free
25 medium or serum-containing medium until a retinal tissue is formed, whereby an aggregate containing a retinal tissue is obtained. In one embodiment, adhesion culture of the aggregate containing retinal progenitor cells is performed until not less than 10% (preferably, not less than 20%, not less than 30%, not
30 less than 40%, not less than 50%, not less than 60%, not less than 70%, not less than 80%, not less than 90%) of the cells express Chx10.

[0221]

By the production method of the present invention, a
35 retinal tissue can be obtained from pluripotent stem cells with

high efficiency. Since the retinal tissue obtained by the production method of the present invention contains neurons (neuronal cell) specific to each of the retinal layers, it is also possible to obtain cells constituting a retinal tissue, 5 such as photoreceptor cell, horizontal cell, bipolar cell, amacrine cell, retinal ganglion cell or a progenitor/precursor cell thereof and the like. Which cell was obtained from the obtained retinal tissue can be confirmed by a method known per se, for example, expression of a cell marker.

10 [0222]

The obtained aggregate containing a retinal tissue may also be directly used as a reagent for evaluating toxicity or efficacy. An aggregate containing a retinal tissue is subjected to a dispersion treatment (e.g., trypsin/EDTA 15 treatment), and the obtained cells are subjected to a selection using FACS or MACS, whereby highly pure retinal tissue-constituting cells, for example, highly pure photoreceptor cells, can also be obtained.

[0223]

20 A retinal tissue containing a ciliary marginal zone-like structure can be produced from a cell aggregate containing the retinal tissue obtained by the production method of the present invention, namely, the aforementioned step (1) - step (3), by a known method (e.g., WO 15/087614), specifically, the following 25 step (A) and step (B).

[0224]

In one embodiment, a ciliary marginal zone-like structure can be produced by the following step (A) and step (B) from a cell aggregate containing the retinal tissue on days 6 - 30, 30 days 10 - 20 (day 10, day 11, day 12, day 13, day 14, day 15, day 16, day 17, day 18, day 19 or day 20) from the start of suspension culture of step (2), which aggregate is a cell aggregate containing the retinal tissue obtained by steps (1) - (3).

35 [0225]

The ciliary marginal zone-like structure in the present specification refers to a structure similar to a ciliary marginal zone. Examples of the "ciliary marginal zone (CMZ)" include a tissue present in the boundary region of a retinal tissue (specifically, neural retina) and retinal pigment epithelium in the retina in vivo, which is a region containing tissue stem cells of retina (retinal stem cells). Ciliary marginal zone is also called a ciliary margin or retinal margin, and the ciliary marginal zone, ciliary margin and retinal margin are equivalent tissues. It is known that the ciliary marginal zone plays an important role in the supply of retinal progenitor cells or differentiated cells to retinal tissues, maintenance of retinal tissue structure and so on. Examples of the marker gene of the ciliary marginal zone include Rdh10 gene (positive), Otx1 gene (positive), Zic1 (positive) and so on.

[0226]

Step (A) comprises culturing a cell aggregate comprising a retinal tissue obtained by the production method of the present invention, i.e., step (1) - step (3), in which Chx10 positive cells are present in a proportion of 20% or more and 100% or less of the retinal tissue, in a serum-free medium or serum-containing medium each containing a Wnt signal pathway activating substance and/or an FGF signal pathway inhibiting substance for only a period before the appearance of a RPE65 gene-expressing cell.

[0227]

As a preferable culture of step (A) here, suspension culture can be mentioned.

[0228]

As a serum-free medium to be used in step (A), a serum-free medium which is a basal medium supplemented with N2 or KSR can be mentioned. More specifically, a serum-free medium which is a DMEM/F-12 medium supplemented with N2 supplement (Life Technologies) can be mentioned. As the serum-containing medium, a serum-containing medium which is a basal medium supplemented

with fetal bovine serum can be mentioned.

[0229]

The culture conditions of step (A) such as culture temperature, CO₂ concentration can be appropriately set. The culture temperature is, for example, in the range of about 30°C to about 40°C, preferably, for example, around about 37°C. The CO₂ concentration is, for example, in the range of about 1% to about 10%, preferably, for example, around about 5%.

[0230]

10 In step (A), when the above-mentioned "cell aggregate comprising a retinal tissue" is cultured in the medium, the Wnt signal transduction pathway activating substance to be contained in a serum-free medium or serum-containing medium is not particularly limited as long as it can enhance signal transduction mediated by Wnt. Specific examples of the Wnt signal transduction pathway activating substance include a protein belonging to Wnt family (e.g., Wnt1, Wnt3a, Wnt7a), Wnt receptor, Wnt receptor agonist, GSK3 β inhibitor (e.g., 6-Bromoindirubin-3'-oxime (BIO), CHIR99021, Kenpaullone) and so
20 on.

[0231]

The concentration of the Wnt signal transduction pathway activating substance to be contained in a serum-free medium or serum-containing medium in step (A) in the case of a common Wnt signal transduction pathway activating substance such as CHIR99021 is, for example, in the range of about 0.1 μ M to about 100 μ M, preferably, for example, in the range of about 1 μ M to about 30 μ M, more preferably, for example, around 3 μ M.

[0232]

30 The FGF signal transduction pathway inhibiting substance to be contained in a serum-free medium or serum-containing medium in step (A) when the above-mentioned "cell aggregate comprising a retinal tissue" is cultured in the medium is not particularly limited as long as it can inhibit signal transduction mediated by FGF. Examples of the FGF signal
35

transduction pathway inhibiting substance include FGF receptor, FGF receptor inhibitor (e.g., SU-5402, AZD4547, BGJ398), MAP kinase cascade inhibiting substance (e.g., MEK inhibitor, MAPK inhibitor, ERK inhibitor), PI3 kinase inhibitor, Akt inhibitor
5 and so on.

[0233]

The concentration of the FGF signal transduction pathway inhibiting substance contained in a serum-free medium or serum-containing medium in step (A) only needs to be a concentration
10 at which differentiation of an aggregate into ciliary marginal zone-like structure can be induced. For example, in the case of SU-5402, it is added to the medium to a concentration of about 0.1 μM to about 100 μM , preferably about 1 μM to about 30 μM , more preferably about 5 μM .

15 [0234]

"Culturing for only a period before the appearance of a RPE65 gene-expressing cell" in step (A) means culturing only in the whole or a part of the period before the appearance of a RPE65 gene-expressing cell. That is, culturing only in the
20 whole or a part of the period (any period) suffices, during which the aforementioned "cell aggregate comprising a retinal tissue" present in the culture system is constituted by cells that do not substantially express RPE65 gene. By employing such culturing, a cell aggregate in which a RPE65 gene-
25 expressing cell does not appear can be obtained.

[0235]

To determine such particular period, the aforementioned "cell aggregate comprising a retinal tissue" is used as a sample, and the presence or absence of expression of RPE65 gene
30 contained in the sample or the level thereof may be measured by a general genetic engineering method or a biochemical method. Specifically, for example, the presence or absence of expression of RPE65 gene or the level thereof can be examined by subjecting a cryosection of the aforementioned "cell
35 aggregate comprising a retinal tissue" to an immunostaining

method using an antibody against RPE65 protein.

[0236]

As a "period before the appearance of a RPE65 gene-expressing cell" in step (A), for example, a period during
5 which the ratio of Chx10 positive cells present in the above-mentioned retinal tissue decreases than that at the time of start of the culture of the aforementioned cell aggregate in a serum-free medium or serum-containing medium each containing a Wnt signal transduction pathway activating substance and/or an
10 FGF signal transduction pathway inhibiting substance, and falls within the range of 30% to 0% can be mentioned. As the "cell aggregate in which a RPE65 gene-expressing cell does not appear", a cell aggregate in which Chx10 positive cells are present in the above-mentioned retinal tissue in a proportion
15 of within 30% to 0% of the tissue can be mentioned.

[0237]

While the number of days of the "period before the appearance of a RPE65 gene-expressing cell" in step (A) varies depending on the kind of the Wnt signal transduction pathway
20 activating substance and/or the FGF signal transduction pathway inhibiting substance, the kind of the serum-free medium or serum-containing medium, other culture conditions and so on, it is, for example, within 14 days. More specifically, when a serum-free medium (e.g., serum-free medium which is a basal
25 medium supplemented with N2) is used, the above-mentioned period is preferably, for example, within 10 days (e.g., within 10 days from about day 20, with the start of step (2) as the starting point), more preferably, for example, 3 days to 6 days. When a serum-containing medium (e.g., serum-containing medium
30 which is a basal medium supplemented with fetal bovine serum) is used, the aforementioned period is preferably, for example, within 12 days (e.g., within 12 days from about day 20, with the start of step (2) as the starting point), more preferably, for example, 6 days to 9 days.

35 [0238]

Then as step (B), the "cell aggregate in which a RPE65 gene-expressing cell does not appear" obtained by culturing as mentioned above is cultured in a serum-free medium or serum-containing medium each free of a Wnt signal transduction
5 pathway activating substance.

[0239]

As a preferable culture in step (B), suspension culture can be mentioned.

[0240]

10 The serum-free medium of step (B) is preferably free of an FGF signal transduction pathway inhibiting substance.

[0241]

As the serum-free medium in step (B), a medium which is a basal medium supplemented with N2 or KSR can be mentioned. As
15 the serum-containing medium, a medium which is a basal medium supplemented with fetal bovine serum can be mentioned. More specifically, a serum-containing medium which is a DMEM/F-12 medium supplemented with fetal bovine serum can be mentioned.

[0242]

20 The above serum-free medium or serum-containing medium in step (B) may contain a known growth factor, an additive and a chemical substance that promote the growth, and so on. Examples of the known growth factor include EGF, FGF, IGF, insulin and so on. Examples of the additive that promotes the
25 growth include N2 supplement (Life Technologies), B27 supplement (Life Technologies), KSR (Life Technologies) and so on. Examples of the chemical substance that promotes the growth include retinoids (e.g., retinoic acid) and taurine.

[0243]

30 A preferable period for the culturing in step (B) is, for example, a period for the culturing during which the ratio of Chx10 positive cells present in the above-mentioned retinal tissue increases than that at the time of start of the culturing of the aforementioned cell aggregate in a serum-free
35 medium or serum-containing medium each free a Wnt signal

transduction pathway activating substance, and reaches 30% or more. Specifically, for example, it is 3 days - 60 days, preferably about 35 days, from day about 3, with the start of step (A) as the starting point.

5 [0244]

The culture conditions such as culture temperature, CO₂ concentration and the like in step (B) can be appropriately set. The culture temperature is, for example, in the range of about 30°C to about 40°C, preferably, for example, around about 37°C. 10 The CO₂ concentration is, for example, in the range of about 1% to about 10%, preferably, for example, around about 5%.

[0245]

While the number of the above-mentioned culture days until "a cell aggregate comprising a ciliary marginal zone-like 15 structure" is obtained in step (B) varies depending on the kind of the serum-free medium or serum-containing medium, other culture conditions and so on, it is, for example, within 100 days. The above number of culture days is preferably, for example, 20 days to 70 days, more preferably, for example, 30 20 days to 60 days.

[0246]

In a "cell aggregate comprising a ciliary marginal zone-like structure" prepared by the aforementioned step (A), (B), a retinal pigment epithelium and a retinal tissue (specifically, 25 neural retina) are present adjacent to the ciliary marginal zone-like structure in the same cell aggregate. The structure can be confirmed by microscopic observation and so on. Specifically, for example, the presence of a ciliary marginal zone-like structure as an epithelial structure having a thick 30 retina side and a thin retinal pigment epithelium side, which is formed between a retinal tissue having high transparency and retinal pigment epithelium showing pigmentation, can be confirmed by microscopic observation. In addition, the presence of ciliary marginal zone-like structure can be 35 confirmed by identifying Rdh10 positive, Otx1 positive, or Zic1

positive cells with immunostaining a cryosection of aggregate.
[0247]

In a further embodiment, differentiation of a retinal tissue (neuroepithelium) contained in the aggregate by the
5 aforementioned steps (A), (B) proceeds, and a mature retinal tissue and the aforementioned cells containing at least one, preferably plural, more preferably all, cells selected from the group consisting of photoreceptor precursor cell, photoreceptor cell, cone photoreceptor cell, rod photoreceptor cell,
10 horizontal cell, interneuron (amacrine cell, ganglion cell etc.) can be produced.
[0248]

A retinal tissue containing a retinal pigment epithelial cell can be produced by the following step (C) from a cell
15 aggregate containing a retinal tissue obtained by the production method, i.e., the aforementioned step (1) - step (3), of the present invention and the like. A retinal pigment epithelial sheet can be produced by the following step (D) from a retinal pigment epithelial cell obtained by the following
20 step (C).
[0249]

The "retinal pigment epithelial cell" in the present invention means an epithelial cell present on the outside of the neural retinal tissue in retina in vivo. Whether it is a
25 retinal pigment epithelial cell can be confirmed by those of ordinary skill in the art based on, for example, expression of a cell marker (RPE65 (matured retinal pigment epithelial cell), Mitf (juvenile or matured retinal pigment epithelial cell) and the like), the presence of melanin granule, characteristic cell
30 morphology of polygon and the like.
[0250]

First, in step (C), a cell aggregate containing a retinal tissue obtained by the production method of the present invention is cultured in suspension in a serum-free medium or
35 serum-containing medium free of a BMP signal transduction

pathway activating substance but containing a Wnt signal transduction pathway activating substance to give an aggregate containing retinal pigment epithelial cells.

[0251]

5 As a serum-free medium to be used in step (C), a serum-free medium which is a basal medium supplemented with N2 or KSR can be mentioned. More specifically, a serum-free medium which is a DMEM/F-12 medium supplemented with N2 supplement (Life Technologies) can be mentioned. As the serum-containing medium,
10 a serum-containing medium which is a basal medium supplemented with fetal bovine serum can be mentioned.

[0252]

 The serum-free medium to be used in step (C) may contain, in addition to the aforementioned Wnt signal transduction
15 pathway activating substance, the aforementioned Nodal/Activin signal transduction pathway activating substance, and/or the aforementioned FGF signal transduction pathway inhibiting substance.

[0253]

20 A preferable culture in step (C) is, for example, suspension culture.

[0254]

 Step (D) in which the aggregate obtained in step (C) of the present invention is dispersed and the obtained cells are
25 cultured in an adhered state is explained.

[0255]

 Step (D) is performed within 60 days, preferably within 30 days, more preferably 3 days, after the start of step (C).

[0256]

30 The medium (basal medium) in step (D) is not particularly limited, and the basal medium described in the above-mentioned definition can be appropriately selected. As a serum-free medium or serum-containing medium to be used for adhesion culture in step (D), the aforementioned medium can be mentioned.
35 To avoid complicated preparation, a serum-free medium

supplemented with an appropriate amount of a commercially available serum alternative such as KSR and the like (e.g., a medium of 1:1 mixture of DMEM/F-12 and Neurobasal supplemented with 1/2 x N2 supplement, 1/2 x B27 supplement and 100 μ M 2-mercaptoethanol) is preferably used. The amount of KSR to be added to the serum-free medium is, for example, generally about 1% to about 20%, preferably about 2% to about 20%, in the case of a cell derived from human iPS cell.

[0257]

10 In step (D), it is preferable to culture cells in the aforementioned serum-free medium or serum-containing medium containing a ROCK inhibiting substance.

[0258]

In step (D), it is more preferable to culture cells in a serum-free medium or serum-containing medium further containing one or more substances selected from the group consisting of a Wnt signal transduction pathway activating substance, an FGF signal transduction pathway inhibiting substance, an Activin signal transduction pathway activating substance and a BMP signal transduction pathway activating substance.

[0259]

The Activin signal transduction pathway activating substance is a substance capable of enhancing a signal mediated by Activin. Examples of the Activin signal transduction pathway activating substance include proteins belonging to the Activin family (e.g., Activin A, Activin B, Activin C, Activin AB and the like), Activin receptor, and Activin receptor agonist.

[0260]

30 The concentration of the Activin signal transduction pathway activating substance to be used in step (D) may be any concentrations as long as a uniformed sheet of retinal pigment epithelial cells can be efficiently formed. For example, Recombinant Human/Mouse/Rat Activin A (R&D systems, #338-AC) is added to a concentration of about 1 ng/ml to about 10 μ g/ml,

preferably about 10 ng/ml to about 1 µg/ml, more preferably about 100 ng/ml. An Activin signal transduction pathway activating substance is added, for example, within 18 days, preferably on day 6, from the start of step (D).

5 [0261]

In step (D), adhesion culture is preferably performed on a culture vessel whose surface is treated with a culture substrate. As a culture substrate to be used for treating culture vessel in step (D), a cell culture substrate enabling
10 adhesion culture of aggregate-derived cells and formation of a retinal pigment epithelial sheet can be mentioned.

[0262]

As mentioned above, the production method of a retinal cell or retinal tissue containing step (A) and step (B) or step
15 (C) and step (D) in addition to the above-mentioned step (1) - step (3) is also within the scope of the present invention. Step (3) may include the above-mentioned step (A) and step (B) or step (C) and step (D) and the like.

[0263]

20 4. Method for evaluating toxicity or efficacy

Since a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention is useful as a material for disease study or drug discovery in a screening
25 for a medicament for treating a disease due to a disorder of a retinal tissue or retinal cells, or in toxicity evaluation, it can be used as a reagent for evaluating toxicity or efficacy of a test substance. For example, iPS cells are produced from a human patient with a disease due to a disorder of a retinal
30 tissue, particularly a hereditary disease, and using the iPS cells, a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) is produced by the method of the present invention. The retinal tissue or retinal cells may reproduce the disorder of the
35 retinal tissue causing the disease of the patient in vitro.

Therefore, the present invention provides a method for evaluating toxicity or efficacy of a test substance, which comprises contacting the test substance with a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention, and detecting an influence of the substance on the cells or tissues.

[0264]

For example, a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) having a particular disorder (e.g., hereditary disorder), which is produced by the production method of the present invention, are cultured in the presence or absence (negative control) of a test substance. Then, the severity of disorder of the retinal tissue or retinal cells treated with the test substance is compared with that of the negative control. As a result, a test substance that reduced the severity of the disorder can be selected as a candidate substance for a medicament for treating the disease resulting from the disorder. For example, a test substance that improves the physiological activity (e.g., survival promotion or maturation) of retinal cells produced by the production method of the present invention can be explored as a candidate substance of a pharmaceutical product. Alternatively, according to the production method of the present invention, retinal cells are prepared by inducing differentiation of the induced pluripotent stem cells which are prepared from a somatic cell having a gene mutation that causes a particular disorder such as a retinal disease and the like. A candidate of a test substance effective as a medicament or prophylactic drug for the disorder can be explored based on whether the retinal cells added with a test substance show the aforementioned disorder, as an index.

[0265]

For toxicity evaluation, a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific

neural cell) produced by the production method of the present invention are cultured in the presence or absence (negative control) of a test substance. Then, the severity of toxicity on the retinal tissue or retinal cells treated with the test
5 substance is compared with that of the negative control. As a result, a test substance that exerted toxicity as compared to the negative control can be judged as a substance having toxicity to the retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell).

10 [0266]

That is, the present invention encompasses a method for evaluating toxicity comprising the following steps:
(step 1) a step of culturing a retinal tissue or retinal cells produced by the production method of the present invention
15 under viable culture conditions for a given time in the presence of a test substance, and measuring the severity of cell injury,
(step 2) a step of culturing a retinal tissue or retinal cells produced by the production method of the present invention
20 under viable culture conditions for a given time in the absence of a test substance or in the presence of a positive control, and measuring the severity of cell injury,
(step 3) a step of evaluating the toxicity of the test substance in step 1, based on the difference in the results
25 measured in (step 1) and (step 2).

As used herein, "in the absence of a test substance" encompasses adding only a medium or a solvent used to dissolve the test substance instead of adding a test substance. In addition, "positive control" means a known compound having
30 toxicity. Examples of the method for measuring the severity of cell injury include a method for measuring the number of viable cells, for example, a method for measuring intracellular ATP amount, a method for measuring the number of viable cells by cell staining (e.g., nucleus staining) and morphology
35 observation and the like.

[0267]

In step 3, as a method for evaluating the toxicity of a test substance, the measurement value in step 1 and the measurement value of the negative control in step 2 are compared, and when the severity of cell injury in step 1 is high, the test substance can be judged to have toxicity. In addition, the measurement value in step 1 and the measurement value of the positive control in step 2 are compared, and when the severity of cell injury in step 1 is the same or above, the test substance can be judged to have toxicity.

[0268]

5. Pharmaceutical composition

The present invention provides a pharmaceutical composition containing an effective amount of a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention.

[0269]

The pharmaceutical composition contains an effective amount of a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention, and a pharmaceutically acceptable carrier.

[0270]

As a pharmaceutically acceptable carrier, a physiological aqueous solvent (saline, buffer, serum-free medium etc.) can be used. Where necessary, in a transplantation therapy, a medicament containing a tissue or cells to be transplanted may contain conventionally used preservative, stabilizer, reducing agent, isotonizing agent and the like.

[0271]

The pharmaceutical composition of the present invention can be produced as a suspension by suspending retinal tissues or retinal cells produced by the production method in an appropriate physiological aqueous solvent. Where necessary,

the composition may be added with a cryopreservative, cryopreserved, thawed when in use, washed with buffer, and used for a transplantation therapy.

[0272]

5 A retinal tissue obtained by the production method of the present invention may also be cut in an appropriate size with tweezers and the like to give a sheet preparation.

[0273]

Cells obtained by the production method of the present
10 invention may also be subjected to adhesion culture in step (3) for differentiation induction to form a sheet-like cells to give a sheet preparation.

[0274]

The pharmaceutical composition of the present invention
15 is useful as a medicament for a disease due to a disorder of a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell).

[0275]

6. Therapeutic drug (Medicament)

20 A retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention is useful for a transplantation therapy for a disease due to a disorder of a retinal tissue or retinal cells. Thus, the present invention
25 provides a medicament for treating a disease due to a disorder of a retinal tissue or retinal cells, which contains a retinal tissue or retinal cells (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention. A retinal tissue or retinal cells
30 (e.g., retinal progenitor cell, retinal layer-specific neural cell) produced by the production method of the present invention can be used as a medicament for treating the disease due to a disorder of a retinal tissue or retinal cells or to complement the corresponding damaged site in a damaged state of
35 a retinal tissue. A disease due to a disorder of a retinal

tissue or retinal cells, and a damaged state of a retinal tissue can be treated by transplanting a retinal tissue or retinal cells produced by the production method of the present invention to a patient with a disease due to a disorder of a
5 retinal tissue or retinal cells, or a damaged state of a retinal tissue, who requires transplantation, to complement the retinal cells or disordered retinal tissue itself. Examples of the disease due to a disorder of a retinal tissue or retinal cell include macular degeneration, age-related macular
10 degeneration, retinal pigment denaturation, cataract, glaucoma, cornea disease, retinopathy and the like which are ophthalmologic diseases.

[0276]

In transplantation therapy, rejection due to the
15 difference in histocompatibility antigens often poses a problem. However, the problem can be solved by using pluripotent stem cells (e.g., induced pluripotent stem cells) established from the somatic cells of the transplantation recipient. That is, in a preferable embodiment, pluripotent stem cells (e.g.,
20 induced pluripotent stem cells) established from the somatic cells of the recipient are used as pluripotent stem cells in the method of the present invention, and a retinal tissue or retinal cells, which is immunologically self for the recipient, are produced and transplanted to the recipient.

25 [0277]

In addition, an allogenic retinal tissue or retinal cell may be produced from a pluripotent stem cell (e.g., induced pluripotent stem cell) established from a somatic cell of others who are immunologically compatible with the recipient
30 (e.g., compatible in HLA type and MHC type), and transplanted to the recipient.

[Examples]

[0278]

The present invention is explained in detail in the
35 following by referring to Examples and Pharmacological Examples,

Examples, which are not to be construed as limitative.

[0279]

Example 1: Production Example of retinal tissue from human iPS cell using Wnt signal transduction pathway inhibiting substance
5 in step 2 and step 3 after step 1 (also called Precondition)

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured under feeder free according to the method described in "Scientific Reports, 4, 3594 (2014)". As a feeder-free medium, StemFit (registered trade mark) medium
10 (AK03N, manufactured by Ajinomoto Co., Inc.) was used, and as a feeder-free scaffold, Laminin 511-E8 (manufactured by Nippi, Inc.) was used.

[0280]

As a specific maintenance culture operation, subconfluent
15 human iPS cells (1231A3 strain) were first washed with PBS, and dispersed into single cells by using TrypLE Select (manufactured by Life Technologies). Thereafter, the aforementioned human iPS cells dispersed into single cells were seeded in a plastic culture dish coated with Laminin 511-E8,
20 and cultured under feeder-free conditions in StemFit (registered trade mark) medium in the presence of Y27632 (ROCK inhibiting substance, 10 μ M). When a 6-well plate (manufactured by Iwaki, for cell culture, culture area 9.4 cm²) was used as the aforementioned plastic culture dish, the number
25 of plated cells for the aforementioned human iPS cells dispersed into single cells was adjusted to 1.0×10^4 . One day after seeding, the entire amount of the medium was changed with StemFit (registered trade mark) medium free of Y27632. Thereafter, once in 1 - 2 days, the entire amount of the medium
30 was changed with StemFit (registered trade mark) medium free of Y27632. Thereafter, the cells were cultured until 6 days after seeding when they became subconfluent (60% of culture area is covered with cells).

[0281]

35 As a specific example of step 1 (Precondition) of the

present production method, the following operation was performed. Subconfluent human iPS cells (1231A3 strain) were first washed with PBS, and dispersed into single cells by using TrypLE Select (manufactured by Life Technologies). Thereafter, 5 the aforementioned human iPS cells dispersed into single cells were seeded in a plastic culture dish (manufactured by Iwaki) coated with Laminin 511-E8, and cultured under feeder-free conditions in StemFit (registered trade mark) medium in the presence of Y27632 (ROCK inhibiting substance, 10 μ M). When a 10 6-well plate (manufactured by Iwaki, culture area 9.4 cm²) was used as the aforementioned plastic culture dish, the number of plated cells for the aforementioned human iPS cells dispersed into single cells was adjusted to 1.0×10^4 . When a 60 mm dish (manufactured by Iwaki, for cell culture, culture area 21 cm²) 15 was used as the aforementioned plastic culture dish, the number of plated cells for the aforementioned human iPS cells dispersed into single cells was adjusted to 2.0×10^4 . One day after seeding, the entire amount of the medium was changed with StemFit (registered trade mark) medium free of Y27632. 20 Thereafter, once in 1 - 2 days, the entire amount of the medium was changed with StemFit (registered trade mark) medium free of Y27632. Thereafter, the cells were cultured until 5 days after seeding, i.e., one day before subconfluence (50% of culture area is covered with cells). Even when cultured for 6 days 25 after seeding, similar results were obtained.

[0282]

The aforementioned feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day in the presence of SB431542 (TGF β signal transduction 30 pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition treatment, Fig. 1 "Precondition: TGF β R-i+Shh") or in the absence thereof (step 1: without Precondition treatment, Fig. 1 "Untreated control") in a Stem 35 Fit (registered trade mark) medium (AK03N; manufactured by

Ajinomoto Co., Inc.). The cultured cells were subjected to bright field observation using an inverted microscope (manufactured by KEYENCE CORPORATION, BIOREVO). As a result, it was found that a treatment with a TGF β signal transduction pathway inhibiting substance (SB431542) and a Sonic hedgehog signal transduction pathway activating substance during the feeder-free culture did not exert a large influence on the morphology of human iPS cells.

[0283]

10 The thus-prepared human iPS cells free of a Precondition treatment and Precondition-treated human iPS cells were treated with a cell dispersion solution by using TrypLE Select (manufactured by Life Technologies), and further dispersed into single cells by a pipetting operation. Thereafter, the above
15 human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium in a non-cell-adhesive 96 well culture plate (PrimeSurface 96V-bottom plate, SUMITOMO BAKELITE) at 1.0×10^4 cells per well, and subjected to suspension culture at 37°C, 5% CO₂. As a serum-free medium
20 (gfCDM+KSR) therefor, a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x chemically defined lipid concentrate was used.

[0284]

At the start of the suspension culture (on day 0 after
25 the start of suspension culture, step 2 start), Y27632 (final concentration 20 μ M) was added to the aforementioned serum-free medium, and the cells were further cultured in a medium containing or not containing a Wnt signal transduction pathway inhibiting substance (IWR-1e, 3 μ M). A cell aggregate was
30 formed under conditions with or without Precondition by day 2 after the start of suspension culture (step 2 completed, and step 3 started).

[0285]

On day 3 after the start of suspension culture, a medium
35 not containing Y27632, containing human recombinant BMP4

(manufactured by R&D) and further containing or not containing IWR-1e was added by 50 µl such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change.

5 Thereafter, a half-medium exchange operation using the aforementioned serum-free medium not containing Y27632, SAG and human recombinant BMP4 and further containing or not containing IWR-1e was performed once every 2 - 4 days to avoid change of the concentration of exogenous IWR-1e.

10 [0286]

The thus-prepared cells on day 17 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti) (Fig. 1). As a result, under conditions without
15 Precondition and without addition of a Wnt signal transduction pathway inhibiting substance, aggregate growth was poor and the shape thereof collapsed (Fig. 1A). In contrast, a cell aggregate not deformed and having a dense interior was obtained under conditions with addition of a Wnt signal transduction
20 pathway inhibiting substance, and it was found that a neural tissue was formed (Fig. 1B). Furthermore, under conditions with Precondition and without addition of a Wnt signal transduction pathway inhibiting substance (Fig. 1C) or with addition of a Wnt signal transduction pathway inhibiting
25 substance (Fig. 1D), an undeformed round cell aggregate with a smooth surface and dense inside of aggregate was obtained, and it was found that a neural tissue was formed.

[0287]

The above cell aggregates on day 17 after the start of
30 suspension culture were each fixed with 4% para-formaldehyde and cryosections were prepared. These cryosections were immunostained for Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers, or Chx10 (anti-Chx10 antibody, manufactured by Exalpha, sheep), which is one
35 of the retinal tissue markers. These immunostained sections

were observed using an inverted fluorescence microscope
(manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 2).

[0288]

As a result, a cell aggregate produced under conditions
5 without Precondition with SB431542 and SAG in step 1, and
without addition of IWR-1e in step 2 and step 3, it was found
that the proportion rate of Rx positive cells to the total
cells was not more than about 3% (Fig. 2A).

In the cell aggregate produced under conditions not
10 including Precondition with SB431542 and SAG in step 1 and
including addition of IWR-1e in step 2 and step 3, the
proportion of Rx positive cells in the total cells was about
40% (Fig. 2B). From the analysis of serial sections, weak
Chx10 positiveness was observed in the cell aggregate produced
15 under these conditions (Fig. 2F).

In the cell aggregate produced under conditions including
Precondition with SB431542 and SAG in step 1 and no addition of
IWR-1e in step 2 and step 3, it was found that the proportion
of Rx positive cells in the total cells was about 40% (Fig. 2C).
20 From the analysis of serial sections, cell aggregate produced
under these conditions, about 40% of Chx10 positive (strong
positive) cells was observed (Fig. 2G).

Furthermore, it was found that, in the cell aggregate
produced under conditions including Precondition with SB431542
25 and SAG in step 1 and including addition of IWR-1e in step 2
and step 3, the proportion of Rx positive cells in the total
cells was about 60% (Fig. 2D). From the analysis of serial
sections, in a cell aggregate produced under these conditions,
about 60% of Chx10 positive (strong positive) cells were
30 observed (Fig. 2H).

[0289]

From these results, it was found that a neural tissue was
hardly formed under conditions not including Precondition and
not including addition of a Wnt inhibitor in step 2 and step 3,
35 whereas a neural tissue can be formed under any other

conditions.

As compared to the conditions without Precondition and without addition of Wnt inhibitor in step 2 and step 3, it was found that retinal tissue formation is somewhat promoted under conditions without Precondition and with addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3 (Fig. 1A, B, Fig. 2A, B, E, F). On the other hand, it was found that a retinal tissue can be highly efficiently formed with Precondition with a TGF β family signal transduction pathway inhibiting substance and a Shh signal transduction pathway activating substance in step 1 and further addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3 (Fig. 2B, D, F, H).

[0290]

Example 2: Production Example of retinal tissue from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and addition of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day in the presence of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition treatment, Fig. 3 "Precondition: Shh") or in the absence thereof (step 1: Precondition-free conditions, Fig. 3 "Untreated control") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

[0291]

The thus-prepared Precondition-treated human iPS cells,

and human iPS cells under Precondition-free conditions were each treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed
5 into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium
10 (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of
15 step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following five Conditions 1 - 5.

.Condition 1. Conditions not including Precondition treatment
20 in step 1 and not including addition of an exogenous Wnt signal transduction pathway inhibiting substance or an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 3A).

25 .Condition 2. Condition including Precondition treatment in step 1 and not including addition of an exogenous Wnt signal transduction pathway inhibiting substance or an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2
30 (Fig. 3B).

.Condition 3. Condition including Precondition treatment in step 1, addition of IWR-1e (3 μ M) as an Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance
35 to the aforementioned serum-free medium at the start of step 2

(Fig. 3C).

.Condition 4. Condition including Precondition treatment in step 1, no addition of an exogenous Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as a
5 Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 3D).

.conditions 5. Condition including Precondition treatment in step 1, addition of IWR-1e (3 μ M) as a Wnt signal transduction
10 pathway inhibiting substance and addition of SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 3E).

By day 2 after the start of suspension culture, cell
15 aggregates were formed under any of Condition 1-5 (step 2 completed, and step 3 started).

[0292]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4
20 (manufactured by R&D) and further containing or not containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change.

Thereafter, a half-medium exchange operation using the
25 aforementioned serum-free medium not containing Y27632, SAG and human recombinant BMP4 and further containing or not containing IWR-1e was performed once every 2 - 4 days to avoid change of the concentration of exogenous IWR-1e.

[0293]

30 The thus-prepared cells on day 17 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti) (Fig. 3). As a result, it was found that an aggregate was not grown and a neural tissue was not formed
35 under Condition 1 without Precondition in step 1 (Fig. 3A),

whereas aggregate was grown and a neural tissue could be formed under Conditions 2 - 5 with a Precondition treatment with a Sonic hedgehog signal transduction pathway activating substance in step 1 (Fig. 3B-E). Particularly, it was found that the formation efficiency of a neural tissue is good under Conditions 3 and 5 wherein a Wnt signal transduction pathway inhibiting substance was allowed to act in step 2 and step 3 (Fig. 3C, E), as compared to Conditions 2 and 4 wherein a Wnt signal transduction pathway inhibiting substance was not allowed to act (Fig. 3B, D).

[0294]

The above cell aggregates on day 17 after the start of suspension culture were each fixed with 4% para-formaldehyde to give cryosections. These cryosections were immunostained for Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers, or Chx10 (anti-Chx10 antibody, manufactured by Exalpha, sheep), which is one of the retinal tissue markers. These immunostained sections were observed using an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 4).

[0295]

As a result, it was found that a neural tissue was not formed under Condition 1 without Precondition, whereas a neural tissue was formed under any of Conditions 2 - 5 with a Precondition treatment with SAG.

Of these, under Condition 2 with Precondition with SAG in step 1 and without addition of an exogenous Wnt signal transduction pathway inhibiting substance in step 2 and step 3, the proportion of Rx positive cells in the total cells was about 40% (Fig. 4A). From the analysis of serial sections, it was found that, in the retinal tissue of the cell aggregate produced under the aforementioned conditions, Rx positive cell was a Chx10 co-positive (Fig. 4E).

On the other hand, it was found that the proportion of Rx positive cells in the total cells is about 90% in a cell

aggregate produced under Condition 3 with Precondition with SAG in step 1, and addition of IWR-1e in step 2 and step 3 (Fig. 4B). From the analysis of serial sections, it was found that, in the retinal tissue of the cell aggregate produced under the
5 aforementioned conditions, the Rx positive cell is a Chx10 co-positive (Fig. 4F).

That is, from the comparison of Condition 2 and Condition 3, it was found that a retinal tissue can be formed with about 90% efficiency by Precondition in step 1 and a further
10 treatment with an Wnt signal transduction pathway inhibiting substance in step 2 and step 3.

In the cell aggregate produced under Condition 4 including Precondition with SAG in step 1 and including Condition 4 with addition of SAG in step 2, the proportion of
15 Rx positive cells in the total cells was about 60% (Fig. 4C). From the analysis of serial sections, it was found that, in the retinal tissue of the cell aggregate produced under the above-mentioned conditions, Rx positive cell is a Chx10 co-positive (Fig. 4G).

20 On the other hand, it was found that, in the cell aggregate produced under conditions including Precondition with SAG in step 1 and Condition 5 with addition of SAG in step 2 and addition of IWR-1e in step 2 and step 3, the proportion of Rx positive cells in the total cells was about 90% (Fig. 4D).
25 From the analysis of serial sections, it was found that, in the retinal tissue of the cell aggregate produced under the aforementioned conditions, Rx positive cell is a Chx10 co-positive (Fig. 4H).

[0296]

30 That is, from the comparison of Condition 4 and Condition 5, it was found that a retinal tissue can be formed with about 90% efficiency by Precondition in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, and a further treatment with a Wnt signal transduction
35 pathway inhibiting substance in step 2 and step 3.

[0297]

Example 3: Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction pathway inhibiting substance in step 1 and addition of Wnt
5 signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade
10 mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under
15 conditions of addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance (step 1: Precondition (TGF β R-i) treatment, Fig. 5 "Precondition: TGF β R-i"), or in the presence of SB431542
20 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 5 "Precondition: TGF β R-i+Shh"), in a Stem Fit (registered trade mark) medium (AK03N;
25 manufactured by Ajinomoto Co., Inc.).

[0298]

The thus-prepared Precondition (TGF β R-i)-treated human iPS cells, and Precondition (TGF β R-i+Shh)-treated human iPS cells were each treated with the cell dispersion solution by
30 using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate,
35 manufactured by SUMITOMO BAKELITE). Thereafter, the cells were

subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 µM 1-monothioglycerol, 1 x Chemically defined
5 lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 µM) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following three
10 Conditions 1 - 3.

•Condition 1. Condition including Precondition (TGFβR-i) treatment in step 1 and addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating
15 substance to the aforementioned serum-free medium at the start of step 2 (Fig. 5A).

•Condition 2. Condition including Precondition (TGFβR-i+Shh) treatment in step 1, and addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and no
20 addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 5B).

•Condition 3. Condition including Precondition (TGFβR-i+Shh) treatment in step 1, and addition of IWR-1e (3 µM) as a Wnt
25 signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 5C).

By day 2 after the start of suspension culture, cell
30 aggregates were formed under any of the aforementioned Condition 1-5 (step 2 completed, and step 3 started).

[0299]

On day 3 after the start of suspension culture, the aforementioned serum-free medium not containing Y27632 or SAG,
35 containing human recombinant BMP4 (manufactured by R&D) and

further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation using
5 the aforementioned serum-free medium not containing Y27632, SAG and human recombinant BMP4 and further containing IWR-1e was performed once every 2 - 4 days to avoid change of the concentration of exogenous IWR-1e.

The thus-prepared cells on day 17 after the start of
10 suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue could be formed under any of Condition 1-3.

[0300]

15 Cell aggregates on day 17 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx
20 antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These immunostained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO). As a result, it was found that a neural tissue was formed and the proportion of Chx10 positive
25 in the neural tissues is about 90% under any of Conditions 1 - 3 (Fig. 5A-C). From the analysis of serial sections, in the retinal tissues, these Chx10 positive cells could be confirmed to be Rx co-positive.

That is, it was found that a retinal tissue can be formed
30 efficiently from human iPS cells under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance in step 1 and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3.

[0301]

35 Example 4: Production Example of retinal tissue from human iPS

cell including Precondition with BMP signal transduction pathway inhibiting substance in step 1 and addition of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

5 Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi,
10 Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions of addition of LDN193189 (BMP signal transduction pathway inhibiting substance (BMPR-i), 100 nM) and no addition
15 of an exogenous Sonic hedgehog signal transduction pathway activating substance (step 1: Precondition (BMPR-i) treatment, Fig. 6 "Precondition: BMPR-i"), or in the presence of LDN193189 (BMP signal transduction pathway inhibiting substance, 100 nM, BMPR-i) and SAG (Sonic hedgehog signal transduction pathway
20 activating substance (Shh), 300 nM) (step 1: Precondition (BMPR-i+Shh) treatment, Fig. 6 "Precondition: BMPR-i+Shh"), in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

[0302]

25 The thus-prepared Precondition (BMPR-i)-treated human iPS cells, and Precondition (BMPR-i+Shh)-treated human iPS cells were each treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS
30 cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-
35 free medium (gfCDM+KSR) therefor, a serum-free medium which is

a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following three Conditions 1 - 3.

•Condition 1. Condition including Precondition (BMPR-i) treatment in step 1 and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 6A).

•Condition 2. Condition including Precondition (BMPR-i+Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 6B).

•Condition 3. Condition including Precondition (BMPR-i+Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 6C).

By day 2 after the start of suspension culture, cell aggregates were formed under any of the aforementioned Condition 1-3 (step 2 completed, and step 3 started).

[0303]

On day 3 after the start of suspension culture, the aforementioned serum-free medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5

nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation using the aforementioned serum-free medium not containing Y27632, SAG and human recombinant BMP4 and further containing IWR-1e was
5 performed once every 2 - 4 days to avoid change of the concentration of exogenous IWR-1e.

The thus-prepared cells on day 17 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation,
10 ECLIPSE Ti). As a result, it was found that a neural tissue could be formed under any of Condition 1-3.

[0304]

Cell aggregates on day 17 after the start of the aforementioned suspension culture were fixed with 4% para-
15 formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These immunostained sections were observed
20 with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO). As a result, it was found that a neural tissue was formed and the proportion of Chx10 positive in the neural tissues is about 90% under any of Conditions 1 - 3 (Fig. 6A-C). From the analysis of serial sections, in the
25 retinal tissues, these Chx10 positive cells could be confirmed to be Rx co-positive.

That is, it was found that a retinal tissue can be formed efficiently from human iPS cells under conditions including Precondition with a BMP signal transduction pathway inhibiting
30 substance in step 1 and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3.

[0305]

Example 5: Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction
35 pathway inhibiting substance and/or a Sonic hedgehog signal

transduction pathway activating substance in step 1, addition
of Wnt signal transduction pathway inhibiting substance in step
2 and step 3, addition of BMP signal transduction pathway
activating substance in step 3 and use of Sonic hedgehog signal
5 transduction pathway activating substance and serum

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a
10 feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including no addition of an exogenous TGF β signal
15 transduction pathway inhibiting substance and addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (Shh) treatment, Fig. 7 "Precondition: Shh"), and conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting
20 substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 7 "Precondition: TGF β R-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

25 [0306]

The thus-prepared Precondition (Shh)-treated human iPS cells or Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells
30 by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to
35 suspension culture at 37°C, 5% CO₂. As the serum-free medium

(gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following four Conditions 1 - 4.

10 .Condition 1. Condition including Precondition (Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 7A).

.Condition 2. Condition including Precondition (Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 7B).

.Condition 3. Condition including Precondition (TGF β R-i+Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 7C).

.Condition 4. Condition including Precondition (TGF β R-i+Shh) treatment in step 1, and addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 7D).

By day 2 after the start of suspension culture, cell aggregates were formed under any of the aforementioned

Condition 1-4 (step 2 completed, and step 3 started).

[0307]

On day 3 after the start of suspension culture, the
aforementioned serum-free medium not containing Y27632 or SAG,
5 containing human recombinant BMP4 (manufactured by R&D) and
further containing IWR-1e was added by 50 μ l such that a final
concentration of exogenous human recombinant BMP4 would be 1.5
nM (55 ng/ml) and the concentration of exogenous IWR-1e would
not change. Thereafter, a half-medium exchange operation using
10 the aforementioned serum-free medium not containing Y27632, SAG
and human recombinant BMP4 and further containing IWR-1e was
performed once every 2 - 4 days to avoid change of the
concentration of exogenous IWR-1e.

Next, on day 10 after the start of suspension culture,
15 using Shh serum-containing medium obtained by adding SAG (100
nM) as a Sonic hedgehog signal transduction pathway activating
substance and fetal calf serum (10%) as a serum to the
aforementioned serum-free medium (Fig. 7 "Shh+serum"), an
operation to exchange about 80% of the medium (80% exchange
20 operation) was repeated 3 times, thereby changing the medium to
the aforementioned Shh serum-containing medium not containing
Y27632, human recombinant BMP4 or IWR1-e, and containing SAG
and the serum. Thereafter, once every 2 - 4 days, a half-
medium exchange operation with the aforementioned Shh serum-
25 containing medium was performed.

The thus-prepared cells on day 17 after the start of
suspension culture were subjected to bright field observation
using an inverted microscope (manufactured by Nikon Corporation,
ECLIPSE Ti). As a result, it was found that a neural tissue
30 could be formed under any of Conditions 1-4.

[0308]

Cell aggregates on day 17 after the start of the
aforementioned suspension culture were fixed with 4% para-
formaldehyde to produce cryosections. These cryosections were
35 immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep),

which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These immunostained sections were observed with an inverted fluorescence microscope (manufactured by
5 KEYENCE CORPORATION, BIOREVO). As a result, it was found that a neural tissue was formed and the proportion of Chx10 positive in the neural tissues is about 80% under any of Conditions 1 - 4 (Fig. 7A-D). From the analysis of serial sections, in the retinal tissues, these Chx10 positive cells could be confirmed
10 to be Rx co-positive.

That is, it was found that a retinal tissue can be formed efficiently from human iPS cells under conditions including Precondition in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and a BMP signal
15 transduction pathway inhibiting substance in step 3, followed by culturing in Shh serum-containing medium.

[0309]

Example 6: Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction
20 pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance and TGF β signal transduction pathway inhibiting substance in step 2 and step 3

25 Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi,
30 Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and
35 SAG (Sonic hedgehog signal transduction pathway activating

substance (Shh), 300 nM) (step 1: Precondition (TGFβR-i+Shh) treatment, Fig. 8 "Precondition: TGFβR-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

5 [0310]

The thus-prepared Precondition (TGFβR-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human
10 iPS cells dispersed into single cells were suspended in 100 μl of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As
15 the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μM 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of
20 suspension culture, start of step 2), Y27632 (final concentration 20 μM) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following four Conditions 1 - 4.

•Condition 1. Condition including addition of IWR-1e (3 μM) as
25 a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 8A).

•Condition 2. Condition including addition of IWR-1e (3 μM) as
30 a Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as a Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 8B).

•Condition 3. Condition including addition of IWR-1e (3 μM) as
35 a Wnt signal transduction pathway inhibiting substance and

SB431542 (5 μ M) as a TGF β signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2
5 (Fig. 8C).

•Condition 4. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SB431542 (5 μ M) as a TGF β signal transduction pathway inhibiting substance and SAG (30 nM) as a Sonic hedgehog signal
10 transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 8D).

By day 2 after the start of suspension culture, cell aggregates were formed under any of Conditions 1-4 (step 2 completed, and step 3 started).

15 [0311]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing or not containing IWR-1e and SB431542 was added by 50 μ l such that a final
20 concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e and SB431542 would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG,
25 human recombinant BMP4 and further containing or not containing IWR-1e and SB431542 to avoid change of the concentration of exogenous IWR-1e and SB431542.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times using the
30 aforementioned serum-free medium so that the concentration of exogenous IWR-1e and/or SB431542 would be 3% or less as compared to that before the medium exchange. Thereafter, once every 2 - 4 days, a half-medium exchange operation was performed using the aforementioned serum-free medium free of
35 Y27632, SAG, human recombinant BMP4, IWR-1e and SB431542.

The thus-prepared cells on day 19 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue
5 could be formed under any of Conditions 1-4.

[0312]

Cell aggregates on day 19 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were
10 immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. As counterstaining, nucleic acid was stained with DAPI. These stained sections were observed with an
15 inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 8).

[0313]

As a result, it was found that a neural tissue was formed under any of Conditions 1 - 4. In Condition 1, the proportion
20 of the Chx10 positive retinal tissue was about 70%, in Condition 2, the proportion of the Chx10 positive retinal tissue was about 70%, in Condition 3, the proportion of the Chx10 positive retinal tissue was about 70%, and in Condition 4, the proportion of the Chx10 positive retinal tissue was about
25 80%. From the analysis of serial sections, these Chx10 positive cells could be confirmed to be Rx co-positive cells.

That is, it was found that efficient differentiation into a retinal tissue is also achieved under conditions including addition of, as in Condition 1 and Condition 2, a Wnt signal
30 transduction pathway inhibiting substance in step 2 and step 3 and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3. Furthermore, it was found that efficient differentiation into a retinal tissue is also achieved under
35 conditions including addition of, as in Conditions 3 and 4, a

Wnt signal transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance in step 2 and step 3.

[0314]

5 Example 7: Production Example of retinal tissue from human iPS cell including Precondition with BMP signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance and TGF β
10 signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade
15 mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under
20 conditions including addition of LDN193189 (BMP signal transduction pathway inhibiting substance (BMPR-i), 100 nM) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (BMPR-i+Shh) treatment, Fig. 9 "Precondition: BMPR-i+Shh") in a Stem Fit
25 (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (BMPR-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into
30 single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were
35 subjected to suspension culture at 37°C, 5% CO₂. As the serum-

free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of
5 suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in the serum-free medium under the following two Conditions 1 - 2.

10 •Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 9A).

15 •Condition 2. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SB431542 (5 μ M) as a TGF β signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to
20 the aforementioned serum-free medium at the start of step 2 (Fig. 9B).

By day 2 after the start of suspension culture, cell aggregates were formed under Condition 1 and Condition 2 (step 2 completed, and step 3 started).

25 [0315]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing or not containing IWR-1e and SB431542 was added by 50 μ l such that a final
30 concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4
35 and further containing or not containing IWR-1e and SB431542 to

avoid change of the concentration of exogenous IWR-1e and SB431542.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times using the
5 aforementioned serum-free medium so that the concentration of exogenous IWR-1e and/or SB431542 would be 3% or less as compared to that before the medium exchange. Thereafter, once every 2 - 4 days, a half-medium exchange operation was performed using the aforementioned serum-free medium free of
10 Y27632, SAG, human recombinant BMP4, IWR-1e and SB431542.

The thus-prepared cells on day 19 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue
15 could be formed under Condition 1 and Condition 2.
[0316]

Cell aggregates on day 19 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were
20 immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. As counterstaining, nucleic acid was stained with DAPI. These stained sections were observed with an
25 inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 9).

[0317]

As a result, it was found that a neural tissue could be formed in Condition 1 and Condition 2. In Condition 1, the
30 proportion of the Chx10 positive retinal tissue was about 80%, and in Condition 2, the proportion of the Chx10 positive retinal tissue was about 90%. From the analysis of serial sections, it could be confirmed that these Chx10 positive cells are Rx co-positive cells.

35 That is, in Condition 1, it was found that efficient

differentiation into a retinal tissue is also achieved under conditions including Precondition with a BMP signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3. Furthermore, in Condition 2, it was found that efficient differentiation into retinal cell is also achieved under conditions including Precondition with a BMPR inhibitor and a Sonic hedgehog signal transduction pathway activating substance in step 1, and addition of a Wnt signal transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance in step 2 and step 3.

In addition, it was found from Example 6 and Example 7 that efficient differentiation into retinal cell is also achieved by the addition of a Wnt signal transduction pathway inhibiting substance step 2 and step 3 when step 1 includes Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance or Precondition with a BMPR inhibitor and a Sonic hedgehog signal transduction pathway activating substance.

[0318]

Example 8: Production Example of retinal tissue using human iPS cell established using Sendaivirus vector as starting material and including Precondition in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2

Human iPS cells (TFH-R1-10-2 strain and TFH-R2-10-F8 strain, established by Sumitomo Dainippon Pharma Co., Ltd.) were established as follows. They were established according to the method described in the published protocol of Life Technologies (iPS 2.0 Sendai Reprogramming Kit, Publication Number MAN0009378, Revision 1.0) and the published protocol of

Kyoto University (establish maintenance culture of human iPS cells, CiRA_Ff-iPSC_protocol_JP_v140310, <http://www.cira.kyoto-u.ac.jp/j/research/protocol.html>), and using peripheral blood mononuclear cells (PBMC) prepared by a well-known method as the starting material, commercially available Sendaivirus vector (4 factors of Oct3/4, Sox2, KLF4, L-Myc, CytoTune kit manufactured by DNAVEC Corporation (currently ID Pharma Co., Ltd.)), and StemFit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.), Laminin511-E8 (manufactured by Nippi, Inc.).

Human iPS cells (TFH-R1-10-2 strain and TFH-R2-10-F8 strain) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 8 "Precondition: TGF β R-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium

which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium, IWR-1e (3 μ M) was added as a Wnt signal transduction pathway inhibiting substance (Fig. 10 "Wnt-i (day 0-17)"), and SAG (30 nM) was added as a Sonic hedgehog signal transduction pathway activating substance (Fig. 10 "Shh day 0") and the cells were cultured. A cell aggregate was formed by day 2 after the start of suspension culture by using human iPS cells of any of TFH-R1-10-2 strain and TFH-R2-10-F8 strain as the starting material (step 2 completed, and step 3 started).

15 [0319]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of IWR-1e.

25

The thus-prepared cells on day 17 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue could be formed by using human iPS cells of any of TFH-R1-10-2 strain and TFH-R2-10-F8 strain as the starting material.

30

[0320]

Cell aggregates on day 17 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were

35

immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These stained sections were observed with an
5 inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 10).

As a result, it was found that a neural tissue could be formed by using human iPS cells of any of TFH-R1-10-2 strain and TFH-R2-10-F8 strain as the starting material. It was also
10 found that the proportion of the Chx10 positive retinal tissue is about 70% when TFH-R1-10-2 strain is the starting material (Fig. 10A). From the analysis of serial sections, it could be confirmed that these Chx10 positive cells were Rx co-positive cells (Fig. 10B). It was found that the proportion of the
15 Chx10 positive retinal tissue is about 90% when TFH-R2-10-F8 strain is the starting material (Fig. 10C). From the analysis of serial sections, it could be confirmed that these Chx10 positive cells were Rx co-positive cells (Fig. 10D).

[0321]

20 From these results, it was found that human iPS cells established using Sendaivirus vector as the starting material also efficiently differentiate into a retinal tissue under conditions including a Precondition treatment in step 1 and addition of a Wnt signal transduction pathway inhibiting
25 substance in step 2 and step 3.

Therefore, it was found that a retinal cell and/or a retinal tissue can be produced by the production method of the present application from iPS cells established using an episomal vector (e.g., 1231A3 strain) and human iPS cell
30 established using a Sendaivirus vector (e.g., TFH-R1-10-2 strain and TFH-R2-10-F8 strain) irrespective of the establishing method of the iPS cells.

[0322]

Example 9: Production Example of retinal cell containing
35 photoreceptor cell from human iPS cell including Precondition

with TGF β signal transduction pathway inhibiting substance and
Sonic hedgehog signal transduction pathway activating substance
in step 1 and use of Wnt signal transduction pathway inhibiting
substance and TGF β signal transduction pathway inhibiting
5 substance in step 2 and step 3

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a
10 feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal
15 transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

20 The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l
25 of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium
30 which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final
35 concentration 20 μ M) was added to the above-mentioned serum-

free medium, IWR-1e (3 μ M) was added as a Wnt signal transduction pathway inhibiting substance, and SAG (30 nM) was added as a Sonic hedgehog signal transduction pathway activating substance, and the cells were cultured. A cell
5 aggregate was formed by day 2 after the start of suspension culture.

[0323]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4
10 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4
15 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of IWR-1e.

The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation
20 using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed.

A part of the cells on day 20 after the start of the aforementioned suspension culture was extracted, fixed with 4%
25 para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These stained sections were observed
30 with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO). As a result, it was found that the cells on day 20 after the start of the aforementioned suspension culture contained a neural tissue and the neural tissue was a Chx10 and Rx co-positive retinal tissue.

35 [0324]

A part of the cells on day 20 after the start of the aforementioned suspension culture was extracted and subjected to a differentiation culture according to the method described in "Nature Communications 6, 6286 (2015)".

5 The cell aggregate on day 20 after the start of suspension culture were transferred to a 90 mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) and cultured in a serum-free medium (DMEM/F12 medium added with 1% N2 supplement) containing a Wnt signal transduction pathway
10 activating substance (CHIR99021, 3 μ M) and an FGF signal transduction pathway inhibiting substance (SU5402, 5 μ M) at 37°C, 5% CO₂ for 3 days, i.e., up to day 23 after the start of suspension culture. During this period, about 50 aggregates were cultured in suspension in a 10 ml serum-free medium
15 containing the aforementioned CHIR99021 and SU5402 per one 90-mm low adhesive culture dish. On day 23 after the start of suspension culture, a thin neuroepithelium was formed, and a retinal pigment epithelial (RPE)-like tissue was formed.
[0325]

20 The cell aggregate on day 23 after the start of suspension culture were cultured in suspension in a 90-mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) in a serum-containing medium (DMEM/F12 medium added with 10% fetal calf serum, 1% N2 supplement, 0.5 μ M retinoic acid,
25 and 100 μ M taurine) not containing a Wnt signal transduction pathway activating substance or an FGF signal transduction pathway inhibiting substance at 37°C, 5% CO₂, atmospheric oxygen concentration (about 20%) up to day 58 after the start of suspension culture (35 days). From day 20 after the start
30 of suspension culture to the completion of the suspension culture, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-containing medium. During this period, about 30 aggregates per one 90-mm low adhesive culture dish were cultured in suspension in 15 ml
35 of the aforementioned serum-containing medium. A neural

retina-like tissue was present from day 35 after the start of suspension culture.

[0326]

The cell aggregates on day 58 after the start of
5 suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that dye deposited retinal pigment epithelial cells could be formed in addition to the neural tissues (Fig. 11, A, shown with an arrowhead).

10 [0327]

The thus-prepared cell aggregates on day 58 after the start of suspension culture were each fixed with 4% para-formaldehyde, and cryosections were prepared. These cryosections were immunostained for Rx (anti-Rax/Rx antibody,
15 manufactured by Takara, Guinea Pig.), which is one of the retinal tissue markers, Chx10 (anti-Chx10 antibody, manufactured by Exalpha, sheep), which is one of the neural retinal progenitor cell markers, Crx (anti-Crx antibody, manufactured by Takara, rabbit), which is one of the
20 photoreceptor precursor cell markers, and observed using an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO).

[0328]

As a result, it was found that about 90% of the Rx
25 positive retinal tissues were formed in the cell aggregate on day 58 after the start of the aforementioned suspension culture (Fig. 11, B). It was further found that the Rx positive retinal tissues contain Chx10 positive neural retinal progenitor cells (Fig. 11, C). Also, it was found that the Rx
30 positive retinal tissues contain Crx positive photoreceptor precursor cells (Fig. 11, D). From morphological observation, it was found that the Rx positive retinal tissues contain a ciliary marginal zone-like structure (Fig. 11, E, shown with an arrow). From morphological observation, it could be confirmed
35 that Rx positive and Chx10 negative and Crx negative inner

layer retina nerve cells (e.g., ganglion cell and amacrine cell) are contained in the inside of the retinal tissue (Fig. 11).

[0329]

5 From these results, it was found that feeder-free cultured human iPS cells as the starting material efficiently differentiate into a retinal tissue under conditions including a Precondition treatment in step 1 and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3.
10 It was further found that retinal cells (or retina layer specific nerve cells), for example, neural retinal progenitor cells, photoreceptor precursor cells, retinal pigment epithelial cells, inner layer retina nerve cells, ciliary marginal zone-like structure can be produced by continuing
15 differentiation culture of the prepared retinal tissues. In addition, it was found that neural retinal progenitor cells, photoreceptor precursor cells, inner layer retina nerve cells form a continuous epithelial structure having a layer structure in the aforementioned retinal tissues.

20 That is, it was found that retinal tissues .retinal cells useful for regenerative medicine .cell transplantation treatment and evaluation study of effectiveness .safety can be produced from feeder-free cultured human pluripotent stem cells by the production method of the present application.

25 [0330]

Example 10: Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of
30 Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade
35 mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a

feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under
5 conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 12 "Precondition: TGF β R-i+Shh") in a Stem Fit
10 (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed
15 into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the
20 cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of
25 the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under three conditions of the following Conditions 1 - 3.
30 •Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 12A-C, "Condition 1").
35 •Condition 2. Condition including addition of IWR-1e (3 μ M) as

a Wnt signal transduction pathway inhibiting substance and addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 12D-F "Condition 2").

•Condition 3. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 12G-I "Condition 3").

A cell aggregate was formed by day 2 after the start of suspension culture under Condition 1, Condition 2 and Condition 3 (step 2 completed, and step 3 started).

[0331]

On day 3 after the start of suspension culture, a medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with medium the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 12 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 20 after the start of

suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition 1, Condition 2 and Condition 3.

5 [0332]

The aggregates on day 20 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep),
10 which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers, or Pax6, which is one of the markers of neural tissues including a retinal tissue. These stained sections were observed with an inverted fluorescence microscope
15 (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 12).
[0333]

As a result, it was found that a neural tissue could be formed in Condition 1, Condition 2 and Condition 3. In Condition 1, the proportion of the Chx10 positive retinal
20 tissue was about 95% (Fig. 12A), in Condition 2, the proportion of the Chx10 positive retinal tissue was about 95% (Fig. 12D), and in Condition 3, the proportion of the Chx10 positive retinal tissue was about 85% (Fig. 12G). From the analysis of serial sections, it could be confirmed that these Chx10
25 positive cells are Rx and Pax6 co-positive cells.

That is, in Condition 1, it was found that efficient differentiation into a retinal tissue is also achieved under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog
30 signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (on
35 day 12 after the start of suspension culture). Furthermore, in

Condition 2 and Condition 3, it was found that efficient differentiation into retinal cell is also achieved under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3. [0334]

10 Example 11: Production Example of retinal tissue from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

15 Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (Shh) treatment, Fig. 13 "Precondition: Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.) (that is, a Sonic hedgehog signal transduction pathway activating substance was allowed to act for two days before start of differentiation).

30 The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-

35

adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is
5 a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 µM 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 µM) was added
10 to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under three conditions of the following Conditions 1 - 3.

•Condition 1. Condition including addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and no
15 addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 13A-C, "Condition 1").

•Condition 2. Condition including addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and
20 addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 13D-F "Condition 2").

•Condition 3. Condition including addition of IWR-1e (3 µM) as
25 a Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 13G-I "Condition 3").

30 A cell aggregate was formed by day 2 after the start of suspension culture under Condition 1, Condition 2 and Condition 3 (step 2 completed, and step 3 started).

[0335]

On day 3 after the start of suspension culture, a medium
35 medium not containing Y27632 or SAG, containing human

recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change.

5 Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 10 after the start of suspension culture, a 80%
10 medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned
15 serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation,
20 ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition 1, Condition 2 and Condition 3.
[0336]

The cell aggregates on day 20 after the start of the aforementioned suspension culture were fixed with 4% para-
25 formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers, or Pax6, which is one of the markers of neural
30 tissues including a retinal tissue. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 13).
[0337]

As a result, it was found that a neural tissue was formed
35 in Condition 1, Condition 2 and Condition 3. In Condition 1,

the proportion of the Chx10 positive retinal tissue was about 95%, in Condition 2, the proportion of the Chx10 positive retinal tissue was about 90%, and in Condition 3, the proportion of the Chx10 positive retinal tissue was about 70%.

5 From the analysis of serial sections, it could be confirmed that these Chx10 positive cells are Rx and Pax6 co-positive cells.

That is, in Condition 1, it was found that efficient differentiation into a retinal tissue is also achieved under
10 conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance
15 to 3% or less in the course of step 3 (on day 10 after the start of suspension culture). Furthermore, in Condition 2 and Condition 3, it was found that efficient differentiation into retinal cell is also achieved under conditions including Precondition with a Sonic hedgehog signal transduction pathway
20 activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3.

[0338]

25 Example 12: Production Example of retinal tissue containing photoreceptor cell from human iPS cell including Precondition with TGF β signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting
30 substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a
35 feeder-free medium and Laminin 511-E8 (manufactured by Nippi,

Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium of the following Condition 1.

•Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2.

A cell aggregate was formed by day 2 after the start of suspension culture under Condition 1, Condition 2 and Condition 3 (step 2 completed, and step 3 started).

[0339]

On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of
5 exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with medium the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further
10 containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of
15 exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

20 The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed.

25 [0340]

A part of the cells on day 20 after the start of the aforementioned suspension culture was extracted and subjected to a differentiation culture according to the method described in "Nature Communications 6, 6286 (2015)".

30 The cell aggregate on day 20 after the start of suspension culture were transferred to a 90 mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) and cultured in a serum-free medium (DMEM/F12 medium added with 1% N2 supplement) containing a Wnt signal transduction pathway
35 activating substance (CHIR99021, 3 μ M) and an FGF signal

transduction pathway inhibiting substance (SU5402, 5 μ M) at 37°C, 5% CO₂ for 3 days, i.e., up to day 23 after the start of suspension culture. During this period, about 50 aggregates were cultured in suspension in a 10 ml serum-free medium containing the aforementioned CHIR99021 and SU5402 per one 90-mm low adhesive culture dish. On day 23 after the start of suspension culture, a thin neuroepithelium was formed, and a retinal pigment epithelial (RPE)-like tissue was formed.

[0341]

10 The cell aggregate on day 23 after the start of suspension culture were cultured in suspension in a 90-mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) in a serum-containing medium (DMEM/F12 medium added with 10% fetal calf serum, 1% N2 supplement, 0.5 μ M retinoic acid, 15 and 100 μ M taurine) not containing a Wnt signal transduction pathway activating substance or an FGF signal transduction pathway inhibiting substance at 37°C, 5% CO₂, atmospheric oxygen concentration (about 20%) up to day 85 after the start of suspension culture (62 days). From day 23 after the start of suspension culture to the completion of the suspension culture, once every 2 - 4 days, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-containing medium. During this period, about 30 aggregates per one 90-mm low adhesive culture dish were cultured in suspension in 15 ml of the aforementioned serum-containing medium. A neural retina-like tissue was present from day 35 after the start of suspension culture.

20 [0342]

 The cell aggregates on day 85 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue having a continuous epithelial structure was formed (Fig. 14A).

30 [0343]

35 The thus-prepared cell aggregates on day 85 after the

start of suspension culture were each fixed with 4% para-
formaldehyde, and cryosections were prepared. These
cryosections were immunostained for Crx (anti-Crx antibody,
manufactured by Takara, rabbit), which is one of the
5 photoreceptor precursor cell markers, Chx10 (anti-Chx10
antibody, manufactured by Exalpha, sheep), which is one of the
neural retinal progenitor cell markers, Rx (anti-Rax/Rx
antibody, manufactured by Takara, Guinea Pig.), which is one of
the retinal tissue markers, Recoverin (anti-Recoverin antibody,
10 manufactured by Proteintech, rabbit), which is one of the
photoreceptor cell markers, NRL (anti-NRL antibody,
manufactured by R and D, goat), which is one of the rod
photoreceptor precursor cell markers, RXR-gamma (anti-RXRG
antibody, manufactured by R and D, goat), which is one of the
15 corn photoreceptor precursor cell markers, N-cadherin (anti-N-
cadherin antibody, manufactured by BD, mouse), which is one of
the neural tissue markers, Aqp1 (anti-Aqp1 antibody,
manufactured by Millipore, rabbit), which is one of the RPE and
ciliary margin markers, and observed using an inverted
20 fluorescence microscope (manufactured by KEYENCE CORPORATION,
BIOREVO). As counterstaining, nucleic acid was stained with
DAPI.

[0344]

As a result, it was found that about 90% of retinal
25 tissues containing Chx10 positive neural retinal progenitor
cells and Crx positive photoreceptor precursor cells were
formed in the aforementioned cell aggregates on day 85 after
the start of suspension culture (Fig. 14, B). From the
analysis of serial sections, it was found that the retinal
30 tissue was an Rx positive retinal tissue and contained
Recoverin positive photoreceptor cell (Fig. 14, C). From the
analysis of serial sections, it was found that the retinal
tissue contained NRL positive rod photoreceptor precursor cells
and RXR-gamma positive corn photoreceptor precursor cells (Fig.
35 14, D). Also, it was found that the retinal tissue was an N-

cadherin positive neural tissue (Fig. 14E).

[0345]

From these results, it was found that feeder-free cultured human iPS cells as the starting material efficiently
5 differentiate into a retinal tissue under conditions including a Precondition treatment with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1 and addition of a Wnt signal transduction pathway inhibiting
10 substance in step 2 and step 3. It was further found that retinal cells (or retina specific nerve cells), for example, neural retinal progenitor cell, photoreceptor precursor cell, photoreceptor cell, corn photoreceptor precursor cell, rod photoreceptor precursor cell can be produced by continuing
15 differentiation culture of the prepared retinal tissues. In addition, it was found that neural retinal progenitor cells, photoreceptor precursor cells, photoreceptor cell form a continuous epithelial structure having a layer structure in the aforementioned retinal tissues.

20 [0346]

Example 13: Production Example of retinal tissue containing photoreceptor cell from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway
25 inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a
30 feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal
35 transduction pathway activating substance (Shh), 300 nM) (step

1: Precondition (Shh) treatment, Fig. 15) in a Stem Fit
(registered trade mark) medium (AK03N; manufactured by
Ajinomoto Co., Inc.) (that is, Sonic hedgehog signal
transduction pathway activating substance was allowed to act
5 for two days before start of differentiation).

The thus-prepared Precondition (Shh)-treated human iPS
cells were treated with the cell dispersion solution by using
TrypLE Select (Life Technologies), further dispersed into
single cells by pipetting operation, and the above human iPS
10 cells dispersed into single cells were suspended in 100 μ l of a
serum-free medium at 1.0×10^4 cells per well of a non-cell-
adhesive 96-well culture plate (PrimeSurface 96V-bottom plate,
manufactured by SUMITOMO BAKELITE). Thereafter, the cells were
subjected to suspension culture at 37°C, 5% CO₂. As the serum-
15 free medium (gfCDM+KSR) therefor, a serum-free medium which is
a 1:1 mixture of F-12 medium and IMDM medium supplemented with
10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined
lipid concentrate was used. At the time of the start of
suspension culture (day 0 after the start of suspension culture,
20 start of step 2), Y27632 (final concentration 20 μ M) was added
to the above-mentioned serum-free medium and the cells were
cultured in a serum-free medium under three conditions of the
following Conditions.

.Condition 1. Condition including addition of IWR-1e (3 μ M) as
25 a Wnt signal transduction pathway inhibiting substance and
addition of SAG (30 nM) as an exogenous Sonic hedgehog signal
transduction pathway activating substance to the aforementioned
serum-free medium at the start of step 2.

A cell aggregate was formed by day 2 after the start of
30 suspension culture under Condition 1, Condition 2 and Condition
3 (step 2 completed, and step 3 started).

[0347]

On day 3 after the start of suspension culture, a medium
medium not containing Y27632 or SAG, containing human
35 recombinant BMP4 (manufactured by R&D) and further containing

IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once
5 every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 9 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the
10 aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant
15 BMP4 and IWR-1e.

The thus-prepared cells on day 19 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue
20 was formed.

[0348]

A part of the cells on day 19 after the start of the aforementioned suspension culture was extracted and subjected to a differentiation culture according to the method described
25 in "Nature Communications 6, 6286 (2015)".

The cell aggregate on day 19 after the start of suspension culture were transferred to a 90 mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) and cultured in a serum-free medium (DMEM/F12 medium added with 1%
30 N2 supplement) containing a Wnt signal transduction pathway activating substance (CHIR99021, 3 μ M) and an FGF signal transduction pathway inhibiting substance (SU5402, 5 μ M) at 37°C, 5% CO₂ for 3 days, i.e., up to day 22 after the start of suspension culture. During this period, about 50 aggregates
35 were cultured in suspension in 10 ml of a serum-free medium

containing the aforementioned CHIR99021 and SU5402 per one 90-mm low adhesive culture dish. On day 22 after the start of suspension culture, a thin neuroepithelium was formed, and a retinal pigment epithelial (RPE)-like tissue was formed.

5 [0349]

The cell aggregate on day 22 after the start of suspension culture were cultured in suspension in a 90-mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) in a serum-containing medium (DMEM/F12 medium added with 10% fetal calf serum, 1% N2 supplement, 0.5 μ M retinoic acid, and 100 μ M taurine) not containing a Wnt signal transduction pathway activating substance or an FGF signal transduction pathway inhibiting substance at 37°C, 5% CO₂, atmospheric oxygen concentration (about 20%) up to day 92 after the start of suspension culture (70 days). From day 20 after the start of suspension culture to the completion of the suspension culture, once every 2 - 4 days, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-containing medium. During this period, about 30 aggregates per one 90-mm low adhesive culture dish were cultured in suspension in 15 ml of the aforementioned serum-containing medium. A neural retina-like tissue was present from day 35 after the start of suspension culture.

[0350]

25 The cell aggregates on day 92 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue having a continuous epithelial structure was formed (Fig. 15A).

30 [0351]

The thus-prepared cell aggregates on day 92 after the start of suspension culture were each fixed with 4% para-formaldehyde, and cryosections were prepared. These cryosections were immunostained for Crx (anti-Crx antibody, manufactured by Takara, rabbit), which is one of the

photoreceptor precursor cell markers, Chx10 (anti-Chx10 antibody, manufactured by Exalpha, sheep), which is one of the neural retinal progenitor cell markers, Rx (anti-Rax/Rx antibody, manufactured by Takara, Guinea Pig.), which is one of the retinal tissue markers, Recoverin (anti-Recoverin antibody, manufactured by Proteintech, rabbit), which is one of the photoreceptor cell markers, NRL (anti-NRL antibody, manufactured by R and D, goat), which is one of the rod photoreceptor precursor cell markers, RXR-gamma (anti-RXRG antibody, manufactured by R and D, goat), which is one of the corn photoreceptor precursor cell markers, N-cadherin (anti-N-cadherin antibody, manufactured by BD, mouse), which is one of the neural tissue markers, Aqp1 (anti-Aqp1 antibody, manufactured by Millipore, rabbit), which is one of the RPE and ciliary margin markers, and observed using an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO). As counterstaining, nucleic acid was stained with DAPI.

[0352]

As a result, it was found that about 80% of retinal tissues containing Chx10 positive neural retinal progenitor cells and Crx positive photoreceptor precursor cells were formed in the aforementioned cell aggregates on day 92 after the start of suspension culture (Fig. 15, B). From the analysis of serial sections, it was found that the retinal tissue was an Rx positive retinal tissue and contained Recoverin positive photoreceptor cell (Fig. 15, C). From the analysis of serial sections, it was found that the retinal tissue contained NRL positive rod photoreceptor precursor cells and RXR-gamma positive corn photoreceptor precursor cells (Fig. 15, D). Also, from the analysis of serial sections, it was found that the retinal tissue was an N-cadherin positive neural tissue (Fig. 15, E). From the analysis of serial sections, it was found that the retinal tissue contained Aqp1 positive ciliary margin-like structure (Fig. 15, E).

[0353]

From these results, it was found that feeder-free cultured human iPS cells as the starting material efficiently differentiate into a retinal tissue under conditions including a Precondition treatment with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3. It was further found that retinal cells (or retina specific nerve cells), for example, neural retinal progenitor cell, photoreceptor precursor cell, photoreceptor cell, cone photoreceptor precursor cell, rod photoreceptor precursor cell, ciliary margin-like structure can be produced by continuing differentiation culture of the prepared retinal tissues. In addition, it was found that neural retinal progenitor cells, photoreceptor precursor cells, photoreceptor cell form a continuous epithelial structure having a layer structure in the aforementioned retinal tissues.

[0354]

Example 14: Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trademark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal

transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 16 "Precondition: TGF β R-i+Shh") in a Stem Fit
5 (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed
10 into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the
15 cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 6% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of
20 the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under the following Condition.

25 •Condition. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (30 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 16, A, B).

30 A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).
[0355]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions A, B.

35 •Condition A. On day 3 after the start of suspension culture,

a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
5 the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

10 .Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
15 the concentration of exogenous IWR-1e would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange operation was performed using the aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and
20 IWR-1e such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further
25 containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

[0356]

After culturing under two conditions of the above-mentioned Conditions A, B, the cells were further cultured as
30 follows.

On day 12 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before
35 medium exchange. Thereafter, a half-medium exchange operation

was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 18 after the start of
5 suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition A and Condition B.

[0357]

10 The aggregates on day 18 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers. These stained
15 sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 16A, B).

[0358]

As a result, it was found that a neural tissue was formed under Condition A and Condition B. In Condition A, the
20 proportion of the Chx10 positive retinal tissue was about 80%, and in Condition B, the proportion of the Chx10 positive retinal tissue was about 80%.

That is, it was found that efficient differentiation into a retinal tissue is also achieved, when the addition condition
25 of exogenous BMP is any condition of Condition A and Condition B, under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway
30 activating substance in step 2, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On day 12 after the start of suspension
35 culture), and the concentration of KSR contained in the

aforementioned serum-free medium of 6%.

[0359]

Example 15: Production Example of retinal tissue from human iPS cell including Precondition with Sonic hedgehog signal

5 transduction pathway activating substance in step 1 and use of
Wnt signal transduction pathway inhibiting substance in step 2
and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method
10 described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before
15 subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (+Shh) treatment, Fig. 16 "Precondition: Shh") in a Stem Fit (registered trade mark) medium (AK03N;
20 manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS
25 cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is
30 a 1:1 mixture of F-12 medium and IMDM medium supplemented with 6% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of
35 step 2), Y27632 (final concentration 20 μ M) was added to the

above-mentioned serum-free medium and the cells were cultured in a serum-free medium under the following Condition.

.Condition. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no
5 addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 16, C, D).

A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).

10 [0360]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions C and D.

.Condition C. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human
15 recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once
20 every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

.Condition D. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human
25 recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e would not change.

Further, On day 6 after the start of suspension culture, a
30 half-medium exchange operation was performed using the aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and IWR-1e such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e
35 would not change. Thereafter, a half-medium exchange operation

was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

5 [0361]

After culturing under two conditions of the above-mentioned Conditions C and D, the cells were cultured as follows.

On day 12 after the start of suspension culture, a 80%
10 medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned
15 serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 18 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation,
20 ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition C and Condition D.

[0362]

The aggregates on day 18 after the start of the aforementioned suspension culture were fixed with 4% para-
25 formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 16C, D).

30 [0363]

As a result, it was found that a neural tissue was formed under Condition C and Condition D. In Condition C, the proportion of the Chx10 positive retinal tissue was about 80%, and in Condition D, the proportion of the Chx10 positive
35 retinal tissue was about 80%.

That is, it was found that efficient differentiation into a retinal tissue is also achieved, when the addition condition of exogenous BMP is any condition of Condition C and Condition D, under conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On day 12 after the start of suspension culture), and the concentration of KSR contained in the aforementioned serum-free medium of 6%.

[0364]

Example 16: Production Example of retinal tissue from human iPS cell including Precondition with BMP signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of LDN193189 (BMP signal transduction pathway inhibiting substance (BMPR-i), 100 nM) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (BMPR-i+Shh) treatment, Fig. 17 "Precondition: BMPR-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (BMPR-i+Shh)-treated human

iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under the following Condition.

.Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 17, A, B).

A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).

[0365]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions A, B.

.Condition A. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-

le to avoid change of the concentration of exogenous IWR-1e.

•Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing
5 IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange operation was performed using the
10 aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and IWR-1e such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation
15 was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

[0366]

20 After culturing under two conditions of the above-mentioned Conditions A, B, the cells were cultured as follows.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of
25 exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

30 The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition A and Condition B.

35 [0367]

The aggregates on day 20 after the start of the
aforementioned suspension culture were fixed with 4% para-
formaldehyde to produce cryosections. These cryosections were
immunostained for Chx10 (anti-Chx10 antibody, Exalpa, sheep),
5 which is one of the retinal tissue markers. These stained
sections were observed with an inverted fluorescence microscope
(manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 17A, B).
[0368]

As a result, it was found that a neural tissue was formed
10 under Condition A and Condition B. In Condition A, the
proportion of the Chx10 positive retinal tissue was about 90%,
and in Condition B, the proportion of the Chx10 positive
retinal tissue was about 90%.

That is, it was found that efficient differentiation into
15 a retinal tissue is also achieved, when the addition condition
of exogenous BMP is any condition of Condition A and Condition
B, under conditions including Precondition with a BMP signal
transduction pathway inhibiting substance and a Sonic hedgehog
signal transduction pathway activating substance in step 1,
20 addition of a Wnt signal transduction pathway inhibiting
substance in step 2 and step 3, and reduction of the
concentration of the exogenous Wnt signal transduction pathway
inhibiting substance to 3% or less in the course of step 3 (On
day 10 after the start of suspension culture).

25 [0369]

Example 17: Production Example of retinal tissue from human iPS
cell including Precondition with TGF β signal transduction
pathway inhibiting substance and Sonic hedgehog signal
transduction pathway activating substance in step 1 and use of
30 Wnt signal transduction pathway inhibiting substance in step 2
and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto
University) were cultured feeder free according to the method
described in Example 1 and using StemFit (registered trade
35 mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a

feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for one day under
5 conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.), whereby human iPS cells one day before subconfluence were prepared. Furthermore, human iPS cells one
10 day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) in a Stem Fit (registered trade mark)
15 medium (AK03N; manufactured by Ajinomoto Co., Inc.) (step 1: Precondition (TGF β R-i 24h+Shh 48h) treatment, Fig. 18).

The thus-prepared Precondition (TGF β R-i 24h+Shh 48h)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further
20 dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO
25 BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was
30 used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under the following Condition.

35 •Condition. Condition including addition of IWR-1e (3 μ M) as a

Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 18, A, B).

5 A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).
[0370]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions A, B.

10 .Condition A. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
15 the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

20 .Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
25 the concentration (3 μ M) of exogenous IWR-1e would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange operation was performed using the aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and
30 IWR-1e such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further
35 containing IWR-1e to avoid change of the concentration of

exogenous IWR-1e.

[0371]

After culturing under two conditions of the above-mentioned Conditions A, B, the cells were further cultured as follows.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition A and Condition B.

[0372]

The aggregates on day 20 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 18A, B).

[0373]

As a result, it was found that a neural tissue was formed under Condition A and Condition B. In Condition A, the proportion of the Chx10 positive retinal tissue was about 80%, and in Condition B, the proportion of the Chx10 positive retinal tissue was about 80%.

That is, it was found that efficient differentiation into a retinal tissue is also achieved, when the addition condition of exogenous BMP is any condition of Condition A and Condition

B, under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On day 10 after the start of suspension culture).

[0374]

10 Example 18:: Production Example of retinal tissue from human iPS cell including Precondition with BMP signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2
15 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for one day under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.), whereby human iPS cells one day before subconfluence were prepared. Furthermore, human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of LDN193189 (BMP signal transduction pathway inhibiting substance (BMPR-i), 100 nM) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.) (step 1: Precondition (BMPR-i 24h+Shh 48h) treatment, Fig. 19).

The thus-prepared Precondition (BMPR-i 24h+Shh 48h)-
treated human iPS cells were treated with the cell dispersion
solution by using TrypLE Select (Life Technologies), further
dispersed into single cells by pipetting operation, and the
5 above human iPS cells dispersed into single cells were
suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells
per well of a non-cell-adhesive 96-well culture plate
(PrimeSurface 96V-bottom plate, manufactured by SUMITOMO
BAKELITE). Thereafter, the cells were subjected to suspension
10 culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR)
therefor, a serum-free medium which is a 1:1 mixture of F-12
medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-
monothioglycerol, 1 x Chemically defined lipid concentrate was
used. At the time of the start of suspension culture (day 0
15 after the start of suspension culture, start of step 2), Y27632
(final concentration 20 μ M) was added to the above-mentioned
serum-free medium and the cells were cultured in the serum-free
medium under the following Condition.

.Condition. Condition including addition of IWR-1e (3 μ M) as a
20 Wnt signal transduction pathway inhibiting substance and no
addition of an exogenous Sonic hedgehog signal transduction
pathway activating substance to the aforementioned serum-free
medium at the start of step 2 (Fig. 19, A, B).

A cell aggregate was formed by day 2 after the start of
25 suspension culture (step 2 completed, and step 3 started).

[0375]

Furthermore, the cells were cultured in a serum-free
medium under two conditions of the following Conditions A, B.

.Condition A. On day 3 after the start of suspension culture,
30 a medium medium not containing Y27632 or SAG, containing human
recombinant BMP4 (manufactured by R&D) and further containing
IWR-1e was added by 50 μ l such that a final concentration of
exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
the concentration (3 μ M) of exogenous IWR-1e would not change.
35 Thereafter, a half-medium exchange operation was performed once

every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

•Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange operation was performed using the aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and IWR-1e such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

[0376]

The cells were cultured as follows under two conditions of the above-mentioned Conditions A, B.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue

was formed under Condition A and Condition B.

[0377]

The aggregates on day 20 after the start of the
aforementioned suspension culture were fixed with 4% para-
5 formaldehyde to produce cryosections. These cryosections were
immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep),
which is one of the retinal tissue markers. These stained
sections were observed with an inverted fluorescence microscope
(manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 19A, B).

10 [0378]

As a result, it was found that a neural tissue was formed
under Condition A and Condition B. In Condition A, the
proportion of the Chx10 positive retinal tissue was about 50%,
and in Condition B, the proportion of the Chx10 positive
15 retinal tissue was about 70%.

That is, it was found that differentiation into a retinal
tissue is also achieved, when the addition condition of
exogenous BMP is any condition of Condition A and Condition B,
under conditions including Precondition with a BMP signal
20 transduction pathway inhibiting substance and a Sonic hedgehog
signal transduction pathway activating substance in step 1,
addition of a Wnt signal transduction pathway inhibiting
substance in step 2 and step 3, and reduction of the
concentration of the exogenous Wnt signal transduction pathway
25 inhibiting substance to 3% or less in the course of step 3 (On
day 10 after the start of suspension culture).

[0379]

Example 19: Production Example of retinal tissue from human iPS
cell including Precondition with TGF β signal transduction
30 pathway inhibiting substance and Sonic hedgehog signal
transduction pathway activating substance in step 1 and use of
Wnt signal transduction pathway inhibiting substance and TGF β
signal transduction pathway inhibiting substance in step 2 and
step 3

35 Human iPS cells (QHJI01s04 strain, obtained from Kyoto

University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 20 "Precondition: TGF β R-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium, and the cells were cultured in the serum-free medium under following Condition.

.Condition. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SB431542 (5 μ M) as TGF β signal transduction pathway inhibiting

substance, and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 20, A, B).

5 A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).
[0380]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions A, B.

10 •Condition A. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e and SB431542 was added by 50 μ l such that a final
15 concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentrations of exogenous IWR-1e (3 μ M) and SB431542 (5 μ M) would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with medium the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e and
20 SB431542 to avoid change of the concentration of exogenous IWR-1e and SB431542.

•Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing
25 IWR-1e and SB431542 was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentrations of exogenous IWR-1e (3 μ M) and SB431542 (5 μ M) would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange
30 operation was performed using the aforementioned serum-free medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and IWR-1e and SB431542 such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e and SB431542 would
35 not change. Thereafter, a half-medium exchange operation was

performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e and SB431542.

5 [0381]

The cells were further cultured as follows after culturing under two conditions of the above-mentioned Conditions A, B.

On day 10 after the start of suspension culture, a 80%
10 medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e and SB431542 would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the
15 aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4, IWR-1e and SB431542.

The thus-prepared cells on day 20 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation,
20 ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition A and Condition B.

[0382]

The aggregates on day 20 after the start of the aforementioned suspension culture were fixed with 4% para-
25 formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 20A, B).

30 [0383]

As a result, it was found that a neural tissue was formed under Condition A and Condition B. In Condition A, the proportion of the Chx10 positive retinal tissue was about 95%, and in Condition B, the proportion of the Chx10 positive
35 retinal tissue was about 95%.

That is, it was found that efficient differentiation into a retinal tissue is also achieved, when the addition condition of exogenous BMP is any condition of Condition A and Condition B, under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On day 10 after the start of suspension culture).

[0384]

Example 20: Production Example of retinal tissue from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance and TGF β signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (Shh) treatment, Fig. 21 "Precondition: Shh 48 h") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using

TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.0×10^4 cells per well of a non-cell-
5 adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with
10 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium, and the cells were
15 cultured in the serum-free medium. A cell aggregate was formed by day 2 after the start of suspension culture.

.Condition. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and SB431542 (5 μ M) as a TGF β signal transduction pathway
20 inhibiting substance, and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 21, A, B).

A cell aggregate was formed by day 2 after the start of
25 suspension culture (step 2 completed, and step 3 started).
[0385]

Furthermore, the cells were cultured in a serum-free medium under two conditions of the following Conditions A, B.
.Condition A. On day 3 after the start of suspension culture,
30 a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e and SB431542 was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentrations of exogenous IWR-1e (3 μ M)
35 and SB431542 (5 μ M) would not change. Thereafter, a half-

medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e and SB431542 to avoid change of the concentration of exogenous IWR-1e and
5 SB431542.

•Condition B. On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e and SB431542 was added by 50 μ l such that a final
10 concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentrations of exogenous IWR-1e (3 μ M) and SB431542 (5 μ M) would not change. Further, On day 6 after the start of suspension culture, a half-medium exchange operation was performed using the aforementioned serum-free
15 medium not containing Y27632 or SAG and containing human recombinant BMP4 (manufactured by R&D) and IWR-1e and SB431542 such that the concentration of exogenous human recombinant BMP4 and the concentration of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once
20 every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e and SB431542.

[0386]

25 The cells cultured under two conditions of the above-mentioned Conditions A, B were cultured as follows.

On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of
30 exogenous IWR-1e and SB431542 would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4, IWR-1e and SB431542.

35 The thus-prepared cells on day 20 after the start of

suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition A and Condition B.

5 [0387]

The aggregates on day 20 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep),
10 which is one of the retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 21A, B).
[0388]

As a result, it was found that a neural tissue was formed
15 under Condition A and Condition B. In Condition A, the proportion of the Chx10 positive retinal tissue was about 80%, and in Condition B, the proportion of the Chx10 positive retinal tissue was about 80%.

That is, it was found that efficient differentiation into
20 a retinal tissue is also achieved, when the addition condition of exogenous BMP is any condition of Condition A and Condition B, under conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway
25 inhibiting substance and a TGF β signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance and a TGF β signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On
30 day 10 after the start of suspension culture).

[0389]

Example 21: Production Example of retinal tissue from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and use of
35 Wnt signal transduction pathway inhibiting substance in step 2

and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade
5 mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for two days under
10 conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (Shh) treatment, Fig. 22) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

15 The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a
20 serum-free medium at 1.3×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is
25 a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added
30 to the above-mentioned serum-free medium, and the cells were cultured in the serum-free medium under the following Condition 1 and Condition 2.

•Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no
35 addition of an exogenous Sonic hedgehog signal transduction

pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 22A, B).

•Condition 2. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and
5 addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 22C, D).

A cell aggregate was formed by day 2 after the start of suspension culture under Condition 1 and Condition 2 (step 2
10 completed, and step 3 started).

[0390]

On day 3 after the start of suspension culture, under Condition 1 and Condition 2, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured
15 by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change.

On day 6 after the start of suspension culture, a 80%
20 medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned
25 serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

[0391]

The thus-prepared cells on day 18 after the start of suspension culture were subjected to bright field observation
30 using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition 1 and Condition 2.

[0392]

The aggregates on day 18 after the start of the
35 aforementioned suspension culture were fixed with 4% para-

formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig), which is one of retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO).

[0393]

As a result, it was found that a neural tissue was formed under Condition 1 and Condition 2. In Condition 1, the proportion of the Chx10 positive retinal tissue was about 90% (Fig. 22A), and in Condition 2, the proportion of the Chx10 positive retinal tissue was about 80% (Fig. 22C). From the analysis of serial sections, it was found that the Chx10 positive cell in Condition 1 and Condition 2 is Rx co-positive cell (Fig. 22 B, D).

% or less after the start of suspension culture That is, it was found that efficient differentiation into a retinal tissue is also achieved, under conditions with or without addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, under conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, reduction of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (On day 6 after the start of suspension culture),

[0394]

Example 22: Production Example of retinal tissue containing photoreceptor precursor cell and ganglion cell from human iPS cell including Precondition with Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (QHJI01s04 strain, obtained from Kyoto

University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, 5 Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells two days before subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 10 1: Precondition (Shh) treatment, Fig. 23) in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using 15 TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 µl of a serum-free medium at 1.3×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, 20 manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 µM 1-monothioglycerol, 1 x Chemically defined 25 lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 µM) was added to the above-mentioned serum-free medium, and the cells were cultured in the serum-free medium under the following Condition 30 1 and Condition 2.

•Condition 1. Condition including addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free 35 medium at the start of step 2 (Fig. 23A-D).

.Condition 2. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 23E-H).

A cell aggregate was formed by day 2 after the start of suspension culture under Condition 1 and Condition 2 (step 2 completed, and step 3 started).

[0395]

On day 3 after the start of suspension culture, under Condition 1 and Condition 2, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration (3 μ M) of exogenous IWR-1e would not change.

On day 6 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

[0396]

The thus-prepared cells on day 11 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed under Condition 1 and Condition 2.

[0397]

A part of the aggregates on day 11 after the start of the aforementioned suspension culture was extracted and fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody,

Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig), which is one of retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by
5 KEYENCE CORPORATION, BIOREVO).

[0398]

As a result, it was found that a neural tissue was formed under Condition 1 and Condition 2. In Condition 1, the proportion of the Chx10 positive retinal tissue was about 90%
10 (Fig. 23A), and in Condition B, the proportion of the Chx10 positive retinal tissue was about 90% (Fig. 23E). From the analysis of serial sections, it was found that the Chx10 positive cell in Condition 1 and Condition 2 is Rx co-positive cell (Fig. 23 B, F).

15 [0399]

Under Condition 1 and Condition 2, a part of the cells on day 11 after the start of the aforementioned suspension culture was extracted and subjected to a differentiation culture according to the method described in "Nature Communications 6,
20 6286 (2015)".

The cell aggregate on day 11 after the start of suspension culture were transferred to a 90 mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) and cultured in a serum-free medium (DMEM/F12 medium added with 1%
25 N2 supplement) containing a Wnt signal transduction pathway activating substance (CHIR99021, 3 μ M) and an FGF signal transduction pathway inhibiting substance (SU5402, 5 μ M) at 37°C, 5% CO₂ for 3 days, i.e., up to day 14 after the start of suspension culture. During this period, about 50 aggregates
30 were cultured in suspension in a 10 ml serum-free medium containing the aforementioned CHIR99021 and SU5402 per one 90-mm low adhesive culture dish. On day 14 after the start of suspension culture, a thin neuroepithelium was formed, and a retinal pigment epithelial (RPE)-like tissue was formed.

35 [0400]

The cell aggregate on day 14 after the start of suspension culture were cultured in suspension in a 90-mm low adhesive culture dish (manufactured by SUMITOMO BAKELITE CO., LTD.) in a serum-containing medium (DMEM/F12 medium added with 10% fetal calf serum, 1% N2 supplement and 100 μ M taurine) not containing a Wnt signal transduction pathway activating substance or an FGF signal transduction pathway inhibiting substance at 37°C, 5% CO₂, atmospheric oxygen concentration (about 20%) up to day 35 after the start of suspension culture (21 days). From day 20 after the start of suspension culture to the completion of the suspension culture, once every 2 - 4 days, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-containing medium. During this period, about 30 aggregates per one 90-mm low adhesive culture dish were cultured in suspension in 15 ml of the aforementioned serum-containing medium.

[0401]

The cell aggregates on day 35 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural retina - like tissue was present and a neural tissue having a continuous epithelial structure can be formed.

[0402]

The thus-prepared cell aggregates on day 35 after the start of suspension culture were each fixed with 4% para-formaldehyde, and cryosections were prepared. These cryosections were immunostained for Crx (anti-Crx antibody, manufactured by Takara, rabbit), which is one of the photoreceptor precursor cell markers or Brn3b (anti-Brn3b antibody, manufactured by Santa Cruz, goat), which is one of the ganglion cell markers, and observed using an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO).

[0403]

As a result, it was found that about 90% of retinal tissues containing Crx positive photoreceptor precursor cells were formed in the aforementioned cell aggregates on day 35 after the start of suspension culture under Condition 1 and Condition 2 (Fig. 23 C, G). From the analysis of serial sections, it was found that the retinal tissue contained Brn3b positive ganglion cells (Fig. 23 D, H).

[0404]

From these results, it was found that feeder-free cultured human iPS cells as the starting material also efficiently differentiate into a retinal tissue under conditions with or without addition of a Sonic hedgehog signal transduction pathway activating substance in step 2 on the conditions including a Precondition treatment with a Sonic hedgehog signal transduction pathway activating substance in step 1 and addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3. It was further found that retinal cells (or retina specific nerve cells), for example, photoreceptor precursor cell and ganglion cell can be produced by continuing differentiation culture of the prepared retinal tissues. In addition, it was found that photoreceptor precursor cells and ganglion cell form a continuous epithelial structure having a layer structure in the aforementioned retinal tissues.

[0405]

Example 23: Production Example of retinal tissue using human iPS cell established using Sendaivirus vector as starting material and including Precondition in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2

Human iPS cells (TFH-HA strain, established by Sumitomo Dainippon Pharma Co., Ltd.) were established as follows. They were established using peripheral blood mononuclear cells (PBMC) prepared by a well-known method as the starting material, commercially available Sendaivirus vector (4 factors of Oct3/4, Sox2, KLF4, L-Myc, CytoTune kit manufactured by ID Pharma Co.,

Ltd.), and StemFit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.), Laminin511-E8 (manufactured by Nippi, Inc.).

Human iPS cells (TFH-HA strain) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.
[0406]

10 The feeder-free cultured human iPS cells one day before subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 24 "Precondition: TGF β R-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.3×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium
25 supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium, and the cells were cultured in a serum-free medium
35

under two conditions of the following Conditions 1 and 2.

•Condition 1. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and no addition of an exogenous Sonic hedgehog signal transduction
5 pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 24A, B, "Condition 1").

•Condition 2. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (10 nM) as an exogenous Sonic hedgehog signal
10 transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 24E, F "Condition 2").

By day 2 after the start of suspension culture, cell aggregates were formed under Condition 1 and Condition 2 (step
15 2 completed, and step 3 started).

[0407]

On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing
20 IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e (3 μ M) would not change. Thereafter, a half-medium exchange operation was performed once
25 every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 12 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the
30 aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant
35 BMP4 and IWR-1e.

The thus-prepared cells on day 9 after the start of suspension culture were subjected to bright field observation using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue
5 was formed under Condition 1 and Condition 2.

[0408]

Cell aggregates on day 19 after the start of the aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were
10 immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig.), which is one of the retinal tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE
15 CORPORATION, BIOREVO) (Fig. 24).

[0409]

As a result, it was found that a neural tissue could be formed in Condition 1 and Condition 2. In Condition 1, the proportion of the Chx10 positive retinal tissue was about 90%
20 (Fig. 24A), and in Condition 2, the proportion of the Chx10 positive retinal tissue was about 90% (Fig. 24E). From the analysis of serial sections, it could be confirmed that these Chx10 positive cells are Rx co-positive cells (Fig. 24B, F).

That is, using human iPS cells established using
25 Sendaivirus vector as the starting material in Condition 1, it was found that efficient differentiation into a retinal tissue is also achieved under conditions including Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance
30 in step 1, no addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less
35 in the course of step 3 (on day 12 after the start of

suspension culture). Furthermore, using human iPS cells established using Sendaivirus vector as the starting material in Condition 2, it was found that efficient differentiation into retinal cell is also achieved under conditions including
5 Precondition with a TGF β signal transduction pathway inhibiting substance and a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, and addition of a Wnt signal transduction pathway inhibiting
10 substance in step 2 and step 3.

[0410]

Example 24: Production Example of retinal tissue using human iPS cell established using Sendaivirus vector as starting material and including Precondition in step 1 and use of Wnt
15 signal transduction pathway inhibiting substance in step 2

Human iPS cells (TFH-HA strain, established by Sumitomo Dainippon Pharma Co., Ltd.) were established as describe in Example 23. Human iPS cells were cultured feeder free according to the method described in Example 1 and using
20 StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

[0411]

The feeder-free cultured human iPS cells 2 days before
25 subconfluence were feeder-free cultured for two days under conditions including addition of SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (Fig. 24 "Precondition: Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

30 The thus-prepared Precondition (Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a
35 serum-free medium at 1.3×10^4 cells per well of a non-cell-

adhesive 96-well culture plate (PrimeSurface 96V-bottom plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is
5 a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 µM 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 µM) was added
10 to the above-mentioned serum-free medium, and the cells were cultured in a serum-free medium under two conditions of the following Conditions 3 and 4.

•Condition 3. Condition including addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and no
15 addition of an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 24C, D, "Condition 3").

•Condition 4. Condition including addition of IWR-1e (3 µM) as a Wnt signal transduction pathway inhibiting substance and
20 addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2 (Fig. 24G, H "Condition 4").

By day 2 after the start of suspension culture, cell
25 aggregates were formed under Condition 3 and Condition 4 (step 2 completed, and step 3 started).

[0412]

On day 3 after the start of suspension culture, a medium medium not containing Y27632 or SAG, containing human
30 recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 µl such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and the concentration of exogenous IWR-1e (3 µM) would not change. Thereafter, a half-medium exchange operation was performed once
35 every 2 - 4 days with the aforementioned serum-free medium not

containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

On day 12 after the start of suspension culture, a 80%
5 medium exchange operation was performed 3 times by using the
aforementioned serum-free medium so that the concentration of
exogenous IWR-1e would be 3% or less as compared to that before
medium exchange. Thereafter, a half-medium exchange operation
was performed once every 2 - 4 days with the aforementioned
10 serum-free medium not containing Y27632, SAG, human recombinant
BMP4 and IWR-1e.

The thus-prepared cells on day 9 after the start of
suspension culture were subjected to bright field observation
using an inverted microscope (manufactured by Nikon Corporation,
15 ECLIPSE Ti). As a result, it was found that a neural tissue
was formed under Condition 3 and Condition 4.

[0413]

Cell aggregates on day 19 after the start of the
aforementioned suspension culture were fixed with 4% para-
20 formaldehyde to produce cryosections. These cryosections were
immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep),
which is one of the retinal tissue markers, or Rx (anti-Rx
antibody, Takara, Guinea Pig.), which is one of the retinal
tissue markers. These stained sections were observed with an
25 inverted fluorescence microscope (manufactured by KEYENCE
CORPORATION, BIOREVO) (Fig. 24).

[0414]

As a result, it was found that a neural tissue could be
formed in Condition 3 and Condition 4. In Condition 3, the
30 proportion of the Chx10 positive retinal tissue was about 60%
(Fig. 24C), and in Condition 4, the proportion of the Chx10
positive retinal tissue was about 50% (Fig. 24G). From the
analysis of serial sections, it could be confirmed that these
Chx10 positive cells are Rx co-positive cells (Fig. 24D, H).

35 That is, using human iPS cells established using

Sendaivirus vector as the starting material in Condition 3, it was found that efficient differentiation into a retinal tissue is also achieved under conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, no addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3 (on day 12 after the start of suspension culture). Furthermore, using human iPS cells established using Sendaivirus vector as the starting material in Condition 4, it was found that efficient differentiation into retinal cell is also achieved under conditions including Precondition with a Sonic hedgehog signal transduction pathway activating substance in step 1, addition of a Sonic hedgehog signal transduction pathway activating substance in step 2, addition of a Wnt signal transduction pathway inhibiting substance in step 2 and step 3, and reduction of the concentration of the exogenous Wnt signal transduction pathway inhibiting substance to 3% or less in the course of step 3.

[0415]

Example 25: : Production Example of retinal tissue from human iPS cell including Precondition with TGF β signal transduction pathway inhibiting substance and Sonic hedgehog signal transduction pathway activating substance in step 1 and use of Wnt signal transduction pathway inhibiting substance in step 2 and step 3

Human iPS cells (1231A3 strain, obtained from Kyoto University) were cultured feeder free according to the method described in Example 1 and using StemFit (registered trade mark) medium (AK03N, manufactured by Ajinomoto Co., Inc.) as a feeder-free medium and Laminin 511-E8 (manufactured by Nippi, Inc.) as a feeder-free scaffold.

The feeder-free cultured human iPS cells one day before

subconfluence were feeder-free cultured for one day under conditions including addition of SB431542 (TGF β signal transduction pathway inhibiting substance (TGF β R-i), 5 μ M) and SAG (Sonic hedgehog signal transduction pathway activating substance (Shh), 300 nM) (step 1: Precondition (TGF β R-i+Shh) treatment, Fig. 25 "Precondition: TGF β R-i+Shh") in a Stem Fit (registered trade mark) medium (AK03N; manufactured by Ajinomoto Co., Inc.).

The thus-prepared Precondition (TGF β R-i+Shh)-treated human iPS cells were treated with the cell dispersion solution by using TrypLE Select (Life Technologies), further dispersed into single cells by pipetting operation, and the above human iPS cells dispersed into single cells were suspended in 100 μ l of a serum-free medium at 1.3×10^4 cells per well of a non-cell-adhesive 96-well culture plate (PrimeSurface 96 slit well plate, manufactured by SUMITOMO BAKELITE). Thereafter, the cells were subjected to suspension culture at 37°C, 5% CO₂. As the serum-free medium (gfCDM+KSR) therefor, a serum-free medium which is a 1:1 mixture of F-12 medium and IMDM medium supplemented with 10% KSR, 450 μ M 1-monothioglycerol, 1 x Chemically defined lipid concentrate was used. At the time of the start of suspension culture (day 0 after the start of suspension culture, start of step 2), Y27632 (final concentration 20 μ M) was added to the above-mentioned serum-free medium and the cells were cultured in a serum-free medium under the following Condition 2.

•Condition 2. Condition including addition of IWR-1e (3 μ M) as a Wnt signal transduction pathway inhibiting substance and addition of SAG (10 nM) as an exogenous Sonic hedgehog signal transduction pathway activating substance to the aforementioned serum-free medium at the start of step 2

A cell aggregate was formed by day 2 after the start of suspension culture (step 2 completed, and step 3 started).

[0416]

On day 4 after the start of suspension culture, a medium

medium not containing Y27632 or SAG, containing human recombinant BMP4 (manufactured by R&D) and further containing IWR-1e was added by 50 μ l such that a final concentration of exogenous human recombinant BMP4 would be 1.5 nM (55 ng/ml) and
5 the concentration (3 μ M) of exogenous IWR-1e would not change. Thereafter, a half-medium exchange operation was performed once every 2 - 4 days with the serum-free medium not containing Y27632, SAG, human recombinant BMP4 and further containing IWR-1e to avoid change of the concentration of exogenous IWR-1e.

10 On day 10 after the start of suspension culture, a 80% medium exchange operation was performed 3 times by using the aforementioned serum-free medium so that the concentration of exogenous IWR-1e would be 3% or less as compared to that before medium exchange. Thereafter, a half-medium exchange operation
15 was performed once every 2 - 4 days with the aforementioned serum-free medium not containing Y27632, SAG, human recombinant BMP4 and IWR-1e.

The thus-prepared cells on day 17 after the start of suspension culture were subjected to bright field observation
20 using an inverted microscope (manufactured by Nikon Corporation, ECLIPSE Ti). As a result, it was found that a neural tissue was formed.

[0417]

The aggregates on day 17 after the start of the
25 aforementioned suspension culture were fixed with 4% para-formaldehyde to produce cryosections. These cryosections were immunostained for Chx10 (anti-Chx10 antibody, Exalpha, sheep), which is one of the retinal tissue markers, or Rx (anti-Rx antibody, Takara, Guinea Pig), which is one of the retinal
30 tissue markers. These stained sections were observed with an inverted fluorescence microscope (manufactured by KEYENCE CORPORATION, BIOREVO) (Fig. 25).

[0418]

As a result, it was found that a neural tissue could be
35 formed. In this condition, the proportion of the Chx10

positive retinal tissue was about 90% (Fig. 25A). From the analysis of serial sections, it could be confirmed that these Chx10 positive cells are Rx co-positive cells (Fig. 25, B).

That is, it was found that efficient differentiation into
5 retinal cell is also achieved under conditions including
Precondition with a TGF β signal transduction pathway inhibiting
substance and a Sonic hedgehog signal transduction pathway
activating substance in step 1, addition of a Sonic hedgehog
signal transduction pathway activating substance and seeding
10 into a slit well plate in step 2, and addition of a Wnt signal
transduction pathway inhibiting substance in step 2 and step 3.
[Industrial Applicability]

[0419]

According to the present invention, retinal cells or
15 retinal tissues, and cell aggregates used to produce these can
be produced with high efficiency from pluripotent stem cells
cultured in the absence of feeder cells.

[0420]

The contents disclosed in any publication stated in the
20 present specification, including patents, patent applications
and scientific literatures, are hereby incorporated in their
entireties by reference, to the extent that they have been
disclosed herein.

[0421]

25 This application is based on a patent application No.
2016-086602 filed in Japan (filing date: April 22, 2016), the
contents of which are incorporated in full herein.

CLAIMS

1. A method for producing a retinal cell or a retinal tissue, comprising the following steps (1) - (3):
 - 5 (1) a first step of culturing pluripotent stem cells in the absence of feeder cells in a medium containing 1) a TGF β family signal transduction pathway inhibiting substance and/or a Sonic hedgehog signal transduction pathway activating substance, and 2) a factor for maintaining undifferentiated state,
 - 10 (2) a second step of culturing the cells obtained in the first step in suspension in a medium containing a Wnt signal transduction pathway inhibiting substance to form a cell aggregate, and
 - (3) a third step of culturing the aggregate obtained in the
15 second step in suspension in the presence or absence of a Wnt signal transduction pathway inhibiting substance in a medium containing a BMP signal transduction pathway activating substance to obtain an aggregate containing a retinal cell or a retinal tissue.
- 20 2. The production method according to claim 1, wherein the pluripotent stem cells are cultured for 0.5 hr - 144 hr in the first step.
- 25 3. The production method according to claim 1 or 2, wherein the culturing in the first step is performed by adhesion culture.
4. The production method according to any of claims 1 to 3, wherein, in the second step, the cells obtained in the first
30 step are dispersed, and the dispersed cells are cultured in suspension.
5. The production method according to any of claims 1 to 4, wherein the factor for maintaining undifferentiated state
35 comprises at least an FGF signal transduction pathway

activating substance.

6. The production method according to claim 5, wherein the FGF signal transduction pathway activating substance is bFGF.

5

7. The production method according to any of claims 1 to 6, wherein the medium used for suspension culture in the second step further comprises a Sonic hedgehog signal transduction pathway activating substance.

10

8. The production method according to claim 7, wherein the pluripotent stem cells are human pluripotent stem cells and, in the second step, the concentration of the Sonic hedgehog signal transduction pathway activating substance in the medium is a
15 concentration corresponding to Sonic hedgehog signal transduction activity of SAG at 10 nM to 700 nM.

9. The production method according to any of claims 1 to 8, wherein the TGF β family signal transduction pathway inhibiting
20 substance is a TGF β signal transduction pathway inhibiting substance or a BMP signal transduction pathway inhibiting substance.

10. The production method according to any of claims 1 to 9,
25 wherein the TGF β family signal transduction pathway inhibiting substance is one or more substances selected from the group consisting of Lefty, SB431542, A-83-01 and LDN193189.

11. The production method according to any of claims 1 to 10,
30 wherein the Sonic hedgehog signal transduction pathway activating substance is one or more substances selected from the group consisting of Shh, SAG and Purmorphamine.

12. The production method according to any of claims 1 to 11,
35 wherein, in the third step, the BMP signal transduction pathway

activating substance is added to the medium between day 1 and day 9 from the start of the second step.

13. The production method according to claim 12, wherein, in
5 the third step, the BMP signal transduction pathway activating substance is added to the medium between day 1 and day 6 from the start of the second step.

14. The production method according to claim 13, wherein, in
10 the third step, the BMP signal transduction pathway activating substance is added to the medium between day 1 and day 3 from the start of the second step.

15. The production method according to any of claims 1 to 14,
15 wherein the BMP signal transduction pathway activating substance is one or more proteins selected from the group consisting of BMP2, BMP4, BMP7 and GDF7.

16. The production method according to any of claims 1 to 15,
20 wherein the BMP signal transduction pathway activating substance is BMP4.

17. The production method according to any of claims 1 to 16,
wherein, in the third step, the concentration of the Sonic
25 hedgehog signal transduction pathway activating substance in the medium is not more than a concentration corresponding to Sonic hedgehog signal transduction activity of SAG at 700 nM.

18. The production method according to any of claims 1 to 17,
30 wherein the culturing is performed in a medium containing a Wnt signal transduction pathway inhibiting substance for 3 days to for 18 days from the start of the second step.

19. The production method according to any of claims 1 to 18,
35 wherein the culturing is performed in a medium containing a Wnt

signal transduction pathway inhibiting substance for 10 days from the start of the second step.

20. The production method according to any of claims 1 to 19,
5 wherein the Wnt signal transduction pathway inhibiting substance is IWR-1-endo.

21. The production method according to any of claims 1 to 20,
wherein the aggregate obtained in the third step comprises one
10 or more cells selected from the group consisting of retinal progenitor cell, neural retinal progenitor cell, photoreceptor precursor cell, photoreceptor cell, rod photoreceptor cell, cone photoreceptor cell, horizontal cell, amacrine cell, interneuron, ganglion cell, bipolar cell, retinal pigment
15 epithelial cell, and ciliary marginal zone cell.

22. The production method according to any of claims 1 to 21,
wherein the pluripotent stem cells are human pluripotent stem cells.

20

23. The production method according to any of claims 1 to 22,
wherein the pluripotent stem cells are induced pluripotent stem cells.

25 24. The production method according to any of claims 1 to 23,
wherein a uniformed aggregate is formed in the second step.

25. The production method according to any of claims 1 to 24,
wherein the suspension culture is performed in the absence of a
30 basement membrane preparation.

26. A reagent for evaluating toxicity or efficacy of a test substance, comprising a retinal cell or a retinal tissue produced by the method according to any of claims 1 to 25.

35

27. A method for evaluating toxicity or efficacy of a test substance, comprising bringing the substance into contact with a retinal cell or a retinal tissue produced by the method according to any of claims 1 to 25, and detecting an influence
5 of the substance on the cell or tissue.

28. A medicament for treating a disease due to a disorder of a retinal cell or a retinal tissue, comprising a retinal cell or a retinal tissue produced by the method according to any of
10 claims 1 to 25.

29. The medicament according to claim 28, wherein the retinal cell or retinal tissue is a retinal progenitor cell, a retinal layer-specific neural cell or a retinal tissue.
15

30. A method for treating a disease due to a disorder of a retinal cell or a retinal tissue, comprising transplanting an effective amount of a retinal cell or a retinal tissue produced by the method according to any of claims 1 to 25 to a subject
20 in need of the transplantation.

31. A retinal cell or a retinal tissue produced by the method according to any of claims 1 to 25 for use in the treatment of a disease due to a disorder of a retinal cell or a retinal
25 tissue.

32. A pharmaceutical composition comprising a retinal cell or a retinal tissue produced by the method according to any of claims 1 to 25 as an active ingredient.