

DESCRIPTION

NITROGEN-CONTAINING HETEROCYCLIC CARBOXAMIDE
DERIVATIVES OR SALTS THEREOF AND ANTIVIRAL
AGENTS COMPRISING THE SAME

TECHNICAL FIELD

The present invention relates to antiviral agents comprising a nitrogen-containing heterocyclic carboxamide derivative or a salt thereof.

5 BACKGROUND ART

Nowadays, antiviral agents are selected and put to use in accordance with the objective viruses. For instance, Acyclovir and Vidarabine are used against herpes viruses; Ganciclovir and Foscarnet are used
10 against cytomegalo virus; and interferon is used against hepatitis viruses.

Influenza virus is a central virus of the cold syndrome, which has attacked human being periodically to cause many deaths amounting to tens
15 millions. Although the number of deaths shows a tendency of decrease in the recent years owing to the improvement in hygienic and nutritive conditions, the prevalence of influenza is repeated every year, and it is apprehended that a new virus may appear to cause a
20 wider prevalence.

For prevention of influenza virus, vaccine is used widely, in addition to which low molecular weight

substances such as Amantadine and Ribavirin are also used.


Amantadine is used for prevention and treatment of influenza. Its function mechanism is said to consist in inhibiting the fusion between influenza virus and cell membrane, and it is effective against A-type influenza virus. Its problems are, however, that it is ineffective against B type influenza virus, its resistant virus appears, and it causes side effects such as nerve disturbance. Although Rimantadine which is a derivative of Amantadine has a more improved antiviral activity, the problem of side effect is not overcome by it. Ribavirin which is a guanosine derivative shows a viral RNA polymerase-inhibitory activity and is effective upon A type and B type influenza viruses. Its internal use, however, brings about no sufficient clinical effect.

The present invention provides an antiviral agent exhibiting a preventive effect and a therapeutic effect against various viruses, especially influenza viruses.


DISCLOSURE OF THE INVENTION

The present inventors have conducted researches and studies on compounds showing an antiviral activity against various viruses, especially influenza viruses. As a result, it has been found that pyrazine carboxamide derivatives have an anti-influenza

virus activity. The inventors conducted further studies to find that nitrogen-containing heterocyclic carboxamide derivatives represented by the following general formula [1]:



5 wherein ring A represents a substituted or unsubstituted pyrazine, pyrimidine, pyridazine or triazine ring; R¹ represents O or OH; R² represents a hydrogen atom, an acyl group or a substituted or unsubstituted carbamoylalkyl or carboxyalkyl group; and the broken
 10 line represents a single bond or a double bond;
 or salts thereof exhibit an excellent antiviral activity against A-, B- and C-type of influenza viruses and other various viruses, these compounds have low cytotoxicity and are useful as an antiviral agents of
 15 high safety, as well as that novel N-containing heterocyclic carboxamide derivatives represented by the following general formula [1a]:



wherein ring A' represents a pyrazine ring substituted with a halogen atom, a hydroxyl group or an oxide group; R¹ represents O or OH; R² represents a hydrogen atom, an acyl group or a substituted or unsubstituted carbamoylalkyl or carboxyalkyl group; and the broken line represents a single bond or a double bond; or salts thereof exhibit an excellent antiviral activity. Based on these findings, the present invention has been accomplished.

10 The present invention will be described in detail below.

 Unless otherwise indicated, the term "halogen atom" used in this specification means a fluorine atom, a chlorine atom, a bromine atom or an iodine atom; the term "alkyl group" means a straight or branched chain C₁₋₆ alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl and the like; the term "alkenyl group" means a straight or branched chain C₂₋₆ alkenyl group such as vinyl, allyl and the like; the term "cycloalkyl group" means a C₃₋₆ cycloalkyl group such as cyclopropyl, cyclopentyl, cyclohexyl and the like; the term "alkoxy group" means a straight or branched chain C₁₋₆ alkyl-O-group such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy and the like; the term "cycloalkyloxy group" means a C₃₋₆ cycloalkyl-O- group such as cyclopropyloxy, cyclopentyloxy, cyclohexyloxy and the like; the term

"alkylthio group" means a straight or branched chain C_{1-6} alkyl-S- group such as methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio, tert-butylthio, pentylthio and the like;

5 the term "alkylamino group" means an amino group substituted with one or more straight or branched chain C_{1-6} alkyl groups such as methylamino, ethylamino, propylamino, butylamino, pentylamino, hexylamino, dimethylamino, diethylamino, methylethylamino,

10 dipropylamino, dibutylamino, dipentylamino and the like; the term "cycloalkylamino group" means a C_{3-6} cycloalkyl-NH- group such as cyclopropylamino, cyclopentylamino, cyclohexylamino and the like; the term "halogenoalkyl group" means a halogen-substituted

15 C_{1-6} alkyl group such as trifluoromethyl, trichloromethyl, chloromethyl and the like; the term "aryl group" means a phenyl group, a naphthyl group and the like; the term "aryloxy group" means an aryl-O- group such as phenyloxy, naphthyloxy and the like; the term

20 "arylthio group" means an aryl-S- group such as phenylthio, naphthylthio and the like; the term "arylamino group" means an aryl-NH- group such as phenylamino, naphthylamino and the like; the term "acyl group" means a C_{2-5} alkanoyl group such as formyl,

25 acetyl, propionyl, butyryl, isobutyryl, valeryl and the like and an aroyl group such as benzoyl, naphthoyl and the like; the term "alkoxycarbonyl group" means a straight or branched chain C_{1-6} alkoxycarbonyl group such

as methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl and the like; the term


- 5 "alkylcarbamoyl group" means a carbamoyl group substituted with one or more straight or branched chain C_{1-6} alkyl groups such as methylcarbamoyl, dimethylcarbamoyl and the like; the term "carbamoylalkyl group" means a straight or branched chain C_{1-6} alkyl group
- 10 substituted with a carbamoyl group such as carbamoylmethyl, carbamoylethyl, carbamoylisopropyl and the like; the term "carboxyalkyl group" means a straight or branched chain C_{1-6} alkyl group substituted with a carboxyl group such as carboxymethyl, carboxylethyl,
- 15 carboxyisopropyl and the like; the term "heterocyclic group" means a 4-, 5- or 6-membered ring or a fused ring thereof having, as the hetero atoms constituting said ring, at least one heteroatoms selected from the group consisting of oxygen atom, nitrogen atom and
- 20 sulfur atom, such as oxetanyl, thietanyl, azetidiny, furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, isothiazolyl, pyrrolidinyl, benzofuranyl, benzothiazolyl, pyridyl, quinolyl, pyrimidinyl and morpholinyl; and the term "oxide group"
- 25 means an oxygen atom linked to a nitrogen atom in a ring. The term "lower" means that the number of carbon atoms is 1 to 6.

The protecting group for carboxyl group

includes any groups which can conventionally be used as a protecting group for carboxyl group. The examples thereof include alkyl groups such as methyl, ethyl, n-propyl, isopropyl, 1,1-dimethylpropyl, n-butyl, tert-butyl and the like; aryl groups such as phenyl, naphthyl and the like; aralkyl groups such as benzyl, diphenylmethyl, trityl, p-nitrobenzyl, p-methoxybenzyl, bis(p-methoxyphenyl)methyl and the like; acylalkyl groups such as acetylmethyl, benzoylmethyl, p-nitrobenzoylmethyl, p-bromobenzoylmethyl, p-methanesulfonylbenzoylmethyl and the like; oxygen-containing heterocyclic groups such as 2-tetrahydropyranyl, 2-tetrahydrofuranyl and the like; halogeno-alkyl groups such as 2,2,2-trichloroethyl and the like; alkylsilylalkyl groups such as 2-(trimethylsilyl)ethyl and the like; acyloxyalkyl groups such as acetoxymethyl, propionyloxymethyl, pivaloyloxymethyl and the like; nitrogen-containing heterocyclic alkyl groups such as phthalimidomethyl, succinimidomethyl and the like; cycloalkyl groups such as cyclohexyl and the like; alkoxyalkyl groups such as methoxymethyl, ethoxymethyl, 2-(trimethylsilyl)ethoxymethyl and the like; aralkoxyalkyl groups such as benzyloxymethyl and the like; alkylthioalkyl groups such as methylthiomethyl, 2-methylthioethyl and the like; arylthioalkyl groups such as phenylthiomethyl and the like; alkenyl groups such as 1,1-dimethyl-2-propenyl, 3-methyl-3-butenyl, allyl and the like;

substituted silyl groups such as trimethylsilyl, triethylsilyl, triisopropylsilyl, diethylisopropylsilyl, tert-butyldimethylsilyl, tert-butyldiphenylsilyl, diphenylmethylsilyl, tert-butylmethoxyphenylsilyl and the like; etc.

In general formula [1], ring A represents a pyrazine ring, a pyrimidine ring, a pyridazine ring or a triazine ring. More specifically, ring A represents any of the following structures:



10 In the structures above, the mark *1 expresses the position of substitution with R^1 , and the mark *2 expresses the position of substitution with $-C(=O)NHR^2$.

The carbamoylalkyl or carboxyalkyl group represented by R^2 may be substituted with at least one
 15 substituent selected from the group consisting of halogen atoms; alkyl groups unsubstituted or substituted with hydroxyl, alkoxy, alkylthio, aryl,

amino or alkylamino groups; halogenoalkyl groups;
 alkenyl groups; cycloalkyl groups; hydroxyl groups;
 alkoxy groups; cycloalkyloxy groups; alkoxycarbonyl
 groups; mercapto groups; alkylthio groups unsubstituted
 5 or substituted with one or more aryl groups; aryl
 groups; aryloxy groups; arylthio groups; arylamino
 groups; cyano groups; nitro groups; amino groups
 unsubstituted or substituted with one or more acyl
 groups; alkylamino groups; cycloalkylamino groups; acyl
 10 groups; hydrazino groups; carboxyl groups; carbamoyl
 groups; thiocarbamoyl groups; alkylcarbamoyl groups and
 heterocyclic groups.

Of ring A represented by general formula [1],
 preferred are pyrazine ring, pyrimidine ring and
 15 triazine ring, and more preferred is pyrazine ring.
 The substituent on ring A includes groups selected from
 the group consisting of halogen atoms; alkyl groups
 unsubstituted or substituted with one or more hydroxyl,
 alkoxy, alkylthio, aryl, amino or alkylamino groups;
 20 halogenoalkyl groups; alkenyl groups; cycloalkyl
 groups; hydroxyl groups; alkoxy groups; cycloalkyloxy
 groups; alkoxycarbonyl groups; mercapto groups;
 alkylthio groups unsubstituted or substituted with one
 or more aryl groups; aryl groups; aryloxy groups;
 25 arylthio groups; arylamino groups; cyano groups; nitro
 groups; amino groups unsubstituted or substituted with
 one or more acyl groups; alkylamino groups;
 cycloalkylamino groups; acyl groups; hydrazino groups;

carboxyl groups; carbamoyl groups; thiocarbamoyl groups, alkylcarbamoyl groups and heterocyclic groups. Ring A can have one or more of the above-mentioned substituents. Further, the substituent on ring A is preferably linked to carbon atom of the ring.

The salt of the compound of general formula [1] includes any usually known salts formed at the site of basic groups such as amino group and salts formed at the site of acidic group such as hydroxyl and carboxyl groups. The salts formed at the site of basic group include salts of mineral acids such as hydrochloric acid, hydrobromic acid, sulfuric acid and the like; salts of organic carboxylic acids such as tartaric acid, formic acid, citric acid, trichloroacetic acid, trifluoroacetic acid and the like, and salts of sulfonic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, mesitylenesulfonic acid, naphthalenesulfonic acid and the like. The salts formed at the site of acidic group include salts of alkali metals such as sodium, potassium and the like; salts of alkaline earth metals such as calcium, magnesium and the like; ammonium salts; and salts of nitrogen-containing organic bases such as trimethylamine, triethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, diethylamine, dicyclohexylamine, procaine, dibenzylamine, N-benzyl- β -phenethylamine, 1-phenamine, N,N'-dibenzylethylenediamine and the like.

Among the salts mentioned above, preferred are pharmacologically acceptable ones.

Typical compounds represented by general formula [1] are shown in Tables 1 to 4.

5 In the tables shown below, the abbreviations have the following meanings:

Me: methyl

Et: ethyl

iPr: isopropyl

10 tBu: tert-butyl,

Ph: phenyl

Ac: acetyl

Bz: benzoyl

R2a: $-\text{CH}(\text{COOH})\text{CH}_2\text{COOH}$

15 R2b: $-\text{CH}(\text{CH}_3)\text{CONHCH}(\text{CH}_3)\text{COOH}$




Table 1

No.	G ₁	G ₂	G ₃	G ₄	R ²
1	N	CH	CH	N	H
2	N	CH	CH	N→O	H
3	N	CH	CH	N	Ac
4	N	CH	CH	N	Bz
5	N	CH	CH	H	C(O)tBu
6	N	CH	CH	H	CH ₂ CONH ₂
7	N	CH	C-Cl	N	H
8	N	CH	C-Br	N	H
9	N	CH	C-OH	N	H
10	N	CH	C-OMe	N	H
11	N	CH	C-OEt	N	H
12	N	CH	C-OiPr	N	H
13	N	CH	C-O-cyclopropyl	N	H
14	N	CH	C-NHMe	N	H
15	N	CH	C-NMe ₂	N	H
16	N	CH	C-NH-cyclopropyl	N	H
17	N	CH	C-NHAc	N	H
18	N	CH	C-NHBz	N	H
19	N	CH	C-NHPh	N	H
20	N	CH	C-SH	N	H
21	N	CH	C-SMe	N	H
22	N	CH	C-COOH	N	H
23	N	CH	C-COOMe	N	H
24	N	CH	C-COOEt	N	H
25	N	CH	C-CONH ₂	N	H
26	N	CH	C-CSNH ₂	N	H
27	N	CH	C-CONHMe	N	H
28	N	CH	C-CONMe ₂	N	H
29	N	CH	C-Et	N	H
30	N	CH	C-iPr	N	H
31	N	CH	C-CCl ₃	N	H
32	N	CH	C-CF ₃	N	H
33	N	CH	C-CH ₂ Cl	N	H
34	N	CH	C-CH ₂ -SMe	N	H
35	N	CH	C-CH ₂ -OMe	N	H
36	N	CH	C-CH ₂ -Ph	N	H
37	N	CH	C-CH ₂ -NH ₂	N	H
38	N	CH	C-CH ₂ -OH	N	H
39	N	CH	C-CH ₂ -NHMe	N	H
40	N	CH	C-CH=CH ₂	N	H

Table 2

No.	G ₁	G ₂	G ₃	G ₄	R ²
41	N	CH	C-Ph	N	H
42	N	CH	C-pyridin-3-yl	N	H
43	N	CH	C-furan-2-yl	N	H
44	N	CH	C-thiophen-2-yl	N	H
45	N	CH	C-thiazol-2-yl	N	H
46	N	CH	C-pyrrolidin-1-yl	N	H
47	N	CH	C-piperidin-1-yl	N	H
48	N	CH	C-morpholin-4-yl	N	H
49	N	CH	C-CN	N	H
50	N	CH	C-NO ₂	N	H
51	N	CH	C-Bz	N	H
52	N	CH	C-Ac	N	H
53	N	C-CONH ₂	C-NH ₂	N	H
54	N	C-NH ₂	C-COOH	N	H
55	N	C-NH ₂	C-COOMe	N	H
56	N	C-Cl	C-COOMe	N	H
57	N	C-OMe	C-Me	N	H
58	N	C-COOH	C-Me	N	H
59	N	C-COOH	C-NHMe	N	H
60	N	C-COOMe	C-Cl	N	H
61	N	C-COOMe	C-piperidin-1-yl	N	H
62	N	C-OMe	C-CN	N	H
63	N	C-Me	C-Me	N	H
64	N	C-Ph	C-Ph	N	H
65	N	C-F	CH	N	H
66	N	C-Cl	CH	N	H
67	N	C-Br	CH	N	H
68	N	C-OH	CH	N	H
69	N	C-OMe	CH	N	H
70	N	C-OET	CH	N	H
71	N	C-O-iPr	CH	N	H
72	N	C-cyclopropyl	CH	N	H
73	N	C-OPh	CH	N	H
74	N	C-NH ₂	CH	N	H
75	N	C-NHMe	CH	N	H
76	N	C-NMe ₂	CH	N	H
77	N	C-cyclopropyl	CH	N	H
78	N	C-NHPh	CH	N	H
79	N	C-NHAc	CH	N	H
80	N	C-NHBz	CH	N	H

Table 3

No.	G ₁	G ₂	G ₃	G ₄	R ²
81	N	C-COOMe	CH	N	H
82	N	C-CF ₃	CH	N	H
83	N	C-Ph	CH	N	H
84	N	C-pyridin-4-yl	CH	N	H
85	N	C-CN	CH	N	H
86	N	C-NO ₂	CH	N	H
87	CH	N	CH	N	H
88	CH	N	C-Me	N	H
89	CH	N	C-Et	N	H
90	CH	N	C-iPr	N	H
91	CH	N	C-F	N	H
92	CH	N	C-Cl	N	H
93	CH	N	C-Br	N	H
94	CH	N	C-Ph	N	H
95	CH	N	C-OH	N	H
96	CH	N	C-OMe	N	H
97	CH	N	C-OEt	N	H
98	CH	N	C-OiPr	N	H
99	CH	N	C-SH	N	H
100	CH	N	C-SMe	N	H
101	CH	N	C-S-CH ₂ Ph	N	H
102	CH	N	C-SPh	N	H
103	CH	N	C-NH ₂	N	H
104	CH	N	C-NHMe	N	H
105	CH	N	C-NMe ₂	N	H
106	CH	N	C-piperidin-1-yl	N	H
107	CH	N	C-NH-Ph	N	H
108	C-OH	N	C-SH	N	H
109	C-OH	N	C-NH ₂	N	H
110	C-Me	N	C-OH	N	H
111	N	CH	N	CH	H
112	N	C-F	N	CH	H
113	N	C-Cl	N	CH	H
114	N	C-F	N	C-Me	H
115	N	C-Cl	N	C-Et	H
116	N	C-OMe	N	C-OH	H
117	N	C-NH ₂	N	C-OH	H
118	N	C-NHAc	N	C-OH	H
119	N	C-SH	N	C-OH	H
120	N	C-SMe	N	C-OH	H


Table 4

No.	G ₁	G ₂	G ₃	G ₄	R ²
121	N	C-Me	N	C-NHMe	H
122	N	C-Ph	N	C-OH	H
123	N	C-Me	N	C-OH	H
124	N	C-Me	N	C-S-Ph	H
125	N	C-Me	N	C-Cl	H
126	N	C-Me	N	C-OMe	H
127	N	C-Me	N	C-OPh	H
128	N	C-Me	N	C-morpholin-4-yl	H
129	CH	CH	N	N	H
130	CH	C-OH	N	N	H
131	CH	C-Me	N	N	H
132	N	CH	N	N	H
133	N	C-Cl	N	N	H
134	N	C-Me	N	N	H
135	N	C-OMe	N	N	H
136	N	C-NH ₂	N	N	H
137	N	C-NHAc	N	N	H
138	N	C-pyridin-4-yl	N	N	H
139	N	N	CH	N	H
140	N	N	C-Br	N	H
141	N	N	C-OH	N	H
142	N	N	C-OiPr	N	H
143	N	N	C-NHPh	N	H
144	N	N	C-thiazol-2-yl	N	H
145	N	N	C-SH	N	H
146	N	N	C-SMe	N	H
147	N	N	C-Cl	N	H
148	N	N	C-NHNNH ₂	N	H
149	N	N	C-Ph	N	H
150	N	N	C-pyridin-2-yl	N	H
151	N	N	C-thiophen-2-yl	N	H
152	N	N	C-NHMe	N	H
153	N	N	C-NMe ₂	N	H
154	N	CH	C-F	N	H
155	N	CH	CH	N	R2a
156	N	CH	CH	N	R2a

The nitrogen-containing heterocyclic carboxamide derivatives represented by general formula [1] or salts thereof are commercially available or can be produced according to any of known processes or
5 analogous processes or by a combination thereof. As the papers describing the production processes thereof, J. Am. Chem. Soc., 71, 78 (1949); J. Am. Chem. Soc., 78, 242-244 (1956); J. Heterocycl. Chem., 15(4), 665-670 (1978); J. Chem. Soc., 1379 (1955); U.S. Patent No.
10 5,597,823; etc. can be referred to.

More specifically, the nitrogen-containing heterocyclic carboxamide derivatives represented by general formula [1] or salts thereof can be produced according to the following Production Processes 1 to 3.

Production Process 1



wherein R^1 , R^2 , ring A and broken line are as defined above, and R^3 represents a protecting group for carboxyl group.

- 5 (1-a) A compound of general formula [1] can be obtained by reacting a compound of general formula [2] with a compound of general formula [4].

The solvent which can be used in this reaction is not particularly limited, so far as it causes no adverse effect on the reaction. The examples of the solvent include alcohols such as methanol, ethanol, isopropyl alcohol and the like; halogenated hydrocarbons such as dichloromethane, chloroform, dichloroethane and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; ethers such as dioxane, tetrahydrofuran, anisole, diethylene glycol diethyl ether, dimethyl cellosolve and the like;

10
15

nitriles such as acetonitrile and the like; amides such as N,N-dimethylformamide, N,N-dimethylacetamide and the like; and sulfoxides such as dimethyl sulfoxide and the like. These solvents can be used in admixture.

5 The compound of general formula [4] is used at least in an equimolar amount to the compound of general formula [2], and preferably in an amount of 1.0-5.0 mole per mole of the compound of general formula [2].

10 This reaction can be carried out usually at 0-100°C and preferably at 20-80°C, for a period of 5-24 hours and preferably for 30 minutes to 10 hours.

(1-b) A compound of general formula [1] can be obtained by subjecting a compound of general formula
15 [3] and a compound of general formula [4] to a dehydrating condensation reaction.

 The solvent which can be used in this reaction is not particularly limited, so far as it causes no adverse effect on the reaction. The examples
20 of the solvent include ethers such as dioxane, tetrahydrofuran, anisole, diethylene glycol diethyl ether, dimethyl cellosolve and the like; halogenated hydrocarbons such as dichloromethane, chloroform, dichloroethane and the like; and amides such as N,N-
25 dimethylformamide, N,N-dimethylacetamide and the like. These solvents can be used in admixture.

 The compound of general formula [4] can be


used at least in an equimolar amount to the compound of general formula [3] and preferably in an amount of 1.0-2.0 mole per mole of the compound of general formula [3].

5 The dehydrating condensing agent which can be used in this reaction includes, for example, 1,3-dicyclohexyl carbodiimide, N,N'-carbonyl diimidazole, 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide and the like.

10 The dehydrating condensing agent can be used at least in an equimolar amount to the compound of general formula [3] and preferably in an amount of 1.0-2.0 mole per mole of the compound of general formula [3].

15 This reaction can be carried out usually at 0-100°C and preferably at 20-60°C, for a period of 5 minutes to 24 hours and preferably for 30 minutes to 10 hours.

Production Process 2




wherein R^1 , R^2 , ring A and broken line are as defined above, and R^4 represents a lower alkyl group.

A compound of general formula [1] can be obtained by subjecting a compound of general formula
5 [5] to alkyl-ether scission.

More specifically, in a case where R^4 is a methyl group, the reaction can be carried out according to the description of PROTECTIVE GROUPS IN ORGANIC SYNTHESIS, Second edition, JOHN WILEY & SONS, pp. 145-
10 199 (1991) or by an analogous method.

Production Process 3



wherein R^1 , ring A and broken line are as defined above, and R^{2a} represents an acyl group.

A compound of general formula [1c] can be
15 obtained by subjecting a compound of general formula [1b] to acylation in the presence of an acid-eliminating agent.

The solvent which can be used in this reaction is not particularly limited, so far as it
20 causes no adverse effect on the reaction. The examples of the solvent include ethers such as dioxane,

tetrahydrofuran, anisole, diethylene glycol diethyl ether, dimethyl cellosolve and the like; aromatic hydrocarbons such as benzene, toluene, xylene and the like; halogenated hydrocarbons such as dichloromethane, 5 chloroform, dichloroethane and the like; amides such as N,N-dimethylformamide, N,N-dimethylacetamide and the like; water; etc. These solvents can be used in admixture.

The acylating agent can be used at least in 10 an equimolar amount to the compound of general formula [1b] and preferably in an amount of 1.0-2.0 mole per mole of the compound of general formula [1b].

The acid-eliminating agent used in this reaction includes, for example, pyridine, triethyl- 15 amine, sodium hydrogen carbonate and the like.

The acid-eliminating agent can be used at least in an equimolar amount to the compound of general formula [1b] and preferably in an amount of 1.0-2.0 mole per mole of the compound of general formula [1b].

20 This reaction can be carried out usually at 0-100°C and preferably at 20-60°C, for a period of 5 minutes to 24 hours and preferably for 30 minutes to 10 hours.

In the Production Processes 1-3, the 25 compounds of general formula [2], [3], [4], [5] and [1b] can be replaced with their salt, respectively. As the salts, the same ones as mentioned for the compound of general formula [1] can be used.

Some of the compounds of general formulas [2], [3], [4], [5] and [1b] and salts thereof may have various isomers such as optical isomers and position isomers, and solvated products. In such cases, any of
5 these isomers and solvates may be used in the present invention.

The compound of general formula [1c] thus obtained can be converted to salts thereof. The salts include the same ones as mentioned for the compound of
10 general formula [1].

The objective viruses of the antiviral agent comprising the nitrogen-containing heterocyclic carboxamide derivative represented by general formula [1] or salt thereof according to the present invention
15 include A-, B- and C-type influenza viruses, papilloma virus, adenovirus, A type hepatitis virus, B type hepatitis virus, C type hepatitis virus, poliovirus, echovirus, Coxsackie virus, enterovirus, rhinovirus, rotavirus, Newcastle disease virus, mumps virus,
20 vesicular stomatitis virus, and Japanese encephalitis virus. The antiviral agent of the present invention exhibits an especially high effect against influenza viruses.

By combining the nitrogen-containing
25 heterocyclic carboxamide derivatives of the present invention represented by general formula [1] or salts thereof with conventional known excipients, adjuvants and additives, pharmaceutical preparations such as

solutions, suspensions, powders, granules, fine granules, tablets, capsules, syrups, elixirs, spirits, troches, gargles, aerosols, etc. can be obtained.

These pharmaceutical preparations can be administered
5 either orally or non-orally, namely by injection, percutaneous administration, intrarectal administration, intranarial administration, etc.

The method of administration, dosage and frequency of administration of the antiviral agent of
10 the present invention can be properly selected depending upon the age, body weight and symptom of the patient. Usually 1 to 10 mg/kg of the nitrogen-containing heterocyclic carboxamide derivative or a salt thereof can be administered to an adult either at
15 once or in several portions.

Next, antiviral activity and cytotoxicity of the nitrogen-containing heterocyclic carboxamide derivatives of the present invention represented by general formula [1] or salts thereof will be explained.

20 Sample: A nitrogen-containing heterocyclic carboxamide derivative represented by general formula [1] or a salt thereof was dissolved in dimethyl sulfoxide to prepare a solution having a concentration of 10 mg/mL. At the time of use, the solution was
25 diluted with a culture medium to a predetermined concentration and then put to use.

As the host cell of influenza virus, MDCK cell (canine kidney cell) was used. For the

cytotoxicity test, Vero cell (monkey kidney cell) was used.

Culture medium: In the multiplication of MDCK cell and Vero cell and in the cytotoxicity test using Vero cell, E'-MEM (product of Nissui) to which 10% fetal bovine serum had been added was used.

In the measurement of antiviral activity, E'-MEM (product of Nissui) to which 1% bovine serum albumin had been added was used.

10 Test Example 1 (Anti-influenza activity)

MDCK cells were seeded to a 6-well plate (product of CORNING) at 5×10^5 cells/well, and cultured overnight at 35°C in 5% CO₂-air atmosphere. Then, the cultured MDCK cells on the plate was treated with influenza virus A/PR/8/34 strain diluted with a serum-free culture medium at the concentration of 200 PFU/mL, at 0.5 mL/well for one hour to achieve inoculation and adsorption. After completion of the inoculation and adsorption, an E'-MEM culture medium containing 0.6% Agar Noble, 1% bovine serum albumin and 3 µg/mL acetyltrypsin and also containing a test compound at a prescribed concentration was added to the cells. After sufficient coagulation, the plate was turned upside down, and cultured for 3 days. After completion of the culture, the alive cells were dyed with 1% Neutral Red. Then, the cells were fixed with 10% formalin. The agar medium was removed therefrom with running water.

Thereafter, the number of plaques was counted. The plaque inhibition rate was expressed in percentage calculated in comparison with control containing no test compound.

- 5 The results are shown in Table 5, wherein the test compound number are the same as those in Tables 1 to 4.

Table 5

No.	Concentration of test compound added ($\mu\text{g/mL}$)	Inhibition rate (%)
1	1	91.9
2	100	32.4
9	100	50.0
41	100	25.0
65	1	100
66	100	39.2
67	100	35.2
83	100	39.8
84	100	39.5
87	100	85.7
95	100	30.5
129	100	28.0
139	100	49.3
141	100	26.3
145	100	36.8
154	100	23.0
155	10	35.5
156	100	36.0

Test Example 2 (Cytotoxic activity)

- 10 A culture medium containing a predetermined concentration of test compound was added to a 96-well

plate (product of CORNING) in 100 μ L/well. Then, Vero cells were adjusted to the concentration of 2×10^4 cells/mL with culture medium. The solution was added to the plate at 100 μ L/well, and cultured at 37°C in 5% CO₂-air atmosphere for 4 days. When the culture was completed, the number of alive cells was counted according to the XTT method [for example, see CANCER RESEARCH, Vol. 48, Pages 4827-4833 (1988)].

As a result, the 50% cell growth inhibitory concentration (IC₅₀) of 3-hydroxy-2-pyrazinecarboxamide (compound No. 1) was 250 μ g/mL or more.

BEST MODE FOR CARRYING OUT THE INVENTION

Next, the compound of the present invention is explained with reference to Referential Examples and Examples. The invention is by no means limited by these examples.

In the eluent, the mixing ratios are all by volume.

The carrier used in column chromatography was Silica Gel BW-127ZH (product of Fuji Silicia Chemical Co.), and the carrier used in the reverse phase column chromatography was LC-SORB SP-B-ODS (product of Chemco Co.).

The symbol mark used in the Referential Examples and Examples has the following meaning:

DMSO-d₆ : Deuterated dimethyl sulfoxide

Referential Example 1

In 6 mL of dichloromethane is suspended 0.30 g of methyl 3-hydroxy-2-pyrazinecarboxylate obtained according to the method described in literature [J. Heterocycl. Chem., 34, 27 (1997)]. To the suspension are successively added 0.54 mL of triethylamine and 0.29 g of glycine methyl ester hydrochloride. The resulting mixture is stirred at ambient temperature for 5 hours. After cooling, the solvent is distilled off under reduced pressure. The residue is purified by reverse phase column chromatography (eluent: water) to obtain 0.16 g of methyl 2-[(3-hydroxy-2-pyrazinyl)-carbonyl]amino}acetate.

IR(KBr) cm^{-1} : 1750, 1735, 1685

15 NMR(DMSO- d_6) δ value: 3.67(3H,s), 4.12(2H,d,J=6Hz), 7.70-8.30(2H,m), 9.60-10.10(1H,m), 13.10(1H,brs)

Referential Example 2

In 100 mL of concentrated sulfuric acid is dissolved 17.00 g of methyl 6-bromo-3-amino-2-pyrazinecarboxylate obtained according to the method described in literature [J. Am. Chem. Soc., 2798-2800 (1949)]. At an ice-cooled temperature, 10.11 g of sodium nitrite is added to the suspension, which is stirred for 30 minutes. The reaction mixture is poured into 920 mL of methanol, and heated under reflux for 5 hours. The reaction mixture is cooled and then concentrated under reduced pressure. The residue thus

obtained is added to a mixture of 500 mL of ice water and 600 mL of chloroform and separated into layers.

The organic layer is washed successively with saturated aqueous solution of sodium hydrogen carbonate, water

5 and saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. Then, the solvent is distilled off under reduced pressure to obtain 6.30 g of methyl 6-bromo-3-methoxy-2-pyrazinecarboxylate.

10 IR(KBr) cm^{-1} : 1734

NMR(CDCl_3) δ value: 3.97(3H,s), 4.06(3H,s), 8.37(1H,s)

Referential Example 3

In a nitrogen atmosphere, 11.38 g of methyl 6-bromo-3-methoxy-2-pyrazinecarboxylate is dissolved in 15 227 mL of toluene. To the solution are successively added 10.32 g of benzophenone-imine, 0.42 g of tris(dibenzylideneacetone) dipalladium, 0.86 g of (s)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and 6.20 g of sodium t-butoxide. The resulting mixture is 20 stirred at 80°C for one hour. The reaction mixture is cooled and filtered. The filtrate is purified by column chromatography (eluent: toluene:ethyl acetate = 20:1). The oily product thus obtained is dissolved in 140 mL of tetrahydrofuran, 7 mL of 2 mol/mL 25 hydrochloric acid is added thereto and the resulting solution is stirred at ambient temperature for 15 minutes. The resulting reaction mixture is added to a

mixture of 200 mL of chloroform and 50 mL of water, and alkalified with 1 mol/mL of sodium hydroxide, and the organic layer is separated. The organic layer is washed with saturated aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and the solvent is distilled off under reduced pressure. The residue thus obtained is purified by column chromatography (eluent: toluene:ethyl acetate = 1:1) to obtain 3.64 g of methyl 6-amino-3-methoxy-2-pyrazinecarboxylate.

IR(KBr) cm^{-1} : 1716, 1670

NMR(DMSO- d_6) δ value: 3.80(3H,s), 3.82(3H,s), 7.20(2H,brs), 7.77(1H,s)

Referential Example 4

In 70 mL of methanol is dissolved 3.50 g of methyl 6-amino-3-methoxy-2-pyrazinecarboxylate. Gaseous ammonia is introduced to saturate the solution with ammonia. The solution is stirred at ambient temperature for 14 hours. The solvent is distilled off from the reaction mixture under reduced pressure to obtain 3.1 g of 6-amino-3-methoxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1684

NMR(DMSO- d_6): 3.79(3H,s), 5.87(2H,brs), 7.30-7.75(3H,m)

Referential Example 5

In a nitrogen atmosphere, 1.50 g of 6-amino-

3-methoxy-2-pyrazinecarboxamide is dissolved in 12 mL of 70% pyridine hydrofluoride under ice-cooling. Then, 0.71 g of sodium nitrite is added thereto at -50°C, and the resulting solution is stirred at 10°C for 1 hour.

- 5 The resulting reaction mixture is stirred at ambient temperature for an additional 1 hour. Thereafter, a mixture of 50 mL of ice water and 100 mL of chloroform is added thereto and the resulting mixture is separated into layers. The organic layer is washed with
- 10 saturated aqueous solution of sodium chloride and saturated aqueous solution of sodium hydrogen carbonate, and dried over anhydrous magnesium sulfate. Then, the solvent is distilled off under reduced pressure to obtain 1.29 g of 6-fluoro-3-methoxy-2-
- 15 pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1706

NMR(DMSO- d_6): 3.95(3H,s), 7.55-8.15(2H,m),
8.39(1H,d,J=8.3Hz)

Referential Example 6

- 20 In 100 mL of methanol is suspended 1.96 g of methyl 5-amino-3-methoxy-2-pyrazinecarboxylate obtained according to the method described in literature (JP-A-50-105675). The resulting suspension is saturated with ammonia by introducing gaseous ammonia thereto at
- 25 -20°C. Then, the solution thus obtained is allowed to react at 95°C for 24 hours in a stainless steel-made closed vessel. After cooling, the solvent is distilled

off under reduced pressure to obtain 1.57 g of 5-amino-3-methoxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1654, 1637

NMR(DMSO- d_6) δ value: 3.82(3H,s), 7.00(3H,brs),
5 7.30(1H,brs), 7.43(1H,s)

Referential Example 7

In a nitrogen atmosphere, 0.5 g of 5-amino-3-methoxy-2-pyrazinecarboxamide is dissolved in 9 mL of 70% pyridine hydrofluoride under ice-cooling. Then,
10 0.23 g of sodium nitrite is added thereto at -70°C , and the resulting solution is heated to a temperature of -10°C in 30 minutes. Further, the solution is stirred at ambient temperature for 30 minutes. A mixture of 30 mL of ice water and 100 mL of chloroform is added
15 thereto and the resulting mixture is separated into layers. The organic layer is washed with saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. The solvent is distilled off under reduced pressure. The residue thus obtained
20 is purified by column chromatography (eluent: chloroform:methanol = 10:1) to obtain 0.37 g of 5-fluoro-3-methoxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1705

NMR(DMSO- d_6) δ value: 3.94(3H,s), 7.65(1H,brs),
25 7.85(1H,brs), 8.12(1H,d,J=8.3Hz)

Example 1

In 3 mL of methanol is suspended 0.6 g of methyl 6-bromo-3-hydroxy-2-pyrazinecarboxylate obtained according to the method described in literature [J. Med. Chem., 1969, 12(2), 285-287]. Then, 6 mL of 25% aqueous ammonia is added thereto and the resulting solution is stirred at ambient temperature for 17 hours. The reaction mixture is adjusted to pH 3 by adding 6 mol/L hydrochloric acid. The solvent is distilled off under reduced pressure. Isopropyl ether and water are added to the residue and filtered to obtain 0.33 g of 6-bromo-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1700, 1665

15 NMR(DMSO- d_6) δ value: 7.50(2H,brs), 8.08(1H,s), 9.95(1H,brs)

Example 2

In 10 mL of dimethylformamide is suspended 0.5 g of 3,5-dihydroxy-1,2,4-triazine-6-carboxylic acid obtained according to the method described in literature (JP-A-54-79292). Then, 2.06 g of N,N'-carbonyldiimidazole is added thereto, and the resulting solution is stirred at ambient temperature for 6 hours. The reaction mixture is cooled with ice, saturated with gaseous ammonia, and then stirred for 15 minutes at the same temperature as above. The deposited crystals are collected by filtration to obtain 0.37 g of 3,5-dioxo-

2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxamide.

IR(KBr) cm^{-1} : 1732, 1710, 1685, 1656

NMR(DMSO- d_6) δ value: 7.75(1H,s), 7.97(1H,s), 12.20-12.80(2H,m)

5 Example 3

In 5 mL of acetic anhydride is suspended 0.5 g of 3-hydroxy-2-pyrazinecarboxamide. The resulting solution is stirred at 110°C for 1 hour. The deposited crystals are collected by filtration to obtain 0.5 g of

10 N^2 -acetyl-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1725, 1695, 1655

NMR(DMSO- d_6) δ value: 2.25(3H,s), 7.53(1H,d,J=4Hz), 7.69(1H,d,J=4Hz), 11.70(1H,brs)

Example 4

15 In 5 mL of 25% aqueous ammonia is suspended 0.25 g of methyl 6-chloro-3-hydroxy-2-pyrazinecarboxylate obtained according to the method described in literature [J. Med. Chem., 285-287 (1969)]. The suspension is stirred at ambient

20 temperature for 1 hour. The deposited crystals are collected by filtration to obtain 0.18 g of 6-chloro-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1652

NMR(DMSO- d_6): 7.22(2H,brs), 7.91(1H,s), 10.40(1H,brs)

Example 5

In 50 mL of tetrahydrofuran is suspended 1.00g of 3-hydroxy-2-pyrazinecarboxamide. Then, 3.5 mL of triethylamine and 1.67 mL of benzoyl chloride are successively added thereto. The resulting solution is stirred at 60°C for 5 hours and cooled. The deposited crystals are collected by filtration. The crystals thus obtained are suspended in a mixture of 8 mL of water and 1 mL of 1 mol/mL hydrochloric acid, stirred at ambient temperature for 30 minutes, and collected by filtration to obtain 0.41 g of N²-benzoyl-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm⁻¹: 1735

NMR(DMSO-d₆) δ value: 7.20-8.40(7H,m), 12.60(1H,brs)

15 According to the same method as above, N²-(2,2-dimethylpropanoyl)-3-hydroxy-2-pyrazinecarboxamide is obtained.

IR(KBr) cm⁻¹: 1725

NMR(DMSO-d₆) δ value: 1.21(9H,s), 7.49(1H,d,J=2Hz),
20 7.95(1H,d,J=2Hz), 14.80(1H,brs)

Example 6

In 0.5 mL of 47% aqueous solution of hydrogen bromide is dissolved 0.05 g of 3-methoxy-2-pyrazinecarboxamide-4-oxide obtained according to the method described in literature [Eur. J. Med. Chem., 15(2), 157-163 (1980)]. The solution is stirred at 45°C

for 2 hours. The deposited crystals are collected by filtration and washed successively with ethanol and diethyl ether to obtain 0.03 g of 3-hydroxy-2-pyrazinecarboxamide-4-oxide.

5 IR(KBr) cm^{-1} : 1695

NMR(DMSO- d_6) δ value: 7.19(1H,d,J=6Hz),
7.56(1H,d,J=6Hz), 7.70(1H,brs), 7.95(1H,brs),
10.75(1H,brs)

Example 7

10 In 4 mL of methanol is suspended 0.19 g of methyl 2-[[[(3-hydroxy-2-pyrazinyl)carbonyl]-amino]acetate. The suspension is saturated with ammonia by introducing gaseous ammonia thereinto under ice-cooling for 30 minutes. The resulting mixture is
15 stirred at the same temperature as above for 1 hour and then at ambient temperature for 15 hours. The solvent is distilled off under reduced pressure. The residue thus obtained is dissolved in a mixture of 4 mL of water and 1 mL of methanol. To the resulting solution
20 is added 0.9 mL of 1 mol/L hydrochloric acid. The deposited crystals are collected by filtration to obtain 0.16 g of N^2 -(2-amino-2-oxoethyl)-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1675

25 NMR(DMSO- d_6) δ value: 3.90(1H,d,J=5Hz), 7.10(1H,brs),
7.40(1H,brs), 7.60-8.40(2H,m), 9.50(1H,brs),
13.0(1H,brs)

Example 8

In a nitrogen atmosphere, 1.51 g of sodium iodide is dissolved in 22 mL of acetonitrile. Then, 1.10 g of trimethylsilyl chloride is added thereto.

5 The resulting solution is stirred at ambient temperature for 20 minutes. Then, 0.43 g of 6-fluoro-3-methoxy-2-pyrazinecarboxamide is added thereto. The resulting solution is stirred at the same temperature as above for 18 hours. Then, a mixture of 10 mL of

10 water and 200 mL of chloroform is added to the reaction mixture, and separated into layers. The organic layer thus obtained is washed successively with 5% aqueous solution of sodium thiosulfate and saturated aqueous solution of sodium chloride, and dried over anhydrous

15 magnesium sulfate. The solvent is distilled off under reduced pressure. The residue thus obtained is purified by column chromatography (eluent: hexane:ethyl acetate = 2:1) to obtain 0.06 g of 6-fluoro-3-hydroxy-2-pyrazinecarboxamide.

20 IR(KBr) cm^{-1} : 1685, 1670, 1656
NMR(CDCl_3): 5.40-7.80 (2H,m), 8.31 (1H,d,J=7.82Hz), 12.33 (1H,s)

Example 9

In 24 mL of concentrated sulfuric acid is

25 suspended 4.00 g of 3-hydroxy-2-pyrazinecarboxamide. To the suspension is added 3.09 g of potassium nitrate under ice-cooling. The resulting mixture is stirred at

40°C for 3 hours. The reaction mixture is poured into 240 mL of water, and the deposited crystals are collected by filtration. The crystals thus obtained are suspended in 80 mL of water and heated under reflux for 30 minutes. After cooling, the crystals are collected by filtration to obtain 2.45 g of 3-hydroxy-6-nitro-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1705, 1685, 1655

NMR(DMSO- d_6) δ value: 8.10(1H,brs), 8.30(1H,brs),
10 8.96(1H,s)

Example 10

In 2.1 mL of water is suspended 0.5 g of 2-aminomalonamide. Under ice-cooling, 0.43 g of ethyl glyoxalate is added to the suspension, which is then stirred for 40 minutes. Then, 0.85 mL of 5 mol/mL sodium hydroxide is added to the resulting suspension, which is then stirred at the same temperature as above for 40 minutes. The reaction mixture is adjusted to pH 12 by adding 1 mol/L sodium hydroxide, and once made into a solution. The solution is then adjusted to pH 2 by adding 6 mol/mL hydrochloric acid. The deposited crystals are collected by filtration and washed successively with water and 50% (w/w) ethanol to obtain 0.15 g of 3,5-dihydroxy-2-pyrazinecarboxamide.

25 IR(KBr) cm^{-1} : 1660

NMR(D_2O) δ value: 6.97(1H,s)

Example 11

In 2.0 mL of water are suspended 0.65 mL of diethyl-2-oxomalonate and 0.5 g of 2-aminomalonamide. Under ice-cooling, 0.85 mL of 5 mol/mL sodium hydroxide is added thereto. The resulting solution is stirred for 40 minutes. Then, at ambient temperature, 2.55 mL of 5 mol/mL sodium hydroxide is added to the solution, which is then stirred for an additional 30 minutes. Ethanol is added to the reaction mixture and the precipitate is collected by filtration to obtain 0.24 g of 3,5-dihydroxy-6-ethoxycarbonyl-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1655, 1735

NMR(D_2O) δ value: 1.17(3H,t,J=7Hz), 4.15(2H,q,J=7Hz)

15 Example 12

In a mixture of 1.0 mL of water and 1.0 mL of ethanol is suspended 0.13 g of 3,5-dihydroxy-6-ethoxycarbonyl-2-pyrazinecarboxamide. At ambient temperature, 0.34 mL of 5 mol/mL sodium hydroxide is added to the suspension, which is then stirred for 16 hours. The reaction mixture is adjusted to pH 2 by adding 1 mol/mL hydrochloric acid. The deposited crystals are collected by filtration and washed with water to obtain 0.07 g of 3,5-dihydroxy-6-carboxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1650

Example 13

In a nitrogen atmosphere, 0.09 g of 5-fluoro-3-methoxy-2-pyrazinecarboxamide is suspended in 3.6 mL of acetonitrile. Then, 0.16 g of sodium iodide and 5 0.11 g of chlorotrimethylsilane are successively added to the suspension, which is then stirred at ambient temperature for 20 hours. Then, 2 mL of water and 40 mL of chloroform are added to the reaction mixture and separated into layers. The organic layer is separated, 10 washed successively with 5% (w/v) aqueous solution of magnesium thiosulfate and saturated aqueous solution of sodium chloride and dried over anhydrous magnesium sulfate. Then, the solvent is distilled off under reduced pressure. The residue thus obtained is 15 purified by column chromatography (eluent: chloroform) to obtain 0.01 g of 5-fluoro-3-hydroxy-2-pyrazinecarboxamide.

IR(KBr) cm^{-1} : 1670

NMR(CDCl_3) δ value: 5.80(1H,s), 7.45(1H,brs),
20 7.93(1H,d,J=7.8Hz), 12.93(1H,s)

Example 14

In 5 mL of ethanol is dissolved 0.1 g of ethyl 5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxylate obtained according to the method 25 described in literature (J. Am. Chem. Soc., 1956, 78, 1258-1259). At ambient temperature, gaseous ammonia is introduced into the solution for 30 minutes to saturate

the solution with ammonia. After allowing the solution to stand at the same temperature as above for 15 hours, the resulting crystals are collected by filtration.

The crystals thus collected are washed with three 5 mL portions of ethanol to obtain 0.05 g of 5-oxo-3-thioxo-2,3,4,5-tetrahydro-1,2,4-triazine-6-carboxamide.

IR(KBr) cm^{-1} : 1654

NMR(DMSO- d_6) δ value: 3.30(4H,brs)

Example 15

10 According to the method described in literature (International Patent Application: WO 98130549), 6-oxo-1,6-dihydro-5-pyrimidinecarboxamide is obtained.

Example 16

15 According to the method described in literature (Chemische Berichte, 1964, 97, 3349-3353), 3-oxo-2,3-dihydro-4-pyridazinecarboxamide is obtained.

Example 17

In 5 mL of ethanol is dissolved 0.06 g of
20 ethyl 5-oxo-4,5-dihydro-1,2,4-triazine-6-carboxylate. At 10°C, the solution is saturated with ammonia by introducing gaseous ammonia thereinto for 20 minutes. The solution is allowed to stand at ambient temperature for 15 hours, and the resulting crystals are collected
25 by filtration. The crystals thus obtained are washed

successively with two 2 mL portions of ethanol and then two 1 mL portion of methanol to obtain 0.03 g of 5-oxo-4,5-dihydro-1,2,4-triazine-6-carboxamide.

IR(KBr) cm^{-1} : 1654

5 NMR(DMSO- d_6) δ value: 3.60(1H,brs), 7.44(1H,brs), 8.39(1H,s), 9.74(1H,brs)

Example 18

In 4 mL of dimethyl sulfoxide is dissolved 0.4 g of methyl 3-hydroxy-2-pyrazinecarboxylate. Then, 10 0.27 g of L-aspartic acid and 0.85 mL of triethylamine are successively added to the solution, which is then stirred at 50°C for 6 hours. The deposited crystals are filtered off, and the filtrate is concentrated under reduced pressure. Then, 2 mL of water and 0.2 mL of 15 methanol are added to the resulting residue. The precipitate formed is collected by filtration to obtain 0.09 g of (2S)-2-([(3-oxo-3,4-dihydro-2-pyrazinyl)-carbonyl]amino)-butanedioic acid.

IR(KBr) cm^{-1} : 1695, 1680, 1665

20 NMR(DMSO- d_6) δ value: 2.83(2H,d,J=5Hz), 4.50-5.00(1H,m), 7.60-8.05(2H,m), 9.95(1H,d,J=9Hz), 12.90(3H,brs)

Example 19

In 5 mL of dimethyl sulfoxide is dissolved 0.42 g of L-alanyl-L-alanine trifluoroacetate. Then, 25 1.07 mL of triethylamine and 0.71 g of methyl 3-hydroxy-2-pyrazinecarboxylate are successively added to

the solution, which is then stirred at 40°C for 17 hours. The solvent is distilled off under reduced pressure, and 2 mL of water was added to the residue thus obtained. The deposited product is collected by
 5 filtration and purified by column chromatography [eluent: chloroform:methanol = 30:1) to obtain 0.035 g of (2S)-2-[(2S)-2-[(3-oxo-3,4-dihydro-2-pyrazinyl)carbonyl]amino}-propanoyl)amino]propanoic acid.

10 IR(KBr) cm^{-1} : 1665, 1675, 1655

NMR(DMSO- d_6) δ value: 1.28(3H,d,J=7Hz),
 1.32(3H,d,J=7Hz), 3.95-4.95(2H,m), 5.1(2H,brs),
 7.71(1H,d,J=3Hz), 7.87(1H,d,J=3Hz), 8.32(1H,d,J=7Hz),
 9.9(1H,brs)


15 INDUSTRIAL APPLICABILITY

An antiviral agent comprising the nitrogen-containing heterocyclic carboxamide derivative represented by general formula [1] or a salt thereof is useful for preventing and treating virus-infections and
 20 especially influenza virus-infections.



CLAIMS


1. An antiviral agent comprising a nitrogen-containing heterocyclic carboxamide derivative represented by the following general formula:



wherein ring A represents a substituted or unsubstituted pyrazine, pyrimidine, pyridazine or triazine ring; R¹ represents O or OH; R² represents a hydrogen atom, an acyl group or a substituted or unsubstituted carbamoylalkyl or carboxyalkyl group; and the broken line represents a single bond or a double bond;

or a salt thereof.


2. An antiviral agent according to Claim 1, wherein said ring A is a substituted or unsubstituted pyrazine, pyrimidine or triazine ring.
3. An antiviral agent according to Claim 2, wherein said ring A is a substituted or unsubstituted pyrazine ring.
4. An antiviral agent according to any one of Claims 1 to 3, wherein the virus is influenza virus, papilloma virus, adenovirus, A type hepatitis virus, B type hepatitis virus, C type hepatitis virus, poliovirus, echovirus, Cocksackie virus, enterovirus,



rhinovirus, rotavirus, Newcastle disease virus, mumps virus, vesicular stomatitis virus or Japanese encephalitis virus.

5. An antiviral agent according to Claim 4, wherein said virus is influenza virus.

6. An nitrogen-containing heterocyclic carboxamide derivative represented by the following general formula:



wherein ring A' represents a pyrazine ring substituted with a halogen atom, a hydroxyl group or an oxido group; R¹ represents O or OH; R² represents a hydrogen atom, an acyl group or a substituted or unsubstituted carbamoylalkyl or carboxyalkyl group; and the broken line represents a single bond or a double bond; or a salt thereof.

7. A nitrogen-containing heterocyclic carboxamide derivative according to Claim 6, wherein said ring A' is a pyrazine ring substituted with a halogen atom; and R² represents a hydrogen atom.