


5

Preparation of hydroxyl-containing compounds from formic esters

10


The invention relates to a process for preparing hydroxyl-containing compounds (1), in particular alcohols, and formic esters of a hydroxyl containing compound (2) by transesterification of a formic ester (also "formate" hereinafter) of a hydroxyl containing compound (1) with a hydroxyl-containing compound (2), in particular alcohol, in the presence of a tertiary amine.

It is known to prepare esters R^1COOR^3 and/or hydroxyl-containing compound, in particular alcohol R^2OH , by transesterification of esters R^1COOR^2 with hydroxyl-containing compound, in particular alcohol R^3OH , according to the following scheme:



25 Transesterification reactions of this type are described, for example, in The Chemistry of Carboxylic Acids and Esters, Ed. S. Patai, 1969, page 103 et seq.; Methoden der organischen Chemie (Houben-Weyl), Volume E5, Georg-Thieme Verlag, Stuttgart 1985, page 702 et seq.
In the case where R^1 is H, the esters are of formic acid.

30



Some selected reactions in which formic esters are produced or result in industrial processes and are subsequently transesterified in accordance with Equation 2 are listed below.

- 5 1. Formic esters are employed as protective group for alcohols in synthetic chemistry. To obtain the unprotected alcohol it is subsequently necessary to remove the protective group quantitatively.

- 10 2. Formic esters occur in industrial processes in which formic acid is used as acidic catalyst, and in which alcohols are produced or are present. Particularly in the case of high-boiling compounds or polymers, these esters are often difficult to remove from the required alcohols by a physical purification process such as recrystallization, extraction or distillation, because their physical properties are very
15 similar to those of the free alcohols. This is why the chemical method of transesterification is used, for example, for purification. The transesterification must take place quantitatively, and catalyst residues must not remain in the product.
 - 20 a) For example, polyoxybutylene glycol formate (polyTHF formate) is formed in the cationic polymerization of tetrahydrofuran (THF) in the presence of formic acid (EP-A-503 386). In order to obtain pure polyoxybutylene glycol (polyTHF), the formate group must be removed quantitatively.
 - 25 b) Another use in polymer chemistry is the polymerization of vinyl formate to poly(vinyl formate). In order to obtain the industrial product poly(vinyl alcohol), the formate groups, which are present in equimolar amounts, must be removed quantitatively.
 - 30 c) In the Prins reaction of olefins with formaldehyde, which is

catalyzed by formic acid, large amounts of the formates of the alcohols which are formed are produced (D.R. Adams, S.P. Bhatnagar, Synthesis (1977) 661-672; J.S. Bajorek, R. Battaglia, G. Pratt, J.K. Sutherland, J. Chem. Soc. Perkin I (1974) 1243-1245). In order to obtain the alcohols, the ester must be cleaved during the workup.

3. In reactions which are carried out in the presence of formaldehyde there is frequently formation, by Cannizzaro reaction of two equivalents of formaldehyde, of one equivalent of methanol and one equivalent of formic acid. Formic acid is also produced by crossed Cannizzaro reaction of aldehydes with formaldehyde. Formic acid produced in this way readily forms, under the reaction conditions, unwanted formic esters as byproduct. In this case too it is necessary to remove the formates either by physical separation methods or by chemical reaction.

The reactions according to Equation 1 and Equation 2 are equilibrium reactions. Starting from the ester R^1COOR^2 , the equilibrium can be shifted in favor of the required product, the ester R^1COOR^3 or the alcohol R^2OH , by either using one initial component in excess or, more preferably, removing one reaction component, the alcohol R^2OH or ester R^1COOR^3 produced, for example by distillation or by crystallization, from the equilibrium.

It is known that addition of a catalyst is necessary to carry out the transesterification according to Equation 1 or Equation 2. Typical transesterification catalysts used in industry include sulfuric acid, p-toluenesulfonic acid, sodium hydroxide solution, sodium alcoholates, aluminum alcoholates, potassium cyanide as well as acidic or basic ion

exchangers. However, these catalysts have a number of disadvantages (see: The Chemistry of Carboxylic Acids and Esters, Ed. S. Patai, 1969, page 103 et seq.; Methoden der organischen Chemie (Houben-Weyl), Volume E5, Georg-Thieme Verlag, Stuttgart 1985, page 702 et seq.).

5

1. The strong bases and acids used as catalyst may lead to numerous unwanted side reactions such as elimination, C-alkylation or polymerization.
- 10 2. If the required product is the alcohol R^2OH , when strongly basic inorganic compounds like sodium methanolate are used as catalyst there are losses of yield, since part of the alcohol remains bound as alcoholate.
- 15 3. Neutralization is necessary for removal of the catalyst and liberation of alcohols from their alcoholates. This produces inorganic salts which have to be removed from the required product. This leads to decomposition of the catalyst. The inorganic salt must be disposed of.
- 20 4. Isolation of the required product is particularly difficult when the required alcohol or ester is high-boiling, crystalline or a polymer. The resulting inorganic salt cannot always be removed without difficulty in this case. Distillation in the presence of an inorganic salt results in decomposition of the required alcohol or ester.
- 25 5. If acidic or basic ion exchangers are employed as catalyst, the reaction cannot be carried out at elevated temperatures because ion exchangers are, as a rule, decomposed at temperatures above
- 30 60-100°C. On the other hand, in this case too, one reaction

component, eg. the alcohol R^2OH , may remain bound to the solid support, resulting in losses of yield. The useful life of ion exchangers is limited. Another disadvantage of heterogeneous catalysts is that they are unsuitable for reaction with compounds of low solubility, especially polymers.

6. If the transesterification is carried out in the presence of water, a competing reaction is observed in the form of hydrolysis of the ester to form alcohol and inorganic salt of the acid, eg. sodium formate in the case of formic acid. Removal of alcohols by distillation in the presence of formic esters is known to lead to losses of yield due to decomposition. In addition, when inorganic alcoholates are used as catalyst, water leads to formation of the free base and thus inactivation of the actual catalyst. For example, sodium hydroxide solution is formed from sodium alcoholate and water and is a far less effective catalyst.

EP-A-0 289 921 describes for example the transesterification of trimethylolalkane formate with methanol. It is possible to use as catalysts alkali metal or alkaline earth metal alcoholates. To obtain the required alcohol TMP it is necessary to remove the catalyst before further workup, either by ion exchanger or by neutralization with an acid. This results in decomposition of the catalyst, and the workup process becomes more costly. An inorganic salt, eg. sodium acetate or chloride, results as byproduct and must be disposed of.

The use of tertiary as transesterification catalysts is known for specific reactions.

DE-A 24 60 039 describes the transesterification of acetoxymethylpyridines

with methanol to hydroxymethylpyridines and methyl acetate in the presence of the tertiary amine triethylamine. The alcohol which is formed, hydroxymethylpyridine, is stabilized by electron-attracting groups. The transesterification disclosed in DE-A 24 60 039 is restricted by the fact
5 that only a selected activated acetic ester with primary alcohol functionality and pyridine substituents can be reacted with methanol.

The low reactivity of non-activated acetates on transesterification with methanol using tertiary amine bases as catalyst has been confirmed
10 experimentally. In the attempt to transesterify a mixture of the mono-, di- and triacetates of 1,1,1-trimethylolpropane (TMP) to TMP and methyl acetate under the conditions described in DE-A 24 60 039 there was no detectable conversion to methyl acetate. The reaction likewise does not succeed on addition of a stronger base with an amidine structure such as
15 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). Only after addition of methanolic tetramethylammonium hydroxide, ie. of OH^- ions, is formation of methyl acetate observed, although with a very unsatisfactory conversion.

20 Another disadvantage of the process described in DE-A 24 60 039 is that the transesterification provides virtually no reaction with ethanol and completely fails with isopropanol.

EP-A-0 168 167 discloses the transesterification of ethyl acetate and
25 propylene carbonate with methanol using a heterogeneous amine-containing catalyst. The catalyst system consists of an organic compound which contains a VA group element, and of an insoluble solid as support. The organic compound containing the VA group element can be an amine or phosphine. The organic compound is, for example, an amidine, guanidine,
30 mono-, di- or trialkylamine.

The yield of methyl acetate from the transesterification of ethyl acetate with methanol after a reaction time of 24 hours is only 27%. No statements are made about the useful life of the heterogeneous support.

- 5 Further disadvantages of the process disclosed in EP-A-0 168 167 are the complicated preparation of the catalyst and the restriction of application, because the reaction is carried out heterogeneously, to specific substrates which are soluble in the reaction medium.
- 10 It is an object of the present invention to provide a process for preparing a hydroxyl-containing compounds, in particular alcohols, and formic esters by transesterification of formic esters without the disadvantages and restrictions described above.
- 15 We have found that this object is achieved by a process for preparing hydroxyl-containing compounds and formic esters by transesterification, wherein formic esters of an organic hydroxyl-containing compound (1) and an organic hydroxyl-containing compound (2) are transesterified in the presence of a tertiary amine as transesterification catalysts to form the
- 20 organic hydroxyl-containing compound (1) and formic esters of the hydroxyl-containing compound (2), where compounds (1) and (2) differ from one another. For example hydroxyl-containing compounds R^2OH and formic esters $HCOOR^3$ are prepared by transesterifying formic esters of the formula $HCOOR^2$ and a compound of the formula R^3OH in the
- 25 presence of a tertiary amine as transesterification catalyst to form formic ester of the formula $HCOOR^3$ and a compound of the formula R^2OH , where R^2 and R^3 are organic radicals which differ from one another. The radicals R^2 and R^3 are organic substituents which are inert under the reaction conditions; they are not critical for the ability of the process to
- 30 be carried out.

Preferred embodiments are described below.

The advantages of the process according to the invention are many.

- 5 1. The reaction takes place with high selectivity and high yield in respect of formic esters.
2. The reaction conditions and temperatures are mild. Very mild bases are used, and thus side reactions are avoided.
- 10 3. There is no loss of catalyst, in contrast to the inorganic and heterogeneous catalysts described above from the prior art. Recovery and recirculation are straightforward.
- 15 4. The reaction mixtures are worked up in a straightforward manner by distillation or crystallization. No neutralization step is necessary.
5. The reaction takes place quantitatively, even in the presence of water.

20

The hydroxyl-containing compound (2) used for the transesterification is preferably added relative to the hydroxyl groups in an amount which is at least stoichiometric relative to the formic ester groups to the formic ester of the hydroxyl-containing compound (1).

25

Transesterification of the formic ester from an organic hydroxyl-containing compound (1), e.g. R^2OH , in particular alcohol, and formic acid relative to the hydroxyl groups with at least equimolar amounts of another organic hydroxyl-containing compound (2) (with respect to the ester group), in
30 particular alcohol R^3OH , preferably in a molar ratio of hydroxyl group to

ester group of 1:10, in particular 1:5, in the presence of a tertiary amine takes place rapidly and with quantitative yield and is not restricted to particular formic esters like the reaction known from EP-A-0 289 921. It is possible, for example, according to one embodiment of the invention to
5 convert the formates of trimethylolpropane with methanol in the presence of triethylamine quantitatively into TMP and methyl formate. Formic esters of secondary, tertiary and aromatic alcohols, such as cyclohexyl formate, tert-butyl formate, phenyl formate, likewise react after addition of methanol and tertiary amine base to give the alcohol and methyl
10 formate.

The organic hydroxyl-containing compound (1) can have one or more hydroxyl groups, preferably 1 to 6, especially 1 to 3 hydroxyl groups. It can be a hydroxyl-containing polymer as well, e.g. a polyvinyl alcohol. It
15 can have the formula R^2OH .

The R^2 radical in the formic ester of the formula $HCOOR^2$ to be transesterified is generally a saturated or unsaturated, branched or unbranched, linear, unsubstituted or substituted hydrocarbon radical which
20 has preferably 1 to 30 carbon atoms, in particular 2 to 20 carbon atoms, particularly preferably 3 to 18 carbon atoms, and which contains, if required, ether group(s), ester group(s), hydroxyl group(s) or ketone group(s), or an aromatic hydrocarbon radical which has preferably 6 to 12 carbon atoms and which is substituted if required, or a cycloalkyl
25 radical which has preferably 3 to 12, in particular 6 to 12, carbon atoms and which is substituted if required, or
a heterocyclic radical which has preferably 3 to 12 carbon atoms and 1 to 3 nitrogen atoms and which is substituted if required, where the substituents are selected from benzyl, phenyl, cyclohexyl, pyridyl,
30 methylpyridyl, hydroxyl, carboxyl, alkyl, the latter with preferably 1 to

12 carbon atoms.

R^2 is particularly preferably propyl, tert-butyl, a trimethylolpropane residue, cyclohexyl, phenyl, a neopentyl glycol hydroxypivalate residue, a
5 polyoxybutylene residue, a ditrimethylolpropane residue or methylpyridyl.

The radical R^2 of the formic ester of the formula $HCOOR^2$ to be transesterified can also be functionalized by other radicals.

10 The tertiary amines used as transesterification catalysts can be compounds of the general structural formula $R^aR^bR^cN$. The radicals R^a , R^b , R^c are identical or different. They are preferably, each independently of the others, a branched or unbranched alkyl radical preferably with 1 to 12, preferably 1 to 6, most preferably 1 to 3, carbon atoms, cycloalkyl
15 radical preferably with 3 to 12, in particular 6 to 12, carbon atoms or aromatic hydrocarbon radical preferably with 6 to 12 carbon atoms, each of which can be substituted if required.

The tertiary amine can also be a heterocyclic tertiary nitrogen compound selected from heterocycloalkane, heterocycloalkene or heteroaromatic
20 compound preferably with 3 to 12 carbon atoms and 1 to 3 nitrogen atoms.

The heterocyclic nitrogen compound preferably used is 1,4-diazabicyclooctane, N,N'-dialkylpiperazine, N-alkylpiperidine,
25 N-alkylaziridine, N-alkylpyrrole, N-methylimidazole or 1,8-diazabicyclo[5.4.0]undec-7-ene.

The alkyl radical in the abovementioned compounds is preferably a branched or unbranched, linear alkyl radical preferably with 1 to 6, most
30 preferably 1 to 4, carbon atoms.

The pK_a of the tertiary amine should be greater than 5, preferably from 7 to 12. The choice of the amine can be adapted to suit the other reaction components without difficulty. The amines which are preferably used are those having a boiling point above the boiling point of the formic ester which is formed and the boiling point of the hydroxyl-containing compound (2) employed, in particular alcohol. If methanol is used for the transesterification, suitable amines have a boiling point above that of methyl formate and that of methanol, in particular trialkylamines such as triethylamine and tributylamine.

10

The transesterification catalyst is preferably unsupported or homogeneous. The transesterification catalyst is preferably added in liquid form.

The tertiary amine can be added in catalytic amounts of from 0.01 to 10 mol%, preferably from 0.1 to 5 mol%, based on formate group as 100 mol%.

An excess of tertiary amine going beyond the amount of formic acid equivalents present has no disadvantages because all the reaction components are recovered after the transesterification is complete. For example, volatile amines such as trimethylamine or triethylamine can easily be removed by distillation from the high-boiling alcohol and be reused. In the event that the amine used is higher-boiling than the alcohol (1) or R^2OH , the procedure can be reversed. The required alcohol (1) or R^2OH is first removed from the tertiary amine by distillation. The remaining tertiary amine need not be worked up because the amine-containing residue can be employed again directly for the reaction. In certain cases, an excess of amine component is advantageous, especially when the amine is suitable as solvent for the reaction.

30

Examples of organic hydroxyl-containing compounds (2) used in the transesterification are alcohols having up to 6 hydroxyl groups, e.g. of the formula R^3OH .

- 5 In a preferred embodiment, the radical R^3 in the alcohol R^3OH used for the transesterification is a saturated or unsaturated, branched or unbranched linear hydrocarbon radical which has preferably 1 to 20 carbon atoms, in particular 1 to 8 carbon atoms, particularly preferably 1 to 3 carbon atoms, an aromatic hydrocarbon radical which has preferably
10 6 to 12 carbon atoms or a cycloalkyl radical which has preferably 3 to 12, in particular 6 to 12, carbon atoms.

Examples of the alcohol R^3OH used for the transesterification are cyclohexanol, methanol, ethanol, propanol, butanol, isopropanol,
15 isobutanol, tert-butanol and benzyl alcohol.

The alcohols R^2OH and R^3OH can be primary, secondary or tertiary. The radicals R^2 and R^3 may carry further OH groups and thus the ester $HCOOR^2$ or $HCOOR^3$ may also have more than one formic ester group.
20

In the following the compound (1) is illustrated by R^2OH and the compound (2) is illustrated by R^1OH . The illustration is valid as well for (1) and (2).

- 25 The range of application of the transesterification depends on the physical properties of the reaction components. It may be advantageous for the alcohol R^2OH and the ester $HCOOR^2$ to be higher-boiling than the alcohol R^3OH used for the transesterification and the ester $HCOOR^3$ produced therefrom, that is to say the boiling points are in the sequence $HCOOR^2$
30 $\approx R^2OH > R^3OH \geq HCOOR^3$.

Alcohols R^3OH suitable to be employed in the transesterification to prepare purer products are those which are more volatile than the hydroxyl-containing compound formed, for example the alcohol R^2OH . R^3 can be a primary, secondary or tertiary functionality. If R^3OH is less
5 volatile than R^2OH , mixtures of products result.

Particularly suitable alcohols are those whose boiling point is higher than that of the corresponding formic ester $HCOOR^3$ which is formed, so that the formic ester formed during the transesterification can be removed
10 from the equilibrium by distillation. It is possible in this way to avoid a large excess of alcohol. Likewise, the alcohol R^3OH should, where possible, not form an azeotrope with the formic ester. However, even in a case of this type, the transesterification is possible by removing the low-boiling alcohol and formic esters together. Alcohols such as methanol,
15 ethanol, propanol, butanol, isopropanol, isobutanol, tert-butanol are particularly suitable. Methanol is preferably employed because in this case there is a large difference in boiling point between alcohol and formic ester. To carry out the reaction quantitatively, R^3OH is employed in amounts which are at least stoichiometric relative to the equivalents of
20 formic acid. It is advantageous in certain cases to increase the amount of the alcohol R^3OH because this then also acts as solvent for the ester $HCOOR^2$ to be transesterified or the alcohol R^2OH which is formed. If the transesterification results in an alcohol R^2OH which is crystalline and of low solubility in the reaction medium, the latter can be isolated after
25 the reaction directly by crystallization from R^3OH or the amine. In the case of a polymeric alcohol, this can be precipitated, for example from R^3OH .

In a preferred embodiment of the invention, a process for preparing
30 alcohols by transesterification of a formic ester in accordance with

Equation 2, where the alcohol R^2OH and the ester $HCOOR^2$ are higher-boiling than the alcohol R^3OH used for the transesterification and the ester $HCOOR^3$ produced therefrom, is made available.

5 The transesterification can be carried out either continuously or batchwise. The temperature used for the reaction depends on the boiling points of the reaction components. The transesterification can be carried out under reduced pressure, under atmospheric pressure or under elevated pressure. In a preferred embodiment with methanol as transesterification reagent and
10 triethylamine as catalyst, the formic ester $HCOOR^2$ is reacted under atmospheric pressure at from 35 to 65°C. In this case, the $HCOOR^2$ /methanol/triethylamine molar ratio for the reaction components employed is 1/1-10/0.0005-0.1, in particular about 1/5/0.05. The amount of methanol can, however, be reduced to the amount corresponding to
15 that of the formate. The methyl formate which is formed is removed continuously in a conventional distillation apparatus. A countercurrent column is suitable for the continuous procedure.

The transesterification can also be carried out in the presence of other
20 inert solvents added to homogenize the reaction components. It is possible to employ, for example, acyclic and cyclic ethers such as tetrahydrofuran, dioxane and diethyl ether or aromatic compounds such as toluene, and esters which are stable to the reaction medium, such as ethyl acetate.

25 Another advantage of the process according to the invention is that the transesterification of formic esters $HCOOR^2$ can also be carried out completely in the presence of water.

In contrast to this, transesterification of formic esters of
30 trimethylolpropane (TMP) with a 1:1 mixture of methanol and water

based on volume with 10 mol% sodium hydroxide solution as catalyst does not take place quantitatively. Sodium formate is formed by hydrolysis, and the maximum conversion observed even on prolonged treatment with methanol is only 75% based on TMP formates. In
5 addition, the sodium formate which is formed interferes with the isolation of TMP by distillation.

The range of applications of the transesterification which has been found is to be illustrated by examples.

10

The transesterification, according to the invention, of the formic esters takes place selectively in respect of the formate functionality in the presence of other ester functionalities. Mono- and diformic esters of neopentyl glycol hydroxypivalate (NHP), which are byproducts of the
15 industrial production of NHP as described in, for example, DE 19 58 463 or DE 22 33 357, can be reacted selectively. Selective transesterification of these mono- and diformic esters of NHP is carried out according to the invention with a mixture of methanol and triethylamine to give NHP and methyl formate without cleavage of the central
20 ester group of the NHP. NHP is obtained without losses after removal of excess methanol/triethylamine. By contrast, when typical "inorganic" transesterification catalysts such as sodium methanolate are used there is found to be formation of methyl hydroxypivalate and neopentyl glycol.

25 Formic esters of polymeric compounds can also be transesterified very well. For example, polytetrahydrofuran (polyTHF) which is esterified by formic acid on the alcoholic end groups (a product of the polymerization of THF with W catalysts and formic acid (EP-A-0 503 394)) can be converted with methanol and triethylamine as catalyst quantitatively into
30 polyTHF by one embodiment. The purified polyTHF is isolated in a

straightforward manner by removing methanol and triethylamine by distillation.

Surprisingly, when the inorganic base sodium hydroxide solution is
5 replaced by a tertiary amine such as triethylamine the formates of TMP
are converted almost quantitatively into TMP. Once again partial
hydrolysis to formic acid, which is present in solution as triethyl-
ammonium formate, is observed. However, the reaction also takes place
quantitatively at a $\text{pH} < 7$. This means that catalysis by a tertiary amine
10 such as triethylamine is superior to catalysis by sodium hydroxide
solution. In addition, the reaction catalyzed by a tertiary amine has
advantages owing to simplified workup because alcohols can be distilled
undecomposed even in the presence of organic formic acid salts.

15 The invention is explained in detail by means of the following examples.

Comparative Example 1

This comparative example shows the low reactivity of unactivated acetates
20 on transesterification with methanol.

132 g (0.75 mol) of a mixture of trimethylolpropane (TMP), TMP
monoacetate, TMP diacetate and TMP triacetate (GC %
areas = 32/50.5/16/1.5) are dissolved in 240 g of methanol (7.5 mol).
25 38.5 g (0.38 mol) of triethylamine are added and the mixture is heated
to reflux. A temperature of 65°C is observed at the top of the column,
and methyl acetate is undetectable. The percentage ratio of acetates in the
bottom remains unchanged.

30 All the methanol and triethylamine are distilled out, and 200 ml of

methanol and 22 g of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) are added to the reaction mixture. No conversion of the acetates is observed after heating to reflux. The temperature at the top of the column is 65°C.

- 5 Finally, 20 g of a 25% strength aqueous tetramethylammonium hydroxide solution are added to the reaction mixture. The boiling point at the top of the column falls, and a mixture of methanol and methyl acetate can be distilled out. The conversion of the TMP acetates emerges from the analysis by gas chromatography: TMP 43%, TMP monoacetate 51%,
10 TMP diacetate 6%, TMP triacetate 1%.

Example 1a)

- 15 Preparation of a mixture of the formic esters of 1,1,1-trimethylolpropane (TMP).
134 g of TMP are refluxed with 46 g of formic acid for 5 hours. The water which has formed is then distilled out. Finally, a mixture of TMP and the mono-, di- and triformates of TMP is distilled out. GC analysis
20 reveals a ratio of 49/40/10/1.

Example 1b)

- 81 g (0.5 mol) of the mixture obtained from a) are dissolved in 64 g of
25 methanol. 4 g of triethylamine are added and the mixture is stirred at an internal temperature of 40°C. GC analysis of the reaction mixture after 15 minutes reveals a composition of 80/18/2/-. The mixture is then heated, and methyl formate is distilled out until conversion is quantitative after one hour.

Example 2

The procedure is as in Example 1b). In place of triethylamine, 4 g of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) are used as catalyst. After 5 minutes, 95% TMP and 5% TMP monoformate are present. Quantitative conversion is achieved by distilling out methyl formate.

Example 3

The procedure is as in Example 1b). The starting materials for the transesterification are 81 g of TMP formate mixture (Example 1a)), 80 g of methanol and 28.5 g of N,N'-dimethylpiperazine. After 2 hours, TMP formates are no longer detectable in the reaction mixture.

15

Example 4

The procedure is as in Example 1b). The starting materials are 81 g of TMP formate mixture (Example 1a)), 80 g of methanol and 28.5 g of N-ethylpiperidine. After 2 hours, TMP formates are no longer detectable in the reaction mixture.

Example 5

25

The procedure is as in Example 1b). The starting materials are 246 g of TMP formate mixture (Example 1a)), 370 g of ethanol and 76 g of triethylamine. The temperature at the top of the column is 54 - 55°C (boiling point of ethyl formate). Distillation is continuous and, after 1 hour, the temperature increases to 78°C. The distillate contains 88.7%

30

ethyl formate, 10.2% ethanol and 1.1% triethylamine. The conversion of TMP formates is 100%.

Example 6

5

The procedure is as in Example 5. The starting materials are 246 g of TMP formate mixture (Example 1a)), 480 g of isopropanol and 76 g of triethylamine. The temperature at the top of the column is initially 72°C. Distillation is continuous and the temperature rises to 82°C within 4
10 hours. The distillate contains 40% isopropyl formate, 35% isopropanol and 25% triethylamine. The conversion of the TMP formates is 100%.

Example 7

15 The procedure is as in Example 5. The starting materials are 66.5 g of TMP formate mixture (Example 1a)), 148 g of tert-butanol and 19 g of triethylamine. The temperature at the top of the column is initially 77°C. Distillation is continuous and the temperature rises to 82°C within 1
hour. The distillate contains 15% tert-butyl formate, 70% tert-butanol and
20 15% triethylamine. The conversion of the TMP formates is 100%.

Example 8

42 g of a mixture of cyclohexanol and cyclohexyl formate (65/35) are
25 dissolved in 64 g of methanol. 20 g of triethylamine are added and the mixture is heated to boiling, and methyl formate is distilled out continuously through a column. The cyclohexanol to cyclohexyl formate ratio after 4 hours is 97.5/2.5.

30 Example 9

The procedure is as in Example 8 but with the following starting materials: 15 g of a mixture of tert-butanol and tert-butyl formate (97.8/1.2), 32 g of methanol, 10 g of triethylamine. The ratio of tert-butanol to tert-butyl formate after 4 hours is 99.1/0.9.

5

Example 10

The procedure is as in Example 8 but with the following starting materials: 82.4 g of a mixture of neopentyl glycol hydroxypivalate (NHP) and neopentyl glycol hydroxypivalate mono- and diformates (94.5/3.2/2.3), 10 65 g of methanol, 4 g of triethylamine. The mixture is stirred at 40°C for 2 hours. The formation of methyl formate is established by gas chromatography. The NHP components are present in a ratio of 98.8/1.2/-. Complete conversion of the NHP formates is achieved by 15 distilling out methyl formate and methanol. No other additional components can be detected.

Comparative Example 2

20 The procedure is as in Example 1b). The starting materials are 243 g of TMP formate mixture (Example 1a)), 240 g of methanol, 135 g of water and 6 g of NaOH. The pH of the solution decreases during the reaction from 9.8 to 5.3. At a bottom temperature of 65°C, methyl formate is distilled out at a temperature at the top of the column of 35°C. The 25 temperature at the top increases to 65°C after 1 hour. The composition of the reaction solution reveals a ratio of TMP and mono-, di- and triformates of 83/16/1/-.

Example 11

30

The procedure is as in Comparative Example 2. The starting materials are 243 g of TMP formate mixture (Example 1a)), 240 g of methanol, 135 g of water and 15 g of triethylamine. The pH of the solution falls during the reaction (3 hours) from 10.2 to 5.3. At a bottom temperature of
5 65°C, methyl formate is distilled out at a temperature at the top of the column of 35°C. The temperature at the top increases to 65°C after 3 hours. The composition of the reaction solution reveals a ratio of TMP and mono-, di- and triformates of 96.5/3.5/-/-.

10 Example 12

PolyTHF monoformate (prepared by polymerization of THF catalyzed by $H_3PW_{12}O_{40}$ in the presence of formic acid as described in EP-A-503 394) employed as precursor had the following characteristics:

15 $M_n = 1030$ (from OH value and SV);
SV = 55 mg KOH/g; OH value = 54 mg KOH/g.

500 g of polyTHF monoformate 1030 (0.5 mol) were dissolved in 480 g of methanol (15 mol) in a distillation apparatus. 5 g of triethylamine
20 (0.05 mol; previously distilled under argon) were added and the reaction mixture was then heated to 67°C (bottom temperature). Methyl formate (boiling point 32°C) which was formed was distilled out through a Vigreux column until the temperature at the top was 65°C. To remove excess methanol, catalyst and low molecular weight oligomers, the residue
25 was concentrated at 40°C under reduced pressure and then subjected to short-path distillation at 230°C/0.5 mbar.

The polyTHF remaining as distillation residue (440 g) had the following characteristics:

30 $M_n = 1170$ (OH value);

OH value = 96 mg KOH/g; SV = 0.3 mg KOG/g;
residual N content = 5 ppm (determination method: chemical
luminescence; detection limit about 1 ppm; ref.: "Quantitative Organische
Elementaranalyse", Friedrich Ehrenberger, VCH Weinheim, 1991, page
s 382 et seq.; apparatus used: DN 1000 from Dohrmann)
The conversion calculated from the SV is 99.5%;
Abbreviations: OH value = hydroxyl value; SV = saponification value;
M_n = number average molecular weight

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



We claim:

1. A process for preparing hydroxyl-containing compounds and formic
5 esters by transesterification, wherein formic esters of an organic
hydroxyl-containing compound (1) and an organic hydroxyl-containing
compound (2) are transesterified in the presence of a tertiary amine
as transesterification catalyst to form the organic hydroxyl-containing
compound (1) and formic esters of the hydroxyl-containing compound
10 (2), where compounds (1) and (2) differ from another.
2. A process as claimed in claim 1, wherein the hydroxyl-containing
compound (2) to be used for the transesterification is added relative
to the hydroxyl groups in at least the stoichiometric amount relative
15 to the formic ester of the hydroxyl-containing compound (1) based on
the formic ester groups.
3. A process as claimed in claim 1, wherein in the formic ester
employed is a hydroxyl-containing compound (1) formic ester of
20 polytetrahydrofuran, in particular the terminal monoformic ester, or
formic ester of trimethylolpropane, or formic ester of neopentyl
glycol hydroxypivalate or formic ester of a polyvinyl alcohol, with
preferably all the formic acid groups being removed.
- 25 4. A process as claimed in any of claims 1 to 3, wherein the tertiary
amine used for the transesterification is

a tertiary amine a) of the general structural formula $R^aR^bR^cN$ where
the radicals R^a , R^b , R^c can be identical or different and are each
30 independently of the others a branched or unbranched alkyl radical





which has preferably 1 to 12, more preferably 1 to 6, in particular 1 to 3, carbon atoms, a cycloalkyl radical which has preferably 3 to 12, in particular 6 to 12 carbon atoms or an aromatic hydrocarbon radical which has preferably 6 to 12 carbon atoms, and which can be substituted if required, or

b) a heterocyclic tertiary nitrogen compound selected from heterocycloalkane, heterocycloalkene or a heteroaromatic compound which has preferably 3 to 12 carbon atoms and 1 to 3 nitrogen atoms.

5. A process as claimed in claim 4, wherein the heterocyclic nitrogen compound used for the transesterification is 1,4-diazabicyclooctane, N,N'-dialkylpiperazine, N-alkylpiperidine, N-alkylaziridine, N-alkylpyrrole, N-methylimidazole or 1,8-diazabicyclo[5.4.0]undec-7-ene.
6. A process as claimed in any of claims 1 to 5, wherein the tertiary amine is employed in an amount of from 0.01 to 10 mol%, preferably in an amount of from 0.1 to 5 mol% based on the formic ester groups as 100 mol%.
7. The process as claimed in any of claims 1 to 6, wherein the hydroxyl-containing compound used for the transesterification is one having 1 to 6 OH groups, preferably monohydric, dihydric or trihydric, in particular monohydric, alcohols.
8. A process as claimed in any of claims 1 to 7, wherein the organic hydroxyl-containing compound (2) used in the transesterification is an alcohol of the formula R^3OH where R^3 is a saturated or unsaturated, branched or unbranched, linear hydrocarbon radical which has



THIS SHEET HAS BEEN
REPLACED

preferably 1 to 20 carbon atoms, in particular 1 to 8 carbon atoms,
particularly preferably 1 to 3 carbon atoms, an aromatic hydrocarbon
radical which has preferably 6 to 12 carbon atoms or a cycloalkyl
radical which has preferably 3 to 12, in particular 6 to 12, carbon
atoms.

9. A process as claimed in any of claims 1 to 8, wherein the hydroxyl-
containing compound (2) used in the transesterification is more
volatile than the hydroxyl-containing compound (1) formed in the
transesterification.

10. A process as claimed in any of claims 1 to 9, wherein the hydroxyl-
containing compound (2) used in the transesterification has a higher
boiling point than the boiling point of the formic ester formed.

This Sheet Has Been Replaced

