


TETRAHYDROISOQUINOLINE SULFONAMIDE DERIVATIVES, THE PREPARATION THEREOF, AND THE USE OF THE SAME IN THERAPEUTICS

5 The present invention relates to sulfonamide derivatives, to the preparation thereof and to the therapeutic use thereof, in particular in the treatment of disorders that are improved by modulation of the histamine H₃ receptor, such as obesity, diabetes and central nervous system diseases such as vigilance and sleep disorders.

10 Consequently, a first subject of the present invention is the compounds corresponding to formula I



in which:

15

n can represent a value between 1 and 6;

-(C)_n- represents a -C₁₋₆ alkylidene group optionally substituted with 1 to 4 substituents chosen from a halogen atom, and a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino or C₁₋₃ alkoxy group;

20

R1 represents

- a hydrogen atom,
- a C₁₋₆ alkyl group;

25

R2 represents

- a hydrogen atom,
- a C₁₋₆ alkyl or C₃₋₆ cycloalkyl group optionally substituted with 1 to 4 substituents chosen from a halogen atom, a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy or C₃₋₆ cycloalkyl group, a monocyclic heteroaryl such as a thienyl, furyl or pyrrolyl, or an aryl, such as a phenyl or a naphthyl; the aryl being optionally substituted with 1 to 4 substituents chosen from a halogen atom, a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl or C₁₋₃ alkoxy group or a C₁₋₃ alkylidenedioxy

30

35

group;

B represents

- **NR₃R₄,**
 - R₃ and R₄ represent, independently of one another, a C₁₋₆ alkyl group, or a hydrogen atom; or
 - R₃ and R₄ together represent a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, a C₁₋₃ alkylidene-O-C₁₋₃ alkylidene group, or a C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene group where R₅ represents a hydrogen atom, or a C₁₋₃ alkyl or C₁₋₆ alkylcarbonyl group, it being possible for these C₁₋₃ alkyl and C₁₋₆ alkylcarbonyl groups to be substituted with a halogen atom, or a hydroxyl, C₁₋₃ alkoxy, nitro, cyano or amino group; or
- an aminocycle, linked via a carbon to the group -NR₁-(C)_n-, such as aziridine, azetidine, pyrrolidine, piperidine or morpholine;

the groups R₃ and R₄ and also the aminocycle being optionally substituted with 1 to 4 substituents chosen from a phenyl, a benzyl, a halogen atom, and a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl or C₁₋₃ alkoxy group; and the nitrogen atom optionally substituted with a C₁₋₃ alkyl.

In the context of the present invention, the term:

- "C_{x-z}" is intended to mean a carbon-based chain that may contain from x to z carbon atoms; for example, C₁₋₃ indicates a carbon-based chain that may contain from 1 to 3 carbon atoms;
- "alkyl" is intended to mean a linear or branched, saturated aliphatic group; for example, a C₁₋₄ alkyl group represents a linear or branched, saturated carbon-based chain containing from 1 to 6 carbon atoms, more particularly a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, etc. radical; the term "C_{x-y} alkylidene" denoting a divalent, linear or branched C_{x-y} alkyl group; the term "C₂₋₈ alkenylidene" denoting a divalent, linear or branched, unsaturated C_{x-y} alkyl group;
- "C_{x-y} alkoxy" is intended to mean an alkyloxy group comprising a linear or branched, saturated aliphatic chain, containing x to y carbon atoms;
- "halogen atom" is intended to mean a fluorine, a chlorine, a bromine or an iodine;
- "C₁₋₃ monoalkylamino" is intended to mean an amino monosubstituted with a C₁₋₃ alkyl group;
- "C₂₋₆ dialkylamino" is intended to mean an amino disubstituted with two C₁₋₃ alkyl groups;

- "C₁₋₂ perhaloalkyl" is intended to mean a C₁₋₂ alkyl group in which all the hydrogen atoms are substituted with halogen atoms;
- "C₁₋₃ haloalkyl" is intended to mean a C₁₋₃ alkyl group in which at least one hydrogen atom is substituted with a halogen atom.

5

The compounds of formula I can contain one or more asymmetrical carbon atoms. They can therefore exist in the form of enantiomers or of diastereoisomers. These enantiomers and diastereoisomers, and also mixtures thereof, including racemic mixtures, are part of the invention.

10

The compounds of general formula I can be in the form of free bases or of addition salts with acids, which are also part of the invention. According to the present invention, these salts comprise those with pharmaceutically acceptable acids, but also those with inorganic or organic acids that allow a suitable separation or

15 crystallization of the compounds of formula I. These salts can be prepared, according to methods known to those skilled in the art, for example, by reaction of the compound of formula I in the form of a base with the acid in an appropriate solvent, such as an alcoholic solution or an organic solvent, and then separation of the media that contains it by evaporation of the solvent or by filtration.

20

The compounds of formula I can also exist in the form of hydrates or of solvates, i.e. in the form of associations or combinations with one or more water molecules or with a solvent. Such hydrates and solvates are also part of the invention.

25

Furthermore, in the context of the present invention, the term "protective group Pg" is intended to mean a group that makes it possible, firstly, to protect a reactive function such as a hydroxyl or an amine during a synthesis and, secondly, to regenerate the intact reactive function at the end of synthesis. Examples of protective groups and also the methods of protection and deprotection are given in "Protective groups in

30 Organic Synthesis 3rd Ed.", Greene and Wuts (John Wiley & Sons, Inc., New York, 1999).


A subject of the present invention is also the compounds chosen from the following subgroups, in which:

35

- n is equal to 2, 3 or 4; and/or
- R₁ represents a hydrogen atom or a C₁₋₂ alkyl group; and/or
- R₂ represents a hydrogen atom, or a C₁₋₄ alkyl or C₅₋₆ cycloalkyl group optionally substituted with 1 to 4 substituents chosen from a phenyl, and a C₃₋₆ cycloalkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl or C₁₋₃ alkoxy group; the phenyl being optionally

substituted with 1 to 4 substituents chosen from a halogen atom, a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl or C₁₋₃ alkoxy group or a C₁₋₃ alkylidenedioxy group; and/or

- 5 - **B** represents • NR₃R₄,
- R₃ and R₄ represent, independently of one another, a C₁₋₄ alkyl group; or
 - when R₃ and R₄ together represent a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, a C₁₋₃ alkylidene-O-C₁₋₃ alkylidene group or a C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene group, **B** represents a group:





or

- an aminocycle linked via a carbon to the group -NR₁-(C)_n, such as aziridine, azetidine, pyrrolidine, piperidine or morpholine;

the groups R₃, R₄ and R₅ and also the aminocycle being optionally substituted; and, more specifically, the subgroup where at the same time n, R₁, R₂ and **B** are as defined above.

More particularly, when **B** represents NR₃R₄ and R₃ and R₄ together form a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, a C₁₋₃ alkylidene-O-C₁₋₃ alkylidene group or a C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene group, or when **B** represents an aminocycle, then **B** is chosen from the following groups:



Another subject of the present invention concerns the following compounds:


1. N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
2. (+/-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
3. N-[3-(diethylamino)propyl]-2-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
4. N-[3-(diethylamino)propyl]-N-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
5. 2-benzyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
6. 2-(cyclopropylmethyl)-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
7. 2-(cyclohexylmethyl)-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
8. (+/-)-N-[3-(2-methylpiperidin-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
9. N-[3-(3,6-dihydropyridin-1(2*H*)-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
10. N-[3-(diethylamino)propyl]-2-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
11. N-[3-(diethylamino)propyl]-2-(2-thienylmethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
12. N-[3-(diethylamino)propyl]-2-(3-thienylmethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
13. N-[3-(2,5-dihydro-1*H*-pyrrol-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
14. 2-cyclohexyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
15. (+/-)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
16. N-[3-(2,5-dihydro-1*H*-pyrrol-1-yl)propyl]-2-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
17. N-[3-(4-benzylpiperazin-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
18. N-(3-pyrrolidin-1-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
19. N-(3-morpholin-4-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
20. N-[3-(dimethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
21. 2-(cyclohexylmethyl)-N-(3-pyrrolidin-1-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
22. (+/-)-2-(cyclopropylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;

23. (+/-)-2-benzyl-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
24. (+/-)-2-(4-isopropylbenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
- 5 25. (+/-)-2-(1,3-benzodioxol-5-ylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
26. (+)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
27. (-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
- 10 28. (+)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
29. (-)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide;
- 15 30. (+/-)-2-(4-bromobenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide
31. (+/-)-2-(2,5-dimethoxybenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide
32. (+/-)-2-(2-methylbutyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide
- 20 33. (+/-)-2-(3-methoxybenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide
- and
34. (+/-)-2-(3,5-dimethyl-benzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide.
- 25

A second subject of the present invention is processes for preparing the compounds of formula I according to the invention.

- 30 Thus, the compounds of formula I can be prepared according to the process represented in Scheme 1.


Scheme 1



According to the process of Scheme 1, the compounds of formula I, in which R2 is other than a hydrogen atom, are prepared by amminative reduction, by reacting a secondary amine of formula I, in which R2 represents H, with an aldehyde or a ketone of formula III, where R6 and R7, after reaction, together form R2 as defined in formula I. The compounds of formula I where R2 represents a hydrogen atom can be obtained by deprotection of the compounds of formula II, according to conventional methods known to those skilled in the art. For example, the compounds of formula II, when Pg is a trifluoroacetyl group, can be deprotected in the presence of a base such as, for example, sodium carbonate, potassium carbonate, ammonia or barium hydroxide in a protic solvent, such as water or methanol or a mixture of these solvents, at a temperature between 0 and 100°C. Alternatively, the deprotection of the compounds of formula II, when Pg is a trifluoroacetyl group, can be carried out in the presence of an acid such as, for example, hydrochloric acid in a protic or aprotic solvent, such as water, methanol, ethanol or ethyl acetate or a mixture of these solvents, at a temperature between 0 and 100°C. Illustrations of the process are given in the examples.

The starting compounds of formula II can be prepared according to Scheme 2 or can be synthesized by conventional methods known to those skilled in the art.

Scheme 2




According to this scheme, the compounds of formula II, in which n, R1 and B are as defined in formula I, can be prepared by reaction of an amine of formula V, in which R1 and B are as defined in formula I, with a sulfonyl chloride of formula IV, in which Pg represents an appropriate protective group such as, for example, a trifluoroacetamide, so as to form a derivative of sulfonamide type of formula II, according to conventional methods known to those skilled in the art, for example, the reaction can be carried out in a protic or aprotic solvent, such as tetrahydrofuran, dichloromethane, ethyl acetate, N,N-dimethylformamide or acetonitrile or a mixture of these solvents, at a temperature of between 0 and 100°C, in the presence of a base

such as, for example, potassium carbonate, sodium carbonate, diisopropylethylamine or triethylamine.

Alternatively, the compounds of formula II can be prepared by a Mitsunobu type reaction, according to Scheme 3.

5

Scheme 3




- 10 According to this alternative, a sulfonamide of formula VII, in which R1 is as defined in formula I, is reacted with an amino alcohol of formula VIII, in which n and B are as defined above. The reaction can be carried out conventionally in the presence of Mitsunobu reagents, such as an azo derivative, for example diethylazodicarboxylate, diisopropylazodicarboxylate, di-tert-butylazodicarboxylate, 1,1'-
- 15 (azodicarbonyl)dipiperidine or *N,N,N',N'*-tetramethylazodicarboxamide, and a phosphine, for example triphenylphosphine or tributylphosphine. The reaction can be carried out in an aprotic solvent, such as tetrahydrofuran or dioxane or a mixture of these solvents, at a temperature between 0 and 100°C, to give the compound of formula II. The sulfonamide of formula VII in which R1 is as defined in formula I can
- 20 be prepared by reaction of an amine of formula VI, in which R1 is as defined in formula I, with a sulfonyl chloride of formula IV, in which Pg represents an appropriate protective group such as, for example, a trifluoroacetamide, according to conventional methods known to those skilled in the art, for example, the reaction can be carried out in a protic or aprotic solvent, such as tetrahydrofuran,
- 25 dichloromethane, ethyl acetate, *l*a *N,N*-dimethylformamide or acetonitrile or a mixture of these solvents, at a temperature between 0 and 100°C, in the presence of a base such as, for example, potassium carbonate, sodium carbonate, diisopropylethylamine or triethylamine.

The starting compounds IV and the amines of formulae V and VIII are directly commercially available, can be synthesized by conventional methods known to those skilled in the art, or are known in the literature.

- 5 For example, the diamines of formula V, in which n is equal to 3, can be prepared according to Scheme 4.

Scheme 4




10

- According to this process, the compounds of formula V, in which n is equal to 3, R1 represents a hydrogen atom and B represents an amine group, can be prepared by means of an addition reaction of an amine of formula X, in which R3 and R4 are as defined above, with the acrylonitrile of formula IX, so as to form a derivative of aminonitrile type of formula XI, according to conventional methods known to those skilled in the art, followed by reduction of the nitrile. The reduction can be carried out according to methods known to those skilled in the art, for example in the presence of diisobutylaluminum hydride at a temperature between -70°C and 40°C in an aprotic solvent such as dichloromethane or toluene or a mixture of these solvents; the reduction can also be carried out in the presence of a reducing agent, such as hydrogen, in the presence of a catalyst such as platinum, palladium or Raney nickel, in a solvent such as methanol or ethyl acetate, so as to give the compound of formula V, in which n is equal to 3, R1 represents a hydrogen atom and B represents an amine group. The compounds of formula V, in which R1 represents a C_{1-6} alkyl group, can be prepared by alkylation of the compound of formula V obtained above, according to conventional methods known to those skilled in the art.
- 15
- 20
- 25

- A subject of the present invention is also the compounds of formula II, when Pg represents a protective group or a hydrogen atom, as intermediates for the preparation of the compound of formula I.
- 30

The following examples illustrate the processes and techniques suitable for the preparation of this invention, without, however, limiting the scope of the claim.

- 35 **Example 1- N-[2-(1-Methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride**



1.1- (+/-)-N-[2-(1-Methylpyrrolidin-2-yl)ethyl]-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride

5

A solution of 5.00 g (0.015 mol) of 2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonyl chloride and 2.00 g (0.015 mol) of (+/-)-2-(1-methyl-pyrrolidin-2-yl)-ethylamine, in 50 ml of dichloromethane, is stirred overnight at ambient temperature. The solution is concentrated to dryness. The solid formed is purified by silica gel column chromatography, with a dichloromethane/methanol (97:3) mixture used as eluant, to give 4.30 g of the desired product in the form of a white solid.

10

Yield: 62%

Mp = amorphous

15

1.2- (+/-)-N-[2-(1-Methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride

A solution of 4.30 g (0.0094 mol) of 2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamido-[2-(1-methylpyrrolidin-2-yl)ethyl] hydrochloride in 50 ml of methanol saturated with hydrogen chloride is heated for twelve hours at 60°C. The mixture is cooled and the solid that has formed is filtered off, washed with methanol and dried. 2.00 g of the desired product are obtained as a white solid.

20

25

Yield: (65%)

Mp = 209-212°C

30

¹H-NMR (DMSO-d₆) δ(ppm): 7.9 (1H, t), 7.7 (2H, d), 7.5 (1H, d), 4.3 (2H, s), 3.6 (5H, m), 3.1 (3H, m), 2.9 (2H, m), 2.7 (3H, m), 2.1 (1H, m), 1.9 (3H, m), 1.6 (2H, m), 2.7 (3H, s); 2.9 (2H, t).

1.3- (+) or (-)-N-[2-(1-Methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride

The compound obtained above in 1.2 is separated by chiral-phase preparative chromatography, to give its enantiomers. Specifically, the separation of 15.00 g of


35

(+/-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride is carried out with a PROCHROM LC50 preparative HPLC system, with a CHIRALPACK AD stationary phase and a mobile phase formed from isohexane/ethanol/methanol (80%:10%:10%)+0.2% of diethylamine, to give 5.39 g of the dextrorotatory product, in the form of a white powder, with a chiral-phase enantiomeric purity of 99.67%, and 4.89 g of the levorotatory product, in the form of a white solid, with a chiral-phase enantiomeric purity of 99.48%. The two products are converted into their corresponding hydrochloride by a treatment with isopropanol saturated with hydrogen chloride.

Dextrorotatory enantiomer: Mp = 114-117°C; $[\alpha]_D^{20} = +16$ (c=0.5, methanol)

Levorotatory enantiomer: Mp = 115-117°C; $[\alpha]_D^{20} = -16$ (c=0.5, methanol)

Example 2- (+/-)-2-(Cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide oxalate (1:2)



0.32 g (0.0003 mol) of palladium-on-charcoal at 10% is added to a solution of 2.02 g (0.0063 mol) of (+/-)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamido-[2-(1-methylpyrrolidin-2-yl)ethyl] and 0.70 g (0.0063 mol) of cyclohexanecarboxaldehyde in 100 ml of methanol. The solution is hydrogenated for 24 hours in a Paar hydrogenator at a pressure of 45 Psi. The catalyst is removed by filtration and the filtered solution is evaporated to dryness. The crude oil obtained (2.90 g) is purified by silica gel column chromatography, with a dichloromethane/methanol (95:5) mixture used as eluant. The desired product (1.62 g; 62%) is obtained in the form of an oil.

The above oil is dissolved in 20 ml of ethanol, and then 0.77 g (0.0086 mol) of oxalic acid dissolved in 15 ml of ethanol is added. The precipitate is filtered off and washed with cold ethanol. 2.01 g of the desired product are obtained in the form of a white solid.

Yield: 86%

Mp = 142-147°C

¹H-NMR (DMSO-d₆) δ(ppm): 7.5 (2H, m), 7.29 (1H, m), 3.84 (2H, s), 3.35 (2H, m), 3.1 (1H, m), 2.9 (1H, m), 2.8 (2H, s), 2.7 (1H, m), 2.6 (3H, s), 2.5 (2H, m), 2.1-1.4

(13H, m), 1.15 (4H, m), 0.8 (2H, m).

Example 3- (+)-2-(Cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide oxalate (1:1.5)

5

According to a process similar to Example 2, with 1.00 g (0.0031 mol) of (+)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide and 0.35 g (0.0031 mol) of cyclohexanecarboxaldehyde in 50 ml of methanol as starting product, 0.46 g of base is obtained, which base is converted into the corresponding

10

Yield: 20%

Mp = 134-140°C

$[\alpha]_D^{20} = +12$ (c=0.5, methanol)

15

Example 4- (-)-2-(Cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide oxalate (1:2)

According to a process similar to Example 2, with 1.00 g (0.0031 mol) of (-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide and 0.35 g (0.0031 mol) of cyclohexanecarboxaldehyde in 50 ml of methanol as starting product, 0.90 g of base is obtained, which base is converted into the corresponding dioxalate hydrate, as a white solid.

20

Yield: 27%


Mp = 133-138°C

$[\alpha]_D^{20} = -8$ (c=0.5, methanol)

25

Example 5- 2-Benzyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide oxalate (1:2)

30



5.1- N-[2-(3-Diethylaminopropyl)-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride

35

A solution of 1.00 g (0.0031 mol) of 2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonyl chloride and 0.61 g (0.0047 mol) of N,N-diethyl- N-

aminopropylamine in 25 ml of dichloromethane is stirred overnight at ambient temperature. The solution is concentrated to dryness and the oil formed is purified by silica gel column chromatography, with a dichloromethane/methanol (97:3) mixture used as eluant. 1.27 g are obtained.

5

Yield: 97%

Mp = Oil

5.2- N-[3-(Diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride

10

A solution of 5.39 g (0.013 mol) of N-[2-(3-diethylaminopropyl)-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride, dissolved in 60 ml of methanol saturated with hydrogen chloride, is heated for twelve hours at 60°C. The mixture is cooled and the solid that has formed is filtered off, washed with methanol and dried. The residue is dissolved in an aqueous solution of sodium hydroxide. The aqueous phase is extracted several times with ethyl ether. The organic phases are combined and dried over anhydrous magnesium sulfate. The oil obtained (2.80 g) is used without additional purification.

15

20

Yield: 67%

Mp = Oil

5.3- 2-Benzyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide oxalate (1:2)

25

1.5 ml of acetic acid are added to a suspension of 0.45 g (0.0014 mol) of N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide and 0.149 g (0.0014 mol) of benzaldehyde in 12 ml of tetrahydrofuran. The suspension is stirred for 1 hour at ambient temperature and then 0.14 g (0.0021 mol) of sodium cyanoborohydride was added. The mixture was stirred overnight. The mixture is concentrated to dryness and the residue is treated with water and washed with ethyl ether. The aqueous phase is basified to pH = 10 and is extracted several times with ethyl ether. The organic phases are combined and dried over anhydrous magnesium sulfate. The oil obtained is purified by silica gel column chromatography, with a dichloromethane/methanol (95:5) mixture used as eluant. The desired product (0.13 g; 22%) is obtained in the form of an oil.

30

35

The above oil is dissolved in 5 ml of ethanol, and then 0.06 g (0.0007 mol) of oxalic acid, dissolved in 5 ml of ethanol, is added. The precipitate is filtered off and washed

with cold ethanol. 0.12 g of the desired product is obtained as a white solid.

Yield: 63%


Mp = 91-103°C

5

¹H-NMR (DMSO-d₆) δ(ppm): 7.5-7.1 (8H, m), 3.52 (2H, d), 2.9-2.5 (14H, m), 1.5 (2H, m), 0.9 (6H, t).

10

Example 6 *N*-[3-(Diethylamino)propyl]-*N*-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide hydrochloride



6.1- N-[2-(3-Diethylaminopropyl)-N-methyl-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide

0.09 g (0.0022 mol) of a dispersion of sodium hydride (60%) in a mineral oil is added
 5 to a solution of 0.91 g (0.0021 mol) of N-[2-(3-diethylaminopropyl)-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide (obtained according to the method described in stage 5.1- of Example 5) in 17 ml of dimethylformamide, cooled to 0°C. The mixture is stirred for one hour and 0.60 g (0.0042 mol) of methyl iodide is added. The mixture is stirred overnight at ambient temperature, and the
 10 solution is concentrated to dryness. The residue is treated with water. The aqueous phase is extracted several times with ethyl acetate. The organic phases are combined and dried over anhydrous magnesium sulfate. The oil obtained after filtration and evaporation of the solvent is purified by silica gel column chromatography, with a dichloromethane/methanol (95:5) mixture used as eluant, to
 15 give 0.31 g of the desired product in the form of an oil.

Yield: 34%

Mp = Oil

6.2- N-[2-(3-Diethylaminopropyl)-N-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide

To a solution of 0.20 g (0.00046 mol) of N-[2-(3-diethylaminopropyl)-N-methyl-2-(2,2,2-trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide dissolved in 10 ml
 25 of methanol saturated with hydrogen chloride is heated for twelve hours at 60°C. The solution is concentrated to dryness and the residue is treated with water. The aqueous phase is extracted several times with ethyl ether. The organic phases are combined and dried over anhydrous magnesium sulfate, and the solution is concentrated to dryness, to give 0.02 g of the desired product in the form of an oil.


30

Yield: 12%

Mp = Oil

¹H-NMR (DMSO) δ (ppm): 9.67 (1H, s, NH), 7.7 (1H, s), 7.68 (1H, d), 7.5 (1H, d), 4.4
 35 (2H, s), 3.5-2.9 (12H, m), 2.7 (3H, s), 1.9 (2H, m), 1.15 (6H, t).

Example 7. 2-Cyclohexyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide ethanodioate (1:1)



- 5 ml of acetic acid are added to a suspension of 1.70 g (0.0050 mol) of N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide (obtained according to the method described in stage 3.2- of Example 3) and 0.98 g (0.0014 mol) of cyclohexanone in 50 ml of tetrahydrofuran. The mixture is stirred for 3 hours at ambient temperature and 0.47 g (0.0075 mol) of sodium cyanoborohydride is added. The mixture is stirred overnight. The mixture is concentrated to dryness.
- 10 The residue is treated with water and washed with ethyl ether. The aqueous phase is basified to pH = 10 and is extracted several times with ethyl ether. The organic phases are combined and dried over anhydrous magnesium sulfate. The oil obtained is purified by silica gel column chromatography, with a dichloromethane/methanol (95:5) mixture used as eluant. The desired product (0.22 g; 11%) is obtained in the
- 15 form of an oil.


The above oil is dissolved in 5 ml of ethanol, and then a solution of 0.11 g (0.0012 mol) of oxalic acid, dissolved in 5 ml of ethanol, is added. The precipitate is filtered off and washed with cold ethanol. 0.32 g of the desired product is obtained as

20 a white solid.

Yield: 97%

Mp = 76-83°C.

- 25 ¹H-NMR (DMSO-d₆) δ(ppm): 7.6 (2H, m), 7.4 (1H, m), 4.2 (2H, s), 3.3 (H, m), 3.0 (H, m), 2.7 (H, m), 2.4 (H, m), 2.0 (H, m), 1.7 (H, m), 1.4 (H, m), 1.1 (H, m).

Example 8- 1-Aminopropylpyrroline**5 8.1- 1-(2-Cyanoethyl)pyrroline**

0.95 ml (0.0014 mol) of acrylonitrile is added, at 0°C, to a solution of 1.00 g (0.0014 mol) of pyrroline in methanol. The mixture is stirred overnight at ambient temperature and is concentrated to dryness, to give 1.65 g of the desired product in the form of an oil.

Yield: 93 %

Mp = Oil

15 8.2- 1-(3-Aminopropyl)pyrroline

0.024 g of 1M diisobutylaluminum hydride in toluene is added to a solution of 1.00 g (0.0089 mol) of 1-(2-cyanoethyl)pyrroline in 20 ml of dichloromethane. The mixture is stirred overnight at ambient temperature. The solution is treated with sodium sulfate decahydrate. The mixture is stirred for half an hour. The inorganic phases are filtered and the filtrate is concentrated to dryness, so as to obtain 1.00 g of a colorless oil.

Yield: 97%


Mp = Oil

¹H-NMR (CD₃Cl) δ (ppm): 5.75 (2H, s), 3.47 (4H, s), 2.75 (2H, t), 2.56 (2H, t), 1.69 (2H, q).



























The table below illustrates the chemical structures and the physical properties of some compounds according to the invention. The elemental microanalyses and the NMR, IR or mass spectra confirm the structures of the compounds obtained.

In the table, for the compounds of formula I, "Mp." corresponds to the melting point.

Table 1



No.	R2	-(C)n-	R1	B	Mp. (°C)	salt
1.	H	-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	97-102	Hydrochloride
2.	H	-(CH ₂) ₂ -	H		209-212	Hydrochloride
3.	CH ₃	-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	58-62	Oxalate
4.	H	-(CH ₂) ₃ -	CH ₃	-N(C ₂ H ₅) ₂	amorphous	Hydrochloride
5.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	91-95	Oxalate
6.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	98-103	Oxalate
7.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	144-147	Oxalate
8.	H	-(CH ₂) ₃ -	H		65-70	Hydrochloride
9.	H	-(CH ₂) ₃ -	H		>250	Hydrochloride
10.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	54-59	Oxalate
11.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	109-115	Oxalate
12.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	89-95	Oxalate
13.	H	-(CH ₂) ₃ -	H		228-231	Hydrochloride
14.		-(CH ₂) ₃ -	H	-N(C ₂ H ₅) ₂	55-60	Oxalate
15.		-(CH ₂) ₂ -	H		142-148	Oxalate
16.	-CH ₃	-(CH ₂) ₃ -	H		99-104	Oxalate
17.	H	-(CH ₂) ₃ -	H		244-249	Hydrochloride
18.	H	-(CH ₂) ₃ -	H		237-241	Hydrochloride

No.	R2	-(C)n-	R1	B	Mp. (°C)	salt
19.	H	-(CH ₂) ₃ -	H		238-245	Hydrochloride
20.	H	-(CH ₂) ₃ -	H	-N (CH ₃) ₂	230-234	Hydrochloride
21.		-(CH ₂) ₃ -	H		130-141	Oxalate
22.		-(CH ₂) ₂ -	H		50-60	Oxalate
23.		-(CH ₂) ₂ -	H		77-82	Oxalate
24.		-(CH ₂) ₂ -	H		103-110	Oxalate
25.		-(CH ₂) ₂ -	H		106-109	Oxalate
26.	H	-(CH ₂) ₂ -	H		114-117	Hydrochloride
27.	H	-(CH ₂) ₂ -	H		115-117	Hydrochloride
28.		-(CH ₂) ₂ -	H		134-140	Oxalate
29.		-(CH ₂) ₂ -	H		133-138	Oxalate
30.		-(CH ₂) ₂ -	H		86-110	Oxalate
31.		-(CH ₂) ₂ -	H		87-90	Oxalate
32.		-(CH ₂) ₂ -	H		66-77	Oxalate
33.		-(CH ₂) ₂ -	H		80-107	Oxalate
34.		-(CH ₂) ₂ -	H		121-125	Oxalate

The compounds of the invention of formula I were subjected to pharmacological assays which showed their advantage as active substances in therapeutics.

More particularly, the compounds of the invention are histamine H₃-receptor antagonists. H₃ receptors are known to those skilled in the art and their advantage in therapeutics has been described in the literature ("Histamine H₃ receptor antagonists" Exp. Opin. Ther. Patents (2000) 10 (7) :1045-1055).

Thus, the compounds of the invention of formula I were subjected to an *in vitro* affinity assay on the native histamine H₃ receptor in an adult rat brain membrane preparation, by specific binding of [³H]-N- α -methylhistamine to this receptor, according to the methods described by A. Korte et al., in Biochem. Biophys. Res. Commun. 168, 979-986(1990) and by R.E. West Jr. et al., in Mol. Pharmacol. 38, 610-613(1990).

The K_i values for the compounds of the invention with respect to H₃ receptors are between 0.1 nM and 5.0 μ M, and, more particularly, (+/-)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide (compound 15; Table 1) has a K_i of 0.3 nM.

The compounds of the invention of formula I were also subjected to a cAMP formation assay, on the human histamine H₃ receptor transfected into CHO cells, by inhibition of the agonism caused by the specific binding of R- α -methylhistamine to this receptor, according to the methods described by T.W. Lovenberg et al., in J. Pharmacol. Exp. Ther. 293, 771-778(2000).

The IC₅₀ values for the compounds of the invention with respect to H₃ receptors are between 0.1 nM and 5.0 μ M.

By way of example, compound 15, included in Table 1, has an IC₅₀<10 nM, using an EIA kit (Amersham) to measure cAMP formation, on the human histamine H₃ receptor transfected into CHO cells, by inhibition of the agonism caused by the specific binding of R- α -methylhistamine to this receptor.

The compounds according to the invention have an activity that is selective for the histamine H₃ receptor. Effectively, the compounds have a K_i of greater than 7.0 μ M in the *in vitro* affinity assay on the native histamine H₁ receptor in an adult rat brain membrane preparation by specific binding of [³H]-pirilamine to this receptor, according to the method described by Y.Q. Liu et al., in J. Pharmacol. Exp. Ther. 268, 959 (1994).

Furthermore, the compounds of the invention of formula I were subjected to *in vivo* tests showing their ability to reduce food intake in rats fasting for 24 h.

The experiments were carried out on Wistar rats. The rats were placed individually in transparent plastic cages 48x26, 5x21.5 cm. These cages were placed in a room insulated against any noise, at a temperature of 20 to 22°C, with a light cycle from 7 o'clock in the morning to 7 o'clock in the evening, the rats having free access to water and to the food.

Before the experiment was carried out, the rats were made to fast for 24 h, with access, however, to water *ad libitum*. On the day of the experiment, the carrier or the compound according to the present invention was administered i.p. or p.o., 15 or 30 minutes before a known amount of food (30 g) was made available. Each hour, for 6 hours, the amount of food ingested by the rat was measured.

It was shown that the AD₅₀ values (mg/kg i.p. or p.o.) for the compounds of the invention with respect to food intake may be less than 10. For example, (+/-)-2-(4-isopropylbenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide (compound 24; Table 1) decreases food intake by 54% during the first hour after the i.p. administration of 10 mg/kg of the product.

The results of the tests show that the compounds of the invention make it possible to reduce food intake in animals. Thus, they make it possible to control weight gain, to treat obesity or to aid weight loss, in animals, but also in humans.

Thus, according to another of its aspects, a subject of the invention is medicaments which comprise a compound of formula I or an addition salt of the latter with a pharmaceutically acceptable acid or else a hydrate or a solvate of the compound of formula I.

These medicaments find their use in therapeutics, in particular in the treatment of pathologies in which a histamine H₃-receptor antagonist provides a therapeutic benefit. In particular, such pathologies are obesity and diabetes. In addition, these compounds may be used in the treatment of central nervous system diseases such as vigilance and sleep disorders, narcolepsy, Alzheimer's disease and other dementias, Parkinson's disease, attention disorders in hyperkinetic children, memory and learning disorders, epilepsy, schizophrenia, moderate cognitive disorders, depression and anxiety. The states of depression and anxiety include, for example, anxieties of anticipatory type (before a surgical procedure, before a dental treatment,

etc), anxiety caused by alcohol or drug dependency or withdrawal, mania, seasonal affective disorders, migraines and nausea. They can also be used in the treatment of sexual dysfunction, dizziness and travel sickness.

The use of the compounds according to the invention for the preparation of a medicament for use in treating the pathologies mentioned above is an integral part of the invention.

According to another of its aspects, the present invention relates to pharmaceutical compositions comprising, as active ingredient, at least one compound according to the invention. These pharmaceutical compositions contain an effective dose of at least one compound according to the invention, or a pharmaceutically acceptable salt, a hydrate or a solvate of said compound, and also at least one or more pharmaceutically acceptable excipients. Said excipients are chosen according to the pharmaceutical form and the method of administration desired, from the usual excipients that are known to those skilled in the art.

In the pharmaceutical compositions of the present invention for oral, sublingual, subcutaneous, intramuscular, intravenous, topical, local, intratracheal, intranasal, transdermal or rectal administration, the active ingredient of formula I above, or possible salt, solvate or hydrate thereof, can be administered in unit administration form, as a mixture with conventional pharmaceutical excipients, to animals or to human beings for the prophylaxis or the treatment of the disorders or of the diseases above.

The appropriate unit administration forms comprise oral administration forms, such as tablets, soft or hard gelatin capsules, powders, granules and oral solutions or suspensions, sublingual, buccal, intratracheal or intranasal administration forms, forms for administration by inhalation, topical, transdermal, subcutaneous, intramuscular or intravenous administration forms, rectal administration forms, and implants. For topical application, the compounds according to the invention can be used in creams, gels, ointments or lotions.

In order to obtain the desired prophylactic or therapeutic effect, the dose of active ingredient can range between 0.1 μg and 50 mg per kg of bodyweight and per day. Each unit dose can contain from 0.1 to 1000 mg, preferably from 1 to 500 mg, of active ingredient in combination with a pharmaceutical excipient. This unit dose can be administered 1 to 5 times a day so as to administer a daily dose of from 0.5 to 5000 mg, preferably from 1 to 2500 mg.

There may be specific cases where higher or lower dosages are appropriate. Such dosages also belong to the invention. According to the usual practice, the dosage appropriate for each patient is determined by the physician according to the method of administration and the weight and response of said patient.

5

By way of example, a unit administration form of a compound according to the invention:


	Compound according to the invention	50.0 mg
10	Mannitol	223.75 mg
	Sodium croscarmellose	6.0 mg
	Corn starch	15.0 mg
	Hydroxypropylmethylcellulose	2.25 mg
	Magnesium stearate	3.0 mg

15

According to another of its aspects, the present invention also relates to a method of treating the pathologies indicated above, which comprises the administration, to a patient, of an effective dose of a compound according to the invention or a pharmaceutically acceptable salt or hydrate or solvate thereof.

Claims:

1. Compound of formula I:



in which:

n may adopt a value from 1 to 6;

-(C)_n- represents a -C₁₋₆ alkylidene group, optionally substituted by 1 to 4 substituents selected from a halogen atom or a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino or C₁₋₃ alkoxy group;

R1 represents

- a hydrogen atom,
- a C₁₋₆ alkyl group;

R2 represents

- a hydrogen atom,
- a C₁₋₆ alkyl or C₃₋₆ cycloalkyl group, each of which is optionally substituted by 1 to 4 substituents selected from a halogen atom, a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy or C₃₋₆ cycloalkyl group, a monocyclic heteroaryl such as a thienyl, furyl or pyrrolyl or an aryl, such as a phenyl or a naphthyl, the aryl being optionally substituted by 1 to 4 substituents selected from a halogen atom or a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy or a C₁₋₃ alkylidenedioxy group;

B represents • **NR₃R₄**,

- where R₃ and R₄ represent, independently of one another, a C₁₋₆ alkyl group or a hydrogen atom; or
- R₃ and R₄ together represent a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, a C₁₋₃ alkylidene-O-C₁₋₃ alkylidene group, or a group C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene in which R₅ represents a hydrogen atom or a C₁₋₃ alkyl or C₁₋₆ alkylcarbonyl group, it being possible for these C₁₋₃ alkyl and C₁₋₆ alkylcarbonyl groups to be substituted by a halogen atom or a hydroxyl, C₁₋₃ alkoxy, nitro, cyano or amino group; or
- a cyclic amino, bonded via a carbon to the group -NR₁-(C)_n-, chosen among aziridine, azetidine, pyrrolidine, piperidine **and** morpholine;


the groups R₃ and R₄ and the cyclic amino being optionally substituted by 1 to 4 substituents selected from a phenyl, a benzyl, a halogen atom or a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl or C₁₋₃ alkoxy group; and the nitrogen atom being optionally substituted by a C₁₋₃ alkyl, in the form of a base or addition salt with an acid, or in the hydrate or solvate form, **other than the compound in which R₁ and R₂ represent hydrogen atoms, B represent a dimethylamino group and -(C)_n- represent an ethylidene group.**

2. Compound according to Claim 1, characterized in

that:

- n is 2, 3 or 4; and
- R1 represents a hydrogen atom or a C₁₋₂ alkyl group; and
- R2 represents a hydrogen atom or a C₁₋₄ alkyl or C₅₋₆ cycloalkyl group each of which is optionally substituted by 1 to 4 substituents selected from a phenyl or a C₃₋₆ cycloalkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl or C₁₋₃ alkoxy group, the phenyl being optionally substituted by 1 to 4 substituents selected from a halogen atom or a hydroxyl, nitro, cyano, amino, C₁₋₃ monoalkylamino, C₂₋₆ dialkylamino, C₁₋₃ alkyl, C₁₋₂ perhaloalkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy or a C₁₋₃ alkylidenedioxy group; and
- B represents• NR₃R₄,
 - where R₃ and R₄ represent, independently of one another, a C₁₋₄ alkyl group; or
 - when R₃ and R₄ together represent a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, C₁₋₃ alkylidene-O-C₁₋₃ alkylidene or C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene,

B represents a group:




- or a cyclic amino bonded via a carbon to the group -NR₁-(C)_n, chosen among aziridine, azetidine, pyrrolidine, piperidine **and** morpholine;

the groups R₃, R₄ and R₅ and the cyclic amino being optionally substituted; in the form of a base or addition salt with an

acid, or in the hydrate or solvate form.

3. Compound according to Claim 2, characterized in that, when B represents NR₃R₄ and R₃ and R₄ together form a C₁₋₆ alkylidene group, a C₂₋₈ alkenylidene group, C₁₋₃ alkylidene-O-C₁₋₃ alkylidene or C₁₋₃ alkylidene-N(R₅)-C₁₋₃ alkylidene or when B represents a cyclic amino, B is selected from the following groups:




4. Compound according to Claim 1, 2 or 3, characterized in that it consists in:

- N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(diethylamino)propyl]-2-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(diethylamino)propyl]-N-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- 2-benzyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- 2-(cyclopropylmethyl)-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- 2-(cyclohexylmethyl)-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-N-[3-(2-methylpiperidin-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;

- N-[3-(3,6-dihydropyridin-1(2H)-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(diethylamino)propyl]-2-isopropyl-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(diethylamino)propyl]-2-(2-thienylmethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(diethylamino)propyl]-2-(3-thienylmethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(2,5-dihydro-1H-pyrrol-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- 2-cyclohexyl-N-[3-(diethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(2,5-dihydro-1H-pyrrol-1-yl)propyl]-2-methyl-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(4-benzylpiperazin-1-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-(3-pyrrolidin-1-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-(3-morpholin-4-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- N-[3-(dimethylamino)propyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- 2-(cyclohexylmethyl)-N-(3-pyrrolidin-1-ylpropyl)-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-2-(cyclopropylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-2-benzyl-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-

- 1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
- (+/-)-2-(4-isopropylbenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(1,3-benzodioxol-5-ylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (-)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (-)-2-(cyclohexylmethyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(4-bromobenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(2,5-dimethoxybenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(2-methylbutyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(3-methoxybenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide;
 - (+/-)-2-(3,5-dimethylbenzyl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulphonamide
- in the form of a base or addition salt with an acid, or in the hydrate or solvate form.

5. Compound of formula II:



in which Pg represents a hydrogen atom or a protective group and, $-(C)_n-$ and R1 are as defined in Claim 1, and B represents:

- NR3R4, where as defined in claim 1; or
 - a cyclic amino, bonded via a carbon to the group -NR1-(C)n-, chosen among aziridine, azetidine, pyrrolidine, piperidine and morpholine;
- the groups R3 and R4 and the cyclic amino being optionally substituted as defined in claim 1.

6. Pharmaceutical composition comprising a compound of formula I according to any one of Claims 1 to 4 or a salt, solvate or hydrate thereof and at least one pharmaceutical excipient.

7. Use of a compound of formula I according to any one of Claims 1 to 4 or of a salt, solvate or hydrate thereof for preparing a medicinal product intended for treating obesity and diabetes.

8. Use of a compound of formula I according to any one of Claims 1 to 4 or of a salt, solvate or hydrate thereof for preparing a medicinal product intended for treating central nervous system diseases such as wakefulness and sleep disorders, narcolepsy, Alzheimer's disease and other dementias, Parkinson's disease, attention disorders in hyperkinetic children, memory and learning disorders, epilepsy, schizophrenia, moderate cognitive disorders,

depression, anxiety, sexual dysfunction, vertigo and travel sickness.