

PYRAZOLOPYRIMIDINE COMPOUNDS

RELATED APPLICATIONS

This application claims the benefit of International Application No.
5 PCT/CA2013/000957, filed November 15, 2013. The entire teachings of the aforementioned application are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Protein kinases have been the subject of extensive study in the search for new
10 therapeutic agents in various diseases, for example, cancer. Protein kinases are known to mediate intracellular signal transduction by effecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is involved in a signaling pathway. There are a number of kinases and pathways through which extracellular and other stimuli cause a variety of cellular responses to occur inside the cell.

15 Human TTK protein kinase (TTK), also known as tyrosine threonine kinase, dual specificity protein kinase TTK, Monopolar Spindle 1 (Mps1) and Phosphotyrosine-Picked Threonine Kinase (PYT), is a conserved multispecific kinase that is capable of phosphorylating serine, threonine and tyrosine residues when expressed in *E. coli* (Mills *et al.*, *J. Biol. Chem.* 22(5): 16000-16006 (1992)). TTK mRNA is not expressed in the majority of physiologically
20 normal tissues in human (*Id.*). TTK mRNA is expressed in some rapidly proliferating tissues, such as testis and thymus, as well as in some tumors (for example, TTK mRNA was not expressed in renal cell carcinoma, was expressed in 50% of breast cancer samples, was expressed in testicular tumors and ovarian cancer samples) (*Id.*). TTK is expressed in some cancer cell lines and tumors relative to normal counterparts (*Id.*; see also WO 02/068444 A1).

25 Therefore, agents which inhibit a protein kinase, in particular TTK, have the potential to treat cancer. There is a need for additional agents which can act as protein kinase inhibitors, in particular TTK inhibitors.

In addition, cancer recurrence, drug resistance or metastasis is one of the major challenges in cancer therapies. Cancer patients who responded favorably to the initial anti-
30 cancer therapy often develop drug resistance and secondary tumors that lead to the relapse of the disease. Recent research evidences suggest that the capability of a tumor to grow and propagate is dependent on a small subset of cells within the tumor. These cells are termed tumor-initiating cells (TICs) or cancer stem cells. It is thought that the TICs are responsible for drug resistance,


cancer relapse and metastasis. Compounds that can inhibit the growth and survival of these tumor-initiating cells can be used to treat cancer, metastasis or prevent recurrence of cancer. Therefore, a need exists for new compounds that can inhibit the growth and survival of tumor-imitating cells.

5

SUMMARY OF THE INVENTION

Applicants have now discovered that certain pyrazolopyrimidine compounds are potent kinase inhibitors, such as TTK protein kinase (*see* Example B). Applicants have also discovered that these compounds have potent anticancer activity against breast cancer, colon cancer, and ovarian cancer cells in cell culture study (*see* Examples C-D). Based on these discoveries, pyrazolopyrimidine compounds, pharmaceutical compositions thereof, and methods of treating cancer with the pyrazolopyrimidine compounds are disclosed herein.

The present teachings are directed, at least in part, to a compound represented by the following structural formula:



15

or a pharmaceutically acceptable salt thereof, wherein:

R¹ is -NH-CH₂-Cy,

Cy is C₃-C₄ cycloalkyl optionally substituted with one or two groups selected from alkyl and hydroxyl;

20

R² is -O-pyridinyl; -NH-(C₂-C₆)hydroxyalkyl optionally substituted with cyclopropyl or isopropyl; or -NH-(C₃-C₆)cycloalkyl optionally substituted with hydroxyl or (C₁-C₂)hydroxyalkyl;

R⁴ is selected from hydrogen, halogen, and (C₁-C₃)alkyl; and

R^d is cyclopropyl. Preferably, R^d is chlorine or methyl.

In one embodiment, the present teachings include a pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a compound represented by structural formula (I) described above or a pharmaceutically acceptable salt thereof.

5 In another embodiment, the present teachings provide a method of treating a subject having cancer comprising administering to the subject an effective amount of a compound of structural formula (I) or a pharmaceutically acceptable salt thereof.

Another embodiment of the present teachings provides a method of inhibiting TTK activity in a subject in need of inhibition of TTK activity, comprising administering to the subject an effective amount of a compound represented by Structural Formula (I) or a
10 pharmaceutically acceptable salt thereof.


Another embodiment of the present teachings includes the use of a compound represented by Structural Formula (I) or a pharmaceutically acceptable salt thereof in therapy. In some embodiments, the therapy is for treating a subject with cancer. Alternatively, the therapy is for inhibiting TTK activity in a subject in need of inhibition of TTK activity.

15 Another embodiment of the present teachings includes the use of a compound represented by Structural Formula (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating a subject with cancer.


Another embodiment of the present teachings includes the use of a compound represented by Structural Formulas (I) or a pharmaceutically acceptable salt thereof for the
20 manufacture of a medicament for inhibiting TTK activity in a subject in need of inhibition of TTK activity.

DETAILED DESCRIPTION OF THE INVENTION


In one embodiment, the present teachings are directed to a compound represented by
25 Structural Formula (I) or a pharmaceutically acceptable salt thereof. The invention also includes the compounds depicted by structure and/or described by name in the Exemplification, and includes both the neutral forms and as well as pharmaceutically acceptable salts thereof. Treatments with and/or uses of these compounds (including neutral forms and pharmaceutically acceptable salts thereof) as described herein are also included in the invention. Specific
30 examples of compounds of the invention are shown below:



or a pharmaceutically acceptable salt thereof.




or a pharmaceutically acceptable salt thereof;



5

or a pharmaceutically acceptable salt thereof; and



or a pharmaceutically acceptable salt thereof.

The term “alkyl” used alone or as part of a larger moiety, such as “hydroxyalkyl”, and
 10 the like, means saturated aliphatic straight-chain or branched monovalent hydrocarbon radical.
 Unless otherwise specified, an alkyl group typically has 1-6 carbon atoms, *i.e.*, (C₁-C₆)alkyl. As

used herein, a "(C₁-C₆)alkyl" group means a radical having from 1 to 6 carbon atoms in a linear or branched arrangement.

"Cycloalkyl" means a saturated aliphatic cyclic hydrocarbon radical optionally containing one or more double bonds. It can be monocyclic, bicyclic (e.g., a bridged bicyclic ring), polycyclic (e.g., tricyclic), or fused. For example, monocyclic (C₃-C₇)cycloalkyl means a
5 radical having from 3-7 carbon atoms arranged in a monocyclic ring. A (C₃-C₇)cycloalkyl includes, but is not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

Certain of the compounds described herein may exist in various stereoisomeric or tautomeric forms. Stereoisomers are compounds which differ only in their spatial arrangement.
10 When a disclosed compound is named or depicted by structure without indicating stereochemistry, it is understood that the name or structure encompasses all possible stereoisomers, tautomers, geometric isomers or a combination thereof.

When a geometric isomer is depicted by name or structure, it is to be understood that the geometric isomeric purity of the named or depicted geometric isomer is at least 60%, 70%, 80%,
15 90%, 99% or 99.9% pure by weight. Geometric isomeric purity is determined by dividing the weight of the named or depicted geometric isomer in the mixture by the total weight of all of the geometric isomers in the mixture.

Racemic mixture means 50% of one enantiomer and 50% of its corresponding enantiomer. The present teachings encompass all enantiomerically-pure, enantiomerically-enriched, diastereomerically pure, diastereomerically enriched, and racemic mixtures, and
20 diastereomeric mixtures of the compounds described herein.

Enantiomeric and diastereomeric mixtures can be resolved into their component enantiomers or stereoisomers by well known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral
25 salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers can also be obtained from diastereomerically- or enantiomerically-pure intermediates, reagents, and catalysts by well known asymmetric synthetic methods.

When a compound is designated by a name or structure that indicates a single enantiomer, unless indicated otherwise, the compound is at least 60%, 70%, 80%, 90%, 99% or
30 99.9% optically pure (also referred to as "enantiomerically pure"). Optical purity is the weight in the mixture of the named or depicted enantiomer divided by the total weight in the mixture of both enantiomers.

When the stereochemistry of a disclosed compound is named or depicted by structure, and the named or depicted structure encompasses more than one stereoisomer (e.g., as in a

diastereomeric pair), it is to be understood that one of the encompassed stereoisomers or any mixture of the encompassed stereoisomers are included. It is to be further understood that the stereoisomeric purity of the named or depicted stereoisomers at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight. The stereoisomeric purity in this case is determined by dividing the total weight in the mixture of the stereoisomers encompassed by the name or structure by the total weight in the mixture of all of the stereoisomers.

Included in the present teachings are pharmaceutically acceptable salts of the compounds disclosed herein. The disclosed compounds have basic amine groups and therefore can form pharmaceutically acceptable salts with pharmaceutically acceptable acid(s). Suitable pharmaceutically acceptable acid addition salts of the compounds described herein include salts of inorganic acids (such as hydrochloric acid, hydrobromic, phosphoric, nitric, and sulfuric acids) and of organic acids (such as, acetic acid, benzenesulfonic, benzoic, methanesulfonic, and p-toluenesulfonic acids). Compounds of the present teachings with acidic groups such as carboxylic acids can form pharmaceutically acceptable salts with pharmaceutically acceptable base(s). Suitable pharmaceutically acceptable basic salts include ammonium salts, alkali metal salts (such as sodium and potassium salts) and alkaline earth metal salts (such as magnesium and calcium salts). Compounds with a quaternary ammonium group also contain a counteranion such as chloride, bromide, iodide, acetate, perchlorate and the like. Other examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, benzoates and salts with amino acids such as glutamic acid.

Compounds described herein can inhibit various kinases, including the TTK. Thus, generally, compounds described herein are useful in the treatment of diseases or conditions associated with such kinases. In some embodiments, compounds described herein can inhibit TTK.

In one embodiment, the compounds described herein are TTK inhibitors, and are useful for treating diseases, such as cancer, associated with such kinase(s).

Another aspect of the present teachings relates to a method of treating a subject with cancer comprising administering to the subject an effective amount of a compound described herein. In one embodiment, the compounds described herein inhibit the growth of a tumor. For example, the compounds described herein inhibit the growth of a tumor that overexpresses TTK.

Cancers that can be treated (including reduction in the likelihood of recurrence) by the methods of the present teachings include lung cancer, breast cancer, colon cancer, brain cancer, neuroblastoma, prostate cancer, melanoma, glioblastoma multiform, ovarian cancer, lymphoma, leukemia, melanoma, sarcoma, paraneoplasia, osteosarcoma, germinoma, glioma and mesothelioma. In one embodiment, the cancer is selected from leukemia, acute myeloid

leukemia, chronic myelogenous leukemia, breast cancer, brain cancer, colon cancer, colorectal cancer, head and neck cancer, hepatocellular carcinoma, lung adenocarcinoma, metastatic melanoma, pancreatic cancer, prostate cancer, ovarian cancer and renal cancer. In one embodiment, the cancer is lung cancer, colon cancer, brain cancer, neuroblastoma, prostate cancer, melanoma, glioblastoma multiforme or ovarian cancer. In another embodiment, the cancer is pancreatic cancer, prostate cancer, lung cancer, melanoma, breast cancer, colon cancer, or ovarian cancer. In yet another embodiment, the cancer is breast cancer, colon cancer and ovarian cancer. In yet another embodiment, the cancer is a breast cancer. In yet another embodiment, the cancer is a basal sub-type breast cancer or a luminal B sub-type breast cancer.

10 In yet another embodiment, the cancer is a basal sub-type breast cancer that overexpresses TTK. In yet another embodiment, the basal sub-type breast cancer is ER (estrogen receptor), HER2 and PR (progesterone receptor) negative breast cancer. In yet another embodiment, the cancer is a soft tissue cancer. A "soft tissue cancer" is an art-recognized term that encompasses tumors derived from any soft tissue of the body. Such soft tissue connects, supports, or surrounds

15 various structures and organs of the body, including, but not limited to, smooth muscle, skeletal muscle, tendons, fibrous tissues, fatty tissue, blood and lymph vessels, perivascular tissue, nerves, mesenchymal cells and synovial tissues. Thus, soft tissue cancers can be of fat tissue, muscle tissue, nerve tissue, joint tissue, blood vessels, lymph vessels, and fibrous tissues. Soft tissue cancers can be benign or malignant. Generally, malignant soft tissue cancers are referred

20 to as sarcomas, or soft tissue sarcomas. There are many types of soft tissue tumors, including lipoma, lipoblastoma, hibernoma, liposarcoma, leiomyoma, leiomyosarcoma, rhabdomyoma, rhabdomyosarcoma, neurofibroma, schwannoma (neurilemoma), neuroma, malignant schwannoma, neurofibrosarcoma, neurogenic sarcoma, nodular tenosynovitis, synovial sarcoma, hemangioma, glomus tumor, hemangiopericytoma, hemangioendothelioma, angiosarcoma,

25 Kaposi sarcoma, lymphangioma, fibroma, elastofibroma, superficial fibromatosis, fibrous histiocytoma, fibrosarcoma, fibromatosis, dermatofibrosarcoma protuberans (DFSP), malignant fibrous histiocytoma (MFH), myxoma, granular cell tumor, malignant mesenchymomas, alveolar soft-part sarcoma, epithelioid sarcoma, clear cell sarcoma, and desmoplastic small cell tumor. In a particular embodiment, the soft tissue cancer is a sarcoma selected from the group consisting

30 of a fibrosarcoma, a gastrointestinal sarcoma, a leiomyosarcoma, a dedifferentiated liposarcoma, a pleomorphic liposarcoma, a malignant fibrous histiocytoma, a round cell sarcoma, and a synovial sarcoma.

In some embodiments, the present teachings provide methods of inhibiting the growth of tumor-initiating cells or reducing the likelihood of recurrence of a cancer in a subject who is

35 undergoing an anti-cancer therapy. The method comprises the steps of:

- a) assessing the subject to determine whether the cancer is in remission; and

b) if the cancer is in remission; then administering to the subject an effective amount of a TTK inhibitor (*e.g.*, a compound represented by Structural Formula (I). If the cancer is not in remission, the method optionally further comprises the step of continuing the anti-cancer therapy until the cancer goes into remission and then the step b) of administering an effective amount of a TTK inhibitor (*e.g.*, a compound represented by Structural Formula (I)).

As used herein, the term “tumor-initiating cells” or “TICs” refer to cells present within some tumors that possess the ability to self-renew and proliferate. These cells are sometimes called cancer stem cells (CSCs) and may be observed to share certain characteristics with normal stem cells, including a stem cell-like phenotype and function. In some embodiments, TICs are characterized by their ability to form tumors after xenotransplantation in immunodeficient mice.

In some embodiments, the present teachings provide methods of inhibiting the growth of tumor-initiating cells or reducing the likelihood of recurrence of a cancer in a subject whose cancer is in remission comprising administering to the subject an effective amount of a TTK inhibitor (*e.g.*, a compound represented by Structural Formula (I)).

In some embodiments, *e.g.*, where the subject is being treated to reduce the likelihood of recurrence of a cancer, the subject has already been treated with an anti-cancer therapy. Alternatively, the subject has already been treated with an anti-cancer therapy and the subject is in remission.

In some embodiments, the present teachings provide methods of treating a subject with a cancer comprising administering to the subject an effective amount of a compound represented by Structural Formula (I) in combination with an effective anti-cancer therapy. In one embodiment, the cancer is a metastatic cancer. A “metastatic cancer” is a cancer that has spread from its primary site to other parts of the body.

In another embodiment, the present teachings are directed to a method of treating a subject with a drug-resistant cancer. A “drug-resistant cancer” is a cancer that is not responsive to one, two, three, four, five or more drugs that are typically used for the treatment of the cancer. In one embodiment, the drug-resistant cancer is mediated by the growth of tumor-initiating cells.

Suitable methods known in the art can be used for assessing a subject to determine whether the cancer is in remission. For example, the size of the tumor and/or tumor markers, usually proteins associated with tumors, can be monitored to determine the state of the cancer. Size of the tumor can be monitored with imaging devices, such as X-ray, MRI, CAT scans, ultrasound, mammography, PET and the like or via biopsy.

For methods described herein, *e.g.*, coadministration methods, the anti-cancer therapy are selected from the group consisting of surgery, radiation therapy, immunotherapy, endocrine

therapy, gene therapy and administration of an anti-cancer agent. Alternatively, the anti-cancer therapy is radiation therapy. In another alternative, the anti-cancer therapy is immunotherapy. In another alternative, the anti-cancer therapy is administration of an anti-cancer agent. In yet another alternative, the anti-cancer therapy is surgery.

5 Radiation therapy is the use of radiation to kill, destroy or treat the cancers. Exemplary radiation therapy includes, but is not limited to, gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and radioisotope therapy (*i.e.*, systemic radioactive isotopes therapy),

 An endocrine therapy is a treatment that adds, blocks or removes hormones. For
10 example, chemotherapeutic agents that can block the production or activity of estrogen have been used for treating breast cancer. In addition, hormonal stimulation of the immune system has been used to treat specific cancers, such as renal cell carcinoma and melanoma. In one embodiment, the endocrine therapy comprises administration of natural hormones, synthetic hormones or other synthetic molecules that may block or increase the production of the body's
15 natural hormones. In another embodiment, the endocrine therapy includes removal of a gland that makes a certain hormone.

 As use herein, a gene therapy is the insertion of genes into a subject's cell and biological tissues to treat diseases, such as cancer. Exemplary gene therapy includes, but is not limited to, a germ line gene therapy and a somatic gene therapy.

20 Immunotherapy (also called biological response modifier therapy, biologic therapy, biotherapy, immune therapy, or biological therapy) is treatment that uses parts of the immune system to fight disease. Immunotherapy can help the immune system recognize cancer cells, or enhance a response against cancer cells. Immunotherapies include active and passive immunotherapies. Active immunotherapies stimulate the body's own immune system while
25 passive immunotherapies generally use immune system components created outside of the body.

 Examples of active immunotherapies include, but are not limited to vaccines including cancer vaccines, tumor cell vaccines (autologous or allogeneic), dendritic cell vaccines, antigen vaccines, anti-idiotypic vaccines, DNA vaccines, viral vaccines, or Tumor-Infiltrating Lymphocyte (TIL) Vaccine with Interleukin-2 (IL-2) or Lymphokine-Activated Killer (LAK)
30 Cell Therapy,.

 Examples of passive immunotherapies include but are not limited to monoclonal antibodies and targeted therapies containing toxins. Monoclonal antibodies include naked antibodies and conjugated monoclonal antibodies (also called tagged, labeled, or loaded antibodies). Naked monoclonal antibodies do not have a drug or radioactive material attached
35 whereas conjugated monoclonal antibodies are joined to, for example, a chemotherapy drug

(chemolabeled), a radioactive particle (radiolabeled), or a toxin (immunotoxin). Examples of these naked monoclonal antibody drugs include, but are not limited to Rituximab (Rituxan), an antibody against the CD20 antigen used to treat, for example, B cell non-Hodgkin lymphoma; Trastuzumab (Herceptin), an antibody against the HER2 protein used to treat, for example, advanced breast cancer; Alemtuzumab (Campath), an antibody against the CD52 antigen used to treat, for example, B cell chronic lymphocytic leukemia (B-CLL); Cetuximab (Erbix), an antibody against the EGFR protein used, for example, in combination with irinotecan to treat, for example, advanced colorectal cancer and head and neck cancers; and Bevacizumab (Avastin) which is an antiangiogenesis therapy that works against the VEGF protein and is used, for example, in combination with chemotherapy to treat, for example, metastatic colorectal cancer. Examples of the conjugated monoclonal antibodies include, but are not limited to Radiolabeled antibody Ibritumomab tiuxetan (Zevalin) which delivers radioactivity directly to cancerous B lymphocytes and is used to treat, for example, B cell non-Hodgkin lymphoma; radiolabeled antibody Tositumomab (Bexxar) which is used to treat, for example, certain types of non-Hodgkin lymphoma; and immunotoxin Gemtuzumab ozogamicin (Mylotarg) which contains calicheamicin and is used to treat, for example, acute myelogenous leukemia (AML). BL22 is a conjugated monoclonal antibody for treating, for example, hairy cell leukemia, immunotoxins for treating, for example, leukemias, lymphomas, and brain tumors, and radiolabeled antibodies such as OncoScint for example, for colorectal and ovarian cancers and ProstaScint for example, for prostate cancers.

Further examples of therapeutic antibodies that can be used include, but are not limited to, HERCEPTIN[®] (Trastuzumab) (Genentech, CA) which is a humanized anti-HER2 monoclonal antibody for the treatment of patients with metastatic breast cancer; REOPRO[®] (abciximab) (Centocor) which is an anti-glycoprotein IIb/IIIa receptor on the platelets for the prevention of clot formation; ZENAPAX[®] (daclizumab) (Roche Pharmaceuticals, Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection; PANOREX[™] which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a murine anti-idiotypic (GD3 epitope) IgG antibody (ImClone System); IMC-C225 which is a chimeric anti-EGFR IgG antibody (ImClone System); VITAXIN[™] which is a humanized anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Campath 1H/LDP-03 which is a humanized anti-CD52 IgG1 antibody (Leukosite); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); RITUXAN[™] which is a chimeric anti-CD20 IgG1 antibody (IDEC Pharm/Genentech, Roche/Zettyaku); LYMPHOCIDE[™] which is a humanized anti-CD22 IgG antibody (Immunomedics); LYMPHOCIDE[™] Y-90 (Immunomedics); Lymphoscan (Tc-99m-labeled; radioimaging; Immunomedics); Nuvion (against CD3; Protein Design Labs); CM3

is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80 antibody (IDEC Pharm/Mitsubishi); ZEVALIN[™] is a radiolabelled murine anti-CD20 antibody (IDEC/Schering AG); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); D2E7 is a humanized anti-TNF- α antibody (CAT/BASF); CDP870 is a humanized anti-TNF- α Fab fragment (Celltech); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CD20-sreptavidin (+biotin-⁹⁰Yttrium 90; NeoRx); CDP571 is a humanized anti-TNF- α IgG4 antibody (Celltech); LDP-02 is a humanized anti- α 4 β 7 antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA[™] is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN[™] is a humanized anti-VLA-4 IgG antibody (Elan); and CAT-152 is a human anti-TGF- β ₂ antibody (Cambridge Ab Tech).

Immunotherapies that can be used in the present teachings include adjuvant immunotherapies. Examples include cytokines, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte-colony stimulating factor (G-CSF), macrophage inflammatory protein (MIP)-1-alpha, interleukins (including IL-1, IL-2, IL-4, IL-6, IL-7, IL-12, IL-15, IL-18, IL-21, and IL-27), tumor necrosis factors (including TNF-alpha), and interferons (including IFN-alpha, IFN-beta, and IFN-gamma); aluminum hydroxide (alum); Bacille Calmette-Guérin (BCG); Keyhole limpet hemocyanin (KLH); Incomplete Freund's adjuvant (IFA); QS-21; DETOX; Levamisole; and Dinitrophenyl (DNP), and combinations thereof, such as, for example, combinations of, interleukins, for example, IL-2 with other cytokines, such as IFN-alpha.

Alternatively, the anti-cancer therapy described herein includes administration of an anti-cancer agent. An "anti-cancer agent" is a compound, which when administered in an effective amount to a subject with cancer, can achieve, partially or substantially, one or more of the following: arresting the growth, reducing the extent of a cancer (*e.g.*, reducing size of a tumor), inhibiting the growth rate of a cancer, and ameliorating or improving a clinical symptom or indicator associated with a cancer (such as tissue or serum components) or increasing longevity of the subject.

The anti-cancer agent suitable for use in the methods described herein includes any anti-cancer agents that have been approved for the treatment of cancer. In one embodiment, the anti-cancer agent includes, but is not limited to, a targeted antibody, an angiogenesis inhibitor, an

alkylating agent, an antimetabolite, a vinca alkaloid, a taxane, a podophyllotoxin, a topoisomerase inhibitor, a hormonal antineoplastic agent and other antineoplastic agents.

Examples of alkylating agents useful in the methods of the present teachings include but are not limited to, nitrogen mustards (*e.g.*, mechloroethamine, cyclophosphamide, chlorambucil, melphalan, *etc.*), ethylenimine and methylmelamines (*e.g.*, hexamethylmelamine, thiotepa), alkyl sulfonates (*e.g.*, busulfan), nitrosoureas (*e.g.*, carmustine, lomustine, semustine, streptozocin, *etc.*), or triazenes (decabazine, *etc.*). Examples of antimetabolites useful in the methods of the present teachings include but are not limited to folic acid analog (*e.g.*, methotrexate), or pyrimidine analogs (*e.g.*, fluorouracil, floxouridine, Cytarabine), purine analogs (*e.g.*, mercaptopurine, thioguanine, pentostatin). Examples of plant alkaloids and terpenoids or derivatives thereof include, but are not limited to, vinca alkaloids (*e.g.*, vincristine, vinblastine, vinorelbine, vindesine), podophyllotoxin, and taxanes (*e.g.*, paclitaxel, docetaxel). Examples of a topoisomerase inhibitor include, but are not limited to, irinotecan, topotecan, amsacrine, etoposide, etoposide phosphate and teniposide. Examples of antineoplastic agents include, but are not limited to, actinomycin, anthracyclines (*e.g.*, doxorubicin, daunorubicin, valrubicin, idarubicin, epirubicin), bleomycin, plicamycin and mitomycin.

In one embodiment, the anti-cancer agents that can be used in the present teachings include Adriamycin, Dactinomycin, Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; broprimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqunone; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; efornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; interleukin II (including recombinant interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1 ; interferon alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride;

lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedapa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone
 5 hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; pipsulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine;
 10 simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricyribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulazole hydrochloride; uracil
 15 mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinat sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride.

Yet other anti-cancer agents/drugs that can be used in the present teachings include, but
 20 are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen;
 25 antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid;
 30 bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflata; broprimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorlins; chloroquinoxaline
 35 sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin;

crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A;
 cyclopentantraquinones; cycloplatam; cypemycin; cytarabine ocfosfate; cytolytic factor;
 cytostatin; dacliximab; decitabine; dehydroidemnin B; deslorelin; dexamethasone;
 dexifosfamide; dexrazoxane; dexverapamil; diaziqune; didemnin B; didox; diethylnorspermine;
 5 dihydro-5-azacytidine; 9- dioxamycin; diphenyl spiromustine; docosanol; dolasetron;
 doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine;
 edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue;
 estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane;
 fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone;
 10 fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine;
 gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors;
 gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide;
 hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat;
 imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor
 15 inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol,
 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide;
 kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate;
 leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon;
 leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine
 20 analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7;
 lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan;
 lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat;
 masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril;
 merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine;
 25 mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues;
 mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene;
 molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid
 A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple
 tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial
 30 cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip;
 naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic
 acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide
 antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone;
 ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin;
 35 oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene;
 parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin;

pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate;
 phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A;
 placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-
 triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone;
 5 prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C
 inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine
 nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin
 polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein
 transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re
 10 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide;
 roquinimex; rubiginone B1; ruboxyl; safinol; saintopin; SarCNU; sarcophytol A; sargramostim;
 Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal
 transduction inhibitors; signal transduction modulators; single chain antigen-binding protein;
 sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin
 15 binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin;
 spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide;
 stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist;
 suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen
 methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase
 20 inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine;
 thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin
 receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine;
 titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors;
 tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine
 25 kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth
 inhibitory factor; urokinase receptor antagonists; vaporeotide; variolin B; vector system,
 erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine;
 vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer. Preferred
 additional anti-cancer drugs are 5-fluorouracil and leucovorin.

30 In one embodiment, the anti-cancer agents that can be used in methods described herein
 are selected from the group consisting of paclitaxel, docetaxel, 5-fluorouracil, trastuzumab,
 lapatinib, bevacizumab, letrozole, goserelin, tamoxifen, cetuximab, panitumumab, gemcitabine,
 capecitabine, irinotecan, oxaliplatin, carboplatin, cisplatin, doxorubicin, epirubicin,
 cyclophosphamide, methotrexate, vinblastine, vincristine, melphalan and a combination thereof.

35 In one embodiment, the anti-cancer agent and the compound represented by Structural
 Formula (I) are administered contemporaneously. When administered contemporaneously, the

anti-cancer agent and the compound can be administered in the same formulation or in different formulations. Alternatively, the compound and the additional anti-cancer agent are administered separately. Alternatively, the compound and the additional anti-cancer agent can be administered sequentially, as separate compositions, within an appropriate time frame (*e.g.*, a cancer treatment session/interval (*e.g.*, about 1.5 to about 5 hours to about 10 hours to about 15 hours to about 20 hours; about 1 day to about 2 days to about 5 days to about 10 days to about 14 days)) as determined by the skilled clinician (*e.g.*, a time sufficient to allow an overlap of the pharmaceutical effects of the therapies). The compound and the additional anti-cancer agent can be administered in a single dose or multiple doses in an order and on a schedule suitable to achieve a desired therapeutic effect (*e.g.*, inhibition of tumor growth).

In one embodiment, the subject in the methods described herein has not been previously treated with a TTK inhibitor (*e.g.*, the compound represented by Structural Formula (I)).

The term “inhibiting the growth of tumor-initiating cells” refers to decreasing the rate of the proliferation and/or survival of the tumor-initiating cells.

As used herein, the term “reducing the likelihood of recurrence of a cancer” means partially or totally inhibiting, delaying the return of a cancer at or near a primary site and/or at a secondary site after a period of remission. It also means that the cancer is less likely to return with treatment described herein than in its absence.

As used herein, the term “remission” refers to a state of cancer, wherein the clinical symptoms or indicators associated with a cancer have disappeared or cannot be detected, typically after the subject has been successfully treated with an anti-cancer therapy.

As used herein, “treatment” is an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of extent of disease, stabilized (*i.e.*, not worsening) state of disease, reducing the likelihood of the spread of the disease, delay or slowing of disease progression, amelioration or palliation of the disease state, whether detectable or undetectable. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment. “Treatment” also includes reducing the likelihood of reoccurrence of the disease.

As used herein, “treating a subject with a cancer” includes achieving, partially or substantially, one or more of the following: arresting the growth, reducing the extent of the cancer (*e.g.*, reducing size of a tumor), inhibiting the growth rate of the cancer, ameliorating or improving a clinical symptom or indicator associated with the cancer (such as tissue or serum components) or increasing longevity of the subject; and reducing the likelihood of recurrence of the cancer.

Generally, an effective amount of a compound taught herein varies depending upon various factors, such as the given drug or compound, the pharmaceutical formulation, the route of administration, the type of disease or disorder, the identity of the subject or host being treated, and the like, but can nevertheless be routinely determined by one skilled in the art. An effective
5 amount of a compound of the present teachings may be readily determined by one of ordinary skill by routine methods known in the art.

The term an "effective amount" means an amount when administered to the subject which results in beneficial or desired results, including clinical results, *e.g.*, inhibits, suppresses or reduces the cancer (*e.g.*, as determined by clinical symptoms or the amount of cancer cells) in
10 a subject as compared to a control.

In an embodiment, an effective amount of a compound taught herein ranges from about 0.1 to about 1000 mg/kg body weight, alternatively about 1 to about 500 mg/kg body weight, and in another alternative, from about 20 to about 300 mg/kg body weight. In another embodiment, an effective amount of a compound taught herein ranges from about 0.5 to about
15 5000 mg/m², alternatively about from 5 to about 2500 mg/m², and in another alternative from about 50 to about 1000 mg/m². The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject suffering from cancer or reduce the likelihood of recurrence of a cancer. These factors include, but are not limited to, the severity of the disease or disorder, previous treatments, the general health and/or age of the subject and
20 other diseases present.

Moreover, for methods described herein (including treating a subject with a cancer or reducing the likelihood of recurrence of a cancer), a "treatment" or dosing regimen of a subject with an effective amount of the compound of the present teachings may consist of a single administration, or alternatively comprise a series of applications. For example, the compound of
25 the present teachings may be administered at least once a week. However, in another embodiment, the compound may be administered to the subject from about one time per week to once daily for a given treatment. The length of the treatment period depends on a variety of factors, such as the severity of the disease, the age of the patient, the concentration and the activity of the compounds of the present teachings, or a combination thereof. It will also be
30 appreciated that the effective dosage of the compound used for the treatment may increase or decrease over the course of a particular treatment regime. Changes in dosage may result and become apparent by standard diagnostic assays known in the art. In some instances, chronic administration may be required.

A "subject" is a mammal, preferably a human, but can also be an animal in need of
35 veterinary treatment, *e.g.*, companion animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*,

cows, sheep, pigs, horses, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like).

The compounds taught herein can be administered to a patient in a variety of forms depending on the selected route of administration, as will be understood by those skilled in the art. The compounds of the present teachings may be administered, for example, by oral, parenteral, buccal, sublingual, nasal, rectal, patch, pump or transdermal administration and the pharmaceutical compositions formulated accordingly. Parenteral administration includes intravenous, intraperitoneal, subcutaneous, intramuscular, transepithelial, nasal, intrapulmonary, intrathecal, rectal and topical modes of administration. Parenteral administration can be by continuous infusion over a selected period of time.

The compounds taught herein can be suitably formulated into pharmaceutical compositions for administration to a subject. The pharmaceutical compositions of the present teachings optionally include one or more pharmaceutically acceptable carriers and/or diluents therefor, such as lactose, starch, cellulose and dextrose. Other excipients, such as flavoring agents; sweeteners; and preservatives, such as methyl, ethyl, propyl and butyl parabens, can also be included. More complete listings of suitable excipients can be found in the Handbook of Pharmaceutical Excipients (5th Ed., Pharmaceutical Press (2005)). A person skilled in the art would know how to prepare formulations suitable for various types of administration routes. Conventional procedures and ingredients for the selection and preparation of suitable formulations are described, for example, in Remington's Pharmaceutical Sciences (2003 - 20th edition) and in The United States Pharmacopeia: The National Formulary (USP 24 NF19) published in 1999. The carriers, diluents and/or excipients are "acceptable" in the sense of being compatible with the other ingredients of the pharmaceutical composition and not deleterious to the recipient thereof.

Typically, for oral therapeutic administration, a compound of the present teachings may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like.

Typically for parenteral administration, solutions of a compound of the present teachings can generally be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, DMSO and mixtures thereof with or without alcohol, and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

Typically, for injectable use, sterile aqueous solutions or dispersion of, and sterile powders of, a compound described herein for the extemporaneous preparation of sterile injectable solutions or dispersions are appropriate.

For nasal administration, the compounds of the present teachings can be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomizing device. Alternatively, the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal after use. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomizer.

For buccal or sublingual administration, the compounds of the present teachings can be formulated with a carrier such as sugar, acacia, tragacanth, or gelatin and glycerine, as tablets, lozenges or pastilles.

For rectal administration, the compounds described herein can be formulated in the form of suppositories containing a conventional suppository base such as cocoa butter.

The compounds of invention may be prepared by methods known to those skilled in the art, as illustrated by the general schemes and procedures below and by the preparative examples that follow. All starting materials are either commercially available or prepared by methods known to those skilled in the art and the procedures described below.

In accordance with another aspect of the present invention, the compounds of the invention can be prepared by processes analogous to those established in the art. General methods to synthesize the claimed compounds are elaborated below in Example A.

EXEMPLIFICATION

Example A: Synthesis

General Methods

Commercially available starting materials, reagents, and solvents were used as received. In general, anhydrous reactions were performed under an inert atmosphere such as nitrogen or Argon. PoraPak[®] Rxn CX refers to a commercial cation-exchange resin available from Waters.


Microwave reactions were performed with a Biotage Initiator microwave reactor. Reaction progress was generally monitored by TLC using Merck silica gel plates with visualization by UV at 254 nm, by analytical HPLC or by LCMS (Bruker Exquire 4000 or Waters Acquity UPLC system). Flash column chromatographic purification of intermediates or final products was performed using 230-400 mesh silica gel 60 from EMD chemicals or Silicycle, or purified using a Biotage Isolera with KP-SIL or HP-SIL silica cartridges, or KP-NH basic modified silica and corresponding samplets. Reverse-phase HPLC purification was performed on a Varian PrepStar model SD-1 HPLC system with a Varian Monochrom 10 μ C-18 reverse-phase column using a of about 5-30% MeCN or MeOH/0.05% TFA-H₂O to 70-90% MeCN or MeOH/0.05% TFA in H₂O over a 20-40-min period at a flow rate of 30-80 mL/min. Reverse phase purification was also performed using a Biotage Isolera equipped with a KP-C18-H column using a between 10-95% MeOH or CH₃CN/ 0.1% TFA in H₂O. Proton NMRs were recorded on a Bruker 400 MHz spectrometer, and mass spectra were obtained using a Bruker Esquire 4000 spectrometer or Waters Acquity UPLC system.

Compound names were generated using the software built into CambridgeSoft-PerkinElmer's ChemBioDraw Ultra version 11.0 or 12.0.

Abbreviations:

20	aq	aqueous
	Ar	argon
	Boc	<i>tert</i> -butoxy carbonyl
	br.	broad
	calcd	calculated
25	d	doublet
	DCM	dichloromethane
	DIPEA	diisopropylethylamine
	DMF	N,N-dimethylformamide
	DMSO	dimethylsulfoxide
30	dppf	1,1'- bis(diphenylphosphino) ferrocene
	h	hour
	HPLC	high performance liquid chromatography
	LC-MS	liquid chromatography coupled to mass spectrometry
	min	minute
35	m	multiplet
	MS ESI	mass spectra, electrospray ionization
	NMR	nuclear magnetic resonance
	O/N	overnight
	PE	petroleum ether
40	PMB	para-methoxybenzyl
	prep	preparative


rt room temperature
 s singlet
 t triplet
 TFA trifluoroacetic acid
 5 THF tetrahydrofuran




Intermediates:

10

4-bromo-N-cyclopropyl-2-methylbenzamide




To a suspension of 4-bromo-2-methylbenzoic acid (43.0 g, 200 mmol) and oxalyl dichloride (30.5 g, 240 mmol) in DCM (300 mL) was added DMF (0.1 mL). The resulting reaction mixture was stirred at rt for 16 h. The reaction turned into a clear yellow solution slowly over 16 h. Solvent was then removed *in vacuo*, and the crude product was used in the next step without further purification. The crude product was redissolved in DCM (300 mL) and cooled to 0 °C. A mixture of TEA (42 mL, 300 mmol) and cyclopropylamine (12.6 g, 220 mmol) in DCM (100 mL) was added slowly over 15 min, and the resulting mixture was stirred at rt for 2 h. The reaction was diluted with DCM (200 mL) and water was added. The resulting mixture was extracted with DCM and the combined organic extracts were dried over MgSO₄ and concentrated to give the desired product as a pale pink solid (50.1 g, 99%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.37 (s, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 7.17 (d, *J* = 8.1 Hz, 1H), 5.93 (br. s, 1H), 2.90-2.85 (m, 1H), 2.40 (s, 3H), 0.90-0.85 (m, 2H), 0.62-0.58 (m, 2H); MS ESI [*M* + *H*]⁺ 253.9, calcd for [C₁₁H₁₂BrNO + *H*]⁺ 254.0.

N-(4-bromo-2-methylphenyl)cyclopropanecarboxamide

In a 100 mL RBF, 4-bromo-2-methylaniline (3.7 g, 20 mmol) and DIPEA (6.95 mL, 40 mmol) were combined with DMF (40 mL).


The reaction was cooled to 0 °C in an ice bath and

- 5 cyclopropanecarbonyl chloride (2.1 g, 20 mmol) was added. The mixture was stirred at 0 °C for 1 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The organic layers were combined, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give the title compound as a white solid (4.76 g, 94%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.85-7.72 (m, 1H), 7.38-7.28 (m, 2H), 7.15-7.02 (m, 1H), 2.27 (s, 3H), 1.57-1.48 (m, 1H), 1.10 (quint, *J* = 3.9 Hz, 2H), 0.92-0.79 (m, 2H); MS ESI [M + H]⁺ 253.9, calcd for [C₁₁H₁₂BrNO + H]⁺ 254.0.

N-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide

To a mixture of 4-bromo-N-cyclopropyl-2-methylbenzamide (3.73 g, 14 mmol), Bis(pinacolato)diboron (5.59 g, 22 mmol), anh KOAc (4.29 g, 43 mmol) in DMF (37 mL) was purged with Ar for 10 min at rt. Then PdCl₂(dppf)·DCM (0.59 g, 5 mol%)

- was added and the reaction was heated at 100 °C in oil bath for 4 h. After reaction completion the reaction mass was diluted with EtOAc (200 mL) and H₂O (100 mL). The combined layer filtered through celite pad and washed it with little EtOAc. The aq. layer further extracted with EtOAc (50 mL) and the combined organic layer washed with brine, dried over Na₂SO₄, filtered, and concentrated to give crude oily residue. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-100%) to give the title compound as a creamy solid (4.15 g, 94%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.66 (s, 1H), 7.63-7.61 (d, *J* = 7.6 Hz, 1H), 7.32-7.30 (d, *J* = 7.6 Hz, 1H), 5.85 (s, 1H), 2.93-2.87 (m, 1H), 2.46 (s, 3H), 1.35 (s, 12H), 0.90-0.86 (m, 2H), 0.63-0.59 (m, 2H); MS ESI [M + H]⁺ 302.2, calcd for [C₁₇H₂₄BNO₃ + H]⁺ 302.2.

3-bromo-5,7-dichloropyrazolo[1,5-a]pyrimidine


To a stirred solution of sodium ethoxide in EtOH, which was prepared from sodium (281.3 g, 12.0 mol) and EtOH (10 L) by the conventional method, were added diethyl malonate (963.7 g, 6.02 mol) at ambient temperature and then compound 1H-pyrazol-3-amine (500 g, 6.02 mol).

The reaction mixture was refluxed for 12 hours. After cooled to room temperature, the precipitates were collected by filtration and dissolved in water. The aqueous solution was acidified with 2 M HCl (pH = 2). The resulting precipitates were collected by filtration and dried under reduced pressure to afford pyrazolo[1,5-a]pyrimidine-5,7(4H,6H)-dione (649 g, 71%) as a yellow solid, which was used for the next reaction without further purification.

A stirred suspension of pyrazolo[1,5-a]pyrimidine-5,7(4H,6H)-dione (265 g, 1.75 mol) and *N,N*-dimethylaniline (335.6 mL) in POCl₃ (2.00 kg, 13.2 mol) was refluxed for 4 hours. After cooled to room temperature, the reaction mixture was poured into ice-water, and stirred for 30 min, neutralized with saturated aqueous sodium carbonate and extracted with EtOAc. The combined organic layers were washed with water, brine and dried over MgSO₄, filtered and evaporated. The residue was purified by column chromatography on silica gel (gradient: EtOAc/PE 1:10) to give 5,7-dichloropyrazolo[1,5-a]pyrimidine (287 g, 87%) as a yellow solid.


To a solution 5,7-dichloropyrazolo[1,5-a]pyrimidine (246.6 g, 1.31 mol) in CH₃CN (1.8 L) was added NBS (245 g, 1.38 mol). The resulting mixture was stirred at room temperature for 2 hours. After removal of the solution, the reaction mixture was purified by column chromatography on silica gel (gradient: EtOAc/PE 1:5) to give the title compound (313.5 g, 89%) as light yellow solid. ¹H NMR (300 MHz, CDCl₃): δ ppm 7.04 (s, 1H), 8.21 (s, 1H); MS ESI [M + H]⁺ 265.9, calcd for [C₆H₂BrCl₂N₃ + H]⁺ 265.9.

3-bromo-5-chloro-N-(cyclopropylmethyl)pyrazolo[1,5-a]pyrimidin-7-amine



To a solution of 3-bromo-5,7-dichloropyrazolo[1,5-a]pyrimidine (5.0 g, 19 mmol) in DCM (50 mL) was added cyclopropylmethanamine (1.48 g, 21 mmol), and DIPEA (6.6 mL, 38 mmol). The reaction was stirred at rt for 2 h. Water and DCM were added to separate the phases and the aqueous phase was extracted with DCM. The combined organic extracts were dried over NaSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-50%) to give the title compound as a yellow solid (5.25 g, 92%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.97 (s, 1H), 6.50 (br. s, 1H), 5.97 (s, 1H), 3.25 (dd, *J* = 7.3, 5.5 Hz, 2H), 1.26-1.13 (m, 1H), 0.74-0.65 (m, 2H), 0.37 (q, *J* = 5.0 Hz, 2H); MS ESI [M + H]⁺ 301.0, calcd for [C₁₀H₁₀BrClN₄ + H]⁺ 301.0.


(1s,3s)-3-(((3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)amino)methyl)-1-methylcyclobutanol



To a solution of 3-bromo-5,7-dichloropyrazolo[1,5-a]pyrimidine (14.6 g, 55.3 mmol) in DCM (200 mL) was added cis-hydroxy-3-methylcyclobutane-1-methylamine (7.0 g, 60.9 mmol), and DIPEA (19.2 mL, 110.6 mmol). The reaction was stirred at rt for 16 h.

Water and DCM were added to separate the phases and the aqueous phase was extracted with DCM. The combined organic extracts were dried over Na₂SO₄, filtered and concentrated to give the title compound as a yellow solid (18.1 g, 95%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.96 (s, 1 H), 6.60-6.49 (m, 1H), 5.99 (s, 1H), 3.47 (t, *J* = 6.0 Hz, 2H), 2.38-2.27 (m, 3H), 1.96-1.85 (m, 2H), 1.43 (s, 3H); MS ESI [M + H]⁺ 345.1, calcd for [C₁₂H₁₄BrClN₄O + H]⁺ 345.0.


tert-butyl (3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)(cyclopropylmethyl)carbamate



To a solution of 3-bromo-5-chloro-*N*-(cyclopropylmethyl)pyrazolo[1,5-a]pyrimidin-7-amine (5.25 g, 17.5 mmol) in DCM (150 mL) was added Boc₂O (5.70 g, 26.2 mmol), DMAP (0.21 g, 1.75 mmol), and TEA (7.3 mL, 52.5 mmol). The

reaction was stirred at rt for 4 h. Water and DCM were added to separate the phases and the aqueous phase was extracted with DCM. The combined organic extracts were dried over MgSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-40 %) to give the title compound as a yellow solid (6.24 g, 89%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.12 (s, 1H), 6.85 (s, 1H), 3.72 (d, *J* = 7.3 Hz, 2H), 1.39 (s, 9H), 1.05-0.94 (m, 1H), 0.46-0.37 (m, 2H), 0.13-0.03 (m, 2H); MS ESI [M – C₄H₈]⁺ 345.0, calcd for [C₁₅H₁₈BrClN₄O₂ – C₄H₈]⁺ 345.0.


tert-butyl (3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)(((1s,3s)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate



To a solution of (1s,3s)-3-(((3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)amino)methyl)-1-methylcyclobutanol (18.1 g, 52.6 mmol), Boc₂O (34.3 g, 158 mmol), and TEA (22 mL, 157.8 mmol) in DCM (200 mL) was added DMAP (1.28


g, 10.5mmol). The reaction was stirred at 40 °C for 16 h. The solvent was removed *in vacuo* and water and DCM were added to separate the phases and the aqueous phase was extracted with DCM. The combined organic extracts were dried over NaSO₄, filtered and concentrated. The crude product was purified by flash chromatography (3 Biotage
 5 100g SiO₂ columns in parallel, gradient: EtOAc/hex 0-30%) to give the title compound as a beige solid (12.7 g, 44%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.11 (s, 1H), 6.74 (s, 1H), 3.88 (d, *J* = 6.5 Hz, 2H), 2.17 (s, 3H), 1.99-1.88 (m, 2H), 1.47 (s, 3H), 1.45 (s, 9H), 1.35 (s, 9H); MS ESI [M + H]⁺ 545.1, calcd for [C₂₂H₃₀BrClN₄O₅ + H]⁺ 545.1.

10 (1*s*,3*s*)-3-(((3-bromo-5-chloropyrazolo[1,5-*a*]pyrimidin-7-yl)(4-methoxybenzyl)amino)methyl)-1-methylcyclobutanol



To a solution of (1*s*,3*s*)-3-(((3-bromo-5-chloropyrazolo[1,5-*a*]pyrimidin-7-yl)amino)methyl)-1-methylcyclobutanol (12.0 g, 34.7 mmol) and 4-methoxybenzyl chloride (5.2 mL, 38.2 mmol) in DMF (50 mL) was added K₂CO₃ (9.6 g, 69.4 mmol). The resulting mixture was stirred at 60 °C for 3 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were washed with water, dried over MgSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 10-100%) using three columns
 20 in parallel. Impure fractions were collected and purified by flash chromatography using the above conditions. Pure fractions were combined and concentrated to give the desired product as a pale yellow solid (13.1 g, 81%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.01 (s, 1H), 7.17 (d, *J* = 8.7 Hz, 2H), 6.67 (d, *J* = 8.7 Hz, 2H), 6.02 (s, 1H), 4.98 (s, 2H), 3.83 (d, *J* = 6.4 Hz, 2H), 3.81 (s, 3H), 2.25-2.16 (m, 3H), 1.76-1.72 (m, 2H), 1.36 (s, 3H); MS
 25 ESI [M + H]⁺ 467.1, calcd for [C₂₀H₂₂BrClN₄O₂ + H]⁺ 467.1.

tert-butyl (3-bromo-5-(((1*S*,2*R*)-2-hydroxycyclohexyl)amino)pyrazolo[1,5-*a*]pyrimidin-7-yl)(cyclopropylmethyl)carbamate




To a solution of using tert-butyl (3-bromo-5-chloropyrazolo[1,5-*a*]pyrimidin-7-yl)(cyclopropylmethyl)carbamate (9.0 g, 22.5 mmol) in NMP (90 mL) was added (1*R*,2*S*)-2-aminocyclohexanol·HCl (4.08 g, 27.0 mmol), and DIPEA (3.47 g, 25.1 mmol). The reaction was divided and sealed into six microwave vials. Each vial

was microwaved for 3 h at 130 °C. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-50%) using three columns in parallel to give the title compound as a beige solid (8.51 g, 79%). ¹H NMR (400 MHz, CDCl₃) δ ppm 7.80 (s, 1H), 6.08 (s, 1H), 5.33-5.19 (m, 1H), 4.26-4.14 (m, 1H), 4.13-4.06 (m, 1H), 3.61 (d, *J* = 7.3 Hz, 2H), 2.51 (br. s, 1H), 1.89-1.62 (m, 8H), 1.55-1.45 (m, 2H), 1.40 (s, 9H), 1.06-0.95 (m, 1H), 0.47-0.38 (m, 2H), 0.17-0.07 (m, 2H); MS ESI [M + H]⁺ 408.3, calcd for [C₂₁H₃₀BrN₅O₃ + H]⁺ 480.2.

10


tert-butyl (3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-7-yl)(cyclopropylmethyl)carbamate



A mixture of tert-butyl (3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)(cyclopropylmethyl)carbamate (400 mg, 1.0 mmol), 3-hydroxypyridine (475 mg, 5.0 mmol), DBU (0.75 mL, 5.0 mmol) in DME (5 mL) was combined and sealed into a microwave vial. The mixture was microwaved for 1 h at 85 °C. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-50%) to give the title compound as a brown oil (367 mg, 80%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.64 (d, *J* = 2.5 Hz, 1H) 8.54 (dd, *J* = 4.5, 1.5 Hz, 1H) 7.97 (s, 1H), 7.78-7.72 (m, 1H), 7.44-7.38 (m, 1H) 6.61 (s, 1H) 3.71 (d, *J* = 7.3 Hz, 2H) 1.39 (s, 9 H), 1.06-1.01 (m, 1H) 0.45 (s, 2H), 0.13-0.09 (m, 2H); MS ESI [M + H]⁺ 460.0, calcd for [C₂₀H₂₂BrN₅O₃ + H]⁺ 460.1.

25


tert-butyl (3-bromo-5-(cyclopentylamino)pyrazolo[1,5-a]pyrimidin-7-yl)((1*s*,3*s*)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate



To a solution of tert-butyl (3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)((1*s*,3*s*)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate (0.45 g, 0.83 mmol) in NMP (4 mL) was added cyclopentylamine (0.085 g, 1.0 mmol), and DIPEA (0.21 g, 1.66 mmol). The reaction was microwaved for 3 h at 130 °C. Water and EtOAc were added to separate the phases and the aqueous phase


was extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 0-60%) to give the title compound as a white solid (0.40 g, 67%).
¹H NMR (400 MHz, CDCl₃) δ ppm 7.79 (s, 1H), 5.94 (s, 1H), 5.08 (br. s, 1H), 4.23 (br.
 5 s, 1H), 3.85-3.72 (m, 2H), 2.28-2.06 (m, 5H), 2.02-1.89 (m, 2H), 1.81-1.61 (m, 5H), 1.48 (s, 4H), 1.44 (s, 9H), 1.36 (s, 9H); MS ESI [M + H]⁺ 594.3, calcd for [C₂₇H₄₀BrN₅O₅ + H]⁺ 594.2.

tert-butyl (3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-7-yl)((1s,3s)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate



A mixture of tert-butyl (3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)((1s,3s)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate (12.7 g, 23.3 mmol), 3-hydroxypyridine (4.43 g, 46.6 mmol), and K₂CO₃ (9.65 g, 69.9 mmol) was combined in DMF (100 mL) and stirred at rt overnight. The reaction mixture was diluted with EtOAc and washed with water, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was triturated with Et₂O and filtered to give the title compound as a beige solid (8.43 g, 14.0 mmol). The mother liquor was concentrated *in vacuo* and purified by flash chromatography (gradient: EtOAc/hex 0-40%) to give the title compound as a light orange solid (4.07 g, 6.74 mmol). The solids obtained through trituration and flash chromatography were combined to give the title compound as a light orange solid (12.50 g, 89%)
 20 ¹H NMR (400 MHz, CDCl₃) δ ppm 8.65 (d, *J* = 2.8 Hz, 1H), 8.55 (dd, *J* = 4.6, 1.4 Hz, 1H), 7.98 (s, 1H), 7.78-7.73 (m, 1H), 7.46-7.40 (m, 1H), 6.50 (s, 1H), 3.89 (d, *J* = 6.8 Hz, 2H),
 25 2.30-2.15 (m, 3H), 2.02-1.93 (m, 2H), 1.50 (s, 3H), 1.45 (s, 9H), 1.38 (s, 9H); MS ESI [M + H]⁺ 604.3, calcd for [C₂₇H₃₄BrN₅O₆ + H]⁺ 604.2.

(1s,3s)-3-(((3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-7-yl)(4-methoxybenzyl)amino)methyl)-1-methylcyclobutanol




To a solution of (1s,3s)-3-(((3-bromo-5-chloropyrazolo[1,5-a]pyrimidin-7-yl)(4-methoxybenzyl)amino)methyl)-1-methylcyclobutanol (13.1 g, 28.2 mmol) and 3-hydroxypyridine (4.0 g, 42.3 mmol) in DMF (70 mL) was added K₂CO₃ (7.8 g, 56.3

mmol). The resulting mixture was stirred at 80 °C for 22 h. A mixture of 60% NaH (0.6 g, 15 mmol) and 3-hydroxypyridine (1.4 g, 14.7 mmol) in DMF was stirred at rt for 30 min before adding to the reaction mixture. The resulting solution was stirred at 100 °C for 20 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were washed with water, dried over MgSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hexanes 10-100%) using three columns in parallel. Impure fractions were collected and purified by flash chromatography (gradient: EtOAc/hexanes 50-100%). Pure fractions were combined and concentrated to give the desired product as a pale yellow solid (9.5 g, 64%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.56 (d, *J* = 4 Hz, 1H), 8.45 (d, *J* = 8 Hz, 1H), 8.00 (s, 1H), 7.69-7.65 (m, 1H), 7.36 (dd, *J* = 4.8, 0.4 Hz, 1H), 7.19 (d, *J* = 8.7 Hz, 2H), 6.85 (d, *J* = 8.7 Hz, 2H), 5.71 (s, 1H), 4.91 (s, 2H), 3.87 (d, *J* = 6.7 Hz, 2H), 3.87 (s, 3H), 2.30-2.15 (m, 3H), 1.77 (t, *J* = 8.1 Hz, 2H), 1.37 (s, 3H); MS ESI [M + H]⁺ 524.2, calcd for [C₂₅H₂₆BrN₅O₃ + H]⁺ 524.1.

15

N-cyclopropyl-4-(7-(((1*s*,3*s*)-3-hydroxy-3-methylcyclobutyl)methyl)(4-methoxybenzyl)amino)-5-(pyridin-3-yloxy)pyrazolo[1,5-*a*]pyrimidin-3-yl)-2-methylbenzamide




20

A mixture of (1*s*,3*s*)-3-(((3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-*a*]pyrimidin-7-yl)(4-methoxybenzyl)amino)methyl)-1-methylcyclobutanol (9.5 g, 18.1 mmol), *N*-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (7.6 g, 25.4 mmol), 2M K₃PO₄ (23 mL, 45.3 mmol) in THF (100 mL) was purged with Ar for 10 min before the addition of PdCl₂dppfDCM (1.5 g, 1.8 mmol). The resulting mixture was heated under reflux in an oil bath at 84 °C for 16 h. The reaction mixture was diluted with DCM and sat. NaHCO₃. The aqueous phase was extracted with DCM and the combined organic extracts were dried over MgSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 50-100%) using three columns in parallel. Impure fractions were collected and purified by flash chromatography using the above conditions. Pure fractions were combined and concentrated to give the desired product as an orange solid (9.6 g, 85%). ¹H NMR (400 MHz, CDCl₃) δ ppm 8.90-8.84 (m, 1H), 8.55 (d, *J* = 4.5 Hz, 1H), 8.29 (s,

1H). 7.88-7.83 (m, 1H), 7.63 (s, 1H), 7.58-7.53 (m, 2H), 7.35-7.31 (m, 1H), 7.23 (d, $J = 8.6$ Hz, 2H), 6.89 (d, $J = 8.6$ Hz, 2H), 6.19 (br. s, 1H), 5.82 (s, 1H), 4.99 (s, 2H), 3.90 (d, $J = 10.0$ Hz, 2H), 3.80 (s, 3H), 2.93-2.90 (m, 1H), 2.40 (s, 3H), 2.36-2.28 (m, 1H), 2.22-2.20 (m, 2H), 1.89-1.88 (m, 1H), 1.80 (t, $J = 8.7$ Hz, 2H), 1.38 (s, 3H), 0.88-0.85 (m, 2H), 0.68-0.64 (m, 2H); MS ESI 619.3 $[M + H]^+$, calcd for $[C_{36}H_{38}N_6O_4 + H]^+$ 619.3.

Preparation of Exemplary Compounds of the Invention

A1: N-cyclopropyl-4-(7-((cyclopropylmethyl)amino)-5-(((1S,2R)-2-hydroxycyclohexyl)amino)pyrazolo[1,5-a]pyrimidin-3-yl)-2-methylbenzamide hydrochloride




A mixture of tert-butyl (3-bromo-5-(((1S,2R)-2-hydroxycyclohexyl)amino)pyrazolo[1,5-a]pyrimidin-7-yl)(cyclopropylmethyl)carbamate (8.48 g, 17.7 mmol), N-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (6.92 g, 23.0 mmol), $PdCl_2dppf$ DCM (1.44 g, 1.76 mmol), and 2M K_3PO_4 (26.6 mL, 106.2 mmol) in THF (60 mL) was divided and sealed into four microwave vials. Each vial was charged with Ar and heated in the microwave at 125 °C for 3 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 20-100%) using two columns in parallel to give a yellow solid.

The above compound was dissolved in DCM (20 mL) and treated with TFA (20 mL) at rt for 1 h. After reaction completion, solvent was removed in vacuo and the crude was redissolved in DCM (20 mL), neutralized with saturated aqueous sodium bicarbonate and extracted with EtOAc. The combined organic extracts were dried over $NaSO_4$, filtered and concentrated. The crude product was purified by flash chromatography (gradient: 50-100% EtOAc/hex then MeOH/DCM 0-10%) and triturated with Et_2O to give the title compound as a free base (white solid). The free base was dissolved in a mixture of DCM (25 mL) and MeOH (50 mL), HCl (1M Et_2O , 2 equiv) was then added slowly. Solvent was removed *in vacuo* to give the title compound as a pale yellow solid in HCl salt (2.09 g, 25% over 2 steps). ^1H NMR (400 MHz,

CD_3OD) δ ppm 8.21 (s, 1H), 7.51-7.37 (m, 3H), 5.67 (br. s, 1H), 4.04-3.82 (m, 2H), 3.46-3.37 (m, 2H), 2.93-2.82 (m, 1H), 2.47 (s, 3H), 1.89-1.64 (m, 6H), 1.61-1.42 (m, 2H), 1.31-1.21 (m, 1H), 0.88-0.78 (m, 2H), 0.70-0.58 (m, 4H), 0.46-0.36 (m, 2H); MS ESI $[M + H]^+$ 475.3, calcd for $[C_{27}H_{34}N_6O_2 + H]^+$ 475.3. HPLC purity: 98% at 254 nm.

5

A2: *N*-cyclopropyl-4-(7-((cyclopropylmethyl)amino)-5-(pyridin-3-yloxy)pyrazolo[1,5-*a*]pyrimidin-3-yl)-2-methylbenzamide hydrochloride




A mixture of *tert*-butyl (3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-*a*]pyrimidin-7-yl)(cyclopropylmethyl)carbamate (367 mg, 0.80 mmol), *N*-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (289 mg, 0.96 mmol), $PdCl_2dppf$ DCM (65 mg, 0.080 mmol), and 2M K_3PO_4 (1.4 mL, 2.8 mmol) in THF (4 mL) was sealed into a microwave vial. The vial was charged with

Ar and heated in the microwave at 130 °C for 4 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by flash chromatography (gradient: EtOAc/hex 20-100%) to give a yellow oil.

The above compound was dissolved in DCM (6 mL) and treated with TFA (2 mL) at rt for 1 h. After reaction completion, solvent was removed *in vacuo* and the crude was redissolved in MeOH (4 mL), filtered, and purified by prep HPLC. The fractions were passed through a PoraPak column and triturated with Et_2O to give the title compound as a free base (white solid, 119 mg). A portion of the free base (67 mg, 0.15 mmol) was then dissolved in MeOH (10 mL), and HCl (1M Et_2O , 2 equiv) was then added slowly. The mixture was concentrated and triturated with Et_2O to give the title compound as a yellow solid in HCl salt (55 mg, 25% yield over 2 steps). ^1H NMR (400 MHz, CD_3OD) δ ppm 9.10 (s, 1H), 8.82 (d, $J = 6.0$ Hz, 1H), 8.73-8.66 (m, 1H), 8.41 (s, 1H), 8.24-8.15 (m, 1H), 7.69 (s, 1H), 7.60 (d, $J = 8.3$ Hz, 1H), 7.23 (d, $J = 8.0$ Hz, 1H), 6.07 (s, 1H), 3.39 (d, $J = 7.0$ Hz, 2H), 2.89-2.78 (m, 1H), 2.29 (s, 3H), 1.37-1.23 (m, 1H), 0.86-0.75 (m, 2H), 0.70-0.56 (m, 4H), 0.45-0.38 (m, 2H); MS ESI $[M + H]^+$ 455.2, calcd for $[C_{26}H_{26}N_6O_2 + H]^+$ 455.2. HPLC purity: 95.9% at 254 nm.

A3: 4-(5-(cyclopentylamino)-7-(((1*s*,3*s*)-3-hydroxy-3-methylcyclobutyl)methyl)amino)pyrazolo[1,5-*a*]pyrimidin-3-yl)-*N*-cyclopropyl-2-methylbenzamide hydrochloride




- 5 A mixture of tert-butyl (3-bromo-5-(cyclopentylamino)pyrazolo[1,5-*a*]pyrimidin-7-yl)((1*s*,3*s*)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate (0.40 g, 0.67 mmol), *N*-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (0.26 g, 0.88 mmol),
- 10 PdCl₂dppfDCM (0.055 g, 0.067 mmol), and 2M K₃PO₄ (1 mL, 2.01 mmol) in THF (4 mL) was charged with Ar and heated in the microwave at 130 °C for 3 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient:
- 15 EtOAc/hex 20-100%) to give a yellow oil.

The above compound was dissolved in DCM (10 mL) and treated with TFA (3 mL) at rt for 3 h. After reaction completion, solvent was removed *in vacuo* and the crude product was dissolved in MeOH (5 mL). The mixture was filtered and purified by prep-HPLC. The compound was passed through a PoraPak cartridge and triturated with Et₂O

20 to give the title compound as a free base (white solid). The free base was dissolved in MeOH (5 mL), and HCl (1M Et₂O, 2 equiv) was then added slowly. Solvent was removed *in vacuo* to give the title compound as a light orange solid in HCl salt (75 mg, 21% over 2 steps). ¹H NMR (400 MHz, CD₃OD) δ ppm 8.18 (s, 1H), 7.42 (s, 3H), 5.52 (br. s, 1H), 4.23-4.11 (m, 1H), 3.68-3.54 (m, 2H), 2.93-2.81 (m, 1H), 2.46 (s, 3H), 2.38-


25 2.27 (m, 1H), 2.26-2.07 (m, 4H), 1.99-1.89 (m, 2H), 1.88-1.59 (m, 6H), 1.35 (s, 3H), 0.86-0.79 (m, 2H), 0.66-0.58 (m, 2H); MS ESI [M + H]⁺ 489.4, calcd for [C₂₈H₃₆N₆O₂ + H]⁺ 489.3. HPLC purity: 99% at 254 nm.

A4: N-cyclopropyl-4-(7-((((1s,3s)-3-hydroxy-3-methylcyclobutyl)methyl)amino)-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-3-yl)-2-methylbenzamide hydrochloride and its free base



- 5 A). Through Boc deprotection: A mixture of tert-butyl (3-bromo-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-7-yl)((1s,3s)-3-((tert-butoxycarbonyl)oxy)-3-methylcyclobutyl)methyl)carbamate (0.23 g, 0.38 mmol), N-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (0.15 g, 0.49 mmol),
10 PdCl₂dppfDCM (0.15 g, 0.49 mmol), and 2M K₃PO₄ (0.57 mL, 1.14 mmol) in THF (4 mL) was charged with Ar and heated in the microwave at 130 °C for 3 h. Water and EtOAc were added to separate the phases and the aqueous phase was extracted with EtOAc. The combined organic extracts were dried over NaSO₄, filtered and concentrated. The crude product was purified by flash chromatography (gradient:
15 EtOAc/hex 20-60%) to give a yellow oil.



The above intermediate was dissolved in DCM (10 mL) and treated with TFA (3 mL) at rt for 3 h. After reaction completion, solvent was removed *in vacuo* and the crude product was dissolved in MeOH (5 mL). The mixture was filtered and purified by prep-HPLC. The compound was passed through a PoraPak cartridge and triturated with Et₂O
20 to give the title compound as a free base (white solid). The free base was dissolved in MeOH (5 mL), and HCl (1 M Et₂O, 2 equiv) was then added slowly. Solvent was removed *in vacuo* to give the title compound as a beige solid in HCl salt (96 mg, 47% over 2 steps). ¹H NMR (400 MHz, CD₃OD) δ ppm 9.14 (br. s, 1H), 8.89-8.82 (m, 1H), 8.79-8.71 (m, 1H), 8.40 (s, 1H), 8.31-8.21 (m, 1H), 7.68 (s, 1H), 7.59 (d, *J* = 9.5 Hz, 1H), 7.23 (d, *J* = 8.0 Hz, 1H), 6.06 (s, 1H), 3.56 (d, *J* = 6.5 Hz, 2H), 2.88-2.79 (m, 1H),
25 2.40-2.31 (m, 1H), 2.29 (s, 3H), 2.26-2.18 (m, 2H), 1.99-1.89 (m, 2H), 1.37 (s, 3H), 0.85-0.76 (m, 2H), 0.63-0.53 (m, 2H); MS ESI [M + H]⁺ 499.3, calcd for [C₂₈H₃₀N₆O₃ + H]⁺ 499.2. HPLC purity: 99.5% at 254 nm.







B). Through PMB deprotection: A mixture of N-cyclopropyl-4-(7-((((1s,3s)-3-hydroxy-3-methylcyclobutyl)methyl)(4-methoxybenzyl)amino)-5-(pyridin-3-yloxy)pyrazolo[1,5-a]pyrimidin-3-yl)-2-methylbenzamide (9.6 g, 15.5 mmol), TFA (50 mL) in DCE

(70 mL) was heated in an oil bath at 50 °C for 4 h. After reaction completion, solvent was removed in vacuo and the crude product was dissolved in a mixture of MeOH/DCM (100 mL/25 mL). 2M Na₂CO₃ (150 mL) was then added and the resulting mixture was stirred at rt for 30 min. The reaction mixture was diluted with DCM and the phases were separated. The aqueous phase was extracted with DCM and the combined organic extracts were washed with water, dried over MgSO₄, filtered and concentrated. The crude product was triturated and sonicated in a mixture of DCM/Et₂O (10 mL/70 mL) to give the title compound as a off white solid in free base (5.9 g, 77%). ¹H NMR (400 MHz, CD₃OD) δ ppm 8.58-8.53 (m, 1H), 8.50-8.46 (m, 1H), 8.36 (s, 1H), 7.86-7.80 (m, 1H), 7.76-7.72 (m, 1H), 7.61-7.55 (m, 2H), 7.18 (d, *J* = 8.0 Hz, 1H), 5.92 (s, 1H), 3.52 (d, *J* = 6.8 Hz, 2H), 2.86-2.77 (m, 1H), 2.38-2.28 (m, 1H), 2.25 (s, 3H), 2.24-2.18 (m, 2H), 1.99-1.88 (m, 2H), 1.37 (s, 3H), 0.84-0.75 (m, 2H), 0.64-0.54 (m, 2H); MS ESI [M + H]⁺ 499.2, calcd for [C₂₈H₃₀N₆O₃ + H]⁺ 499.2. HPLC purity: 96.1% at 235 nm.

The following final compounds were synthesized similarly to the syntheses of A1-4.

Structure	Example number MS calcd; MS ESI [M+H] ⁺ Salt Form; HPLC purity	¹ H NMR
	A5 [C ₂₈ H ₃₀ N ₆ O ₃ +H] ⁺ 499.2; 499.3; free base; 98.9% at 254 nM	(400 MHz, CD ₃ OD) δ ppm 8.55 (dd, <i>J</i> = 2.8, 0.4 Hz, 1H), 8.47 (dd, <i>J</i> = 4.8, 1.6 Hz, 1H), 8.33 (s, 1 H), 7.83-7.80 (m, 1H), 7.72 (s, 1H), 7.58-7.55 (m, 2H), 7.18 (d, <i>J</i> = 8.0 Hz, 1H), 5.89 (s, 1H), 3.49 (d, <i>J</i> = 7.6 Hz, 2H), 2.87-2.78 (m, 2H), 2.28-2.23 (m, 5H), 1.97-1.92 (m, 2H), 1.36 (s, 3H), 0.82-0.77 (m, 2H), 0.62-0.58 (m, 2H).
	A6 [C ₂₇ H ₂₈ N ₆ O ₃ +H] ⁺ 485.2; 485.3; HCl salt; 98.8% at 254 nm	(400 MHz, CD ₃ OD) δ ppm 9.13 (d, <i>J</i> = 2.3 Hz, 1H), 8.86 (d, <i>J</i> = 5.8 Hz, 1H), 8.77-8.69 (m, 1H), 8.40 (s, 1H), 8.26 (dd, <i>J</i> = 8.8, 5.8 Hz, 1H), 7.67 (s, 1H), 7.59 (d, <i>J</i> = 8.0 Hz, 1H), 7.24 (d, <i>J</i> = 8.0 Hz, 1H), 6.07 (s, 1H), 4.53-4.39 (m, 1H), 3.57 (d, <i>J</i> = 8.0 Hz, 2H), 2.89-2.79 (m, 1H), 2.76-2.61 (m, 1H), 2.30 (s, 3H), 2.29-2.21 (m, 2H), 2.21-2.08 (m, 2H), 0.85-0.78 (m, 2H), 0.65-0.58 (m, 2H).

	A8 [C ₂₇ H ₃₄ N ₆ O ₂ +H] ⁺ 475.3; 475.4; HCl salt; 99.3% at 254 nm	(400 MHz, CD ₃ OD) δ ppm 8.22 (s, 1H), 7.51-7.36 (m, 3H), 5.68 (s, 1H), 3.77 (s, 2H), 3.37-3.33 (m, 2H), 2.93-2.84 (m, 1H), 2.47 (s, 3H), 2.07-1.95 (m, 4H), 1.93-1.72 (m, 4H), 1.34-1.19 (m, 1H), 0.88-0.78 (m, 2H), 0.69-0.59 (m, 4H), 0.45-0.37 (m, 2H).
	A11 [C ₂₇ H ₂₈ N ₆ O ₃ + H] ⁺ 485.2; 485.2; HCl salt; 98.6% at 254 nm	(400 MHz, CD ₃ OD) δ ppm 9.14 (s, 1H), 8.85 (d, J = 5.5 Hz, 1H), 8.79-8.72 (m, 1H), 8.40 (s, 1H), 8.30-8.21 (m, 1H), 7.68 (s, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.23 (d, J = 8.0 Hz, 1H), 6.06 (s, 1H), 4.17-4.06 (m, 1H), 3.54 (d, J = 6.5 Hz, 2H), 2.88-2.78 (m, 1H), 2.53-2.43 (m, 2H), 2.29 (s, 4H), 1.81-1.68 (m, 2H), 0.84-0.77 (m, 2H), 0.62-0.56 (m, 2H).
	A12 [C ₂₇ H ₃₄ N ₆ O ₂ +H] ⁺ 475.3; 475.4; HCl salt; 97.5% at 254 nm	(400 MHz, CD ₃ OD) δ ppm 8.18 (s, 1H), 7.55-7.29 (m, 3H), 5.54 (br, s, 1H), 4.50-4.37 (m, 1H), 4.26-4.07 (m, 1H), 3.62 (d, J = 6.8 Hz, 2H), 2.92-2.83 (m, 1H), 2.75-2.61 (m, 1H), 2.46 (s, 3H), 2.31-2.02 (m, 6H), 1.92-1.56 (m, 6H), 0.91-0.75 (m, 2H), 0.71-0.52 (m, 2H).
	A13 [C ₂₈ H ₃₆ N ₆ O ₂ +H] ⁺ 489.3; 489.4 TFA salt; 98.2% at 254 nm	(400 MHz, CD ₃ OD) δ ppm 8.18 (s, 1H), 7.48-7.36 (m, 3H), 5.52 (s, 1H), 4.23-4.10 (m, 1H), 3.65-3.57 (m, 2H), 2.93-2.75 (m, 2H), 2.46 (s, 3H), 2.31-2.21 (m, 2H), 2.20-2.11 (m, 3H), 2.06-1.89 (m, 2H), 1.89-1.50 (m, 6H), 1.37 (s, 3H), 0.87-0.79 (m, 2H), 0.68-0.56 (m, 2H).

Example B: TTK Inhibition Assay

Active TTK was purchased from Invitrogen as an amino terminal GST fusion of full length human TTK. Amino terminal 6 histidine, sumo tagged human TTK (residues 1-275) was expressed in *E. coli*, and purified to >95% homogeneity by Ni²⁺ agarose, gel filtration, and ion exchange chromatography.

TTK activity was measured using an indirect ELISA detection system. GST-TTK (0.68 nM) was incubated in the presence of either 16 μM ATP (Sigma cat# A7699) or 100 μM ATP, 50mM Hepes pH 7.2, 1mM EGTA, 10mM MgCl₂, and 0.1% Pluronic in a 96 well microtitre plate pre-coated with amino terminal 6 histidine, sumo tagged TTK (amino acid residues 1-275).

The reaction was allowed to proceed for 30 minutes, followed by 5 washes of the plate with Wash Buffer (phosphate buffered saline supplemented with 0.2% Tween 20), and incubation for 30 minutes with a 1:3000 dilution of primary antibody (Cell Signaling cat# 9381). The plate was washed 5 times with Wash Buffer, incubated for 30

minutes in the presence of secondary antibody coupled to horse radish peroxidase (BioRad cat# 1721019, 1:3000 concentration), washed an additional 5 times with Wash Buffer, and incubated in the presence of TMB substrate (Sigma cat# T0440). The colourimetric reaction was allowed to continue for 5 minutes, followed by addition of
5 stop solution (0.5 N sulphuric acid), and quantified by detection at 450 nm with either a monochromatic or filter based plate reader (Molecular Devices M5 or Beckman DTX880, respectively).

Compound inhibition was determined at either a fixed concentration (10 μ M) or at a variable inhibitor concentration (typically 0.5 μ M to 0.001 μ M in a 10 point dose
10 response titration). Compounds were pre-incubated in the presence of enzyme for 5 minutes prior to addition of ATP and the activity remaining quantified using the above described activity assay. The % Inhibition of a compound was determined using the following formula; % Inhibition = $100 \times (1 - (\text{experimental value} - \text{background value}) / (\text{high activity control} - \text{background value}))$. The IC_{50} value was determined using
15 a non-linear 4 point logistic curve fit (XLfit4, IDBS) with the formula; $(A + (B / (1 + ((x/C)^D))))$, where A = background value, B = range, C = inflection point, D = curve fit parameter.

In Table 1 below, IC_{50} value ranges for exemplary compounds are given using 100 μ M ATP. The IC_{50} ranges are indicated as "A," "B," and "C," for values less than or equal to 0.1 μ M;
20 those greater than 0.1 μ M and less than or equal to 0.5 μ M; and those greater than 0.5 μ M, respectively. IC_{50} ranges denoted with an asterisk indicated that 16 μ M ATP (Sigma cat# A7699) was used in the assay.

Example C: Cancer Cell Line Data on Exemplary Compounds of the Invention

Breast cancer cells (MDA-MB-468), colon cancer cells (HCT116) and ovarian cancer
25 cells (OVCAR-3) were seeded (1000 to 4000 in 80 μ l per well depending on the cell growth rate) into 96 well plates 24 hours before compound overlay. Compounds were prepared as 10 mM stock solutions in 100% DMSO which were diluted with DMEM (Dulbecco's Modified Eagle's Medium) cell growth Medium (Invitrogen, Burlington, ON, Canada) containing 10% FBS (Fetal Bovine Serum) to concentrations ranging from 50 nM to 250 μ M. Aliquots (20 μ l)
30 from each concentration were overlaid to 80 μ l of the pre-seeded cells in the 96 well plates to make final concentrations of 10 nM to 50 μ M. The cells were cultured for 5 days before the Sulforhodamine B assay (SRB) was performed to determine the compound's cell growth inhibition activity.

Sulforhodamine B (purchased from Sigma, Oakville, ON, Canada) is a water-soluble dye that binds to the basic amino acids of the cellular proteins. Thus, colorimetric measurement of the bound dye provides an estimate of the total protein mass that is related to the cell number. the cells are fixed in situ by gently aspirating off the culture media and adding 50 µl ice cold
5 10% Trichloroacetic Acid (TCA) per well and incubate at 4°C for 30-60 min. The plates are washed with H₂O five times and allowed to air dry for 5 min. Addition of 50 µl 0.4%(w/v) SRB solution in 1% (v/v) acetic acid to each well and incubation for 30 min at RT completes the staining reaction. Following staining, plates are washed four times with 1% acetic acid to remove unbound dye and then allowed to air dry for 5 min. The stain is solubilized with 100 µl
10 of 10 mM Tris pH 10.5 per well. Absorbance is read at 570 nm.

The percentage (%) of relative growth inhibition was calculated by comparing to DMSO treated only cells (100%). GI₅₀'s were determined for compounds with cytotoxic activity. The GI₅₀ was calculated using GraphPad PRISM software (GraphPad Software, Inc., San Diego, CA, USA). GI₅₀ (growth inhibition) is the compound concentration that causes 50% inhibition of cell
15 growth.

In Table 1 below, GI₅₀ value ranges for compound examples against breast cancer cell lines (MDA-MB-468), colon cancer cell lines (HCT116) and ovarian cancer cell lines (OVCAR-3) are given. The example compounds demonstrated varying growth inhibition/cell killing activity against cells of breast cancer, colon cancer, and ovarian cancer. The GI₅₀ ranges are
20 indicated as "A," "B," and "C," for values less than or equal to 0.1 µM; those greater than 0.1 µM and less than or equal to 0.5 µM; and those greater than 0.5 µM, respectively.

Example D: Colon and Ovarian Cancer Tumor-Initiating Cell Data of Exemplary Compounds

Materials and Methods: Non-tissue or tissue cultured treated T-75 flask and 96-well plates were purchased from VWR. Vitamin B-27 supplement, MEM NEAA (minimum essential medium non-essential amino acids), sodium pyruvate, L-glutamine, N2 supplement, penicillin-streptomycin and fungizone/amphotericin B were obtained from Invitrogen. Lipid mixture, heparin and EGF were purchased from Sigma; bFGF
30 from BD Biosciences. Tumor Initiating Cells (TICs) from colon were routinely maintained using non-tissue cultured treated T-75 flasks in DMEM:F12 medium containing 0.2XB-27 supplement, 4ug/ml heparin, 1XMEM NEAA, 1Xsodium pyruvate, 1 mM glutamine, 10 pg/ul bFGF, 20 pg/ul EGF, 1X N2 supplement, lipid mixture, penicillin-streptomycin and fungizone/amphotericin B. Ovarian TICs were
35 routinely maintained using tissue cultured treated T-75 flasks in DMEM:F12 medium

containing 1XB-27 supplement, 4ug/ml heparin, 20 pg/ul bFGF, 20 pg/ul EGF and penicillin-streptomycin.

Assay Protocol: Compounds described herein were dissolved in DMSO and further diluted in cell culture medium for GI₅₀ determination. Colon TICs were trypsinized and seeded into non-tissue cultured treated 96-well plates with 4,000 cells/well. After 24 h, compound was added into the cell culture at different concentrations, and the final concentration of DMSO was adjusted to 0.1%. Cells were then cultured at 37°C for 9 days. Ovarian TICs were trypsinized and seeded into tissue cultured treated 96-well plates with 1,000 cells/well. After 24 h, compound was added into the cell culture at different concentrations, and the final concentration of DMSO was adjusted to 0.1%. Cells were then cultured at 37°C for 6 days. Cell viability was assessed by Alamar Blue assay: 10 ul of Alamar Blue was added into each well. After 4 hours incubation at 37°C, fluorescence was recorded at excitation 544 and emission 590. GI₅₀ (Growth inhibition) was calculated using GraphPad Prism 4.0 software. Cell growth inhibition data for compounds described herein is tabulated below.

In Table 1 below, GI₅₀ value ranges for compound examples against TICs (Colon 12 and Ovarian 2393A) are given. The GI₅₀ ranges are indicated as "A," "B," and "C," for values less than or equal to 0.1 µM; those greater than 0.1 µM and less than or equal to 0.5 µM; and those greater than 0.5 µM, respectively.


Table 1: *In vitro* activity of Compound Examples

Example #	TTK IC ₅₀ Range	Cancer Cell Line GI ₅₀ Range			Tumor Initiating Cell GI ₅₀ Range	
		MDA-MB-468	HCT116	OVCAR-3	Ovarian 2393A	Colon 12
A1	A	A	A	A	A	A
A2	A	A	A	A	ND	ND
A3	A	A	A	A	ND	ND
A4	A	A	A	A	A	ND
A5	A	A	A	B	ND	ND
A6	A	A	A	A	A	A
A8	A	B	B	C	ND	ND
A11	A	A	A	A	ND	ND
A12	A	A	A	A	ND	ND
A13	A	B	A	B	ND	ND

ND – not determined

CLAIMS

1. A compound represented by the following structural formula:



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is $-NH-CH_2-Cy$;


Cy is cyclopropyl or cyclobutyl, wherein the cyclopropyl or cyclobutyl is optionally substituted with one or two groups selected from alkyl and hydroxyl;

R^2 is $-O$ -pyridinyl; $-NH-(C_2-C_6)$ hydroxyalkyl optionally substituted with cyclopropyl or isopropyl; or $-NH-(C_3-C_6)$ cycloalkyl optionally substituted with hydroxyl or (C_1-C_2) hydroxyalkyl;

R^4 is selected from hydrogen, halogen, and (C_1-C_3) alkyl; and


R^d is cyclopropyl.

2. The compound of claim 1, wherein R^4 is chlorine or methyl.
3. The compound of claim 1, wherein the compound is represented by the following structural formula:




or a pharmaceutically acceptable salt thereof.

4. The compound of claim 1, wherein the compound is represented by the following structural formula:




or a pharmaceutically acceptable salt thereof.

5. The compound of claim 1, wherein the compound is represented by the following structural formula:



or a pharmaceutically acceptable salt thereof.

6. The compound of claim 1, wherein the compound is represented by the following structural formula:



or a pharmaceutically acceptable salt thereof.

7. A pharmaceutical composition comprising a compound of any one of the preceding claims and a pharmaceutically acceptable carrier or diluent.