

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

(A)

Before administration
(Aug-13)



(B)

After administration
(Jan-15)




Fig. 7





Fig. 8

(A)

Before administration
(Apr-14)



(B)

After administration
(Mar-15)



Fig. 9





Fig. 10

(A)

Before administration
(Jul-14)



(B)

After administration
(Nov-14)



Fig. 11




Fig. 12

(A)

Before administration
(Aug-13)



(B)

After administration
(May-14)



Fig. 13




Fig. 14

(A)

Before administration
(Apr-14)



(B)

After administration
(Feb-15)




Fig. 15




Fig. 16





Fig. 17

(A)

Before administration
(Dec-14)



(B)

After administration
(Feb-15)




Fig. 18





Fig. 19

(A)

Before administration
(Jun-14)



(B)

After administration
(Jun-15)





Fig. 20



- G1: PBS (control)
- ◆ G2: gefitinib 10 mg/kg
- ▲ G3: Bestatin 30 mg/kg
- G4: gefitinib 10 mg/kg + Bestatin 0.1 mg/kg
- * G5: gefitinib 10 mg/kg + Bestatin 0.3 mg/kg
- G6: gefitinib 10 mg/kg + Bestatin 1 mg/kg
- ◇ G7: gefitinib 10 mg/kg + Bestatin 3 mg/kg
- △ G8: gefitinib 10 mg/kg + Bestatin 10 mg/kg
- G9: gefitinib 10 mg/kg + Bestatin 30 mg/kg