

WO 2012/102875

PCT/US2012/021181

-1-

ANALGESIC COMPOUNDS, METHODS, AND FORMULATIONS

Opiates, a class of centrally acting compounds, are the most frequently used agents for pain control and alleviation, and which act upon one or more of the human or mammal opiate receptors. Technically, opiates are the natural alkaloids found in the resin of the opium poppy, but current usage of the term includes synthetic variations, termed opioids. Opiates are narcotic agonistic analgesics and are drugs including or derived from opium, such as morphine, codeine, and many synthetic congeners of morphine, with morphine and hydrocodone preparations being the most widely used opiates. Opiates are natural and synthetic drugs with morphine-like actions, and are subject to control under U.S. Federal narcotics law (scheduled drugs) and the laws of most other nations and international organizations because of their addicting properties and the subsequent destructive toll exacted on the abusers and those with any connection to them.

Tramadol is a synthetic analog of the phenanthrene alkaloid codeine and, as such, is an opioid and also a prodrug (codeine is metabolized to morphine, tramadol is converted to M-1 also called O-desmethyltramadol). Tramadol, like the opiates, is associated with adverse effects, such as physical and psychological dependence, severe withdrawal symptoms, as well as other somewhat less serious side-effects, including nausea, vomiting, sweating, constipation, and drowsiness.

While the opiates and related drugs clearly serve useful purposes, alternative therapies and compounds for alleviation of pain are desirable due to the known problems of the opiates. Particularly, compounds which are not scheduled, allow for lower dosing frequency, and/or do not exhibit the side-effects either at all or to the degree associated with the opiates and related drugs, would provide such alternative therapies.

WO2002/26694 relates to O-substituted 6-methyl-tramadol derivatives, processes for their production, medicaments containing the derivatives, and uses of the derivatives for treating pain.

EP112787 relates to new esters derived from substituted phenyl-cyclohexyl compounds, which are derived from tramadol.

WO2003/048113 relates to tramadol analogs useful for the treatment of CNS-related disorders, including pain, anxiety, depression and attention deficit disorder.

EP2022778 relates to a (R,R)-tramadol-(S)-naproxene salt, a crystalline form of (R,R)-tramadol-(S)-naproxene salt, their processes for preparation and uses as medicament for the treatment of pain.

GB997399 relates to phenol ethers that contain basic groups and to methods for preparing them.

US5564100 relates to cycloalkene compounds and processes for their manufacture.


US3652589 relates to cycloalkanol-substituted phenol ethers having basic groups, to pharmaceutical compositions containing the compounds, processes for manufacturing the compounds, and methods of using the compounds.

Arzneimittel-Forschung/Drug Research, Vol. 28, No. 1a (1978), pages 107-113 relates to phenyl substituted aminomethylcycloalkanol compounds tested for analgesic properties.


Journal of the Mexican Chemical Society, Vol. 49, No. 4, (2005), pages 324-327 relates to tramadol and derivatives of tramadol, and processes for preparing them.

Summary

In a first aspect, there is provided a compound, or salt thereof, of formula I:



wherein A is



R^1 is hydrogen, C_1-C_5 alkyl, C_1-C_5 haloalkyl, C_1-C_5 alkanol, $-(C_1-C_5$ alkyl)phenyl, or phenyl, or a group of the formula $-C(O)-R^{12}$, where R^{12} is C_1-C_5 alkyl, C_1-C_5 alkoxy, C_1-C_5 haloalkyl, C_1-C_5 alkanol, $-(C_1-C_5$ alkyl)phenyl, or phenyl;

R^2 is hydrogen, C_1-C_5 alkyl, C_1-C_5 alkoxy, halogen, C_1-C_5 haloalkyl, or C_1-C_5 haloalkoxy;

R^3 is hydrogen, C_1-C_5 alkyl, C_1-C_5 alkoxy, halogen, C_1-C_5 haloalkyl, or C_1-C_5 haloalkoxy;

R^4 is hydrogen, C_1-C_5 alkyl, or $-(C_1-C_5$ alkyl)phenyl;

R^5 is hydrogen, C_1-C_5 alkyl, or $-(C_1-C_5$ alkyl)phenyl;

R^6 is hydrogen, hydroxy, or is absent;

R^7 is hydrogen;

R^8 is hydrogen or methyl;

R^9 is hydrogen or methyl;

R^{10} is hydrogen;

R^{11} is hydrogen or C_1-C_5 alkyl;

provided that one of I), II), and III) is present, wherein I), II) and III) are:

I) R^7 and R^{10} combine to form $-CH_2-$ or $-(CH_2)_2-$;

II) R^8 and R^9 combine to form a cyclopropyl group with the carbon to which they are attached; and

III) R^{10} and R^{11} combine to form $-\text{CH}_2-$ or $-(\text{CH}_2)_3-$.


In a second aspect, there is provided a pharmaceutical formulation comprising a compound as defined herein and one or more pharmaceutically acceptable carriers.

In a third aspect, there is provided the use of an effective amount of a compound as defined herein in the manufacture of a medicament for controlling pain in a mammal.

In a fourth aspect, there is provided a compound or salt as defined herein for use in therapy.


Detailed Disclosure

Provided are analgesic compounds, and salts thereof, of formula I:



wherein A is

-2-



R^1 is hydrogen, C_1 - C_5 alkyl, C_1 - C_5 alkoxy, C_1 - C_5 haloalkyl, C_1 - C_5 alkanol, $-(C_1$ - C_5 alkyl)phenyl, or phenyl, or a group of the formula $-C(O)-R^{12}$, where R^{12} may be C_1 - C_5 alkyl, C_1 - C_5 alkoxy, C_1 - C_5 haloalkyl, C_1 - C_5 alkanol, $-(C_1$ - C_5 alkyl)phenyl, or phenyl;

R^2 is hydrogen, C_1 - C_5 alkyl, C_1 - C_5 alkoxy, halogen, C_1 - C_5 haloalkyl, or C_1 - C_5 haloalkoxy;

R^3 is hydrogen, C_1 - C_5 alkyl, C_1 - C_5 alkoxy, halogen, C_1 - C_5 haloalkyl, or C_1 - C_5 haloalkoxy;

R^4 is hydrogen, C_1 - C_5 alkyl, or $-(C_1$ - C_5 alkyl)phenyl;

R^5 is hydrogen, C_1 - C_5 alkyl, or $-(C_1$ - C_5 alkyl)phenyl;

R^6 is hydrogen, hydroxy, or is absent;

R^7 is hydrogen;

R^8 is hydrogen or methyl;

R^9 is hydrogen or methyl;

R^{10} is hydrogen;

R^{11} is hydrogen or C_1 - C_5 alkyl;


or R^7 and R^{10} combine to form $-CH_2-$ or $-(CH_2)_2-$;

or R^8 and R^9 combine to form a cyclopropyl group with the carbon to which they are


attached;

or R^{10} and R^{11} combine to form $-CH_2-$ or $-(CH_2)_3-$.


More particularly, A may be:



-3-



Of the first four definitions immediately above for A, the following are preferred:



C_1 – C_5 alkyl refers to straight chain and branched alkyls having one to five carbon atoms, and includes methyl, ethyl, propyl, n-butyl, iso-butyl, pentyl, isopentyl, and neopentyl.

C_1 – C_5 alkoxy refers to straight chain and branched alkoxy groups having one to five carbon atoms, and includes methoxy, ethoxy, propoxy, n-butoxy, iso-butoxy, pentoxy, isopentoxy, and neopentoxy.

Halogen or halo refers to fluorine, bromine, chlorine, and iodine.



Haloalkyl as used herein refers to an alkyl (as noted above) substituted with one or more halo atoms. Such groups include trifluoromethyl, methylchloride, dichloromethyl, pentylchloride, butyl chloride, and isopropyl chloride.

Haloalkoxy refers to an alkoxy group, as described herein, which is substituted with one to six halo groups. Examples of fluoroalkoxy groups include fluoromethoxy, difluoromethoxy, trifluoromethoxy, pentafluoroethoxy, and trifluoroethoxy.

C_1 – C_5 alkanols refer to methanol, ethanol, propanol, or methoxyethanol.


In the definition of A:

-4-



the portion

5



PivCl refers to pivaloyl chloride.

Controlling pain refers to either suppressing, inhibiting, ameliorating, reducing, or eliminating pain, its severity, and/or duration. As such, the invention is applicable to the alleviation of existing pain as well as the suppression or inhibition of pain which would otherwise ensue from an imminent pain-causing event. The pain being alleviated can be chronic or acute.

Mammal includes both human or non-human mammals. Non-human mammals include domestic animals, such as livestock animals and companion animals. Livestock animals include cattle, camellids, pigs, sheep, goats, and horses. Companion animals include dogs, rabbits, cats, and other pets owned and maintained in close association with humans as part of the human-animal bond.

Effective amount refers to the amount of a compound of formula I, or a salt thereof, sufficient to control or alleviate pain in a mammal in need thereof, and as such will depend upon several factors. Ranges for a compound of formula I, or a salt thereof, in the methods include from 0.01 to 1000 mg/kg and more desirably, 0.1 to 100 mg/kg of the animal's body weight. The frequency of the administration will also be dependent upon several factors, and can be a single dose administered once a day or once a week for a duration determined by the attending

-5-

doctor or veterinarian. The dose can be also be split into two or more smaller doses given in a timeframe to result in the control or alleviation of pain.

Pharmaceutically acceptable as used in this application, for example with reference to salts and formulation components such as carriers, includes “veterinarily acceptable”, and thus includes both human and animal applications independently.

Pharmaceutically acceptable salts, and common methodology for preparing them, are known in the art. See, e.g., P. Stahl, *et al.*, HANDBOOK OF PHARMACEUTICAL SALTS: PROPERTIES, SELECTION AND USE, (VCHA/Wiley-VCH, 2002); S.M. Berge, *et al.*, “Pharmaceutical Salts,” *Journal of Pharmaceutical Sciences*, Vol. 66, No. 1, January 1977. A preferred salt is the hydrochloride salt.

The compounds of formula I and their salts may be formulated as pharmaceutical compositions for systemic administration. Such pharmaceutical compositions and processes for making the same are known in the art for both humans and non-human mammals. See, e.g., REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY, (A. Gennaro, *et al.*, eds., 19th ed., Mack Publishing Co., 1995). Additional active ingredients may be included in the formulation containing a compound of formula I or a salt thereof.

Carrier is used herein to describe any ingredient other than the active component(s) in a formulation. The choice of carrier will to a large extent depend on factors such as the particular mode of administration, the effect of the carrier on solubility and stability, and the nature of the dosage form.


The compounds of the invention may be made by the following described procedures, as well as the procedures described in Selnick, H. C.; Bourgeois, M. L.; Butcher, J. W.; Radzilowski, E. M. *Tetrahedron Lett.* **1993**, 34, 2043; Alvarado, C; Guzmán, A.; Díaz, E.; Patino, R. *J. Mex. Chem. Soc.* **2005**, 49, 324; Evans, G. R.; Paloma, Fernández, D.; Henshilwood, J. A., Lloyd, S.; Nicklin, C. *Org. Process Res. Dev.* **2002**, 6, 729; and Mohacsi, E.; O’Brien, J. P.; Blount, J. F. *J.Heterocycl.Chem.* 1990, 27, 1623.

The invention provides a pharmaceutical formulation comprising a compound of formula I, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers. The pharmaceutical formulation may further comprise at least one additional active ingredient. A pharmaceutical formulation may be a human pharmaceutical formulation or a veterinary pharmaceutical formulation.


-6-

The invention provides a method of controlling pain in a mammal in need thereof comprising administering an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, to said mammal. The method may further provide administering at least one other active ingredient to said mammal. The mammal may be a human or non-human mammal, and further may be a companion animal, such as a dog or cat.

For compounds of the formula Ia, below, Schemes A-C and Preparations and/or Examples 1-76 illustrate methods of preparing them.




Scheme A

Preparation 1

Synthesis of 2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-one


-7-



Stir a mixture of bicyclo[3.2.1]octan-3-one (5.2 g, 41.9 mmol), (HCHO)_n (1.51 g, 50.3 mmol), dimethylamine hydrochloride (3.42 g, 41.9 mmol) and 0.5 mL of *conc.* HCl in MeCN (50 mL) at 80 °C for 2 hours. After removal of the solvent under vacuum, dissolve the residue in H₂O (20 mL) and wash with EtOAc (20 mL x 3). Basify the aqueous solution with NaOH to pH = 10. Extract the resultant aqueous mixture with EtOAc (60 mL x 3). The combined organic layers are washed with brine (10 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum to give crude 2-dimethylaminomethyl- bicyclo[3.2.1]octan-3-one as brown oil (5.9 g, yield: 78.7%). MS (m/z): 182 (M+1).

Example 2

Synthesis of 2-dimethylaminomethyl-3-(3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol.








Add dropwise a solution of *t*-BuLi (19.2 mL, 25.0 mmol) in hexane *via* syringe to a solution of 1-bromo-3-methoxy-benzene (3.74 g, 20.0 mmol) in THF (60 mL) at -78 °C under N₂. After being stirred at -78 °C for 1 hour, add dropwise a solution of 2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-one (1.81 g, 10.0 mmol) in THF (10 mL) to the reaction mixture and stir the resultant mixture at -78 °C for additional 1 hour. Quench the reaction with saturated aqueous NH₄Cl solution (20 mL). Extract the aqueous mixture with EtOAc (50 mL x 3). The combined organic layers are washed with brine (15 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by silica gel chromatography (CH₂Cl₂:MeOH = 30:1) to afford






-8-

2-dimethylaminomethyl-3-(3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol as white solid (1.49 g, yield: 51%). MS (m/z): 290 (M+1).

The following compounds may be prepared essentially by the method of Example 2.

Ex. No.	Chemical name	Structure	Physical data
3	2-Dimethylaminomethyl-3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 308 (M+1).
4	2-Dimethylaminomethyl-3-(5-methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 374 (M+1).
5	2-Dimethylaminomethyl-3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 308 (M+1).
6	2-Dimethylaminomethyl-3-(4-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 308 (M+1).
7	2-Dimethylaminomethyl-3-(2-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 308 (M+1).
8	2-Dimethylaminomethyl-3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 304 (M+1).


-9-

Ex. No.	Chemical name	Structure	Physical data
9	3-(2-Chloro-5-methoxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol		MS (m/z): 324 (M+1).
10	3-(3-Chloro-5-methoxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol		MS (m/z): 324 (M+1).
11	2-Dimethylaminomethyl-3-(5-methoxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 304 (M+1).
12	2-Dimethylaminomethyl-3-(3-methoxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 304 (M+1).
13	2-Dimethylaminomethyl-3-(3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 276 (M+1).

Example 14




Synthesis of 2-dimethylaminomethyl-3-(3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride.




-10-





Add H₂O (100 mg, 5.56 mmol) and TMSCl (361 mg, 3.34 mmol) to a solution of 2-dimethylaminomethyl-3-(3-methoxy-phenyl)-bicyclo [3.2.1] octan-3-ol (877 mg, 3.03 mmol) in 2-butanone (60 mL). Stir the mixture at ambient temperature for 12 hours. Concentrate the mixture under vacuum to give 2-dimethylaminomethyl-3-(3-methoxy- phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride as white solid (986 mg, Yield: 100%). ¹H NMR (400 MHz, D₂O) δ 7.23-7.27 (t, *J* = 16.4, 1H), 7.00-7.02 (d, *J* = 8.0, 1H), 6.95-6.96 (t, *J* = 4.0, 1H), 6.78-6.80 (d, *J* = 10.4, 1H), 3.71 (s, 3H), 3.05-3.19 (m, 1H), 2.52-2.58 (m, 4H), 2.14-2.26 (m, 4H), 2.09-2.14 (m, 2H), 2.05-2.06 (d, *J* = 2.4, 1H), 1.89-1.92 (m, 1H), 1.78-1.80 (m, 1H), 1.51-1.68 (m, 4H), 1.40-1.52 (m, 1H).

The following compounds may be prepared essentially by the method of Example 14.



Ex. No.	Chemical name	Structure	Physical data
15	2-Dimethylaminomethyl-3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.92 (s, 1H), 6.88-6.85 (d, J = 12.0, 1H), 6.62-6.59 (d, J = 12.0, 1H), 3.82 (s, 3H), 2.80 (s, 3H), 2.74-2.70 (d, J = 16.0, 1H), 2.51-2.48 (m, 4H), 2.40 (s, 1H), 2.30-2.25 (m, 3H), 2.27-2.24 (d, J = 12.0, 2H), 2.17-2.15 (m, 1H), 1.81-1.64 (m, 4H).
16	2-Dimethylaminomethyl-3-(5-methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, D_2O) δ 7.24-7.26 (d, J = 8.4, 1H), 7.13 (s, 1H), 6.86-6.88 (d, J = 8.8, 1H), 3.73 (s, 3H), 3.11-3.17 (m, 1H), 2.65 (s, 3H), 2.48-2.55 (m, 2H), 2.18-2.39 (m, 6H), 1.92-1.94 (m, 1H), 1.60-1.75 (m, 5H), 1.48-1.56 (m, 1H).
17	2-Dimethylaminomethyl-3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, D_2O) δ 6.98-6.70 (m, 2H), 6.79-6.81 (m, 1H), 3.70 (s, 3H), 3.08-3.11 (t, J = 12.8, 1H), 2.55-2.62 (m, 4H), 2.25-2.48 (m, 6H), 2.17 (s, 1H), 1.85-1.94 (m, 1H), 1.49-1.57 (m, 6H).

18	2-Dimethylaminomethyl-3-(4-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.12-7.06 (m, 2H), 7.08 (s, 1H), 3.82 (s, 3H), 2.64-2.57 (m, 3H), 2.35 (m, 2H), 2.31 (s, 1H), 2.21-2.11 (m, 2H), 2.12 (s, 6H), 1.87-1.74 (m, 2H), 1.66-1.48 (m, 3H).
19	2-Dimethylaminomethyl-3-(2-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.11-7.01 (m, 3H), 3.77 (s, 3H), 3.14-3.09 (m, 1H), 2.63-2.57 (m, 4H), 2.45-2.26 (m, 5H), 2.18 (s, 1H) 2.06 (s, 1H), 1.95-1.90 (m, 1H), 1.79-1.74 (m, 1H), 1.69-1.53 (m, 4H), 1.51-1.47 (m, 1H).
20	2-Dimethylaminomethyl-3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 6.86 (s, 1H), 6.76 (s, 1H), 6.65 (s, 1H), 3.69 (s, 3H), 3.03-3.09 (m, 1H), 2.60 (s, 3H), 2.52-2.55 (d, <i>J</i> = 13.2, 1H), 2.33 (s, 3H), 2.14-2.24 (m, 6H), 2.04-2.08 (d, <i>J</i> = 14.4, 1H), 1.90- 1.94 (m, 1H), 1.76-1.80 (m, 1H), 1.58-1.66 (m, 4H), 1.45-1.49 (m, 1H).


-13-

21	3-(2-Chloro-5-methoxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.44-7.45 (d, <i>J</i> = 3.2, 1H), 7.32-7.34 (d, <i>J</i> = 8.8, 1H), 6.86-6.89 (m, 1H), 3.73 (s, 3H), 3.27-3.30 (t, <i>J</i> = 13.6, 1H), 3.05-3.07 (d, <i>J</i> = 9.2, 1H), 2.90-2.94 (m, 1H), 2.81 (s, 3H), 2.71-2.75 (d, <i>J</i> = 13.6, 1H), 2.47 (m, 5H), 2.29-2.35 (m, 1H), 2.03-2.12 (m, 2H), 1.59-1.79 (m, 4H).
22	3-(3-Chloro-5-methoxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.12 (s, 1H), 7.02 (t, <i>J</i> = 2.0, 1H), 6.85-6.86 (t, <i>J</i> = 4.0, 1H), 3.82 (s, 3H), 3.36 (m, 1H), 2.68-2.71 (m, 7H), 2.36-2.37 (m, 2H), 2.25-2.28 (m, 2H), 2.14-2.18 (m, 1H), 2.03-2.09 (t, <i>J</i> = 23.6, 1H), 1.70-1.86 (m, 4H), 1.59-1.67 (m, 1H).


-14-

23	2-Dimethylaminomethyl-3-(5-methoxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400MHz, D ₂ O) δ 7.26 (s, 1H), 7.21-7.19 (d, <i>J</i> = 8.0, 1H), 6.88-6.86 (d, <i>J</i> = 8.0, 1H), 3.81 (s, 3H), 3.23-3.17 (m, 1H), 3.71-3.66 (m, 3H), 2.59-2.51 (m, 6H), 2.41 (s, 4H), 2.32 (s, 1H), 2.04-2.01 (m, 1H), 1.90-1.88 (m, 2H), 1.75-1.67 (m, 2H), 1.63-1.59 (m, 2H).
24	2-Dimethylaminomethyl-3-(3-methoxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.23-7.21 (d, <i>J</i> = 4.0, 1H), 7.05 (s, 1H), 7.03 (s, 1H), 3.85 (s, 3H), 3.22-3.18 (m, 1H), 2.71-2.64 (m, 4H), 2.42 (s, 3H), 2.36-2.27 (m, 3H), 2.20-2.16 (m, 4H), 2.08-2.02 (m, 1H), 1.94-1.90 (m, 1H), 1.88-1.75 (m, 4H), 1.60-1.56 (m, 1H).

-15-


25	2-Dimethylaminomethyl-3-(3-hydroxy-phenoxy)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.14-7.18 (t, <i>J</i> = 15.6, 1H), 6.91-6.93 (d, <i>J</i> = 7.2, 1H), 6.85 (s, 1H), 6.64-6.66 (d, <i>J</i> = 6.8, 1H), 3.00-3.06 (t, <i>J</i> = 22.8, 1H), 2.53-2.57 (m, 4H), 2.28 (s, 3H), 2.22 (s, 1H), 2.12-2.14 (m, 2H), 2.01-2.05 (d, <i>J</i> = 13.6, 1H), 1.90 (m, 1H), 1.76 (m, 1H), 1.60-1.65 (m, 4H), 1.45 (m, 1H).
----	--	---	---

Scheme B



Preparation 26






Synthesis of (3-bromo-4-fluorophenoxy)-tert-butyl-dimethyl-silane










-16-

Add TBSCl (3.37 g, 22.37 mmol) and imidazole (1.9 g, 27.96 mmol) to a solution of 3-bromo-4-fluoro-phenol (3.56 g 18.64 mmol) in CH₂Cl₂ (60 ml). Stir the solution at ambient temperature for 3 hours. Quench the reaction with water (30 mL). Extract the aqueous layer with CH₂Cl₂ (30 mL x 3). The combined organic layers are washed with brine, dried over Na₂SO₄, concentrated under vacuum. The residue is purified by silica gel chromatography (petroleum ether) to give (3-bromo-4-fluoro-phenoxy)-tert-butyl-dimethyl-silane as colorless oil (5.81 g, 99%), MS (m/z): 303 (M-1).

10 The following compounds may be prepared essentially by the method of Preparation 26.

Prep No.	Chemical name	Structure	Physical data
27	(3-Bromo-5-fluoro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 303 (M-1).
28	(3-Bromo-5-trifluoromethyl-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 353 (M-1).
29	(3-Bromo-4-trifluoromethoxy-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 369 (M-1).
30	(5-Bromo-2-trifluoromethyl-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 353 (M-1).
31	(5-Bromo-2-fluoro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 303 (M-1).


-17-

Prep No.	Chemical name	Structure	Physical data
32	(3-Bromo-2-fluoro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 303 (M-1).
33	(3-Bromo-5-methyl-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 299 (M-1).
34	(3-Bromo-4-chloro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 319(M-1).
35	(3-Bromo-5-chloro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 319 (M-1).
36	(3-Bromo-4-methyl-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 299 (M-1).
37	(5-Bromo-2-methyl-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 299 (M-1).
38	(5-Bromo-2,3-difluoro-phenoxy)-tert-butyl-dimethyl-silane		MS (m/z): 321 (M-1).

Example 39

Synthesis of 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan -3-ol

-18-










Cool a solution of (3-bromo-4-fluoro-phenoxy)-tert-butyl-dimethyl-silane (5.81 g, 19.03 mmol) in THF (100 mL) to -78 °C under N₂. Then add dropwise a solution of *t*-BuLi (14.6 mL, 19.03 mmol) in hexane *via* syringe to the reaction solution. After being stirred at -78 °C for 1 hour, add dropwise a solution of 2-dimethylaminomethyl-bicyclo [3.2.1]octan-3-one (2.87 g, 15.86 mmol) in THF (1.5 mL) to the reaction mixture and stir the mixture at -78 °C for additional 1 hour. Quench the reaction with 60 mL of diluted HCl (2N), and stir at ambient temperature for 2 hours. Basify the resultant mixture with K₂CO₃ to pH = 9, and extract with EtOAc (100 mL X 3).

The combined organic layers are washed with brine, dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by silica gel chromatography (CH₂Cl₂ to CH₂Cl₂:MeOH = 10:1) to afford 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol as white solid (2.38 g, yield: 51.2%). MS (m/z): 294 (M+1).

The following compounds may be prepared essentially by the method of Example 39.

Ex. No.	Chemical name	Structure	Physical data
40	2-Dimethylaminomethyl-3-(3-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 294 (M+1).
41	2-Dimethylaminomethyl-3-(3-hydroxy-5-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 344 (M+1).

Ex. No.	Chemical name	Structure	Physical data
42	2-Dimethylaminomethyl-3-(5-hydroxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 360 (M+1).
43	2-Dimethylaminomethyl-3-(3-hydroxy-4-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 344 (M+1).
44	2-Dimethylaminomethyl-3-(4-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 294 (M+1).
45	2-Dimethylaminomethyl-3-(2-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 294 (M+1).
46	2-Dimethylaminomethyl-3-(3-hydroxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 290 (M+1).
47	3-(2-Chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol		MS (m/z): 310 (M+1).
48	3-(3-Chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol		MS (m/z): 310 (M+1).


-20-

Ex. No.	Chemical name	Structure	Physical data
49	2-Dimethylaminomethyl-3-(5-hydroxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 290 (M+1).
50	2-Dimethylaminomethyl-3-(3-hydroxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol		MS (m/z): 290 (M+1).
51	3-(3,4-Difluoro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol		MS (m/z): 312 (M+1).

Example 52

Synthesis of 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol

5




10



Add H₂O (175.1 mg, 9.73 mmol) and TMSCl (1.057 g, 9.73 mmol) to a solution of 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol (2.38 g, 8.11 mmol) in 2-butanone (60 mL). Stir the reaction mixture at ambient temperature for 3 hours. After removal solvent by evaporation, wash the residue with EtOAc to give 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride as white solid (2.23 g, Yield: 83.5%). ¹H NMR (400 MHz, D₂O) δ 6.89-6.93 (m, 2H), 6.66-6.70 (m, 1H), 3.10-3.14 (m, 1H), 2.59-2.65 (m, 4H), 2.42-2.44 (d, *J* = 8.0, 1H), 2.36-2.41 (d, *J* = 20.0, 3H), 2.30-2.33 (d, *J* = 12.0, 2H), 2.18 (s, 1H), 1.89-1.95 (m, 1H), 1.74-1.78 (m, 1H), 1.51-1.67 (m, 4H), 1.47-1.50 (m, 1H).



The following compounds may be prepared essentially by the method of Example 52.



Ex. No.	Chemical name	Structure	Physical data
53	2-Dimethylaminomethyl-3-(3-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.74-6.79 (m, 2H), 6.39-6.43 (d, $J = 14.4$, 1H), 3.26-3.23 (t, $J = 14.0$ Hz, 1H), 2.74-2.79 (m, 4H), 2.50 (s, 3H), 2.38 (s, 2H), 2.29-2.31 (m, 1H), 2.16-2.27 (m, 2H), 2.03-2.09 (m, 1H), 1.71-1.85 (m, 4H), 1.60-1.67 (m, 1H).
54	2-Dimethylaminomethyl-3-(3-hydroxy-5-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, D_2O) δ 7.40-7.42 (d, $J = 4.0$, 1H), 7.08-7.10 (d, $J = 4.0$, 1H), 6.93 (s, 1H), 2.41-2.53 (m, 6H), 2.24 (s, 4H), 2.18 (s, 1H), 2.10 (s, 1H), 1.83-1.91 (m, 1H), 1.66-1.72 (m, 2H), 1.39-1.55 (m, 4H).



-22-

Ex. No.	Chemical name	Structure	Physical data
55	2-Dimethylaminomethyl-3-(5-hydroxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.08-7.11 (d, <i>J</i> ₁ = 8.8, <i>J</i> ₂ = 1.2, 1H), 6.96-6.97 (d, <i>J</i> = 3.2, 1H), 6.65-6.68 (d, <i>J</i> = 12.0, 1H), 3.02-3.08 (t, <i>J</i> = 23.2 Hz, 1H), 2.46-2.57 (m, 4H), 2.20-2.41 (m, 6H), 2.10 (s, 1H), 1.83-1.87 (m, 1H), 1.48-1.67 (m, 5H), 1.38-1.41 (m, 1H).


Ex. No.	Chemical name	Structure	Physical data
56	2-Dimethylaminomethyl-3-(3-hydroxy-4-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.45-7.47 (d, <i>J</i> = 8.0, 1H), 6.98-7.01 (m, 2H), 3.03-3.09 (t, <i>J</i> = 23.6, 1H), 2.60 (s, 3H), 2.48-2.51 (d, <i>J</i> = 13.2, 1H), 2.30 (s, 3H), 2.23 (s, 1H), 2.12-2.15 (d, <i>J</i> = 8.4, 2H), 2.00-2.04 (m, 1H), 1.87-1.91 (m, 1H), 1.73-1.77 (m, 1H), 1.57-1.67 (m, 4H), 1.44-1.47 (m, 1H).
57	2-Dimethylaminomethyl-3-(4-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.01-7.06 (m, 2H), 6.90-6.91 (m, 1H), 3.03-3.07 (t, <i>J</i> = 13.6, 1H), 2.57-2.62 (m, 4H), 2.05-2.32 (m, 7H), 1.88-1.92 (m, 1H), 1.72-1.79 (m, 1H), 1.62-1.68 (m, 4H), 1.43-1.51 (m, 1H).


Ex. No.	Chemical name	Structure	Physical data
58	2-Dimethylaminomethyl-3-(2-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.00-7.96 (m, 2H), 6.88(s, 1H), 3.10-3.18 (m, 1H), 2.78-2.79 (d, <i>J</i> = 4.0, 4H), 2.27-2.47 (m, 6H), 2.18 (s, 1H), 1.80-1.99 (m, 1H), 1.76 (s, 1H), 1.66-1.68 (d, <i>J</i> = 8.0, 4H), 1.49 (s, 1H).
59	2-Dimethylaminomethyl-3-(3-hydroxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400MHz, D ₂ O) δ 6.80 (s, 1H), 6.69 (s, 1H), 6.53 (s, 1H), 3.03-3.06 (t, <i>J</i> = 13.2, 1H), 2.55-2.61 (m, 4H), 2.32 (s, 3H), 2.24 (s, 1H), 2.04-2.18 (m, 6H), 1.77-1.90 (m, 1H), 1.24-1.33 (m, 1H), 1.55-1.69 (m, 4H), 1.41-1.52 (m, 1H).

Ex. No.	Chemical name	Structure	Physical data
60	3-(2-Chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.34-7.35 (d, <i>J</i> = 2.4, 1H), 7.20-7.21 (d, <i>J</i> = 4.8, 1H), 6.69-6.72 (m, 1H), 3.25-3.31 (m, 1H), 3.02-3.05 (d, <i>J</i> = 9.2, 1H), 2.90-2.94 (m, 1H), 2.73-2.77 (m, 4H), 2.41 (m, 5H), 2.28 (m, 1H), 1.95-2.13 (m, 2H), 1.62-1.78 (m, 4H).
61	3-(3-Chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.01 (t, <i>J</i> = 3.6, 1H), 6.88-6.89 (t, <i>J</i> = 3.6, 1H), 6.69-6.70 (t, <i>J</i> = 3.6, 1H), 3.28-3.31 (t, <i>J</i> = 13.6, 1H), 2.72-2.81 (m, 4H), 2.51 (s, 3H), 2.14-2.40 (m, 5H), 2.04-2.10 (m, 1H), 1.73-1.85 (m, 4H), 1.64-1.67 (m, 1H).

Ex. No.	Chemical name	Structure	Physical data
62	2-Dimethylaminomethyl-3-(5-hydroxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.06 (s, 1H), 7.00-7.02 (d, <i>J</i> = 8.0, 1H), 6.62-6.64 (m, 1H), 3.04-3.11 (m, 1H), 2.60 (s, 4H), 2.37-2.48 (m, 5H), 2.29 (s, 4H), 2.09 (s, 1H), 1.90-1.93 (m, 1H), 1.75-1.78 (m, 2H), 1.59-1.63 (m, 2H), 1.48-1.54 (m, 2H).
63	2-Dimethylaminomethyl-3-(3-hydroxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.06-7.08 (d, <i>J</i> = 4.0, 1H), 6.86 (s, 1H), 6.85 (s, 1H), 3.03-3.07 (m, 1H), 2.54-2.59 (m, 4H), 2.22-2.29 (m, 4H), 2.12-2.14 (m, 2H), 2.01-2.07 (m, 4H), 1.91-1.94 (m, 1H), 1.76-1.88 (m, 1H), 1.55-1.66 (m, 4H), 1.43-1.49 (m, 1H).

-27-

Ex. No.	Chemical name	Structure	Physical data
64	3-(3,4-Difluoro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.75-6.80 (m, 2H), 3.20-3.21 (m, 1H), 2.36-2.72 (m, 7H), 2.26-2.27 (m, 1H), 2.13-2.20 (m, 2H), 2.02-2.08 (m, 2H), 1.91-1.97 (m, 1H), 1.52-1.69 (m, 4H), 1.50-1.51 (m, 1H).




5

Example 65




Synthesis of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester

-28-





Stir a mixture of 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo
 [3.2. 1]octan-3-ol (587 mg, 2.0 mmol), 2,2-dimethyl-propionyl chloride (361.5 mg, 3.0 mmol) and
 5 triethyl amine (606 mg, 6.0 mmol) in CH₂Cl₂ (60 mL) at ambient temperature for 12 hours.
 Concentrate the mixture under vacuum, and purify the residue by preparative TLC to give
 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-
 hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester (368 mg, Yield: 48.8%). MS (m/z): 378
 (M+1). ¹H NMR (400 MHz, Methanol-d₄) δ 7.29-7.32 (d, *J*₁ = 6.8, *J*₂ = 2.8, 1H), 7.14-7.19 (d, *J*₁ =
 10 11.6, *J*₂ = 8.8, 1H), 7.01-7.05 (m, 1H), 3.19-3.25 (m, 1H), 2.64-2.67 (d, *J* = 13.6, 1H), 2.51 (s, 6H),
 2.46-2.48 (m, 2H), 2.36-2.40 (m, 2H), 2.25-2.30 (m, 1H), 1.99-2.06 (m, 1H), 1.78-1.85 (m, 4H),
 1.61-1.65 (m, 1H), 1.36 (s, 9H).

The following compounds may be prepared essentially by the method of Example 65.

Ex. No.	Chemical name	Structure	Physical data
66	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluorophenyl ester		MS (m/z): 398(M+1), ¹ H NMR (400 MHz, Methanol-d ₄) δ 8.19-8.21 (m, 2H), 7.72-7.76 (t, <i>J</i> = 14.8, 1H), 7.58-7.62 (t, <i>J</i> = 15.6, 2H), 7.47-7.50 (m, 1H), 7.20-7.22 (m, 2H), 3.15-3.18 (m, 1H), 2.62-2.66 (d, <i>J</i> = 13.2, 1H), 2.40-2.55 (m, 10H), 2.28-2.31 (m, 1H), 2.03 (m, 1H), 1.76-1.83 (m, 4H), 1.41-1.50 (m, 1H).
67	2,2-Dimethyl-propionic acid 3-(2-dimethylamino methyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl ester		MS (m/z): 360 (M+1).
68	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl ester		MS (m/z): 380 (M+1).


-30-

Ex. No.	Chemical name	Structure	Physical data
69	2,2-Dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester		MS (m/z): 378 (M+1).
70	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluorophenyl ester		MS (m/z): 398 (M+1), ¹ H NMR (400 MHz, Methanol-d ₄) δ 8.07-8.09 (m, 2H), 7.59-7.61 (t, <i>J</i> = 7.6, 1H), 7.45-7.49 (t, <i>J</i> = 15.6, 2H), 7.10-7.15 (m, 2H), 6.81-6.83 (d, <i>J</i> = 8.8, 1H), 2.71-2.77 (m, 1H), 2.27-2.28 (m, 2H), 2.07-2.13 (m, 7H), 2.04-2.07 (m, 2H), 1.94-2.03 (m, 2H), 1.59-1.74 (m, 4H), 1.45-1.48 (m, 1H).

Example 71



Synthesis of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester hydrochloride

-31-




Add H₂O (9 mg, 0.5mmol) and TMSCl (43 mg, 0.397 mmol) to a solution of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester (150 mg, 0.397 mmol) in 2-butanone (50 mL). Stir the reaction mixture at 0 °C for 2 hours. Evaporate the mixture under vacuum to give 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl- 3-hydroxy-bicyclo [3.2.1]oct-3-yl)-4-fluoro- phenyl ester hydrochloride as white solid (164mg, Yield: 100%). ¹H NMR (400 MHz, Methanol-d₄) δ 7.30-7.32 (d, *J*₁ = 6.8, *J*₂ = 2.8, 1H), 7.15-7.20 (d, *J*₁ = 11.6, *J*₂ = 8.8, 1H), 7.03-7.06 (m, 1H), 3.28-3.31 (m, 1H), 2.77-2.81 (m, 4H), 2.37-2.52 (m, 7H), 2.27 (m, 1H), 2.03 (m, 1H), 1.79-1.86 (m, 4H), 1.63-1.65 (m, 1H), 1.38 (s, 9H).


The following compounds may be prepared essentially by the method of Example 71.

Ex. No.	Chemical name	Structure	Physical data
72	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluorophenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 8.19-8.21 (m, 2H), 7.72-7.76 (t, <i>J</i> = 14.8, 1H), 7.58-7.62 (t, <i>J</i> = 15.6, 2H), 7.49-7.51 (t, <i>J</i> = 8.0, 1H), 7.22-7.24 (m, 2H), 2.42-2.87 (m, 12H), 2.25-2.28 (m, 1H), 2.01-2.03 (m, 1H), 1.79-1.85 (m, 4H), 1.64-1.67 (m, 1H).
73	2,2-Dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl esterhydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.43-7.44 (d, <i>J</i> = 4.4, 2H), 7.24 (s, 1H), 6.94-6.95 (m, 1H), 3.21-3.22 (m, 1H), 2.37-2.81 (m, 9H), 2.21-2.30 (m, 3H), 2.09-2.11 (m, 1H), 1.75-1.88 (m, 4H), 1.63-1.65 (m, 1H), 1.38 (s, 9H).


-33-

Ex. No.	Chemical name	Structure	Physical data
74	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 8.20-8.22 (t, <i>J</i> = 8.4, 2H), 7.72-7.74 (m, 1H), 7.58-7.62 (t, <i>J</i> = 15.2, 2H), 7.50-7.51 (d, <i>J</i> = 5.2, 2H), 7.43 (s, 1H), 7.14 (m, 1H), 3.33 (m, 1H), 2.74-2.86 (m, 4H), 2.43-2.48 (m, 4H), 2.24-2.42 (m, 4H), 2.01-2.10 (m, 1H), 1.76-1.92 (m, 4H), 1.65-1.66 (m, 1H).

-34-


Ex. No.	Chemical name	Structure	Physical data
75	2,2-Dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.22-7.25 (m, 1H), 7.07 (s, 1H), 6.78-6.80 (m, 1H), 3.25-3.28 (m, 1H), 2.72-2.78 (m, 4H), 2.48 (s, 3H), 2.38-2.39 (d, <i>J</i> = 4.0, 2H), 2.17-2.37 (m, 3H), 2.03 (m, 1H), 1.71-1.87 (m, 4H), 1.64-1.67 (m, 1H), 1.38 (s, 9H).

-35-


Ex. No.	Chemical name	Structure	Physical data
76	Benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluorophenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 8.19-8.21 (t, J = 8.4, 2H), 7.72-7.74 (t, J = 8.4, 1H), 7.58-7.62 (t, J = 15.6, 2H), 7.30-7.33 (d, J_1 = 10.4, J_2 = 1.6, 1H), 7.27 (s, 1H), 6.98-7.00 (d, J_1 = 8.8, J_2 = 2.0, 1H), 2.55-2.81 (m, 8H), 2.41 (m, 1H), 2.36 (m, 1H), 2.21-2.35 (m, 3H), 2.08-2.10 (m, 1H), 1.74-1.91 (m, 4H), 1.63-1.67 (m, 1H).

-36-

For compounds of the formula Ib, below, Schemes D and E and Examples 77- 113 illustrate methods of preparing them.




Scheme D



Example 77

Synthesis of 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol













Add TsOH (5.0 g, 29.1 mmol) to a solution of 2-dimethylaminomethyl-3-(3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol (4.8 g, 17.5 mmol) in toluene (150 mL). Heat the reaction mixture to reflux for 2 hours and then quench the reaction by addition of saturated

-37-




aqueous K_2CO_3 (20 mL). Extract the aqueous layer with EtOAc (60 mL x 3). The combined organic layers are washed with brine, dried over Na_2SO_4 and evaporated under vacuum. Purify the residue by preparative HPLC to yield 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol as white solid (2.27 g, 50.2 %). MS (m/z): 258 (M+1).

The following compounds may be prepared essentially by the method of Example 77.

Ex. No.	Chemical name	Structure	Physical data
78	[3-(3-Methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine		MS (m/z): 272 (M+1).
79	[3-(5-Methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine		MS (m/z): 356 (M+1).
80	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-trifluoromethoxy-phenol		MS (m/z): 342 (M+1).
81	[3-(3-Fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine		MS (m/z): 290 (M+1).


Ex. No.	Chemical name	Structure	Physical data
82	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluoro-phenol		MS (m/z): 276 (M+1).
83	[3-(2-Fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-yl methyl]-dimethyl-amine		MS (m/z): 290 (M+1).
84	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluoro-phenol		MS (m/z): 276 (M+1).
85	[3-(3-Methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-yl methyl]-dimethyl-amine		MS (m/z): 286 (M+1).
86	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-methyl-phenol		MS (m/z): 272 (M+1).
87	3-Chloro-5-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol		MS (m/z): 292 (M+1).

-39-

Ex. No.	Chemical name	Structure	Physical data
88	[3-(5-Methoxy-2-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-yl methyl]-dimethyl-amine		MS (m/z): 286 (M+1).
89	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-methyl-phenol		MS (m/z): 272 (M+1).
90	4-Chloro-3-(4-dimethylamino methyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol		MS (m/z): 292 (M+1).

Example 91

Synthesis of 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol hydrochloride



5



Add H₂O (392 mg, 21.8 mmol) and TMSCl (1.4 g, 12.9 mmol) to a solution of 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol (2.8 g, 10.9 mmol) in 2-butanone (200 mL). Stir the reaction mixture at ambient temperature for 12 hours. Collect the precipitate by filter, and wash with EtOAc (30 mL x 2), dry under vacuum to give



10



3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol hydrochloride as white solid (2.41 g, Yield: 75.5%). ^1H NMR(400 MHz, D_2O) δ 7.20-7.24 (t, $J = 15.6$, 1H), 6.76-6.81 (m, 2H), 6.72 (s, 1H), 6.15 (d, $J = 6.4$, 1H), 3.58-3.61 (d, $J = 12.8$, 1H), 3.08-3.14 (t, $J = 26.0$, 1H), 2.89-2.92 (m, 1H), 2.88 (s, 3H), 2.76 (s, 3H), 2.56 (m, 1H), 2.45 (m, 1H), 1.72-1.83 (m, 6H).



5



The following compounds may be prepared essentially by the method of Example 91.



Ex. No.	Chemical name	Structure	Physical data
92	[3-(3-Methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine hydrochloride		^1H NMR (400 MHz, D_2O) δ 7.24-7.28 (t, $J = 16.0$, 1H), 6.82-6.87 (t, $J = 20.4$, 2H), 6.78 (s, 1H), 6.12-6.14 (d, $J = 7.2$, 1H), 3.77 (s, 3H), 3.56-3.59 (m, 1H), 3.04-3.11 (m, 1H), 2.77-2.86 (m, 6H), 2.54 (m, 1H), 2.42 (m, 1H), 2.15 (s, 2H), 1.77-1.80 (m, 1H), 1.65-1.75 (m, 4H).
93	[3-(5-Methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine hydrochloride		^1H NMR (400 MHz, D_2O) δ 7.15-7.18 (d, $J = 9.6$, 1H), 6.81-6.84 (m, 1H), 6.69-6.70 (d, $J = 5.6$, 1H), 6.03-6.01 (d, $J = 4.0$, 1H), 3.68 (s, 3H), 3.46-3.47 (d, $J = 4.0$, 1H), 3.01-3.06 (m, 1H), 2.64-2.70 (d, $J = 9.6$, 6H), 2.32-2.46 (m, 4H), 1.16-1.80 (m, 6H).

Ex. No.	Chemical name	Structure	Physical data
94	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-trifluoromethoxy-phenol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.02-7.04 (d, J = 8.8, 1H), 6.68-6.71 (d, J = 11.6, 1H), 6.56 (d, J = 3.2, 1H), 5.92-5.93 (d, J = 6.8, 1H), 3.32-3.37 (d, J = 18.8, 1H), 2.58-2.59 (m, 2H), 2.37-2.54 (m, 1H), 2.18 (s, 6H), 1.71-1.88 (m, 7H).
95	[3-(3-Fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethylamine hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 6.35-6.38 (m, 3H), 6.01-6.12 (d, J = 6.4, 1H), 3.69-3.70 (d, J = 3.2, 3H), 3.48-3.50 (m, 1H), 3.02-3.09 (m, 1H), 2.68-2.82 (m, 7H), 2.49 (s, 1H), 2.36 (s, 1H), 1.63-1.74 (m, 6H).


Ex. No.	Chemical name	Structure	Physical data
96	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluorophenol hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 6.44-6.49 (m, 2H), 6.41-6.42 (d, <i>J</i> = 2.0, 1H), 6.18-6.19 (d, <i>J</i> = 6.8, 1H), 3.60-3.63 (d, <i>J</i> = 14.0, 1H), 3.16-3.22 (t, <i>J</i> = 25.6, 1H), 2.99-3.00 (d, <i>J</i> = 3.6, 1H), 2.90-2.93 (d, <i>J</i> = 14.8, 6H), 2.64 (s, 1H), 2.57 (s, 1H), 1.92-1.95 (d, <i>J</i> = 10.8, 1H), 1.85 (m, 5H).
97	[3-(2-Fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethylamine hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 6.91-6.96 (m, 1H), 6.77-6.79 (m, 1H), 6.66-6.68 (m, 1H), 6.07-6.08 (d, <i>J</i> = 4.0, 1H), 3.66-3.67 (d, <i>J</i> = 4.0, 3H), 3.47 (s, 1H), 3.03 (m, 1H), 2.60-2.69 (m, 6H), 2.48 (s, 1H), 2.33 (s, 1H), 1.62-1.78 (m, 7H).

Ex. No.	Chemical name	Structure	Physical data
98	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluorophenol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 6.83-6.88 (m, 1H), 6.63-6.67 (m, 1H), 6.54-6.57 (m, 1H), 6.03-6.05 (d, <i>J</i> = 8.0, 1H), 3.44-3.48 (d, <i>J</i> = 16.0, 1H), 3.01-3.09 (m, 1H), 2.61-2.69 (m, 7H), 2.48 (s, 1H), 2.32 (s, 1H), 1.76-1.79 (d, <i>J</i> = 12.0, 1H), 1.70-1.71 (d, <i>J</i> = 4.0, 1H), 1.61-1.65 (m, 4H).
99	[3-(3-Methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethylamine hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 6.67-6.69 (m, 2H), 6.57 (s, 1H), 6.13-6.14 (d, <i>J</i> = 6.8, 1H), 3.80 (s, 3H), 3.64-3.67 (d, <i>J</i> = 12.4, 1H), 3.14-3.26 (m, 1H), 2.86-2.95 (m, 7H), 2.57-2.63 (m, 2H), 2.33 (s, 3H), 1.94-1.97 (m, 1H), 1.79-1.92 (m, 5H).


Ex. No.	Chemical name	Structure	Physical data
100	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-methyl-phenol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 6.63-6.53 (d, <i>J</i> = 1.2, 2H), 6.44 (s, 1H), 6.11-6.13 (d, <i>J</i> = 6.4, 1H), 3.59-3.63 (m, 1H), 3.13-3.19 (m, 1H), 2.99-3.00 (m, 1H), 2.89-2.97 (m, 6H), 2.55-2.62 (m, 2H), 2.18 (s, 3H), 1.80-1.95 (m, 6H).
101	3-Chloro-5-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol hydrochloride		¹ H NMR (400MHz, Methanol-d ₄) δ 6.71-6.73 (m, 2H), 6.58-6.59 (t, <i>J</i> = 3.6, 1H), 6.16-6.18 (d, <i>J</i> = 6.4, 1H), 3.62-3.68 (m, 1H), 3.16-3.23 (t, <i>J</i> = 26.0, 1H), 2.90-2.96 (m, 7H), 2.56-2.64 (m, 2H), 1.83-1.95 (m, 6H).

Ex. No.	Chemical name	Structure	Physical data
102	[3-(5-Methoxy-2-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethylamine hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.10 (d, J = 6.4, 1H), 6.75 (d, J_1 = 2.4, J_2 = 6.4, 1H), 6.58 (m, 1H), 5.78 (s, 1H), 3.71 (s, 3H), 3.35 (d, J = 12.0, 1H), 3.12 (t, J = 12.4, 1H), 2.62 (s, 6H), 2.48 (s, 2H), 2.35 (m, 1H), 2.06 (s, 3H), 1.84 (m, 1H), 1.73 (m, 1H), 1.65 (m, 4H).
103	3-(4-Dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-methyl-phenol hydrochloride		¹ H NMR (400 MHz, D ₂ O) δ 7.21-7.23 (d, J = 4.0, 1H), 6.82-6.84 (d, J = 4.0, 1H), 6.69 (s, 1H), 6.96-6.97 (d, J = 4.0, 1H), 3.50-3.52 (m, 1H), 3.25-3.27 (m, 1H), 2.85 (s, 6H), 2.66-2.71 (m, 2H), 2.54 (s, 1H), 2.23 (s, 3H), 1.87-2.05 (m, 6H).

-46-

Ex. No.	Chemical name	Structure	Physical data
104	4-Chloro-3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.18-7.20 (d, J = 8.8, 1H), 6.70-6.73 (d, J_1 = 8.8, J_2 = 3.2, 1H), 6.60 (s, 1H), 5.97-5.98 (d, J = 6.0, 1H), 3.71-3.76 (m, 1H), 3.32-3.34 (t, J = 6.4, 1H), 2.87 (s, 6H), 2.63-2.66 (s, 3H), 2.03-2.05 (d, J = 10.8, 1H), 1.83-1.94 (m, 5H).

Scheme E




5

Example 105

Synthesis of 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester



-47-





Stir a mixture of 3-(4-(dimethylaminomethyl)-bicyclo[3.2.1]oct-2-en-3-yl)-phenol (200 mg, 0.778 mmol), 2,2-dimethyl-propionyl chloride (111 mg, 0.927 mmol) and triethyl amine (234 mg, 2.316 mmol) in CH₂Cl₂ (40mL) at ambient temperature for 12 hours. Concentrate the mixture under vacuum, and purify the residue by Preparative TLC to give 2,2-dimethyl-propionic acid 3-(4-(dimethylaminomethyl)-bicyclo [3.2.1]oct-2-en-3-yl)-phenyl ester as white solid (232 mg, Yield: 87.5%). MS (m/z): 342 (M+1). ¹H NMR (400 MHz, Methanol-d₄) δ 7.29-7.32 (t, *J* = 15.6Hz, 1H), 7.06-7.08 (d, *J* = 7.6, 1H), 6.88-6.90 (d, *J*₁ = 8.0, *J*₂ = 1.6, 1H), 6.84-6.85 (t, *J* = 3.6, 1H), 6.05-6.07 (d, *J* = 6.8, 1H), 3.22-3.32 (m, 1H), 2.52-2.60 (m, 1H), 2.62-2.63 (m, 1H), 2.37-2.41 (m, 1H), 2.17 (s, 6H), 2.07-2.14 (m, 1H), 1.71-1.85 (m, 6H), 1.37 (s, 9H).

The following compounds may be prepared essentially by the method of Example 105.

Ex. No.	Chemical name	Structure	Physical data
106	Benzoic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester		MS (m/z): 362 (M+1), ¹ H NMR (400 MHz, Methanol-d ₄) δ 8.19-8.21 (d, <i>J</i> = 8.0, 2H), 7.70-7.73 (t, <i>J</i> = 15.2 Hz, 1H), 7.57-7.61 (t, <i>J</i> = 19.6, 2H), 7.36-7.40 (t, <i>J</i> = 15.6, 1H), 7.14-7.16 (d, <i>J</i> = 8.0, 1H), 7.07-7.09 (m, 2H), 6.11-6.13 (d, <i>J</i> = 6.8, 1H), 3.31-3.33 (m, 2H), 2.59-2.64 (d, <i>J</i> = 20.4, 2H), 2.42-2.45 (t, <i>J</i> = 13.2, 1H), 2.18-2.20 (m, 6H), 1.75-1.88 (m, 6H).

Ex. No.	Chemical name	Structure	Physical data
107	2,2-Dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluorophenyl ester		MS (m/z): 360 (M+1), ¹H NMR (400 MHz, Methanol-d₄) δ 6.94-6.97 (m, 1H), 6.80-6.81 (m, 2H), 6.25-6.28 (d, <i>J</i> = 9.2, 1H), 3.68-3.71 (d, <i>J</i> = 12.4, 1H), 3.18-3.25 (t, <i>J</i> = 26.0, 1H), 2.90-2.93 (m, 7H), 2.62-2.66 (m, 2H), 1.85-1.96 (m, 6H), 1.37 (s, 9H).
108	Isobutyric acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluorophenyl ester		MS (m/z): 346 (M+1). ¹H NMR (400 MHz, Methanol-d₄) δ 6.86-6.89 (m, 1H), 6.76-6.79 (m, 2H), 6.14-6.15 (d, <i>J</i> = 6.8, 1H), 3.36-3.39 (m, 1H), 2.82-2.85 (m, 1H), 2.54-2.63 (m, 3H), 2.37 (s, 6H), 2.25-2.29 (m, 1H), 1.74-1.87 (m, 6H), 1.31-1.32 (d, <i>J</i> = 6.8, 6H).


-49-

Ex. No.	Chemical name	Structure	Physical data
109	2,2-Dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluorophenyl ester		MS (m/z): 360 (M+1), ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.10 (t, <i>J</i> = 9.6, 1H), 6.97 (m, 1H), 6.88 (m, 1H), 6.12 (d, <i>J</i> = 9.6, 1H), 3.48 (m, 1H), 2.85 (t, <i>J</i> = 12.4, 1H), 2.61 (s, 2H), 2.52 (s, 6H), 2.33 (m, 1H), 1.94 (m, 1H), 1.83(m, 5H), 1.38 (s, 9H).
110	Isobutyric acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluorophenyl ester		MS (m/z): 346 (M+1). ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.10 (t, <i>J</i> = 9.6, 1H), 7.01 (m, 1H), 6.90 (m, 1H), 6.11 (d, <i>J</i> = 6.4, 1H), 3.50 (m, 1H), 2.85 (m, 2H), 2.62 (s, 2H), 2.55 (s, 6H), 2.48 (m, 1H), 1.95 (m, 1H), 1.76 (m, 5H), 1.31 (d, <i>J</i> = 6.8, 6H).

Example 111

Synthesis of 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester hydrochloride



-50-



Add H₂O (18 mg, 1mmol) and TMSCl (75 mg, 0.69 mmol) to a solution of 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)- phenyl ester (197 mg, 0.58 mmol) in 2-butanone (70mL). Stir the reaction mixture at ambient temperature for 2 hours. Evaporate the mixture under vacuum to afford 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl -bicyclo[3.2.1]oct-2-en-3-yl)- phenyl ester hydrochloride as white solid (219 mg, Yield: 100%). ¹H NMR (400 MHz, Methanol-d₄) δ 7.38-7.42 (t, *J* = 16.4, 1H), 7.13-7.15 (d, *J*₁ = 7.2, *J*₂ = 0.8, 1H), 6.96-6.98 (m, 2H), 6.21-6.23 (d, *J* = 6.8, 1H), 3.68-3.71 (d, *J* = 12.0, 1H), 3.16-3.23 (t, *J* = 25.6, 1H), 2.99-3.00 (m, 1H), 2.88-2.92 (d, *J* = 14.8, 6H), 2.65 (m, 1H), 2.60 (m, 1H), 1.95-1.98 (m, 1H), 1.82-1.86 (m, 5H), 1.37 (s, 9H).

The following compounds may be prepared essentially by the method of Example 111.


-51-

Ex. No.	Chemical name	Structure	Physical data
112	Benzoic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 8.19-8.21 (t, J = 8.0, 2H), 7.71-7.74 (t, J = 14.8, 1H), 7.57-7.61 (t, J = 15.6, 2H), 7.43-7.72 (t, J = 15.6, 1H), 7.13-7.22 (m, 3H), 6.25-6.27 (d, J = 6.8, 1H), 3.71-3.74 (d, J = 12.0, 1H), 3.19-3.26 (t, J = 25.6, 1H), 3.02-3.06 (m, 1H), 2.90-2.94 (d, J = 14.8, 6H), 2.65 (s, 2H), 1.95-1.98 (m, 1H), 1.83-1.87 (m, 5H).
113	2,2-Dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluoro-phenyl ester hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.94-6.96 (d, J_1 = 3.6, J_2 = 2.0, 1H), 6.81-6.83 (m, 2H), 6.25-6.27 (d, J = 6.8, 1H), 3.65-3.68 (d, J = 10.8, 1H), 3.15-3.21 (t, J = 26.0, 1H), 2.88-2.93 (m, 7H), 2.66 (m, 1H), 2.58 (m, 1H), 1.82-1.96 (m, 6H), 1.37 (s, 9H).

For compounds of the formula Ic, below, Schemes F and G and Preparations and/or


5 Examples 114-140 illustrate methods of preparing them.

-52-




Ic

5

Scheme FExample 114

10

Synthesis of 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol








Add Pd/C (80 mg) to a solution of 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol (250 mg, 0.97 mmol) in MeOH (40 mL). Then stir the mixture at 25 °C for 12 hours under 45 PSI of H₂. After filtration, Concentrate the solution under vacuum to afford

15



-53-

3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol as white solid (252 mg, Yield: 100 %).
MS (m/z): 260 (M+1).

The following compounds may be prepared essentially by the method of Example 114.


Ex. No.	Chemical name	Structure	Physical data
115	[3-(3-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine		MS (m/z): 274 (M+1).
116	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenol		MS (m/z): 278 (M+1).
117	[3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine		MS (m/z): 292 (M+1).
118	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenol		MS (m/z): 278 (M+1).
119	[3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine		MS (m/z): 292 (M+1).

-54-

Ex. No.	Chemical name	Structure	Physical data
120	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-methyl-phenol		MS (m/z): 274 (M+1).
121	[3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine		MS (m/z): 288 (M+1).

Example 122




Synthesis of 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol hydrochloride






5


Add H₂O (70 mg, 3.89 mmol) and TMSCl (126 mg, 1.17 mmol) to a solution of 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol (252 mg, 0.97 mmol) in 2-butanone (30ml). Then stir the reaction mixture at ambient temperature for 12 hours. Evaporate the mixture under vacuum to give 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol hydrochloride as white solid (286 mg, Yield: 99.8%). ¹H NMR (400 MHz, D₂O) δ 7.04-7.07 (t, J = 10.4, 1H), 6.58-6.72 (m, 3H), 2.80-2.91 (m, 1H), 2.52-2.54 (m, 3H), 2.44-2.48 (m, 3H), 2.33-2.41 (m, 1H), 2.20-2.29 (m, 1H), 2.11 (s, 1H), 1.99 (s, 2H), 1.36-1.47 (m, 8H).

15 The following compounds may be prepared essentially by the method of Example 122.


Ex. No.	Chemical name	Structure	Physical data
123	[3-(3-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine hydrochloride		^1H NMR (400 MHz, D_2O) δ 7.23-7.27 (t, $J = 15.6$, 1H), 6.80-6.90 (m, 3H), 3.73 (s, 3H), 2.95-3.01 (m, 1H), 2.54-2.62 (m, 6H), 2.38-2.42 (m, 2H), 2.22 (s, 1H), 2.12-2.14 (m, 2H), 1.48-1.67 (m, 8H).
124	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenol hydrochloride		^1H NMR (400 MHz, D_2O) δ 6.55-6.49 (m, 2H), 6.41-6.38 (d, $J = 12.0$, 1H), 2.97-2.91 (m, 1H), 2.60-2.56 (m, 3H), 2.52 (s, 3H), 2.46-2.33 (m, 3H), 2.16 (s, 1H), 2.08-1.99 (m, 3H), 1.62-1.59 (m, 1H), 1.45-1.37 (m, 4H), 1.37-1.35 (m, 1H).
125	[3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine hydrochloride		^1H NMR (400 MHz, D_2O) δ 6.61-6.59 (d, $J = 8.0$, 2H), 6.52-6.50 (m, 1H), 3.66 (s, 3H), 2.94-2.90 (m, 1H), 2.58-2.52 (m, 6H), 2.37-2.33 (m, 2H), 2.16 (s, 1H), 2.06-2.03 (m, 2H), 1.61-1.36 (m, 8H).

Ex. No.	Chemical name	Structure	Physical data
126	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluorophenol hydrochloride		^1H NMR (400 MHz, D_2O) δ 6.91-6.83 (m, 1H), 6.74 (s, 1H), 6.62-6.49 (m, 1H), 3.03-2.94 (m, 1H), 2.49 (s, 4H), 2.42-2.35 (m, 2H), 2.19-2.08 (m, 6H), 1.63-1.40 (m, 7H).
127	[3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethylamine hydrochloride		^1H NMR (400 MHz, D_2O) δ 6.98-6.86 (m, 1H), 6.76 (s, 1H), 6.75-6.74 (m, 1H), 3.68 (s, 3H), 3.04-2.98 (m, 1H), 2.81 (s, 1H), 2.64-2.55 (m, 6H), 2.36-2.39 (d, $J = 12.0$, 1H), 2.21 (s, 1H), 2.06-1.98 (m, 2H), 1.69-1.42 (m, 8H).
128	3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-methylphenol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.59 (s, 1H), 6.51 (s, 2H), 3.06-3.12 (m, 1H), 2.67-2.80 (m, 6H), 2.53-2.54 (m, 1H), 2.33-2.38 (m, 3H), 2.28 (s, 3H), 2.17-2.21 (t, $J = 12.0$, 1H), 1.52-1.87 (m, 8H).

-57-

Ex. No.	Chemical name	Structure	Physical data
129	[3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethylamine hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.71 (s, 1H), 6.64-6.65 (d, J = 5.6, 2H), 3.80 (s, 3H), 3.13-3.34 (m, 1H), 2.82-2.83 (d, J = 2.4, 3H), 2.72 (d, J = 2.4, 3H), 2.51-2.55 (m, 1H), 2.35-2.44 (m, 6H), 2.23-2.26 (t, J = 11.2, 1H), 1.60-1.85 (m, 8H).

Scheme G




5

Example 130

Synthesis of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenyl ester



-58-





Stir a mixture of 3-(2-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol (200 mg, 0.772 mmol), 2,2-dimethyl-propionyl chloride (111.6 mg, 0.927 mmol) and triethyl amine (234 mg, 2.316 mmol) in CH₂Cl₂ (40 mL) at ambient temperature for 12 hours. Concentrate the mixture under reduced pressure, and purify the residue by preparative TLC to afford 2,2-dimethyl-propionic acid 3-(2-(2-dimethylaminomethyl-bicyclo[3.2.1] oct-3-yl)-phenyl ester as white solid (217 mg, Yield: 81.9%). MS (m/z): 344 (M+1). ¹H NMR (400 MHz, Methanol-d₄) δ 7.36-7.40 (t, *J* = 13.6, 1H), 7.19-7.21 (d, *J* = 7.2, 1H), 7.03 (s, 1H), 6.92-6.94 (d, *J* = 8.0, 1H), 2.97-3.03 (t, *J* = 23.6, 1H), 2.49-2.68 (m, 6H), 2.40-2.44 (d, *J* = 13.2, 1H), 2.35 (s, 2H), 2.17-2.21 (m, 2H), 1.60-1.84 (m, 8H), 1.37 (s, 9H).

The following compounds are prepared essentially by the method of Example 130.

Ex.	Chemical name	Structure	Physical data
131	benzoic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenyl ester		MS (m/z): 364 (M+1), ¹ H NMR (400 MHz, CDCl ₃) δ 8.19-8.22 (m, 2H), 7.63-7.64 (t, <i>J</i> = 7.6, 1H), 7.50-7.52 (t, <i>J</i> = 7.6, 2H), 7.32-7.36 (t, <i>J</i> = 16.4, 1H), 7.04-7.10 (m, 3H), 2.46 (s, 1H), 2.19-2.39 (m, 3H), 2.10 (s, 6H), 1.80-1.90 (t, <i>J</i> = 37.2, 1H), 1.52-1.71 (m, 9H).

Ex.	Chemical name	Structure	Physical data
132	2,2-dimethyl-propionic acid 3-(2-dimethylaminoethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester		MS (m/z): 362 (M+1), 1H NMR (400 MHz, Methanol-d4) δ 6.96-6.98 (d, $J=10.0$, 1H), 6.84 (s, 1H), 6.73-6.76 (m, 1H), 2.66-2.72 (m, 1H), 2.49-2.54 (m, 1H), 2.30-2.37 (m, 8H), 2.04-2.07 (m, 2H), 1.56-1.81 (m, 8H), 1.36 (s, 9H).
133	isobutyric acid 3-(2-dimethylaminoethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester		MS (m/z): 348 (M+1). 1H NMR (400 MHz, Methanol-d4) δ 6.82-6.85 (d, $J=9.6$, 1H), 6.73 (s, 1H), 6.62-6.65 (m, 1H), 2.68-2.75 (m, 1H), 2.34-2.46 (m, 3H), 2.15-2.28 (m, 1H), 2.15 (s, 6H), 1.76-1.81 (m, 2H), 1.44-1.69 (m, 8H), 1.19-1.20 (d, $J=7.2$, 6H).


-60-

Ex.	Chemical name	Structure	Physical data
134	2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluorophenyl ester		MS (m/z): 362 (M+1). ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.10 (t, <i>J</i> = 8.8, 2H), 6.90 (m, 1H), 2.92 (s, 1H), 2.43 (s, 1H), 2.33 (m, 2H), 2.03 (s, 6H), 1.90 (s, 1H), 1.50-1.82 (m, 9H), 1.45 (s, 9H).
135	isobutyric acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluorophenyl ester		MS (m/z): 348 (M+1). ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.13 (t, <i>J</i> = 8.4, 2H), 6.90 (m, 1H), 2.85 (m, 2H), 2.40 (s, 1H), 2.33 (m, 2H), 2.06 (s, 6H), 1.50-1.88 (m, 10H), 1.31 (d, <i>J</i> = 6.4, 6H).

Example 136

Synthesis of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenyl ester hydrochloride

5







-61-

Add H₂O (9 mg, 0.5mmol) and TMSCl (39mg, 0.357mmol) to a solution of 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenyl ester (102mg, 0.297mmol) in 2-butanone (50mL). Then stir the reaction mixture at ambient temperature for 4 hours. Evaporate the mixture under vacuum to give 2,2-dimethyl-propionic acid

- 5 3-(2-dimethylaminomethyl -bicyclo[3.2.1]oct-3-yl)-phenyl ester hydrochloride as white solid (112mg, Yield: 100%). ¹H NMR (400 MHz, Methanol-d₄) δ 7.38-7.42 (t, *J* = 15.6, 1H), 7.20-7.22 (d, *J* = 7.6, 1H), 7.05 (s, 1H), 6.94-6.96 (m, 1H), 3.08-3.14 (m, 1H), 2.79 (s, 3H), 2.66 (s, 3H), 2.53-2.57 (m, 2H), 2.36 (s, 2H), 2.23 (m, 1H), 1.61-1.85 (m, 8H), 1.39 (s, 9H).


- 10 The following compounds may be prepared essentially by the method of Example 135.

Ex. No.	Chemical name	Structure	Physical data
137	2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.04-7.06 (d, <i>J</i> = 9.6, 1H), 6.92 (s, 1H), 6.78-6.81 (m, 1H), 3.10-3.16 (m, 1H), 2.70-2.81 (m, 6H), 2.52-2.58 (m, 2H), 2.36 (s, 2H), 2.24 (m, 1H), 1.58-1.86 (m, 8H), 1.37 (s, 9H).
138	isobutyric acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester hydrochloride		¹ H NMR (400 MHz, Methanol-d ₄) δ 7.03-7.06 (d, <i>J</i> = 9.6, 1H), 6.95 (s, 1H), 6.79-6.83 (m, 1H), 3.10-3.16 (m, 1H), 2.82-2.90 (m, 1H), 2.81 (s, 3H), 2.69 (s, 3H), 2.52-2.58 (m, 2H), 2.37-2.38 (m, 2H), 2.22-2.27 (m, 1H), 1.56-1.85 (m, 8H), 1.31 (s, 6H).


Ex. No.	Chemical name	Structure	Physical data
139	2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.22 (m, 2H), 6.95 (s, 1H), 3.12 (t, $J = 11.2$, 1H), 2.98 (s, 1H), 2.72 (s, 6H), 2.45 (d, $J = 12.8$, 1H), 2.39 (s, 2H), 2.25 (s, 1H), 1.84 (m, 1H), 1.52-1.81 (m, 7H), 1.47 (s, 9H).
140	isobutyric acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.26 (s, 1H), 7.15 (m, 1H), 7.01 (s, 1H), 3.16 (t, $J = 10.8$, 1H), 2.98 (s, 1H), 2.84 (m, 3H), 2.70 (s, 3H), 2.53 (m, 1H), 2.41 (s, 1H), 2.38 (s, 1H), 2.27 (s, 1H), 1.53-1.92 (m, 9H), 1.29 (d, $J = 6.8$, 6H).

-63-


For compounds of the formula Id, below, Schemes H-J and Preparations and/or Examples 147-164 illustrate methods of preparing them.



Id

Scheme H**Preparation 141**


Synthesis of bicyclo[4.1.0]heptan-2-one



-64-


Add trimethylsulfoxonium iodide (126.3 g, 0.572 mol) to a suspension of sodium hydride (22.9 g, 60% in oil dispersion, 0.572 mol) in dry DMSO (550 mL) at ambient temperature within 10 minutes. Stir the mixture for 1 hour, and then add dropwise a solution of cyclohex-2-enone (51.2 g, 0.52 mol) in dry DMSO (100 mL) to the reaction mixture. Stir the reaction mixture at 50 °C for additional 2 hours. Pour the reaction mixture into ice-water, and extract the aqueous mixture with CH₂Cl₂ (300 mL x 3). The combined organic layers were washed with brine, dried over MgSO₄, concentrated in vacuo. Purify the residue by distillation (55~60 °C, 5 mmHg) to afford bicyclo[4.1.0] heptan-2-one as colorless oil (38.45 g, 66.6 %), MS (m/z): 109 (M-1).

The following compound may be prepared essentially by the method of Preparation 141.

Prep No.	Chemical name	Structure	Physical data
142	5,5-Dimethyl-bicyclo[4.1.0]heptan-2-one		MS (m/z): 137 (M-1).

Preparation 143

Synthesis of (bicyclo[4.1.0]hept-2-en-2-yloxy)-trimethyl-silane




Add dropwise bicyclo[4.1.0]heptan-2-one (43.3 g, 0.393 mol) in dry THF (50 mL) to a solution of LDA (237 mL, 0.472 mol) in THF (500 mL). Stir the reaction mixture for 30 min and then add dropwise TMSCl (64.2 g, 0.59 mol) to the reaction mixture. Stir the resultant solution for additional 1 hour. Pour the mixture into ice water (92 mL) and extract the aqueous mixture with EtOAc (100 mL x 3). The combined organic layers are washed with brine, dried over MgSO₄,

-65-

filtered and concentrated in vacuo to afford crude (bicyclo[4.1.0]hept-2-en-2-yloxy)-trimethyl-silane (68.0 g, 95.0 %) as yellowish oil, MS (m/z): 181 (M-1).


The following compound may be prepared essentially by the method of Preparation 143.

Prep No.	Chemical name	Structure	Physical data
144	(5,5-Dimethyl-bicyclo[4.1.0]hept-2-en-2-yloxy)-trimethyl-silane		MS (m/z): 209 (M-1).

5

Preparation 145

Synthesis of 3-dimethylaminomethyl-bicyclo[4.1.0]heptan-2-one




10

Add (bicyclo[4.1.0]hept-2-en-2-yloxy)-trimethyl-silane (72.9 g, 0.40 mol) to a cooled suspension of N,N-dimethylmethyleiminium chloride (48.6 g, 0.52 mol) in CH₂Cl₂ (400 mL) at 0 °C. After being stirred for overnight at ambient temperature, dilute the reaction mixture with 2 N HCl (200 mL). After removal of organic layers, wash the aqueous layer with EtOAc (50 mL x 3) and then basify with NaOH to PH = 10. Extract the resultant mixture with EtOAc (100 mL x 4). The combined organic layers are washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo. Purify the residue by distillation (80 ~ 85 °C, 5 mmHg) to afford 3-dimethylaminomethyl-bicyclo [4.1.0]heptan-2-one (16.0 g, 24.0 %) as colorless oil, MS (m/z): 168 (M+1).

20


The following compound may be prepared essentially by the method of Preparation 145.

-66-

Prep No.	Chemical name	Structure	Physical data
146	3-Dimethylaminomethyl-5,5-dimethyl-bicyclo[4.1.0]heptan-2-one		MS (m/z): 196 (M+1).

Example 147





Synthesis of 3-Dimethylaminomethyl-2-(3-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol




5

- Add a solution of *t*-BuLi (5.8 mL, 8.68 mmol) *via* syringe to a solution of 3-bromo-phenol (752 mg, 4.34 mmol) in THF (60 mL) at -78°C under N₂. After being stirred at -78 °C for 1 hour, add a solution of 3-dimethylaminomethyl-bicyclo[4.1.0] heptan-2-one (300 mg, 2.17mmol) in THF (2 mL) to the reaction mixture and stir the reaction mixture at -78 °C for additional 2 hours. Quench the reaction mixture with saturated NH₄Cl solution (30 mL). Extract the aqueous resultant mixture with EtOAc (60 mL x 3). The combined organic layers are washed with brine (30 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by Preparative HPLC to give
- 3-dimethylaminomethyl-2-(3-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol as white solid (73 mg, yield: 12.9%). MS (m/z): 262 (M+1).

The following compounds may be prepared essentially by the method of Example 147.

Ex. No.	Chemical name	Structure	Physical data
148	3-Dimethylaminomethyl-2-(3-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 276 (M+1).
149	3-Dimethylaminomethyl-2-(3-fluoro-5-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 294 (M+1).
150	3-Dimethylaminomethyl-2-(5-methoxy-2-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 290 (M+1).
151	3-Dimethylaminomethyl-2-(3-methoxy-phenyl)-5,5-dimethyl-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 304 (M+1) ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.31 (t, J = 8.0, 1H), 7.13 (m, 2H), 6.82 (d, J ₁ = 6.4, J ₂ = 2.4, 1H), 3.82 (s, 3H), 2.32 (m, 1H), 2.18 (s, 1H), 1.96 (s, 7H), 1.83 (m, 1H), 1.38 (m, 1H), 1.28 (m, 3H), 1.22 (m, 2H), 1.10 (s, 3H), 1.02 (m, 1H), 0.61 (m, 1H), 0.50 (m, 1H).


-68-

Ex. No.	Chemical name	Structure	Physical data
152	3-Dimethylaminomethyl-2-(3-hydroxy-phenyl)-5,5-dimethyl-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 290 (M+1), ¹ H NMR (Methanol-d ₄ , 400 MHz) δ 7.29 (t, J = 8.0, 1H), 7.03 (m, 2H), 6.64 (m, 1H), 2.35 (m, 1H), 1.96 (s, 7H), 1.83 (m, 1H), 1.38 (m, 1H), 1.28 (s, 3H), 1.22 (m, 2H), 1.10 (s, 3H), 1.02 (m, 1H), 0.61 (m, 1H), 0.50 (m, 1H).

Example 153

Synthesis of 3-dimethylaminomethyl-2-(3-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride

5





Add H₂O (10 mg, 0.560 mmol) and TMSCl (33 mg, 0.308 mmol) to a solution of 3-dimethylaminomethyl-2-(3-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol (73 mg, 0.280 mmol) in 2-butanone (10 mL). Then stir the mixture at room temperature for 2 hours. Evaporate the mixture under vacuum to give 3-dimethylaminomethyl-2-(3-hydroxyl -phenyl)-bicyclo[4.1.0]heptan-2-ol


10

hydrochloride as white solid (83 mg, Yield: 100%). ^1H NMR (400 MHz, Methanol- d_4) δ 7.21-7.25 (t, $J=16.0$, 1H), 7.17-7.18 (t, $J=4.4$, 1H), 7.11-7.13 (m, 1H), 6.76-6.79 (d, $J=10.0$, 1H), 3.23-3.25 (m, 1H), 2.68-2.69 (m, 7H), 2.14-2.17 (m, 1H), 2.04-2.09 (m, 1H), 1.76-1.81 (m, 1H), 1.62-1.64 (m, 1H), 1.47-1.49 (m, 1H), 1.11-1.16 (m, 1H), 0.93-1.05 (m, 1H), 0.73-0.84 (m, 1H), 0.50-0.53 (m, 1H).


The following compounds may prepared essentially by the method of Example 153.

Ex. No.	Chemical name	Structure	Physical data
154	3-Dimethylaminomethyl-2-(3-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.23 (m, 1H), 7.17 (m, 1H), 7.11 (m, 1H), 6.78 (m, 1H), 3.72 (s, 3H), 2.95 (m, 1H), 2.60-2.71 (m, 7H), 2.16 (m, 1H), 2.05 (m, 1H), 1.80 (m, 1H), 1.65 (m, 1H), 1.48 (m, 1H), 1.15 (m, 1H), 1.01 (m, 1H), 0.67 (m, 1H), 0.50 (m, 1H).
155	3-Dimethylaminomethyl-2-(3-fluoro-5-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.08 (d, $J=1.6$, 1H), 7.00 (m, 1H), 6.62 (m, 1H), 3.82 (s, 3H), 3.02 (m, 1H), 2.75 (m, 7H), 2.23 (m, 1H), 2.10 (m, 1H), 1.81 (m, 1H), 1.73 (m, 1H), 1.62 (m, 1H), 1.26 (m, 1H), 1.12 (m, 1H), 0.85 (m, 1H), 0.64 (m, 1H).

-70-

Ex. No.	Chemical name	Structure	Physical data
156	3-Dimethylaminomethyl-2-(5-methoxy-2-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.10-7.12 (d, J = 8.0, 1H), 6.92 (s, 1H), 6.75-6.77 (d, J = 8.0, 1H), 3.79 (s, 3H), 3.12 (s, 2H), 2.85 (s, 6H), 2.59 (s, 3H), 2.47 (s, 1H), 2.15-2.16 (m, 1H), 1.63-1.80 (m, 3H), 1.41-1.44 (m, 1H), 1.25-1.29 (m, 1H), 0.70-0.71 (m, 1H), 0.25-0.26 (m, 1H).

Scheme I




5

Example 157

Synthesis of 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0] heptan-2-ol

-71-



Add a solution of *t*-BuLi (19.4 mL, 25.2 mmol) *via* syringe to a solution of (3-bromo-5-fluoro-phenoxy)-tert-butyl-dimethyl-silane (6.99 g, 22.9 mmol) in THF (100 mL) at -78 °C under N₂. After being stirred at -78 °C for 1 hour, add dropwise a solution of 3-dimethylaminomethyl-bicyclo[4.1.0]heptan-2-one (2.55 g, 15.3 mmol) to the reaction mixture and stir the reaction mixture at -78 °C for additional 2 hours. Quench the reaction with saturated NH₄Cl solution (30 mL). Extract the resultant aqueous mixture with EtOAc (100 mL x 3). The combined organic layers are washed with brine (30 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Treat the residue with TBAF (6.00 g, 22.9 mmol) in CH₂Cl₂ (80mL) at ambient temperature for 4 hours. Concentrate the mixture, and purify the residue by preparative HPLC to give 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0] heptan-2-ol as white solid (592 mg, yield: 13.9%). MS (m/z): 280 (M+1).


The following compounds may be prepared essentially by the method of Example 157.

Ex. No.	Chemical name	Structure	Physical data
158	3-Dimethylaminomethyl-2-(2-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 280 (M+1).
159	3-Dimethylaminomethyl-2-(3-hydroxy-5-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol		MS (m/z): 276 (M+1).

-72-

Example 160

Synthesis of 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0] heptan-2-ol hydrochloride





5

Add H₂O (18 mg, 1.0 mmol) and TMSCl (92 mg, 0.857 mmol) to a solution of 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0] heptan-2-ol (218 mg, 0.781 mmol) in 2-butanone (70 mL). Then stir the reaction mixture at room temperature for 2
10 hours. After removal the solvent by evaporation, wash the residue with EtOAc (3 mL x 2) to give 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)- bicyclo[4.1.0]heptan-2-ol hydrochloride as white solid (72 mg, Yield: 67.8%). ¹H NMR (400 MHz, D₂O) δ 6.93 (s, 1H), 6.87-6.89 (d, *J* = 9.6, 1H), 6.42-6.58 (m, 1H), 3.32-3.33 (t, *J* = 3.2, 1H), 2.98-3.04 (m, 1H), 2.77 (m, 6H), 2.22-2.23 (m, 1H), 2.06 (m, 1H), 1.81-1.85 (m, 2H), 1.61-1.71 (m, 1H), 1.17-1.21 (m, 1H), 1.08-1.10 (m,
15 1H), 0.83-0.86 (m, 1H), 0.56-0.60 (m, 1H).


The following compounds may be prepared essentially by the method of Example 160.

-73-


Ex. No.	Chemical name	Structure	Physical data
161	3-Dimethylamino methyl-2-(2-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 7.27-7.30 (m, 1H), 6.91 (m, 1H), 6.69-6.73 (m, 1H), 3.06-3.11 (m, 1H), 2.76-2.86 (m, 7H), 2.46-2.54 (m, 1H), 2.20-2.26 (m, 1H), 1.83-1.87 (m, 1H), 1.64-1.71 (m, 1H), 1.52-1.57 (m, 1H), 1.19-1.21 (m, 1H), 1.02-1.14 (m, 1H), 0.73-0.79 (m, 1H), 0.59-0.63 (m, 1H).
162	3-Dimethylamino methyl-2-(3-hydroxy-5-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol hydrochloride		^1H NMR (400 MHz, Methanol- d_4) δ 6.95 (s, 1H), 6.91 (s, 1H), 6.56 (s, 1H), 2.97-3.04 (m, 1H), 2.78-2.82 (d, $J_1=13.2$, $J_2=2.0$, 1H), 2.74 (s, 6H), 2.31 (s, 3H), 2.20-2.27 (m, 1H), 2.05-2.10 (m, 1H), 1.80-1.87 (m, 1H), 1.68-1.76 (m, 1H), 1.55-1.60 (m, 1H), 1.17-1.21 (m, 1H), 1.06-1.12 (m, 1H), 0.86 (m, 1H), 0.58 (m, 1H).

Scheme J

-74-

Example 163


Synthesis of 2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo
[4.1.0]hept-2-yl)-5-fluoro-phenyl ester



Stir a mixture of 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo
[4. 1.0]heptan-2-ol (295 mg, 1.06 mmol), 2,2-dimethyl-propionyl chloride (153 mg, 1.27 mmol)
and triethyl-amine (321 mg, 3.18 mmol) in CH_2Cl_2 (40 mL) at ambient temperature for 5 hours.
Quench the mixture with 10 mL of H_2O . Separate the aqueous layer off, and concentrate the
organic layer under reduced pressure. Purify the residue by preparative TLC to give
2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2- hydroxy-bicyclo
[4.1.0]hept-2-yl)-5-fluoro-phenyl ester as white solid (228 mg, Yield: 59.2%). MS (m/z): 364
(M+1). ^1H NMR (400 MHz, Methanol- d_4) δ 7.25-7.28 (d, $J_1 = 10.4$, $J_2 = 1.6$, 1H), 7.14 (s, 1H),
6.76-6.78 (d, $J_1 = 9.2$, $J_2 = 2.4$, 1H), 2.37-2.46 (m, 1H), 2.15-2.19 (m, 7H), 1.99-2.03 (d, $J = 13.2$,
1H), 1.82 (m, 2H), 1.53-1.68 (m, 2H), 1.37 (s, 9H), 1.17 (m, 1H), 1.01-1.06 (m, 1H), 0.75-0.81 (m,
1H), 0.48-0.50 (m, 1H).


The following compound may be prepared essentially by the method of Example 163.

-75-

Ex. No.	Chemical name	Structure	Physical data
164	2,2-Dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo[4.1.0]hept-2-yl)-4-fluoro-phenyl ester		MS (m/z): 364 (M+1) ¹ H NMR (400 MHz, Methanol-d ₄) δ 7.50-7.53 (m, 1H), 7.06-7.15 (m, 1H), 6.96-7.00 (m, 1H), 2.41-2.44 (m, 1H), 2.17-2.22 (m, 2H), 2.02 (s, 5H), 1.98-2.02 (m, 2H), 1.80-1.85 (m, 1H), 1.52-1.64 (m, 2H), 1.38 (s, 9H), 1.13-1.22 (m, 1H), 1.02-1.06 (m, 1H), 0.70-0.72 (m, 1H), 0.59-0.636 (m, 1H).


-76-

For compounds of the formula Ie, below, Scheme K and Preparations and/or Examples 165-176 illustrate methods of preparing them.



5


Scheme K



10

Preparation 165

(1R,2R,5S)-2,6,6-trimethylbicyclo[3.1.1]heptan-3-one




-77-

Jones reagent: add a mixture of 9 ml of H_2SO_4 and 40 ml of H_2O to a solution of Cr_2O_3 (15.0 g, 0.15 mol) in H_2O (20 mL) at 0 °C.

Add dropwise the above Jones reagent to a solution of (1R,2R,3R,5S)-2,6,6-trimethylbicyclo[3.1.1]heptan-3-ol (23.1 g, 0.15 mol) in acetone (100 mL) over a period for 2 hours at 0 °C. Stir the resultant mixture for additional 1 hour. Dilute the mixture with 100 mL of water and then extract the aqueous mixture with ether (50 mL X3). The combined organic layers are washed with brine (20 mL), dried over Na_2SO_4 , filtered, and concentrated under vacuum to give crude (1R,2R,5S)-2,6,6-trimethylbicyclo[3.1.1]heptan-3-one as pale yellow oil (21.3 g, yield: 93.4%), MS (m/z): 151 (M-1).

10


The following compound may be prepared essentially by the method of Preparation 165.

Prep No.	Chemical name	Structure	Physical data
166	(1S,2S,5R)-2,6,6-trimethylbicyclo[3.1.1]heptan-3-one		MS (m/z): 151 (M-1).

Preparation 167

(1R,2R,4S,5S)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-one

15




Stir a mixture of (1R,2R,5S)-2,6,6-trimethylbicyclo[3.1.1]heptan-3-one (7.6 g, 50.0 mmol), $(\text{HCHO})_n$ (1.8 g, 60.0 mmol), N-methyl-2-phenylethanamine (6.75 g, 50.0 mmol) and

-78-

5.0 mL of *conc.* HCl in MeCN (50 mL) at 70 °C for 3 hours. After removal of the solvent under vacuum, dissolve the residue in H₂O (30 mL) and washed with EtOAc (20 mL X 2). Basify the aqueous solution with K₂CO₃ to pH = 9 and extract the resultant mixture with EtOAc (30 mL×3). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum to give crude product as brown oil, which was further purified by silica gel chromatography to give (1R,2R,4S,5S)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-one as pale yellow oil (9.24 g, yield: 61.8%), MS (m/z): 300 (M+1).


10 The following compound may be prepared essentially by the method of Preparation 167.

Prep No.	Chemical name	Structure	Physical data
168	(1S,2S,4R,5R)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-one		MS (m/z): 300 (M+1).

Example 169

(1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol



15



-79-

Add a solution of *t*-BuLi (20 mL, 26.0 mmol) *via* syringe to a solution of 3-bromo-phenol (2.06 g, 12 mmol) in THF (20 mL) at -78°C under N₂. After being stirred at -78 °C for 1 hour, add a solution of (1R,2R,4S,5S)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-one (2.99 g, 10 mmol) in THF (2 mL) to the reaction mixture and stir the reaction mixture at -78 °C for additional 2 hours. Quench the reaction with saturated NH₄Cl solution (30 mL). Extract the aqueous mixture with EtOAc (60 mL x 3). The combined organic layers are washed with brine (30 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by Preparative HPLC to give (1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol as pale yellow oil (0.27 g, yield: 6.9%), MS (m/z): 394 (M+1).

The following compound may be prepared essentially by the method of Example 169.

Ex. No.	Chemical name	Structure	Physical data
170	(1R,2R,3S,4S,5S)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol		MS (m/z): 408 (M+1).
171	(1S,2S,3R,4R,5R)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol		MS (m/z): 394 (M+1).


-80-

Ex. No.	Chemical name	Structure	Physical data
172	(1S,2S,3R,4R,5R)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol		MS (m/z): 408 (M+1).

Example 173

(1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol hydrochloride

5



Add H₂O (4 mg, 0.25 mmol) and TMSCl (27 mg, 0.25 mmol) to a solution of (1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol (100 mg, 0.25 mmol) in 2-butanone (5 mL). Stir the reaction mixture at room temperature for 3 hours. After removal solvent by evaporation, wash the residue with EtOAc (1 mL X 2) and dry under vacuum to give (1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol hydrochloride (93 mg, Yield: 85.3%) as white solid. ¹H NMR (400 MHz, D₂O) δ 7.21-7.32 (m, 7H), 7.97-7.99 (d, J = 7.2, 1H), 6.71-6.78 (dd, J₁ = 8.0, J₂ = 22.0, 1H), 2.86-3.33 (m, 5H), 2.43-2.68 (m, 7H), 2.11-2.20 (m, 2H),

-81-

1.73-1.75 (m, 1H), 1.23-1.25 (d, $J = 5.6$, 3H), 1.04-1.05 (d, $J = 4.4$, 3H), 0.69-0.71 (d, $J = 5.6$, 3H). $[\alpha]_D^{20} = -8.6$ ($c = 1.4$ mg/mL, MeOH).


The following compounds may be prepared essentially by the method of Example 173.

Ex. No.	Chemical name	Structure	Physical data
174	(1R,2R,3S,4S,5S)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol hydrochloride		$^1\text{H NMR}$ (400 MHz, D_2O) δ 7.02-7.26 (m, 7H), 6.74-6.86 (m, 2H), 3.64-3.66 (d, $J = 10.4$, 3H), 2.76-3.22 (m, 5H), 2.32-2.53 (m, 7H), 2.02-2.14 (m, 2H), 1.64-1.67 (m, 1H), 1.14-1.56 (d, $J = 6.8$, 3H), 0.95-0.97 (d, $J = 6.4$, 3H), 0.63 (s, 3H). $[\alpha]_D^{20} = -16.1$ ($c = 6.4$ mg/mL, MeOH)
175	(1S,2S,3R,4R,5R)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol hydrochloride		MS (m/z): 394 ($M+1$). $[\alpha]_D^{20} = 9.3$ ($c = 4.5$ mg/mL, MeOH)

-82-


Ex. No.	Chemical name	Structure	Physical data
176	(1S,2S,3R,4R,5R)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol hydrochloride		MS (m/z): 408 (M+1). [α] _D ²⁰ = 14.3 (c = 6.0 mg/mL, MeOH)

For compounds of the formula If, below, Scheme L and Preparations and/or Examples 177-180 illustrate methods of preparing them.




Scheme L

-83-

Preparation 177

Synthesis of (2,3,3a,6,7,7a-hexahydro-1H-inden-4-yloxy)trimethylsilane

5




Add dropwise hexahydro-1H-inden-4(2H)-one (1.6 g, 11.6 mmol) in dry THF (5 mL) to a solution of LDA (23.2 mL, 23.2 mmol) in THF (30 mL). Stir the reaction mixture for 30 min and then add dropwise TMSCl (3.5 g, 32.0 mmol) to the reaction mixture. Stir the resultant solution for additional 1 hour. Pour the mixture into ice water (50 mL) and extract the aqueous mixture with EtOAc (50 mL x 3). The combined organic layers are washed with brine, dried over MgSO₄, filtered and concentrated in vacuo to afford crude (2,3,3a,6,7,7a-hexahydro-1H-inden-4-yloxy) trimethylsilane (2.45 g, 100 %) as yellowish oil, MS (m/z): 209 (M-1).

Preparation 178

5-((dimethylamino)methyl)hexahydro-1H-inden-4(2H)-one


-84-



Add (2,3,3a,6,7,7a-hexahydro-1H-inden-4-yloxy) trimethylsilane (2.45 g, 11.6 mmol) to a cooled suspension of N,N-dimethylmethyleiminium iodide (2.78 g, 15.0 mmol) in CH₂Cl₂ (60 mL) at 0 °C. After being stirred for overnight at ambient temperature, dilute the reaction mixture with 2 N HCl (20 mL). After removal of organic layers, wash the aqueous layer with EtOAc (30 mL x 3) and then basify with NaOH to PH = 10. Extract the resultant mixture with EtOAc (50 mL x 4). The combined organic layers are washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to afford 5-((dimethylamino)methyl)hexahydro-1H-inden-4(2H)-one (623 mg, 27.5 %) as oil, MS (m/z): 196 (M+1).

Example 179

5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol



15

Add a solution of *t*-BuLi (2.1 mL, 3.07 mmol) *via* syringe to a solution of 3-bromo-phenol (569 mg, 3.076 mmol) in THF (50 mL) at -78°C under N₂. After being stirred at -78 °C for 1 hour, add a solution of 5-((dimethylamino)methyl) hexahydro-1H-inden-4(2H)-one (300 mg, 1.54 mmol) in THF (2 mL) to the reaction mixture and stir the reaction mixture at -78 °C for additional 2 hours. Quench the reaction mixture with saturated NH₄Cl solution (30 mL). Extract the aqueous resultant mixture with EtOAc (50 mL x 3). The combined organic layers


-85-

are washed with brine (30 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by Preparative HPLC to give 5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol (92 mg, yield: 19.7%). MS (m/z): 290 (M+1).

5

Example 180

5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol hydrochloride





10 Add H₂O (10 mg, 0.54 mmol) and TMSCl (35 mg, 0.33 mmol) to a solution of 5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol (78 mg, 0.27 mmol) in 2-butanone (10 mL). Then stir the mixture at room temperature for 12 hours. Evaporate the mixture under vacuum to give 5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol hydrochloride as a mixture of 4 diastereoisomers
15 (91 mg, Yield: 100%).

20

Example 181

(+)-(1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl)bicyclo[3.2.1]octan-3-ol and
25 (-)-(1R,2S,3S,5S)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl)bicyclo[3.2.1]octan-3-ol.

-86-




5 Separate 0.5 g of 2-((dimethylamino)methyl)-3-(3-hydroxyphenyl)bicyclo[3.2.1]
 octan-3-ol (1.82 mmol) by SFC to afford (+)-(1S,2R,3R,5R)-2-((dimethylamino) methyl)-3-
 (3-hydroxyphenyl)bicyclo[3.2.1]octan-3-ol (8 mg; Yield: 3.2%; MS (m/z): 276 (M+1)) and
 (-)-(1R,2S,3S,5S)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl) bicyclo[3.2.1]octan-3-ol
 10 (135 mg, Yield: 54.0%; MS (m/z): 276 (M+1)) as white solids. The relative configuration of two
 enantiomers is confirmed by 2D NMR and the absolute configuration of the two
 enantiomers is determined by x-ray analysis. (+)-Enantiomers show better biological
 activity than (-)-enantiomer in bioassay(s).

15

Example 182

(S)-3-((1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl
 2-phenylpropanoate hydrochloride

-87-




- 5 Stir a solution of (S)-2-phenylpropanoic acid (300 mg, 2.0 mmol) in 5 mL of oxalyl chloride for 3 hours at room temperature. After removal of solvent under vacuum, (S)-2-phenylpropanoyl chloride is obtained as pale yellow oil. Dissolve the oil in 10 mL of CH₂Cl₂. Add (1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl) bicyclo[3.2.1]octan-3-ol (400 mg, 1.45 mmol) and Et₃N (293 mg, 2.9 mmol) to the solution. Stir the resultant
- 10 mixture at room temperature for additional 3 hours. After addition of 20 mL of water, basify the mixture with K₂CO₃ to pH = 10, and extract the aqueous mixture with EtOAc (20 mL X 3). The combined organic layers are washed with brine (20 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by silica gel chromatography (CH₂Cl₂:MeOH = 30:1) to afford (S)-3-((1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl) phenyl 2-phenylpropanoate (500 mg, 84.7%; MS (m/z): 408 (M+1)) as yellow oil. Stir
- 15 (S)-3-((1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl 2-phenylpropanoate (250 mg, 0.61 mmol) with H₂O (11 mg, 0.61 mmol) and TMSCl (66 mg, 0.61 mmol) in 2-butanone (5 mL) for 3 hours. Collect the precipitate by filtration to afford

-88-

(S)-3-((1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl 2-phenylpropanoate hydrochlorid (215 mg, Yield: 79.3%) as white solid. The absolute configuration of its hydrochloride salt form was confirmed by x-ray analysis.

¹H NMR (400 MHz, DMSO) δ 9.81 (br, 1H), 7.31-7.41 (m, 7H), 7.14 (s, 1H), 6.87-6.90 (m, 1H), 5.01 (s, 1H), 4.08-4.10 (m, 1H), 3.02-3.08 (m, 1H), 2.42-2.58 (m, 4H), 2.12-2.26 (m, 6H), 1.90-1.99 (m, 2H), 1.75-1.80 (m, 2H), 1.51-1.59 (m, 6H).


The following compound may be prepared essentially by the method of Example 182.

Ex. No	Chemical name	Structure	Physical data
183	(S)-3-((1R,2S,3S,5S)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl 2-phenylpropanoate hydrochloride		¹ H NMR (400 MHz, DMSO) δ 9.78 (br, 1H), 7.34-7.47 (m, 7H), 7.23 (s, 1H), 6.91-6.93 (m, 1H), 5.14 (s, 1H), 4.13-4.18 (m, 1H), 3.09-3.15 (m, 1H), 2.48-2.65 (m, 4H), 2.17-2.32 (m, 6H), 1.97-2.06 (m, 2H), 1.81-1.86 (m, 2H), 1.49-1.66 (m, 6H).

10

Example 184


3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol and 3-((1S,5R)-2-((dimethylamino)methyl)bicyclo[3.2.1]oct-2-en-3-yl)phenol




-89-

Add TsOH (5.0 g, 29.1 mmol) to a solution of 2-dimethylaminomethyl-3-(3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol (4.8 g, 17.5 mmol) in toluene (150 mL). Heat the reaction mixture to reflux for 2 hours and then quench the reaction by addition of saturated aqueous K₂CO₃ (20 mL). Extract the aqueous layer with EtOAc (60 mL x 3). The combined organic layers are washed with brine, dried over Na₂SO₄ and evaporated under vacuum. Purify the residue by preparative HPLC to yield 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol (2.27 g, 50.2 %; MS (m/z): 258 (M+1)) and 3-(2-((dimethylamino)methyl)bicyclo[3.2.1]oct-2-en-3-yl) phenol (517 mg, 11.5 %; MS (m/z): 258 (M+1)) as white solid. Two isomers show comparable biological activity in bioassay.


For compounds of the formula Ig, below, Scheme M and Preparations and/or Examples 185-187 illustrate methods of preparing them.

**Scheme M**

-90-

Preparation 185

5-dimethylaminomethyl-spiro[2.5]octan-6-one




Stir a mixture of spiro[2.5]octan-6-one (252 mg, 2.03 mmol), (HCHO)_n (61 mg, 2.03 mmol), dimethylamine hydrochloride (166 mg, 2.03 mmol) and 0.3 mL of *conc.* HCl in MeCN (30 mL) at 60 °C for 6 hours. Quench the reaction with saturated aqueous NH₄Cl solution (15 mL). Basify the aqueous solution with K₂CO₃ to pH ≈ 9. Extract the aqueous mixture with EtOAc (30 mL x 2). The combined organic layers are washed with brine (15 mL), dried over Na₂SO₄, filtered, and concentrated under vacuum. Purify the residue by silica gel chromatography (CH₂Cl₂:MeOH = 30:1) to afford 5-Dimethylaminomethyl-spiro[2.5]octan-6-one as brown oil (142mg, yield: 38.4%). MS (m/z): 182 (M+1).

Example 186

5-dimethylaminomethyl-6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan -6-ol


-91-



Cool a solution of (3-bromo-4-fluoro-phenoxy)-tert-butyl-dimethyl-silane (472 mg, 1.547 mmol) in THF (40 mL) to -78°C under N_2 . Then add dropwise a solution of *n*-BuLi (0.63 mL, 1.547 mmol) in hexane *via* syringe to the reaction solution. After being stirred at -78°C for 2 hour, add dropwise a solution of 5-Dimethylaminomethyl-spiro[2.5]octan-6-one (70 mg, 0.387 mmol) in THF (1 mL) to the reaction mixture and stir the mixture at -78°C for additional 2 hours. Quench the reaction with 20 mL of diluted HCl (2N), and stir at ambient temperature for 2 hours. Basify the resultant mixture with K_2CO_3 to pH = 9, and extract with EtOAc (30 mL X 2). The combined organic layers are washed with brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum. Purify the residue by silica gel chromatography (CH_2Cl_2 to $\text{CH}_2\text{Cl}_2:\text{MeOH} = 10:1$) to afford 5-dimethylaminomethyl -6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan-6-ol as white solid (41mg, yield: 36.3%). MS (m/z): 294 ($M+1$).

Example 187

15 5-dimethylaminomethyl-6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan -6-ol hydrochloride



Add H_2O (9 mg, 0.500 mmol) and TMSCl (18 mg, 0.167 mmol) to a solution of 5-dimethylaminomethyl-6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan-6-ol (41 mg, 0.139 mmol) in 2-butanone (30 mL). Stir the mixture at 0°C for 2 hours. Concentrate the mixture under vacuum to give 5-dimethylaminomethyl-6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan-6-ol hydrochloride as white solid (46 mg, Yield: 100%). ^1H NMR (400 MHz, D_2O) δ 7.13-7.15 (d, $J_1 = 6.8$, $J_2 = 3.2$, 1H), 6.93-6.98 (d, $J_1 = 12.0$, $J_2 = 8.8$, 1H), 6.69-6.73 (m, 1H), 3.05-3.10 (m, 1H),

-92-

2.65-2.76 (m, 8H), 2.44-2.45 (m, 1H), 2.27-2.35 (m, 2H), 1.70-1.74 (m, 1H), 0.96-0.98 (m, 1H), 0.79-0.82 (m, 1H), 0.45 (m, 1H).

Mu opioid agonists, such as morphine, are well known to control pain (Tschenkte *et al.*, Tapentadol Hydrochloride - Analgesic, Mu-opioid receptor agonist, Noradrenaline reuptake inhibitor. *E. Drugs Fut* 2006, 31(12): 1053). However, mu opioid agonists also can cause several undesired side effects such as nausea, emesis, constipation, mental clouding, and respiratory depression. In addition, use of opioids for an extended period of time (as is needed in chronic pain) can result in physical dependence and addiction.

Tramadol, a mu-opioid agonist, has not been associated with clinically significant side effects such as respiratory depression, constipation, or sedation (*Id.*). In addition, extended use of Tramadol does not lead to tolerance, dependence, and addiction (*Id.*). Tramadol is believed to exert its chronic pain relief through a combination of three mechanisms of action; mu opioid agonism and serotonin and norepinephrine reuptake inhibition (Raffa *et al.*, Opioid and nonopioid components independently contribute to the mechanism of action of tramadol, an 'atypical' opioid analgesic. *J Pharmacol Exp Ther.* 1992 Jan; 260(1):275-85). Because Tramadol relieves pain through a combination of mechanisms of action it is able to relieve pain even though it is a much less potent mu opioid receptor agonist compared to morphine. The reason for the better side effect profile of Tramadol as compared to morphine is believed to be a result of Tramadol's lower affinity for the mu opioid receptor. A molecule that, like Tramadol, controls chronic pain by multiple mechanisms of action is desired, and a compound which possessed a favorable side effect profile, particularly as compared to morphine, and that also was longer acting was sought. Preferably, a molecule that needed to be administered at most 1 time per day for extended pain relief is sought.

Molecules that are mu opioid agonists and also norepinephrine and/or serotonin reuptake inhibitors are preferably selected for evaluation in the rat tail flick model. In order to be selected for the rat tail flick model the compound needs to demonstrate an EC₅₀ of less than 50 micromolar in a functional assay for mu opioid receptor activation. In addition, the active species needs to demonstrate an IC₅₀ of less than 500 micromolar in a functional assay for inhibition of norepinephrine and/or serotonin reuptake. Activity of the corresponding phenols are used to select ether and ester prodrugs for evaluation in the rat tail flick model.

-93-

The Tail Flick Assay, based upon the methods of D'Amour & Smith (J. Pharmacol. Exp. Therap., 72: 74-79, 1941), is used to measure loss of pain sensation in the rat tail, allowing the experimenter to screen drugs for analgesic effect. Since then, the method has been used in a number of publications. For example, it was used to evaluate the analgesic effect of O-alkyl derivatives of tramadol by Liming Shao and Michael Hewitt etc. (Bioorganic & Medicinal Chemistry Letters, 18:1674-1680, 2008).

A Tail Flick Unit is used to measure tail flick latency in response to heat. The unit consists of an infrared (I.R.) source (50W bulb) with adjustable intensity that is focused through a flush mounted window on the upper panel of the unit and onto the rat tail. Once I.R. source is focused on the rat tail and turned on, a timer starts. When the thermal threshold for pain is reached and the rat flicks its tail, the I.R. source automatically shuts off and the timer stop, displaying the latency time.

On the day of experiment, the animals are tested to determine baseline latency. Each rat is given one test to determine latency to tail flick with a cut off 10 seconds. Baselines are recorded and those animals with a baseline latency of 2-5 seconds are used in the experiment. Animals are divided into several groups according to the baseline latency. The Tail-flick latencies are measured at different timepoints for up to 180 minutes after administration of vehicle or test articles. If treatment group continues to display pain control after 180 minutes, additional measurements are taken every 60 minutes until pain control is no longer observed.

I.R. Source is set at 40 units (determined to elicit tail flick response in desired 2-5 seconds in naïve animals). A cut off time of 10 seconds is set to avoid tissue damage, in the event that the animal does not flick its tail.

Test articles or vehicle are administered by intravenous (IV) injection through the tail vein or by oral gavage at time 0. Animals are gently restrained and held on top of a tail flick unit. The tail is then wiped clean with a cotton square and placed over the I.R. source so that the beam is focused at approximately midway the length of the tail. After the rat is in position, the I.R. source is turned on. When the thermal threshold for pain is reached, the rat flicked its tail and the I.R. source automatically shuts off. Latency time (in seconds) is then recorded.

Data are recorded and graphed as %MPE (Maximum Possible Effect). % MPE is calculated using the following formula:

-94-

% MPE = [(Test latency – Baseline latency) / (Cut off latency (10 sec.'s) – Baseline latency)] *100

Compounds were administered by the IV route in 20%

2-hydroxypropyl-beta-cyclodextrin (20% HP-β-CD). The dose levels of test article were 2.5, 5,
5 or 20 mg/kg. Examples 2, 3, 13, 40, 77, 78, 114, and Tramadol provided at least 50% MPE at 20
mg/kg. Compounds 5, 8, 17, 105, 107, 148, 158, and 159 provided at least 50% MPE at 5 mg/kg.
Compounds 82, 84, and 116 provided at least 50% MPE at 2.5 mg/kg.


Compounds were administered by oral gavage in 0.5% methylcellulose at 20 or 30
mg/kg. Examples 2, 3, 13, 115, and Tramadol provided at least 50% MPE at 30 mg/kg.

10 Compounds 5, 39, 65, 106, and 115 provided at least 50% MPE at 20 mg/kg.


The data above supports the contention the compounds are useful in controlling
pain.

WE CLAIM:

1. A compound, or salt thereof, of formula I:



wherein A is



R^1 is hydrogen, C_1-C_5 alkyl, C_1-C_5 haloalkyl, C_1-C_5 alkanol, $-(C_1-C_5$ alkyl)phenyl, or phenyl, or a group of the formula $-C(O)-R^{12}$, where R^{12} is C_1-C_5 alkyl, C_1-C_5 alkoxy, C_1-C_5 haloalkyl, C_1-C_5 alkanol, $-(C_1-C_5$ alkyl)phenyl, or phenyl;

R^2 is hydrogen, C_1-C_5 alkyl, C_1-C_5 alkoxy, halogen, C_1-C_5 haloalkyl, or C_1-C_5 haloalkoxy;

R^3 is hydrogen, C_1-C_5 alkyl, C_1-C_5 alkoxy, halogen, C_1-C_5 haloalkyl, or C_1-C_5 haloalkoxy;

R^4 is hydrogen, C_1-C_5 alkyl, or $-(C_1-C_5$ alkyl)phenyl;

R^5 is hydrogen, C_1-C_5 alkyl, or $-(C_1-C_5$ alkyl)phenyl;

R^6 is hydrogen, hydroxy, or is absent;

R^7 is hydrogen;

R^8 is hydrogen or methyl;

R^9 is hydrogen or methyl;

R^{10} is hydrogen;

R^{11} is hydrogen or C_1-C_5 alkyl;


provided that one of I), II), and III) is present, wherein I), II) and III) are:

I) R^7 and R^{10} combine to form $-CH_2-$ or $-(CH_2)_2-$;


II) R^8 and R^9 combine to form a cyclopropyl group with the carbon to which they are attached; and

III) R^{10} and R^{11} combine to form $-\text{CH}_2-$ or $-(\text{CH}_2)_3-$.

2. The compound, or salt thereof, of claim 1 wherein A is selected from



3. The compound of claim 2 of the formula Ia:



Ia,

or a salt thereof.

4. The compound of claim 2 wherein it is

2-dimethylaminomethyl-3-(3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;

2-dimethylaminomethyl-3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;


2-dimethylaminomethyl-3-(5-methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-

-ol; 2-dimethylaminomethyl-3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(4-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(2-fluoro-3-methoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 3-(2-chloro-5-methoxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol;
 3-(3-chloro-5-methoxyphenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(5-methoxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-methoxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-hydroxy-5-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(5-hydroxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-hydroxy-4-trifluoromethyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(4-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(2-fluoro-3-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-hydroxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 3-(2-chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol;
 3-(3-chloro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(5-hydroxy-2-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(3-hydroxy-4-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol;
 3-(3,4-difluoro-5-hydroxy-phenyl)-2-dimethylaminomethyl-bicyclo[3.2.1]octan-3-ol;
 2-dimethylaminomethyl-3-(2-fluoro-5-hydroxy-phenyl)-bicyclo[3.2.1]octan-3-ol;
 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester;
 benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester;
 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl ester;
 benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-phenyl ester;

2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester; or

benzoic acid 3-(2-dimethylaminomethyl-3-hydroxy-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester; or a salt thereof.

5. The compound of claim 2 of the formula Ib:



Ib,

or a salt thereof.

6. The compound of claim 2 wherein it is

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol;

[3-(3-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;

[3-(5-methoxy-2-trifluoromethoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-trifluoromethoxy-phenol;

[3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluoro-phenol;

[3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluoro-phenol;

[3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-methyl-phenol;

3-chloro-5-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol;

[3-(5-methoxy-2-methyl-phenyl)-bicyclo[3.2.1]oct-3-en-2-ylmethyl]-dimethyl-amine;


3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-methyl-phenol;

4-chloro-3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol;

2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester;

benzoic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenyl ester;
 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluoro-phenyl ester;
 isobutyric acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-5-fluoro-phenyl ester;
 2,2-dimethyl-propionic acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl) -4-fluoro-phenyl ester; or
 isobutyric acid 3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-4-fluoro-phenyl ester;
 or a salt thereof.

7. The compound of claim 2 wherein it is of the formula Ic:




or a salt thereof.

8. The compound of claim 2 wherein it is:

3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenol;
 [3-(3-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine;
 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenol;
 [3-(3-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine;
 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenol;
 [3-(2-fluoro-5-methoxy-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine;
 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-methyl-phenol;
 [3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]oct-2-ylmethyl]-dimethyl-amine;
 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct -3-yl)-phenyl ester;
 benzoic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-phenyl ester;

2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester;
 isobutyric acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-5-fluoro-phenyl ester;
 2,2-dimethyl-propionic acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester;
 or isobutyric acid 3-(2-dimethylaminomethyl-bicyclo[3.2.1]oct-3-yl)-4-fluoro-phenyl ester;
 or a salt thereof.

9. The compound of claim 2 wherein it is of the formula Id:




Id

or a salt thereof.

10. The compound of claim 2 wherein it is

3-dimethylaminomethyl-2-(3-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-fluoro-5-methoxy-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(5-methoxy-2-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-methoxy-phenyl)-5,5-dimethyl-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-hydroxy-phenyl)-5,5-dimethyl-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(2-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 3-dimethylaminomethyl-2-(3-hydroxy-5-methyl-phenyl)-bicyclo[4.1.0]heptan-2-ol;
 2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo[4.1.0]hept-2-yl)-5-fluoro-phenyl ester; or
 2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo[4.1.0]hept-2-yl)-4-fluoro-phenyl ester; or a salt thereof.

11. The compound of claim 2 wherein it is of the formula Ie:



Ie,

or a salt thereof.

12. The compound of claim 11 wherein it is

(1R,2R,3S,4S,5S)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-

((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol;

(1R,2R,3S,4S,5S)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-

((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol;


(1S,2S,3R,4R,5R)-3-(3-hydroxyphenyl)-2,6,6-trimethyl-4-

((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol;

(1S,2S,3R,4R,5R)-3-(3-methoxyphenyl)-2,6,6-trimethyl-4-

((methyl(phenethyl)amino)methyl)bicyclo[3.1.1]heptan-3-ol; or a salt thereof.

13. The compound of claim 2 wherein it is of the formula If:



If,

or a salt thereof.

14. The compound of claim 13 wherein it is 5-((dimethylamino)methyl)-4-(3-hydroxyphenyl)octahydro-1H-inden-4-ol, or a salt thereof.

15. The compound of claim 2 wherein it is

(+)-(1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl)bicyclo[3.2.1]octan-3-ol;

(-)-(1R,2S,3S,5S)-2-((dimethylamino)methyl)-3-(3-hydroxyphenyl)bicyclo[3.2.1]octan-3-

ol;


(S)-3-((1S,2R,3R,5R)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl 2-phenylpropanoate hydrochloride;

(S)-3-(1R,2S,3S,5S)-2-((dimethylamino)methyl)-3-hydroxybicyclo[3.2.1]octan-3-yl)phenyl 2-phenylpropanoate hydrochloride;

3-(4-dimethylaminomethyl-bicyclo[3.2.1]oct-2-en-3-yl)-phenol;

3-((1S,5R)-2-((dimethylamino)methyl)bicyclo[3.2.1]oct-2-en-3-yl)phenol; or a salt thereof.

16. A compound of claim 2 wherein it is of the formula Ig:



Ig,

or a salt thereof.

17. The compound of claim 16 where it is 5-dimethylaminomethyl-6-(2-fluoro-5-hydroxy-phenyl)-spiro[2.5]octan-6-ol, or a salt thereof.

18. The compound of claim 4 wherein it is 2-dimethylaminomethyl-3-(3-methoxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol, or a salt thereof.

19. The compound of claim 4 wherein it is 2-dimethylaminomethyl-3-(3-hydroxy-5-methyl-phenyl)-bicyclo[3.2.1]octan-3-ol, or a salt thereof.

20. The compound of claim 10 wherein it is 3-dimethylaminomethyl-2-(3-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol, or a salt thereof.

21. The compound of claim 10 wherein it is 3-dimethylaminomethyl-2-(2-fluoro-5-hydroxy-phenyl)-bicyclo[4.1.0]heptan-2-ol, or a salt thereof.
22. The compound of claim 10 wherein it is 2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo[4.1.0]hept-2-yl)-5-fluoro-phenyl ester, or a salt thereof.
23. The compound of claim 10 wherein it is 2,2-dimethyl-propionic acid 3-(3-dimethylaminomethyl-2-hydroxy-bicyclo[4.1.0]hept-2-yl)-4-fluoro-phenyl ester, or a salt thereof.
24. The compound of any one of claims 18-23, wherein it is the hydrochloride salt thereof.
25. A pharmaceutical formulation comprising a compound of any of claims 1-24 and one or more pharmaceutically acceptable carriers.
26. The pharmaceutical formulation of claim 25 wherein it further comprises at least one additional active ingredient.
27. The pharmaceutical formulation of claim 25 wherein it is a human pharmaceutical formulation.
28. The pharmaceutical formulation of claim 25 wherein it is a veterinary pharmaceutical formulation.
29. Use of an effective amount of a compound of any of claims 1 through 24 in the manufacture of a medicament for controlling pain in a mammal.

30. The use of claim 29 wherein the medicament comprises at least one other active ingredient.
31. The use of claim 29 or claim 30, wherein said mammal is a human.
32. The use of claim 29 or claim 30, wherein said mammal is a companion animal.
33. The use of claim 32, wherein said companion animal is a dog or cat.
34. A compound or salt according to any one of claims 1 to 24 for use in therapy.
35. The compound or salt according to claim 34, wherein the use in therapy comprises controlling pain.