

KINASE INHIBITORS

This invention relates to compounds and compositions that are p38 MAPK inhibitors, useful as anti-inflammatory agents in the treatment of, *inter alia*, diseases of the respiratory tract.

Background to the invention

5 Mitogen activated protein kinases (MAPK) constitute a family of proline-directed serine/threonine kinases that activate their substrates by dual phosphorylation. There are four known human isoforms of p38 MAP kinase, p38 α , p38 β , p38 γ and p38 δ . The p38 kinases, which are also known as cytokine suppressive anti-inflammatory drug binding proteins (CSBP), stress
10 activated protein kinases (SAPK) and RK, are responsible for phosphorylating (Stein *et al.*, Ann. Rep. Med Chem., 1996, 31, 289-298) and activating transcription factors (such as ATF-2, MAX, CHOP and C/ERPB) as well as other kinases (such as MAPKAP-K2/3 or MK2/3), and are themselves activated by physical and chemical stress (e.g. UV, osmotic stress),
15 pro-inflammatory cytokines and bacterial lipopolysaccharide (LPS) (Herlaar E. & Brown Z., Molecular Medicine Today, 1999, 5, 439-447). The products of p38 phosphorylation have been shown to mediate the production of inflammatory cytokines, including tumor necrosis factor alpha (TNF α) and interleukin- (IL)-1, and cyclooxygenase-2 (COX-2). IL-1 and TNF α are also
20 known to stimulate the production of other proinflammatory cytokines such as IL-6 and IL-8.

IL-1 and TNF α are biological substances produced by a variety of cells, such as monocytes or macrophages. IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation
25 and other physiological conditions such as inflammation (e.g. Dinarello *et al.*, Rev. Infect. Disease, 1984, 6, 51). Excessive or unregulated TNF production

(particularly TNF α) has been implicated in mediating or exacerbating a number of diseases, and it is believed that TNF can cause or contribute to the effects of inflammation in general. IL-8 is a chemotactic factor produced by several cell types including mononuclear cells, fibroblasts, endothelial cells, and keratinocytes. Its production from endothelial cells is induced by IL-1, TNF, or lipopolysaccharide (LPS). IL-8 stimulates a number of functions *in vitro*. It has been shown to have chemoattractant properties for neutrophils, T-lymphocytes and basophils. Increase in IL-8 production is also responsible for chemotaxis of neutrophils into the inflammatory site *in vivo*.

Inhibition of signal transduction *via* p38, which in addition to IL-1, TNF and IL-8 described above is also required for the synthesis and/or action of several additional pro-inflammatory proteins (e.g., IL-6, GM-CSF, COX-2, collagenase and stromelysin), is expected to be a highly effective mechanism for regulating the excessive and destructive activation of the immune system. This expectation is supported by the potent and diverse anti-inflammatory activities described for p38 kinase inhibitors (Badger *et al.*, J. Pharm. Exp. Thera., 1996, 279, 1453 -1461; Griswold *et al*, Pharmacol. Comm., 1996, 7, 323-229). In particular, p38 kinase inhibitors have been described as potential agents for treating rheumatoid arthritis. In addition to the links between p38 activation and chronic inflammation and arthritis, there is also data implicating a role for p38 in the pathogenesis of airway diseases in particular COPD and asthma. Stress stimuli (including tobacco smoke, infections or oxidative products) can cause inflammation within the lung environment. Inhibitors of p38 have been shown to inhibit LPS and ovalbumin induced airway TNF- α , IL-1 β , IL-6, IL-4, IL-5 and IL-13 (Haddad *et al*, Br. J. Pharmacol., 2001, 132 (8), 1715-1724; Underwood *et al*, Am. J. Physiol. Lung Cell. Mol. 2000, 279, 895-902; Duan *et al.*, 2005 Am. J. Respir. Crit. Care Med., 171, 571-578; Escott *et al* Br. J. Pharmacol., 2000, 131, 173-176;


Underwood *et al.*, J. Pharmacol. Exp. Ther. 2000, 293, 281-288). Furthermore, they significantly inhibit neutrophilia and the release of MMP-9 in LPS, ozone or cigarette smoke animal models. There is also a significant body of preclinical data highlighting the potential benefits of inhibition of the p38 kinase that could be relevant in the lung (Lee *et al.*, Immunopharmacology, 2000, 47, 185-200). Thus, therapeutic inhibition of p38 activation may be important in the regulation of airway inflammation.

The implication of the p38MAPK pathway in various diseases has been reviewed by P. Chopra *et al.* (Expert Opinion on Investigational Drugs, 2008, 17(10), 1411-1425). It is believed that the compounds of the present invention can be used to treat p38 mediated diseases such as: chronic obstructive pulmonary disease (COPD), asthma, chronic or acute bronchoconstriction, bronchitis, acute lung injury and bronchiectasis, pulmonary artery hypertension, tuberculosis, lung cancer, inflammation generally (e.g. inflammatory bowel disease), arthritis, neuroinflammation, pain, fever, fibrotic diseases, pulmonary disorders and diseases (e.g., hyperoxic alveolar injury), cardiovascular diseases, post -ischemic reperfusion injury and congestive heart failure, cardiomyopathy, stroke, ischemia, reperfusion injury, renal reperfusion injury, brain edema, neurotrauma and brain trauma, neurodegenerative disorders, central nervous system disorders, liver disease and nephritis, gastrointestinal conditions, ulcerative diseases, Crohn's disease, ophthalmic diseases, ophthalmological conditions, glaucoma, acute injury to the eye tissue and ocular traumas, diabetes, diabetic nephropathy, skin-related conditions, myalgias due to infection, influenza, endotoxic shock, toxic shock syndrome, autoimmune disease, graft rejection, bone resorption diseases, multiple sclerosis, psoriasis, eczema, disorders of the female reproductive system, pathological (but non -malignant) conditions, such as hemangiomas, angiofibroma of the nasopharynx, and avascular necrosis of bone, benign and

malignant tumors/neoplasia including cancer, leukaemia, lymphoma, systemic lupus erythematosus (SLE), angiogenesis including neoplasia, haemorrhage, coagulation, radiation damage, and/or metastasis. Chronic release of active TNF can cause cachexia and anorexia, and TNF can be lethal. TNF has also
5 been implicated in infectious diseases. These include, for example, malaria, mycobacterial infection and meningitis. These also include viral infections, such as HIV, influenza virus, and herpes virus, including herpes simplex virus type-1 (HSV-1), herpes simplex virus type-2 (HSV-2), cytomegalovirus (CMV), varicella-zoster virus (VZV), Epstein-Barr virus, human herpes virus-
10 6 (HHV-6), human herpesvirus-7 (HHV7), human herpesvirus-8 (HHV-8), pseudorabies and rhinotracheitis, among others.

Known P38 kinase inhibitors have been reviewed by G. J. Hanson (Expert Opinions on Therapeutic Patents, 1997, 7, 729-733) J Hynes *et al.* (Current Topics in Medicinal Chemistry, 2005, 5, 967-985), C. Dominguez *et al.*
15 *al* (Expert Opinions on Therapeutics Patents, 2005, 15, 801-816) and L. H. Pettus & R. P. Wurtz (Current Topics in Medicinal Chemistry, 2008, 8, 1452-1467). P38 kinase inhibitors containing a triazolopyridine motif are known in the art, for example WO07/091152, WO04/072072, WO06/018727.

International Patent Application WO2010/094956 discloses
20 triazolopyridine derivatives of formula (I) as being p38 MAP Kinase inhibitors:



In such compounds, A represents an optionally substituted divalent arylene radical, an heteroarylene radical, a (C₃-C₆) divalent cycloalkylene
25 radical having 5 or 6 ring atoms or a piperidinylene radical.

The compounds are said to be useful in as anti-inflammatory agents in the treatment of diseases of the respiratory tract.

The object of the present invention is to identify novel and potent p38 mitogen activated protein kinase inhibitors which are useful in the treatment
5 of inflammatory and obstructive diseases of the respiratory tract.


It is another object of the present invention, to identify novel potent p38 mitogen activated protein kinase inhibitors which show an appropriate developability profile on inhalatory administration to effectively treat respiratory obstructive or inflammatory diseases. It is to be understood that
10 such profile may be achieved in a number of different ways by modulation of specific properties; by way of example, it could be achieved by administration of a low effective dose of the drug thus limiting side effects or via a long duration of action in the lungs which may reduce the frequency of administration.

15 **Brief Description of the Invention**

The compounds of the present invention are inhibitors of p38 mitogen activated protein kinase ("p38 MAPK", "p38 kinase" or "p38"), including p38 α kinase, and are inhibitors of cytokine and chemokine production including TNF α and IL-8 production. They have a number of therapeutic
20 applications, in the treatment of inflammatory diseases, particularly allergic and non-allergic airways diseases, more particularly obstructive or inflammatory airways diseases such as chronic obstructive pulmonary disease ("COPD") and asthma. They are therefore particularly suited for pulmonary delivery, by inhalation by nose or mouth.

25 **Description of the invention**

According to the invention there is provided a compound of formula (I), or a pharmaceutically acceptable salt thereof:



wherein;

W is a heteroatom selected from N or O, wherein N is substituted with hydrogen, C₁-C₆ alkyl or C₃-C₅ cycloalkyl;


Y is selected in the group consisting of: a group -S(O)_p- wherein **p** is 0, 1 or 2; a group -O(CR³R⁴)_n-; a group -(CR⁵R⁶)_n-; a group -NR⁷-; a group -OC(O)-; a group -OC(O)NH-; and a group -OC(O)O-;

R³, R⁴, R⁵ and R⁶ are each independently hydrogen, fluorine or C₁-C₆ alkyl, or, respectively, **R³ and R⁴**, or **R⁵ and R⁶** may form together with the carbon atom to which they are attached a saturated 3-6 membered carbocyclic monocyclic ring optionally substituted by a group C₁-C₆ alkyl, hydroxyl or halo;


n is 0, 1, 2 or 3;

R⁷ is hydrogen, C₁-C₆ alkyl, or C₃-C₇ cycloalkyl wherein such C₁-C₆ alkyl or C₃-C₇ cycloalkyl are optionally substituted by a group C₁-C₃ alkyl, C₃-C₆ cycloalkyl, hydroxyl, cyano or halo;


R¹ is a group selected from (IIa) - (IIc):



(IIa)



(IIb)



(IIc)

R⁸ and R⁹ are each independently hydrogen or C₁-C₆ alkyl, or **R⁸ and R⁹** may form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or a fused or spiro bicyclic ring system optionally containing a further heteroatom which is oxygen or nitrogen, said nitrogen atom being optionally substituted by C₁-C₆ alkyl; wherein such C₁-C₆

alkyl groups may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₆ cycloalkyl, hydroxyl or halo;

X¹, X², X³, X⁴ and **X⁵** are each independently a carbon atom, a nitrogen atom, a group -(CH)- or a group -NH-; such that each combination thereof
 5 forms an aromatic ring system;

R¹⁰ is selected from a group consisting of: Hydrogen, -CN, -NR^AR^B,
 -N(R^C)(C₂-C₆alkylene)-NR^AR^B, -N(R^C)(C₃-C₇cycloalkylene)-NR^AR^B,
 -(C₁-C₆alkylene)-NR^AR^B, -(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₂-C₆alkylene)-
 NR^AR^B, -O-(C₃-C₇cycloalkylene)-NR^AR^B, -S-(C₂-C₆alkylene)-NR^AR^B, -S-(C₃-
 10 C₇cycloalkylene)-NR^AR^B, -N(R^C)C(O)-(C₁-C₆alkylene)-NR^AR^B, -N(R^C)C(O)-
 (C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B,
 -C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-OR^D,
 -C(O)N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)C(O)NR^AR^B, -C(O)NR^AR^B,
 -N(R^C)C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -N(R^C)C(O)N(R^C)-(C₃-
 15 C₇cycloalkylene)-NR^AR^B, -(C₂-C₆alkylene)-OR^D, -(C₃-C₇cycloalkylene)-OR^D,
 -O-(C₂-C₆alkylene)-OR^D, -O-(C₃-C₇cycloalkylene)-OR^D, -S-(C₂-C₆alkylene)-
 OR^D, -S-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₁-C₆alkylene)-NR^AR^B,
 -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-
 NR^AR^B, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-
 20 C₆alkylene)-OR^D, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₂-
 C₆alkylene)-OR^D, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-OR^D, -S(O)₂N(R^AR^B),
 -N(R^C)S(O)₂R^D, -N(R^C)C(O)R^C, -OR^C, -SR^C, -(C₃-C₇heterocycloalkyl),
 (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), (C₅-C₇heterocycloalkyl)(C₃-
 C₆cycloalkyl)-, and C₃-C₇ heterocycloalkylcarbonyl; wherein any of such C₁-
 25 C₆alkyl, C₃-C₆cycloalkyl, -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-
 C₇cycloalkylene)-, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆
 alkyl), (C₅-C₇ heterocycloalkyl)-(C₃-C₆ cycloalkyl) and (C₃-
 C₇heterocycloalkyl)carbonyl portion in the above listed groups may be

optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, hydroxyl or halo;

R¹¹ is linked to X⁴ and is selected from a group consisting of:

Hydrogen; -CN; C₁-C₆ alkyl which is substituted by a group selected
 5 from -CN, -OR^C, -SR^C, halo; C₃-C₆cycloalkyl which is substituted by a group selected from C₁-C₄ alkyl, -CN, -OR^C, -SR^D, halo; -NR^AR^B, -N(R^C)(C₂-C₆alkylene)-NR^AR^B, -N(R^C)(C₃-C₇cycloalkylene)-NR^AR^B, -(C₁-C₆alkylene)-NR^AR^B, -(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₂-C₆alkylene)-NR^AR^B, -O-(C₃-C₇cycloalkylene)-NR^AR^B, -S-(C₂-C₆alkylene)-NR^AR^B, -S-(C₃-C₇cycloalkylene)-NR^AR^B, -N(R^C)C(O)-(C₁-C₆alkylene)-NR^AR^B, -N(R^C)C(O)-(C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-OR^D, -C(O)N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)C(O)N(R^AR^B), -C(O)N(R^AR^B), -N(R^C)C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -N(R^C)C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₂-C₆alkylene)-OR^D, -O-(C₃-C₇cycloalkylene)-OR^D, -S-(C₂-C₆alkylene)-OR^D, -S-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₁-C₆alkylene)-NR^AR^B, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-OR^D, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₂-C₆alkylene)-OR^D, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-OR^D, -S(O)₂N(R^AR^B), -N(R^C)S(O)₂R^D, -N(R^C)C(O)R^C, OR^C, SR^C, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), (C₅-C₇heterocycloalkyl)(C₃-C₆cycloalkyl) and (C₃-C₇heterocycloalkyl)carbonyl, wherein any of such C₁-C₆alkyl, C₃-C₆cycloalkyl, -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-C₇cycloalkylene)-, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl), (C₅-C₇heterocycloalkyl)-(C₃-C₆ cycloalkyl) and (C₃-C₇heterocycloalkyl)carbonyl
 10
 15
 20
 25 portion in the above listed groups may be optionally substituted by one, two or

three groups R^{25} which are independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; or

- R^{11} is linked to X^4 and is phenyl or 5- or 6-membered monocyclic heteroaryl, wherein such phenyl or 5- or 6-membered monocyclic heteroaryl is substituted by a group selected in the list consisting of: C_1 - C_6 alkyl which is substituted by a group -CN; C_3 - C_6 cycloalkyl which is substituted by a group selected from: -CN, $-OR^C$, $-SR^C$ or halo; $-N(R^C)(C_2$ - C_6 alkylene)- $NR^A R^B$, $-N(R^C)(C_3$ - C_7 cycloalkylene)- $NR^A R^B$, $-(C_1$ - C_6 alkylene)- $NR^A R^B$, $-(C_3$ - C_7 cycloalkylene)- $NR^A R^B$, $-O$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-S$ -(C_2 - C_6 alkylene)- $NR^A R^B$, $-S$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-N(R^C)C(O)$ -(C_1 - C_6 alkylene)- $NR^A R^B$, $-N(R^C)C(O)$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-C(O)N(R^C)$ -(C_2 - C_6 alkylene)- $NR^A R^B$, $-C(O)N(R^C)$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-C(O)N(R^C)$ -(C_2 - C_6 alkylene)- OR^D , $-C(O)N(R^C)$ -(C_3 - C_7 cycloalkylene)- OR^D , $-N(R^C)C(O)N(R^C)$ -(C_2 - C_6 alkylene)- $NR^A R^B$, $-N(R^C)C(O)N(R^C)$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-O$ -(C_3 - C_7 cycloalkylene)- OR^D , $-S$ -(C_3 - C_7 cycloalkylene)- OR^D , $-N(R^C)S(O)_2$ -(C_1 - C_6 alkylene)- $NR^A R^B$, $-N(R^C)S(O)_2$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-S(O)_2N(R^C)$ -(C_2 - C_6 alkylene)- $NR^A R^B$, $-S(O)_2N(R^C)$ -(C_3 - C_7 cycloalkylene)- $NR^A R^B$, $-S(O)_2N(R^C)$ -(C_2 - C_6 alkylene)- OR^D , $-S(O)_2N(R^C)$ -(C_3 - C_7 cycloalkylene)- OR^D , $-N(R^C)S(O)_2$ -(C_2 - C_6 alkylene)- OR^D , $-N(R^C)S(O)_2$ -(C_3 - C_7 cycloalkylene)- OR^D , $-N(R^C)S(O)_2R^D$, $-(C_3$ - C_7 heterocycloalkyl), $(C_5$ - C_7 heterocycloalkyl)-(C₁-C₆alkyl), $(C_5$ - C_7 heterocycloalkyl)(C₃-C₆cycloalkyl) and $(C_3$ - C_7 heterocycloalkyl)carbonyl, wherein any of such C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $-(C_1$ - C_6 alkylene)-, $-(C_2$ - C_6 alkylene)-, $-(C_3$ - C_7 cycloalkylene)-, $-(C_3$ - C_7 heterocycloalkyl), $(C_5$ - C_7 heterocycloalkyl)-(C₁-C₆ alkyl), $(C_5$ - C_7 heterocycloalkyl)-(C₃-C₆ cycloalkyl) and $(C_3$ - C_7 heterocycloalkyl)carbonyl portion in the above listed groups may be optionally substituted by one, two or

three groups R^{25} which are independently selected in the group consisting of: C_1 - C_6 alkyl, (C_1-C_3) haloalkyl, (C_1-C_4) hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo;

R^A and R^B are at each occurrence independently hydrogen, C_1 - C_6 alkyl
5 or C_3 - C_7 cycloalkyl, such C_1 - C_6 alkyl and C_3 - C_7 cycloalkyl being optionally substituted by a group C_1 - C_3 alkyl, C_3 - C_7 cycloalkyl, $-OR^D$, $-CN$ or halo; alternatively, R^A and R^B , may form together with the nitrogen atom to which they are attached a 5-11- membered saturated heterocyclic monocyclic or bicyclic ring system which is optionally substituted by one or more group
10 $-OR^D$, $-CN$, halo, C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl, such C_1 - C_6 alkyl and C_3 - C_7 cycloalkyl being optionally substituted by a group C_1 - C_3 alkyl, C_3 - C_7 cycloalkyl, $-OR^D$, $-CN$ or halo; and which 5-11- membered saturated heterocyclic monocyclic or bicyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally
15 substituted by C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, wherein any of such C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, $-OR^D$, $-CN$, or halo; and/or R^A and R^B may be linked to one carbon atom of the $-(C_1-C_6\text{alkylene})-$, $-(C_2-C_6\text{alkylene})-$ or $-(C_3-C_7\text{cycloalkylene})-$ portion of the group linked to the nitrogen to which they are
20 connected to form a saturated cycle of up to 6 ring atoms;

R^C is at each occurrence independently hydrogen, C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, such C_1 - C_6 alkyl and C_3 - C_6 cycloalkyl being optionally substituted by a group C_1 - C_3 alkyl, $-OR^D$, $-CN$ or halo;

R^D is at each occurrence independently hydrogen, $-CH_3$ or $-C_2H_5$;


25 R^{12} and R^{13} are independently hydrogen, C_1 - C_6 alkyl, or halogen;

A is a divalent cycloalkylene radical having 5, 6 or 7 ring atoms; said cycloalkylene ring being attached to W and Y , and fused to a phenyl ring or to a monocyclic heteroaryl ring having 5 or 6 ring atoms, such phenyl or


heteroaryl ring being optionally substituted by one or two groups R^{24} ;

R^{24} is at each occurrence independently selected from the group consisting of: C_1 - C_6 alkyl, halogen and cyano;


R^2 is a radical of formula (IIIa), (IIIb), (IIIc), or (IIId):




(IIIa)



(IIIb)



(IIIc)



(IIId)


wherein

R^{14} is selected in the group consisting of: -F, -CH₃, -C₂H₅, -CH₂OH, -CH₂OMe, -CF₂CF₃, -CH₂SCH₃, -SCH₃ and -SC₂H₅;

R^{15} and R^{16} are independently -CH₃ or -C₂H₅;

R^{17} is selected from the group consisting of: lone electron pair, hydrogen, -CF₃, -NR^ER^F, -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), aryl or heteroaryl wherein any of such -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), aryl or heteroaryl may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, or halo; or

R^{17} is a group of general formula (IV)



(IV)

wherein

R^{20} is selected in the group consisting of: -F, -CH₃, -C₂H₅, -CH₂OH, -CH₂OMe, -CF₂CF₃, -CH₂SCH₃, -SCH₃ and -SC₂H₅;

R^{21} is $-CH_3$ or $-C_2H_5$;

or

R^{20} and R^{21} as defined above may form together with the carbon atom to which they are attached a saturated 3-7- membered monocyclic ring;

- 5 R^E and R^F are each independently C_1 - C_6 alkyl, optionally substituted by a group C_1 - C_3 alkyl, $-OR^G$, $-CN$ or halo; alternatively, R^E and R^F may form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more groups $-OR^G$, $-CN$, halo, C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl, such C_1 - C_6 alkyl and C_3 - C_7 cycloalkyl being optionally substituted
10 by a group C_1 - C_3 alkyl, C_3 - C_7 cycloalkyl, $-OR^G$, $-CN$ or halo; and which 5-11- membered saturated monocyclic or bicyclic heterocyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, wherein any of such
15 C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl may be optionally substituted by a group C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl;

R^G is hydrogen, $-CH_3$ or $-C_2H_5$;

- R^{18} is selected in the group consisting of: lone electron pair, hydrogen, aryl, heteroaryl, $-(C_1-C_6\text{alkyl})$, $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$,
20 $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$ and $(C_5-C_7\text{heterocycloalkyl})-(C_3-C_6\text{cycloalkyl})$, wherein any of such aryl, heteroaryl, $-(C_1-C_6\text{alkyl})$, $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$ and $(C_5-C_7\text{heterocycloalkyl})-(C_3-C_6\text{cycloalkyl})$ may be optionally substituted by a group $-CN$, $-OH$, halo, $-COOR^M$, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $-O-(C_1$ -
25 C_6 alkyl), $-O-(C_3-C_6\text{cycloalkyl})$, $-S-(C_1-C_6\text{alkyl})$, $-S-(C_3-C_6\text{cycloalkyl})$, $-NR^H R^J$, $-N(R^L)(C_2-C_6\text{alkylene})-NR^H R^J$, $-N(R^L)(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-(C_1-C_6\text{alkylene})-NR^H R^J$, $-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-O-(C_2-C_6\text{alkylene})-NR^H R^J$, $-O-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-S-(C_2-C_6\text{alkylene})-NR^H R^J$, $-S-(C_3-$

C_7 cycloalkylene)- $NR^H R^J$, $-N(R^L)C(O)-(C_1-C_6$ alkylene)- $NR^H R^J$, $-N(R^L)C(O)-(C_3-C_7$ cycloalkylene)- $NR^H R^J$, $-C(O)N(R^L)-(C_2-C_6$ alkylene)- $NR^H R^J$, $-C(O)N(R^L)-(C_3-C_7$ cycloalkylene)- $NR^H R^J$, $-C(O)N(R^L)-(C_2-C_6$ alkylene)- OR^M , $-C(O)N(R^L)-(C_3-C_7$ cycloalkylene)- OR^M , $-N(R^L)C(O)N(R^H R^J)$, $-C(O)N(R^H R^J)$,
 5 $-N(R^L)C(O)N(R^L)-(C_2-C_6$ alkylene)- $NR^H R^J$, $-N(R^L)C(O)N(R^L)-(C_3-C_7$ cycloalkylene)- $NR^H R^J$, $-O-(C_2-C_6$ alkylene)- OR^M , $-O-(C_3-C_7$ cycloalkylene)- OR^M , $-S-(C_2-C_6$ alkylene)- OR^M , $-S-(C_3-C_7$ cycloalkylene)- OR^M , $-N(R^L)S(O)_2-(C_1-C_6$ alkylene)- $NR^H R^J$, $-N(R^L)S(O)_2-(C_3-C_7$ cycloalkylene)- $NR^H R^J$, $-S(O)_2N(R^L)-(C_2-C_6$ alkylene)- $NR^H R^J$, $-S(O)_2N(R^L)-(C_3-C_7$ cycloalkylene)- $NR^H R^J$,
 10 $-S(O)_2N(R^L)-(C_2-C_6$ alkylene)- OR^M , $-S(O)_2N(R^L)-(C_3-C_7$ cycloalkylene)- OR^M , $-N(R^L)S(O)_2-(C_2-C_6$ alkylene)- OR^M , $-N(R^L)S(O)_2-(C_3-C_7$ cycloalkylene)- OR^M , $-S(O)_2N(R^H R^J)$, $-N(R^L)S(O)_2R^L$, $-N(R^L)C(O)R^L$, OR^L , SR^L , $-(C_3-C_7$ heterocycloalkyl), $(C_5-C_7$ heterocycloalkyl)- $(C_1-C_6$ alkyl) and $(C_5-C_7$ heterocycloalkyl)- $(C_3-C_6$ cycloalkyl), wherein any of such C_1-C_6 alkyl, C_3-C_6 cycloalkyl, $-(C_1-C_6$ alkylene)-, $-(C_2-C_6$ alkylene)-, $-(C_3-C_7$ cycloalkylene)-, $-(C_3-C_7$ heterocycloalkyl), $(C_5-C_7$ heterocycloalkyl)- $(C_1-C_6$ alkyl) and $(C_5-C_7$ heterocycloalkyl)- $(C_3-C_6$ cycloalkyl) portion in the above listed groups may be optionally substituted by a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, $-OR^L$ or halo;

R^H and R^J , are at each occurrence independently hydrogen, C_1-C_6 alkyl or C_3-C_6 cycloalkyl, such C_1-C_6 alkyl or C_3-C_6 cycloalkyl being optionally substituted by a group C_1-C_3 alkyl, $-OR^M$, CN or halo;

alternatively, R^H and R^J may also together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more groups $-OR^M$, $-CN$, halo, C_1-C_6 alkyl or C_3-C_7 cycloalkyl, such C_1-C_6 alkyl and C_3-C_7 cycloalkyl being optionally substituted by a group C_1-C_3 alkyl, C_3-C_7 cycloalkyl, $-OR^M$, CN or halo; and which 5-11- membered saturated monocyclic or bicyclic heterocyclic ring optionally contains a further

heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl, wherein any of such C₁-C₆ alkyl or C₃-C₆ cycloalkyl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, -OR^M, CN, or halo;

5 and/or R^H and R^J may be linked to one carbon atom of the -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)- or -(C₃-C₇cycloalkylene)- portion of the group linked to the nitrogen to which they are connected to form a saturated cycle of up to 6 ring atoms;


R^L is at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl or C₃-C₆ cycloalkyl being optionally substituted
10 by a group C₁-C₃ alkyl, -OR^M, -CN or halo;

R^M is at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl or C₃-C₆ cycloalkyl being optionally substituted by a group hydroxyl, -CN or halo;

15 z¹, z², z³, and z⁴ are independently selected in the group consisting of: C, N, S, O, a group -CH-, and a group -NH-, in such a combination that the resulting ring formed is an aromatic system;

R¹⁹ is selected from the group consisting of: hydrogen, -CF₃, -NR^ER^F, -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), aryl or heteroaryl wherein any
20 of such -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), aryl or heteroaryl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, or halo; or

R¹⁹ is a group of general formula (V)



25 wherein R²⁰, R²¹, R^E and R^F are as above defined;

T is -N= or -CR²³=;


R^{23} is H, halo, $-CH_3$, or $-CN$;

R^{22} is H, halo, $-CH_3$, or $-CN$;

q is 0, 1, 2 or 3;

with the proviso that when Y is a group $-O(CR^3R^4)_n-$, n is 1 and R^{10} is
 5 $-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$,
 $-N(R^C)C(O)N(R^A R^B)$, $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$,
 $-N(R^C)C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, or
 $-N(R^C)C(O)R^C$, then X_1 is nitrogen.

In one embodiment, there is provided a compound of formula (IP), or a
 10 pharmaceutically acceptable salt thereof:



wherein;

W is a heteroatom selected from N or O, wherein N is substituted with
 15 hydrogen or C_1-C_6 alkyl or C_3-C_5 cycloalkyl;


Y is selected in the group consisting of: a group $-S(O)_p-$ wherein p is 0, 1 or 2; a group $-O(CR^3R^4)_n-$; a group $-(CR^5R^6)_n-$; a group $-NR^7-$; a group $-OC(O)-$; a group $-OC(O)NH-$; and a group $-OC(O)O-$;

R^3 , R^4 , R^5 and R^6 are each independently hydrogen, fluorine or C_1-C_6
 20 alkyl, or, respectively, R^3 and R^4 , or R^5 and R^6 may form together with the carbon atom to which they are attached a saturated 3-6 membered carbocyclic monocyclic ring optionally substituted by a group C_1-C_6 alkyl, hydroxyl or halo.

n is 0, 1, 2 or 3

R^7 is hydrogen, C_1-C_6 alkyl, or C_3-C_7 cycloalkyl wherein such C_1-C_6
 25 alkyl or C_3-C_7 cycloalkyl are optionally substituted by a group C_1-C_3 alkyl, C_3-C_6 cycloalkyl, hydroxyl, cyano or halo;

R^1 is a group selected from (IIa) - (IIc);



R^8 and R^9 are each independently hydrogen or C_1 - C_6 alkyl, or R^8 and R^9 may form together with the nitrogen atom to which they are attached a 5-11-membered saturated monocyclic or a fused or spiro bicyclic ring system optionally containing a further heteroatom which is oxygen or nitrogen, said nitrogen atom being optionally substituted by C_1 - C_6 alkyl; wherein such C_1 - C_6 alkyl groups may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, hydroxyl or halo;

X^1 , X^2 , X^3 , X^4 and X^5 are each independently a carbon atom, a nitrogen atom, a group $-(CH)-$ or a group $-NH-$; such that each combination thereof forms an aromatic ring system;

R^{10} is selected from a group consisting of: Hydrogen, $-NR^A R^B$, $-N(R^C)(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-(C_1-C_6\text{alkylene})-NR^A R^B$, $-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-O-(C_2-C_6\text{alkylene})-NR^A R^B$, $-O-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S-(C_2-C_6\text{alkylene})-NR^A R^B$, $-S-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-OR^D$, $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)C(O)NR^A R^B$, $-C(O)NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-(C_2-C_6\text{alkylene})-OR^D$, $-(C_3-C_7\text{cycloalkylene})-OR^D$, $-O-(C_2-C_6\text{alkylene})-OR^D$, $-O-(C_3-C_7\text{cycloalkylene})-OR^D$, $-S-(C_2-C_6\text{alkylene})-OR^D$, $-S-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)S(O)_2-(C_1-C_6\text{alkylene})-NR^A R^B$,

$-N(R^C)S(O)_2-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-$
 $NR^A R^B$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_2-$
 $C_6\text{alkylene})-OR^D$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)S(O)_2-(C_2-$
 $C_6\text{alkylene})-OR^D$, $-N(R^C)S(O)_2-(C_3-C_7\text{cycloalkylene})-OR^D$, $-S(O)_2N(R^A R^B)$,
 5 $-N(R^C)S(O)_2R^D$, $-N(R^C)C(O)R^C$, OR^C , SR^C , $-(C_3-C_7\text{heterocycloalkyl})$,
 $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$,

$(C_5-C_7\text{heterocycloalkyl})(C_3-C_6\text{cycloalkyl})-$, and C_3-C_7
 heterocycloalkylcarbonyl, wherein any of such alkylene, cycloalkylene,
 heterocycloalkyl, heterocycloalkyl-(C_1-C_6 alkyl), heterocycloalkyl-(C_3-
 10 $C_6\text{cycloalkyl})$ and heterocycloalkylcarbonyl may be optionally substituted by
 a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, hydroxyl or halo.

R^{11} is linked to X^4 and is selected from a group consisting of:

Hydrogen; $-CN$; C_1-C_6 alkyl which is substituted by a group selected
 from $-CN$, $-OR^C$, $-SR^C$, halo; $C_3-C_6\text{cycloalkyl}$ which is substituted by a group
 15 selected from C_1-C_4 alkyl, $-CN$, $-OR^C$, $-SR^D$, halo; $-NR^A R^B$, $-N(R^C)(C_2-$
 $C_6\text{alkylene})-NR^A R^B$, $-N(R^C)(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-(C_1-C_6\text{alkylene})-$
 $NR^A R^B$, $-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-O-(C_2-C_6\text{alkylene})-NR^A R^B$, $-O-(C_3-$
 $C_7\text{cycloalkylene})-NR^A R^B$, $-S-(C_2-C_6\text{alkylene})-NR^A R^B$, $-S-(C_3-$
 $C_7\text{cycloalkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-$
 20 $(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$,
 $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-OR^D$,
 $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)C(O)N(R^A R^B)$, $-C(O)N(R^A R^B)$,
 $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_3-$
 $C_7\text{cycloalkylene})-NR^A R^B$, $-O-(C_2-C_6\text{alkylene})-OR^D$, $-O-(C_3-C_7\text{cycloalkylene})-$
 25 OR^D , $-S-(C_2-C_6\text{alkylene})-OR^D$, $-S-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)S(O)_2-$
 $(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)S(O)_2-(C_3-C_7\text{cycloalkylene})-NR^A R^B$,
 $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-$
 $NR^A R^B$, $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-OR^D$, $-S(O)_2N(R^C)-(C_3-$

C_7 cycloalkylene)-OR^D, $-N(R^C)S(O)_2-(C_2-C_6$ alkylene)-OR^D, $-N(R^C)S(O)_2$
 $(C_3-C_7$ cycloalkylene)-OR^D, $-S(O)_2N(R^A R^B)$, $-N(R^C)S(O)_2R^D$, $-N(R^C)C(O)R^C$,
OR^C, SR^C, $-(C_3-C_7$ heterocycloalkyl), $(C_5-C_7$ heterocycloalkyl)-(C₁-C₆alkyl),
 $(C_5-C_7$ heterocycloalkyl)(C₃-C₆cycloalkyl), C₃-C₇ heterocycloalkylcarbonyl,
5 wherein any of such alkylene, cycloalkylene, heterocycloalkyl,
heterocycloalkyl-(C₁-C₆alkyl), heterocycloalkyl-(C₃-C₆cycloalkyl) and
heterocycloalkylcarbonyl may be optionally substituted by a group C₁-C₆
alkyl, C₃-C₇ cycloalkyl, hydroxyl or halo; or

R¹¹ is linked to X⁴ and is phenyl or 5- or 6-membered monocyclic
10 heteroaryl, wherein such phenyl or 5- or 6-membered monocyclic heteroaryl is
substituted by a group selected in the list consisting of: C₁-C₆ alkyl which is
substituted by a group -CN; C₃-C₆ cycloalkyl which is substituted by a group
selected from: -CN, -OR^C, -SR^C or halo; $-N(R^C)(C_2-C_6$ alkylene)-NR^AR^B,
 $-N(R^C)(C_3-C_7$ cycloalkylene)-NR^AR^B, $-(C_1-C_6$ alkylene)-NR^AR^B, $-(C_3-$
15 C_7 cycloalkylene)-NR^AR^B, $-O-(C_3-C_7$ cycloalkylene)-NR^AR^B,
 $-S-(C_2-C_6$ alkylene)-NR^AR^B, $-S-(C_3-C_7$ cycloalkylene)-NR^AR^B, $-N(R^C)C(O)-$
 $(C_1-C_6$ alkylene)-NR^AR^B, $-N(R^C)C(O)-(C_3-C_7$ cycloalkylene)-NR^AR^B,
 $-C(O)N(R^C)-(C_2-C_6$ alkylene)-NR^AR^B, $-C(O)N(R^C)-(C_3-C_7$ cycloalkylene)-
NR^AR^B, $-C(O)N(R^C)-(C_2-C_6$ alkylene)-OR^D, $-C(O)N(R^C)-(C_3-$
20 C_7 cycloalkylene)-OR^D, $-N(R^C)C(O)N(R^C)-(C_2-C_6$ alkylene)-NR^AR^B,
 $-N(R^C)C(O)N(R^C)-(C_3-C_7$ cycloalkylene)-NR^AR^B, $-O-(C_3-C_7$ cycloalkylene)-
OR^D, $-S-(C_3-C_7$ cycloalkylene)-OR^D, $-N(R^C)S(O)_2-(C_1-C_6$ alkylene)-NR^AR^B,
 $-N(R^C)S(O)_2-(C_3-C_7$ cycloalkylene)-NR^AR^B, $-S(O)_2N(R^C)-(C_2-C_6$ alkylene)-
NR^AR^B, $-S(O)_2N(R^C)-(C_3-C_7$ cycloalkylene)-NR^AR^B, $-S(O)_2N(R^C)-(C_2-$
25 C_6 alkylene)-OR^D, $-S(O)_2N(R^C)-(C_3-C_7$ cycloalkylene)-OR^D, $-N(R^C)S(O)_2-(C_2-$
 C_6 alkylene)-OR^D, $-N(R^C)S(O)_2-(C_3-C_7$ cycloalkylene)-OR^D, $-N(R^C)S(O)_2R^D$,
 $-(C_3-C_7$ heterocycloalkyl), $(C_5-C_7$ heterocycloalkyl)-(C₁-C₆alkyl),
 $(C_5-C_7$ heterocycloalkyl)(C₃-C₆cycloalkyl), C₃-C₇heterocycloalkylcarbonyl,

wherein any of such alkylene, cycloalkylene, heterocycloalkyl, heterocycloalkyl-(C₁-C₆alkyl), heterocycloalkyl-(C₃-C₆cycloalkyl) and heterocycloalkylcarbonyl may be optionally substituted by a group C₁-C₆alkyl, C₃-C₇cycloalkyl, hydroxyl or halo;

- 5 **R^A** and **R^B** are at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^D, -CN or halo; alternatively, **R^A** and **R^B**, may form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic ring
- 10 system which is optionally substituted by one or more group OR^D, CN, halo, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, OR^D, CN or halo; and which 5-11- membered saturated monocyclic or bicyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said
- 15 nitrogen atom optionally substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl wherein any of such alkyl or cycloalkyl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, -OR^D, -CN, or halo; and/or **R^A** and **R^B** may be linked to one carbon atom of the alkylene or cycloalkylene portion of the group linked to the nitrogen to which they are connected to form a
- 20 saturated cycle of up to 6 ring atoms;

R^C is at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl and C₃-C₆ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, -OR^D, -CN or halo;

R^D is at each occurrence independently hydrogen, -CH₃ or -C₂H₅;


- 25 **R¹²** and **R¹³** are independently hydrogen, C₁-C₆ alkyl, or halogen;

A is a divalent cycloalkylene radical having 5, 6 or 7 ring atoms; said cycloalkylene ring being attached to W and Y, and fused to a phenyl ring or to a monocyclic heteroaryl ring having 5 or 6 ring atoms, such phenyl or


heteroaryl ring being optionally substituted by one or two groups R^{24} ;

R^{24} is at each occurrence independently selected from the group consisting of: C_1 - C_6 alkyl, halogen and cyano;


R^2 is a radical of formula (IIIa), (IIIb), (IIIc), or (IIId):




(IIIa)



(IIIb)



(IIIc)



(IIId)

5


wherein

R^{14} is selected in the group consisting of: -F, -CH₃, -C₂H₅, -CH₂OH, -CH₂OMe, -CF₂CF₃, -CH₂SCH₃, -SCH₃ and -SC₂H₅;

10 R^{15} and R^{16} are independently -CH₃ or -C₂H₅;

R^{17} is selected from the group consisting of: lone electron pair, hydrogen, -CF₃, -NR^ER^F, -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), aryl or heteroaryl wherein any of such cycloalkyl, heterocycloalkyl, aryl or heteroaryl may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, or
15 halo; or

R^{17} is a group of general formula (IV)



(IV)

wherein

20 R^{20} is selected in the group consisting of: -F, -CH₃, -C₂H₅, -CH₂OH, -CH₂OMe, -CF₂CF₃, -CH₂SCH₃, -SCH₃ and -SC₂H₅;

R^{21} is $-CH_3$ or $-C_2H_5$;

or;

R^{20} and R^{21} as defined above may form together with the carbon atom to which they are attached a saturated 3-7- membered monocyclic ring;

- 5 R^E and R^F are each independently C_1 - C_6 alkyl, optionally substituted by a group C_1 - C_3 alkyl, $-OR^G$, $-CN$ or halo; alternatively, R^E and R^F may form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic ring system which is optionally substituted by one or more groups $-OR^G$, $-CN$, halo, C_1 - C_6 alkyl or C_3 - C_7 cycloalkyl, such
- 10 C_1 - C_6 alkyl and C_3 - C_7 cycloalkyl being optionally substituted by a group C_1 - C_3 alkyl, C_3 - C_7 cycloalkyl, $-OR^G$, $-CN$ or halo; and which 5-11- membered saturated monocyclic or bicyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl wherein any of such alkyl or cycloalkyl may
- 15 be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl;

R^G is hydrogen, $-CH_3$ or $-C_2H_5$;

- R^{18} is selected in the group consisting of: lone electron pair, hydrogen, aryl, heteroaryl, $-(C_1-C_6\text{alkyl})$, $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$ or $(C_5-C_7\text{heterocycloalkyl})-(C_3-C_6$
- 20 $\text{cycloalkyl})$, wherein any of such aryl, heteroaryl, alkyl, cycloalkyl, heterocycloalkyl, heterocycloalkyl- $(C_1-C_6\text{ alkyl})$, or heterocycloalkyl- $(C_3-C_6\text{ cycloalkyl})$ may be optionally substituted by a group $-CN$, $-OH$, halo, $-COOR^M$, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $-O-(C_1-C_6\text{alkyl})$, $-O-(C_3-C_6\text{cycloalkyl})$, $-S-(C_1-C_6\text{alkyl})$, $-S-(C_3-C_6\text{cycloalkyl})$, $-NR^H R^J$, $-N(R^L)(C_2-C_6\text{alkylene})-NR^H R^J$, $-N(R^L)(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-(C_1-C_6\text{alkylene})-$
- 25 $NR^H R^J$, $-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-O-(C_2-C_6\text{alkylene})-NR^H R^J$, $-O-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-S-(C_2-C_6\text{alkylene})-NR^H R^J$, $-S-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-N(R^L)C(O)-(C_1-C_6\text{alkylene})-NR^H R^J$, $-N(R^L)C(O)-$

- (C₃-C₇cycloalkylene)-NR^HR^J, -C(O)N(R^L)-(C₂-C₆alkylene)-NR^HR^J,
 -C(O)N(R^L)-(C₃-C₇cycloalkylene)-NR^HR^J, -C(O)N(R^L)-(C₂-C₆alkylene)-OR^M,
 -C(O)N(R^L)-(C₃-C₇cycloalkylene)-OR^M, -N(R^L)C(O)N(R^HR^J), -C(O)N(R^HR^J),
 -N(R^L)C(O)N(R^L)-(C₂-C₆alkylene)-NR^HR^J, -N(R^L)C(O)N(R^L)-(C₃-
 5 C₇cycloalkylene)-NR^HR^J, -O-(C₂-C₆alkylene)-OR^M, -O-(C₃-C₇cycloalkylene)-
 OR^M, -S-(C₂-C₆alkylene)-OR^M, -S-(C₃-C₇cycloalkylene)-OR^M, -N(R^L)S(O)₂-
 (C₁-C₆alkylene)-NR^HR^J, -N(R^L)S(O)₂-(C₃-C₇cycloalkylene)-NR^HR^J,
 -S(O)₂N(R^L)-(C₂-C₆alkylene)-NR^HR^J, -S(O)₂N(R^L)-(C₃-C₇cycloalkylene)-
 NR^HR^J, -S(O)₂N(R^L)-(C₂-C₆alkylene)-OR^M, -S(O)₂N(R^L)-(C₃-
 10 C₇cycloalkylene)-OR^M, -N(R^L)S(O)₂-(C₂-C₆alkylene)-OR^M, -N(R^L)S(O)₂-(C₃-
 C₇cycloalkylene)-OR^M, -S(O)₂N(R^HR^J), -N(R^L)S(O)₂R^L, -N(R^L)C(O)R^L, OR^L,
 SR^L, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl), and (C₅-
 C₇ heterocycloalkyl)-(C₃-C₆ cycloalkyl), wherein any of such alkyl,
 cycloalkyl, alkylene, cycloalkylene, heterocycloalkyl, heterocycloalkyl-(C₁-C₆
 15 alkyl), and (heterocycloalkyl)-(C₃-C₆ cycloalkyl) may be optionally
 substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, OR^L or halo;

R^H and **R^J**, are at each occurrence independently hydrogen, C₁-C₆ alkyl
 or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl or C₃-C₆ cycloalkyl being optionally
 substituted by a group C₁-C₃ alkyl, OR^M, CN or halo; alternatively, R^H and R^J
 20 may also together with the nitrogen atom to which they are attached a 5-11-
 membered saturated monocyclic or bicyclic ring system which is optionally
 substituted by one or more group OR^M, CN, halo, C₁-C₆ alkyl or C₃-C₇
 cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted
 by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, OR^M, CN or halo; and which
 25 5-11- membered saturated monocyclic or bicyclic ring optionally contains a
 further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally
 substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl wherein any of such alkyl or
 cycloalkyl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇

cycloalkyl, $-\text{OR}^{\text{M}}$, $-\text{CN}$, or halo; and/or R^{H} and R^{J} may be linked to one carbon atom of the alkylene or cycloalkylene portion of the group linked to the nitrogen to which they are connected to form a saturated cycle of up to 6 ring atoms;


5 R^{L} is at each occurrence independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, such $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl being optionally substituted by a group $\text{C}_1\text{-C}_3$ alkyl, OR^{M} , CN or halo;

R^{M} is at each occurrence independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, such $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl being optionally substituted
10 by a group hydroxyl, CN or halo;

z^1 , z^2 , z^3 , and z^4 are independently selected in the group consisting of: C, N, S, O, a group $-\text{CH}-$, and a group $-\text{NH}-$, in such a combination that the resulting ring formed is an aromatic system;

R^{19} is selected from the group consisting of: hydrogen, $-\text{CF}_3$, $-\text{NR}^{\text{E}}\text{R}^{\text{F}}$,
15 $-(\text{C}_3\text{-C}_7\text{cycloalkyl})$, $-(\text{C}_3\text{-C}_7\text{heterocycloalkyl})$, aryl or heteroaryl wherein any of such cycloalkyl, heterocycloalkyl, aryl or heteroaryl may be optionally substituted by a group $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, or halo or;

R^{19} is a group of general formula (V)



20

wherein R^{20} , R^{21} , R^{E} and R^{F} are as above defined;

T is $-\text{N}=\text{}$ or $-\text{CR}^{23}=\text{}$;

R^{23} is H, halo, $-\text{CH}_3$, or $-\text{CN}$;

R^{22} is H, halo, $-\text{CH}_3$, or $-\text{CN}$;

25 q is 0, 1, 2 or 3;

with the proviso that when Y is a group $-\text{O}(\text{CR}^3\text{R}^4)_n-$, n is 1 and R^{10} is

$-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^A R^B)$, $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, or $-N(R^C)C(O)R^C$, then X_1 is nitrogen.

5 In another aspect, the invention includes pharmaceutical compositions comprising a compound of the invention, together with one or more pharmaceutically acceptable carriers and/or excipients. Particularly preferred are compositions adapted for inhalation for pulmonary administration.

In another aspect, the invention includes the use of a compound of the
 10 invention for the treatment of diseases or conditions which benefit from inhibition of p38 MAP kinase activity. The treatment of obstructive or inflammatory airways diseases is a preferred use. All forms of obstructive or inflammatory airways diseases are potentially treatable with the compounds of the present invention, in particular an obstructive or inflammatory airways
 15 disease that is a member selected from the group consisting of chronic eosinophilic pneumonia, asthma, COPD, COPD that includes chronic bronchitis, pulmonary emphysema or dyspnea associated or not associated with COPD, COPD that is characterized by irreversible, progressive airways obstruction, adult respiratory distress syndrome (ARDS), exacerbation of
 20 airways hyper-reactivity consequent to other drug therapy and airways disease that is associated with pulmonary hypertension, chronic inflammatory diseases including cystic fibrosis, bronchiectasis and pulmonary fibrosis (Idiopathic). Efficacy is anticipated when p38 kinase inhibitors are administered either locally to the lung (for example by inhalation and intranasal delivery) or via
 25 systemic routes (for example, oral, intravenous and subcutaneous delivery).

Terminology

As used herein, the terms "halogen" or "halo" include fluorine, chlorine, bromine and iodine atoms.

As used herein, the term "C_x-C_yalkyl" wherein x and y are integers, refers to a straight or branched chain alkyl radical having from x to y carbon atoms. Thus when x is 1 and y is 6, for example, the term includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

As used herein, the term "C_x-C_yhaloalkyl" refers to the above "C_x-C_yalkyl" group wherein one or more hydrogen atoms are replaced by one or more halogen atoms.

As used herein, the term "C_x-C_yhydroxyalkyl" refers to the above "C_x-C_yalkyl" group wherein one hydrogen atom is replaced by one hydroxyl group.


As used herein, the term "C_x-C_yalkylene" wherein x and y are integers, refers to a C_x-C_yalkyl radical having in total two unsatisfied valencies, such as a divalent methylene radical.

As used herein, the term "carbocyclic" refers to a mono-, bi- or tricyclic radical having up to 16 ring atoms, all of which are carbon, and includes aryl and cycloalkyl.

As used herein, the term "C_z-C_kcycloalkyl" wherein z and k are integers refers to a monocyclic saturated carbocyclic radical having from z to k carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. Comprised within the scope of the term "C_z-C_kcycloalkyl", are those radicals having two unsatisfied valencies on the same carbon atom which will link to any C_x-C_yalkyl, C_x-C_yalkylene C_z-C_kcycloalkyl C_z-C_kcycloalkylene, C_z-C_kheterocycloalkyl, C_z-C_kheterocycloalkylC_x-C_yalkyl, C_z-C_kheterocycloalkylC_z-C_kcycloalkyl or (C_z-C_k)heterocycloalkylcarbonyl group by replacement of two hydrogen atoms placed on the same carbon. In such circumstances, this radical forms a gem-disubstituted or spiro system together with the C_x-C_yalkyl, C_x-C_yalkylene C_z-

C_k cycloalkyl C_z - C_k cycloalkylene, C_z - C_k heterocycloalkyl, C_z - C_k heterocycloalkyl C_x - C_y alkyl, C_z - C_k heterocycloalkyl C_z - C_k cycloalkyl or (C_z - C_k)heterocycloalkylcarbonyl group it is linked to.

The term " C_z - C_k cycloalkylene radical" refers to a C_z - C_k cycloalkyl
5 radical having two unsatisfied valencies on different cycle carbon atoms as follows:



As used herein, the unqualified term "aryl" refers to a mono- or bi-
10 cyclic carbocyclic aromatic radical, and includes radicals having two monocyclic carbocyclic aromatic rings which are directly linked by a covalent bond. Illustrative of such radicals are phenyl, biphenyl and naphthyl.

As used herein, the unqualified term "heteroaryl" refers to a mono- or
bi-cyclic aromatic radical containing one or more heteroatoms selected from
15 S, N and O, and includes radicals having two such monocyclic rings, or one such monocyclic ring and one monocyclic aryl ring, which are fused through a common bond. Illustrative examples of 5,6-membered heteroaryl are: are thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, pyrazolyl, oxazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl,
20 pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl. Illustrative examples of 8,10-membered heteroaryl are: benzothienyl, benzofuryl, benzimidazolyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, benzotriazolyl, indolyl and indazolyl.

As used herein, the unqualified term "heterocyclyl" or "heterocyclic"
25 and relates to a saturated mono-, bi- or tri-cyclic non-aromatic radical

containing one or more heteroatoms selected from S, N and O. In the case of bicyclic heterocyclic systems, included within the scope of the term are fused, spiro and bridged bicyclic systems. In particular, the term “C_z-C_kheterocycloalkyl” refers to monocyclic (C_z-C_k)cycloalkyl groups, in which
5 at least one ring carbon atom is replaced by a heteroatom (e.g. N, NH, S or O). Examples of (C_z-C_k)heterocycloalkyl include pyrrolidinyl, thiazolidinyl, piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl.

By analogy, the term “C_z-C_kheterocycloalkylene”, refers to a divalent C_z-C_kheterocycloalkyl radical, wherein C_z-C_kheterocycloalkyl is as above
10 defined.

The term “C_z-C_kheterocycloalkylC_x-C_yalkyl” refers to the above “C_x-C_yalkyl” group wherein one or more hydrogen atoms are replaced by one or more “C_z-C_kheterocycloalkyl” groups. Comprised within the scope of the term “C_z-C_kheterocycloalkylC_x-C_yalkyl” are systems where two hydrogen
15 atoms linked to the same carbon atom in “C_x-C_yalkyl” group are replaced by one “C_z-C_kheterocycloalkyl” group. Such radical thus form a gem-disubstituted “C_z-C_kheterocycloalkylC_x-C_yalkyl” system, such as a 1,2-dimethyl-pyrrolidin-2-yl radical.

The term “C_z-C_kheterocycloalkylC_z-C_kcycloalkyl” refers to the above
20 “C_z-C_kcycloalkyl” group wherein one or more hydrogen atoms are replaced by one or more “C_z-C_kheterocycloalkyl” groups.

The expression “(C_z-C_k)cycloalkylcarbonyl” refers to (C_z-C_k)cycloalkyl-CO- groups wherein the group “(C_z-C_k)cycloalkyl” has the meaning above defined.

25 The expression “(C_z-C_k)heterocycloalkylcarbonyl” refers to (C_z-C_k)heterocycloalkyl-CO- groups wherein the group “(C_z-C_k)heterocycloalkyl” has the meaning above defined.

Compounds of the invention may exist in one or more geometrical,

optical, enantiomeric, diastereomeric and tautomeric forms, including but not limited to *cis*- and *trans*-forms, *E*- and *Z*-forms, *R*-, *S*- and *meso*-forms, keto-, and enol-forms. Unless otherwise stated a reference to a particular compound includes all such isomeric forms, including racemic and other mixtures thereof. Where appropriate such isomers can be separated from their mixtures by the application or adaptation of known methods (e.g. chromatographic techniques and recrystallisation techniques). Where appropriate such isomers may be prepared by the application of adaptation of known methods (e.g. asymmetric synthesis).

Throughout the specification the use of an asterisk “*” in the definition of a structural formula, indicates the point of attachment for the radical group to the rest of the molecule.

As used herein the term “salt” includes base addition, acid addition and ammonium salts. As briefly mentioned above compounds of the invention which are acidic can form salts, including pharmaceutically acceptable salts, with bases such as alkaline metal hydroxides, e.g. sodium and potassium hydroxides; alkaline earth metal hydroxides e.g. calcium, barium and magnesium hydroxides; with organic bases e.g. *N*-methyl-*D*-glucamine, choline tris(hydroxymethyl)amino-methane, *L*-arginine, *L*-lysine, *N*-ethyl piperidine, dibenzylamine and the like. Those compounds of the invention which are basic can form salts, including pharmaceutically acceptable salts with inorganic acids, e.g. with hydrohalic acids such as hydrochloric or hydrobromic acids, sulphuric acid, nitric acid or phosphoric acid and the like, and with organic acids e.g. with acetic, formic, trifluoroacetic, tartaric, succinic, fumaric, maleic, malic, salicylic, citric, methanesulphonic, p-toluenesulphonic, benzoic, benzenesulfonic, glutamic, lactic, and mandelic acids and the like. Those compounds (I) which have a basic nitrogen can also form quaternary ammonium salts with a pharmaceutically acceptable counter-

ion such as ammonium, chloride, bromide, acetate, formate, p-toluenesulfonate, succinate, hemi-succinate, naphthalene-bis sulfonate, methanesulfonate, trifluoroacetate, xinafoate, and the like. For a review on salts, see Handbook of Pharmaceutical Salts: Properties, Selection, and Use by
5 Stahl and Wermuth (Wiley-VCH, Weinheim, Germany, 2002).

It is expected that compounds of the invention may be prepared in the form of hydrates, and solvates. Any reference herein, including the claims herein, to “compounds with which the invention is concerned” or “compounds of the invention” or “the present compounds”, and the like, includes reference
10 to salts hydrates, and solvates of such compounds. The term ‘solvate’ is used herein to describe a molecular complex comprising the compound of the invention and a stoichiometric amount of one or more pharmaceutically acceptable solvent molecules, for example, ethanol. The term ‘hydrate’ is employed when said solvent is water.

15 Individual compounds of the invention may exist in several polymorphic forms and may be obtained in different crystal or co-crystal habits, and they are intended to be included within the meaning of the term “compounds of the invention”.

The compounds may also be administered in the form of prodrugs
20 thereof. Thus certain derivatives of the compounds which may be active in their own right or may have little or no pharmacological activity themselves can, when administered into or onto the body, be converted into compounds of the invention having the desired activity, for example, by hydrolytic cleavage. Such derivatives are referred to as ‘prodrugs’. Further information on the use
25 of prodrugs may be found in Pro-drugs as Novel Delivery Systems, Vol. 14, ACS Symposium Series (T. Higuchi and V.J. Stella) and Bioreversible Carriers in Drug Design, Pergamon Press, 1987 (ed. E. B. Roche, American Pharmaceutical Association; C.S. Larsen and J. Østergaard, Design and


application of prodrugs, In Textbook of Drug Design and Discovery, 3rd Edition, 2002, Taylor and Francis).

Prodrugs in accordance with the invention can, for example, be produced by replacing appropriate functionalities present in the compounds of formula (I) with certain moieties known to those skilled in the art as 'pro-moieties' as described, for example, in Design of Prodrugs by H. Bundgaard (Elsevier, 1985). Such examples could be a prodrug of a carboxyl group (such as $-\text{CO}-\text{O}-\text{CH}_2-\text{O}-\text{CO}-\text{tBu}$ as used in the pivampicillin prodrug of ampicillin), an amide ($-\text{CO}-\text{NH}-\text{CH}_2-\text{NAlk}_2$) or an amidine ($-\text{C}(=\text{N}-\text{O}-\text{CH}_3)-\text{NH}_2$).

Embodiments of the Invention

It is to be understood that all preferred groups or embodiments described herebelow for compounds of formula (I) may be combined among each other and apply as well to compounds of formula (Ia), (Ib), (Ic), (IA), (IB), (IC), (ID), (IE) or (IF) as below defined *mutatis mutandis*.


In one embodiment, compounds of formula (Ia) are provided, which are compounds of formula (I) as above defined wherein carbon stereogenic center on the cycloalkylene portion of ring A which is linked to group W and identified with number (1) herebelow, possess the absolute configuration herebelow represented:



(Ia)

In another embodiment, compounds of formula (Ib) are provided, which are compounds of formula (I) as above defined wherein carbon stereogenic center on the cycloalkylene portion of ring A which are linked to group W and


Y and identified, respectively, with numbers (1) and (2) herebelow, possess the absolute configuration herebelow represented:



(Ib)

5 In a further embodiment, compound of formula (Ic) are provided, which are compounds of formula (I) as above defined wherein carbon stereogenic center on the cycloalkylene portion of ring A which are linked to group W and Y and identified, respectively, with numbers (1) and (2) herebelow, possess the absolute configuration herebelow represented:

10



(Ic)

In one embodiment, W is NH or O. In a further embodiment, W is NH.

15 In one embodiment, Y is a group $-S(O)_p-$, a group $-O(CR^3R^4)_n-$, a group $-(CR^5R^6)_n-$, or a group $-NR^7-$; p is zero and n is 0, 1 or 2. In another embodiment, Y is a group $-S(O)_p-$ or a group $-O(CR^3R^4)_n$ or; p is zero and n is 0 or 1.

In a further embodiment, Y is a group $-O(CR^3R^4)_n-$ and n is 0.

20 In one embodiment, R^3 , R^4 , R^5 and R^6 are each independently hydrogen, fluorine or C_1 - C_6 alkyl. In another embodiment, R^3 , R^4 , R^5 and R^6 are hydrogen.


In one embodiment, R^7 is hydrogen, C_1 - C_6 alkyl, or C_3 - C_7 cycloalkyl.

In one embodiment, R^7 is hydrogen.


25 In one embodiment, A is a divalent cycloalkylene radical having 5 or 6 ring atoms; said cycloalkylene ring being attached to W and Y, and fused to a

phenyl ring or to a monocyclic heteroaryl ring having 5 or 6 ring atoms, such phenyl or heteroaryl ring being optionally substituted by one or two groups R^{24} .

In a further embodiment, A is group selected in the group consisting of:




In a still further embodiment, A is group:



In an additional embodiment, A is group:


10



15 In one embodiment, R^{24} is not present or, if present, is at each occurrence independently selected from the group consisting of: C_1 - C_2 alkyl, -F, -Cl and cyano; in a further embodiment, R^{24} is not present or, if present, is at each occurrence independently Methyl or -F. In a further embodiment, R^{24} is not present.


In one embodiment, R^1 is a group of formula (IIa):

33



(IIa)


In a further embodiment, R¹ is a group of formula (IIa):



(IIa)

- 5 and **R⁸** and **R⁹** form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or a fused or spiro bicyclic ring system optionally containing a further heteroatom which is oxygen or nitrogen, said nitrogen atom being optionally substituted by C₁-C₆ alkyl; wherein such C₁-C₆ alkyl groups may be optionally substituted by a group C₁-
 10 C₆ alkyl, C₃-C₆ cycloalkyl, hydroxyl or halo.

In an additional embodiment, R¹ is a group of formula (IIa):




(IIa)

- and **R⁸** and **R⁹** form together with the nitrogen atom to which they are attached a 5 to 7- membered saturated monocyclic ring system optionally
 15 containing a further heteroatom which is oxygen or nitrogen, said nitrogen atom being optionally substituted by C₁-C₆ alkyl. In a still further embodiment, such saturated monocyclic ring system is a morpholine ring.

In another embodiment, R¹ is a group of formula (IIb):

34




(IIb)

In one embodiment, X^1 is a group $-(CH)-$ or a nitrogen atom. In another embodiment, X^1 is a group $-(CH)-$.

In one embodiment, R^{10} is selected from a group consisting of: $-CN$,
 5 $-C(O)N(R^A R^B)$, and $-N(R^C)C(O)R^C$.

In one embodiment, R^{12} is hydrogen, C_1 - C_6 alkyl, or halogen.


In a further embodiment, R^1 is a group of formula (IIb):



(IIb)


Wherein X^1 is a group $-(CH)-$, R^{10} is selected from a group consisting
 10 of: $-CN$, $-C(O)N(R^A R^B)$, and $-N(R^C)C(O)R^C$; and R^{12} is hydrogen.

In a further embodiment, R^1 is a group of formula (IIc):




(IIc)

In one embodiment, the group (IIc) is a group of formula (IIca) or (IIcb)
 which is connected to the group Y through one of the carbons as below
 15 indicated:




(IIca)



(IIcb)

In another embodiment, the group (IIc) is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X₂



5 In one embodiment, X⁴ is a carbon atom.

In one embodiment, X⁵ is a nitrogen atom.


In another embodiment, X⁴ is a carbon atom, X⁵ is a nitrogen atom, X³ is a nitrogen atom and X² is nitrogen.

10 In another embodiment, X⁴ is a carbon atom, X⁵ is a nitrogen atom, X³ is a nitrogen atom and X² is group -CH-.

In another embodiment, X⁴ is a nitrogen atom, X⁵ is a group -CH- atom, X³ is a carbon atom and X² is group -CH-.

In one embodiment, R¹³ is hydrogen, C₁-C₆ alkyl, or halogen.

15 In a further embodiment, the group (IIc) is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X₂



And wherein X⁴ is a carbon atom, X⁵ is a nitrogen atom, X³ is a nitrogen atom and X² is a group -CH-, and R¹³ is hydrogen.

20 In one embodiment, R¹¹ is selected from a group consisting of: -NR^AR^B, (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), -(C₃-C₇heterocycloalkyl), wherein any

of such (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) or -(C₃-C₇heterocycloalkyl) may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, hydroxyl or halo.

In another embodiment, R¹¹ is selected from a group consisting of:
 5 -NR^AR^B, -(C₁-C₆alkylene)-NR^AR^B, (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl),
 -(C₃-C₇heterocycloalkyl), wherein any of such (C₅-C₇heterocycloalkyl)-(C₁-
 C₆alkyl) or -(C₃-C₇heterocycloalkyl) may be optionally substituted by one,
 two or three groups R²⁵ which are independently selected in the list consisting
 of: C₁-C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl,
 10 hydroxyl and halo.

In another embodiment, R¹¹ is Phenyl or 5- or 6-membered monocyclic
 heteroaryl which is substituted by a group selected from:
 (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), -(C₃-C₇heterocycloalkyl), wherein any
 of such (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) or -(C₃-C₇heterocycloalkyl) may
 15 be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, hydroxyl
 or halo.

In another embodiment, R¹¹ is Phenyl or 5- or 6-membered monocyclic
 heteroaryl which is substituted by (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) or
 -(C₃-C₇heterocycloalkyl), wherein any (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl),
 20 -(C₃-C₇heterocycloalkyl), may be optionally substituted by one, two or three
 groups R²⁵ which are independently selected in the list consisting of: C₁-C₆
 alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and
 halo.

In one embodiment, R²⁵ is one, two or three groups independently
 25 selected in the list consisting of: C₁-C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-
 C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and halo.


In one embodiment, R^A and R^B are at each occurrence independently
 hydrogen, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇

cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, OR^D, CN or halo.

In another embodiment, **R^A** and **R^B** form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more group -OR^D, CN, halo, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^D, -CN or halo.

In a still further embodiment, **R^A** and **R^B** form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more group -OR^D, -CN, halo, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^D, -CN or halo; and which 5-11- membered saturated monocyclic or bicyclic heterocyclic ring contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl, wherein any of such C₁-C₆ alkyl or C₃-C₆ cycloalkyl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, OR^D, CN, or halo.


In one embodiment, **R¹¹** is a group:



Wherein **R²⁵** is optionally present and represents one, two, or three substituents independently selected in the list consisting of: C₁-C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group **R¹¹** to the


rest of the molecule via X^4 .

In further embodiment, R^{11} is a group:



- 5 Wherein R^{25} represents one or two C_1 - C_6 alkyl substituents; and wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .


In one embodiment, R^{11} is a group:



10


- Wherein R^{25} is optionally present and represents one, two or three substituents independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R^{11} to the
- 15 rest of the molecule via X^4 .

In a further embodiment, R^{11} is a group:




- Wherein R^{25} represents one, two or three substituents independently
- 20 selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .

In a further embodiment, R^{11} is a group:




- Wherein R^{25} represents one or two C_1 - C_6 alkyl substituents; and
 5 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .

In a further embodiment, R^{11} is a group:




- 10 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .

In a further embodiment, R^{11} is a group:




- 15 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .

In a still further embodiment, R^{11} is a group:



- 20 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 .

In one embodiment, R^2 is a radical of formula (IIIa):




(IIIa)

In one embodiment, R^{14} is selected in the group consisting of: $-\text{CH}_3$, $-\text{CH}_2\text{OH}$, or $-\text{CH}_2\text{SCH}_3$; in another embodiment, R^{14} is $-\text{CH}_3$.

In one embodiment, R^{15} and R^{16} are independently $-\text{CH}_3$ or $-\text{C}_2\text{H}_5$; in
5 another embodiment, R^{15} and R^{16} are $-\text{CH}_3$.


In another embodiment, R^2 is a radical of formula (IIIa):



(IIIa)

R^{14} is $-\text{CH}_3$, and R^{15} and R^{16} are $-\text{CH}_3$.

10 In another embodiment, R^2 is a radical of formula (IIIb):




(IIIb)

In one embodiment, R^{17} is selected from the group consisting of: lone electron pair, hydrogen, $-\text{CF}_3$, $-\text{NR}^{\text{E}}\text{R}^{\text{F}}$, $-(\text{C}_3\text{-C}_6\text{cycloalkyl})$,

-(C₄-C₆heterocycloalkyl), aryl or heteroaryl wherein any of such -(C₃-C₆cycloalkyl), -(C₄-C₆heterocycloalkyl), aryl or heteroaryl may be optionally substituted by a group methyl, isopropyl or halo. In another embodiment, R¹⁷ is selected from the group consisting of: lone electron pair, hydrogen, -CF₃,
 5 morpholine, cyclohexyl, phenyl or pyridyl.

In another embodiment, R¹⁷ is a group of general formula (IV)



In one embodiment, R²⁰ is selected in the group consisting of: F, -CH₃;
 10 -CH₂OH, -CH₂OMe, -CH₂SCH₃; in another embodiment, R²⁰ is selected in the group consisting of: -CH₃; -CH₂OH, -CH₂OMe. In another embodiment, R²⁰ is -CH₃.

In one embodiment, R²¹ is -CH₃.

In another embodiment, R²⁰ and R²¹ as defined above may form
 15 together with the carbon atom to which they are attached a cyclohexane or cyclopropyl ring; in a further embodiment, R²⁰ and R²¹ as defined above may form together with the carbon atom to which they are attached a cyclopropyl ring.


In one embodiment, R¹⁸ is phenyl or heteroaryl which is optionally
 20 substituted by a group -CN, -OH, halo, -COOR^M, C₁-C₆alkyl, C₃-C₆cycloalkyl, -O-(C₁-C₆alkyl), -O-(C₃-C₆cycloalkyl), -S-(C₁-C₆alkyl), -S-(C₃-C₆cycloalkyl), -NR^HR^J, -N(R^L)(C₂-C₆alkylene)-NR^HR^J, -N(R^L)(C₃-C₇cycloalkylene)-NR^HR^J, -(C₁-C₆alkylene)-NR^HR^J, -(C₃-C₇cycloalkylene)-NR^HR^J, -O-(C₂-C₆alkylene)-NR^HR^J, -O-(C₃-C₇cycloalkylene)-NR^HR^J, -S-(C₂-C₆alkylene)-NR^HR^J, -S-(C₃-
 25 C₇cycloalkylene)-NR^HR^J, -N(R^L)C(O)-(C₁-C₆alkylene)-NR^HR^J, -N(R^L)C(O)-(C₃-C₇cycloalkylene)-NR^HR^J, -C(O)N(R^L)-(C₂-C₆alkylene)-NR^HR^J,

$-C(O)N(R^L)-(C_3-C_7\text{cycloalkylene})-NR^H R^J$, $-C(O)N(R^L)-(C_2-C_6\text{alkylene})-OR^M$,
 $-C(O)N(R^L)-(C_3-C_7\text{cycloalkylene})-OR^M$, $-N(R^L)C(O)N(R^H R^J)$, $-C(O)N(R^H R^J)$,
 $-N(R^L)C(O)N(R^L)-(C_2-C_6\text{alkylene})-NR^H R^J$, $-N(R^L)C(O)N(R^L)-(C_3-$
 $C_7\text{cycloalkylene})-NR^H R^J$, $-O-(C_2-C_6\text{alkylene})-OR^M$, $-O-(C_3-C_7\text{cycloalkylene})-$
5 OR^M , $-S-(C_2-C_6\text{alkylene})-OR^M$, $-S-(C_3-C_7\text{cycloalkylene})-OR^M$, $-N(R^L)S(O)_2-$
 $(C_1-C_6\text{alkylene})-NR^H R^J$, $-N(R^L)S(O)_2-(C_3-C_7\text{cycloalkylene})-NR^H R^J$,
 $-S(O)_2N(R^L)-(C_2-C_6\text{alkylene})-NR^H R^J$, $-S(O)_2N(R^L)-(C_3-C_7\text{cycloalkylene})-$
 $NR^H R^J$, $-S(O)_2N(R^L)-(C_2-C_6\text{alkylene})-OR^M$, $-S(O)_2N(R^L)-(C_3-$
 $C_7\text{cycloalkylene})-OR^M$, $-N(R^L)S(O)_2-(C_2-C_6\text{alkylene})-OR^M$, $-N(R^L)S(O)_2-(C_3-$
10 $C_7\text{cycloalkylene})-OR^M$, $-S(O)_2N(R^H R^J)$, $-N(R^L)S(O)_2R^L$, $-N(R^L)C(O)R^L$, OR^L ,
 SR^L , $-(C_3-C_7\text{heterocycloalkyl})$, $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{ alkyl})$, and
 $(C_5-C_7\text{ heterocycloalkyl})-(C_3-C_6\text{ cycloalkyl})$, wherein any of such alkyl,
cycloalkyl, alkylene, cycloalkylene, heterocycloalkyl, heterocycloalkyl- $(C_1-C_6$
alkyl), heterocycloalkyl- $(C_3-C_6\text{ cycloalkyl})$ and heterocycloalkylcarbonyl may
15 be optionally substituted by a group $C_1-C_6\text{ alkyl}$, $C_3-C_7\text{ cycloalkyl}$, $-OR^L$ or
halo. In a further embodiment, R^{18} is phenyl substituted by $-(C_1-C_6\text{alkyl})$.

In one embodiment, R^{18} is $-(C_1-C_6\text{alkyl})$ or $-(C_3-C_7\text{cycloalkyl})$.


In one embodiment, $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N; in another
 embodiment, $z^1 = O$, $z^2 = C$, z^3 and z^4 are N; in a further embodiment, $z^1 =$
 20 $-CH-$, z^2 and z^3 are N and z^4 is $-CH-$; in an additional embodiment, $z^1 = N$, z^2
 is C, z^3 is N and z^4 is O; in a still further embodiment, $z^1 = N$, z^2 is C, z^3 is O
 and z^4 is N.

In an additional embodiment, R^2 is a radical of formula (IIIb):



(IIIb) ;


Wherein $z^1 = -CH-$, $z^2 = C$, z^3 is oxygen, z^4 is N, R^{18} is a lone pair, and R^{17} is a group of general formula (IV)



5


Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$.

In a further embodiment, R^2 is a radical of formula (IIIb):




(IIIb) ;

10 Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$.

In another embodiment, R^2 is a radical of formula (IIIb):




(IIIb) ;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general
 5 formula (IV)




Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and wherein R^{18} is
 phenyl, which may be optionally substituted by a group $-CN$, $-OH$,
 10 $-COOR^M$, C_1-C_6 alkyl, $-N(R^L)(C_2-C_6$ alkylene) $-NR^H R^J$, $-(C_1-C_6$ alkylene) $-$
 $NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-OR^M$, $-S-(C_2-$
 C_6 alkylene) $-OR^M$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl), and $(C_5-C_7$
 heterocycloalkyl) $-(C_3-C_6$ cycloalkyl), wherein any of such C_1-C_6 alkyl, $-(C_1-$
 C_6 alkylene) $-$, $-(C_2-C_6$ alkylene) $-$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl) and
 15 $(C_5-C_7$ heterocycloalkyl) $-(C_3-C_6$ cycloalkyl) may be optionally substituted by a
 group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, OR^L or halo.

In a further embodiment, R^2 is a radical of formula (IIIb):



(IIIb) ;


Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)



5

Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is phenyl, which is substituted in the para position by a group C_1-C_6 alkyl.


In an additional embodiment, R^2 is a radical of formula (IIIb):



(IIIb) ;

10


Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)



Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is phenyl, which is substituted in the meta position by a group $-O-(C_2-C_6\text{alkylene})-NR^H R^J$ or $-(C_1-C_6\text{alkylene})-NR^H R^J$.


In another embodiment, R^2 is a radical of formula (IIIb):

5



(IIIb) ;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




10

Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is a 5 or 6-membered heteroaryl which is optionally substituted by $C_1-C_6\text{alkyl}$, $-N(R^L)(C_2-C_6\text{alkylene})-NR^H R^J$, or $-(C_1-C_6\text{alkylene})-NR^H R^J$, wherein any of such $C_1-C_6\text{alkyl}$, $-(C_1-C_6\text{alkylene})-$, $-(C_2-C_6\text{alkylene})-$, $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$ and $(C_5-C_7\text{heterocycloalkyl})-(C_3-C_6\text{cycloalkyl})$ may be optionally substituted by a group $C_1-C_6\text{alkyl}$, $C_3-C_7\text{cycloalkyl}$, OR^L or halo. In one embodiment, for such compounds, R^{18} is an imidazole ring which is optionally substituted by $C_1-C_6\text{alkyl}$, $-N(R^L)(C_2-C_6\text{alkylene})-NR^H R^J$, or $-(C_1-C_6\text{alkylene})-NR^H R^J$.


20

In another embodiment, R^2 is a radical of formula (IIIb):



(IIIb) ;


Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)



5

Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is a group - $(C_1-C_6\text{alkyl})$, optionally substituted by a group $-OH$ or $-NR^H R^J$, or a group $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$ which may be optionally substituted by a group C_1-C_6 alkyl.

10 In a further embodiment, R^2 is a radical of formula (IIIc):



(IIIc)

;


In one embodiment, R^{19} is selected from the group consisting of: hydrogen, $-CF_3$, $-NR^E R^F$, $-(C_3-C_6\text{cycloalkyl})$, $-(C_3-C_6\text{heterocycloalkyl})$, aryl or heteroaryl wherein any of such cycloalkyl, heterocycloalkyl, aryl or heteroaryl may be optionally substituted by a group C_1-C_2 alkyl, C_3-C_5 cycloalkyl, or halo. In another embodiment, R^{19} is selected from the group consisting of:

15

hydrogen, $-\text{CF}_3$, morpholine, cyclohexyl, phenyl or pyridyl wherein any of such morpholine, cyclohexyl, phenyl or pyridyl may be optionally substituted by a group methyl, $-\text{F}$ or $-\text{Cl}$.

In another embodiment, R^{19} is a group of general formula (V)

5




In one embodiment, T is $-\text{N}=\text{}$. In another embodiment, T is $-\text{CR}^{23}=\text{}$.

In one embodiment, R^{22} is H, F, $-\text{Cl}$, $-\text{CH}_3$, or $-\text{CN}$; in another embodiment, R^{22} is H or F.

10

In one embodiment, R^{23} is H, F, $-\text{Cl}$, $-\text{CH}_3$, or $-\text{CN}$; in another embodiment, R^{23} is $-\text{Cl}$.


In another embodiment, R^2 is a radical of formula (IIIc):



;

Wherein R^{19} is $-(\text{C}_3\text{-C}_6\text{heterocycloalkyl})$, which is optionally substituted by a group $\text{C}_1\text{-C}_2$ alkyl, $\text{C}_3\text{-C}_5$ cycloalkyl, or halo; wherein T is $-\text{CR}^{23}=\text{}$, R^{22} is H or F.


In a still further embodiment, R^2 is a radical of formula (IIIId):




In one embodiment, q is 0, 1 or 2; in another embodiment, q is 0 or 1. In

a further embodiment, q is zero.

In one embodiment, compounds of formula (IA) are provided wherein W is NH, Y is a group $-O(CR^3R^4)_n-$ and n is 0, A is group:




Wherein R^1 is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X^2




10 And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a nitrogen atom and X^2 is a group $-CH-$, and R^{13} is hydrogen;

Wherein R^{11} is a group:




15 Wherein R^{25} is optionally present and represents one, two or three substituents independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 ;

Wherein R^2 is a radical of formula (IIIb):



(IIIb) ;


Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)



5


Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and wherein R^{18} is phenyl, which may be optionally substituted by a group $-CN$, $-OH$, $-COOR^M$, C_1-C_6 alkyl, $-N(R^L)(C_2-C_6$ alkylene) $-NR^H R^J$, $-(C_1-C_6$ alkylene) $-NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-OR^M$, $-S-(C_2-C_6$ alkylene) $-OR^M$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl), and $(C_5-C_7$ heterocycloalkyl) $-(C_3-C_6$ cycloalkyl), wherein any of such C_1-C_6 alkyl, $-(C_1-C_6$ alkylene) $-$, $-(C_2-C_6$ alkylene) $-$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl) and $(C_5-C_7$ heterocycloalkyl) $-(C_3-C_6$ cycloalkyl) may be optionally substituted by a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, OR^L or halo.

15 In one embodiment, compounds of formula (IB) are provided wherein W is NH, Y is a group $-O(CR^3 R^4)_n-$ and n is 0, A is group:




Wherein R^1 is a group of formula (IIca) as above defined which is

connected to the group Y through the carbon adjacent to X²




And wherein X⁴ is a carbon atom, X⁵ is a nitrogen atom, X³ is a nitrogen atom and X² is a group -CH-, and R¹³ is hydrogen;

5 Wherein **R¹¹** is a group:




Wherein **R²⁵** is optionally present and represents one, two or three substituents independently selected in the list consisting of: C₁-C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and halo; and
 10 wherein the asterisk represents the point of attachment for group R¹¹ to the rest of the molecule via X⁴;

wherein R² is a radical of formula (IIIb):




(IIIb)

15 Wherein **z¹** = -CH-, **z²** = C, **z³** and **z⁴** are N and R¹⁷ is a group of general formula (IV)




Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and wherein R^{18} is phenyl, which may be optionally substituted by a group $-CN$, $-OH$, $-COOR^M$, C_1-C_6 alkyl, $-N(R^L)(C_2-C_6$ alkylene) $-NR^H R^J$, $-(C_1-C_6$ alkylene) $-NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-NR^H R^J$, $-O-(C_2-C_6$ alkylene) $-OR^M$, $-S-(C_2-C_6$ alkylene) $-OR^M$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl), and $(C_5-C_7$ heterocycloalkyl) $-(C_3-C_6$ cycloalkyl), wherein any of such wherein any of such C_1-C_6 alkyl, $-(C_1-C_6$ alkylene) $-$, $-(C_2-C_6$ alkylene) $-$, $(C_5-C_7$ heterocycloalkyl) $-(C_1-C_6$ alkyl) and $(C_5-C_7$ heterocycloalkyl) $-(C_3-C_6$ cycloalkyl) may be optionally substituted by a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, OR^L or halo.

In one embodiment, compounds of formula (IC) are provided wherein W is NH, Y is a group $-O(CR^3 R^4)_n-$ and n is 0, A is group:




Wherein R^1 is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X^2




And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a nitrogen atom and X^2 is a group $-CH-$, and R^{13} is hydrogen;

R^{11} is a group:




Wherein R^{25} is optionally present and represents one, two or three substituents independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and
 5 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 ;

R^2 is a radical of formula (IIIb):




(IIIb) ;

10 Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is a 5 or 6-
 15 membered heteroaryl, which is optionally substituted by C_1 - C_6 alkyl, $-N(R^L)(C_2$ - C_6 alkylene)- $NR^H R^J$, or $-(C_1$ - C_6 alkylene)- $NR^H R^J$, wherein any of such C_1 - C_6 alkyl, $-(C_1$ - C_6 alkylene)-, $-(C_2$ - C_6 alkylene)-, (C_5 - C_7 heterocycloalkyl)-(C_1 - C_6 alkyl) and (C_5 - C_7 heterocycloalkyl)-(C_3 - C_6 cycloalkyl) may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7
 20 cycloalkyl, OR^L or halo.

In one embodiment, compounds of formula (ID) are provided wherein W is NH, Y is a group $-O(CR^3R^4)_n-$ and n is 0, A is group:




- 5 R1 is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X₂




And wherein X⁴ is a carbon atom, X⁵ is a nitrogen atom, X³ is a nitrogen atom and X² is a group -CH-, and R¹³ is hydrogen;

- 10 Wherein R¹¹ is a group:




- Wherein R²⁵ is optionally present and represents one two or three substituents independently selected in the list consisting of: C₁-C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R¹¹ to the rest of the molecule via X⁴;

Wherein R² is a radical of formula (IIIb):



(IIIb) ;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




5

Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is a 5 or 6-membered heteroaryl, which is optionally substituted by C_1-C_6 alkyl, $-N(R^L)(C_2-C_6$ alkylene)- $NR^H R^J$, or $-(C_1-C_6$ alkylene)- $NR^H R^J$, wherein any of such C_1-C_6 alkyl, $-(C_1-C_6$ alkylene)-, $-(C_2-C_6$ alkylene)-, $(C_5-$

10 C_7 heterocycloalkyl)- $(C_1-C_6$ alkyl) and $(C_5-C_7$ heterocycloalkyl)- $(C_3-C_6$ cycloalkyl) may be optionally substituted by a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, OR^L or halo.


In one embodiment, compounds of formula (IE) are provided wherein W is NH, Y is a group $-O(CR^3 R^4)_n-$ and n is 0, A is group:

15



Wherein R^1 is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X_2

56




(IIca)

And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a nitrogen atom and X^2 is a group $-CH-$, and R^{13} is hydrogen;


wherein R^{11} is a group:

5



Wherein R^{25} is optionally present and represents one two or three substituents independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and
 10 wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 ;


Wherein R^2 is a radical of formula (IIIb):



(IIIb)


;

15 Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$ and R^{18} is a group -
 (C₁-C₆alkyl), optionally substituted by a group $-OH$ or $-NR^H R^J$, or a group
 (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) which may be optionally substituted by
 5 a group C₁-C₆ alkyl.

In one embodiment, compounds of formula (IF) are provided wherein
 W is NH, Y is a group $-O(CR^3 R^4)_n-$ and n is 0, A is group:




10 Wherein R^1 is a group of formula (IIca) as above defined which is
 connected to the group Y through the carbon adjacent to X_2



And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a
 nitrogen atom and X^2 is a group $-CH-$, and R^{13} is hydrogen;

15 Wherein R^{11} is a group:




Wherein R^{25} is optionally present and represents one two or three
 substituents independently selected in the list consisting of: C₁-C₆ alkyl, (C₁-

C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R¹¹ to the rest of the molecule via X⁴;


Wherein R² is a radical of formula (IIIb):

5



(IIIb) ;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R¹⁷ is a group of general formula (IV)



10

Wherein R²⁰ is -CH₃ or -CH₂OH, and R²¹ is -CH₃ and R¹⁸ is a group - (C₁-C₆alkyl), optionally substituted by a group -OH or -NR^HR^J, or a group (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) which may be optionally substituted by a group C₁-C₆ alkyl.

15 In one embodiment, a compound of formula (I) is selected in the list consisting of:

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(2-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

20

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[2-(4-methyl-piperazin-1-yl)-ethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-

tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

5 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-

10 tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((S)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

15 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((S)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-isopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

20 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(6-cyano-pyridin-3-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

N-(4-{(1S,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy}-pyridin-2-yl)-2-methoxy-acetamide;

25 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[1-(2-hydroxy-ethyl)-1H-indazol-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((R)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

N-(4-{{(1R,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy}-pyridin-2-yl)-2-methoxy-acetamide;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(4-methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-morpholin-4-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-pyrrolidin-1-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[6-(morpholine-4-carbonyl)-pyridin-3-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(4-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(3-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(2-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(6-morpholin-4-ylmethyl-pyridin-3-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(1-methyl-piperidin-4-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[1-(2,2-difluoro-ethyl)-piperidin-4-ylmethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxypiperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[(2-hydroxy-ethyl)-methyl-amino]-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

10 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

15 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

20 1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

25 3-[3-tert-Butyl-5-(3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido)-pyrazol-1-yl]-benzoic acid ethyl ester;

1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-

{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(2-morpholin-4-yl-ethoxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

5 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-(2-Hydroxy-1,1-dimethyl-ethyl)-2-p-tolyl-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,3S)-3-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-indan-1-yl}-urea;

20 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

25 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
5 {[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[(2-dimethylamino-ethyl)-methyl-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

10 1-[5-tert-Butyl-2-(3-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[methyl-(2-morpholin-4-yl-ethyl)-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-
15 tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{[(1S,4R)-4-[3-((R)-1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{[(1S,4R)-4-[3-((S)-1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
20 tetrahydro-naphthalen-1-yl]}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{[(1S,4R)-4-[3-((S)-1,2-dimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]}-urea;

25 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxymethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-

[1,4]oxazepan-4-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-{3-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(2-tert-Butyl-5-p-tolyl-3H-imidazol-4-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(4-hydroxy-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(4-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4S)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(3-hydroxymethyl-4-methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-(4-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-1-isopropyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-dimethylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

5 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-ethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

10 tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

15 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-3-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

20 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

25 tetrahydro-naphthalen-1-yl}-urea;

1-[(1S,4R)-4-(3-Azepan-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-piperazine-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl} urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-
5 [1,4]diazepan-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl} urea;

1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

10 1-[5-tert-Butyl-2-(4-cyano-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4S)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
15 tetrahydro-naphthalen-1-yl} -urea;

1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl} -urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-pyrrolidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

25 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1-dimethylamino-1-methyl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl} -urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-3-

hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxy-4-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1-methyl-1-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-4-methyl-morpholin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[(S)-1-(3-hydroxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-hydroxymethyl-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1,4-dimethyl-piperazin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1,4,4-trimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

5 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1-methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-1-methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

10 tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-1-methyl-pyrrolidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

15 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((2S,4R)-4-fluoro-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-3-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

20 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-3-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-

25 tetrahydro-naphthalen-1-yl)-urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl)-urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-(cis-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 {[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]}-urea;

1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-
10 urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1-dimethylamino-cyclopentyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[(1S,4R)-4-(3-Amino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea;

1-[5-tert-Butyl-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

25 N-(5-tert-Butyl-2-methoxy-3-{3-[(1S,4R)-4-(3-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-phenyl)-methanesulfonamide;

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[8-methyl-3-

((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-(3-tert-Butyl-1'-methyl-1'H-[1,4']bipyrazolyl-5-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

N-[5-tert-Butyl-2-methoxy-3-(3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-phenyl]-methanesulfonamide;

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-isoxazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[3-tert-Butyl-1'-(3-dimethylamino-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[3-tert-Butyl-1'-(3-morpholin-4-yl-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-

1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-{(1S,4R)-4-[3-((2S,6R)-2,6-Dimethyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-
(3-fluoro-5-morpholin-4-yl-phenyl)-urea;

5 1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-Cyclopropyl-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
10 urea;

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-
((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-
1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[1-(2-dimethylamino-ethyl)-1H-imidazol-4-yl]-2H-
15 pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea;

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-
((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
tetrahydro-naphthalen-1-yl}-urea;

20 1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{(1S,4R)-4-[3-(8-Aza-bicyclo[3.2.1]oct-8-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-
25 (2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-
3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-
5 3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-
methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
tetrahydronaphthalen-1-yl]-urea;

10 1-[5-tert-Butyl-2-(2-dimethylamino-ethyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(2-piperidin-1-yl-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-
4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-
15 1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[2-(4-methyl-piperazin-1-yl)-ethyl]-2H-pyrazol-3-yl}-
3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
20 {(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-morpholin-4-yl-ethyl)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

25 1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethyl)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethyl]-phenyl}-2H-

pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethyl)-phenyl]-2H-pyrazol-3-
5 yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-piperidin-1-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-
3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

10 1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethyl]-phenyl}-2H-
pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-{5-tert-Butyl-2-[3-(2-[1,4]oxazepan-4-yl-ethyl)-phenyl]-2H-pyrazol-
15 3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-{3-[2-(4-methyl-[1,4]diazepan-1-yl)-ethyl]-phenyl}-
2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
20 urea;

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-
((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-{2-[(2-dimethylamino-ethyl)-methyl-amino]-
25 pyrimidin-4-yl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-
yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-

yl}-3-((1S,4R)-4-[3-((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-
 5 ((1S,4R)-4-[3-((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

10 1-[5-tert-Butyl-2-(3-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

15 1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

20 1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-{(1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea;

25 1-[5-tert-Butyl-2-(3-morpholin-4-yl-methyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-

6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-{3-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-
2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
5 urea;

1-{5-tert-Butyl-2-[3-((R)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-
2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

10 1-{5-tert-Butyl-2-[3-((S)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-
2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-[2-[3-(2-Dimethylamino-ethoxy)-phenyl]-5-(2-hydroxy-1,1-dimethyl-
15 ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
20 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-diethylamino-ethoxy)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-
25 yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-
yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-

a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-{3-[2-(4-fluoropiperidin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

10 1-(5-tert-Butyl-2-{3-[2-(4-methyl-[1,4]-diazepan-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

15 1-{5-tert-Butyl-2-[3-(2-[1,4]oxazepan-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

20 1-(2-{3-[2-(8-Aza-bicyclo[3.2.1]oct-8-yl)-ethoxy]-phenyl}-5-tert-butyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

25 1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-(3-{2-[(2-methoxy-ethyl)-methyl-amino]-ethoxy}-phenyl)-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-

urea;

1-(5-tert-Butyl-2-{3-[2-(4-methoxy-piperidin-1-yl-ethoxy)]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-

5 urea;

1-(5-tert-Butyl-2-{3-[2-(3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

10 1-{5-tert-Butyl-2-[3-(4-methoxy-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

15 1-[5-tert-Butyl-2-(3-[(2-methoxy-ethyl)-methyl-amino]-methyl)-phenyl]-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-[5-tert-Butyl-2-[3-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-[5-tert-Butyl-2-(3-dimethylaminomethyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

25 1-[5-tert-Butyl-2-(3-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea;

1-((1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-

a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea;

1-[5-tert-Butyl-2-(3-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 {((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[3-(4-methyl-[1,4]diazepan-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-
pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

10 1-{5-tert-Butyl-2-[3-((1S,4S)-5-methyl-2,5-diaza-bicyclo[2.2.1]hept-2-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(4-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
15 {((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(4-dimethylaminomethyl-phenyl)-2H-pyrazol-3-yl]-3-
{((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

20 1-{5-tert-Butyl-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[4-(4-methyl-[1,4]diazepan-1-ylmethyl)-phenyl]-2H-
25 pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-{5-tert-Butyl-2-[4-(4-methoxy-piperidin-1-ylmethyl)-phenyl]-2H-

pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-[5-tert-Butyl-2-(4-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 {((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

1-[5-tert-Butyl-2-(4-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
{((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea;

10 1-{5-tert-Butyl-2-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-2H-
pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-(5-tert-Butyl-2-{4-[(ethyl-methyl-amino)-methyl]-phenyl}-2H-
15 pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;


1-[5-tert-Butyl-2-(4-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-
phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
20 [1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea;

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-
yl}-3-((1S,4S)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-
6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea;

25 And pharmaceutically acceptable salts thereof.


According to another aspect of the invention there is provided a
compound of formula (J), or a pharmaceutically acceptable salt thereof:

80



Wherein R^2 , W, A and Y have the same meaning as provided above for compounds of formula (I); and wherein

R^1 is a group (IIc) wherein R^{13} , X^2 , X^3 , X^4 and X^5 have the same
5 meaning as provided above for compounds of formula (I);



And wherein R^{11} is phenyl or 5- or 6-membered monocyclic heteroaryl, wherein such phenyl or 5- or 6-membered monocyclic heteroaryl is substituted by a group $-O-(C_2-C_6\text{alkylene})-NR^R R^S$, wherein such $C_2-C_6\text{alkylene}$ may be
10 optionally substituted by a group $C_1-C_6\text{alkyl}$, $C_3-C_7\text{cycloalkyl}$, hydroxyl or halo;

R^R and R^S are at each occurrence independently $C_1-C_6\text{ alkyl}$ or $C_3-C_7\text{ cycloalkyl}$, such $C_1-C_6\text{ alkyl}$ and $C_3-C_7\text{ cycloalkyl}$ being substituted by one or more group OR^T , CN or halo; alternatively, R^R and R^S , may form together with
15 the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic ring system which is substituted by one or more group OR^T , CN, halo, $C_1-C_6\text{ alkyl}$ or $C_3-C_7\text{ cycloalkyl}$, such $C_1-C_6\text{ alkyl}$ and $C_3-C_7\text{ cycloalkyl}$ being optionally substituted by a group $C_1-C_3\text{ alkyl}$, $C_3-C_7\text{cycloalkyl}$, OR^D , CN or halo; and which 5-11- membered saturated
20 monocyclic or bicyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by $C_1-C_6\text{ alkyl}$ or $C_3-C_6\text{ cycloalkyl}$ wherein any of such alkyl or cycloalkyl may be optionally substituted by a group $C_1-C_6\text{ alkyl}$, $C_3-C_7\text{ cycloalkyl}$, OR^D , CN, or halo; and

R^T is at each occurrence independently hydrogen, -CH₃ or -C₂H₅.

Where applicable, all preferred groups or embodiments described herebelow for compounds of formula (I), (Ia), (Ib) and (Ic) may be combined among each other and apply as well to compounds of formula (J) as above defined *mutatis mutandis*.

Utility

As mentioned above the compounds of the invention are p38MAPK inhibitors, and thus may have utility for the treatment of diseases or conditions which benefit from inhibition of the p38 enzyme. Such diseases and conditions are known from the literature and several have been mentioned above. However, the compounds are generally of use as anti-inflammatory agents, particularly for use in the treatment of respiratory disease. In particular, the compounds may be used in the treatment of chronic obstructive pulmonary disease (COPD), chronic bronchitis, lung fibrosis, pneumonia, acute respiratory distress syndrome (ARDS), pulmonary emphysema, or smoking-induced emphysema, intrinsic (non-allergic asthma and extrinsic (allergic) asthma, mild asthma, moderate asthma, severe asthma, steroid resistant asthma, neutrophilic asthma, bronchitic asthma, exercise induced asthma, occupational asthma and asthma induced following bacterial infection, cystic fibrosis, pulmonary fibrosis and bronchiectasis.

The present invention provides the use of the compounds of the invention for the prevention and/or treatment of any disease or condition which benefit from inhibition of the p38 enzyme.

In a further aspect the present invention provides the use of compounds of the invention for the preparation of a medicament for the prevention and/or treatment of any disease or condition which benefit from inhibition of the p38 enzyme.

Moreover the present invention provides a method for prevention and/or

treatment of any disease which benefit from inhibition of the p38 enzyme, said method comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of the invention.

Compositions

5 As mentioned above, the compounds with which the invention is concerned are p38 kinase inhibitors, and are useful in the treatment of several diseases for example inflammatory diseases of the respiratory tract. Examples of such diseases are referred to above, and include asthma, rhinitis, allergic airway syndrome, bronchitis and chronic obstructive pulmonary disease.

10 It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing treatment.

15 Optimum dose levels and frequency of dosing will be determined by clinical trial, as is required in the pharmaceutical art. In general, the daily dose range for oral administration will lie within the range of from about 0.001 mg to about 100 mg per kg body weight of a human, often 0.01 mg to about 50 mg per kg, for example 0.1 to 10 mg per kg, in single or divided doses. In general,

20 the daily dose range for inhaled administration will lie within the range of from about 0.1 µg to about 1mg per kg body weight of a human, preferably 0.1 µg to 50 µg per kg, in single or divided doses. On the other hand, it may be necessary to use dosages outside these limits in some cases. For the purpose of the invention, inhaled administration is preferred.

25 The compounds with which the invention is concerned may be prepared for administration by any route consistent with their pharmacokinetic properties. Orally administrable compositions may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral,

topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example
5 lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or acceptable wetting agents such as sodium lauryl sulfate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in
10 the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible
15 fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

20 For topical application to the skin, the drug may be made up into a cream, lotion or ointment. Cream or ointment formulations which may be used for the drug are conventional formulations well known in the art, for example as described in standard textbooks of pharmaceuticals such as the British Pharmacopoeia.

25 The active ingredient may also be administered parenterally in a sterile medium. Depending on the vehicle and concentration used, the drug can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as a local anaesthetic, preservative and buffering agents can be dissolved in the

vehicle.

However, for treatment of an inflammatory disease of the respiratory tract, compounds of the invention may also be formulated for inhalation, for example as a nasal spray, or dry powder or aerosol inhalers. For delivery by
5 inhalation, the active compound is preferably in the form of microparticles. They may be prepared by a variety of techniques, including spray-drying, freeze-drying and micronisation. Aerosol generation can be carried out using, for example, pressure-driven jet atomizers or ultrasonic atomizers, preferably using propellant-driven metered aerosols or propellant-free administration of
10 micronized active compounds from, for example, inhalation capsules or other “dry powder” delivery systems.

By way of example, a composition of the invention may be prepared as a suspension for delivery from a nebuliser or as an aerosol in a liquid propellant, for example for use in a pressurised metered dose inhaler (PMDI).
15 Propellants suitable for use in a PMDI are known to the skilled person, and include CFC-12, HFA-134a, HFA-227, HCFC-22 (CCl_2F_2) and HFA-152 (CH_3F and isobutane).

In a preferred embodiment of the invention, a composition of the invention is in dry powder form, for delivery using a dry powder inhaler
20 (DPI). Many types of DPI are known.

Microparticles for delivery by administration may be formulated with excipients that aid delivery and release. For example, in a dry powder formulation, microparticles may be formulated with large carrier particles that aid flow from the DPI into the lung. Suitable carrier particles are known, and
25 include lactose particles; they may have a mass median aerodynamic diameter of greater than $90\text{ }\mu\text{m}$.

In the case of an aerosol-based formulation, an example is:

Compound of the invention 24 mg / canister

Lecithin, NF Liq. Conc. 1.2 mg / canister
Trichlorofluoromethane, NF 4.025 g / canister
Dichlorodifluoromethane, NF 12.15 g / canister.

The active compounds may be dosed as described depending on the
5 inhaler system used. In addition to the active compounds, the administration
forms may additionally contain excipients, such as, for example, propellants
(e.g. Frigen in the case of metered aerosols), surface-active substances,
emulsifiers, stabilizers, preservatives, flavorings, fillers (e.g. lactose in the
case of powder inhalers) or, if appropriate, further active compounds.

10 For the purposes of inhalation, a large number of systems are available
with which aerosols of optimum particle size can be generated and
administered, using an inhalation technique which is appropriate for the
patient. In addition to the use of adaptors (spacers, expanders) and pear-
shaped containers (e.g. Nebulator®, Volumatic®), and automatic devices
15 emitting a puffer spray (Autohaler®), for metered aerosols, in particular in the
case of powder inhalers, a number of technical solutions are available (e.g.
Diskhaler®, Rotadisk®, Turbohaler® or the inhalers for example as described
EP-A-0505321). Additionally, compounds of the invention may be delivered
in multi-chamber devices thus allowing for delivery of combination agents.

20 *Combinations*

Other compounds may be combined with compounds with which the
invention is concerned for the prevention and treatment of inflammatory
diseases, in particular respiratory diseases. Thus the present invention is also
concerned with pharmaceutical compositions comprising a therapeutically
25 effective amount of a compound of the invention and one or more other
therapeutic agents. Suitable therapeutic agents for a combination therapy with
compounds of the invention include, but are not limited to: (1) corticosteroids,
such as fluticasone propionate, fluticasone furoate, mometasone furoate,

beclometasone dipropionate, ciclesonide, budesonide, GSK 685698, GSK 870086, QAE 397, QMF 149, TPI-1020; (2) β 2-adrenoreceptor agonists such as salbutamol, albuterol, terbutaline, fenoterol, and long acting β 2-adrenoreceptor agonists such as salmeterol, indacaterol, formoterol (including formoterol fumarate), arformoterol, carmoterol, GSK 642444, GSK 159797, GSK 159802, GSK 597501, GSK 678007, AZD3199; (3) corticosteroid/long acting β 2 agonist combination products such as salmeterol/fluticasone propionate (Advair/Seretide), formoterol/budesonide (Symbicort), formoterol/fluticasone propionate (Flutiform), formoterol/ciclesonide, formoterol/mometasone furoate, formoterol/ beclometasone dipropionate, indacaterol/mometasone furoate, Indacaterol/QAE 397, GSK 159797/GSK 685698, GSK 159802/GSK 685698, GSK 642444/GSK 685698, GSK 159797/GSK 870086, GSK 159802/GSK 870086, GSK 642444/GSK 870086, arformoterol/ciclesonide;(4) anticholinergic agents, for example muscarinic-3 (M3) receptor antagonists such as ipratropium bromide, tiotropium bromide, Aclidinium (LAS-34273), NVA-237, GSK 233705, Darotropium, GSK 573719, GSK 961081, QAT 370, QAX 028, EP-101; (5) dual pharmacology M3-anticholinergic/ β 2-adrenoreceptor agonists such as GSK961081, AZD2115 and LAS190792; (6) leukotriene modulators, for example leukotriene antagonists such as montelukast, zafirulast or pranlukast or leukotriene biosynthesis inhibitors such as Zileuton or BAY-1005, or LTB4 antagonists such as Amelubant, or FLAP inhibitors such as GSK 2190914, AM-103; (7) phosphodiesterase-IV (PDE-IV) inhibitors (oral or inhaled), such as roflumilast, cilomilast, Oglemilast, ONO-6126, Tetomilast, Tofimilast, UK 500,001, GSK 256066; (8) antihistamines, for example selective histamine-1 (H1) receptor antagonists, such as fexofenadine, cetirizine, loratidine or astemizole or dual H1/H3 receptor antagonists such as GSK 835726, GSK 1004723, or selective histamine-4 (H4) receptor antagonists, such as

ZPL3893787; (9) antitussive agents, such as codeine or dexamorphan; (10) a mucolytic, for example N acetyl cysteine or fudostein; (11) a expectorant/mucokinetic modulator, for example ambroxol, hypertonic solutions (e.g. saline or mannitol) or surfactant; (12) a peptide mucolytic, for example recombinant human deoxyribonoclease I (dornase-alfa and rhDNase) or helicidin; (13) antibiotics, for example azithromycin, tobramycin and aztreonam; (14) non-selective COX-1 / COX-2 inhibitors, such as ibuprofen or ketoprofen; (15) COX-2 inhibitors, such as celecoxib and rofecoxib; (16) VLA-4 antagonists, such as those described in WO97/03094 and WO97/02289; (17) TACE inhibitors and TNF- α inhibitors, for example anti-TNF monoclonal antibodies, such as Remicade and CDP-870 and TNF receptor immunoglobulin molecules, such as Enbrel; (18) inhibitors of matrix metalloprotease, for example MMP-12; (19) human neutrophil elastase inhibitors, such as ONO-6818 or those described in WO2005/026124, WO2003/053930 and WO06/082412; (20) A2b antagonists such as those described in WO2002/42298; (21) modulators of chemokine receptor function, for example antagonists of CCR3 and CCR8; (22) compounds which modulate the action of other prostanoid receptors, for example a thromboxane A₂ antagonist; DP1 antagonists such as MK-0524, CRTH2 antagonists such as ODC9101 and OC000459 and AZD1981 and mixed DP1/CRTH2 antagonists such as AMG 009 and AMG853; (23) PPAR agonists including PPAR alpha agonists (such as fenofibrate), PPAR delta agonists, PPAR gamma agonists such as Pioglitazone, Rosiglitazone and Balaglitazone; (24) methylxanthines such as theophylline or aminophylline and methylxanthine/corticosteroid combinations such as theophylline/budesonide, theophylline/fluticasone propionate, theophylline/ciclesonide, theophylline/mometasone furoate and theophylline/beclometasone dipropionate; (25) A2a agonists such as those described in EP1052264 and EP1241176; (26) CXCR2 or IL-8 antagonists

such as SCH 527123 or GSK 656933; (27) IL-R signalling modulators such as kineret and ACZ 885; (28) MCP-1 antagonists such as ABN-912.

The invention is also directed to a kit comprising the pharmaceutical compositions of compounds of the invention alone or in combination with or
5 in admixture with one or more pharmaceutically acceptable carriers and/or excipients and a device which may be a single- or multi-dose dry powder inhaler, a metered dose inhaler or a nebulizer.

Methods of synthesis

In one aspect of the present invention, a process for the preparation of
10 compounds of the invention is provided, according to general synthetic routes described in this section. In the following reaction schemes, unless otherwise indicated, the groups mentioned assume the same meaning as those reported for compounds of formula (I).

The skilled person may introduce, where appropriate, suitable variations
15 to the conditions specifically described in the experimentals in order to adapt the synthetic routes to the provision of further compounds of the invention. Such variations may include, but are not limited to, use of appropriate starting materials to generate different compounds, changes in the solvent and temperature of reactions, replacements of reactives with analogous chemical
20 role, introduction or removal of protection/deprotection stages of functional groups sensitive to reaction conditions and reagents, as well as introduction or removal of specific synthetic steps oriented to further functionalisation of the chemical scaffold.

Processes which can be used and are described and reported in
25 Examples and Schemes, should not be viewed as limiting the scope of the synthetic methods available for the preparation of the compounds of the invention.

The process described is particularly advantageous as it is susceptible

of being properly modulated, through any proper variant known to the skilled person, so as to obtain any of the desired compounds of the invention. Such variants are comprised within the scope of the present invention.

From all of the above, it should be clear to the skilled person that any of the described groups may be present as such or in any properly protected form.

In particular, functional groups present in the intermediate and compounds and which could generate unwanted side reaction and by-products, need to be properly protected before the alkylation, acylation, coupling or sulfonylation takes place. Likewise, subsequent deprotection of those same protected groups may follow upon completion of the said reactions.

In the present invention, unless otherwise indicated, the term "protecting group" designates a protective group adapted to preserve the function of the group it is bound to. Typically, protective groups are used to preserve amino, hydroxyl, or carboxyl functions. Appropriate protecting groups may thus include, for example, benzyl, benzyloxycarbonyl, t-butoxycarbonyl, alkyl or benzyl esters or the like, which are well known to those skilled in the art [see, for a general reference, T.W. Green; Protective Groups in Organic Synthesis (Wiley, N.Y. 1981)].

Likewise, selective protection and deprotection of any of the said groups, for instance including carbonyl, hydroxyl or amino groups, may be accomplished according to very well known methods commonly employed in organic synthetic chemistry.


Optional salification of the compounds of formula (I) or N-oxides on the pyridine ring thereof may be carried out by properly converting any of the free acidic or amino groups into the corresponding pharmaceutically acceptable salts. In this case too, the operative conditions being employed for the optional salification of the compounds of the invention are all within the

ordinary knowledge of the skilled person.

From all of the above, it should be clear that the above process, comprehensive of any variant thereof for the preparation of suitable compounds of the invention, may be conveniently modified so that to adapt
 5 the reaction conditions to the specific needs, for instance by choosing appropriate condensing agents, solvents and protective groups, as the case may be.

For example compounds of the invention of formula (I) may be prepared according to the route illustrated in Scheme 1.

10



Scheme 1


Compounds of general formula (I) may be prepared from compounds of general formula (1a) by reaction with a compound of general formula (1b1) or
 15 (1b2) wherein R^2 is as defined in general formula (I), in a suitable solvent such as dimethyl sulfoxide, 1,4-dioxane, N,N-dimethylformamide or acetonitrile, in the presence of a base such as diisopropylethylamine at a range of temperatures, preferably between room temperature and 100°C.

Compounds of general formula (1b1) and (1b2) are either known in the
 20 literature or may be prepared from amines of general formula (1c) according to known literature procedures (e.g. see for reference WO2006009741, EP1609789).

Compounds of general formula (1c) are either known in the literature or

may be synthesised by one skilled in the art by adapting appropriate literature methods (e.g WO2010077836, WO2006009741, WO2008125014, J. Med Chem., 2007, 50, 4016, Bulletin des Societes Chimiques Belges, 1987, 96, 675-709, Organic & Biomolecular Chemistry, 2006, 4, 4158-4164).


- 5 Compounds of general formula (1ca), i.e. compounds of formula (1c) wherein R^2 is a group of formula (IIIb) and R^{17} , R^{18} , z^1 , z^2 , z^3 and z^4 are as defined above can be prepared from compounds of formula (1e),



- 10 using a suitable reducing agent such as tin (II) chloride, iron, or hydrogen gas with a suitable catalyst such as palladium on carbon, in a suitable solvent such as methanol, ethanol or acetic acid, at a range of temperatures, preferably between room temperature and 100°C.


- Compounds of general formula (1e) are known in the literature or may
15 be prepared by those skilled in the art using literature methods (e.g. WO2008034008, WO20110189167, WO2010068258).

- Alternatively, compounds of general formula (1ca) as above defined can be prepared from compounds of formula (1f), wherein R^{17} , R^{18} , z^1 , z^2 , z^3 and z^4 are as defined above and wherein PG is a suitable compatible protecting
20 group known to those skilled in the art, such as benzyl, benzyl carbamate or tert-butyl carbamate,



using suitable deprotection conditions such as hydrochloric acid, trifluoroacetic acid, or hydrogen catalysed by for example palladium on carbon, in a suitable solvent such as dichloromethane, methanol, ethanol or acetic acid, at a range of temperatures, preferably between 0°C and 100°C.


- 5 Compounds of general formula (1f) can be prepared by reaction of compounds of formula (1g), wherein R^{17} , R^{18} , z^1 , z^2 , z^3 and z^4 are as defined above



- with compounds of formula (1h) as above reported wherein PG is a
- 10 suitable protecting group known to those skilled in the art, such as benzyl, benzyl carbamate or tert-butyl carbamate, using suitable conditions such as in the presence of a base such as potassium carbonate or diisopropylethyl amine or under Buchwald conditions (with a catalyst such as $Pd(OAc)_2$, a ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and base such as sodium tert-
- 15 butoxide), in a suitable solvent such as toluene or tetrahydrofuran, at a range of temperatures, preferably between room temperature and 150°C.

- Compounds of general formula (1g) and (1h) are known in the literature or may be prepared by those skilled in the art by adapting appropriate literature methods (e.g. WO2011042389, Chemistry-A European Journal,
- 20 2011, 17, 6606-6609, S6606/1-S6606/38).

Compounds of general formula (1a) may be prepared according to the route illustrated in Scheme 2.




Scheme 2

Compounds of general formula (1a) may be prepared from compounds of general formula (2a) by reaction with a compound of general formula (2b), (2c) or (2d). Wherein G is a suitable chemical group known to those skilled in the art selected such that it can facilitate a suitable coupling reaction such as nucleophilic displacement or metal catalysed cross coupling: For example in cases such that when Y is -O-, -S- or -NR⁷-, examples of G may include halogen or a suitable leaving group such as mesylate or triflate either directly linked or attached via a group -(CR³R⁴)_n-. Examples of the coupling conditions used may include using a base such as sodium hydride or potassium *tert*-butoxide and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone in a suitable solvent such as N,N-dimethylformamide, toluene, 1,4-dioxane or acetonitrile at a range of temperatures, preferably between room temperature and 150°C. For example in cases such that when Y is -O- and G is -OH or -SH a method to perform this coupling may involve Mitsunobu conditions (diethylazodicarboxylate / triphenylphosphine) in a suitable solvent such as tetrahydrofuran or 1,4-dioxane at a range of temperatures preferably between -10°C and 100°C. For example in cases such as when Y is -O-, -S- or -NR⁷- and G is a group such as halogen, triflate or boronic acid/ester a method to perform this coupling may be under metal (for example palladium or copper)

catalysed coupling conditions in the presence of a suitable ligand such as Xantphos or 1,10-phenanthroline in the presence of a base such as caesium carbonate in a suitable solvent such as tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably between -10°C and 150°C. For example in cases such as when Y is -O- and G is a group such as -COOMe, -COOH, isocyanate, -OCOCl or -NHCOOCH₂CCl₃ examples of conditions to perform this coupling may involve the use of a base such as sodium hydride or triethylamine or a coupling reagent such as HATU in a suitable solvent such as dichloromethane, tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably between -10°C and 150°C.


Compounds of formula (2b) are commercially available, are known in the literature or may be synthesised from compounds of formula (2e), wherein R⁸ and R⁹ are as defined for compounds of formula (I), by adapting appropriate literature methods (e.g. WO 2006133006) or using methods known to those skilled in the art such as by reacting (2e) with a suitable alkylating agent such as dibromoethane or bromoethanol in the presence of a suitable base such as sodium hydride or potassium carbonate in a suitable solvent such as tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably between -10°C and 150°C, or by reacting (2e) with a suitable aldehyde in the presence of a reducing agent such as sodium triacetoxyborohydride in a suitable solvent such as dichloroethane or tetrahydrofuran at a range of temperatures preferably between -10°C and 100°C.



Compounds of formula (2e) are commercially available, are known in

the literature or may be synthesised by those skilled in the art using literature methods.

Compounds of formula (2c) may be synthesised from compounds of formula (2f).




Wherein X₁ and R¹² are defined as for compounds of formula (I), G is a group such as halogen, -O-PG or -S-PG wherein PG represents a protecting group such as triisopropylsilyl or *tert*-butyldimethylsilyl (methods for whose introduction and removal are well known by those skilled in the art) and J may represent groups such as halogen, -NH₂, -OH, -SH, -COOH, -SO₂Cl which can be modified using literature methods to introduce an appropriate group R¹⁰ by those skilled in the art. For example in cases such as when J is halogen, a method such as nucleophilic substitution with a suitable alcohol, amine or thiol may be used in the presence of a suitable base such as sodium hydride, triethylamine or potassium carbonate in a suitable solvent such as tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably between -10°C and 150°C. For example in cases such as when J is -NH₂, -OH or -SH, a method such as alkylation may be used with a suitable alkylating agent such as an alkyl halide in the presence of a suitable base such as sodium hydride or potassium carbonate in a suitable solvent such as tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably between -10°C and 150°C. For example in cases such as where J is -COOH or -SO₂Cl a method such as reaction with a suitable amine in the presence of a suitable base such as triethylamine or a coupling reagent such as HATU in a suitable solvent such as tetrahydrofuran, 1,4-dioxane or N,N-dimethylformamide at a range of temperatures preferably

between -10°C and 150°C.

Compounds of formula (2da), i.e. compounds of formula(2a) wherein $X^4 = C$ may be prepared according to the routes described in Scheme 3 herebelow:

5



Scheme 3

Compounds of general formula (2da) as above defined may be prepared from compounds of general formula (3e) using a suitable oxidant such as chloramine T, lead tetracetate or phenyl iodine(III) diacetate, in a suitable solvent such as dichloromethane or ethanol at a range of temperatures, preferably between room temperature and 100°C.

Compounds of general formula (3e) may be prepared from compounds of general formula (3a) by reaction with an aldehyde of general formula (3c) above reported.

in a suitable solvent such as ethanol or tetrahydrofuran at a range of temperatures, preferably between room temperature and 80°C.

Compounds of formula (3c) are commercially available, known in the literature or may be prepared using literature methods by those skilled in the art.

Alternatively, compounds of formula (2da) may be prepared from


compounds of formula (3d) using a suitable dehydrating agent such as Burgess' reagent, triphenyl phosphine and hexachloroethane, phosphorus oxychloride, acetic acid or Mitsunobu conditions (diethylazodicarboxylate/triphenylphosphine/trimethylsilylazide), in the
5 absence or presence of a suitable solvent such as tetrahydrofuran, toluene or NMP, at a range of temperatures, preferably between room temperature and 150°C.

Compounds of formula (3d) may be prepared from compounds of formula (3a) by reaction with a compound of general formula (3b1) using a
10 suitable acylating/dehydrating agent such as triphenylphosphine/trichloroacetonitrile in the presence of a base such as diisopropylethylamine, in a suitable solvent such as dichloromethane or acetonitrile, at a range of temperatures, preferably between room temperature and 150°C.

15 Or by reaction with a compound of general formula (3b2) in the presence of a base such as diisopropylethylamine, in a suitable solvent such as dichloromethane or THF at a range of temperatures preferably between -10°C and the boiling point of the solvent.

Compounds of formulae (3b1) and (3b2) are commercially available,
20 known in the literature or may be prepared by literature methods by those skilled in the art.

Alternatively, compounds of formula (2da) as above defined may be prepared according to the route in Scheme 4:



Scheme 4

Compounds of general formula (2da) may be prepared from compounds of general formula (4c). Wherein G² may represent groups such as halogen, -CHO, -COOH, -COOEt and SO₂Cl.

For example, compounds of general formula (2da) may be prepared from compounds of general formula (4c), wherein G² represents halogen, using methods such as a metal (for example palladium) catalysed coupling with a suitable R¹¹G⁵ derivative wherein G⁵ is a group such as boronate acid/ester or stannane in a suitable solvent such as tetrahydrofuran or 1,4-dioxane at a range of temperatures preferably between ambient temperature and 150°C. An alternative method may involve displacement of said halogen with a suitable group R¹¹H (such as that containing an -NH, -OH or -SH group) in the presence of a base such as sodium hydride, potassium *tert*-butoxide or N,N-diethylisopropylamine in a suitable solvent such as N,N-Dimethylformamide, toluene, 1,4-dioxane or acetonitrile at a range of temperatures, preferably between room temperature and 150°C.

The group G² may be also transformed from groups such as halogen to groups such as -CHO, -COOH, -COOEt and SO₂Cl by means of metal insertion methods known to those skilled in the art such as palladium catalysis, Grignard formation or lithium halogen exchange.

Compounds of general formula (2da) wherein R^{11} is a group such as -
 $CH_2-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_3-$
 $C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-OR^D$, $-C(O)N(R^C)-$
 $(C_3-C_7\text{cycloalkylene})-OR^D$, $-C(O)N(R^A R^B)$, $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-$
5 $NR^A R^B$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_2-$
 $C_6\text{alkylene})-OR^D$ or $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$ may be prepared
from compounds of general formula (4c), wherein G^2 represents -CHO,
-COOH, -COOEt and -SO₂Cl, by reaction with a suitable amine such as
HNR^AR^B etc using methods such as reductive amination (using a reagent such
10 as sodium triacetoxyborohydride) or amide/sulphonamide formation in the
presence of suitable reagents such as HATU with a base such as
N,N-diethylisopropylamine or trimethylaluminium in a suitable solvent such
as dichloromethane, N,N-dimethylformamide, toluene, 1,4-dioxane or
acetonitrile at a range of temperatures, preferably between room temperature
15 and 150°C.

Compounds of general formula (4c) wherein G^2 is a group such as
-COOEt, may be synthesised from compounds of general formula (4a) by
reaction with a compound such as diethyloxalate in the presence of an acid
such as acetic acid at a range of temperatures, preferably between room
20 temperature and 120°C.


Compounds of general formula (4c) wherein G^2 is a group such as
bromine or chlorine, may be synthesised from compounds of general formula
(4b) by reaction with a compound such as N-chlorosuccinimide or
N-bromosuccinimide in a solvent such as chloroform at a range of
25 temperatures, preferably between -10°C and room temperature.

Compounds of general formula (4b) may be synthesised from
compounds of general formula (4a) by reaction with a compound such as
diethoxymethylacetate at a range of temperatures, preferably between room

temperature and 100°C.

Compounds of general formula (2db), i.e. compounds of formula (2d) wherein X_4 is nitrogen, may be prepared from compounds of general formula (4b) wherein $X^4 = NH$, by reaction with a suitable alkylating agent R^{11} in the presence of a base such as caesium carbonate in a suitable solvent such as N,N-dimethylformamide at a range of temperatures, preferably between room temperature and 150°C.

Alternatively, compounds of general formula (1aa), i.e. compounds of formula (1a) wherein X^4 is CH may be prepared according to the route illustrated in Scheme 5.



Scheme 5

Compounds of general formula (1aa) may be prepared from compounds of general formula (5e) wherein PG is a suitable protecting group known in the art such as Boc by using the appropriate deprotection conditions such as trifluoroacetic acid in a solvent such as dichloromethane at a range of temperatures, preferably between -10°C and room temperature.

Compounds of general formula (5e) may be prepared from compounds of general formula (5d) using a suitable dehydrating agent such as Burgess' reagent, triphenyl phosphine and hexachloroethane, phosphorus oxychloride, acetic acid or Mitsunobu conditions.

(diethylazodicarboxylate/triphenylphosphine/trimethylsilylazide), in the absence or presence of a suitable solvent such as tetrahydrofuran, toluene or NMP, at a range of temperatures, preferably between room temperature and 120°C.

5 Compounds of general formula (5d) may be prepared from compounds of general formula (5c) by reaction with a compound of general formula (3b1) as above defined using a suitable acylating/dehydrating agent such as triphenylphosphine/trichloroacetonitrile in the presence of a base such as diisopropylethylamine, in a suitable solvent such as dichloromethane or
10 acetonitrile, at a range of temperatures, preferably between room temperature and 150°C.

 or by reaction with a compound of general formula (3b2) as above defined in the presence of a base such as diisopropylethylamine, in a suitable solvent such as dichloromethane or THF at a range of temperatures preferably
15 between -10°C and the boiling point of the solvent.


 Compounds of general formula (5c) may be prepared from compounds of general formula (5b) wherein G³ is a suitable leaving group such as halogen, by reaction with a reagent such as hydrazine monohydrate in a suitable solvent such as ethanol at a range of temperatures preferably between
20 room temperature and 100°C.

 Compounds of general formula (5b) may be prepared from compounds of general formula (5a) by reaction with a suitable protecting group reagent known in the art such as Boc anhydride in the presence of a base such as triethylamine in a suitable solvent such as dichloromethane or tetrahydrofuran
25 at a range of temperatures preferably between room temperature and 100°C.

 Compounds of general formula (5a) can be synthesised by the methods described above for the synthesis of (1a).

 Compounds of general formula (2aa), i.e. compounds of formula (2a)

wherein $Y = O, W = NH$ and PG is a suitable protective group such as trifluoroacetate may be prepared according to the route illustrated in scheme 6:



Scheme 6


Compounds of general formula (2aa) may be prepared from compounds of general formula (6b) and (6c) by removal of the protecting group PG using methods known in the art such as aqueous sodium hydroxide in a solvent such as methanol at a range of temperatures preferably between room temperature and 100°C.

Compounds of general formula (6b), wherein PG is a protecting group, preferably trifluoroacetamide, and the group $-OH$ is placed on the cycloalkylene portion of ring A may be prepared from compounds of general formula (6a) by using a chiral reductive method such as using formic acid and $RuCl[S,S-Tsdpn(p-cymene)]$ in the presence of a base such as triethylamine in a solvent such as N,N -dimethylformamide at a range of temperatures preferably between room temperature and 150°C. It will be recognised that compounds of formula (6a) may be homochiral as illustrated or be the opposite enantiomer or racemic.

It will be realised by those skilled in the art that any combination of stereocentres in (2aa) can be prepared using both enantiomers of (6a) and


using $\text{RuCl}[\text{R,R-Tsdpen}(\text{p-cymene})]$ or $\text{RuCl}[\text{S,S-Tsdpen}(\text{p-cymene})]$. Compound (2a) is drawn with no defined stereocentres but any combination can be obtained as illustrated in Scheme 2.

Compounds of formula (6a) can be prepared from compounds of
5 formula (6d)



using a suitable oxidant such as potassium permanganate and magnesium sulfate in a suitable solvent methanol/water at a range of temperatures preferably between room temperature and the boiling point of
10 the solvent. It will be recognised that compounds of formula (6d) may be homochiral as illustrated or be the opposite enantiomer or racemic.

Compounds of formula (6d) can be prepared from compounds of formula (6e) where PG is a suitable protecting group such as trifluoroacetate or tert-butyl carbonate:




15 using ethyl trifluoroacetate or di-tert-butyl dicarbonate in the presence of base such as triethylamine or diisopropylethylamine in a solvent such as methanol or dichloromethane at a range of temperatures preferably between 0°C and the boiling point of the solvent. It will be recognised that compounds
20 of formula (6e) may be homochiral as illustrated or be the opposite enantiomer or racemic.

Compounds of formula (6e) are known in the literature and may be prepared by those skilled in the art by adapting literature methods (e.g. for S-(+)-1-amino-1,2,3,4-tetrahydronaphthalene see Journal of the Chemical
25 Society, Perkin Transactions 1: 1985, 2039-44, for (S)-(+)-8-amino-5,6,7,8-

tetrahydroquinoline see Journal of Organic Chemistry, 2007, 72, 669-671 and for 1-aminoindan see Tetrahedron Letters, 2011, 52, 1310-1312).

Compounds of general formula (2ab), i.e. compounds of formula (2a) wherein $Y = NR^7$ and $W = NH$, may be prepared according to the route 5 illustrated in scheme 7:



Scheme 7

Compounds of general formula (2ab) may be prepared from compounds 10 of general formula (7a) by removal of the protecting group PG using methods known in the art such as aqueous sodium hydroxide in a solvent such as methanol or trifluoroacetic acid in dichloromethane at a range of temperatures preferably between room temperature and 100°C.

Compounds of formula (7a) may be prepared from compounds of 15 general formula (6a) and amines (7b) by reaction under reductive amination conditions, using a reducing agent such as sodium triacetoxyborohydride and a solvent such as 1,2-dichloroethane at a range of temperatures preferably between room temperature and 100°C.


Compounds for formula (7b) are known in the literature and may be 20 prepared by those skilled in the art using literature procedures. Compounds of

formula (6a) can be prepared as described above.

Alternatively, compounds of formula (7a) may be prepared from compounds of general formula (7c) wherein G^4 is a suitable chemical group known to those skilled in the art selected such that it can facilitate a reaction such as a nucleophilic substitution. For example G is a suitable leaving group such as halogen or mesylate which can react with a suitable amine (7b) in the presence of a suitable base such as sodium hydride or potassium *tert*-butoxide and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone in a suitable solvent such as N,N-dimethylformamide, toluene, 1,4-dioxane or acetonitrile at a range of temperatures, preferably between room temperature and 150°C.

Compounds of formula (7c) can be prepared from compounds of formulae (6b) or (6c) using halogenating conditions such as carbon tetrabromide and triphenylphosphine in dichloromethane or activation conditions such as methane sulfonyl chloride in dichloromethane in the presence of base such as diisopropylamine.

Alternatively, compounds of general formula (Id), i.e. compounds of formula (I) wherein $Y = S$ and $W = NH$ may be prepared according to the route illustrated in scheme 8:



Scheme 8

Compounds of general formula (Id) may be prepared from compounds of general formula (1ab), i.e. compounds of formula (1a) wherein $Y = S$ and

W = NH: using compounds of formula (1b1) or (1b2) in a suitable solvent such as dimethyl sulfoxide, 1,4-dioxane, N,N-dimethylformamide or acetonitrile, in the presence of a base such as diisopropylethylamine at a range of temperatures, preferably between room temperature and 100°C.

- 5 Compounds of formula (1ab) as above defined may be prepared from compounds of formula (8g) using deprotection conditions such as hydrazine in methanol at a range of temperatures preferably between room temperature and the boiling point of the solvent.

 Compounds of formula (8g) wherein Y = S, can be prepared from
10 compounds of formula (8f) by reaction with compounds of formulae (2b), (2c) or (2d). Examples of the coupling conditions used may include using a base such as sodium hydride or potassium *tert*-butoxide and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone in a suitable solvent such as N,N-dimethylformamide, toluene, 1,4-dioxane or acetonitrile at a range of
15 temperatures, preferably between room temperature and 150°C. Alternative methods to perform this coupling may involve Mitsunobu conditions (diethylazodicarboxylate / triphenylphosphine) or metal (for example palladium) catalysed coupling conditions in a suitable solvent such as tetrahydrofuran or 1,4-dioxane at a range of temperatures preferably between
20 -10°C and 150°C and starting from the appropriate derivative of formula (2b), (2c), or (2d).

 Compounds of formula (8f) can be prepared from compounds of formula (8e) using dithiothreitol, monopotassium phosphate, potassium carbonate in a solvent such as methanol in the presence of acetic acid at a
25 range of temperatures preferably between room temperature and the boiling point of the solvent.

 Compounds of formula (8e) can be prepared from compounds of formula (8d) using 2-nitrobenzenesulfonyl chloride in acetic acid at a range of

temperatures preferably between room temperature and 100°C.


Compounds of formula (8d) can be prepared from compounds of formula (8c) using phthalimide, triphenylphosphine and diisopropyl azodicarboxylate in a solvent such as tetrahydrofuran at a range of
5 temperature preferably between 0°C and the boiling point of the solvent.

Compounds of formula (8c) can be prepared from compounds of formula (8b) using a reducing agent such as sodium borohydride in a solvent such as methanol at a range of temperatures preferably between 0°C and the boiling point of the solvent.

10 Compounds of formula (8b) can be prepared from compounds of formula (8a) using *tert*-butanethiol in the presence of a base such as diisopropylethylamine in a solvent such as tetrahydrofuran at a range of temperatures preferably between 0°C and the boiling point of the solvent.

Compounds of formula (8a) are known in the literature and can be
15 prepared by those skilled in the art using literature methods (e.g. 3-bromo-indan-1-one see WO 2010108058).

Alternatively, compounds of general formula (1ab), i.e. compounds of formula (1a) wherein $Y = CH_2$, $W = NH$ and PG is a suitable protective group such as trifluoroacetate may be prepared according to the route illustrated in
20 scheme 9:



Scheme 9

Compounds of general formula (1ab) may be prepared from compounds of general formula (9b) by removal of the protecting group PG using methods known in the art such as aqueous sodium hydroxide in a solvent such as methanol at a range of temperatures preferably between room temperature and 100°C.

Compounds of formula (9b) may be prepared from compounds of general formula (9a) by reaction with a suitable reduction agent for example hydrogen gas in the presence of a suitable catalyst such as palladium on activated charcoal in a suitable solvent such as ethanol at a range of temperatures between room temperature and 70°C and pressures between atmospheric and 4 Barr.

Compounds of formula (9a) may be prepared from compounds of general formula (6a) by means of a reaction such as a Wittig (or one of the closely related variants such as the Horner-Wadsworth-Emmons) with a suitable substrate such as $\text{R}^1\text{-CH}_2\text{-P(O)(OMe)}_2$ in the presence of a suitable base such as sodium hydride in a suitable solvent such as tetrahydrofuran at a range of temperatures preferably between -10°C and 100°C.


Compounds such as $\text{R}^1\text{-CH}_2\text{-P(O)(OMe)}_2$ may be synthesised from compounds of the general formula $\text{R}^1\text{-CH}_2\text{-Hal}$ wherein Hal represents a

halogen such as -Br or -Cl by reaction with a compound such as trimethylphosphite at a range of temperatures preferably between 0°C and 100°C.

Compounds such as $R^1\text{-CH}_2\text{-Hal}$ may be synthesised from compounds
5 of formula $R^1\text{-CH}_3$ by means of a reaction such as a radical halogenation using
a reagent such as N-bromosuccinimide in the presence of a catalyst such as
AIBN in a suitable solvent such as carbon tetrachloride at a range of
temperatures preferably between 0°C and 80°C. Compounds such as $R^1\text{-CH}_2\text{-}$
Hal may also be synthesised from compounds formula $R^1\text{-CH}_2\text{-OH}$ by means
10 of using halogenating conditions such as carbon tetrabromide and
triphenylphosphine in dichloromethane or activation conditions such as
methane sulfonyl chloride in dichloromethane in the presence of base such as
diisopropylamine.

Compounds such as $R^1\text{-CH}_3$ and $R^1\text{-CH}_2\text{-OH}$ may be prepared by
15 methods outlined above for compounds (2b), (2c) and (2d).

Compounds of the invention of formula (1ac), i.e. compounds of
formula (1a) where $Y = (\text{CR}^5\text{R}^6)_n$ and $W = \text{NH}$, may be prepared according to
the route illustrated in Scheme 10.

Scheme 10

Compounds of formula (1ac) may be prepared from compounds of formula (10c) wherein PG is a suitable protecting group known to those skilled in the art, such as trifluoroacetamide, tert-butyl carbamate and benzyl carbamate by using suitable de-protection conditions such as, sodium hydroxide in methanol, trifluoroacetic acid in dichloromethane or hydrogen gas catalysed by for example palladium on carbon in ethanol, at a range of temperatures, preferably between 0°C and 100°C.

Compounds of formula (10c) may be prepared from compounds of formula (10b) wherein PG is a suitable protecting group known to those skilled in the art, such as trifluoroacetamide, tert-butyl carbamate and benzyl carbamate by using hydrogen gas in the presence of a catalyst such as palladium on carbon, in a suitable solvent such as methanol or ethanol, in the presence or absence of an acid such as HCl, at a range of temperatures, preferably between 0°C and 100°C.

Compounds of formula (10b) may be prepared from compounds of formula (10a) and (10f) by a reaction such as a cross-coupling using a suitable

catalyst such as tetrakis(triphenylphosphine)palladium (0) or palladium acetate, and a base such as diisopropylethylamine, sodium tert-butoxide or caesium carbonate in a suitable solvent such as NMP, toluene or DMF, at a range of temperatures, preferably between 0°C and 100°C. Alternatively (10b) may be prepared by adapting literature procedures (e.g those reported in WO2009022633).

Compounds of formula (10f) are known in the literature or may be prepared by those skilled in the art by adapting literature procedures (e.g WO2008063287).

Compounds of formula (10a) may be prepared from compounds of formula (6a) using a triflating agent such as triflic anhydride, in the presence of a suitable base such as pyridine or 2,6-bis(tert-butyl)-4-methylpyridine, in a solvent such as dichloromethane or chloroform at a range of temperatures, preferably between 0°C and boiling point of the solvent. Alternatively (10a) may be prepared by adapting literature procedures (e.g those described in WO2009022633).

General experimental details

Abbreviations used in the experimental section: AcOH = acetic acid; aq. = aqueous; DCM = dichloromethane; DIAD = Diisopropyl azodicarboxylate; DIPEA = diisopropylethylamine; DMAP = N,N- dimethylaminopyridine; DMF = N,N- dimethylformamide; DMSO = dimethyl sulfoxide; EDC = 1-ethyl-3-(3'- dimethylaminopropyl)carbodiimide Hydrochloride; EtOAc = ethyl acetate; EtOH = ethanol; Et₂O = diethyl ether; Et₃N = triethylamine; EtNiPr₂ = diisopropylethylamine; FCC = flash column chromatography; h = hour; HATU = 2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate; HOBt = 1-hydroxy-benzotriazole; HPLC = high performance liquid chromatography; IMS = Industrial Methylated Spirits; LCMS = liquid chromatography mass spectrometry; NaOH = sodium hydroxide; MeCN =

acetonitrile; MeOH = MeOH; min = minutes; NH₃ = ammonia; NMR = nuclear magnetic resonance; RT = room temperature; Rt = retention time; sat. = saturated; SCX-2 = strong cation exchange chromatography; TFA = trifluoroacetic acid; THF = Tetrahydrofuran; H₂O = water; IMS = industrial methylated spirit; Xantphos =
5 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene; X-Select = Waters X-select HPLC column; IPA = propan-2-ol; LDA = lithium diisopropylamide; MDAP = mass-directed auto-purification; MeOH = methanol; Ph₃P = triphenylphosphine; TBAF = tetrabutylammonium fluoride.

In the procedures that follow, after each starting material, reference to a
10 Intermediate/Example number is usually provided. This is provided merely for assistance to the skilled chemist. The starting material may not necessarily have been prepared from the batch referred to.

When reference is made to the use of a “similar” or “analogous” procedure, as will be appreciated by those skilled in the art, such a procedure
15 may involve minor variations, for example reaction temperature, reagent/solvent amount, reaction time, work-up conditions or chromatographic purification conditions.

The nomenclature of structures was assigned using Autonom 2000 Name software from MDL Inc. When the nomenclature of structures could not
20 be assigned using Autonom, ACD/Name software utility part of the ACD/Labs Release 12.00 Product Version 12.5 (Build 45133, 16 Dec 2010) was used. Stereochemical assignments of compounds are based on comparisons with data reported in WO2008/043019 for key intermediates. All reactions were carried out under anhydrous conditions and an atmosphere of nitrogen or
25 argon unless specified otherwise. Unless otherwise stated all transformations were carried at ambient temperature (room temperature).

NMR spectra were obtained on a Varian Unity Inova 400 spectrometer with a 5mm inverse detection triple resonance probe operating at 400 MHz or

on a Bruker Avance DRX 400 spectrometer with a 5 mm inverse detection triple resonance TXI probe operating at 400 MHz or on a Bruker Avance DPX 300 spectrometer with a standard 5mm dual frequency probe operating at 300 MHz. Shifts are given in ppm relative to tetramethylsilane ($\delta = 0$ ppm). J values are given in Hz through-out. NMR spectra were assigned using DataChord Spectrum Analyst Version 4.0.b21 or SpinWorks version 3.

Where products were purified by flash column chromatography, 'flash silica' refers to silica gel for chromatography, 0.035 to 0.070 mm (220 to 440 mesh) (e.g. Fluka silica gel 60), and an applied pressure of nitrogen up to 10 p.s.i accelerated column elution or use of the CombiFlash® Companion purification system or use of the Biotage SP1 purification system. All solvents and commercial reagents were used as received.

Compounds purified by preparative HPLC were purified using a C18-reverse-phase column (100 × 22.5 mm i.d. Genesis column with 7 μ m particle size), or a Phenyl-Hexyl column (250 x 21.2 mm i.d. Gemini column with 5 μ m particle size), UV detection between 220 - 254 nm, flow 5-20 mL/min), eluting with gradients from 100-0 to 0-100% water/acetonitrile (containing 0.1% TFA or 0.1% formic acid) or water/MeOH (containing 0.1% TFA or 0.1% formic acid), or a C18-reverse-phase column (19 × 250 mm, XBridge OBD, with 5 μ m particle size), eluting with gradients from 100-0 to 0-100% water/acetonitrile (containing 0.1% NH_4OH); or a ChiralPak IC column (10 × 250 mm i.d., with 5 μ m particle size), unless otherwise indicated. Fractions containing the required product (identified by LCMS analysis) were pooled, the organic solvent removed by evaporation, and the remaining aqueous residue lyophilised, to give the final product. Products purified by preparative HPLC were isolated as free base, formate or TFA salts, unless otherwise stated.

The Liquid Chromatography Mass Spectroscopy (LCMS) and HPLC systems used are:

Method 1

Waters Platform LC Quadrupole mass spectrometer with a C18-reverse-phase column (30 × 4.6 mm Phenomenex Luna 3 µm particle size), elution with A: water + 0.1% formic acid; B: acetonitrile + 0.1% formic acid.

5 Gradient:

	Gradient - Time	flow mL/min	%A	%B
	0.00	2.0	95	5
	0.50	2.0	95	5
	4.50	2.0	5	95
10	5.50	2.0	5	95
	6.00	2.0	95	5

Detection - MS, ELS, UV (200 µL split to MS with in-line HP1100 DAD detector). MS ionization method - Electrospray (positive and negative ion).

15 **Method 2**

Waters ZMD quadrupole mass spectrometer with a C18-reverse-phase column (30 × 4.6 mm Phenomenex Luna 3 µm particle size), elution with A: water + 0.1% formic acid; B: acetonitrile + 0.1% formic acid. Gradient:

	Gradient - Time	flow mL/min	%A	%B
20	0.00	2.0	95	5
	0.50	2.0	95	5
	4.50	2.0	5	95
	5.50	2.0	5	95
	6.00	2.0	95	5

25 Detection - MS, ELS, UV (200 µL split to MS with in-line Waters 996 DAD detector). MS ionization method - Electrospray (positive and negative ion).

Method 3

Waters ZMD quadrupole mass spectrometer with a C18-reverse-phase column (30 × 4.6 mm Phenomenex Luna 3 µm particle size), elution with A: water + 0.1% formic acid; B: acetonitrile + 0.1% formic acid. Gradient:

5	Gradient - Time	flow mL/min	%A	%B
	0.00	2.0	95	5
	0.50	2.0	95	5
	4.50	2.0	5	95
	5.50	2.0	5	95
10	6.00	2.0	95	5

Detection - MS, ELS, UV (200 µL split to MS with in-line HP1100 DAD detector). MS ionization method - Electrospray (positive and negative ion).

Method 4

15 VG Platform II quadrupole spectrometer with a C18-reverse-phase column (30 × 4.6 mm Phenomenex Luna 3 µm particle size, elution with A: water + 0.1% formic acid; B: acetonitrile + 0.1% formic acid. Gradient:

	Gradient - Time	flow mL/min	%A	%B
	0.00	2.0	95	5
20	0.30	2.0	95	5
	4.30	2.0	5	95
	5.30	2.0	5	95
	5.80	2.0	95	5
	6.00	2.0	95	5

25 Detection - MS, ELS, UV (200 µl/min split to the ESI source with inline HP1050 DAD detector). MS ionization method - Electrospray (positive and negative ion).

Method 5

Waters micromass ZQ2000 quadrupole mass spectrometer with an Acquity BEH C18 1.7um 100 x 2.1mm, Acquity BEH Shield RP18 1.7um 100 x 2.1mm or Acquity HSST3 1.8um 100 x 2.1mm, maintained at 40°C. Elution
 5 with A: water + 0.1% formic acid; B: acetonitrile + 0.1% formic acid.
 Gradient:

	Gradient - Time	flow mL/min	%A	%B
	0.00	0.4	95	5
	0.40	0.4	95	5
10	6.00	0.4	5	95
	6.80	0.4	5	95
	7.00	0.4	95	5
	8.00	0.4	95	5

Detection - MS, UV PDA. MS ionization method - Electrospray
 15 (positive and negative ion).

Method 6

Phenomenex Gemini C18-reverse-phase column (250 × 21.20 mm 5 µm particle size), elution with A: water + 0.1% formic acid; B: CH₃CN + 0.1% formic acid. Gradient - 90% A/10% B to 2% A/98% B over 20 min - flow rate
 20 18 mL/min. Detection - In-line UV detector set at 254 nM wavelength.

Method 7

Agilent 1260 infinity purification system. Column: XSELECT CSH Prep C18 OBD, particle size 5µm, 30 × 150mm, RT. Elution with A: water + 0.1% formic acid; B: CH₃CN + 0.1% formic acid. Gradient - 90% A/10% B to
 25 2% A/95% B over 22 min - flow rate 60 mL/min. Detection - In-line Agilent 6100 series single Quadrupole LC/MS.

Method 8

Agilent 1260 infinity purification system. Column: XBridge Prep C18 OBD, particle size 5 μ m, 30 $\times$ 150mm, RT. Elution with A: water + 0.1% ammonia; B: CH₃CN + 0.1% ammonia. Gradient - 90% A/10% B to 2% A/98% B over 22 min - flow rate 60 mL/min. Detection - In-line Agilent 6100 series single Quadrupole LC/MS.


Differential Scanning Calorimetry (DSC): it should be recognized that the endotherm peak as measured is dependent on a number of factors including the machine used, the rate of heating, the calibration standard, humidity and the purity of the sample used.

Melting points reported in the experimentals are estimated on the basis of the onset of endotherm peaks registered during DSC analysis.


It is to be understood by the skilled person that, where the expression “partial formate salt” is used, it is to be intended as identifying derivatives where only part of the basic compound has been converted into formate salt and thus containing less than one equivalent of formate counterion. Exact salt/free base ratio is provided by associated NMS analysis.

Intermediate A

(1*R*,4*S*)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-ol




a. **2,2,2-Trifluoro-*N*-(*S*)-1,2,3,4-tetrahydro-naphthalen-1-yl-acetamide (Intermediate Aa)**



5 Ethyl trifluoroacetate (24.2 mL, 204 mmol) was added dropwise to a solution of (*S*)-(1,2,3,4-tetrahydro-naphthalen-1-yl)amine (Alfa Aesar; 25.0 g, 170 mmol) and triethylamine (35.5 mL, 255 mmol) in MeOH (250 mL) at RT and stirred for 18 h. The mixture was concentrated to approximately 1/3 of its volume and then partitioned between DCM (200 mL) and water (200 mL). The
10 aqueous layer was extracted into DCM (3 x) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to yield the *title compound* (41.1 g, 169 mmol, 99%). ¹H NMR (400 MHz, CDCl₃): 1.80-1.95 (3H, m), 2.05-2.15 (1H, m), 2.75-2.90 (2H, m), 5.18-5.25 (1H, q, J 5.0 Hz), 6.38-6.48 (1H, br s), 7.12-7.16 (1H, m), 7.20-7.26 (3H, m).

15 b. **2,2,2-Trifluoro-*N*-((*S*)-4-oxo-1,2,3,4-tetrahydro-naphthalen-1-yl)-acetamide (Intermediate Ab)**




Magnesium sulfate monohydrate (46.6 g, 338 mmol) in water (500 mL)
20 was added to an ice cold solution of Intermediate Aa (41.1 g, 169 mmol) in acetone (1.0 L). Potassium permanganate (80.1 g, 507 mmol) was added portionwise (10.0 g portions) over a period of 45 min. The mixture was then stirred for 18 h. Sodium thiosulfate pentahydrate (126 g, 510 mmol) in water (400 mL) was added and the reaction stirred for 30 min. The mixture was
25 concentrated to ~300 mL, then water (1.0 L), Celite (60 g) and EtOAc (1.0 L)

were sequentially added. The mixture was thoroughly stirred, and then filtered through a pad of Celite. The aqueous layer was extracted into EtOAc (3 x) and the combined organic layers washed with brine, dried (MgSO₄) and concentrated *in vacuo* to yield the *title compound* (36.6 g, 142 mmol, 84%).

5 ¹H NMR (400 MHz, CDCl₃): 2.20-2.30 (1H, dddd, J 13.3, 10.0, 8.8, 4.5 Hz), 2.43-2.52 (1H, dddd, J 13.3, 7.2, 4.6, 4.6 Hz), 2.67-2.77 (1H, ddd, J 17.4, 10.1, 4.6 Hz), 2.78-2.88 (1H, ddd, J 17.4, 7.1, 4.6 Hz), 5.39-5.47 (1H, td, 8.5, 4.5 Hz), 7.32-7.37 (1H, d, J 7.7 Hz), 7.44-7.49 (1H, t, J 7.6 Hz), 7.59-7.64 (1H, td, J 7.6, 1.4 Hz), 8.03-8.07 (1H, dd, J 7.7, 1.4 Hz).

10 c. **2,2,2-Trifluoro-N-((1S,4R)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-acetamide (Intermediate Ac)**



Degassed DMF (argon sparged, 100 mL) was added to Intermediate Ab
15 (8.00 g, 31.3 mmol) and [N-[(1R,2R)-2-(amino-κN)-1,2-diphenylethyl]-4-methylbenzenesulfonamidato-κN]chloro[(1,2,3,4,5,6-η)-1-methyl-4-(1-methylethyl)benzene]-ruthenium (Strem Chemicals Inc.; 594 mg, 0.93 mmol). Triethylamine (8.66 mL, 62.6 mmol) was added slowly to ice cold formic acid (2.34 mL, 62.6 mmol) and stirred for 20 min, this was then added to the DMF
20 solution. The reaction was heated to 60°C for 18 h. After cooling, the mixture was partitioned between DCM (200 mL) and water (600 mL). The aqueous layer was extracted DCM (3 x) and the combined organic layers washed with brine, dried (MgSO₄) and concentrated *in vacuo*. Purification by FCC, using 0-100% EtOAc in cyclohexane, afforded the *title compound* (7.10 g, 27.4
25 mmol, 88%). ¹H NMR (400 MHz, CDCl₃): 1.88-1.92 (1H, d, J 4.8 Hz), 1.98-2.18 (4H, m), 4.80-4.88 (1H, m), 5.165-5.24 (1H, m), 6.70-6.80 (1H, br s),


7.25-7.30 (1H, m), 7.30-7.40 (2H, m), 7.45-7.50 (1H, m).

**d. (1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-ol
(Intermediate A)**


Sodium hydroxide (2.10 g, 53.0 mmol) was added to an ice cold
5 solution of Intermediate Ac (3.43 g, 13.2 mmol) in MeOH/water (2:1, 50 mL)
and stirred for 3.5 h. The mixture was loaded on to a SCX-2 cartridge, eluting
with MeOH then 2M NH₃ in MeOH, to yield the *title compound* (2.30 g, 13.2
mmol, 99%). ¹H NMR (400 MHz, d₆-DMSO): 1.66-1.90 (4H, m), 3.71-3.77
(1H, t, J 5.4 Hz), 4.46-4.54 (1H, t, J 5.4 Hz), 7.14-7.22 (2H, m), 7.32-7.38
10 (1H, m), 7.40-7.46 (1H, m).

Intermediate B

(1S,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-ol



15 **a. 2,2,2-Trifluoro-N-((1S,4S)-4-hydroxy-1,2,3,4-tetrahydro-
naphthalen-1-yl)-acetamide (Intermediate Ba)**



Argon was bubbled through a solution of Intermediate Ab (8.00 g, 31.1
20 mmol) and [N-[(1S,2S)-2-(amino-κN)-1,2-diphenylethyl]-4-
methylbenzenesulfonamidato-κN]chloro[(1,2,3,4,5,6-η)-1-methyl-4-(1-
methylethyl)benzene]-ruthenium (Strem Chemicals Inc.; 0.06 g, 0.93 mmol) in
dry DMF (100 mL) for 10 min. A premixed combination of formic acid

(2.4 mL, 62.2 mmol) and Et₃N (8.60 mL, 62.2 mmol) was added and the mixture stirred at 50°C for 24 h. After cooling, the mixture was concentrated to ~25 mL. Water (70 mL) was added and the resulting precipitate filtered, and washed with DCM (3 × 30 mL) and diethyl ether (30 mL) to leave a solid
5 (4.75 g). The filtrate was decanted to leave a dark solid. Subsequent purification by FCC, using 0-30% EtOAc in cyclohexane, gave a solid. This was combined with the first obtained solid to give the *title compound* as a beige solid (5.93 g, 74%). ¹H NMR (400 MHz, d₆-DMSO): 1.60-1.83 (2H, m), 2.06-2.17 (2H, m), 4.60 (1H, m), 5.08 (1H, m), 5.28 (1H, d), 7.07 (1H, m),
10 7.25 (1H, ddd), 7.28 (1H, ddd), 7.50 (1H, dd), 9.78 (1H, d).


b. (1S,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-ol (Intermediate B)

To a grey solution of Intermediate Ba (5.55 g, 21.4 mmol) in MeOH (50 mL), NaOH (1.28 g, 32.1 mmol) in water (15 mL) was added and the mixture stirred at RT for 3 days. NaOH (1.28 g, 32.1 mmol) was added and
15 the brown solution was stirred for 5 h. The solution was loaded on to a SCX-2 cartridge, eluting with MeOH then 2M NH₃ in MeOH, to leave a grey solid. The solid was suspended in DCM (50 mL), sonicated, filtered and dried under vacuum to leave the *title compound* as a pale grey solid (2.93 g, 84%). ¹H NMR (400 MHz, d₆-DMSO): 1.41-1.64 (2H, m), 2.02-2.13 (2H, m), 3.82 (1H, dd), 4.55 (1H, dd), 5.08 (1H, br s), 7.13-7.22 (2H, m), 7.35-7.49 (2H, m).
20


Example 1

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(2-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

5



a. 3-Pyrrolidin-1-yl-propionic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 1a)




10

5-fluoro-2-hydrazinopyridine (285 mg, 2.24 mmol) and 3-(pyrrolidin-1-yl)propanoic acid hydrochloride (400 mg, 2.24 mmol) were dissolved in DMF (15 mL). EDC (516 mg, 2.69 mmol), HOBt (30.0 mg, 0.22 mmol) and triethylamine (374 μ L, 2.69 mmol) were added and the reaction stirred for 18 h. The mixture was loaded onto an SCX-2 cartridge, which was washed with MeOH then with 2M NH_3 in MeOH. The basic fractions were evaporated *in vacuo* then purified by FCC using 0-10% [2M NH_3 in MeOH] in DCM to give the *title compound* contaminated with several impurities (400 mg). The product was used in the next step without further purification.


20

b. 6-Fluoro-3-(2-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 1b)



Intermediate 1a (400 mg) was dissolved in THF (20 mL) and cooled in
5 an ice/water bath. Triphenylphosphine (643 mg, 2.45 mmol) was added
followed by triethylamine (682 μ L, 4.90 mmol) and hexachloroethane
(581 mg, 2.45 mmol). The reaction mixture was stirred for 18 h and then
loaded onto an SCX-2 cartridge, which was washed with MeOH then eluted
with 2M NH_3 in MeOH. The resulting residue was purified by FCC using
10 0-10% [2M NH_3 in MeOH] in DCM to give the *title compound* (292 mg, 1.16
mmol, 52%). ^1H NMR (400 MHz, CDCl_3): 1.80-1.87 (4H, m), 2.60-2.68 (4H,
m), 3.00-3.10 (2H, t, J 7.3), 3.24-3.34 (2H, t, J 7.3), 7.13-7.21 (1H, ddd, J 9.9,
7.6, 2.3), 7.70-7.77 (1H, dd, J 9.8, 4.5), 8.00-8.04 (1H, m).

c. (1S,4R)-4-[3-(2-Pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 1c)



Intermediate A (189 mg, 1.16 mmol) was added to a suspension of
sodium hydride (60% in mineral oil, 139 mg, 3.48 mmol) in DMF (5 mL). The
20 reaction was stirred for 20 mins, then Intermediate 1b (292 mg, 1.16 mmol)
was added in DMF (5 mL) and the reaction heated to 60°C for 2 h. The
mixture was cooled and quenched by dropwise addition of MeOH. The

solution was loaded onto an SCX-2 cartridge, which was washed with MeOH and product eluted with 2M NH₃ in MeOH. The residue was purified by FCC using 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* (270 mg, 0.74 mmol, 64%). ¹H NMR (400 MHz, d₄-MeOD): 1.80-1.86 (4H, m), 1.92-2.02 (1H, m), 2.05-2.18 (2H, m), 2.30-2.40 (1H, m), 2.62-2.70 (4H, m), 2.99-3.05 (2H, t, J 7.8), 3.28-3.36 (2H, t, J 7.8), 3.98-4.04 (1H, dd, J 8.6, 5.2), 5.46-5.50 (1H, t, J 4.3), 7.22-7.40 (4H, m), 7.54-7.58 (1H, d, J 7.8), 7.63-7.68 (1H, dd, J 9.8, 0.7), 7.97 (1H, s), 8.09-8.11 (1H, d, J 1.6).


d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(2-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 1)

Intermediate 1c (270 mg, 0.74 mmol) was dissolved in 1,4-dioxane (6 mL) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 329 mg, 0.81 mmol) and DIPEA (257 µL, 1.48 mmol) were added. The reaction was heated to 60°C for 18 h. After cooling, the mixture was partitioned between EtOAc (50 mL) and water (50 mL), and then extracted into EtOAc (3 x). The combined organic extracts were washed with brine, dried (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, then further purified by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% formic acid) to give the *title compound* (78.0 mg, 0.12 mmol, 17%). LCMS (Method 5): Rt 3.67 mins, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.90-2.06 (6H, m), 2.07-2.18 (1H, m), 2.22-2.30 (1H, m), 2.39 (3H, s), 3.08-3.16 (4H, m), 3.42-3.50 (4H, m), 4.87-4.92 (1H, dd, J 9.0, 5.6), 5.43-5.47 (1H, t, J 4.2), 6.33 (1H, s), 7.20-7.36 (10H, m), 7.62-7.66 (1H, d, J 9.7), 8.07-8.09 (1H, d, J 1.6), 8.45 (0.6H, s).


Example 2

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[2-(4-

**methyl-piperazin-1-yl)-ethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-
1,2,3,4-tetrahydro-naphthalen-1-yl)-urea formate salt**




**a. 3-(4-Methyl-piperazin-1-yl)-propionic acid N'-(5-fluoro-pyridin-
5 2-yl)-hydrazide (Intermediate 2a)**



5-fluoro-2-hydrazinopyridine (295 mg, 2.32 mmol) and 3-(4-methyl-1-
piperazinyl)propionic acid (400 mg, 2.32 mmol) were dissolved in DCM
10 (15 mL). EDC (536 mg, 2.79 mmol) and HOBt (31.0 mg, 0.23 mmol) were
added and the reaction stirred for 18 h. The mixture was loaded onto an SCX-
2 cartridge, which was washed with MeOH product eluted with 2M NH₃ in
MeOH. Further purification by FCC, using 0-10% [2M NH₃ in MeOH] in
DCM, afforded the *title compound* (458 mg, 1.63 mmol, 73%). ¹H NMR (400
15 MHz, CDCl₃): 2.32 (3H, s), 2.48-2.54 (2H, t, J 6.0), 2.54-2.70 (8H, br s),
2.69-2.75 (2H, t, J 6.0), 6.60-6.65 (1H, dd, J 8.9, 3.3), 6.75-6.78 (1H, d, J 4.4),
7.25-7.34 (1H, m), 8.02-8.05 (1H, d, J 3.0), 10.50 (1H, s).


b. 6-Fluoro-3-[2-(4-methyl-piperazin-1-yl)-ethyl]-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 2b)



5 Intermediate 2a (458 mg, 1.63 mmol) was dissolved in THF (15 mL) and cooled in an ice/water bath. Triphenylphosphine (854 mg, 3.26 mmol) was added followed by triethylamine (907 μ L, 6.52 mmol) and hexachloroethane (773 mg, 3.26 mmol). The reaction was stirred for 18 h and then loaded onto an SCX-2 cartridge, which was washed with MeOH and product eluted with
10 2M NH_3 in MeOH. Further purification by FCC, using 0-10% [2M NH_3 in MeOH] in DCM, afforded the *title compound* (310 mg, 1.18 mmol, 72%). LCMS (Method 4): Rt 0.28 min, m/z 264 [MH^+].

c. (1S,4R)-4-{3-[2-(4-Methyl-piperazin-1-yl)-ethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 2c)

15



Intermediate A (180 mg, 1.11 mmol) was added to a suspension of sodium hydride (60% in mineral oil, 133 mg, 3.33 mmol) in DMF (5 mL). The
20 reaction was stirred for 20 min, then Intermediate 2b (310 mg, 1.11 mmol) was added in DMF (5 mL) and the resulting mixture heated to 60°C for 3 h.

After cooling, the reaction was quenched by dropwise addition of MeOH. The solution was loaded onto an SCX-2 cartridge, which was washed with MeOH and product eluted with 2M NH₃ in MeOH. The residue was purified by FCC, using 2-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* (200 mg, 0.47 mmol, 43%). LCMS (Method 4): Rt 0.28 min, m/z 407 [MH⁺].


d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[2-(4-methyl-piperazin-1-yl)-ethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea formate salt (Example 2)

Intermediate 2c (85.0 mg, 0.21 mmol) was dissolved in 1,4-dioxane (2 mL) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 96.0 mg, 0.24 mmol) and DIPEA (75.0 μL, 0.43 mmol) were added. The reaction was heated to 60°C for 18 h. After cooling, the mixture was partitioned between EtOAc (50 mL) and water (50 mL), and extracted into EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, then further purified by HPLC (C18 X-select column, 10-40% MeCN in H₂O, 0.1% formic acid) to give the *title compound* mainly as formic acid salt (43.0 mg, 0.065 mmol, 31%). LCMS (Method 5) Rt 3.54 mins, m/z 662 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.90-2.05 (2H, m), 2.06-2.15 (1H, m), 2.20-2.30 (1H, m), 2.38 (3H, s), 2.40 (3H, s), 2.55-2.75 (8H, br s), 2.90-2.95 (2H, t, J 7.4), 3.29-3.32 (2H, t, J 7.4), 4.86-4.92 (1H, dd, J 8.8, 5.6), 5.43-5.47 (1H, t, J 4.2), 6.33 (1H, s), 7.21-7.35 (11H, m), 7.60-7.64 (1H, d, J 9.9), 8.07-8.09 (1H, d, J 1.9), 8.40-8.50 (0.4H, br s, formate).

Example 3

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-


1-yl]-urea



a. Piperidine-1-carboxylic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide


(Intermediate 3a)

5



1-Piperidine carbonyl chloride (348 mg, 0.30 mL, 2.36 mmol) was added dropwise to a solution of 5-fluoro-2-hydrazinyl-pyridine (see for reference WO2010022076; 0.30 g, 2.36 mmol) and DIPEA (1.2 mL, 7.08 mmol) in DCM (10 mL) at RT under nitrogen and the mixture stirred for 2 h. The solution was washed with water (2 x 15 mL) and dried (Na₂SO₄). The solvent was evaporated and the residue triturated (diethyl ether) and filtered to afford the *title compound* as an off-white solid (475 mg, 84%). LCMS (Method 1): Rt 1.82 min, m/z 239 [MH⁺].


b. 6-Fluoro-3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 3b)



Hexachloroethane (826 mg, 3.92 mmol) was added portionwise to a

solution of Intermediate 3a (466 mg, 1.95 mmol), triphenylphosphine (1.03g, 3.92 mmol) and triethylamine (1.1 mL, 7.83 mmol) in dry THF (30 mL) at RT, and the mixture stirred for 2 h. The resulting precipitate was filtered off and the filtrate evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH, to give the *title compound* as pale orange coloured gum (206 mg, 48%). LCMS (Method 1): Rt 2.44 min, m/z 221 [MH⁺].

c. (1S,4R)-4-(3-Piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-cis-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 3c)



10

Intermediate A (100 mg, 0.61 mmol) was added portionwise to a suspension of sodium hydride (60% in mineral oil, 73 mg, 1.84 mmol) in dry DMF (2 mL) at RT, and the mixture stirred for 15 min. Intermediate 3b (135 mg, 0.61 mmol) was then added in one portion and the mixture heated at 60°C for 3 h. After cooling, saturated NH₄Cl (*ca.* 0.2 mL) added. The mixture was then partitioned between water (15 mL) and ethyl acetate (3 x 15 mL) and the combined organic extracts washed with brine (2 x 15 mL) and dried (Na₂SO₄). The solvent was evaporated and the residue purified by SCX-2, eluting with MeOH followed by NH₃ in MeOH, to give the *title compound* as brown coloured gum (133 mg, 60%). LCMS (Method 1): Rt 1.95 min, m/z 364 [MH⁺].

20

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 3)

25

The *title compound* was prepared starting from Intermediate 3c and [5-


tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009) by using an analogous procedure to that described in Example 1 step d. LCMS (Method 5): Rt 4.79 min, m/z 916 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.32 (9H, s), 1.60-1.77 (6H, m), 1.86-1.95 (1H, m), 2.01-2.11 (2H, m), 2.20-2.27 (1H, m), 2.36 (3H, t), 3.15 (4H, m), 5.08 (1H, m), 5.18 (1H, m), 5.41 (1H, d, J 9.20), 6.28 (1H, s), 6.48 (1H br s), 6.96 (1H, dd, J 2.12, 9.75), 7.21 (2H, d, J 8.13), 7.24-7.33 (6H, m), 7.38-7.41 (2H, m), 7.45-7.48 (1H, d, J 9.65).

e. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Crystallization of Example 3)**

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (3.974g, Example 3) was dissolved in hot tert-butyl alcohol (~130ml) and then dried by lyophilisation overnight. The solid material was then slurried in iso-propyl acetate (120ml) in a maturation chamber which cycled between ambient and 50°C with four hours spent under each condition. After 3 days the reaction was cooled to RT and then stirred at RT for four days. The resulting off-white solid was isolated by filtration and dried at 40°C / 0.5mbar. Yield = 3.67g (92%). Mp = 253°C.


Example 4

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



5


a. 4-[N'-(5-fluoro-pyridin-2-yl)-hydrazinocarbonyl]-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 4a)



EDC (543 mg, 2.83 mmol) was added portionwise to a stirred solution of 5-fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 0.30 g, 2.36 mmol), piperidine-1,4-dicarboxylic acid mono-tert-butyl ester (Aldrich, 649 mg, 2.83 mmol) and HOBt (32 mg, 0.24 mmol) in dry DCM (20 mL). The mixture was stirred at RT for 18 h. The solution was washed with water (2 x 20 mL), dried (Na₂SO₄) and evaporated. The residue was triturated (diethyl ether) to give the *title compound* as an off-white solid (713 mg, 82%). LCMS (Method 1): Rt 2.76 min, m/z 339 [MH⁺].


20

b. 4-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 4b)



Hexachloroethane (990 mg, 4.18 mmol) was added portionwise to a solution of Intermediate 4a (707 mg, 2.09 mmol), triphenylphosphine (1.103g, 4.18 mmol) and triethylamine (1.2 mL, 8.36 mmol) in dry THF (30 mL) at RT. The mixture was stirred for 2 h. The resulting precipitate was filtered off and the filtrate evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH, to give a pale orange coloured solid. This was triturated (diethyl ether) to give the *title compound* as a buff-coloured solid (540 mg, 80%). LCMS (Method 1): Rt 2.79 min, m/z 321 and 221(-Boc) [MH⁺].

c. 4-[6-((1S,4R)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester. (Intermediate 4c)




Intermediate A (100 mg, 0.61 mmol) was added portionwise to a suspension of sodium hydride (60% in mineral oil, 73 mg, 1.84 mmol) in dry DMF (2 mL) at RT. The mixture was then stirred for 15 min. Intermediate 4b (193 mg, 0.61 mmol) was added in one portion and the mixture heated at

60°C for 3 h. After cooling, saturated NH₄Cl (*ca.* 0.2 mL) added. The mixture was partitioned between water (15 mL) and ethyl acetate (3 x 15 mL) and the combined organic extracts washed with brine (2 x 15 mL), dried (Na₂SO₄) and evaporated. The residue was purified by SCX-2, eluting with MeOH followed
5 by 2M NH₃ in MeOH, to give the *title compound* as brown coloured foam (261 mg, 92%). LCMS (Method 1): Rt 2.19 min, m/z 464 [MH⁺].

d. 4-(6-[(1S,4R)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy]-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester.

10 **(Intermediate 4d)**



The *title compound* was prepared starting from Intermediate 4c and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009) by using an
15 analogous procedure to that described in Example 1 step d. LCMS (Method 1): Rt 3.89 min, m/z 719 [MH⁺].


e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 4)

20 HCl (4M in dioxane, 1mL) was added dropwise to a solution of Intermediate 4d (272 mg, 0.37 mmol) in MeOH (1 mL) and the mixture stirred at RT for 5 h. The reaction mixture was passed down a SCX-2 cartridge, eluting with MeOH followed by 2M NH₃ in MeOH, to afford a brown foam.


Further purification by HPLC (50-95% MeCN in H₂O) gave the *title compound* as a off-white powder (37 mg). LCMS (Method 5): Rt 3.65 min, m/z 519 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.33 (9H, s), 1.87-1.97 (5H, m), 2.05-2.13 (2H, m), 2.22-2.29 (1H, m), 2.37 (3H, s), 2.71-2.81 (1H, m), 3.10 (1H, m), 3.19 (2H, m), 5.08 (1H, m), 5.22 (1H, m), 5.44 (1H, m), 6.29 (1H, s), 7.06 (1H, dd, J 2.13, 9.80), 7.23-7.32 (7H, m), 7.39-7.45 (3H, m), 7.63 (1H, d, J 9.75).

Example 5

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. (S)-1-Methyl-pyrrolidine-2-carboxylic acid [N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 5a)



EDC (271 mg, 1.41 mmol) was added portionwise to a solution of 5-fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 0.15 g, 1.18 mmol), N-methyl-L-proline monohydrate (0.20 g, 1.36 mmol) and HOBt (16 mg, 0.12 mmol) in dry DCM (5 mL) at RT and stirred for 16 h. The solution was diluted with DCM (15 mL), washed with water (150 mL), dried (Na₂SO₄) and evaporated to give the *title compound* as a pale yellow gum

(189 mg, 67%). LCMS (Method 1): Rt 0.31 min, m/z 239 [MH⁺].


b. 6-Fluoro-3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 5b)



5

Hexachloroethane (375 mg, 1.59 mmol) was added portionwise to a solution of Intermediate 5a (189 mg, 0.79 mmol), triphenylphosphine (416 mg, 1.59 mmol) and triethylamine (0.44 mL, 3.17 mmol) in dry THF (10 mL) at RT and stirred for 4 h. The resulting precipitate was filtered off and the filtrate evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH gave the *title compound* as a brown foam (136 mg, 78%). LCMS (Method 1): Rt 0.45 min, m/z 221 [MH⁺].

c. (1S,4R)-4-[3-((S)-1-Methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine.

15 **(Intermediate 5c)**

Intermediate A (128 mg, 0.77 mmol) was added portionwise to a suspension of sodium hydride (60% in mineral oil, 92 mg, 2.30 mmol) in dry DMF (3 mL) at RT and stirred for 15 mins. Intermediate 5b (169 mg, 0.77 mmol) was then added in one portion and the mixture heated at 60°C for 4 h. After cooling, saturated NH₄Cl (*ca.* 0.2 mL) was added. The mixture was partitioned between water (10 mL) and ethyl acetate (3 x 10 mL). The aqueous


phase was concentrated *in vacuo* and the residue purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH, to give the *title compound* as brown coloured foam (103 mg, 36%). LCMS (Method 1): Rt 1.34 min, m/z 364 [MH⁺].

5 **d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 5)**


The *title compound* was prepared with Intermediate 5c and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester
10 (Synthetic Communications, 2009, 39, 3999-4009) using an analogous procedure to that described in Example 1 step d. LCMS (Method 5): Rt 3.76 min, m/z 319 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.33 (9H, s), 1.88-2.12 (6H, m), 2.21 (3H, s), 2.21-2.30 (2H, m), 2.33-2.39 (1H, m), 2.37 (3H, s), 3.19-3.24 (1H, m), 3.98-4.02 (1H, m), 5.05-5.12 (1H, m), 5.17 (1H, t, J 4.0),
15 5.25 (1H, d, J 8.7), 6.25-6.27 (2H, m), 7.03 (1H, dd, J 2.0, 9.8), 7.22-7.34 (6H, m), 7.40 (2H, d, J 8.5), 7.61 (1H, d, J 9.8), 8.27 (1H, m).

Example 6

20 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((S)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea**




a. (S)-2-[N'-(5-fluoro-pyridin-2-yl)-hydrazinocarbonyl]-pyrrolidine-1-carboxylic acid tert-butyl ester (Intermediate 6a)




5 EDC (543 mg, 2.83 mmol) was added portionwise to a stirred solution of 5-fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 0.30 g, 2.36 mmol), N-tert-butylcarbonyl-L-proline (609 mg, 2.83 mmol) and HOBt (32 mg, 0.24 mmol) in dry DCM (15 mL) at RT and stirred for 18 h. The solution was washed with water (2 x 20 mL) and dried
10 (Na_2SO_4) and evaporated to give the *title compound* as a pale yellow foam (767 mg, 100%). LCMS (Method 1): Rt 2.69 min, m/z 325 [MH^+].

b. (S)-2-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester (Intermediate 6b)




15 Hexachloroethane (1.34 g, 5.65 mmol) was added portionwise to a solution of Intermediate 6a (917 mg, 2.83 mmol), triphenylphosphine (1.48 g, 5.65 mmol) and triethylamine (1.6 mL, 11.3 mmol) in dry THF (15 mL) at RT and stirred for 4 h. The resulting precipitate was filtered off and the filtrate
20 evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH_3 in MeOH, to give the *title compound* as a buff-coloured foam (669 mg, 77%). LCMS (Method 4): Rt 2.48 min, m/z 307 [MH^+].

c. (S)-2--[6-((1S,4R)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidine-1-carboxylic acid

tert-butyl ester. (Intermediate 6c)

Intermediate A (150 mg, 0.92 mmol) was added portionwise to a
 5 suspension of sodium hydride (60% in mineral oil, 110 mg, 2.76 mmol) in dry
 DMF (2 mL) at RT and stirred for 15 mins. Intermediate 6b (281 mg, 0.92
 mmol) was then added in one portion and the mixture heated at 60°C for 3 h.
 After cooling, saturated NH₄Cl (*ca.* 0.2 mL) was added. The mixture was
 partitioned between water (15 mL) and ethyl acetate (3 x 15 mL). The
 10 combined organic extracts were washed with brine (20 mL), dried (Na₂SO₄)
 and evaporated to give the *title compound* as brown coloured gum (74 mg,
 18%). LCMS (Method 1): Rt 2.08 min, m/z 450 [MH⁺].

**d. (S)-2-(6-[(1S,4R)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-
 ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy]-[1,2,4]triazolo[4,3-
 15 a]pyridin-3-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester.
 (Intermediate 6d)**



The *title compound* was prepared starting from Intermediate 6c and
 20 [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl
 ester (Synthetic Communications, 2009, 39, 3999-4009) by using an


analogous procedure to that described in Example 1 step d. LCMS (Method 1): Rt 3.88 min, m/z 705 [MH⁺].

e. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((S)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 6)**


HCl (4M in dioxane; 1 mL) was added to a solution of Intermediate 6d (225 mg, 0.32mmol) in MeOH (0.5 mL) at RT and stirred for 8 h. The solution was loaded on to a SCX-2 (10g) cartridge, eluting with MeOH followed by 2M NH₃ in MeOH, to give a brown foam. Further purification by HPLC (50-10 85% MeCN in H₂O (0.1%NH₃)) gave the *title compound* as an off-white powder after freeze drying (7 mg, 4%). LCMS (Method 5): Rt 3.73 min, m/z 605 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.29 (9H, s), 1.84-2.07 (5H, m), 2.18-2.31 (3H, m), 2.35 (3H, s), 2.96-3.07 (2H, m), 4.64 (1H, t, J 7.5), 5.01-5.08 (1H, m), 5.19 (1H, t, J 4.0), 5.27 (1H, d, J 8.8), 6.24 (2H, s), 7.02 (1H, 15 dd, J 2.2, 10.0), 7.19-7.28 (6H, m), 7.37 (2H, d, J 8.4), 7.58 (1H, d, J 10.0), 7.98 (1H, m).

Example 7

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperazin-1-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea




a. **4-[N'-(5-fluoro-pyridin-2-yl)-hydrazinocarbonylmethyl]-piperazine-1-carboxylic acid tert-butyl ester (Intermediate 7a)**




5 EDC (543 mg, 2.83 mmol) was added portionwise to a solution of 2-(4-Boc-1-piperazinyl)acetic acid (576 mg, 2.36 mmol), 5-fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 0.30 g, 2.36 mmol) and HOBt (32 mg, 0.24 mmol) in dry DCM (15 mL) at RT and stirred for 16 h. The solution was diluted with DCM (20 mL), washed with water (2 x 20 mL)
10 and dried (Na₂SO₄). The solvent was evaporated to give the title compound as an off-white solid (667 mg, 80%). LCMS (Method 1): Rt 1.99 min, m/z 354 [MH⁺].

b. **4-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-piperazine-1-carboxylic acid tert-butyl ester (Intermediate 7b)**



15 Hexachloroethane (893 mg, 3.77 mmol) was added portionwise to a solution of Intermediate 7a (667 mg, 1.89 mmol), triphenylphosphine (990 mg, 3.77 mmol) and triethylamine (1.05 mL, 7.55 mmol) in dry THF (20 mL) at RT and stirred for 4 h. The resulting precipitate was filtered off and the filtrate evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH, to give an off-white solid. This was triturated (diethyl ether) to give the *title compound* as a colourless solid LCMS (Method
20 1): Rt 2.33 min, m/z 336 [MH⁺].


c. 4-[6-((1S,4R)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-piperazine-1-carboxylic acid tert-butyl ester. (Intermediate 7c)



5

The *title compound* was prepared with Intermediate 7b and Intermediate A using an analogous procedure to that described in Example 6 step c. LCMS (Method 1): Rt 1.98 min, m/z 479 [MH⁺].

d. 4-(6-{(1S,4R)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy}-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-piperazine-1-carboxylic acid tert-butyl ester. (Intermediate 7d)



15 The *title compound* was prepared starting from Intermediate 7c and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009) by using an analogous procedure to that described in Example 1 step d. LCMS (Method 1): Rt 3.59 min, m/z 734 [MH⁺].


20 e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperazin-1-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-

tetrahydro-naphthalen-1-yl]-urea (Example 7)

HCl (4M in dioxane; 1 mL) was added to a solution of Intermediate 7d (92 mg, 0.12mmol) in MeOH (1 mL) at Rt and stirred for 4.5 h. The solution was loaded on to a SCX-2 cartridge, eluting with MeOH followed by 2M NH₃ in MeOH, to give a pale yellow foam. Further purification by HPLC (40-80% MeCN in H₂O (0.1%NH₃)) gave the *title compound* as an off-white powder after freeze drying (28 mg, 35%). LCMS (Method 5): Rt 3.59 min, m/z 634 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.33 (9H, s), 1.89-1.98 (1H, m), 2.06-2.14 (2H, m), 2.25-2.32 (1H, m), 2.38 (3H, s), 2.38-2.48 (4H, m), 2.80-2.83 (4H, m), 4.03 (2H, s), 5.06-5.12 (1H, m), 5.20-5.24 (2H, m), 6.28 (1H, s), 6.34 (1H, br s), 7.08 (1H, dd, J 9.8, 2.0), 7.23-7.36 (6H, m), 7.40 (2H, d, J 7.7), 7.63 (1H, d, J 9.5), 8.14 (1H, m).


Example 8

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-isopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



a. 2-(5-Fluoropyridin-2-yl)-N-(propan-2-yl)hydrazinecarboxamide (Intermediate 8a)


20



Isopropyl isocyanate (0.25 mL, 2.60 mmol) was added dropwise, over 2


min, to a solution of 5-fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 0.30 g, 2.36 mmol) in dry DCM (10 mL) at RT and stirred for 3 h. The solvent was evaporated and the residue triturated (diethyl ether) and filtered to give the *title compound* as a colourless solid (464 mg, 93%). LCMS (Method 1): Rt 1.92 min, m/z 213 [MH⁺].

b. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-isopropyl-amine. (Intermediate 8b)



Hexachloroethane (1.77 g, 7.75 mmol) was added portionwise to a solution of Intermediate 8a 822 mg, 3.87 mmol), triphenylphosphine (2.03 g, 7.75 mmol) and triethylamine (2.2 mL, 15.49 mmol) in dry THF (15 mL) at RT and stirred for 16 h. The resulting precipitate was filtered off and the filtrate evaporated. The residue was purified by SCX-2, eluting with MeOH followed by 2M NH₃ in MeOH, to give the *title compound* as a buff-coloured foam (618 mg, 82%). LCMS (Method 1): Rt 1.55 min, m/z 195 [MH⁺].

c. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-isopropyl-carbamic acid tert-butyl ester. (Intermediate 8c)




Di-tert-butyl dicarbonate (472 mg, 2.16 mmol) in dry DCM (1 mL) was added to a solution of Intermediate 8b (168 mg, 0.86 mmol) in DCM (2 mL). 4-(1-Pyrrolidinyl)pyridine (*ca.* 5 mg) was then added at RT and stirred for 5 h.

The solvent was evaporated and the residue purified by SCX-2, eluting with MeOH then 2M NH₃ in MeOH, affording a dark coloured foam. Further purification by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, afforded the *title compound* (112 mg). LCMS (Method 4): Rt 2.90 min, m/z 295 [MH⁺] /

5 589 [2MH⁺]


d. [6-((1S,4R)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-isopropyl-carbamic acid tert-butyl ester (Intermediate 8d)



10 Sodium hydride (60% in mineral oil, 45 mg, 1.14 mmol) was added portionwise to a solution of Intermediate A (62 mg, 0.38 mmol) in dry DMF (1 mL) at RT and stirred for 15 min. Intermediate 8c (112 mg, 0.38 mmol) was added and the mixture heated at 60°C for 4 h. After cooling, the mixture was partitioned between water (10 mL) and EtOAc (3 x 10 mL) and the combined

15 organic extracts washed with brine (2 x 15 mL), dried (Na₂SO₄) and evaporated. The residue was purified by SCX-2, eluting with MeOH then 2M NH₃ in MeOH, affording a dark gum. Further purification by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, afforded the *title compound* (75 mg, 45%). LCMS (Method 1): Rt 2.20 min, m/z 438 [MH⁺].

e. (6-((1S,4R)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-isopropyl-carbamic acid tert-butyl ester. (Intermediate 8e)



5

The *title compound* was prepared starting from Intermediate 8d and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009) by using an analogous procedure to that described in Example 1 step d. LCMS (Method 4): Rt 3.98 min, m/z 693 [MH⁺].

10

f. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-isopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 8)

HCl (4.0 M in dioxane; 1 mL) was added to a solution of Intermediate 8e (110 mg, 0.16 mmol) in MeOH (1 mL) at RT and stirred for 5 h. The solution was loaded on to a SCX-2 cartridge, eluting with MeOH then 2M NH₃ in MeOH, to afford a dark gum. Further purification by HPLC (eluting with 50-90% MeCN in H₂O (0.1%NH₃)) gave the *title compound* as a colourless powder after freeze drying (10 mg, 11%). LCMS (Method 5): Rt 4.15 min, m/z 593 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.21-1.27 (15H, m), 1.87 (1H, m), 2.00 (2H, m), 2.10 (1H, m), 2.31 (3H, s), 3.85 (1H, br s), 3.94 (1H, m), 5.02 (1H, m), 5.11 (1H, t, J 4.13), 5.59 (1H, d), 6.24 (1H, s), 6.54 (1H, s), 6.84 (1H, dd, J 9.93, 2.05), 7.14 (3H, d, J 8.13), 7.20-7.26 (3H, m),


20

7.30-7.34 (3H, m).


Example 9

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(6-cyano-pyridin-3-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

5



a. 5-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-pyridine-2-carbonitrile (Intermediate 9a)



10


Intermediate A (300 mg, 1.84 mmol) was added to an ice cold stirred suspension of sodium hydride (60% in mineral oil, 221 mg, 5.52 mmol) in DMF (15 mL) and stirred for 15 min. 2-Cyano-5-fluoropyridine (224 mg, 1.84 mmol) was added and the reaction warmed to RT. After 90 min, the reaction was quenched by dropwise addition of water and the mixture partitioned between EtOAc (75 mL) and water (150 mL). The aqueous layer was then extracted with EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give the *title compound* (255 mg, 0.96 mmol, 52%). LCMS (Method 4): Rt 0.28, 1.73, m/z 266.1 [MH⁺].

b. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(6-cyano-pyridin-3-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea. (Example 9)

The *title compound* was prepared starting from Intermediate 9a and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009) using an analogous procedure to that described in Example 1 step d. LCMS (Method 5): Rt 5.25 mins, m/z 521 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.29 (9H, s), 1.76-1.88 (1H, m), 2.00-2.20 (3H, m), 2.36 (3H, s), 4.94-4.98 (1H, d, J 8.8), 5.00-5.08 (1H, m), 5.38-5.42 (1H, t, J 3.6), 6.03 (1H, s), 6.22 (1H, s), 7.16-7.20 (1H, m), 7.21-7.26 (3H, m), 7.26-7.32 (3H, m), 7.34-7.38 (2H, d, J 8.3), 7.63-7.66 (1H, d, J 8.8), 8.36-8.38 (1H, d, J 2.7).


Example 10

N-(4-{(1S,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy}-pyridin-2-yl)-2-methoxy-acetamide



a. ((1S,4S)-4-Hydroxy-1,2,3,4-tetrahydro-naphthalen-1-yl)-carbamic acid tert-butyl ester (Intermediate 10a)


20



Intermediate B (1.62 g, 9.94 mmol) was suspended in acetonitrile


(35 mL) then di-tert-butyl-dicarbonate (2.40 g, 11 mmol) was added. Mixture stirred at RT for 16 h. Some insoluble material was still present, so mixture filtered through Celite, washing with DCM. Filtrate was evaporated to give an off-white solid and then purified by FCC, eluting with 0-80% ethyl acetate in cyclohexane, to give the *title compound* as a white solid (2.38 g, 91%). ¹H NMR (300 MHz, d₆-DMSO): 1.42 (9H, s), 1.54-1.70 (2H, m), 1.92-2.18 (2H, m), 4.46-4.73 (2H, m), 5.17 (1H, d, J 6.3), 7.10-7.25 (4H, m), 7.39-7.46 (1H, m).

b. [(1S,4S)-4-(2-Chloro-pyridin-4-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-carbamic acid tert-butyl ester (Intermediate 10b)




Sodium hydride (60% in mineral oil, 0.32 g, 8.00 mmol) was suspended in dry DMF (15 mL) under argon. To this was added Intermediate 10a (1.05 g, 4.00 mmol) followed by 2-chloro-4-nitropyridine (0.64 g, 4.02 mmol). The dark coloured mixture was stirred at RT under argon for 30 min. The reaction mixture was diluted with water and extracted with DCM (3 x 30 mL). The combined organic layers were washed with brine, dried (Na₂SO₄), filtered, concentrated *in vacuo*. Purification by FCC, eluting with 0-70% ethyl acetate in cyclohexane, gave the *title compound* as a white foam (1.43 g, 95%). LCMS (Method 3): Rt 4.22 min, m/z 373.1 [MH⁺].

c. **{(1S,4S)-4-[2-(2-Methoxy-acetyl-amino)-pyridin-4-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-carbamic acid tert-butyl ester (Intermediate 10c)**



5 Intermediate 10b (562 mg, 1.50 mmol), 2-methoxyacetamide (334 mg, 3.75 mmol), Xantphos (173 mg, 0.30 mmol) and potassium carbonate (518 mg, 3.75 mmol) were suspended in 1,4-dioxane (20 mL), warmed, degassed, and placed under argon. To this mixture palladium acetate (34.0 mg, 0.15 mmol) was added. The mixture was degassed and then heated at reflux
10 for 17 h. After cooling, the mixture was filtered through Celite, washing with DCM, and evaporated to a yellow gum. Purification by FCC, eluting with 0-100% ethyl acetate in cyclohexane, gave the *title compound* as a white foam (281 mg, 44%). LCMS (Method 3): Rt 3.27 min, m/z 450.2 [MNa⁺].

d. **N-[4-((1S,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-pyridin-2-yl]-2-methoxy-acetamide (Intermediate 10d)**



To a solution of Intermediate 10c (272 mg, 0.64 mmol) in DCM (6 mL) was added TFA (2 mL), and the mixture stirred at RT for 30 min. The mixture
20 was concentrated *in vacuo*. The residue was purified on an Isolute SCX-2 cartridge, eluting with MeOH then 0.4-1M NH₃ in MeOH, to give the *title*


compound as a colourless gum (208 mg, 100%). LCMS (Method 3): Rt 0.44 min, m/z 350.2 [MNa⁺].

e. N-(4-((1S,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy}-pyridin-2-yl)-2-methoxy-acetamide (Example 10)


To a solution of Intermediate 10d (204 mg, 0.62 mmol) and DIPEA (0.127 mL, 0.80 mmol) in 1,4-dioxane (6 mL), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 315 mg, 0.78 mmol) was added. The mixture was heated at 80°C under argon for 17 h, and then evaporated to dryness. The residue was purified by FCC, eluting with 0-100% ethyl acetate in cyclohexane, to give the slightly impure product (0.262 g). Further purification by HPLC (Method 6) gave the *title compound* as a white solid (102 mg, 28%). LCMS (Method 5): Rt 4.51 min, m/z 583 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.26 (9H, s), 1.69-1.81 (1H, m), 1.90-2.24 (3H, m), 2.36 (3H, s), 3.36 (3H, s), 4.05 (2H, s), 4.86-4.94 (1H, m), 5.62-5.67 (1H, m), 6.32 (1H, s), 6.91 (1H, dd, J 2.5, 5.9), 7.03 (1H, d, J 8.4), 7.26-7.37 (8H, m), 7.60 (1H, br d, J 2.0), 8.01 (1H, s), 8.16 (1H, d, J 5.7), 9.90 (1H, s).

Example 11

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[1-(2-hydroxy-ethyl)-1H-indazol-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. **6-Fluoro-1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-indazole**
(Intermediate 11a)



5 A mixture of 6-fluoro-1H-indazole (2.0 g, 14.7 mmol), and 2-(2-bromo-ethoxy)-tetrahydro-pyran (3.38 g, 16.2 mmol) in DMF (25 mL) was treated with caesium carbonate (6.1 g, 18.7 mmol) and stirred at RT for 18 h. The solvent volume was reduced *in vacuo*, and the residue partitioned between EtOAc (100 mL) and water (100 mL). The aqueous layer was extracted into
10 EtOAc (3 x). The combined organic layers were washed with saturated aqueous sodium chloride solution, dried (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-50% cyclohexane in EtOAc, to afford the *title product* and a yellow oil. LCMS (Method 1): Rt 3.42 min, m/z 181 [MH⁺] (M-THP). ¹H NMR (300 MHz; CDCl₃) 1.52-1.56 (6 H, m), 3.39-3.49 (1 H, m), 3.57-3.68 (1 H, m), 3.89-3.91 (1 H, m), 4.19-4.21 (1 H, m),
15 4.56-4.57 (3 H, m), 6.88 (1 H, td, J 9.15 and 2.26), 7.30 (1 H, m), 7.62 (1 H, m), 8.04 (1 H, s).


b. **2-[6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-indazol-1-yl]-ethanol** (Intermediate 11b)



Intermediate A (212 mg, 1.3 mmol) was added portionwise to a suspension of sodium hydride (60% in mineral oil, 120 mg, 3.0 mmol) in DMF (3 mL) at RT and stirred for 20 min. A solution of Intermediate 11a (264.3 mg, 1.0 mmol) in DMF (1 mL) was then added dropwise and the
5 resulting mixture stirred at 60°C for 2.5 h. After cooling, the mixture was diluted with EtOAc (50 mL) and poured onto ice water. The organic layer was washed with saturated aqueous sodium chloride solution, dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was loaded on to a SCX-2 cartridge, eluting with MeOH then 2M NH₃ in MeOH. Further purification by
10 FCC, using 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* as a brown oil. LCMS (Method 1): Rt 1.80 min, m/z 323 [MH⁺].


c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[1-(2-hydroxy-ethyl)-1H-indazol-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 11)

15 The *title compound* was prepared starting from Intermediate 11b and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 315 mg, 0.78 mmol) by using an analogous procedure to that described in Example 1 step d. LCMS (Method 5): Rt 4.99 min, m/z 578 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO):
20 1.22 (9 H, s), 1.81-1.92 (2 H, m), 2.02-2.06 (2 H, m), 2.31 (3 H, s), 3.74-3.75 (2 H, m), 4.34 (2 H, t, J 5.75), 4.78 (2 H, m), 5.53 (1 H, t, J 4.8), 6.28 (1 H, s), 6.74 (1 H, dd, J 8.75 and 2.03), 7.08 (1 H, d, J 8.53), 7.27-7.28 (9 H, m), 7.57 (1 H, d, J 8.75), 7.89 (1 H, s), 7.96 (1 H, s).

Example 12

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-((R)-3-pyrrolidin-2-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea

The *title compound* was prepared starting from N-tert-butylcarbonyl-D-proline using analogous procedures to those described in Example 6. LCMS (Method 5): Rt 3.69 min, m/z 605 [MH⁺]; ¹H NMR (400 MHz, CDCl₃): 1.33 (9H, s), 1.86-2.39 (8H, m), 2.36 (3H, s), 2.96-3.08 (2H, m), 4.51 (1H, t, J 7.4 Hz), 5.04-5.10 (1H, m), 5.24 (1H, t, J 4.0 Hz), 5.52 (1H, d, J 8.1 Hz), 6.30 (1H, s), 6.57 (1H, br s), 7.03 (1H, dd, J 2.2, 10.0 Hz), 7.20 (2H, d, J 7.9 Hz), 7.23-7.34 (4H, m), 7.38 (2H, d, J 7.4 Hz), 7.56 (1H, d, J 9.8 Hz), 7.92 (1H, m).

Example 13


N-(4-((1R,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydronaphthalen-1-yloxy)-pyridin-2-yl)-2-methoxyacetamide

The *title compound* was prepared starting from Intermediate A by using analogous procedures to those described in Example 10. LCMS (Method 5):


Rt 4.54 min, m/z 583 [MH⁺]; ¹H NMR (400 MHz, d₆-DMSO): 1.21 (9H, s), 1.66-2.11 (4H, m), 2.31 (3H, s), 3.31 (3H, s), 4.0 (2H, s), 4.72-4.81 (1H, m), 5.49-5.58 (1H, m), 6.26 (1H, s), 6.86 (1H, dd, J 5.8, 2.2 Hz), 7.07 (1H, d, J 8.5 Hz), 7.19-7.33 (8H, m), 7.68 (1H, d, J 2.2 Hz), 7.96 (1H, s), 8.12 (1H, d, J 5.8 Hz), 9.86 (1H, br s).

Example 14

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




a. (4-Methyl-piperazin-1-yl)-acetic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 14a)



5-Fluoro-2-hydrazinyl-pyridine (for reference procedure see WO2010022076; 500 mg, 3.94 mmol) and 4-methyl-1-piperazin-1-yl acetic acid (684 mg, 4.33 mmol) were dissolved in DMF (10.0 mL). EDC (831 mg, 4.33 mmol) and HOBt (53.0 mg, 0.39 mmol) were added and the reaction stirred for 18 h. The mixture was loaded onto an SCX-2 cartridge, which was washed with MeOH then eluted with 2M NH₃ in MeOH. The resulting residue was purified by FCC, using 4-20% [2M NH₃ in MeOH] in DCM, to give the


title compound (570 mg, 54%). ¹H NMR (400 MHz, CDCl₃): 2.32 (3H, s), 2.47-2.56 (4H, br s), 2.62-2.71 (4H, br s), 3.18 (2H, s), 6.56-6.61 (1H, br s), 6.60-6.66 (1H, dd, J 8.9, 3.4 Hz), 7.26-7.32 (1H, m), 8.02-8.05 (1H, d, J 2.9 Hz), 8.93 (1H, br s).

5 **b. 6-Fluoro-3-(4-methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 14b)**



Intermediate 14a (570 mg, 2.13 mmol) was dissolved in THF (20.0 mL) and cooled in an ice/water bath. Triphenylphosphine (1.12 g, 4.27 mmol) was added followed by triethylamine (1.19 mL, 8.54 mmol) and hexachloroethane (1.01 g, 4.27 mmol). The reaction was stirred for 18 h. The mixture was loaded onto an SCX-2 cartridge, washing with MeOH and eluting with 2M NH₃ in MeOH. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (470 mg, 1.89 mmol, 89%). LCMS (Method 1): Rt 0.36 min, m/z 250 [MH⁺].

c. (1S,4R)-4-[3-(4-Methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 14c)




20 Intermediate A (200 mg, 1.22 mmol) was added to a suspension of sodium hydride (60% in mineral oil, 146 mg, 3.66 mmol) in DMF (6.0 mL) and stirred for 20 min. Intermediate 14b (305 mg, 1.22 mmol) was added and the reaction heated to 60°C for 90 min. The mixture was cooled and quenched by dropwise addition of MeOH. The solution was diluted with MeOH and

loaded onto an SCX-2 cartridge, washing with MeOH and eluting with 2M NH₃ in MeOH. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* (240 mg, 0.61 mmol, 50%). LCMS (Method 4): Rt 0.29 min, m/z 393 [MH⁺].

5 **d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-piperazin-1-yl)methyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 14)**

Intermediate 14c (120 mg, 0.31 mmol) was dissolved in 1,4-dioxane (2.0 mL) and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 124
10 mg, 0.31 mmol) and diisopropylethylamine (106 µL, 0.61 mmol) were added. The reaction was heated to 60°C for 20 h. After cooling, the mixture was partitioned between EtOAc (50 mL) and water (50 mL), and extracted into EtOAc (3 x). The combined organic layers were washed with brine, dried
15 (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, then further purified by HPLC (C18 X-select column, 20-70% MeCN in H₂O, 0.1% formic acid) to give the *title compound* as the formic acid salt (103 mg, 0.16 mmol, 52%). LCMS (Method 5): Rt 3.63 min, m/z = 648 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.90-1.99 (1H, m), 1.99-2.07 (1H, m), 2.10-2.19 (1H, m), 2.21-2.30 (1H, m), 2.37 (3H, s), 2.38 (3H, s), 2.52-2.67 (8H, br s), 4.05-4.09 (1H, d, J 14.3 Hz), 4.11-4.15 (1H, d, J 14.3 Hz), 4.88-4.93 (1H, dd, J 8.9, 5.6 Hz), 5.41-5.44 (1H, t, J 4.3 Hz), 6.33 (1H, s), 7.22-7.36 (11H, m), 7.63-7.67 (1H, d, J 10.0 Hz), 8.21-8.22 (1H, d, J 1.9 Hz), 8.44-8.45 (0.25H, s).


Example 15

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-morpholin-4-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

The *title compound* was prepared starting from 1-morpholinoacetic acid by using analogous procedures to those described for Example 14. LCMS (Method 5): Rt 4.02 min, $m/z = 635$ [MH^+]. ^1H NMR (400 MHz, d_4 -MeOD): 1.30 (9H, s), 1.90-1.97 (1H, m), 1.97-2.07 (1H, m), 2.10-2.19 (1H, m), 2.23-2.31 (1H, m), 2.38 (3H, s), 2.42-2.53 (4H, m), 3.60-3.64 (4H, m), 4.01-4.05 (1H, d, J 14.3 Hz), 4.08-4.12 (1H, d, J 14.3 Hz), 4.88-4.93 (1H, dd, J 9.0, 5.6 Hz), 5.42-5.45 (1H, t, J 4.3 Hz), 6.33 (1H, s), 7.22-7.36 (11H, m), 7.63-7.66 (1H, d, J 10.0 Hz), 8.24-8.26 (1H, d, J 2.0 Hz), 8.48-8.52 (0.25H, s).

Example 16

15



1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-pyrrolidin-1-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea


20

The *title compound* was prepared starting from 1-pyrrolidin-1-yl-acetic


acid hydrochloride (for reference procedure see US5756533) by using analogous procedures to those described for Example 14. LCMS (Method 5): Rt 3.72 min, $m/z = 619$ $[MH^+]$. ^1H NMR (400 MHz, d_4 -MeOD): 1.30 (9H, s), 1.80-1.85 (4H, m), 1.90-2.05 (2H, m), 2.07-2.15 (1H, m), 2.21-2.30 (1H, m), 2.38 (3H, s), 2.65-2.70 (4H, m), 4.21-4.26 (1H, d, J 14.3 Hz), 4.26-4.31 (1H, d, J 14.3 Hz), 4.87-4.92 (1H, dd, J 8.9, 5.6 Hz), 5.40-5.43 (1H, t, J 4.3 Hz), 6.33 (1H, s), 7.21-7.36 (11H, m), 7.63-7.67 (1H, d, J 9.9 Hz), 8.23-8.24 (1H, d, J 1.9 Hz), 8.29-8.31 (0.3H, br s).

Example 17

10 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[6-(morpholine-4-carbonyl)-pyridin-3-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**




15 **a. (5-Fluoro-pyridin-2-yl)-morpholin-4-yl-methanone (Intermediate 17a)**



20 2-Cyano-5-fluoropyridine (1.00 g, 8.19 mmol) was dissolved in hydrochloric acid (37% aqueous, 1.0 mL) and heated to 60°C for 18 h and then evaporated *in vacuo*. The residue was suspended in DMF (40.0 mL) and EDC (1.89 g, 9.83 mmol), HOBt (111 mg, 0.82 mmol), morpholine (787 μ L, 9.00 mmol) and triethylamine (1.25 mL, 9.00 mmol) were added and the

reaction stirred for 18 h. The reaction was partitioned between water (200 mL) and EtOAc (100 mL), and extracted into EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-8% [2M NH₃ in MeOH] in DCM, to give the *title compound* (350 mg), contaminated with a single impurity (~10% by NMR integration). The product was used in the next step without further purification. LCMS (Method 1): Rt 1.88 min, m/z 211 [MH⁺].

b. [5-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-pyridin-2-yl]-morpholin-4-yl-methanone (Intermediate 17b)



Intermediate A (150 mg, 0.92 mmol) was added to a suspension of sodium hydride (60% in mineral oil, 147 mg, 3.66 mmol) in DMF (3.0 mL). The reaction was stirred for 20 min, then Intermediate 17a (280 mg) in DMF (3.0 mL) was added and the reaction heated to 60°C for 90 min. The mixture was cooled and quenched by dropwise addition of MeOH. The solution was diluted with MeOH and loaded onto an SCX-2 cartridge, which was washed sequentially with MeOH and 2M NH₃ in MeOH. The basic fractions were evaporated *in vacuo* then purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (234 mg, 0.66 mmol, 72%). LCMS (Method 4): Rt 1.60 min, m/z 354 [MH⁺].

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[6-(morpholine-4-carbonyl)-pyridin-3-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 17)


Intermediate 17b (234 mg, 0.66 mmol) was dissolved in 1,4-dioxane (7.0 mL) and [5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-

trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 268 mg, 0.66 mmol) and diisopropylethylamine (229 μ L, 1.32 mmol) were added. The reaction mixture was heated to 60°C for 20 h, then cooled, partitioned between EtOAc (50 mL) and water (50 mL), and extracted into EtOAc (3 x).

- 5 The combined organic layers were washed with brine, dried (MgSO_4) and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH_3 in MeOH] in DCM, to give the *title compound* (120 mg, 0.20 mmol, 30%). LCMS (Method 5): Rt 4.76 min, m/z = 609 [MH^+]. ^1H NMR (400 MHz, d_4 -MeOD): 1.30 (9H, s), 1.85-2.05 (2H, m), 2.05-2.18 (2H, m), 2.38 (3H, s),
 10 3.57-3.76 (8H, br m), 4.86-4.92 (1H, dd, J 8.6, 5.5 Hz), 5.54-5.57 (1H, t, J 4.3 Hz), 6.33 (1H, s), 7.20-7.34 (9H, m), 7.58-7.65 (2H, m), 8.24-8.26 (1H, dd, J 2.4, 0.8 Hz).


Example 18

- 15 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea**



- a. **4-Morpholin-4-ylmethyl-benzoic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 18a)**


20



HOBt (53 mg, 0.39 mmol) was added to (5-fluoro-pyridin-2-yl)-

hydrazine (For reference procedure see WO2010022076; 500 mg, 3.94 mmol), 4-morpholin-4-ylmethyl-benzoic acid (1.04 g, 4.72 mmol) and EDC (907 mg, 4.72 mmol) in DCM (5.0 mL) and the reaction stirred for 4 h. The reaction was partitioned between DCM (75 mL) and saturated aqueous NaHCO₃ (75 mL) and the aqueous layer extracted with DCM (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* then purified by FCC using [0.5-7.5% 2M NH₃ in MeOH] in DCM to give the *title compound* (980 mg, 2.97 mmol, 75%). LCMS (Method 4): Rt 0.27, m/z 331 [MH⁺].

b. 6-Fluoro-3-(4-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 18b)




To an ice cold solution of Intermediate 18a (980 mg, 2.97 mmol) in THF (15 mL) was added triphenylphosphine (1.56 g, 5.94 mmol), triethylamine (1.65 mL, 11.9 mmol) and hexachloroethane (1.40 g, 5.94 mmol). The reaction was stirred for 90 min then partitioned between EtOAc (75 mL) and water (75 mL) and the aqueous layer extracted with EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* then purified by SCX-2, washing with MeOH and eluting with 2M NH₃ in MeOH to give the *title compound* (640 mg, 2.05 mmol, 69%). LCMS (Method 2): Rt 0.28, 1.32, m/z 313 [MH⁺].

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(4-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-


1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 18)

The *title compound* was prepared starting from Intermediate 18b using analogous procedures to those described for Example 14. LCMS (Method 5): Rt 3.79 mins, $m/z = 711$ $[MH^+]$. ^1H NMR (400 MHz, $CDCl_3$): 1.29 (9H, s),
5 1.80-1.90 (1H, m), 1.96-2.08 (2H, m), 2.16-2.24 (1H, m), 2.33 (3H, s), 2.44-2.49 (4H, t, J 4.2 Hz), 3.56 (2H, s), 3.68-3.71 (4H, t, J 4.6 Hz), 5.00-5.06 (1H, td, J 8.8, 5.2 Hz), 5.11-5.17 (2H, m), 6.22 (2H, s), 7.06-7.10 (1H, dd, J 9.9, 2.1 Hz), 7.17-7.22 (3H, m), 7.24-7.30 (3H, m), 7.33-7.36 (2H, d, J 8.4 Hz), 7.49-7.53 (2H, d, J 8.1 Hz), 8.68-7.71 (2H, d, J 8.2 Hz), 7.71 (1H, s), 7.81-7.82 (1H, d, J 1.6 Hz).

Example 19

**1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(3-
15 morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-
1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**


The *title compound* was prepared starting from 3-morpholin-4-ylmethyl-benzoic acid by using analogous procedures to those described for Example 18. LCMS (Method 5): Rt 3.81 mins, m/z 711.3 $[MH^+]$; ^1H NMR (400 MHz, d_4 -MeOD): 1.29 (9H, s), 1.89-2.10 (3H, m), 2.20-2.28 (1H, m), 2.37 (3H, s), 2.45-2.50 (4H, t, J 4.3 Hz), 3.61 (2H, s), 3.61-3.65 (4H, t, J 4.6 Hz), 4.84-4.88 (1H, dd, J 8.9, 5.7 Hz), 5.36-5.39 (1H, t, J 4.1 Hz), 6.31 (1H, s), 7.18-7.38 (9H, m), 7.53-7.59 (2H, m), 7.70-7.74 (1H, dt, J 6.4, 2.3 Hz), 7.72-7.75 (1H, d, J 9.9 Hz), 7.80 (1H, s), 8.02-8.04 (1H, d, J 1.7 Hz).

Example 20

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(2-morpholin-4-ylmethyl-phenyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

The *title compound* was prepared starting from 2-morpholin-4-ylmethyl-benzoic acid by using analogous procedures to those described for Example 18. LCMS (Method 5): Rt 4.00 mins, m/z 711.4 [MH⁺]; ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.82-2.02 (3H, m), 2.04-2.18 (5H, m), 2.38 (3H, s), 3.08-3.16 (4H, br s), 3.50-3.54 (1H, d, J 13.5 Hz), 3.54-3.58 (1H, d, J 13.5 Hz), 4.81-4.86 (1H, dd, J 8.8, 5.6 Hz), 5.29-5.33 (1H, t, J 4.0 Hz), 6.31 (1H, s), 7.17-7.35 (9H, m), 7.35-7.39 (1H, dd, J 9.9, 2.0 Hz), 7.49-7.59 (6H, m), 7.74-7.78 (1H, d, J 9.9 Hz).

Example 21




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(6-morpholin-4-ylmethyl-pyridin-3-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

Borane dimethylsulfide complex (2M in THF, 99 μL, 0.20 mmol) was added to a solution of Example 17 (60.0 mg, 0.099 mmol) in THF (3.00 mL). The reaction stirred for 20 min then heated to 60°C overnight. The reaction was cooled


and further borane dimethylsulfide complex (2M in THF, 99 μ L, 0.20 mmol) added. After stirring for 20 min at RT the reaction was heated to 60°C overnight. The reaction was cooled and further borane dimethylsulfide complex (2M in THF, 99 μ L, 0.20 mmol) added. After stirring for 20 min at RT the reaction was heated to 60°C overnight. The reaction was cooled and quenched by dropwise addition of MeOH, then evaporated *in vacuo*. The residue was then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by HPLC (C18 X-select column, 10-60% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying (11 mg, 19%). LCMS (Method 5): Rt 3.78 min, m/z 595.2 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.86-2.15 (4H, m), 2.38 (3H, s), 2.46 (4H, t, J 9.3), 3.57 (2H, s), 3.66 (4H, t, J 9.4), 4.89 (1H, dd, J 8.7, 5.7), 5.44 (1H, t, J 8.7), 6.33 (1H, s), 7.19-7.34 (8H, m), 7.44 (1H, d, J 8.5), 7.51 (1H, dd, J 8.7, 2.9), 8.15 (1H, d, J 2.8).

Example 22

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1-methyl-piperidin-4-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 4-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 22a)



A dark brown solution of (5-fluoro-pyridin-2-yl)-hydrazine (549 mg, 4.32


mmol) and *N*-Boc-4-piperidine acetaldehyde (Aldrich, 982 mg, 4.32 mmol) in EtOH (10 mL) was stirred at reflux for 30 min, then cooled to 0°C, diluted with DCM (25 mL) and then (diacetoxyiodo)benzene (1.67 g, 5.18 mmol) was added portionwise over 1 min. The purple solution was stirred at RT for 30 min, then aqueous NaOH (1M, 20 mL) was added and the mixture shaken. The aqueous layer was extracted with DCM (2 × 20 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave an orange solid. FCC, using 3% MeOH in DCM, gave the *title compound* as a pale orange solid (1.53 g, 90%). LCMS (Method 3): Rt 3.15, m/z 235 [M-CO₂C₄H₉+H⁺].

b. **4-[6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl]-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 22b)**




To a solution of Intermediate A (392 mg, 2.40 mmol) in dry DMF (5 mL) was added NaH (60% dispersion in mineral oil, 240 mg, 6.00 mmol) and the resulting brown suspension was stirred at RT for 45 min (CARE: gas evolution). Intermediate 22a (669 mg, 2.00 mmol) was added and the dark brown solution stirred at 60°C for 2 h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (5 mL), applied to an SCX-2 cartridge (20 g) and washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* left a dark brown residue. FCC, using 4-9% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown oil (421 mg, 44%). LCMS (Method 3): Rt 2.45 min, m/z 478 [MH⁺].

c. 4-(6-((1R,4S)-4-[3-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-ureido]-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 22c)



5 A dark brown solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 162 mg, 0.401 mmol), Intermediate 22b (174 mg, 0.364 mmol) and DIPEA (0.079 mL, 0.455 mmol) in DMF (5 mL) was stirred at 100°C for 3 h. The solution was cooled to RT, concentrated *in vacuo*, suspended in water (10 mL) and extracted
10 with DCM (2 × 10 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. FCC, using 2-6% MeOH in DCM, gave the *title compound* as an off-white solid (155 mg, 58%). LCMS (Method 3): Rt 4.16 min, m/z 733 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-4-ylmethyl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 22d)




An orange solution of Intermediate 22c (155 mg, 0.211 mmol) and TFA (0.157 mL, 2.11 mmol) in DCM (3 mL) was stirred at RT for 3 h. The solution was
20 concentrated *in vacuo*, redissolved in MeOH (1 mL), applied to an SCX-2 cartridge (2 g) and washed with MeOH (15 mL). The product was eluted with 2M NH₃ in MeOH (15 mL); concentration *in vacuo* left the *title compound* as a pale

brown solid (116 mg, 87%). LCMS (Method 3): Rt 2.90 min, m/z 633 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(1-methyl-piperidin-4-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 22)

To a suspension of Intermediate 22d (58 mg, 0.0917 mmol) and formaldehyde (37%wt in water, 0.074 mL, 0.917 mmol) in DCM-MeOH (4:1, 2.5 mL), were added AcOH (0.0105 mL, 0.183 mmol) and NaBH(OAc)₃ (38.8 mg, 0.183 mmol) sequentially, then the solution stirred at RT for 2.5 h. The solution was concentrated *in vacuo* to ~0.5 mL volume, diluted with MeOH (0.5 mL), then applied to an SCX-2 cartridge and washed with MeOH (15 mL). The product was eluted with 2M NH₃ in MeOH (15 mL); concentration *in vacuo* left a pale brown solid. HPLC (XBridge C18, 40-98% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white solid after freeze-drying (15.0 mg, 25%). LCMS (Method 5): Rt 3.66 min, m/z 647 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.22 (9H, s), 1.26 (2H, m), 1.58 (2H, d, J 12.6), 1.75 (2H, t, J 11.4), 1.78-1.92 (3H, m), 2.03 (2H, m), 2.07 (3H, s), 2.31 (3H, s), 2.68 (2H, d, J 10.7), 2.97 (2H, d, J 7.0), 4.78 (1H, m), 5.48 (1H, t, J 4.5), 6.27 (1H, s), 7.07 (1H, d, J 8.5), 7.11 (1H, dd, J 9.5, 2.0), 7.23-7.35 (8H, m), 7.63 (1H, d, J 9.8), 8.03 (1H, s), 8.17 (1H, s).

Example 23




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[1-(2,2-difluoro-ethyl)-piperidin-4-ylmethyl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea

To an orange solution of Intermediate 22d (58 mg, 0.0917 mmol) and DIPEA (0.0319 mL, 0.183 mmol) in DCM-MeOH (4:1, 2.5 mL), was added 2,2-


difluoroethyl trifluoromethane-sulfonate (Fluorochem, 29.4 mg, 0.138 mmol) and the solution stirred at RT for 2 h. DIPEA (0.0319 mL, 0.183 mmol) and 2,2-difluoroethyl trifluoromethanesulfonate (29.4 mg, 0.138 mmol) were added sequentially, and the pale green solution stirred at RT for 1 h. Water (2 mL) was added and the mixture extracted with DCM (2 × 3 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a pale green-brown solid. HPLC (XBridge C18, 50-98% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white solid after freeze-drying (11.5 mg, 18%). LCMS (Method 5): Rt 3.75 min, m/z 697 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.22 (9H, s), 1.24-1.33 (2H, m), 1.59 (2H, d, J 13.0), 1.75-1.92 (3H, m), 2.01-2.10 (4H, m), 2.31 (3H, s), 2.62 (2H, td, J 15.8, 4.8), 2.82 (2H, d, J 11.4), 2.98 (2H, d, J 7.2), 4.78 (1H, m), 5.48 (1H, t, J 4.7), 6.04 (1H, tt, J 55.7, 4.4), 6.27 (1H, s), 7.07 (1H, d, J 8.9), 7.11 (1H, dd, J 10.0, 2.2), 7.21-7.35 (8H, m), 7.63 (1H, d, J 9.8), 8.03 (1H, s), 8.17 (1H, d, J 2.0).

Example 24

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxypiperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 6-Fluoro-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 24a)



(5-Fluoro-pyridin-2-yl)-hydrazine (500 mg, 3.93 mmol) in diethoxymethyl acetate (5 mL) was stirred at RT for 2 h. The resulting precipitate was diluted with cyclohexane (5 mL) and filtered to give the *title compound* (379 mg, 70%). ¹H


NMR (400 MHz, CDCl₃): 7.25 (1H, m), 7.84 (1H, m), 8.09 (1H, t), 8.84 (1H, s).

b. 3-Chloro-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 24b)




A solution of Intermediate 24a (789 mg, 5.98 mmol) and N-chlorosuccinimide (878 mg, 6.57 mmol) in chloroform (15 mL) was heated at 65°C overnight. The cooled mixture was washed with sat. aq. NaHCO₃ solution (2 × 15 mL) and dried (Na₂SO₄). The solvent was evaporated, then the residue suspended in diethyl ether (10 mL) and filtered to give the *title compound* (730 mg, 76%). LCMS (Method 1): Rt 1.83 min, m/z 172 [MH⁺].

c. 1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-4-ol (Intermediate 24c)



A brown solution of Intermediate 24b (855 mg, 4.98 mmol) and 4-hydroxypiperidine (2.02 g, 19.9 mmol) in DMA (15 mL) was irradiated to 175°C in the microwave for 3 h. The cooled solution was concentrated *in vacuo*, then the residue diluted with water (20 mL) and brine (20 mL). The mixture was washed with diethyl ether (2 × 50 mL), then extracted with EtOAc (2 × 50 mL). The combined EtOAc layers were washed with brine (50 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave a yellow oil. The aqueous was further extracted with DCM (3 × 50 mL). The combined DCM layers were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown oil. The two oils were combined. FCC, using 4-5% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a yellow crystalline solid (514 mg, 44%). LCMS (Method 3): Rt 1.96 min, m/z 237 [MH⁺].


d. 6-Fluoro-3-(4-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 24d)



Triisopropylsilyl trifluoromethanesulfonate (395 mg, 1.29 mmol) was added dropwise to a solution of Intermediate 24c (254 mg, 1.07 mmol) and Et₃N (0.20 mL, 1.40 mmol) in DMF (3 mL) at RT under N₂. The mixture was stirred for 1 h then Et₃N (0.10 mL, 0.70 mmol) and triisopropylsilyl trifluoromethanesulfonate (200 mg, 0.65 mmol) were added sequentially, and the mixture stirred for 1 h. The solvent was evaporated, the residue applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left the *title compound* (339 mg, 80%). LCMS (Method 1): Rt 4.74 min, m/z 393 [MH⁺].

e. (1S,4R)-4-[3-(4-Triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3a]pyridin-6-yloxy]1,2,3,4-tetrahydro-naphthalen-1-ylamine.


(Intermediate 24e)



Intermediate A (81 mg, 0.497 mmol) was added dropwise to a suspension of NaH (60% dispersion in oil, 59 mg, 1.49 mmol) in DMF (3 mL) at RT under N₂. The mixture was stirred for 15 min then Intermediate 24d (150 mg, 0.382 mmol) was added and the mixture stirred at 60°C under N₂ for 3 h. Sat. aq. NH₄Cl solution (0.2 mL) was added to the cooled mixture which was then diluted with water (15 mL) and extracted with EtOAc (3 × 15 mL). The combined organics were washed with brine (15 mL) and dried (Na₂SO₄). The solvent was evaporated,

the residue applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left the *title compound* (180 mg, 90%). LCMS (Method 1): Rt 3.05 min, m/z 536 [MH⁺].

f. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-triisopropylsilyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea. (Intermediate 24f)



A mixture of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 140 mg, 0.346 mmol), Intermediate 24e (185 mg, 0.346 mmol) and DIPEA (0.18 mL, 1.03 mmol) in DMF (3 mL) was heated at 60°C for 4 h under N₂. The cooled mixture was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown gum. FCC, using 0.6-6% [2M NH₃ in MeOH] in DCM, gave the *title compound* (124 mg, 45%). LCMS (Method 4): Rt 5.15 min, m/z 791 [MH⁺].


g. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxypiperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 24)

TBAF (1M in THF, 0.235 mL, 0.235 mmol) was added dropwise to a solution of Intermediate 24f (124 mg, 0.157 mmol) in dry THF (3 mL) at -30°C for 4 h under N₂. The mixture was allowed to warm to RT, then stirred for 5 h. The solution was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown foam. HPLC (Gemini C18; 40-90% MeCN in H₂O, 0.1% NH₄OH) gave the *title*


compound as a white powder (77 mg, 77%). LCMS (Method 5): Rt 4.18 min, m/z 635 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.33 (9H, s), 1.71-1.80 (2H, m), 1.87-1.96 (1H, m), 2.01-2.12 (4H, m), 2.25 (1H, m), 2.36 (3H, s), 3.04-3.11 (2H, m), 3.37-3.44 (2H, m), 3.93 (1H, m), 5.09 (1H, td, J 8.9, 5.3), 5.19 (1H, t, J 4.0), 5.44 (1H, d, J 8.8), 6.28 (1H, s), 6.46 (1H, br s), 6.99 (1H, dd, J 9.9, 2.1), 7.21 (2H, d, J 8.1), 7.25-7.33 (5H, m), 7.39 (2H, d, J 8.1), 7.49 (1H, d, J 9.9).

Example 25

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[(2-hydroxy-ethyl)-methyl-amino]-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 2-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-methyl-amino]ethanol and 2-[(3-chloro-[1,2,4]triazolo[4,3-a]pyridin-6-yl)-methyl-amino]ethanol (Intermediate 25a)




A solution of Intermediate 24b (300 mg, 1.75 mmol) and 2-methylamino ethanol (660 mg, 8.77 mmol) in NMP (2 ml) was heated at 165°C for 2 h in the microwave. The cooled mixture was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown gum. FCC, using 3-6% [2M NH₃ in MeOH] in DCM, gave a mixture of the *title compounds* (240 mg, 65%). LCMS (Method 4): Rt 1.05 min, m/z 211 [MH⁺] and Rt 1.76 min, m/z 227 [MH⁺].

b. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-methyl-(2-triisopropylsilyloxy-ethyl)-amine. (Intermediate 25b)



Triisopropylsilyl trifluoromethanesulfonate (454 mg, 1.48 mmol) was added dropwise to a solution of Intermediate 25a (240 mg, 1.14 mmol) and Et₃N (0.24 mL, 1.71 mmol) in DMF (2 mL) at RT under N₂ and the mixture was stirred for 3 h. The solvent was evaporated, the residue applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown gum. FCC, using 0-3% [2M NH₃ in MeOH] in DCM, gave the *title compound* (160 mg, 38%). LCMS (Method 4): Rt 4.25 min, m/z 367 [MH⁺].


c. [6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl]-methyl-(2-triisopropylsilyloxy-ethyl)-amine. (Intermediate 25c)



Intermediate A (93 mg, 0.568 mmol) was added dropwise to a suspension of NaH (60% dispersion in oil, 68 mg, 1.71 mmol) in DMF (3 mL) at RT under N₂. The mixture was stirred for 15 min then Intermediate 25b (160 mg, 0.437 mmol) was added and the mixture stirred at 60°C under N₂ for 3 h. Sat. aq. NH₄Cl solution (0.2 mL) was added to the cooled mixture which was then diluted with water (15 mL) and extracted with EtOAc (3 × 15 mL). The combined organics were washed with brine (15 mL) and dried (Na₂SO₄). The solvent was evaporated, the residue applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown foam. FCC, using 0-6% [2M NH₃ in MeOH] in DCM, gave the *title compound* (64 mg,

29%). LCMS (Method 1): Rt 2.71 min, m/z 510 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[methyl-(2-triisopropylsilanyloxy-ethyl-amino)]-1-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea. (Intermediate 25d)



5

A mixture of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 71 mg, 0.175 mmol), Intermediate 25c (64 mg, 0.125 mmol) and DIPEA (0.093 mL, 0.524 mmol) in DMF (2 mL) was heated at 60°C for 4 h under N₂. The cooled mixture was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a brown gum. FCC, using 3-6% [2M NH₃ in MeOH] in DCM, gave the *title compound* (65 mg, 68%). LCMS (Method 1): Rt 4.78 min, m/z 765 [MH⁺].

10

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-{3-[(2-hydroxy-ethyl)-methyl-amino]-1-[1,2,4]triazolo[4,3-a]pyridine-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 25)

15

TBAF (1M in THF; 0.127 mL, 0.235 mmol) was added dropwise to a solution of Intermediate 25d (65 mg, 0.085 mmol) in dry THF (0.3 mL) at -30°C under N₂. The mixture was allowed to warm to RT, then stirred for 1 h. The solution was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* left a pale yellow oil. Further purification by HPLC. (Gemini C18; 40-100% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white powder (32 mg, 62%). LCMS (Method 5): Rt 4.11 min, m/z 609 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.32 (9H, s), 1.90-1.99 (1H, m), 2.02-2.12 (2H, m), 2.23-2.30 (1H, m), 2.37 (3H, s), 2.94

20


25

(3H, s), 3.11 (1H, ddd, J 14.3, 6.9, 3.5), 3.18 (1H, ddd, J 14.3, 5.6, 3.2), 3.68-3.79 (2H, m), 5.06 (1H, td, J 8.6, 5.5), 5.20 (1H, t, J 4.2), 5.50 (1H, d, J 8.7), 6.31 (1H, s), 6.42 (1H, s), 6.98 (1H, dd, J 9.9, 2.0), 7.18-7.30 (6H, m), 7.39 (2H, d, J 8.2), 7.49 (1H, d, J 9.9), 7.58 (1H, d, J 2.0).

5

Example 26


1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



10

a. (S)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidin-3-ol

(Intermediate 26a)




15

A mixture of Intermediate 24b (300 mg, 1.74 mmol) and (S)-3-hydroxypyrrolidine (600 mg, 9.96 mmol) in NMP (6 mL) was heated in the microwave at 160°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-8% [2M NH₃ in MeOH] in DCM, gave the *title compound* (150 mg, 38%). LCMS (Method 1): Rt 1.45 min, m/z 223 [MH⁺].


20

b. 6-Fluoro-3-((S)-3-triisopropylsilanyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 26b)




Triisopropylsilyl trifluoromethanesulfonate (250 mg, 0.81 mmol) was added to a solution of Intermediate 26a (150 mg, 0.67 mmol) and Et₃N (101 mg, 1.00 mmol) in DMF (2 mL) and the mixture stirred at RT for 1 h. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (220 mg, 86%). LCMS (Method 4): Rt 4.15 min, m/z 379 [MH⁺].

c. (1S,4R)-4-[3-((S)-3-Triisopropylsilanyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 26c)



To a solution of Intermediate A (104 mg, 0.640 mmol) in DMF (2 mL) was added NaH (60% in oil, 70 mg, 1.74 mmol) and the mixture stirred at RT for 20 min, before Intermediate 26b (220 mg, 0.582 mmol) was added. This mixture was stirred thermally at 60°C for 4 h, then at 60°C in the microwave for 3 h. The cooled reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM gave the *title compound* as a yellow gum (70 mg, 23%). LCMS (Method 1): Rt 2.89, m/z 522 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-3-triisopropylsilyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 26d)



5 A solution of Intermediate 26c (70 mg, 0.134 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 81 mg, 0.201 mmol) and DIPEA (70 mg, 0.54 mmol) in DMF (2 mL) was stirred at 60°C for 30 min. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was
10 eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as an off-white foam (19 mg, 23%). LCMS (Method 1): Rt 4.70, m/z 777 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 26)


15

To a solution of Intermediate 26d (19 mg, 0.024 mmol) in THF (1 mL) at -30°C was added TBAF (1M in THF, 36 μL, 0.036 mmol) and the mixture was allowed to warm to RT over 1 h. The reaction mixture was applied to an SCX-2 cartridge (2 g) and washed with MeOH. The product was eluted with 2M NH₃ in
20 MeOH; concentration *in vacuo* gave a residue. Purification by HPLC (C6-Ph column, 35-75% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as an off-white powder after freeze-drying (10.0 mg, 60%). LCMS (Method 5): Rt 4.00 min, m/z 621 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.29 (9H, s), 1.89-2.09 (4H, m), 2.13-2.29 (2H, m), 2.34 (3H, s), 3.35 (1H, td, J 9.2, 4.6), 3.45 (1H, d, J 11.1), 3.50


(1H, dd, J 11.1, 4.3), 3.60 (1H, m), 4.51 (1H, m), 5.04 (1H, td, J 8.5, 5.3), 5.15 (1H, t, J 4.1), 5.57 (1H, d, J 8.6), 6.26 (1H, s), 6.67 (1H, br s), 6.89 (1H, dd, J 9.9, 1.7), 7.15 (2H, d, J 7.1), 7.24-7.29 (4H, m), 7.30-7.38 (4H, m).

Example 27

5 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**




a. **[(R)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidin-2-yl]-methanol (Intermediate 27a)**



A mixture of Intermediate 24b (300 mg, 1.74 mmol) and (R)-(-)-2-(hydroxymethyl)-pyrrolidine (704 mg, 9.96 mmol) in NMP (6 mL) was heated in the microwave at 160°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (75 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (220 mg, 53%). LCMS (Method 4): Rt 1.50, m/z 237 [MH⁺].


b. **6-Fluoro-3-((S)-2-triisopropylsilyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 27b)**



Triisopropylsilyl trifluoromethanesulfonate (430 mg, 1.40 mmol) was added


to a solution of Intermediate 27a (220 mg, 0.93 mmol) and Et₃N (190 mg, 1.86 mmol) in a DMF (2 mL) and the mixture stirred at RT for 1 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (170 mg, 47%). LCMS (Method 1): Rt 4.45, m/z 393 [MH⁺].

c. (1S,4R)-4-[3-((R)-2-Triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 27c)



To a solution of Intermediate A (60 mg, 0.364 mmol) in DMF (2 mL) was added NaH (60% in oil, 40 mg, 0.99 mmol) and the mixture stirred at RT for 20 min, before Intermediate 27b (130 mg, 0.331 mmol) was added. This mixture was heated at 60°C in the microwave for 1.5 h. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown oil (50 mg, 28%). LCMS (Method 4): Rt 2.36, m/z 536 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-2-triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 27d)



A solution of Intermediate 27c (50 mg, 0.093 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic


Communications, 2009, 39, 3999-4009; 56 mg, 0.140 mmol) and DIPEA (70 mg, 0.54 mmol) in DMF (2 mL) was stirred at 60°C for 1 h. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using
5 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as an off-white foam (27 mg, 36%). LCMS (Method 4): Rt 4.58, m/z 791 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 27)

10 To a solution of Intermediate 27d (27 mg, 0.034 mmol) in THF (1 mL) at -30°C was added TBAF (1M in THF, 36 µL, 0.036 mmol) and the mixture was allowed to warm to RT over 1 h. The reaction mixture was applied to an SCX-2 cartridge (2 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. Purification by HPLC (C6-Ph
15 column, 10-70% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (10.0 mg, 45%). LCMS (Method 5): Rt 4.18 min, m/z 635 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.25 (9H, s), 1.63-1.75 (1H, m), 1.81-1.95 (3H, m), 1.97-2.01 (3H, m), 2.26 (1H, m), 2.31 (3H, s), 3.17-3.26 (1H, m), 3.32-3.47 (2H, m), 3.57 (1H, dd, J 11.6, 3.2), 4.07-4.15 (1H, m), 4.99-5.07 (1H,
20 m), 5.15 (1H, t, J 4.1), 5.57 (1H, d, J 8.9), 6.25 (1H, s), 6.61 (1H, s), 6.91 (1H, d, J 10.2), 7.16 (2H, d, J 8.3), 7.20-7.29 (4H, m), 7.32 (2H, d, J 8.0), 7.35-7.41 (2H, m).


Example 28

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




5

a. [(S)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidin-2-yl]-methanol (Intermediate 28a)



A mixture of Intermediate 24b (300 mg, 1.74 mmol) and L-prolinol (704 mg, 9.96 mmol) in NMP (4 mL) was heated in the microwave at 160°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (210 mg, 50%). LCMS (Method 4): Rt 1.50 min, m/z 237 [MH⁺].

b. 6-Fluoro-3-((S)-2-triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 28b)




Triisopropylsilyl trifluoromethanesulfonate (327 mg, 1.06 mmol) was added to a solution of Intermediate 28a (210 mg, 0.89 mmol) and Et₃N (135 mg, 1.33 mmol) in a DMF (3 mL) and the mixture stirred at RT for 1 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue.

20


FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (110 mg, 31%). LCMS (Method 1): Rt 4.45 min, m/z 393 [MH⁺].

c. (1S,4R)-4-[3-((S)-2-Triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 28c)



To a solution of Intermediate A (50 mg, 0.309 mmol) in DMF (2 mL) was added NaH (60% in oil, 33 mg, 0.80 mmol) and the mixture stirred at RT for 20 min, before Intermediate 28b (110 mg, 0.280 mmol) was added. This mixture was heated at 60°C in the microwave for 1.25 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-7% [2M NH₃ in MeOH] in DCM gave the *title compound* as a viscous yellow oil (42 mg, 28%). LCMS (Method 4): Rt 2.55 min, m/z 536 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 28d)



A solution of Intermediate 28c (40 mg, 0.074 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 45 mg, 0.112 mmol) and DIPEA (38 mg, 0.296 mmol) in DMF (2 mL) was stirred at 60°C for 30 min. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was


eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a viscous yellow oil (38 mg, 65%). LCMS (Method 1): Rt 4.72 min, m/z 791 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 28)


To a solution of Intermediate 28d (38 mg, 0.048 mmol) in THF (1 mL) at -30°C was added TBAF (1M in THF, 72 µL, 0.072 mmol) and the mixture was allowed to warm to RT over 1 h. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. Purification by HPLC (C6-Ph column, 10-70% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (12 mg, 39%). LCMS (Method 5): Rt 4.20 min, m/z 635 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.29 (9H, s), 1.78-2.28 (8H, m), 2.38 (3H, s), 3.36-3.46 (1H, m), 3.53 (2H, dd, J 5.0, 1.6), 3.64-3.73 (1H, m), 4.04-4.12 (1H, m), 4.85-4.91 (1H, m), 5.36 (1H, t, J 4.1), 6.32 (1H, s), 7.11 (1H, dd, J 10.0, 2.1), 7.20-7.35 (8H, m), 7.44 (1H, d, J 9.9), 7.99-8.01 (1H, m).

Example 29

1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




**a. 3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-benzoic acid ethyl ester
(Intermediate 29a)**




A solution of 3-hydrazino benzoic acid (5.53 g, 36.4 mmol) and 4,4-
5 dimethyl-3-oxo pentanenitrile (5.00 g, 40.0 mmol) and concentrated sulfuric acid
(2 mL) in EtOH (72 mL) was stirred at reflux for 19 h. The cooled mixture was
concentrated *in vacuo*, was diluted with 1N NaOH solution (15 mL) and extracted
with EtOAc. The combined organics were dried and concentrated *in vacuo*. The
residue was purified by FCC, using 0-40% EtOAc in cyclohexane, to give the *title*
10 *compound* as an off-white powder (5.92 g, 56%). LCMS (Method 3): Rt 3.04 min,
m/z 288 [MH⁺].

**b. [3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenyl]-methanol
(Intermediate 29b)**



15 To a solution of Intermediate 29a (1.00 g, 3.48 mmol) and Et₃N (265 μ L,
1.91 mmol) in EtOH (35 mL) was added NaBH₄ (198 mg, 5.23 mmol) and the
suspension stirred at RT for 1.5 h. NaBH₄ (198 mg, 5.23 mmol) was added and the
suspension stirred for a further 19 h. NaBH₄ (1.31 g, 34.8 mmol) was added and
the suspension stirred for a further 24 h, then diluted with water and extracted with
20 DCM. The combined organics were dried and concentrated *in vacuo*. The residue
was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title*
compound as a white solid (740 mg, 87%). LCMS (Method 3): Rt 2.05 min, m/z
246 [MH⁺].

c. [5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 29c)




To a bi-phasic mixture of Intermediate 29b (737 mg, 3.00 mmol) in EtOAc
5 (22.5 mL) and 1N NaOH solution (8.11 mL, 8.11 mmol) at 0°C was added 2,2,2-trichloroethyl chloroformate (0.45 mL, 3.30 mmol) and the mixture stirred for 1.25 h. The layers were separated and the organic layer was washed with brine, dried and concentrated *in vacuo* to give the *title compound* as an off-white solid (1.26 g, 99%). LCMS (Method 3): Rt 4.00 min, m/z 420, 422 [MH⁺].

10 **d. 1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 29)**

A solution of Intermediate 29c (509 mg, 1.21 mmol) and Intermediate 3c
15 (400 mg, 1.10 mmol) and DIPEA (0.58 mL, 3.30 mmol) in THF (11 mL) was stirred at reflux for 15 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the product. A 50 mg portion of this was further purified by HPLC (XBridge C18 column, 10-98% MeCN in H₂O, 0.1% NH₄OH) to give the *title compound*
20 as a white powder after freeze-drying (10 mg, 20%). LCMS (Method 5): Rt 4.23 min, m/z 635 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.35 (9H, s), 1.57-1.74 (6H, m), 1.89 (1H, t, J 10.7), 2.01-2.08 (2H, m), 2.14-2.22 (1H, m), 3.06 (4H, t, J 5.1), 4.56 (2H, s), 5.09 (1H, m), 5.18 (1H, t, J 4.3), 5.92 (1H, d, J 8.6), 6.40 (1H, s), 6.88 (1H, dd, J 9.9, 2.1), 7.14-7.17 (3H, m), 7.30-7.32 (5H, m), 7.42
25 (1H, d, J 8.1), 7.50 (1H, s).


Example 30

3-[3-tert-Butyl-5-(3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzoic acid ethyl ester



5

a. 3-[3-tert-Butyl-5-(2,2,2-trichloro-ethoxycarbonylamino)-pyrazol-1-yl]-benzoic acid ethyl ester (Intermediate 30a)



The *title compound* was prepared from Intermediate 29a using an analogous procedure to that described for Intermediate 29c. LCMS (Method 3): Rt 4.67 min, m/z 462, 464 [MH⁺].

b. 3-[3-tert-Butyl-5-(3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzoic acid ethyl ester (Example 30)


A mixture of Intermediate 5c (163 mg, 0.45 mmol), Intermediate 30a (208 mg, 0.45 mmol) in 1,4-dioxane (3 mL) and DIPEA (119 μ L, 0.68 mmol) was stirred at 90°C for 3 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-12% MeOH in DCM, to give the product (291 mg, 96%). A 64 mg portion of this was further purified by HPLC (C18 X-select column, 30-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying powder (38 mg). LCMS (Method 5): Rt 3.85 min, m/z 677.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.26-1.33 (12H, m), 1.80-

15

20



2.27 (11H, m), 2.30-2.39 (1H, m), 3.09-3.16 (1H, m), 3.99 (1H, br t, J 8.2), 4.32 (2H, q, J 7.1), 4.74-4.83 (1H, m), 5.36-5.41 (1H, m), 6.33 (1H, s), 7.11 (1H, d, J 8.6), 7.19-7.37 (5H, m), 7.66 (1H, t, J 7.9), 7.75 (1H, d, J 9.9), 7.80-7.84 (1H, m), 7.93-7.98 (1H, m), 8.07-8.09 (1H, m), 8.22-8.28 (2H, m).

5

Example 31


**1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea, partial formate salt**

10 A solution of Example 30 (225 mg, 0.333 mmol) and sodium borohydride (31.5 mg, 0.833 mmol) in ethanol (3 mL) was stirred at RT for 2.5 h. Sodium borohydride (31.5 mg, 0.833 mmol) was added and the solution stirred for 90 min. Sodium borohydride (31.5 mg, 0.833 mmol) was added and the solution stirred for 2.5 h. Sodium borohydride (31.5 mg, 0.833 mmol) was added and the solution
 15 stirred for a further 15.5 h. Water was added followed by sat. aq. NH_4Cl solution. The mixture was then extracted with DCM (4 x 20 mL). The combined organic extracts were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the product (150 mg). This was further purified by HPLC (C18 X-select column, 30-98% MeCN in H_2O , 0.1% HCO_2H) to
 20 give the *title compound* as a white powder after freeze-drying (97 mg, 46%). LCMS (Method 5): Rt 3.29 min, m/z 635.2 $[\text{MH}^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 1.28 (9H, s), 1.82-2.26 (11H, m), 2.30-2.39 (1H, m), 3.09-3.16 (1H, m), 3.99 (1H, t, J 8.1), 4.57 (2H, s), 4.78-4.86 (1H, m), 5.39 (1H, t, J 4.3), 6.33 (1H, s), 7.11 (1H, d, J 8.4), 7.24-7.38 (7H, m), 7.42-7.48 (2H, m), 7.75 (1H, d, J 9.9), 8.1
 25 (1H, s), 8.20 (0.5H, s), 8.24 (1H, m).

Example 32**1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(2-morpholin-4-yl-ethoxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea****5 a. 2-((1S,4R)-4-Hydroxy-1,2,3,4-tetrahydro-naphthalen-1-yl)-isoindole-1,3-dione (Intermediate 32a)**

A solution of Intermediate A (150 mg, 0.92 mmol) and phthalic anhydride (143 mg, 0.97 mmol) in toluene (9 mL) was stirred and heated at reflux for 19.5 h.

10 After cooling, the mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-50% EtOAc in cyclohexane, to give the *title compound* as white powder (215 mg, 79%). LCMS (Method 3): Rt 3.36 min, m/z 316 [MNa⁺].


b. 2-[(1S,4R)-4-(2-Morpholin-4-yl-2-oxo-ethoxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-isoindole-1,3-dione (Intermediate 32b)

15 A solution of Intermediate 32a (115 mg, 0.39 mmol) in dry THF (4 mL) was added NaH (60% in mineral oil, 23 mg, 0.59 mmol) at RT and stirred for 15 min. 4-(Chloroacetyl)morpholine (56 µL, 0.43 mmol) was then added and the mixture heated at reflux for 3.5 h. After cooling, the dark brown mixture was

20 diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% MeOH in DCM, to give the *title compound* as light brown foam (103 mg, 62%). LCMS


(Method 3): Rt 3.44 min, m/z 443 [MNa⁺].

c. 2-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-1-morpholin-4-yl-ethanone (Intermediate 32c)



5 A solution of Intermediate 32b (100 mg, 0.23 mmol) and hydrazine hydrate (74 μ L, 2.4 mmol) in dry MeOH (6 mL) was stirred at RT for 5 h. The reaction mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* as an off white foam (50 mg, 72%). LCMS (Method 3): Rt 1.70 min, m/z 291 [MH⁺].

10 **d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(2-morpholin-4-yl-2-oxo-ethoxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 32d)**




A stirred solution of Intermediate 32c (50 mg, 0.17 mmol) and 5-tert-butyl-
15 2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 84 mg, 0.21 mmol) and DIPEA (90 μ L, 0.52 mmol) in THF (1.7 mL) was heated at reflux for 21 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-
20 5% [2M NH₃ in MeOH] in DCM, to give impure product. This residue purified by HPLC (Gemini C18 column, 30-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying (37 mg, 39%). LCMS (Method 5): Rt 4.73 min, m/z 546 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(2-morpholin-4-yl-ethoxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 32)

To a solution of Intermediate 32d (50 mg, 62.3 μ mol) in THF (1.7 mL) was added borane (1M in THF, 0.12 mL, 0.12 mmol) and the mixture stirred at 60°C. After 23 h, further borane (1M in THF, 0.31 mL, 0.31 mmol) was added. After 26 h, further borane (1M in THF, 0.31 mL, 0.31 mmol) was added. After 3 d, the cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by HPLC (Gemini C18 column, 20-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying (15 mg, 47%). LCMS (Method 5): Rt 3.64 min, m/z 532 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.32 (9H, s), 1.93-1.96 (3H, m), 2.02-2.07 (1H, m), 2.32 (4H, t, J 4.5), 2.38 (3H, s), 2.47 (1H, dt, J 13.0, 5.3), 2.61 (1H, ddd, J 13.0, 7.3, 5.4), 3.56-3.58 (5H, m), 3.67-3.68 (1H, m), 4.35-4.37 (1H, m), 4.98-5.02 (1H, m), 5.50 (1H, d, J 8.8), 6.27 (2H, s), 7.18-7.28 (5H, m), 7.30-7.32 (1H, m), 7.36 (2H, d, J 8.2).

Example 33


1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}urea



20

25

a. [5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 33a)




To a suspension of [4-(5-amino-3-tert-butyl-pyrazol-1-yl)-phenyl]-methanol
 5 (WO2011/070368; 3.05 g, 12.4 mmol) in aq. NaOH solution (1 M, 31 mL, 31 mmol) and EtOAc (30 mL) at RT was added 2,2,2-trichloroethyl chloroformate (1.88 mL, 13.7 mmol) over 3 min (CARE: exotherm to ~35°C) and the mixture stirred at RT for 1 h. The aqueous layer was extracted with EtOAc (20 mL), then the combined organics washed with brine (25 mL), dried (Na₂SO₄), filtered and
 10 concentrated *in vacuo* to leave a pale orange solid. Recrystallisation from hot cyclohexane-EtOAc (3:1, 30 mL) and drying *in vacuo* gave the *title compound* as a flocculent off-white solid (3.87 g, 74%). LCMS (Method 3): Rt 4.00 min, m/z 420, 422 [MH⁺].

**b. 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
 15 {(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}urea (Example 33)**

A mixture of Intermediate 5c (109 mg, 0.300 mmol) and Intermediate 33a (126 mg, 0.300 mmol) in 1,4-dioxane (3 mL) and DIPEA (78 µL, 0.45 mmol) was stirred at 80°C for 3 h, and then at 95°C for 2 h. The cooled mixture was
 20 concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the *title compound* as an off-white powder after freeze-drying (95 mg, 50%). LCMS (Method 5): Rt 3.26 min, m/z 635.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.81-2.26 (11H, m), 2.31-2.40 (1H, m) 3.10-3.17 (1H, m), 3.99 (1H, br t, J 8.1), 4.56 (2H, d, J 5.6), 4.78-4.87 (1H, m), 5.29 (1H, t, J 5.7),
 25 5.37-5.42 (1H, m), 6.33 (1H, s), 7.11 (1H, d, J 8.7), 7.24-7.47 (9H, m), 7.75 (1H, d,

J 9.7), 8.07 (1H, s), 8.24 (1H, br d).

Example 34




**1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 [(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-
tetrahydro-naphthalen-1-yl]-urea**

A mixture of Intermediate 3c (109 mg, 0.300 mmol) and Intermediate 33a (126 mg, 0.300 mmol) in 1,4-dioxane (3 mL) and DIPEA (78 μ L, 0.45 mmol) was stirred at 80°C for 3 h, and then at 95°C for 2 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the *title compound* as an off-white powder after freeze-drying (95 mg, 50%). LCMS (Method 5): Rt 4.21 min, m/z 635.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.56-2.17 (10H, m), 3.11-3.17 (4H, m), 4.56 (2H, d, J 5.7), 4.77-4.86 (1H, m), 5.29 (1H, t, J 5.7), 5.52-5.57 (1H, br t), 6.33 (1H, s), 7.09 (1H, d, J 8.4), 7.16 (1H, dd, J 9.7, 2.2), 7.25-7.47 (8H, m), 7.58-7.64 (2H, m), 8.07 (1H, s).

Example 35

**1-{5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-
20 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**




a.

Di-tert-butyl

1-{3-[(2-{[tert-

butyl(dimethyl)silyl|oxy}ethyl)sulfanyl|phenyl}hydrazine-1,2-dicarboxylate


(Intermediate 35a)



A mixture of 3-bromothiophenol (1.00 g, 5.29 mmol),
5 bromoethoxydimethylsilyl ether (1.36 mL, 6.35 mmol) and K_2CO_3 (1.46 g, 10.6 mmol) in acetone (15 mL) was stirred at RT overnight. The mixture was filtered, evaporated, and the residue re-dissolved in dry THF (15 mL) and cooled to $-78^\circ C$. nBuLi (1.6M in hexanes, 4.5 mL, 7.28 mmol) was added dropwise and the mixture stirred for 10 min. Di-tert-butyl azodicarboxylate (1.54 g, 6.68 mmol) was added
10 in one portion at $-78^\circ C$ and the mixture stirred for 20 min. The mixture was then allowed to warm to RT over 2 h. The reaction was quenched with sat. aq. NH_4Cl solution (15 mL), then extracted with EtOAc (3×15 mL). The combined organic extracts were washed with brine (20 mL), dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% EtOAc in pentane, to give
15 the *title compound* as a pale yellow oil (1.68 g, 64%). ^1H NMR (400 MHz, $CDCl_3$): 0.04 (6H, s), 0.84 (9H, s), 1.48 (18H, m), 3.04 (2H, t), 3.79 (2H, t), 7.14-7.28 (3H, m), 7.42 (1H, s).

b. 2-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl)phenylsulfanyl]-ethanol.


(Intermediate 35b)



A mixture of Intermediate 35a (1.68 g, 3.37 mmol), pivaloyl acetonitrile (0.42 g, 3.37 mmol) and concentrated HCl solution (1.7 mL) in ethanol (10 mL) was heated under reflux for 3 h. After cooling, the pH was adjusted to ~7 (using

sat. aq. NaHCO₃ solution) and the mixture diluted with water (20 mL) and extracted with EtOAc (3 × 20 mL). The combined organic extracts were washed with brine (20mL), dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, eluting with 20-80% EtOAc in pentane, to give the *title compound* as a pale yellow oil (458 mg, 47%). LCMS (Method 1): Rt 2.35 min, m/z 292 [MH⁺].

c. {5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-carbamic acid 2,2,2-trichloro-ethyl ester. (Intermediate 35c)




2,2,2-Trichloroethyl chloroformate (0.10 mL, 0.78 mmol) was added to a solution of Intermediate 35b (176 mg, 0.60 mmol) and DIPEA (0.31 mL, 1.81 mmol) in THF (10 mL) and the mixture stirred at RT for 3 h. The mixture was diluted with water (15 mL), extracted with EtOAc (3 × 20 mL), then the combined organic extracts dried (Na₂SO₄) and concentrated *in vacuo*. The residue was suspended in cyclohexane, and filtered to give the *title compound* as a yellow solid (280 mg, 100%). LCMS (Method 1): Rt 3.93 min, m/z 466, 468 [MH⁺]

d. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 35)

A mixture of Intermediate 5c (109 mg, 0.3 mmol) and Intermediate 35c (140 mg, 0.3 mmol) in 1,4-dioxane (3 mL) and DIPEA (78 μL, 0.45 mmol) was stirred at 90°C for 4 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the *title compound* as an off-white powder after freeze-drying (115 mg, 56%). LCMS (Method 5): Rt 3.47 min, m/z 681 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.83-2.26

(11H, m), 2.31-2.39 (1H, m), 3.04 (2H, t, J 6.8), 3.09 (1H, m), 3.59 (2H, q, J 6.0), 3.99 (1H, t, J 8.2), 4.78-4.86 (1H, m), 4.95 (1H, t, J 5.6), 5.36-5.41 (1H, m), 6.33 (1H, s), 7.08 (1H, d, J 8.4), 7.24-7.46 (9H, m), 7.75 (1H, d, J 9.6), 8.12 (1H, s), 8.25 (1H, br d).

5

Example 36

1-[5-(2-Hydroxy-1,1-dimethyl-ethyl)-2-p-tolyl-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

10

The *title compound* was prepared as an off-white solid (120 mg, 68%) using Intermediate 3c (100 mg, 0.28 mmol) and [5-(2-hydroxy-1,1-dimethyl-ethyl)-2-p-tolyl-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (WO2009/015000; 138 mg, 0.34 mmol) in a similar manner to Example 34. LCMS (Method 5): Rt 4.02 min, m/z 635 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.21 (6H, s), 1.58-1.64 (2H, m), 1.69-1.75 (4H, m), 1.81-1.97 (2H, m), 2.00-2.16 (2H, m), 2.36 (3H, s), 3.14 (4H, t, J 5.4), 3.43 (2H, s), 4.55 (1H, br s), 4.79-4.85 (1H, m), 5.54 (1H, t, J 4.3), 6.32 (1H, s), 7.08 (1H, d, J 8.6), 7.16 (1H, dd, J 4.9, 2.3), 7.26-7.39 (8H, m), 7.59-7.63 (2H, m), 8.07 (1H, s).


15

20


Example 37

**1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,3S)-3-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-indan-1-yl}-urea**

5



**a. (1S,3S)-3-[3-((S)-1-Methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-indan-1-ylamine (Intermediate 37a)**



(1S,3S)-3-Amino-indan-1-ol (110 mg, 0.74 mmol) was added to a
10 suspension of NaH (60% in mineral oil, 87 mg, 2.17 mmol) in anhydrous DMF (5
mL) at RT and stirred for 20 min. Intermediate 5b (160 mg, 0.72 mmol) was then
added in one portion and the mixture heated at 60°C for 3 h. After cooling the
mixture was diluted with water (20 mL) and extracted with EtOAc (3 × 50 mL).
The combined organics were washed with brine (50 mL), dried (MgSO₄) and
15 evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in
DCM, to give the *title compound* as give a foam (132 mg, 52%). LCMS (Method
1): Rt 0.34 min, m/z 350 [MH⁺]. ¹H NMR (300 MHz, CDCl₃): 1.95-2.16 (6H, m),
2.24 (3H, s), 2.26-2.45 (2H, m), 2.77 (1H, ddd, J 14.1, 7.0, 2.0), 3.23-3.30 (1H, m),
4.02-4.09 (1H, m), 4.71 (1H, t, J 6.8), 5.68 (1H, dd, J 6.2, 1.9), 7.04 (1H, dd, J 9.9,
20 2.2), 7.29-7.50 (4H, m), 7.64 (1H, dd, J 9.9, 0.9), 8.35 (1H, d, J 2.2).


**b. 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,3S)-3-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-**

ylxy]-indan-1-yl}-urea (Example 37)

A solution of Intermediate 33a (190 mg, 0.454 mmol), Intermediate 37a (132 mg, 0.378 mmol) and DIPEA (132 μ L, 0.756 mmol) in anhydrous DMF (3 mL) was stirred at 100°C for 3 h. After cooling, the mixture was partitioned between EtOAc (50 mL) and water (50 mL), and then extracted into EtOAc (3 $\times$ 50 mL). The combined organics were washed with brine (50 mL), dried (MgSO₄) and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* as give an off-white solid after freeze-drying (143 mg, 61%). LCMS (Method 5): Rt 3.23, m/z 621 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.22 (9H, s), 1.95-1.97 (3H, m), 2.07 (3H, s), 2.20-2.23 (3H, m), 2.52-2.53 (1H, m), 3.10 (1H, m), 3.92 (1H, t, J 8.1), 4.51 (2H, d, J 5.6), 5.25-5.27 (2H, m), 5.83 (1H, d, J 5.8), 6.25 (1H, s), 6.99 (1H, d, J 7.9), 7.18 (1H, dd, J 9.9, 2.2), 7.24-7.32 (2H, m), 7.35-7.41 (6H, m), 7.67 (1H, d, J 9.9), 8.10 (1H, s), 8.20 (1H, s).


Example 38

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



a. (R)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidin-3-ol


(Intermediate 38a)



A mixture of Intermediate 24b (300 mg, 1.74 mmol) and (R)-3-hydroxypyrrolidine (607 mg, 9.96 mmol) in NMP (4 mL) was heated in the


microwave at 160°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (170 mg, 44%). LCMS (Method 1): Rt 0.39 min, m/z 223 [MH⁺].

b. 6-Fluoro-3-((R)-3-triisopropylsilyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 38b)



Triisopropylsilyl trifluoromethanesulfonate (280 mg, 0.919 mmol) was added to a solution of Intermediate 38a (170 mg, 0.766 mmol) and Et₃N (116 mg, 1.14 mmol) in a DMF (3 mL) and the mixture stirred at RT for 1 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (260 mg, 90%). LCMS (Method 3): Rt 4.26 min, m/z 379 [MH⁺].


c. (1S,4R)-4-[3-((R)-3-Triisopropylsilyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 38c)



To a solution of Intermediate A (123 mg, 0.755 mmol) in DMF (4 mL) was added NaH (60% in oil, 82 mg, 2.06 mmol) and the mixture stirred at RT for 20 min, before Intermediate 38b (260 mg, 0.687 mmol) was added. This mixture was heated at 60°C in the microwave for 3 h. The reaction mixture was applied to an

SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a viscous yellow oil (100 mg, 23%). LCMS (Method 1): Rt 2.90 min, m/z 522 [MH⁺].

5 **d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-3-triisopropylsilyloxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 38d)**



A solution of Intermediate 38c (100 mg, 0.191 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 115 mg, 0.286 mmol) and DIPEA (100 mg, 0.764 mmol) in DMF (2 mL) was stirred at 60°C for 1 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a viscous yellow oil (80 mg, 53%). LCMS (Method 4): Rt 4.55 min, m/z 777 [MH⁺].


15 **e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-3-hydroxy-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 38)**

20 To a solution of Intermediate 38d (80 mg, 0.10 mmol) in THF (1 mL) at -30°C was added TBAF (1M in THF, 150 µL, 0.150 mmol) and the mixture was allowed to warm to RT over 1 h. The reaction mixture was applied to an SCX-2 cartridge (5 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. Purification by HPLC (C6-Ph


column, 35-75% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (22 mg, 35%). LCMS (Method 5): Rt 4.01 min, m/z 621 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.29 (9H, s), 1.86-2.13 (4H, m), 2.14-2.28 (2H, m), 2.38 (3H, s), 3.40-3.45 (1H, m), 3.49-3.56 (1H, m), 3.66-3.71 (1H, m), 3.74-3.81 (1H, m), 4.50-4.55 (1H, m), 4.85-4.91 (1H, m), 5.37 (1H, t J 4.2), 6.33 (1H, s), 7.10 (1H, dd, J 9.8, 2.2), 7.19-7.36 (8H, m), 7.43 (1H, d, J 9.8), 7.76 (1H, d, J 1.6).

Example 39

10 **1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea**



a. **5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-ylamine (Intermediate 39a)**




15

DIAD (847 μL, 4.32 mmol) was added slowly to a solution of 3-(5-amino-3-tert-butyl-1H-pyrazol-1-yl)phenol (US2006/35922; 500 mg, 2.16 mmol), 2-(tetrahydro-pyran-2-yloxy)-ethanol (439 μL, 3.25 mmol) and Ph₃P (1.13 g, 4.32 mmol) in THF (10.0 mL) and the mixture stirred for 72 h. The reaction mixture was partitioned between EtOAc (75 mL) and H₂O (75 mL) and the aqueous layer extracted with EtOAc (3 ×). The combined organic layers were dried (MgSO₄), filtered and evaporated *in vacuo*. Purification by FCC, using 5-60% EtOAc in

20

cyclohexane, gave the *title compound* (1.26 g). LCMS (Method 4): Rt 2.77, m/z 360 [MH⁺].

b. (5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 39b)




5

The *title compound* was prepared starting from 2,2,2-trichloroethylchloroformate and Intermediate 39a by using an analogous procedure to that described for Intermediate 35c. LCMS (Method 4): Rt 3.85, m/z 536 [MH⁺].

10

c. 1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 39c)



A solution of Intermediate 39b (287 mg, 0.53 mmol), Intermediate 3c (177 mg, 0.49 mmol) and DIPEA (256 μ L, 1.46 mmol) in THF (5 ml) was stirred at 60°C for 16.5 h. Water was added and the mixture extracted with DCM (3 $\times$ 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as an off-white powder (361 mg, 99%). LCMS (Method 3): Rt 4.05 min, m/z 749 [MH⁺].

20


d. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-

[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 39)


Pyridinium p-toluenesulfonate (362 mg, 1.44 mmol) was added to a solution of Intermediate 39c (360 mg, 0.48 mmol) in MeOH (5 mL). The solution was stirred at 40°C for 19 h, then diluted with water and extracted with DCM (3 × 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-7.5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a white powder after freeze-drying (218 mg, 68%). LCMS (Method 5): Rt 4.22 min, m/z 665 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.60-1.63 (2H, m), 1.70-1.75 (4H, m), 1.86-1.92 (2H, m), 2.01-2.06 (1H, m), 2.10-2.16 (1H, m), 3.14 (4H, t, J 5.2), 3.71 (2H, q, J 5.1), 4.03 (2H, t, J 5.0), 4.79-4.82 (1H, m), 4.87 (1H, t, J 5.5), 5.54 (1H, t, J 4.3), 6.33 (1H, s), 6.96 (1H, m), 7.11-7.22 (4H, m), 7.34-7.46 (5H, m), 7.61-7.64 (2H, m), 8.11 (1H, s).

Example 40

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea partial formate salt




a. Methanesulfonic acid 2-[3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-phenoxy]-ethyl ester (Intermediate 40a)



Methane sulfonyl chloride (46.0 μ L, 0.59 mmol) was added to an ice cold solution of Example 39 (187 mg, 0.28 mmol) and DIPEA (122 μ L, 0.70 mmol) in DCM (3.0 mL). After 2.5 h the reaction was partitioned between DCM and water. The aqueous layer was then extracted with DCM (3 $\times$). The combined organic layers were washed with brine, dried (MgSO_4), filtered and evaporated *in vacuo* to give the *title compound* (208 mg, 99%). LCMS (Method 2): Rt 3.38 min, m/z 743 [MH^+].

b. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea, partial formate salt (Example 40)

Dimethylamine (2M in MeOH, 1.12 mL, 2.24 mmol) was added to a solution of Intermediate 40a (208 mg, 0.28 mmol) in THF (3.0 mL). The reaction was heated to 50°C in a sealed environment overnight. Further dimethylamine (2M in MeOH, 250 μ L, 0.50 mmol) was added and heating continued for 8 h then the mixture was cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO_4), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH_3 in MeOH] in DCM. Further purification by HPLC (C18 X-select column, 20-98% MeCN in H_2O , 0.1% HCO_2H) gave the *title compound* as a white powder after freeze-drying (110 mg, 57%). LCMS (Method 5): Rt 3.47 min, m/z 692.5 [MH^+]. ^1H NMR (400 MHz, d_4 -MeOD): 1.30 (9H, s), 1.63-1.69 (2H, m), 1.73-1.80 (4H, m), 1.88-2.04 (2H, m), 2.04-2.14 (1H, m), 2.22-2.32 (1H, m), 2.60 (6H, s), 3.08-3.14 (4H, m), 3.17 (2H, t, J 5.4), 4.23 (2H, t, J 5.4), 4.89 (1H, dd, J 8.7, 5.8), 5.41 (1H, t, J 8.2), 6.34 (1H, s), 7.03-7.07 (1H, ddd, J 8.5, 2.4, 0.6), 7.09-7.13 (2H, m), 7.18-7.25 (3H, m), 7.26-7.31 (2H, m), 7.39-7.44 (1H, t, J 7.8), 7.50-7.54 (2H, m), 8.45 (0.7H, br s).


Example 41

1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

To a solution of Example 29 (100 mg, 0.15 mmol) and Et₃N (65 μ L, 0.47 mmol) in THF (7 mL) at 0°C was added mesyl chloride (19 μ L, 0.19 mmol), and the mixture stirred for 30 min. To the solution was added 1-methyl piperazine (52 μ L, 0.47 mmol) and the solution heated at reflux for 35 min. Water was added and the mixture extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by HPLC (Gemini C18 column, 30-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying (32 mg, 28%). LCMS (Method 5): Rt 3.52 min, m/z 717 [MH⁺]. ¹H NMR (400 MHz, CDCl₃): 1.34 (9H, s), 1.66 (2H, q, J 5.4), 1.72-1.76 (4H, m), 1.90-1.99 (1H, m), 2.04-2.13 (3H, m), 2.23 (3H, s), 2.39-2.55 (10H, m), 3.16 (4H, t, J 5.2), 3.60 (1H, d, J 13.4), 3.62 (1H, d, J 13.4), 5.13 (1H, td, J 8.7, 5.3), 5.21 (1H, t, J 4.4), 6.00 (1H, br s), 6.37 (1H, s), 6.91 (1H, br s), 6.97-6.98 (1H, m), 7.23 (1H, d, J 7.5), 7.29-7.54 (9H, m).


Example 42

**1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
 {(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**



5

**a. 1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-
 2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-
 [1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
 urea (Intermediate 42a)**



10

A mixture of Intermediate 5c (150 mg, 0.410 mmol) and Intermediate 39b (230 mg, 0.430 mmol) in 1,4-dioxane (4 mL) and DIPEA (113 μ L, 0.65 mmol) was stirred at 90°C for 4.5 h. The mixture was concentrated *in vacuo*. The residue was purified by FCC twice, using 0-10% MeOH in DCM, to give the *title compound* as a pale yellow foam (234 mg, 76%). LCMS (Method 3): Rt 3.03 min, m/z 749.2 [MH⁺].

15

**b. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
 {(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 42)**


20

A mixture of Intermediate 42a (230 mg, 0.300 mmol) and pyridinium p-toluenesulfonate (226 mg, 0.900 mmol) in MeOH (3 mL) was stirred at RT for 15


h. Pyridinium p-toluenesulfonate (113 mg, 0.450 mmol) was added and the mixture stirred at 50-55°C for 8 h. The cooled mixture was diluted with sat. aq. NaHCO₃ solution and extracted with DCM (3 × 15 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the *title compound* as a white powder after freeze-drying (163 mg, 82%). LCMS (Method 5): Rt 3.33 min, m/z 665.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.82-2.27 (11H, m), 2.31-2.39 (1H, m), 3.10-3.17 (1H, m), 3.71 (2H, q, J 5.1), 3.96-4.05 (3H, m), 4.79-4.89 (2H, m), 5.37-5.41 (1H, m), 6.34 (1H, s), 6.95-6.99 (1H, m), 7.05-7.15 (3H, m), 7.24-7.43 (6H, m), 7.75 (1H, d, J 9.9), 8.11 (1H, s), 8.25 (1H, br d).

Example 43

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[(2-dimethylamino-ethyl)-methyl-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea




a. N-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-N,N',N'-trimethylethane-1,2-diamine (Intermediate 43a)



A mixture of Intermediate 24b (300 mg, 1.74 mmol) and N,N,N-trimethylethane-1,2-diamine (900 mg, 8.77 mmol) in NMP (2 mL) was heated in the microwave at 170°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown oil (190 mg, 46%). LCMS

(Method 4): Rt 0.38 min, m/z 238 [MH⁺].

b. N-[6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl]-N,N',N'-trimethyl-ethane-1,2-diamine (Intermediate 43b)



5

To a solution of Intermediate A (143 mg, 0.88 mmol) in DMF (3 mL) was added NaH (60% in oil, 96 mg, 2.4 mmol) and the mixture stirred at RT for 20 min, before Intermediate 43a (190 mg, 0.80 mmol) was added. This mixture was heated at 60°C in the microwave for 1 h. The reaction mixture was applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-20% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a pale brown oil (140 mg, 46%). LCMS (Method 1): Rt 1.34 min, m/z 381 [MH⁺].

10

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[(2-dimethylamino-ethyl)-methyl-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Example 43)

15


A solution of Intermediate 43b (140 mg, 0.36 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 218 mg, 0.54 mmol) and DIPEA (186 mg, 1.44 mmol) in DMF (6 mL) was stirred at 60°C for 1 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave a viscous yellow gum. HPLC (C6-Ph column, 10-40% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as an off-white powder after freeze-drying (80 mg, 35%). LCMS (Method 5): Rt 3.67 min, m/z 636.3 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.30 (9H, s), 1.86-2.15 (3H,

20

25

m), 2.19-2.30 (1H, m), 2.38 (3H, s), 2.57 (6H, s), 2.92 (3H, s), 3.00 (2H, t J 6.2), 3.46-3.60 (2H, m), 4.89 (1H, m), 5.42 (1H, t, J 3.9), 6.33 (1H, s), 7.16-7.36 (9H, m), 7.52 (1H, d, J 10.1), 7.90 (1H, s).

Example 44



5

1-[5-tert-Butyl-2-(3-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

To a solution of Example 29 (100 mg, 0.15 mmol) and Et₃N (65 μ L, 0.47 mmol) in THF (7 mL) at 0°C was added mesyl chloride (19 μ L, 0.19 mmol), and the mixture stirred for 0.5 h. To the solution was added piperidine (47 μ L, 0.47 mmol), and the mixture heated at reflux for 16 h. NaI (24 mg, 0.15 mmol) was added and the solution heated at reflux for 1 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by HPLC (XBridge C18 column, 20-98% MeCN in H₂O, 0.1% HCO₂H) to give the impure product. The residue was re-purified by HPLC (Gemini C18 column, 20-98% MeCN in H₂O, 0.1% HCO₂H), the product containing fractions were partitioned between DCM and saturated NaHCO₃ solution. The combined organics were dried and concentrated *in vacuo* to give the *title compound* as a white powder (14 mg, 12%). LCMS (Method 5): Rt 3.60 min, m/z 702 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.34 (9H, s), 1.39-1.44 (2H, m), 1.52-1.55 (4H, m), 1.67-1.71 (2H, m), 1.77-1.80 (4H, m), 2.03-2.08 (3H, m), 2.24-2.33 (1H, m), 3.20 (4H, t, J 5.3), 3.55 (2H, s), 4.93 (1H, dd, J 8.6, 5.6), 5.44 (1H, t, J 4.2), 6.38 (1H, s), 7.21-7.33 (5H, m), 7.40-

10


15

20

7.56 (6H, m).


Example 45

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[methyl-(2-morpholin-4-yl-ethyl)-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea




a. (2-Morpholin-4-yl-ethyl)-carbamic acid tert-butyl ester

(Intermediate 45a)



4-(2-Aminoethyl)morpholine (5.00 g, 38.5 mmol) was added to a stirred mixture of di-*tert*-butyl dicarbonate (8.38 g, 38.5 mmol) and indium(III)chloride (85 mg, 0.39 mmol). This mixture was stirred for 1 min before being diluted with EtOAc (200 mL) and washed with water ($\times$ 2). The organic layer was dried (MgSO₄) and was evaporated *in vacuo* to give the *title compound* as a clear oil (8.14 g, 92%). ¹H NMR (400 MHz, CDCl₃): 1.45 (9H, s), 2.40-2.50 (6H, m), 3.23 (2H, q, J 5.9), 3.70 (4H, t, J 4.7), 4.96 (1H, br s).


b. Methyl-(2-morpholin-4-yl-ethyl)-amine (Intermediate 45b)



To a solution of Intermediate 45a (6.0 g, 24.6 mmol) in THF (123 mL) was added lithium aluminium hydride (2.33 g, 61.5 mmol) portionwise, then the mixture was stirred at reflux for 4 h. A further portion of lithium aluminium hydride (1.28 g, 33.6 mmol) was added to the cooled solution, then the mixture was stirred at reflux for a further 8 h. Water (3.75 mL) was added dropwise, followed by 4N aqueous NaOH (11.3 mL) and water (3.75 mL). The mixture was


filtered, washed with diethyl ether, dried (MgSO₄) and concentrated *in vacuo* to give the *title compound* as a yellow oil (3.67 g, 99%). ¹H NMR (400 MHz, CDCl₃): 2.40-2.52 (9H, m), 2.67 (2H, t, J 2.7), 3.70 (4H, t, J 4.5).

c. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-methyl-(2-morpholin-4-yl-ethyl)-amine (Intermediate 45c)



A mixture of Intermediate 24b (300 mg, 1.74 mmol) and Intermediate 45b (1.26 g, 8.77 mmol) in NMP (2 mL) was heated in the microwave at 170°C for 2 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as an orange oil (350 mg, 71%). ¹H NMR (400 MHz, CDCl₃): 2.41-2.52 (4H, m), 2.58 (2H, t, J 5.6), 3.03 (3H, s), 3.27 (2H, t, J 6.0), 3.62 (4H, t, J 5.2), 7.04-7.08 (1H, m), 7.58-7.65 (1H, m), 8.39 (1H, t, J 2.4).

d. [6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl]-methyl-(2-morpholin-4-yl-ethyl)-amine (Intermediate 45d)



To a solution of Intermediate A (225 mg, 1.38 mmol) in DMF (3 mL) was added NaH (60% in oil, 150 mg, 3.75 mmol) and the mixture stirred at RT for 20 min, before Intermediate 45c (350 mg, 1.25 mmol) was added. This mixture was stirred at 60°C for 7 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-20% [2M NH₃ in MeOH] in DCM, gave the *title compound* as an orange oil (270 mg, 51%). LCMS (Method

4): Rt 0.22 min, m/z 423 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[methyl-(2-morpholin-4-yl-ethyl)-amino]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Example 45)


5 A solution of Intermediate 45d (270 mg, 0.63 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 380 mg, 0.95 mmol) and DIPEA (325 mg, 2.52 mmol) in DMF (8 mL) was stirred at 60°C for 1 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was
10 eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave a stiff yellow gum which was further purified by HPLC (C18 column, 10-45% MeCN in H₂O, 0.1% HCO₂H). The residue was recrystallized from boiling EtOAc to give the *title compound* as a white powder (60.0 mg, 14%). LCMS (Method 5): Rt 3.65 min, m/z 678.3 [MH⁺].
15 ¹H NMR (400 MHz, d₄-MeOH): 1.30 (9H, s), 1.87-2.15 (3H, m), 2.18-2.27 (1H, m), 2.32 (4H, t, J 4.4), 2.38 (3H, s), 2.55 (2H, t, J 5.6), 2.96 (3H, s), 3.31-3.40 (6H, m), 4.89 (1H, m), 5.40 (1H, t, J 4.2), 6.33 (1H, s), 7.19-7.36 (9H, m), 7.53 (1H, d, J 10.0), 7.92 (1H, d, J 1.6).

20


25

Example 46

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea and 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (1:1 mixture of diastereomers)




a. 2-Morpholin-4-yl-propionic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide
(Intermediate 46a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (1.62 g, 12.8 mmol), 2-(morpholin-4-yl)propanoic acid (Enamine, 2.50 g, 12.8 mmol), HOBt.H₂O (196 mg, 1.28 mmol) and Et₃N (2.14 mL, 15.3 mmol) in DCM (50 mL) was added EDC (2.94 g, 15.3 mmol) and the brown solution stirred at RT for 16 h. Water (20 mL) and brine (20 mL) were added, and the mixture shaken. The aqueous layer was extracted with DCM (20 mL), then the combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to give the *title compound* as a brown foam (3.43 g, >99%). LCMS (Method 3): Rt 0.44 min, m/z 269 [MH⁺].


b. 6-Fluoro-3-(1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridine
(Intermediate 46b)



To a solution of Intermediate 46a (3.43 g, 12.8 mmol), Ph₃P (6.71 g, 25.6

mmol) and Et₃N (7.14 mL, 51.2 mmol) in THF (75 mL) at 0°C was added hexachloroethane (6.06 g, 25.6 mmol) and the solution stirred at RT for 4 h. The brown suspension was filtered and the solid washed with THF (10 mL). The combined organics were applied to an SCX-2 cartridge, which was washed with DCM-MeOH (1:1, 100 mL) and MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave an off-white solid. FCC, using 1-5% [2M NH₃ in MeOH] in DCM, gave the *title compound* as an off-white solid (1.70 g, 53%). ¹H NMR (300 MHz, CDCl₃): 1.66 (3H, d, J 6.8), 2.45-2.60 (4H, m), 3.63-3.76 (4H, m), 4.28 (1H, q, J 6.8), 7.21 (1H, ddd, J 10.0, 7.5, 2.3), 7.76 (1H, ddd, J 10.0, 4.9, 0.9), 8.50 (1H, ddd, J 3.9, 2.3, 0.9).

c. (1S,4R)-4-[3-((R)-1-Morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine and (1S,4R)-4-[3-((S)-1-Morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylam (1:1 mixture diastereomers, Intermediate 46c)



15

A suspension of Intermediate A (343 mg, 2.10 mmol) and NaH (60% dispersion in oil, 240 mg, 6.00 mmol) in dry DMF (10 mL) at RT under Ar was stirred for 30 min (CARE: gas evolution), then Intermediate 46b (501 mg, 2.00 mmol) was added and the brown mixture stirred at 60°C under Ar for 1 h. The solution was concentrated *in vacuo*, redissolved in MeOH (4 mL) and AcOH (0.60 mL, 10.0 mmol), then applied to an SCX-2 cartridge (20 g) and washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (50 mL); concentration *in vacuo* gave a dark brown foam. FCC, using 2-8% [2M NH₃ in MeOH] in DCM, gave the *title compounds* as a light brown foam (558 mg, 71%). LCMS (Method 3): Rt 0.44 min, m/z 394 [MH⁺].

25


d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-1-

morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea and 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(S)-1-morpholin-4-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (1:1 mixture of diastereomers). (Example 46)


A yellow solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 223 mg, 0.550 mmol), Intermediate 46c (197 mg, 0.500 mmol) and DIPEA (0.109 mL, 0.625 mmol) in dioxane (10 mL) was stirred at 60°C for 18 h. The cooled solution was concentrated *in vacuo*, suspended in water (10 mL) and extracted with DCM (2 × 10 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a yellow-brown gum. FCC, using 2-7% [2M NH₃ in MeOH] in DCM, gave the *title compounds* as a pale yellow solid (192 mg, 59%). LCMS (Method 5): two peaks in 1:1 ratio, Rt 4.13 and 4.15 min, m/z 649 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9 H, s), 1.52 (1.5 H, d, J 6.8), 1.53 (1.5 H, d, J 6.8), 1.83-1.99 (2 H, m), 2.10-2.17 (2 H, m), 2.36 (3 H, s), 2.43 (2 H, t, J 4.5), 2.51-2.55 (2 H, m), 3.47-3.55 (4 H, m), 4.46 (0.5 H, q, J 6.8), 4.49 (0.5 H, q, J 6.8), 4.83 (1 H, m), 5.45 (0.5 H, t, J 4.5), 5.52 (0.5 H, t, J 4.5), 6.33 (1 H, s), 7.10 (1 H, d, J 8.5), 7.21-7.44 (9 H, m), 7.73 (1 H, dd, J 9.9, 2.6), 8.03 (1 H, s), 8.32 (0.5 H, d, J 2.1), 8.37 (0.5 H, d, J 2.1).

Example 47

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(S)-1,2-dimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (S)-1,2-Dimethyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 47a)




To a suspension of (S)-1,2-dimethyl-pyrrolidine-2-carboxylic acid (660 mg, 4.61 mmol), (5-fluoro-pyridin-2-yl)-hydrazine (488 mg, 3.84 mmol) and Et₃N (1.1 mL, 7.7 mmol) in DCM (20 mL) were added HOBt.H₂O (51 mg, 0.38 mmol) and EDCI.HCl (884 mg, 4.61 mmol) and the mixture stirred at RT overnight. The mixture was partitioned between EtOAc/water and extracted with EtOAc. The combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo*. FCC, using 0-10% MeOH in DCM, gave the *title compound* as a brown solid (668 mg, 69%). LCMS (Method 4): Rt 0.40, m/z 253 [MH⁺].

b. 3-((S)-1,2-Dimethyl-pyrrolidin-2-yl)-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 47b)



To a solution of Intermediate 47a (668 mg, 2.64 mmol), Et₃N (1.5 mL, 11 mmol) and Ph₃P (1.4 g, 5.3 mmol) in THF (20 mL) at 0°C was added hexachloroethane (1.25 g, 5.29 mmol). The mixture was stirred at 0°C for 10 min, then at RT for 1 h. The mixture was partitioned between EtOAc/water and the aqueous extracted with EtOAc. The combined organics were dried (Na₂SO₄), filtered and concentrated *in vacuo*. Purification by FCC, using 0-10% MeOH in EtOAc, gave the *title compound* as a pale brown solid (300 mg, 49%). LCMS (Method 4): Rt 0.39, m/z 235 [MH⁺].


c. (1S,4R)-4-[3-((S)-1,2-Dimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 47c)



5 To a solution of Intermediate 47b (150 mg, 0.64 mmol) and Intermediate A (126 mg, 0.64 mmol) in DMF (3 mL) was added NaH (60% in mineral oil, 90 mg, 2.24 mmol) portionwise. The mixture was stirred at 60°C for 1.5 h then allowed to cool to RT. The mixture was carefully quenched by pouring into MeOH (10 mL), then applied to an SCX-2 cartridge and washed with MeOH. The product was
 10 eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown solid (180 mg, 75%). LCMS (Method 4): Rt 0.49, m/z 378 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1,2-dimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 47)

The *title compound* was prepared as an off-white solid (80 mg, 53%) using (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 80 mg, 0.20 mmol) and Intermediate 47c (90 mg, 0.24 mmol) in a similar manner to Example 1, step d.
 20 LCMS (Method 5): Rt 3.83 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.51 (3H, s), 1.78-1.99 (5H, m), 2.02 (3H, s), 2.03-2.22 (3H, m), 2.36 (3H, s), 2.65 (1H, q, J 8.6), 3.11-3.17 (1H, m), 4.79-4.86 (1H, m), 5.32 (1H, t, J 4.2), 6.32 (1H, s), 7.11 (1H, d, J 8.6), 7.25-7.39 (9H, m), 7.75 (1H, d, J 9.8), 8.05 (1H, s), 8.43 (1H, d, J 2.2).


Example 48

1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxymethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

To a solution of Example 29 (204 mg, 0.32 mmol) and Et₃N (134 μ L, 0.97 mmol) in DCM (3.2 mL) at 0°C was added mesyl chloride (39 μ L, 0.39 mmol), and the mixture stirred for 1 h. The solution was washed with water, dried, and concentrated *in vacuo* to give methanesulfonic acid 3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-benzyl ester. This was immediately dissolved in dioxane (1.5 mL) and ethylene glycol (1.5 mL) and the solution stirred at 85°C for 1 h. The cooled reaction mixture was suspended in water and filtered to leave a solid. Purification by HPLC (XBridge C18 column, 40-98% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (59 mg, 27%). LCMS (Method 5): Rt 4.27 min, m/z 679 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.60-1.64 (2H, m), 1.70-1.75 (4H, m), 1.86-1.93 (2H, m), 2.03 (1H, m), 2.11 (1H, m), 3.14 (4H, t, J 5.2), 3.54-3.48 (4H, m), 4.56 (2H, s), 4.64 (1H, s), 4.80 (1H, m), 5.54 (1H, t, J 4.3), 6.34 (1H, s), 7.07 (1H, d, J 8.6), 7.16 (1H, dd, J 9.8, 2.2), 7.37-7.49 (8H, m), 7.60-7.63 (2H, m), 8.11 (1H, s).


Example 49

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-[1,4]oxazepan-4-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



5


a. 6-Fluoro-3-[1,4]oxazepan-4-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 49a)



A solution of Intermediate 24b (429 mg, 2.50 mmol) and homomorpholine (939 mg, 9.30 mmol) in NMP (10 mL) was heated in the microwave at 170°C for 10 h. The cooled mixture was applied to an SCX-2 cartridge (70 g), washing with methanol then eluting basic components with 0.4–2 M NH₃ in MeOH. Product containing fractions were combined and concentrated *in vacuo*. The residue was purified by FCC, using 0–12% [2M NH₃ in MeOH] in DCM, to give impure product. Further purified by FCC, using 0–12% MeOH in EtOAc gave the *title compound* as a pale brown gum (147 mg, 25%). LCMS (Method 3): Rt 2.11 min, m/z 237 [MH⁺].

15

b. (1S,4R)-4-(3-[1,4]Oxazepan-4-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 49b)




20

To a solution of Intermediate A (145 mg, 0.614 mmol) in DMF (3 mL) was

added NaH (60% dispersion in oil, 74 mg, 1.84 mmol). The mixture was stirred at RT for 15 min, then a solution of Intermediate 49a (100 mg, 0.614 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 2.25 h. The cooled mixture was diluted with water and extracted with DCM (4 × 25 mL). The combined
5 organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-12% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale brown gum (93 mg, 40%). LCMS (Method 3): Rt 0.43 min, m/z 380 [MH⁺].

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-[1,4]oxazepan-4-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 49)
10

A solution of Intermediate 49b (90.0 mg, 0.237 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 97 mg, 0.24 mmol) in 1,4-dioxane (3 mL) and DIPEA (63 µL, 0.36 mmol) was stirred at 90°C for 4 h, then more DIPEA (63
15 µL, 0.36 mmol) was added and mixture stirred at 100-110°C for 3.5 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-12% [2M NH₃ in MeOH] in DCM, to give the product (31 mg, 21%). This was further purified by HPLC (C18 X-select column, 35-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as a white powder after freeze-drying (20 mg, 13%).
20 LCMS (Method 5): Rt 4.40 min, m/z 635.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.80-2.18 (6H, m), 2.36 (3H, s), 3.44-3.51 (4H, m), 3.78-3.85 (4H, m), 4.77-4.85 (1H, m), 5.51 (1H, t, J 4.4), 6.32 (1H, s), 7.10-7.16 (2H, m), 7.25-7.40 (8H, m), 7.60 (1H, d, J 9.6), 7.65 (1H, d, J 1.5), 8.10 (1H, s).


Example 50

1-(5-tert-Butyl-2-{3-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea


To a solution of Example 29 (150 mg, 0.24 mmol) and Et₃N (98 μ L, 0.71 mmol) in DCM (7 mL) at 0°C was added mesyl chloride (28 μ L, 0.28 mmol), and the mixture stirred for 1 h. The solution was washed with water, dried, and concentrated *in vacuo* to give methanesulfonic acid 3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-benzyl ester. This was immediately dissolved in THF (2.5 mL) and DIPEA (83 μ L, 0.47 mmol) then 1-(2-hydroxyethyl) piperazine (87 μ L, 0.71 mmol) added and the solution stirred at reflux for 1 h. Water was added and the mixture extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by HPLC (XBridge C18 column, 20-98% MeCN in H₂O, 0.1% NH₄OH) to give the *title compound* as a white powder after freeze-drying (45 mg, 25%). LCMS (Method 5): Rt 3.42 min, m/z 747 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.60-1.64 (2H, m), 1.69-1.75 (4H, m), 1.83-1.94 (2H, m), 2.03-2.12 (2H, m), 2.29-2.45 (8H, m), 2.33 (2H, t, J 6.4), 3.14 (4H, t, J 5.2), 3.44 (2H, m), 3.49 (2H, s), 4.31 (1H, s), 4.80 (1H, m), 5.54 (1H, t, J 4.4), 6.33 (1H, s), 7.03 (1H, d, J 8.6), 7.16 (1H, dd, J 9.8, 2.2), 7.34-7.46 (8H, m), 7.60-7.63 (2H, m), 8.10 (1H, s).

Example 51

1-(2-tert-Butyl-5-p-tolyl-3H-imidazol-4-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




5

a. 2-tert-Butyl-5-p-tolyl-1H-imidazole (Intermediate 51a)

A red suspension of 2,2-dimethylpropionamide hydrochloride (Atlantic, 1.00 g, 7.32 mmol), 2-bromo-4'-methylacetophenone (Aldrich, 1.56 g, 7.32 mmol) and K_2CO_3 (1.52 g, 11.0 mmol) in DMF (20 mL) was stirred at 75°C for 3 h. The mixture was cooled to RT, concentrated *in vacuo*, suspended in water (25 mL) and extracted with DCM (2 × 25 mL). The combined organics were passed through a hydrophobic frit, concentrated *in vacuo*, redissolved in MeOH (5 mL), applied to an SCX-2 cartridge (20 g) and washed with MeOH (100 mL). The product was eluted with 2M NH_3 in MeOH (60 mL); concentration *in vacuo* left a yellow foam (1.26 g). FCC, using 1-5% MeOH in DCM, gave the *title compound* as a pale yellow solid (770 mg, 49%). LCMS (Method 3): Rt 2.26 min, m/z 215 $[MH^+]$.

10

15


b. 2-tert-Butyl-4-nitroso-5-p-tolyl-1H-imidazole (Intermediate 51b)

20

A yellow solution of Intermediate 51a (390 mg, 1.82 mmol) and isopentyl nitrite (0.294 mL, 2.18 mmol) in THF (10 mL) was stirred at 50°C for 8 h, then cooled to RT and concentrated *in vacuo* to leave a dark green oil. FCC, using 0-


25% EtOAc in cyclohexane, gave the *title compound* as a green solid (154 mg, 35%). LCMS (Method 3): Rt 3.82 min, m/z 244 [MH⁺].

c. 2-tert-Butyl-5-p-tolyl-1H-imidazol-4-ylamine (Intermediate 51c)



5 A suspension of Intermediate 51b (150 mg, 0.617 mmol) and Pd/C (10%, 15 mg) in EtOH (10 mL) was stirred at RT under H₂ for 2 h. The flask was evacuated and purged with N₂ thrice, then the mixture filtered through Celite, and the filter-cake washed with EtOH (10 mL). The combined organics were concentrated *in vacuo* to leave a red-brown gum. FCC, using 1-6% MeOH in
10 DCM, gave the *title compound* as a yellow-orange gum (96.3 mg, 68%). LCMS (Method 3): Rt 2.26 min, m/z 230 [MH⁺].

d. (2-tert-Butyl-5-p-tolyl-1H-imidazol-4-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 51d)




15 To a solution of Intermediate 51c (45.3 mg, 0.197 mmol) in EtOAc (1 mL) and aqueous NaOH (1M, 0.49 mL) was added 2,2,2-trichloroethyl chloroformate (0.0326 mL, 0.237 mmol) and the resulting mixture stirred at RT for 45 min. The layers were separated and the aqueous extracted with EtOAc (2 mL). The combined organics were washed with brine (2 mL), dried (Na₂SO₄), filtered and
20 concentrated *in vacuo* to leave an orange-red gum. FCC, using 0-50% EtOAc in cyclohexane, gave the *title compound* as a pale orange film (44.1 mg, 55%). LCMS (Method 3): Rt 2.87 min, m/z 404, 406 [MH⁺].

e. 1-(2-tert-Butyl-5-p-tolyl-3H-imidazol-4-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-

yl]-urea. (Example 51)

An orange solution of Intermediate 51d (65.4 mg, 0.162 mmol), Intermediate 3c (55.9 mg, 0.153 mmol) and DIPEA (0.042 mL, 0.242 mmol) in dioxane (2 mL) was stirred at 60°C for 18 h. After cooling, the mixture was concentrated *in vacuo*, suspended in water (5 ml) and extracted with DCM (2 × 5 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave an orange-brown gum. FCC, using 2-8% MeOH in DCM, gave a pale yellow solid (50.5 mg). Further purification by HPLC (C18 XBridge column, 25-75% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white solid after freeze-drying (33.3 mg, 33%). LCMS (Method 5): Rt 3.35 min, m/z 619 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 3:1 ratio of tautomers: 1.30 (6.75H, s), 1.31 (2.25H, s), 1.61 (2H, m), 1.72 (4H, m), 1.83-2.00 (2H, m), 2.05 (1H, m), 2.15 (1H, m), 2.30 (0.75H, s), 2.32 (2.25H, s), 3.13 (4H, q, J 5.2), 4.87 (1H, m), 5.54 (1H, t, J 4.3), 7.14-7.19 (2.5H, m), 7.21 (1.5H, d, J 8.2), 7.25-7.40 (4H, m), 7.54 (1.5H, d, J 8.3), 7.60-7.71 (3.5H, m), 11.56 (0.75H, s), 11.68 (0.25H, s).


Example 52

1-{5-tert-Butyl-2-[3-(4-hydroxy-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

To a solution of Example 29 (150 mg, 0.24 mmol) and Et₃N (98 μL, 0.71 mmol) in DCM (7 mL) at 0°C was added mesyl chloride (28 μL, 0.28 mmol), and

the mixture stirred for 1 h. The solution was washed with water, dried, and concentrated *in vacuo* to give methanesulfonic acid 3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-benzyl ester. This was immediately dissolved in THF (2.5 mL) and DIPEA (83 μ L, 0.47 mmol) then 4-hydroxypiperidine (83 μ L, 0.71 mmol) added and the solution stirred at reflux for 1 h. Water was added and the mixture extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the product. This was purified further by HPLC (XBridge C18 column, 20-98% MeCN in H₂O, 0.1% NH₄OH) to give the *title compound* as a white powder after freeze-drying (30 mg, 18%). LCMS (Method 5): Rt 3.41 min, m/z 718 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.34-1.40 (2H, m), 1.60-1.75 (7H, m), 1.81-1.96 (3H, m), 2.00-2.15 (4H, m), 2.64-2.69 (2H, m), 3.14 (4H, t, J 5.2), 3.40-3.45 (1H, m), 3.47 (2H, s), 4.51 (1H, d, J 3.9), 4.82 (1H, m), 5.54 (1H, t, J 4.4), 6.33 (1H, s), 7.04 (1H, d, J 8.5), 7.16 (1H, dd, J 9.8, 2.2), 7.26-7.33 (4H, m), 7.35-7.46 (4H, m), 7.60-7.63 (2H, m), 8.10 (1H, s).

Example 53




1-{5-tert-Butyl-2-[3-(2-hydroxy-ethylsulfanyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

A solution of Intermediate 3c (107 mg, 0.29 mmol) and Intermediate 53c (151 mg, 0.32 mmol) in THF (3 mL) and DIPEA (169 μ L, 0.97 mmol) was stirred at reflux for 20.5 h. The cooled reaction mixture was diluted with water and


extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give the product. This residue purified further by HPLC (XBridge C18 column, 30-98% MeCN in H₂O, 0.1% NH₄OH) to give the *title compound* as a white powder after freeze-drying (21 mg, 10%). LCMS (Method 5): Rt 4.42 min, m/z 681 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.60-1.63 (2H, m), 1.69-1.74 (4H, m), 1.86-1.93 (2H, m), 2.03 (1H, m), 2.13 (1H, m), 3.09 (2H, t, J 6.8), 3.14 (4H, t, J 5.2), 3.59 (2H, q, J 6.2), 4.80 (1H, m), 4.95 (1H, t, J 5.6), 5.54 (1H, t, J 4.3), 6.33 (1H, s), 7.07 (1H, d, J 8.6), 7.16 (1H, dd, J 9.8, 2.1), 7.35-7.47 (8H, m), 7.60-7.64 (2H, m), 8.14 (1H, s).

Example 54

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(4-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. [1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-4-yl]-methanol (Intermediate 54a)



A solution of Intermediate 24b (593 mg, 3.45 mmol) and 4-piperidinemethanol (1.59 g, 13.8 mmol) in NMP (10 mL) was heated in the microwave at 170°C for 3 h. The cooled mixture was applied to an SCX-2 cartridge (70 g), washing with methanol then eluting basic components with 0.4 –


2 M NH_3 in MeOH. Product containing fractions were combined and concentrated *in vacuo*. The residue was purified by FCC, using 0-15% MeOH in EtOAc, to give the *title compound* as a brown gum (481 mg, 56%). LCMS (Method 3): Rt 2.12 min, m/z 251 $[\text{MH}^+]$.

5 **b. 6-Fluoro-3-(4-triisopropylsilanyloxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin (Intermediate 54b)**



To a solution of Intermediate 54a (470 mg, 1.88 mmol) and Et_3N (390 μL , 2.82 mmol) in DCM (5 mL) was added triisopropylsilyltrifluoromethane sulfonate
10 (607 μL , 2.26 mmol) and the mixture stirred at RT for 0.5 h. The mixture was washed with sat. aq. NaHCO_3 solution, dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane, then 10% MeOH in EtOAc, to give the *title compound* as a pale yellow solid (565 mg, 74%). LCMS (Method 3): Rt 5.21 min, m/z 407 $[\text{MH}^+]$.


15 **c. (1S,4R)-4-[3-(4-Triisopropylsilanyloxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 54c)**



To a solution of Intermediate A (223 mg, 1.37 mmol) in DMF (3 mL) was
20 added NaH (60% dispersion in oil, 168 mg, 4.20 mmol) and the mixture stirred at RT for 15 min. A solution of Intermediate 54b (555 mg, 1.37 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 1.75 h. The cooled mixture was

diluted with water and extracted with DCM (5×20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% [2M NH_3 in MeOH] in DCM, to give the *title compound* as a brown gum (344 mg, 46%). LCMS (Method 3): Rt 3.29 min, m/z 550 [MH^+].

5 **d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-triisopropylsilyloxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 54d)**



A solution of Intermediate 54c (171 mg, 0.311 mmol) and (5-tert-butyl-2-p-
10 tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 132 mg, 0.327 mmol) in 1,4-dioxane (3 mL) and DIPEA (87 μL , 0.50 mmol) was stirred at 90°C for 3 h, and then at 100°C for 2.5 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-8% MeOH in DCM, to give the *title compound* as a pale brown
15 solid (215 mg, 86%). LCMS (Method 3): Rt 5.70 min, m/z 805.4 [MH^+].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 54)


A solution of Intermediate 54d (210 mg, 0.261 mmol) and TBAF (1M in
20 THF, 0.31 mL, 0.31 mmol) in THF (5 mL) was stirred at RT for 1.25 h. The mixture was diluted with water and extracted with DCM (3×15 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-16% MeOH in DCM, to give the *title compound* as an off-white powder after freeze-drying (134 mg, 79%). LCMS (Method 5): Rt 4.24 min, m/z

649.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.38-2.16 (9H, m), 2.36 (3H, s), 2.85-2.97 (2H, m), 3.28-3.49 (4H, m), 4.52 (1H, t, J 5.2), 4.78-4.85 (1H, m), 5.55 (1H, t, J 4.3), 6.32 (1H, s), 7.07 (1H, d, J 8.6), 7.15 (1H, dd, J 9.7, 2.1), 7.25-7.41 (8H, m), 7.60 (1H, d, J 9.7), 7.64 (1H, d, J 1.6), 8.04 (1H, s).

5


Example 55

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4S)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



10

a. (1S,4S)-4-(3-Piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 55a)



Intermediate B (74.0 mg, 454 μmol) was added to a mixture of NaH (60% in mineral oil, 27.2 mg, 681 μmol) in DMF (2.5 mL) at RT and stirred for 15 min.

15 Intermediate 3b (100 mg, 454 μmol) was then added and the resulting mixture heated to 60°C for 1 h. After cooling, the reaction was quenched with sat. aq. NH₄Cl solution. The mixture was diluted with water and extracted with EtOAc. The combined organic extracts were concentrated *in vacuo*. The resulting residue was purified by FCC, using 0 to 10% [2M NH₃ in MeOH] in DCM, to afford the
20 *title compound* as an orange residue (122 mg, 74%). LCMS (Method 3): Rt 2.13 min, m/z 364 [MH⁺].


c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4S)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-

urea (Example 55)


A mixture of Intermediate 55a (120 mg, 0.33 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 147 mg, 0.36 mmol), DIPEA (86.0 μ L, 0.50 mmol) and 1,4-dioxane (2.5 mL) were heated to 60°C for 18 h. After cooling the solvent was evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give a pale yellow foam. This was lyophilised to provide the *title compound* as an off-white solid (28 mg, 20%). LCMS (Method 5) Rt 4.76 min, m/z 619 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 1.26 (9H, s), 1.61 (2H, m), 1.72 (5H, m), 2.02-2.17 (3H, m), 2.35 (3H, s), 3.13 (4H, m), 4.90 (1H, m), 5.58 (1H, m), 6.31 (1H, s), 6.99 (1H, d, J 7.8), 7.19 (1H, dd, J 9.7, 2.2), 7.26-7.43 (8H, m), 7.58-7.64 (2H, m), 7.98 (1H, s).

Example 56

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(3-hydroxymethyl-4-methyl-piperazin-1-yl)methyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



a. (3-Hydroxymethyl-4-methyl-piperazin-1-yl)-acetic acid ethyl ester (Intermediate 56a)




20

A mixture of (1-methyl-2-piperazinyl)methanol dihydrochloride (500 mg, 2.46 mmol), ethyl bromoacetate (410 mg, 2.46 mmol), K_2CO_3 (1.02 g, 7.38 mmol) and MeCN (15 mL) were heated to reflux for 18 h. After cooling, the mixture was

filtered and concentrated *in vacuo*. The *title compound* was isolated as a colourless oil (525 mg, 98%) and used without further purification. ¹H NMR (300 MHz, CDCl₃): 1.25 (3H, t, J 7.0), 2.35 (3H, s), 2.35-2.59 (4H, m), 2.76-2.90 (3H, m), 3.19 (2H, m), 3.44 (1H, dd, J 11.3, 2.0), 3.87 (1H, dd, J 11.3, 4.0), 4.15 (2H, q, J 7.1).


b. Sodium (4-methyl-3-triisopropylsilanyloxymethyl-piperazin-1-yl)-acetate (Intermediate 56b)



To a mixture of Intermediate 56a (520 mg, 2.40 mmol), Et₃N (485 mg, 4.80 mmol) and DCM (15 mL), triisopropylsilyl trifluoromethanesulfonate (774 μL, 2.88 mmol) was added. The mixture was stirred for 1 h. The mixture was diluted with DCM and washed with water, sat. aq. NaHCO₃ solution and brine, and concentrated *in vacuo*. The *title compound* was isolated as a pale yellow oil (882 mg, quant.) and used without further purification.

A mixture of crude silyl-protected product (440 mg, 1.18 mmol), NaOH (47.0 mg, 1.18 mmol), water (10 mL) and THF (10 mL) were stirred for 18 h. The mixture was concentrated *in vacuo* to give the *title compound* as a pink solid (393 mg, 91%). LCMS (Method 3): Rt 2.81 min, m/z 345 [M-Na+2H⁺].


c. (4-Methyl-3-triisopropylsilanyloxymethyl-piperazin-1-yl)-acetic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 56c)



To a mixture of Intermediate 56b (390 mg, 1.06 mmol), (5-fluoro-pyridin-2-yl)-hydrazine (135 mg, 1.06 mmol) and HOBt.H₂O (14.4 mg, 106 μmol) in DCM (15 mL) was added EDC (245 mg, 1.27 mmol) and stirred for 18 h. The mixture was washed with water, sat. aq. NaHCO₃ and brine, and concentrated *in vacuo* to


leave the *title compound* as a brown residue (412 mg, 86%). LCMS (Method 3): Rt 3.00 min, m/z 454 [MH⁺].

d. 6-Fluoro-3-(4-methyl-3-triisopropylsilyloxymethyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 56d)



Hexachloroethane (426 mg, 1.80 mmol) was added portion-wise to a stirred mixture of Intermediate 56c (410 mg, 0.90 mmol), Ph₃P (472 mg, 1.80 mmol) and Et₃N (500 μL, 3.60 mmol) in THF (15 mL) at RT. The reaction mixture was stirred for 18 h, then filtered and concentrated *in vacuo*. The resulting residue was
10 purified by FCC, using 0-25% MeOH in EtOAc, to give the *title compound* (237 mg, 60%). LCMS (Method 3): Rt 2.95 min, m/z 436 [MH⁺].


e. (1S,4R)-4-[3-(4-Methyl-3-triisopropylsilyloxymethyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-ylamine (Intermediate 56e)



Intermediate A (115.4 mg, 0.71 mmol) was added to a mixture of NaH (60% in mineral oil, 87.2 mg, 2.18 mmol) in DMF (4 mL) at RT and stirred for 15 min. Intermediate 56d (237 mg, 0.54 mmol) was then added and the resulting mixture heated to 60°C for 2 h. After cooling, the reaction was quenched with sat.
20 aq. NH₄Cl solution. The mixture was diluted with water and extracted with EtOAc. The combined organic extracts were washed with sat. aq. NaHCO₃ solution and brine, and concentrated *in vacuo*. The resulting residue was loaded onto an SCX-2 cartridge, washed with MeOH then eluted with 1M NH₃ in MeOH; concentration

in vacuo gave the *title compound* as a brown residue (91 mg, 29%). LCMS (Method 3): Rt 3.08 min, m/z 579 [MH⁺].

- f. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(4-methyl-3-triisopropylsilanyloxymethyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 56f)**



- A mixture of Intermediate 56e (87.0 mg, 0.15 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 66.8 mg, 0.17 mmol), DIPEA (39.2 μ L, 0.23 mmol) and 1,4-dioxane (2.5 mL) were heated to 60°C for 18 h. After cooling the solvent was evaporated *in vacuo* and the residue purified by FCC, using 0-10% MeOH in DCM, to provide the *title compound* as a pale yellow foam (64 mg, 51%). LCMS (Method 3): Rt 3.63 min, m/z 834 [M⁺].

- g. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(3-hydroxymethyl-4-methyl-piperazin-1-ylmethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 56)**


- A mixture of Intermediate 56f (60.0 mg, 72.0 μ mol) and TBAF (1M in THF, 144 μ L, 144 μ mol) in THF (3 mL) was stirred at RT for 1 h. The mixture was diluted with water and extracted with EtOAc (3 $\times$). The combined organic extracts were washed with brine, and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to afford the *title compound* as an off-white solid (24.9 mg, 51%). LCMS (Method 5): Rt 3.54 min, m/z 678 [MH⁺].

¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.82-2.01 (4H, m), 2.01-2.25 (7H, m), 2.53-2.88 (3H, m), 3.21 (1H, m), 3.54 (1H, m), 3.96-4.11 (2H, m), 4.37 (1H, dt, J 12.5, 5.4), 4.84 (1H, m), 5.47 (1H, m), 6.32 (1H, s), 7.10 (1H, d, J 8.4), 7.22-7.44 (9H, m), 7.72 (1H, d, J 10.3), 8.02 (1H, s), 8.30 (1H, m).

5


Example 57

**1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-(4-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**



10

**a. 3-(4-Allyloxy-piperidin-1-yl)-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine
(Intermediate 57a)**




15

20


To an opaque pale yellow solution of Intermediate 24c (510 mg, 2.16 mmol) in dry THF at 0°C under Ar was added NaH (60% dispersion in oil, 216 mg, 5.40 mmol) (CARE: gas evolution) and the resulting suspension was stirred at 0°C for 30 min. Allyl bromide (0.467 mL, 5.40 mmol) was added and the suspension stirred at RT under Ar for 1 h and at 70°C for 18 h. Water (25 mL) was added to the cooled solution, then the mixture extracted with EtOAc (3 × 25 mL). The combined organics were washed with brine (25 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave a dark brown oil. FCC, using 0-8% MeOH in EtOAc, gave the *title compound* as a brown oil (171 mg, 29%). LCMS (Method 3): Rt 2.81 min, m/z 277 [MH⁺].

b. (1S,4R)-4-[3-(4-Allyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 57b)



A suspension of Intermediate A (108 mg, 0.661 mmol) and NaH (60% dispersion in oil, 72.1 mg, 180 mmol) in dry DMF (5 mL) at RT under Ar was stirred for 1 h (CARE: gas evolution). A solution of Intermediate 57a (166 mg, 0.601 mmol) in dry DMF (3 mL) was added and the dark brown solution stirred at 60°C for 90 min. The solution was concentrated *in vacuo*, and the residue redissolved in MeOH (5 mL) and AcOH (0.171 mL, 3.00 mmol). The solution was applied to an SCX-2 cartridge (10 g) and was washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* left a dark brown solid. FCC, using 2-8% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a dark brown oil (204 mg, 81%). LCMS (Method 3): Rt 2.34 min, m/z 420 [MH⁺].

c. 1-[(1S,4R)-4-[3-(4-Allyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-[5-tert-butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-urea (Intermediate 57c)



A yellow-brown solution of Intermediate 33a (99.6 mg, 0.237 mmol), Intermediate 57b (90.3 mg, 0.215 mmol) and DIPEA (0.047 mL, 0.269 mmol) in dioxane (3 mL) was stirred at 60°C for 18 h. The solution was concentrated *in*

vacuo, then the residue suspended in water (5 mL) and extracted with DCM (2 × 5 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo*. The residue was purified by FCC, using 2-8% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale yellow solid (96.4 mg, 65%).


5 LCMS (Method 3): Rt 3.54 min, m/z 691 [MH⁺].

**d. 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-(4-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea. (Example 57)**

Ar was bubbled through a solution of Intermediate 57c (93.0 mg, 0.135
10 mmol) and 1,3-dimethyl barbituric acid (63.0 mg, 0.404 mmol) in DCM (5 mL) at
RT under Ar for 30 min, then Pd(PPh₃)₄ (15.6 mg, 0.0135 mmol) was added and
the orange solution was stirred at RT for 1 h, and then at reflux for 17 h. The
suspension was concentrated *in vacuo*, the residue redissolved in DCE (10 mL)
and Ar bubbled through the mixture for 30 min. Pd(PPh₃)₄ (15.6 mg, 0.0135
15 mmol) was added and the opaque orange-red solution stirred at 75°C under Ar for
2 h. The cooled solution was concentrated *in vacuo* to ~3 mL volume, then applied
to an SCX-2 cartridge (5 g) and washed with MeOH (25 mL). The product was
eluted with 2M NH₃ in MeOH (20 mL); concentration *in vacuo* left an orange
solid. FCC, using 5-15% MeOH in DCM, gave a pale orange solid (46.0 mg).
20 Further purification by HPLC (XBridge C18 column, 20-80% MeCN in H₂O,
0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (27.1
mg, 31%). LCMS (Method 5): Rt 3.59 min, m/z 651 [MH⁺]. ¹H NMR (400 MHz,
d₆-DMSO): 1.23 (9H, s), 1.56-1.67 (2H, m), 1.76-1.92 (4H, m), 1.95-2.11 (2H, m),
2.91-2.97 (2H, m), 3.31-3.35 (2H, m), 3.64-3.71 (1H, m), 4.51 (2H, d, J 4.2), 4.70
25 (1H, d, J 3.9), 4.77 (1H, m), 5.25 (1H, t, J 5.3), 5.50 (1H, t, J 4.4), 6.29 (1H, s),
7.05 (1H, d, J 8.5), 7.10 (1H, dd, J 9.9, 2.1), 7.21-7.25 (2H, m), 7.28-7.34 (2H, m),
7.39 (2H, d, J 9.1), 7.40 (2H, d, J 9.1), 7.56 (1H, dd, J 9.9, 0.8), 7.61 (1H, dd, J 2.1,
0.9), 8.04 (1H, s).


Example 58

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1*S*,4*R*)-4-[3-((*S*)-1-isopropyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea, partial formate salt




5

a. (*S*)-1-Isopropyl-pyrrolidine-2-carboxylic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 58a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (200 mg, 1.57 mmol) in DMF (15.0 mL) was added (*S*)-1-isopropyl-pyrrolidine-2-carboxylic acid (*Chem. Commun.* **2006**, 14, 1482; 247 mg, 1.57 mmol), EDC (332 mg, 1.73 mmol) and HOBT.H₂O (20.0 mg, 0.16 mmol). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* (346 mg, 83%). LCMS (Method 1): Rt 0.37 min, m/z 267.1 [MH⁺].

b. 6-Fluoro-3-((*S*)-1-isopropyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-*a*]pyridine (Intermediate 58b)




20

To a solution of Intermediate 58a (346 mg, 1.30 mmol), Ph₃P (681 mg, 2.60 mmol) and Et₃N (723 μL, 5.20 mmol) in THF (10.0 mL) at 0°C was added

hexachloroethane (616 mg, 2.60 mmol). The reaction was stirred at RT overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was dissolved in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (220 mg, 68%). LCMS (Method 4): Rt 0.32 min, m/z 249.1 [MH⁺].

c. **(1*S*,4*R*)-4-[3-((*S*)-1-Isopropyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine** (Intermediate 58c)



To a suspension of NaH (60% in mineral oil, 142 mg, 3.56 mmol) in DMF (2.00 mL) was added (1*R*,4*S*)-4-amino-1,2,3,4-tetrahydro-naphthalen-1-ol Intermediate A (145 mg, 0.88 mmol) and the reaction stirred for 20 min. A solution of Intermediate 58b (220 mg, 0.88 mmol) in DMF (2.00 mL) was added and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 2-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (145 mg, 44%). LCMS (Method 4): Rt 0.29 min, m/z 392.2 [MH⁺].


d. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1*S*,4*R*)-4-[3-((*S*)-1-isopropyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea, partial formate salt** (Example 58)

To a solution of Intermediate 58c (140 mg, 0.37 mmol) in 1,4-dioxane (3.00 mL) was added DIPEA (130 μL, 0.75 mmol) and (5-tert-butyl-2-p-tolyl-2H-

pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 151 mg, 0.37 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-5% MeOH in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (60 mg, 25%). LCMS (Method 5): Rt 3.84 min, m/z 647.3 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 0.99 (6H, d, J 6.5), 1.29 (9H, s), 1.90-2.06 (5H, m), 2.11 (1H, m), 2.22-2.36 (2H, m), 2.38 (3H, s), 2.73-2.84 (2H, m), 3.25 (1H, m), 4.59 (1H, t, J 7.3), 4.90 (1H, dd, J 9.1, 5.7), 5.29 (1H, t, J 4.1), 6.33 (1H, s), 7.20-7.36 (9H, m), 7.65 (1H, d, J 10.0), 8.25 (0.5H, br s), 8.42 (1H, d, J 1.8).


Example 59

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-dimethylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea



a. 2-(5-Fluoropyridin-2-yl)-N,N-dimethylhydrazine carboxamide


(Intermediate 59a)



A solution of (5-fluoro-pyridin-2-yl)-hydrazine (500 mg, 3.93 mmol), dimethylcarbonyl chloride (505 mg, 4.72 mmol) and DIPEA (1.01 g, 7.86 mmol) in DCM (20 mL) was stirred at reflux for 16 h. The reaction mixture was applied


to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* and then trituration with diethyl ether gave the *title compound* (600 mg, 77%). ¹H NMR (400 MHz, CDCl₃): 2.99 (6H, s), 6.46 (2H, m), 6.75 (1H, dd, J 9.1, 3.5), 7.22-7.32 (1H, m), 8.03 (1H, d, J 2.7).

5 **b. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-dimethyl-amine**
(Intermediate 59b)



To a solution of Intermediate 59a (590 mg, 2.98 mmol), Ph₃P (1.56 g, 5.96 mmol) and Et₃N (1.20 g, 11.9 mmol) in THF (40 mL) was added hexachloroethane
10 (1.41 g, 5.96 mmol) and the mixture stirred at 60°C for 9 h. The mixture was diluted with EtOAc (100 mL), washed with water, brine, dried (MgSO₄) and then concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, then trituated with diethyl ether to give the *title compound* (78 mg, 14%). LCMS (Method 1): Rt 2.24 min, m/z 181 [MH⁺].

15 **c. [6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-**
[1,2,4]triazolo[4,3-a]pyridin-3-yl)-dimethyl-amine (Intermediate 59c)




To a solution of Intermediate A (75 mg, 0.458 mmol) in DMF (2 mL) was added NaH (60% in oil, 50 mg, 1.25 mmol) and the mixture stirred at RT for 20
20 min, before Intermediate 59b (75 mg, 0.416 mmol) was added. This mixture was stirred at 60°C for 1 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM gave the *title compound* as a yellow gum (82 mg, 61%). LCMS (Method 4):
25 Rt 1.49 min, m/z 324 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-dimethylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 59)


A solution of Intermediate 59c (80 mg, 0.25 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 150 mg, 0.370 mmol) and DIPEA (129 mg, 1.00 mmol) in DMF (2 mL) was stirred at 60°C for 30 min. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-5% [2M NH₃ in MeOH] in DCM, gave the *title* compound as an off-white solid (75 mg, 52%). LCMS (Method 5): Rt 4.39 min, m/z 579.1 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.29 (9H, s), 1.83-2.17 (3H, m), 2.17-3.00 (1H, m), 2.38 (3H, s), 2.92 (6H, s), 4.88 (1H, dd, J 8.5, 5.4), 5.41 (1H, t, J 3.0), 6.33 (1H, s), 7.14-7.36 (9H, m), 7.50 (1H, d, J 9.7), 7.70 (1H, d, J 1.4).

15 **Example 60**

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(R)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



a. **6-Fluoro-3-((R)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-**
a]pyridine (Intermediate 60a)




A mixture of (5-fluoro-pyridin-2-yl)-hydrazine (889 mg, 7.00 mmol), N-

methyl-D-proline hydrochloride (1.274 g, 7.7 mmol) and HOBt.H₂O (95 mg, 0.70 mmol) in DCM (20 mL) was treated with EDC (1.494 g, 7.8 mmol), and the mixture stirred at RT for 6.5 h. The mixture was then diluted with sat. aq. NaHCO₃ solution, NaCl added, and the mixture extracted with DCM (8 × 20 mL). The combined organics were dried and concentrated *in vacuo* to give (R)-1-methyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide as a pale orange solid (1.09 g).

This was dissolved in THF (20 mL) then Ph₃P (2.40 g, 9.16 mmol), Et₃N (2.6 mL, 18.3 mmol), and hexachloroethane (2.17 g, 9.16 mmol) added sequentially. The mixture was stirred at RT for 4 h, then was filtered, the filter-cake washed with diethyl ether, and the combined organics concentrated *in vacuo*. The residue was purified on a SCX-2 cartridge (50 g), washing with methanol then eluting basic components with 0.2 – 2 M ammonia in methanol. Product containing fractions were combined and concentrated *in vacuo*. The residue was further purified by FCC, using 0-100% EtOAc in cyclohexane, then 8-10% MeOH in EtOAc, to give the *title compound* as a pale red oil (876 mg, 57%). LCMS (Method 3): Rt 0.55 min, m/z 221 [MH⁺].

b. (1S,4R)-4-[3-((R)-1-Methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 60b)



To a solution of Intermediate A (245 mg, 1.50 mmol) in DMF (4 mL) was added NaH (60% dispersion in oil, 180 mg, 4.50 mmol) and the mixture stirred at RT for 0.5 h. A solution of Intermediate 60a (330 mg, 1.50 mmol) in DMF (5 mL) was added and the mixture stirred at 60°C for 2 h. The cooled solution was diluted with water and extracted with DCM (5 × 20 mL). The combined organics were


dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-15% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a brown solid after freeze-drying (408 mg, 74%). LCMS (Method 3): Rt 0.44 min, m/z 364 [MH⁺].

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 60)


A solution of Intermediate 60b (123 mg, 0.339 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 140 mg, 0.346 mmol) in 1,4-dioxane (4 mL) and DIPEA (91 µL, 0.52 mmol) was stirred at 93°C for 3.25 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the impure product. Further purification by HPLC (C18 X-select column, 25-70% MeCN in H₂O, 0.1% formic acid) gave the *title compound* as a white powder after freeze-drying (118 mg, 56%). LCMS (Method 5): Rt 3.67 min, m/z 619.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.80-2.27 (11H, m), 2.31-2.40 (4H, m), 3.11-3.18 (1H, m), 3.98 (1H, t, J 8.2), 4.78-4.86 (1H, m), 5.42 (1H, t, J 4.3), 6.32 (1H, s), 7.09 (1H, d, J 8.5), 7.26-7.42 (9H, m), 7.75 (1H, dd, J 9.7, 1.0), 8.03 (1H, s), 8.28 (1H, d, J 1.0).

Example 61

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1-ethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (S)-1-Ethyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 61a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (178 mg, 1.39 mmol) in DMF (10.0 mL) was added (S)-1-ethyl-pyrrolidine-2-carboxylic acid (200 mg, 1.39 mmol), EDC (293 mg, 1.53 mmol) and HOBt.H₂O (18.0 mg, 0.14 mmol). The reaction was stirred for 4 h then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (190 mg, 54%). LCMS (Method 4): Rt 0.29 min, m/z 253.1 [MH⁺].


b. 6-Fluoro-3-((S)-1-ethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 61b)



To a solution of Intermediate 61a (190 mg, 0.75 mmol), Ph₃P (395 mg, 1.51 mmol) and Et₃N (419 μL, 3.01 mmol) in THF (6.00 mL) at 0°C was added hexachloroethane (357 mg, 1.51 mmol). The reaction was stirred at RT overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was dissolved in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (157 mg, 89%). LCMS (Method 2): Rt 0.28 min, m/z 235.2 [MH⁺].

c. (1S,4R)-4-[3-((S)-1-Ethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-

***a*[pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 61c)**



To a suspension of NaH (60% in mineral oil, 107 mg, 2.68 mmol) in DMF
 5 (2.00 mL) was added Intermediate A (110 mg, 0.67 mmol) and the reaction stirred
 for 20 min. Intermediate 61b (150 mg, 0.67 mmol) was added in DMF (2.00 mL)
 and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by
 dropwise addition of methanol, before being diluted with methanol and loaded
 onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted
 10 with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 2-10%
 [2M NH₃ in MeOH] in DCM, gave the *title compound* (72.0 mg, 28%). LCMS
 (Method 4): Rt 0.29 min, m/z 378.2 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1-ethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 61)


To a solution of Intermediate 61c (72.0 mg, 0.19 mmol) in 1,4-dioxane
 (2.00 mL) was added DIPEA (66 μL, 0.38 mmol) and (5-tert-butyl-2-p-tolyl-2H-
 pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic
 Communications, 2009, 39, 3999-4009; 77.0 mg, 0.19 mmol). The reaction was
 20 heated to 60°C overnight then cooled and partitioned between EtOAc and water.
 The aqueous layer was then extracted with EtOAc (3 ×). The combined organic
 layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*.
 The residue was purified by FCC, using 0-7.5% MeOH in DCM. Further
 purification by HPLC (C18 X-select column, 20-70% MeCN in H₂O, 0.1%
 25 HCO₂H) gave the *title compound* as a white powder after freeze-drying (37 mg,
 31%). LCMS (Method 5): Rt 3.73 min, m/z 633.2 [MH⁺]. ¹H NMR (400 MHz, d₄-

MeOD): 0.94 (3H, t, J 7.2), 1.30 (9H, s), 1.89-2.16 (6H, m), 2.22-2.31 (2H, m), 2.31-2.38 (2H, m), 2.38 (3H, s), 2.49 (1H, m), 3.32 (1H, m), 4.11 (1H, t, J 8.0), 4.90 (1H, dd, J 9.0, 5.6), 5.28 (1H, t, J 4.1), 6.33 (1H, s), 7.19-7.36 (9H, m), 7.65 (1H, dd, J 10.0, 0.7), 8.37 (1H, d, J 1.9).

5


Example 62

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1*S*,4*R*)-4-[3-[(*S*)-1-methyl-piperidin-2-yl]-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




10

a. (*S*)-1-Methyl-piperidine-2-carboxylic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 62a)




To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (250 mg, 1.96 mmol) in DMF (20.0 mL) was added (*S*)-1-methyl-piperidine-2-carboxylic acid (281 mg, 1.96 mmol), EDC (416 mg, 2.16 mmol) and HOBT.H₂O (25.0 mg, 0.20 mmol). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (331 mg, 67%). LCMS (Method 1): Rt 0.35 min, m/z 253.2 [MH⁺].

b. 6-Fluoro-3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 62b)



To a solution of Intermediate 62a (331 mg, 1.31 mmol), Ph_3P (688 mg, 2.63 mmol) and Et_3N (731 μL , 5.25 mmol) in THF (13.0 mL) at 0°C was added hexachloroethane (622 mg, 2.63 mmol). The reaction was stirred at RT overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO_4), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH_3 in MeOH; concentration *in vacuo* gave the *title compound* (250 mg, 81%). LCMS (Method 4): R_t 0.31 min, m/z 235.1 [MH^+].

c. (1S,4R)-4-[3-((S)-1-Methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 62c)



To a suspension of NaH (60% in mineral oil, 171 mg, 4.27 mmol) in DMF (4.00 mL) was added Intermediate A (175 mg, 1.07 mmol) and the reaction stirred for 20 min. Intermediate 62b (250 mg, 1.07 mmol) was added in DMF (2.00 mL) and the reaction heated to 60°C for one hour. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH and product eluted with 2M NH_3 in MeOH. The resulting residue was purified by FCC, using 2-10% [2M NH_3 in MeOH] in DCM, to give the *title compound* (287 mg, 71%).

LCMS (Method 1): Rt 1.64, m/z 378.2 [MH⁺].


d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 62)

To a solution of Intermediate 62c (140 mg, 0.37mmol) in 1,4-dioxane (4.00 mL) was added DIPEA (129 μ L, 0.74 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 150 mg, 0.37 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water.


The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-7.5% MeOH in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (73 mg, 31%). LCMS (Method 5): Rt 3.75 min, m/z 633.2 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.45 (1H, m), 1.60-1.83 (3H, m), 1.83-1.93 (2H, m), 1.93-2.07 (5H, m), 2.11 (1H, m), 2.20-2.33 (2H, m), 2.38 (3H, s), 3.05 (1H, dt, J 11.8, 2.8), 3.79 (1H, dd, J 10.9, 2.8), 4.91 (1H, dd, J 9.0, 5.7), 5.32 (1H, t, J 4.0), 6.33 (1H, s), 7.19-7.36 (9H, m), 7.64 (1H, d, J 10.0), 8.43 (1H, d, J 1.7).

Example 63

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. **(S)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-3-ol**
(Intermediate 63a)




A mixture of Intermediate 24b (394 mg, 2.30 mmol), (S)-3-hydroxypiperidine hydrochloride (1.00 g, 7.27 mmol) and DIPEA (1.27 mL, 7.30 mmol) in DMA (8 mL) was heated in the microwave at 170-180°C for 9.5 h. The cooled mixture was concentrated *in vacuo* and applied to an SCX-2 cartridge (70 g), washing with methanol then eluting basic components with 0.4 – 2 M ammonia in methanol. Product containing fractions were combined and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the *title compound* as a pale brown solid (204 mg, 38%). LCMS (Method 3): Rt 2.02 min, m/z 237 [MH⁺].

b. **6-Fluoro-3-((S)-3-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 63b)**




To a solution of Intermediate 63a (204 mg, 0.864 mmol) and Et₃N (180 µL, 1.3 mmol) in DCM (5 mL) was added triisopropylsilyltrifluoromethane sulfonate (279 µL, 1.04 mmol) and the mixture stirred at RT for 1 h. The mixture was washed with sat. aq. NaHCO₃ solution, dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane, then 10% MeOH in EtOAc, to give the *title compound* as a pale brown gum (290 mg, 86%). LCMS (Method 3): Rt 4.99 min, m/z 393 [MH⁺].

c. (1S,4R)-4-[3-((S)-3-Triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 63c)



5 To a solution of Intermediate A (116 mg, 0.714 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 44 mg, 1.1 mmol) and the mixture stirred at RT for 20 min. A solution of Intermediate 63b (280 mg, 0.714 mmol) in DMF (3 mL) was added and the mixture was stirred at 60°C for 25 min, and up to 150°C over 20 min. The cooled mixture was diluted with water and extracted with DCM (4 ×
10 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified twice by FCC, using 0-12% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a dark brown gum (141 mg, 37%). LCMS (Method 3): Rt 3.12 min, m/z 536 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-3-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 63d)




A solution of Intermediate 63c (140 mg, 0.261 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic
20 Communications, 2009, 39, 3999-4009; 106 mg, 0.261 mmol) in 1,4-dioxane (3 mL) and DIPEA (70 μL, 0.4 mmol) was stirred at 90°C for 3 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-10%

MeOH in DCM, to give the *title compound* as a pale brown gum (124 mg, 60%).
LCMS (Method 3): Rt 5.43 min, m/z 791 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-3-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 63)

A solution of Intermediate 63d (123 mg, 0.156 mmol) and TBAF (1M in THF, 195 μ L, 0.195 mmol) in THF (4 mL) was stirred at RT for 1 h. The mixture was diluted with water and extracted with DCM (4 $\times$ 15 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-12% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a white powder after freeze-drying (82 mg, 83%). LCMS (Method 5): Rt 4.27 min, m/z 635.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.38-1.47 (1H, m), 1.62-1.74 (1H, m), 1.79-1.98 (4H, m), 1.99-2.16 (2H, m), 2.36 (3H, s), 2.78-2.86 (1H, m), 2.89-2.98 (1H, m), 3.20-3.36 (2H, m, under water signal), 3.77-3.86 (1H, m), 4.77-4.86 (1H, m), 4.90 (1H, d, J 5.2), 5.53 (1H, t, J 4.3), 6.32 (1H, s), 7.08 (1H, d, J 8.7), 7.16 (1H, dd, J 10.1, 2.1), 7.26-7.41 (8H, m), 7.61 (1H, d, J 10.0), 7.69 (1H, d, J 1.4), 8.03 (1H, s).

Example 64




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-3-hydroxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

The *title compound* was prepared starting from (R)-3-hydroxypiperidine hydrochloride using analogous procedures to those described for Example 63.
LCMS (Method 5): Rt 4.27 min, m/z 635.2 [MH⁺]. ¹H NMR (400 MHz, d₆-

DMSO): 1.27 (9H, s), 1.36-1.47 (1H, m), 1.65-1.76 (1H, m), 1.77-1.98 (4H, m), 1.99-2.17 (2H, m), 2.36 (3H, s), 2.78-2.85 (1H, m), 2.87-2.95 (1H, m), 3.20-3.37 (2H, m, under water signal), 3.75-3.84 (1H, m), 4.78-4.85 (1H, m), 4.90 (1H, d, J 5.0), 5.51 (1H, t, J 4.3), 6.32 (1H, s), 7.08 (1H, d, J 8.4), 7.17 (1H, dd, J 10.0, 2.2),
5 7.26-7.41 (8H, m), 7.61 (1H, d, J 10.0), 7.69 (1H, d, J 1.4), 8.03 (1H, s).

Example 65


**1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1*S*,4*R*)-4-[3-((*S*)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea, partial formate salt**



10 To a solution of Intermediate 62c (117 mg, 0.31 mmol) in 1,4-dioxane (3.00 mL) was added DIPEA (108 μ L, 0.62 mmol) and Intermediate 33a (130 mg, 0.31 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered
15 and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (80 mg, 39%). LCMS (Method 5): Rt 3.31 min, m/z 649.3 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.31 (9H, s), 1.49 (1H, m), 1.60-2.05 (7H, m), 2.05 (3H, s), 2.10
20 (1H, m), 2.25-2.35 (2H, m), 3.09 (1H, m), 3.87 (1H, dd, J 10.9, 2.4), 4.65 (2H, s), 4.91 (1H, m), 5.33 (1H, t, J 4.0), 6.34 (1H, s), 7.19-7.33 (5H, m), 7.42-7.51 (4H, m), 7.65 (1H, d, J 9.6), 8.26 (0.5H, br s), 8.43 (1H, d, J 1.7).


Example 66

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-(3-(4-hydroxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea



5


a. [1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-4-yl]-ethanol (Intermediate 66a)



A mixture of Intermediate 24b (542 mg, 3.17 mmol) and 4-piperidine-ethanol (1.54 g, 11.9 mmol) in DMA (10 mL) was heated in the microwave at 170°C for 3 h. The cooled mixture was applied to an SCX-2 cartridge (50 g), washing with methanol then eluting basic components with 0.4 – 2 M ammonia in methanol. Product containing fractions were combined and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the *title compound* as a pale brown solid (324 mg, 41%). LCMS (Method 3): Rt 2.28 min, m/z 265 [MH⁺].

15

b. 6-Fluoro-3-(4-triisopropylsilanyloxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin (Intermediate 66b)




20

To a solution of Intermediate 66a (319 mg, 1.21 mmol) and Et₃N (252 μL,

1.82 mmol) in DCM (10 mL) was added triisopropylsilyltrifluoromethane sulfonate (319 μ L, 1.45 mmol) and the mixture stirred at RT for 1.5 h. The mixture was washed with sat. aq. NaHCO_3 solution, dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* as an orange solid (465 mg, 92%). LCMS (Method 3): Rt 5.35 min, m/z 421 $[\text{MH}^+]$.

c. (1S,4R)-4-[3-(4-Triisopropylsilyloxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 66c)




10

To a solution of Intermediate A (122 mg, 0.75 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 60 mg, 1.5 mmol) and the mixture stirred at RT for 25 min. A solution of Intermediate 66b (315 mg, 0.75 mmol) in DMF (2.5 mL) was added and the mixture stirred at 60°C for 1.5 h. The cooled mixture was diluted with water and extracted with DCM (5×25 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% [2M NH_3 in MeOH] in DCM, to give the *title compound* as a dark brown gum (180 mg, 42%). LCMS (Method 3): Rt 3.35 min, m/z 564 $[\text{MH}^+]$.

20

25

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-triisopropylsilyloxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 66d)



5 A mixture of Intermediate 66c (178 mg, 0.316 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 128 mg, 0.316 mmol) in 1,4-dioxane (3 mL) and DIPEA (83 μ L, 0.474 mmol) was stirred at 95°C for 4 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* as a brown foam (208 mg, 81%). LCMS (Method 3): Rt 5.96 min, m/z 819 [MH⁺].


e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxyethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 66)

15 A solution of Intermediate 66d (207 mg, 0.253 mmol) and TBAF (1M in THF, 0.303 mL, 0.303 mmol) in THF (4 mL) was stirred at RT for 2.25 h. The mixture was diluted with water and extracted with DCM (4 $\times$ 15 mL). The combined organic extracts were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-15% MeOH in DCM, to give the product. This was further purified by HPLC (C18 X-select column, 30-98% MeCN in H₂O, 0.1% formic acid) to give the *title compound* as an off-white powder after freeze-drying (114 mg, 68%). LCMS (Method 5): Rt 4.29 min, m/z 663.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.38-1.51 (4H, m), 1.57-1.68 (1H, m), 1.72-1.97 (4H, m), 1.99-2.16 (2H, m), 2.36 (3H, s), 2.85-2.95 (2H, m), 3.30-3.52 (4H, m,


under water signal), 4.38 (1H, s), 4.77-4.85 (1H, m), 5.55 (1H, t, J 4.3), 6.32 (1H, s), 7.09 (1H, d, J 8.6), 7.16 (1H, dd, J 9.9, 2.2), 7.26-7.40 (8H, m), 7.61 (1H, d, J 9.9), 7.65 (1H, d, J 1.4), 8.06 (1H, s).

Example 67

5 **1-[(1S,4R)-4-(3-Azepan-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea**




10 **a. 6-Fluoro-3-azepan-1-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 67a)**



A solution of Intermediate 24b (451 mg, 2.50 mmol) and azapane (990 mg, 10.0 mmol) in DMA (10 mL) was heated in the microwave at 175-180°C for 6 h. The mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give impure product. Further purification by FCC, using 0-12% MeOH in DCM, gave impure product. Further purification by FCC, using 0-6% MeOH in EtOAc, gave the *title compound* as a green gum (260 mg, 44%). LCMS (Method 3): Rt 2.76 min, m/z 235 [MH⁺].

20 **b. (1S,4R)-4-(3-Azepan-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 67b)**




To a solution of Intermediate A (137 mg, 0.84 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 68 mg, 1.7 mmol) and the mixture stirred at RT for 25 min. A solution of Intermediate 67a (197 mg, 0.84 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 1.25 h. The cooled mixture was diluted with water and extracted with DCM (5 × 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a dark brown gum (175 mg, 55%). LCMS (Method 3): Rt 2.22 min, m/z 378 [MH⁺].

c. 1-[(1S,4R)-4-(3-Azepan-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea (Example 67)

A solution of Intermediate 67b (173 mg, 0.458 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 186 mg, 0.458 mmol) in 1,4-dioxane (3 mL) and DIPEA (122 µL, 0.700 mmol) was stirred at 95°C for 3.5 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-14% [2M NH₃ in MeOH] in DCM. Further purification by FCC, using 0-10% MeOH in DCM, gave impure product. This was further purified by HPLC (C18 X-select column, 30-98% MeCN in H₂O, 0.1% HCO₂H) to give impure product. Further purified by HPLC (XBridge column, 35-98% MeCN in H₂O, 0.1% NH₄OH) to give an off-white powder after freeze-drying (50 mg). This material was further purified by MDAP (Method 7) to give the *title compound* as a white solid (28 mg, 10%). LCMS (Method 5): Rt 4.77 min, m/z 633.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.64-1.71 (4H, m), 1.74-1.97 (6H, m), 1.99-2.18 (2H, m), 2.36 (3H, s), 3.38-3.43 (4H, m, under water signal), 4.77-4.85 (1H, m), 5.47 (1H, t, J 4.3), 6.32 (1H, s), 7.07-7.14 (2H, m), 7.25-7.40 (8H, m), 7.55-7.59 (2H, m), 8.06 (1H, s).

Example 68

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-piperazin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea




5

a. 6-Fluoro-3-(4-methyl-piperazin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 68a)



A solution of Intermediate 24b (451 mg, 2.50 mmol) and N-methyl-piperazine (1.11 mL, 10.0 mmol) in DMA (10 mL) was heated in the microwave at 170°C for 8 h. The mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* as an orange gum (190 mg, 32%). LCMS (Method 3): Rt 0.43 min, m/z 236 [MH⁺].

b. (1S,4R)-4-[3-(4-Methyl-piperazin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-amine (Intermediate 68b)



To a solution of Intermediate A (125 mg, 0.766 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 62 mg, 1.54 mmol) and the mixture stirred at RT for 0.5 h. A solution of Intermediate 68a (180 mg, 0.766 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 1.75 h. The cooled mixture was

20


diluted with water and extracted with DCM (5×25 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale brown gum (225 mg, 78%). LCMS (Method 3): Rt 0.44 min, m/z 379 [MH⁺].

5 **c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-piperazine-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}urea (Example 68)**

A solution of Intermediate 68b (215 mg, 0.560 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic
10 Communications, 2009, 39, 3999-4009; 226 mg, 0.560 mmol) in 1,4-dioxane (3 mL) and DIPEA (146 μ L, 0.840 mmol) was stirred at 65°C for 18 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 10% MeOH in DCM, then 0-14% [2M NH₃ in MeOH] in DCM, to give impure product. Further purification by HPLC (XBridge C18 column, 35-95% MeCN in H₂O,
15 0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (205 mg, 58%). LCMS (Method 5): Rt 3.58 min, m/z 634.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.79-1.98 (2H, m), 1.99-2.15 (2H, m), 2.26 (3H, s), 2.36 (3H, s), 2.52-2.57 (4H, m), 3.15-3.24 (4H, m), 4.78-4.85 (1H, m), 5.56 (1H, t, J 4.4), 6.33 (1H, s), 7.07 (1H, d, J 8.6), 7.15 (1H, dd, J 9.9, 2.1), 7.26-7.40 (8H, m),
20 7.62 (1H, d, J 9.9), 7.68 (1H, d, J 1.8), 8.04 (1H, s).


Example 69

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(4-methyl-[1,4]diazepan-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl}urea




5

a. 6-Fluoro-3-(4-methyl-[1,4]diazepan-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 69a)



A solution of Intermediate 24b (451 mg, 2.50 mmol) and 1-methyl-[1,4]diazepane (1.14 g, 10.0 mmol) in DMA (10 mL) was heated in the microwave at 175°C for 6 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a brown oil (480 mg, 77%). LCMS (Method 3): Rt 0.44 min, m/z 250 [MH⁺].

b. (1S,4R)-4-[3-(4-Methyl-[1,4]diazepan-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-ylamine (Intermediate 69b)



To a solution of Intermediate A (183 mg, 1.12 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 149 mg, 2.24 mmol) and the mixture stirred at

20

RT for 0.5 h. A solution of Intermediate 69a (400 mg, 1.12 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 1.5 h. The cooled mixture was diluted with water and extracted with DCM (3 × 20 mL) and DCM-MeOH (4:1, 2 × 25 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-25% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a dark brown gum (179 mg, 41%). LCMS (Method 3): Rt 0.44 min, m/z 393 [MH⁺].


c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-1,4-diazepan-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]urea (Example 69)

A solution of Intermediate 69b (170 mg, 0.43 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 174 mg, 0.43 mmol) in 1,4-dioxane (3 mL) and DIPEA (113 µL, 0.65 mmol) was stirred at 65°C for 18 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 10% MeOH in DCM, then 0-15% [2M NH₃ in MeOH] in DCM, to give impure product. Further purification by HPLC (XBridge C18 column, 35-95% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (186 mg, 67%). LCMS (Method 5): Rt 3.57 min, m/z 648.2 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.79-1.98 (4H, m), 2.00-2.18 (2H, m), 2.31 (3H, s), 2.36 (3H, s), 2.62-2.67 (2H, m), 2.68-2.73 (2H, m), 3.44-3.52 (4H, m), 4.78-4.85 (1H, m), 5.49 (1H, t, J 4.4), 6.32 (1H, s), 7.08 (1H, d, J 8.5), 7.11 (1H, dd, J 9.7, 2.0), 7.26-7.40 (8H, m), 7.58 (1H, d, J 9.7), 7.63 (1H, d, J 1.9), 8.03 (1H, s).


Example 70

1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

5



a. 1-{5-tert-Butyl-2-[4-(tert-butyl-dimethyl-silanyloxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 70a)



10

The *title compound* was synthesised from 5-tert-butyl-2-[4-(tert-butyl-dimethyl-silanyloxy)-phenyl]-2H-pyrazol-3-ylamine (ref: US2006/035922) and Intermediate 3c using analogous procedures to those used in the preparation of Example 30. Beige solid. LCMS (Method 3) Rt 4.78 min, m/z 735 [MH⁺].

15

b. 1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 70)


20

A mixture of Intermediate 70a (124 mg, 169 μmol) and TBAF (1M in THF, 0.34 mL, 0.34 mmol) in THF (5 mL) was stirred for 1 h. The mixture was diluted with water and extracted with EtOAc (3 ×). The combined organic extracts were washed with brine, and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, then further purified by HPLC (30-98% MeCN in


H₂O, 0.1% HCO₂H) to give the *title compound* as an off-white solid (18.4 mg, 18%). LCMS (Method 5) Rt 4.22 min, m/z 621 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.25 (9H, s), 1.61 (2H, m), 1.72 (4H, m), 1.79-1.98 (2H, m), 1.99-2.15 (2H, m), 3.13 (4H, m), 4.81 (1H, m), 5.54 (1H, t, J 4.5), 6.29 (1H, s), 6.85 (2H, m), 7.09 (1H, d, J 8.7), 7.14 (1H, dd, J 9.7, 2.0), 7.20-7.39 (6H, m), 7.59-7.64 (2H, m), 7.95 (1H, s), 9.77 (1H, br s).

Example 71

1-[5-tert-Butyl-2-(4-cyano-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 4-(5-Amino-3-tert-butyl-pyrazol-1-yl)-benzonitrile (Intermediate 71a)



A cream suspension of 4-cyanophenylhydrazine hydrochloride (1.70 g, 10.0 mmol) and 4,4-dimethyl-3-oxopentanenitrile (1.31 g, 10.5 mmol) in EtOH (25 mL) was stirred at reflux for 4 h, then at RT for 64 h, and again at reflux for 24 h. The solution was cooled to RT, concentrated *in vacuo*, and partitioned between water (50 mL) and EtOAc (75 mL). The organics were washed with brine (50 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave an orange solid (2.28 g, 95%). LCMS (Method 3): Rt 3.45 min, m/z 241 [MH⁺].

b. [5-tert-Butyl-2-(4-cyano-phenyl)-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 71b)



To a suspension of Intermediate 71a (2.28 g, 9.49 mmol) in EtOAc (25 mL) and aqueous NaOH (1M, 23.7 mL, 23.7 mmol) was added 2,2,2-trichloroethyl chloroformate (1.57 mL, 11.4 mmol) dropwise over 2 min. A precipitate formed which redissolved after 15 min, then the orange solution was stirred at RT for 90 min. 2,2,2-Trichloroethyl chloroformate (0.391 mL, 2.85 mmol) was added and the orange mixture stirred at RT for 16 h. The layers were separated and the aqueous extracted with EtOAc (25 mL). The combined organics were washed with brine (25 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave an orange-red oil. Recrystallisation from cyclohexane gave an off-white solid (3.12 g, 79%). LCMS (Method 3): Rt 4.46 min, m/z 415, 417 [MH⁺].


c. 1-[5-tert-Butyl-2-(4-cyano-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 71)

A brown solution of Intermediate 71b (114 mg, 0.275 mmol), Intermediate 3c (91.0 mg, 0.250 mmol) and DIPEA (0.054 mL, 0.313 mmol) in dioxane (3 mL) was stirred at 60°C for 16 h. The cooled solution was concentrated *in vacuo*, the residue suspended in water (4 mL) and extracted with DCM (2 × 4 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo*. FCC, using 3% [2M NH₃ in MeOH] in DCM, gave a pale yellow solid (99.5 mg, 63%). LCMS (Method 5): Rt 4.63 min, m/z 630 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.29 (9H, s), 1.60-1.64 (2H, m), 1.70-1.76 (4H, m), 1.86-1.94 (2H, m), 1.99-2.07 (1H, m), 2.11-2.18 (1H, m), 3.14 (4H, t, J 5.2), 4.79 (1H, m),


5.54 (1H, t, J 4.3), 6.38 (1H, s), 7.12 (1H, d, J 8.6), 7.16 (1H, dd, J 9.8, 2.2), 7.25 (1H, d, J 7.7), 7.27-7.37 (2H, m), 7.39 (1H, d, J 7.6), 7.60-7.63 (2H, m), 7.78 (2H, d, J 8.6), 7.98 (2H, d, J 8.6), 8.30 (1H, s).

Example 72

5 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4S)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**



a. **(1S,4S)-4-[3-((S)-1-Methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-amine. (Intermediate 72a)**



A solution of Intermediate B (179 mg, 1.10 mmol) in dry DMF (2.5 mL) was added NaH (60% in mineral oil, 333 mg, 5.00 mmol) at RT and stirred for 15 min. Intermediate 5b (220 mg, 1.00 mmol) in DMF (2.5 mL) was then added and the mixture heated at 60°C for 1 h. After cooling, the resulting dark brown mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as light brown powder after freeze drying (224 mg, 61%). LCMS (Method 3): Rt 1.34 min, m/z 364 [MH⁺].


b. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4S)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-**

tetrahydro-naphthalen-1-yl}-urea (Example 72)


A stirred solution of Intermediate 72a (219 mg, 0.60 mmol) and 5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 269 mg, 0.66 mmol) and
5 DIPEA (317 μ L, 1.14 mmol) in THF (6 mL) was heated at reflux for 23 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give impure product. This residue was purified further by HPLC (ChiralPak IC column, 50% IPA in
10 heptanes) to give the *title compound* as a white powder after freeze-drying (225 mg, 60%). LCMS (Method 5): Rt 3.65 min, m/z 619 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.26 (9H, s), 1.74 (1H, m), 1.93-1.99 (3H, m), 2.08 (3H, s), 2.13-2.20 (4H, m), 2.32 (1H, m), 2.36 (3H, s), 3.14 (1H, td, J 8.1, 2.5), 3.95 (1H, t, J 8.1), 4.90 (1H, m), 5.48 (1H, t, J 4.0), 6.32 (1H, s), 7.00 (1H, d, J 8.2), 7.33-7.45 (9H, m), 7.74 (1H, dd, J 9.9, 0.8), 7.98 (1H, s), 8.24 (1H, d, J 2.1).
15

Example 73

1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



- a. 1-{5-tert-Butyl-2-[4-(tert-butyl-dimethyl-silanyloxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-[(S)-1-methyl-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 73a)




The *title compound* was synthesised from 5-tert-butyl-2-[4-(tert-butyl-dimethyl-silanyloxy)-phenyl]-2H-pyrazol-3-ylamine (ref: US2006/035922) and Intermediate 5c using analogous procedures to those used in the preparation of Example 30. Yellow powder. LCMS (Method 3): Rt 3.59 min, m/z 735 [MH⁺].

- b. 1-[5-tert-Butyl-2-(4-hydroxy-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-[(S)-1-methyl-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 73)

A solution of Intermediate 73a (160 mg, 0.21 mmol) and TBAF (1M in THF, 0.26 mL, 0.26 mmol) in THF (2 mL) was stirred at RT for 30 min, then diluted with water and extracted with DCM (3 × 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a white powder after freeze-drying (107 mg, 79%). LCMS (Method 5): Rt 3.24 min, m/z 621 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.26 (9H, s), 1.81-2.25 (11H, m), 2.34-2.36 (1H, m), 3.12-3.14 (1H, m), 3.99 (1H, t, J 8.2), 4.81-4.84 (1H, m), 5.39 (1H, t, J 4.3), 6.29 (1H, s), 6.85-6.88 (2H, m), 7.09 (1H, d, J 8.6), 7.28-7.39 (7H, m), 7.75 (1H, dd, J 9.9, 0.8), 7.92 (1H, s), 8.24 (1H, d, J 2.1), 9.72 (1H, s).


Example 74

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea




5

a. 2-[1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-4-yl]-propan-2-ol (Intermediate 74a)




A brown solution of Intermediate 24b (666 mg, 3.88 mmol) and 2-piperidinyl propan-2-ol (TCI, 1.39 g, 9.71 mmol) in DMA (10 mL) was irradiated to 175°C for 3 h in the microwave. The cooled solution was concentrated *in vacuo*, suspended in water (10 mL) and brine (10 mL), then extracted with DCM (2 × 20 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown oil (1.3 g). FCC, using 5% MeOH in DCM, gave the *title compound* as a pale yellow solid (484 mg, 45%). LCMS (Method 3): Rt 2.42 min, m/z 279 [MH⁺].

b. 6-Fluoro-3-[4-(1-methyl-1-triisopropylsilyloxy-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 74b)



A solution of Intermediate 74a (480 mg, 1.72 mmol), triisopropylsilyl trifluoromethane-sulfonate (0.579 mL, 2.16 mmol) and Et₃N (0.361 mL, 2.59 mmol) in dry DCM (5 mL) under N₂ was stirred at reflux for 2 h. The mixture was cooled to RT, then Et₃N (0.361 mL, 2.59 mmol) and triisopropylsilyl trifluoromethanesulfonate (0.579 mL, 2.16 mmol) were added and the orange solution stirred at reflux for 3 h. To the cooled solution, water (5 mL) was added and the mixture shaken. The aqueous was extracted with DCM (5 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave an orange oil. FCC, using 1-3% MeOH in DCM, gave the *title compound* as a pale yellow solid (592 mg, 79%). LCMS (Method 3): Rt 5.66 min, m/z 435 [MH⁺].

c. (1S,4R)-4-{3-[4-(1-Methyl-1-triisopropylsilanyloxy-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 74c)



15

To a solution of Intermediate A (129 mg, 0.788 mmol) in dry DMF (2 mL) at RT under Ar was added NaH (60% dispersion in oil, 45.0 mg, 1.13 mmol) (CARE: gas evolution) and the resulting opaque brown solution stirred at RT for 45 min. A solution of Intermediate 74b (326 mg, 0.750 mmol) in dry DMF (2 mL) was added and the resulting dark brown solution stirred at 60°C for 2.5 h. The solution was concentrated *in vacuo*, redissolved in MeOH (2 mL) and AcOH (0.100 mL), applied to an SCX-2 cartridge and washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* gave a viscous dark brown oil. FCC, using 1-6% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a viscous brown oil (131 mg, 30%). LCMS (Method 3): Rt 3.53

25

min, m/z 578 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(1-hydroxy-1-methyl-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Example 74)


5 A brown solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 89.6 mg, 0.221 mmol), Intermediate 74c (128 mg, 0.221 mmol) and DIPEA (0.048 mL, 0.277 mmol) in dioxane (3 mL) was stirred at 60°C for 16 h, and at 80°C for 6 h. The cooled solution was concentrated *in vacuo*, suspended in water (4 mL) and
10 extracted with DCM (2 × 4 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a viscous brown oil. FCC, using 1-4% [2M NH₃ in MeOH] in DCM) gave 1-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(1-methyl-1-triisopropylsilanyloxy-ethyl)-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-
15 naphthalen-1-yl)-urea as a yellow solid (125 mg).

The solid was dissolved in THF (3 mL), treated with TBAF (1M in THF, 0.161 mL, 0.161 mmol) and the solution stirred at RT for 2 h, and then at reflux for 1 h. The solution was cooled to ~40°C, then TBAF (1M in THF, 0.161 mL, 0.161 mmol) added and the solution stirred at reflux for 16 h. The cooled solution was
20 concentrated *in vacuo*, suspended in water (10 mL) and extracted with EtOAc-MeOH (19:1, 2 × 10 mL). The combined organics were washed with brine (10 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave a dark green residue. FCC, using 2-7% MeOH in DCM, gave the *title compound* as a pale yellow solid (60.6 mg, 41%). LCMS (Method 5): Rt 4.42 min, m/z 677 [MH⁺]. ¹H
25 NMR (400 MHz, d₆-DMSO): 1.09 (6H, s), 1.27 (9H, s), 1.37-1.44 (1H, m), 1.49-1.61 (2H, m), 1.80 (2H, d, J 12.6), 1.83-1.96 (2H, m), 2.01-2.10 (2H, m), 2.36 (3H, s), 2.81-2.90 (2H, m), 3.49 (2H, t, J 11.2), 4.16 (1H, s), 4.81 (1H, td, J 8.5, 5.5), 5.56 (1H, t, J 4.4), 6.32 (1H, s), 7.08 (1H, d, J 8.6), 7.14 (1H, dd, J 9.9, 2.1), 7.27-


7.40 (8H, m), 7.61 (1H, d, J 9.9), 7.66 (1H, d, J 2.0), 8.04 (1H, s).

Example 75

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-pyrrolidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 6-Fluoro-3-pyrrolidin-1-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 75a)



A mixture of Intermediate 24b (515 mg, 3.00 mmol), pyrrolidine (0.85 g, 12.0 mmol) and DMA (10 mL) was heated to 175°C for 4 h using microwave irradiation. After cooling, the solvent was removed *in vacuo* and the resulting residue purified by FCC, using 0 to 15% MeOH in DCM, to provide the *title compound* as a brown gum (285 mg, 26%). LCMS (Method 3): Rt 1.95 min, m/z 207 [MH⁺].

b. (1S,4R)-4-(3-Pyrrolidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 75b)



Intermediate A (222 mg, 1.36 mmol) was added to a mixture of NaH (60% in mineral oil, 163 mg, 4.08 mmol) in DMF (15 mL) at RT and stirred for 30 min. Intermediate 75a (280 mg, 1.36 mmol) was then added and the resulting mixture heated to 60°C for 1 h. After cooling, the reaction was quenched with sat. aq.


NH₄Cl solution. The mixture was diluted with water and extracted with EtOAc. The combined organic extracts were washed with sat. aq. NaHCO₃ solution and brine, and concentrated *in vacuo*. The resulting residue was purified by FCC, using 0 to 15% [2M NH₃ in MeOH] in DCM, to afford the *title compound* as a brown solid (65.0 mg, 14%). LCMS (Method 3): Rt 1.89 min, m/z 350 [MH⁺].

c. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-pyrrolidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 75)


A mixture of Intermediate 75b (62.0 mg, 0.18 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 78.9 mg, 0.195 mmol), DIPEA (46.3 μL, 0.27 mmol) and 1,4-dioxane (2.5 mL) were heated to 60°C for 18 h. After cooling the solvent was evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, then further purified by HPLC (30-98% MeCN in H₂O, 0.1% HCO₂H) to give the *title compound* as an off-white solid (22 mg, 20%). LCMS (Method 5) Rt 4.30 min, m/z 605 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.79-2.15 (8H, m), 2.36 (3H, s), 3.49 (4H, m), 4.81 (1H, m), 5.51 (1H, t, J 4.5), 6.30 (1H, s), 7.03 (1H, dd, J 10.1, 2.2), 7.09 (1H, d, J 8.5), 7.25-7.40 (8H, m), 7.51 (1H, m), 7.83 (1H, m), 8.07 (1H, s).

Example 76

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(1-dimethylamino-1-methyl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt




a. 2-Dimethylamino-2-methyl-propionic acid N'-(5-fluoro-pyridin-2-yl)-

hydrazide (Intermediate 76a)

To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (200 mg, 1.57 mmol) in DMF (10.0 mL) was added 2-dimethylamino-2-methyl-propionic acid (206 mg, 1.57 mmol), EDC (332 mg, 1.73 mmol) and HOBt.H₂O (21.0 mg, 0.16 mmol). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (195 mg, 51%). ¹H NMR (300 MHz, CDCl₃): 1.25 (6H, s), 2.28 (6H, s), 6.44 (1H, br s), 6.62 (1H, dd, J 9.1, 3.6), 7.27 (1H, ddd, J 9.1, 8.0, 3.1), 8.03 (1H, d, J 2.9), 9.04 (1H, br s).


b. [1-(6-Fluoro-[1,2,4]triazolo[4,3-*a*]pyridin-3-yl)-1-methyl-ethyl]-dimethyl-amine (Intermediate 76b)



To a solution of Intermediate 76a (195 mg, 0.81 mmol), Ph₃P (426 mg, 1.62 mmol) and Et₃N (452 μL, 3.25 mmol) in THF (9.00 mL) at 0°C was added hexachloroethane (385 mg, 1.62 mmol). The reaction was stirred at RT overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was dissolved in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH. The reaction showed 50% conversion so was re-submitted to the reaction conditions overnight at 50°C. The workup and purification procedures were repeated to give the *title compound* (153 mg, 85%). ¹H NMR (300 MHz, CDCl₃): 1.60 (6H, s), 2.20 (6H, s), 7.16 (1H, ddd, J 9.9, 7.4, 2.4), 7.71 (1H,

ddd, J 9.9, 5.0, 0.8), 8.89 (1H, ddd, J 4.4, 2.4, 0.8).

c. (1*S*,4*R*)-4-[3-(1-Dimethylamino-1-methyl-ethyl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 76c)



5

To a suspension of NaH (60% in mineral oil, 112 mg, 2.80 mmol) in DMF (2.50 mL) was added Intermediate A (114 mg, 0.70 mmol) and the reaction stirred for 20 min. Intermediate 76b (153 mg, 0.70 mmol) was added in DMF (2.50 mL) and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 2-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (120 mg, 47%). LCMS (Method 4): Rt 0.31, m/z 366 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1*S*,4*R*)-4-[3-(1-dimethylamino-1-methyl-ethyl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea, partial formate salt (Example 76)

To a solution of Intermediate 76c (120 mg, 0.33 mmol) in 1,4-dioxane (3.00 mL) was added DIPEA (117 μL, 0.66 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 133 mg, 0.33 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-8% MeOH in DCM. Further purification by HPLC (C18 X-select column, 25-60% MeCN in H₂O, 0.1% HCO₂H) gave the


20

25


title compound as a white powder after freeze-drying (21 mg, 10%). LCMS (Method 5): Rt 3.86 min, m/z 621.1 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.55 (3H, s), 1.56 (3H, s), 1.88-2.04 (2H, m), 2.10 (1H, m), 2.17 (6H, s), 2.26 (1H, m), 2.38 (3H, s), 4.90 (1H, dd, J 5.7, 8.7), 5.30 (1H, t, J 4.1), 6.33 (1H, s), 7.20-7.36 (9H, m), 7.60-7.64 (1H, d, J 9.9), 8.51 (0.5H, br s), 8.66 (1H, d, J 1.9).

Example 77

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-3-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. [(R)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidin-3-yl]-methanol (Intermediate 77a)




A brown solution of Intermediate 24b (565 mg, 3.29 mmol) and R-3-hydroxymethyl piperidine (Chess GmbH, 948 mg, 8.23 mmol) in DMA (10 mL) was irradiated at 175°C for 3 h in the microwave. The cooled solution was concentrated *in vacuo*, suspended in water (10 mL) and brine (10 mL), and then extracted with DCM (2 × 20 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown oil. FCC, using 4-5% MeOH in DCM, gave the *title compound* as a viscous yellow oil (237 mg, 29%). LCMS (Method 3): Rt 2.11 min, m/z 251 [MH⁺].

b. 6-Fluoro-3-((R)-3-triisopropylsilanyloxymethyl-piperidin-1-yl)-

[1,2,4]triazolo[4,3-a]pyridine (Intermediate 77b)

To a solution of Intermediate 77a (233 mg, 0.931 mmol) and Et₃N (0.195 mL, 1.40 mmol) in dry DCM (5 mL) at RT under N₂, was added triisopropylsilyl trifluoromethanesulfonate (0.313 mL, 1.16 mmol) and the yellow solution stirred at RT for 30 min. Water (5 mL) was added and the mixture shaken. The aqueous was extracted with DCM (5 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a yellow oil. FCC, using 1-5% MeOH in DCM, gave the *title compound* as a yellow oil (333 mg, 88%). LCMS (Method 3): Rt 5.23 min, m/z 407 [MH⁺].


c. (1S,4R)-4-[3-((R)-3-Triisopropylsilanyloxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 77c)



To a solution of Intermediate A (140 mg, 0.860 mmol) in dry DMF (2 mL) at RT under Ar was added NaH (60% dispersion in oil, 49.0 mg, 1.23 mmol) (CARE: gas evolution) and the resulting brown suspension was stirred at RT for 45 min. A solution of Intermediate 77b (333 mg, 0.819 mmol) in dry DMF (2 mL) was added and the resulting dark brown solution stirred at 60°C under Ar for 2.5 h. The solution was concentrated *in vacuo*, redissolved in MeOH (2 mL) and AcOH (0.100 mL), then applied to and SCX-2 cartridge, and washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* left a viscous dark brown oil. FCC, using 2-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a yellow-brown viscous oil (257 mg, 57%).

LCMS (Method 3): Rt 3.27 min, m/z 550 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-3-triisopropylsilanyloxy-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 77d)



5

An orange-brown solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 103 mg, 0.255 mmol), Intermediate 77c (140 mg, 0.255 mmol) and DIPEA (0.055 mL, in dioxane (3 mL) was stirred at 60°C for 16 h and then at 80°C for 2 h. The cooled solution was concentrated *in vacuo*, suspended in water (4 mL) and extracted with DCM (2 × 4 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to ~0.5 mL volume. FCC, using 2-3% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a yellow solid (136 mg, 66%). LCMS (Method 3): Rt 5.65 min, m/z 805 [MH⁺].

10

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-3-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 77)

A brown solution of Intermediate 77d (135 mg, 0.168 mmol) and TBAF (1M in THF, 0.184 mL, 0.184 mmol) in THF (3 mL) was stirred at RT for 1 h. The solution was concentrated *in vacuo*, suspended in water (10 mL) and extracted with EtOAc-MeOH (19:1, 2 × 10 mL). The combined organics were washed with brine (10 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo* to leave a brown solid. FCC, using 5-9% MeOH in DCM, gave a pale brown solid. Further purification by HPLC (XBridge C18 column, 25-75% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (63.9 mg,


20

25

59%). LCMS (Method 5): Rt 4.33 min, m/z 649 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9 H, s), 1.29-1.35 (1H, m), 1.71-1.77 (3H, m), 1.82-1.96 (3H, m), 2.01-2.17 (2H, m), 2.36 (3H, s), 3.02-3.07 (2H, m), 3.20 (1H, dd, J 11.9, 3.3), 3.25-3.28 (1H, m), 3.42 (1H, dt, J 10.5, 5.0), 3.52-3.58 (1H, m), 4.69 (1H, t, J 5.1), 4.81 (1H, td, J 8.6, 5.4), 5.48 (1H, t, J 4.3), 6.32 (1H, s), 7.09 (1H, d, J 8.3), 7.12 (1H, dd, J 9.9, 2.1), 7.27-7.39 (8H, m), 7.60 (1H, d, J 9.9), 7.80 (1H, d, J 2.1), 8.04 (1H, s).

Example 78



1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-3-hydroxymethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



The *title compound* was prepared starting from (S)-3-hydroxymethyl piperidine (Chess GmbH) using analogous procedures to those described in Example 77. LCMS (Method 5): Rt 4.33 min, m/z 649 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.27 (1H, m), 1.69-1.79 (3H, m), 1.82-1.96 (3H, m), 2.03-2.14 (2H, m), 2.36 (3H, s), 2.96 (1H, dd, J 11.9, 8.2), 3.02-3.08 (1H, m), 3.24 (1H, dd, J 11.9, 3.3), 3.29 (1H, m), 3.43 (1H, dt, J 10.6, 5.1), 3.47-3.54 (1H, m), 4.65 (1H, t, J 5.2), 4.81 (1H, td, J 8.5, 5.4), 5.50 (1H, t, J 4.4), 6.32 (1H, s), 7.09 (1H, d, J 8.5), 7.13 (1H, dd, J 9.9, 2.1), 7.27-7.40 (8H, m), 7.60 (1H, d, J 9.9), 7.78 (1H, s), 8.05 (1H, s).

Example 79


1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(4-hydroxy-4-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-

tetrahydro-naphthalen-1-yl}-urea**a. 4-Hydroxy-4-methyl-piperidine-1-carboxylic acid tert-butyl ester****(Intermediate 79a)**

5


To a solution of 1-Boc-4-piperidone (10.0 g, 50.0 mmol) in diethyl ether (100 mL) at 0°C was added methylmagnesium bromide (3.0 M in Et₂O, 22.3 mL, 67.0 mmol) maintaining the temperature below +10°C. The reaction mixture was allowed to warm to RT over 1 h. The reaction was quenched by addition of sat. aq. NH₄Cl solution, then the mixture was extracted with diethyl ether (2 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane, to give the *title compound* (6.76 g, 62%). ¹H NMR (400 MHz, CDCl₃): 1.26 (3H, s), 1.45 (9H, s), 1.48-1.62 (4H, m), 3.15-3.32 (2H, m), 3.70 (2H, d, J 12.5).

15

b. 4-Methyl-piperidin-4-ol (Intermediate 79b)


A solution of Intermediate 79a (5.50 g, 25.6 mmol) in TFA (20 mL) and DCM (40 mL) was stirred at RT for 1 h. The reaction mixture was applied to SCX-2 cartridges (2 × 70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (3.19 g, 99%). ¹H NMR (400 MHz, CDCl₃): 1.24 (3H, s), 1.51-1.61 (4H, m), 2.75-2.87 (2H, m), 2.89-3.02 (2H, m).

20

c. 1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-4-methyl-piperidin-4-ol**(Intermediate 79c)**


A mixture of Intermediate 24b (400 mg, 2.32 mmol) and Intermediate 79b
5 (1.33 g, 11.6 mmol) in NMP (5 mL) was heated in the microwave at 170°C for 3 h. The reaction mixture was applied to an SCX-2 cartridge (25 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-8% [2M NH₃ in MeOH] in DCM, gave the *title compound* (190 mg, 32%). LCMS (Method 1): Rt 2.01 min, m/z 251 [MH⁺].

10 **d. 6-Fluoro-3-(4-methyl-4-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 79d)**




Triisopropylsilyl trifluoromethanesulfonate (550 mg, 1.81 mmol) was added to a solution of Intermediate 79c (150 mg, 0.60 mmol) and Et₃N (242 mg, 2.40
15 mmol) in a DCM (5 mL) and the mixture stirred at reflux for 1 h. The reaction mixture was diluted with DCM (30 mL), washed with water (2 ×), brine, dried (MgSO₄), and then concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* (235 mg, 96%). ¹H NMR (400 MHz, CDCl₃): 1.00-1.20 (21H, m), 1.40 (3H, s), 1.76-1.93 (4H, m), 3.18-3.29 (2H, m), 3.44-3.58 (2H, m), 7.03-7.12 (1H, m), 7.58-7.69 (2H, m).
20

e. (1S,4R)-4-[3-(4-Methyl-4-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 79e)



5 To a solution of Intermediate A (111 mg, 0.680 mmol) in DMF (2 mL) was added NaH (60% in oil, 68 mg, 1.70 mmol) and the mixture stirred at RT for 20 min, before Intermediate 79d (230 mg, 0.566 mmol) was added. This mixture was stirred at 60°C for 1 h. The cooled reaction mixture was diluted with EtOAc (100 mL), washed with water (2 ×), brine, dried (MgSO₄) and concentrated *in vacuo*.
 10 The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (160 mg, 51%). LCMS (Method 1): Rt 3.06 min, m/z 550 [MH⁺].

f. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-methyl-4-triisopropylsilanyloxy-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 79f)



A solution of Intermediate 79e (160 mg, 0.291 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 176 mg, 0.437 mmol) and DIPEA (150
 20 mg, 1.16 mmol) in DMF (4 mL) was stirred at 50°C for 1 h. The reaction mixture was diluted with EtOAc and washed with water (2 ×), brine, dried (MgSO₄),


filtered and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (210 mg, 93%). LCMS (Method 4): Rt 5.32 min, m/z 805 [MH⁺].

g. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(4-hydroxy-4-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 79)


To a solution of Intermediate 79f (210 mg, 0.260 mmol) in THF (4 mL) at -30°C was added TBAF (1M in THF, 390 µL, 0.390 mmol) and the mixture was allowed to warm to RT over 1 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, and then recrystallisation from boiling EtOAc (~40 mL) gave the *title compound* as a white powder (67 mg, 40%). LCMS (Method 5): Rt 4.31 min, m/z 649.1 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.30 (3H, s), 1.33 (9H, s), 1.71-1.77 (2H, m), 1.84-1.93 (2H, m), 1.95-2.17 (3H, m), 2.25-2.32 (1H, m), 2.41 (3H, s), 3.20-3.28 (2H, m), 3.36-3.45 (2H, m), 4.72-4.92 (1H, m), 5.47 (1H, t, J 4.4), 6.36 (1H, s), 6.88 (1H, d, J 9.3), 7.21-7.40 (8H, m), 7.58 (1H, d, J 9.7), 7.62 (1H, d, J 1.7).

Example 80

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1-methyl-1-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 2-Methyl-2-pyrrolidin-1-yl-propionic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 80a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (200 mg, 1.57 mmol) in
5 DMF (10.0 mL) was added 2-methyl-2-pyrrolidin-1-yl-propionic acid (246 mg, 1.57 mmol), EDC (332 mg, 1.73 mmol) and HOBt.H₂O (21.0 mg, 0.16 mmol). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The
10 residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (260 mg, 62%). ¹H NMR (300 MHz, CDCl₃): 1.30 (6H, s), 1.76-1.84 (4H, m), 2.64-2.72 (4H, m), 6.46 (1H, br s), 6.62 (1H, dd, J 9.0, 3.6), 7.28 (1H, ddd, J 9.0, 8.0, 2.9), 8.03 (1H, d, J 2.9), 9.06 (1H, br s).


**b. 6-Fluoro-3-(1-methyl-1-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-
15 a]pyridine (Intermediate 80b)**



To a solution of Intermediate 80a (260 mg, 0.98 mmol), Ph₃P (511 mg, 1.95 mmol) and Et₃N (544 μL, 3.91 mmol) in THF (10.0 mL) at 0°C was added hexachloroethane (462 mg, 1.95 mmol). The reaction was stirred at RT overnight
20 then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH. The reaction showed 50% conversion so was re-submitted
25 to the reaction conditions overnight at 50°C. The workup and purification procedures were repeated to give the *title compound* (209 mg, 86%). ¹H NMR (300

MHz, CDCl₃): 1.64 (6H, s), 1.74-1.82 (4H, m), 2.52-2.57 (4H, m), 7.16 (1H, ddd, J 9.8, 7.4, 2.3), 7.69 (1H, m), 8.93 (1H, ddd, J 4.5, 2.3, 0.7).

c. (1*S*,4*R*)-4-[3-(1-Methyl-1-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 80c)



To a suspension of NaH (60% in mineral oil, 135 mg, 3.37 mmol) in DMF (4.50 mL) was added Intermediate A (137 mg, 0.84 mmol) and the reaction stirred for 20 min. Intermediate 80b (209 mg, 0.84 mmol) was added in DMF (1.50 mL) and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH and product eluted with 2M NH₃ in MeOH. The resulting residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (180 mg, 55%). LCMS (Method 4): Rt 0.31, m/z 392.2 [MH⁺].


d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1*S*,4*R*)-4-[3-(1-methyl-1-pyrrolidin-1-yl-ethyl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 80)

To a solution of Intermediate 80c (180 mg, 0.46 mmol) in 1,4-dioxane (5.00 mL) was added DIPEA (160 μL, 0.92 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 185 mg, 0.46 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-8% MeOH in DCM. Further purification


by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (58 mg, 20%). LCMS (Method 5): Rt 3.84 min, m/z 647.2 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.60 (6H, s), 1.70-1.76 (4H, m), 1.86-2.06 (2H, m), 2.11 (1H, m), 2.20 (1H, m), 2.38 (3H, s), 2.45-2.52 (2H, m), 2.55-2.62 (2H, m), 4.89 (1H, dd, J 5.6, 8.9), 5.29 (1H, t, J 4.2), 6.33 (1H, s), 7.20-7.36 (9H, m), 7.62 (1H, d, J 9.9), 8.77 (1H, d, J 2.0).

Example 81

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (S)-2-Methyl-piperidine-1-carbonyl chloride (Intermediate 81a)



To a pale yellow solution of triphosgene (742 mg, 2.50 mmol) and pyridine (0.404 mL, 5.00 mmol) in dry DCM (10 mL) at 10°C under N₂ was added (S)-2-methyl piperidine (Aldrich, 0.603 mL, 5.00 mmol) cautiously over 2 min. The vivid orange solution was stirred at RT for 18 h, then pyridine (0.404 mL, 5.00 mmol) and triphosgene (742 mg, 2.50 mmol) were added sequentially (CARE: exotherm on addition of triphosgene) and the orange solution stirred at RT for 4 h. Aqueous HCl solution (1M, 10 mL) was added and the mixture stirred vigorously until gas evolution ceased (30 min). The aqueous layer was extracted with DCM (10 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a dark red oil (745 mg, 92%).

¹H NMR (300 MHz, CDCl₃): 1.25 (3H, d, J 7.0), 1.42-1.77 (6H, m), 3.07 (1H, br s), 4.17 (1H, dd, J 13.8, 4.0), 4.62 (1H, apparent quin, J 6.3).


b. (S)-2-Methyl-piperidine-1-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 81b)



5

To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (582 mg, 4.58 mmol) and DIPEA (0.997 mL, 5.72 mmol) in dry DCM (20 mL) was added Intermediate 81a (740 mg, 4.58 mmol) and the resulting red solution stirred at RT for 66 h, and at reflux for 4 h. To the cooled solution was added water (20 mL) and the mixture
10 shaken. The aqueous was extracted with DCM (20 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a pale brown solid. FCC, using 0-5% MeOH in DCM, gave the *title compound* as a pale yellow solid (657 mg, 57%). ¹H NMR (300 MHz, CDCl₃): 1.22 (3H, d, J 6.9), 1.40-1.77 (6H, m), 2.96 (1H, td, J 13.0, 2.9), 3.85 (1H, ddd, J 13.3, 4.5, 2.2), 4.31
15 (1H, m), 6.50 (1H, s), 6.59 (1H, br s), 6.75 (1H, ddd, J 9.0, 3.6, 0.7), 7.27 (1H, ddd, J 9.1, 8.1, 3.1), 8.02 (1H, d, J 2.9).


c. 6-Fluoro-3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 81c)



To a solution of Intermediate 81b (657 mg, 2.60 mmol), Ph₃P (1.37 g, 5.21 mmol) and Et₃N (1.45 mL, 10.4 mmol) in THF (20 mL) at 0°C was added hexachloroethane (1.23 g, 5.21 mmol) and the resulting yellow suspension was stirred at RT for 2 h, and at reflux for 19 h. The cooled solution was filtered and concentrated *in vacuo*. The brown oil was redissolved in MeOH (3 mL) and
20 applied to an SCX-2 cartridge (20 g), washing with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* gave a brown

oil. FCC, using 1-5% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown oil (542 mg, 89%). LCMS (Method 3): Rt 3.00 min, m/z 235 [MH⁺].

d. (1S,4R)-4-[3-((S)-2-Methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 81d)




To a solution of Intermediate A (374 mg, 2.29 mmol) in dry DMF (3 mL) under N₂ was added NaH (60% dispersion in oil, 183 mg, 4.58 mmol) and the resulting opaque brown solution was stirred at RT for 45 min (CARE: gas evolution). A solution of Intermediate 81c (537 mg, 2.29 mmol) in dry DMF (7 mL) was added and the dark brown solution stirred at 60°C under N₂ for 90 min. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (3 mL) and AcOH (0.10 mL) and then applied to an SCX-2 cartridge (20 g), washing with MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (50 mL); concentration *in vacuo* gave a dark brown solid. FCC, using 2-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a pale brown foam (505 mg, 58%). LCMS (Method 3): Rt 2.29 min, m/z 378 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea. (Example 81)

An orange-brown solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 106 mg, 0.263 mmol), Intermediate 81d (94.4 mg, 0.250 mmol) and DIPEA (0.054 mL, 0.313 mmol) in dry dioxane (3 mL) was stirred at 70°C for 16 h. The cooled solution was concentrated *in vacuo*, suspended in water (5 mL) and extracted with DCM (2 × 5 mL). The combined organics were passed through a

hydrophobic frit and concentrated *in vacuo* to leave a brown gum. FCC, using 2-6% MeOH in DCM, gave a pale yellow solid (123 mg). Further purification by HPLC (XBridge C18 column, 40-90% MeCN in H₂O, 0.1% NH₄OH) gave the *title compound* as a white powder after freeze-drying (72.4 mg, 46%). LCMS (Method 5): Rt 5.04 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J 6.3), 1.27 (9H, s), 1.48-1.55 (2H, m), 1.63-1.71 (2H, m), 1.75-1.97 (4H, m), 2.00-2.16 (2H, m), 2.36 (3H, s), 2.90 (1H, ddd, J 12.1, 9.0, 3.9), 3.16 (1H, dt, J 12.1, 4.3), 3.29 (1H, m), 4.82 (1H, td, J 8.6, 5.5), 5.51 (1H, t, J 4.4), 6.32 (1H, s), 7.08 (1H, d, J 8.6), 7.19 (1H, dd, J 9.9, 2.2), 7.25-7.38 (8H, m), 7.64 (1H, dd, J 9.8, 0.8), 7.69 (1H, d, J 2.1), 8.04 (1H, s).

Example 82




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

The *title compound* was prepared starting from (R)-2-methyl piperidine (ABCR) using analogous procedures to those described in Example 81. LCMS (Method 5): Rt 5.03 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.88 (3H, d, J 6.3), 1.27 (9H, s), 1.48-1.55 (2H, m), 1.65-1.71 (2H, m), 1.76-1.98 (4H, m), 2.01-2.14 (2H, m), 2.36 (3H, s), 2.94 (1H, ddd, J 12.2, 8.1, 4.8), 3.17 (1H, dt, J 12.1, 4.3), 3.26-3.30 (1H, m), 4.82 (1H, td, J 8.6, 5.4), 5.54 (1H, t, J 4.4), 6.32 (1H, s), 7.08 (1H, d, J 8.6), 7.18 (1H, dd, J 9.8, 2.2), 7.26-7.38 (8H, m), 7.64 (1H, d, J 9.9), 7.70 (1H, d, J 2.1), 8.04 (1H, s).


Example 83

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1*S*,4*R*)-4-[3-[(*S*)-4-methyl-morpholin-3-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea, partial formate salt



5


a. (*R*)-3-[*N'*-(5-Fluoro-pyridin-2-yl)-hydrazinocarbonyl]-morpholine-4-carboxylic acid tert-butyl ester (Intermediate 83a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (143 mg, 1.13 mmol) in DCM (10.0 mL) was added (*R*)-morpholine-3,4-dicarboxylic acid 4-tert-butyl ester (260 mg, 1.13 mmol), EDC (238 mg, 1.24 mmol) and HOBt.H₂O (15.0 mg, 0.11 mmol). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (375 mg, 98%). LCMS (Method 1): Rt 2.69, m/z 341.2 [MH⁺].

10

b. (*S*)-3-(6-Fluoro-[1,2,4]triazolo[4,3-*a*]pyridin-3-yl)-morpholine-4-carboxylic acid tert-butyl ester (Intermediate 83b)




20

To a solution of Intermediate 83a (375 mg, 1.10 mmol), Ph₃P (578 mg, 2.20 mmol) and Et₃N (614 μL, 4.41 mmol) in THF (11.0 mL) at 0°C was added

hexachloroethane (523 mg, 2.20 mmol). The reaction was stirred at RT overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (297 mg, 84%). LCMS (Method 1): Rt 2.73, m/z 323.2 [MH⁺].


c. 6-Fluoro-3-((S)-4-methyl-morpholin-3-yl)-[1,2,4]triazolo[4,3-*a*]pyridine (Intermediate 83c)



To a solution of Intermediate 83b (297 mg, 0.92 mmol) in DCM (8.0 mL) at 0°C was added TFA (400 µL) and the reaction stirred at RT overnight. Further TFA (600 µL) was added and the reaction stirred for 1 h then partitioned between DCM and sat. aq. NaHCO₃ solution. The aqueous layer was then extracted with DCM (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in DCM (4.0 ml) and MeOH (3 drops) added. Formaldehyde solution (37 wt% in water, 162 µL, 2.00 mmol) was added followed by sodium triacetoxyborohydride (372 mg, 2.40 mmol). The reaction was stirred overnight then loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* (75.0 mg, 34%). ¹H NMR (400 MHz, d₄-MeOD): 2.09 (3H, s), 2.49 (1H, td, J 11.4, 3.8), 2.98 (1H, dt, J 11.9, 2.0), 3.79-3.90 (3H, m), 3.94 (1H, m), 4.01 (1H, dd, J 10.1, 4.0), 7.50 (1H, ddd, J 10.0, 7.7, 2.2), 7.81 (1H, ddd, J 10.1, 4.9, 0.8), 8.98 (1H, ddd, J 4.0, 2.3, 0.8).

d. (1S,4R)-4-[3-((S)-4-Methyl-morpholin-3-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate

83d)



To a suspension of NaH (60% in mineral oil, 51.0 mg, 1.27 mmol) in DMF (1.50 mL) was added Intermediate A (51.0 mg, 0.32 mmol) and the reaction stirred
 5 for 20 min. Intermediate 83c (75.0 mg, 0.32 mmol) was added in DMF (1.50 mL) and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10%
 10 [2M NH₃ in MeOH] in DCM, gave the *title compound* (31 mg, 26%). LCMS (Method 4): Rt 0.32, m/z 380.0 [MH⁺].


e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-4-methyl-morpholin-3-yl)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea, partial formate salt (Example 83)

15 To a solution of Intermediate 83d (31.0 mg, 0.082 mmol) in 1,4-dioxane (2.00 mL) was added DIPEA (28.0 μL, 0.16 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 33.0 mg, 0.082 mmol). The reaction was heated to 60°C overnight then cooled and partitioned between EtOAc and water.
 20 The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (10 mg,
 25 19%). LCMS (Method 5): Rt 3.90 min, m/z 635.2 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.86-2.06 (3H, m), 2.11 (1H, m), 2.08 (3H, s), 2.25 (1H, m),


2.38 (3H, s), 2.46 (1H, td, J 11.7, 3.5), 2.92 (1H, dt, J 11.8, 1.7), 3.73-3.91 (3H, m), 3.97 (1H, dd, J 9.9, 4.0), 4.90 (1H, dd, J 8.6, 5.6), 5.39 (1H, t, J 4.1), 6.33 (1H, s), 7.21-7.36 (9H, m), 7.65 (1H, d, J 9.9), 8.24 (0.4H, br s), 8.55 (1H, d, J 1.4).

Example 84

5 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[(S)-1-(3-hydroxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea**




a. **(S)-1-(2-tert-Butoxycarbonyl-ethyl)-pyrrolidine-2-carboxylic acid benzyl ester (Intermediate 84a)**




Tert-butyl acrylate (5.00 mL) was added to a solution of L-proline benzyl ester hydrochloride salt (1.10 g, 4.54 mmol) and Et₃N (632 μL, 4.54 mmol) in tert-butanol (5.00 mL). The reaction was heated to 80°C for 3 h then and evaporated *in vacuo*. The residue was purified by FCC, using 0-50% EtOAc in cyclohexane, to give the *title compound* (1.20 g, 79%). ¹H NMR (300 MHz, CDCl₃): 1.43 (9H, s), 1.72-2.00 (3H, m), 2.08 (1H, m), 2.39-2.48 (3H, m), 2.72 (1H, ddd, J 12.2, 8.4, 6.7), 3.00 (1H, dt, J 12.3, 7.8), 3.13 (1H, m), 3.28 (1H, dd, J 8.6, 5.4), 5.13 (1H, d, J 12.3), 5.18 (1H, d, J 12.3), 7.29-7.39 (5H, m).

b. (S)-1-(2-tert-Butoxycarbonyl-ethyl)-pyrrolidine-2-carboxylic acid (Intermediate 84b)




A solution of Intermediate 84a (1.10 g, 3.29 mmol) in IMS (25.0 mL) was added to palladium on charcoal (10 wt%, 110 mg) and the reaction stirred under H₂ for 1 h. The mixture was filtered through Celite and evaporated *in vacuo* to give the *title compound* (840 mg, 99%). ¹H NMR (300 MHz, d₄-MeOD): 1.48 (9H, s), 1.91 (1H, m), 2.02-2.24 (2H, m), 2.41 (1H, m), 2.74 (2H, t, J 7.1), 3.13 (1H, ddd, J 11.1, 10.2, 7.0), 3.32-3.53 (2H, m), 3.72 (1H, ddd, J 11.0, 7.2, 3.5), 3.89 (1H, dd, J 9.5, 5.3).

c. 3-{(S)-2-[N'-(5-Fluoro-pyridin-2-yl)-hydrazinocarbonyl]-pyrrolidin-1-yl}-propionic acid tert-butyl ester (Intermediate 84c)




HOBt.H₂O (695 mg, 3.62 mmol) was added to a suspension of (5-fluoro-pyridin-2-yl)-hydrazine (418 mg, 3.29 mmol), Intermediate 84b (840 mg, 3.29 mmol), and EDC (44.0 mg, 0.33 mmol) in DMF (25.0 mL). The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (1.16 g, 99%). LCMS (Method 1): Rt 2.06 mins, m/z 353.3 [MH⁺].

d. 3-[(S)-2-(6-Fluoro-[1,2,4]triazolo[4,3-*a*]pyridin-3-yl)-pyrrolidin-1-yl]-propionic acid tert-butyl ester (Intermediate 84d)




Hexachloroethane (1.56 g, 6.58 mmol) was added to a solution of
5 Intermediate 84c (1.16 g, 3.29 mmol), Ph₃P (1.72 g, 6.58 mmol) and Et₃N (1.80 mL, 13.1 mmol) in THF (30.0 mL) at 0°C. The reaction was stirred overnight then partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC,
10 using EtOAc then 10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (1.06 g, 99%). LCMS (Method 4): Rt 1.97 min, m/z 335.1 [MH⁺].

e. 3-[(S)-2-(6-Fluoro-[1,2,4]triazolo[4,3-*a*]pyridin-3-yl)-pyrrolidin-1-yl]-propan-1-ol (Intermediate 84e)




15 Lithium aluminium hydride solution (2M in THF, 1.64 mL, 3.28 mmol) was added to a solution of Intermediate 84d (550 mg, 1.64 mmol) in THF (15.0 mL) at -10°C. The reaction was stirred for 1 h then quenched by dropwise addition of water (125 µL), sodium hydroxide (3M aqueous, 125 µL) and water (375 µL). The reaction was stirred for 10 min then partitioned between EtOAc and water. The
20 aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 2-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (145 mg, 33%). LCMS (Method 4): Rt 0.32 mins, m/z 265.2 [MH⁺].

f. 6-Fluoro-3-[(S)-1-(3-triisopropylsilanyloxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-*a*]pyridine (Intermediate 84f)



Triisopropylsilyl chloride (266 μ L, 1.24 mmol) was added to a solution of
5 Intermediate 84e (218 mg, 0.82 mmol) and Et₃N (266 μ L, 1.24 mmol) in DCM
(8.00 mL). The reaction was stirred overnight then 4-(dimethylamino)pyridine
(10.0 mg, 0.082 mmol) added. The reaction was stirred for 3 h then partitioned
between EtOAc and water. The aqueous layer was then extracted with EtOAc (3
 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered
10 and evaporated *in vacuo*. The residue was purified by FCC, using 0-7.5% [2M
NH₃ in MeOH] in DCM, to give the *title compound* (317 mg, 92%). ¹H NMR (300
MHz, CDCl₃): 0.92 (21H, s), 1.56-1.68 (2H, m), 1.89-2.10 (3H, m), 2.21-2.40 (3H,
m), 2.56 (1H, dt, J 12.1, 7.9), 3.40 (1H, m), 3.46-3.64 (2H, m), 4.22 (1H, t, J 8.4),
7.16 (1H, ddd, J 9.9, 7.5, 2.3), 7.71 (1H, ddd, J 10.0, 4.9, 0.6), 8.64 (1H, m).


15 **g. (1S,4R)-4-{3-[(S)-1-(3-Triisopropylsilanyloxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 84g)**



To a suspension of NaH (60% in mineral oil, 121 mg, 3.02 mmol) in DMF
20 (2.00 mL) was added Intermediate A (122 mg, 0.75 mmol) and the reaction stirred
for 20 min. Intermediate 84f (317 mg, 0.75 mmol) was added in DMF (2.00 mL)
and the reaction heated to 60°C for 1 h. The cooled reaction was quenched by
dropwise addition of methanol, before being diluted with methanol and loaded
onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted

with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 2-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* (140 mg, 33%). LCMS (Method 4): Rt 2.37, m/z 564.2 [MH⁺].

h. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[(S)-1-(3-triisopropylsilyloxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Intermediate 84h)**



To a solution of Intermediate 84g (140 mg, 0.25 mmol) in 1,4-dioxane (3.00 mL) was added DIPEA (86.0 μL, 0.50 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 100 mg, 0.25 mmol). The reaction was heated to 60°C overnight then further (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (50.0 mg, 0.12 mmol) was added and heating at 60°C continued overnight. The reaction was cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (70.0 mg, 34%). LCMS (Method 1): Rt 3.46, m/z 820.4 [MH⁺].


i. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[(S)-1-(3-hydroxy-propyl)-pyrrolidin-2-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Example 84)**

TBAF (1M in THF, 100 μL, 0.10 mmol) was added to a solution of Intermediate 84h (70.0 mg, 0.085 mmol) in THF (2.00 mL). The reaction was stirred for 2 h then partitioned between EtOAc and water. The aqueous layer was


then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (29 mg, 52%). LCMS (Method 5): Rt 3.63 min, m/z 663.3 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.30 (9H, s), 1.54-1.62 (2H, quin, J 7.0), 1.88-2.15 (6H, m), 2.20-2.37 (4H, m), 2.38 (3H, s), 2.55 (1H, dt, J 12.0, 8.1), 3.32-3.48 (3H, m), 4.11 (1H, t, J 8.0), 4.90 (1H, dd, J 9.1, 5.7), 5.31 (1H, t, J 4.0), 6.33 (1H, s), 7.20-7.36 (9H, m), 7.65 (1H, d, J 9.9), 8.32 (1H, d, J 1.7).

Example 85

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-hydroxymethyl-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (3R,7aS)-3-Trichloromethyl-tetrahydro-pyrrolo[1,2-c]oxazol-1-one (Intermediate 85a)



A suspension of L-proline (10.0 g, 87.0 mmol) and chloral hydrate (21.6 g, 130 mmol) in chloroform (100 mL) was heated at reflux for 6 h under a soxlet adapter charged with sodium sulfate. The cooled mixture was washed with water (2 ×), dried (MgSO₄) and concentrated *in vacuo*. The product was recrystallised from boiling absolute ethanol to give the *title compound* as white needles (13.3 g, 63%). ¹H NMR (400 MHz, CDCl₃): 1.65-1.83 (1H, m), 1.86-2.01 (1H, m), 2.04-

2.31 (2H, m), 3.06-3.18 (1H, m), 3.36-3.49 (1H, m), 4.12 (1H, dd, J 4.6, 4.0), 5.16 (1H, s).

b. (3R,7aR)-7a-Benzylloxymethyl-3-trichloromethyl-tetrahydropyrrolo[1,2-c]oxazol-1-one (Intermediate 85b)




5

To a solution of LDA (531 mL, 0.109M, 58.1 mmol) at -78°C was added a solution of Intermediate 85a (10.0 g, 41.2 mmol) in THF (200 mL) over 30 min maintaining the temperature below -70°C . The mixture was stirred at -78°C for 2 h, then a solution of benzyloxymethyl chloride (11.6 g, 74.1 mmol) in THF (50 mL) added, maintaining the temperature below -70°C . The mixture was allowed to warm to -30°C over 3 h before being quenched with dropwise addition of water (300 mL). The aqueous was extracted with EtOAc (2 $\times$). The combined organic layers were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by FCC, using 0-60% diethyl ether in cyclohexane, to give the *title compound* as a clear oil (6.39 g, 42%). ^1H NMR (400 MHz, CDCl_3): 1.63-1.77 (1H, m), 1.94-2.04 (1H, m), 2.08-2.17 (1H, m), 2.26-2.36 (1H, m), 3.22-3.29 (1H, m), 3.30-3.38 (1H, m), 3.74, (2H, s), 4.62 (2H, s), 4.99 (1H, s), 7.26-7.37 (5H, m).

15

c. (R)-2-Benzylloxymethyl-pyrrolidine-2-carboxylic acid methyl ester (Intermediate 85c)




20

A solution of Intermediate 85b (6.39 g, 17.9 mmol) and sodium methoxide (580 mg, 10.8 mmol) in MeOH (100 mL) was stirred at RT for 18 h. The mixture was cooled to 0°C , then acetyl chloride (28.4 g, 359 mmol) added. This mixture was stirred at reflux for 1 h, then cooled and concentrated *in vacuo*. The residue was divided, applied to SCX-2 cartridges (3 $\times$ 70 g) and washed with MeOH. The product was eluted with 2M NH_3 in MeOH; concentration *in vacuo* gave the *title*

25


compound as a pale yellow oil (3.92 g, 87%). ¹H NMR (400 MHz, CDCl₃): 1.62-1.86 (3H, m), 1.07-2.13 (1H, m), 2.48 (1H, br s), 2.93-3.11 (2H, m), 3.69-3.78 (4H, m), 4.53 (1H, d, J 12.9), 4.55 (1H, d, J 12.9), 7.21-7.37 (5H, m).

d. (R)-2-Benzyloxymethyl-1-methyl-pyrrolidine-2-carboxylic acid methyl ester (Intermediate 85d)




A solution of Intermediate 85c (3.37 g, 13.5 mmol) and formaldehyde (37% aqueous, 4.38 mL, 54.1 mmol) in DCM (140 mL) was stirred at RT for 30 min, then sodium triacetoxyborohydride (5.74 g, 27.1 mmol) was added. This mixture was stirred at RT for 18 h then washed with sat. aq. NaHCO₃ solution, dried (MgSO₄) and concentrated in *vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a clear oil (2.59 g, 72%). LCMS (Method 4): Rt 1.85 min, m/z 263 [MH⁺].

e. (R)-2-Hydroxymethyl-1-methyl-pyrrolidine-2-carboxylic acid methyl ester (Intermediate 85e)




A suspension of Intermediate 85d (2.59 g, 9.84 mmol), palladium hydroxide on charcoal (10%, 250 mg) and glacial acetic acid (1 mL) in IMS (20 mL) under an atmosphere of hydrogen (1 atm) was stirred at RT for 5 d. The mixture was filtered and concentrated in *vacuo* to give the *title compound* as a clear oil (1.70 g, 99%), as a mixture of methyl and ethyl esters. Data for major product, methyl ester: ¹H NMR (400 MHz, CDCl₃): 1.75-1.95 (2H, m), 2.01-2.18 (2H, m), 2.36 (3H, s), 2.86-2.99 (2H, m), 3.65 (3H, s), 4.19 (2H, m), 4.91 (1H, br s).

f. (R)-1-Methyl-2-triisopropylsilanyloxymethyl-pyrrolidine-2-carboxylic acid methyl ester (Intermediate 85f)




Triisopropylsilyl trifluoromethanesulfonate (11.7 g, 38.1 mmol) was added
5 to a solution of Intermediate 85e (2.20 g, 12.7 mmol) and Et₃N (12.8 g, 127 mmol)
in DCM (100 mL) at 0°C, then the mixture stirred at RT for 30 min. The mixture
was diluted with DCM and washed with 10% aq. K₂CO₃ solution, water, brine,
dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by FCC, using
0-5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale yellow
10 solid (2.51 g, 65%). ¹H NMR (400 MHz, CDCl₃): 0.97-1.15 (21H, m), 1.72-1.96
(2H, m), 2.23-2.35 (2H, m), 2.46 (3H, s), 2.89-2.99 (2H, m), 3.69 (3H, m), 4.12-
4.22 (2H, m).

g. (R)-1-Methyl-2-triisopropylsilanyloxymethyl-pyrrolidine-2-carboxylic acid (Intermediate 85g)




15 A solution of Intermediate 85f (2.50 g, 7.93 mmol) and lithium hydroxide
monohydrate (3.33 g, 79.3 mmol) in MeOH and water (50mL) was stirred at reflux
for 2 h. The cooled solution was acidified with 1M HCl to pH 5. The MeOH was
removed *in vacuo* and the resulting aqueous layer was extracted with EtOAc (3 ×).
20 The combined organic layers were dried (MgSO₄) and concentrated *in vacuo* to
give the *title compound* as a white foam (2.21 g, 92%). ¹H NMR (400 MHz,
CDCl₃): 1.00-1.21 (21H, m), 1.93-2.04 (2H, m), 2.32-2.42 (1H, m), 2.43-2.56 (1H,
m), 2.96-3.09 (4H, m), 3.90-4.00 (1H, m), 4.20 (1H, d, J 10.9), 4.22 (1H, d, J
10.9).

25 **h. (R)-1-Methyl-2-triisopropylsilanyloxymethyl-pyrrolidine-2-**

carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 85h)


To a solution of Intermediate 85g (1.08 g, 3.58 mmol), (5-fluoro-pyridin-2-yl)-hydrazine (500 mg, 3.93 mmol) and DIPEA (1.38 g, 10.7 mmol) in DCM (120 mL) was added HATU (2.04 g, 5.37 mmol) and the mixture stirred at RT for 2 h. The solution was washed with 10% aq. K₂CO₃ solution, brine, dried (MgSO₄), and concentrated *in vacuo*. The residue was purified by FCC, using 0-6% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale yellow oil (1.37 g, 90%). LCMS (Method 4): Rt 2.74 min, m/z 425 [MH⁺].

i. 6-Fluoro-3-((S)-1-methyl-2-triisopropylsilyloxymethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 85i)




To a solution of Intermediate 85h (1.37 g, 2.08 mmol), Ph₃P (1.09 g, 4.16 mmol) and Et₃N (840 mg, 8.32 mmol) in THF (60 mL) was added hexachloroethane (980 mg, 4.16 mmol) and the mixture stirred at RT for 20 h. The mixture was diluted with DCM, washed with 10% aq. K₂CO₃ solution, brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane, to give the *title compound* as a clear oil (350 mg, 41%). LCMS (Method 4): Rt 3.07 min, m/z 407 [MH⁺].

j. (1S,4R)-4-[3-((S)-1-Methyl-2-triisopropylsilanyloxymethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 85j)



5 To a solution of Intermediate A (170 mg, 1.03 mmol) in DMF (4 mL) was added NaH (60% in oil, 103 mg, 2.59 mmol) and the mixture stirred at RT for 20 min, before Intermediate 85i (350 mg, 0.862 mmol) was added. This mixture was stirred at 60°C for 4 h. The cooled mixture was diluted with EtOAc, washed with water (3 ×), brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was
 10 purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a clear viscous oil (230 mg, 48%). LCMS (Method 1): Rt 2.28 min, m/z 550 [MH⁺].

k. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1-methyl-2-triisopropylsilanyloxymethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-
 15 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 85k)



A solution of Intermediate 85j (230 mg, 0.418 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic
 20 Communications, 2009, 39, 3999-4009; 253 mg, 0.628 mmol) and DIPEA (215 mg, 1.67 mmol) in DMF (4 mL) was stirred at 50°C for 1 h. The mixture was


diluted with EtOAc, washed with water (2 ×), brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a white foam (250 mg, 48%). LCMS (Method 3): Rt 3.70 min, m/z 805 [MH⁺].

5 **1. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-hydroxymethyl-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 85)**


To a solution of Intermediate 85k (250 mg, 0.310 mmol) in THF (10 mL) at -30°C was added TBAF (1M in THF, 620 µL, 0.620 mmol) and the mixture was
10 allowed to warm to RT over 2 h. The reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave a crude product. Recrystallisation from hot diethyl ether (50 mL), EtOAc (10 mL) and cyclohexane (20 mL) gave the *title compound* as a white
15 powder (113 mg, 56%). LCMS (Method 5): Rt 3.68 min, m/z 649.1 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.29 (9H, s), 1.86-2.34 (11H, m), 2.38 (3H, s), 3.02 (1H, m), 3.09 (1H, dt, J 8.8, 3.5), 4.07 (1H, d, J 11.0), 4.25 (1H, d, J 4.3), 4.52 (1H, br s), 4.90 (1H, m), 5.25 (1H, t, J 4.3), 6.33 (1H, s), 7.18-7.36 (9H, m), 7.62 (1H, d J 7.6), 8.44 (1H, s).

20 **Example 86**

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-(1,4-dimethyl-piperazin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




a. **2-[N'-(5-Fluoro-pyridin-2-yl)-hydrazinocarbonyl]-piperazine-1,4-dicarboxylic acid di-tert-butyl ester (Intermediate 86a)**



To a solution of piperazine-1,2,4-tricarboxylic acid 1,4-di-tert-butyl ester
 5 (780 mg, 2.4 mmol) and (5-fluoro-pyridin-2-yl)-hydrazine (250 mg, 2.0 mmol) and
 Et₃N (0.83 mL, 6 mmol) in DCM (20 mL) were added HOBt.H₂O (27 mg, 0.2
 mmol) and EDC (542 mg, 2.4 mmol) sequentially, and then the mixture stirred at
 RT overnight. The mixture was partitioned between EtOAc/water and extracted
 with EtOAc. The combined organics were dried over Na₂SO₄, filtered and
 10 concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in
 DCM, to give the *title compound* as a brown solid (550 mg, 63%). LCMS (Method
 4): Rt 3.28, m/z 440 [MH⁺].

b. **6-Fluoro-3-piperazin-2-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 86b)**




15 To a solution of Intermediate 86a (550 mg, 1.25 mmol), Et₃N (0.69 mL, 5
 mmol) and Ph₃P (657 mg, 2.5 mmol) in THF (20 mL) at 0°C was added
 hexachloroethane (590 mg, 2.5 mmol) and the mixture stirred for 10 min. The
 solution was allowed to warm to RT, stirred for 1 h, then partitioned between
 20 EtOAc/water and extracted with EtOAc. The combined organics were dried over
 Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by FCC,
 using 0-100% EtOAc in cyclohexane, to give crude 2-(6-fluoro-[1,2,4]triazolo[4,3-
 a]pyridin-3-yl)-piperazine-1,4-dicarboxylic acid di-tert-butyl ester, contaminated
 with triphenylphosphine oxide, as a pale brown solid (500 mg).

25 The solid (500 mg) was treated with a 4M HCl solution in dioxane (30 mL)


and the mixture stirred at RT for 2 h. The suspension was diluted with MeOH, then applied onto an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* as a pale yellow solid (100 mg, 23% over 2 steps). LCMS (Method 4): Rt 0.26, m/z 222 [MH⁺].

c. 3-(1,4-Dimethyl-piperazin-2-yl)-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 86c)



To a suspension of Intermediate 86b (100 mg) in 1,2-dichloroethane (10 mL) were added formaldehyde (30% in water, 104 μ L, 1.13 mmol) and NaBH(OAc)₃ (240 mg, 1.13 mmol) sequentially. The mixture was stirred at RT for 2 h, then applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave the *title compound* as a pale yellow solid (100 mg, 89%). LCMS (Method 4): Rt 0.49, m/z 250 [MH⁺].

d. (1S,4R)-4-[3-(1,4-Dimethyl-piperazin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 86d)



To a solution of Intermediate 86c (100 mg, 0.40 mmol) and Intermediate A (79 mg, 0.40 mmol) in DMF (3 mL) was added NaH (60% in mineral oil, 56 mg, 1.4 mmol) portion-wise. The mixture was stirred at 60°C for 1.5 h then allowed to cool to RT. The mixture was carefully quenched by pouring into MeOH (10 mL) then applied to an SCX-2 cartridge and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using


0-20% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a brown oil (104 mg, 66%). LCMS (Method 4): Rt 0.26, m/z 393 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-(1,4-dimethyl-piperazin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 86)


The *title compound* was prepared as an off white solid (70 mg, 42%) using (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 107 mg, 0.26 mmol) and Intermediate 86d (104 mg, 0.26 mmol) in a similar manner to Example 1, step d. LCMS (Method 5): Rt 3.64 min, m/z 648 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.83-1.98 (5H, m), 2.05-2.25 (6H, m), 2.30-2.42 (5H, m), 2.75 (2H, d, J 10.7), 2.89-2.94 (1H, m), 3.92-3.95 (1H, m), 4.80-4.86 (1H, m), 5.38-5.46 (1H, m), 6.32 (1H, s), 7.10 (1H, d, J 8.6), 7.26-7.42 (9H, m), 7.75 (1H, d, J 9.7), 8.04 (1H, s), 8.16 (1H, s), 8.46-8.50 (1H, m).

Example 87

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1,4,4-trimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (S)-1,4,4-Trimethyl-pyrrolidine-2-carboxylic acid (Intermediate 87a)



A suspension of 4,4-dimethyl-L-proline hydrochloride (900 mg, 5.00 mmol), formaldehyde (37% aqueous, 450 μL), DIPEA (645 mg, 5.00 mmol) and palladium on carbon (10%, 400 mg) in IMS (50 mL) was stirred under an


atmosphere of hydrogen (1 atm) for 4 h. The suspension was filtered and concentrated *in vacuo* to give the *title compound* as a viscous pink oil (780 mg, 99%). ¹H NMR (400 MHz, CDCl₃): 1.47 (3H, s), 1.49 (3H, s), 2.04-2.16 (1H, m), 2.23-2.34 (1H, m), 2.93 (3H, s), 3.10 (2H, m), 3.78-3.89 (1H, m), 4.86 (1H, br s).

5 **b. (S)-1,4,4-Trimethyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 87b)**



To a solution of Intermediate 87a (1.70 g, 5.00 mmol), (5-fluoro-pyridin-2-yl)-hydrazine (600 mg, 5.00 mmol) and DIPEA (1.29 g, 10.0 mmol) in DCM was
10 added HATU (2.28 g, 6.00 mmol) and the mixture stirred at RT for 1 h. The reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a yellow oil (950 mg, 71%). ¹H NMR (400 MHz, CDCl₃): 1.13 (3H,
15 s), 1.15 (3H, s), 1.74 (1H, dd, J 12.5, 6.2), 2.12 (1H, dd, J 11.6, 10.0), 2.29 (1H, d, J 8.9), 2.46 (3H, s), 2.88 (1H, d, J 8.9), 3.15 (1H, dd, J 9.7, 6.5), 6.56-6.65 (2H, m), 7.28 (1H dt, J 8.6, 2.9), 8.03 (1H, d, J 2.5), 9.09 (1H, br s).

c. 6-Fluoro-3-((S)-1,4,4-trimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 87c)




20

To a solution of Intermediate 87b (950 mg, 3.57 mmol), Ph₃P (1.87 g, 7.14 mmol) and Et₃N (1.44 g, 14.28 mmol) in THF (40 mL) was added hexachloroethane (1.65 g, 7.14 mmol) and the mixture stirred at RT for 20 h. The reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with
25 MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo*

gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM gave the *title compound* as a clear oil (750 mg, 84%). ¹H NMR (400 MHz, CDCl₃): 1.22 (3H, s), 1.24 (3H, s), 1.84 (1H, dd, J 12.7, 10.8), 2.04 (1H, dd, J 12.4, 7.4), 2.16 (3H, s), 2.25 (1H, d, J 9.1), 3.04 (1H, d, J 9.3), 4.22 (1H, dd, J 10.0, 7.6), 7.14-7.23 (1H, m), 7.69-7.77 (1H, m), 8.75-8.80 (1H, m).

d. (1S,4R)-4-[3-((S)-1,4,4-Trimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 87d)



To a solution of Intermediate A (552 mg, 3.89 mmol) in DMF (25 mL) was added NaH (60% in oil, 338 mg, 8.46 mmol) and the mixture stirred at RT for 20 min, before Intermediate 87c (700 mg, 2.82 mmol) was added. This mixture was stirred at 60°C for 1 h. The cooled reaction mixture was applied to an SCX-2 cartridge (70 g) and washed with MeOH. The product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-10% [2M NH₃ in MeOH] in DCM gave the *title compound* (820 mg, 74%). LCMS (Method 4): Rt 1.53 min, m/z 392 [MH⁺].


e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-1,4,4-trimethyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 87)

A solution of Intermediate 87d (150 mg, 0.383 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 232 mg, 0.575 mmol) and DIPEA (198 mg, 1.53 mmol) in DMF (4 mL) was stirred at 50°C for 1 h. The cooled reaction mixture was applied to an SCX-2 cartridge (10 g) and washed with MeOH. The


product was eluted with 2M NH₃ in MeOH; concentration *in vacuo* gave a residue. FCC, using 0-6% [2M NH₃ in MeOH] in DCM, then trituration with diethyl ether gave the *title compound* as an off-white powder (170 mg, 68%). LCMS (Method 5): Rt 3.80 min, m/z 647.5 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOH): 1.15 (3H, s), 1.24 (3H, s), 1.30 (9H, s), 1.85-2.05 (4H, m), 2.14 (3H, s), 2.19-2.31 (2H, m), 2.38 (3H, s), 2.97 (1H, d, J 10.3), 3.45 (1H, dd, J 7.5, 6.1), 4.06 (1H, t, J 9.3), 4.90 (1H, m), 5.30 (1H, t, J 4.2), 6.33 (1H, s), 7.20-7.36 (9H, m), 7.65 (1H, d, J 10.8), 8.46 (1H, d, J 1.1).

Example 88

10 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-1-methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**




a. **(S)-3-[N'-(5-Fluoro-pyridin-2-yl)-hydrazinocarbonyl]-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 88a)**



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (1.27 g, 10.0 mmol), (S)-1-BOC piperidine-3-carboxylic acid (Alfa Aesar, 2.29 g, 10.0 mmol) and HOBt.H₂O (153 mg, 1.00 mmol) in DCM (50 mL) was added EDC (2.30 g, 12.0 mmol) and the resulting orange solution stirred at RT for 16 h. Water (25 mL) was added and the mixture shaken. The aqueous was extracted with DCM (25 mL) then the combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a pale brown foam. FCC, using 0-5% MeOH in DCM, gave the *title compound* as a pale orange foam (3.21 g, 95%). LCMS (Method 3): Rt 3.14 min,


m/z 361 [MNa⁺].

b. (S)-3-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester (Intermediate 88b)




5 To a solution of Intermediate 88a (3.21 g, 9.49 mmol), Ph₃P (4.98 g, 19.0 mmol) and Et₃N (5.29 mL, 37.9 mmol) in THF (100 mL) at 0°C was added hexachloroethane (4.49 g, 19.0 mmol) and the resulting opaque orange solution stirred vigorously at RT for 2 h. The suspension was filtered, and the filter-cake washed with THF (20 mL). The combined organics were concentrated *in vacuo*,
10 redissolved in MeOH (5 mL), applied to an SCX-2 cartridge (70 g) and washed with MeOH (150 mL). The product was eluted with 2M NH₃ in MeOH (150 mL); concentration *in vacuo* gave a pale orange solid (2.62 g). FCC, using 2-5% MeOH in DCM, gave the *title compound* as a pale yellow solid (2.01 g, 66%). LCMS (Method 3): Rt 3.09 min, m/z 221 [M-C₅H₉O₂+H⁺].

15 **c. 6-Fluoro-3-(S)-piperidin-3-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 88c)**




A yellow solution of Intermediate 88b (2.01 g, 6.27 mmol) and TFA (11.0 mL, 148 mmol) in DCM (50 mL) was stirred at reflux for 30 min. The cooled
20 solution was concentrated *in vacuo*, redissolved in MeOH (3 mL), applied to an SCX-2 cartridge (50 g) and washed with MeOH (150 mL). The product was eluted with 2M NH₃ in MeOH (100 mL); concentration *in vacuo* left the *title compound* as a pale orange solid (1.26 g, 91%). LCMS (Method 3): Rt 0.35 min, m/z 221 [MH⁺].

25 **d. 6-Fluoro-3-((S)-1-methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-**

a]pyridine (Intermediate 88d)

To a yellow solution of Intermediate 88c (220 mg, 1.00 mmol), formaldehyde (37% in water, 0.811 mL, 10.0 mmol) and AcOH (0.057 mL, 1.00 mmol) in DCM-MeOH (5:1, 12 mL) was added NaBH(OAc)₃ (424 mg, 2.00 mmol) (CARE: gas evolution) and the solution stirred at RT for 2 h. The solution was concentrated *in vacuo* to ~5 mL volume, then applied to an SCX-2 cartridge (10 g) and washed with MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (50 mL); concentration *in vacuo* gave the *title compound* as a pale yellow gum, that became a white solid on standing (234 mg, quant.). LCMS (Method 3): Rt 0.45 min, m/z 235 [MH⁺].

e. (1S,4R)-4-[3-((S)-1-Methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 88e)




To a solution of Intermediate A (196 mg, 1.20 mmol) in dry DMF (3 mL) at RT under N₂ was added NaH (60% dispersion in oil, 80 mg, 2.00 mmol) and the resulting opaque brown solution stirred for 45 min. A solution of Intermediate 88d (234 mg, 1.00 mmol) in dry DMF (2 mL) was added and the resulting dark brown solution stirred at 60°C for 90 min. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL) and AcOH (0.1 mL), applied to and SCX-2 cartridge and washed with MeOH (75 mL). The product was eluted with 2M NH₃ in MeOH (60 mL); concentration *in vacuo* left a dark brown solid. FCC, using 3-13% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a light brown solid after

freeze-drying (280 mg, 74%). LCMS (Method 3): Rt 0.44 min, m/z 378 [MH⁺].

f. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((S)-1-methyl-piperidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea. (Example 88)

5 An orange-brown solution (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 243 mg, 0.600 mmol), Intermediate 88e (151 mg, 0.400 mmol) and DIPEA (0.122 mL, 0.700 mmol) in dry dioxane (5 mL) was stirred at 65°C for 15 h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL),
10 applied to an SCX-2 cartridge (10 g) and washed with MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (30 mL); concentration *in vacuo* left a brown solid. FCC, using 2-7% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a pale yellow solid (188 mg, 74%). LCMS (Method 5): Rt 3.63 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.55 (1H, qd, J 11.9, 5.3), 1.68-1.79 (2H, m), 1.83-2.03 (4H, m), 2.06-2.14 (2H, m), 2.22 (1H, t, J 10.9), 2.22 (3H, s), 2.36 (3H, s), 2.83 (1H, br d, J 10.9), 3.01 (1H, ddd, J 11.1, 3.6, 1.6), 3.54 (1H, tt, J 11.1, 3.7), 4.83 (1H, td, J 8.4, 5.5), 5.57 (1H, t, J 4.5), 6.33 (1H, s), 7.09 (1H, d, J 8.5), 7.15 (1H, dd, J 9.8, 2.0), 7.27-7.42 (8H, m), 7.68 (1H, d, J 9.9), 8.03 (1H, s), 8.32 (1H, d, J 2.0).

20 **Example 89**




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(R)-1-methyl-piperidin-3-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea


The *title compound* was prepared starting from (R)-1-BOC-piperidine-3-carboxylic acid (Alfa Aesar) using analogous procedures to those described in Example 88. LCMS (Method 5): Rt 3.62 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.55 (1H, qd, J 12.0, 5.5), 1.67-1.79 (2H, m), 1.82-2.16 (6H, m), 2.22 (3H, s), 2.23 (1H, t, J 11.0), 2.36 (3H, s), 2.83 (1H, br d, J 11.0), 2.99 (1H, br d, J 11.2), 3.53 (1H, tt, J 11.1, 3.7), 4.83 (1H, td, J 8.4, 5.5), 5.57 (1H, t, J 4.5), 6.33 (1H, s), 7.09 (1H, d, J 8.5), 7.16 (1H, dd, J 9.8, 2.1), 7.35-7.33 (8H, m), 7.68 (1H, d, J 9.9), 8.04 (1H, s), 8.29 (1H, d, J 2.0).

Example 90

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(R)-1-methyl-pyrrolidin-3-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. (R)-3-[N'-(5-Fluoro-pyridin-2-yl)-hydrazinocarbonyl]-pyrrolidine-1-carboxylic acid tert-butyl ester (Intermediate 90a)



To a brown solution of (5-fluoro-pyridin-2-yl)-hydrazine (590 mg, 4.64 mmol), (R)-1-N-BOC-beta proline (Manchester Organics, 1.00 g, 4.64 mmol) and HOBt.H₂O (71.1 mg, 0.464 mmol) in DCM (20 mL) at RT was added EDC (1.07 g, 5.57 mmol) (CARE: exotherm to ~35°C) and the resulting solution stirred at RT for 4 h. Water (20 mL) was added and the mixture shaken. The aqueous was

extracted with DCM (20 mL), then the combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to ~1 mL volume. FCC, using 1-10% MeOH in DCM, gave the *title compound* as a pale brown foam (1.37 g, 91%). LCMS (Method 3): Rt 2.93 min, m/z 347 [MNa⁺].


5 **b. (R)-3-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester (Intermediate 90b)**



To a brown solution Intermediate 90a (1.37 g, 4.22 mmol), Ph₃P (2.22 g, 8.45 mmol) and Et₃N (2.35 mL, 16.9 mmol) in THF (25 mL) at 0°C was added
10 hexachloroethane (2.00 g, 8.45 mmol) in 3 portions at 1 min intervals. The solution was stirred at 0°C for 30 min and at RT for 3.5 h. The suspension was filtered and the filter-cake washed with THF (10 mL). The combined organics were applied to an SCX-2 cartridge (50 g), then washed with DCM-MeOH (1:1, 50 mL) and MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (75 mL);
15 concentration *in vacuo* gave a brown oil that solidified on standing.


The solid was dissolved in DCM (10 mL) then Et₃N (0.580 mL, 4.16 mmol) and BOC anhydride (454 mg, 2.08 mmol) added sequentially. The solution was stirred at RT for 45 min, then water (10 mL) was added and the aqueous extracted with DCM (5 mL). The combined organics were passed through a hydrophobic frit
20 and concentrated *in vacuo* to leave a brown gum. FCC, using 2.5-3% MeOH in DCM, gave the *title compound* as a pale brown solid (933 mg, 73%). LCMS (Method 3): Rt 2.94 min, m/z 207 [MH⁺-C₅H₉O₂].

c. 6-Fluoro-3-(R)-pyrrolidin-3-yl-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 90c)




A mixture of Intermediate 90b (335 mg, 1.09 mmol) and TFA (0.812 mL, 10.9 mmol) in DCM (5 mL) was stirred at reflux for 45 min. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL), applied to an SCX-2 cartridge (10 g) and washed with MeOH (50 mL). The product was eluted with 2M
5 NH₃ in MeOH; concentration *in vacuo* left the *title compound* as a pale yellow solid (225 mg, >99%). LCMS (Method 3): Rt 0.44 min, m/z 207 [MH⁺].

d. 6-Fluoro-3-((R)-1-methyl-pyrrolidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 90d)



10 To a solution of Intermediate 90c (225 mg, 1.09 mmol), formaldehyde (37% in water, 0.885 mL, 10.9 mmol) and AcOH (0.0625 mL, 1.09 mmol) in DCM (20 mL) and MeOH (1 mL) was added NaBH(OAc)₃ (462 mg, 2.18 mmol) (CARE: gas evolution) and the pale yellow solution stirred at RT for 16 h. The solution was concentrated to ~2 mL volume, then was applied to an SCX-2 cartridge and was
15 washed with MeOH (25 mL). The product was eluted with 2M NH₃ in MeOH (25 mL); concentration *in vacuo* left the *title compound* as a pale brown viscous oil (228 mg, 95%). LCMS (Method 3): Rt 0.44 min, m/z 221 [MH⁺].

e. (1S,4R)-4-[3-((R)-1-Methyl-pyrrolidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 90e)



To a solution of Intermediate A (203 mg, 1.24 mmol) in dry DMF (3 mL) under N₂ was added NaH (60% dispersion in oil, 83.0 mg, 2.07 mmol) (CARE: gas evolution) and the resulting opaque brown solution stirred at RT for 45 min. A


solution of Intermediate 90d (228 mg, 1.03 mmol) in dry DMF (2 mL) was added and the dark brown solution stirred at 60°C under N₂ for 90 min. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL) and AcOH (0.1 mL), applied to and SCX-2 cartridge (20 g) and washed with MeOH (75 mL). The product was eluted with 2M NH₃ in MeOH (50 mL); concentration *in vacuo* gave a dark brown solid. FCC, using 5-15% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a light brown powder after freeze-drying (299 mg, 80%). LCMS (Method 3): Rt 0.44 min, m/z 364 [MH⁺].

f. **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((R)-1-methyl-pyrrolidin-3-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea. (Example 90)**

A brown solution of (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 227 mg, 0.560 mmol), Intermediate 90e (145 mg, 0.400 mmol) and DIPEA (0.111 mL, 0.640 mmol) in dry dioxane (5 mL) was stirred at 65°C for 15 h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL), applied to an SCX-2 cartridge (10 g) and washed with MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (30 mL); concentration *in vacuo* gave a brown solid. FCC, using 2-7% [2M NH₃ in MeOH] in DCM, gave a pale brown solid (195 mg). Further purification by MDAP (Method 8) gave the *title compound* as a white powder after freeze-drying (10.0 mg, 4%). LCMS (Method 5): Rt 3.62 min, m/z 619 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.83-1.98 (2H, m), 2.08-2.20 (3H, m), 2.32 (3H, s), 2.33-2.40 (1H, m), 2.36 (3H, s), 2.59-2.70 (2H, m), 2.75 (1H, dd, J 9.2, 6.1), 2.96 (1H, t, J 8.7), 3.93-4.01 (1H, m), 4.83 (1H, td, J 8.4, 5.4), 5.46 (1H, t, J 4.6), 6.32 (1H, s), 7.10 (1H, d, J 8.5), 7.18 (1H, dd, J 9.9, 2.1), 7.27-7.40 (8H, m), 7.69 (1H, dd, J 9.9, 0.8), 8.04 (1H, s), 8.44 (1H, d, J 2.0).

Example 91


1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[3-((2S,4R)-4-fluoro-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



5

a. (2S,4R)-4-Fluoro-1-methyl-pyrrolidine-2-carboxylic acid acid


(Intermediate 91a)



A solution of (2S,4R)-4-fluoropyrrolidine-2-carboxylic acid (1.00 g, 7.52 mmol), formaldehyde (37%wt in water, 1.00 mL, 30.8 mmol), 12N HCl (0.5 mL) in IMS (38 mL) was purged with argon. To the solution was added Pd/C (100 mg), then the flask was evacuated and filled with H₂ the resulting suspension was stirred at RT for 3 d. The reaction mixture was passed through a pad of Celite and the mixture was concentrated *in vacuo* to give the *title compound* as a pale yellow gum (600 mg, 54%). ¹H NMR (300 MHz, d₆-DMSO): 2.29-2.37 (1H, m), 2.63-2.67 (1H, m), 2.97 (3H, s), 3.44-3.53 (1H, m), 4.00-4.04 (1H, m), 4.58 (1H, dd, J 11.6, 7.11), 5.41-5.45 (1H, m).

15

b. (2S,4R)-4-Fluoro-1-methyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 91b)




20

A solution of (5-fluoro-pyridin-2-yl)-hydrazine (378 mg, 2.98 mmol),

Intermediate 91a (600 mg, 4.08 mmol) and HOBt.H₂O (44 mg, 0.12 mmol) in DCM (10 mL) and DMF (3 mL) was added EDC (628 mg, 3.28 mmol) portionwise at RT and the mixture stirred at RT for 17 h. The solution was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-7.5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as light orange powder (366 mg, 35%). LCMS (Method 3): Rt 0.56 min, m/z 257 [MH⁺].

c. 6-Fluoro-3-((2S,4R)-4-fluoro-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 91c)




10

A solution of Intermediate 91b (189 mg, 0.79 mmol), Ph₃P (416 mg, 1.59 mmol) and Et₃N (0.44 mL, 3.17 mmol) in THF (10 mL) at RT was added hexachloroethane (375 mg, 1.59 mmol) portion wise and the mixture stirred for 4 h. The solution was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* as light yellow powder (214 mg, 63%). LCMS (Method 3): Rt 0.42 min, m/z 239 [MH⁺].

15

d. (1S,4R)-4-[3-((2S,4R)-4-Fluoro-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine. (Intermediate 91d)

20



To a solution of Intermediate A (161 mg, 0.99 mmol) in dry DMF (2.5 mL) at RT was added NaH (60% in mineral oil, 108 mg, 2.70 mmol) and the mixture stirred for 15 min. Intermediate 91c (214 mg, 0.89 mmol) in DMF (2.5 mL) was


then added and the mixture heated at 60°C for 1 h. After cooling, the resulting dark brown mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give crude product. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as light brown foam (175 mg, 51%). LCMS (Method 3): Rt 0.43 min, m/z 382 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(2S,4R)-4-fluoro-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 91)

10 A solution of Intermediate 91d (164 mg, 0.43 mmol) and 5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 191 mg, 0.47 mmol) and DIPEA (225 µL, 1.14 mmol) in THF (5 mL) was stirred at reflux for 18.5 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics
15 were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give impure product. This residue was purified further by HPLC (XBridge C18 column, 30-98% MeCN in H₂O, 0.1% NH₄OH) to give the *title compound* as a white powder after freeze-drying (63 mg, 23%). LCMS (Method 5): Rt 3.88 min, m/z 637 [MH⁺]. ¹H NMR (400 MHz, d₆-
20 DMSO): 1.27 (9H, s), 1.88-1.93 (2H, m), 2.06-2.10 (2H, m), 2.17 (3H, s), 2.30-2.41 (2H, m), 2.36 (3H, s), 2.61-2.64 (2H, m), 3.49 (1H, ddd, J 25.6, 11.5, 5.5), 4.54 (1H, t, J 7.7), 4.80-4.83 (1H, m), 5.49 (1H, t, J 4.3), 6.32 (1H, s), 7.10 (1H, d, J 8.5), 7.30-7.42 (9H, m), 7.75 (1H, dd, J 9.9, 0.8), 8.05 (1H, s), 8.35 (1H, d, J 2.1).


Example 92

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-3-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




5

a. [(S)-1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-pyrrolidin-3-yl]-methanol (Intermediate 92a)



A solution of Intermediate 24b (485 mg, 2.83 mmol) and (S)-3-(hydroxymethyl)pyrrolidine (1.00 g, 9.90 mmol) in DMA (10 mL) was heated in the microwave at 175°C for 2.5 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a brown gum (670 mg, quant.). LCMS (Method 3): Rt 0.44 min, m/z 237 [MH⁺].

b. 6-Fluoro-3-((S)-3-triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 92b)




To a solution of Intermediate 92a (667 mg, 2.83 mmol) and Et₃N (630 μL, 4.53 mmol) in DCM (5 mL) was added triisopropylsilyltrifluoromethane sulfonate (950 μL, 3.54 mmol) and the mixture was stirred at RT for 1 h. The mixture was washed with sat. sodium hydrogen carbonate solution then brine, dried and

20


concentrated *in vacuo*. The residue was purified by FCC, 2-6% MeOH in DCM, to give the *title compound* as a pale green viscous oil (1.11 g, quant.). LCMS (Method 3): Rt 4.76 min, m/z 393 [MH⁺].

c. (1S,4R)-4-[3-((S)-3-Triisopropylsilanyloxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 92c)



To a solution of Intermediate A (179 mg, 1.10 mmol) in DMF (3 mL) was added NaH (60% dispersion in oil, 120 mg, 3.00 mmol) and the mixture stirred at RT for 25 min. A solution of Intermediate 92b (958 mg, 1.10 mmol) in DMF (3 mL) was added and the mixture stirred at 60°C for 1.5 h. The solution was cooled, water added, and the mixture extracted with DCM (4 × 25 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-16% MeOH in DCM, to give the *title compound* as a brown gum (216 mg, 37%). LCMS (Method 3): Rt 3.08 min, m/z 536 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-3-triisopropylsilanyloxy-methyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 92d)




A solution of Intermediate 92c (210 mg, 0.39 mmol) and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 159 mg, 0.390 mmol) in 1,4-dioxane (4

mL) and DIPEA (105 μ L, 0.600 mmol) was stirred at 65°C for 15.5 h and at 90°C for 1.5 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* as a pale brown solid (242 mg, 79%). LCMS (Method 3): Rt 5.18 min, m/z 791 [MH⁺].

5 **e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-3-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 92)**

To a solution of Intermediate 92d (240 mg, 0.300 mmol) and TBAF (1M in THF, 0.400 mL, 0.400 mmol) in THF (3 mL) was stirred at RT for 1 h. Water was
10 added and the mixture extracted with DCM (4 $\times$ 20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-14% MeOH in DCM, to give the *title compound* as an off-white powder after freeze-drying (154 mg, 81%). LCMS (Method 5): Rt 4.01 min, m/z 635.3 [MH⁺].
¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.68-1.79 (1H, m), 1.79-1.97 (2H, m), 1.98-2.14 (3H, m), 2.36 (3H, s), 2.40-2.48 (1H, m), 3.23-3.30 (1H, m), 3.41-
15 3.64 (5H, m), 4.71 (1H, t, J 5.3), 4.77-4.85 (1H, m), 5.51 (1H, br t, J 4.4), 6.33 (1H, s), 7.03-7.10 (2H, m), 7.25-7.40 (8H, m), 7.53 (1H, d, J 9.9), 7.82 (1H, d, J 1.8), 8.03 (1H, s).

Example 93



20


1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((R)-3-hydroxymethyl-pyrrolidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

The *title compound* was prepared starting from (R)-3-
25 (hydroxymethyl)pyrrolidine using analogous procedures to those described in

Example 92. LCMS (Method 5): Rt 4.01 min, m/z 635.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.67-1.77 (1H, m), 1.79-1.97 (2H, m), 1.98-2.15 (3H, m), 2.37 (3H, s), 2.40-2.48 (1H, m), 3.25-3.30 (1H, m), 3.41-3.62 (5H, m), 4.71 (1H, t, J 5.3), 4.77-4.85 (1H, m), 5.51 (1H, br t, J 4.3), 6.33 (1H, s), 7.04-7.10 (2H, m), 7.25-7.40 (8H, m), 7.53 (1H, d, J 9.9), 7.82 (1H, d, J 1.6), 8.04 (1H, s).


Example 94

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea



10


a. 2-[4-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-piperazin-1-yl]-ethanol (Intermediate 94a)



A solution of Intermediate 24b (485 mg, 2.82 mmol) and 1-(2-hydroxyethyl)-piperazine (1.39 mL, 11.3 mmol) in DMA (10 mL) was heated in the microwave at 170°C for 8 h. The cooled mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% [2M NH₃ in MeOH] in DCM, to give impure product. Further purified by FCC, using 0-7% [2M NH₃ in MeOH] in DCM gave the *title compound* as a pale brown gum (175 mg, 23%). LCMS (Method 3): Rt 0.43 min, m/z 266 [MH⁺].


20

b. 6-Fluoro-3-[4-(2-triisopropylsilyloxy-ethyl)-piperazin-1-yl]-

[1,2,4]triazolo[4,3-a]pyridine (Intermediate 94b)


To a solution of Intermediate 94a (169 mg, 0.64 mmol) and Et₃N (265 μ L, 1.91 mmol) in DCM (6 mL) was added triisopropylsilyl trifluoromethanesulfonate (189 μ L, 0.70 mmol). The solution was stirred for 2.5 h, then diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-5% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale brown gum (242 mg, 90%). LCMS (Method 3): Rt 2.93 min, m/z 422 [MH⁺].

10 **c. (1S,4R)-4-{3-[4-(2-Triisopropylsilyloxy-ethyl)-piperazin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 94c)**



To a solution of Intermediate A (102 mg, 0.63 mmol) in DMF (1.5 mL) was added NaH (60% dispersion in oil, 68 mg, 1.71 mmol). The mixture was stirred at RT for 5 min, then a solution of Intermediate 94b (240 mg, 0.57 mmol) in DMF (1.5 mL) was added and the mixture stirred at 60°C for 1.25 h. The cooled mixture was diluted with water and extracted with DCM (4 $\times$ 25 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* as a pale brown gum (219 mg, 68%). LCMS (Method 3): Rt 2.65 min, m/z 565 [MH⁺].

d. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(2-triisopropylsilyloxy-ethyl)-piperazin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Intermediate 94d)



5 A solution of Intermediate 94c (170 mg, 0.42 mmol), 5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Synthetic Communications, 2009, 39, 3999-4009; 170 mg, 0.42 mmol) and DIPEA (199 μ L, 1.14 mmol) in THF (4 mL) was stirred at reflux for 15.5 h. The cooled reaction mixture was diluted with water and extracted with DCM. The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-7.5% [2M NH_3 in MeOH] in DCM, to give the *title compound* as a yellow powder
10 (293 mg, 93%). LCMS (Method 3): Rt 3.70 min, m/z 820 [MH^+].


e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy}-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea (Example 94)
15

A solution of Intermediate 94d (243 mg, 0.29 mmol) and TBAF (1M in THF, 0.53 mL, 0.53 mmol) in THF (3.5 mL) was stirred at RT for 45 min, then diluted with water and extracted with DCM (3×20 mL). The combined organics were dried and concentrated *in vacuo*. The residue was purified by FCC, using 0-7.5% [2M NH_3 in MeOH] in DCM, to give the *title compound* as a white powder
20 after freeze-drying (135 mg, 58%). LCMS (Method 5): Rt 3.52 min, m/z 664 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 1.27 (9H, s), 1.86-1.92 (2H, m), 2.04-2.11 (2H, m), 2.36 (3H, s), 2.48 (2H, q, J 5.8), 2.66 (4H, t, J 4.8), 3.19 (4H, t, J 4.5),


3.55 (2H, q, J 5.8), 4.44 (1H, t, J 5.4), 4.79-4.82 (1H, m), 5.57 (1H, t, J 4.4), 6.32 (1H, s), 7.07 (1H, d, J 8.5), 7.15 (1H, dd, J 9.9, 2.1), 7.32-7.44 (8H, m), 7.60-7.63 (1H, m), 7.68 (1H, d, J 2.0), 8.04 (1H, s).

Example 95

5 **1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea**



a. **3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenol (Intermediate 95a)**




10

A mixture of 5-tert-butyl-2H-pyrazol-3-ylamine (1 g, 7.18 mmol), 3-iodophenol (1.74 g, 7.90 mmol), copper (I) iodide (68 mg, 0.36 mmol), (1S,2S)-N,N'-dimethyl cyclohexane-1,2-diamine (204 mg, 1.44 mmol) and potassium carbonate (2.08 g, 15.09 mmol) in toluene (8 mL) was de-gassed and flushed with argon (3x). The reaction mixture was then treated with microwave irradiation, at 150°C for 1 h. The mixture was diluted with EtOAc (20 mL) and washed with water (20 mL). The aqueous layer was extracted with a further 20 mL EtOAc and the combined organic layers were washed with brine (20 mL), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue obtained was purified by FCC, using 0-100% EtOAc in cyclohexane to afford the *title compound* as a brown gum (1.42 g, 86%). LCMS (Method 3): Rt 2.21 min, m/z 232 [MH⁺].

15


20

b. 5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-ylamine (Intermediate 95b)



A solution of intermediate 95a (1.42 g, 6.15 mmol) and triphenylphosphine (3.22 g, 12.29 mmol) in THF (50 mL), under an atmosphere of argon was treated with 2-(tetrahydro-pyran-2-yloxy)-ethanol (1.25 mL, 9.22 mmol), followed by the dropwise addition of diethyl azodicarboxylate (1.94 mL, 12.29 mmol). The reaction mixture was then stirred at RT for 1 h. Water (0.5 mL) was added and the mixture was concentrated *in vacuo*. The residue was taken up in EtOAc (8 mL) and cyclohexane was added until triphenylphosphine oxide had precipitated out of solution. This was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane to afford the *title compound* as an orange/brown oil (2.98 g, >100%). LCMS (Method 3): Rt 3.20 min, m/z 360 [MH⁺].


c. (5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 95c)



A solution of intermediate 95b (2.21 g, 6.16 mmol) in EtOAc (20 mL) was treated with aqueous NaOH (1M, 11.08 mmol), followed by 2,2,2-trichloroethyl chloroformate (1.01 mL, 7.39 mmol) and the reaction mixture was stirred at RT for 1 h. The mixture was partitioned between EtOAc (50 mL) and

water (50 mL). The layers were separated and the aqueous layer was extracted with a further 50 mL EtOAc. The combined organic layers were dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane to afford the *title compound* as a dark orange oil
5 (2.39 g, 73%). LCMS (Method 3): Rt 4.70 min, m/z 534/536 [MH⁺].

d. 1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 95d)




10

A mixture of intermediate 95c (200 mg, 0.37 mmol), intermediate 81d (141 mg, 0.37 mmol) and DIPEA (98 μ L, 0.56 mmol) in dioxane (1.5 mL) was heated at 70°C for 20 h. The reaction mixture was cooled to RT, diluted with DCM (5 mL) and washed with water (2 x 5 mL). The organic layer was passed
15 through a hydrophobic frit and concentrated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM to afford the *title compound* as a yellow glass (158 mg, 55%). LCMS (Method 3): Rt 4.25 min, m/z 763 [MH⁺].

20

e. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 95e)



5 A mixture of intermediate 95d (158 mg, 0.21 mmol) and pyridinium
 p-toluene sulfonate (156 mg, 0.62 mmol) in methanol was heated at 45°C for 18 h.
 The mixture was cooled to RT and was partitioned between DCM (5 mL) and
 saturated aqueous NaHCO₃ (5 mL). The layers were separated and the organic
 layer was passed through a hydrophobic frit and concentrated *in vacuo*. The
 10 residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM to afford
 the *title compound* as a beige solid (105 mg, 74%). LCMS (Method 3): Rt 3.64
 min, m/z 679 [MH⁺].


f. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-
 yl-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-
 15 6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 95)

To a solution of intermediate 95e (105 mg, 0.15 mmol) and DIPEA (81 µL,
 0.46 mmol) in DCM (2 mL) was added methanesulfonyl chloride (16 µL, 0.20
 mmol). The reaction mixture was stirred at RT for 30 min. The mixture was
 diluted with DCM (5 mL) and washed with water (2 x 5 mL). The organic layer
 20 was passed through a hydrophobic frit and concentrated *in vacuo*. The residue was
 taken up in THF (2 mL) and dimethylamine solution in THF (2M, 1.55 mL) was
 added. The reaction mixture was heated at 60°C in a sealed tube for 20 h. The
 reaction mixture was cooled to RT and concentrated *in vacuo*. The residue was
 triturated with Et₂O (1 mL) and the solid obtained was purified by MDAP to afford


the *title compound* as a glassy solid (44 mg, 40%). LCMS (Method 5): Rt 3.64 min, m/z 706.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J 6.7 Hz), 1.28 (9H, s), 1.46-1.54 (2H, m), 1.63-1.72 (2H, m), 1.75-1.76 (4H, m), 2.00-2.18 (2H, m), 2.22 (6H, s), 2.66 (2H, t, J 5.8 Hz), 2.85-2.95 (1H, m), 3.12-3.20 (1H, m, obscured by water), 3.27-3.35 (1H, m, obscured by water), 4.10 (2H, t, J 5.4 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J 4.0 Hz), 6.33 (1H, s), 6.95-6.99 (1H, m), 7.06-7.14 (3H, m), 7.19 (1H, dd, J 10.0 Hz), 7.25-7.43 (5H, m), 7.64 (1H, d, J 10.6 Hz), 7.69 (1H, d, J 1.8 Hz), 8.12 (1H, s).

Example 96

10 **1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-(*cis*-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**




15 **a. *cis*-2,6-Dimethyl-piperidine-1-carboxylic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 96a)**



Pyridine (2.40 mL, 29.7 mmol) was added dropwise to an ice cold suspension of triphosgene (4.42 g, 14.9 mmol) in DCM (30.0 mL). *cis*-2,6-Dimethylpiperidine (2.00 mL, 14.9 mmol) was added and the reaction stirred for 3 hours, then quenched by dropwise addition of HCl (1M aqueous, 30 mL). The mixture was extracted into DCM (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The product was used in the next reaction without purification.

cis-2,6-Dimethylpiperidine-carbamoylchloride (1.79 g, 10.2 mmol) was dissolved in DCM (90.0 mL) and DIPEA (2.42 mL, 13.9 mmol) was added followed by (5-fluoro-pyridin-2-yl)-hydrazine (WO2010022076, 1.18 g, 9.30 mmol) and the reaction heated to 45°C overnight. The reaction was cooled and quenched into water. The mixture was extracted into DCM (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC using 0-6% [2M NH₃ in MeOH] in DCM to give the *title compound* (1.38 g, 56%). ¹H NMR (300 MHz, CDCl₃): 1.29 (6H, d, J 7.1 Hz), 1.48-1.90 (6H m), 4.26 (2H, qn, J 6.9 Hz), 6.39 (1H, d, J 2.3 Hz), 6.50 (1H, d, J 2.3 Hz), 6.74 (1H, dd, J 9.0, 3.6 Hz), 7.27 (1H, ddd, J 8.9, 8.0, 2.9 Hz), 8.03 (1H, d, J 2.9 Hz).


b. 3-(*cis*-2,6-Dimethyl-piperidin-1-yl)-6-fluoro-[1,2,4]triazolo[4,3-*a*]pyridine (Intermediate 96b)



To an ice cold solution of Intermediate 96a (1.38 g, 5.19 mmol) in THF (50.0 mL) was added sequentially triphenylphosphine (2.72 g, 10.4 mmol), Et₃N (2.89 mL, 20.8 mmol) and hexachloroethane (2.46 g, 10.4 mmol). The cooling bath was removed and the reaction was heated to 55°C overnight. The reaction was cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH. The residue was concentrated *in vacuo* then resubmitted to the reaction conditions overnight. After analogous workup the residue was concentrated *in vacuo* to give the *title compound* (1.03 g, 80%). ¹H NMR (300 MHz, CDCl₃): 0.68 (6H, d, J 6.2 Hz), 1.36-1.68 (3H m), 1.74-1.90 (3H,


m), 3.35 (2H, m), 7.14 (1H, ddd, J 9.8, 7.5, 2.3 Hz), 7.65 (1H, ddd, J 9.9, 4.7, 0.8 Hz), 8.02 (1H, ddd, J 3.2, 2.3, 0.8 Hz).

c. (1*S*,4*R*)-4-[3-(*cis*-2,6-Dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 96c)



To a suspension of sodium hydride (60% in mineral oil, 323 mg, 8.06 mmol) in DMF (10.0 mL) was added (1*R*,4*S*)-4-amino-1,2,3,4-tetrahydro-naphthalen-1-ol (329 mg, 2.02 mmol) and the reaction stirred for 20 min. Intermediate 96b (500 mg, 2.02 mmol) was added in DMF (2.00 mL) and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (453 mg, 57%). LCMS (Method 4): Rt 2.13, m/z 392 [MH⁺].


d. 1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1*S*,4*R*)-4-[3-(*cis*-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 96d)



To a solution of Intermediate 96c (100 mg, 0.25 mmol) in 1,4-dioxane (3.00

mL) was added DIPEA (89.0 μ L, 0.51 mmol) and Intermediate 39b (133 mg, 0.25 mmol). The reaction was heated to 70°C overnight then cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC using 0-8% MeOH in DCM to give the *title compound*. LCMS (Method 4): Rt 4.09 min, m/z 777 [MH⁺].

e. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-(*cis*-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 96e)




10

Pyridinium p-toluenesulfonate (65.0 mg, 0.26 mmol) was added to a solution of Intermediate 96d in MeOH (2.0 mL). The reaction was heated to 60°C for 3.5 h then cooled and solvent volume reduced *in vacuo*. The residue was partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give the *title compound* (89.0 mg, 99%). LCMS (Method 4): Rt 3.49 min, m/z 693 [MH⁺].

20

25

f. Methanesulfonic acid 2-{3-[3-tert-butyl-5-(3-{(1*S*,4*R*)-4-[3-((2*S*,6*R*)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenoxy}-ethyl ester (Intermediate 96f)



5

Methane sulfonyl chloride (11.0 μ L, 0.13 mmol) was added to an ice cold solution of Intermediate 96e (89.0 mg, 0.13 mmol) and DIPEA (27.0 μ L, 0.15 mmol) in DCM (1.5 mL). After 1 hour additional DIPEA (27.0 μ L, 0.15 mmol) and methane sulfonyl chloride (11.0 μ L, 0.13 mmol) was added. After a further 1 hour additional DIPEA (27.0 μ L, 0.15 mmol) and methane sulfonyl chloride (11.0 μ L, 0.13 mmol) was added. After 1 hour the reaction was partitioned between DCM and water. The aqueous layer was then extracted with DCM (3 $\times$). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give the *title compound* (100 mg, 99%). LCMS (Method 1): Rt 3.73 min, m/z 771 [MH⁺].

15

g. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1*S*,4*R*)-4-[3-(*cis*-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 96)


20

Dimethylamine (2M in MeOH, 520 μ L, 1.04 mmol) was added to a solution of Intermediate 96f (100 mg, 0.13 mmol) in THF (1.5 mL). The reaction was heated to 50°C in a sealed environment overnight. The mixture was cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 $\times$). The combined organic layers were washed with brine, dried


(MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM. Further purification by HPLC (C18 X-select column, 10-98% MeCN in H₂O, 0.1% HCO₂H) gave the *title compound* as a white powder after freeze-drying (50 mg, 54%). LCMS (Method 5): Rt 3.79 min, m/z 720.5 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 0.65 (3H, d, J 6.0 Hz), 0.68 (3H, d, J 6.0 Hz), 1.31 (9H, s), 1.40-1.50 (2H, m), 1.53-1.65 (1H, m), 1.73-1.87 (3H, m), 1.92-2.05 (2H, m), 2.05-2.15 (1H, m), 2.19-2.26 (1H, m), 2.43 (6H, s), 2.92 (2H, t, J 5.2 Hz), 3.24 (2H, m), 4.18 (2H, t, J 5.3 Hz), 4.90 (1H, dd, J 8.7, 5.8), 5.42 (1H, t, J 8.2), 6.34 (1H, s), 7.02-7.06 (1H, ddd, J 8.5, 2.4, 0.6), 7.08-7.13 (2H, m), 7.17-7.25 (3H, m), 7.26-7.31 (2H, m), 7.38-7.44 (1H, t, J 7.8), 7.56-7.60 (1H, dd, J 9.8, 0.7 Hz), 7.86 (1H, d, J 1.8 Hz), 8.45 (0.3H, br s).

Example 97

**1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**



**a. 1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 97a)**




20

A brown solution of Intermediate 29c (177 mg, 0.420 mmol), Intermediate 81d (151 mg, 0.400 mmol) and DIPEA (0.087 mL, 0.50 mmol) in dry dioxane (5

mL) was stirred at 100°C for 3 h. The cooled solution was concentrated *in vacuo*, suspended in water (10 mL) and extracted with DCM (2 × 10 mL). The combined organics were passed through a hydrophobic frit and concentrated under vacuum to leave a dark brown gum. FCC, using 3-8% MeOH in DCM, gave the *title compound* as an orange solid (188 mg, 72%). LCMS (Method 4): Rt 3.63 min, m/z 649 [MH⁺].

b. Methanesulfonic acid 3-[3-tert-butyl-5-(3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzyl ester and 1-[5-tert-Butyl-2-(3-chloromethyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (1:1 mixture). (Intermediate mixture 97b)



To a solution of Intermediate 97a (187 mg, 0.288 mmol) and DIPEA (0.151 mL, 0.865 mmol) in DCM (5 mL) at 0°C was added mesyl chloride (0.056 mL, 0.721 mmol) and the resulting orange solution stirred at 0°C for 1 h. Water (5 mL) was added and the mixture shaken. The aqueous layer was extracted with DCM (5 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the mixture of *title compounds* as a brown gum. LCMS (Method 4): 1:1 ratio, Rt 3.87 min, m/z 727 [MH⁺] and Rt 4.11, m/z 667 [MH⁺].

c. 1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 97)


A brown solution of Intermediate mixture 97b (0.144 mmol) and

morpholine (0.063 mL, 0.72 mmol) in dry DMF (3 mL) was stirred at 75°C for 2

h. The cooled solution was concentrated *in vacuo*, suspended in water (10 mL) and extracted with DCM (2 × 10 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. FCC, 3-7% MeOH in DCM, gave a pale yellow solid (58.6 mg). MDAP (Method 7) gave the *title compound* as a pale yellow solid (31 mg, 28%). LCMS (Method 5): Rt 3.59 min, m/z 718.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J 6.3), 1.28 (9H, s), 1.48-1.55 (2H, m), 1.63-1.71 (2H, m), 1.75-1.97 (4H, m), 2.00-2.16 (2H, m), 2.37 (4H, t, J 4.2), 2.90 (1H, ddd, J 12.2, 9.0, 4.0), 3.15 (1H, dt, J 12.0, 4.2), 3.51 (2H, s), 3.55 (4H, t, J 4.4), 4.82 (1H, td, J 8.6, 5.6), 5.51 (1H, t, J 4.3), 6.33 (1H, s), 7.04 (1H, d, J 8.6), 7.19 (1H, dd, J 9.9, 2.2), 7.25-7.29 (2H, m), 7.31-7.48 (6H, m), 7.64 (1H, dd, J 9.8, 0.8), 7.69 (1H, d, J 2.1), 8.10 (1H, s), 8.16 (1H, s). One 1H signal under solvent peak.

Example 98

1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




The *title compound* was prepared starting from N-methyl piperazine using analogous procedures to those described in Example 97. LCMS (Method 5): Rt 3.61 min, m/z 731.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J 6.3), 1.28 (9H, s), 1.48-1.56 (2H, m), 1.63-1.71 (2H, m), 1.75-1.96 (4H, m), 2.00-2.16 (2H, m), 2.12 (3H, s), 2.26-2.43 (8H, m), 2.90 (1H, ddd, J 11.8, 9.2, 4.2), 3.16 (1H, dt, J 12.0, 4.2), 3.28-3.34 (1H, m), 3.50 (2H, s), 4.82 (1H, td, J 8.6, 5.5), 5.52 (1H, t, J 4.3), 6.33 (1H, s), 7.04 (1H, d, J 8.6), 7.19 (1H, dd, J 9.9, 2.2), 7.25-7.47 (8H,


m), 7.64 (1H, dd, J 9.8, 0.8), 7.69 (1H, d, J 2.1), 8.11 (1H, s), 8.18 (1H, s).

Example 99

1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{{(1S,4R)-4-[3-(1-dimethylamino-cyclopentyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. 1-Dimethylamino-cyclopentanecarboxylic acid (Intermediate 99a)



A solution of 1-amino-cyclopentanecarboxylic acid (2.0 g, 15.5 mmol) and formaldehyde (37% aqueous solution, 3 mL) in formic acid (4 mL) was heated at 80 °C for 2.5 h. The volatiles were concentrated *in vacuo* and the resulting residue was loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH to afford the *title compound* (1.82 g, 75%).
¹H NMR (400 MHz, CD₃OD): 1.71-2.02 (6H, m), 2.16-2.29 (2H, m), 2.77 (6H, s).


b. 1-Dimethylamino-cyclopentanecarboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 99b)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (For reference procedure see WO2010022076; 200 mg, 1.57 mmol) in DMF (10 mL) were added Intermediate 99a (247 mg, 1.57 mmol), EDC (333 mg, 1.73 mmol) and HOBt (21 mg, 0.16 mmol). The reaction mixture was stirred at RT for 18 h and then partitioned between EtOAc and water. The aqueous phase was further extracted


with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-8% [2M NH₃ in MeOH] in DCM, to give the *title compound* (282 mg, 67%). ¹H NMR (400 MHz, CDCl₃): 1.54-1.73 (4H, m), 1.76-2.04 (4H, m), 2.25 (6H, s), 6.46 (1H, s), 6.64 (1H, dd, J = 9.0, 3.5 Hz), 7.23-7.33 (1H, m), 8.04 (1H, d, J = 3.0 Hz), 8.94 (1H, s).

c. [1-(6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-cyclopentyl]-dimethylamine (Intermediate 99c)



Hexachloroethane (503 mg, 2.12 mmol) was added portionwise over 5 minutes at RT to a stirred mixture of Intermediate 99b (282 mg, 1.06 mmol), triethylamine (590 μ L, 4.24 mmol) and triphenylphosphine (555 mg, 2.12 mmol) in THF (11 mL). The reaction mixture was stirred at 50 $^{\circ}$ C for 2.5 h and then partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC using SCX-2 cartridge. The cartridge was washed with MeOH and the product was eluted with 2M NH₃ in MeOH to give the *title compound* (207 mg, 79%). ¹H NMR (400 MHz, CDCl₃): 1.67-1.78 (4H, m), 2.10-2.22 (8H, m), 2.24-2.34 (2H, m), 7.15 (1H, ddd, J = 10.0, 7.4, 2.3 Hz), 7.69 (1H, dd, J = 10.0, 5.0 Hz), 8.65-8.68 (1H, m).

d. (1S,4R)-4-[3-(1-Dimethylamino-cyclopentyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 99d)




5 To a suspension of sodium hydride (60% in mineral oil, 133 mg, 3.34 mmol) in DMF (5 mL) was added Intermediate A (136 mg, 0.83 mmol) and the reaction mixture was stirred for 20 min. A solution of Intermediate 99c (207 mg, 0.83 mmol) in DMF (3 mL) was added and the reaction mixture was heated at 60 °C for 90 min. After cooling, the reaction mixture was quenched by careful
10 addition of MeOH and then loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (211 mg, 65%). LCMS (Method 4): Rt 1.49 min, m/z
15 392 [MH⁺].

e. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-(1-dimethylamino-cyclopentyl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 99)

A mixture of Intermediate 99d (108 mg, 0.27 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference
20 procedure see Synthetic Communications, 2009, 39, 3999-4009; 105 mg, 0.27 mmol) and DIPEA (93 µL, 0.54 mmol) in dioxane (3 mL) was stirred at 60 °C for 18 h and then at 80 °C for 4 h. The reaction mixture was diluted with EtOAc and then poured into water. The aqueous phase was extracted with EtOAc (x 3) and the
25 combined organic layers were washed with brine, dried (MgSO₄) and concentrated

in vacuo. The resultant residue was purified by MDAP (Method 7) to afford the *title compound* (23 mg, 13%). LCMS (Method 5): Rt 3.90 min, m/z 647 [MH⁺]. ¹H NMR (400 MHz, CD₃OD): 1.34 (9H, s), 1.59-1.85 (4H, m), 1.91-2.15 (3H, m), 2.18 (6H, s), 2.21-2.36 (5H, m), 2.42 (3H, s), 4.91-4.96 (1H, m), 5.34 (1H, t, J = 4.1 Hz), 6.37 (1H, s), 7.22-7.40 (9H, m), 7.65 (1H, d, J = 9.9 Hz), 8.46 (1H, d, J = 2.2 Hz).

Example 100




1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

The *title compound* was prepared starting from Intermediate 96c and (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference procedure see Synthetic Communications, 2009, 39, 3999-4009) using analogous procedures to those described in Example 3. LCMS (Method 5): Rt 5.33 min, m/z 647 [MH⁺]. ¹H NMR (400 MHz, CD₃OD): 0.64-0.74 (6H, m), 1.34 (9H, s), 1.41-1.72 (3H, m), 1.75-2.19 (6H, m), 2.21-2.31 (1H, m), 2.42 (3H, s), 3.22-3.35 (2H, m), 4.90-4.98 (1H, m), 5.45 (1H, t, J = 4.1 Hz), 6.37 (1H, s), 7.19-7.40 (9H, m), 7.62 (1H, dd, J = 9.9, 0.8 Hz), 7.89 (1H, d, J = 2.1 Hz).


Example 101

1-[(1S,4R)-4-(3-Amino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea



5


a. 2-(5-Fluoropyridin-2-yl)-N,N-di(prop-2-en-1-yl)hydrazinecarboxamide (Intermediate 101a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (For reference procedure see WO2010022076; 2.54 g, 20.0 mmol) in DCM (150 mL) and DIPEA (3.87 g, 30.0 mmol) was added N,N-di-2-propen-1-yl-carbamic chloride (For reference procedure see for example Tetrahedron **1996**, 52, 13739-13750; 4.13 g, 26.0 mmol) and the reaction mixture was stirred at RT for 18 h. The reaction mixture was diluted with MeOH (50 mL) and loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The compound containing fractions were concentrated *in vacuo* and the resulting residue was purified by FCC on silica, using a gradient of 0-6% [2M NH₃ in MeOH] in DCM, to afford the *title compound* (2.30 g, 46%) as a yellow solid. LCMS (Method 1): Rt 2.26 min, m/z 251 [MH⁺].


20

b. **Diallyl-(6-fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-amine**
(Intermediate 101b)



5 Hexachloroethane (4.34 g, 18.4 mmol) was added portionwise over 5 min at RT to a stirred mixture of Intermediate 101a (2.30 g, 9.2 mmol), triethylamine (3.71 g, 36.8 mmol) and triphenylphosphine (4.82 g, 18.4 mmol) in THF (90 mL). The reaction mixture was stirred at RT for 1 h, then diluted with MeOH and purified by FCC using SCX-2 cartridge. The cartridge was washed with MeOH and the product was eluted with 2M NH₃ in MeOH. The compound containing
10 fractions were combined and concentrated *in vacuo*. The resultant residue was subjected to the follow purification procedure; FCC on silica (0-6% [2M NH₃ in MeOH] in DCM), then SCX-2 cartridge (conditions described above) and finally further FCC on silica (0-2% MeOH in EtOAc), to afford the *title compound* (1.34
15 g, 62%). ¹H NMR (400 MHz, CDCl₃): 3.87 (4H, d, J = 6.2 Hz), 5.16-5.35 (4H, m), 5.85-6.02 (2H, m), 7.11 (1H, ddd, J = 10.0, 7.6, 2.3 Hz), 7.58-7.67 (1H, m), 7.75-7.80 (1H, m).

c. **[6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl]-bis-((E)-propenyl)-amine**
20 **(Intermediate 101c)**



To a suspension of sodium hydride (40% in mineral oil, 690 mg, 17.31 mmol) in DMF (20 mL) was added Intermediate A (1.12 g, 6.92 mmol) and the

reaction mixture was stirred for 20 min at RT. A solution of Intermediate 101b (1.34 g, 5.77 mmol) in DMF (20 mL) was added and the reaction mixture was stirred at RT for 1 h. The reaction mixture was quenched by careful addition of MeOH and then loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (1.33 g, 61%) as an orange gum. ¹H NMR (400 MHz, CDCl₃): 0.92-1.13 (1H, m), 1.39 (6H, dd, J = 7.2, 1.7 Hz), 1.82-2.16 (4H, m), 2.26-2.38 (1H, m), 3.93-3.99 (1H, m), 4.99-5.10 (2H, m), 5.18 (1H, t, J = 4.4 Hz), 6.10 (2H, dd, J = 8.2, 2.0 Hz), 7.09 (1H, dd, J = 9.9, 2.2 Hz), 7.25-7.44 (3H, m), 7.48-7.53 (1H, m), 7.57-7.66 (2H, m).


d. 1-[(1S,4R)-4-(3-Amino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-3-(5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-urea
(Example 101)

A mixture of Intermediate 101c (1.33 g, 3.54 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference procedure see Synthetic Communications, 2009, 39, 3999-4009; 2.0 g, 4.96 mmol) and DIPEA (1.82, 14.16 mmol) in DMF (20 mL) was stirred at RT for 20 h. The reaction mixture was diluted with MeOH and loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by recrystallization from 5% MeOH in DCM (few drops of cyclohexane added to start recrystallization), to give the *title compound* (760 mg, 39%) as an off white dust. LCMS (Method 5): Rt 3.81 min, m/z 551 [MH⁺]. ¹H NMR (400 MHz, CD₃OD): 1.34 (9H, s), 1.86-2.08 (2H, m), 2.09-2.19 (1H, m), 2.22-2.34 (1H, m), 2.42 (3H, s), 4.89-4.96 (1H, m), 5.32 (1H, t, J = 4.4


Hz), 6.37 (1H, s), 7.05 (1H, dd, $J = 10.0, 2.1$ Hz), 7.23-7.44 (9H, m), 7.83 (1H, d, $J = 2.0$ Hz).

Example 102

1-[5-tert-Butyl-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydronaphthalen-1-yl]-urea



a. [5-tert-Butyl-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 102a)



10

To a mixture of 5-tert-butyl-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-ylamine (For reference procedure see for example WO2003072569; 833 mg, 3.62 mmol) in NaOH (1M in water, 5.4 mL, 4.4 mmol) and EtOAc (5 mL) was added 2,2,2-trichloroethyl chloroformate (548 μ L, 3.98 mmol) and the reaction mixture was stirred at RT for 1 h. Additional NaOH (1M in water, 5.4 mL) and 2,2,2-trichloroethyl chloroformate (548 μ L) were added and stirring at RT was continued for 1 h. Additional NaOH (1M in water, 5.4 mL) and 2,2,2-trichloroethyl chloroformate (548 μ L) were added and stirring at RT was continued for 2 h. Additional NaOH (1M in water, 10.8 mL) and 2,2,2-trichloroethyl chloroformate (1.10 mL) were added and stirring at RT was continued for 24 h. The aqueous layer was extracted with EtOAc (x 3) and the

15

20


combined organic layers were washed with brine, dried (Na_2SO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 10-50% EtOAc in cyclohexane, to afford the *title compound* (550 mg, 37%) as a white foam. LCMS (Method 3): Rt 3.87 min, m/z 405 [MH^+].

5 **b. 1-[5-tert-Butyl-2-(6-methyl-pyridin-3-yl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 102)**

A mixture of Intermediate 102a (84 mg, 0.206 mmol), Intermediate 3c (75 mg, 0.206 mmol) and DIPEA (45 μL , 0.258 mmol) in dioxane (3 mL) was stirred
10 at 65 °C for 16 h. After cooling, the volatiles were concentrated *in vacuo* and the resultant residue was partitioned between water and DCM. The organic layer was dried through a separator phase and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 2-7% MeOH in DCM followed by HPLC (XBridge C18 column, 25-75% MeCN in H_2O , 0.1% NH_4OH), to afford
15 the *title compound* (41 mg, 32%) as a white solid. LCMS (Method 5): Rt 4.06 min, m/z 620 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 1.27 (9H, s), 1.56-1.66 (2H, m), 1.67-1.77 (4H, m), 1.79-1.96 (2H, m), 1.96-2.20 (2H, m), 2.53 (3H, s), 3.13 (4H, t, $J = 5.2$ Hz), 4.75-4.84 (1H, m), 5.54 (1H, t, $J = 4.3$ Hz), 6.35 (1H, s), 7.06 (1H, d, $J = 8.6$ Hz), 7.16 (1H, dd, $J = 9.8, 2.2$ Hz), 7.23-7.43 (5H, m), 7.58-7.64 (2H, m),
20 7.81 (1H, dd, $J = 8.3, 2.6$ Hz), 8.18 (1H, s), 8.59 (1H, d, $J = 2.6$ Hz).


Example 103

1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



5


a. 2-(5-Fluoropyridin-2-yl)-N,N-di(propan-2-yl)hydrazinecarboxamide (Intermediate 103a)



To a solution of (5-fluoro-pyridin-2-yl)-hydrazine (For reference procedure
 10 see WO2010022076; 200 mg, 1.57 mmol) in DCM (15mL) was added triethylamine (328 μ L, 2.35 mmol) followed by *N,N*-diisopropylcarbamoyl chloride (282 mg, 1.73 mmol). The reaction mixture was heated at 45 $^{\circ}$ C for 4 h then poured into water and extracted with DCM (x 3). The combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The
 15 resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane, to afford the *title compound* (200 mg, 50%). ^1H NMR (400 MHz, CDCl_3): 1.30 (12H, d, J = 6.8 Hz), 3.80-4.03 (2H, m), 6.17 (1H, s), 6.53 (1H, s), 6.67-6.81 (1H, m), 7.24-7.36-(1H, m), 8.02 (1H, d, J = 3.0 Hz).

b. (6-Fluoro-[1,2,4]triazolo[4,3-a]pyridin-3-yl)-diisopropyl-amine (Intermediate 103b)


20



The *title compound* was prepared starting from Intermediate 103a using


analogous procedures to those described in Intermediate 99c. ¹H NMR (400 MHz, CDCl₃): 1.09 (12H, d, J = 6.5 Hz), 3.60-3.79 (2H, m), 7.11-7.24 (1H, m), 7.44-7.62 (1H, m), 7.98-8.05 (1H, m).

c. **[6-((1R,4S)-4-Amino-1,2,3,4-tetrahydro-naphthalen-1-yloxy)-[1,2,4]triazolo[4,3-a]pyridin-3-yl]-diisopropyl-amine (Intermediate 103c)**




To a suspension of sodium hydride (60% in mineral oil, 136 mg, 3.39 mmol) in DMF (5 mL) was added Intermediate A (138 mg, 0.85 mmol) and the reaction mixture was stirred for 15 min at RT. A solution of Intermediate 103b (200 mg, 0.85 mmol) in DMF (3 mL) was added and the reaction mixture was heated at 60 °C for 1 h. The reaction mixture was quenched by careful addition of MeOH and then loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (220 mg, 68%). LCMS (Method 4): Rt 2.07 min, m/z 380 [MH⁺].

d. **1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 103d)**




The *title compound* was prepared starting from Intermediate 103c and Intermediate 95c using analogous procedures to those described in Intermediate 95d. LCMS (Method 4): Rt 3.92 min, m/z 765 [MH⁺].

e. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
5 [(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-
tetrahydro-naphthalen-1-yl]-urea (Intermediate 103e)



To a solution of Intermediate 103d (100 mg, 0.13 mmol) in MeOH (2 mL) was added pyridinium p-toluenesulfonate (65 mg, 0.26 mmol) and the reaction mixture was heated to reflux temperature for 3 h. The volatiles were concentrated *in vacuo* and the resultant residue was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (97 mg, 99%). LCMS (Method 2): Rt 3.39 min, m/z 681 [MH⁺].

15 f. Methanesulfonic acid 2-[3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-pyrazol-1-yl)-phenoxy]-ethyl ester (Intermediate 103f)



20 To an ice-bath cooled solution of Intermediate 103e (97 mg, 0.13 mmol) in DCM (2 mL) was added DIPEA (58 μL, 0.32 mmol) followed by methanesulfonyl


chloride (22 μ L, 0.27 mmol). The reaction mixture was stirred for 2 h then partitioned between DCM and water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (130 mg, quantitative). LCMS (Method 1): Rt 3.69 min, m/z 759 [MH^+].

g. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-diisopropylamino-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 103)

To a solution of Intermediate 103f (130 mg, 0.13 mmol) in THF (1.5 mL) was added dimethylamine (2M in MeOH, 520 μ L, 1.04 mmol). The reaction mixture was heated at 50 $^{\circ}\text{C}$ for 18 h, then cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM followed by HPLC (C18 X-select column, 10-98% gradient MeCN in H_2O , 0.1% HCO_2H), to give the *title compound* (37 mg, 40%). LCMS (Method 5): Rt 3.70 min, m/z 708 [MH^+]. ^1H NMR (400 MHz, CD_3OD): 1.05 (12H, dd, J = 6.4, 1.8 Hz), 1.33 (9H, s), 2.18-2.30 (4H, m), 2.43 (6H, s), 2.91 (2H, t, J = 5.2 Hz), 3.62-3.74 (2H, m), 4.19 (2H, t, J = 5.3 Hz), 4.88-4.97 (1H, m), 5.42 (1H, t, J = 4.1 Hz), 6.37 (1H, s), 7.04-7.13 (3H, m), 7.19-7.34 (5H, m), 7.44 (1H, t, J = 8.3 Hz), 7.60 (1H, d, J = 9.8 Hz), 7.87 (1H, d, J = 2.1 Hz).


Example 104

N-(5-tert-Butyl-2-methoxy-3-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-phenyl)-methanesulfonamide



5

a. (5-tert-Butyl-3-methanesulfonylamino-2-methoxy-phenyl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 104a)



NaOH (199 mg, 4.97 mmol) was added to an ice-bath cooled solution of N-
 10 [3-amino-5-(1,1-dimethylethyl)-2-methoxyphenyl]-methanesulfonamide (For
 reference procedure see for example WO2010026096; 500 mg, 1.84 mmol). The
 ice bath was removed and complete dissolution of reagents occurred. 2,2,2-
 Trichloroethyl chloroformate (380 μ L, 2.76 mmol) was added and the reaction
 mixture was warmed to RT and stirred for 90 min. The reaction mixture was
 15 partitioned between EtOAc and water. The aqueous phase was extracted with
 EtOAc (x 3) and the combined organic layers were washed with brine, dried
 (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on
 silica, using a gradient of 0-60% EtOAc in cyclohexane, to afford the *title*
compound (620 mg, 75%) as a white solid. LCMS (Method 1): Rt 3.98 min, m/z
 20 445, 447, 449 [MH^+].


b. N-(5-tert-Butyl-2-methoxy-3-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-phenyl)-methanesulfonamide (Example 104)

A mixture of Intermediate 3c (50 mg, 0.14 mmol), Intermediate 104a (62


mg, 0.14 mmol) and DIPEA (48 μ L, 0.27 mmol) in dioxane (1.5 mL) was stirred at 70 $^{\circ}$ C for 18 h and then at 80 $^{\circ}$ C for 48 h. The reaction mixture was diluted with EtOAc and then poured into water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-8% MeOH in DCM and then by HPLC (C18 X-select column, 10-98% gradient MeCN in H_2O , 0.1% HCO_2H), to afford the *title compound* (19 mg, 21%). LCMS (Method 5): Rt 4.47 min, m/z 662 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 1.25 (9H, s), 1.57-1.66 (2H, m), 1.69-1.78 (4H, m), 1.83-1.95 (1H, m), 1.98-2.24 (3H, m), 3.05 (3H, s), 3.15 (4H, t, $J = 5.3$ Hz), 3.69 (3H, s), 4.87-4.96 (1H, m), 5.59 (1H, t, $J = 4.5$ Hz), 6.94 (1H, d, $J = 2.3$ Hz), 7.19 (1H, dd, $J = 9.9$, 2.1 Hz), 7.31 (1H, dd, $J = 8.2$, 6.7 Hz), 7.35-7.45 (4H, m), 7.60-7.69 (2H, m), 8.07 (1H, s), 8.19 (1H, d, $J = 2.4$ Hz).

Example 105

15 **1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[8-methyl-3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**




a. **N-Benzhydrylidene-N'-(5-fluoro-3-methyl-pyridin-2-yl)-hydrazine (Intermediate 105a)**



A mixture of 2-bromo-5-fluoro-3-methyl-pyridine (190 mg, 1.0 mmol), benzhydrylidene-hydrazine (216 mg, 1.1 mmol), phenylboronic acid (5 mol%, 6


mg) and potassium *tert*-butoxide (157 mg, 1.4 mmol) in anhydrous toluene (6 mL) was degassed with a stream of argon, treated with Pd(OAc)₂ (2 mol%, 5 mg) and racemic (1,1'-binaphthalene-2,2'-diyl)bis(diphenylphosphine) (2 mol%, 12 mg) and stirred at 80 °C for 1 h. After cooling to RT, the reaction mixture was diluted
5 with EtOAc and washed with water followed by a saturated aqueous solution of NaHCO₃. The organic layer was dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-10% MeOH in DCM followed by 100% DCM, to afford the *title compound* (196 mg, 64%). LCMS (Method 4): Rt 3.44 min, m/z 306 [MH⁺].

10 **b. (5-Fluoro-3-methyl-pyridin-2-yl)-hydrazine (Intermediate 105b)**



A solution of Intermediate 105a (194 mg, 0.635 mmol) in toluene (13 mL) was treated with HCl (37%, 3.5 mL) and heated at 110 °C for 1.5 h. After cooling to RT, the reaction mixture was diluted with water and extracted with toluene (x
15 3). The aqueous layer was cooled to 0 °C, neutralised with 5N NaOH and extracted with DCM (x 3). The combined organic layers were dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (78 mg, 86%) as a silver/green solid. LCMS (Method 4): Rt 0.29 min, m/z 142 [MH⁺].


20 **c. (S)-1-Methyl-pyrrolidine-2-carboxylic acid N'-(5-fluoro-3-methyl-pyridin-2-yl)-hydrazide (Intermediate 105c)**



A solution of Intermediate 105b (76 mg, 0.537 mmol), (S)-1-methyl-pyrrolidine-2-carboxylic acid (55 mg, 0.72 mmol) and HOBt (8 mg, 0.72 mmol) in anhydrous DCM (5 mL) was treated with EDC (138 mg, 0.717 mmol) and
25 stirred at RT under a nitrogen atmosphere for 18 h. The reaction mixture was diluted with DCM and washed with brine. The organic layer was dried (MgSO₄)


and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-10% MeOH in DCM, to afford the *title compound* (76 mg, 56%). ¹H NMR (400 MHz, CDCl₃): 1.74-2.05 (3H, m), 2.15-2.33 (4H, m), 2.34-2.46 (1H, m), 2.52 (3H, s), 3.02-3.12 (1H, m), 3.15-3.25 (1H, m), 6.70 (1H, s), 7.08-7.18 (1H, m), 7.90 (1H, d, J = 2.8 Hz), 9.52 (1H, br s).

d. 6-Fluoro-8-methyl-3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 105d)



Hexachloroethane (141 mg, 0.587 mmol) was added portionwise at RT to a stirred mixture of Intermediate 105c (74 mg, 0.293 mmol), triethylamine (166 μ L, 1.17 mmol) and triphenylphosphine (154 mg, 0.587 mmol) in THF (5 mL). The reaction mixture was stirred at 80 $^{\circ}$ C for 2 h, then cooled to RT and filtered. The filtrate was concentrated *in vacuo* and the resultant residue was purified by SCX cartridge. The cartridge was washed with 10% MeOH in DCM and the product eluted with 10% (2M NH₃ in MeOH) in DCM. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 1-10% MeOH in DCM, to afford the *title compound* (59 mg, 86%) as a colourless oil. ¹H NMR (400 MHz, CDCl₃): 1.88-2.12 (3H, m), 2.15-2.53 (5H, m), 2.69 (3H, s), 3.28 (1H, t, J = 7.8 Hz), 4.01-4.15 (1H, m), 6.97 (1H, d, J = 8.4 Hz), 8.50 (1H, s).

e. (1S,4R)-4-[8-Methyl-3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 105e)



A mixture of Intermediate A (44 mg, 0.267 mmol) and sodium hydride (60% in mineral oil, 29 mg, 0.729 mmol) in anhydrous DMF (1 mL) was stirred at RT for 20 min under argon atmosphere. To the reaction mixture was added a solution of Intermediate 105d (57 mg, 0.243 mmol) in anhydrous DMF (2 mL) and stirring at 60 °C was continued for 2.5 h. After cooling to RT, the reaction mixture was diluted with EtOAc and poured into ice-water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by SCX cartridge by washing the cartridge with 10% MeOH in DCM and eluting the product with 10% (2M NH₃ in MeOH) in DCM and then by FCC on silica, using 10% MeOH in DCM followed by 10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (53 mg, 58%). LCMS (Method 1): Rt 0.38, 1.54 min, m/z 378 [MH⁺].


f. 1-(5-tert-Butyl-2-p-tolyl-2H-pyrazol-3-yl)-3-((1S,4R)-4-[8-methyl-3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 105)

A mixture of Intermediate 105e (53 mg, 0.14 mmol), (5-tert-butyl-2-p-tolyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference procedure see Synthetic Communications, 2009, 39, 3999-4009; 67 mg, 0.168 mmol) and DIPEA (49 µL, 0.28 mmol) in DMF (2 mL) was stirred at 100 °C for 1.5 h under argon atmosphere. After cooling to RT, the reaction mixture was diluted with EtOAc and washed with brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane followed by 10% [2M NH₃ in MeOH] in EtOAc. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by HPLC (C18 Phenomenex Gemini column, 5-98% gradient MeCN in H₂O, 0.1% NH₄OH), to afford the *title compound* (52 mg, 74%). LCMS (Method 5): Rt 3.76 min, m/z 633 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.23 (9H, s), 1.75-2.30 (13H, m), 2.32 (3H, s), 2.48


(3H, s), 3.19 (1H, br s), 4.72-4.83 (1H, m), 5.30-5.38 (1H, m), 6.28 (1H, s), 7.03-7.11 (2H, m), 7.20-7.36 (8H, m), 8.00 (1H, s), 8.11 (1H, s).

Example 106

1-{5-tert-Butyl-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea




a. 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 106a)



A mixture of Intermediate 3c (100 mg, 0.28 mmol), Intermediate 33a (116 mg, 0.28 mmol) and DIPEA (96 μ L, 0.55 mmol) in dioxane (3 mL) was stirred at 80 $^{\circ}$ C for 16 h. After cooling, the reaction mixture was partitioned between water and EtOAc. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-10% MeOH in DCM, to afford the *title compound* (156 mg, 88%). LCMS (Method 4): Rt 3.07 min, m/z 635 [MH^+].

b. 1-[5-tert-Butyl-2-(4-chloromethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-

tetrahydro-naphthalen-1-yl]-urea (Intermediate 106b)


To an ice-bath cooled solution of Intermediate 106a (150 mg, 0.25 mmol) in DCM (3 mL) was added DIPEA (107 μ L, 0.61 mmol) followed by methanesulfonyl chloride (29 μ L, 0.37 mmol). The reaction mixture was stirred for 2 h. After warming to RT, additional amount of DIPEA (65 μ L) and methanesulfonyl chloride (30 μ L) were added and stirring was continued for 2 h. The reaction mixture was partitioned between DCM and water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.59 min, m/z 653 [MH⁺].

c. 1-{5-tert-Butyl-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 106)

To a solution of Intermediate 106b (0.25 mmol) in THF (2.5 mL) was added DIPEA (130 μ L, 0.75 mmol) followed by N-methylpiperazine (83 μ L, 0.75 mmol). The reaction mixture was heated at 50 °C for 18 h in a sealed tube, then cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM followed by HPLC (C18 X-select column, 10-98% gradient MeCN in H₂O, 0.1% HCO₂H), to give the *title compound* (13 mg, 7%). LCMS (Method 5): Rt 3.44 min,

m/z 717 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.57-1.66 (2H, m), 1.68-1.77 (4H, m), 1.79-1.95 (2H, m), 1.98-2.19 (5H, m), 2.22-2.46 (8H, m), 3.14 (4H, t, J = 5.3 Hz), 3.49 (2H, s), 4.76-4.85 (1H, m), 5.54 (1H, t, J = 4.4 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.17 (1H, dd, J = 9.8, 2.2 Hz), 7.25-7.47 (8H, m),
5 7.59-7.65 (2H, m), 8.11 (1H, s).

Example 107




1-(3-tert-Butyl-1'-methyl-1'H-[1,4']bipyrazolyl-5-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-
10 **naphthalen-1-yl]-urea**

To a solution of Intermediate 3c (100 mg, 0.27 mmol) in dioxane (3 mL) was added DIPEA (90 μL, 0.55 mmol) followed by (3-tert-butyl-1'-methyl-1'H-[1,4']bipyrazolyl-5-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference procedure see for example US6492529; 109 mg, 0.27 mmol). The reaction mixture
15 was heated at 60 °C for 18 h, then cooled and poured into water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% MeOH in DCM followed by MDAP (Method 7) purification, to give the *title compound* (100 mg,
20 61%). LCMS (Method 5): Rt 4.05 min, m/z 609 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.24 (9H, s), 1.54-1.66 (2H, m), 1.67-1.76 (4H, m), 1.81-2.18 (4H, m), 3.13 (4H, t, J = 5.2 Hz), 3.86 (3H, s), 4.78-4.87 (1H, m), 5.55 (1H, t, J = 4.1 Hz), 6.26 (1H, s), 7.10-7.18 (2H, m), 7.25-7.41 (4H, m), 7.58-7.64 (3H, m), 7.98-8.02 (2H, m).


Example 108

1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



5


a. 3-tert-Butyl-1'-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1'H-[1,4']bipyrazolyl-5-ylamine (Intermediate 108a)



To a mixture of 4-iodo-1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-pyrazole
 10 (For reference procedure see for example WO2010139731; 1.54 g, 4.81 mmol), 5-tert-butyl-2H-pyrazol-3-ylamine (668 mg, 4.81 mmol), copper (I) iodide (45 mg, 0.24 mmol) and K₂CO₃ (1.39 g, 10.1 mmol) was added a solution of toluene (5 mL), previously degassed by using a stream of argon. (R,R)-(-)-N,N'-Dimethyl-1,2-cyclohexanediamine (151 μL, 0.96 mmol) was then added and the reaction
 15 mixture was heated at 150 °C for 2.5 h under microwave irradiation. The crude reaction mixture was poured into water and extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane, to give the *title compound* (1.29 g, 81%). LCMS

(Method 1): Rt 2.31 min, m/z 334 [MH⁺].

b. {3-tert-Butyl-1'-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1'H-[1,4']bipyrazolyl-5-yl}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 108b)




5

To a mixture of Intermediate 108a (1.29 g, 3.87 mmol) in water (5 mL) and EtOAc (13 mL) was added NaOH (310 mg, 7.74 mmol). After 10 min stirring, 2,2,2-trichloroethyl chloroformate (640 μ L, 4.65 mmol) was added and the reaction mixture was stirred at RT for 1 h. The aqueous layer was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-75% EtOAc in cyclohexane, to afford the *title compound* (1.24 g, 63%). LCMS (Method 4): Rt 3.92 min, m/z 508, 510 [MH⁺].

15

20


c. 1-{3-tert-Butyl-1'-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1'H-[1,4']bipyrazolyl-5-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 108c)



5

The *title compound* was prepared starting from Intermediate 3c and Intermediate 108b using analogous procedures to those described in Intermediate 106a. LCMS (Method 4): Rt 3.27 min, m/z 723.5 [MH⁺].

d. 1-[3-tert-Butyl-1'-(2-hydroxy-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 108d)




To a solution of Intermediate 108c (170 mg, 0.23 mmol) in MeOH (2.5 mL) was added pyridinium p-toluenesulfonate (118 mg, 0.47 mmol) and the reaction mixture was heated at 60°C for 2 h. The resultant mixture was poured into water and a saturated aqueous solution of NaHCO₃ was added. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified

15

by FCC on silica, using a gradient of 0-10% MeOH in DCM, to give the *title compound* (105 mg, 72%). LCMS (Method 1): Rt 2.96 min, m/z 639 [MH⁺].

- e. Methanesulfonic acid 2-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-[1,4']bipyrzoly-1'-yl)-ethyl ester (Intermediate 108e)**




To an ice-bath cooled solution of Intermediate 108d (105 mg, 0.16 mmol) in DCM (2 mL) was added DIPEA (43 μ L, 0.25 mmol) followed by methanesulfonyl chloride (15 μ L, 0.20 mmol). The reaction mixture was stirred for 2 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.09 min, m/z 717 [MH⁺].

- f. 1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrzoly-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 108)**

To a solution of Intermediate 108e (0.20 mmol) in THF (2 mL) was added dimethylamine (2M in MeOH, 0.8 mL, 1.6 mmol) and stirring at RT was continued for 48 h. The crude reaction mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM followed by MDAP (Method 7) purification, to give

the *title compound* (20 mg, 15%). LCMS (Method 5): Rt 3.34 min, m/z 666 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.24 (9H, s), 1.54-1.65 (2H, m), 1.67-1.77 (4H, m), 1.81-2.13 (4H, m), 2.16 (6H, s), 2.67 (2H, t, J = 6.5 Hz), 3.13 (4H, t, J = 5.3 Hz), 4.20 (2H, t, J = 6.5 Hz), 4.79-4.87 (1H, m), 5.55 (1H, t, J = 4.5 Hz), 6.26 (1H, s), 7.11-7.19 (2H, m), 7.26-7.41 (4H, m), 7.58-7.64 (3H, m), 7.99-8.05 (2H, m).

Example 109




N-[5-tert-Butyl-2-methoxy-3-(3-((1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-ureido)-phenyl]-methanesulfonamide formate salt

A mixture of Intermediate 5c (122 mg, 0.33 mmol), Intermediate 104a (150 mg, 0.33 mmol) and DIPEA (87 μL, 0.50 mmol) in dioxane (2 mL) was stirred at 60°C for 18 h and then at 85°C for 48 h. The volatiles were concentrated *in vacuo* and the resultant residue was purified by MDAP (Method 7) to afford the *title compound* (88 mg, 40%). LCMS (Method 5): Rt 3.41 min, m/z 662 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.24 (9H, s), 1.81-2.27 (10H, m), 2.28-2.40 (1H, m), 3.04 (3H, s), 3.11-3.18 (2H, m), 3.67 (3H, s), 3.99 (1H, t, J = 8.3 Hz), 4.83-4.99 (1H, m), 5.43 (1H, t, J = 4.5 Hz), 6.93 (1H, d, J = 2.3 Hz), 7.25-7.44 (7H, m), 7.76 (1H, d, J = 10.3 Hz), 8.05 (1H, s), 8.15 (1H, s), 8.18 (1H, d, J = 2.8 Hz), 8.27 (1H, d, J = 2.1 Hz).


Example 110

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




5

a. 2-(5-Amino-3-tert-butyl-pyrazol-1-yl)-ethanol (Intermediate 110a)



2-Hydrazino-ethanol (2.98 mL, 44.0 mmol) was added to a stirred solution of 4,4-dimethyl-3-oxo-pentanenitrile (5.0 g, 40.0 mmol) in EtOH (IMS grade, 40 mL) at RT followed by concentrated HCl (0.4 mL). The reaction mixture was heated at reflux temperature for 18 h then cooled and concentrated *in vacuo*. The resultant residue was triturated with pentane to give the title compound (7.4 g, quantitative) as a yellow solid. LCMS (Method 3): Rt 0.81 min, m/z 184 [MH⁺].

b. [5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 110b)



2,2,2-Trichloroethyl chloroformate (3.24 mL, 23.5 mmol) was added dropwise over 5 min to a solution of Intermediate 110a (4.11 g, 22.4 mmol) in NaOH (1M in water, 33.6 mL, 33.6 mmol) and EtOAc (35 mL) cooled at 0 °C under argon atmosphere. The reaction mixture was stirred at RT for 5 h. Additional

20


2,2,2-trichloroethyl chloroformate (462 μ L) was added and stirring at RT was continued for 16 h. Additional NaOH (1M in water, 15 mL) and 2,2,2-trichloroethyl chloroformate (462 μ L) were added and stirring at RT was continued for 1 h. The aqueous layer was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (Na_2SO_4) and concentrated *in vacuo*. The resultant residue was dissolved in cyclohexane (100 mL) and left to stand for 5 days. The resulting suspension was filtered and the solid collected by filtration washing with additional cyclohexane affording *the title compound* (3.64 g, 45%) as a white solid. The filtrate was concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a mixture of 50% EtOAc in cyclohexane, then partitioned between DCM and water. The organic layer was dried through a phase separator and concentrated *in vacuo* to afford additional *title compound* (1.43 g, 18%) as an orange gum. LCMS (Method 3): Rt 3.72 min, m/z 358 [MH^+].

c. 1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 110)

A mixture of Intermediate 110b (143 mg, 0.40 mmol), Intermediate 81d (1151mg, 0.40 mmol) and DIPEA (82 μ L, 0.50 mmol) in dioxane (5 mL) was stirred at 100 $^{\circ}\text{C}$ for 4 h. After cooling, the reaction mixture was concentrated *in vacuo* and then partitioned between water and DCM. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried through a phase separator and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 3-15% MeOH in DCM followed by MDAP (Method 7) purification, to afford the *title compound* (36 mg, 15%) as a glassy white solid. LCMS (Method 5): Rt 4.02 min, m/z 587 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.92 (3H, d, $J = 6.3$ Hz), 1.21 (9H, s), 1.44-1.57 (2H, m), 1.60-1.74 (2H, m), 1.74-1.87 (2H, m), 1.88-2.22 (4H, m), 2.86-2.96 (1H, m),

3.12-3.20 (2H, m), 3.63-3.71 (2H, m), 3.95 (2H, t, $J = 6.0$ Hz), 4.82-4.91 (1H, m), 5.00 (1H, br s), 5.54 (1H, t, $J = 4.3$ Hz), 6.07 (1H, s), 7.10 (1H, d, $J = 8.6$ Hz), 7.22 (1H, dd, $J = 9.8, 2.2$ Hz), 7.26-7.32 (1H, m), 7.35-7.42 (3H, m), 7.65 (1H, d, $J = 10.8$ Hz), 7.71 (1H, d, $J = 2.1$ Hz), 8.19-8.23 (1H, m).

5


Example 111

1-(5-tert-Butyl-isoxazol-3-yl)-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea

10

The *title compound* was prepared starting from Intermediate 81d and (5-tert-butyl-isoxazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (For reference procedure see WO2006091671) using analogous procedures to those described in Example 110 step c. LCMS (Method 5): R_t 4.67 min, m/z 544 $[MH^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 0.91 (3H, d, $J = 6.3$ Hz), 1.28 (9H, s), 1.44-1.57 (2H, m), 1.60-1.73 (2H, m), 1.74-1.86 (2H, m), 1.88-2.22 (4H, m), 2.86-2.94 (1H, m), 3.12-3.20 (2H, m), 4.86-4.95 (1H, m), 5.54 (1H, t, $J = 4.2$ Hz), 6.40 (1H, s), 7.00 (1H, d, $J = 8.6$ Hz), 7.21-7.26 (1H, m), 7.27-7.33 (1H, m), 7.34-7.42 (3H, m), 7.64 (1H, dd, $J = 9.8, 0.8$ Hz), 7.71 (1H, d, $J = 2.10$ Hz), 9.31 (1H, s).

15

Example 112


20

1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt


The *title compound* was prepared starting from Intermediate 108e and morpholine using analogous procedures to those described in Example 108. LCMS (Method 5): Rt 3.37 min, m/z 708 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.24 (9H, s), 1.55-1.65 (2H, m), 1.67-1.76 (4H, m), 1.79- 2.19 (4H, m), 2.41 (4H, d, J = 4.7 Hz), 2.72 (2H, t, J = 6.7 Hz), 3.13 (4H, t, J = 5.3 Hz), 3.53 (4H, t, J = 4.6 Hz), 4.23 (2H, t, J = 6.6 Hz), 4.78-4.87 (1H, m), 5.55 (1H, t, J = 4.4 Hz), 6.25 (1H, s), 7.11-7.18 (2H, m), 7.25-7.41 (4H, m), 7.58-7.64 (3H, m), 8.00 (1H, s), 8.05 (1H, s), 8.16 (1H, s).

Example 113

1-[3-tert-Butyl-1'-(3-dimethylamino-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt




a. 4-Iodo-1-[3-(tetrahydro-pyran-2-yloxy)-propyl]-1H-pyrazole (Intermediate 113a)



To a mixture of 4-iodo-1H-pyrazole (2.0 g, 10.3 mmol) and Cs₂CO₃ (5.04 g, 15.5 mmol) in MeCN (28 mL) was added 2-(3-bromopropoxy)tetrahydro-2H-pyran (1.84 mL, 10.8 mmol) and the mixture stirred overnight. The crude reaction

mixture was poured into water and extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-80% EtOAc in cyclohexane, to give the *title compound* (2.95 g, 81%). ¹H NMR (300 MHz, CDCl₃): 1.50-1.58 (4H, m), 1.65-1.90 (2H, m), 2.12 (2H, qn, J = 6.4 Hz), 3.35 (1H, dt, J = 10.2, 5.9 Hz), 3.46-3.54 (1H, m), 3.73 (1H, dt, J = 10.2, 5.9 Hz), 3.80-3.88 (1H, m), 4.26 (2H, td, J = 6.9, 1.5 Hz), 4.54 (1H, dd, J = 4.5, 3.1 Hz), 7.46 (1H, s), 7.50 (1H, s).


b. 3-tert-Butyl-1'-[3-(tetrahydro-pyran-2-yloxy)-propyl]-1'H-[1,4']bipyrazolyl-5-ylamine (Intermediate 113b)



To a mixture of intermediate 113a (1.50 g, 4.46 mmol), 5-tert-butyl-2H-pyrazol-3-ylamine (620 mg, 4.46 mmol), copper (I) iodide (42 mg, 0.22 mmol) and K₂CO₃ (1.29 g, 9.37 mmol) was added a solution of toluene (4.6 mL), previously degassed by using a stream of argon. (R,R)-(-)-N,N'-Dimethyl-1,2-cyclohexanediamine (141 µL, 0.89 mmol) was then added and the reaction mixture was heated at 140 °C for 2.5 h under microwave irradiation. The crude reaction mixture was poured into water and extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane, to give the *title compound* (1.14 g, 73%). LCMS (Method 4): Rt 2.34 min, m/z 348 [MH⁺].


c. {3-tert-Butyl-1'-[3-(tetrahydro-pyran-2-yloxy)-propyl]-1'H-[1,4']bipyrazolyl-5-yl}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate

113c)




To a stirred mixture of Intermediate 113b (1.14 g, 3.28 mmol) in water (6 mL) and EtOAc (12 mL) was added NaOH (263 mg, 6.57 mmol). After 10 min, 2,2,2-trichloroethyl chloroformate (543 μ L, 3.94 mmol) was added and the reaction mixture was stirred at RT for 1 h. The aqueous layer was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane, to afford the *title compound* (1.57 g, 91%). LCMS (Method 4): Rt 3.99 min, m/z 522, 524 [MH⁺].

d. 1-{3-tert-Butyl-1'-[3-(tetrahydro-pyran-2-yloxy)-propyl]-1'H-[1,4']bipyrazolyl-5-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 113d)




The *title compound* was prepared starting from Intermediate 3c and Intermediate 113c using analogous procedures to those described in Intermediate 106a. LCMS (Method 4): Rt 3.34 min, m/z 737 [MH⁺].

e. 1-[3-tert-Butyl-1'-(3-hydroxy-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-

tetrahydro-naphthalen-1-yl]-urea (Intermediate 113e)

To a solution of Intermediate 113d (263 mg, 0.36 mmol) in MeOH (2.5 mL) was added pyridinium p-toluenesulfonate (179 mg, 0.71 mmol) and the reaction mixture was heated at 60 °C for 3 h. The resultant mixture was poured into water and a saturated aqueous solution of NaHCO₃ was added. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 3-10% MeOH in DCM, to give the *title compound* (163 mg, 70%). LCMS (Method 4): Rt 3.00 min, m/z 653 [MH⁺].

f. Methanesulfonic acid 3-(3-tert-butyl-5-{3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-ureido}-[1,4']bipyrazolyl-1'-yl)-propyl ester (Intermediate 113f)




To an ice-bath cooled solution of Intermediate 113e (163 mg, 0.25 mmol) in DCM (3 mL) was added DIPEA (174 μL, 1.0 mmol) followed by methanesulfonyl chloride (39 μL, 0.50 mmol). The reaction mixture was stirred for 1 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.12 min, m/z 731 [MH⁺].

g. 1-[3-tert-Butyl-1'-(3-dimethylamino-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt (Example 113)

To a solution of Intermediate 113f (0.125 mmol) in THF (1.5 mL) was added dimethylamine (2M in MeOH, 0.5 mL, 1.0 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (10 mg, 12%). LCMS (Method 5): Rt 3.36 min, m/z 680 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.21 (9H, s), 1.54-1.65 (2H, m), 1.67-1.77 (4H, m), 1.81-1.95 (4H, m), 1.99-2.07 (4H, m) 2.08 (6H, s), 2.16 (2H, t, J = 6.9 Hz), 3.10 (4H, t, J = 5.2 Hz), 4.09 (2H, t, J = 7.1 Hz), 4.79 (1H, td, J = 8.4, 5.5 Hz), 5.51 (1H, t, J = 4.5 Hz), 6.21 (1H, s), 7.09-7.14 (2H, m), 7.22-7.37 (4H, m), 7.55-7.62 (3H, m), 7.97-8.01 (2H, m), 8.22 (0.4H, br s).

Example 114




1-[3-tert-Butyl-1'-(3-morpholin-4-yl-propyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

The *title compound* was prepared starting from Intermediate 113f and morpholine using analogous procedures to those described in Example 113. LCMS (Method 5): Rt 3.36 min, m/z 722 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.20


(9H, s), 1.54-1.60 (2H, m), 1.65-1.72 (4H, m), 1.80-1.96 (4H, m), 1.98-2.10 (4H, m) 2.22 (2H, t, J = 7.0 Hz), 2.24-2.30 (4H, m), 3.10 (4H, t, J = 5.2 Hz), 3.50 (4H, t, J = 4.6 Hz), 4.09 (2H, t, J = 7.0 Hz), 4.79 (1H, td, J = 8.4, 5.5 Hz), 5.51 (1H, t, J = 4.5 Hz), 6.21 (1H, s), 7.10-7.15 (2H, m), 7.22-7.36 (4H, m), 7.55-7.62 (3H, m),
 5 7.97-8.01 (2H, m), 8.22 (0.3H, br s).

Example 115

1-{(1S,4R)-4-[3-((2S,6R)-2,6-Dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-(3-fluoro-5-morpholin-4-yl-phenyl)-urea



a. (3-Fluoro-5-morpholin-4-yl-phenyl)-carbamic acid 2,2,2-trichloroethyl ester (Intermediate 115a)



2,2,2-Trichloroethyl chloroformate (1.05 mL, 7.65 mmol) was added to an
 15 ice cold solution of 3-fluoro-5-morpholin-4-ylaniline (reference procedure see EP1102743; 1.0 g, 5.10 mmol) and triethylamine (1.77 mL, 10.2 mmol) in THF (25 mL). The reaction mixture was stirred at RT for 1.5 h, then quenched with water. The aqueous layer was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*
 20 to afford the *title compound* (1.89 g, 99%) as a clear oil. LCMS (Method 3): Rt 3.75 min, m/z 371, 373 [MH⁺].


b. 1-{(1S,4R)-4-[3-((2S,6R)-2,6-Dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-

(3-fluoro-5-morpholin-4-yl-phenyl)-urea (example 115)

The *title compound* was prepared starting from Intermediate 96c and intermediate 115a using analogous procedures to those described in Example 99.

LCMS (Method 5): Rt 4.70 min, m/z 614 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO):

5 0.55 (3H, d, J = 6.3 Hz), 0.59 (3H, d, J = 6.3 Hz), 1.35-1.53 (3H, m), 1.63-1.69 (2H, m), 1.72-1.78 (1H, m), 1.82-1.99 (2H, m), 2.00-2.13 (2H, m), 3.03 (4H, t, J = 4.8 Hz), 3.08-3.20 (2H, m), 3.65-3.69 (2H, t, J = 4.8 Hz), 4.80-4.88 (1H, td, J = 8.7, 5.5 Hz), 5.50 (1H, t, J = 4.0 Hz), 6.31 (1H, dt, J = 12.4, 2.0 Hz), 6.68 (1H, br s), 6.71 (1H, d, J = 8.9 Hz), 6.81 (1H, dt, J = 11.3, 1.9 Hz), 7.18-7.25 (2H, m),
10 7.29-7.37 (3H, m), 7.62 (1H, dd, J = 9.8, 0.6 Hz), 7.86 (1H, d, J = 2.0 Hz), 8.51 (1H, s).


Example 116

15 **1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea**

A solution of Example 110 (80 mg, 0.136 mmol), DIPEA (71 μ L, 0.409 mmol) and methanesulfonyl chloride (26 μ L, 0.34 mmol) in DCM (3 mL) was stirred at RT for 90 min. Additional DIPEA (71 μ L) and methanesulfonyl chloride (26 μ L) were added and stirring was continued for 1 h. Water (1 mL) and a
20 saturated aqueous solution of NaHCO₃ (2 mL) were added and the reaction mixture was vigorously stirred for 30 min. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were dried through a phase separator and concentrated *in vacuo*. The resultant residue was dissolved in DMF (2 mL)

and morpholine (36 μ L, 0.41 mmol) and heated at 75 $^{\circ}$ C for 18 h. After cooling, the volatiles were concentrated *in vacuo* and the resultant residue was loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH_3 in MeOH. The product containing fractions were combined and concentrated *in vacuo* and the resultant residue was purified by MDAP (Method 7) to afford the title compound (23 mg, 26%) as a white solid. LCMS (Method 5): Rt 3.48 min, m/z 656 $[\text{MH}^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 0.92 (3H, d, J = 6.3 Hz), 1.21 (9H, s), 1.46-1.58 (2H, m), 1.63-1.73 (2H, m), 1.74-1.87 (2H, m), 1.89-2.02 (2H, m), 2.02-2.24 (2H, m), 2.35-2.43 (4H, m), 2.63 (2H, t, J = 7.1 Hz), 2.87-2.97 (1H, m), 3.14-3.22 (2H, m), 3.55 (4H, t, J = 4.5 Hz), 4.01 (2H, t, J = 7.1 Hz), 4.82-4.92 (1H, m), 5.51-5.57 (1H, m), 6.04 (1H, s), 7.10 (1H, d, J = 8.6 Hz), 7.23 (1H, dd, J = 9.8, 2.2 Hz), 7.27-7.42 (4H, m), 7.65 (1H, d, J = 9.9 Hz), 7.72 (1H, s), 8.37-8.45 (1H, m).

Example 117




1-Cyclopropyl-3-[(1S,4R)-4-[3-[(2S,6R)-2,6-dimethyl-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

The *title compound* was prepared starting from Intermediate 96c and 4-nitrophenyl cyclopropylcarbamate (For reference procedure see WO2007/72158) using analogous procedures to those described in Example 99. LCMS (Method 5): Rt 4.70 min, m/z 614 $[\text{MH}^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 0.31-0.35 (2H, m), 0.52-0.60 (8H, m), 1.33-1.56 (3H, m), 1.63-1.70 (2H, m), 1.73-1.88 (3H, m), 1.94-2.11 (2H, m), 3.08-3.20 (2H, m), 4.78 (1H, td, J = 8.6, 6.8 Hz), 5.47 (1H, t, J = 3.8 Hz), 6.07 (1H, d, J = 2.1 Hz), 6.24 (1H, d, J = 8.9 Hz), 7.15-7.22 (2H, m), 7.27-

7.32 (3H, m), 7.62 (1H, d, J = 9.8 Hz), 7.84 (1H, d, J = 1.7 Hz), 8.12 (0.6H, br s).

Example 118




**1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-
5 ((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-
1,2,3,4-tetrahydro-naphthalen-1-yl]-urea**

A mixture of Intermediate 96c (160 mg, 0.41 mmol), Intermediate 110b (142 mg, 0.41 mmol) and DIPEA (142 μ L, 0.82 mmol) in dioxane (5 mL) was stirred at 80 °C overnight. After cooling, the reaction mixture was partitioned
10 between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-8% MeOH in DCM followed by MDAP (Method 7) purification, to afford the *title compound* (86 mg, 32%) as a glassy white solid. LCMS (Method
15 5): Rt 4.32 min, m/z 601 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.2 Hz), 0.58 (3H, d, J = 6.2 Hz), 1.17 (9H, s), 1.35-1.55 (3H, m), 1.63-1.69 (2H, m), 1.73-1.79 (1H, m), 1.82-2.00 (2H, m), 2.00-2.13 (2H, m), 3.09-3.19 (2H, m), 3.62 (2H, br s), 3.90 (2H, t, J = 5.9 Hz), 4.82 (1H, td, J = 8.6, 5.7 Hz), 4.92-4.98 (1H, br s), 5.50 (1H, t, J = 4.2 Hz), 6.02 (1H, s), 7.04 (1H, d, J = 8.6 Hz), 7.18 (1H,
20 dd, J = 9.8, 2.3 Hz), 7.20-7.25 (1H, m), 7.30-7.36 (3H, m), 7.62 (1H, dd, J = 9.7, 0.6 Hz), 7.86 (1H, d, J = 2.1 Hz), 8.14 (1H, s).


Example 119

1-{5-tert-Butyl-2-[1-(2-dimethylamino-ethyl)-1H-imidazol-4-yl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea



5

a. 4-Iodo-1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazole and 5-Iodo-1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazole (Intermediate 119a)




10 To an ice cold mixture of 4-iodo-1H-imidazole (1.0 g, 5.13 mmol) and sodium hydroxide (226 mg, 5.64 mmol) in DMF (10 mL) was added 2-(2-bromothoxy)tetrahydro-2H-pyran (1.18 mL, 5.64 mmol) and the mixture stirred overnight. The crude reaction mixture was poured into water and extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried (MgSO₄)

15 and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 20-100% EtOAc in cyclohexane, to give the *title compound* (1.50 g, 91%) as a 2:1 mixture of regioisomers. LCMS (Method 4): Rt 2.03 min, m/z 323 [MH⁺].


20

b. 5-tert-Butyl-2-{1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazol-4-yl}-2H-pyrazol-3-ylamine (Intermediate 119b)



To a mixture of Intermediate 119a (1.50 g, 4.66 mmol), 5-tert-butyl-2H-pyrazol-3-ylamine (648 mg, 4.66 mmol), copper(I)iodide (44 mg, 0.23 mmol) and K_2CO_3 (1.35 g, 9.78 mmol) was added a solution of toluene (10 mL), previously degassed by using a stream of argon. (R,R)-(-)-N,N'-Dimethyl-1,2-cyclohexanediamine (147 μ L, 0.93 mmol) was then added and the reaction mixture was heated at 150 $^{\circ}C$ for 3 h under microwave irradiation. The crude reaction mixture was poured into water and extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried ($MgSO_4$) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-100% EtOAc in cyclohexane, to give a product consisting of the *title compound* (550 mg) in mixture with the un-reacted 5-iodo-1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazole isomer. LCMS (Method 4): Rt 2.32 min, m/z 334 [MH^+].


c. (5-tert-Butyl-2-{1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazol-4-yl}-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 119c)



To a mixture of Intermediate 119b (550 mg) in water (3 mL) and EtOAc (6


mL) was added NaOH (132 mg, 3.30 mmol). After 10 min stirring, 2,2,2-trichloroethyl chloroformate (273 μ L, 1.98 mmol) was added and the reaction mixture was stirred at RT for 2 h. The aqueous layer was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-80% EtOAc in cyclohexane, to afford the *title compound* (175 mg). LCMS (Method 4): Rt 4.54 min, m/z 508, 510 [MH⁺].

d. **1-(5-tert-Butyl-2-{1-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1H-imidazol-4-yl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 119d)**



The *title compound* was prepared starting from Intermediate 3c and Intermediate 119c using analogous procedures to those described in Intermediate 106a. LCMS (Method 4): Rt 5.52 min, m/z 723 [MH⁺].

e. **1-{5-tert-Butyl-2-[1-(2-hydroxy-ethyl)-1H-imidazol-4-yl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 119e)**



To a solution of intermediate 119d (135 mg, 0.19 mmol) in MeOH (2 mL)

CC1(C)C2=CN3=C(C=C2N3CCN3CCOS(=O)(=O)C)NC(=O)N[C@@H]4C5=CC=CC=C5[C@H](O4)c6ccc7nnc(N7C8CCNCC8)c6

20 g. 1-{5-tert-Butyl-2-[1-(2-dimethylamino-ethyl)-1H-imidazol-4-yl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-(3-piperidin-1-yl-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 119)


To a solution of Intermediate 119f (0.10 mmol) in THF (2 mL) was added

dimethylamine (2M in MeOH, 0.4 mL, 0.8 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*.


- 5 The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM followed by HPLC (XBridge C18 column, 30-98% MeCN in H₂O, 0.1% NH₄OH), to give the *title compound* (12 mg, 18%). LCMS (Method 5): Rt 3.43 min, m/z 666 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.21 (9H, s), 1.54-1.60 (2H, m), 1.65-1.71 (4H, m), 1.86-1.94 (2H, m), 1.95-2.05 (1H, m) 2.13 (6H, s), 2.13-2.16 (1H, m), 2.55 (2H, t, J = 6.3 Hz), 3.10 (4H, t, J = 5.2 Hz), 4.05 (2H, t, J = 7.1 Hz), 4.86 (1H, td, J = 8.4, 5.5 Hz), 5.51 (1H, t, J = 4.5 Hz), 6.32 (1H, s), 7.12 (1H, dd, J = 10.0, 1.9 Hz), 7.20-7.25 (2H, m), 7.28-7.36 (3H, m), 7.56-7.59 (2H, m), 7.63 (1H, d, J = 1.4 Hz), 7.90 (1H, d, J = 8.6 Hz), 9.63 (1H, br s).

15 **Example 120**

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




- 20 **a. (S)-1-Methyl-piperidine-2-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 120a)**



To a solution of (S)-1-methyl-piperidine-2-carboxylic acid (250 mg, 1.74 mmol) and (5-fluoro-pyridin-2-yl)-hydrazine (244 mg, 1.92 mmol) in DCM (20


mL) was added HATU (790 mg, 2.09 mmol) and DIPEA (606 μ L, 3.48 mmol) and the reaction stirred for 18 h. The reaction was then heated to 45°C for 1 h then cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with
5 brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM to give the *title compound* (272 mg, 62%). LCMS (Method 4): Rt 0.87 min, m/z 253 [MH^+].

b.6-Fluoro-3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridine
10 **(Intermediate 120b)**



Hexachloroethane (505 mg, 2.14 mmol) was added portionwise over 5 min at RT to a stirred mixture of Intermediate 120a (270 mg, 1.07 mmol), triethylamine (595 μ L, 4.28 mmol) and triphenylphosphine (560 mg, 2.14 mmol) in THF (10
15 mL). The reaction mixture was stirred for 1.5 h and then diluted with methanol and loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH_3 in MeOH and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-8% [2M NH_3 in MeOH] in EtOAc to give the *title compound* (105 mg, 42%). ^1H NMR (400 MHz, CDCl_3): 1.34-1.51 (1H, m), 1.62-1.92 (5H, m), 2.00 (3H, s), 2.20 (1H, td, J = 11.4,
20 3.7 Hz), 3.06 (1H, m), 3.87 (1H, dd, J = 10.9, 3.3 Hz), 7.18 (1H, ddd, J = 9.8, 7.6, 2.3 Hz), 7.71 (1H, ddd, J = 9.9, 5.0, 0.7 Hz), 8.75-7.79 (1H, m).

c.(1S,4R)-4-[3-((S)-1-Methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 120c)



5 To a suspension of sodium hydride (60% in mineral oil, 54 mg, 1.34 mmol) in DMF (1 mL) was added Intermediate A (88 mg, 0.54 mmol) and the reaction mixture was stirred for 20 min. A solution of Intermediate 120b (105 mg, 0.45 mmol) in DMF (1 mL) was added and the reaction mixture was heated at 60 °C for 90 min. After cooling, the reaction mixture was quenched by careful addition of

10 MeOH and then loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The product containing fractions were concentrated *in vacuo* and the resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (77 mg, 45%). ¹H NMR (400 MHz, CDCl₃): 1.38-1.51 (1H, m), 1.52-1.90 (6H, m), 1.90-2.05 (1H, m), 2.03 (3H, s), 2.04-2.15 (1H, m), 2.15-2.22 (1H, m), 2.33-2.44 (1H, m), 2.99-3.06 (1H, m), 3.82 (1H, dd, J = 11.3, 3.2 Hz), 3.99 (1H, dd, J = 7.9, 5.1 Hz), 5.23 (1H, t, J = 4.5 Hz), 7.11 (1H, dd, J = 9.9, 2.2 Hz), 7.23-7.30 (1H, m), 7.34-7.41 (2H, m), 7.60 (1H, d, J = 7.7 Hz), 7.65 (1H, dd, J = 9.9, 0.7 Hz), 8.42 (1H, d, J = 2.2 Hz).

20 **d. 1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-piperidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Example 120)**

A mixture of Intermediate 120c (35 mg, 0.10 mmol), Intermediate 110b (54 mg, 0.15 mmol) and DIPEA (53 μL, 0.30 mmol) in dioxane (1 mL) was stirred at


25 50 °C overnight. After cooling, the reaction mixture was loaded onto an SCX

cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH. The resultant residue was purified by FCC on silica, using a gradient of 1-8% MeOH in DCM followed by MDAP (Method 7) purification, to afford the *title compound* (4 mg, 7%) as a glassy white solid. LCMS (Method 5):


5 Rt 3.02 min, m/z 586 [MH⁺]. ¹H NMR (400 MHz, d₄-MeOD): 1.20-1.30 (2H, m), 1.25 (9H, s), 1.40-1.52 (1H, m), 1.63-1.83 (3H, m), 1.85-1.92 (2H, m), 2.01 (3H, s), 2.06-2.20 (3H, m), 2.24 (1H, td, J = 11.7, 3.1 Hz), 2.31-2.38 (1H, m), 3.04-3.09 (1H, m), 3.76-3.81 (1H, m), 3.81 (2H, t, J = 5.4 Hz), 4.08 (2H, t, J = 5.4 Hz), 4.98 (1H, dd, J = 8.1, 6.5 Hz), 5.36 (1H, t, J = 3.8 Hz), 6.12 (1H, s), 7.24 (1H, t, J = 7.4 Hz), 7.28-7.37 (3H, m), 7.44 (1H, d, J = 7.8 Hz), 7.66 (1H, dd, J = 9.9, 0.6 Hz), 8.46 (1H, d, J = 1.9 Hz).

Example 121

1-**[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-**-(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea****



a. **Methanesulfonic acid 2-[3-tert-butyl-5-(3-**-(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-ethyl ester (Intermediate 121a)****





To an ice-bath cooled solution of Example 118 (43 mg, 0.07 mmol) in DCM (1.5 mL) was added DIPEA (49 μ L, 0.18 mmol) followed by methanesulfonyl chloride (11 μ L, 0.11 mmol). The reaction mixture was stirred for 1 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.38 min, m/z 679 [MH⁺].

**b. 1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 121)**

To a solution of Intermediate 121a (0.07 mmol) in THF (2 mL) was added DIPEA (50 μ L, 0.29 mmol) and morpholine (25 μ L, 0.29 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM followed by MDAP (Method 7) purification, to give the *title compound* (19 mg, 39%). LCMS (Method 5): Rt 3.65 min, m/z 670 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.56 (3H, d, J = 6.2 Hz), 0.59 (3H, d, J = 6.2 Hz), 1.16 (9H, s), 1.37-1.55 (3H, m), 1.63-1.69 (2H, m), 1.73-1.79 (1H, m), 1.85-2.00 (2H, m), 2.04-2.10 (2H, m), 2.34 (4H, t, J = 4.5 Hz), 2.57 (2H, t, J = 7.2 Hz), 3.09-3.19 (2H, m), 3.50 (4H, t, J = 4.5 Hz), 3.95 (2H, t, J = 7.2 Hz), 4.82 (1H, td, J = 8.6, 5.7 Hz), 5.50 (1H, t, J = 4.2 Hz), 5.99 (1H, s), 6.92 (1H, d, J = 8.6 Hz), 7.18 (1H, dd, J = 9.8, 2.3 Hz), 7.20-7.26 (1H, m), 7.30-7.36 (3H, m), 7.62 (1H, dd, J = 9.7, 0.6 Hz), 7.86 (1H, d, J = 2.1 Hz), 8.19 (1H, s).


Example 122

1-{(1S,4R)-4-[3-(8-Aza-bicyclo[3.2.1]oct-8-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-

dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea formate salt**a. 8-Aza-bicyclo[3.2.1]octane-8-carbonyl chloride (Intermediate 122a)**

5 To a cooled (10°C) solution 8-aza-bicyclo[3.2.1]octane (295 mg, 2.00 mmol) and pyridine (486 μ L, 6.02 mmol) in DCM (10 mL) was added triphosgene (595 mg, 2.00 mmol) portionwise. The reaction was allowed to reach room temperature and stirred overnight. The mixture was quenched with HCl (1M aqueous, 10 mL) and stirred for 30 min. The aqueous phase was extracted with

10 DCM (x 3) and the combined organic layers were washed with HCl (1M aqueous), brine, dried (MgSO₄) and concentrated *in vacuo*, to give the *title compound* (244 mg, 70%). ¹H NMR (400 MHz, CDCl₃): 1.46-1.70 (4H, m), 1.72-1.92 (5H, m), 2.04-2.11 (2H, m), 4.39-4.44 (2H, m).


b. 8-Aza-bicyclo[3.2.1]octane-8-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 122b)

To a solution of Intermediate 122a (240 mg, 1.38 mmol) and (5-fluoro-pyridin-2-yl)-hydrazine (176 mg, 1.38 mmol) in DCM (10 mL) was added DIPEA (721 μ L, 4.15 mmol) and the reaction stirred for 5 days. The reaction was then

20 partitioned between DCM and water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a

gradient of 0-10% MeOH in DCM to give the *title compound* (170 mg, 47%).
LCMS (Method 3): Rt 2.31 min, m/z 265 [MH⁺].

**c. 3-(8-Aza-bicyclo[3.2.1]oct-8-yl)-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine
(Intermediate 122c)**



5


Hexachloroethane (296 mg, 1.25 mmol) was added portionwise over 5 min at RT to a stirred mixture of Intermediate 122b (165 mg, 0.62 mmol), triethylamine (350 μ L, 2.50 mmol) and triphenylphosphine (327 mg, 1.25 mmol) in THF (10 mL). The reaction mixture was stirred for 3.5 h and then heated to 60°C overnight. The temperature was increased to 70°C for 5 days, then cooled and concentrated *in vacuo*. The residue was taken up in dioxane, and re-treated with hexachloroethane (150 mg, 0.63 mmol) and triethylamine (200 μ L, 1.43 mmol) at 115°C overnight. The mixture was cooled and concentrated *in vacuo*. The residue was taken up in methanol and loaded onto an SCX cartridge. The cartridge was washed with MeOH and the product eluted with 2M NH₃ in MeOH and concentrated *in vacuo* to give the *title compound* (117 mg, 76%). LCMS (Method 3): Rt 2.88 min, m/z 247 [MH⁺].

10

15

**d. (1S,4R)-4-[3-(8-Aza-bicyclo[3.2.1]oct-8-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine
(Intermediate 122d)**

20



To a suspension of sodium hydride (60% in mineral oil, 51 mg, 1.3 mmol) in DMF (1.5 mL) was added Intermediate A (73 mg, 0.45 mmol) followed by intermediate 122c (110 mg, 0.45 mmol) and the reaction mixture was heated at 60


°C for 1 h. After cooling, the reaction mixture was quenched by careful addition of NH₄Cl (saturated aqueous) and the aqueous phase was extracted with EtOAc (x 3). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a
5 gradient of 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* (98 mg, 56%). LCMS (Method 3): Rt 2.26 min, m/z 390 [MH⁺].

e. **1-{(1S,4R)-4-[3-(8-Aza-bicyclo[3.2.1]oct-8-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea formate salt (Example
10 **122)****


The titled compound was prepared using Intermediate 95c and Intermediate 122d following the procedures described in Example 95. LCMS (Method 5): Rt 3.55 min, m/z 318.5 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.27 (9H, s), 1.46-1.58 (3H, m), 1.63-1.79 (3H, m), 1.87-2.16 (8H, m), 2.19 (6H, s), 2.61 (2H, t, J =
15 5.9 Hz), 3.99 (2H, m), 4.08 (2H, t, J = 5.9 Hz), 4.81 (1H, m), 5.52 (1H, m), 6.32 (1H, s), 6.96 (1H, m), 7.05-7.17 (4H, m), 7.25-7.42 (5H, m), 7.56-7.61 (2H, m), 8.14 (1H, s), 8.19 (1H, s).

Example 123

**1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt
20**



a. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
 {(1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 123a)



5 The *title compound* was prepared from (R)-2-methyl piperidine using analogous procedures to those described for the preparation of Intermediate 95e. LCMS (Method 3): Rt 3.63 min, m/z 679 [MH⁺].

b. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-
 3-yl}-3-[(1S,4R)-4-[3-((R)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
 10 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt (Example 123)


To a solution of Intermediate 123a (108 mg, 0.159 mmol) and DIPEA (0.083 mL, 0.48 mmol) in DCM (4 mL) was added methanesulfonyl chloride (36.5 mg, 0.318 mmol) and the resulting yellow solution stirred at RT for 20 min. Water
 15 (2 mL) and sat. aq. NaHCO₃ solution (2 mL) were added, then the aqueous extracted with DCM (4 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a yellow solid.

The solid was dissolved in THF (2 mL), then dimethylamine (2 M in THF, 1.6 mL, 3.2 mmol) added, and the mixture stirred at 60°C in a sealed vial for 18 h.
 20 The cooled solution was concentrated *in vacuo*, redissolved in MeOH (1 mL), applied to an SCX-2 cartridge (5 g), washed with MeOH (20 mL) and eluted with 2 M NH₃ in MeOH (20 mL); concentration *in vacuo* gave a glassy yellow solid (98 mg). MDAP (Method 7) gave a pale yellow solid (63 mg). Additional purification by HPLC (Gemini C18, 15-65% MeCN in water, 0.1% HCO₂H, 20 min, × 2) gave
 25 the title compound as a white solid (18.6 mg, 17%). LCMS (Method 5): Rt 3.62


min, m/z 706.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.88 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.48-1.56 (2H, m), 1.65-1.71 (2H, m), 1.76-1.96 (4H, m), 2.00-2.14 (2H, m), 2.17 (6H, s), 2.59 (2H, t, J = 5.8 Hz), 2.95 (1H, ddd, J = 12.2, 8.1, 4.8 Hz), 3.18 (1H, dt, J = 12.1, 4.3 Hz), 3.26-3.29 (1H, m), 4.07 (2H, t, J = 5.8 Hz), 4.81 (1H, td, J = 8.6, 5.6 Hz), 5.53 (1H, t, J = 4.3 Hz), 6.31 (1H, s), 6.95 (1H, dd, J = 8.4, 2.2 Hz), 7.09 (2H, m), 7.18 (1H, dd, J = 9.8, 2.2 Hz), 7.25-7.41 (6H, m), 7.64 (1H, d, J = 9.9 Hz), 7.70 (1H, d, J = 2.1 Hz), 8.34 (1H, s), 8.50 (0.3H, s).

Example 124

1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




a. 1-{3-tert-Butyl-1'-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1'H-[1,4']bipyrazolyl-5-yl}-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 124a)



A mixture of Intermediate 96c (200 mg, 0.51 mmol), Intermediate 108b (261 mg, 0.51 mmol) and DIPEA (178 μ L, 1.02 mmol) in dioxane (5 mL) was


stirred at 80 °C overnight. After cooling, the reaction mixture was partitioned between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a
5 gradient of 1-10% MeOH in DCM to afford the *title compound* (364 mg, 95%) as a glassy white solid%). LCMS (Method 4): Rt 3.69 min, m/z 751 [MH⁺].

b. 1-[3-tert-Butyl-1'-(2-hydroxy-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-
{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate
10 124b)



To a solution of Intermediate 124a (239 mg, 0.32 mmol) in MeOH (3.5 mL) was added pyridinium *p*-toluenesulfonate (160 mg, 0.64 mmol) and the reaction mixture was heated at 60 °C for 3 h. The resultant mixture was poured into water and a saturated aqueous solution of NaHCO₃ was added. The aqueous phase was
15 extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 3-10% MeOH in DCM, to give the *title compound* (174 mg, 82%). LCMS (Method 4): Rt 3.22 min, m/z 667 [MH⁺].

c. Methanesulfonic acid 2-[3-tert-butyl-5-(3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-[1,4']bipyrzoly-1'-yl]-ethyl ester (Intermediate 124c)



5

To an ice-bath cooled solution of Intermediate 124b (174 mg, 0.26 mmol) in DCM (3 mL) was added DIPEA (182 μ L, 1.0 mmol) followed by methanesulfonyl chloride (41 μ L, 0.52 mmol). The reaction mixture was stirred for 1 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.47 min, m/z 746 [MH^+].

10

d. 1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrzoly-5-yl]-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 124)


15

To a solution of Intermediate 124c (0.13 mmol) in THF (2.5 mL) was added DIPEA (90 μ L, 0.52 mmol) and morpholine (45 μ L, 0.52 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM, followed by MDAP (Method 7)

20

purification, to give the *title compound* (26 mg, 26%). LCMS (Method 5): Rt 3.70 min, m/z 736 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.2 Hz), 0.58 (3H, d, J = 6.2 Hz), 1.20 (9H, s), 1.34-1.54 3H, m), 1.63-1.68 (2H, m), 1.72-1.86 (2H, m), 1.89-1.97 (1H, m), 2.01-2.07 (2H, m), 2.37 (4H, t, J = 4.4 Hz), 2.69 (2H, t, J = 6.7 Hz), 3.08-3.18 (2H, m), 3.49 (4H, t, J = 4.6 Hz), 4.19 (2H, t, J = 6.7 Hz), 4.80 (1H, dt, J = 8.6, 5.6 Hz), 5.49 (1H, t, J = 4.3 Hz), 6.22 (1H, s), 7.11 (1H, d, J = 8.6 Hz), 7.16 (1H, dd, J = 9.8, 2.2 Hz), 7.19-7.25 (1H, m), 7.28-7.33 (3H, m), 7.59 (1H, s), 7.62 (1H, d, J = 9.6 Hz), 7.84 (1H, d, J = 1.9 Hz), 7.96 (1H, s), 8.02 (1H, s), 8.14 (0.6H, br s).

Example 125




1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea

To a solution of Intermediate 124c (0.13 mmol) in THF (2.5 mL) was added dimethylamine (2M in MeOH, 1.3 mL, 2.6 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 2-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (26 mg, 29%). LCMS (Method 5): Rt 3.68 min, m/z 694 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.2 Hz), 0.58 (3H, d, J = 6.2 Hz), 1.20 (9H, s), 1.34-1.54 3H, m), 1.63-1.68 (2H, m), 1.72-1.86 (2H, m),


1.89-1.97 (1H, m), 2.01-2.07 (2H, m), 2.23 (6H, br s), 2.77 (2H, br s), 3.08-3.18 (2H, m), 4.22 (2H, br s), 4.80 (1H, dt, J = 8.6, 5.6 Hz), 5.49 (1H, t, J = 4.3 Hz), 6.22 (1H, s), 7.11 (1H, d, J = 8.6 Hz), 7.16 (1H, dd, J = 9.8, 2.2 Hz), 7.19-7.25 (1H, m), 7.28-7.33 (3H, m), 7.60 (1H, s), 7.62 (1H, d, J = 9.6 Hz), 7.84 (1H, d, J = 1.9 Hz), 7.98 (1H, s), 8.02 (1H, s).

Example 126

1-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea



a. (5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 126a)




A solution of 5-tert-butyl-2-methyl-2H-pyrazol-3-ylamine (0.5 g, 3.26 mmol) in EtOAc (10 mL) was treated with aqueous NaOH (1M, 5.87 mmol), followed by 2,2,2-trichloroethyl chloroformate (0.54 mL, 3.92 mmol) and the reaction mixture was stirred at RT for 1 h. The mixture was partitioned between EtOAc (10 mL) and water (2 x 10 mL). The organic layer was dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by FCC, using 0-100% EtOAc in cyclohexane to afford the *title compound* as a pale orange gum (0.915 g, 86%). LCMS (Method 3): Rt 3.88 min, m/z 328/330 [MH⁺].

b. 1-(5-tert-Butyl-2-methyl-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(S)-2-methyl-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 126)


A solution of intermediate 126a (70 mg, 0.21 mmol), intermediate 81d (80 mg, 0.21 mmol) and DIPEA (55 μ L, 0.32 mmol) in dioxane (1 mL) was heated at 70°C for 18 h. The mixture was triturated with diethyl ether (1 mL) and the solid obtained was purified by MDAP (Method 7) to afford the *title compound* as a glassy solid (25 mg, 21%). LCMS (Method 5): Rt 4.19 min, m/z 557.3 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.1 Hz), 1.20 (9H, s), 1.44-1.57 (2H, m), 1.61-1.73 (2H, m), 1.74-1.87 (2H, m), 1.87-2.03 (2H, m), 2.03-2.22 (2H, m), 2.87-2.95 (1H, m), 3.13-3.20 (1H, m), 3.39 (1H, m, obscured by water), 3.57 (3H, s), 4.83-4.91 (1H, m), 5.54 (1H, t, J = 4.0 Hz), 6.02 (1H, s), 6.93 (1H, d, J = 8.9 Hz), 7.22 (1H, dd, J = 9.8 Hz), 7.26-7.32 (1H, m), 7.35-7.42 (3H, m), 7.65 (1H, dd, J = 9.7, 1.0 Hz), 7.70-7.73 (1H, m), 8.28 (1H, s).

Example 127

1-[5-tert-Butyl-2-(2-dimethylamino-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-[(S)-2-methyl-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt



a. Methanesulfonic acid 2-[3-tert-butyl-5-(3-{{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-ethyl ester (Intermediate 127a)



5

To a solution of Example 110 (176 mg, 0.300 mmol) and DIPEA (0.157 mL, 0.900 mmol) in DCM (5 mL) at 0°C was added methanesulfonyl chloride (0.047 mL, 0.600 mmol) and the resulting orange solution stirred at 0°C for 30 min. Water (2 mL) and sat. aq. NaHCO₃ solution (2 mL) were added and the mixture shaken. The aqueous was extracted with DCM (5 mL) then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a yellow gum (199 mg, 100%). LCMS (Method 3): Rt 3.59 min, m/z 665 [MH⁺].


b. 1-[5-tert-Butyl-2-(2-dimethylamino-ethyl)-2H-pyrazol-3-yl]-3-{{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt. (Example 127)

A yellow solution of Intermediate 127a (0.100 mmol) and dimethylamine (2M in THF, 1.0 mL, 2.0 mmol) in THF (1 mL) was stirred in a sealed vial at 60°C for 16 h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL), applied to an SCX-2 cartridge (5 g) and washed with MeOH (15 mL). The product was eluted with 2M NH₃ in MeOH (15 mL); concentration *in vacuo* left a yellow film. MDAP (Method 7) gave the *title compound* as an off-white solid (42 mg, 62%). LCMS (Method 5): Rt 3.43 min, m/z 614.4 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.21 (9H, s), 1.46-1.56 (2H, m), 1.63-1.72 (2H, m), 1.76-1.85 (2H, m), 1.92-2.01 (2H, m), 2.03-2.11 (1H, m), 2.15-2.19 (1H,

25

m), 2.21 (6H, s), 2.61 (2H, t, $J = 6.9$ Hz), 2.91 (1H, ddd, $J = 12.2, 9.2, 4.0$ Hz), 3.17 (1H, dt, $J = 12.2, 4.3$ Hz), 3.28-3.36 (1H, m), 3.99 (2H, t, $J = 6.9$ Hz), 4.87 (1H, td, $J = 8.5, 6.2$ Hz), 5.53 (1H, t, $J = 4.1$ Hz), 6.03 (1H, s), 6.97 (1H, d, $J = 8.7$ Hz), 7.21 (1H, dd, $J = 9.8, 2.2$ Hz), 7.29 (1H, td, $J = 7.3, 2.0$ Hz), 7.34-7.41 (3H, m), 7.65 (1H, dd, $J = 9.8, 0.8$ Hz), 7.71 (1H, d, $J = 2.1$ Hz), 8.17 (1.4H, s), 8.48 (1H, s).

Example 128




10 **1-[5-tert-Butyl-2-(2-piperidin-1-yl-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt**

A yellow solution of Intermediate 127a (0.100 mmol) and piperidine (0.049 mL, 0.50 mmol) in dry DMF (2 mL) was stirred at 75°C for 2.5 h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL), applied to an SCX-2 cartridge (2 g) and washed with MeOH (20 mL). The product was eluted with 2M NH₃ in MeOH (15 mL); concentration *in vacuo* left a yellow solid. MDAP (Method 7) gave the *title compound* as an off-white solid (37 mg, 52%). LCMS (Method 5): Rt 3.68 min, m/z 654.5 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, $J = 6.3$ Hz), 1.20 (9H, s), 1.33-1.38 (2H, m), 1.45-1.55 (6H, m), 1.63-1.72 (2H, m), 1.75-1.86 (2H, m), 1.89-2.02 (2H, m), 2.04-2.12 (1H, m), 2.14-2.21 (1H, m), 2.40 (4H, t, $J = 4.7$ Hz), 2.61 (2H, t, $J = 7.1$ Hz), 2.91 (1H, ddd, $J = 12.2, 9.2, 4.2$ Hz), 3.16 (1H, dt, $J = 12.0, 4.3$ Hz), 3.32 (1H, m), 3.99 (2H, t, $J = 7.1$ Hz), 4.87 (1H, td, $J = 8.5, 5.7$ Hz), 5.54 (1H, t, $J = 4.3$ Hz), 6.04 (1H, s), 6.95 (1H, d, $J = 8.6$ Hz), 7.22 (1H, dd, $J = 9.8, 2.2$ Hz), 7.29 (1H, td, $J = 7.2, 1.8$ Hz),

7.34-7.41 (3H, m), 7.65 (1H, dd, J = 9.8, 0.8 Hz), 7.71 (1H, d, J = 2.1 Hz), 8.16 (1.4H, s), 8.29 (1H, s).

Example 129




1-{5-tert-Butyl-2-[2-(4-methyl-piperazin-1-yl)-ethyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A yellow solution of Intermediate 127a (0.100 mmol) and N-methyl
10 piperazine (0.056 mL, 0.50 mmol) in dry DMF (2 mL) was stirred at 75°C for 2.5
h. The cooled solution was concentrated *in vacuo*, redissolved in MeOH (2 mL),
applied to an SCX-2 cartridge (5 g) and washed with MeOH (20 mL). The product
was eluted with 2M NH₃ in MeOH (15 mL); concentration *in vacuo* left an orange
solid. MDAP (Method 7) gave the *title compound* as an off-white solid (12 mg,
15 17%). LCMS (Method 5): Rt 3.36 min, m/z 669.5 [MH⁺]. ¹H NMR (400 MHz, d₆-
DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.20 (9H, s), 1.49-1.56 (2H, m), 1.63-1.72 (2H,
m), 1.75-1.85 (2H, m), 1.91-2.02 (2H, m), 2.04-2.10 (1H, m), 2.12 (3H, s), 2.14-
2.21 (1H, m), 2.29 (4H, br s), 2.40 (4H, br s), 2.60 (2H, t, J = 7.1 Hz), 2.91 (1H,
ddd, J = 12.2, 9.0, 4.0 Hz), 3.17 (1H, dt, J = 12.0, 4.3 Hz), 3.28-3.36 (1H, m), 3.97
20 (2H, t, J = 7.1 Hz), 4.87 (1H, td, J = 8.5, 5.7 Hz), 5.54 (1H, t, J = 4.3 Hz), 6.04
(1H, s), 6.95 (1H, d, J = 8.6 Hz), 7.22 (1H, dd, J = 9.8, 2.1 Hz), 7.29 (1H, td, J =
7.2, 1.7 Hz), 7.34-7.41 (3H, m), 7.65 (1H, dd, J = 9.8, 0.8 Hz), 7.71 (1H, d, J = 2.1
Hz), 8.20 (1H, s), 8.27 (1H, s).


Example 130

**1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**



5

**a. 1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
 a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate
 130a)**




10

A mixture of Intermediate 96c (150 mg, 0.38 mmol), Intermediate 29c (161 mg, 0.38 mmol) and DIPEA (133 μ L, 0.77 mmol) in dioxane (4 mL) was stirred at 80 $^{\circ}$ C for 7 hours. After cooling, the reaction mixture was partitioned between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 1-10% MeOH in DCM to afford the *title compound* (364 mg, 95%) as a glassy white solid (%). LCMS (Method 4): Rt 3.50 min, m/z 663 [MH^+].

15

20

b. Methanesulfonic acid 3-[3-tert-butyl-5-(3-{{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzyl ester (Intermediate 130b)



5

To an ice-bath cooled solution of Intermediate 130a (175 mg, 0.26 mmol) in DCM (2.5mL) was added DIPEA (184 μ L, 1.0 mmol) followed by methanesulfonyl chloride (41 μ L, 0.52 mmol). The reaction mixture was stirred for 1 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.72 min, m/z 741 [MH^+].

10

c. 1-[5-tert-Butyl-2-(3-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-{{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 130)

15


To a solution of Intermediate 130b (0.17 mmol) in THF (2 mL) was added DIPEA (122 μ L, 0.70 mmol) and morpholine (62 μ L, 0.70 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM, followed by MDAP (Method 7)

20

purification, to give the *title compound* (25 mg, 19%). LCMS (Method 5): Rt 3.75 min, m/z 732 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.2 Hz), 0.58 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.35-1.53 (3H, m), 1.63-1.68 (2H, m), 1.72-1.86 (2H, m), 1.86-1.92 (1H, m), 2.01-2.07 (2H, m), 2.32 (4H, t, J = 4.1 Hz), 3.08-3.20 (2H, m), 3.46 (2H, s), 3.50 (4H, t, J = 4.5 Hz), 4.78 (1H, td, J = 8.6, 5.9 Hz), 5.47 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 6.98 (1H, d, J = 8.6 Hz), 7.15 (1H, dd, J = 9.8, 2.1 Hz), 7.17-7.23 (2H, m), 7.25-7.31 (3H, m), 7.32-7.44 (3H, m), 7.61 (1H, d, J = 9.7 Hz), 7.83 (1H, d, J = 1.8 Hz), 8.05 (1H, s), 8.12 (1.6H, s).


Example 131

10 **1-{5-tert-Butyl-2-[3-(2-morpholin-4-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**



a. **2-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl-phenol)]-ethanol**


15 **(Intermediate 131a)**



A mixture of 5-tert-butyl-2H-pyrazol-3-ylamine (629 mg, 4.52 mmol), 2-(3-bromophenyl)-ethanol (1g, 4.52 mmol), copper (I) iodide (43 mg, 0.23 mmol), (1S,2S)-N,N'-dimethyl cyclohexane-1,2-diamine (129 mg, 0.90 mmol) and potassium carbonate (1.31 g, 9.50 mmol) in toluene (8 mL) was de-gassed and flushed with argon (3x). The reaction mixture was then treated with

microwave irradiation, at 150°C for 3 h and then at 160°C for 16 h. The mixture was diluted with EtOAc (10 mL) and washed with water (10 mL). The aqueous layer was extracted with a further 10 mL of EtOAc and the combined organic layers were washed with brine (10 mL), dried (MgSO₄), filtered and concentrated *in vacuo*. The residue obtained was purified by FCC, using 0-100% EtOAc in cyclohexane to afford the *title compound* as a brown gum (364 mg, 31%). LCMS (Method 3): Rt 0.43/2.00/2.24 min, m/z 260 [MH⁺].

b. {5-tert-Butyl-2-[3-[2-(hydroxyethyl)-phenyl]-2H-pyrazol-3-yl]}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 131b)




10

A solution of Intermediate 131a (364 mg, 1.40 mmol) in EtOAc (5 mL) was treated with aqueous NaOH (1M, 2.53 mmol), followed by 2,2,2-trichloroethyl chloroformate (231 µL, 1.68 mmol) and the reaction mixture was stirred at RT for 1 h. The mixture was partitioned between EtOAc (5 mL) and water (5 mL). The layers were separated and the aqueous layer was extracted with a further 5 mL EtOAc. The combined organic layers were dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by FCC, using 0-50% EtOAc in cyclohexane to afford the *title compound* as a red gum (516 mg, 85%). LCMS (Method 3): Rt 4.03 min, m/z 434/436 [MH⁺].

20


25

c. 1-{5-tert-Butyl-2-[3-(2-(hydroxyethyl)-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 131c)



5 A mixture of Intermediate 131b (328 mg, 0.75 mmol), Intermediate 81d (285 mg, 0.75 mmol) and DIPEA (197 μ L, 1.13 mmol) in dioxane (4.5 mL) was heated at 70°C for 20 h. The reaction mixture was cooled to RT, diluted with DCM (10 mL) and washed with water (2 x 10 mL). The organic layer was passed through a hydrophobic frit and concentrated *in vacuo*. The residue was purified by
 10 FCC, using 0-10% [2M NH₃ in MeOH] in DCM to afford the *title compound* as a beige solid (358 mg, 72%). LCMS (Method 3): Rt 3.67 min, m/z 663 [MH⁺].

d. Methanesulfonic acid 2-{3-[3-tert-butyl-5-(3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenyl}-ethyl ester
 15 (Intermediate 131d)




To a yellow solution of Intermediate 131c (358 mg, 0.540 mmol) and DIPEA (0.28 mL, 1.6 mmol) in DCM at 0°C was added mesyl chloride (0.084 mL, 1.1 mmol) dropwise over 30 seconds, then the solution stirred at 0°C for 30 min.
 20 Water (10 mL) and sat. aq. NaHCO₃ (10 mL) were added and the mixture shaken.

The aqueous was extracted with DCM (10 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. LCMS (Method 3): Rt 3.91 min, m/z 741 [MH⁺].

e. 1-{5-tert-Butyl-2-[3-(2-morpholin-4-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 131)

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and morpholine (25.9 μ L, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (12 mg, 27%). LCMS (Method 5): Rt 3.64 min, m/z 732.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.2 Hz), 1.28 (9H, s), 1.50 (2H, m), 1.66 (2H, m), 1.76-2.18 (6H, overlapped m), 2.40 (4H, m), 2.53 (m, obscured by solvent), 2.80 (2H, m), 2.90 (1H, m), 3.15 (1H, m, obscured by solvent), 3.31 (m, obscured by solvent), 3.54 (4H, m), 4.82 (1H, m), 5.51 (1H, t, J = 4.0 Hz), 6.32 (1H, s), 7.09 (1H, d, J = 8.7 Hz), 7.19 (1H, dd, J = 9.8, 2.4 Hz), 7.24-7.44 (8H, m), 7.63 (1H, m), 7.69 (1H, m), 8.10 (1H, s), 8.20 (1H, s).

Example 132




1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and pyrrolidine (24.8 μ L, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The

mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (15 mg, 35%). LCMS (Method 5): Rt 3.70 min, m/z 716.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.6 Hz), 1.28 (9H, s), 1.50 (2H, m), 1.61-1.72 (6H, m), 1.76-2.18 (6H, overlapped m), 2.53 (m, obscured by solvent), 2.72 (2H, m), 2.82 (2H, m), 2.90 (1H, m), 3.16 (1H, m, obscured by solvent), 3.31 (m, obscured by solvent), 4.82 (1H, m), 5.52 (1H, t, J = 4.2 Hz), 6.33 (1H, s), 7.10 (1H, d, J = 8.4 Hz), 7.19 (1H, dd, J = 10.0, 2.0 Hz), 7.24-7.44 (8H, m), 7.64 (1H, m), 7.69 (1H, m), 8.14 (1H, s), 8.19 (1H, s).

Example 133




1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethyl]-phenyl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and N-ethylmethylamine (25.8 μ L, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (10 mg, 24%). LCMS (Method 5): Rt 3.68 min, m/z 704.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.2 Hz), 0.96 (3H, t, J = 7.5 Hz), 1.28 (9H, s), 1.50 (2H, m), 1.66 (2H, m), 1.76-2.17 (6H, overlapped m), 2.20 (3H, s), 2.42 (2H, q, J = 7.5 Hz), 2.60 (2H, m), 2.78 (2H, m), 2.90 (1H, m), 3.16 (1H, m, obscured by solvent), 3.31 (m, obscured by solvent), 4.82 (1H, m), 5.52 (1H, t, J = 4.4 Hz), 6.33 (1H, s), 7.10 (1H, d, J = 8.5 Hz), 7.19 (1H, dd, J = 9.7, 2.2 Hz), 7.24-7.43 (8H, m), 7.64


(1H, d, J = 9.6 Hz), 7.69 (1H, m), 8.12 (1H, s), 8.20 (1H, s).

Example 134




**1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethyl)-phenyl]-2H-pyrazol-3-yl}-
3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt**

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and dimethylamine solution (2M in THF, 0.6 mL, 1.2 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (18 mg, 41%). LCMS (Method 5): Rt 3.63 min, m/z 690.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.47-1.55 (2H, m), 1.63-1.71 (2H, m), 1.75-1.97 (4H, m), 2.00-2.16 (2H, m), 2.19 (6H, s), 2.53 (2H, t, J = 8.1 Hz), 2.78 (2H, t, J = 7.6 Hz), 2.90 (1H, ddd, J = 12.2, 9.0, 4.0 Hz), 3.16 (1H, dt, J = 12.1, 4.2 Hz), 3.31 (1H, m), 4.82 (1H, td, J = 8.6, 5.5 Hz), 5.52 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.24-7.42 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, d, J = 2.1 Hz), 8.12 (1H, s), 8.19 (1H, s).

Example 135

**1-{5-tert-Butyl-2-[3-(2-piperidin-1-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-
3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
5 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**


A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and piperidine (30 μ L, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (26 mg, 55%). LCMS (Method 10 5): Rt 3.75 min, m/z 730.7 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.31-1.37 (2H, m), 1.43-1.55 (6H, m), 1.62-1.71 (2H, m), 1.75-1.97 (4H, m), 2.00-2.17 (2H, m), 2.41 (4H, t, J = 4.4 Hz), 2.52-2.56 (2H, m), 2.79 (2H, t, J = 7.9 Hz), 2.90 (1H, ddd, J = 12.1, 9.1, 4.0 Hz), 3.16 (1H, dt, J = 11.8, 4.1 Hz), 3.28-3.35 (1H, m), 4.82 (1H, td, J = 8.6, 5.5 Hz), 5.52 (1H, t, J = 4.3 15 Hz), 6.32 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.9, 2.1 Hz), 7.24-7.42 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, d, J = 2.1 Hz), 8.11 (1H, s), 8.19 (1.4H, s).

Example 136

1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethyl]-phenyl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and N-methylpiperazine (33 μ L, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (16 mg, 33%). LCMS (Method 5): Rt 3.54 min, m/z 745.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.47-1.55 (2H, m), 1.63-1.71 (2H, m), 1.75-1.97 (4H, m), 2.00-2.08 (1H, m), 2.11 (3H, s), 2.12-2.17 (1H, m), 2.29 (4H, br s), 2.42 (4H, br s), 2.51-2.55 (2H, m), 2.76-2.80 (2H, m), 2.90 (1H, ddd, J = 12.1, 9.1, 4.1 Hz), 3.16 (1H, dt, J = 11.9, 4.2 Hz), 3.28-3.34 (1H, m), 4.82 (1H, td, J = 8.6, 5.6 Hz), 5.52 (1H, t, J = 4.3 Hz), 6.32 (1H, s), 7.08 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.24-7.42 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, d, J = 2.1 Hz), 8.10 (1H, s), 8.19 (1.4H, s).

Example 137




1-{5-tert-Butyl-2-[3-(2-[1,4]oxazepan-4-yl-ethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and homomorpholine (30.4 mg, 0.30 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified

by MDAP (Method 7). The *title product* was isolated as an off-white solid (15 mg, 31%). LCMS (Method 5): Rt 3.66 min, m/z 746.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.47-1.55 (2H, m), 1.63-1.71 (2H, m), 1.73-1.97 (6H, m), 2.00-2.17 (2H, m), 2.66-2.81 (8H, m), 2.90 (1H, ddd, J = 12.2, 9.0, 4.0 Hz), 3.16 (1H, dt, J = 12.2, 4.4 Hz), 3.28-3.34 (1H, m), 3.56-3.59 (2H, m), 3.63 (2H, t, J = 6.0 Hz), 4.82 (1H, td, J = 8.6, 5.5 Hz), 5.52 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.08 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.24-7.42 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, dd, J = 2.1, 0.9 Hz), 8.09 (1H, s), 8.18 (1.3H, s).

10 **Example 138**




1-(5-tert-Butyl-2-{3-[2-(4-methyl-[1,4]diazepan-1-yl)-ethyl]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

15 A solution of Intermediate 131d (44.5 mg, 0.06 mmol) and N-methylhomopiperazine (34.3 mg, 0.300 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give an off-white solid (22 mg). Prep HPLC (Gemini C18, 20-60% MeCN in water, 0.1% HCO₂H, 20 min) and concentration
20 of the desired fractions removed the MeCN. The aqueous was washed with DCM (30 mL), then basified with aq. NaOH solution (1M, 0.5 mL) and extracted with DCM (2 × 10 mL). The combined organics were passed through a hydrophobic


frit, then HCO₂H (0.01 mL) added and the mixture concentrated *in vacuo* to leave the *title compound* as a clear film that became a solid on standing (10.6 mg, 23%). LCMS (Method 5): Rt 3.19 min, m/z 759.7 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.48-1.55 (2H, m), 1.64-1.74 (4H, m), 1.77-1.96 (4H, m), 2.00-2.16 (2H, m), 2.29 (3H, s), 2.58-2.62 (4H, m), 2.69-2.80 (8H, m), 2.91 (1H, ddd, J = 12.4, 9.3, 4.3 Hz), 3.16 (1H, dt, J = 11.9, 4.3 Hz), 3.28-3.34 (1H, m), 4.82 (1H, td, J = 9.0, 5.7 Hz), 5.52 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.24-7.42 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, dd, J = 2.2, 0.9 Hz), 8.10 (1H, s), 8.19 (1.4H, s).

Example 139

1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea



a. (*S*)-3-Methyl-morpholine-4-carboxylic acid *N'*-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 139a)




Pyridine (1.60 mL, 19.8 mmol) was added dropwise to an ice cold suspension of triphosgene (2.94 g, 9.90 mmol) in DCM (20.0 mL). (*S*)-3-Methylmorpholine (1.00 mL, 9.90 mmol) was added and the reaction stirred for 3 hours, then quenched by dropwise addition of HCl (1M aqueous, 20 mL). The mixture was extracted into DCM (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo* to give (*S*)-3-

methylmorpholine-carbamoylchloride (1.25 g, 77%). The product was used in the next reaction without purification.

(S)-3-Methylmorpholine-carbamoylchloride (766 mg, 4.33 mmol) was dissolved in DCM (35.0 mL) and DIPEA (1.03 mL, 5390 mmol) was added followed by (5-fluoro-pyridin-2-yl)-hydrazine (WO2010022076, 500 mg, 3.94 mmol) and the reaction heated to 45°C overnight. The reaction was cooled and quenched into water. The mixture was extracted into DCM (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC using 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* (574 mg, 57%). ¹H NMR (300 MHz, CDCl₃): 1.36 (3H, d, J = 6.9 Hz), 3.30 (1H, td, J = 12.5, 3.9 Hz), 3.47-3.74 (4H, m), 3.89-4.03 (2H, m), 6.41 (1H, br s), 6.45 (1H, br s), 6.73 (1H, dd, J = 9.0, 3.5 Hz), 7.29 (1H, ddd, J = 9.0, 7.9, 2.9 Hz), 8.04 (1H, d, J = 2.9 Hz).


b. 6-Fluoro-3-((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 139b)



To an ice cold solution of Intermediate 139a (574 mg, 2.26 mmol) in THF (23.0 mL) was added sequentially triphenylphosphine (1.18 g, 4.52 mmol), triethylamine (1.26 mL, 9.04 mmol) and hexachloroethane (1.07 g, 4.52 mmol). The cooling bath was removed and the reaction was heated to 55°C overnight. The reaction was cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 x). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was taken up in MeOH and loaded onto an SCX-2 cartridge, which was washed with MeOH and eluted with 2M NH₃ in MeOH. The residue was concentrated *in vacuo* and then re-submitted to the above reaction conditions overnight. The reaction was

cooled and partitioned between EtOAc and water. The aqueous layer was then extracted with EtOAc (3 ×). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated *in vacuo*. The residue was purified by FCC using 0-10% [2M NH₃ in MeOH] in DCM to give the *title compound* (154 mg, 29%). ¹H NMR (300 MHz, CDCl₃): 0.94 (3H, d, J = 6.3 Hz), 3.13 (1H, dt, J = 12.2, 3.1 Hz), 3.32 (1H, ddd, J = 12.4, 9.5, 3.2 Hz), 3.49 (1H, dd, J = 11.1, 8.6 Hz), 3.56-3.68 (1H, m), 3.84 (1H, ddd, J = 12.1, 9.4, 2.7 Hz), 3.92-4.01 (2H, m), 7.16 (1H, ddd, J = 9.9, 7.5, 2.3 Hz), 7.68 (1H, ddd, J = 10.0, 4.8, 0.8 Hz), 7.85 (1H, m).

c. **(1*S*,4*R*)-4-[3-((*S*)-3-Methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine** (Intermediate 139c)



To a suspension of sodium hydride (60% in mineral oil, 187 mg, 4.68 mmol) in DMF (6.0 mL) was added Intermediate A (191 mg, 1.17 mmol) and the reaction stirred for 20 min. Intermediate 139b (276 mg, 1.17 mmol) was added in DMF (2.00 mL) and the reaction heated to 60°C for 1 h. The reaction was cooled and quenched by dropwise addition of methanol, before being diluted with methanol and loaded onto an SCX-2 cartridge, which was washed with MeOH. The product was eluted with 2M NH₃ in MeOH. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to give the *title compound* (250 mg, 56%). LCMS (Method 4): Rt 1.83, m/z 380 [MH⁺].

d. 1-[5-tert-Butyl-2-(2-hydroxy-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 139)


A mixture of Intermediate 139c (125 mg, 0.33 mmol), Intermediate
5 110b (115 mg, 0.33 mmol) and DIPEA (115 μ L, 0.66 mmol) in dioxane (3 mL)
was stirred at 80°C overnight. After cooling, the reaction mixture was partitioned
between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and
the combined organic layers were washed with brine, dried (MgSO₄) and
concentrated *in vacuo*. The resultant residue was purified by FCC, using a gradient
10 of 0-10% MeOH in DCM, to afford the *title compound* (94 mg, 48%). A sample of
this (40.0 mg) was further purified by MDAP (Method 7) purification to afford the
title compound (20 mg) as a white solid. LCMS (Method 5): Rt 3.64 min, m/z 589
[MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.83 (3H, d, J = 5.6 Hz), 1.16 (9H, s),
1.83-1.96 (2H, m), 1.97-2.07 (1H, m), 2.08-2.17 (1H, m), 2.95 (1H, ddd, J = 12.0,
15 7.2, 4.9 Hz), 3.17 (1H, dt, J = 12.2, 3.3 Hz), 3.36-3.44 (2H, m), 3.62 (2H, q, J = 5.5
Hz), 3.72-3.80 (2H, m), 3.81-3.88 (1H, m), 3.88-3.92 (2H, t, J = 6.0 Hz), 4.81 (1H,
td, J = 8.5, 6.1 Hz), 4.96 (1H, t, J = 5.0 Hz), 5.52 (1H, t, J = 4.2 Hz), 6.02 (1H, s),
7.05 (1H, d, J = 8.6 Hz), 7.18 (1H, dd, J = 9.9, 2.1 Hz), 7.22-7.27 (1H, m), 7.30-
7.37 (3H, m), 7.62 (1H, d, J = 9.9 Hz), 7.79 (1H, d, J = 1.9 Hz), 8.16 (1H, s).

20


25

Example 140

1-(5-tert-Butyl-2-{2-[(2-dimethylamino-ethyl)-methyl-amino]-pyrimidin-4-yl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. 5-tert-Butyl-2-(2-chloro-pyrimidin-4-yl)-2H-pyrazol-3-ylamine (Intermediate 140a)




A mixture of 2,4-dichloropyrimidine (1.0 g, 6.7 mmol), 3-(tert-butyl)-1H-pyrazol-5-amine (1.03 g, 7.4 mmol) and sodium carbonate (1.42 g, 13.4 mmol) in 1,4-dioxane was treated with 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (xantphos) (58 mg, 0.10 mmol) followed by tris(dibenzylideneacetone)dipalladium (31 mg, 0.03 mmol). The mixture was degassed then heated to 70°C under N₂ for 5h. After cooling to room temperature the mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was partitioned between EtOAc and water and the phases were separated. The aqueous layer was extracted with EtOAc (x2) and the combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-15% EtOAc in cyclohexane, to give the *title compound* (0.59 g, 35%). LCMS (Method 3): Rt 3.97 min, m/z 252.2 [MH⁺].

b. N-[4-(5-Amino-3-tert-butyl-pyrazol-1-yl)-pyrimidin-2-yl]-N,N',N'-

trimethyl-ethane-1,2-diamine (Intermediate 140b)

A solution of Intermediate 140a (250 mg, 0.99 mmol) and N,N,N'-trimethylethylenediamine (0.32 mL, 2.48 mmol) in IPA (2.5 mL) was heated to
5 120°C for 10 mins under microwave irradiation. The cooled reaction mixture was concentrated *in vacuo* and the residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was extracted with EtOAc (x2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to give the *title compound* (0.31 g, quant.) as a golden oil
10 that was used in the next step without purification. LCMS (Method 3): Rt 2.51 min, m/z 318.3 [MH⁺].

c. (5-tert-Butyl-2-{2-[(2-dimethylamino-ethyl)-methyl-amino]-pyrimidin-4-yl}-2H-pyrazol-3-yl)-carbamic acid phenyl ester (Intermediate 140c)



15

A solution of Intermediate 140b (129 mg, 0.41 mmol) in DCM (4.1 mL) was treated with pyridine (0.046 mL, 0.57 mmol) and the mixture was cooled 0°C. Phenyl chloroformate (0.066 mL, 0.53 mmol) was added and the resulting mixture was stirred at RT for 16 h. The mixture was cooled to 0°C and treated with
20 pyridine (0.013 mL, 0.16 mmol) then phenyl chloroformate (0.015 mL, 0.12 mmol) and stirred at RT for a further 1.75 h. Once again the mixture was cooled to 0°C and treated with pyridine (0.013 mL, 0.16 mmol) then phenyl chloroformate

(0.015 mL, 0.12 mmol) and stirred at RT for 1.75 h. The mixture was diluted with water and the phases were separated. The aqueous layer was extracted with DCM (x 2) and the combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to give a yellow gum (236 mg). Trituration with ether and
5 filtration of the resulting solid gave the *title compound* as a white solid (75 mg, 39%). LCMS (Method 3): Rt 3.21 min, m/z 438.4 [MH⁺].


**d. 1-(5-tert-Butyl-2-{2-[(2-dimethylamino-ethyl)-methyl-amino]-pyrimidin-4-yl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
10 urea (Example 140)**

A solution of Intermediate 81d (25 mg, 0.066 mmol) in 1,4-dioxane (0.7 mL) was treated with Intermediate 140c (32 mg, 0.073 mmol) and DIPEA (0.014 mL, 0.083 mmol) and the mixture was stirred at 70°C for 18 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and
15 water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. FCC, using 0-8% MeOH in DCM, gave a colourless glass (27 mg). The glass was purified further by HPLC (Phenomenex Gemini C18 column, 5-95% MeCN in H₂O, 0.1% HCO₂H) over 30 mins.
20 Concentration of the fractions *in vacuo* gave a colourless glass that was dissolved in ether. A solid was precipitated out of solution by addition of cyclohexane. The solid was filtered off to give the *title compound* as a white solid (8 mg, 17%). LCMS (Method 5): Rt 4.07 min, m/z 721.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO, 353K): 0.92 (3H, d, J = 6.4 Hz), 1.29 (9H, s), 1.49-1.58 (2H, m), 1.65-1.72 (2H, m), 1.77-1.86 (2H, m), 1.99 (6H, s), 2.00-2.14 (4H, m), 2.20-2.29 (1H, m), 3.12-3.19 (5H, m), 3.30-3.39 (1H, m), 3.58-3.72 (2H, m), 4.94-5.02 (1H, m), 5.42 (1H, t, J = 3.9 Hz), 6.55 (1H, s), 7.03 (1H, d, J = 5.5 Hz), 7.15 (1H, dd, J = 9.8, 2.3 Hz), 7.22-7.27 (1H, m), 7.31-7.37 (2H, m), 7.41-7.45 (1H, m), 7.56-7.61 (2H, m),


7.70 (1H, br s), 8.35 (1H, d, $J = 5.5$ Hz), 10.18 (1H, br s).

Example 141

**1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-
5 *a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**




**a. 1-(5-tert-Butyl-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-
2H-pyrazol-3-yl)-3-{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-
[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
10 urea (Intermediate 141a)**



A mixture of Intermediate 139c (125 mg, 0.33 mmol), Intermediate 39b (177 mg, 0.33 mmol) and DIPEA (115 μ L, 0.66 mmol) in dioxane (3 mL) was stirred at 80°C overnight. After cooling, the reaction mixture was partitioned
15 between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% MeOH in DCM, to afford the *title compound* (214 mg, 85%).

LCMS (Method 4): Rt 3.65 min, m/z 765 [MH⁺].


**b. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-
{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 141b)**



5

To a solution of Intermediate 141a (214 mg, 0.28 mmol) in MeOH (3.0 mL) was added pyridinium p-toluenesulfonate (141 mg, 0.56 mmol) and the reaction mixture was heated at 60°C for 3 h. The resultant mixture was poured into water and a saturated aqueous solution of NaHCO₃ was added. The aqueous phase was
10 extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% MeOH in DCM, to give the *title compound* (132 mg, 69%). LCMS (Method 4): Rt 3.04 min, m/z 681 [MH⁺].

**c. Methanesulfonic acid 2-{3-[3-tert-butyl-5-(3-{(1*S*,4*R*)-4-[3-((*S*)-3-
15 methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-
tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenoxy}-ethyl ester (Intermediate 141c)**



To an ice-bath cooled solution of Intermediate 141b (132 mg, 0.19 mmol) in


DCM (2.0 mL) was added DIPEA (135 μ L, 0.78 mmol) followed by methanesulfonyl chloride (30 μ L, 0.39 mmol). The reaction mixture was stirred for 2 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the subsequent step without further purification. LCMS (Method 4): Rt 3.30 min, m/z 759 [MH^+].

d. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 141)

To a solution of Intermediate 141c (0.19 mmol) in THF (2.0 mL) was added dimethylamine (2M in MeOH, 1.9 mL, 3.8 mmol) and the reaction was heated to 50°C in a sealed tube overnight. The crude reaction mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (50 mg, 34%). LCMS (Method 5): Rt 3.33 min, m/z 708 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.83 (3H, d, $J = 5.7$ Hz), 1.23 (9H, s), 1.78-1.92 (2H, m), 1.94-2.04 (1H, m), 2.04-2.12 (1H, m), 2.16 (6H, s), 2.59 (2H, t, $J = 5.7$ Hz), 2.94 (1H, ddd, $J = 12.2, 7.3, 4.7$ Hz), 3.16 (1H, dt, $J = 12.2, 3.3$), 3.35-3.43 (2H, m), 3.72-3.80 (2H, m), 3.81-3.87 (1H, m), 4.04 (2H, t, $J = 5.7$ Hz), 4.77 (1H, td, $J = 8.5, 5.9$ Hz), 5.50 (1H, t, $J = 4.1$ Hz), 6.28 (1H, s), 6.90-6.94 (1H, m), 7.01-7.09 (3H, m), 7.14-7.18 (1H, dd, $J = 9.8, 2.1$ Hz), 7.20-7.26 (2H, m), 7.26-7.30 (1H, m), 7.30-7.38 (2H, m), 7.62 (1H, d, $J = 9.8$ Hz), 7.78 (1H, d, $J = 1.8$ Hz), 8.09 (1H, s), 8.12 (1.4H, s).


Example 142

1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt



5

a. Methanesulfonic acid 2-[3-tert-butyl-5-(3-{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-ethyl ester (Intermediate 142a)



10

To an ice-bath cooled solution of Example 139 (54 mg, 0.092 mmol) in DCM (2.0 mL) was added DIPEA (65 μ L, 0.37 mmol) followed by methanesulfonyl chloride (14 μ L, 0.18 mmol). The reaction mixture was stirred for 1 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.04 min, m/z 667 [MH⁺].


15

b. 1-[5-tert-Butyl-2-(2-morpholin-4-yl-ethyl)-2H-pyrazol-3-yl]-3-{(1*S*,4*R*)-4-[3-((*S*)-3-methyl-morpholin-4-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-

20

ylxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 142)

To a solution of Intermediate 142a (0.092 mmol) in THF (2 mL) was added DIPEA (64 μ L, 0.37 mmol) and morpholine (21 μ L, 0.37 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned
 5 between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (21 mg, 34%). LCMS (Method 5): Rt 3.16
 10 min, m/z 658 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.83 (3H, d, J = 5.6 Hz), 1.16 (9H, s), 1.83-1.96 (2H, m), 1.97-2.07 (1H, m), 2.08-2.17 (1H, m), 3.34 (4H, t, J = 4.5 Hz), 2.57 (2H, t, J = 7.1 Hz), 2.95 (1H, ddd, J = 12.0, 7.2, 4.9 Hz), 3.17 (1H, dt, J = 12.2, 3.3 Hz), 3.36-3.43 (2H, m), 3.50 (4H, t, J = 4.5 Hz), 3.72-3.80 (2H, m), 3.81-3.88 (1H, m), 3.92-3.98 (2H, t, J = 7.1 Hz), 4.82 (1H, td, J = 8.5, 6.1
 15 Hz), 5.52 (1H, t, J = 4.2 Hz), 5.99 (1H, s), 6.89 (1H, d, J = 8.6 Hz), 7.18 (1H, dd, J = 9.9, 2.1 Hz), 7.22-7.27 (1H, m), 7.30-7.38 (3H, m), 7.62 (1H, d, J = 9.9 Hz), 7.79 (1H, d, J = 1.9 Hz), 8.16 (1H, s), 8.16 (0.2H, s).


Example 143

20 **1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 130b (0.25 mmol) in THF (2.5 mL) was added

N-methylpiperazine (277 μ L, 2.50 mmol) and the reaction stirred at 50°C for 24 h. After cooling, the mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (37 mg, 19%). LCMS (Method 5): Rt 3.77 min, *m/z* 745 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, *J* = 6.2 Hz), 0.58 (3H, d, *J* = 6.2 Hz), 1.23 (9H, s), 1.35-1.54 (3H, m), 1.63-1.68 (2H, m), 1.72-1.86 (2H, m), 1.86-1.92 (1H, m), 2.01-2.06 (2H, m), 2.07 (3H, s), 2.26 (4H, br s), 2.34 (4H, br s), 3.08-3.20 (2H, m), 3.46 (2H, s), 4.78 (1H, td, *J* = 8.6, 5.9 Hz), 5.47 (1H, t, *J* = 4.2 Hz), 6.28 (1H, s), 7.00 (1H, d, *J* = 8.6 Hz), 7.15 (1H, dd, *J* = 9.8, 2.1 Hz), 7.18-7.25 (2H, m), 7.25-7.31 (3H, m), 7.32-7.37 (2H, m), 7.41 (1H, t, *J* = 7.6 Hz), 7.61 (1H, d, *J* = 9.7 Hz), 7.83 (1H, d, *J* = 1.8 Hz), 8.05 (1H, s), 8.12 (1.1H, s).

Example 144




**1-[5-tert-Butyl-2-(3-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 130b (0.32 mmol) in THF (3.0 mL) was added pyrrolidine (530 μ L, 6.4 mmol) and the reaction stirred at 50°C for 24 h. After cooling, the mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed

with brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (60 mg, 24%). LCMS (Method 5): Rt 3.79 min, m/z 716 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.2 Hz), 0.58 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.35-1.54 (3H, m), 1.60-1.68 (6H, m), 1.72-1.85 (2H, m), 1.86-1.92 (1H, m), 2.01-2.06 (2H, m), 2.38-2.44 (4H, m), 3.08-3.20 (2H, m), 3.58 (2H, s), 4.77 (1H, td, J = 8.6, 5.9 Hz), 5.48 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 7.00 (1H, d, J = 8.6 Hz), 7.15 (1H, dd, J = 9.8, 2.1 Hz), 7.18-7.25 (2H, m), 7.26-7.35 (4H, m), 7.37 (1H, s), 7.41 (1H, t, J = 7.6 Hz), 7.61 (1H, d, J = 9.7 Hz), 7.83 (1H, d, J = 1.8 Hz), 8.05 (1H, s), 8.13 (1.2H, s).

Example 145




1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

Pyrrolidine (36 μ L, 0.43 mmol) was added to a solution of Intermediate 96f (111 mg, 0.14 mmol) in THF (2 mL). The reaction was heated to 60°C in a sealed vial overnight. The mixture was cooled, evaporated *in vacuo* and the residue was purified by FCC, using 0-15% MeOH in DCM, then triturated with Et₂O to give the *title compound* (27 mg, 25%) as an off-white solid. LCMS (Method 5): Rt 3.87 min, m/z 746.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6.0 Hz), 0.58 (3H, d, J 6.0 Hz), 1.23 (9H, s), 1.32-1.54 (3H, m), 1.53-1.78 (7H, m), 1.79-1.95 (2H, m), 1.96-2.10 (2H, m), 2.49-2.60 (2H, m), 2.68-2.94 (2H, m), 3.06-3.20

(3H, m), 3.27 (1H, m, obscured by water), 4.03-4.14 (2H, m), 4.74-4.83 (1H, m), 5.49 (1H, t, $J = 4.0$ Hz), 6.29 (1H, s), 6.90-6.96 (1H, m), 7.01-7.09 (3H, m), 7.16 (1H, dd, $J = 9.7, 2.0$ Hz), 7.19-7.25 (2H, m), 7.26-7.32 (2H, m), 7.33-7.39 (1H, m), 7.62 (1H, d, $J = 9.7$ Hz), 7.84 (1H, m), 8.06 (1H, s).

5

Example 146

1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((2S,6R)-2,6-dimethyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea


10

1-Methylpiperazine (48 μ L, 0.43 mmol) was added to a solution of Intermediate 96f (111 mg, 0.14 mmol) in THF (2 mL). The reaction was heated to 60°C in a sealed vial overnight. 1-Methylpiperazine (48 μ L, 0.43 mmol) was added and the reaction continued for a further 24 h. The mixture was cooled, evaporated *in vacuo* and the residue was purified by FCC, using 0-15% MeOH in DCM, then triturated with Et₂O to give the *title compound* (42 mg, 38%) as an off-white solid. LCMS (Method 5): Rt 3.76 min, m/z 775.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.60 (3H, d, $J = 6.0$ Hz), 0.63 (3H, d, $J = 6.0$ Hz), 1.28 (9H, s), 1.36-1.62 (3H, m), 1.65-1.75 (2H, m), 1.76-1.99 (3H, m), 2.03-2.12 (2H, m), 2.14-2.26 (3H, m), 2.34-2.61 (5H, m, obscured by solvent), 2.63-2.74 (3H, m), 3.11-3.24 (2H, m), 3.28-3.38 (2H, m), 4.1 (2H, t, $J = 5.8$ Hz), 4.78-4.87 (1H, m), 5.50-5.56 (1H, m), 6.33 (1H, s), 6.93-7.00 (1H, m), 7.05-7.13 (3H, m), 7.20 (1H, dd, $J = 9.9, 2.4$ Hz), 7.23-7.31 (2H, m), 7.34 (2H, t, $J = 8.0$ Hz), 7.40 (1H, t, $J = 8.0$ Hz),

20

7.67 (1H, d, J = 9.9 Hz), 7.87-7.91 (1H, m), 8.11 (1H, s).

Example 147




1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-[(2S,6R)-2,6-dimethyl-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt


N-Ethylmethylamine (37 μ L, 0.43 mmol) was added to a solution of Intermediate 96f (111 mg, 0.14 mmol) in THF (2 mL). The reaction was heated to 60°C in a sealed vial overnight. *N*-Ethylmethylamine (37 μ L, 0.43 mmol) was added and the reaction continued for a further 24 h. The mixture was cooled, evaporated *in vacuo* and the residue was purified by FCC, using 0-15% MeOH in DCM, then triturated with Et₂O. Further purification by HPLC (Gemini C18 column, 10-98% MeCN in H₂O, 0.1% formic acid) followed by evaporation of the fractions and trituration with Et₂O gave the *title compound* (12 mg, 11%) as an off-white solid. LCMS (Method 5): Rt 3.85 min, m/z 734.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.55 (3H, d, J = 6 Hz), 0.58 (3H, d, J = 6 Hz), 0.89 (3H, t, J = 7.0 Hz), 1.23 (9H, m), 1.33-1.57 (3H, m), 1.62-1.70 (2H, m), 1.71-1.95 (3H, m), 1.96-2.09 (2H, m), 2.13 (3H, s), 2.35 (2H, q, J = 7.2 Hz), 2.63 (2H, t, J = 5.7 Hz), 3.06-3.20 (3H, m), 3.21-3.36 (2H, m, obscured by solvent), 4.03 (2H, t, J = 5.7 Hz), 4.74-4.83 (1H, m), 5.48 (1H, t, J = 5.5 Hz), 6.28 (1H, s), 6.88-6.94 (1H, m), 7.00-7.10 (3H, m), 7.16 (1H, dd, J = 9.8, 2.0 Hz), 7.18-7.26 (2H, m), 7.26-7.32 (2H, m), 7.35 (1H, t, J = 8.0 Hz), 7.62 (1H, d, J = 9.8 Hz), 7.81-7.86 (1H, m), 8.09 (1H, s), 8.37 (0.2H, s).

Example 148

1-{(1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea formate salt




5

a. 4-Aza-spiro[2.5]octane-4-carbonyl chloride (Intermediate 148a)

To a vigorously stirred pale yellow solution of 4-azaspiro[2.5]octane hydrochloride (ABCR, 443 mg, 3.00 mmol), pyridine (0.36 mL, 4.5 mmol) and DIPEA (0.52 mL, 3.0 mmol) in DCM (10 mL) at 0°C was added a solution of triphosgene (594 mg, 2.00 mmol) in DCM (5 mL) dropwise over 5 min. The resulting orange solution was stirred at 0°C for 30 min, and at RT for 66 h. Aq. HCl solution (1M, 15 mL) was added and the mixture stirred for 30 min. The aqueous was extracted with DCM (4 × 15 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a red-brown oil (100% yield). ¹H NMR (300 MHz, CDCl₃): ~2:1 ratio of rotamers, 0.71-0.76 (1.33H, m), 0.82 (0.67H, br s), 1.00-1.04 (1.33H, m), 1.09 (0.67H, br s), 1.53-1.59 (2H, m), 1.64-1.72 (2H, m), 1.75-1.83 (2H, m), 3.61-3.74 (2H, m).

b. 4-Aza-spiro[2.5]octane-4-carboxylic acid N'-(5-fluoro-pyridin-2-yl)-hydrazide (Intermediate 148b)




A red solution of (5-fluoro-pyridin-2-yl)-hydrazine (477 mg, 3.75 mmol), Intermediate 148a (3.00 mmol) and DIPEA (0.78 mL, 4.5 mmol) in DCM (20 mL)

20


was stirred at reflux for 40 h. To the cooled solution was added water (20 mL) and the mixture shaken. The aqueous was extracted with DCM (2×10 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo*. Flash chromatography (silica 40g, 1-5% [2M NH_3 in MeOH] in DCM) gave the
5 *title compound* as a yellow solid (587 mg, 74%). LCMS (Method 3): Rt 2.66 min, m/z 265 [MH^+].

c. 3-(4-Aza-spiro[2.5]oct-4-yl)-6-fluoro-[1,2,4]triazolo[4,3-a]pyridine (Intermediate 148c)



10 To a solution of Intermediate 148b (587 mg, 2.22 mmol), triphenyl phosphine (1.17 g, 4.44 mmol) and triethylamine (1.24 mL, 8.88 mmol) in dioxane (25 mL) at RT was added hexachloroethane (1.05 g, 4.44 mmol). The resulting opaque solution was stirred at RT for 30 min, and at reflux for 16 h. The cooled solution was filtered, and the filter-cake washed with THF (5 mL). The combined
15 organics were applied to an SCX-2 cartridge, which was washed with MeOH (100 mL). The product was eluted with 2M NH_3 in MeOH (75 mL); concentration *in vacuo* left a brown oil. Flash chromatography (silica 40 g, 1.5-4.5% MeOH in DCM) gave the *title compound* as a brown gum (82%). LCMS (Method 3): Rt 2.92 min, m/z 247 [MH^+].


20 **d. (1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate 148d)**



To an opaque brown solution of Intermediate A (371 mg, 2.28 mmol) in dry DMF (5 mL) under N_2 was added sodium hydride (60% dispersion in mineral oil,


273 mg, 6.83 mmol) and the resulting mixture stirred at RT for 30 min. (CARE: gas evolution) A solution of Intermediate 148c (448 mg, 1.82 mmol) in dry DMF (5 mL) was added and the resulting dark brown solution stirred at 60 °C for 1 h. The solution was cooled to RT, then water (0.2 mL) added (CARE: gas evolution) and the mixture concentrated *in vacuo*. The residue was redissolved in MeOH (10 mL), applied to an SCX-2 cartridge (25 g) and washed with MeOH (100 mL). The product was eluted with 2M NH₃ in MeOH (75 mL); concentration *in vacuo* left a dark brown gum. Flash chromatography (silica 40 g, 4-7% [2M NH₃ in MeOH] in DCM) gave the *title compound* as a pale brown foam (446 mg, 63%). LCMS (Method 3): Rt 2.31 min, m/z 390 [MH⁺].

e. {5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 148e)




A colourless solution of Intermediate 39b (120 mg, 0.224 mmol) and pyridinium p-toluenesulfonate (169 mg, 0.673 mmol) in MeOH (5 mL) was stirred at 40°C for 90 min. The solution was concentrated *in vacuo*, suspended in water (3 mL) and sat. aq. NaHCO₃ solution (3 mL), then extracted with DCM (2 × 5 mL). The combined organics were passed through a hydrophobic frit, concentrated *in vacuo* and azeotroped with toluene (2 × 5 mL) to leave the *title compound* as a white solid (100 mg, 99%). LCMS (Method 3): Rt 4.01 min, m/z 450, 452 [MH⁺].

f. 1-{(1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea (Intermediate 148f)



5 An orange-brown solution of Intermediate 148d (87.2 mg, 0.224 mmol), Intermediate 148e (100 mg, 0.224 mmol) and DIPEA (0.049 mL, 0.28 mmol) in dry dioxane (3 mL) was stirred at 75°C for 16 h. The cooled solution was concentrated *in vacuo*, suspended in water (5 mL) and extracted with DCM (2 × 5 mL). The combined organics were passed through a hydrophobic frit and
10 concentrated *in vacuo* to leave a brown gum. Flash chromatography (silica 12g, 4-8% MeOH in DCM) gave the *title compound* as a pale yellow solid (138 mg, 89%). LCMS (Method 3): Rt 3.57 min, m/z 691 [MH⁺].

g. Methanesulfonic acid 2-{3-[5-(3-{(1S,4R)-4-[3-(4-aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-3-tert-butyl-pyrazol-1-yl]-phenoxy}-ethyl ester (Intermediate 148g)
15



A yellow-orange solution of Intermediate 148f (137 mg, 0.198 mmol), DIPEA (0.10 mL, 0.60 mmol) and methanesulfonyl chloride (45 mg, 0.40 mmol) in DCM (5 mL) was stirred at 0°C for 30 min, and at RT for 30 min. DIPEA
20 (0.052 mL, 0.30 mmol) and MsCl (23 mg, 0.20 mmol) were added and the solution stirred at RT for 30 min. Water (2 mL) and sat. aq. NaHCO₃ solution (2 mL) were


added and the mixture shaken. The aqueous was extracted with DCM (5 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a yellow solid (100%). LCMS (Method 3): Rt 3.80 min, m/z 769 [MH⁺].

5 **h. 1-{(1S,4R)-4-[3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-3-{5-tert-butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea formate salt. (Example 148)**


A brown solution of Intermediate 148g (0.099 mmol) and dimethylamine
10 (2M in THF, 0.99 mL, 1.98 mmol) in THF (1 mL) was stirred in a sealed vial at 60°C for 16 h. The solution was decanted and concentrated. MDAP (Method 7) gave the *title compound* as a pale yellow solid (50 mg, 65%). LCMS (Method 5): Rt 3.62 min, m/z 718.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.42-0.53 (4H, m), 1.27 (9H, s), 1.53-1.61 (2H, m), 1.74-1.96 (6H, m), 1.99-2.15 (2H, m), 2.20 (6H, s), 2.63
15 (2H, t, J = 5.8 Hz), 3.22-3.29 (2H, m), 4.09 (2H, t, J = 5.8 Hz), 4.81 (1H, td, J = 8.6, 5.5 Hz), 5.48 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 6.95-6.98 (1H, m), 7.06-7.12 (3H, m), 7.14 (1H, dd, J = 9.9, 2.2 Hz), 7.25-7.42 (5H, m), 7.56 (1H, d, J = 2.1 Hz), 7.61 (1H, dd, J = 9.8, 0.8 Hz), 8.12 (1H, s), 8.17 (1.4H, s).

Example 149

20 **1-[5-tert-Butyl-2-(3-morpholin-4-yl-methyl-phenyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl}-urea formate salt**




a. 1-[5-tert-butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-{1,2,4}triazolo[4,3-a]pyridin-6-
 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 149a)



5 A solution of Intermediate 5c (1.60 g, 4.40 mmol) and Intermediate 29c
 (1.85 g, 4.40 mmol) in dioxane (20 mL) was treated with DIPEA (1.15 mL, 6.60
 mmol) and the reaction mixture was heated at 70°C for 36 h. The mixture was
 concentrated *in vacuo* and the residue was dissolved in DCM (150 mL) and
 washed with water (2 x 75 mL) and brine (75 mL). The organic layer was dried
 10 (MgSO₄), filtered and concentrated *in vacuo* and the foam obtained purified by
 FCC, using 0-5% MeOH in DCM then 0-5% [2M NH₃ in MeOH] in DCM to
 afford the *title compound* as a yellow foamy solid (0.94 g, 85%). LCMS (Method
 3): Rt 2.76 min, m/z 635 [MH⁺].

b. Methanesulfonic acid 3-[3-tert-butyl-5-(3-{(1S,4R)-4-[3-((S)-1-
 15 methyl-pyrrolidin-2-yl)-{1,2,4}triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-
 tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzyl ester (Intermediate
 149b)



A solution of Intermediate 149a (200 mg, 0.32 mmol) and DIPEA (164 μL,

0.95 mmol) in DCM (4 mL) was treated with methanesulfonyl chloride (32 μ L, 0.41 mmol) and the reaction mixture was stirred at RT for 0.5 h. The mixture was diluted with DCM (5 mL) and washed with water (10 mL) and brine (10 mL). The layers were separated and the organic layer was passed through a phase separator and concentrated *in vacuo* to afford the *title compound* as a pale yellow solid (208 mg, 92%). LCMS (Method 3): Rt 3.02 min, m/z 713 [MH⁺].


c. 1-[5-tert-Butyl-2-(3-morpholin-4-yl-methyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea formate salt (Example 149)

A mixture of Intermediate 149b (104 mg, 0.15 mmol) and morpholine (64 μ L, 0.73 mmol) in THF (2 mL) was stirred at 60 °C for 18 h. The reaction mixture was cooled to RT, diluted with DCM (5 mL) and washed with water (5 mL) and brine (5 mL). The organic layer was passed through a phase separator and concentrated *in vacuo* and the resultant residue was purified by prep. HPLC, using 5-50% CH₃CN in water, buffered with formic acid, on a Gemini C₁₈ column, over 20 mins to afford the *title compound* as an off white solid (22 mg, 21%). LCMS (Method 5): Rt 2.66 min, m/z 704.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.84-2.11 (6H, m), 2.13 (3H, s), 2.15-2.26 (2H, m), 2.31-2.40 (5H, m), 3.10-3.17 (1H, m, obscured by water), 3.51 (2H, s, obscured by water), 3.55 (4H, t, J = 4.6 Hz), 3.99 (1H, t, J = 8.0 Hz), 4.78-4.86 (1H, m), 5.39 (1H, t, J = 4.2 Hz), 6.33 (1H, s), 7.06 (1H, d, J = 8.6 Hz), 7.24-7.49 (9H, m), 7.75 (1H, d, J = 9.7 Hz), 8.10 (1H, s), 8.19 (0.6H, br s), 8.24 (1H, d, J = 1.8 Hz).

Example 150


1-(5-tert-Butyl-2-{3-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-

urea formic acid salt



a. N-(3-Bromo-phenyl)-N,N',N'-trimethyl-ethane-1,2-diamine


(Intermediate 150a)



5

A solution of N'-(3-bromo-phenyl)-N,N-dimethyl-ethane-1,2-diamine (400 mg, 1.6 mmol) was formed in 1,2-dichloroethane (10 mL). Formaldehyde (250 μ L, 37% in water, 3.3 mmol) was added and the mixture stirred for 10 minutes at RT. Sodium triacetoxyborohydride (700 mg, 3.3 mmol) was added and the mixture stirred for 2 h at RT. Further formaldehyde (250 μ L, 37% in water, 3.3 mmol) and sodium triacetoxyborohydride (700 mg, 3.3 mmol) were added and the mixture stirred at RT for 16 h. Mixture quenched with sat. NaHCO_3 (aq) and extracted with DCM. The combined organic phases were dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the title compound (200 mg, 47%). LCMS (Method 3): Rt 2.35 min, m/z 257, 259 $[\text{MH}^+]$.

b. N-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenyl]-N,N',N'-trimethyl-ethane-1,2-diamine (Intermediate 150b)




20

A solution of Intermediate 150a (200 mg, 0.78 mmol), 3-(tert-butyl)-1H-pyrazol-5-amine (130 mg, 0.93 mmol) and trans-N,N -dimethylcyclohexane

diamine (22mg, 0.16 mmol) was formed in toluene (2 mL). Potassium carbonate (226 mg, 1.6 mmol) was added and the mixture degassed by bubbling nitrogen through it. Copper (I) iodide (15 mg, 0.08 mmol) was added and the mixture was heated to 150°C for 24 h under microwave irradiation. The cooled reaction mixture was concentrated *in vacuo* and the residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was extracted with EtOAc. The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the *title compound* (50 mg, 20%). LCMS (Method 3): Rt 1.84 min, m/z 316 [MH⁺].

c. (5-tert-Butyl-2-{3-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-2H-pyrazol-3-yl)-carbamic acid phenyl ester (Intermediate 150c)



A solution of Intermediate 150b (50 mg, 0.16 mmol) in DCM (2 mL) was treated with pyridine (26 µL, 0.32 mmol) and phenyl chloroformate (32 µL, 0.25 mmol). The resulting mixture was stirred at RT for 2 h. Further pyridine (26 µL, 0.32 mmol) and phenyl chloroformate (32 µL, 0.25 mmol) were added and the mixture stirred at RT for 16 h. The mixture was diluted with water and the phases separated. The aqueous layer was extracted with DCM and the combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, to give the *title compound* (30 mg, 43%). LCMS (Method 3): Rt 2.97 min, m/z 436 [MH⁺].


d. 1-(5-tert-Butyl-2-{3-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-

urea formic acid salt (Example 150)


A solution of Intermediate 81d (26 mg, 0.07 mmol) in 1,4-dioxane (2 mL) was treated with Intermediate 150c (30 mg, 0.07 mmol) and DIPEA (24 μ L, 0.13 mmol) and the mixture was stirred at 70°C for 18 h. After cooling to RT the mixture was concentrated *in vacuo*. The residue was purified by FCC, using 0-20% MeOH in DCM, then by prep HPLC (C18 column, 10-98% MeCN in H₂O, 0.1% HCO₂H) over 20 mins. Concentration of the fractions *in vacuo* gave pale yellow glass. Trituration with Et₂O gave the *title compound* as an off white solid (20 mg, 40%). LCMS (Method 5): Rt 3.70 min, m/z 719.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.4 Hz), 1.27 (9H, s), 1.47-1.53 (2H, m), 1.64-1.71 (2H, m), 1.77-1.97 (4H, m), 2.00-2.15 (2H, m), 2.16 (3H, s), 2.39 (2H, t, J = 7.2 Hz), 2.87-2.92 (1H, m), 2.93 (3H, s), 3.13-3.21 (4H, m), 3.35-3.46 (4H, m), 4.80-4.87 (1H, m), 5.52 (1H, t, J = 4.3 Hz), 6.32 (1H, s), 6.68-6.72 (3H, m), 7.13 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.7, 2.2 Hz), 7.24-7.38 (5H, m), 7.64 (1H, d, J = 9.7 Hz), 7.69 (1H, m), 8.11 (1H, s).

Example 151

1-{5-tert-Butyl-2-[3-((R)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




a. **2-[3-((R)-2-Benzyloxy-1-methyl-ethoxy)-phenyl]-5-tert-butyl-2H-pyrazol-3-ylamine (Intermediate 151a)**



A mixture of Intermediate 95a (0.50 g, 2.2 mmol) in dry THF (17 mL) was
5 treated with triphenylphosphine (1.13 g, 4.32 mmol) and (S)-(+)-1-Benzyloxy-2-propanol (0.52 mL, 3.24 mmol). The mixture was cooled to 0°C then diisopropylazodicarboxylate (0.85 mL, 4.32 mmol) was added dropwise. After stirring at ambient temperature for 1.75 h the mixture was treated with water (0.2 mL) then applied to a pre-conditioned (with MeOH) SCX-2 cartridge. The
10 cartridge was eluted with MeOH then 2N NH₃ in MeOH. The product containing fractions were combined and concentrated *in vacuo* to a viscous brown oil. The oil was purified by FCC, using 0-30% EtOAc in cyclohexane, to give the *title compound* as a golden oil (0.37 g, 45%). LCMS (Method 3): Rt 3.71 min, m/z 380.3 [MH⁺].


15 b. **(R)-2-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenoxy]-propan-1-ol (Intermediate 151b)**



A solution of Intermediate 151a (0.37 g, 0.97 mmol) in ethanol (IMS grade, 8 mL) was treated with water (1 mL) and ammonium formate (0.61 g, 9.7 mmol).
20 The mixture was purged with N₂ then 10% Pd/C (0.31g, 0.29 mmol Pd) was added. The mixture was heated to reflux for 16 h then filtered through celite and the filtrate was concentrated *in vacuo*. The residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was extracted with


EtOAc (x2). The combined organic phase was washed with saturated aqueous sodium bicarbonate solution and brine, dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by FCC, using 0-75% EtOAc in cyclohexane to give the *title compound* (0.22 g, 79%) as a pink gum. LCMS (Method 3): Rt 2.44 min, m/z 290.3 $[\text{MH}^+]$.

c. {5-tert-Butyl-2-[3-((R)-2-hydroxy-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 151c)



A solution of Intermediate 151b (112 mg, 0.39 mmol) in EtOAc (2.1 mL) was treated with aqueous 1N NaOH (0.7 mL, 0.70 mmol) then 2,2,2-trichloroethyl chloroformate (0.056 mL, 0.41 mmol) and the mixture was stirred at RT for 2.5 h. The mixture was treated with another portion of 2,2,2-trichloroethyl chloroformate (0.015 mL, 0.11 mmol) and stirred at RT for a further 1 h. The mixture was diluted with EtOAc and water and the phases were separated. The aqueous layer was extracted with EtOAc (x 2) and the combined organic phase was washed with brine, dried (Na_2SO_4), and concentrated *in vacuo* to give a pale red gum (195 mg). The gum was purified by FCC, using 0-40% EtOAc in cyclohexane, to give the *title compound* as a pale golden gum (134 mg, 74%). LCMS (Method 3): Rt 4.15 min, m/z 464.2, 466.2 $[\text{MH}^+]$.

d. 1-{5-tert-Butyl-2-[3-((R)-2-hydroxy-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 151d)




5

A solution of Intermediate 151c (102 mg, 0.22 mmol) in 1,4-dioxane (2.0 mL) was treated with Intermediate 81d (75 mg, 0.20 mmol) and DIPEA (0.043 mL, 0.25 mmol) and the mixture was stirred at 70°C for 7.5 h then at 50°C for 64 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. The gum was purified by FCC, using 0-10% MeOH in DCM to give the *title compound* as a yellow glass (84 mg, 61%). LCMS (Method 3): Rt 3.75 min, m/z 693.5 [MH⁺].

15

e. Methanesulfonic acid (R)-2-{3-[3-tert-butyl-5-(3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenoxy}-propyl urea (Intermediate 151e)



20

A solution of Intermediate 151d (81 mg, 0.12 mmol) in DCM (2.7 mL) was treated with DIPEA (0.061 mL, 0.35 mmol) then methanesulfonyl chloride (0.012

mL, 0.15 mmol). After 30 mins the mixture was treated with methanesulfonyl chloride (0.012 mL, 0.15 mmol) and then, after another 30 mins, with DIPEA (0.030 mL, 0.18 mmol). The mixture was stirred at RT for 16 h. Another portion of methanesulfonyl chloride (0.012 mL, 0.15 mmol) was added and then after 30 mins another portion of methanesulfonyl chloride (0.012 mL, 0.15 mmol) and DIPEA (0.030 mL, 0.18 mmol) were added. After 30 mins the mixture was diluted with DCM and saturated aqueous sodium bicarbonate solution. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with water, saturated aqueous sodium bicarbonate solution and brine, dried (Na₂SO₄) and concentrated *in vacuo* to give the *title compound* as a brown glass (92 mg, quant.). LCMS (Method 3): Rt 3.97 min, m/z 771.5 [MH⁺].


f. 1-{5-tert-Butyl-2-[3-((R)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea formate salt (Example 151)

A solution of Intermediate 151e (89 mg, 0.12 mmol) in THF (0.8 mL) was treated with a 2M solution of dimethylamine in THF (1.2 mL, 2.3 mmol) and the mixture was stirred at 60°C for 42 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown glass. The glass was purified by HPLC (Phenomenex Gemini C18 column, 5-95% MeCN in H₂O, 0.1% HCO₂H) over 25 mins. Concentration of the fractions *in vacuo* gave a pale yellow glass. The glass was triturated with ether to give a solid that was filtered off to give the *title compound* as a fawn coloured solid (16 mg, 19%). LCMS (Method 5): Rt 3.68 min, m/z 720.5 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.5 Hz), 1.24 (3H, d, J = 6.1 Hz), 1.27


(9H, s), 1.46-1.56 (2H, m), 1.61-1.72 (2H, m), 1.74-1.98 (4H, m), 1.99-2.16 (2H, m), 2.17 (6H, s), 2.32-2.39 (1H, m), 2.86-2.95 (1H, m), 3.13-3.19 (1H, m), 3.29-3.34 (1H, m), 4.56-4.65 (1H, m), 4.79-4.86 (1H, m), 5.52 (1H, t, $J = 4.4$ Hz), 6.32 (1H, s), 6.94-6.99 (1H, m), 7.03-7.12 (3H, m), 7.19 (1H, dd, $J = 9.9, 2.0$ Hz), 7.24-7.41 (5H, m), 7.64 (1H, d, $J = 10.2$ Hz), 7.68-7.70 (1H, m), 8.15 (1H, s), 8.20 (1H, s).

Example 152

1-**{5-tert-Butyl-2-[3-((S)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}**-3-**{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}**-urea formate salt



a. **2-[3-((S)-2-Benzoyloxy-1-methyl-ethoxy)-phenyl]-5-tert-butyl-2H-pyrazol-3-ylamine (Intermediate 152a)**




15

A mixture of Intermediate 95a (0.50 g, 2.2 mmol) in dry THF (17 mL) was treated with triphenylphosphine (1.13 g, 4.32 mmol) and (R)-(-)-1-benzyloxy-2-propanol (0.52 mL, 3.24 mmol) then diisopropylazodicarboxylate (0.85 mL, 4.32 mmol) was added dropwise [the reaction mixture became very warm]. After stirring at ambient temperature for 1.75 h the mixture was treated with water (0.2 mL) then concentrated *in vacuo* to a viscous orange oil. The oil was purified by FCC, using 0-7% EtOAc in DCM. The resulting oil was purified further by FCC,

20


using 0-30% EtOAc in cyclohexane. The product was applied to a pre-conditioned (with MeOH) SCX-2 cartridge and eluted with MeOH then 2N NH₃ in MeOH. The product containing fractions were combined and concentrated *in vacuo* to give the *title compound* as a viscous golden oil (0.39 g, 48%). LCMS (Method 3): Rt 3.77 min, m/z 380.4 [MH⁺].

b. (S)-2-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenoxy]-propan-1-ol (Intermediate 152b)




A solution of Intermediate 152a (0.39 g, 1.0 mmol) in ethanol (IMS grade, 10 mL) was purged with N₂ then 10% Pd/C (0.11g, 0.1 mmol Pd) was added. The mixture was purged with H₂ then stirred under an atmosphere of H₂ (balloon) at ambient temperature for 2 h, then at 50°C for 1.5 h and 45°C for 30 mins. Another portion of Pd/C (0.11 g, 0.1 mmol Pd) was added and the mixture was stirred at 45°C for 16 h. The cooled reaction mixture was purged with N₂ and treated with water (2 mL) then ammonium formate (0.65 g, 10.3 mmol) and heated to reflux for 5. Another portion of ammonium formate (0.65 g, 10.3 mmol) and Pd/C (0.11 g, 0.1 mmol) were added and the mixture was stirred at reflux for 16 h. The mixture was filtered through celite and the filtrate was concentrated *in vacuo*. The residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was extracted with EtOAc (x2). The combined organic phase was washed with saturated aqueous sodium bicarbonate solution and brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-75% EtOAc in cyclohexane to give the *title compound* (0.25 g, 83%) as a pink gum. LCMS (Method 3): Rt 2.44 min, m/z 290.3 [MH⁺].

c. {5-tert-Butyl-2-[3-((S)-2-hydroxy-1-methyl-ethoxy)-phenyl]-2H-

pyrazol-3-yl}-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 152c)

A solution of Intermediate 152b (69 mg, 0.24 mmol) in EtOAc (1.2 mL) was treated with aqueous 1N NaOH (0.4 mL, 0.43 mmol) then 2,2,2-trichloroethyl chloroformate (0.034 mL, 0.25 mmol) and the mixture was stirred at RT for 2.5 h. The mixture was treated with another portion of 2,2,2-trichloroethyl chloroformate (0.010 mL, 0.07 mmol) and stirred at RT for a further 0.75 h. The mixture was diluted with EtOAc and water and the phases were separated. The aqueous layer was extracted with EtOAc (x 2) and the combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to give a pale red gum (121 mg). The gum was purified by FCC, using 0-40% EtOAc in cyclohexane, to give the *title compound* as a pale golden gum (62 mg, 56%). LCMS (Method 3): Rt 4.15 min, m/z 464.2, 466.2 [MH⁺].


d. 1-{5-tert-Butyl-2-[3-((S)-2-hydroxy-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 152d)



A solution of Intermediate 152c (59 mg, 0.13 mmol) in 1,4-dioxane (1.3 mL) was treated with Intermediate 81d (44 mg, 0.12 mmol) and DIPEA (0.025 mL, 0.14 mmol) and the mixture was stirred at 70°C for 6.75 h then at 50°C for 64 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned

between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. The gum was purified by FCC, using 0-10% MeOH in DCM to give the *title compound* as a pale yellow glass (49 mg, 61%). LCMS (Method 3): Rt 3.74 min, m/z 693.5 [MH⁺].

e. Methanesulfonic acid (S)-2-{3-[3-tert-butyl-5-(3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenoxy}-propyl urea (Intermediate 152e)



10

A solution of Intermediate 152d (46 mg, 0.066 mmol) in DCM (1.5 mL) was treated with DIPEA (0.035 mL, 0.20 mmol) then methanesulfonyl chloride (0.0067 mL, 0.086 mmol). After 30 mins the mixture was treated with methanesulfonyl chloride (0.0067 mL, 0.086 mmol) and then, after another 30 mins, with DIPEA (0.018 mL, 0.10 mmol). The mixture was stirred at RT for 16 h. Another portion of methanesulfonyl chloride (0.0067 mL, 0.086 mmol) was added and then after 30 mins another portion of methanesulfonyl chloride (0.0067 mL, 0.086 mmol) and DIPEA (0.018 mL, 0.10 mmol) were added. After 30 mins the mixture was diluted with DCM and saturated aqueous sodium bicarbonate solution. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with water, saturated aqueous sodium bicarbonate solution and brine, dried (Na₂SO₄) and concentrated *in vacuo* to give the *title compound* as a brown glass (54 mg, quant.). LCMS (Method 3): Rt 3.99 min, m/z 771.5 [MH⁺].


20

f. 1-{5-tert-Butyl-2-[3-((S)-2-dimethylamino-1-methyl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 152)


- 5 A solution of Intermediate 152e (51 mg, 0.066 mmol) in THF (0.4 mL) was treated with a 2M solution of dimethylamine in THF (0.66 mL, 1.32 mmol) and the mixture was stirred at 60°C for 42 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined
10 organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown glass. The glass was purified by HPLC (Phenomenex Gemini C18 column, 5-95% MeCN in H₂O, 0.1% HCO₂H) over 25 mins. Concentration of the fractions *in vacuo* gave a pale yellow glass. The glass was triturated with diethyl ether to give *title compound* as a fawn coloured solid (2.5 mg, 5%). LCMS
15 (Method 5): Rt 3.68 min, m/z 720.5 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.5 Hz), 1.24 (3H, d, J = 6.5 Hz), 1.27 (9H, s), 1.46-1.56 (2H, m), 1.61-1.72 (2H, m), 1.74-1.98 (4H, m), 1.99-2.16 (2H, m), 2.18 (6H, s), 2.32-2.38 (1H, m), 2.86-2.95 (1H, m), 3.12-3.19 (1H, m), 4.56-4.65 (1H, m), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.1 Hz), 6.32 (1H, s), 6.94-6.99 (1H, m), 7.03-7.11 (3H, m), 7.19
20 (1H, dd, J = 9.6, 2.4 Hz), 7.24-7.41 (5H, m), 7.64 (1H, d, J = 10.0 Hz), 7.68-7.70 (1H, m), 8.15 (1H, s), 8.21 (1H, s).

Example 153

1-[2-[3-(2-Dimethylamino-ethoxy)-phenyl]-5-(2-hydroxy-1,1-dimethyl-ethyl)-2H-pyrazol-3-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt




a. 3-(tert-Butyl-diphenyl-silanyloxy)-2,2-dimethyl-propionic acid methyl ester (Intermediate 153a)




- 10 A solution of methyl 2,2-dimethyl-3-hydroxypropionate (5.0 g, 37.8 mmol) in DMF (75 mL) was cooled to 0°C and treated with imidazole (3.86 g, 56.7 mmol) then tert-butyl diphenylchlorosilane (11.8 mL, 45.4 mmol). The mixture was stirred at RT for 22 h then concentrated *in vacuo*. The residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was
- 15 extracted with EtOAc (x2). The combined organic phase was washed with 10% aqueous citric acid solution, saturated aqueous sodium bicarbonate solution and brine, dried (Na₂SO₄) and concentrated *in vacuo* to give the *title compound* as a golden oil (15.93 g, quant.). LCMS (Method 3): Rt 5.32 min, m/z 393.3 [M+Na⁺].

**b. 5-(tert-Butyl-diphenyl-silanyloxy)-4,4-dimethyl-3-oxo-pentanenitrile
(Intermediate 153b)**



A suspension of sodium hydride (60% dispersion in oil) (2.12 g, 52.9 mmol) in toluene (65 mL) at reflux under N₂ was added over 55 mins a solution of Intermediate 153a (15.93 g, assumed 37.8 mmol) and acetonitrile (2.87 mL, 54.9 mmol) in toluene (31 mL). The mixture was heated at reflux for 67 h. The cooled reaction mixture was slowly acidified to pH 5 with aqueous 1N HCl and extracted with EtOAc. The aqueous layer was extracted with EtOAc (x2) and the combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-50% ether in pentane, to give the *title compound* (3.52 g, 25%) as a yellow oil. LCMS (Method 3): Rt 4.94 min, m/z 402.3 [M+Na⁺].


c. 2-(3-Benzyloxy-phenyl)-5-[2-(tert-butyl-diphenyl-silanyloxy)-1,1-dimethyl-ethyl]-2H-pyrazol-3-ylamine (Intermediate 153c)



A solution of Intermediate 153b (1.00 g, 2.63 mmol) in ethanol (IMS grade, 20 mL) was treated with 3-benzyloxyphenylhydrazine hydrochloride (0.66 mL, 2.63 mmol) then DIPEA (0.46 mL, 2.63 mmol) and the mixture was heated to reflux for 114 h. The cooled reaction mixture was diluted with EtOAc and a saturated aqueous sodium bicarbonate solution and the mixture was filtered and the


phases separated. The aqueous layer was extracted with EtOAc (x 2) and the combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to give an orange oil (1.62 g). The gum was purified by FCC, using 0-20% EtOAc in cyclohexane, to give the *title compound* as a yellow oil (1.21 g, 80%). LCMS (Method 3): Rt 5.40 min, m/z 576.5 [MH⁺].

d. 3-{5-Amino-3-[2-(tert-butyl-diphenyl-silanyloxy)-1,1-dimethyl-ethyl]-pyrazol-1-yl}-phenol (Intermediate 153d)



A solution of Intermediate 153c (1.21 g, 2.1 mmol) in ethanol (IMS grade, 20 mL) was treated with water (1 mL) and ammonium formate (1.33 g, 21 mmol). The mixture was purged with N₂ then 10% Pd/C (0.67g, 0.63 mmol Pd) was added. The mixture was heated to reflux for 1 h then filtered through celite and the filtrate was concentrated *in vacuo*. The residue was partitioned between EtOAc and water. The phases were separated and the aqueous layer was extracted with EtOAc (x2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by FCC, using 0-40% EtOAc in cyclohexane, to give the *title compound* (0.53 g, 41% over 2 steps) as a golden gum. LCMS (Method 3): Rt 4.55 min, m/z 486.5 [MH⁺].


e. **5-[2-(tert-Butyl-diphenyl-silanyloxy)-1,1-dimethyl-ethyl]-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-ylamine**
(Intermediate 153e)



5

A mixture of Intermediate 153d (0.53 g, 1.09 mmol) in dry THF (8 mL) was treated with triphenylphosphine (0.57 g, 2.18 mmol) and 2-(tetrahydro-2H-pyran-2-yloxy)ethanol (0.22 mL, 1.64 mmol). The mixture was cooled to 0°C then diisopropylazodicarboxylate (0.43 mL, 2.18 mmol) was added dropwise. After stirring at ambient temperature for 20 h the mixture was treated with water (0.1 mL) and concentrated *in vacuo* to a yellow oil. The oil was purified by FCC, using 0-30% EtOAc in cyclohexane, to give the *title compound* as an orange oil (0.41 g, 61%). LCMS (Method 3): Rt 5.25 min, m/z 614.6 [MH⁺].


f. **(5-[2-(tert-Butyl-diphenyl-silanyloxy)-1,1-dimethyl-ethyl]-2-{3-[2-(tetrahydro-pyran-2-yloxy)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 153f)**



15

A solution of Intermediate 153e (0.41 g, 0.67 mmol) in EtOAc (3.6 mL) was treated with aqueous 1N NaOH (1.2 mL, 1.2 mmol) then 2,2,2-trichloroethyl chloroformate (0.10 mL, 0.74 mmol) and the mixture was stirred at RT for 4 h. The mixture was diluted with EtOAc and water and the phases were separated. The aqueous layer was extracted with EtOAc (x 2) and the combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to give a pale red oil (0.67 g). The oil was purified by FCC, using 0-25% EtOAc in cyclohexane to give the *title compound* as a pale golden gum (0.38 g, 72%). LCMS (Method 3): Rt 5.86 min, m/z 788.3, 790.3 [MH⁺].


g. 1-[3-(1-([tert-Butyl(diphenyl)silyl]oxy)-2-methylpropan-2-yl)-1-{3-[2-(tetrahydro-2H-pyran-2-yloxy)ethoxy]phenyl}-1H-pyrazol-5-yl]-3-[(1S,4R)-4-({3-[(2S)-2-methylpiperidin-1-yl][1,2,4]triazolo[4,3-a]pyridin-6-yl}oxy)-1,2,3,4-tetrahydronaphthalen-1-yl]urea (Intermediate 153g)



A solution of Intermediate 153f (377 mg, 0.48 mmol) in 1,4-dioxane (4.8 mL) was treated with Intermediate 81d (164 mg, 0.43 mmol) and DIPEA (0.095 mL, 0.54 mmol) and the mixture was stirred at 70°C for 30 mins then at 50°C for 64 h then at 70°C for 22 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. The gum was purified by FCC, using 0-7% MeOH in DCM to give the *title compound* as a brown glass (320 mg, 72%). LCMS (Method 3): Rt 5.66 min, m/z 1017.7


[MH⁺].

- h. 1-{3-(1-{[tert-Butyl(diphenyl)silyl]oxy}-2-methylpropan-2-yl)-1-[3-(2-hydroxyethoxy)phenyl]-1H-pyrazol-5-yl}-3-[(1S,4R)-4-({3-[(2S)-2-methylpiperidin-1-yl][1,2,4]triazolo[4,3-a]pyridin-6-yl}oxy)-1,2,3,4-tetrahydronaphthalen-1-yl]urea (Intermediate 153h)



A solution of Intermediate 153g (317 mg, 0.31 mmol) in MeOH (3.1 mL) was treated with pyridinium para-toluenesulfonate (235 mg, 0.93 mmol) and the mixture was stirred at 45°C for 19 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and a saturated aqueous sodium bicarbonate solution. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with water, a saturated aqueous sodium bicarbonate solution, and brine, dried (Na₂SO₄) and concentrated *in vacuo* to a cream coloured foam. The foam was purified by FCC, using 0-10% MeOH in DCM, to give the *title compound* as a colourless glass (217 mg, 75%). LCMS (Method 3): Rt 5.06 min, m/z 933.7 [MH⁺].

i. 2-{3-[3-(1-{[tert-Butyl(diphenyl)silyl]oxy}-2-methylpropan-2-yl)-5-(([(1S,4R)-4-({3-[(2S)-2-methylpiperidin-1-yl][1,2,4]triazolo[4,3-a]pyridin-6-yl]oxy)-1,2,3,4-tetrahydronaphthalen-1-yl]carbamoyl}amino)-1H-pyrazol-1-yl]phenoxy}ethyl methanesulfonate (Intermediate 153i)




5

A solution of Intermediate 153h (214 mg, 0.23 mmol) in DCM (2.3 mL) was treated with DIPEA (0.12 mL, 0.69 mmol) then methanesulfonyl chloride (0.037 mL, 0.48 mmol). After 30 mins the mixture was diluted with DCM and saturated aqueous sodium bicarbonate solution. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with water, saturated aqueous sodium bicarbonate solution and brine, dried (Na₂SO₄) and concentrated *in vacuo* to an orange gum. The gum was purified by FCC, using 0-8% MeOH in DCM, to give the *title compound* as a pale brown glass (134 mg, 58%). LCMS (Method 3): Rt 5.18 min, m/z 1011.7 [MH⁺].

10

15

j. 1-[3-(1-{[tert-Butyl(diphenyl)silyl]oxy}-2-methylpropan-2-yl)-1-{3-[2-(dimethylamino)ethoxy]phenyl}-1H-pyrazol-5-yl]-3-[(1S,4R)-4-({3-[(2S)-2-methylpiperidin-1-yl][1,2,4]triazolo[4,3-a]pyridin-6-yl]oxy)-1,2,3,4-tetrahydronaphthalen-1-yl]urea (Example 153j)



A solution of Intermediate 153i (131 mg, 0.13 mmol) in THF (0.8 mL) was treated with a 2M solution of dimethylamine in THF (1.3 mL, 2.6 mmol) and the mixture was stirred at 60°C for 18 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and water. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. The gum was purified by FCC, using 0-8% (2N NH₃/MeOH) in DCM, to give the *title compound* as a white foam (93 mg, 75%). LCMS (Method 3): Rt 3.91 min, m/z 960.6 [MH⁺].


1-[2-[3-(2-Dimethylamino-ethoxy)-phenyl]-5-(2-hydroxy-1,1-dimethyl-ethyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt (Example 153)

A solution of Intermediate 153j (90 mg, 0.09 mmol) in THF (0.5 mL) was treated with a 1M solution of tetrabutylammonium fluoride in THF (0.5 mL, 0.5 mmol) and the reaction mixture was stirred at RT for 1 h then at 50°C for 16 h. The cooled solution was concentrated *in vacuo*, and the residue was partitioned between DCM and a saturated aqueous sodium bicarbonate solution. The phases were separated and the aqueous layer was extracted with DCM (x 2). The combined organic phase was washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to a brown gum. The gum was purified by FCC, using 0-8% [2M NH₃ in MeOH] in DCM, and then purified further by HPLC (Phenomenex Gemini C18 column, 5-95% MeCN in H₂O, 0.1% HCO₂H) over 25 mins to give a colourless glass. The glass was triturated with diethyl ether to give the *title compound* as a white solid (11 mg, 16%). LCMS (Method 5): Rt 3.17 min, m/z 722.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.21 (6H, s), 1.44-1.57 (2H, m), 1.62-1.73 (2H, m), 1.74-1.98 (4H, m), 1.99-2.17 (2H, m), 2.18 (6H, s), 2.61 (2H, t, J = 6.0 Hz), 2.86-2.95 (1H, m), 3.12-3.20 (1H, m), 3.27-3.36 (1H, m), 3.44


(2H, s), 4.08 (2H, t, J = 6.0 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.32 (1H, s), 6.94-6.99 (1H, m), 7.06-7.14 (3H, m), 7.19 (1H, dd, J = 10.1, 2.0 Hz), 7.24-7.43 (5H, m), 7.64 (1H, d, J = 9.7 Hz), 7.68-7.71 (1H, m), 8.15 (1H, s), 8.20 (1H, s).

5 **Example 154**

1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt



10 **Methanesulfonic acid 2-{3-[3-tert-butyl-5-(3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-phenoxy}-ethyl ester (Intermediate 154a)**




15 Methanesulfonyl chloride (81 μ L, 1.05 mmol) was added to a stirred solution of Intermediate 95e (545 mg, 803 mmol) and DIPEA (420 μ L, 2.41 mmol) in DCM (20 mL), under argon and the reaction mixture was stirred at RT for 1.5 h. A further 25 μ L, 0.32 mmol of methanesulfonyl chloride was added and

the reaction mixture continued to stir at RT for 2 h. Water (20 mL) was added and the two layers were shaken thoroughly before being separated. The aqueous layer was extracted with DCM (2 x 20 mL) and the combined organics were passed through a phase separator and concentrated *in vacuo* to afford the *title compound* as an orange glass (690 mg, >100%). LCMS (Method 3): Rt 3.89 min, m/z 757 [MH⁺].


1-{5-tert-Butyl-2-[3-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 154)

A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and pyrrolidine (27 µL, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (29 mg, 59%). LCMS (Method 5): Rt 3.72 min, m/z 732.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 8.0 Hz), 1.28 (9H, s), 1.46-1.56 (2H, m), 1.61-1.72 (6H, m), 1.75-1.98 (4H, m), 1.99-2.18 (2H, m), 2.52 (4H, m, obscured by solvent), 2.80 (2H, t, J = 8.0 Hz), 2.86-2.95 (1H, m), 3.12-3.19 (1H, m, obscured by water), 3.27-3.35 (1H, m, obscured by water), 4.11 (2H, t, J = 8.0 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.33 (1H, s), 6.97 (1H, dd, J = 8.0 Hz, 4.0 Hz), 7.05-7.14 (3H, m), 7.19 (1H, dd, J = 8.0 Hz, 4.0 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 8.0 Hz), 7.69 (1H, d, J = 4.0 Hz), 8.14 (1H, s), 8.18 (1.3H, s).

Example 155


**1-{5-tert-Butyl-2-[3-(2-diethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-
3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
5 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and diethylamine (34 μ L, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. Additional diethylamine (102 μ L, 0.99 mmol) was added and the mixture subsequently stirred at 60°C in a sealed tube for 24 h. The mixture was
10 concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (22 mg, 45%). LCMS (Method 5): Rt 3.74 min, m/z 734.6 $[MH^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 0.89-0.96 (9H, m), 1.28 (9H, s), 1.46-1.56 (2H, m), 1.62-1.73 (2H, m), 1.74-1.98 (4H, m), 1.99-2.18 (2H, m), 2.52 (4H, m, obscured by solvent), 2.76 (2H, t, $J = 8.0$ Hz), 2.87-2.94
15 (1H, m), 3.12-3.19 (1H, m, obscured by water), $\sim$ 3.30 (1H, m, obscured by water), 4.05 (2H, t, $J = 8.0$ Hz), 4.77-4.86 (1H, m), 5.52 (1H, t, $J = 4.0$ Hz), 6.33 (1H, s), 6.95 (1H, dd, $J = 8.0$ Hz, 4.0 Hz), 7.05-7.13 (3H, m), 7.19 (1H, dd, $J = 8.0$ Hz, 4.0 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, $J = 8.0$ Hz), 7.69 (1H, d, $J = 4.0$ Hz), 8.13 (1H, s), 8.19 (1H, s).

Example 156


1-{5-tert-Butyl-2-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt

A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and morpholine (29 μ L, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. Additional morpholine (29 μ L, 0.33 mmol) was added and the mixture subsequently stirred at 60°C in a sealed tube for 24 h. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (25 mg, 50%). LCMS (Method 5): Rt 3.66 min, m/z 748.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.0 Hz), 1.27 (9H, s), 1.46-1.56 (2H, m), 1.61-1.73 (2H, m), 1.75-1.98 (4H, m), 1.99-2.19 (2H, m), 2.42 (4H, t, J = 4.0 Hz), 2.67 (2H, t, J = 8.0 Hz), 2.86-2.95 (1H, m), 3.12-3.20 (1H, m, obscured by water), ~3.30 (1H, m, obscured by water), 3.53 (4H, t, J = 8.0 Hz), 4.12 (2H, t, J = 8.0 Hz), 4.77-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.33 (1H, s), 6.95-6.99 (1H, m), 7.07-7.13 (3H, m), 7.19 (1H, dd, J = 8.0 Hz, 4.0 Hz), 7.25-7.43 (5H, m), 7.64 (1H, d, J = 8.0 Hz), 7.69 (1H, d, J = 4.0 Hz), 8.13 (1H, s), 8.21 (0.5H, s).

Example 157

1-{5-tert-Butyl-2-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt


A solution of Intermediate 154a (66 mg, 88 μmol) and piperidine (44 μL , 0.45 mmol) in THF (2 mL) was heated at 60°C overnight in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give the *title compound* as a white solid (31 mg, 46%). LCMS (Method 5): Rt 3.77 min, m/z 746 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.91 (3H, d, $J = 6.2$ Hz), 1.24-1.34 (11H, m), 1.39-1.55 (6H, m), 1.61-2.18 (8H, m), 2.34-2.43 (4H, m), 2.63 (2H, t, $J = 6.1$ Hz), 2.86-2.94 (1H, m), 3.11-3.19 (1H, m), 3.27-3.36 (1H, m), 4.09 (2H, t, $J = 5.9$ Hz), 4.77 (1H, m), 5.52 (1H, t, $J = 4.0$ Hz), 6.32 (1H, s), 6.93-6.98 (1H, m), 7.04-7.13 (3H, m), 7.19 (1H, dd, $J = 9.5$ Hz, 2.6 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, $J = 9.9$ Hz), 7.69 (1H, d, $J = 1.8$ Hz), 8.12 (1H, s), 8.17 (1.5H, s).

Example 158

1-(5-tert-Butyl-2-{3-[2-(4-methyl-piperazin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and 1-methylpiperazine (37 μ L, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (28 mg, 55%). LCMS (Method 5): Rt 3.62 min, m/z 761.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 8.0 Hz), 1.28 (9H, s), 1.47-1.56 (2H, m), 1.61-1.73 (2H, m), 1.74-1.97 (4H, m), 2.00-2.19 (5H, m), 2.21—2.35 (4H, m), 2.36-2.49 (4H, m), 2.66 (2H, t, J = 8.0 Hz), 2.86-2.95 (1H, m), 3.12-3.19 (1H, m, obscured by water), 3.27-3.35 (1H, m, obscured by water), 4.10 (2H, t, J = 8.0 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.33 (1H, s), 6.94-6.98 (1H, m), 7.05-7.13 (3H, m), 7.19 (1H, dd, J = 8.0 Hz, 4.0 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 8.0 Hz), 7.69 (1H, d, J = 4.0 Hz), 8.13 (1H, s), 8.17 (1.4H, s).

Example 159




1-{5-tert-Butyl-2-{3-[2-(4-fluoropiperidin-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and 4-fluoropiperidine (34 mg, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in

a sealed tube. Additional 4-fluoropiperidine (34 mg, 0.33 mmol) was added and the mixture subsequently stirred at 60°C in a sealed tube for 24 h. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (24 mg, 47%). LCMS (Method 5): Rt 3.75 min, m/z 764.6 $[MH^+]$. ^1H NMR (400 MHz, d_6 -DMSO): 0.91 (3H, d, $J = 8.0$ Hz), 1.27 (9H, s), 1.44-1.64 (2H, m), 1.59-1.72 (4H, m), 1.72-1.98 (6H, m), 1.99-2.19 (2H, m), 2.31-2.40 (2H, m), 2.55-2.64 (2H, m), 2.68 (2H, t, $J = 8.0$ Hz), 2.87-2.95 (1H, m), 3.12-3.19 (1H, m, obscured by water), ~3.30 (1H, m, completely obscured by water), 4.10 (2H, t, $J = 8.0$ Hz), 4.51-4.59 (0.5H, m), 4.64-4.71 (0.5H, m), 4.78-4.86 (1H, m), 5.52 (1H, t, $J = 4.0$ Hz), 6.33 (1H, s), 6.95-6.99 (1H, m), 7.06-7.13 (3H, m), 7.19 (1H, dd, $J = 8.0$ Hz, 4.0 Hz), 7.25-7.42 (5H, m), 7.64 (1H, d, $J = 8.0$ Hz), 7.69 (1H, d, $J = 4.0$ Hz), 8.12 (1H, s), 8.18 (1H, s).

Example 160




1-(5-tert-Butyl-2-{3-[2-(4-methyl-[1,4]-diazepan-1-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

A solution of Intermediate 154a (50.5 mg, 0.066 mmol) and N-methyl homopiperazine (39 mg, 0.34 mmol) in THF (1 mL) was heated at 60°C overnight in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give a white solid of the *title compound* (28 mg, 54%). LCMS (Method 5): Rt 3.28 min, m/z 775 $[MH^+]$. ^1H

NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.0 Hz), 1.27 (9H, s), 1.47-1.56 (2H, m), 1.62-1.73 (4H, m), 1.75-2.18 (6H, m), 2.28 (3H, s), 2.56-2.63 (4H, m), 2.70-2.77 (4H, m), 2.85 (2H, t, J = 6.0 Hz), 2.87-2.95 (1H, m), 3.12-3.20 (1H, m), 3.27-3.35 (1H, m), 4.07 (2H, t, J = 5.6 Hz), 4.77-4.86 (1H, m), 5.52 (1H, t, J = 3.8 Hz), 6.32 (1H, s), 6.93-6.98 (1H, m), 7.04-7.10 (2H, m), 7.14 (1H, d, J = 8.9 Hz), 7.19 (1H, dd, J = 9.8 Hz, 2.2 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 4.9 Hz), 7.69 (1H, d, J = 1.8 Hz), 8.18 (1H, s), 8.20 (2H, s).

Example 161




1-{5-tert-Butyl-2-[3-(2-[1,4]oxazepan-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 154a (50.5 mg, 66 μ mol) and [1,4]oxazepine (70 mg, 0.68 mmol) in THF (1 mL) was heated at 60°C for 48 h in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give the *title compound* as a white solid (27 mg, 53%). LCMS (Method 5): Rt 3.68 min, m/z 762 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.2 Hz), 1.27 (9H, s), 1.46-1.54 (2H, m), 1.62-2.17 (12H, m), 2.67-2.72 (3H, m), 2.86 (2H, t, J = 6.0 Hz), 2.88-2.94 (1H, m), 3.12-3.19 (1H, m), 3.28-3.34 (1H, m, obscured by water peak), 3.53-3.58 (2H, m), 3.61 (2H, t, J = 6.0 Hz), 4.09 (2H, t, J = 6.1 Hz), 4.77-4.86 (1H,

m), 5.52 (1H, t, $J = 4.3$ Hz), 6.33 (1H, s), 6.94-6.99 (1H, m), 7.05-7.12 (2H, m), 7.19 (1H, dd, $J = 8.6$ Hz, 2.0 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, $J = 9.8$ Hz), 7.69 (1H, d, $J = 2.0$ Hz), 8.17 (1H, s), 8.18 (1H, s).

Example 162

5




1-(2-{3-[2-(8-Aza-bicyclo[3.2.1]oct-8-yl)-ethoxy]-phenyl}-5-tert-butyl-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

10


A solution of Intermediate 154a (50.5 mg, 66 μ mol) and 8-aza-bicyclo[3.2.1]octane (38 mg, 0.34 mmol) in THF (1 mL) was heated at 60°C overnight in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give the *title compound* as a white solid (32 mg, 62%). LCMS (Method 5): Rt 3.84 min, m/z 772 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.91 (3H, d, $J = 6.4$ Hz), 1.21-1.36 (11H, m), 1.37-1.54 (4H, m), 1.58-1.72 (4H, m), 1.74-2.18 (8H, m), 2.72 (2H, d, $J = 6.4$ Hz), 2.86-2.95 (1H, m), 3.11-3.20 (1H, m), 3.22-3.36 (5H, m), 4.09 (2H, t, $J = 6.1$ Hz), 4.77-4.86 (1H, m), 5.52 (1H, t, $J = 4.1$ Hz), 6.33 (1H, s), 6.94-6.99 (1H, m), 7.04-7.11 (2H, m), 7.14 (1H, d, $J = 8.6$ Hz), 7.18 (1H, dd, $J = 10.2$ Hz, 2.1 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, $J = 10.2$ Hz), 7.69 (1H, d, $J = 2.0$ Hz), 8.17 (1H, s), 8.19 (1.6H, s).

20

Example 163

**1-(5-tert-Butyl-2-{3-[2-(ethyl-methyl-amino)-ethoxy]-phenyl}-2H-
 5 pyrazol-3-yl)-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
 [1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-
 urea formate salt**


A solution of Intermediate 154a (50.5 mg, 66 μ mol) and N-ethylmethanamine (40 mg, 0.68 mmol) in THF (1 mL) was heated at 60°C for
 10 48 h in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give the *title compound* as a white solid (24 mg, 50%). LCMS (Method 5): Rt 3.69 min, m/z 720 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 0.94 (3H, t, J = 7.2 Hz), 1.27 (9H, s), 1.46-1.55 (2H, m), 1.64-1.72 (2H, m), 1.76-2.09 (6H, m), 2.19
 15 (3H, s), 2.41 (2H, quart, J = 7.0 Hz), 2.68 (2H, t, J = 4.7 Hz), 2.86-2.94 (1H, m), 3.12-3.19 (1H, m), 3.29-3.34 (1H, m, obscured by water peak), 4.08 (2H, t, J = 6.1 Hz), 4.77-4.86 (1H, m), 5.52 (1H, t, J = 4.4 Hz), 6.33 (1H, s), 6.94-6.98 (1H, m), 7.05-7.13 (3H, m), 7.19 (1H, dd, J = 9.6 Hz, 2.2 Hz), 7.23-7.42 (5H, m), 7.64 (1H, d, J = 10.2 Hz), 7.69 (1H, d, J = 2.2 Hz), 8.13 (1H, s), 8.18
 20 (1H, s).

Example 164

1-{5-tert-Butyl-2-(3-{2-[(2-methoxy-ethyl)-methyl-amino]-ethoxy}-phenyl)-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
5 [1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
urea formate salt


A solution of Intermediate 154a (50.5 mg, 0.067 mmol) and N-(2-methoxyethyl)methylamine (35 μ L, 0.33 mmol) in THF (1 mL) was stirred at 60°C for 20 h in a sealed tube. Additional N-(2-methoxyethyl)methylamine (35 μ L, 0.33 mmol) was added and the mixture subsequently stirred at 60°C in a sealed tube for 24 h. The mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7). The *title product* was isolated as an off-white solid (24 mg, 48%).

LCMS (Method 5): Rt 3.72 min, m/z 750.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 8.0 Hz), 1.27 (9H, s), 1.44-1.56 (2H, m), 1.61-1.73 (2H, m), 1.73-1.98 (4H, m), 1.98-2.22 (2H, m), 2.24 (3H, s), 2.55 (2H, t, J = 8.0 Hz), 2.74 (2H, t, J = 8.0 Hz), 2.86-2.95 (1H, m), 3.12-3.18 (1H, m, obscured by water), 3.19 (3H, s, obscured by water), ~3.30 (1H, m, obscured by water), 3.38 (2H, t, J = 8.0 Hz, obscured by water), 4.07 (2H, t, J = 8.0 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.33 (1H, s), 6.94-6.99 (1H, m), 7.05-7.14 (3H, m), 7.19 (1H, dd, J = 8.0 Hz, 4.0 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 8.0 Hz), 7.69 (1H, d, J = 4.0 Hz), 8.13 (1H, s), 8.19 (1H, s).

Example 165

**1-(5-tert-Butyl-2-{3-[2-(4-methoxy-piperidin-1-yl)-ethoxy]-phenyl}-
 5 2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-
 [1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-
 urea formate salt**

A solution of Intermediate 154a (50.5 mg, 0.066 mmol) and 4-methoxypiperidine (39 μ L, 0.34 mmol) in THF (1 mL) was heated at 60°C
 10 overnight in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give a white solid of the *title compound* (31 mg, 60%). LCMS (Method 5): Rt 3.75 min, m/z 776 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.2 Hz), 1.27 (9H, s), 1.30-1.42 (2H, m), 1.45-1.57 (2H, m), 1.61-2.18 (13H, m), 2.66 (2H, t, J = 5.7 Hz), 2.68-
 15 2.76 (1H, m), 2.86-2.95 (1H, m), 3.04-3.19 (2H, m), 3.16 (3H, s), 3.28-3.34 (1H, m), 4.09 (2H, t, J = 5.9 Hz), 4.77-4.85 (1H, m), 5.52 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 6.93-6.98 (1H, m), 7.04-7.13 (3H, m), 7.18 (1H, dd, J = 10.3 Hz, 2.3 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 9.5 Hz), 7.69 (1H, d, J = 2.3 Hz), 8.12 (1H, s), 8.16 (1.7H, s).


Example 166

1-(5-tert-Butyl-2-{3-[2-(3-oxa-8-aza-bicyclo[3.2.1]oct-8-yl)-ethoxy]-phenyl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

A solution of Intermediate 154a (50.5 mg, 66 μ mol) and 3-oxa-8-aza-bicyclo[3.2.1]octane (76 mg, 0.68 mmol) in THF (1 mL) was heated at 60°C for 48 h in a sealed vessel. The reaction mixture was concentrated *in vacuo* and the residue purified by MDAP (Method 7) to give the *title compound* as a white solid (24 mg, 46%). LCMS (Method 5): Rt 3.72 min, m/z 774 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.28 (9H, s), 1.46-1.54 (2H, m), 1.64-1.71 (4H, m), 1.75-2.17 (8H, m), 2.59 (2H, t, J = 5.9 Hz), 2.86-2.95 (1H, m), 3.07-3.11 (2H, m), 3.12-3.19 (1H, m), 3.28-3.34 (3H, m, obscured by water peak), 3.49 (2H, d, J = 9.6 Hz), 4.09 (2H, t, J = 6.3 Hz), 4.78-4.85 (1H, m), 5.52 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 6.94-6.99 (1H, m), 7.05-7.14 (3H, m), 7.19 (1H, dd, J = 9.6 Hz, 2.4 Hz), 7.24-7.42 (5H, m), 7.64 (1H, d, J = 10.0 Hz), 7.69 (1H, d, J = 2.2 Hz), 8.15 (1H, s), 8.20 (1H, s).


Example 167

1-{5-tert-Butyl-2-[3-(4-methoxy-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt



5

a. 1-[5-tert-Butyl-2-(3-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea (Intermediate 167a)




10

A mixture of Intermediate 81d (319 mg, 0.85 mmol), Intermediate 29c (335 mg, 0.85 mmol) and DIPEA (294 μ L, 1.69 mmol) in dioxane (10 mL) was stirred at 80°C for 18 hours. After cooling, the reaction mixture was partitioned between water and DCM. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% MeOH in DCM, to afford the *title compound* (362 mg, 66%). LCMS (Method 4): Rt 3.29 min, m/z 649 [MH^+].

15

20

b. Methanesulfonic acid 3-[3-tert-butyl-5-(3-{(1*S*,4*R*)-4-[3-((*S*)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-pyrazol-1-yl]-benzyl ester (Intermediate 167b)



5

To an ice-bath cooled solution of Intermediate 167a (316 mg, 0.49 mmol) in DCM (5.0 mL) was added DIPEA (339 μ L, 1.95 mmol) followed by methanesulfonyl chloride (76 μ L, 0.97 mmol). The reaction mixture was stirred for 3 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (Quantitative, assumed 0.49 mmol). The isolated product was used in the following step without further purification. LCMS (Method 4): Rt 3.61 min, m/z 727 [MH^+].


c. 1-{5-tert-Butyl-2-[3-(4-methoxy-piperidin-1-yl)methyl]-phenyl]-2H-pyrazol-3-yl}-3-{(1*S*,4*R*)-4-[3-((*S*)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-*a*]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 167)

To a solution of Intermediate 167b (0.19 mmol) in THF (2 mL) was added DIPEA (130 μ L, 0.75 mmol) and 4-methoxypiperidine (86 mg, 0.75 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM, followed by

20

MDAP (Method 7) purification, to give the *title compound* (20 mg, 14%). LCMS (Method 5): Rt 3.66 min, m/z 746 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.30-1.40 (2H, m), 1.42-1.50 (2H, m), 1.58-1.67 (2H, m), 1.70-1.83 (5H, m), 1.83-1.93 (1H, m), 1.96-2.12 (4H, m), 2.56-2.64 (2H, m), 2.86 (1H, ddd, J = 12.7, 9.0, 3.9 Hz), 3.05-3.12 (3H, m), 3.13 (3H, s), 3.45 (2H, s), 4.77 (1H, td, J = 8.4, 5.7 Hz), 5.47 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 7.00 (1H, d, J = 8.6 Hz), 7.14 (1H, dd, J = 9.8, 2.1 Hz), 7.20-7.30 (4H, m), 7.30-7.37 (3H, m), 7.40 (1H, t, J = 7.7 Hz), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.07 (1H, s), 8.23 (0.3H, s).

10 **Example 168**




1-[5-tert-Butyl-2-(3-([(2-methoxy-ethyl)-methyl-amino]-methyl)-phenyl)-2H-pyrazol-3-yl]-3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl)-urea formate salt

To a solution of Intermediate 167b (0.19 mmol) in THF (2 mL) was added DIPEA (130 μ L, 0.75 mmol) and *N*-(methoxyethyl)methylamine (81 μ L, 0.75 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (20 mg,

14%). LCMS (Method 5): Rt 3.65 min, m/z 720 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.42-1.48 (2H, m), 1.58-1.67 (2H, m), 1.72-1.92 (4H, m), 1.96-2.12 (2H, m), 2.12 (3H, s), 2.49 (2H, t, J = 6.0 Hz), 2.86 (1H, ddd, J = 12.7, 9.0, 3.9 Hz), 3.12 (1H, dt, J = 12.1, 4.3 Hz), 3.16 (3H, s), 3.39 (2H, t, J = 6.0 Hz), 3.50 (2H, s), 4.77 (1H, td, J = 8.4, 5.7 Hz), 5.47 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 7.02 (1H, d, J = 8.6 Hz), 7.14 (1H, dd, J = 9.8, 2.1 Hz), 7.20-7.25 (2H, m), 7.25-7.30 (2H, m), 7.30-7.35 (2H, m), 7.38 (1H, s), 7.40 (1H, t, J = 7.7 Hz), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.07 (1H, s), 8.28 (0.2H, s).

Example 169




1-{5-tert-Butyl-2-[3-(4-fluoropiperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methylpiperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

To a solution of Intermediate 167b (0.19 mmol) in THF (2 mL) was added DIPEA (260 μ L, 1.5 mmol) and 4-fluoropiperidine hydrochloride (104 mg, 0.75 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (25 mg, 16%). LCMS (Method 5): Rt 3.68 min, m/z 734 [MH⁺]. ¹H NMR (400 MHz, d₆-

DMSO): 0.86 (3H, d, $J = 6.2$ Hz), 1.23 (9H, s), 1.42-1.50 (2H, m), 1.58-1.70 (4H, m), 1.70-1.92 (6H, m), 1.95-2.12 (2H, m), 2.23-2.31 (2H, m), 2.86 (1H, ddd, $J = 12.7, 9.0, 3.9$ Hz), 3.11 (1H, dt, $J = 12.2, 4.3$ Hz), 3.24-3.30 (2H, m), 3.48 (2H, s), 4.60 (1H, dsp, $J = 48.9, 3.5$ Hz), 4.77 (1H, td, $J = 8.4, 5.7$ Hz), 5.47 (1H, t, $J = 4.2$ Hz), 6.28 (1H, s), 6.98 (1H, d, $J = 8.6$ Hz), 7.14 (1H, dd, $J = 9.8, 2.1$ Hz), 7.20-7.25 (2H, m), 7.25-7.34 (3H, m), 7.35-7.38 (2H, m), 7.40 (1H, t, $J = 7.7$ Hz), 7.59 (1H, d, $J = 10.0$ Hz), 7.62 (1H, d, $J = 1.8$ Hz), 8.05 (1H, s), 8.10 (1.8H, s).

Example 170




**1-[5-tert-Butyl-2-(3-dimethylaminomethyl-phenyl)-2H-pyrazol-3-yl]-3-
{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 167b (0.16 mmol) in THF (2.5 mL) was added dimethylamine solution (2M THF, 1.6 mL, 3.27 mmol) and the reaction stirred at 50°C for 24 h in a sealed vessel. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (20 mg, 16%). LCMS (Method 5): Rt 3.58 min, m/z 676 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, $J = 6.2$ Hz), 1.23 (9H, s), 1.42-1.50 (2H, m), 1.58-1.67 (2H, m), 1.70-1.90

(4H, m), 1.94-2.03 (1H, m), 2.04-2.12 (2H, m), 2.11 (6H, s), 2.86 (1H, ddd, $J = 12.7, 9.0, 3.9$ Hz), 3.12 (1H, dt, $J = 11.9, 4.1$ Hz), 3.39 (2H, s), 4.77 (1H, td, $J = 8.4, 5.7$ Hz), 5.46 (1H, t, $J = 4.2$ Hz), 6.27 (1H, s), 7.02 (1H, d, $J = 8.6$ Hz), 7.15 (1H, dd, $J = 9.8, 2.1$ Hz), 7.20-7.29 (4H, m), 7.29-7.34 (1H, m), 7.35-7.41 (3H, m), 7.59 (1H, d, $J = 10.0$ Hz), 7.64 (1H, d, $J = 1.8$ Hz), 8.07 (1H, s), 8.25 (0.3H, s).

Example 171




**1-[5-tert-Butyl-2-(3-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
 10 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 167b (0.16 mmol) in THF (2 mL) was added pyrrolidine (273 μ L, 3.27 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH_3 in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (26 mg, 22%). LCMS (Method 5): Rt 3.64 min, m/z 702 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.86 (3H, d, $J = 6.2$ Hz), 1.23 (9H, s), 1.42-1.50 (2H, m), 1.58-1.66 (6H, m), 1.70-1.84 (3H, m), 1.84-1.92 (1H, m), 1.95-2.12 (2H, m), 2.36-2.42 (4H, m), 2.86 (1H, ddd, $J = 12.7, 9.0, 3.9$ Hz), 3.12 (1H, dt, $J = 11.9, 4.1$ Hz), 3.58 (2H, s), 4.77 (1H, td, $J = 8.4, 5.7$ Hz), 5.46 (1H, t, $J = 4.2$ Hz), 6.28 (1H, s), 7.01 (1H, d, $J = 8.6$ Hz), 7.15 (1H, dd, $J = 9.8, 2.1$ Hz), 7.20-7.25 (2H, m), 7.26-


7.34 (4H, m), 7.35 (1H, s), 7.39 (1H, t, J = 7.8 Hz), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.07 (1H, s), 8.19 (0.7H, s).

Example 172




1-((1S,4R)-4-(3-(4-Aza-spiro[2.5]oct-4-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy)-1,2,3,4-tetrahydro-naphthalen-1-yl)-3-{5-tert-butyl-2-[3-(2-morpholin-4-yl-ethoxy)-phenyl]-2H-pyrazol-3-yl}-urea formate salt

A brown solution of Intermediate 148g (0.099 mmol) and morpholine (43 mg, 0.50 mmol) in THF (1 mL) was stirred in a sealed vial at 60°C for 18 h. The solution was decanted and concentrated. MDAP (Method 7) gave the *title compound* as an off-white solid (44 mg, 54%). LCMS (Method 5): Rt 3.64 min, m/z 760.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.41-0.54 (4H, m), 1.27 (9H, s), 1.52-1.62 (2H, m), 1.72-1.81 (3H, m), 1.83-1.96 (3H, m), 1.99-2.15 (2H, m), 2.42 (4H, t, J = 4.4 Hz), 2.66 (2H, t, J = 5.7 Hz), 3.22-3.29 (2H, m), 3.52 (4H, t, J = 4.58 Hz), 4.11 (2H, t, J = 5.7 Hz), 4.81 (1H, td, J = 8.6, 5.5 Hz), 5.48 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 6.95-6.98 (1H, m), 7.06-7.12 (3H, m), 7.14 (1H, dd, J = 9.9, 2.2 Hz), 7.25-7.42 (5H, m), 7.57 (1H, dd, J = 2.1, 0.9 Hz), 7.61 (1H, dd, J = 9.8, 0.8 Hz), 8.11 (1H, s), 8.16 (1.3H, s).

Example 173


**1-[5-tert-Butyl-2-(3-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 167b (0.22 mmol) in THF (2.5 mL) was added piperidine (424 μ L, 4.3 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (44 mg, 25%). LCMS (Method 5): Rt 3.69 min, m/z 716 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.28-1.33 (2H, m), 1.38-1.48 (6H, m), 1.58-1.66 (2H, m), 1.70-1.92 (4H, m), 1.95-2.12 (2H, m), 2.26-2.34 (4H, m), 2.86 (1H, ddd, J = 12.7, 9.0, 3.9 Hz), 3.11 (1H, dt, J = 11.9, 4.1 Hz), 3.23-3.29 (1H, m), 3.44 (2H, s), 4.77 (1H, td, J = 8.4, 5.7 Hz), 5.46 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 6.99 (1H, d, J = 8.6 Hz), 7.15 (1H, dd, J = 9.8, 2.1 Hz), 7.20-7.25 (2H, m), 7.25-7.29 (2H, m), 7.29-7.36 (3H, m), 7.39 (1H, t, J = 7.8 Hz), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.06 (1H, s), 8.12 (1.8H, s).

Example 174

1-{5-tert-Butyl-2-[3-(4-methyl-[1,4]diazepan-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

To a solution of Intermediate 167b (0.22 mmol) in THF (2.5 mL) was added 1-methylhomopiperazine (534 μ L, 4.3 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, followed by MDAP (Method 7) purification, to give the *title compound* (28 mg, 15%). LCMS (Method 5): Rt 3.27 min, m/z 745 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.42-1.50 (2H, m), 1.58-1.66 (2H, m), 1.68-1.84 (5H, m), 1.84-1.92 (1H, m), 1.95-2.05 (1H, m), 2.05-2.12 (1H, m), 2.32 (3H, s), 2.58-2.66 (6H, m), 2.69 (2H, t, J = 5.7 Hz), 2.86 (1H, ddd, J = 12.7, 9.0, 3.9 Hz), 3.11 (1H, dt, J = 11.9, 4.1 Hz), 3.23-3.29 (1H, m), 3.62 (2H, s), 4.77 (1H, td, J = 8.4, 5.7 Hz), 5.47 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 7.04 (1H, d, J = 8.6 Hz), 7.14 (1H, dd, J = 9.8, 2.1 Hz), 7.20-7.25 (2H, m), 7.26-7.36 (4H, m), 7.38-7.43 (2H, m), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.12 (1H, s), 8.16 (2.4H, s).

Example 175


1-{5-tert-Butyl-2-[3-((1S,4S)-5-methyl-2,5-diaza-bicyclo[2.2.1]hept-2-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

To a solution of Intermediate 167b (0.15 mmol) in THF (20 mL) was added (1S,4S)-2-methyl-2,5-diaza-bicyclo[2.2.1]heptane (366 mg, 3.3 mmol) and the reaction stirred at 50°C for 24 h. The crude reaction mixture was cooled and partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM, then further purified by HPLC (C18 X-select column, 10-75% MeCN in H₂O, 0.1% formic acid) to give the *title compound* (11 mg, 9%). LCMS (Method 5): Rt 3.45 min, m/z 743 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.86 (3H, d, J = 6.2 Hz), 1.23 (9H, s), 1.42-1.50 (2H, m), 1.54-1.66 (4H, m), 1.72-1.84 (3H, m), 1.84-1.92 (1H, m), 1.95-2.15 (2H, m), 2.22 (3H, s), 2.46-2.48 (1H, m), 2.52 (1H, dd, J = 9.6, 2.4 Hz), 2.59 (1H, d, J = 9.7 Hz), 2.71 (1H, d, J = 9.6 Hz), 2.86 (1H, ddd, J = 12.7, 9.0, 3.9 Hz), 3.09-3.13 (2H, m), 3.22 (1H, s), 3.28-3.32 (1H, m), 3.63 (1H, d, J = 13.9 Hz), 3.70 (1H, d, J = 13.9 Hz), 4.77 (1H, td, J = 8.4, 5.7 Hz), 5.47 (1H, t, J = 4.2 Hz), 6.28 (1H, s), 7.04 (1H, d, J = 8.6 Hz), 7.14 (1H, dd, J = 9.8, 2.1 Hz), 7.20-


7.25 (2H, m), 7.26-7.36 (4H, m), 7.36-7.41 (2H, m), 7.59 (1H, d, J = 10.0 Hz), 7.64 (1H, d, J = 1.8 Hz), 8.11 (1H, s), 8.25 (0.7H, s).

Example 176

1-[5-tert-Butyl-2-(4-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
 5 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea




a. 1-[5-tert-Butyl-2-(4-hydroxymethyl-phenyl)-2H-pyrazol-3-yl]-3-
 10 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea dichloromethane solvate
 (Intermediate 176a)



A red-brown solution of Intermediate 33a (883 mg, 2.10 mmol), Intermediate 81d (755 mg, 2.00 mmol) and DIPEA (0.44 mL, 2.5 mmol) in dry dioxane (20 mL) was stirred at 70°C for 16 h, and at 80°C for 5 h. The cooled solution was concentrated *in vacuo*, suspended in water (15 mL) and extracted with DCM (2 × 15 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown oil. Flash chromatography (silica 80 g, 4-8% MeOH in DCM) gave the *title compound* as a
 20 pale yellow foam (1.27 g, 87%). LCMS (Method 3): Rt 3.61 min, m/z 649 [MH⁺].

b. 1-[5-tert-Butyl-2-(4-formyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 176b)



5 To a solution of Intermediate 176a (1.10 g, 1.70 mmol) in DCM (25 mL) at 0°C was added Dess-Martin periodinane (791 mg, 1.86 mmol) and the resulting red-brown solution stirred at 0°C for 45 min. Sodium metabisulfite (920 mg), water (10 mL) and sat. aq. NaHCO₃ solution (10 mL) were added and the mixture stirred at RT until gas evolution ceased (15 min). The mixture was diluted with
10 water (25 mL) and extracted with DCM (2 × 25 mL). The combined organics were passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a glassy orange solid (1.10 g, 100%). LCMS (Method 3): Rt 3.91 min, m/z 647 [MH⁺].

c. 1-[5-tert-Butyl-2-(4-morpholin-4-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Example 176)

To a solution of Intermediate 176b (0.10 mmol) and morpholine (17.4 mg, 0.200 mmol) in DCM (3 mL) was added sodium triacetoxyborohydride (31.8 mg, 0.15 mmol) and the resulting yellow solution stirred at RT for 1 h. Water (3 mL)
20 was added and the mixture shaken. The aqueous was extracted with DCM (3 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a vivid yellow gum. Flash chromatography (silica 12g, 3.5-4.5% [2M NH₃ in MeOH] in DCM) gave a vivid yellow gum (70 mg). Prep HPLC (Gemini C18, 25-75% MeCN in water, 0.1% HCO₂H, 20 min) and concentration


of the desired fractions removed the MeCN. The aqueous was extracted with DCM (3 × 15 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a glassy orange solid (43.7 mg). Prep HPLC (Gemini C18, 25-50% MeCN in water, 20 min) and concentration of the desired

5 fractions removed the MeCN. The aqueous was extracted with DCM (3 × 5 mL), then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave the *title compound* as a white solid (25.4 mg, 35%). LCMS (Method 5): Rt 3.61 min, m/z 718.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.27 (9H, s), 1.47-1.55 (2H, m), 1.63-1.71 (2H, m), 1.76-1.95 (4H, m),

10 2.00-2.16 (2H, m), 2.37 (4H, t, J = 4.3 Hz), 2.90 (1H, ddd, J = 12.1, 9.0, 3.9 Hz), 3.16 (1H, dt, J = 12.2, 4.3 Hz), 3.29-3.34 (1H, m, obscured by water signal), 3.50 (2H, s), 3.56 (4H, t, J = 4.5 Hz), 4.81 (1H, td, J = 8.6, 5.6 Hz), 5.51 (1H, t, J = 4.3 Hz), 6.34 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.9, 2.2 Hz), 7.25-7.37 (4H, m), 7.43 (2H, d, J = 8.7 Hz), 7.45 (2H, d, J = 8.7 Hz), 7.64 (1H, dd, J =

15 9.8, 0.8 Hz), 7.69 (1H, dd, J = 2.2, 0.9 Hz), 8.08 (1H, s).


Example 177



1-[5-tert-Butyl-2-(4-dimethylaminomethyl-phenyl)-2H-pyrazol-3-yl]-3-
 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
 20 yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt


To an orange solution of Intermediate 176b (0.188 mmol) and dimethylamine solution (2M in THF, 0.19 mL, 0.38 mmol) in DCM (3 mL) was added sodium triacetoxyborohydride (60 mg, 0.28 mmol) and the mixture stirred at

RT for 1 h. Water (3 mL) was added and the mixture shaken. The aqueous was extracted with DCM (3 mL) then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. Flash chromatography (silica 12 g, 4-6.5% [2M NH₃ in MeOH] in DCM) gave a yellow solid (88 mg). Prep HPLC (Gemini C18, 25-50% MeCN in water, 0.1% HCO₂H, 20 min) and concentration *in vacuo* of the relevant fractions gave a pale yellow film. Trituration with diethyl ether gave the *title compound* as an off-white solid (48.4 mg, 36%). LCMS (Method 3): Rt 2.99 min, m/z 676 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.27 (9H, s), 1.45-1.56 (2H, m), 1.62-1.71 (2H, m), 1.75-1.96 (4H, m), 2.00-2.15 (2H, m), 2.17 (6H, s), 2.90 (1H, dd, J = 12.3, 9.1, 3.9), 3.16 (1H, dt, J = 12.3, 4.3 Hz), 3.28-3.34 (1H, m obscured by water peak), 3.44 (2H, s), 4.82 (1H, td, J = 8.6, 5.7 Hz), 5.51 (1H, t, J = 4.3 Hz), 6.34 (1H, s), 7.10 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.25-7.47 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, dd, J = 2.2, 0.9 Hz), 8.10 (1H, s), 8.16 (0.7H, s).

Example 178


1-{5-tert-Butyl-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4R)-4-[3-[(S)-2-methyl-piperidin-1-yl]-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt

5 To an orange solution of Intermediate 176b (0.188 mmol) and N-methyl piperazine (38 mg, 0.38 mmol) in DCM (3 mL) was added sodium triacetoxyborohydride (60 mg, 0.28 mmol) and the mixture stirred at RT for 1 h. Water (3 mL) was added and the mixture shaken. The aqueous was extracted with DCM (3 mL) then the combined organics passed through a hydrophobic frit and
10 concentrated *in vacuo* to leave a brown gum. Flash chromatography (silica 12 g, 5-8% [2M NH₃ in MeOH] in DCM) gave a yellow solid (47 mg). Prep HPLC (Gemini C18, 25-50% MeCN in water, 0.1% HCO₂H, 20 min) and concentration *in vacuo* of the relevant fractions gave a pale yellow film. Trituration with diethyl ether gave the *title compound* as an off-white solid (20.6 mg, 14%). LCMS
15 (Method 3): Rt 2.98 min, m/z 731 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.27 (9H, s), 1.45-1.56 (2H, m), 1.62-1.71 (2H, m), 1.75-1.95 (4H, m), 1.99-2.16 (2H, m), 2.14 (3H, s), 2.26-2.45 (8H, m), 2.90 (1H, ddd, J = 12.2, 9.0, 4.0 Hz), 3.16 (1H, dt, J = 12.2, 4.2 Hz), 3.28-3.34 (1H, m, obscured by water signal), 3.49 (2H, s), 4.81 (1H, td, J = 8.6, 5.6 Hz), 5.51 (1H, t, J = 4.3 Hz),
20 6.33 (1H, s), 7.10 (1H, d, J = 8.6 Hz), 7.20 (1H, dd, J = 9.9, 2.1 Hz), 7.25-7.47 (8H, m), 7.64 (1H, dd, J = 9.8, 0.8 Hz), 7.69 (1H, dd, J = 2.2, 0.9 Hz), 8.10 (1H, s), 8.17 (0.6H, s).

Example 179

1-{5-tert-Butyl-2-[4-(4-methyl-[1,4]diazepan-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea

To an orange solution of Intermediate 176b (0.188 mmol) and N-methyl homopiperazine (43 mg, 0.38 mmol) in DCM (3 mL) was added sodium triacetoxyborohydride (60 mg, 0.28 mmol) and the mixture stirred at RT for 1 h. Water (3 mL) was added and the mixture shaken. The aqueous was extracted with DCM (3 mL) then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. Flash chromatography (silica 12 g, 7-11% [2M NH₃ in MeOH] in DCM) gave the *title compound* as a yellow solid (102 mg, 73%). LCMS (Method 3): Rt 2.84 min, m/z 745 [MH⁺]. ¹H NMR (300 MHz, CDCl₃): 0.91 (3H, d, J = 6.2 Hz), 1.34 (9H, s), 1.40-1.55 (2H, m), 1.62-1.99 (7H, m), 2.15-2.02-2.14 (2H, m), 2.21-2.30 (1H, m), 2.33 (3H, s), 2.55-2.69 (8H, m), 2.90 (1H, ddd, J = 12.1, 9.3, 4.1), 3.04 (1H, dt, J = 12.1, 4.2 Hz), 3.23-3.32 (1H, m), 3.58 (2H, s), 5.09 (1H, td, J = 8.9, 5.2 Hz), 5.19 (1H, t, J = 3.9 Hz), 6.04 (1 H, br d, J = 8.6 Hz), 6.33 (1 H, s), 6.94 (1H, dd, J = 9.9, 2.1 Hz), 7.22 (1H, br s), 7.25-7.39 (7H, m), 7.43 (1H, d, J = 2.2 Hz), 7.47 (2H, m).

Example 180

1-{5-tert-Butyl-2-[4-(4-methoxy-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt

To an orange solution of Intermediate 176b (0.188 mmol) and 4-methoxypiperidine (43 mg, 0.38 mmol) in DCM (3 mL) was added sodium triacetoxyborohydride (60 mg, 0.28 mmol) and the mixture stirred at RT for 1 h.

10 Water (3 mL) was added and the mixture shaken. The aqueous was extracted with DCM (3 mL) then the combined organics passed through a hydrophobic frit and concentrated *in vacuo* to leave a brown gum. Flash chromatography (silica 12 g, 3.5-6% [2M NH₃ in MeOH] in DCM) gave a yellow solid (140 mg). Prep HPLC (Gemini C18, 25-50% MeCN in water, 0.1% HCO₂H, 20 min, × 2) and


15 concentration *in vacuo* of the relevant fractions gave a pale yellow film. Trituration with diethyl ether gave the *title compound* as an off-white solid (70.5 mg, 48%).

LCMS (Method 5): Rt 3.69 min, m/z 746.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.27 (9H, s), 1.37-1.56 (4H, m), 1.62-1.71 (2H, m), 1.75-1.95 (6H, m), 2.00-2.16 (4H, m), 2.62-2.69 (2H, m), 2.90 (1H, ddd, J =

20 12.2, 9.1, 4.1 Hz), 3.12-3.18 (2H, m), 3.20 (3H, s), 3.29-3.34 (1H, m), 3.49 (2H, s), 4.82 (1H, td, J = 8.6, 5.6 Hz), 5.51 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.10 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.8, 2.2 Hz), 7.25-7.46 (8H, m), 7.64 (1H, dd, J = 9.8,


0.8 Hz), 7.69 (1H, dd, $J = 2.2, 0.9$ Hz), 8.10 (1H, s), 8.16 (0.75H, s).

Example 181




**1-[5-tert-Butyl-2-(4-pyrrolidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
5 {(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-
yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 176b (0.188 mmol) and pyrrolidine (31.1 μ L, 0.38 mmol) in DCM (3 mL), sodium triacetoxyborohydride (59.8 mg, 0.28 mmol) was added. The mixture was stirred at RT for 1 h. Water was added and the
10 mixture extracted with DCM. The combined organics were passed through a phase separator and concentrated to dryness. The resulting residue was purified by RP-HPLC (C18, 18 mL/min, 20 – 85% MeCN in H₂O + 0.1% HCO₂H) and the relevant fractions combined and concentrated to dryness. The *title product* was isolated as a light orange solid (49.5 mg). LCMS (Method 5): Rt 3.66 min, m/z
15 702.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, $J = 6.3$ Hz), 1.27 (9H, s), 1.50 (2H, m), 1.69 (6H, m), 1.75-1.95 (4H, m), 1.98-2.17 (2H, m), 2.47 (4H, m, obscured by solvent), 2.90 (1H, m), 3.15 (1H, m, obscured by solvent), 3.63 (2H, s, obscured by solvent), 4.81 (1H, m), 5.51 (1H, t, $J = 4.3$ Hz), 6.33 (1H, s), 7.09 (1H, d, $J = 8.9$ Hz), 7.19 (1H, dd, $J = 9.6, 2.2$ Hz), 7.24-7.38 (4H, m), 7.43 (4H, m),
20 7.64 (1H, d, $J = 9.6$ Hz), 7.69 (1H, d, $J = 1.3$ Hz), 8.09 (1H, s), 8.16 (1.4H, s).

Example 182


**1-[5-tert-Butyl-2-(4-piperidin-1-ylmethyl-phenyl)-2H-pyrazol-3-yl]-3-
 5 { (1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**

To a solution of Intermediate 176b (0.188 mmol) and piperidine (37.2 μ L, 0.38 mmol) in DCM (3 mL), sodium triacetoxyborohydride (59.8 mg, 0.28 mmol) was added. The mixture was stirred at RT for 1 h. Water was added and the
 10 mixture extracted with DCM. The combined organics were passed through a phase separator and concentrated to dryness. The resulting residue was purified by RP-HPLC (C18, 18 mL/min, 20 – 85% MeCN in H₂O + 0.1% HCO₂H) and the relevant fractions combined and concentrated to dryness. The *title product* was isolated as an off white solid (14 mg). LCMS (Method 5): Rt 3.71 min, m/z 716.6
 15 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.4 Hz), 1.27 (9H, s), 1.36 (2H, m), 1.47 (6H, m), 1.67 (2H, m), 1.76-1.96 (4H, m), 1.98-2.16 (2H, m), 2.33 (4H, m), 2.90 (1H, m), 3.15 (1H, m, obscured by solvent), 3.46 (2H, s, obscured by solvent), 4.81 (1H, m), 5.51 (1H, t, J = 4.1 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 10.0, 2.2 Hz), 7.24-7.46 (8H, m), 7.64 (1H, d, J = 10.0 Hz), 7.69 (1H, d, J = 1.5 Hz), 8.09 (1H, s), 8.17 (1H, s).
 20

Example 183


1-{5-tert-Butyl-2-[4-(4-fluoro-piperidin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

To a solution of Intermediate 176b (0.188 mmol) and 4-fluoropiperidine (38.8 mg, 0.38 mmol) in DCM (3 mL), sodium triacetoxyborohydride (59.8 mg, 0.28 mmol) was added. The mixture was stirred at RT for 1 h. Water was added and the mixture extracted with DCM. The combined organics were passed through a phase separator and concentrated to dryness. The resulting residue was purified by RP-HPLC (C18, 18 mL/min, 20 – 85% MeCN in H₂O + 0.1% HCO₂H) and the relevant fractions combined and concentrated to dryness. The *title product* was isolated as an off white solid (31 mg). LCMS (Method 5): Rt 3.71 min, m/z 734.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.1 Hz), 1.27 (9H, s), 1.50 (2H, m), 1.67 (4H, m), 1.76-1.96 (6H, m), 1.98-2.17 (2H, m), 2.31 (2H, m), 2.54 (m, obscured by solvent), 2.90 (1H, m), 3.16 (1H, m, obscured by solvent), 3.51 (2H, s, obscured by solvent), 4.66 (1H, dm, J = 48 Hz), 4.82 (1H, m), 5.51 (1H, t, J = 4.2 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.7 Hz), 7.19 (1H, dd, J = 9.8, 2.3 Hz), 7.24-7.50 (8H, m), 7.63 (1H, d, J = 10.2 Hz), 7.69 (1H, d, J = 1.4 Hz), 8.10 (1H, s), 8.16 (1H, s).

Example 184

1-(5-tert-Butyl-2-{4-[(ethyl-methyl-amino)-methyl]-phenyl}-2H-pyrazol-3-yl)-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

To a solution of Intermediate 176b (0.188 mmol) and N-ethylmethylaniline (32.3 μ L, 0.38 mmol) in DCM (3 mL), sodium triacetoxyborohydride (59.8 mg, 0.28 mmol) was added. The mixture was stirred at RT for 1 h. Water was added and the mixture extracted with DCM. The combined organics were passed through a phase separator and concentrated to dryness. The resulting residue was purified by RP-HPLC (C18, 18 mL/min, 20 – 85% MeCN in H₂O + 0.1% HCO₂H) and the relevant fractions combined and concentrated to dryness. The *title product* was isolated as an off white solid (26.1 mg). LCMS (Method 5): Rt 3.65 min, m/z 690.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.4 Hz), 1.02 (3H, t, J = 7.1 Hz), 1.27 (9H, s), 1.50 (2H, m), 1.66 (2H, m), 1.74-1.97 (4H, m), 1.98-2.11 (2H, m), 2.12 (3H, s), 2.40 (2H, q, J = 7.1 Hz), 2.90 (1H, m), 3.16 (1H, m, obscured by solvent), 3.49 (2H, s, obscured by solvent), 4.81 (1H, m), 5.51 (1H, t, J = 4.6 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.6 Hz), 7.19 (1H, dd, J = 9.6, 2.1 Hz), 7.24-7.47 (8H, m), 7.64 (1H, d, J = 10.2 Hz), 7.69 (1H, d, J = 1.6 Hz), 8.09 (1H, s), 8.17 (1H, s).


Example 185

1-[5-tert-Butyl-2-(4-[(2-methoxy-ethyl)-methyl-amino]-methyl)-phenyl]-2H-pyrazol-3-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea formate salt

To a solution of Intermediate 176b (0.188 mmol) and N-(2-methoxyethyl)methyl amine (32.7 mg, 0.38 mmol) in DCM (3 mL), sodium triacetoxyborohydride (59.8 mg, 0.28 mmol) was added. The mixture was stirred at RT for 1 h. Water was added and the mixture extracted with DCM. The combined organics were passed through a phase separator and concentrated to dryness. The resulting residue was purified by RP-HPLC (C18, 18 mL/min, 20 – 85% MeCN in H₂O + 0.1% HCO₂H) and the relevant fractions combined and concentrated to dryness. The *title product* was isolated as an off white solid (23.9 mg). LCMS (Method 5): Rt 3.68 min, m/z 720.5 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.3 Hz), 1.27 (9H, s), 1.50 (2H, m), 1.66 (2H, m), 1.74-1.96 (4H, m), 1.98-2.16 (2H, m), 2.17 (3H, s), 2.54 (2H, t, J = 6.0 Hz, obscured by solvent), 2.90 (1H, m), 3.16 (1H, m, obscured by solvent), 3.23 (3H, s, obscured by solvent), 3.45 (2H, t, J = 6.0 Hz, obscured by solvent), 3.54 (2H, s, obscured by solvent), 4.82 (1H, m), 5.51 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.09 (1H, d, J = 8.8 Hz), 7.19 (1H, dd, J = 9.7, 2.2 Hz), 7.24-7.50 (8H, m), 7.64 (1H, d, J = 10.0 Hz), 7.69 (1H, d, J = 1.5 Hz), 8.09 (1H, s), 8.17 (1H, s).

Example 186


1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl-3-[(1S,4S)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea



5


a. 2-[3-(5-Amino-3-tert-butyl-pyrazol-1-yl)-phenoxy]-ethanol

(Intermediate 186a)




A solution of Intermediate 95a (1.2 g, 5.2 mmol) and triphenyl phosphine (2.72 g, 10.4 mmol) in THF (30 mL), under an atmosphere of argon was treated with 2-(tetrahydro-pyran-2-yloxy)-ethanol (1.05 mL, 7.78 mmol), followed by the dropwise addition of diisopropylazodicarboxylate (1.64 mL, 10.4 mmol). The reaction mixture was then stirred at RT for 1 h. The reaction mixture was eluted on a SCX-2 cartridge using MeOH and 2M NH₃ in MeOH. The basic fractions were evaporated under reduced pressure and the product was purified by FCC, using EtOAc/DCM 0-60% to afford the *title compound* as a brown gum (1.0 g, 80%). LCMS (Method 3): Rt 2.27 min, m/z 276 [MH⁺].

b. {5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-
carbamic acid 2,2,2-trichloro-ethyl ester (Intermediate 148e)



A solution of Intermediate 186a (1.0 g, 3.63 mmol) in EtOAc (20 mL) was
5 treated with aqueous NaOH (1M, 6.53 mmol), followed by 2,2,2-trichloroethyl
chloroformate (0.529 mL, 3.85 mmol) and the reaction mixture was stirred at RT
for 1 h. The mixture was partitioned between EtOAc (20 mL) and water (20 mL).
The layers were separated and the aqueous layer was extracted with a further 20
mL EtOAc. The combined organic layers were dried (Na₂SO₄), filtered and
10 concentrated *in vacuo*. The residue was purified by FCC, using 0-40% EtOAc in
DCM. The combined fractions were evaporated under reduced pressure and the
product was triturated with DCM. The solid was filtered and washed with Diethyl
ether to afford the *title compound* as a white solid (0.930 g, 58%). LCMS (Method
3): Rt 4.01 min, m/z 450 [M].


15 c. (1S,4S)-4-[3-((S)-2-Methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-
a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-ylamine (Intermediate
186b)



To a solution of Intermediate B (232 mg, 1.6 mmol) in dry DMF (10 mL)
20 under N₂ was added sodium (60% dispersion in oil, 320 mg, 8.0 mmol) and the
resulting opaque brown solution was stirred at RT for 45 min (CARE: gas
evolution). A solution of Intermediate 81c (342 mg, 1.46 mmol) in dry DMF (7
mL) was added and the dark brown solution stirred at 60°C under N₂ for 90 min.

The cooled solution was concentrated *in vacuo*, redissolved in MeOH (3 mL) and then applied to an SCX-2 cartridge (20 g), washing with MeOH (50 mL). The product was eluted with 2M NH₃ in MeOH (50 mL); concentration *in vacuo* gave a dark brown solid. FCC, using 0-10% [2M NH₃ in MeOH] in DCM, gave the *title compound* as a pale brown foam (262 mg, 45%). LCMS (Method 3): Rt 2.30 min, m/z 378 [MH⁺].

d. 1-{5-tert-Butyl-2-[3-(2-hydroxy-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4S)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl]-urea(Intermediate 186c)



10

A mixture of Intermediate 148e (313 mg, 0.7 mmol), Intermediate 186b (262 mg, 0.7 mmol) and DIPEA (182 μ L, 1.05 mmol) in dioxane (4 mL) was heated at 70°C for 20 h. The reaction mixture was cooled to RT and eluted directly on SCX-2 using MeOH and 2M NH₃ in MeOH. The basic fractions were evaporated under reduced pressure. The residue was purified by FCC, using 0-10% [2M NH₃ in MeOH] in DCM, to afford the *title compound* as a yellow glass (293 mg). LCMS (Method 3): Rt 3.63 min, m/z 679 [MH⁺].

15

e. 1-{5-tert-Butyl-2-[3-(2-dimethylamino-ethoxy)-phenyl]-2H-pyrazol-3-yl}-3-[(1S,4S)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydronaphthalen-1-yl]-urea (Example 186)

20


To a solution of Intermediate 186c (293 mg, 0.44 mmol) and DIPEA (304 μ L, 1.76 mmol) in DCM (10 mL) was added methanesulfonyl chloride (103 μ L, 1.33 mmol). The reaction mixture was stirred at RT for 1 hour. The mixture was concentrated *in vacuo*. The residue was taken up in THF (2 mL) and

dimethylamine solution in THF (2M, 3.3 mL) was added. The reaction mixture was heated at 60°C in a sealed tube for 72 h. The reaction mixture was cooled to RT and concentrated *in vacuo*. The product was purified by HPLC using a Gemini C18 column and using a gradient of 20-35% +0.1% Formic acid over 10 minutes.


- 5 The combined fractions were evaporated under reduced pressure and crystallised by MeCN/Et₂O to afford the *title compound* as a white solid (90 mg, 30%). LCMS (Method 3): Rt 3.01 min, m/z 706 [MH⁺]. ¹H NMR (300 MHz, d₆-DMSO): 0.91 (3H, d, J = 6.7 Hz), 1.23 (9H, s), 1.46-1.54 (2H, m), 1.63-1.72 (2H, m), 1.75-1.76 (2H, m), 2.00-2.18 (2H, m), 2.43 (6H, s), 2.66 (2H, t, J 5.8 Hz), 2.85-2.95 (2H, m),
10 3.12-3.20 (4H, m), 4.10 (2H, t, J = 5.4 Hz), 4.78-4.86 (1H, m), 5.52 (1H, t, J = 4.0 Hz), 6.33 (1H, s), 6.95-6.99 (2H, m), 7.06-7.14 (2H, m), 7.19 (1H, dd, J = 10.0Hz), 7.25-7.32 (3H, m), 7.33-7.42 (2H, m), 7.58 (1H, d, J = 10.6 Hz), 7.63 (1H, d, J = 1.8 Hz), 8.17 (1H, s).

Example 187

- 15 **1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-[(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt**



a. 1-{3-tert-Butyl-1'-[2-(tetrahydro-pyran-2-yloxy)-ethyl]-1'H-[1,4']bipyrazolyl-5-yl}-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 187a)




5

A mixture of Intermediate 81d (240 mg, 0.64 mmol), Intermediate 108b (325 mg, 0.64 mmol) and DIPEA (221 μ L, 1.27 mmol) in dioxane (6 mL) was stirred at 80°C overnight. After cooling, the reaction mixture was partitioned between water and EtOAc. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% MeOH in DCM to afford the *title compound* (377 mg, 80%). LCMS (Method 4): Rt 3.48 min, m/z 737 [MH^+].

10

b. 1-[3-tert-Butyl-1'-(2-hydroxy-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea (Intermediate 187b)

15




20

To a solution of Intermediate 187a (377 mg, 0.51 mmol) in MeOH (5 mL) was added pyridinium p-toluenesulfonate (257 mg, 1.02 mmol) and the reaction mixture was heated at 55°C for 3 h. The resultant mixture was poured

into water and a saturated aqueous solution of NaHCO_3 was added. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 3-10% MeOH in DCM, to give the *title compound* (250 mg, 75%). LCMS (Method 1): Rt 3.09 min, m/z 653 [MH^+].

c. **Methanesulfonic acid 2-[3-tert-butyl-5-(3-((1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-ureido)-[1,4']bipyrazolyl-1'-yl]-ethyl ester**
(Intermediate 187c)




To an ice-bath cooled solution of Intermediate 187b (250 mg, 0.38 mmol) in DCM (4 mL) was added DIPEA (267 μL , 1.53 mmol) followed by methanesulfonyl chloride (60 μL , 0.77 mmol). The reaction mixture was stirred for 2 h and then quenched with water. The aqueous phase was extracted with DCM (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo* to afford the *title compound* (Quantitative). Product used in the following step without further purification. LCMS (Method 4): Rt 3.25 min, m/z 732 [MH^+].

d. 1-[3-tert-Butyl-1'-(2-dimethylamino-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt (Example 187)

5 To a solution of Intermediate 187c (0.19 mmol) in THF (3 mL) was added dimethylamine (2M in MeOH, 1.9 mL, 3.8 mmol) and the reaction stirred at 50°C in a sealed vial overnight. The crude reaction mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried
10 (MgSO₄) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 0-10% [2M NH₃ in MeOH] in DCM followed by MDAP (Method 7) purification, to give the *title compound* (46 mg, 33%). LCMS (Method 5): Rt 3.50 min, m/z 680 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 0.87 (3H, d, J = 6.4 Hz), 1.20 (9H, s), 1.43-1.51 (2H, m), 1.58-1.69 (2H, m), 1.70-1.88 (3H, m), 1.88-1.96 (1H, m), 1.96-2.12 (2H, m), 2.13 (6H, s), 2.63 (2H, t, J = 6.5 Hz), 2.87 (1H, ddd, J = 12.6, 9.1, 3.9 Hz), 3.12 (1H, dt, J = 12.1, 4.2 Hz), 4.16 (2H, t, J = 6.5 Hz), 4.80 (1H, td, J = 8.5, 5.8 Hz), 5.48 (1H, t, J = 4.3 Hz), 6.22 (1H, s), 7.12 (1H, d, J = 8.5 Hz), 7.15 (1H, dd, J = 9.9, 2.2 Hz), 7.21-7.26 (1H, m), 7.28-7.36 (3H, m), 7.58 (1H, s),
15 7.60 (1H, d, J = 9.5 Hz), 7.66 (1H, d, J = 2.1 Hz), 7.96 (1H, s), 8.00 (1H, s), 8.13 (1H, s).


Example 188



1-[3-tert-Butyl-1'-(2-morpholin-4-yl-ethyl)-1'H-[1,4']bipyrazolyl-5-yl]-3-{(1S,4R)-4-[3-((S)-2-methyl-piperidin-1-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea partial formate salt

5 To a solution of Intermediate 187c (0.19 mmol) in THF (3 mL) was added morpholine (166 μ L, 1.9 mmol) and the reaction stirred at 50°C overnight. The crude reaction mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc (x 3) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated *in vacuo*. The resultant residue was purified by FCC on silica, using a gradient of 2.5-10% [2M NH_3 in MeOH] in DCM followed by MDAP (Method 7) purification, to give the *title compound* (49 mg, 35%). LCMS (Method 5): Rt 3.53 min, m/z 722 [MH^+]. ^1H NMR (400 MHz, d_6 -DMSO): 0.87 (3H, d, J = 6.4 Hz), 1.20 (9H, s), 1.43-1.51 (2H, m), 1.58-1.69 (2H, m), 1.70-1.88 (3H, m), 1.88-1.96 (1H, m), 1.96-2.12 (2H, m), 2.37 (4H, t, J = 4.5 Hz), 2.69 (2H, t, J = 6.5 Hz), 2.87 (1H, ddd, J = 12.6, 9.1, 3.9 Hz), 3.12 (1H, dt, J = 12.1, 4.2 Hz), **1H UNDER SOLVENT**, 3.50 (4H, t, J = 4.6 Hz), 4.20 (2H, t, J = 6.5 Hz), 4.80 (1H, td, J = 8.5, 5.8 Hz), 5.48 (1H, t, J = 4.3 Hz), 6.22 (1H, s), 7.11 (1H, d, J = 8.5 Hz), 7.15 (1H, dd, J = 9.9, 2.2 Hz), 7.21-7.26 (1H, m), 7.28-7.35 (3H, m), 7.60 (1H, d, J = 9.8 Hz), 7.60 (1H, s), 7.66 (1H, d, J = 2.1 Hz), 7.96 (1H, s), 8.02 (1H, s), 8.12 (0.5H, s).

Example 189



1-{5-tert-Butyl-2-[3-(4-methyl-piperazin-1-ylmethyl)-phenyl]-2H-pyrazol-3-yl}-3-{(1S,4R)-4-[3-((S)-1-methyl-pyrrolidin-2-yl)-[1,2,4]triazolo[4,3-a]pyridin-6-yloxy]-1,2,3,4-tetrahydro-naphthalen-1-yl}-urea formate salt

5 A mixture of Intermediate 149b (87 mg, 0.12 mmol) and 1-methylpiperazine (68 μ L, 0.61 mmol) in THF (2 mL) was stirred at 60°C for 18 h. The reaction mixture was cooled to RT, diluted with DCM (5 mL) and washed with water (5 mL). The aqueous layer was extracted with DCM (5 mL). The combined organic extracts were washed with water (2 x 5 mL) and passed through a phase separator. The filtrate was concentrated *in vacuo* and the resultant residue was purified by MDAP (Method 7) to afford the *title compound* (23 mg, 27%). LCMS (Method 5): Rt 2.69 min, m/z 717.6 [MH⁺]. ¹H NMR (400 MHz, d₆-DMSO): 1.28 (9H, s), 1.84-2.11 (6H, m), 2.13 (6H, br s), 2.15-2.26 (2H, m), 2.26-2.46 (8H, m), 3.13 (1H, m, obscured by water), 3.50 (2H, s, obscured by water), 3.99 (1H, t, J = 8.4 Hz), 4.82 (1H, m), 5.39 (1H, t, J = 4.3 Hz), 6.33 (1H, s), 7.06 (1H, d, J = 8.8 Hz), 7.24-7.48 (9H, m), 7.75 (1H, d, J = 9.7 Hz), 8.10 (1H, s), 8.19 (1.9H, br s), 8.24 (1H, d, J = 1.7 Hz).

Biological assays

20 P38alpha enzyme inhibition assay

The inhibitory activity of compounds was determined using an Alphascreen® (Perkin Elmer) based kinase activity assay. Kinase reactions consisted of 25mM HEPES pH 7.5, 10mM MgCl₂, 100 μ M Na₃VO₄, 2mM DTT, 0.05mg/ml Tween 20, 100pM p38alpha (Invitrogen, PV3304), 1% DMSO and 0.3 μ g/ml ATF-2 fusion protein (New England Biolabs, 9224). Compounds were incubated under these conditions for 2 hours, at 25°C, prior to the initiation of the kinase activity by the addition of the 250 μ M ATP. Reaction volumes were 20 μ L. After 1hr at 25°C reactions were stopped by the

adding 10uL of 25mM HEPES pH 7.5 containing 62.5mM EDTA, 0.05% Triton X-100, 10% BSA and 0.83ng/uL anti-phospho-ATF2 antibody (Abcam, ab28812). Detection was performed by measuring luminescence following the addition of Alphascreen Donor beads (Perkin Elmer 6765300) and Protein A

5 Alphascreen Acceptor beads (Perkin Elmer 6760137), both at a final concentration of 20ug/ml. IC₅₀ values were determined from concentration-response curves.

Results are shown in the following Table:

Example	p38 α inhibition
1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 148, 154, 155, 156, 157, 158, 159, 160, 162, 163, 165, 166, 167, 168, 169, 170, 171, 176, 178, 181, 182, 183, 184, 185	++++
21	+++
32	++

10 In the table above, p38 α binding potencies (IC₅₀ values) are indicated as follows: 7000-500nM ‘+’; 500-100nM ‘++’; 100-10nM ‘+++’; <10nM ‘++++’.

LPS-stimulated PBMC TNF α release assay

Peripheral Blood Mononuclear Cells (PBMCs) were isolated from

15 healthy human volunteer blood using a standard density gradient

centrifugation technique. Citrated blood was placed onto Histopaque™ and centrifuged. The PBMCs were removed from the density gradient interface and washed in phosphate buffered saline (PBS). The PBMCs were suspended in RPMI 1640 medium (without serum), dispensed into a 96-well plate and

5 incubated at 37°C for 3h in a humidified incubator. After incubation, the medium was replaced (with medium containing 1% foetal bovine serum) and the plate incubated at 37°C, for 1h, in the presence of test compound or the appropriate vehicle. LPS (10ng/ml), or an appropriate vehicle control, was then added to the cells and the plate returned to the incubator for 18h.

10 Cell-free supernatants were removed and assayed for TNFα levels using an ELISA kit from R&D Systems.

A dose response curve to each test compound was performed and the effect of compound in each experiment was expressed as a percentage inhibition of the control TNFα release. Dose response curves were plotted and

15 compound potency (IC₅₀) was determined. Compounds were tested in at least three separate experiments.

Results are shown in the following Table:

Example numbers	p38α inhibition
3, 5, 8, 10, 12, 13, 15, 16, 18, 19, 20, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 46, 47, 48, 49, 50, 52, 53, 54, 55, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 92, 93, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 120, 121, 122, 123, 124, 125, 126, 127, 128, 130	++++
1, 2, 11, 14, 21, 57, 94, 119, 129	+++
4, 7, 9, 22	++

In the table above, p38α potencies (IC₅₀ values) are indicated as follows:

>1000 nM ‘+’; 1000-100nM ‘++’; 100-10nM ‘+++’; <10nM ‘++++’. All compounds tested exhibited IC₅₀ values <1000nM.

Pre-clinical mouse model of COPD – Tobacco smoke induced pulmonary inflammation.

5 Previous studies have established that the number of inflammatory cells recovered by bronchoalveolar lavage (BAL) is significantly elevated 24h following the final of four consecutive daily tobacco smoke (TS) exposures. This timepoint was used in the studies reported here.

10 Protocols for the exposure of mice to TS, obtaining bronchoalveolar lavage fluid (BALF) and preparation of cytopsin slides for differential cell counts are as outlined below.

Daily exposure of mice to TS for 4 consecutive days

15 In this exposure protocol, mice were exposed in groups of 5 in individual clear polycarbonate chambers (27 cm x 16 cm x 12 cm). The TS from the cigarettes was allowed to enter the exposure chambers at a flow rate of 100 ml/min. In order to minimise any potential problems caused by repeated exposure to a high level of TS, the exposure of the mice to TS was increased gradually over the exposure period to a maximum of 6 cigarettes. The exposure schedule used over the four days was as follows:

20 Day 1: 5 cigarettes (approximately 25min exposure)
Day 2: 7 cigarettes (approximately 35min exposure)
Day 3: 9 cigarettes (approximately 45min exposure)
Day 4: 9 cigarettes (approximately 45min exposure)

25 A further group of mice were exposed to air on a daily basis for equivalent lengths of time as controls (no TS-exposure).

Bronchoalveolar lavage (BAL) analysis

Bronchoalveolar lavage was performed as follows: the trachea was cannulated using a 10mm long Luer-fitting stainless steel cannula. Phosphate

buffered saline (PBS) was used as the lavage fluid. A volume of 0.4ml was gently instilled and withdrawn 3 times, using a 1ml syringe and then placed in an Eppendorf tube and kept on ice prior to subsequent determinations.

Cell counts:

5 Lavage fluid was separated from cells by centrifugation and the supernatant decanted and frozen for subsequent analyses. The cell pellet was re-suspended in a known volume of PBS and total cell numbers calculated by counting a stained (Turks stain) aliquot under a microscope using a haemocytometer.

10 Differential cell counts were performed as follows:

 The residual cell pellet was diluted to approximately 10^5 cells per ml. A volume of 500 μ l was placed in the funnel of a cytopsin slide and centrifuged for 6min at 800rpm, RCF = 72.26 x g (Shandon Cytospin 3). The slide was air-dried and stained using Wrights/Giemsa stain as per the proprietary
15 instructions. When dried and cover-slipped, differential cell counts were performed using light microscopy. Approximately four hundred cells were counted by an unbiased operator using light microscopy. Cells were differentiated using standard morphometric techniques.

Drug Treatment

20 Rodents such as mice and rats are obligate nose breathers, thus oral delivery of test materials (such as therapeutic agents) for inhalation will not produce good lung exposure. As a consequence, delivery of therapeutic agents to the lungs in rodents is generally achieved by intranasal, intratracheal or inhalation by either nose-only or whole body aerosol exposure.

25 Nose-only or whole body aerosol exposure methods utilise large amounts of test material and are generally reserved for inhalation toxicology studies rather than more routine pharmacological efficacy studies. Intratracheal administration is a very efficient delivery method as almost all of

the test material is delivered to the lungs but is an invasive technique. For studies in the mouse particularly, it is also technically demanding as the diameter of the trachea is small. The intranasal route is less invasive than the intratracheal route and so is particularly suitable for repeat dosing studies such as the four day mouse model described. Following intranasal administration, ~50% of the dose administered is delivered to the lungs (Eyles JE, Williamson ED and Alpar HO. 1999, Int J Pharm, 189(1):75-9).

As a surrogate route for oral inhalation, mice were dosed intra-nasally with vehicle (0.2% tween 80 in saline) containing test compound. The control groups of mice received vehicle 1hr prior to being exposed to air or TS.

Data management and statistical analysis:

All results are presented as individual data points for each animal and the mean value was calculated for each group. Since tests for normality were positive, the data were subjected to a one-way analysis of variance test (ANOVA), followed by a Bonferroni correction for multiple comparisons in order to test for statistically significant differences between treatment groups. A “p” value of < 0.05 was considered to be statistically significant. Percentage inhibitions were automatically calculated within Excel spreadsheets for the cell data using the formula below:

$$\% \text{ Inhibition} = \left(1 - \frac{\text{Treatment group result} - \text{air group result}}{\text{TS vehicle group result} - \text{air group result}} \right) \times 100$$

Inhibition data for other parameters were calculated manually using the above formula.

Example 5 of the invention was tested in the above reported model and results are herebelow reported.

As illustrated in Figure 1, Example 5 significantly inhibited the BAL cell influx induced by TS at 3, 30 or 100 µg/kg when administered by the intranasal route. Similar findings were observed with BAL neutrophils (Figure

2). The results demonstrate a clear anti-inflammatory effect in the lungs of mice exposed to TS.

Brief description of the figures


Figure 1. is a graph that illustrates the effect of intranasal
5 administration to laboratory mice with vehicle (0.2% tween 80 in saline),
Example 5 (3 µg/kg), Example 5 (30 µg/kg) or Example 5 (100 µg/kg) on the
number of BAL cells induced by tobacco smoke 24 hours post the final
exposure.

Figure 2. is a graph that illustrates the effect of intranasal
10 administration to laboratory mice with vehicle (0.2% tween 80 in saline),
Example 5 (3 µg/kg), Example 5 (30 µg/kg) or Example 5 (100 µg/kg) on the
number of BAL neutrophils induced by tobacco smoke 24 hours post the final
exposure.

CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:

5



wherein;

W is a heteroatom selected from N or O, wherein N is substituted with hydrogen, C₁-C₆ alkyl or C₃-C₅ cycloalkyl;


10 **Y** is selected in the group consisting of: a group -S(O)_p- wherein **p** is 0, 1 or 2; a group -O(CR³R⁴)_n-; a group; -(CR⁵R⁶)_n-; a group -NR⁷-; a group -OC(O)-; a group -OC(O)NH-; and a group -OC(O)O-;

R³, R⁴, R⁵ and R⁶ are each independently hydrogen, fluorine or C₁-C₆ alkyl, or, respectively, **R³** and **R⁴**, or **R⁵** and **R⁶** may form together with the
15 carbon atom to which they are attached a saturated 3-6 membered carbocyclic monocyclic ring optionally substituted by a group C₁-C₆ alkyl, hydroxyl or halo;


n is 0, 1, 2 or 3;

R⁷ is hydrogen, C₁-C₆ alkyl, or C₃-C₇ cycloalkyl wherein such C₁-C₆
20 alkyl or C₃-C₇ cycloalkyl are optionally substituted by a group C₁-C₃ alkyl, C₃-C₆ cycloalkyl, hydroxyl, cyano or halo;


R¹ is a group selected from (IIa) - (IIc):



(IIa)



(IIb)



(IIc)

R^8 and R^9 are each independently hydrogen or C_1 - C_6 alkyl, or R^8 and R^9 may form together with the nitrogen atom to which they are attached a 5-11-membered saturated monocyclic or a fused or spiro bicyclic ring system optionally containing a further heteroatom which is oxygen or nitrogen, said nitrogen atom being optionally substituted by C_1 - C_6 alkyl; wherein such C_1 - C_6 alkyl groups may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, hydroxyl or halo;

X^1 , X^2 , X^3 , X^4 and X^5 are each independently a carbon atom, a nitrogen atom, a group $-(CH)-$ or a group $-NH-$; such that each combination thereof forms an aromatic ring system;

R^{10} is selected from a group consisting of: Hydrogen, $-NR^A R^B$, $-N(R^C)(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-(C_1-C_6\text{alkylene})-NR^A R^B$, $-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-O-(C_2-C_6\text{alkylene})-NR^A R^B$, $-O-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S-(C_2-C_6\text{alkylene})-NR^A R^B$, $-S-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-C(O)N(R^C)-(C_2-C_6\text{alkylene})-OR^D$, $-C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)C(O)NR^A R^B$, $-C(O)NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-(C_2-C_6\text{alkylene})-OR^D$, $-(C_3-C_7\text{cycloalkylene})-OR^D$, $-O-(C_2-C_6\text{alkylene})-OR^D$, $-O-(C_3-C_7\text{cycloalkylene})-OR^D$, $-S-(C_2-C_6\text{alkylene})-OR^D$, $-S-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)S(O)_2-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)S(O)_2-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-S(O)_2N(R^C)-(C_2-C_6\text{alkylene})-OR^D$, $-S(O)_2N(R^C)-(C_3-C_7\text{cycloalkylene})-OR^D$, $-N(R^C)S(O)_2-(C_2-C_6\text{alkylene})-OR^D$, $-N(R^C)S(O)_2-(C_3-C_7\text{cycloalkylene})-OR^D$, $-S(O)_2N(R^A R^B)$, $-N(R^C)S(O)_2R^D$, $-N(R^C)C(O)R^C$, $-OR^C$, $-SR^C$, $-(C_3-C_7\text{heterocycloalkyl})$, $(C_5-C_7\text{heterocycloalkyl})-(C_1-C_6\text{alkyl})$, $(C_5-C_7\text{heterocycloalkyl})(C_3-$

C₆cycloalkyl)-, and C₃-C₇ heterocycloalkylcarbonyl; wherein any of such C₁-C₆alkyl, C₃-C₆cycloalkyl, -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-C₇cycloalkylene)-, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), (C₅-C₇ heterocycloalkyl)-(C₃-C₆ cycloalkyl) and (C₃-C₇heterocycloalkyl)carbonyl portion in the above listed groups may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, hydroxyl or halo;

R¹¹ is linked to X⁴ and is selected from a group consisting of:

Hydrogen; -CN; C₁-C₆ alkyl which is substituted by a group selected from -CN, -OR^C, -SR^C, halo; C₃-C₆cycloalkyl which is substituted by a group selected from C₁-C₄ alkyl, -CN, -OR^C, -SR^D, halo; -NR^AR^B, -N(R^C)(C₂-C₆alkylene)-NR^AR^B, -N(R^C)(C₃-C₇cycloalkylene)-NR^AR^B, -(C₁-C₆alkylene)-NR^AR^B, -(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₂-C₆alkylene)-NR^AR^B, -O-(C₃-C₇cycloalkylene)-NR^AR^B, -S-(C₂-C₆alkylene)-NR^AR^B, -S-(C₃-C₇cycloalkylene)-NR^AR^B, -N(R^C)C(O)-(C₁-C₆alkylene)-NR^AR^B, -N(R^C)C(O)-(C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-OR^D, -C(O)N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)C(O)N(R^AR^B), -C(O)N(R^AR^B), -N(R^C)C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -N(R^C)C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₂-C₆alkylene)-OR^D, -O-(C₃-C₇cycloalkylene)-OR^D, -S-(C₂-C₆alkylene)-OR^D, -S-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₁-C₆alkylene)-NR^AR^B, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-OR^D, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₂-C₆alkylene)-OR^D, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-OR^D, -S(O)₂N(R^AR^B), -N(R^C)S(O)₂R^D, -N(R^C)C(O)R^C, OR^C, SR^C, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl), (C₅-C₇heterocycloalkyl)(C₃-C₆cycloalkyl) and (C₃-C₇

heterocycloalkyl)carbonyl, wherein any of such C₁-C₆alkyl, C₃-C₆cycloalkyl, -
 (C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-C₇cycloalkylene)-, -(C₃-
 C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl), (C₅-C₇
 heterocycloalkyl)-(C₃-C₆ cycloalkyl) and (C₃-C₇heterocycloalkyl)carbonyl
 5 portion in the above listed groups may be optionally substituted by one, two or
 three groups **R**²⁵ which are independently selected in the list consisting of: C₁-
 C₆ alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇ cycloalkyl, hydroxyl
 and halo; or

R¹¹ is linked to X⁴ and is phenyl or 5- or 6-membered monocyclic
 10 heteroaryl, wherein such phenyl or 5- or 6-membered monocyclic heteroaryl is
 substituted by a group selected in the list consisting of: C₁-C₆ alkyl which is
 substituted by a group -CN; C₃-C₆ cycloalkyl which is substituted by a group
 selected from: -CN, -OR^C, -SR^C or halo; -N(R^C)(C₂-C₆alkylene)-NR^AR^B,
 -N(R^C)(C₃-C₇cycloalkylene)-NR^AR^B, -(C₁-C₆alkylene)-NR^AR^B, -(C₃-
 15 C₇cycloalkylene)-NR^AR^B, -O-(C₃-C₇cycloalkylene)-NR^AR^B,
 -S-(C₂-C₆alkylene)-NR^AR^B, -S-(C₃-C₇cycloalkylene)-NR^AR^B, -N(R^C)C(O)-
 (C₁-C₆alkylene)-NR^AR^B, -N(R^C)C(O)-(C₃-C₇cycloalkylene)-NR^AR^B,
 -C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B, -C(O)N(R^C)-(C₃-C₇cycloalkylene)-
 NR^AR^B, -C(O)N(R^C)-(C₂-C₆alkylene)-OR^D, -C(O)N(R^C)-(C₃-
 20 C₇cycloalkylene)-OR^D, -N(R^C)C(O)N(R^C)-(C₂-C₆alkylene)-NR^AR^B,
 -N(R^C)C(O)N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -O-(C₃-C₇cycloalkylene)-
 OR^D, -S-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₁-C₆alkylene)-NR^AR^B,
 -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-C₆alkylene)-
 NR^AR^B, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-NR^AR^B, -S(O)₂N(R^C)-(C₂-
 25 C₆alkylene)-OR^D, -S(O)₂N(R^C)-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂-(C₂-
 C₆alkylene)-OR^D, -N(R^C)S(O)₂-(C₃-C₇cycloalkylene)-OR^D, -N(R^C)S(O)₂R^D,
 -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl),
 (C₅-C₇heterocycloalkyl)(C₃-C₆cycloalkyl) and (C₃-

C₇heterocycloalkyl)carbonyl, wherein any of such C₁-C₆alkyl, C₃-C₆cycloalkyl, -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-C₇cycloalkylene)-, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl), (C₅-C₇heterocycloalkyl)-(C₃-C₆ cycloalkyl) and (C₃-C₇heterocycloalkyl)carbonyl
 5 portion in the above listed groups may be optionally substituted by one, two or three groups **R**²⁵ which are independently selected in the group consisting of: C₁-C₆alkyl, (C₁-C₃) haloalkyl, (C₁-C₄)hydroxyalkyl, C₃-C₇cycloalkyl, hydroxyl and halo;

R^A and **R**^B are at each occurrence independently hydrogen, C₁-C₆ alkyl
 10 or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^D, -CN or halo; alternatively, **R**^A and **R**^B, may form together with the nitrogen atom to which they are attached a 5-11- membered saturated heterocyclic monocyclic or bicyclic ring system which is optionally substituted by one or more group
 15 -OR^D, -CN, halo, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^D, -CN or halo; and which 5-11- membered saturated heterocyclic monocyclic or bicyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally
 20 substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl, wherein any of such C₁-C₆ alkyl or C₃-C₆ cycloalkyl may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, -OR^D, -CN, or halo; and/or **R**^A and **R**^B may be linked to one carbon atom of the -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)- or -(C₃-C₇cycloalkylene)- portion of the group linked to the nitrogen to which they are
 25 connected to form a saturated cycle of up to 6 ring atoms;

R^C is at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl and C₃-C₆ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, -OR^D, -CN or halo;


R^D is at each occurrence independently hydrogen, $-CH_3$ or $-C_2H_5$;

R^{12} and R^{13} are independently hydrogen, C_1 - C_6 alkyl, or halogen;

A is a divalent cycloalkylene radical having 5, 6 or 7 ring atoms; said cycloalkylene ring being attached to W and Y , and fused to a phenyl ring or to
 5 a monocyclic heteroaryl ring having 5 or 6 ring atoms, such phenyl or heteroaryl ring being optionally substituted by one or two groups R^{24} ;

R^{24} is at each occurrence independently selected from the group consisting of: C_1 - C_6 alkyl, halogen and cyano;

R^2 is a radical of formula (IIIa), (IIIb), (IIIc), or (IIId):



10


wherein

R^{14} is selected in the group consisting of: $-F$, $-CH_3$, $-C_2H_5$, $-CH_2OH$, $-CH_2OMe$, $-CF_2CF_3$, $-CH_2SCH_3$, $-SCH_3$ and $-SC_2H_5$;

R^{15} and R^{16} are independently $-CH_3$ or $-C_2H_5$;

15 R^{17} is selected from the group consisting of: lone electron pair, hydrogen, $-CF_3$, $-NR^E R^F$, $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, aryl or heteroaryl wherein any of such $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, aryl or heteroaryl may be optionally substituted by a group C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, or halo; or

20 R^{17} is a group of general formula (IV)



wherein

R^{20} is selected in the group consisting of: -F, -CH₃, -C₂H₅, -CH₂OH, -CH₂OMe, -CF₂CF₃, -CH₂SCH₃, -SCH₃ and -SC₂H₅;

5 R^{21} is -CH₃ or -C₂H₅;

or

R^{20} and R^{21} as defined above may form together with the carbon atom to which they are attached a saturated 3-7- membered monocyclic ring;

R^{E} and R^{F} are each independently C₁-C₆ alkyl, optionally substituted by
 10 a group C₁-C₃ alkyl, -OR^G, -CN or halo; alternatively, R^{E} and R^{F} may form together with the nitrogen atom to which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more groups -OR^G, -CN, halo, C₁-C₆ alkyl or C₃-C₇ cycloalkyl, such C₁-C₆ alkyl and C₃-C₇ cycloalkyl being optionally substituted
 15 by a group C₁-C₃ alkyl, C₃-C₇cycloalkyl, -OR^G, -CN or halo; and which 5-11- membered saturated monocyclic or bicyclic heterocyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C₁-C₆ alkyl or C₃-C₆ cycloalkyl, wherein any of such C₁-C₆ alkyl or C₃-C₆ cycloalkyl may be optionally substituted by a group C₁-
 20 C₆ alkyl or C₃-C₇ cycloalkyl;

R^{G} is hydrogen, -CH₃ or -C₂H₅;

R^{18} is selected in the group consisting of: lone electron pair, hydrogen, aryl, heteroaryl, -(C₁-C₆alkyl), -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl) and (C₅-C₇heterocycloalkyl)-(C₃-C₆
 25 cycloalkyl), wherein any of such aryl, heteroaryl, -(C₁-C₆alkyl), -(C₃-C₇cycloalkyl), -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆alkyl)

and (C₅-C₇heterocycloalkyl)-(C₃-C₆ cycloalkyl) may be optionally substituted by a group -CN, -OH, halo, -COOR^M, C₁-C₆alkyl, C₃-C₆cycloalkyl, -O-(C₁-C₆alkyl), -O-(C₃-C₆cycloalkyl), -S-(C₁-C₆alkyl), -S-(C₃-C₆cycloalkyl), -NR^HR^J, -N(R^L)(C₂-C₆alkylene)-NR^HR^J, -N(R^L)(C₃-C₇cycloalkylene)-NR^HR^J,
5 - (C₁-C₆alkylene)-NR^HR^J, - (C₃-C₇cycloalkylene)-NR^HR^J, -O-(C₂-C₆alkylene)-NR^HR^J, -O-(C₃-C₇cycloalkylene)-NR^HR^J, -S-(C₂-C₆alkylene)-NR^HR^J, -S-(C₃-C₇cycloalkylene)-NR^HR^J, -N(R^L)C(O)-(C₁-C₆alkylene)-NR^HR^J, -N(R^L)C(O)-(C₃-C₇cycloalkylene)-NR^HR^J, -C(O)N(R^L)-(C₂-C₆alkylene)-NR^HR^J, -C(O)N(R^L)-(C₃-C₇cycloalkylene)-NR^HR^J, -C(O)N(R^L)-(C₂-C₆alkylene)-OR^M,
10 -C(O)N(R^L)-(C₃-C₇cycloalkylene)-OR^M, -N(R^L)C(O)N(R^HR^J), -C(O)N(R^HR^J), -N(R^L)C(O)N(R^L)-(C₂-C₆alkylene)-NR^HR^J, -N(R^L)C(O)N(R^L)-(C₃-C₇cycloalkylene)-NR^HR^J, -O-(C₂-C₆alkylene)-OR^M, -O-(C₃-C₇cycloalkylene)-OR^M, -S-(C₂-C₆alkylene)-OR^M, -S-(C₃-C₇cycloalkylene)-OR^M, -N(R^L)S(O)₂-(C₁-C₆alkylene)-NR^HR^J, -N(R^L)S(O)₂-(C₃-C₇cycloalkylene)-NR^HR^J,
15 -S(O)₂N(R^L)-(C₂-C₆alkylene)-NR^HR^J, -S(O)₂N(R^L)-(C₃-C₇cycloalkylene)-NR^HR^J, -S(O)₂N(R^L)-(C₂-C₆alkylene)-OR^M, -S(O)₂N(R^L)-(C₃-C₇cycloalkylene)-OR^M, -N(R^L)S(O)₂-(C₂-C₆alkylene)-OR^M, -N(R^L)S(O)₂-(C₃-C₇cycloalkylene)-OR^M, -S(O)₂N(R^HR^J), -N(R^L)S(O)₂R^L, -N(R^L)C(O)R^L, OR^L, SR^L, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl) and (C₅-C₇ heterocycloalkyl)-(C₃-C₆ cycloalkyl), wherein any of such C₁-C₆alkyl, C₃-C₆cycloalkyl, -(C₁-C₆alkylene)-, -(C₂-C₆alkylene)-, -(C₃-C₇cycloalkylene)-, -(C₃-C₇heterocycloalkyl), (C₅-C₇heterocycloalkyl)-(C₁-C₆ alkyl) and (C₅-C₇ heterocycloalkyl)-(C₃-C₆ cycloalkyl) portion in the above listed groups may be optionally substituted by a group C₁-C₆ alkyl, C₃-C₇ cycloalkyl, -OR^L or halo;
25 **R^H** and **R^J**, are at each occurrence independently hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl, such C₁-C₆ alkyl or C₃-C₆ cycloalkyl being optionally substituted by a group C₁-C₃ alkyl, -OR^M, CN or halo;

alternatively, R^H and R^J may also together with the nitrogen atom to

which they are attached a 5-11- membered saturated monocyclic or bicyclic heterocyclic ring system which is optionally substituted by one or more groups $-OR^M$, $-CN$, halo, C_1-C_6 alkyl or C_3-C_7 cycloalkyl, such C_1-C_6 alkyl and C_3-C_7 cycloalkyl being optionally substituted by a group C_1-C_3 alkyl, C_3-
 5 C_7 cycloalkyl, $-OR^M$, CN or halo; and which 5-11- membered saturated monocyclic or bicyclic heterocyclic ring optionally contains a further heteroatom which is oxygen or nitrogen, said nitrogen atom optionally substituted by C_1-C_6 alkyl or C_3-C_6 cycloalkyl, wherein any of such C_1-C_6 alkyl or C_3-C_6 cycloalkyl may be optionally substituted by a group C_1-C_6
 10 alkyl, C_3-C_7 cycloalkyl, $-OR^M$, CN , or halo;

and/or R^H and R^J may be linked to one carbon atom of the $-(C_1-C_6\text{alkylene})-$, $-(C_2-C_6\text{alkylene})-$ or $-(C_3-C_7\text{cycloalkylene})-$ portion of the group linked to the nitrogen to which they are connected to form a saturated cycle of up to 6 ring atoms;


15 R^L is at each occurrence independently hydrogen, C_1-C_6 alkyl or C_3-C_6 cycloalkyl, such C_1-C_6 alkyl or C_3-C_6 cycloalkyl being optionally substituted by a group C_1-C_3 alkyl, $-OR^M$, $-CN$ or halo;

R^M is at each occurrence independently hydrogen, C_1-C_6 alkyl or C_3-C_6 cycloalkyl, such C_1-C_6 alkyl or C_3-C_6 cycloalkyl being optionally substituted
 20 by a group hydroxyl, $-CN$ or halo;

z^1 , z^2 , z^3 , and z^4 are independently selected in the group consisting of: C, N, S, O, a group $-CH-$, and a group $-NH-$, in such a combination that the resulting ring formed is an aromatic system;

R^{19} is selected from the group consisting of: hydrogen, $-CF_3$, $-NR^E R^F$,
 25 $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, aryl or heteroaryl wherein any of such $-(C_3-C_7\text{cycloalkyl})$, $-(C_3-C_7\text{heterocycloalkyl})$, aryl or heteroaryl may be optionally substituted by a group C_1-C_6 alkyl, C_3-C_7 cycloalkyl, or halo; or

R^{19} is a group of general formula (V)



wherein R^{20} , R^{21} , R^E and R^F are as above defined;

T is $-N=$ or $-CR^{23}=$;


R^{23} is H, halo, $-CH_3$, or $-CN$;

5 R^{22} is H, halo, $-CH_3$, or $-CN$;

q is 0, 1, 2 or 3;

with the proviso that when Y is a group $-O(CR^3R^4)_n-$, n is 1 and R^{10} is $-NR^A R^B$, $-N(R^C)C(O)-(C_1-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^A R^B)$, $-N(R^C)C(O)N(R^C)-(C_2-C_6\text{alkylene})-NR^A R^B$, $-N(R^C)C(O)N(R^C)-(C_3-C_7\text{cycloalkylene})-NR^A R^B$, or $-N(R^C)C(O)R^C$, then X_1 is nitrogen.


2. A compound of formula (I) according to claim 1 which is a compound of formula (Ia) wherein carbon stereogenic center on the cycloalkylene portion of ring A which is linked to group W and identified with number (1) herebelow, possess the absolute configuration herebelow represented:



Or a pharmaceutically acceptable salt thereof.

20 3. A compound of formula (I), according to claim 1 or 2, which is a compound of formula (Ib) wherein carbon stereogenic centers on the cycloalkylene portion of ring A which are linked to group W and Y and identified, respectively, with numbers (1) and (2) herebelow, possess the absolute configuration herebelow represented:


511



(Ib)


Or a pharmaceutically acceptable salt thereof.

4. A compound of formula (I) according to anyone of claims 1 to 3
 5 wherein A is group selected in the group consisting of:



Or a pharmaceutically acceptable salt thereof.

5. A compound of formula (I) according to anyone of claims 1 to 4
 10 wherein W is NH or O, or a pharmaceutically acceptable salt thereof.
6. A compound of formula (I) according to anyone of claims 1 to 5
 wherein Y is $-S(O)_p-$, a group $-O(CR^3R^4)_n-$, $-(CR^5R^6)_n-$, or a group $-NR^7-$; p is
 zero and n is 0,1 or 2, or a pharmaceutically acceptable salt thereof.
7. A compound of formula (I) according to anyone of claims 1 to 6,
 15 wherein R^1 is a group of formula (IIc):




(IIc)


or a pharmaceutically acceptable salt thereof.

8. A compound of formula (I) according to anyone of claims 1 to 7,

wherein W is NH, Y is a group $-O(CR^3R^4)_n-$ and n is 0, A is group:




Wherein R^1 is a group of formula (IIca) as above defined which is connected
5 to the group Y through the carbon adjacent to X^2



And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a nitrogen atom
and X^2 is a group $-CH-$, and R^{13} is hydrogen;


Wherein R^{11} is a group:

10



Wherein R^{25} is optionally present and represents one, two or three substituents
independently selected in the list consisting of: C_1 - C_6 alkyl, $(C_1$ - $C_3)$ haloalkyl,
 $(C_1$ - $C_4)$ hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and wherein the
15 asterisk represents the point of attachment for group R^{11} to the rest of the
molecule via X^4 ;


Wherein R^2 is a radical of formula (IIIb):



(IIIb)

;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)




5


And Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$.

9. A compound of formula (I) according to anyone of claims 1 to 7, wherein W is NH, Y is a group $-O(CR^3R^4)_n-$ and n is 0, A is group:

10




Wherein R^1 is a group of formula (IIca) as above defined which is connected to the group Y through the carbon adjacent to X^2



15 And wherein X^4 is a carbon atom, X^5 is a nitrogen atom, X^3 is a nitrogen atom and X^2 is a group $-CH-$, and R^{13} is hydrogen;


Wherein R^{11} is a group:



Wherein R^{25} is optionally present and represents one, two or three substituents
 5 independently selected in the list consisting of: C_1 - C_6 alkyl, (C_1 - C_3) haloalkyl, (C_1 - C_4)hydroxyalkyl, C_3 - C_7 cycloalkyl, hydroxyl and halo; and wherein the asterisk represents the point of attachment for group R^{11} to the rest of the molecule via X^4 ;

Wherein R^2 is a radical of formula (IIIb):


10



(IIIb)

;

Wherein $z^1 = -CH-$, $z^2 = C$, z^3 and z^4 are N and R^{17} is a group of general formula (IV)



(IV)

15

And Wherein R^{20} is $-CH_3$ or $-CH_2OH$, and R^{21} is $-CH_3$.

10. A pharmaceutical composition comprising a compound as claimed in any of the preceding claims 1 to 9, together with one or more pharmaceutically acceptable carriers.

11. A compound as claimed in any of claims 1 to 9 for use in the treatment of diseases or conditions which benefit from inhibition of p38 MAP kinase activity.

12. Compounds according to anyone of claims 1 to 9 for use in the
5 treatment of diseases or conditions which benefit from inhibition of p38 MAP kinase activity according to claim 11 wherein the diseases or conditions is chronic eosinophilic pneumonia, asthma, COPD, adult respiratory distress syndrome (ARDS), exacerbation of airways hyper-reactivity consequent to other drug therapy or airways disease that is associated with pulmonary
10 hypertension.

13. A method of treating diseases or conditions in a human subject which benefit from inhibition of p38 MAP kinase activity comprising administering to such subject in need thereof a compound as claimed in any of claims 1 to 9.

14. A method according to claim 13, wherein the disease or condition is
15 chronic eosinophilic pneumonia, asthma, COPD, adult respiratory distress syndrome (ARDS), exacerbation of airways hyper-reactivity consequent to other drug therapy or airways disease that is associated with pulmonary hypertension.

15. The use of a compound as claimed in any of claims 1 to 9 in the
20 manufacture of a medicament for the treatment of diseases or conditions which benefit from inhibition of p38 MAP kinase activity.

16. The use according to claim 15 wherein the disease or condition is chronic eosinophilic pneumonia, asthma, COPD, adult respiratory distress syndrome (ARDS), exacerbation of airways hyper-reactivity consequent to
25 other drug therapy or airways disease that is associated with pulmonary hypertension.