

**COMPOUNDS HAVING MUSCARINIC RECEPTOR ANTAGONIST AND BETA2
ADRENERGIC RECEPTOR AGONIST ACTIVITY**

FIELD OF THE INVENTION

The present invention relates to compounds of general formula I, acting both as muscarinic receptor antagonists and beta2 adrenergic receptor agonists, the processes for the preparation thereof, compositions comprising them, the therapeutic uses and combinations with other pharmaceutical active ingredients.

BACKGROUND OF THE INVENTION

Pulmonary disorders, such as asthma and chronic obstructive pulmonary disease (COPD), are commonly treated with bronchodilators. A well-known class of bronchodilators consists of beta-2 adrenergic receptor agonists, such as salbutamol, fenoterol, formoterol and salmeterol. These compounds are generally administered by inhalation.

Another well-known class of bronchodilators consists of muscarinic receptor antagonists (anticholinergic compounds), such as ipratropium and tiotropium. These compounds are also typically administered by inhalation.

Inhaled formulations of both beta-2 agonists and muscarinic receptor antagonists are valuable agents in the treatment of asthma and COPD, with both classes of agents providing symptomatic relief due to their ability to relax constricted airways. Observations that the bronchodilator effects of the two classes of agents were additive, prompted studies with combinations of the two agents. In 1975, it was shown that that beneficial effects could be achieved by combining two ingredients such as fenoterol and ipratropium bromide in a single aerosol. This prompted the development of fixed dose combinations of ipratropium bromide firstly with fenoterol (Berodual, introduced in 1980), and then with salbutamol (Combivent, introduced in 1994).

More recently the availability of both long-acting muscarinic antagonists and long-acting beta-2 agonists prompted to the development of combinations of these agents.

For example, WO 00/69468 discloses medicament compositions containing a muscarinic receptor antagonist, such as tiotropium bromide, and beta-2 adrenergic receptor agonists, such as formoterol fumarate or salmeterol, and WO 2005/115467 discloses a combination which comprises a beta-2 agonist and an antagonist of M3 muscarinic receptors which is a
5 salt of 3(R)-(2-hydroxy-2,2-dithien-2-ylacetoxy)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane.

An alternative approach to the development of fixed dose combinations is the identification of molecules that combine both activities of muscarinic antagonism and beta-2 agonism. In fact compounds possessing both beta-2 adrenergic receptor agonist and
10 muscarinic receptor antagonist activity are highly desirable since such bifunctional compounds would provide bronchodilation through two independent mechanisms of action while having a single molecule pharmacokinetics.

Such kind of compounds was described in some patent applications, such as WO 2004/074246, WO 2004/074812, WO 2005/051946, WO 2006/023457,
15 WO 2006/023460, WO 2010/123766, WO 2011/048409 and co-pending patent applications WO 2012/168349, WO 2012/168359, WO2014/086924, WO 2014/086927.


It has now been found that some particular aryl or heteroaryl hydroxyacetic ester derivatives, besides possessing both beta-2 adrenergic receptor agonist and muscarinic receptor antagonist activity, possess elevated affinity for the M3 muscarinic receptors and
20 long lasting bronchodilating activity.

SUMMARY OF THE INVENTION

The present invention relates to compounds of general formula **I**, acting both as muscarinic receptor antagonists and beta2 adrenergic receptor agonists, the processes for the preparation thereof, compositions comprising them, the therapeutic uses and
25 combinations with other pharmaceutical active ingredients among which are, for instance, those currently used in the treatment of respiratory disorders, e.g. corticosteroids, P38 MAP kinase inhibitors, IKK2, HNE inhibitors, PDE4 inhibitors, leukotriene modulators, NSAIDs and mucus regulators.


DETAILED DESCRIPTION OF THE INVENTION

In particular, the invention is directed to compounds of general formula




wherein


Q is a group of formula



Y is selected from **Y2** and **Y1** which are divalent groups of formula



or








wherein


A1 and **A2** are independently absent or selected from (C₁-C₁₂)alkylene, (C₃-C₈)cycloalkylene and (C₃-C₈)heterocycloalkylene optionally substituted by one or more substituents selected from (C₁-C₆)alkyl, aryl(C₁-C₆)alkyl and heteroaryl(C₁-C₆)alkyl;

B is absent or is selected from (C₃-C₈)cycloalkylene, (C₃-C₈)heterocycloalkylene, arylene or heteroarylene optionally substituted by one or more groups selected from -OH, halogens, -CN, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)haloalkyl, (C₁-C₆)haloalkoxy and aryl(C₁-C₆)alkyl;


- 5 **C** is absent or is selected from -O-, -C(O)-, -OC(O)-, -(O)CO-, -S-, -S(O)-, -S(O)₂- and -N(R₇)-, or is one of the following groups **C1-C23**




5




C14




C15




C16




C17




C18




C19




C20



C21



C22



C23

wherein **R**₇ is in each occurrence independently H or selected from linear or branched
 (C₁-C₈)alkyl, aryl(C₁-C₆)alkyl, arylsulfanyl, arylsulfinyl, arylsulfonyl,
 5 (C₃-C₈)cycloalkyl, (C₃-C₈)heterocycloalkyl, aryl, and heteroaryl;

D is absent or is selected from (C₁-C₁₂)alkylene, (C₂-C₁₂)alkenylene, (C₂-C₆)alkynylene,
 arylene, heteroarylene, (C₃-C₈)cycloalkylene, (C₃-C₈)heterocycloalkylene; said
 arylene, heteroarylene, (C₃-C₈)cycloalkylene and (C₃-C₈)heterocycloalkylene being

optionally substituted by one or more groups selected from $-\text{OH}$, halogen, $-\text{CN}$, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_1\text{-C}_6)\text{haloalkyl}$, $(\text{C}_1\text{-C}_6)\text{haloalkoxy}$ and $\text{aryl}(\text{C}_1\text{-C}_6)\text{alkyl}$;

n is at each occurrence independently 0 or an integer from 1 to 3;

5 **m** is at each occurrence independently an integer from 1 to 3;

E is absent or is selected from $-\text{O}-$, $-\text{NR}_7-$, $-\text{NR}_7\text{-C(O)-}$, $-\text{C(O)-NR}_7-$, $-\text{OC(O)-}$, $-\text{C(O)-}(\text{CH}_2)_n\text{-O-}$; $-\text{NR}_7\text{-C(O)-}(\text{CH}_2)_n\text{-O-}$, $-\text{NR}_7\text{-C(O)-NR}_7-$ and $-\text{S-}$;

10 **G** is arylene or heteroarylene, optionally substituted by one or more substituents selected from halogen atoms, $-\text{OH}$, oxo ($=\text{O}$), $-\text{SH}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{CON}(\text{R}_6)_2$, $-\text{NH}_2$, $-\text{NHCOR}_6$, $-\text{CO}_2\text{R}_6$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfanyl}$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfinyl}$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfonyl}$, $(\text{C}_1\text{-C}_{10})\text{alkyl}$, aryl , haloaryl , heteroaryl and $(\text{C}_1\text{-C}_{10})\text{alkoxy}$;


15 **R₁** is selected from $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_8)\text{heterocycloalkyl}$, aryl , heteroaryl , $\text{aryl}(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkyl}$ and $(\text{C}_3\text{-C}_8)\text{cycloalkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$, optionally substituted by one or more group selected independently from halogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, and $(\text{C}_1\text{-C}_{10})\text{alkoxy}$;

s is 0 or an integer from 1 to 3;


R₂ is a nitrogen containing group which may be selected from:

- a group (a) which is $-\text{NR}_3\text{R}_4$ wherein R_3 and R_4 are independently hydrogen or $(\text{C}_1\text{-C}_4)\text{alkyl}$; and

20 - a group (b) of formula J1, J2, J3, J4 or J5



R₅ is a group of formula **K**

**K**

wherein **p** is 0 or an integer from 1 to 4; **q** is 0 or an integer from 1 to 4;

P is absent or is selected from the divalent group consisting of O, S, SO, SO₂, CO, NR₆, CH=CH, N(R₆)SO₂, N(R₆)COO, N(R₆)C(O), SO₂N(R₆), OC(O)N(R₆) and C(O)N(R₆);

W is selected from H, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, aryl and heteroaryl, optionally substituted by one or more substituents selected independently from halogen atoms, -OH, oxo (=O), -SH, -NO₂, -CN, -CON(R₆)₂, -NH₂, -NHCOR₆, -CO₂R₆, (C₁-C₁₀)alkylsulfanyl, (C₁-C₁₀)alkylsulfinyl, (C₁-C₁₀)alkylsulfonyl, (C₁-C₁₀)alkyl and (C₁-C₁₀)alkoxy;

R₆ is at each occurrence independently H or selected from (C₁-C₁₀)alkyl, (C₁-C₆)haloalkyl, (C₂-C₆)alkynyl, (C₂-C₆)alkenyl, (C₃-C₈)cycloalkyl, heteroaryl and aryl optionally substituted by one or more substituents selected from halogen atoms, -OH, oxo (=O), -SH, -NO₂, -CN, -CONH₂, -COOH, (C₁-C₁₀)alkoxycarbonyl, (C₁-C₁₀)alkylsulfanyl, (C₁-C₁₀)alkylsulfinyl, (C₁-C₁₀)alkylsulfonyl, (C₁-C₁₀)alkyl and (C₁-C₁₀)alkoxy;

and pharmaceutically acceptable salts or solvates thereof.

The term “pharmaceutically acceptable salts”, as used herein, refers to compounds according to the invention obtained by converting any of the free acid or basic group, if present, into the corresponding addition salt with any base or acid conventionally intended as being pharmaceutically acceptable.

Suitable examples of said salts may thus include mineral or organic acid addition salts of basic residues such as amino groups, as well as mineral or organic basic addition salts of acid residues such as carboxylic groups.

Cations of inorganic bases which can be suitably used to prepare salts within the invention comprise ions of alkali or alkaline earth metals such as potassium, sodium,

calcium or magnesium.

Salts obtained by reacting the main compound, functioning as a base, with an inorganic or organic acid to form a salt comprise, for example, salts of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methane sulfonic acid, camphor sulfonic acid, acetic acid, oxalic acid, maleic acid, fumaric acid, succinic acid and citric acid.

Many organic compounds can form complexes with solvents in which they are reacted or from which they are precipitated or crystallized, known as "solvates". Pharmaceutically acceptable solvates of compound of the invention are within the scope of the invention.

Included within the scope of the present invention are also polymorphs and crystalline forms of compounds of formula (I), or of pharmaceutically acceptable salts, or solvates thereof.

The term "halogen", "halogens" or "halogen atoms" as used herein includes fluorine, chlorine, bromine, and iodine, preferably chlorine or fluorine, referring independently to one or more of these atoms.

The expression "(C₁-C_x)alkyl" refers to straight or branched chain alkyl groups wherein the number of carbon atoms is 1 to x, preferably 1 to 6 thus referring to (C₁-C₆)alkyl. Examples are methyl, ethyl, n-propyl, isopropyl, t-butyl, pentyl, hexyl, octyl, nonyl, decyl, undecyl, dodecyl and the like.

The expression "(C₁-C_x)alkylene" refers to divalent groups wherein the number of carbon atoms is from 1 to x, preferably from 1 to 6 thus referring to (C₁-C₆)alkylene, such as methylene, ethylene, n-propylene, isopropylene, n-butylene, t-butylene, pentylene, hexylene, octylene, nonylene, decylene, undecylene, dodecylene and the like. The above divalent groups can also be referred to as methanediyl, ethanediyl, n-propanediyl, propan1,2diyl and the like.

The expression "(C₁-C₆)haloalkyl" refers to the above "(C₁-C₆)alkyl" group wherein one or more hydrogen atoms are replaced by one or more halogen atoms, which can be the same or different from each other. Examples include halogenated, poly-halogenated and

fully halogenated alkyl groups wherein one or more of the hydrogen atoms are replaced by halogen atoms, e.g. trifluoromethyl group.

The expression “hydroxy(C₁-C₆)alkyl” likewise refers to -alkyl-OH groups.

The expressions “(C₁-C₁₀)alkylsulfanyl”, “(C₁-C₁₀)alkylsulfinyl” or
5 “(C₁-C₁₀)alkylsulfonyl” refer, respectively, to alkyl-S-, alkyl-SO- or alkyl-SO₂- groups.

The expression “(C₂-C_x)alkenyl” refers to straight or branched carbon chains with one or more double bonds, wherein the number of carbon atoms is from 1 to x. Examples comprise ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl and the like.

10 The expression “(C₂-C_x)alkenylene” refers to divalent groups, such as ethenylene, propenylene, butenylene, pentenylene, hexenylene, heptenylene, octenylene, nonenylene, decenylene, undecenylene, dodecenylene and the like.

The expression “(C₂-C_x)alkynyl” refers to straight or branched carbon chains with one or more triple bonds, wherein the number of carbon atoms is from 1 to x. Examples
15 comprise ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like.

The expression “(C₂-C₆)alkynylene” refers to divalent groups, such as ethynylene, propynylene, butynylene, pentynylene, hexynylene and the like; otherwise commonly referred to as ethynediyl, propynediyl, butyndiyl and the like.

The expression “(C₁-C_x)alkoxy” refers to alkyl-oxy (i.e. alkoxy) groups, being the
20 alkyl portion as above defined, wherein the number of carbon atoms is from 1 to x. Examples comprise methoxy (i.e. CH₃O-), ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentoxy, hexoxy and the like.

The expression “(C₁-C₆)haloalkoxy” refers to the above “(C₁-C₆)alkoxy” group wherein one or more hydrogen atoms are replaced by one or more halogen atoms, which
25 can be the same or different from each other.

Examples include halogenated, poly-halogenated and fully halogenated alkoxy groups wherein one or more of the hydrogen atoms are replaced by halogen atoms, e.g. trifluoromethoxy group.

The expression “(C₁-C₁₀)alkoxycarbonyl” refers to (C₁-C₁₀)alkoxyC(O)- groups. Examples include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isopropoxycarbonyl, and the like.

The expression “(C₃-C₈)cycloalkyl” refers to mono or bi-cycloaliphatic hydrocarbon groups with 3 to 8 carbon atoms. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]hept-2-yl and the like.

The expression “(C₃-C₈)heterocycloalkyl” refers to saturated or partially saturated monocyclic (C₃-C₈)cycloalkyl groups, in which at least one ring carbon atom is replaced by a heteroatom or heteroaromatic group (e.g. N, NH, S or O). Examples include quinuclidinyl, pyrrolidinyl, piperidinyl, azabicyclo[3.2.1]octan-3-yl and azoniabicyclo[2.2.2]octanyl, [1.2.3.6]tetrahydropyridin-1yl and the like.

The expressions “(C₃-C₈)cycloalkylene” and “(C₃-C₈)heterocycloalkylene” herewith refer to divalent groups. The term cycloalkylene refers to saturated cycloalkane-diyl and partially saturated monocyclic groups such as cycloalkene-diyl. Examples are divalent groups, such as, respectively, cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, cycloheptylene, bicyclo[2.2.1]hept-2-ylene and quinuclidinylene, pyrrolidinylene, piperidinylene, azabicyclo[3.2.1]octan-3-ylene, azoniabicyclo[2.2.2]octanylene, [1.2.3.6]tetrahydropyridin-[1.4]diyl and the like. The above divalent groups can also be referred to as cyclopropanediyl, cyclobutanediyl, cyclopentanediy, cyclohexanediy, cycloheptanediy, bicyclo[2.2.1]heptaendiy and quinuclidinediy, pyrrolidinediy, piperidinediy, azabicyclo[3.2.1]octandiy, azoniabicyclo[2.2.2]octandiy, [1.2.3.6]tetrahydropyridine-[1.4]diyl and the like.

The expression “aryl” refers to mono, bi- or tricyclic ring systems having 5 to 20, preferably 5 to 15, more preferably from 5 to 8 ring atoms, and wherein at least one ring is aromatic.

The expression “heteroaryl” refers to mono, bi- or tri-cyclic systems with 5 to 20 ring atoms, preferably 5 to 15, in which at least one ring is aromatic and in which at least one carbon ring atom is a heteroatom or heteroaromatic group (e.g. N, NH, S or O).

Examples of suitable aryl or heteroaryl monocyclic systems include, for instance, thiophene, benzene, pyrrole, pyrazole, imidazole, isoxazole, oxazole, isothiazole, thiazole, pyridine, imidazolidine, furan radicals and the like.

Examples of suitable aryl or heteroaryl bicyclic systems include naphthalene, biphenylene, purine, pteridine, benzotriazole, quinoline, isoquinoline, indole, isoindole, benzothiophene, dihydrobenzo dioxin, dihydro-indene, dihydrobenzo dioxepin, benzo oxazine radicals and the like.

Examples of suitable aryl or heteroaryl tricyclic systems include fluorene radicals as well as benzocondensed derivatives of the aforementioned heteroaryl bicyclic systems.

The expressions “arylene” and “heteroarylene” refer to divalent groups, such a phenylene, biphenylene and thienylene. Such groups are also commonly named as “arenediyl” or “heteroarenediyl” groups. e.g. ortho-phenylene is also named benzene-1,2-diyl, para-phenylene is also named benzene-1,4-diyl, meta-phenylene is also named benzene-1,3-diyl.

The expressions “aryl(C₁-C₆)alkyl”, “heteroaryl(C₁-C₆)alkyl” and “(C₃-C₈)cycloalkyl(C₁-C₆)alkyl” refer to a “(C₁-C₆)alkyl” respectively substituted by one or more aryl, heteroaryl or (C₃-C₈)cycloalkyl groups, as defined above.

Examples of aryl(C₁-C₆)alkyl include triphenylmethyl.

The expressions “arylsulfanyl”, “arylsulfinyl” or “arylsulfonyl” refer, respectively, to aryl-S-, aryl-SO- or aryl-SO₂- groups. Preferred aryl groups are phenyl-S-, phenyl-SO- or phenyl-SO₂-.

The expression “haloaryl” refers to the above “aryl” group wherein one or more hydrogen atoms are replaced by one or more halogen atoms, which can be the same or different from each other.


As used herein an oxo moiety is represented by (O) as an alternative to other common representations, e.g. (=O). Thus, in terms of general formula, the carbonyl group is herein preferably represented as -C(O)- as an alternative to the other common representations such as -CO-, -(CO)- or -C(=O)-. In general the bracketed group is a

lateral group, not included into the chain, and brackets are used, when deemed useful, to help disambiguating linear chemical formulas; e.g. the sulfonyl group $\text{-SO}_2\text{-}$ might be also represented as $\text{-S(O)}_2\text{-}$ to disambiguate e.g. with respect to the sulfinic group -S(O)O- .

Whenever basic amino or quaternary ammonium groups are present in the compounds of formula I, as above said, physiological acceptable anions, selected among
 5 chloride, bromide, iodide, trifluoroacetate, formate, sulfate, phosphate, methanesulfonate, nitrate, maleate, acetate, citrate, fumarate, tartrate, oxalate, succinate, benzoate, p-toluenesulfonate, pamoate and naphthalene disulfonate may be present. Likewise, in the presence of acidic groups such as -COOH groups, corresponding physiological cation salts
 10 may be present as well, for instance including alkaline or alkaline earth metal ions.

Compounds of general formula (I) contain at least two stereogenic centers. Therefore, the invention also includes any of the optical stereoisomers, diastereoisomers and mixtures thereof, in any proportion.


In particular, the carbon atom (2) linked to R_1 , -OH , G and C=O groups, and the
 15 carbon atom (1) linked to Q , -OH and the rest of the molecule, represent stereogenic centers.



Thus, compounds according to the invention, having at least two stereogenic
 20 centers, may accordingly exist as at least four diastereoisomers. Where the compounds according to the invention possess more than two stereogenic centers, they will exist as 2^n diastereoisomers (wherein n here refers to the number of stereogenic centers). It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed

within the scope of the present invention.

In a preferred embodiment, the invention is directed to compounds of formula (I)', which are compounds of formula (I) as above defined where the absolute configuration of carbon (1) is that shown herebelow:





The absolute configuration for a chiral carbon is determined via X ray and assigned on the basis of Cahn-Ingold-Prelog nomenclature based on groups' priorities.

10 In a preferred embodiment, for compounds of formula (I), absolute configuration at carbon (1) is (R).

As already mentioned, compounds of formula (I) may exist as at least four diastereoisomers (Ia), (Ib), (Ic) and (Id) herebelow represented, which are comprised within the scope of the present invention; each diastereoisomer (Ia), (Ib), (Ic), (Id) may be constituted by a mixture of corresponding epimers when a third stereogenic center is

15 present in the molecule.






A further preferred embodiment is directed to compounds of formula (Ia) or (Ib), which are respectively compounds of formula (I) as above defined wherein the absolute configuration at carbon (1) is (R) and at carbon (2) is (R); or wherein the absolute configuration at carbon (1) is (R) and at carbon (2) is (S).

It is to be understood that all preferred groups or embodiments described herebelow and hereabove for compounds of formula (I) may be combined among each other and apply to compounds of formula (Ia), (Ib), (Ic), (Id) and (I)' as well *mutatis mutandis*.

A first preferred embodiment is directed to group of compounds of general formula **I** wherein **R₂** is a group of formula **J3**:



J3

R₅ is a group of formula **K**, wherein **p** is 0 or 1, **P** is absent or is CO, **q** is absent or is 1 and **W** is H or is selected from (C₁-C₆)alkyl and aryl, and all the other variables are as defined above.

In a more preferred embodiment, **R₂** is a group of formula **J3**, **R₅** is methyl or benzyl, and all the other variables are as defined above.

In another preferred embodiment, **G** is arylene and **R₁** is aryl, optionally substituted by one or more group independently selected from halogen, (C₁-C₈)alkyl and

(C₁-C₁₀)alkoxy.

In a further preferred embodiment **A1** and **A2** are independently absent or selected from methylene, ethylene, propylene, butylene, pentylene, hexylene, heptylene, octylene and nonylene, **G** is phenylene, and **R₁** is phenyl, optionally substituted by one or more
5 group selected independently from halogen, (C₁-C₈)alkyl and (C₁-C₁₀)alkoxy, and all the other variables are as defined above.

In this first preferred embodiment when **R₁** is a phenyl group substituted by one or more groups independently selected from halogen, (C₁-C₈)alkyl and (C₁-C₁₀)alkoxy; it is preferred that **R₁** is substituted in para and/or meta position.

10 In another preferred embodiment **E** is -O- or -C(O)-(CH₂)_n-O- or -NR₇-C(O)-(CH₂)_n-O-; **G** is phenylene wherein **E** is linked to the phenyl ring **G** in meta position, and **R₁** is phenyl, optionally substituted by one or more group selected from halogen, (C₁-C₈)alkyl or (C₁-C₁₀)alkoxy groups.

Also in this other preferred embodiment, when **R₁** is phenyl substituted by one or
15 more group selected from halogen, (C₁-C₈)alkyl or (C₁-C₁₀)alkoxy groups, it is preferred that **R₁** is substituted in para and/or meta position.

Most preferred compounds in this other preferred embodiment are those compounds of formula (I) wherein **Y** is **Y2**;

A2 is absent and **A1** is independently selected from (C₁-C₁₂)alkylene which is
20 methylene, ethylene, n-propylene, butylene, pentylene, hexylene, heptylene, octylene;

B is absent or is selected from (C₃-C₈)heterocycloalkylene which is piperidinylen, arylene which is phenylene and heteroarylene which is pyridine-diyl; **B** being optionally substituted by one or more groups selected independently from -OH, halogen which is fluorine, chlorine, bromine, (C₁-C₆)alkoxy which is methoxy, ethoxy, isopropoxy,
25 (C₁-C₆)haloalkyl which is trifluoromethyl and (C₁-C₆)haloalkoxy which is trifluoromethoxy;

C is absent or is selected from -C(O)-, or is one of the following groups **C1**, **C2**, **C7**, **C8**, **C9**, **C10**, **C12**, **C13**, **C14**, **C15**, **C16**, **C17**, **C18**, **C19**, **C20**, **C21**, **C23**; wherein **R₇**

is H;

D is absent or is selected from arylene which is para-phenylene or meta-phenylene, (C₃-C₈)cycloalkylene which is cyclohexanediyl, (C₃-C₈)heterocycloalkylene which is piperidindiyl, pyrrolidindiyl, azetidindiyl;

5 **n** is at each occurrence independently 0 or an integer from 1, 2 or 3;

m is at each occurrence independently an integer from 1, 2 or 3;

E is absent or is selected from -O-, -C(O)-(CH₂)_n-O- which is -C(O)-CH₂-O-; -NR₇-C(O)-(CH₂)_n-O- which is -NH-C(O)-CH₂-O-;

G is arylene which is meta-phenylene;


10 **R₁** is selected from thiophenyl, cyclohexyl or phenyl optionally substituted in para and/or meta position by one or more groups selected independently from fluorine, methyl, ethyl;

s is 1;

R₂ is J3 wherein **R₅** is benzyl

15 and pharmaceutically acceptable salts or solvates thereof.

A second preferred group of compounds is that of general formula I wherein **R₂** is a nitrogen containing group selected from J1, J2, or J5




20 and all the other variables as defined above.

Particularly preferred compounds of formula (I) are those

wherein

Y is a divalent groups of formula



Y2

A2 is absent and **A1** is independently selected from (C₁-C₁₂)alkylene which is methylene, ethylene, n-propylene, isopropylene, butylene, pentylene, hexylene, heptylene,
5 octylene;

B is absent or is selected from (C₃-C₈)heterocycloalkylene which is piperidinylene, arylene which is phenylene, or heteroarylene which is pyridine-diyl, pyrazole-diyl; **B** being optionally substituted by one or more groups selected from -OH, halogen which is fluorine, chlorine, bromine, -CN, (C₁-C₆)alkyl which is methyl, (C₁-C₆)alkoxy which is methoxy,
10 ethoxy, isopropoxy, (C₁-C₆)haloalkyl which is trifluoromethyl and (C₁-C₆)haloalkoxy which is trifluoromethoxy;

C is absent or is selected from -O-, -C(O)-, or is one of the following groups **C1**, **C2**, **C4**, **C7**, **C8**, **C9**, **C10**, **C12**, **C13**, **C14**, **C15**, **C16**, **C17**, **C18**, **C19**, **C20**, **C21** **C22** **C23**;

wherein **R₇** is in each occurrence independently H or selected from (C₁-C₈)alkyl
15 which is methyl, ethyl and aryl(C₁-C₆)alkyl which is benzyl;

D is absent or is selected from arylene which is para-phenylene or meta-phenylene, (C₃-C₈)cycloalkylene which is cyclohexanediyl, (C₃-C₈)heterocycloalkylene which is piperidindiyl, pyrrolidindiyl, azetidindiyl;

n is at each occurrence independently 0 or an integer from 1 to 3;

20 **m** is at each occurrence independently an integer from 1 to 3;

E is absent or is selected from -O-, -NR₇- which is -NH-, -NR₇-C(O)- which is -NH-C(O)-, -C(O)-NR₇- which is -C(O)-NH-, -C(O)-(CH₂)_n-O- which is -C(O)-CH₂-O-; -NR₇-C(O)-(CH₂)_n-O- which is -NH-C(O)-CH₂-O-;

G is arylene which is meta-phenylene or para-phenylene;

25 **R₁** is selected from thiophenyl, cyclohexyl, cyclopentyl and phenyl optionally substituted by one or more group selected independently from fluorine, methyl, ethyl and methoxy;

s is 0, 1 or 2

R₂ is a nitrogen containing group which may be selected from:

- a group (a) which is -NR₃R₄ wherein R₃ and R₄ are methyl; and
- a group (b) of formula J1, J2, J3, J4 or J5 wherein

R₅ is methyl or benzyl

5 and pharmaceutically acceptable salts or solvates thereof.

The present invention is also directed to a process for the preparation of the compounds of general formula **I**.

The present invention also provides pharmaceutical compositions of compounds of formula **I** alone or in combination with or in admixture with one or more pharmaceutically acceptable carriers and/or excipients.

10

The present invention also provides the use of compounds of formula **I** for preparing a medicament.

In a further aspect, the invention provides the use of compounds of formula **I** for the prevention and/or treatment of any broncho-obstructive or inflammatory disease, preferably asthma or chronic bronchitis or chronic obstructive pulmonary disease (COPD).

15

In a further aspect, the invention provides the use of compounds of formula **I** for the manufacture of a medicament for the prevention and/or treatment of any broncho-obstructive or inflammatory disease, preferably asthma or chronic bronchitis or chronic obstructive pulmonary disease (COPD).

20 The present invention further provides a method for prevention and/or treatment of any broncho-obstructive or inflammatory disease, preferably asthma or chronic bronchitis or chronic obstructive pulmonary disease (COPD), which comprises administering to a subject in need thereof a therapeutically effective amount of a compound of general formula **I**.

25 The present invention also provides pharmaceutical compositions suitable for administration by inhalation.

Inhalable preparations include inhalable powders, propellant-containing metering aerosols or propellant-free inhalable formulations.

The invention is also directed to a device which may be a single- or multi-dose dry powder inhaler, a metered dose inhaler and a soft mist nebulizer comprising the compounds of formula I.

The invention is also directed to a kit comprising the pharmaceutical compositions of compounds of formula I alone or in combination with or in admixture with one or more pharmaceutically acceptable carriers and/or excipients and a device which may be a single- or multi-dose dry powder inhaler, a metered dose inhaler and a soft mist nebulizer comprising the said combination or admixture.

According to specific embodiments, the present invention provides the compounds reported below:

No	STRUCTURE_NAME
1	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
2	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(p-tolyl)acetate
3	(1-benzylpiperidin-4-yl)methyl 2-(4-fluorophenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
4	(1-benzylpiperidin-4-yl)methyl 2-(3-fluorophenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
5	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(m-tolyl)acetate
6	(1-benzylpiperidin-4-yl)methyl 2-(2-chlorophenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
7	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(o-tolyl)acetate
8	(1-benzylpiperidin-4-yl)methyl 2-(2-ethylphenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate

(continued)

9	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(thiophen-2-yl)acetate
10	(1-benzylpiperidin-4-yl)methyl 2-cyclohexyl-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
11	(1-benzylpiperidin-4-yl)methyl 2-(3-ethylphenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
12	(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(4-methoxyphenyl)acetate
12A	(R)-quinuclidin-3-yl 2-cyclopentyl-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate
13	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
14	(R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
15	(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
16	(R)-(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
17	(S)-2-(dimethylamino)ethyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
18	(R)-2-(dimethylamino)ethyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
19	(S)-(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
20	(R)-(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
21	(S)-1-methylpiperidin-4-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
22	(R)-1-methylpiperidin-4-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate

(continued)

23	(S)-(1-methylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
24	(R)-(1-methylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
25	(S)-(R)-1-methylpiperidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
26	(S)-1-methylazetidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
27	(R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)phenyl)-2-phenylacetate
28	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((7-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)heptyl)oxy)phenyl)-2-phenylacetate
29	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)phenyl)-2-phenylacetate
30	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((8-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)octyl)oxy)phenyl)-2-phenylacetate
31	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
32	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
32A	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)propoxy)phenyl)-2-phenylacetate
32B	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)propoxy)phenyl)-2-phenylacetate
32C	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)nicotinamido)propoxy)phenyl)-2-phenylacetate
32D	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-ethoxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32E	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethoxy)benzamido)propoxy)phenyl)-2-phenylacetate

(continued)

32F	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32G	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-isopropoxybenzamido)propoxy)phenyl)-2-phenylacetate
32H	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32I	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32J	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32K	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)picolinamido)propoxy)phenyl)-2-phenylacetate
32L	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2,3-difluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate
32M	(S)-(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethyl)benzamido)propoxy)phenyl)-2-phenylacetate
33	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
33A	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(4-(2-(((R)-2-hydroxy-2-(4-hydroxy-6-oxo-5,6-dihydronaphthalen-1-yl)ethyl)amino)ethyl)piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
34	(R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethoxy)phenyl)-2-phenylacetate
35	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
36	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
36A	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

37	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)propoxy)phenyl)-2-phenylacetate
38	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethoxy)phenyl)-2-phenylacetate
39	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)phenyl)-2-phenylacetate
40	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)phenyl)-2-phenylacetate
41	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-benzoyl)azetidin-3-yl)methoxy)phenyl)-2-phenylacetate
42	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)azetidin-3-yl)methoxy)phenyl)-2-phenylacetate
42A	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxyphenyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
42B	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-((2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxyphenyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
42C	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
42D	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
42E	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
42F	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
43	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoxy)benzyl)oxy)phenyl)-2-phenylacetate

(continued)

44	(R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate
45	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-((3-(benzyl(3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate
46	(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-phenylacetate
47	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate
48	(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate
49	(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate
50	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)carbamoyl)phenyl)-2-phenylacetate
51	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
52	(R)-quinuclidin-3-yl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate
53	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
54	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)carbonyl)amino)phenyl)-2-phenylacetate
55	(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentanamido)phenyl)-2-phenylacetate
56	(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)ureido)phenyl)-2-phenylacetate
57	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

58	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperazin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
59	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
60	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)(methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
61	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-((4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
62	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
63	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
64	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
65	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-4-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
66	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
67	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
68	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
69	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

70	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
71	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
72	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
73	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
74	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aR,6aR)-5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-1(2H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
75	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
76	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
77	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aS,6aS)-5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-1(2H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
78	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
79	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
80	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
81	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

82	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
83	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(ethyl(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
84	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)(methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
85	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)propyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
86	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperazin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
87	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(benzyl(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
88	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
89	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-((4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
90	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
91	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
92	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
93	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.3]heptan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

94	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(7-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
95	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
96	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonan-7-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
97	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
98	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.5]nonan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
99	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,9-diazaspiro[5.5]undecan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
100	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-1-oxa-4,9-diazaspiro[5.5]undecan-4-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
101	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
102	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
103	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
104	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
105	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

106	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
107	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
108	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
109	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
110	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
111	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aR,6aR)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
112	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aS,6aS)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
113	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
114	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)carbamoxy)benzyl)oxy)phenyl)-2-phenylacetate
115	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-3,9-diazaspiro[5.5]undecane-3-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
116	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.3]heptane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
117	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(7-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate

(continued)

118	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperazine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
119	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonane-7-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
120	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.5]nonane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
121	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-1-oxa-4,9-diazaspiro[5.5]undecane-4-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
122	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,9-diazaspiro[5.5]undecane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate
123	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate
124	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate
125	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate
126	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)ethyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate
127	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(3-hydroxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
128	(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(2-(((1R,3S)-3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
129	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methylbenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

130	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
131	(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
132	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
133	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
134	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2,3-difluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
135	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
136	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-bromo-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
137	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
138	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
139	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
140	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-ethoxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
141	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethoxy)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate

(continued)

142	(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate
143	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-isopropoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
144	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)picolinamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
145	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)nicotinamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
146	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)phenyl)acetamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
147	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
148	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate
149	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
150	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidin-1-yl)-2-oxoethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate
151	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-(1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidine-4-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
152	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
153	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate


(continued)


154	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
155	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
156	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-(5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)thiophene-2-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
157	(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)cyclohexane-1-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
158	1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
159	((R)-1-benzylpyrrolidin-3-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
160	((S)-1-benzylpyrrolidin-3-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
161	(1-cyclobutylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
162	(1-methylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
163	(R)-1-benzylpyrrolidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
164	(S)-1-benzylpyrrolidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
165	1-benzylazetidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate
166	1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)phenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
167	1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxyphenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate
168	1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxyphenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate

The compounds of the invention can be prepared from readily available starting materials using the following general methods and procedures or by using other information readily available to those of ordinary skill in the art. Although a particular embodiment of the present invention may be shown or described herein, those skilled in the art will
5 recognize that all embodiments or aspects of the present invention can be prepared using the methods described herein or by using other known methods, reagents and starting materials. When typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. While the optimum reaction conditions may vary
10 depending on the particular reactants or solvent used, such conditions can be readily determined by one skilled in the art by routine optimisation procedures.

Compounds of general formula **I** may be prepared according to the following synthetic **Scheme**.

Scheme





General procedure for the preparation of compounds of formula I

Compounds of formula **I** are compounds in which **Y** is a divalent group of formula **Y1** or **Y2**. Although groups **Y1** and **Y2** are different the approach to be considered for the synthesis of compounds of formula **I** in which **Y** is **Y1** or **Y2** is similar and mainly depends on the functional group present in the linker **Y**. The following synthesis described for **Y2** can be extended with minor modifications to **Y1**.

Compounds **XII** represents a compound wherein **A1** is alkylene substituted with oxo, leading to an aldehyde or ketone protected as cyclic acetal. The cyclic acetal-protecting group (PG) can be removed leading to a compound of formula **XIII**.

The synthesis of compounds of formula **I** may require the protection of potential reactive functionalities in addition to those methods already described. Examples of compatible protecting groups (PG) and their particular methods of protection and deprotection are described in "Protecting groups in organic Synthesis" by T.W. Green and P. Wutz (Wiley-Interscience publication, 1999). Compounds **I** can be prepared for example by reaction of a compound of general formula **XIII** with a compound of formula **XIV**. This reductive amination reaction can be performed following several different protocols described in the literature. For example, it can be performed in solvent such as methanol, ethanol, tetrahydrofuran (THF) or dichloromethane (DCM) using a reducing agent such as NaBH₄, NaCNBH₃ or NaB(AcO)₃H. It could be useful to preform the imine before adding the reducing agent. The reaction proceeds smoothly at room temperature (RT) over 1 to 12 hours.

The intermediate **XII** can be easily prepared by reaction of a compound **XI** with a compound of formula HO-(CH₂)_s-R₂ under the known condensation conditions for the preparation of esters. The reaction occurs smoothly in an aprotic polar solvent such as DCM, THF or DMF at room or higher temperature in the presence of a condensing agent such as EDC, DCC, HATU. Alternatively the acid **XI** can be converted into the corresponding chloride (e.g. with COCl₂ in DMC) or imidazolide (with CDI in DCM or DMF) and then treated with HO-(CH₂)_s-R₂.

Intermediate **XI** can be easily obtained by saponification of ester **VIII** using alkaline hydroxides in polar solvent, such as MeOH, EtOH, and THF mixed with water in a suitable ratio. The reaction occurs at RT, but higher temperature can speed the reaction up that can complete in a time ranging from 30 minutes to overnight.

5 Synthesis of compound of formula **VIII** can be achieved in many different ways, all depending on functional groups present in G. For example, it can be obtained by reaction of a compound **VII** with a compound **III**. In the case n is greater than zero and E is -O- the reaction is the well-known alkylation of phenols. The leaving group LG can be easily displaced reacting the two compounds in a polar solvent such as for example, but not
10 limited to, ACN or DMF, for one or more hours at room or higher temperature.

The same reaction conditions can be used for the conversion of a compound **II** into **IV** using a compound of formula **III**.

Compound of formula **VII** can be obtained using the addition of Grignard reagent (or other metal derivatives) to keto-containing compounds such as **VI** or **II**. The selection
15 of one of the two possible reactions depends on the availability of suitable Grignard's reagent or a precursor for its generation. The conversion of **VI** to **VII** can be performed in a solvent such as Et₂O or THF at a temperature below 0°C. The reaction is normally smooth and completes over a period ranging from one to overnight standing at room temperature. The same reaction conditions can be used for the reaction of **IV** with **V** to give **VIII**.

20 In another embodiment of the present invention, intermediate **VIII** can be used for the preparation of **I** using a different approach.

First **VIII** is deprotected to give an aldehyde that can be easily reacted with a compound of formula **XIV** under the reductive amination conditions described above for the conversion of **XIII** into **I**. The ester of formula **IX** obtained is then hydrolysed, under
25 acidic or basic aqueous condition as described for the preparation of **XI**, to give **X** that can be reacted under ester formation condition with a compound of formula HO-(CH₂)_s-R₂.

The presence of OH and NH moiety, for example in a compound of formula **X**, can affect the formation of the ester **I** as these functional groups can compete with the OH in

HO-(CH₂)_s-R₂. For this reason, it is worth considering performing such described reaction, for the conversion of **X** into **I**, using an intermediate in which the OH and NH are protected with suitable protecting group. For the purpose silyl ethers for the OH and carbamates for the NH are well known protecting groups, whose selection should not be limited to them
5 as it often depends on the complexity of the molecule and the presence of other functional groups in **Y** that can be not compatible with the reaction conditions required for introduction or removal of protecting groups.

A similar consideration applies to the moiety **Q** that contains a phenolic OH. In addition, in this case its reactivity can suggests its protection to avoid any competitive
10 reaction and formation of by-products. Benzyl is a suitable protecting group. The specific sequence of reaction for the preparation of **I** might be different from case to case, those skilled in the art will readily recognize the appropriate sequence, that must be evaluated considering the structure of linker **Y**.

In another embodiment of the invention, compounds **I** bearing an ester moiety in
15 the linker **Y**, can be prepared by treating a compound **XXVIII** with a compound **XXIX**, wherein **A2** is functionalized with OH, under the condensation reaction condition for the preparation of esters. A compound of formula **I**, wherein **C** is equal to **C1**, can be prepared reacting a compound **XXVIII** with a compound **XXIX** wherein **A2** is substituted with -NR₇ under the known reaction conditions for the preparation of amide starting from
20 carboxylic acid and amines. Using a similar approach it is possible to synthesize a compound of formula **I** wherein **C** has one of the other meanings cited, using intermediates featuring the correct functional groups that can react together to form one of meaning of **C**.

Compound **XXVIII** can be obtained by a reaction of a compound **XXV** with a suitable intermediate whose formula depends on the structure of the linker **Y**. An example
25 is the preparation of a compound of formula **I** wherein **E** is -O- which can be easily obtained treating a compound **XXV** with a compound **XXVII** where LG is leaving group such as chlorine, bromine or sulfonates. The alkylation reaction can be performed in polar solvents such as DCM, ACN or DMF at room or higher temperature in the presence of base that can

catalyse the reaction.

Compound **XXV** can be obtained from **XXX** via a reaction for ester formation already described for the conversion of **XI** into **XII**. Compound **XXX** is easily obtained from compound **VII** by mean of a standard reaction for the hydrolysis of esters.

5 Compounds **VII**, **XXX** and **XXV** are all suitable intermediates for the preparation of single stereoisomers of compound of formula **I**. For example, the mixtures of isomers of these three intermediates can be separated by mean of chiral chromatographic separation. In case, for example, **R2** is enantiomerically pure **J1**, **J2** or **J5**, a compound **XXV** is a mixture of diastereoisomers that can be separated by mean of a normal chromatographic
10 separation.

The absolute configuration of the stereogenic centre present in **VII** is conserved unmodified within the synthetic pathway applied for the conversion of **VII** into the final compound **I**.

In another embodiment, compounds **I** can be prepared following a different
15 synthetic approach in which a compound **XX** is reacted with a compound **XXII** under the transition metal catalysed cross-coupling reaction conditions, followed by reduction of the double bond $-(CH)_2-$, leading to a compound of formula **I** wherein $n = 2$. Alternatively, it can be prepared by reaction of a compound **XVII** with a compound **XXIII** under the condition described above for the reaction of a compound **II** with a compound **III**.

20 Intermediates **XX** and **XVII** can be prepared by reaction of compound **XIV** under reductive amination conditions, described above for the reaction of compound **XIII** with **XIV**, starting from compound **XIX** and **XVIII** respectively. Alternatively, compounds **XX** and **XVII** can be prepared by alkylation of compound **XIV** with compound **XXI** and **XX** respectively under alkylation condition described above for the preparation of compound
25 **VIII** by reaction of compound **VII** with compound **III**.


Compounds **XIV** can be obtained by simple reduction of the azide **XV**. The reaction can be accomplished by mean of a catalytic hydrogenation in the presence Palladium catalyst. The reaction occurs, in polar solvent such as methanol or ethanol, under hydrogen

atmosphere or under hydrogen transfer condition, using for example 1,4-cyclohexadiene or 1-methyl-1,4-cyclohexadiene as source of hydrogen. The reaction proceeds at room temperature. In case it is performed under hydrogen transfer conditions higher temperature can be required. Alternatively, the conversion can be accomplished under Staudinger
5 reaction conditions.

The azide **XV** can be easily prepared from **XXIV** by the well-known nucleophilic substitution of alkyl bromide with alkaline azide. The reaction proceeds at a temperature ranging from 50 to 80°C and in a polar solvent such as for example DMF or NMP and can be accelerated by the presence of alkaline iodide.

10 In another embodiment, compounds **XVII** can be prepared by reacting an intermediate **XXIV** with an amine **XXV**. This reaction is a common alkylation of amine in which the leaving group LG (normally chlorine, bromine or sulphate) is displaced by a nucleophile like the amine **XXV** as such or protected at the aminic moiety. Several methods to perform a reaction that normally occurs in a polar solvent at a temperature higher than
15 room temperature are described in the literature.

Compounds of formula **I** wherein s is 0 and **R2** is **J1**, **J2** or **J5** contain three stereogenic centres, as indicated below (wherein e.g. J= J2) with the symbol *. Therefore the structure of formula **I** is characterized by different stereoisomers.



20 Each diastereoisomer can be obtained theoretically by chromatographic separation of the mixture obtained by reacting racemic mixtures of the required intermediates. It is clear that this approach it is not convenient and that it can be used only for the separation of mixtures containing few diastereoisomers.

25 In a more convenient approach, the synthesis of each single stereoisomer can be accomplished using, in the reactions described above, only enantiomerically pure

intermediates.

The enantiomerically pure alcohol required for the preparation of compounds **I** wherein **R2** is **J1**, **J2** or **J5** are commercially available.

The preparation of single enantiomerically pure compounds **XXIV** wherein LG is
5 bromine are described in WO2005/092861 (cited by WO2007/107228).

The invention also provides pharmaceutical compositions of compounds **I** in admixture with one or more pharmaceutically acceptable carriers, for example those described in Remington's Pharmaceutical Sciences Handbook, XVII Ed., Mack Pub., N.Y., U.S.A.

10 Administration of the compounds of the invention may be accomplished according to patient needs, for example, orally, nasally, parenterally (subcutaneously, intravenously, intramuscularly, intrasternally and by infusion), by inhalation, rectally, vaginally, topically, locally, transdermally, and by ocular administration.

Various solid oral dosage forms can be used for administering compounds of the
15 invention including such solid forms as tablets, gelcaps, capsules, caplets, granules, lozenges and bulk powders. The compounds of the invention can be administered alone or combined with various pharmaceutically acceptable carriers, diluents (such as sucrose, mannitol, lactose, starches) and known excipients, including suspending agents, solubilizers, buffering agents, binders, disintegrants, preservatives, colorants, flavorants,
20 lubricants and the like. Time release capsules, tablets and gels are also advantageous in administering the compounds of the present invention.

Various liquid oral dosage forms can also be used for administering compounds of the invention, including aqueous and non-aqueous solutions, emulsions, suspensions, syrups, and elixirs. Such dosage forms can also contain suitable known inert diluents such
25 as water and suitable known excipients such as preservatives, wetting agents, sweeteners, flavorants, as well as agents for emulsifying and/or suspending the compounds of the invention. The compounds of the invention may be injected, for example, intravenously, in the form of an isotonic sterile solution. Other preparations are also possible.

Suppositories for rectal administration of the compounds of the invention can be prepared by mixing the compound with a suitable excipient such as cocoa butter, salicylates and polyethylene glycols.

Formulations for vaginal administration can be in the form of cream, gel, paste, foam, or spray formula containing, in addition to the active ingredient, such as suitable carriers, are also known.

For topical administration the pharmaceutical composition can be in the form of creams, ointments, liniments, lotions, emulsions, suspensions, gels, solutions, pastes, powders, sprays, and drops suitable for administration to the skin, eye, ear or nose. Topical administration may also involve transdermal administration via means such as transdermal patches.

For the treatment of the diseases of the respiratory tract, the compounds of the invention are preferably administered by inhalation.

Inhalable preparations include inhalable powders, propellant-containing metering aerosols or propellant-free inhalable formulations.

For administration as a dry powder, known single- or multi-dose inhalers may be utilized. In that case the powder may be filled in gelatine, plastic or other capsules, cartridges or blister packs or in a reservoir.

A diluent or carrier, generally non-toxic and chemically inert to the compounds of the invention, e.g. lactose or any other additive suitable for improving the respirable fraction may be added to the powdered compounds of the invention.

Inhalation aerosols containing propellant gas such as hydrofluoroalkanes may contain the compounds of the invention either in solution or in dispersed form. The propellant-driven formulations may also contain other ingredients such as co-solvents, stabilizers and optionally other excipients.

The propellant-free inhalable formulations comprising the compounds of the invention may be in form of solutions or suspensions in an aqueous, alcoholic or hydroalcoholic medium and they may be delivered by jet or ultrasonic nebulizers known

from the prior art or by soft-mist nebulizers such as Respimat®.

The compounds of the invention may be administered as the sole active agent or in combination with other pharmaceutical active ingredients including those currently used in the treatment of respiratory disorders, e.g. corticosteroids, P38 MAP kinase inhibitors, IKK2, HNE inhibitors, PDE4 inhibitors, leukotriene modulators, NSAIDs and mucus regulators.

The dosages of the compounds of the invention depend upon a variety of factors including the particular disease to be treated, the severity of the symptoms, the route of administration, the frequency of the dosage interval, the particular compound utilized, the efficacy, toxicology profile, and pharmacokinetic profile of the compound.

Advantageously, the compounds of formula I can be administered for example, at a dosage comprised between 0.001 and 1000 mg/day, preferably between 0.1 and 500 mg/day.

When the compounds of formula I are administered by inhalation route, they are preferably given at a dosage comprised between 0.001 and 500 mg/day, preferably between 0.1 and 200 mg/day.

The compounds of formula I may be administered for the prevention and/or treatment of broncho-obstructive or inflammatory diseases, such as asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), bronchial hyperreactivity, cough, emphysema or rhinitis; urological disorders such as urinary incontinence, pollakiuria, cystospasm, chronic cystitis and overactive bladder (OAB); gastrointestinal disorders such as bowel syndrome, spastic colitis, diverticulitis, peptic ulceration, gastrointestinal motility or gastric acid secretion; dry mouth; mydriasis, tachycardia; ophthalmic interventions cardiovascular disorders such as vagally induced sinus bradycardia.

The present invention will now be further described by way of the following examples.

The intermediate compounds for the synthesis of final compounds of general

formula (I) were obtained through the preparations herebelow described.

PREPARATIONS OF INTERMEDIATES AND EXAMPLES

Chemical Names of the compounds were generated with Structure To Name Enterprise 10.0 CambridgeSoft for all final compounds names or with ChemDraw Professional 15 for intermediates names.

Abbreviations

Et₂O = diethyl ether; Et₃N = triethyl amine; DCE = 1,2-dichloroethane; TEA = triethyl amine; DCC = N,N'-Dicyclohexylcarbodiimide; HOBt = Hydroxybenzotriazole; HATU = (Dimethylamino)-N,N-dimethyl(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yloxy)methaniminium hexafluorophosphate; EDC = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; DMAP = 4-dimethylaminopyridine; DMF = dimethylformamide; EtOAc = Ethyl acetate; RT = room temperature; THF = tetrahydrofuran; DCM = dichloromethane; MeOH = methyl alcohol; EtOH = ethylic alcohol; LHMDs = Lithium bis(trimethylsilyl)amide; m-CPBA = meta-Chloroperoxybenzoic acid; TFA = Trifluoroacetic acid; LC-MS = Liquid Chromatography/Mass Spectrometry; HPLC = high pressure liquid chromatography; MPLC = medium pressure liquid chromatography; SFC = Supercritical Fluid Chromatography

General Experimental details

NMR characterization:

¹H-NMR spectra were performed on a Varian MR-400 spectrometer operating at 400 MHz (proton frequency), equipped with: a self-shielded z-gradient coil 5 mm 1H/nX broad band probehead for reverse detection, deuterium digital lock channel unit, quadrature digital detection unit with transmitter offset frequency shift.

Alternatively a Bruker instrument was used (either Bruker Avance 400MHz or Bruker Avance III 400 MHz) operating at 400 MHz using the stated solvent at around room temperature unless otherwise stated.

In all cases, NMR data were consistent with the proposed structures. Chemical shift

are reported as δ values in ppm relative to trimethyl silane (TMS) as an internal standard. Coupling constants (J values) are given in hertz (Hz) and multiplicities are reported using the following abbreviation (s= singlet, d=doublet, t=triplet, q=quartet, m=multiplet, br=broad, nd=not determined).

5 LC/UV/MS Analytical Methods

LC/MS retention times are estimated to be affected by an experimental error of ± 0.5 min.

Methods 1 – 2 – 3 - 4:

LC-UV-MS instrument (Waters Acquity UPLC system) equipped with UV (PDA
10 detector) and mass spectrometer (Acquity QDa Detector). UV acquisition range 210 - 400 nm. MS acquisition range 110 – 1200 amu.

LC-UV

Eluent A: water/ACN 95/5 (0.05% HCOOH), Eluent B: ACN/water 95/5 (0.05%
HCOOH)

15 Flow 1 mL/min

Column Temperature: 40°C

Gradient 1:

Time (min)	%A	%B
0.00	99	1
1.50	0.1	99.9
1.90	0.1	99.9
2.00	99	1

Gradient 2:

Time (min)	%A	%B
0.00	99	1
3.50	0.1	99.9
3.90	0.1	99.9
4.00	99	1

Methods:

Method	Gradient	Column
1	Gradient 1	Acquity UPLC BEH C18 1.7 um 50x2.1 mm
2	Gradient 1	Acquity UPLC CSH C18 1.7 um 50x2.1 mm
3	Gradient 2	Kinetex UPLC C8 1.7 um 50x2.1 mm
4	Gradient 2	Acquity UPLC CSH C18 1.7 um 50x2.1 mm

MS instrument: Waters ZQ (or equivalent)Polarity ES

Capillary (kV) 3.20

5 Cone (V) 25.00

Extractor (V) 3.00

RF Lens (V) 0.1

Polarity ES-

Capillary (kV) 3.40

10 Cone (V) 24.00

Extractor (V) 2.00

RF Lens (V) 0.2

Source Temperature (°C) 130

Desolvation Temperature (°C) 400

15 Cone Gas Flow (L/Hr) 80

Desolvation Gas Flow (L/Hr) 800

Mass range: 60 to 1200

Scan time (sec): 0.4

Method 5

20 LC-UV-MS instrument (Waters Acquity UPLC system) equipped with UV (PDA detector) and mass spectrometer (Waters TQS Detector). UV acquisition range 210 - 400 nm. MS acquisition range 100 – 1000 amu ESI+ and ESI-

Solvents:A% HCOONH₄ buffer 0,025 M pH 3

B% ACN + 0.1% HCOOH

Flow (ml/min) 0.5 mL/min

Stop time (mins) 10 min

Column Temperature 55°C

5 UV acquisition range (nm): 254

Injection volume (μl): 2

Column: Kinetex 1.7u C8 100A 100 x 2.1 mm

Gradient:

Time (min)	%A	%B
0.00	99	1
0.5	99	1
3	70	30
6.5	50	50
7.5	20	80
8	20	80
8.1	99	1
10	99	1

10 Method 6

10cm_Formic_ACE 3 C18 AR_HPLC_CH3CN

HPLC Setup

Solvents: - Acetonitrile (Far UV grade) with 0.1% (V/V) formic acid

Water (High purity via PureLab Ultra unit) with 0.1% formic acid

15 Column: - Hichrom ACE 3 C18-AR mixed mode column 100x4.6mm

Flow Rate: - 1 ml/min

Gradient:

Time (min)	%A	%B
0.00	98	2
3	98	2
12	0	100
15.4	0	100
15.5	98	2
17	98	2

Injectons 0.2-10ul

Maximum pressure setting 400 bar.

5 Instrument: Agilent 1100, Binary Pump, Agilent Sampler and Agilent DAD detector

Diode array detection: (300nm, Band Width 200nm; Ref. 450nm, Band Width 100nm)

Method 7

10 15cm_Formic_Ascentic_HPLC_CH3CN

HPLC Setup

Solvents: - Acetonitrile (Far UV grade) with 0.1% (V/V) formic acid

Water (High purity via PureLab Ultra unit) with 0.1% formic acid

Column: - Supelco, Ascentis® Express C18 or Hichrom Halo C18, 2.7µm C18,

15 150 x 4.6mm.

Flow Rate: - 1ml/min

Gradient:

Time (min)	%A	%B
0.00	96	4
3	96	4
9	0	100
13.6	0	100
13.7	96	4
15	96	4

Injectons 0.2-10 ul

Maximum pressure setting 400 bar.

Instrument: Agilent 1100, Binary Pump, Agilent Sampler and Agilent DAD detector

Diode array detection: (300nm, Band Width 200nm; Ref. 450nm, Band Width
5 100 nm)

Method 8

10cm_ESCI_Formic_MeCN

HPLC Setup

Solvents: - Acetonitrile (Far UV grade) with 0.1% (V/V) formic acid

10 Water (High purity via PureLab Option unit) with 0.1% formic acid

Column: - Phenomenex Luna 5 μ C18 (2), 100 x 4.6mm. (Plus guard cartridge)

Flow Rate: - 2ml/min

Gradient:

Time (min)	%A	%B
0.00	95	5
3.5	5	95
5.5	5	95
5.6	95	5
6.5	95	5

Injections 2-7 μ l (concentration ~ 0.2 -1mg/ml).

15 UV detection via HP or Waters DAD

Start Range (nm) 210 End Range (nm) 400 Range interval (nm)

4.0

Other wavelength traces are extracted from the DAD data.

Optional ELS detection using Polymer Labs ELS-1000.

20 MS detection: Micromass ZQ, single quadrupole LC-MS or Quattro Micro LC-MS-MS.

Flow splitter gives approximately 300 μ l/min to mass spec

Scan range for MS Data (m/z)

Start (m/z) 100

End (m/z) 650 or 1500 when required

With +ve / -ve switching

Ionisation is routinely ESCI an option which gives both ESI and APCI data from a single run.

5 Typical ESI voltages and temperatures are:

Source 120-150C 3.5KV capillary 25V cone

Typical APCI voltages and temperatures are:

Source 140-160C 17uA corona 25V cone

Method 9. HPLC-AB

10 - HPLC with UV-Vis or DAD detector

- column: Waters Symmetry C18, 3.9x150mm, 5.0µm

Eluents:

- (A) 0,1% formic acid -water solution

- (B) 0,1% formic acid -ACN solution

15 - injection volume: 3µL

- flow: 1.0 ml/min

Gradient:

Time (min)	%A	%B
0.00	95	5
3	95	5
9	83	17
15	83	17
25	20	80
27	20	80
28	95	5
30	95	5

Column compartment:

- column temperature: 25°C

20 - time of analysis: 30min

Detector: - wave length: 220 nm

Method 10. Chiral-HPLC

Equipment:

- HPLC with UV-Vis or DAD detector
- column: Chiralpak IC, 4,6mmx250mmx 5um

5

Eluents:

- (A)IPA + 0.1% TFA
- (B)Hexane + 0.1% TFA

Autosampler:

- injectionvolume:3µL

10

Pump:

- flow: 1.0ml/min
- 90% B

Column compartment:

- column oven temperature: 25 °C

15

- time of analysis: 30 min

Detector:

- wave length: 220 nm

Method 11

10cm_Formic_AQ

20

UPLC Setup

Solvents: - B Acetonitrile (Far UV grade) with 0.1% (V/V) formic acid

A Water (High purity via PureLab Option unit) with 0.1% formic

acid

Column: - Acquity UPLC HSS C18 1.8um 100 x 2.1mm. (Plus guard cartridge)

25

Flow Rate: - 0.5 ml/min

Gradient:

Time (min)	%A	%B
0.00	95	5
1.2	95	5
3.5	0	100
4.9	0	100
5	95	5
6	95	5

Injectons 0.5-2ul

UV detection via Waters DAD

5 Start Range (nm) 210 End Range (nm) 400 Resolution (nm)

1.2

MS detection: Waters SQD2, single quadrapole UPLC-MS

Scan range for MS Data (m/z)

Start (m/z) 100

10 End (m/z) 700 or 1500 when required

With +ve / -ve switching

Ionisation is ESI.

ESI voltages and temperatures are:

Source 150C 3.5KV capillary 25V cone

15 **Method 12**

10cm_Bicarb_AQ

UPLC Setup

Solvents: - Acetonitrile (Far UV grade)

20 Water (High purity via PureLab Option unit) with 10mM ammonium bicarbonate
(ammonium hydrogen carbonate)

Column: - Acquity UPLC BEH Shield RP18 1.7um 100 x 2.1mm. (Plus guard cartridge)

Flow Rate: - 0.5ml/min

Gradient: - A: Water / Basic B: MeCN / Basic

Time	A%	B%
0.00	95	5
1.20	95	5
3.5	0	100
4.90	0	100
5.00	95	5
6.00	95	5

Typical Injections 0.5-2ul (concentration ~ 0.2 -1mg/ml).

UV detection via Waters DAD

Start Range (nm) 210 End Range (nm) 400 Resolution (nm)

5 1.2

Other wavelength traces are extracted from the DAD data.

MS detection: Waters SQD2, single quadrupole UPLC-MS

Flow splitter gives approximately 300ul/min to mass spec

Scan range for MS Data (m/z)

10 Start (m/z) 100

End (m/z) 700 or 1500 when required

With +ve / -ve switching

Preparative reverse-phase HPLC conditions

Preparative HPLC

15 Waters Micromass ZQ/Sample manager 2767

Photodiode array detector 2996;

Column: XTerra Prep MS C18 Column (5 µm, 19 x 150 mm, Waters)

Flow rate: 20 ml/min with MS detection

UV wavelength: 254 nm.

20 Mobile phase: Solvent A (water:MeCN:HCOOH 95:5:0.05); Solvent B
(water:MeCN:HCOOH 5:95:0.05)

Gradient:

Time (min)	%A	%B
0.00	100.0	0.00
1.00	100	0.00
10.00	0.00	100.0
11.00	0.00	100.0
12.00	100.0	0.00

Where the preparation of starting materials is not described, these are commercially available, known in the literature, or readily obtainable by those skilled in the art using standard procedures.

5 Flash chromatography is carried out using an Isolera MPLC system (manufactured by Biotage) using pre-packed silica gel or reverse-phase cartridges (supplied by Biotage).


Many of the compounds described in the following Examples have been prepared from stereochemically pure starting materials, for example 95% ee.


10 The stereochemistry of the compounds in the Examples, where indicated, has been assigned on the assumption that absolute configuration at resolved stereogenic centers of starting materials is maintained throughout any subsequent reaction conditions.

In the procedures that follow, after each starting material, reference to a compound number is sometimes provided. This is provided merely for assistance to the skilled chemist. The starting material may not necessarily have been prepared from the batch
15 referred to.

When reference is made to the use of a “similar” or “analogous” procedure, as will be appreciated by those skilled in the art, such a procedure may involve minor variations, for example reaction temperature, reagent/solvent amount, reaction time, work-up conditions or chromatographic purification conditions.


20 **Preparation of (*R*)-5-(2-Amino-1-hydroxyethyl)-8-hydroxyquinolin-2(1*H*)-one hydrochloride**



Step 1; 8-(Benzyloxy)-5-(2-bromoacetyl)quinolin-2(1H)-one

A suspension of 5-acetyl-8-(benzyloxy)quinolin-2(1H)-one (19.4 g, 66.4 mmol) in anhydrous THF (240 mL) and anhydrous methanol (165 mL) was added with a solution of *tetra*-n-butylammonium tribromide (54.5 g, 113.0 mmol) in anhydrous THF (130 mL) drop-wise over 1.5 hours. The resulting solution was stirred at room temperature overnight before concentrating under reduced pressure without heating. The residue was re-dissolved in methanol (200 mL). Saturated aqueous ammonium chloride solution (390 mL) was added with ice-cooling. The resulting suspension was filtered and the solid washed with water and air-dried under vacuum. The solid was suspended in DCM and methanol (1:1 v/v, 100 mL) for 90 minutes. The solid was collected by filtration, washed with DCM and air-dried to afford the title compound (18.0 g, 73%).

¹H NMR (400 MHz, DMSO-d₆): δ 11.07 (s, 1 H), 8.51 (d, *J* = 10.0 Hz, 1 H), 7.94-7.83 (m, 1 H), 7.60 (d, *J* = 7.5 Hz, 2 H), 7.44-7.27 (m, 4 H), 6.79-6.65 (m, 1 H), 5.53-5.39 (s, 2 H), 4.93 (s, 2 H)


Step 2; (R)-8-(Benzyloxy)-5-(2-bromo-1-hydroxyethyl)quinolin-2(1H)-one

8-(Benzyloxy)-5-(2-bromoacetyl)quinolin-2(1H)-one (26.0 g, 69.9 mmol) and (*R*)-3,3-diphenyl-1-methyltetrahydro-3*H*-pyrrolo[1,2-*c*][1,3,2]oxazaborole (21.3 g, 76.8 mmol) were azeotroped with toluene (× 3) then suspended in anhydrous THF (400 mL)

under an atmosphere of nitrogen. The suspension was cooled to -20 °C (external temperature) and borane dimethyl sulfide complex solution (45.4 mL, 90.8 mmol, 2.0 M solution in THF) was added by syringe pump over 3 hours. After complete addition the reaction mixture was stirred for one hour before quenching with methanol (25 mL). The reaction was warmed to room temperature over 20 minutes. The mixture was concentrated under reduced pressure and the residue was suspended in aqueous hydrochloric acid (500 mL, 1 M solution) and stirred at room temperature for 18 hours. After this time the solid was collected by filtration and washed with water (3 × 100 mL). The solid was partially dissolved in ethyl acetate and heated at reflux for 2 hours. The remaining solid was removed by hot filtration and the filtrate was evaporated to afford the title compound. The solid collected from the hot ethyl acetate was again partially dissolved in ethyl acetate and heated at reflux for 2 hours then filtered to give filtrate containing pure product. This process was repeated four more times. The combined solid was recrystallised from ethyl acetate and petroleum ether to afford the title compound (20.0 g, 76%).

¹H NMR (400 MHz, DMSO-d₆): δ 10.68 (s, 1 H), 8.19 (d, *J* = 9.9 Hz, 1 H), 7.58 (d, *J* = 7.5 Hz, 2 H), 7.41-7.36 (m, 2 H), 7.34-7.29 (m, 1 H), 7.23-7.19 (m, 2 H), 6.57 (d, *J* = 9.8 Hz, 1 H), 5.94 (d, *J* = 4.7 Hz, 1 H), 5.31 (s, 2 H); 5.25-5.19 (m, 1 H), 3.71-3.58 (m, 2 H).

Step 3; (R)-8-(Benzyloxy)-5-(2-bromo-1-((*tert*-butyldimethylsilyl)oxy)ethyl)quinolin-2(1*H*)-one




2,6-Lutidine (6.9 mL, 59.5 mmol) was added to a solution of (R)-8-(benzyloxy)-5-(2-bromo-1-hydroxyethyl)quinolin-2(1*H*)-one (10.1 g, 27.0 mmol) in DCM (100 mL) at 0 °C. The reaction mixture was stirred for 5 minutes then *tert*-butyldimethylsilyl

trifluoromethanesulfonate (13.0 mL, 56.8 mmol) was added dropwise over 15 minutes. The mixture was stirred at 0°C for 30 minutes, followed by room temperature overnight. After this time the reaction was quenched with saturated aqueous sodium bicarbonate solution and extracted with DCM (x 3). The combined organic extracts were dried (magnesium sulfate), filtered and concentrated under reduced pressure. *Iso*-hexane (500 mL) was added to the crude material and the resulting solid collected by filtration. The solid was recrystallised from ethyl acetate and petroleum ether (40 : 60) to afford the title compound (11.3 g, 85%).

¹H NMR (400 MHz, CDCl₃): δ 9.19 (s, 1 H), 8.23 (dd, *J* = 9.9, 4.4 Hz, 1 H), 7.43 (d, *J* = 4.6 Hz, 5 H), 7.17 (dd, *J* = 8.3, 4.5 Hz, 1 H), 7.03 (dd, *J* = 8.2, 4.4 Hz, 1 H), 6.71 (dd, *J* = 9.9, 3.7 Hz, 1 H), 5.18 (d, *J* = 4.5 Hz, 3 H), 3.63-3.56 (m, 1 H), 3.49 (dd, *J* = 10.4, 4.8 Hz, 1 H), 0.88 (t, *J* = 4.4 Hz, 9 H), 0.14 (d, *J* = 4.4 Hz, 3 H), -0.11 (d, *J* = 4.4 Hz, 3 H).

Step 4; (R)-5-(2-Azido-1-((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one



15

(R)-8-(Benzyloxy)-5-(2-bromo-1-((*tert*-butyldimethylsilyl)oxy)ethyl)-quinolin-2(1H)-one (10.0 g, 20.5 mmol) was dissolved in DMF (180 mL) and water (20 mL). Sodium iodide (3.39 g, 22.6 mmol) and sodium azide (1.47 g, 22.6 mmol) were added sequentially. The reaction mixture was stirred at RT until all the solid was in solution. The solution was heated at 80°C for 40 hours then cooled to RT and diluted with ethyl acetate (300 mL). The mixture was washed with water, brine (x2) and the organic extract was dried (magnesium sulfate), filtered and concentrated under reduced pressure. The crude residue was triturated with *iso*-hexane to afford the desired compound (8.16 g, 88%). Used without further purification in the next step.

¹H NMR (400 MHz, CDCl₃): δ 9.19 (s, 1 H), 8.18 (d, *J* = 9.9 Hz, 1 H), 7.45-7.36 (m, 5 H), 7.20 (d, *J* = 8.3 Hz, 1 H), 7.04 (d, *J* = 8.3 Hz, 1 H), 6.70 (dd, *J* = 9.9, 2.2 Hz, 1 H), 5.19-5.13 (m, 3 H), 3.48 (dd, *J* = 12.7, 8.1 Hz, 1 H), 3.26 (dd, *J* = 12.7, 3.8 Hz, 1 H), 0.89 (s, 9 H), 0.14 (s, 3 H), -0.11 (s, 3 H).


5 **Step 5; (R)-5-(2-Amino-1-hydroxyethyl)-8-hydroxyquinolin-2(1H)-one hydrochloride**



A solution of (R)-5-(2-azido-1-((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (4.50 g, 10.0 mmol) in ethanol (50 mL) was added with
10 10% palladium on charcoal (4.50 g) followed by 1-methyl-1,4-cyclohexadiene (11.0 mL, 97.9 mmol). The reaction was warmed to 60 °C and then stirred at 60 °C for 2 hours. The reaction mixture was allowed to cool and filtered through a pad of celite. The filtercake was washed with further ethanol and the filtrate was evaporated under reduced pressure. The residue was evaporated from *iso*-propanol (x2) and dissolved in *iso*-propanol (30 mL).
15 HCl-dioxane (4M, 50 mL, 200 mmol) was added and the reaction mixture stirred at room temperature for 18 hours. The resulting suspension was filtered, the filtercake washed with ether and the solid dried under vacuum in the presence of P₂O₅ to afford the title compound (1.65 g, 62%).


¹H NMR (400 MHz, MeOD): δ 7.71 (d, *J* = 9.8 Hz, 1 H), 6.57 (d, *J* = 8.2 Hz, 1 H),
20 6.31 (d, *J* = 8.2 Hz, 1 H), 6.02 (dd, *J* = 9.8, 6.5 Hz, 1 H), 4.58 (dd, *J* = 9.6, 3.5 Hz, 1 H), 2.47-2.31 (m, 2 H).

General scheme for the preparation of compounds of example 1 – Scheme 1




Example 1

- (1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-((*R*)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate (Compound 1)



Step 1; Ethyl 2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetate




- A stirred solution of ethyl benzoylformate (1.78 g, 10.0 mmol) in THF (30 mL) at -78 °C was added drop-wise with a solution of 3-benzyloxyphenyl magnesium bromide (1.0M solution in THF, 11.0 mL, 11.0 mmol) over 20 minutes. The reaction mixture was allowed to warm slowly to room temperature and stirred at room temperature for 18 hours. The reaction mixture was quenched with 10% aqueous HCl and then extracted with ethyl acetate. The ethyl acetate extract was washed with brine, dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified

by flash column chromatography (eluent – 100% *iso*-hexane to 12:1 *iso*-hexane/ethyl acetate) to afford the title compound (2.23 g, 62 %).

¹H NMR (400MHz, CDCl₃); δ 7.44 – 7.23 (m, 11 H), 7.13 – 7.09 (m, 1 H), 7.05 – 7.04 (m, 1 H), 6.95 – 6.92 (m, 1 H), 5.04 (s, 2H), 4.34 – 4.32 (m, 2 H), 4.23 (s, 1 H), 1.28 – 1.25 (m, 3 H).


Step 2; Ethyl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate



10% Pd-C (2.0 g) was added to a solution of ethyl 2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetate (2.22 g, 6.13 mmol) in ethanol (20 mL). After 5 minutes 1-methyl-1,4-cyclohexadiene (3.5 mL, 31.1 mmol) was added and the reaction mixture carefully heated to 70 °C [CARE – Vigorous evolution of gas]. The reaction mixture was heated at this temperature for 30 minutes and the reaction mixture was allowed to cool. The suspension was filtered and the filter-cake washed with further ethanol. The filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 4:1 *iso*-hexane/ethyl acetate) to afford the title compound (1.20 g, 72 %).

¹H NMR (400MHz, CDCl₃); δ 7.44 – 7.42 (m, 2 H), 7.36 – 7.29 (m, 3 H), 7.22 – 7.19 (m, 1 H), 7.02 – 6.99 (m, 1 H), 6.93 - 6.92 (m, 1 H), 6.80 – 6.78 (m, 1 H), 4.81 (s, 1 H), 4.35 – 4.27 (m, 2 H), 4.23 (s, 1 H), 1.30 – 1.26 (m, 3 H).

Step 3; Ethyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate




A stirred solution of ethyl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate (1.19 g,

4.38 mmol) in DMF (15 mL) was added with potassium carbonate (0.907 g, 6.57 mmol). The reaction mixture was stirred at room temperature for 10 minutes and then 2-(4-chlorobutyl)-1,3-dioxolane (0.78 mL, 5.25 mmol) was added. The reaction mixture was heated at 80 °C for 42 hours. 10% Aqueous potassium hydrogen sulfate was added to the reaction mixture which was then extracted with ethyl acetate. The organic phase was washed with brine, dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 4:1 *iso*-hexane/ethyl acetate) to afford the title compound (1.07 g, 61 %).

¹H NMR (400MHz, CDCl₃); δ 7.44 – 7.42 (m, 2 H), 7.34 – 7.31 (m, 3 H), 7.25 – 7.21 (m, 1 H), 7.02 – 6.98 (m, 2 H), 6.85 – 6.82 (m, 1 H), 4.88 – 4.86 (m, 1 H), 4.35 – 4.30 (m, 2 H), 4.22 (s, 1 H), 3.97 – 3.83 (m, 6 H), 1.83 – 1.71 (m, 4 H), 1.61 – 1.59 (m, 2H), 1.28 – 1.22 (m, 3H).

Step 4; 2-(3-(4-(1,3-Dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetic

acid




A stirred solution of ethyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate (1.06 g, 2.65 mmol) in ethanol (10 mL) was added with 2M aqueous sodium hydroxide (10 mL). The reaction mixture was stirred at room temperature for 72 hours. The reaction mixture was treated with 10% aqueous potassium hydrogen sulfate and extracted with DCM (x2) and ethyl acetate. The combined organic phases were dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure to afford the title compound (1.09 g, >100%).

¹H NMR (400MHz, CDCl₃); δ 7.44 – 7.42 (m, 2 H), 7.34 – 7.31 (m, 3 H), 7.25 – 7.21 (m, 1 H), 7.02 – 6.98 (m, 2 H), 6.85 – 6.82 (m, 1 H), 4.88 – 4.86 (m, 1H), 4.10 – 3.83

(m, 6H), 1.83 – 1.71 (m, 4 H), 1.61 – 1.55 (m, 2H).


Step 5; *tert*-Butyl 4-(tosyloxymethyl)piperidine-1-carboxylate



A stirred solution of *tert*-butyl 4-(hydroxymethyl)piperidine-1-carboxylate (5.0 g, 23.2 mmol) in anhydrous pyridine (18.5 mL) at 0 °C under nitrogen was added with *p*-toluenesulfonyl chloride (4.87 g, 25.55 mmol) in one portion. The reaction was stirred at 0 °C for 100 minutes before warming to room temperature. After 18 hours the reaction mixture was poured into water and extracted with ethyl acetate (x3). The combined organic extracts were washed with aqueous 1M hydrochloric acid (x2), brine, dried (magnesium sulfate), filtered and evaporated under reduced pressure to yield the title compound as a yellow solid (7.87 g, 91%).

¹H NMR (400 MHz, CDCl₃): δ 7.78 (d, J=8.4 Hz, 2H), 7.35 (d, J=8.0 Hz, 2H), 4.15 - 4.07 (m, 2H), 3.85 (d, J=6.5 Hz, 2H), 2.68 - 2.60 (m, 2H), 2.46 (s, 3H), 1.88 - 1.78 (m, 1H), 1.66 - 1.59 (m, 2H), 1.44 (s, 9H), 1.16 - 1.04 (m, 2H).

Step 6; *tert*-Butyl 4-((2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate




A stirred solution of 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetic acid (1.0 g, 2.69 mmol) in DMF (10 mL) was added with potassium carbonate (0.558 g, 4.04 mmol). After 10 minutes *tert*-butyl 4-(tosyloxymethyl)piperidine-1-carboxylate (1.29 g, 3.50 mmol) was added and the resulting mixture heated at 80 °C for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (twice). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by flash column

chromatography (eluent – 100% *iso*-hexane to 4:1 *iso*-hexane/ethyl acetate) to afford the title compound (1.42 g, 93 %).


¹H NMR (400MHz, CDCl₃); δ 7.44 – 7.42 (m, 2 H), 7.34 – 7.31 (m, 3 H), 7.25 – 7.21 (m, 1 H), 7.02 – 6.98 (m, 2 H), 6.85 – 6.82 (m, 1 H), 4.88 – 4.86 (m, 1 H), 4.19 – 3.83 (m, 11H), 2.60 (m, 2H), 1.88 – 1.71 (m, 4H), 1.61 – 1.58 (m, 3H), 1.54 - 1.46 (m, 11H), 1.26 -1.24 (m, 2H).

Step 7; Piperidin-4-ylmethyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate hydrochloride



A solution of HCl in dioxane (4M, 10 mL) was added to *tert*-butyl 4-((2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetoxymethyl)piperidine-1-carboxylate (1.40 g, 2.46 mmol) and the mixture stirred at room temperature for 2 hours. The solvent was evaporated under reduced pressure and the residue used directly in the next step with no purification.

Step 8; (1-Benzylpiperidin-4-yl)methyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate



A stirred solution of piperidin-4-ylmethyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate hydrochloride in acetonitrile (15 mL) was added with triethylamine (0.683 mL, 4.91 mmol). The reaction mixture was stirred at room temperature for 20 minutes. Benzaldehyde (0.325 mL, 3.20 mmol) was added and the mixture stirred for a further 30 minutes. This mixture was added with sodium triacetoxyborohydride (1.04 g, 4.91 mmol) followed by acetic acid (0.562 mL, 9.82 mmol).

The resulting mixture was stirred at room temperature for 72 hours. The reaction mixture was diluted with ethyl acetate and washed with 10% aqueous potassium carbonate and brine. The organic phase was dried and the filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% DCM to 25:1 DCM/methanol) to afford the title compound (0.804 g, 58 %).

^1H NMR (400MHz, CDCl_3); δ 7.42 – 7.38 (m, 2H), 7.35 – 7.19 (m, 9H), 7.00 – 6.94 (m, 2H), 6.83 (ddd, $J = 8.2, 2.5, 0.9$ Hz, 1H), 5.30 (s, 2H), 4.88 – 4.83 (m, 1H), 4.22 (s, 1H), 4.15 – 4.06 (m, 2H), 4.01 – 3.80 (m, 6H), 3.45 (s, 2H), 2.81 (d, $J = 11.3$ Hz, 2H), 1.92 – 1.66 (m, 6H), 1.70 – 1.46 (m, 3H), 1.28 – 1.13 (m, 2H).




Step 9; (Compound 1)




A stirred solution of (1-benzylpiperidin-4-yl)methyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate (0.390 g, 0.70 mmol) in THF (4 mL) was added with 10% aqueous hydrochloric acid (8 mL). The resulting mixture was stirred at room temperature for 2 hours. The mixture was added with 10% aqueous potassium carbonate and then extracted with ethyl acetate (twice). The combined organic phases were dried and the filtrate was evaporated under reduced pressure. The residue was dissolved in methanol




(3 mL) and added to a pre-stirred (10 minutes) mixture of (*R*)-5-(2-amino-1-hydroxyethyl)-8-hydroxyquinolin-2(1*H*)-one hydrochloride (0.193 g, 0.60 mmol, 80% purity) and triethylamine (0.167 mL, 1.20 mmol) in methanol (3 mL). This mixture was stirred at room temperature for 1 hour and then sodium triacetoxyborohydride (0.254 g, 1.20 mmol) followed by acetic acid (0.137 mL, 2.39 mmol) were added. The reaction mixture was stirred for a further 1 hour. The reaction mixture was diluted with *iso*-butanol and washed with water. The aqueous phase was extracted with further *iso*-butanol. The combined *iso*-butanol extracts were evaporated under reduced pressure. The residue was purified by reverse phase preparative HPLC to afford the title compound (1).

The following compounds were prepared as described in Example 1 with the corresponding commercially available substituted ethyl benzoylformate replacing ethyl

benzoylformate in Step 1. If the required substituted ethyl benzoylformate was not available it was prepared according to Scheme 2 and Procedure 1 using the appropriate Grignard reagent.


	R_t (min)	Structure	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
1	8.46		6	(DMSO-d ₆ /D ₂ O); δ 8.19 (d, J = 9.9 Hz, 1H); 7.45 (s, 5H); 7.30 (d, J = 7.6 Hz, 6H); 7.22 (d, J = 8.1 Hz, 1H); 7.16 (d, J = 8.2 Hz, 1H); 7.00 (d, J = 8.2 Hz, 1H); 6.86-6.79 (m, 3H); 6.60 (d, J = 9.8 Hz, 1H); 5.31-5.24 (m, 1H); 4.20 (s, 3H); 4.05-3.71 (m, 4H); 3.32-3.24 (m, 2H); 3.07 (s, 2H); 2.96 (s, 2H); 2.87 (s, 2H); 1.85 (s, 1H); 1.76-1.61 (m, 5H); 1.45-1.27 (m, 3H)	TFA
2	7.29		7	(DMSO-d ₆); δ 8.31 (s, 2H), 8.19 (d, J=9.9 Hz, 1H), 7.32 - 7.18 (m, 9H), 7.14 - 7.09 (m, 3H), 6.96 (d, J=8.2 Hz, 1H), 6.88 - 6.83 (m, 3H), 6.53 (d, J=9.9 Hz, 1H), 5.20 (dd, J=4.5, 8.2 Hz, 1H), 4.01 - 3.96 (m, 2H), 3.89 (dd, J=6.3, 6.3 Hz, 2H), 3.40 (s, 2H), 2.89 - 2.69 (m, 6H), 2.28 (s, 3H), 1.83 (dd, J=9.5, 11.8 Hz, 2H), 1.72 - 1.63 (m, 2H), 1.60 - 1.37 (m, 7H), 1.17 - 1.04 (m, 2H).	Formate
3	7.23		7	(DMSO-d ₆); δ 10.54 - 10.49 (m, 2H), 8.60 - 8.59 (m, 2H), 7.48 (s, 5H), 7.38 - 7.33 (m, 2H), 7.25 (dd, J=8.0, 8.0 Hz, 1H), 7.18 - 7.13 (m, 3H), 6.99 (d, J=8.2 Hz, 1H), 6.90 - 6.85 (m, 3H), 6.58 (d, J=9.9 Hz, 1H), 6.33-6.01 (s, 1H), 5.32 (dd, J=2.2, 9.6 Hz, 1H), 4.31 - 4.20 (m, 2H), 4.01 (d, J=6.4 Hz, 2H), 3.91 (dd, J=6.3, 6.3 Hz, 2H), 3.34 (d, J=14.1 Hz, 2H), 3.12 - 2.86 (m, 6H), 1.85 - 1.83 (m, 1H), 1.77 (s, 1H), 1.72 - 1.64 (m, 6H), 1.48 - 1.30 (m, 4H).	TFA

4	7.23		7	(DMSO-d ₆); δ 10.52 (s, 2H), 9.49 - 9.49 (m, 1H), 8.60 - 8.59 (m, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.48 (s, 6H), 7.42 - 7.35 (m, 1H), 7.26 (dd, J=7.9, 7.9 Hz, 1H), 7.19 - 7.10 (m, 4H), 6.99 (d, J=8.2 Hz, 1H), 6.91 - 6.86 (m, 4H), 6.58 (d, J=9.9 Hz, 1H), 6.14 - 6.14 (m, 1H), 5.30 (dd, J=2.2, 9.7 Hz, 1H), 4.31 - 4.21 (m, 3H), 4.02 (d, J=6.5 Hz, 2H), 3.91 (dd, J=6.2, 6.2 Hz, 2H), 3.34 (d, J=11.8 Hz, 2H), 3.10 - 2.86 (m, 5H), 1.89 - 1.84 (m, 1H), 1.77 - 1.64 (m, 5H), 1.48 - 1.30 (m, 4H).	TFA
5	7.28		7	(DMSO-d ₆); δ 10.52 - 10.46 (2H, m), 9.49 - 9.36 (1H, m), 8.57 - 8.56 (2H, m), 8.16 (1H, d, J=7.3 Hz), 7.48 (5H, s), 7.26 - 7.19 (2H, m), 7.15 (2H, d, J=8.2 Hz), 7.09 (2H, dd, J=8.3, 8.3 Hz), 6.99 (1H, d, J=8.2 Hz), 6.89 - 6.83 (3H, m), 6.58 (2H, dd, J=1.9, 9.9 Hz), 6.17 - 6.16 (1H, m), 5.31 (1H, d, J=8.9 Hz), 4.26 (2H, d, J=4.9 Hz), 4.01 (2H, d, J=6.4 Hz), 3.91 (2H, dd, J=6.3, 6.3 Hz), 3.34 (2H, d, J=11.5 Hz), 3.11 - 2.86 (6H, m), 2.27 (3H, s), 1.84 (1H, d, J=3.1 Hz), 1.76 - 1.64 (6H, m), 1.48 - 1.34 (4H, m);	TFA
6	8.12		7	(MeOD); δ 8.38 (d, J=9.7 Hz, 1H), 7.54 - 7.47 (m, 5H), 7.44 (d, J=7.8 Hz, 1H), 7.35 - 7.28 (m, 3H), 7.27 (s, 1H), 7.19 - 7.10 (m, 2H), 7.05 (d, J=8.2 Hz, 1H), 6.98 - 6.91 (m, 2H), 6.70 (d, J=9.9 Hz, 1H), 5.41 (t, J=6.9 Hz, 1H), 4.29 (s, 2H), 4.12 - 4.02 (m, 4H), 3.50 - 3.44 (m, 2H), 3.25 (d, J=6.4 Hz, 2H), 3.12 (t, J=9.1 Hz, 2H), 2.97 (t, J=12.2 Hz, 2H), 2.06 - 2.00 (m, 1H), 1.93 - 1.82 (m, 6H), 1.63 (t, J=17.3 Hz, 2H), 1.57 - 1.42 (m, 2H).	TFA

7	8.15		7	(MeOD); δ 8.38 (d, J=9.7 Hz, 1H), 7.53 - 7.48 (m, 5H), 7.32 - 7.19 (m, 4H), 7.13 (s, 1H), 7.07 - 6.99 (m, 3H), 6.93 - 6.86 (m, 2H), 6.70 (d, J=9.8 Hz, 1H), 5.43 - 5.38 (m, 1H), 4.28 (s, 2H), 4.10 (d, J=6.5 Hz, 2H), 4.01 (dd, J=6.1, 6.1 Hz, 2H), 3.51 - 3.45 (m, 2H), 3.25 (d, J=6.5 Hz, 2H), 3.15 - 3.09 (m, 2H), 2.97 (dd, J=10.8, 13.1 Hz, 2H), 2.29 - 2.21 (m, 4H), 2.01 - 1.96 (m, 1H), 1.89 - 1.80 (m, 5H), 1.66 - 1.58 (m, 2H), 1.50 - 1.36 (m, 2H).	TFA
8	7.36		7	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.51 - 7.48 (m, 5H), 7.33 - 7.23 (m, 4H), 7.10 (s, 1H), 7.07 - 7.03 (m, 2H), 6.99 (d, J=7.8 Hz, 1H), 6.92 - 6.84 (m, 2H), 6.70 (d, J=9.9 Hz, 1H), 5.41 (t, J=8.5 Hz, 1H), 4.38 - 4.24 (m, 2H), 4.09 (d, J=6.5 Hz, 2H), 4.00 (t, J=7.0 Hz, 2H), 3.51 - 3.45 (m, 2H), 3.25 (d, J=6.4 Hz, 2H), 3.16 - 3.08 (m, 2H), 2.96 (t, J=12.7 Hz, 2H), 2.64 - 2.56 (m, 2H), 2.06 - 1.95 (m, 1H), 1.86 - 1.78 (m, 6H), 1.65 - 1.59 (m, 2H), 1.49 - 1.24 (m, 2H), 1.08 (t, J=8.2 Hz, 3H).	TFA
9	7.15		7	(DMSO-d ₆); δ 10.49 (d, J=10.4 Hz, 2H), 9.52 - 9.52 (m, 1H), 8.59 - 8.58 (m, 2H), 8.17 (d, 9.9 Hz, 1H), 7.48 (s, 6H), 7.25 (dd, J=7.9, 7.9 Hz, 1H), 7.15 (d, J=8.2 Hz, 1H), 7.06 - 7.02 (m, 2H), 7.00 - 6.94 (m, 4H), 6.86 (dd, J=2.2, 8.0 Hz, 1H), 6.59 (dd, J=1.6, 9.9 Hz, 1H), 6.17 - 6.17 (m, 1H), 5.30 (d, J=9.5 Hz, 1H), 4.26 (d, J=4.9 Hz, 2H), 4.01 (d, J=6.4 Hz, 2H), 3.91 (dd, J=6.2, 6.2 Hz, 2H), 3.37 - 3.32 (m, 2H), 3.10 - 2.86 (m, 6H), 1.90 - 1.84 (m, 1H), 1.79 - 1.66 (m, 6H), 1.47 - 1.33 (m, 4H).	TFA


10	7.36		7	(DMSO-d ₆ , 90 °C); δ 8.15 (d, J=9.9 Hz, 1H), 7.50 – 7.39 (m, 5H), 7.20 (dd, J=7.8, 7.8 Hz, 1H), 7.12 (dd, J=8.2, 8.2 Hz, 3H), 6.98 (d, J=8.2 Hz, 1H), 6.80 – 6.76 (m, 1H), 6.54 (d, J=9.8 Hz, 1H), 5.31 (dd, J=4.4, 8.3 Hz, 1H), 4.20 (s, 2H), 3.95 (dd, J=6.3, 6.3 Hz, 4H), 3.58 – 3.19 (m, 5H), 3.15 – 3.07 (m, 2H), 3.02 (dd, J=7.7, 7.7 Hz, 2H), 2.88 (s, 1H), 2.13 (dd, J=10.9, 10.9 Hz, 1H), 1.76 – 1.68 (m, 6H), 1.58 (d, J=9.3 Hz, 2H), 1.52 – 1.33 (m, 6H), 1.23 (d, J=13.1 Hz, 1H), 1.14 – 0.98 (m, 4H).	TFA
11	7.35		7	(DMSO-d ₆); δ 10.52 (s, 1H), 10.48 (s, 1H), 9.43 (s, 1H), 8.57 (s, 2H), 8.15 (d, J=9.9 Hz, 1H), 7.47 (s, 4H), 7.24 (dd, J=7.4, 7.4 Hz, 2H), 7.19 – 7.13 (m, 3H), 7.10 (d, J=7.8 Hz, 1H), 6.99 (d, J=8.2 Hz, 1H), 6.91 – 6.84 (m, 3H), 6.61 – 6.56 (m, 2H), 6.54 (s, 1H), 6.17 (d, J=3.3 Hz, 1H), 5.30 (d, J=9.4 Hz, 1H), 4.30 – 4.25 (m, 2H), 4.01 (d, J=5.6 Hz, 1H), 3.91 (dd, J=6.2, 6.2 Hz, 2H), 3.16 – 2.81 (m, 9H), 2.61 – 2.53 (m, 2H), 1.83 (s, 1H), 1.74 – 1.64 (m, 6H), 1.47 – 1.30 (m, 4H), 1.13 (dd, J=7.5, 7.5 Hz, 3H).	TFA

Scheme 2



5 Procedure 1

Preparation of ethyl 3-methylbenzoylformate




A stirred solution of diethyl oxalate (5 g, 32.2 mmol) in THF (100 mL) was cooled to -78 °C and then a solution of 3-methylphenyl magnesium bromide (0.5M in THF, 75 mL, 37.5 mmol) was added drop-wise over 25 minutes. The reaction mixture was held at -78 °C for 30 minutes and then the mixture warmed to room temperature. The reaction mixture was stirred at room temperature for 3 hours and then cooled to 0 °C. 10% Aqueous hydrochloric acid was added followed by ethyl acetate. The mixture was stirred vigorously for 30 minutes and the organic phase removed. The aqueous phase was extracted with further ethyl acetate and the combined organic extracts washed with brine, and dried over anhydrous magnesium sulfate. The filtrate was evaporated under reduced pressure and the residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 5:1 *iso*-hexane/ethyl acetate) to afford the title compound (3.0 g, 49 %).


¹H NMR (400MHz, CDCl₃); δ 7.81 -7.79 (m, 2H), 7.48 – 7.46 (m, 1H), 7.45 – 7.37 (m, 1H), 4.44 (m, 2H), 2.42 (s, 3H), 1.42 (m, 3H).

Example 2

(*R*)-Quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((*R*)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(4-methoxyphenyl)acetate (Compound 12)




Step 1; 2-(3-(4-(1,3-Dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-(4-methoxyphenyl)acetic acid



The title compound was prepared as described in Example 1 Steps 1 to 4 with ethyl 4-methoxybenzoylformate replacing ethyl benzoylformate in Step 1.

¹H NMR (400 MHz, DMSO-d₆); δ 7.34 (d, J=8.9 Hz, 2H), 7.16 (dd, J=8.2, 8.2 Hz, 1H), 6.98 (d, J=7.2 Hz, 2H), 6.82 (d, J=8.8 Hz, 2H), 6.80 - 6.75 (m, 1H), 4.78 (dd, J=4.7, 4.7 Hz, 1H), 3.87 - 3.72 (m, 9H), 1.75 - 1.57 (m, 4H), 1.51 - 1.44 (m, 2H).

Step 2; (*R*)-Quinuclidin-3-yl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-(4-methoxyphenyl)acetate



A solution of 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-(4-methoxyphenyl)acetic acid (0.200 g, 0.50 mmol) in THF (20 mL) was added with carbonyldiimidazole (0.105 g, 0.65 mmol) and the reaction mixture heated under reflux for

2 hours. The reaction mixture was allowed to cool to room temperature and (*R*)-quinuclidinol (0.215 g, 1.69 mmol) added. The reaction mixture was heated under reflux for 72 hours. The reaction mixture was diluted with ethyl acetate and washed sequentially with 10% aqueous sodium carbonate and brine, and dried over anhydrous magnesium sulfate. The filtrate was evaporated under reduced pressure and the crude material used in the next step with no purification.


Step 3; (Compound 12)

The title compound (12) was prepared as described in Example 1 Step 9.


N	R _t (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
12	6.96	7	(MeOD); δ 8.53 (s, 2H), 8.38 (d, J=9.9 Hz, 1H), 7.35 - 7.25 (m, 4H), 7.06 - 6.99 (m, 3H), 6.97 - 6.89 (m, 3H), 6.72 - 6.68 (m, 1H), 5.40 (t, J=9.7 Hz, 1H), 5.13 - 5.11 (m, 1H), 4.02 - 3.96 (m, 2H), 3.81 (s, 3H), 3.52 - 3.45 (m, 1H), 3.23 (d, J=6.4 Hz, 2H), 3.10 (t, J=9.7 Hz, 2H), 3.06 - 2.99 (m, 3H), 2.88 - 2.73 (m, 2H), 2.19 (d, J=1.6 Hz, 1H), 1.90 - 1.77 (m, 6H), 1.65 - 1.52 (m, 4H).	Formate

Example 3

(*R*)-Quinuclidin-3-yl 2-cyclopentyl-2-hydroxy-2-(3-((5-(((*R*)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)-phenyl)acetate
(Compound 12A)




Step 1; Ethyl 2-(3-(benzyloxy)phenyl)-2-oxoacetate



The title compound was prepared as described in Procedure 1.

^1H NMR (400 MHz, DMSO- d_6); δ 7.57 - 7.52 (m, 3H), 7.49 - 7.39 (m, 5H), 7.37 - 7.32 (m, 1H), 5.21 (s, 2H), 4.42 (q, $J=7.1$ Hz, 2H), 1.33 (dd, $J=7.2$, 7.2 Hz, 3H).


Step 2; 2-(3-(4-(1,3-Dioxolan-2-yl)butoxy)phenyl)-2-cyclopentyl-2-hydroxyacetic acid



The title compound was prepared as described in Example 1 Steps 1 to 4 with ethyl 2-(3-(benzyloxy)phenyl)-2-oxoacetate and cyclopentyl magnesium bromide replacing ethyl benzoylformate and 3-benzyloxyphenyl magnesium bromide respectively in Step 1.

^1H NMR (400 MHz, DMSO- d_6); δ 7.34 (d, $J=8.9$ Hz, 2H), 7.16 (dd, $J=8.2$, 8.2 Hz, 1H), 6.98 (d, $J=7.2$ Hz, 2H), 6.82 (d, $J=8.8$ Hz, 2H), 6.80 - 6.75 (m, 1H), 4.78 (dd, $J=4.7$, 4.7 Hz, 1H), 3.87 - 3.72 (m, 9H), 1.75 - 1.57 (m, 4H), 1.51 - 1.44 (m, 2H).


Step 3; (R)-Quinuclidin-3-yl 2-cyclopentyl-2-hydroxy-2-(3-((5-((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate (Compound 12A)



The title compound was prepared as described in Example 2 Step 2 and 3 to afford the title compound (12A).

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
12A	7.05	6	(MeOD) δ 8.39 (d, J=9.9 Hz, 1H), 7.32 - 7.21 (m, 4H), 7.05 (d, J=8.2 Hz, 1H), 6.88 - 6.84 (m, 1H), 6.70 (d, J=9.9 Hz, 1H), 5.42 (t, J=6.7 Hz, 1H), 5.14 (d, J=10.0 Hz, 1H), 4.04 (t, J=5.0 Hz, 2H), 3.78 - 3.66 (m, 1H), 3.37 - 3.20 (m, 6H), 3.18 - 3.10 (m, 2H), 3.09 - 2.97 (m, 1H), 2.37 - 2.30 (m, 1H), 2.08 - 1.96 (m, 2H), 1.94 - 1.80 (m, 6H), 1.72 - 1.62 (m, 8H), 1.58 - 1.47 (m, 1H), 1.43 - 1.33 (m, 2H)	TFA


**General scheme for the preparation of enantiopure intermediate 1 and 2 –
scheme 3**




Procedure 2

Preparation of (*S*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid and (*R*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid

5




Step 1; (3-Bromophenoxy)(*tert*-butyl)dimethylsilane



A solution of 3-bromophenol (100.00 g, 580 mmol) and imidazole (78.70 g, 1.16 mmol) in DCM (1.2 L) was added drop-wise to a cooled (ice/NaCl bath) solution of *tert*-butyldimethylsilyl chloride (95.83 g, 630 mmol) in DCM (100 mL). The reaction mixture was stirred at room temperature for 16 hours. The reaction mixture was washed with water, brine and the organic phase was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 5:1 *iso*-hexane/ethyl acetate) to afford the title product (163.7 g, 99%).

^1H NMR (400 MHz, CDCl_3) δ 7.12 – 7.10 (m, 2H), 7.03 (ddd, $J = 2.2, 1.4, 0.8$ Hz, 1H), 6.81 – 6.76 (m, 1H), 1.00 (s, 9H), 0.22 (s, 6H).

Step 2; (*R*)-Quinuclidin-3-yl 2-oxo-2-phenylacetate




Benzoylformic acid (50.00 g, 0.33 mol) was dissolved in chloroform (400 mL) and oxalyl chloride (40.02 mL, 0.47 mol) was added drop-wise at room temperature. The reaction mixture stirred for 1 hour and the solvent evaporated under reduced pressure. The residue was dissolved in chloroform (305 mL) and cooled to 0 °C. A solution of 3-(*R*)-

quinuclidinol (50.83 g, 0.40 mmol) in chloroform (200 mL) was added drop-wise. The reaction mixture was allowed warm to room temperature and stirred for 1 hour. The reaction mixture was cooled to 0 °C and quenched with 10% aqueous potassium carbonate solution. The organic layer was separated and washed with water (2x), brine (3x) and dried over anhydrous magnesium sulfate. The filtrate was evaporated under reduced pressure to afford the title compound (77.2 g, 89%).

¹H NMR (400 MHz, DMSO-d₆) δ 8.01 – 7.95 (m, 2H), 7.84 – 7.77 (m, 1H), 7.68 – 7.61 (m, 2H), 5.15 – 5.05 (m, 1H), 3.26 (dd, J = 14.7, 8.2 Hz, 1H), 2.79 – 2.58 (m, 5H), 2.13 – 2.04 (m, 1H), 1.72 – 1.53 (m, 3H), 1.42 – 1.33 (m, 1H).

Step 3; (*R*)-Quinuclidin-3-yl 2-(3-((*tert*-butyldimethylsilyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



Magnesium turnings (8.58 g, 0.36 mol, 1.5 eq) and iodine (catalytic) were suspended in anhydrous THF (460 mL) under argon and a solution of (3-bromophenoxy)(*tert*-butyl)dimethylsilane (88.01g, 0.306 mol, 1.2 eq) in anhydrous THF (175 mL) added drop-wise. The reaction mixture was stirred at reflux for 2 hours. The reaction mixture was cooled to room temperature and then added drop-wise *via* a syringe pump to a solution of (*R*)-quinuclidin-3-yl 2-oxo-2-phenylacetate (66.2 g, 0.26 mol, 1 eq) in anhydrous THF (660 mL) at -75 °C over 2 hours. The reaction mixture was allowed to warm to room temperature and then stirred overnight. The reaction mixture was cooled to 0 °C and quenched with saturated aqueous ammonium chloride (500 mL). The reaction mixture was extracted with ethyl acetate and the organic phase was washed with brine, dried over anhydrous magnesium sulfate and the filtrate evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% DCM to 20:1 DCM / methanol) to afford the title product (66.2 g, 78.5%).

^1H NMR (400 MHz, DMSO- d_6) δ 7.37 – 7.28 (m, 5H), 7.25 – 7.20 (m, 1H), 7.01 – 6.95 (m, 1H), 6.83 – 6.76 (m, 2H), 6.60 (s, 1H), 4.84 – 4.78 (m, 1H), 3.12 – 3.06 (m, 1H), 2.65 – 2.56 (m, 3H), 2.48 – 2.37 (m, 2H), 1.90 – 1.80 (m, 1H), 1.59 – 1.42 (m, 2H), 1.36 – 1.30 (m, 1H), 1.21 – 1.12 (m, 1H), 0.92 – 0.89 (m, 9H), 0.12 (d, J = 1.0 Hz, 6H).

5 **Step 4; (*R*)-Quinuclidin-3-yl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate**




(*R*)-Quinuclidin-3-yl 2-(3-((*tert*-butyldimethylsilyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (93.72 g, 200 mmol) was dissolved in THF (937 mL) and TBAF*3H₂O (69.54 g, 11.5 mmol) was added and the reaction mixture was stirred at room temperature for 1 hour. THF was evaporated and the resulting crude was dissolved in ethyl acetate and washed with 0.5M hydrochloric acid (3 x 350 mL). The combined aqueous layers were neutralized with sodium hydrogen carbonate (to pH 8). The resulting precipitate was filtered to give 26.05 g HPLC 100% (51.9:48.1). Aqueous layer (filtrate) was washed with ethyl acetate (3 x 1 L) and the combined organic layers were concentrated to give desired product (38.64 g, yield 55.6, 75:25 ratio).

Method 10. Chiral-HPLC

First diastereoisomer rt 12.18 min

Second diastereoisomer rt 12.87 min

20 **Step 5; (*S,R*)- and (*R,R*)-Quinuclidin-3-yl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate**



(*R*)-Quinuclidin-3-yl 2-(3-((*tert*-butyldimethylsilyl)oxy)phenyl)-2-hydroxy-2-

phenylacetate (26 g, 1:1 ratio) was purified via flash column chromatography (25 g SiO₂/1 g crude, eluent: DCM; Acetone; TEA; 7:2.5:0.5) leading to

i) **(S)-(R)-quinuclidin-3-yl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate** (7.00 g, de=99.2%).


5 Method 9 HPLC-AB: t_R 12.1 min.

ii) **(R)-(R)-quinuclidin-3-yl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate** (1.7 g, de=95%).

Method 9 HPLC-AB: t_R 12.5 min.

The absolute configuration of the unknown stereogenic center was assigned by
10 single crystal X-ray diffraction.

Step 6; (S)-2-Hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid




A solution of **(S)-(R)-quinuclidin-3-yl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate** (28.24 g, 79.91 mmol) in methanol (282.40 mL, 10 vol at 0°C, was added
15 drop-wise with . The reaction mixture was heated to reflux and stirred for 30 min. The solvent was evaporated and the reaction mixture adjusted to pH-8 with 1M hydrochloric acid and washed with ethyl acetate (2 x 100 mL). The aqueous layer was acidified with 1M hydrochloric acid to pH 4-5 and extracted with ethyl acetate (3 x 250 mL). The combined organic extracts were dried (magnesium sulfate), concentrated and then dried under vacuum
20 for 3 day at 35°C to give the title compound (17.58 g, 90% yield).

¹H NMR (400 MHz, DMSO-d₆) δ 13.12 (bs, 1H), 9.33 (bs, 1H), 7.41 – 7.25 (m, 5H), 7.14 – 7.07 (m, 1H), 6.84 – 6.78 (m, 2H), 6.69 – 6.64 (m, 1H).

Method 10 Chiral HPLC: t_R 12.95 min,


General scheme for the preparation of enantiopure intermediate 3 and 4 – scheme 4



5 Procedure 3

Preparation of (*S*)-tert-butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxymethyl)piperidine-1-carboxylate and (*R*)-tert-butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxymethyl)piperidine-1-carboxylate

Step 1; 2-(3-(Benzyloxy)phenyl)-2-hydroxy-2-phenylacetic acid



10


A stirred solution of ethyl 2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetate (as described in Example 1, Step 1) (20.3 g, 56.1 mmol) in THF (100 mL) was added with aqueous 2M sodium hydroxide (20 mL). The reaction mixture was stirred at room temperature for 2 hours and then the solvent removed under reduced pressure. The pH was adjusted to 1 with aqueous 1M hydrochloric acid and then extracted with ethyl acetate (x2). The combined organic extracts were washed with brine, dried over anhydrous magnesium sulfate and the filtrate evaporated under reduced pressure to afford the title product (18.4 g, 98%).

15

¹H NMR (400 MHz, DMSO-d₆); δ 7.45 - 7.23 (m, 11H), 7.02 (dd, J=2.1, 2.1 Hz,

1H), 6.98 - 6.93 (m, 2H), 5.05 (s, 2H), 3.34 (s, 1H).


Step 2; *tert*-Butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate



5 A stirred solution of 2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetic acid (10.1 g, 30.2 mmol) in DMF (100 mL) was added with potassium carbonate (5.0 g, 36.2 mmol). After 10 minutes *tert*-butyl 4-(tosyloxymethyl)piperidine-1-carboxylate (as described in Example 1, Step 5) (11.1 g, 30.2 mmol) was added and the resulting mixture heated at 80 °C for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the
10 filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 4:1 *iso*-hexane/ethyl acetate) to afford the title compound (9.51 g, 59 %).

¹H NMR (400 MHz, CDCl₃); δ 7.42 - 7.24 (m, 11H), 7.08 (dd, J=2.1, 2.1 Hz, 1H),
15 6.99 (d, J=7.8 Hz, 1H), 6.93 (dd, J=2.4, 8.4 Hz, 1H), 5.02 (s, 2H), 4.21 (s, 1H), 4.12 - 4.03 (m, 4H), 2.63 - 2.57 (m, 2H), 1.79 - 1.69 (m, 1H), 1.50 - 1.40 (m, 11H), 1.09 - 0.89 (m, 2H).

Step 3; (*S*)-*tert*-Butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate and (*R*)-*tert*-butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate
20



tert-Butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate was separated by chiral SFC.

SFC method:

Instrument: Waters Prep 100


Column: YMC Amylose-C

Eluant: 30 / 70 *iso*-propyl alcohol / CO₂ at a flow rate of 100 ml/min 120 bar and

5 40°C.


General scheme for the preparation of enantiopure compounds of example 4

scheme 5




Example 4

(*S*)-(1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((*R*)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate (Compound 13)




Step 1; (*S*)-4-(1,3-Dioxolan-2-yl)butyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate



A stirred solution of (*S*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid (1.90 g, 7.78 mmol) in DMF (30 mL) was added with potassium carbonate (3.22 g, 23.3 mmol). This mixture was added with 2-(4-chlorobutyl)-1,3-dioxolane (2.89 mL, 19.45 mmol) and the reaction mixture heated at 85 °C for 72 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 1:1 *iso*-hexane/ethyl acetate) to afford the title compound (2.8 g, 72 %).

¹H NMR (400 MHz, DMSO-*d*₆); δ 7.33 (d, *J*=4.1 Hz, 6H), 6.89 - 6.85 (m, 3H), 6.57 (s, 1H), 4.78 (dd, *J*=4.8, 4.8 Hz, 1H), 4.69 (dd, *J*=4.8, 4.8 Hz, 1H), 4.14 (dd, *J*=6.4, 6.4 Hz, 2H), 3.93 - 3.71 (m, 10H), 1.74 - 1.44 (m, 10H), 1.32 - 1.16 (m, 2H).


Step 2; (*S*)-2-(3-(4-(1,3-Dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetic acid



A stirred solution of (*S*)-4-(1,3-dioxolan-2-yl)butyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate (2.8 g, 5.59 mmol) in THF (25 mL) was added with aqueous with 2M sodium hydroxide (25 mL). The reaction mixture was stirred at room temperature for 1 hour. The organic solvent was evaporated under reduced pressure. The aqueous residue was washed with ether and the pH of the aqueous phase adjusted to 5 with aqueous 1M hydrochloric acid. The mixture was extracted with ethyl acetate and the organic phase washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 1:1 *iso*-hexane/ethyl acetate) to afford the title compound (1.46 g, 70 %).

¹H NMR (400 MHz, DMSO-d₆); δ 7.45 - 7.25 (m, 6H), 7.01 - 6.96 (m, 2H), 6.91 - 6.88 (m, 1H), 6.40 (br s, 1H), 4.83 (dd, J=4.8, 4.8 Hz, 1H), 3.98 - 3.78 (m, 6H), 1.81 - 1.62 (m, 4H), 1.57 - 1.49 (m, 2H).

Step 3; (*S*)-(1-Benzylpiperidin-4-yl)methyl 2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate



A stirred solution of (*S*)-2-(3-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetic acid (0.119 g, 0.32 mmol) in DMF (2 mL) was added with carbonyl diimidazole (0.078 g, 0.48 mmol) and the mixture heated at 50 °C for 30 minutes. A solution of N-benzyl piperidine-4-methanol (0.098 g, 0.32 mmol) in DMF (0.5 mL) was added and the mixture heated at 50 °C for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by reverse phase column chromatography to afford the title compound (0.075 g, 38 %).

¹H NMR (400MHz, CDCl₃); δ 7.42 – 7.38 (m, 2H), 7.35 – 7.19 (m, 9H), 7.00 –

6.94 (m, 2H), 6.83 (ddd, $J = 8.2, 2.5, 0.9$ Hz, 1H), 5.30 (s, 2H), 4.88 – 4.83 (m, 1H), 4.22 (s, 1H), 4.15 – 4.06 (m, 2H), 4.01 – 3.80 (m, 6H), 3.45 (s, 2H), 2.81 (d, $J = 11.3$ Hz, 2H), 1.92 – 1.66 (m, 6H), 1.70 – 1.46 (m, 3H), 1.28 – 1.13 (m, 2H).

Step 4; (Compound 13)

5 The title compound (13) was prepared as described in Example 1 Step 9










	R_t (min)	Method	NMR data (400 MHz)	Salt 2 eq unless stated
13	2.32	8	(DMSO-d ₆ , 90°C); δ 8.15 (d, $J=9.2$ Hz, 1H), 7.48 (m, 4H), 7.35 - 7.31 (m, 5H), 7.24 (dd, $J=7.9, 7.9$ Hz, 1H), 7.15 (d, $J=8.2$ Hz, 1H), 6.99 (d, $J=8.2$ Hz, 1H), 6.89 - 6.85 (m, 3H), 6.63 - 6.57 (m, 2H), 5.30 (d, $J=9.8$ Hz, 1H), 4.31 - 4.21 (m, 3H), 4.01 (d, $J=6.4$ Hz, 2H), 3.90 (dd, $J=6.2, 6.2$ Hz, 2H), 3.34 (d, $J=11.3$ Hz, 2H), 3.10 - 2.97 (m, 5H), 2.90 (d, $J=12.4$ Hz, 2H), 1.85 (m, 1H), 1.71 - 1.65 (m, 5H), 1.47 - 1.30 (m, 4H).	TFA

The following compounds were prepared as described in Example 4 with the relevant (*R*) or (*S*) stereoisomer of 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid used in Step 1 and the required commercially available amine replacing N-benzyl piperidine-4-methanol

10 in Step 3.

Compound number	Relevant stereoisomer	Amine	Structure
14			
15			
16			
17			
18			
19			
20			
21			
22			
23			

(continued)

24			
25			
26			

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
14	7.19	7	(DMSO-d ₆); δ 10.54 (d, J=15.7 Hz, 2H), 9.44 (s, 1H), 8.61 - 8.60 (m, 2H), 8.20 (d, J=11.6 Hz, 1H), 7.53 (s, 5H), 7.40 - 7.36 (m, 6H), 7.20 (d, J=8.3 Hz, 1H), 7.04 (d, J=8.1 Hz, 1H), 6.95 - 6.90 (m, 3H), 6.69 - 6.62 (m, 2H), 6.21 (s, 1H), 5.37 - 5.34 (m, 1H), 4.37 - 4.26 (m, 3H), 4.06 (d, J=6.3 Hz, 2H), 3.95 (dd, J=6.2, 6.2 Hz, 2H), 3.38 (d, J=10.9 Hz, 2H), 3.16 - 2.90 (m, 5H), 1.93 - 1.88 (m, 1H), 1.83 - 1.71 (m, 6H), 1.53 - 1.34 (m, 4H)	TFA
15	6.94	7	(DMSO-d ₆); δ 10.54 (d, J=15.9 Hz, 2H), 9.66 (s, 1H), 8.60 - 8.60 (m, 2H), 8.21 (d, J=10.4 Hz, 1H), 7.41 (d, J=4.5 Hz, 4H), 7.40 - 7.35 (m, 1H), 7.31 (dd, J=8.0, 8.0 Hz, 1H), 7.20 (d, J=8.1 Hz, 1H), 7.05 - 6.91 (m, 4H), 6.81 (s, 1H), 6.64 (dd, J=1.9, 10.0 Hz, 1H), 6.20 (d, J=2.0 Hz, 1H), 5.35 (d, J=9.6 Hz, 1H), 5.21 - 5.19 (m, 1H), 3.97 (dd, J=6.2, 6.2 Hz, 2H), 3.78 - 3.70 (m, 1H), 3.28 - 2.98 (m, 9H), 2.27 (d, J=2.5 Hz, 1H), 1.95 - 1.87 (m, 2H), 1.79 - 1.61 (m, 6H), 1.53 - 1.44 (m, 2H)	TFA
16	6.95	7	(DMSO-d ₆); δ 10.54 (d, J=12.9 Hz, 2H), 9.74 - 9.71 (m, 1H), 8.61 - 8.60 (m, 2H), 8.21 (d, J=9.0 Hz, 1H), 7.44 - 7.30 (m, 6H), 7.20 (d, J=8.1 Hz, 1H), 7.02 (dd, J=8.1, 14.9 Hz, 2H), 6.95 - 6.91 (m, 2H), 6.81 (s, 1H), 6.64 (dd, J=1.9, 10.0 Hz, 1H), 6.22 (s, 1H), 5.35 (d, J=9.6 Hz, 1H), 5.23 - 5.18 (m, 1H), 3.97 (t, J=6.7 Hz, 2H), 3.74 (td, J=5.2, 17.9 Hz, 1H), 3.27 - 2.98 (m, 9H), 2.27 (d, J=3.0 Hz, 1H), 1.96 - 1.86 (m, 2H), 1.79 - 1.61 (m, 6H), 1.53 - 1.43 (m, 2H)	TFA













(continued)

17	2.14	8	(DMSO-d ₆); δ 10.50 (d, J=18.1 Hz, 2H), 9.59 (s, 1H), 8.56 - 8.56 (m, 2H), 8.15 (d, J=10.7 Hz, 1H), 7.35 (m, 6H), 7.15 (d, J=8.3 Hz, 1H), 6.98 (d, J=8.2 Hz, 1H), 6.94 (d, J=8.2 Hz, 1H), 6.89 - 6.87 (m, 2H), 6.68 (s, 1H), 6.59 (dd, J=2.1, 9.9 Hz, 1H), 6.17 (d, J=3.4 Hz, 1H), 5.31 - 5.28 (m, 1H), 4.47 (dd, J=5.0, 5.0 Hz, 2H), 3.97 - 3.88 (m, 2H), 3.14 - 2.98 (m, 6H), 2.69 (s, 6H), 1.74 - 1.64 (m, 4H), 1.48 - 1.40 (m, 2H).	TFA
18	6.89	7	(DMSO-d ₆); δ 10.49 (d, J=14.1 Hz, 2H), 9.62 (s, 1H), 8.57 - 8.56 (m, 2H), 8.15 (d, J=9.9 Hz, 1H), 7.35 (d, J=4.3 Hz, 4H), 7.33 - 7.23 (m, 2H), 7.15 (d, J=8.2 Hz, 1H), 6.98 (d, J=8.2 Hz, 1H), 6.93 (d, J=8.0 Hz, 1H), 6.90 - 6.87 (m, 2H), 6.67 (s, 1H), 6.59 (dd, J=1.8, 9.9 Hz, 1H), 6.17 - 6.16 (m, 1H), 5.30 (d, J=9.4 Hz, 1H), 4.47 (dd, J=5.1, 5.1 Hz, 2H), 3.91 (dd, J=6.3, 6.3 Hz, 2H), 3.40 (s, 2H), 3.14 - 3.04 (m, 2H), 3.01 - 2.95 (m, 2H), 2.69 (s, 6H), 1.75 - 1.64 (m, 4H), 1.48 - 1.40 (m, 2H).	TFA
19	6.90	7	(DMSO-d ₆); δ 10.54 (d, J=13.9 Hz, 2H), 10.17 (s, 1H), 8.62 - 8.60 (m, 2H), 8.20 (d, J=11.5 Hz, 1H), 7.41-7.28 (m, 6H), 7.20 (d, J=10.2 Hz, 1H), 7.05 - 6.97 (m, 2H), 6.93 (m, 2H), 6.64 (d, J=8.9 Hz, 2H), 6.21 (s, 1H), 5.50 (d, J=16.6 Hz, 1H), 5.36 (d, J=8.9 Hz, 1H), 3.97 (dd, J=6.2, 6.2 Hz, 2H), 3.61 (m, 4H), 3.18 - 3.01 (m, 4H), 2.95 - 2.72 (m, 3H), 2.5-2.3 (m, 1H), 2.1-1.9 (m, 1H), 1.81 - 1.69 (m, 4H), 1.53 - 1.44 (m, 2H).	TFA
20	6.90	7	(DMSO-d ₆); δ 10.49 (d, J=13.2 Hz, 2H), 10.15 (s, 1H), 8.57 (s, 2H), 8.15 (d, J=9.9 Hz, 1H), 7.35 (d, J=3.3 Hz, 4H), 7.33 - 7.23 (m, 2H), 7.15 (d, J=8.2 Hz, 1H), 7.00 - 6.92 (m, 2H), 6.88 (d, J=7.5 Hz, 2H), 6.59 (dd, J=1.8, 9.8 Hz, 2H), 6.17 - 6.17 (m, 1H), 5.44 (d, J=22.5 Hz, 1H), 5.31 (d, J=9.2 Hz, 1H), 3.92 (t, J=5.4 Hz, 2H), 3.59 (m, 2H), 3.41 - 3.40 (m, 0.5H), 3.13 - 2.97 (m, 5H), 2.77 (d, J=63.6 Hz, 4H), 2.25 (m, 0.5H), 2.09 - 2.08 (m, 0.5H), 1.91 (m, 0.5H), 1.74 - 1.64 (m, 4H), 1.48 - 1.40 (m, 2H).	TFA
21	6.89	7	(DMSO-d ₆ , 90°C); δ 8.18 (d, J=9.9 Hz, 1H), 7.42 - 7.24 (m, 6H), 7.15 (d, J=8.2 Hz, 1H), 7.03 - 6.94 (m, 3H), 6.89 (dd, J=2.2, 8.1 Hz, 1H), 6.56 (d, J=9.8 Hz, 1H), 5.35 (dd, J=4.4, 8.4 Hz, 1H), 5.09 (s, 1H), 3.95 (dd, J=6.3, 6.3 Hz, 2H), 3.19 - 3.00 (m, 8H), 2.71 (s, 3H), 2.11 - 2.06 (m, 2H), 1.78 - 1.70 (m, 6H), 1.53 - 1.46 (m, 2H).	TFA

(continued)

22	6.90	7	(DMSO-d ₆ , 90°C); δ 8.19 (d, J=9.9 Hz, 1H), 7.42 - 7.22 (m, 6H), 7.15 (d, J=8.2 Hz, 1H), 7.02 (d, J=8.2 Hz, 1H), 6.96 (d, J=8.0 Hz, 2H), 6.90 - 6.85 (m, 1H), 6.55 (d, J=9.9 Hz, 1H), 5.36 (dd, J=4.5, 8.3 Hz, 1H), 5.07 (d, J=1.9 Hz, 1H), 3.95 (dd, J=6.3, 6.3 Hz, 2H), 3.19 - 3.00 (m, 8H), 2.70 (s, 3H), 2.05 (d, J=8.9 Hz, 2H), 1.86 - 1.84 (m, 2H), 1.79 - 1.67 (m, 4H), 1.53 - 1.45 (m, 2H).	TFA
23	6.96	7	(DMSO-d ₆ , 90°C); δ 8.19 (d, J=9.9 Hz, 1H), 7.40 - 7.22 (m, 6H), 7.15 (d, J=8.2 Hz, 1H), 7.02 (d, J=8.0 Hz, 1H), 6.96 - 6.94 (m, 2H), 6.88 - 6.86 (m, 1H), 6.56 (d, J=9.9 Hz, 1H), 6.26 - 6.26 (m, 1H), 5.35 (dd, J=4.5, 8.4 Hz, 1H), 4.09 (d, J=6.5 Hz, 2H), 3.96 (dd, J=6.4, 6.4 Hz, 2H), 3.31 (d, J=11.0 Hz, 2H), 3.16 - 2.96 (m, 6H), 2.73 (s, 3H), 1.92 - 1.85 (m, 1H), 1.78 - 1.71 (m, 6H), 1.54 - 1.42 (m, 4H).	TFA
24	6.97	7	(DMSO-d ₆); δ 8.30 (s, 2H), 8.19 (d, J=9.6 Hz, 1H), 7.33 - 7.31 (m, 6H), 7.11 (d, J=8.3 Hz, 1H), 6.96 (d, J=8.2 Hz, 1H), 6.90 - 6.85 (m, 3H), 6.53 (d, J=9.8 Hz, 1H), 5.20 (dd, J=4.3, 8.4 Hz, 1H), 3.99 (d, J=6.0 Hz, 2H), 3.90 (dd, J=6.3, 6.3 Hz, 2H), 2.90 - 2.68 (m, 5H), 2.13 (s, 3H), 1.84 - 1.77 (m, 2H), 1.74 - 1.64 (m, 2H), 1.61 - 1.39 (m, 8H), 1.18 - 1.09 (m, 2H).	Formate
25	6.95	7	(DMSO-d ₆ , 90°C); δ 8.18 (d, J=10.5 Hz, 1H), 7.43 - 7.31 (m, 5H), 7.25 (dd, J=8.0, 8.0 Hz, 1H), 7.15 (d, J=8.2 Hz, 1H), 7.03 - 6.94 (m, 3H), 6.87 (dd, J=2.4, 8.3 Hz, 1H), 6.57 (d, J=9.8 Hz, 1H), 5.33 (dd, J=4.5, 8.5 Hz, 1H), 5.16 (m, 1H), 4.00 - 3.93 (m, 2H), 3.17 - 3.01 (m, 10H), 2.74 (s, 3H), 1.78 - 1.69 (m, 6H), 1.53 - 1.45 (m, 2H).	TFA
26	6.87	7	(DMSO-d ₆ , 90°C); δ 8.19 (d, J=9.9 Hz, 1H), 7.42 (dd, J=1.5, 8.3 Hz, 2H), 7.38 - 7.24 (m, 4H), 7.15 (d, J=8.2 Hz, 1H), 7.03 - 6.95 (m, 3H), 6.88 (dd, J=2.3, 8.0 Hz, 1H), 6.56 (d, J=9.9 Hz, 1H), 5.37 - 5.31 (m, 2H), 4.43 - 4.37 (m, 2H), 3.99 - 3.88 (m, 4H), 3.22 - 3.11 (m, 2H), 3.07 - 3.01 (m, 2H), 2.80 (s, 3H), 1.78 - 1.69 (m, 4H), 1.53 - 1.44 (m, 2H).	TFA



The following compounds were prepared as described in Example 4 with the relevant (*R*) or (*S*) stereoisomer of 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid used in Step 1 and the appropriate alkylating agent replacing 2-(4-chlorobutyl)-1,3-dioxolane in Step 1.

N	Relevant stereoisomer	Alkylating Agent	Structure
27			
28			
29			
30			


N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
27	7.25	7	(DMSO-d ₆); δ 10.54 (s, 1H), 10.49 (s, 1H), 9.56 (s, 1H), 8.61 - 8.61 (m, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.47 (s, 5H), 7.35 - 7.31 (m, 6H), 7.15 (d, J=8.0 Hz, 1H), 7.03 - 6.97 (m, 1H), 6.90 - 6.83 (m, 3H), 6.62 (s, 1H), 6.58 (d, J=9.9 Hz, 1H), 6.18 - 6.18 (m, 1H), 5.31 (d, J=9.2 Hz, 1H), 4.25 - 4.21 (m, 2H), 4.01 (d, J=6.4 Hz, 2H), 3.90 (dd, J=6.3, 6.3 Hz, 2H), 3.08 (m, 4H), 2.97 - 2.88 (m, 4H), 1.75 - 1.61 (m, 7H), 1.36 (dd, J=11.5, 11.5 Hz, 6H).	TFA
28	7.34	77	(DMSO-d ₆); δ 10.50 (m, 2H), 9.51 (s, 1H), 8.58 (s, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.47 (s, 5H), 7.33 (s, 5H), 7.23 (dd, J=8.0, 8.0 Hz, 1H), 7.15 (d, J=8.2 Hz, 1H), 6.99 (d, J=8.2 Hz, 1H), 6.90 - 6.85 (m, 3H), 6.64 - 6.57 (m, 2H), 6.18 (d, J=2.5 Hz, 1H), 5.31 (d, J=9.0 Hz, 1H), 4.27 - 4.23 (m, 2H), 4.01 (d, J=6.3 Hz, 2H), 3.90 (dd, J=6.4, 6.4 Hz, 2H), 2.98 - 2.98 (m, 10H), 1.68 - 1.64 (m, 5H), 1.38 - 1.31 (m, 8H).	TFA
29	7.44	77	(DMSO-d ₆); δ 10.51 (s, 2H), 9.56 (s, 1H), 8.61 (s, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.48 (s, 5H), 7.33 - 7.22 (m, 6H), 7.15 (d, J=8.2 Hz, 1H), 6.99 (d, J=8.2 Hz, 1H), 6.90 - 6.83 (m, 3H), 6.59 (t, J=11.1 Hz, 2H), 6.18 (s, 1H), 5.31 (d, J=8.8 Hz, 1H), 4.28 - 4.21 (m, 2H), 4.01 (d, J=6.3 Hz, 2H), 3.90 (dd, J=6.1, 6.1 Hz, 2H), 3.14 - 2.85 (m, 7H), 1.75 - 1.62 (m, 7H), 1.40 - 1.34 (m, 7H).	TFA

(continued)

30	7.43	77	(DMSO-d ₆); δ 10.53 (d, J=21.3 Hz, 2H), 9.79 (s, 1H), 8.76 (s, 1H), 8.61 (s, 1H), 8.19 (d, J=12.0 Hz, 1H), 7.49 - 7.46 (m, 5H), 7.35 - 7.31 (m, 6H), 7.15 (d, J=8.3 Hz, 1H), 7.00 (d, J=8.2 Hz, 1H), 6.87 (d, J=8.4 Hz, 3H), 6.62 (s, 1H), 6.57 (d, J=9.8 Hz, 1H), 6.19 (s, 1H), 5.34 (d, J=8.0 Hz, 1H), 4.29 - 4.20 (m, 2H), 4.00 (d, J=6.4 Hz, 2H), 3.89 (dd, J=6.4, 6.4 Hz, 2H), 3.11 - 2.83 (m, 7H), 1.83 - 1.64 (m, 8H), 1.43 - 1.23 (m, 10H).	TFA
----	------	----	--	-----


Procedure 4**Preparation of (*R*)-4-(((*tert*-butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-****2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid**

5

Step 1; (*R*)-5-(2-amino-1-(((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one

A solution of (*R*)-5-(2-azido-1-(((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1*H*)-one (3.83 g, 8.5 mmol) in THF / water (30 mL / 1 mL) was added with polymer supported triphenylphosphine (7.3 g, 1.4 mmol/g loading, 10.2 mmol). The reaction mixture was heated at reflux with no stirring for 18 hours. The suspension was filtered, the filter cake was washed with ethyl acetate and the filtrate was evaporated under reduced pressure to afford the title compound. The material was used directly in the next step with no purification.


Step 2; (*R*)-Methyl 4-(((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-

((*tert*-butyldimethylsilyl)oxy)ethyl)amino)methyl)benzoate

A stirred mixture of (*R*)-5-(2-amino-1-((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (2.9 g, 6.83 mmol) and methyl 4-formylbenzoate (1.23 g, 7.49 mmol) in DCM (30 mL) was added with anhydrous magnesium sulfate and the mixture stirred at room temperature for 18 hours. The suspension was filtered through a plug of anhydrous magnesium sulfate, the filter cake was washed with further DCM and the filtrate was evaporated under reduced pressure. The residue was dissolved in methanol (30 mL) and the mixture cooled to 0°C. Sodium borohydride (0.517 g, 13.7 mmol) was added portion wise and the reaction mixture stirred at 0 °C for 30 minutes. The coolant was removed and the mixture stirred at room temperature for a further 2 hours. The mixture was quenched with 10% aqueous potassium carbonate and extracted with DCM (x2). The combined DCM extracts were washed with brine, dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% DCM to 25:1 DCM/methanol) to afford the title compound (3.26 g, 83 %).

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.51 (s, 1H), 8.20 (d, *J*=9.9 Hz, 1H), 7.81 (d, *J*=8.3 Hz, 2H), 7.52 (d, *J*=7.2 Hz, 2H), 7.36 - 7.31 (m, 4H), 7.29 - 7.25 (m, 1H), 7.15 (d, *J*=8.3 Hz, 1H), 7.07 (d, *J*=8.4 Hz, 1H), 6.46 (d, *J*=9.9 Hz, 1H), 5.23 (s, 2H), 5.13 (dd, *J*=4.7, 7.2 Hz, 1H), 3.80 - 3.78 (m, 3H), 3.73 (s, 2H), 2.72 (dd, *J*=7.8, 12.0 Hz, 1H), 2.63 - 2.55 (m, 1H), 0.77 (s, 9H), 0.00 (s, 3H), -0.21 (s, 3H).


Step 3; (R)-Methyl 4-(((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)methyl)benzoate



A stirred solution of (R)-methyl 4-(((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)amino)methyl)benzoate (3.25 g, 5.67 mmol) in DCM (25 mL) was added with a solution of di-*tert*-butyl dicarbonate (1.49 g, 6.83 mmol) in DCM (5 mL). The reaction mixture was stirred at room temperature for 18 hours. The solvent was evaporated under reduced pressure and the residue purified by flash column chromatography (eluent – 100% *iso*-hexane to 3:2 *iso*-hexane/ethyl acetate) to afford the title compound (2.84 g, 75 %).

LCMS (Method 11); Rt 4.33 min; M+1 673.4

Step 4; (R)-4-(((2-(8-(Benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)methyl)benzoic acid




A stirred solution of (R)-methyl 4-(((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)-amino)methyl)benzoate (2.84 g, 4.22 mmol) in methanol (10 mL) was added with aqueous 2M sodium hydroxide (10 mL). The reaction mixture was then stirred at room temperature for 18 hours. The solvent was evaporated under reduced pressure and the residue partitioned between DCM and 10% aqueous potassium hydrogen sulfate. The organic phase was removed and the aqueous phase extracted with further DCM. The combined DCM extracts were washed with brine, dried over anhydrous magnesium sulfate and the filtrate was evaporated under

reduced pressure to afford the title compound (2.58 g, 93%).

LCMS (Method 11); Rt 4.05 min; M+1 659

Step 5; (R)-4-(((tert-Butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid




5

10% Palladium on carbon (2 g) was added to a stirred solution of (R)-4-(((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)methyl)benzoic acid (2.06 g, 3.12 mmol) in ethanol (25 mL) under an inert nitrogen atmosphere. 1-Methyl-1,4-cyclohexadiene (1.75 mL, 15.6 mmol) was added and the reaction mixture was carefully heated to reflux [Care – vigorous evolution of gas]. The reaction mixture was heated under reflux for 1 hour. The suspension was filtered, the filter cake washed with further ethanol and the filtrate was evaporated under reduced pressure to afford the title compound (1.66 g, 94%).

LCMS (Method 11); Rt 3.61 min; M+1 569.5.

Also prepared in the same fashion were;


(R)-4-(((tert-Butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoic acid



Starting from methyl 4-formyl-3-methoxybenzoate

LCMS Method 11; Rt 3.75 min; ES⁺ 599.4

(R)-4-(((tert-Butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoic acid




Starting from methyl 4-formyl-2-methoxybenzoate

20

LCMS Method 11; Rt 3.61; ES⁺ 599.4


(R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-fluorobenzoic acid



5 Starting from methyl 2-fluoro-4-formylbenzoate

LCMS Method 11; Rt 3.64; ES⁺ 587.3


(R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-fluorobenzoic acid



10 Starting from methyl 3-fluoro-4-formylbenzoate

LCMS Method 11; Rt 3.70; ES⁺ 587.4


(R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methylbenzoic acid



15 Starting from methyl 4-formyl-3-methylbenzoate

LCMS Method 11; Rt 3.70; ES⁺ 587.4


(R)-2-(4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)phenyl)acetic acid



20 Starting from methyl 2-(4-formylphenyl)acetate.

LCMS Method 12; Rt 2.93; ES⁺ 583.4

(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethyl)benzoic acid



Starting from methyl 3-trifluoromethyl-4-formylbenzoate

5 LCMS Method 11; Rt 3.79; ES⁺ 637.3


Synthesis of methyl 3-trifluoromethyl-4-formylbenzoate

A solution of 4-methyl-3-trifluoromethylbenzoic acid (1.71 g, 8.37 mmol) in DMF (20 mL) was added with potassium carbonate (1.39 g, 10.1 mmol) and the reaction mixture was stirred at room temperature for five minutes. Iodomethane (0.78 mL, 12.5 mmol) was added and the reaction mixture stirred at room temperature for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure to afford methyl 4-methyl-3-trifluoromethylbenzoate. The material was dissolved in carbon tetrachloride (10 mL) and treated sequentially with N-bromosuccinimide (2.74 g, 15.4 mmol) and benzoyl peroxide (catalytic) and heated at 80 °C for 18 hours. The mixture allowed to cool and diluted with water. The mixture was poured through a hydrophobic frit and the solvent evaporated at reduced pressure to afford methyl 4-(dibromomethyl)-3-trifluoromethylbenzoate. The material was dissolved in acetone / water (25 mL / 5 mL) and silver nitrate (2.38 g, 14.0 mmol) added. The reaction mixture was stirred at room temperature for 72 hours. The suspension was filtered through a pad of celite and the filtrate diluted with ethyl acetate. The solution was washed with water and brine. The organic phase was dried over anhydrous magnesium sulfate, filtered and evaporated. The residue was dissolved acetone / water (25 mL / 5 mL) and silver nitrate (2.38 g, 14.0 mmol) added. The reaction mixture was stirred at room temperature for 24 hours. The suspension was filtered through a pad of celite and the filtrate diluted with ethyl acetate. The solution was washed with water and brine. The organic phase was dried over

anhydrous magnesium sulfate, filtered and evaporated. The residue was purified by flash column chromatography (eluent – 100% i-hexane to 8:1 i-hexane/ethyl acetate) to afford the title compound (0.537 g, 33%).

¹H NMR (400 MHz, CDCl₃); δ 10.46 - 10.45 (m, 1H), 8.46 (s, 1H), 8.35 (d, J=8.2 Hz, 1H), 8.20 (d, J=8.2 Hz, 1H), 4.00 (s, 3H).


(R)-4-(((tert-Butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2,3-difluorobenzoic acid



Starting from methyl 2,3-difluoro-4-formylbenzoate (prepared as described for methyl 3-trifluoromethyl-4-formylbenzoate).

LCMS (10cm_Formic_Aq); Rt 3.66; ES⁺ 605 / 607.

(R)-4-(((tert-Butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-ethoxybenzoic acid



Starting from ethyl 3-ethoxy-4-formylbenzoate.


LCMS Method 11; Rt 3.76; ES⁺ 613.7.

Synthesis of ethyl 3-ethoxy-4-formylbenzoate.

A solution of 4-formyl-3-hydroxybenzoic acid (1.0 g, 6.02 mmol) in DMF (20 mL) was added with potassium carbonate (2.49 g, 18.1 mmol) and ethyl iodide (2.82 g, 18.1 mmol). The reaction mixture was stirred at room temperature for 72 hours. The reaction mixture was concentrated at reduced pressure and the residue partitioned between ethyl acetate and water. The organic extract was washed with brine (x2), poured through a hydrophobic frit and the solvent evaporated at reduced pressure. The residue was purified by flash column chromatography (eluent – 100% i-hexane to 4:1 i-hexane/ethyl acetate) to afford the title compound (1.41 g, 100%).

¹H NMR (400 MHz, CDCl₃); δ 10.55 (s, 1H), 7.87 (d, J=8.3 Hz, 1H), 7.66 (d, J=8.3 Hz, 2H), 4.40 (q, J=7.2 Hz, 2H), 4.23 (q, J=7.0 Hz, 2H), 1.51 (dd, J=6.9, 6.9 Hz, 3H), 1.41 (dd, J=7.1, 7.1 Hz, 3H).


(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-isopropoxybenzoic acid



Starting from isopropoxy 3-isopropoxy-4-formylbenzoate (prepared as ethyl 3-ethoxy-4-formylbenzoate).

LCMS Method 11; Rt 3.82; ES⁺ 627.6.


(R)-6-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)nicotinic acid



Starting from methyl 6-formylnicotinate.

LCMS Method 12; Rt 2.83; ES⁺ 570.4

(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-fluoro-5-methoxybenzoic acid



Starting from methyl 2-fluoro-4-formyl-5-methoxybenzoate.

LCMS Method 11; Rt 3.66; ES⁺ 617.6.


Synthesis of methyl 2-fluoro-4-formyl-5-methoxybenzoate.

A solution of methyl 2-fluoro-5-methoxy-4-methylbenzoate (0.576 g, 2.91 mmol) in carbon tetrachloride (20 mL) was added with N-bromosuccinimide (0.57 g, 3.2 mmol) and benzoyl peroxide (cat). The reaction mixture was heated at 70 °C for 6 hours and then allowed to cool. The suspension was filtered and the filtrate evaporated at reduced pressure.

The residue was dissolved in DCM and washed with water, 1M aqueous sodium thiosulfate and the organic phase passed through a hydrophobic frit. The filtrate was concentrated at reduced pressure and the residue purified by flash column chromatography (eluent – 100% i-hexane to 4:1 i-hexane/ethyl acetate) to afford methyl 4-(bromomethyl)-2-fluoro-5-methoxybenzoate (0.591 g, 2.13 mmol). This material was dissolved in acetonitrile (20 mL) and treated with pyridine N-oxide (0.203 g, 2.13 mmol) and silver (I) oxide (0.247 g, 1.07 mmol) and the mixture stirred at room temperature for 18 hours. The suspension was filtered through a pad of celite and the filter pad washed with further acetonitrile. The filtrate was evaporated and the residue dissolved in DCM. The DCM solution was washed with water and then passed through a hydrophobic frit and the solvent was evaporated. The residue purified by flash column chromatography (eluent – 100% i-hexane to 4:1 i-hexane/ethyl acetate) to afford the title compound (0.230 g, 37%).

¹H NMR (400 MHz, CDCl₃); δ 10.45 – 10.44 (m, 1H), 7.39 – 7.52 (m, 2H), 3.98 (s, 3H), 3.97 (s, 3H).


***tert*-Butyl (R)-(4-amino-2-methoxybenzyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate**



From 2-methoxy-4-nitrobenzaldehyde


LCMS Method 11; Rt 3.57 min; ES⁺ 570.4

***tert*-Butyl (R)-(4-amino-5-fluoro-2-methoxybenzyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate**




From 5-fluoro-2-methoxy-4-nitrobenzaldehyde

LCMS Method 12; Rt 3.60 min; ES⁺ 588.5

Procedure 4A

(R)-5-(((*tert*-Butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)picolinic acid



5

A solution of *tert*-butyl (R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl((6-bromopyridin-3-yl)methyl)carbamate (prepared as described in Procedure 4 using 2-bromo-5-formylpyridine in Step 2) (1.47 g, 2.12 mmol) in toluene (50 mL) was added with phenyl formate (1.03 g, 8.47 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.294 g, 0.24 mmol), triethylamine (0.59 mL, 4.24 mmol) was added and the mixture was de-gassed for 15 minutes with nitrogen. Palladium acetate (0.057 g, 0.25 mmol) was added and the mixture was heated at 80 °C for 18 hours. Further phenyl formate (1.03 g, 8.47 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.294 g, 0.24 mmol), triethylamine (0.59 mL, 4.24 mmol) and palladium acetate (0.057 g, 0.25 mmol) was added and the reaction heated at 80 °C for a further 24 hours. The reaction mixture was evaporated at reduced pressure to about 1/3 of the initial volume and then diluted with ethyl acetate. The organic phase was washed with water, brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 100% ethyl acetate) to afford the major component. This material was dissolved in methanol (20 mL) and 2M aqueous sodium hydroxide added (2 mL) added. The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was concentrated and water added. The aqueous was washed with ether and then the


20

aqueous pH adjusted to 5. The aqueous was extracted with DCM (x3), the combined DCM extracts were passed through a hydrophobic frit and evaporated at reduced pressure. The residue was dissolved in ethanol (30 mL) and 10% Pd-C (0.064 g) added followed by 1-methyl-1,4-cyclohexadiene (0.282 g, 3.00 mmol). The reaction mixture was heated to reflux and heated at reflux for three hours. The reaction mixture was filtered and the filtrate evaporated at reduced pressure to afford the title compound (0.314 g, 26%).

LCMS Method 11; Rt 3.37 min; ES⁺ 570.6

Also prepared by this method;


(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethoxy)benzoic acid



Starting from 4-bromo-2-(trifluoromethoxy)benzaldehyde.


LCMS Method 12; Rt 3.02 min; ES⁺ 653.5.

(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-fluoro-3-methoxybenzoic acid




Starting from 4-bromo-3-fluoro-2-methoxybenzaldehyde.

LCMS Method 12; Rt 3.02 min; ES⁺ 653.5.


Procedure 4B

(R)-4-(((*tert*-Butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-chlorobenzoic acid



5

Step 1; (R)-5-(2-Amino-1-(((*tert*-butyldimethylsilyl)oxy)ethyl)-8-hydroxyquinolin-2(1H)-one




A solution of (R)-5-(2-azido-1-(((*tert*-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (8.5 g, 18.9 mmol) in ethanol (120 mL) was added with 10% palladium on charcoal (8.5 g) followed by 1-methyl-1,4-cyclohexadiene (21 mL, 186.9 mmol) and the reaction mixture heated to 60 °C for one hour. The reaction was allowed to cool to room temperature and the suspension was filtered. The filter cake was washed with further ethanol and the filtrate evaporated at reduced pressure. The residue was triturated with i-hexane, the solid recovered by filtration and subsequently dried under vacuum to afford the title compound (5.55 g, 88%).

^1H NMR (400 MHz, CDCl_3); δ 8.22 (d, $J=9.8$ Hz, 1H), 7.04 (d, $J=8.0$ Hz, 1H), 6.91 (d, $J=7.9$ Hz, 1H), 6.63 (d, $J=9.3$ Hz, 1H), 5.03 (dd, $J=4.9$, 4.9 Hz, 1H), 3.04 - 2.93 (m, 2H), 0.92 - 0.85 (m, 9H), 0.20 (s, 3H), -0.18 (s, 3H).

20

Step 2; (R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-chlorobenzoic acid




The title compound was prepared as in Preparation 4 using Step 3 and 5.

5 LCMS Method 11; Rt 3.76; ES⁺ 603.3 / 605.4.

Also prepared in the same fashion were;


(R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-chlorobenzoic acid



10 Starting from methyl 2-chloro-4-formylbenzoate

LCMS Method 11; Rt 3.70; ES⁺ 603.3 / 605.4

(R)-4-(((tert-Butoxycarbonyl)(2-(((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-chloro-5-methoxybenzoic acid



15 Starting from phenyl 2-chloro-4-formyl-5-methoxybenzoate

LCMS Method 11; Rt 3.88; ES⁺ 633.5 / 635.6

Synthesis of phenyl 2-chloro-4-formyl-5-methoxybenzoate


A stirred solution of 4-bromo-2-methoxybenzaldehyde (1.0 g, 4.56 g) in acetonitrile (15 mL) was added with N-chlorosuccinimide (0.731 g, 5.48 mmol) and the reaction mixture was heated at 80 °C for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate, filtered and the solvent evaporated at reduced pressure. The residue was dissolved in toluene (20 mL) and phenyl formate (2 mL, 18.3 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.642 g, 1.11 mmol), triethylamine (1.27 mL, 9.13 mmol) was added and the mixture was de-gassed for 15 minutes with nitrogen.

Palladium acetate (0.122 g, 0.54 mmol) was added and the mixture was heated at 80 °C for 18 hours. The reaction mixture was evaporated at reduced pressure to about 1/3 of the initial volume and then diluted with ethyl acetate. The organic phase was washed with water, brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was purified by flash column chromatography (eluent – 100% *iso*-hexane to 100% ethyl acetate) to afford the title compound (0.554 g, 42%).


¹H NMR (400 MHz, CDCl₃); δ 10.47 (s, 1H), 7.93 (s, 1H), 7.58 (s, 1H), 7.46 - 7.31 (m, 5H), 4.02 (s, 3H).

Procedure 5

10 **Preparation of (*R*)-4-(2-((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoic acid**



15 **Step 1; (*R*)-8-(Benzyloxy)-5-(1-((*tert*-butyldimethylsilyl)oxy)-2-((4-hydroxyphenethyl)amino)ethyl)quinolin-2(1H)-one**




A mixture of (*R*)-8-(benzyloxy)-5-(2-bromo-1-((*tert*-butyldimethylsilyl)oxy)ethyl)quinolin-2(1H)-one (1.0 g, 2.05 mmol) in NMP (2 mL) was added with tyramine (1.41 g, 10.2 mmol). The mixture was heated at 80 °C for 18 hours. The mixture was diluted with ethyl acetate and washed sequentially with 10% aqueous potassium hydrogensulfate and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure to afford the title compound (2.58 g, 93%).

¹H NMR (400 MHz, CDCl₃); δ 9.30 - 9.30 (m, 1H), 8.27 (d, J=9.9 Hz, 1H), 7.46 -

7.38 (m, 6H), 7.08 (d, J=8.3 Hz, 1H), 6.97 (dd, J=4.8, 8.4 Hz, 3H), 6.71 (d, J=8.5 Hz, 2H), 6.65 (d, J=9.9 Hz, 1H), 5.16 (s, 2H), 5.14 - 5.07 (m, 1H), 2.97 - 2.82 (m, 3H), 2.78 - 2.68 (m, 3H), 1.65 (br s, 1H), 0.82 (s, 9H), 0.00 (s, 3H), -0.22 (s, 3H).


Step 2; (R)-tert-Butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(4-hydroxyphenethyl)carbamate



A stirred solution of **(R)-8-(benzyloxy)-5-(1-((tert-butyldimethylsilyl)oxy)-2-((4-hydroxyphenethyl)amino)ethyl)quinolin-2(1H)-one** (6.0 g, 11.02 mmol) in DCM (75 mL) was added with a solution of di-*tert*-butyldicarbonate (2.18 g, 16.5 mmol) in DCM (15 mL). The reaction mixture was stirred at room temperature for 2 hours. The solvent was evaporated under reduced pressure and the residue purified by flash column chromatography (eluent – 100% *iso*-hexane to 3:2 *iso*-hexane/ethyl acetate) to afford the title compound (6.30 g, 89 %).

¹H NMR (400 MHz, DMSO-d₆, 100 °C); δ 9.94 - 9.92 (m, 1H), 8.74 (s, 1H), 8.29 (d, J=9.9 Hz, 1H), 7.56 (d, J=7.2 Hz, 2H), 7.44 - 7.35 (m, 3H), 7.24 (d, J=8.3 Hz, 1H), 7.14 (d, J=8.4 Hz, 1H), 6.92 (d, J=8.4 Hz, 2H), 6.70 - 6.67 (m, 2H), 6.56 (d, J=9.9 Hz, 1H), 5.31 (s, 3H), 3.37 - 3.23 (m, 4H), 2.66 - 2.55 (m, 2H), 1.39 (s, 9H), 0.86 (s, 9H), 0.04 (s, 3H), -0.13 (s, 3H).

Step 3; (R)-4-(2-((2-(8-(Benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)ethyl)phenyl trifluoromethanesulfonate




A cooled solution of **(R)-tert-butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-**

5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(4-hydroxyphenethyl)carbamate (6.3 g, 9.78 mmol) in DCM (100 mL) at 0 °C was added with triethylamine (2.72 mL, 19.56 mmol) followed by 2-[N,N-bis(trifluoromethanesulfonyl)amino]-5-chloropyridine (4.21 g, 10.76 mmol) and the mixture was stirred at 0 °C for 2 hours. The coolant was removed and 2M aqueous sodium hydroxide (25 mL) was added to the reaction mixture, and the mixture was stirred for 15 minutes. The organic phase was separated, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated under reduced pressure to afford the title compound (7.3 g, 96%).

¹H NMR (400 MHz, DMSO-d₆, 100 °C); δ 8.74 (s, 1H), 8.29 (d, J=9.9 Hz, 1H), 7.56 (d, J=7.2 Hz, 2H), 7.44 - 7.35 (m, 3H), 7.24 (d, J=8.3 Hz, 1H), 7.14 (d, J=8.4 Hz, 1H), 6.92 (d, J=8.4 Hz, 2H), 6.70 - 6.67 (m, 2H), 6.56 (d, J=9.9 Hz, 1H), 5.31 (s, 3H), 2.96 (s, 4H), 2.66 - 2.55 (m, 2H), 1.39 (s, 9H), 0.86 (s, 9H), 0.04 (s, 3H), -0.13 (s, 3H).

Step 4; (*R*)-Phenyl 4-(2-((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(*tert*-butoxycarbonyl)amino)ethyl)benzoate



15


A solution of (*R*)-4-(2-((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(*tert*-butoxycarbonyl)amino)ethyl)phenyl trifluoromethanesulfonate (7.29 g, 9.4 mmol) in toluene (75 mL) was added with phenyl formate (4 mL, 37.57 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (1.3 g, 2.25 mmol), triethylamine (2.61 mL, 18.81 mmol) and the mixture was de-gassed for 15 minutes with nitrogen. Palladium acetate (0.242 g, 1.12 mmol) was added and the mixture was heated at 80 °C for 6 hours. The reaction mixture was evaporated under reduced pressure to about 1/3 of the initial volume and then diluted with ethyl acetate. The organic phase was washed with water, brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated under reduced pressure. The residue was purified by flash column

25

chromatography (eluent – 100% *iso*-hexane to 100% ethyl acetate) to afford the title compound (4.5 g, 64%).

¹H NMR (400 MHz, DMSO-d₆, 100 °C); δ 9.98 - 9.96 (m, 1H), 8.29 (d, J=9.9 Hz, 1H), 8.05 (d, J=8.3 Hz, 2H), 7.56 (d, J=7.2 Hz, 2H), 7.51 - 7.46 (m, 2H), 7.43 - 7.27 (m, 8H), 7.24 (s, 1H), 7.17 (d, J=8.3 Hz, 1H), 6.57 (d, J=9.9 Hz, 1H), 5.35 (d, J=1.9 Hz, 1H), 5.32 (s, 2H), 3.49 - 3.34 (m, 4H), 2.90 - 2.80 (m, 2H), 1.39 (s, 9H), 0.87 (s, 9H), 0.06 (s, 3H), -0.12 (s, 3H).

Step 5; (R)-4-(2-((2-(8-(Benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(*tert*-butoxycarbonyl)amino)ethyl)benzoic acid




10

A solution of (*R*)-phenyl 4-(2-((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(*tert*-butoxycarbonyl)amino)ethyl)benzoate (4.5 g, 6.0 mmol) in THF (30 mL) was added with 2M aqueous sodium hydroxide (30mL) and the mixture was stirred at room temperature for 16 hours. The solvent was evaporated under reduced pressure to half the initial volume and then diluted with water (25 mL). The resulting mixture was acidified to pH 3 using 2M aqueous hydrochloric acid and then extracted with ethyl acetate (x3). The combined organic phases were washed sequentially with water, brine (25ml), dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated under reduced pressure to afford the title compound (4 g, 100 %). Material used directly in the next step.

20

Step 6; (R)-4-(2-((*tert*-Butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoic acid




A solution of (*R*)-4-(2-((2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(*tert*-butoxycarbonyl)amino)ethyl)benzoic acid (4 g, 5.95 mmol) in ethanol (100 mL) was added with 10% Pd-C (2 g) and 1-methyl-1,4-cyclohexadiene (3.29 mL, 29.76 mmol) and the mixture was heated at 80 °C for 3.5 hours.


- 5 The reaction mixture was filtered through celite and the celite washed with further ethanol. The filtrate was evaporated under reduced pressure to afford the title compound (3.5 g, 100%).

Procedure 6

Preparation of (*R*)-*tert*-butyl (2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)(piperidin-4-ylmethyl)carbamate



Step 1; (*R*)-8-(Benzyloxy)-5-(2-(((1-benzylpiperidin-4-yl)methyl)amino)-1-((*tert*-butyldimethylsilyl)oxy)ethyl)quinolin-2(1H)-one




- 15 A mixture of (*R*)-8-(benzyloxy)-5-(2-bromo-1-((*tert*-butyldimethylsilyl)oxy)ethyl)quinolin-2(1H)-one (2.39 g, 4.90 mmol) in NMP (4 mL) was added with (1-benzylpiperidin-4-yl)methanamine (5 g, 24.5 mmol). The mixture was heated at 80 °C for 18 hours. The mixture was diluted with ethyl acetate and washed sequentially with water (x2) and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated. The residue was purified by flash column chromatography (eluent – 100% DCM to 40:1 DCM / 7M NH₃/MeOH) to afford the title compound (2.65 g, 88%).

¹H NMR (400 MHz, CDCl₃); δ 9.19 - 9.15 (m, 1H), 8.30 - 8.28 (s, 1H), 7.43 - 7.41 (m, 5H), 7.31 (d, J=4.4 Hz, 5H), 7.12 (d, J=8.3 Hz, 1H), 7.00 (d, J=8.3 Hz, 1H), 6.66 (d,

J=9.9 Hz, 1H); 5.16 (s, 2H), 5.15 – 5.10 (m, 1H), 3.49 (s, 2H), 2.95 - 2.84 (m, 3H), 2.71 (dd, J=4.0, 12.2 Hz, 1H), 2.56 - 2.43 (m, 2H), 1.94 (dd, J=11.7, 11.7 Hz, 2H), 1.51 - 1.39 (m, 2H), 1.30 - 1.22 (m, 3H), 0.87 (s, 9H), 0.06 (s, 3H), -0.19 (s, 3H).


Step 2; (R)-tert-Butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)((1-benzylpiperidin-4-yl)methyl)carbamate



A stirred solution of (R)-8-(benzyloxy)-5-(2-(((1-benzylpiperidin-4-yl)methyl)amino)-1-((tert-butyldimethylsilyl)oxy)ethyl)quinolin-2(1H)-one (2.65 g, 4.33 mmol) in DCM (25 mL) was added with a solution of di-tert-butyl dicarbonate (1.13 g, 5.18 mmol) in DCM (5 mL). The reaction mixture was stirred at room temperature for 16 hours. The solvent was evaporated under reduced pressure and the residue purified by flash column chromatography (eluent – 100% DCM to 30:1 DCM / 7M NH₃/MeOH) to afford the title compound (2.83 g, 92 %).

¹H NMR (400 MHz, DMSO-d₆, 100 °C); δ 9.96 (s, 1H), 8.30 (d, J=9.9 Hz, 1H), 7.55 (d, J=7.3 Hz, 2H), 7.42 - 7.21 (m, 9H), 7.14 (d, J=8.3 Hz, 1H), 6.55 (d, J=9.9 Hz, 1H), 5.40 - 5.33 (m, 1H), 5.30 (s, 2H), 3.44 (s, 2H), 3.38 (d, J=6.1 Hz, 2H), 3.12 - 2.97 (m, 2H), 2.95 (s, 2H), 2.78 - 2.73 (m, 2H), 2.00 - 1.88 (m, 2H), 1.40 (s, 10H), 1.22 - 1.09 (m, 2H), 0.87 (s, 9H), 0.06 (s, 3H), -0.13 (s, 3H).

Step 3; (R)-tert-Butyl (2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)(piperidin-4-ylmethyl)carbamate




A stirred solution of (R)-tert-butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)((1-benzylpiperidin-4-yl)methyl)carbamate (2.80

g, 4.33 mmol) in ethanol (30 mL) was added with 10% Pd/C (2.8 g) and 1-methyl-1,4-cyclohexadiene (4.86 mL, 43.3 mmol). The reaction mixture was heated to reflux [**Care – vigorous evolution of gas**] and heated under reflux for 1 hour. The suspension was filtered and the filtrate evaporated under reduced pressure to afford the title compound (2.26 g, 5 98%).

¹H NMR (400 MHz, DMSO-d₆, 100 °C); δ 8.26 (d, J=9.9 Hz, 1H), 7.05 (d, J=8.2 Hz, 1H), 7.01 (d, J=8.0 Hz, 1H), 6.53 - 6.49 (m, 1H), 5.37 - 5.35 (m, 1H), 3.22 (d, J=12.8 Hz, 2H), 3.08 (ddd, J=7.0, 14.3, 17.8 Hz, 2H), 2.83 - 2.72 (m, 2H), 1.84 - 1.76 (m, 1H), 1.71 - 1.61 (m, 2H), 1.43 - 1.41 (m, 14H), 0.87 - 0.85 (m, 9H), 0.02 (s, 3H), -0.14 (d, J=2.1 10 Hz, 3H).


Also prepared by this method (R)-tert-butyl (2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)(2-(piperidin-4-yl)ethyl)carbamate




¹H NMR (400 MHz, DMSO-d₆, 90 °C); δ 8.26 (d, J=9.9 Hz, 1H), 7.05 (d, J=8.2 Hz, 1H), 7.01 (d, J=8.0 Hz, 1H), 6.53 - 6.49 (m, 1H), 5.37 - 5.35 (m, 1H), 3.40 - 3.30 (m, 15 2H), 3.25 - 3.10 (m, 2H), 3.00 - 2.90 (m, 2H), 2.50 - 2.40 (m, 2H), 1.60 - 1.55 (m, 2H), 1.45 (s, 9H), 1.40 - 1.30 (m, 2H), 1.20 - 0.95 (m, 3H), 0.87 - 0.85 (m, 9H), 0.02 (s, 3H), -0.14 (d, J=2.1 Hz, 3H).

Procedure 7

20 **Preparation of (S)-4-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoic acid hydrochloride**




Step 1; (S)-tert-Butyl 4-((2-hydroxy-2-(3-hydroxyphenyl)-2-

phenylacetoxymethyl)piperidine-1-carboxylate

A stirred solution of (*S*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid (10.69 g, 43.77 mmol) in DMF (30 mL) was added with potassium hydrogen carbonate (8.76 g, 87.5 mmol) and the mixture stirred at room temperature for 10 minutes. A solution of *tert*-butyl 4-(tosyloxymethyl)piperidine-1-carboxylate (as described in Example 1, Step 5) (16.17 g, 43.77 mmol) in DMF (55 mL) was added and the reaction mixture heated at 60 °C for 27 hours. Diluted with ethyl acetate and washed with aqueous sodium hydrogen carbonate (x2), brine and dried over anhydrous magnesium sulfate. The filtrate was evaporated under reduced pressure and the residue purified by flash column chromatography (eluent – 100% *iso*-hexane to 2:3 *iso*-hexane/ethyl acetate) to afford the title compound (13.42 g, 69 %).

¹H NMR (400 MHz, DMSO-*d*₆); δ 9.35 (s, 1H), 7.33 (d, *J*=4.4 Hz, 4H), 7.31 - 7.26 (m, 1H), 7.12 (dd, *J*=7.8, 7.8 Hz, 1H), 6.77 - 6.73 (m, 2H), 6.69 - 6.66 (m, 1H), 6.50 (s, 1H), 4.07 - 3.99 (m, 2H), 3.87 (d, *J*=11.4 Hz, 2H), 2.68 - 2.63 (m, 2H), 1.76 - 1.67 (m, 1H), 1.48 (d, *J*=12.0 Hz, 2H), 1.38 (s, 9H), 1.10 - 0.83 (m, 2H).


Step 2; (*S*)-Piperidin-4-ylmethyl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate hydrochloride

A solution of (*S*)-*tert*-butyl 4-((2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetoxymethyl)piperidine-1-carboxylate (7.07 g, 16.01 mmol) in 1,4-dioxane (20 mL) was added with a solution of HCl-dioxane (4M, 80 mL) and the reaction mixture was stirred at room temperature for 1 hour 30 minutes. The solvent was evaporated under

reduced pressure to afford the title compound (5.9 g, 97%).

¹H NMR (400 MHz, DMSO-d₆); δ 9.38 (s, 1H), 8.69 - 8.68 (m, 1H), 8.39 (s, 1H), 7.34 (d, J=4.0 Hz, 4H), 7.32 - 7.27 (m, 1H), 7.13 (dd, J=7.8, 7.8 Hz, 1H), 6.78 - 6.74 (m, 2H), 6.70 - 6.67 (m, 1H), 6.52 (s, 1H), 4.02 (d, J=6.5 Hz, 2H), 3.21 (d, J=12.4 Hz, 2H),
5 2.86 - 2.77 (m, 2H), 1.92 - 1.84 (m, 1H), 1.68 (d, J=13.9 Hz, 2H), 1.35 - 1.25 (m, 2H).


Step 3; (*S*)-(1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate



A stirred suspension of (*S*)-piperidin-4-ylmethyl 2-hydroxy-2-(3-hydroxyphenyl)-
10 2-phenylacetate hydrochloride (5.9 g, 15.61 mmol) in DCM (160 mL) was added with benzaldehyde (2.38 mL, 23.4 mmol) and stirred at 30 °C for 1 hour. Sodium triacetoxyborohydride (6.62 g, 31.2 mmol) was added and the reaction mixture stirred at 30 °C for 2 hours. Further sodium triacetoxyborohydride (1.65 g, 7.8 mmol) was added and the mixture stirred at 30 °C for 18 hours. The reaction mixture was quenched with
15 saturated aqueous sodium hydrogen carbonate and the organic phase removed. The aqueous phase was extracted with further DCM. The combined organic phases were washed with brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated under reduced pressure. The residue was purified by flash column chromatography (eluent – 100% DCM to 9:1 DCM/methanol) to afford the title compound
20 (13.42 g, 69 %).

¹H NMR (400 MHz, DMSO-d₆); δ 9.33 (s, 1H), 7.34 - 7.25 (m, 9H), 7.11 (dd, J=7.8, 7.8 Hz, 1H), 6.76 - 6.73 (m, 2H), 6.68 - 6.65 (m, 1H), 6.48 (s, 1H), 5.77 (s, 1H), 3.99 (d, J=6.3 Hz, 2H), 3.40 (s, 2H), 2.72 (d, J=11.4 Hz, 2H), 1.86 - 1.79 (m, 2H), 1.54 - 1.43 (m, 3H), 1.12 (q, J=11.8 Hz, 2H).


25 **Step 4; (*S*)-*tert*-Butyl 4-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoate**



A stirred solution of (*S*)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetate (1.23 g, 2.84 mmol) in DMF (15 mL) was added with potassium carbonate (0.785 g, 5.69 mmol) and the mixture stirred at room temperature for 5 minutes. *tert*-Butyl 4-(bromomethyl)benzoate (0.924 g, 6.83 mmol) was added and the reaction mixture was heated at 40 °C for 8 hours. Diluted with ethyl acetate and washed with aqueous sodium hydrogen carbonate (x2), brine and dried over anhydrous magnesium sulfate. The filtrate was evaporated under reduced pressure and the residue purified by flash column chromatography (eluent – 100% DCM to 7:3 DCM / methanol) to afford the title compound (1.89 g, 100 %).

¹H NMR (400 MHz, DMSO-*d*₆); δ 7.89 (d, *J*=8.4 Hz, 2H), 7.50 (d, *J*=8.4 Hz, 2H), 7.30 - 7.23 (m, 11H), 6.97 - 6.91 (m, 3H), 6.60 (s, 1H), 5.15 (s, 2H), 3.98 (d, *J*=6.3 Hz, 2H), 3.39 (s, 2H), 2.74 - 2.68 (m, 2H), 1.85 - 1.77 (m, 2H), 1.55 (s, 9H), 1.53 - 1.43 (m, 3H), 1.18 - 1.06 (m, 2H).

Step 5; (*S*)-4-((3-(2-((1-Benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoic acid hydrochloride




A stirred solution of (*S*)-*tert*-butyl 4-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoate (1.89 g, 3.04 mmol) in 1,4-dioxane (5 mL) was added with a solution of HCl-dioxane (4M, 15 mL) and the reaction mixture stirred at room temperature for 18 hours. The solvent was evaporated under

reduced pressure to afford the title compound (1.83 g, 100%).

^1H NMR (400 MHz, DMSO- d_6); δ 7.95 (d, $J=8.3$ Hz, 2H), 7.52 (d, $J=8.3$ Hz, 4H), 7.48 - 7.44 (m, 3H), 7.32 (dd, $J=3.4$, 3.4 Hz, 6H), 6.99 - 6.90 (m, 3H), 6.64 (s, 1H), 5.15 (s, 2H), 4.24 - 4.19 (m, 2H), 4.00 (d, $J=6.4$ Hz, 2H), 3.28 (s, 1H), 3.06 (s, 1H), 2.90 - 2.88 (m, 2H), 1.82 (s, 1H), 1.71 (d, $J=13.2$ Hz, 2H), 1.44 (d, $J=11.7$ Hz, 2H).

Procedure 8


Preparation of (*S*)-3-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoic acid hydrochloride



The title compound was prepared as in Procedure 7 with *tert*-butyl 3-(bromomethyl)benzoate replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

Procedure 9


Preparation of (*S*)-2-(3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)acetic acid hydrochloride



The title compound was prepared as in Procedure 7 with *tert*-butyl bromoacetate replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

Procedure 10


Preparation of (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-phenyl-2-(3-(piperidin-4-ylmethoxy)phenyl)acetate hydrochloride




The title compound was prepared as in Procedure 7 with *tert*-butyl 4-(tosyloxymethyl)piperidine-1-carboxylate (as described in Example 1, Step 5) replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

Procedure 11

Preparation of (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(azetidin-3-ylmethoxy)phenyl)-2-hydroxy-2-phenylacetate 2,2,2-trifluoroacetate




Step 1; (S)-*tert*-Butyl 3-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)azetidine-1-carboxylate



The title compound was prepared as in Procedure 7 with *tert*-butyl 3-(bromomethyl)azetidine-1-carboxylate replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

¹H NMR (400 MHz, DMSO-d₆); δ 7.35 – 7.21 (6H), 6.89 (dd, J=6.3, 6.3 Hz, 3H), 6.60 (s, 1H), 4.06 - 3.89 (m, 6H), 3.82 (d, J=6.8 Hz, 2H), 3.71 (d, J=7.2 Hz, 2H), 3.65 - 3.54 (m, 4H), 3.40 (s, 2H), 2.95 - 2.86 (m, 1H), 2.71 (d, J=11.8 Hz, 2H), 1.83 (dd, J=11.4, 11.4 Hz, 2H), 1.53 - 1.43 (m, 2H), 1.38 (s, 9H), 1.22 - 1.08 (m, 2H).


Step 2; (S)-(1-Benzylpiperidin-4-yl)methyl 2-(3-(azetidin-3-ylmethoxy)phenyl)-2-hydroxy-2-phenylacetate 2,2,2-trifluoroacetate



A solution of (S)-*tert*-butyl 3-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)azetidine-1-carboxylate (0.388 g, 0.65 mmol) in DCM (2 mL) was added with trifluoroacetic acid (2 mL). The reaction mixture was stirred at room temperature for 1 hour. The solvent was evaporated under reduced pressure and the title compound was used without further purification in subsequent steps.

Procedure 12


Preparation of (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-aminopropoxy)phenyl)-2-hydroxy-2-phenylacetate hydrochloride



The title compound was prepared as in Procedure 7 with *tert*-butyl (3-bromopropyl)carbamate replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

Procedure 13


Preparation of (*R*)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-aminoethoxy)phenyl)-2-hydroxy-2-phenylacetate hydrochloride




The title compound was prepared as in Procedure 7 with (*R*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid replacing (*S*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid in Step 1 and *tert*-butyl (2-chloroethyl)carbamate replacing *tert*-butyl 4-(bromomethyl)benzoate in **Step 4**.

Procedure 14

Preparation of (1-benzylpiperidin-4-yl)methyl 2-(3-(2-aminoethyl)phenyl)-2-hydroxy-2-phenylacetate



Step 1; *tert*-Butyl 4-((2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetoxy)methyl)piperidine-1-carboxylate




A stirred solution of *tert*-butyl 4-((2-(3-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate (18.4 g, 34.6 mmol) in ethanol (300 mL) was added with 10% Pd/C (4.6 g) and 1-methyl-1,4-cyclohexadiene (19.6 mL, 173.2 mmol). The reaction mixture was heated to reflux for two hours. The suspension was filtered through celite and the filtrate evaporated under reduced pressure to afford the title

product (15.5 g, 100%).


¹H NMR (400 MHz, CDCl₃); δ 7.41 - 7.30 (m, 6H), 7.20 (dd, J=8.0, 8.0 Hz, 1H), 7.00 (s, 1H), 6.87 (s, 1H), 6.80 (dd, J=2.1, 8.0 Hz, 1H), 4.22 (s, 1H), 4.10 (d, J=5.9 Hz, 2H), 4.08 - 4.02 (m, 2H), 3.75 - 3.69 (m, 1H), 2.68 - 2.57 (m, 2H), 1.83 - 1.74 (m, 1H), 1.69 - 1.55 (m, 1H), 1.49 (d, J=12.7 Hz, 1H), 1.40 (s, 9H), 1.21 - 1.04 (m, 1H).

Step 2; *tert*-Butyl 4-((2-hydroxy-2-phenyl-2-(3-(((trifluoromethyl)sulfonyl)oxy)phenyl)acetoxy)methyl)piperidine-1-carboxylate



The title compound was prepared as described in Procedure 5 Step 3 with *tert*-butyl 4-((2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetoxy)methyl)piperidine-1-carboxylate replacing (*R*)-*tert*-Butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((*tert*-butyldimethylsilyl)oxy)ethyl)(4-hydroxyphenethyl)carbamate. The product was used directly in the next step.


Step 3; Piperidin-4-ylmethyl 2-hydroxy-2-phenyl-2-(3-(((trifluoromethyl)sulfonyl)oxy)phenyl)acetate hydrochloride



The title compound was prepared as described in Procedure 7 Step 2.

¹H NMR (400 MHz, DMSO-*d*₆); δ 7.57 (dd, J=8.0, 8.0 Hz, 1H), 7.49 - 7.46 (m, 2H), 7.41 - 7.31 (m, 6H), 7.05 (s, 1H), 4.06 (d, J=6.7 Hz, 2H), 3.21 (d, J=12.4 Hz, 2H), 2.85 - 2.75 (m, 2H), 1.93 - 1.85 (m, 1H), 1.69 - 1.60 (m, 2H), 1.36 - 1.25 (m, 2H).


Step 4; (1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-phenyl-2-(3-

(((trifluoromethyl)sulfonyl)oxy)phenyl)acetate

The title compound was prepared as described in Procedure 7 Step 3.


¹H NMR (400 MHz, CDCl₃); δ 7.52 (d, J=7.9 Hz, 1H), 7.44 - 7.39 (m, 2H), 7.37 - 7.21 (m, 11H), 4.33 (s, 1H), 4.12 (d, J=6.8 Hz, 2H), 3.49 (s, 2H), 3.46 (s, 2H), 2.82 (d, J=11.5 Hz, 2H), 1.92 - 1.84 (m, 2H), 1.52 - 1.44 (m, 1H), 1.27 - 1.15 (m, 2H).

Step 5; (1-Benzylpiperidin-4-yl)methyl 2-(3-(2-((tert-butoxycarbonyl)amino)ethyl)phenyl)-2-hydroxy-2-phenylacetate



A mixture of (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-phenyl-2-(3-(((trifluoromethyl)sulfonyl)oxy)phenyl)acetate (1.56 g, 2.77 mmol), potassium *tert*-butyl N-[2-(trifluoroboranuidyl)ethyl]carbamate (0.7 g, 2.79 mmol), palladium (II) acetate (0.09 g, 0.41 mmol), RuPhos (0.39 g, 0.83 mmol) and cesium carbonate (2.92 g, 8.3 mmol) in toluene / water (30 mL, 3 to 1) was thoroughly degassed with nitrogen and then heated at 95 °C for 16 hours. The reaction mixture was partitioned between DCM and brine. The organic phase passed through a hydrophobic frit and the filtrate evaporated under reduced pressure. The residue purified by flash column chromatography (eluent – 100% hexane to 100% ethyl acetate) to afford the title compound (1.21 g, 78 %).


Step 6: (1-Benzylpiperidin-4-yl)methyl 2-(3-(2-aminoethyl)phenyl)-2-hydroxy-

2-phenylacetate hydrochloride

The title compound was prepared as in Procedure 7 Step 2.

Procedure 15


5 Preparation of 1-Benzylpiperidin-4-yl (S)-2-(3-((4-(aminomethyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate dihydrochloride



The title compound was prepared as in Procedure 7 with *tert*-butyl (4-(bromomethyl)benzyl)carbamate replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

10 Example 5

(S)-(1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(2-((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate (Compound 31)















- 15 A stirred solution of (*R*)-4-(2-((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoic acid (0.13 g, 0.23 mmol) and *N,N*-diisopropylethylamine (0.066 mL, 0.38 mmol) in DMF (2 mL) was added with HATU (0.11 g, 0.29 mmol) and the mixture stirred at room temperature for 30 minutes. A solution of (*S*)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-phenyl-2-(3-(piperidin-4-ylmethoxy)phenyl)acetate hydrochloride (0.11 g, 0.19 mmol) in DMF (1 mL) was added
- 20

and the reaction mixture stirred at room temperature for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with water and brine (x2). The organic phase was dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure. The residue was treated with a solution of HCl-dioxane (4M, 15 mL) and the
5 reaction mixture stirred at room temperature for 18 hours. The solvent was evaporated under reduced pressure and the residue purified by reverse phase preparative HPLC. To afford the title compound (31).

The following compounds were prepared as described in Example 4 with the relevant acid coupled to the appropriate amine.

N	Relevant acid	Relevant Amine	Structure
32			
32A			
32B			
32C			
32D			

(continued)

32E			
32F			
32G			
32H			

(continued)

			32I
			32J
			32K
			32L

(continued)

32M			
33			
33A			
34			










(continued)

35		
36		
36A		










(continued)

			37
			38
			39

(continued)

		
		
		
40	41	42

(continued)

42A			
42B			
42C			

(continued)

42D	
42E	
42F	

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
31	7.32	7	(DMSO-d ₆ , 90 °C); δ 8.25 (s, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.33 - 7.22 (m, 15H), 7.08 (d, J=8.2 Hz, 1H), 6.93 (d, J=8.0 Hz, 1H), 6.89 - 6.86 (m, 3H), 6.52 (d, J=9.9 Hz, 1H), 5.13 - 5.08 (m, 1H), 4.55 - 4.42 (m, 1H), 4.02 - 3.99 (m, 2H), 3.79 (d, J=6.1 Hz, 2H), 3.67 - 3.55 (m, 1H), 3.39 (s, 2H), 2.95 - 2.85 (m, 4H), 2.84 - 2.76 (m, 3H), 2.75 - 2.67 (m, 2H), 2.04 - 1.95 (m, 1H), 1.87 - 1.67 (m, 4H), 1.53 - 1.42 (m, 3H), 1.24 - 1.07 (m, 4H).	mono Formate
32	7.18	7	(MeOD); δ 8.50 (s, 2H), 8.29 (d, J=9.9 Hz, 1H), 7.87 (d, J=8.4 Hz, 2H), 7.56 (d, J=8.4 Hz, 2H), 7.40 (s, 6H), 7.39 - 7.32 (m, 4H), 7.28 - 7.23 (m, 2H), 7.03 - 6.96 (m, 3H), 6.90 (dd, J=2.4, 8.2 Hz, 1H), 6.64 (d, J=9.8 Hz, 1H), 5.36 (dd, J=4.4, 8.8 Hz, 1H), 4.21 (s, 2H), 4.11 - 4.03 (m, 4H), 3.94 (s, 2H), 3.57 (dd, J=6.9, 6.9 Hz, 2H), 3.22 - 3.09 (m, 4H), 2.55 (dd, J=10.4, 12.3 Hz, 2H), 2.05 (s, 3H), 1.84 - 1.78 (m, 1H), 1.70 (d, J=12.9 Hz, 2H), 1.36 (q, J=12.5 Hz, 2H).	Formate
32A	2.47	11	(MeOD); δ 8.47 (s, 2H), 8.26 (d, J=9.9 Hz, 1H), 7.54 (s, 1H), 7.51 - 7.25 (m, 14H), 7.04 (d, J=8.2 Hz, 1H), 7.00 - 6.96 (m, 2H), 6.91 (dd, J=2.2, 8.0 Hz, 1H), 6.65 (d, J=9.8 Hz, 1H), 5.46 - 5.38 (m, 1H), 4.34 (s, 2H), 4.13 - 4.04 (m, 6H), 3.97 (s, 3H), 3.59 (dd, J=6.9, 6.9 Hz, 2H), 3.29 (d, J=12.7 Hz, 2H), 3.22 - 3.18 (m, 2H), 2.77 - 2.67 (m, 2H), 2.15 - 2.07 (m, 2H), 1.93 - 1.82 (m, 1H), 1.74 (d, J=13.6 Hz, 2H), 1.48 - 1.37 (m, 2H).	Formate
32B	2.51	11	(MeOD); δ 8.53 (s, 1H), 8.30 (d, J=9.9 Hz, 1H), 7.90 (d, J=7.9 Hz, 1H), 7.41 - 7.23 (m, 13H), 7.14 (dd, J=1.2, 7.9 Hz, 1H), 7.04 - 6.98 (m, 3H), 6.93 (dd, J=2.0, 8.2 Hz, 1H), 6.64 (d, J=9.9 Hz, 1H), 5.37 (dd, J=4.8, 8.3 Hz, 1H), 4.17 (s, 2H), 4.11 - 4.05 (m, 4H), 3.93 (s, 2H), 3.89 (s, 3H), 3.60 (t, J=6.7 Hz, 2H), 3.20 - 3.09 (m, 4H), 2.53 (dd, J=10.2, 12.2 Hz, 2H), 2.13 - 2.06 (m, 2H), 1.86 - 1.73 (m, 1H), 1.71 - 1.63 (m, 2H), 1.42 - 1.29 (m, 2H).	mono-formate

(continued)

32C	2.46	11	(MeOD); δ 8.99 (d, J=1.5 Hz, 1H), 8.50 (s, 1H), 8.36 (d, J=9.9 Hz, 1H), 8.20 (dd, J=2.3, 8.1 Hz, 1H), 7.54 (d, J=8.3 Hz, 1H), 7.40 - 7.31 (m, 12H), 7.02 (d, J=8.1 Hz, 1H), 6.99 - 6.95 (m, 2H), 6.91 - 6.88 (m, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.39 (dd, J=5.2, 7.7 Hz, 1H), 4.28 (s, 2H), 4.12 - 4.04 (m, 4H), 3.88 (s, 2H), 3.59 (dd, J=6.9, 6.9 Hz, 2H), 3.18 - 3.11 (m, 4H), 2.53 - 2.43 (m, 2H), 2.14 - 2.05 (m, 2H), 1.83 - 1.76 (m, 1H), 1.71 - 1.64 (m, 2H), 1.40 - 1.30 (m, 2H).	mono-formate
32D	2.53	11	(MeOD); δ 8.52 (s, 1H), 8.26 (d, J=9.9 Hz, 1H), 7.46 (s, 1H), 7.42 (s, 2H), 7.41 - 7.35 (m, 7H), 7.35 - 7.30 (m, 3H), 7.24 (dd, J=8.3, 8.3 Hz, 2H), 7.03 - 6.94 (m, 3H), 6.90 (d, J=8.6 Hz, 1H), 6.62 (d, J=9.6 Hz, 1H), 5.33 (dd, J=6.6, 6.6 Hz, 1H), 4.21 - 4.14 (m, 4H), 4.11 - 4.03 (m, 4H), 3.80 (s, 2H), 3.57 (dd, J=6.8, 6.8 Hz, 2H), 3.14 - 3.04 (m, 4H), 2.37 (dd, J=11.9, 11.9 Hz, 2H), 2.12 - 2.06 (m, 2H), 1.77 - 1.70 (m, 1H), 1.68 - 1.60 (m, 2H), 1.44 (t, J=6.8 Hz, 3H), 1.37 - 1.27 (m, 2H).	mono-formate
32E	2.55	11	(MeOD); δ 8.39 (s, 2H), 8.33 (d, J=9.9 Hz, 1H), 7.85 - 7.81 (m, 2H), 7.69 (d, J=8.1 Hz, 1H), 7.45 (s, 5H), 7.41 - 7.30 (m, 5H), 7.28 - 7.22 (m, 2H), 7.03 - 6.95 (m, 3H), 6.90 (dd, J=2.4, 8.5 Hz, 1H), 6.63 (d, J=9.9 Hz, 1H), 5.32 (dd, J=4.3, 8.6 Hz, 1H), 4.16 - 4.10 (m, 6H), 4.05 (t, J=6.1 Hz, 2H), 3.57 (t, J=6.9 Hz, 2H), 3.35 (br s, 2H), 3.10 - 3.00 (m, 2H), 2.86 - 2.77 (m, 2H), 2.13 - 2.06 (m, 2H), 1.95 - 1.88 (m, 1H), 1.76 (d, J=13.4 Hz, 2H), 1.49 - 1.38 (m, 2H).	formate
32F	2.51	11	(MeOD); δ 8.43 (s, 2H), 8.31 (d, J=9.6 Hz, 1H), 7.45 (s, 5H), 7.42 - 7.24 (m, 9H), 7.05 - 6.96 (m, 3H), 6.92 (dd, J=2.5, 8.3 Hz, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.42 - 5.37 (m, 1H), 4.27 (s, 2H), 4.14 - 4.04 (m, 9H), 3.58 (dd, J=6.8, 6.8 Hz, 2H), 3.32 - 3.27 (m, 2H), 3.14 (d, J=6.8 Hz, 2H), 2.83 - 2.73 (m, 2H), 2.13 - 2.04 (m, 2H), 1.95 - 1.86 (m, 1H), 1.76 (d, J=12.6 Hz, 2H), 1.48 - 1.38 (m, 2H).	formate
32G	2.55	11	(MeOD); δ 8.36 (s, 2H), 8.14 (d, J=9.9 Hz, 1H), 7.39 (s, 1H), 7.35 - 7.18 (m, 12H), 7.15 - 7.10 (m, 2H), 6.92 - 6.83 (m, 3H), 6.78 (dd, J=2.1, 8.0 Hz, 1H), 6.52 (d, J=9.9 Hz, 1H), 5.23 (dd, J=4.5, 8.6 Hz, 1H), 4.71 - 4.64 (m, 1H), 4.12 (d, J=2.8 Hz, 2H), 4.00 - 3.91 (m, 4H), 3.82 (s, 2H), 3.45 (dd, J=6.9, 6.9 Hz, 2H), 3.09 - 3.02 (m, 4H), 2.43 (dd, J=11.0, 12.0 Hz, 2H), 2.01 - 1.95 (m, 2H), 1.74 - 1.67 (m, 1H), 1.58 (d, J=13.1 Hz, 2H), 1.28 - 1.23 (m, 8H).	formate

(continued)

32H	2.49	11	(MeOD); δ 8.46 (s, 2H), 8.32 (d, J=9.9 Hz, 1H), 7.58 (d, J=1.3 Hz, 1H), 7.46 - 7.23 (m, 14H), 7.04 - 6.96 (m, 3H), 6.92 (dd, J=2.1, 8.0 Hz, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.34 (dd, J=4.2, 9.0 Hz, 1H), 4.14 - 4.04 (m, 8H), 3.56 (dd, J=6.7, 6.7 Hz, 2H), 3.28 - 3.22 (m, 2H), 3.06 - 3.01 (m, 2H), 2.75 - 2.65 (m, 2H), 2.09 - 2.05 (m, 2H), 1.91 - 1.83 (m, 1H), 1.77 - 1.71 (m, 2H), 1.45 - 1.35 (m, 2H).	formate
32I	2.48	11	(MeOD) d 8.46 (s, 2H), 8.30 (d, J=9.9 Hz, 1H), 7.43 - 7.31 (m, 12H), 7.28 - 7.23 (m, 2H), 7.04 - 6.96 (m, 3H), 6.91 (dd, J=2.1, 8.0 Hz, 1H), 6.65 (d, J=9.9 Hz, 1H), 5.36 (t, J=6.6 Hz, 1H), 4.17 (s, 2H), 4.13 - 4.04 (m, 4H), 4.00 (s, 2H), 3.89 (s, 3H), 3.59 (dd, J=6.7, 6.7 Hz, 2H), 3.23 (d, J=12.1 Hz, 2H), 3.10 (d, J=6.1 Hz, 2H), 2.67 - 2.58 (m, 2H), 2.10 - 2.05 (m, 2H), 1.90 - 1.82 (m, 1H), 1.72 (d, J=12.6 Hz, 2H), 1.43 - 1.32 (m, 2H).	formate
32J	2.48	11	(MeOD); δ 8.50 (s, 2H), 8.30 (d, J=9.0 Hz, 1H), 7.45 (s, 1H), 7.41 - 7.32 (m, 12H), 7.09 (s, 1H), 7.04 - 6.96 (m, 3H), 6.92 (dd, J=2.0, 8.1 Hz, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.34 (dd, J=5.2, 8.0 Hz, 1H), 4.13 - 4.06 (m, 6H), 3.94 (s, 2H), 3.87 (s, 3H), 3.57 (t, J=6.8, 6.8 Hz, 2H), 3.22 - 3.09 (m, 2H), 3.08 - 3.04 (m, 2H), 2.57 (dd, J=11.7 Hz, 2H), 2.14 - 2.05 (m, 2H), 1.88 - 1.80 (m, 1H), 1.68 (d, J=12.1 Hz, 2H), 1.42 - 1.32 (m, 2H).	formate
32K	2.47	11	(MeOD); δ 8.66 (d, J=1.5 Hz, 1H), 8.42 (s, 2H), 8.31 (d, J=9.9 Hz, 1H), 8.10 (d, J=8.1 Hz, 1H), 8.02 (dd, J=2.1, 8.0 Hz, 1H), 7.44 - 7.37 (m, 6H), 7.37 - 7.22 (m, 6H), 7.10 (dd, J=2.0, 2.0 Hz, 1H), 7.02 (d, J=8.3 Hz, 1H), 6.97 - 6.91 (m, 2H), 6.63 (d, J=9.9 Hz, 1H), 5.34 (dd, J=4.5, 8.3 Hz, 1H), 4.18 (s, 2H), 4.13 - 4.05 (m, 6H), 3.62 (t, J=6.6, 6.6 Hz, 2H), 3.29 - 3.23 (m, 2H), 3.12 - 3.06 (m, 2H), 2.69 (dd, J=11.5, 11.5 Hz, 2H), 2.14 - 2.06 (m, 2H), 1.86 - 1.80 (m, 1H), 1.72 (d, J=13.6 Hz, 2H), 1.43 - 1.33 (m, 2H).	formate
32L	2.49	11	(MeOD); δ 8.50 (s, 1H), 8.34 (d, J=9.9 Hz, 1H), 7.47 - 7.42 (m, 2H), 7.42 - 7.37 (m, 6H), 7.36 - 7.21 (m, 6H), 7.02 - 6.96 (m, 3H), 6.92 - 6.89 (m, 1H), 6.64 (d, J=9.9 Hz, 1H), 5.27 (dd, J=4.0, 8.6 Hz, 1H), 4.13 - 4.01 (m, 6H), 3.95 (s, 2H), 3.57 (t, J=6.8 Hz, 2H), 3.22 - 3.15 (m, 2H), 2.99 - 2.86 (m, 2H), 2.57 (dd, J=10.0, 12.5 Hz, 2H), 2.12 - 2.05 (m, 2H), 1.88 - 1.79 (m, 1H), 1.69 (d, J=12.4 Hz, 2H), 1.42 - 1.32 (m, 2H).	mono-formate

(continued)

32M	2.53	11	(MeOD) d 8.48 (s, 1H), 8.33 (d, J=9.9 Hz, 1H), 8.17 (d, J=1.3 Hz, 1H), 8.04 (dd, J=1.4, 8.0 Hz, 1H), 7.80 (d, J=8.1 Hz, 1H), 7.43 - 7.36 (m, 7H), 7.35 - 7.29 (m, 3H), 7.27 - 7.22 (m, 2H), 7.02 - 6.96 (m, 3H), 6.91 - 6.88 (m, 1H), 6.63 (d, J=9.9 Hz, 1H), 5.27 (dd, J=4.2, 8.5 Hz, 1H), 4.14 - 4.09 (m, 4H), 4.05 (t, J=6.0 Hz, 2H), 4.01 (s, 2H), 3.58 (t, J=6.8 Hz, 2H), 3.24 (d, J=12.1 Hz, 2H), 3.01 - 2.89 (m, 2H), 2.68 - 2.60 (m, 2H), 2.13 - 2.05 (m, 2H), 1.90 - 1.82 (m, 1H), 1.71 (d, J=12.9 Hz, 2H), 1.44 - 1.34 (m, 2H).	mono-formate
33	7.28	7	(DMSO-d ₆ , 90 °C); δ 8.23 (s, 2H), 8.19 (d, J=10.1 Hz, 1H), 7.46 (d, J=8.3 Hz, 2H), 7.38 - 7.25 (m, 13H), 7.10 (d, J=8.2 Hz, 1H), 6.98 - 6.91 (m, 4H), 6.53 (d, J=9.9 Hz, 1H), 5.19 - 5.13 (m, 1H), 5.09 (s, 2H), 4.53 - 4.40 (m, 1H), 4.00 (d, J=6.1 Hz, 2H), 3.62 - 3.50 (m, 1H), 3.40 (s, 2H), 3.06 - 2.95 (m, 1H), 2.84 (d, J=6.4 Hz, 2H), 2.76 - 2.63 (m, 4H), 1.88 - 1.61 (m, 5H), 1.56 - 1.43 (m, 3H), 1.17 - 1.09 (m, 4H).	Formate
33A	2.52	1	(DMSO-d ₆); δ 8.35 (s, 2H), 8.26 - 8.24 (m, 1H), 7.51 (d, J=8.1 Hz, 2H), 7.18 - 7.26 (m, 15 H), 7.19 - 7.12 (m, 1H), 7.05 - 6.96 (m, 3H), 6.61 - 6.59 (m, 1H), 5.31 (d, J=6.6 Hz, 1H), 5.14 (s, 2H), 4.54 - 4.53 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.62 - 3.58 (m, 1H), 3.23 (s, 2H), 3.04 - 2.78 (m, 8H), 1.93 - 1.84 (m, 2H), 1.81 - 1.76 (m, 1H), 1.59 - 1.52 (m, 6H), 1.24 - 1.13 (m, 4H).	Formate
34	7.39	7	(DMSO-d ₆ , 90 °C); δ 8.63 (t, J=7.2 Hz, 1H), 8.22 (s, 2H), 8.17 (d, J=9.9 Hz, 1H), 7.80 (d, J=8.3 Hz, 2H), 7.34 - 7.20 (m, 11H), 7.18 - 7.14 (m, 1H), 7.10 (d, J=8.3 Hz, 1H), 6.95 (d, J=8.2 Hz, 1H), 6.92 - 6.89 (m, 3H), 6.75 (d, J=8.3 Hz, 2H), 6.53 (d, J=9.9 Hz, 1H), 5.15 (t, J=6.3 Hz, 1H), 4.08 - 3.97 (m, 4H), 3.64 - 3.58 (m, 2H), 3.00 - 2.90 (m, 2H), 2.86 (d, J=6.5 Hz, 4H), 2.71 (d, J=11.4 Hz, 2H), 1.84 - 1.78 (m, 2H), 1.53 - 1.44 (m, 3H), 1.16 - 1.06 (m, 2H).	Formate
35	2.50	11	(DMSO--d ₆); δ 10.54 - 10.48 (m, 2H), 9.57 - 9.47 (m, 1H), 8.64 - 8.64 (m, 2H), 8.18 (d, J=9.9 Hz, 1H), 7.52 - 7.41 (m, 8H), 7.36 - 7.24 (m, 6H), 7.16 (d, J=8.2 Hz, 1H), 7.02 - 6.95 (m, 3H), 6.92 (d, J=7.7 Hz, 1H), 6.73 - 6.63 (m, 1H), 6.59 (d, J=9.9 Hz, 1H), 6.25 - 6.14 (m, 1H), 5.39 - 5.33 (m, 1H), 5.10 (s, 2H), 4.54 - 4.42 (m, 1H), 4.26 (dt, J=6.0, 18.4 Hz, 2H), 4.00 (d, J=6.4 Hz, 2H), 3.54 - 3.54 (m, 1H), 3.37 - 3.29 (m, 2H), 3.14 - 3.05 (m, 3H), 3.00 - 2.85 (m, 4H), 2.10 - 2.04 (m, 1H), 1.90 - 1.79 (m, 2H), 1.77 - 1.64 (m, 3H), 1.40 - 1.31 (m, 2H), 1.27 - 1.11 (m, 3H).	TFA

(continued)

36	2.38	11	(DMSO-d ₆ , 100°C); δ 8.24 (d, J=9.9 Hz, 1H), 7.53 - 7.49 (m, 5H), 7.44 - 7.25 (m, 6H), 7.20 (d, J=8.3 Hz, 1H), 7.07 - 7.01 (m, 3H), 6.93 - 6.89 (m, 1H), 6.61 (d, J=9.9 Hz, 1H), 5.43 (dd, J=4.7, 8.3 Hz, 1H), 4.75 (s, 2H), 4.28 (s, 2H), 4.14 - 4.10 (m, 3H), 3.40 - 3.30 (m, 2H), 3.23 - 3.19 (m, 2H), 3.05 - 2.88 (m, 6H), 2.14 - 2.05 (m, 1H), 2.00 - 1.76 (m, 6H), 1.59 - 1.47 (m, 2H), 1.33 - 1.18 (m, 2H).	TFA
36A	2.53	11	(DMSO-d ₆); δ 8.35 (s, 2H), 8.26 (d, J=9.9 Hz, 1H), 7.39 - 7.28 (m, 12H), 7.18 (d, J=8.3 Hz, 1H), 7.03 (d, J=8.1 Hz, 1H), 6.96 (d, J=8.3 Hz, 1H), 6.93 - 6.87 (m, 2H), 6.60 (d, J=9.9 Hz, 1H), 5.31 (dd, J=3.5, 8.8 Hz, 1H), 4.78 (d, J=4.3 Hz, 2H), 4.32 (d, J=12.1 Hz, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.83 (d, J=14.1 Hz, 1H), 3.23 (s, 2H), 3.05 - 2.86 (m, 5H), 2.78 (d, J=11.4 Hz, 2H), 1.93 - 1.85 (m, 2H), 1.69 (d, J=12.9 Hz, 2H), 1.58 - 1.48 (m, 7H), 1.23 - 1.12 (m, 3H), 1.00 - 0.97 (m, 1H).	Formate
37	2.48	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.83 (d, J=8.3 Hz, 2H), 7.52 - 7.48 (m, 6H), 7.44 - 7.23 (m, 9H), 7.05 (d, J=8.2 Hz, 1H), 6.99 - 6.94 (m, 2H), 6.91 (dd, J=2.3, 8.2 Hz, 1H), 6.70 (d, J=9.9 Hz, 1H), 5.42 (dd, J=5.5, 8.0 Hz, 1H), 4.29 (dd, J=6.6, 24.6 Hz, 2H), 4.12 - 4.00 (m, 4H), 3.56 (dd, J=6.9, 6.9 Hz, 2H), 3.50 - 3.35 (m, 4H), 3.31 - 3.28 (m, 2H), 3.20 - 3.10 (m, 2H), 2.97 (dd, J=10.7, 12.7 Hz, 2H), 2.11 - 2.04 (m, 2H), 1.94 (dd, J=3.1, 8.9 Hz, 1H), 1.83 (d, J=14.6 Hz, 2H), 1.47 - 1.40 (m, 2H).	TFA
38	2.46	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.83 (d, J=8.3 Hz, 2H), 7.52 - 7.45 (m, 6H), 7.44 - 7.39 (m, 3H), 7.39 - 7.25 (m, 5H), 7.06 - 6.93 (m, 4H), 6.70 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.6, 7.9 Hz, 1H), 4.26 (s, 2H), 4.16 - 4.09 (m, 4H), 3.75 (dd, J=5.7, 5.7 Hz, 2H), 3.50 - 3.35 (m, 5H), 3.18 - 3.10 (m, 2H), 2.96 (dd, J=12.5, 12.5 Hz, 2H), 2.00 - 1.78 (m, 4H), 1.50 - 1.35 (m, 2H).	TFA
39	2.41	11	(MeOD); δ 8.28 (d, J=9.9 Hz, 1H), 7.86 (d, J=8.2 Hz, 2H), 7.63 (d, J=8.4 Hz, 2H), 7.48 (s, 5H), 7.39 - 7.26 (m, 10H), 7.03 (d, J=8.3 Hz, 1H), 6.65 (dd, J=1.7, 9.9 Hz, 1H), 5.43 (dd, J=5.8, 7.8 Hz, 1H), 4.39 (s, 2H), 4.28 (s, 2H), 4.08 (d, J=3.5 Hz, 2H), 3.62 - 3.55 (m, 2H), 3.49 (d, J=11.8 Hz, 2H), 3.27 - 3.23 (m, 2H), 3.01 - 2.89 (m, 4H), 1.95 (s, 1H), 1.82 (d, J=14.1 Hz, 2H), 1.49 - 1.38 (m, 1H).	TFA

(continued)



40	2.47	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.78 (d, J=8.4 Hz, 2H), 7.48 (s, 5H), 7.43 - 7.35 (m, 5H), 7.35 - 7.25 (m, 7H), 7.05 (d, J=8.2 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.43 (dd, J=6.7, 6.7 Hz, 1H), 4.27 (s, 2H), 4.10 (dd, J=1.5, 6.3 Hz, 2H), 3.60 - 3.53 (m, 2H), 3.49 - 3.43 (m, 2H), 3.41-3.35 (m, 2H), 3.21 - 3.10 (m, 3H), 3.02 - 2.88 (m, 5H), 1.99 - 1.92 (m, 1H), 1.85 - 1.75 (m, 2H), 1.50 - 1.37 (m, 2H).	TFA
41	2.47	11	(DMSO-d ₆) δ 10.51 (s, 2H), 9.58 (s, 1H), 9.19 (d, J=27.2 Hz, 2H), 8.10 (d, J=9.9 Hz, 1H), 7.69 (d, J=7.3 Hz, 2H), 7.61 (d, J=8.3 Hz, 2H), 7.48 (s, 5H), 7.32 (s, 6H), 7.28 - 7.24 (m, 1H), 7.13 (d, J=8.0 Hz, 1H), 7.01 - 6.96 (m, 1H), 6.93 - 6.88 (m, 2H), 6.57 (d, J=9.7 Hz, 1H), 6.19 (s, 1H), 5.36 (d, J=10.4 Hz, 1H), 4.41 (t, J=8.6 Hz, 1H), 4.31 - 4.23 (m, 5H), 4.12 (d, J=7.2 Hz, 3H), 4.01 (d, J=6.3 Hz, 2H), 3.89 - 3.85 (m, 1H), 3.05 - 3.02 (m, 4H), 2.90 - 2.89 (m, 2H), 1.75 (d, J=14.8 Hz, 3H), 1.37 (d, J=13.3 Hz, 2H), 1.29 - 1.21 (m, 2H).	TFA
42	2.46	11	(MeOD) δ 8.37 (d, J=9.9 Hz, 1H), 7.69 - 7.62 (m, 2H), 7.50 (dd, J=4.0, 4.0 Hz, 6H), 7.43 (dd, J=3.9, 8.2 Hz, 3H), 7.39 - 7.27 (m, 5H), 7.05 (d, J=8.2 Hz, 1H), 7.03 - 6.91 (m, 3H), 6.70 (d, J=9.8 Hz, 1H), 5.45 - 5.40 (m, 1H), 4.53 - 4.48 (m, 1H), 4.28 (s, 4H), 4.15 - 4.09 (m, 5H), 3.49 - 3.35 (m, 3H), 3.29 (d, J=8.0 Hz, 2H), 3.17 - 3.09 (m, 4H), 3.01 - 2.94 (m, 2H), 2.13 - 1.72 (m, 4H), 1.50 - 1.38 (m, 2H).	TFA
42A	2.43	11	(MeOD); δ 8.53 (s, 1H), 8.20 (d, J=9.9 Hz, 1H), 7.52 (d, J=1.9 Hz, 1H), 7.41 - 7.24 (m, 14H), 7.13 (dd, J=2.1, 2.1 Hz, 1H), 7.09 (d, J=7.9 Hz, 1H), 7.04 - 7.01 (m, 2H), 6.63 (d, J=9.9 Hz, 1H), 5.37 (dd, J=6.5, 6.5 Hz, 1H), 4.68 (s, 2H), 4.23 (s, 2H), 4.08 (d, J=6.4 Hz, 2H), 3.89 (s, 3H), 3.77 (s, 2H), 3.14 (d, J=6.5 Hz, 2H), 3.05 (d, J=11.8 Hz, 2H), 2.39 - 2.31 (m, 2H), 1.79 - 1.71 (m, 1H), 1.61 (d, J=12.7 Hz, 2H), 1.38 - 1.26 (m, 2H).	mono-Formate
42B	2.49	11	(MeOD); δ 8.55 (s, 1H), 8.27 (d, J=9.9 Hz, 1H), 7.75 (d, J=6.4 Hz, 1H), 7.39 - 7.29 (m, 11H), 7.23 (d, J=8.7 Hz, 2H), 7.13 - 7.09 (m, 2H), 7.05 - 7.00 (m, 2H), 6.63 (d, J=9.8 Hz, 1H), 5.31 (dd, J=5.6, 7.5 Hz, 1H), 4.73 (s, 2H), 4.10 - 4.05 (m, 4H), 3.83 (s, 3H), 3.68 (s, 2H), 3.07 - 2.96 (m, 4H), 2.24 (dd, J=11.5, 11.5 Hz, 2H), 1.76 - 1.67 (m, 1H), 1.58 (d, J=12.0 Hz, 2H), 1.34 - 1.27 (m, 2H).	mono-Formate

(continued)

42C	2.57	11	(DMSO-d ₆ , 90 °C) δ 8.69 (t, J=5.9 Hz, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.88 (d, J=8.3 Hz, 2H), 7.46 (s, 5H), 7.38 - 7.24 (m, 12H), 7.16 (d, J=8.3 Hz, 1H), 7.02 (d, J=8.2 Hz, 2H), 6.98 - 6.93 (m, 2H), 6.58 (d, J=9.9 Hz, 1H), 5.36 (dd, J=4.4, 8.7 Hz, 1H), 5.03 (s, 2H), 4.51 (d, J=6.0 Hz, 2H), 4.19 - 4.15 (m, 2H), 4.08 - 4.03 (m, 2H), 3.35 - 3.29 (m, 4H), 3.24 - 3.18 (m, 3H), 2.86 - 2.86 (m, 2H), 1.94 - 1.82 (m, 1H), 1.73 - 1.72 (m, 2H), 1.42 - 1.42 (m, 2H).	TFA
42D	2.64	11	(DMSO-d ₆) δ 10.53 (m, 2H), 9.45 - 9.40 (m, 1H), 9.17 (m, 3H), 8.13 (d, J=9.9 Hz, 1H), 8.00 (d, J=8.1 Hz, 2H), 7.68 (d, J=8.3 Hz, 2H), 7.52 (m, 5H), 7.42 - 7.35 (m, 10H), 7.17 (d, J=8.1 Hz, 1H), 7.04 - 6.93 (m, 4H), 6.67 (s, 1H), 6.63 (dd, J=1.8, 9.9 Hz, 1H), 6.23 (d, J=1.0 Hz, 1H), 5.39 (d, J=8.8 Hz, 1H), 5.06 (s, 2H), 4.54 (d, J=5.6 Hz, 2H), 4.38 - 4.25 (m, 4H), 4.08 - 4.01 (m, 2H), 3.38 (m, 2H), 3.17 - 3.04 (m, 2H), 2.99 - 2.90 (m, 2H), 1.83 - 1.71 (m, 3H), 1.44 - 1.35 (m, 2H).	TFA
42E	2.60	11	(MeOD); δ 8.38 (s, 2H), 8.19 (d, J=9.9 Hz, 1H), 7.81 (d, J=7.9 Hz, 1H), 7.36 - 7.12 (m, 17H), 7.05 (d, J=8.0 Hz, 1H), 6.93 - 6.83 (m, 4H), 6.54 (d, J=9.8 Hz, 1H), 5.31 - 5.26 (m, 1H), 4.94 (s, 2H), 4.51 (s, 2H), 4.12 (s, 2H), 3.97 (dd, J=1.8, 6.6 Hz, 2H), 3.91 (s, 2H), 3.88 (s, 3H), 3.14 - 3.02 (m, 4H), 2.54 (dd, J=11.0, 11.0 Hz, 2H), 1.78 - 1.66 (m, 1H), 1.57 (d, J=12.2 Hz, 2H), 1.31 - 1.19 (m, 2H).	formate
42F	2.59	11	(MeOD); δ 8.51 (s, 1H), 8.28 (d, J=9.9 Hz, 1H), 7.58 (s, 1H), 7.56 - 7.24 (m, 16H), 7.05 - 6.92 (m, 5H), 6.66 (d, J=9.9 Hz, 1H), 5.45 - 5.37 (m, 1H), 5.05 (s, 2H), 4.61 (s, 2H), 4.60 (s, 1H), 4.31 (s, 2H), 4.09 (dd, J=1.3, 6.4 Hz, 2H), 3.98 (s, 5H), 3.82 (s, 1H), 3.23 - 3.16 (m, 4H), 2.64 - 2.55 (m, 2H), 1.87 - 1.79 (m, 1H), 1.67 (d, J=12.9 Hz, 2H), 1.37 (q, J=11.9 Hz, 2H).	mono-formate

Example 6


(S)-(1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate (Compound 43)

**Step 1; 4-(Chloromethyl)-N-(3,3-diethoxypropyl)benzamide**

A solution of 4-(chloromethyl)-benzoyl chloride (1.17 g, 6.21 mmol) in DCM (10 mL) was added with a solution of 3,3-diethoxypropan-1-amine (0.914 g, 6.21 mmol) and triethylamine (1.04 mL, 7.77 mmol) in DCM (10 mL). The reaction mixture was stirred at room temperature for 3 hours. The solvent was evaporated under reduced pressure and the residue dissolved in ethyl acetate. The organic phase was washed with 1M aqueous sodium hydroxide, brine and dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure to afford the title compound (1.83 g, 99%).

¹H NMR (400 MHz, DMSO-d₆); δ 8.45 (dd, J=5.5, 5.5 Hz, 1H), 7.83 (d, J=8.4 Hz, 2H), 7.52 (d, J=8.3 Hz, 2H), 4.81 (s, 2H), 4.56 (dd, J=5.5, 5.5 Hz, 1H), 3.59 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.45 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.29 (dd, J=6.0, 7.4 Hz, 2H), 1.79 (dd, J=6.4, 13.6 Hz, 2H), 1.12 (dd, J=7.0, 7.0 Hz, 6H).

Step 2; (2S)-(1-Benzylpiperidin-4-yl)methyl 2-(3-((4-((3-ethoxy-3-propoxypropyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



The title compound was prepared as described in Procedure 7 Step 4 with

4-(chloromethyl)-N-(3,3-diethoxypropyl)benzamide replacing *tert*-butyl
4-(bromomethyl)benzoate.

¹H NMR (400 MHz, DMSO-d₆); δ 8.43 (dd, J=5.5, 5.5 Hz, 1H), 7.83 (d, J=8.4 Hz, 2H), 7.47 (d, J=8.4 Hz, 2H), 7.26 – 7.20 (m, 11H), 6.97 - 6.90 (m, 3H), 6.60 (s, 1H), 5.11
5 (s, 2H), 4.56 (dd, J=5.5, 5.5 Hz, 1H), 3.98 (d, J=6.3 Hz, 2H), 3.58 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.48 - 3.38 (m, 5H), 2.73 - 2.67 (m, 2H), 1.85 - 1.76 (m, 3H), 1.52 - 1.44 (m, 3H), 1.12 (dd, J=7.0, 7.0 Hz, 6H).

Step 3; Compound (43)


The title compound (43) was prepared as described in Example 1 Step 9.


10

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
43	7.26	7	(DMSO-d ₆ , 90°C); δ 8.37 (dd, J=5.6, 5.6 Hz, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.85 (d, J=8.2 Hz, 2H), 7.51 - 7.44 (m, 8H), 7.38 - 7.24 (m, 6H), 7.16 (d, J=8.2 Hz, 1H), 7.04 (d, J=2.0 Hz, 1H), 7.02 - 6.95 (m, 3H), 6.56 (d, J=9.9 Hz, 1H), 5.34 (dd, J=4.5, 8.5 Hz, 1H), 5.12 (s, 2H), 4.18 (s, 2H), 4.07 (s, 2H), 3.42 - 3.37 (m, 4H), 3.18 - 3.08 (m, 4H), 2.88 - 2.88 (m, 2H), 2.01 - 1.93 (m, 2H), 1.87 (s, 1H), 1.74 (d, J=13.2 Hz, 2H), 1.42 - 1.42 (m, 2H).	TFA

Example 7


(*R*)-(1-Benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((*R*)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbonyl)benzyl)oxy)phenyl)-2-phenylacetate (Compound
15 44)



Step 1; 4-(Chloromethyl)-N-(3,3-diethoxypropyl)benzamide

A solution of 4-(chloromethyl)-benzoyl chloride (1.17 g, 6.21 mmol) in DCM (10 mL) was added with a solution of 3,3-diethoxypropan-1-amine (0.914 g, 6.21 mmol) and triethylamine (1.04 mL, 7.77 mmol) in DCM (10 mL). The reaction mixture was stirred at room temperature for 3 hours. The solvent was evaporated under reduced pressure and the residue dissolved in ethyl acetate. The organic phase was washed with 1M aqueous sodium hydroxide, brine and dried over anhydrous magnesium sulfate and the filtrate was evaporated under reduced pressure to afford the title compound (1.83 g, 99%).

¹H NMR (400 MHz, DMSO-d₆); δ 8.45 (dd, J=5.5, 5.5 Hz, 1H), 7.83 (d, J=8.4 Hz, 2H), 7.52 (d, J=8.3 Hz, 2H), 4.81 (s, 2H), 4.56 (dd, J=5.5, 5.5 Hz, 1H), 3.59 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.45 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.29 (dd, J=6.0, 7.4 Hz, 2H), 1.79 (dd, J=6.4, 13.6 Hz, 2H), 1.12 (dd, J=7.0, 7.0 Hz, 6H).

Step 2; (2R)-(1-Benzylpiperidin-4-yl)methyl 2-(3-((4-((3-ethoxy-3-propoxypropyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate

The title compound was prepared as described in Procedure 7 with (*R*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid replacing (*S*)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid in Step 1 and 4-(chloromethyl)-N-(3,3-diethoxypropyl)benzamide replacing *tert*-butyl 4-(bromomethyl)benzoate in Step 4.

¹H NMR (400 MHz, DMSO-d₆); δ 8.43 (dd, J=5.5, 5.5 Hz, 1H), 7.83 (d, J=8.4 Hz, 2H), 7.47 (d, J=8.4 Hz, 2H), 7.26 – 7.20 (m, 11H), 6.97 - 6.90 (m, 3H), 6.60 (s, 1H), 5.11 (s, 2H), 4.56 (dd, J=5.5, 5.5 Hz, 1H), 3.98 (d, J=6.3 Hz, 2H), 3.58 (ddd, J=7.1, 9.5, 14.1 Hz, 2H), 3.48 - 3.38 (m, 5H), 2.73 - 2.67 (m, 2H), 1.85 - 1.76 (m, 3H), 1.52 - 1.44 (m, 3H),

1.12 (dd, J=7.0, 7.0 Hz, 6H).


Step 3; (Compound 44)

The title compound (44) was prepared as described in Example 1 Step 9.

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
44	7.25	6	(DMSO-d ₆ , 90°C); δ 8.36 (dd, J=5.5, 5.5 Hz, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.85 (d, J=7.7 Hz, 2H), 7.48 (m, J=5.5 Hz, 8H), 7.38 - 7.24 (m, 6H), 7.16 (d, J=8.2 Hz, 1H), 7.05 (s, 1H), 7.02 - 6.96 (m, 3H), 6.57 (d, J=9.8 Hz, 1H), 5.34 (dd, J=4.5, 8.6 Hz, 1H), 5.12 (s, 2H), 4.22 (s, 2H), 4.06 (s, 2H), 3.43 - 3.37 (m, 4H), 3.18 - 3.08 (m, 4H), 2.87 (s, 2H), 2.05 - 1.94 (m, 2H), 1.88 (s, 1H), 1.77 - 1.74 (m, 2H), 1.42 - 1.42 (m, 2H).	TFA


5 Example 8

(S)-(1-Benzylpiperidin-4-yl)methyl 2-(3-((3-(benzyl(3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbonyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (Compound 45)



10


Step 1; N-(2-(1,3-Dioxolan-2-yl)ethyl)-N-benzyl-3-(chloromethyl)benzamide



The title compound was prepared as described in Example 5 Step 1 with *N*-benzyl-2-(1,3-dioxolan-2-yl)ethanamine replacing 3,3-diethoxypropan-1-amine and

3-(chloromethyl)benzoyl chloride replacing 4-(chloromethyl)benzoyl chloride.

Step 2; (S)-3-((2-(1,3-Dioxolan-2-yl)ethyl)(benzyl)carbamoyl)benzyl 2-(3-((2-(1,3-dioxolan-2-yl)ethyl)(benzyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



The title compound was prepared as described in Example 4 Step 1 with N-(2-(1,3-dioxolan-2-yl)ethyl)-N-benzyl-3-(chloromethyl)benzamide replacing 2-(4-chlorobutyl)-1,3-dioxolane.


Step 3; (Compound 45)

The title compound (45) was prepared as described in Example 4 Step 2 through Step 4.


N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
45	7.50	6	(DMSO-d ₆ , 90°C); δ 8.15 (d, J=9.4 Hz, 1H), 7.45 (d, J=8.0 Hz, 8H), 7.38 - 7.22 (m, 12H), 7.14 (d, J=8.2 Hz, 1H), 7.04 - 6.92 (m, 4H), 6.55 (d, J=9.9 Hz, 1H), 5.30 (dd, J=4.8, 7.8 Hz, 1H), 5.07 (s, 2H), 4.57 (s, 2H), 4.07 - 4.07 (m, 2H), 3.43 - 3.40 (m, 2H), 3.16 - 3.08 (m, 7H), 2.99 (s, 2H), 1.96 (dd, J=7.5, 7.5 Hz, 3H), 1.73 - 1.69 (m, 2H), 1.40 - 1.40 (m, 2H).	TFA

Example 9

(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-phenylacetate diformate (Compound 46)




5

Step 1; Methyl 4-(4-(1,3-dioxolan-2-yl)butoxy)benzoate

A solution of methyl 4-hydroxybenzoate (1 g, 6.57 mmol) in DMF (0.509 ml) K_2CO_3 (1.045 g, 7.56 mmol) and 2-(4-bromobutyl)-1,3-dioxolane (1.649 g, 7.89 mmol) were added. The mixture was stirred at RT overnight to get to completion. The mixture was poured onto saturated $NaCl_{aq}$ (25 ml) and the aqueous phase was extracted with DCM (10ml x 3). The combined organic layers were dried over Na_2SO_4 and evaporated in vacuo to give a crude that was purified by flash chromatography (silica gel, eluents: 100% n-Hex to 100% AcOEt) to afford methyl 4-(4-(1,3-dioxolan-2-yl)butoxy)benzoate (1.6 g, 5.71 mmol, 87 % yield).

15

UPLC-MS: 1.07 min, 249 [(M+H)-MeOH]⁺, method 2.

Step 2; 4-(4-(1,3-dioxolan-2-yl)butoxy)benzoic acid


20

A solution of methyl 4-(4-(1,3-dioxolan-2-yl)butoxy)benzoate (1.13 g, 4.03 mmol) in THF (24 ml, 293 mmol), LiOH 1M (1.209 ml, 1.209 mmol) was added with

and the mixture stirred at RT overnight. LiOH 1M (1.5 ml) was added and the mixture stirred at RT over the weekend to get to completion. NaHCO₃ was added to pH=6-7 and the aqueous phase extracted with AcOEt (10 ml x 3). The organic phase was dried over Na₂SO₄ and evaporated in vacuo to give a crude that was used without further purifications for next steps.

UPLC-MS: 0.86 min, 267 [(M+H)]+, method 2.


Step 3; (R)-quinuclidin-3-yl 2-oxo-2-phenylacetate



A solution of 2-oxo-2-phenylacetic acid (1 g, 6.66 mmol) in DCM (25 ml, 389 mmol), oxalyl chloride (0.816 ml, 9.33 mmol) was added dropwise at RT, and the mixture stirred for 20 min. Then (R)-quinuclidin-3-ol (0.847 g, 6.66 mmol) was added and the mixture stirred at RT for 2hrs. DCM (50 ml) was added and the organic phase washed with NaHCO₃ saturated solution (30 ml X 3). The organic layer was evaporated in vacuo to give (R)-quinuclidin-3-yl 2-oxo-2-phenylacetate (870 mg, 3.36 mmol, 50.4 % yield) as an oily crude that was used without further purifications for next steps.

UPLC-MS: 0.37 min, 260 [M+H]⁺, method 2.

Step 4; (R)-quinuclidin-3-yl 2-(3-((tert-butoxycarbonyl)amino)phenyl)-2-hydroxy-2-phenylacetate




A solution of (R)-quinuclidin-3-yl 2-oxo-2-phenylacetate (960 mg, 3.70 mmol) in THF (20 ml, 244 mmol) at -15deg, was added dropwise with (3-(bis(trimethylsilyl)amino)phenyl)magnesium chloride 1M in THF (3.70 ml, 3.70 mmol)

and the mixture stirred for 15 min at 0deg, then allowed to warm to RT and stirred for 3 hrs at RT. The reaction was quenched by addition of MeCN (2ml) and partitioned between AcOEt (200mL) and water (200mL). Organic layer was washed twice with water, once with sat NaCl, dried over Na₂SO₄ and dried under reduced pressure to give the desylilated
5 compound. The crude was dissolved in DCM (20 ml) and di-tert-butyl dicarbonate (1.117 ml, 4.81 mmol), DMAP (90 mg, 0.740 mmol) and N-ethyl-N-isopropylpropan-2-amine (0.838 ml, 4.81 mmol) were added and the mixture stirred at RT overnight. DCM was evaporated in vacuo to give a crude that was purified by flash chromatography (silica gel, eluents: DCM to 1/1 DCM/EtOH) to afford (R)-quinuclidin-3-yl 2-(3-((tert-butoxycarbonyl)amino)phenyl)-2-hydroxy-2-phenylacetate (580 mg, 1.282 mmol, 34.6 %
10 yield)

UPLC-MS: 0.64 min, 454 [M+H]⁺, method 2.

Step 5; (R)-quinuclidin-3-yl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate dihydrochloride




15

A solution of (R)-quinuclidin-3-yl 2-(3-((tert-butoxycarbonyl)amino)phenyl)-2-hydroxy-2-phenylacetate (500 mg, 1.105 mmol) in acetonitrile (2 ml) was added with HCl 4M in dioxane (1 ml, 32.9 mmol) and the mixture stirred at RT for 2hrs. The solvent was evaporated in vacuo to give (R)-quinuclidin-3-yl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate dihydrochloride (493 mg, 1.160 mmol, 105 % yield) that was used without
20 further purifications for next steps.

UPLC-MS: 0.50 min, 426 [M+H]⁺, method 1.

25


Step 6; (R)-quinuclidin-3-yl 2-(3-(4-(4-(1,3-dioxolan-2-yl)butoxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate



A solution of 4-(4-(1,3-dioxolan-2-yl)butoxy)benzoic acid (324 mg, 1.215 mmol) in DMF (5 ml), HATU (504 mg, 1.326 mmol) and DIPEA (212 μ l, 1.215 mmol) were added. The mixture was stirred at RT for 30 min, then (R)-quinuclidin-3-yl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate dihydrochloride (470 mg, 1.105 mmol) in DMF (5 ml) and DIPEA (386 μ l, 2.210 mmol) were added to the mixture and stirred at RT on. The mixture was purified by reverse phase chromatography (C-18 silica gel, eluents: 100% H₂O/Acetonitrile 95/5+HCOOH 0.1% to 100% H₂O/ACN 5/95+HCOOH 0.1%) to afford (R)-quinuclidin-3-yl 2-(3-(4-(4-(1,3-dioxolan-2-yl)butoxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate (272 mg, 0.453 mmol, 41.0 % yield).

UPLC-MS: 0.72 min, 601 [M+H]⁺, method 2.


Step 7; (R)-quinuclidin-3-yl 2-hydroxy-2-(3-(4-((5-oxopentyl)oxy)benzamido)phenyl)-2-phenylacetate



A solution of (R)-quinuclidin-3-yl 2-(3-(4-(4-(1,3-dioxolan-2-yl)butoxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate (272 mg, 0.453 mmol) in Acetonitrile (5.723 ml) HCl 2M (5.6 ml, 11.32 mmol) was added and the mixture stirred at RT for 2hrs. NaHCO₃ saturated solution was added to pH=8 and the aqueous phase
 5 extracted with AcOEt (50 ml X 2). The organic phase was dried over Na₂SO₄ and evaporated in vacuo to give the title compound (195 mg, 0.350 mmol, 77 % yield) that was used without further purifications for next steps.

UPLC-MS: 0.67 min, 557 [M+H]⁺, method 2.

Step 8; (R)-quinuclidin-3-yl 2-(3-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyl
 10 **dimethylsilyl)oxy)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate**




A solution of (R)-quinuclidin-3-yl 2-hydroxy-2-(3-(4-((5-oxopentyl)oxy)benzamido)phenyl)-2-phenylacetate (195 mg, 0.350 mmol) in 1/1 solution DCM/EtOH (4 ml) (R)-5-(2-amino-1-((tert-butyl dimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (223 mg, 0.525 mmol) and acetic acid (0.040 ml, 0.701 mmol) were added. Then a spatula of sodium sulfate was added and the mixture stirred at RT for 15 min. sodium triacetoxyborohydride (371 mg, 1.752 mmol) was added
 15 to the mixture and stirred at RT for 2hrs. HCl 2M (2ml) was added, the mixture filtered and evaporated in vacuo to give a crude that was purified by reverse phase chromatography (C-18 silica gel, eluents: 100% H₂O/Acetonitrile 95/5+HCOOH 0.1% to 100% H₂O/ACN 5/95+HCOOH 0.1%) to afford (R)-quinuclidin-3-yl 2-(3-(4-((5-(((R)-2-(8-(benzyloxy)-2-

oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyl dimethylsilyl)oxy)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate (150 mg, 0.155 mmol, 44.4 % yield).

UPLC-MS: 1.43 min, 966 [M+H]⁺, method 2.

5 **Step 9; (R)-quinuclidin-3-yl 2-(3-(4-((5-(((R)-2-((tert-butyldimethylsilyl)oxy)-**
2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-
yl)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate



(R)-quinuclidin-3-yl 2-(3-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-

10 dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyloxy)ethylamino)pentyl)oxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate (150 mg, 0.155 mmol) and Pd-C 5% Enghelhart (50% water) (165 mg, 1.554 μ mol) were dissolved in MeOH (5 ml) and stirred at RT under H₂ atmosphere for 2hrs. The reaction was filtered over a small pad of celite and the filtrate dried under reduced
15 pressure to give the title compound (70 mg, 0.080 mmol, 51.5 % yield).

UPLC-MS: 1.17 min, 876 [M+H]⁺, method 4.

Step 10 (Compound 46)


A solution of (R)-quinuclidin-3-yl 2-(3-(4-((5-(((R)-2-((tert-butyl)dimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-hydroxy-2-phenylacetate (70 mg, 0.080 mmol) in THF (2 ml) HCl 4M in dioxane (800 μ L, 3.20 mmol) was added and the mixture stirred at RT for 6hrs , then kept at -4 deg overnight and then stirred again at RT for 6 hrs. Et₂O (20ml) was added and removed under vacuum twice, then the residual solvent was evaporated under reduced pressure without warming to give a crude that was

purified via reverse phase preparative HPLC to afford the title compound (46) (12 mg, 0.014 mmol, 17.59 % yield).


N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
46	3.99	761,0	5	(MeOH-d ₄) δ 8.50 (br. s., 2 H), 8.36 (d, J=9.87 Hz, 1 H), 7.89 (dd, J=8.88, 7.23 Hz, 2 H), 7.64 - 7.16 (m, 9 H), 7.01 (dd, J=14.63, 8.39 Hz, 3 H), 6.68 (d, J=9.87 Hz, 1 H), 5.46 - 5.34 (m, 1 H), 5.16 (br. s., 1 H), 4.10 (t, J=6.08 Hz, 2 H), 3.54 - 3.44 (m, 1 H), 3.26 - 2.73 (m, 8 H), 2.34 - 2.12 (m, 1 H), 2.02 - 1.73 (m, 5 H), 1.72 - 1.46 (m, 6 H)	Formate

Example 10

5 (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate diformate (Compound 47)



Step 1; methyl 2-(3-formylphenyl)-2-hydroxy-2-phenylacetate



10


A solution of methyl 2-oxo-2-phenylacetate (4.9 mL, 34.7 mmol) in 133 mL of dry THF were added dropwise, at -20°C, 40 mL of 1M (3-(diethoxymethyl)phenyl)magnesium bromide in THF (40.0 mmol) and the mixture stirred overnight at RT. Reaction was quenched by the addition of saturated NH₄Cl_{aq} and extracted with EtOAc. The organic layer was washed with aqueous saturated NaCl_{aq} and evaporated under reduced pressure;

15

the residue was dissolved in a mixture of 20 mL 2M HCl_{aq} and 20 mL of MeCN. The reaction was stirred for 2h at RT, then diluted with AcOEt and 1M HCl_{aq}, organic layer washed with saturated NaCl_{aq} and dried over Na₂SO₄. After evaporation, the residue was purified by flash chromatography (silica gel, eluents: hexane to 6/4 AcOEt/hexane) to afford the title compound (6.8 g, 25.2 mmol, 75 % yield).

UPLC-MS: 0.89 min, 253.1 [(M+H)-H₂O]⁺, method 1. a20140911_06


Step 2; 3-(1-hydroxy-2-methoxy-2-oxo-1-phenylethyl)benzoic acid



Methyl 2-(3-formylphenyl)-2-hydroxy-2-phenylacetate (3.8 g, 14.06 mmol), K₂H₂PO₄ (3.83 g, 28.1 mmol), 2-methyl-2-butene (14.89 ml, 141 mmol) and NaClO₂ (2.54 g, 28.1 mmol) were reacted in 36 mL of t-BuOH/water 1/1 from 0°C to RT for 4h. Reaction was quenched by the addition of 1M HCl_{aq} until pH 6-7, diluted with water and extracted with AcOEt. Organic layer was dried over Na₂SO₄ and evaporated to dryness. The crude was submitted to flash chromatography (silica gel, eluents: 100% DCM + 0.2% AcOH to 80/20 DCM/EtOH + 0.2% AcOH) to afford the title compound (4.1 g). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.79 min, 269.2 [(M+H)-H₂O]⁺, method 1.

Step 3; methyl 2-hydroxy-2-(3-((4-oxobutyl)carbamoyl)phenyl)-2-phenylacetate




3-(1-hydroxy-2-methoxy-2-oxo-1-phenylethyl)benzoic acid 2 (2.6 g, 9.08 mmol) and HATU (3.45 g, 9.08 mmol) were reacted for 30 min at 0°C, then 4,4-diethoxybutan-1-amine (2.354 ml, 13.62 mmol) and DIEA (2.379 ml, 13.62 mmol) were added and the

mixture stirred at RT for 2h. Reaction was quenched by the addition of 100 mL of 1M HCl_{aq} and stirred for 1h RT, then extracted with AcOEt, organic layer dried over Na₂SO₄ and evaporated to dryness. The crude oil was purified by flash chromatography (silica gel, eluent: hexane to AcOEt) to give the title compound (2 g, 5.63 mmol, 62.0 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.73 min, 356.1 [M+H]⁺, method 1.

Step 4; methyl 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate




1(R)-5-(2-amino-1-((tert-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (2.390 g, 5.63 mmol) and methyl 2-hydroxy-2-(3-((4-oxobutyl)carbamoyl)phenyl)-2-phenylacetate (2 g, 5.63 mmol) were reacted for 10 min at RT in 30 mL of DCM with a spatula tip of Na₂SO₄, then 0.3 mL of AcOH and NaB(OAc)₃H (2.39 g, 11.30 mmol) were added and the mixture stirred at RT for 2h. The reaction was filtered, dried to half of the initial volume and diluted with 10 mL of THF and 10 mL of saturated NaHCO₃ aq and added with Boc₂O (2.46 g, 11.30 mmol). The mixture was stirred vigorously for 0.5h at RT, then diluted with AcOEt and washed with saturated NaHCO₃ aq, saturated NaCl_{aq}, dried over Na₂SO₄ and evaporated to dryness. The crude was purified by flash chromatography (silica gel, DCM to AcOEt) to afford the title compound (0.9 g, 1.042 mmol, 18.51 % yield).

UPLC-MS: 1.56 min, 864.5 [M+H]⁺, method 1.

Step 5; 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-


((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid



Methyl 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (0.9 g, 1.042 mmol) and LiOH (0.125 g, 5.21 mmol) were reacted in 7 mL of THF/water 1/1 for 1h at RT. The reaction was quenched by the addition of 0.1M HCl_{aq} and extracted with AcOEt, organic layer washed twice with 0.1M HCl_{aq}, once with saturated NaCl_{aq}, dried over Na₂SO₄ and evaporated to dryness to give the title compound (869 mg, 1.022 mmol, 98 % yield).

UPLC-MS: 1.50 min, 850.3 [M+H]⁺, method 1.

Step 6; benzyl 4-(hydroxymethyl)piperidine-1-carboxylate




Piperidin-4-ylmethanol (5.2 g, 45.1 mmol) was dissolved in 60 mL of THF and 60 mL of saturated NaHCO₃ aq and then benzyl (2,5-dioxopyrrolidin-1-yl) carbonate (11.25 g, 45.1 mmol) added portionwise. The mixture was stirred for 1h at RT, partitioned between AcOEt and water, washed twice with 1M HCl_{aq} and saturated NaCl_{aq}. Organic layer was dried over Na₂SO₄, evaporated under reduced pressure to afford the title compound (10.77 g, 43.2 mmol, 96 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.80 min, 250.2 [M+H]⁺, method 1.

Step 7; benzyl 4-((2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-


yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetoxymethyl)piperidine-1-carboxylate



2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid (400 mg, 0.471 mmol) and CDI (229 mg, 1.412 mmol) were reacted for 30 min at RT, then benzyl 4-(hydroxymethyl)piperidine-1-carboxylate (469 mg, 1.882 mmol) added and the reaction stirred for 2h at 60°C, then RT overnight. The crude was partitioned between saturated NaHCO₃ aq and AcOEt, organic layer dried over Na₂SO₄ and evaporated to dryness. The crude was purified by flash chromatography (silica gel, eluents: DCM to AcOEt) to afford the title compound (110 mg, 0.102 mmol, 21.62 % yield).

UPLC-MS: 1.68 min, 1082.6 [M+H]⁺, method 1.

Step 8; piperidin-4-ylmethyl 2-(3-((4-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate formate




Benzyl 4-((2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetoxymethyl)piperidine-1-carboxylate (110 mg, 0.102 mmol) was

dissolved in 1 mL of MeOH, added of 7.80 μ L of HCOOH, Pd-C 10% wet (10.83 mg, 5.09 μ mol) and hydrogenated for 3h at RT under balloon pressure of hydrogen. Reaction mixture was filtered and reduced to dryness to give the title compound (120 mg). The product obtained was used in the following steps without further purifications.

5 UPLC-MS: 0.94 min, 857.6 [M+H]⁺, method 2.

Step 9; (1-benzylpiperidin-4-yl)methyl 2-(3-((4-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



10 piperidin-4-ylmethyl 2-(3-((4-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate formate (92 mg, 0.102 mmol) and benzaldehyde (13.45 μ L, 0.132 mmol) were dissolved in 1 mL of DCM and stirred for 1h at RT, then 7.58 μ L of AcOH and Na(OAc)₃H (28.1 mg, 0.132 mmol) were and the mixture stirred overnight at RT. Further 1 eq. of aldehyde and Na(OAc)₃H were needed to bring reaction to completion. Reaction mixture was partitioned between AcOEt and saturated NaHCO₃ aq, washed twice with saturated NaHCO₃ aq and saturated NaCl_{aq}, organic layer dried with Na₂SO₄ and evaporated to dryness to give the title compound (100 mg). The product obtained was used in the following steps without further purifications.

20 purifications.

UPLC-MS: 1.04 min, 946.6 [M+H]⁺, method 2.

Step 10; (Compound 47)


(1-benzylpiperidin-4-yl)methyl 2-(3-((4-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (100 mg, 0.106 mmol) was dissolved in 0.5 mL of MeCN and added with 2 mL of 5 M HCl_{aq}. The mixture was stirred for 1 h at RT and then submitted to reversed phase flash chromatography (C18 silica gel, eluents: from 100% A to 100% B, A: water/MeCN 95/5 + 0.1% HCOOH, B: MeCN/water 95/5 + 0.1% HCOOH) to afford the title compound (47) (68 mg, 0.082 mmol, 78 % yield).


N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
47	3.83	733,1	5	(DMSO-d ₆) δ 10.56 - 10.00 (bs, 1 H), 8.46 (s, 1 H), 8.13 (m, 1 H), 7.94 - 7.86 (m, 1 H), 7.74 (dt, J=7.48, 1.36 Hz, 1 H), 7.50 - 7.117 (m, 13 H), 7.09 (d, J=7.89 Hz, 1 H), 6.94 (d, J=7.89 Hz, 1 H), 6.70 (br. s., 1 H), 6.52 (d, J=9.87 Hz, 1 H), 5.15 (br. s., 1 H), 4.45 - 4.19 (m, 1 H), 4.00 (m, 2 H), 3.23 (m, 7 H), 2.91 - 2.63 (m, 6 H), 1.82 (t, J=11.02 Hz, 2 H), 1.53 (d, J=2.30 Hz, 5 H), 1.45 (d, J=13.15 Hz, 2 H).	Formate

Example 11

(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate diformate (compound 48)




Step 1; (R)-quinuclidin-3-yl 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



5 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid as described in Example 10, step 5 (150 mg, 0.176 mmol) and CDI (57.2 mg, 0.353 mmol) were reacted for 30 min at RT, then (R)-quinuclidin-3-ol (67.3 mg, 0.529 mmol) added and the reaction stirred overnight. The crude was partitioned
10 between saturated NaHCO_3 aq and AcOEt, organic layer was washed twice with water, saturated NaCl aq, dried over Na_2SO_4 and evaporated to dryness. The resulting crude was submitted to flash chromatography (silica gel, eluents: from 100% AcOEt to 100% AcOEt/7N NH_3 in MeOH 9/1) to afford the title compound (93 mg, 0.097 mmol, 54.9 % yield).

15 UPLC-MS: 1.13 min, 959.7 $[\text{M}+\text{H}]^+$, method 2.

Step 2; (R)-quinuclidin-3-yl 2-(3-((4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate formate



20 (R)-quinuclidin-3-yl 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-

butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (90 mg, 0.094 mmol), Pd/C (9.98 mg, 4.69 μ mol) and HCOOH (7.20 μ l, 0.188 mmol) were dissolved in 0.94 mL of MeOH and hydrogenated under balloon pressure of H₂ for 2h at RT until complete debenzylation. Reaction mixture was filtered on PTFE membrane and
 5 evaporated to dryness to give the title compound (95 mg). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.95 min, 869.6 [M+H]⁺, method 2.

Step 3; (compound 48)


The title example was made in a similar way as that of the compound of example
 10 10, step 10 (86 mg, 0.094 mmol) to give the title compound (48) 60 mg, 0.080 mmol, 85 % yield).

UPLC-MS: 3.30 min (43.2%) - 3.32 min (56.8%), 655.1 [M+H]⁺, Method 5

N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
48	3.31	655.1	5	(DMSO-d ₆) δ ppm 10.44 (s, 2 H), 10.25 (br. s., 1 H), 9.22 - 8.92 (m, 1 H), 8.75 - 8.48 (m, 2 H), 8.32 - 8.16 (m, 1 H), 7.95 (dt, J=5.59, 1.64 Hz, 1 H), 7.80 (t, J=8.06 Hz, 1 H), 7.68 - 7.30 (m, 7 H), 7.15 (d, J=8.22 Hz, 1 H), 6.99 (d, J=8.22 Hz, 1 H), 6.95 - 6.80 (m, 1 H), 6.55 (d, J=9.87 Hz, 1 H), 6.01 - 6.25 (m, 1 H), 5.41 (d, J=9.87 Hz, 1 H), 5.05 - 5.26 (m, 1 H), 3.59 - 3.79 (m, 2 H), 2.78 - 3.33 (m, 12 H), 2.22 (br. s., 1 H), 1.32 - 2.00 (m, 8 H)	Formate


Example 12

(R)-1-methylpyrrolidin-3-yl 2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate diformate (Compound 49)



5


Step 1; (R)-1-methylpyrrolidin-3-yl 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



10 The title compound was made in similar way as that of Example 11 Step 1 from 2-(3-((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid (150 mg, 0.176 mmol) to give the desired product (93 mg, 0.100 mmol, 56.5 % yield).

15 UPLC-MS: 1.12 min, 933.0 [M+H]⁺, method 2.

Step 2; (R)-1-methylpyrrolidin-3-yl 2-(3-((4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



The title compound was made in similar way as that of Example 11, step 2 from (R)-1-methylpyrrolidin-3-yl 2-(3-(((4-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (90 mg, 0.092 mmol), Pd-C 10% wet (9.78 mg, 4.60 μ mol) and HCOOH (7.05 μ l, 0.184 mmol) to give the desired product (90 mg). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.93min, 843.7 [M+H]⁺, method 2.

Step 3; (Compound 49)


The title example was made in a similar way as that of the compound of example 10, step 10 from (R)-1-methylpyrrolidin-3-yl 2-(3-(((4-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate formate (90 mg, 0.101 mmol) to give the desired compound (40 mg, 0.055 mmol, 54.8 % yield).

N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
49	3.08	629.0	5	(DMSO-d ₆) δ 10.10 - 10.50 (m, 1 H), 8.47 (s, 1 H), 8.21 (s, 1 H), 8.16 (d, J=9.87 Hz, 1 H), 7.90 (q, J=1.97 Hz, 1 H), 7.75 (dd, J=7.73, 1.48 Hz, 1 H), 7.24 - 7.50 (m, 7 H), 7.09 (d, J=8.22 Hz, 1 H), 6.94 (d, J=7.89 Hz, 1 H), 6.68 (br. s., 1 H), 6.52 (d, J=9.87 Hz, 1 H), 5.08 - 5.28 (m, 2 H), 2.88 - 2.80 (m, 2 H), 2.76 (d, J=6.58 Hz, 2 H), 2.65 (dd, J=10.69, 6.08 Hz, 1 H), 2.54 (s, 1 H), 2.47 (d, J=10.85 Hz, 1 H), 2.25 (d, J=7.23 Hz, 1 H), 2.18 (d, J=3.29 Hz, 4 H), 1.64 (td, J=6.74, 2.63 Hz, 1 H), 1.54 (br. s., 4 H)	Formate


Example 13

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

yl)ethyl)amino)methyl)benzamido)propyl)carbamoyl)phenyl)-2-phenylacetate diformate (Compound 50)




Step 1; Methyl 2-(3-((3-((tert-butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



3-(1-Hydroxy-2-methoxy-2-oxo-1-phenylethyl)benzoic acid obtained as in Example 10, step 2 (500 mg, 1.747 mmol) and TBTU (561 mg, 1.747 mmol) were reacted for 5 min at RT in 5 mL DMF, then DIEA (0.305 ml, 1.747 mmol) and tert-butyl (3-aminopropyl)carbamate (335 mg, 1.921 mmol) were added and the resulting mixture stirred at RT for 30 min. Reaction was partitioned between 0.5M HCl_{aq} and AcOEt, organic phase washed three times with 0.5M HCl_{aq} and once with saturated NaHCO₃ aq, once with saturated NaCl_{aq} and dried over Na₂SO₄ to give the title compound (0.78 g, 1.763 mmol, quantitative yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.99 min, 465 [M+Na]⁺, method 1.

Step 2; 2-(3-((3-((tert-butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid




20 Methyl 2-(3-((3-((tert-butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-

hydroxy-2-phenylacetate (0.77 g, 1.740 mmol) and LiOH (0.208 g, 8.70 mmol) were reacted in 11 mL of THF/water 1/1 for 1h at RT, then reaction quenched by the addition of 0.5M HCl_{aq} and extracted with AcOEt. Organic phase was washed with saturated NaCl_{aq}, dried over Na₂SO₄ and evaporated to dryness to give the title compound (0.61 g, 1.424 mmol, 82 % yield).

UPLC-MS: 0.86 min, 451.1 [M+Na]⁺, method 1.


Step 3; (1-benzylpiperidin-4-yl)methyl 2-(3-((3-((tert-butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



2-(3-((3-((tert-butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetic acid (300 mg, 0.700 mmol) and CDI (227 mg, 1.400 mmol) were reacted in 4.7 mL of DMF at RT for 5 min, then (1-benzylpiperidin-4-yl)methanol (431 mg, 2.100 mmol) added and the mixture stirred at 45°C overnight. Reaction was partitioned between saturated NaHCO₃ and AcOEt, and organic phase after evaporation was submitted to flash chromatography (silica gel, eluents: from 100% AcOEt to 80/20 AcOEt/NH₃ 7N in MeOH) to afford the title compound (310 mg, 0.503 mmol, 71.9 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 0.75 min, 616.2 [M+H]⁺, method 2.

Step 4; (1-benzylpiperidin-4-yl)methyl 2-(3-((3-aminopropyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate




(1-Benzylpiperidin-4-yl)methyl

2-(3-((3-((tert-

5 butoxycarbonyl)amino)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (310 mg, 0.503 mmol) was stirred for 1h at RT in 4M HCl dioxane (8.00 mmol) diluted with further 2.5 mL of dioxane, then dried under reduce pressure and the residue triturated in Et₂O to give the title compound (0.22 g, 0.374 mmol, 74.2 % yield) used in the next steps without further purifications.

10 UPLC-MS: 0.38 min, 516.1 [M+H]⁺, method 2.

Step 5; (1-benzylpiperidin-4-yl)methyl 2-(3-((3-(4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)-carbamoyl)phenyl)-2-hydroxy-2-phenylacetate



15

(R)-4-(((tert-butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid, obtained as for Procedure 4 Steps 1-5, (63 mg, 0.111 mmol) and TBTU (35.6 mg, 0.111 mmol) was reacted for 5 min in 1.1 mL of DMF, (1-benzylpiperidin-4-yl)methyl 2-(3-((3-aminopropyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate diformate (85 mg,

20

0.144 mmol) and DIEA (58.0 μ l, 0.332 mmol) were added and the mixture stirred for 1h at 60°C. Reaction was partitioned between AcOEt and saturated NaHCO₃, washed twice with water, saturated NaCl_{aq}, organic layer dried over Na₂SO₄ and evaporated to dryness. The resulting crude (110 mg, 0.103 mmol, 93 % yield) was used in the next step without further purifications.

UPLC-MS: 1.06 min, 1066.2 [M+H]⁺, method 2.


Step 6; (Compound 50)

The title example was made in a similar way as that of the compound of example 10, step 10, from (1-benzylpiperidin-4-yl)methyl 2-(3-((3-(4-(((tert-butoxycarbonyl)((R)-2-((tert-butyl)dimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (108 mg, 0.101 mmol) to give the title compound (50) (22 mg, 0.023 mmol, 23.01 % yield).


N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
50	3.91	852.1	5	(DMSO-d ₆) δ ppm 10.54 - 9.76 (bs, 2 H), 8.66 - 8.38 (m, 2 H), 8.23 - 8.05 (m, 2 H), 7.90 (s, 1 H), 7.80 (d, J=7.94 Hz, 3 H), 7.52 - 7.14 (m, 15 H), 7.07 (d, J=8.38 Hz, 1 H), 6.91 (d, J=8.38 Hz, 1 H), 6.73 (s, 1 H), 6.48 (d, J=9.70 Hz, 1 H), 5.70 - 5.35 (m, 1 H), 5.53 - 4.94 (m, 1 H), 4.23 - 4.45 (m, 1 H), 4.00 (d, J=6.62 Hz, 2 H), 3.88 (br. s., 2 H), 3.38 (m, 8 H), 2.72 (br. s., 4 H), 1.75 (s, 4 H), 1.64 - 1.43 (m, 3 H)	Formate

Example 14

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate diformate (Compound 51)




5

Step 1; methyl 2-(4-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetate ()

A solution of methyl 2-oxo-2-phenylacetate (12.83 ml, 91 mmol) in 350 mL of THF, was added dropwise with 100 mL of a solution 1M (4-(benzyloxy)phenyl)magnesium bromide (100 mmol) in THF at 0°C over 30 min and stirred overnight at RT. Reaction was partitioned between AcOEt and saturated NaCl_{aq}, organic layer dried over Na₂SO₄ and evaporated to dryness. The residue was crystallised in Et₂O to afford the title compound (15 g, 43.1 mmol, 47.4 % yield).

UPLC-MS: 1.24 min, 331.2 [(M+H)-H₂O]⁺, method 1.

Step 2; methyl 2-hydroxy-2-(4-hydroxyphenyl)-2-phenylacetate


15

Methyl 2-(4-(benzyloxy)phenyl)-2-hydroxy-2-phenylacetate (5g, 14.35 mmol) was dissolved in 100 mL of a mixture ethanol/AcOEt 6/4, added with Pd-C 5% wet (3.05 g, 0.718 mmol) and stirred under 25 psi of hydrogen for 2h at RT. Reaction mixture was

filtered on PTFE membrane and evaporated to dryness to give the crude, that was recrystallized in iPr₂O to afford the title compound (2.95 g, 11.43 mmol, 79.6 % yield).

UPLC-MS: 0.76 min, 241.1 [(M+H)-H₂O]⁺, method 1.


Step 3; methyl 2-(4-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate (343-28-1)



methyl 2-hydroxy-2-(4-hydroxyphenyl)-2-phenylacetate (1 g, 3.87 mmol) and 2-(4-bromobutyl)-1,3-dioxolane (0.616 ml, 4.07 mmol) were reacted in 15.5 mL of at 80°C for 8h and RT overnight. Reaction mixture was partitioned between water and AcOEt, washed twice with water and once with saturated NaCl_{aq}. Organic layer was dried over Na₂SO₄ and evaporate to dryness to give the title compound (1.54 g, 3.99 mmol, 103 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 1.12 min, 369.3 [(M+H)-H₂O]⁺, method 1.


Step 4; methyl 2-hydroxy-2-(4-((5-oxopentyl)oxy)phenyl)-2-phenylacetate



Methyl 2-(4-(4-(1,3-dioxolan-2-yl)butoxy)phenyl)-2-hydroxy-2-phenylacetate (1.50 g, 3.88 mmol) was dissolved in 19.4 mL of a mixture ACN / 1M HCl_{aq} and stirred for 6h at RT. Reaction was partitioned in AcOEt and washed three times with water and once with saturated NaCl_{aq}. Organic layer was dried over Na₂SO₄ and evaporated to dryness to give the title compound (1.12 g, 3.27 mmol, 84 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 1.04 min, 325.2 [(M+H)-H₂O]⁺, method 1.


Step 5; methyl 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-

butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate

(R)-5-(2-amino-1-((tert-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (1.637 g, 3.86 mmol) and methyl 2-hydroxy-2-(4-((5-oxopentyl)oxy)phenyl)-2-phenylacetate (1.10 g, 3.21 mmol) were reacted for 1h at RT in 16 mL of DCM with a spatula tip of Na₂SO₄, then AcOH (0.184 ml, 3.21 mmol) and Na(OAc)₃H (1.362 g, 6.43 mmol) were added in sequence and the mixture stirred for 3h at RT. The reaction was cooled to 0°C before addition of TEA (0.448 ml, 3.21 mmol) and di-tert-butyl dicarbonate (0.746 ml, 3.21 mmol), then the reaction was stirred for further 2h at RT. The mixture was partitioned between AcOEt and washed three times with 0.2 M HCl_{aq}, twice with water and once with saturated NaCl_{aq}. Organic layer was dried over Na₂SO₄ and evaporated to dryness to give a crude that after purification by flash chromatography (silica gel – eluents: hexane to hexane/THF) afforded the title compound (1.33 g, 1.563 mmol, 48.6 % yield).

UPLC-MS: 1.78 min, 873.7 [M+Na]⁺, method 2.


Step 6; 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetic acid



The title compound was made in a similar way as that of the Example 10, step 5 from methyl 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (1.33 g, 1.563 mmol) and LiOH (0.187 g, 7.81 mmol) to give the desired product (1.1 g, 1.314 mmol, 84 % yield).

UPLC-MS: 1.68 min, 838.2 [M+H]⁺, method 1. 20141106_16


Step 7; benzyl 4-((2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate



The title compound was made in a similar way as that of the compound of Example 10, step 7 from 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetic acid (200 mg, 0.239 mmol) to give the desired product (98 mg, 0.092 mmol, 38.4 % yield).

UPLC-MS: 3.25 min, 1068.5 [M+H]⁺, method 3.


Step 8; piperidin-4-ylmethyl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyltrimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



5 The title compound was made in a similar way as that of the Example 10, step 8 from benzyl 4-((2-(4-((5-((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyltrimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetoxy)methyl)piperidine-1-carboxylate (98 mg, 0.092 mmol), Pd-C 10% wet (9.76 mg, 4.59 μ mol), and HCOOH (7.04 μ l, 0.183 mmol) to give the desired product
10 (73 mg, 0.082 mmol, 89 % yield). The product obtained was used in the following steps without further purifications.

UPLC-MS: 1.08 min, 844.7 [M+H]⁺, method 2.

Step 9; (1-benzylpiperidin-4-yl)methyl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyltrimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



The title compound was made in a similar way as that of the Example 10, step 9 from (1-benzylpiperidin-4-yl)methyl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyltrimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (73 mg, 0.082 mmol), benzaldehyde (26.1 mg, 0.246 mmol), AcOH (9.39 μ l, 0.164 mmol) and NaB(OAc)₃H (52.1 mg, 0.246 mmol) to give the desired product (77 mg). The product obtained was used in the following steps without further purifications.

5 UPLC-MS: 1.17 min, 934.7 [M+H]⁺, method 2.

Step 10; (Compound 51)


The title example was made in a similar way as that of the example 10, step 10 from (1-benzylpiperidin-4-yl)methyl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyl

10 yl)dimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (77 mg, 0.082 mmol) to give the title product (51) (28 mg, 0.037 mmol, 44.4 % yield).


N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
51	4.15	720	5	(DMSO-d ₆) δ ppm 10.51 - 10.21 (bs, 1 H), 8.27 - 8.10 (m, 2 H), 7.40 - 7.18 (m, 12 H), 7.11 (d, J=8.22 Hz, 1 H), 6.95 (d, J=8.22 Hz, 1 H), 6.90 - 6.82 (m, 2 H), 6.54 (d, J=9.87 Hz, 1 H), 6.42 (br. s., 1 H), 5.19 (br. s., 1 H), 4.02 - 3.90 (m, 4 H), 3.41 - 3.35 (m, 2 H), 2.68 - 2.96 (m, 6 H), 1.91 - 1.39 (m, 10 H), 1.15 - 1.07 (m, 2 H)	Formate

Example 15

15 (R)-quinuclidin-3-yl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate diformate (Compound 52)




Step 1; (R)-quinuclidin-3-yl 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



5 The title compound was made in a similar way as that of the Example 14, step 7 to give the desired product (138 mg, 0.146 mmol, 81 % yield).

UPLC-MS: 1.27 min, 946.7 [M+H]⁺, method 2.

Step 2; (R)-quinuclidin-3-yl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



The title compound was made in a similar way as that of the Example 14, step 8 from (R)-quinuclidin-3-yl 2-(4-((5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (138 mg, 0.146 mmol), Pd-C 10% wet (15.52 mg, 7.29 μmol) and HCOOH (11.19 μl, 0.292 mmol) to give the desired product (150 mg). The product obtained was used in the following step without further purifications.

UPLC-MS: 1.09 min, 856.7 [M+H]⁺, method 2.


Step 3; (compound 52)

The title example was made in a similar way as that of the compound of example 10, step 10, from (R)-quinuclidin-3-yl 2-(4-((5-((tert-butoxycarbonyl)((R)-2-((tert-butyl)dimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-hydroxy-2-phenylacetate formate (132 mg, 0.146 mmol) to give the title product (52) (68 mg, 0.099 mmol, 67.6 % yield).


N	Rt (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
52	3.64	642.0	5	(DMSO-d ₆) δ ppm 10.59 - 10.24 (bs, 1 H), 8.19 (d, J=10.19 Hz, 1 H), 8.15 (s, 1 H), 7.38 - 7.27 (m, 5 H), 7.25 (d, J=8.88 Hz, 2 H), 7.14 (d, J=8.22 Hz, 1 H), 6.98 (d, J=8.22 Hz, 1 H), 6.92 - 6.83 (m, 2 H), 6.56 (d, J=9.87 Hz, 1 H), 6.45 (br. s., 1 H), 5.32 (dd, J=9.87, 2.63 Hz, 1 H), 4.92 - 4.71 (m, 1 H), 4.44 - 4.21 (m, 1 H), 4.05 - 3.89 (m, 2 H), 3.18 (dd, J=13.98, 8.39 Hz, 2 H), 3.10 - 2.88 (m, 5 H), 2.77 - 2.58 (m, 3 H), 1.92 (br. s., 1 H), 1.76 - 1.34 (m, 8 H)	Formate

Example 16

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate diformate (Compound 53)




Step 1; methyl 2-(4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



Methyl 2-hydroxy-2-(4-hydroxyphenyl)-2-phenylacetate obtained as in Example 14, step 2 (0.5 g, 1.936 mmol), tert-butyl 4-(bromomethyl)benzylcarbamate (0.581 g, 1.936 mmol) and K₂CO₃ (0.268 g, 1.936 mmol) were reacted in 9.7 mL DMF at 80°C for 3h. Reaction was partitioned between water and AcOEt, washed three times with water and saturated NaCl_{aq}, dried over Na₂SO₄ and evaporated to dryness. The residue was purified by flash column chromatography (silica gel; eluents: hexane to 50/50 hexane/AcOEt) to afford the title compound (1.42 g, 93 %).

UPLC-MS: 1.30 min, 460.0 [(M+H)-H₂O]⁺, method 2.


Step 2; 2-(4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetic acid



The title compound was made in a similar way as that of the example 10, step 5 from methyl 2-(4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (0.65 g, 1.361 mmol) and LiOH (0.163 g, 6.81 mmol) to give the desired product (673mg, 1.452 mmol, 107 % yield).

UPLC-MS: 2.28 min, 446.0 [(M+H)-H₂O]⁺, method 4.


Step 3; 2 (1-benzylpiperidin-4-yl)methyl 2-(4-((4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



The title intermediate was made in a similar way as that of the compound of Example 10, step 7 from 2-(4-((4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetic acid (630 mg, 1.359 mmol), CDI (661 mg, 4.08 mmol) and (1-benzylpiperidin-4-yl)methanol (1116 mg, 5.44 mmol) to give the desired product (520 mg, 0.799 mmol, 58.8 % yield).

UPLC-MS: 0.92 min, 651.1 [M+H]⁺, method 2.


Step 4; (1-benzylpiperidin-4-yl)methyl 2-(4-((4-(aminomethyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate bis hydrochloride



The title compound was made in a similar way as that of the Example 10, step 10 from 2 (1-benzylpiperidin-4-yl)methyl 2-(4-((4-(((tert-butoxycarbonyl)amino)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (520 mg, 0.799 mmol) to give the title compound (357 mg, 0.572 mmol, 71.6 % yield).

UPLC-MS: 0.49 min, 551.1 [M+H]⁺, method 2.

Step 5; (1-benzylpiperidin-4-yl)methyl 2-(4-((4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate



5

(R)-4-(((tert-butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid, obtained as described in Procedure 4, steps 1-5, (50 mg, 0.264 mmol), (1-benzylpiperidin-4-yl)methyl 2-(4-((4-(aminomethyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate bis hydrochloride (164 mg, 0.264 mmol), 1H-benzo[d][1,2,3]triazol-1-ol (46.3 mg, 0.343 mmol), DMAP (4 mg, 0.033 mmol) and DIEA (0.138 ml, 0.791 mmol) were stirred for 5 min in 3 mL of DMF, then EDC (65.7 mg, 0.343 mmol) added and the mixture stirred overnight at RT. Reaction mixture was partitioned between saturated NaHCO_3 aq and AcOEt, organic layer washed three times with water and saturated NaCl aq, dried over Na_2SO_4 and evaporated to dryness to give the crude (292 mg). The crude obtained was used in the following step without further purifications.

15

UPLC-MS: 1.14 min, 1102.1 $[\text{M}+\text{H}]^+$, method 2.

Step 6; (Compound 53)



The title example was made in a similar way as that of the example 10, step 10 from 1-benzylpiperidin-4-yl)methyl 2-(4-((4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (290 mg, 0.263 mmol) to give the title product (53) (62 mg, 0.063 mmol, 24.05 % yield).

20

N	R _t (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt 2 eq unless stated
53	4.72	887.0	5	(DMSO-d ₆) δ ppm 10.33 (br. s., 1 H), 9.00 (t, J=5.95 Hz, 1 H), 8.22 - 8.01 (m, 2 H), 7.85 (d, J=7.94 Hz, 2 H), 7.49 - 6.83 (m, 22 H), 6.60 - 6.34 (m, 2 H), 5.20 - 4.97 (m, 3 H), 4.47 (d, J=5.73 Hz, 2 H), 4.34 (br. s., 1 H), 3.97 (d, J=6.17 Hz, 2 H), 3.89 (s, 2 H), 3.59 - 3.37 (m, 5 H), 2.84 - 2.59 (m, 4 H), 1.84 (t, J=11.03 Hz, 2 H), 1.58 - 1.35 (m, 3 H), 1.19 - 0.97 (m, 4 H)	Formate

Example 17

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)carbonyl)amino)phenyl)-2-phenylacetate bis
5 hydrochloride (Compound 54)


**Step 1; Methyl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate**

Methyl 2-oxo-2-phenylacetate 16.4 mmol, 4.22 g), was dissolved (in DMF (5 ml,
10 21.32 mmol) and the solution was cooled to -15 C°, then (3-(bis(trimethylsilyl)amino)phenyl)magnesium chloride (21.32 ml, 21.32 mmol) was added dropwise and the mixture left to reach rt. Methyl 2-oxo-2-phenylacetate (3.5 g, 21.32 mmol) was dissolved in DMF dry (10 ml) under argon and the solution was cooled

down to -20 C°, then (3-(bis(trimethylsilyl)amino)phenyl)magnesium chloride 1.0 M in THF (21.32 ml, 21.32 mmol) was added dropwise and the mixture left to reach rt and stirred for further 1.5 h. The reaction was quenched by the addition of Acetonitrile (20 ml) and partitioned between AcOEt (80 ml) and water (80 ml). Organic layer was washed twice
5 with water, once with sat NaCl, dried over Na₂SO₄ and dried under reduced pressure. The crude was purified by flash chromatography on silica gel (eluent –from 95/5 Hexane/AcOEt to 100% AcOEt) to afford the title compound (3.73 g, 14.50 mmol, 68 % yield) .

UPLC-MS: 0.89 min, 258.1 [(M+H)]+, method 4.

10 **Step 2; (R)-tert-butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(5-hydroxypentyl)carbamate**




(R)-8-(benzyloxy)-5-(2-bromo-1-((tert-butyldimethylsilyl)oxy)ethyl)quinolin-
2(1H)-one (500 mg, 1.024 mmol) and 5-aminopentan-1-ol (106 mg, 1.024 mmol) were
15 dissolved in NMP (3 ml) and the solution was stirred under nitrogen at 85 °C overnight. The reaction mixture was partitioned between AcOEt and water, the organic phase washed twice with water and the organic layers were dried over Na₂SO₄ and evaporated under reduced pressure to give a crude that was dissolved in 1/1 THF and NaHCO₃ sat. sol. (6 ml), di-tert-butyl dicarbonate (67.1 mg, 0.308 mmol) was added and the mixture heated at rt
20 for 1h. The reaction mixture was partitioned between AcOEt (20 ml) and water and the organic phase washed twice with water, once with HCl 0.5 M(20 ml) and once with brine. Organic layers were dried over Na₂SO₄ and dried under reduced pressure. The crude was purified by flash chromatography on silica gel (eluent - from 100 % Hexane to

20%/80%Hexane/Ethyl acetate 80 %.) to give the title compound (76 mg, 0.124 mmol, 12.16 % yield).

UPLC-MS: 2.86 min, 611 [(M+H)]+, method 4.


Step 3; methyl 2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate



Methyl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate (1 g, 3.89 mmol) was dissolved in Acetonitrile (12 ml) and trichloromethyl carbonochloridate (1.922 g, 9.72 mmol) was added. The solution was stirred at rt for 7 hours then it was evaporated under vacuum, added with 1 ml of DCM and evaporated again. The residue was dissolved in DCM (5 ml), added with (R)-tert-butyl (2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(5-hydroxypentyl)carbamate (2.374 g, 3.89 mmol), and the reaction was stirred at rt for 6 hours. The solvent was evaporated and the residue purified by flash chromatography on silica gel (eluent – Hexane/Ethyl acetate from 0 to 90 %) to afford the title compound (344 mg, 9.9 %).

UPLC-MS: 1.64 min, 895.4[(M+H)]+, method 2.

Step 4; 2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxa-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetic acid




5 Methyl 2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxa-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate (143 mg, 0.160 mmol) was dissolved in DMF (0.7 ml) and LiOH 1 N (0.320 ml, 0.320 mmol) was dropped inside; the mixture was stirred at rt for 2 h. The pH of mixture was adjust with a 1 N HCl aq solution to pH= 5.

10 Reaction mixture was partitioned between AcOEt (5 ml) and water and the organic phase washed twice with water then with brine. Finally the organic solution was dried over sodium sulphate, evaporated under vacuum to afford the title compound (120 mg, 85 %), that was used as such for the next steps.

UPLC-MS: 1.59 min, 880.2 [(M+H)]+, method 2.

15 **Step 5; (1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-8-(tert-butoxycarbonyl)-10-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)-12,12,13,13-tetramethyl-2,11-dioxa-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate**



2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxa-8-aza-12-silatetradecan-1-


20

oyl)amino)phenyl)-2-hydroxy-2-phenylacetic acid (120 mg, 0.136 mmol) was dissolved in DMF (1.5 ml, 0.136 mmol) and di(1H-imidazol-1-yl)methanone, CDI (33.2 mg, 0.205 mmol) was added and the mixture was stirred at rt for 1 h. Finally (1-benzylpiperidin-4-yl)methanol (33.6 mg, 0.164 mmol) was added and the reaction was left to stir at rt for 3 h.

5 Reaction mixture was partitioned between AcOEt (20 ml) and water; the organic phase was washed twice with water, HCl 0.5 M(20 ml) and brine. Finally the solution was dried over sodium sulphate and evaporated under vacuum. The residue was submitted to flash chromatography on silica gel (eluent- from 100% AcOEt to 80/20 AcOEt/(NH₃ 7N in MeOH) to afford the title compound (75 mg, 0.070 mmol, 51.5 % yield) as yellowish foam.

10 UPLC-MS: 2.51 min, 1066.6 [(M+H)]⁺, method 4.

Step 6; (1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-8-(tert-butoxycarbonyl)-10-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate



15 (1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate (150 mg, 0.281 mmol) was dissolved in methanol (2 ml, 0.281 mmol) under nitrogen then the suspension was added with Pd/BaSO₄ (40 mg, 0.281 mmol) and hydrogenated for 2h at rt under balloon pressure of hydrogen. Reaction mixture was filtered and dried to give the crude title compound (75 mg, 0.153 mmol, 54.6 % yield) as a yellowish oil, used as is in the next step without further purification.

20 UPLC-MS: 2.59 min, 977.53 [(M+H)]⁺, method 4.


Step 7; (Compound 54)

(1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-8-(tert-butoxycarbonyl)-10-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate (30 mg, 0.031 mmol) was dissolved in HCl 37 % (0.5 ml) and the solution was stirred at room temperature for 5 min. with complete disappearing of starting material in UPLC. Evaporation of solvent was performed using V10 Biotage. The residue was triturated in acetone and filtrate on a cartridge to afford the title compound (54) (11 mg, 47 %).


N	R _t (min)	[M+H] ⁺	Method	NMR data (600 MHz)	Salt 2 eq unless stated
54	4.19	763.1	5	(DMSO-d ₆) δ ppm 10.44 (br. s., 2 H), 9.58 (s, 1 H), 9.16 (br. s., 1 H), 8.66 (br. s., 1 H), 8.26 (d, J=9.87 Hz, 1 H), 7.69 - 6.82 (m, 15 H), 6.55 (d, J=9.87 Hz, 1 H), 6.26 - 5.95 (m, 1 H), 5.44 (dd, J=10.03, 2.14 Hz, 1 H), 4.22 (d, J=4.93 Hz, 2 H), 4.06 - 3.87 (m, 4 H), 3.28 (d, J=11.84 Hz, 2 H), 3.12 - 2.73 (m, 6 H), 1.86 - 1.25 (m, 10 H)	HCl

10 **Example 18**

(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentanamido)phenyl)-2-phenylacetate bis hydrochloride (Compound 55)




15 **Step 1; Methyl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)amino)pentanamido)phenyl)-2-hydroxy-2-**

phenylacetate

Methyl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate (500 mg, 1.943 mmol) obtained as in Example 17, step 1, was dissolved in DCM (2 ml) and 5-bromopentanoyl chloride (504 mg, 2.53 mmol) was added and the reaction was stirred at rt for 2 hours. The mixture was diluted with DCM (2 ml) and the organic phase was washed with NaHCO₃ saturated solution, then with 1 M HCl and the twice with brine. The solution was dried over sodium sulphate and evaporated under vacuum to give the title compound as a pale brown oil (490 mg, 60%).

UPLC-MS: 1.82 min, 404.01 [(M+H)-H₂O]⁺, method 4.

Step 2; Methyl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate




(R)-5-(2-amino-1-((tert-butyldimethylsilyl)oxy)ethyl)-8-(benzyloxy)quinolin-2(1H)-one (376 mg, 0.885 mmol) was dissolved in DMF dry (2 ml). 4-aminobutan-1-ol (0.082 ml, 0.884 mmol) was added and the solution stirred under nitrogen at 95 °C for 4 hrs. Reaction mixture was partitioned between AcOEt and water and the organic phase washed twice with water. The organic layer was dried over Na₂SO₄ and evaporated under reduced pressure to give an pale yellow oil. The residue was purified by reversed phase

flash column chromatography (eluent – from 100%A to 100%B in 15CV; A: water/MeCN 95/5 + 0.01%HCOOH; B: water/MeCN 5/95 + 0.01%HCOOH) to afford the title compound (464 mg, 0.607 mmol, 82 % yield).

UPLC-MS: 2.13 min, 764.01 [(M+H)]⁺, method 4.


5 **Step 3; Methyl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate**



Methyl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-
10 butyldimethylsilyl)oxy)ethyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate
(170 mg, 0.223 mmol) was dissolved in THF (1 ml), the di-tert-butyl dicarbonate (72.8 mg, 0.334 mmol) and NaHCO₃ saturated solution (1 ml, 4.45 mmol) were added and the mixture was stirred at rt for 1h. Reaction mixture was partitioned between AcOEt (20 ml) and water and the organic phase washed twice with water, HCl 0.5 M (20 ml) and brine.
15 Finally the organic solution was dried over sodium sulphate, evaporated under vacuum to give the title compound in quantitative yield (192 mg). No purification was needed.

UPLC-MS: 1.57 min, 864.37 [(M+H)]⁺, method 2.


Step 4; 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyl)dimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetic acid



5 Methyl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyl)dimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate (170 mg, 0.197 mmol) was dissolved in DMF (1 ml) and LiOH 1 N (0.393 ml, 0.393 mmol) was added and the mixture was stirred at rt for 1h. The pH of mixture was adjust with a 1 N HCl aq solution to pH= 5. Reaction mixture was partitioned
10 between AcOEt (5 ml) and water and the organic phase washed twice with water then with brine. Finally the organic solution was dried over sodium sulphate, evaporated under vacuum to afford the title compound (170 mg, 90 %). No purification was needed.

UPLC-MS: 1.50 min, 850.01 [(M+H)]⁺, method 2.

Step 5; (R)-quinuclidin-3-yl 2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-butyl)dimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate




2-(3-(5-(((R)-2-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-2-((tert-

butyldimethylsilyl)oxy)ethyl)(tert-butoxycarbonyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetic acid (170 mg, 0.20 mmol) was dissolved in DMF (1.3 ml) and di(1H-imidazol-1-yl)methanone, CDI (48.6 mg, 0.30 mmol) was added and the mixture was stirred at rt for 1 h. Finally (R)-quinuclidin-3-ol (30.5 mg, 0.240 mmol) was added and the reaction was left to stir at rt for 3 h. Reaction mixture was partitioned between AcOEt (20 ml) and water; the organic phase was washed twice with water, HCl 0.5 M (20 ml) and brine. Finally the solution was dried over sodium sulphate and evaporated under vacuum. The residue was purified by flash column chromatography (eluent – 100% AcOEt to 80/20 AcOEt/NH₃ 7N in MeOH) to afford the title compound (105 mg, 54.7 %).

UPLC-MS: 1.30 min, 959.52 [(M+H)]⁺, method 2.

Step 6; (R)-quinuclidin-3-yl 2-(3-(5-((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate



(1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-10-(8-(benzyloxy)-2-oxo-1,2-dihydroquinolin-5-yl)-8-(tert-butoxycarbonyl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate (102 mg, 0.106 mmol) was dissolved in Methanol (1.5 ml) under nitrogen then the suspension was added with Pd/BaSO₄ (40 mg, 0.281 mmol) and hydrogenated for 2h at rt under balloon pressure of hydrogen. Reaction mixture was filtered and dried to give the crude (1-benzylpiperidin-4-yl)methyl 2-(3-(((R)-8-(tert-butoxycarbonyl)-10-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)-12,12,13,13-tetramethyl-2,11-dioxo-8-aza-12-silatetradecan-1-oyl)amino)phenyl)-2-hydroxy-2-phenylacetate (75 mg, 0.086 mmol, 81.1 % yield) as a yellowish oil, used as is in the next step without further purification.

UPLC-MS: 2.29 min, 435.1 [(M+H)/2]⁺, method 4.

Step 7; (Compound 55)


(R)-quinuclidin-3-yl 2-(3-(5-(((tert-butoxycarbonyl)((R)-2-((tert-butyl)dimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentanamido)phenyl)-2-hydroxy-2-phenylacetate (45 mg, 0.052 mmol) was dissolved in HCl 35 % (0.7 ml) and stirred at rt for 5 min. with complete disappearing of starting material in UPLC. Evaporation of solvent was performed using V10 Biotage. The residue was triturated in acetone and filtrate on a cartridge to afford the title compound (55) (34 mg, 88 %).

10

N	R _t (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt 2 eq unless stated
55	3.38/3.41	655.1	5	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 10.60 - 10.37 (3 s, 3 H each), 9.94 - 5.17 (m, 21 H), 5.21 - 5.08 (m, 1 H), 3.75 - 1.75 (m, 17 H);	HCl


Example 19

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)ureido)phenyl)-2-phenylacetate diformate (Compound 56)



15


Step 1; methyl 2-(3-(3-(3-((tert-butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetate



Methyl 2-(3-aminophenyl)-2-hydroxy-2-phenylacetate (500 mg, 1.943 mmol) obtained as in Example 17, step 1, was dissolved in Acetonitrile (1.5 ml, 1.943 mmol) and trichloromethyl carbonochloridate (961 mg, 4.86 mmol) was added. The solution was stirred at rt for 7 hours then it was evaporated under vacuum, added with 1 ml of DCM and evaporated again. The residue was dissolved in DCM (3 ml) and added with tert-butyl (3-aminopropyl)carbamate (Aldrich, 339 mg, 1.943 mmol). The reaction was stirred at rt 1 hour, just after a few minutes abundant precipitation of salt. The solvent was filtered and the filtrate was evaporated to give a pale yellow oil of title compound (800 mg, 1.749 mmol, 90 %). No further purification was needed.

UPLC-MS: 1.15 min, 457.1 ([M+H]⁺), method 2.

Step 2; 2-(3-(3-(3-((tert-butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetic acid




Methyl 2-(3-(3-(3-((tert-butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetate (200 mg, 0.437 mmol))

and LiOH (52.3 mg, 2.186 mmol) were dissolved in Tetrahydrofuran (Volume: 1.5)/Water (Volume: 1.5) for 2h at rt, then the mixture partitioned between AcOEt and HCl 0.2 M. Organic layer was washed with sat NaCl aq, dried over Na₂SO₄ dry and evaporated under reduced pressure to give 2-(3-(3-(3-((tert-

butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetic acid (174 mg, 0.393 mmol, 90 % yield) as a white foam. No further purification was needed.

UPLC-MS: 0.93 min, 444.1 [M+H]⁺, method 2.


Step 3; (1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-aminopropyl)ureido)phenyl)-2-hydroxy-2-phenylacetate



2-(3-(3-(3-((tert-butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetic acid (200 mg, 0.451 mmol) was reacted with di(1H-imidazol-1-yl)methanone (146 mg, 0.902 mmol) for 30 min. at rt, then (1-benzylpiperidin-4-yl)methanol (278 mg, 1.353 mmol) was added and the reaction stirred for 2h at 60 °C, then at rt overnight. The crude was partitioned between sat NaHCO₃ and AcOEt. Organic layer was dried over Na₂SO₄ and dried under reduced pressure. The crude was purified over silica gel by flash chromatography (silica gel, eluents: from 100% AcOEt to 80/20 AcOEt/NH₃ 7N in MeOH) to afford the title compound (138 mg, 0.219 mmol, 48.5 % yield)

UPLC-MS: 1.29 min, 630.55 [M+H]⁺, method 4.

Step 4; (1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-aminopropyl)ureido)phenyl)-2-hydroxy-2-phenylacetate bis hydrochloride




(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-((tert-butoxycarbonyl)amino)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetate (138 mg, 0.219 mmol) was stirred for 1 h at rt in Dioxane/HCl 4 N (1.5 ml) diluted with further 1.5 mL of

dioxane; then the mixture was dried under reduce pressure and the residue triturated in Et₂O to give (1-benzylpiperidin-4-yl)methyl 2-(3-((3-aminopropyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate, 2HCl (0.22 g, 0.374 mmol, 74.2 % yield) and (1-benzylpiperidin-4-yl)methyl 2-(3-((3-aminopropyl)carbamoyl)phenyl)-2-hydroxy-2-phenylacetate (0.090 g, 0.170 mmol, 77.6 % yield) both as white solid used in the next steps without further purifications.

UPLC-MS: 0.51 min, 530.86 [M+H]⁺, method 4.

Step 5; (1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-(4-(((tert-butoxycarbonyl)((R)-2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetate



(R)-4-(((tert-butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid, obtained as described in Procedure 4, steps 1-5 (65.2 mg, 0.115 mmol) and TBTU (36.6 mg, 0.115 mmol) were reacted for 5 min in 1.1 mL of DMF, then (1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-aminopropyl)ureido)phenyl)-2-hydroxy-2-phenylacetate (90 mg, 0.149 mmol) and DIEA (58.0 µL, 0.332 mmol) were added and the mixture stirred for 1h at 60 °C. Reaction was partitioned between AcOEt and saturated NaHCO₃, organic layer was washed twice with water, saturated NaCl aq, then dried over Na₂SO₄ and evaporated to dryness. The orange resulting oil (43.4 mg, 0.040 mmol, 35 % yield) was used in the next step without further purifications.

UPLC-MS: 1.14 min, 1081.24 [M+H]⁺, method 2.


Step 6; (Compound 56)

(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-(4-(((tert-butoxycarbonyl)((R)-2-((tert-


butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)ureido)phenyl)-2-hydroxy-2-phenylacetate (50 mg, 0.046 mmol) was dissolved in acetonitrile (1 ml) and HCl 5 N (1 ml) were added. The mixture was stirred at rt for 15 min. then submitted to reverse phase flash chromatography (C18 silica gel, eluents: from 100% A to 100% B, A: water/MeCN 95/5 + 0.1% HCOOH, B: MeCN/water 95/5 + 0.1% HCOOH) to afford the title compound (56) (17 mg, 0.020 mmol, 43.5 % yield).

N	R _t (min)	[M+H] ⁺	Method	NMR data (400 MHz)	Salt 2 eq unless stated
56	3.94	867,0	5	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 10.45 - 10.11 (m, 2 H), 8.60 - 6.12 (m, 36 H), 3.78 - 0.82 (m, 18 H)	Formate

10 Example 20




(1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 57)



5

Step 1; (1-Benzylpiperidin-4-yl)methyl (S)-2-(3-(2-((tert-butoxycarbonyl)amino)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate




10 A stirred solution of (S)-2-(3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)acetic acid hydrochloride (0.20 g, 0.38 mmol) in DMF (1.9 mL) was added with DIPEA (0.132 mL, 0.76 mmol) and HATU (0.173 g, 0.46 mmol) and the mixture stirred at room temperature for 30 minutes. To this mixture was added a solution of N-(tert-butoxycarbonyl)ethylenediamine HCl (0.061 g, 0.38 mmol) and DIPEA 15 (0.13 mL, 0.76 mmol) in DMF (1.9 mL). The reaction mixture diluted with ethyl acetate and washed with 2M aqueous sodium hydroxide and brine. The organic phase was dried over magnesium sulfate, filtered and the filtrate evaporated at reduced pressure to afford the title compound (0.265 g, >100%).

¹H NMR (400 MHz, CDCl₃); δ 7.39 (dd, J=1.9, 7.7 Hz, 2H), 7.34 – 7.17 (m, 10H), 7.07 - 7.04 (m, 2H), 6.87 (dd, J=2.0, 7.9 Hz, 1H), 5.07 (s, 1H), 4.42 (s, 2H), 4.14 – 4.07 (m, 4H), 3.48 (s, 2H), 3.42 – 3.38 (m, 3H), 3.29 – 3.22 (m, 2H), 2.97 (s, 2H), 2.85 – 2.81 (m, 2H), 1.96 – 1.88 (m, 2H), 1.67 – 1.56 (m, 1H), 1.40 (s, 9H).

20

Step 2; (1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 57)



5

A stirred solution of (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-((tert-butoxycarbonyl)amino)ethyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate (0.265 g, 0.42 mmol) was added with a solution of HCl in dioxan (4M, 15 mL) and the mixture stirred at room temperature for 1 hour. The solvent evaporated under reduced pressure. The residue was dissolved in DMF (3 mL) and added to a pre-stirred (15 minutes) mixture of (R)-4-(((tert-butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid (0.20 g, 0.35 mmol), DIPEA (0.214 mL, 1.23 mmol) and HATU (0.16 g, 0.42 mmol) in DMF (2 mL). The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture dissolved in ethyl acetate and washed with saturated aqueous sodium carbonate and brine. The organic phase was dried with magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was dissolved in dioxan (1 mL) and a solution of HCl – dioxan (2 mL) added. The reaction mixture was stirred at room temperature for 1 hour. The solvent was evaporated at reduced pressure and the residue purified by reverse preparative HPLC.

15
20











¹H NMR (400 MHz, DMSO-d₆); δ 10.50 (s, 2H), 9.44 (s, 1H), 9.16 (s, 2H), 8.61 (dd, J=5.3, 5.3 Hz, 1H), 8.30 (dd, J=5.4, 5.4 Hz, 1H), 8.08 (d, J=9.9 Hz, 1H), 7.88 (d, J=8.3 Hz, 2H), 7.61 (d, J=8.4 Hz, 2H), 7.47 (s, 5H), 7.33 (d, J=4.5 Hz, 3H), 7.32 - 7.23 (m, 3H), 7.12 (d, J=8.3 Hz, 1H), 7.00 - 6.93 (m, 2H), 6.89 (dd, J=2.3, 8.0 Hz, 2H), 6.67 (s, 1H), 6.57 (d, J=9.9 Hz, 1H), 6.19 (s, 1H), 5.36 - 5.32 (m, 1H), 4.43 (s, 2H), 4.30 - 4.23 (m, 4H), 4.01

25













(d, J=5.0 Hz, 2H), 3.36 - 3.28 (m, 5H), 3.09 - 3.08 (m, 2H), 3.04 - 2.94 (m, 1H), 2.93 - 2.88 (m, 2H), 1.78 - 1.66 (m, 3H), 1.40 - 1.31 (m, 2H).

The following compounds were prepared in a similar fashion using the appropriate amine;













5

N	Appropriate amine	Structure
58		
59		
60		
61		
62		







(continued)

63		
64		
65		
66		
67		
68		

(continued)

69		
70		
71		
72		
73		
74		

(continued)

75		
76		
77		

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
57	2.38	11	(DMSO-d ₆); δ 10.50 (s, 2H), 9.44 (s, 1H), 9.16 (s, 2H), 8.61 (dd, J=5.3, 5.3 Hz, 1H), 8.30 (dd, J=5.4, 5.4 Hz, 1H), 8.08 (d, J=9.9 Hz, 1H), 7.88 (d, J=8.3 Hz, 2H), 7.61 (d, J=8.4 Hz, 2H), 7.47 (s, 5H), 7.33 (d, J=4.5 Hz, 3H), 7.32 - 7.23 (m, 3H), 7.12 (d, J=8.3 Hz, 1H), 7.00 - 6.93 (m, 2H), 6.89 (dd, J=2.3, 8.0 Hz, 2H), 6.67 (s, 1H), 6.57 (d, J=9.9 Hz, 1H), 6.19 (s, 1H), 5.36 - 5.32 (m, 1H), 4.43 (s, 2H), 4.30 - 4.23 (m, 4H), 4.01 (d, J=5.0 Hz, 2H), 3.36 - 3.28 (m, 5H), 3.09 - 3.08 (m, 2H), 3.04 - 2.94 (m, 1H), 2.93 - 2.88 (m, 2H), 1.78 - 1.66 (m, 3H), 1.40 - 1.31 (m, 2H).	TFA
58	2.43	11	(DMSO-d ₆); δ 10.52 (s, 2H), 9.40 (s, 1H), 9.12 - 9.12 (m, 2H), 8.07 (d, J=9.9 Hz, 1H), 7.62 (d, J=8.3 Hz, 2H), 7.51 (d, J=8.3 Hz, 2H), 7.47 (s, 6H), 7.33 (d, J=4.4 Hz, 4H), 7.31 - 7.21 (m, 2H), 7.15 - 7.11 (m, 1H), 6.98 (d, J=8.2 Hz, 1H), 6.93 - 6.88 (m, 3H), 6.58 (d, J=9.7 Hz, 1H), 6.20 (s, 1H), 5.35 (dd, J=2.1, 9.9 Hz, 1H), 4.82 (s, 2H), 4.29 - 4.21 (m, 6H), 4.04 - 3.97 (m, 5H), 3.34 (d, J=12.1 Hz, 4H), 3.12 - 2.98 (m, 3H), 2.92 - 2.88 (m, 2H), 1.77 - 1.67 (m, 3H), 1.41 - 1.33 (m, 2H).	TFA

(continued)

59	2.45	11	(DMSO-d ₆); δ 8.24 (d, J=7.7 Hz, 1H), 8.12 (d, J=9.9 Hz, 1H), 7.79 (d, J=8.3 Hz, 2H), 7.38 (d, J=8.3 Hz, 2H), 7.32 (d, J=4.4 Hz, 4H), 7.30 - 7.21 (m, 7H), 7.06 (d, J=8.2 Hz, 1H), 6.91 (d, J=7.8 Hz, 3H), 6.85 (dd, J=1.9, 7.3 Hz, 1H), 6.61 (s, 1H), 6.46 (d, J=9.9 Hz, 1H), 5.36 (s, 1H), 5.06 (dd, J=4.1, 7.8 Hz, 1H), 4.80 - 4.71 (m, 2H), 4.28 (d, J=12.5 Hz, 1H), 4.04 (dd, J=3.2, 7.2 Hz, 1H), 4.00 (d, J=6.3 Hz, 2H), 3.85 - 3.81 (m, 1H), 3.79 (s, 2H), 3.13 (dd, J=11.7, 11.7 Hz, 1H), 2.76 - 2.59 (m, 5H), 1.86 - 1.77 (m, 4H), 1.54 - 1.44 (m, 6H), 1.15 - 1.06 (m, 2H).	Free Base
60	2.39	11	(DMSO-d ₆); δ 10.33 (s, 1H), 8.65 and 8.46 (s, 1H), 8.12 (dd, J=4.6, 10.0 Hz, 1H), 7.77 (dd, J=8.3, 10.3 Hz, 2H), 7.40 (dd, J=5.7, 7.8 Hz, 2H), 7.34 - 7.23 (m, 9H), 7.14 (dd, J=8.0, 8.0 Hz, 1H), 7.05 (dd, J=8.4, 8.4 Hz, 2H), 6.92 - 6.78 (m, 3H), 6.68 - 6.57 (m, 1H), 6.47 (d, J=9.8 Hz, 1H), 5.41 (s, 1H), 5.10 - 5.06 (m, 1H), 4.74 (d, J=12.4 Hz, 2H), 4.01 - 3.96 (m, 2H), 3.81 (s, 2H), 3.46 (s, 2H), 3.38 (s, 3H), 3.00 (s, 2H), 2.85 (s, 2H), 2.74 - 2.66 (m, 5H), 1.85 - 1.79 (m, 2H), 1.51 - 1.44 (m, 2H), 1.15 - 1.06 (m, 2H).	Free Base
61	2.50	11	(DMSO-d ₆); δ 10.51(s, 1H), 10.49 (s, 1H), 9.40 (s, 1H), 9.15 (s, 2H), 8.58 (dd, J=5.7, 5.7 Hz, 1H), 8.07 (d, J=7.9 Hz, 1H), 7.91 (d, J=8.0 Hz, 2H), 7.62 (d, J=8.4 Hz, 2H), 7.46 (s, 5H), 7.34 - 7.32 (m, 3H), 7.32 - 7.29 (m, 1H), 7.23 (dd, J=8.0, 8.0 Hz, 1H), 7.12 (d, J=8.2 Hz, 1H), 6.97 (d, J=8.2 Hz, 1H), 6.90 (d, J=7.9 Hz, 1H), 6.86 (s, 1H), 6.83 (d, J=8.3 Hz, 1H), 6.63 (s, 1H), 6.58 - 6.53 (m, 2H), 6.19 (d, J=2.5 Hz, 1H), 5.37 - 5.36 (m, 1H), 4.75 (d, J=3.5 Hz, 2H), 4.29 - 4.28 (m, 6H), 4.02 (d, J=6.1 Hz, 2H), 3.83 - 3.83 (m, 1H), 3.15 - 3.14 (m, 5H), 3.02 (s, 4H), 1.82 - 1.81 (m, 1H), 1.71 (s, 6H), 1.35 - 1.35 (m, 3H).	TFA

(continued)

62	2.50	11	(DMSO-d ₆); δ 10.51 (s, 1H), 10.47 (s, 1H), 9.38 (s, 1H), 9.13 (s, 2H), 8.29 (d, J=7.8 Hz, 1H), 8.07 (d, J=10.1 Hz, 1H), 7.96 (d, J=8.0 Hz, 1H), 7.89 (d, J=8.3 Hz, 2H), 7.60 (d, J=8.4 Hz, 2H), 7.48 (m, 5H), 7.34 - 7.32 (m, 6H), 7.12 (d, J=8.3 Hz, 1H), 7.00 - 6.93 (m, 3H), 6.88 (dd, J=2.4, 8.2 Hz, 1H), 6.66 (s, 1H), 6.57 (dd, J=2.1, 9.9 Hz, 1H), 6.19 - 6.19 (m, 1H), 5.33 (d, J=10.0 Hz, 1H), 4.41 (s, 2H), 4.31 - 4.21 (m, 4H), 4.04 - 3.97 (m, 2H), 3.75 - 3.74 (m, 1H), 3.61 - 3.61 (m, 1H), 3.33 (d, J=11.8 Hz, 2H), 3.09 - 3.08 (m, 2H), 3.01 - 2.87 (m, 3H), 1.88 - 1.67 (m, 6H), 1.43 - 1.34 (m, 6H).	TFA
63	2.43	11	(DMSO-d ₆); δ 10.51 (s, 1H), 10.47 (s, 1H), 9.38 (s, 1H), 9.13 (s, 2H), 8.79 (d, J=7.0 Hz, 1H), 8.51 (d, J=7.4 Hz, 1H), 8.08 (d, J=10.0 Hz, 1H), 7.93 (d, J=8.4 Hz, 2H), 7.62 (d, J=8.3 Hz, 2H), 7.48 (s, 5H), 7.35 - 7.26 (m, 6H), 7.12 (d, J=8.3 Hz, 1H), 7.02 - 6.93 (m, 3H), 6.89 (dd, J=2.6, 7.9 Hz, 1H), 6.67 - 6.66 (m, 1H), 6.58 (d, J=9.3 Hz, 1H), 6.19 - 6.19 (m, 1H), 5.36 - 5.34 (m, 1H), 4.43 (s, 4H), 4.31 - 4.21 (m, 5H), 4.03 (d, J=6.5 Hz, 2H), 3.34 (d, J=11.5 Hz, 2H), 3.11 - 2.87 (m, 3H), 2.35 (dd, J=6.9, 6.9 Hz, 4H), 1.77 - 1.67 (m, 3H), 1.40 - 1.31 (m, 2H).	TFA
64	2.46	11	(DMSO-d ₆); δ 10.60 (s, 1H), 10.50 (s, 1H), 9.40 (s, 1H), 9.10 (s, 2H), 8.06 (d, J=10.4 Hz, 1H), 7.60 (d, J=8.2 Hz, 2H), 7.48 (s, 5H), 7.45 (d, J=8.2 Hz, 2H), 7.36 - 7.27 (m, 5H), 7.23 (t, J=7.9 Hz, 1H), 7.13 (d, J=8.3 Hz, 1H), 6.98 (d, J=8.2 Hz, 1H), 6.91 - 6.87 (m, 2H), 6.86 - 6.81 (m, 1H), 6.64 (s, 1H), 6.59 - 6.55 (m, 1H), 6.15 (s, 1H), 5.35 - 5.33 (m, 1H), 4.76 (s, 2H), 4.32 - 4.22 (m, 4H), 4.04 - 3.97 (m, 2H), 3.62 (s, 2H), 3.44 - 3.26 (m, 8H), 3.10 - 3.02 (m, 3H), 2.95 - 2.86 (m, 2H), 1.76 - 1.66 (m, 2H), 1.50 (s, 4H), 1.42 - 1.33 (m, 6H).	TFA

(continued)

65	2.43	11	(DMSO-d ₆); δ 10.55 (s, 1H), 10.45 (s, 1H), 9.40 (s, 1H), 8.75 (s, 2H), 8.08 (d, J=9.9 Hz, 1H), 8.04 (d, J=7.4 Hz, 1H), 7.61 (d, J=8.2 Hz, 2H), 7.47 (s, 5H), 7.44 (d, J=7.4 Hz, 2H), 7.32 (d, J=4.4 Hz, 4H), 7.30 - 7.23 (m, 2H), 7.13 (d, J=8.3 Hz, 1H), 7.00 - 6.92 (m, 3H), 6.87 (dd, J=2.3, 8.1 Hz, 1H), 6.65 (s, 1H), 6.58 (d, J=9.9 Hz, 1H), 6.15 (s, 1H), 5.35 - 5.33 (m, 1H), 4.42 (s, 2H), 4.32 - 4.20 (m, 4H), 4.03 - 3.97 (m, 2H), 3.93 - 3.88 (m, 1H), 3.48 - 3.48 (m, 1H), 3.34 (d, J=11.9 Hz, 2H), 3.11 - 3.02 (m, 4H), 2.94 - 2.86 (m, 3H), 1.85 - 1.84 (m, 1H), 1.76 - 1.66 (m, 4H), 1.39 - 1.30 (m, 4H).	TFA
66	2.39	11	(MeOD); δ 8.29 (d, J=9.8 Hz, 1H), 7.75 (d, J=7.9 Hz, 2H), 7.63 (d, J=6.9 Hz, 2H), 7.53 - 7.48 (m, 4H), 7.39 (d, J=7.2 Hz, 2H), 7.34 - 7.27 (m, 6H), 7.09 (s, 1H), 7.04 (d, J=8.3 Hz, 2H), 6.98 (dd, J=2.2, 8.0 Hz, 1H), 6.66 (d, J=9.3 Hz, 1H), 5.43 (dd, J=6.7, 6.7 Hz, 1H), 4.76 - 4.69 (m, 1H), 4.60 (dd, J=7.9, 7.9 Hz, 1H), 4.52 (s, 2H), 4.46 (dd, J=9.1, 9.1 Hz, 1H), 4.28 (s, 3H), 4.12 (d, J=6.4 Hz, 3H), 3.51 - 3.45 (m, 2H), 3.27 (d, J=6.7 Hz, 2H), 3.01 - 2.91 (m, 2H), 2.68 (s, 3H), 1.83 (d, J=14.1 Hz, 2H), 1.42 (d, J=12.9 Hz, 2H).	TFA
67	2.42	11	(MeOD); δ 8.52 (s, 2H), 8.29 (d, J=9.9 Hz, 1H), 7.91 (d, J=8.0 Hz, 2H), 7.56 (d, J=8.3 Hz, 2H), 7.42 - 7.38 (m, 6H), 7.37 - 7.23 (m, 6H), 7.09 - 7.05 (m, 1H), 7.03 - 6.97 (m, 2H), 6.93 (dd, J=2.5, 8.2 Hz, 1H), 6.63 (d, J=9.8 Hz, 1H), 5.35 (dd, J=4.3, 8.8 Hz, 1H), 4.84 - 4.76 (m, 1H), 4.70 - 4.59 (m, 3H), 4.41 - 4.27 (m, 2H), 4.18 - 4.10 (m, 5H), 3.94 (d, J=2.3 Hz, 2H), 3.22 - 3.04 (m, 4H), 2.56 (dd, J=12.4, 12.4 Hz, 2H), 1.85 - 1.79 (m, 1H), 1.71 - 1.64 (m, 2H), 1.43 - 1.30 (m, 2H).	Formate
68	2.38	11	(DMSO-d ₆ , 100 °C) δ 8.12 (d, J=9.8 Hz, 1H), 7.99 (d, J=6.3 Hz, 1H), 7.58 (q, J=8.6 Hz, 4H), 7.50 - 7.47 (m, 5H), 7.41 - 7.24 (m, 6H), 7.14 (d, J=8.3 Hz, 1H), 7.04 - 7.00 (m, 3H), 6.90 (dd, J=2.2, 8.0 Hz, 1H), 6.56 (d, J=9.9 Hz, 1H), 5.39 (dd, J=4.6, 8.3 Hz, 1H), 4.44 (s, 2H), 4.34 - 4.30 (m, 3H), 4.26 (s, 2H), 4.13 - 4.06 (m, 2H), 3.76 - 3.70 (m, 2H), 3.37 - 3.29 (m, 4H), 3.23 - 3.15 (m, 2H), 2.95 - 2.91 (m, 2H), 2.19 - 2.10 (m, 1H), 1.97 - 1.87 (m, 2H), 1.80 - 1.76 (m, 2H), 1.49 - 1.49 (m, 2H).	TFA

(continued)

69	2.46	11	(DMSO-d ₆ , 100 °C) δ 8.19 - 8.15 (m, 3H), 7.57 (d, J=7.5 Hz, 1H), 7.40 - 7.23 (m, 15H), 7.10 (d, J=8.3 Hz, 1H), 7.06 - 7.00 (m, 2H), 6.96 (d, J=8.2 Hz, 1H), 6.90 (dd, J=2.3, 8.0 Hz, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.09 (dd, J=4.6, 7.7 Hz, 1H), 4.41 (d, J=2.8 Hz, 2H), 4.05 (d, J=6.3 Hz, 2H), 3.92 (d, J=14.0 Hz, 1H), 3.81 (s, 2H), 3.72 (d, J=13.4 Hz, 1H), 3.44 (s, 2H), 3.13 - 3.02 (m, 3H), 2.88 - 2.67 (m, 4H), 1.97 - 1.86 (m, 3H), 1.71 - 1.49 (m, 6H), 1.28 - 1.13 (m, 2H).	Formate
70	2.42	11	(DMSO-d ₆ , 100 °C) δ 8.19 - 8.15 (m, 3H), 7.92 (d, J=6.4 Hz, 1H), 7.45 - 7.23 (m, 15H), 7.09 (d, J=8.0 Hz, 1H), 7.04 - 7.00 (m, 2H), 6.96 (d, J=8.0 Hz, 1H), 6.90 (dd, J=2.4, 8.1 Hz, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.08 (dd, J=4.7, 7.6 Hz, 1H), 4.43 (s, 2H), 4.40 - 4.30 (m, 1H), 4.05 (d, J=6.4 Hz, 2H), 3.82 (s, 2H), 3.70 (dd, J=6.7, 11.4 Hz, 1H), 3.62 - 3.47 (m, 2H), 3.45 (s, 2H), 3.36 (dd, J=5.3, 11.5 Hz, 1H), 2.88 - 2.71 (m, 4H), 2.20 - 2.07 (m, 1H), 1.98 - 1.85 (m, 3H), 1.62 - 1.49 (m, 3H), 1.28 - 1.13 (m, 2H).	Formate
71	2.41	11	(DMSO-d ₆ , 100 °C) δ 8.11 (d, J=9.9 Hz, 1H), 7.66 (d, J=7.7 Hz, 1H), 7.60 (d, J=8.2 Hz, 2H), 7.46 (d, J=8.5 Hz, 7H), 7.41 - 7.24 (m, 6H), 7.14 (d, J=8.2 Hz, 1H), 7.06 - 7.00 (m, 3H), 6.90 (dd, J=2.3, 8.2 Hz, 1H), 6.56 (d, J=9.9 Hz, 1H), 5.38 (dd, J=4.7, 8.2 Hz, 1H), 4.42 (s, 2H), 4.31 (d, J=2.4 Hz, 2H), 4.23 (s, 2H), 4.10 (d, J=5.6 Hz, 2H), 4.00 - 3.95 (m, 1H), 3.85 - 3.74 (m, 2H), 3.50 - 3.25 (m, 2H), 3.23 - 2.98 (m, 5H), 2.57 - 2.53 (m, 1H), 1.92 - 1.87 (m, 2H), 1.81 - 1.71 (m, 3H), 1.66 - 1.47 (m, 4H).	TFA
72	2.43	11	(DMSO-d ₆ , 100 °C) δ 8.19 - 8.12 (m, 2H), 7.86 (d, J=6.5 Hz, 1H), 7.78 (d, J=8.0 Hz, 2H), 7.39 (d, J=7.9 Hz, 4H), 7.35 - 7.20 (m, 8H), 7.09 (d, J=8.2 Hz, 1H), 7.01 - 6.94 (m, 3H), 6.90 - 6.86 (m, 1H), 6.48 (d, J=9.8 Hz, 1H), 5.11 (dd, J=4.6, 7.5 Hz, 1H), 4.72 (s, 2H), 4.04 (d, J=6.4 Hz, 3H), 3.85 (br s, 4H), 3.46 (s, 2H), 2.89 - 2.71 (m, 5H), 2.54 - 2.40 (m, 1H), 2.06 - 1.92 (m, 3H), 1.81 - 1.77 (m, 1H), 1.69 - 1.57 (m, 2H), 1.52 - 1.50 (m, 3H), 1.24 - 1.14 (m, 2H).	mono-Formate










(continued)

73	2.45	11	(DMSO-d ₆ , 100 °C) δ 8.17 (d, J=7.9 Hz, 3H), 7.79 (d, J=8.3 Hz, 2H), 7.41 - 7.36 (m, 4H), 7.35 - 7.21 (m, 9H), 7.09 (d, J=8.2 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.89 (dd, J=1.9, 8.2 Hz, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.08 (dd, J=4.9, 7.7 Hz, 1H), 4.64 (s, 2H), 4.54 - 4.45 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.85 - 3.21 (m, 3H), 3.65 - 3.59 (m, 1H), 3.46 - 3.43 (m, 3H), 2.87 - 2.67 (m, 4H), 2.56 - 2.47 (m, 1H under the DMSO), 2.17 - 2.17 (m, 1H), 2.06 - 2.00 (m, 1H), 1.98 - 1.90 (m, 2H), 1.62 - 1.49 (m, 3H), 1.25 - 1.13 (m, 2H).	mono-Formate
74	2.40	11	(DMSO-d ₆ , 100 °C) δ 8.17 (d, J=9.9 Hz, 1H), 8.15 (s, 1H), 7.40 - 7.20 (m, 15H), 7.10 (d, J=8.3 Hz, 1H), 7.01 - 6.94 (m, 3H), 6.87 - 6.84 (m, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.09 (dd, J=4.6, 7.7 Hz, 1H), 4.63 (s, 2H), 4.43 - 4.34 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.81 (s, 2H), 3.76 - 3.49 (m, 4H), 3.45 - 3.39 (m, 3H), 2.88 - 2.68 (m, 5H), 2.57 - 2.51 (m, 1H), 2.06 - 1.99 (m, 1H), 1.97 - 1.90 (m, 2H), 1.78 - 1.77 (m, 1H), 1.62 - 1.49 (m, 3H), 1.28 - 1.13 (m, 2H).	mono-Formate
75	2.43	11	(DMSO-d ₆ , 100 °C) δ 8.21 (s, 1H), 8.17 (d, J=9.9 Hz, 1H), 7.86 (d, J=6.3 Hz, 1H), 7.77 (d, J=8.3 Hz, 2H), 7.41 - 7.20 (m, 13H), 7.09 (d, J=8.2 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.91 - 6.87 (m, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.08 (dd, J=4.7, 7.6 Hz, 1H), 4.73 (s, 2H), 4.05 (d, J=6.4 Hz, 3H), 3.87 - 3.78 (m, 4H), 3.44 (s, 2H), 2.87 - 2.67 (m, 6H), 2.57 - 2.46 (m, 1H), 2.11 - 1.89 (m, 3H), 1.82 - 1.56 (m, 3H), 1.56 - 1.49 (m, 2H), 1.24 - 1.13 (m, 2H).	mono-Formate
76	2.40	11	(DMSO-d ₆ , 100 °C) δ 8.19 - 8.14 (m, 3H), 7.82 (d, J=8.3 Hz, 2H), 7.46 - 7.38 (m, 4H), 7.35 - 7.23 (m, 9H), 7.10 (d, J=8.2 Hz, 1H), 7.00 - 6.95 (m, 3H), 6.90 (dd, J=2.0, 8.2 Hz, 1H), 6.49 (d, J=9.9 Hz, 1H), 6.07 (s, 1H), 5.16 (dd, J=4.6, 7.8 Hz, 1H), 4.64 (s, 2H), 4.54 - 4.44 (m, 1H), 4.06 (d, J=6.3 Hz, 2H), 3.93 (s, 2H), 3.74 - 3.60 (m, 2H), 3.51 (s, 2H), 3.46 - 3.39 (m, 1H), 2.90 - 2.77 (m, 4H), 2.57 - 2.47 (m, 1H), 2.25 - 2.12 (m, 1H), 2.04 - 1.97 (m, 3H), 1.65 - 1.51 (m, 3H), 1.28 - 1.16 (m, 2H).	mono-Formate

(continued)

77	2.40	11	(DMSO-d ₆ , 100 °C) δ 8.19 - 8.15 (m, 2H), 7.40 - 7.20 (m, 15H), 7.10 (d, J=8.3 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.85 (dd, J=1.9, 8.0 Hz, 1H), 6.47 (d, J=9.9 Hz, 1H), 5.09 (dd, J=4.6, 7.7 Hz, 1H), 4.63 (s, 2H), 4.44 - 4.37 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.82 (s, 2H), 3.79 - 3.73 (m, 1H), 3.72 - 3.49 (m, 3H), 3.45 - 3.42 (m, 3H), 2.95 - 2.71 (m, 5H), 2.57 - 2.50 (m, 1H), 2.08 - 2.01 (m, 1H), 2.03 - 1.89 (m, 2H), 1.80 - 1.74 (m, 1H), 1.62 - 1.49 (m, 3H), 1.24 - 1.14 (m, 2H).	mono- Formate
----	------	----	--	------------------

The following compounds were prepared using the appropriate amine and protected acid in the final coupling step.

N	Appropriate amine /	protected acid	Structure
78			
79			
80			

(continued)

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
78	2.39	11	(MeOD); δ 8.46 (s, 2H), 8.26 (dd, J=4.0, 9.9 Hz, 1H), 7.48 (d, J=7.8 Hz, 1H), 7.44 (s, 5H), 7.41 - 7.37 (m, 2H), 7.36 - 7.24 (m, 7H), 7.09 - 7.02 (m, 3H), 6.99 - 6.95 (m, 1H), 6.64 (d, J=9.6 Hz, 1H), 5.44 - 5.38 (m, 1H), 4.77 - 4.68 (m, 1H), 4.62 (t, J=8.6 Hz, 1H), 4.52 (s, 2H), 4.50 - 4.42 (m, 1H), 4.33 - 4.26 (m, 3H), 4.18 - 4.10 (m, 3H), 4.07 (s, 2H), 3.95 (s, 3H), 3.27 (d, J=12.4 Hz, 2H), 3.19 (d, J=8.6 Hz, 2H), 2.72 (ddd, J=12.6, 12.6, 2.5 Hz, 2H), 1.94 - 1.82 (m, 1H), 1.74 (d, J=12.9 Hz, 2H), 1.47 - 1.36 (m, 2H).	Formate
79	2.47	11	(MeOD); δ 8.55 (s, 1H), 8.28 (d, J=9.9 Hz, 1H), 7.45 - 7.23 (m, 13H), 7.08 (d, J=1.3 Hz, 1H), 7.06 - 6.99 (m, 4H), 6.93 (dq, J=0.7, 3.6 Hz, 1H), 6.65 (d, J=9.9 Hz, 1H), 5.34 (dd, J=5.2, 8.0 Hz, 1H), 4.77 (s, 2H), 4.18 (s, 2H), 4.10 (d, J=6.3 Hz, 2H), 3.91 (s, 3H), 3.77 (s, 4H), 3.61 - 3.50 (m, 4H), 3.42 (s, 2H), 3.10 - 3.02 (m, 4H), 2.33 (dd, J=10.2, 11.7 Hz, 2H), 1.80 - 1.69 (m, 1H), 1.63 (d, J=15.7 Hz, 6H), 1.57 - 1.47 (m, 4H), 1.38 - 1.24 (m, 2H).	mono-formate
80	2.44	11	(DMSO-d ₆); δ 10.32 (s, 2H), 9.23 - 9.19 (m, 1H), 8.72 - 8.67 (m, 2H), 8.45 (dd, J=5.7, 5.7 Hz, 1H), 7.89 (d, J=11.9 Hz, 1H), 7.35 (d, J=2.8 Hz, 3H), 7.30 (s, 6H), 7.16 (d, J=4.3 Hz, 7H), 6.97 (d, J=8.3 Hz, 1H), 6.81 (d, J=8.1 Hz, 1H), 6.75 - 6.69 (m, 2H), 6.65 (dd, J=2.4, 8.2 Hz, 1H), 6.40 (d, J=9.9 Hz, 1H), 5.18 (dd, J=4.5, 7.6 Hz, 1H), 4.59 (d, J=3.3 Hz, 2H), 4.12 - 4.03 (m, 6H), 3.87 - 3.79 (m, 4H), 3.67 - 3.63 (m, 1H), 3.16 (d, J=11.4 Hz, 2H), 3.06 - 2.98 (m, 2H), 2.92 - 2.68 (m, 5H), 1.67 - 1.64 (m, 2H), 1.61 - 1.48 (m, 5H), 1.24 - 1.13 (m, 2H), 1.03 - 0.97 (m, 1H), 0.84 (dd, J=6.8, 12.1 Hz, 1H).	TFA
81	2.49	11	(MeOD); δ 8.54 (s, 1H), 8.32 (d, J=9.9 Hz, 1H), 7.45 - 7.26 (m, 13H), 7.23 (d, J=4.5 Hz, 1H), 7.16 - 7.10 (m, 1H), 7.07 - 7.02 (m, 2H), 7.00 (s, 1H), 6.94 (dd, J=2.3, 8.2 Hz, 1H), 6.66 (d, J=9.8 Hz, 1H), 5.38 (dd, J=4.6, 8.4 Hz, 1H), 4.78 (s, 2H), 4.19 (s, 2H), 4.12 (dd, J=2.6, 6.2 Hz, 2H), 3.96 (s, 2H), 3.90 (s, 3H), 3.81 - 3.74 (m, 2H), 3.58 - 3.52 (m, 4H), 3.28 - 3.08 (m, 6H), 2.57 (dd, J=11.4, 11.4 Hz, 2H), 1.87 - 1.79 (m, 1H), 1.70 - 1.31 (m, 12H).	mono-formate

82	2.41	11	(MeOD); δ 8.50 (s, 2H), 8.33 (d, J=9.8 Hz, 1H), 7.44 (s, 5H), 7.41 - 7.25 (m, 9H), 7.14 (d, J=7.4 Hz, 1H), 7.10 - 7.02 (m, 3H), 6.97 (dd, J=2.4, 8.2 Hz, 1H), 6.67 (d, J=9.7 Hz, 1H), 5.41 (dd, J=6.2, 6.2 Hz, 1H), 4.73 - 4.65 (m, 1H), 4.53 (s, 2H), 4.46 - 4.39 (m, 1H), 4.24 (s, 2H), 4.23 - 4.18 (m, 1H), 4.13 (d, J=6.2 Hz, 2H), 4.11 - 4.07 (m, 1H), 4.05 (s, 2H), 4.00 - 3.95 (m, 1H), 3.93 (s, 3H), 3.26 (d, J=12.2 Hz, 2H), 3.16 (d, J=6.3 Hz, 2H), 2.69 (dd, J=11.6, 11.6 Hz, 2H), 1.89 - 1.85 (m, 1H), 1.75 (d, J=12.8 Hz, 2H), 1.48 - 1.30 (m, 2H).	formate
----	------	----	---	---------

The following compounds were prepared as described in **Example 20** with **(R)-4-(2-((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoic acid** (prepared in **Procedure 5**) replacing **(R)-4-(((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid** and the use of the appropriate amine replacing N-(*tert*-butoxycarbonyl)ethylenediamine HCl in **Step 1**.










N	Appropriate amine	Structure
83		
84		
85		

(continued)

(continued)

89	90	91

(continued)

92			
93			
94			

(continued)









 <chem>COC(=O)N1CCN(C1Cc2ccc(cc2)C(=O)N3CCN(CC3Cc4cc(O)c5c(c4)nc6c(nc7c6cnc7C(=O)N8CCCCN8Cc9ccccc9)cc5)cc4)CCN(C1)Cc2ccccc2</chem>	 <chem>COC(=O)N1CCN(C1Cc2ccc(cc2)C(=O)N3CCN(C3Cc4cc(O)c5c(c4)nc6c(nc7c6cnc7C(=O)N8CCCCN8Cc9ccccc9)cc5)cc4)CCN(C1)Cc2ccccc2</chem>	 <chem>COC(=O)N1CCN(C1Cc2ccc(cc2)C(=O)N3CCN(C3Cc4cc(O)c5c(c4)nc6c(nc7c6cnc7C(=O)N8CCCCN8Cc9ccccc9)cc5)cc4)CCN(C1)Cc2ccccc2</chem>
 <chem>CC(C)(C)OC(=O)N1CCN(C1C2CCNCC2)C3CCNCC3</chem>	 <chem>CC(C)(C)OC(=O)N1CCN(C1C2CCNCC2)C3CCNCC3</chem>	 <chem>CC(C)(C)OC(=O)N1CCN(C1C2CCNCC2)C3CCNCC3</chem>
95	96	97

(continued)




[illegible]

(continued)


(continued)

106		
107		
108		
109		

(continued)

		
110	111	112

(continued)



N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
83	2.45	11	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.82 (d, J=8.3 Hz, 1H), 7.75 (d, J=8.3 Hz, 1H), 7.48 (s, 5H), 7.43 - 7.29 (m, 9H), 7.23 - 7.15 (m, 1H), 7.04 (dd, J=8.3, 8.3 Hz, 2H), 6.93 (dd, J=2.2, 2.2 Hz, 1H), 6.89 - 6.81 (m, 1H), 6.70 (d, J=9.9 Hz, 1H), 5.42 (dd, J=6.7, 6.7 Hz, 1H), 4.80 (ddd, J=12.7, 12.7, 12.7 Hz, 2H), 4.28 (t, J=9.7 Hz, 2H), 4.13 - 4.06 (m, 2H), 3.59 - 3.43 (m, 7H), 3.37 - 3.35 (m, 2H), 3.29 (d, J=5.8 Hz, 2H), 3.16 - 3.08 (m, 3H), 3.01 - 2.92 (m, 2H), 1.96 - 1.95 (m, 1H), 1.81 - 1.76 (m, 2H), 1.42 - 1.30 (m, 2H), 1.25 - 1.12 (m, 3H).	TFA
84	2.40	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.81 (d, J=8.3 Hz, 1H), 7.74 (d, J=8.2 Hz, 1H), 7.51 - 7.46 (m, 6H), 7.41 - 7.29 (m, 8H), 7.15 (q, J=7.9 Hz, 1H), 7.05 (d, J=8.3 Hz, 1H), 7.00 (d, J=8.2 Hz, 1H), 6.93 - 6.91 (m, 1H), 6.82 (ddd, J=5.7, 10.6, 17.0 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.45 - 5.40 (m, 1H), 4.84 - 4.76 (m, 2H), 4.26 (d, J=8.0 Hz, 2H), 4.11 - 4.05 (m, 2H), 3.61 - 3.35 (m, 7H), 3.29 (d, J=7.5 Hz, 2H), 3.12 (s, 3H), 3.08 (d, J=7.8 Hz, 1H), 3.00 (s, 1H), 2.99 - 2.92 (m, 2H), 1.95 - 1.89 (m, 1H), 1.80 - 1.69 (m, 3H), 1.43 - 1.30 (m, 2H).	TFA
85	2.43	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.84 (d, J=8.4 Hz, 2H), 7.52 - 7.48 (m, 6H), 7.40 (dd, J=8.6, 8.6 Hz, 4H), 7.35 - 7.28 (m, 6H), 7.07 - 7.04 (m, 3H), 6.98 (dd, J=2.1, 8.8 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.45 - 5.40 (m, 1H), 4.51 (s, 2H), 4.27 (s, 2H), 4.11 (d, J=6.3 Hz, 2H), 3.49 - 3.35 (m, 6H), 3.32 - 3.28 (m, 3H), 3.19 - 3.10 (m, 3H), 2.97 (dd, J=10.9, 12.7 Hz, 2H), 2.01 - 1.93 (m, 1H), 1.84 - 1.77 (m, 4H), 1.47 - 1.37 (m, 2H).	TFA
86	2.43	11	(MeOD); δ 8.38 (1H, d, J=9.9 Hz), 7.53 - 7.49 (6H, m), 7.46 (4H, d, J=4.3 Hz), 7.41 - 7.26 (6H, m), 7.06 (2H, d, J=8.2 Hz), 6.97 (1H, s), 6.94 (1H, d, J=7.2 Hz), 6.71 (1H, d, J=9.9 Hz), 5.43 (1H, dd, J=5.6, 7.8 Hz), 4.84 (2H, s), 4.28 (2H, s), 4.17 - 4.08 (1H, m), 3.69 (6H, m), 3.51 - 3.37 (4H, m), 3.30 (2H, s), 3.20 - 3.11 (3H, m), 3.01 - 2.93 (2H, m), 1.98 - 1.92 (1H, m), 1.84 - 1.77 (4H, m), 1.47 - 1.33 (2H, m)	TFA

(continued)

87	2.57	11	(MeOD); δ 8.37 (dd, J=2.1, 9.9 Hz, 1H), 7.81 (d, J=8.4 Hz, 1H), 7.75 (d, J=8.2 Hz, 1H), 7.42 - 7.23 (m, 16H), 7.22 - 7.14 (m, 1H), 7.07 - 6.95 (m, 4H), 6.87 (dd, J=1.9, 7.3 Hz, 1H), 6.79 (dd, J=2.3, 8.1 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.42 (dd, J=6.3, 6.3 Hz, 1H), 5.02 - 4.97 (m, 1H), 4.81 (t, J=4.4 Hz, 1H), 4.70 (d, J=5.9 Hz, 2H), 4.26 (s, 2H), 4.19 (s, 2H), 3.63 - 3.49 (m, 5H), 3.16 - 3.10 (m, 2H), 2.93 - 2.93 (m, 3H), 1.79 (dt, J=15.4, 39.6 Hz, 5H), 1.31 (s, 3H).	TFA
88	2.45	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.86 (d, J=8.4 Hz, 2H), 7.53 - 7.48 (m, 5H), 7.43 (d, J=8.4 Hz, 2H), 7.41 - 7.29 (m, 7H), 7.10 - 7.04 (m, 3H), 6.98 (dd, J=2.0, 8.2 Hz, 1H), 6.71 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.3, 8.2 Hz, 1H), 4.60 - 4.43 (m, 5H), 4.28 (s, 2H), 4.13 (d, J=6.4 Hz, 2H), 3.50 - 3.41 (m, 4H), 3.20 - 3.11 (m, 3H), 3.01 - 2.95 (m, 2H), 2.54 - 2.42 (m, 4H), 2.05 - 1.75 (m, 3H), 1.50 - 1.32 (m, 2H).	TFA
89	2.46	11	(MeOD); δ 8.54 (d, J=1.9 Hz, 1H), 8.37 (d, J=9.9 Hz, 1H), 7.84 (d, J=8.3 Hz, 2H), 7.53 - 7.48 (m, 5H), 7.43 (d, J=8.3 Hz, 2H), 7.41 - 7.29 (m, 6H), 7.29 - 7.24 (m, 1H), 7.05 (d, J=8.2 Hz, 2H), 6.95 - 6.89 (m, 2H), 6.70 (d, J=9.9 Hz, 1H), 5.42 (dd, J=5.8, 7.8 Hz, 1H), 4.83 - 4.72 (m, 2H), 4.42 (s, 1H), 4.27 (s, 2H), 4.18 - 4.10 (m, 2H), 3.94 (d, J=12.8 Hz, 1H), 3.49 - 3.35 (m, 7H), 3.20 - 3.07 (m, 4H), 2.97 (dd, J=12.5, 12.5 Hz, 2H), 2.67 (s, 1H), 1.95 (d, J=3.0 Hz, 2H), 1.78 (dd, J=13.9, 15.6 Hz, 4H), 1.45 - 1.31 (m, 2H), 1.25 - 1.11 (m, 2H).	TFA
90	2.51	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.83 (d, J=8.3 Hz, 2H), 7.53 - 7.48 (m, 6H), 7.44 - 7.28 (m, 8H), 7.09 - 7.04 (m, 3H), 6.96 (dd, J=2.3, 8.3 Hz, 1H), 6.71 (d, J=9.9 Hz, 1H), 5.42 (dd, J=5.3, 8.2 Hz, 1H), 4.48 (s, 2H), 4.29 (s, 2H), 4.13 (d, J=6.3 Hz, 2H), 3.90 - 3.72 (m, 2H), 3.38 (d, J=6.3 Hz, 5H), 3.30 (s, 1H), 3.16 - 3.10 (m, 3H), 3.01 - 2.94 (m, 2H), 2.01 (s, 2H), 1.95 - 1.81 (m, 5H), 1.52 - 1.41 (m, 5H).	TFA

(continued)

91	2.46	11	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.53 - 7.48 (m, 6H), 7.43 (s, 4H), 7.40 - 7.27 (m, 6H), 7.09 - 7.03 (m, 3H), 6.96 (dd, J=2.3, 8.1 Hz, 1H), 6.71 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.3, 8.1 Hz, 1H), 4.58 - 4.44 (m, 2H), 4.35 - 4.27 (m, 3H), 4.28 (s, 2H), 4.12 (d, J=6.3 Hz, 2H), 4.05 - 3.98 (m, 1H), 3.73 - 3.68 (m, 1H), 3.51 - 3.40 (m, 2H), 3.24 - 3.09 (m, 3H), 3.06 - 2.93 (m, 3H), 2.06 - 1.67 (m, 6H), 1.61 - 1.36 (m, 5H);	TFA
92	2.44	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.85 (d, J=6.8 Hz, 2H), 7.53 - 7.49 (m, 4H), 7.45 - 7.25 (m, 11H), 7.08 - 6.91 (m, 3H), 6.70 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.6, 8.0 Hz, 1H), 4.85 - 4.77 (m, 2H), 4.52 - 4.43 (m, 1H), 4.28 (s, 2H), 4.19 - 4.08 (m, 2H), 4.01 - 3.97 (m, 1H), 3.50 - 3.35 (m, 6H), 3.30 - 3.08 (m, 4H), 3.04 - 2.83 (m, 3H), 2.08 - 1.95 (m, 3H), 1.79 (s, 1H), 1.65 - 1.50 (m, 2H), 1.42 - 1.31 (m, 2H).	TFA
93	2.53	11	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.66 (d, J=8.3 Hz, 2H), 7.51 - 7.50 (m, 5H), 7.43 (d, J=8.2 Hz, 2H), 7.41 - 7.35 (m, 6H), 7.33 - 7.26 (m, 1H), 7.09 - 7.03 (m, 2H), 6.97 - 6.89 (m, 2H), 6.70 (d, J=9.9 Hz, 1H), 5.43 (dd, J=5.8, 7.6 Hz, 1H), 4.54 - 4.41 (m, 4H), 4.30 (d, J=14.1 Hz, 4H), 4.19 - 4.10 (m, 4H), 3.47 - 3.35 (m, 4H), 3.29 (d, J=7.3 Hz, 2H), 3.20 - 3.10 (m, 3H), 3.01 - 2.88 (m, 2H), 1.95 (d, J=3.8 Hz, 1H), 1.80 - 1.78 (m, 2H), 1.49 - 1.29 (m, 3H)	TFA
94	2.54	11	(MeOD); δ 8.37 (d, J=9.9 Hz, 1H), 7.69 (d, J=8.3 Hz, 2H), 7.53 - 7.48 (m, 5H), 7.44 (d, J=8.2 Hz, 4H), 7.40 - 7.25 (m, 5H), 7.05 (d, J=8.2 Hz, 2H), 6.94 - 6.90 (m, 2H), 6.71 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.1, 8.3 Hz, 1H), 4.79 (dd, J=4.1, 4.1 Hz, 2H), 4.28 (s, 2H), 4.14 - 4.10 (m, 4H), 3.95 (s, 2H), 3.59 - 3.37 (m, 6H), 3.17 - 3.10 (m, 3H), 3.02 - 2.94 (m, 2H), 1.96 (s, 1H), 1.85 - 1.74 (m, 9H), 1.46 - 1.35 (m, 2H).	TFA

(continued)

95	2.60	11	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.53 - 7.47 (m, 6H), 7.43 (s, 4H), 7.41 - 7.25 (m, 6H), 7.05 (d, J=8.2 Hz, 2H), 6.97 - 6.90 (m, 2H), 6.71 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.1, 8.3 Hz, 1H), 4.79 - 4.77 (m, 2H), 4.28 (s, 2H), 4.16 - 4.09 (m, 2H), 3.77 - 3.77 (m, 2H), 3.57 - 3.36 (m, 10H), 3.30 (d, J=5.3 Hz, 2H), 3.19 - 3.10 (m, 3H), 3.01 - 2.94 (m, 2H), 1.96 - 1.92 (m, 1H), 1.79 (dd, J=12.9, 12.9 Hz, 2H), 1.70 - 1.47 (m, 7H), 1.44 - 1.31 (m, 2H).	TFA
96	2.53	11	(MeOD); δ 8.38 (dd, J=5.6, 8.4 Hz, 1H), 7.54 - 7.48 (m, 6H), 7.43 (s, 4H), 7.40 - 7.27 (m, 6H), 7.06 (dd, J=7.7, 7.7 Hz, 2H), 6.96 (s, 1H), 6.92 (dd, J=2.3, 8.0 Hz, 1H), 6.71 (d, J=9.9 Hz, 1H), 5.42 (dd, J=5.0, 8.4 Hz, 1H), 4.62 (s, 2H), 4.28 (s, 2H), 4.13 (d, J=6.1 Hz, 2H), 4.09 - 4.01 (m, 2H), 3.75 (s, 3H), 3.51 - 3.35 (m, 5H), 3.30 (d, J=5.0 Hz, 2H), 3.16 - 3.10 (m, 3H), 3.02 - 2.94 (m, 2H), 2.01 - 1.96 (m, 1H), 1.80 - 1.80 (m, 7H), 1.47 - 1.42 (m, 2H).	TFA
97	2.49	11	(MeOD); δ 8.38 (d, J=9.9 Hz, 1H), 7.53 - 7.49 (m, 5H), 7.48 (s, 1H), 7.42 - 7.29 (m, 10H), 7.05 (d, J=8.2 Hz, 3H), 6.96 (dd, J=2.0, 8.2 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.3, 8.3 Hz, 1H), 4.65 - 4.50 (m, 3H), 4.28 (s, 2H), 4.12 (d, J=6.3 Hz, 2H), 3.75 - 3.62 (m, 1H), 3.51 - 3.38 (m, 5H), 3.18 - 3.08 (m, 5H), 3.02 - 2.94 (m, 4H), 2.01 - 1.94 (m, 1H), 1.85 - 1.77 (m, 4H), 1.60 (s, 1H), 1.48 - 1.29 (m, 2H), 1.16 - 1.16 (m, 2H).	TFA
98	2.47	11	(DMSO-d ₆) δ 8.17 (d, J=9.9 Hz, 1H), 7.32 - 7.22 (m, 16H), 7.05 (d, J=8.3 Hz, 1H), 6.94 - 6.87 (m, 3H), 6.84 - 6.84 (m, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.01 (dd, J=4.5, 7.8 Hz, 1H), 4.58 - 4.47 (m, 2H), 4.03 - 3.97 (m, 2H), 3.85 - 3.83 (m, 2H), 3.60 - 3.52 (m, 2H), 3.30 (m, 3H), 2.84 - 2.67 (m, 8H), 1.87 - 1.72 (m, 5H), 1.53 - 1.40 (m, 6H), 1.16 - 1.05 (m, 3H)	Free Base

(continued)

99	2.50	11	(MeOD); δ 8.52 (s, 2H), 8.37 (d, J=9.9 Hz, 1H), 7.39 (s, 9H), 7.37 - 7.22 (m, 6H), 7.04 - 6.93 (m, 5H), 6.69 (d, J=9.9 Hz, 1H), 5.38 (dd, J=4.3, 9.0 Hz, 1H), 4.89 - 4.74 (m, 2H), 4.05 (d, J=5.9 Hz, 2H), 3.86 - 3.83 (m, 2H), 3.61 - 3.04 (m, 19H), 2.39 (s, 1H), 1.60 - 1.56 (m, 6H), 1.43 - 1.42 (m, 1H), 1.35 - 1.29 (m, 4H).	Formate
100	2.44	11	(DMSO-d ₆ , 100 °C) δ 8.18 (1H, d, J=9.9 Hz), 7.47 (5H, s), 7.39 - 7.22 (10H, m), 7.16 (1H, d, J=8.2 Hz), 7.04 - 6.97 (3H, m), 6.89 (1H, dd, J=2.1, 7.8 Hz), 6.58 (1H, d, J=9.8 Hz), 5.36 (1H, dd, J=4.9, 8.3 Hz), 4.74 (2H, s), 4.21 (3H, m), 4.06 (3H, m), 3.67 (5H, dd, J=5.0, 5.0 Hz), 3.48 (2H, dd, J=4.8, 4.8 Hz), 3.41 - 3.19 (8H, m), 3.10 - 3.04 (2H, m), 2.98 - 2.77 (2H, m), 2.56 - 2.54 (1H, m), 1.89 (1H, s), 1.78 - 1.74 (3H, m), 1.57 - 1.40 (3H, m);	TFA
101	2.42	11	(MeOD); δ 8.47 (s, 2H), 8.37 (d, J=9.9 Hz, 1H), 7.87 (d, J=7.9 Hz, 2H), 7.45 - 7.28 (m, 14H), 7.07 - 7.03 (m, 2H), 7.01 - 6.91 (m, 2H), 6.70 (d, J=9.8 Hz, 1H), 5.41 (dd, J=5.5, 8.0 Hz, 1H), 4.84 - 4.76 (m, 1H), 4.69 - 4.63 (m, 1H), 4.62 (d, J=2.1 Hz, 2H), 4.41 - 4.27 (m, 2H), 4.14 - 4.07 (m, 6H), 3.29 - 3.23 (m, 4H), 3.16 - 3.08 (m, 3H), 2.78 - 2.67 (m, 2H), 1.94 - 1.85 (m, 1H), 1.76 - 1.72 (m, 2H), 1.45 - 1.34 (m, 2H)	Formate
102	2.43	11	(MeOD); δ 8.47 (s, 2H), 8.37 (d, J=9.9 Hz, 1H), 7.87 (d, J=7.9 Hz, 2H), 7.45 - 7.28 (m, 14H), 7.07 - 7.03 (m, 2H), 7.01 - 6.91 (m, 2H), 6.70 (d, J=9.8 Hz, 1H), 5.41 (dd, J=5.5, 8.0 Hz, 1H), 4.84 - 4.76 (m, 1H), 4.69 - 4.63 (m, 1H), 4.62 (d, J=2.1 Hz, 2H), 4.41 - 4.27 (m, 2H), 4.14 - 4.07 (m, 6H), 3.29 - 3.23 (m, 4H), 3.16 - 3.08 (m, 3H), 2.78 - 2.67 (m, 2H), 1.94 - 1.85 (m, 1H), 1.76 - 1.72 (m, 2H), 1.45 - 1.34 (m, 2H)	Formate
103	2.41	11	(MeOD); δ 8.41 (d, J=9.9 Hz, 1H), 7.78 (d, J=8.3 Hz, 2H), 7.50 (s, 5H), 7.42 - 7.24 (m, 9H), 7.06 (d, J=8.3 Hz, 3H), 6.97 - 6.93 (m, 1H), 6.72 (d, J=9.8 Hz, 1H), 5.45 (dd, J=6.8, 6.8 Hz, 1H), 4.50 (s, 2H), 4.28 (s, 2H), 4.10 (d, J=6.4 Hz, 2H), 3.54 - 3.37 (m, 9H), 3.21 - 3.10 (m, 3H), 3.02 - 2.93 (m, 2H), 2.01 - 1.93 (m, 1H), 1.84 - 1.74 (m, 2H), 1.49 - 1.37 (m, 2H).	TFA

(continued)

104	2.39	11	(DMSO, 100 °C); δ 8.21 (d, J=9.9 Hz, 1H), 8.17 (s, 1H), 7.92 - 7.92 (m, 1H), 7.39 - 7.24 (m, 15H), 7.08 (d, J=8.2 Hz, 1H), 7.04 - 7.00 (m, 2H), 6.95 (d, J=8.2 Hz, 1H), 6.90 (dd, J=2.4, 8.0 Hz, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.04 (dd, J=5.0, 7.5 Hz, 1H), 4.43 (s, 2H), 4.37 (br s, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.71 (dd, J=6.7, 11.4 Hz, 1H), 3.60 - 3.49 (m, 2H), 3.44 (s, 2H), 3.36 (dd, J=5.3, 11.4 Hz, 2H), 2.91 - 2.72 (m, 6H), 2.17 - 2.07 (m, 1H), 1.97 - 1.86 (m, 3H), 1.62 - 1.48 (m, 3H), 1.28 - 1.13 (m, 2H).	mono Formate
105	2.44	11	(DMSO-d ₆ , 100 °C); δ 8.21 (d, J=9.9 Hz, 1H), 8.16 (s, 2H), 7.57 (d, J=8.4 Hz, 1H), 7.40 - 7.21 (m, 15H), 7.10 - 7.04 (m, 2H), 7.01 (d, J=8.0 Hz, 1H), 6.95 (d, J=8.2 Hz, 1H), 6.91 (dd, J=2.3, 8.2 Hz, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.06 - 5.02 (m, 1H), 4.41 (s, 2H), 4.04 (d, J=6.3 Hz, 2H), 3.93 - 3.69 (m, 4H), 3.43 (s, 2H), 3.13 - 3.04 (m, 4H), 2.91 - 2.68 (m, 7H), 1.98 - 1.90 (m, 2H), 1.69 - 1.48 (m, 4H), 1.24 - 1.13 (m, 2H).	Formate
106	2.47	11	(DMSO-d ₆ , 100 °C); δ 8.21 (d, J=9.9 Hz, 1H), 8.17 (s, 1H), 7.57 (d, J=7.4 Hz, 1H), 7.41 - 7.37 (m, 2H), 7.32 - 7.22 (m, 12H), 7.10 - 7.00 (m, 3H), 6.95 (d, J=8.0 Hz, 1H), 6.90 (dd, J=2.3, 8.2 Hz, 1H), 6.50 (d, J=9.9 Hz, 1H), 5.04 (dd, J=4.9, 7.4 Hz, 1H), 4.41 (d, J=2.5 Hz, 2H), 4.05 (d, J=6.3 Hz, 2H), 3.92 (d, J=12.5 Hz, 1H), 3.85 - 3.77 (m, 1H), 3.72 (d, J=11.8 Hz, 1H), 3.44 (s, 2H), 3.13 - 3.02 (m, 2H), 2.91 - 2.72 (m, 8H), 2.57 - 2.47 (m, 1H), 1.97 - 1.85 (m, 3H), 1.72 - 1.48 (m, 6H), 1.28 - 1.13 (m, 2H).	mono Formate
107	2.40	11	(DMSO-d ₆ , 100 °C); δ 8.20 (d, J=9.9 Hz, 1H), 7.97 (d, J=6.1 Hz, 1H), 7.51 - 7.47 (m, 6H), 7.41 - 7.24 (m, 9H), 7.17 (d, J=8.3 Hz, 1H), 7.04 - 7.00 (m, 3H), 6.90 (dd, J=2.1, 8.2 Hz, 1H), 6.58 (d, J=9.8 Hz, 1H), 5.38 (dd, J=4.8, 8.3 Hz, 1H), 4.43 (s, 2H), 4.33 (dd, J=6.3, 12.4 Hz, 1H), 4.23 (s, 2H), 4.10 (d, J=5.3 Hz, 2H), 3.72 (dd, J=6.8, 11.4 Hz, 1H), 3.61 - 3.48 (m, 4H), 3.14 - 3.05 (m, 6H), 2.99 - 2.95 (m, 2H), 2.57 - 2.55 (m, 1H), 2.17 - 2.05 (m, 1H), 1.94 - 1.85 (m, 2H), 1.79 - 1.75 (m, 2H), 1.45 - 1.44 (m, 2H).	TFA

(continued)


108	2.44	11	(DMSO-d ₆ , 100 °C); δ 8.21 - 8.19 (m, 2H), 7.85 - 7.80 (m, 1H), 7.75 (d, J=8.2 Hz, 2H), 7.41 - 7.38 (m, 2H), 7.33 - 7.22 (m, 11H), 7.08 (d, J=8.2 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.90 - 6.87 (m, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.03 (dd, J=5.0, 7.5 Hz, 1H), 4.73 (s, 2H), 4.06 - 4.03 (m, 3H), 3.88 - 3.87 (m, 2H), 3.74 - 3.68 (m, 1H), 3.44 (s, 2H), 3.13 - 2.97 (m, 1H), 2.93 - 2.71 (m, 8H), 2.57 - 2.47 (m, 1H), 2.05 - 1.89 (m, 3H), 1.82 - 1.76 (m, 1H), 1.72 - 1.46 (m, 4H), 1.24 - 1.13 (m, 2H).	mono-formate
109	2.41	11	(DMSO, 100 °C); δ 8.21 - 8.17 (m, 3H), 8.12 (s, 1H), 7.76 (d, J=8.2 Hz, 2H), 7.39 (dd, J=1.4, 8.3 Hz, 2H), 7.35 - 7.22 (m, 11H), 7.08 (d, J=8.2 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.89 (dd, J=1.9, 8.2 Hz, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.03 (dd, J=5.0, 7.5 Hz, 1H), 4.64 (s, 2H), 4.53 (s, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.88 - 3.58 (m, 2H), 3.43 (s, 2H), 2.91 - 2.71 (m, 9H), 2.56 - 2.45 (m, 1H), 2.17 (br s, 1H), 2.05 (br s, 1H), 1.98 - 1.90 (m, 2H), 1.63 - 1.49 (m, 3H), 1.28 - 1.13 (m, 2H).	formate
110	2.43	11	(DMSO-d ₆ , 100 °C); δ 8.20 (d, J=9.9 Hz, 1H), 8.16 - 8.11 (m, 2H), 7.77 (d, J=8.3 Hz, 2H), 7.39 (dd, J=1.4, 8.3 Hz, 2H), 7.35 - 7.21 (m, 11H), 7.09 (d, J=8.3 Hz, 1H), 7.00 - 6.95 (m, 3H), 6.89 (dd, J=2.0, 8.2 Hz, 1H), 6.50 (d, J=9.9 Hz, 1H), 6.10 (s, 1H), 5.09 (dd, J=4.9, 7.5 Hz, 1H), 4.64 (s, 2H), 4.54 - 4.45 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.72 - 3.59 (m, 2H), 3.47 - 3.40 (m, 3H), 3.00 - 2.73 (m, 8H), 2.57 - 2.47 (m, 1H), 2.17 (br s, 1H), 2.04 - 1.91 (m, 3H), 1.63 - 1.55 (m, 1H), 1.52 (d, J=13.2 Hz, 2H), 1.25 - 1.14 (m, 2H).	mono-formate
111	2.41	11	(DMSO-d ₆); δ 8.21 (d, J=9.8 Hz, 1H), 8.16 (s, 2H), 7.40 - 7.20 (m, 15H), 7.08 (d, J=8.4 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.85 (dd, J=1.9, 8.0 Hz, 1H), 6.50 (d, J=9.0 Hz, 1H), 5.04 (dd, J=5.0, 7.4 Hz, 1H), 4.63 (s, 2H), 4.43 - 4.37 (m, 1H), 4.04 (d, J=6.4 Hz, 2H), 3.79 - 3.73 (m, 1H), 3.72 - 3.49 (m, 4H), 3.46 - 3.39 (m, 3H), 2.95 - 2.73 (m, 8H), 2.57 - 2.47 (m, 1H), 2.06 - 2.00 (m, 1H), 1.97 - 1.89 (m, 2H), 1.81 - 1.76 (m, 1H), 1.62 - 1.49 (m, 3H), 1.28 - 1.13 (m, 2H).	formate

(continued)


112	2.42	11	(DMSO-d ₆ , 100 °C); δ 8.21 (d, J=9.9 Hz, 1H), 8.16 (s, 1H), 7.41 - 7.20 (m, 15H), 7.08 (d, J=8.4 Hz, 1H), 7.00 - 6.94 (m, 3H), 6.87 - 6.84 (m, 1H), 6.50 (d, J=9.9 Hz, 1H), 5.04 (dd, J=5.0, 7.4 Hz, 1H), 4.63 (s, 2H), 4.44 - 4.36 (m, 1H), 4.05 (d, J=6.3 Hz, 2H), 3.81 - 3.75 (m, 1H), 3.72 - 3.49 (m, 4H), 3.46 - 3.40 (m, 3H), 2.94 - 2.72 (m, 8H), 2.58 - 2.50 (m, 1H), 2.08 - 2.01 (m, 1H), 1.97 - 1.89 (m, 2H), 1.80 - 1.77 (m, 1H), 1.62 - 1.49 (m, 3H), 1.28 - 1.13 (m, 2H).	mono-formate
113	2.45	11	(DMSO-d ₆ , 100 °C); δ 8.22 - 8.18 (m, 2H), 7.84 (d, J=5.4 Hz, 1H), 7.74 (d, J=8.2 Hz, 2H), 7.39 (dd, J=1.3, 8.3 Hz, 2H), 7.33 - 7.22 (m, 11H), 7.08 (d, J=8.2 Hz, 1H), 7.01 - 6.94 (m, 3H), 6.91 - 6.87 (m, 1H), 6.49 (d, J=9.9 Hz, 1H), 5.03 (dd, J=5.0, 7.5 Hz, 1H), 4.73 (s, 2H), 4.05 (d, J=6.4 Hz, 3H), 3.91 - 3.84 (m, 2H), 3.43 (s, 2H), 3.10 - 3.00 (m, 2H), 2.91 - 2.68 (m, 7H), 2.57 - 2.46 (m, 1H), 2.00 - 1.89 (m, 3H), 1.83 - 1.49 (m, 6H), 1.24 - 1.13 (m, 2H).	mono-formate

Example 21

(1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate (Compound 114)




Step 1; (1-Benzylpiperidin-4-yl)methyl (S)-2-(3-((4-(((1R,3S)-3-((tert-butoxycarbonyl)amino)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate



A stirred solution of (S)-4-((3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)methyl)benzoic acid hydrochloride (0.20 g, 0.33 mmol) in DMF (1.9 mL) was added with DIPEA (0.132 mL, 0.76 mmol) and HATU (0.173 g, 0.46 mmol) and the mixture stirred at room temperature for 30 minutes. This mixture was added with a solution of *tert*-butyl *trans*-(3-aminocyclobutyl)carbamate hydrochloride (0.068 g, 0.38 mmol) and DIPEA (0.13 mL, 0.76 mmol) in DMF (1.9 mL). The reaction mixture diluted with ethyl acetate and washed with 2M aqueous sodium hydroxide and brine. The organic phase was dried over magnesium sulfate, filtered and the filtrate evaporated at reduced pressure to afford the title compound (0.247 g, >100%).













¹H NMR (400 MHz, DMSO-d₆); δ 8.67 (d, J=6.9 Hz, 1H), 7.86 (d, J=8.3 Hz, 2H), 7.47 (d, J=8.3 Hz, 2H), 7.33 - 7.24 (m, 12H), 6.96 (d, J=5.9 Hz, 2H), 6.92 (dd, J=8.1, 8.1 Hz, 1H), 6.60 (s, 1H), 5.12 (s, 2H), 4.41 - 4.39 (m, 1H), 4.15 - 4.06 (m, 1H), 4.01 - 3.97 (m, 2H), 3.39 (s, 2H), 2.34 - 2.22 (m, 5H), 1.82 (dd, J=11.0, 11.0 Hz, 2H), 1.46 (d, J=12.9 Hz, 3H), 1.39 (s, 9H), 1.37 (s, 1H), 1.17 - 1.05 (m, 2H).

Step 2; (1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate (Compound 114)







A stirred solution of (1-benzylpiperidin-4-yl)methyl (S)-2-(3-((4-(((1*r*,3*S*)-3-((tert-butoxycarbonyl)amino)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate (0.247 g, 0.34 mmol) was added with a solution of HCl in dioxan (4M, 15 mL) and the mixture stirred at room temperature for 1 hour. The solvent evaporated under reduced pressure. The residue was dissolved in DMF (3 ml) and added to a pre-stirred (15

minutes) mixture of (R)-4-(2-((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoic acid (0.196 g, 0.35

N	Appropriate amine	Structure
115		
116		
117		
118		
119		
120		

(continued)

121		
122		

mmol), DIPEA (0.214 mL, 1.23 mmol) and HATU (0.16 g, 0.42 mmol) in DMF (2 mL). The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture dissolved in ethyl acetate and washed with saturated aqueous sodium carbonate and brine. The organic phase was dried with magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was dissolved in dioxan (1 mL) and a solution of HCl – dioxan (2 mL) added. The reaction mixture was stirred at room temperature for 1 hour. The solvent was evaporated at reduced pressure and the residue purified by reverse preparative HPLC.

¹H NMR (400 MHz, DMSO-d₆); δ 8.81 (d, J=7.1 Hz, 1H), 8.74 (d, J=7.1 Hz, 1H), 8.28 (s, 1H), 8.22 (d, J=9.5 Hz, 1H), 7.94 (d, J=8.3 Hz, 2H), 7.87 (d, J=8.3 Hz, 2H), 7.54 (d, J=8.3 Hz, 2H), 7.37 - 7.28 (m, 13H), 7.13 (d, J=8.1 Hz, 1H), 7.02 - 6.95 (m, 4H), 6.57 (d, J=9.9 Hz, 1H), 5.18 (s, 2H), 5.14 (dd, J=5.2, 7.5 Hz, 1H), 4.66 - 4.57 (m, 2H), 4.04 (d, J=6.3 Hz, 2H), 3.44 (s, 2H), 2.97 - 2.83 (m, 6H), 2.77 (d, J=11.1 Hz, 2H), 2.49 (t, J=7.1 Hz, 4H), 1.88 (dd, J=9.9, 11.6 Hz, 2H), 1.57 - 1.47 (m, 3H), 1.21 - 1.12 (m, 2H).

The following compounds were prepared in a similar fashion using the appropriate amine;

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
114	2.57	11	(DMSO-d ₆); δ 8.81 (d, J=7.1 Hz, 1H), 8.74 (d, J=7.1 Hz, 1H), 8.28 (s, 1H), 8.22 (d, J=9.5 Hz, 1H), 7.94 (d, J=8.3 Hz, 2H), 7.87 (d, J=8.3 Hz, 2H), 7.54 (d, J=8.3 Hz, 2H), 7.37 - 7.28 (m, 13H), 7.13 (d, J=8.1 Hz, 1H), 7.02 - 6.95 (m, 4H), 6.57 (d, J=9.9 Hz, 1H), 5.18 (s, 2H), 5.14 (dd, J=5.2, 7.5 Hz, 1H), 4.66 - 4.57 (m, 2H), 4.04 (d, J=6.3 Hz, 2H), 3.44 (s, 2H), 2.97 - 2.83 (m, 6H), 2.77 (d, J=11.1 Hz, 2H), 2.49 (t, J=7.1 Hz, 4H), 1.88 (dd, J=9.9, 11.6 Hz, 2H), 1.57 - 1.47 (m, 3H), 1.21 - 1.12 (m, 2H).	TFA
115	2.58	11	(DMSO-d ₆); δ 8.23 (s, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.46 (d, J=8.2 Hz, 2H), 7.37 (d, J=8.2 Hz, 2H), 7.33 - 7.25 (m, 15H), 7.08 (d, J=8.3 Hz, 1H), 6.98 - 6.91 (m, 4H), 6.52 (d, J=9.8 Hz, 1H), 5.12 (d, J=5.9 Hz, 1H), 5.09 (s, 2H), 4.00 (d, J=6.3 Hz, 2H), 3.60 (s, 2H), 3.39 (s, 2H), 3.29 (s, 6H), 2.97 - 2.87 (m, 2H), 2.83 - 2.77 (m, 4H), 2.72 (d, J=11.3 Hz, 2H), 1.87 - 1.79 (m, 2H), 1.53 - 1.45 (m, 10H), 1.17 - 1.08 (m, 2H).	Formate
116	2.57	11	(DMSO-d ₆ , 90 °C); δ 8.18 (d, J=9.9 Hz, 1H), 7.65 - 7.59 (m, 3H), 7.48 (d, J=7.2 Hz, 5H), 7.39 - 7.24 (m, 8H), 7.16 (d, J=8.3 Hz, 1H), 7.04 - 6.95 (m, 3H), 6.58 (d, J=9.9 Hz, 1H), 5.36 (dd, J=4.5, 8.5 Hz, 1H), 5.12 (s, 2H), 4.35 - 4.33 (m, 8H), 4.22 (s, 2H), 4.06 (s, 2H), 3.34 - 3.29 (m, 4H), 3.21 - 3.17 (m, 4H), 3.10 - 3.04 (m, 5H), 1.97 - 1.71 (m, 4H), 1.42 - 1.42 (m, 2H).	TFA
117	2.57	11	(DMSO-d ₆); δ 8.23 (s, 1H), 8.18 (d, J=9.9 Hz, 1H), 7.65 (d, J=8.3 Hz, 2H), 7.48 (d, J=8.3 Hz, 2H), 7.32 - 7.24 (m, 15H), 7.08 (d, J=8.3 Hz, 1H), 6.98 - 6.90 (m, 4H), 6.61 (s, 1H), 6.52 (d, J=9.9 Hz, 1H), 5.12 - 5.08 (m, 3H), 4.06 (s, 2H), 3.99 (d, J=6.3 Hz, 2H), 3.81 (s, 2H), 3.48 (m, 1H), 3.39 (s, 2H), 3.28 (m, 3H), 2.94 - 2.85 (m, 2H), 2.83 - 2.77 (m, 4H), 2.72 (d, J=11.4 Hz, 2H), 1.87 - 1.72 (m, 6H), 1.53 - 1.45 (m, 3H), 1.16 - 1.07 (m, 2H).	mono-Formate







(continued)

118	2.55	11	(DMSO-d ₆); δ 10.55 (s, 1H), 10.45 (s, 1H), 9.40 (s, 1H), 8.75 (s, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.52 - 7.41 (m, 11H), 7.36 - 7.25 (m, 8H), 7.17 (d, J=8.3 Hz, 1H), 7.01 - 6.95 (m, 3H), 6.92 (d, J=7.9 Hz, 1H), 6.65 (s, 1H), 6.60 (d, J=9.9 Hz, 1H), 6.15 (s, 1H), 5.34 (d, J=8.0 Hz, 1H), 5.10 (s, 2H), 4.25 (dt, J=6.3, 18.9 Hz, 2H), 4.00 (d, J=6.5 Hz, 2H), 3.63 - 3.26 (m, 10H), 3.18 - 2.99 (m, 5H), 2.94 - 2.85 (m, 2H), 1.84 - 1.83 (m, 1H), 1.75 - 1.66 (m, 3H), 1.35 (q, J=11.9 Hz, 2H).	TFA
119	2.55	11	(DMSO-d ₆) δ 10.55 (s, 1H), 10.45 (s, 1H), 9.40 (s, 1H), 8.75 (s, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.63 (d, J=8.3 Hz, 2H), 7.48 (m, 7H), 7.40 - 7.24 (m, 10H), 7.16 (d, J=8.3 Hz, 1H), 7.01 - 6.96 (m, 3H), 6.92 (d, J=8.2 Hz, 1H), 6.65 (s, 1H), 6.60 (dd, J=2.0, 9.8 Hz, 1H), 6.15 (s, 1H), 5.33 (d, J=8.4 Hz, 1H), 5.09 (s, 2H), 4.26 (dt, J=5.5, 18.7 Hz, 2H), 4.04 - 3.97 (m, 4H), 3.81 (s, 2H), 3.56 - 3.56 (m, 2H), 3.34 - 3.24 (m, 5H), 3.18 - 2.97 (m, 5H), 2.89 (m, 2H), 1.75 - 1.66 (m, 7H), 1.36 (dd, J=12.3, 12.3 Hz, 2H).	TFA
120	2.57	11	(DMSO-d ₆); δ 10.49 (m, 2H), 9.50 (s, 1H), 8.70 (m, 2H), 8.18 (d, J=9.7 Hz, 1H), 7.64 - 7.64 (m, 2H), 7.52 - 7.46 (m, 7H), 7.36 - 7.31 (m, 10H), 7.16 (d, J=8.3 Hz, 1H), 7.01 - 6.96 (m, 3H), 6.92 (d, J=7.5 Hz, 1H), 6.60 (s, 1H), 6.59 (d, J=9.8 Hz, 1H), 6.20 (s, 1H), 5.35 (d, J=9.4 Hz, 1H), 5.11 (s, 2H), 4.30 - 4.19 (m, 2H), 4.02 - 3.96 (m, 2H), 3.84 - 3.51 (m, 6H), 3.34 - 3.00 (m, 10H), 2.90 - 2.85 (m, 2H), 1.83 - 1.82 (m, 3H), 1.74 (m, 2H), 1.48 - 1.32 (m, 4H).	TFA
121	2.56	11	(DMSO-d ₆); δ 10.51 - 10.46 (m, 2H), 9.40 (s, 1H), 8.76 - 8.76 (m, 2H), 8.17 (d, J=9.9 Hz, 1H), 7.51 - 7.47 (m, 7H), 7.35 - 7.31 (m, 12H), 7.17 (d, J=8.3 Hz, 1H), 7.01 - 6.95 (m, 3H), 6.92 (d, J=8.0 Hz, 1H), 6.65 (s, 1H), 6.60 (d, J=9.8 Hz, 1H), 6.21 - 6.20 (m, 1H), 5.34 (d, J=8.0 Hz, 1H), 5.10 (s, 2H), 4.32 - 4.19 (m, 2H), 4.10 - 4.10 (m, 1H), 4.03 - 3.96 (m, 2H), 3.63 - 3.62 (m, 5H), 3.34 (d, J=12.0 Hz, 4H), 3.18 - 2.99 (m, 6H), 2.90 - 2.89 (m, 4H), 1.75 - 1.66 (m, 7H), 1.39 - 1.33 (m, 2H).	TFA







(continued)

122	2.63	11	(DMSO-d ₆) δ 10.52 - 10.46 (m, 2H), 9.40 (s, 1H), 8.75 (s, 2H), 8.16 (d, J=9.9 Hz, 1H), 7.50 - 7.46 (m, 7H), 7.38 - 7.24 (m, 12H), 7.16 (d, J=8.3 Hz, 1H), 7.01 - 6.94 (m, 3H), 6.91 (d, J=7.7 Hz, 1H), 6.65 (s, 1 H), 6.60 (dd, J=1.8, 9.9 Hz, 1H), 6.20 (m, 1H), 5.33 (d, J=8.5 Hz, 1H), 5.09 (s, 2H), 4.25 (dt, J=5.7, 18.7 Hz, 2H), 4.02 - 3.96 (m, 2H), 3.36 - 3.26 (m, 9H), 3.18 (s, 2H), 3.10 - 3.08 (m, 2H), 3.01 (d, J=7.9 Hz, 3H), 2.94 - 2.85 (m, 2H), 1.83 (s, 1H), 1.75 - 1.65 (m, 2H), 1.50 (m, 6H), 1.34 (m, 4H).	TFA
-----	------	----	---	-----


The following compounds were prepared as described in **Example 20** with the appropriate acid replacing **(R)-4-(((*tert*-butoxycarbonyl)(2-(((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid** and the use of the appropriate amine replacing *tert*-butyl *trans*-(3-aminocyclobutyl)carbamate hydrochloride in **Step 1**.

N	Appropriate amine /	appropriate acid	Structure
123			
124			


(continued)

	
	
	
125	126


N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
123	2.55	11	(MeOD); δ 8.49 (s, 2H), 8.33 (d, J=9.9 Hz, 1H), 7.95 (d, J=8.4 Hz, 2H), 7.87 (d, J=8.3 Hz, 2H), 7.63 (d, J=8.3 Hz, 2H), 7.51 (d, J=8.4 Hz, 2H), 7.45 (s, 5H), 7.41 - 7.26 (m, 7H), 7.06 - 6.97 (m, 4H), 6.68 (d, J=9.9 Hz, 1H), 5.42 (dd, J=5.5, 7.7 Hz, 1H), 5.15 (s, 2H), 4.70 - 4.61 (m, 2H), 4.31 (s, 2H), 4.11 - 4.06 (m, 4H), 3.29 - 3.18 (m, 4H), 2.75 - 2.65 (m, 2H), 2.60 (t, J=6.9 Hz, 4H), 1.90 - 1.80 (m, 1H), 1.71 (d, J=13.1 Hz, 2H), 1.45 - 1.33 (m, 2H).	formate
124	2.55	11	(MeOD); δ 8.50 (s, 2H), 8.29 (d, J=9.9 Hz, 1H), 7.87 (d, J=8.3 Hz, 2H), 7.59 - 7.50 (m, 5H), 7.44 (s, 5H), 7.41 - 7.26 (m, 7H), 7.07 - 6.98 (m, 4H), 6.67 (d, J=9.8 Hz, 1H), 5.43 (dd, J=5.6, 7.7 Hz, 1H), 5.15 (s, 2H), 4.69 - 4.62 (m, 2H), 4.36 (s, 2H), 4.10 (d, J=6.2 Hz, 2H), 4.05 (s, 2H), 4.02 (s, 3H), 3.28 - 3.19 (m, 4H), 2.69 - 2.58 (m, 6H), 1.89 - 1.81 (m, 1H), 1.70 (d, J=13.1 Hz, 2H), 1.38 (q, J=12.4 Hz, 2H).	formate
125	2.35	11	(MeOD); δ 8.53 (s, 1H), 8.29 (d, J=10.0 Hz, 1H), 7.88 (d, J=8.3 Hz, 2H), 7.82 (d, J=8.4 Hz, 2H), 7.55 (d, J=8.3 Hz, 2H), 7.47 (d, J=8.3 Hz, 2H), 7.42 - 7.24 (m, 12H), 7.04 - 6.96 (m, 4H), 6.64 (d, J=9.8 Hz, 1H), 5.36 (dd, J=4.6, 8.7 Hz, 1H), 5.12 (s, 2H), 4.19 (s, 2H), 4.06 (d, J=6.5 Hz, 2H), 3.90 (s, 2H), 3.66 (s, 4H), 3.17 - 3.09 (m, 4H), 2.49 (dd, J=10.3, 12.2 Hz, 2H), 1.83 - 1.71 (m, 1H), 1.68 - 1.61 (m, 2H), 1.39 - 1.27 (m, 2H).	mono-formate
126	2.54	11	(MeOD); δ 8.53 (s, 2H), 8.24 (d, J=9.9 Hz, 1H), 7.83 (d, J=8.3 Hz, 2H), 7.50 - 7.44 (m, 5H), 7.41 - 7.23 (m, 12H), 7.04 - 6.95 (m, 4H), 6.63 (d, J=9.8 Hz, 1H), 5.37 (dd, J=6.6, 6.6 Hz, 1H), 5.12 (s, 2H), 4.23 (s, 2H), 4.06 (d, J=6.5 Hz, 2H), 3.92 (s, 3H), 3.86 (s, 2H), 3.67 (s, 4H), 3.15 - 3.07 (m, 4H), 2.44 (dd, J=10.2, 12.5 Hz, 2H), 1.81 - 1.72 (m, 1H), 1.62 (d, J=12.0 Hz, 2H), 1.37 - 1.27 (m, 2H).	formate

Example 22

(1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(3-hydroxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 127)




Step 1; (1-Benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1r,3S)-3-aminocyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate dihydrochloride











The title compound was prepared as described in Example 20 Step 1 and 2.

Step 2;

(1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(3-hydroxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 127)



A solution of 4-formyl-3-hydroxybenzaldehyde (0.103 g, 0.62 mmol) and DIPEA (0.24 mL, 1.4 mmol) in DMF (2 mL) was added with HATU (0.255 g, 0.67 mmol) followed by a solution of (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-aminocyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate dihydrochloride (0.177 g, 0.28 mmol) in DMF (3 mL) and the reaction stirred at room temperature for 18 hours. The reaction mixture was dissolved in ethyl acetate and washed with saturated aqueous sodium hydrogencarbonate (x2) and brine. The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was dissolved in methanol (4 mL) and triethylamine (0.056 mL, 0.4 mmol) and (R)-5-(2-amino-1-hydroxyethyl)-8-hydroxyquinolin-2(1H)-one hydrochloride (0.061 g, 0.19 mmol) added. The reaction mixture was stirred at room temperature for 1.5 hours then sodium triacetoxyborohydride (0.136 g, 0.64 mmol) and acetic acid (0.034 mL) added. The reaction mixture was stirred for a further 2 hours. The reaction mixture was diluted with *iso*-butanol and washed with water. The aqueous phase was extracted with further *iso*-butanol. The combined *iso*-butanol extracts were evaporated under reduced pressure. The residue was purified by reverse phase preparative HPLC to afford the title compound.

135	Example 22	
136	Example 22	
137	Example 22	
138	Example 20	
139	Example 20	
140	Example 20	
141	Example 20	
142	Example 20	

(continued)

[illegible]

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
127	2.45	11	(DMSO-d ₆); δ 10.53 - 10.46 (m, 3H), 9.35 (s, 1H), 8.88 (s, 1H), 8.78 (s, 1H), 8.73 (d, J=6.7 Hz, 1H), 8.49 (d, J=7.4 Hz, 1H), 8.07 (d, J=9.3 Hz, 1H), 7.49 (m, 6H), 7.38 - 7.32 (m, 8H), 7.12 (t, J=7.2 Hz, 1H), 7.01 - 6.92 (m, 3H), 6.88 (dd, J=2.3, 8.2 Hz, 1H), 6.67 (s, 1H), 6.56 (dd, J=2.0, 9.9 Hz, 1H), 6.19 (d, J=3.8 Hz, 1H), 5.37 - 5.33 (m, 1H), 4.43 (m, 4H), 4.30 - 4.24 (m, 4H), 4.02 (d, J=6.3 Hz, 2H), 3.04 (m, 4H), 2.92 - 2.86 (m, 2H), 2.33 (dd, J=6.8, 6.8 Hz, 4H), 1.75 (m, 3H), 1.35 (m, 2H).	TFA

(continued)

128	2.43	11	(DMSO-d ₆); δ 10.50 (m, 2H), 9.34 (s, 1H), 9.17 (s, 2H), 8.77 (d, J=7.3 Hz, 1H), 8.50 (d, J=7.4 Hz, 1H), 8.10 (d, J=9.6 Hz, 1H), 7.64 (dd, J=7.7, 7.7 Hz, 1H), 7.49 (s, 6H), 7.42 (d, J=8.0 Hz, 1H), 7.34 (m, 6H), 7.14 - 7.10 (m, 1H), 7.01 - 6.93 (m, 3H), 6.88 (dd, J=2.4, 8.0 Hz, 1H), 6.66 (s, 1H), 6.60 (dd, J=2.1, 9.9 Hz, 1H), 6.21 (d, J=2.6 Hz, 1H), 5.35 (d, J=9.5 Hz, 1H), 4.42 (s, 4H), 4.31 - 4.21 (m, 4H), 4.04 - 3.97 (m, 2H), 3.35 (d, J=12.3 Hz, 2H), 3.09 - 2.87 (m, 4H), 2.37 - 2.29 (m, 4H), 1.77 - 1.66 (m, 3H), 1.41 - 1.30 (m, 2H).	TFA
129	2.43	11	(DMSO-d ₆); δ 10.50 (m, 2H), 9.42 (s, 1H), 9.08 (s, 2H), 8.76 (d, J=7.2 Hz, 1H), 8.51 (d, J=7.3 Hz, 1H), 8.15 (d, J=8.2 Hz, 1H), 7.76 (d, J=7.3 Hz, 2H), 7.58 (d, J=8.5 Hz, 1H), 7.48 (m, 5H), 7.35 - 7.26 (m, 6H), 7.16 (d, J=8.3 Hz, 1H), 7.02 - 6.97 (m, 2H), 6.94 (d, J=8.5 Hz, 1H), 6.89 (dd, J=2.3, 8.0 Hz, 1H), 6.67 (s, 1H), 6.59 (dd, J=2.1, 9.9 Hz, 1H), 6.22 - 6.20 (m, 1H), 5.41 (d, J=9.8 Hz, 1H), 4.43 (m, 4H), 4.35 - 4.21 (m, 4H), 4.03 (d, J=5.4 Hz, 2H), 3.35 (d, J=12.2 Hz, 2H), 3.18 - 3.13 (m, 2H), 2.95 - 2.86 (m, 2H), 2.43 (s, 3H), 2.34 (dd, J=6.8, 6.8 Hz, 4H), 1.86 - 1.84 (m, 1H), 1.76 - 1.67 (m, 2H), 1.36 (q, J=12.0 Hz, 2H).	TFA
130	2.43	11	(DMSO-d ₆); δ 10.53 - 10.47 (m, 2H), 9.39 (s, 1H), 9.16 (s, 2H), 8.50 (d, J=7.4 Hz, 1H), 8.42 (d, J=7.0 Hz, 1H), 8.08 (d, J=9.9 Hz, 1H), 7.65 (d, J=7.8 Hz, 1H), 7.48 (m, 5H), 7.37 - 7.24 (m, 7H), 7.16 - 7.11 (m, 2H), 7.02 - 6.93 (m, 3H), 6.89 (dd, J=8.0 Hz, 1H), 6.66 (s, 1H), 6.58 (dd, J=2.0, 9.9 Hz, 1H), 6.22 - 6.22 (m, 1H), 5.36 (d, J=9.0 Hz, 1H), 4.49 - 4.44 (m, 1H), 4.43 (s, 2H), 4.37 (dd, J=6.7, 13.4 Hz, 1H), 4.25 (t, J=4.7 Hz, 4H), 4.01 (d, J=5.4 Hz, 2H), 3.89 (s, 3H), 3.34 (d, J=9.4 Hz, 2H), 3.11 - 2.86 (m, 4H), 2.32 (dd, J=6.8, 6.8 Hz, 4H), 1.85 - 1.84 (m, 1H), 1.77 - 1.67 (m, 2H), 1.40 - 1.31 (m, 2H).	TFA
131	2.47	11	(DMSO-d ₆); δ 10.48 (m, 2H), 9.37 (s, 1H), 8.87 - 8.86 (m, 2H), 8.79 (d, J=6.9 Hz, 1H), 8.52 (d, J=7.3 Hz, 1H), 8.08 (d, J=8.4 Hz, 1H), 7.53 (m, 3H), 7.48 (m, 6H), 7.35 - 7.26 (m, 6H), 7.13 (dd, J=8.2, 8.2 Hz, 1H), 7.02 - 6.93 (m, 3H), 6.89 (dd, J=2.4, 8.0 Hz, 1H), 6.58 (dd, J=1.8, 9.9 Hz, 1H), 6.20 (s, 1H), 5.38 - 5.33 (m, 1H), 4.44 (m, 4H), 4.30 - 4.21 (m, 4H), 4.03 (d, J=7.0 Hz, 2H), 3.97 (s, 3H), 3.33 (d, J=11.3 Hz, 2H), 3.07 (m, 2H), 2.95 - 2.87 (m, 2H), 2.36 (dd, J=7.2, 7.2 Hz, 4H), 1.86 - 1.85 (m, 1H), 1.77 - 1.67 (m, 2H), 1.41 - 1.30 (m, 2H).	TFA

(continued)

132	2.46	11	(DMSO-d ₆); δ 10.52 - 10.46 (m, 2H), 9.37 (s, 1H), 9.18 (s, 2H), 8.88 (d, J=6.9 Hz, 1H), 8.51 (d, J=7.4 Hz, 1H), 8.14 (d, J=11.8 Hz, 1H), 7.81 - 7.71 (m, 3H), 7.48 (m, 6H), 7.34 (m, 6H), 7.15 (d, J=8.2 Hz, 1H), 7.02 - 6.93 (m, 3H), 6.89 (dd, J=2.4, 8.0 Hz, 1H), 6.59 (dd, J=1.7, 9.9 Hz, 1H), 6.22 (s, 1H), 5.38 - 5.36 (m, 1H), 4.43 (m, 4H), 4.36 - 4.21 (m, 4H), 4.05 - 3.97 (m, 2H), 3.35 (d, J=11.8 Hz, 2H), 3.18 - 3.09 (m, 2H), 2.95 - 2.86 (m, 2H), 2.35 (dd, J=6.8, 6.8 Hz, 4H), 1.85 - 1.84 (m, 1H), 1.77 - 1.67 (m, 2H), 1.40 - 1.30 (m, 2H).	TFA
133	2.46	11	(MeOD); δ 8.51 (s, 2H), 8.34 (d, J=9.9 Hz, 1H), 7.55 (s, 1H), 7.47 (d, J=7.9 Hz, 1H), 7.42 - 7.40 (m, 7H), 7.39 - 7.29 (m, 4H), 7.24 (d, J=8.4 Hz, 1H), 7.09 - 7.06 (m, 2H), 7.03 - 6.96 (m, 2H), 6.67 (d, J=9.9 Hz, 1H), 5.31 (dd, J=3.9, 8.9 Hz, 1H), 4.53 - 4.45 (m, 4H), 4.13 (d, J=6.4 Hz, 2H), 4.03 (s, 2H), 3.92 (s, 2H), 3.35 - 3.33 (m, 2H), 3.17 (d, J=11.9 Hz, 2H), 3.03 - 2.91 (m, 2H), 2.53 - 2.43 (m, 5H), 1.88 - 1.80 (m, 1H), 1.69 (dd, J=1.8, 13.6 Hz, 2H), 1.43 - 1.30 (m, 2H)	Formate
134	2.44	11	(DMSO-d ₆); δ 10.55 (d, J=8.8 Hz, 2H), 9.45 - 9.41 (m, 1H), 9.33 - 9.30 (m, 2H), 8.99 (d, J=7.1 Hz, 1H), 8.55 (d, J=7.6 Hz, 1H), 8.20 (d, J=9.8 Hz, 1H), 7.58 - 7.50 (m, 6H), 7.43 - 7.37 (m, 4H), 7.37 - 7.30 (m, 2H), 7.21 (d, J=8.3 Hz, 1H), 7.08 - 7.02 (m, 2H), 7.00 (d, J=7.8 Hz, 1H), 6.95 (dd, J=2.4, 8.2 Hz, 1H), 6.71 (s, 1H), 6.66 (dd, J=1.8, 9.9 Hz, 1H), 6.58 (s, 1H), 6.30 - 6.27 (m, 1H), 5.42 (d, J=8.8 Hz, 1H), 4.51 - 4.42 (m, 7H), 4.38 - 4.27 (m, 2H), 4.08 (d, J=6.3 Hz, 2H), 3.24 - 3.11 (m, 3H), 3.01 - 2.92 (m, 2H), 2.46 - 2.33 (m, 4H), 1.93 - 1.72 (m, 3H), 1.47 - 1.35 (m, 2H)	TFA
135	2.64	11	(DMSO-d ₆); δ 10.22 (s, 1H), 8.73 (d, J=7.1 Hz, 1H), 8.41 (d, J=7.3 Hz, 1H), 8.10 (d, J=8.6 Hz, 2H), 7.85 (d, J=1.8 Hz, 1H), 7.73 (dd, J=1.6, 8.0 Hz, 1H), 7.51 (d, J=7.8 Hz, 1H), 7.28 - 7.16 (m, 10H), 7.00 (d, J=8.3 Hz, 1H), 6.94 (dd, J=2.1, 2.1 Hz, 1H), 6.89 - 6.80 (m, 3H), 6.55 (s, 1H), 6.42 (d, J=9.9 Hz, 1H), 5.34 (s, 1H), 5.01 (dd, J=4.7, 7.7 Hz, 1H), 4.35 (s, 4H), 3.94 (d, J=6.3 Hz, 2H), 3.80 (s, 2H), 3.33 (s, 2H), 2.72 - 2.59 (m, 3H), 2.27 (dd, J=6.8, 6.8 Hz, 4H), 1.81 - 1.73 (m, 2H), 1.50 - 1.36 (m, 3H), 1.11 - 1.00 (m, 2H).	mono-formate

(continued)


136	2.51	11	(DMSO-d ₆); δ 10.51 (d, J=12.8 Hz, 2H), 9.35 (s, 1H), 9.15 (s, 2H), 8.87 (d, J=6.8 Hz, 1H), 8.49 (d, J=7.6 Hz, 1H), 8.11 (d, J=11.2 Hz, 1H), 7.88 (d, J=1.3 Hz, 1H), 7.58 (dd, J=1.1, 8.0 Hz, 1H), 7.52 - 7.47 (m, 5H), 7.38 - 7.24 (m, 6H), 7.16 - 7.11 (m, 2H), 7.03 - 6.98 (m, 3H), 6.95 (d, J=8.1 Hz, 1H), 6.90 (dd, J=2.4, 8.5 Hz, 1H), 6.62 (dd, J=5.7, 5.7 Hz, 1H), 6.23 (s, 1H), 5.36 (d, J=7.6 Hz, 1H), 4.47 - 4.36 (m, 4H), 4.32 - 4.22 (m, 3H), 4.03 (d, J=7.1 Hz, 2H), 3.36 (d, J=12.8 Hz, 2H), 3.13 - 2.88 (m, 5H), 2.38 - 2.29 (m, 4H), 1.90 - 1.82 (m, 1H), 1.78 - 1.66 (m, 2H), 1.41 - 1.33 (m, 2H).	TFA
137	2.45	11	(DMSO-d ₆); δ 10.52 - 10.50 (m, 2H), 9.39 (s, 1H), 8.93 (s, 2H), 8.87 (d, J=6.8 Hz, 1H), 8.50 (d, J=7.4 Hz, 1H), 8.12 (d, J=9.9 Hz, 1H), 7.61 (s, 1H), 7.49 - 7.48 (m, 5H), 7.38 - 7.25 (m, 6H), 7.15 (dd, J=4.1, 4.1 Hz, 2H), 7.02 - 6.94 (m, 3H), 6.89 (dd, J=2.4, 7.9 Hz, 1H), 6.69 (s, 1H), 6.61 (d, J=9.9 Hz, 1H), 6.22 (s, 1H), 5.37 (dd, J=3.5, 8.9 Hz, 1H), 4.46 - 4.36 (m, 4H), 4.32 - 4.22 (m, 4H), 4.03 (d, J=6.4 Hz, 2H), 3.88 (s, 3H), 3.34 (d, J=11.5 Hz, 2H), 3.07 - 3.07 (m, 2H), 2.96 - 2.87 (m, 2H), 2.39 - 2.30 (m, 4H), 1.77 - 1.67 (m, 3H), 1.41 - 1.31 (m, 2H).	TFA
138	2.48	11	(DMSO-d ₆); δ 10.26 - 10.21 (s, 1H), 8.87 (d, J=7.1 Hz, 1H), 8.43 (d, J=7.3 Hz, 1H), 8.12 - 8.08 (m, 3H), 8.05 (d, J=8.1 Hz, 1H), 7.79 (d, J=8.1 Hz, 1H), 7.29 - 7.17 (m, 11H), 7.01 (d, J=8.3 Hz, 1H), 6.95 (dd, J=2.0, 2.0 Hz, 1H), 6.90 - 6.81 (m, 3H), 6.56 (s, 1H), 6.42 (d, J=9.9 Hz, 1H), 5.40 (s, 1H), 5.02 (dd, J=4.7, 7.7 Hz, 1H), 4.46 - 4.37 (m, 2H), 4.36 (s, 2H), 3.97 - 3.87 (m, 4H), 3.45 (s, 2H), 2.72 - 2.60 (m, 4H), 2.29 (t, J=6.8, 6.8 Hz, 4H), 1.83 - 1.73 (t, 2H), 1.49 - 1.37 (m, 3H), 1.12 - 1.00 (m, 2H).	formate
139	2.43	11	(DMSO-d ₆); δ 10.51 (s, 2H), 9.40 (s, 1H), 9.04 - 9.04 (m, 2H), 8.83 (d, J=6.9 Hz, 1H), 8.51 (d, J=7.4 Hz, 1H), 8.13 (d, J=9.8 Hz, 1H), 7.49 - 7.48 (m, 5H), 7.42 - 7.24 (m, 8H), 7.15 (d, J=8.3 Hz, 1H), 7.02 - 6.93 (m, 3H), 6.89 (dd, J=2.4, 8.1 Hz, 1H), 6.67 (s, 1H), 6.60 (d, J=9.9 Hz, 1H), 6.23 (s, 1H), 5.38 (dd, J=3.0, 9.3 Hz, 1H), 4.48 - 4.38 (m, 4H), 4.36 - 4.26 (m, 4H), 4.04 (d, J=5.2 Hz, 2H), 3.96 (d, J=2.0 Hz, 3H), 3.34 (d, J=11.5 Hz, 2H), 3.15 - 3.05 (m, 2H), 2.96 - 2.87 (m, 2H), 2.38 - 2.29 (m, 4H), 1.84 - 1.66 (m, 3H), 1.41 - 1.31 (m, 2H).	TFA

(continued)


140	2.46	11	(MeOD); δ 8.54 (s, 1H), 8.29 (d, J=9.9 Hz, 1H), 7.49 (d, J=6.7 Hz, 2H), 7.46 - 7.31 (m, 12H), 7.24 (d, J=8.3 Hz, 1H), 7.11 - 7.07 (m, 2H), 7.04 - 6.98 (m, 2H), 6.64 (d, J=9.9 Hz, 1H), 5.36 - 5.31 (m, 1H), 4.59 - 4.48 (m, 4H), 4.23 - 4.17 (m, 4H), 4.12 (d, J=6.9 Hz, 2H), 3.79 (s, 2H), 3.13 - 3.04 (m, 4H), 2.53 - 2.46 (m, 4H), 2.36 (dd, J=11.6, 11.6 Hz, 2H), 1.82 - 1.74 (m, 1H), 1.67 - 1.64 (m, 2H), 1.46 (dd, J=7.0, 7.0 Hz, 3H), 1.39 - 1.29 (m, 2H).	mono-formate
141	2.49	11	(MeOD); δ 8.50 (s, 1H), 8.34 (d, J=9.9 Hz, 1H), 7.87 (dd, J=1.6, 7.9 Hz, 1H), 7.83 - 7.82 (m, 1H), 7.67 (d, J=8.0 Hz, 1H), 7.45 - 7.40 (m, 7H), 7.39 - 7.30 (m, 4H), 7.24 (d, J=8.2 Hz, 1H), 7.10 - 7.08 (m, 2H), 7.03 - 6.98 (m, 2H), 6.64 (d, J=9.8 Hz, 1H), 5.28 (dd, J=4.3, 8.4 Hz, 1H), 4.60 - 4.45 (m, 4H), 4.14 (d, J=6.3 Hz, 2H), 4.04 (s, 2H), 3.97 (s, 2H), 3.21 (d, J=12.6 Hz, 2H), 3.00 - 2.88 (m, 2H), 2.64 - 2.57 (m, 2H), 2.53 - 2.44 (m, 4H), 1.91 - 1.83 (m, 1H), 1.74 - 1.70 (m, 2H), 1.44 - 1.33 (m, 2H).	mono-formate
142	2.47	11	(MeOD); δ 8.49 (s, 2H), 8.32 (d, J=9.9 Hz, 1H), 7.45 - 7.40 (m, 7H), 7.39 - 7.25 (m, 7H), 7.10 - 7.07 (m, 2H), 7.04 (d, J=8.2 Hz, 1H), 7.01 - 6.97 (m, 1H), 6.68 (d, J=9.9 Hz, 1H), 5.40 - 5.35 (m, 1H), 4.60 - 4.45 (m, 4H), 4.18 (s, 2H), 4.15 (d, J=6.2 Hz, 2H), 4.02 (s, 2H), 3.93 (s, 3H), 3.24 (d, J=12.0 Hz, 2H), 3.12 (d, J=6.1 Hz, 2H), 2.65 (dd, J=11.5, 13.1 Hz, 2H), 2.48 (dd, J=6.9, 6.9 Hz, 4H), 1.89 - 1.84 (m, 1H), 1.73 (d, J=12.0 Hz, 2H), 1.47 - 1.33 (m, 2H).	formate
143	2.52	11	(MeOD); δ 8.50 (s, 2H), 8.29 (d, J=9.9 Hz, 1H), 7.54 (s, 1H), 7.51 - 7.47 (m, 2H), 7.44 - 7.40 (m, 7H), 7.39 - 7.30 (m, 4H), 7.26 (d, J=8.3 Hz, 1H), 7.11 - 7.07 (m, 2H), 7.04 (d, J=7.9 Hz, 1H), 7.01 - 6.97 (m, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.37 (dd, J=4.6, 8.7 Hz, 1H), 4.87 - 4.82 (m, 1H), 4.61 - 4.46 (m, 4H), 4.26 (d, J=2.8 Hz, 2H), 4.14 (d, J=6.4 Hz, 2H), 3.96 (s, 2H), 3.24 - 3.17 (m, 4H), 2.61 - 2.46 (m, 6H), 1.90 - 1.83 (m, 1H), 1.76 - 1.70 (m, 2H), 1.45 - 1.32 (m, 8H).	formate

(continued)


144	2.46	11	(MeOD); δ 8.71 (d, J=1.5 Hz, 1H), 8.45 (s, 2H), 8.35 (d, J=9.9 Hz, 1H), 8.10 (d, J=8.0 Hz, 1H), 8.03 (dd, J=2.0, 8.2 Hz, 1H), 7.48 - 7.41 (m, 7H), 7.39 - 7.31 (m, 4H), 7.27 (d, J=7.6 Hz, 1H), 7.11 - 7.07 (m, 2H), 7.05 - 6.98 (m, 2H), 6.66 (d, J=9.9 Hz, 1H), 5.35 (dd, J=4.3, 8.7 Hz, 1H), 4.65 - 4.57 (m, 1H), 4.51 (s, 2H), 4.50 - 4.43 (m, 1H), 4.18 - 4.13 (m, 4H), 4.11 (s, 2H), 3.34 - 3.27 (m, 2H), 3.11 - 3.06 (m, 2H), 2.80 - 2.74 (m, 2H), 2.59 - 2.45 (m, 4H), 1.94 - 1.88 (m, 1H), 1.76 (d, J=14.2 Hz, 2H), 1.49 - 1.36 (m, 2H).	formate
145	2.44	11	(MeOD); δ 9.12 (d, J=1.8 Hz, 1H), 9.02 (d, J=5.5 Hz, 1H), 8.56 (d, J=7.6 Hz, 1H), 8.39 (d, J=9.9 Hz, 1H), 8.31 (dd, J=2.2, 8.2 Hz, 1H), 7.59 (d, J=8.3 Hz, 1H), 7.52 - 7.51 (m, 5H), 7.43 - 7.32 (m, 7H), 7.11 - 7.05 (m, 3H), 7.00 (dd, J=2.1, 7.2 Hz, 1H), 6.71 (d, J=9.8 Hz, 1H), 5.55 (dd, J=5.2, 8.2 Hz, 1H), 4.63 - 4.48 (m, 6H), 4.29 (s, 2H), 4.14 (d, J=6.3 Hz, 2H), 3.52 - 3.42 (m, 2H), 3.41 - 3.36 (m, 2H), 2.99 (dd, J=10.6, 13.2 Hz, 2H), 2.50 (dd, J=6.8, 6.8 Hz, 4H), 2.02 - 1.96 (m, 1H), 1.84 (d, J=14.7 Hz, 2H), 1.49 - 1.39 (m, 2H)	TFA
146	2.49	11	(MeOD); δ 8.52 (s, 1H), 8.26 (d, J=9.9 Hz, 1H), 7.47 - 7.43 (m, 2H), 7.42 - 7.38 (m, 9H), 7.37 - 7.25 (m, 5H), 7.09 - 7.06 (m, 2H), 7.03 (d, J=8.2 Hz, 1H), 6.99 - 6.96 (m, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.37 (dd, J=4.7, 8.7 Hz, 1H), 4.49 - 4.41 (m, 3H), 4.34 - 4.25 (m, 1H), 4.22 (s, 2H), 4.12 (d, J=6.4 Hz, 2H), 3.90 (s, 2H), 3.56 (s, 2H), 3.18 - 3.13 (m, 4H), 2.50 (dd, J=11.5, 11.5 Hz, 2H), 2.42 - 2.27 (m, 4H), 1.86 - 1.78 (m, 1H), 1.68 (d, J=12.2 Hz, 2H), 1.42 - 1.30 (m, 2H)	mono-formate

Example 23

- (1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 147).




Step 1; Benzyl (R)-3-(4-(((tert-butoxycarbonyl)(2-((tert-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidine-1-carboxylate



A solution of 1-((benzyloxy)carbonyl)azetidine-3-carboxylic acid (0.306 g, 1.3 mmol) and DIPEA (0.266 mL, 1.3 mmol) in DMF (10 mL) was added with HATU (0.494 g, 0.494 mmol) and the reaction mixture stirred at room temperature for 30 minutes. *(R)*-*Tert*-butyl (2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)(piperidin-4-ylmethyl)carbamate (0.531 g, 1.0 mmol) was added and the mixture stirred for a further 18 hours. The mixture was diluted with ethyl acetate and washed with water, 10% aqueous potassium carbonate and brine (x2). The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure to afford the title compound (0.759 g, 100%).

LCMS Method 11; Rt 3.80 min; ES⁺ 749.6.


Step 2; *tert*-Butyl (R)-((1-(azetidine-3-carbonyl)piperidin-4-yl)methyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate



A solution of benzyl (R)-3-(4-(((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidine-1-carboxylate (0.759 g, 1.0 mmol) in ethanol (10 mL) was added with 10% palladium on carbon (0.8 g) and 1-methyl-1,4-cyclohexadiene (0.562 mL). The reaction mixture was heated to reflux and heated at reflux for one hour. The suspension was filtered and the filtrate evaporated at reduced pressure to afford the title compound (0.516 g, 84%).

LCMS Method 11; Rt 3.02 min; ES⁺ 615.

Step 3; (1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate (Compound 147).



5

A stirred solution of *tert*-butyl (R)-((1-(azetidine-3-carbonyl)piperidin-4-yl)methyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)carbamate (0.250 g, 0.41 mmol) and (S)-2-(3-(2-((1-benzylpiperidin-4-yl)methoxy)-1-hydroxy-2-oxo-1-phenylethyl)phenoxy)acetic acid hydrochloride (0.235 g, 0.45 mmol) in DMF (3 mL) was added with EDC (0.11 g, 0.57 mmol), HOBt (0.078 g, 0.58 mmol) and 4-DMAP (catalytic). The reaction mixture was stirred at room temperature for 72 hours. The reaction mixture was diluted with ethyl acetate and washed with water, 10% aqueous potassium carbonate and brine (x2). The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was dissolved in MeCN (2 mL) and a solution of HCl – dioxan (4 mL) added. The reaction mixture was stirred at room temperature for 1 hour. The solvent was evaporated at reduced pressure and the residue purified by reverse preparative HPLC.

15

The following compounds were prepared by this method;

20


247

N	Structure
147	
148	
149	
150	


N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
147	2.40	11	(MeOD); δ 8.43 - 8.38 (m, 1H), 7.54 - 7.48 (m, 6H), 7.41 - 7.28 (m, 6H), 7.10 - 7.05 (m, 2H), 7.00 - 6.98 (m, 1H), 6.93 (dd, J=2.3, 8.1 Hz, 1H), 6.73 - 6.68 (m, 1H), 5.46 (dd, J=5.6, 8.0 Hz, 1H), 4.61 (d, J=4.1 Hz, 3H), 4.53 - 4.40 (m, 2H), 4.29 (s, 2H), 4.26 - 4.16 (m, 1H), 4.14 (d, J=6.0 Hz, 3H), 3.89 - 3.80 (m, 1H), 3.72 (d, J=14.6 Hz, 1H), 3.52 - 3.45 (m, 2H), 3.29 (d, J=7.2 Hz, 2H), 3.19 - 2.95 (m, 6H), 2.76 (dd, J=12.7, 12.7 Hz, 1H), 2.13 - 2.09 (m, 1H), 2.02 - 1.72 (m, 4H), 1.42 (dd, J=12.9, 12.9 Hz, 2H), 1.32 - 1.24 (m, 2H).	TFA

(continued)

148	2.40	11	(MeOD); δ 8.40 (d, J=9.9 Hz, 1H), 7.54 - 7.47 (m, 6H), 7.42 - 7.25 (m, 6H), 7.06 (d, J=8.2 Hz, 2H), 6.99 (s, 1H), 6.92 (dd, J=2.1, 8.3 Hz, 1H), 6.71 (d, J=9.9 Hz, 1H), 5.46 (t, J=7.3 Hz, 1H), 4.83 - 4.72 (m, 3H), 4.61 (d, J=14.6 Hz, 1H), 4.49 - 4.33 (m, 1H), 4.28 (s, 2H), 4.20 - 4.10 (m, 3H), 3.98 (d, J=15.6 Hz, 1H), 3.51 - 3.42 (m, 2H), 3.30 - 3.15 (m, 5H), 3.10 - 2.94 (m, 5H), 2.82 (t, J=11.9 Hz, 1H), 2.70 (t, J=11.9 Hz, 1H), 2.14 - 2.10 (m, 1H), 1.98 - 1.93 (m, 2H), 1.84 - 1.78 (m, 4H), 1.70 - 1.55 (m, 1H), 1.46 - 1.20 (m, 4H).	TFA
149	2.43	11	(MeOD) d 8.41 (d, J=9.8 Hz, 1H), 7.95 (d, J=6.6 Hz, 1H), 7.51 (s, 5H), 7.40 - 7.27 (m, 6H), 7.11 - 7.05 (m, 2H), 7.01 (s, 1H), 6.96 (dd, J=2.1, 8.0 Hz, 1H), 6.71 (d, J=9.9 Hz, 1H), 5.45 (t, J=6.4 Hz, 1H), 4.61 (d, J=13.6 Hz, 1H), 4.47 (s, 2H), 4.28 (s, 2H), 4.17 - 4.08 (m, 3H), 3.72 - 3.67 (m, 1H), 3.50 - 3.43 (m, 2H), 3.28 (d, J=7.3 Hz, 2H), 3.21 - 3.13 (m, 2H), 3.10 - 2.93 (m, 5H), 2.71 - 2.62 (m, 2H), 2.13 - 2.09 (m, 1H), 1.97 - 1.91 (m, 4H), 1.86 - 1.76 (m, 4H), 1.64 - 1.47 (m, 2H), 1.43 - 1.19 (m, 6H).	TFA
150	2.42	11	(MeOD); δ 8.40 (d, J=9.9 Hz, 1H), 7.53 - 7.48 (m, 5H), 7.42 - 7.29 (m, 7H), 7.11 - 7.04 (m, 3H), 7.00 (dd, J=2.3, 8.1 Hz, 1H), 6.71 (d, J=9.8 Hz, 1H), 5.45 (dd, J=6.0, 7.8 Hz, 1H), 4.59 - 4.53 (m, 3H), 4.28 (s, 2H), 4.25 - 4.09 (m, 4H), 3.96 (d, J=13.2 Hz, 1H), 3.51 - 3.45 (m, 2H), 3.28 (d, J=7.8 Hz, 2H), 3.21 - 3.14 (m, 2H), 3.09 - 2.94 (m, 4H), 2.74 (t, J=13.2 Hz, 1H), 2.15 - 2.08 (m, 1H), 1.99 - 1.80 (m, 5H), 1.47 - 1.20 (m, 3H).	TFA

Example 24**(1-Benzylpiperidin-4-yl)methyl****(S)-2-hydroxy-2-(3-((1-(1-(4-(2-((R)-2-****hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-**

**5 yl)ethyl)amino)ethyl)benzoyl)piperidine-4-carbonyl)piperidin-4-yl)methoxy)phenyl)-
2-phenylacetate (Compound 151)**



The title compound was prepared using a similar route to that described in Example 22 using previously described compounds (see Procedures section). An acidic deprotection
10 step (as described in Example 20 Step 2) was utilized prior to the final coupling step.

The following compounds were prepared by this method;


250

[illegible]


N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
151	2.55	11	(DMSO-d ₆ , 90 °C); δ 8.20 (d, J=9.9 Hz, 1H), 7.49 - 7.43 (m, 5H), 7.35 (dd, J=7.6, 7.6 Hz, 9H), 7.27 - 7.21 (m, 1H), 7.16 (d, J=8.3 Hz, 1H), 7.02 (d, J=8.2 Hz, 1H), 6.96 - 6.92 (m, 2H), 6.89 - 6.86 (m, 1H), 6.57 (d, J=9.9 Hz, 1H), 5.38 (dd, J=4.3, 8.6 Hz, 1H), 4.19 - 4.19 (m, 4H), 4.11 - 3.97 (m, 3H), 3.82 (d, J=6.3 Hz, 2H), 3.31 (t, J=8.2 Hz, 2H), 3.28 - 3.08 (m, 9H), 3.01 - 2.75 (m, 5H), 2.02 - 1.97 (m, 1H), 1.81 - 1.53 (m, 11H), 1.28 - 1.18 (m, 2H).	TFA
152	2.53	11	(MeOD); δ 8.50 (s, 1H), 8.37 (d, J=9.9 Hz, 1H), 7.88 (d, J=8.3 Hz, 2H), 7.44 - 7.33 (m, 12H), 7.32 - 7.24 (m, 2H), 7.04 (d, J=8.2 Hz, 1H), 7.03 - 6.98 (m, 2H), 6.90 (dd, J=2.1, 7.9 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.39 (dd, J=4.8, 8.7 Hz, 1H), 4.57 (d, J=12.2 Hz, 1H), 4.29 (dd, J=15.6, 38.7 Hz, 2H), 4.12 (d, J=6.7 Hz, 2H), 4.01 (d, J=13.4 Hz, 1H), 3.92 (s, 2H), 3.84 (d, J=6.0 Hz, 2H), 3.28 - 3.07 (m, 9H), 2.79 - 2.67 (m, 1H), 2.52 - 2.52 (m, 2H), 2.12 - 2.06 (m, 1H), 1.95 - 1.79 (m, 3H), 1.70 (d, J=13.6 Hz, 2H), 1.48 - 1.28 (m, 4H).	mono-formate
153	2.53	11	¹ H NMR (400 MHz, MeOD) d 8.40 (s, 2H), 8.14 (d, J=9.9 Hz, 1H), 7.41 (s, 1H), 7.39 - 7.35 (m, 1H), 7.32 - 7.19 (m, 11H), 7.14 - 7.08 (m, 2H), 6.89 - 6.82 (m, 3H), 6.77 - 6.74 (m, 1H), 6.51 (d, J=9.9 Hz, 1H), 5.20 (dd, J=5.3, 7.8 Hz, 1H), 4.43 (d, J=12.8 Hz, 1H), 4.24 - 4.08 (m, 2H), 4.04 (s, 2H), 3.98 - 3.85 (m, 3H), 3.81 (s, 3H), 3.70 (d, J=6.0 Hz, 2H), 3.58 (s, 2H), 3.03 - 2.84 (m, 5H), 2.66 - 2.53 (m, 1H), 2.13 (dd, J=11.5, 11.5 Hz, 2H), 1.96 - 1.91 (m, 1H), 1.83 - 1.70 (m, 2H), 1.63 - 1.54 (m, 1H), 1.48 (d, J=13.3 Hz, 2H), 1.36 - 1.24 (m, 1H), 1.17 (q, J=12.5 Hz, 3H).	formate
154	2.50	11	(MeOD); δ 8.48 (s, 2H), 8.32 (d, J=9.9 Hz, 1H), 7.97 (d, J=8.3 Hz, 2H), 7.62 (d, J=8.3 Hz, 2H), 7.45 (s, 5H), 7.43 - 7.34 (m, 5H), 7.27 (dd, J=7.8, 7.8 Hz, 2H), 7.04 (d, J=8.2 Hz, 1H), 6.99 (d, J=2.3 Hz, 2H), 6.91 (dd, J=1.9, 8.2 Hz, 1H), 6.67 (d, J=9.8 Hz, 1H), 5.44 - 5.39 (m, 1H), 4.61 - 4.60 (m, 1H), 4.35 - 4.27 (m, 4H), 4.16 - 4.08 (m, 4H), 4.02 (d, J=13.7 Hz, 1H), 3.85 (d, J=6.0 Hz, 2H), 3.28 (d, J=12.3 Hz, 2H), 3.20 (d, J=7.4 Hz, 3H), 2.79 - 2.68 (m, 4H), 2.11 - 2.06 (m, 1H), 1.96 - 1.86 (m, 2H), 1.76 (d, J=13.2 Hz, 2H), 1.51 - 1.39 (m, 3H), 1.37 - 1.26 (m, 1H).	formate

(continued)

155	2.52	11	(DMSO-d ₆); d 10.50 (s, 2H), 9.44 - 9.44 (m, 1H), 9.20 - 9.20 (m, 2H), 8.74 (dd, J=4.3, 4.3 Hz, 1H), 8.10 (d, J=9.9 Hz, 1H), 7.98 (d, J=7.9 Hz, 1H), 7.48 (s, 5H), 7.42 (s, 1H), 7.34 (d, J=3.8 Hz, 4H), 7.32 - 7.21 (m, 3H), 7.14 (d, J=8.3 Hz, 1H), 6.99 (d, J=8.2 Hz, 1H), 6.89 (d, J=7.8 Hz, 3H), 6.64 (s, 1H), 6.58 (d, J=9.9 Hz, 1H), 6.34 - 6.12 (m, 1H), 5.39 - 5.35 (m, 1H), 4.44 (d, J=12.5 Hz, 1H), 4.31 - 4.19 (m, 4H), 4.02 (d, J=6.4 Hz, 2H), 3.86 - 3.77 (m, 4H), 3.34 (d, J=11.8 Hz, 2H), 3.11 - 3.01 (m, 4H), 2.96 - 2.87 (m, 2H), 2.70 - 2.64 (m, 1H), 2.03 - 2.00 (m, 1H), 1.82 - 1.67 (m, 6H), 1.42 - 1.24 (m, 3H), 1.20 - 1.11 (m, 1H).	TFA
------------	------	----	--	-----

Example 25

(1-Benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-(5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)thiophene-2-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate
 5
 (Compound 156)




The title compound was prepared using a similar route to that described in Example 23 using previously described compounds (see Procedures section). A base promoted deprotection step (as described in Example 1 Step 4) was utilized prior to the final coupling step.

5 The following compounds were prepared by this method;

N	Structure
157	

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
156	2.53	11	(MeOD); δ 8.53 (s, 1H), 8.41 (d, J=9.9 Hz, 1H), 7.40 - 7.25 (m, 14H), 7.05 (d, J=8.2 Hz, 1H), 6.99 - 6.97 (m, 2H), 6.90 (dd, J=2.1, 8.0 Hz, 1H), 6.70 (d, J=9.8 Hz, 1H), 5.42 (dd, J=5.0, 8.5 Hz, 1H), 4.44 (s, 4H), 4.11 (dd, J=2.8, 6.4 Hz, 2H), 3.87 - 3.81 (m, 4H), 3.23 - 3.05 (m, 7H), 3.00 (d, J=6.9 Hz, 3H), 2.39 (t, J=12.9 Hz, 2H), 2.15 - 2.09 (m, 2H), 1.98 - 1.88 (m, 4H), 1.80 - 1.74 (m, 1H), 1.67 (d, J=12.8 Hz, 2H), 1.45 - 1.29 (m, 6H).	mono-formate
157	2.55	11	(DMSO-d ₆ , 100 °C); δ 8.20 (d, J=9.9 Hz, 1H), 7.48 - 7.46 (m, 5H), 7.41 - 7.21 (m, 6H), 7.16 (d, J=8.3 Hz, 1H), 7.02 (d, J=8.2 Hz, 1H), 6.97 - 6.93 (m, 2H), 6.89 - 6.86 (m, 1H), 6.58 (d, J=9.9 Hz, 1H), 5.38 (dd, J=4.8, 8.2 Hz, 1H), 4.26 - 4.06 (m, 8H), 3.83 (d, J=6.3 Hz, 2H), 3.20 - 3.14 (m, 2H), 3.01 - 2.98 (m, 2H), 2.93 - 2.76 (m, 7H), 2.61 - 2.60 (m, 2H), 2.10 - 1.99 (m, 2H), 1.80 - 1.69 (m, 12H), 1.62 - 1.48 (m, 6H), 1.29 - 1.11 (m, 4H).	TFA

Example 26




1-Benzylpiperidin-4-yl

(S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-


hydroxy-2-oxo-1,2-dihydroquinolin-5-

5 **yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate (Compound**

158)




Step 1; Methyl (S)-2-(3-(3-((tert-butoxycarbonyl)amino)propoxy)phenyl)-2-hydroxy-2-phenylacetate



A stirred solution of methyl (S)-2-hydroxy-2-(3-hydroxyphenyl)-2-phenylacetic acid (3.0 g, 11.62 mmol) in DMF (11 mL) was added with potassium carbonate (3.21 g, 23.23 mmol) followed by a solution of *tert*-butyl (3-bromopropyl)carbamate (3.46 g, 14.52 mmol) in DMF (11 mL). The reaction mixture was heated at 60 °C for 18 hours. The reaction mixture was dissolved in ethyl acetate and washed with aqueous 1M sodium hydroxide (x2), water and brine. The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was purified by flash column chromatography (eluent 100% i-hexane to 60% ethyl acetate / i-hexane) to afford the title compound (4.71 g, 98%).

¹H NMR (400 MHz, DMSO-d₆); δ 7.38 – 7.27(m, 6H), 6.95 - 6.88 (m, 4H), 6.70 (s, 1H), 3.96 (dd, J=6.3, 6.3 Hz, 2H), 3.77 (s, 3H), 3.11 (q, J=6.5 Hz, 2H), 1.90 - 1.81 (m, 2H), 1.42 (s, 9H).

Step 2; (S)-2-(3-(3-((tert-Butoxycarbonyl)amino)propoxy)phenyl)-2-hydroxy-2-phenylacetic acid




A stirred solution of methyl (S)-2-(3-(3-((tert-butoxycarbonyl)amino)propoxy)phenyl)-2-hydroxy-2-phenylacetate (4.71 g, 11.34 mmol) in THF / methanol (57 mL / 57 mL) was added with aqueous 2M sodium hydroxide (57 mL). The reaction mixture was stirred at room temperature for 18 hours. The solvent was concentrated under reduced pressure to 1/3 volume. The solution was washed with ether and

the pH adjusted to pH2 and the solution was extracted with ethyl acetate. The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure to afford the title compound (4.1 g, 90%).

¹H NMR (400 MHz, DMSO-d₆); δ 7.45 - 7.41 (m, 2H), 7.39 - 7.26 (m, 4H),
5 7.01 - 6.96 (m, 2H), 6.94 - 6.87 (m, 2H), 6.37 (br s), 3.96 (dd, J=6.3, 6.3 Hz, 2H), 3.14 - 3.07 (m, 2H), 1.89 - 1.81 (m, 2H), 1.42 (s, 9H).


Step 3; 1-Benzylpiperidin-4-yl (S)-2-(3-(3-aminopropoxy)phenyl)-2-hydroxy-2-phenylacetate dihydrochloride



10 Carbonyldiimidazole (0.242 g, 1.50 mmol) was added to a solution of (S)-2-(3-(3-
((*tert*-butoxycarbonyl)amino)propoxy)phenyl)-2-hydroxy-2-phenylacetic acid (0.20 g, 0.5
mmol) in DMF (2 mL) and the mixture stirred at room temperature for 15 mins. 1-
Benzylpiperidin-4-ol (0.381 g, 1.99 mmol) in DMF (0.5 mL) was added and the reaction
mixture heated at 60 °C for 18 hours. The reaction mixture was diluted with ethyl acetate
15 and washed with saturated aqueous sodium carbonate (x2) and brine. The organic phase
was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced
pressure. The residue was purified by flash column chromatography (eluent 100% DCM
to 5% MeOH / DCM) to afford major product. This material was dissolved in dioxane (1
mL) and a solution of 4M HCl in dioxane (2 mL) added. The reaction mixture stirred at
20 room temperature for 1.5 hours. The solvent was evaporated at reduced pressure to afford
the title compound (0.157 g, 57%).

LCMS Method 11; Rt 2.35; ES⁺ 475.

Step 4; 1-Benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate (Compound 158)



5





The title compound was prepared as described in Example 5.

The following compounds were prepared according to Example 26;

N	Structure
159	
160	

(continued)

258

161	
162	
163	
164	

(continued)

[illegible]

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
158	2.42	11	(MeOD); δ 8.44 (s, 2H), 8.30 (d, J=9.9 Hz, 1H), 7.89 (d, J=8.2 Hz, 2H), 7.60 (d, J=8.2 Hz, 2H), 7.45 - 7.34 (m, 10H), 7.29 (dd, J=8.1, 8.1 Hz, 2H), 7.05 - 6.93 (m, 4H), 6.66 (d, J=9.8 Hz, 1H), 5.42 (t, J=6.6 Hz, 1H), 5.12 - 5.04 (m, 1H), 4.33 (s, 2H), 4.09 (t, J=6.0 Hz, 2H), 3.82 (s, 2H), 3.59 (t, J=6.8 Hz, 2H), 3.21 (d, J=6.7 Hz, 2H), 2.85 - 2.76 (m, 2H), 2.62 - 2.62 (m, 2H), 2.16 - 2.07 (m, 2H), 2.01 - 1.96 (m, 2H), 1.90 - 1.82 (m, 2H).	formate
159	2.44	11	(DMSO- d_6); δ 10.54 (m, 2H), 9.96 - 9.96 (m, 1H), 9.18 (s, 2H), 8.66 (dd, J=5.4, 5.4 Hz, 1H), 8.13 (d, J=8.9 Hz, 1H), 7.95 (d, J=8.1 Hz, 2H), 7.66 (d, J=8.1 Hz, 2H), 7.52 (d, J=5.3 Hz, 6H), 7.41 - 7.38 (m, 2H), 7.36 (s, 3H), 7.33 - 7.25 (m, 1H), 7.18 (d, J=8.3 Hz, 1H), 7.03 (d, J=8.1 Hz, 1H), 6.97 - 6.91 (m, 3H), 6.63 (d, J=9.9 Hz, 1H), 6.25 (s, 1H), 5.39 (d, J=7.8 Hz, 1H), 4.43 - 4.15 (m, 6H), 4.07 - 4.02 (m, 3H), 3.47 - 3.47 (m, 4H), 3.17 - 3.02 (m, 3H), 2.88 - 2.73 (m, 1H), 2.19 - 2.19 (m, 1H), 2.06 - 1.98 (m, 2H), 1.84 - 1.76 (m, 1H).	TFA
160	2.44	11	(DMSO- d_6); δ 10.54 (m, 2H), 10.01 (s, 1H), 9.18 (s, 2H), 8.66 (dd, J=5.6, 5.6 Hz, 1H), 8.13 (d, J=9.9 Hz, 1H), 7.95 (d, J=8.1 Hz, 2H), 7.66 (d, J=8.3 Hz, 2H), 7.52 (m, 5H), 7.40 - 7.34 (m, 5H), 7.30 (dd, J=8.0, 8.0 Hz, 1H), 7.18 (d, J=8.1 Hz, 1H), 7.03 (d, J=8.1 Hz, 1H), 6.98 - 6.90 (m, 3H), 6.78 - 6.71 (m, 1H), 6.63 (d, J=9.9 Hz, 1H), 6.24 (s, 1H), 5.40 (d, J=9.3 Hz, 1H), 4.41 - 4.15 (m, 7H), 4.08 - 4.01 (m, 2H), 3.48 (dd, J=6.9, 11.1 Hz, 4H), 3.17 - 3.05 (m, 4H), 2.87 - 2.77 (m, 1H), 2.06 - 1.98 (m, 2H), 1.83 - 1.74 (m, 1H).	TFA


(continued)

161	2.41	11	(DMSO-d ₆); δ 8.54 - 8.47 (m, 1H), 8.20 (s, 2H), 8.13 (dd, J=3.1, 9.2 Hz, 1H), 7.79 (d, J=8.3 Hz, 2H), 7.41 (d, J=8.3 Hz, 2H), 7.33 (d, J=4.4 Hz, 4H), 7.32 - 7.22 (m, 2H), 7.07 (d, J=8.2 Hz, 1H), 6.94 - 6.87 (m, 4H), 6.48 (d, J=9.8 Hz, 1H), 5.10 (dd, J=4.3, 8.0 Hz, 1H), 3.99 (dd, J=4.0, 5.9 Hz, 4H), 3.84 (s, 2H), 3.40 (td, J=7.6, 19.3 Hz, 2H), 2.77 - 2.61 (m, 5H), 1.99 - 1.88 (m, 4H), 1.77 - 1.47 (m, 9H), 1.14 - 1.05 (m, 2H).	formate
162	2.46	11	(DMSO-d ₆); δ 8.50 (t, J=7.1 Hz, 1H), 8.22 (s, 2H), 8.13 (d, J=9.9 Hz, 1H), 7.79 (d, J=8.2 Hz, 2H), 7.41 (d, J=8.3 Hz, 2H), 7.33 (d, J=4.4 Hz, 4H), 7.32 - 7.22 (m, 2H), 7.07 (d, J=8.2 Hz, 1H), 6.94 - 6.87 (m, 4H), 6.48 (d, J=9.9 Hz, 1H), 5.10 (dd, J=4.3, 8.0 Hz, 1H), 4.02 - 3.96 (m, 4H), 3.83 (s, 2H), 3.44 - 3.37 (m, 2H), 2.79 - 2.64 (m, 4H), 2.17 (s, 3H), 1.99 - 1.87 (m, 4H), 1.53 - 1.45 (m, 3H), 1.21 - 1.10 (m, 2H).	formate
163	2.41	11	(MeOD); δ 8.43 (s, 2H), 8.30 (d, J=9.8 Hz, 1H), 7.90 (d, J=8.3 Hz, 2H), 7.61 (d, J=8.3 Hz, 2H), 7.43 (dd, J=1.7, 8.0 Hz, 2H), 7.38 - 7.24 (m, 10H), 7.05 - 6.99 (m, 3H), 6.92 (dd, J=2.0, 8.2 Hz, 1H), 6.66 (d, J=9.9 Hz, 1H), 5.46 - 5.39 (m, 2H), 4.34 (s, 2H), 4.06 (dd, J=6.0, 6.0 Hz, 2H), 3.90 - 3.77 (m, 2H), 3.58 (dd, J=6.8, 6.8 Hz, 2H), 3.22 (d, J=6.7 Hz, 2H), 3.15 (dd, J=5.5, 12.4 Hz, 1H), 3.01 - 2.82 (m, 3H), 2.38 - 2.28 (m, 1H), 2.10 (ddd, J=8.2, 8.2, 8.2 Hz, 2H), 2.00 - 1.93 (m, 1H).	formate
164	2.41	11	(MeOD); δ 8.43 (s, 2H), 8.30 (d, J=9.8 Hz, 1H), 7.90 (d, J=8.2 Hz, 2H), 7.61 (d, J=8.2 Hz, 2H), 7.44 - 7.24 (m, 12H), 7.05 - 7.00 (m, 3H), 6.92 (dd, J=2.3, 8.3 Hz, 1H), 6.66 (d, J=9.8 Hz, 1H), 5.41 (q, J=6.5 Hz, 2H), 4.33 (s, 2H), 4.06 (dd, J=6.0, 6.0 Hz, 2H), 3.81 (q, J=12.5 Hz, 2H), 3.58 (dd, J=6.8, 6.8 Hz, 2H), 3.22 (d, J=6.7 Hz, 2H), 3.13 (dd, J=5.7, 12.5 Hz, 1H), 2.97 - 2.88 (m, 2H), 2.88 - 2.79 (m, 1H), 2.38 - 2.27 (m, 1H), 2.14 - 2.06 (m, 2H), 1.98 - 1.91 (m, 1H).	formate
165	2.41	11	(MeOD); δ 8.46 (s, 2H), 8.31 (d, J=9.9 Hz, 1H), 7.89 (d, J=8.3 Hz, 2H), 7.60 (d, J=8.3 Hz, 2H), 7.44 - 7.41 (m, 2H), 7.38 - 7.24 (m, 10H), 7.05 - 6.99 (m, 3H), 6.93 (dd, J=2.1, 8.2 Hz, 1H), 6.66 (d, J=9.8 Hz, 1H), 5.41 (dd, J=5.7, 7.7 Hz, 1H), 5.25 - 5.18 (m, 1H), 4.31 (s, 2H), 4.08 (t, J=6.0 Hz, 2H), 3.85 (dd, J=6.8, 9.7 Hz, 2H), 3.77 (s, 2H), 3.59 (t, J=6.8 Hz, 2H), 3.35 - 3.30 (m, 2H), 3.23 - 3.19 (m, 2H), 2.15 - 2.07 (m, 2H).	formate

Example 27

1-Benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)phenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate

5 (Compound 166)



A suspension of (R)-4-(((*tert*-butoxycarbonyl)(2-((*tert*-butyldimethylsilyl)oxy)-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoic acid (0.337 g, 0.59 mmol) in toluene (5 mL) was added with triethylamine (0.108 mL, 0.77 mmol) and diphenyl phosphorylazide (0.153 mL, 0.71 mmol). The reaction mixture was heated at 90 °C for 3.5 hours. A solution of 1-benzylpiperidin-4-yl (S)-2-(3-((4-(aminomethyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate dihydrochloride (0.37 g, 0.59 mmol) and triethylamine (0.165 mL, 1.19 mmol) in DMF (2 mL) was added and the reaction mixture heated at 90 °C for 18 hours. The reaction mixture was diluted with ethyl acetate and washed with saturated aqueous sodium hydrogen carbonate and brine. The organic phase was dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated at reduced pressure. The residue was dissolved in MeCN (1 mL) and a solution of HCl – dioxan (4 mL) added. The reaction mixture was stirred at room temperature for 1 hour. The solvent was evaporated at reduced pressure and the residue purified by reverse preparative HPLC.

The following compounds were prepared by this method;

N	Structure
167	
168	

N	Rt (min)	Method	NMR data (400 MHz)	Salt (2 eq unless stated)
166	2.60	11	(MeOD); δ 8.57 (s, 1H), 8.26 (d, J=9.8 Hz, 1H), 7.47 (d, J=8.5 Hz, 2H), 7.39 - 7.32 (m, 16H), 7.29 - 7.23 (m, 2H), 7.04 - 6.94 (m, 4H), 6.65 (d, J=9.8 Hz, 1H), 5.34 (dd, J=4.3, 9.0 Hz, 1H), 5.05 (s, 2H), 4.42 (s, 2H), 4.11 (s, 2H), 4.10 - 4.03 (m, 2H), 3.64 (s, 2H), 3.16 - 3.06 (m, 2H), 2.95 (d, J=11.8 Hz, 2H), 2.17 (dd, J=9.9, 12.2 Hz, 2H), 1.71 - 1.65 (m, 1H), 1.57 (d, J=13.2 Hz, 2H), 1.33 - 1.18 (m, 2H).	TFA
167	2.57	11	(MeOD); δ 8.57 (s, 1H), 8.22 (d, J=9.9 Hz, 1H), 7.40 - 7.18 (m, 18H), 7.03 - 6.95 (m, 4H), 6.86 (dd, J=1.9, 8.1 Hz, 1H), 6.64 (d, J=9.9 Hz, 1H), 5.32 (dd, J=5.0, 8.4 Hz, 1H), 5.05 (s, 2H), 4.43 (s, 2H), 4.09 (s, 2H), 4.06 (dd, J=3.5, 6.5 Hz, 2H), 3.85 (s, 3H), 3.53 (s, 2H), 3.09 - 3.05 (m, 2H), 2.91 - 2.86 (m, 2H), 2.08 - 1.99 (m, 2H), 1.65 - 1.62 (m, 1H), 1.54 (d, J=13.2 Hz, 2H), 1.32 - 1.19 (m, 2H).	mono-formate

(continued)

168	2.58	11	(MeOD); δ 8.57 (s, 1H), 8.25 (d, J=9.9 Hz, 1H), 8.08 (d, J=8.3 Hz, 1H), 7.37 - 7.24 (m, 18H), 7.08 - 6.94 (m, 4H), 6.64 (d, J=9.9 Hz, 1H), 5.32 (dd, J=4.4, 8.7 Hz, 1H), 5.05 (s, 2H), 4.93 (d, J=2.1 Hz, 1H), 4.42 (s, 2H), 4.09 - 4.04 (m, 4H), 3.91 (s, 3H), 3.60 (s, 2H), 3.11 - 3.05 (m, 2H), 2.95 - 2.88 (m, 2H), 2.11 (t, J=11.4 Hz, 1H), 1.71 - 1.60 (m, 1H), 1.59 - 1.53 (m, 2H), 1.24 (ddd, J=6.3, 12.3, 18.3 Hz, 2H).	mono-formate
-----	------	----	---	--------------

Biological characterization

Example 28

M3 Receptor radioligand binding assay

Human M3 receptor membranes (15ug/well) from Perkin Elmer were incubated with 0.52nM Scopolamine Methyl Chloride, [N-methyl-3H] with or without test compounds, or a saturating concentration of Atropine (5 μ M) for the determination of non-specific binding. The assay was carried out in 96-well polypropylene plates in a volume of 250ul. The assay buffer used was 50mM Tris-HCl, 154mM NaCl (pH 7.4). The final assay concentration of DMSO was 0.5% (v/v). The plates were sealed and incubated for 2h at room temperature on an orbital shaker (slow speed). Membranes were harvested onto 96-well unifilter GF/C filter plates pre-treated with 0.5% polyethyleneimine (v/v), using a filter manifold, washed four times with 200ul of assay buffer. The plates were dried before addition of 50ul of microscint-0, sealed then read in a Trilux Microbeta scintillation counter. IC₅₀ values are determined from competition curves using a non-linear curve fitting program. K_i values were calculated from IC₅₀ values by the Cheng and Prusoff equation.

The M3 K_i values of the compounds according to the invention are less than 50 nM, most of them even less than 10 nM. The preferred compounds according to the invention have K_i value less than 4 nM or even less than 3 nM for the enantiomeric mixture and at least for one of the enantiomeric pure form (S or R) on the stereogenic center (2).

Example 29 **β_2 adrenoceptor radioligand binding assay**

Human β_2 adrenoceptor membranes (7.5 ug/well) from Perkin Elmer were incubated with 0.3 nM ¹²⁵-I Cyanopindolol with or without test compounds, or a saturating concentration of s-propranolol (2 μ M) for the determination of non-specific binding. The assay was carried out in 96-well polypropylene plates in a volume of 200 μ l. The assay buffer used was 25 mM HEPES, 0.5% BSA (w/v), 1 mM EDTA, 0.02% ascorbic acid (v/v), (pH 7.4). The final assay concentration of DMSO was 0.5% (v/v). The plates were sealed and incubated for 1h at room temperature on an orbital shaker (slow speed). Membranes were harvested onto 96-well unifilter GF/C filter plates pre-treated with 0.5% polyethyleneimine (v/v), using a filter manifold, washed six times with 200 μ l of wash buffer containing 10 mM HEPES and 500 mM NaCl. The plates were dried before addition of 50 μ l of microscint-0, sealed then read in a Trilux Microbeta scintillation counter. IC₅₀ values are determined from competition curves using a non-linear curve fitting program. Ki values were calculated from IC₅₀ values by the Cheng and Prusoff equation.

The β_2 Ki values of the compounds according to the invention are less than 50 nM, most of them even less than 10 nM.


In the following table the compounds tested are classified in terms of binding affinity according to the following ranges:

- +++ : IC₅₀ < 0.3 nM
++ : IC₅₀ in the range 0.3-3 nM
+ : IC₅₀ > 3 nM

No	M3	B2	No	M3	B2	No	M3	B2	No	M3	B2	No	M3	B2
1	++	+++	32J	++	+++	62	+++	+++	104	+++	++	146	++	+
2	++	++	32K	++	+++	63	+++	+++	105	++	++	147	++	+++
3	++	+++	32L	++	+	64	+++	++	106	++	++	148	++	+++
4	++	+++	32M	++	+	65	+++	++	107	++	+++	149	+++	++
5	++	++	33	+++	+++	66	++	+++	108	++	++	150	+++	+++
6	+	+++	33A	++	+++	67	+++	+++	109	++	++	151	+++	+++
7	+	++	34	+	++	68	++	++	110	++	++	152	+++	++
8	+	++	35	+++	+++	69	++	++	111	++	++	153	+++	+++
9	++	+++	36	+++	+++	70	++	++	112	++	+++	154	+++	+++
10	++	++	36A	++	++	71	++	+	113	++	++	155	++	+++
11	+++	+++	37	+++	+++	72	++	+	114	++	++	156	+++	+++
12	+	+	38	+++	+++	73	++	+++	115	++	++	157	++	+
12A	++	+++	39	++	+++	74	++	++	116	++	+++	158	++	+++
13	+++	+++	40	++	++	75	++	+	117	++	+++	159	++	+++
14	+	+++	41	++	++	76	++	++	118	+++	++	160	++	+++
15	+++	+++	42	+++	+++	77	++	++	119	+++	++	161	++	+++
16	+	++	42A	+++	++	78	+++	+++	120	+++	+++	162	++	+++
17	+	++	42B	+++	+++	79	+++	+++	121	+++	+++	163	++	+++
18	+	+++	42B	++	+++	80	++	+++	122	++	+	164	++	++
19	++	+++	42C	++	+++	81	++	++	123	++	++	165	++	+++
20	+	+++	42D	++	+++	82	++	++	124	++	++	166	++	++
21	++	+++	42E	++	+++	83	+	++	125	++	++	167	++	++
22	+	+++	42F	++	+++	84	+	++	126	++	++	168	++	++
23	++	++	43	+++	+++	85	++	+++	127	+++	+++			
24	+	+++	44	++	+++	86	++	+++	128	++	+++			
25	++	+++	45	++	++	87	++	++	129	++	+++			
26	++	+++	46	++	+++	88	+++	++	130	++	++			
27	+	++	47	++	+++	89	++	+++	131	++	+++			
28	++	++	48	+	+++	90	++	++	132	++	+++			
29	+++	+++	49	++	+++	91	++	+++	133	+++	+++			
30	++	++	50	++	+	92	+++	+++	134	++	+++			
31	+++	+++	51	++	+++	93	++	+++	135	++	+++			
32	++	+++	52	++	+++	94	+++	+++	136	++	+++			
32A	++	+++	53	+	+	95	+++	+++	137	++	+++			
32B	+++	+++	54	++	+++	96	+++	+++	138	++	++			
32C	++	+++	55	++	+	97	++	+++	139	++	+++			
32D	++	+++	56	+	+	98	++	++	140	++	++			
32E	++	++	57	++	++	99	++	++	141	++	++			
32F	++	+++	58	++	++	100	+++	++	142	++	+++			
32G	++	++	59	++	+++	101	+++	+++	143	++	++			
32H	++	+++	60	++	++	102	+++	+++	144	++	+++			
32I	+++	+++	61	++	+++	103	+++	++	145	++	++			


CLAIMS

1. A Compound of general formula I




wherein


Q is a group of formula



Y is selected from **Y2** and **Y1** which are divalent groups of formula



or











wherein


A1 and **A2** are independently absent or selected from (C₁-C₁₂)alkylene, (C₃-C₈)cycloalkylene and (C₃-C₈)heterocycloalkylene optionally substituted by one or more substituents selected from (C₁-C₆)alkyl, aryl(C₁-C₆)alkyl and heteroaryl(C₁-C₆)alkyl;

B is absent or is selected from (C₃-C₈)cycloalkylene, (C₃-C₈)heterocycloalkylene, arylene or heteroarylene optionally substituted by one or more groups selected from -OH, halogens, -CN, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)haloalkyl, (C₁-C₆)haloalkoxy and aryl(C₁-C₆)alkyl;


- 5 **C** is absent or is selected from -O-, -C(O)-, -OC(O)-, -(O)CO-, -S-, -S(O)-, -S(O)₂- and -N(R₇)-, or is one of the following groups **C1-C23**




268




C14




C15




C16




C17




C18




C19




C20



C21



C22



C23

wherein **R**₇ is in each occurrence independently H or selected from linear or branched

(C₁-C₈)alkyl, aryl(C₁-C₆)alkyl, arylsulfanyl, arylsulfinyl, arylsulfonyl,

5 (C₃-C₈)cycloalkyl, (C₃-C₈)heterocycloalkyl, aryl, and heteroaryl;

D is absent or is selected from (C₁-C₁₂)alkylene, (C₂-C₁₂)alkenylene, (C₂-C₆)alkynylene, arylene, heteroarylene, (C₃-C₈)cycloalkylene, (C₃-C₈)heterocycloalkylene; said arylene, heteroarylene, (C₃-C₈)cycloalkylene and (C₃-C₈)heterocycloalkylene being

optionally substituted by one or more groups selected from $-\text{OH}$, halogen, $-\text{CN}$, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, $(\text{C}_1\text{-C}_6)\text{haloalkyl}$, $(\text{C}_1\text{-C}_6)\text{haloalkoxy}$ and $\text{aryl}(\text{C}_1\text{-C}_6)\text{alkyl}$;

n is at each occurrence independently 0 or an integer from 1 to 3;

5 **m** is at each occurrence independently an integer from 1 to 3;

E is absent or is selected from $-\text{O}-$, $-\text{NR}_7-$, $-\text{NR}_7\text{-C(O)-}$, $-\text{C(O)-NR}_7-$, $-\text{OC(O)-}$, $-\text{C(O)-}(\text{CH}_2)_n\text{-O-}$; $-\text{NR}_7\text{-C(O)-}(\text{CH}_2)_n\text{-O-}$, $-\text{NR}_7\text{-C(O)-NR}_7-$ and $-\text{S-}$;


G is arylene or heteroarylene, optionally substituted by one or more substituents selected from halogen atoms, $-\text{OH}$, oxo ($=\text{O}$), $-\text{SH}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{CON}(\text{R}_6)_2$,
 10 $-\text{NH}_2$, $-\text{NHCOR}_6$, $-\text{CO}_2\text{R}_6$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfanyl}$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfinyl}$, $(\text{C}_1\text{-C}_{10})\text{alkylsulfonyl}$, $(\text{C}_1\text{-C}_{10})\text{alkyl}$, aryl, haloaryl, heteroaryl and $(\text{C}_1\text{-C}_{10})\text{alkoxy}$;

R₁ is selected from $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_3\text{-C}_8)\text{heterocycloalkyl}$, aryl, heteroaryl, $\text{aryl}(\text{C}_1\text{-C}_6)\text{alkyl}$, $\text{heteroaryl}(\text{C}_1\text{-C}_6)\text{alkyl}$ and $(\text{C}_3\text{-C}_8)\text{cycloalkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$,
 15 optionally substituted by one or more group selected independently from halogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, and $(\text{C}_1\text{-C}_{10})\text{alkoxy}$;

s is 0 or an integer from 1 to 3;


R₂ is a nitrogen containing group which is selected from:

- a group (a) which is $-\text{NR}_3\text{R}_4$ wherein R_3 and R_4 are independently hydrogen or
 20 $(\text{C}_1\text{-C}_4)\text{alkyl}$; and
 - a group (b) of formula J1, J2, J3, J4 or J5



R₅ is a group of formula **K**

270

**K**

wherein **p** is 0 or an integer from 1 to 4; **q** is 0 or an integer from 1 to 4;


P is absent or is selected from the divalent group consisting of O, S, SO, SO₂, CO, NR₆, CH=CH, N(R₆)SO₂, N(R₆)COO, N(R₆)C(O), SO₂N(R₆), OC(O)N(R₆) and C(O)N(R₆);

W is selected from H, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, aryl and heteroaryl, optionally substituted by one or more substituents selected independently from halogen atoms, -OH, oxo (=O), -SH, -NO₂, -CN, -CON(R₆)₂, -NH₂, -NHCOR₆, -CO₂R₆, (C₁-C₁₀)alkylsulfanyl, (C₁-C₁₀)alkylsulfinyl, (C₁-C₁₀)alkylsulfonyl, (C₁-C₁₀)alkyl and (C₁-C₁₀)alkoxy;

R₆ is at each occurrence independently H or selected from (C₁-C₁₀)alkyl, (C₁-C₆)haloalkyl, (C₂-C₆)alkynyl, (C₂-C₆)alkenyl, (C₃-C₈)cycloalkyl, heteroaryl and aryl optionally substituted by one or more substituents selected from halogen atoms, -OH, oxo (=O), -SH, -NO₂, -CN, -CONH₂, -COOH, (C₁-C₁₀)alkoxycarbonyl, (C₁-C₁₀)alkylsulfanyl, (C₁-C₁₀)alkylsulfinyl, (C₁-C₁₀)alkylsulfonyl, (C₁-C₁₀)alkyl and (C₁-C₁₀)alkoxy;

and pharmaceutically acceptable salts or solvates thereof.


2. A compound according to claim 1, wherein **R₂** is a group of formula **J3**:

**J3**

R₅ is a group of formula **K**, wherein **p** is 0 or 1, **P** is absent or is CO, **q** is absent or is 1 and


W is H or is selected from (C₁-C₆)alkyl and aryl.

3. A compound according to claim 2 wherein **R₅** is methyl or benzyl.
4. A compound according to claim 1 wherein **G** is arylene and **R₁** is aryl, optionally substituted by one or more group independently selected from halogen, (C₁-C₈)alkyl and (C₁-C₁₀)alkoxy.
5. A compound according to claim 4 wherein **A1** and **A2** are independently absent or selected from methylene, ethylene, propylene, butylene, pentylene, hexylene, heptylene, octylene and nonylene, **G** is phenylene, and **R₁** is phenyl, optionally substituted by one or more group independently selected from halogen, (C₁-C₈)alkyl and (C₁-C₁₀)alkoxy.
6. A compound according to claim 1 wherein **E** is -O- or -C(O)-(CH₂)_n-O- or -NR₇-C(O)-(CH₂)_n-O-; **G** is phenylene wherein **E** is linked to the phenyl ring **G** in meta position, and **R₁** is phenyl, optionally substituted by one or more group selected from halogen, (C₁-C₈)alkyl or (C₁-C₁₀)alkoxy groups.
7. A compound according to claim 1, wherein **R₂** is a nitrogen containing group selected from J1, J2, or J5



and all the other variables are as defined in claim 1.

8. A compound according to claim 1 wherein the absolute configuration of carbon (1) is that shown hereinbelow:




R₁, R₂, s, Q, Y and G are as defined in claim 1, and pharmaceutically acceptable salt or solvate thereof.

9. A compound according to claim 1 wherein

Y is a divalent groups of formula

5



Y2

A₂ is absent and A₁ is independently selected from (C₁-C₁₂)alkylene which is methylene, ethylene, n-propylene, isopropylene, butylene, pentylene, hexylene, octylene;

10 B is absent or is selected from (C₃-C₈)heterocycloalkylene which is piperidinylene, arylene which is phenylene, and heteroarylene which is pyridine-diyl, or pyrazole-diyl; B being optionally substituted by one or more groups selected from -OH, halogen which is fluorine, chlorine, bromine, -CN, (C₁-C₆)alkyl which is methyl, (C₁-C₆)alkoxy which is methoxy, ethoxy, isopropoxy, (C₁-C₆)haloalkyl which is trifluoromethyl and
15 (C₁-C₆)haloalkoxy which is trifluoromethoxy;

C is absent or is selected from -O-, -C(O)-, or is one of the following groups C1, C2, C4, C7, C8, C9, C10, C12, C13, C14, C15, C16, C17, C18, C19, C20, C21 C22 C23

wherein R₇ is in each occurrence independently H or selected from (C₁-C₈)alkyl which is methyl, ethyl and aryl(C₁-C₆)alkyl which is benzyl;

20 D is absent or is selected from arylene which is para-phenylene or metha-phenylene, (C₃-C₈)cycloalkylene which is cyclohexanediyl, (C₃-C₈)heterocycloalkylene which is piperidindiyl, pyrrolidindiyl or azetidindiyl;

n is at each occurrence independently 0 or an integer from 1 to 3;

m is at each occurrence independently an integer from 1 to 3;

25 E is absent or is selected from -O-, -NR₇- which is -NH-, -NR₇-C(O)- which is -NH-C(O)-, -C(O)-NR₇- which is -C(O)-NH-, -C(O)-(CH₂)_n-O- which is -C(O)-CH₂-O- and -NR₇-C(O)-(CH₂)_n-O- which is -NH-C(O)-CH₂-O-;

G is arylene which is metha-phenylene or para-phenylene;

R₁ is selected from thiophenyl, cyclohexyl, cyclopentyl and phenyl optionally substituted by one or more group selected independently from fluorine, methyl, ethyl and methoxy;

s is 0, 1 or 2

5 **R₂** is a nitrogen containing group which may be selected from:

- a group (a) which is -NR₃R₄ wherein R₃ and R₄ are methyl; and

- a group (b) of formula J1, J2, J3, J4 or J5 wherein

R₅ is methyl or benzyl

and pharmaceutically acceptable salts or solvates thereof.

10 10. A compound according to claim 1 which is selected in the list consisting of:

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(p-tolyl)acetate;

15 (1-benzylpiperidin-4-yl)methyl 2-(4-fluorophenyl)-2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate;

(1-benzylpiperidin-4-yl)methyl 2-(3-fluorophenyl)-2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

20 yl)ethyl)amino)pentyl)oxy)phenyl)acetate;

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(m-tolyl)acetate;

(1-benzylpiperidin-4-yl)methyl 2-(2-chlorophenyl)-2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

25 yl)ethyl)amino)pentyl)oxy)phenyl)acetate;

(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(o-tolyl)acetate;

(1-benzylpiperidin-4-yl)methyl 2-(2-ethylphenyl)-2-hydroxy-2-(3-(((5-(((R)-2-

- hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate;
(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(thiophen-2-yl)acetate;
5 (1-benzylpiperidin-4-yl)methyl 2-cyclohexyl-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate;
(1-benzylpiperidin-4-yl)methyl 2-(3-ethylphenyl)-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate;
10 (R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-(4-methoxyphenyl)acetate;
15 (R)-quinuclidin-3-yl 2-cyclopentyl-2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)acetate;
(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
20 (R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
25 (R)-(R)-quinuclidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
(S)-2-(dimethylamino)ethyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;

- (R)-2-(dimethylamino)ethyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- (S)-(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- 5 (R)-(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- (S)-1-methylpiperidin-4-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- (R)-1-methylpiperidin-4-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- 10 (S)-(1-methylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- (R)-(1-methylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- 15 (S)-(R)-1-methylpiperidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- (S)-1-methylazetidin-3-yl 2-hydroxy-2-(3-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
- 20 (R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((7-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)heptyl)oxy)phenyl)-2-phenylacetate;
- 25 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)hexyl)oxy)phenyl)-2-phenylacetate;

phenylacetate;

(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((8-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)octyl)oxy)phenyl)-2-phenylacetate;

5 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;

(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;

10 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;

(R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethoxy)phenyl)-2-phenylacetate;

15 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;

20 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)propoxy)phenyl)-2-phenylacetate;

25 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)phenyl)-2-phenylacetate;
- 5 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-benzoyl)azetidin-3-yl)methoxy)phenyl)-2-phenylacetate;
- 10 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)azetidin-3-yl)methoxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
- 15 (R)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-((3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-((3-(benzyl(3-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)propyl)carbamoyl)benzyl)oxy)phenyl)-2-hydroxy-2-phenylacetate;
- 20 (R)-quinuclidin-3-yl 2-hydroxy-2-(3-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)benzamido)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate;
- (R)-quinuclidin-3-yl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-

- dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate;
(R)-1-methylpyrrolidin-3-yl 2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)butyl)carbamoyl)phenyl)-2-phenylacetate;
5 (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)carbamoyl)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
10 (R)-quinuclidin-3-yl 2-hydroxy-2-(4-((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;
15 (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(((5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentyl)oxy)carbonyl)amino)phenyl)-2-phenylacetate;
(R)-quinuclidin-3-yl 2-hydroxy-2-(3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)pentanamido)phenyl)-2-phenylacetate;
20 (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propyl)ureido)phenyl)-2-phenylacetate;
(S)- (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)propoxy)phenyl)-2-phenylacetate;
25 (S)- (1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)propoxy)phenyl)-2-phenylacetate;

- (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)nicotinamido)propoxy)phenyl)-2-phenylacetate;
- 5 (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(3-ethoxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 10 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethoxy)benzamido)propoxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 15 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-isopropoxybenzamido)propoxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 20 (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 25 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

- yl)ethyl)amino)methyl)picolinamido)propoxy)phenyl)-2-phenylacetate;
(S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(3-(2,3-difluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-hydroxy-2-phenylacetate;
5 (S)-(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethyl)benzamido)propoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(4-(2-(((R)-2-hydroxy-2-(4-hydroxy-6-oxo-5,6-dihydronaphthalen-1-yl)ethyl)amino)ethyl)piperidine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
(S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxyphenyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-((2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxyphenyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;
25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-

- (8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperazin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)(methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-4-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aR,6aR)-5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-1(2H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aS,6aS)-5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-1(2H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((1-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(ethyl(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)(methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)propyl)amino)-2-oxoethoxy)phenyl)-2-

phenylacetate;

(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperazin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

5 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(benzyl(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;

10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;

(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)methyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;

20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;

(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-

- diazaspiro[3.3]heptan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(7-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonan-7-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-4-yl)methyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.5]nonan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,9-diazaspiro[5.5]undecan-2-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-1-oxa-4,9-diazaspiro[5.5]undecan-4-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)azetidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)ethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((R)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)pyrrolidin-3-yl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)pyrrolidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aR,6aR)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((3aS,6aS)-1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((R)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzamido)cyclobutyl)carbamoxy)benzyl)oxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-3,9-diazaspiro[5.5]undecane-3-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.3]heptane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(7-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-

- diazaspiro[3.5]nonane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(4-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperazine-1-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(2-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,7-diazaspiro[3.5]nonane-7-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(6-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,6-diazaspiro[3.5]nonane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-1-oxa-4,9-diazaspiro[5.5]undecane-4-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(9-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)-2,9-diazaspiro[5.5]undecane-2-carbonyl)benzyl)oxy)phenyl)-2-phenylacetate;
15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)ethyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
25

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)ethyl)carbamoyl)benzyl)oxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(3-hydroxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 10 (S)-(1-benzylpiperidin-4-yl)methyl 2-(3-(2-(((1R,3S)-3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methylbenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (S)-(1-benzylpiperidin-4-yl)methyl 2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 25

- (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2,3-difluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-bromo-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-chloro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethyl)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(3-ethoxy-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;

- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-(trifluoromethoxy)benzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 5 (1-benzylpiperidin-4-yl)methyl (S)-2-(3-(2-(((1R,3S)-3-(2-fluoro-4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-5-methoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-hydroxy-2-phenylacetate;
- 10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-isopropoxybenzamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(5-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)picolinamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(6-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)nicotinamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,3S)-3-(2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)phenyl)acetamido)cyclobutyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)azetidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
- 25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-

- carbonyl)piperidin-1-yl)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-(((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)cyclohexyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
5 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(2-((2-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidin-1-yl)-2-oxoethyl)amino)-2-oxoethoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-(1-(4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)piperidine-4-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
10 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(2-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)ethyl)benzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
15 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxybenzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
20 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
(1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxybenzoyl)glycyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
25 (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-(5-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-

- 1-carbonyl)thiophene-2-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
- (1-benzylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-((1-((1R,4S)-4-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)piperidine-1-carbonyl)cyclohexane-1-carbonyl)piperidin-4-yl)methoxy)phenyl)-2-phenylacetate;
- 5 1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- 10 ((R)-1-benzylpyrrolidin-3-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- ((S)-1-benzylpyrrolidin-3-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- 15 (1-cyclobutylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- (1-methylpiperidin-4-yl)methyl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- 20 (R)-1-benzylpyrrolidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- (S)-1-benzylpyrrolidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;
- 25 1-benzylazetidin-3-yl (S)-2-hydroxy-2-(3-(3-(4-(((R)-2-hydroxy-2-(8-hydroxy-

2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)benzamido)propoxy)phenyl)-2-phenylacetate;

1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-

5 yl)ethyl)amino)methyl)phenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;

1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-3-methoxyphenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;

1-benzylpiperidin-4-yl (S)-2-hydroxy-2-(3-((4-((3-(4-(((R)-2-hydroxy-2-(8-
10 hydroxy-2-oxo-1,2-dihydroquinolin-5-yl)ethyl)amino)methyl)-2-methoxyphenyl)ureido)methyl)benzyl)oxy)phenyl)-2-phenylacetate;

and pharmaceutically acceptable salt and solvate thereof.

11. A pharmaceutical composition comprising a compound of formula I as defined in
15 any one of the claims 1-9, with one or more pharmaceutically acceptable carriers and/or excipients.

12. A pharmaceutical composition according to claim 11 to be administered by
inhalation, such as inhalable powders, propellant-containing metering aerosols or
20 propellant-free inhalable formulations.

13. A compound of formula I as defined in claims 1-10 for use in the prevention
and/or treatment of broncho-obstructive or inflammatory diseases, preferably asthma or
chronic bronchitis or chronic obstructive pulmonary disease (COPD).

25

14. A combination of a compound of formula I as defined in any one of the
claims 1-10 with one or more active ingredients selected from the classes
consisting of corticosteroids, P38 MAP kinase inhibitors, IKK2 inhibitors, HNE
inhibitors, PDE4 inhibitors, leukotriene modulators, NSAIDs and mucus
30 regulators.

15. A device comprising the pharmaceutical composition according to claim 14,

which may be a single- or multi-dose dry powder inhaler, a metered dose inhaler and a soft mist nebulizer.

16. Use of a compound of formula **I** as defined in claims 1-10 in the manufacture of
5 a medicament for the prevention and/or treatment of broncho-obstructive or inflammatory diseases, preferably asthma or chronic bronchitis or chronic obstructive pulmonary disease (COPD).