

This invention relates to nonsteroidal progesterone receptor modulators, a process for their production, the use of progesterone receptor modulators for the production of pharmaceutical agents as well as pharmaceutical compositions that contain these compounds.

The steroid hormone progesterone regulates the reproductive process in the female organism in a decisive way. During the cycle and in pregnancy, progesterone is secreted in large amounts from the ovary or the placenta. By interaction with estrogens, progesterone produces cyclic changes of the uterine mucous membrane (endometrium) in the menstrual cycle. Under the influence of elevated progesterone levels after ovulation, the uterine mucous membrane is converted into a state that allows the nidation of an embryo (blastocyte). In pregnancy, progesterone controls the relaxation of the myometrium and retains the function of the decidual tissue.

In addition, it is known that progesterone inhibits the endometrial proliferation by the suppression of the estrogen-mediated mitosis in the uterus tissue (K. Chwalisz, R. M. Brenner, U. Fuhrmann, H. Hess-Stumpp, W. Elger, Steroids 65, 2000, 741-751).

An important role of the progesterone and the progesterone receptors is also known in pathophysiological processes. Progesterone receptors are detected in foci of endometriosis, but also in tumors of the uterus, the breast and the CNS. In addition, it is known that uterus leiomyomas grow in a progesterone-dependent manner.

The actions of progesterone in the tissues of genital organs and in other tissues are carried out by interactions with progesterone receptors, which are responsible for the cellular effects.

Progesterone receptor modulators are either pure agonists or partially or completely inhibit the action of progesterone. Consequently, substances are defined as pure agonists, partial agonists (SPRMS) and pure antagonists.

According to the ability of the progesterone receptor modulators to influence the action of the progesterone receptor, these compounds have a considerable potential as therapeutic agents for gynecological and oncological indications as well as for obstetrics and birth control.

Pure progesterone receptor antagonists completely inhibit the action of progesterone in the progesterone receptor. They have antioviulatory properties as well as the ability to inhibit estrogen effects in the endometrium up to full atrophy. They are therefore especially suitable for intervening in the female reproductive process, e.g., in post-ovulation, to prevent nidation; in pregnancy, to increase the reactivity of the uterus to prostaglandins or oxytocin or to ensure the opening and softening ("maturation") of the cervix as well as to make the myometrium highly prepared for labor.

In foci of endometriosis or in tumor tissue, which are (is) equipped with progesterone receptors, an advantageous influence of the disease process is expected after application of pure progesterone receptor antagonists. Special advantages for influencing pathologic conditions, such as endometriosis or uterus leiomyomas, could then be given if in addition an inhibition of the ovulation can be achieved by the progesterone receptor antagonists. When ovulation is inhibited, a portion of the ovarian hormone production and thus the stimulative effect that is due to this portion are also due to the pathologically altered tissue.

A large number of analogs with varying degrees of progesterone receptor-antagonistic activity followed the first described progesterone receptor antagonist RU 486 (also mifepristone). While RU 486, in addition to the progesterone receptor-

antagonistic action, also shows an antiglucocorticoidal action, compounds synthesized later are distinguished primarily by a more selective action than progesterone receptor antagonists.

From the literature, in addition to steroidal compounds such as onapristone or lilopristone, which are distinguished from progesterone-receptor-antagonistic action to antiglucocorticoidal action relative to RU 486 by a better dissociation of action, various nonsteroidal structures, whose antagonistic action on the progesterone receptor is examined, are also known [see, e.g., S. A. Leonhardt and D. P. Edwards, *Exp. Biol. Med.* 227: 969-980 (2002) and R. Winneker, A. Fensome, J. E. Wrobel, Z. Zhang, P. Zhang, *Seminars in Reproductive Medicine*, Volume 23: 46-57 (2005)]. Previously known compounds, however, have only moderately antagonistic activity compared to the known steroidal structures. The most effective nonsteroidal compounds are described as having *in vitro* activities of 10% of the activity of RU 486.

The antiglucocorticoidal activity is disadvantageous for a therapeutic application in which the inhibition of the progesterone receptors is a primary focus of therapy. An antiglucocorticoidal activity causes undesirable side effects in the case of therapeutically necessary dosages. This can prevent the application of a therapeutically useful dose or lead to termination of the treatment.


The partial or complete reduction of the antiglucocorticoidal properties is therefore an important requirement for the therapy with progesterone receptor antagonists, in particular for those indications that require a treatment lasting weeks or months.

In contrast to the pure antagonists, progesterone receptor partial agonists (SPRMs) show a residual agonistic property, which can be strongly pronounced to different degrees. This leads to the fact that these substances show potential agonistic

actions of the progesterone receptor in specific organ systems (D. DeManno, W. Elger, R. Garg, R. Lee, B. Schneider, H. Hess-Stumpp, G. Schuber, K. Chwalisz, *Steroids* 68, 2003, 1019-1032). Such an organ-specific and dissociated action can be of therapeutic use for the indications described.

It is therefore the object of this invention to make available additional nonsteroidal progesterone receptor modulators. These compounds are to have a reduced antiglucocorticoidal action and are therefore suitable for the therapy and prophylaxis of gynecological diseases such as endometriosis, leiomyomas of the uterus, dysfunctional bleeding and dysmenorrhea. In addition, the compounds according to the invention are to be suitable for the therapy and prophylaxis of hormone-dependent tumors, for example breast, endometrial, ovarian and prostate cancers. In addition, the compounds are to be suitable for use in female birth control and for female hormone replacement therapy.

The object is achieved according to this invention by the preparation of non-steroidal compounds of general formula I



in which

R^1 and R^2 , independently of one another, mean a hydrogen atom, a straight or nonstraight, branched or unbranched C_1 - C_5 -alkyl group, also together with the C atom of the chain forming a ring with a total of 3-7 members,

R^3 means a radical $C\equiv C-R^a$, whereby

R^a means a hydrogen or a C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl, or heterocycloalkyl that optionally is substituted in one or more places, in the same way or differently, with K, or an aryl or heteroaryl that optionally is substituted in one or more places, in the same way or differently, with L,

K is a cyano, halogen, hydroxy, nitro, $-C(O)R^b$, CO_2R^b , $-O-R^b$, $-S-R^b$, $SO_2NR^cR^d$, $-C(O)-NR^cR^d$, $-OC(O)-NR^cR^d$, or $-C=NOR^b-NR^cR^d$ or a C_3 - C_{10} -cycloalkyl that optionally is substituted in one or more places, in the same way or differently, with M, heterocycloalkyl, or aryl or heteroaryl that optionally is substituted in one or more places with L,

L means C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_1 - C_6 -perfluoroalkyl, C_1 - C_6 -perfluoroalkoxy, C_1 - C_6 -alkoxy- C_1 - C_6 -alkoxy, $(CH_2)_p$ - C_3 - C_{10} -cycloalkyl, $(CH_2)_p$ -heterocycloalkyl, $(CH_2)_pCN$, $(CH_2)_pHal$, $(CH_2)_pNO_2$, $(CH_2)_p$ - C_6 - C_{12} -aryl, $(CH_2)_p$ -heteroaryl, $-(CH_2)_pPO_3(R^b)_2$, $-(CH_2)_pNR^cR^d$, $-(CH_2)_pNR^cCOR^b$, $-(CH_2)_pNR^cCSR^b$, $-(CH_2)_pNR^cS(O)R^b$, $-(CH_2)_pNR^cS(O)_2R^b$, $-(CH_2)_pNR^cCONR^cR^d$, $-(CH_2)_pNR^cCOOR^b$, $-(CH_2)_pNR^cC(NH)NR^cR^d$, $-(CH_2)_pNR^cCSNR^cR^d$, $-(CH_2)_pNR^cS(O)NR^cR^d$, $-(CH_2)_pNR^cS(O)_2NR^cR^d$, $-(CH_2)_pCOR^b$, $-(CH_2)_pCSR^b$, $-(CH_2)_pS(O)R^b$, $-(CH_2)_pS(O)(NH)R^b$, $-(CH_2)_pS(O)_2R^b$,

$-(\text{CH}_2)_p\text{S}(\text{O})_2\text{NR}^c\text{R}^d$, $-(\text{CH}_2)_p\text{SO}_2\text{OR}^b$,
 $-(\text{CH}_2)_p\text{CO}_2\text{R}^b$, $-(\text{CH}_2)_p\text{CONR}^c\text{R}^d$,
 $-(\text{CH}_2)_p\text{CSNR}^c\text{R}^d$, $-(\text{CH}_2)_p\text{OR}^b$, $-(\text{CH}_2)_p\text{SR}^b$,
 $-(\text{CH}_2)_p\text{CR}^b(\text{OH})-\text{R}^e$, $-(\text{CH}_2)_p-\text{C}=\text{NOR}^b$,
 $-\text{O}-(\text{CH}_2)_n-\text{O}-$, $-\text{O}-(\text{CH}_2)_n-\text{CH}_2-$, $-\text{O}-\text{CH}=\text{CH}-$
 or $-(\text{CH}_2)_{n+2}-$, whereby $n = 1$ or 2 , and the terminal
 oxygen atoms and/or carbon atoms are linked to
 directly adjacent ring-carbon atoms,

M means C_1 - C_6 -alkyl or a group $-\text{COR}^b$, CO_2R^b , $-\text{OR}^b$, or
 $-\text{NR}^c\text{R}^d$, whereby

R^b means a hydrogen or a C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl,
 C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{12} -aryl or C_1 -
 C_3 -perfluoroalkyl, and

R^c and R^d , independently of one another, mean a hydrogen,
 C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -
 cycloalkyl, C_6 - C_{12} -aryl, $\text{C}(\text{O})\text{R}^b$ or a hydroxy
 group, whereby if

R^c is a hydroxy group, R^d can be only one hydrogen, a
 C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -
 cycloalkyl or C_6 - C_{12} -aryl and vice versa, and

R^e means a hydrogen, C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl,
 C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl or C_6 - C_{12} -aryl,
 and

p can be a number from 0- 6,

or

R^3 is a radical $C = C-R^gR^h$, whereby

R^g and R^h , independently of one another, are a hydrogen or a C_1 - C_8 -alkyl,

C_2 - C_8 -alkenyl or C_2 - C_8 -alkinyl that optionally is substituted in one or more places, in the same way or differently, with X, in which

X is a cyano, halogen, hydroxy, nitro, $-C(O)R^b$, CO_2R^b ,

$-O-R^b$, $-C(O)-NR^cR^d$, $-NR^cR^d$ with the meanings already

further mentioned above for R^b , R^c and R^d , and

R^{4a} and R^{4b} , independently of one another, mean a hydrogen atom, a C_1 - C_4 -alkyl,

a C_2 - C_4 -alkenyl or together with the ring-carbon atom forming a 3- to 6-membered ring,

A means a monocyclic or bicyclic, carbocyclic or heterocyclic aromatic ring, which optionally can be substituted in one or more places with C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_1 - C_6 -perfluoroalkyl, C_1 - C_6 -perfluoroalkoxy, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkoxy, $(CH_2)_p$ - C_3 - C_{10} -cycloalkyl, $(CH_2)_p$ -heterocycloalkyl, $(CH_2)_p$ CN, $(CH_2)_p$ Hal, $(CH_2)_p$ NO₂, $(CH_2)_p$ - C_6 - C_{12} -aryl, $(CH_2)_p$ -heteroaryl, $-(CH_2)_pPO_3(R^b)_2$, $-(CH_2)_pNR^cR^d$, $-(CH_2)_pNR^cCOR^b$, $-(CH_2)_pNR^cCSR^b$, $-(CH_2)_pNR^cS(O)R^b$, $-(CH_2)_pNR^cS(O)_2R^b$, $-(CH_2)_pNR^cCONR^cR^d$, $-(CH_2)_pNR^cCOOR^b$, $-(CH_2)_pNR^cC(NH)NR^cR^d$, $-(CH_2)_pNR^cCSNR^cR^d$, $-(CH_2)_pNR^cS(O)NR^cR^d$, $-(CH_2)_pNR^cS(O)_2NR^cR^d$, $-(CH_2)_pCOR^b$, $-(CH_2)_pCSR^b$, $-(CH_2)_pS(O)R^b$, $-(CH_2)_pS(O)(NH)R^b$, $-(CH_2)_pS(O)_2R^b$, $-(CH_2)_pS(O)_2NR^cR^d$, $-(CH_2)_pSO_2OR^b$, $-(CH_2)_pCO_2R^b$, $-(CH_2)_pCONR^cR^d$, $-(CH_2)_pCSNR^cR^d$, $-(CH_2)_pOR^b$, $-(CH_2)_pSR^b$, $-(CH_2)_pCR^b(OH)-R^d$, $-(CH_2)_pC=NOR^b$, $-O-(CH_2)_n-O-$, $-O-(CH_2)_n-CH_2-$, $-O-CH=CH-$ or

$-(\text{CH}_2)_{n+2}-$, whereby $n = 1$ or 2 , and the terminal oxygen atoms and/or carbon atoms are linked to directly adjacent ring-carbon atoms, or

A means a radical $-\text{CO}_2\text{R}^b$, $\text{C}(\text{O})\text{NR}^c\text{R}^d$, COR^b ,

or

A means an alkenyl group $-\text{CR}^5 = \text{CR}^6\text{R}^7$, whereby

R^5 , R^6 and R^7 are the same or different and, independently of one

another, mean hydrogen atoms, halogen atoms, aryl radicals or an unsubstituted or partially or completely fluorinated C_1 - C_5 -alkyl group, or

A means an alkynyl group $-\text{C}\equiv\text{CR}^5$, with the meaning cited above for R^5 , and

B means a carbonyl group or a CH_2 group

as well as their pharmaceutically acceptable salts.

The compounds of general formula I according to the invention can be present as different stereoisomers because of the presence of asymmetry centers. Both the racemates and the separately present stereoisomers are part of the subject of this invention.


In addition, this invention comprises the new compounds as pharmaceutical active ingredients, their production, their therapeutic application and pharmaceutical dispensing forms that contain the new substances.

The compounds of general formula (I) according to the invention or their pharmaceutically acceptable salts can be used for the production of a pharmaceutical agent, especially for treatment and prophylaxis of gynecological diseases, such as endometriosis, leiomyomas of the uterus, dysfunctional bleeding and dysmenorrhea. In addition, the compounds according to the invention can be used for the treatment and

prophylaxis of hormone-dependent tumors, such as, for example, for breast, prostate and endometrial cancers.

The compounds of general formula (I) according to the invention or their pharmaceutically acceptable salts are suitable for use for female birth control or for female hormone replacement therapy.

A process for the production of the compounds of general formula (I), moreover, is also a subject of this invention. Substituent R^3 is introduced to a keto group by selective addition reaction of organometallic compounds such as lithium alkynylene or magnesium haloalkynylene. This results, either directly or after implementing additional modifications, in the compounds of general formula (I) according to the invention.



The production of the compounds according to the invention is carried out by selective addition of organometallic compounds to ketoamides, which were described in, e.g., laid-open specifications WO 200375915 and WO 9854159. The organometallic compounds can be, for example, lithium alkynyl compounds or magnesium haloalkynyl compounds. The latter are produced by, e.g., reaction of the corresponding alkynes with butyllithium or Grignard compounds. Analogously to this, the corresponding

organometallic alkenyl compounds can also be produced. The reactivity of the keto group in comparison to amidocarbonyl or to phthalide is in this case significantly higher, such that with suitable selection of the reaction conditions, a selective addition is achieved. As an alternative, the alkynyl or alkenyl radicals that are introduced as R^3 can also be further modified later. For these modifications, reactions that have become known to one skilled in the art, such as oxidation, reduction, substitution, alkylation, or palladium-catalyzed reaction, are suitable. Optionally present protective groups are cleaved off at a suitable time.

The nonsteroidal compounds of general formula I according to the invention have a strongly antagonistic or strongly partially agonistic action on the progesterone receptor. They exhibit a strong dissociation of action with respect to their bonding strength on the progesterone receptor and on the glucocorticoid receptor. While known progesterone receptor antagonists, such as Mifepristone (RU 486), in addition to the desired high binding affinity for the progesterone receptor likewise show a high affinity for the glucocorticoid receptor, the compounds according to the invention are distinguished by a very low glucocorticoid receptor bond with simultaneously present high progesterone receptor affinity.

The substituents of the compounds of general formula I according to the invention that are defined as groups can have the meanings below in each case:

C_1-C_5 -, C_1-C_6 - or C_1-C_8 -alkyl groups are defined as straight or nonstraight, branched or unbranched alkyl radicals. In this case, for example, this is a methyl, ethyl, n-propyl, iso-propyl, n-, iso-, tert-butyl, an n-pentyl, 2,2-dimethylpropyl, 3-methylbutyl, hexyl, heptyl or octyl group.

In terms of R^a , in this case, the methyl, ethyl, n-propyl or n-butyl group as well as an n-pentyl group are preferred.

In terms of R^1 and R^2 , methyl or ethyl is preferred.

According to the invention, a hydrogen is preferred for $R4^a$ and $R4^b$.

Alkenyl is defined as straight or nonstraight, branched or unbranched alkenyl radicals. In terms of the invention, a C_2 – C_8 -alkenyl group is defined, for example, as follows: vinyl, allyl, 3-buten-1-yl- or 2,3-dimethyl-2-propenyl. If aromatic compound A is substituted with a C_2 – C_8 -alkenyl radical, this is preferably a vinyl group.

Alkynyl is defined as straight or nonstraight, branched or unbranched alkynyl radicals. For example, an ethynyl, propynyl, butynyl, pentynyl, hexynyl or octynyl group, but preferably an ethynyl or propynyl group, is to stand for a C_2 – C_8 -alkynyl radical.

For C_3 – C_{10} -cycloalkyl, for example, cyclopropane, cyclobutane, cyclopentane and cyclohexane can be mentioned. Cyclopropyl, cyclopentyl and cyclohexyl are preferred.

In terms of R^a , K or L, heterocycloalkyl is defined as 3- to 8-membered heterocycloalkyl radicals. Examples of heterocycloalkyl are morpholine, tetrahydrofuran, pyran, piperazine, piperidine, pyrrolidine, oxirane, oxetane, aziridine, dioxolane and dioxane. In this case, the position of the heteroatom in relation to the point of linkage can be any chemically possible position.

For example, methoxymethoxy, ethoxymethoxy or 2-methoxyethoxy can stand for a C_1 – C_6 -alkoxyl- C_1 – C_6 -alkoxy group.

In terms of the invention, a radical OR^b is a hydroxy, methoxy, ethoxy, n-propoxy, iso-propoxy, n-, iso-, or tert-butoxy group, or an n-pentoxo, 2,2-dimethylpropoxy or 3-methylbutoxy group. Hydroxy, methoxy and ethoxy are preferred.

For a partially or completely fluorinated C_1 – C_5 -alkyl group, the perfluorinated alkyl groups that appear above are considered. Of the latter, primarily the

trifluoromethyl group or the pentafluoroethyl group as well as as partially fluorinated alkyl groups, for example the 5,5,4,4-pentafluoropentyl group or the 5,5,5,4,4,3,3-heptafluoropentyl group, are preferred.

A fluorine, chlorine, bromine or iodine atom can stand for a halogen atom.

Preferred here is fluorine, chlorine or bromine.

If R^1 and R^2 together with the C atom of the chain form a 3- to 7-membered ring, this is, for example, a cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl ring. The cyclopropyl ring as well as the cyclopentyl ring are preferred.

The monocyclic or bicyclic carbocyclic aromatic ring A, which can be substituted in several places, is a carbocyclic or heterocyclic aryl radical.

In the first case, it is, for example, a phenyl or naphthyl radical, preferably a phenyl radical.

As a heterocyclic radical, for example, a monocyclic heterocyclic radical, for example the thienyl, furyl, pyranlyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, thiazolyl, oxazolyl, furazanyl, pyrrolinyl, imidazolinyl, pyrazolinyl, thiazolinyl, triazolyl, or tetrazolyl radical, and specifically all possible isomers relative to the positions of the heteroatoms, can be used.

In terms of R^3 , an aryl radical is an optionally substituted phenyl, 1- or 2-naphthyl radical, whereby the phenyl radical is preferred. Examples of a heteroaryl radical are the 2-, 3- or 4-pyridinyl radical, the 2- or 3-furyl radical, the 2- or 3-thienyl radical, the 2- or 3-pyrrolyl radical, the 2-, 4- or 5-imidazolyl radical, the pyrazinyl radical, the 2-, 4- or 5-pyrimidinyl radical or the 3- or 4-pyridazinyl radical.













The number p for the $(CH_2)_p$ radical can be a number from 0 to 6, preferably 0 to


2. "Radicals" are defined according to the invention as all functional groups that are presented in connection with $(CH_2)_p$.







In the case that the compounds of general formula I ($B = -CH_2-$) are present as salts, this can be, for example, in the form of hydrochloride, sulfate, nitrate, tartrate, citrate, fumarate, succinate or benzoate.








If the compounds according to the invention are present as racemic mixtures, they can be separated into the pure, optically active forms according to methods of racemate separation that are familiar to one skilled in the art. For example, the racemic mixtures can be separated into pure isomers by chromatography on an even optically active carrier material (CHIRALPAK AD[®]). It is also possible to esterify the free hydroxy group in a racemic compound of general formula I with an optically active acid and to separate the diastereomeric esters that are obtained by fractionated crystallization or by chromatography and to saponify the separated esters in each case to form the optically pure isomers. As an optically active acid, for example, mandelic acid, camphorsulfonic acid or tartaric acid can be used.


The compounds that are mentioned below as well as the use thereof are preferred according to the invention:

No.	Racemic or Enantiomer	R3
1	rac	
2	+	
3	-	
4	rac	
5	+	
6	-	
7	rac	
8	+	
9	-	
10	rac	
11	+	
12	-	
13	rac	
14	+	
15	-	
16	rac	
17	+	
18	-	
19	rac	
20	+	
21	-	
22	rac	
23	+	
24	-	
25	rac	
26	+	
27	-	
28	rac	
29	+	
30	-	
31	rac	
32	+	
33	-	
34	rac	
35	+	
36	-	
37	rac	
38	+	
39	-	
40	rac	
41	+	




42	-	
43	rac	
44	+	
45	-	
46	rac	
47	+	
48	-	
49	rac	
50	+	
51	-	
52	rac	
53	+	
54	-	
55	rac	
56	+	
57	-	
58	rac	
59	+	
60	-	















No.	Racemic or Enantiomer	R3
61	rac	
62	+	
63	-	
64	rac	
65	+	
66	-	
67	rac	
68	+	
69	-	
70	rac	
71	+	
72	-	
73	rac	
74	+	
75	-	
76	rac	
77	+	
78	-	
79	rac	
80	+	
81	-	
82	rac	
83	+	
84	-	


















85	rac	
86	+	
87	-	
88	rac	
89	+	
90	-	
91	rac	
92	+	
93	-	
94	rac	
95	+	
96	-	
97	rac	
98	+	
99	-	
100	rac	
101	+	
102	-	
103	rac	
104	+	
105	-	
106	rac	
107	+	
108	-	
109	rac	
110	+	
111	-	
112	rac	
113	+	
114	-	
115	rac	
116	+	
117	-	
118	rac	
119	+	
120	-	


No.	Racemic or Enantiomer	R3
121	rac	
122	+	
123	-	
124	rac	
125	+	
126	-	
127	rac	











128	+	
129	-	
130	rac	
131	+	
132	-	
133	rac	
134	+	
135	-	
136	rac	
137	+	
138	-	
139	rac	
140	+	
141	-	
142	rac	
143	+	
144	-	
145	rac	
146	+	
147	-	
148	rac	
149	+	
150	-	
151	rac	
152	+	
153	-	
154	rac	
155	+	
156	-	
157	rac	
158	+	
159	-	
160	rac	
161	+	
162	-	
163	rac	
164	+	
165	-	
166	rac	
167	+	
168	-	
169	rac	
170	+	
171	-	
172	rac	
173	+	
174	-	


175	rac	
176	+	
177	-	
178	rac	
179	+	
180	-	













No.	Racemic or Enantiomer	R3
181	rac	
182	+	
183	-	
184	rac	
185	+	
186	-	
187	rac	
188	+	
189	-	
190	rac	
191	+	
192	-	
193	rac	
194	+	
195	-	
196	rac	
197	+	
198	-	
199	rac	
200	+	
201	-	
202	rac	
203	+	
204	-	
205	rac	
206	+	
207	-	
208	rac	
209	+	
210	-	
211	rac	
212	+	
213	-	
214	rac	
215	+	
216	-	
217	rac	
218	+	





219	-	
220	rac	
221	+	
222	-	
223	rac	
224	+	
225	-	
226	rac	
227	+	
228	-	
229	rac	
230	+	
231	-	
232	rac	
233	+	
234	-	
235	rac	
236	+	
237	-	
238	rac	
239	+	
240	-	
















No.	Racemic or Enantiomer	R3
241	rac	
242	+	
243	-	
244	rac	
245	+	
246	-	
247	rac	
248	+	
249	-	
250	rac	
251	+	
252	-	
253	rac	
254	+	
255	-	
256	rac	
257	+	
258	-	
259	rac	
260	+	















261	-	
262	rac	
263	+	
264	-	
265	rac	
266	+	
267	-	
268	rac	
269	+	
270	-	
271	rac	
272	+	
273	-	
274	rac	
275	+	
276	-	
277	rac	
278	+	
279	-	
280	rac	
281	+	
282	-	
283	rac	
284	+	
285	-	
286	rac	
287	+	
288	-	
289	rac	
290	+	
291	-	
292	rac	
293	+	
294	-	
295	rac	
296	+	
297	-	
298	rac	
299	+	
300	-	


No.	Racemic or Enantiomer	R3
301	rac	
302	+	
303	-	

















304	rac	
305	+	
306	-	
307	rac	
308	+	
309	-	
310	rac	
311	+	
312	-	
313	rac	
314	+	
315	-	
316	rac	
317	+	
318	-	
319	rac	
320	+	
321	-	
322	rac	
323	+	
324	-	
325	rac	
326	+	
327	-	
328	rac	
329	+	
330	-	
331	rac	
332	+	
333	-	
334	rac	
335	+	
336	-	
337	rac	
338	+	
339	-	
340	rac	
341	+	
342	-	
343	rac	
344	+	
345	-	
346	rac	
347	+	
348	-	
349	rac	
350	+	


351	-	
352	rac	
353	+	
354	-	
355	rac	
356	+	
357	-	
358	rac	
359	+	
360	-	
















No.	Racemic or Enantiomer	R3
361	rac	
362	+	
363	-	
364	rac	
365	+	
366	-	
367	rac	
368	+	
369	-	
370	rac	
371	+	
372	-	
373	rac	
374	+	
375	-	
376	rac	
377	+	
378	-	
379	rac	
380	+	
381	-	
382	rac	
383	+	
384	-	
385	rac	
386	+	
387	-	
388	rac	
389	+	
390	-	














391	rac	
392	+	
393	-	
394	rac	
395	+	
396	-	
397	rac	
398	+	
399	-	
400	rac	
401	+	
402	-	
403	rac	
404	+	
405	-	
406	rac	
407	+	
408	-	
409	rac	
410	+	
411	-	
412	rac	
413	+	
414	-	
415	rac	
416	+	
417	-	
418	rac	
419	+	
420	-	





No.	Racemic or Enantiomer	R3
421	rac	
422	+	
423	-	
424	rac	
425	+	
426	-	
427	rac	
428	+	
429	-	
430	rac	
431	+	
432	-	
433	rac	













434	+	
435	-	
436	rac	
437	+	
438	-	
439	rac	
440	+	
441	-	
442	rac	
443	+	
444	-	
445	rac	
446	+	
447	-	
448	rac	
449	+	
450	-	
451	rac	
452	+	
453	-	
454	rac	
455	+	
456	-	
457	rac	
458	+	
459	-	
460	rac	
461	+	
462	-	
463	rac	
464	+	
465	-	
466	rac	
467	+	
468	-	
		
469	rac	
470	+	
471	-	
472	rac	
473	+	
474	-	
475	rac	
476	+	
477	-	
478	rac	
479	+	
480	-	











No.	Racemic or Enantiomer	R3
481	rac	
482	+	
483	-	
484	rac	
485	+	
486	-	
487	rac	
488	+	
489	-	
490	rac	
491	+	
492	-	
493	rac	
494	+	
495	-	
496	rac	
497	+	
498	-	
499	rac	
500	+	
501	-	
502	rac	
503	+	
504	-	
505	rac	
506	+	
507	-	
508	rac	
509	+	
510	-	
511	rac	
512	+	
513	-	
214	rac	
515	+	
516	-	
517	rac	
518	+	
519	-	
520	rac	
521	+	
522	-	
523	rac	
524	+	
525	-	
526	rac	






227	+	
528	-	
529	rac	
530	+	
531	-	
532	rac	
533	+	
534	-	
535	rac	
536	+	
537	-	
538	rac	
539	+	
540	-	

















No.	Racemic or Enantiomer	R3
541	rac	
542	+	
543	-	
544	rac	
545	+	
546	-	
547	rac	
548	+	
549	-	
550	rac	
551	+	
552	-	
553	rac	
554	+	
555	-	
556	rac	
557	+	
558	-	
559	rac	
560	+	
561	-	
562	rac	
563	+	
564	-	
565	rac	
566	+	
567	-	
568	rac	
569	+	
570	-	


















571	rac	
572	+	
573	-	
574	rac	
575	+	
576	-	
577	rac	
578	+	
579	-	
580	rac	
581	+	
582	-	
583	rac	
584	+	
585	-	
586	rac	
587	+	
588	-	
589	rac	
590	+	
591	-	
592	rac	
593	+	
594	-	
595	rac	
596	+	
597	-	
598	rac	
599	+	
600	-	

No.	Racemic or Enantiomer	R3
601	rac	
602	+	
603	-	
604	rac	
605	+	
606	-	
607	rac	
608	+	
609	-	
610	rac	
611	+	
612	-	




613	rac	
614	+	
615	-	
616	rac	
617	+	
618	-	
619	rac	
620	+	
621	-	
622	rac	
623	+	
624	-	
625	rac	
626	+	
627	-	
628	rac	
629	+	
630	-	
631	rac	
632	+	
633	-	
634	rac	
635	+	
636	-	
637	rac	
638	+	
639	-	
640	rac	
641	+	
642	-	
643	rac	
644	+	
645	-	
646	rac	
647	+	
648	-	
649	rac	
650	+	
651	-	
652	rac	
653	+	
654	-	
655	rac	
656	+	
657	-	
658	rac	
659	+	
660	-	












No.	Racemic or Enantiomer	R3
661	rac	
662	+	
663	-	
664	rac	
665	+	
666	-	
667	rac	
668	+	
669	-	
670	rac	
671	+	
672	-	
673	rac	
674	+	
675	-	
676	rac	
677	+	
678	-	
679	rac	
680	+	
681	-	
682	rac	
683	+	
684	-	
685	rac	
686	+	
687	-	
688	rac	
689	+	
690	-	
691	rac	
692	+	
693	-	
694	rac	
695	+	
696	-	
697	rac	
698	+	
699	-	
700	rac	
701	+	
702	-	
703	rac	
704	+	
705	-	







706	rac	
707	+	
708	-	
709	rac	
710	+	
711	-	
712	rac	
713	+	
714	-	
715	rac	
716	+	
717	-	
718	rac	
719	+	
720	-	
















No.	Racemic or Enantiomer	R3
721	rac	
722	+	
723	-	
724	rac	
725	+	
726	-	
727	rac	
728	+	
729	-	
730	rac	
731	+	
732	-	
733	rac	
734	+	
735	-	
736	rac	
737	+	
738	-	
739	rac	
740	+	
741	-	
742	rac	
743	+	
744	-	
745	rac	
746	+	














747	-	
748	rac	
749	+	
750	-	
751	rac	
752	+	
753	-	
754	rac	
755	+	
756	-	
757	rac	
758	+	
759	-	
760	rac	
761	+	
762	-	
763	rac	
764	+	
765	-	
766	rac	
767	+	
768	-	
769	rac	
770	+	
771	-	
772	rac	
773	+	
774	-	
775	rac	
776	+	
777	-	
778	rac	
779	+	
780	-	


No.	Racemic or Enantiomer	R3
781	rac	
782	+	
783	-	
784	rac	
785	+	
786	-	
787	rac	
788	+	















789	-	
790	rac	
791	+	
792	-	
793	rac	
794	+	
795	-	
796	rac	
797	+	
798	-	
799	rac	
800	+	
801	-	
802	rac	
803	+	
804	-	
805	rac	
806	+	
807	-	
808	rac	
809	+	
810	-	
811	rac	
812	+	
813	-	
814	rac	
815	+	
816	-	
817	rac	
818	+	
819	-	
820	rac	
821	+	
822	-	
823	rac	
824	+	
825	-	
826	rac	
827	+	
828	-	
829	rac	
830	+	
831	-	
832	rac	
833	+	
834	-	
835	rac	
836	+	
837	-	


838	rac	
839	+	
840	-	












No.	Racemic or Enantiomer	R3
841	rac	
842	+	
843	-	
844	rac	
845	+	
846	-	
847	rac	
848	+	
849	-	
850	rac	
851	+	
852	-	
853	rac	
854	+	
855	-	
856	rac	
857	+	
858	-	
859	rac	
860	+	
861	-	
862	rac	
863	+	
864	-	
865	rac	
866	+	
867	-	
868	rac	
869	+	
870	-	
871	rac	
872	+	
873	-	
874	rac	
875	+	
876	-	
877	rac	
878	+	
879	-	





880	rac	
881	+	
882	-	
883	rac	
884	+	
885	-	
886	rac	
887	+	
888	-	
889	rac	
890	+	
891	-	
892	rac	
893	+	
894	-	
895	rac	
896	+	
897	-	
898	rac	
899	+	
900	-	
















No.	Racemic or Enantiomer	R3
901	rac	
902	+	
903	-	
904	rac	
905	+	
906	-	
907	rac	
908	+	
909	-	
910	rac	
911	+	
912	-	
913	rac	
914	+	
915	-	
916	rac	
917	+	
918	-	
919	rac	
920	+	
921	-	














922	rac	
923	+	
924	-	
925	rac	
926	+	
927	-	
928	rac	
929	+	
930	-	
931	rac	
932	+	
933	-	
934	rac	
935	+	
936	-	
937	rac	
938	+	
939	-	
940	rac	
941	+	
942	-	
943	rac	
944	+	
945	-	
946	rac	
947	+	
948	-	
949	rac	
950	+	
951	-	
952	rac	
953	+	
954	-	
955	rac	
956	+	
957	-	
958	rac	
959	+	
960	-	



No.	Racemic or Enantiomer	R3
961	rac	
962	+	
















963	-	
964	rac	
965	+	
966	-	
967	rac	
968	+	
969	-	
970	rac	
971	+	
972	-	
973	rac	
974	+	
975	-	
976	rac	
977	+	
978	-	
979	rac	
980	+	
981	-	
982	rac	
983	+	
984	-	
		
985	rac	
986	+	
987	-	
988	rac	
989	+	
990	-	
991	rac	
992	+	
993	-	
994	rac	
995	+	
996	-	
997	rac	
998	+	
999	-	
1000	rac	
1001	+	
1002	-	
1003	rac	
1004	+	
1005	-	
1006	rac	
1007	+	
1008	-	
1009	rac	


1010	+	
1011	-	
1012	rac	
1013	+	
1014	-	
1015	rac	
1016	+	
1017	-	
1018	rac	
1019	+	
1020	-	

No.	Racemic or Enantiomer	R3
1021	rac	
1022	+	
1023	-	
1024	rac	
1025	+	
1026	-	
1027	rac	
1028	+	
1029	-	
1030	rac	
1031	+	
1032	-	
1033	rac	
1034	+	
1035	-	
1036	rac	
1037	+	
1038	-	
1039	rac	
1040	+	
1041	-	
1042	rac	
1043	+	
1044	-	
1045	rac	
1046	+	
1047	-	
1048	rac	
1049	+	
1050	-	
1051	rac	




1052	+	
1053	-	
1054	rac	
1055	+	
1056	-	
1057	rac	
1058	+	
1059	-	
1060	rac	
1061	+	
1062	-	
1063	rac	
1064	+	
1065	-	
1066	rac	
1067	+	
1068	-	
1069	rac	
1070	+	
1071	-	
1072	rac	
1073	+	
1074	-	
1075	rac	
1076	+	
1077	-	
1078	rac	
1079	+	
1080	-	





No.	Racemic or Enantiomer	R3
1081	rac	
1082	+	
1083	-	
1084	rac	
1085	+	
1086	-	
1087	rac	
1088	+	
1089	-	
1090	rac	
1091	+	
1092	-	
1093	rac	
1094	+	











1095	-	
1096	rac	
1097	+	
1098	-	
1099	rac	
1100	+	
1101	-	
1102	rac	
1103	+	
1104	-	
1105	rac	
1106	+	
1107	-	
1108	rac	
1109	+	
1110	-	
1111	rac	
1112	+	
1113	-	
1114	rac	
1115	+	
1116	-	
1117	rac	
1118	+	
1119	-	
1120	rac	
1121	+	
1122	-	
1123	rac	
1124	+	
1125	-	
1126	rac	
1127	+	
1128	-	
1129	rac	
1130	+	
1131	-	
1132	rac	
1133	+	
1134	-	
1135	rac	
1136	+	
1137	-	
1138	rac	
1139	+	
1140	-	











No.	Racemic or Enantiomer	R3
1141	rac	
1142	+	
1143	-	
1144	rac	
1145	+	
1146	-	
1147	rac	
1148	+	
1149	-	
1150	rac	
1151	+	
1152	-	
1153	rac	
1154	+	
1155	-	
1156	rac	
1157	+	
1158	-	
1159	rac	
1160	+	
1161	-	
1162	rac	
1163	+	
1164	-	
1165	rac	
1166	+	
1167	-	
1168	rac	
1169	+	
1170	-	
1171	rac	
1172	+	
1173	-	
1174	rac	
1175	+	
1176	-	
1177	rac	
1178	+	
1179	-	
1180	rac	
1181	+	
1182	-	
1183	rac	
1184	+	
1185	-	
1186	rac	





1187	+	
1188	-	
1189	rac	
1190	+	
1191	-	
1192	rac	
1193	+	
1194	-	
1195	rac	
1196	+	
1197	-	
1198	rac	
1199	+	
1200	-	

















No.	Racemic or Enantiomer	R3
1201	rac	
1202	+	
1203	-	
1204	rac	
1205	+	
1206	-	
1207	rac	
1208	+	
1209	-	
1210	rac	
1211	+	
1212	-	
1213	rac	
1214	+	
1215	-	
1216	rac	
1217	+	
1218	-	
1219	rac	
1220	+	
1221	-	
1222	rac	
1223	+	
1224	-	
1225	rac	
1226	+	
1227	-	















1228	rac	
1229	+	
1230	-	
1231	rac	
1232	+	
1233	-	
1234	rac	
1235	+	
1236	-	
1237	rac	
1238	+	
1239	-	
1240	rac	
1241	+	
1242	-	
1243	rac	
1244	+	
1245	-	
1246	rac	
1247	+	
1248	-	
1249	rac	
1250	+	
1251	-	
1252	rac	
1253	+	
1254	-	
1255	rac	
1256	+	
1257	-	
1258	rac	
1259	+	
1260	-	


No.	Racemic or Enantiomer	R3
1261	rac	
1262	+	
1263	-	
1264	rac	
1265	+	
1266	-	
1267	rac	
1268	+	














1269	-	
1270	rac	
1271	+	
1272	-	
1273	rac	
1274	+	
1275	-	
1276	rac	
1277	+	
1278	-	
1279	rac	
1280	+	
1281	-	
1282	rac	
1283	+	
1284	-	
1285	rac	
1286	+	
1287	-	
1288	rac	
1289	+	
1290	-	
1291	rac	
1292	+	
1293	-	
1294	rac	
1295	+	
1296	-	
1297	rac	
1298	+	
1299	-	
1300	rac	
1301	+	
1302	-	
1303	rac	
1304	+	
1305	-	
1306	rac	
1307	+	
1308	-	
1309	rac	
1310	+	
1311	-	
1312	rac	
1313	+	
1314	-	
1315	rac	
1316	+	
1317	-	


1318	rac	
1319	+	
1320	-	















No.	Racemic or Enantiomer	R3
1321	rac	
1322	+	
1323	-	
1324	rac	
1325	+	
1326	-	
1327	rac	
1328	+	
1329	-	
1330	rac	
1331	+	
1332	-	
1333	rac	
1334	+	
1335	-	
1336	rac	
1337	+	
1338	-	
1339	rac	
1340	+	
1341	-	
1342	rac	
1343	+	
1344	-	
1345	rac	
1346	+	
1347	-	
1348	rac	
1349	+	
1350	-	
1351	rac	
1352	+	
1353	-	
1354	rac	
1355	+	
1356	-	
1357	rac	
1358	+	
















1359	-	
1360	rac	
1361	+	
1362	-	
1363	rac	
1364	+	
1365	-	
1366	rac	
1367	+	
1368	-	
1369	rac	
1370	+	
1371	-	
1372	rac	
1373	+	
1374	-	
1375	rac	
1376	+	
1377	-	
1378	rac	
1379	+	
1380	-	

No.	Racemic or Enantiomer	R3
1381	rac	
1382	+	
1383	-	
1384	rac	
1385	+	
1386	-	
1387	rac	
1388	+	
1389	-	
1390	rac	
1391	+	
1392	-	
1393	rac	
1394	+	
1395	-	
1396	rac	
1397	+	




1398	-	
1399	rac	
1400	+	
1401	-	
1402	rac	
1403	+	
1404	-	
1405	rac	
1406	+	
1407	-	
1408	rac	
1409	+	
1410	-	
1411	rac	
1412	+	
1413	-	
1414	rac	
1415	+	
1416	-	
1417	rac	
1418	+	
1419	-	
1420	rac	
1421	+	
1422	-	
1423	rac	
1424	+	
1425	-	
1426	rac	
1427	+	
1428	-	
1429	rac	
1430	+	
1431	-	
1432	rac	
1433	+	
1434	-	
1435	rac	
1436	+	
1437	-	
1438	rac	
1439	+	
1440	-	











No.	Racemic or Enantiomer	R3
1441	rac	
1442	+	
1443	-	
1444	rac	
1445	+	
1446	-	
1447	rac	
1448	+	
1449	-	
1450	rac	
1451	+	
1452	-	
1453	rac	
1454	+	
1455	-	
1456	rac	
1457	+	
1458	-	
1459	rac	
1460	+	
1461	-	
1462	rac	
1463	+	
1464	-	
1465	rac	
1466	+	
1467	-	
1468	rac	
1469	+	
1470	-	
1471	rac	
1472	+	
1473	-	
1474	rac	
1475	+	
1476	-	
1477	rac	
1478	+	
1479	-	
1480	rac	
1481	+	
1482	-	
1483	rac	
1484	+	







1485	-	
1486	rac	
1487	+	
1488	-	
1489	rac	
1490	+	
1491	-	
1492	rac	
1493	+	
1494	-	
1495	rac	
1496	+	
1497	-	
1498	rac	
1499	+	
1500	-	















No.	Racemic or Enantiomer	R3
1501	rac	
1502	+	
1503	-	
1504	rac	
1505	+	
1506	-	
1507	rac	
1508	+	
1509	-	
1510	rac	
1511	+	
1512	-	
1513	rac	
1514	+	
1515	-	
1516	rac	
1517	+	
1518	-	
1519	rac	
1520	+	
1521	-	
1522	rac	
1523	+	
1524	-	
1525	rac	
1526	+	
1527	-	
1528	rac	




1529	+	
1530	-	
1531	rac	
1532	+	
1533	-	
1534	rac	
1535	+	
1536	-	
1537	rac	
1538	+	
1539	-	
1540	rac	
1541	+	
1542	-	
1543	rac	
1544	+	
1545	-	
1546	rac	
1547	+	
1548	-	
		
1549	rac	
1550	+	
1551	-	
1552	rac	
1553	+	
1554	-	
1555	rac	
1556	+	
1557	-	
1558	rac	
1559	+	
1560	-	

No.	Racemic or Enantiomer	R3
1561	rac	
1562	+	
1563	-	
1564	rac	
1565	+	
1566	-	
1567	rac	
1568	+	
1569	-	
1570	rac	



1571	+	
1572	-	
1573	rac	
1574	+	
1575	-	
1576	rac	
1577	+	
1578	-	
1579	rac	
1580	+	
1581	-	
1582	rac	
1583	+	
1584	-	
1585	rac	
1586	+	
1587	-	
1588	rac	
1589	+	
1590	-	
1591	rac	
1592	+	
1593	-	
1594	rac	
1595	+	
1596	-	
1597	rac	
1598	+	
1599	-	
1600	rac	
1601	+	
1602	-	
1603	rac	
1604	+	
1605	-	
1606	rac	
1607	+	
1608	-	
1609	rac	
1610	+	
1611	-	
1612	rac	
1613	+	
1614	-	
1615	rac	
1616	+	
1617	-	

1618	rac	
1619	+	
1620	-	

Biological Characterization of the Compounds According to the Invention

The identification of progesterone receptor modulators can be performed using simple methods, test programs that are known to one skilled in the art. To this end, for example, a compound that is to be tested can be incubated together with a gestagen in a test system for progesterone receptors, and it can be examined whether the progesterone-mediated action in this test system is altered in the presence of modulators.

The substances of general formula I according to the invention were tested in the following models:

Progesterone Receptor Binding Test

Measurement of the Receptor Binding Affinity:

The receptor binding affinity was determined by competitive binding of a specifically binding ^3H -labeled hormone (tracer) and the compound to be tested on receptors in the cytosol from animal target organs. In this case, receptor saturation and reaction equilibrium were sought.

The tracers and increasing concentrations of the compound to be tested (competitor) were co-incubated with the receptor-containing cytosol fraction at 0-4°C over 18 hours. After separation of the unbonded tracer with carbon-dextran suspension, the receptor-bonded tracer portion was measured for each concentration, and the IC_{50} was determined from the concentration sequence. As a quotient of the IC_{50} values of the reference substance and the compound to be tested ($\times 100\%$), the relative molar binding affinity (RBA) was calculated (RBA of the reference substance = 100%).

For the receptor types, the following incubation conditions were selected:

Progesterone Receptor:

Uterus cytosol of the estradiol-primed rabbit, homogenized in TED buffer (20 mmol of Tris/HCl, pH 7.4; 1 mmol of ethylenediamine tetraacetate, 2 mmol of dithiothreitol) with 250 mmol of saccharose; stored at -30°C. Tracer: ^3H -ORG 2058, 5 nmol; reference substance: progesterone.

Glucocorticoid Receptor:

Thymus cytosol of the adrenalectomized rat, thymi stored at -30°C; buffer: TED. Tracer: ^3H -Dexamethasone, 20 nmol; reference substance: dexamethasone.

The relative receptor binding affinities (RBA values) of the compounds of general formula (I) according to the invention on the progesterone receptor are between 3 and 100% relative to the progesterone. On the glucocorticoid receptor, the RBA values are in the range of 3 to 30% relative to dexamethasone.

The compounds according to the invention accordingly have a high affinity to the progesterone receptor but only a low affinity to the glucocorticoid receptor.

Antagonism of Progesterone Receptor PR-B

The transactivation assay is performed as described in WO 02/054064.

Agonism of Progesterone Receptor PR-B

The transactivation assay is performed as described in Fuhrmann et al. (Fuhrmann, U.; Hess-Stump, H.; Cleve, A.; Neef, G.; Schwede, W.; Hoffmann, J.; Fritzemeier, K.-H.; Chwalisz, K.; Journal of Medicinal Chem., 43, 26, 2000, 5010-5016).

No.	Antagonistic activity		Agonistic activity	
	IC ₅₀ [nM]	Efficacy [%]	EC ₅₀ [nM]	Efficacy [%]
9	3	96	n.b.	3
13	3	92	n.b.	7
3b	0,4	97	n.b.	2
7	4	82	2	11

Dosage

For use according to the invention, the progesterone receptor modulators can be administered orally, enterally, parenterally or transdermally.

In general, satisfactory results can be expected in the treatment of the above-mentioned indications if the daily doses encompass a range of 1 µg to 500 mg of the compound according to the invention.

Suitable dosages of the compounds according to the invention in humans for the treatment of endometriosis, leiomyomas of the uterus and dysfunctional bleeding as well as for use in birth control as well as for hormone replacement therapy are 50 µg to 500 mg per day, depending on age and constitution of the patient, whereby the necessary daily dose can be administered one or more times.

For treatment of breast cancer, the dosage range for the compounds according to the invention comprises 10 mg to 1000 mg daily.

The formulation of the pharmaceutical preparations based on the new compounds is carried out in a way that is known in the art, by the active ingredient being processed with the vehicles, fillers, substances that influence decomposition, binding agents, moisturizers, lubricants, absorbents, diluents, flavoring correctives, dyes, etc., that are commonly used in galenicals and being converted into the desired form of administration.

In this case, reference is made to Remington's Pharmaceutical Science, 15th Ed. Mack Publishing Company, East Pennsylvania (1980).

For oral administration, in particular tablets, film tablets, coated tablets, capsules, pills, powders, granulates, lozenges, suspensions, emulsions or solutions are suitable.

For parenteral administration, injection and infusion preparations are possible.

For intraarticular injection, correspondingly prepared crystal suspensions can be used.

For intramuscular injection, aqueous and oily injection solutions or suspensions and corresponding depot preparations can be used.

For rectal administration, the new compounds can be used in the form of suppositories, capsules, solutions (e.g., in the form of enemas) and ointments both for systemic therapy and for local therapy.

In addition, agents for vaginal application can also be mentioned as preparations.

For pulmonary administration of the new compounds, the latter can be used in the form of aerosols and inhalants.

For transdermal administration, patches are possible, or for topical application, formulations in gels, ointments, fatty ointments, creams, pastes, powders, milk and tinctures are possible. The dosage of the compounds of general formula I should be 0.01% - 20% in these preparations to achieve an adequate pharmacological action.

Corresponding tablets can be obtained by, for example, mixing active ingredient with known adjuvants, for example inert diluents such as dextrose, sugar, sorbitol, mannitol, polyvinylpyrrolidone, explosives such as corn starch or alginic acid, binders such as starch or gelatin, lubricants such as magnesium stearate or talc and/or agents for achieving a depot effect such as carboxylpolymethylene, carboxyl methyl cellulose,

cellulose acetate phthalate or polyvinyl acetate. The tablets can also consist of several layers.

Accordingly, coated tablets can be produced by coating cores, produced analogously to the tablets, with agents that are commonly used in tablet coatings, for example polyvinylpyrrolidone or shellac, gum arabic, talc, titanium oxide or sugar. In this case, the coated tablet shell can also consist of several layers, whereby the adjuvants that are mentioned above in the tablets can be used.

Solutions or suspensions of the compounds of general formula I according to the invention can contain additional taste-improving agents such as saccharine, cyclamate or sugar, as well as, e.g., flavoring substances, such as vanilla or orange extract. In addition, they can contain suspending adjuvants such as sodium carboxy methyl cellulose or preservatives such as p-hydroxybenzoates.

The capsules that contain compounds of general formula I can be produced by, for example, the compound(s) of general formula I being mixed with an inert vehicle such as lactose or sorbitol and encapsulated in gelatin capsules.

Suitable suppositories can be produced by, for example, mixing with vehicles that are provided for this purpose, such as neutral fats or polyethylene glycol, or derivatives thereof.

The compounds of general formula (I) according to the invention or their pharmaceutically acceptable salts can be used based on their antagonistic or partial agonistic action for the production of a pharmaceutical agent, in particular for treatment and prophylaxis of gynecological diseases, such as endometriosis, leiomyomas of the uterus, dysfunctional bleeding and dysmenorrhea. In addition, they can be used to counteract hormonal irregularities, to trigger menstruation and alone or in combination with prostaglandins and/or oxytocin to induce birth.

In addition, the compounds of general formula (I) according to the invention or their pharmaceutically acceptable salts are suitable for the production of preparations for contraception for women (see also WO 93/23020, WO 93/21927).

In addition, the compounds according to the invention or their pharmaceutically acceptable salts can be used alone or in combination with a Selective Estrogen Receptor Modulator (SERM) for female hormone replacement therapy.

In addition, the above-mentioned compounds exert an antiproliferative action in hormone-dependent tumors. They are therefore suitable for the therapy of hormone-dependent carcinomas, such as, for example, for breast, prostate or endometrial carcinomas.

The compounds according to the invention or their pharmaceutically acceptable salts can be used for the treatment of hormone-dependent carcinomas, both in first-line therapy and in second-line therapy, in particular after tamoxifen failure.

The compounds of general formula (I) according to the invention that have an antagonistic or partial agonistic action or their pharmaceutically acceptable salts can also be used in combination with compounds that have an antiestrogenic action (estrogen receptor antagonists or aromatase inhibitors) or Selective Estrogen Receptor Modulators (SERM) for the production of pharmaceutical preparations for treating hormone-dependent tumors. For the treatment of endometriosis or leiomyomas of the uterus, the compounds according to the invention can also be used in combination with SERMs or an antiestrogen (estrogen receptor antagonists or aromatase inhibitors). In the treatment of hormone-dependent tumors, the progesterone receptor modulator and the antiestrogen (estrogen receptor antagonists or aromatase inhibitors) or the SERM can be provided for simultaneous or else for sequential administration. In sequential administration, preferably first the antiestrogen (estrogen receptor antagonists or aromatase inhibitor) or SERM is administered, and then the progesterone receptor modulator is administered.

In this case, in the combination with the nonsteroidal progesterone receptor modulators according to the invention, for example, the following antiestrogens (estrogen receptor antagonists or aromatase inhibitors) or SERMs are considered:

tamoxifen, 5-(4-{5-[(RS)-(4,4,5,5,5-pentafluoropentyl)sulfinyl]-pentyloxy}phenyl)-6-phenyl-8,9-dihydro-7H-benzocyclohepten-2-ol (WO 00/03979), ICI 182 780 (7alpha-[9-(4,4,5,5-pentafluoropentyl)sulfinyl]nonyl]estra-1,3,5(10)-triene-3,17-beta-diol), 11beta-fluoro-7alpha-[5-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17beta-diol (WO98/07740), 11beta-fluoro-7alpha-{5-[methyl(7,7,8,8,9,9,10,10,10-nonafluorodecyl)amino]pentyl}estra-1,3,5(10)-triene-3,17beta-diol (WO 99/33855), 11beta-fluoro-17alpha-methyl-7alpha-{5-[methyl(8,8,9,9,9-pentafluorononyl)-amino]pentyl}estra-1,3,5(10)-triene-3,17beta-diol (WO 03/045972), clomifene, raloxifene as well as other antiestrogenically active compounds, and aromatase inhibitors, such as, for example, fadrozole, formestane, letrozole, anastrozole or atamestane.

Finally, this invention also relates to the use of the compounds of general formula I, optionally together with an antiestrogen or SERM, for the production of a pharmaceutical agent.

This invention also relates to pharmaceutical compositions that contain at least one compound according to the invention, optionally in the form of a pharmaceutically/pharmacologically compatible salt, without or together with pharmaceutically compatible adjuvants and/or vehicles.

These pharmaceutical compositions and pharmaceutical agents can be provided for oral, rectal, vaginal, subcutaneous, percutaneous, intravenous or intramuscular

administration. In addition to commonly used vehicles and/or diluents, they contain at least one compound according to the invention.

The pharmaceutical agents of the invention are produced in a known way with the commonly used solid or liquid vehicles or diluents and the usually used pharmaceutical-technical adjuvants corresponding to the desired type of administration with a suitable dosage. The preferred preparations exist in a dispensing form that is suitable for oral administration. Such dispensing forms are, for example, tablets, film tablets, coated tablets, capsules, pills, powders, solutions or suspensions or else depot forms.

The pharmaceutical compositions that contain at least one of the compounds according to the invention are preferably administered orally.


Parenteral preparations, such as injection solutions, are also considered.

In addition, for example, suppositories and agents for vaginal application can also be mentioned as preparations.

The following examples are used for a more detailed explanation of the subject of the invention, without intending that it be limited to these examples.

The production of the starting compound 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide is described in Patent WO 200375915, and the production of 5-{3-[1-phenyl-cyclopropyl]-2-oxopropionylamino}phthalide is described in WO 9854159.


rac-5-{2-Ethynyl-2-hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-propionylamino}phthalide 1



Ethynyl magnesium bromide (6 ml, 0.5 M in tetrahydrofuran) was added to an ice-cold solution that consists of 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide (632 mg) in THF (4 ml). The reaction solution under argon was allowed to come to room temperature within 3 hours. Then, the reaction mixture was poured into ice-cold, saturated ammonium chloride solution. It was extracted with ethyl acetate. The combined organic phases were washed with saturated sodium chloride solution and dried on sodium sulfate. The crude product that is obtained was chromatographed on silica gel. 2.2 g of product was obtained.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.83 (1H), 0.92-1.10 (2H), 2.37 (1H), 2.56 (1H), 2.59 (1H), 3.10 (1H), 5.28 (2H), 7.02 (1H), 7.31 (1H), 7.37 (1H), 7.58 (1H), 7.86 (1H), 7.94 (1H), 8.70 (1H).


rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-propinyl-propionylamino}phthalide 2



Analogously to Example 1, 145 mg of product was obtained from 1-propynylmagnesium bromide (2 ml of 0.5 M solution in tetrahydrofuran) and 210 mg of 6-[4-(2-chloro-4-fluorophenyl)-4-methyl-2-oxovaleroylamino]-4-methyl-2,3-benzoxazin-1-one.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.86 (1H), 0.90-1.05 (3H), 1.72 (3H), 2.35 (1H), 2.49 (1H), 2.96 (1H), 5.27 (2H), 7.03 (1H), 7.30 (1H), 7.36 (1H), 7.58 (1H), 7.85 (1H), 7.98 (1H), 8.73 (1H).

(+)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-phenyl-propionylamino}phthalide 3a and
(-)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(phenylethynyl)-propionylamino}phthalide 3b



n-Butyllithium (625 μ l, 1.6 M in hexane) was added at -78°C to a solution of 110 μ l of phenylacetylene in tetrahydrofuran. Stirring was allowed to continue at this temperature for 30 minutes, and then a solution of 5-{3-[1-(2-fluoro-5-

trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide (210 mg) in 5 ml of tetrahydrofuran was added in drops. Then, it was allowed to come to 23°C over about 3 hours and then stirred for 10 more hours. Then, the reaction mixture was poured into ice-cold, saturated ammonium chloride solution. It was extracted with ethyl acetate. The combined organic phases were washed with saturated sodium chloride solution and dried on sodium sulfate. The crude product was chromatographed on silica gel. The racemic mixture obtained was then separated by preparative chiral HPLC (Chiralpak AD column, 250x10 mm) into enantiomers 3a (46 mg) and 3b (47 mg).

3a and 3b:


¹H-NMR (ppm, CDCl₃, 300 MHz): 0.88 (1H), 0.95-1.11 (3H), 2.46 (1H), 2.65 (1H), 3.10 (1H), 5.27 (2H), 7.00 (1H), 7.24-7.42 (7H), 7.61 (1H), 7.84 (1H), 7.98 (1H), 8.80 (1H).

3a : $[\alpha]_{20}^D$: + 12.9° (CHCl₃, 1.06 g/100 ml; λ=589 nm)

3b : $[\alpha]_{20}^D$: - 14.4° (CHCl₃, 1.03 g/100 ml; λ=589 nm)

Analogously to Example 3, compounds 4 and 5 were produced from 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide and the respective lithium aryl acetylide.

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-trifluoromethylphenyl)ethynyl]propionylamino}phthalide 4



$^1\text{H-NMR}$ (ppm, CDCl_3 , 300 MHz): 0.92 (1H), 0.99-1.16 (3H), 2.55 (1H), 2.68 (1H), 3.27 (1H), 5.30 (2H), 7.03 (1H), 7.30-7.52 (4H), 7.55-7.62 (2H), 6.67 (1H), 7.99 (1H), 8.03 (1H), 8.84 (1H).

(+)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-trifluoromethylphenyl)ethynyl]propionylamino}phthalide **4a** and


(-)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-trifluoromethylphenyl)ethynyl]propionylamino}phthalide **4b**

The racemic mixture (150 mg) which was described in example **4** was separated by preparative chiral HPLC (column Chiralpak AD 250x10 mm) into the enantiomers **4a** (51 mg) and **4b** (62 mg).

4a : $[\alpha]_{20}^D$: + 6.3° (CHCl_3 , 1.07 g/100 ml; $\lambda=589$ nm)

4b : $[\alpha]_{20}^D$: - 5.3° (CHCl_3 , 1.09 g/100 ml; $\lambda=589$ nm)

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-methylphenyl)ethynyl]propionylamino}phthalide **5**



$^1\text{H-NMR}$ (ppm, CDCl_3 , 300 MHz): 0.87 (1H), 0.93-1.15 (3H), 2.38 (3H), 2.45 (1H), 2.66 (1H), 3.11 (1H), 5.25 (2H), 6.99 (1H), 7.10 (2H), 7.18-7.38 (4H), 7.61 (1H), 7.86 (1H), 8.00 (1H), 8.80 (1H).


(+)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-methylphenyl)ethynyl]propionylamino}phthalide 5a and
(-)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-methylphenyl)ethynyl]propionylamino}phthalide 5b

The racemic mixture (109 mg) which was described in example 5 was separated by preparative chiral HPLC (column Chiralpak AD 250x10 mm) into the enantiomers 5a (41 mg) and 5b (28 mg).

5a : $[\alpha]_{20}^D$: + 14.8° (CHCl_3 , 1.07 g/100 ml; λ =589 nm)

5b : $[\alpha]_{20}^D$: - 16.3° (CHCl_3 , 1.13 g/100 ml; λ =589 nm)

rac-5-{2-[(4-Acetoxyphenyl)ethynyl]-2-hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-propionylamino}phthalide 6




A suspension of the compound (104 mg) described under Example 1, triphenylphosphine (12.2 mg), copper iodide (8.9 mg), 4-iodophenyl acetate (92 mg), palladium acetate (5.3 mg) in THF (5 ml) and triethylamine (5 ml) was reacted for 1 hour in an ultrasound bath at 25°C under argon. Then, it was poured into saturated, aqueous ammonium chloride solution. It was extracted with ethyl acetate and washed with water and saturated sodium chloride solution. The combined organic phases were

dried on sodium sulfate. After column chromatography of the crude product on silica gel, 55 mg of product was obtained.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.88 (1H), 0.95-1.10 (3H), 2.29 (3H), 2.45 (1H), 2.63 (1H), 3.17 (1H), 5.29 (2H), 6.97-7.07 (3H), 7.28-7.37 (4H), 7.60 (1H), 7.84 (1H), 7.98 (1H), 8.80 (1H).


rac-5-{2-Hydroxy-2-[(4-hydroxyphenyl)ethynyl]-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-propionylamino}phthalide 7



A solution of the compound described under 6 (45 mg) in 5 ml of methanol was mixed with sodium bicarbonate (130 mg). Stirring was continued for 2 more hours at 23°C. Then, the reaction mixture was diluted with ethyl acetate. Then, it was washed twice with saturated sodium chloride solution. After drying on sodium sulfate, the crude product was purified on silica gel by column chromatography. 38 mg of product was obtained.

¹H-NMR (ppm, CDCl₃, 300 MHz): 0.87 (1H), 0.92-1.11 (3H), 2.43 (1H), 2.64 (1H), 3.11 (1H), 5.27 (2H), 5.67 (1H), 6.73 (2H), 6.98 (1H), 7.14 (2H), 7.28-7.38 (2H), 7.60 (1H), 7.85 (1H), 7.97 (1H), 8.84 (1H).


rac-5-{2-[(4-Carboxyphenyl)ethynyl]-2-hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-propionylamino}phthalide 8



Analogously to Example 6, compound 8 was produced from the compound described under Example 1 and 4-iodobenzoic acid.

¹H-NMR (ppm, CDCl₃/MeOD (5%), 400 MHz): 0.82 (1H), 0.89-1.05 (3H), 2.37 (1H), 2.65 (1H), 5.24 (2H), 6.97 (1H), 7.35 (1H), 7.44 (2H), 7.50-7.65 (2H), 7.72 (1H), 7.80 (1H), 7.92 (2H).

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(pentin-1-yl)-propionylamino}phthalide 9



A solution that consists of 1-pentyne (0.94 ml) in THF (9 ml) was mixed at -78°C with nBuLi (0.6 ml, 1.6 M in hexane). It was allowed to stir for 30 minutes at -78°C , and then a solution of 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide (200 mg) in 3 ml of tetrahydrofuran was added. Then, it was allowed to come to 23°C over about 3 hours, and it was stirred for 10 more hours at this temperature. Then, the reaction mixture was poured into ice-cold, saturated ammonium chloride solution. It was extracted with ethyl acetate. The combined organic phases were washed with saturated sodium chloride solution and dried on

sodium sulfate. The crude product was chromatographed on silica gel. 130 mg of product was obtained.

$^1\text{H-NMR}$ (ppm, CDCl_3 , 400 MHz): 0.82 (1H), 0.92-1.07 (6H), 1.45 (2H), 2.08 (2H), 2.30 (1H), 2.53 (1H), 2.83 (1H), 5.27 (2H), 7.02 (1H), 7.29 (1H), 7.36 (1H), 7.57 (1H), 7.84 (1H), 7.96 (1H), 8.72 (1H).


(+)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(pentin-1-yl)-propionylamino}phthalide 9a and
(-)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(pentin-1-yl)-propionylamino}phthalide 9b

The racemic mixture (120 mg) which was described in example 9 was separated by preparative chiral HPLC (column Chiralpak AD 250x10 mm) into the enantiomers 9a (46 mg) and 9b (47 mg).

9a : $[\alpha]_{20}^{\text{D}}$: + 10.9° (CHCl_3 , 1.01 g/100 ml; λ =589 nm)

9b : $[\alpha]_{20}^{\text{D}}$: - 10.6° (CHCl_3 , 1.08 g/100 ml; λ =589 nm)


rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(hexin-1-yl)-propionylamino}phthalide 10



Compound 10 was synthesized analogously to Example 9.

$^1\text{H-NMR}$ (ppm, CDCl_3 , 400 MHz): 0.80-1.06 (7H), 1.30-1.50 (2H), 1.59 (2H), 2.10 (2H), 2.30 (1H), 2.52 (1H), 2.82 (1H), 5.28 (2H), 7.02 (1H), 7.30 (1H), 7.36 (1H), 7.57 (1H), 7.84 (1H), 7.95 (1H), 8.72 (1H).

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-hydroxy)butin-1-yl]-propionylamino}phthalide 11




Stage A: Reaction of 4-(tert-butyldimethylsilyloxy)but-1-yne (175 mg), nBuLi (0.59 ml, 1.6 M in hexane) 5-{3-[1-(2-Fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}-phthalide (200 mg) in tetrahydrofuran analogously to the process described under Example 9 yielded 165 mg of product.

Stage B: The product obtained under stage A (160 mg) was dissolved in 5 ml of tetrahydrofuran. At 0°C, 270 µl of a 1 molar solution of tetrabutylammonium fluoride in tetrahydrofuran was added and stirred for one hour at 0°C and for another 2 hours at 23°C. Then, the reaction mixture was poured into saturated, aqueous sodium bicarbonate solution. It was extracted several times with ethyl acetate. The combined organic phases were washed with saturated sodium chloride solution and dried on sodium sulfate. After column chromatography on silica gel, 77 mg of product was obtained.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.83 (1H), 0.90-1.03 (3H), 2.20-2.40 (3H), 2.50 (1H), 3.39 (1H), 3.68 (2H), 5.25 (2H), 7.01 (1H), 7.32 (2H), 7.57 (1H), 7.82 (1H), 7.93 (1H), 8.91 (1H).


Analogously to Example 11, compounds 12 and 13 were produced from 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide:

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(5-hydroxy)pentin-1-yl]-propionylamino}phthalide 12



¹H-NMR (ppm, CDCl₃, 400 MHz): 0.83 (1H), 0.90-1.03 (3H), 1.70 (2H), 2.24 (2H), 2.33 (1H), 2.50 (1H), 3.09 (1H), 3.71 (2H), 5.26 (2H), 7.02 (1H), 7.35 (2H), 7.57 (1H), 7.83 (1H), 7.97 (1H), 8.82 (1H).

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(3-hydroxy)propin-1-yl]-propionylamino}phthalide 13



¹H-NMR (ppm, CDCl₃, 400 MHz): 0.84 (1H), 0.90-1.03 (3H), 2.37 (1H), 2.52 (1H), 3.25 (1H), 4.17 (2H), 5.27 (2H), 7.02 (1H), 7.30-7.40 (2H), 7.58 (1H), 7.83 (1H), 7.91 (1H), 8.77 (1H).


**(+)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(3-hydroxy)propin-1-yl]-propionylamino}phthalide 13a and
(-)-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(3-hydroxy)propin-1-yl]-propionylamino}phthalide 13b**

The racemic mixture (80 mg) which was described in example **13** was separated by preparative chiral HPLC (column Chiralpak AD 250x10 mm) into the enantiomers **13a** (35 mg) and **13b** (37 mg).

13a : $[\alpha]_{20}^D$: + 28.3° (CHCl₃, 1.01 g/100 ml; λ =589 nm)

13b : $[\alpha]_{20}^D$: - 29.3° (CHCl₃, 1.08 g/100 ml; λ =589 nm)

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-[(4-hydroxy-3-methyl)butin-1-yl]-propionylamino}phthalide **14**




Stage A: Analogously to Example 11, 300 mg of 5-{3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-oxopropionylamino}phthalide and 282 mg of tert-butyl-(1,1-dimethylprop-2-ynyl-oxy)-dimethylsilane are reacted. 15 mg of product A is obtained.

Stage B: 70 mg of the compound that is obtained under A was dissolved in 1 ml of dichloromethane. 650 μ l of trifluoroacetic acid (20% in dichloromethane) was added at 0°C, and it was stirred for 3.5 hours at 0°C. Then, it was evaporated to the dry state in a vacuum, and the residue was purified by column chromatography on silica gel. 27 mg of product was obtained.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.82 (1H), 0.90-1.00 (2H), 1.04 (1H), 1.47 (6H), 2.28 (1H), 2.58 (1H), 3.08 (1H), 5.27 (2H), 7.03 (1H), 7.30 (1H), 7.36 (1H), 7.59 (1H), 7.83 (1H), 7.91 (1H), 8.78 (1H).


rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(2-(tert-butylcarboxy)ethin-1-yl)-propionylamino}phthalide 15



Compound 15 was synthesized analogously to Example 9.

¹H-NMR (ppm, CDCl₃, 400 MHz): 0.87 (1H), 0.93-1.05 (3H), 1.46 (9H), 2.42 (1H), 2.59 (1H), 3.39 (1H), 5.28 (2H), 7.03 (1H), 7.30-7.42 (2H), 7.57 (1H), 7.85 (1H), 7.92 (1H), 8.68 (1H).

rac-5-{2-Hydroxy-3-[1-(2-fluoro-5-trifluoromethylphenyl)-cyclopropyl]-2-(2-carboxyethin-1-yl)-propionylamino}phthalide 16




50 mg of the compound that is described under Example 15 was dissolved in 5 ml of dichloromethane. 100 µl of trifluoroacetic acid was added, and it was stirred for 12 more hours at 23°C. Then, it was evaporated to the dry state in a vacuum, and the residue was purified by column chromatography on silica gel. 36 mg of product was obtained.

¹H-NMR (ppm, DMSO-D₆, 300 MHz): 0.48 (1H), 0.78 (1H), 0.86 (1H), 1.09 (1H), 1.73 (1H), 2.89 (1H), 5.27 (2H), 6.62 (1H), 7.13 (1H), 7.31 (1H), 7.41 (1H), 7.53 (1H), 7.62 (1H), 7.68 (1H), 9.85 (1H).

Analogously to Example 3, compound 17 was produced from 5-{3-[1-phenyl-cyclopropyl]-2-oxopropionylamino}phthalide:


rac-5-{2-Hydroxy-3-[1-phenyl-cyclopropyl]-2-(phenyl-ethinyl)propionylamino}-phthalide 17



¹H-NMR (ppm, CDCl₃, 400 MHz): 0.78 (1H), 0.90 (1H), 1.10-1.21 (2H), 2.38 (1H), 2.72 (1H), 2.77 (1H), 5.28 (2H), 7.18 (1H), 7.25-7.42 (6H), 7.41-7.52 (4H), 7.82 (1H), 8.06 (1H), 8.79 (1H).

Claims

1. Compounds of general formula I



in which

R^1 and R^2 , independently of one another, mean a hydrogen atom, a straight or nonstraight, branched or unbranched C_1 - C_5 -alkyl group, also together with the C atom of the chain forming a ring with a total of 3-7 members,

R^3 means a radical $C\equiv C-R^a$, whereby

R^a means a hydrogen or a C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl, or heterocycloalkyl that optionally is substituted in one or more places, in the same way or differently, with K, or an aryl or heteroaryl that optionally is substituted in one or more places, in the same way or differently, with L,

K is a cyano, halogen, hydroxy, nitro, $-C(O)R^b$, CO_2R^b , $-O-R^b$, $-S-R^b$, $SO_2NR^cR^d$, $-C(O)-NR^cR^d$, $-OC(O)-NR^cR^d$, or $-C=NOR^b-NR^cR^d$ or a C_3 - C_{10} -cycloalkyl that optionally is substituted in one or more places, in the same way or differently, with M, heterocycloalkyl, or aryl or heteroaryl that optionally is substituted in one or more places with L,

L means C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_1 - C_6 -

perfluoroalkyl, C₁-C₆-perfluoroalkoxy, C₁-C₆-alkoxy-C₁-C₆-alkoxy, (CH₂)_p-C₃-C₁₀-cycloalkyl, (CH₂)_p-heterocycloalkyl, (CH₂)_pCN, (CH₂)_pHal, (CH₂)_pNO₂, (CH₂)_p-C₆-C₁₂-aryl, (CH₂)_p-heteroaryl, -(CH₂)_pPO₃(R^b)₂, -(CH₂)_pNR^cR^d, -(CH₂)_pNR^eCOR^b, -(CH₂)_pNR^eCSR^b, -(CH₂)_pNR^eS(O)R^b, -(CH₂)_pNR^eS(O)₂R^b, -(CH₂)_pNR^eCONR^cR^d, -(CH₂)_pNR^eCOOR^b, -(CH₂)_pNR^eC(NH)NR^cR^d, -(CH₂)_pNR^eCSNR^cR^d, -(CH₂)_pNR^eS(O)NR^cR^d, -(CH₂)_pNR^eS(O)₂NR^cR^d, -(CH₂)_pCOR^b, -(CH₂)_pCSR^b, -(CH₂)_pS(O)R^b, -(CH₂)_pS(O)(NH)R^b, -(CH₂)_pS(O)₂R^b, -(CH₂)_pS(O)₂NR^cR^d, -(CH₂)_pSO₂OR^b, -(CH₂)_pCO₂R^b, -(CH₂)_pCONR^cR^d, -(CH₂)_pCSNR^cR^d, -(CH₂)_pOR^b, -(CH₂)_pSR^b, -(CH₂)_pCR^b(OH)-R^e, -(CH₂)_p-C=NOR^b, -O-(CH₂)_n-O-, -O-(CH₂)_n-CH₂-, -O-CH=CH- or -(CH₂)_{n+2}-, whereby n = 1 or 2, and the terminal oxygen atoms and/or carbon atoms are linked to directly adjacent ring-carbon atoms,

M means C₁-C₆-alkyl or a group -COR^b, CO₂R^b, -O-R^b, or -NR^cR^d, whereby

R^b means a hydrogen or a C₁-C₆-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkinyl, C₃-C₁₀-cycloalkyl, C₆-C₁₂-aryl or C₁-C₃-perfluoroalkyl and

R^c and R^d , independently of one another, mean a hydrogen,

C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{12} -aryl, $C(O)R^b$ or a hydroxy group, whereby if

R^c is a hydroxy group, R^d can be only one hydrogen, a C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl or C_6 - C_{12} -aryl and vice versa, and

R^c means a hydrogen, C_1 - C_6 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_3 - C_{10} -cycloalkyl or C_6 - C_{12} -aryl, and

p can be a number from 0- 6,

or

R^3 is a radical $C = C-R^gR^h$, whereby

R^g and R^h , independently of one another, are a hydrogen or a C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl or C_2 - C_8 -alkinyl that optionally is substituted in one or more places, in the same way or differently, with X, in which

X is a cyano, halogen, hydroxy, nitro, $-C(O)R^b$, CO_2R^b , $-O-R^b$, $-C(O)-NR^cR^d$, $-NR^cR^d$ with the meanings, already further mentioned above, for R^b , R^c and R^d , and

R^{4a} and R^{4b} , independently of one another, mean a hydrogen atom, a C_1 - C_4 -alkyl, a C_2 - C_4 -alkenyl or together with the ring-carbon atom forming a 3- to 6-membered ring,

A means a monocyclic or bicyclic, carbocyclic or heterocyclic aromatic ring, which optionally can be substituted in one or more places with C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkinyl, C_1 - C_6 -perfluoroalkyl, C_1 - C_6 -

perfluoroalkoxy, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy,
 (CH₂)_p-C₃-C₁₀-cycloalkyl, (CH₂)_p-heterocycloalkyl, (CH₂)_pCN,
 (CH₂)_pHal, (CH₂)_pNO₂, (CH₂)_p-C₆-C₁₂-aryl, (CH₂)_p-heteroaryl,
 -(CH₂)_pPO₃(R^b)₂, -(CH₂)_pNR^cR^d, -(CH₂)_pNR^cCOR^b, -(CH₂)_pNR^cCSR^b,
 -(CH₂)_pNR^cS(O)R^b, -(CH₂)_pNR^cS(O)₂R^b, -(CH₂)_pNR^cCONR^cR^d,
 -(CH₂)_pNR^cCOOR^b, -(CH₂)_pNR^cC(NH)NR^cR^d, -(CH₂)_pNR^cCSNR^cR^d,
 -(CH₂)_pNR^cS(O)NR^cR^d, -(CH₂)_pNR^cS(O)₂NR^cR^d, -(CH₂)_pCOR^b,
 -(CH₂)_pCSR^b, -(CH₂)_pS(O)R^b, -(CH₂)_pS(O)(NH)R^b, -(CH₂)_pS(O)₂R^b,
 -(CH₂)_pS(O)₂NR^cR^d, -(CH₂)_pSO₂OR^b, -(CH₂)_pCO₂R^b, -(CH₂)_pCONR^cR^d,
 -(CH₂)_pCSNR^cR^d, -(CH₂)_pOR^b, -(CH₂)_pSR^b, -(CH₂)_pCR^b(OH)-R^d,
 -(CH₂)_p-C=NOR^b, -O-(CH₂)_n-O-, -O-(CH₂)_n-CH₂-, -O-CH=CH- or
 -(CH₂)_{n+2}-, whereby n = 1 or 2, and the terminal oxygen atoms and/or
 carbon atoms are linked to directly adjacent ring-carbon atoms, or

A means a radical -CO₂R^b, C(O)NR^cR^d, COR^b,

or

A means an alkenyl group -CR⁵=CR⁶R⁷, whereby

R⁵, R⁶ and R⁷ are the same or different and, independently of one another, mean

hydrogen atoms, halogen atoms, aryl radicals or an unsubstituted or

partially or completely fluorinated C₁-C₅-alkyl group, or

A means an alkynyl group -C≡CR⁵, with the meaning cited above for R⁵, and

B means a carbonyl group or a CH₂ group







as well as their pharmaceutically acceptable salts.


2. Compounds according to claim 1, in which R¹ and R² preferably mean a hydrogen atom, a methyl group or an ethyl group.









3. Compounds according to claim 1, in which R^1 and R^2 preferably together with the C atom of the chain form a ring with a total of 3 to 7 links.
4. Compounds according to claims 1 to 3, in which R^3 preferably means alkenyl, alkynyl, arylalkynyl, heteroarylalkynyl, cycloalkylalkynyl, or heterocycloalkylalkynyl.
5. Compounds according to one of the preceding claims, in which R^3 preferably means a vinyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, hydroxypropynyl, hydroxybutynyl, 3-hydroxy-3-methylbutynyl, hydroxypentynyl, carboxypropynyl, t-butylcarboxypropynyl, phenylethynyl, (hydroxyphenyl)ethynyl, (methoxyphenyl)ethynyl, (dimethylaminophenyl)ethynyl, (methylphenyl)ethynyl, (cyanophenyl)ethynyl, (trifluoromethyl)ethynyl, (diphenyl)ethynyl, (nitrophenyl)ethynyl, (tert-butylphenyl)ethynyl, (acetylphenyl)ethynyl, (acetoxypheynyl)ethynyl, (carboxyphenyl)ethynyl or a benzylethynyl group.
6. Compounds according to one of the preceding claims, in which A is preferably an aromatic ring.
7. Compounds according to one of the preceding claims, in which A is preferably a phenyl or naphthyl radical.
8. Compounds according to claim 7, in which A preferably is an unsubstituted phenyl radical or optionally a phenyl radical that is substituted in one or more places.
9. Compounds according to claim 8, whereby the phenyl radical is preferably substituted with one or two halogen atoms or a trifluoromethyl group.
10. Compounds according to claim 9, in which the halogen atoms are preferably chlorine and/or fluorine.
11. Compounds according to claims 1-8, in which A preferably is an $-O-(CH_2)_n-$ O- or $-O-(CH_2)_n-CH_2-$ substituted phenyl ring, whereby the respectively directly adjacent ring-carbon atoms are linked.






12. Compounds according to one of the preceding claims, in which R^{4a} and R^{4b} , independently of one another, in each case are a hydrogen atom.




13. Compounds according to claims 1 to 12, namely:


No.	Racemic or Enantiomer	R3
1	rac	
2	+	
3	-	
4	rac	
5	+	
6	-	
7	rac	
8	+	
9	-	
10	rac	
11	+	
12	-	
13	rac	
14	+	
15	-	
16	rac	
17	+	
18	-	
19	rac	
















20	+	
21	-	
22	rac	
23	+	
24	-	
25	rac	
26	+	
27	-	
28	rac	
29	+	
30	-	
31	rac	
32	+	
33	-	
34	rac	
35	+	
36	-	
37	rac	
38	+	
39	-	
40	rac	
41	+	
42	-	
43	rac	


44	+	
45	-	
46	rac	
47	+	
48	-	
49	rac	
50	+	
51	-	
52	rac	
53	+	
54	-	
55	rac	
56	+	
57	-	
58	rac	
59	+	
60	-	


No.	Racemic or Enantiomer	R3
61	rac	
62	+	
63	-	
64	rac	
65	+	
66	-	
67	rac	
68	+	
69	-	
70	rac	
71	+	
72	-	
73	rac	
74	+	
75	-	
76	rac	
77	+	
78	-	
79	rac	
80	+	
81	-	

















82	rac	
83	+	
84	-	
85	rac	
86	+	
87	-	
88	rac	
89	+	
90	-	
91	rac	
92	+	
93	-	
94	rac	
95	+	
96	-	
97	rac	
98	+	
99	-	
100	rac	
101	+	
102	-	
103	rac	
104	+	
105	-	




106	rac	
107	+	
108	-	
109	rac	
110	+	
111	-	
112	rac	
113	+	
114	-	
115	rac	
116	+	
117	-	
118	rac	
119	+	
120	-	



No.	Racemic or Enantiomer	R3
121	rac	
122	+	
123	-	
124	rac	












125	+	
126	-	
127	rac	
128	+	
129	-	
130	rac	
131	+	
132	-	
133	rac	
134	+	
135	-	
136	rac	
137	+	
138	-	
139	rac	
140	+	
141	-	
142	rac	
143	+	
144	-	
145	rac	
146	+	









147	-	
148	rac	
149	+	
150	-	
151	rac	
152	+	
153	-	
154	rac	
155	+	
156	-	
157	rac	
158	+	
159	-	
160	rac	
161	+	
162	-	
163	rac	
164	+	
165	-	
166	rac	
167	+	
168	-	
169	rac	
170	+	


171	-	
172	rac	
173	+	
174	-	
175	rac	
176	+	
177	-	
178	rac	
179	+	
180	-	




No.	Racemic or Enantiomer	R3
181	rac	
182	+	
183	-	
184	rac	
185	+	
186	-	
187	rac	
188	+	
189	-	












190	rac	
191	+	
192	-	
193	rac	
194	+	
195	-	
196	rac	
197	+	
198	-	
199	rac	
200	+	
201	-	
202	rac	
203	+	
204	-	
205	rac	
206	+	
207	-	
208	rac	
209	+	
210	-	
211	rac	
212	+	
213	-	







214	rac	
215	+	
216	-	
217	rac	
218	+	
219	-	
220	rac	
221	+	
222	-	
223	rac	
224	+	
225	-	
226	rac	
227	+	
228	-	
229	rac	
230	+	
231	-	
232	rac	
233	+	
234	-	
235	rac	
236	+	
237	-	









238	rac	
239	+	
240	-	


No.	Racemic or Enantiomer	R3
241	rac	
242	+	
243	-	
244	rac	
245	+	
246	-	
247	rac	
248	+	
249	-	
250	rac	
251	+	
252	-	
253	rac	
254	+	
255	-	
256	rac	















257	+	
258	-	
259	rac	
260	+	
261	-	
262	rac	
263	+	
264	-	
265	rac	
266	+	
267	-	
268	rac	
269	+	
270	-	
271	rac	
272	+	
273	-	
274	rac	
275	+	
276	-	
277	rac	
278	+	
279	-	
280	rac	


281	+	
282	-	
283	rac	
284	+	
285	-	
286	rac	
287	+	
288	-	
289	rac	
290	+	
291	-	
292	rac	
293	+	
294	-	
295	rac	
296	+	
297	-	
298	rac	
299	+	
300	-	


No.	Racemic or Enantiomer	R3
301	rac	
302	+	
303	-	
304	rac	
305	+	
306	-	
307	rac	
308	+	
309	-	
310	rac	
311	+	
312	-	
313	rac	
314	+	
315	-	
316	rac	
317	+	
318	-	
319	rac	
320	+	
321	-	
322	rac	


















323	+	
324	-	
325	rac	
326	+	
327	-	
328	rac	
329	+	
330	-	
331	rac	
332	+	
333	-	
334	rac	
335	+	
336	-	
337	rac	
338	+	
339	-	
340	rac	
341	+	
342	-	
343	rac	
344	+	




345	-	
346	rac	
347	+	
348	-	
349	rac	
350	+	
351	-	
352	rac	
353	+	
354	-	
355	rac	
356	+	
357	-	
358	rac	
359	+	
360	-	


No.	Racemic or Enantiomer	R3
361	rac	
362	+	
363	-	












364	rac	
365	+	
366	-	
367	rac	
368	+	
369	-	
370	rac	
371	+	
372	-	
373	rac	
374	+	
375	-	
376	rac	
377	+	
378	-	
		
379	rac	
380	+	
381	-	
382	rac	
383	+	
384	-	
385	rac	









386	+	
387	-	
388	rac	
389	+	
390	-	
391	rac	
392	+	
393	-	
394	rac	
395	+	
396	-	
397	rac	
398	+	
399	-	
400	rac	
401	+	
402	-	
403	rac	
404	+	
405	-	
406	rac	
407	+	
408	-	
409	rac	


410	+	
411	-	
412	rac	
413	+	
414	-	
415	rac	
416	+	
417	-	
418	rac	
419	+	
420	-	




No.	Racemic or Enantiomer	R3
421	rac	
422	+	
423	-	
424	rac	
425	+	
426	-	
427	rac	
428	+	
429	-	












430	rac	
431	+	
432	-	
433	rac	
434	+	
435	-	
436	rac	
437	+	
438	-	
439	rac	
440	+	
441	-	
442	rac	
443	+	
444	-	
445	rac	
446	+	
447	-	
448	rac	
449	+	
450	-	
451	rac	
452	+	
453	-	








454	rac	
455	+	
456	-	
457	rac	
458	+	
459	-	
460	rac	
461	+	
462	-	
463	rac	
464	+	
465	-	
466	rac	
467	+	
468	-	
469	rac	
470	+	
471	-	
472	rac	
473	+	
474	-	
475	rac	
476	+	
477	-	









		
478	rac	
479	+	
480	-	


No.	Racemic or Enantiomer	R3
481	rac	
482	+	
483	-	
484	rac	
485	+	
486	-	
487	rac	
488	+	
489	-	
490	rac	
491	+	
492	-	
493	rac	
494	+	















495	-	
496	rac	
497	+	
498	-	
499	rac	
500	+	
501	-	
502	rac	
503	+	
504	-	
505	rac	
506	+	
507	-	
508	rac	
509	+	
510	-	
511	rac	
512	+	
513	-	
214	rac	
515	+	
516	-	
517	rac	
518	+	



519	-	
520	rac	
521	+	
522	-	
523	rac	
524	+	
525	-	
526	rac	
227	+	
528	-	
529	rac	
530	+	
531	-	
532	rac	
533	+	
534	-	
535	rac	
536	+	
537	-	
538	rac	
539	+	
540	-	


No.	Racemic or Enantiomer	R3
541	rac	
542	+	
543	-	
544	rac	
545	+	
546	-	
547	rac	
548	+	
549	-	
550	rac	
551	+	
552	-	
553	rac	
554	+	
555	-	
556	rac	
557	+	
558	-	
559	rac	
560	+	
561	-	
562	rac	


















563	+	
564	-	
565	rac	
566	+	
567	-	
568	rac	
569	+	
570	-	
571	rac	
572	+	
573	-	
574	rac	
575	+	
576	-	
577	rac	
578	+	
579	-	
580	rac	
581	+	
582	-	
583	rac	
584	+	
585	-	
586	rac	




587	+	
588	-	
589	rac	
590	+	
591	-	
592	rac	
593	+	
594	-	
595	rac	
596	+	
597	-	
598	rac	
599	+	
600	-	





No.	Racemic or Enantiomer	R3
601	rac	
602	+	
603	-	
604	rac	
605	+	












606	-	
607	rac	
608	+	
609	-	
610	rac	
611	+	
612	-	
613	rac	
614	+	
615	-	
616	rac	
617	+	
618	-	
		
619	rac	
620	+	
621	-	
622	rac	
623	+	
624	-	
625	rac	
626	+	
627	-	








628	rac	
629	+	
630	-	
631	rac	
632	+	
633	-	
634	rac	
635	+	
636	-	
637	rac	
638	+	
639	-	
640	rac	
641	+	
642	-	
643	rac	
644	+	
645	-	
646	rac	
647	+	
648	-	
649	rac	
650	+	
651	-	


652	rac	
653	+	
654	-	
655	rac	
656	+	
657	-	
658	rac	
659	+	
660	-	






No.	Racemic or Enantiomer	R3
661	rac	
662	+	
663	-	
664	rac	
665	+	
666	-	
667	rac	
668	+	
669	-	
670	rac	












671	+	
672	-	
673	rac	
674	+	
675	-	
676	rac	
677	+	
678	-	
679	rac	
680	+	
681	-	
682	rac	
683	+	
684	-	
685	rac	
686	+	
687	-	
688	rac	
689	+	
690	-	
691	rac	
692	+	
693	-	
694	rac	








695	+	
696	-	
697	rac	
698	+	
699	-	
700	rac	
701	+	
702	-	
703	rac	
704	+	
705	-	
706	rac	
707	+	
708	-	
709	rac	
710	+	
711	-	
712	rac	
713	+	
714	-	
715	rac	
716	+	







717	-	
718	rac	
719	+	
720	-	


No.	Racemic or Enantiomer	R3
721	rac	
722	+	
723	-	
724	rac	
725	+	
726	-	
727	rac	
728	+	
729	-	
730	rac	
731	+	
732	-	
733	rac	
734	+	
735	-	
















736	rac	
737	+	
738	-	
739	rac	
740	+	
741	-	
742	rac	
743	+	
744	-	
745	rac	
746	+	
747	-	
748	rac	
749	+	
750	-	
751	rac	
752	+	
753	-	
754	rac	
755	+	
756	-	
757	rac	
758	+	
759	-	


760	rac	
761	+	
762	-	
763	rac	
764	+	
765	-	
766	rac	
767	+	
768	-	
769	rac	
770	+	
771	-	
772	rac	
773	+	
774	-	
775	rac	
776	+	
777	-	
778	rac	
779	+	
780	-	


No.	Racemic or Enantiomer	R3
781	rac	
782	+	
783	-	
784	rac	
785	+	
786	-	
787	rac	
788	+	
789	-	
790	rac	
791	+	
792	-	
793	rac	
794	+	
795	-	
796	rac	
797	+	
798	-	
799	rac	
800	+	
801	-	


















802	rac	
803	+	
804	-	
805	rac	
806	+	
807	-	
808	rac	
809	+	
810	-	
811	rac	
812	+	
813	-	
814	rac	
815	+	
816	-	
817	rac	
818	+	
819	-	
820	rac	
821	+	
822	-	
823	rac	
824	+	
825	-	




826	rac	
827	+	
828	-	
829	rac	
830	+	
831	-	
832	rac	
833	+	
834	-	
835	rac	
836	+	
837	-	
838	rac	
839	+	
840	-	





No.	Racemic or Enantiomer	R3
841	rac	
842	+	
843	-	
844	rac	



















845	+	
846	-	
847	rac	
848	+	
849	-	
850	rac	
851	+	
852	-	
853	rac	
854	+	
855	-	
856	rac	
857	+	
858	-	
859	rac	
860	+	
861	-	
862	rac	
863	+	
864	-	
865	rac	
866	+	
867	-	

868	rac	
869	+	
870	-	
871	rac	
872	+	
873	-	
874	rac	
875	+	
876	-	
877	rac	
878	+	
879	-	
880	rac	
881	+	
882	-	
883	rac	
884	+	
885	-	
886	rac	
887	+	
888	-	
889	rac	
890	+	
891	-	







892	rac	
893	+	
894	-	
895	rac	
896	+	
897	-	
898	rac	
899	+	
900	-	


No.	Racemic or Enantiomer	R3
901	rac	
902	+	
903	-	
904	rac	
905	+	
906	-	
907	rac	
908	+	
909	-	
910	rac	








911	+	
912	-	
913	rac	
914	+	
915	-	
916	rac	
917	+	
918	-	
919	rac	
920	+	
921	-	
922	rac	
923	+	
924	-	
925	rac	
926	+	
927	-	
928	rac	
929	+	
930	-	
931	rac	
932	+	
933	-	
934	rac	








935	+	
936	-	
937	rac	
938	+	
939	-	
940	rac	
941	+	
942	-	
943	rac	
944	+	
945	-	
946	rac	
947	+	
948	-	
949	rac	
950	+	
951	-	
952	rac	
953	+	
954	-	
955	rac	
956	+	
957	-	
958	rac	






959	+	
960	-	


No.	Racemic or Enantiomer	R3
961	rac	
962	+	
963	-	
964	rac	
965	+	
966	-	
967	rac	
968	+	
969	-	
970	rac	
971	+	
972	-	
973	rac	
974	+	
975	-	
976	rac	
977	+	















978	-	
979	rac	
980	+	
981	-	
982	rac	
983	+	
984	-	
985	rac	
986	+	
987	-	
988	rac	
989	+	
990	-	
991	rac	
992	+	
993	-	
994	rac	
995	+	
996	-	
997	rac	
998	+	
999	-	


1000	rac	
1001	+	
1002	-	
1003	rac	
1004	+	
1005	-	
1006	rac	
1007	+	
1008	-	
1009	rac	
1010	+	
1011	-	
1012	rac	
1013	+	
1014	-	
1015	rac	
1016	+	
1017	-	
1018	rac	
1019	+	
1020	-	


No.	Racemic or Enantiomer	R3
1021	rac	
1022	+	
1023	-	
1024	rac	
1025	+	
1026	-	
1027	rac	
1028	+	
1029	-	
1030	rac	
1031	+	
1032	-	
1033	rac	
1034	+	
1035	-	
1036	rac	
1037	+	
1038	-	
1039	rac	
1040	+	
1041	-	

















1042	rac	
1043	+	
1044	-	
1045	rac	
1046	+	
1047	-	
1048	rac	
1049	+	
1050	-	
1051	rac	
1052	+	
1053	-	
1054	rac	
1055	+	
1056	-	
1057	rac	
1058	+	
1059	-	
1060	rac	
1061	+	
1062	-	
1063	rac	
1064	+	




1065	-	
1066	rac	
1067	+	
1068	-	
1069	rac	
1070	+	
1071	-	
1072	rac	
1073	+	
1074	-	
1075	rac	
1076	+	
1077	-	
1078	rac	
1079	+	
1080	-	




No.	Racemic or Enantiomer	R3
1081	rac	
1082	+	
1083	-	












1084	rac	
1085	+	
1086	-	
1087	rac	
1088	+	
1089	-	
1090	rac	
1091	+	
1092	-	
1093	rac	
1094	+	
1095	-	
1096	rac	
1097	+	
1098	-	
1099	rac	
1100	+	
1101	-	
1102	rac	
1103	+	
1104	-	
1105	rac	
1106	+	
1107	-	









1108	rac	
1109	+	
1110	-	
1111	rac	
1112	+	
1113	-	
1114	rac	
1115	+	
1116	-	
1117	rac	
1118	+	
1119	-	
1120	rac	
1121	+	
1122	-	
1123	rac	
1124	+	
1125	-	
1126	rac	
1127	+	
1128	-	
1129	rac	
1130	+	
1131	-	

1132	rac	
1133	+	
1134	-	
1135	rac	
1136	+	
1137	-	
1138	rac	
1139	+	
1140	-	




No.	Racemic or Enantiomer	R3
1141	rac	
1142	+	
1143	-	
1144	rac	
1145	+	
1146	-	
1147	rac	
1148	+	
1149	-	
1150	rac	
1151	+	












1152	-	
1153	rac	
1154	+	
1155	-	
1156	rac	
1157	+	
1158	-	
1159	rac	
1160	+	
1161	-	
1162	rac	
1163	+	
1164	-	
1165	rac	
1166	+	
1167	-	
1168	rac	
1169	+	
1170	-	
1171	rac	
1172	+	
1173	-	
1174	rac	
1175	+	







1176	-	
1177	rac	
1178	+	
1179	-	
1180	rac	
1181	+	
1182	-	
1183	rac	
1184	+	
1185	-	
1186	rac	
1187	+	
1188	-	
1189	rac	
1190	+	
1191	-	
1192	rac	
1193	+	
1194	-	
1195	rac	
1196	+	
1197	-	
1198	rac	
1199	+	








1200	-	
------	---	--


No.	Racemic or Enantiomer	R3
1201	rac	
1202	+	
1203	-	
1204	rac	
1205	+	
1206	-	
1207	rac	
1208	+	
1209	-	
1210	rac	
1211	+	
1212	-	
1213	rac	
1214	+	
1215	-	
1216	rac	
1217	+	
1218	-	
















1219	rac	
1220	+	
1221	-	
1222	rac	
1223	+	
1224	-	
1225	rac	
1226	+	
1227	-	
1228	rac	
1229	+	
1230	-	
1231	rac	
1232	+	
1233	-	
1234	rac	
1235	+	
1236	-	
1237	rac	
1238	+	
1239	-	
1240	rac	


1241	+	
1242	-	
1243	rac	
1244	+	
1245	-	
1246	rac	
1247	+	
1248	-	
1249	rac	
1250	+	
1251	-	
1252	rac	
1253	+	
1254	-	
1255	rac	
1256	+	
1257	-	
1258	rac	
1259	+	
1260	-	


No.	Racemic or Enantiomer	R3
1261	rac	
1262	+	
1263	-	
1264	rac	
1265	+	
1266	-	
1267	rac	
1268	+	
1269	-	
1270	rac	
1271	+	
1272	-	
1273	rac	
1274	+	
1275	-	
1276	rac	
1277	+	
1278	-	
1279	rac	
1280	+	
1281	-	


















		
1282	rac	
1283	+	
1284	-	
1285	rac	
1286	+	
1287	-	
1288	rac	
1289	+	
1290	-	
1291	rac	
1292	+	
1293	-	
1294	rac	
1295	+	
1296	-	
1297	rac	
1298	+	
1299	-	
1300	rac	
1301	+	
1302	-	
1303	rac	





1304	+	
1305	-	
1306	rac	
1307	+	
1308	-	
1309	rac	
1310	+	
1311	-	
1312	rac	
1313	+	
1314	-	
1315	rac	
1316	+	
1317	-	
1318	rac	
1319	+	
1320	-	



No.	Racemic or Enantiomer	R3
1321	rac	
1322	+	












1323	-	
1324	rac	
1325	+	
1326	-	
1327	rac	
1328	+	
1329	-	
1330	rac	
1331	+	
1332	-	
1333	rac	
1334	+	
1335	-	
1336	rac	
1337	+	
1338	-	
1339	rac	
1340	+	
1341	-	
1342	rac	
1343	+	
1344	-	








1345	rac	
1346	+	
1347	-	
1348	rac	
1349	+	
1350	-	
1351	rac	
1352	+	
1353	-	
1354	rac	
1355	+	
1356	-	
1357	rac	
1358	+	
1359	-	
1360	rac	
1361	+	
1362	-	
1363	rac	
1364	+	
1365	-	
1366	rac	
1367	+	
1368	-	




1369	rac	
1370	+	
1371	-	
1372	rac	
1373	+	
1374	-	
1375	rac	
1376	+	
1377	-	
1378	rac	
1379	+	
1380	-	



No.	Racemic or Enantiomer	R3
1381	rac	
1382	+	
1383	-	
1384	rac	
1385	+	










1386	-	
1387	rac	
1388	+	
1389	-	
1390	rac	
1391	+	
1392	-	
1393	rac	
1394	+	
1395	-	
1396	rac	
1397	+	
1398	-	
1399	rac	
1400	+	
1401	-	
1402	rac	
1403	+	
1404	-	
1405	rac	
1406	+	
1407	-	
1408	rac	
1409	+	









1410	-	
1411	rac	
1412	+	
1413	-	
1414	rac	
1415	+	
1416	-	
1417	rac	
1418	+	
1419	-	
1420	rac	
1421	+	
1422	-	
1423	rac	
1424	+	
1425	-	
1426	rac	
1427	+	
1428	-	
1429	rac	
1430	+	
1431	-	


1432	rac	
1433	+	
1434	-	
1435	rac	
1436	+	
1437	-	
1438	rac	
1439	+	
1440	-	




No.	Racemic or Enantiomer	R3
1441	rac	
1442	+	
1443	-	
1444	rac	
1445	+	
1446	-	
1447	rac	
1448	+	
1449	-	
1450	rac	












1451	+	
1452	-	
1453	rac	
1454	+	
1455	-	
1456	rac	
1457	+	
1458	-	
1459	rac	
1460	+	
1461	-	
1462	rac	
1463	+	
1464	-	
1465	rac	
1466	+	
1467	-	
1468	rac	
1469	+	
1470	-	
1471	rac	
1472	+	
1473	-	








1474	rac	
1475	+	
1476	-	
1477	rac	
1478	+	
1479	-	
1480	rac	
1481	+	
1482	-	
1483	rac	
1484	+	
1485	-	
1486	rac	
1487	+	
1488	-	
1489	rac	
1490	+	
1491	-	
1492	rac	
1493	+	
1494	-	
1495	rac	
1496	+	
1497	-	






1498	rac	
1499	+	
1500	-	


No.	Racemic or Enantiomer	R3
1501	rac	
1502	+	
1503	-	
1504	rac	
1505	+	
1506	-	
1507	rac	
1508	+	
1509	-	
1510	rac	
1511	+	
1512	-	
1513	rac	
1514	+	















1515	-	
1516	rac	
1517	+	
1518	-	
1519	rac	
1520	+	
1521	-	
1522	rac	
1523	+	
1524	-	
1525	rac	
1526	+	
1527	-	
1528	rac	
1529	+	
1530	-	
1531	rac	
1532	+	
1533	-	
1534	rac	
1535	+	
1536	-	
1537	rac	
1538	+	

1539	-	
1540	rac	
1541	+	
1542	-	
1543	rac	
1544	+	
1545	-	
1546	rac	
1547	+	
1548	-	
1549	rac	
1550	+	
1551	-	
1552	rac	
1553	+	
1554	-	
1555	rac	
1556	+	
1557	-	
1558	rac	
1559	+	
1560	-	

No.	Racemic or Enantiomer	R3
1561	rac	
1562	+	
1563	-	
1564	rac	
1565	+	
1566	-	
1567	rac	
1568	+	
1569	-	
1570	rac	
1571	+	
1572	-	
1573	rac	
1574	+	
1575	-	
1576	rac	
1577	+	
1578	-	
1579	rac	
1580	+	
1581	-	
1582	rac	



1583	+	
1584	-	
1585	rac	
1586	+	
1587	-	
1588	rac	
1589	+	
1590	-	
1591	rac	
1592	+	
1593	-	
1594	rac	
1595	+	
1596	-	
1597	rac	
1598	+	
1599	-	
1600	rac	
1601	+	
1602	-	
1603	rac	
1604	+	
1605	-	
1606	rac	

1607	+	
1608	-	
1609	rac	
1610	+	
1611	-	
1612	rac	
1613	+	
1614	-	
1615	rac	
1616	+	
1617	-	
1618	rac	
1619	+	
1620	-	

14. Pharmaceutical composition that contains at least one compound of general formula I according to one of claims 1 to 13 and optionally at least one additional active ingredient together with pharmaceutically compatible adjuvants and/or vehicles.

15. Pharmaceutical composition according to claim 14, wherein the additional active ingredient is a SERM (selective estrogen receptor modulator), an aromatase inhibitor, an antiestrogen, or a prostaglandin.

16. Pharmaceutical composition according to claim 14, whereby the active

ingredients can be tamoxifen, 5-(4-{5-[(RS)-(4,4,5,5,5-pentafluoropentyl)sulfinyl]-pentyloxy}phenyl)-6-phenyl-8,9-dihydro-7H-benzocyclohepten-2-ol, ICI 182 780 (7alpha-[9-(4,4,5,5-pentafluoropentyl)sulfinyl]nonyl]estra-1,3,5(10)-triene-3,17-beta-diol), 11beta-fluoro-7alpha-[5-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17beta-diol, 11beta-fluoro-7alpha-{5-[methyl(7,7,8,8,9,9,10,10,10-nonafluorodecyl)amino]pentyl}estra-1,3,5(10)-triene-3,17beta-diol, 11beta-fluoro-17alpha-methyl-7alpha-{5-[methyl(8,8,9,9,9-pentafluorononyl)amino]pentyl}estra-1,3,5(10)-triene-3,17beta-diol, clomifene, raloxifene, fadrozole, formestane, letrozole, anastrozole or atamestane.

17. Use of compounds according to one of claims 1 to 13 for the production of a pharmaceutical agent.

18. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of gynecological diseases such as endometriosis, leiomyomas of the uterus, dysfunctional bleeding and dysmenorrhea.

19. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of hormone-dependent tumors.

20. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of breast cancer.

21. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of endometrial carcinoma.

22. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of ovarian cancer.

23. Use of compounds according to claim 17 for the production of a pharmaceutical agent for therapy and prophylaxis of prostate cancer.

24. Use of compounds according to claim 17 for the production of a pharmaceutical agent for female hormone replacement therapy.
25. Use of compounds according to claim 17 for female birth control.
26. Process for the selective addition of lithium alkynyl compounds and magnesium haloalkynyl compounds to ketoamide.