

table providing the SEQ ID NO for each VL and CDR sequence illustrated in FIG. 2C. FIG. 2E shows a sequence alignment of example anti-BCMA binding molecules comprising a VH domain. Complementarity determining regions (CDRs) and framework regions (FRs) are shown, as determined by Chothia numbering (FIG. 2E). SEQ ID NOS 253-260, CDR1 in order from top to bottom; SEQ ID NOS 261-268, CDR2 from top to bottom, SEQ ID NOS 269-276, CDR3 from top to bottom. FIG. 2F is a table providing the SEQ ID NO for each VH and CDR illustrated in FIG. 2E. FIG. 2G shows a sequence alignment of example anti-BCMA binding molecules comprising a VL domain. CDRs and FRs are shown, as determined by Chothia numbering (FIG. 2G). FIG. 2H is a table providing the SEQ ID NO for each VL and CDR sequence illustrated in FIG. 2G. SEQ ID NOS 37-44, CDR1 in order from top to bottom; SEQ ID NOS 45-52, CDR2 from top to bottom, SEQ ID NOS 277-284, CDR3 from top to bottom.

[0056] FIG. 3 depicts CAR expression in primary human T cells electroporated with mRNA encoding for various CARs. Data obtained from CAR having a complete hinge domain (“CHD”) is shown and data obtained from CAR having a truncated hinge domain (“THD”) is shown.

[0057] FIGs. 4A-4X show IFN γ , IL-2, and TNF α production by electroporated anti-FLT3 CAR T cells following 16 hours of co-culture with the indicated target cell lines. FIGs. 4A-4B, 4G-4H, 4M-4N, and 4S-4T show IFN γ production following co-culture with Namalwa, EoL-1, HL60, and MV4;11 target cells, respectively. FIGs. 4C-4D, 4I-4J, 4O-4P, and 4U-4V show IL-2 production following co-culture with Namalwa, EoL-1, HL60, and MV4;11 target cells, respectively. FIGs. 4E-4F, 4K-4L, 4Q-4R, and 4W-4X show TNF α production following co-culture with Namalwa, EoL-1, HL60, and MV4;11 target cells, respectively.

[0058] FIGs. 5A-5H show cytolytic activity of electroporated anti-FLT3 CAR T cells against Namalwa (FIGs. 5A-5B), EoL1 (FIGs. 5C-5D), HL60 (FIGs. 5E-5F), and MV4;11 (FIGs. 5G-5H) target cell lines following 16 hours of co-culture.

[0059] FIGs. 6A-6B depict CAR expression in lentivirus transduced primary human T cells from two healthy donors.

[0060] FIGs. 7A-7F show IFN γ (FIGs. 7A-7B), TNF α (FIGs. 7C-7D), and IL-2 (FIGs. 7E-7F) production by lentivirus transduced CAR T cells from two healthy donors following 16 hours of co-culture with the indicated target cell lines