

Novel carboxylic acid derivatives with amide side chains, their preparation and use as endothelin receptor antagonists

- 5 The present invention relates to novel carboxylic acid derivatives, their preparation and use

Endothelin is a peptide which is composed of 21 amino acids and is synthesized and released by vascular endothelium. Endothelin
10 exists in three isoforms, ET-1, ET-2 and ET-3. "Endothelin" or "ET" hereinafter refers to one or all isoforms of endothelin. Endothelin is a potent vasoconstrictor and has a strong effect on vessel tone. It is known that this vasoconstriction is caused by binding of endothelin to its receptor (Nature, 332, 411-415,
15 1988; FEBS Letters, 231, 440-444, 1988 and Biochem. Biophys. Res. Commun., 154, 868-875, 1988).

Elevated or abnormal release of endothelin causes persistent vasoconstriction in peripheral, renal and cerebral blood vessels,
20 which may result in disorders. As reported in the literature, endothelin is involved in a number of disorders. These include: hypertension, acute myocardial infarct, pulmonary hypertension, Raynaud's syndrome, cerebral vasospasms, stroke, benign prostate hypertrophy, atherosclerosis and asthma (J. Vascular Med. Biology
25 2, 207 (1990), J. Am. Med. Association 264, 2868 (1990), Nature 344, 114 (1990), N. Engl. J. Med. 322, 205 (1989), N. Engl. J. Med. 328, 1732 (1993), Nephron 66, 373 (1994), Stroke 25, 904 (1994), Nature 365, 759 (1993), J. Mol. Cell. Cardiol. 27, A234 (1995); Cancer Research 56, 663 (1996)).


30

At least two endothelin receptor subtypes, ET_A and ET_B receptors, are currently described in the literature (Nature 348, 730 (1990), Nature 348, 732 (1990)). Accordingly, substances which inhibit the binding of endothelin to the two receptors ought to
35 antagonize the physiological effects of endothelin and therefore represent valuable drugs.

Mixed ET_A/ET_B receptor antagonists have been described in DE Patent Application 19636046.3. The spacer Q (see formula II) is
40 important for these compounds, corresponds in length to a C₂-C₄-alkyl chain and has the function of producing a defined distance between R⁶ and W in compounds of the formula II.


45

5



Furthermore, in the patent application WO 97/38980, the following compounds of the formula VII are described as endothelin receptor antagonists:

15



20 An advantage of these compounds is said to be the low plasma binding.

We have now found, surprisingly, that the receptor affinity and selectivity can be influenced using the spacer $\text{Q} = \text{R}^6\text{CR}^7\text{R}^8$ (see 25 formula I), as a function of $\text{R}^6 = \text{amide}$. Thus, either ET_A -selective, ET_B -selective or else mixed receptor antagonists [sic] can be prepared.

The antagonists referred to here as ET_A (ET_B)-specific antagonists 30 are those whose affinity for the ET_A (ET_B) receptor is at least ten times higher than their affinity for the ET_B (ET_A) receptor. Preferred compounds are those whose difference in affinity for the two receptors is at least twenty.

35 Mixed endothelin receptor antagonists are those compounds which bind with approximately the same affinity to the ET_A and ET_B receptors. Approximately the same affinity for receptors exists when the ratio of the affinities is greater than 0.05 (preferably 0.1) and less than 20 (preferably 10).


40

It is an object of the present invention to identify compounds which belong to one of the three selectivity groups.

45

3

We have found that this object is achieved by carboxylic acid derivatives of the formula I



10

[lacuna] R¹ is tetrazole [sic] or a group



15 where R has the following meaning:

a) a radical OR^9 where R^9 is:

hydrogen, the cation of an alkali metal, the cation of an alkaline earth metal, a physiologically tolerated organic ammonium ion such as tertiary C₁-C₄-alkylammonium or the ammonium ion;

25 C₃-C₈-cycloalkyl, C₁-C₈-alkyl, CH₂-phenyl, each of which can be substituted by one or more of the following radicals:
halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto, C₁-C₄-alkylthio, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂;

30 a C₃-C₆-alkenyl - or a C₃-C₆-alkynyl group, it being possible
for these groups in turn to carry one to five halogen atoms;


35 R⁹ can also be a phenyl radical which may carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, mercapto, C₁-C₄-alkylthio, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂;

b) a 5-membered heteroaromatic system which is linked via a nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and triazolyl, which may carry one or two halogen atoms, or one or two C₁-C₄-alkyl or one or two C₁-C₄-alkoxy groups.

c) a group

45

4



5

where k can assume the values 0, 1 and 2, p can assume the values 1, 2, 3 and 4, and R¹⁰ is

10 C₁-C₄-alkyl, C₃-C₈-cycloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or phenyl, which can be substituted by one or more, eg. one to three, of the following radicals:

15 halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, mercapto, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂.

d) a radical



where R¹¹ is:

25

C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl, it being possible for these radicals to carry a C₁-C₄-alkoxy, C₁-C₄-alkylthio and/or a phenyl radical as specified under c);

30

phenyl, which can be substituted by one to three of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, mercapto, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂

35

The other substituents have the following meanings:

R² hydrogen, hydroxyl, NH₂, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-hydroxyalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, or CR² is linked to CR¹⁰ as indicated under Z to give a 5- or 6-membered ring.

X nitrogen or methine.

45

Y nitrogen or methine.

5

- Z nitrogen or CR^{12} , where R^{12} is hydrogen, halogen or C_1 - C_4 -alkyl, or CR^{12} forms together with CR^2 or CR^3 a 5- or 6-membered alkylene or alkenylene ring which can be substituted by one or two C_1 - C_4 -alkyl groups and in which in each case one or more methylene groups can be replaced by oxygen, sulfur, $-NH$ or $N(C_1$ - C_4 -alkyl).

At least one of the ring members X, Y or Z is nitrogen.

- 10 R^3 hydrogen, hydroxyl, NH_2 , $NH(C_1$ - C_4 -alkyl), $N(C_1$ - C_4 -alkyl) $_2$, halogen, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -hydroxyalkyl, C_1 - C_4 -alkylthio, or CR^3 is linked to CR^{12} as indicated under Z to give a 5- or 6-membered ring.

15

- R^4 and R^5 (which can be identical or different):

phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, mercapto, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, phenoxy, carboxyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, amino, $NH(C_1$ - C_4 -alkyl), $N(C_1$ - C_4 -alkyl) $_2$ or phenyl which can be substituted one or more times, eg. one to three times, by halogen, nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy or C_1 - C_4 -alkylthio; or


25

phenyl or naphthyl which are connected together in ortho positions by a direct linkage, a methylene, ethylene, or ethenylene group, an oxygen or sulfur atom or an SO_2 , NH or N -alkyl group;


30

C_3 - C_8 -cycloalkyl.

- 35 R^6 a group



or



40

- R^{13} and R^{14} (which can be identical or different):

hydrogen, with the proviso that R^{13} and R^{14} must not simultaneously be hydrogen,

45

C_1 - C_8 -alkyl, C_3 - C_8 -alkenyl or C_3 - C_8 -alkynyl, it being possible for each of these radicals to be substituted one or more

- times by: halogen, hydroxyl, mercapto, carboxyl, nitro, amino, carboxamide [sic], cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl,
- 5 C₃-C₈-alkylcarbonylalkyl, C₃-C₈-cycloalkyl, 1-indanyl, 2-indanyl, 1-tetrahydronaphthyl, 2-tetrahydronaphthyl, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, phenoxy or phenyl, it being possible for said aryl radicals in turn to be substituted one or more times, eg. one to three times, by halogen, hydroxyl,
- 10 mercapto, carboxyl, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, or C₁-C₄-alkylthio;
- C₃-C₈-cycloalkyl, it being possible for each of these radicals
- 15 to be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, nitro, cyano, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy;
- phenyl or naphthyl, each of which can be substituted by one
- 20 or more of the following radicals: halogen, nitro, carboxamide [sic], mercapto, carboxyl, cyano, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-alkenyloxy, C₁-C₄-haloalkyl, C₃-C₆-alkynyloxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy,
- 25 C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, dioxomethylene [sic], dioxoethylene [sic] or phenyl, which can be substituted one or more times, eg. one to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or
- 30 C₁-C₄-alkylthio;
- or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain which is closed to a ring and which can be substituted one or more times by C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-alkoxy,
- 35 C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, and in which one alkylene [sic] group can be replaced by oxygen, sulfur, nitrogen or N(C₁-C₄-alkyl), such as -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-, -(CH₂)₂-S-(CH₂)₂-, -(CH₂)₂-NH-(CH₂)₂-, -(CH₂)₃-, -(CH₂)₂-N(CH₃)-(CH₂)₂-;
- 40 or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain or C₄-C₇-alkenylene chain, which is closed to a ring and each of which can be substituted one to three times by C₁-C₄-alkyl and to each of which is fused a phenyl ring which can be
- 45 substituted one or more times by halogen, C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, hydroxyl, carboxyl, amino, carboxamide

[sic].

R⁷ and R⁸ (which can be identical or different):

5 hydrogen, C₁-C₄-alkyl.

R¹⁵ C₁-C₄-alkyl, C₁-C₄-alkylthio, C₁-C₄-alkoxy, each of which
 carry one of the following radicals: hydroxyl, carboxyl,
 amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, carboxamide [sic] or
 10 CON(C₁-C₄-alkyl)₂.

R¹⁸ hydrogen;

C₁-C₈-alkyl, C₃-C₈-alkenyl or C₃-C₈-alkynyl, where each of
 15 these radicals can be substituted one or more times by:
 halogen, carboxyl, cyano, C₁-C₄-alkoxy, C₁-C₄-alkylthio,
 C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl,
 C₃-C₈-cycloalkyl, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂,
 phenoxy or phenyl, where the abovementioned aryl radicals for
 20 their part can be substituted one or more times, for example
 one to three times by halogen, hydroxyl, mercapto, carboxyl,
 nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy,
 C₁-C₄-haloalkoxy, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, or
 C₁-C₄-alkylthio;

25 C₃-C₈-cycloalkyl, where each of these radicals can be
 substituted one or more times by: halogen, C₁-C₄-alkyl;

phenyl or naphthyl, each of which can be substituted by one
 30 or more of the following radicals: halogen, nitro, mercapto,
 carboxyl, cyano, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl,
 C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl,
 C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy,
 C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, NH(C₁-C₄-alkyl),
 35 N(C₁-C₄-alkyl)₂, dioxomethylene [sic], dioxoethylene [sic] or
 phenyl which can be substituted one or more times, for
 example one to three times by halogen, nitro, cyano,
 C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or
 C₁-C₄-alkylthio;

40 C₁-C₈-alkylcarbonyl, C₂-C₈-alkenylcarbonyl or C₂-C₈-alkynyl-
 carbonyl, where each of these radicals can be substituted one
 or more times by: halogen, C₁-C₄-alkoxy, C₁-C₄-alkylthio,
 C₁-C₄-haloalkoxy, C₁-C₄-alkoxycarbonyl, C₃-C₈-cycloalkyl,
 45 phenoxy or phenyl, where the abovementioned aryl radicals for
 their part can be substituted one or more times, for example
 one to three times by halogen, hydroxyl, mercapto, carboxyl,

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, or C₁-C₄-alkylthio;

- 5 benzyloxycarbonyl, C₃-C₈-cycloalkylcarbonyl, where each of these radicals can be substituted one or more times by: halogen, C₁-C₄-alkyl;

- 10 phenylcarbonyl or naphthylcarbonyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, mercapto, carboxyl, cyano, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-alkenyl-oxy, C₁-C₄-haloalkyl, C₃-C₆-alkynyloxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, 15 C₁-C₄-alkylthio, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, dioxo-methylene [sic], dioxoethylene [sic] or phenyl which can be substituted one or more times, for example one to three times by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

- 20 C₁-C₈-alkylsulfonyl, C₃-C₈-alkenylsulfonyl or C₃-C₈-alkynyl-sulfonyl, where each of these radicals can be substituted one or more times by: halogen, C₁-C₄-alkoxy, phenyl, where the abovementioned aryl radical for its part can be substituted 25 one or more times, for example one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, or C₁-C₄-alkylthio;

- 30 C₃-C₈-cycloalkylsulfonyl;

- 35 phenylsulfonyl or naphthylsulfonyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-alkenyloxy, C₁-C₄-halo-alkyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted one or more times, for example one to three 40 times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio.

- 40 R²⁰ hydrogen;

- 45 C₁-C₄-alkyl, where each of these radicals can be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, amino, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkoxycarbonyl, C₃-C₈-cycloalkyl, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, indolyl, phenoxy or phenyl, where the

- abovementioned aryl radicals for their part can be substituted one or more times, for example one to three times by halogen, hydroxyl, mercapto, carboxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, amino,
- 5 NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂ or C₁-C₄-alkylthio.

R²¹ hydrogen, C₁-C₄-alkyl.

W sulfur or oxygen.

10

In these cases and hereinafter, the following definitions apply:

An alkali metal is, for example, lithium, sodium, potassium;

- 15 An alkaline earth metal is, for example, calcium, magnesium, barium;

C₃-C₈-cycloalkyl is, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl;

20

C₁-C₄-haloalkyl can be linear or branched such as fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl, dichlorofluoromethyl, trichloromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl,

25 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl or pentafluoroethyl;

C₁-C₄-haloalkoxy can be linear or branched such as difluoromethoxy, trifluoromethoxy, chlorodifluoromethoxy,

- 30 1-fluoroethoxy, 2,2-difluoroethoxy, 1,1,2,2-tetrafluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-1,1,2-trifluoroethoxy, 2-fluoroethoxy or pentafluoroethoxy;

C₁-C₄-alkyl can be linear or branched such as methyl, ethyl,

- 35 1-propyl, 2-propyl, 2-methyl-2-propyl, 2-methyl-1-propyl, 1-butyl or 2-butyl;

C₂-C₄-alkenyl can be linear or branched such as ethenyl,

1-propen-3-yl, 1-propen-2-yl, 1-propen-1-yl, 2-methyl-1-propenyl,

- 40 1-butenyl or 2-butenyl;

C₂-C₄-alkynyl can be linear or branched such as ethynyl,

1-propyn-1-yl, 1-propyn-3-yl, 1-butyne-4-yl or 2-butyne-4-yl;

- 45 C₁-C₄-alkoxy can be linear or branched such as methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy

or 1,1-dimethylethoxy;

C₃-C₆-alkenyloxy can be linear or branched such as allyloxy, 2-buten-1-yloxy or 3-buten-2-yloxy;

5

C₃-C₆-alkynyloxy can be linear or branched such as 2-propyn-1-yloxy, 2-butyn-1-yloxy or 3-butyn-2-yloxy;

C₁-C₄-alkylthio can be linear or branched such as methylthio, ethylthio, propylthio, 1-methylethylthio, butylthio, 1-methylpropylthio, 2-methylpropylthio or 1,1-dimethylethylthio;

10

C₁-C₄-alkylcarbonyl can be linear or branched such as acetyl, ethylcarbonyl or 2-propylcarbonyl, 1-propylcarbonyl, 1-butylcarbonyl;

15

C₁-C₄-alkoxycarbonyl can be linear or branched such as methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, i-propoxycarbonyl or n-butoxycarbonyl;

20

C₃-C₈-alkylcarbonylalkyl can be linear or branched such as 2-oxo-1-propyl, 3-oxo-1-butyl or 3-oxo-2-butyl

C₁-C₈-alkyl can be linear or branched such as C₁-C₄-alkyl, pentyl, hexyl, heptyl or octyl;

25

C₁-C₈-alkylcarbonyl can be linear or branched such as C₁-C₄-alkylcarbonyl, 1-pentylcarbonyl, 1-hexylcarbonyl, 1-heptylcarbonyl or 1-octylcarbonyl;

30

C₂-C₈-alkenylcarbonyl can be linear or branched such as ethenylcarbonyl, 1-propen-3-ylcarbonyl, 1-propen-2-ylcarbonyl, 1-propen-1-ylcarbonyl, 2-methyl-1-propenylcarbonyl, 1-buten-1-ylcarbonyl, 1-penten-1-ylcarbonyl, 1-octen-1-ylcarbonyl;

35

C₂-C₈-alkynylcarbonyl can be linear or branched such as ethynylcarbonyl, 1-propyn-3-ylcarbonyl, 1-propyn-1-ylcarbonyl, 1-butyn-1-ylcarbonyl, 1-pentyn-1-ylcarbonyl, 1-octyn-1-ylcarbonyl;

40

C₃-C₈-cycloalkylcarbonyl, cyclopropylcarbonyl, cyclobutylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, 4-methylcyclohex-1-ylcarbonyl, cycloheptylcarbonyl or cyclooctylcarbonyl;

45

C₁-C₄-alkylsulfonyl can be linear or branched such as methylsulfonyl, ethylsulfonyl or 2-propylsulfonyl, 1-propylsulfonyl, 2-methyl-1-propylsulfonyl, 1-butylsulfonyl;

- 5 C₁-C₈-alkylsulfonyl can be linear or branched such as C₁-C₄-alkylsulfonyl, 1-pentylsulfonyl, 1-hexylsulfonyl, 1-heptylsulfonyl or 1-octylsulfonyl;

- C₃-C₈-alkenylsulfonyl can be linear or branched such as
10 1-propen-3-ylsulfonyl, 1-propen-2-ylsulfonyl, 1-propen-1-ylsulfonyl, 2-methyl-1-propen-1-ylsulfonyl, 1-buten-1-ylsulfonyl, 1-penten-1-ylsulfonyl, 1-octen-1-ylsulfonyl;

- C₃-C₈-alkynylsulfonyl can be linear or branched such as
15 1-propyn-3-ylsulfonyl, 1-propyn-1-ylsulfonyl, 1-butyn-1-ylsulfonyl, 1-pentyn-1-ylsulfonyl, 1-octyn-1-ylsulfonyl;

- C₃-C₈-cycloalkylsulfonyl is, for example, cyclopropylsulfonyl, cyclobutylsulfonyl, cyclopentylsulfonyl, cyclohexylsulfonyl,
20 4-methylcyclohex-1-ylsulfonyl, cycloheptylsulfonyl or cyclooctylsulfonyl;

halogen is, for example, fluorine, chlorine, bromine, iodine.

- 25 The invention further relates to those compounds from which the compounds of the formula I can be liberated (called prodrugs).


- Preferred prodrugs are those for which the liberation takes place under conditions like those prevailing in certain compartments of
30 the body, eg. in the stomach, intestine, bloodstream, liver.

- The compounds I and the intermediates for preparing them, such as III, IV and V, may have one or more asymmetrically substituted carbon atoms. Compounds of this type can exist as pure
35 enantiomers or pure diastereomers or as mixture thereof. The use of an enantiomerically pure compound as active ingredient is preferred.

- The invention further relates to the use of the abovementioned
40 carboxylic acid derivatives for producing drugs, in particular for producing inhibitors for ET_A and ET_B receptors. The compounds according to the invention are suitable as selective and as mixed antagonists as have been defined at the outset.

- 45 Compound of the formula V where W is sulfur or oxygen can be prepared as described in WO 96/11914.

12





Compounds of the formula V can be obtained in enantiomerically pure form by starting from enantiomerically pure compounds of the formula III and reacting them with compounds of the formula IV as described in WO 96/11914.

It is additionally possible to obtain enantiomeric compounds of the formula V by subjecting racemic or diastereomeric compounds of the formula V to a conventional racemate resolution using suitable enantiomerically pure bases. Examples of suitable bases of this type are 4-chlorophenylethylamine and the bases mentioned in WO 96/11914.

It is moreover possible to obtain enantiomerically pure compounds of the formula V by acid-catalyzed transesterification as described in DE 19636046.3.


The preparation of compounds of the formula III has been described in WO 96/11914, whereas compounds of the formula IVa (R⁶ = amide) or IVb (R⁶ = sulfonamide/amide) either are known or can be synthesized by generally known methods, such as :



40

The compounds according to the invention in which the substituents have the meanings stated under formula I can be prepared, for example, by reacting the carboxylic acid derivatives of the formula V in which the substituents have the stated meanings with compounds of the formula VI.

13



In formula VI, R^{16} is halogen or $\text{R}^{17}\text{-SO}_2\text{-}$, where R^{17} can be $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-haloalkyl}$ or phenyl. In addition, at least one
 10 of the ring members X or Y or Z is nitrogen. The reaction preferably takes place in an inert solvent or diluent with the addition of a suitable base, ie. a base which deprotonates the intermediate V, at a temperature in the range from room temperature to the boiling point of the solvent.

15

Compounds of type I with $\text{R}^1 = \text{COOH}$ can be obtained directly in this way if the intermediate V where R^1 is COOH is deprotonated with two equivalents of a suitable base and reacted with
 20 compounds of the formula V. This reaction also takes place in an inert solvent and at a temperature in the range from room temperature to the boiling point of the solvent. Examples of such solvents or diluents are aliphatic, alicyclic and aromatic hydrocarbons, each of which may be chlorinated, such as hexane, cyclohexane, petroleum ether, naphtha, benzene, toluene, xylene,
 25 methylene chloride, chloroform, carbon tetrachloride, ethyl chloride and trichloroethylene, ethers, such as diisopropyl ether, dibutyl ether, methyl tert-butyl ether, propylene oxide, dioxane and tetrahydrofuran, nitriles such as acetonitrile and propionitrile, amides such as dimethylformamide,
 30 dimethylacetamide and N-methylpyrrolidone, sulfoxides and sulfones, for example dimethyl sulfoxide and sulfolane.

Compounds of the formula VI are known, some of them can be bought or they can be prepared in a generally known manner.

35

It is possible to use as base an alkali metal or alkaline earth metal hydride such as sodium hydride potassium hydride or calcium hydride, a carbonate such as alkali metal carbonate, eg. sodium or potassium carbonate, an alkali metal or alkaline earth metal
 40 hydroxide such as sodium or potassium hydroxide, an organometallic compound such as butyllithium or an alkali metal amide such as lithium diisopropylamide or lithium amide.

Compounds of the formula I can also be prepared by starting from
 45 the corresponding carboxylic acids, ie. compounds of the formula I where R^1 is COOH , and first converting them in a conventional way into an activated form such as an acid halide, an anhydride

or imidazolidine, and then reacting the latter with an appropriate hydroxyl compound HOR⁹. This reaction can be carried out in conventional solvents and often requires addition of a base, in which case those mentioned above are suitable. These two steps
 5 can also be simplified, for example, by allowing the carboxylic acid to act on the hydroxyl compound in the presence of a dehydrating agent such as a carbodiimide.

It is additionally possible to prepare compounds of the formula I
 10 also by starting from salts of the corresponding carboxylic acids, ie. from compounds of the formula I where R¹ is a group COOM where M can be an alkali metal cation or the equivalent of an alkaline earth metal cation. These salts can be reacted with many compounds of the formula R-A, where A is a conventional
 15 nucleofugic leaving group, for example halogen such as chlorine, bromine, iodine or unsubstituted or halogen-, alkyl- or haloalkyl-substituted aryl- or alkylsulfonyl such as toluenesulfonyl and methylsulfonyl, or another equivalent leaving group. Compounds of the formula R-A having a reactive substituent
 20 A are known or can easily be obtained with general expert knowledge. This reaction can be carried out in conventional solvents and is advantageously undertaken with addition of a base, in which case those mentioned above are suitable.

25 It is necessary in some cases for preparing the compounds I according to the invention to use generally known protective group techniques. If, for example, R¹³ is 4-hydroxyphenyl, the hydroxyl group can be initially protected as benzyl ether, which is then cleaved at a suitable stage in the reaction sequence.

30 Compounds of the formula I where R¹ is tetrazole [sic] can be prepared as described in WO 96/11914.

With a view to the biological effect, preferred carboxylic acid
 35 derivatives of the formula I, both as pure enantiomers and pure diastereomers and as mixture thereof, are those where the substituents have the following meanings:


R² hydrogen, hydroxyl, halogen, N(C₁-C₄-alkyl)₂, C₁-C₄-alkyl,
 40 C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-hydroxyalkyl, or CR² is linked to CR¹² as indicated under Z to give a 5- or 6-membered ring;

X nitrogen or methine;
 45


Y nitrogen or methine;

15

- Z nitrogen or CR¹², where R¹² is hydrogen, fluorine or C₁-C₄-alkyl, or CR¹² forms together with CR² or CR³ a 5- or 6-membered alkylene or alkenylene ring which can be substituted by one or two methyl groups and in each of which a methylene group can be replaced by oxygen or sulfur, such as -CH₂-CH₂-O-, -CH₂-CH₂-CH₂-O-, -CH=CH-O-, -CH=CH-CH₂O-, -CH(CH₃)-CH(CH₃)-O-, -CH=C(CH₃)-O-, -C(CH₃)=C(CH₃)-O-, or -C(CH₃)=C(CH₃)-S;
- 5
- 10 At least one of the ring members X, Y or Z is nitrogen.
- R³ hydrogen, hydroxyl, halogen, N(C₁-C₄-alkyl)₂, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkyl, C₁-C₄-hydroxyalkyl, C₁-C₄-haloalkoxy, or CR³ is linked to CR¹⁰ as indicated under Z to give a 5- or 6-membered ring;
- 15
- R⁴ and R⁵ (which may be identical or different):
- 20 phenyl or naphthyl, each of which may be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, mercapto, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, NH(C₁-C₄-alkyl) or N(C₁-C₄-alkyl)₂ or phenyl which may be substituted one or more times, eg. one to three times, by halogen, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio; or
- 25
- 30 phenyl or naphthyl which are connected together in ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂-, NH- or N-alkyl group
- C₃-C₈-cycloalkyl;
- 35 R⁶ a group



or



40

where the molecular weight of the groups R¹³ and R¹⁴ taken together must be at least 60.

- 45 R¹³ and R¹⁴ (which may be identical or different):

16

hydrogen, C₁-C₈-alkyl, C₃-C₈-alkenyl or C₃-C₈-alkynyl, it being possible for each of these radicals to be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, amino, cyano, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₃-C₈-cycloalkyl, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, phenoxy or phenyl, it being possible for said aryl radicals in turn to be substituted once to three times by halogen, hydroxyl, carboxyl, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, N(C₁-C₄-alkyl)₂, or C₁-C₄-alkylthio;

C₃-C₈-cycloalkyl, it being possible for each of these radicals to be substituted one or more times by: halogen, hydroxyl, mercapto, carboxyl, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy;

phenyl or naphthyl, each of which may be substituted by one or more of the following radicals: halogen, carboxyl, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxy, phenoxy, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted once to three times by halogen; C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;

or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain which is closed to a ring and which may be substituted one or more times by C₁-C₄-alkyl, C₁-C₄-haloalkyl, and in which one alkylene [sic] group can be replaced by oxygen or sulfur, such as -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-, -(CH₂)₂-S-(CH₂)₂-;

or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain or C₄-C₇-alkenylene chain which is closed to a ring and to which a phenyl ring is fused, such as 7-azabicyclo[4.2.0]octa-1,3,5-triene, 2,3-dihydroindole, indole, 1,3-dihydroisoindole, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, it being possible for the phenyl ring in each case to be substituted once to three times by halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, hydroxyl, carboxyl.

The molecular weight of the groups R¹³ and R¹⁴ taken together must be at least 46.

R⁷ and R⁸ (which may be identical or different):

hydrogen, C₁-C₄-alkyl.

- 5 R¹⁵ C₁-C₄-alkyl, C₁-C₄-alkoxy, each of which may carry one of the following radicals: hydroxyl, carboxyl, amino, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, carboxamide [sic] or CON(C₁-C₄-alkyl)₂.
- 10 R¹⁸ hydrogen;
- C₁-C₄-alkyl, C₃-C₄-alkenyl or C₃-C₄-alkynyl, where each of these radicals can be substituted one or more times by: halogen, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy,
- 15 C₃-C₈-cycloalkyl, phenoxy or phenyl, where the abovementioned aryl radicals for their part can be substituted one or more times, for example one to three times by halogen, hydroxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio;
- 20 C₃-C₈-cycloalkyl, where each of these radicals can be substituted one or more times by: C₁-C₄-alkyl;
- phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, hydroxyl, R¹⁵,
- 25 C₁-C₄-alkyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted one or more times, for example one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy;
- 30 R¹⁹ C₁-C₄-alkylcarbonyl, C₂-C₄-alkenylcarbonyl or C₂-C₄-alkynylcarbonyl, where each of these radicals can be substituted one or more times by: halogen, C₁-C₄-alkoxy, C₃-C₈-cycloalkyl, phenoxy or phenyl, where the abovementioned aryl radicals for
- 35 their part can be substituted one or more times, for example one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio;
- C₃-C₈-cycloalkylcarbonyl, where each of these radicals can be
- 40 substituted one or more times by: C₁-C₄-alkyl;
- phenylcarbonyl or naphthylcarbonyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, hydroxyl, R¹⁵, C₁-C₄-alkyl, C₁-C₄-haloalkyl,
- 45 C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy, phenoxy, C₁-C₄-alkylthio, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted one or more times,

18

for example one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, or C₁-C₄-alkylthio;

- 5 C₁-C₄-alkylsulfonyl, where each of these radicals can be substituted one or more times by: halogen, C₁-C₄-alkoxy, phenyl, where the abovementioned aryl radical for its part can be substituted one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio;
- 10 C₃-C₈-cycloalkylsulfonyl; phenylsulfonyl or naphthylsulfonyl, each of which can be substituted by one or more of the following radicals: halogen, cyano, R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxy, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted
- 15 one to three times by halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio;

R²⁰ hydrogen;

- 20 C₁-C₄-alkyl, where each of these radicals can be monosubstituted by: hydroxyl, mercapto, carboxyl, amino, C₃-C₈-cycloalkyl, indolyl, phenoxy or phenyl, where the abovementioned aryl radicals for their part can be substituted one to three times by halogen, hydroxyl,
- 25 mercapto, carboxyl, C₁-C₄-alkyl, C₁-C₄-alkoxy, amino or C₁-C₄-alkylthio.

R²¹ hydrogen, C₁-C₄-alkyl.

- 30 W sulfur or oxygen;

Particularly preferred compounds of the formula I, both as pure enantiomers and pure diastereomers or as mixture thereof, are those where the substituents have the following meanings:

- 35 R² trifluoromethyl, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, hydroxymethyl, or CR² is linked to CR¹² as indicated under Z to give a 5- or 6-membered ring;
- 40 X nitrogen or methine;
- Y nitrogen or methine;
- Z nitrogen or CR¹² where R¹² are [sic] hydrogen, fluorine or
- 45 C₁-C₄-alkyl, or CR¹² forms together with CR² or CR³ einen 5- or 6-membered alkylene or alkenylene ring which may be substituted by one or two methyl groups, and in each of which

19

one methylene group can be replaced by oxygen or sulfur, such as $-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{CH}=\text{CH}-\text{O}-$, $-\text{CH}(\text{CH}_3)-\text{CH}(\text{CH}_3)-\text{O}-$, $-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-\text{O}-$;

5 at least one of the ring members X, Y or Z is nitrogen

R^3 trifluoromethyl, $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_4\text{-alkylthio}$, hydroxymethyl, or CR^3 is linked to CR^{12} as indicated under Z to give a 5- or 6-membered ring;

10

R^4 and R^5 (which may be identical or different):


phenyl or naphthyl, each of which may be substituted by one or more of the following radicals: halogen, hydroxyl,

15 $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-haloalkyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_4\text{-haloalkoxy}$, $\text{NH}(\text{C}_1\text{-C}_4\text{-alkyl})$, $\text{N}(\text{C}_1\text{-C}_4\text{-alkyl})_2$, phenoxy or phenyl, which may be substituted once to three times by halogen, $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-haloalkyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$ or $\text{C}_1\text{-C}_4\text{-haloalkoxy}$; or

20 phenyl or naphthyl which are connected together in ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an $\text{SO}_2\text{-}$, NH- or N-alkyl group

25 $\text{C}_5\text{-C}_7\text{-cycloalkyl}$;

R^6 a group



35 where the molar weight of the groups R^{13} and R^{14} taken together must be at least 60.

R^{13} and R^{14} (which may be identical or different):

40 hydrogen, $\text{C}_1\text{-C}_5\text{-alkyl}$, $\text{C}_3\text{-C}_5\text{-alkenyl}$ or $\text{C}_3\text{-C}_5\text{-alkynyl}$, it being possible for each of these radicals to be substituted one or more times by: halogen, hydroxyl, carboxyl, amino, $\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_4\text{-haloalkoxy}$, $\text{C}_5\text{-C}_6\text{-cycloalkyl}$, $\text{NH}(\text{C}_1\text{-C}_4\text{-alkyl})$, $\text{N}(\text{C}_1\text{-C}_4\text{-alkyl})_2$, phenoxy or phenyl, it being possible for said aryl radicals in turn to be substituted one
45 to three times by halogen, hydroxyl, $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-haloalkyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_4\text{-haloalkoxy}$, $\text{N}(\text{C}_1\text{-C}_4\text{-alkyl})_2$;

20

- C₃-C₈-cycloalkyl, it being possible for each of these radicals to be substituted one or more times by: halogen, carboxyl, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy; phenyl which can be substituted one to three times by:
- 5 halogen, carboxyl, hydroxyl, amino, R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-haloalkoxy, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, dioxomethylene [sic], dioxoethylene [sic] or phenyl which can be substituted once
- 10 to three times by halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio;
- or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain which is closed to a ring and which may be substituted one or more times by C₁-C₄-alkyl and in which one alkylene [sic] group can
- 15 be replaced by oxygen or sulfur, such as -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-, -(CH₂)₂-S-(CH₂)₂-;
- or R¹³ and R¹⁴ together form a C₃-C₇-alkylene chain which is
- 20 closed to a ring and to which [lacuna] phenyl ring is fused, such as 2,3-dihydroindole, indole, 1,3-dihydroisoindole, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, it being possible for the phenyl ring in each case to be
- substituted one to three times by halogen, C₁-C₄-alkyl,
- 25 C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, hydroxyl, carboxyl.

The groups R¹³ and R¹⁴ taken together must contain at least 5 carbon atoms.

- 30 R⁷ and R⁸ (which may be identical or different):

hydrogen, C₁-C₄-alkyl.

- 35 R¹⁵ C₁-C₄-alkyl, C₁-C₄-alkoxy, each of which may carry one of the following radicals: hydroxyl, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, carboxamide [sic] or CON(C₁-C₄-alkyl)₂.

R¹⁸ hydrogen;

- 40 C₁-C₄-alkyl, where each of these radicals can be substituted one to three times by: halogen, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₈-cycloalkyl, phenoxy or phenyl, where the abovementioned aryl radicals for their part can be substituted one to three
- 45 times by: halogen, C₁-C₄-alkyl or C₁-C₄-alkoxy;

C₃-C₈-cycloalkyl;

21

- phenyl which can be substituted one to three times by:
 halogen, hydroxyl, R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxycarbonyl,
 C₁-C₄-alkoxy, dioxomethylene [sic], dioxoethylene [sic] or
 phenyl which can be substituted one to three times by:
 5 halogen, C₁-C₄-alkyl, trifluoromethyl, C₁-C₄-alkoxy;
- R¹⁹ C₁-C₄-alkylcarbonyl, where each of these radicals can be
 substituted one to three times by: halogen, C₁-C₄-alkoxy,
 C₃-C₈-cycloalkyl, phenyl which can be substituted for its part
 10 one to three times by: halogen, C₁-C₄-alkyl or C₁-C₄-alkoxy;
- C₃-C₈-cycloalkylcarbonyl;
- phenylcarbonyl or naphthylcarbonyl, each of which can be
 15 substituted by one or more of the following radicals:
 halogen, R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxy, phenoxy,
 dioxomethylene [sic], dioxoethylene [sic] or phenyl which can
 be substituted one to three times by: halogen, C₁-C₄-alkyl or
 C₁-C₄-alkoxy;
- 20 C₁-C₄-alkylsulfonyl, where each of these radicals can be
 substituted one to three times by: halogen, C₁-C₄-alkoxy,
 phenyl which for its part can be substituted one to three
 times by: halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy or
 25 C₁-C₄-alkylthio;
- C₃-C₈-cycloalkylsulfonyl;
- phenylsulfonyl or naphthylsulfonyl, where each of these
 30 radicals can be substituted one to three times by: halogen,
 R¹⁵, C₁-C₄-alkyl, C₁-C₄-alkoxy, dioxomethylene [sic],
 dioxoethylene [sic] or phenyl;
- R²⁰ hydrogen, C₁-C₄-alkyl.
 35 R²¹ hydrogen, C₁-C₄-alkyl.
- W sulfur or oxygen;
- 40 The compounds of the present invention provide a novel
 therapeutic potential for the treatment of hypertension,
 pulmonary hypertension, myocardial infarct, angina pectoris,
 arrhythmia, acute/chronic kidney failure, chronic cardiac
 insufficiency, renal insufficiency, cerebral vasospasms, cerebral
 45 ischemia, subarachnoid hemorrhages, migraine, asthma,
 atherosclerosis, endotoxic shock, endotoxin-induced organ
 failure, intravascular coagulation, restenosis after angioplasty

and bypass operations, benign prostate hyperplasia, ischemic and intoxication-induced kidney failure or hypertension, metastasis and growth of mesenchymal tumors, contrast agent-induced kidney failure, pancreatitis, gastrointestinal ulcers

5

The invention furthermore provides combinations of endothelin receptor antagonists of the formula I and inhibitors of the renin-angiotensin system. Inhibitors of the renin-angiotensin system are renin inhibitors, angiotensin-II antagonists and

10 angiotensin converting enzyme (ACE) inhibitors. Preference is given to combinations of endothelin receptor antagonists of the formula I and ACE inhibitors.

The invention furthermore provides combinations of endothelin receptor antagonists of the formula I and beta blockers.

15 The invention furthermore provides combinations of endothelin receptor antagonists of the formula I and diuretics.

The invention furthermore provides combinations of endothelin receptor antagonists of the formula I and substances which block the action of VEGF (vascular endothelial growth factor). Such

20 substances are, for example, antibodies directed against VEGF, or specific binding proteins, or else low-molecular-weight substances which are able to specifically inhibit VEGF release or receptor binding.

The abovementioned combinations can be administered

25 simultaneously or sequentially. They can be employed both in a single pharmaceutical formulation or else in separate formulations. The application form can also vary, for example, the endothelin receptor antagonists can be administered orally and the VEGF inhibitors parenterally.


30 These combination products are particularly suitable for the treatment and prevention of hypertension and its sequelae, and for the treatment of cardiac insufficiency.

The good action of the compounds can be shown in the following experiments: [sic]

35

The invention furthermore provides a structural fragment of the formula

40




45 in which the radicals R¹, R⁴, R⁵, R⁶, R⁷, R⁸ and W are as defined above.

23

Such structural fragments are suitable as structural components of endothelin receptor antagonists.


The invention furthermore provides endothelin receptor antagonists comprising a structural fragment of the formula



in which the radicals R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , W , X , Y and Z are as defined above, covalently linked to a group which has a molecular weight of at least 30, preferably 40.


The invention furthermore provides

endothelin receptor antagonists comprising a structural fragment of the formula



in which the radicals R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , R^{20} , R^{21} , W , X , Y and Z are as defined in claim 1, covalently linked via a nitrogen atom to a group which has a molecular weight of at least 58.

The invention furthermore provides compounds of the formula Ia



in which the radicals R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , R^{20} , R^{21} , W , X , Y and Z are as defined in claim 1.

Receptor binding studies

Cloned human ET_A or ET_B receptor-expressing CHO cells were employed for binding studies.

Membrane preparation

The ET_A or ET_B receptor-expressing CHO cells were grown in DmeM NUT MIX F₁₂-medium (Gibco, No. 21331-020) with 10% fetal calf
5 serum (PAA Laboratories GmbH, Linz, No. A15-022), 1 mM glutamine (Gibco No. 25030-024), 100 U/ml penicillin and 100 µg/ml streptomycin (Gibco [sic], Sigma No. P-0781). After 48 hours, the cells were washed with PBS and incubated with 0.05% trypsin-containing PBS at 37°C for 5 minutes. This was followed by
10 neutralization with medium, and the cells were collected by centrifugation at 300 x g.

For the membrane preparation, the cells were adjusted to a concentration of 10⁸ cells/ml of buffer (50 mM tris-HCL [sic]
15 buffer, pH 7.4) and then disintegrated by ultrasound Branson Sonifier 250, 40-70 seconds/constant/output [sic] 20).

Binding assays

20 For the ET_A and ET_B receptor binding assay, the membranes were suspended in incubation buffer (50 mM tris-HCL, pH 7.4 with 5 mM MnCl₂, 40 mg/ml Bacitracin and 0.2% BSA) in a concentration of 50 µg of protein per assay mixture and incubated with 25 pM [125I]-ET₁ [sic] (ET_A receptor assay) or 25 pM [125I]-ET₃ [sic]
25 (ET_B receptor assay) at 25°C in the presence and absence of test substance. The nonspecific binding was determined with 10⁻⁷ M ET₁. After 30 min, filtration through GF/B glass fiber filters (Whatman, England) in a Skatron cell harvester (Skatron, Lier, Norway) was carried out to separate free and bound radioligand,
30 and the filters was washed with ice-cold tris-HCL buffer, pH 7.4 with 0.2% BSA. The radioactivity collected on the filters was quantified using a Packard 2200 CA liquid scintillation counter.

In vivo testing of the ET antagonists:

35 Male ST rats weighing 250-300 g were anesthetized with amobarbital, artificially ventilated, vagotomized and pithed. The carotid artery and the jugular vein were cathetized [sic].
40 In control animals, intravenous administration of 1 µg/kg ET₁ results in a distinct rise in blood pressure which persists for a lengthy period.

The test animals received i.v. injections of the test compounds
45 (1 ml/kg) 30 min before the ET₁ administration. To determine the ET antagonistic properties, the changes in the blood pressure of

0050/48493

25

the test animals were compared with those of the control animals.

Oral testing of the mixed ET_A and ET_B antagonists:

- 5 Male normotensive rats (Sprague Dawley, Janvier) weighing 250-350 g are pretreated orally with the test substances. 80 minutes later, the animals are anesthetized with urethane and the carotid artery (for measuring the blood pressure) and the jugular vein (administration of big endothelin/endothelin-1) are
10 catheterized.

- After a stabilization period, big endothelin (20 µg/kg, administration volume 0.5 ml/kg) or ET1 (0.3 µg/kg, administration volume 0.5 ml/kg) is administered intravenously.
15 The blood pressure and heart rate are recorded continuously for 30 minutes. The distinct and long-lasting changes in blood pressure are calculated as area under the curve (AUC). To determine the antagonistic effect of the test substances, the AUC of the substance-treated animals is compared with the AUC of the
20 control animals.

- The compounds according to the invention can be administered orally or parenterally (subcutaneously, intravenously, intramuscularly, intraperitoneally) in a conventional way.
25 Administration can also take place with vapors or sprays through the nasopharyngeal space.

- The dosage depends on the age, condition and weight of the patient and on the mode of administration. As a rule, the daily
30 dose of active ingredient is about 0.5-50 mg/kg of bodyweight on oral administration and about 0.1-10 mg/kg of bodyweight on parenteral administration.

- The novel compounds can be used in conventional solid or liquid
35 pharmaceutical forms, eg. as uncoated or (film-)coated tablets, capsules, powders, granules, suppositories, solutions, ointments, creams or sprays. These are produced in a conventional way. The active ingredients can for this purpose be processed with conventional pharmaceutical auxiliaries such as tablet binders,
40 bulking agents, preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, release-slowing agents, antioxidants and/or propellant gases (cf. H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1991). The administration
45 forms obtained in this way normally contain from 0.1 to 90% by weight of active ingredient.

Synthesis examples

Example 1:

5 N,N-di-n-Butyl-2-benzyloxyacetamide

3 g of N-methylmorpholine and 4 g of isobutyl chloroformate were successively added dropwise to 5 g of 2-benzyloxyacetic acid in 50 ml of THF at -10°C. The mixture was stirred for 10 minutes and
10 then 5 ml of di-n-butylamine and a further 3 g of N-methylmorpholine were added. After one hour, the mixture was added to 500 ml of water and extracted several times with ether. The collected organic phases were dried over magnesium sulfate and, after removal of the solvent by distillation, 7 g of an oil
15 were isolated and were immediately employed further.

Example 2:

N,N-di-n-Butyl-2-hydroxyacetamide

20

4 g of N,N-di-n-butyl-2-benzyloxyacetamide were dissolved in 50 ml of ethanol, and a spatula tip of Pd/carbon was added. The mixture was stirred under a hydrogen atmosphere for 16 hours, and then the catalyst was filtered off and the solvent was distilled
25 off. 3 g of an oil were isolated and were immediately reacted further.

Example 3:

30 Methyl 2-hydroxy-3-(N,N-di-n-butylcarbamoylmethoxy)-3,3-diphenylpropionate

1.3 g of N,N-di-n-butyl-2-hydroxyacetamide and 1.8 g of methyl 2,3-epoxy-3,3-diphenylpropionate were dissolved in 30 ml of
35 methylene chloride and, while cooling in ice, a catalytic amount of p-toluenesulfonic acid was added. The mixture was stirred at room temperature for 24 hours and then added to sodium bicarbonate solution, the organic phase was separated off and dried over magnesium sulfate, and the solvent was distilled off.
40 The residue was purified by chromatography, and 1.4 g of an oil were isolated and were immediately reacted further.

Example 4:

2-Hydroxy-3-(N,N-dibutylcarbamoylmethoxy)-3,3-diphenylpropionic acid

5

1.42 g of methyl 2-hydroxy-3-(N,N-dibutylcarbamoylmethoxy)-3,3-diphenylpropionate were dissolved in 10 ml of dioxane and 4.8 ml of 1N sodium hydroxide solution and stirred at room temperature for 3 hours. Water was then added to the mixture, and the aqueous phase was extracted with ether. The aqueous phase was acidified with hydrochloric acid and extracted with ethyl acetate, and the organic phase was dried over magnesium sulfate. After removal of the solvent by distillation, 1.1 g of oil were isolated and were immediately employed further.

15

Example 5:

2-(4,6-Dimethyl-2-pyrimidinyl-2-oxo)-3-(N,N-dibutylcarbamoylmethoxy)-3,3-diphenylpropionic acid (I-347)

20

560 mg of 2-hydroxy-3-(N,N-dibutylcarbamoylmethoxy)-3,3-diphenylpropionic acid were introduced into THF, and 63 mg of lithium amide and, 10 minutes later, 256 mg of 2-methylsulfone-4,6-dimethylpyrimidine [sic] were added. The mixture was stirred at 50°C for 5 hours and then water was added. The aqueous phase was acidified with citric acid and extracted with ethyl acetate. The organic phase was dried, the solvent was distilled off, and the residue was purified by chromatography. The isolated product was crystallized from ether/n-hexane.

30

¹H-NMR (200 MHz): 7.30-7.20 ppm (10 H, m), 6.75 (1 H, s), 6.15 (1 H, s), 4.50 (1 H, d), 4.20 (1 H, d), 3.30 (2 H, dd), 2.95 (2 H, dd), 2.35 (6 H, s), 1.55-1.00 (8 H, m), 0.95 (3 H, tr), 0.80 (3 H, tr).

35

ESI-MS: M⁺ = 533

Example 6:

40 N-Propyl-N-(2-hydroxyethyl)benzenesulfonamide

At 0°C, 5.16 g (50 mmol) of N-propylethanolamine were introduced into 70 ml of methylene chloride, and 9.7 g (55 mmol) of benzenesulfonyl chloride and 7.6 g (75 mmol) of triethylamine were added successively. After 2 hours of stirring at 0°C, the mixture was allowed to warm to room temperature, stirring was continued for a further hour and the mixture was then extracted

28

with 1M hydrochloric acid and subsequently with 2M aqueous sodium hydroxide solution. The organic phase was dried over Na_2SO_4 , filtered, concentrated and the resulting residue (13.2 g) was chromatographed over silica gel (methylene chloride/methanol 5 19:1). Yield: 7.4 g as an oil which was directly reacted further.

Example 7:

Methyl 2-hydroxy-3-(2-(N-propyl-N-benzenesulfonylamino)ethoxy)-
10 3,3-diphenylpropionate

7.3 g (30 mmol) of N-propyl-N-(2-hydroxyethyl)benzenesulfonamide and 7.6 g (30 mmol) of methyl 2,3-epoxy-3,3-diphenylpropionate were dissolved in 40 ml of methylene chloride and, with
15 ice-cooling, 0.57 g (3 mmol) of p-toluenesulfonic acid was added. The reaction mixture was stirred at room temperature for 24 hours and then diluted with methylene chloride and extracted with 2M aqueous sodium hydroxide solution, the organic phase was separated off and dried over sodium sulfate and the solvent was
20 distilled off. The residue (12.0 g of an oil) was directly reacted further.

Example 8:

25 2-Hydroxy-3-(2-(N-propyl-N-benzenesulfonylamino)ethoxy)-
3,3-diphenylpropionic acid

6.0 g of methyl
2-hydroxy-3-(2-(N-propoyl-N-benzenesulfonylamino)-
30 ethoxy)-3,3-diphenylpropionate [sic] (from Example 7) were dissolved in 70 ml of dioxane and mixed with 36 ml of 1M KOH and stirred at room temperature overnight. The reaction mixture was subsequently mixed with water and the aqueous phase was extracted with ether. The aqueous phase was acidified with hydrochloric
35 acid and extracted with ether, the organic phase was dried over sodium sulfate and the solvent was distilled off. The residue (3.3 g) was chromatographed over silica gel (methylene chloride/methanol 9:1), giving 2.6 g of product.

40 m.p.: 144-146°C (from ether)

Example 9:

2-(4-Methyl-6-methoxypyrimidin-2-yloxy)-3-(2-(N-propyl-N-benzene-
45 sulfonylamino)ethoxy)-3,3-diphenylpropionic acid (II-2)

29

135 mg (5.6 mmol) of lithium amide (95%) were suspended in 5 ml of dimethylformamide, cooled to 0°C, admixed with 0.9 g (1.9 mmol) of 2-hydroxy-3-(2-(N-propoyl-N-benzenesulfonylamino)ethoxy)-3,3-diphenylpropionic [sic] acid, dissolved in 4 ml of
 5 dimethylformamide and stirred at 0°C for 30 min. 0.56 g (2.8 mmol) of 2-methylsulfone-4-methyl-6-methoxypyrimidine [sic] were then added and the mixture was stirred at room temperature overnight and then mixed with water. The aqueous phase was extracted with ether, the resulting organic phase was discarded and the aqueous
 10 phase was adjusted to pH 1 using hydrochloric acid and extracted with ether. The organic phase was dried over sodium sulfate, the solvent was distilled off and the residue (1.26 g) was triturated in ether/heptane. Yield: 0.9 g of a white solid.

15 ESI-MS: 606 (M+H)⁺

¹H-NMR (270 MHz, CDCl₃): 7.70-7.85 ppm (2 H, m); 7.20-7.55 (13 H, m); 6.25 (1 H, s); 6.15 (1 H, s); 3.9 (3 H, s); 3.50-3.75 (2 H, m); 3.20-3.50 (2 H, m); 3.00-3.15 (2 H, m); 2.30 (3 H, s);
 20 1.35-1.55 (2 H, m); 0.75 (3 H, tr).

Example 10:

Methyl 2-hydroxy-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenyl-
 25 propionate

9.8 g (50 mmol) of benzyl (2-hydroxyethyl)carbamate and 12.7 g (50 mmol) of methyl 2,3-epoxy-3,3-diphenylpropionate were dissolved in 80 ml of methylene chloride and, with ice-cooling,
 30 0.95 g (5 mmol) of p-toluenesulfonic acid was added. The reaction mixture was stirred at room temperature for 24 hours and then diluted with methylene chloride and extracted with 2M aqueous sodium hydroxide solution, the organic phase was separated off and dried over sodium sulfate and the solvent was distilled off.
 35 The residue (22.2 g of an oil) was directly reacted further.

Example 11:

2-Hydroxy-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenyl-
 40 propionic acid

22.2 g of methyl 2-hydroxy-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenylpropionate (from Example 10) were dissolved in 300 ml of dioxane, mixed with 148 ml of 1M KOH and stirred at room
 45 temperature overnight. The reaction mixture was subsequently mixed with water and the aqueous phase was extracted with ether. The aqueous phase was acidified with hydrochloric acid and

30

extracted with ether, the organic phase was dried over sodium sulfate and the solvent was distilled off. The residue (17.5 g) was directly used further.

5 Example 12:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenylpropionic acid (II-32)

- 10 2.5 g (103 mmol) of lithium amide (95%) were suspended in 60 ml of dimethylformamide, cooled to 0°C, admixed with 15 g (34.4 mmol) of 2-hydroxy-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenylpropionic acid, dissolved in 60 ml of dimethylformamide, and stirred at 0°C for 30 min. 8.34 g (44.7 mmol) of 2-methyl-
- 15 sulfone-4-methyl-6-methoxypyrimidine [sic] in 30 ml of dimethylformamide were then added and the mixture was stirred at room temperature for 3 days and then mixed with water. The aqueous phase was extracted with ether, the resulting organic phase was discarded and the aqueous phase was then adjusted to
- 20 pH 1 using hydrochloric acid and extracted with ether. The organic phase was dried over sodium sulfate, the solvent was distilled off and the residue was chromatographed over silica gel (methylene chloride/methanol 9:1). Yield: 14.0 g of a white foam.
- 25 ¹H-NMR (270 MHz, DMSO): 12.0-13.0 ppm (1H, brd); 7.10-7.45 (16 H, m); 6.95 (1 H, s); 6.20 (1 H, s); 5.0 (2 H, s); 3.80-3.95 (2 H, m); 3.55-3.70 (2 H, m); 3.20-3.40 (2 H, m); 2.30 (6 H, s).

Example 13:

30

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-aminoethoxy)-3,3-diphenylpropionic acid

- A solution of 13.1 g (24.2 mmol) of 2-(4,6-dimethylpyrimidin-
- 35 2-yloxy)-3-(2-benzyloxycarbonylaminoethoxy)-3,3-diphenylpropionic acid in 200 ml of methanol was hydrogenated with hydrogen overnight, under atmospheric pressure and at room temperature, using 800 mg of palladium on activated carbon (10%). The reaction mixture was diluted with methanol to dissolve precipitated
- 40 product, filtered and concentrated. Yield: 9.6 g of a white solid.

- ¹H-NMR (270 MHz, DMSO): 7.10-7.40 ppm (10 H, m); 6.90 (1 H, s); 6.00 (1 H, s); 3.60-3.75 (2 H, m); 2.90-3.00 (2 H, m); 2.25 (6 H, s).
- 45 s).

Example 14:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(3,4-dimethoxybenzoyl-amino)ethoxy)-3,3-diphenylpropionic acid (II-62)

5

A solution of 1.0 g (2.5 mmol) of 2-(4,6-dimethylpyrimidin-2-yl-oxy)-3-(2-aminoethoxy)-3,3-diphenylpropionic acid in 10 ml of methylene chloride was successively admixed at room temperature with 0.35 g (2.7 mmol) of N-ethyldiisopropylamine, 0.03 g

10 (0.2 mmol) of dimethylaminopyridine and 0.54 g (2.7 mmol) of 3,4-dimethoxybenzoyl chloride. The mixture was stirred at room temperature for 4 days and then diluted with diethyl ether and extracted with 1M hydrochloric acid and 1M aqueous sodium hydroxide solution, and the combined alkaline phases were made
15 acidic and extracted with ether. The organic phase was dried over sodium sulfate, the solvent was distilled off and the residue (0.9 g) was chromatographed over silica gel (methylene chloride/methanol 9:1). Yield: 280 mg of a white foam.

20 ESI-MS: 571 (M+H)⁺

¹H-NMR (360 MHz, DMSO): 7.10-7.55 ppm (12 H, m); 7.00 (1 H, d); 6.90 (1 H, s); 6.20 (1 H, s); 3.65-4.00 (2 H, m); 3.80 (3 H, s); 3.75 (3 H, s); 3.45-3.55 (2 H, m); 2.30 (6 H, s).

25

Example 15:

Benzyl (S)-5,5-diphenyl-2-oxo-1,4-dioxane-6-carboxylate

30 38 g (100 mmol) of the benzyl (S)-2-hydroxy-3-methoxy-3,3-diphenylpropionate were added to 9.8 g (130 mmol) of glycolic acid, and the mixture was stirred with 300 mg of anhydrous para-toluenesulfonic acid at 70°C on a rotary evaporator under reduced pressure for 20 minutes. The content of the flask was
35 taken up in dichloromethane, the acid was separated off using sodium hydrogen sulfate solution, the organic phase was separated off and dried and the solvent was distilled off. The residue was recrystallized from ether, and 21 g (54 mmol) of product were isolated.

40

$[\alpha]_D = +283^\circ$ at 20°C in ethanol

Example 16:

45 (S)-(1,1-Diphenyl-2-hydroxy-2-benzyloxycarbonylethoxy)acetic acid

14 g (36 mmol) of benzyl (S)-5,5-diphenyl-2-oxo-1,4-dioxane-6-carboxylate were dissolved in 50 ml of DMF and, with ice-cooling, 43 ml of 1 N NaOH solution were added. After ten minutes, the mixture was diluted with 300 ml of water and 5 neutralized with 43 ml of 1 N hydrochloric acid and the aqueous phase was extracted with ether. The ether phase was dried, the solvent was distilled off and the residue (8.8 g, 21 mmol of an oil) was directly reacted further.

10 Example 17:

(S)-(1,1-Diphenyl-2-(4,6-dimethylpyrimidin-2-yloxy)-2-benzyloxy-carbonylethoxy)acetic acid

15 6.6 g (15 mmol) of (S)-(1,1-diphenyl-2-hydroxy-2-benzyloxy-carbonylethoxy)acetic acid were introduced into 75 ml of DMF, and 1.4 g of NaH (30 mmol, 50% suspension) were added a little at a time with ice-cooling. 3.6 g (19.5 mmol) of 4,6-dimethyl-2-methylsulfonepyrimidine [sic] were subsequently 20 added and the mixture was stirred for a quarter of an hour and then warmed to room temperature. After 45 minutes, the reaction was complete and the reaction solution was poured into 500 ml of ice-water. The aqueous phase was extracted with ethyl acetate, the combined organic phases were dried and the solvent was 25 distilled off. The oily residue was stirred with ether/hexane, and 6.4 g of crystals could be isolated.

Example 18:

30 Benzyl (S)-2-(4,6-dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(3-methylphenyl)carbamoylmethoxy-3,3-diphenylpropionate [sic]

Under protective gas and at -10°C, 512 mg (1 mmol) of S-(1,1-diphenyl-2-(4,6-dimethylpyrimidin-2-yloxy)-2-benzyloxy-35 carbonylethoxy)acetic acid were dissolved in 20 ml of dichloromethane, and 121 mg (1 mmol) of N-(3-methylphenyl)-N-methylamine, 92 ml (1 mmol) of ethyldiisopropylamine and 191 mg (1 mmol) of N-(3-dimethylaminopropyl)-N-ethylcarbodiimide were added successively. After one hour, the reaction mixture was 40 warmed to room temperature and stirred for a further 16 hours. The mixture was subsequently diluted with dichloromethane to 100 ml and washed with citric acid and water. The organic phase was dried and the solvent was distilled off. For further purification, the residue was subjected to flash chromatography 45 (ethyl acetate/cyclohexane 1/1), and 290 mg of product were isolated and were immediately employed further.

Example 19:

(S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(3-methylphenyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

5

260 mg of benzyl (S)-2-(4,6-dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(3-methylphenyl)carbamoylmethoxy)-3,3-diphenylpropionate were dissolved in 50 ml of ethyl acetate in ethyl acetate [sic], and a spatula tip of Pd/C was added. The mixture was stirred under an atmosphere of hydrogen for 2 hours. The Pd/C was subsequently filtered off and the ethyl acetate was distilled off. The residue was stirred with ether/hexane, and 127 mg of crystals could be isolated.

15 $[\alpha]_D = + 90^\circ$ at 20°C in ethanol

$^1\text{H-NMR}$ (200 MHz): 7.40-7.00 ppm (14 H, m), 6.75 (1 H, s), 6.05 (1 H, s), 4.15 (1 H, d), 3.75 (1 H, d), 3.25 (3 H, s), 2.40 (6 H, s), 2.20 (3 H, s).

20

ESI-MS: $M^+ = 525$

The following compounds were prepared in a similar way to the examples mentioned above

25

Example 20:

2-(4-Methyl-6-methoxy-2-pyrimidinyloxy)-3-(N,N-dibutylcarbamoylmethoxy)-3,3-diphenylpropionic acid (I-349)

30

$^1\text{H-NMR}$ (200 MHz): 7.30-7.20 ppm (10 H, m), 6.25 (1 H, s), 6.00 (1 H, s), 4.50 (1 H, d), 4.25 (1 H, d), 3.95 (3 H, s), 3.30 (2 H, dd), 2.95 (2 H, dd), 2.25 (3 H, s), 1.55-1.00 (8 H, m), 0.95 (3 H, tr), 0.80 (3 H, tr).

35

ESI-MS: $M^+ = 549$

Example 21:

40 2-(4,6-Dimethyl-2-pyrimidinyloxy)-3-(N-methyl-N-phenylcarbamoylmethoxy)-3,3-diphenylpropionic acid (I-109)

ESI-MS: $M^+ = 511$

45 $^1\text{H-NMR}$ (200 MHz): 7.40-7.20 ppm (15 H, m), 6.80 (1 H, s), 6.15 (1 H, s), 4.15 (1 H, d), 3.8 (1 H, d), 3.30 (3 H, s), 2.35 (6 H, s).

Example 22:

2-(4-methyl-6-methoxy-2-pyrimidinyloxy)-3-(N-methyl-N-phenyl-carbamoylmethoxy)-3,3-diphenylpropionic acid (I-111)

5

¹H-NMR (200 MHz): 7.40-7.20 ppm (15 H, m), 6.30 (1 H, s), 6.00 (1 H, s), 4.20 (1 H, d), 3.80 (3 H, s), 3.75 (1 H, d), 3.25 (3 H, s), 2.30 (3 H, s).

10 ESI-MS: M⁺ = 527

Example 23:

2-(4,6-Dimethyl-2-pyrimidinyloxy)-3-(2-oxo-2-(1,2,3,4-tetrahydro-15 isoquinolin-2-yl)ethoxy)-3,3-diphenylpropionic acid (I-307)

¹H-NMR (200 MHz): 7.40-7.10 ppm (14 H, m), 6.60 (1 H, s), 6.05 (1 H, s), 4.75-4.25 (4 H, m), 3.85 (1 H, m), 3.50-3.25 (1 H, m), 3.00-2.75 (2 H, m), 2.25 (3 H, s), 2.10 (3 H, s).

20

ESI-MS: M⁺ = 537

Example 24:

25 2-(4-Methyl-6-methoxy-2-pyrimidinyloxy)-3-(2-oxo-2-(1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy)-3,3-diphenylpropionic acid (I-309)

¹H-NMR (200 MHz): 7.40-7.10 ppm (14 H, m), 6.20 (1 H, s), 6.00 (1 H, s), 4.75-4.25 (4 H, m), 3.85 (1 H, m), 3.75 (3 H, s), 3.40 (1 H, m), 3.00-2.75 (2 H, m), 2.10 (3 H, s).

30

ESI-MS: M⁺ = 553

Example 25:

35

2-(4,6-Dimethyl-2-pyrimidinyloxy)-3-(N-ethoxymethylene-N-(2,6-diethylphenyl)carbamoylmethoxy)-3,3-diphenylpropionic [sic] acid (I-325)

40 ¹H-NMR (200 MHz): 7.40-7.10 ppm (13 H, m), 6.75 (1 H, s), 6.15 (1 H, s), 5.10 (1 H, d), 4.90 (1 H, d), 4.00-3.70 (4 H, m), 2.70-2.30 (4 H, m), 2.40 (6 H, s), 1.25 (6 H, m), 1.10 (3 H, tr).

ESI-MS: M⁺ = 611.

45

Example 26:

2-(4,6-Dimethyl-2-pyrimidinyloxy)-3-(N-isopropyl-N-phenyl-carbamoylmethoxy)-3,3-diphenylpropionic acid (I-271)

5

¹H-NMR (200 MHz): 7.30-7.10 ppm (15 H, m), 6.70 (1 H, s), 6.10 (1 H, s), 5.10 (1 H, m), 4.00 (1 H, d), 3.60 (1 H, d), 2.30 (6 H, s), 1.10 (6 H, m).

10 ESI-MS: M⁺ = 539.

Example 27:

2-(4,6-Dimethyl-2-pyrimidinyloxy)-3-(N-methoxymethylene-N-(2,6-
15 diisopropylphenyl)carbamoylmethoxy)-3,3-diphenylpropionic [sic] acid (I-334)

¹H-NMR (200 MHz): 7.40-7.10 ppm (13 H, m), 6.75 (1 H, s), 6.15 (1 H, s), 5.10 (1 H, d), 4.90 (1 H, d), 4.10 (1 H, d), 3.75 (1 H, d), 3.50 (3 H, s), 3.30 (1 H, m), 2.9 (1 H, m), 2.30 (6 H, s),
20 1.20 (9 H, m), 0.6 (3 H, d).

ESI-MS: M⁺ = 625.

25 Example 28:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(N-propyl-N-benzene-sulfonylamino)ethoxy)-3,3-diphenylpropionic acid (II-48)

30 ESI-MS: 590 (M+H)⁺

¹H-NMR (270 MHz, CDCl₃): 7.75-7.85 ppm (2 H, m); 7.20-7.55 (13 H, m); 6.70 (1 H, s); 6.25 (1 H, s); 3.55-3.75 (2 H, m); 3.20-3.50 (2 H, m); 3.00-3.15 (2 H, m); 2.35 (6 H, s); 1.35-1.50 (2 H, m);
35 0.75 (3 H, tr).

Example 29:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(N-butyl-N-benzene-sulfonylamino)ethoxy)-3,3-diphenylpropionic acid (II-20)
40

ESI-MS: 604 (M+H)⁺

36

¹H-NMR (200 MHz, CDCl₃): 7.75-7.85 ppm (2 H, m); 7.20-7.55 (13 H, m); 6.70 (1 H, s); 6.20 (1 H, s); 3.20-3.75 (4 H, m); 3.00-3.15 (2 H, m); 2.35 (6 H, s); 1.35-1.50 (2 H, m); 1.10-1.30 (2 H, m); 0.75 (3 H, tr).

5

Example 30:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(4-methoxyphenyl)-carbamoylmethoxy)-3,3-diphenylpropionic acid (I-37)

10

¹H-NMR (200 MHz, DMSO): 9.75 ppm (NH), 7.50-7.10 (12 H, m), 6.90 (1 H, s), 6.80 (2 H, d), 6.10 (1 H, s), 4.25 (1 H, d), 4.10 (1 H, d), 3.75 (3 H, s), 2.25 (6 H, s).

15 ESI-MS: M⁺ = 527

Example 31:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-phenylcarbamoylmethoxy)-3,3-diphenylpropionic acid (I-19)

¹H-NMR (200 MHz, DMSO): 9.90 ppm (NH), 7.70-7.20 (14 H, m), 7.10 (1 H, tr), 6.80 (1 H, s), 6.20 (1 H, s), 4.30 (1 H, d), 4.20 (1 H, d), 2.30 (6 H, s).

25

ESI-MS: M⁺ = 497

Example 32:

30 2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(4-methylphenyl)-carbamoylmethoxy)-3,3-diphenylpropionic acid (I-28)

¹H-NMR (200 MHz, DMSO): 9.80 ppm (NH), 7.50-7.20 (12 H, m), 7.10 (2 H, d), 6.80 (1 H, s), 6.10 (1 H, s), 4.25 (1 H, d), 4.05 (1 H, d), 2.30 (6 H, s), 2.20 (3 H, s).

35

ESI-MS: M⁺ = 511

Example 33:

40

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-butyl-N-phenylcarbamoylmethoxy)-3,3-diphenylpropionic acid (I-190)

¹H-NMR (200 MHz): 7.25-7.10 ppm (15 H, m), 6.70 (1 H, s), 6.10 (1 H, s), 4.20 (1 H, d), 3.7 (2 H, m), 2.25 (6 H, s), 1.5-1.1 (4 h, m), 0.8 (3 H, tr).

45

ESI-MS: $M^+ = 553$

Example 34:

- 5 2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-oxo-2-(6,7-dimethoxy-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy)-3,3-diphenylpropionic acid

ESI-MS: $M^+ = 597$

10

m.p.: 145-148°C

Example 35:

- 15 2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-oxo-2-(4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-2-yl)ethoxy)-3,3-diphenylpropionic acid

ESI-MS: $M^+ = 565$

20

m.p.: 185-187°C

Example 36:

- 25 (S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(3-methylphenyl)-carbamoylmethoxy)-3,3-diphenylpropionic acid

$^1\text{H-NMR}$ (200 MHz): 9.10 ppm (NH), 7.50-7.25 (12 H, m), 7.10 (1 H, tr), 6.80 (1 H, d), 6.60 (1 H, s), 6.20 (1 H, s), 4.10 (1 H, d),
30 3.80 (1 H, d), 2.30 (6 H, s), 2.25 (3 H, s).

ESI-MS: $M^+ = 511$

Example 37:

35

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2-naphth-2-ylethyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

$^1\text{H-NMR}$ (200 MHz): 8.20 ppm (1 H, m), 7.90-7.70 (3 H, m), 7.50-7.15
40 (14 H, m), 6.60/6.65 (1 H, s, rotamers), 6.20/6.15 (1 H, s, rotamers), 4.50 (1 H, d, rotamers), 4.25 (1 H, d, rotamers), 3.9 (1 H, m), 3.50-3.20 (3 H, m), 3.05/2.70 (3 H, s, rotamers)
2.30/2.25 (6 H, s, rotamers).

45 ESI-MS: $M^+ = 589$

Example 38:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2-(4-methoxyphenyl)butyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

5

¹H-NMR (200 MHz): 7.50-7.05 (12 H, m), 6.95-6.60 (3 H, m), 6.05 (1 H, s, rotamers), 4.50-4.00 (2 H, m, rotamers), 3.75 (3 H, d, rotamers), 3.2-2.8 (3 H, m, rotamers), 2.9 (3 H, s, rotamers), 2.30 (6 H, s, rotamers), 1.70-1.50 (2 H, m), 0.70-0.60 (3 H, m,

10 rotamers).

ESI-MS: M⁺ = 597

Example 39:

15

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2-isopropyl-2-(3,4-dimethoxyphenyl)-3-methylbutyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

20 ¹H-NMR (200 MHz): 7.30-7.20 (10 H, m), 6.95-6.60 (4 H, m), 6.20 (1 H, s), 4.40 (2 H, m), 4.05 (1 H, d) 3.85 (7 H, m), 2.5 (3 H, s), 2.3 (6 H, s), 2.30-2.20 (2 H, m), 1.00-0.70 (12 H).

ESI-MS: M⁺ = 683

25

Example 40:

(S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-benzyl-carbamoylmethoxy)-3,3-diphenylpropionic acid

30

¹H-NMR (200 MHz): 7.30-7.10 ppm (15 H, m), 6.75 (1 H, s), 6.20 (1 H, s), 4.75-4.20 (4 H, m, rotamers), 3.00/2.60 (3 H, s, rotamers), 2.35/2.30 (6 H, s, rotamers).

35 ESI-MS: M⁺ = 525

Example 41:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(2,6-diethylphenyl)-carbamoylmethoxy)-3,3-diphenylpropionic acid (I-82)

40

¹H-NMR (200 MHz): 8.30 ppm (NH), 7.50-7.00 (13 H, m), 6.75 (1 H, s), 6.25 (1 H, s), 4.25 (1 H, d), 3.90 (1 H, d), 2.60 (4 H, q), 2.30 (6 H, s), 1.20 (6 H, tr).

45

ESI-MS: M⁺ = 533

Example 42:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(4-chlorophenyl)carbamoyl-methoxy)-3,3-diphenylpropionic acid (I-46)

5

¹H-NMR (200 MHz): 10.00 ppm (NH), 7.70 (2 H, d), 7.50-7.10 (12 H, m), 6.75 (1 H, s), 6.20 (1 H, s), 4.20 (1 H, d), 3.80 (1 H, d), 2.30 (6 H, s).

10 ESI-MS: M⁺ = 531

Example 43:

2-(4,6-Diethylpyrimidin-2-yloxy)-3-(N-methyl-N-phenylcarbamoyl-methoxy)-3,3-diphenylpropionic acid

15

¹H-NMR (200 MHz): 7.50-7.10 ppm (15 H, m), 6.80 (1 H, s), 6.10 (1 H, s), 4.20 (1 H, d), 3.30 (1 H, s), 2.70 (4 H, q), 1.20 (6 H, tr).

20

ESI-MS: M⁺ = 539

Example 44:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-(3-methoxyphenyl)-carbamoylmethoxy)-3,3-diphenylpropionic acid

25

¹H-NMR (200 MHz): 9.80 ppm (NH), 7.50-7.10 (13 H, m), 6.75 (1 H, s), 6.60 (1 H, dtr), 6.20 (1 H, s), 4.10 (1 H, d), 3.80 (1 H, d), 3.75 (3 H, s), 2.30 (6 H, s).

30

ESI-MS: M⁺ = 527

Example 45:

35

(S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-benzylcarbamoyl-methoxy)-3,3-diphenylpropionic acid

¹H-NMR (200 MHz): 7.50-7.10 ppm (15 H, m), 6.75 (1 H, s), 6.20 (1 H, s), 4.45 (1 H, dd), 4.40 (1 H, dd), 4.10 (1 H, d), 3.90 (1 H, d), 2.40 (6 H, s).

40

ESI-MS: M⁺ = 511

45

Example 46:

(S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(4-methoxybenzyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

5

¹H-NMR (200 MHz): 7.50-7.10 ppm (13 H, m), 6.75 (3 H, m, rotamers), 6.20 (1 H, s, rotamers), 4.70-4.00 (4 H, m, rotamers), 3.75 (3 H, s), 3.00/2.70 (3 H, s, rotamers), 2.40/2.35 (6 H, s, rotamers).

10

ESI-MS: M⁺ = 555

Example 47:

15 (S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-ethyl-N-benzylcarbamoylmethoxy)-3,3-diphenylpropionic acid

¹H-NMR (200 MHz): 7.50-7.20 ppm (15 H, m), 6.70 (1 H, s), 6.20 (1 H, s, rotamers), 4.75-4.10 (4 H, m, rotamers), 3.70/3.30/3.00 (20 H, m, rotamers), 2.35/2.30 (6 H, s, rotamers), 1.10/1.00 (3 H, tr, rotamers).

ESI-MS: M⁺ = 539

25 Example 48:

(S)-2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2,6-dichlorobenzyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

30 ESI-MS: M⁺ = 593

m.p.: 105-110°C

Example 49:

35

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2-phenylethyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

¹H-NMR (200 MHz): 7.50-7.20 ppm (14 H, m), 6.75 (1 H, m), 6.70 (1 H, s, rotamers), 6.15/6.10 (1 H, s, rotamers), 4.50-4.00 (2 H, d, rotamers), 3.70 (1 H, m), 3.50 (1 H, m), 3.20/2.70 (5 H, m, rotamers), 2.35/2.30 (6 H, s, rotamers).

ESI-MS: M⁺ = 539

45

Example 50:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(N-methyl-N-(2-(3,4-dimethoxyphenyl)ethyl)carbamoylmethoxy)-3,3-diphenylpropionic acid

5

¹H-NMR (200 MHz): 7.50-7.25 ppm (10 H, m), 6.80-6.70 (3 H, m), 6.35 (1 H, m), 4.50-4.00 (2 H, m, rotamers), 3.75 (3 H, s, rotamers), 3.50-2.70 (5 H, m, rotamers), 2.30/2.25 (6 H, s, rotamers).

10

ESI-MS: M⁺ = 599

Example 51:

15 2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(3,4-dimethoxybenzoyl-N-methylamino)ethoxy)-3,3-diphenylpropionic acid (II-78)

¹H-NMR (200 MHz): 7.30-7.00 ppm (10 H, m), 7.00-6.80 (3 H, m), 6.60 (1 H, s), 6.20 (1 H, s), 3.90 (6 H, s), 3.90-3.50 (4 H, m),
20 3.10 (3 H, s), 2.30 (6 H, s).

ESI-MS: M⁺ = 585

Example 52:

25

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(2,6-dimethoxybenzoyl-N-methylamino)ethoxy)-3,3-diphenylpropionic acid (II-88)

¹H-NMR (200 MHz): 7.50-7.00 ppm (10 H, m), 6.70-6.40 (4 H, m),
30 6.30/6.20 (1 H, s, rotamers), 4.10-3.30 (4 H, m), 3.80/3.75/3.65/3.60 (6 H, s, rotamers), 3.10/2.80 (3 H, s), 2.35/2.30 (6 H, s).

ESI-MS: M⁺ = 585

35

Example 53:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(3,4-dichlorobenzoyl-amino)ethoxy)-3,3-diphenylpropionic acid (II-115)

40

ESI-MS: 580 (M+H)⁺

¹H-NMR (270 MHz, DMSO): 12.0-13.0 ppm (1 H, brd); 8.80 (1 H, t); 7.15-7.65 (13 H, m); 6.95 (1 H, s); 6.20 (1 H, s); 3.85 (1 H, m);
45 3.65-3.80 (1 H, m); 3.45-3.60 (2 H, m); 2.30 (6 H, s).

Example 54:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(2,6-dimethoxybenzoyl-amino)ethoxy)-3,3-diphenylpropionic acid (II-122)

5

ESI-MS: 572 (M+H)⁺

¹H-NMR (270 MHz, CDCl₃): 7.45-7.55 ppm (2 H, m); 7.20-7.40 (10 H, m); 6.65 (1 H, s); 6.55 (1 H, d); 6.35 (1 H, t); 6.25 (1 H, s);
10 3.60-3.90 (4 H, m); 3.80 (6 H, s); 2.35 (6 H, s).

Example 55:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(2,4,6-trimethylbenzoyl-
15 amino)ethoxy)-3,3-diphenylpropionic acid (II-169)

ESI-MS: 554 (M+H)⁺

¹H-NMR (270 MHz, CDCl₃): 7.15-7.55 ppm (10 H, m); 6.90 (1 H, s);
20 6.80 (1 H, s); 6.70 (1 H, s); 6.60 (1 H, tr); 6.25 (1 H, s);
3.60-3.80 (2 H, m); 2.30 (6 H, s); 2.20 (6 H, s); 2.15 (3 H, s).

Example 56:

25 2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(2,3-dimethylbenzoyl-amino)ethoxy)-3,3-diphenylpropionic acid (II-190)

ESI-MS: 540 (M+H)⁺

30 ¹H-NMR (200 MHz, DMSO): 8.30 ppm (1 H, t); 7.10-7.55 ppm (13 H, m); 6.95 (1 H, s); 6.15 (1 H, s); 3.85-4.00 (1 H, m); 3.65-3.80 (1 H, m); 3.45-3.60 (2 H, m); 2.35 (6 H, s); 2.30 (3 H, s); 2.25 (3 H, s).

35 Example 57:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(3,5-dichlorobenzoyl-amino)ethoxy)-3,3-diphenylpropionic acid (II-205)

40 ESI-MS: 580 (M+H)⁺

¹H-NMR (200 MHz, DMSO): 12.4-13.0 ppm (1 H, brd); 8.80 (1 H, tr);
7.80 (2 H, m); 7.75 (1 H, m); 7.10-7.45 (10 H, m); 6.90 (1 H, s);
6.15 (1 H, s); 3.80-4.00 (1 H, m); 3.60-3.80 (1 H, m); 3.45-3.60
45 (2 H, m); 2.30 (6 H, s).

Example 58:

2-(4,6-Dimethylpyrimidin-2-yloxy)-3-(2-(1-naphthoylamino)ethoxy)-
3,3-diphenylpropionic acid (II-210)

5

ESI-MS: 562 (M+H)⁺

¹H-NMR (200 MHz, DMSO): 12.4-13.0 ppm (1 H, brd); 8.70 (1 H, tr);
8.20-8.30 (1 H, m); 7.85-8.80 (2 H, m); 7.10-7.60 (14 H, m); 6.90
10 (1 H, s); 6.15 (1 H, s); 3.80-4.00 (1 H, m); 3.65-3.80 (1 H, m);
3.50-3.60 (2 H, m); 2.30 (3 H, s).

The compounds in Table I can be prepared in a similar way or as
described in the general part.

15

20

25


30

35

40

45

Table I



No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-1	COOH	Phenyl	Me ₂ N	Me	Me	Me	Me	CH	N	N	O
I-2	COOH	Phenyl	Me ₂ N	H	H	OMe	OMe	CH	N	N	S
I-3	COOH	Phenyl	Me ₂ N	H	H	OMe	Me	CH	N	N	O
I-4	COOH	Phenyl	Me ₂ N	H	H	CH ₂ OH	Me	CH	N	N	O
I-5	COOMe	Phenyl	Me ₂ N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-6	COOH	Phenyl	Me ₂ N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-7	COOH	Phenyl	Me ₂ N	H	H	Ethyl	Ethyl	CH	N	N	O
I-8	COOH	Phenyl	Me ₂ N	H	H	CF ₃	Me	CH	N	N	S
I-9	COOH	Phenyl	Me ₂ N	H	H	OMe	CF ₃	CH	N	N	O
I-10	COOH	Phenyl	Butyl-HN	H	H	Me	Me	CH	N	N	O
I-11	COOH	Phenyl	Butyl-HN	Me	H	OMe	OMe	CH	N	N	S
I-12	COOH	Phenyl	Butyl-HN	H	H	OMe	Me	CH	N	N	O
I-13	COOH	Phenyl	Butyl-HN	Butyl	H	CH ₂ OH	Me	CH	N	N	O
I-14	COOH	Phenyl	Butyl-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-15	COOH	Phenyl	Butyl-HN	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-16	COOH	Phenyl	Butyl-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-17	COOH	4-F-Phenyl	Butyl-HN	H	H	CF ₃	Me	CH	N	N	S
I-18	COOH	Phenyl	Butyl-HN	H	H	OMe	CF ₃	CH	N	N	O
I-19	COOH	Phenyl	Phenyl-HN	H	H	Me	Me	CH	N	N	O
I-20	COOH	Phenyl	Phenyl-HN	H	H	OMe	OMe	CH	N	N	S
I-21	COOH	Phenyl	Phenyl-HN	H	H	OMe	Me	CH	N	N	O
I-22	COOH	Phenyl	Phenyl-HN	H	H	CH ₂ OH	Me	CH	N	N	O
I-23	COOH	4-F-Phenyl	Phenyl-HN	Me	Me	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
I-24	COOH	Phenyl	Phenyl-HN	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-25	COOH	Phenyl	Phenyl-HN	H	H	Ethyl	Ethyl	CH	N	N	S
I-26	COOH	4-F-Phenyl	Phenyl-HN	H	H	CF ₃	Me	CH	N	N	O
I-27	COOH	4-F-Phenyl	Phenyl-HN	H	H	OMe	CF ₃	CH	N	N	O
I-28	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	Me	Me	CH	N	N	O
I-29	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	OMe	OMe	CH	N	N	O
I-30	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	OMe	Me	CH	N	N	O
I-31	COOMe	Phenyl	(4-Methoxyphenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	O
I-32	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
I-33	COOH	Phenyl	(4-Methoxyphenyl)-HN	Me	Me	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-34	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	S
I-35	COOH	Phenyl	(4-Methoxyphenyl)-HN	H	H	CF ₃	Me	CH	N	N	O
I-36	COOH	4-F-Phenyl	(4-Methoxyphenyl)-HN	H	H	OMe	CF ₃	CH	N	N	O
I-37	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	Me	Me	CH	N	N	O
I-38	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	OMe	OMe	CH	N	N	S
I-39	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	OMe	Me	CH	N	N	O
I-40	COOH	4-F-Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	CH ₂ OH	Me	CH	N	N	O
I-41	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-42	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-43	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-44	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	H	H	CF ₃	Me	CH	N	N	O
I-45	COOH	Phenyl	(4-Methoxyphenyl)-HN [sic]	Ethyl	H	OMe	CF ₃	CH	N	N	O
I-46	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	Me	Me	CH	N	N	O
I-47	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	OMe	OMe	CH	N	N	O
I-48	COOH	Phenyl	(4-Chlorophenyl)-HN	Me	H	OMe	Me	CH	N	N	O
I-49	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	S
I-50	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-51	COOMe	Phenyl	(4-Chlorophenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-52	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-53	COOH	Phenyl	(4-Chlorophenyl)-HN	H	H	CF ₃	Me	CH	N	N	O
I-54	COOH	4-F-Phenyl	(4-Chlorophenyl)-HN	H	H	OMe	CF ₃	CH	N	N	O
I-55	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	Me	Me	CH	N	N	O
I-56	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	OMe	OMe	CH	N	N	S
I-57	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	OMe	Me	CH	N	N	O
I-58	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	O
I-59	COOH	4-F-Phenyl	(3,4-Dichlorophenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-60	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	Me	Me	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-61	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-62	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	Me	Me	CF ₃	Me	CH	N	N	O
I-63	COOH	Phenyl	(3,4-Dichlorophenyl)-HN	H	H	OMe	CF ₃	CH	N	N	O
I-64	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	Me	Me	CH	N	N	O
I-65	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	OMe	OMe	CH	N	N	O
I-66	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	Me	H	OMe	Me	CH	N	N	O
I-67	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-68	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH ₂ -CH ₂ -CH ₂ -C	N	N	O
I-69	COOH	4-F-Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C	O-CH ₂ -CH ₂ -C	N	N	S
I-70	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-71	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	CF ₃	Me	CH	N	N	O
I-72	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	OMe	CF ₃	CH	N	N	O
I-73	COOH	Phenyl	(3,4-Dimethoxyphenyl)-HN	H	H	Me	Me	CH	N	N	O
I-74	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	Me	Me	OMe	OMe	CH	N	N	O
I-75	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	OMe	Me	CH	N	N	O
I-76	COOH	4-F-Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	O
I-77	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH ₂ -CH ₂ -CH ₂ -C	N	N	S
I-78	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C	O-CH ₂ -CH ₂ -C	N	N	O
I-79	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-80	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	CF ₃	Me	CH	N	N	O
I-81	COOH	Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	OMe	CF ₃	CH	N	N	S
I-82	COOMe	Phenyl	(2,6-Diethylphenyl)-HN	H	H	Me	Me	CH	N	N	O
I-83	COOH	Phenyl	(2,6-Diethylphenyl)-HN	H	H	OMe	OMe	CH	N	N	O
I-84	COOH	4-F-Phenyl	(2,6-Diethylphenyl)-HN	H	H	OMe	Me	CH	N	N	O
I-85	COOH	Phenyl	(2,6-Diethylphenyl)-HN	Me	Me	CH ₂ OH	Me	CH	N	N	O
I-86	COOH	Phenyl	(2,6-Diethylphenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH ₂ -CH ₂ -CH ₂ -C	N	N	O
I-87	COOH	Phenyl	(2,6-Diethylphenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C	O-CH ₂ -CH ₂ -C	N	N	O
I-88	COOH	Phenyl	(2,6-Diethylphenyl)-HN	Et	H	Ethyl	Ethyl	CH	N	N	O
I-89	COOH	Phenyl	(2,6-Diethylphenyl)-HN	H	H	CF ₃	Me	CH	N	N	O
I-90	COOH	Phenyl	(2,6-Diethylphenyl)-HN	H	H	OMe	CF ₃	CH	N	N	O
I-91	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	Me	Me	CH	N	N	O
I-92	COOH	4-F-Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	OMe	OMe	CH	N	N	O
I-93	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	OMe	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-94	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	CH ₂ OH	Me	CH	N	N	O
I-95	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	Et	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-96	COOMe	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-97	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-98	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	H	H	CF ₃	Me	CH	N	N	S
I-99	COOH	Phenyl	(2,6-Diisopropylphenyl)-HN	Me	Me	OMe	CF ₃	CH	N	N	O
I-100	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	Me	Me	CH	N	N	O
I-101	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	O
I-102	COOH	4-F-Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-103	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-104	COOH	Phenyl	(N-Butyl-N-Me)-N	Butyl	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-105	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-106	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-107	COOH	4-F-Phenyl	(N-Butyl-N-Me)-N	H	H	CF ₃	Me	CH	N	N	O
I-108	COOH	Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-109	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	Me	Me	CH	N	N	O
I-110	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	O
I-111	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-112	COOH	Phenyl	(N-Phenyl-N-Me)-N	Ethyl	H	CH ₂ OH	Me	CH	N	N	O
I-113	COOH	4-F-Phenyl	(N-Phenyl-N-Me)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-114	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-115	COOH	Phenyl	(N-Phenyl-N-Me)-N	Ethyl	H	Ethyl	Ethyl	CH	N	N	O
I-116	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	CF ₃	Me	CH	N	N	O
I-117	COOH	Phenyl	(N-Phenyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-118	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	Me	Me	CH	N	N	O
I-119	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	OMe	OMe	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-120	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	OMe	Me	CH	N	N	S
I-121	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	Me	Me	CH ₂ OH	Me	CH	N	N	O
I-122	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-123	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-124	COOMe	4-F-Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-125	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-126	COOH	Phenyl	(N-4-Methylphenyl-N-Methyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-127	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	Me	Me	CH	N	N	O
I-128	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	OMe	OMe	CH	N	N	O
I-129	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	OMe	Me	CH	N	N	O
I-130	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	Me	Me	CH ₂ OH	Me	CH	N	N	O
I-131	COOH	4-F-Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-132	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-133	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-134	COOH	4-F-Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	CF ₃	Me	CH	N	N	O
I-135	COOH	Phenyl	(N-4-Methoxyphenyl-N-Me)-N [sic]	H	H	OMe	CF ₃	CH	N	N	O
I-136	COOMe	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	Me	Me	CH	N	N	O
I-137	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	Butyl	H	OMe	OMe	CH	N	N	O
I-138	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	Me	CH	N	N	O
I-139	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	CH ₂ OH	Me	CH	N	N	O
I-140	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-141	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-142	COOMe	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-143	COOH	4-F-Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	CF ₃	Me	CH	N	N	O
I-144	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	CF ₃	CH	N	N	O
I-145	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	Me	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-146	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	O
I-147	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	Me	Me	OMe	Me	CH	N	N	O
I-148	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-149	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-150	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-151	COOH	4-F-Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	Me	Me	Ethyl	Ethyl	CH	N	N	S
I-152	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	CF ₃	Me	CH	N	N	O
I-153	COOH	Phenyl	(N-3,4-Dichlorophenyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-154	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	Ethyl	H	Me	Me	CH	N	N	O
I-155	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	S
I-156	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-157	COOH	4-F-Phenyl	(N-4-Chlorophenyl-N-Me)-N	Ethyl	H	CH ₂ OH	Me	CH	N	N	O
I-158	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-159	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-160	COOH	4-F-Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-161	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	Me	Me	CF ₃	Me	CH	N	N	O
I-162	COOH	Phenyl	(N-4-Chlorophenyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-163	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	Me	Me	CH	N	N	S
I-164	COOH	4-F-Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	OMe	CH	N	N	O
I-165	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	Me	H	OMe	Me	CH	N	N	O
I-166	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	CH ₂ OH	Me	CH	N	N	O
I-167	COOH	4-F-Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-168	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-169	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-170	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	CF ₃	Me	CH	N	N	O
I-171	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Me)-N [sic]	H	H	OMe	CF ₃	CH	N	N	O









No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-172	COOMe	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	Me	Me	CH	N	N	S
I-173	COOH	4-F-Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	O
I-174	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-175	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-176	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	Ethyl	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
I-177	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-178	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-179	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	CF ₃	Me	CH	N	N	O
I-180	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-181	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	Me	Me	CH	N	N	O
I-182	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	OMe	OMe	CH	N	N	O
I-183	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-184	COOMe	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-185	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
I-186	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	Me	Me	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-187	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	Ethyl	Ethyl	CH	N	N	S
I-188	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	CF ₃	Me	CH	N	N	O
I-189	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Me)-N	H	H	OMe	CF ₃	CH	N	N	O
I-190	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-191	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	OMe	OMe	CH	N	N	O
I-192	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-193	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	Me	Me	CH ₂ OH	Me	CH	N	N	O
I-194	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	S
I-195	COOH	4-F-Phenyl	(N-Phenyl-N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-196	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-197	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	Me	H	CF ₃	Me	CH	N	N	O








No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-198	COOH	Phenyl	(N-Phenyl-N-Butyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-199	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-200	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	Me	H	OMe	OMe	CH	N	N	O
I-201	COOMe	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-202	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-203	COOH	4-F-Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-204	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-205	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-206	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	Me	Me	CF ₃	Me	CH	N	N	O
I-207	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-208	COOH	Phenyl	(N-4-Methylphenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-209	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	Me	Me	OMe	OMe	CH	N	N	O
I-210	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-211	COOH	4-F-Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-212	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	Me	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-213	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-214	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-215	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-216	COOH	Phenyl	(N-4-Methoxyphenyl-N-Butyl)-N	Bu	H	OMe	CF ₃	CH	N	N	O
I-217	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-218	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	Propyl	H	OMe	OMe	CH	N	N	O
I-219	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-220	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-221	COOH	4-F-Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-222	COOH	Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-223	COOH	4-F-Phenyl	(N-3,4-Dimethoxyphenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-224	COOH	Phenyl	(N-3,4-Dimethoxyphenyl)-(N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	S
I-225	COOH	Phenyl	(N-3,4-Dimethoxyphenyl)-(N-Butyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-226	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-227	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	Me	Me	OMe	OMe	CH	N	N	O
I-228	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-229	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	S
I-230	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-231	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-232	COOMe	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-233	COOH	4-F-Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-234	COOH	Phenyl	(N-3,4-Dichlorophenyl)-(N-Butyl)-N	Propyl	H	OMe	CF ₃	CH	N	N	O
I-235	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-236	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	OMe	OMe	CH	N	N	O
I-237	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	Me	Me	OMe	Me	CH	N	N	O
I-238	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	S
I-239	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-240	COOH	4-F-Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-241	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-242	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-243	COOH	Phenyl	(N-4-Chlorophenyl)-(N-Butyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-244	COOH	Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	H	H	Me	Me	CH	N	N	O
I-245	COOH	Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	Butyl	H	OMe	OMe	CH	N	N	S
I-246	COOH	4-F-Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	H	H	OMe	Me	CH	N	N	O
I-247	COOH	Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	H	H	CH ₂ OH	Me	CH	N	N	O
I-248	COOH	Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-249	COOH	Phenyl	(N-2,6-Dimethoxyphenyl)-(N-Butyl)-N [sic]	Butyl	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O





No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-250	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Butyl)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-251	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Butyl)-N [sic]	H	H	CF ₃	Me	CH	N	N	O
I-252	COOH	Phenyl	(N-2,6-Dimethoxyphenyl-N-Butyl)-N [sic]	H	H	OMe	CF ₃	CH	N	N	O
I-253	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	O
I-254	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	OMe	OMe	CH	N	N	O
I-255	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	Me	Me	OMe	Me	CH	N	N	O
I-256	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-257	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-258	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	Ethyl	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-259	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-260	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-261	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Butyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-262	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	Me	Me	CH	N	N	S
I-263	COOMe	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	OMe	OMe	CH	N	N	O
I-264	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	OMe	Me	CH	N	N	O
I-265	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	CH ₂ OH	Me	CH	N	N	O
I-266	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-267	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
I-268	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	Ethyl	Ethyl	CH	N	N	O
I-269	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-270	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Butyl)-N	Butyl	H	OMe	CF ₃	CH	N	N	O
I-271	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	Me	Me	CH	N	N	O
I-272	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	OMe	OMe	CH	N	N	O
I-273	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	OMe	Me	CH	N	N	O
I-274	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	Me	Me	CH ₂ OH	Me	CH	N	N	O
I-275	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-276	COOH	4-F-Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	OMe		O-CH ₂ -CH ₂ -C	N	N	O
I-277	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	Ethyl	Ethyl	CH	N	N	S
I-278	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	CF ₃	Me	CH	N	N	O
I-279	COOH	Phenyl	(N-Phenyl-N-Isopropyl)-N	H	H	OMe	CF ₃	CH	N	N	O
I-280	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	Me	Me	CH	N	N	O
I-281	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	OMe	CH	N	N	O
I-282	COOH	4-F-Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	Me	CH	N	N	⊖
I-283	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	CH ₂ OH	Me	CH	N	N	O
I-284	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	Propyl	H	OMe		CH ₂ -CH ₂ -CH ₂ -C	N	N	O
I-285	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe		O-CH ₂ -CH ₂ -C	N	N	O
I-286	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	Ethyl	Ethyl	CH	N	N	S
I-287	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	CF ₃	Me	CH	N	N	O
I-288	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	Me	Me	OMe	CF ₃	CH	N	N	O
I-289	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	Me	Me	CH	N	N	O
I-290	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	OMe	CH	N	N	O
I-291	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	Me	CH	N	N	O
I-292	COOH	4-F-Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	CH ₂ OH	Me	CH	N	N	O
I-293	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe		CH ₂ -CH ₂ -CH ₂ -C	N	N	O
I-294	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe		O-CH ₂ -CH ₂ -C	N	N	O
I-295	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	Me	H	Ethyl	Ethyl	CH	N	N	S
I-296	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	CF ₃	Me	CH	N	N	O
I-297	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	CF ₃	CH	N	N	O
I-298	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	Me	Me	CH	N	N	O
I-299	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	OMe	CH	N	N	O
I-300	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	OMe	Me	CH	N	N	S
I-301	COOH	4-F-Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -N)	H	H	CH ₂ OH	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-302	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -)N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-303	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -)N	Propyl	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-304	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -)N	H	H	Ethyl	Ethyl	CH	N	N	O
I-305	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -)N	H	H	CF ₃	Me	CH	N	N	O
I-306	COOH	Phenyl	(-CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -)N	H	H	OMe	CF ₃	CH	N	N	S
I-307	COOH	Phenyl		H	H	Me	Me	CH	N	N	O
I-308	COOH	Phenyl		H	H	OMe	OMe	CH	N	N	O
I-309	COOH	Phenyl		H	H	OMe	Me	CH	N	N	O
I-310	COOH	Phenyl		H	H	CH ₂ OH	Me	CH	N	N	S
I-311	COOH	4-F-Phenyl		H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-312	COOH	Phenyl		Me	Me	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-313	COOH	Phenyl		H	H	Ethyl	Ethyl	CH	N	N	O
I-314	COOH	Phenyl		H	H	CF ₃	Me	CH	N	N	O
I-315	COOH	Phenyl		H	H	OMe	CF ₃	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-316	COOH	Phenyl		H	H	Me	Me	CH	N	N	O
I-317	COOH	4-F-Phenyl		H	H	OMe	OMe	CH	N	N	O
I-318	COOH	Phenyl		H	H	OMe	Me	CH	N	N	O
I-319	COOH	Phenyl		Ethyl	H	CH ₂ OH	Me	CH	N	N	O
I-320	COOH	Phenyl		H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	S
I-321	COOH	Phenyl		H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
I-322	COOH	Phenyl		H	H	Ethyl	Ethyl	CH	N	N	O
I-323	COOH	Phenyl		H	H	CF ₃	Me	CH	N	N	O
I-324	COOH	Phenyl		H	H	OMe	CF ₃	CH	N	N	O
I-325	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	Me	Me	CH	N	N	O
I-326	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	OMe	OMe	CH	N	N	O
I-327	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	OMe	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-328	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	CH ₂ OH	Me	CH	N	N	O
I-329	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-330	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-331	COOH	4-F-Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	S
I-332	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	CF ₃	Me	CH	N	N	O
I-333	COOH	Phenyl	(N-2,6-Diethylphenyl-N-Ethoxymethylene)-N [sic]	H	H	OMe	CF ₃	CH	N	N	O
I-334	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	Me	Me	CH	N	N	O
I-335	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	OMe	OMe	CH	N	N	O
I-336	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	Me	Me	OMe	Me	CH	N	N	O
I-337	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	CH ₂ OH	Me	CH	N	N	S
I-338	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-339	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-340	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	Ethyl	Ethyl	CH	N	N	O
I-341	COOH	Phenyl	(N-2,6-Diisopropylphenyl-N-Methoxymethylene)-N [sic]	H	H	CF ₃	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-342	COOH	Phenyl		H	H	Me	Me	CH	N	N	O
I-343	COOH	Phenyl		H	H	OMe	OMe	CH	N	N	O
I-344	COOH	Phenyl		H	H	OMe	Me	CH	N	N	O
I-345	COOH	Phenyl		H	H	CH ₂ OH	Me	CH	N	N	S
I-346	COOH	Phenyl		H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-347	COOH	Phenyl	Bu ₂ N	H	H	Me	Me	CH	N	N	O
I-348	COOH	Phenyl	Bu ₂ N	Me	H	OMe	OMe	CH	N	N	S
I-349	COOH	Phenyl	Bu ₂ N	H	H	OMe	Me	CH	N	N	O
I-350	COOH	Phenyl	Bu ₂ N	Butyl	H	CH ₂ OH	Me	CH	N	N	O
I-351	COOH	Phenyl	Bu ₂ N	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-352	COOH	Phenyl	Bu ₂ N	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-353	COOH	Phenyl	Bu ₂ N	H	H	Ethyl	Ethyl	CH	N	N	O
I-354	COOH	4-F-Phenyl	Bu ₂ N	H	H	CF ₃	Me	CH	N	N	S
I-355	COOH	Phenyl, 4 Cl-Phenyl	Me ₂ N	Me	Me	Me	Me	CH	N	N	O
I-356	COOH	4-Cl-Phenyl, 4-F-Phenyl	Phenyl-HN	H	H	OMe	OMe	CH	N	N	S
I-357	COOH	4-F-Phenyl, Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-358	COOH	4-Me-Phenyl, Naphthyl	(N-Phenyl-N-Me)-N	H	H	CH ₂ OH	Me	CH	N	N	O



No.	R ¹	R ⁴ , R ⁵	R ²²	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
I-359	COOMe	2-F-Phenyl, Phenyl	(4-Chlorophenyl)-HN	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
I-360	COOH	2-F-Phenyl, 4-Me-Phenyl	(2,6-Dimethoxyphenyl)-HN	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
I-361	COOH	Naphthyl, Phenyl	(2,6-Diethylphenyl)-HN	H	H	Ethyl	Ethyl	CH	N	N	O
I-362	COOH	Phenyl, 4 Cl Phenyl		H	H	CF ₃	Me	CH	N	N	S
I-363	COOH	4-Cl-Phe- nyl, 4-F- Phenyl		H	H	OMe	CF ₃	CH	N	N	O
I-364	COOH	Naphthyl, Naphthyl	Me ₂ N	Me	Me	Me	Me	CH	N	N	O
I-365	COOH	Naphthyl, Naphthyl	Phenyl-HN	H	H	OMe	OMe	CH	N	N	S
I-366	COOH	4-F-Phenyl, 4 Cl Phenyl	(N-Butyl-N-Me)-N	H	H	OMe	Me	CH	N	N	O
I-367	COOH	4-F-Phenyl, Phenyl	Bu ₂ N	H	H	Me	Me	CH	N	N	O

Table II



No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-1	COOH	Phenyl	(N-(2-OMe-Phenyl(CO)-N-Propyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-2	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Propyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-3	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Propyl)-N-CH ₂ -	H	H	OMe	OMe	CF	N	N	O
II-4	COOMe	Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH ₂ -CH ₂ -CH ₂ -C	N	N	O
II-5	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	O-CH ₂ -CH ₂ -C	N	N	O
II-6	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-7	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	S
II-8	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-9	COOH	Phenyl, 4 Cl	(MeCO-N-Me)-N-CH ₂ -	Me	Me	Me	Me	CH	N	N	O
II-10	COOH	Phenyl	(N-Phenyl(CO-N-Butyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-11	COOH	Phenyl	(N-Phenyl(CO-N-Propyl)-N-CH ₂ -	Me	H	OMe	OMe	CH	N	N	S
II-12	COOH	Phenyl	(N-Phenyl(CO-N-Propyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-13	COOH	Phenyl	(N-Phenyl(CO-N-Me)-N-CH ₂ -	Butyl	H	CH ₂ OH	Me	CH	N	N	O
II-14	COOH	Phenyl	(N-Phenyl(CO-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH ₂ -CH ₂ -CH ₂ -C	N	N	O
II-15	COOH	4-Cl-Phenyl, 4-F-Phenyl	(N-Phenyl(SO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	S

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-16	COOH	Phenyl	(N-Phenyl(CO-N-Me)-N-CH ₂ -	H	H	OMe		O-CH ₂ -CH ₂ -C	N	N	O
II-17	COOH	Phenyl	(N-Phenyl(CO-N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-18	COOH	4-F-Phenyl	(N-(4-OMe-Phenyl(CO)-N-Butyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	S
II-19	COOH	Phenyl	(N-(3-OMe-Phenyl(CO)-N-Propyl)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-20	COOH	Phenyl	(N-Phenyl(SO ₂ -N-Butyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-21	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	S
II-22	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-23	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-24	COOH	4-F-Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	Me	Me	OMe		CH ₂ -CH ₂ -CH ₂ -C	N	N	O
II-25	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe		O-CH ₂ -CH ₂ -C	N	N	O
II-26	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	S
II-27	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-28	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-29	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	Me	Me	N	CH	N	O
II-30	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CF	N	N	O
II-31	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-32	COOH	Phenyl	Phenyl-CH ₂ -O-CO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-33	COOH	4-F-Phenyl, Phenyl	(N-(3-OMe-Phenyl(CO)-N-Propyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-34	COOMe	Phenyl	2,6-Di-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-35	COOH	Phenyl	2,6-Di-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	OMe		CH ₂ -CH ₂ -CH ₂ -C	N	N	O
II-36	COOH	Phenyl	2,6-Di-OMe-Phenyl(CO)-HN-CH ₂ -	Me	Me	OMe		O-CH ₂ -CH ₂ -C	N	N	O
II-37	COOH	Phenyl	2,6-Di-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	S
II-38	COOH	Phenyl	2,5-Di-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-39	COOH	4-F-Phenyl	2,4-Di-OMe-Phenyl(CO)-HN-CH ₂ (Me)-	H	H	OMe	CF ₃	CH	N	N	O
II-40	COOH	Phenyl	2,3-Di-Me-Phenyl(CO)-HN-CH ₂ -	H	H	Me	Me	CF	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-41	COOH	Phenyl	2,3-Di-Me-PhenylCO-HN-CH ₂ -	H	H	OMe	OMe	CH	N	N	S
II-42	COOH	4-Me-Phenyl, Naphthyl	(N-PhenylCO-N-Butyl)-N-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-43	COOH	Phenyl	3,4-Di-Me-PhenylCO-HN-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-44	COOH	4-F-Phenyl	3,5-Di-OMe-PhenylCO-HN-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-45	COOH	Phenyl	3,4-Di-OMe-PhenylCO-HN-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-46	COOH	Phenyl	(N-PhenylISO ₂ -N-Propyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-47	COOH	Phenyl	(N-PhenylISO ₂ -N-Propyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-48	COOH	Phenyl	(N-PhenylISO ₂ -N-Propyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-49	COOH	Phenyl	(N-PhenylISO ₂ -N-Propyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-50	COOH	Phenyl	(N-PhenylISO ₂ -N-Ethyl)-N-CH ₂ -	Ethyl	H	OMe	CF ₃	CH	N	N	O
II-51	COOH	Phenyl	(N-PhenylISO ₂ -N-Ethyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-52	COOH	Phenyl	(N-PhenylISO ₂ -N-Ethyl)-N-CH ₂ -	H	H	H	OMe	CH	N	N	O
II-53	COOH	Phenyl	(N-PhenylISO ₂ -N-Ethyl)-N-CH(Me)-	Me	H	OMe	Me	CH	N	N	O
II-54	COOH	Phenyl	(N-PhenylISO ₂ -N-Ethyl)-N-CH ₂ -	H	H	CH ₂ F	Me	CH	N	N	S
II-55	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-56	COOMe	Phenyl	(N-(3-Cl-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-57	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-58	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-59	COOH	4-F-Phenyl	(N-(3-Cl-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-60	COOH	Phenyl	(N-(3-Cl-4-Me-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-61	COOH	Phenyl	(N-(3,4-Di-Cl-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	S
II-62	COOH	Phenyl	3,4-Di-OMe-PhenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-63	COOH	Phenyl	(N-(3,4-Di-Cl-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-64	COOH	Phenyl	(N-(3,4-Di-Cl-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	H	Me	C-Me	N	N	O
II-65	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-PhenylISO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-66	COOH	Phenyl	(N-(2,6-Di-OMe-PhenylSO ₂)-N-Ethyl)-N-CH ₂ -	Me	Me	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-67	COOMe	2-F-Phenyl, Phenyl	(N-(3-Cl-4-Me-PhenylSO ₂)-N-Ethyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-68	COOH	Phenyl	(N-(2,6-Di-OMe-PhenylSO ₂)-N-Ethyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	N	CH	N	O
II-69	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	Me	Me	CF ₃	Me	CH	N	N	O
II-70	COOH	Phenyl	(N-(3-Cl-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-71	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-72	COOH	Phenyl	(N-(3-OMe-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-73	COOH	Phenyl	(N-(3-Cl-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	Me	H	OMe	Me	CH	N	N	O
II-74	COOH	Phenyl	(N-(3-Cl-4-Me-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	H	OMe	C-Me	N	N	O
II-75	COOH	Phenyl	(N-(3,4-Di-Cl-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-76	COOH	4-F-Phenyl	(N-(3,4-Di-Cl-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-77	COOH	Phenyl	(N-(2,6-Di-OMe-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-78	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl)-CO)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-79	COOH	Phenyl	(N-(2,6-Di-OMe-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-80	COOH	Phenyl	(N-(2,6-Di-OMe-PhenylSO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-81	COOH	Phenyl	PhenylSO ₂ -HN-CH(Benzyl)-	H	H	Me	Me	CH	N	N	O
II-82	COOH	Phenyl	PhenylSO ₂ -HN-CH ₂ -	Me	Me	OMe	OMe	CH	N	N	O
II-83	COOH	2-F-Phenyl, 4-Me-Phenyl	3-HOOCCH ₂ O-4-OMe-PhenylCO-HN-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-84	COOH	Phenyl	PhenylSO ₂ -HN-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-85	COOH	4-F-Phenyl	PhenylSO ₂ -HN-CH ₂ -	H	H	H	OMe	C-OMe	N	N	O
II-86	COOH	Phenyl	PhenylSO ₂ -HN-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	S
II-87	COOH	Phenyl	(N-ButylSO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-88	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl)-CO)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-89	COOH	Phenyl	(N-ButylSO ₂ -N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-90	COOH	Phenyl	(N-ButylSO ₂ -N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-91	COOH	Phenyl	(N-MeSO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	S
II-92	COOMe	Phenyl	(N-MeSO ₂ -N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-93	COOH	Phenyl	(N-MeSO ₂ -N-Me)-N-CH(iso-Propyl)-	H	H	OMe	OMe	CH	N	N	O
II-94	COOH	4-F-Phenyl	(N-MeSO ₂ -N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-95	COOH	Phenyl	(N-MeSO ₂ -N-Me)-N-CH ₂ -	Me	Me	H	Me	CH	N	N	O
II-96	COOH	Phenyl	(N-MeSO ₂ -N-Ethyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-97	COOH	Naphthyl, Phenyl	2,3-Di-Me-PhenylCO-HN-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-98	COOH	Phenyl	(N-MeSO ₂ -N-Ethyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
II-99	COOH	Phenyl	(N-MeSO ₂ -N-Ethyl)-N-CH ₂ -	Et	H	Ethyl	Ethyl	CH	N	N	O
II-100	COOH	Phenyl	(N-MeSO ₂ -N-Ethyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-101	Tetrazole [sic]	Phenyl	(N-MeSO ₂ -N-Ethyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-102	COOH	Phenyl	(N-ButylSO ₂ -N-Propyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-103	COOH	4-F-Phenyl	(N-PhenylCO-N-Ethyl)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-104	COOH	Phenyl	(N-PhenylCO-N-Ethyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-105	COOH	Phenyl	(N-(4-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	H	H	Me	Me	CF	N	N	O
II-106	COOH	Phenyl	(N-MeCO-N-(4-OMe-3-Me-Phenyl))-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-107	COOH	Phenyl	(N-(3-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	Et	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-108	COOMe	Phenyl	(N-(2-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	S
II-109	COOH	Phenyl	(N-(3,4-Di-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-110	COOH	Phenyl	(N-(3,4-Di-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	S
II-111	COOH	Phenyl	(N-(3,4-Di-OMe-PhenylCO)-N-Ethyl)-N-CH ₂ -	Me	Me	OMe	CF ₃	CH	N	N	O
II-112	COOH	Phenyl	(N-(3-H ₂ NCOCH ₂ -PhenylCO)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-113	COOH	Phenyl	(N-(3-H ₂ NCOCH ₂ -PhenylCO)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-114	COOH	4-F-Phenyl	(N-(3-H ₂ NCOCH ₂ -4-Me-Phe-nyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-115	COOH	Phenyl	3,4-Di-Cl-Phenyl(CO)-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-116	COOH	Phenyl	(N-(3-H ₂ NCOCH ₂ -4-Me-Phe-nyl(CO)-N-Me)-N-CH ₂ -	H	H	H	OMe	C-Me	N	N	O
II-117	COOH	Phenyl	(N-(3-HOOCCH ₂ -4-OMe-Phe-nyl(CO)-N-Me)-N-CH ₂ -	Butyl	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-118	COOH	Phenyl	Me-CH=CH-CO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-119	COOH	Phenyl	(N-(3-HOOCCH ₂ -4-OMe-Phe-nyl(CO)-N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-120	COOH	Phenyl	(N-(3-HOOCCH ₂ -4-Cl-Phe-nyl(CO)-N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-121	COOH	4-F-Phenyl	(N-(3-HOOCCH ₂ -4-Cl-Phe-nyl(CO)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-122	COOH	Phenyl	2,6-Di-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-123	COOH	Phenyl	3-HOOCCH ₂ -4-Cl-Phenyl(CO)-HN-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-124	COOH	Phenyl	3-HOOCCH ₂ -4-Cl-Phenyl(CO)-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-125	COOH	Phenyl	3-HOOCCH ₂ -4-Cl-Phenyl(CO)-HN-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-126	COOH	Phenyl	3-HOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-127	COOH	Phenyl	3-HOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	Ethyl	H	H	OMe	CH	N	N	O
II-128	COOH	4-F-Phenyl	3-HOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-129	COOH	Phenyl	(N-MeCO-N-(4-OMe-3-Me-Phenyl))-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-130	COOH	Phenyl	3-HOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-131	COOH	Naphthyl, Naphthyl	2,6-Diethylphenyl(CO)-HN-CH ₂ -	Me	Me	Me	Me	CH	N	N	O
II-132	COOH	Phenyl	3-HOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	Ethyl	H	Ethyl	Ethyl	CH	N	N	O
II-133	COOH	Phenyl	3-MeOOCCH ₂ O-4-OMe-Phenyl(CO)-HN-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-134	COOH	Phenyl	3-MeOOCCH ₂ O-4-OMe-PhenylCO-HN-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-135	COOH	Phenyl	3-MeOOCCH ₂ O-4-OMe-PhenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-136	COOH	Phenyl	Me-CH=CH-CO-HN-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-137	COOH	Phenyl	4-MeOOCCH ₂ O-3-OMe-PhenylCO-HN-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-138	COOH	Phenyl	4-MeOOCCH ₂ O-3-OMe-PhenylCO-HN-CH ₂ -	H	H	OMe	Me	CH	N	N	S
II-139	COOH	Phenyl	4-MeOOCCH ₂ O-3-OMe-PhenylCO-HN-CH ₂ -	Me	Me	Ethyl	Me	N	CH	N	O
II-140	COOH	Phenyl	(N-(4-OMe-Phenyl)-CH ₂ -CO)-N-Butyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-141	COOH	Phenyl	(N-EthylCO-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-142	COOH	Phenyl	(N-(3-OMe-Phenyl)-CH ₂ -CO)-N-Pro- pyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
II-143	COOMe	4-F-Phenyl	(N-(2-OMe-Phenyl)-CH ₂ -CO)-N-Pro- pyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-144	COOH	Phenyl	(N-(3,4-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-145	COOH	Phenyl	(N-(3,4-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-146	COOH	Phenyl	(N-(3,4-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-147	COOH	Phenyl	(N-(3,4-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-148	COOH	Phenyl	(N-(3,4-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-149	COOH	Phenyl	(N-(2,6-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	Me	Me	CH ₂ OH	Me	CH	N	N	O
II-150	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-Phe- nyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-151	COOH	Phenyl	(N-MeCO-N-(4-Cl-Phenyl))-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-152	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-153	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-154	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-155	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-156	COOMe	Phenyl	(N-(3,4-Di-OMe-Phenyl-CH ₂ -CO)-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-157	COOH	Phenyl	Iso-Propyl(CO)-HN-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-158	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Benzyl)-N-CH ₂ -	Butyl	H	OMe	OMe	CH	N	N	O
II-159	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Benzyl)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-160	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Benzyl)-N-CH ₂ -	H	H	H	OMe	CMe	N	N	O
II-161	COOH	Phenyl	(N-(3,4-Di-OMe-Phenyl(CO)-N-Benzyl)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-162	COOH	4-F-Phenyl	(N-MeCO-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	S
II-163	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	S
II-164	COOH	Naphthyl, Naphthyl	(N-(3-Cl-4-Me-Phenyl(SO ₂)-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	S
II-165	COOMe	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-166	COOH	4-F-Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-167	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-168	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-169	COOH	Phenyl	2,4,6-Tri-Me-Phenyl(CO)-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-170	COOH	Phenyl	(N-(2,6-Di-OMe-Phenyl)-CH ₂ -CO)-N-Ethyl)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-171	COOH	Phenyl	(N-(2-Me-3-Cl-4-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	Me	Me	OMe	Me	CH	N	N	O
II-172	COOH	Phenyl	(N-(3-Me-2-Cl-4-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	H	H	H	Me	C-Me	N	N	O
II-173	COOH	Phenyl	(N-(3-Me-4-Cl-5-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-174	COOH	Phenyl	(N-(3-Me-4-Cl-5-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-175	COOH	4-F-Phenyl	(N-(3,5-Di-Me-4-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	Me	Me	Ethyl	Ethyl	CH	N	N	S
II-176	COOH	Phenyl	(N-(3,5-Di-Me-4-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-177	COOH	Phenyl	(N-(3,5-Di-Me-4-OMe-Phe-nylCO)-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-178	COOH	Phenyl	(N-PhenylCO)-N-MeOMe)-N-CH ₂ -	Ethyl	H	Me	Me	CH	N	N	O
II-179	COOH	Phenyl	(N-PhenylCO)-N-MeOMe)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	S
II-180	COOH	Phenyl	(N-(4-OMe-PhenylCO)-N-MeOMe)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-181	COOH	4-F-Phenyl	(N-(3-OMe-PhenylCO)-N-MeOMe)-N-CH ₂ -	Ethyl	H	H	OMe	CH	N	N	O
II-182	COOH	4-F-Phenyl, 4 Cl Phenyl	(N-PhenylCO)-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-183	COOH	Phenyl	(N-(2-OMe-PhenylCO)-N-MeOMe)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C		N	N	O
II-184	COOH	Phenyl	(N-MeCO)-N-Phenyl)-N-CH ₂ -	Me	Me	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-185	COOH	Phenyl	(N-(3,4-Di-OMe-Phe-nylCO)-N-MeOMe)-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C		N	N	O
II-186	COOH	4-F-Phenyl	(N-(3,4-Di-OMe-PhenylCO)-N-MeOMe-nyl)-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-187	COOH	Phenyl	(N-(3,4-Di-OMe-PhenylCO)-N-MeOBu-tyl)-N-CH ₂ -	Me	Me	CF ₃	Me	CH	N	N	O
II-188	COOH	Phenyl	(N-(3,4-Di-OMe-Phe-nylCO)-N-MeOMe)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-189	COOH	Phenyl	(N-(3,4-Di-OMe-Phe-nylCO)-N-MeOMe)-N-CH ₂ -	H	H	Me	Me	CH	N	N	S
II-190	COOH	Phenyl	2,3-Di-Me-PhenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-191	COOH	4-F-Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl-CH ₂))-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-192	COOH	Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl-CH ₂))-N-CH ₂ -	Me	H	OMe	Me	CH	N	N	O
II-193	COOH	Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl-CH ₂))-N-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-194	COOH	4-F-Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl-CH ₂))-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-195	COOH	4-F-Phenyl	PropylCO-HN-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	S
II-196	COOH	Phenyl	(N-PhenylCO)-N-(3-OMe-Phenyl-CH ₂))-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	S
II-197	COOH	Phenyl	(N-PhenylCO)-N-(2-OMe-Phenyl-CH ₂))-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O
II-198	COOH	Phenyl	(N-PhenylCO)-N-(3-Me-Phenyl-CH ₂))-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-199	COOH	Phenyl	(N-PhenylCO)-N-(4-Me-Phenyl-CH ₂))-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-200	COOH	Phenyl	(N-EthylCO)-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-201	COOMe	Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	Me	Me	CH	N	N	S
II-202	COOH	4-F-Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-203	COOH	Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-204	COOH	Phenyl	(N-PhenylCO)-N-(4-OMe-Phenyl))-N-CH ₂ -	H	H	H	OMe	C-Me	N	N	O
II-205	COOH	Phenyl	3,5-Di-Cl-PhenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-206	COOH	Phenyl	(N-PhenylCO)-N-(3-OMe-Phenyl))-N-CH ₂ -	Ethyl	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-207	COOH	Phenyl	MeCO-HN-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-208	COOH	Phenyl	(N-PhenylCO)-N-(2-OMe-Phenyl))-N-CH ₂ -	H	H	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
II-209	COOH	Phenyl	(N-PhenylCO)-N-(3-Me-Phenyl))-N-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	O

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ⁷	R ⁸	R ²	R ³	Z	X	Y	W
II-210	COOH	Phenyl	Naphthyl-1-CO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-211	COOH	Phenyl	(N-Phenyl)CO-N-(4-Me-Phenyl)-N-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-212	COOH	Phenyl	CyclohexylCO-HN-CH ₂ -	Me	Me	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
II-213	COOH	Phenyl	(N-2,6-Diethylphenyl)CO-N-Me)-N-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-214	COOH	4-F-Phenyl	(N-2,6-Diisopropylphenyl)CO-N-Me)-N-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-215	COOH	Phenyl	(N-2,6-Diisopropylphenyl)CO-N-Me)-N-CH ₂ -	H	H	OMe	OMe	CH	N	N	O
II-216	COOH	Phenyl	(N-MeCO-N-Me)-N-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-217	COOH	Phenyl	(N-2,6-Diethylphenyl)CO-N-Me)-N-CH ₂ -	H	H	OMe	Me	CH	N	N	O
II-218	COOH	4-F-Phenyl, Phenyl	2,4,6-Tri-Me-PhenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O
II-219	COOMe	Phenyl	(N-2,6-Diethylphenyl)CO-N-Me)-N-CH ₂ -	H	H	CH ₂ OH	Me	CH	N	N	O
II-220	COOH	Phenyl	2,6-DiethylphenylCO-HN-CH ₂ -	H	H	OMe	CH ₂ -CH ₂ -CH ₂ -C	CH	N	N	O
II-221	COOH	Phenyl	2,6-DiethylphenylCO-HN-CH ₂ -	Me	Me	OMe	O-CH ₂ -CH ₂ -C	CH	N	N	O
II-222	COOH	4-F-Phenyl	2,6-DiethylphenylCO-HN-CH ₂ -	H	H	Ethyl	Ethyl	CH	N	N	S
II-223	COOH	Phenyl	2,6-DiethylphenylCO-HN-CH ₂ -	H	H	CF ₃	Me	CH	N	N	O
II-224	COOH	Phenyl	2,6-DimethylphenylCO-HN-CH ₂ -	H	H	OMe	CF ₃	CH	N	N	O
II-225	COOH	Phenyl	2,6-DimethylphenylCO-HN-CH ₂ -	H	H	Me	Me	CH	N	N	O

Example 59:

Receptor binding data were measured by the binding assay
5 described above for the compounds listed below.

The results are shown in Table 3.

Table 3

10

Receptor binding data (K_i values)

15

Compound	ET _A [nM/l] [sic]	ET _B [nM/l] [sic]
I-109	0,4	142
I-111	0,3	109
I-347	3,8	155
I-349	3,0	142
20 I-307	1,6	10
I-309	1	12

25

30

35

40


45



We claim:

1. A carboxylic acid derivative of the formula I

5



where R¹ is tetrazole [sic] or a group

15



where R has the following meaning:

20 a) a radical OR⁹ where R⁹ is:

hydrogen, the cation of an alkali metal, the cation of an alkaline earth metal or a physiologically tolerated organic ammonium ion;

25

C₃-C₈-cycloalkyl, C₁-C₈-alkyl,

CH₂-phenyl, unsubstituted or substituted,

30

C₃-C₆-alkenyl or a C₃-C₆-alkynyl group, unsubstituted or substituted, or


phenyl, unsubstituted or substituted.

35

b) a 5-membered heteroaromatic system linked via a nitrogen atom.

c) a group

40




where k can assume the values 0, 1 and 2, p can assume the values 1, 2, 3 and 4, and R¹⁰ is C₁-C₄-alkyl, C₃-C₈-cycloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl.

45

d) a radical

5



where R^{11} is:

10

C_1 - C_4 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, it being possible for these radicals to carry a C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio and/or a phenyl radical;

15

phenyl, unsubstituted or substituted.

20

R^2 is hydrogen, hydroxyl, NH_2 , $\text{NH}(\text{C}_1$ - C_4 -alkyl), $\text{N}(\text{C}_1$ - C_4 -alkyl) $_2$, halogen, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy or C_1 - C_4 -alkylthio, or CR^2 is linked to CR^{12} as indicated under Z to give a 5- or 6-membered ring;

25

X is nitrogen or methine;

Y is nitrogen or methine;

30

Z is nitrogen or CR^{12} , where R^{12} is hydrogen, halogen or C_1 - C_4 -alkyl, or CR^{12} forms together with CR^2 or CR^3 a 5- or 6-membered alkylene or alkenylene ring which can be unsubstituted or substituted and in which in each case one or more methylene groups can be replaced by oxygen, sulfur, $-\text{NH}$ or $-\text{N}(\text{C}_1$ - C_4 -alkyl);

35

R^3 is hydrogen, hydroxyl, NH_2 , $\text{NH}(\text{C}_1$ - C_4 -alkyl), $\text{N}(\text{C}_1$ - C_4 -alkyl) $_2$, halogen, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio; or CR^3 is linked to CR^{12} as indicated under Z to give a 5- or 6-membered ring;

40

R^4 and R^5 (which can be identical or different) are:

45

phenyl or naphthyl, unsubstituted or substituted, or

phenyl or naphthyl which are connected together in ortho positions by a direct linkage, a methylene, ethylene or


75

ethenylene group, an oxygen or sulfur atom or an SO₂, NH or N-alkyl group,

C₃-C₈-cycloalkyl, unsubstituted or substituted;

5

R⁶ is a group



where R¹³ and R¹⁴ can be identical or different and have the following meanings:

15

hydrogen, with the proviso that R¹³ and R¹⁴ must not simultaneously be hydrogen,

20

C₁-C₈-alkyl, C₃-C₈-cycloalkyl, C₃-C₈-alkenyl, C₃-C₈-alkynyl, benzyl, phenyl, naphthyl, each unsubstituted or substituted,

25

or R¹³ and R¹⁴ together form an unsubstituted or substituted C₃-C₇-alkylene chain which is closed to a ring and in which one alkylene group can be replaced by oxygen, sulfur or nitrogen,

30

or R¹³ and R¹⁴ together form an unsubstituted or substituted C₃-C₇-alkylene chain or C₃-C₇-alkenylene chain which is closed to a ring and to which an unsubstituted or substituted phenyl ring is fused;

35

R⁷ and R⁸ (which can be identical or different) are: hydrogen, C₁-C₄-alkyl;

R¹⁸ is hydrogen;

40

C₁-C₈-alkyl, C₃-C₈-alkenyl or C₃-C₈-alkynyl, phenyl, naphthyl, C₃-C₈-cycloalkyl, where these radicals can be unsubstituted or substituted;

45

R¹⁹ is C₁-C₈-alkylcarbonyl, C₂-C₈-alkenylcarbonyl, C₂-C₈-alkynylcarbonyl, benzyloxycarbonyl, C₃-C₈-cycloalkylcarbonyl, phenylcarbonyl or naphthylcarbonyl, where said radicals can be unsubstituted or substituted;

C₁-C₈-alkylsulfonyl, C₃-C₈-alkenylsulfonyl or
C₃-C₈-alkynylsulfonyl, phenylsulfonyl or naphthyl-
sulfonyl, in each case unsubstituted or substituted;
C₃-C₈-cycloalkylsulfonyl;

5

R²⁰ is hydrogen, C₁-C₄-alkyl, unsubstituted or substituted.

R²¹ is hydrogen, C₁-C₄-alkyl.

10


W is sulfur or oxygen.



and the physiologically tolerated salts and the
enantiomerically pure and diastereomerically pure forms.


- 15 2. A drug preparation for oral, parenteral [sic] use comprising
at least one carboxylic acid derivative I as claimed in claim
1, in addition to conventional medicinal auxiliaries, per
single dose.
- 20 3. The use of a carboxylic acid derivative as claimed in claim 1
for treating diseases.
4. The use of a compound I as claimed in claim 1 as endothelin
receptor antagonist.
- 25 5. The use of a carboxylic acid derivative I as claimed in
claim 1 for producing drugs for treating diseases in which
elevated endothelin levels occur.
- 30 6. The use of a carboxylic acid derivative I as claimed in
claim 1 for producing drugs for treating diseases in which
endothelin contributes to cause and/or progression.
7. The use of a carboxylic acid derivative I as claimed in
35 claim 1 for treating chronic heart failure, restenosis, high
blood pressure, pulmonary hypertension, acute/chronic kidney
failure, cerebral ischemia, benign prostate hyperplasia and
prostate cancer.
- 40 8. A combination of carboxylic acid derivatives of the formula I
as claimed in claim 1 and one or more active substances,
selected from inhibitors of the renin-angiotensin system such
as renin inhibitors, angiotensin II antagonists, angiotensin
converting enzyme (ACE) inhibitors, mixed ACE/neutral
45 endopeptidase (NEP) inhibitors, β -blockers, diuretics,
calcium antagonists and VEGF-blocking substances.

9. The use of compounds of the formula V



10 where the radicals R^1 , R^4 , R^5 , R^6 , R^7 , R^8 and W have the meanings stated in claim 1, as starting material for synthesizing endothelin receptor antagonists.


10. A structural fragment of the formula



20 where the radicals R^1 , R^4 , R^5 , R^6 , R^7 , R^8 and W have the meanings stated in claim 1.

25 11. The use of a structural fragment as claimed in claim 10 as a structural component of an endothelin [sic] receptor antagonist.

12. An endothelin receptor antagonist, comprising a structural fragment of the formula




40 in which the radicals R^1 , R^2 , R^3 , R^4 , R^5 , R^7 , R^8 , W, X, Y and Z are as defined in claim 1, covalently linked to a group which has a molecular weight of at least 30.

13. An endothelin receptor antagonist, comprising a structural fragment of the formula


45

78



in which the radicals R¹, R², R³, R⁴, R⁵, R⁷, R⁸, R²⁰, R²¹, W, X, Y and Z are as defined in claim 1, covalently linked via a nitrogen atom to a group which has a molecular weight of at least 58.

14. A compound of the formula Ia



Ia

in which the radicals R¹, R², R³, R⁴, R⁵, R⁷, R⁸, R²⁰, R²¹, W, X, Y and Z are as defined in claim 1.

25

30

35

40


45

Novel carboxylic acid derivatives with amide side chains, their preparation and use as endothelin receptor antagonists

5 Abstract

Carboxylic acid derivatives of the formula I

10



15

where the substituents have the meanings explained in the description are used as endothelin receptor antagonists.

20

25

30

35

40

45