METHODS AND SYSTEMS FOR CONVERTING PRECURSOR CELLS INTO GASTRIC TISSUES THROUGH DIRECTED DIFFERENTIATION

Statement Regarding Federally-Sponsored Research

[0001] This invention was made with government support under DK080823, DK092456, and GM063483 awarded by the National Institutes of Health. The government has certain rights to the invention.

Priority Claim

[0002] This application claims priority to and benefit of US Provisional Patent Application No. 62/003,719, to Wells et al, filed on May 28th, 2014, entitled "Methods and Systems for Converting Precursor Cells into Gastric Tissues through Directed Differentiation" for all purposes.

Field of the Invention

[0003] Disclosed herein are methods and systems relating to converting stem cells into specific tissue(s) or organ(s) through directed differentiation. In particular, disclosed are methods and systems for promoting definitive endoderm formation from human pluripotent stem cells. Also disclosed are methods and systems for promoting gastric organoids or tissue formations from differentiated definitive endoderm.

Background

[0004] Stomach function and architecture vary widely between mammalian species, to accommodate a wide variety of habitats and diets. Consequently, non-human models of gastric development and disease have significant limitations. For example, the bacterium *Helicobacter Pylori* infects 50% of the world's

population, with 10% developing peptic ulcer disease and 1-2%[1-3] developing gastric cancer. Gastric diseases, including peptic ulcer disease and gastric cancer, affect 10% of the world's population and are largely due to chronic *H. pylori* infection. Current models of *H. pylori*-induced disease rely upon animal models that do not exhibit the same pathophysiological features as the human response to infection[4], and gastric cell lines lack the cellular and architectural complexity of the gastric epithelium *in vivo*. Thus there is no adequate model to study the effects of *H. pylori* infection as it occurs in humans. While recent advances using adult gastric stem cells allow for growth of rodent gastric epithelium *in vitro*[5], obtaining these cells from human patients would require surgery. Moreover, such a method could not be used to model embryonic development of the human stomach or stromal-epithelial interactions. Species differences in embryonic development and architecture of the adult stomach make murine models suboptimal for the study of organogenesis and pathogenesis of this organ. Thus, there is a need for robust *in vitro* systems for elucidating the mechanisms underlying human stomach development and disease and for the identification of novel treatments useful for human treatment of such diseases.

[0005] What is needed in the art are methods and systems for accurately controlling the destination of a precursor cell such as a human pluripotent stem cell, in order to create the specific type of tissue or organism desired, in particular, gastric tissues that can be used for one or more of the aforementioned purposes.

Brief Summary

[0006] Disclosed are methods of inducing formation of a gastric cells and/or a gastric tissue, such as in the form of a gastric organoid. The formation of gastric cells and/or tissue may be carried out by

the activating and/or inhibiting of one or more signaling pathways within a precursor cell. Also disclosed are methods for using the disclosed gastric cells, gastric tissues, and or gastric organoids derived from precursor cells.

[0006a] In one embodiment, there is provided a method of inducing formation of a gastric tissue, comprising the steps of:

a) activating a combination of a WNT signaling pathway and a FGF signaling pathway within a definitive endoderm derived from an induced pluripotent stem cell, an embryonic stem cell, and/or an endoderm cell, wherein said WNT signalling pathway is activated using an agent selected from one or more molecules selected from the group consisting of Wnt1, Wnt2, Wnt2b, Wnt4, Wnt5a, Wnt5b, Wnt6, Wnt7a, Wnt7b, Wnt8a, Wnt9a, Wnt10a, Wnt10b, Wnt11, Wnt16, a GSK3β inhibitor such as Chiron, and small molecule modulators that activate the WNT signalling pathway, and wherein said FGF signalling pathway is activated using an agent selected from one or more molecules selected from the group consisting of FGF1, FGF2, FGF3, FGF4, FGF10, FGF11, FGF12, FGF13, FGF14, FGF15, FGF16, FGF17, FGF18, FGF19, FGF20, FGF21, FGF22, FGF23 and small molecule modulators that activate the FGF signalling pathway,

b) inhibiting a BMP signaling pathway within said definitive endoderm derived from an induced pluripotent stem cell, embryonic stem cell, and/or endoderm cell by contacting said definitive endoderm derived from an induced pluripotent stem cell, embryonic stem cell, and/or endoderm cell with an agent that inhibits the BMP pathway, wherein said agent is selected from Noggin, Dorsomorphin, LDN189, DMH-1, or small molecule modulators that inhibit the BMP signaling pathway; and

c) administering retinoic acid to said definitive endoderm;

wherein steps a) through c) are carried out for a period of time sufficient to cause formation of a foregut spheroid, further

comprising the step of contacting said foregut spheroid with EGF, RA, and a BMP inhibitor.

[0006b] In one embodiment, there is provided method of generating a gastric organoid comprising fundus tissue comprising the steps of

a) generating a definitive endoderm (DE) from a pluripotent stem cell;

b) contacting said definitive endoderm with an agent that activates the FGF pathway as defined herein, an agent that activates the WNT pathway as defined in herein, an agent that inhibits the BMP pathway as defined herein, and retinoic acid to generate a posterior foregut; and

c) contacting said posterior foregut with EGF, retinoic acid, an agent that activates the WNT pathway as defined herein, and an agent that inhibits the BMP pathway as defined herein, to generate a fundus tissue.

[0006c] In one embodiment, there is provided a method of generating a gastric organoid comprising antrum tissue comprising the steps of:

a) generating a definitive endoderm (DE) from a pluripotent stem cell;

b) contacting said definitive endoderm with an agent that activates the FGF pathway as defined herein, an agent that activates the WNT pathway as defined herein, an agent that inhibits the BMP pathway as defined herein, and retinoic acid to generate a posterior foregut; and

c) contacting said posterior foregut with EGF, retinoic acid, and an agent that inhibits the BMP pathway as defined herein, to generate an antrum tissue.

Brief Description of the Drawings

[0007] Those of skill in the art will understand that the drawings, described below, are for illustrative purposes only. The drawings are not intended to limit the scope of the present teachings in any way.

[0008] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application

[0009] publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[0010] **FIG 1** depicts expression of Sox2/Cdx2/B-catenin in gastric spheroids and the effect of RA.

[0011] **FIG 2A-2E** depicts schematic representation of the *in vitro* culture system used to direct the differentiation of hPSCs into three-dimensional gastric organoids (FIG 2A), the defining markers of developing posterior foregut organs by wholemount immunofluorescent staining of mouse E10.5 embryos with Sox2, Pdx1 and Cdx2 (FIG 2B), PDX1 expression in the presence and absence of RA (FIG 2C), stereomicrographs showing morphological changes during growth of posterior foregut spheroids into hGOs (FIG 2D), and a comparison of developing mouse antrum at E14.5 and E18.5 and comparable stages of hGO development (FIG 2E).

[0012] **FIG 3A-3D** depictsMuc5AC, TFF2, GSII UEAI, and CHGA expression in P12 Antrum, E18.5 Antrum and d34 Organoid (FIG

3A), a schematic representation of the different roles for EGF in the growth, morphogenesis, and cell type specification during development of hGOs (FIG 3B), expression of gastrin, ghrelin, 5-HT, and ChrA in gastric organoids with and without DOX (FIG 3C), and relative expression of NEUROG3 at multiple concentrations of EGF (FIG 3D).

[0012] **FIG 4A-4D** depicts SOX9 Ki67 expression in d34 Organoid, E18.5 Antrum, and P12 Antrum (FIG 4A), H.Pylori infection of organoids visualized using brightfield microscopy and immunofluorescent staining (FIG 4B), immunoprecipitation for the oncogene c-Met (FIG 4C), and cell proliferation in the hGO epithelium, measured by EdU incorporation (FIG 4D).

[0013] **FIG 5A-5D** depicts Sox2 and Cdx2 expression in the presence of GSK3β inhibitor CHIR99021 and recombinant WNT3A in the presence and absence of noggin (FIG 5A), CHIR induced gut tube morphogenesis and spheroid production visualized using bright field microscopy (FIG 5B), immunofluorescent staining of monolayer cultures to assess CDX2 induction in CHIR/FGF-treated endoderm and SOX2 induction in noggin- and CHIR/FGF/noggin-treated endoderm (FIG 5C), qPCR analysis of BMP target genes MSX1/2 (FIG 5D), and SOX2 and CDX2 expression in the presence and absence of BMP2

[0014] **FIG 6A-6G** depicts a table comparing spheroid formation and characteristics between two hESC lines (H1 and H9) and one iPSC line (72.3) (FIG 6A), immunofluorescent staining of day 34 hGOs derived from H1 and iPSC 72.3 cell lines (FIG 6B), organ epithelial cell type quantification in day 34 hGOs (FIG 6C), characterization of the induced pluripotent stem cell line iPSC 72.3 (FIG 6D-G).

[0015] **FIG 7A-7D** depicts a schematic illustrating of foregut patterning experiments (FIG 7A), Brightfield images that show that RA increases the number of spheroids that are produced from foregut monolayer cultures (FIG 7B), a lower power image of FIG1d showing immunofluorescent image of a 14 somite stage embryo with Hnf1β protein localized to the posterior portion of the foregut (FIG 7C), qPCR analysis of gene expression in foregut spheroids treated with RA (FIG 7D).

[0016] **FIG 8** depicts brightfield images and immunostaining at late stages of hGO differentiation.

[0017] **FIG 9** depicts transcription factor expression during development of the mouse antrum and human gastric organoids during four embryonic stages (E12.5, E14.5, E16.5 and E18.5) and one postnatal stage (P12) of *in vivo* antrum development.

[0018] **FIG 10** depicts pHH3/E-Cad/DAPI expression and aPCC/E-CAD/DAPI expression in E12.5 Antrum and Day 13 hGO.

[0019] **FIG 11A-11C** depicts expression of the antral mesenchyme transcription factor BAPX1 (FIG 11A) and staining for mesenchymal cell type markers (FIG 11C).

[0020] **FIG 12** depicts gastric antrum endocrine cell development *in vivo*.

[0021] **FIG 13A-13B** depicts staining for the pan-endocrine marker CHGA (FIG 13A) and expression of endocrine markers CHGA, GASTRIN, GHRELIN, and SOMATOSTATIN (FIG 13B).

[0022] **FIG 14** shows a summary of methods for the directed differentiation of gastric organoids. Each step in the differentiation process is indicated, along with representative stereomicrographs.

[0023] **FIG 15** depicts a schematic of the mouse stomach and measurement of known regional markers in the forestomach, fundus, antrum, and duodenum.

[0024] **FIG 16** depicts measurement of new regional markers in the forestomach, fundus, antrum, and duodenum.

[0025] **FIG 17** depicts the fundus specification protocol and measurement of GAPDH, Gata4, Axin2, Sox2, Pdx1, and Cdx2 in control, Wnt100, Wnt500 and CHIR treated cells. The y axis represents relative gene expression.

[0026] **FIG 18** depicts measurements of Axin2, IRX2, IRX3, Pitx1, and IRX4 in The fundus protocol. The y axis represents relative gene expression.

[0027] **FIG 19** is a schematic depicting the formation of intestine tissue, fundus tissue, and antrum tissue from definitive endoderm.

Detailed Description

[0028] Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

[0029] As used herein, the term "totipotent stem cells" (also known as omnipotent stem cells) are stem cells that can differentiate into embryonic and extra-embryonic cell types. Such cells can construct a complete, viable, organism. These cells are produced from the fusion of an egg and sperm cell. Cells produced by the first few divisions of the fertilized egg are also totipotent.

[0030] As used herein, the term "pluripotent stem cells (PSCs)" encompasses any cells that can differentiate into nearly all cell types of the body, i.e., cells derived from any of the three germ layers (germinal epithelium), including endoderm (interior stomach

lining, gastrointestinal tract, the lungs), mesoderm (muscle, bone, blood, urogenital), and ectoderm (epidermal tissues and nervous system). PSCs can be the descendants of inner cell mass cells of the primplantation blastocyst or obtained through induction of a non-pluripotent cell, such as an adult somatic cell, by forcing the expression of certain genes. Pluripotent stem cells can be derived from any suitable source, as will be readily understood by one of skill in the art. Examples of sources of pluripotent stem cells include mammalian sources, including human, rodent, porcine, bovine, but are not so limited.

[0031] As used herein, the term "induced pluripotent stem cells (iPSCs)," also commonly abbreviated as iPS cells, refers to a type of pluripotent stem cells artificially derived from a normally non-pluripotent cell, such as an adult somatic cell, by inducing a "forced" expression of certain genes.

[0032] As used herein, the term "embryonic stem cells (ESCs)," also commonly abbreviated as ES cells, refers to cells that are pluripotent and derived from the inner cell mass of the blastocyst, an early-stage embryo. For purpose of the present invention, the term "ESCs" is used broadly sometimes to encompass the embryonic germ cells as well.

[0033] As used herein, the term "precursor cell" encompasses any cells that can be used in methods described herein, through which one or more precursor cells acquire the ability to renew itself or differentiate into one or more specialized cell types. In some embodiments, a precursor cell is pluripotent or has the capacity to becoming pluripotent. In some embodiments, the precursor cells are subjected to the treatment of external factors (e.g., growth factors) to acquire pluripotency. In some embodiments, a precursor cell can be a totipotent (or omnipotent) stem cell; a pluripotent

stem cell (induced or non-induced); a multipotent stem cell; an oligopotent stem cells and a unipotent stem cell. In some embodiments, a precursor cell can be from an embryo, an infant, a child, or an adult. In some embodiments, a precursor cell can be a somatic cell subject to treatment such that pluripotency is conferred via genetic manipulation or protein/peptide treatment.

[0034] In developmental biology, cellular differentiation is the process by which a less specialized cell becomes a more specialized cell type. As used herein, the term "directed differentiation" describes a process through which a less specialized cell becomes a particular specialized target cell type. The particularity of the specialized target cell type can be determined by any applicable methods that can be used to define or alter the destiny of the initial cell. Exemplary methods include but are not limited to genetic manipulation, chemical treatment, protein treatment, and nucleic acid treatment.

[0035] As used herein, the term "cellular constituents" are individual genes, proteins, mRNA expressing genes, and/or any other variable cellular component or protein activities such as the degree of protein modification (e.g., phosphorylation), for example, that is typically measured in biological experiments (e.g., by microarray or immunohistochemistry) by those skilled in the art. Significant discoveries relating to the complex networks of biochemical processes underlying living systems, common human diseases, and gene discovery and structure determination can now be attributed to the application of cellular constituent abundance data as part of the research process. Cellular constituent abundance data can help to identify biomarkers, discriminate disease subtypes and identify mechanisms of toxicity.

[0036] Stem cells are found in all multi cellular organisms. They are characterized by the ability to renew themselves through mitotic cell division and differentiate into a diverse range of specialized cell types. The two broad types of mammalian stem cells are 1) embryonic stem cells that are isolated from the inner cell mass of blastocysts, and 2) adult stem cells that are found in adult tissues. In a developing embryo, stem cells can differentiate into all of the specialized embryonic tissues. In adult organisms, stem cells and progenitor cells act as a repair system for the body, replenishing specialized cells, but also maintaining the normal turnover of regenerative organs, such as blood, skin, or gastric tissues.

[0037] Stem cells can now be grown and transformed into specialized cells with characteristics consistent with cells of various tissues such as muscles or nerves through cell culture. Highly plastic adult stem cells from a variety of sources, including umbilical cord blood and bone marrow, are routinely used in medical therapies. Embryonic cell lines and autologous embryonic stem cells generated through therapeutic cloning have also been proposed as promising candidates for future therapies.

[0038] The classical definition of a stem cell is typically indicative of two properties: self-renewal, the ability to go through numerous cycles of cell division while maintaining the undifferentiated state, and potency, the capacity to differentiate into specialized cell types. In some embodiments, stem cells are either totipotent or pluripotent, *i.e.* they are able to give rise to any mature cell type, although multipotent or unipotent progenitor cells may sometimes referred to as stem cells.

[0039] Potency specifies the differentiation potential (the potential to differentiate into different cell types) of the stem cell. Totipotent stem cells (also known as omnipotent stem cells) can differentiate

into embryonic and extraembryonic cell types. These cells can construct a complete, viable, organism. The cells are produced from the fusion of an egg and sperm cell. Cells produced by the first few divisions of the fertilized egg are also totipotent. Pluripotent stem cells (PSCs) are the descendants of totipotent cells and can differentiate into nearly all cells, *i.e.,* cells derived from any of the three germ layers, including endoderm (interior stomach lining, gastrointestinal tract, the lungs), mesoderm (muscle, bone, blood, urogenital), and ectoderm (epidermal tissues and nervous system). Multipotent stem cells can differentiate into a number of cells, but only those of a closely related family of cells. Oligopotent stem cells can differentiate into only a few cells, such as lymphoid or myeloid stem cells. Unipotent cells can produce only one cell type, their own, but have the property of self-renewal which distinguishes them from non-stem cells (*e.g.,* muscle stem cells).

[0040] Embryonic and induced pluripotent stem cells have had an unprecedented impact on the ability to study human diseases and to generate replacement tissues that are therapeutically effective in animal models.

[0041] In developmental biology, cellular differentiation is the process by which a less specialized cell becomes a more specialized cell type. Most successful efforts to direct the differentiation of human PSCs into therapeutic cell types have been based on studies of embryonic organ development. Examples include the generation of liver hepatocytes and pancreatic endocrine cells, which have shown functional potential in animal models of liver disease and diabetes. Similarly, differentiation of PSCs into intestine may provide therapeutic benefit for diseases such as necrotizing enterocolitis, inflammatory bowel diseases and short gut syndromes.

[0042] As discussed above, a pluripotent stem cell has the potential to differentiate into any of the three germ layers: endoderm (interior stomach lining, gastrointestinal tract, the lungs), mesoderm (muscle, bone, blood, urogenital), and ectoderm (epidermal tissues and nervous system). As such, pluripotent stem cells can give rise to any fetal or adult cell type. However, the fate of the particular pluripotent stem cells is controlled by numerous cellular signaling pathway and numerous factors. Further, the pluripotent stem cells alone cannot develop into a fetal or adult animal because they lack the potential to contribute to extraembryonic tissue, such as the placenta.

[0043] To date, gastric tissues have not been generated from human pluripotent stem cells (hPSCs). Successful efforts to differentiate PSCs into lung, liver, pancreas and intestinal cells have depended on a sound molecular understanding of the embryonic development of these organs.[6-10] Unfortunately, a problem in the art is the many gaps in understanding of stomach development subsequent to endoderm formation. Therefore, to direct differentiation of hPSCs into gastric tissue, the signaling pathways that regulate several critical stages of early stomach development including foregut specification and patterning, gastric specification, and lastly, gastric epithelial growth and differentiation were identified by Applicant. In addition, to generate more functional, complex three-dimensional tissue, Applicant aimed to induce several morphogenetic processes that occur during stomach development including morphogenesis of the foregut tube and the formation of gastric epithelial structures including glands and pits.

[0044] As described herein, methods and systems are established using a temporal series of growth factor manipulations to mimic embryonic gastric tissue development in culture. In particular, methods and

systems are established to direct *in vitro* the differentiation of PSCs, both human embryonic stem cells (hESC) and induced pluripotent stem cells (iPSC), into gastric tissue. These factors directed human intestinal development in vitro in stages that approximate fetal gut development: activin-induced definitive endoderm (DE) formation; FGF/Wnt/BMP induced posterior foregut pattering, and finally a pro-gastric culture system obtained by modulation of retinoic acid and EFG signaling that promoted gastric tissue growth, morphogenesis and cytodifferentiation into functional gastric cell types and morphology including gastric glands and pits, proliferative zones, surface and antral mucous cells, and endocrine cells expressing gastrin, ghrelin, and somatostatin.

[0045] Applicant has identified novel embryonic signaling pathways that allow for efficient step-wise differentiation of human PSCs into gastric cells, gastric tissues, and/or three-dimensional gastric tissue (hGOs) with complex architecture and cellular composition. Applicant has further found that the developing hGOs undergo molecular and morphological stages of differentiation that are nearly identical to the developing antrum of the mouse, and that the resulting gastric organoids may contain an array of mucous, endocrine, and progenitor cells that constitute the normal antral epithelium and a three-dimensional organization comparable to the fetal/postnatal stomach.

[0046] The disclosed human gastric cells, gastric tissue and/or gastric organoids (hGOs) may be used as an *in vitro* system to identify new mechanisms of human stomach development, physiology, and may be used as a model of the pathophysiological response of the gastric epithelium to *H. pylori*. The disclosed gastric cells, gastric tissue and/or gastric hGOs and methods present new opportunities

for drug discovery and modeling of early stages of gastric cancer. Moreover, disclosed herein is the first three-dimensional production of a human embryonic foregut, which is a promising starting point for the generation of other foregut organ tissues including lungs and pancreas.

[0047] In one aspect, a method of inducing formation of a gastric cell, gastric tissue, and or gastric hGO from a precursor cell is disclosed. The method may comprise the step of a) activating one or more signaling pathways within a precursor cell, wherein the one or more signaling pathways are selected from the WNT signaling pathway, the WNT/FGF signaling pathway, and the FGF signaling pathway to obtain a gastric cell, gastric tissue and/or gastric hGO descended from the precursor cell. The method may further comprise a step b) of inhibiting one or more signaling pathways within a precursor cell. The one or more signaling pathways that are inhibited may comprise a BMP signaling pathway.

[0048] The method may further comprise the step of contacting the precursor cell with retinoic acid. The contacting of a precursor cell with retinoic acid may occur after the activating and inhibiting steps above.

[0049] The method may further comprise the step of contacting a gastric organoid to EGF at a concentration and/or length of time sufficient to increase the diameter of the gastric organoid to greater than about 1 mm in diameter, or greater than about 2 mm in diameter, or greater than about 3 mm in diameter, or greater than about 4 mm in diameter.

[0050] In one aspect, the one or more signaling pathways may be selected from a Wnt signaling pathway, Wnt/beta-catenin signaling, Wnt/APC signaling, and Wnt/PCP pathway signaling.

[0051] In one aspect, the step of activating a Wnt signaling pathway may comprise contacting a precursor cell with one or more molecules selected from the group consisting of Wnt1, Wnt2, Wnt2b, Wnt3, Wnt3a, Wnt4, Wnt5a, Wnt5b, Wnt6, Wnt7a, Wnt7b, Wnt8a, Wnt8b, Wnt9a, Wnt9b, Wnt10a, Wnt10b, Wnt11, and Wnt16.

[0052] In one aspect, the step of activating the FGF signaling pathway may comprise contacting a precursor cell with one or more molecules selected from the group consisting of FGF1, FGF2, FGF3, FGF4, FGF5, FGF6, FGF7 FGF8, FGF9, FGF10, FGF11, FGF12, FGF13, FGF14, FGF16, FGF17, FGF18, FGF19, FGF20, FGF21, FGF22, and FGF23.

[0053] In one aspect, the step of inhibiting a BMP signaling pathway may comprise contacting the precursor cell with a BMP inhibitor. In one aspect, the BMP inhibitor may be selected from Dorsomorphin, LDN189, DMH-1, Noggin and combinations thereof. In one aspect, the BMP inhibitor may be Noggin.

[0054] In one aspect, the activating step may comprise contacting a precursor cell with Wnt3a, FGF4, and a BMP inhibitor over a specified period which is referred to as an incubation period. The contacting steps may occur simultaneously, or, in other aspects, the contacting steps may occur subsequently.

[0055] In one aspect, a precursor cell, which may comprise definitive endoderm, may be contacted by a signaling agent that may comprise 1) Wnt3a or a GSK-inhibitor (for example, CHIRON) in combination with 2) FGF4, during a first incubation period. The first incubation period may further comprise a BMP inhibitor. Following the first incubation period, the precursor cells may be subjected to a second incubation period wherein the precursor cells are contacted with retinoic acid (RA). In one aspect, the first

incubation period and the second incubation period overlap. In some embodiments, the first incubation period and the second incubation period do not overlap.

[0056] In one aspect, the first and/or second incubation period, and/or the totality of the first and second incubation period may be between 24 and 120 hours, or from about 36 to about 108 hours, or from about 48 to about 96 hours, or from about 60 to about 84 hours. In one aspect, the first incubation period may be at least about 24 hours.

[0057] In one aspect, the second incubation period (wherein the precursor cells may be contacted with RA) begins about 72 hours after the first incubation period. In a further aspect, the second incubation period begins after cultures have formed foregut spheroids from the precursor cells. The foregut spheroids may then be transferred to a 3-dimentional matrix under growth conditions suitable for formation of a gastric organoid, for example, by application of the foregut spheroids to Matrigel™ (Corning, BD Bioscience). Following transfer to Matrigel, the foregut spheroids are contacted with RA during a third incubation period in which continued 3D growth may occur. The spheroids may then be contacted with EGF during a fourth incubation period, which may overlap with the third incubation period. The third incubation period may be about 24 hours.

[0058] In one aspect, the precursor cell may be contacted with Wnt3a at a concentration between 50-1500 ng/ml, or from about 100 to about 1200 ng/ml, or from about 200 to about 1000 ng/ml, or from about 300 to about 900 ng/ml, or from about 400 to about 800 ng/ml, or from about 500 to about 700 ng/ml.

[0059] In one aspect, the precursor cell may be selected from an embryonic stem cell, an embryonic germ cell, an induced pluripotent stem cell, a mesoderm cell, a definitive endoderm cell, a posterior endoderm cell, and a hindgut cell.

[0060] In one aspect, the precursor cell may be a definitive endoderm cell derived from a pluripotent stem cell.

[0061] In one aspect, the precursor cell may be a pluripotent stem cell such as an embryonic stem cell, an embryonic stem cell, or an induced pluripotent stem cell.

[0062] In one aspect, the definitive endoderm cell may be derived by contacting the pluripotent stem cell with one or more molecules selected from of Activin, the BMP subgroups of the TGF-beta superfamily of growth factors; Nodal, Activin A, Activin B, BMP4, Wnt3a, and a combination thereof.

[0063] In one aspect, a gastric tissue may be produced *in vitro* from one or more precursor cells.

[0064] In one aspect, the one or more precursor cells may be selected from an embryonic stem cell, a mesoderm cell, a definitive endoderm cell, a posterior endoderm cell, an anterior endoderm cell, a foregut cell, and a hindgut cell.

[0065] In one aspect, the pluripotent stem cell may be a mammalian pluripotent stem cell, including but not limited to human pluripotent stem cell, or a mouse pluripotent stem cell.

[0066] In one aspect, the human pluripotent stem cell may be selected from a human embryonic stem cell, a human embryonic germ cell, and an induced human pluripotent stem cell.

[0067] In one aspect, a kit comprising a gastric cell, tissue, or organoid produced *in vitro* from one or more precursor cells is provided.

[0068] In one aspect, a method for identifying the absorption effect of gastric cells or tissues is provided. The method may comprise the steps of contacting a gastric cell, tissue, or organoid derived from a precursor cell with a compound; and detecting a level of absorption of a compound by said gastric cells or tissues.

[0069] In one aspect, a method for identifying the toxicity of a compound on a gastric cell or tissue is provided. The method may comprise the steps of contacting a gastric cell, tissue, or organoid derived from a precursor cell with a compound; and detecting a level of absorption of a compound by said gastric cells or tissues.

[0070] In one aspect, compositions comprising three-dimensional human gastric organoids (hGOs) generated *de novo*, and methods of making same through directed differentiation of human pluripotent stem cells (hPSCs) are disclosed. Such hGOs may be used these to model stomach development as well as the early events that occur during *H. pylori* infection.

[0071] In one aspect, methods for generating an hGO *in vitro* through the directed differentiation of human pluripotent stem cells (hPSCs) are disclosed. This human gastric tissue may be used to model human stomach development and disease. Methods for inducing definitive endoderm (DE) to form 3-dimensional gut tube structures are also disclosed. In one aspect, this may be carried out by activating FGF and WNT signaling, while a foregut fate may be promoted by simultaneously inhibiting BMP signaling. Foregut spheroids may then be directed into a posterior foregut and gastric fate by manipulation of retinoic acid and EGF signaling, resulting in hGOs.

[0072] Developing hGOs may undergo molecular and morphogenetic changes nearly identical to the developing mouse antrum, forming

gastric glands and pits, proliferative zones, surface and antral mucous cells, and endocrine cells expressing Gastrin, Ghrelin and Somatostatin. Using hGOs to model human stomach development it has been determined that EGF signaling represses endocrine cell development upstream of the transcription factor NEUROGENIN 3. Applicant has further found that hGOs faithfully recapitulate early stages of gastric disease initiated by *H. pylori*, including rapid activation of c-Met signaling and epithelial proliferation. Together, these studies describe a novel and robust *in vitro* system for elucidating the mechanisms underlying human stomach development and disease.

[0073] Pluripotent Stem Cells Derived from Embryonic Cells

[0074] In one aspect, the methods may include the step of obtaining stem cells that are pluripotent or can be induced to become pluripotent. In some embodiments, pluripotent stem cells are derived from embryonic stem cells, which are in turn derived from totipotent cells of the early mammalian embryo and are capable of unlimited, undifferentiated proliferation *in vitro*. Embryonic stem cells are pluripotent stem cells derived from the inner cell mass of the blastocyst, an early-stage embryo. Methods for deriving embryonic stem cells from blastocytes are well known in the art. For example, while certain cell types are exemplified herein, it would be understood by one of skill in the art that the methods and systems described herein are applicable to any stem cells.

[0075] Additional stem cells that can be used in embodiments in accordance with the present invention include but are not limited to those provided by or described in the database hosted by the National Stem Cell Bank (NSCB), Human Embryonic Stem Cell Research Center at the University of California, San Francisco (UCSF); WISC cell Bank at the Wi Cell Research Institute; the

University of Wisconsin Stem Cell and Regenerative Medicine Center (UW-SCRMC); Novocell, Inc. (San Diego, Calif.); Cellartis AB (Goteborg, Sweden); ES Cell International Pte Ltd (Singapore); Technion at the Israel Institute of Technology (Haifa, Israel); and the Stem Cell Database hosted by Princeton University and the University of Pennsylvania. Exemplary embryonic stem cells that can be used in embodiments in accordance with the present invention include but are not limited to SA01 (SA001); SA02 (SA002); ES01 (HES-1); ES02 (HES-2); ES03 (HES-3); ES04 (HES-4); ES05 (HES-5); ES06 (HES-6); BG01 (BGN-01); BG02 (BGN-02); BG03 (BGN-03); TE03 (13); TE04 (14); TE06 (16); UC01 (HSF1); UC06 (HSF6); WA01 (H1); WA07 (H7); WA09 (H9); WA13 (H13); WA14 (H14).

[0076] In some embodiments, the stem cells may be further modified to incorporate additional properties. Exemplary modified cell lines include, but are not limited to, H1 OCT4-EGFP; H9 Cre-LoxP; H9 hNanog-pGZ; H9 hOct4-pGZ; H9 in GFPhES; and H9 Syn-GFP.

[0077] More details on embryonic stem cells can be found in, for example, Thomson et al., 1998, "Embryonic Stem Cell Lines Derived from Human Blastocysts," Science 282 (5391):1145-1147; Andrews et al., 2005, "Embryonic stem (ES) cells and embryonal carcinoma (EC) cells: opposite sides of the same coin," Biochem Soc Trans 33:1526-1530; Martin 1980, "Teratocarcinomas and mammalian embryogenesis,". Science 209 (4458):768-776; Evans and Kaufman, 1981, "Establishment in culture of pluripotent cells from mouse embryos," Nature 292(5819): 154-156; Klimanskaya et al., 2005, "Human embryonic stem cells derived without feeder cells," Lancet 365 (9471): 1636-1641; each of which is hereby incorporated herein in its entirety.

[0078] Alternative, pluripotent stem cells can be derived from embryonic germ cells (EGCs), which are the cells that give rise to the gametes of organisms that reproduce sexually. EGCs are derived from primordial germ cells found in the gonadal ridge of a late embryo, have many of the properties of embryonic stem cells. The primordial germ cells in an embryo develop into stem cells that in an adult generate the reproductive gametes (sperm or eggs). In mice and humans it is possible to grow embryonic germ cells in tissue culture under appropriate conditions. Both EGCs and ESCs are pluripotent. For purpose of the present invention, the term "ESCs" is used broadly sometimes to encompass EGCs.

[0079] Induced Pluripotent Stem Cells (iPSCs)

[0080] In some embodiments, iPSCs are derived by transfection of certain stem cell-associated genes into non-pluripotent cells, such as adult fibroblasts. Transfection is typically achieved through viral vectors, such as retroviruses. Transfected genes include the master transcriptional regulators Oct-3/4 (Pouf51) and Sox2, although it is suggested that other genes enhance the efficiency of induction. After 3-4 weeks, small numbers of transfected cells begin to become morphologically and biochemically similar to pluripotent stem cells, and are typically isolated through morphological selection, doubling time, or through a reporter gene and antibiotic selection. As used herein, iPSCs may include, but are not limited to, first generation iPSCs, second generation iPSCs in mice, and human induced pluripotent stem cells. In some embodiments, a retroviral system may used to transform human fibroblasts into pluripotent stem cells using four pivotal genes: Oct3/4, Sox2, Klf4, and c-Myc. In alternative embodiments, a lentiviral system is used to transform somatic cells with OCT4, SOX2, NANOG, and LIN28. Genes whose expression may be induced in iPSCs include

but are not limited to Oct-3/4 (e.g., Pou5f1); certain members of the Sox gene family (e.g., Sox1, Sox2, Sox3, and Sox15); certain members of the Klf family (e.g., Klf1, Klf2, Klf4, and Klf5), certain members of the Myc family (e.g., C-myc, L-myc, and N-myc), Nanog, and LIN28.

[0081] In some embodiments, non-viral based technologies may be employed to generate iPSCs. In some embodiments, an adenovirus can be used to transport the requisite four genes into the DNA of skin and liver cells of mice, resulting in cells identical to embryonic stem cells. Since the adenovirus does not combine any of its own genes with the targeted host, the danger of creating tumors is eliminated. In some embodiments, reprogramming can be accomplished via plasmid without any virus transfection system at all, although at very low efficiencies. In other embodiments, direct delivery of proteins is used to generate iPSCs, thus eliminating the need for viruses or genetic modification. In some embodiment, generation of mouse iPSCs is possible using a similar methodology: a repeated treatment of the cells with certain proteins channeled into the cells via poly-arginine anchors was sufficient to induce pluripotency. In some embodiments, the expression of pluripotency induction genes can also be increased by treating somatic cells with FGF2 under low oxygen conditions.

[0082] More details on embryonic stem cells can be found in, for example, Kaji et al., 2009, "Virus free induction of pluripotency and subsequent excision of reprogramming factors," Nature 458:771-775; Woltjen et al., 2009, "piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells," Nature 458:766-770; Okita et al., 2008, "Generation of Mouse Induced Pluripotent Stem Cells Without Viral Vectors," Science 322(5903):949-953; Stadtfeld et al., 2008, "Induced Pluripotent Stem Cells Generated

without Viral Integration," Science 322(5903):945-949; and Zhou et al., 2009, "Generation of Induced Pluripotent Stem Cells Using Recombinant Proteins," Cell Stem Cell 4(5):381-384; each of which is hereby incorporated herein in its entirety.

[0083] In some embodiments, exemplary iPS cell lines include but are not limited to iPS-DF19-9; iPS-DF19-9; iPS-DF4-3; iPS-DF6-9; iPS(Foreskin); iPS(IMR90); and iPS(IMR90).

[0084] It has been shown that iPSCs were capable of differentiation in a fashion similar to ESCs into fully differentiated tissues. For example, iPSCs were differentiated into neurons, expressing βIII-tubulin, tyrosine hydroxylase, AADC, DAT, ChAT, LMX1B, and MAP2. The presence of catecholamine-associated enzymes may indicate that iPSCs, like hESCs, may be differentiable into dopaminergic neurons. Stem cell-associated genes were shown to be down-regulated after differentiation. It has also been shown that iPSCs may be differentiated into cardiomyocytes that spontaneously began beating. Cardiomyocytes expressed TnTc, MEF2C, MYL2A, MYHCβ, and NKX2.5. Stem cell-associated genes were down-regulated after differentiation.

[0085] Gastric Organ and Development

[0086] Prior to Applicant's invention, no systems were available for converting precursor cells such as embryonic stem cells and/or iPSCs into gastric tissues.

[0087] In some embodiments, PSCs, such as ESCs and iPSCs, undergo directed differentiation in a step-wise manner first into definitive endoderm (DE) then into three dimensional gut tube structures (foregut spheroids) then into three dimensional gastric organoid (hGO) via formation of a posterior foregut/gastric tissue.

[0088] In some embodiments, PSCs, such as ESCs and iPSCs, undergo directed differentiation in a non step-wise manner where molecules (e.g., growth factors, ligands) for promoting DE formation and those for subsequent tissue formation are added at the same time.

[0089] Definitive Endoderm

[0090] The epithelium of the stomach is derived from a simple sheet of cells called the definitive endoderm (DE). The anterior DE forms the foregut and its associated organs including the lungs, esophagus, stomach, liver and pancreas and the posterior DE forms the midgut and hindgut, which forms the small and large intestines and parts of the genitourinary system. The DE gives rise to the epithelia of the gastrointestinal and respiratory tracts *in vivo*. Studies using mouse, chick and frog embryos suggest that establishing the anterior-posterior pattern in DE at the gastrula stage is a prerequisite for subsequent foregut and hindgut development. In some embodiments, PSCs, such as ESCs and iPSCs, undergo directed differentiation in a step-wise manner first into definitive endoderm (DE) then into anterior/foregut epithelium (e.g., foregut spheroids), and then into gastric tissue. The BMP, Wnt and FGF signaling pathways are believed to be critical for this process. Activation of WNT and FGF act to promote gut tube morphogenesis and inhibition of BMP signaling promotes a foregut fate. The simple cuboidal epithelium of the foregut first develops into a pseudostratified columnar epithelium, then into glands and pits containing gastric epithelium and a proliferative zone at the base of the villi, which corresponds with the presumptive progenitor domain.

[0091] A robust and efficient process is established to direct the differentiation of DE into gastric tissue *in vitro*. In some embodiments, directed differentiation is achieved by selectively

activating certain signaling pathways in the iPSCs and/or DE cells. In some embodiments, the signaling pathways are those active in gastric tissue development, including but not limited to the Wnt signaling pathway, Wnt/APC signaling pathway, FGF signaling pathway, TGF-beta signaling pathway, BMP signaling pathway; EGF signaling pathway, and Retinoic Acid signaling pathway.

[0092] More details on the functions of signaling pathways relating to DE development and/or intestinal development in general can be found in, for example, Zorn and Wells, 2009, "Vertebrate endoderm development and organ formation," Annu Rev Cell Dev Biol 25:221-251; Dessimoz et al., 2006, "FGF signaling is necessary for establishing gut tube domains along the anterior-posterior axis in vivo," Mech Dev 123:42-55; McLin et al., 2007, "Repression of Wnt/{beta}-catenin signaling in the anterior endoderm is essential for liver and pancreas development. Development," 134:2207-2217; Wells and Melton, 2000, Development 127:1563-1572; de Santa Barbara et al., 2003, "Development and differentiation of the intestinal epithelium," Cell Mol Life Sci 60(7): 1322-1332; Sancho et al., 2004, "Signaling Pathways in Intestinal Development and Cancer," Annual Review of Cell and Developmental Biology 20:695-723; Logan and Nusse, 2004, "The Wnt Signaling Pathway in Development and Disease," Annual Review of Cell and Developmental Biology 20:781-810; Taipalel and Beachyl, 2001, "The Hedgehog and Wnt signalling pathways in cancer," Nature 411:349-354; Gregorieff and Clevers, 2005, "Wnt signaling in the intestinal epithelium: from endoderm to cancer," Genes & Dev. 19: 877-890; each of which is hereby incorporated herein in its entirety.

[0093] Any methods for producing definitive endoderm from pluripotent cells (e.g., iPSCs or ESCs) are applicable to the methods described

herein. In some embodiments, pluripotent cells are derived from a morula. In some embodiments, pluripotent stem cells are stem cells. Stem cells used in these methods can include, but are not limited to, embryonic stem cells. Embryonic stem cells can be derived from the embryonic inner cell mass or from the embryonic gonadal ridges. Embryonic stem cells or germ cells can originate from a variety of animal species including, but not limited to, various mammalian species including humans. In some embodiments, human embryonic stem cells are used to produce definitive endoderm. In some embodiments, human embryonic germ cells are used to produce definitive endoderm. In some embodiments, iPSCs are used to produce definitive endoderm.

[0094] In some embodiments, one or more growth factors are used in the differentiation process from pluripotent stem cells to DE cells. The one or more growth factors used in the differentiation process can include growth factors from the TGF-beta superfamily. In such embodiments, the one or more growth factors comprise the Nodal/Activin and/or the BMP subgroups of the TGF-beta superfamily of growth factors. In some embodiments, the one or more growth factors are selected from the group consisting of Nodal, Activin A, Activin B, BMP4, Wnt3a or combinations of any of these growth factors.

[0095] In some embodiments, the embryonic stem cells or induced pluripotent cells and iPSCs are treated with the one or more growth factors for 6 or more hours; 12 or more hours; 18 or more hours; 24 or more hours; 36 or more hours; 48 or more hours; 60 or more hours; 72 or more hours; 84 or more hours; 96 or more hours; 120 or more hours; 150 or more hours; 180 or more hours; or 240 or more hours.

[0096] In some embodiments, the embryonic stem cells or germ cells and iPSCs are treated with the one or more growth factors at a concentration of 10 ng/ml or higher; 20 ng/ml or higher; 50 ng/ml or higher; 75 ng/ml or higher; 100 ng/ml or higher; 120 ng/ml or higher; 150 ng/ml or higher; 200 ng/ml or higher; 500 ng/ml or higher; 1,000 ng/ml or higher; 1,200 ng/ml or higher; 1,500 ng/ml or higher; 2,000 ng/ml or higher; 5,000 ng/ml or higher; 7,000 ng/ml or higher; 10,000 ng/ml or higher; or 15,000 ng/ml or higher. In some embodiments, concentration of the growth factor is maintained at a constant level throughout the treatment. In other embodiments, concentration of the growth factor is varied during the course of the treatment. In some embodiments, the growth factor is suspended in media that include fetal bovine serine (FBS) with varying HyClone concentrations. One of skill in the art would understand that the regimen described herein is applicable to any known growth factors, alone or in combination. When two or more growth factors are used, the concentration of each growth factor may be varied independently.

[0097] In some embodiments, populations of cells enriched in definitive endoderm cells are used. In some embodiments, the definitive endoderm cells are isolated or substantially purified. In some embodiments, the isolated or substantially purified definitive endoderm cells express the SOX17, FOXA2, and/or the CXRC4 marker to a greater extent than the OCT4, AFP, TM, SPARC and/or SOX7 markers.

[0098] Methods for enriching a cell population with definitive endoderm are also contemplated. In some embodiments, definitive endoderm cells can be isolated or substantially purified from a mixed cell population by contacting the cells with a reagent that binds to a molecule that is present on the surface of definitive endoderm cells

but which is not present on the surface of other cells in the mixed cell population, and then isolating the cells bound to the reagent. In certain embodiments, the cellular constituent that is present on the surface of definitive endoderm cells is CXCR4.

[0099] Still other embodiments of the present invention relate to CXCR4 antibodies, SDF-1 ligands or other ligands for CXCR4 can be used to obtain definitive endoderm cells in an enriched, isolated or substantially purified form. For example, a CXCR4 antibody, an SDF-1 ligand or another ligand for CXCR4 can be used as a reagent in a method, such as affinity-based separation or magnetic-based separation, to enrich, isolate or substantially purify preparations of definitive endoderm cells that bind to the reagent.

[00100] In some embodiments, definitive endoderm cells and hESCs are treated with one or more growth factors. Such growth factors can include growth factors from the TGF-beta superfamily. In such embodiments, the one or more growth factors comprise the Nodal/Activin and/or the BMP subgroups of the TGF-beta superfamily of growth factors. In some embodiments, the one or more growth factors are selected from the group consisting of Nodal, Activin A, Activin B, BMP4, Wnt3a or combinations of any of these growth factors.

[00101] Additional methods for obtaining or creating DE cells that can be used in the present invention include but are not limited to those described in U.S. Pat. No. 7,510,876 to D'Amour et al.; U.S. Pat. No. 7,326,572 to Fisk et al.; Kubol et al., 2004, "Development of definitive endoderm from embryonic stem cells in culture," Development 131:1651-1662; D'Amour et al., 2005, "Efficient differentiation of human embryonic stem cells to definitive endoderm," Nature Biotechnology 23:1534-1541; and Ang et al., 1993, "The formation and maintenance of the definitive endoderm

lineage in the mouse: involvement of HNF3/forkhead proteins," Development 119:1301-1315; each of which is hereby incorporated by reference herein in its entirety.

[00102] Directed Differentiation of Posteriorized DE

[00103] In some embodiments, activin-induced definitive endoderm (DE) can further undergo FGF/Wnt/Noggin induced anterior endoderm patterning, foregut specification and morphogenesis, and finally a pro-gastric culture system to promote gastric tissue growth, morphogenesis and cytodifferentiation into functional gastric cell types including surface mucous cells, mucous gland cells, endocrine, and progenitor cells. In some embodiments, human PSCs are efficiently directed to differentiate *in vitro* into gastric epithelium that includes mucous, endocrine, and progenitor cell types. It will be understood that molecules such as growth factors can be added to any stage of the development to promote a particular type of gastric tissue formation.

[00104] In some embodiments, anteriorized endoderm cells of the DE are further developed into one or more specialized cell types.

[00105] In some embodiments, soluble FGF and Wnt ligands and BMP antagonists are used to mimic early foregut specification in culture to convert, through directed differentiation, DE developed from iPSCs or ESCs into foregut epithelium that efficiently gives rise to all the major antrum gastric cell types. In human, directed differentiation of DE is achieved through selective activating certain signaling pathways that are important to gastric development.

[00106] Human stomach/gastric development *in vitro* occurs in stages that approximate fetal gut development; endoderm formation, anterior endoderm patterning, foregut morphogenesis, fetal gastric, antral

and fundic development, epithelial morphogenesis, formation of a presumptive progenitor domain, and differentiation into functional cell types of the stomach.

[00107] It will be understood by one of skill in the art that altering the expression of any Wnt signaling protein in combination with any FGF ligand can give rise to directed differentiation in accordance of the present invention. In some embodiments, the alteration is over-expression of Wnt3, in particular Wnt3a. In some embodiments, the alternation is over-expression of Wnt1 or other Wnt ligands..

[00108] It will be understood by one of skill in the art that altering the signaling activity of the Wnt signaling pathway in combination with altering the signaling activity of the FGF signaling pathway can give rise to directed differentiation in accordance of the present invention. In some embodiments, the alteration is through the use of small molecule modulators that activate the aforementioned pathways. For example, Small molecule modulators of the Wnt pathway included, but is not limited to Lithium Chloride; 2-amino-4,6-disubstituted pyrimidine (hetero) arylpyrimidines; IQ1; QS11; NSC668036; DCA beta-catenin; 2-amino-4-[3,4-(methylenedioxy)-benzyl-amino]-6-(3-methoxyphenyl) pyrimidine.

[00109] In alternative embodiments, cellular constituents associated with the Wnt and/or FGF signaling pathways, for example, natural inhibitors or antagonist of the pathways can be inhibited to result in activation of the Wnt and/or FGF signaling pathways.

[00110] In some embodiment, the cellular constituents are inhibited by other cellular constituents or extrinsic molecules. Exemplary natural inhibitors of Wnt signaling include but are not limited to Dkk1, SFRP proteins and FrzB. In some embodiments, the

extrinsic molecules may include, but are not limited to, small molecules such as WAY-316606; SB-216763; or BIO (6-bromoindirubin-3′-oxime).

[00111] More details are found, for example, in Liu et al., "A small-molecule agonist of the Wnt signaling pathway," Angew Chem Int Ed Engl. 44(13):1987-1990 (2005); Miyabayashi et al., "Wnt/beta-catenin/CBP signaling maintains long-term murine embryonic stem cell pluripotency," Proc Natl Acad Sci USA. 104(13):5668-5673 (2007); Zhang et al., "Small-molecule synergist of the Wnt/beta-catenin signaling pathway," Proc Natl Acad Sci U S A. 104(18):7444-7448 (2007); Neiiendam et al., "An NCAM-derived FGF-receptor agonist, the FGL-peptide, induces neurite outgrowth and neuronal survival in primary rat neurons," J. Neurochem. 91(4):920-935 (2004); Shan et al., "Identification of a specific inhibitor of the dishevelled PDZ domain," Biochemistry 44(47):15495-15503 (2005); Coghlan et al., "Selective small molecule inhibitors of glycogen synthase kinase-3 modulate glycogen metabolism and gene transcription," Chem. Biol. 7(10):793-803 (2000); Coghlan et al., "Selective small molecule inhibitors of glycogen synthase kinase-3 modulate glycogen metabolism and gene transcription," Chemistry & Biology 7(10):793-803; and Pai et al., "Deoxycholic acid activates beta-catenin signaling pathway and increases colon cell cancer growth and invasiveness," Mol Biol Cell. 15(5):2156-2163 (2004); each of which is hereby incorporated by reference in its entirety.

[00112] In some embodiments, siRNA and/or shRNA targeting cellular constituents associated with the Wnt and/or FGF signaling pathways are used to activate these pathways. It would be understood by one of skill in the art that the target cellular

constituents may include, but are not limited to, SFRP proteins; GSK3, Dkk1, and FrzB.

[00113] More details about RNAi based technologies can be found, for example, in Couzin, 2002, Science 298:2296-2297; McManus et al., 2002, Nat. Rev. Genet. 3, 737-747; Hannon, G. J., 2002, Nature 418, 244-251; Paddison et al., 2002, Cancer Cell 2, 17-23; Elbashir et al., 2001. EMBO J. 20:6877-6888; Tuschl et al., 1999, Genes Dev. 13:3191-3197; Hutvagner et al., Sciencexpress 297:2056-2060; each of which is hereby incorporated by reference in its entirety.

[00114] Fibroblast growth factors (FGFs) are a family of growth factors involved in angiogenesis, wound healing, and embryonic development. The FGFs are heparin-binding proteins and interactions with cell-surface associated heparan sulfate proteoglycans have been shown to be essential for FGF signal transduction. FGFs are key players in the processes of proliferation and differentiation of wide variety of cells and tissues. In humans, 22 members of the FGF family have been identified, all of which are structurally related signaling molecules. Members FGF1 through FGF10 all bind fibroblast growth factor receptors (FGFRs). FGF1 is also known as acidic, and FGF2 is also known as basic fibroblast growth factor. Members FGF11, FGF12, FGF13, and FGF14, also known as FGF homologous factors 1-4 (FHF1-FHF4), have been shown to have distinct functional differences compared to the FGFs. Although these factors possess remarkably similar sequence homology, they do not bind FGFRs and are involved in intracellular processes unrelated to the FGFs. This group is also known as "iFGF." Members FGF16 through FGF23 are newer and not as well characterized. FGF15 is the mouse ortholog of human FGF19 (hence there is no human

FGF15). Human FGF20 was identified based on its homology to Xenopus FGF-20 (XFGF-20). In contrast to the local activity of the other FGFs, FGF15/FGF19, FGF21 and FGF23 have more systemic effects.

[00115] In some embodiments, it will be understood by one of skill in the art that any of the FGFs can be used in conjunction with a protein from the Wnt signaling pathway. In some embodiments, soluble FGFs may include, but are not limited to, FGF4, FGF2, and FGF3.

[00116] In some embodiment, the cellular constituents of the FGF signaling pathway are inhibited by other cellular constituents or extrinsic molecules. Exemplary natural inhibitors of FGF signaling may include, but are not limited to, the Sprouty family of proteins and the Spred family of proteins. As discussed above, proteins, small molecules, nucleic acids can be used to activating the FGF signaling pathway.

[00117] It will be understood by one of skill in the art that the methods and compositions described herein in connection with the Wnt and FGF signaling pathways are provided by way of examples. Similar methods and compositions are applicable to other signaling pathways disclosed herein.

[00118] In some embodiments, DE culture may be treated with the one or more molecules of a signaling pathway described herein for 6 or more hours; 12 or more hours; 18 or more hours; 24 or more hours; 36 or more hours; 48 or more hours; 60 or more hours; 72 or more hours; 84 or more hours; 96 or more hours; 120 or more hours; 150 or more hours; 180 or more hours; 200 or more hours, 240 or more hours; 270 or more hours; 300 or more hours; 350 or more hours; 400 or more hours; 500 or more hours; 600 or more hours; 700 or

more hours; 800 or more hours; 900 or more hours; 1,000 or more hours; 1,200 or more hours; or 1,500 or more hours.

[00119] In some embodiments, DE culture is treated with the one or more molecules of a signaling pathway described herein at a concentration of 10 ng/ml or higher; 20 ng/ml or higher; 50 ng/ml or higher; 75 ng/ml or higher; 100 ng/ml or higher; 120 ng/ml or higher; 150 ng/ml or higher; 200 ng/ml or higher; 500 ng/ml or higher; 1,000 ng/ml or higher; 1,200 ng/ml or higher; 1,500 ng/ml or higher; 2,000 ng/ml or higher; 5,000 ng/ml or higher; 7,000 ng/ml or higher; 10,000 ng/ml or higher; or 15,000 ng/ml or higher. In some embodiments, concentration of signaling molecule is maintained at a constant throughout the treatment. In other embodiments, concentration of the molecules of a signaling pathway is varied during the course of the treatment. In some embodiments, a signaling molecule in accordance with the present invention is suspended in media comprising DMEM and fetal bovine serine (FBS). The FBS can be at a concentration of 2% and more; 5% and more; 10% or more; 15% or more; 20% or more; 30% or more; or 50% or more. One of skill in the art would understand that the regiment described herein is applicable to any known molecules of the signaling pathways described herein, alone or in combination, including but not limited to any molecules in the Wnt and FGF signaling pathways.

[00120] In embodiments where two or more signaling molecules are used to treat the DE culture, the signaling molecules can be added simultaneously or separately. When two or more molecules are used, the concentration of each may be varied independently.

[00121] Differentiation of PSCs into DE culture and subsequently into various intermediate mature gastric cell types can be determined by the presence of stage-specific cell markers. In some embodiments,

expression of representative cellular constituents is used to determine DE formation. The representative cellular constituents may include, but are not limited to, CMKOR1, CXCR4, GPR37, RTN4RL1, SLC5A9, SLC40A1, TRPA1, AGPAT3, APOA2, C20orf56, C21orf129, CALCR, CCL2, CER1, CMKOR1, CRIP1, CXCR4, CXorf1, DIO3, DIO30S, EB-1, EHHADH, ELOVL2, EPSTI1, FGF17, FLJ10970, FLJ21195, FLJ22471, FLJ23514, FOXA2, FOXQ1, GATA4, GPR37, GSC, LOC283537, MYL7, NPPB, NTN4, PRSS2, RTN4RL1, SEMA3E, SIAT8D, SLC5A9, SLC40A1, SOX17, SPOCK3, TMOD1, TRPA1, TTN, AW166727, AI821586, BF941609, AI916532, BC034407, N63706 and AW772192.

[00122] Additional cellular constituents suitable for detecting DE formation can be found in, for example, in U.S. patent application Ser. No. 11/165,305, filed Jun. 23, 2005; U.S. patent application Ser. No. 11/317,387, filed Dec. 22, 2005; U.S. patent Ser. No. 11/021,618, filed Dec. 23, 2004; U.S. patent application Ser. No. 11/021,618, 11/115,868 filed on Apr. 26, 2005; U.S. patent application Ser. No. 11/317,387, filed on Dec. 22, 2005; U.S. patent application Ser. No. 11/474,211, filed on Jun. 23, 2006; U.S. patent application Ser. No. 11/165,305, filed on Jun. 23, 2005; U.S. patent application Ser. No. 11/587,735 filed on Aug. 29, 2008; U.S. patent application Ser. No. 12/039,701, filed on Feb. 28, 2008; U.S. patent application Ser. No. 12/414,482, filed on Mar. 30, 2009; U.S. patent application Ser. No. 12/476,570, filed on Jun. 2, 2009; U.S. patent application Ser. No. 12/093,590 filed on Jul. 21, 2008; U.S. patent application Ser. No. 12/582,600 filed on Oct. 20, 2009; each of which is hereby incorporated by reference herein in its entirety.

[00123] In some embodiments, expression of SOX2 is used to reveal tendency of foregut formation after DE have been incubated with

FGF4 and Wnt3a plus Noggin for a period of time, for example, for 12 hours or longer; 18 hours or longer; 24 hours or longer; 36 hours or longer; 48 hours or longer; 60 hours or longer; or 90 hours or longer. In some embodiments, longer periods of incubation are needed to achieve a stable anterior endoderm phenotype as measured by prolonged expressed of CDX2. In such embodiments, the periods of incubation can be for 60 hours or longer; 72 hours or longer; 84 hours or longer; 96 hours or longer; 108 hours or longer; 120 hours or longer; 140 hours or longer; 160 hours or longer; 180 hours or longer; 200 hours or longer; 240 hours or longer; or 300 hours or longer.

[00124] Alternatively, in some embodiments, the absence of cellular constituents, such as hindgut markers such as CDX2 can be used to reveal directed foregut formation. In some embodiments, gastric transcription factors PDX1, KLF5, and SOX9 can be used to represent gastric development. In some embodiments, GATA4 and/or GATA6 protein expression can be used to represent gastric development. In these embodiments, the periods of incubation can be for 12 hours or longer; 18 hours or longer; 24 hours or longer; 36 hours or longer; 48 hours or longer; 60 hours or longer; or 90 hours or longer. Alternatively, the periods of incubation can be for 60 hours or longer; 72 hours or longer; 84 hours or longer; 96 hours or longer; 108 hours or longer; 120 hours or longer; 140 hours or longer; 160 hours or longer; 180 hours or longer; 200 hours or longer; 240 hours or longer; or 300 hours or longer.

[00125] In some embodiments, abundance data of cellular constituents, for example, protein and/or gene expression levels, are determined by immunohistochemistry using primary and/or secondary antibodies targeting molecules in the relevant signaling pathways. In other embodiments, abundance data of cellular constituents, for example,

protein and/or gene expression levels, are determined by microarray analyses.

[00126] Still alternatively, morphological changes can be used to represent the progress of directed differentiation. In some embodiments, foregut spheroids may be further subject to 3-dimensional culture conditions for further maturation. Additionally, gastric organoids can be observed in 6 days or longer; 7 days or longer; 9 days or longer; 10 days or longer; 12 days or longer; 15 days or longer; 20 days or longer; 25 days or longer; 28 days or longer; 32 days or longer; 36 days or longer; 40 days or longer; 45 days or longer; 50 days or longer; or 60 days or longer.

[00127] Directed Differentiation of Pluripotent Stem Cells

[00128] In some embodiments, pluripotent stem cells are converted into gastric cell types via a "one step" process. For example, one or more molecules that can differentiate pluripotent stem cells into DE culture (e.g., ActivinA) are combined with additional molecules that can promote directed differentiation of DE culture (e.g., Wnt3a/FGF4 activators and BMP inhibitors) to directly treat pluripotent stem cells.

[00129] Utilities and Kits Embodiments

[00130] In some embodiments, gastric tissue or related cell types described herein can be used to screen drugs for gastric uptake and/or mechanisms of transport and/or treatment of H.Pylori. For example, this can be done in a high throughput manner to screen for the most readily absorbed or effective drugs, and can augment Phase 1 clinical trials that are done to study drug gastric uptake and gastric toxicity. This may include pericellular and intracellular transport mechanisms of small molecules, peptides, metabolites, salts. The gastric tissues disclosed herein may further be used to

assess compatibility with any agent and/or device that is intended to come into contact with the gastric tissues to assess biocompatibility.

[00131] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to identify the molecular basis of normal human gastric development.

[00132] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to identify the molecular basis of congenital defects affecting human gastric development.

[00133] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to correct gastric congenital defects caused by genetic mutations. In particular, mutation affecting human gastric development can be corrected using iPSC technology and genetically normal gastric tissue or related cell types described herein. In some embodiments, gastric tissue or related cell types described herein can be used to generate replacement tissue. Examples of genetic diseases include but are not limited to Neurog3 mutations and Enteric anendocrinosis, PTF1a mutations and neonatal diabetes, PDX1 mutations that effect enteroendocrine cells of the stomach.

[00134] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to generate replacement gastric tissue for diseases or conditions such as peptic ulcer disease, Ménétrier's disease, or for gastric cancer patients.

[00135] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to study microbiotic interactions with the human host epithelium and host immunity.

[00136] In some embodiments, gastric tissue or related cell types described herein, in particular the enteroendocrine cells can be used to study hormonal regulation of feeding behavior, metabolism, mediated by gastric endocrine.

[00137] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein, in particular the enteroendocrine cells that produce the hormone gastrin or ghrelin can be used to study and improve, for example, metabolic control in patients with obesity, metabolic syndrome, or Type 2 diabetes.

[00138] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to replace any damaged or removed gastric tissue in a subject in need thereof.

[00139] In some embodiments, a gastric cell, gastric tissue and/or gastric hGO described herein can be used to screen for toxicity and efficacy of any drug that acts on the gastric tissues.

[00140] In some embodiments where a gastric cell, gastric tissue and/or gastric hGO described herein are used to determine the absorption level of a compound, the compound will be contacted with the gastric cell, gastric tissue and/or gastric hGO with a compound; and a level of absorption of the compound by the gastric cell, gastric tissue and/or gastric hGO can be quantified. In some embodiments, the compound may be labeled with a radio-isotope, a fluorescent label and or a primary or secondary visible marker.

[00141] In some embodiments, a diagnostic kit or package is developed to include the gastric cell, gastric tissue and/or gastric hGO described herein and based on one or more of the aforementioned utilities.

[00142] Having described the invention in detail, it will be apparent that modifications, variations, and equivalent embodiments are possible

without departing the scope of the invention defined in the appended claims. Furthermore, it should be appreciated that all examples in the present disclosure are provided as non-limiting examples.

EXAMPLES

[00143] The following non-limiting examples are provided to further illustrate embodiments of the invention disclosed herein. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches that have been found to function well in the practice of the invention, and thus can be considered to constitute examples of modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

[00144] **Pluripotent stem cell culture**

[00145] Human embryonic stem cell lines WA01 (H1) and WA09 (H9) were obtained from WiCell. ESC and iPSC lines were maintained as colonies in feeder-free conditions on HESC-qualified Matrigel (BD Biosciences) in mTesR1 media (Stem Cell Technologies). Cells were routinely passaged every four days with dispase (Invitrogen).

[00146] **DE induction.**

[00147] (Summarized in FIG. 14.) Human ES and iPS cells were plated as single cells in mTesR1 media plus ROCK inhibitor Y27632 (10 μM; Stemgent) in a Matrigel (BD Biosciences)-coated 24-well plate at 150,000 cells per well. The ROCK inhibitor enhances the

survival of stem cells after plating for differentiation. Beginning the next day, cells were treated with Activin A (100 ng ml-1; Cell Guidance Systems) for three days in RPMI 1640 (Invitrogen) containing increasing concentrations of 0%, 0.2%, and 2.0% define fetal bovine serum (dFBS; Invitrogen).

[00148] **Differentiation of Definitive Endoderm (DE)**

[00149] For differentiation, PSCs were plated as single cells in a Matrigel-coated 24-well dish using accutase (Stem Cell Technologies), at a density of 150,000 cells per well in mTesR1 with ROCK inhibitor Y-27632 (10 μM; Stemgent). On the following day, PSCs were differentiated to DE as previously described[11, 35]. Cells were exposed to Activin A (100 ng ml-1; Cell Guidance Systems) for three days in RPMI 1640 media (Invitrogen) containing increasing concentrations of 0%, 0.2%, and 2.0% define fetal bovine serum (dFBS; Invitrogen). In addition, BMP4 (50 ng ml-1; R&D Systems) was added on the first day of DE induction.

[00150] **Endoderm patterning and gut tube morphogenesis.**

[00151] Following DE induction, cells were treated for three days with growth factors/antagonists in RPMI 1640 with 2.0% dFBS. To generate posterior foregut spheroids, DE was treated for three days with noggin (200 ng ml-1; R&D Systems), FGF4 (500 ng ml-1; R&D Systems), and either WNT3A (500 ng ml-1; R&D Systems) or CHIR99021 (2 μM; Stemgent). CHIR99021 is a small molecule that stimulates the Wnt signaling pathway. RA (2 μM; Sigma Aldrich) is added on the final day. Three-dimensional growth and antral specification. Posterior foregut spheroids were embedded in Matrigel (BD Biosciences) as previously described[10, 12] and subsequently grown in Advanced DMEM/F12 (Invitrogen) supplemented with N2 (Invitrogen), B27 (Invitrogen), L-glutamine,

10 μM HEPES, penicillin/streptomycin, and EGF (100 ng ml-1; R&D Systems). For antral specification, RA and noggin were added for the first three days of three-dimensional growth. For endocrine cell specification, the EGF concentration is lowered to 10 ng ml^{-1} at day 30.

[00152] **Endoderm patterning and foregut spheroid generation**

[00153] Following DE induction, cells were cultured in RPMI 1640 media with 2.0% dFBS and growth factors: WNT3A (500 ng ml-1; R&D Systems), CHIR99021 (2 μM; Stemgent); FGF4 (500 ng ml-1; R&D Systems), and Noggin (200 ng ml-1; R&D Systems). The media was changed every day. After three days, the combination of WNT3A (or CHIR99021), FGF4, and Noggin resulted in floating foregut spheroids in the culture wells. To posteriorize the foregut endoderm, RA (2 μM; Sigma Aldrich) was added on the third day of WNT/FGF/Noggin treatment.

[00154] **Three-dimensional culture of gastric organoids**

[00155] The spheroids were transferred to a three-dimensional in vitro culture system as previously described5,10,12. Briefly, spheroids were collected, resuspended in 50 μl Matrigel (BD Biosciences), and plated in a three-dimensional droplet. After Matrigel was allowed to solidify for 10-15 minutes in a tissue culture incubator, spheroids were overlaid with gut media: Advanced DMEM/F12 with N2 (Invitrogen), B27 (Invitrogen), L-glutamine, 10 μM HEPES, penicillin/streptomycin, and EGF (100 ng ml-1; R&D Systems). For the first three days, RA and Noggin were added to the gut media. Media was replaced every 3-4 days, as necessary. At day 20, organoids were collected and re-plated in fresh Matrigel at dilution of ~1:12.

[00156] **Generation of dox-inducible hNEUROG3 hESC line**

[00157] To generate over-expression construct, hNEUROG3 cDNA (Dana-Farber/Harvard Cancer Center DNA Resource Core; clone HsCD00345898) was cloned into pInducer20 lentiviral vector (gift from T. Westbrook36) using Gateway Cloning (Invitrogen) methods. High-titer lentiviral particles were produced by the CCHMC Viral Vector Core. H1 hESCs were dissociated with Accutase, plated as a single cell suspension in mTesR1 with 10 μM Y-27632, and exposed to lentivirus for four hours. mTesR1 was replaced daily and after two days, G418 (200 μg ml-1) was added to the media to select for integrating clones. G418-resistant cells were maintained in antibiotic indefinitely, but were otherwise cultured and passaged normally.

[00158] **Generation and characterization of iPSC lines**

[00159] Primary human foreskin fibroblasts (HFFs) were cultured from neonatal human foreskin tissue and obtained from 2 donors through the Department of Dermatology, University of Cincinnati, and were a kind gift from Susanne Wells PhD. HFFs were cultured in Fibroblast Media consisting of DMEM (Invitrogen) supplemented with 10% FCS (Hyclone) and used for reprogramming between passages 5 and 8. EBNA1/OriP-based episomal plasmids pCLXE-hOct3/4-shp53, pCLXE-hSox2-Klf4, pCLXE-hLmyc-Lin28, and pCLXE-GFP used for this study were previously described37 and obtained from Addgene (ID #s: 27077, 27078, 27080, and 27082 respectively). The optimized Human Dermal Fibroblast Nucleofector Kit (VPD-1001; Lonza) was used for transfection of HFFs with episomal plasmids. Briefly, for each transfection 1 x 106 HFFs were pelleted by centrifugation at 200xg for 10 minutes at room temperature, resuspended in 100 μl room temperature Nucleofector Solution and nucleofected with 1.25 μg each episomal plasmid (program U20). Cells from 2 transfections (2 x

106 total cells) were replated in a 10 cm tissue culture plate in Fibroblast Media, and cultured at 37°C/5% CO2. Six days post-transfection, 4.5 x 105 HFFs were replated in Fibroblast Media in a gelatin-coated 10cm dish containing 1.07 x 106 irradiated mouse embryonic fibroblasts (MEFs). Starting on day 7 post-transfection, cells were fed daily with DMEM/F12 media supplemented with 20% knockout serum replacement, 1 mM L-glutamine, 0.1 mM β-mercaptoethanol, 0.1 mM non-essential amino acids, and 4 ng ml-1 basic FGF (all from Invitrogen). Approximately 2 weeks later, discrete colonies with hESC-like morphology were manually excised and replated in mTeSR1 media (Stem Cell Technologies) in tissue culture dishes coated with hESC-qualified matrigel (Becton Dickinson). Following adaptation to mTeSR1/matrigel culture, iPSCs that maintained robust proliferation and hESC-like morphology with minimal spontaneous differentiation were expanded for cryopreservation and characterization.

[00160] Standard metaphase spreads and G-banded karyotypes were determined by the CCHMC Cytogenetics Laboratory. For teratoma formation, iPSCs from 3 wells of a 6-well dish were combined and gently resuspended in ice-cold DMEM/F12. Immediately before injection, matrigel was added to a final concentration of ~33% and cells were injected subcutaneously into immune-compromised NOD/SCID GAMMA C-/- mice. Tumors formed within 6-12 weeks. Excised teratomas were fixed, embedded in paraffin, and sections were stained with hematoxylin and eosin for histological examination.

[00161] **Exemplary Protocol for Gastric Organoid**

[00162] The following Table illustrates an exemplary treatment protocol for the development of a gastric organoid from a precursor cell. **Basal Gut Media = Advanced DMEM/F12 + B27 + N2 + l-

glutamine + HEPES; mTesR1 is available from StemCell Technologies.

Day	Media	Treatment	Stage of Development/Outcome
-1	mTesR1	Y27632 (10 uM)	
0	RPMI	Activin A (100 ng/mL) + BMP4 (50 ng/mL)	Monolayer hPSCs (ready to differentiate)
1	RPMI + 0.2% dFCS	Activin A (100 ng/mL)	
2	RPMI + 2.0% dFCS	Activin A (100 ng/mL)	
3	RPMI + 2.0% dFCS	FGF4 (500 ng/mL) + Wnt3a (500 ng/mL) OR CHIR99021 (2 uM)	Definitive Endoderm
4	RPMI + 2.0% dFCS	FGF4 (500 ng/mL) + Wnt3a (500 ng/mL) OR CHIR99021 (2 uM)	
5	RPMI + 2.0% dFCS	RA (2 uM) + FGF4 (500 ng/mL) + Wnt3a (500 ng/mL) OR CHIR99021 (2 uM)	
6	Basal Gut Media	EGF (100 ng/mL) + RA (2 uM) + Noggin (200 ng/mL)	3-D Posterior Foregut Spheroid
7	Basal Gut Media**	EGF (100 ng/mL) + RA (2 uM) + Noggin (200 ng/mL)	
8	Basal Gut Media**	EGF (100 ng/mL) + RA (2 uM) + Noggin (200 ng/mL)	
9	Basal Gut Media**	EGF (100 ng/mL) + Noggin (200 ng/mL)	Presumptive Antral Epithelium (Pdx1/Sox2)
10	Basal Gut Media**	EGF (100 ng/mL) + Noggin (200 ng/mL)	
11	Basal Gut Media**	EGF (100 ng/mL) + Noggin (200 ng/mL)	
12	Basal Gut Media**	EGF (100 ng/mL) + Noggin (200 ng/mL)	
13	Basal Gut Media**	EGF (100 ng/mL)	
14	Basal Gut Media**	EGF (100 ng/mL)	
15	Basal Gut Media**	EGF (100 ng/mL)	
16	Basal Gut Media**	EGF (100 ng/mL)	
17	Basal Gut Media**	EGF (100 ng/mL)	
18	Basal Gut Media**	EGF (100 ng/mL)	

19	Basal Gut Media**	EGF (100 ng/mL)	
20	Basal Gut Media**	EGF (100 ng/mL)	
21	Basal Gut Media**	EGF (100 ng/mL)	
22	Basal Gut Media**	EGF (100 ng/mL)	
23	Basal Gut Media**	EGF (100 ng/mL)	
24	Basal Gut Media**	EGF (100 ng/mL)	
25	Basal Gut Media**	EGF (100 ng/mL)	
26	Basal Gut Media**	EGF (100 ng/mL)	
27	Basal Gut Media**	EGF (100 ng/mL)	
28	Basal Gut Media**	EGF (100 ng/mL)	
29	Basal Gut Media**	EGF (100 ng/mL)	
30	Basal Gut Media**	EGF (100 ng/mL)	
31	Basal Gut Media**	EGF (10 ng/mL)	
32	Basal Gut Media**	EGF (10 ng/mL)	
33	Basal Gut Media**	EGF (10 ng/mL)	
34	Basal Gut Media**	EGF (10 ng/mL)	Gastric Organoids with Differentiated Cell Types

[00163] **Fundus Specification Protocol**

[00164] Applicant first sought to identify genes specifically expressed in the fundus, but not in the antrum, at embryonic stages. The digestive tracts of E14.5 mouse embryos were micro-dissected and separated into four regions: forestomach (including esophagus), fundus, antrum, and duodenum. See FIG 15. These regions were then analyzed by qPCR for markers of regionalization. FIG 15 shows expression of control genes known to be expressed in different regions. The fundus and antrum can be distinguished from the forestomach and duodenum by their high expression of Sox2 and Gata4, and absence of P63 and Cdx2. Importantly, Pdx1

(marker of antrum) is expressed at much higher levels in the antrum tissue than in the fundus, indicating accurate dissection.

[00165] Bioinformatics analysis of published microarray datasets from embryonic mouse endoderm and adult human stomach tissue were used to generate a list of candidate genes that may be preferentially expressed in the fundus but not antrum. Expression of these putative markers in the E14.5 mouse segments was examined by qPCR. Irx1, Irx2, Irx3, Irx5, and Pitx1 are indeed expressed at higher levels in the fundus than in the antrum. Thus these markers may be used as indicators of fundus specification in the hPSC-derived foregut cultures. See FIG 16.

[00166] Next, the function of Wnt signaling in the regulation of fundus-antrum patterning at days 6-9 of the gastric organoid differentiation protocol was tested. Addition of Wnt3a (at 100 ng/mL and 500 ng/mL) had no effect on spheroid gene expression, but it also did induce expression of the Wnt target gene Axin2. Thus, the effects of the small molecule CHIR99021 (CHIR; 2 uM) were tested. CHIR99021 stimulates Wnt signaling in a receptor-independent fashion. Exposure to CHIR resulted in a robust repression of Pdx1 expression levels, consistent with fundus specification. CHIR did not induce expression of the intestinal marker Cdx2. See FIG 17. FIG 18 shows that, consistent with repression of Pdx1, exposure to CHIR induced high levels of expression of fundus-specific markers IRX3 and IRX5.

[00167] **H. pylori infection**

[00168] H. pylori strain G2738 and a mutant G27 strain lacking CagA (ΔCagA)39 were grown on blood agar plates consisting of Columbia Agar Base (Fisher Scientific), 5% horse blood (Colorado Serum Company), 5 μg ml-1, vancomycin and 10 μg ml-1

trimethoprim as described previously 40. For organoid injections, H. pylori were resuspended in brucella broth at a concentration of 1 x 109 bacteria ml-1 and loaded onto the Nanoject II (Drummond) microinjector apparatus. Approximately 200 nl (containing 2 x 105 bacteria) were injected directly in the lumen of each organoid, and injected organoids were cultured for 24 hours. Brucella broth was injected as a negative control.

[00169] **Materials and Methods**

[00170] **Immunofluorescent staining**

[00171] All tissues were fixed in 4% paraformaldehyde for either one hour at room temperature for frozen processing or overnight at 4°C for paraffin processing. For frozen sections, tissue was protected in 30% sucrose overnight at 4°C, then embedded in OCT (Tissue-Tek), and cut at 10 μm. For paraffin sections, tissue was processed through a graded ethanol series, followed by xylene, and then embedded in paraffin and cut at 7 μm. Tissue culture cells were fixed for 15 minutes at room temperature and stained directly. For staining, frozen slides were thawed to room temperature and rehydrated in PBS, while paraffin slides were deparaffinized and subjected to antigen retrieval. Slides were blocked in 5% normal donkey serum (Jackson Immuno Research) in PBS plus 0.5% Triton-X for 30 minutes at room temperature. Primary antibodies (listed in Methods Table 1) were diluted in blocking buffer and incubated overnight at 4°C. Slides were washed in PBS and incubated with secondary antibody for one hour at room temperature, and coverslips were mounted using Fluoromount-G (Southern Biotech). Confocal images were captured on a Nikon A1Rsi inverted confocal microscope.

[00172] **RNA isolation and qPCR**

[00173] Total RNA was isolated from tissues using the Nucleospin RNA II kit (Machery-Nagel). Reverse transcription was performed from 100 ng RNA using Superscript VILO cDNA Synthesis Kit (Invitrogen) according to manufacturer's protocol. qPCR was done using Quantitect SybrGreen Master Mix (Qiagen) on a CFX-96 Real-time PCR Detection System (BioRad). Analysis was performed using the ΔΔCT method. PCR primers were designed using sequences from qPrimerDepot (http://primerdepot.nci.nih.gov) and are listed in Table 2.

[00174] **Immunoprecipitation and Western blot analysis**

[00175] H. pylori-infected organoids were harvested from Matrigel in ice-cold PBS and centrifuged at 150 g for 5 minutes. Tissue was lysed in M-PER Mammalian Protein Extract Reagent (Thermo Scientific) supplemented with protease inhibitors (Roche). 10 µg total protein from the cell lysates were immunoprecipitated with anti-c-Met antibody (2 µg; Cell Signaling 4560) at 4°C for 16 hours. Protein A/G agarose beads (20 µl; Santa Cruz Biotechnology) were then added and the samples were incubated at 4°C for 16 hours. Immunoprecipitates were washed 3 times in PBS and then resuspended in Laemmli Loading Buffer containing β-mercaptoethanol (40 µl; BioRad). Samples were run on a 4-20% Tris-Glycine Gradient Gel (Invitrogen) and run at 80 V for 3.5 hours. Gels were transferred to nitrocellulose membranes (Whatman Protran, 0.45 µm) at 105 V for 1.5 hours. Membranes were blocked in KPL Detector Block Solution (Kirkeaard & Perry Laboratories) for one hour at room temperature and then incubated with primary antibody overnight at 4°C. Primary antibodies used: anti-phosphotyrosine (Santa Cruz, sc-7020; 1:100), anti-c-Met (Abcam, ab59884; 1:100), and anti-H. pylori CagA (Abcam, ab90490; 1:100). Membranes were washed and incubated in Alexa

Fluor anti-mouse 680 (Invitrogen; 1:1000) secondary antibody. Blots were imaged using the Odyssey Infrared Imaging Software System (Licor).

[00176] **Discussion**

[00177] hPSCs were differentiated into definitive endoderm (DE) [11], which gives rise to the epithelia of the gastrointestinal and respiratory tracts in vivo. The next two essential events in development of all endoderm organs are patterning of DE along the anterior-to-posterior (A-P) axis and gut tube morphogenesis, resulting in formation of the Sox2+ foregut in the anterior and the Cdx2+ mid-hindgut in the posterior (as highlighted in the E8.5, 14 somite stage mouse embryo, FIG. 1A). This morphogenesis and the tissue interactions between endoderm and mesoderm appear to be critical for proper organogenesis both *in vivo* and *in vitro*. WNT3A and FGF4 have been previously demonstrated to synergize to do three things: posteriorize hPSC-derived DE, promote the expansion of mesenchyme, and induce the assembly of gut tube-like structures expressing the mid-hindgut marker CDX2[10,12]. FIG 1A shows that Sox2 protein marks foregut endoderm and Cdx2 protein marks mid/hindgut endoderm in e8.5 (14 somite stage) mouse embryos. FIG 1B shows that inhibiting BMP repressed mid/hindgut fate and promoted expression of the foregut marker SOX2. PCR analysis of patterning markers in hPSC-DE cultures exposed to three days in media alone (control) or with the indicated growth factors/antagonists. The combined activity of WNT and FGF induced CDX2 expression as previously reported[10] whereas the BMP antagonist noggin repressed CDX2 expression and was sufficient to induce high levels of the foregut marker SOX2. *, $p<0.05$ compared to control. **, $p<0.005$ compared to WNT/FGF. FIG1C depicts that foregut spheroids generated with

Wnt/FGF/Noggin have high levels of SOX2 protein by wholemount immunofluorescence staining and mRNA as compared to spheroids generated with Wnt and FGF alone, which have high levels of CDX2. *, p<1.0x10-6. FIG 1D depicts that the posterior foregut in an e8.5, 14-somite stage mouse embryo gives rise to the stomach and pancreas and has high levels of Hnf1β protein. FIG 1E depicts that exposing cultures to RA on the final day of the spheroid generation step induces expression of HNF1β in SOX2-expressing epithelium, resulting in the formation of posterior foregut spheroids. *, p<0.005. FIG 1F depicts a lineage diagram that summarizes the patterning effects of noggin and RA in the formation of both anterior and posterior foregut endoderm. Scale bars, 100 μm. Error bars represent standard deviation.

[00178] Table 1. Primary Antibodies.

Antibody	Species	Company	Product No.	Dilution
alpha-SM-actin	rabbit	GeneTex	GTX100034	1:200
aPKC	rabbit	Santa Cruz	sc216	1:200
B-catenin	rabbit	Santa Cruz	sc7199	1:100
Cdx2	mouse	Biogenex	MU392A-UC	1:500
ChrA	rabbit	Immunostar	20086	1:500
Desmin	goat	Santa Cruz	sc7559	1:200
E-Cadherin	mouse	BD Biosciences	610182	1:500
E-Cadherin	goat	R&D Systems	AF648	1:500
FoxF1	goat	R&D Systems	AF4798	1:500
Gastrin	rabbit	Dako	A0568	1:1000
Gata4	mouse	Santa Cruz	sc25310	1:200
Ghrelin	goat	Santa Cruz	sc10368	1:200
H. pylori	rabbit	Abcam	ab80519	1:1000
Hnf1B	mouse	BD Biosciences	612504	1:500
Hnf1B	goat	Santa Cruz	sc4711	1:500
Ki67	rabbit	Abcam	ab833	1:200
Ki67	rat	Dako	m7249	1:100
Klf5	rat	Dr. Ryozo Nagai	Shindo et al., 2002	1:2000
Muc5AC	mouse	Abcam	ab3649	1:500
Nanog	rabbit	Abcam	ab21624	1:500
Oct3/4	mouse	Santa Cruz	sc5279	1:500

Pdx1	goat	Abcam	ab47383	1:5000
pHH3	rabbit	Cell Signaling	9701	1:500
Serotonin (5-HT)	rabbit	Immunostar	20080	1:1000
Somatostatin	goat	Santa Cruz	sc7819	1:100
Sox2	goat	Santa Cruz	sc17320	1:500
Sox2	rabbit	Seven Hills Bioreagents	WRAB-1236	1:1000
Tff2	goat	Santa Cruz	sc23558	1:500
Vimentin	goat	Santa Cruz	sc7557	1:200

[00179] Table 2. qPCR Primer Sequences.

Target (Human)	Forward Primer	Reverse Primer
ATP4A	TGGTAGTAGCCAAAGCAGCC	TGCCATCCAGGCTAGTGAG
ATP4B	ACCACGTAGAAGGCCACGTA	TGGAGGAGTTCCAGCGTTAC
AXIN2	CTGGTGCAAAGACATAGCCA	AGTGTGAGGTCCACGGAAAC
BAPX1	CAACACCGTCGTCCTCG	CCGCTTCCAAAGACCTAGAG
CDX2	CTGGAGCTGGAGAAGGAGTTTC	ATTTTAACCTGCCTCTCAGAGAGC
CHGA	TGACCTCAACGATGCATTTC	CTGTCCTGGCTCTTCTGCTC
GAPDH	CCCATCACCATCTTCCAGGAG	CTTCTCCATGGTGGTGAAGACG
GAST	CAGAGCCAGTGCAAAGATCA	AGAGACCTGAGAGGCACCAG
GATA4	TCCAAACCAGAAAACGGAAGC	GCCCGTAGTGAGATGACAGG
GHRL	GCTGGTACTGAACCCCTGAC	GATGGAGGTCAAGCAGAAGG
GKN1	AGCTAGGGCAGGAGCTAGAAA	GCTTGCCTACTCCTCTGTCC
HNF1B	TCACAGATACCAGCAGCATCAGT	GGGCATCACCAGGCTTGTA
HNF6	TGTTGCCTCTATCCTTCCCA	GGAGGATGTGGAAGTGGCT
IRX1	CCGTAGGGGTAATAAGCCG	ATCTCAGCCTCTTCTCGCAG
IRX2	GTGGTGTGCGCGTCGTA	GGCGTTCAGCCCCTACC
IRX3	GGAGAGAGCCGATAAGACCA	AGTGCCTTGGAAGTGGAGAA
IRX5	GGTGTGTGGTCGTAGGGAGA	GCTACAACTCGCACCTCCA
MIST1	TGCTGGACATGGTCAGGAT	CGGACAAGAAGCTCTCCAAG
MSX1	GGTTCGTCTTGTGTTTGCG	CCCGAGAAGCCCGAGAG
MSX2	GGTCTTGTGTTTCCTCAGGG	AAATTCAGAAGATGGAGCGG
MUC2	TGTAGGCATCGCTCTTCTCA	GACACCATCTACCTCACCCG
MUC5AC	CCAAGGAGAACCTCCCATAT	CCAAGCGTCATTCCTGAG
MUC6	CAGCAGGAGGAGATCACGTTCAAG	GTGGGTGTTTTCCTGTCTGTCATC
NEUROG3	CTTCGTCTTCCGAGGCTCT	CTATTCTTTTGCGCCGGTAG
PDX1	CGTCCGCTTGTTCTCCTC	CCTTTCCCATGGATGAAGTC
PITX1	TTCTTGGCTGGGTCGTCT	TCGTCTGACACGGAGCTG
PTF1a	AGAGAGTGTCCTGCTAGGGG	CCAGAAGGTCATCATCTGCC

SST	GCGCTGTCCATCGTCCTGGCCC	AGCCGGGTTTGAGTTAGCAGAT
SOX2	GCTTAGCCTCGTCGATGAAC	AACCCCAAGATGCACAACTC
TFF1	AATTCTGTCTTTCACGGGGG	GGAGAACAAGGTGATCTGCG
TFF2	TCTGAGACCTCCATGACGC	ATGGATGCTGTTTCGACTCC
TFF3	CACTCCTTGGGGGTGACA	CTCCAGCTCTGCTGAGGAGT

[00180] To promote the formation of foregut structures in the hPSC-derived DE, Applicant sought to separate the ability of WNT/FGF to stimulate gut tube morphogenesis from their role in promoting a posterior endoderm fate. Based on *in vivo* studies from developmental model organisms[13, 14], the function of BMP signaling in regulating A-P patterning was tested and Applicant determined that WNT/FGF require BMP activity to initiate the hindgut program. Specifically, inhibition of BMP signaling with the antagonist Noggin repressed CDX2 and induced the foregut marker SOX2 in DE cultures after three days, even in the presence of WNT/FGF (FIG 1B-C and FIG 5). Importantly, inhibition of BMP signaling had no effect on the ability of WNT/FGF to promote mesenchyme expansion and assembly of gut tube structures, thus resulting in the formation of $SOX2^+$ foregut spheroids.

[00181] FIG 5 shows that BMP signaling is required in parallel with activation of WNT and FGF to promote a posterior fate. FIG 5A shows that the GSK3β inhibitor CHIR99021 (CHIR; 2 μM) induced the same posteriorizing effects as recombinant WNT3A and these can be blocked by BMP inhibition. FIG 5B shows that CHIR induced gut tube morphogenesis and spheroid production occurs in a similar manner to WNT3A. FIG 5C depicts immunofluorescent staining of monolayer cultures which confirms the high efficiency of CDX2 induction in CHIR/FGF-treated endoderm and SOX2 induction in noggin- and CHIR/FGF/noggin-treated endoderm. FIG 5D shows qPCR analysis of BMP target

genes MSX1/2, which indicates that BMP activity is not increased in response to Wnt/FGF, but target genes are suppressed in response to noggin, demonstrating the presence of endogenous BMP signaling. FIG 5E shows that addition of BMP2 (100 ng mL-1) did not substitute for or augment the ability of Wnt/FGF to posteriorize endoderm. These data indicate that the posteriorizing effect of Wnt/FGF is not mediated by up-regulation of BMP signaling but does require endogenous BMP activity. Scale bars, 1 mm in FIG 5B; 100 μm in FIG 5C. Error bars represent standard deviation.

[00182] Spheroid morphogenesis is a robust process in both hESC and hiPSC lines (FIG 6A) and >90% of spheroid cells express SOX2 (FIG 1C), indicating efficient specification into the foregut lineage. Thus, a new epistatic relationship between WNT, FGF and BMP has been identified by Applicant in which all three pathways cooperate to promote a mid-hindgut fate, whereas WNT and FGF act separately from BMP to drive assembly of endoderm and mesoderm into gut tube structures.

[00183] FIGS 2A-2G depicts gastric organoid differentiation is an efficient and cell line-independent process. FIG 2A, Table comparing spheroid formation and characteristics between two hESC lines (H1 and H9) and one iPSC line (72.3). FIG 2B, Immunofluorescent staining of day 34 hGOs derived from H1 and iPSC 72.3 cell lines. iPSC-derived organoids exhibit the same morphological and molecular features of those derived from hESCs. FIG 2C. Organ epithelial cell type quantification in day 34 hGOs. Greater than 90% of the epithelium is antral, indicated by PDX1 expression and lack of PTF1A expression, whereas less than 5% express markers associated with other organs derived from endoderm including CDX2 (intestine), albumin (liver) and p63 (squamous epithelium).

d-g, Characterization of induced pluripotent stem cell line iPSC 72.3. FIG 2D, iPSC 72.3 exhibits normal morphological characteristics of pluripotent stem cell colonies, as compared to the H1 hESC line and FIG 2E, has a normal 46;XY karyotype. FIG 2F, iPSC 72.3 expresses pluripotent markers OCT3/4 and NANOG, and FIG 2G, demonstrates pluripotency by differentiation into endoderm, mesoderm, and ectoderm lineages in an in vivo teratoma assay. Scale bars, 100 μm. Error bars represent standard deviation.

[00184] *In vivo*, both the fundic and antral domains of the stomach arise from the posterior segment of $Sox2^+$ foregut endoderm, along with the pancreas, liver and duodenum. To direct $SOX2^+$ foregut spheroids into the gastric lineage, Applicant sought to identify signaling pathways that promote posterior foregut fate. Applicant focused on retinoic acid (RA) signaling given its role in development of posterior foregut-derived organs.[15-17] *In vivo*, the posterior foregut is marked by expression of Hnf1β (FIG 1D. Applicant identified that a 24-hour exposure to RA on the final day (days 5-6) of the patterning/spheroid generation stage (FGF4/WNT3A/Noggin) results in robust activation of posterior foregut markers and the formation of $SOX2/HNF1\beta^+$ posterior foregut spheroids (*FIG 1E and FIG 7*). *Thus, the precise temporal* and combinatorial manipulation of RA, WNT, FGF, and BMP signaling pathways allowed for the generation of three-dimensional posterior foregut spheroids.

[00185] FIGS 7A-7D shows that retinoic acid posteriorizes foregut endoderm. FIG 7A depicts a schematic illustrating of foregut patterning experiments. DE cultures were treated with Wnt(CHIR)/FGF/noggin for three days to generate Sox2-positive foregut spheroids, and RA is added for 24 hours on the third day of

patterning. FIG 7B depicts Brightfield images that show that RA increases the number of spheroids that are produced from foregut monolayer cultures. FIG 7C depicts a lower power image of FIG1D showing immunofluorescent image of a 14 somite stage embryo with *Hnf1β* protein localized to the posterior portion of the foregut. Boxed region of embryo is shown in FIG 1D. FIG 7D shows qPCR analysis of gene expression in foregut spheroids treated with RA. Posterior foregut markers HNF1β and HNF6 are robustly induced by 24-hour exposure to RA. *, $p<0.05$. Scale bars, 1 mm in FIG 7B; 100 μm in FIG 7C. Error bars represent standard deviation.

[00186] The molecular mechanisms that direct the posterior foregut into distinct organ lineages are poorly understood. Early in development, presumptive organ domains are marked by distinct gene expression patterns: $Sox2^+$ fundus, $Sox2/Pdx1^+$ antrum, $Pdx1/Ptf1\alpha^+$ pancreas, and $Pdx1/Cdx2^+$ duodenum (FIG 2B). Applicant used these molecular markers to identify signaling pathways that direct posterior foregut spheroid cultures into the gastric lineage. Following transfer of spheroids to three-dimensional culture conditions, further treatment with RA for 72 hours (days 6-9) caused a >100-fold increase in *PDX1* mRNA levels while maintaining high *SOX2* expression (FIG 2C). Importantly, the RA treatment did not promote a pancreatic fate as observed by others[9], since expression of the pancreas-specific marker *PTF1α* was not induced. These data demonstrate that the combination of RA signaling with three-dimensional growth efficiently direct posterior foregut spheroids into a $SOX2/PDX1^+$ epithelium indicative of an early antrum fate.

[00187] FIG 2 generally depicts the specification and growth of human antral gastric organoids. Error bars represent standard deviation.

FIG2A depicts schematic representation of the *in vitro* culture system used to direct the differentiation of hPSCs into three-dimensional gastric organoids, FIG2B depicts the defining markers of developing posterior foregut organs by wholemount immunofluorescent staining of mouse E10.5 embryos with Sox2, Pdx1 and Cdx2. Co-expression of Sox2 and Pdx1 is unique to the distal portion of the gastric epithelium, the presumptive antrum (a), Sox2 expression marks the fundus (f), Pdx1 (and Ptf1a) expression marks the dorsal (dp) and ventral (vp) pancreas, and Pdx1/Cdx2 co-expression marks the duodenum (d). FIG2C shows that the posterior foregut spheroids cultured in three-dimensional matrix for three days in the presence of RA (2 μM) co-expressed high levels of PDX1 and SOX2 and did not express the pancreatic marker PTF1α, similar to the developing antrum *, $p<0.05$. FIG2D depicts stereomicrographs showing morphological changes during growth of posterior forgut spheroids into gastric organoids. By four weeks, the epithelium of hGOs exhibited a complex glandular architecture, scale bar, 500 μm. FIG2E depicts a comparison of developing mouse antrum at E14.5 and E18.5 and comparable stages of hGO development. Sox2 and Pdx1 are co-expressed in the early pseudostratified epithelia in both mouse antrum and hGOs. At later stages, Sox2 is down-regulated as the epithelia transform into more mature glandular structures. Pdx1 is maintained in the antrum throughout adulthood *in vivo* and at all stages examined in hGOs, scale bars 100 μm in FIG2E.

[00188] Applicant used SOX2/PDX1$^+$ spheroids to identify pathways that promote growth and morphogenesis of the early gastric epithelium and found that high concentrations of EGF (100 ng mL^{-1}) were sufficient to promote robust outgrowth of human antral gastric organoids (hGOs). Over the course of 3-4 weeks, spheroids that were <100 μm in diameter grew into organoids that were 2-4 mm

in diameter. At the later stages of culture (~ day 27), the hGO epithelium underwent a series of morphogenetic changes reminiscent of the late stages of embryonic gastric development, during which a simple, flat pseudostratified epithelium transitions into an elaborate, convoluted glandular epithelium (FIG 2D). The initial outgrowth of foregut spheroids is dependent on EGF (data not shown); moreover, epithelial expansion and morphogenesis into glands does not occur when EGF is removed from the media at day 27 (FIG 8). These results support published findings that indicate an important role for EGF in promoting proper growth of the gastric mucosa[19, 20].

[00189] FIG 8 shows that EGF is required for glandular morphogenesis in gastric organoids. Brightfield images and immunostaining demonstrate the requirement for EGF for epithelial morphogenesis and gland formation at late stages of hGO differentiation. When EGF is removed from the growth medium at day 27, prior to glandular morphogenesis, the hGO epithelium retains a simple, cuboidal structure that fails to form glands. Scale bars, 100 μm.

[00190] A comparison of hGO growth with development of the embryonic mouse stomach revealed that hGO development is astonishingly similar to *in vivo* stomach organogenesis. At early stages (E12-14 in mouse and 13-day hGOs), both epithelia are pseudostratified and contain mitotic cells that are concentrated toward the luminal face (FIG 9 and FIG 10), indicating an interkinetic nuclear migration process[21]. The early hGOs are appropriately polarized and contain secondary lumina that are outlined by expression of the apical marker aPKC[22] (FIG 10).

[00191] Between E16.5 and early postnatal stages, the antrum transforms into a simple columnar epithelium exhibiting a highly structured organization consisting of glands and pits (FIG 2E and FIG 9).

Between 13 and 34 days *in vitro*, the hGO epithelium undergoes similar transitions to form a tall columnar epithelium with a glandular structure similar to the late fetal antrum (FIG 2E). Analysis of expression of the transcription factors Sox2, Pdx1, Gata4 and Klf5 revealed a stereotypic temporospatial expression pattern that accompany these morphogenetic processes both *in vivo* and *in vitro* (FIG 9). At early stages these factors are all co-expressed in the immature, pseudostratified epithelium. However at later stages, Sox2 expression is down-regulated as the epithelium forms early glands and pits, whereas the expression of the other factors is maintained indefinitely. Based on these data, it is estimated that 13-day hGOs represent a developmental stage similar to the E12-14 mouse antrum, whereas 34-day hGOs are more comparable to the late fetal-early postnatal antrum. Further, it is concluded that hGOs recapitulate normal embryonic development and that the molecular and morphogenetic processes that occur during antrum development are conserved between rodents and humans.

[00192] FIG 9 shows a comparison of transcription factor expression during development of the mouse antrum and human gastric organoids. Four embryonic stages (E12.5, E14.5, E16.5 and E18.5) and one postnatal stage (P12) of *in vivo* antrum development were analyzed for transcription factor expression: Sox2, Pdx1, Gata4, Klf5, and FoxF1. The same markers were analyzed at two stages (day 13 and day 34) of *in vitro* hGO development and revealed that organoid development parallels what occurs in vivo. At early stages of antrum development the epithelial marker Sox2 is expressed ubiquitously but at later stages it is down-regulated, while other epithelial transcription factors, Pdx1, Gata4 and Klf5, exhibit persistent expression throughout development. Both early and late stage hGOs contain FoxF1-positive mesenchymal cells surrounding

the epithelium. Scale bars, 100 μm. FIG 10 shows that early stage human gastric organoids exhibit stereotypic architecture and nuclear behavior. At 13 days, hGOs contain pseudostratified epithelia that display apicobasal polarity marked by the apical marker aPKC and the basolateral marker E-Cadherin, similar to the E12.5 mouse antrum. Further, secondary lumina lined by apical membrane (white arrows) are seen within the organoid epithelium. Both the E12.5 mouse antrum and day 7 hGOs appear to undergo interkinetic nuclear migration, indicated by the presence of mitotic nuclei, pHH3, in only the apical portions of cells. Scale bars, 50 μm.

[00193] The foregut spheroids contained a mesenchyme component similar to the mid-hindgut spheroids that were previously described[10]. During the differentiation into gastric organoids, the mesenchyme expands and expresses key transcription factors associated with antral mesenchyme development, including FOXF1 and BAPX1 (FIG 10 and FIG 11). At later stages, hGO mesenchyme largely consists of VIMENTIN$^+$ submucosal fibroblasts and a smaller number of ACTA2$^+$ subepithelial myofibroblasts (FIG 11), indicative of immature gastric mesenchyme. hGOs do not form differentiated layers of smooth muscle as occurs *in vivo*. Given that there is such robust epithelial morphogenesis in the absence of any exogenous factors except EGF, it seems likely that the mesenchyme plays a role in epithelial development. It is therefore surprising that the epithelium does not appear to promote robust differentiation of the mesenchyme. This suggests that other stimuli, possibly mechanical, play a role in gastric mesenchyme differentiation.

[00194] FIG 11 shows mesenchymal differentiation in gastric organoids. FIG 11A shows temporal expression analysis of the antral

mesenchyme transcription factor BAPX1. Similar to its known embryonic expression pattern, BAPX1 is up-regulated during the earlier stages of hGO differentiation and then down-regulated coincident with functional cell type marker expression. FIG 11B shows that staining for mesenchymal cell type markers reveals that day 34 hGOs contain FOXF1/VIMENTIN-positive submucosal fibroblasts and a small number of VIMENTIN/ALPHA-SM-ACTIN (SMA)-expressing subepithelial fibroblasts. hGOs lack a robust smooth muscle layer, indicated by SMA/Desmin-positive cells in the *in vivo* antrum. Scale bars, 100 μm. Error bars represent standard deviation.

[00195] The principal functional cell types found in the antrum are mucous cells, which secrete the protective mucus layers that line the gastric epithelium, and endocrine cells that secrete hormones to regulate gastrointestinal physiology and metabolic homeostasis[24]. By day 34, hGOs contain surface mucous cells (MUC5AC/UEAI$^+$) that secrete mucus into the lumen and have the same tall columnar morphology as their *in vivo* counterpart. hGOs also contain TFF2/GSII$^+$ antral gland cells, indicating appropriate differentiation in the antral mucous lineages (FIG 3A). In addition, hGOs develop a progenitor cell niche, indicated by basally located zones of proliferation and SOX9 expression (FIG 4A), although the proliferative index of the epithelium is variable and ranges between 1-10%. Thus, the *in vitro* hGOs contain a physiological gastric epithelium that comprises both progenitor and differentiated cell types.

[00196] FIG 4 shows that human gastric organoids exhibit acute responses to H. pylori infection. FIG 4A shows that Day 28 hGOs contained proliferative cells, marked by Ki67, and SOX9+ progenitor cells that were restricted toward the bottoms of the early glands, similar

to the late embryonic and postnatal mouse antrum. FIG 4B shows that hGOs were used to model human-specific disease processes of H. pylori infection. Bacteria were microinjected into the lumen of hGOs and bacteria were visualized in the lumen 24 hours post-injection by brightfield microscopy (black arrow) and immunofluorescent staining. FIG 4C depicts immunoprecipitation for the oncogene c-Met and demonstrates that H. pylori induced a robust activation (tyrosine phosphorylation) of c-Met, and that this is a CagA-dependent process. Further, CagA directly interacts with c-Met in human gastric epithelial cells. FIG 4D shows that within 24 hours, H. pylori infection caused a two-fold increase in the number of proliferating cells in the hGO epithelium, measured by EdU incorporation. *, $p<0.05$. Scale bars, 100 μm in a; 25 μm in b. Error bars represent s.e.m.

[00197] FIG3 demonstrates that human gastric organoids contain normal differentiated antral cell types and can be used to model human stomach development. FIG 3A demonstrates that hGOs contain all the major antral cell lineages. The 34-day hGOs have surface mucous cells (Muc5AC) and mucous gland cells (TFF2), as well as lectin staining that distinguishes surface mucous, UEAI, and mucous gland cells, GSII. hGOs also contain endocrine cells as marked by ChromograninA (CHGA). FIG 3B is a schematic representation of the different roles for EGF in the growth, morphogenesis, and cell type specification during development of hGOs. High levels of EGF were required at early developmental stages for gland formation, however it repressed endocrine differentiation at late stages of development; thus, the EGF concentration was reduced at day 30 to allow for endocrine cell development. FIG 3C shows that all major endocrine hormones are expressed in hGOs upon withdrawal of EGF including gastrin, ghrelin, and serotonin (5-HT). FIG 3D shows that high levels of

EGF repress NEUROG3 expression. A reduction in EGF concentration at day 30 resulted in a significant increase in NEUROG3 expression measured at day 34 by qPCR, indicating that EGF acts upstream of NEUROG3 in endocrine specification. *, $p<0.05$. FIG 3E shows that NEUROG3 acts downstream of EGF to induce endocrine cell fate. Forced expression of NEUROG3 using a dox-inducible system was sufficient to override the endo*crine-re*pressing effects of high EGF (100 ng mL-1). hGOs were exposed to dox (1 μg mL-1) for 24 hours at day 30 and analyzed at day 34. Dox-treated organoids exhibited a robust induction of ChrA-expressing endocrine cells. Scale bars, 100 μm. Error bars represent standard deviation.

[00198] There is also an abundance of CHROMOGRANIN-A $(CHGA)^+$ endocrine cells in 34-day hGOs, including the four main endocrine cell types in the antrum expressing gastrin, ghrelin, somatostatin, and serotonin (FIG 3C and FIG 12). Interestingly, we observed that high levels of EGF repress endocrine cell formation such that 100 ng ml^{-1} resulted in <1 endocrine cell per organoid. In contrast, hGOs cultured in lower levels of EGF (10 ng ml^{-1}) from day 30-34 developed an abundance of endocrine cells (FIG 13). Moreover, high EGF also inhibited expression of the proendocrine transcription factor NEUROG3 (FIG 3D), which has been extensively studied in the pancreas and intestine[25-28] and is required in the formation of most gastric endocrine lineages[29, 30]. These data suggest a novel inhibitory effect of EGFR signaling in gastric endocrine cell specification upstream of NEUROG3. To test this model, Applicant used a doxycycline-inducible hNEUROG3 over-expressing hESC line, and found that NEUROG3 expression was sufficient to overcome the endocrine inhibitory effect of high EGF (100 ng ml^{-1}), resulting in robust formation of $CHGA^+$ endocrine cells (FIG 3E and FIG 13). From these findings we concluded that

EGF inhibits the formation of endocrine progenitor cells through repression of NEUROG3, and that NEUROG3 is sufficient for the specification of human gastric endocrine cells.

[00199] FIG 12 shows gastric antrum endocrine cell development *in vivo*. Endocrine cell differentiation in the antrum is first evident at E18.5, but is more definitive at postnatal stages (P12 shown). At early stages, all expected gastric endocrine subtypes are evident, including gastrin, ghrelin, somatostatin, and serotonin (5-HT). Scale bars, 100 μm. FIG 13 shows that EGF signaling represses a NEUROG3-dependent gastric endocrine specification program. FIG 13A shows that hGOs maintained in high concentrations of EGF (100 ng mL-1) had very few endocrine cells at day 34, shown by staining for the pan-endocrine marker CHGA. A reduction of EGF concentration (10 ng mL-1) at day 24 resulted in more physiologic numbers of endocrine cells in the gastric epithelium. FIG 13B shows generation of hGOs from a hESC line stably transfected with a dox-inducible NEUROG3-overexpressing transgene, to test whether EGF repression of endocrine differentiation occurs upstream of NEUROG3. hGOs were maintained in high EGF (100 ng mL-1) then at day 30 were treated with doxycycline (1 μg mL-1) for 24 hours and then analyzed at day 34. Dox-treated hGOs show robust activation of endocrine markers CHGA, GASTRIN, GHRELIN, and SOMATOSTATIN, and they contain CHGA- (FIG 3A), GHRELIN-, and SOMATOSTATIN-positive cells with endocrine morphology. *, $p<0.05$. Scale bars, 100 μm. Error bars represent standard deviation.

[00200] Clinical evidence indicates that predominant colonization of the antrum has an important role in *H. pylori*-mediated disease[31,32]. Thus, Applicant tested whether hGOs could be used to model the

pathophysiologic response of human stomach to the pathogen *H. pylori*. To mimic the normal host-pathogen interface, we introduced *H. pylori* directly to the luminal surface of the epithelium by microinjection into the lumen of the organoids and measured epithelial signaling and proliferation (FIG 4). Bacteria were observed tightly associated with the hGO epithelium by immunofluorescence (FIG 4B). Within 24 hours, Applicant observed significant epithelial responses to *H. pylori* including robust activation of the gastric oncogene c-Met[33] and a 2-fold increase in epithelial cell proliferation. The *H. pylori* virulence factor CagA plays a pivotal role in the etiology of disease. Consistent with published studies[34], Applicant demonstrated that CagA translocates into the organoid epithelial cells and forms a complex with c-Met (FIG 4C). Furthermore, the epithelial response was abrogated when hGOs were injected with a non-pathogenic strain of *H. pylori* lacking CagA, reinforcing the importance of this factor in *H. pylori*-mediated human pathogenesis. Thus, the pathophysiological response of hGOs to *H. pylori* makes them an unprecedented model for elucidating the initiating events of human gastric disease mediated by *H. pylori*.

[00201] **Additional References**

[00202] 1. Wen, S. & Moss, S. F. Helicobacter pylori virulence factors in gastric carcinogenesis. Cancer Lett. 282, 1–8 (2009).

[00203] 2. Yuan, Y., Padol, I. T. & Hunt, R. H. Peptic ulcer disease today. Nat Clin Pract Gastroenterol Hepatol 3, 80–89 (2006).

[00204] 3. Parkin, D. M. The global health burden of infection-associated cancers in the year 2002. Int. J. Cancer 118, 3030–3044 (2006).

[00205] 4. Peek, R. M. Helicobacter pylori infection and disease: from humans to animal models. Dis Model Mech 1, 50–55 (2008).

[00206] 5. Barker, N. et al. Lgr5(+ve) stem cells drive self-renewal in the stomach and build long-lived gastric units *in vitro*. Cell Stem Cell 6, 25–36 (2010).

[00207] 6. Longmire, T. A. et al. Efficient Derivation of Purified Lungand Thyroid Progenitors from Embryonic Stem Cells. Stem Cell 10, 398–411 (2012).

[00208] 7. Mou, H. et al. Generation of Multipotent Lung and Airway Progenitors from Mouse ESCs and Patient-Specific Cystic Fibrosis iPSCs. Stem Cell 10, 385–397 (2012).

[00209] 8. Si-Tayeb, K. et al. Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. Hepatology 51, 297–305 (2010).

[00210] 9. D'Amour, K. A. et al. Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells. Nat Biotechnol 24, 1392–1401 (2006).

[00211] 10. Spence, J. R. et al. Directed differentiation of human pluripotent stem cells into intestinal tissue *in vitro*. Nature 470, 105–109 (2011).

[00212] 11. D'Amour, K. A. et al. Efficient differentiation of human embryonic stem cells to definitive endoderm. Nat Biotechnol 23, 1534–1541 (2005).

[00213] 12. McCracken, K. W., Howell, J. C., Spence, J. R. & Wells, J. M. Generating human intestinal tissue from pluripotent stem cells *in vitro*. Nature Protocols 6, 1920–1928 (2011).

[00214] 13. Kumar, M., Jordan, N., Melton, D. & Grapin-Botton, A. Signals from lateral plate mesoderm instruct endoderm toward a pancreatic fate. Dev Biol 259, 109–122 (2003).

[00215] 14. Tiso, N., Filippi, A., Pauls, S., Bortolussi, M. & Argenton, F. BMP signalling regulates anteroposterior endoderm patterning in zebrafish. Mech Dev 118, 29–37 (2002).

[00216] 15. Wang, Z., Dollé, P., Cardoso, W. V. & Niederreither, K. Retinoic acid regulates morphogenesis and patterning of posterior foregut derivatives. Dev Biol 297, 433–445 (2006).

[00217] 16. Martín, M. et al. Dorsal pancreas agenesis in retinoic acid-deficient Raldh2 mutant mice. Dev Biol 284, 399–411 (2005).

[00218] 17. Molotkov, A., Molotkova, N. & Duester, G. Retinoic acid generated by Raldh2 in mesoderm is required for mouse dorsal endodermal pancreas development. Dev Dyn 232, 950–957 (2005).

[00219] 18. Kawaguchi, Y. et al. The role of the transcriptional regulator Ptf1a in converting intestinal to pancreatic progenitors. Nat Genet 32, 128–134 (2002).

[00220] 19. Johnson, L. R. & Guthrie, P. D. Stimulation of rat oxyntic gland mucosal growth by epidermal growth factor. Am. J. Physiol. 238, G45–9 (1980).

[00221] 20. Majumdar, A. P. Postnatal undernutrition: effect of epidermal growth factor on growth and function of the gastrointestinal tract in rats. J. Pediatr. Gastroenterol. Nutr. 3, 618–625 (1984).

[00222] 21. Spear, P. C. & Erickson, C. A. Interkinetic nuclear migration: A mysterious process in search of a function. Develop. Growth Differ. 54, 306–316 (2012).

[00223] 22. Grosse, A. S. et al. Cell dynamics in fetal intestinal epithelium: implications for intestinal growth and morphogenesis. Development 138, 4423–4432 (2011).

[00224] 23. Verzi, M. P. et al. Role of the homeodomain transcription factor Bapx1 in mouse distal stomach development. Gastroenterology 136, 1701–1710 (2009).

[00225] 24. Mills, J. C. & Shivdasani, R. A. Gastric Epithelial Stem Cells. Gastroenterology 140, 412–424 (2011).

[00226] 25. Gradwohl, G., Dierich, A., LeMeur, M. & Guillemot, F. neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. Proc Natl Acad Sci USA 97, 1607–1611 (2000).

[00227] 26. Jenny, M. et al. Neurogenin3 is differentially required for endocrine cell fate specification in the intestinal and gastric epithelium. EMBO J 21, 6338–6347 (2002).

[00228] 27. Johansson, K. A. et al. Temporal control of neurogenin3 activity in pancreas progenitors reveals competence windows for the generation of different endocrine cell types. Dev Cell 12, 457–465 (2007).

[00229] 28. López-Díaz, L. et al. Intestinal Neurogenin 3 directs differentiation of a bipotential secretory progenitor to endocrine cell rather than goblet cell fate. Dev Biol 309, 298–305 (2007).

[00230] 29. Schonhoff, S. E., Giel-Moloney, M. & Leiter, A. B. Neurogenin 3-expressing progenitor cells in the gastrointestinal tract differentiate into both endocrine and non-endocrine cell types. Dev Biol 270, 443–454 (2004).

[00231] 30. Lee, C. S., Perreault, N., Brestelli, J. E. & Kaestner, K. H. Neurogenin 3 is essential for the proper specification of gastric enteroendocrine cells and the maintenance of gastric epithelial cell identity. Genes Dev 16, 1488–1497 (2002).

[00232] 31. Olbe, L., Hamlet, A., Dalenbäck, J. & Fändriks, L. A mechanism by which Helicobacter pylori infection of the antrum contributes to the development of duodenal ulcer. Gastroenterology 110, 1386–1394 (2001).

[00233] 32. Xia, H. H. et al. Antral-type mucosa in the gastric incisura, body, and fundus (antralization): a link between Helicobacter pylori infection and intestinal metaplasia? Am. J. Gastroenterol. 95, 114–121 (2000).

[00234] 33. Churin, Y. et al. Helicobacter pylori CagA protein targets the c-Met receptor and enhances the motogenic response. J. Cell Biol. 161, 249–255 (2003).

[00235] 34. Peek, R. M. et al. Helicobacter pylori cagA+ strains and dissociation of gastric epithelial cell proliferation from apoptosis. J. Natl. Cancer Inst. 89, 863–868 (1997).

[00236] 35. Teo, A. K. K. et al. Activin and BMP4 Synergistically Promote Formation of Definitive Endoderm in Human Embryonic Stem Cells. Stem Cells 30, 631–642 (2012).

[00237] 36. Meerbrey, K. L. et al. The pINDUCER lentiviral toolkit for inducible RNA interference in vitro and in vivo. Proc Natl Acad Sci USA 108, 3665–3670 (2011).

[00238] 37. Okita, K. et al. An efficient nonviral method to generate integration-free human-induced pluripotent stem cells from cord blood and peripheral blood cells. Stem Cells 31, 458–466 (2013).

[00239] 38. Covacci, A. et al. Molecular characterization of the 128-kDa immunodominant antigen of Helicobacter pylori associated with cytotoxicity and duodenal ulcer. Proc Natl Acad Sci USA 90, 5791–5795 (1993).

[00240] 39. Amieva, M. R., Salama, N. R., Tompkins, L. S. & Falkow, S. Helicobacter pylori enter and survive within multivesicular vacuoles of epithelial cells. Cell. Microbiol. 4, 677–690 (2002).

[00241] 40. Schumacher, M. A. et al. Gastric Sonic Hedgehog acts as a macrophage chemoattractant during the immune response to Helicobacter pylori. Gastroenterology 142, 1150–1159.e6 (2012).

CLAIMS

1. A method of inducing formation of a gastric tissue, comprising the steps of:

a) activating a combination of a WNT signaling pathway and a FGF signaling pathway within a definitive endoderm derived from an induced pluripotent stem cell, an embryonic stem cell, and/or an endoderm cell, wherein said WNT signalling pathway is activated using an agent selected from one or more molecules selected from the group consisting of Wnt1, Wnt2, Wnt2b, Wnt4, Wnt5a, Wnt5b, Wnt6, Wnt7a, Wnt7b, Wnt8a, Wnt9a, Wnt10a, Wnt10b, Wnt11, Wnt16, a GSK3β inhibitor such as Chiron, and small molecule modulators that activate the WNT signalling pathway, and wherein said FGF signalling pathway is activated using an agent selected from one or more molecules selected from the group consisting of FGF1, FGF2, FGF3, FGF4, FGF10, FGF11, FGF12, FGF13, FGF14, FGF15, FGF16, FGF17, FGF18, FGF19, FGF20, FGF21, FGF22, FGF23 and small molecule modulators that activate the FGF signalling pathway,

b) inhibiting a BMP signaling pathway within said definitive endoderm derived from an induced pluripotent stem cell, embryonic stem cell, and/or endoderm cell by contacting said definitive endoderm derived from an induced pluripotent stem cell, embryonic stem cell, and/or endoderm cell with an agent that inhibits the BMP pathway, wherein said agent is selected from Noggin, Dorsomorphin, LDN189, DMH-1, or small molecule modulators that inhibit the BMP signaling pathway; and

c) administering retinoic acid to said definitive endoderm;

wherein said steps a) through c) are carried out for a period of time sufficient to cause formation of a foregut spheroid, further comprising the step of contacting said foregut spheroid with EGF, RA, and a BMP inhibitor.

2. The method of claim 1, wherein said definitive endoderm is derived from contacting a pluripotent stem cell with one or more molecules selected from Activin, the BMP subgroups of the TGF-beta superfamily of growth factors; Nodal, Activin A, Activin B, BMP4, Wnt3a, and combinations thereof.

3. The method of claim 1 or 2, wherein said small molecule modulator that activate the WNT signalling pathway is selected from the group consisting of : Lithium Chloride; 2-amino-4,6-disubstituted pyrimidine (hetero) arylpyrimidines; IQ1; QS11; NSC668036; DCA beta-catenin; and 2-amino-4-[3,4-(methylenedioxy)-benzyl-amino]-6-(3-methoxyphenyl) pyrimidine.

4. The method of any one of claims 1 to 3, wherein said FGF signalling pathway is activated using FGF4, more preferably FGF4 at a concentration between about 50 to about 1500 ng/ml.

5. The method of any one of claims 1 to 4, wherein said WNT signalling pathway is activated using Wnt3a, preferably Wnt3a at a concentration between about 50 to about 1500 ng/ml.

6. The method of any one of claims 1 to 5, wherein said activating step comprises contacting said definitive endoderm with both Wnt3a and FGF4 or CHIR99021 and FGF4 over a specified activation period.

7. The method of any one of claims 1 to 6, wherein said definitive endoderm is contacted with a BMP inhibitor at a concentration between about 50 to about 1500 ng/ml.

8. The method of any one of claims 1 to 7, wherein the retinoic acid of step c) is administered in an amount of between about 0.2 uM to about 20 uM, preferably about 2.0 uM, preferably wherein step c) is carried out until expression of Pdx1/Sox1 is detected in said foregut spheroids.

9. The method of any one of claims 1 to 8, wherein said exposure to EGF is at a concentration and length of time sufficient to increase said gastric spheroids to greater than 1 mm in diameter, or greater than 2 mm in diameter, or greater than 3 mm in diameter, or greater than 4 mm in diameter.

10. The method of any one of claims 1 to 9, further comprising the step of culturing said activated definitive endoderm *in vitro* to form a 3-dimensional tissue structure.

11. The method of any one of claims 1 to 10, wherein said activating and inhibiting steps are conducted *in vitro*.

12. The method of any one of claims 1 to 11, wherein said activation period and said inhibition period overlap.

13. The method of any one of claims 1 to 11, wherein said activation period and said inhibition period do not overlap.

14. The method of any one of claims 1 to 13, wherein said activation period is between about 24 and about 120 hours.

15. A method of generating a gastric organoid comprising fundus tissue comprising the steps of

a) generating a definitive endoderm (DE) from a pluripotent stem cell;

b) contacting said definitive endoderm with an agent that activates the FGF pathway as defined in claim 1, an agent that activates the WNT pathway as defined in claim 1 or claim 3, an agent that inhibits the BMP pathway as defined in claim 1, and retinoic acid to generate a posterior foregut; and

c) contacting said posterior foregut with EGF, retinoic acid, an agent that activates the WNT pathway as defined in claim 1 or claim 3, and an agent that inhibits the BMP pathway as defined in claim 1, to generate a fundus tissue.

16. A method of generating a gastric organoid comprising antrum tissue comprising the steps of:

a) generating a definitive endoderm (DE) from a pluripotent stem cell;

b) contacting said definitive endoderm with an agent that activates the FGF pathway as defined in claim 1, an agent that activates the WNT pathway as defined in claim 1 or claim 3, an agent that inhibits the BMP pathway as defined in claim 1, and retinoic acid to generate a posterior foregut; and

c) contacting said posterior foregut with EGF, retinoic acid, and an agent that inhibits the BMP pathway as defined in claim 1, to generate an antrum tissue.