RNA PREPARATIONS COMPRISING PURIFIED MODIFIED RNA FOR REPROGRAMMING CELLS

The present application claims priority to U.S. Provisional Application Serial Number 61/267,312 filed December 7, 2009, which is herein incorporated by reference in its entirety.

The present application also claims priority to U.S. Application Serial Number 11/990,646, filed March 27, 2009, which is a U.S. National Entry of PCT/US06/32372 filed August 21, 2006, which claims priority to U.S. Provisional Application 60/710,164, filed August 23, 2005, all of which are herein incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to compositions and methods for changing or reprogramming the state of differentiation of eukaryotic cells, including human or other animal cells, by contacting the cells with purified RNA preparations comprising or consisting of one or more different single-strand mRNA molecules that each encode a reprogramming factor (e.g., an iPS cell induction factor). The purified single-stranded mRNA molecules preferably comprise at least one modified nucleoside (e.g., selected from the group consisting of a pseudouridine (abbreviated by the Greek letter "psi" or "Ψ"), 5-methylcytosine (m^5C), 5-methyluridine (m^5U), 2'-O-methyluridine (Um or $m^{2'\text{-}O}U$), 2-thiouridine (s^2U), and N^6-methyladenosine (m^6A)) in place of at least a portion of the corresponding unmodified canonical nucleoside (e.g., in place of substantially all of the corresponding unmodified A, C, G, or T canonical nucleoside). In addition, the single-stranded mRNA molecules are preferably purified to be substantially free of RNA contaminant molecules that would activate an unintended response, decrease expression of the single-stranded mRNA, and/or activate RNA sensors in the cells. In certain embodiments, the purified RNA preparations are substantially free of RNA contaminant molecules that are: shorter or longer than the full-length single-stranded mRNA molecules, double-stranded, and/or uncapped RNA.

BACKGROUND

In 2006, it was reported (Takahashi and Yamanaka 2006) that the introduction of genes encoding four protein factors (OCT4 (Octamer-4; POU class 5 homeobox 1), SOX2 (SRY (sex

determining region Y)-box 2), KLF4 (Krueppel-like factor 4), and c-MYC) into differentiated mouse somatic cells induced those cells to become pluripotent stem cells, (referred to herein as "induced pluripotent stem cells," "iPS cells," or "iPSCs"). Following this original report, pluripotent stem cells were also induced by transforming human somatic cells with genes encoding the similar human protein factors (OCT4, SOX2, KLF4, and c-MYC) (Takahashi et al. 2007), or by transforming human somatic cells with genes encoding human OCT4 and SOX2 factors plus genes encoding two other human factors, NANOG and LIN28 (Lin-28 homolog A) (Yu et al. 2007). All of these methods used retroviruses or lentiviruses to integrate genes encoding the reprogramming factors into the genomes of the transformed cells and the somatic cells were reprogrammed into iPS cells only over a long period of time (e.g., in excess of a week).

The generation iPS cells from differentiated somatic cells offers great promise as a possible means for treating diseases through cell transplantation. The possibility to generate iPS cells from somatic cells from individual patients also may enable development of patient-specific therapies with less risk due to immune rejection. Still further, generation of iPS cells from disease-specific somatic cells offers promise as a means to study and develop drugs to treat specific disease states (Ebert et al. 2009, Lee et al. 2009, Maehr et al. 2009).

Viral delivery of genes encoding protein reprogramming factors (or "iPSC factors") provides a highly efficient way to make iPS cells from somatic cells, but the integration of exogenous DNA into the genome, whether random or non-random, creates unpredictable outcomes and can ultimately lead to cancer (Nakagawa et al. 2008). New reports show that iPS cells can be created (at lower efficiency) by using other methods that do not require genome integration. For example, repeated transfections of expression plasmids containing genes for OCT4, SOX2, KLF4 and c-MYC into mouse embryonic fibroblasts to generate iPS cells was demonstrated (Okita et al. 2008). Induced pluripotent stem cells were also generated from human somatic cells by introduction of a plasmid that expressed genes encoding human OCT4, SOX2, c-MYC, KLF4, NANOG and LIN28 (Yu et al. 2009). Other successful approaches for generating iPS cells include treating somatic cells with: recombinant protein reprogramming factors (Zhou et al. 2009); non-integrating adenoviruses (Stadtfeld et al. 2008); or piggyBac transposons (Woltjen et al. 2009) to deliver reprogramming factors. Presently, the generation of iPS cells using these non-viral delivery techniques to deliver reprogramming factors is extremely

inefficient. Future methods for generating iPS cells for potential clinical applications will need to increase the speed and efficiency of iPS cell formation while maintaining genome integrity.

SUMMARY OF THE INVENTION

The present invention provides compositions and methods for reprogramming the state of differentiation of eukaryotic cells, including human or other animal cells, by contacting the cells with purified RNA preparations comprising or consisting of one or more different single-strand mRNA molecules that each encode a reprogramming factor (e.g., an iPS cell induction factor). The purified single-stranded mRNA molecules preferably comprise at least one modified nucleoside (e.g., selected from the group consisting of a pseudouridine (Ψ), 5-methylcytosine (m^5C), 5-methyluridine (m^5U), 2'-O-methyluridine (Um or $m^{2'\text{-}O}U$), 2-thiouridine (s^2U), and N^6-methyladenosine (m^6A)) in place of at least a portion (e.g., including substantially all) of the corresponding unmodified canonical nucleoside of the corresponding unmodified A, C, G, or T canonical nucleoside. In addition, the single-stranded mRNA molecules are preferably purified to be substantially free of RNA contaminant molecules that would activate an unintended response, decrease expression of the single-stranded mRNA, and/or activate RNA sensors (e.g., double-stranded RNA-dependent enzymes) in the cells. In certain embodiments, the purified RNA preparations are substantially free of RNA contaminant molecules that are: shorter or longer than the full-length single-stranded mRNA molecules, double-stranded, and/or uncapped RNA. In some preferred embodiments, the invention provides compositions and methods for reprogramming differentiated eukaryotic cells, including human or other animal somatic cells, by contacting the cells with purified RNA preparations comprising or consisting of one or more different single-strand mRNA molecules that each encode an iPS cell induction factor.

In some embodiments, the present invention provides methods for changing the state of differentiation of a somatic cell comprising: introducing an mRNA encoding an iPS cell induction factor into a somatic cell to generate a reprogrammed dedifferentiated cell, wherein the mRNA comprises at least one 5-methylcytidine (or other modified based described herein).

In certain embodiments, the present invention provides methods for reprogramming a cell that exhibits a first differentiated state or phenotype to a cell that exhibits a second differentiated state or phenotype comprising: introducing into the cell that exhibits a first differentiated state a purified RNA preparation comprising modified mRNA molecules that encode at least one

reprogramming factor and culturing the cell under conditions wherein the cell exhibits a second differentiated state. In certain embodiments, the modified mRNA molecules contain at least one modified nucleoside selected from the group consisting of psuedouridine or 5-methylcytidine. In certain embodiments, the cell is from a human or animal. In further embodiments, the purified RNA preparation: i) comprises first single-stranded mRNAs encoding a first iPS cell induction factor, wherein substantially all of the first single-stranded complete mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of RNA contaminant molecules which are able to activate RNA sensors in said somatic cell. In certain embodiments, the RNA contaminant molecules are selected from the group consisting of: partial mRNAs encoding only a portion of said iPS cell induction factor, RNA molecules that are smaller than the full-length mRNA, RNA molecules that are larger than the full-length mRNA, double-stranded mRNA molecules, and un-capped mRNA molecules.

In some embodiments, the present invention provides methods for reprogramming a somatic cell (e.g., dedifferentiating or transdifferentiating) comprising: contacting a somatic cell with a purified RNA preparation to generate a reprogrammed cell, wherein the purified RNA preparation: i) comprises first single-stranded mRNAs encoding a first iPS cell induction factor, wherein substantially all of the first single-stranded complete mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of contaminant molecules (e.g., RNA contaminant molecules) which are able to activate RNA sensors in the somatic cell. In particular embodiments, the RNA contaminant molecules comprise: partial mRNAs encoding only a portion of the iPS cell induction factor, single-stranded run-on mRNAs encoding the iPS cell induction factor and encoding at least an additional portion of the iPS cell induction factor, double-stranded mRNA molecules, and un-capped mRNA molecules. In certain embodiments, the first single-stranded mRNAs do not also encoding an additional portion of the first iPS cell induction factor.

In some embodiments, the reprogrammed cell is a dedifferentiated cell (e.g., stem cell or stem cell-like cell). In other embodiments, the reprogrammed cell is a transdifferentiated cells (e.g., a skin cells is reprogrammed into a neuronal cell, or other type of change). In further embodiments, the first single-stranded mRNAs encode the complete first iPS induction factor (e.g, the mRNA encodes the entire coding sequence for a particular iPS induction factor). In other embodiments, the contacting further comprises contacting the somatic cell with a growth

factor and/or cytokine (e.g,. after a period of time). In further embodiments, the contact further comprises contacting the somatic cell with an immune response inhibitor.

In certain embodiments, all or nearlly all of the uridine nucleosides in the first single-stranded mRNA are replaced by pseudouridine nucleosides. In other embodiments, all or nearly all of the cytidine nucleosides in the first single-stranded mRNA are replaced by 5-methylcytidine nucleosides or another base recited herein.

In particular embodiments, the present invention provides methods for generating a reprogrammed cell comprising: contacting a somatic cell with a purified RNA preparation to generate a reprogrammed cell that is able to survive in culture for at least 10 days (e.g., at least 10 days ... at least 13 days ... at least 16 days .... at least 20 days ... at least 40 days ... or is able to form a cell-line), wherein the purified RNA preparation comprises first single-stranded mRNAs encoding an iPS cell induction factor, and wherein a majority of the first single-stranded mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue.

In certain embodiments, the purified RNA preparation is free of an amount of RNA contaminant molecules that would activate an immune response in the somatic cell sufficient to prevent the reprogrammed cell from surviving at least 10 days in culture (e.g., at least 10 days ... at least 15 days ... at least 20 days ... at least 40 days, or longer). In other embodiments, the RNA contaminant molecules include: partial mRNAs encoding only a portion of the iPS cell induction factor, single-stranded run-on mRNAs fully encoding the iPS cell induction factor and encoding at least an additional portion of the iPS cell induction factor, double-stranded mRNA molecules, un-capped mRNA molecules, and mixtures thereof. In certain embodiments, the reprogrammed cell that is generated is able to form a reprogrammed cell-line. In other embodiments, the purified RNA preparation is free of an amount of RNA contaminant molecules that would activate an immune response in the somatic cell sufficient to prevent generation of the reprogrammed cell-line.

In particular embodiments, the RNA contaminant molecules are selected from the group consisting of: partial mRNAs encoding only a portion of the iPS cell induction factor, single-stranded run-on mRNAs encoding the iPS cell induction factor and encoding at least an additional portion of the iPS cell induction factor, double-stranded mRNA molecules, un-capped mRNA molecules, and mixtures thereof.

In some embodiments, the present invention provides methods for generating a reprogrammed cell-line comprising: a) contacting a somatic cell with a purified RNA preparation to generate a reprogrammed cell, wherein the purified RNA preparation comprises mRNAs encoding an iPS cell induction factor, and wherein a majority of the mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and b) culturing the dedifferentiated cell to generate a reprogrammed cell-line. In other embodiments, the purified RNA preparation is free of an amount of contaminant molecules that would activate an immune response in the somatic cell sufficient to prevent generation of the reprogrammed cell-line. In certain embodiments, the immune response involves activation of RNA sensors in the somatic cell.

In some embodiments, the present invention provides methods for reprogramming a somatic cell comprising: contacting a somatic cell with a purified RNA preparation to generate a reprogrammed cell, wherein the purified RNA preparation: i) comprises first single-stranded mRNAs encoding a first iPS cell induction factor, wherein substantially all of the first single-stranded mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of: a) partial mRNAs encoding only a portion of the first iPS cell induction factor, and b) double-stranded mRNA molecules. In further embodiments, the first single-stranded mRNA do not also encode an additional portion of the first iPS cell induction factor. In particular embodiments, the first single-stranded mRNA fully encode the first iPS cell induction factor. In other embodiments, the purified RNA preparation is also substantially free (or essentially free or virtually free or free) of single-stranded run-on mRNAs encoding the first iPS cell induction factor and encoding at least an additional portion of the first iPS cell induction factor. In other embodiments, the substantially all of the first single-stranded complete mRNAs are 5' capped. In other embodiments, the purified RNA preparation is also substantially free of un-capped mRNA molecules. In some embodiments, substantially all of the first single-stranded mRNAs comprise at least one psuedouridine residue. In additional embodiments, tsubstantially all of the first single-stranded mRNAs comprise at least one 5-methylcytidine residue. In other embodiments, substantially all of the first single-stranded mRNAs comprise at least one psuedouridine residue and at least one 5-methycytidine residue.

In certain embodiments, the purified RNA preparation comprises a transfection reagent. In other embodiments, the purified RNA preparation is obtained by HPLC purification of an

RNA sample that contains a substantial amount of the partial mRNAs and the double-stranded mRNAs. In further embodiments, the purified RNA preparation is essentially free of the partial mRNAs and the single-stranded run-on mRNAs. In some embodiments, the purified RNA preparation is essentially free or virtually free or free of double-stranded mRNA molecules. In other embodiments, the purified RNA preparation is essentially free or virtually free or free of un-capped mRNA molecules. In some embodiments, substantially all of the first single-stranded mRNAs are polyadenylated. In other embodiments, the first single-stranded complete mRNAs are capped with 7-methylguanosine.

In some embodiments, the first iPS cell induction factor is selected from the group consisting of KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2. In other embodiments, the purified RNA preparation: i) further comprises second single-stranded mRNAs encoding a second iPS cell induction factor, wherein the second single-stranded mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is further substantially free of: a) partial mRNAs encoding only a portion of the second iPS cell induction factor, and b) double-stranded mRNAs. In other embodiments, the purified RNA preparation is further substantially free of single-stranded run-on mRNAs encoding a second iPS cell induction factor and encoding at least an additional portion of the second iPS cell induction factor. In some embodiments, the second iPS cell induction factor is selected from the group consisting of KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2. In certain embodiments, the somatic cell is a fibroblast. In other embodiments, the reprogrammed cell is a pluripotent stem cell. In other embodiments, the dedifferentiated cell expresses NANOG and TRA-1-60. In some embodiments, the cell is in vitro. In further embodiments, the cell resides in culture. In particular embodiments, the cell resides in MEF-conditioned medium.

In some embodiments, the present invention provides compositions comprising a purified RNA preparation, wherein the purified RNA preparation: i) comprises first single-stranded mRNAs encoding a first iPS cell induction factor, wherein the first single-stranded mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of RNA contaminant molecules, which are able to activate RNA sensors in a somatic cell. In certain embodiments, the present invention provides compositions comprising a purified RNA preparation, wherein the purified RNA preparation: i) comprises first single-stranded mRNAs encoding a first iPS cell induction factor, wherein the first single-stranded

complete mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of: a) partial mRNAs encoding only a portion of the first iPS cell induction factor, and b) double-stranded RNA.

In certain embodiments, the purified RNA preparation is also substantially free of single-stranded run-on mRNAs encoding the first iPS cell induction factor and encoding at least an additional portion of the first iPS cell induction factor. In some embodiments, the purified RNA preparation: i) further comprises second single-stranded mRNAs encoding a second iPS cell induction factor, wherein the second single-stranded complete mRNAs comprise at least one pseudouridine residue and/or at least one 5-methylcytidine residue, and ii) is substantially free of: a) partial mRNAs encoding only a portion of the second iPS cell induction factor, and b) single-stranded run-on mRNAs encoding second first iPS cell induction factor and encoding at least an additional portion of the second iPS cell induction factor.

In some embodiments, the present invention provides compositions comprising an in vitro-synthesized mRNA encoding the MYC gene, wherein the in vitro-synthesized mRNA comprises at least one pseudouridine residue and/or at least one 5-methylcytidine residue. In certain embodiments, the compositions are substantially free of RNA contaminant molecules which are able to activate RNA sensors in a somatic cell.

In particular embodiments, the present invention provides methods for inducing a mammalian cell to produce the MYC protein comprising: contacting a mammalian cell with an in vitro-synthesized mRNA encoding the MYC gene, wherein the in vitro-synthesized mRNA comprises at least one pseudouridine residue and/or at least one 5-methylcytidine residue, thereby inducing the mammalian cell to produce the MYC protein. In other embodiments, the mammalian cell is a dendritic cell. In other embodiments, the mammalian cell is an alveolar cell, an astrocyte, a microglial cell, or a neuron.

In some embodiments, the present invention provides methods of treating a subject comprising contacting a subject with the MYC protein producing mammalian cell described above and herein.

In additional embodiments, the present invention provides methods of synthesizing an in vitro-transcribed RNA molecule encoding the MYC gene comprising: combining an isolated RNA polymerase, a tempalte nucleic acid sequence encoding the MYG gene, unmodified nucleotides, and pseudouridine or 5-methylcytidine modified nucleotides under conditions such

that an in vitro-transcribed RNA molecule encoding the MYC gene is generated that comprises at least one pseudouridine or 5-methylcytidine residue.

Experiments conducted during the development of embodiments of the present invention demonstrated that mRNA molecules can be administered to cells and induce a dedifferentiation process to generate dedifferentiated cells—including pluripotent stem cells. Thus, the present invention provides compositions and methods for generating iPS cells. Surprisingly, the administration of mRNA can provide highly efficient generation of iPS cells.

The present invention also provides RNA, oligoribonucleotide, and polyribonucleotide molecules comprising pseudouridine or a modified nucleoside, gene therapy vectors comprising same, methods of synthesizing same, and methods for gene replacement, gene therapy, gene transcription silencing, and the delivery of therapeutic proteins to tissue *in vivo,* comprising the molecules. The present invention also provides methods of reducing the immunogenicity of RNA, oligoribonucleotide, and polyribonucleotide molecules.

In some embodiments, the present invention provides methods for dedifferentiating a somatic cell comprising: introducing mRNA encoding one or more iPSC induction factors into a somatic cell to generate a dedifferentiated cell.

In some embodiments, the present invention provides methods for dedifferentiating a somatic cell comprising: introducing mRNA encoding one or more iPSC induction factors into a somatic cell and maintaining the cell under conditions wherein the cell is viable and the mRNA that is introduced into the cell is translated in sufficient amount and for sufficient time to generate a dedifferentiated cell. In some preferred embodiments, the dedifferentiated cell is an induced pluripotent stem cell (iPSC).

In some embodiments, the present invention provides methods for changing the state of differentiation (or differentiated state) of a eukaryotic cell comprising: introducing mRNA encoding one or more reprogramming factors into a cell and maintaining the cell under conditions wherein the cell is viable and the mRNA that is introduced into the cell is translated in sufficient amount and for sufficient time to generate a cell that exhibits a changed state of differentiation compared to the cell into which the mRNA was introduced.

In some embodiments, the present invention provides methods for changing the state of differentiation of a eukaryotic cell comprising: introducing mRNA encoding one or more reprogramming factors into a cell and maintaining the cell under conditions wherein the cell is

viable and the mRNA that is introduced into the cell is translated in sufficient amount and for sufficient time to generate a cell that exhibits a changed state of differentiation compared to the cell into which the mRNA was introduced. In some embodiments, the changed state of differentiation is a dedifferentiated state of differentiation compared to the cell into which the mRNA was introduced. For example, in some embodiments, the cell that exhibits the changed state of differentiation is a pluripotent stem cell that is dedifferentiated compared to a somatic cell into which the mRNA was introduced (e.g., a somatic cell that is differentiated into a fibroblast, a cardiomyocyte, or another differentiated cell type). In some embodiments, the cell into which the mRNA is introduced is a somatic cell of one lineage, phenotype, or function, and the cell that exhibits the changed state of differentiation is a somatic cell that exhibits a lineage, phenotype, or function that is different than that of the cell into which the mRNA was introduced; thus, in these embodiments, the method results in transdifferentiation (Graf and Enver 2009).

The methods of the invention are not limited with respect to a particular cell into which the mRNA is introduced. In some embodiments of any of the above methods, the cell into which the mRNA is introduced is derived from any multi-cellular eukaryote. In some embodiments of any of the above methods, the cell into which the mRNA is introduced is selected from among a human cell and another animal cell. Although the work presented herein was performed using cells of humans or other animals, the applicants further claim that the methods of the present invention comprising reprogramming human and animal cells by contacting the cells with a purified RNA preparation that consists of one or more purified single-stranded mRNA molecules, each of which encodes a protein reprogramming factor (e.g., a transcription factor) also pertains to reprogramming of other eukaryotic cells (e.g., plant cells and a fungal cells). In some embodiments of any of the above methods, the cell into which the mRNA is introduced is a normal cell that is from an organism that is free of a known disease. In some embodiments of any of the above methods, the cell into which the mRNA is introduced is a cell from an organism that has a known disease. In some embodiments of any of the above methods, the cell into which the mRNA is introduced is a cell that is free of a known pathology. In some embodiments of any of the above methods, the cell into which the mRNA is introduced is a cell that exhibits a disease state or a known pathology (e.g., a cancer cell, or a pancreatic beta cell that exhibits metabolic properties characteristic of a diabetic cell).

The invention is not limited to the use of a specific cell type (e.g., to a specific somatic cell type) in embodiments of the methods comprising introducing mRNA encoding one or more iPSC cell induction factors in order to generate a dedifferentiated cell (e.g., an iPS cell). Any cell that is subject to dedifferentiation using iPS cell induction factors is contemplated. Such cells include, but are not limited to, fibroblasts, keratinocytes, adipocytes, lymphocytes, T-cells, B-Cells, cells in mononuclear cord blood, buccal mucosa cells, hepatic cells, HeLa, MCF-7 or other cancer cells. In some embodiments, the cells reside *in vitro* (e.g., in culture) or *in vivo*. In some embodiments, when generated in culture, a cell-free conditioned medium (e.g., MEF-conditioned medium) is used. As demonstrated below, such a medium provided enhanced iPS cell generation. The invention is not limited, however, to the culturing conditions used. Any culturing condition or medium now known or later identified as useful for the methods of the invention (e.g., to generate iPS cells from somatic cells and maintain said cells) is contemplated for use with the invention. For example, although not preferred, in some embodiments of the method, a feeder cell layer is used instead of conditioned medium for culturing the cells that are treated using the method.

In some embodiments of any of these methods, the step of introducing mRNA comprises delivering the mRNA into the cell (e.g., a human or other animal somatic cell) with a transfection reagent (e.g., TRANSIT™ mRNA transfection reagent, MirusBio, Madison, WI).

However, the invention is not limited by the nature of the transfection method utilized. Indeed, any transfection process known, or identified in the future that is able to deliver mRNA molecules into cells *in vitro* or *in vivo*, is contemplated, including methods that deliver the mRNA into cells in culture or in a life-supporting medium, whether said cells comprise isolated cells or cells comprising a eukaryotic tissue or organ, or methods that deliver the mRNA *in vivo* into cells in an organism, such as a human, animal, plant or fungus. In some embodiments, the transfection reagent comprises a lipid (e.g., liposomes, micelles, etc.). In some embodiments, the transfection reagent comprises a nanoparticle or nanotube. In some embodiments, the transfection reagent comprises a cationic compound (e.g., polyethylene imine or PEI). In some embodiments, the transfection method uses an electric current to deliver the mRNA into the cell (e.g., by electroporation). In some embodiments, the transfection method uses a bolistics method to deliver the mRNA into the cell (e.g., a "gene gun" or biolistic particle delivery system.)

The data presented herein shows that, with respect to the mRNA introduced into the cell, certain amounts of the mRNAs used in the EXAMPLES described herein resulted in higher efficiency and more rapid induction of pluripotent stem cells from the particular somatic cells used than other amounts of mRNA. However, the methods of the present invention are not limited to the use of a specific amount of mRNA to introduce into the cell. For example, in some embodiments, a total of three doses, with each dose comprising 18 micrograms of each of six different mRNAs, each encoding a different human reprogramming factor, was used to introduce the mRNA into approximately 3 x 10^5 human fibroblast cells in a 10-cm plate (e.g., delivered using a lipid-containing transfection reagent), although in other embodiments, higher or lower amounts of the mRNAs were used to introduce into the cells.

The invention is not limited to a particular chemical form of the mRNA used, although certain forms of mRNA may produce more efficient results. However, in some preferred embodiments, the mRNA comprises at least one modified nucleoside (e.g., selected from the group consisting of a pseudouridine (Ψ), 5-methylcytosine (m^5C), 5-methyluridine (m^5U), 2'-O-methyluridine (Um or $m^{2'\text{-}O}U$), 2-thiouridine (s^2U), and N^6-methyladenosine (m^6A)) in place of at least a portion of the corresponding unmodified canonical nucleoside (e.g., in some preferred embodiments, at least one modified nucleoside in place of substantially all of the corresponding unmodified A, C, G, or T canonical nucleoside). In some embodiments, the mRNA is polyadenylated. In some preferred embodiments, the mRNA is prepared by polyadenylation of an in vitro-transcribed (IVT) RNA, the method comprising contacting the IVT RNA using a poly(A) polymerase (e.g., yeast RNA polymerase or *E. coli* poly(A) polymerase). In some embodiments, the mRNA is polyadenylated during IVT by using a DNA template that encodes the poly(A) tail. Regardless of whether the RNA is polyadenylated using a poly(A) polymerase or during IVT of a DNA template, in some preferred embodiments, the mRNA comprises a poly-A tail (e.g., a poly-A tail having 50-200 nucleotides, e.g., preferably 100-200, 150-200 nucleotides, or greater than 150 nucleotides), although in some embodiments, a longer or a shorter poly-A tail is used. In some embodiments, the mRNA used in the methods is capped. To maximize efficiency of expression in the cells, it is preferred that the majority of mRNA molecules contain a cap. In some preferred embodiments, the mRNA molecules used in the methods are synthesized *in vitro* by incubating uncapped primary RNA in the presence a capping enzyme system. In some preferred embodiments, the primary RNA used in the capping enzyme

reaction is synthesized by *in vitro* transcription (IVT) of a DNA molecule that encodes the RNA to be synthesized. The DNA that encodes the RNA to be synthesized contains an RNA polymerase promoter, to which, an RNA polymerase binds and initiates transcription therefrom. It is also known in the art that mRNA molecules often have regions of differing sequence located before the translation start codon and after the translation stop codon that are not translated. These regions, termed the five prime untranslated region (5' UTR) and three prime untranslated region (3' UTR), respectively, can affect mRNA stability, mRNA localization, and translational efficiency of the mRNA to which they are joined. Certain 5' and 3' UTRs, such as those for alpha and beta globins are known to improve mRNA stability and expression of mRNAs. Thus, in some preferred embodiments, the mRNAs that encode reprogramming factors (e.g., iPSC induction factors) exhibit a 5' UTR and/or a 3' UTR that results in greater mRNA stability and higher expression of the mRNA in the cells (e.g., an alpha globin or a beta globin 5' UTR and/or 3' UTR; e.g., a Xenopus or human alpha globin or a beta globin 5' UTR and/or 3' UTR, or, e.g., a tobacco etch virus (TEV) 5' UTR).

The IVT can be performed using any RNA polymerase as long as synthesis of the mRNA from the DNA template that encodes the RNA is specifically and sufficiently initiated from a respective cognate RNA polymerase promoter and full-length mRNA is obtained. In some preferred embodiments, the RNA polymerase is selected from among T7 RNA polymerase, SP6 RNA polymerase and T3 RNA polymerase. In some other embodiments, capped RNA is synthesized co-transcriptionally by using a dinucleotide cap analog in the IVT reaction (e.g., using an AMPLICAP™ T7 Kit or a MESSAGEMAX™ T7 ARCA-CAPPED MESSAGE Transcription Kit; EPICENTRE or CellScript, Madison, WI, USA). If capping is performed co-transcriptionally, preferably the dinucleotide cap analog is an anti-reverse cap analog (ARCA). However, use of a separate IVT reaction, followed by capping with a capping enzyme system, which results in approximately 100% of the RNA being capped, is preferred over co-transcriptional capping, which typically results in only about 80% of the RNA being capped. Thus, in some preferred embodiments, a high percentage of the mRNA molecules used in a method of the present invention are capped (e.g., greater than 80%, greater than 90%, greater than 95%, greater than 98%, greater than 99%, greater than 99.5%, or greater than 99.9% of the population of mRNA molecules are capped). In some preferred embodiments, the mRNA used in the methods of the present invention has a cap with a cap1 structure, meaning that the 2'

hydroxyl of the ribose in the penultimate nucleotide with respect to the cap nucleotide is methylated. However, in some embodiments, mRNA used in the methods has a cap with a cap0 structure, meaning that the 2' hydroxyl of the ribose in the penultimate nucleotide with respect to the cap nucleotide is not methylated. With some but not all transcripts, transfection of eukaryotic cells with mRNA having a cap with a cap1 structure results in a higher level or longer duration of protein expression in the transfected cells compared to transfection of the same cells with the same mRNA but with a cap having a cap0 structure. In some embodiments, the mRNA used in the methods of the present invention has a modified cap nucleotide. In some experiments performed prior to the experiments presented in the EXAMPLES herein, the present applicants found that, when 1079 or IMR90 human fibroblast cells were transfected with OCT4 mRNA that contained either uridine, or pseudouridine in place of uridine, the pseudouridine-containing mRNA was translated at a higher level or for a longer duration than the mRNA that contained uridine. Therefore, in some preferred embodiments, one or more or all of the uridines contained in the mRNA(s) used in the methods of the present invention is/are replaced by pseudouridine (e.g., by substituting pseudouridine-5'-triphosphate in the IVT reaction to synthesize the RNA in place of uridine-5'-triphosphate). However, in some embodiments, the mRNA used in the methods of the invention contains uridine and does not contain pseudouridine. In some preferred embodiments, the mRNA comprises at least one modified nucleoside (e.g., selected from the group consisting of a pseudouridine (Ψ), 5-methylcytosine (m^5C), 5-methyluridine (m^5U), 2'-O-methyluridine (Um or $m^{2'-O}U$), 2-thiouridine (s^2U), and N^6-methyladenosine (m^6A)) in place of at least a portion of the corresponding unmodified canonical nucleoside (e.g., in place of substantially all of the corresponding unmodified A, C, G, or T canonical nucleoside). In some preferred embodiments, the mRNA comprises at least one modified nucleoside selected from the group consisting of a pseudouridine (Ψ) and 5-methylcytosine (m^5C). In some preferred embodiments, the mRNA comprises both pseudouridine (Ψ) and 5-methylcytosine (m^5C). In addition, in order to accomplish specific goals, a nucleic acid base, sugar moiety, or internucleotide linkage in one or more of the nucleotides of the mRNA that is introduced into a eukaryotic cell in any of the methods of the invention may comprise a modified nucleic acid base, sugar moiety, or internucleotide linkage.

The invention is also not limited with respect to the source of the mRNA that is delivered into the eukaryotic cell in any of the methods of the invention. In some embodiments, such as those described in the EXAMPLES, the mRNA is synthesized by in vitro transcription of a DNA template comprising a gene cloned in a linearized plasmid vector or by in vitro transcription of a DNA template that is synthesized by PCR or RT-PCR (i.e., by IVT of a PCR amplification product), capping using a capping enzyme system or by co-transcriptional capping by incorporation of a dinucleotide cap analog (e.g., an ARCA) during the IVT, and polyadenylation using a poly(A) polymerase. In some preferred embodiments, the mRNA is synthesized by IVT of a DNA template comprising a gene cloned in a linearized plasmid vector or a PCR or RT-PCR amplification product, wherein the DNA template encodes a 3'poly(A) tail. In some other embodiments, the mRNA that is delivered into the eukaryotic cell in any of the methods of the invention is derived directly from a cell or a biological sample. For example, in some embodiments, the mRNA derived from a cell or biological sample is obtained by amplifying the mRNA from the cell or biological sample using an RNA amplification reaction, and capping the amplified mRNA using a capping enzyme system or by co-transcriptional capping by incorporation of a dinucleotide cap analog (e.g., an ARCA) during the IVT, and, if the amplified mRNA does not already contain a template-encoded poly(A) tail from the RNA amplification reaction, polyadenylating the amplified mRNA using a poly(A) polymerase.

With respect to the methods comprising introducing mRNA encoding one or more iPS cell induction factors in order to generate a dedifferentiated cell (e.g., an iPS cell), the invention is not limited by the nature of the iPS cell induction factors used. Any mRNA encoding one or more protein induction factors now known, or later discovered, that find use in dedifferentiation, are contemplated for use in the present invention. In some embodiments, one or more mRNAs encoding for KLF4, LIN28, c-MYC, NANOG, OCT4, or SOX2 are employed. Oct-3/4 and certain members of the Sox gene family (Sox1, Sox2, Sox3, and Sox15) have been identified as transcriptional regulators involved in the induction process. Additional genes, however, including certain members of the Klf family (Klf1, Klf2, Klf4, and Klf5), the Myc family (C-myc, L-myc, and N-myc), Nanog, and LIN28, have been identified to increase the induction efficiency. Any one or more such factors may be used as desired.

While the compositions and methods of the invention may be used to generate iPS cells, the invention is not limited to the generation of such cells. For example, in some embodiments,

mRNA encoding one or more reprogramming factors is introduced into a cell in order to generate a cell with a changed state of differentiation compared to the cell into which the mRNA was introduced. For example, in some embodiments, mRNA encoding one or more iPS cell induction factors is used to generate a dedifferentiated cell that is not an iPS cells. Such cells find use in research, drug screening, and other applications.

In some embodiments, the present invention further provides methods employing the dedifferentiated cells generated by the above methods. For example, such cells find use in research, drug screening, and therapeutic applications in humans or other animals. For example, in some embodiments, the cells generated find use in the identification and characterization of iPS cell induction factors as well as other factors associated with differentiation or dedifferentiation. In some embodiments, the generated dedifferentiated cells are transplanted into an organism or into a tissue residing *in vitro* or *in vivo*. In some embodiments, an organism, tissue, or culture system housing the generated cells is exposed to a test compound and the effect of the test compound on the cells or on the organism, tissue, or culture system is observed or measured.

In some other embodiments, a dedifferentiated cell generated using the above methods (e.g., an iPS cell) is further treated to generate a differentiated cell that has the same state of differentiation or cell type compared to the somatic cell from which the dedifferentiated cell was generated. In some other embodiments, the dedifferentiated cell generated using the above methods (e.g., an iPS cell) is further treated to generate a differentiated cell that has a different state of differentiation or cell type compared to the somatic cell from which the dedifferentiated cell was generated. In some embodiments, the differentiated cell is generated from the generated dedifferentiated cell (e.g., the generated iPS cell) by introducing mRNA encoding one or more reprogramming factors into the generated iPS cell during one or multiple treatments and maintaining the cell into which the mRNA is introduced under conditions wherein the cell is viable and is differentiated into a cell that has a changed state of differentiation or cell type compared to the generated dedifferentiated cell (e.g., the generated iPS cell) into which the mRNA encoding the one or more reprogramming factors is introduced. In some of these embodiments, the generated differentiated cell that has the changed state of differentiation is used for research, drug screening, or therapeutic applications (e.g., in humans or other animals). For example, the generated differentiated cells find use in the identification and characterization

of reprogramming factors associated with differentiation. In some embodiments, the generated differentiated cells are transplanted into an organism or into a tissue residing *in vitro* or *in vivo*. In some embodiments, an organism, tissue, or culture system housing the generated differentiated cells is exposed to a test compound and the effect of the test compound on the cells or on the organism, tissue, or culture system is observed or measured.

In some preferred embodiments of the method comprising introducing mRNA encoding one or more iPSC induction factors into a somatic cell and maintaining the cell under conditions wherein the cell is viable and the mRNA that is introduced into the cell is expressed in sufficient amount and for sufficient time to generate a dedifferentiated cell (e.g., wherein the dedifferentiated cell is an induced pluripotent stem cell), the sufficient time to generate a dedifferentiated cell is less than one week. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 50 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 100 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 150 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 200 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 300 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 400 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 500 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 600 dedifferentiated cells per 3 x 10^5 input cells (e.g., iPSCs) into which the mRNA is

introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 700 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 800 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 900 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. In some preferred embodiments of this method, the reprogramming efficiency for generating dedifferentiated cells is greater than or equal to 1000 dedifferentiated cells (e.g., iPSCs) per 3 x 10^5 input cells into which the mRNA is introduced. Thus, in some preferred embodiments, this method was greater than 2-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 5-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 10-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 20-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 25-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 30-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 35-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector). In some preferred embodiments, this method was greater than 40-fold more efficient than the published protocol comprising delivery of reprogramming factors with a viral vector (e.g., a lentivirus vector).

The present invention further provides compositions (systems, kits, reaction mixtures, cells, mRNA) used or useful in the methods and/or generated by the methods described herein.

For example, in some embodiments, the present invention provides an mRNA encoding an iPS cell induction factor, the mRNA having pseudouridine in place of uridine.

The present invention further provides compositions comprising a transfection reagent and an mRNA encoding an iPS cell induction factor (e.g., a mixture of transfection reagent and mRNA). In some embodiments, the compositions comprise mRNA encoding a plurality (e.g., 2 or more, 3 or more, 4 or more, 5 or more, or 6) of iPS cell induction factors, including, but not limited to, KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2.

The compositions may further comprise any other reagent or component sufficient, necessary, or useful for practicing any of the methods described herein. Such reagents or components include, but are not limited to, transfection reagents, culture medium (e.g., MEF-condition medium), cells (e.g., somatic cells, iPS cells), containers, boxes, buffers, inhibitors (e.g., RNase inhibitors), labels (e.g., fluorescent, luminescent, radioactive, etc.), positive and/or negative control molecules, reagents for generating capped mRNA, dry ice or other refrigerants, instructions for use, cell culture equipment, detection/analysis equipment, and the like.

This invention provides RNA, oligoribonucleotide, and polyribonucleotide molecules comprising pseudouridine or a modified nucleoside, gene therapy vectors comprising same, gene therapy methods and gene transcription silencing methods comprising same, methods of reducing an immunogenicity of same, and methods of synthesizing same.

In one embodiment, the present invention provides a messenger RNA comprising a pseudouridine residue. In another embodiment, the present invention provides an RNA molecule encoding a protein of interest, said RNA molecule comprising a pseudouridine residue. In another embodiment, the present invention provides an *in vitro-transcribed* RNA molecule, comprising a pseudouridine or a modified nucleoside. In another embodiment, the present invention provides an *in vitro-synthesized* oligoribonucleotide, comprising a pseudouridine or a modified nucleoside, wherein the modified nucleoside is m^5C, m^5U, m^6A, s^2U, Ψ, or 2'-O-methyl-U. In another embodiment, the present invention provides a gene-therapy vector, comprising an *in vitro-synthesized* polyribonucleotide molecule, wherein the polyribonucleotide molecule comprises a pseudouridine or a modified nucleoside.

In another embodiment, the present invention provides a double-stranded RNA (dsRNA) molecule containing, as part of its sequence, a pseudouridine or a modified nucleoside and further comprising an siRNA or shRNA. In another embodiment, the dsRNA molecule is greater

than 50 nucleotides in length. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for inducing a mammalian cell to produce a recombinant protein, comprising contacting the mammalian cell with an *in vitro*-synthesized RNA molecule encoding the recombinant protein, the *in vitro*-synthesized RNA molecule comprising a pseudouridine or a modified nucleoside, thereby inducing a mammalian cell to produce a recombinant protein.

In another embodiment, the present invention provides a method for treating anemia in a subject, comprising contacting a cell of the subject with an *in vitro-synthesized* RNA molecule, the *in vitro*-synthesized RNA molecule encoding erythropoietin, thereby treating anemia in a subject.

In another embodiment, the present invention provides a method for treating a vasospasm in a subject, comprising contacting a cell of the subject with an *in vitro-synthesized* RNA molecule, the *in vitro-synthesized* RNA molecule encoding inducible nitric oxide synthase (iNOS), thereby treating a vasospasm in a subject.

In another embodiment, the present invention provides a method for improving a survival rate of a cell in a subject, comprising contacting the cell with an *in vitro*-synthesized RNA molecule, the *in vitro-synthesized* RNA molecule encoding a heat shock protein, thereby improving a survival rate of a cell in a subject.

In another embodiment, the present invention provides a method for decreasing an incidence of a restenosis of a blood vessel following a procedure that enlarges the blood vessel, comprising contacting a cell of the blood vessel with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a heat shock protein, thereby decreasing an incidence of a restenosis in a subject.

In another embodiment, the present invention provides a method for increasing a hair growth from a hair follicle is a scalp of a subject, comprising contacting a cell of the scalp with an *in vitro* synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a telomerase or an immunosuppressive protein, thereby increasing a hair growth from a hair follicle.

In another embodiment, the present invention provides a method of inducing expression of an enzyme with antioxidant activity in a cell, comprising contacting the cell with an *in vitro*-

synthesized RNA molecule, the *in vitro-synthesized* RNA molecule encoding the enzyme, thereby inducing expression of an enzyme with antioxidant activity in a cell.

In another embodiment, the present invention provides a method for treating cystic fibrosis in a subject, comprising contacting a cell of the subject with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding Cystic Fibrosis Transmembrane Conductance Regulator (CFTR), thereby treating cystic fibrosis in a subject.

In another embodiment, the present invention provides a method for treating an X-linked agammaglobulinemia in a subject, comprising contacting a cell of the subject with an *in vitro* synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a Bruton's tyrosine kinase, thereby treating an X-linked agammaglobulinemia.

In another embodiment, the present invention provides a method for treating an adenosine deaminase severe combined immunodeficiency (ADA SCID) in a subject, comprising contacting a cell of the subject with an *in vitro-synthesized* RNA molecule, the *in vitro*-synthesized RNA molecule encoding an ADA, thereby treating an ADA SCID.

In another embodiment, the present invention provides a method for producing a recombinant protein, comprising contacting an *in vitro* translation apparatus with an *in vitro-synthesized* polyribonucleotide, the *in vitro*-synthesized polyribonucleotide comprising a pseudouridine or a modified nucleoside, thereby producing a recombinant protein.

In another embodiment, the present invention provides a method of synthesizing an *in vitro*-transcribed RNA molecule comprising a modified nucleotide with a pseudouridine modified nucleoside, comprising contacting an isolated polymerase with a mixture of unmodified nucleotides and the modified nucleotide.

In another embodiment, the present invention provides an *in vitro* transcription apparatus, comprising: an unmodified nucleotide, a nucleotide containing a pseudouridine or a modified nucleoside, and a polymerase. In another embodiment, the present invention provides an *in vitro* transcription kit, comprising: an unmodified nucleotide, a nucleotide containing a pseudouridine or a modified nucleoside, and a polymerase. Each possibility represents a separate embodiment of the present invention.

DESCRIPTION OF THE FIGURES

The following figures form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these figures in combination with the detailed description of specific embodiments presented herein.

Figure 1 shows that mRNAs encoding each of the six human reprogramming factors, prepared as described in the EXAMPLES, are translated and localized to the predicted subcellular locations after transfection into human newborn 1079 fibroblasts. Untreated human 1079 fibroblasts: Photos A, E, I, M, Q, and U show phase contrast images of the untreated human 1079 fibroblasts which were not transfected with an mRNA encoding a reprogramming factor and photos B, F, J, N, R, and V show fluorescent images of the same fields after the cells were stained with an antibody specific for each reprogramming factor; these results show that there was little or none of these endogenous reprogramming factor proteins in untreated human 1079 fibroblasts. Treated human 1079 fibroblasts: Photos C, G, K, O, S, and W show phase contrast images of the human 1079 fibroblasts which were transfected with an mRNA encoding the indicated reprogramming factor, and photos D, H, L, P, T, and X show fluorescent images of the same fields after the cells were stained with an antibody specific for each reprogramming factor 24 hours after transfection. These results show that each of the reprogramming factor proteins was expressed in the human 1079 fibroblast cells 24 hours after transfection with the respective reprogramming factor-encoding mRNAs and that the reprogramming factor proteins were localized in the predicted subcellular locations. A-T are at 20x magnification. U-X are at 10x magnification.

Figure 2 shows that mRNA encoding human reprogramming factors (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) produce iPS cells in human somatic cells. Figure 2 shows bright-field (A,C) and immunofluorescent (B,D) images of an iPS cell colony at 12 days after the final transfection with mRNA encoding reprogramming factors. NANOG staining is observed in colony #1 (B, D). Images A and B are at 10x magnification. C and D are at 20x magnification.

Figure 3 shows that iPS colonies derived from human 1079 and IMR90 somatic cells are positive for NANOG and TRA-1-60. Figure 3 shows phase contrast (A,D,G) and immunofluorescent (B,C,E,F,H,I) images of iPS colonies derived from 1079 cells (A, D) and IMR90 cells (G). The same iPS colony shown in (A) is positive for both NANOG (B) and TRA-

1-60 (C). The iPS colony shown in (D) is NANOG-positive (E) and TRA-1-60-positive (F). The iPS colony generated from IMR90 fibroblasts (G) is also positive for both NANOG (H) and TRA-1-60 (I). All images are at 20x magnification.

Figure 4 shows that rapid, enhanced-efficiency iPSC colony formation is achieved by transfecting cells with mRNA encoding reprogramming factors in MEF-conditioned medium. Over 200 colonies were detected 3 days after the final transfection; in the 10-cm dish, IMR90 cells were transfected three times with 36 μg of each reprogramming mRNA (i.e., encoding KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2). Representative iPSC colonies are shown at 4x (A, B), 10x (C-E) and 20x magnification (F). Eight days after the final mRNA transfection with mRNAs encoding the six reprogramming factors, more than 1000 iPSC colonies were counted in IMR90 cells transfected with 18 μg (G, I) or 36 μg (H) of each of the six mRNAs. Representative colonies are shown at 4x magnification (G-H) and at 10x magnification (I).

Figure 5 shows that 1079- and IMR90-derived iPSC colonies are positive for both NANOG and TRA-1-60. Eight days after the final mRNA transfection with 36 μg of mRNA for each of the six reprogramming factors, the 1079-derived iPSC colonies (shown in A, D, and G) are positive for NANOG (B, E, and H) and TRA-1-60 (C, F, and I). Eight days after the final mRNA transfection with 18 μg (J-L) or 36 μg (M-O) of mRNA for each of the six reprogramming factors, IMR90-derived iPS colonies are also positive for NANOG (K, N) and TRA-1-60 (L, O).

Figure 6. Production of TNF-α by MDDCs transfected with natural RNA, demonstrating that unmodified *in vitro-synthesized* RNA and bacterial RNA and mammalian mitochondrial RNA is highly immunogenic, while other mammalian RNA is weakly immunogenic. Human MDDCs were incubated with Lipofectin® alone, or complexed with R-848 (1 μg/ml), or RNA (5 μg/ml) from 293 cells (total, nuclear and cytoplasmic RNAs), mouse heart (polyA+ mRNA), human platelet mitochondrial RNA, bovine tRNA, bacterial tRNA and total RNA *(E. coli)* with or without RNase digestion. After 8 h, TNF-α was measured in the supernatants by ELISA. Mean values ± SEM are shown. Results are representative of 3 independent experiments.

Figure 7. TLR-dependent activation by RNA demonstrates that m6A and s2U modification blocks TLR3 signaling, while all modifications block TLR7 and TLR8 signaling, and that less modified bacterial RNA and unmodified *in vitro*-transcribed RNA activates all three

TLR. (A) Aliquots (1 μg) of *in vitro*-transcribed RNA-1571 without (none) or with m^5C, m^6A, Ψ, m^5U or S^2U nucleoside modifications were analyzed on denaturing agarose gel followed by ethidium bromide-staining and UV illumination. (B) 293 cells expressing human TLR3, TLR7, TLR8 and control vectors were treated with Lipofectin® alone, Lipofectin®-R-848 (1 μg/ml) or RNA *(5 μg*/ml). Modified nucleosides present in RNA-730 and RNA-1571 are noted. (C) CpG ODN-2006 (5 μg/ml), LPS (1.0 μg/ml) and RNA isolates were obtained from rat liver, mouse cell line (TUBO) and human spleen (total), human platelet mitochondrial RNA, or from two different E. *coli* sources. 293-hTLR9 cells served as control. After 8 h, IL-8 was measured in the supernatants by ELISA. Mean values ± SEM are shown. Cell lines containing hTLR3-targeted siRNA are indicated with asterisk. The results are representative of four independent experiments.

Figure 8. Cytokine production by RNA-transfected DC demonstrates that all modifications block activation of cytokine generated DC, while only uridine modifications block blood-derived DC activation. MDDC generated with GM-CSF/IL-4 (A, C) or GM-CSF/IFN-α MDDCs (B), and primary DC1 and DC2 (D) were treated for 8 to 16 h with Lipofectin® alone, Lipofectin®-R-848 (1 μg/ml) or RNA (5 μg/ml). Modified nucleosides present in RNA-1571 are noted. TNF-α, IL-12(p70) and IFN-α were measured in the supernatant by ELISA. Mean values ± SEM are shown. The results are representative of 10 (A and C), 4 (B), and 6 (D) independent experiments. E. Activation of DC by RNA. MDDC were treated for 20 h with Lipofectin® alone or complexed with 1 μg/ml poly(I):(C) or R-848 as positive controls (top panel) or Lipofectin® complexed with the indicated RNA (5 μg/ml; bottom panel). Modified nucleosides present in RNA-1886 are noted. Expression of CD83, CD80, and HLA-DR was determined by flow cytometry.

Figure 9. Activation of DC by RNA demonstrates that all nucleoside modification inhibits the RNA-mediated DC activation. MDDC were treated for 20 h with Lipofectin® alone, Lipofectin®-R-848 (1 μg/ml) or RNA-1571, modified as indicated (5 μg/ml). (A) CD83 and HLA-DR staining. (B) TNF-α levels in the supernatants and mean fluorescence of CD80 andCD86 in response to incubation with RNA. The volume of medium was increased 30-fold for flow cytometry, as indicated by the asterisk. Data are representative of four independent experiments.

Figure 10. Capped RNA-1571 containing different amounts (0, 1, 10, 50, 90, 99 and 100% of modified nucleoside, relative to the corresponding unmodified NTP) were transcribed, and it was found that modification of only a few nucleosides resulted in an inhibition of activation of DC. A. All transcripts were digested to monophosphates and analyzed by reversed-phase HPLC to determine the relative amount of modified nucleoside incorporation. Representative absorbance profiles obtained at the indicated (Ψ:U) ratios are shown. Elution times are noted for 3'-monophosphates of pseudouridine (Ψ), cytidine (C), guanosine (G), uridine (U), 7-methylguanosine ("m7G") and adenosine ("A"). (B) Modified nucleoside content of RNA-1571. The expected percentage of m^6A, Ψ (pseudouridine), or m^5C in RNA-1571 was calculated based on the relative amount of modified NTP in the transcription reaction and the nucleoside composition of RNA-1571 (A: 505, U: 451, C: 273, G: 342). Values for measured modified nucleoside content were determined based on quantitation of the HPLC chromatograms. Notes: A: values (%) for m^6ATP, ΨTP and m^5CTP relative to ATP, UTP and CTP, respectively. B: values for m^6A, Ψ and m^5C monophosphates relative to all NMPs. (C) MDDC were transfected with Lipofectin® complexed capped RNA-1571 (5 μg/ml) containing the indicated amount of m^6A, Ψ or m^5C. After 8 h, TNF-α was measured in the supernatants. Data expressed as relative inhibition of TNF-α. Mean values ± SEM obtained in 3 independent experiments are shown.

Figure 11. TNF-α expression by oligoribonucleotide-transfected DCs demonstrates that as few as one modified nucleoside reduces DC activation. (A) Sequences of oligoribonucleotides (ORN) synthesized chemically (ORN1-4) (SEQ ID NOs: 6-9) or transcribed *in vitro* (ORN5-6) (SEQ ID NOs: 10-11) are shown. Positions of modified nucleosides Um (2'-O-methyluridine), m^5C and Ψ are highlighted. Human MDDC were transfected with Lipofectin® alone (medium), R-848 (1 μg/ml) or Lipofectin® complexed with RNA (5 μg/ml). Where noted, cells were treated with 2.5 μg/ml cycloheximide (CHX). (B). After 8 h incubation, TNF-α was measured in the supernatant. (C) RNA from the cells was analyzed by Northern blot. Representative mean values ± SEM of 3 independent experiments are shown.

Figure 12. A. Ψ-modified mRNA does not stimulate pro-inflammatory cytokine production *in vivo*. Serum samples (6 h after injection) were analyzed by ELISA and revealed that 3 μg of unmodified mRNA induced a higher level of IFN-α than did 3 μg of Ψ-modified mRNA ($P < 0.001$). Levels of IFN-α induced by 3 μg of Ψ-modified mRNA were similar to those obtained when animals were injected with uncomplexed lipofectin. Values are expressed as the mean ± s.e.m. (n = 3 or 5 animals/group). B. Similar results were observed with TNF-α.

Figure 13. mRNA containing pseudouridine (Ψ) does not activate PKR. Ψ: pseudouridine. Control: unmodified RNA. m5C: mRNA with m^5C modification.

Figure 14. Increased expression of luciferase from pseudouridine-containing mRNA in rabbit reticulocyte lysate. Luc-Y: mRNA with pseudouridine modification; luc-C: unmodified RNA. Data is expressed by normalizing luciferase activity to unmodified luciferase RNA.

Figure 15. Increased expression of renilla from pseudouridine-containing mRNA in cultured cells. A. 293 cells. B. Murine primary, bone marrow-derived mouse dendritic cells. renilla-Y: mRNA with pseudouridine modification; renilla -C:unmodified RNA.RNA was modified with m^5C, m^6A,and m^5U as noted.

Figure 16. A. Additive effect of 3' and 5' elements on translation efficiency of Ψ-modified mRNA. 293 cells were transfected with firefly luciferase conventional and Ψ-modified mRNAs that had 5' cap (capLuc), 50 nt-long 3' polyA-tail (TEVlucA50), both or neither of these elements (capTEVlucA50 and Luc, respectively). Cells were lysed 4 h later and luciferase activities measured in aliquots (1/20th) of the total lysates. B. Ψ-modified mRNA is more stable than unmodified mRNA. 293 cells transfected with capTEVlucA_n containing unmodified or Ψ-modified nucleosides were lysed at the indicated times following transfection. Aliquots (1/20th) of the lysates were assayed for luciferase. Standard errors are too small to be visualized with error bars. C. Expression of β-galactosidase is enhanced using Ψ-modified mRNA compared with conventional mRNA. 293 cells seeded in 96-well plates were transfected with lipofectin-complexed mRNAs (0.25 μg/well) encoding bacterial β-galactosidase (lacZ). The transcripts had cap and 3' polyA-tail that were either 30 nt-long (caplacZ) or ~200 nt-long (caplacZ-An).

Constructs made using conventional U or Ψ nucleosides were tested. Cells were fixed and stained with X-gal, 24 h post-transfection. Images were taken by inverted microscopy (40 and 100X magnification) from representative wells.

Figure 17. A. Expression of renilla following intracerebral injection of modified or unmodified encoding mRNA. Rat brain cortex was injected at 8 sites/animals. One hemisphere was injected with capped, renilla-encoding RNA with pseudouridine modification (capRenilla-Y), while the corresponding hemisphere with capped RNA with no nucleoside modification (capRenilla-C). Data from 2 animals (6 injection sites) are shown. BG; lower level of detection of the assay. B. Intravenously-delivered Ψ-modified mRNA is expressed in spleen. Lipofectin-complexed ψmRNA (0.3 μg capTEVlucAn/mouse) was administered by tail vein injection. Animals were sacrificed at 2 and 4 h post-injection and luciferase activities measured in aliquots (1/10th) of organs homogenized in lysis buffer. Values represent luciferase activities in the whole organs. C. Ψ-modified mRNA exhibits greater stability and translation *in vivo*. Lipofectin-complexed capTEVlucAn (0.3 μg/60 μl/animal) with or without Ψ modifications was delivered i.v. to mice. Animals were sacrificed at 1, 4 and 24 h post-injection, and 1/2 of their spleens were processed for luciferase enzyme measurements (left panel) and the other half for RNA analyses (right panel). Luciferase activities were measured in aliquots (1/5th) of the homogenate made from half of the spleens. Plotted values represent luciferase activities in the whole spleen and are expressed as the mean ± s.e.m. (n = 3 or 4/point). D. Expression of firefly luciferase following intratracheal injection of mRNA. capTEVluc-Y: capped, firefly luciferase encoding pseudouridine-modified RNA. CapTEVluc-C: capped RNA with no nucleoside modification.

Figure 18. Protein production is dependent on the amount of mRNA delivered intravenously in mice. The indicated amounts of lipofectin-complexed nucleic acids, capTEVlucAn mRNA with or without Ψ constituents and pCMVluc plasmid DNA in a volume of 60 μl/animal were delivered by i.v. injection into mice. Animals injected with mRNA or plasmid DNA were sacrificed at 6 h or 24 h post-injection, respectively, and luciferase activities were measured in aliquots (1/10th) of their spleens homogenized in lysis buffer. The value from each animal is shown, and short horizontal lines indicate the mean; N.D., not detectable.

Figure 19. Expression of firefly luciferase following intratracheal delivery of encoding mRNA. mRNA were complexed to lipofectin (or PEI, as noted) and animals were injected with 0.3 μg firefly luciferase-encoding mRNA with or without Ψ modification, then sacrificed 3 hours later. Lungs were harvested and homogenized, and luciferase activity was measured in aliquots of the lysed organs.

Figure 20. Ψ-modified mRNA does not induce inflammatory mediators after pulmonary delivery. Induction of TNF-α and IFN-α in serum following intratracheal delivery of luciferase-encoding unmodified mRNA or Ψ-modified mRNA. Serum levels of TNF-α and IFN-α were determined by ELISA 24 hours after mRNA delivery.

Figure 21 shows the results from Example 35: Firefly or Renilla luciferase encoding mRNA with the indicated modifications were complexed to lipofectin and delivered to murine dendritic (A) and HEK293T (B) cells. Human DC were transfected with firefly or renilla luciferase-encoding mRNA complexed with TransIT with the indicated modifications (C). Data is expressed as the fold-change compared to unmodified mRNA.

Figure 22 shows the results from Example 36: T7 polymerase transcription reactions used for the generation of mRNA results in large quantities of RNA of the correct size, but also contains contaminants. This is visualized by application of RNA to a reverse phase HPLC column that separates RNA based on size under denaturing conditions. Ψ-modified TEV-luciferase-A51 RNA was applied to the HPLC column in 38% Buffer B and subjected to a linear gradient of increasing Buffer B to 55%. The profile demonstrated both smaller than expected and larger than expected contaminants.

Figure 23 shows the results from Example 37: (**A**) EPO encoding mRNA with the indicated modifications and with or without HPLC purification were delivered to murine DCs and EPO levels in the supernatant were measured 24 hr later. While m5C/Ψ-modified mRNA had the highest level of translation prior to HPLC purification, Ψ-modified mRNA had the highest translation after HPLC purification. (**B**) Human DCs were transfected with renilla encoding mRNA with the indicated modifications with or without HPLC purification.

Figure 24 shows the results from Example 38: (**A**) Human DCs were transfected with RNA complexed to TransIT with the indicated modifications with or without HPLC purification.

IFN-α levels were measured after 24 hr. HPLC purification increased the immunogenicity of unmodified RNA, which is dependent of the sequence, as other unmodified RNAs had similar levels of IFN-α or reduced levels after HPLC purification. Ψ-modified RNA had unmeasurable levels of IFN-α, similar to control treated DCs. **(B)** Ψ-modified RNA before (-) and after HPLC purification (P1 and P2) was analyzed for dsRNA using dot blotting with a monoclonal antibody specific for dsRNA (J2). Purification of RNA removed dsRNA contamination. (**C**) Ψ-modified RNA encoding iPS factors are immunogenic, which is removed by HPLC purification of the RNA.

Figure 25 provides the mRNA coding sequence for KLF4 (SEQ ID NO:12) and LIN28 (SEQ ID NO:13).

Figure 26 provides the mRNA coding sequence for cMYC (SEQ ID NO:14) and NANOG (SEQ ID NO:15).

Figure 27 provides the mRNA coding sequence for OCT4 (SEQ ID NO:16) and SOX2 (SEQ ID NO:17).

Figure 28 shows that mRNA encoding human reprogramming factors (KLF4, c-MYC, OCT4, and SOX2) produce iPS cells in primary human keratinocyte cells. Figure 28 shows phase contrast images of HEKn cells at 2 days (A) and iPS colony formation at 11 days (B) and 20 days (C) after the final transfection with mRNA encoding 4 reprogramming factors. Images are at 10x magnification.

Figure 29 shows that mRNA encoding human reprogramming factors (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) produce iPS cells in human keratinocytes that are positive for known iPS cell markers. Figure 29 shows phase contrast images of colonies derived from HEKn cells (A, D, and G). The same iPS colony shown in (A) is positive for both KLF4 (B) and LIN28 (C). The iPS colony shown in (D) is SSEA4-positive (E) and TRA-1-60-positive (F). The iPS colony shown in (G) is NANOG-positive (H). All images are at 20x magnification.

Figure 30 shows increases in the expression of 3 iPS-associated messages in HEKn cells transfected with 4 reprogramming mRNAs (KLF4, c-MYC, OCT4, and SOX2) which did not include the reprogramming factor NANOG. Increased expression of the messages was detected by qPCR and is normalized to GAPDH expression. The expression level of each message is depicted in relation to the level found in the original cell line.

DEFINITIONS

The present invention will be understood and interpreted based on terms as defined below.

As used herein "substantially all," in reference to single-stranded complete mRNAs comprising a pseudouridine or 5-methylcytidine residue, means that of all the single-stranded complete mRNAs present in a sample, at least 95% have either a pseudouridine or 5-methylcytidine residue.

As used herein "essentially all," in reference to single-stranded complete mRNAs comprising a pseudouridine or 5-methylcytidine residue, means that of all the single-stranded complete mRNAs present in a sample, at least 99% have either a pseudouridine or 5-methylcytidine residue.

As used herein "RNA contaminant molecules" are molecules that comprise RNA residues and that can at least partially activate an immune response when transfected into a cell (e.g., by activating RNA sensors such as RNA-dependent protein kinase (PKR), retinoic acid-inducible gene-I (RIG-I), Toll-like receptor (TLR)3, TLR7, TLR8, and oligoadenylate synthetase (OAS), or RNA molecules that can at least partially activate an RNA interference (RNAi) response (e.g., including a response to large double-stranded RNA molecules or to small double-stranded RNA molecules (siRNAs)) in the cell. Exemplary RNA contaminant molecules include, but are not limited to: partial or non-full-length mRNAs encoding only a portion of a reprogramming factor (e.g., a non-full-length iPS cell induction factor); single-stranded mRNAs that are greater than the full-length mRNA that encodes a reprogramming factor (e.g., an iPS cell induction factor), e.g., without being bound by theory, by "run-on IVT" or other mechanisms; double-stranded large or small mRNA molecules; and uncapped mRNA molecules.

As used herein, a purified RNA preparation is "substantially free" of RNA contaminant molecules (or a particular recited RNA contaminant), when less than 0.5% of the total RNA in the purified RNA preparation consists of RNA contaminant molecules (or a particularly recited RNA contaminant). The amounts and relative amounts of non-contaminant mRNA molecules and RNA contaminant molecules (or a particular RNA contaminant) may be determined by HPLC or other methods used in the art to separate and quantify RNA molecules.

As used herein, a purified RNA preparation is "essentially free" of RNA contaminant molecules (or a particular recited RNA contaminant), when less than 1.0% of the total RNA in the purified RNA preparation consists of RNA contaminant molecules (or a particularly recited RNA contaminant). The amounts and relative amounts of non-contaminant mRNA molecules and RNA contaminant molecules (or a particular RNA contaminant) may be determined by HPLC or other methods used in the art to separate and quantify RNA molecules.

As used herein, a purified RNA preparation is "virtually free" of RNA contaminant molecules (or a particular recited RNA contaminant), when less than 0.1% of the total RNA in the purified RNA preparation consists of RNA contaminant molecules (or a particularly recited RNA contaminant). The amounts and relative amounts of non-contaminant mRNA molecules and RNA contaminant molecules (or a particular RNA contaminant) may be determined by HPLC or other methods used in the art to separate and quantify RNA molecules.

As used herein, a purified RNA preparation is "free" of RNA contaminant molecules (or a particular recited RNA contaminant), when less than 0.01% of the total RNA in the purified RNA preparation consists of RNA contaminant molecules (or a particularly recited RNA contaminant). The amounts and relative amounts of non-contaminant mRNA molecules and RNA contaminant molecules (or a particular RNA contaminant) may be determined by HPLC or other methods used in the art to separate and quantify RNA molecules.

The terms "comprising", "containing", "having", "include", and "including" are to be construed as "including, but not limited to" unless otherwise noted. The terms "a," "an," and "the" and similar referents in the context of describing the invention and, specifically, in the context of the appended claims, are to be construed to cover both the singular and the plural unless otherwise noted. The use of any and all examples or exemplary language ("for example", "e.g.", "such as") is intended merely to illustrate aspects or embodiments of the invention, and is not to be construed as limiting the scope thereof, unless otherwise claimed.

With respect to the use of the word "derived", such as for an RNA (including mRNA) or a polypeptide that is "derived" from a sample, biological sample, cell, tumor, or the like, it is meant that the RNA or polypeptide either was present in the sample, biological sample, cell, tumor, or the like, or was made using the RNA in the sample, biological sample, cell, tumor, or the like by a process such as an in vitro transcription reaction, or an RNA amplification reaction, wherein the RNA or polypeptide is either encoded by or a copy of all or a portion of the RNA or

polypeptide molecules in the original sample, biological sample, cell, tumor, or the like. By way of example, such RNA can be from an in vitro transcription or an RNA amplification reaction, with or without cloning of cDNA, rather than being obtained directly from the sample, biological sample, cell, tumor, or the like, so long as the original RNA used for the in vitro transcription or an RNA amplification reaction was from the sample, biological sample, cell, tumor, or the like. The terms "sample" and "biological sample" are used in their broadest sense and encompass samples or specimens obtained from any source that contains or may contain eukaryotic cells, including biological and environmental sources. As used herein, the term "sample" when used to refer to biological samples obtained from organisms, includes bodily fluids (e.g., blood or saliva), feces, biopsies, swabs (e.g., buccal swabs), isolated cells, exudates, and the like. The organisms include fungi, plants, animals, and humans. However, these examples are not to be construed as limiting the types of samples or organisms that find use with the present invention. In addition, in order to perform research or study the results related to use of a method or composition of the invention, in some embodiments, a "sample" or "biological sample" comprises fixed cells, treated cells, cell lysates, and the like. In some embodiments, such as embodiments of the method wherein the mRNA is delivered into a cell from an organism that has a known disease or into a cell that exhibits a disease state or a known pathology, the "sample" or "biological sample" also comprises bacteria or viruses.

As used herein, the term "incubating" and variants thereof mean contacting one or more components of a reaction with another component or components, under conditions and for sufficient time such that a desired reaction product is formed.

As used herein, a "nucleoside" consists of a nucleic acid base (e.g., the canonical nucleic acid bases: guanine (G), adenine (A), thymine (T), uracil (U), and cytosine (C)); or a modified nucleic acid base (e.g., 5-methylcytosine (m^5C)), that is covalently linked to a pentose sugar (e.g., ribose or 2'-deoxyribose), whereas and a "nucleotide" or "mononucleotide" consists of a nucleoside that is phosphorylated at one of the hydroxyl groups of the pentose sugar. Linear nucleic acid molecules are said to have a "5' terminus" (5' end) and a "3' terminus" (3' end) because, except with respect to capping or adenylation (e.g., adenylation by a ligase), mononucleotides are joined in one direction via a phosphodiester linkage to make oligonucleotides or polynucleotides, in a manner such that a phosphate on the 5' carbon of one mononucleotide sugar moiety is joined to an oxygen on the 3' carbon of the sugar moiety of its

neighboring mononucleotide. Therefore, an end of a linear single-stranded oligonucleotide or polynucleotide or an end of one strand of a linear double-stranded nucleic acid (RNA or DNA) is referred to as the "5' end" if its 5' phosphate is not joined or linked to the oxygen of the 3' carbon of a mononucleotide sugar moiety, and as the "3' end" if its 3' oxygen is not joined to a 5' phosphate that is joined to a sugar of another mononucleotide. A terminal nucleotide, as used herein, is the nucleotide at the end position of the 3' or 5' terminus.

In order to accomplish specific goals, a nucleic acid base, sugar moiety, or internucleoside (or internucleotide) linkage in one or more of the nucleotides of the mRNA that is introduced into a eukaryotic cell in any of the methods of the invention may comprise a modified base, sugar moiety, or internucleoside linkage. For example, in addition to the other modified nucleotides discussed elsewhere herein for performing the methods of the present invention, one or more of the nucleotides of the mRNA can also have a modified nucleic acid base comprising or consisting of: xanthine; allyamino-uracil; allyamino-thymidine; hypoxanthine; 2-aminoadenine; 5-propynyl uracil; 5-propynyl cytosine; 4-thiouracil; 6-thioguanine; an aza or deaza uracil; an aza or deaza thymidine; an aza or deaza cytosines; an aza or deaza adenine; or an aza or deaza guanines; or a nucleic acid base that is derivatized with a biotin moiety, a digoxigenin moiety, a fluorescent or chemiluminescent moiety, a quenching moiety or some other moiety in order to accomplish one or more specific other purposes; and/or one or more of the nucleotides of the mRNA can have a sugar moiety, such as, but not limited to: 2'-fluoro-2'-deoxyribose or 2'-O-methyl-ribose, which provide resistance to some nucleases; or 2'-amino-2'-deoxyribose or 2'-azido-2'-deoxyribose, which can be labeled by reacting them with visible, fluorescent, infrared fluorescent or other detectable dyes or chemicals having an electrophilic, photoreactive, alkynyl, or other reactive chemical moiety.

In some embodiments of the invention, one or more of the nucleotides of the mRNA comprises a modified internucleoside linkage, such as a phosphorothioate, phosphorodithioate, phosphoroselenate, or phosphorodiselenate linkage, which are resistant to some nucleases, including in a dinucleotide cap analog (Grudzien-Nogalska et al. 2007) that is used in an IVT reaction for co-transcriptional capping of the RNA, or in the poly(A) tail (e.g., by incorporation of a nucleotide that has the modified phosphorothioate, phosphorodithioate, phosphoroselenate, or phosphorodiselenate linkage during IVT of the RNA or, e.g., by incorporation of ATP that contains the modified phosphorothioate, phosphorodithioate, phosphoroselenate, or

phosphorodiselenate linkage into a poly(A) tail on the RNA by polyadenylation using a poly(A) polymerase). The invention is not limited to the modified nucleic acid bases, sugar moieties, or internucleoside linkages listed, which are presented to show examples which may be used for a particular purpose in a method.

As used herein, a "nucleic acid" or a "polynucleotide" or an "oligonucleotide" is a covalently linked sequence of nucleotides in which the 3' position of the sugar moiety of one nucleotide is joined by a phosphodiester bond to the 5' position of the sugar moiety of the next nucleotide (i.e., a 3' to 5' phosphodiester bond), and in which the nucleotides are linked in specific sequence; i.e., a linear order of nucleotides. In some embodiments, the nucleic acid or polynucleotide or oligonucleotide consists of or comprises 2'-deoxyribonucleotides (DNA). In some embodiments, the oligonucleotide consists of or comprises ribonucleotides (RNA).

The terms "isolated" or "purified" when used in relation to a polynucleotide or nucleic acid, as in "isolated RNA" or "purified RNA" refers to a nucleic acid that is identified and separated from at least one contaminant with which it is ordinarily associated in its source. Thus, an isolated or purified nucleic acid (e.g., DNA and RNA) is present in a form or setting different from that in which it is found in nature, or a form or setting different from that which existed prior to subjecting it to a treatment or purification method. For example, a given DNA sequence (e.g., a gene) is found on the host cell chromosome together with other genes as well as structural and functional proteins, and a specific RNA (e.g., a specific mRNA encoding a specific protein), is found in the cell as a mixture with numerous other RNAs and other cellular components. The isolated or purified polynucleotide or nucleic acid may be present in single-stranded or double-stranded form.

A "cap" or a "cap nucleotide" means a nucleoside-5'-triphosphate that, under suitable reaction conditions, is used as a substrate by a capping enzyme system and that is thereby joined to the 5'-end of an uncapped RNA comprising primary RNA transcripts or RNA having a 5'-diphosphate. The nucleotide that is so joined to the RNA is also referred to as a "cap nucleotide" herein. A "cap nucleotide" is a guanine nucleotide that is joined through its 5' end to the 5' end of a primary RNA transcript. The RNA that has the cap nucleotide joined to its 5' end is referred to as "capped RNA" or "capped RNA transcript" or "capped transcript." A common cap nucleoside is 7-methylguanosine or N^7-methylguanosine (sometimes referred to as "standard cap"), which has a structure designated as "m^7G," in which case the capped RNA or "m^7G-capped RNA" has a

structure designated as $m^7G(5')ppp(5')N_1(pN)_x$-OH(3'), or more simply, as $m^7GpppN_1(pN)_x$ or $m^7G[5']ppp[5']N$, wherein m^7G represents the 7-methylguanosine cap nucleoside, ppp represents the triphosphate bridge between the 5' carbons of the cap nucleoside and the first nucleotide of the primary RNA transcript, $N_1(pN)_x$-OH(3') represents the primary RNA transcript, of which N_1 is the most 5'-nucleotide, "p" represents a phosphate group, "G" represents a guanosine nucleoside, "m^7" represents the methyl group on the 7-position of guanine, and "[5']" indicates the position at which the "p" is joined to the ribose of the cap nucleotide and the first nucleoside of the mRNA transcript ("N"). In addition to this "standard cap," a variety of other naturally-occurring and synthetic cap analogs are known in the art. RNA that has any cap nucleotide is referred to as "capped RNA." The capped RNA can be naturally occurring from a biological sample or it can be obtained by *in vitro* capping of RNA that has a 5' triphosphate group or RNA that has a 5' diphosphate group with a capping enzyme system (e.g., vaccinia capping enzyme system or *Saccharomyces cerevisiae* capping enzyme system). Alternatively, the capped RNA can be obtained by *in vitro* transcription (IVT) of a DNA template that contains an RNA polymerase promoter, wherein, in addition to the GTP, the IVT reaction also contains a dinucleotide cap analog (e.g., a m^7GpppG cap analog or an N^7-methyl, 2'-O-methyl-GpppG ARCA cap analog or an N^7-methyl, 3'-O-methyl-GpppG ARCA cap analog) using methods known in the art (e.g., using an AMPLICAP™ T7 capping kit or a MESSAGEMAX™ T7 ARCA-CAPPED MESSAGE Transcription Kit, EPICENTRE or CellScript).

Capping of a 5'-triphosphorylated primary mRNA transcript *in vivo* (or using a capping enzyme system *in vitro*) occurs via several enzymatic steps (Higman et al. 1992, Martin et al. 1975, Myette and Niles 1996).

The following enzymatic reactions are involved in capping of eukaryotic mRNA:

(1) RNA triphosphatase cleaves the 5'-triphosphate of mRNA to a diphosphate,

$pppN_1(p)N_x$-OH(3') → $ppN_1(pN)_x$-OH(3') + Pi; and then

(2) RNA guanyltransferase catalyzes joining of GTP to the 5'-diphosphate of the most 5' nucleotide (N_1) of the mRNA,

$ppN_1(pN)_x$-OH(3') + GTP → $G(5')ppp(5')N_1(pN)_x$-OH(3') + PPi; and finally,

(3) guanine-7-methyltransferase, using S-adenosyl-methionine (AdoMet) as a co-factor, catalyzes methylation of the 7-nitrogen of guanine in the cap nucleotide,

$G(5')ppp(5')N_1(pN)_x\text{-}OH(3') + AdoMet \rightarrow m^7G(5')ppp(5')N_1(pN)_x\text{-}OH(3') + AdoHyc.$

RNA that results from the action of the RNA triphosphatase and the RNA guanyltransferase enzymatic activities, as well as RNA that is additionally methylated by the guanine-7-methyltransferase enzymatic activity, is referred to herein as "5' capped RNA" or "capped RNA", and a "capping enzyme system" or, more simply, a "capping enzyme" herein means any combination of one or more polypeptides having the enzymatic activities that result in "capped RNA." Capping enzyme systems, including cloned forms of such enzymes, have been identified and purified from many sources and are well known in the art (Banerjee 1980, Higman et al. 1992, Higman et al. 1994, Myette and Niles 1996, Shuman 1995, Shuman 2001, Shuman et al. 1980, Wang et al. 1997). Any capping enzyme system that can convert uncapped RNA that has a 5' polyphosphate to capped RNA can be used to provide a capped RNA for any of the embodiments of the present invention. In some embodiments, the capping enzyme system is a poxvirus capping enzyme system. In some preferred embodiments, the capping enzyme system is vaccinia virus capping enzyme. In some embodiments, the capping enzyme system is *Saccharomyces cerevisiae* capping enzyme. Also, in view of the fact that genes encoding RNA triphosphatase, RNA guanyltransferase and guanine-7-methyltransferase from one source can complement deletions in one or all of these genes from another source, the capping enzyme system can originate from one source, or one or more of the RNA triphosphatase, RNA guanyltransferase, and/or guanine-7-methyltransferase activities can comprise a polypeptide from a different source.

A "modified cap nucleotide" of the present invention means a cap nucleotide wherein the sugar, the nucleic acid base, or the internucleoside linkage is chemically modified compared to the corresponding canonical 7-methylguanosine cap nucleotide. Examples of a modified cap nucleotide include a cap nucleotide comprising: (i) a modified 2'- or 3'- deoxyguanosine-5'-triphosphate (or guanine 2'- or 3'- deoxyribonucleic acid-5'-triphosphate) wherein the 2'- or 3'-deoxy position of the deoxyribose sugar moiety is substituted with a group comprising an amino group, an azido group, a fluorine group, a methoxy group, a thiol (or mercapto) group or a methylthio (or methylmercapto) group; or (ii) a modified guanosine-5'-triphosphate, wherein the O^6 oxygen of the guanine base is methylated; or (iii) 3'-deoxyguanosine. For the sake of clarity, it will be understood herein that an "alkoxy-substituted deoxyguanosine-5'-triphosphate" can also be referred to as an "O-alkyl-substituted guanosine-5'-triphosphate"; by way of example, but

without limitation, 2'-methoxy-2'-deoxyguanosine-5'-triphosphate (2'-methoxy-2'-dGTP) and 3'-methoxy-3'-deoxyguanosine-5'-triphosphate (3'-methoxy-3'-dGTP) can also be referred to herein as 2'-O-methylguanosine-5'-triphosphate (2'-OMe-GTP) and 3'-O-methylguanosine-5'-triphosphate (3'-OMe-GTP), respectively. Following joining of the modified cap nucleotide to the 5'-end of the uncapped RNA comprising primary RNA transcripts (or RNA having a 5'-diphosphate), the portion of said modified cap nucleotide that is joined to the uncapped RNA comprising primary RNA transcripts (or RNA having a 5'-diphosphate) may be referred to herein as a "modified cap nucleoside" (i.e., without referring to the phosphate groups to which it is joined), but sometimes it is referred to as a "modified cap nucleotide".

A "modified-nucleotide-capped RNA" is a capped RNA molecule that is synthesized using a capping enzyme system and a modified cap nucleotide, wherein the cap nucleotide on its 5' terminus comprises the modified cap nucleotide, or a capped RNA that is synthesize co-transcriptionally in an in vitro transcription reaction that contains a modified dinucleotide cap analog wherein the dinucleotide cap analog contains the chemical modification in the cap nucleotide. In some embodiments, the modified dinucleotide cap analog is an anti-reverse cap analog or ARCA (Grudzien et al. 2004, Jemielity et al. 2003, Grudzien-Nogalska et al. 2007, Peng et al. 2002, Stepinski et al. 2001).

A "primary RNA" or "primary RNA transcript" means an RNA molecule that is synthesized by an RNA polymerase *in vivo* or *in vitro* and which RNA molecule has a triphosphate on the 5'-carbon of its most 5' nucleotide.

An "RNA amplification reaction" or an "RNA amplification method" means a method for increasing the amount of RNA corresponding to one or multiple desired RNA sequences in a sample. For example, in some embodiments, the RNA amplification method comprises: (a) synthesizing first-strand cDNA complementary to the one or more desired RNA molecules by RNA-dependent DNA polymerase extension of one or more primers that anneal to the desired RNA molecules; (b) synthesizing double-stranded cDNA from the first-strand cDNA using a process wherein a functional RNA polymerase promoter is joined thereto; and (c) contacting the double-stranded cDNA with an RNA polymerase that binds to said promoter under transcription conditions whereby RNA corresponding to the one or more desired RNA molecules is obtained. Unless otherwise stated related to a specific embodiment of the invention, an RNA amplification reaction according to the present invention means a sense RNA amplification reaction, meaning

an RNA amplification reaction that synthesizes sense RNA (e.g., RNA having the same sequence as an mRNA or other primary RNA transcript, rather than the complement of that sequence). Sense RNA amplification reactions known in the art, which are encompassed within this definition include, but are not limited to, the methods which synthesize sense RNA described in Ozawa et al. (Ozawa et al. 2006) and in U.S. Patent Application Nos. 20090053775; 20050153333; 20030186237; 20040197802; and 20040171041. The RNA amplification method described in U.S. Patent Application No. 20090053775 is a preferred method for obtaining amplified RNA derived from one or more cells, which amplified RNA is then used to make mRNA for use in the methods of the present invention.

A "poly-A polymerase" ("PAP") means a template-independent RNA polymerase found in most eukaryotes, prokaryotes, and eukaryotic viruses that selectively uses ATP to incorporate AMP residues to 3'-hydroxylated ends of RNA. Since PAP enzymes that have been studied from plants, animals, bacteria and viruses all catalyze the same overall reaction (Edmonds 1990) are highly conserved structurally (Gershon 2000)and lack intrinsic specificity for particular sequences or sizes of RNA molecules if the PAP is separated from proteins that recognize AAUAAA polyadenylation signals (Wilusz and Shenk 1988), purified wild-type and recombinant PAP enzymes from any of a variety of sources can be used for the present invention. In some embodiments, a PAP enzyme from *Saccharomyces* (e.g., from *S. cerevisiae*) is used for polyadenylation to make purified RNA preparations comprising or consisting of one or more modified mRNAs, each of which encodes a reprogramming factor (e.g., an iPS cell induction factor). In some embodiments, a PAP enzyme from *E. coli* is used for polyadenylation to make purified RNA preparations comprising or consisting of one or more modified mRNAs, each of which encodes a reprogramming factor (e.g., an iPS cell induction factor).

A "reprogramming factor" means a protein, polypeptide, or other biomolecule that, when used alone or in combination with other factors or conditions, causes a change in the state of differentiation of a cell in which the reprogramming factor is introduced or expressed. In some preferred embodiments of the methods of the present invention, the reprogramming factor is a protein or polypeptide that is encoded by an mRNA that is introduced into a cell, thereby generating a cell that exhibits a changed state of differentiation compared to the cell in which the mRNA was introduced. In some preferred embodiments of the methods of the present invention,

the reprogramming factor is a transcription factor. One embodiment of a reprogramming factor used in a method of the present invention is an "iPS cell induction factor."

An "iPS cell induction factor" or "iPSC induction factor" is a protein, polypeptide, or other biomolecule that, when used alone or in combination with other reprogramming factors, causes the generation of iPS cells from somatic cells. Examples of iPS cell induction factors include OCT4, SOX2, c-MYC, KLF4, NANOG and LIN28. iPS cell induction factors include full length polypeptide sequences or biologically active fragments thereof. Likewise an mRNA encoding an iPS cell induction factor may encode a full length polypeptide or biologically active fragments thereof. The mRNA coding seqeunce for exemplary iPS induction factors are shown in Figures 25 (KLF4 and LIN28), 26 (cMYC and NANOG), and 27 (OCT4 and SOX2). In certain emboidments, the present invention employs the sequences or similar sequences shown in these figures, including mRNA molecules that additionally comprise, joined to these mRNA sequences, oligoribonucleotides which exhibit any of the 5' and 3' UTR sequences, Kozak sequences, IRES sequences, cap nucleotides, and/or poly(A) sequences used in the experiments described herein, or which are generally known in the art and which can be used in place of those used herein by joining them to these protein-coding mRNA sequences for the purpose of optimizing translation of the respective mRNA molecules in the cells and improving their stability in the cell in order to accomplish the methods described herein.

"Differentiation" or "cellular differentiation" means the naturally occurring biological process by which a cell that exhibits a less specialized state of differentiation or cell type (e.g., a fertilized egg cell, a cell in an embryo, or a cell in a eukaryotic organism) becomes a cell that exhibits a more specialized state of differentiation or cell type. Scientists, including biologists, cell biologists, immunologists, and embryologists, use a variety of methods and criteria to define, describe, or categorize different cells according to their "cell type," "differentiated state," or "state of differentiation." In general, a cell is defined, described, or categorized with respect to its "cell type," "differentiated state," or "state of differentiation" based on one or more phenotypes exhibited by that cell, which phenotypes can include shape, a biochemical or metabolic activity or function, the presence of certain biomolecules in the cell (e.g., based on stains that react with specific biomolecules), or on the cell (e.g., based on binding of one or more antibodies that react with specific biomolecules on the cell surface). For example, in some embodiments, different

cell types are identified and sorted using a cell sorter or fluorescent-activated cell sorter (FACS) instrument. "Differentiation" or "cellular differentiation" can also occur to cells in culture.

The term "reprogramming" as used herein means differentiation or cellular differentiation that occurs in response to delivery of one or more reprogramming factors into the cell, directly (e.g., by delivery of protein or polypeptide reprogramming factors into the cell) or indirectly (e.g., by delivery of the purified RNA preparation of the present invention which comprises one or more mRNA molecules, each of which encodes a reprogramming factor) and maintaining the cells under conditions (e.g., medium, temperature, oxygen and CO_2 levels, matrix, and other environmental conditions) that are conducive for differentiation. The term "reprogramming" when used herein is not intended to mean or refer to a specific direction or path of differentiation (e.g., from a less specialized cell type to a more specialized cell type) and does not exclude processes that proceed in a direction or path of differentiation than what is normally observed in nature. Thus, in different embodiments of the present invention, "reprogramming" means and includes any and all of the following:

(1) "Dedifferentiation", meaning a process of a cell that exhibits a more specialized state of differentiation or cell type (e.g., a mammalian fibroblast, a keratinocyte, a muscle cell, or a neural cell) going to a cell that exhibits a less specialized state of differentiation or cell type (e.g., an iPS cell);

(2) "Transdifferentiation", meaning a process of a cell that exhibits a more specialized state of differentiation or cell type (e.g., a mammalian fibroblast, a keratinocyte, or a neural cell) going to another more specialized state of differentiation or cell type (e.g., from a fibroblast or keratinocyte to a muscle cell); and

(3) "Redifferentiation" or "Expected Differentiation" or Natural Differentiation", meaning a process of a cell that exhibits any particular state of differentiation or cell type going to another state of differentiation or cell type as would be expected in nature if the cell was present in its natural place and environment (e.g., in an embryo or an organism), whether said process occurs *in vivo* in an organism or in culture (e.g., in response to one or more reprogramming factors).

DESCRIPTION OF THE INVENTION

The present invention provides compositions and methods for reprogramming the state of differentiation of eukaryotic cells, including human or other animal cells, by contacting the cells with purified RNA preparations comprising or consisting of one or more different single-strand mRNA molecules that each encode a reprogramming factor (e.g., an iPS cell induction factor). The purified single-stranded mRNA molecules preferably comprise at least one modified nucleoside selected from the group consisting of a pseudouridine (Ψ), 5-methylcytosine (m^5C), 5-methyluridine (m^5U), 2'-O-methyluridine (Um or $m^{2'\text{-}O}U$), 2-thiouridine (s^2U), and N^6-methyladenosine (m^6A) in place of at least a portion (e.g., including substantially all) of the corresponding unmodified canonical nucleoside of the corresponding unmodified A, C, G, or T canonical nucleoside. In addition, the single-stranded mRNA molecules are preferably purified to be substantially free of RNA contaminant molecules that would activate an unintended response, decrease expression of the single-stranded mRNA, and/or activate RNA sensors in the cells. In certain embodiments, the purified RNA preparations are substantially free of RNA contaminant molecules that are: shorter or longer than the full-length single-stranded mRNA molecules, double-stranded, and/or uncapped RNA. In some preferred embodiments, the invention provides compositions and methods for reprogramming differentiated eukaryotic cells, including human or other animal somatic cells, by contacting the cells with purified RNA preparations comprising or consisting of one or more different single-strand mRNA molecules that each encode an iPS cell induction factor.

In certain embodiments, the mRNA used in the purified RNA preparations is purified to remove substantially, essentially, or virtually all of the contaminants, including substantially, essentially, or virtually all of the RNA contaminants. The present invention is not limited with respect to the purification methods used to purify the mRNA, and the invention includes use of any method that is known in the art or developed in the future in order to purify the mRNA and remove contaminants, including RNA contaminants, that interfere with the intended use of the mRNA. For example, in preferred embodiments, the purification of the mRNA removes contaminants that are toxic to the cells (e.g., by inducing an innate immune response in the cells, or, in the case of RNA contaminants comprising double-stranded RNA, by inducing RNA interference (RNAi), e.g., via siRNA or long RNAi molecules) and contaminants that directly or indirectly decrease translation of the mRNA in the cells). In some embodiments, the mRNA is

purified by HPLC using a method described herein, including in the Examples. In certain embodiments, the mRNA is purified using on a polymeric resin substrate comprising a C18 derivatized styrene-divinylbenzene copolymer and a triethylamine acetate (TEAA) ion pairing agent is used in the column buffer along with the use of an acetonitrile gradient to elute the mRNA and separate it from the RNA contaminants in a size-dependent manner; in some embodiments, the mRNA purification is performed using HPLC, but in some other embodiments a gravity flow column is used for the purification. In some embodiments, the mRNA is purified using a method described in the book entitled "RNA Purification and Analysis" by Douglas T. Gjerde, Lee Hoang, and David Hornby, published by Wiley-VCH, 2009, herein incorporated by reference. In some embodiments, the mRNA purification is carried out in a non-denaturing mode (e.g., at a temperature less than about 50 degrees C, e.g., at ambient temperature). In some embodiments, the mRNA purification is carried out in a partially denaturing mode (e.g., at a temperature less than about 50 degrees C and 72 degrees C). In some embodiments, the mRNA purification is carried out in a denaturing mode (e.g., at a temperature greater than about 72 degrees C). Of course, those with knowledge in the art will know that the denaturing temperature depends on the melting temperature (Tm) of the mRNA that is being purified as well as on the melting temperatures of RNA, DNA, or RNA/DNA hybrids which contaminate the mRNA. In some other embodiments, the mRNA is purified as described by Mellits KH et al. (Removal of double-stranded contaminants from RNA transcripts: synthesis of adenovirus VA RNA1 from a T7 vector. Nucleic Acids Research 18: 5401-5406, 1990, herein incorporated by reference in its entirety). These authors used a three step purification to remove the contaminants which may be used in embodiments of the present invention. Step 1 was 8% polyacrylamide gel electrophoresis in 7M urea (denaturing conditions). The major RNA band was excised from the gel slice and subjected to 8% polyacrylamide gel electrophoresis under nondenaturing condition (no urea) and the major band recovered from the gel slice. Further purification was done on a cellulose CF-11 column using an ethanol-salt buffer mobile phase which separates double stranded RNA from single stranded RNA (Franklin RM. 1966. Proc. Natl. Acad. Sci. USA 55: 1504-1511; Barber R. 1966. Biochem. Biophys. Acta 114:422; and Zelcer A et al. 1982. J. Gen. Virol. 59: 139-148, all of which are herein incorporated by reference) and the final purification step was cellulose chromatography. In some other embodiments, the mRNA is purified using an hydroxylapatite (HAP) column under either non-

denaturing conditions or at higher temperatures (e.g., as described by Pays E. 1977. Biochem. J. 165: 237-245; Lewandowski LJ et al. 1971. J. Virol. 8: 809-812; Clawson GA and Smuckler EA. 1982. Cancer Research 42: 3228-3231; and/or Andrews-Pfannkoch C et al. 2010. Applied and Environmental Microbiology 76: 5039-5045, all of which are herein incorporated by reference). In some other embodiments, the mRNA is purified by weak anion exchange liquid chromatography under non-denaturing conditions (e.g., as described by Easton LE et al. 2010. RNA 16: 647-653 to clean up *in vitro* transcription reactions, herein incorporated by reference). In some embodiments, the mRNA is purified using a combination of any of the above methods or another method known in the art or developed in the future. In still another embodiment, the mRNA used in the compositions and methods of the present invention is purified using a process which comprises treating the mRNA with an enzyme that specifically acts (e.g., digests) one or more contaminant RNA or contaminant nucleic acids (e.g., including DNA), but which does not act on (e.g., does not digest) the desired mRNA. For example, in some embodiments, the mRNA used in the compositions and methods of the present invention is purified using a process which comprises treating the mRNA with a ribonuclease III (RNase III) enzyme (e.g., *E. coli* RNase III) and the mRNA is then purified away from the RNase III digestion products. A ribonuclease III (RNase III) enzyme herein means an enzyme that digests double-stranded RNA greater than about twelve basepairs to shore double-stranded RNA fragments. In some embodiments, the mRNA used in the compositions and methods of the present invention is purified using a process which comprises treating the mRNA with one or more other enzymes that specifically digest one or more contaminant RNAs or contaminant nucleic acids (e.g., including DNA).

This invention provides RNA, oligoribonucleotide, and polyribonucleotide molecules comprising pseudouridine or a modified nucleoside, gene therapy vectors comprising same, gene therapy methods and gene transcription silencing methods comprising same, methods of reducing an immunogenicity of same, and methods of synthesizing same. These modified sequences are preferably present in the purified RNA preparations described herein.

In one embodiment, the present invention provides a messenger RNA comprising a pseudouridine residue. In another embodiment, the messenger RNA encodes a protein of interest. Each possibility represents a separate embodiment of the present invention. In another embodiment, the present invention provides an RNA molecule encoding a protein of interest, said RNA molecule comprising a pseudouridine residue. In another embodiment, the present

invention provides *in vitro*-transcribed RNA molecule, comprising a pseudouridine. In another embodiment, the present invention provides an *in vitro*-transcribed RNA molecule, comprising a modified nucleoside.

As provided herein, the present invention provides methods for synthesizing *in vitro*-transcribed RNA molecules, comprising pseudouridine and/or modified nucleosides. In another embodiment, the present invention provides a messenger RNA molecule comprising a pseudouridine residue.

In another embodiment, an *in vitro*-transcribed RNA molecule of methods and compositions of the present invention is synthesized by T7 phage RNA polymerase. In another embodiment, the molecule is synthesized by SP6 phage RNA polymerase. In another embodiment, the molecule is synthesized by T3 phage RNA polymerase. In another embodiment, the molecule is synthesized by a polymerase selected from the above polymerases. In another embodiment, the *in vitro*-transcribed RNA molecule is an oligoribonucleotide. In another embodiment, the *in vitro*-transcribed RNA molecule is a polyribonucleotide. Each possibility represents a separate embodiment of the present invention. In another embodiment, the present invention provides an *in vitro*-synthesized oligoribonucleotide, comprising a pseudouridine or a modified nucleoside, wherein the modified nucleoside is m^5C, m^5U, m^6A, s^2U, Ψ, or 2'-O-methyl-U. In another embodiment, the present invention provides an *in* vitro-synthesized polyribonucleotide, comprising a pseudouridine or a modified nucleoside, wherein the modified nucleoside is m^5C, m^5U, m^6A, s^2U, Ψ, or 2'-O-methyl-U.

In another embodiment, the *in vitro*-synthesized oligoribonucleotide or polyribonucleotide is a short hairpin (sh)RNA. In another embodiment, the *in vitro*-synthesized oligoribonucleotide is a small interfering RNA (siRNA). In another embodiment, the *in vitro*-synthesized oligoribonucleotide is any other type of oligoribonucleotide known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention further comprises an open reading frame that encodes a functional protein. In another embodiment, the RNA molecule or oligoribonucleotide molecule functions without encoding a functional protein (e.g. in transcriptional silencing), as an RNzyme, etc. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule further comprises a poly-A tail. In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule does not comprise a poly-A tail. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule further comprises an m7GpppG cap. In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule does not comprise an m7GpppG cap. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule further comprises a cap-independent translational enhancer. In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule molecule does not comprise a cap-independent translational enhancer. In another embodiment, the cap-independent translational enhancer is a tobacco etch virus (TEV) cap-independent translational enhancer. In another embodiment, the cap-independent translational enhancer is any other cap-independent translational enhancer known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a gene-therapy vector, comprising an *in vitro*-synthesized polyribonucleotide molecule, wherein the polyribonucleotide molecule comprises a pseudouridine or a modified nucleoside.

In another embodiment, an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention comprises a pseudouridine. In another embodiment, the RNA molecule or oligoribonucleotide molecule comprises a modified nucleoside. In another embodiment, the RNA molecule or oligoribonucleotide molecule is an *in vitro*-synthesized RNA molecule or oligoribonucleotide. Each possibility represents a separate embodiment of the present invention.

"Pseudouridine" refers, in another embodiment, to $m^1acp^3\Psi$ (1-methyl-3-(3-amino-3-carboxypropyl) pseudouridine. In another embodiment, the term refers to $m^1\Psi$ (1--methylpseudouridine). In another embodiment, the term refers to Ψm (2'-O-methylpseudouridine. In another embodiment, the term refers to m^5D (5-methyldihydrouridine). In another embodiment, the term refers to $m^3\Psi$ (3-methylpseudouridine). In another

embodiment, the term refers to a pseudouridine moiety that is not further modified. In another embodiment, the term refers to a monophosphate, diphosphate, or triphosphate of any of the above pseudouridines. In another embodiment, the term refers to any other pseudouridine known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention is a therapeutic oligoribonucleotide.

In another embodiment, the present invention provides a method for delivering a recombinant protein to a subject, the method comprising the step of contacting the subject with an RNA, oligoribonucleotide, polyribonucleotide molecule, or a gene-therapy vector of the present invention, thereby delivering a recombinant protein to a subject.

In another embodiment, the present invention provides a double-stranded RNA (dsRNA) molecule comprising a pseudouridine or a modified nucleoside and further comprising an siRNA or short hairpin RNA (shRNA). In another embodiment, the dsRNA molecule is greater than 50 nucleotides in length. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the pseudouridine or a modified nucleoside is within the siRNA sequence. In another embodiment, the pseudouridine or a modified nucleoside is outside the siRNA sequence. In another embodiment, 1 or more pseudouridine and/or a modified nucleoside residues are present both within and outside the siRNA sequence. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the siRNA or shRNA is contained internally in the dsRNA molecule. In another embodiment, the siRNA or shRNA is contained on one end of the dsRNA molecule. In another embodiment, one or more siRNA or shRNA is contained on one end of the dsRNA molecule, while another one or more is contained internally. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the length of an RNA, oligoribonucleotide, or polyribonucleotide molecule (e.g. a single-stranded RNA (ssRNA) or dsRNA molecule) of methods and compositions of the present invention is greater than 30 nucleotides in length. In another embodiment, the RNA molecule or oligoribonucleotide is greater than 35 nucleotides in length. In another embodiment, the length is at least 40 nucleotides. In another embodiment, the length is at least 45 nucleotides. In another embodiment, the length is at least 55 nucleotides. In another embodiment, the length is at least 60 nucleotides. In another embodiment, the length is at least

60 nucleotides. In another embodiment, the length is at least 80 nucleotides. In another embodiment, the length is at least 90 nucleotides. In another embodiment, the length is at least 100 nucleotides. In another embodiment, the length is at least 120 nucleotides. In another embodiment, the length is at least 140 nucleotides. In another embodiment, the length is at least 160 nucleotides. In another embodiment, the length is at least 180 nucleotides. In another embodiment, the length is at least 200 nucleotides. In another embodiment, the length is at least 250 nucleotides. In another embodiment, the length is at least 300 nucleotides. In another embodiment, the length is at least 350 nucleotides. In another embodiment, the length is at least 400 nucleotides. In another embodiment, the length is at least 450 nucleotides. In another embodiment, the length is at least 500 nucleotides. In another embodiment, the length is at least 600 nucleotides. In another embodiment, the length is at least 700 nucleotides. In another embodiment, the length is at least 800 nucleotides. In another embodiment, the length is at least 900 nucleotides. In another embodiment, the length is at least 1000 nucleotides. In another embodiment, the length is at least 1100 nucleotides. In another embodiment, the length is at least 1200 nucleotides. In another embodiment, the length is at least 1300 nucleotides. In another embodiment, the length is at least 1400 nucleotides. In another embodiment, the length is at least 1500 nucleotides. In another embodiment, the length is at least 1600 nucleotides. In another embodiment, the length is at least 1800 nucleotides. In another embodiment, the length is at least 2000 nucleotides. In another embodiment, the length is at least 2500 nucleotides. In another embodiment, the length is at least 3000 nucleotides. In another embodiment, the length is at least 4000 nucleotides. In another embodiment, the length is at least 5000 nucleotides. Each possibility represents a separate embodiment of the present invention.

In another embodiment, a dsRNA molecule of methods and compositions of the present invention is manufactured by *in vitro*-transcription. In another embodiment, the step of *in vitro*-transcription utilizes T7 phage RNA polymerase. In another embodiment, the *in vitro*-transcription utilizes SP6 phage RNA polymerase. In another embodiment, the *in vitro*-transcription utilizes T3 phage RNA polymerase. In another embodiment, the *in vitro*-transcription utilizes an RNA polymerase selected from the above polymerases. In another embodiment, the *in vitro*-transcription utilizes any other RNA polymerase known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the dsRNA molecule is capable of being processed by a cellular enzyme to yield the siRNA or shRNA. In another embodiment, the cellular enzyme is an endonuclease. In another embodiment, the cellular enzyme is Dicer. Dicer is an RNase III-family nuclease that initiates RNA interference (RNAi) and related phenomena by generation of the small RNAs that determine the specificity of these gene silencing pathways (Bernstein E, Caudy AA et al, Role for a bidentate ribonuclease in the initiation step of RNA interference. Nature 2001; 409(6818): 363-6). In another embodiment, the cellular enzyme is any other cellular enzyme known in the art that is capable of cleaving a dsRNA molecule. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the dsRNA molecule contains two siRNA or shRNA. In another embodiment, the dsRNA molecule contains three siRNA or shRNA. In another embodiment, the dsRNA molecule contains more than three siRNA or shRNA. In another embodiment, the siRNA and/or shRNA are liberated from the dsRNA molecule by a cellular enzyme. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for administering an siRNA or shRNA to a cell, comprising administering a dsRNA molecule of the present invention, wherein the cell processes the dsRNA molecule to yield the siRNA or shRNA, thereby administering a siRNA or shRNA to a cell.

In another embodiment, the nucleoside that is modified in an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention is uridine (U). In another embodiment, the modified nucleoside is cytidine (C). In another embodiment, the modified nucleoside is adenosine (A). In another embodiment the modified nucleoside is guanosine (G). Each possibility represents a separate embodiment of the present invention.

In another embodiment, the modified nucleoside of methods and compositions of the present invention is m^5C (5-methylcytidine). In another embodiment, the modified nucleoside is m^5U (5-methyluridine). In another embodiment, the modified nucleoside is m^6A (N^6-methyladenosine). In another embodiment, the modified nucleoside is s^2U (2-thiouridine). In another embodiment, the modified nucleoside is Ψ (pseudouridine). In another embodiment, the modified nucleoside is Um (2'-O-methyluridine).

In other embodiments, the modified nucleoside is m^1A (1-methyladenosine); m^2A (2-methyladenosine); Am(2'-O-methyladenosine); ms^2m^GA (2-methylthio-N^6-methyladenosine);

i^6A(~isopentenyladenosine); ms^2i6A (2-methylthio-N^6isopentenyladenosine); io^6A (N^6-(cis-hydroxyisopentenyl)adenosine); ms^2io^6A (2-methylthio-N^6-(cis-hydroxyisopentenyl)adenosine); g^6A (N^6-glycinylcarbamoyladenosine); t^6A (N^6-threonylcarbamoyladenosine); ms^2t^6A (2-methylthio-N^6-threonylcarbamoyladenosine); m^6t^6A (N^6-methyl-N^6-threonylcarbamoyladenosine); hn^6A(N^6hydroxynorvalylcarbamoyladenosine); ms^2hn^6A (2-methylthio-N^6-hydroxynorvalyl carbamoyladenosine); Ar(p) (2'-O-ribosyladenosine (phosphate)); I (inosine); m^1I (1-methylinosine); m^1Im (1,2'-O-dimethylinosine); m^3C (3-methylcytidine); Cm (2'-O-methylcytidine); S^2C (2thiocytidine); ac^4C (N^4-acetylcytidine); f^5C (5-formylcytidine); m^5Cm (5,2'-O-dimethylcytidine); ac^4Cm (N^4-acetyl-2'-O-methylcytidine); k^2C (lysidine); m^1G (1-methylguanosine); m^2G (N^2-methylguanosine); m^7G (7-methylguanosine); Gm (2'-O-methylguanosine); $m^2{}_2G$ (N^2,N^2-dimethylguanosine); m^2Gm (N^2,2'-O-dimethylguanosine); $m^2{}_2Gm$ (N^2,N^2,2'-O-trimethylguanosine); Gr(p) (2'-O-ribosylguanosine (phosphate)); yW (wybutosine); o_2yW (peroxywybutosine); OHyW (hydroxywybutosine); OHyW* (undermodified hydroxywybutosine); imG (wyosine); mimG (methylwyosine); Q (queuosine); oQ (epoxyqueuosine); galQ (galactosyl-queuosine); manQ (mannosylqueuosine); $preQ_0$ (7-cyano-7-deazaguanosine); $preQ_1$ (7-aminomethyl-7-deazaguanosine); G^+ (archaeosine); D (dihydrouridine); m^5Um (5,2'-O-dimethyluridine); S^4U (4-thiouridine); m^5s^2U (5-methyl-2-thiouridine); s^2Um (2-thio-2'-O-methyluridine); acp^3U (3-(3-amino-3-carboxypropyl)uridine); ho^5U (5-hydroxyuridine); mo^5U (5-methoxyuridine); cmo^5U (uridine 5-oxyacetic acid); $mcmo^5U$ (uridine 5-oxyacetic acid methyl ester); chm^5U (5-(carboxyhydroxymethyl)uridine)); $mchm^5U$ (5(carboxyhydroxymethyl)uridine methyl ester); mcm^5U (5-methoxycarbonylmethyluridine); mcm^5Um (5-methoxycarbonylmethyl-2'-O-methyluridine); mcm^5s^2U (5-methoxycarbonylmethyl-2-thiouridine); nm^5s^2U (5-aminomethyl-2-thiouridine); mnm^5U (5-methylaminomethyluridine); mnm^5s^2U (5-methylaminomethyl-2-thiouridine); $mnmse^2U$ (5-methylaminomethyl-2-selenouridine); ncm^5U (5-carbamoylmethyluridine); ncm^5Um (5-carbamoylmethyl-2'-O-methyluridine); $cmnm^5U$ (5-carboxymethylaminomethyluridine); $cmnm^5Um$ (5-carboxymethylaminomethyl-2'--methyluridine); $cmnm^5s^2U$ (5-carboxymethylaminomethyl-2-thiouridine); $m^6{}_2A$ (N^6,N^6-dimethyladenosine); Im (2'-O-methylinosine); m^4C (N^4-methylcytidine); m^4Cm (N^4,2'-O-dimethylcytidine); hm^5C (5-hydroxymethylcytidine); m^3U (3-methyluridine); cm^5U (5-carboxymethyluridine); m^6Am (N^6,2'-Odimethyladenosine); $m^6{}_2Am$ (N^6,N^6, 2'-O-trimethyladenosine); $m^{2,7}G$(N^2,7-

dimethylguanosine); $m^{2,2,7}G$ (N^2,N^2,7-trimethylguanosine); m^3Um (3,2'-O-dimethyluridine); m^5D (5-methyldihydrouridine); f^5Cm (5-formyl-2'-O-methylcytidine); m^1Gm (1,2'-O-dimethylguanosine); m^1Am (1,2'-O-dimethyladenosine); τm^5U (5-taurinomethyluridine); $\tau m5s2U$ (5-taurinomethyl-2-thiouridine)); imG-14 (4-demethylwyosine); imG2 (isowyosine); or ac^6A (N^6-acetyladenosine). Each possibility represents a separate embodiment of the present invention.

In another embodiment, an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention comprises a combination of 2 or more of the above modifications. In another embodiment, the RNA molecule or oligoribonucleotide molecule comprises a combination of 3 or more of the above modifications. In another embodiment, the RNA molecule or oligoribonucleotide molecule comprises a combination of more than 3 of the above modifications. Each possibility represents a separate embodiment of the present invention.

In another embodiment, between 0.1% and 100% of the residues in the RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention are modified (e.g. either by the presence of pseudouridine or a modified nucleoside base). In another embodiment, 0.1 % of the residues are modified. In another embodiment, 0.2%. In another embodiment, the fraction is 0.3%. In another embodiment, the fraction is 0.4%. In another embodiment, the fraction is 0.5%. In another embodiment, the fraction is 0.6%. In another embodiment, the fraction is 0.8%. In another embodiment, the fraction is 1%. In another embodiment, the fraction is 1.5%. In another embodiment, the fraction is 2%. In another embodiment, the fraction is 2.5%. In another embodiment, the fraction is 3%. In another embodiment, the fraction is 4%. In another embodiment, the fraction is 5%. In another embodiment, the fraction is 6%. In another embodiment, the fraction is 8%. In another embodiment, the fraction is 10%. In another embodiment, the fraction is 12%. In another embodiment, the fraction is 14%. In another embodiment, the fraction is 16%. In another embodiment, the fraction is 18%. In another embodiment, the fraction is 20%. In another embodiment, the fraction is 25%. In another embodiment, the fraction is 30%. In another embodiment, the fraction is 35%. In another embodiment, the fraction is 40%. In another embodiment, the fraction is 45%. In another embodiment, the fraction is 50%. In another

embodiment, the fraction is 60%. In another embodiment, the fraction is 70%. In another embodiment, the fraction is 80%. In another embodiment, the fraction is 90%. In another embodiment, the fraction is 100%.

In another embodiment, the fraction is less than 5%. In another embodiment, the fraction is less than 3%. In another embodiment, the fraction is less than 1%. In another embodiment, the fraction is less than 2%. In another embodiment, the fraction is less than 4%. In another embodiment, the fraction is less than 6%. In another embodiment, the fraction is less than 8%. In another embodiment, the fraction is less than 10%. In another embodiment, the fraction is less than 12%. In another embodiment, the fraction is less than 15%. In another embodiment, the fraction is less than 20%. In another embodiment, the fraction is less than 30%. In another embodiment, the fraction is less than 40%. In another embodiment, the fraction is less than 50%. In another embodiment, the fraction is less than 60%. In another embodiment, the fraction is less than 70%.

In another embodiment, 0.1% of the residues of a given nucleotide (uridine, cytidine, guanosine, or adenine) are modified. In another embodiment, the fraction of the nucleotide is 0.2%. In another embodiment, the fraction is 0.3%. In another embodiment, the fraction is 0.4%. In another embodiment, the fraction is 0.5%. In another embodiment, the fraction is 0.6%. In another embodiment, the fraction is 0.8%. In another embodiment, the fraction is 1%. In another embodiment, the fraction is 1.5%. In another embodiment, the fraction is 2%. In another embodiment, the fraction is 2.5%. In another embodiment, the fraction is 3%. In another embodiment, the fraction is 4%. In another embodiment, the fraction is 5%. In another embodiment, the fraction is 6%. In another embodiment, the fraction is 8%. In another embodiment, the fraction is 10%. In another embodiment, the fraction is 12%. In another embodiment, the fraction is 14%. In another embodiment, the fraction is 16%. In another embodiment, the fraction is 18%. In another embodiment, the fraction is 20%. In another embodiment, the fraction is 25%. In another embodiment, the fraction is 30%. In another embodiment, the fraction is 35%. In another embodiment, the fraction is 40%. In another embodiment, the fraction is 45%. In another embodiment, the fraction is 50%. In another embodiment, the fraction is 60%. In another embodiment, the fraction is 70%. In another embodiment, the fraction is 80%. In another embodiment, the fraction is 90%. In another embodiment, the fraction is 100%.

In another embodiment, the fraction of the given nucleotide is less than 8%. In another embodiment, the fraction is less than 10%. In another embodiment, the fraction is less than 5%. In another embodiment, the fraction is less than 3%. In another embodiment, the fraction is less than 1%. In another embodiment, the fraction is less than 2%. In another embodiment, the fraction is less than 4%. In another embodiment, the fraction is less than 6%. In another embodiment, the fraction is less than 12%. In another embodiment, the fraction is less than 15%. In another embodiment, the fraction is less than 20%. In another embodiment, the fraction is less than 30%. In another embodiment, the fraction is less than 40%. In another embodiment, the fraction is less than 50%. In another embodiment, the fraction is less than 60%. In another embodiment, the fraction is less than 70%.

In another embodiment, the terms "ribonucleotide," "oligoribonucleotide," and "polyribonucleotide" refers to a string of at least 2 base-sugar-phosphate combinations. The term includes, in another embodiment, compounds comprising nucleotides in which the sugar moiety is ribose. In another embodiment, the term includes both RNA and RNA derivatives in which the backbone is modified. "Nucleotides" refers, in another embodiment, to the monomeric units of nucleic acid polymers. RNA may be, in another embodiment, in the form of a tRNA (transfer RNA), snRNA (small nuclear RNA), rRNA (ribosomal RNA), mRNA (messenger RNA), anti-sense RNA, small interfering RNA (siRNA), micro RNA (miRNA) and ribozymes. The use of siRNA and miRNA has been described (Caudy AA et al, Genes & Devel 16: 2491-96 and references cited therein). In addition, these forms of RNA may be single, double, triple, or quadruple stranded. The term also includes, in another embodiment, artificial nucleic acids that may contain other types of backbones but the same bases. In another embodiment, the artificial nucleic acid is a PNA (peptide nucleic acid). PNA contain peptide backbones and nucleotide bases and are able to bind, in another embodiment, to both DNA and RNA molecules. In another embodiment, the nucleotide is oxetane modified. In another embodiment, the nucleotide is modified by replacement of one or more phosphodiester bonds with a phosphorothioate bond. In another embodiment, the artificial nucleic acid contains any other variant of the phosphate backbone of native nucleic acids known in the art. The use of phosphothiorate nucleic acids and PNA are known to those skilled in the art, and are described in, for example, Neilsen PE, Curr Opin Struct Biol 9:353-57~and Raz NK et al Biochem Biophys Res Commun. 297:1075-84. The production and use of nucleic acids is known to those skilled in art and is described, for example,

in Molecular Cloning, (2001), Sambrook and Russell, eds. and Methods in Enzymology: Methods for molecular cloning in eukaryotic cells (2003) Purchio and G. C. Fareed. Each nucleic acid derivative represents a separate embodiment of the present invention

In another embodiment, the term "oligoribonucleotide" refers to a string comprising fewer than 25 nucleotides (nt). In another embodiment, "oligoribonucleotide" refers to a string of fewer than 24 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 23 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 22 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 21 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 20 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 19 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 18 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 17 nucleotides. In another embodiment, "oligoribonucleotide" refers to a string of fewer than 16 nucleotides. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the term "polyribonucleotide" refers to a string comprising more than 25 nucleotides (nt). In another embodiment, "polyribonucleotide" refers to a string of more than 26 nucleotides. In another embodiment, "polyribonucleotide" refers to a string of more than 28 nucleotides. In another embodiment, "the term" refers to a string of more than 30 nucleotides. In another embodiment, "the term" refers to a string of more than 32 nucleotides. In another embodiment, "the term" refers to a string of more than 35 nucleotides. In another embodiment, "the term" refers to a string of more than 40 nucleotides. In another embodiment, "the term" refers to a string of more than 50 nucleotides. In another embodiment, "the term" refers to a string of more than 60 nucleotides. In another embodiment, "the term" refers to a string of more than 80 nucleotides. In another embodiment, "the term" refers to a string of more than 100 nucleotides. In another embodiment, "the term" refers to a string of more than 120 nucleotides. In another embodiment, "the term" refers to a string of more than150 nucleotides. In another embodiment, "the term" refers to a string of more than 200 nucleotides. In another embodiment, "the term" refers to a string of more than 300 nucleotides. In another embodiment, "the term" refers to a string of more than 400 nucleotides. In another embodiment, "the term" refers to a string of more than 500 nucleotides. In another embodiment, "the term" refers to a string of more

than600 nucleotides. In another embodiment, "the term" refers to a string of more than 800 nucleotides. In another embodiment, "the term" refers to a string of more than 1000 nucleotides. In another embodiment, "the term" refers to a string of more than 1200 nucleotides. In another embodiment, "the term" refers to a string of more than 1400 nucleotides. In another embodiment, "the term" refers to a string of more than 1600 nucleotides. In another embodiment, "the term" refers to a string of more than1800 nucleotides. In another embodiment, "the term" refers to a string of more than 2000 nucleotides. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for inducing a mammalian cell to produce a protein of interest, comprising contacting the mammalian cell with an *in vitro*-synthesized RNA molecule encoding the recombinant protein, the *in vitro*-synthesized RNA molecule comprising a pseudouridine or a modified nucleoside, thereby inducing a mammalian cell to produce a protein of interest. In another embodiment, the protein of interest is a recombinant protein. Each possibility represents a separate embodiment of the present invention.

"Encoding" refers, in another embodiment, to an RNA molecule that encodes the protein of interest. In another embodiment, the RNA molecule comprises an open reading frame that encodes the protein of interest. In another embodiment, one or more other proteins is also encoded. In another embodiment, the protein of interest is the only protein encoded. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method of inducing a mammalian cell to produce a recombinant protein, comprising contacting the mammalian cell with an *in vitro*-transcribed RNA molecule encoding the recombinant protein, the *in vitro*-transcribed RNA molecule further comprising a pseudouridine or a modified nucleoside, thereby inducing a mammalian cell to produce a recombinant protein.

In another embodiment, an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention is translated in the cell more efficiently than an unmodified RNA molecule with the same sequence. In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule exhibits enhanced ability to be translated by a target cell. In another embodiment, translation is enhanced by a factor of 2-fold relative to its unmodified counterpart. In another embodiment, translation is enhanced by a 3-fold factor. In

another embodiment, translation is enhanced by a 5-fold factor. In another embodiment, translation is enhanced by a 7-fold factor. In another embodiment, translation is enhanced by a 10-fold factor. In another embodiment, translation is enhanced by a 15-fold factor. In another embodiment, translation is enhanced by a 20-fold factor. In another embodiment, translation is enhanced by a 50-fold factor. In another embodiment, translation is enhanced by a 100-fold factor. In another embodiment, translation is enhanced by a 200-fold factor. In another embodiment, translation is enhanced by a 500-fold factor. In another embodiment, translation is enhanced by a 1000-fold factor. In another embodiment, translation is enhanced by a 2000-fold factor. In another embodiment, the factor is 10-1000-fold. In another embodiment, the factor is 10-100-fold. In another embodiment, the factor is 10-200-fold. In another embodiment, the factor is 10-300-fold. In another embodiment, the factor is 10-500-fold. In another embodiment, the factor is 20-1000-fold. In another embodiment, the factor is 30-1000-fold. In another embodiment, the factor is 50-1000-fold. In another embodiment, the factor is 100-1000-fold. In another embodiment, the factor is 200-1000-fold. In another embodiment, translation is enhanced by any other significant amount or range of amounts. Each possibility represents a separate embodiment of the present invention.

Methods of determining translation efficiency are well known in the art, and include, e.g. measuring the activity of an encoded reporter protein (e.g. luciferase or renilla [Examples herein] or green fluorescent protein [Wall AA, *Phillips AM et al, Effective translation of the second* cistron in two Drosophila dicistronic transcripts is determined by the absence of in-frame AUG codons in the first cistron. J Biol Chem 2005;280(30): 27670-8]), or measuring radioactive label incorporated into the translated protein (Ngosuwan J, Wang NM et al, Roles of cytosolic Hsp70 and Hsp40 molecular chaperones in post-translational translocation of presecretory proteins into the endoplasmic reticulum. J Biol Chem 2003;278(9): 7034-42). Each method represents a separate embodiment of the present invention.

In some expression studies provided herein, translation was measured from RNA complexed to Lipofectin® (Gibco BRL, Gaithersburg, MD, USA) and injected into the tail vein of mice. In the spleen lysates, pseudouridine-modified RNA was translated significantly more efficiently than unmodified RNA (Figure 17B). Under the conditions utilized herein, efficiency of transfection-based methods of the present invention correlates with the ability of the transfection reagent to penetrate into tissues, providing an explanation for why the effect was

most pronounced in spleen cells. Splenic blood flow is an open system, with blood contents directly contacting red and white pulp elements including lymphoid cells.

In another experiment, *in vitro* phosphorylation assays were performed using recombinant human PKR and its substrate, eIF2α in the presence of capped, renilla-encoding mRNA (0.5 and 0.05 ng/μl). mRNA containing pseudouridine (Ψ) did not activate PKR, as detected by lack of both self-phosphorylation of PKR and phosphorylation of eIF2α, while RNA without nucleoside modification and mRNA with m5C modification activated PKR. Phosphorylated eIF2α is known to block initiation of mRNA translation, therefore lack of phosphorylation enables, in another embodiment, enhanced translation of the mRNA containing pseudouridine (Ψ).

In another embodiment, the enhanced translation is in a cell (relative to translation in the same cell of an unmodified RNA molecule with the same sequence; Examples 13-14). In another embodiment, the enhanced translation is *in vitro* (e.g. in an *in vitro* translation mix or a reticulocyte lysate; Examples 13-14. In another embodiment, the enhanced translation is *in vivo* (Example 13). In each case, the enhanced translation is relative to an unmodified RNA molecule with the same sequence, under the same conditions. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention is significantly less immunogenic than an unmodified *in vitro*-synthesized RNA molecule with the same sequence. In another embodiment, the modified RNA molecule is 2-fold less immunogenic than its unmodified counterpart. In another embodiment, immunogenicity is reduced by a 3-fold factor. In another embodiment, immunogenicity is reduced by a 5-fold factor. In another embodiment, immunogenicity is reduced by a 7-fold factor. In another embodiment, immunogenicity is reduced by a 10-fold factor. In another embodiment, immunogenicity is reduced by a 15-fold factor. In another embodiment, immunogenicity is reduced by a 20-fold factor. In another embodiment, immunogenicity is reduced by a 50-fold factor. In another embodiment, immunogenicity is reduced by a 100-fold factor. In another embodiment, immunogenicity is reduced by a 200-fold factor. In another embodiment, immunogenicity is reduced by a 500-fold factor. In another

embodiment, immunogenicity is reduced by a 1000-fold factor. In another embodiment, immunogenicity is reduced by a 2000-fold factor. In another embodiment, immunogenicity is reduced by another fold difference.

In another embodiment, "significantly less immunogenic" refers to a detectable decrease in immunogenicity. In another embodiment, the term refers to a fold decrease in immunogenicity (e.g. one of the fold decreases enumerated above). In another embodiment, the term refers to a decrease such that an effective amount of the RNA, oligoribonucleotide, or polyribonucleotide molecule can be administered without triggering a detectable immune response. In another embodiment, the term refers to a decrease such that the RNA, oligoribonucleotide, or polyribonucleotide molecule can be repeatedly administered without eliciting an immune response sufficient to detectably reduce expression of the recombinant protein. In another embodiment, the decrease is such that the RNA, oligoribonucleotide, or polyribonucleotide molecule can be repeatedly administered without eliciting an immune response sufficient to eliminate detectable expression of the recombinant protein.

"Effective amount" of the RNA, oligoribonucleotide, or polyribonucleotide molecule refers, in another embodiment, to an amount sufficient to exert a therapeutic effect. In another embodiment, the term refers to an amount sufficient to elicit expression of a detectable amount of the recombinant protein. Each possibility represents a separate embodiment of the present invention.

Reduced immunogenicity of RNA, oligoribonucleotide, and polyribonucleotide molecules of the present invention is demonstrated herein (Examples 4-11).

Methods of determining immunogenicity are well known in the art, and include, e.g. measuring secretion of cytokines (e.g. IL-12, IFN-α, TNF-α, RANTES, MIP-1α or β, IL-6, IFN-β, or IL-8; Examples herein), measuring expression of DC activation markers (e.g. CD83, HLA-DR, CD80 and CD86; Examples herein), or measuring ability to act as an adjuvant for an adaptive immune response. Each method represents a separate embodiment of the present invention.

In another embodiment, the relative immunogenicity of the modified nucleotide and its unmodified counterpart are determined by determining the quantity of the modified nucleotide required to elicit one of the above responses to the same degree as a given quantity of the

unmodified nucleotide. For example, if twice as much modified nucleotide is required to elicit the same response, than the modified nucleotide is two-fold less immunogenic than the unmodified nucleotide.

In another embodiment, the relative immunogenicity of the modified nucleotide and its unmodified counterpart are determined by determining the quantity of cytokine (e.g. IL-12, IFN-α, TNF-α, RANTES, MIP-1α or β, IL-6, IFN-β, or IL-8) secreted in response to administration of the modified nucleotide, relative to the same quantity of the unmodified nucleotide. For example, if one-half as much cytokine is secreted, than the modified nucleotide is two-fold less immunogenic than the unmodified nucleotide. In another embodiment, background levels of stimulation are subtracted before calculating the immunogenicity in the above methods. Each possibility represents a separate embodiment of the present invention.

In another embodiment, a method of present invention further comprises mixing the RNA, oligoribonucleotide, or polyribonucleotide molecule with a transfection reagent prior to the step of contacting. In another embodiment, a method of present invention further comprises administering the RNA, oligoribonucleotide, or polyribonucleotide molecule together with the transfection reagent. In another embodiment, the transfection reagent is a cationic lipid reagent (Example 6).

In another embodiment, the transfection reagent is a lipid-based transfection reagent. In another embodiment, the transfection reagent is a protein-based transfection reagent. In another embodiment, the transfection reagent is a polyethyleneimine based transfection reagent. In another embodiment, the transfection reagent is calcium phosphate. In another embodiment, the transfection reagent is Lipofectin® or Lipofectamine®. In another embodiment, the transfection reagent is any other transfection reagent known in the art.

In another embodiment, the transfection reagent forms a liposome. Liposomes, in another embodiment, increase intracellular stability, increase uptake efficiency and improve biological activity. In another embodiment, liposomes are hollow spherical vesicles composed of lipids arranged in a similar fashion as those lipids which make up the cell membrane. They have, in another embodiment, an internal aqueous space for entrapping water soluble compounds and range in size from 0.05 to several microns in diameter. In another embodiment, liposomes can deliver RNA to cells in a biologically active form.

Each type of transfection reagent represents a separate embodiment of the present invention.

In another embodiment, the target cell of methods of the present invention is an antigen--presenting cell. In another embodiment, the cell is an animal cell. In another embodiment, the cell is a dendritic cell (Example 14). In another embodiment, the cell is a neural cell. In another embodiment, the cell is a brain cell (Example 16). In another embodiment, the cell is a spleen cell. In another embodiment, the cell is a lymphoid cell. In another embodiment, the cell is a lung cell (Example 16). In another embodiment, the cell is a skin cell. In another embodiment, the cell is a keratinocyte. In another embodiment, the cell is an endothelial cell. In another embodiment, the cell is an astrocyte, a microglial cell, or a neuron (Example 16). In another embodiment, the cell is an alveolar cell (Example 16). In another embodiment, the cell is a surface alveolar cell (Example 16). In another embodiment, the cell is an alveolar macrophage. In another embodiment, the cell is an alveolar pneumocyte. In another embodiment, the cell is a vascular endothelial cell. In another embodiment, the cell is a mesenchymal cell. In another embodiment, the cell is an epithelial cell. In another embodiment, the cell is a hematopoietic cell. In another embodiment, the cell is colonic epithelium cell. In another embodiment, the cell is a lung epithelium cell. In another embodiment, the cell is a bone marrow cell.

In other embodiments, the target cell is a Claudius' cell, Hensen cell, Merkel cell, Muller cell, Paneth cell, Purkinje cell, Schwann cell, Sertoli cell, acidophil cell, acinar cell, adipoblast, adipocyte, brown or white alpha cell, amacrine cell, beta cell, capsular cell, cementocyte, chief cell, chondroblast, chondrocyte, chromaffin cell, chromophobic cell, corticotroph, delta cell, Langerhans cell, follicular dendritic cell, enterochromaffin cell, ependymocyte, epithelial cell, basal cell, squamous cell, endothelial cell, transitional cell, erythroblast, erythrocyte, fibroblast, fibrocyte, follicular cell, germ cell, gamete, ovum, spermatozoon, oocyte, primary oocyte, secondary oocyte, spermatid, spermatocyte, primary spermatocyte, secondary spermatocyte, germinal epithelium, giant cell, glial cell, astroblast, astrocyte, oligodendroblast, oligodendrocyte, glioblast, goblet cell, gonadotroph, granulosa cell, haemocytoblast, hair cell, hepatoblast, hepatocyte, hyalocyte, interstitial cell,' juxtaglomerular cell, keratinocyte, keratocyte, lemmal cell, leukocyte, granulocyte, basophil, eosinophil, neutrophil, lymphoblast, B-lymphoblast, T-lymphoblast, lymphocyte, B-lymphocyte, T-lymphocyte, helper induced T-lymphocyte, Th1 T-lymphocyte, Th2 T-lymphocyte, natural killer cell, thymocyte, macrophage,

Kupffer cell, alveolarmacrophage, foam cell, histiocyte, luteal cell, lymphocytic stem cell, lymphoid cell, lymphoid stem cell, macroglial cell, mammotroph, mast cell, medulloblast, megakaryoblast, megakaryocyte, melanoblast, melanocyte, mesangial cell, mesothelial cell, metamyelocyte, monoblast, monocyte, mucous neck cell, muscle cell, cardiac muscle cell, skeletal muscle cell, smooth muscle cell, myelocyte, myeloid cell, myeloid stem cell, myoblast, myoepithelial cell, myofibrobast, neuroblast, neuroepithelial cell, neuron, odontoblast, osteoblast, osteoclast, osteocyte, oxyntic cell, parafollicular cell, paraluteal cell, peptic cell, pericyte, peripheral blood mononuclear cell, phaeochromocyte, phalangeal cell, pinealocyte, pituicyte, plasma cell, platelet, podocyte, proerythroblast, promonocyte, promyeloblast, promyelocyte, pronormoblast, reticulocyte, retinal pigment epithelial cell, retinoblast, small cell, somatotroph, stem cell, sustentacular cell, teloglial cell, or zymogenic cell. Each possibility represents a separate embodiment of the present invention.

A variety of disorders may be treated by employing methods of the present invention including, *inter alia,* monogenic disorders, infectious diseases, acquired disorders, cancer, and the like. Exemplary monogenic disorders include ADA deficiency, cystic fibrosis, familial-hypercholesterolemia, hemophilia, chronic ganulomatous disease, Duchenne muscular dystrophy, Fanconi anemia, sickle-cell anemia, Gaucher's disease, Hunter syndrome, X-linked SCID, and the like. In another embodiment, the disorder treated involves one of the proteins listed below. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the recombinant protein encoded by an RNA, oligoribonucleotide, or polyribonucleotide molecule of methods and compositions of the present invention is ecto-nucleoside triphosphate diphosphohydrolase.

In another embodiment, the recombinant protein is erythropoietin (EPO). In other embodiments, the encoded recombinant protein is ABCA4; ABCD3; ACADM; AGL; AGT; ALDH4AI; ALPL; AMPD1; APOA2; AVSD1; BRCD2; C1QA; C1QB; C1QG; C8A; C8B; CACNA1S; CCV; CD3Z; CDC2L1; CHML; CHS1; CIAS1; CLCNKB; CMD1A; CMH2; CMM; COL11A1; COL8A2; COL9A2; CPT2; CRB1; CSE; CSF3R; CTPA; CTSK; DBT; DIO1; DISC1; DPYD;EKV; ENO1; ENO1P; EPB41; EPHX1; F13B; F5; FCGR2A; FCGR2B; FCGR3A; FCHL; FH; FMO3; FMO4; FUCA1; FY; GALE; GBA; GFND; GJA8; GJB3; GLC3B; HF1; HMGCL; HPC1; HRD; HRPT2; HSD3B2; HSPG2; KCNQ4; KCS; KIF1B; LAMB3; LAMC2; LGMD1B; LMNA; LOR; MCKD1; MCL1; MPZ; MTHFR; MTR; MUTYH;

MYOC; NB; NCF2; NEM1; NPHS2; NPPA; NRAS; NTRK1; OPTA2; PBX1; PCHC; PGD; PHA2A; PHGDH; PKLR; PKP1; PLA2G2A; PLOD; PPOX; PPT1; PRCC; PRG4; PSEN2; PTOS1; REN; RFX5; RHD; RMD1; RPE65; SCCD; SERPINC1; SJS1; SLC19A2; SLC2A1; SPG23; SPTA1; TAL1; TNFSF6; TNNT2; TPM3; TSHB; UMPK; UOX; UROD; USH2A; VMGLOM; VWS; WS2B; ABCB11; ABCG5; ABCG8; ACADL; ACP1; AGXT; AHHR; ALMS1; ALPP; ALS2; APOB; BDE; BDMR; BJS; BMPR2; CHRNA1; CMCWTD; CNGA3; COL3A1; COL4A3; COL4A4; COL6A3; CPSI; CRYGA; CRYGEP1; CYP1B1; CYP27A1; DBI; DES; DYSF; EDAR; EFEMPI; EIF2AK3; ERCC3; FSHR; GINGF; GLC1B; GPD2; GYPC; HADHA; HADHB; HOXD13; HPE2; IGKC; IHH; IRSI; ITGA6; KHK; KYNU; LCT; LHCGR; LSFC; MSH2; MSH6; NEB; NMTC; NPHP1; PAFAH1P1; PAX3; PAX8; PMS1; PNKD; PPH1; PROC; REG1A; SAG; SFTPB; SLC11A1; SLC3A1; SOS1; SPG4; SRD5A2; TCL4; TGFA; TMD; TPO; UGT1A@; UV24; WSS; XDH; ZAP70; ZFHX1B; ACAA1; AGS1; AGTR1; AHSG; AMT; ARMET; BBS3; BCHE; BCPM; BTD; CASR; CCR2; CCR5; CDL1; CMT2B; COL7A1; CP; CPO; CRV; CTNNB1; DEM; ETM1; FANCD2; FIH; FOXL2; GBE1; GLB1; GLC1C; GNAI2; GNATI; GP9; GPX1; HGD; HRG; ITIH1; KNG; LPP; LRS1; MCCCI; MDS1; MHS4; MITF; MLH1; MYL3; MYMY; OPA1; P2RY12; PBXP1; PCCB; POU1F1; PPARG; PROS1; PTHR1; RCA1; RHO; SCA7; SCLC1; SCN5A; SI; SLC25A20; SLC2A2; TF; TGFBR2; THPO; THRB; TKT; TM4SF1; TRH; UMPS; UQCRC1; USH3A; VHL; WS2A; XPC; ZNF35; ADH1B; ADH1C; AFP; AGA; AIH2; ALB; ASMD; BFHD; CNGA1; CRBM; DCK; DSPP; DTDP2; ELONG; ENAM; ETFDH; EVC; F11; FABP2; FGA; FGB; FGFR3; FGG; FSHMD1A; GC; GNPTA; GNRHR; GYPA; HCA; HCL2; HD; HTN3; HVBS6; IDUA; IF; JPD; KIT; KLKB1; LQT4; MANBA; MLLT2; MSX1; MTP; NR3C2; PBT; PDE6B; PEE1; PITX2; PKD2; QDPR; SGCB; SLC25A4; SNCA; SOD3; STATH; TAPVR1; TYS; WBS2; WFS1; WHCR; ADAMTS2; ADRB2; AMCN; AP3BI; APC; ARSB; B4GALT7; BHR1; C6; C7; CCAL2; CKN1; CMDJ; CRHBP; CSF1R; DHFR; DIAPH1; DTR; EOS; EPD; ERVR; F12; FBN2; GDNF; GHR; GLRA1; GM2A; HEXB; HSD17B4; ITGA2; KFS; LGMD1A; LOX; LTC4S; MAN2A1; MCC; MCCC2; MSH3; MSX2; NR3C1; PCSK1; PDE6A; PFBI; RASAI; SCZDI; SDHA; SGCD; SLC22A5; SLC26A2; SLC6A3; SM1; SMA@; SMN1; SMN2; SPINK5; TCOF1; TELAB1; TGFBI; ALDH5A1; ARG1; AS; ASSP2; BCKDHB; BF; C2; C4A; CDKN1A; COL10A1; COL11A2; CYP21A2; DYX2; EJM1; ELOVL4; EPM2A; ESR1; EYA4; F13A1; FANCE; GCLC; GJA1; GLYS1; GMPR; GSE; HCR; HFE; HLA-A; HLA-DPBI; HLA-

DRA; HPFH; ICS1; IDDM1; IFNGR1; IGAD1; IGF2R; ISCW; LAMA2; LAP; LCA5; LPA; MCDR1; MOCS1; MUT; MYB; NEU1; NKS1; NYS2; OA3; ODDD; OFC1; PARK2; PBCA; PBCRA1; PDB1; PEX3; PEX6; PEX7; PKHD1; PLA2G7; PLG; POLH; PPAC; PSORS1; PUJO; RCD1; RDS; RHAG; RP14; RUNX2; RWS; SCA1; SCZD3; SIASD; SOD2; ST8; TAP1; TAP2; TFAP2B; TNDM; TNF; TPBG; TPMT; TULP1; WISP3; AASS; ABCB1; ABCB4; ACHE; AQP1; ASL; ASNS; AUTS1; BPGM; BRAF; C7orf2; CACNA2D1; CCM1; CD36; CFTR; CHORDOMA; CLCN1; CMH6; CMT2D; COL1A2; CRS; CYMD; DFNA5; DLD; DYT11; EEC1; ELN; ETV1; FKBP6; GCK; GHRHR; GHS; GLI3; GPDS1; GUSB; HLXB9; HOXA13; HPFH2; HRX; IAB; IMMP2L; KCNH2; LAMB1; LEP; MET; NCF1; NM; OGDH; OPN1SW; PEX1; PGAM2; PMS2; PON1; PPP1R3A; PRSS1; PTC; PTPN12; RP10; RP9; SERPINE1; SGCE; SHFM1; SHH; SLC26A3; SLC26A4; SLOS; SMAD1; TBXAS1; TWIST; ZWS1; ACHM3; ADRB3; ANK1; CA1; CA2; CCAL1; CLN8; CMT4A; CNGB3; COH1; CPP; CRH; CYP11B1; CYP11B2; DECR1; DPYS; DURS1; EBS1; ECA1; EGI; EXT1; EYA1; FGFR1; GNRH1; GSR; GULOP; HR; KCNQ3; KFM; KWE; LGCR; LPL; MCPH1; MOS; MYC; NAT1; NAT2; NBS1; PLAT; PLEC1; PRKDC; PXMP3; RP1; SCZD6; SFIPC; SGM1; SPG5A; STAR; TG; TRPS1; TTPA; VMD1; WRN; ABCA1; ABL1; ABO; ADAMTS13; AK1; ALAD; ALDH1A1; ALDOB; AMBP; AMCD1; ASS; BDMF; BSCL; C5; CDKN2A; CHAC; CLA1; CMD1B; COL5A1; CRAT; DBH; DNAI1; DYS; DYT1; ENG; FANCC; FBP1; FCMD; FRDA; GALT; GLDC; GNE; GSM1; GSN; HSD17B3; HSN1; IBM2; INVS; JBTS1; LALL; LCCS1; LCCS; LGMD2H; LMX1B; MLLT3; MROS; MSSE; NOTCH1; ORM1; PAPPA; PIP5K1B; PTCH; PTGS1; RLN1; RLN2; RMRP; ROR2; RPD1; SARDH; SPTLC1; STOM; TDFA; TEK; TMC1; TRIM32; TSC1; TYRP1; XPA; CACNB2; COL17A1; CUBN; CXCL12; CYP17; CYP2C19; CYP2C9; EGR2; EMX2; ERCC6; FGFR2; HK1; HPS1; IL2RA; LGI1; LIPA; MAT1A; MBL2; MKI67; MXI1; NODAL; OAT; OATL3; PAX2; PCBD; PEO1; PHYH; PNLIP; PSAP; PTEN; RBP4; RDPA; RET; SFTPA1; SFTPD; SHFM3; SIAL; THC2; TLX1; TNFRSF6; UFS; UROS; AA; ABCC8; ACAT1; ALX4; AMPD3; ANC; APOA1; APOA4; APOC3; ATM; BSCL2; BWS; CALCA; CAT; CCND1; CD3E; CD3G; CD59; CDKN1C; CLN2; CNTF; CPT1A; CTSC; DDB1; DDB2; DHCR7; DLAT; DRD4; ECB2; ED4; EVR1; EXT2; F2; FSHB; FTH1; G6PT1; G6PT2; GIF; HBB; HBBP1; HBD; HBE1; HBG1; HBG2; HMBS; HND; HOMG2; HRAS; HVBS1; IDDM2; IGER; INS; JBS; KCNJ11; KCNJ1; KCNQ1; LDHA; LRP5; MEN1; MLL; MYBPC3; MYO7A; NNO1; OPPG; OPTB1; PAX6; PC;

PDX1; PGL2; PGR; PORC; PTH; PTS; PVRL1; PYGM; RAG1; RAG2; ROM1; RRAS2; SAA1; SCA5; SCZD2; SDHD; SERPING1; SMPD1; TCIRG1; TCL2; TECTA; TH; TREH; TSG101; TYR; USH1C; VMD2; VRNI; WT1; WT2; ZNF145; A2M; AAAS; ACADS; ACLS; ACVRL1; ALDH2; AMHR2; AOM; AQP2; ATD; ATP2A2; BDC; C1R; CD4; CDK4; CNA1; COL2A1; CYP27B1; DRPLA; ENUR2; FEOM1; FGF23; FPF; GNB3; GNS; HAL; HBP1; HMGA2; HMN2; HPD; IGF1; KCNA1; KERA; KRAS2; KRT1; KRT2A; KRT3; KRT4; KRT5; KRT6A; KRT6B; KRTHB6; LDHB; LYZ; MGCT; MPE; MVK; MYL2; OAP; PAH; PPKB; PRB3; PTPN11; PXR1; RLS; RSN; SAS; SAX1; SCA2; SCNN1A; SMAL; SPPM; SPSMA; TBX3; TBX5; TCF1; TPI1; TSC3; ULR; VDR; VWF; ATP7B; BRCA2; BRCD1; CLN5; CPB2; ED2; EDNRB; ENUR1; ERCC5; F10; F7; GJB2; GJB6; IPF1; MBS1; MCOR; NYS4; PCCA; RB1; RHOK; SCZD7; SGCG; SLC10A2; SLC25A15; STARP1; ZNF198; ACHM1; ARVD1; BCH; CTAA1; DAD1; DFNB5; EML1; GALC; GCH1; IBGC1; IGH@; IGHC group; IGHG1; IGHM; IGHR; IV; LTBP2; MCOP; MJD; MNG1; MPD1; MPS3C; MYH6; MYH7; NP; NPC2; PABPN1; PSEN1; PYGL; RPGRIP1; SERPINA1; SERPINA3; SERPINA6; SLC7A7; SPG3A; SPTB; TCL1A; TGMI; TITF1; TMIP; TRA@; TSHR; USH1A; VP; ACCPN; AHO2; ANCR; B2M; BBS4; BLM; CAPN3; CDAN1; CDAN3; CLN6; CMH3; CYP19; CYP1A1; CYP1A2; DYX1; EPB42; ETFA; EYCL3; FAH; FBN1; FES; HCVS; HEXA; IVD; LCS1; LIPC; MY05A; OCA2; OTSC1; PWCR; RLBP1; SLC12A1; SPG6; TPM1; UBE3A; WMS; ABCC6; ALDOA; APRT; ATP2A1; BBS2; CARD15; CATM; CDH1; CETP; CHST6; CLN3; CREBBP; CTH; CTM; CYBA; CYLD; DHS; DNASE1; DPEP1; ERCC4; FANCA; GALNS; GAN; HAGH; HBA1; HBA2; HBHR; HBQ1; HBZ; HBZP; HP; HSD11B2; IL4R; LIPB; MC1R; MEFV; MHC2TA; MLYCD; MMVP1; PHKB; PHKG2; PKD1; PKDTS; PMM2; PXE; SALL1; SCA4; SCNN1B; SCNN1G; SLC12A3; TAT; TSC2; VDI; WT3; ABR; ACACA; ACADVL; ACE; ALDH3A2; APOH; ASPA; AXIN2; BCL5; BHD; BLMH; BRCA1; CACD; CCA1; CCZS; CHRNB1; CHRNE; CMT1A; COL1A1; CORD5; CTNS; EPX; ERBB2; G6PC; GAA; GALK1; GCGR; GFAP; GH1; GH2; GP1BA; GPSC; GUCY2D; ITGA2B; ITGB3; ITGB4; KRT10; KRT12; KRT13; KRT14; KRT14L1; KRT14L2; KRT14L3; KRT16; KRT16L1; KRT16L2; KRT17; KRT9; MAPT; MDB; MDCR; MGI; MHS2; MKS1; MPO; MYO15A; NAGLU; NAPB; NF1; NME1; P4HB; PAFAH1B1; PECAM1; PEX12; PHB; PMP22; PRKAR1A; PRKCA; PRKWNK4; PRP8; PRPF8; PTLAH; RARA; RCV1; RMSA1; RP17; RSS; SCN4A; SERPINF2; SGCA; SGSH; SHBG; SLC2A4; SLC4A1; SLC6A4; SMCR;

SOST; SOX9; SSTR2; SYM1; SYNSI; TCF2; THRA; TIMP2; TOC; TOP2A; TP53; TRIM37; VBCH; ATP8B1; BCL2; CNSN; CORD11; CYB5; DCC; F5F8D; FECH; FEO; LAMA3; LCFS2; MADH4; MAFD1; MC2R; MCL; MYP2; NPC1; SPPK; TGFBRE; TGIF; TTR; AD2; AMH; APOC2; APOE; ATHS; BAX; BCKDHA; BCL3; BFIC; C3; CACNA1A; CCO; CEACAM5; COMP; CRX; DBA; DDU; DFNA4; DLL3; DM1; DMWD; E11S; ELA2; EPOR; ERCC2; ETFB; EXT3; EYCLI; FTL; FUT1; FUT2; FUT6; GAMT; GCDH; GPI; GUSM; HB1; HCL1; HHC2; HHC3; ICAM3; INSR; JAK3; KLK3; LDLR; LHB; LIG1; LOH19CR1; LYL1; MAN2B1; MCOLN1; MDRV; MLLT1; NOTCH3; NPHS1; OFC3; OPA3; PEPD; PRPF31; PRTN3; PRX; PSG1; PVR; RYR1; SLC5A5; SLC7A9; STK11; TBXA2R; TGFB1; TNNI3; TYROBP; ADA; AHCY; AVP; CDAN2; CDPD1; CHED1; CHED2; CHRNA4; CST3; EDN3; EEGV1; FTLL1; GDF5; GNAS; GSS; HNF4A; JAG1; KCNQ2; MKKS; NBIA1; PCK1; PI3; PPCD; PPGB; PRNP; THBD; TOP1; AIRE; APP; CBS; COL6A1; COL6A2; CSTB; DCR; DSCR1; FPDMM; HLCS; HPE1; ITGB2; KCNE1; KNO; PRSS7; RUNX1; SOD1; TAM; ADSL; ARSA; BCR; CECR; CHEK2; COMT; CRYBB2; CSF2RB; CTHM; CYP2D6;CYP2D7P1; DGCR; DIA1; EWSR1; GGT1; MGCR; MN1; NAGA; NE2; OGS2; PDGFB; PPARA; PRODH; SCO2; SCZD4; SERPIND1; SLC5A1; SOXI0; TCN2; TIMP3; TST; VCF; ABCD1; ACTL1; ADFN; AGMX2; AHDS; AIC; AIED; AIH3; ALAS2; AMCD; AMELX; ANOP1; AR; ARAF1; ARSC2; ARSE; ARTS; ARX; ASAT; ASSP5; ATP7A; ATRX; AVPR2; BFLS; BGN; BTK; BZX; C1HR; CACNA1F; CALB3; CBBM; CCT; CDR1; CFNS; CGF1; CHM; CHR39C; CIDX; CLA2; CLCN5; CLS; CMTX2; CMTX3; CND; COD1; COD2; COL4A5; COL4A6; CPX; CVD1; CYBB; DCX; DFN2; DFN4; DFN6; DHOF; DIAPH2; DKC1; DMD; DSS; DYT3; EBM; EBP; ED1; ELK1; EMD; EVR2; F8; F9; FCP1; FDPSL5; FGD1; FGS1; FMR1; FMR2; G6PD; GABRA3; GATA1; GDI1; GDXY; GJB1; GK; GLA; GPC3; GRPR; GTD; GUST; HMS1; HPRT1; HPT; HTC2; HTR2C; HYR; IDS; IHG1; IL2RG; INDX; IP1; IP2; JMS; KAL1; KFSD; LICAM; LAMP2; MAA; MAFD2; MAOA; MAOB; MCF2; MCS; MEAX; MECP2; MF4; MGCI; MIC5; MID1; MLLT7; MLS; MRSD; MRX14; MRX1; MRX20; MRX2; MRX3; MRX40; MRXA; MSD; MTMI; MYCL2; MYPI; NDP; NHS; NPHLI; NROBI; NSX; NYSI; NYX; OAI; OASD; OCRL; ODTI; OFD1; OPA2; OPD1; OPEM; OPN1LW; OPN1MW; OTC; P3; PDHA1; PDR; PFC; PFKFB1; PGK1; PGK1P1; PGS; PHEX; PHKA1; PHKA2; PHP; PIGA; PLP1; POF1; POLA; POU3F4; PPMX; PRD; PRPSI; PRPS2; PRS; RCCP2; RENBP; RENS1; RP2; RP6; RPGR; RPS4X; RPS6KA3;

RS1; S11; SDYS; SEDL; SERPINA7; SH2D1A; SHFM2; SLC25A5; SMAX2; SRPX; SRS; STS; SYN1; SYP; TAF1; TAZ; TBX22; TDD; TFE3; THAS; THC; TIMM8A; TIMP1; TKCR; TNFSF5; UBE1; UBE2A; WAS; WSN; WTS; WWS; XIC; XIST; XK; XM; XS; ZFX; ZIC3; ZNF261 ; ZNF41; ZNF6; AMELY; ASSP6; AZFI; AZF2; DAZ; GCY; RPS4Y; SMCY; SRY; ZFY; ABAT; AEZ; AFA; AFD1; ASAH1; ASD1; ASMT; CCAT; CECR9; CEPA; CLA3; CLN4; CSF2RA; CTSI; DF; DIH1; DWS; DYT2; DYT4; EBR3; ECT; EEF1A1L14; EYCL2; FANCB; GCSH; GCSL; GIP; GTS; HHG; HMI; HOAC; HOKPP2; HRPT1; HSD3B3; HTC1; HV1S; ICHQ; ICR1; ICR5; IL3RA; KAL2; KMS; KRT18; KSS; LCAT; LHON; LIMM; MANBB; MCPH2; MEB; MELAS; MIC2; MPFD; MS; MSS; MTATP6; MTCOI; MTCO3; MTCYB; MTND1; MTND2; MTND4; MTND5; MTND6; MTRNR1; MTRNR2; MTTE; MTTG; MTTI; MTTK; MTTL1; MTTL2; MTTN; MTTP; MTTS1; NAMSD; OCD1; OPD2; PCK2; PCLD; PCOS1; PFKM; PKD3; PRCA1; PRO1; PROP1; RBS; RFXAP; RP; SHOX; SLC25A6; SPG5B; STO; SUOX; THM; or TTD. Each recombinant protein represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for treating anemia in a subject, comprising contacting a cell of the subject with an *in vitro*-synthesized RNA molecule, the *in vitro* synthesized RNA molecule encoding erythropoietin, thereby treating anemia in a subject. In another embodiment, the *in vitro*-synthesized RNA molecule further comprises a pseudouridine or a modified nucleoside. Each possibility represents a separate embodiment of the present invention. In another embodiment, the cell is a subcutaneous tissue cell. In another embodiment, the cell is a lung cell. In another embodiment, the cell is a fibroblast. In another embodiment, the cell is a lymphocyte. In another embodiment, the cell is a smooth muscle cell. In another embodiment, the cell is any other type of cell known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for treating a vasospasm in a subject, comprising contacting a cell of the subject with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding inducible nitric oxide synthase (iNOS), thereby treating a vasospasm in a subject.

In another embodiment, the present invention provides a method for improving a survival rate of a cell in a subject, comprising contacting the cell with an *in vitro*-synthesized RNA

molecule, the *in vitro*-synthesized RNA molecule encoding a heat shock protein, thereby improving a survival rate of a cell in a subject.

In another embodiment, the cell whose survival rate is improved is an ischemic cell. In another embodiment, the cell is not ischemic. In another embodiment, the cell has been exposed to an ischemic environment. In another embodiment, the cell has been exposed to an environmental stress. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for decreasing an incidence of a restenosis of a blood vessel following a procedure that enlarges the blood vessel, comprising contacting a cell of the blood vessel with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a heat shock protein, thereby decreasing an incidence of a restenosis in a subject.

In another embodiment, the procedure is an angioplasty. In another embodiment, the procedure is any other procedure known in the art that enlarges the blood vessel. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for increasing a hair growth from a hair follicle is a scalp of a subject, comprising contacting a cell of the scalp with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a telomerase or an immunosuppressive protein, thereby increasing a hair growth from a hair follicle.

In another embodiment, the immunosuppressive protein is α-melanocyte-stimulating hormone (α-MSH). In another embodiment, the immunosuppressive protein is transforming growth factor-β 1 (TGF- β 1). In another embodiment, the immunosuppressive protein is insulin-like growth factor-I (IGF-I). In another embodiment, the immunosuppressive protein is any other immunosuppressive protein known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method of inducing expression of an enzyme with antioxidant activity in a cell, comprising contacting the cell with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding the enzyme, thereby inducing expression of an enzyme with antioxidant activity in a cell.

In one embodiment, the enzyme is catalase. In another embodiment, the enzyme is glutathione peroxidase. In another embodiment, the enzyme is phospholipid hydroperoxide glutathione peroxidase. In another embodiment, the enzyme is superoxide dismutase-1. In another embodiment, the enzyme is superoxide dismutase-2. In another embodiment, the enzyme is any other enzyme with antioxidant activity that is known in the art. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method for treating cystic fibrosis in a subject, comprising contacting a cell of the subject with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding Cystic Fibrosis Transmembrane Conductance Regulator (CFTR), thereby treating cystic fibrosis in a subject.

In another embodiment, the present invention provides a method for treating an X-linked agammaglobulinemia in a subject, comprising contacting a cell of the subject with an *in vitro* synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding a Bruton's tyrosine kinase, thereby treating an X-linked agammaglobulinemia.

In another embodiment, the present invention provides a method for treating an adenosine deaminase severe combined immunodeficiency (ADA SCID) in a subject, comprising contacting a cell of the subject with an *in vitro*-synthesized RNA molecule, the *in vitro*-synthesized RNA molecule encoding an ADA, thereby treating an ADA SCID.

In another embodiment, the present invention provides a method for reducing immune responsiveness of the skin and improving skin pathology, comprising contacting a cell of the subject with an *in vitro-synthesized* RNA molecule, the *in vitro*-synthesized RNA molecule encoding an ecto-nucleoside triphosphate diphosphohydrolase, thereby reducing immune responsiveness of the skin and improving skin pathology.

In another embodiment, an RNA molecule or ribonucleotide molecule of the present invention is encapsulated in a nanoparticle. Methods for nanoparticle packaging are well known in the art, and are described, for example, in Bose S, et al (Role of Nucleolin in Human Parainfluenza Virus Type 3 Infection of Human Lung Epithelial Cells. J. Virol. 78:8146. 2004); Dong Y et al. Poly(d,l-lactide-co-glycolide)/montmorillonite nanoparticles for oral delivery of anticancer drugs. Biomaterials 26:6068. 2005); Lobenberg R. et al (Improved body distribution of 14C-labelled AZT bound to nanoparticles in rats determined by radioluminography. J Drug Target 5:171. 1998); Sakuma SR et al (Mucoadhesion of polystyrene nanoparticles having

surface hydrophilic polymeric chains in the gastrointestinal tract. Int J Pharm 177:161. 1999); Virovic L et al. Novel delivery methods for treatment of viral hepatitis: an update. Expert Opin Drug Deliv 2:707.2005); and Zimmermann E et al., Electrolyte-and pH-stabilities of aqueous solid lipid nanoparticle (SLN) dispersions in artificial gastrointestinal media. Eur J Pharm Biopharm 52:203.2001). Each method represents a separate embodiment of the present invention.

Various embodiments of dosage ranges of compounds of the present invention can be used in methods of the present invention. In one embodiment, the dosage is in the range of 1-10 μg/day. In another embodiment, the dosage is 2-10 μg/day. In another embodiment, the dosage is 3-10 μg/day. In another embodiment, the dosage is 5-10 μg/day. In another embodiment, the dosage is 2-20 μg/day. In another embodiment, the dosage is 3-20 μg/day. In another embodiment, the dosage is 5-20 μg/day. In another embodiment, the dosage is 10-20 μg/day. In another embodiment, the dosage is 3-40 μg/day. In another embodiment, the dosage is 5-40 μg/day. In another embodiment, the dosage is 10-40 μg/day. In another embodiment, the dosage is 20-40 μg/day. In another embodiment, the dosage is 5-50 μg/day. In another embodiment, the dosage is 10-50 μg/day. In another embodiment, the dosage is 20-50 μg/day. In one embodiment, the dosage is 1-100 μg/day. In another embodiment, the dosage is 2-100 μg/day. In another embodiment, the dosage is 3-100 μg/day. In another embodiment, the dosage is 5-100 μg/day. In another embodiment the dosage is 10-100 μg/day. In another embodiment the dosage is 20-100 μg/day. In another embodiment the dosage is 40-100 μg/day. In another embodiment the dosage is 60-100 μg/day.

In another embodiment, the dosage is 0.1 μg/day. In another embodiment, the dosage is 0.2 μg/day. In another embodiment, the dosage is 0.3 μg/day. In another embodiment, the dosage is 0.5 μg/day. In another embodiment, the dosage is 1 μg/day. In another embodiment, the dosage is 2 mg/day. In another embodiment, the dosage is 3 μg/day. In another embodiment, the dosage is 5 μg/day. In another embodiment, the dosage is 10 μg/day. In another embodiment, the dosage is 15 μg/day. In another embodiment, the dosage is 20 μg/day. In another embodiment, the dosage is 30 μg/day. In another embodiment, the dosage is 40 μg/day. In another embodiment, the dosage is 60 μg/day. In another embodiment, the dosage is 80 μg/day. In another embodiment, the dosage is 100 μg/day.

In another embodiment, the dosage is 10 μg/dose. In another embodiment, the dosage is 20 μg/dose. In another embodiment, the dosage is 30 μg/dose. In another embodiment, the dosage is 40 μg/dose. In another embodiment, the dosage is 60 μg/dose. In another embodiment, the dosage is 80 μg/dose. In another embodiment, the dosage is 100 μg/dose. In another embodiment, the dosage is 150 μg/dose. In another embodiment, the dosage is 200 μg/dose. In another embodiment, the dosage is 300 μg/dose. In another embodiment, the dosage is 400 μg/dose. In another embodiment, the dosage is 600 μg/dose. In another embodiment, the dosage is 800 μg/dose. In another embodiment, the dosage is 1000 μg/dose. In another embodiment, the dosage is 1.5 mg/dose. In another embodiment, the dosage is 2 mg/dose. In another embodiment, the dosage is 3 mg/dose. In another embodiment, the dosage is 5 mg/dose. In another embodiment, the dosage is 10 mg/dose. In another embodiment, the dosage is 15 mg/dose. In another embodiment, the dosage is 20 mg/dose. In another embodiment, the dosage is 30 mg/dose. In another embodiment, the dosage is 50 mg/dose. In another embodiment, the dosage is 80 mg/dose. In another embodiment, the dosage is 100 mg/dose.

In another embodiment, the dosage is 10-20 μg/dose. In another embodiment, the dosage is 20-30 μg/dose. In another embodiment, the dosage is 20-40 μg/dose. In another embodiment, the dosage is 30-60 μg/dose. In another embodiment, the dosage is 40-80 μg/dose. In another embodiment, the dosage is 50-100 μg/dose. In another embodiment, the dosage is 50-150 μg/dose. In another embodiment, the dosage is 100-200 μg/dose. In another embodiment, the dosage is 200-300 μg/dose. In another embodiment, the dosage is 300-400 μg/dose. In another embodiment, the dosage is 400-600 μg/dose. In another embodiment, the dosage is 500-800 μg/dose. In another embodiment, the dosage is 800-1000 μg/dose. In another embodiment, the dosage is 1000-1500 μg/dose. In another embodiment, the dosage is 1500-2000 μg/dose. In another embodiment, the dosage is 2-3 mg/dose. In another embodiment, the dosage is 2-5 mg/dose. In another embodiment, the dosage is 2-10 mg/dose. In another embodiment, the dosage is 2-20 mg/dose. In another embodiment, the dosage is 2-30 mg/dose. In another embodiment, the dosage is 2-50 mg/dose. In another embodiment, the dosage is 2-80 mg/dose. In another embodiment, the dosage is 2-100 mg/dose. In another embodiment, the dosage is 3-10 mg/dose. In another embodiment, the dosage is 3-20 mg/dose. In another embodiment, the dosage is 3-30 mg/dose. In another embodiment, the dosage is 3-50 mg/dose. In another embodiment, the dosage is 3-80 mg/dose. In another embodiment, the dosage is 3-100 mg/dose.

In another embodiment, the dosage is 5-10 mg/dose. In another embodiment, the dosage is 5-20 mg/dose. In another embodiment, the dosage is 5-30 mg/dose. In another embodiment, the dosage is 5-50 mg/dose. In another embodiment, the dosage is 5-80 mg/dose. In another embodiment, the dosage is 5-100 mg/dose. In another embodiment, the dosage is 10-20 mg/dose. In another embodiment, the dosage is 10-30 mg/dose. In another embodiment, the dosage is 10-50 mg/dose. In another embodiment, the dosage is 10-80 mg/dose. In another embodiment, the dosage is 10-100 mg/dose.

In another embodiment, the dosage is a daily dose. In another embodiment, the dosage is a weekly dose. In another embodiment, the dosage is a monthly dose. In another embodiment, the dosage is an annual dose. In another embodiment, the dose is one is a series of a defined number of doses. In another embodiment, the dose is a one-time dose. As described below, in another embodiment, an advantage of RNA, oligoribonucleotide, or polyribonucleotide molecules of the present invention is their greater potency, enabling the use of smaller doses.

In another embodiment, the present invention provides a method for producing a recombinant protein, comprising contacting an *in vitro* translation apparatus with an *in vitro*-synthesized oligoribonucleotide, the *in vitro*-synthesized oligoribonucleotide comprising a pseudouridine or a modified nucleoside, thereby producing a recombinant protein.

In another embodiment, the present invention provides a method for producing a recombinant protein, comprising contacting an *in vitro* translation apparatus with an *in vitro*-transcribed RNA molecule of the present invention, the *in vitro*-transcribed RNA molecule comprising a pseudouridine or a modified nucleoside, thereby producing a recombinant protein.

In another embodiment, the present invention provides an *in vitro* transcription apparatus, comprising: an unmodified nucleotide, a nucleotide containing a pseudouridine or a modified nucleoside, and a polymerase. In another embodiment, the present invention provides an *in vitro* transcription kit, comprising: an unmodified nucleotide, a nucleotide containing a pseudouridine or a modified nucleoside, and a polymerase. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the *in vitro* translation apparatus comprises a reticulocyte lysate. In another embodiment, the reticulocyte lysate is a rabbit reticulocyte lysate.

In another embodiment, the present invention provides a method of reducing an immunogenicity of an oligoribonucleotide molecule or RNA molecule, the method comprising

the step of replacing a nucleotide of the oligoribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby reducing an immunogenicity of an oligoribonucleotide molecule or RNA molecule.

In another embodiment, the present invention provides a method of reducing an immunogenicity of a gene-therapy vector comprising a polyribonucleotide molecule or RNA molecule, the method comprising the step of replacing a nucleotide of the polyribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby reducing an immunogenicity of a gene-therapy vector.

In another embodiment, the present invention provides a method of enhancing *in vitro* translation from an oligoribonucleotide molecule or RNA molecule, the method comprising the step of replacing a nucleotide of the oligoribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby enhancing *in vitro* translation from an oligoribonucleotide molecule or RNA molecule.

In another embodiment, the present invention provides a method of enhancing *in vivo* translation from a gene-therapy vector comprising a polyribonucleotide molecule or RNA molecule, the method comprising the step of replacing a nucleotide of the polyribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby enhancing *in vivo* translation from a gene-therapy vector.

In another embodiment, the present invention provides a method of increasing efficiency of delivery of a recombinant protein by a gene therapy vector comprising a polyribonucleotide molecule or RNA molecule, the method comprising the step of replacing a nucleotide of the polyribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby increasing efficiency of delivery of a recombinant protein by a gene therapy vector.

In another embodiment, the present invention provides a method of increasing *in vivo* stability of gene therapy vector comprising a polyribonucleotide molecule or RNA molecule, the method comprising the step of replacing a nucleotide of the polyribonucleotide molecule or RNA molecule with a modified nucleotide that contains a modified nucleoside or a pseudouridine, thereby increasing *in vivo* stability of gene therapy vector.

In another embodiment, the present invention provides a method of synthesizing an *in vitro*-transcribed RNA molecule comprising a pseudouridine nucleoside, comprising contacting

an isolated polymerase with a mixture of unmodified nucleotides and the modified nucleotide (Examples 5 and 10).

In another embodiment, *in vitro* transcription methods of the present invention utilize an extract from an animal cell. In another embodiment, the extract is from a reticulocyte or cell with similar efficiency of in vitro transcription. In another embodiment, the extract is from any other type of cell known in the art. Each possibility represents a separate embodiment of the present invention.

Any of the RNA molecules or oligoribonucleotide molecules of the present invention may be used, in another embodiment, in any of the methods of the present invention.

In another embodiment, the present invention provides a method of enhancing an immune response to an antigen, comprising administering the antigen in combination with mitochondrial (mt) RNA (Examples 4 and 8).

In another embodiment, the present invention provides a method of reducing the ability of an RNA molecule to stimulate a dendritic cell (DC), comprising modifying a nucleoside of the RNA molecule by a method of the present invention (e.g., see EXAMPLES).

In another embodiment, the DC is a DC1 cell. In another embodiment, the DC is a DC2 cell. In another embodiment, the DC is a subtype of a DC1 cell or DC2 cell. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a method of reducing the ability of an RNA molecule to stimulate signaling by TLR3, comprising modifying a nucleoside of the RNA molecule by a method of the present invention. In another embodiment, the present invention provides a method of reducing the ability of an RNA molecule to stimulate signaling by TLR7, comprising modifying a nucleoside of the RNA molecule by a method of the present invention. In another embodiment, the present invention provides a method of reducing the ability of an RNA molecule to stimulate signaling by TLR8, comprising modifying a nucleoside of the RNA molecule by a method of the present invention. Each possibility represents a separate embodiment of the present invention.

In another embodiment, all of the internucleoside or internucleotide linkages in the RNA, oligoribonucleotide, or polyribonucleotide molecule are phosphodiester. In another embodiment, the inter-nucleotide linkages are predominantly phosphodiester. In another embodiment, most of the internucleotide linkages are phosphorothioate. In another embodiment, most the inter-

nucleotide linkages are phosphodiester. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the percentage of the inter-nucleotide linkages in the RNA, oligoribonucleotide, or polyribonucleotide molecule that are phosphodiester is above 50%. In another embodiment, the percentage is above 10%. In another embodiment, the percentage is above 15%. In another embodiment, the percentage is above 20%. In another embodiment, the percentage is above 25%. In another embodiment, the percentage is above 30%. In another embodiment, the percentage is above 35%. In another embodiment, the percentage is above 40%. In another embodiment, the percentage is above 45%. In another embodiment, the percentage is above 55%. In another embodiment, the percentage is above 60%. In another embodiment, the percentage is above 65%. In another embodiment, the percentage is above 70%. In another embodiment, the percentage is above75%. In another embodiment, the percentage is above 80%.In another embodiment, the percentage is above 85%. In another embodiment, the percentage is above 90%. In another embodiment, the percentage is above 95%.

In another embodiment, a method of the present invention comprises increasing the number, percentage, or frequency of modified nucleosides in the RNA molecule to decrease immunogenicity or increase efficiency of translation. As provided herein (e.g., see EXAMPLES), the number of modified residues in an RNA, oligoribonucleotide, or polyribonucleotide molecule determines, in another embodiment, the magnitude of the effects observed in the present invention.

In another embodiment, the present invention provides a method for introducing a recombinant protein into a cell of a subject, comprising contacting the subject with an *in vitro*-transcribed RNA molecule encoding the recombinant protein, the *in vitro*-transcribed RNA molecule further comprising a pseudouridine or another modified nucleoside, thereby introducing a recombinant protein into a cell of a subject.

In another embodiment, the present invention provides a method for decreasing TNF-α production in response to a gene therapy vector in a subject, comprising the step of engineering the vector to contain a pseudouridine or a modified nucleoside base, thereby decreasing TNF-α production in response to a gene therapy vector in a subject.

In another embodiment, the present invention provides a method for decreasing IL-12 production in response to a gene therapy vector in a subject, comprising the step of engineering the vector to contain a pseudouridine or a modified nucleoside base, thereby decreasing IL-12 production in response to a gene therapy vector in a subject.

In another embodiment, the present invention provides a method of reducing an immunogenicity of a gene therapy vector, comprising introducing a modified nucleoside into said gene therapy vector, thereby reducing an immunogenicity of a gene therapy vector.

As provided herein, findings of the present invention show that primary DC have an additional RNA signaling entity that recognizes m5C-and m6A-modified RNA and whose signaling is inhibited by modification of U residues.

In another embodiment, an advantage of an RNA, oligoribonucleotide, and polyribonucleotide molecules of the present invention is that RNA does not incorporate to the genome (as opposed to DNA-based vectors). In another embodiment, an advantage is that translation of RNA, and therefore appearance of the encoded product, is instant. In another embodiment, an advantage is that the amount of protein generated from the mRNA can be regulated by delivering more or less RNA. In another embodiment, an advantage is that repeated delivery of purified pseudouridine or other modified RNA, oligoribnucleotide, or polyribonucleotide molecules does not induce an immune response, whereas repeated delivery of unmodified RNA could induce signaling pathways though RNA sensors.

In another embodiment, an advantage is lack of immunogenicity, enabling repeated delivery without generation of inflammatory cytokines. In another embodiment, stability of RNA is increased by circularization, decreasing degradation by exonucleases.

In another embodiment, the present invention provides a method of treating a subject with a disease that comprises an immune response against a self-RNA molecule, comprising administering to the subject an antagonist of a TLR-3 molecule, thereby treating a subject with a disease that comprises an immune response against a self-RNA molecule.

In another embodiment, the present invention provides a method of treating a subject with a disease that comprises an immune response against a self-RNA molecule, comprising administering to the subject an antagonist of a TLR-7 molecule, thereby treating a subject with a disease that comprises an immune response against a self-RNA molecule.

In another embodiment, the present invention provides a method of treating a subject with a disease that comprises an immune response against a self-RNA molecule, comprising administering to the subject an antagonist of a TLR-8 molecule, thereby treating a subject with a disease that comprises an immune response against a self-RNA molecule.

In another embodiment, the disease that comprises an immune response against a self-RNA molecule is an auto-immune disease. In another embodiment, the disease is systemic lupus erythematosus (SLE). In another embodiment, the disease is another disease known in the art that comprises an immune response against a self-RNA molecule. Each possibility represents a separate embodiment of the present invention.

In another embodiment, the present invention provides a kit comprising a reagent utilized in performing a method of the present invention. In another embodiment, the present invention provides a kit comprising a composition, tool, or instrument of the present invention.

In another embodiment, the present invention provides a kit for measuring or studying signaling by a TLR-3, TLR-7 and TLR-8 receptor, as exemplified in Example 7.

In another embodiment, a treatment protocol of the present invention is therapeutic. In another embodiment, the protocol is prophylactic. Each possibility represents a separate embodiment of the present invention.

In one embodiment, the phrase "contacting a cell" or "contacting a population" refers to a method of exposure, which can be direct or indirect. In one method such contact comprises direct injection of the cell through any means well known in the art, such as microinjection. In another embodiment, supply to the cell is indirect, such as via provision in a culture medium that surrounds the cell, or administration to a subject, or via any route known in the art. In another embodiment, the term "contacting" means that the molecule of the present invention is introduced into a subject receiving treatment, and the molecule is allowed to come in contact with the cell *in vivo*. Each possibility represents a separate embodiment of the present invention.

Methods for quantification of reticulocyte frequency and for measuring EPO biological activity are well known in the art, and are described, for Example, in Ramos, AS et al (Biological evaluation of recombinant human erythropoietin in pharmaceutical products. Braz J Med Biol Res 36:1561). Each method represents a separate embodiment of the present invention.

Compositions of the present invention can be, in another embodiment, administered to a subject by any method known to a person skilled in the art, such as parenterally, paracancerally,

transmucosally, transdermally, intramuscularly, intravenously, intra-dermally, subcutaneously, intraperitoneally, intraventricularly, intra-cranially, intra-vaginally or intra-tumorally.

In another embodiment of methods and compositions of the present invention, the compositions are administered orally, and are thus formulated in a form suitable for oral administration, i.e. as a solid or a liquid preparation. Suitable solid oral formulations include tablets, capsules, pills, granules, pellets and the like. Suitable liquid oral formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In another embodiment of the present invention, the active ingredient is formulated in a capsule. In accordance with this embodiment, the compositions of the present invention comprise, in addition to the active compound and the inert carrier or diluent, a hard gelating capsule.

In other embodiments, the pharmaceutical compositions are administered by intravenous, intra-arterial, or intra-muscular injection of a liquid preparation. Suitable liquid formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In another embodiment, the pharmaceutical compositions are administered intravenously and are thus formulated in a form suitable for intravenous administration. In another embodiment, the pharmaceutical compositions are administered intra-arterially and are thus formulated in a form suitable for intra-arterial administration. In another embodiment, the pharmaceutical compositions are administered intra-muscularly and are thus formulated in a form suitable for intra-muscular administration.

In another embodiment, the pharmaceutical compositions are administered topically to body surfaces and are thus formulated in a form suitable for topical administration. Suitable topical formulations include gels, ointments, creams, lotions, drops and the like. For topical administration, the compositions or their physiologically tolerated derivatives are prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a pharmaceutical carrier.

In another embodiment, the composition is administered as a suppository, for example a rectal suppository or a urethral suppository. In another embodiment, the pharmaceutical composition is administered by subcutaneous implantation of a pellet. In another embodiment, the pellet provides for controlled release of agent over a period of time.

In another embodiment, the active compound is delivered in a vesicle, e.g. a liposome (see Langer, Science 249: 1527-1533 (1990); Treat et al., in Liposomes in the Therapy of

Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, ibid., pp. 317-327; see generally ibid).

As used herein "pharmaceutically acceptable carriers or diluents" are well known to those skilled in the art. The carrier or diluent may be may be, in various embodiments, a solid carrier or diluent for solid formulations, a liquid carrier or diluent for liquid formulations, or mixtures thereof.

In another embodiment, solid carriers/diluents include, but are not limited to, a gum, a starch (e.g. com starch, pregeletanized starch), a sugar (e.g., lactose, mannitol, sucrose, dextrose), a cellulosic material (e.g. microcrystalline cellulose), an acrylate (e.g. polymethylacrylate), calcium carbonate, magnesium oxide, talc, or mixtures thereof.

In other embodiments, pharmaceutically acceptable carriers for liquid formulations may be aqueous or non-aqueous solutions, suspensions, emulsions or oils. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Examples of oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, mineral oil, olive oil, sunflower oil, and fish-liver oil.

Parenteral vehicles (for subcutaneous, intravenous, intra-arterial, or intramuscular injection) include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's and fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose, and the like. Examples are sterile liquids such as water and oils, with or without the addition of a surfactant and other pharmaceutically acceptable adjuvants. In general, water, saline, aqueous dextrose and related sugar solutions, and glycols such as propylene glycols or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions. Examples of oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, mineral oil, olive oil, sunflower oil, and fish-liver oil.

In another embodiment, the compositions further comprise binders (e.g. acacia, cornstarch, gelatin, carbomer, ethyl cellulose, guar gum, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, povidone), disintegrating agents (e.g. cornstarch, potato starch, alginic acid, silicon dioxide, croscarmelose sodium, crospovidone, guar gum, sodium starch glycolate),

buffers (e.g., Tris-HCI., acetate, phosphate) of various pH and ionic strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), protease inhibitors, surfactants (e.g. sodium lauryl sulfate), permeation enhancers, solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite, butylated hydroxyanisole), stabilizers (e.g. hydroxypropyl cellulose, hyroxypropylmethyl cellulose), viscosity increasing agents(e.g. carbomer, colloidal silicon dioxide, ethyl cellulose, guar gum), sweeteners (e.g. aspartame, citric acid), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), lubricants (e.g. stearic acid, magnesium stearate, polyethylene glycol, sodium lauryl sulfate), flow-aids (e.g. colloidal silicon dioxide), plasticizers (e.g. diethyl phthalate, triethyl citrate), emulsifiers (e.g. carbomer, hydroxypropyl cellulose, sodium lauryl sulfate), polymer coatings (e.g., poloxamers or poloxamines), coating and film forming agents (e.g. ethyl cellulose, acrylates, polymethacrylates) and/or adjuvants. Each of the above excipients represents a separate embodiment of the present invention.

In another embodiment, the pharmaceutical compositions provided herein are controlled-release compositions, i.e. compositions in which the compound is released over a period of time after 15 administration. Controlled-or sustained-release compositions include formulation in lipophilic depots (e.g. fatty acids, waxes, oils). In another embodiment, the composition is an immediate-release composition, i.e. a composition in which the entire compound is released immediately after administration.

In another embodiment, molecules of the present invention are modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or polyproline. The modified compounds are known to exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds (Abuchowski et al., 1981; Newmark et al., 1982; and Katre et al., 1987). Such modifications also increase, in another embodiment, the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the immunogenicity and reactivity of the compound. As a result, the desired *in vivo* biological activity may be achieved by the administration of such polymer-compound abducts less frequently or in lower doses than with the unmodified compound.

An active component is, in another embodiment, formulated into the composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (e.g., formed with the free amino groups of a polypeptide or antibody molecule), which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

Each of the above additives, excipients, formulations and methods of administration represents a separate embodiment of the present invention.

EXAMPLES

The following experimental protocols were employed in the Examples provided below, unless indicated otherwise.

MATERIALS AND METHODS FOR EXAMPLES 1-3

Cell Culture. Newborn human foreskin fibroblast 1079 cells (Cat# CRL-2097, ATCC, Manassas, VA) and human IMR90 cells (Cat# CCL-186, ATCC) were cultured in Advanced MEM Medium (Invitrogen, Carlsbad, CA) supplemented with 10% heat-inactivated fetal bovine serum (FBS, Hyclone Laboratories, Logan, UT), 2mM Glutamax (Invitrogen), 0.1mM β-mercaptoethanol (Sigma, St. Louis, MO), and Penicillin/Streptomycin (Invitrogen). All cells were grown at 37^0C and 5% CO_2. In some experiments, human iPS cells that were induced using methods described herein were maintained on irradiated mouse embryonic fibroblasts (MEFs) (R&D Systems, Minneapolis, MN) on 10-cm plates pre-coated with 0.1% gelatin (Millipore, Phillipsburg, NJ) in DMEM/F12 medium supplemented with 20% KnockOut serum replacer, 0.1mM L-glutamine (all from Invitrogen), 0.1mM β-mercaptoethanol (Sigma) and 100ng/ml basic fibroblast growth factor (Invitrogen). In some experiments, human iPS cells that were induced using methods described herein were maintained in MEF-conditioned medium that had been collected as previously described (Xu et al. 2001).

Constructions of Vectors. The cDNAs for the open reading frames (ORFs) of KLF4, LIN28, NANOG, and OCT4 were PCR amplified from cDNA clones (Open Biosystems,

Huntsville, AL), cloned into a plasmid vector downstream of a T7 RNA polymerase promoter (Mackie 1988, Studier and Moffatt 1986) (e.g., various pBluescript™, Agilent, La Jolla, CA or pGEM™, Promega, Madison, WI, vectors) and sequenced. The ORF of SOX2 was PCR amplified from a cDNA clone (Invitrogen) and the ORF of c-MYC was isolated by RT-PCR from HeLa cell total RNA. Both SOX2 and c-MYC ORF were also cloned into a plasmid vector downstream of a T7 RNA polymerase promoter and sequenced.

Alternative plasmid vectors containing human open reading frames of (KLF4, LIN28, c-MYC, NANOG, OCT4 and SOX2) were cloned into pBluescriptII. These pBluescriptII vectors where constructed by ligating the above open reading frames into the EcoRV (cMyc) or EcoRV/SpeI (KLF4, LIN28, NANOG, OCT4, and SOX2) sites between the 5' and 3' Xenopus laevis beta-globin untranslated regions described (Krieg and Melton 1984).

mRNA Production. The T7 RNA polymerase promoter-containing plasmid contructs (pT7-KLF4, pT7-LIN28, pT7-c-MYC, pT7-OCT4, pT7-SOX2, or pT7-XBg-KLF4, pT7-XBg-LIN28, pT7-XBg-c-MYC, pT7-XBg-OCT4, and pT7-XBg-SOX2) were linearized with BamHI and pT7-NANOG and pT7-XBg-NANOG were linearized with Xba I. The mSCRIPT™ mRNA production system (EPICENTRE or CellScript, Madison, WI, USA) was used to produce mRNA with a 5' Cap1 structure and a 3' Poly (A) tail (e.g., with approximately 150 A residues), except that pseudouridine-5'-triphosphate (TRILINK, San Diego, CA) was used in place of uridine-5'-triphosphate in the T7 RNA polymerase in vitro transcription reactions.

mRNA Purification and Analysis. In some experimental embodiments, the mRNA was purified by HPLC, column fractions were collected, and the mRNA fractions were analyzed for purity an immunogenicity as described in "Materials and Methods for Examples 35-38" and/or as described and shown for Figures 22-24. In some experimental embodiments, purified RNA preparations comprising or consisting of mRNAs encoding one or more reprogramming factors which exhibited little or no immunogenicity were used for the experiments for reprogramming human somatic cells to iPS cells.

Reprogramming of Human Somatic Cells on MEFs. 1079 fibroblasts were plated at 1 x 10^5 cells/well of a 6-well dish pre-coated with 0.1% gelatin (Millipore) and grown overnight. The 1079 fibroblasts were transfected with equal amounts of each reprogramming factor mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) using TransIT mRNA transfection reagent (MirusBio, Madison, WI). A total of three transfections were performed, with one transfection

being performed every other day, with media changes the day after the first and second transfection. The day after the third transfection, the cells were trypsinized and 3.3 x 10^5 cells were plated in 1079 medium onto 0.1% gelatin pre-coated 10-cm plate seeded with 7.5 x 10^5 MEFs the day before. The day after plating the transfected 1079 fibroblasts onto MEFs, the medium was changed to iPS cell medium. The iPS cell medium was changed every day. Eight days after plating the transfected cells onto MEFs, MEF-conditioned medium was used. MEF conditioned medium was collected as previously described (Xu et al. 2001). Plates were screened every day for the presence of colonies with an iPS morphology using an inverted microscope.

Alternative protocols for reprogramming 1079 and IMR90 fibroblasts on MEFs were also used. MEFs were plated at 1.25 x 10^5 cells/well of a 0.1% gelatin pre-coated 6 well dish and incubated overnight in complete fibroblast media. 1079 or IMR90 fibroblasts were plated at 3 x 10^4 cells/well of a 6 well dish seeded with MEFs the previous day and grown overnight at 37°C/5%CO_2. The mScript Kit was then used to generate Cap1/poly-adenylated mRNA from the following vectors (pT7-Xβg-KLF4, pT7-Xβg-LIN28, pT7-Xβg-c-MYC, pT7-Xβg-NANOG, pT7-Xβg-OCT4, and pT7-Xβg-SOX2) for use in these daily transfections. All six reprogramming mRNAs were diluted to 100 ng/μl of each mRNA. Equal molarity of each mRNA was added together using the following conversion factors (OCT4 is set at 1 and all of the other mRNAs are multiplied by these conversion factors to obtain equal molarity in each mRNA mix). KLF=1.32, LIN28=0.58, c-MYC=1.26, NANOG=0.85, OCT4=1, and SOX2=0.88. To obtain equal molarity of each factor 132 μl of KLF4, 58 μl of LIN28, 126 μl of c-MYC, 85 μl of NANOG, 100 μl of OCT4 and 88 μl of SOX2 mRNA (each at 100ng/μl) would be added together. A 600μg total dose for transfections would mean that 100ng (using molarity conversions above) of each of six reprogramming mRNAs was used. Trans-IT mRNA transfection reagent was used to transfect these mRNA doses. For all transfections, mRNA pools were added to 250μl of either DMEM/F12 media without additives or Advanced MEM media without additives. 5μl of mRNA boost reagent and 5μl of TransIT transfection reagent was added to each tube and incubated at room temp for two minutes before adding the transfection mix to 2.5mls of either Advanced MEM media with 10% FBS + 100ng/ml of hFGFb or iPS media containing 100ng/ml of hFGFb. Transfections were repeated everyday for 10-16 days. The media was changed 4 hours after each transfection. In some experiments, the cells were

trypsinized and replated onto new MEF plates between 5-8 days after the inititial transfection. 1079 cells were split 1/6 or 1/12 onto new MEF plates while IMR90 cells were split 1/3 or 1/6 onto new MEF plates.

Reprogramming of Human Somatic Cells in MEF-Conditioned Medium. 1079 or IMR90 fibroblasts were plated at 3 x 10^5 cells per 10cm dishes pre-coated with 0.1% gelatin (Millipore) and grown overnight. The 1079 or IMR90 fibroblasts were transfected with equal amounts of reprogramming factor mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) using TransIT mRNA transfection reagent (MirusBio, Madison, WI). For each transfection, either 6 μg, 18 μg, or 36 μg of each reprogramming mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) was used per 10-cm dish. A total of three transfections were performed, with one transfection being performed every other day with the medium being changed the day after each of the first and second transfections. All transfections were performed in MEF-conditioned medium. The day after the third transfection, the cells were trypsinized and 3 x 10^5 cells were plated on new 10-cm dishes pre-coated with 0.1% gelatin (Millipore). The cells were grown in MEF-conditioned medium for the duration of the experiment.

Similar daily mRNA transfections were also performed as described in the previous section with the only difference being that MEFs were not used as feeder layers, only MEF conditioned media was used.

Immunoflourescence. The 1079 cells or 1079-derived iPS cell plates were washed with PBS and fixed in 4% paraformaldehyde in PBS for 30 minutes at room temperature. The iPS cells were then washed 3 times for 5 minutes each wash with PBS followed by three washes in PBS + 0.1% Triton X-100. The iPS cells were then blocked in blocking buffer (PBS + 0.1% Triton, 2% FBS, and 1% BSA) for 1 hour at room temperature. The cells were then incubated for 2 hours at room temperature with the primary antibody (mouse anti-human OCT4 Cat# sc-5279, Santa Cruz Biotechnology, Santa Cruz, CA), (rabbit anti-human NANOG Cat #3580, rabbit anti-human KLF4 Cat # 4038, mouse anti-human LIN28 Cat# 5930, rabbit anti-human c-MYC Cat# 5605, rabbit anti-human SOX2 Cat# 3579, and mouse anti-TRA-1-60 all from Cell Signaling Technology, Beverly, MA) at a 1:500 dilution in blocking buffer. After washing 5 times in PBS + 0.1% Triton X-100, the iPS cells were incubated for 2 hours with the anti-rabbit Alexa Fluor 488 antibody (Cat # 4412, Cell Signaling Technology), anti-mouse FITC secondary (Cat# F5262, Sigma), or an anti-mouse Alexa Fluor 555 (Cat# 4409, Cell Signaling Technology)

at 1:1000 dilutions in blocking buffer. Images were taken on a Nikon TS100F inverted microscope (Nikon, Tokyo, Japan) with a 2-megapixel monochrome digital camera (Nikon) using NIS-elements software (Nikon).

EXAMPLE 1

This Example describes tests to determine if transfections with mRNA encoding KLF4, LIN28, c-MYC, NANOG, OCT4 and SOX2 resulted in expression and proper subcellular localization of each respective protein product in newborn fetal foreskin 1079 fibroblasts. The mRNAs used in the experiments were made with pseudouridine-5'-triphosphate substituting for uridine-5'-triphosphate (Kariko et al. 2008). The 1079 fibroblasts were transfected with 4 μg of each mRNA per well of a 6-well dish and immunofluorescence analysis was performed 24 hours post-transfection. Endogenous KLF4, LIN28, NANOG, OCT4 and SOX2 protein levels were undetectable by immunoflourescence in untransfected 1079 cells (Fig.1: B, F, N, R, V). Endogenous levels of c-MYC were relatively high in untransfected 1079 cells (Fig. 1J). Transfections with mRNAs encoding the transcription factors, KLF4, c-MYC, NANOG, OCT4, and SOX2 all resulted in primarily nuclear localization of each protein 24 hours after mRNA transfections (Figure 1: D, L, P, T, X). The cytoplasmic mRNA binding protein, LIN28, was localized to the cytoplasm (Fig. 1: H).

EXAMPLE 2

Having demonstrated efficient mRNA transfection and proper subcellular localization of the reprogramming proteins, this Example describes development of a protocol for iPS cell generation from somatic fibroblasts. Equal amounts (by weight) of KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2 mRNAs were transfected into 1079 fibroblasts three times (once every other day). The day after the third transfection, the cells were plated onto irradiated MEF feeder cells and grown in iPS cell medium. Six days after plating the 1079 fibroblasts onto irradiated MEFs, two putative iPS cell colonies became apparent on the 10-cm plate transfected with 3 μg of each reprogramming factor mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2). The colonies were allowed to grow until 12 days after the last transfection before they were fixed for immunofluorescence analysis. The inner cell mass-specific marker NANOG is often used to assay whether iPS cell colonies are truly iPS colonies (Gonzalez et al. 2009,

Huangfu et al. 2008). NANOG expression arising from the mRNAs that were transfected 12 days earlier would be negligible based on previous reports on the duration of mRNA stability and expression (Kariko et al. 2008). Staining for NANOG showed that both of the two iPS cell colonies were NANOG positive (Fig. 2 B, D, and not shown). The surrounding fibroblasts that were not part of the iPS cell colony were NANOG negative, suggesting that they were not reprogrammed into iPS cells.

In a subsequent experiment using the same protocol, both 1079 fibroblasts and human IMR90 fibroblasts were transfected with the same reprogramming mRNAs. Multiple colonies were detected as early as 4 days after plating the transfected cells on irradiated MEFs. When 6 μg of each mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) were used in transfections in 6-well dishes, 3 putative iPS cell colonies were later detected in both cell lines after plating on MEFs in 10-cm plates (Figure 3). In addition to analyzing these colonies for expression of NANOG, TRA-1-60, a more stringent marker of fully reprogrammed iPS cells (Chan et al. 2009), was also used for immunofluorescence analysis. iPS colonies generated from 1079 fibroblasts (Fig. 3 A-F) and from IMR90 fibroblasts (Fig. 3 G-I) were positive for both NANOG and TRA-1-60, indicating that these colonies are fully reprogrammed type III iPS cell colonies. This protocol comprising three transfections of mRNAs encoding all six reprogramming factors and then plating onto MEF feeder cells resulted in a similar reprogramming efficiency (3-6 iPS colonies per 1 x 10^6 input cells) as was previously reported by protocols comprising delivery of the same reprogramming factors by transfection of an expression plasmid (Aoi et al. 2008).

EXAMPLE 3

This Example describes attempts to improve the efficiency of reprogramming differentiated cells using mRNA. In one approach, a protocol was used that comprised transfecting 1079 or IMR90 fibroblasts three times (once every other day) with the mRNAs encoding the six reprogramming factors in MEF-conditioned medium rather than in fibroblast medium and then growing the treated 1079 fibroblasts in MEF-conditioned medium rather than plating them on a MEF feeder layer after the treatments. At the highest transfection dose utilized (36 μg of each reprogramming factor per 10-cm dish), 208 iPS cell colonies were detected three days after the final transfection (Figure A-F). Interestingly no iPS cell colonies were detected in

the dishes transfected with either 6 or 18 μg of each of the reprogramming factors at the 3-day timepoint, suggesting that a dose above 18 μg was important, under these conditions, for iPS cell colony formation to occur within 3 days in MEF-conditioned medium. IMR90 cells showed an even higher number of iPS cell colonies, with around 200 colonies 8 days after the last transfection in the plate transfected with three 6-μg doses of each of the six reprogramming factor mRNAs and >1000 colonies in IMR90 cells transfected three times with 18-μg or 36-μg doses of each of the six reprogramming mRNAs (Figure 4 G-I). Colonies were visible 3 days after the final transfection in 1079 cells, whereas colonies only became visible 6-7 days after the final transfection in IMR90 cells. Therefore, the more mature colonies derived from the 1079 cells were larger and denser and were darker in bright-field images compared to the IMR90 colonies (Figure 4). All of the colonies on the 1079 plate transfected three times with 36 μg of each reprogramming mRNA were positive for both NANOG and TRA-1-60 8 days after the final mRNA transfection (Figure5 A-I). All of the more immature IMR90 iPS colonies were also positive for both NANOG and TRA-1-60 (Figure 5 J-O), but showed less robust staining for both markers due to their less dense cellular nature compared to the more mature 1079 colonies (Figure 5 A-I). The present protocol comprising delivery of the mRNAs into 1079 or IMR90 cells in MEF-conditioned medium had a reprogramming efficiency of 200 to >1000 colonies per 3×10^5 input cells. This protocol for inducing iPS cells was faster and almost 2-3 orders of magnitude more efficient than published protocols comprising transfecting fibroblasts with DNA plasmids encoding these same six reprogramming factors in fibroblast medium (Aoi et al. 2008). Still further, this protocol was over 7-40 times more efficient than the published protocol comprising delivery of reprogramming factors with lentiviruses, based on the published data that lentiviral delivery of reprogramming factors into 1079 newborn fibroblasts, which resulted in approximately 57 iPS cell colonies per 6×10^5 input cells (Aoi et al. 2008). This protocol is also much faster than the published methods.

EXAMPLE 4

NATURALLY OCCURRING RNA MOLECULES EXHIBIT DIFFERENTIAL ABILITIES TO ACTIVATE DENDRITIC CELLS MATERIALS AND EXPERIMENTAL METHODS

Plasmids and Reagents

Plasmids pT7T3D-MART-1 and pUNO-hTLR3 were obtained from the ATCC (Manassas, VA) and InvivoGen (San Diego, CA), respectively. pTEVluc was obtained from Dr Daniel Gallie (UC Riverside),contains pT7-TEV (the leader sequence of the tobacco etch viral genomic RNA)-luciferase-A50, and is described in Gallie, DR et al, 1995. The tobacco etch viral 5' leader and poly(A) tail are functionally synergistic regulators of translation. Gene 165:233) pSVren was generated from p2luc (Grentzmann G, Ingram JA, et al, A dual-luciferase reporter system for studying recoding signals. RNA 1998;4(4): 479-86) by removal of the firefly luciferase coding sequence with BamHI and NotI digestions, end-filling, and religation.

Human TLR3-specific siRNA, pTLR3-sh was constructed by inserting synthetic ODN encoding shRNA with 20-nt-long homology to human TLR3 (nt 703-722, accession: NM_003265) into plasmid pSilencer 4.1-CMV-neo (Ambion, Austin, TX). pCMV-hTLR3 was obtained by first cloning hTLR3-specific PCR product (nt 80-2887; Accession NM_003265) into pCRII-TOPO (Invitrogen, Carlsbad, CA), then released with Nhe I-Hind III cutting and subcloning to the corresponding sites of pcDNA3.1 (Invitrogen). LPS (E. coli 055:B5) was obtained from Sigma Chemical Co, St. Louis, MO. CpG ODN2006 and R-848 were obtained from InvivoGen.

Cells and cell culture

Human embryonic kidney 293 cells (ATCC) were propagated in DMEM supplemented with glutamine (Invitrogen) and 10% FCS (Hyclone, Ogden, UT) (complete medium). In all cases herein, "293 cells" refers to human embryonic kidney (HEK) 293 cells. 293-hTLR3 cell line was generated by transforming 293 cells with pUNO-hTLR3. Cell lines 293-hTLR7, 293-hTLR8 and 293-hTLR9 (InvivoGen) were grown in complete medium supplemented with blasticidin (10 μg/ml) (Invivogen). Cell lines 293-ELAM-luc and TLR7-293 (M. Lamphier, Eisai Research Institute, Andover MA), and TLR3-293 cells were cultured as described (Kariko et al, 2004, mRNA is an endogenous ligand for Toll-like receptor 3. J Biol Chem 279: 12542-

12550). Cell lines 293, 293-hTLR7 and 293-hTLR8 were stably transfected with pTLR3-sh and selected with G-418 (400 μg/ml) (Invitrogen). Neo-resistant colonies were screened and only those that did not express TLR3, determined as lack of IL-8 secretion loin response to poly(I):(C), were used in further studies. Leukopheresis samples were obtained from HIV-uninfected volunteers through an IRB-approved protocol.

Murine DC generation

Murine DCs were generated by collecting bone marrow cells from the tibia and femurs of 6-8-week-old C57BL/6 mice and lysing the red blood cells. Cells were seeded in 6-well plates at 10^6 cells/well in 2 ml DMEM + 10% FCS and 20 ng/ml muGM-CSF (R & D Systems). On day 3, 2 ml of fresh medium with muGM-CSF was added. On day 6, 2 ml medium/well was collected, and cells were pelleted and resuspended in fresh medium with muGM-CSF. On day 7 of the culture, the muDC were harvested, washed.

Natural RNA

Mitochondria were isolated from platelets obtained from the University of Pennsylvania Blood Bank using a fractionation lyses procedure (Mitochondria isolation kit; Pierce, Rockford, IL). RNA was isolated from the purified mitochondria, cytoplasmic and nuclear fractions of 293 cells, un-fractioned 293 cells, rat liver, mouse cell line TUBO, and DH5alpha strain of *E. coli* by Master Blaster® (BioRad, Hercules, CA). Bovine tRNA, wheat tRNA, yeast tRNA, E. coli tRNA, poly(A)+ mRNA from mouse heart and poly(I):(C) were purchased from Sigma, total RNA from human spleen and *E. coli* RNA were purchased from Ambion. Oligoribonucleotide-5'-monophosphates were synthesized chemically (Dharmacon, Lafayette, CO).

Aliquots of RNA samples were incubated in the presence of Benzonase nuclease (1 U per 5 μl of RNA at 1 microgram per microliter (μg/μl) for 1 h) (Novagen, Madison, WI). Aliquots of RNA-730 were digested with alkaline phosphatase (New England Biolabs). RNA samples were analyzed by denaturing agarose or polyacrylamide gel electrophoresis for quality assurance. Assays for LPS in RNA preparations using the Limulus Amebocyte Lysate gel clot assay were negative with a sensitivity of 3 picograms per milliliter (pg/ml) (University of Pennsylvania, Core Facility).

HPLC analysis

Nucleoside monophosphates were separated and visualized via HPLC. To release free nucleoside 3'-monophosphates, 5 μg aliquots of RNA were digested with 0.1 U RNase T2

(Invitrogen) in 10 µl of 50 mM NaOAc and 2 mM EDTA buffer (pH 4.5) overnight, then the samples were injected into an Agilent 1100 HPLC using a Waters Symmetry C 18 column (Waters, Milford, MA). At a flow rate of 1 mL/min, a gradient from 100% buffer A (30 mM KH_2P0_4 and 10 mM tetraethylammonium phosphate [PicA reagent, Waters], pH 6.0) to 30% buffer B (acetonitrile) was run over 60 minutes. Nucleotides were detected using a photodiode array at 254 nm. Identities were verified by retention times and spectra.

Dendritic cell assays

Dendritic cells in 96-well plates (approximately 1.1 x 10^5 cells/well) were treated with R-848, Lipofectin®, or Lipofectin®-RNA for 1 h, then the medium was changed. At the end of 8 h (unless otherwise indicated), cells were harvested for either RNA isolation or flow cytometry, while the collected culture medium was subjected to cytokine ELISA. The levels of IL-12 (p70) (BD Biosciences Pharmingen, San Diego, CA), IFN-α, TNF-α, and IL-8 (Biosource International, Camarillo, CA) were measured in supernatants by sandwich ELISA. Cultures were performed in triplicate or quadruplicate and measured in duplicate.

Northern blot analysis

RNA was isolated from MDDCs after an 8 h incubation following treatment as described above. Where noted, cells were treated with 2.5 µg/ml cycloheximide (Sigma) 30 min prior to the stimulation and throughout the entire length of incubation. RNA samples were processed and analyzed on Northern blots as described (Kariko et al, 2004, ibid) using human TNF-α and GAPDH probes derived from plasmids (pE4 and pHcGAP, respectively) obtained from ATCC.

RESULTS

To determine the immuno-stimulatory potential of different cellular RNA subtypes, RNA was isolated from different subcellular compartments-i.e. cytoplasm, nucleus and mitochondria. These RNA fractions, as well as total RNA, tRNA and polyA-tail-selected mRNA, all from mammalian sources, were complexed to Lipofectin® and added to MDDC. While mammalian total, nuclear and cytoplasmic RNA all stimulated MDDC, as evidenced by detectable TNF-α secretion, the TNF-α levels were much lower than those induced by *in vitro*-synthesized mRNA (Figure 6). Moreover, mammalian tRNA did not induce any detectable level of TNF-α, while mitochondrial (mt) RNA induced much more TNF-α than the other mammalian RNA subtypes.

Bacterial total RNA was also a potent activator of MDDC; by contrast, bacterial tRNA induced only a low level of TNF-α. tRNA from other sources (yeast, wheat germ, bovine) were non-stimulatory. Similar results were observed when RNA from other mammalian sources was tested. When RNA samples were digested with Benzonase, which cleaves ssRNA and dsRNA, RNA signaling was abolished in MDDC, verifying that TNF-α secretion was due to the RNA in the preparations. The activation potentials of the RNA types tested exhibited an inverse correlation with the extent of nucleoside modification. Similar results were obtained in the experiments described in this Example for both types of cytokine-generated DC.

These findings demonstrate that the immunogenicity of RNA is affected by the extent of nucleoside modification, with a greater degree of modification tending to decrease immunogenicity.

EXAMPLE 5

IN VITRO SYNTHESIS OF RNA MOLECULES WITH MODIFIED NUCLEOSIDES

MATERIALS AND EXPERIMENTAL METHODS

In vitro-transcribed RNA

Using *in vitro* transcription assays (MessageMachine and MegaScript kits; Ambion,) the following long RNAs were generated by T7 RNA polymerase (RNAP) as described (Kariko et al, 1998, Phosphate-enhanced transfection of cationic lipid-complexed mRNA and plasmid DNA. Biochim Biophys Acta 1369, 320-334) (Note: the names of templates are indicated in parenthesis; the number in the name of the RNA specifies the length): RNA-1866 (Nde I-linearized pTEVluc) encodes firefly luciferase and a 50 nt-long polyA-tail. RNA-1571 (Ssp I-linearized pSVren) encodes Renilla luciferase. RNA-730 (Hind III-linearized pT7T3D-MART-l) encodes the human melanoma antigen MART-I. RNA-713 (EcoR I-linearized pTIT3D-MART-l) corresponds to antisense sequence of MART-1, RNA497 (Bgl II-linearized pCMV-hTLR3) encodes a partial 5' fragment of hTLR3. Sequences of the RNA molecules are as follows:

RNA-I866:

ggaauucucaacacaacauauacaaaacaaacgaaucucaagcaaucaagcauucuacuucuauugcagcaauuuaaaucauuu

cuuuuaaagcaaaagcaauuuucugaaaauuuucaccauuuacgaacgauagccauggaagacgccaaaaacauaaagaaagg

cccggcgccauucuauccucuagaggauggaaccgcuggagagcaacugcauaaggcuaugaagagauacgcccugguuccu
ggaacaauugcuuuuacagaugcacauaucgaggugaacaucacguacgcggaauacuucgaaauguccguucgguuggcag
aagcuaugaaacgauaugggcugaauacaaaucacagaaucgucguaugcagugaaaacucucuucaauucuuuaugccggu
guugggcgcguuauuuaucggaguugcaguugcgcccgcgaacgacauuuauaaugaacgugaauugcucaacaguaugaac
auuucgcagccuaccguaguguuuguuuccaaaaagggguugcaaaaaauuuugaacgugcaaaaaaaauuaccaauaaucca
gaaaauuauuaucauggauucuaaaacggauuaccagggauuucagucgauguacacguucgucacaucucaucuaccuccc
gguuuuaaugaauacgauuuuguaccagaguccuuugaucgugacaaaacaauugcacugauaaugaauuccucuggaucua
cuggguuaccuaagggugugggcccuucccgcauagaacugccugcgucagauucucgcaugccagagauccuauuuuuggcaa
ucaaaucauucccggauacugcgauuuuaaguguuguuccauuccaucacgguuuuggaauguuuacuacacucggauauuu
gauauguggauuucgagucgucuuaauguauagauuugaagaagagcuguuuuuacgaucccuucaggauuacaaaauucaa
agugcguugcuaguaccaacccuauuuucauucuucgccaaaagcacucugauugacaaauacgauuuaucuaauuuacacga
aauugcuucuggggcgcaccucuuucgaaagaagucggggaagcgguugcaaaacgcuuccaucuuccagggauacgacaa
ggauaugggcucacugagacuacaucagcuauucugauuacacccgagggggaugauaaaccgggcgcggucgguaaaguug
uuccauuuuuuugaagcgaagguuguggaucuggauaccgggaaaacgcugggcguuaaucagagaggcgaauuauguguca
gaggaccuaugauuauguccgguuauguaaacaauccggaagcgaccaacgccuugauugacaaggauggauggcuacauuc
uggagacauagcuuacugggacgaagacgaacacuucuucauaguugaccgcuugaagucuuuaauuaaauacaaaggauau
caggugggccccgcugaauuggaaucgauauuguuacaacaccccaacaucuucgacgcgggcguggcaggucuucccgacg
augacgccggugaacuucccgccgccguuguuguuuuggagcacggaaagacgaugacggaaaaagagaucguggauuacgu
ggccagucaaguaacaaccgcgaaaaaguugcgcggaggaguuguguuuguggacgaaguaccgaaaggucuuaccggaaaa
cucgacgcaagaaaaaucagagagauccucauaaaggccaagaagggcggaaaguccaaauuguaaaauguaaacucuagagga
ucccaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaaca (SEQ ID No: 1).

RNA-1571:

ggcuagccaccaugacuucgaaaguuuaugauccagaacaaaggaaacggaugauaacuggucgcaguggugggccagaug
uaaacaaaugaaauguucuugauucauuuauuaauuauuaugauucagaaaaacaugcagaaaaugcuguuauuuuuuuacau
gguaacgcggccucuucuuauuuauggcgacauguugugccacauauugagccaguagcgcgguguauuauaccagaccuua
uuugguaugggcaaaucaggcaaaucugguaaugguucuuauagguuacuugaucauuacaaauaucuuacugcaugguuug
aacuucuuaaauuuaccaaagaagaucauuuuugucggccaugauuggggugcuuguuuggcauuucauuauagcuaugagc
aucaagauaagaucaaagcaauaguucacgcugaaaguguaguagaugugauugaaucaugggaugaauggccugauauuga
agaagauauugcguugaucaaaucugaagaaggagaaaaaaugguuuuggagaauaacuucuucguggaaaccauguugcca
ucaaaaaucaugagaaaguuagaaccagaagaauuugcagcauaucuugaaccauucaaagagaaaggugaaguucgucgucc

aacauuaucauggccucgugaaaucccguuaguaaaaggugguaaaccugacguuguacaaauuguuaggaauuauaaaugcu
uaucuacgugcaagugaugauuuaccaaaaauguuuauugaaucggacccaggauucuuuuccaaugcuauuguugaaggug
ccaagaaguuuccuaauacugaauuugucaaaguaaaaggucuucauuuuucgcaagaagaugcaccugaugaaaugggaaa
auauaucaaaucguucguugagcgaguucucaaaaaugaacaaaugucgacgggggccccuaggaauuuuuuuagggaagauc
uggccuuccuacaagggaaggccagggaauuuucuucagagcagaccagagccaacagccccaccagaagagagcuucagguc
uggggguagagacaacaacucccccucagaagcaggagccgauagacaaggaacuguauccuuuaacuucccucagaucacucu
uuggcaacgaccccucgucacaauaaagauaggggggcaacuaaagggaucggccgcuucgagcagacaugauaagauacauu
gaugaguuuggacaaaccacaacuagaaugcagugaaaaaaaugcuuuauuuugugaaauuugugaugcuauugcuuuauuug
uaaccauuauaagcugcaauaaacaaguuaacaacaacaauugcauucauuuuauguuucagguucagggggaggugugggga
gguuuuuuaaagcaaguaaaaccucuacaaaugugguaaaaucgauaaguuuaaacagauccagguggcacuuuucggggaa
augugcgcggaaccccuauuuguuuauuuuucuaaauacauucaaauauguaucccgcucaugagacaauaacccugauaaau
gcuucaauaau (SEQ ID No: 2).

RNA-730:

gggaauuuggcccucgaggccaagaauucggcacgaggcacgcggccagccagcagacagaggacucucauuaaggaaggug
uccugugcccugacccuacaagaugccaagagaagaugcucacuucaucuaugguuaccccaagaaggggcacggccacucuu
acaccacggcugaagaggccgcugggaucggcauccugacagugauccugggagucuuacugcucaucggcuguugguauug
uagaagacgaaauggauacagagccuugauggauaaaagucuucauguuggcacucaaugugccuuaacaagaagaugcccac
aagaaggguuugaucaucgggacagcaaagugucucuucaagagaaaaacugugaaccugugguucccaaugcuccaccugc
uuaugagaaacucucugcagaacagucaccaccaccuuauucaccuuaagagccagcgagacaccugagacaugcugaaauua
uuucucucacacuuuugcuugaauuuaauacagacaucuaauguucuccuuuggaaugguguaggaaaaaugcaagccaucu
cuaauaauaagucaguguuaaaauuuuuaguaggucccgcuagcaguacuaaucaugugaggaaaugaugagaaauauuaaauu
gggaaaacuccaucaauaaauguugcaaugcaugauaaaaaaaaaaaaaaaaaaaaaacugcggccgca (SEQ ID No: 3).

RNA-713

gggaauaagcuugcggccgcaguuuuuuuuuuuuuuuuuuuuuuuuaucaugcauugcaacauuuauugauggaguuuucccaau
uuaaauauuucucaucauuuccucacaugauuaguacugcuagcggaccuacuaaaauuuuuaacacugacuuauuauuagaga
uggcuugcauuuuuccuacaccauuccaaaggagaacauuagaugucuguauaaaauucaagcaaaagugugagagaaauaau
uucagcaugucucaggugucucgcuggcucuuaaggugaauaaggugguggugacuguucugcagagaguuucucauaagc
agguggagcauugggaaccacagguucacaguuuuucucuugaagagacacuuugcugucccgaugaucaaacccuucuugu
gggcaucuucuuguuaaggcacauugagugccaacaugaagacuuuuauccaucaaggcucuguauccauuucgucuucuac

aauaccaacagccgaugagcaguaagacucccaggaucacugucaggaugccgaucccagcggccucuucagccgugguguaa gaguggccgugccccuucuuuggguaaccauagaugaagugagcaucuucucuuggcaucuuguagggucagggcacagga caccuuccuuaaugagaguccucugucugcuggcuggccgcgugccucgugccgaauu (SEQ ID No: 4).

RNA-497:

gggagacccaagcuggcuagcagucauccaacagaaucaugagacagacuuugccuuguaucuacuuuuggggggccuuuu gcccuuugggaugcugugugcauccuccaccaccaagugcacuguuagccaugaaguugcugacugcagccaccugaaguug acucagguacccgaugaucuacccacaaacauaacaguguugaaccuuacccauaaucaacucagaagauuaccagccgccaac uucacaaggguauagccagcuaacuagcuuggauguaggauuuaacaccaucucaaaacuggagccagaauugugccagaaacu ucccauguuaaaaguuuugaaccuccagcacaaugagcuaucucaacuuucugauaaaaccuuugccuucugcacgaauuuga cugaacuccaucucauguccaacucaauccagaaaauuaaaaauaauccccuuugucaagcagaagaauuuaaucacauua (SEQ ID No: 5).

To obtain modified RNA, the transcription reaction was assembled with the replacement of one (or two) of the basic NTPs with the corresponding triphosphate-derivative(s) of the modified nucleotide 5-methylcytidine, 5-methyluridine, 2-thiouridine, N^6-methyladenosine or pseudouridine (TriLink, San Diego, CA). In each transcription reaction, all 4 nucleotides or their derivatives were present at 7.5 millimolar (mM) concentration. In selected experiments, as indicated, 6 mM m7GpppG cap analog (New England BioLabs, Beverly, MA) was also included to obtain capped RNA. ORN5 and ORN6 were generated using DNA oligodeoxynucleotide templates and T7 RNAP (Silencer® siRNA construction kit, Ambion).

RESULTS

To further test the effect of nucleoside modifications on immunogenicity, an *in vitro* system was developed for producing RNA molecules with pseudouridine or modified nucleosides. *In vitro* transcription reactions were performed in which 1 or 2 of the 4 nucleotide triphosphates (NTP) were substituted with a corresponding nucleoside-modified NTP. Several sets of RNA with different primary sequences ranging in length between 0.7-1.9 kb, and containing either none, 1 or 2 types of modified nucleosides were transcribed. Modified RNAs were indistinguishable from their non-modified counterparts in their mobility in denaturing gel electrophoresis, showing that they were intact and otherwise unmodified (Figure 7A). This

procedure worked efficiently with any of T7, SP6, and T3 phage polymerases, and therefore is generalizable to a wide variety of RNA polymerases.

These findings provide a novel *in vitro* system for production of RNA molecules with modified nucleosides.

EXAMPLE 6

IN VITRO-TRANSCRIBED RNA STIMULATES HUMAN TLR3, AND NUCLEOSIDE MODIFICATIONS REDUCE THE IMMUNOGENICITY OF RNA MATERIALS AND EXPERIMENTAL METHODS

Parental-293; 293-hTLR7 and 293-hTLR8 cells, all expressing TLR3-specific siRNA, and 293hTLR9, TLR3-293 were seeded into 96-well plates (5 x 10^4 cells/well) and cultured without antibiotics. On the subsequent day, the cells were exposed to R-848 or RNA complexed to Lipofectin® (Invitrogen) as described (Kariko et al, 1998, ibid). RNA was removed after one hour (h), and cells were further incubated in complete medium for 7 h. Supernatants were collected for IL-8 measurement.

RESULTS

To determine whether modification of nucleosides influences the RNA-mediated activation of TLRs, human embryonic kidney 293 cells were stably transformed to express human TLR3. The cell lines were treated with Lipofectin®-complexed RNA, and TLR activation was monitored as indicated 10 by interleukin (IL)-8 release. Several different RNA molecules were tested. Unmodified, *in vitro* transcribed RNA elicted a high level of IL-8 secretion. RNA containing m6A or s2U nucleoside modifications, but contrast, did not induce detectable IL-8 secretion (Figure 7B). The other nucleoside modifications tested (i.e. m5C, m5U, Ψ, and m5C/Ψ) had a smaller suppressive effect on TLR3 stimulation (Figure 7B). "Ψ" refers to pseudouridine.

Thus, nucleoside modifications such as m^6A S^2U, m^5C, m^5U, Ψ, reduce the immunogenicity of RNA as mediated by TLR3 signaling.

EXAMPLE 7

IN VITRO-TRANSCRIBED RNA STIMULATES HUMAN TLR7 AND TLR8, AND NUCLEOSIDE MODIFICATIONS REDUCE THE IMMUNOGENICITY OF RNA

To test the possibility that 293 express endogenous TLR3 that interfere with assessing effects of RNA on specific TLR receptors, expression of endogenous TLR3 was eliminated from the 293-TLR8 cell line by stably transfecting the cells with a plasmid expressing TLR3-specific short hairpin (sh)RNA (also known as siRNA). This cell line was used for further study, since it did not respond to poly(I):(C), LPS, and CpG-containing oligodeoxynucleotides (ODNs), indicating the absence ofTLR3, TLR4 and TLR9, but did respond to R-848, the cognate ligand of human TLR8 (Figure 7B). When the 293-hTLR8 cells expressing TLR3-targeted shRNA (293-hTLR8 shRNA-TLR3 cells) were transfected with *in vitro* transcribed RNA, they secreted large amounts of IL-8. By contrast, RNA containing most of the nucleoside modifications (m^5C, m^5U, Ψ, and m^5C/Ψ, S^2U) eliminated stimulation (no more IL-8 production than the negative control, i.e. empty vector). m6A modification had a variable effect, in some cases eliminating and in other cases reducing IL-8 release (Figure 7B).

The results of this Example and the previous Example show that (a) RNA with natural phosphodiester inter-nucleotide linkages (e.g. *in vitro*-transcribed RNA) stimulates human TLR3, TLR7 and TLR8; and (b) nucleoside modifications such as m6A, m5C, m5U, s2U and Ψ, alone and in combination, reduce the immunogenicity of RNA as mediated by TLR3, TLR7 and TLR8 signaling. In addition, these results provide a novel system for studying signaling by specific TLR receptors.

EXAMPLE 8

NUCLEOSIDE MODIFICATIONS REDUCE THE IMMUNOGENICITY OF RNA AS MEDIATED BY TLR7 AND TLR8 SIGNALING

The next set of experiments tested the ability of RNA isolated from natural sources to stimulate TLR3, TLR7 and TLR8. RNA from different mammalian species were transfected into the TLR3, TLR7 and TLR8-expressing 293 cell lines described in the previous Example. None of the mammalian RNA samples induced IL-8 secretion above the level of the negative control. By contrast, bacterial total RNA obtained from two different *E. coli* sources induced robust IL-8

secretion in cells transfected with TLR3, TLR7 and TLR8, but not TLR9 (Figure 7C). Neither LPS nor unmethylated DNA (CpG ODN) (the potential contaminants in bacterial RNA isolates) activated the tested TLR3, TLR7 or TLR8. Mitochondrial RNA isolated from human platelets stimulated human TLR8, but not TLR3 or TLR7.

These results demonstrate that unmodified *in vitro*-transcribed and bacterial RNA are activators ofTLR3, TLR7 and TLR8, and mitochondrial RNA stimulates TLR8. In addition, these results confirm the finding that nucleoside modification of RNA decreases its ability to stimulate TLR3, TLR7 and TLR8.

EXAMPLE 9

NUCLEOSIDE MODIFICATIONS REDUCE THE CAPACITY OF RNA TO INDUCE CYTOKINE SECRETION AND ACTIVATION MARKER EXPRESSION BY DC

MATERIALS AND EXPERIMENTAL METHODS

DC *stimulation assays*

After 20 h of incubation with RNA, DCs were stained with CD83-phycoerythrin mAb (Research Diagnostics Inc, Flanders, NJ), HLA-DR-Cy5PE, and CD80 or CD86-fluorescein isothiocyanate mAb and analyzed on a FACScalibur® flow cytometer using CellQuest® software (BD Biosciences). Cell culture supernatants were harvested at the end of a 20 h incubation and subjected to cytokine ELISA. The levels of IL-12 (p70) (BD Biosciences Pharmingen, San Diego, CA), IFN-α, and TNF-α (Biosource International, Camarillo, CA) were measured in supernatants by ELISA. Cultures were performed in triplicate or quadruplicate, and each sample was measured in duplicate.

RESULTS

The next experiments tested the ability of RNA containing modified or unmodified nucleosides to stimulate cytokine-generated MDDC. Nucleoside modifications reproducibly diminished the ability of 5 RNA to induce TNF-α and IL-12 secretion by both GM-CSF/IL-4-generated MDDC and (GMCSF)/IFN-α-generated MDDC, in most cases to levels no greater than the negative control (Figures 8A and B). Results were similar when other sets of RNA with the same base modifications but different primary sequences and lengths were tested, or when

the RNA was further modified by adding a 5' cap structure and/or 3'-end polyA-tail or by removing the 5' triphosphate moiety. RNAs of different length and sequence induced varying amounts of TNF-α from DC, typically less than a two-fold difference (Figure 8C).

Next, the assay was performed on primary DC1 and DC2. Primary monocytoid (DCI, BDCA1$^+$) and plasmacytoid (DC2, BDCA4$^+$) DC were purified from peripheral blood. Both cell types produced TNF-α when exposed to R-848, but only DC1 responded to poly(I):(C), at a very low level, indicating an absence of TLR3 activity in DC2. Transfection of *in vitro* transcripts induced TNF-α secretion in both DC1 and DC2, while m5U, Ψ or s2U-modified transcripts were not stimulatory (Figure 8D). In contrast to the cytokine-generated DC, m5C and m6A modification of RNA did not decrease its stimulatory capacity in the primary DC1 and DC2. Transcripts with m6A/Ψ double modification were non-stimulatory, while a mixture of RNA molecules with single type of modification (m6A+Ψ) was a potent cytokine inducer. Thus, uridine modification exerted a dominant suppressive effect on an RNA molecule *in cis* in primary DC. These results were consistent among all donors tested.

These findings show that *in vitro*-transcribed RNA stimulates cytokine production by DC. In addition, since DC2 do not express TLR3 or TLR8, and m5C and m6A modification of RNA decreased its stimulatory capacity of TLR7, these findings show that primary DC have an additional RNA signaling entity that recognizes m5C- and m6A-modified RNA and whose signaling is inhibited by modification of U residues.

As additional immunogenicity indicators, cell surface expression of CD80, CD83, CD86 and MHC class II molecules, and secretion of TNF-α were measured by FACS analysis of MDDC treated with RNA-1571 and its modified versions. Modification of RNA with pseudouridine and modified nucleosides (m5C, m6A, s2U and m6A/Ψ) decreased these markers (Figure 9), confirming the previous findings.

In summary, RNA's capacity to induce DCs to mature and secrete cytokines depends on the subtype of DC as well as on the characteristics of nucleoside modification present in the RNA. An increasing amount of modification decreases the immunogenicity of RNA.

EXAMPLE 10

SUPPRESSION OF RNA-MEDIATED IMMUNE STIMULATION IS PROPORTIONAL TO THE NUMBER OF MODIFIED NUCLEOSIDES PRESENT IN RNA

MATERIALS AND EXPERIMENTAL METHODS

Human DC

For cytokine-generated DC, monocytes were purified from PBMC by discontinuous Percoll gradient centrifugation. The low density fraction (monocyte enriched) was depleted of B, T, and, NK cells using magnetic beads (Dynal, Lake Success, NY) specific for CD2, CD16, CD19, and CD56, yielding highly purified monocytes as determined by flow cytometry using anti-CD14 (>95%) or antiCD11c (>98%) mAb.

To generate immature DC, purified monocytes were cultured in AIM V serum-free medium (Life Technologies), supplemented with GM-CSF (50 ng/ml) + IL-4 (100 ng/ml) (R & D Systems, Minneapolis, MN) in AIM V medium (Invitrogen) for the generation of monocyte-derived DC (MDDC) as described (Weissman, D et al, 2000. J Immunol 165: 4710-4717). DC were also generated by treatment with GM-CSF (50 ng/ml) + IFN-α (1,000 V/ml) (R & D Systems) to obtain IFN-α MDDC (Santini et al., 2000. Type I interferon as a powerful adjuvant for monocyte-derived dendritic cell development and activity in vitro and in Hu-PBL-SCID mice. J Exp Med 191: 1777-178).

Primary myeloid and plasmacytoid DCs (DC1 and DC2) were obtained from peripheral blood using BDCA-1 and BDCA-4 cell isolation kits (Miltenyi Biotec Auburn, CA), respectively.

RESULTS

Most of the nucleoside-modified RNA utilized thus far contained one type of modification occurring in approximately 25% of the total nucleotides in the RNA (e.g. all the uridine bases). To define the minimal frequency of particular modified nucleosides that is sufficient to reduce immunogenicity under the conditions utilized herein, RNA molecules with limited numbers of modified nucleosides were generated. In the first set of experiments, RNA was transcribed *in vitro* in the presence of varying ratios of m6A, Ψ (pseudouridine) or m5C to their corresponding unmodified NTPs. The amount of incorporation of modified nucleoside phosphates into RNA was expected to be proportional to the ratio contained in the transcription

reaction, since RNA yields obtained with T7 RNAP showed the enzyme utilizes NTPs of m6A, Ψ or m5C almost as efficiently as the basic NTPs. To confirm this expectation, RNA transcribed in the presence of UTP:Ψ in a 50:50 ratio was digested and found to contain UMP and Ψ in a nearly 50:50 ratio (Figure 10A).

RNA molecules with increasing modified nucleoside content were transfected into MDDC, and TNF-α secretion was assessed. Each modification (m6A, Ψ and m5C) inhibited TNF-α secretion proportionally to the fraction of modified bases. Even the smallest amounts of modified bases tested (0.2-0.4%, corresponding to 3-6 modified nucleosides per 1571 nt molecule), was sufficient to measurably inhibit cytokine secretion (Figure 10B). RNA with of 1.7-3.2% modified nucleoside levels (14-29 modifications per molecule) exhibited a 50% reduction in induction of TNF-α expression. In TLR-expressing 293 cells, a higher percentage (2.5%) of modified nucleoside content was required to inhibit RNA-mediated signaling events.

Thus, pseudouridine and modified nucleosides reduce the immunogenicity of RNA molecules, even when present as a small fraction of the residues.

In additional experiments, 21-mer oligoribonucleotides (ORN) with phosphodiester inter-nucleotide linkages were synthesized wherein modified nucleosides (m5C, Ψ or 2'-O-methyl-U [Um]) were substituted in a particular position (Figure 11A). While the unmodified ORN induced TNF-α secretion, this effect was abolished by the presence of a single nucleoside modification (Figure 11B). Similar results were obtained with TLR-7 and TLR-8-transformed 293 cells expressing TLR3-targeted siRNA.

The above results were confirmed by measuring TNF-α mRNA levels in MDDC by Northern blot assay, using both the above 21-mer ORN (ORN1) and 31-mer *in vitro*-synthesized transcripts (ORN5 and ORN6). To amplify the signal, cycloheximide, which blocks degradation of selected mRNAs, was added to some samples, as indicated in the Figure. The unmodified ODN increased TNF-α mRNA levels, while ORNs containing a single modified nucleoside were significantly less stimulatory; ORN2-Um exhibited the greatest decrease TNF-α production (Figure 11C). Similar results were observed in mouse macrophage-like RAW cells and in human DC.

In summary, each of the modifications tested (m6A, m5C, m5U, s2U, Ψ and 2'-O-methyl) suppressed RNA mediated immune stimulation, even when present as a small fraction of the residues. Further suppression was observed when the proportion of modified nucleosides was increased.

EXAMPLE 11

PSEUDOURIDINE-MODIFICATION OF RNA REDUCES ITS IMMUNOGENICITY *IN VIVO*

To determine the effect of pseudouridine modification on immunogenicity of RNA *in vivo*, 0.25 μg RNA) was complexed to Lipofectin® and injected intra-tracheally into mice, mice were bled 24 h later, and circulating levels of TNF-α and IFN-α were assayed from serum samples. Capped, pseudouridine-modified mRNA induced significantly less TNF-α. and IFN-α mRNA than was elicited by unmodified mRNA (Figure 12A-B).

These results provide further evidence that pseudouridine-modified mRNA is significantly less immunogenic *in vivo* than unmodified RNA.

EXAMPLE 12

PSEUDOURIDINE-CONTAINING RNA EXHIBITS DECREASED ABILITY TO ACTIVATE PRK

MATERIALS AND EXPERIMENTAL METHODS

PKR phosphorylation assays

Aliquots of active PKR agarose (Upstate) were incubated in the presence of magnesium/ATP coctail (Upstate), kinase buffer and [gamma^{32}p] ATP mix and RNA molecules for 30 min at 30°C. Unmodified RNA and RNA with nucleoside modification (m5C, pseudouridine, m6A, m5U) and dsRNA were tested. Human recombinant eIF2a (BioSource) was added, and samples were further incubated for 5 min, 30°C. Reactions were stopped by adding NuPage LDS sample buffer with reducing reagent (Invitrogen), denatured for 10 min, 70° C, and analyzed on 10% PAGE. Gels were dried and exposed to film. Heparin (1 U/μl), a PKR activator, was used as positive control.

RESULTS

To determine whether pseudouridine-containing mRNA activates dsRNA-dependent protein kinase (PKR), *in vitro* phosphorylation assays were performed using recombinant human PKR and its substrate, eIF2α (eukaryotic initiation factor 2 alpha) in the presence of capped, renilla-encoding mRNA (0.5 and 0.05 ng/μl). mRNA containing pseudouridine (Ψ) did not activate PKR, as detected by lack of both self-phosphorylation of PKR and phosphorylation of eIF2α, while RNA without nucleoside modification and mRNA with m^5C modification activated PKR (Figure 13). Thus, pseudouridine modification decreases RNA immunogenicity.

EXAMPLE 13

ENHANCED TRANSLATION OF PROTEINS FROM PSEUDOURIDINE AND m^5C-CONTAINING RNA *IN VITRO*

MATERIALS AND EXPERIMENTAL METHODS

In vitro translation of mRNA in rabbit reticulocyte lysate

In vitro-translation was performed in rabbit reticulocyte lysate (Promega, Madison WI). A 9-μl aliquot of the lysate was supplemented with 1 μl (1 μg) mRNA and incubated for 60 min at 30°C. One μl aliquot was removed for analysis using firefly and renilla assay systems (Promega, Madison WI), and a LUMAT LB 950 luminometer (Berthold/EG&G Wallac, Gaithersburg, MD) with a 10 sec measuring time.

RESULTS

To determine the effect of pseudouridine modification on RNA translation efficiency *in vitro,* (0.1 μg/μl) uncapped mRNA modified with pseudouridine encoding firefly luciferase was incubated in rabbit reticulocyte lysate for 1 h at 30°C, and luciferase activity was determined. mRNA containing pseudouridine was translated more than 2-fold more efficiently than RNA without pseudouridine in rabbit reticulocyte lysates, but not in wheat extract or *E. coli* lysate (Figure 14), showing that pseudouridine modification increases RNA translation efficiency. Similar results were obtained with m^5C-modified RNA. When a polyA tail was added to pseudouridine-containing mRNA, a further 10-fold increase in translation efficiency was observed. (Example 10).

Thus, pseudouridine and m^5C modification increases RNA translation efficiency, and addition of a polyA tail to pseudouridine-containing mRNA further increases translation efficiency.

EXAMPLE 14

ENHANCED TRANSLATION OF PROTEINS FROM PSEUDOURIDINECONTAINING RNA IN CULTURED CELLS

MATERIALS AND EXPERIMENTAL METHODS

Translation assays in cells

Plates with 96 wells were seeded with 5 x 10^4 cells per well 1 day before transfection. Lipofectin®-mRNA complexes were assembled and added directly to the cell monolayers after removing the culture medium (0.2 μg mRNA-0.8 μg lipofectin in 50 μl per well). Cells were incubated with the transfection mixture for 1 h at 37°C, 5% CO_2 incubator, then the mixture was replaced with fresh, pre-warmed medium containing 10% FCS, then cells were analyzed as described in the previous Example.

RESULTS

To determine the effect of pseudouridine modification on RNA translation in cultured cells, 293 cells were transfected with *in vitro*-transcribed, nucleoside-modified, capped mRNA encoding the reporter protein renilla. Cells were lysed 3 h after initiation of transfection, and levels of renilla were measured by enzymatic assays. In 293 cells, pseudouridine-and m5C-modified DNA were translated almost 10 times and 4 times more efficiently, respectively, than unmodified mRNA (Figure 15A).

Next, the experiment was performed with primary, bone marrow-derived mouse DC, in this case lysing the cells 3 h and 8 h after transfection. RNA containing the pseudouridine modification was translated 15-30 times more efficiently than unmodified RNA (Figure 15B).

Similar expression results were obtained using human DC and other primary cells and established cell lines, including CHO and mouse macrophage-like RAW cells. In all cell types, pseudouridine modification produced the greatest enhancement of the modifications tested.

Thus, pseudouridine modification increased RNA translation efficiency in all cell types tested, including different types of both professional antigen-presenting cells and non-

professional antigen-presenting cells, providing further evidence that pseudouridine modification increases the efficiency of RNA translation.

EXAMPLE 15

5' AND 3' ELEMENTS FURTHER ENHANCE THE TRANSLATION OF ψmRNA IN MAMMALIAN CELLS

To test the effect of additional RNA structural elements on enhancement of translation by pseudouridine modification, a set of firefly luciferase-encoding ψmRNAs were synthesized that contained combinations of the following modifications: 1) a unique 5' untranslated sequence (TEV, a cap independent translational enhancer), 2) cap and 3) polyA-tail. The ability of these modifications to enhance translation of ψmRNA or conventional mRNA was assessed (Figure 16A). These structural elements additively enhanced translational efficiency of both conventional and ψmRNA, with ψmRNA exhibiting greater protein production from all constructs.

Ability of protein expression from the most efficient firefly luciferase ψmRNA construct, capTEVlucA50 (containing TEV, cap, and an extended poly(A) tail) was next examined over 24 hours in 293 cells (Figure 16B). ψmRNA produced more protein at every time point tested and conferred more persistent luciferase expression than equivalent conventional mRNA constructs, showing that ψ-modifications stabilize mRNA.

To test whether ψ-modification of mRNA improved translation efficiency in mammalian cells *in situ,* caplacZ-ψmRNA constructs with or without extended polyA-tails (A_n) and encoding β-galactosidase (lacZ) were generated and used to transfect 293 cells. 24 h after mRNA delivery, significant increases in β-galactosidase levels were detected by X-gal visualization, in both caplacZ and caplacZ-A_n, compared to the corresponding control (conventional) transcripts (Figure 16C). This trend was observed when either the number of cells expressing detectable levels of β-galactosidase or the signal magnitude in individual cells was analyzed.

EXAMPLE 13

ENHANCED TRANSLATION OF PROTEINS FROM PSEUDOURIDINE CONTAINING RNA *IN VIVO* MATERIALS AND EXPERIMENTAL METHODS

Intracerebral RNA injections

All animal procedures were in accordance with the NIH Guide for Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee. Male Wistar rats (Charles River Laboratories, Wilmington, MA) were anesthetized by intraperitoneal injection of sodium pentobarbital (60 mg/kg body weight). Heads were placed in a stereotaxic frame, and eight evenly spaced 1.5 mm diameter burr holes were made bilaterally (coordinates relative to bregma: anterior/posterior +3,0, -3, -6 mm; lateral ±2.5 mm) leaving the dura intact. Intra-cerebral injections were made using a 25 µl syringe (Hamilton, Reno, NV) with a 30 gauge, 1 inch sterile needle (Beckton 25 Dickinson Labware, Franklin Lakes, NJ) which was fixed to a large probe holder and stereotactic arm. To avoid air space in the syringe, the needle hub was filled with 55 µl complex before the needle was attached, and the remainder of the sample was drawn through the needle. Injection depth (2 mm) was determined relative to the surface of the dura, and 4 µl complex (32 ng mRNA) was administered in a single, rapid bolus infusion. 3 hours (h) later, rats were euthanized with halothane, and brains were removed into chilled phosphate buffered saline.

Injection of RNA into mouse tail vein

Tail veins of female BALB/c mice (Charles River Laboratories) were injected (bolus) with 60 µl Lipofectin®-complexed RNA (0.26 µg). Organs were removed and homogenized in luciferase or Renilla lysis buffer in microcentrifuge tubes using a pestle. Homogenates were centrifuged, and supernatants were analyzed for activity.

Delivery of RNA to the lung

Female BALB/c mice were anaesthetized using ketamine (100 mg/kg) and xylasine (20 mg/kg). Small incisions were made in the skin adjacent to the trachea. When the trachea was exposed, 501-11 of Lipofectin®-complexed RNA (0.2 µg) was instilled into the trachea towards the lung. Incisions were closed, and animals allowed to recover. 3 hours after RNA delivery, mice were sacrificed by cervical dislocation and lungs were removed, homogenized in luciferase

or Renilla lysis buffer (250 µl), and assayed for activity. In a different set of animals, blood samples (100 µl/animal) were collected from tail veins, clotted, and centrifuged. Serum fractions were used to determine levels of TNF and IFNα by ELISA as described in the Examples above, using mouse-specific antibodies.

RESULTS

To determine the effect of pseudouridine modification on RNA translation *in vivo,* each hemisphere of rat brain cortexes was injected with either capped, renilla-encoding pseudouridine modified RNA or unmodified RNA, and RNA translation was measured. Pseudouridine-modified RNA was translated significantly more efficiently than unmodified RNA (Figure 17A).

Next, expression studies were performed in mice. Firefly luciferase-encoding mRNAs because no endogenous mammalian enzyme interferes with its detection. Transcripts (unmodified and ψmRNA) were constructed with cap, TEV (capTEVA$_{50}$) and extended (-200 nt) poly(A) tails. 0.25 µg RNA Lipofectin®-complexed was injected into mice (intravenous (i.v.) tail vein). A range of organs were surveyed for luciferase activity to determine the optimum measurement site. Administration of 0.3 µg capTEVlucAn ψmRNA induced high luciferase expression in spleen and moderate expression in bone marrow, but little expression in lung, liver, heart, kidney or brain (Figure 17B). In subsequent studies, spleens were studied.

Translation efficiencies of conventional and ψmRNA (0.015 mg/kg; 0.3 µg/animal given intravenously) were next compared in time course experiments. Luciferase activity was readily detectable at 1 h, peaked at 4 h and declined by 24 h following administration of either conventional or ψmRNA, but at all times was substantially greater in animals given ψmRNA (Figure 17C, left panel). By 24 h, only animals injected with ψmRNA demonstrated detectable splenic luciferase activity (4-fold above background). A similar relative pattern of expression (between modified and unmodified mRNA) was obtained when mRNAs encoding *Renilla* luciferase (capRen with or without ψ modifications) were injected into the animals instead of firefly luciferase, or when isolated mouse splenocytes were exposed to mRNA in culture.

In the next experiment, 0.25 µg mRNA-Lipofectin® was delivered to mouse lungs by intratracheal injection. Capped, pseudouridine-modified RNA was translated more efficiently than capped RNA without pseudouridine modification (Figure 17D).

Thus, pseudouridine modification increases RNA translation efficiency *in vitro,* in cultured cells, and *in vivo*-in multiple animal models and by multiple routes of administration, showing its widespread application as a means of increasing the efficiency of RNA translation.

EXAMPLE 17

PSEUDOURIDINE MODIFICATION ENHANCES RNA STABILITY IN VIVO

Northern analyses of splenic RNA at 1 and 4 h post injection in the animals from the previous Example revealed that the administered mRNAs, in their intact and partially degraded forms, were readily detectable (Figure 17C, right panel). By contrast, at 24 h, unmodified capTEVlucAn mRNA was below the level of detection, while capTEVlucAn ψmRNA, though partially degraded, was still clearly detectable. Thus, ψmRNA is more stably preserved *in vivo* than control mRNA.

To test whether *in vivo* protein production is quantitatively dependent on the concentration of intravenously-delivered mRNA, mRNAs were administered to mice at 0.015-0.150 mg/kg (0.3-3.0 μg capTEVlucAn per animal) and spleens were analyzed 6 hours later as described above. Luciferase expression correlated quantitatively with the amount of injected RNA (Figure 18) and at each concentration.

These findings confirm the results of Example 15, demonstrating that ψmRNA is more stable than unmodified RNA. Further immunogenicity of ψ-mRNA was less than unmodified RNA, as described herein above (Figure 12 and Figure 17C, right panel).

To summarize Examples 16-17, the 3 advantages of ψ-mRNA compared with conventional mRNA (enhanced translation, increased stability and reduced immunogenicity) observed *in vitro* are also observed *in vivo*.

EXAMPLE 18

ψmRNA DELIVERED VIA THE RESPIRATORY TRACT BEHAVES SIMILARLY TO INTRAVENOUSLY ADMINISTERED mRNA

To test the ability of ψmRNA to be delivered by inhalation, Lipofectin®-or PEI-complexed mRNAs encoding firefly luciferase were delivered to mice by the intratracheal route,

wherein a needle was placed into the trachea and mRNA solution sprayed into the lungs. Similar to intravenous delivery, significantly greater luciferase expression was observed with ψmRNA compared to unmodified mRNA (Figure 19), although significantly less protein was produced with the intratracheal as compared to the intravenous routes. Unmodified mRNA administered by the intratracheal route was associated with significantly higher concentrations of inflammatory cytokines (IFN-α and TNF-α) compared with vehicle controls, while ψmRNA was not (Figure 19).

Thus, ψmRNA can be delivered by inhalation without activating the innate immune response.

EXAMPLE 19

DELIVERY OF EPO-ψmRNA TO 293 CELLS

ψmRNA was generated from a plasmid containing the human EPO cDNA. When 0.25 μg of EPO-ψmRNA was transfected into 10^6 cultured 293 cells, greater than 600 mU/ml of EPO protein was produced. Thus, modified RNA molecules of the present invention are efficacious at delivering recombinant proteins to cells.

EXAMPLE 20

PREPARATION OF IMPROVED EPO-ENCODING ψmRNA CONSTRUCTS

MATERIALS AND EXPERIMENTAL METHODS

The EPO coding sequence is cloned using restriction enzyme techniques to generate 2 new plasmids, pTEV-EPO and pT7TS-EPO, that are used as templates for EPO-ψmRNA production. EPO-ψmRNAs are produced from these templates by *in vitro* transcription (MessageMachine® and MegaScript® kits; Ambion,) using T7 RNA polymerase (RNAP), incorporating nucleosides at equimolar (7.5mM) concentrations. To incorporate the nucleoside-modifications, ψ triphosphate (TriLink, San Diego, CA) replaces UTP in the transcription reaction. To ensure capping of the ψmRNA, a non-reversible cap-analog, 6 mM 3'-O-Me-

m7GpppG (New England BioLabs, Beverly, MA) is also included. The ψmRNAs are poly(A)-tailed in a reaction of ~1.5 μg/μl RNA,5 mM ATP, and 60 U/μl yeast poly(A) polymerase (USB, Cleveland, OH) mixed at 30°C for 3 to 24 h. Quality of ψmRNAs is assessed by denaturing agarose gel electrophoresis. Assays for LPS in mRNA preparations using the Limulus Amebocyte Lysate gel clot assay with a sensitivity of 3 pg/ml are also performed.

RESULTS

The proximal 3'-untranslated region (3'UTR) of EPO-ψmRNA preserves a ~90 nt-long pyrimidine-rich stabilizing element from the nascent EPO mRNA, which stabilizes EPO mRNA by specific association with a ubiquitous protein, erythropoietin mRNA-binding protein (ERBP). To maximize the stability of EPO-ψmRNA, 2 alterations are incorporated into the EPO plasmid to improve the stability and translational efficiency of the transcribed mRNA: 1) A 5'UTR sequence of the tobacco etch virus (TEV) is incorporated upstream of the EPO coding sequence to generate pTEV-EPO. 2) A plasmid, pT7TS-EPO, is generated, wherein the EPO cDNA is flanked by sequences corresponding to 5' and 3' UTRs of Xenopus beta-globin mRNA.

In addition, the length of the poly(A) tail during the production of ψmRNA from these plasmid templates is extended, by increasing the incubation period of the poly(A) polymerase reaction. The longer poly(A) tail diminishes the rate at which ψmRNA degrades during translation.

These improvements result in enhanced translation efficiency *in vivo,* thus minimizing the therapeutic dose of the final product.

EXAMPLE 21

IN VITRO ANALYSIS OF PROTEIN PRODUCTION FROM EPO mRNA CONSTRUCTS

MATERIALS AND EXPERIMENTAL METHODS

Preparation of mammalian cells.

Human embryonic kidney 293 cells (ATCC) are propagated in DMEM supplemented with glutamine (Invitrogen) and 10% FCS (Hyclone, Ogden, UT) (complete medium). Leukopheresis samples are obtained from HIV-uninfected volunteers through an IRB-approved protocol. DCs are produced as described above and cultured with GM-CSF (50 ng/ml) + IL-4 (100 ng/ml) (R & D Systems) in AIM V medium® (Invitrogen).

Murine spleen cells and DC are obtained by published procedures. Briefly, spleens from BALB/c mice are aseptically removed and minced with forceps in complete medium. Tissue fragments are sedimented by gravity and the single cell suspension washed and lysed with AKC lysis buffer (Sigma). Murine DCs are derived from bone marrow cells collected from femurs and tibia of 6-9-weekold BALB/c mice. Cells are cultured in DMEM containing 10% FCS (Invitrogen) and 50 ng/ml muGM-CSF (R&D) and used on day 7.

Transfection of cells and detection of EPO and pro-inflammatory cytokines

Transfections are performed with Lipofectin in the presence of phosphate buffer, an effective delivery method for splenic and *in vitro* cell expression. EPO-ψmRNA (0.25 μg/well; 100,000 cells) is added to each cell type in triplicate for 1 hour, and supernatant replaced with fresh medium. 24 hours later, supernatant is collected for ELISA measurement of EPO, IFN-α or β and TNF-α.

RESULTS

To evaluate the impact of unique UTRs on enhancement of ψmRNA translational efficiency, EPO-ψmRNA containing, or not containing, each improvement (5' TEV element, (β-globin 5' and 3' UTRs) with long poly(A) tails are tested for *in vitro* protein production and *in vitro* immune activation using EPO mRNA containing conventional nucleosides as controls. Efficiency of protein production from each mRNA is assessed in mammalian cell lines, (HEK293, CHO), human and murine primary DCs, and spleen cells for each mRNA. Measurement of total EPO produced in all cell types and immunogenicity (supernatant-associated proinflammatory cytokines) in primary cells is evaluated. The mRNA construct that demonstrates the optimum combination of high EPO production (in 1 or more cell types) and

low cytokine elicitation is used in subsequent studies. Improvements in 5' and 3'UTRs of EPO-ψmRNA and longer poly(A) tails result in an estimated 2-10-fold enhancement in translation efficiency, with no increase in immunogenicity.

EXAMPLE 22

CHARACTERIZATION OF EPO PRODUCTION AND BIOLOGICAL RESPONSE TO EPO-ψmRNA *IN VIVO*

MATERIALS AND EXPERIMENTAL METHODS

Administration of EPO-ψmRNA to mice.

All animal studies described herein are performed in accordance with the NIH Guide for Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee of the University of Pennsylvania. Female BALB/c mice (n=5 per experimental condition; 6 weeks, 18-23 g; Charles River Laboratories) are anesthetized using 3.5% halothane in a mixture of N_2O and O_2 (70:30), then halothane reduced to 1% and anesthesia maintained using a nose mask. Animal body temperatures are maintained throughout the procedure using a 37°C warmed heating pad. EPO-ψmRNA-lipofectin complexes (constructed by mixing varying amounts of nucleic acid with 1 µl lipofectin in 60 µl final volume are injected into the lateral tail vein. Blood samples are collected 3 times a day for 3 days post mRNA injection during the time-course study, at 1 optimal time point in dose-response studies, and daily from days 2-6 in studies for reticulocytosis.

Determination of reticulocytes by flow cytometry.

Whole blood samples are stained using Retic-COUNT reagent (BD Diagnostics) and data events acquired on a FACScan flow cytometer. Red blood cells (RBCs) are selected by forward and side scatter properties and analyzed for uptake of Thiazole Orange. Cells stained with Retic-COUNT reagent are detected by fluorescence and reticulocytes expressed as the percentage of total RBC. At least 50,000 events are counted per sample.

RESULTS

To optimize production of biologically functional human EPO protein (hEPO) in response to EPO-encoding mRNA, the following studies are performed:

Time course of EPO production after a single injection of EPO-ψmRNA. Following intravenous administration of 1 μg EPO-ψmRNA, hEPO is measured serially from 1-96 h after EPO-ψmRNA administration by ELISA, to determine the half-life of EPO protein in the serum. This half-life is a result of both the half-life of EPO protein and the functional half-life of the EPO-ψmRNA. The resulting optimal time point for measuring EPO protein after EPO-ψmRNA administration is utilized in subsequent studies.

Dose-response of EPO production after a single injection of EPO-ψmRNA. To determine the correlation between the amount of EPO protein produced and the amount of EPO-ψmRNA administered, increasing concentrations of EPO-ψmRNA (0.01 to 1 μg/animal) are administered and EPO is measured at the optimal time point.

Relationship between hEPO production and reticulocytosis. To measure the effect of EPO-ψmRNA on a biological correlate of EPO activity, flow cytometry is used to determine reticulocyte frequency in blood). Flow cytometry has a coefficient of variation of < 3%. Mice receive a single dose of EPO-ψmRNA, and blood is collected from mice daily from days 2-6. The relationship between EPO-ψmRNA dose and reticulocyte frequency is then evaluated at the time point of maximal reticulocytosis. The dose of EPO-ψmRNA that leads to at least a 5% increase in reticulocyte count is used in subsequent studies. Serum hEPO concentrations in mice of an estimated 50 mU/ml and/or an increase in reticulocyte frequency of an estimated 5% are obtained.

EXAMPLE 23

MEASURING IMMUNE RESPONSES TO EPO-ψmRNA *IN VIVO*

MATERIALS AND EXPERIMENTAL METHODS

Detection of cytokines in plasma.

Serum samples obtained from blood collected at different times during and after 7 daily lipofectin-complexed mRNA administrations are analyzed for mouse IFN-α, TNF-α, and IL-12 using ELISA kits.

Northern blot analysis.

Aliquots (2.0 μg) of RNA samples isolated from spleen are separated by denaturing 1.4% agarose gel electrophoresis, transferred to charged membranes (Schleicher and Schuell) and hybridized in MiracleHyb® (Stratagene). Membranes are probed for TNF-α, down-stream IFN signaling molecules (e.g. IRF7, IL-12 p35 and p40, and GAPDH) and other markers of immune activation. Specificity of all probes is confirmed by sequencing. To probe the membranes, 50 ng of DNA is labeled using Redivue [alpha-^{32}P] dCTP® (Amersham) with a random prime labeling kit (Roche). Hybridized membranes are exposed to Kodak BioMax MS film using an MS intensifier screen at -70°C.

Histopathology.

Spleens from EPO-ψmRNA-treated and positive and negative control-treated mice are harvested, fixed, sectioned, stained with hematoxylin and eosin and examined by a veterinary pathologist for signs of immune activation.

RESULTS

To confirm the reduced immunogenicity of RNA molecules of the present invention, mice (n = 5) receive daily doses of EPO-ψmRNA for 7 days, then are evaluated for immune-mediated adverse events, as indicated by serum cytokine concentrations, splenic expression of mRNAs encoding inflammatory proteins, and pathologic examination. Maximum administered doses are 3 μg or 5 x the effective single dose as determined above. Unmodified mRNA and Lipofectin® alone are used as positive and negative controls, respectively.

These studies confirm the reduced immunogenicity of RNA molecules of the present invention.

EXAMPLE 24

FURTHER IMPROVEMENT OF EPO-ψmRNA DELIVERY METHODS

Nanoparticle complexing.

Polymer and ψmRNA solutions are mixed to form complexes. Various formulation conditions are tested and optimized: (1) sub-22 nm polyethylenimine (PEI)/mRNA complexes are made by addition of 25 volumes of mRNA to 1 volume of PEI in water with no mixing for 15 minutes. (2) The rod-like poly-L-lysine -polyethylene glycol (PLL-PEG) with average dimensions of 12x150 nm is synthesized by slow addition of9 volumes of mRNA to 1 volume of CK_{30}-PEG_{10k} in acetate counterion buffer while vortexing. (3)For synthesis of biodegradable gene carrier polymer, polyaspartic anhydride co-ethylene glycol (PAE) is synthesized by ring opening polycondensation of N-(Benzyloxycarbonyl)L-aspartic anhydride and ethylene glycol. Then, the pendent amine of aspartic acid is deprotected and protonated by acidification with hydrogen chloride and condensed with mRNA. (4) For latest generation of nanoparticles, aliquot stock $CK_{30}PEG_{10k}$ as ammonium acetate (1.25mL; 6.4mg/mL) is added to siliconized Eppendorf tubes. Then mRNA is added slowly to $CK_{30}PEG_{10k}$ (2.5mg in 11.25mL RNase free H_2O) over 1-2 mins. After 15 mins, it is diluted 1:2 in RNase-free H_2O.

Intratracheal delivery.

Mice are anesthetized with 3% halothane (70% N_2O + 30% O_2) in an anesthetic chamber and maintained with 1% halothane (70% N_20 + 30% O_2) during operation using a nose cone. Trachea os exposed, and 50 µl of mRNA complex is infused with 150 µl air into the lung through the trachea using 250 µl Hamilton syringe (Hamilton, Reno, NV) with a 27 G 1/2" needle.

RESULTS

To improve efficiency of delivery and expression of ψmRNA administered via the intratracheal (i.t.) route, ψmRNA is encapsulated in nanoparticles. Nanoparticle packaging involves condensing and encapsulating DNA (for example) into particles that are smaller than the pore of the nuclear membrane, using chemicals including poly-L-lysine and polyethylene glycol. RNA is packaged into 4 different nanoparticle formulations (PEI, PLL,PAE, and $CK_{30}PEG_{10k}$), and efficiency of ψmRNA delivery is compared for luciferase-encoding ψmRNA

compare the (Luc-ψmRNA). Delivery kinetics and dose response are then characterized using EPO-ψmRNA.

EXAMPLE 25

PREVENTION OF RESTENOSIS BY DELIVERY TO THE CAROTID ARTERY OF RECOMBINANT HEAT SHOCK PROTEIN-ENCODING, MODIFIED mRNA

MATERIALS AND EXPERIMENTAL METHODS

Experimental design

RNA is administered to the carotid artery of rats by intra-arterial injection near the time of balloon angioplasty, after which blood flow is reinstated. Rats are sacrificed 3 h following injection, carotid artery sections are excised, vascular endothelial cells are harvested and homogenized, and luciferase activity is determined as described in above Examples.

RESULTS

Luciferase-encoding pseudouridine-modified RNA is administered to rat carotid arteries. 3 hours later, luciferase RNA can be detected at the delivery site but not the adjacent sites.

Next, this protocol is used to prevent restenosis of a blood vessel following balloon angioplasty in an animal restenosis model, by delivery of modified RNA encoding a heat shock protein, e.g. HSP70; a growth factor (e.g. platelet-derived growth factor (PDGF), vascular endothelial growth factor (V-EGF), or insulin-like growth factor (IGF); or a protein that down-regulates or antagonizes growth factor signaling. Administration of modified RNA reduces incidence of restenosis.

EXAMPLE 26

TREATMENT OF CYSTIC FIBROSIS BY DELIVERY OF CFTR-ENCODING MODIFIED mRNA MOLECULES TO RESPIRATORY EPITHELIUM

CFTR-encoding pseudouridine- or nucleoside-modified RNA is delivered, as described in Example 16, to the lungs of a cystic fibrosis animal model, and its effect on the disease is assessed as described in Scholte BJ, et al (Animal models of cystic fibrosis. J Cyst Fibros 2004; 3 Suppl2: 183-90) or Copreni E, et al, Lentivirus-mediated gene transfer to the respiratory

epithelium: a promising approach to gene therapy of cystic fibrosis. GeneTher 2004; 11 Suppl 1: S67-75). Administration of the RNA ameliorates cystic fibrosis.

In additional experiments, modified mRNA molecules of the present invention are used to deliver to the lungs, other recombinant proteins of therapeutic value, e.g. via an inhaler that delivers RNA.

EXAMPLE 27

TREATMENT OF XLA BY DELIVERY OF ADA-ENCODING MODIFIED mRNA MOLECULES TO HEMATOPOIETIC CELLS

ADA-encoding pseudouridine-or nucleoside-modified RNA is delivered to the hematopoietic cells of an X-linked agammaglobulinemia animal model, and its effect on the disease is assessed as described in Tanaka M, Gunawan F, et al, Inhibition of heart transplant injury and graft coronary artery disease after prolonged organ ischemia by selective protein kinase C regulators. J Thorac Cardiovasc Surg 2005;129(5): 1160-7) or Zonta S, Lovisetto F, et al, Uretero-neocystostomy in a swine model of kidney transplantation: a new technique. J Surg Res. 2005 Apr;124(2):250-5). Administration of the RNA is found to improve XLA.

EXAMPLE 28

PREVENTION OF ORGAN REJECTION BY DELIVERY OF IMMUNO-MODULATORY PROTEIN-ENCODING MODIFIED mRNA MOLECULES TO A TRANSPLANT SITE

Pseudouridine-or nucleoside-modified RNA encoding a cytokine, a chemokine, or an interferon IS (e.g. IL-4, IL-13, IL-I0, or TGF-β) is delivered to the transplant site of an organ transplant rejection animal model, and its effect on the incidence of rejection is assessed as described in Yu PW, Tabuchi R S et al, Sustained correction of B-cell development and function in a murine model of X-linked agammaglobulinemia (XLA) using retroviral-mediated gene transfer. Blood. 2004 104(5): 1281-90) or Satoh M, Mizutani A et al, X-linked immunodeficient mice spontaneously produce lupus-related anti20 RNA helicase A autoantibodies, but are resistant to pristane-induced lupus. Int Immunol 2003, 15(9):1117-24). Administration of the RNA reduces incidence of transplant rejection.

EXAMPLE 29

TREATMENT OF NIEMANN-PICK DISEASE, MUCOPOLYSACCHARIDOSIS, AND OTHER INBORN METABOLIC ERRORS BY DELIVERY OF MODIFIED mRNA TO BODY TISSUES

Sphingomyelinase-encoding pseudouridine-or nucleoside-modified RNA is delivered to the lung, brain, or other tissue of Niemann-Pick disease Type A and B animal models, and its effect on the disease is assessed as described in Passini MA, Macauley SL, et al, AAV vector-mediated correction of brain pathology in a mouse model of Niemann-Pick A disease. Mol Ther 2005;11(5): 754-62) or Buccoliero R, Ginzburg L, et al, Elevation of lung surfactant phosphatidylcholine in mouse models of Sandhoff and of Niemann-Pick A disease. J Inherit Metab Dis 2004;27(5): 641-8). Administration of the RNA is found to improve the disease.

Pseudouridine-or nucleoside-modified RNA encoding alpha-L-iduronidase, iduronate-2--sulfatase, or a related enzyme is delivered to the body tissues of a mucopolysaccharidosis animal model of, and its effect on the disease is assessed as described in Simonaro CM, D'Angelo M, et al, Joint and bone disease in mucopolysaccharidoses VI and VII: identification of new therapeutic targets and biomarkers using animal models. Pediatr Res 2005;57(5 Pt 1): 701-7) or McGlynn R, Dobrenis K, et al, Differential subcellular localization of cholesterol, gangliosides, and glycosaminoglycans in murine models of mucopolysaccharide storage disorders. J Comp Neurol 2004 20;480(4): 415-26). Administration of the RNA ameliorates the disease.

In additional experiments, modified mRNA molecules of the present invention are used to provide clotting factors (e.g. for hemophiliacs). In additional experiments, modified mRNA molecules of the present invention are used to provide acid-b-glucosidase for treating Gaucher's. In additional experiments, modified mRNA molecules of the present invention are used to provide alpha-galactosidase A for treating Fabry's diseases. In additional experiments, modified mRNA molecules of the present invention are used to provide cytokines for treatment of infectious diseases.

In additional experiments, modified mRNA molecules of the present invention are used to correct other inborn errors of metabolism, by administration of mRNA molecules encoding, e.g. ABCA4; ABCD3; ACADM; AGL; AGT; ALDH4Al; ALPL; AMPD1; APOA2; AVSD1; BRCD2; C1QA; C1QB; C1QG; C8A; C8B; CACNA1S; CCV; CD3Z; CDC2L1; CHML; CHS1; CIAS1; CLCNKB; CMD1A; CMH2; CMM; COL11AI; COL8A2; COL9A2; CPT2; CRB1;

CSE; CSF3R; CTPA; CTSK; DBT; DIO1; DISC1; DPYD; EKV; ENO1; ENO1P; EPB41; EPHX1; F13B; F5; FCGR2A; FCGR2B; FCGR3A; FCHL; FH; FMO3; FMO4; FUCA1; FY; GALE; GBA; GFND; GJA8; GJB3; GLC3B; HF1; HMGCL; HPC1; HRD; HRPT2; HSD3B2; HSPG2; KCNQ4; KCS; KIF1B; LAMB3; LAMC2; LGMD1B; LMNA; LOR; MCKD1; MCL1; MPZ; MTHFR; MTR; MUTYH; MYOC; NB; NCF2; NEM1; NPHS2; NPPA; NRAS; NTRK1; OPTA2; PBX1; PCHC; PGD; PHA2A; PHGDH; PKLR; PKP1; PLA2G2A; PLOD; PPOX; PPT1; PRCC; PRG4; PSEN2; PTOS1; REN; RFX5; RHD; RMD1; RPE65; SCCD; SERPINC1; SJS1; SLC19A2; SLC2A1; SPG23; SPTA1; TAL1; TNFSF6; TNNT2; TPM3; TSHB; UMPK; UOX; UROD; USH2A; VMGLOM; VWS; WS2B; ABCB11; ABCG5; ABCG8; ACADL; ACP1; AGXT; AHHR; ALMS1; ALPP; ALS2; APOB; BDE; BDMR; BJS; BMPR2; CHRNA1; CMCWTD; CNGA3; COL3A1; COL4A3; COL4A4; COL6A3; CPS1; CRYGA; CRYGEP1; CYP1B1; CYP27A1; DBI; DES; DYSF; EDAR; EFEMP1; EIF2AK3; ERCC3; FSHR; GINGF; GLC1B; GPD2; GYPC; HADHA; HADHB; HOXD13; HPE2; IGKC; IHH; IRS1; ITGA6; KHK; KYNU; LCT; LHCGR; LSFC; MSH2; MSH6; NEB; NMTC; NPHP1; PAFAH1P1; PAX3; PAX8; PMS1; PNKD; PPH1; PROC; REGIA; SAG; SFTPB; SLC11A1; SLC3A1; SOS1; SPG4; SRD5A2; TCL4; TGFA; TMD; TPO; UGT1A@; UV24; WSS; XDH; ZAP70; ZFHX1B; ACAA1; AGS1; AGTR1; AHSG; AMT; ARMET; BBS3; BCHE; BCPM; BTD; CASR; CCR2; CCR5; CDL1; CMT2B; COL7A1; CP; CPO; CRY; CTNNB1; DEM; ETM1; FANCD2; F1H; FOXL2; GBE1; GLB1; GLC1C; GNAI2; GNAT1; GP9; GPX1; HGD; HRG; ITIH1; KNG; LPP; LRS1; MCCC1; MDS1; MHS4; MITF; MLH1; MYL3; MYMY; OPA1; P2RY12; PBXPI; PCCB; POU1FI; PPARG; PROS1; PTHR1; RCA1; RHO; SCA7; SCLC1; SCN5A; SI; SLC25A20; SLC2A2; TF; TGFBR2; THPO; THRB; TKT; TM4SF1; TRH; UMPS; UQCRC1; USH3A; VHL; WS2A; XPC; ZNF35; ADH1B; ADH1C; AFP; AGA; AIH2; ALB; ASMD; BFHD; CNGA1; CRBM; DCK; DSPP; DTDP2; ELONG; ENAM; ETFDH; EVC; F11; FABP2; FGA;FGB; FGFR3; FGG; FSHMD1A; GC; GNPTA; GNRHR; GYPA; HCA; HCL2; HD; HTN3; HVBS6; IDUA; IF; JPD; KIT; KLKB1; LQT4; MANBA; MLLT2; MSX1; MTP; NR3C2; PBT; PDE6B; PEE1; PITX2; PKD2; QDPR; SGCB; SLC25A4; SNCA; SOD3; STATH; TAPVR1; TYS; WBS2; WFS1; WHCR; ADAMTS2; ADRB2; AMCN; AP3BI; APC; ARSB; B4GALT7; BHR1; C6; C7; CCAL2; CKN1; CMDJ; CRHBP; CSF1R; DHFR; DIAPH1; DTR; EOS; EPD; ERVR; F12; FBN2; GDNF; GHR; GLRA1; GM2A; HEXB; HSD17B4; ITGA2; KFS; LGMD1A; LOX; LTC4S; MAN2A1; MCC; MCCC2; MSH3; MSX2; NR3C1;

PCSK1; PDE6A; PFBI; RASA1; SCZD1; SDHA; SGCD; SLC22A5; SLC26A2; SLC6A3; SM1; SMA@; SMN1; SMN2; SPINK5; TCOF1; TELAB1; TGFBI; ALDH5A1; ARG1; AS; ASSP2; BCKDHB; BF; C2; C4A; CDKN1A; COL10A1; COL11A2; CYP21A2; DYX2; EJM1; ELOVL4; EPM2A; ESR1; EYA4; F13A1; FANCE; GCLC; GJA1; GLYS1; GMPR; GSE; HCR; HFE; HLA-A; HLA-DPB1; HLA-DRA; HPFH; ICS1; IDDM1; IFNGR1; IGAD1; IGF2R; ISCW; LAMA2; LAP; LCA5; LPA; MCDR1; MOCS1; MUT; MYB; NEU1; NKS1; NYS2; OA3; OODD; OFC1; PARK2; PBCA; PBCRA1; PDB1; PEX3; PEX6; PEX7; PKHD1; PLA2G7; PLG; POLH; PPAC; PSORS1; PUJO; RCD1; RDS; RHAG; RP14; RUNX2; RWS; SCA1; SCZD3; SIASD; SOD2; ST8; TAP1; TAP2; TFAP2B; TNDM; TNF; TPBG; TPMT; TULP1; WISP3; AASS; ABCB1; ABCB4; ACHE; AQP1; ASL; ASNS; AUTS1; BPGM; BRAF; C7orf2; CACNA2D1; CCM1; CD36; CFTR; CHORDOMA; CLCN1; CMH6; CMT2D; COL1A2; CRS; CYMD; DFNA5; DLD; DYT11; EEC1; ELN; ETV1; FKBP6; GCK; GHRHR; GHS; GLI3; GPDS1; GUSB; HLXB9; HOXA13; HPFH2; HRX; IAB; IMMP2L; KCNH2; LAMB1; LEP; MET; NCF1;NM; OGDH; OPN1SW; PEX1; PGAM2; PMS2; PON1; PPP1R3A; PRSS1; PTC; PTPN12; RP10; RP9; SERPINE1; SGCE; SHFM1; SHH; SLC26A3; SLC26A4; SLOS; SMAD1; TBXAS1; TWIST; ZWS1; ACHM3; ADRB3; ANKI; CA1; CA2; CCAL1; CLN8; CMT4A; CNGB3; COH1; CPP; CRH; CYP11B1; CYP11B2; DECR1; DPYS; DURS1; EBS1; ECA1; EGI; EXT1; EYA1; FGFR1; GNRH1; GSR; GULOP; HR; KCNQ3; KFM; KWE; LGCR; LPL; MCPH1; MOS; MYC; NAT1; NAT2; NBS1; PLAT; PLEC1; PRKDC; PXMP3; RP1; SCZD6; SFTPC; SGM1; SPG5A; STAR; TG; TRPS1; TTPA; VMD1; WRN; ABCA1; ABL1; ABO; ADAMTS13; AK1; ALAD; ALDH1A1; ALDOB; AMBP; AMCD1; ASS; BDMF; BSCL; C5; CDKN2A; CHAC; CLA1; CMD1B; COL5A1; CRAT; DBH; DNAI1; DYS; DYT1; ENG; FANCC; FBP1; FCMD; FRDA; GALT; GLDC; GNE; GSM1; GSN; HSD17B3; HSN1; IBM2; INVS; JBTS1; LALL; LCCS1; LCCS; LGMD2H; LMX1B; MLLT3; MROS; MSSE; NOTCH1; ORM1; PAPPA; PIP5K1B; PTCH; PTGS1; RLN1; RLN2; RMRP; ROR2; RPD1; SARDH; SPTLC1; STOM; TDFA; TEK; TMC1; TRIM32; TSC1; TYRP1; XPA; CACNB2; COL17A1; CUBN; CXCL12; CYP17; CYP2C19; CYP2C9; EGR2; EMX2; ERCC6; FGFR2; HK1; HPSI; IL2RA; LGI1; LIPA; MAT1A; MBL2; MKI67; MXI1; NODAL; OAT; OATL3; PAX2; PCBD; PEO1; PHYH; PNL1P; PSAP;PTEN; RBP4; RDPA; RET; SFTPA1; SFTPD; SHFM3; SIAL; THC2; TLX1; TNFRSF6; UFS; UROS; AA; ABCC8; ACAT1; ALX4; AMPD3; ANC; APOA1; APOA4; APOC3; ATM; BSCL2; BWS; CALCA; CAT; CCND1;

CD3E; CD3G; CD59; CDKN1C; CLN2; CNTF; CPT1A; CTSC; DDB1; DDB2; DHCR7; DLAT; DRD4; ECB2; ED4; EVR1; EXT2; F2; FSHB; FTH1; G6PT1; G6PT2; GIF; HBB; HBBP1; HBD; HBE1; HBG1; HBG2; HMBS; HND; HOMG2; HRAS; HVBS1; IDDM2; IGER; INS; JBS; KCNJ11; KCNJ1; KCNQ1; LDHA; LRP5; MEN1; MLL; MYBPC3; MYO7A; NNO1; OPPG; OPTB1; PAX6; PC; PDX1; PGL2; PGR; PORC; PTH; PTS; PVRL1; PYGM; RAG1; RAG2; ROM1; RRAS2; SAA1; SCA5; SCZD2; SDHD; SERPING1; SMPD1; TCIRG1; TCL2; TECTA; TH; TREH; TSG101; TYR; USH1C; VMD2; VRN1; WT1; WT2; ZNF145; A2M; AAAS; ACADS; ACLS; ACVRL1; ALDH2; AMHR2; AOM; AQP2; ATD; ATP2A2; BDC; C1R; CD4; CDK4; CNA1; COL2A1; CYP27B1; DRPLA; ENUR2; FEOM1; FGF23; FPF; GNB3; GNS; HAL; HBP1; HMGA2; HMN2; HPD; IGF1; KCNA1; KERA; KRAS2; KRT1; KRT2A; KRT3; KRT4; KRT5; KRT6A; KRT6B; KRTHB6; LDHB; LYZ; MGCT; MPE; MVK; MYL2; OAP; PAH; PPKB; PRB3; PTPN11; PXR1; RLS; RSN; SAS; SAX1; SCA2; SCNN1A; SMAL; SPPM; SPSMA; TBX3; TBX5; TCF1; TPI1; TSC3; ULR; VDR; VWF; ATP7B; BRCA2; BRCD1; CLN5; CPB2; ED2; EDNRB; ENUR1; ERCC5; F10; F7; GJB2; GJB6; IPF1; MBS1; MCOR; NYS4; PCCA; RB1; RHOK; SCZD7; SGCG; SLC10A2; SLC25A15; STARP1; ZNF198; ACHM1; ARVDI; BCH; CTAA1; DAD1; DFNB5; EML1; GALC; GCH1; IBGC1; IGH@; IGHC group; IGHG1; IGHM; IGHR; IV; LTBP2; MCOP; MJD; MNG1 MPD1; MPS3C; MYH6; MYH7; NP; NPC2; PABN1; PSEN1 PYGL; RPGRIP1; SERPINA1; SERPINA3; SERPINA6; SLC7A7; SPG3A; SPTB; TCL1A; TGMI; TITF1; TMIP; TRA@; TSHR; USH1A; VP; ACCPN; AHO2; ANCR; B2M; BBS4; BLM; CAPN3; CDAN1; CDAN3; CLN6; CMH3; CYP19; CYP1A1; CYP1A2; DYX1; EPB42; ETFA; EYCL3; FAH; FBN1; FES; HCVS; HEXA; IVD; LCS1; LIPC; MYO5A; OCA2; OTSC1; PWCR; RLBP1; SLC12A1; SPG6; TPM1; UBE3A; WMS; ABCC6; ALDOA; APRT; ATP2A1; BBS2; CARD15; CATM; CDH1; CETP; CHST6; CLN3; CREBBP; CTH; CTM; CYBA; CYLD; DHS; DNASE1; DPEP1; ERCC4; FANCA; GALNS; GAN; HAGH; HBA1; HBA2; HBHR; HBQ1; HBZ; HBZP; HP; HSD11B2; IL4R; LIPB; MC2R; MEFV; MHC2TA; MLYCD; MMVP1; PHKB; PHKG2; PKD1; PKDTS; PMM2; PXE; SALL1; SCA4; SCNN1B; SCNN1G; SLC12A3; TAT; TSC2; VDI; WT3; ABR; ACACA; ACADVL; ACE; ALDH3A2; APOH; ASPA; AXIN2; BCL5; BHD; BLMH; BRCA1; CACD; CCA1; CCZS; CHRNB1; CHRNE; CMT1A; COL1A1; CORD5; CTNS; EPX; ERBB2; G6PC; GAA; GALK1; GCGR; GFAP; GH1; GH2; GP1BA; GPSC; GUCY2D; ITGA2B; ITGB3; ITGB4; KRT10; KRT12; KRT13;

KRT14; KRT14L1; KRT14L2; KRT14L3; KRT16; KRT16L1; KRT16L2; KRT17; KRT9; MAPT; MDB; MDCR; MGI; MHS2; MKS1; MPO; MYO15A; NAGLU; NAPB; NF1; NME1; P4HB; PAFAH1B1; PECAM1; PEX12; PHB; PMP22; PRKAR1A; PRKCA; PRKWNK4; PRP8; PRPF8; PTLAH; RARA; RCV1; RMSA1; RP17; RSS; SCN4A; SERPINF2; SGCA; SGSH; SHBG; SLC2A4; SLC4A1; SLC6A4; SMCR; SOST; SOX9; SSTR2; SYM1; SYNS1; TCF2; THRA; TIMP2; TOC; TOP2A; TP53; TRIM37; VBCH; ATP8B1; BCL2; CNSN; CORD1; CYB5; DCC; F5F8D; FECH; PEO; LAMA3; LCFS2; MADH4; MAFD1; MC2R; MCL; MYP2; NPC1; SPPK; TGFBRE; TGIF; TTR; AD2; AMH; APOC2; APOE; ATHS; BAX; BCKDHA; BCL3; BFIC; C3; CACNA1A; CCO; CEACAM5; COMP; CRX; DBA; DDU; DFNA4; DLL3; DM1; DMWD; E11S; ELA2; EPOR; ERCC2; ETFB; EXT3; EYCL1; FTL; FUT1; FUT2; FUT6; GAMT; GCDH; GPI; GUSM; HB1; HCL1; HHC2; HHC3; ICAM3; INSR; JAK3; KLK3; LDLR; LHB; LIG1; LOH19CR1; LYL1; MAN2B1; MCOLN1; MDRV; MLLT1; NOTCH3; NPHS1; OFC3; OPA3; PEPD; PRPF31; PRTN3; PRX; PSG1; PVR; RYR1; SLC5A5; SLC7A9; STK11; TBXA2R; TGFB1; TNNI3; TYROBP; ADA; AHCY; AVP; CDAN2; CDPD1; CHED1; CHED2; CHRNA4; CST3; EDN3; EEGV1; FTLL1; GDF5; GNAS; GSS; HNF4A; JAG1; KCNQ2; MKKS; NBIA1; PCK1; PI3; PPCD; PPGB; PRNP; THBD; TOP1; AIRE; APP; CBS; COL6A1; COL6A2; CSTB; DCR; DSCR1; FPDMM; HLCS; HPE1; ITGB2; KCNE1; KNO; PRSS7; RUNX1; SOD1; TAM; ADSL; ARSA; BCR; CECR; CHEK2; COMT; CRYBB2; CSF2RB; CTHM; CYP2D6; CYP2D7P1; DGCR; DIA1; EWSR1; GGT1; MGCR; MN1; NAGA; NF2; OGS2; PDGFB; PPARA; PRODH; SCO2; SCZD4; SERPIND1; SLC5A1; SOX10; TCN2; TIMP3; TST; VCF; ABCD1; ACTL1; ADFN; AGMX2; AHDS; AIC; AIED; AIH3; ALAS2; AMCD; AMELX; ANOP1; AR; ARAF1; ARSC2; ARSE; ARTS; ARX; ASAT; ASSP5; ATP7A; ATRX; AVPR2; BFLS; BGN; BTK; BZX; C1HR; CACNA1F; CALB3; CBBM; CCT; CDR1; CFNS; CGF1; CHM; CHR39C; CIDX; CLA2; CLCN5; CLS; CMTX2; CMTX3; CND; COD1; COD2; COL4A5; COL4A6; CPX; CVD1; CYBB; DCX; DFN2; DFN4; DFN6; DHOF; DIAPH2; DKC1; DMD; DSS; DYT3; EBM; EBP; ED1; ELK1; EMD; EVR2; F8; F9; FCP1; FDPSL5; FGD1; FGS1; FMR1; FMR2; G6PD; GABRA3; GATA1; GDI1; GDXY; GJB1; GK; GLA; GPC3; GRPR; GTD; GUST; HMS1; HPRT1; HPT; HTC2; HTR2C; HYR; IDS; IHG1; IL2RG; INDX; IP1; IP2; JMS; KAL1; KFSD; L1CAM; LAMP2; MAA; MAFD2; MAOA; MAOB; MCF2; MCS; MEAX; MECP2; MF4; MGC1; MIC5; MID1; MLLT7; MLS; MRSD; MRX14; MRX1; MRX20; MRX2; MRX3; MRX40;

MRXA; MSD; MTM1; MYCL2; MYP1; NDP; NHS; NPHL1; NR0B1; NSX; NYS1; NYX; OA1; OASD; OCRL; ODT1; OFD1; OPA2; OPD1; OPEM; OPN1LW; OPN1MW; OTC; P3; PDHA1; PDR; PFC; PFKFB1; PGK1; PGK1P1; PGS; PHEX; PHKA1; PHKA2; PHP; PIGA; PLP1; POF1; POLA; POU3F4; PPMX; PRD; PRPS1; PRPS2; PRS; RCCP2; RENBP; RENS1; RP2; RP6; RPGR; RPS4X; RPS6KA3; RS1; S11; SDYS; SEDL; SERPINA7; SH2D1A; SHFM2; SLC25A5; SMAX2; SRPX; SRS; STS; SYN1; SYP; TAF1; TAZ; TBX22; TDD; TFE3; THAS; THC; TIMM8A; TIMP1; TKCR; TNFSF5; UBE1; UBE2A; WAS; WSN; WTS; WWS; XIC; XIST; XK; XM; XS; ZFX; ZIC3; ZNF261; ZNF41; ZNF6; AMELY; ASSP6; AZF1; AZF2; DAZ; GCY; RPS4Y; SMCY; SRY; ZFY; ABAT; AEZ; AFA; AFD1; ASAH1; ASD1; ASMT; CCAT; CECR9; CEPA; CLA3; CLN4; CSF2RA; CTS1; DF; DIH1; DWS; DYT2; DYT4; EBR3; ECT; EEF1A1L14; EYCL2; FANCB; GCSH; GCSL; GIP; GTS; HHG; HMI; HOAC; HOKPP2; HRPT1; HSD3B3; HTC1; HV1S; ICHQ; ICR1; ICR5; IL3RA; KAL2; KMS; KRT18; KSS; LCAT; LHON; LIMM; MANBB; MCPH2; MEB; MELAS; MIC2; MPFD; MS; MSS; MTATP6; MTCO1; MTCO3; MTCYB; MTND1; MTND2; MTND4; MTND5; MTND6; MTRNR1; MTRNR2; MTTE; MTTG; MTTI; MTTK; MTTL1; MTTL2; MTTN; MTTP; MTTS1; NAMSD; OCD1; OPD2; PCK2; PCLD; PCOS1; PFKM; PKD3; PRCA1; PRO1; PROP1; RBS; RFXAP; RP; SHOX; SLC25A6; SPG5B; STO; SUOX; THM; or TTD.

EXAMPLE 30

TREATMENT OF VASOSPASM BY DELIVERY OF iNOS-ENCODING MODIFIED mRNA MOLECULES TO BODY TISSUES

Inducible nitric oxide synthase (iNOS)-encoding pseudouridine-or nucleoside-modified RNA is delivered to the vascular endothelium of vasospasm animals models (e.g. subarachnoid hemorrhage), and its effect on the disease is assessed as described in Pradilla G, Wang PP, et al, Prevention of vasospasm by anti-CD11/CD18 monoclonal antibody therapy following subarachnoid hemorrhage in rabbits. J Neurosurg 2004;101(1): 88-92) or Park S, Yamaguchi M, et al, Neurovascular protection reduces early brain injury after subarachnoid hemorrhage. Stroke 2004;35(10): 2412-7). Administration of the RNA ameliorates the disease.

EXAMPLE 31

RESTORATION OF HAIR GROWTH BY DELIVERY OF MODIFIED mRNA ENCODING AN IMMUNOSUPPRESSIVE PROTEIN

Pseudouridine-or nucleoside-modified RNA encoding a telomerase or an immunosuppressive protein (e.g. α-MSH, TGF-β 1, or IGF-I is delivered to hair follicles of animals used as models of hair loss or balding, and its effect on hair growth is assessed as described in Jiang J, Tsuboi R, et al, Topical application of ketoconazole stimulates hair growth in C3H/HeN mice. J Dermatol 2005;32(4): 243-7) or McElwee KJ, Freyschmidt-Paul P, et al, Transfer of CD8(+) cells induces localized hair loss whereas CD4(+)/CD25(-) cells promote systemic alopecia areata and CD4(+)/CD25(+) cells blockade disease onset in the C3H/HeJ mouse model. J Invest Dermatol 2005;124(5): 947-57). Administration of the RNA restores hair growth.

EXAMPLE 32

SYNTHESIS OF AN *IN VITRO*-TRANSCRIBED RNA MOLECULE WITH ALTERED NUCLEOSIDES CONTAINING AN siRNA

A double-stranded RNA (dsRNA) molecule comprising pseudouridine or a modified nucleoside and further comprising a small interfering RNA (siRNA) or short hairpin RNA (shRNA) is synthesized by the following procedure: Complementary RNA strands with the desired sequence containing uridine or 1 or more modified nucleosides are synthesized by *in vitro* transcription (e.g. by T7, SP6, or T3 phage RNA polymerase) as described in Example 5. dsRNA molecules exhibit reduced immunogenicity. In other experiments, the dsRNA molecules are designed to be processed by a cellular enzyme to yield the desired siRNA or shRNA. Because dsRNA molecules of several hundred nucleotides are easily synthesized, each dsRNA may also be designed to contain several siRNA or shRNA molecules, to facilitate delivery of multiple siRNA or shRNA to a single target cell.

EXAMPLE 33

USE OF AN IN VITRO-TRANSCRIBED RNA MOLECULE WITH ALTERED NUCLEOSIDES TO DELIVER siRNA

The dsRNA molecule of the previous Example is complexed with a transfection reagent (e.g a cationic transfection reagent, a lipid-based transfection reagent, a protein-based transfection reagent, a polyethyleneimine based transfection reagent, or calcium phosphate) and delivered to a target cell of interest. Enzymes in or on the surface of the target cell degrade the dsRNA to the desired siRNA or shRNA molecule(s). This method effectively silences transcription of 1 or more cellular genes corresponding to the siRNA or shRNA sequence(s).

EXAMPLE 34

TESTING THE EFFECT OF ADDITIONAL NUCLEOSIDE MODIFICATIONS ON RNA IMMUNOGENICITY AND EFFICIENCY OF TRANSLATION

Additional nucleoside modifications are introduced into *in vitro*-transcribed RNA, using the methods described above in Examples 5 and 10, and their effects on immunogenicity translation efficiency are tested as described in Examples 4-11 and 12-18, respectively. Certain additional modifications are found to decrease immunogenicity and enhance translation. These modifications are additional embodiments of methods and compositions ofthe present invention.

Modifications tested include, e.g.:

m^1A; m^2A; Am; ms^2m^6A; i^6A; ms^2i6A; io^6A; ms^2i0^6A; g^6A; t^6A; ms^2t^6A; m^6t^6A; hn^6A; ms^2hn^6A; Ar(p); I; m^1I; m^1Im; m^3C; Cm; S^2C; ac^4C; f^5c; m^5Cm; ac^4Cm; k^2c; m^1G; m^2G; m^7G; Gm; $m^2{}_2G$; m^2Gm; $m^2{}_2Gm$; Gr(p); yW; o_2yW; OHyW; OHyW*; imG; mimG; Q; oQ; galQ; manQ; $preQ_0$; $preQ_1$; G^+; D; m^5Um', $m^1\Psi$; Ψm; S^4U; ·m^5s^2U', S^2Um, ' acp^3U·, ho^5u., mo^5U·; cmo^5U·; $mcmo^5U$; chm^5U; $mchm^5U$·; mcm^5U; mcm^5Um; mcm^5s^2U; nm^5s^2U; mnm^5U; mnm^5s^2U; mnm^5se^2U; ncm^5U; ncm^5Um; $cmnm^5U$; $cmnm^5Um$; $cmnm^5s^2U$; $m^6{}_2A$; Im; m^4C; m^4Cm; hm^5C; m^3U; $m^1acp3\Psi$; cm^5U; m^6Am; $m^6{}_2Am$; $m^{2,7}G$; $m^{2,2,7}G$; m^3Um; m^5D; $m^3\Psi$; f^5Cm; m^1Gm; m^1Am; Tm^5U; Tm^5s^2U; imG-14; imG2; and ac^6A.

Materials and Methods for Examples 35-38

HPLC purification of RNA: mRNA produced by T7 polymerase transcription was purified by HPLC using a column matrix of alkylated nonporous polystyrene-divinylbenzene (PS-DVB) copolymer microspheres (2.1 μm) (21 mm x 100 mm column) and a buffer system of Triethylammonium acetate (TEAA) with an acetonitrile gradient. Buffer A contained 0.1 M TEAA and Buffer B contained 0.1 M TEAA and 25% acetonitrile. Columns were equilibrated with 38% Buffer B in Buffer A, loaded with RNA, and then run with a linear gradient to 55% Buffer B over 30 minutes at 5 ml/minute. Fractions corresponding to the desired peak were collected. RNA analyses were performed with the same column matrix and buffer system, but using a 7.8 mm x 50 mm column at 1.0 ml/min and a gradient duration of 25 minutes.

RNA isolation from column fractions: Collected fractions were combined, and first their RNA content was concentrated using Amicon *Ultra-15 centrifugal filter units with 30K* membrane (Millipore). The filter device was filled with 15 ml sample and spin at 4,000x g for 10 min (4°C) in a Thermo Scientific Sorvall ST16R centrifuge using swinging bucket rotor. Under these conditions, ~98% of the solvent volume can be removed. When the collected fractions had a volume over 15 ml, the filter unit was reused by filling up with additional column fractions and centrifuging again until all of the RNA was in one tube. To remove the salt and solvent from the concentrated RNA, nuclease free water was added (up to 15 ml) and the filter device was spun again. The process of "washing out" was repeated until the concentration of the acetonitrile was <0.001%. The desalted and solvent-free sample was removed from the filtering device and the RNA was recovered by overnight precipitation at -20 °C in NaOAc (0.3 M, pH 5.5), isopropanol (1 volume) and glycogen (3 μl). The precipitated RNA was collected, washed twice with ice-cold 75% ethanol and reconstituted in water.

dsRNA dot blot: RNA (25-100 ng) was blotted onto a nitrocellulose membrane, allowed to dry, blocked with 5% non-fat dried milk in TBS buffer supplemented with 0.05% Tween-20 (TBS-T), and incubated with dsRNA-specific mAb J2 or K1 (English & Scientific Consulting) for 60 minutes. Membranes were washed 6 times with TBS-T and then reacted with HRP-conjugated donkey anti-mouse antibody (Jackson Immunology). After washing 6 times, dsRNA was

detected with the addition of SuperSignal West Pico Chemiluminescent substrate (Pierce) and image capture for 30 seconds to 2 minutes on a Fujifilm LAS1000 digital imaging system.

Dendritic cell generation: Monocytopheresis samples were obtained from normal volunteers through an IRB-approved protocol. Human DCs were produced by treating monocytes with GM-CSF (50 ng/ml) + IL-4 (100 ng/ml) (R&D Systems) in AIM V medium (Invitrogen) for 7 days. On days 3 and 6, a 50% volume of new medium with cytokines was added.

Murine DC were generated by isolating bone marrow mononuclear cells from Balb/c mice and culturing in RPMI + 10% FBS medium supplemented with murine GM-CSF (20 ng/ml, Peprotech). On days 3 and 6, a 50% volume of new medium with GM-CSF was added. Non-adherent cells were used after 7 days of culture.

Lipofectin complexing of RNA: Stock phosphate buffer was added to serum-free DMEM to give final concentrations of 20 mM potassium phosphate and 100 ng/ml BSA, pH 6.4. For 3 wells of a 96-well plate, lipofectin complexed RNA was prepared in the following ratios: 2.4 µl of lipofectin was added to 21.3 µl serum-free DMEM medium with phosphate buffer and incubated at room temperature for 10 minutes. Then, 0.75 µg of RNA in 9.9 µl serum-free DMEM was added and the mixture was incubated for 10 additional minutes at room temperature. Lastly, 116.4 ml serum-free DMEM was added to bring up the final volume to 150 ml. The mixture was vortexed.

TransIT complexing of RNA: For each well of a 96-well plate, 0.25 µg of RNA was added to 17.3 µl of serum-free DMEM on ice. TransIT mRNA reagent (0.3 ul) was added with vortexing followed by 0.2 µl of mRNA boost reagent and vortexing. Complexed RNA was added within 5 minutes of formation.

Cell transfections: For lipofectin complexed RNA, 50 µl (0.25 µg RNA/well) was added directly to cells, 5 x 10^5 per well. Transfected cells were incubated for 1 h at 37°C in a 5% CO2 incubator. The lipofectin-RNA mixture was removed and replaced with 200 µl pre-warmed serum containing medium. For TransIT complexed RNA, 17 µl of complex was added to cells, 5 x 10^5 per well, in 200 µl of serum containing medium. Cells were lysed in specific lysis media, 3

to 24 hr after transfection and firefly or renilla luciferase activity was measured with specific substrates in a luminometer.

RNA immunogenicity analysis: DCs (murine or human) in 96-well plates (5 x 10^5 cells/well) were treated with medium, or lipofectin or TransIT complexed RNA. Supernatant was harvested after 24 hr and subjected to analysis. The levels of IFN-α (TransIT delivered RNA) or TNF-α (Lipofectin delivered RNA) (Biosource International, Camarillo, CA) were measured in supernatants by ELISA. Cultures were performed in triplicate to quadruplicate and measured in duplicate.

EXAMPLE 35

This example examines the sequence and cell type dependency of translation of Ψ-, m5C, and Ψ/m5C-modified mRNA relative to the unmodified (U) RNA. Firefly or Renilla luciferase encoding mRNA with the indicated modifications were complexed to lipofectin and delivered to murine dendritic (A) and HEK293T (B) cells. Human DC were transfected with firefly or renilla luciferase-encoding mRNA with the indicated modifications complexed with TransIT (**C**). The data demonstrates that depending on the sequence of the RNA and the type of cell translating it, the optimal modification varies. It also shows that the enhancement caused by incorporation of modified nucleosides is substantially greater for primary cells compared to transformed cell lines. Enzyme activity in lysed cells was measured using specific substrates and measurement of light produced in a luminometer and expressed as fold increase compared to unmodified (U) RNA.

EXAMPLE 36

Phage polymerase transcription reactions used for the generation of mRNA results in large quantities of RNA of the correct size, but also contains contaminants. This is visualized by application of RNA to a reverse phase HPLC column that separates RNA based on size under denaturing conditions. Y-modified TEV-luciferase-A51 RNA was applied to the HPLC column in 38% Buffer B and subjected to a linear gradient of increasing Buffer B to 55%. The profile

demonstrated both smaller than expected and larger than expected contaminants. These results are shown in Figure 22.

EXAMPLE 37

HPLC purification increases translation of all types of modified or unmodified RNA, but Ψ-modified mRNA is translated best. The results are shown in Figure 23: (**A**) EPO encoding mRNA with the indicated modifications and with or without HPLC purification were delivered to murine DCs and EPO levels in the supernatant were measured 24 hr later. While m5C/Ymodified mRNA had the highest level of translation prior to HPLC purification, Ψ-modified mRNA had the highest translation after HPLC purification. (**B**) Human DCs were transfected with renilla encoding mRNA with the indicated modifications with or without HPLC purification. Similar to murine DCs and EPO mRNA, after HPLC purification, Y-modified mRNA had the highest level of translation.

EXAMPLE 38

Ψ, m5C, and Ψ/m5C-modified mRNA have low levels of immunogenicity that is reduced to control levels with HPLC purification. The results are shown in Figure 24: (**A**) Human DCs were transfected with RNA complexed to TransIT with the indicated modifications with or without HPLC purification. IFN-α levels were measured after 24 hr. HPLC purification increased the immunogenicity of unmodified RNA, which is dependent of the sequence, as other unmodified RNAs had similar levels of IFN-α or reduced levels. Ψ-modified RNA had unmeasurable levels of IFN-α similar to control treated DCs. (**B**) Ψ-modified RNA before (-) and after HPLC purification (P1 and P2) was analyzed for dsRNA using dot blotting with a monoclonal antibody specific for dsRNA (J2). Purification of RNA removed dsRNA contamination. (**C**) Ψ-modified RNA encoding iPS factors are immunogenic, which is removed by HPLC purification of the RNA.

MATERIALS AND METHODS FOR EXAMPLES 39-41

Cell Culture. Neonatal human epidermal keratinocytes (HEKn) cells (Invitrogen) were cultured in EpiLife Medium supplemented with keratinocyte growth supplement and Penicillin/Streptomycin (Invitrogen). All cells were grown at 37°C and 5% CO_2. The human iPS cells that were induced using methods described herein were transferred to hESC-qualified matrigel matrix (BD Biosciences) coated 6-well plates after transfection.

Constructions of Vectors. Generally the same as Examples 1-3.

mRNA Production. Generally the same as for Examples 1-3.

mRNA Purification and Analysis. In some experimental embodiments, the mRNA was purified by HPLC, column fractions were collected, and the mRNA fractions were analyzed for purity an immunogenicity as described in "Materials and Methods for Examples 35-38" and/or as described and shown for Figures 22-24. In some preferred experimental embodiments, purified RNA preparations comprising or consisting of mRNAs encoding one or more reprogramming factors which exhibited little or no immunogenicity were used for the experiments for reprogramming human somatic cells to iPS cells.

Reprogramming of Primary Keratinocytes. HEKn cells were plated at 1 x 10^5 cells/well of a 6-well dish in EpiLife medium and grown overnight. The cells were transfected with equal amounts of each reprogramming factor mRNA (KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2) or a subset of the factors using TransIT™ mRNA transfection reagent (MirusBio, Madison, WI). Three transfections were performed, every other day, with media changes every day. The day after the third transfection, the cells were trypsinized and plated in mTeSR1 medium (StemCell Technologies) onto matrigel-coated 6-well plates. The mTeSR cell medium was changed daily. Cells were maintained at 37°C with 5% CO_2. Plates were screened for morphology changes using an inverted microscope.

HEKn cells were also reprogrammed by a single transfection by electroporation with equal amounts of each reprogramming factor mRNA. The cells were plated directly onto matrigel-coated plates at a density of 1 x 10^5 cells per 6-well dish or 7.5 x 10^5 cells per 10 cm dish in mTeSR1 medium which was changed daily.

Immunofluorescence. Generally the same as Examples 1-3

Quantitative RT-PCR (qPCR) Cellular RNA was reverse-transcribed using standard methods and an oligo $d(T)_{21}$ primer from equivalent amounts of cellular RNA. Three messages

were amplified using gene-specific primers and real-time PCR using SYBR green detection and GAPDH normalization. Expression levels were determined in relation to the level of expression in the original HEKn cell line, and depicted as changes in cycle threshold (C_T) level.

EXAMPLE 39

This example describes development of a protocol for iPS cell generation from somatic keratinocytes. Equal amounts (by weight) of KLF4, c-MYC, OCT4, and SOX2 mRNAs were transfected into HEKn cells three times (once every other day) with TransIT™ mRNA Reagent. Medium was changed daily. The day after the third transfection, the cells were plated onto matrigel-coated dishes and grown in mTeSR1 cell medium. By 11 days after the first transfection the reprogrammed cell morphology began to appear (Fig. 28).

EXAMPLE 40

This example describes the characterization of cells resulting from transfection of primary keratinocytes with equal amounts of KLF4, LIN28, c-MYC, NANOG, OCT4, and SOX2 mRNAs. One million HEKn cells were electoporated once with 5 micrograms of each mRNA and plated on matrigel-coated 10 cm dishes in mTeSR1 cell medium. 15 Days after transfection the cells were fixed for immunofluorescence analysis. The resulting colonies were positive for the iPS markers KLF4, LIN28, SSEA4, TRA-1-60, and NANOG (Fig. 29).

EXAMPLE 41

This example describes expression differences between primary keratinocytes and keratinocytes reprogrammed with equal amounts of KLF4, c-MYC, OCT4, and SOX2 mRNAs. 7.5×10^5 HEKn cells were electoporated once with 3 or 5 micrograms of each mRNA and plated on matrigel-coated 10 cm dishes in mTeSR1 medium. Medium was changed daily. 13 days after transfection the cells were transferred to freshly-coated matrigel plates. 21 days after transfection, total cellular RNA was purified from untransfected HEKn cells and two wells of reprogrammed cells. An equal amounts of each cellular RNA was converted to cDNA and analyzed by qPCR. Increased levels of NANOG, CRIPTO and REX1 were detected by qPCR using message-specific primers (Fig. 30). These three messages have been shown to be elevated in iPS cells (Aasen T et al. 2008. Nature Biotech 26: 1276). None of these factors was

introduced into the cells by transfection; therefore the changes in expression are due to the influence of the reprogramming factors that were introduced.

EXAMPLE 42

Transdifferentiating Cells with mRNAs

Cells can be transdifferentiated using the purified mRNA preparations described herein, or the modified mRNAs described herein, or the purified mRNA preparations containing modified mRNAs described herein. In this Example, a purified RNA preparation containing OCT4 mRNA that has at least one pseudouridine or one 5-methycytidine is employed. Such purified and modified OCT4 mRNAs are substituted for the OCT4 encoding vectors in the protocol described by Szabo et al. (Nature 468: 521-528, 2010, which is herein incorporated by reference in its entirety as if fully set forth herein) and in the protocol described in Racila et al. (Gene Therapy, 1-10, 2010, herein incorporated by reference in its entirety as if fully set forth herein). In one embodiment of each of these methods, the purified RNA preparation comprises or consists of OCT 4 mRNA, wherein all of the uridine nucleosides are replaced by pseudouridine nucleosides. In one embodiment of each of these methods, the purified RNA preparation comprises or consists of OCT 4 mRNA, wherein all of the cytidine nucleosides are replaced by 5-methylcytidine nucleosides. In one embodiment of each of these methods, the purified RNA preparation comprises or consists of OCT 4 mRNA, wherein all of the uridine nucleosides are replaced by pseudouridine nucleosides and all of the cytidine nucleosides are replaced by 5-methylcytidine nucleosides. In preferred embodiments, the OCT4 mRNA is purified to be free of contaminating RNAs. The Racilla et al. reference describes a system in which human keratinocytes were transdifferentiated by being redirected to an alternative differentiation pathway. In particular, transient transfection of human skin keratinocytes with the transcription factor OCT4 was employed. After 2 days, these transfected cells displayed expression of endogenous embryonic genes and showed reduced genomic methylation. It was shown that such cells could be converted into neuronal and contractile mesenchymal cell types.

The Szabo et al. reference demonstrated the ability to generate progenitors and mature cells of the haematopoietic fate directly from human dermal fibroblasts without establishing pluripotency. In particular, ectopic expression of OCT4 activated haematopoietic transcription factors, together with specific cytokine treatment, allowed generation of cells expressing the pan-

leukocyte marker CD45. These unique fibroblast-derived cells gave rise to granulocytic, monocytic, megakaryocytic and erythroid lineages, and demonstrated in vivo engraftment capacity.

Besides the use of OCT4, both of these protocols also employed cytokines or growth factors, such as tranforming growth factor (TGF), PDGF-BB, stem cell factor (SCF), and FMS-like tyrosine kinase 3 ligand (Flt3L). Other growth factors and cytokines could be used, such as granulocyte-colony stimulating factor (G-CSF), IL-3, IL-6, erythropoietin, basic fibroblast growth factor (bFGF), insulin-like growth factor 2 (IGFII), and bone morphogenetic protein 4 (BMP-4). As such, in certain embodiments, the Racilla et al. or Szabo et al. protocols are repeated with the substitution of modified OCT4 mRNA (e.g., psuedouridine-modified and/or 5-methycytidine-modified), along with the use of the growth factors or cytokines recited above. In some embodiments, the cells are contacted with the cytokine and/or growth factor proteins that are used. In some other embodiments, the cells are contacted with modified mRNAs (e.g., modified mRNAs as described in the present application, e.g., e.g., psuedouridine-modified and/or 5-methycytidine-modified) that encode one or more of the cytokines and/or growth factors that are used in the transdifferentiation protocol. It will be clear from this description, that the present invention includes contacting a human or animal cell with a purified RNA preparation comprising or consisting of mRNA that encodes a reprogramming factor in order to transdifferentiate a cell having a first state of differentiation or phenotype to a cell having a second state of differentiation or phenotype.

REFERENCES

Aoi T, Yae K, Nakagawa M, Ichisaka T, Okita K, Takahashi K, Chiba T, Yamanaka S. 2008. Generation of pluripotent stem cells from adult mouse liver and stomach cells. Science 321: 699-702.

Banerjee AK. 1980. 5'-terminal cap structure in eucaryotic messenger ribonucleic acids. Microbiol Rev 44: 175-205.

Chan EM, et al. 2009. Live cell imaging distinguishes bona fide human iPS cells from partially reprogrammed cells. Nat Biotechnol 27: 1033-1037.

Ebert AD, Yu J, Rose FF, Jr., Mattis VB, Lorson CL, Thomson JA, Svendsen CN. 2009. Induced pluripotent stem cells from a spinal muscular atrophy patient. Nature 457: 277-280.

Edmonds M. 1990. Polyadenylate polymerases. Methods Enzymol 181: 161-170.

Feng, R et al. 2008. PU.1 and C/EBPa/b convert fibroblasts into macrophage-like cells. Proc. Natl Acad. Sci. USA 105: 6057–6062.

Gershon PD. 2000. (A)-tail of two polymerase structures. Nat Struct Biol 7: 819-821.

Gonzalez F, Barragan Monasterio M, Tiscornia G, Montserrat Pulido N, Vassena R, Batlle Morera L, Rodriguez Piza I, Izpisua Belmonte JC. 2009. Generation of mouse-induced pluripotent stem cells by transient expression of a single nonviral polycistronic vector. Proc Natl Acad Sci U S A 106: 8918-8922.

Graf T, Enver T. 2009. Forcing cells to change lineages. Nature 462: 587-594.

Grudzien E, Stepinski J, Jankowska-Anyszka M, Stolarski R, Darzynkiewicz E, Rhoads RE. 2004. Novel cap analogs for in vitro synthesis of mRNAs with high translational efficiency. RNA 10: 1479-1487.

Grudzien-Nogalska E, Jemielty J, Kowalska J, Darzynkiewicz E, Rhoads R. 2007. Phosphorothioate cap analogs stabilize mRNA and increase translational efficiency in mammalian cells. RNA 13: 1745-1755.

Higman MA, Bourgeois N, Niles EG. 1992. The vaccinia virus mRNA (guanine-N7-)-methyltransferase requires both subunits of the mRNA capping enzyme for activity. J Biol Chem 267: 16430-16437.

Higman MA, Christen LA, Niles EG. 1994. The mRNA (guanine-7-)methyltransferase domain of the vaccinia virus mRNA capping enzyme. Expression in Escherichia coli and structural and kinetic comparison to the intact capping enzyme. J Biol Chem 269: 14974-14981.

Huangfu D, Osafune K, Maehr R, Guo W, Eijkelenboom A, Chen S, Muhlestein W, Melton DA. 2008. Induction of pluripotent stem cells from primary human fibroblasts with only Oct4 and Sox2. Nat Biotechnol 26: 1269-1275.

Ieda, M et al. 2010. Direct reprogramming of fibroblasts into functional cardiomyocytes by defined factors. Cell 142: 375–386.

Jemielity J, Fowler T, Zuberek J, Stepinski J, Lewdorowicz M, Niedzwiecka A, Stolarski R, Darzynkiewicz E, Rhoads RE. 2003. Novel "anti-reverse" cap analogs with superior translational properties. RNA 9: 1108-1122.

Kariko K, Muramatsu H, Welsh FA, Ludwig J, Kato H, Akira S, Weissman D. 2008. Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability. Mol Ther 16: 1833-1840.

Krieg PA, Melton DA. 1984. Functional messenger RNAs are produced by SP6 in vitro transcription of cloned cDNAs. Nucleic Acids Res 12: 7057-7070.

Lee G, et al. 2009. Modelling pathogenesis and treatment of familial dysautonomia using patient-specific iPSCs. Nature 461: 402-406.

Mackie GA. 1988. Vectors for the synthesis of specific RNAs in vitro. Biotechnology 10: 253-267.

Maehr R, Chen S, Snitow M, Ludwig T, Yagasaki L, Goland R, Leibel RL, Melton DA. 2009. Generation of pluripotent stem cells from patients with type 1 diabetes. Proc Natl Acad Sci U S A 106: 15768-15773.

Martin SA, Paoletti E, Moss B. 1975. Purification of mRNA guanylyltransferase and mRNA (guanine-7-) methyltransferase from vaccinia virions. J Biol Chem 250: 9322-9329.

Myette JR, Niles EG. 1996. Domain structure of the vaccinia virus mRNA capping enzyme. Expression in Escherichia coli of a subdomain possessing the RNA 5'-triphosphatase and guanylyltransferase activities and a kinetic comparison to the full-size enzyme. J Biol Chem 271: 11936-11944.

Nakagawa M, Koyanagi M, Tanabe K, Takahashi K, Ichisaka T, Aoi T, Okita K, Mochiduki Y, Takizawa N, Yamanaka S. 2008. Generation of induced pluripotent stem cells without Myc from mouse and human fibroblasts. Nat Biotechnol 26: 101-106.

Okita K, Nakagawa M, Hyenjong H, Ichisaka T, Yamanaka S. 2008. Generation of mouse induced pluripotent stem cells without viral vectors. Science 322: 949-953.

Ozawa T, Kishi H, Muraguchi A. 2006. Amplification and analysis of cDNA generated from a single cell by 5'-RACE: application to isolation of antibody heavy and light chain variable gene sequences from single B cells. Biotechniques 40: 469-470, 472, 474 passim.

Peng ZH, Sharma V, Singleton SF, Gershon PD. 2002. Synthesis and application of a chain-terminating dinucleotide mRNA cap analog. Org Lett 4: 161-164.

Racila D et al. 2010. Transient expression of OCT 4 IS sufficient to allow human keratinocytes to change their differentiation pathway. Gene Therapy advance online publication, (October 28, 2010; doi:10.1038/gt.2010.148).

Shuman S. 1995. Capping enzyme in eukaryotic mRNA synthesis. Prog Nucleic Acid Res Mol Biol 50: 101-129.

Shuman. 2001. Structure, mechanism, and evolution of the mRNA capping apparatus. Prog Nucleic Acid Res Mol Biol 66: 1-40.

Shuman S, Surks M, Furneaux H, Hurwitz J. 1980. Purification and characterization of a GTP-pyrophosphate exchange activity from vaccinia virions. Association of the GTP-pyrophosphate exchange activity with vaccinia mRNA guanylyltransferase. RNA (guanine-7-)methyltransferase complex (capping enzyme). J Biol Chem 255: 11588-11598.

Stadtfeld M, Nagaya M, Utikal J, Weir G, Hochedlinger K. 2008. Induced pluripotent stem cells generated without viral integration. Science 322: 945-949.

Stepinski J, Waddell C, Stolarski R, Darzynkiewicz E, Rhoads RE. 2001. Synthesis and properties of mRNAs containing the novel "anti-reverse" cap analogs 7-methyl(3'-O-methyl)GpppG and 7-methyl (3'-deoxy)GpppG. RNA 7: 1486-1495.

Studier FW, Moffatt BA. 1986. Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. J Mol Biol 189: 113-130.

Szabo E, Rampalli S, Risueno RM, Schnerch A, Mitchell R, Fiebig-Comyn A, Levadoux-Martin M, Bhatia M. 2010. Direct conversion of human fibroblasts to multilineage blood progenitors. Nature. 468: 521-526.

Takahashi K, Yamanaka S. 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. Cell 126: 663-676.

Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. 2007. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell 131: 861-872.

Vierbuchen T et al. 2010. Direct conversion of fibroblasts to functional neurons by defined factors. Nature 463: 1035–1041.

Wang SP, Deng L, Ho CK, Shuman S. 1997. Phylogeny of mRNA capping enzymes. Proc Natl Acad Sci U S A 94: 9573-9578.

Wilusz J, Shenk T. 1988. A 64 kd nuclear protein binds to RNA segments that include the AAUAAA polyadenylation motif. Cell 52: 221-228.

Woltjen K, et al. 2009. piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. Nature 458: 766-770.

Xu C, Inokuma MS, Denham J, Golds K, Kundu P, Gold JD, Carpenter MK. 2001. Feeder-free growth of undifferentiated human embryonic stem cells. Nat Biotechnol 19: 971-974.

Yu J, Hu K, Smuga-Otto K, Tian S, Stewart R, Slukvin, II, Thomson JA. 2009. Human induced pluripotent stem cells free of vector and transgene sequences. Science 324: 797-801.

Yu J, et al. 2007. Induced pluripotent stem cell lines derived from human somatic cells. Science 318: 1917-1920.

Zhou H, et al. 2009. Generation of induced pluripotent stem cells using recombinant proteins. Cell Stem Cell 4: 381-384.

CLAIMS

1. A method for reducing the immunogenicity for mammalian cells of *in vitro*-synthesized RNA molecules comprising:

purifying *in vitro*-synthesized RNA molecules comprising a modified nucleoside selected from Ψ, $m^1Ψ$, m^5U, mo^5U, and s^2U in place of U using a process that removes RNA contaminant molecules that are immunogenic and toxic for mammalian cells by inducing an innate immune response, such that a purified RNA preparation comprising said *in vitro*-synthesized RNA molecules lacks immunogenicity,

wherein said *in vitro*-synthesized RNA molecules are obtained by *in vitro* transcription and encode a recombinant protein,

wherein said lack of immunogenicity is detectable using an *in vitro* MDDC immunogenicity assay by measuring no increase in the amount of IFN-α or TNF-α secreted by human or murine monocyte-derived dendritic cells (MDDCs) transfected with said purified RNA preparation that is greater than the amount of IFN-α or TNF-α secreted by MDDCs transfected with negative controls that do not contain said *in vitro*-synthesized RNA molecules; and

wherein said purification step enables repeated delivery of said purified RNA preparation into a mammalian cell without generation of inflammatory cytokines.

2. The method according to claim 1, wherein said *in vitro*-synthesized RNA molecules further comprise or consist of at least one of the following:

- they further comprise a poly-A tail;
- greater than 80% of said molecules further comprise a 5' cap;
- they are mRNA (messenger RNA) molecules; or
- they further comprise the modified nucleoside 5-methylcytidine (m^5C).

3. The method of any one of claims 1-2, wherein said reduced immunogenicity said purified RNA preparation is detectable using an *in vitro* assay comprising measuring decreased secretion of a cytokine selected from the group consisting of: IL-12, IFN-α, TNF-α, RANTES, MIP-1α, MIP-1β, IL-6, IFN-β and IL-8, wherein preferably said cytokine which is measured is selected from IL-12, IFN-α and TNF-α from mammalian cells contacted with said purified RNA

preparation than from said mammalian cells transfected with the same amount of the unpurified *in vitro*-synthesized molecules.

4. The method of any one of claims 1-3, wherein the reduced immunogenicity of said purified RNA preparation can be detected using an *in vitro* dendritic cell assay by measuring secretion of an amount of IFN-α or TNF-α cytokine from dendritic cells transfected with said purified RNA preparation that is less than the amount of secretion of said cytokine from dendritic cells transfected with the same amount of said *in vitro*-synthesized molecules not subjected to said purifying.

5. The method of any one of claims 1-4, wherein the recombinant protein is selected from the group consisting of: erythropoietin (EPO); a detectable enzyme selected from firefly luciferase, Renilla luciferase, bacterial beta-galactosidase (lacZ), and green fluorescent protein (GFP); a transcription factor selected from MYC and SRY or MCOP; a growth factor or cytokine selected from the group consisting of platelet-derived growth factor (PDGF), vascular endothelial growth factor (VEGF), transforming growth factor-beta1 (TGF-beta1), insulin-like growth factor (IGF), alpha-melanocyte-stimulating hormone (alpha-MSH); insulin-like growth factor-I (IGF-I); IL-4; IL-13; and IL-10; inducible nitric oxide synthase (iNOS); a heat shock protein; Cystic Fibrosis Transmembrane Conductance Regulator (CFTR); an enzyme with antioxidant activity selected from among catalase, phospholipid hydroperoxide glutathione peroxidase, superoxide dismutase-1, and superoxide dismutase-2; Bruton's tyrosine kinase; adenosine deaminase; ecto-nucleoside triphosphate diphosphydrolase; ABCA4; ABCD3; ACADM; AGL; AGT; ALDH4A1; ALPL; AMPD1; APOA2; AVSD1; BRCD2; C1QA; C1QB; C1QG; C8A; C8B; CACNA1S; CCV; CD3Z; CDC2L1; CHML; CHS1; CIAS1; CLCNKB; CMD1A; CMH2; CMM; COL11A1; COL8A2; COL9A2; CPT2; CRB1; CSE; CSF3R; CTPA; CTSK; DBT; DIO1; DISC1; DPYD; EKV; ENO1; ENO1P; EPB41; EPHX1; F13B; F5; FCGR2A; FCGR2B; FCGR3A; FCHL; FH; FMO3; FMO4; FUCA1; FY; GALE; GBA; GFND; GJA8; GJB3; GLC3B; HF1; HMGCL; HPC1; HRD; HRPT2; HSD3B2; HSPG2; KCNQ4; KCS; KIF1B; LAMB3; LAMC2; LGMD1B; LMNA; LOR; MCKD1; MCL1; MPZ; MTHFR; MTR; MUTYH; MYOC; NB; NCF2; NEM1; NPHS2; NPPA; NRAS; NTRK1; OPTA2; PBX1; PCHC; PGD; PHA2A; PHGDH; PKLR; PKP1; PLA2G2A; PLOD; PPOX; PPTO; PRCC; PRG4;

PSEN2; PTOS1; REN; RFX5; RHD; RMD1; RPE65; SCCD; SERPINC1; SJS1; SLC19A2; SLC2A1; SPG23; SPTA1; TAL1; TNFSF6; TNNT2; TPM3; TSHB; UMPK; UOX; UROD; USH2A; VMGLOM; VWS; WS2B; ABCB11; ABCG5; ABCG8; ACADL; ACP1; AGXT; AHHR; ALMS1; ALPP; ALS2; APOB; BDE; BDMR; BJS; BMPR2; CHRNA1; CMCWTD; CNGA3; COL3A1; COLAA3; COL4A4; COL6A3; CPS1; CRYGA; CRYGEP1; CYP1B1; CYP27A1; DBI; DES; DYSF; EDAR; EFEMP1; EIF2AK3; ERCC3; FSHR; GINGF; GLC1B; GPD2; GYPC; HADHA; HADHB; HOXD13; HPE2; IGKC; IHH; IRS1; ITGA6; KHK; KYNU; LCT; LHCGR; LSFC; MSH2; MSH6; NEB; NMTC; NPHP1; PAFAH1P1; PAX3; PAX8; PMS1; PNKD; PPH1; PROC; REG1A; SAG; SFTPB; SLC11A1; SLC3A1; SOS1; SPG4; SRD5A2; TCL4; TGFA; TMD; TPO; UGT1A; UV24; WSS; XDH; ZAP70; ZFHX1B; ACAA1; AGS1; AGTR1; AHSG; AMT; ARMET; BBS3; BCHE; BCPM; BTD; CASR; CCR2; CCR5; CDL1; CMT2B; COL7A1; CP; CPO; CRV; CTNNB1; DEM; ETM1; FANCD2; FIH; FOXL2; GBE1; GLB1; GLCLC; GNAI2; GNAT1; GP9; GPX1; HGD; HRG; ITIH1; KNG; LPP; LRS1; MCCC1; MDS1; MHS4; MITF; MLH1; MYL3; MYMY; OPA1; P2RY12; PBXP1; PCCB; POU1F1; PPARG; PROS1; PTHR1; RCA1; RHO; SCA7; SCLC1; SCN5A; SI; SLC25A20; SLC2A2; TF; TGFBR2; THPO; THRB; TKT; TM4SF1; TRH; UMPS; UQCRC1; USH3A; VHL; WS2A; XPC; ZNF35; ADH1B; ADH1C; AFP; AGA; AIH2; ALB; ASMD; BFHD; CNGA1; CRBM; DCK; DSPP; DTDP2; ELONG; ENAM; ETFDH; EVC; F11; FABP2; FGA; FGB; FGFR3; FGG; FSHMD1A; GC; GNPTA; GNRHR; GYPA; HCA; HCL2; HD; HTN3; HVBS6; IDUA; IF; JPD; KIT; KLKB1; LQT4; MANBA; MLLT2; MSX1; MTP; NR3C2; PBT; PDE6B; PEE1; PITX2; PKD2; QDPR; SGCB; SLC25A4; SNCA; SOD3; STATH; TAPVR1; TYS; WBS2; WFS1; WHCR; ADAMTS2; ADRB2; AMCN; AP3B1; APC; ARSB; B4GALT7; BHR1; C6; C7; CCAL2; CKN1; CMDJ; CRHBP; CSF1R; DHFR; DIAPH1; DTR; EOS; EPD; ERVR; F12; FBN2; GDNF; GHR; GLRA1; GM2A; HEXB; HSD17B4; ITGA2; KFS; LGMDLA; LOX; LTC4S; MAN2A1; MCC; MCCC2; MSH3; MSX2; NR3C1; PCSK1; PDE6A; PFBI; RASA1; SCZD1; SDHA; SGCD; SLC22A5; SLC26A2; SLC6A3; SM1; SMA; SMN1; SMN2; SPINK5; TCOF1; TELAB1; TGFBI; ALDH5A1; ARG1; AS; ASSP2; BCKDHB; BF; C2; C4A; CDKN1A; COL10A1; COL11A2; CYP21A2; DYX2; EJM1; ELOVL4; EPM2A; ESR1; EYA4; F13A1; FANCE; GCLC; GJA1; GLYS1; GMPR; GSE; HCR; HFE; HLA-A; HLA-DPB1; HLA-DRA; HPFH; ICS1; IDDM1; IFNGR1; IGAD1; IGF2R; ISCW; LAMA2; LAP; LCA5; LPA; MCDR1; MOCS1; MUT; MYB; NEU1; NKS1; NYS2;

OA3; ODDD; OFC0; PARK2; PBCA; PBCRA1; PDB1; PEX3; PEX6; PEX7; PKHD1; PLA2G7; PLG; POLH; PPAC; PSORS1; PUJO; RCD1; RDS; RHAG; RP14; RUNX2; RWS; SCA1; SCZD3; SIASD; SOD2; ST8; TAP1; TAP2; TFAP2B; TNDM; TNF; TPBG; TPMT; TULP1; WISP3; AASS; ABCB1; ABCB4; ACHE; AQP1; ASL; ASNS; AUTS1; BPGM; BRAF; C7orf2; CACNA2D1; CCM1; CD36; CFTR; CHORDOMA; CLCN1; CMH6; CMT2D; COL1A2; CRS; CYMD; DFNA5; DLD; DYT11; EEC1; ELN; ETV1; FKBP6; GCK; GHRHR; GHS; GLI3; GPDS1; GUSB; HLXB9; HOXA13; HPFH2; HRX; IAB; IMMP2L; KCNH2; LAMBI; LEP; MET; NCF1; NM; OGDH; OPN1SW; PEX1; PGAM2; PMS2; PON1; PPP1R3A; PRSS1; PTC; PTPN12; RP10; RP9; SERPINE1; SGCE; SHFM1; SHH; SLC26A3; SLC26A4; SLOS; SMAD1; TBXAS1; TWIST; ZWS1; ACHM3; ADRB3; ANK1; CA1; CA2; CCAL1; CLN8; CMT4A; CNGB3; COH1; CPP; CRH; CYP11B1; CYP11B2; DECR1; DPYS; DURS1; EBS1; ECA1; EGI; EXT1; EYA1; FGFR1; GNRH1; GSR; GULOP; HR; KCNQ3; KFM; KWE; LGCR; LPL; MCPH1; MOS; MYC; NAT1; NAT2; NBS1; PLAT; PLEC1; PRKDC; PXMP3; RP1; SCZD6; SFTPC; SGM1; SPG5A; STAR; TG; TRPS1; TTPA; VMD1; WRN; ABCA1; ABL1; ABO; ADAMTS13; AK1; ALAD; ALDH1A1; ALDOB; AMBP; AMCD1; ASS; BDMF; BSCL; C5; CDKN2A; CHAC; CLA1; CMD1B; COL5A1; CRAT; DBH; DNAI1; DYS; DYT1; ENG; FANCC; FBP1; FCMD; FRDA; GALT; GLDC; GNE; GSM1; GSN; HSD17B3; HSN1; IBM2; INVS; JBTS1; LALL; LCCS1; LCCS; LGMD2H; LMX1B; MLLT3; MROS; MSSE; NOTCH1; ORM1; PAPPA; PIP5K1B; PTCH; PTGS1; RLN1; RLN2; RMRP; ROR2; RPD1; SARDH; SPTLC1; STOM; TDFA; TEK; TMC1; TRIM32; TSC1; TYRP1; XPA; CACNB2; COL17A1; CUBN; CXCL12; CYP17; CYP2C19; CYP2C9; EGR2; EMX2; ERCC6; FGFR2; HK1; HPS1; IL2RA; LGI1; LIPA; MAT1A; MBL2; MKI67; MXI1; NODAL; OAT; OATL3; PAX2; PCBD; PEO1; PHYH; PNLIP; PSAP; PTEN; RBP4; RDPA; RET; SFTPA1; SFTPD; SHFM3; SIAL; THC2; TLX1; TNFRSF6; UFS; UROS; AA; ABCC8; ACAT1; ALX4; AMPD3; ANC; APOAL; APOA4; APOC3; ATM; BSCL2; BWS; CALCA; CAT; CCND1; CD3E; CD3G; CD59; CDKNLC; CLN2; CNTF; CPT1A; CTSC; DDB1; DDB2; DHCR7; DLAT; DRD4; ECB2; ED4; EVR1; EXT2; F2; FSHB; FTH1; G6PT1; G6PT2; GIF; HBB; HBBP1; HBD; HBE1; HBG1; HBG2; HMBS; HND; HOMG2; HRAS; HVBS1; IDDM2; IGER; INS; JBS; KCNJ11; KCNJ1; KCNQ1; LDHA; LRP5; MEN1; MLL; MYBPC3; MYO7A; NNO1; OPPG; OPTB1; PAX6; PC; PDX1; PGL2; PGR; PORC; PTH; PTS; PVRL1; PYGM; RAG1; RAG2; ROM1; RRAS2; SAA1; SCA5; SCZD2; SDHD; SERPING1; SMPD1; TCIRG1;

TCL2; TECTA; TH; TREH; TSG101; TYR; USH1C; VMD2; VRNI; WT1; WT2; ZNF145; A2M; AAAS; ACADS; ACLS; ACVRL1; ALDH2; AMHR2; AOM; AQP2; ATD; ATP2A2; BDC; CIR; CD4; CDK4; CNA1; COL2A1; CYP27B1; DRPLA; ENUR2; FEOM1; FGF23; FPF; GNB3; GNS; HAL; HBP1; HMGA2; HMN2; HPD; IGF1; KCNA1; KERA; KRAS2; KRT1; KRT2A; KRT3; KRT4; KRT5; KRT6A; KRT6B; KRTHB6; LDHB; LYZ; MGCT; MPE; MVK; MYL2; OAP; PAH; PPKB; PRB3; PTPN11; PXR1; RLS; RSN; SAS; SAX1; SCA2; SCNN1A; SMAL; SPPM; SPSMA; TBX3; TBX5; TCF1; TPI1; TSC3; ULR; VDR; VWF; ATP7B; BRCA2; BRCD1; CLN5; CPB2; ED2; EDNRB; ENUR1; ERCC5; F10; F7; GJB2; GJB6; IPF1; MBS1; MCOR; NYS4; PCCA; RB1; RHOK; SCZD7; SGCG; SLC10A2; SLC25A15; STARP1; ZNF198; ACHM1; ARVD1; BCH; CTAA1; DAD1; DFNB5; EML1; GALC; GCH1; IBGC1; IGH; IGHC group; IGHG1; IGHM; IGHR; IV; LTBP2; MJD; MNG1; MPD1; MPS3C; MYH6; MYH7; NP; NPC2; PABPN1; PSEN1; PYGL; RPGRIP1; SERPINA1; SERPINA3; SERPINA6; SLC7A7; SPG3A; SPTB; TCL1A; TGM1; TITF1; TMIP; TRA; TSHR; USHLA; VP; ACCPN; AHO2; ANCR; B2M; BBS4; BLM; CAPN3; CDAN1; CDAN3; CLN6; CMH3; CYP19; CYP1A1; CYP1A2; DYX1; EPB42; ETFA; EYCL3; FAH; FBN1; FES; HCVS; HEXA; IVD; LCS1; LIPC; MYO5A; OCA2; OTSC1; PWCR; RLBP1; SLC12A1; SPG6; TPM1; UBE3A; WMS; ABCC6; ALDOA; APRT; ATP2A1; BBS2; CARD15; CATM; CDH1; CETP; CHST6; CLN3; CREBBP; CTH; CTM; CYBA; CYLD; DHS; DNASE1; DPEP1; ERCC4; FANCA; GALNS; GAN; HAGH; HBA1; HBA2; HBHR; HBQ1; HBZ; HBZP; HP; HSD11B2; IL4R; LIPB; MC1R; MEFV; MHC2TA; MLYCD; MMVP1; PHKB; PHKG2; PKD1; PKDTS; PMM2; PXE; SALL1; SCA4; SCNN1B; SCNN1G; SLC12A3; TAT; TSC2; VDI; WT3; ABR; ACACA; ACADVL; ACE; ALDH3A2; APOH; ASPA; AXIN2; BCL5; BHD; BLMH; BRCA1; CACD; CCA1; CCZS; CHRNB1; CHRNE; CMT1A; COL1A1; CORD5; CTNS; EPX; ERBB2; G6PC; GAA; GALK1; GCGR; GFAP; GH1; GH2; GP1BA; GPSC; GUCY2D; ITGA2B; ITGB3; ITGB4; KRT10; KRT12; KRT13; KRT14; KRT14L1; KRT14L2; KRT14L3; KRT16; KRT16L1; KRT16L2; KRT17; KRT9; MAPT; MDB; MDCR; MGI; MHS2; MKS1; MPO; MYO15A; NAGLU; NAPB; NF1; NME1; P4HB; PAFAH1B1; PECAM1; PEX12; PHB; PMP22; PRKAR1A; PRKCA; PRKWNK4; PRP8; PRPF8; PTLAH; RARA; RCV1; RMSA1; RP17; RSS; SCN4A; SERPINF2; SGCA; SGSH; SHBG; SLC2A4; SLC4A1; SLC6A4; SMCR; SOST; SOX9; SSTR2; SYM1; SYNS1; TCF2; THRA; TIMP2; TOC; TOP2A; TP53; TRIM37; VBCH; ATP8B1; BCL2; CNSN; CORD1; CYB5; DCC;

F5F8D; FECH; FEO; LAMA3; LCFS2; MADH4; MAFD1; MC2R; MCL; MYP2; NPC1; SPPK; TGFBRE; TGIF; TTR; AD2; AMH; APOC2; APOE; ATHS; BAX; BCKDHA; BCL3; BFIC; C3; CACNA1A; CCO; CEACAM5; COMP; CRX; DBA; DDU; DFNA4; DLL3; DM1; DMWD; E11S; ELA2; EPOR; ERCC2; ETFB; EXT3; EYCL1; FTL; FUT1; FUT2; FUT6; GAMT; GCDH; GPI; GUSM; HB1; HCL1; HHC2; HHC3; ICAM3; INSR; JAK3; KLK3; LDLR; LHB; LIG1; LOH19CR1; LYL1; MAN2B1; MCOLN1; MDRV; MLLT1; NOTCH3; NPHS1; OFC3; OPA3; PEPD; PRPF31; PRTN3; PRX; PSG1; PVR; RYR1; SLC5A5; SLC7A9; STK11; TBXA2R; TGFB1; TNNI3; TYROBP; ADA; AHCY; AVP; CDAN2; CDPD1; CHED1; CHED2; CHRNA4; CST3; EDN3; EEGV1; FTLL1; GDF5; GNAS; GSS; HNF4A; JAG1; KCNQ2; MKKS; NBIA1; PCK1; PI3; PPCD; PPGB; PRNP; THBD; TOP1; AIRE; APP; CBS; COL6A1; COL6A2; CSTB; DCR; DSCR1; FPDMM; HLCS; HPE1; ITGB2; KCNE1; KNO; PRSS7; RUNX1; SOD1; TAM; ADSL; ARSA; BCR; CECR; CHEK2; COMT; CRYBB2; CSF2RB; CTHM; CYP2D6; CYP2D7P1; DGCR; DIA1; EWSR1; GGT1; MGCR; MN1; NAGA; NE2; OGS2; PDGFB; PPARA; PRODH; SCO2; SCZD4; SERPIND1; SLC5A1; SOX10; TCN2; TIMP3; TST; VCF; ABCD1; ACTL1; ADFN; AGMX2; AHDS; AIC; AIED; AIH3; ALAS2; AMCD; AMELX; ANOP1; AR; ARAF1; ARSC2; ARSE; ARTS; ARX; ASAT; ASSP5; ATP7A; ATRX; AVPR2; BFLS; BGN; BTK; BZX; C1HR; CACNA1F; CALB3; CBBM; CCT; CDR1; CFNS; CGF1; CHM; CHR39c; CIDX; CLA2; CLCN5; CLS; CMTX2; CMTX3; CND; COD1; COD2; COL4A5; COL4A6; CPX; CVD1; CYBB; DCX; DFN2; DFN4; DFN6; DHOF; DIAPH2; DKC1; DMD; DSS; DYT3; EBM; EBP; ED1; ELK1; EMD; EVR2; F8; F9; FCP1; FDPSL5; FGD1; FGS1; FMR1; FMR2; G6PD; GABRA3; GATA1; GDI1; GDXY; GJB1; GK; GLA; GPC3; GRPR; GTD; GUST; HMS1; HPRT1; HPT; HTC2; HTR2c; HYR; IDS; IHG1; IL2RG; INDX; IP1; IP2; JMS; KAL1; KFSD; L1CAM; LAMP2; MAA; MAFD2; MAOA; MAOB; MCF2; MCS; MEAX; MECP2; MF4; MGC1; MIC5; MID1; MLLT7; MLS; MRSD; MRX14; MRX1; MRX20; MRX2; MRX3; MRX40; MRXA; MSD; MTM1; MYCL2; MYP1; NDP; NHS; NPHL1; NROB1; NSX; NYS1; NYX; OA1; OASD; OCRL; ODT1; OFD1; OPA2; OPD1; OPEM; OPN1LW; OPN1MW; OTC; P3; PDHA1; PDR; PFC; PFKFB1; PGK1; PGK1P1; PGS; PHEX; PHKA1; PHKA2; PHP; PIGA; PLP1; POF1; POLA; POU3F4; PPMX; PRD; PRPS1; PRPS2; PRS; RCCP2; RENBP; RENS1; RP2; RP6; RPGR; RPS4X; RPS6KA3; RS1; S11; SDYS; SEDL; SERPINA7; SH2D1A; SHFM2; SLC25A5; SMAX2; SRPX; SRS; STS; SYN1; SYP; TAF1; TAZ; TBX22; TDD; TFE3; THAS;

THC; TIMM8A; TIM1; TKCR; TNFSF5; UBE1; UBE2A; WAS; WSN; WTS; WWS; XIC; XIST; XK; XM; XS; ZFX; ZIC3; ZNF261; ZNF41; ZNF6; AMELY; ASSP6; AZF1; AZF2; DAZ; GCY; RPS4Y; SMCY; ZFY; ABAT; AEZ; AFA; AFD1; ASAH1; ASD1; ASMT; CCAT; CECR9; CEPA; CLA3; CLN4; CSF2RA; CTS1; DF; DIH1; DWS; DYT2; DYT4; EBR3; ECT; EEF1A1L14; EYCL2; FANCB; GCSH; GCSL; GIP; GTS; HHG; HMI; HOAC; HOKPP2; HRPT1; HSD3B3; HTC1; HV1S; ICHQ; ICR1; ICR5; IL3RA; KAL2; KMS; KRT18; KSS; LCAT; LHON; LIMM; MANBB; MCPH2; MEB; MELAS; MIC2; MPFD; MS; MSS; MTATP6; MTCO1; MTC03; MTCYB; MTND1; MTND2; MTND4; MTND5; MTND6; MTRNR1; MTRNR2; MTTE; MTTG; MTTI; MTTK; MTTL1; MTTL2; MTTN; MTTP; MTTS1; NAMSD; OCD1; OPD2; PCK2; PCLD; PCOS1; PFKM; PKD3; PRCA1; PRO1; PROP1; RBS; RFXAP; RP; SHOX; SLC25A6; SPG5B; STO; SUOX; THM; and TTD.

6. The method of any one of claims 1-5, wherein said purified RNA preparation, comprises at least one of the following features:

- it can be repeatedly administered to a mammalian cell without eliciting an immune response sufficient to detectably reduce expression of said recombinant protein; or

- it can be repeatedly introduced into mammalian cells without eliciting an immune response sufficient to eliminate detectable expression of said recombinant protein.

7. Use of said purified RNA preparation made using the method of any one of claims 1-6 in the preparation of a pharmaceutical composition for treatment of a monogenic disorder due to an inborn error of metabolism,

wherein said pharmaceutical composition is to be administered to a subject with said disorder,

wherein said treatment is a therapeutic treatment or a prophylactic treatment; and

wherein said recombinant protein encoded by said *in vitro*-synthesized RNA molecules is: EPO; iNOS; a heat shock protein; CFTR; Bruton's tyrosine kinase; ADA; ecto-nucleoside triphosphate diphosphohydrolase; Sphingomyelinase; alpha-L-iduronidase, iduronate-2--sulfatase; acid beta-glucosidase; alpha-galactosidase A; ABCA4; ABCD3; ACADM; AGL; AGT; ALDH4Al; ALPL; AMPD1; APOA2; AVSD1; BRCD2; C1QA; C1QB; C1QG; C8A;

C8B; CACNA1S; CCV; CD3Z; CDC2L1; CHML; CHS1; CIAS1; CLCNKB; CMD1A; CMH2; CMM; COL11AI; COL8A2; COL9A2; CPT2; CRB1; CSE; CSF3R; CTPA; CTSK; DBT; DIO1; DISC1; DPYD; EKV; ENO1; ENO1P; EPB41; EPHX1; F13B; F5; FCGR2A; FCGR2B; FCGR3A; FCHL; FH; FMO3; FMO4; FUCA1; FY; GALE; GBA; GFND; GJA8; GJB3; GLC3B; HF1; HMGCL; HPC1; HRD; HRPT2; HSD3B2; HSPG2; KCNQ4; KCS; KIF1B; LAMB3; LAMC2; LGMD1B; LMNA; LOR; MCKD1; MCL1; MPZ; MTHFR; MTR; MUTYH; MYOC; NB; NCF2; NEM1; NPHS2; NPPA; NRAS; NTRK1; OPTA2; PBX1; PCHC; PGD; PHA2A; PHGDH; PKLR; PKP1; PLA2G2A; PLOD; PPOX; PPT1; PRCC; PRG4; PSEN2; PTOS1; REN; RFX5; RHD; RMD1; RPE65; SCCD; SERPINC1; SJS1; SLC19A2; SLC2A1; SPG23; SPTA1; TAL1; TNFSF6; TNNT2; TPM3; TSHB; UMPK; UOX; UROD; USH2A; VMGLOM; VWS; WS2B; ABCB11; ABCG5; ABCG8; ACADL; ACP1; AGXT; AHHR; ALMS1; ALPP; ALS2; APOB; BDE; BDMR; BJS; BMPR2; CHRNA1; CMCWTD; CNGA3; COL3A1; COL4A3; COL4A4; COL6A3; CPS1; CRYGA; CRYGEP1; CYP1B1; CYP27A1; DBI; DES; DYSF; EDAR; EFEMP1; EIF2AK3; ERCC3; FSHR; GINGF; GLC1B; GPD2; GYPC; HADHA; HADHB; HOXD13; HPE2; IGKC; IHH; IRS1; ITGA6; KHK; KYNU; LCT; LHCGR; LSFC; MSH2; MSH6; NEB; NMTC; NPHP1; PAFAH1P1; PAX3; PAX8; PMS1; PNKD; PPH1; PROC; REGIA; SAG; SFTPB; SLC11A1; SLC3Al; SOS1; SPG4; SRD5A2; TCL4; TGFA; TMD; TPO; UGT1A; UV24; WSS; XDH; ZAP70; ZFHX1B; ACAA1; AGS1; AGTR1; AHSG; AMT; ARMET; BBS3; BCHE; BCPM; BTD; CASR; CCR2; CCR5; CDL1; CMT2B; COL7A1; CP; CPO; CRY; CTNNB1; DEM; ETM1; FANCD2; F1H; FOXL2; GBE1; GLB1; GLC1C; GNAI2; GNAT1; GP9; GPX1; HGD; HRG; ITIH1; KNG; LPP; LRS1; MCCC1; MDS1; MHS4; MITF; MLH1; MYL3; MYMY; OPA1; P2RY12; PBXPI; PCCB; POU1FI; PPARG; PROS1; PTHR1; RCA1; RHO; SCA7; SCLC1; SCN5A; SI; SLC25A20; SLC2A2; TF; TGFBR2; THPO; THRB; TKT; TM4SF1; TRH; UMPS; UQCRC1; USH3A; VHL; WS2A; XPC; ZNF35; ADH1B; ADH1C; AFP; AGA; AIH2; ALB; ASMD; BFHD; CNGA1; CRBM; DCK; DSPP; DTDP2; ELONG; ENAM; ETFDH; EVC; F11; FABP2; FGA;FGB; FGFR3; FGG; FSHMD1A; GC; GNPTA; GNRHR; GYPA; HCA; HCL2; HD; HTN3; HVBS6; IDUA; IF; JPD; KIT; KLKB1; LQT4; MANBA; MLLT2; MSX1; MTP; NR3C2; PBT; PDE6B; PEE1; PITX2; PKD2; QDPR; SGCB; SLC25A4; SNCA; SOD3; STATH; TAPVR1; TYS; WBS2; WFS1; WHCR; ADAMTS2; ADRB2; AMCN; AP3BI; APC; ARSB; B4GALT7; BHR1; C6; C7; CCAL2; CKN1; CMDJ; CRHBP; CSF1R; DHFR; DIAPH1;

DTR; EOS; EPD; ERVR; F12; FBN2; GDNF; GHR; GLRA1; GM2A; HEXB; HSD17B4; ITGA2; KFS; LGMD1A; LOX; LTC4S; MAN2A1; MCC; MCCC2; MSH3; MSX2; NR3C1; PCSK1; PDE6A; PFBI; RASA1; SCZD1; SDHA; SGCD; SLC22A5; SLC26A2; SLC6A3; SM1; SMA; SMN1; SMN2; SPINK5; TCOF1; TELAB1; TGFBI; ALDH5Al; ARG1; AS; ASSP2; BCKDHB; BF; C2; C4A; CDKN1A; COL10A1; COL11A2; CYP21A2; DYX2; EJM1; ELOVL4; EPM2A; ESR1; EYA4; F13A1; FANCE; GCLC; GJA1; GLYS1; GMPR; GSE; HCR; HFE; HLA-A; HLA-DPB1; HLA-DRA; HPFH; ICS1; IDDM1; IFNGR1; IGAD1; IGF2R; ISCW; LAMA2; LAP; LCA5; LPA; MCDR1; MOCS1; MUT; MYB; NEU1; NKS1; NYS2; OA3; OODD; OFC1; PARK2; PBCA; PBCRA1; PDB1; PEX3; PEX6; PEX7; PKHD1; PLA2G7; PLG; POLH; PPAC; PSORS1; PUJO; RCD1; RDS; RHAG; RP14; RUNX2; RWS; SCA1; SCZD3; SIASD; SOD2; ST8; TAP1; TAP2; TFAP2B; TNDM; TNF; TPBG; TPMT; TULP1; WISP3; AASS; ABCB1; ABCB4; ACHE; AQP1; ASL; ASNS; AUTS1; BPGM; BRAF; C7orf2; CACNA2D1; CCM1; CD36; CFTR; CHORDOMA; CLCN1; CMH6; CMT2D; COL1A2; CRS; CYMD; DFNA5; DLD; DYT11; EEC1; ELN; ETV1; FKBP6; GCK; GHRHR; GHS; GLI3; GPDS1; GUSB; HLXB9; HOXA13; HPFH2; HRX; IAB; IMMP2L; KCNH2; LAMB1; LEP; MET; NCF1;NM; OGDH; OPN1SW; PEX1; PGAM2; PMS2; PON1; PPP1R3A; PRSS1; PTC; PTPN12; RP10; RP9; SERPINE1; SGCE; SHFM1; SHH; SLC26A3; SLC26A4; SLOS; SMAD1; TBXAS1; TWIST; ZWS1; ACHM3; ADRB3; ANKI; CA1; CA2; CCAL1; CLN8; CMT4A; CNGB3; COH1; CPP; CRH; CYP11B1; CYP11B2; DECR1; DPYS; DURS1; EBS1; ECA1; EGI; EXT1; EYA1; FGFR1; GNRH1; GSR; GULOP; HR; KCNQ3; KFM; KWE; LGCR; LPL; MCPH1; MOS; MYC; NAT1; NAT2; NBS1; PLAT; PLEC1; PRKDC; PXMP3; RP1; SCZD6; SFTPC; SGM1; SPG5A; STAR; TG; TRPS1; TTPA; VMD1; WRN; ABCA1; ABL1; ABO; ADAMTS13; AK1; ALAD; ALDH1A1; ALDOB; AMBP; AMCD1; ASS; BDMF; BSCL; C5; CDKN2A; CHAC; CLA1; CMD1B; COL5A1; CRAT; DBH; DNAI1; DYS; DYT1; ENG; FANCC; FBP1; FCMD; FRDA; GALT; GLDC; GNE; GSM1; GSN; HSD17B3; HSN1; IBM2; INVS; JBTS1; LALL; LCCS1; LCCS; LGMD2H; LMX1B; MLLT3; MROS; MSSE; NOTCH1; ORM1; PAPPA; PIP5K1B; PTCH; PTGS1; RLN1; RLN2; RMRP; ROR2; RPD1; SARDH; SPTLC1; STOM; TDFA; TEK; TMC1; TRIM32; TSC1; TYRP1; XPA; CACNB2; COLl7A1; CUBN; CXCL12; CYP17; CYP2C19; CYP2C9; EGR2; EMX2; ERCC6; FGFR2; HK1; HPSI; IL2RA; LGI1; LIPA; MAT1A; MBL2; MKI67; MXI1; NODAL; OAT; OATL3; PAX2; PCBD; PEO1; PHYH; PNL1P; PSAP;PTEN; RBP4; RDPA; RET; SFTPA1;

SFTPD; SHFM3; SIAL; THC2; TLX1; TNFRSF6; UFS; UROS; AA; ABCC8; ACAT1; ALX4; AMPD3; ANC; APOA1; APOA4; APOC3; ATM; BSCL2; BWS; CALCA; CAT; CCND1; CD3E; CD3G; CD59; CDKN1C; CLN2; CNTF; CPT1A; CTSC; DDB1; DDB2; DHCR7; DLAT; DRD4; ECB2; ED4; EVR1; EXT2; F2; FSHB; FTH1; G6PT1; G6PT2; GIF; HBB; HBBP1; HBD; HBE1; HBG1; HBG2; HMBS; HND; HOMG2; HRAS; HVBS1; IDDM2; IGER; INS; JBS; KCNJ11; KCNJ1; KCNQ1; LDHA; LRP5; MEN1; MLL; MYBPC3; MYO7A; NNO1; OPPG; OPTB1; PAX6; PC; PDX1; PGL2; PGR; PORC; PTH; PTS; PVRL1; PYGM; RAG1; RAG2; ROM1; RRAS2; SAA1; SCA5; SCZD2; SDHD; SERPING1; SMPD1; TCIRG1; TCL2; TECTA; TH; TREH; TSG101; TYR; USH1C; VMD2; VRN1; WT1; WT2; ZNF145; A2M; AAAS; ACADS; ACLS; ACVRL1; ALDH2; AMHR2; AOM; AQP2; ATD; ATP2A2; BDC; C1R; CD4; CDK4; CNA1; COL2A1; CYP27B1; DRPLA; ENUR2; FEOM1; FGF23; FPF; GNB3; GNS; HAL; HBP1; HMGA2; HMN2; HPD; IGF1; KCNA1; KERA; KRAS2; KRT1; KRT2A; KRT3; KRT4; KRT5; KRT6A; KRT6B; KRTHB6; LDHB; LYZ; MGCT; MPE; MVK; MYL2; OAP; PAH; PPKB; PRB3; PTPN11; PXR1; RLS; RSN; SAS; SAX1; SCA2; SCNN1A; SMAL; SPPM; SPSMA; TBX3; TBX5; TCF1; TPI1; TSC3; ULR; VDR; VWF; ATP7B; BRCA2; BRCD1; CLN5; CPB2; ED2; EDNRB; ENUR1; ERCC5; F10; F7; GJB2; GJB6; IPF1; MBS1; MCOR; NYS4; PCCA; RB1; RHOK; SCZD7; SGCG; SLC10A2; SLC25A15; STARP1; ZNFl98; ACHM1; ARVDI; BCH; CTAA1; DAD1; DFNB5; EML1; GALC; GCH1; IBGC1; IGH; IGHC group; IGHG1; IGHM; IGHR; IV; LTBP2; MCOP; MJD; MNG1 MPD1; MPS3C; MYH6; MYH7; NP; NPC2; PABN1; PSEN1 PYGL; RPGRIP1; SERPINA1; SERPINA3; SERPINA6; SLC7A7; SPG3A; SPTB; TCL1A; TGMI; TITF1; TMIP; TRA; TSHR; USH1A; VP; ACCPN; AHO2; ANCR; B2M; BBS4; BLM; CAPN3; CDAN1; CDAN3; CLN6; CMH3; CYP19; CYP1A1; CYP1A2; DYX1; EPB42; ETFA; EYCL3; FAH; FBN1; FES; HCVS; HEXA; IVD; LCS1; LIPC; MYO5A; OCA2; OTSC1; PWCR; RLBP1; SLC12A1; SPG6; TPM1; UBE3A; WMS; ABCC6; ALDOA; APRT; ATP2A1; BBS2; CARD15; CATM; CDH1; CETP; CHST6; CLN3; CREBBP; CTH; CTM; CYBA; CYLD; DHS; DNASE1; DPEP1; ERCC4; FANCA; GALNS; GAN; HAGH; HBA1; HBA2; HBHR; HBQ1; HBZ; HBZP; HP; HSD11B2; IL4R; LIPB; MC2R; MEFV; MHC2TA; MLYCD; MMVP1; PHKB; PHKG2; PKD1; PKDTS; PMM2; PXE; SALL1; SCA4; SCNN1B; SCNN1G; SLC12A3; TAT; TSC2; VDI; WT3; ABR; ACACA; ACADVL; ACE; ALDH3A2; APOH; ASPA; AXIN2; BCL5; BHD; BLMH; BRCA1; CACD; CCA1; CCZS; CHRNB1; CHRNE;

CMT1A; COL1A1; CORD5; CTNS; EPX; ERBB2; G6PC; GAA; GALK1; GCGR; GFAP; GH1; GH2; GP1BA; GPSC; GUCY2D; ITGA2B; ITGB3; ITGB4; KRT10; KRT12; KRT13; KRT14; KRT14L1; KRT14L2; KRT14L3; KRT16; KRT16L1; KRT16L2; KRT17; KRT9; MAPT; MDB; MDCR; MGI; MHS2; MKS1; MPO; MYO15A; NAGLU; NAPB; NF1; NME1; P4HB; PAFAH1B1; PECAM1; PEX12; PHB; PMP22; PRKAR1A; PRKCA; PRKWNK4; PRP8; PRPF8; PTLAH; RARA; RCV1; RMSA1; RP17; RSS; SCN4A; SERPINF2; SGCA; SGSH; SHBG; SLC2A4; SLC4A1; SLC6A4; SMCR; SOST; SOX9; SSTR2; SYM1; SYNS1; TCF2; THRA; TIMP2; TOC; TOP2A; TP53; TRIM37; VBCH; ATP8B1; BCL2; CNSN; CORD1; CYB5; DCC; F5F8D; FECH; PEO; LAMA3; LCFS2; MADH4; MAFD1; MC2R; MCL; MYP2; NPC1; SPPK; TGFBRE; TGIF; TTR; AD2; AMH; APOC2; APOE; ATHS; BAX; BCKDHA; BCL3; BFIC; C3; CACNA1A; CCO; CEACAM5; COMP; CRX; DBA; DDU; DFNA4; DLL3; DM1; DMWD; E11S; ELA2; EPOR; ERCC2; ETFB; EXT3; EYCL1; FTL; FUT1; FUT2; FUT6; GAMT; GCDH; GPI; GUSM; HB1; HCL1; HHC2; HHC3; ICAM3; INSR; JAK3; KLK3; LDLR; LHB; LIG1; LOH19CR1; LYL1; MAN2B1; MCOLN1; MDRV; MLLT1; NOTCH3; NPHS1; OFC3; OPA3; PEPD; PRPF31; PRTN3; PRX; PSG1; PVR; RYR1; SLC5A5; SLC7A9; STK11; TBXA2R; TGFB1; TNNI3; TYROBP; ADA; AHCY; AVP; CDAN2; CDPD1; CHED1; CHED2; CHRNA4; CST3; EDN3; EEGV1; FTLL1; GDF5; GNAS; GSS; HNF4A; JAG1; KCNQ2; MKKS; NBIA1; PCK1; PI3; PPCD; PPGB; PRNP; THBD; TOP1; AIRE; APP; CBS; COL6A1; COL6A2; CSTB; DCR; DSCR1; FPDMM; HLCS; HPE1; ITGB2; KCNE1; KNO; PRSS7; RUNX1; SOD1; TAM; ADSL; ARSA; BCR; CECR; CHEK2; COMT; CRYBB2; CSF2RB; CTHM; CYP2D6; CYP2D7P1; DGCR; DIA1; EWSR1; GGT1; MGCR; MN1; NAGA; NF2; OGS2; PDGFB; PPARA; PRODH; SCO2; SCZD4; SERPIND1; SLC5A1; SOX10; TCN2; TIMP3; TST; VCF; ABCD1; ACTL1; ADFN; AGMX2; AHDS; AIC; AIED; AIH3; ALAS2; AMCD; AMELX; ANOP1; AR; ARAF1; ARSC2; ARSE; ARTS; ARX; ASAT; ASSP5; ATP7A; ATRX; AVPR2; BFLS; BGN; BTK; BZX; C1HR; CACNA1F; CALB3; CBBM; CCT; CDR1; CFNS; CGF1; CHM; CHR39C; CIDX; CLA2; CLCN5; CLS; CMTX2; CMTX3; CND; COD1; COD2; COL4A5; COL4A6; CPX; CVD1; CYBB; DCX; DFN2; DFN4; DFN6; DHOF; DIAPH2; DKC1; DMD; DSS; DYT3; EBM; EBP; ED1; ELK1; EMD; EVR2; F8; F9; FCP1; FDPSL5; FGD1; FGS1; FMR1; FMR2; G6PD; GABRA3; GATA1; GDI1; GDXY; GJB1; GK; GLA; GPC3; GRPR; GTD; GUST; HMS1; HPRT1; HPT; HTC2; HTR2C; HYR; IDS; IHG1; IL2RG; INDX; IP1; IP2; JMS; KAL1; KFSD; L1CAM;

LAMP2; MAA; MAFD2; MAOA; MAOB; MCF2; MCS; MEAX; MECP2; MF4; MGC1; MIC5; MID1; MLLT7; MLS; MRSD; MRX14; MRX1; MRX20; MRX2; MRX3; MRX40; MRXA; MSD; MTM1; MYCL2; MYP1; NDP; NHS; NPHL1; NR0B1; NSX; NYS1; NYX; OA1; OASD; OCRL; ODT1; OFD1; OPA2; OPD1; OPEM; OPN1LW; OPN1MW; OTC; P3; PDHA1; PDR; PFC; PFKFB1; PGK1; PGK1P1; PGS; PHEX; PHKA1; PHKA2; PHP; PIGA; PLP1; POF1; POLA; POU3F4; PPMX; PRD; PRPS1; PRPS2; PRS; RCCP2; RENBP; RENS1; RP2; RP6; RPGR; RPS4X; RPS6KA3; RS1; S11; SDYS; SEDL; SERPINA7; SH2D1A; SHFM2; SLC25A5; SMAX2; SRPX; SRS; STS; SYN1; SYP; TAF1; TAZ; TBX22; TDD; TFE3; THAS; THC; TIMM8A; TIMP1; TKCR; TNFSF5; UBE1; UBE2A; WAS; WSN; WTS; WWS; XIC; XIST; XK; XM; XS; ZFX; ZIC3; ZNF261; ZNF41; ZNF6; AMELY; ASSP6; AZF1; AZF2; DAZ; GCY; RPS4Y; SMCY; SRY; ZFY; ABAT; AEZ; AFA; AFD1; ASAH1; ASD1; ASMT; CCAT; CECR9; CEPA; CLA3; CLN4; CSF2RA; CTS1; DF; DIH1; DWS; DYT2; DYT4; EBR3; ECT; EEF1A1L14; EYCL2; FANCB; GCSH; GCSL; GIP; GTS; HHG; HMI; HOAC; HOKPP2; HRPT1; HSD3B3; HTC1; HV1S; ICHQ; ICR1; ICR5; IL3RA; KAL2; KMS; KRT18; KSS; LCAT; LHON; LIMM; MANBB; MCPH2; MEB; MELAS; MIC2; MPFD; MS; MSS; MTATP6; MTCO1; MTCO3; MTCYB; MTND1; MTND2; MTND4; MTND5; MTND6; MTRNR1; MTRNR2; MTTE; MTTG; MTTI; MTTK; MTTL1; MTTL2; MTTN; MTTP; MTTS1; NAMSD; OCD1; OPD2; PCK2; PCLD; PCOS1; PFKM; PKD3; PRCA1; PRO1; PROP1; RBS; RFXAP; RP; SHOX; SLC25A6; SPG5B; STO; SUOX; THM; or TTD.

8. The use of claim 7, wherein the pharmaceutical composition further comprises at least one of the following:

- a pharmaceutically acceptable carrier or diluent; and/or

- a transfection reagent, wherein said transfection reagent is preferably selected from the group consisting of a lipid-based transfection reagent such as a cationic lipid reagent, a protein-based transfection reagent, a polyethyleneimine-based transfection reagent and calcium phosphate.

9. The use of any one of claims 7-8, wherein said composition is to be administered to said subject parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously, intra-dermally, subcutaneously, intra-peritonealy, intra-ventricularly, intra-

cranially, intra-vaginally or intra-tumorally, and/or, wherein said composition is to be administered at a dose selected from among a one-time dose, a daily dose, a weekly dose, a monthly dose and a series of a defined number of doses.

10. Use of the purified RNA preparation encoding a recombinant protein of any one of claims 1-6 in the preparation of a medicament for the treatment of a disorder due to an inborn error of metabolism,

wherein said treatment is a therapeutic treatment or a prophylactic treatment;

wherein said medicament is to be administered as per a single dose to a subject with said disorder, wherein said recombinant protein is produced in cells of said subject after said administration,

wherein said recombinant protein is:

EPO; iNOS; a heat shock protein; CFTR; Bruton's tyrosine kinase; ADA; ecto-nucleoside triphosphate diphosphohydrolase; Sphingomyelinase; alpha-L-iduronidase, iduronate-2--sulfatase; acid beta-glucosidase; alpha-galactosidase A; ABCA4; ABCD3; ACADM; AGL; AGT; ALDH4Al; ALPL; AMPD1; APOA2; AVSD1; BRCD2; C1QA; C1QB; C1QG; C8A; C8B; CACNA1S; CCV; CD3Z; CDC2L1; CHML; CHS1; CIAS1; CLCNKB; CMD1A; CMH2; CMM; COL11AI; COL8A2; COL9A2; CPT2; CRB1; CSE; CSF3R; CTPA; CTSK; DBT; DIO1; DISC1; DPYD; EKV; ENO1; ENO1P; EPB41; EPHX1; F13B; F5; FCGR2A; FCGR2B; FCGR3A; FCHL; FH; FMO3; FMO4; FUCA1; FY; GALE; GBA; GFND; GJA8; GJB3; GLC3B; HF1; HMGCL; HPC1; HRD; HRPT2; HSD3B2; HSPG2; KCNQ4; KCS; KIF1B; LAMB3; LAMC2; LGMD1B; LMNA; LOR; MCKD1; MCL1; MPZ; MTHFR; MTR; MUTYH; MYOC; NB; NCF2; NEM1; NPHS2; NPPA; NRAS; NTRK1; OPTA2; PBX1; PCHC; PGD; PHA2A; PHGDH; PKLR; PKP1; PLA2G2A; PLOD; PPOX; PPT1; PRCC; PRG4; PSEN2; PTOS1; REN; RFX5; RHD; RMD1; RPE65; SCCD; SERPINC1; SJS1; SLC19A2; SLC2A1; SPG23; SPTA1; TAL1; TNFSF6; TNNT2; TPM3; TSHB; UMPK; UOX; UROD; USH2A; VMGLOM; VWS; WS2B; ABCB11; ABCG5; ABCG8; ACADL; ACP1; AGXT; AHHR; ALMS1; ALPP; ALS2; APOB; BDE; BDMR; BJS; BMPR2; CHRNA1; CMCWTD; CNGA3; COL3A1; COL4A3; COL4A4; COL6A3; CPS1; CRYGA; CRYGEP1; CYP1B1; CYP27A1; DBI; DES; DYSF; EDAR; EFEMP1; EIF2AK3; ERCC3; FSHR; GINGF; GLC1B; GPD2; GYPC; HADHA; HADHB; HOXD13; HPE2; IGKC; IHH; IRS1; ITGA6; KHK; KYNU; LCT;

LHCGR; LSFC; MSH2; MSH6; NEB; NMTC; NPHP1; PAFAH1P1; PAX3; PAX8; PMS1; PNKD; PPH1; PROC; REGIA; SAG; SFTPB; SLC11A1; SLC3Al; SOS1; SPG4; SRD5A2; TCL4; TGFA; TMD; TPO; UGT1A; UV24; WSS; XDH; ZAP70; ZFHX1B; ACAA1; AGS1; AGTR1; AHSG; AMT; ARMET; BBS3; BCHE; BCPM; BTD; CASR; CCR2; CCR5; CDL1; CMT2B; COL7A1; CP; CPO; CRY; CTNNB1; DEM; ETM1; FANCD2; F1H; FOXL2; GBE1; GLB1; GLC1C; GNAI2; GNAT1; GP9; GPX1; HGD; HRG; ITIH1; KNG; LPP; LRS1; MCCC1; MDS1; MHS4; MITF; MLH1; MYL3; MYMY; OPA1; P2RY12; PBXPI; PCCB; POU1FI; PPARG; PROS1; PTHR1; RCA1; RHO; SCA7; SCLC1; SCN5A; SI; SLC25A20; SLC2A2; TF; TGFBR2; THPO; THRB; TKT; TM4SF1; TRH; UMPS; UQCRC1; USH3A; VHL; WS2A; XPC; ZNF35; ADH1B; ADH1C; AFP; AGA; AIH2; ALB; ASMD; BFHD; CNGA1; CRBM; DCK; DSPP; DTDP2; ELONG; ENAM; ETFDH; EVC; F11; FABP2; FGA;FGB; FGFR3; FGG; FSHMD1A; GC; GNPTA; GNRHR; GYPA; HCA; HCL2; HD; HTN3; HVBS6; IDUA; IF; JPD; KIT; KLKB1; LQT4; MANBA; MLLT2; MSX1; MTP; NR3C2; PBT; PDE6B; PEE1; PITX2; PKD2; QDPR; SGCB; SLC25A4; SNCA; SOD3; STATH; TAPVR1; TYS; WBS2; WFS1; WHCR; ADAMTS2; ADRB2; AMCN; AP3BI; APC; ARSB; B4GALT7; BHR1; C6; C7; CCAL2; CKN1; CMDJ; CRHBP; CSF1R; DHFR; DIAPH1; DTR; EOS; EPD; ERVR; F12; FBN2; GDNF; GHR; GLRA1; GM2A; HEXB; HSD17B4; ITGA2; KFS; LGMD1A; LOX; LTC4S; MAN2A1; MCC; MCCC2; MSH3; MSX2; NR3C1; PCSK1; PDE6A; PFBI; RASA1; SCZD1; SDHA; SGCD; SLC22A5; SLC26A2; SLC6A3; SM1; SMA; SMN1; SMN2; SPINK5; TCOF1; TELAB1; TGFBI; ALDH5Al; ARG1; AS; ASSP2; BCKDHB; BF; C2; C4A; CDKN1A; COL10A1; COL11A2; CYP21A2; DYX2; EJM1; ELOVL4; EPM2A; ESR1; EYA4; F13A1; FANCE; GCLC; GJA1; GLYS1; GMPR; GSE; HCR; HFE; HLA-A; HLA-DPB1; HLA-DRA; HPFH; ICS1; IDDM1; IFNGR1; IGAD1; IGF2R; ISCW; LAMA2; LAP; LCA5; LPA; MCDR1; MOCS1; MUT; MYB; NEU1; NKS1; NYS2; OA3; OODD; OFC1; PARK2; PBCA; PBCRA1; PDB1; PEX3; PEX6; PEX7; PKHD1; PLA2G7; PLG; POLH; PPAC; PSORS1; PUJO; RCD1; RDS; RHAG; RP14; RUNX2; RWS; SCA1; SCZD3; SIASD; SOD2; ST8; TAP1; TAP2; TFAP2B; TNDM; TNF; TPBG; TPMT; TULP1; WISP3; AASS; ABCB1; ABCB4; ACHE; AQP1; ASL; ASNS; AUTS1; BPGM; BRAF; C7orf2; CACNA2D1; CCM1; CD36; CFTR; CHORDOMA; CLCN1; CMH6; CMT2D; COL1A2; CRS; CYMD; DFNA5; DLD; DYT11; EEC1; ELN; ETV1; FKBP6; GCK; GHRHR; GHS; GLI3; GPDS1; GUSB; HLXB9; HOXA13; HPFH2; HRX; IAB; IMMP2L; KCNH2;

LAMB1; LEP; MET; NCF1;NM; OGDH; OPN1SW; PEX1; PGAM2; PMS2; PON1; PPP1R3A; PRSS1; PTC; PTPN12; RP10; RP9; SERPINE1; SGCE; SHFM1; SHH; SLC26A3; SLC26A4; SLOS; SMAD1; TBXAS1; TWIST; ZWS1; ACHM3; ADRB3; ANKI; CA1; CA2; CCAL1; CLN8; CMT4A; CNGB3; COH1; CPP; CRH; CYP11B1; CYP11B2; DECR1; DPYS; DURS1; EBS1; ECA1; EGI; EXT1; EYA1; FGFR1; GNRH1; GSR; GULOP; HR; KCNQ3; KFM; KWE; LGCR; LPL; MCPH1; MOS; MYC; NAT1; NAT2; NBS1; PLAT; PLEC1; PRKDC; PXMP3; RP1; SCZD6; SFTPC; SGM1; SPG5A; STAR; TG; TRPS1; TTPA; VMD1; WRN; ABCA1; ABL1; ABO; ADAMTS13; AK1; ALAD; ALDH1A1; ALDOB; AMBP; AMCD1; ASS; BDMF; BSCL; C5; CDKN2A; CHAC; CLA1; CMD1B; COL5A1; CRAT; DBH; DNAI1; DYS; DYT1; ENG; FANCC; FBP1; FCMD; FRDA; GALT; GLDC; GNE; GSM1; GSN; HSD17B3; HSN1; IBM2; INVS; JBTS1; LALL; LCCS1; LCCS; LGMD2H; LMX1B; MLLT3; MROS; MSSE; NOTCH1; ORM1; PAPPA; PIP5K1B; PTCH; PTGS1; RLN1; RLN2; RMRP; ROR2; RPD1; SARDH; SPTLC1; STOM; TDFA; TEK; TMC1; TRIM32; TSC1; TYRP1; XPA; CACNB2; COLl7A1; CUBN; CXCL12; CYP17; CYP2C19; CYP2C9; EGR2; EMX2; ERCC6; FGFR2; HK1; HPSI; IL2RA; LGI1; LIPA; MAT1A; MBL2; MKI67; MXI1; NODAL; OAT; OATL3; PAX2; PCBD; PEO1; PHYH; PNL1P; PSAP;PTEN; RBP4; RDPA; RET; SFTPA1; SFTPD; SHFM3; SIAL; THC2; TLX1; TNFRSF6; UFS; UROS; AA; ABCC8; ACAT1; ALX4; AMPD3; ANC; APOA1; APOA4; APOC3; ATM; BSCL2; BWS; CALCA; CAT; CCND1; CD3E; CD3G; CD59; CDKN1C; CLN2; CNTF; CPT1A; CTSC; DDB1; DDB2; DHCR7; DLAT; DRD4; ECB2; ED4; EVR1; EXT2; F2; FSHB; FTH1; G6PT1; G6PT2; GIF; HBB; HBBP1; HBD; HBE1; HBG1; HBG2; HMBS; HND; HOMG2; HRAS; HVBS1; IDDM2; IGER; INS; JBS; KCNJ11; KCNJ1; KCNQ1; LDHA; LRP5; MEN1; MLL; MYBPC3; MYO7A; NNO1; OPPG; OPTB1; PAX6; PC; PDX1; PGL2; PGR; PORC; PTH; PTS; PVRL1; PYGM; RAG1; RAG2; ROM1; RRAS2; SAA1; SCA5; SCZD2; SDHD; SERPING1; SMPD1; TCIRG1; TCL2; TECTA; TH; TREH; TSG101; TYR; USH1C; VMD2; VRN1; WT1; WT2; ZNF145; A2M; AAAS; ACADS; ACLS; ACVRL1; ALDH2; AMHR2; AOM; AQP2; ATD; ATP2A2; BDC; C1R; CD4; CDK4; CNA1; COL2A1; CYP27B1; DRPLA; ENUR2; FEOM1; FGF23; FPF; GNB3; GNS; HAL; HBP1; HMGA2; HMN2; HPD; IGF1; KCNA1; KERA; KRAS2; KRT1; KRT2A; KRT3; KRT4; KRT5; KRT6A; KRT6B; KRTHB6; LDHB; LYZ; MGCT; MPE; MVK; MYL2; OAP; PAH; PPKB; PRB3; PTPN11; PXR1; RLS; RSN; SAS; SAX1; SCA2; SCNN1A; SMAL; SPPM; SPSMA; TBX3; TBX5; TCF1; TPI1; TSC3; ULR; VDR;

VWF; ATP7B; BRCA2; BRCD1; CLN5; CPB2; ED2; EDNRB; ENUR1; ERCC5; F10; F7; GJB2; GJB6; IPF1; MBS1; MCOR; NYS4; PCCA; RB1; RHOK; SCZD7; SGCG; SLC10A2; SLC25A15; STARP1; ZNFl98; ACHM1; ARVDI; BCH; CTAA1; DAD1; DFNB5; EML1; GALC; GCH1; IBGC1; IGH; IGHC group; IGHG1; IGHM; IGHR; IV; LTBP2; MCOP; MJD; MNG1 MPD1; MPS3C; MYH6; MYH7; NP; NPC2; PABN1; PSEN1 PYGL; RPGRIP1; SERPINA1; SERPINA3; SERPINA6; SLC7A7; SPG3A; SPTB; TCL1A; TGMI; TITF1; TMIP; TRA; TSHR; USH1A; VP; ACCPN; AHO2; ANCR; B2M; BBS4; BLM; CAPN3; CDAN1; CDAN3; CLN6; CMH3; CYP19; CYP1A1; CYP1A2; DYX1; EPB42; ETFA; EYCL3; FAH; FBN1; FES; HCVS; HEXA; IVD; LCS1; LIPC; MYO5A; OCA2; OTSC1; PWCR; RLBP1; SLC12A1; SPG6; TPM1; UBE3A; WMS; ABCC6; ALDOA; APRT; ATP2A1; BBS2; CARD15; CATM; CDH1; CETP; CHST6; CLN3; CREBBP; CTH; CTM; CYBA; CYLD; DHS; DNASE1; DPEP1; ERCC4; FANCA; GALNS; GAN; HAGH; HBA1; HBA2; HBHR; HBQ1; HBZ; HBZP; HP; HSD11B2; IL4R; LIPB; MC2R; MEFV; MHC2TA; MLYCD; MMVP1; PHKB; PHKG2; PKD1; PKDTS; PMM2; PXE; SALL1; SCA4; SCNN1B; SCNN1G; SLC12A3; TAT; TSC2; VDI; WT3; ABR; ACACA; ACADVL; ACE; ALDH3A2; APOH; ASPA; AXIN2; BCL5; BHD; BLMH; BRCA1; CACD; CCA1; CCZS; CHRNB1; CHRNE; CMT1A; COL1A1; CORD5; CTNS; EPX; ERBB2; G6PC; GAA; GALK1; GCGR; GFAP; GH1; GH2; GP1BA; GPSC; GUCY2D; ITGA2B; ITGB3; ITGB4; KRT10; KRT12; KRT13; KRT14; KRT14L1; KRT14L2; KRT14L3; KRT16; KRT16L1; KRT16L2; KRT17; KRT9; MAPT; MDB; MDCR; MGI; MHS2; MKS1; MPO; MYO15A; NAGLU; NAPB; NF1; NME1; P4HB; PAFAH1B1; PECAM1; PEX12; PHB; PMP22; PRKAR1A; PRKCA; PRKWNK4; PRP8; PRPF8; PTLAH; RARA; RCV1; RMSA1; RP17; RSS; SCN4A; SERPINF2; SGCA; SGSH; SHBG; SLC2A4; SLC4A1; SLC6A4; SMCR; SOST; SOX9; SSTR2; SYM1; SYNS1; TCF2; THRA; TIMP2; TOC; TOP2A; TP53; TRIM37; VBCH; ATP8B1; BCL2; CNSN; CORD1; CYB5; DCC; F5F8D; FECH; PEO; LAMA3; LCFS2; MADH4; MAFD1; MC2R; MCL; MYP2; NPC1; SPPK; TGFBRE; TGIF; TTR; AD2; AMH; APOC2; APOE; ATHS; BAX; BCKDHA; BCL3; BFIC; C3; CACNA1A; CCO; CEACAM5; COMP; CRX; DBA; DDU; DFNA4; DLL3; DM1; DMWD; E11S; ELA2; EPOR; ERCC2; ETFB; EXT3; EYCL1; FTL; FUT1; FUT2; FUT6; GAMT; GCDH; GPI; GUSM; HB1; HCL1; HHC2; HHC3; ICAM3; INSR; JAK3; KLK3; LDLR; LHB; LIG1; LOH19CR1; LYL1; MAN2B1; MCOLN1; MDRV; MLLT1; NOTCH3; NPHS1; OFC3; OPA3; PEPD; PRPF31; PRTN3; PRX; PSG1; PVR; RYR1;

SLC5A5; SLC7A9; STK11; TBXA2R; TGFB1; TNNI3; TYROBP; ADA; AHCY; AVP; CDAN2; CDPD1; CHED1; CHED2; CHRNA4; CST3; EDN3; EEGV1; FTLL1; GDF5; GNAS; GSS; HNF4A; JAG1; KCNQ2; MKKS; NBIA1; PCK1; PI3; PPCD; PPGB; PRNP; THBD; TOP1; AIRE; APP; CBS; COL6A1; COL6A2; CSTB; DCR; DSCR1; FPDMM; HLCS; HPE1; ITGB2; KCNE1; KNO; PRSS7; RUNX1; SOD1; TAM; ADSL; ARSA; BCR; CECR; CHEK2; COMT; CRYBB2; CSF2RB; CTHM; CYP2D6; CYP2D7P1; DGCR; DIA1; EWSR1; GGT1; MGCR; MN1; NAGA; NF2; OGS2; PDGFB; PPARA; PRODH; SCO2; SCZD4; SERPIND1; SLC5A1; SOX10; TCN2; TIMP3; TST; VCF; ABCD1; ACTL1; ADFN; AGMX2; AHDS; AIC; AIED; AIH3; ALAS2; AMCD; AMELX; ANOP1; AR; ARAF1; ARSC2; ARSE; ARTS; ARX; ASAT; ASSP5; ATP7A; ATRX; AVPR2; BFLS; BGN; BTK; BZX; C1HR; CACNA1F; CALB3; CBBM; CCT; CDR1; CFNS; CGF1; CHM; CHR39C; CIDX; CLA2; CLCN5; CLS; CMTX2; CMTX3; CND; COD1; COD2; COL4A5; COL4A6; CPX; CVD1; CYBB; DCX; DFN2; DFN4; DFN6; DHOF; DIAPH2; DKC1; DMD; DSS; DYT3; EBM; EBP; ED1; ELK1; EMD; EVR2; F8; F9; FCP1; FDPSL5; FGD1; FGS1; FMR1; FMR2; G6PD; GABRA3; GATA1; GDI1; GDXY; GJB1; GK; GLA; GPC3; GRPR; GTD; GUST; HMS1; HPRT1; HPT; HTC2; HTR2C; HYR; IDS; IHG1; IL2RG; INDX; IP1; IP2; JMS; KAL1; KFSD; L1CAM; LAMP2; MAA; MAFD2; MAOA; MAOB; MCF2; MCS; MEAX; MECP2; MF4; MGC1; MIC5; MID1; MLLT7; MLS; MRSD; MRX14; MRX1; MRX20; MRX2; MRX3; MRX40; MRXA; MSD; MTM1; MYCL2; MYP1; NDP; NHS; NPHL1; NR0B1; NSX; NYS1; NYX; OA1; OASD; OCRL; ODT1; OFD1; OPA2; OPD1; OPEM; OPN1LW; OPN1MW; OTC; P3; PDHA1; PDR; PFC; PFKFB1; PGK1; PGK1P1; PGS; PHEX; PHKA1; PHKA2; PHP; PIGA; PLP1; POF1; POLA; POU3F4; PPMX; PRD; PRPS1; PRPS2; PRS; RCCP2; RENBP; RENS1; RP2; RP6; RPGR; RPS4X; RPS6KA3; RS1; S11; SDYS; SEDL; SERPINA7; SH2D1A; SHFM2; SLC25A5; SMAX2; SRPX; SRS; STS; SYN1; SYP; TAF1; TAZ; TBX22; TDD; TFE3; THAS; THC; TIMM8A; TIMP1; TKCR; TNFSF5; UBE1; UBE2A; WAS; WSN; WTS; WWS; XIC; XIST; XK; XM; XS; ZFX; ZIC3; ZNF261; ZNF41; ZNF6; AMELY; ASSP6; AZF1; AZF2; DAZ; GCY; RPS4Y; SMCY; SRY; ZFY; ABAT; AEZ; AFA; AFD1; ASAH1; ASD1; ASMT; CCAT; CECR9; CEPA; CLA3; CLN4; CSF2RA; CTS1; DF; DIH1; DWS; DYT2; DYT4; EBR3; ECT; EEF1A1L14; EYCL2; FANCB; GCSH; GCSL; GIP; GTS; HHG; HMI; HOAC; HOKPP2; HRPT1; HSD3B3; HTC1; HV1S; ICHQ; ICR1; ICR5; IL3RA; KAL2; KMS; KRT18; KSS; LCAT; LHON; LIMM; MANBB; MCPH2; MEB; MELAS; MIC2; MPFD;

MS; MSS; MTATP6; MTCO1; MTCO3; MTCYB; MTND1; MTND2; MTND4; MTND5; MTND6; MTRNR1; MTRNR2; MTTE; MTTG; MTTI; MTTK; MTTL1; MTTL2; MTTN; MTTP; MTTS1; NAMSD; OCD1; OPD2; PCK2; PCLD; PCOS1; PFKM; PKD3; PRCA1; PRO1; PROP1; RBS; RFXAP; RP; SHOX; SLC25A6; SPG5B; STO; SUOX; THM; or TTD.

11. The use of any one of claims 7-10, wherein said disorder and said recombinant protein, respectively, are selected from the group consisting of: anemia in a subject and said recombinant protein is erythropoietin (EPO); vasospasm in a subject and said recombinant protein is inducible nitric oxide synthase (iNOS); restenosis of a blood vessel following a procedure that enlarges the blood vessel and said recombinant protein is a heat shock protein, thereby decreasing an incidence of a restenosis in a subject; cystic fibrosis in a subject and said recombinant protein is Cystic Fibrosis Transmembrane Conductance Regulator (CFTR), thereby treating cystic fibrosis in a subject; X-linked agammaglobulinemia in a subject and said recombinant protein is Bruton's tyrosine kinase, thereby treating an X-linked agammaglobulinemia; adenosine deaminase severe combined immunodeficiency (ADA SCID) in a subject and said recombinant protein is adenosine deaminase (ADA), thereby treating an ADA SCID; immune responsiveness due to a skin pathology and said recombinant protein is ecto-nucleoside triphosphate diphosphohydrolase, thereby reducing immune responsiveness of the skin and improving skin pathology; Niemann-Pick disease and said recombinant protein is Sphingomyelinase, thereby treating Niemann-Pick disease; mucopolysaccharidosis and said recombinant protein is alpha-L-iduronidase, iduronate-2-sulfatase, thereby a treating mucopolysaccharide storage disorder; Gaucher's disease and said recombinant protein is acid beta-glucosidase, thereby treating a Gaucher's disease; and Fabry's disease and said recombinant protein is alpha-galactosidase A, thereby treating a Fabry's disease.

12. A method for reducing immunogenicity of a preparation composed of *in vitro*-synthesized RNA molecules comprising:

purifying a preparation of *in vitro*-synthesized RNA molecules obtained by *in vitro* transcription (IVT) that comprises a modified nucleoside selected from Ψ, $m^1\Psi$, m^5U, mo^5U and s^2U in place of uridine (U) using a process that removes RNA contaminant molecules that are immunogenic and toxic to the cell by inducing an innate immune response, to generate a

purified RNA preparation that is free of RNA contaminant molecules, including double-stranded RNA molecules, and that has reduced immunogenicity for mammalian cells;

wherein said *in vitro*-synthesized RNA is mRNA that encodes a recombinant protein;

wherein said reduced immunogenicity is detectable using an *in vitro* MDDC immunogenicity assay by measuring an amount of IFN-α or TNF-α cytokine secreted by human or murine monocyte-derived dendritic cells (MDDCs) transfected with said purified RNA preparation that is at least 5-fold or at least 10-fold less than the amount of said cytokine that is secreted from MDDCs transfected with said preparation of *in vitro*-synthesized RNA molecules not subjected to said purifying;

wherein the decrease in immunogenicity of said purified RNA preparation comprising said in vitro-synthesized RNA molecules is such that that an effective amount can be administered without triggering a detectable immune response.

13. The method of claim12, wherein said purifying comprises the step of treating the RNA preparation with E. *coli* RNase III or another RNase III enzyme that digests double-stranded RNA greater than about twelve basepairs to short double-stranded RNA fragments such that short RNase III digestion products are generated, and separating said short RNase III digestion products from said purified RNA preparation such that said purified RNA preparation is free from said short RNase III digestion products.

14. The method of claim 13, wherein said IVT is performed in a reaction mixture where all or a desired fraction of the UTP is replaced by the 5'-triphosphate of a modified nucleoside selected from Ψ, $m^1Ψ$, m^5U, mo^5U and s^2U.

15. The method of claim 14, wherein said IVT further comprises replacing all or a desired fraction of the CTP with the 5'-triphosphate-derivative of m^5C.

16. The method of any of claims 12-15, wherein at least one of the following apply to said *in vitro*-synthesized RNA molecules:

a) they further comprise a poly-A tail having 50-100 or greater than 150 A nucleotides;

b) greater than 98% of said in vitro-synthesized RNA molecules comprise a 5' cap;

c) approximately 100% of said RNA molecules further comprise a 5' cap with a cap0 or a cap1 structure;

d) they further comprise at least one particular sequence selected from the group consisting of a heterologous 5' UTR sequence, a Kozak sequence, an IRES sequence, and 3' UTR sequence;

e) they further comprise a 5' UTR sequence or 3' UTR sequence from a Xenopus or human alpha globin or beta globin mRNA, or wherein said 5' UTR sequence is a sequence from a tobacco etch virus (TEV) RNA;

f) they are mRNA (messenger RNA) molecules;

g) they comprise an open reading frame that encodes a functional protein;

h) they have a length comprising at least 120 nucleotides;

i) they have a length comprising at least 250 nucleotides;

j) they comprise s^2U in place of U and the fraction of said uridine nucleotides that are replaced by s^2U is selected from: less than 20%, less than 40% and less than 70%: or

k) they encode a detectable enzyme selected from firefly luciferase, Renilla luciferase, bacterial beta-galactosidase (lacZ), and green fluorescent protein (GFP)

l) they lack immunogenicity, enabling repeated delivery into a mammalian cell without generation of inflammatory cytokines; or

m) they are mRNA (messenger RNA) molecules.

17. Use of a purified RNA preparation obtained using the method of any of claims 1-6 or the method of any of claims 12-16 in the preparation of a medicament for treatment of a disorder selected from the group consisting of an inborn error of metabolism, a monogenic disorder, an infectious disease, an acquired disorder and a cancer in a subject, wherein said medicament is to be administered to said subject.

18. The use of claim 17, wherein said treatment is a therapeutic treatment or a prophylactic treatment.

19. Use of any one of claims 17-18, wherein a single dose of said purified preparation is to be administered to said subject without triggering a detectable immune response, or

wherein said purified preparation is to be administered in a daily dose, a weekly dose, a monthly dose or a series of a defined number of doses to a subject, without triggering a detectable immune response.

20. The use of any of claims 17-19, wherein the medicament further comprises at least one of the following:

- a pharmaceutically acceptable carrier or diluent; and/or

- a transfection reagent, wherein said transfection reagent is selected from the group consisting of a lipid-based transfection reagent such as a cationic lipid reagent, a protein-based transfection reagent, a polyethyleneimine-based transfection reagent and calcium phosphate.

21. The use of any one of claims 17 - 20, wherein said medicament is to be administered to said subject parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously, intra-dermally, subcutaneously, intra-peritonealy, intra-ventricularly, intra-cranially, intra-vaginally or intra-tumorally, and/or, wherein said medicament is to be administered at a dose selected from among a one-time dose, a daily dose, a weekly dose, a monthly dose and a series of a defined number of doses.

22. Use of said purified RNA preparation made using the method of any one of claims 1-6 or 12-16 in the preparation of a pharmaceutical composition for the treatment of a disorder selected from the group consisting of a monogenic disorder, an infectious disease, an acquired disorder and a cancer, wherein said pharmaceutical composition is to be administered to a subject with said disorder, without triggering a detectable immune response, and wherein the length of said RNA molecules is at least 250 nucleotides.

23. The use of claims 22, wherein said treatment is a therapeutic treatment or a prophylactic treatment.

SEQUENCE LISTING

<110> Dahl, Gary
Person, Anthony D.
Kariko, Katalin
Weissman, Drew

<120> RNA Containing Modified Nucleosides and Methods of Use Thereof

<130> EPICE-31302/WO-2/ORD

<150> US 61/267,312
<151> 2009-12-07

<150> US 11/990,646
<151> 2009-03-27

<150> PCT/US 2006/032372
<151> 2006-08-21

<150> US 60/710,164
<151> 2005-08-23

<160> 17

<170> PatentIn version 3.5

<210> 1
<211> 1864
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 1
ggaauucuca acacaacaua uacaaaacaa acgaaucuca agcaaucaag cauucuacuu 60

cuauugcagc aauuuaaauc auuucuuuua aagcaaaagc aauuuucuga aaauuuucac 120

cauuuacgaa cgauagccau ggaagacgcc aaaaacauaa agaaaggccc ggcgccauuc 180

uauccucuag aggauggaac cgcuggagag caacugcaua aggcuaugaa gagauacgcc 240

cugguuccug gaacaauugc uuuuacagau gcacauaucg aggugaacau cacguacgcg 300

gaauacuucg aaauguccgu ucgguuggca gaagcuauga aacgauaugg gcugaauaca 360

aaucacagaa ucgucguaug cagugaaaac ucucuucaau ucuuuaugcc gguguugggc 420

gcguuauuua ucggaguugc aguugcgccc gcgaacgaca uuuauaauga acgugaauug 480

cucaacagua ugaacauuuc gcagccuacc guaguguuug uuuccaaaaa gggguugcaa 540

aaaauuuuga acgugcaaaa aaaauuacca auaauccaga aaauuauuau cauggauucu 600

aaaacggauu accagggauu ucagucgaug uacacguucg ucacaucuca ucuaccuccc 660

gguuuuaaug aauacgauuu uguaccagag uccuuugauc gugacaaaac aauugcacug 720

auaaugaauu ccucuggauc uacuggguua ccuaagggug uggcccuucc gcauagaacu 780

gccugcguca gauucucgca ugccagagau ccuauuuuug gcaaucaaau cauuccggau 840

acugcgauuu uaaguguugu uccauuccau cacgguuuug gaauguuuac uacacucgga 900

uauuugauau guggauuucg agucgucuua auguauagau uugaagaaga gcuguuuuua 960

cgaucccuuc aggauuacaa aauucaaagu gcguugcuag uaccaacccu auuuucauuc 1020

uucgccaaaa gcacucugau ugacaaauac gauuuaucua auuuacacga aauugcuucu 1080

gggggcgcac cucuuucgaa agaagucggg gaagcgguug caaaacgcuu ccaucuucca 1140

gggauacgac aaggauaugg gcucacugag acuacaucag cuauucugau uacacccgag 1200

ggggaugaua aaccgggcgc ggucgguaaa guuguuccau uuuuugaagc gaagguugug 1260

gaucuggaua ccgggaaaac gcugggcguu aaucagagag gcgaauuaug ugucagagga 1320

ccuaugauua uguccgguua uguaaacaau ccggaagcga ccaacgccuu gauugacaag 1380

gauggauggc uacauucugg agacauagcu uacugggacg aagacgaaca cuucuucaua 1440

guugaccgcu ugaagucuuu aauuaaauac aaaggauauc agguggcccc cgcugaauug 1500

gaaucgauau uguuacaaca ccccaacauc uucgacgcgg gcguggcagg ucuucccgac 1560

gaugacgccg gugaacuucc cgccgccguu guuguuuugg agcacggaaa gacgaugacg 1620

gaaaaagaga ucguggauua cguggccagu caaguaacaa ccgcgaaaaa guugcgcgga 1680

ggaguugugu uuguggacga aguaccgaaa ggucuuaccg gaaaacucga cgcaagaaaa 1740

aucagagaga uccucauaaa ggccaagaag ggcggaaagu ccaaauugua aaauguaacu 1800

cuagaggauc cccaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1860

aaca 1864

<210> 2
<211> 1571
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 2
ggcuagccac caugacuucg aaaguuuaug auccagaaca aaggaaacgg augauaacug 60

guccgcagug gugggccaga uguaaacaaa ugaauguucu ugauucauuu auuaauuauu 120

augauucaga aaaacaugca gaaaaugcug uuauuuuuuu acaugguaac gcggccucuu 180

cuuauuuaug gcgacauguu gugccacaua uugagccagu agcgcggugu auuauaccag 240

accuuauugg uaugggcaaa ucaggcaaau cugguaaugg uucuuauagg uuacuugauc 300

auuacaaaua ucuuacugca ugguuugaac uucuuaauuu accaaagaag aucauuuuug 360

ucggccauga uuggggugcu uguuuggcau uucauuauag cuaugagcau caagauaaga 420

ucaaagcaau aguucacgcu gaaaguguag uagaugugau ugaaucaugg gaugaauggc 480

cugauauuga agaagauauu gcguugauca aaucugaaga aggagaaaaa augguuuugg 540

agaauaacuu cuucguggaa accauguugc caucaaaaau caugagaaag uuagaaccag 600

aagaauuugc agcauaucuu gaaccauuca aagagaaagg ugaaguucgu cguccaacau 660

uaucauggcc ucgugaaauc ccguuaguaa aaggugguaa accugacguu guacaaauug 720

uuaggaauua uaaugcuuau cuacgugcaa gugaugauuu accaaaaaug uuuauugaau 780

cggacccagg auucuuuucc aaugcuauug uugaaggugc caagaaguuu ccuaauacug 840

aauuugucaa aguaaaaggu cuucauuuuu cgcaagaaga ugcaccugau gaaaugggaa 900

aauauaucaa aucguucguu gagcgaguuc ucaaaaauga acaaaugucg acgggggccc 960

cuaggaauuu uuuagggaag aucuggccuu ccuacaaggg aaggccaggg aauuuucuuc 1020

agagcagacc agagccaaca gccccaccag aagagagcuu caggucuggg guagagacaa 1080

caacucccccc ucagaagcag gagccgauag acaaggaacu guauccuuua acuucccuca 1140

gaucacucuu uggcaacgac cccucgucac aauaaagaua gggggcaac uaaagggauc 1200

ggccgcuucg agcagacaug auaagauaca uugaugaguu uggacaaacc acaacuagaa 1260

ugcagugaaa aaaaugcuuu auuugugaaa uuugugaugc uauugcuuua uuuguaacca 1320

uuauaagcug caauaaacaa guuaacaaca acaauugcau ucauuuuaug uuucagguuc 1380

agggggaggu gugggagguu uuuuaaagca aguaaaaccu cuacaaaugu gguaaaaucg 1440

auaaguuuaa acagauccag guggcacuuu ucggggaaau gugcgcggaa ccccuauuug 1500

uuuauuuuuc uaaauacauu caaauaugua uccgcucaug agacaauaac ccugauaaau 1560

gcuucaauaa u 1571

<210> 3
<211> 726
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 3
gggaauuugg cccucgaggc caagaauucg gcacgaggca cgcggccagc cagcagacag 60

aggacucuca uuaaggaagg uguccugugc ccugacccua caagaugcca agagaagaug 120

cucacuucau cuaugguuac cccaagaagg ggcacggcca cucuuacacc acggcugaag 180

aggccgcugg gaucggcauc cugacaguga uccugggagu cuuacugcuc aucggcuguu 240

gguauuguag aagacgaaau ggauacagag ccuugaugga uaaaagucuu cauguuggca 300

cucaaugugc cuuaacaaga agaugcccac aagaaggguu ugaucaucgg gacagcaaag 360

ugucucuuca agagaaaaac ugugaaccug ugguucccaa ugcuccaccu gcuuaugaga 420

aacucucugc agaacaguca ccaccaccuu auucaccuua agagccagcg agacaccuga 480

gacaugcuga aauuauuucu cucacacuuu ugcuugaauu uaauacagac aucuaauguu 540

cuccuuugga augguguagg aaaaaugcaa gccaucucua auaauaaguc aguguuaaaa 600

uuuuaguagg uccgcuagca guacuaauca ugugaggaaa ugaugagaaa uauuaaauug 660

ggaaaacucc aucaauaaau guugcaaugc augauaaaaa aaaaaaaaaa aaaaacugcg 720

gccgca 726

<210> 4
<211> 712
<212> RNA

<213> Artificial Sequence

<220>
<223> Synthetic

<400> 4
gggaauaagc uugcggccgc aguuuuuuuu uuuuuuuuuu uuaucaugca uugcaacauu 60

uauugaugga guuuucccaa uuuaauauuu cucaucauuu ccucacauga uuaguacugc 120

uagcggaccu acuaaaauuu uaacacugac uuauuauuag agauggcuug cauuuuuccu 180

acaccauucc aaaggagaac auuagauguc uguauaaauu caagcaaaag ugugagagaa 240

auaauuucag caugucucag gugucucgcu ggcucuuaag gugaauaagg uggugugac 300

uguucugcag agaguuucuc auaagcaggu ggagcauugg gaaccacagg uucacaguuu 360

uucucuugaa gagacacuuu gcugucccga ugaucaaacc cuucuugugg gcaucuucuu 420

guuaaggcac auugagugcc aacaugaaga cuuuuaucca ucaaggcucu guauccauuu 480

cgucuucuac aauaccaaca gccgaugagc aguaagacuc ccaggaucac ugucaggaug 540

ccgaucccag cggccucuuc agccguggug uaagaguggc cgugccccuu cuuggguaa 600

ccauagauga agugagcauc uucucuuggc aucuuguagg gucagggcac aggacaccuu 660

ccuuaaugag aguccucugu cugcuggcug gccgcgugcc ucgugccgaa uu 712

<210> 5
<211> 494
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 5
gggagaccca agcuggcuag cagucaucca acagaaucau gagacagacu uugccuugua 60

ucuacuuuug ggggggccuu uugcccuuug ggaugcugug ugcauccucc accaccaagu 120

gcacuguuag ccaugaaguu gcugacugca gccaccugaa guugacucag guacccgaug 180

aucuacccac aaacauaaca guguugaacc uuacccauaa ucaacucaga agauuaccag 240

ccgccaacuu cacaagguau agccagcuaa cuagcuugga uguaggauuu aacaccaucu 300

caaaacugga gccagaauug ugccagaaac uucccauguu aaaaguuuug aaccuccagc 360

acaaugagcu aucucaacuu ucugauaaaa ccuuugccuu cugcacgaau uugacugaac 420

uccaucucau guccaacuca auccagaaaa uuaaaaauaa ucccuuuguc aagcagaaga 480

auuuaaucac auua 494

<210> 6
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 6

uggauccggc uuugagaucu u 21

<210> 7
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> modified_base
<222> (11)..(11)
<223> um

<400> 7
uggauccggc uuugagaucu u 21

<210> 8
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> modified_base
<222> (6)..(6)
<223> m5c

<400> 8
uggauccggc uuugagaucu u 21

<210> 9
<211> 21
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<220>
<221> modified_base
<222> (12)..(12)
<223> p

<400> 9
uggauccggc uuugagaucu u 21

<210> 10
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 10
gggagacagg gguguccgcc auuuccaggu u 31

<210> 11
<211> 31
<212> RNA
<213> Artificial Sequence

<220>
<223> Synthetic

<400> 11
gggagacagg cuauaacuca cauaauguau u 31

<210> 12
<211> 1440
<212> RNA
<213> Homo sapiens

<400> 12
augaggcagc caccuggcga gucugacaug gcugucagcg acgcgcugcu cccaucuuuc 60

uccacguucg cgucuggccc ggcgggaagg gagaagacac ugcgucaagc aggugccccg 120

aauaaccgcu ggcgggagga gcucucccac augaagcgac uucccccagu gcuucccggc 180

cgccccuaug accuggcggc ggcgaccgug gccacagacc uggagagcgg cggagccggu 240

gcggcuugcg gcgguagcaa ccuggcgccc cuaccucgga gagagaccga ggaguucaac 300

gaucuccugg accuggacuu uauucucucc aauucgcuga cccauccucc ggagucagug 360

gccgccaccg uguccucguc agcgucagcc uccucuucgu cgucgccguc gagcagcggc 420

ccugccagcg cgcccuccac cugcagcuuc accuauccga uccgggccgg gaacgacccg 480

ggcguggcgc cgggcggcac gggcggaggc cuccucuaug gcagggaguc cgcucccccu 540

ccgacggcuc ccuucaaccu ggcggacauc aacgacguga gccccucggg cggcuucgug 600

gccgagcucc ugcggccaga auuggacccg guguacauuc cgccgcagca gccgcagccg 660

ccagguggcg ggcugauggg caaguucgug cugaaggcgu cgcugagcgc cccuggcagc 720

gaguacggca gcccgucggu caucagcguc agcaaaggca gcccugacgg cagccacccg 780

gugguggugg cgcccuacaa cggcgggccg ccgcgcacgu gccccaagau caagcaggag 840

gcggucucuu cgugcaccca cuugggcgcu ggacccccuc ucagcaaugg ccaccggccg 900

gcugcacacg acuucccccu ggggcggcag cuccccagca ggacuacccc gacccugggu 960

cuugaggaag ugcugagcag cagggacugu cacccugccc ugccgcuucc ucccggcuuc 1020

caucccccacc cggggcccaa uuacccaucc uuccugcccg aucagaugca gccgcaaguc 1080

ccgccgcucc auuaccaaga gcucaugcca cccgguuccu gcaugccaga ggagcccaag 1140

ccaaagaggg gaagacgauc guggccccgg aaaaggaccg ccacccacac uugugauuac 1200

gcgggcugcg gcaaaaccua cacaaagagu ucccaucuca aggcacaccu gcgaacccac 1260

acaggugaga aaccuuacca cugugacugg gacggcugug gauggaaauu cgcccgcuca 1320

gaugaacuga ccaggcacua ccguaaacac acggggcacc gcccguucca gugccaaaaa 1380

ugcgaccgag cauuuuccag gucggaccac cucgccuuac acaugaagag gcauuuuuaa 1440

<210> 13
<211> 529
<212> RNA
<213> Homo sapiens

<400> 13
augggcuccg uguccaacca gcaguuugca gguggcugcg ccaaggcggc agaagaggcg 60

cccgaggagg cgccggagga cgcggcccgg gcggcggacg agccucagcu gcugcacggu 120

gcgggcaucu guaaggguuu caacgugcgc auggggguucg gcuuccuguc caugaccgcc 180

cgcgccgggg ucgcgcucga ccccccagug gaugucuuug ugcaccagag uaagcugcac 240

auggaagggu uccggagcuu gaaggagggu gaggcagugg aguucaccuu uaagaaguca 300

gccaaggguc uggaauccau ccgugucacc ggaccuggug gaguauucug uauugggagu 360

gagaggcggc caaaaggaaa gagcaugcag aagcgcagau caaaaggaga caggugcuac 420

aacuguggag gucuagauca ucaugccaag gaaugcaagc ugccacccca gcccaagaag 480

ugccacuucu gccagagcau cagccauaug guagccucau guccgcuga 529

<210> 14
<211> 1365
<212> RNA
<213> Homo sapiens

<400> 14
auggauuuuu uucggguagu ggaaaaccag cagccucccg cgacgaugcc ccucaacguu 60

agcuucacca acaggaacua ugaccucgac uacgacucgg ugcagccgua uuucuacugc 120

gacgaggagg agaacuucua ccagcagcag cagcagagcg agcugcagcc cccggcgccc 180

agcgaggaua ucuggaagaa auucgagcug cugcccaccc cgccccuguc cccuagccgc 240

cgcuccgggc ucugcucgcc cuccuacguu gcggucacac ccuucucccu ucggggagac 300

aacgacggcg guggcgggag cuucuccacg gccgaccagc uggagauggu gaccgagcug 360

cugggaggag acauggugaa ccagaguuuc aucugcgacc cggacgacga gaccuucauc 420

aaaaacauca ucauccagga cuguaugugg agcggcuucu cggccgccgc caagcucguc 480

ucagagaagc uggccuccua ccaggcugcg cgcaaagaca gcggcagccc gaaccccgcc 540

cgcggccaca gcgucugcuc caccuccagc uuguaccugc aggaucugag cgccgccgcc 600

ucagagugca ucgaccccuc gguggucuuc cccuacccuc ucaacgacag cagcucgccc 660

aaguccugcg ccucgcaaga cuccagcgcc uucucuccgu ccucggauuc ucugcucucc 720

ucgacggagu ccucccccgca gggcagcccc gagccccugg ugcuccauga ggagacaccg 780

cccaccacca gcagcgacuc ugaggaggaa caagaagaug aggaagaaau cgauguuguu 840

ucuguggaaa agaggcaggc uccuggcaaa aggucagagu cuggaucacc uucugcugga 900

ggccacagca aaccuccuca cagcccacug guccucaaga ggugccacgu cuccacacau 960

cagcacaacu acgcagcgcc ucccuccacu cggaaggacu auccugcugc caagaggguc 1020

aaguuggaca gugucagagu ccugagacag aucagcaaca accgaaaaug caccagcccc 1080

agguccucgg acaccgagga gaaugucaag aggcgaacac acaacgucuu ggagcgccag 1140

aggaggaacg agcuaaaacg gagcuuuuuu gcccugcgug accagauccc ggaguuggaa 1200

aacaaugaaa aggcccccaa gguaguuauc cuuaaaaaag ccacagcaua cauccugucc 1260

guccaagcag aggagcaaaa gcucauuucu gaagaggacu uguugcggaa acgacgagaa 1320

caguugaaac acaaacuuga acagcuacgg aacucuugug cguaa 1365

<210> 15
<211> 918
<212> RNA
<213> Homo sapiens

<400> 15
augagugugg auccagcuug uccccaaagc uugccuugcu uugaagcauc cgacuguaaa 60

gaaucuucac cuaugccugu gauuuguggg ccugaagaaa acuauccauc cuugcaaaug 120

ucuucugcug agaugccuca cacagagacu gucucuccuc uuccuuccuc cauggaucug 180

cuuauucagg acagcccuga uucuuccacc agucccaaag gcaaacaacc cacuucugca 240

gagaauagug ucgcaaaaaa ggaagacaag gucccgguca agaaacagaa gaccagaacu 300

guguucucuu ccacccagcu guguguacuc aaugauagau uucagagaca gaaauaccuc 360

agccuccagc agaugcaaga acucuccaac auccugaacc ucagcuacaa acaggugaag 420

accugguucc agaaccagag aaugaaaucu aagagguggc agaaaaacaa cuggccgaag 480

aauagcaaug gugugacgca gaaggccuca gcaccuaccu accccagccu cuacucuucc 540

uaccaccagg gaugccuggu gaacccgacu gggaaccuuc caauguggag caaccagacc 600

uggaacaauu caaccuggag caaccagacc cagaacaucc aguccuggag caaccacucc 660

uggaacacuc agaccuggug cacccaaucc uggaacaauc aggccuggaa cagucccuuc 720

uauaacugug gagaggaauc ucugcagucc ugcaugcacu uccagccaaa uucuccugcc 780

agugacuugg aggcugccuu ggaagcugcu ggggaaggcc uuaauguaau acagcagacc 840

acuagguauu uuaguacucc acaaaccaug gauuuauucc uaaacuacuc caugaacaug 900

caaccugaag acguguga 918

<210> 16
<211> 1083
<212> RNA
<213> Homo sapiens

<400> 16
auggcgggac accuggcuuc agauuuugcc uucucgcccc cuccaggugg uggaggugau 60

gggccagggg ggccggagcc gggcuggguu gauccucgga ccuggcuaag cuuccaaggc 120

ccuccuggag ggccaggaau cgggccgggg guugggccag gcucugaggu gugggggauu 180

cccccaugcc ccccgccgua ugaguucugu ggggggaugg cguacugugg gccccagguu 240

ggaguggggc uagugcccca aggcggcuug gagaccucuc agccugaggg cgaagcagga 300

gucggggugg agagcaacuc cgauggggcc uccccggagc ccugcaccgu caccccuggu 360

gccgugaagc uggagaagga gaagcuggag caaaacccgg aggaguccca ggacaucaaa 420

gcucugcaga aagaacucga gcaauuugcc aagcuccuga agcagaagag gaucacccug 480

ggauauacac aggccgaugu ggggcucacc cuggggguuc uauuugggaa gguauucagc 540

caaacgacca ucugccgcuu ugaggcucug cagcuuagcu ucaagaacau guguaagcug 600

cggcccuugc ugcagaagug gguggaggaa gcugacaaca augaaaaucu ucaggagaua 660

ugcaaagcag aaacccucgu gcaggcccga aagagaaagc gaaccaguau cgagaaccga 720

gugagaggca accuggagaa uuuguuccug cagugcccga aacccacacu gcagcagauc 780

agccacaucg cccagcagcu ugggcucgag aaggaugugg uccgagugug guucuguaac 840

cggcgccaga agggcaagcg aucaagcagc gacuaugcac aacgagagga uuuugaggcu 900

gcuggucuc cuuucucagg gggaccagug uccuuuccuc uggccccagg gccccauuuu 960

gguaccccag gcuaugggag cccucacuuc acugcacugu acuccucggu cccuuucccu 1020

gaggggaag ccuuuccccc ugucucuguc accacucugg gcucucccau gcauucaaac 1080

uga 1083

<210> 17
<211> 954
<212> RNA
<213> Homo sapiens

<400> 17
auguacaaca ugauggagac ggagcugaag ccgccgggcc cgcagcaaac uucgggggc 60

ggcggcggca acuccaccgc ggcggcggcc ggcggcaacc agaaaaacag cccggaccgc 120

gucaagcggc ccaugaaugc cuucauggug uggucccgcg ggcagcggcg caagauggcc 180

caggagaacc ccaagaugca caacucggag aucagcaagc gccugggcgc cgaguggaaa 240

cuuuugucgg agacggagaa gcggccguuc aucgacgagg cuaagcggcu gcgagcgcug 300

cacaugaagg agcacccgga uuauaaauac cggccccggc ggaaaaccaa gacgcucaug 360

aagaaggaua aguacacgcu gcccggcggg cugcuggccc ccggcggcaa uagcauggcg 420

agcggggucg ggguggggcgc cggccugggc gcgggcguga accagcgcau ggacaguuac 480

gcgcacauga acggcuggag caacggcagc uacagcauga ugcaggacca gcugggcuac 540

ccgcagcacc cgggccucaa ugcgcacggc gcagcgcaga ugcagcccau gcaccgcuac 600

gacgugagcg cccugcagua caacuccaug accagcucgc agaccuacau gaacggcucg 660

cccaccuaca gcauguccua cucgcagcag ggcaccccug gcauggcucu uggcuccaug 720

gguucggugg ucaaguccga ggccagcucc agccccccug ugguuaccuc uuccucccac 780

uccagggcgc ccugccaggc cggggaccuc cgggacauga ucagcaugua ucucccggc 840

gccgaggugc cggaacccgc cgcccccagc agacuucaca ugucccagca cuaccagagc 900

ggcccggugc ccggcacggc cauuaacggc acacugcccc ucucacacau guga 954

RNA PREPARATIONS COMPRISING PURIFIED MODIFIED RNA FOR REPROGRAMMING CELLS

ABSTRACT

The present invention provides compositions and methods for reprogramming somatic cells using purified RNA preparations comprising single-strand mRNA encoding an iPS cell induction factor. The purified RNA preparations are preferably substantially free of RNA contaminant molecules that: i) would activate an immune response in the somatic cells, ii) would decrease expression of the single-stranded mRNA in the somatic cells, and/or iii) active RNA sensors in the somatic cells. In certain embodiments, the purified RNA preparations are substantially free of partial mRNAs, double-stranded RNAs, un-capped RNA molecules, and/or single-stranded run-on mRNAs.

(FIG. 1)