

IAP BIR DOMAIN BINDING COMPOUNDS

FIELD OF THE INVENTION

- 5 The present invention concerns bridged compounds that bind to IAP BIR domains, and which are useful for treating proliferative disorders and disorders of dysregulated apoptosis, such as cancer.

BACKGROUND OF THE INVENTION

- 10 Apoptosis, or programmed cell death, typically occurs in the normal development and maintenance of healthy tissues in multicellular organisms. It is a complex process which results in the removal of damaged, diseased or developmentally redundant cells, in the absence of signs of inflammation or necrosis.
- 15 Intrinsic apoptotic pathways are known to be dysregulated, most particularly in cancer and lymphoproliferative syndromes, as well as autoimmune disorders such as multiple sclerosis, in neurodegenerative diseases and in inflammation. As well, alterations in a host apoptotic response have been described in the development or maintenance of viral and bacterial infections.
- 20 The caspases are a family of proteolytic enzymes from the class of cysteine proteases which are known to initiate and execute apoptosis. In normal cells, the caspases are present as inactive zymogens, which are catalytically activated following external signals, for example those resulting from ligand driven Death Receptor activation, such as
- 25 cytokines or immunological agents, or by release of mitochondrial factors, such as cytochrome C following genotoxic, chemotoxic, or radiation-induced cellular injury. The Inhibitors of Apoptosis Proteins (IAPs) constitute a family of proteins which are capable of binding to and inhibiting the caspases, thereby suppressing cellular apoptosis. Because of their central role in regulating Caspase activity, the IAPs are capable of inhibiting
- 30 programmed cell death from a wide variety of triggers, which include loss of homeostatic, or endogenous cellular growth control mechanisms, as well as chemotherapeutic drugs and irradiation.

The IAPs contain one to three homologous structural domains known as baculovirus IAP repeat (BIR) domains. They may also contain a RING zinc finger domain at the C-terminus, with a capability of inducing ubiquitylation of IAP-binding molecules via its E3 ligase function. The human IAPs, XIAP, HIAP1 (also referred to as cIAP2), and HIAP2 (cIAP1) each have three BIR domains, and a carboxy terminal RING zinc finger. Another IAP, NAIP, has three BIR domains (BIR1, BIR2 and BIR3), but no RING domain, whereas Livin, TsIAP and MLIAP have a single BIR domain and a RING domain. The X chromosome-linked inhibitor of apoptosis (XIAP) is an example of an IAP which can inhibit the initiator caspase, known as caspase-9, and the effector caspases, Caspase-3 and Caspase-7, by direct binding. It can also induce the removal of caspases through the ubiquitylation-mediated proteasome pathway via the E3 ligase activity of a RING zinc finger domain. It is via the BIR3 domain that XIAP binds to and inhibits caspase-9. The linker-BIR2 domain of XIAP inhibits the activity of caspases-3 and -7. The BIR domains have also been associated with the interactions of IAPs with tumor necrosis factor-receptor associated factor (TRAFs)-1 and -2, and to TAB1, as adaptor proteins effecting survival signaling through NFkB activation. The IAPs thus function as a direct brake on the apoptosis cascade, by preventing the action of, or inhibiting active caspases and by re-directing cellular signaling to a pro-survival mode.

Progress in the cancer field has led to a new paradigm in cancer biology wherein neoplasia may be viewed as a failure of cancer cells to execute normal pathways of apoptosis. Normal cells receive continuous feedback from their environment through various intracellular and extracellular factors, and "commit suicide" if removed from this context. This induction of apoptosis is achieved by activation of the caspase cascade. Cancer cells, however, gain the ability to overcome or bypass this apoptosis regulation and continue with inappropriate proliferation. The majority of treatments for cancer induce at least a partial apoptotic response in the cancer target cell, resulting in remission or initiation of tumor regression. In many cases, however, residual cells which are apoptosis-resistant are capable of escaping therapy and continuing the process of oncogenic/genetic change, resulting in the emergence of highly drug-resistant, metastatic disease which overcomes our ability to effectively treat the disease. Furthermore, most cancer therapies, including radiation therapy and traditional chemotherapy do induce apoptosis in cancer cells, but cause additional cellular injury, due to their lack of specificity in inducing apoptosis solely in cancer cells. The need to improve the specificity/potency of pro-apoptosis agents used to treat cancer, and indeed other proliferative disorders, is

important because of the benefits in decreasing the side effects associated with administration of these agents. Therefore, finding novel means of inducing apoptosis in cancer cells is a highly desired medical need and its solution offers the possibility of entirely new treatments for cancer.

5

A growing body of data indicates that cancer cells may avoid apoptosis by the sustained over-expression of one or more members of the IAP family of proteins, as documented in many primary tumor biopsy samples, as well as most established cancer cell lines.

Epidemiological studies have demonstrated that over-expression of the various IAPs is associated with poor clinical prognosis and survival. For XIAP this is shown in cancers as diverse as leukemia and ovarian cancer. Over expression of HIAP1 and HIAP2 resulting from the frequent chromosome amplification of the 11q21-q23 region, which encompasses both, has been observed in a variety of malignancies, including medulloblastomas, renal cell carcinomas, glioblastomas, and gastric carcinomas. (X)IAP negative regulatory molecules such as XAF, appear to be tumor suppressors, which are very frequently lost in clinical cancers. Thus, by their ability to suppress the activation and execution of the intrinsic mediators of apoptosis, the caspases, the IAPs may directly contribute to tumor progression and resistance to pharmaceutical intervention. Induction of apoptosis in cancer cells by the use of potent small molecules which bind to specific IAP domains is the subject of this invention.

15
20

We and others have demonstrated the critical importance of the individual BIR domains for affecting the antiapoptotic function of the IAPs. We have proposed that antagonists of the IAPs, which may bind to the individual BIR domains, would disrupt the antiapoptotic function of the IAPs. Indeed, individual BIRs serve as critical binding sites for the N-terminal Ser-Gly-Val-Asp, Ser-Gly-Pro-Ile and Ala-Thre-Pro-Ile residues of the Caspases 3, 7, and 9, respectively, and such binding is imperative for the Caspase-inhibitory function of the IAPs. The binding of N-terminal AxPy tetra-peptide residues to XIAP results in the release of the active caspases 3, 7 and 9. In the case of the other IAPs, such as c-IAP1 and c-IAP2, the functions of the BIRs, when ligand-bound, appear to direct the activation of the ubiquitin ligase RING function of the IAPs to a bound target, or individual IAPs themselves, to cause proteosomal loss. In either case, small molecule antagonists of the IAPs should be excellent pro-apoptotic agents, with potential uses in cancer, various proliferative disorders and inflammation.

30
35

A mammalian mitochondrial protein, namely Second Mitochondria-derived Activator of Caspases (SMAC) which antagonizes IAP function, binds mainly to the BIR 3 or 2 sites on respective IAPs via an AxPy amino-terminal tetrapeptide. Four *Drosophila* death-inducing proteins, Reaper, HID, Grim, and Sickie, which antagonize the ability of the *Drosophila* IAPs to inhibit caspases, also bind the BIR domains of the analogous *Drosophila* IAPs via a short AxPy amino-terminal tetrapeptide, a sequence that fits into the BIR binding pocket and disrupts IAP-caspase interactions.

The overall topology of individual BIR domains is highly conserved between the human IAPs and between individual BIR domains of the human IAPs, each BIR being a zinc finger polypeptide domain, locked into a coordinated Zn atom by two cysteines and a histidine residue. The X-ray crystallographic structures of XIAP BIR2 and BIR3 reveal a critical binding pocket for an AXPY motif on the surface of each BIR domain. There are alterations in the intervening amino acid sequences that form the binding pocket and groove in both BIR2 and BIR3. Likewise, we have described homologous domains in the BIRs of other IAPs cIAP1 and cIAP2. This opens the possibility of obtaining various classes of natural and synthetic binding compounds which will have different specificity and binding affinities between each of the BIR domains for each of the IAPs. Discerning the way in which such compounds will affect the biological function of the IAPs in cancer cells vs normal cells is a major new challenge in the discovery of novel mechanism agents to treat cancer and other proliferative disorders where dysregulated IAP function is observed. It is our finding that certain classes of BIR binding compounds may bind to IAP BIRs, with unexpected selectivity and potency, resulting in distinct therapeutic advantages for certain structural classes, potentially resulting from either IAP loss of function or loss of cellular IAP protein, or both.

A number of peptidic AxPy-like and heterocyclic modified AxPy peptidic compounds have been described which activate cellular Caspase 3 by reportedly binding to XIAP BIR3. For a recent reviews, see Elmore et al., Annual Reports in Medicinal Chemistry, **40** (2006) 245-262; Sun et al., Bioorg. Med. Chem. Lett. **15** (2005) 793-797; Oost et al., J. Med. Chem., 2004, **47**(18), 4417-4426; Park et al., Bioorg. Med. Chem. Lett. **15** (2005) 771-775; Franklin et al., Biochemistry, Vol. **42**, No. 27, 2003, 8223-8231; Kip et al., Biochemistry 2002, **41**, 7344-7349; Wu et al., Chemistry and Biology, Vol. **10**, 759-767 (2003); Glover et al., Analytical Biochemistry, **320** (2003) 157-169; United States published patent application number 20020177557; and United States published patent application

number 20040180828; United States published patent application number US2006/0025347A1; United States published patent application number US2005/0197403A1; and United States published patent application number US2006/0194741A1.

5

The aforesaid compounds have been shown to target an isolated BIR3 domain of XIAP via displacement of a fluorescently-labeled probe and they appear to induce an apoptotic event in a select set of cancer cell lines with potency in the low micromolar-nanomolar range. These compounds displayed poor in-vivo activity, likely due to limited bioavailability and may therefore have limited therapeutic application.

10

Thus, IAP BIR domains represent an attractive target for the discovery and development of novel therapeutic agents, especially for the treatment of proliferative disorders such as cancer.

15

SUMMARY OF THE INVENTION



The inventors have previously disclosed a series of compounds which bind to the BIR units of the IAPs and induce apoptosis in various cancer cell lines (US published patent application number 20060264379). A characteristic of these compounds is the presence of a central pyrrolidine unit. We now herein disclose that the linkage of two BIR binding units via a substituted pyrrolidine, with preference for the site, orientation and chemical nature of the linkage, provides novel and distinctly advantageous classes of compounds with up to 1000 fold increase in potency, resulting from induction of apoptosis, against various cancer cell lines, over their corresponding non-bridged BIR binding compounds and that these compounds display the requisite potency, stability and pharmaceutical properties for the treatment of human cancers. Advantageously, the chemical nature of the bridging group can be chosen to cause the translation of the high intrinsic (sub-nanomolar) cellular potency to microgram/kg potency in inhibiting and/or suppressing IAPs in several *in-vivo* xenograft models of human cancers. Furthermore, the compounds described have pharmaceutically acceptable stability in a range of mammalian tissues and fluids and have pharmaceutical properties which ensure adequate solubility and bioavailability using various routes of administration, suitable for clinical use. Such administration results in sustained in vivo effects in mammals as measured in normal and tumor tissues.

20

25

30

In one embodiment of the present invention, there is provided an isomer, enantiomer, diastereoisomer or tautomer of a compound represented by Formula I or II:



wherein

m is 0, 1 or 2;

10 Y is NH, O or S;

BG is

1) $-X-L-X^1-$;

15 X and X^1 are independently selected from

1) O,

2) NR^{13} ,


3) S,

4) $-C_1-C_6$ alkyl-,

20 5) $-C_1-C_6$ alkyl-O-,

6) $-C_1-C_6$ alkyl- NR^{13} -,



7) $-C_1-C_6$ alkyl-S-,



10 L is selected from:

- 1) -C₁-C₂₀ alkyl-,
- 2) -C₂-C₆ alkenyl-,
- 3) -C₂-C₄ alkynyl-,
- 4) -C₃-C₇ cycloalkyl-,
- 5) -aryl-,
- 6) -biphenyl-,
- 7) - heteroaryl-,
- 8) - heterocyclyl-,
- 9) -C₁-C₆ alkyl-(C₂-C₆ alkenyl)-C₁-C₆ alkyl-,
- 10) -C₁-C₆ alkyl-(C₂-C₄ alkynyl)-C₁-C₆ alkyl-,
- 11) -C₁-C₆ alkyl-(C₃-C₇ cycloalkyl)-C₁-C₆ alkyl-,
- 12) -C₁-C₆ alkyl- aryl-C₁-C₆ alkyl-,

- 13) $-C_1-C_6$ alkyl-biphenyl- $-C_1-C_6$ alkyl-,
 14) $-C_1-C_6$ alkyl-heteroaryl- $-C_1-C_6$ alkyl-,
 15) $-C_1-C_6$ alkyl-heterocycyl- $-C_1-C_6$ alkyl-,
 16) $-C_1-C_6$ alkyl-Y- $-C_1-C_6$ alkyl-,
 5 17) -aryl-Y-aryl-,
 18) -heteroaryl-Y-heteroaryl-,
 19) -heterocyclyl-Y-heterocyclyl-,



- 10 wherein the alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more R^6 substituents, and the aryl, biphenyl, heteroaryl, and heterocyclyl are optionally substituted with one or more R^{10} substituents;

Q and Q^1 are independently selected from

- 15 1) NR^4R^5 ,
 2) OR^{11} , or
 3) $S(O)_mR^{11}$; or

Q and Q^1 are independently selected from

- 20 1) aryl, or
 2) heteroaryl, the aryl and the heteroaryl being optionally substituted with one or more R^{10} substituents;

A and A^1 are independently selected from

- 25 1) $-CH_2-$,
 2) $-CH_2CH_2-$,
 3) $-CH(C_1-C_6 \text{ alkyl})-$,
 4) $-CH(C_3-C_7 \text{ cycloalkyl})-$,
 5) $-C_3-C_7 \text{ cycloalkyl}-$,
 30 6) $-CH(C_1-C_6 \text{ alkyl}-C_3-C_7 \text{ cycloalkyl})-$,
 7) $-C(O)-$, or

8) $-\text{C}(\text{O})\text{OR}^{13}$;

R^1 and R^{100} are independently selected from

- 1) H, or
- 5 2) $\text{C}_1\text{--C}_6$ alkyl optionally substituted with one or more R^6 substituents;

R^2 and R^{200} are independently selected from

- 1) H, or
- 10 2) $\text{C}_1\text{--C}_6$ alkyl optionally substituted with one or more R^6 substituents;

R^3 and R^{300} are independently $\text{C}_1\text{--C}_6$ alkyl optionally substituted with one or more R^6 substituents;

R^4 and R^5 are each independently selected from

- 15 1) H,
- 2) haloalkyl,
- 3) $\leftarrow\text{C}_1\text{--C}_6$ alkyl,
- 4) $\leftarrow\text{C}_2\text{--C}_6$ alkenyl,
- 5) $\leftarrow\text{C}_2\text{--C}_4$ alkynyl,
- 20 6) $\leftarrow\text{C}_3\text{--C}_7$ cycloalkyl,
- 7) $\leftarrow\text{C}_3\text{--C}_7$ cycloalkenyl,
- 8) $\leftarrow\text{aryl}$,
- 9) $\leftarrow\text{heteroaryl}$,
- 10) $\leftarrow\text{heterocyclyl}$,
- 25 11) $\leftarrow\text{heterobicycyl}$,
- 12) $\leftarrow\text{C}(\text{O})\text{--R}^{11}$,
- 13) $\leftarrow\text{C}(\text{O})\text{O--R}^{11}$,
- 14) $\leftarrow\text{C}(=\text{Y})\text{NR}^8\text{R}^9$, or
- 15) $\leftarrow\text{S}(\text{O})_2\text{--R}^{11}$,

30 wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicycyl is optionally substituted with one or more R^{10} substituents;

R^6 is

- 35 1) halogen,

- 2) NO₂,
- 3) CN,
- 4) haloalkyl,
- 5) C₁-C₆ alkyl,
- 5 6) C₂-C₆ alkenyl,
- 7) C₂-C₄ alkynyl,
- 8) C₃-C₇ cycloalkyl,
- 9) C₃-C₇ cycloalkenyl,
- 10) aryl,
- 10 11) heteroaryl,
- 12) heterocyclyl,
- 13) heterobicyclyl,
- 14) OR⁷,
- 15) S(O)_mR⁷,
- 15 16) NR⁸R⁹,
- 17) NR⁸S(O)₂R¹¹,
- 18) COR⁷,
- 19) C(O)OR⁷,
- 20) CONR⁸R⁹,
- 20 21) S(O)₂NR⁸R⁹,
- 22) OC(O)R⁷,
- 23) OC(O)Y-R¹¹,
- 24) SC(O)R⁷, or
- 25) NC(Y)NR⁸R⁹,
- 25 wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

R⁷ is

- 1) H,
- 30 2) haloalkyl,
- 3) C₁-C₆ alkyl,
- 4) C₂-C₆ alkenyl,
- 5) C₂-C₄ alkynyl,
- 6) C₃-C₇ cycloalkyl,
- 35 7) C₃-C₇ cycloalkenyl,

- 8) aryl,
- 9) heteroaryl,
- 10) heterocyclyl,
- 11) heterobicyclyl,
- 5 12) $R^8R^9NC(=Y)$, or
- 13) C_1-C_6 alkyl- C_2-C_4 alkenyl, or
- 14) C_1-C_6 alkyl- C_2-C_4 alkynyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 10 heterobicyclyl is optionally substituted with one or more R^{10} substituents;

R^8 and R^9 are each independently

- 1) H,
- 2) haloalkyl,
- 15 3) C_1-C_6 alkyl,
- 4) C_2-C_6 alkenyl,
- 5) C_2-C_4 alkynyl,
- 6) C_3-C_7 cycloalkyl,
- 7) C_3-C_7 cycloalkenyl,
- 20 8) aryl,
- 9) heteroaryl,
- 10) heterocyclyl,
- 11) heterobicyclyl,
- 12) $C(O)R^{11}$,
- 25 13) $C(O)Y-R^{11}$, or
- 14) $S(O)_2-R^{11}$,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 30 heterobicyclyl is optionally substituted with one or more R^{10} substituents;

or R^8 and R^9 together with the nitrogen atom to which they are bonded form a five, six or seven membered heterocyclic ring optionally substituted with one or more R^6 substituents;

R^{10} is

- 35 1) halogen,

- 2) NO₂,
- 3) CN,
- 4) B(OR¹³)(OR¹⁴),
- 5) C₁-C₆ alkyl,
- 5 6) C₂-C₆ alkenyl,
- 7) C₂-C₄ alkynyl,
- 8) C₃-C₇ cycloalkyl,
- 9) C₃-C₇ cycloalkenyl,
- 10) haloalkyl,
- 10 11) OR⁷,
- 12) NR⁸R⁹,
- 13) SR⁷,
- 14) COR⁷,
- 15) C(O)OR⁷,
- 15 16) S(O)_mR⁷,
- 17) CONR⁸R⁹,
- 18) S(O)₂NR⁸R⁹,
- 19) aryl,
- 20) heteroaryl,
- 20 21) heterocyclyl, or
- 22) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, and cycloalkenyl is optionally substituted with one or more R⁶ substituents;

25 R¹¹ is

- 1) haloalkyl,
- 2) C₁-C₆ alkyl,
- 3) C₂-C₆ alkenyl,
- 4) C₂-C₄ alkynyl,
- 30 5) C₃-C₇ cycloalkyl,
- 6) C₃-C₇ cycloalkenyl,
- 7) aryl,
- 8) heteroaryl,
- 9) heterocyclyl, or
- 35 10) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

5 R^{12} is

- 1) haloalkyl,
- 2) C_1-C_6 alkyl,
- 3) C_2-C_6 alkenyl,
- 4) C_2-C_4 alkynyl,
- 10 5) C_3-C_7 cycloalkyl,
- 6) C_3-C_7 cycloalkenyl,
- 7) aryl,
- 8) heteroaryl,
- 9) heterocyclyl,
- 15 10) heterobicyclyl,
- 11) $C(O)-R^{11}$,
- 12) $C(O)O-R^{11}$,
- 13) $C(O)NR^8R^9$,
- 14) $S(O)_m-R^{11}$, or
- 20 15) $C(=Y)NR^8R^9$,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

25 R^{13} and R^{14} are each independently

- 1) H, or
- 2) C_1-C_6 alkyl; or


R^{13} and R^{14} are combined to form a heterocyclic ring or a heterobicyclyl ring;

30 R^{20} is

- 1) H,
- 2) NH_2 , or
- 3) $NHFmoc$;

or a prodrug; or the compound of Formula I or II is labeled with a detectable label or an affinity tag.


In another aspect of the present invention, there is provided an intermediate compound
5 represented by Formula 1-iv:



wherein PG³, R¹, R², R³, R⁴, and R⁵ are as defined herein.

10


In another aspect of the present invention, there is provided an intermediate compound
represented by Formula 2-iv:



wherein PG⁴, R¹, R², R³, R⁴, and R⁵ are as defined herein.

15


In another aspect of the present invention, there is provided an intermediate compound
represented by Formula 3-ii:



wherein PG^4 , R^1 , R^2 , R^3 , A, and Q are as defined herein, and L is $-(CH_2)_r$, $-(CH_2)_r$ -Y- $(CH_2)_r$, -alkyl-aryl-alkyl-, -alkyl-heteroaryl-alkyl-, cycloalkyl, aryl or heteroaryl.

5


In another aspect of the present invention, there is provided an intermediate compound represented by Formula 4(i):



wherein PG^4 , PG^{400} , L, R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A, A', Q and Q' are as defined herein.


10

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 6-ii:




wherein PG^4 , PG^{400} , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound
5 represented by Formula 7-v:




wherein PG^4 , PG^{400} , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound
10 represented by Formula 8-ii:



wherein PG^4 , r , L , R^{100} , R^{200} , A^1 , and Q^1 are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 8-iii:




8-iii

5

wherein PG^4 , r , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , A , A^1 , Q and Q^1 are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 17-i:




17-i

10


wherein PG^4 , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 18-i:



wherein PG⁴, L, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, A, A¹, Q and Q¹ are as defined herein.


In another aspect of the present invention, there is provided an intermediate compound
 5 represented by Formula 19-2:



wherein PG¹, PG², L, X are as defined herein.


10

In another aspect of the present invention, there is provided an intermediate compound
 represented by Formula 19-3:



wherein PG², L, and X, are as defined herein.


- 5 In another aspect of the present invention, there is provided an intermediate compound represented by Formula 19-8:



wherein PG⁴, L, X, X¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, R⁴, R⁴⁰⁰, R⁵ and R⁵⁰⁰ are as defined herein.


10

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 20-1a:




wherein PG¹, L, X, X¹, R⁴ and R⁵ are as defined herein.

- 5 In another aspect of the present invention, there is provided an intermediate compound represented by Formula 20-2:




- 10 wherein PG⁴, L, X, and X¹, are as defined herein.

In another aspect of the present invention, there is provided an intermediate compound represented by Formula 20-4:



wherein PG^4 , L , X , X^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , R^4 , R^{400} , R^5 , and R^{500} are as defined herein.


- 5 In another aspect of the present invention, there is provided a process for producing compounds represented by Formula I, described hereinabove, the process comprising:
a) coupling two intermediates represented by Formula 1-iv:



- 10 wherein PG^3 , R^1 , R^2 , R^3 , A , and Q are as defined herein,
in a solvent; and
b) removing the protecting groups so as to form compounds of Formula 1.

- In another aspect of the present invention, there is provided a process for producing
15 compounds represented by Formula I, described hereinabove, the process comprising:

- a) coupling an intermediate represented by Formula 2-iv:



wherein PG⁴ is a protecting group, and R¹, R², R³, A, and Q are as defined herein.

and an activated diacid (0.5 equiv) in a solvent; and

b) removing the protecting groups so as to form compounds of Formula I.

5

In another aspect of the present invention, there is provided a method for the preparation of a pharmaceutically acceptable salt of compound of formula I and II, by the treatment of a compound of formula I or II with 1 to 2 equiv of a pharmaceutically acceptable acid, as defined herein.

10

In another aspect of the present invention, there is provided a pharmaceutical composition comprising a compound, as described above, mixed with a pharmaceutically acceptable carrier, diluent or excipient.

15

In another aspect of the present invention, there is provided a pharmaceutical composition adapted for administration as an agent for treating a proliferative disorder in a subject, comprising a therapeutically effective amount of a compound, as described above.

20

In another aspect of the present invention, there is provided a pharmaceutical composition comprising a compound of Formula I in combination with one or more death receptor agonists, for example, an agonist of TRAIL receptor.

25

In another aspect of the present invention, there is provided a pharmaceutical composition comprising a compound of formula I in combination with any therapeutic agent that increases the response of one or more death receptor agonists, for example cytotoxic cytokines such as interferons.

30

In another aspect of the present invention, there is provided a method of preparing a pharmaceutical composition, the method comprising: mixing a compound, as described above, with a pharmaceutically acceptable carrier, diluent or excipient.

35

In another aspect of the present invention, there is provided a method of treating a disease state characterized by insufficient apoptosis, the method comprising: administering to a subject in need thereof, a therapeutically effective amount of a pharmaceutical composition, as described above, so as to treat the disease state.

In another aspect of the present invention, there is provided a method of modulating IAP function, the method comprising: contacting a cell with a compound of the present invention so as to prevent binding of a BIR binding protein to an IAP BIR domain thereby
5 modulating the IAP function.

In another aspect of the present invention, there is provided a method of treating a proliferative disease, the method comprising: administering to a subject in need thereof, a therapeutically effective amount of the pharmaceutical composition, as described above,
10 so as to treat the proliferative disease.

In another aspect of the present invention, there is provided a method of treating cancer, the method comprising: administering to a subject in need thereof, a therapeutically effective amount of the pharmaceutical composition, as described above, so as to treat
15 the cancer.

In another aspect of the present invention, there is provided a method of treating cancer, the method comprising: administering to the subject in need thereof, a therapeutically effective amount of a pharmaceutical composition, as described above, in combination or
20 sequentially with an agent selected from:

- a) an estrogen receptor modulator,
- b) an androgen receptor modulator,
- c) retinoid receptor modulator,
- d) a cytotoxic agent,
- 25 e) an antiproliferative agent,
- f) a prenyl-protein transferase inhibitor,
- g) an HMG-CoA reductase inhibitor,
- h) an HIV protease inhibitor,
- i) a reverse transcriptase inhibitor,
- 30 k) an angiogenesis inhibitor,
- l) a PPAR- γ agonist,
- m) a PPAR- δ agonist,
- n) an inhibitor of inherent multidrug resistance,
- o) an anti-emetic agent,
- 35 p) an agent useful in the treatment of anemia,

q) agents useful in the treatment of neutropenia,

r) an immunologic-enhancing drug.

s) a proteasome inhibitor;

t) an HDAC inhibitor;

5 u) an inhibitor of the chemotrypsin-like activity in the proteasome; or

v) E3 ligase inhibitors;

w) a modulator of the immune system such as, but not limited to, interferon-alpha, Bacillus Calmette-Guerin (BCG), and ionizing radiation (UVB) that can induce the release of cytokines, such as the interleukins, TNF, or induce release of death receptor ligands such

10 as TRAIL;

x) a modulator of death receptors TRAIL and TRAIL agonists such as the humanized antibodies HGS-ETR1 and HGS-ETR2;

or in combination or sequentially with radiation therapy, so as to treat the cancer.

15 In another aspect of the present invention, there is provided a method for the treatment or prevention of a proliferative disorder in a subject, the method comprising: administering to the subject a therapeutically effective amount of the composition, described above.

In another aspect of the present invention, the method further comprises administering to
20 the subject a therapeutically effective amount of a chemotherapeutic agent prior to, simultaneously with or after administration of the composition.

In yet another aspect, the method further comprises administering to the subject a therapeutically effective amount of a death receptor agonist prior to, simultaneously with
25 or after administration of the composition. The death receptor agonist is TRAIL or the death receptor agonist is a TRAIL antibody. The death receptor agonist is typically administered in an amount that produces a synergistic effect.

In yet another aspect, there is provided use of the compound as described above for the
30 manufacture of a medicament for treating or preventing a disease state characterized by insufficient apoptosis.

In yet another aspect, there is provided use of the compound as described above for the
35 manufacture of a medicament for treating or preventing a proliferative disorder.

In yet another aspect, there is provided use of the compound as described above in combination with an agent for the manufacture of a medicament for treating or preventing a proliferative disorder, wherein the agent is selected from:

- a) an estrogen receptor modulator,
- 5 b) an androgen receptor modulator,
- c) retinoid receptor modulator,
- d) a cytotoxic agent,
- e) an antiproliferative agent,
- f) a prenyl-protein transferase inhibitor,
- 10 g) an HMG-CoA reductase inhibitor,
- h) an HIV protease inhibitor,
- i) a reverse transcriptase inhibitor,
- k) an angiogenesis inhibitor,
- l) a PPAR- γ agonist,
- 15 m) a PPAR- δ agonist,
- n) an inhibitor of inherent multidrug resistance,
- o) an anti-emetic agent,
- p) an agent useful in the treatment of anemia,
- q) agents useful in the treatment of neutropenia,
- 20 r) an immunologic-enhancing drug.
- s) a proteasome inhibitor;
- t) an HDAC inhibitor;
- u) an inhibitor of the chemotrypsin-like activity in the proteasome; or
- v) E3 ligase inhibitors;
- 25 w) a modulator of the immune system such as, but not limited to, interferon-alpha, Bacillus Calmette-Guerin (BCG), and ionizing radiation (UVB) that can induce the release of cytokines, such as the interleukins, TNF, or induce release of death receptor ligands such as TRAIL;
- x) a modulator of death receptors TRAIL and TRAIL agonists such as the humanized
- 30 antibodies HGS-ETR1 and HGS-ETR2;
- or in combination or sequentially with radiation therapy.

In yet another aspect, there is provided use of the compound as described above in combination with a death receptor agonist for the manufacture of a medicament the

- 35 treatment or prevention of a proliferative disorder in a subject.

In yet another aspect, there is provided a pharmaceutical composition comprising the compound as described above, mixed with a pharmaceutically acceptable carrier, diluent or excipient, for treating or preventing a disease state characterized by insufficient apoptosis.

5

In yet another aspect, there is provided a pharmaceutical composition comprising the compound as described above in combination with any compound that increases the circulating level of one or more death receptor agonists for preventing or treating a proliferative disorder.

10

In yet another aspect, there is provided a method of preparing a pharmaceutical composition, the method comprising: mixing the compound as described above, with a pharmaceutically acceptable carrier, diluent or excipient.

15

In another aspect of the present invention, there is provided a probe, the probe being a compound of Formula I or II above, the compound being labeled with a detectable label or an affinity tag.

20

In another aspect of the present invention, there is provided a method of identifying compounds that bind to an IAP BIR domain, the assay comprising:

- a) contacting an IAP BIR domain with a probe to form a probe:BIR domain complex, the probe being displaceable by a test compound;
 - b) measuring a signal from the probe so as to establish a reference level;
 - c) incubating the probe:BIR domain complex with the test compound;
 - 25 d) measuring the signal from the probe;
 - e) comparing the signal from step d) with the reference level, a modulation of the signal being an indication that the test compound binds to the BIR domain,
- wherein the probe is a compound of Formula I or II labeled with a detectable label or an affinity label.

30

In another aspect of the present invention, there is provided a method of detecting loss of function or suppression of IAPs in vivo, the method comprising: a) administering to a subject, a therapeutically effective amount of a pharmaceutical composition, as defined above; b) isolating a tissue sample from the subject; and c) detecting a loss of function or suppression of IAPs from the sample.

35

BRIEF DESCRIPTION OF THE FIGURES

Further aspects and advantages of the present invention will become better understood
5 with reference to the description in association with the following Figures, wherein:

FIGURE 1 depicts SKOV-3 Human Ovarian Cancer Cell Line Xenograph Study with compound 3. Female CD-1 nude mice (approximately 20-25 g) were subcutaneously
10 injected 5×10^6 SKOV-3 human ovarian tumor cells in 50% matrigel subcutaneously in the right flank. On day 55, when tumors were approximately 100 mm^3 , treatment was initiated with compound 3 treating with compound for 5 consecutive days followed by 2 days with no drug treatment for the duration of the experiment. Tumor size was measured with digital calipers and calculated as $V = (a \times b^2)/2$, wherein, a is the longest dimension and b
15 is the width. Tumor regression was observed at 1 mg/kg while tumor stasis was observed to 0.3 mg/kg.

FIGURE 2 depicts MDA-MB-231 Human Mammary Cancer Cell Line Xenograph Study with compound 3. Female CD-1 nude mice (approximately 20-25 g) were subcutaneously
20 injected 1×10^6 MDA-MB-231 human mammary tumor cells in the right flank. On day 71, when tumors were approximately 90 mm^3 , treatment was initiated with compound 3 treating with compound for 5 consecutive days followed by 2 days with no drug treatment for the duration of the experiment. Tumor size was measured with digital calipers and calculated as $V = (a \times b^2)/2$, wherein, a is the longest dimension and b is the width. Tumor
25 regression was observed at 1 mg/kg.

FIGURE 3 demonstrates that compound 3 induces a loss of cIAP-1 in HCT-116 cells in vitro. PC3, SKOV3, MDA-MB-231, HCT-116 cells were treated with various concentrations of compound 3 and incubated at 37°C for 5 hours. Cells were collected
30 and the level of cIAP-1 and actin (loading control-not shown) were detected by western blot. Results indicated that compound 3 induced cIAP-1 loss in human cancer cells in a time dependent manner (not shown). Using a similar method as described in Figure 3, compound 3 induced a loss of loss of c-IAP1 from ES2 and 4T1 cell lines (not shown) and a loss of cIAP2 from PC3 cells (not shown).

35

FIGURE 4 demonstrates the in vitro modulation of IAP's in mice white cells. CD-1 mice whole blood was incubated in vitro with various concentrations of compound 3 for 3 hours. White blood cells were isolated from the treated whole blood via Ficoll gradient. Protein was isolated from the white blood cells and the relative amount of cIAP-1 and tubulin (loading control) was revealed by western blotting. In vitro results indicated that compound 3 induces cIAP-1 loss in mice blood.

FIGURE 5 demonstrates in vivo modulation of cIAP-1 in mice white cells. Compound 3 was administered to CD-1 mice by i.v. bolus injection as the indicated dose. After 1 to 48 hours the animals were sacrificed, the blood collected, the white cells were isolated on Ficoll gradient and the protein extracted. The relative amount of cIAP-1 and tubulin (loading control) was revealed by western blotting (shown below at 3 hr time point). Results indicated that compound 3 induces cIAP-1 down regulation in mice white cells, using ex-vivo detection methods.

FIGURE 6 depicts a MDA-MB-231 Human Breast Cancer Cell Line Xenograft Study with compound 24 (1 mg/kg). Female CD-1 nude mice (approximately 20-25 g) were subcutaneously injected with 1×10^6 MDA-MB-231 human breast tumor cells in 50% matrigel subcutaneously in the right flank. On day 55, when tumors were approximately 100 mm³, treatment was initiated with compound 24 treating with compound for 5 consecutive days followed by 2 days with no drug treatment for the duration of the experiment. Tumor size was measured with digital calipers and calculated as $V = (a \times b^2)/2$, wherein, a is the longest dimension and b is the width. Circles – 20 % HPCD control; diamonds – compound 24.

Figure 7 demonstrates in vivo modulation of cIAP-1 in rat white cells. Compound 24 was administered to rats by i.v. bolus injection. After 1 to 48 hours, the blood was collected, the white cells were isolated on Ficoll gradient and the protein extracted. The relative amount of cIAP-1 and tubulin (loading control) was revealed by western blotting (shown below at 3 hr time point). Results indicated that compound 24 induces cIAP-1 loss in mice white cells, using ex-vivo detection methods.

DETAILED DESCRIPTION OF THE INVENTION


In many cancer and other diseases, an up-regulation of IAPs induced by gene defects or by chemotherapeutic agents has been correlated to an increased resistance to apoptosis. Conversely, our results show that cells decreased in IAP levels are more sensitive to chemotherapeutic agents and to death receptor agonists such as TRAIL. It is believed that a small molecule, which will antagonize IAP function, or a loss of IAPs from diseased cells, will be useful as a therapeutic agent. We report herein that compounds of the instant invention can directly bind to IAPs, cause a down regulation of IAP proteins in cells, and induce apoptosis in cancer cells. Furthermore, compounds of the instant invention have demonstrated synergistic effects in combination with clinically relevant agents used in the treatment of cancer.

We have discovered a novel series of bridged compounds, which bind to the intact cellular IAPs and results in profound, sustained IAP protein down-modulation and enhanced cellular apoptosis of cancer cells through enhanced release of active Caspase 3. This biological response has been observed in various cell lines derived from human breast, pancreatic, colon, lung, ovarian cancers and primary human leukemia and lymphoma cells. The compounds were found to be highly synergistic with Death Receptor Agonist-mediated killing, such as TRAIL, TRAIL Receptor Monoclonal Antibodies and TNF- α , in a large and comprehensive range of cancer cells. Based upon these findings, the compounds will find application in treatment of many cancer types, such as solid tumors and tumors originating from the hematopoietic system. Moreover, the compounds of the present invention may also find application in preventing metastatic cancer cell invasion, inflammation, and in other diseases characterized by cells that are resistant to apoptosis via upregulation of any one of the IAPs. As shown in Figure 3, compound 3 is capable of inducing the complete loss of c-IAP1/2 proteins from multiple tumor cell lines at concentrations of less than 10 nM. Other compounds of the instant invention were shown to display a similar ability to induce time dependent IAP loss from cancer cells. This loss in IAP proteins strongly correlate to the ED₅₀ in SKOV3 cells.

The 'bridging' of two IAP BIR binding units, M1 and M2, described in more detail below, using an appropriate 'bridging unit', linked to one of the pyrrolidine rings, provides bridged IAP BIR binding compounds, which demonstrate significantly increased anti-cancer activity (10-1000 fold), as compared to their monomeric units. This improved activity results from an improved ability to bind to the BIR domains of the intact IAPs, and results in the induction of apoptosis in various cancer cell lines.


Various factors influence the in vitro proapoptotic character of the compounds of the present invention. Specifically, these include i) the point of attachment of the linker/pyrrolidine bond, ii) the stereochemistry at the linker/pyrrolidine bond, iii) the linker
 5 moieties themselves, including stereochemistry, regiochemistry, and the rigidity of the linker system, iv) alkyl substitution at R^1 and R^{100} , and v) the substitution pattern at R^4 , R^{400} , R^5 , and R^{500} .


For ease of description, throughout the description, the compounds of Formula I and
 10 Formula II, may also include the use of the terms P1, P2, P3, P4 and P5. These terms refer to the amino acids or modified amino acids within either of Formula I or II. The following illustrates the use of the terms:



15 wherein the waved line represents a covalent bond to another BIR binding unit.

The compounds of the present invention may also be represented by Formula 3 or Formula 4 in which M1 and M2 represent independent BIR binding domains.





wherein R^1 , R^2 , R^{100} , R^{200} , R^3 , R^{300} , R^{20} , A , A^1 , Q , Q^1 , and BG as defined herein, and the dotted line represents a hypothetical dividing line for comparing the substituents associated with $M1$ and $M2$.

In one subset of Formula 3, $M1$ is the same as $M2$ and the dotted line denotes a line of symmetry. In another subset, $M1$ is different from $M2$.

- 10 In one subset, compounds of Formula 4 are asymmetrical about the dotted line. In another subset the substituents on $M1$ and $M2$ are the same. In another subset, the substituents on $M1$ and $M2$ are different.

One skilled in the art will recognize that when $M1$ and $M2$ are the same, the R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{13} , R^{14} , r , m , Y , A , Q , and X substituents in $M1$ have the same meaning as the R^{100} , R^{200} , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{13} , R^{14} , r , m , Y , A^1 , Q^1 , and X^1 substituents respectively in $M2$. When $M1$ and $M2$ are different, at least one R^1 , R^2 , R^{100} , R^{200} , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{13} , R^{14} , r , m , Y , A , A^1 , Q , Q^1 , X , and X^1 substituent is different in either of $M1$ or $M2$.

Alternatively the substituents in $M1$ can be defined as R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{13} , R^{14} , r , m , p , Y , A , Q , and X , and those in $M2$ can be defined as R^{100} , R^{200} , R^{400} , R^{500} , R^{600} , R^{700} , R^{800} , R^{900} , R^{1000} , R^{1100} , R^{1300} , R^{1400} , r^1 , m^1 , p^1 , Y^1 , A^1 , Q^1 and X^1 respectively. In the case where $M1$ and $M2$ are the same, the R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{13} , R^{14} , r , m , Y , A , Q , and X substituents in $M1$ have the same meanings as R^{100} , R^{200} , R^{400} , R^{500} , R^{600} , R^{700} , R^{800} , R^{900} , R^{1000} , R^{1100} , R^{1300} , R^{1400} , r^1 , m^1 , Y^1 , A^1 , Q^1 and X^1 respectively in $M2$. In the case where $M1$ and $M2$ are different, at least one of the aforesaid substituents is different.

The compounds of the present invention are useful as BIR domain binding compounds in mammalian IAPs and are represented by either Formula I or Formula II. The following are embodiments, groups and substituents of the compounds according to Formula I and

5 Formula II, which are described hereinafter in detail.

A and A¹:

In one subset of compounds of Formula I or II, A and A¹ are both CH₂.

10 In an alternative subset of compounds of Formula I or II, A and A¹ are both C=O.

In another alternative subset of compounds of Formula I or II, A is CH₂ and A¹ is C=O.




In another alternative subset of compounds of Formula I or II, A and A¹ are both
15 C(O)OCH₃.

In another alternative subset of compounds of Formula I or II, A and A¹ are both C(O)OH.

20 Any and each individual definition of A and A¹ as set out herein may be combined with any and each individual definition of Core, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, Q, Q¹, and BG as set out herein.

Core:

25 Therefore, for compounds of Formula I, the present invention comprises compounds of Formula 1A through 1C:





wherein BG, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, and R³⁰⁰ are as defined hereinabove and hereinafter.

In one example, the present invention comprises compounds of Formula 1A.

In an alternative example, the present invention comprises compounds of Formula 1B.

In another alternative example, the present invention comprises compounds of Formula 1C.

Alternatively, compounds of Formula II comprise compounds of Formula 2A and 2B:



wherein BG, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰ and R²⁰ are as defined hereinabove and hereinafter.


In one example, the present invention comprises compounds of Formula 2A.

Any and each individual definition of Core as set out herein may be combined with any and each individual definition of A, A¹, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, R²⁰, Q, Q¹, and BG as set out herein.



BG:

In one subset of the aforesaid compounds, BG is -X-L-X¹-.

In one subset, for compounds of Formula I in which BG is $-X-L-X'$, the invention comprises compounds of Formula 1a through 1c:





5




wherein L, X, X¹, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, and R³⁰⁰ are as defined hereinabove and hereinafter.

One further subset of the aforesaid compounds comprises compounds of Formula 1.1a

5 through 1.1c:






wherein L, X, X¹, A, A¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, R⁴, R⁴⁰⁰, R⁵ and R⁵⁰⁰ are as defined hereinabove and hereinafter.

5

In one subset, for compounds of Formula II in which BG is -X-L-X¹-, the invention comprises compounds of Formula 2a:








10 wherein L, X, X¹, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³ and R²⁰ are as defined hereinabove and hereinafter.

Any and each individual definition of BG as set out herein may be combined with any and each individual definition of Core, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, A, A¹, Q, and Q¹ as set out
 15 herein.

X and X¹:



In one subset of the aforesaid compounds, X and X¹ are independently selected from

- 1) O,
- 2) NR¹³,
- 3) S,
- 4) C₁-C₆ alkyl-O-,
- 5) C₁-C₆ alkyl,



In one example X and X¹ are independently selected from:

- 1) O,





Any and each individual definition of X and X¹ as set out herein may be combined with any and each individual definition of Core, L, A, A¹, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, R²⁰, Q, Q¹, and BG as set out herein.

L:


In one subset of the aforesaid compounds, L is selected from:


- 1) $-C_1-C_{20}$ alkyl-,
- 2) $-C_3-C_7$ cycloalkyl-,
- 3) -aryl-,
- 4) -biphenyl-,
- 5) - heteroaryl-,
- 6) $-C_1-C_6$ alkyl- $(C_2-C_4$ alkynyl)- C_1-C_6 alkyl-,
- 7) $-C_1-C_6$ alkyl- aryl- C_1-C_6 alkyl-,
- 8) -aryl-Y-aryl-,



wherein the alkyl and cycloalkyl are optionally substituted with one or more R^6 substituents, and the aryl, biphenyl and heteroaryl are optionally substituted with one or more R^{10} substituents.

Typical examples of L include





Any and each individual definition of L as set out herein may be combined with any and
 5 each individual definition of Core, A, A¹, r, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, R²⁰, X, X¹, Q, or Q¹,
 as set out herein.


r:

In the aforesaid aspect, r is an integer of 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10.





10






Any and each individual definition of r as set out herein may be combined with any and
 each individual definition of Core, A, L, A¹, R¹, R², R¹⁰⁰, R²⁰⁰, R³, R³⁰⁰, R²⁰, Q, Q¹, X and
 X¹ as set out herein.






15 More explicitly, the invention comprises compounds of Formula 1.1 through 1.18:






1.1











wherein r, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³ and R³⁰⁰ are as defined hereinabove.

10 Alternatively more explicitly, the invention comprises compounds of Formula 2.1 and 2.2:



5 wherein A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³ and R²⁰ are as defined hereinabove.

R¹ and R¹⁰⁰:

In one subset of the aforesaid compounds R¹ and R¹⁰⁰ are both H.

10 In one subset of the aforesaid compounds R¹ and R¹⁰⁰ are both C₁-C₆ alkyl.

In one example, R¹ and R¹⁰⁰ are both CH₃.

Any and each individual definition of R^1 and R^{100} as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^2 , R^{200} , R^3 , R^{300} , Q, Q^1 , B, B^1 , and BG as set out herein.

5 **R^2 and R^{200} :**

In one subset of the aforesaid compounds R^2 and R^{200} are both C_1 - C_6 alkyl optionally substituted with OH.

In one example, R^2 and R^{200} are both CH_3 .

In another example, R^2 is CH_2OH and R^{200} is CH_3 .

10 In another example, R^2 and R^{200} are both CH_2OH .

In another example, R^2 and R^{200} are both CH_2CH_3 .

Any and each individual definition of R^2 and R^{200} as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^3 , R^{300} , Q, Q^1 , and BG as set out herein.

R^3 and R^{300} :

In one subset of compounds of Formula I, R^3 and R^{300} are both C_1 - C_6 alkyl.

In one example, R^3 and R^{300} are both $C(CH_3)_3$.

20

In subset of compounds of Formula II, R^3 is C_1 - C_6 alkyl. In one example, R^3 is $C(CH_3)_3$.

Any and each individual definition of R^3 and R^{300} as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , Q, Q^1 , and BG as set out herein.

25

Q and Q^1 :

In one subset of the aforesaid compounds, Q and Q^1 are both NR^4R^5 , wherein R^4 and R^5 are as defined herein.

30

Any and each individual definition of Q and Q^1 as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} and BG as set out herein.

35 **R^4 and R^5 :**

In one subset of the aforesaid compounds in which A and A¹ are both C=O, R⁴ is H and R⁵ is selected from

- 1) ←C₁–C₆ alkyl
- 2) ←C₂–C₆ alkenyl,
- 5 3) ←C₂–C₄ alkynyl,
- 4) ←C₃–C₇ cycloalkyl,
- 5) ←C₃–C₇ cycloalkenyl,
- 6) ←aryl,
- 7) ←heteroaryl,
- 10 8) ←heterocyclyl, or
- 9) ←heterobicyclyl,


wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;


15 wherein R⁶ and R¹⁰ are as defined herein.

In another subset of the above compounds, R⁴ is H and R⁵ is selected from:


- 1) ←C₁–C₆ alkyl, or
- 2) ←aryl,
- 20 wherein the alkyl is optionally substituted with one or two R⁶ substituents; and wherein the aryl is optionally substituted with one R¹⁰ substituent;
- wherein R⁶ and R¹⁰ are as defined herein.

25 Examples of the aforesaid subset include, R⁴ is H and R⁵ is selected from the group








Therefore, when A and A¹ are both C=O, then Q and Q¹ are both





10


In another example, Q and Q¹ are both





In another example, Q is  and Q¹ is .


In another example, Q and Q¹ are both .


In another example, Q and Q¹ are both .


In another example, Q and Q¹ are both .


5 In another example, Q and Q¹ are both .

In another example, Q and Q¹ are both .


In another example, Q and Q¹ are both .

In another example, Q and Q¹ are both .


In another example, Q and Q¹ are both .

10 In another example, Q and Q¹ are both .


In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




5 In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




In another example, Q and Q¹ are both




In another example, Q and Q¹ are both







In another example, Q and Q¹ are both



In another example, Q and Q¹ are both



In another example, Q and Q¹ are both



In another example, Q and Q¹ are both

5

In an alternative subset of the aforesaid compounds in which A and A¹ are both CH₂, then R⁴ and R⁵ are each independently

- 1) haloalkyl,
- 2) ←C₁–C₆ alkyl,
- 10 3) ←C₂–C₆ alkenyl,
- 4) ←C₂–C₄ alkynyl,
- 5) ←C₃–C₇ cycloalkyl,
- 6) ←C₃–C₇ cycloalkenyl,
- 7) ←aryl,
- 15 8) ←heteroaryl,
- 9) ←heterocyclyl,
- 10) ←heterobicyclyl,
- 11) ←C(O)–R¹¹,
- 12) ←C(O)O–R¹¹,
- 20 13) ←C(=Y)NR⁸R⁹, or
- 14) ←S(O)₂–R¹¹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents; wherein Y, R^6 , R^8 , R^9 , R^{10} and R^{11} are as defined herein.

5


In another subset of the above compounds, R^4 and R^5 are independently selected from

- 1) C_1 - C_6 alkyl,
- 2) $\leftarrow C(O)-R^{11}$,
- 3) $\leftarrow C(O)O-R^{11}$, or
- 4) $\leftarrow S(O)_2-R^{11}$,


10


wherein the alkyl is substituted with an R^6 substituent;


wherein R^6 and R^{11} are as defined herein.

In one subset of the aforesaid compounds, R^4 is $S(O)_2CH_3$ and R^5 is  or


15




In another subset of the aforesaid compounds, R^4 is $C(O)CH_3$ and R^5 is  or




In another subset of the aforesaid compounds, R^4 is



and R^5 is



or



20

Any and each individual definition of R^4 and R^5 as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , and BG as set out herein.

25

R^{11} :

In one subset of the aforesaid compounds,

R¹¹ is

1) C₁–C₆ alkyl, or

2) aryl,

- 5 wherein the alkyl is optionally substituted with one or more R⁶ substituents; and wherein the aryl is optionally substituted with one or more R¹⁰ substituents; wherein R⁶ and R¹⁰ are as defined herein.

- 10 In one subset of the aforesaid compounds, R¹¹ is

1) C₁–C₆ alkyl optionally substituted with one or two R⁶ substituents, or

2) phenyl optionally substituted with one R¹⁰ substituent;

wherein the R⁶ and the R¹⁰ substituents are as defined herein.

- 15 Any and each individual definition of R¹¹ as set out herein may be combined with any and each individual definition of Core, A, A¹, R¹, R¹⁰⁰, R², R²⁰⁰, R⁴, R⁵, R³, R³⁰⁰ and BG as set out herein.

R⁶:

- 20 In one subset of the aforesaid compounds, R⁶ is

1) halogen,

2) NO₂,

3) CN,

4) aryl,

- 25 5) heteroaryl,

6) heterocyclyl,

7) heterobicyclyl,

8) OR⁷,

9) SR⁷, or

- 30 10) NR⁸R⁹,

wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

wherein R⁷, R⁸, R⁹ and R¹⁰ are as defined herein.

- 35 In another subset of the aforesaid compounds, R⁶ is

- 1) halogen,
- 2) aryl, or
- 3) NR^8R^9 ,

wherein the aryl is optionally substituted with one R^{10} substituent;

5 wherein R^8 , R^9 and R^{10} are as defined herein.

In one subset of the aforesaid compounds, R^6 is

- 1) halogen,
- 2) phenyl, or
- 10 3) NR^8R^9 ,

wherein the phenyl is optionally substituted with one R^{10} substituent;

wherein R^8 and R^9 are as defined herein.

15 Any and each individual definition of R^6 as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , R^4 , R^5 , R^3 , R^{300} and BG as set out herein.

R^8 and R^9 :

In one subset of the aforesaid compounds, R^8 and R^9 are each independently

- 20 1) H,
- 2) haloalkyl,
- 3) $\text{C}_1\text{--C}_6$ alkyl,
- 4) $\text{C}_2\text{--C}_6$ alkenyl,
- 5) $\text{C}_2\text{--C}_4$ alkynyl,
- 25 6) $\text{C}_3\text{--C}_7$ cycloalkyl, or
- 7) $\text{C}_3\text{--C}_7$ cycloalkenyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents;

wherein the R^6 substituents are as defined herein.

30 In another subset of the aforesaid compounds, R^8 and R^9 are each independently

- 1) H, or
- 2) $\text{C}_1\text{--C}_6$ alkyl,

wherein the alkyl is optionally substituted with an aryl.

Any and each individual definition of R^8 and R^9 as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , R^4 , R^5 , R^3 , R^{300} and BG as set out herein.

5 **R^{10} :**

In one aspect of the aforesaid compounds, R^{10} is

- 1) halogen,
- 2) NO_2 ,
- 3) CN,
- 10 4) haloalkyl,
- 5) OR^7 ,
- 6) NR^8R^9 , or
- 7) SR^7 ;

wherein R^7 , R^8 , and R^9 are as defined herein.


15


In another aspect of the aforesaid compounds, R^{10} is

- 1) halogen, or
- 2) OC_1-C_6 alkyl.

- 20 Any and each individual definition of R^{10} as set out herein may be combined with any and each individual definition of Core, A, A^1 , R^1 , R^{100} , R^2 , R^{200} , R^4 , R^5 , R^3 , R^{300} and BG as set out herein.

- 25 Alternatively, the invention provides an isomer, enantiomer, diastereoisomer or tautomer of a compound represented by Formula I or Formula 2:





wherein

n is 0 or 1;

5 m is 0, 1 or 2;

p is 1 or 2;

Y is NH, O or S;

LG is

10 2) $-X-L-X^1-$;

X and X¹ are independently selected from

1) O,

2) NR¹³,

15 3) S,




4) C₁-C₆ alkyl-O-,



5) C₁-C₆ alkyl-NR¹³-,

6) C₁-C₆ alkyl-S-,

7) C₁-C₆ alkyl-aryl-O-

20





L is selected from:


- 5 1) $-C_1-C_{20}$ alkyl-,
- 2) $-C_2-C_6$ alkenyl-,
- 3) $-C_2-C_4$ alkynyl-,
- 4) $-C_3-C_7$ cycloalkyl-,
- 5) -phenyl-,
- 10 6) -biphenyl-,
- 7) -heteroaryl-,
- 8) -heterocycyl-,
- 9) $-C_1-C_6$ alkyl- $(C_2-C_6$ alkenyl)- C_1-C_6 alkyl-,
- 10) $-C_1-C_6$ alkyl- $(C_2-C_4$ alkynyl)- C_1-C_6 alkyl,
- 15 11) $-C_1-C_6$ alkyl- $(C_3-C_7$ cycloalkyl)- C_1-C_6 alkyl,
- 12) $-C_1-C_6$ alkyl-phenyl- C_1-C_6 alkyl,
- 13) $-C_1-C_6$ alkyl-biphenyl- C_1-C_6 alkyl,
- 14) $-C_1-C_6$ alkyl-heteroaryl- C_1-C_6 alkyl,
- 15) $-C_1-C_6$ alkyl heterocycyl- C_1-C_6 alkyl,
- 20 16) $-C_1-C_6$ alkyl-O- C_1-C_6 alkyl,
- 17) $-C(O)$ -aryl- $C(O)$ -,
- 18) $-C(O)$ -heteroaryl- $C(O)$ -,
- 19) $-C(O)$ -(C_1-C_6 alkyl)-aryl-(C_1-C_6 alkyl)- $C(O)$ -,
- 20) $-C(O)$ -(C_1-C_6 alkyl)-heteroaryl-(C_1-C_6 alkyl)- $C(O)$ -, or
- 25 21) $-C(O)$ -(C_1-C_6 alkyl)- $(C_3-C_7$ cycloalkyl)-(C_1-C_6 alkyl)- $C(O)$ -;

Q and Q¹ are independently selected from

- 1) NR^4R^5 ,
- 2) OR^{11} , or
- 30 3) $S(O)_mR^{11}$; or

Q and Q¹ are independently selected from aryl or heteroaryl, being optionally substituted with R¹² substituents; or

Q and Q¹ are independently



5

wherein G is a 5, 6 or 7 membered ring which optionally incorporates one or more heteroatoms selected from S, N or O, and which is optionally substituted with one or more R¹² substituents, the ring being optionally fused with an aryl or a heteroaryl, the aryl and the heteroaryl being optionally substituted with one or more R¹² substituents;

10

A and A¹ are independently selected from

- 1) -CH₂-,
- 2) -CH₂CH₂-,
- 3) -CH(C₁-C₆ alkyl)-,
- 4) -CH(C₃-C₇ cycloalkyl)-,
- 5) -C₃-C₇ cycloalkyl-,
- 6) -CH(C₁-C₆ alkyl-C₃-C₇ cycloalkyl)-, or
- 7) -C(O)-;

15

20 R¹ and R¹⁰⁰ are independently selected from

- 3) H, or
- 4) C₁-C₆ alkyl optionally substituted with one or more R⁶ substituents;

R² and R²⁰⁰ are independently H or C₁-C₆ alkyl optionally substituted with one or more R⁶ substituents;

25

R⁴ and R⁵ are independently selected from:

- 1) H,
- 2) C₁-C₆ alkyl→,
- 3) C₃-C₇ cycloalkyl→,
- 4) haloalkyl→,

30

- 5) aryl→,
- 6) biphenyl→,
- 7) heteroaryl-aryl→,
- 8) aryl-heteroaryl→,
- 5 9) aryl-heterocyclyl→,
- 10) heterocyclyl→,
- 11) heteroaryl→,
- 12) heterocyclyl→,
- 13) C₁-C₆ alkyl-O_nC(O)→,
- 10 14) haloalkyl-O_nC(O)→,
- 15) C₃-C₇ cycloalkyl-O_nC(O)→,
- 16) aryl-O_nC(O)→,
- 17) heteroaryl-O_nC(O)→,
- 18) heterocyclyl-O_nC(O)→,
- 15 19) R⁸R⁹NC(=Y)→,
- 20) C₁-C₆ alkyl-S(O)₂→,
- 21) C₃-C₇ cycloalkyl-S(O)₂→,
- 22) aryl-S(O)₂→,
- 23) heteroaryl-S(O)₂→,
- 20 24) heterocyclyl-S(O)₂→,
- 25) fused aryl-C₃-C₇ cycloalkyl→,
- 26) fused heteroaryl-C₃-C₇ cycloalkyl→,
- 27) fused aryl-heterocyclyl→,
- 28) fused heteroaryl-heterocyclyl→,
- 25 29) fused aryl-C₃-C₇ cycloalkyl-O_nC(O)→,
- 30) fused heteroaryl-C₃-C₇ cycloalkyl-O_nC(O)→,
- 31) fused aryl-heterocyclyl-O_nC(O)→, or
- 32) fused heteroaryl-heterocyclyl-O_nC(O)→,

wherein the alkyl and the cycloalkyl are optionally substituted with one or more R⁶ substituents, and the aryl, the heteroaryl and the heterocyclyl are optionally substituted with one or more R¹⁰ substituents;

R⁶ is independently selected from:

- 1) halogen,
- 35 2) C₁-C₆ alkyl,

- 3) C₃–C₇ cycloalkyl,
- 4) haloalkyl,
- 5) aryl,
- 6) heteroaryl,
- 5 7) heterocyclyl,
- 8) OR⁷,
- 9) S(O)_mR⁷,
- 10) NR⁸R⁹,
- 11) COR⁷,
- 10 12) C(O)OR⁷,
- 13) OC(O)R⁷,
- 14) SC(O)R⁷,
- 15) CONR⁸R⁹,
- 16) S(O)₂NR⁸R⁹, or
- 15 17) N(=Y)NR⁸R⁹,

wherein the aryl, the heteroaryl and the heterocyclyl are optionally substituted with one or more R¹⁰ substituents;

R⁷ is independently selected from:

- 20 1) H,
- 2) C₁–C₆ alkyl,
- 3) C₃–C₇ cycloalkyl,
- 4) haloalkyl,
- 5) aryl,
- 25 6) heteroaryl,
- 7) heterocyclyl,
- 8) C(=Y)NR⁸R⁹, or
- 9) C₁–C₆ alkyl-C₂–C₄ alkynyl,

wherein the aryl, heteroaryl, and heterocyclyl are optionally substituted with one or more R¹⁰;

R⁸ and R⁹ are independently selected from:

- 1) H,
- 2) C₁–C₆ alkyl,
- 35 3) C₃–C₇ cycloalkyl,

- 4) haloalkyl,
- 5) aryl,
- 6) heteroaryl,
- 7) heterocyclyl,
- 5 8) $\text{COC}_1\text{-C}_6$ alkyl,
- 9) $\text{COC}_3\text{-C}_3$ cycloalkyl
- 10) CO-aryl,
- 11) CO-heteroaryl,
- 12) CO-heterocyclyl,
- 10 13) $\text{C(O)Y-C}_1\text{-C}_6$ alkyl,
- 14) $\text{C(O)Y-C}_3\text{-C}_3$ cycloalkyl
- 15) C(O)Y-aryl ,
- 16) C(O)Y-heteroaryl , or
- 17) $\text{C(O)Y heterocyclyl}$,
- 15 wherein the aryl, the heteroaryl and the heterocyclyl are optionally substituted with one or more R^{10} substituents;
or R^8 and R^9 together with the nitrogen atom to which they are bonded form a five, six or seven membered heterocyclic ring optionally substituted with one or more R^6 substituents;
- 20 R^{10} is independently selected from:
 - 1) halogen,
 - 2) NO_2 ,
 - 3) CN ,
 - 4) $\text{C}_1\text{-C}_6$ alkyl,
 - 25 5) haloalkyl,
 - 6) $\text{C}_3\text{-C}_7$ cycloalkyl,
 - 7) OR^7 ,
 - 8) NR^8R^9 ,
 - 9) SR^7 ,
 - 30 10) COR^7 ,
 - 11) CO_2R^7 ,
 - 12) $\text{S(O)}_m\text{R}^7$,
 - 13) CONR^8R^9 , or
 - 14) $\text{S(O)}_2\text{NR}^8\text{R}^9$,
 - 35 wherein the alkyl is optionally substituted with one or more R^6 substituents;

R¹¹ is independently selected from

- 1) C₁–C₆ alkyl→,
- 2) C₃–C₇ cycloalkyl→,
- 5 3) aryl→,
- 4) heteroaryl→, or
- 5) heterocyclyl→,

wherein the alkyl and the cycloalkyl are optionally substituted with one or more R⁶ substituents, and the aryl, the heteroaryl and the heterocyclyl are optionally substituted
 10 with one or more R¹⁰ substituents;

R¹² is independently selected from

- 1) C₁–C₆ alkyl→,
- 2) C₃–C₇ cycloalkyl→,
- 15 3) haloalkyl→,
- 4) aryl→,
- 5) heteroaryl→,
- 6) heterocyclyl→,
- 7) C₁–C₆ alkyl–O_nC(O)→,
- 20 8) haloalkyl–O_nC(O)→,
- 9) C₃–C₇ cycloalkyl–O_nC(O)→,
- 10) aryl–O_nC(O)→,
- 11) heteroaryl–O_nC(O)→,
- 12) heterocyclyl–O_nC(O)→,
- 25 13) R⁸R⁹NC(O)→,
- 14) C₁–C₆ alkyl–S(O)_m→,
- 15) C₃–C₇ cycloalkyl–S(O)_m→,
- 16) aryl–S(O)_m→,
- 17) heteroaryl–S(O)_m→,
- 30 18) heterocyclyl–S(O)_m→,
- 19) fused aryl–C₃–C₇ cycloalkyl,
- 20) fused heteroaryl–C₃–C₇ cycloalkyl, or
- 21) C(=Y)–NR⁸R⁹,

wherein the alkyl and the cycloalkyl are optionally substituted with one or more R^6 substituents, and the aryl, the heteroaryl and the heterocyclyl are optionally substituted with one or more R^{10} substituents;

5 R^{13} is

- 1) H,
- 2) C_1-C_6 alkyl→,
- 3) C_3-C_7 cycloalkyl→,
- 4) haloalkyl→,
- 10 5) aryl→,
- 6) heteroaryl→,
- 7) heterocyclyl→,
- 8) C_1-C_6 alkyl- $O_nC(O)$ →,
- 9) haloalkyl- $O_nC(O)$ →,
- 15 10) C_3-C_7 cycloalkyl- $O_nC(O)$ →,
- 11) aryl- $O_nC(O)$ →,
- 12) heteroaryl- $O_nC(O)$ →, or
- 13) heterocyclyl- $O_nC(O)$ →;

20 or a prodrug, or a pharmaceutically acceptable salt, or labeled with a detectable label or an affinity tag thereof.

If any variable, such as R^6 , R^{600} , R^{10} , R^{1000} and the like, occurs more than one time in any constituent structure, the definition of the variable at each occurrence is independent at every other occurrence. If a substituent is itself substituted with one or more substituents, it is to be understood that the one or more substituents may be attached to the same carbon atom or different carbon atoms. Combinations of substituents and variables defined herein are allowed only if they produce chemically stable compounds.

30

One skilled in the art will understand that substitution patterns and substituents on compounds of the present invention may be selected to provide compounds that are chemically stable and can be readily synthesized using the chemistry set forth in the examples and chemistry techniques well known in the art using readily available starting materials.

35

It is to be understood that many substituents or groups described herein have functional group equivalents, which means that the group or substituent may be replaced by another group or substituent that has similar electronic, hybridization or bonding properties.

5

Definitions

Unless otherwise specified, the following definitions apply:

10 The singular forms "a", "an" and "the" include corresponding plural references unless the context clearly dictates otherwise.

As used herein, the term "comprising" is intended to mean that the list of elements following the word "comprising" are required or mandatory but that other elements are optional and may or may not be present .

15

As used herein, the term "consisting of" is intended to mean including and limited to whatever follows the phrase "consisting of". Thus the phrase "consisting of" indicates that the listed elements are required or mandatory and that no other elements may be present.

20 As used herein, the term "alkyl" is intended to include both branched and straight chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, for example, C₁-C₆ as in C₁-C₆ - alkyl is defined as including groups having 1, 2, 3, 4, 5 or 6 carbons in a linear or branched arrangement, and C₁-C₄ as in C₁-C₄ alkyl is defined as including groups having 1, 2, 3, or 4 carbons in a linear or branched arrangement, and for
25 example, C₁-C₂₀ as in C₁-C₂₀ - alkyl is defined as including groups having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 carbons in a linear or branched arrangement, Examples of C₁-C₆-alkyl and C₁-C₄ alkyl as defined above include, but are not limited to, methyl, ethyl, n-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *i*-butyl, pentyl and hexyl.

30 As used herein, the term, "alkenyl" is intended to mean unsaturated straight or branched chain hydrocarbon groups having the specified number of carbon atoms therein, and in which at least two of the carbon atoms are bonded to each other by a double bond, and having either E or Z regeochemistry and combinations thereof. For example, C₂-C₆ as in C₂-C₆ alkenyl is defined as including groups having 2, 3, 4, 5, or 6 carbons in a linear or
35 branched arrangement, at least two of the carbon atoms being bonded together by a

double bond. Examples of C₂-C₆ alkenyl include ethenyl (vinyl), 1-propenyl, 2-propenyl, 1-butenyl and the like.

As used herein, the term "alkynyl" is intended to mean unsaturated, straight chain hydrocarbon groups having the specified number of carbon atoms therein and in which at least two carbon atoms are bonded together by a triple bond. For example C₂-C₄ as in C₂-C₄ alkynyl is defined as including groups having 2, 3, or 4 carbon atoms in a chain, at least two of the carbon atoms being bonded together by a triple bond. Examples of such alkynyls include ethynyl, 1-propynyl, 2-propynyl and the like.


As used herein, the term "cycloalkyl" is intended to mean a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms therein, for example, C₃-C₇ as in C₃-C₇ cycloalkyl is defined as including groups having 3, 4, 5, 6, or 7 carbons in a monocyclic arrangement. Examples of C₃-C₇ cycloalkyl as defined above include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

As used herein, the term "cycloalkenyl" is intended to mean a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms therein, for example, C₃-C₇ as in C₃-C₇ cycloalkenyl is defined as including groups having 3, 4, 5, 6, or 7 carbons in a monocyclic arrangement. Examples of C₃-C₇ cycloalkenyl as defined above include, but are not limited to, cyclopentenyl, and cyclohexenyl.

As used herein, the term "halo" or "halogen" is intended to mean fluorine, chlorine, bromine and iodine.


As used herein, the term "haloalkyl" is intended to mean an alkyl as defined above, in which each hydrogen atom may be successively replaced by a halogen atom. Examples of haloalkyls include, but are not limited to, CH₂F, CHF₂ and CF₃.

As used herein, the term "aryl", either alone or in combination with another radical, means a carbocyclic aromatic monocyclic group containing 6 carbon atoms which may be further fused to a second 5- or 6-membered carbocyclic group which may be aromatic, saturated or unsaturated. Aryl includes, but is not limited to, phenyl, indanyl, 1-naphthyl, 2-naphthyl and tetrahydronaphthyl. The aryls may be connected to another group either at a suitable position on the cycloalkyl ring or the aromatic ring. For example:



5 Arrowed lines drawn from the ring system indicate that the bond may be attached to any of the suitable ring atoms.

As used herein, the term "biphenyl" is intended to mean two phenyl groups bonded together at any one of the available sites on the phenyl ring. For example:




10

As used herein, the term "heteroaryl" is intended to mean a monocyclic or bicyclic ring system of up to ten atoms, wherein at least one ring is aromatic, and contains from 1 to 4 hetero atoms selected from the group consisting of O, N, and S. The heteroaryl substituent may be attached either via a ring carbon atom or one of the heteroatoms.

15 Examples of heteroaryl groups include, but are not limited to thienyl, benzimidazolyl, benzo[b]thienyl, furyl, benzofuranyl, pyranal, isobenzofuranyl, chromenyl, xanthenyl, 2H-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, indolyl, indazolyl, purinyl, 4H-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnoliny, pteridinyl, isothiazolyl, isochromanyl, chromanyl, isoxazolyl, furazanyl, indoliny, isoindoliny, thiazolo[4,5-b]-pyridine, and


20

fluorescein derivatives such as:



or

As used herein, the term "heterocycle", "heterocyclic" or "heterocyclyl" is intended to mean a 5, 6, or 7 membered non-aromatic ring system containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Examples of heterocycles include, but are not limited to pyrrolidinyl, tetrahydrofuranyl, piperidyl, pyrrolinyl, piperazinyl,



5 imidazolidinyl, morpholinyl, imidazolyl, pyrazolidinyl, pyrazolyl, and

As used herein, the term "heterobicycle" either alone or in combination with another radical, is intended to mean a heterocycle as defined above fused to another cycle, be it a heterocycle, an aryl or any other cycle defined herein. Examples of such heterobicycles include, but are not limited to, coumarin, benzo[d][1,3]dioxole, 2,3-

10 dihydrobenzo[b][1,4]dioxine and 3,4-dihydro-2H-benzo[b][1,4]dioxepine.

As used herein, the term "heteroaryl" is intended to mean a monocyclic or bicyclic ring system of up to ten atoms, wherein at least one ring is aromatic, and contains from 1 to 4

15 hetero atoms selected from the group consisting of O, N, and S. The heteroaryl substituent may be attached either via a ring carbon atom or one of the heteroatoms. Examples of heteroaryl groups include, but are not limited to thienyl, benzimidazolyl, benzo[b]thienyl, furyl, benzofuranyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, 2H-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl,

20 indolizyl, isoindolyl, 3H-indolyl, indolyl, indazolyl, purinyl, 4H-quinolizyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, isothiazolyl, isochromanyl, chromanyl, isoxazolyl, furazanyl, indolinyl, and isoindolinyl,

As used herein, the term "heterocycle", "heterocyclic" or "heterocyclyl" is intended to mean

25 a 5, 6, or 7 membered non-aromatic ring system containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Examples of heterocycles include, but are not limited to pyrrolidinyl, tetrahydrofuranyl, piperidyl, pyrrolinyl, piperazinyl, imidazolidinyl, morpholinyl, imidazolyl, pyrazolidinyl, and pyrazolyl,

30 As used herein, the term "heteroatom" is intended to mean O, S or N.

As used herein, the term "activated diacid" is intended to mean a diacid wherein the carboxylic acid moieties have been transformed to, for example, but not limited to, acid halides, a succinate esters, or HOBt esters, either in situ or in a separate synthetic step. For example, succinyl chloride and terephthaloyl chloride are examples of "diacid chlorides". HOBt esters can be formed in situ by the treatment of a diacid with a
5 dehydrating agent such as DCC, EDC, HBTU, or others, a base such as DIPEA, and HOBt in an appropriate solvent. The reaction of an activated diacid with an amine will result in the conversion of the acid functionality to an amide functionality.

10 As used herein, the term "detectable label" is intended to mean a group that may be linked to a compound of the present invention to produce a probe or to an IAP BIR domain, such that when the probe is associated with the BIR domain, the label allows either direct or indirect recognition of the probe so that it may be detected, measured and quantified. As used herein, the term "affinity tag" is intended to mean a ligand or group, which is
15 linked to either a compound of the present invention or to an IAP BIR domain to allow another compound to be extracted from a solution to which the ligand or group is attached.

As used herein, the term "probe" is intended to mean a compound of Formula I which is
20 labeled with either a detectable label or an affinity tag, and which is capable of binding, either covalently or non-covalently, to an IAP BIR domain. When, for example, the probe is non-covalently bound, it may be displaced by a test compound. When, for example, the probe is bound covalently, it may be used to form cross-linked adducts, which may be quantified and inhibited by a test compound.

25 As used herein, the term "optionally substituted with one or more substituents" or its equivalent term "optionally substituted with at least one substituent" is intended to mean that the subsequently described event or circumstances may or may not occur, and that the description includes instances where the event or circumstance occurs and instances
30 in which it does not. The definition is intended to mean from zero to five substituents.

If the substituents themselves are incompatible with the synthetic methods of the present invention, the substituent may be protected with a suitable protecting group (PG) that is stable to the reaction conditions used in these methods. The protecting group may be
35 removed at a suitable point in the reaction sequence of the method to provide a desired

intermediate or target compound. Suitable protecting groups and the methods for protecting and de-protecting different substituents using such suitable protecting groups are well known to those skilled in the art; examples of which may be found in T. Greene and P. Wuts, *Protecting Groups in Chemical Synthesis* (3rd ed.), John Wiley & Sons, NY (1999), which is incorporated herein by reference in its entirety. Examples of protecting groups used throughout include, but are not limited to Fmoc, Bn, Boc, CBz and COCF₃. In some instances, a substituent may be specifically selected to be reactive under the reaction conditions used in the methods of this invention. Under these circumstances, the reaction conditions convert the selected substituent into another substituent that is either useful in an intermediate compound in the methods of this invention or is a desired substituent in a target compound.

Abbreviations for α -amino acids used throughout are as follows:

Amino acid	Abbreviation
α -Amino butyric acid	Abu
Alanine	Ala
Arginine	Arg
Aspartic acid	Asp
Asparagine	Asn
Cysteine	Cys
Glutamic acid	Glu
Glutamine	Gln
Glycine	Gly
Isoleucine	Ile
Histidine	His
Leucine	Leu
Lysine	Lys
Methionine	Met
Phenylalanine	Phe
Proline	Pro
Serine	Ser
Threonine	Thr
Tryptophan	Trp

Amino acid	Abbreviation
Tyrosine	Tyr
Valine	Val

As used herein, the term "residue" when referring to α -amino acids is intended to mean a radical derived from the corresponding α -amino acid by eliminating the hydroxyl of the carboxy group and one hydrogen of the α -amino group. For example, the terms Gln, Ala, Gly, Ile, Arg, Asp, Phe, Ser, Leu, Cys, Asn, and Tyr represent the residues of L-glutamine, L-alanine, glycine, L-isoleucine, L-arginine, L-aspartic acid, L-phenylalanine, L-serine, L-leucine, L-cysteine, L-asparagine, and L-tyrosine, respectively.

As used herein, the term "subject" is intended to mean humans and non-human mammals such as primates, cats, dogs, swine, cattle, sheep, goats, horses, rabbits, rats, mice and the like.

As used herein, the term "prodrug" is intended to mean a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound of the present invention. Thus, the term "prodrug" refers to a precursor of a compound of the invention that is pharmaceutically acceptable. A prodrug may be inactive or display limited activity when administered to a subject in need thereof, but is converted in vivo to an active compound of the present invention. Typically, prodrugs are transformed in vivo to yield the compound of the invention, for example, by hydrolysis in blood or other organs by enzymatic processing. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in the subject (see, Bundgard, H., Design of Prodrugs (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam). The definition of prodrug includes any covalently bonded carriers which release the active compound of the invention in vivo when such prodrug is administered to a subject. Prodrugs of a compound of the present invention may be prepared by modifying functional groups present in the compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or in vivo, to a parent compound of the invention.

As used herein, the term "pharmaceutically acceptable carrier, diluent or excipient" is intended to mean, without limitation, any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent,

dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, emulsifier, or encapsulating agent, such as a liposome, cyclodextrins, encapsulating polymeric delivery systems or polyethyleneglycol matrix, which is acceptable for use in the subject, preferably humans.

5

As used herein, the term "pharmaceutically acceptable salt" is intended to mean both acid and base addition salts.

10

As used herein, the term "pharmaceutically acceptable acid addition salt" is intended to mean those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

15

20

As used herein, the term "pharmaceutically acceptable base addition salt" is intended to mean those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like.

25

30

35

As used herein, the term "BIR domain binding" is intended to mean the action of a compound of the present invention upon an IAP BIR domain, which blocks or diminishes

the binding of IAPs to BIR binding proteins or is involved in displacing BIR binding proteins from an IAP. Examples of BIR binding proteins include, but are not limited to, caspases and mitochondrially derived BIR binding proteins such as Smac, Omi/WTR2A and the like.

- 5 As used herein, the term "insufficient apoptosis" is intended to mean a state wherein a disease is caused or continues because cells deleterious to the subject have not apoptosed. This includes, but is not limited to, cancer cells that survive in a subject without treatment, cancer cells that survive in a subject during or following anti-cancer treatment, or immune cells whose action is deleterious to the subject, and includes,
10 neutrophils, monocytes and auto-reactive T-cells.

- As used herein, the term "therapeutically effective amount" is intended to mean an amount of a compound of Formula I or II which, when administered to a subject is sufficient to effect treatment for a disease-state associated with insufficient apoptosis. The amount of
15 the compound of Formula I will vary depending on the compound, the condition and its severity, and the age of the subject to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

- As used herein, the term "treating" or "treatment" is intended to mean treatment of a
20 disease-state associated with insufficient apoptosis, as disclosed herein, in a subject, and includes: (i) preventing a disease or condition associated with insufficient apoptosis from occurring in a subject, in particular, when such mammal is predisposed to the disease or condition but has not yet been diagnosed as having it; (ii) inhibiting a disease or condition associated with insufficient apoptosis, i.e., arresting its development; or (iii) relieving a
25 disease or condition associated with insufficient apoptosis, i.e., causing regression of the condition.

- As used herein, the term "treating cancer" is intended to mean the administration of a pharmaceutical composition of the present invention to a subject, preferably a human,
30 which is afflicted with cancer to cause an alleviation of the cancer by killing, inhibiting the growth, or inhibiting the metastasis of the cancer cells.

- As used herein, the term "preventing disease" is intended to mean, in the case of cancer, the post-surgical, post-chemotherapy or post-radiotherapy administration of a
35 pharmaceutical composition of the present invention to a subject, preferably a human,

which was afflicted with cancer to prevent the regrowth of the cancer by killing, inhibiting the growth, or inhibiting the metastasis of any remaining cancer cells. Also included in this definition is the prevention of pro-survival conditions that lead to diseases such as asthma, MS and the like.

5

As used herein, the term "synergistic effect" is intended to mean that the effect achieved with the combination of the compounds of the present invention and either the chemotherapeutic agents or death receptor agonists of the invention is greater than the effect which is obtained with only one of the compounds, agents or agonists, or
 10 advantageously the effect which is obtained with the combination of the above compounds, agents or agonists is greater than the addition of the effects obtained with each of the compounds, agents or agonists used separately. Such synergy enables smaller doses to be given.

15 As used herein, the term "apoptosis" or "programmed cell death" is intended to mean the regulated process of cell death wherein a dying cell displays a set of well-characterized biochemical hallmarks that include cell membrane blebbing, cell soma shrinkage, chromatin condensation, and DNA laddering, as well as any caspase-mediated cell death.

20 As used herein, the term "BIR domain" or "BIR" are used interchangeably throughout and are intended to mean a domain which is characterized by a number of invariant amino acid residue including conserved cysteines and one conserved histidine residue within the sequence Cys-(Xaa1)₂Cys-(Xaa1)₁₆His-(Xaa1)₆₋₈Cys. Typically, the amino acid sequence of the consensus sequence is: Xaa1-Xaa1-Xaa1-Arg-Leu-Xaa1-Thr-Phe-Xaa1-Xaa1-Trp -
 25 Pro-Xaa2-Xaa1-Xaa1-Xaa2-Xaa2-Xaa1-Xaa1-Xaa1-Xaa1-Leu-Ala-Xaa1-Ala-Gly-Phe-Tyr-
 Tyr-Xaa1-Gly-Xaa1-Xaa1-Asp-Xaa1-Val-Xaa1-Cys-Phe-Xaa1-Cys-Xaa1-Xaa1-Xaa1-
 Xaa1-Xaa1-Xaa1-Trp-Xaa1-Xaa1-Xaa1-Asp-Xaa1-Xaa1-Xaa1-Xaa1-Xaa1-His-Xaa1-
 Xaa1-Xaa1-Xaa1-Pro-Xaa1-Cys-Xaa1-Phe-Val, wherein Xaa1 is any amino acid and Xaa2 is any amino acid or is absent. Preferably the sequence is substantially identical to
 30 one of the BIR domain sequences provided for XIAP, HIAP1, or HIAP2 herein.
 The BIR domain residues are listed below (see Genome Biology (2001) 1-10):

	XIAP	HIAP-1	HIAP-2
BIR1	21-93	41-113	24-96

BIR2	159-230	179-250	164-235
BIR3	258-330	264-336	250-322
Seq. #	P98170	XP-006266	XP-006267

As used herein, the term "ring zinc finger" or "RZF" is intended to mean a domain having the amino acid sequence of the consensus sequence: Glu-Xaa1-Xaa1-Xaa1-Xaa1-Xaa1-Xaa- 1-Xaa2-Xaa1-Xaa1-Xaa1-Cys-Lys-Xaa3-Cys-Met-Xaa1-Xaa1-Xaa1-Xaa1-Xaa1-
5 Xaa3-X- aa1-Phe-Xaa1-Pro-Cys-Gly-His-Xaa1-Xaa1-Xaa1-Cys-Xaa1-Xaa1-Cys-Ala-Xaa1-Xaa- 1-Xaa1-Xaa1-Xaa1-Cys-Pro-Xaa1-Cys, wherein Xaa1 is any amino acid, Xaa2 is Glu or Asp, and Xaa3 is Val or Ile.

As used herein, the term "IAP" is intended to mean a polypeptide or protein, or fragment
10 thereof, encoded by an IAP gene. Examples of IAPs include, but are not limited to human or mouse NAIP (Birc 1), HIAP-1 (clAP2, Birc 3), HIAP-2 (clAP1, Birc 2), XIAP (Birc 4), survivin (Birc 5), livin (ML-IAP, Birc 7), ILP-2 (Birc 8) and Apollon/BRUCE (Birc 6) (see for example US Patent Numbers 6,107,041; 6,133,437; 6,156,535; 6,541,457; 6,656,704; 6,689,562; Deveraux and Reed, Genes Dev. 13, 239-252, 1999; Kasof and Gomes, J.
15 Biol. Chem., 276, 3238-3246, 2001; Vucic et al., Curr. Biol. 10, 1359-1366, 2000; Ashab et al. FEBS Lett., 495, 56-60, 2001, the contents of which are hereby incorporated by reference).

As used herein, the term "IAP gene" is intended to mean a gene encoding a polypeptide
20 having at least one BIR domain and which is capable of modulating (inhibiting or enhancing) apoptosis in a cell or tissue. The IAP gene is a gene having about 50% or greater nucleotide sequence identity to at least one of human or mouse NAIP (Birc 1), HIAP-1 (clAP2, Birc 3), HIAP-2 (clAP1, Birc 2), XIAP (Birc 4), survivin (Birc 5), livin (ML-IAP, Birc 7), ILP-2 (Birc 8) and Apollon/BRUCE (Birc 6). The region of sequence over
25 which identity is measured is a region encoding at least one BIR domain and a ring zinc finger domain. Mammalian IAP genes include nucleotide sequences isolated from any mammalian source.

As used herein, the term "IC₅₀" is intended to mean an amount, concentration or dosage
30 of a particular compound of the present invention that achieves a 50% inhibition of a

maximal response, such as displacement of maximal fluorescent probe binding in an assay that measures such response.

As used herein, the term "EC₅₀" is intended to mean an amount, concentration or dosage of a particular compound of the present invention that achieves a 50% inhibition of cell survival.

As used herein, the term "modulate" or "modulating" is intended to mean the treatment, prevention, suppression, enhancement or induction of a function or condition using the compounds of the present invention. For example, the compounds of the present invention can modulate IAP function in a subject, thereby enhancing apoptosis by significantly reducing, or essentially eliminating the interaction of activated apoptotic proteins, such as caspase-3, 7 and 9, with the BIR domains of mammalian IAPs or by inducing the loss of XIAP protein in a cell.

As used herein, the term "enhancing apoptosis" is intended to mean increasing the number of cells that apoptose in a given cell population either in vitro or in vivo. Examples of cell populations include, but are not limited to, ovarian cancer cells, colon cancer cells, breast cancer cells, lung cancer cells, pancreatic cancer cells, or T cells and the like. It will be appreciated that the degree of apoptosis enhancement provided by an apoptosis-enhancing compound of the present invention in a given assay will vary, but that one skilled in the art can determine the statistically significant change in the level of apoptosis that identifies a compound that enhances apoptosis otherwise limited by an IAP.

Preferably "enhancing apoptosis" means that the increase in the number of cells undergoing apoptosis is at least 25%, more preferably the increase is 50%, and most preferably the increase is at least one-fold. Preferably the sample monitored is a sample of cells that normally undergo insufficient apoptosis (i.e., cancer cells). Methods for detecting the changes in the level of apoptosis (i.e., enhancement or reduction) are described in the Examples and include methods that quantitate the fragmentation of DNA, methods that quantitate the translocation phosphatoylserine from the cytoplasmic to the extracellular side of the membrane, determination of activation of the caspases and methods quantitate the release of cytochrome C and the apoptosis inhibitory factor into the cytoplasm by mitochondria.

As used herein, the term "proliferative disease" or "proliferative disorder" is intended to mean a disease that is caused by or results in inappropriately high levels of cell division, inappropriately low levels of apoptosis, or both. For example, cancers such as lymphoma, leukemia, melanoma, ovarian cancer, breast cancer, pancreatic cancer, and lung cancer, and autoimmune disorders are all examples of proliferative diseases.

As used herein, the term "death receptor agonist" is intended to mean an agent capable of stimulating by direct or indirect contact the pro apoptotic response mediated by the death-receptors. For example, an agonist TRAIL receptor Antibody would bind to TRAIL receptor (S) and trigger an apoptotic response. On the other hand, other agent such as interferon- α could trigger the release of endogenous TRAIL and/or up regulate the TRAIL receptors in such a way that the cell pro-apoptotic response is amplified.

The compounds of the present invention, or their pharmaceutically acceptable salts may contain one or more asymmetric centers, chiral axes and chiral planes and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms and may be defined in terms of absolute stereochemistry, such as (R)- or (S)- or, as (D)- or (L)- for amino acids. The present invention is intended to include all such possible isomers, as well as, their racemic and optically pure forms. Optically active (+) and (-), (R)- and (S)-, or (D)- and (L)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, such as reverse phase HPLC. The racemic mixtures may be prepared and thereafter separated into individual optical isomers or these optical isomers may be prepared by chiral synthesis. The enantiomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts which may then be separated by crystallization, gas-liquid or liquid chromatography, selective reaction of one enantiomer with an enantiomer specific reagent. It will also be appreciated by those skilled in the art that where the desired enantiomer is converted into another chemical entity by a separation technique, an additional step is then required to form the desired enantiomeric form. Alternatively specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts, or solvents or by converting one enantiomer to another by asymmetric transformation.

Certain compounds of the present invention may exist in Zwitterionic form and the present invention includes Zwitterionic forms of these compounds and mixtures thereof.

Utilities

The compounds of the present invention are useful as IAP BIR domain binding compounds and as such the compounds, compositions and method of the present invention include application to the cells or subjects afflicted with or having a predisposition towards developing a particular disease state, which is characterized by insufficient apoptosis. Thus, the compounds, compositions and methods of the present invention are used to treat cellular proliferative diseases/disorders, which include, but are not limited to, i) cancer, ii) autoimmune disease, iii) inflammatory disorders, iv) proliferation induced post medical procedures, including, but not limited to, surgery, angioplasty, and the like.

The compounds of the present invention may also be useful in the treatment of diseases in which there is a defect in the programmed cell-death or the apoptotic machinery (TRAIL, FAS, apoptosome), such as multiple sclerosis, arteriosclerosis, inflammation, autoimmunity and the like.

The treatment involves administration to a subject in need thereof a compound of the present invention or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof. In particular, the compounds, compositions and methods of the present invention are useful for the treatment of cancer including solid tumors such as skin, breast, brain, lung, testicular carcinomas, and the like. Cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to the following:

Tissue	Example
Adrenal gland	neuroblastoma
Bone	osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondromatous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid

Tissue	Example
	osteoma and giant cell tumors
Cardiac	sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma
Gastrointestinal	esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma)
Genitourinary tract	kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma)
Gynecological	uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma)
Hematologic	blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]

Tissue	Example
Liver	hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma
Lung	bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma
Nervous system	skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma)
Skin	malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids

The compounds of the present invention, or their pharmaceutically acceptable salts or their prodrugs, may be administered in pure form or in an appropriate pharmaceutical composition, and can be carried out via any of the accepted modes of Galenic
5 pharmaceutical practice.

The pharmaceutical compositions of the present invention can be prepared by mixing a compound of the present invention with an appropriate pharmaceutically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid,
10 liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Typical routes of administering such pharmaceutical compositions include, without limitation, oral, topical, transdermal, inhalation, parenteral (subcutaneous injections, intravenous,
intramuscular, intrasternal injection or infusion techniques), sublingual, ocular, rectal,
15 vaginal, and intranasal. Pharmaceutical compositions of the present invention are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a subject. Compositions that will be administered to a

subject or patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a compound of the present invention in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see

5 Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pa., 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state as described above.

10 A pharmaceutical composition of the present invention may be in the form of a solid or liquid. In one aspect, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral syrup, injectable liquid or an aerosol, which is useful in, for example inhalatory administration.

15 For oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

20 As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or
25 gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent.

30 When the pharmaceutical composition is in the form of a capsule, e.g., a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil such as soybean or vegetable oil.

35 The pharmaceutical composition may be in the form of a liquid, e.g., an elixir, syrup,

solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid pharmaceutical compositions of the present invention, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; encapsulating agents such as cyclodextrins or functionalized cyclodextrins, including, but not limited to, α , β , or δ -hydroxypropylcyclodextrins or Captisol; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediamine tetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. An injectable pharmaceutical composition is preferably sterile.

A liquid pharmaceutical composition of the present invention used for either parenteral or oral administration should contain an amount of a compound of the present invention such that a suitable dosage will be obtained. Typically, this amount is at least 0.01% of a compound of the present invention in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition. For parenteral usage, compositions and preparations according to the present invention are prepared so that a parenteral dosage unit contains between 0.01 to 10% by weight of the compound of the present invention. Pharmaceutical compositions may be further diluted at the time of administration; for example a parenteral formulation may be further diluted with a sterile, isotonic solution for injection such as 0.9 % saline, 5 wt % dextrose (D5W), Ringer's solution, or others.

The pharmaceutical composition of the present invention may be used for topical

administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of the compound of the present invention from about 0.1 to about 10% w/v (weight per unit volume).

The pharmaceutical composition of the present invention may be used for rectal administration to treat for example, colon cancer, in the form, e.g., of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

The pharmaceutical composition of the present invention may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

The pharmaceutical composition of the present invention in solid or liquid form may include an agent that binds to the compound of the present invention and thereby assists in the delivery of the compound. Suitable agents that may act in this capacity include, but are not limited to, a monoclonal or polyclonal antibody, a protein or a liposome.

The pharmaceutical composition of the present invention may consist of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols of compounds of the present invention may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, subcontainers, and the like, which together may form a kit. One skilled

in the art, without undue experimentation may determine preferred aerosols.

The pharmaceutical compositions of the present invention may be prepared by methodology well known in the pharmaceutical art. For example, a pharmaceutical composition intended to be administered by injection can be prepared by admixing a compound of the present invention with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the compound of the present invention so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

The compounds of the present invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. Generally, a therapeutically effective daily dose may be from about 0.1 mg to about 40 mg/kg of body weight per day or twice per day of a compound of the present invention, or a pharmaceutically acceptable salt thereof.

Combination therapy

The compounds of the present invention, or pharmaceutically acceptable salts thereof, may also be administered simultaneously with, prior to, or after administration of one or more of the therapeutic agents described below. Such combination therapy may include administration of a single pharmaceutical dosage formulation which contains a compound of the present invention and one or more additional agents given below, as well as administration of the compound of the present invention and each of additional agent in its own separate pharmaceutical dosage formulation. For example, a compound of the present invention and a chemotherapeutic agent, such as taxol (paclitaxel), taxotere, etoposide, cisplatin, vincristine, vinblastine, and the like, can be administered to the patient either together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations or via intravenous injection. Where separate dosage formulations are used, the compounds of the present invention and one or more additional agents can be administered at essentially the same time, i.e.,

concurrently, or at separately staggered times, i.e., sequentially; combination therapy is understood to include all these regimens. In addition, these compounds may synergize with molecules that may stimulate the death receptor apoptotic pathway through a direct or indirect manner, as for example, the compounds of the present invention may be used
5 in combination with soluble TRAIL any agent or procedures that can cause an increase in circulating level of TRAIL, such as interferon-alpha or radiation.

Thus, the present invention also encompasses the use of the compounds of the present invention in combination with radiation therapy or one or more additional agents such as
10 those described in WO 03/099211 (PCT/US03/15861), which is hereby incorporated by reference.

Examples of such additional agents include, but are not limited to the following:

- a) an estrogen receptor modulator,
- b) an androgen receptor modulator,
- 15 c) retinoid receptor modulator,
- d) a cytotoxic agent,
- e) an antiproliferative agent,
- f) a prenyl-protein transferase inhibitor,
- g) an HMG-CoA reductase inhibitor,
- 20 h) an HIV protease inhibitor,
- i) a reverse transcriptase inhibitor,
- k) an angiogenesis inhibitor,
- l) a PPAR- γ agonist,
- m) a PPAR- δ agonist,
- 25 n) an inhibitor of inherent multidrug resistance,
- o) an anti-emetic agent,
- p) an agent useful in the treatment of anemia,
- q) agents useful in the treatment of neutropenia,
- r) an immunologic-enhancing drug.
- 30 s) a proteasome inhibitor such as Velcade and MG132 (7-Leu-Leu-aldehyde) (see He et al. in Oncogene (2004) 23, 2554-2558);
- t) an HDAC inhibitor, such as sodium butyrate, phenyl butyrate, hydroamic acids, cyclin tetrapeptide and the like (see Rosato et al., Molecular Cancer Therapeutics 2003, 1273-1284);
- 35 u) an inhibitor of the chemotrypsin-like activity in the proteasome;

- v) E3 ligase inhibitors;
- w) a modulator of the immune system such as interferon-alpha and ionizing radiation (UVB) that can induce the release of cytokines, such as the interleukins, TNF, or induce release of Death receptor Ligands such as TRAIL;
- 5 x) a modulator of death receptors TRAIL and TRAIL agonists such as the humanized antibodies HGS-ETR1 and HGS-ETR2; and
- or in combination or sequentially with radiation therapy, so as to treat the cancer.

Additional combinations may also include agents which reduce the toxicity of the aforesaid
10 agents, such as hepatic toxicity, neuronal toxicity, nephrotoxicity and the like.

In one example, co-administration of one of the compounds of Formula I of the present invention with a death receptor agonist such as TRAIL, such as a small molecule or an antibody that mimics TRAIL may cause an advantageous synergistic effect. Moreover,
15 the compounds of the present invention may be used in combination with any compounds that cause an increase in circulating levels of TRAIL.

Vinca Alkaloids and Related Compounds

Vinca alkaloids that can be used in combination with the nucleobase oligomers of the
20 invention to treat cancer and other neoplasms include vincristine, vinblastine, vindesine, vinflunine, vinorelbine, and anhydrovinblastine.

Dolastatins are oligopeptides that primarily interfere with tubulin at the vinca alkaloid binding domain. These compounds can also be used in combination with the compounds
25 of the invention to treat cancer and other neoplasms. Dolastatins include dolastatin-10 (NCS 376128), dolastatin-15, ILX651, TZT-1027, symplostatin 1, symplostatin 3, and LU103793 (cemadotin).

Cryptophycins (e.g., cryptophycin 1 and cryptophycin 52 (LY355703)) bind tubulin within
30 the vinca alkaloid-binding domain and induce G2/M arrest and apoptosis. Any of these compounds can be used in combination with the compounds of the invention to treat cancer and other neoplasms.

Other microtubule disrupting compounds that can be used in conjunction with the
35 compounds of the invention to treat cancer and other neoplasms are described in U.S.

Pat. Nos. 6,458,765; 6,433,187; 6,323,315; 6,258,841; 6,143,721; 6,127,377; 6,103,698; 6,023,626; 5,985,837; 5,965,537; 5,955,423; 5,952,298; 5,939,527; 5,886,025; 5,831,002; 5,741,892; 5,665,860; 5,654,399; 5,635,483; 5,599,902; 5,530,097; 5,521,284; 5,504,191; 4,879,278; and 4,816,444, and U.S. patent application Publication Nos. 2003/0153505 A1; 5 2003/0083263 A1; and 2003/0055002 A1, each of which is hereby incorporated by reference.

Taxanes and Other Microtubule Stabilizing Compounds

10 Taxanes such as paclitaxel, doxorubicin, RPR 109881A, SB-T-1213, SB-T-1250, SB-T-101187, BMS-275183, BRT 216, DJ-927, MAC-321, IDN5109, and IDN5390 can be used in combination with the compounds of the invention to treat cancer and other neoplasms. Taxane analogs (e.g., BMS-184476, BMS-188797) and functionally related non-taxanes (e.g., epothilones (e.g., epothilone A, epothilone B (EPO906), deoxyepothilone B, and epothilone B lactam (BMS-247550)), eleutherobin, discodermolide, 2-epi-discodermolide, 15 2-des-methyldiscodermolide, 5-hydroxymethyldiscodermolide, 19-des-aminocarbonyldiscodermolide, 9(13)-cyclodiscodermolide, and laulimalide) can also be used in the methods and compositions of the invention.

Other microtubule stabilizing compounds that can be used in combination with the 20 compounds of the invention to treat cancer and other neoplasms are described in U.S. Pat. Nos. 6,624,317; 6,610,736; 6,605,599; 6,589,968; 6,583,290; 6,576,658; 6,515,017; 6,531,497; 6,500,858; 6,498,257; 6,495,594; 6,489,314; 6,458,976; 6,441,186; 6,441,025; 6,414,015; 6,387,927; 6,380,395; 6,380,394; 6,362,217; 6,359,140; 6,306,893; 6,302,838; 6,300,355; 6,291,690; 6,291,684; 6,268,381; 6,262,107; 6,262,094; 6,147,234; 6,136,808; 25 6,127,406; 6,100,411; 6,096,909; 6,025,385; 6,011,056; 5,965,718; 5,955,489; 5,919,815; 5,912,263; 5,840,750; 5,821,263; 5,767,297; 5,728,725; 5,721,268; 5,719,177; 5,714,513; 5,587,489; 5,473,057; 5,407,674; 5,250,722; 5,010,099; and 4,939,168; and U.S. patent application Publication Nos. 2003/0186965 A1; 2003/0176710 A1; 2003/0176473 A1; 2003/0144523 A1; 2003/0134883 A1; 2003/0087888 A1; 2003/0060623 A1; 30 2003/0045711 A1; 2003/0023082 A1; 2002/0198256 A1; 2002/0193361 A1; 2002/0188014 A1; 2002/0165257 A1; 2002/0156110 A1; 2002/0128471 A1; 2002/0045609 A1; 2002/0022651 A1; 2002/0016356 A1; 2002/0002292 A1, each of which is hereby incorporated by reference.

Other chemotherapeutic agents that may be administered with a compound of the present invention are listed in the following Table:

Alkylating agents	cyclophosphamide lomustine busulfan procarbazine ifosfamide altretamine melphalan estramustine phosphate hexamethylmelamine	mechlorethamine thiotepa streptozocin chlorambucil temozolomide dacarbazine semustine carmustine
Platinum agents	cisplatin carboplatinum oxaliplatin ZD-0473 (AnorMED) spiroplatinum lobaplatin (Aeterna) carboxyphthalatoplatinum satraplatin (Johnson Matthey)	tetraplatin BBR-3464 (Hoffmann-La Roche) Orniplatin SM-11355 (Sumitomo) iproplatin AP-5280 (Access)
Antimetabolites	azacytidine tomudex gemcitabine trimetrexate capecitabine deoxycoformycin 5-fluorouracil fludarabine floxuridine pentostatin 2-chlorodeoxyadenosine raltitrexed	6-mercaptopurine hydroxyurea 6-thioguanine decitabine (SuperGen) cytarabin clofarabine (Bioenvision) 2-fluorodeoxy cytidine irofulven (MGI Pharma) methotrexate DMDC (Hoffmann-La Roche) idatrexate ethynylcytidine (Taiho)
Topoisomerase inhibitors	amsacrine rubitecan (SuperGen) epirubicin exatecan mesylate (Daiichi) etoposide quinamed (ChemGenex) teniposide or mitoxantrone gimatecan (Sigma-Tau) irinotecan (CPT-11) diflomotecan (Beaufour-Ipsen) 7-ethyl-10-hydroxy-camptothecin	TAS-103 (Taiho) Topotecan elsamitrucin (Spectrum) dexrazoxanet (TopoTarget) J-107088 (Merck & Co) pixantrone (Novuspharma) BNP-1350 (BioNumerik) rebeccamycin analogue (Exelixis) CKD-602 (Chong Kun Dang) BBR-3576 (Novuspharma) KW-2170 (Kyowa Hakko)

Antitumor antibiotics	dactinomycin (actinomycin D) amonafile doxorubicin (adriamycin) azonafile deoxyrubicin anthrapyrazole valrubicin oxantrazole daunorubicin (daunomycin) losoxantrone epirubicin bleomycin sulfate (blenoxane) therarubicin	bleomycinic acid idarubicin bleomycin A rubidazone bleomycin B plicamycinp mitomycin C porfiromycin MEN-10755 (Menarini) cyanomorpholinodoxorubicin GPX-100 (Gem Pharmaceuticals) mitoxantrone (novantrone)
Antimitotic agents	paclitaxel SB 408075 (GlaxoSmithKline) docetaxel E7010 (Abbott) Colchicines PG-TXL (Cell Therapeutics) vinblastine IDN 5109 (Bayer) Vincristine A 105972 (Abbott) Vinorelbine A 204197 (Abbott) Vindesine LU 223651 (BASF) dolastatin 10 (NCI) D 24851 (ASTAMedica) rhizoxin (Fujisawa) ER-86526 (Eisai) mivobulin (Warner-Lambert) combretastatin A4 (BMS) cemadotin (BASF) isohomohalichondrin-B (PharmaMar)	RPR 109881A (Aventis) ZD 6126 (AstraZeneca) TXD 258 (Aventis) PEG-paclitaxel (Enzon) epothilone B (Novartis) AZ10992 (Asahi) T 900607 (Tularik) IDN-5109 (Indena) T 138067 (Tularik) AVLB (Prescient NeuroPharma) cryptophycin 52 (Eli Lilly) azaepothilone B (BMS) vinflunine (Fabre) BNP-7787 (BioNumerik) auristatin PE (Teikoku Hormone) CA-4 prodrug (OXiGENE) BMS 247550 (BMS) dolastatin-10 (NIH) BMS 184476(BMS) CA-4 (OXiGENE) BMS 188797 (BMS) taxoprexin (Protarga)
Aromatase inhibitors	Aminoglutethimide Exemestane Letrozole atamestane (BioMedicines)	anastrozole YM-511 (Yamanouchi) formestane
Thymidylate synthase inhibitors	pemetrexed (Eli Lilly) nolatrexed (Eximias)	ZD-9331 (BTG) CoFactor™ (BioKeys)
DNA antagonists	trabectedin (PharmaMar) mafosfamide (Baxter International) glufosfamide (Baxter International) apaziquone (Spectrum Pharmaceuticals)	albumin + 32P (Isotope Solutions) O6 benzyl guanine (Paligent) thymectacin (NewBiotics) edotreotide (Novartis)

Farnesyltransferase inhibitors	arglabin (NuOncology Labs) tipifarnib (Johnson & Johnson) lonafarnib (Schering-Plough)	perillyl alcohol (DOR BioPharma) BAY-43-9006 (Bayer)
Pump inhibitors	CBT-1 (CBA Pharma) zosuquidar trihydrochloride (Eli Lilly)	tariquidar (Xenova) biricodar dicitrate (Vertex) MS-209 (Schering AG)
Histone acetyltransferase inhibitors	tacedinaline (Pfizer) pivaloyloxymethyl butyrate (Titan) SAHA (Aton Pharma)	depsipeptide (Fujisawa) MS-275 (Schering AG)
Metalloproteinase inhibitors	Neovastat (Aeterna Laboratories) CMT-3 (CollaGenex)	marimastat (British Biotech) BMS-275291 (Celltech)
Ribonucleoside reductase inhibitors	gallium maltolate (Titan) tezacitabine (Aventis)	triapine (Vion) didox (Molecules for Health)
TNF alpha agonists/antagonists	virulizin (Lorus Therapeutics) revimid (Celgene)	CDC-394 (Celgene)
Endothelin A receptor antagonist	atrasentan (Abbott) YM-598 (Yamanouchi)	ZD-4054 (AstraZeneca)
Retinoic acid receptor agonists	fenretinide (Johnson & Johnson) alitretinoin (Ligand)	LGD-1550 (Ligand)
Immuno-modulators	Interferon dexosome therapy (Anosys) oncophage (Antigenics) pentrix (Australian Cancer Technology) GMK (Progenics) ISF-154 (Trogen) adenocarcinoma vaccine (Biomira) cancer vaccine (Intercell) CTP-37 (A VI BioPharma)	norelin (Biostar) IRX-2 (Immuno-Rx) BLP-25 (Biomira) PEP-005 (Peplin Biotech) MGV (Progenics) synchrovax vaccines (CTL Immuno) beta.-alethine (Dovetail) melanoma vaccine (CTL Immuno) CLL therapy (Vasogen) p21 RAS vaccine (GemVax)

Hormonal and antihormonal agents	estrogens Prednisone conjugated estrogens methylprednisolone ethinyl estradiol prednisolone chlortrianisen aminoglutethimide idenestrol leuprolide hydroxyprogesterone caproate goserelin medroxyprogesterone leuporelin testosterone	bicalutamide testosterone propionate; fluoxymesterone flutamide methyltestosterone octreotide diethylstilbestrol nilutamide megestrol mitotane tamoxifen P-04 (Novogen) Toremofine 2-methoxyestradiol (EntreMed) dexamethasone arzoxifene (Eli Lilly)
Photodynamic agents	talaporfin (Light Sciences) Pd-bacteriopheophorbide (Yeda) Theralux (Theratechnologies) lutetium texaphyrin (Pharmacyclics)	motexafin gadolinium (Pharmacyclics) hypericin
Tyrosine Kinase Inhibitors	imatinib (Novartis) kahalide F (PharmaMar) leflunomide (Sugen/Pharmacia) CEP-701 (Cephalon) ZD1839 (AstraZeneca) CEP-751 (Cephalon) erlotinib (Oncogene Science) MLN518 (Millenium) canertinib (Pfizer) PKC412 (Novartis) squalamine (Genaera) phenoxodiol () SU5416 (Pharmacia) trastuzumab (Genentech) SU6668 (Pharmacia)	C225 (ImClone) ZD4190 (AstraZeneca) rhu-Mab (Genentech) ZD6474 (AstraZeneca) MDX-H210 (Medarex) vatalanib (Novartis) 2C4 (Genentech) PKI166 (Novartis) MDX-447 (Medarex) GW2016 (GlaxoSmithKline) ABX-EGF (Abgenix) EKB-509 (Wyeth) IMC-1C11 (ImClone) EKB-569 (Wyeth)

Miscellaneous agents	
SR-27897 (CCK A inhibitor, Sanofi-Synthelabo) BCX-1777 (PNP inhibitor, BioCryst) tocladesine (cyclic AMP agonist, Ribapharm) ranpirnase (ribonuclease stimulant, Alfacell) alvocidib (CDK inhibitor, Aventis) galarubicin (RNA synthesis inhibitor, Dong-A) CV-247 (COX-2 inhibitor, Ivy Medical) tirapazamine (reducing agent, SRI International) P54 (COX-2 inhibitor, Phytopharm) N-acetylcysteine (reducing agent, Zambon) CapCell™ (CYP450 stimulant, Bavarian Nordic) R-flurbiprofen (NF-kappaB inhibitor, Encore) GCS-100 (gal3 antagonist, GlycoGenesys) 3CPA (NF-kappaB inhibitor, Active Biotech) G17DT immunogen (gastrin inhibitor, Aphton) seocalcitol (vitamin D receptor agonist, Leo) efaproxiral (oxygenator, Allos Therapeutics) 131-I-TM-601 (DNA antagonist, TransMolecular) PI-88 (heparanase inhibitor, Progen) eflornithine (ODC inhibitor, ILEX Oncology) tesmilifene (histamine antagonist, YM BioSciences) minodronic acid (osteoclast inhibitor, Yamanouchi) histamine (histamine H2 receptor agonist, Maxim) indisulam (p53 stimulant, Eisai) tiazofurin (IMPDH inhibitor, Ribapharm) aplidine (PPT inhibitor, PharmaMar) cilengitide (integrin antagonist, Merck KGaA) rituximab (CD20 antibody, Genentech) SR-31747 (IL-1 antagonist, Sanofi-Synthelabo)	gemtuzumab (CD33 antibody, Wyeth Ayerst) CCI-779 (mTOR kinase inhibitor, Wyeth) PG2 (hematopoiesis enhancer, Pharmagenesis) exisulind (PDE V inhibitor, Cell Pathways) Immunol™ (triclosan oral rinse, Endo) CP-461 (PDE V inhibitor, Cell Pathways) triacyluridine (uridine prodrug, Wellstat) AG-2037 (GART inhibitor, Pfizer) SN-4071 (sarcoma agent, Signature BioScience) WX-UK1 (plasminogen activator inhibitor, Wilex) TransMID-107 .TM. (immunotoxin, KS Biomedix) PBI-1402 (PMN stimulant, ProMetic LifeSciences) PCK-3145 (apoptosis promotor, Procyon) bortezomib (proteasome inhibitor, Millennium) doranidazole (apoptosis promotor, Pola) SRL-172 (T cell stimulant, SR Pharma) CHS-828 (cytotoxic agent, Leo) TLK-286 (glutathione S transferase inhibitor, Telik) trans-retinoic acid (differentiator, NIH) PT-100 (growth factor agonist, Point Therapeutics) MX6 (apoptosis promotor, MAXIA) midostaurin (PKC inhibitor, Novartis) apomine (apoptosis promotor, ILEX Oncology) bryostatin-1 (PKC stimulant, GPC Biotech) urocidin (apoptosis promotor, Bioniche) CDA-II (apoptosis promotor, Everlife) Ro-31-7453 (apoptosis promotor, La Roche) SDX-101 (apoptosis promotor, Salmedix) brostallicin (apoptosis promotor, Pharmacia) ceflatonin (apoptosis promotor, ChemGenex)

Additional combinations may also include agents which reduce the toxicity of the aforesaid agents, such as hepatic toxicity, neuronal toxicity, nephrotoxicity and the like.

5 Screening assays

The compounds of the present invention may also be used in a method to screen for other compounds that bind to an IAP BIR domain. Generally speaking, to use the compounds of the invention in a method of identifying compounds that bind to an IAP BIR domain, the IAP is bound to a support, and a compound of the invention is added to the assay.

- 10 Alternatively, the compound of the invention may be bound to the support and the IAP is added.

There are a number of ways in which to determine the binding of a compound of the present invention to the BIR domain. In one way, the compound of the invention, for example, may be fluorescently or radioactively labeled and binding determined directly.

- 5 For example, this may be done by attaching the IAP to a solid support, adding a detectably labeled compound of the invention, washing off excess reagent, and determining whether the amount of the detectable label is that present on the solid support. Numerous blocking and washing steps may be used, which are known to those skilled in the art.

10

In some cases, only one of the components is labeled. For example, specific residues in the BIR domain may be labeled. Alternatively, more than one component may be labeled with different labels; for example, using I¹²⁵ for the BIR domain, and a fluorescent label for the probe.

- 15 The compounds of the invention may also be used as competitors to screen for additional drug candidates or test compounds. As used herein, the terms "drug candidate" or "test compounds" are used interchangeably and describe any molecule, for example, protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, and the like, to be tested for bioactivity. The compounds may be capable of directly or indirectly altering the
- 20 IAP biological activity.

- Drug candidates can include various chemical classes, although typically they are small organic molecules having a molecular weight of more than 100 and less than about 2,500 Daltons. Candidate agents typically include functional groups necessary for structural
- 25 interaction with proteins, for example, hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group. The drug candidates often include cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more functional groups.

- 30 Drug candidates can be obtained from any number of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or

readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means.

Competitive screening assays may be done by combining an IAP BIR domain and a probe to form a probe:BIR domain complex in a first sample followed by adding a test compound from a second sample. The binding of the test is determined, and a change or difference in binding between the two samples indicates the presence of a test compound capable of binding to the BIR domain and potentially modulating the IAP's activity.

In one case, the binding of the test compound is determined through the use of competitive binding assays. In this embodiment, the probe is labeled with a fluorescent label. Under certain circumstances, there may be competitive binding between the test compound and the probe. Test compounds which displace the probe, resulting in a change in fluorescence as compared to control, are considered to bind to the BIR region.

In one case, the test compound may be labeled. Either the test compound, or a compound of the present invention, or both, is added first to the IAP BIR domain for a time sufficient to allow binding to form a complex.

Formation of the probe:BIR domain complex typically require incubations of between 4 °C and 40 °C for between 10 minutes to about 1 hour to allow for high-throughput screening. Any excess of reagents are generally removed or washed away. The test compound is then added, and the presence or absence of the labeled component is followed, to indicate binding to the BIR domain.

In one case, the probe is added first, followed by the test compound. Displacement of the probe is an indication the test compound is binding to the BIR domain and thus is capable of binding to, and potentially modulating, the activity of IAP. Either component can be labeled. For example, the presence of probe in the wash solution indicates displacement by the test compound. Alternatively, if the test compound is labeled, the presence of the probe on the support indicates displacement.

In one case, the test compound may be added first, with incubation and washing, followed by the probe. The absence of binding by the probe may indicate the test compound is bound to the BIR domain with a higher affinity. Thus, if the probe is detected on the

support, coupled with a lack of test compound binding, may indicate the test compound is capable of binding to the BIR domain.

Modulation is tested by screening for a test compound's ability to modulate the activity of IAP and includes combining a test compound with an IAP BIR domain, as described above, and determining an alteration in the biological activity of the IAP. Therefore in this case, the test compound should both bind to the BIR domain (although this may not be necessary), and alter its biological activity as defined herein.

Positive controls and negative controls may be used in the assays. All control and test samples are performed multiple times to obtain statistically significant results. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound probe determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

Typically, the signals that are detected in the assay may include fluorescence, resonance energy transfer, time resolved fluorescence, radioactivity, fluorescence polarization, plasma resonance, or chemiluminescence and the like, depending on the nature of the label. Detectable labels useful in performing screening assays in this invention include a fluorescent label such as Fluorescein, Oregon green, dansyl, rhodamine, tetramethyl rhodamine, texas red, Eu^{3+} ; a chemiluminescent label such as luciferase; colorimetric labels; enzymatic markers; or radioisotopes such as tritium, I^{125} and the like

Affinity tags, which may be useful in performing the screening assays of the present invention include be biotin, polyhistidine and the like.


SYNTHESIS AND METHODOLOGY

General methods for the synthesis of the compounds of the present invention are shown below and are disclosed merely for the purpose of illustration and are not meant to be interpreted as limiting the processes to make the compounds by any other methods. Those skilled in the art will readily appreciate that a number of methods are available for the preparation of the compounds of the present invention.

General Procedures

Several methods for preparing symmetrically or non-symmetrically bridged compounds represented by formula I and formula II may be envisioned. General methods are illustrated in Schemes 1 to 8 and Schemes 17 to 20, while specific examples are
5 illustrated in schemes 9 to 16 and Schemes 21 to 26.

Scheme 1 illustrates a general procedure for the preparation of bis-alkynyl bridged compounds of formula I. N-PG¹-2-hydroxyproline is deprotonated with NaH and treated with propargyl bromide to provide the proline intermediate 1-i. Activation of the carboxylic
10 acid of 1-i with peptide coupling agents and treatment with a primary or secondary amine, and deprotection of PG¹ provides the amide intermediate 1-ii. Peptide coupling of PG²(H)N(R³)CHCO₂H with 1-ii is effected by activation of the carboxylic acid of PG²(H)N(R³)CHCO₂H with peptide coupling agents, followed by the addition of 1-ii to provide the fully protected amide, which may be further deprotected at
15 PG² to provide amide 1-iii. Activation of the carboxylic acid of PG³(R¹)N(R²)CHCO₂H with peptide coupling agents, followed by the addition of 1-iii to provide the amide intermediate 1-iv. The bis-alkynyl bridging moiety is prepared by homo-coupling of the alkyne moieties of 1-iv using an appropriate catalyst system, and subsequent deprotection of PG³, to provide compound 1-v.




Scheme 1

- 5 As illustrated in Scheme 2, intermediate 2-i is prepared via a typical amide coupling/deprotection scheme. As such, the carboxylic acid moiety of PG^1 -cis-2-amino-Pro(PG^2)-OH is activated with amino acid coupling reagents, treated with an amine to


provide the corresponding amide, followed by the removal of PG^1 , under appropriate reaction conditions, to provide intermediate 2-i. In a similar manner, $\text{PG}^3(\text{H})\text{N}(\text{R}^3)\text{HCCO}_2\text{H}$ is coupled with 2-i, followed by deprotection of PG^3 , to provide 2-ii. $\text{PG}^4(\text{R}^1)\text{N}(\text{R}^2)\text{HCCO}_2\text{H}$ is coupled to 2-ii to provide 2-iii. Deprotection of PG^2 provides 2-iv.

5



Scheme 2


10 Scheme 3 illustrates that amide bridged compound of Formula I, may be prepared by the treatment of 3-i with an appropriately activated diacid in the presence of a base to give 3-ii. Deprotection of PG^4 provides compounds of general formula 3-iii. Activation of the diacid may include the use of active esters, acid chlorides, acid bromides, succinamide esters, HOBt esters, and the use of other reagents used in the formation of amide bonds.



L = $-(CH_2)_r$, $-(CH_2)_r-Y-(CH_2)_r$, -alkyl-aryl-alkyl-, -alkyl-heteroaryl-alkyl-, cycloalkyl, aryl or heteroaryl, wherein r is 1-10


Scheme 3

Scheme 4 illustrates alkyl bridged compounds, which may be prepared using the methods described herein. Treatment of 3-i with 0.5 equiv of an alkyl chain containing two leaving groups, such as 1,5-dibromopentane, 1,10-dibromodecane, and the like, provides intermediate 4-i. Alternatively, reductive amination of 3-i with a dialdehyde may yield intermediate 4-i. Deprotection of PG⁴ yields compounds of formula 4-ii.




Scheme 4

- 5 Scheme 5 illustrates one method wherein two BIR binding units may be bridged via a bis-acetylenic bridging unit. Coupling of 5-i with 5-ii provides a mixture of the symmetrically bridged intermediates 5-iii and 5-v, and the asymmetrical intermediate 5-iv. Separation of 5-ii, 5-iii, and 5-v, may be afforded by methods such as chromatography or recrystallization. Deprotection of intermediates 5-iii, 5-iv, and 5-v, either independently or
- 10 as a combined mixture, provides compounds 5-vi, 5-vii, and 5-viii.



Scheme 5: Coupling of intermediates 5-i and 5-ii.

An additional method for the preparation of asymmetrically bridged compounds is illustrated in Scheme 6. Coupling of 5-i with 6-i will provide a mixture of intermediates 6-ii, 5-iii and 6-iii. Separation of 6-ii, 5-iii, and 6-iii, may be afforded by methods such as chromatography or recrystallization. Deprotection of PG^4 of intermediates 6-iii provides compound 6-iv.




Scheme 6: Coupling of intermediates 5-i and 6-i.

- 5 An alternative strategy involves bridging two BIR binding units via a bis-amide bridging group such as illustrated in Schemes 7 and 8.

The mono-protected bridging group $\text{HO}_2\text{C-L-CO}_2\text{PG}^5$ is activated with peptide coupling agents and subsequently treated with intermediate 3-i to provide intermediate 7-ii.

- 10 Deprotection of PG^5 provides intermediate 7-iii. Treatment of 7-iii with peptide coupling

agents, followed by 7-iv, provides 7-v. Deprotection of PG⁴ and PG⁴⁰⁰ provides compound 7-vi.




5

Scheme 7: P3-P3 bis-amide bridged compounds

A similar process may be applied to the preparation of asymmetrically bridged BIR binding units, which have been bridged between P2 and P3, as illustrated in scheme 8.


10

Treatment of 8-i with a cyclic anhydride, such as succinic or glutaric anhydride, provides intermediate 8-ii. Treatment of 8-ii with amide coupling agents followed by 3-i provides intermediate 8-iii. Deprotection of PG⁴ provides compound 8-iv.



Scheme 8: P2-P3 bis-amide bridged compounds

- 10 Scheme 9 illustrates the synthesis of compound 1. Boc-cis-2-hydroxy-L-proline was treated with NaH in DMF, followed by propargyl bromide to provide intermediate 9-1. Amide coupling with (R)-1,2,3,4-tetrahydro-1-naphthylamine, and Boc deprotection with TFA provided intermediate 9-2. Amide coupling of Boc-tert-BuGly-OH with 9-2, followed by Boc deprotection with TFA provided intermediate 9-3. Amide coupling of Boc-MeAla-OH with intermediate 9-3 provided intermediate 9-4. Homo-coupling of the acetylene groups of 9-4 using a CuI/TMEDA catalyst under O₂, yielded intermediate 9-5, which was deprotected using 4N HCl in 1,4-dioxane, to provide compound 1•2HCl.
- 15




Scheme 9

5

Intermediate 10-6 was used in the preparation of compound 2 and 3 (see Schemes 10 to 12). Intermediate 10-5, was prepared using a similar procedure as described for


intermediate 9-4, using Fmoc-AMPC(2S, 4S)-OH in the first step. Removal of the Fmoc protection group using a base such as 20% morpholine, in a solvent such as THF, provided intermediate 10-6.



Scheme 10


Treatment of intermediate 10-6 with 0.5 equiv of sebacoyl chloride in THF provided 11-1.

- 10 Removal of the Boc protecting groups using 4N HCl in 1,4-dioxane provided compound 2·2HCl (Scheme 11).




Scheme 11


- 5 Similarly, treatment of intermediate 10-6 with 0.5 equiv of terephthaloyl chloride in THF provided 12-1. Removal of the Boc protecting groups using 1N HCl in 1,4-dioxane provided compound 3•2HCl (Scheme 12).



Scheme 12


- 5 Scheme 13 illustrates the preparation of compounds 4 and 5. Intermediates 9-4 and 13-1 were coupled using a CuCl and TMEDA catalysts system, in acetone, under an oxygen atmosphere, to provide a mixture of intermediates 9-5, 13-2, and 13-3. Separation by silica gel chromatography provided the individual intermediates. Intermediates 13-2 and 13-3 were separately deprotected by treatment with 4N HCl in 1,4-dioxane, to provide
- 10 compounds 4•2HCl and 5•2HCl, respectively.








Scheme 13: Synthesis of compounds 4 and 5.

Scheme 14 illustrates the preparation of compound 6. Intermediates 9-4 and 14-1 were coupled using a CuCl and TMEDA catalyst system, in acetone, under an oxygen atmosphere. Intermediate 14-2 was isolated from the resulting mixture by silica gel chromatography. Intermediate 14-2 was deprotected by treatment with 4N HCl in 1,4-dioxane, to provide compound 6•2HCl.

**Scheme 14:**

- Scheme 15 illustrates the preparation of P2-P3 bridged compounds 7 and 8. Intermediate 15-1 was treated with glutaric anhydride to provide intermediate 15-2. Treatment of 15-2 with HBTU, HOBT, and DIPEA, followed by the addition of 10-6 in DMF, provided intermediate 15-3. Removal of the Cbz protection group using H_2 over Pd/C yielded intermediate 15-4. Acylation of 15-4 with Boc-N-MeAla-OH provided intermediate 15-5, which was Boc deprotected using 4N HCl in $1,4\text{-dioxane}$, provided compound 7•2HCl.






Scheme 15

Removal of the Fmoc protection group from compound 7·2HCl, using polystyrene supported N-methylpiperazine in DMF provided compound 8. Removal of the Boc


protection group from intermediate 15-3, using 4N HCl in 1,4-dioxane, provided compound 9•HCl (see Scheme 16).




5

Scheme 16

Scheme 17 illustrates the syntheses of compounds bridged by sulfonamide linkages. Treatment of 3-i with a disulfonyl chloride reagent provides intermediate 17-i. Deprotection of PG⁴ provides compound 17-ii.


**Scheme 17**

Similarly, treatment of 3-i with a bis-isocyanate provides intermediate 18-i, which after
 5 deprotection of PG^4 yields compound 18-ii.




Scheme 18

- Schemes 19a and 19b describe an alternate route to compounds 3-iii, 17-ii or 18-ii, where
- 5 A=CO and Q=NR⁴R⁵. Protected amino-proline derivative 19-1 is treated with LG-L-LG to provide intermediate 19-2 which is then deprotected at PG¹ to yield intermediate 19-3. Intermediate 19-3 is converted intermediate 19-5 by an amino acid coupling/deprotection sequences as described earlier. A third amino acid coupling step converts intermediate 19-5 to intermediate 19-6. Deprotection of 19-6 at PG² yields the diacid intermediate 19-7.
 - 10 Treatment of 19-7 with amino acid coupling reagents, followed R⁴R⁵NH, yields intermediate 19-8, which upon deprotection of PG⁴ provided compound 19-9.




Scheme 19a



Scheme 19b

- 5 This method can be applied to the synthesis of bridged bis-proline amides, sulfonamids and ureas. For example, in the case where the bridging group includes an Ar moiety, X-L-X = -NHS(O)₂-Ar-S(O)₂NH-, NHC(O)-Ar-C(O)NH-, or -NHC(O)NH-Ar-NHC(O)NH-, -O-CH₂ArCH₂-O-.
- 10 Alternatively, the proline derivative 19-2 may be deprotected at PG² and converted to the amide intermediate 20-3. Following a similar procedure as described above, 20-3 can be converted to compound 19-9.





Scheme 20

- 5 The synthesis of compound 20 is illustrated in Scheme 21. N-Boc-cis-4-amino-L-proline methyl ester, 21-1, was treated with terephthaloyl chloride to provide intermediate 21-2, which was further deprotected with TFA to yield intermediate 21-3. Intermediate 21-3 was coupled to Boc-tert-BuGyl-OH, 21-4, using HBTU and HOBt, followed by TFA deprotection, to provide intermediate 21-6. Intermediate 21-6 was coupled to Boc-N-
- 10 MeAla-OH, 21-7, using HBTU and HOBt to provide intermediate 21-8. Saponification of


the methyl ester using LiOH provided intermediate 21-9. Coupling of intermediate 21-9 with phenethylamine using HBTU and HOBT provided intermediate 21-10 which was deprotected using HCl in 1,4-dioxane to provide compound 20•2HCl.

5




**Scheme 21**

This method was used for the preparation of a number of proline amide derivatives such as compounds 19 to 24 and 46 to 51. This is shown in Scheme 22 wherein, intermediate 21-9 was coupled to 2-propylamide, followed by HCl deprotection to yield compound 22•2HCl, while coupling of intermediate 21-9 to 1,1-diphenylmethylethylamine followed by deprotection with HCl provided compound 24•2HCl.


**Scheme 22**

The preparation of pyrrolidine derivatives are described in Scheme 23. The ester moiety
 5 of intermediate 21-8 was reduced to alcohol 23-1 which was subsequently oxidized to
 aldehyde 23-2. Reductive amination using phenethylamine provided intermediate 23-4.
 Acylation of 23-4 with either acetyl chloride or benzoyl chloride provided 23-5 and 23-6,
 respectively. Deprotection with 1N HCl in 1,4-dioxane provided compounds 25•2HCl and

27•2HCl.





Scheme 23a

**Scheme 23b**

Sulfonylation of 23-4 with methanesulfonyl chloride, followed by deprotection with 1N HCl
 5 in 1,4-dioxane provided compound 26 $\cdot 2\text{HCl}$.


Intermediate 24-1 provided an useful template for the preparation of ether bridged
 compounds. Intermediate 24-1 was prepared from N-Boc-cis-4-hydroxy-L-proline and
 α,α' -dibromo-p-xylene as described below. Amide coupling and deprotection of the Boc
 10 protecting groups using TFA provided intermediate 24-3. Two sequential amino acid
 coupling and deprotection steps, as described above, provided compound 29 $\cdot 2\text{HCl}$.






Scheme 24

Intermediate 10-6 was bridged using 1,3-benzenedisulfonyl chloride to provide
 5 intermediate 25-1. Deprotection with 4N HCl in 1,4-dioxane provided compound 33 as its
 bis-HCl salt.



Scheme 25

Similarly, intermediate 10-6 was bridged using 1,4-phenylene diisocyanate to provide intermediate 26-1, which upon deprotection with TFA provided compound 44 as is bis-TFA salt.




5

Scheme 26


The preparation of compound 35 is illustrated in Scheme 27. Intermediate 10-1 was deprotected by the treatment with piperide in THF to provide intermediate 27-1. Coupling of 27-1 with Cbz-Gly-OH followed by Boc deprotection provided intermediate 27-3•TFA. Coupling of 27-3 with Boc-t-BuGly-OH followed by Boc deprotection provided intermediate 27-5•TFA. Coupling of 27-5•2TFA with Boc-NMeAla-OH provided intermediate 27-6. Removal of the Cbz group using H₂ and Pd/C yielded 27-7 which was bridged using terephthaloyl chloride to provide intermediate 27-8. Deprotection of intermediate 27-8 using 4N HCl in 1,4-dioxane provided compounds 35•2HCl.

10

15




Scheme 27a



Scheme 27b

- Various chiral amines were prepared from optically active (R)-2-hydroxy-1-phenylethylamine, 28-1. Intermediate 28-1 was Boc protected to provide 28-2. The alcohol moiety of 28-2 was alkylated with various alkyl halides to provide intermediates 28-3, 28-4, and 28-5. Deprotection of intermediates 28-3, 28-4 and 28-5 with HCl yielded the chiral amines 28-6, 28-7, and 28-8.




Scheme 28

Intermediates 28-6, 28-7, and 28-8 were coupled with intermediate 21-9 in a manner
 5 similar to that illustrated for compounds 22 and 24 (see Scheme Scheme 22) to provide
 compounds 63, 64 and 65, respectively.

Other chiral amines may be obtained commercially or prepared via a number of methods
 including the conversion or interconversion of ketones, alcohols or azides to amines, using
 10 chiral or achiral chemistries, and the chiral resolution of isomers via methods known in the
 art, such as chromatography or crystallization.


For example, Scheme 29 illustrates the enantioselective synthesis of optically enriched,
 asymmetric, 1,1-diphenylmethanamines prepared by the addition of aryl boronic acids to
 15 chiral sulfinimines, followed by hydrolysis of the sulfinamines with acid to provide the
 optically enriched amine intermediates characterized by intermediates 29-4 and 29-5
 (Bolshan, Y.; Batey, R. A., Org. Letters 2005, 7, 1481-1484).



Scheme 29

- 5 Intermediates 29-4 and 29-5 were coupled with intermediate 21-9 in a manner similar to that illustrated for compounds 22 and 24 (see Scheme Scheme 22) to provide compounds 66 and 67, respectively.

Several methods exist for the conversion of tetralones to chiral 1,2,3,4-tetrahydronaphthyl-
 10 1-amines, for example, the method report by R. A. Stalker, et al., Tetrahedron 2002, 58, 4837, as summarized below:



- 15 These chiral amines can be incorporated into compounds of the instant invention using similar methods as illustrated in Schemes 21 and 22.

The above Schemes are applicable to both symmetrical compounds and unsymmetrical compounds of the present invention. The substituents A¹, A, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, and the like are as defined herein. LG is a leaving group such as, for example, Cl, Br, I, OTs, OSu or OMs.

20

EXAMPLES

The following abbreviations are used throughout:

5	Boc:	<i>t</i> -butoxycarbonyl;
	CBz:	benzyloxycarbonyl;
	DCM:	dichloromethane, CH ₂ Cl ₂ ;
	DIPEA:	diisopropylethylamine;
10	DMAP:	4-(dimethylamino)pyridine;
	DMF:	N,N-dimethylformamide;
	DTT:	dithiothreitol;
	EDC:	3-dimethylaminopropyl-3-ethylcarbodiimide hydrochloride;
	EDTA:	ethylenediaminetetracetic acid;
15	Fmoc:	N-(9-fluorenylmethoxycarbonyl);
	HBTU:	O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate;
	HCl:	hydrochloric acid;
	HOAc:	acetic acid;
	HOBt:	1-hydroxybenzotriazole;
20	HPLC:	high performance liquid chromatography;
	LCMS:	liquid chromatography-mass spectrometer;
	MeOH:	methanol;
	MgSO ₄ :	magnesium sulfate;
	MS:	mass spectrum;
25	NaHCO ₃ :	sodium hydrogen carbonate;
	Pd/C:	palladium on carbon;
	TEA:	triethylamine;
	THF:	tetrahydrofuran; and
	TMEDA:	N,N,N,N-tetramethylethylenediamine.

30

SYNTHETIC METHODS

Synthesis of compound 1

35 Step 1: Intermediate 9-1

NaH (440 mg, 11.0 mmol) was suspended in dry DMF (5 mL) under N₂ at 0°C. Boc-cis-2-hydroxyl-L-proline (1.00 g, 4.0 mmol) was dissolved in dry DMF (15 mL) and added

dropwise to the suspension of NaH. The mixture was left to stir at 0 °C for 10 min. Propargyl bromide (560 µL, 5.0 mmol) was added dropwise to the solution. The mixture was stirred at 0 °C for 1 h then quenched with H₂O. The contents were added to a separatory funnel along with EtOAc and 10% citric acid (until pH ~2). The organic layer
5 was collected, dried and concentrated under reduced pressure. Flash chromatography (silica, hexanes/THF) yielded intermediate 9-1 as clear oil. MS (m/z) M+Na =292.

Step 2: Intermediate 9-2

Intermediate 9-1 (560 mg, 2.1 mmol), HOBt (444 mg, 2.9 mmol), EDC (563 mg, 2.9
10 mmol) and DIPEA (1.46 mL, 8.4 mmol) were dissolved in dry dichloromethane (10 mL) under N₂ and stirred for 10 min at room temperature. 1,2,3,4-(R)-Tetrahydronaphthyl-1-amine (368 µL, 2.5mmol) was then added and the solution was left to stir for 24 h at RT. The contents were then added to a separatory funnel along with EtOAc and washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine. The organic layer was collected,
15 dried and concentrated under reduced pressure. The product was treated with 5 ml of 50% CH₂Cl₂/TFA for 1 hr at room temperature. Volatiles were removed in vacuo and the residue triturated with diethyl ether to provide intermediate 9-2·TFA. MS (m/z) M+1 = 299.

Step Three: Intermediate 9-3

Boc-t-Bu-Gly-OH (484 mg, 2.1 mmol), HOBt (375 mg, 2.4 mmol), HBTU (910 mg, 2.4
20 mmol) and DIPEA (1.20 mL, 7 mmol) were dissolved in dry DMF (10 mL) under N₂ and stirred for 10 min at room temperature. **Intermediate 9-2** (720 mg, 1.75 mmol) was then added and the solution was left to stir for 24 h at room temperature. The contents were then added to a separatory funnel along with EtOAc and washed with 10% citric acid (2x),
25 saturated NaHCO₃ (2x) and brine. The organic layer was collected, dried and concentrated under reduced pressure. The resulting product was treated with 10 mL of 50% CH₂Cl₂/TFA for 1 hr at room temperature. Volatiles were removed in vacuo and the residue triturated with diethyl ether to provide intermediate 9-3·TFA. MS (m/z) M+1 = 412.

Step 4: Intermediate 9-4

Boc-N-Me-Ala-OH (278 mg, 1.37 mmol), HOBt (227 mg, 1.49 mmol), EDC (293 mg, 1.49
30 mmol) and DIPEA (796 µL, 4.6 mmol) were dissolved in dry dichloromethane (10 mL) under N₂ and stirred for 10 min at RT. **Intermediate 9-3·TFA** (600 mg, 1.14 mmol) was then added and the solution was left to stir for 24 h at RT. EtOAc was added and the

organic layer was washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine, dried over anhydrous MgSO₄, filtered and volatiles removed under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100 % hexanes/THF, to provide intermediate 9-4. MS (m/z) M+Na=619.

5

Step 5: Compound 1

Intermediate 9-4 (100 mg, 0.17 mmol), CuCl (25 mg, 0.25 mmol) and N,N,N,N-tetramethylethylenediamine (38 μ L, 0.25mmol) were suspended in dry acetone (5 ml) and stirred at room temperature under an O₂ atmosphere for 72 h. EtOAc was added and the organic layer was washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine, dried over anhydrous MgSO₄, filtered and volatiles removed under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100 % hexanes/THF, to provide intermediate 9-5. Treatment of 9-5 with 4N HCl in 1,4-dioxane at 0°C, for 2 hrs, and removal of volatiles under reduced pressure provided compound 1 as its bis-HCl salt. MS (m/z) M+1=991.7.

10

15

Synthesis of intermediate 10-6**Step 1: Intermediate 10-1**

Fmoc-AMPC(2S, 4S)-OH (900 mg, 2.0 mmol), HBTU (1.14 g, 3.0 mmol), and HOBt (415 mg, 3.0 mmol) were dissolved in dry DMF (10 mL) and treated with DIPEA (1050 μ L, 6.0 mmol). The mixture was stirred for 10 minutes before the addition of (R)-1,2,3,4-tetrahydronaphthyl-1-amine (330 μ L, 2.2 mmol). The reaction was stirred overnight before being diluted with ethyl acetate (100 mL) and 10% citric acid (50 mL). The organic layer was separated and washed with 10% citric acid (2 x 50 mL), saturated sodium bicarbonate (3 x 25 mL) and brine (1 x 20 mL), before being dried over anhydrous magnesium sulphate, filtered, and concentrated under reduced pressure to provide crude 10-1 a white solid. This material was 95% pure by LCMS and was advanced to the next step without further purification.

20

25

30

Step 2: Intermediate 10-2

Intermediate 10-1 was dissolved in methylene chloride (10 mL) and treated with TFA (10 mL). The solution was stirred at room temperature for 2 hrs before the volatiles were removed under reduced pressure. The resulting oil was stirred with diethyl ether (25 mL) to provide a light brown solid which was filtered and washed with diethyl ether (2 x 5 mL), to provide compound 10-2 •TFA as a light brown solid (1.17 g).

35

Step 3: Intermediate 10-3

Boc-t-BuGly-OH (460 mg, 2.0 mmol), HBTU (760 mg, 2.0 mmol), and HOBt (270 mg, 2.0 mmol) and DIPEA (765 μ L, 7.5 mmol) were dissolved in dry DMF (10 mL) and the reaction
5 was stirred at room temperature for 10 minutes before intermediate 10-2 (867 mg, 1.5 mmol) was added. The mixture was stirred overnight before being diluted with ethyl acetate (200 mL) and 10% citric acid (50 mL). The organic layer was separated and washed with 10% citric acid (2 x 50 mL), saturated sodium bicarbonate (3 x 25 mL) and brine (1 x 20 mL), before being dried over anhydrous magnesium sulphate, filtered, and
10 concentrated under reduced pressure to provide crude 10-3 a white solid (920 mg, > 95% pure by LCMS) and was advanced to the next step without further purification.

Step 4: Intermediate 10-4

Intermediate 10-3 was dissolved in methylene chloride (10 mL) and treated with TFA (10
15 mL). The solution was stirred at room temperature for 2 hrs before the volatiles were removed under reduced pressure. The resulting oil was stirred with diethyl ether (20 mL) to provide a light brown solid which was filtered, wash with diethyl ether (2 x 5 mL), to provide compound 10-4•TFA a white solid.

Step 5: Intermediate 10-5

Boc-MeAla-OH (308 mg, 1.52 mmol), HBTU (450 mg, 1.91 mmol), and HOBt (260 mg, 1.91 mmol) were dissolved in dry DMF (10 mL). DIPEA (886 μ L, 5.0 mmol) was added and the reaction was stirred at room temperature for 10 minutes before intermediate 10-4
(900 mg, 1.27 mmol) was added. The mixture was stirred overnight before being diluted
25 with ethyl acetate (200 mL) and 10% citric acid (50 mL). The organic layer was separated and washed with 10% citric acid (2 x 50 mL), saturated sodium bicarbonate (3 x 25 mL) and brine (1 x 20 mL), before being dried over anhydrous magnesium sulphate, filtered, and concentrated under reduced pressure to provide crude 10-5 a white solid (0.87 mg, 95.5% pure by LCMS) and was advanced to the next step without further purification.

Step 6: Intermediate 10-6

Intermediate 10-5 was dissolved in 20 % morpholine/THF (10 mL) and the solution was stirred at room temperature for 16 hrs. Volatiles were removed under reduced pressure to provide compound 10-6 a white solid. Further purification by silica gel chromatography
35 provide 10-6 which was 95 % pure LCMS (MS (m/z) M+1 = 617.4.

Synthesis of compound 2**Step 1: Intermediate 11-1**

Crude 10-6 (200 mg, 0.251 mmol) was dissolved in THF (5 mL) and cooled to in an ice
5 bath. DIPEA (50 μ L, 0.275 mmol) and sebacoyl chloride (26 μ L, 0.125 mmol) were added
and the reaction was stirred for 4 hours at room temperature before being diluted with
ethyl acetate (20 mL) and saturated sodium bicarbonate. The organic layer was
separated, washed with brine, dried over anhydrous magnesium sulphate, filtered and the
solvent removed under reduced pressure. The resulting crude solid was purified by silica
10 gel chromatography, eluting with a 10-100% hexane/THF gradient, to provide 11-1 as a
white solid (55 mg).

Step 2: Compound 2

Intermediate 11-1 (50 mg) was treated with 4N HCl in 1,4-dioxane (3 mL) and stirred for 3
15 hrs. Volatiles were removed under reduced pressure and the resulting solid was triturated
with diethyl ether (3 x 5 mL). The resulting solid was dried under reduced pressure to
provide compound 2•2HCl as an off white solid (30 mg). MS (m/z) (M+2)/2=541.4.

Compound 3**Step 1: Intermediate 12-1**

Crude 10-6 (200 mg, 0.251 mmol) was dissolved in THF (5 mL) and cooled to 4 °C on an
ice bath. DIPEA (50 μ L, 0.275 mmol) was added. Terephthaloyl chloride (26 μ L, 0.125
mmol) was added and the reaction was stirred for 16 hours before being diluted with ethyl
25 acetate (20 mL) and saturated sodium bicarbonate. The organic layer was separated,
washed with brine, dried over anhydrous magnesium sulphate, filtered and the solvent
removed under reduced pressure. The resulting crude solid was purified by silica gel
chromatography, eluting with a 25-100% hexanes/THF gradient, to provide 12-1 as a
white solid (90 mg).

Step 2: Compound 3

Intermediate 12-1 (90 mg) was treated with 4N HCl in 1,4-dioxane (3 mL) and stirred for 3
hrs. Volatiles were removed under reduced pressure and the resulting solid was triturated
with diethyl ether, filtered and washed with diethyl ether (3 x 5 mL). The resulting solid
was dried under reduced pressure to provide compound 3•2HCl as an off white solid (40
35 mg). MS (m/z) (M+2)/2=523.4.

Compounds 4 and 5

Intermediate 9-4 (130 mg, 0.21 mmol) and intermediate 13-1 (130 mg, 0.22 mmol), CuCl (60 mg, 0.6mmol) and tetramethylethylenediamine (90 μ L, 0.6mmol) were suspended in dry acetone (15 mL) and stirred at room temperature under an O₂ atmosphere for 120 h. EtOAc was added and the organic layer was washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine, dried over anhydrous Mg₂SO₄, filtered, and concentrated under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100% hexanes/THF, to provide intermediates 9-5, 13-2 and 13-3. Intermediates 13-2 and 13-3 were independently treated with 4N HCl in 1,4-dioxane. Volatiles were removed and the resulting solids triturated with diethyl ether, filtered and washed with diethyl ether, to provide compounds 4 and 5, respectively, as their bis-HCl salts.

Compound 4·2HCl: MS (m/s) M+1=993.5.

Compound 5·2HCl: MS (m/z) M+1=995.6.

Compound 6

Intermediate 9-4 (250 mg, 0.400 mmol), intermediate 14-1 (560 mg, 0.900 mmol), CuCl (267 mg, 2.7 mmol) and tetramethylethylenediamine (405 μ L, 2.7 mmol) were suspended in dry acetone (10 mL) and stirred at room temperature under an O₂ atmosphere for 72 h. Celite was added and the mixture filtered through a pad of celite. EtOAc was added to the filtrate and the resulting organic layer was washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine, dried over anhydrous Mg₂SO₄, filtered, and concentrated under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100% hexanes/THF, to provide intermediate 14-2. Intermediate 14-2 was treated with 4N HCl in 1,4-dioxane. Volatiles were removed and the resulting solids triturated with diethyl ether, filtered and washed with diethyl ether, to provide compound 6 as its bis-HCl salts. MS (m/s) (M+2)/2=514.4.

Compound 7**Step 1:**

Intermediate 15-1 (1.0 g, 1.2 mmol), glutaric anhydride (190 mg, 1.7 mmol) and DIPEA (836 μ L, 4.8 mmol) were dissolved in dichloromethane (20 mL) under N₂ at 0 °C. A catalytic amount of DMAP was added and the solution was stirred for 30 min on ice and 24 hrs at room temperature. EtOAc was added to the filtrate and the resulting organic layer was washed with 10% citric acid (2x), saturated NaHCO₃ (2x) and brine, dried over

anhydrous Mg_2SO_4 , filtered, and concentrated under reduced pressure to provide intermediate 15-2 as a white solid.

Step 2:

- 5 Intermediate 15-2 (420 mg, 0.5 mmol), HOBt (85 mg, 0.63 mmol), HBTU (222 mg, 0.60 mmol) and DIPEA (313 μL , 1.8 mmol) were dissolved in dry DMF (5 ml) under N_2 and stirred for 10 min at RT. Intermediate 10-6 (250 mg, 0.45 mmol) was added and the solution was left to stir for 24 hrs at room temperature. EtOAc was added and the resulting organic layer was washed with 10% citric acid (2x), saturated NaHCO_3 (2x) and brine,
10 dried over anhydrous Mg_2SO_4 , filtered, and concentrated under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100% hexanes/THF, to provide intermediate 15-3.

Step 3:

- 15 Intermediate 15-3 (240 mg, 0.17 mmol), 10 wt% Pd/C (50 mg, 50% H_2O), were suspended in MeOH (10 mL) and stirred for 24 h at room temperature under a H_2 atmosphere (1 atm). The contents were then filtered on celite and washed with MeOH. The filtrate was concentrated under reduced pressure to provide the intermediate 15-4 as a white solid.

Step 4:

- 20 BocNMe-Ala-OH (57 mg, 0.28 mmol), HOBt (44 mg, 0.33 mmol), HBTU (116 mg, 0.31 mmol) and DIPEA (90 μL , 0.52 mmol) were dissolved in dry DMF (5 mL) under N_2 and treated with intermediate 15-4 (160 mg, 0.13 mmol). After stirring for 24 h at room temperature EtOAc was added and the resulting organic layer was washed with 10% citric
25 acid (2x), saturated NaHCO_3 (2x) and brine, dried over anhydrous Mg_2SO_4 , filtered, and concentrated under reduced pressure. The product was purified by silica gel chromatography, eluting with 10-100% hexanes/THF, to provide intermediate 15-5. Intermediate 15-5 was stirred with 4N HCl in 1,4-dioxane for 1 hr at room temperature. Volatiles were removed under reduced pressure to provide compound 7 as its HCl salt.
30 MS (M+2)/2 = 618.0.

Compound 8:

- Compound 7•HCl (100 mg, 0.08 mmol) is dissolved in DMF (1 mL) and added to a CH_2Cl_2 (5 mL) suspension of piperazinomethyl polystyrene resin (0.86 mmol/g, 356 mg, 0.3
35 mmol). After shaking at room temperature for 48 hrs an additional 200 mg of resin was

added. The mixture was shaken for 20 days at room temperature before being, filtered, washing with MeOH. Volatiles were removed under reduced pressure to provide compound 8 a white solid. MS (m/z) $M+1=1012$.

5 **Compound 9:**

Intermediate 15-3 (10 mg) was treated with 4N HCl in 1,4-dioxane for 1 hr. Volatiles were removed under reduced pressure to provide compounds 9 as its HCl salt. MS (m/s) $(M+2)/2 = 642$.

10 Compounds 10, 11, 12, 13, 14, 15, 16, 39, 40, 41, 42, 43, 52, 55, 56, 57, and 58 and were prepared in a similar manner to that described for compound 2, wherein the corresponding sulfonyl chloride or activated diacid was substituted for sebacoyl chloride in Step 1. MS characterization of these compounds is summarized in Table 1.

15 Compound 17 was prepared in a similar manner to that described for compound 29 using 1,2,3,4-(R)-tetrahydronaphthyl-1-amine in place of phenethylamide. MS (m/s) $(M+2)/2 = 510.4$.

Compound 18:

20 Intermediate 21-8 was stirred in 4N HCl in 1,4-dioxane for 2 hrs. Volatiles were removed in vacuo and the residue was triturated with diethyl ether to provide compound 18•2HCl as a white solid. MS (m/z) $M+1=815.4$.

25 Compounds 19, 21, 22, 23, 24, 31, 32, 46, 47, 48, 49, 50, 51, 54, 59, 60, 61, 63, 64, 65, and 67 were prepared in a similar manner to that described for compound 20, wherein phenethylamine was substituted by the corresponding amine in Step 6. MS characterization of these compounds is summarized in Table 1.

Compound 20

30 **Step 1: Intermediate 21-2**

To a solution of N-Boc-cis-4-amino-L-proline methyl ester, 21-1, (10.0 g, 35.61 mmol) in CH_2Cl_2 cooled to 0 °C were sequentially added TEA (14.88 mL, 106.80 mmol), DMAP (10 mg) and terephthaloyl chloride (3.61 g, 17.80 mmol) and the reaction was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was
35 separated, washed with 10 % citric acid, aqueous NaHCO_3 and brine, dried over

anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 21-2 as a white solid.

Step 2: Intermediate 21-3•2TFA

5 Intermediate 21-2 (4.80 g, 7.76 mmol) was dissolved in a mixture of CH_2Cl_2 (40 mL) and TFA (40 mL) at 0 °C. The solution was stirred for 4 hrs at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 21-3•2TFA as a white solid. MS (m/z) $M+1=419.2$

10 **Step 3: Intermediate 21-5**

To a solution of Boc- α -tBuGly-OH, 21-4, (3.95 g, 17.07 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (13.5 mL, 77.6 mmol), HOBt (2.62 g, 19.4 mmol) and HBTU (7.36 g, 19.4 mmol). After stirring for 10 minutes intermediate 21-3•2TFA (5.02 g, 7.76 mmol) was added and the reaction mixture was stirred overnight at room temperature.

15 Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO_3 and brine, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 21-5 as a white solid.

20 **Step 4: Intermediate 21-6•2TFA**

Intermediate 21-5 (6.55 g, 7.76 mmol) was dissolved in a mixture of CH_2Cl_2 (40 mL) and TFA (40 mL) at 0 °C. The solution was stirred for 3 hrs at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 21-6•2TFA as a white solid. MS (m/z) $M+1=645.4$

25

Step 4: Intermediate 21-8

To a solution of Boc-NMe-Ala-OH, 21-7, (3.15 g, 15.51 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (12.3 mL, 70.5 mmol), HOBt (2.38 g, 17.63 mmol) and HBTU (6.69 g, 17.63 mmol). After stirring for 10 minutes intermediate 21-6•2TFA (6.15 g, 7.05 mmol) was added and the reaction mixture was stirred overnight at room temperature.

30 Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO_3 and brine, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 21-8 as a white solid.

35

Step 5: Intermediate 21-9

To a solution of intermediate 21-8 (6.20 g, 6.11 mmol) in THF (80 mL) and MeOH (8 mL) cooled to 0 °C was added 1N aqueous LiOH (30.5 mL) and the reaction was stirred overnight at room temperature. PH was adjusted to 6 with 10% citric acid and ethyl acetate was added, the organic layer was separated, washed with brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo to provide intermediate 21-9 as a white solid.

Step 6: Intermediate 21-10

To a solution of intermediate 21-9 (150 mg, 0.15 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (265 uL, 1.52 mmol), HOBt (51 mg, 0.38 mmol) and HBTU (144 mg, 0.38 mmol). After stirring for 10 minutes phenethylamine (42 uL, 0.33 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 21-10 as a white solid.

Step 7: Compound 20•2HCl

4N HCl in 1,4-dioxane (3.0 mL) was added to intermediate 21-10 (145 mg, 0.12 mmol) and the solution was stirred for 1 hr at 0 °C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 20•2HCl as a white solid. MS (m/z) (M+2)/2=497.6.

Compound 22•2HCl**Step 1: Intermediate 22-1**

To a solution of intermediate 21-9 (75 mg, 0.08 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (135 uL, 0.76 mmol), HOBt (26 mg, 0.19 mmol) and HBTU (72 mg, 0.19 mmol). After stirring for 10 minutes isopropylamine (14 uL, 0.17 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 22-1 as a white solid.

Step 2: Compound 22•2HCl

4N HCl in 1,4 dioxane (2.0 mL) was added to intermediate 22-1 (58 mg, 0.05 mmol) and the solution was stirred for 2 hrs at 0 °C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 22•2HCl as a white solid. MS (m/z) (M+2)/2=435.4.

Compound 24•2HCl**Step 1: Intermediate 22-2**

To a solution of intermediate 21-9 (600 mg, 0.61 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (1.0 mL, 6.08 mmol), HOBt (205 mg, 1.52 mmol) and HBTU (576 mg, 1.52 mmol). After stirring for 10 minutes aminodiphenylmethane (230 uL, 1.34 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 22-2 as a white solid.

Step 2: Compound 24•2HCl

4N HCl in 1,4 dioxane (1.9 mL) was added to intermediate 22-2 (660 mg, 0.50 mmol) and the solution was stirred for 1 hr at 0 °C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 24•2HCl as a white solid. MS (m/z) (M+2)/2=435.4.

Compound 25**Step 1: Intermediate 23-1**

To a solution of intermediate 21-8 (1.10 g, 1.08 mmol) in THF cooled to 0 °C was added lithium borohydride (118 mg, 4.86 mmol) and the reaction mixture was stirred for 3 hrs at room temperature. Ethyl acetate and 10% citric acid were added. The organic layer was separated, washed with aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 23-1 as a white solid. MS (m/z) M+1=959.6

Step 2: Intermediate 23-2

To a solution of intermediate 23-1 (284 mg, 0.29 mmol) in CH₂Cl₂ was added Dess Martin periodinane (314 mg, 0.74 mmol) and the reaction was stirred for 5 hrs at room

temperature. Aqueous NaHCO_3 was added, the organic layer was separated, washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 23-2 as a white solid.

5 **Step 3: Intermediate 23-4**

To a solution of intermediate 23-2 (282 mg, 0.29 mmol) in CH_2Cl_2 was added phenethylamine (82 μL , 0.65 mmol). After stirring for 30 min at room temperature sodium triacetoxyborohydride (280 mg, 1.32 mmol) was added and the reaction mixture was stirred for 2 hrs. Saturated aqueous NaHCO_3 was added, the organic layer was separated, washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 23-4 as a white solid. MS (m/z) $M+1=1165.8$

15 **Step 4: Intermediate 23-5**

To a solution of intermediate 23-4 (100 mg, 0.08 mmol) in CH_2Cl_2 cooled to 0°C were sequentially added triethylamine (60 μL , 0.43 mmol) and acetyl chloride (14 μL , 0.19 mmol). The reaction was stirred for 2 hrs at room temperature before water and ethyl acetate were added. The organic layer was separated, washed with 10 % citric acid, aqueous NaHCO_3 and brine, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 23-5 as a white solid.

25 **Step 5: Compound 25•2HCl**

4N HCl in 1,4-dioxane (3 mL) was added to intermediate 23-5 (80 mg, 0.06 mmol) and the solution was stirred for 1 hr at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 25•2HCl as a white solid. MS (m/z) $(M+2)/2=525.6$.

Compound 26•2HCl

30 **Step 1: Intermediate 23-5**

To a solution of intermediate 23-4 (100 mg, 0.08 mmol) in CH_2Cl_2 cooled to 0°C were sequentially added TEA (60 μL , 0.43 mmol) and methanesulfonyl chloride (15 μL , 0.19 mmol) and the reaction was stirred for 2 hrs at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous

NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 23-5 as a white solid.

Step 2: Compound 26•2HCl

- 5 4N HCl in 1,4-dioxane (3 mL) was added to intermediate 23-5 (79 mg, 0.06 mmol) and the solution was stirred for 1 hr at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 26•2HCl as a white solid. MS (m/z) (M+2)/2=561.6.
- 10 Compounds 27, 28 and 30 were prepared in a similar manner to that described for compound 26, wherein for compound 27 and 28 acetyl chloride and benzoyl chloride, respectively, were used in place of methanesulfonyl chloride.
- Compound 27- MS (m/z) (M+2)/2=587.6.
- Compound 28- MS (m/z) (M+2)/2=543.6.
- 15 Compound 30- MS (m/z) (M+2)/2=579.6.

Compound 29

Step 1: Intermediate 24-1

- To a 1.0 M solution of NaHMDS in THF (21.6 mL, 21.6 mmol) cooled to 0 °C was slowly
20 added a solution of N-Boc-cis-4-hydroxy-L-proline (2.50 g, 10.8 mmol) in DMF. After stirring for 20 minutes at 0 °C α,α'-dibromo-p-xylene (1.37 g, 5.0 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, dried over MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 24-1 as a white
25 solid.

Step 2: Intermediate 24-2

- To a solution of intermediate 24-1 (200 mg, 0.35 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (365 uL, 2.1 mmol), HOBt (132 mg, 0.98 mmol) and HBTU
30 (345 mg, 0.91 mmol). After stirring for 10 minutes phenethylamine (107 uL, 0.85 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 24-2 as a white
35 solid.

Step 3: Intermediate 24-3•2TFA

Intermediate 24-2 (260 mg, 0.35 mmol) was dissolved in a mixture of CH₂Cl₂ (3 mL) and TFA (3 mL). The solution was stirred for 1 hr at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 24-3•2TFA as a white solid. MS (m/z) M+1=571.4

Step 4: Intermediate 24-4

To a solution of Boc- α -¹BuGly-OH (256 mg, 1.10 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (361 μ L, 2.10 mmol), HOBt (175 mg, 1.3 mmol) and HBTU (455 mg, 1.2 mmol). After stirring for 10 minutes intermediate 24-3•2TFA (230 mg, 0.46 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 24-4 as a white solid.

Step 5: Intermediate 24-5•2TFA

Intermediate 24-4 (458 mg, 0.46 mmol) was dissolved in a mixture of CH₂Cl₂ (3 mL) and TFA (3 mL). The solution was stirred for 30 minutes at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 24-5•2TFA as a white solid. MS (m/z) M+1=797.6

Step 6: Intermediate 24-6

To a solution of Boc-NMe-Ala-OH (119 mg, 0.58 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (209 μ L, 1.2 mmol), HOBt (91 mg, 0.67 mmol) and HBTU (236 mg, 0.62 mmol). After stirring for 10 minutes intermediate 24-5•2TFA (250 mg, 0.24 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 24-6 as a white solid.

Step 7: Compound 29•2HCl

4N HCl in 1,4-dioxane (1.0 mL) was added to intermediate 24-6 (280 mg, 0.24 mmol) and the solution was stirred for 1 hr at 0 °C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 29•2HCl as a white solid. MS (m/z) M+1=967.6.

5

Compound 33**Step 1: Intermediate 25-1**

To a solution of intermediate 10-6 (258 mg, 0.50 mmol) in DMF were sequentially added DIPEA (217 uL, 1.25 mmol) and 1,3-benzenedisulfonyl chloride (69 mg, 0.25 mmol) and the reaction was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10% citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 25-1 as a white solid.

15 **Step 2: Compound 33•2HCl**

4N HCl in 1,4-dioxane (2 mL) was added to intermediate 25-1 (143 mg, 0.11 mmol) and the solution was stirred for 1 hr at 0°C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 33•2HCl as a white solid. MS (m/z) (M+2)/2=559.5.

20

Compound 34:

Compound 34 was prepared in a similar manner to compound 20, wherein N-Boc-cis-4-amino-L-proline methyl ester was substituted with N-Boc-trans-4-amino-L-proline methyl ester in step 1 and phenethylamine was substituted with 1,2,3,4-(R)-tetrahydronaphthyl-1-amine in Step 6. MS (m/z) M+1=1045.8.

25

Compound 35•2HCl**Step 1: Intermediate 27-1**

Intermediate 10-1 (3.90 g, 6.68 mmol) was dissolved in THF (100 mL) and treated with piperidine (5.0 mL, 68.2 mmol). The solution was stirred for 16 hrs at room temperature before volatiles were removed under reduced pressure to provide a semi-solid which was suspended in MeOH (5 mL) and filtered. The filtrate was concentrated, suspended in MeOH (5 mL), filtered and the filtrate concentrated to provide intermediate 27-1 as a semi-solid (80 % pure).

35

Step 2: Intermediate 27-2

To a solution of carbobenzyloxyglycine (699 mg, 3.34 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (2.40 mL, 13.9 mmol), HOBt (488 mg, 3.61 mmol) and HBTU (1.37 g, 3.61 mmol). After stirring for 10 minutes intermediate 27-1 (1.00 g, 2.78 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 27-2 as a white solid.

Step 3: Intermediate 27-3•TFA

Intermediate 27-2 (1.42 g, 2.58 mmol) was dissolved in a mixture of CH₂Cl₂ (15 mL) and TFA (15 mL). The solution was stirred for 1 hr at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 27-3•TFA as a white solid. MS (m/z) M+1=451.4

Step 4: Intermediate 27-4

To a solution of Boc- α -tBuGly-OH (668 mg, 2.89 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (2.1 mL, 12.0 mmol), HOBt (488 mg, 3.61 mmol) and HBTU (1.37 g, 3.61 mmol). After stirring for 10 minutes intermediate 27-3•TFA (1.36 g, 2.41 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 27-4 as a white solid.

Step 5: Intermediate 27-5•TFA

Intermediate 27-4 (1.38 g, 2.08 mmol) was dissolved in a mixture of DCM (10 mL) and TFA (10 mL). The solution was stirred for 4 hrs at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide intermediate 27-5•TFA as a white solid. MS (m/z) M+1=564.4

Step 6: Intermediate 27-6

To a solution of Boc-NMe-Ala-OH (902 mg, 4.44 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (3.50 mL, 20.2 mmol), HOBt (682 mg, 5.05 mmol) and HBTU

(1.92 g, 5.05 mmol). After stirring for 10 minutes intermediate 27-5•TFA (1.14 g, 1.68 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 27-6 as a white solid.

Step 7: Intermediate 27-7

To a solution of intermediate 27-6 (925 mg, 1.24 mmol) in anhydrous MeOH (25 mL) and stirred under N₂ was added 10% Pd/C (125 mg). The reaction mixture was purged with H₂ and stirred for 1hr. The reaction was then filtered through celite and the filtrate was concentrated in vacuo. Purification by silica gel chromatography provided intermediate 27-7 as a white solid. MS (m/z) M+1=615.4

Step 8: Intermediate 27-8

To a solution of intermediate 27-7 (150 mg, 0.25 mmol) in DCM cooled to 0 °C were sequentially added TEA (100 uL, 0.73 mmol) and terephthaloyl chloride (25.0 mg, 0.12 mmol) and the reaction was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by silica gel chromatography provided intermediate 27-8 as a white solid.

Step 9: Compound 35•2HCl

4N HCl in 1,4-dioxane (2 mL) was added to intermediate 27-8 (166 mg, 0.12 mmol) and the solution was stirred for 1 hr at room temperature. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 35•2HCl as a white solid. MS (m/z) (M+2)/2=580.6.

Compounds 36, 37, and 38 were prepared from intermediate 27-7, in a similar manner to that described for compound 35, using oxalyl chloride, isophthaloyl dichloride and 4,4'-biphenyldicarbonyl chloride, respectively, in place of phthaloyl chloride in Step 8

Compound 36- MS (m/z) (M+2)/2=542.6.

Compound 37- MS (m/z) (M+2)/2=580.6.

Compound 38- MS (m/z) (M+2)/2=618.6.

Compound 44**Step 1: Intermediate 26-1**

To a solution of intermediate 10-6 (150 mg, 0.27 mmol) in THF were sequentially added
5 TEA (112 μ L, 0.81 mmol) and 1,4-phenylene diisocyanate (43 mg, 0.27 mmol) and the
reaction was stirred at room temperature for 4 hrs. Volatiles were removed under reduced
pressure and the residue purified by silica gel chromatography to provide intermediate 26-
1 as a white solid.

Step 2: Compound 44•2TFA

Intermediate 26-1 (148 mg, 0.12 mmol) was dissolved in a mixture of CH_2Cl_2 (1.5 mL) and
TFA (0.4 mL). The solution was stirred for 1 hr at room temperature. Volatiles were
removed under reduced pressure and the residue was triturated with diethyl ether to
provide compound 44•2TFA as a white solid. MS (m/z) (M+2)/2=538.4.

Compound 62:

Intermediate 21-9 was treated with 4N HCl in 1,4-dioxane for 2 hrs. Volatiles were
removed under reduced pressure and the residue was triturated with diethyl ether to
provide compounds 62•HCl as an off white solid. MS (m/z) M+1=787.6.

Intermediate 28-6**Step 1: Intermediate 28-2**

(S)-(+)-2-Phenylglycinol, 28-1 (1.64 g, 12.0 mmol) was dissolved in CH_2Cl_2 (90 mL).
Boc₂O (2.84 g, 13.0 mmol) and DMAP (34 mg, 0.02 mmol) were added and stirred for 1
25 hour at room temperature. The reaction mixture was diluted with diethyl ether (200 mL)
and 1N HCl (100 mL). The organic layer was washed with 1M HCl (2 x 100 mL), dried
over anhydrous MgSO_4 , filtered and volatiles removed under reduced pressure. The
resulting residue was purified by silica gel chromatography provided an oil to provide
intermediate 28-2 as white solid (2.7 g, 95% yield). MS (m/z) M+1=238.2.


Step 2: Intermediate 28-6

Intermediate 28-2 (420 mg, 1.77mmol) and iodomethane (330 μ L, 5.29 mmol) were
dissolved in anhydrous DMF (25 mL). The mixture was cooled to 0 °C, and then NaH
(60% dispersion in oil, 103 mg, 2.58 mmol) was added. After 2 hours, the reaction mixture
35 was diluted with ethyl acetate (200 mL) and 1M HCl (100 mL). The organic layer was

washed with 1M HCl (2 x 100 mL), dried over anhydrous MgSO_4 , filtered and volatiles removed under reduced pressure. The residue was purified by silica gel chromatography to provide intermediate 28-3 as an oil. Intermediate 28-3 was then chilled to 0 °C and treated with 4M HCl in 1,4-dioxane (5 mL). After stirring for 90 minutes volatiles were removed under reduced pressure and the resulting solid was washed with diethyl ether, to provide intermediate 28-6·HCl as white solid (237 mg, 69% yield). MS (m/z) $M+1=152.2$.

A similar procedure was used for the preparation of intermediates 28-7 and 28-8, wherein methyl iodide was replaced with benzyl bromide for intermediate 28-7 and iodoacetamide for intermediate 28-8.

Compound 66



Step 1: Intermediate 29-1

To a solution of benzaldehyde (840 μL , 8.25 mmol) in CH_2Cl_2 (150 mL) was added (S)-2-methylpropane-2-sulfinamide (1.0 g, 8.25 mmol) and titanium ethoxide (3.5 mL, 16.50 mmol). The reaction mixture was refluxed for 5 hours and cooled to room temperature. Water was added and the mixture was vigorously stirred, then filtered through celite. The aqueous layer was extracted with CH_2Cl_2 (3x) and the combined organic extracts were

washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated in *vacuo*. Purification by silica gel chromatography provided intermediate 29-1 as a yellow oil.

Step 2: Intermediate 29-2

- 5 To a suspension of (S,E)-N-benzylidene-2-methylpropane-2-sulfonamide, 29-11, (110 mg, 0.526 mmol), $[\text{Rh}(\text{cod})(\text{CH}_3\text{CN})_2]\text{BF}_4$ (20.1 mg, 0.053 mmol), para-tolylboronic acid (143 mg, 1.052 mmol) and Et_3N (147 μL , 1.052 mmol) in dioxane (1.2 mL) was added H_2O (2.4 mL). The resulting brown suspension was stirred for 2 days at room temperature. The aqueous layer was extracted with EtOAc (3x) and the combined organic extracts were
- 10 washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated in *vacuo*. Purification by silica gel chromatography provided intermediate 29-2 as a white solid (d.e.= 81%).

Step 3: Intermediate 29-4

- 15 To a solution of (S)-2-methyl-N-((R)-phenyl(p-tolyl)methyl)propane-2-sulfonamide (82 mg, 0.27 mmol), 29-2, in MeOH (270 μL) was added 4N HCl in 1,4-dioxane 4 (140 μL , 0.54 mmol). The solution was stirred for 1 hr at room temperature, then Et_2O was added and a white precipitate was formed. The precipitate was filtered and washed with Et_2O to provide intermediate 29-4•HCl as white solid.

20

Step 4: Intermediate 29-5

- To a solution of intermediate 21-9 (122 mg, 0.123 mmol) in DMF cooled to 0 °C were sequentially added DIPEA (193 μL , 1.107 mmol), HOBt (42 mg, 0.308 mmol) and HBTU (117 mg, 0.308 mmol). After stirring for 10 minutes intermediate 29-4•HCl (59 mg, 0.253
- 25 mmol) was added and the reaction mixture was stirred overnight at room temperature. Water and ethyl acetate were added, the organic layer was separated, washed with 10 % citric acid, aqueous NaHCO_3 and brine, dried over anhydrous MgSO_4 , filtered and concentrated in *vacuo*. Purification by silica gel chromatography provided intermediate 66-1 as a pink solid.

30



Step 5: Compound 66•2HCl

- 4N HCl in 1,4-dioxane (1 mL) was added to intermediate 66-1 (135 mg, 0.12 mmol). The solution was stirred for 1 hour at 0 °C. Volatiles were removed under reduced pressure and the residue was triturated with diethyl ether to provide compound 6•2HCl as a white
- 35 solid. MS (m/z) $(\text{M}+2)/2=573.6$.



Representative compounds of the present invention were prepared according to the above procedures and are illustrated in Table 1:



5

TABLE 1




COMPOUND	STRUCTURE	MS
1		M+1=991.7 [M+2]/2=496.4
2		M+1=1081.8 [M+2]/2=541.4




COMPOUND	STRUCTURE	MS
3		$[M+2]/2=523.4$
4		$M+1=993.5.$




COMPOUND	STRUCTURE	MS
5		$M+1=995.6$ $(M+2)/2=498.5$
6		$(M+2)/2=514.4$




COMPOUND	STRUCTURE	MS
7	 Chemical structure of compound 7: A complex molecule featuring a central chain with multiple amide and ester linkages. It includes a fluorenylmethyl (Fmoc) protecting group on the left, a dimethylamino group on the right, and a fluorenylmethyl group at the bottom. The structure is highly branched and contains several chiral centers.	$(M+2)/2=618.0$
8	 Chemical structure of compound 8: A complex molecule similar to compound 7, but with a different substitution pattern. It features a fluorenylmethyl group on the left, a dimethylamino group on the right, and a fluorenylmethyl group at the bottom. The structure is highly branched and contains several chiral centers.	$M+1=1012$




COMPOUND	STRUCTURE	MS
9		$(M+2)/2 = 642$
10		$[M+2]/2=561.6$




COMPOUND	STRUCTURE	MS
11		M+1=997.6 [M+2]/2=499.4
12		M+1=1067.6 [M+2]/2=534.5
13		[M+2]/2=520.4




COMPOUND	STRUCTURE	MS
14		$[M+2]/2=506.4$
15		$M+1=1073.6$ $[M+2]/2=537.4$
16		$[M+2]/2=569.4$




COMPOUND	STRUCTURE	MS
17		M+1=1019.6 [M+2]/2=510.4
18		M+1=815.4 [M+2]/2=408.2
19		M+1=1029.6 [M+2]/2=515.4

COMPOUND	STRUCTURE	MS
20		$[M+2]/2=497.6$
21		$M+1=1021.6$ $[M+2]/2=511.6$
22		$M+1=869.6$ $[M+2]/2=435.4$





COMPOUND	STRUCTURE	MS
23		M+1=1045.8 [M+2]/2=523.4
24		[M+2]/2=559.4
25		[M+2]/2=525.6





COMPOUND	STRUCTURE	MS
26		$[M+2]/2=561.4$
27		$[M+2]/2=587.6$
28		$[M+2]/2=543.6$




COMPOUND	STRUCTURE	MS
29		M+1=967.6 [M+2]/2=484.5
30		M+1=1157.6 [M+2]/2=579.6
31		[M+2]/2=573.6




COMPOUND	STRUCTURE	MS
32		$[M+2]/2=533.6$
33		$M+1=1117.6$ $[M+2]/2=559.5$
34		$M+1=1045.8$ $[M+2]/2=523.4$




COMPOUND	STRUCTURE	MS
35		$[M+2]/2=580.6$
36		$[M+2]/2=542.6$
37		$[M+2]/2=580.6$
38		$[M+2]/2=618.6$

COMPOUND	STRUCTURE	MS
39		$[M+2]/2=597.4$
40		$M+1=1045.6$ $[M+2]/2=523.4$
41		$M+1=1046.6$ $[M+2]/2=523.8$
42		$[M+2]/2=548.4$

COMPOUND	STRUCTURE	MS
43		M+1=1046.6 [M+2]/2=523.8
44		[M+2]/2=538.4
45		M+1=1018.8 [M+2]/2=509.6
46		M+1=965.6 [M+2]/2=483.4




COMPOUND	STRUCTURE	MS
47		M+1=1113.8 [M+2]/2=557.6
48		M+1=1117.8 [M+2]/2=559.6
49		M+1=993.8 [M+2]/2=497.6

COMPOUND	STRUCTURE	MS
50		M+1=993.8 [M+2]/2=497.6
51		[M+2]/2=619.9
52		M+1=1051.8 [M+2]/2=526.6


COMPOUND	STRUCTURE	MS
53		$M+1=1046.8$ $[M+2]/2=523.4$
54		$M+1=1025.8$ $[M+2]/2=513.5$
55		$M+1=1046.8$ $[M+2]/2=524.2$

COMPOUND	STRUCTURE	MS
56		[M+2]/2=584.6
57		M+1=1095.8 [M+2]/2=548.6
58		M+1=1167.8 [M+2]/2=584.6

COMPOUND	STRUCTURE	MS
59		M+1=1025.8 [M+2]/2=513.6
60		M+1=1117.8 [M+2]/2=559.6
61		M+1=1017.8 [M+2]/2=509.6


COMPOUND	STRUCTURE	MS
62		$M+1=787.6$ $[M+2]/2=394.4$
63		$[M+1]=1053.8$ $(M+2)/2=527.8$
64		$[M+1]=1206.0$ $(M+2)/2=603.6$

COMPOUND	STRUCTURE	MS
65		[M+1]=1139.8 (M+2)/2=570.6
66		(M+2)/2=573.6


COMPOUND	STRUCTURE	MS
67		

Representative compounds of the present invention which can be prepared by simple modification of the above procedures are illustrated in Tables 2 to 6:

TABLE 2



5

wherein A and A¹ are both CH₂ or both C(O), and

R ⁴	R ⁵
H	
H	
H	
H	-CH ₃
H	
H	
H	
H	





TABLE 3

M1-BG-M2

5













Formula 1A

BG is



10

M1	M2

M1	M2
	
	
	
	
	
	

















TABLE 4

5

M1-BG-M2**Formula 1B****BG is**

10

M1	M2

M1	M2
	
	
	
	
	
	






TABLE 5



5

R^1 , R^2 , R^3 , R^{100} , R^{200} and R^{300} are defined as hereinabove,

-X-L-X'- is chosen from:






10

R^4 and R^{400} are H ;


15

R^5 and R^{500} are chosen from:




wherein the aryl moieties may be substituted by R¹⁰ and wherein, R¹⁰ and R^{10'} are independently defined as R¹⁰ hereinabove, and wherein the alkyl may be further substituted by R⁶ as defined hereinabove.

Table 6






$R^1, R^2, R^3, R^{100}, R^{200}$ and R^{300} are defined as hereinabove,

5 -X-L-X'- is chosen from:



10 R^4 and R^{400} are H ;

R^5 and R^{500} are chosen from:



wherein the aryl moieties may be substituted by R^{10} and wherein, R^{10} and $R^{10'}$ are independently defined as R^{10} hereinabove, and wherein the alkyl may be further substituted by R^6 as defined hereinabove.

10

Assays

Molecular constructs for expression

GST-XIAP BIR3RING: XIAP coding sequence amino acids 246-497 cloned into PGEX2T1 via BamH1 and AVA I. The plasmid was transformed into E. coli DH5 α for use in protein expression and purification.

15

GST-HIAP2 (cIAP-1) BIR 3: HIAP2 coding sequence from amino acids 251-363 cloned into PGex4T3 via BamH1 and XhoI. The plasmid was transformed into E. coli DH5 α for use in protein expression and purification.

20

GST-HIAP1(cIAP-2) BIR 3: HIAP1 coding sequence from amino acids 236-349, cloned into PGex4T3 via BamH1 and XhoI. The plasmid was transformed into *E. coli* DH5 α for use in protein expression and purification.

- 5 GST- linker BIR 2 BIR3Ring: XIAP coding sequence from amino acids 93-497 cloned into PGex4T1 via BamH1 and XhoI. Amino acids 93-497 were amplified from full length XIAP in pGex4t3, using the primers: TTAATAGGATCCATCAACGGCTTTTATC and GCTGCATGTGTGTCAGAGG, using standard PCR conditions. The PCR fragment was TA cloned into pCR-2.1 (invitrogen). Linker BIR 2 BIR 3Ring was subcloned into pGex4T1
10 by BamHI/XhoI digestion. The plasmid was transformed into *E. coli* DH5 α for use in protein expression and purification.

- Full-length human XIAP, AEG plasmid number 23. XIAP coding sequence amino acids 1-497 cloned into GST fusion vector, PGEX4T1 via BamH1 and Xho I restriction sites. The
15 plasmid was transformed into *E. coli* DH5 α for use in protein purification.

- GST-XIAP linker BIR 2: XIAP linker BIR 2 coding sequence from amino acids 93-497 cloned into pGex4T3 via BamHI and XhoI. The plasmid was transformed into *E. coli* DH5 α for use in protein expression and purification.
20

Expression and purification of recombinant proteins

A. Expression of Recombinant Proteins

- Glutathione S-transferase (GST) tagged proteins were expressed in *Escherichia coli* strains DH5-alpha. For expression full length XIAP, individual or combinations of XIAP-
25 BIR domains, cIAP-1, cIAP-2 and Livin transformed bacteria were cultured overnight at 37°C in Luria Broth (LB) medium supplemented with 50 ug/ml of ampicillin. The overnight culture was then diluted 25 fold into fresh LB ampicillin supplemented media and bacteria were grown up to $A_{600} = 0.6$ then induced with 1 mM isopropyl-D-1-thiogalactopyranoside for 3 hours. Upon induction, cells were centrifuged at 5000 RPM for 10 minutes and the
30 media was removed. Each pellet obtained from a 1 liter culture received 10 ml of lysis buffer (50 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 1 mM PMSF, 2 mg/ml of lysosyme, 100 μ g/ml), was incubated at 4°C with gentle shaking. After 20 minutes of incubation, the cell suspension was placed at -80°C overnight or until needed.


- 35 *B. Purification of recombinant proteins*

For purification of recombinant proteins, the IPTG-induced cell lysate was thawed vortexed and then disrupted by flash freezing in liquid nitrogen two times with vortexing after each thaw. The cells were disrupted further by passing the extract four times through a Bio-Neb Cell disruptor device (Glas-col) set at 100 psi with Nitrogen gas. The extract
5 was clarified by centrifugation at 4 °C at 15000 RPM in a SS-34 Beckman rotor for 30 minutes. The resulting supernatant was then mixed with 2 ml of glutathione-Sepharose beads (Pharmacia) per 500 ml cell culture (per 1000ml culture for full length XIAP) for 1 hour at 4C. Afterwards, the beads were washed 3 times with 1X Tris-Buffered Saline (TBS) to remove unbound proteins. The retained proteins were eluted with 2 washes of 2
10 ml of 50 mM TRIS pH 8.0 containing 10 mM reduced glutathione. The eluted proteins were pooled and precipitated with 604g/liter of ammonium sulfate and the resulting pellet re-suspended into an appropriate buffer. As judged by SDS-PAGE the purified proteins were >90% pure. The protein concentration of purified proteins was determined from the Bradford method.

His-tag proteins were expressed in the E. Coli strain in E. coli AD494 cells using a pet28ACPP32 construct. The soluble protein fraction was prepared as described above. For protein purification, the supernatant was purified by affinity chromatography using chelating-Sepharose (Pharmacia) charged with NiSO₄ according to the manufacturer's
20 instructions. Purity of the eluted protein was >90% pure as determined by SDS-PAGE. The protein concentration of purified proteins was determined from the Bradford assay.

Synthesis of fluorescent probe P1

A fluorescent peptide probe, Fmoc-Ala-Val-Pro-Phe-Tyr(t-Bu)-Leu-Pro-Gly(t-Bu)-Gly-OH
25 was prepared using standard Fmoc chemistry on 2-chlorotrityl chloride resin (Int. J. Pept. Prot. Res. 38:555-561, 1991). Cleavage from the resin was performed using 20% acetic acid in dichloromethane (DCM), which left the side chain still blocked. The C-terminal protected carboxylic acid was coupled to 4'-(aminomethyl)fluorescein (Molecular Probes, A-1351; Eugene, Oreg.) using excess diisopropylcarbodiimide (DIC) in dimethylformamide
30 (DMF) at room temperature and was purified by silica gel chromatography (10% methanol in DCM). The N-terminal Fmoc protecting group was removed using piperidine (20%) in DMF, and purified by silica gel chromatography (20% methanol in DCM, 0.5% HOAc). Finally, the t-butyl side chain protective groups were removed using 95% trifluoroacetic acid containing 2.5% water and 2.5% triisopropyl silane, to provide probe P1 (>95% pure,
35 HPLC).

Probe P2**5 Binding assay****Fluorescence polarization-based competition assay**

For all assays, the fluorescence and fluorescence-polarization was evaluated using a Tecan Polarion instrument with the excitation filter set at 485 nm and the emission filter set at 535 nm. For each assay, the concentration of the target protein was first established by titration of the selected protein in order to produce a linear dose-response signal when incubated alone in the presence of the fluorescent probe P1 or P2. Upon establishing these conditions, the compounds potency (IC_{50}) and selectivity, was assessed in the presence of a fix defined- amount of target protein and fluorescent probe and a 10 point serial dilution of the selected compounds. For each IC_{50} curve, the assays were run as followed: 25 μ L/well of diluted compound in 50 mM MES buffer pH 6.5 were added into a black 96 well plate then 25 μ L/well of bovine serum albumin (BSA) at 0.5 mg/ml in 50 mM MES pH 6.5. Auto-fluorescence for each compound was first assessed by performing a reading of the compound/BSA solution alone. Then 25 μ L of the fluorescein probe diluted into 50 mM MES containing 0.05 mg/ml BSA were added and a reading to detect quenching of fluorescein signal done. Finally 25 μ L/well of the target or control protein (GST- BIRs) diluted at the appropriate concentration in 50 mM MES containing 0.05 mg/ml BSA were added and the fluorescence polarization evaluated.

Determination of IC₅₀ and Inhibitory constants

For each assay the relative polarization-fluorescence units were plotted against the final concentrations of compound and the IC₅₀ calculated using the Grad pad prism software and/or Cambridge soft. The k_i value were derived from the calculated IC₅₀ value as described above and according to the equation described in Nikolovska-Coleska, Z. (2004) Anal Biochem 332, 261-273.

Fluorescence polarization competition assay

The k_i of various compounds in the BIR2-BIR3-ring FP assay, using probe P2, was determined as described above. For example, compound 3 displayed a k_i of less than 100 nM.

Caspase-3 full length XIAP, linker BIR2 or Linker- BIR2- BIR3-RING derepression assay

In order to determine the relative activity of the selected compound against XIAP-Bir2, we setup an in vitro assay where caspase-3 was inhibited by GST fusion proteins of XIAP linker-Bir2, XIAP Linker Bir2-Bir3-RING or full-length XIAP. Caspase 3 (0.125ul) and 12.25-34.25nM (final concentration) of GST-XIAP fusion protein (GST-Bir2, GST-Bir2Bir3RING or full-length XIAP) were co-incubated with serial dilutions of compound (200 uM-5 pM). Caspase 3 activity was measured by overlaying 25 uL of a 0.4mM DEVD-AMC solution. Final reaction volume was 100 uL. All dilutions were performed in caspase buffer (50mM Hepes pH 7.4, 100mM NaCl, 10% sucrose, 1mM EDTA, 10mM DTT, 0.1% CHAPS (Stennicke, H.R., and Salvesen, G.S. (1997). Biochemical characteristics of caspase-3, -6, -7, and -8. J. Biol. Chem. 272, 25719–25723).

The fluorescent AMC released from the caspase-3 hydrolysis of the substrate was measured in a TECAN spectrophotometer at 360 nm excitation and 444 nm emission, after 15 minutes of incubation at room temperature. IC₅₀ values were calculated on a one or two-site competition model using GraphPad v4.0, using the fluorescence values after 15 minutes of incubation plotted against the log₁₀ concentration of compound.

IC₅₀ values of preferred compounds were shown to correlate with EC₅₀ values against SKOV3s and were typically less than 1 uM.

Cell-free assay**Caspase de-repression assay using cellular extracts (apoptosome)**

100 μ g of 293 cell S100 extract and 0.25 μ M-2 μ M of GST-XIAP fusion protein (XIAP-Bir3RING, XIAP-Bir2Bir3RING, or full-length XIAP) were co-incubated with serial dilutions of compound (40 μ M-5 pM). Caspases present in the extracts were activated by adding
5 1mM dATP, 0.1mM ALLN, 133 μ g Cytochrome C (final concentrations), and incubating at 37°C for 25 minutes. All reactions and dilutions used S100 buffer (50 mM Pipes pH 7.0, 50mM KCl, 0.5mM EGTA pH 8.0, 2mM MgCl₂ supplemented with 1/1000 dilutions of 2 mg/ml Cytochalisin B, 2 mg/ml Chymotstatin, Leupeptin, Pepstatin, Antipain , 0.1M PMSF,
10 1M DTT). Final reaction volume was 30 μ l. Caspase-3 activity was measured by overlaying 30 μ l of a 0.4mM DEVD-AMC solution. Released AMC cleavage was measured in a TECAN spectrophotometer at 360 nm excitation and 444nm emission, on a kinetic cycle of 1 hour with readings taken every 5 minutes. Caspase activity was calculated as V_o of AMC fluorescence/sec. Caspase de-repression by our compounds was compared to fully
15 activated extract and activated extract repressed by the presence of XIAP fusion protein.

IC₅₀ values of preferred compounds were shown to correlate with EC₅₀ values against SKOV3s and were typically less than 1 μ M.

20 Cell Culture and Cell Death Assays*A. Cell culture*

MDA-MD-231 (breast) and H460 (lung) cancer cells were cultured in RPMI1640 media supplemented with 10% FBS and 100 units/mL of Penicillin and Steptomycin.

25 B. Assays

Survival assays were performed on various cell lines including MDA-MB-231, SKOV3, H460, PC3, HCT-116, and SW480 cells. Cells were seeded in 96 well plates at a respective density of 5000 and 2000 cells per well and incubated at 37 °C in presence of 5% CO₂ for 24 hours. Selected compounds were diluted into the media at various
30 concentration ranging from 0.01 μ M up to 100 μ M. Diluted compounds were added onto the MDA-MB-231 cells. For the MDA-MB-231 SKOV3, H460, PC3, HCT-116, and SW480 cells, the compounds were added either alone or in presence of 1-3 ng/ml of TRAIL. After 72 hours cellular viability was evaluated by MTS based assays. A solution of [3-(4,5-dimethylthiazol-2-yl)-5-(3- carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium, inner

salt; MTS] was added onto cells for a period of 1 to 4 hours. Upon incubation the amount of converted MTS was evaluated using a Tecan spectrophotometer set at 570 nm.

MDA-MB-231, SKOV3, and PC3 cells were treated with selected compounds of the present invention and found to have EC₅₀s of 100nM or less. When the above cell lines were treated with the compounds of the present invention in the presence of TRAIL, they were shown to have EC₅₀s of 50 nM or less.

Survival MTT assay

One day prior the treatment with compound, 2000 to 4000 cells per well were plated in a tissue culture treated 96 well format dish with 100ul of media and incubated at 37 °C, 5% CO₂. On the day of compound treatment, compounds were diluted with cell culture media to a working stock concentration of 2X. 100 uL of diluted compound were then added to each well. The treated plate was incubated for 72h at 37 °C, 5% CO₂. Upon incubation, the cell viability was assessed as followed 20 uL of MTT reagent at 5 mg/ml were added per well to cell plate. The plate was incubated for 2h at 37 °C in presence of 5% CO₂. The supernatant was then removed from the plate and 100 uL of isopropanol were added. The absorbance was measured in a TECAN spectrophotometer at 570 nm. The percentage of viability was expressed in percentage of the signal obtained with non treated cells.

As seen in Table 7, compounds represented in Table 1 hereinabove generally displayed EC₅₀ values against MDA-MB-231 and SKOV-3 cells of <1μM. Select compounds had EC₅₀ of <50 nM.

TABLE 7

Compound	MDA-MB231 EC ₅₀ (nM)	SKOV-3 EC ₅₀ (nM)
1	A	A
2	A	A
3	A	A
5	C	
10	A	A
11		A
12		A
13		A
14		A
15		A
16		A
17		A

Compound	MDA-MB231 EC ₅₀ (nM)	SKOV-3 EC ₅₀ (nM)
18		D
19		A
20		A
21		A
22		B
23		A
24		A
25		A
26		B
27		B
28		A
29		B
30		C
31		B
32		B
33		A
34		B
35		B
36		C
37		C
38		B
39		A
40		A
41		A
42		A
43		B
44		A
45		D
46		B
47		A
48		B
49		A
50		B
51		B
52		A
53		D
54		D
55		A
56		A
57		A
58		B
59		B

Compound	MDA-MB231 EC ₅₀ (nM)	SKOV-3 EC ₅₀ (nM)
60		A
61		A
64		A

A - EC₅₀ less than 50 nM
 B - EC₅₀ less than 250 nM
 C - EC₅₀ more than 1000 nM
 5 D- EC₅₀ more than 1000 nM

Apoptosis Assay: Measurement of caspase-3 activity from cultured cells.

- 10 One day, prior to the treatment, 10,000 cells per well were plated in a white tissue culture treated 96 well plate with 100 uL of media. On the day of compound treatment, compounds were diluted with cell culture media to a working stock concentration of 2X and 100ul of diluted compound were added to each well and the plate was incubated for 5h at 37 °C in presence of 5 % CO₂. Upon incubation, the plate was washed twice with
- 15 200 uL of cold TRIS Buffered Saline (TBS) buffer. Cells were lysed with 50ul of Caspase assay buffer (20mM Tris-HCl pH 7.4, 0.1% NP-40, 0.1% Chaps, 1mM DTT, 0.1 mM EDTA, 0.1 mM PMSF, 2 mg/ml Chymostatin, Leupeptin, Pepstatin, Antipapin) then incubated at 4 °C with shaking for 30minutes. 45ul of Caspase assay buffer and 5 uL of Ac-DEVD-AMC at 1 mg/ml were added to each well, the plate shaken and incubated for
- 20 16h at 37 °C. The amount of release AMC was measured in a TECAN spectrophotometer at with the excitation and emission filter set at 360 nm and 444 nm. The percentage of Caspase-3 activity was expressed in comparison of the signal obtained with the non-treated cells.
- 25 IC₅₀ values of preferred compounds were shown to correlate with EC₅₀ values against SKOV3s and were typically less that 1 uM.

Cellular biochemistry:

A. Detection of XIAP and PARP/Caspase-3/Caspase-9

- 30 Detection of cell expressed XIAP and PARP were done by western blotting. Cells were plated at 300 000 cells/well in a 60 mm wells (6 wells plate dish). The next day the cells were treated with selected compound at the indicated concentration. 24 hours later cells the trypsinized cells, pelleted by centrifugation at 1800rpm at 4 °C. The resulting pellet was rinsed twice with cold TBS. The final washed pellet of cells was the lysed with 250 uL

Lysis buffer (NP-40, glycerol, 1% of a protease inhibitor cocktail (Sigma)), placed at 4°C for 25min with gentle shaking. The cells extract was centrifuged at 4 °C for 10min at 10,000 rpm. Both the supernatant and the pellet were kept for western blotting analysis as described below. From the supernatant, the protein content was evaluated and about 50
5 ug of protein was fractionated onto a 10% SDS-PAGE. Pellets were washed with the lysis buffer and re-suspend into 50ul of Lamelli buffer 1X, boiled and fractionated on SDS-PAGE. Upon electrophoresis each gel was electro-transferred onto a nitrocellulose membrane at 0.6A for 2 hours. Membrane non-specific sites were blocked for 1 hours with 5% Skim milk in TBST (TBS containing 0.1% (v/v) Tween-20) at RT. For protein immuno-
10 detection, membranes were incubated overnight with primary antibodies raised against XIAP clone 48 obtained from Becton-Dickison) or PARP: obtained from Cell signal or caspase-3 or caspase-9 primary antibodies were incubated at 4°C with shaking at dilutions as follows:

15 XIAP clone 80 (Becton-Dickinson).....1/2500
PARP (Cell Signal).....1/2500
Caspase 3 (Sigma).....1/1500
Caspase 9 (Upstate).....1/1000

20 Upon overnight incubation, the membranes received three washes of 15 min in TBST then were incubated for 1 hour at room temperature in the presence of a secondary antibody coupled with HRP-enzyme (Chemicon) and diluted at 1/5 000. Upon incubation each membrane were washed three times with TBST and the immunoreactive bands were detected by addition of a luminescent substrate (ECL kit Amersham) and capture of signal on a X-RAY film for various time of exposure.

25 Certain exemplified compounds were shown to induce the cleavage of PARP near concentrations which correlate with EC₅₀ values against SKOV3s and were typically less than 1 uM.

30 Hollow fiber model

Hollow fiber in vivo model were used to demonstrate in vivo efficacy of selected compounds against selected cell lines as single agent therapy or in combination with selected cytotoxic agents. At day 1, selected cell lines were cultured and the fiber filled at a cell density of about 40,000 cells/fiber. At the day of operation (day 4), three fibers are
35 implanted sub-cutaneous into 28-35 Nu/Nu CD-1 male mice. On day 5, mice start to

receive daily injection via sub-cutaneous route of control vehicle or vehicle containing the selected compound at the appropriate concentration and/or injection of cytotoxic agent via intra-peritoneal route. Upon 3-7 days of consecutive drug treatments, the animals are sacrificed, each fiber is removed and the metabolic viability of the remaining cells
5 determined by MTT assay. Efficacy of the compound is define as the difference between the vehicle-treated animal and the animal treated with the compound alone or the compound given in combination of the cytotoxic agent.

MDA-MB-231 cells were implanted on day 1. Compound 3 was administered for 4
10 consecutive days via IV bolus (tail vein) injections at 1, 3, and 10 mg/kg (2 mg/mL in 20 % aqueous HPCD). Complete suppression of cell growth, as compared to 20% HPCD control, was observed for compound 3 at drug concentrations of 3 mg/kg.

SKOV-3 Human Ovarian Cancer Cell Line Xenograft Study with Compound 3

15 Female CD-1 nude mice (approximately 20-25 g) were subcutaneously injected 5×10^6 SKOV-3 human ovarian tumor cells in 50% matrigel subcutaneously in the right flank. On day 55, when tumors were approximately 100 mm^3 , treatment was initiated with compound 3 on a 5 on/2 off treatment schedule for the duration of the experiment. Tumor size was measured with digital calipers and calculated as $V = (a \times b^2)/2$, wherein, a is the
20 longest dimension and b is the width.

Tumor regression was observed while dosing compound 3 at 1 mg/kg while tumor stasis was observed while dosing compound 3 at 0.3 mg/kg (see Figure 1).

MDA-MB-231 Human Mammary Cancer Cell Line Xenograph Study with Compound 3

25 Female CD-1 nude mice (approximately 20-25g) were subcutaneously injected 1×10^6 MDA-MB-231 human mammary tumor cells in the right flank. On day 71, when tumors were approximately 90 mm^3 , treatment was initiated with compound 3 on a 5 on/2 off
30 treatment schedule for the duration of the experiment. Tumor size was measured with digital calipers and calculated as $V = (a \times b^2)/2$, wherein, a is the longest dimension and b is the width.

Tumor regression was observed while dosing compound 3 at 1 mg/kg (see Figure 2).

Pharmacokinetic studies

Selected compounds were dissolved into normal saline and given at various doses using different route of administration, including intravenous bolus, intravenous infusion, oral and subcutaneous injection.

5

Compound of the present invention demonstrated acceptable pharmacokinetics via various routes of administration.

In vitro potency

- 10 Compounds of the instant invention were shown to kill SKOV3 (ovarian), MDA-MB-231 (breast), BT549 (breast), HL-60 (acute promyelocytic leukemia) and PANC-1 (pancreatic), cell lines in vitro, demonstrating EC₅₀s ranging of 0.1 nM to 1000 nM (see Table 7).

- 15 The SAR of these compounds was mapped using SKOV3s and several interesting trends emerged. Changing the stereochemistry at the pyrrolidine bridging site effected the potency of the compounds as seen in the EC₅₀s of the cis- and trans-proline derivatives 3 and 29 (EC₅₀=1 nM, 88 nM, respectively. Amide bridging units provided active compounds, however, a significant range in potency against SKOV3 cells was observed by varying the components of the bridging units. Small, conformationally constrained
- 20 bridging units including, but not limited to, 1,4-phenyl dicarboxamides (terephthaloylamides), 1,3-phenyl dicarboxamides, 2,6-naphthyl dicarboxamides, 1,4-cyclohexyl dicarboxamides, 3,5-pyridyl dicarboxamides, or C₂-C₁₀ aliphatic dicarboxamides provide highly active compounds. Bridging units containing bis-glycine amides such as compounds 30 provide less active compounds (EC₅₀=188 nM).

25

Ether, urea and sulfonamide bridging units provide compounds which were active against SKOV3 cells, although the sulfonamide bridged compounds were generally less active.

- Significant shifts in potency against SKOV3 cells were observed by varying the P4
- 30 substitution, wherein (R)-stereochemistry at the R⁴/R⁴⁰⁰ amide provides compounds which are 5-10 times more potent than the corresponding (S)-isomer. The introduction of hydrophilic moieties close to the amide provide less active compounds such as compound 49.

Additionally, alkylation of the N-terminal alanine moiety provides compounds which are up to 100 fold more potent than the corresponding unsubstituted N-terminal alanine derivative.

5 A subset of cancer cell lines, however, were not innately sensitive to compounds of the instant invention, with EC_{50} s greater than 1000 nM. We have demonstrated that IAP BIR binding compounds demonstrate synergistic killing of various cancer cell lines with death receptor agonists such as TRAIL, agonist TRAIL receptor anti-bodies, TNF- α , and others. We herein disclose that compound of formula I and II also demonstrate synergistic killing
10 of various cancer cell lines with death receptor agonists such as TRAIL. When these cells were treated with compound and TRAIL, or antagonist TRAIL antibody, these cell lines were highly sensitive to compound with EC_{50} s generally less than 100 nM.

These cancer cell lines include HELA (cervical), HCT116 (colon), PC3 (prostate), OVCAR-
15 3 (ovarian), HEY (ovarian), and H460 (lung). In the presence of TRAIL (1-3 ng/mL) and varying concentrations of compound 3 the EC_{50} s for the above cell lines were less than 1000 nM.

20 **In vivo potency**

Compound 3 was tested in SKOV3 and MDA-MB-231 xenograph tumour models (see Figures 1 and 2). In both cases tumour regression was observed at 1 mg/kg when given 5 days on and 2 days off. Tumour stasis was observed in SKOV3 xenograph at 0.1 mg/kg.

25 **Discussion**

The above results suggest that the IAP BIR binding compounds of the instant invention are highly potent agents, both in vitro and in vivo, wherein a mechanistic link between IAP binding and IAP modulations can be correlated to anti-cancer efficacy. We have demonstrated that compounds of the instant invention bind to the BIR domains of the IAPs
30 with high affinity resulting in the release of active caspases 3 and 9. Further, these compounds result in the induction of apoptosis in cancer cells while synergistically sensitizing cancer cell lines to death receptor agonists such as TRAIL. Moreover, when tumour bearing animals were treated with compounds of the instant invention demonstrated tumour stasis and/or tumour regression at pharmaceutically relevant
35 doses..

Compounds of the instant invention demonstrated acceptable pharmacokinetics via several routes of administration.

5 Other Embodiments

From the foregoing description, it will be apparent to one of ordinary skill in the art that variations and modifications may be made to the invention described herein to adapt it to various usages and conditions. Such embodiments are also within the scope of the present invention.

10 All publications mentioned in this specification are hereby incorporated by reference.


CLAIMS

5


We claim:

1. An isomer, enantiomer, diastereoisomer or tautomer of a compound represented by Formula I or II

10



I



II

15

wherein

m is 0, 1 or 2;

Y is NH, O or S;

BG is

20









1) $-X-L-X^1-$;

X and X¹ are independently selected from

1) O,

2) NR^{13} ,

3) S,

4) $-\text{C}_1-\text{C}_6$ alkyl-,5) $-\text{C}_1-\text{C}_6$ alkyl-O,5 6) $-\text{C}_1-\text{C}_6$ alkyl- NR^{13} -,7) $-\text{C}_1-\text{C}_6$ alkyl-S-,

15



L is selected from:

1) $-\text{C}_1-\text{C}_{20}$ alkyl-,2) $-\text{C}_2-\text{C}_6$ alkenyl-,3) $-\text{C}_2-\text{C}_4$ alkynyl-,20 4) $-\text{C}_3-\text{C}_7$ cycloalkyl-,

5) -aryl-,

6) -biphenyl-,

- 7) – heteroaryl–,
 8) – heterocyclyl–,
 9) –C₁–C₆ alkyl–(C₂–C₆ alkenyl)–C₁–C₆ alkyl–,
 10) –C₁–C₆ alkyl–(C₂–C₄ alkynyl)–C₁–C₆ alkyl–
 5 11) –C₁–C₆ alkyl–(C₃–C₇ cycloalkyl)–C₁–C₆ alkyl–,
 12) –C₁–C₆ alkyl– aryl–C₁–C₆ alkyl–,
 13) –C₁–C₆ alkyl–biphenyl–C₁–C₆ alkyl–,
 14) –C₁–C₆ alkyl–heteroaryl–C₁–C₆ alkyl–,
 15) –C₁–C₆ alkyl–heterocyclyl–C₁–C₆ alkyl–,
 10 16) –C₁–C₆ alkyl–Y–C₁–C₆ alkyl–,
 17) –aryl–Y–aryl–,
 18) –heteroaryl–Y–heteroaryl–,
 19) –heterocyclyl–Y–heterocyclyl–,



wherein the alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more R⁶ substituents, and the aryl, biphenyl, heteroaryl, and heterocyclyl are optionally substituted with one or more R¹⁰ substituents;

20 Q and Q¹ are independently selected from

- 1) NR⁴R⁵,
- 2) OR¹¹, or
- 3) S(O)_mR¹¹; or

25 Q and Q¹ are independently selected from

- 1) aryl, or
- 2) heteroaryl, the aryl and the heteroaryl being optionally substituted with one or more R¹⁰ substituents;

30 A and A¹ are independently selected from

- 1) –CH₂–,

- 2) $-\text{CH}_2\text{CH}_2-$,
 3) $-\text{CH}(\text{C}_1-\text{C}_6 \text{ alkyl})-$,
 4) $-\text{CH}(\text{C}_3-\text{C}_7 \text{ cycloalkyl})-$,
 5) $-\text{C}_3-\text{C}_7 \text{ cycloalkyl}-$,
 5 6) $-\text{CH}(\text{C}_1-\text{C}_6 \text{ alkyl}-\text{C}_3-\text{C}_7 \text{ cycloalkyl})-$,
 7) $-\text{C}(\text{O})-$, or
 8) $-\text{C}(\text{O})\text{OR}^{13}$;

R^1 and R^{100} are independently selected from

- 10 1) H, or
 2) C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

R^2 and R^{200} are independently selected from

- 1) H, or
 15 2) C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

R^3 and R^{300} are independently C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

20 R^4 and R^5 are each independently selected from

- 1) H,
 2) haloalkyl,
 3) $\leftarrow\text{C}_1-\text{C}_6 \text{ alkyl}$,
 4) $\leftarrow\text{C}_2-\text{C}_6 \text{ alkenyl}$,
 25 5) $\leftarrow\text{C}_2-\text{C}_4 \text{ alkynyl}$,
 6) $\leftarrow\text{C}_3-\text{C}_7 \text{ cycloalkyl}$,
 7) $\leftarrow\text{C}_3-\text{C}_7 \text{ cycloalkenyl}$,
 8) $\leftarrow\text{aryl}$,
 9) $\leftarrow\text{heteroaryl}$,
 30 10) $\leftarrow\text{heterocyclyl}$,
 11) $\leftarrow\text{heterobicyclyl}$,
 12) $\leftarrow\text{C}(\text{O})-\text{R}^{11}$,
 13) $\leftarrow\text{C}(\text{O})\text{O}-\text{R}^{11}$,
 14) $\leftarrow\text{C}(=\text{Y})\text{NR}^8\text{R}^9$, or
 35 15) $\leftarrow\text{S}(\text{O})_2-\text{R}^{11}$,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

5 R^6 is

- 1) halogen,
- 2) NO_2 ,
- 3) CN,
- 4) haloalkyl,
- 10 5) C_1-C_6 alkyl,
- 6) C_2-C_6 alkenyl,
- 7) C_2-C_4 alkynyl,
- 8) C_3-C_7 cycloalkyl,
- 9) C_3-C_7 cycloalkenyl,
- 15 10) aryl,
- 11) heteroaryl,
- 12) heterocyclyl,
- 13) heterobicyclyl,
- 14) OR^7 ,
- 20 15) $S(O)_mR^7$,
- 16) NR^8R^9 ,
- 17) $NR^8S(O)_2R^{11}$,
- 18) COR^7 ,
- 19) $C(O)OR^7$,
- 25 20) $CONR^8R^9$,
- 21) $S(O)_2NR^8R^9$,
- 22) $OC(O)R^7$,
- 23) $OC(O)Y-R^{11}$,
- 24) $SC(O)R^7$, or
- 30 25) $NC(Y)NR^8R^9$,

wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

R^7 is

- 35 1) H,

- 2) haloalkyl,
- 3) C₁-C₆ alkyl,
- 4) C₂-C₆ alkenyl,
- 5) C₂-C₄ alkynyl,
- 5 6) C₃-C₇ cycloalkyl,
- 7) C₃-C₇ cycloalkenyl,
- 8) aryl,
- 9) heteroaryl,
- 10) heterocyclyl,
- 10 11) heterobicyclyl,
- 12) R⁸R⁹NC(=Y), or
- 13) C₁-C₆ alkyl-C₂-C₄ alkenyl, or
- 14) C₁-C₆ alkyl-C₂-C₄ alkynyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with
 15 one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

R⁸ and R⁹ are each independently

- 1) H,
- 20 2) haloalkyl,
- 3) C₁-C₆ alkyl,
- 4) C₂-C₆ alkenyl,
- 5) C₂-C₄ alkynyl,
- 6) C₃-C₇ cycloalkyl,
- 25 7) C₃-C₇ cycloalkenyl,
- 8) aryl,
- 9) heteroaryl,
- 10) heterocyclyl,
- 11) heterobicyclyl,
- 30 12) C(O)R¹¹,
- 13) C(O)Y-R¹¹, or
- 14) S(O)₂-R¹¹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with
 one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 35 heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

or R^8 and R^9 together with the nitrogen atom to which they are bonded form a five, six or seven membered heterocyclic ring optionally substituted with one or more R^6 substituents;

5 R^{10} is

- 1) halogen,
- 2) NO_2 ,
- 3) CN,
- 4) $B(OR^{13})(OR^{14})$,
- 10 5) C_1-C_6 alkyl,
- 6) C_2-C_6 alkenyl,
- 7) C_2-C_4 alkynyl,
- 8) C_3-C_7 cycloalkyl,
- 9) C_3-C_7 cycloalkenyl,
- 15 10) haloalkyl,
- 11) OR^7 ,
- 12) NR^8R^9 ,
- 13) SR^7 ,
- 14) COR^7 ,
- 20 15) $C(O)OR^7$,
- 16) $S(O)_mR^7$,
- 17) $CONR^8R^9$,
- 18) $S(O)_2NR^8R^9$,
- 19) aryl,
- 25 20) heteroaryl,
- 21) heterocyclyl, or
- 22) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, and cycloalkenyl is optionally substituted with one or more R^6 substituents;

30

R^{11} is

- 1) haloalkyl,
- 2) C_1-C_6 alkyl,
- 3) C_2-C_6 alkenyl,
- 35 4) C_2-C_4 alkynyl,

- 5) C₃–C₇ cycloalkyl,
6) C₃–C₇ cycloalkenyl,
7) aryl,
8) heteroaryl,
5 9) heterocyclyl, or
10) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

10 R¹² is

- 1) haloalkyl,
2) C₁–C₆ alkyl,
3) C₂–C₆ alkenyl,
15 4) C₂–C₄ alkynyl,
5) C₃–C₇ cycloalkyl,
6) C₃–C₇ cycloalkenyl,
7) aryl,
8) heteroaryl,
20 9) heterocyclyl,
10) heterobicyclyl,
11) C(O)–R¹¹,
12) C(O)O–R¹¹,
13) C(O)NR⁸R⁹,
25 14) S(O)_m–R¹¹, or
15) C(=Y)NR⁸R⁹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

30 R¹³ and R¹⁴ are each independently

- 1) H, or
2) C₁–C₆ alkyl; or

R¹³ and R¹⁴ are combined to form a heterocyclic ring or a heterobicyclyl ring;

R^{20} is

- 1) H,
- 2) NH_2 , or
- 3) $NHFmoc$;

5

or a prodrug; or the compound of Formula I or II is labeled with a detectable label or an affinity tag.

2. The compound, according to claim 1, in which A and A^1 are both CH_2 .

10

3. The compound, according to claim 1, in which A and A^1 are both $C=O$.

4. The compound, according to claim 1, in which A is CH_2 and A^1 is $C=O$.

15


5. The compound, according to claim 1, in which A and A^1 are both $C(O)OCH_3$.



6. The compound, according to claim 1, in which A and A^1 are both $C(O)OH$.

7. The compound, according to claim 1, comprises compounds of Formula 1A

20

through 1C:







5

wherein BG, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, and R³⁰⁰ are as defined in claim 1.

8. The compound, according to claim 1, comprise compounds of Formula 2A and 2B:




10





wherein BG, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, and R²⁰ are as defined in claim 1.

9. The compound, according to claim 1, in which BG is -X-L-X¹-.

10. The compound, according to claim 1, comprises compounds of Formula 1a through 1c:







5

wherein L, X, X¹, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, and R³⁰⁰ are as defined in claim 1.


11. The compound, according to claim 1, comprises compounds of Formula 1.1a through 1.1c:



1.1a




1.1b



1.1c

wherein L, X, X¹, A, A¹, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, R⁴, R⁴⁰⁰, R⁵ and R⁵⁰⁰ are as defined in claim 1.






12. The compound, according to claim 1, comprises compounds of Formula 2a:



wherein L, X, X¹, A, A¹, B, Q, Q¹, R¹, R¹⁰⁰, R², R²⁰⁰, R³ and R²⁰ are as defined in claim 1.



- 5 13. The compound, according to claim 1, in which X and X¹ are independently selected from

- 1) O,
 2) NR¹³,
 3) S,
 4) -C₁-C₆ alkyl-O-,
 5) -C₁-C₆ alkyl,



14. The compound, according to claim 13, in which X and X¹ are independently selected from:

1) O,



15. The compound, according to claim 1, in which L is selected from:

1) -C₁-C₂₀ alkyl-,

2) -C₃-C₇ cycloalkyl-,

3) -aryl-,



4) -biphenyl-,

5) -heteroaryl-,

6) -C₁-C₆ alkyl-(C₂-C₄ alkynyl)-C₁-C₆ alkyl-


7) -C₁-C₆ alkyl-aryl-C₁-C₆ alkyl-,

8) -aryl-Y-aryl-,




wherein the alkyl and cycloalkyl are optionally substituted with one or more R⁶ substituents, and the aryl, biphenyl and heteroaryl are optionally substituted with one or more R¹⁰ substituents.

16. The compound, according to claim 15, in which L is selected from the group consisting of:






17. The compound, according to claim 16, in which r is an integer of 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10.





10 18. The compound, according to claim 1, comprises compounds of Formula 1.1 through 1.18:




1.1


15









1.9




1.10




1.11



1.12




1.13




218


1.14




1.15



1.16





1.17



1.18

wherein r is an integer of 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10, and A , A^1 , Q , Q^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 and R^{300} are as defined in claim 1.

19. The compound, according to claim 1, comprises compounds of Formula 2.1 and
5 2.2:



wherein A , A^1 , Q , Q^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} and R^{20} are as defined in claim 1.

20. The compound, according to claim 1, in which R^1 and R^{100} are both H.
21. The compound, according to claim 1, in which R^1 and R^{100} are both C_1 - C_6 alkyl.
- 5 22. The compound, according to claim 21, in which R^1 and R^{100} are both CH_3 .
23. The compound, according to claim 1, in which R^2 and R^{200} are both C_1 - C_6 alkyl optionally substituted with OH.
- 10 24. The compound, according to claim 23, in which R^2 and R^{200} are both CH_3 .
25. The compound, according to claim 23, in which R^2 is CH_2OH and R^{300} is CH_3 .
26. The compound, according to claim 23, in which R^2 and R^{200} are both CH_2OH .
- 15 27. The compound, according to claim 23, in which R^2 and R^{200} are both CH_2CH_3 .
28. The compound, according to claim 1, in which R^3 and R^{300} are both C_1 - C_6 alkyl.
- 20 29. The compound, according to claim 28, in which R^3 and R^{300} are both $C(CH_3)_3$.
30. The compound, according to claim 1, in which Q and Q^1 are both NR^4R^5 .
31. The compound, according to claim 1, in which A and A^1 are both $C=O$, Q and Q^1
 25 are both NR^4R^5 , R^4 is H and
 R^5 is selected from
- 1) $\leftarrow C_1$ - C_6 alkyl,
 - 2) $\leftarrow C_2$ - C_6 alkenyl,
 - 3) $\leftarrow C_2$ - C_4 alkynyl,
 - 30 4) $\leftarrow C_3$ - C_7 cycloalkyl,
 - 5) $\leftarrow C_3$ - C_7 cycloalkenyl,
 - 6) $\leftarrow$ aryl,
 - 7) $\leftarrow$ heteroaryl,
 - 8) $\leftarrow$ heterocyclyl, or
 - 35 9) $\leftarrow$ heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents; wherein R^6 and R^{10} are as defined herein.

5

32. The compound, according to claim 31, in which R^4 is H and R^5 is selected from:

1) $\leftarrow C_1-C_6$ alkyl, or

2) $\leftarrow$ aryl,

wherein the alkyl is optionally substituted with one or two R^6 substituents; and wherein the

10


aryl is optionally substituted with one R^{10} substituent;


wherein R^6 and R^{10} are as defined herein.

33. The compound, according to claim 32, in which R^4 is H and R^5 is selected from the


15

group consisting of:







34. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




35. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




10 36. The compound, according to claim 1, in which A and A¹ are both C=O, and Q




37. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




38. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




39. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




40. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




41. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




10 42. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




43. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




44. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




45. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




46. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




47. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and



10 48. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




49. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




15


50. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




51. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




52. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




53. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




10 54. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




55. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and




15 56. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and
Q¹ are both



57. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and Q¹ are both



58. The compound, according to claim 1, in which A and A¹ are both C=O, and Q and Q¹ are both



59. The compound, according to claim 1, in which A and A¹ are both CH₂ and Q and Q¹ are both NR⁴R⁵, then R⁴ and R⁵ are each independently

1) haloalkyl,

2) ←C₁–C₆ alkyl,

3) ←C₂–C₆ alkenyl,

4) ←C₂–C₄ alkynyl,

5) ←C₃–C₇ cycloalkyl,

6) ←C₃–C₇ cycloalkenyl,

7) ←aryl,

8) ←heteroaryl,

9) ←heterocyclyl,

10) ←heterobicyclyl,

11) ←C(O)–R¹¹,

12) ←C(O)O–R¹¹,

13) ←C(=Y)NR⁸R⁹, or

14) ←S(O)₂–R¹¹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicycyl is optionally substituted with one or more R^{10} substituents; wherein Y, R^6 , R^8 , R^9 , R^{10} and R^{11} are as defined herein.


5

60. The compound, according to claim 59, in which R^4 and R^5 are independently selected from


- 1) C_1-C_6 alkyl,
- 2) $\leftarrow C(O)-R^{11}$,
- 10 3) $\leftarrow C(O)O-R^{11}$, or
- 4) $\leftarrow S(O)_2-R^{11}$,

wherein the alkyl is substituted with an R^6 substituent;
wherein R^6 and R^{11} are as defined herein.

15 61. The compound, according to claim 60, in which R^4 is $S(O)_2CH_3$ and R^5 is





62. The compound, according to claim 60, in which R^4 is $C(O)CH_3$ and R^5 is



20

63. The compound, according to claim 60, in which R^4 is

and R^5 is

25 64. The compound, according to claim 1, in which R^{11} is

- 1) C_1-C_6 alkyl, or
- 2) aryl,

wherein the alkyl is optionally substituted with one or more R^6 substituents; and wherein the aryl is optionally substituted with one or more R^{10} substituents; wherein R^6 and R^{10} are as defined herein.

- 5 65. The compound, according to claim 64, in which R^{11} is
- 1) C_1-C_6 alkyl optionally substituted with one or two R^6 substituents, or
 - 2) phenyl optionally substituted with one R^{10} substituent;
- wherein the R^6 and the R^{10} substituents are as defined herein.

- 10 66. The compound, according to claim 1, in which R^6 is

- 1) halogen,
- 2) NO_2 ,
- 3) CN,
- 4) aryl,
- 15 5) heteroaryl,
- 6) heterocyclyl,
- 7) heterobicyclyl,
- 8) OR^7 ,
- 9) SR^7 , or
- 20 10) NR^8R^9 ,

wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents; wherein R^7 , R^8 , R^9 and R^{10} are as defined herein.

- 25 67. The compound, according to claim 66, in which R^6 is

- 1) halogen,
- 2) aryl, or
- 3) NR^8R^9 ,

wherein the aryl is optionally substituted with one R^{10} substituent;

- 30 wherein R^8 , R^9 and R^{10} are as defined herein.

68. The compound, according to claim 1, in which R^8 and R^9 are each independently

- 1) H,
- 2) haloalkyl,
- 35 3) C_1-C_6 alkyl,

- 4) C₂-C₆ alkenyl,
- 5) C₂-C₄ alkynyl,
- 6) C₃-C₇ cycloalkyl, or
- 7) C₃-C₇ cycloalkenyl,

5 wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R⁶ substituents;
wherein the R⁶ substituents are as defined herein.


69. The compound, according to claim 1, in which R¹⁰ is




- 10 1) halogen,
- 2) NO₂,
- 3) CN,
- 4) haloalkyl,
- 5) OR⁷,
- 15 6) NR⁸R⁹, or
- 7) SR⁷;



wherein R⁷, R⁸, and R⁹ are as defined herein.




70. A compound, according to claim 1, selected from the group consisting of:




20

Cpd	STRUCTURE
1	





Cpd	STRUCTURE
2	
3	
4	




Cpd	STRUCTURE
5	 Chemical structure of compound 5, a complex molecule featuring a central linker with two tetrahydrofuran rings connected by a diene system. The linker is substituted with various functional groups, including amide bonds, hydroxyl groups, and a bicyclic amine moiety.
10	 Chemical structure of compound 10, a complex molecule featuring a central linker with two tetrahydrofuran rings connected by a diene system. The linker is substituted with various functional groups, including amide bonds, hydroxyl groups, and a bicyclic amine moiety.

Cpd	STRUCTURE
11	 Chemical structure of compound 11, a complex molecule featuring two 1,2,3,4-tetrahydronaphthalen-1-yl groups, two 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl groups, and a central 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl group. The structure is highly symmetrical and includes multiple amide and ester linkages.
12	 Chemical structure of compound 12, a complex molecule featuring two 1,2,3,4-tetrahydronaphthalen-1-yl groups, two 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl groups, and a central 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl group. The structure is highly symmetrical and includes multiple amide and ester linkages.
13	 Chemical structure of compound 13, a complex molecule featuring two 1,2,3,4-tetrahydronaphthalen-1-yl groups, two 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl groups, and a central 2,2,4,4-tetramethyl-5-oxo-1,3-dioxane-3-carbonyl group. The structure is highly symmetrical and includes multiple amide and ester linkages.




Cpd	STRUCTURE
14	 Chemical structure of compound 14: A complex molecule featuring a central pyrrolidine ring substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group, a 4-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group, and a 1-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group. The molecule also contains a 1,2,3,4-tetrahydronaphthalen-1-yl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group.
15	 Chemical structure of compound 15: A complex molecule featuring a central pyrrolidine ring substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group, a 4-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group, and a 1-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group. The molecule also contains a 1,2,3,4-tetrahydronaphthalen-1-yl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group.
16	 Chemical structure of compound 16: A complex molecule featuring a central pyrrolidine ring substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group, a 4-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group, and a 1-(4-oxo-4,5-dihydro-1H-benzofuro[3,2-c]pyridin-2-yl)butan-1-yl group. The molecule also contains a 1,2,3,4-tetrahydronaphthalen-1-yl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group.





Cpd	STRUCTURE
17	
18	
19	





Cpd	STRUCTURE
20	 Chemical structure of compound 20: A symmetrical molecule featuring a central benzene ring connected via amide bonds to two 4-membered pyrrolidine rings. Each pyrrolidine ring is substituted with a tert-butyl group, a (S)-1-amino-2-methylpropan-2-yl group, and a 2-phenylethyl group.
21	 Chemical structure of compound 21: A symmetrical molecule featuring a central benzene ring connected via amide bonds to two 4-membered pyrrolidine rings. Each pyrrolidine ring is substituted with a tert-butyl group, a (S)-1-amino-2-methylpropan-2-yl group, and a (S)-1-phenylethyl group.
22	 Chemical structure of compound 22: A symmetrical molecule featuring a central benzene ring connected via amide bonds to two 4-membered pyrrolidine rings. Each pyrrolidine ring is substituted with a tert-butyl group, a (S)-1-amino-2-methylpropan-2-yl group, and an isopropyl group.
23	 Chemical structure of compound 23: A symmetrical molecule featuring a central benzene ring connected via amide bonds to two 4-membered pyrrolidine rings. Each pyrrolidine ring is substituted with a tert-butyl group, a (S)-1-amino-2-methylpropan-2-yl group, and a 1,2,3,4-tetrahydronaphthalen-1-yl group.





Cpd	STRUCTURE
24	
25	
26	




Cpd	STRUCTURE
27	
28	
29	




Cpd	STRUCTURE
30	 Chemical structure of compound 30: A symmetrical molecule featuring two 4-fluorophenyl groups connected via a central benzene ring. The structure includes two piperidine rings, each substituted with a tert-butyl group and a methyl sulfonamide group. The piperidine rings are linked to the central benzene ring via amide bonds. The molecule also contains two chiral centers, each bearing a methyl group and a methyl sulfonamide group.
31	 Chemical structure of compound 31: A symmetrical molecule featuring two biphenyl groups connected via a central benzene ring. The structure includes two piperidine rings, each substituted with a tert-butyl group and a methyl sulfonamide group. The piperidine rings are linked to the central benzene ring via amide bonds. The molecule also contains two chiral centers, each bearing a methyl group and a methyl sulfonamide group.
32	 Chemical structure of compound 32: A symmetrical molecule featuring two naphthalenyl groups connected via a central benzene ring. The structure includes two piperidine rings, each substituted with a tert-butyl group and a methyl sulfonamide group. The piperidine rings are linked to the central benzene ring via amide bonds. The molecule also contains two chiral centers, each bearing a methyl group and a methyl sulfonamide group.




Cpd	STRUCTURE
33	 Chemical structure of compound 33: A complex molecule featuring a central piperidine ring. The piperidine ring is substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 2-position, a 4-sulfamoylphenyl group at the 3-position, and a 2-((2S,3S)-2-((2S,3S)-2-((2S)-1-aminopropan-2-yl)-3-oxo-3-(propan-2-ylamino)butanamide)butanamide)propanamide group at the 4-position. The piperidine ring is also substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 5-position.
34	 Chemical structure of compound 34: A complex molecule featuring a central piperidine ring. The piperidine ring is substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 2-position, a 4-sulfamoylphenyl group at the 3-position, and a 2-((2S,3S)-2-((2S,3S)-2-((2S)-1-aminopropan-2-yl)-3-oxo-3-(propan-2-ylamino)butanamide)butanamide)propanamide group at the 4-position. The piperidine ring is also substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 5-position.
35	 Chemical structure of compound 35: A complex molecule featuring a central piperidine ring. The piperidine ring is substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 2-position, a 4-sulfamoylphenyl group at the 3-position, and a 2-((2S,3S)-2-((2S,3S)-2-((2S)-1-aminopropan-2-yl)-3-oxo-3-(propan-2-ylamino)butanamide)butanamide)propanamide group at the 4-position. The piperidine ring is also substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 5-position.
36	 Chemical structure of compound 36: A complex molecule featuring a central piperidine ring. The piperidine ring is substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 2-position, a 4-sulfamoylphenyl group at the 3-position, and a 2-((2S,3S)-2-((2S,3S)-2-((2S)-1-aminopropan-2-yl)-3-oxo-3-(propan-2-ylamino)butanamide)butanamide)propanamide group at the 4-position. The piperidine ring is also substituted with a 1,2,3,4-tetrahydronaphthalen-1-yl group at the 5-position.




Cpd	STRUCTURE
37	
38	
39	
40	

Cpd	STRUCTURE
41	
42	
43	
44	

Cpd	STRUCTURE
45	 Chemical structure of compound 45, a complex molecule featuring multiple amide bonds, a central benzene ring, and a terminal amine group. The structure includes a 1,2,3,4-tetrahydronaphthalene moiety at the top, a central benzene ring, and a terminal amine group. The molecule is highly branched and contains several chiral centers indicated by wedged and dashed bonds.
46	 Chemical structure of compound 46, a complex molecule featuring multiple amide bonds, a central benzene ring, and a terminal amine group. The structure includes a 1,2,3,4-tetrahydronaphthalene moiety at the top, a central benzene ring, and a terminal amine group. The molecule is highly branched and contains several chiral centers indicated by wedged and dashed bonds.
47	 Chemical structure of compound 47, a complex molecule featuring multiple amide bonds, a central benzene ring, and a terminal amine group. The structure includes a 1,2,3,4-tetrahydronaphthalene moiety at the top, a central benzene ring, and a terminal amine group. The molecule is highly branched and contains several chiral centers indicated by wedged and dashed bonds.




Cpd	STRUCTURE
48	 Chemical structure of compound 48: A complex molecule featuring two piperidine rings connected by a central benzoyl group. The left piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The right piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The central benzoyl group is connected to the piperidine rings via amide bonds. The structure also includes a biphenyl group attached to the right piperidine ring.
49	 Chemical structure of compound 49: A complex molecule featuring two piperidine rings connected by a central benzoyl group. The left piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The right piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The central benzoyl group is connected to the piperidine rings via amide bonds. The structure also includes a biphenyl group attached to the right piperidine ring.
50	 Chemical structure of compound 50: A complex molecule featuring two piperidine rings connected by a central benzoyl group. The left piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The right piperidine ring is substituted with a (S)-1-aminopropan-2-yl group and a (S)-1-((2-aminopropan-2-ylideneamino)oxy)propan-2-yl group. The central benzoyl group is connected to the piperidine rings via amide bonds. The structure also includes a biphenyl group attached to the right piperidine ring.



Cpd	STRUCTURE
51	 Chemical structure of compound 51, a complex molecule featuring multiple amide bonds, a central benzene ring, and several substituted phenyl groups, including a 4-methoxyphenyl group and a 4-methoxy-2-methylphenyl group.
52	 Chemical structure of compound 52, a complex molecule featuring multiple amide bonds, a central benzene ring, and several substituted phenyl groups, including a 4-methoxyphenyl group and a 4-methoxy-2-methylphenyl group.
53	 Chemical structure of compound 53, a complex molecule featuring multiple amide bonds, a central benzene ring, and several substituted phenyl groups, including a 4-methoxyphenyl group and a 4-methoxy-2-methylphenyl group.

Cpd	STRUCTURE
54	 Chemical structure of compound 54, a complex molecule featuring a central pyrrolidine ring substituted with a 4-aminobenzoyl group, a 2-((S)-1-hydroxy-1-phenylethyl)amino group, and a 2-((S)-1-hydroxy-1-phenylethyl)amino group. The molecule also includes a 2-((S)-1-hydroxy-1-phenylethyl)amino group and a 2-((S)-1-hydroxy-1-phenylethyl)amino group.
55	 Chemical structure of compound 55, a complex molecule featuring a central pyrrolidine ring substituted with a 4-aminobenzoyl group, a 2-((S)-1-hydroxy-1-phenylethyl)amino group, and a 2-((S)-1-hydroxy-1-phenylethyl)amino group. The molecule also includes a 2-((S)-1-hydroxy-1-phenylethyl)amino group and a 2-((S)-1-hydroxy-1-phenylethyl)amino group.
56	 Chemical structure of compound 56, a complex molecule featuring a central pyrrolidine ring substituted with a 4-aminobenzoyl group, a 2-((S)-1-hydroxy-1-phenylethyl)amino group, and a 2-((S)-1-hydroxy-1-phenylethyl)amino group. The molecule also includes a 2-((S)-1-hydroxy-1-phenylethyl)amino group and a 2-((S)-1-hydroxy-1-phenylethyl)amino group.

[illegible]

Cpd	STRUCTURE
60	
61	
62	
63	



Cpd	STRUCTURE
	 Chemical structure of compound 63, a complex molecule featuring multiple amide and ester linkages, a central benzene ring, and various side chains including a methoxy-substituted benzyl group and a dimethylamino-substituted side chain.
64	 Chemical structure of compound 64, which is identical to compound 63 but with the methoxy group on the benzyl side chain replaced by a benzyl group (Ph).
65	 Chemical structure of compound 65, a complex molecule featuring multiple amide and ester linkages, a central benzene ring, and various side chains including a benzyl group, a dimethylamino-substituted side chain, and a carboxamide group (H₂N).

Cpd	STRUCTURE
66	 Chemical structure of compound 66: A complex molecule featuring a central benzamide core. The benzamide is substituted with a 4-methylphenyl group and a 4-methoxyphenyl group. The amide nitrogen is part of a 5-membered ring system, which is further substituted with a 4-methylphenyl group and a 4-methoxyphenyl group. The structure includes multiple amide and ester linkages, and a central benzamide moiety.
67	 Chemical structure of compound 67: A complex molecule featuring a central benzamide core. The benzamide is substituted with a 4-methylphenyl group and a 4-methoxyphenyl group. The amide nitrogen is part of a 5-membered ring system, which is further substituted with a 4-methylphenyl group and a 4-methoxyphenyl group. The structure includes multiple amide and ester linkages, and a central benzamide moiety.


71. A compound, according to claim 1, selected from the group consisting of:

5


10

Cpd	Structure
6	 Chemical structure of compound 6, a complex molecule featuring a central alkyne linker. The alkyne is connected via ether linkages to two distinct peptide-like structures. The upper structure includes a benzyl group, a benzamide, and a dimethylamino group. The lower structure includes a fluorenyl group, a tert-butyl group, and a dimethylamino group. Stereochemistry is indicated with wedges and dashes.
7	 Chemical structure of compound 7, a complex molecule featuring a central amide linker. The amide is connected via ether linkages to two distinct peptide-like structures. The upper structure includes a fluorenyl group, a tert-butyl group, and a dimethylamino group. The lower structure includes a fluorenyl group, a tert-butyl group, and a dimethylamino group. Stereochemistry is indicated with wedges and dashes.

; and

Cpd	Structure
8	


72. An intermediate compound represented by Formula 1-iv:



5


wherein PG³, R¹, R², R³, A, and Q are as defined in claim 1.

73. An intermediate compound represented by Formula 2-iv:



wherein PG^4 , R^1 , R^2 , R^3 , A, and Q are as defined in claim 1.


- 5 74. An intermediate compound represented by Formula 3-ii:



wherein PG^4 , PG^{400} , R^1 , R^2 , R^3 , A, and Q are as defined in claim 1, and L is $-(CH_2)_r-$, $-(CH_2)_r-Y-(CH_2)_r-$, -alkyl-aryl-alkyl-, -alkyl-heteroaryl-alkyl-, cycloalkyl, aryl or heteroaryl.

10


75. An intermediate compound represented by Formula 4-i:



4-i

wherein PG⁴, PG⁴⁰⁰, L, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, A, A¹, X, X¹, Q and Q¹ are as defined in claim 1.


- 5 76. An intermediate compound represented by Formula 6-ii:



6-ii


wherein PG⁴, PG⁴⁰⁰, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, A, A¹, Q and Q¹ are as defined in claim 1.

- 10 77. An intermediate compound represented by Formula 7-v:




wherein PG⁴, PG⁴⁰⁰, L, R¹, R¹⁰⁰, R², R²⁰⁰, R³, R³⁰⁰, A, A¹, Q and Q¹ are as defined in claim 1.

- 5 78. An intermediate compound represented by Formula 8-ii:




wherein PG⁴, r, L, R¹⁰⁰, R²⁰⁰, A¹, and Q¹ are as defined in claim 1.

79. An intermediate compound represented by Formula 8-iii:



wherein PG^4 , PG^{400} , r , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined in claim 1.


80. An intermediate compound represented by Formula 17-i:



5

wherein PG^4 , PG^{400} , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined in claim 1.


81. An intermediate compound represented by Formula 18-i:



10


wherein PG^4 , PG^{400} , L , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , A , A^1 , Q and Q^1 are as defined in claim 1.

82. An intermediate compound represented by Formula 19-8:



wherein PG^4 , PG^{400} , L , X , X^1 , R^1 , R^{100} , R^2 , R^{200} , R^3 , R^{300} , R^4 , R^{400} , R^5 and R^{500} are as defined in claim 1.


- 5 83. An intermediate compound represented by Formula 19-3:



wherein PG^2 , L , and X are as defined in claim 1.


10

84. An intermediate compound represented by Formula 20-2:



wherein L, X, X¹, R⁴ and R⁵ are as defined in claim 1.


85. An intermediate compound represented by Formula 20-4:



wherein L, X, X¹, R³, R⁴, and R⁵ are as defined in claim 1.

86. A process for producing compounds represented by Formula I, described hereinabove, the process comprising:

a) coupling two intermediates represented by Formula 1(iv):




wherein PG³, R¹, R², R³, A, and Q are as defined herein,
in a solvent; and

b) removing the protecting groups so as to form compounds of Formula I.

87. A process for producing compounds represented by Formula I, described hereinabove, the process comprising:

5

a) coupling an intermediate represented by Formula 2(iv):




wherein PG⁴ is a protecting group, and R¹, R², R³, A, and Q are as defined herein.


10 and an activated diacid (0.5 equiv) in a solvent; and

b) removing the protecting groups so as to form compounds of Formula I.

88. Use of a compound represented by Formula I or II:



15



or a salt thereof, for the manufacture of a medicament for treating or preventing a disease state characterized by insufficient apoptosis,

wherein:

m is 0, 1 or 2;

5 Y is NH, O or S;

BG is

1) $-X-L-X^1-$;

10 X and X^1 are independently selected from

1) O,

2) NR^{13} ,








3) S,


4) $-C_1-C_6$ alkyl-,

15 5) $-C_1-C_6$ alkyl-O,

6) $-C_1-C_6$ alkyl- NR^{13} -,



7) $-C_1-C_6$ alkyl-S-,





L is selected from:

- 1) $-C_1-C_{20}$ alkyl-,
- 2) $-C_2-C_6$ alkenyl-,
- 3) $-C_2-C_4$ alkynyl-,
- 4) $-C_3-C_7$ cycloalkyl-,
- 5) -aryl-,
- 6) -biphenyl-,
- 7) -heteroaryl-,
- 8) -heterocyclyl-,
- 9) $-C_1-C_6$ alkyl- $(C_2-C_6$ alkenyl)- C_1-C_6 alkyl-,
- 10) $-C_1-C_6$ alkyl- $(C_2-C_4$ alkynyl)- C_1-C_6 alkyl-,
- 11) $-C_1-C_6$ alkyl- $(C_3-C_7$ cycloalkyl)- C_1-C_6 alkyl-,
- 12) $-C_1-C_6$ alkyl-aryl- C_1-C_6 alkyl-,
- 13) $-C_1-C_6$ alkyl-biphenyl- C_1-C_6 alkyl-,
- 14) $-C_1-C_6$ alkyl-heteroaryl- C_1-C_6 alkyl-,
- 15) $-C_1-C_6$ alkyl-heterocyclyl- C_1-C_6 alkyl-,
- 16) $-C_1-C_6$ alkyl-Y- C_1-C_6 alkyl-,
- 17) -aryl-Y-aryl-,
- 18) -heteroaryl-Y-heteroaryl-,
- 19) -heterocyclyl-Y-heterocyclyl-,



- 25 wherein the alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more R^6 substituents, and the aryl, biphenyl, heteroaryl, and heterocyclyl are optionally substituted with one or more R^{10} substituents;

Q and Q^1 are independently selected from

- 1) NR^4R^5 ,
- 2) OR^{11} , or
- 3) $\text{S}(\text{O})_m\text{R}^{11}$; or

5 Q and Q^1 are independently selected from

- 1) aryl, or
- 2) heteroaryl, the aryl and the heteroaryl being optionally substituted with one or more R^{10} substituents;

10 A and A^1 are independently selected from

- 1) $-\text{CH}_2-$,
- 2) $-\text{CH}_2\text{CH}_2-$,
- 3) $-\text{CH}(\text{C}_1-\text{C}_6 \text{ alkyl})-$,
- 4) $-\text{CH}(\text{C}_3-\text{C}_7 \text{ cycloalkyl})-$,
- 15 5) $-\text{C}_3-\text{C}_7 \text{ cycloalkyl}-$,
- 6) $-\text{CH}(\text{C}_1-\text{C}_6 \text{ alkyl}-\text{C}_3-\text{C}_7 \text{ cycloalkyl})-$,
- 7) $-\text{C}(\text{O})-$, or
- 8) $-\text{C}(\text{O})\text{OR}^{13}$;

20 R^1 and R^{100} are independently selected from

- 1) H, or
- 2) C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

R^2 and R^{200} are independently selected from

- 25 1) H, or
- 2) C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

R^3 and R^{300} are independently C_1-C_6 alkyl optionally substituted with one or more R^6 substituents;

30

R^4 and R^5 are each independently selected from

- 1) H,
- 2) haloalkyl,
- 3) $\leftarrow\text{C}_1-\text{C}_6 \text{ alkyl}$,
- 35 4) $\leftarrow\text{C}_2-\text{C}_6 \text{ alkenyl}$,

- 5) $\leftarrow$ C₂-C₄ alkynyl,
 6) $\leftarrow$ C₃-C₇ cycloalkyl,
 7) $\leftarrow$ C₃-C₇ cycloalkenyl,
 8) $\leftarrow$ aryl,
 5 9) $\leftarrow$ heteroaryl,
 10) $\leftarrow$ heterocyclyl,
 11) $\leftarrow$ heterobicyclyl,
 12) $\leftarrow$ C(O)-R¹¹,
 13) $\leftarrow$ C(O)O-R¹¹,
 10 14) $\leftarrow$ C(=Y)NR⁸R⁹, or
 15) $\leftarrow$ S(O)₂-R¹¹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

15

R⁶ is

- 1) halogen,
 2) NO₂,
 3) CN,
 20 4) haloalkyl,
 5) C₁-C₆ alkyl,
 6) C₂-C₆ alkenyl,
 7) C₂-C₄ alkynyl,
 8) C₃-C₇ cycloalkyl,
 25 9) C₃-C₇ cycloalkenyl,
 10) aryl,
 11) heteroaryl,
 12) heterocyclyl,
 13) heterobicyclyl,
 30 14) OR⁷,
 15) S(O)_mR⁷,
 16) NR⁸R⁹,
 17) NR⁸S(O)₂R¹¹,
 18) COR⁷,
 35 19) C(O)OR⁷,

- 20) CONR^8R^9 ,
- 21) $\text{S(O)}_2\text{NR}^8\text{R}^9$
- 22) OC(O)R^7 ,
- 23) OC(O)Y-R^{11} ,
- 5 24) SC(O)R^7 , or
- 25) $\text{NC(Y)NR}^8\text{R}^9$,

wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

10 R^7 is

- 1) H,
- 2) haloalkyl,
- 3) $\text{C}_1\text{--C}_6$ alkyl,
- 4) $\text{C}_2\text{--C}_6$ alkenyl,
- 15 5) $\text{C}_2\text{--C}_4$ alkynyl,
- 6) $\text{C}_3\text{--C}_7$ cycloalkyl,
- 7) $\text{C}_3\text{--C}_7$ cycloalkenyl,
- 8) aryl,
- 9) heteroaryl,
- 20 10) heterocyclyl,
- 11) heterobicyclyl,
- 12) $\text{R}^8\text{R}^9\text{NC(=Y)}$, or
- 13) $\text{C}_1\text{--C}_6$ alkyl- $\text{C}_2\text{--C}_4$ alkenyl, or
- 14) $\text{C}_1\text{--C}_6$ alkyl- $\text{C}_2\text{--C}_4$ alkynyl,

25 wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and heterobicyclyl is optionally substituted with one or more R^{10} substituents;

R^8 and R^9 are each independently

- 30 1) H,
- 2) haloalkyl,
- 3) $\text{C}_1\text{--C}_6$ alkyl,
- 4) $\text{C}_2\text{--C}_6$ alkenyl,
- 5) $\text{C}_2\text{--C}_4$ alkynyl,
- 35 6) $\text{C}_3\text{--C}_7$ cycloalkyl,

- 7) C₃-C₇ cycloalkenyl,
- 8) aryl,
- 9) heteroaryl,
- 10) heterocyclyl,
- 5 11) heterobicyclyl,
- 12) C(O)R¹¹,
- 13) C(O)Y-R¹¹, or
- 14) S(O)₂-R¹¹,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with
 10 one or more R⁶ substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 heterobicyclyl is optionally substituted with one or more R¹⁰ substituents;

or R⁸ and R⁹ together with the nitrogen atom to which they are bonded form a five, six or
 seven membered heterocyclic ring optionally substituted with one or more R⁶ substituents;

15 R¹⁰ is

- 1) halogen,
- 2) NO₂,
- 3) CN,
- 20 4) B(OR¹³)(OR¹⁴),
- 5) C₁-C₆ alkyl,
- 6) C₂-C₆ alkenyl,
- 7) C₂-C₄ alkynyl,
- 8) C₃-C₇ cycloalkyl,
- 25 9) C₃-C₇ cycloalkenyl,
- 10) haloalkyl,
- 11) OR⁷,
- 12) NR⁸R⁹,
- 13) SR⁷,
- 30 14) COR⁷,
- 15) C(O)OR⁷,
- 16) S(O)_mR⁷,
- 17) CONR⁸R⁹,
- 18) S(O)₂NR⁸R⁹,
- 35 19) aryl,

- 20) heteroaryl,
- 21) heterocyclyl, or
- 22) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, and cycloalkenyl is optionally substituted
 5 with one or more R^6 substituents;

R^{11} is

- 1) haloalkyl,
- 2) C_1-C_6 alkyl,
- 10 3) C_2-C_6 alkenyl,
- 4) C_2-C_4 alkynyl,
- 5) C_3-C_7 cycloalkyl,
- 6) C_3-C_7 cycloalkenyl,
- 7) aryl,
- 15 8) heteroaryl,
- 9) heterocyclyl, or
- 10) heterobicyclyl,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with
 one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and
 20 heterobicyclyl is optionally substituted with one or more R^{10} substituents;

R^{12} is

- 1) haloalkyl,
- 2) C_1-C_6 alkyl,
- 25 3) C_2-C_6 alkenyl,
- 4) C_2-C_4 alkynyl,
- 5) C_3-C_7 cycloalkyl,
- 6) C_3-C_7 cycloalkenyl,
- 7) aryl,
- 30 8) heteroaryl,
- 9) heterocyclyl,
- 10) heterobicyclyl,
- 11) $C(O)-R^{11}$,
- 12) $C(O)O-R^{11}$,
- 35 13) $C(O)NR^8R^9$,

14) $S(O)_m-R^{11}$, or

15) $C(=Y)NR^8R^9$,

wherein the alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl is optionally substituted with one or more R^6 substituents; and wherein the aryl, heteroaryl, heterocyclyl, and

5 heterobicyclyl is optionally substituted with one or more R^{10} substituents;

R^{13} and R^{14} are each independently

1) H, or

2) C_1-C_6 alkyl; or

10 R^{13} and R^{14} are combined to form a heterocyclic ring or a heterobicyclyl ring; and

R^{20} is

1) H,

2) NH_2 , or

15 3) $NHFmoc$.

89. Use of the compound of any one of claims 1 to 71 for the manufacture of a medicament for treating or preventing a disease state characterized by insufficient apoptosis.

20

90. Use, according to claims 88 or 89, in which the disease state is cancer.

91. Use of the compound of any one of claims 1 to 71 for the manufacture of a medicament for treating or preventing a proliferative disorder.

25

92. Use of the compound of any one of claims 1 to 71 in combination with an agent for the manufacture of a medicament for treating or preventing a proliferative disorder, wherein the agent is selected from:

a) an estrogen receptor modulator,

30 b) an androgen receptor modulator,

c) retinoid receptor modulator,

d) a cytotoxic agent,

e) an antiproliferative agent,

f) a prenyl-protein transferase inhibitor,

- g) an HMG-CoA reductase inhibitor,
- h) an HIV protease inhibitor,
- i) a reverse transcriptase inhibitor,
- k) an angiogenesis inhibitor,
- 5 l) a PPAR- γ agonist,
- m) a PPAR- δ agonist,
- n) an inhibitor of inherent multidrug resistance,
- o) an anti-emetic agent,
- p) an agent useful in the treatment of anemia,
- 10 q) agents useful in the treatment of neutropenia,
- r) an immunologic-enhancing drug.
- s) a proteasome inhibitor;
- t) an HDAC inhibitor;
- u) an inhibitor of the chemotrypsin-like activity in the proteasome; or
- 15 v) E3 ligase inhibitors;
- w) a modulator of the immune system such as, but not limited to, interferon-alpha, Bacillus Calmette-Guerin (BCG), and ionizing radiation (UVB) that can induce the release of cytokines, such as the interleukins, TNF, or induce release of death receptor ligands such as TRAIL;
- 20 x) a modulator of death receptors TRAIL and TRAIL agonists such as the humanized antibodies HGS-ETR1 and HGS-ETR2;
- or in combination or sequentially with radiation therapy.

93. Use of the compound of any one of claims 1 to 71 in combination with a death
25 receptor agonist for the manufacture of a medicament the treatment or prevention of a proliferative disorder in a subject.

94. Use, according to claim 93, in which the death receptor agonist is TRAIL.

30 95. Use, according to claim 93, in which the death receptor agonist is a TRAIL antibody.

96. Use, according to claim 93, in which the death receptor agonist is in an amount that produces a synergistic effect.


97. Use, according to claims 92 or 93, in which the proliferative disorder is cancer.

98. A pharmaceutical composition comprising the compound of any one of claims 1 to 71, mixed with a pharmaceutically acceptable carrier, diluent or excipient, for treating or preventing a disease state characterized by insufficient apoptosis.

99. A pharmaceutical composition comprising the compound of any one of claims 1 to 71 in combination with any compound that increases the circulating level of one or more death receptor agonists for preventing or treating a proliferative disorder.


100. A method of preparing a pharmaceutical composition, the method comprising: mixing the compound of any one of claims 1 to 71, with a pharmaceutically acceptable carrier, diluent or excipient.

101. An intermediate compound represented by Formula 19-2:



wherein PG¹, PG², L, X are as defined herein

102. An intermediate compound represented by Formula 20-1a:



wherein PG¹, L, X, X¹, R⁴ and R⁵ are as defined herein.

- 5 103. A method for the preparation of a pharmaceutically acceptable salt of compound of Formula I and II, by the treatment of a compound of Formula I or II with 1 to 2 equiv of a pharmaceutically acceptable acid, as defined herein.
- 10 104. A method of detecting loss of function or suppression of IAPs in vivo, the method comprising: a) administering to a subject, a therapeutically effective amount of a pharmaceutical composition, according to claim 88; b) isolating a tissue sample from the subject; and c) detecting a loss of function or suppression of IAPs from the sample.
- 15 105. The method, according to claim 104, wherein the IAP is XIAP.
106. The method, according to claim 104, wherein the IAP is c-IAP1.
107. The method, according to claim 104, wherein the IAP is c-IAP2.
- 20 108. The method, according to claim 104, wherein the tissue is a tumor sample.
109. The method according to claim 104, wherein the tissue is normal tissue.
- 25 110. The method, according to claim 104, wherein the tissue is isolated from the hematopoietic system.

111. The method, according to claim 104, wherein mononuclear cells are isolated from blood, and IAP suppression or loss is measured in the mononeuclear cells.