

ANTIVIRAL PRODRUGS, INTERMEDIATE- AND LONG-ACTING FORMULATIONS, AND METHODS

Cross-reference to Related Applications

5 This application claims priority to U.S. Provisional Patent Application No. 63/303,376, filed January 26, 2022, which is incorporated by reference herein.

Government Sponsorship

This invention was made with government support under grant no. AI131424 and
10 grant no. AI1161348 awarded by the National Institutes of Health. The government has
certain rights in the invention.

Technical Field

The present disclosure relates to antiviral prodrugs, methods for producing
15 antiviral prodrugs, methods of treatment, and pharmaceutical formulations, such as long-
or intermediate-acting formulations, of antiviral prodrugs for treatment of various
diseases, such as SARS CoV-2 infections or human immunodeficiency infections, in
mammals.

20 **Background**

Over the past 18 years, spillover events have introduced the highly transmissible beta-coronavirus strains SARS CoV, MERS CoV, SARS CoV-2 into the human population (*see, e.g.,* Zhong N.S., et. al. *Lancet* 2003;362:1353-1358; Zaki, A.M., et. al., *N Engl J Med* 2012;367:1814-1820; and Zhu N., et. al. *N Engl J Med.* 2020;382:727-733).

25 Although case fatality ratios have varied, each has demonstrated the ability to induce substantial morbidity and mortality – especially among those over 55 and/or those with underlying co-morbid medical conditions (*see, e.g.,* Wu C., et. al., *JAMA Intern Med* 2020; 180: 934-943; and Zhou F. et. al. *Lancet* 2020;395:1054-1062). Although SARS

30 CoV and MERS CoV were largely contained by epidemiological interventions, the current outbreak evolved into a global pandemic responsible for hundreds of millions of infections and over five million deaths (*see, e.g.,* Johns Hopkins Coronavirus Resource Center <https://coronavirus.jhu.edu/map.html> (accessed January 16, 2022). Intensive efforts to develop COVID vaccines have resulted in the development of a number of highly

effective vaccines that provide substantial levels of protection from severe disease and death (*see, e.g.*, Polack F.P. et.al. *N Engl J Med* 2021;383: 2603-2615, and Baden L.R. *N Engl J Med.* 2021;384:403-416). In the case of even the most effective vaccines, immunity begins to wane within 5 months (*see, e.g.*, Keehner J. et.al, *N Engl J Med.* 2021;384:1774-1775). Despite their effectiveness, vaccine hesitancy is widespread and up to 50% of people in many parts of the world have declined vaccination (*see, e.g.*, Daniel C.N. et. al. *Vaccine.* 2022:S0264-410X(21)01667-4). Substantial subgroups of the population are immunocompromised and/or have underlying conditions that limit vaccine effectiveness. Thus, despite major successes in vaccine development, the SARS CoV-2 infection rate remains high and COVID-related morbidity and mortality should be anticipated to persist in the years to come without effective antiviral therapeutics.

Antiviral therapeutic efforts have included both monoclonal antibodies directed at neutralizing epitopes on the viral spike protein and small molecules directed at the viral RNA-dependent RNA polymerase or the virus' 3-CL like protease enzyme. Remdesivir nucleoside triphosphate (RVn triphosphate) potently inhibits enzymatic activity of the polymerase of every coronavirus tested thus far, including SARS CoV-2. Remdesivir also inhibits the polymerases of a number of other pathogenic RNA viruses, including Ebola virus, Nipah virus and respiratory syncytial virus (*see, e.g.*, Wang M., et. al., *Cell Res* 2020;30:269-271; Yan V.C, e.g., *ACS Med Chem Lett.* 2020;11:1361-1366; and Mulangu S. et. al, *N Engl J Med* 2019;381:2293-2303).

Clinical trials of antiviral agents for COVID-19 (including remdesivir) have demonstrated maximal benefit when employed prior to hospitalization (*see, e.g.*, Gottlieb RL *New Engl J Med*, 2021 Dec 22: NEJMoa2116846). Remdesivir can reduce hospitalization or death by 85-90% when administered as three daily intravenous infusions prior to hospitalization. This approach, while highly effective, is extremely challenging to take to scale since it requires three sequential days of intravenous infusion of SARS CoV-2 infected patients. The approach, nonetheless, is a proof of concept that early remdesivir can have a substantial impact on the disease if employed early enough in the illness. The availability of a remdesivir prodrug that could be administered orally for a week or intramuscularly on a single occasion to outpatients with SARS CoV-2 who are at risk for severe COVID-19 disease would greatly extend the clinical reach of this agent in the treatment of COVID-19 and, potentially, for the treatment of Ebola, Nipah, and RSV

infection. Sustained release injectable formulations of remdesivir prodrugs are of interest because RDV is not highly bioavailable following oral administration and must be administered intravenously, thereby, in many instances, severely complicating its administration to high-risk patients prior to hospitalization with relatively advanced disease. A long-acting formulation of remdesivir nucleoside monophosphate, which might enable plasma antiviral activity above the 90% effective concentration for at least about 5 to about 10 days after a single intramuscular or subcutaneous dose, would be particularly useful.

Antiretroviral therapy (ART) has resulted in enormous reductions in HIV-1 related morbidity and mortality in the US and globally (*see, e.g.,* Walensky R.P. et. al., *J Infect Dis* 2006; 194:11–19; and April M.D., et. al. *J Infect Dis*.2014;209:491-9). By 2021, 28 million of the 38 million people with HIV infection were receiving antiretroviral therapy (*see, e.g.,* <https://www.unaids.org/en/resources/fact-sheet> (accessed January 19, 2022)). Of those in antiretroviral therapy, it is estimated that as many as 90% may be fully virally suppressed in resource rich settings. Since plasma HIV-1 RNA levels are not routinely measured in sub-Saharan Africa, the fraction of the HIV-1 infected population currently on fully effective therapy is not known but is likely substantially lower. Prevention of transmission of HIV-1 is also an urgent global priority. With the demonstration that the likelihood of transmission of HIV is directly related to plasma (and by extension genital) levels of HIV-1 (*see, e.g.* Quinn T.C., et.al., *N Engl J Med* 2000;342:921-929), there has been increasing optimism that the epidemic might be ended by an aggressive global expansion of antiretroviral therapy used for both treatment and prevention (*see, e.g.,* Granich R.M., et.al., *N Engl J Med* 2000;342:921-929). Indeed, reductions in incident infections related to treatment of HIV-1 infected individuals been demonstrated both in clinical trials and at the population level (*see, e.g.,* Cohen M.S., et.al., *N Engl J Med*. 2011;365:493-505; Montaner J.S., et.al., *Lancet* 2010; 376: 532–539; Montaner J.S., et.al., *PLoS One* 2014; 9: e87872; and Tanser J.F., et. al., *Science* 2013. 339: 966–971). One and a half million people were newly infected in 2020. Despite early optimism about the development of an effective AIDS vaccine, there are few real insights into how to elicit protective responses with scalable commercial vaccine products. Treatment of high risk, HIV-1 uninfected persons (termed pre-exposure prophylaxis or “PrEP”) has been effective in some populations, but not in others with protection rates ranging from 0 – 62% (*see, e.g.,* Grant

R.M. *N Engl J Med* 2010;363:2587-2599; Baeten J.M., et.al., *N Engl J Med* 2012;367:399-410; Thigpen M.C. *N Engl J Med*. 2012;367:423-434; Choopanya K., et.al., *Lancet* 2013;381:2083-90; Van Damme L., et. al., *N Engl J Med*. 2012;367:411-22; and Marrazzo J.M., et. al., *N Engl J Med* 2015;372:509-18). PrEP is effective if adherence is

5 high but persons at the highest risk for HIV-1 infection are often those facing the greatest barriers to adherence. The failure of the large FEM PrEP and VOICE trials to demonstrate any degree of protection despite concurrent assessments of 88 – 90% adherence by self-report and returned product counts has only underlined the adherence challenges posed by oral PrEP (*see, e.g.*, Van Damme L., et. al., *N Engl J Med*. 2012;367:411-22; and Marrazzo

10 J.M., et. al., *N Engl J Med* 2015;372:509-18). Nonetheless, the results from the oral and topical PrEP trials involving TFV were perhaps more consistent than they might seem: high levels of protection can be achieved when active concentrations of drug are present at the time of exposure. The dual benefits of ART that include improving the health of HIV-1 infected persons receiving therapy and reducing the likelihood of viral transmission have

15 made rapid expansion of antiretroviral therapy a major global priority (*see, e.g.*, UNAIDS. 90-90-90 An Ambitious Treatment Target to help end the AIDS epidemic. Geneva, Switzerland 2014. Available at: http://www.unaids.org/sites/default/files/media_asset/90-90-90_en_0.pdf).

With mitigation of many of the acute and chronic toxicities of first generation

20 ARVs by several currently available once-a-day single tablet regimens (STR), treatment adherence has emerged as the greatest barrier to long-term treatment success. Although most patients can adhere to daily regimens, many of the most vulnerable populations (for prevention and therapy) include those whose adherence is challenged by mental health issues, substance use, and stigma. The development of effective, inexpensive, well-

25 tolerated, convenient modalities that maintain active concentrations at sites of viral replication and exposure for prolonged periods of time would represent a major advance for ART for both treatment and prevention. Long-acting injectable products such as cabotegravir, rilpivavirine and islatravir have provided a proof of concept of this approach (*see, e.g.*, Margolis D.A., et.al., *Lancet Infect Dis*. 2015;15:1145-55; Verloes R., et.al., *HIV*

30 *Med*. 2015;16:477-84; and Matthers R.P., et.al., *Nature Med*. 2021; 27, 1712-1717). The recent decision of the FDA to place trials of islatravir on clinical hold because of leucopenia in a subset of patients underlines the need to develop agents within multiple


classes if long-acting ART is to be of widespread use in patient populations vulnerable to different toxicities and with viruses of differing drug susceptibility.

There remains a need for pharmaceutical formulations that can be delivered in various ways, such as intramuscularly. There also remains a need for pharmaceutical formulations that are long-acting, such as formulations that can provide effective plasma levels for extended periods after infrequent administration, such as once monthly.


Summary

Provided herein are compounds, such as antiviral prodrugs, and pharmaceutical formulations that overcome one or more of the disadvantages of currently used drugs and/or formulations. For example, examples of the compounds and pharmaceutical formulations provided herein include orally useful antiviral prodrugs that may specifically target organs where viral replication is maximal and be conveniently administered at scale in any disease stage. For oral use and enhanced lung exposure, some examples of the prodrugs of RVn or formulations provided herein can accomplish one or more of the following: 1) kinase bypass of the first nucleoside phosphorylation, 2) provide increased oral bioavailability, 3) deliver antivirally significant concentrations to lung and gastrointestinal tract and formulations which may provide for sustained levels in plasma for 5 to 30 days following a single injection. Also provided herein are methods for the synthesis and antiviral evaluation of the compounds, including novel lipophilic prodrugs of RVn-monophosphate that are substantially more active than remdesivir in Vero E6 cells infected with SARS-CoV-2. Not wishing to be bound by any particular theory, examples of the compounds herein are prodrugs that may allow earlier and/or more effective treatment at the time of diagnosis of SARS-CoV-2 infection. The prodrugs herein may represent an approach that may be able to target the antiviral to the lung and away from the liver where remdesivir's major dose limiting is directed.

Compounds, including antiviral prodrugs, are provided herein. The compounds have a structure according to formula (I):



wherein Nuc is selected from the group consisting of an antiviral nucleoside and an antiviral nucleoside analog; Y is independently selected from the group consisting of hydrogen, a C₁-C₃₀ hydrocarbyl, a pharmaceutically acceptable cation, and a covalent bond to a carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog; x is 0 or 1; L is independently a C₁-C₃₀ hydrocarbyl (such as a C₁-C₆ hydrocarbyl); and R is independently selected from the group consisting of a C₁₀-C₃₀ hydrocarbyl and a substituent of formula (A);




wherein R¹ and R² are independently selected from the group consisting of hydrogen and a C₁-C₃₀ hydrocarbyl.


Pharmaceutical formulations are provided. The pharmaceutical formulations include an oil and one or more compounds described herein. The pharmaceutical formulations may be formulated for injection, such as intramuscular injection or subcutaneous injection. The pharmaceutical formulations may be orally bioavailable.

Methods of treatment are provided, such as methods for treating a virus (e.g., coronavirus), including virus infections in mammals. The methods include administering an effective amount of a compound described herein, or a pharmaceutical formulation described herein.

Methods of producing a compound, such as a prodrug, are provided. The methods include (i) providing a compound of formula (a) –




(ii) providing a compound of formula (b) –




formula (b);

(iii) contacting the compound of formula (a) and the compound of formula (b) to form a compound of formula (c) -




formula (c);

5 and (iv) contacting the compound of formula (c) with an acid to form a compound of formula (d)-



formula (d);


wherein Het is a C₁-C₃₀ hydrocarbyl including at least one heteroatom; Y is selected from the group consisting of hydrogen, a C₁-C₃₀ hydrocarbyl, and a pharmaceutically acceptable cation; x is 0 or 1; L is a C₁-C₃₀ hydrocarbyl (such as a C₁-C₆ hydrocarbyl); and R is
 10 selected from the group consisting of a C₁₀-C₃₀ hydrocarbyl and a substituent of formula (A);



wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and a C_1 - C_{30} hydrocarbyl. The methods may include performing an intramolecular esterification reaction of a product, such as a phosphodiester, to form a cyclic phosphate, such as a 3',5'-cyclic phosphate.

Methods of producing a drug triphosphate also are provided. The methods include providing a plurality of cells, contacting the plurality of cells with an amount of a drug, incubating the plurality of cells and the amount of the drug for period effective to form the drug triphosphate.

In one aspect, the present disclosure provides a compound of the following formula or a pharmaceutically acceptable salt thereof:



Additional aspects will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the aspects described herein. The advantages described herein may be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive.

Brief Description of the Drawings

FIG. 1A—FIG. 1F depict concentration-response curves for ODBG-P-RVn, ODE-P-RVn, and HDP-P-RVn (**4b**), remdesivir (RDV), and remdesivir nucleoside (RVn) for SARS-CoV-2 infection in Vero E6 cells in two separate experiments performed in duplicate.

FIG. 2 depicts relative viabilities of embodiments of treated Vero E6 cells, as measured by CellTiter-Glo luminescent cell viability assay.

FIG. 3 depicts LC/MS/MS analysis of cells that were centrifuged at 1200 rpm for 10 minutes, diluent aspirated, and resuspended in 250 μ L of methanol/distilled water (70/30).

FIG. 4A and **FIG. 4B** depict plasma levels of ODBG-P-RVn (**FIG. 4A**) and RVn (**FIG. 4B**) in a seven day oral pharmacokinetic study in Syrian hamsters.

FIG. 5A and **FIG. 5B** depict the stability of ODE-P-RVn and ODBG-P-RVn in human plasma with either K₂EDTA (**FIG. 5A**) or sodium heparin (**FIG. 5B**) as anticoagulants.

FIG. 6A and **FIG. 6B** depict pharmacokinetics of ODE-Bn-TFV and ODE-TFV, respectively, after intramuscular administration of 100 mg/kg in formulation F3 to rats.

FIG. 7A and **FIG. 7B** are plots of the effects of 4 different formulations on the pharmacokinetics of ODE-Bn-TFV and ODE-TFV, respectively, after intramuscular administration of 100 mg/kg to rats.

FIG. 8 depicts a comparison of 96 mg/kg ODE-Bn-TFV pharmacokinetics in rats using formulation F3 versus F8.

FIG. 9 depicts rat plasma levels of ODE-Bn-TFV and ODE-TFV during a 3-month exposure to monthly intramuscular doses of ODE-Bn-TFV in formulation F3.

FIG. 10 depicts plasma levels of ODE-Bn-TFV and ODE-TFV in beagle dogs treated with 100 mg/kg with ODE-Bn-TFV in formulation F3.

FIG. 11A and **FIG. 11B** depict the TFVpp persistence in HFF cells of ODE-TFV (**FIG. 11A**) and ODE-Et-TFV (**FIG. 11B**).

FIG. 12A and **FIG. 12B** depict plasma concentrations of an embodiment of formulations containing ODE-Bn-TFV (**FIG. 12A**) or ODE-TFV (**FIG. 12B**) (ODE-Bn-TFV ED₉₀ 10.9 ng/mL; ODE-TFV ED₉₀ 6.3 nm/mL).

FIG. 12C and **FIG. 12D** depict plasma concentrations of an embodiment of formulations containing ODE-Bn-TFV (**FIG. 12C**) and ODE-TFV (**FIG. 12D**).

FIG. 12E and **FIG. 12F** depict plasma concentrations of an embodiment of formulations containing ODE-Bn-TFV (**FIG. 12E**) and ODE-TFV (**FIG. 12F**).

FIG. 12G and **FIG. 12H** depict plasma concentrations of an embodiment of formulations containing ODE-Bn-TFV (**FIG. 12G**) and ODE-TFV (**FIG. 12H**).

FIG. 12I depicts mean plasma concentrations of an embodiment of formulations.

FIG. 13 depicts the nanograms/mL of an embodiment of active ingredients present in rat plasma after administration of an embodiment of formulations herein.

FIG. 14 depicts amounts of an embodiment of a disphosphate in PBMCs after administration of an embodiment of a formulation herein.

Detailed Description

In one aspect, pharmaceutical formulations are provided herein. The pharmaceutical formulations may include (i) an oil, and (ii) a compound of formula (I), as described herein. In some embodiments, the pharmaceutical formulations also include
5 benzyl alcohol, benzyl benzoate, ethyl alcohol, or a combination thereof.

As used herein, the term "oil" refers to a non-polar compound or a mixture of two or more non-polar compounds that is insoluble in water (e.g., a solubility of less than 1 g per liter of water), such as a non-polar compound that includes an at least partially saturated carbon chain (e.g., hydrocarbons, fatty acids, etc.).

10 Any known oil may be used in the pharmaceutical formulations provided herein. In some embodiments, an oil is selected from the group consisting of sesame oil, triglycerides (e.g., medium chain triglycerides), and a combination thereof.

An oil may be present at any effective concentration in the pharmaceutical formulations provided herein. For example, an oil may be present in a pharmaceutical
15 formulation at an amount of about 60 % to about 95 %, about 65 % to about 95 %, about 70 % to about 95 %, about 75 % to about 95 %, about 80 % to about 95 %, or about 85 % to about 95 %, by weight, based on the weight of the pharmaceutical formulation.

In some embodiments, the pharmaceutical formulations also include benzyl alcohol, benzyl benzoate, ethyl alcohol, or a combination thereof. When present, benzyl
20 alcohol, ethyl alcohol, benzyl benzoate or the combination thereof may be present at a total amount of about 0.01 % to about 25 %, about 0.01 % to about 20 %, about 0.01 % to about 15 %, about 0.01 % to about 10 %, or about 5 % to about 10 %, by weight, based on the weight of the pharmaceutical formulation.

A compound of formula (I) may be present at any effective concentration in the
25 pharmaceutical formulations provided herein. For example, a compound of formula (I) may be present at an amount of about 0.01 % to about 40 %, about 0.01 % to about 30 %, about 0.01 % to about 20 %, about 0.01 % to about 10 %, about 0.01 % to about 5 %, by weight, based on the weight of the pharmaceutical formulation.

The pharmaceutical formulations provided herein may be configured for any route
30 of administration. In some embodiments, the pharmaceutical formulations are configured for oral administration. In some embodiments, the pharmaceutical formulations are configured for injection, such as intramuscular injection or subcutaneous injection.

The pharmaceutical formulations provided herein may be configured to be administered as a single dose. For example, a pharmaceutical formulation may be configured to be administered as a single, long-acting treatment, as described herein. The single dose may provide an effective amount of the pharmaceutical formulation, such as a compound of formula (I) that is present in the pharmaceutical formulation. The pharmaceutical formulations provided herein may be configured to be administered as a series of two or more doses, wherein each dose is administered at least 7 days, at least 14 days, at least 21 days, or at least 28 days after the preceding dose. For example, each of the two or more doses may be a long-acting treatment, as described herein.


In some embodiments, the pharmaceutical formulations include one or more of the components of any of formulations F3, F4, F7, or F8:

Components	F3	F4	F7	F8
Sesame oil	92.65	73.67	-	-
Medium chain triglycerides	-	-	92.65	73.67
Benzyl alcohol	1.22	10.67	1.22	10.67
Benzyl benzoate	-	15.67	-	15.67
Ethyl alcohol	6.12	-	6.12	-

In some embodiments, the pharmaceutical formulation is a long-acting treatment. As used herein, the phrase “long-acting treatment”, “long-acting dose”, and the like refers to a treatment or dose, respectively, that facilitates plasma antiviral activity above the 90% effective concentration for about 5 days to about 30 days, about 5 days to about 28 days, about 5 days to about 21 days, about 5 days to about 14 days, or about 5 days to about 7 days after the administration of a single dose.

The pharmaceutical formulations described herein may include one or more additives that do not undesirably affect one or more features of the pharmaceutical formulations, such as the activity of a compound of formula (I), bioavailability, etc. Non-limiting examples of additives include excipients, coloring agents, flavoring agents, buffers, preservatives, surfactants, other active ingredients, such as anti-inflammatory agents, pain reducing agents, etc.

In one aspect, compounds are provided herein, including compounds of formula (I):




formula (I).

The “Nuc” of formula (I) may be any suitable nucleoside. The nucleoside may be bonded to a compound in any manner. For example, a 5'-hydroxyl of a nucleoside may be joined to a phosphate moiety as an ester bond.

- 5 The nucleoside, in some embodiments, is an antiviral nucleoside. The antiviral nucleoside may be an antiviral ribonucleoside. The nucleoside, in some embodiments, is an antiviral nucleoside analog. The antiviral nucleoside analog may be an antiviral ribonucleoside analog.

- 10 In some embodiments, Nuc is RVn (GS-441524), beta-D-N⁴-hydroxycytidine (NHC), or (2'R)-2-amino-2'-deoxy-2'-fluoro-N,2'-dimethyladenosine (CAS # is 1998705-62-6). In some embodiments, Nuc is GS-441524, and the compound of formula (I) has the following structure:




- 15 Other antivirals for coronavirus infection can also be modified in the manner provided herein. For example, N⁴-hydroxy-cytidine (NHC) is an antiviral candidate entering clinical Phase I evaluation. Other nucleoside analogs known to inhibit RNA viruses are also suitable for modification according to this disclosure.

The “Y” of formula (I) may be any of the substituents described herein. In some embodiments, Y is hydrogen, a C₁-C₃₀ hydrocarbyl, a pharmaceutically acceptable cation,

or a covalent bond to a carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog.


When Y is a covalent bond to a carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog, the covalent bond may be a
 5 covalent bond to any carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog (e.g., the 1' carbon, the 2' carbon, the 3' carbon, or the 4' carbon). In other words, the covalent bond may be a covalent bond between (i) the oxygen to which Y is bonded in formula (I), and (ii) any carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog (e.g., the 1'
 10 carbon, the 2' carbon, the 3' carbon, or the 4' carbon). For example, Nuc may be GS-441524; the covalent bond may be between the oxygen to which Y is bonded in formula (I), and the 3' carbon of five-carbon sugar moiety of GS-441524, and the compound of formula (I) has the following structure:



15 When Y is a pharmaceutically acceptable cation, the pharmaceutically acceptable cation may be Na⁺.

In some embodiments, Y is a C₁-C₂₀ hydrocarbyl, a C₁-C₁₀ hydrocarbyl, or a C₁-C₆ hydrocarbyl. In some embodiments, Y is a C₁-C₆ alkyl, which may be unsubstituted. In some embodiments, Y includes at least one cyclic moiety. The at least one cyclic moiety
 20 may be a monocyclic moiety or a multicyclic moiety, e.g., a bicyclic moiety, a spiro moiety, etc. In some embodiments, Y is aryl, arylalkyl, heteroaryl, heteroarylalkyl, or heterocycloalkyl, each of which may be unsubstituted or substituted. In some embodiments, Y is an unsubstituted or substituted pyridinyl. In some embodiments, Y is


an unsubstituted or substituted benzyl. The unsubstituted or substituted benzyl may have a structure according to formula (B):




formula (B),

wherein R^3 , R^4 , R^5 , R^6 , and R^7 are independently selected from the group consisting of
 5 hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycl, aryl(alkyl), heteroaryl(alkyl), (heterocycl)alkyl, hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, azido, silyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy,
 10 trihalomethanesulfonyl, trihalomethanesulfonamido, amino, mono-substituted amino, and di-substituted amino. In some embodiments, R^3 , R^4 , R^5 , R^6 , and R^7 are hydrogen. In some embodiments, at least two of R^3 , R^4 , R^5 , R^6 , and R^7 are hydrogen. In some embodiments, at least three of R^3 , R^4 , R^5 , R^6 , and R^7 are hydrogen. In some embodiments, at least four of R^3 , R^4 , R^5 , R^6 , and R^7 are hydrogen.

15 When Y is an unsubstituted or substituted benzyl of formula (B), the compound of formula (I) has the following structure:




In formula (I), x may be 1 or 0. When x is 1, the “-O-L-” moiety is present in the compounds of formula (I). When x is 0, R is bonded directed to the oxygen of the phosphonate moiety, as shown in the following structure:



- 5 When “L” is present in the compounds of formula (I), the “L” may be selected from any of the substituents described herein. In some embodiments, L is a C₁-C₃₀ hydrocarbyl, a C₁-C₂₀ hydrocarbyl, a C₁-C₁₀ hydrocarbyl, a C₁-C₆ hydrocarbyl, a C₁-C₅ hydrocarbyl, a C₁-C₄ hydrocarbyl, a C₁-C₃ hydrocarbyl, or a C₁-C₂ hydrocarbyl. In some
10 embodiments, L is an ethyl, which may be unsubstituted. In some embodiments, L is a methyl, which may be unsubstituted. In some embodiments, L is a propyl, which may be unsubstituted.


- The “R” of formula (I) may be selected from any of the substituents described herein. In some embodiments, R is a C₁-C₃₀ hydrocarbyl, a C₅-C₃₀ hydrocarbyl, a C₁₀-C₃₀ hydrocarbyl, a C₁₂-C₂₄ hydrocarbyl, a C₁₃-C₂₉ hydrocarbyl, a C₁₅-C₂₄ hydrocarbyl, or a
15 C₂₀-C₂₄ hydrocarbyl. R, in some embodiments, is a heteroalkyl. R may include 0 to 6 unsaturated bonds, 1 to 6 unsaturated bonds, 2 to 6 unsaturated bonds, 3 to 6 unsaturated bonds, or 4 to 6 unsaturated bonds. The “unsaturated bonds” described herein may include any non-single bond, and when more than one unsaturated bond is present, the two
20 or more unsaturated bonds may be selected independently from a double bond or a triple bond. When one or more double bonds are present, the one or more double bonds may be *cis*-, *trans*-, or a combination thereof. R may include a cyclopropyl moiety, such as a terminal cyclopropyl moiety.

In some embodiments, R is -




- 25 wherein a is 1 to 29. In some embodiments, a is 15 to 25. In some embodiments, a is 18 to 22. In some embodiments, a is 19. In some embodiments, a is 6 to 10. In some embodiments, a is 8.

In some embodiments R is -




wherein b is 1 to 29, c is 0 to 28, and a sum of b and c is 29 or less. In some
embodiments, b is 1 to 4 and c is 15 to 20. In some embodiments, b is 3 and c is 15. In
5 some embodiments, b is 2 and c is 17.

In some embodiments, R is a substituent of formula (A);




wherein R¹ and R² are hydrogen or a C₁-C₃₀ hydrocarbyl, such as a C₁₀-C₃₀ hydrocarbyl,
or a C₁₂-C₂₄ hydrocarbyl. R¹, R², or both R¹ and R² may include at least one cyclic
10 moiety, which may be a monocyclic moiety or a multicyclic moiety, e.g., a bicyclic
moiety, a spiro moiety, etc. R¹, R², or both R¹ and R² may include 0 to 6 unsaturated
bonds, 1 to 6 unsaturated bonds, 2 to 6 unsaturated bonds, 3 to 6 unsaturated bonds, or 4 to
6 unsaturated bonds. When one or more double bonds are present, the one or more double
bonds may be *cis*-, *trans*-, or a combination thereof. R¹, R², or both R¹ and R² may
15 include a branched hydrocarbyl, such as a penultimate branched hydrocarbyl. In some
embodiments, at least one of R¹ and R² are hydrogen. In some embodiments, both R¹ and
R² are independently selected from a C₁-C₃₀ hydrocarbyl.

In some embodiments, R¹, R², or both R¹ and R² are independently selected from
the group consisting of aryl, arylalkyl, heteroaryl, heteroarylalkyl, and heterocycloalkyl,
20 each of which may be unsubstituted or substituted. The arylalkyl may be an unsubstituted
or substituted benzyl. The unsubstituted or substituted benzyl may have a structure
according to formula (C):




- wherein R^8 , R^9 , R^{10} , R^{11} , and R^{12} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl), (heterocyclyl)alkyl, hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, azido, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, amino, mono-substituted amino, and di-substituted amino. In some embodiments, each of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen. In some embodiments, at least two of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen.
- 10 In some embodiments, at least three of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen. In some embodiments, at least four of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen. In some embodiments, at least five of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen.

In some embodiments, R^1 is –




- 15 wherein d is 1 to 29. In some embodiments, d is 5 to 29, 10 to 29, 15 to 29, 20 to 29, 25 to 29, 1 to 25, 1 to 20, 1 to 15, 1 to 10, or 1 to 5.


In some embodiments, R^1 is



wherein e is 1 to 27, f is 0 to 26, and a sum of e and f is 27 or less.


- 20 In some embodiments, R^2 is selected from the group consisting of –






In some embodiments, g is 5 to 10. In some embodiments, g is 7.


The substituent of formula (A) may be a racemate, an sn-1 stereoisomer, or an sn-3 stereoisomer. Throughout this disclosure, when a formula, such as formula (A), is depicted with no indication(s) of spatial orientation, then the formula reads on all isomers, e.g., stereoisomers, of the compounds of the formula. For example, in some embodiments, a compound may have a structure according to formula (I), wherein x is 0, and R is a substituent of formula (A):





This formula lacks any indication of spatial orientation, and therefore reads on the sn-3 isomer thereof, the sn-1 isomer thereof, and mixtures of the sn-3 and sn-1 isomers, including racemic mixtures thereof:





Further non-limiting embodiments of compounds of formula (I) are provided at the following table:





Octadecyloxyethyl benzyl tenofovir (ODE-Bn-TFV)











Octadecyloxyethyl benzyl 9-(*R*)-[(2-phosphonomethoxy)propyl]-2,6-diaminopurine (ODE-Bn-PMPDAP)Oleyloxyethyl benzyl 9-(*R*)-[(2-phosphonomethoxy)propyl]-2,6-diaminopurine (OLE-Bn-PMPDAP)







Octadecyloxyethyl-*phospho*-emtricitabine (ODE-P-FTC)





Octadecyloxyethyl benzyl-phospho-emtricitabine (ODE-Bn-P-FTC)

Oleyloxyethyl benzyl 9-(*R*)-[2-phosphono-methoxy]propyl]-2,6-diaminopurine (OLE-Bn-PMPDAP)


Oleyloxyethyl 9-(*R*)-[(2-phosphonomethoxy)propyl]-2,6-diaminopurine (OLE-(*R*)-PMPDAP)

 Eicosyl-phospho-RVn	 Hexadecyloxypropyl-phospho-RVn
 Octadecyloxyethyl-benzyl-phospho-RVn	 1-O-tetradecyl-2-O-benzyl-<i>sn</i>-glyceryl-phospho-RVn
 1-O-hexadecyl-2-O-benzyl-<i>sn</i>-glyceryl-phospho-RVn	 1-O-hexadecyl-2-O-(3-fluoro, 4-methoxybenzyl)-<i>sn</i>-glyceryl-phospho-RVn
 1-O-octadecyl-2-O-benzyl-<i>rac</i>-glyceryl-phospho-RVn	 1-O-octadecyl-2-O-benzyl-<i>sn</i>-glyceryl-phospho-RVn
 1-O-octadecyl-2-O-octyl-<i>sn</i>-glyceryl-phospho-RVn	 1-O-octadecyl-2-O-(methylcyclohexyl)-<i>sn</i>-glyceryl-phospho-RVn






<i>phospho-RVn</i>  1-<i>O</i>-octadecyl-2-<i>O</i>-(3-fluorobenzyl)-<i>sn</i>-glyceryl-<i>phospho-RVn</i>	 1-<i>O</i>-octadecyl-2-<i>O</i>-(4-methoxybenzyl)-<i>sn</i>-glyceryl-<i>phospho-RVn</i>
 1-<i>O</i>-octadecyl-2-<i>O</i>-benzyl-benzyl-<i>sn</i>-glyceryl-<i>phospho-RVn</i>	
 1-<i>O</i>-octadecyl-2-<i>O</i>-(3-fluoro, 4-methoxybenzyl)-<i>sn</i>-glyceryl-<i>phospho-RVn</i>	
 1-<i>O</i>-octadecyl-2-<i>O</i>-(methylpyridinyl)-<i>sn</i>-glyceryl-<i>phospho-RVn</i>	
 1-<i>O</i>-octadecyl-2-<i>O</i>-(methylpyridinyl)-<i>sn</i>-glyceryl-<i>phospho-RVn</i>	

1-*O*-oleyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn1-*O*-oleyl-2-*O*-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn1-*O*-octadecyl-*sn*-glyceryl-*phospho*-RVnGS-441524-3',5'-cyclic monophosphate, 1-*O*-octadecyl-2-*O*-benzyl-*sn*-glyceryl ester

ODBG-benzyl-phospho-emtricitabine



ODBG-phospho-emtricitabine

1-*O*-octadecyl-2-*O*-(3-trifluoromethyl)-*sn*-glyceryl-*phospho*-RVn1-*O*-octadecyl-2-*O*-(4-trifluoromethyl)-*sn*-glyceryl-*phospho*-RVn1-*O*-octadecyl-2-*O*-(4-cyano)-*sn*-glyceryl-*phospho*-RVn1-*O*-octadecyl-2-*O*-(2-cyanobenzyl)-*sn*-glyceryl-*phospho*-RVn1-*O*-octadecyl-2-*O*-(3-cyanobenzyl)-*sn*-glyceryl-*phospho*-RVn

When used herein with regard to the selection of a substituent, the term “independently” indicates that (i) a substituent at a particular location may be the same or different for each molecule of a formula (e.g., (i) a compound of formula (I) may include two molecules of formula (I), with each molecule having the same or a different C₁-C₃₀ hydrocarbyl selected for R; and/or (ii) two differently labeled substituents selected from the same pool of substituents may be the same or different (e.g., R and Y of a molecule of a compound of formula (I) may both be selected from “a C₁-C₃₀ hydrocarbyl”, and the C₁-C₃₀ hydrocarbyls selected for R and Y may be the same or different)).

The phrases “C₁-C₃₀ hydrocarbyl,” “C₁₀-C₃₀ hydrocarbyl”, and the like, as used herein, generally refer to aliphatic, aryl, or arylalkyl groups containing 1 to 30 carbon atoms, or 10 to 30 carbon atoms, respectively, including unsubstituted groups and substituted derivatives thereof, which, as explained herein, may include, but are not limited to, heteroaryl, heteroarylalkyl, heterocycloalkyl groups, etc. Therefore, when a substituent herein is characterized, for example, as a “C₁-C₃₀ hydrocarbyl”, the phrase “C₁-C₃₀ hydrocarbyl” refers to and includes unsubstituted and substituted C₁-C₃₀ hydrocarbyls, unless expressly noted otherwise. Examples of aliphatic groups, in each instance, include, but are not limited to, an alkyl group, a cycloalkyl group, an alkenyl group, a cycloalkenyl group, an alkynyl group, an alkadienyl group, a cyclic group, and the like, and includes all substituted, unsubstituted, branched, and/or linear analogs or derivatives thereof, in each instance having 1 to 30 total carbon atoms or 10 to 30 total carbon atoms for a “C₁-C₃₀ hydrocarbyl ” and “C₁₀-C₃₀ hydrocarbyl”, respectively. Examples of alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl and dodecyl. Cycloalkyl moieties may be monocyclic or multicyclic, and examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and adamantyl, including any heteroatom substituted derivative thereof. Additional examples of alkyl moieties have linear, branched and/or cyclic portions (e.g., 1-ethyl-4-methyl-cyclohexyl). Representative alkenyl moieties include vinyl, allyl, 1-butenyl, 2-butenyl, isobutenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 2-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl and 3-decenyl. Representative alkynyl moieties include acetylenyl, propynyl, 1-butyne, 2-butyne, 1-

pentynyl, 2-pentynyl, 3-methyl-1-butynyl, 4-pentynyl, 1-hexynyl, 2-hexynyl, 5-hexynyl, 1-heptynyl, 2-heptynyl, 6-heptynyl, 1-octynyl, 2-octynyl, 7-octynyl, 1-nonynyl, 2-nonynyl, 8-nonynyl, 1-decynyl, 2-decynyl and 9-decynyl. Examples of aryl or arylalkyl moieties include, but are not limited to, anthracenyl, azulenyl, biphenyl, fluorenyl, indan, indenyl, naphthyl, phenanthrenyl, phenyl, 1,2,3,4-tetrahydro-naphthalene, anthracenyl, tolyl, xylyl, mesityl, benzyl, and the like, including any heteroatom substituted derivative thereof.

Unless otherwise indicated, the term "substituted," when used to describe a chemical structure or moiety, refers to a derivative of that structure or moiety wherein (i) a multi-valent non-carbon atom (e.g., oxygen, nitrogen, sulfur, phosphorus, etc.) is bonded to one or more carbon atoms of the chemical structure or moiety (e.g., a "substituted" C₄ hydrocarbyl may include, but is not limited to, a pyrimidinyl moiety, a pyridinyl moiety, a dioxanyl moiety, a diethyl ether moiety, a methyl propionate moiety, an *N,N*-dimethylacetamide moiety, a butoxy moiety, *etc.*, and a "substituted" aryl C₁₂ hydrocarbyl may include, but is not limited to, an oxydibenzene moiety, a benzophenone moiety, *etc.*) or (ii) one or more of its hydrogen atoms (e.g., chlorobenzene may be characterized generally as an aryl C₆ hydrocarbyl "substituted" with a chlorine atom) is substituted with a chemical moiety or functional group such as acyl, alcohol, alkoxy, alkanoyloxy, alkoxycarbonyl, alkenyl, alkyl (e.g., methyl, ethyl, propyl, *t*-butyl), alkynyl, alkylcarbonyloxy (-OC(O)alkyl), amide (-C(O)NH-alkyl- or -alkylNHC(O)alkyl), primary, secondary, and tertiary amino (such as alkylamino, arylamino, arylalkylamino), aryl, arylalkyl, aryloxy, azo, azido, carbamoyl (-NHC(O)O-alkyl- or -OC(O)NH-alkyl), carbamyl (*e.g.*, CONH₂, as well as CONH-alkyl, CONH-aryl, and CONH-arylalkyl), carboxyl, carboxylic acid, cyano, cycloalkyl, cycloalkenyl, ester, ether (*e.g.*, methoxy, ethoxy), halo, haloalkyl (*e.g.*, -CCl₃, -CF₃, -C(CF₃)₃), haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, heteroalkyl, heterocycloalkyl, heteroaryl, heteroarylalkyl, isocyanate, isothiocyanate, nitrile, nitro, oxo, phosphodiester, silyl, sulfide, sulfonamido (*e.g.*, SO₂NH₂), sulfone, sulfonyl, sulfinyl, sulfonyl (including alkylsulfonyl, arylsulfonyl and arylalkylsulfonyl), sulfoxide, thiocarbonyl, thiocarbamyl, thiocyanato, thiol (*e.g.*, sulphydryl, thioether) or urea (-NHCONH-alkyl-).

Methods of Treatment

Also provided herein are methods of treatment, including methods of treating a virus infection, such as a coronavirus infection. The virus infection may be an infection in a mammal.

- 5 In some embodiments, the methods include administering to a mammal an effective amount of a compound described herein or a pharmaceutical formulation described herein.

The virus infection may be an RNA virus infection. In some embodiments, the RNA virus infection is caused by a RNA virus of a viral family selected from the group
10 consisting of *Filoviridae*, *Orthomyxoviridae*, *Paramyxoviridae*, *Pneumoviridae*, *Phenuiviridae*, *Nairoviridae*, *Arenaviridae*, *Flaviviridae*, and *Coronaviridae*.

- Compounds, including prodrugs, provided herein may be screened for inhibitory activity against SARS-CoV-2 and related coronaviruses (or other viruses), using conventional techniques for evaluating anti-coronavirus activity and cytotoxicity.
15 Typically, compounds are first screened for inhibition of coronavirus *in vitro*, and those showing significant antiviral activity are then screened for efficacy *in vivo*.


- Non-limiting examples of potentially useful *in vitro* assays including the following: a) Using the OC43 beta-coronavirus strain (ATCC 1558) in the human adenocarcinoma cell line, HCT-8 (ATCC CCL-244), or using coronavirus 229E in MRC-5
20 human lung fibroblasts. Endpoints can include semiquantitative RT-PCR and pfu as determined by triplicate serial dilution. b) The activity of compounds may be studied using laboratory and clinical isolates of SARS-CoV-2 in Vero E6 cells, Caco-2, Calu-3, HPSC human lung cells, or Huh7.5 cells. Initial SARS CoV-2 growth inhibition assays can quantify plaque reduction on Vero cells grown in 12 well plates using a commercial
25 murine anti-SARS CoV-2 spike protein detection antibody (Item 40021-MM07, SinoBiological.com). Virus can also be quantified in culture supernatants by serial dilution on Vero cell lawns and by RT-PCR. Laboratory strains that can be obtained, for example, from BEI Resources (Strains NR52281 and NR52282), and clinical strains that can be isolated from patients participating in clinical trials can be used. Cytotoxicity can
30 be measured by commercially MTT or Cell Titer Glo. Compounds with the lowest 90 % inhibitory concentrations and that require the highest concentrations to induce cellular cytotoxicity may be selected for further evaluation. Anti-coronavirus compounds may

also be evaluated in a lung explant model for SARS CoV infection. To determine activity in primary cells from the organ most clinically impacted by the virus, candidate molecules with the highest therapeutic indices in Vero E6 cells can be advanced to studies in human lung explants.

5 *Methods of Producing a Compound*


Also provided herein are methods of producing a compound, such as a compound described herein, which may be a prodrug.

The compounds provided herein may be prepared by a variety of processes, including the processes described herein. In some embodiments, protected analogs of
 10 remdesivir nucleoside, RVn, 2 are prepared and then coupled to suitable alkoxyalkyl phosphates to form phosphodiesters. Removal of the protecting groups can afford compounds of Formula (I).




15 In some embodiments, 2-C-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2,5-anhydro-D-altronitrile (RVn, 2) is first converted to its 2', 3'-isopropylidene derivative. Mixtures of alkoxyalkyl phosphates and protected RVn may then be treated with *N,N*-dicyclohexylcarbodiimide (DCC) and *N,N*-dimethylaminopyridine (DMAP) under
 20 conditions suitable to prepare the phosphodiesters. Removal of the isopropylidene protecting group by treatment with dilute HCl or other suitable acid may provide compounds of Formula (I) in suitable yield and purity.

In some embodiments, the methods include providing a compound of formula (a) –



wherein x, R, L, and Y are as defined herein.


In some embodiments, the methods include providing a compound of formula (b) –




formula (b),

wherein Het is as defined herein. In some embodiments, Het is selected from the group

5 consisting of –




Also, as explained herein, formula (b) does not include any stereochemical indication(s), and therefore, reads on at least the following stereoisomer of formula (b):



10


In some embodiments, the methods include contacting a compound of formula (a) and a compound of formula (b) to form a compound of formula (c) -



formula (c).

The contacting of a compound of formula (a) and a compound of formula (b) may occur at any temperature or pressure, and may occur in the presence of any suitable liquid. The liquid may include a C₁-C₃₀ hydrocarbyl, such as a C₁-C₃₀ hydrocarbyl that includes at least one cyclic moiety, at least one heteroatom, such as nitrogen, or a combination thereof. In some embodiments, the liquid is *N,N*-dicyclohexylcarbodiimide, 4-dimethylaminopyridine, or a combination thereof.

In some embodiments, the methods contacting a compound of formula (c) with an acid to form a compound of formula (d) -



formula (d).

The acid may include any acid that is capable of facilitating the formation of a compound of formula (d). The acid may be an organic acid or inorganic acid. The acid may include a hydrogen halide, such as hydrogen chloride. The contacting of a compound of formula (c) with an acid may occur in the presence of any suitable liquid. The liquid may be a C₁-C₃₀ hydrocarbyl, such as a C₁-C₃₀ hydrocarbyl including at least one cyclic moiety, at least one heteroatom, or a combination thereof. In some embodiments, the liquid is tetrahydrofuran.

In some embodiments, the methods include performing an intramolecular esterification reaction of a compound of formula (d) to form a cyclic phosphate, such as a 3',5'-cyclic phosphate.

Methods of Producing a Drug Triphosphate

Also provided herein are methods of producing a drug triphosphate. In some embodiments, the methods include providing a plurality of cells, contacting the plurality of cells with an amount of a drug, incubating the plurality of cells and the amount of the drug for period effective to form the drug triphosphate. The plurality of cells may include any suitable cells. The plurality of cells, in some embodiments, includes Vero E6 cells, Calu-2 cells, Caco-2 cells, MRC5 human lung fibroblasts, Huh7.5 cells and PSC human lung cells. In some embodiments, the drug includes remdesivir or the remdesivir nucleoside (GS441524).

All referenced publications are incorporated herein by reference in their entirety. Furthermore, where a definition or use of a term in a reference, which is incorporated by reference herein, is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply.

Unless defined otherwise, all technical and scientific terms and any acronyms used herein have the same meanings as commonly understood by one of ordinary skill in the art in the field of the invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the exemplary methods, devices, and materials are described herein.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, Molecular Cloning: A Laboratory Manual, 2nd ed. (Sambrook et al., 1989); Oligonucleotide Synthesis (M. J. Gait, ed., 1984); Animal Cell Culture (R. I. Freshney, ed., 1987); Methods in Enzymology (Academic Press, Inc.); Current Protocols in Molecular Biology (F. M. Ausubel et al., eds., 1987, and periodic updates); PCR: The Polymerase Chain Reaction (Mullis et al., eds., 1994); Remington, The Science and Practice of Pharmacy, 20th ed., (Lippincott, Williams & Wilkins 2003), and Remington, The Science and Practice of Pharmacy, 22th ed., (Pharmaceutical Press and Philadelphia College of Pharmacy at University of the Sciences 2012).

While certain aspects of conventional technologies have been discussed to facilitate disclosure of various embodiments, applicants in no way disclaim these technical aspects, and it is contemplated that the present disclosure may encompass one or more of the conventional technical aspects discussed herein.

5 The present disclosure may address one or more of the problems and deficiencies of known methods and processes. However, it is contemplated that various embodiments may prove useful in addressing other problems and deficiencies in a number of technical areas. Therefore, the present disclosure should not necessarily be construed as limited to addressing any of the particular problems or deficiencies discussed herein.

10 In this specification, where a document, act or item of knowledge is referred to or discussed, this reference or discussion is not an admission that the document, act or item of knowledge or any combination thereof was at the priority date, publicly available, known to the public, part of common general knowledge, or otherwise constitutes prior art under the applicable statutory provisions; or is known to be relevant to an attempt to solve
15 any problem with which this specification is concerned.

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains,” “containing,” “characterized by,” or any other variation thereof, are intended to encompass a non-exclusive inclusion, subject to any limitation explicitly indicated otherwise, of the recited components. For example, a fusion protein, a
20 pharmaceutical composition, and/or a method that “comprises” a list of elements (e.g., components, features, or steps) is not necessarily limited to only those elements (or components or steps), but may include other elements (or components or steps) not expressly listed or inherent to the fusion protein, pharmaceutical composition and/or method.

25 As used herein, the transitional phrases “consists of” and “consisting of” exclude any element, step, or component not specified. For example, “consists of” or “consisting of” used in a claim would limit the claim to the components, materials or steps specifically recited in the claim except for impurities ordinarily associated therewith (i.e., impurities within a given component). When the phrase “consists of” or “consisting of” appears in a
30 clause of the body of a claim, rather than immediately following the preamble, the phrase “consists of” or “consisting of” limits only the elements (or components or steps) set forth in that clause; other elements (or components) are not excluded from the claim as a whole.

As used herein, the transitional phrases “consists essentially of” and “consisting essentially of” are used to define a compound, pharmaceutical formulations, and/or method that includes materials, steps, features, components, or elements, in addition to those literally disclosed, provided that these additional materials, steps, features,
5 components, or elements do not materially affect the basic and novel characteristic(s) of the claimed invention. The term “consisting essentially of” occupies a middle ground between “comprising” and “consisting of”.

It is understood that aspects and embodiments of the invention described herein include “consisting” and/or “consisting essentially of” aspects and embodiments.

10 The terms “a,” “an,” and “the” are intended to include plural alternatives, e.g., at least one. For instance, the disclosure of “a compound”, “a pharmaceutical formulation”, “an acid”, and the like, is meant to encompass one, or mixtures or combinations of more than one compound, pharmaceutical formulation, acid, and the like, unless otherwise specified.

15 The term “and/or” when used in a list of two or more items, means that any one of the listed items can be employed by itself or in combination with any one or more of the listed items. For example, the expression “A and/or B” is intended to mean either or both of A and B, i.e. A alone, B alone or A and B in combination. The expression “A, B and/or C” is intended to mean A alone, B alone, C alone, A and B in combination, A and C in
20 combination, B and C in combination or A, B, and C in combination.

Various numerical ranges may be disclosed herein. When Applicant discloses or claims a range of any type, Applicant’s intent is to disclose or claim individually each possible number that such a range could reasonably encompass, including end points of the range as well as any sub-ranges and combinations of sub-ranges encompassed therein,
25 unless otherwise specified. Moreover, all numerical end points of ranges disclosed herein are approximate. As a representative example, Applicant discloses, in some embodiments, that “a is 15 to 25”. This range should be interpreted as encompassing 15 and 25, and further encompasses each of 16, 17, 18, 19, 20, 21, 22, 23, and 24, including any ranges and sub-ranges between any of these values.

30 When such values or ranges are expressed, other embodiments disclosed include the specific value recited, from the one particular value, and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent

“about,” it will be understood that the particular value forms another embodiment. It will be further understood that there are a number of values disclosed therein, and that each value is also herein disclosed as “about” that particular value in addition to the value itself. In embodiments, “about” can be used to mean, for example, within 10% of the recited value, within 5% of the recited value, or within 2% of the recited value.

As used herein the term “pharmaceutical composition” refers to pharmaceutically acceptable compositions, wherein the composition comprises a pharmaceutically active agent, and in some embodiments further comprises a pharmaceutically acceptable carrier. In some embodiments, the pharmaceutical composition may be a combination of pharmaceutically active agents and carriers.

The term “combination” refers to either a fixed combination in one dosage unit form, or a kit of parts for the combined administration where one or more active compounds and a combination partner (e.g., another drug as explained below, also referred to as “therapeutic agent” or “co-agent”) may be administered independently at the same time or separately within time intervals. In some circumstances, the combination partners show a cooperative, e.g., synergistic effect. The terms “co-administration” or “combined administration” or the like as utilized herein are meant to encompass administration of the selected combination partner to a single subject in need thereof (e.g., a patient), and are intended to include treatment regimens in which the agents are not necessarily administered by the same route of administration or at the same time. The term “pharmaceutical combination” as used herein means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term “fixed combination” means that the active ingredients, e.g., a compound and a combination partner, are both administered to a patient simultaneously in the form of a single entity or dosage. The term “non-fixed combination” means that the active ingredients, e.g., a compound and a combination partner, are both administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific time limits, wherein such administration provides therapeutically effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g., the administration of three or more active ingredients.

As used herein the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S.

Pharmacopoeia, other generally recognized pharmacopoeia in addition to other formulations that are safe for use in animals, and more particularly in humans and/or non-
5 human mammals.

As used herein the term “pharmaceutically acceptable carrier” refers to an excipient, diluent, preservative, solubilizer, emulsifier, adjuvant, and/or vehicle with which demethylation compound(s), is administered. Such carriers may be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin,
10 such as peanut oil, soybean oil, mineral oil, sesame oil and the like, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents. Antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; and agents for the adjustment of tonicity such as sodium chloride or dextrose may also be a carrier. Methods for producing
15 compositions in combination with carriers are known to those of skill in the art. In some embodiments, the language “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. See, e.g.,
20 Remington, The Science and Practice of Pharmacy, 20th ed., (Lippincott, Williams & Wilkins 2003). Except insofar as any conventional media or agent is incompatible with the active compound, such use in the compositions is contemplated.

As used herein, “therapeutically effective amount” refers to an amount of a pharmaceutically active compound(s) that is sufficient to treat or ameliorate, or in some
25 manner reduce the symptoms associated with diseases and medical conditions. When used with reference to a method, the method is sufficiently effective to treat or ameliorate, or in some manner reduce the symptoms associated with diseases or conditions. For example, an effective amount in reference to diseases is that amount which is sufficient to block or prevent onset; or if disease pathology has begun, to palliate, ameliorate, stabilize, reverse
30 or slow progression of the disease, or otherwise reduce pathological consequences of the disease. In any case, an effective amount may be given in single or divided doses.

As used herein, the terms "treat," "treatment," or "treating" embraces at least an amelioration of the symptoms associated with diseases in the patient, where amelioration is used in a broad sense to refer to at least a reduction in the magnitude of a parameter, e.g. a symptom associated with the disease or condition being treated. As such, "treatment" also includes situations where the disease, disorder, or pathological condition, or at least symptoms associated therewith, are completely inhibited (e.g. prevented from happening) or stopped (e.g. terminated) such that the patient no longer suffers from the condition, or at least the symptoms that characterize the condition.

As used herein, and unless otherwise specified, the terms "prevent," "preventing" and "prevention" refer to the prevention of the onset, recurrence or spread of a disease or disorder, or of one or more symptoms thereof. In certain embodiments, the terms refer to the treatment with or administration of a compound or dosage form provided herein, with or without one or more other additional active agent(s), prior to the onset of symptoms, particularly to subjects at risk of disease or disorders provided herein. The terms encompass the inhibition or reduction of a symptom of the particular disease. In certain embodiments, subjects with familial history of a disease are potential candidates for preventive regimens. In certain embodiments, subjects who have a history of recurring symptoms are also potential candidates for prevention. In this regard, the term "prevention" may be interchangeably used with the term "prophylactic treatment."

As used herein, and unless otherwise specified, a "prophylactically effective amount" of a compound is an amount sufficient to prevent a disease or disorder, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of therapeutic agent, alone or in combination with one or more other agent(s), which provides a prophylactic benefit in the prevention of the disease. The term "prophylactically effective amount" can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent. As used herein, and unless otherwise specified, the term "subject" is defined herein to include animals such as mammals, including, but not limited to, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, and the like. In specific embodiments, the subject is a human. The terms "subject" and "patient" are used interchangeably herein in reference, for example, to a mammalian subject, such as a human.

As used herein, and unless otherwise specified, a compound described herein is intended to encompass all possible stereoisomers, unless a particular stereochemistry is specified. Where structural isomers of a compound are interconvertible via a low energy barrier, the compound may exist as a single tautomer or a mixture of tautomers. This can
5 take the form of proton tautomerism; or so-called valence tautomerism in the compound, e.g., that contain an aromatic moiety.

“Nucleic acid” or “nucleic acid molecule” refers to a multimeric compound comprising two or more covalently bonded nucleosides or nucleoside analogs having nitrogenous heterocyclic bases, or base analogs, where the nucleosides are linked together
10 by phosphodiester bonds or other linkages to form a polynucleotide. Nucleic acids include RNA, DNA, or chimeric DNA-RNA polymers or oligonucleotides, and analogs thereof. A nucleic acid backbone can be made up of a variety of linkages, including one or more of sugar-phosphodiester linkages, peptide-nucleic acid bonds, phosphorothioate linkages, methylphosphonate linkages, or combinations thereof. Sugar moieties of the nucleic acid
15 can be ribose, deoxyribose, or similar compounds having known substitutions (e.g. 2'-methoxy substitutions and 2'-halide substitutions). Nitrogenous bases can be conventional bases (A, G, C, T, U) or analogs thereof (e.g., inosine, 5-methylisocytosine, isoguanine). A nucleic acid can comprise only conventional sugars, bases, and linkages as found in RNA and DNA, or can include conventional components and substitutions (e.g.,
20 conventional bases linked by a 2'-methoxy backbone, or a nucleic acid including a mixture of conventional bases and one or more base analogs). Nucleic acids can include “locked nucleic acids” (LNA), in which one or more nucleotide monomers have a bicyclic furanose unit locked in an RNA mimicking sugar conformation, which enhances hybridization affinity toward complementary sequences in single-stranded RNA (ssRNA),
25 single-stranded DNA (ssDNA), or double-stranded DNA (dsDNA). Nucleic acids can include modified bases to alter the function or behavior of the nucleic acid (e.g., addition of a 3'-terminal dideoxynucleotide to block additional nucleotides from being added to the nucleic acid). Synthetic methods for making nucleic acids in vitro are well known in the art although nucleic acids can be purified from natural sources using routine techniques.
30 Nucleic acids can be single-stranded or double-stranded.

A nucleic acid is typically single-stranded or double-stranded and will generally contain phosphodiester bonds, although in some cases, as outlined, herein, nucleic acid

analogues are included that may have alternate backbones, including, for example and without limitation, phosphoramidate (Beaucage et al. (1993) *Tetrahedron* 49(10):1925 and references therein; Letsinger (1970) *J. Org. Chem.* 35:3800; Sprinzl et al. (1977) *Eur. J. Biochem.* 81:579; Letsinger et al. (1986) *Nucl. Acids Res.* 14: 3487; Sawai et al. (1984) *Chem. Lett.* 805; Letsinger et al. (1988) *J. Am. Chem. Soc.* 110:4470; and Pauwels et al. (1986) *Chemica Scripta* 26: 1419, which are each incorporated by reference), phosphorothioate (Mag et al. (1991) *Nucleic Acids Res.* 19:1437; and U.S. Pat. No. 5,644,048, which are both incorporated by reference), phosphorodithioate (Briu et al. (1989) *J. Am. Chem. Soc.* 111:2321, which is incorporated by reference), O-methylphosphoroamidite linkages (see Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press (1992), which is incorporated by reference), and peptide nucleic acid backbones and linkages (see, Egholm (1992) *J. Am. Chem. Soc.* 114:1895; Meier et al. (1992) *Chem. Int. Ed. Engl.* 31:1008; Nielsen (1993) *Nature* 365:566; and Carlsson et al. (1996) *Nature* 380:207, which are each incorporated by reference). Other analog nucleic acids include those with positively charged backbones (Denpcy et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6097, which is incorporated by reference); non-ionic backbones (U.S. Pat. Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; *Angew (1991) Chem. Intl. Ed. English* 30: 423; Letsinger et al. (1988) *J. Am. Chem. Soc.* 110:4470; Letsinger et al. (1994) *Nucleoside & Nucleotide* 13:1597; Chapters 2 and 3, *ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research"*, Ed. Y. S. Sanghvi and P. Dan Cook; Mesmaeker et al. (1994) *Bioorganic & Medicinal Chem: Lett.* 4: 395; Jeffs et al. (1994) *J. Biomolecular NMR* 34:17; and *Tetrahedron Lett.* 37:743 (1996), which are each incorporated by reference) and non-ribose backbones, including those described in U.S. Pat. Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, *ASC Symposium Series 580, Carbohydrate Modifications in Antisense Research*, Ed. Y. S. Sanghvi and P. Dan Cook, which references are each incorporated by reference. Nucleic acids containing one or more carbocyclic sugars are also included within the definition of nucleic acids (see Jenkins et al. (1995) *Chem. Soc. Rev.* pp 169-176, which is incorporated by reference). Several nucleic acid analogues are also described in, e.g., Rawls, *C & E News* Jun. 2, 1997 page 35, which is incorporated by reference. These modifications of the ribose-phosphate backbone

may be done to facilitate the addition of additional moieties such as labels, or to alter the stability and half-life of such molecules in physiological environments.

In addition to these naturally occurring heterocyclic bases that are typically found in nucleic acids (e.g., adenine, guanine, thymine, cytosine, and uracil), nucleic acid
5 analogs also include those having non-naturally occurring heterocyclic or modified bases, many of which are described, or otherwise referred to, herein. In particular, many non-naturally occurring bases are described further in, e.g., Seela et al. (1991) *Helv. Chim. Acta* 74:1790, Grein et al. (1994) *Bioorg. Med. Chem. Lett.* 4:971-976, and Seela et al. (1999) *Helv. Chim. Acta* 82:1640, which are each incorporated by reference. To further
10 illustrate, certain bases used in nucleotides that act as melting temperature (TO) modifiers are optionally included. For example, some of these include 7-deazapurines (e.g., 7-deazaguanine, 7-deazaadenine, etc.), pyrazolo[3,4-d]pyrimidines, propynyl-dN (e.g., propynyl-dU, propynyl-dC, etc.), and the like. See, e.g., U.S. Pat. No. 5,990,303, entitled "SYNTHESIS OF 7-DEAZA-2'-DEOXYGUANOSINE NUCLEOTIDES," which issued
15 Nov. 23, 1999 to Seela, which is incorporated by reference. Other representative heterocyclic bases include, e.g., hypoxanthine, inosine, xanthine; 8-aza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 7-deaza-8-aza derivatives of adenine, guanine, 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 6-
20 azacytosine; 5-fluorocytosine; 5-chlorocytosine; 5-iodocytosine; 5-bromocytosine; 5-methylcytosine; 5-propynylcytosine; 5-bromovinyluracil; 5-fluorouracil; 5-chlorouracil; 5-iodouracil; 5-bromouracil; 5-trifluoromethyluracil; 5-methoxymethyluracil; 5-ethynyluracil; 5-propynyluracil, and the like.

Examples of modified bases and nucleotides are also described in, e.g., U.S. Pat.
25 No. 5,484,908, entitled "OLIGONUCLEOTIDES CONTAINING 5-PROPYNYL PYRIMIDINES," issued Jan. 16, 1996 to Froehler et al., U.S. Pat. No. 5,645,985, entitled "ENHANCED TRIPLE-HELIX AND DOUBLE-HELIX FORMATION WITH OLIGOMERS CONTAINING MODIFIED PYRIMIDINES," issued Jul. 8, 1997 to Froehler et al., U.S. Pat. No. 5,830,653, entitled "METHODS OF USING OLIGOMERS
30 CONTAINING MODIFIED PYRIMIDINES," issued Nov. 3, 1998 to Froehler et al., U.S. Pat. No. 6,639,059, entitled "SYNTHESIS OF [2.2.1]BICYCLO NUCLEOSIDES," issued Oct. 28, 2003 to Kochkine et al., U.S. Pat. No. 6,303,315, entitled "ONE STEP

SAMPLE PREPARATION AND DETECTION OF NUCLEIC ACIDS IN COMPLEX BIOLOGICAL SAMPLES,” issued Oct. 16, 2001 to Skouv, and U.S. Pat. Application Pub. No. 2003/0092905, entitled “SYNTHESIS OF [2.2.1]BICYCLO NUCLEOSIDES,” by Kochkine et al. that published May 15, 2003, which are each incorporated by reference.

5 An “oligonucleotide” or “oligomer” refers to a nucleic acid that includes at least two nucleic acid monomer units (e.g., nucleotides), typically more than three monomer units, and more typically greater than ten monomer units. The exact size of an oligonucleotide generally depends on various factors, including the ultimate function or use of the oligonucleotide. Oligonucleotides are optionally prepared by any suitable
10 method, including, but not limited to, isolation of an existing or natural sequence, DNA replication or amplification, reverse transcription, cloning and restriction digestion of appropriate sequences, or direct chemical synthesis by a method such as the phosphotriester method of Narang et al. (1979) Meth. Enzymol. 68:90-99; the phosphodiester method of Brown et al. (1979) Meth. Enzymol. 68:109-151; the
15 diethylphosphoramidite method of Beaucage et al. (1981) Tetrahedron Lett. 22:1859-1862; the triester method of Matteucci et al. (1981) J. Am. Chem. Soc. 103:3185-3191; automated synthesis methods; or the solid support method of U.S. Pat. No. 4,458,066, or other methods known in the art. All of these references are incorporated by reference.

Compounds of the present disclosure and methods of use for inhibiting RNA
20 viruses include the following viral families: Filoviridae, Orthomyxoviridae, Paramyxoviridae, Pneumoviridae, Phenuiviridae, Nairoviridae, Arenaviridae, Flaviviridae and Coronaviridae. The names of exemplary viruses in each family are included in the below table.

RNA Virus Families which may be inhibited by compounds of the disclosure


Virus Family	Virus
<i>Filoviridae</i>	Ebola virus
	Sudan virus
	Bundibugyo virus
	Bombali virus
	Reston virus
	Marburg virus
	Ravn virus
<i>Orthomyxoviridae</i>	Influenza viruses
<i>Paramyxoviridae</i>	Nipah virus
	Hendra virus
	Human Parainfluenza viruses

Virus Family	Virus
	Measles virus
	Mumps virus
	Sosuga virus
<i>Pneumoviridae</i>	Respiratory syncytial viruses
	Human metapneumovirus
<i>Phenuiviridae</i>	Rift Valley Fever virus
	Punta Toro phlebovirus
<i>Nairoviridae</i>	Crimean Congo Hemorrhagic Fever virus
	Dugbe virus
<i>Arenaviridae</i>	Lassa virus
	Junin virus
	Lymphocytic choriomeningitis virus
	Guanarito virus
	Machupo virus
<i>Flaviviridae</i>	Kyasanur Forest Disease virus
	Omsk Hemorrhagic Fever virus
	Yellow Fever virus
	Japanese Encephalitis virus
	Hepatitis C Virus
	Zika Virus
	Dengue Viruses
	West Nile Virus
	Tick Borne encephalitis virus
	Murray Valley Fever encephalitis Virus
	Kunjin Virus
	Saint Louis Encephalitis Virus
	Bovine viral diarrhea virus
<i>Coronaviridae</i>	SARS-CoV-2
	MERS
	SARS CoV
	OC43
	229E
	NL43
	Evolving Zoonotic and Human Coronaviruses
	Feline Infectious Peritonitis virus

EMBODIMENTS

Embodiments of the compounds, pharmaceutical formulations, and methods described herein are provided at the following listing:


- 5 *Embodiment 1.* A compound of formula (I) or a pharmaceutically acceptable salt thereof:



formula (I);

wherein Nuc is selected from the group consisting of an antiviral nucleoside and an antiviral nucleoside analog; Y is independently selected from the group consisting of

- 5 hydrogen, a C₁-C₃₀ hydrocarbyl, a pharmaceutically acceptable cation, and a covalent bond to a carbon atom of a five-carbon sugar moiety of the antiviral nucleoside or the antiviral nucleoside analog; x is 0 or 1; L is independently a C₁-C₃₀ hydrocarbyl; and R is independently selected from the group consisting of a C₁₀-C₃₀ hydrocarbyl and a substituent of formula (A);



formula (A),

- 10 wherein R¹ and R² are independently selected from the group consisting of hydrogen and a C₁-C₃₀ hydrocarbyl;

wherein optionally the compound of formula (I) (i) achieves kinase bypass of the first nucleoside phosphorylation, (ii) provides increased oral bioavailability, (iii) delivers antivirally significant concentrations to lung and gastrointestinal tract and formulations,




15 (iv) provides sustained levels in plasma for at least about 5 to about 30 days following administration of a single dose, such as intravascularly, or (v) a combination thereof.


Embodiment 2. The compound of Embodiment 1, wherein the antiviral nucleoside or the antiviral nucleoside analog is an antiviral ribonucleoside or an antiviral ribonucleoside analog, respectively.

- 20 *Embodiment 3.* The compound of any one of the previous Embodiments, wherein Nuc is selected from the group consisting of GS-441524, beta-D-N⁴-hydroxycytidine (NHC), and (2'R)-2-amino-2'-deoxy-2'-fluoro-N,2'-dimethyladenosine.

Embodiment 4. The compound of any one of the previous Embodiments, wherein Nuc is -

- 25 (i) GS-441524:





Embodiment 5. The compound of any one of the previous Embodiments, wherein Y is an unsubstituted C₁-C₆ alkyl, a C₁-C₂₀ hydrocarbyl, a C₁-C₁₀ hydrocarbyl, a C₁-C₆ hydrocarbyl, or Na⁺.


5 *Embodiment 6.* The compound of any one of the previous Embodiments, wherein Y comprises at least one cyclic moiety.

Embodiment 7. The compound of any one of the previous Embodiments, wherein Y is selected from the group consisting of aryl, arylalkyl, heteroaryl, heteroarylalkyl, and heterocycloalkyl, each of which is unsubstituted or substituted.

10 *Embodiment 8.* The compound of any one of the previous Embodiments, wherein the heteroaryl is an unsubstituted or substituted pyridinyl.

Embodiment 9. The compound of any one of the previous Embodiments, wherein the arylalkyl is an unsubstituted or substituted benzyl.

15 *Embodiment 10.* The compound of any one of the previous Embodiments, wherein the unsubstituted or substituted benzyl has a structure according to formula (B):



formula (B),



wherein R^3 , R^4 , R^5 , R^6 , and R^7 are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl), (heterocyclyl)alkyl, hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-
 5 amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, azido, silyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, amino, mono-substituted amino, and di-substituted amino.

Embodiment 11. The compound of any one of the previous Embodiments, wherein
 10 at least two of R^3 , R^4 , R^5 , R^6 , and R^7 are hydrogen.

Embodiment 12. The compound of any one of the previous Embodiments, wherein R (i) is an unsubstituted or substituted C_{12} - C_{24} hydrocarbyl, (ii) comprises 0 to 6 unsaturated bonds, (iii) comprises a cyclopropyl moiety, or (iv) a combination thereof.

Embodiment 13. The compound of any one of the previous Embodiments, wherein
 15 R (i) is an unsubstituted or substituted C_{13} - C_{29} heteroalkyl, (ii) comprises 0 to 6 unsaturated bonds, or (iii) a combination thereof.

Embodiment 14. The compound of any one of the previous Embodiments, wherein R is selected from the group consisting of –



Embodiment 15. The compound of any one of the previous Embodiments, wherein (i) a is 15 to 25, or (ii) b is 1 to 4 and c is 15 to 20.

Embodiment 16. The compound of any one of the previous Embodiments, wherein
 25 (i) a is 19, (ii) b is 3 and c is 15, or (iii) b is 2 and c is 17.

Embodiment 17. The compound of any one of the previous Embodiments, wherein a is 8.


Embodiment 18. The compound of any one of the previous Embodiments, wherein R^1 and/or R^2 , independently, (i) is an unsubstituted or substituted C_{12} - C_{24} hydrocarbyl, (ii)
 30 comprises 0 to 6 unsaturated bonds, or (iii) a combination thereof.

Embodiment 19. The compound of any one of the previous Embodiments, wherein (i) R^1 , (ii) R^2 , or (iii) both R^1 and R^2 are independently selected from a C_1 - C_{30} hydrocarbyl comprising at least one cyclic moiety.

Embodiment 20. The compound of any one of the previous Embodiments, wherein
 5 (i) R^1 , (ii) R^2 , or (iii) both R^1 and R^2 are independently selected from the group consisting of aryl, arylalkyl, heteroaryl, heteroarylalkyl, and heterocycloalkyl, each of which is unsubstituted or substituted.

Embodiment 21. The compound of any one of the previous Embodiments, wherein the arylalkyl is an unsubstituted or substituted benzyl.

10 *Embodiment 22.* The compound of any one of the previous Embodiments, wherein the unsubstituted or substituted benzyl has a structure according to formula (C):



formula (C),



wherein R^8 , R^9 , R^{10} , R^{11} , and R^{12} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocyclyl,
 15 aryl(alkyl), heteroaryl(alkyl), (heterocyclyl)alkyl, hydroxy, alkoxy, acyl, cyano, halogen, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, azido, silyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, amino, mono-substituted amino, and
 20 di-substituted amino.

Embodiment 23. The compound of any one of the previous Embodiments, wherein at least two of R^8 , R^9 , R^{10} , R^{11} , and R^{12} are hydrogen.


Embodiment 24. The compound of any one of the previous Embodiments, wherein the substituent of formula (A) is a racemate, an *sn*-1 stereoisomer (e.g., glyceryl-*sn*-1-phospho), or an *sn*-3 stereoisomer (e.g., glyceryl-*sn*-3-phospho).
 25

Embodiment 25. The compound of any one of the previous Embodiments, wherein –

(i) R^1 and/or R^2 , independently, is selected from the group consisting of –



(ii) R^1 and/or R^2 , independently, is selected from the group consisting of –




Embodiment 26. The compound of any one of the previous Embodiments, wherein g is 5 to 10.


15 *Embodiment 27.* The compound of any one of the previous Embodiments, wherein g is 7.

Embodiment 28. The compound of any one of the previous Embodiments, wherein x is 1, and L is an unsubstituted or substituted C₁-C₃ hydrocarbyl.


Embodiment 29. The compound of any one of the previous Embodiments, wherein L is selected from the group consisting of an unsubstituted methyl, an unsubstituted ethyl and an unsubstituted propyl.

- Embodiment 30.* The compound of any one of the previous Embodiments, wherein
- 5 the compound of formula (I) is at least one of the following compounds or a pharmaceutically acceptable salt thereof:







Oleyloxyethyl benzyl 9-(*R*)-[(2-phosphonomethoxy)propyl]-2,6-diaminopurine (OLE-Bn-PMPDAP)










Octadecyloxyethyl-*phospho*-emtricitabine (ODE-P-FTC)











Octadecyloxyethyl benzyl-phospho-emtricitabine (ODE-Bn-P-FTC)




Oleyloxyethyl benzyl 9-(*R*)-[2-phosphono-methoxy]propyl]-2,6-diaminopurine (OLE-Bn-PMPDAP)


 Oleyloxyethyl 9-(R)-[(2-phosphonomethoxy)propyl]-2,6-diaminopurine (OLE-(R)-PMPDAP)	
 Eicosyl-phospho-RVn	 Hexadecyloxypropyl-phospho-RVn
 Octadecyloxyethyl-benzyl-phospho-RVn	 1-O-tetradecyl-2-O-benzyl-<i>sn</i>-glyceryl-phospho-RVn
 1-O-hexadecyl-2-O-benzyl-<i>sn</i>-glyceryl-phospho-RVn	 1-O-hexadecyl-2-O-(3-fluoro, 4-methoxybenzyl)-<i>sn</i>-glyceryl-phospho-RVn

 1-<i>O</i>-octadecyl-2-<i>O</i>-benzyl-<i>rac</i>-glyceryl-<i>phospho</i>-RVn	 1-<i>O</i>-octadecyl-2-<i>O</i>-benzyl-<i>sn</i>-glyceryl-<i>phospho</i>-RVn
 1-<i>O</i>-octadecyl-2-<i>O</i>-octyl-<i>sn</i>-glyceryl-<i>phospho</i>-RVn	 1-<i>O</i>-octadecyl-2-<i>O</i>-(methylcyclohexyl)-<i>sn</i>-glyceryl-<i>phospho</i>-RVn
 1-<i>O</i>-octadecyl-2-<i>O</i>-(3-fluorobenzyl)-<i>sn</i>-glyceryl-<i>phospho</i>-RVn	 1-<i>O</i>-octadecyl-2-<i>O</i>-(4-methoxybenzyl)-<i>sn</i>-glyceryl-<i>phospho</i>-RVn
 1-<i>O</i>-octadecyl-2-<i>O</i>-benzyl-benzyl-<i>sn</i>-glyceryl-<i>phospho</i>-RVn	
	


1-*O*-octadecyl-2-*O*-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn




1-*O*-octadecyl-2-*O*-(methylpyridinyl)-*sn*-glyceryl-*phospho*-RVn





1-*O*-oleyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn



1-*O*-oleyl-2-*O*-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn














1-*O*-octadecyl-*sn*-glyceryl-*phospho*-RVn


GS-441524-3',5'-cyclic monophosphate, 1-O-octadecyl-2-O-benzyl-*sn*-glyceryl ester


ODBG-benzyl-phospho-emtricitabine


ODBG-phospho-emtricitabine

1-O-octadecyl-2-O-(3-trifluoromethyl)-*sn*-glyceryl-*phospho*-RVn1-O-octadecyl-2-O-(4-trifluoromethyl)-*sn*-glyceryl-*phospho*-RVn

1-O-octadecyl-2-O-(4-cyano)-*sn*-glyceryl-phospho-RVn1-O-octadecyl-2-O-(2-cyanobenzyl)-*sn*-glyceryl-phospho-RVn1-O-octadecyl-2-O-(3-cyanobenzyl)-*sn*-glyceryl-phospho-RVn







Embodiment 31. A pharmaceutical formulation comprising –

(i) an oil, and

(ii) the compound of any one of Embodiments 1 to 30.

5 *Embodiment 32.* The pharmaceutical formulation of Embodiment 31, wherein the oil is selected from the group consisting of sesame oil, triglycerides (e.g., medium chain triglycerides), and a combination thereof.

Embodiment 33. The pharmaceutical formulation of Embodiment 31 or 32, wherein (A) the oil is present in the pharmaceutical formulation at an amount of about 60
10 % to about 95 %, about 65 % to about 95 %, about 70 % to about 95 %, about 75 % to about 95 %, about 80 % to about 95 %, or about 85 % to about 95 %, by weight, based on the weight of the pharmaceutical formulation, (B) the compound of formula (I) is present at an amount of about 0.01 % to about 40 %, about 0.01 % to about 30 %, about 0.01 % to about 20 %, about 0.01 % to about 10 %, about 0.01 % to about 5 %, by weight, based on
15 the weight of the pharmaceutical formulation, or (C) a combination thereof.

Embodiment 34. The pharmaceutical formulation of any one of Embodiments 31 to 33, wherein the pharmaceutical formulation further comprises benzyl alcohol, benzyl benzoate, ethyl alcohol, or a combination thereof; wherein, optionally, the benzyl alcohol, ethyl alcohol, benzyl benzoate, or the combination thereof is present at a total amount of
20 about 0.01 % to about 25 %, about 0.01 % to about 20 %, about 0.01 % to about 15 %, about 0.01 % to about 10 %, or about 5 % to about 10 %, by weight, based on the weight of the pharmaceutical formulation.

Embodiment 35. The pharmaceutical formulation of any one of Embodiments 31 to 34, wherein the pharmaceutical formulation (A) is formulated for injection, such as
25 intramuscular or subcutaneous injection, (B) is configured to achieve kinase bypass of the first nucleoside phosphorylation, (C) delivers antivirally significant concentrations to lungs and/or gastrointestinal tract, (D) provides sustained levels of active ingredient in plasma for about 5 to about 30 days, about 5 days to about 28 days, about 5 days to about 21 days, about 5 days to about 14 days, or about 5 days to about 7 days following
30 administration of a single dose, such as by intramuscular injection, or (E) a combination thereof.

Embodiment 36. A method for treating coronavirus infection in a mammal (e.g., a human), the method comprising administering to the mammal an effective amount of the pharmaceutical formulation of any one of Embodiments 31 to 35.

Embodiments 37. The method of Embodiment 36, wherein the effective amount of the pharmaceutical formulation is administered as a single dose.

Embodiment 38. The method of Embodiment 36 or 37, wherein the pharmaceutical formulation comprises an intermediate acting oil formulation, such as formulation F8 or formulation F7, as described herein.

Embodiment 39. A method for treating HIV infection in a mammal, the method comprising administering to the mammal an effective amount of the pharmaceutical formulation of any one of Embodiments 31 to 35.


Embodiment 40. The method of Embodiment 39, wherein the pharmaceutical formulation is a long-acting treatment.

Embodiment 41. The method of Embodiment 39 or 40, wherein the pharmaceutical formulation comprises formulation F3 or formulation F4, as described herein.


Embodiment 42. A method for treating and/or inhibiting the replication of respiratory syncytial virus (RSV), the method comprises administering to the mammal an effective amount of the pharmaceutical formulation of any one of Embodiments 31 to 35.

Embodiment 43. A method for treating a virus infection in a mammal, the method comprising administering to the mammal an effective amount of the pharmaceutical formulation of any one of Embodiments 31 to 35, wherein the virus is a RNA virus of a viral family selected from the group consisting of *Filoviridae*, *Orthomyxoviridae*, *Paramyxoviridae*, *Pneumoviridae*, *Phenuiviridae*, *Nairoviridae*, *Arenaviridae*, *Flaviviridae*, and *Coronaviridae*.

Embodiment 44. A method for producing a prodrug, the method comprising:
(i) providing a compound of formula (a) –




(ii) providing a compound of formula (b) –




formula (b);

(iii) contacting the compound of formula (a) and the compound of formula (b) to form a compound of formula (c) -




formula (c); and

5 (iv) contacting the compound of formula (c) with an acid to form a compound of formula (d) -



formula (d);

10 wherein Het is a C₁-C₃₀ hydrocarbyl comprising at least one heteroatom; Y is selected from the group consisting of hydrogen, a C₁-C₃₀ hydrocarbyl, and a pharmaceutically acceptable cation; x is 0 or 1; L is independently a C₁-C₃₀ hydrocarbyl; and R is selected from the group consisting of a C₁₀-C₃₀ hydrocarbyl and a substituent of formula (A);



wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and a C_1 - C_{30} hydrocarbyl.

Embodiment 45. The method for producing a prodrug of any of the previous

- 5 Embodiments, wherein the contacting of the compound of formula (a) and the compound of formula (b) occurs in the presence of *N,N*-dicyclohexylcarbodiimide, 4-dimethylaminopyridine, or a combination thereof.

Embodiment 46. The method for producing a prodrug of any of the previous

Embodiments, wherein the acid comprises HCl.

- 10 *Embodiment 47.* The method for producing a prodrug of any of the previous Embodiments, wherein the contacting of formula (c) with the acid occurs in the presence of tetrahydrofuran (THF).

15 EXAMPLES


The present disclosure is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other aspects, embodiments, modifications, and equivalents thereof which, after reading the description

20 herein, may suggest themselves to one of ordinary skill in the art without departing from the spirit of the present disclosure or the scope of the appended claims. Thus, other aspects of this disclosure will be apparent to those skilled in the art from consideration of the specification and practice of the disclosure disclosed herein.

Example 1 – Preparation of Compounds

- 25 In this example, several general methods were used for producing various products and/or intermediates, but other known synthesis techniques may be used.

A. Synthesis of Alkyl and Alkoxyalkyl Esters of GS-441524 5'-monophosphate



Scheme 1. Synthesis of Alkyl and Alkoxyalkyl Esters of GS-441524 5'-Monophosphate.
 Reagents: a) POCl₃, TEA, THF; b) GS-441524 acetonide, DCC/DMAP or DIC/NMI, pyridine; c) formic acid, rt or con. HCl/THF; d) Py BOP, DIEA, DMF

Synthesis of Alkyl and Alkoxyalkyl Phosphates (Scheme 1, **2a-c**)

General Method A. Long-chain alcohols **1a-c** were phosphorylated to afford phosphates **2a-c** as previously described (Ruiz, J., Beadle, J. R., Aldern, K. A., Keith, K., Hartline, C., Kern, E., Hostetler, K. Y. (2007). Synthesis and antiviral evaluation of alkoxyalkyl-phosphate conjugates of cidofovir and adefovir. *Antiviral Res.*, 75, 87-90). Briefly, a solution of the long-chain alcohol (1 eq.) and triethylamine (2 eq.) in anhydrous tetrahydrofuran (THF) was added dropwise to a solution of phosphorus oxychloride (1.5 eq.) in THF with stirring while the temperature was maintained below 20 °C. Stirring was continued for an additional hour at 0 °C, then water was added and the stirring continued overnight followed by extraction with ethyl ether. The crude solid from the ether layer was recrystallized from hexanes to afford phosphates **2a-c**.

2a Eicosyl dihydrogen phosphate¹H (400 MHz, Chloroform-*d*) δ 8.30 (s, 1H), 3.98 (t, 2H), 1.61 (m, 1H), 1.26 (br s, 16H), 0.86 (t, 1H).ESI-MS 650.38 [M-H]⁺

2b 3-(Hexadecyloxy)propyl dihydrogen phosphate ¹H NMR (400 MHz, Chloroform-*d*) δ 4.03 (dt, 2H), 3.49 (t, 2H), 3.40 (t, 2H), 1.94 (p, 2H), 1.59 – 1.55 (m, 2H), 1.26 (br s, 18 H), 0.86 (t, 3H).

2c 2-(Octadecyloxy)ethyl dihydrogen phosphate ¹H NMR (400 MHz, Chloroform-*d*) δ 4.12 (dt, 2H), 3.77 (t, 2H), 3.42 (t, 2H), 1.29 (br s, 20H), 0.94 – 0.85 (t, 3H).

20 Coupling of Phosphates **2a -c** to GS-441524 acetonide (Remdesivir nucleoside, RVn acetonide)

General Method B. *N,N*-Dicyclohexylcarbodiimide (DCC, 1.5 eq) was added to a mixture of GS-441524 acetonide (1 eq, CAS #1191237-80-5, purchased from Ontario Chemicals), a long-chain dihydrogen phosphate (1.0 eq), and 4-dimethylaminopyridine (DMAP, 1.0 eq) in dry pyridine, and then the mixture was heated to 90 °C and stirred for 24h. Water was added to quench the reaction and pyridine was evaporated under vacuum. The residue was adsorbed onto silica gel and purified by flash column chromatography on silica gel 60. Gradient elution (CH₂Cl₂/methanol 10-20%) afforded the protected phosphodiester compound.

30 **General Method C.** *N,N*-Diisopropylcarbodiimide (DIC, 3.3 mmol) was added to a mixture of GS-441524 acetonide (1.65 mmol), lipid phosphate (1.65 mmol), and 1-methylimidazole (NMI, 406 mg, 4.95 mmol) in dry pyridine (30 mL), and then the

mixture was stirred for 48 hours at room temperature until analysis of the reaction mixture by TLC indicated substantial formation of coupled product. Water (5 mL) was then added, and the mixture was concentrated on a rotary evaporator. The residue was adsorbed onto silica gel and purified by flash column chromatography on silica gel 60. Gradient elution (100% CH₂Cl₂ to CH₂Cl₂/20% methanol) afforded the protected phosphodiester analogs.

3a Eicosyl-*phospho*-RVn acetonide. GS-441524 acetonide was coupled to **2a** according to General Method C. The structure was confirmed by ESI-MS 690.50 [M-H]⁻.

3b 3-(Hexadecyloxy)propyl-*phospho*-RVn acetonide. GS-441524 acetonide was coupled to **2b** according to General Method B. N,N-Dicyclohexylcarbodiimide (DCC, 619 mg, 3 mmol) was added to a mixture of GS-441524 acetonide (300 mg, 0.91 mmol), 3-(hexadecyloxy)propyl phosphate (**2b**, 414 mg, 1.10 mmol), and 4-dimethylaminopyridine (DMAP, 122 mg, 1.0 mmol) in 25 mL of dry pyridine, and then the mixture was heated to 90 °C and stirred for 24 hours. Pyridine was then evaporated and the residue was purified by flash column chromatography on silica gel 60. Gradient elution (CH₂Cl₂/methanol 10-20%) afforded 423 mg (67% yield) of Compound **3b**. ¹H NMR (500 MHz, chloroform-d) δ 8.42 (s, 1H), 7.98 (s, 1H), 7.70 (s, 2H), 6.22 (d, J = 6.0 Hz, 1H), 5.68 (d, J = 6.2 Hz, 1H), 5.15 (d, J = 1.0 Hz, 1H), 4.70 (dd, J = 3.8, 0.9 Hz, 1H), 4.48 – 4.42 (m, 1H), 4.26 (ddd, J = 11.2, 8.5, 2.6 Hz, 1H), 4.15 (ddd, J = 11.1, 8.5, 2.6 Hz, 1H), 4.02 (dt, J = 8.5, 6.3 Hz, 2H), 3.49 (t, J = 6.1 Hz, 2H), 3.40 (t, J = 6.1 Hz, 2H), 1.95 (p, J = 6.2 Hz, 2H), 1.54 (tt, J = 7.4, 6.1 Hz, 2H), 1.31 (s, 3H), 1.32 – 1.24 (m, 26H), 0.94 – 0.85 (m, 3H). ESI-MS 691.6 [M-H]⁻.

3c 2-(Octadecyloxy)ethyl-*phospho*-RVn acetonide. GS-441524 acetonide was coupled to **2c** according to General Method B. N,N-Dicyclohexylcarbodiimide (DCC, 0.3 g, 1.4 mmol) was added to a mixture of GS-441524 acetonide (0.23 g, 0.7 mmol), phosphate **2c** (0.27 g, 0.68 mmol), and 4-dimethylaminopyridine (DMAP, 0.07 g, 0.6 mmol) in 10 mL of dry pyridine, and then the mixture was heated to 90 °C and stirred for 3 days. Pyridine was then evaporated, and the residue was purified by flash column chromatography on silica gel 60. Gradient elution (CH₂Cl₂/methanol 10-20%) afforded 0.22 g (45% yield) of phosphodiester **3c**.

30

Synthesis of 4a-c: Removal of the acetonide protecting group

General Method D. (HCl/THF) Concentrated HCl (0.1 mL) in tetrahydrofuran (THF, 1 mL) was added to a stirred solution of acetonide-protected (2',3'-isopropylidene) phosphodiester (0.25 mmol) in THF (10 mL) at room temperature. The mixture was stirred for 3 hours and then sodium bicarbonate (50 mg) and water (2 mL) were added. After stirring an additional 15 minutes the solvents were evaporated and cold water (10 mL) was added to the residue. The crude product was collected by vacuum filtration and dried under vacuum. Purification by flash column chromatography (100% CH₂Cl₂ to CH₂Cl₂/35% methanol) yielded pure phosphodiester analogs.

General Method E. Acetonide analogs (1 mmol) were added to formic acid (25 mL) at room temperature and stirred. The reaction was monitored by TLC until deprotection was complete at about 4 hours. Formic acid was removed by rotary evaporation and the residue was co-evaporated with EtOH (2 x 25 mL), then adsorbed onto silica gel and purified by flash column chromatography. Gradient elution (100% CH₂Cl₂ to CH₂Cl₂/35% methanol) afforded products.

4a Eicosyl-phospho-RVn – Prepared from **3a** according to General Method E. The structure was confirmed by ESI-MS 650.38 [M-H]⁻.

4b 3-(Hexadecyloxy)propyl-phospho-RVn. Prepared from **3b** according to General Method D. Concentrated HCl (0.1 mL) in tetrahydrofuran (THF, 1 mL) was added to a stirred solution of **3b** (100 mg, 0.14 mmol) in THF (10 mL) at room temperature. The mixture was stirred for 3 hours and then sodium bicarbonate (50 mg) and water (2 mL) were added. After stirring an additional 15 minutes the solvents were evaporated and cold water (10 mL) was added to the residue. The solid product was collected by vacuum filtration and dried under vacuum to yield **4b** (79 mg, 87% yield) as an off-white solid. ¹H NMR (500 MHz, CDCl₃-methanol-*d*₄) δ 8.42 (s, 1H), 7.98 (s, 1H), 7.70 (s, 1H), 6.22 (d, *J* = 6.0 Hz, 1H), 5.70 (d, *J* = 6.0 Hz, 1H), 5.12 (d, *J* = 4.2 Hz, 1H), 4.55 (ddd, *J* = 5.5, 2.7, 0.9 Hz, 1H), 4.40 (dtd, *J* = 6.8, 2.6, 0.8 Hz, 1H), 4.33 – 4.27 (m, 2H), 4.25 (ddd, *J* = 11.1, 8.4, 2.6 Hz, 1H), 4.16 (ddd, *J* = 11.3, 8.5, 2.6 Hz, 1H), 4.02 (dt, *J* = 8.5, 6.3 Hz, 2H), 3.49 (t, *J* = 6.1 Hz, 2H), 3.40 (t, *J* = 6.1 Hz, 2H), 1.95 (p, *J* = 6.2 Hz, 2H), 1.59 – 1.50 (m, 1H), 1.34 – 1.24 (m, 23H), 0.94 – 0.85 (m, 3H). ESI MS: 652.39 [M-H]⁻. Purity by HPLC: 99.7%

4c 2-(Octadecyloxy)ethyl-phospho-RVn Prepared from **3c** according to General Method D. Concentrated HCl (0.3 mL) was added slowly to a stirred solution of **3c** (0.2 g, 0.28 mmol) in THF (10 mL) at 0 °C. The mixture was allowed to warm to room temperature overnight and then was diluted with water (2 mL) and adjusted to pH = 8 by adding saturated sodium bicarbonate. The product was extracted with chloroform (3 x 30 mL) and the organic layer was concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel. Elution with 20% MeOH/CH₂Cl₂ gave 0.10 g (55% yield) of compound **4c**. ¹H NMR (400 MHz, CDCl₃-methanol-*d*₄) δ ppm 7.89 (s, 1 H), 6.94 (d, *J*=4.65 Hz, 1H), 6.89 (d, *J*=4.65 Hz, 1H), 4.40 (d, *J*=4.65 Hz, 2H), 4.21 - 4.28 (m, 1H), 4.12 - 4.20 (m, 1H), 4.04 - 4.12 (m, 1H), 3.91 (d, *J*=4.89 Hz, 2H), 3.46 - 3.57 (m, 2H), 3.42 (td, *J*=6.85, 1.96 Hz, 2H), 3.34 (dt, *J*=3.18, 1.59 Hz, 2H), 1.53 (d, *J*=6.85 Hz, 2H), 1.20 - 1.37 (m, 30H), 0.89 (t, *J*=6.97 Hz, 3H). ESI MS: 666.43 [M-H]⁻. Purity by HPLC 98.4%.

B. Synthesis of 2-(Octadecyloxy)ethyl benzyl phospho-RVn (long-acting formulation) (Scheme 1, 6c)

Compound **3c** (160 mg, 0.22 mmol), benzyl alcohol (48 mg, 0.45 mmol), diisopropylethylamine (DIEA, 58 mg, 0.45 mmol), and (1H-benzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 230 mg, 0.45 mmol) in dry DMF (5 mL) were stirred at room temperature for 3 h. DMF was then evaporated, and the residue was dissolved in ethyl acetate (50 mL) and washed with saturated NaHCO₃ (3 x 10 mL). The organic layer was dried over MgSO₄ and concentrated. The residue was purified by column chromatography on silica gel, eluting with chloroform/methanol (0-15%) to yield **5c** (60 mg, 35% yield). ¹H NMR (400 MHz, CDCl₃-CD₃OD) δ ppm 7.87 (d, *J*=4.03 Hz, 1H), 7.27-7.37 (m, 5H), 6.91-6.95 (m, 1H), 6.83-6.89 (m, 1H), 5.41 (d, *J*=6.97 Hz, 1H), 4.92-5.06 (m, 3 H), 4.54-4.60 (m, 1H), 4.24-4.31 (m, 2H), 4.07-4.15 (m, 2H), 3.53-3.60 (m, 2H), 3.38 - 3.51 (m, 2H), 3.32-3.37 (m, 2H), 1.78 - 1.96 (m, 2H), 1.75 (s, 3H), 1.50-1.60 (m, 2H), 1.42 (s, 3H), 1.15 - 1.38 (m, 30H), 0.89(t, *J*=6.54 Hz, 3H). ESI MS: 798.51 [M+H]⁺, 820.56 [M+Na]⁺.


To a solution of **5c** (60 mg, 0.075mmol) in THF (2 mL), concentrated HCl (0.1 mL) was added at 0 °C. After 20 minutes, the ice bath was removed and the reaction was monitored by TLC. After 3 hours, the ice bath was returned and the mixture was

neutralized with sat. NaHCO₃. The mixture was concentrated under vacuum and the residue was purified by column chromatography (silica gel, dichloromethane/methanol 10-20%) to give 35 mg (62% yield) of **6c**. ¹H NMR (400 MHz, CDCl₃+methanol d₄) δ ppm 7.84-7.90 (m, 1H), 7.29-7.38 (m, 5H), 6.89-6.93 (m, 1H), 6.82-6.86 (m, 1H), 5.03 (d, *J*=11.36 Hz, 2H), 4.76-4.81 (m, 1H), 4.40-4.45 (m, 1H), 4.30-4.37 (m, 1H), 4.17 - 4.31 (m, 2H), 4.06-4.14 (m, 2H), 3.54-3.60 (m, 2H), 3.39-3.47 (m, 2H), 3.33-3.37 (m, 2H), 3.12-3.18 (m, 2H), 1.82-1.91 (m, 2H), 1.49-1.59 (m, 2H), 1.20-1.37 (m, 30H), 0.89 (t, *J*=6.60 Hz, 3H). ESI MS: 758.32 [M+H]⁺, 780.43 [M+Na]⁺.

10 C. Synthesis of 1-O-alkyl-2-O-substituted-*sn*-glyceryl esters of GS-441524 5'-monophosphate

The following scheme (Scheme 2) depicts embodiments of synthesis methods that were used to produce the following embodiments of 1-O-alkyl-2-O-substituted-*sn*-glyceryl esters of GS-441524 5'-monophosphate.

15



Scheme 2. Synthesis of 1-*O*-Alkyl-2-*O*-substituted-*sn*-glycerols. *Reagents:* a) R₁ bromides or methanesulfonates, NaH, DMF; b) 80% CH₃COOH, reflux; c) trityl chloride, TEA, DMAP, CH₂Cl₂; d) R₂ bromides or methanesulfonates, NaH, DMF e) acidic deprotection.

5 Synthesis of 1-*O*-Alkyl-*sn*-glycerols (Scheme 2, 10 a-d)

General Method F. In this embodiment, alkylation of 2,3-isopropylideneglycerol using an alkylmethanesulfonate was performed as described in the literature (Fernández, D.M.; Contreras, L.J.; Moreno, B.M.; Silva, E.G.; Mayorga, H.W. Enantiomeric synthesis of natural alkylglycerols and their antibacterial and antibiofilm activities. *Nat. Prod. Res.* 10 **2019**, 1–7.).

 Sodium hydride and DMF were stirred in a flask. Isopropylidene glycerol was added slowly, due to expected hydrogen evolution, while cooling, if necessary, to keep the temperature less than 35 °C. After stirring an additional 30 minutes, alkyl methanesulfonate was added all at once and stirred vigorously for 5 hours. The reaction 15 mix was poured onto crushed ice and stirred gently. The solid was collected on a frit funnel, and then washed with water. To achieve deprotection, the filter cake was added to 80% acetic acid and heated at 80 °C for 1 hour. The flask was cooled and the product crystallized, and was collected via vacuum filtration and dried. Crude product was recrystallized in hexanes or purified by flash column chromatography on silica gel 60.

20 **General Method G.** (Alkylation using a 1-bromoalkane/alkene as described in the literature: (Halldorsson, A., et al. *Tetrahedron: Asymmetry*, **2004**, 15, 2893-2899)).

 Isopropylideneglycerol (1 eq), 1-bromoalkane/alkene (1 eq) and tetrabutylammonium bromide (0.2 eq) were stirred vigorously in a round-bottomed flask. Ground potassium hydroxide (2 eq) was added slowly, and the mixture stirred for about 15 25 hours at 35–40 °C in an oil bath. The alkylated product was extracted into hexanes and the organic phase was washed with H₂O, then evaporated to yield the 1-*O*-alkyl-2,3-isopropylidene-*sn*-glycerol.

Deprotection: The products were refluxed overnight with p-toluenesulfonic acid (10 mol%) in THF/water. After concentration under vacuum, the residue was dissolved in 30 diethyl ether, washed with water and brine solution, dried over anhydrous magnesium sulfate, and solvent removal was achieved *in vacuo* on a rotary evaporator to afford the 1-*O*-alkyl-*sn*-glycerol.

10a. 1-*O*-Tetradecyl-*sn*-glycerol. Synthesized according to General Method F. Analytical data was consistent with literature values (Barragán, C.A.; Silva, E.G.; Moreno, B.M.; Mayorga, H.W. Inhibition of quorum sensing by compounds from two *Eunicea* species and synthetic saturated alkylglycerols. *Vitae* **2018**, 25, 92–103.)

5 **10b.** 1-*O*-Hexadecyl-*sn*-glycerol was purchased from Bachem America

10c. 1-*O*-Octadecyl-*sn*-glycerol was purchased from Bachem America

10d. 1-*O*-Oleoyl-*sn*-glycerol. Synthesized according to General Method G. A mixture of oleyl bromide (541 mg, 1.63 mmol), Bu₄NBr (0.2 eq), 2,3-isopropylidene-*sn*-glycerol (1 eq), and KOH (powder, 2.5 eq) was stirred at 40 °C overnight. A subsequent
10 work up gave 583 mg crude **9d** as an oil. Crude **9d** was treated with p-TsOH.H₂O (0.15 eq.) in refluxing THF (6 mL) and H₂O (2.5 mL) overnight. Purification of the crude oil (540 mg) by flash column chromatography (MeOH in DCM 0–8%) afforded 420 mg 1-*O*-oleyl-*sn*- glycerol **10d** as an oil. Yield 75% (two steps). ¹H NMR (CDCl₃) δ 5.36–5.33 (m, 2H), 3.86–3.85 (m, 1H), 3.72 (dd, 1H), 3.62 (dd, 1H), 3.52 (dd, 1H), 3.46 (dd, 1H),
15 3.50–3.42(m, 2H), 2.02–1.99 (m, 4H), 1.59–1.55 (quintet, 2H), 1.35–1.26 (m, 22H), 0.88 (t, 3H) ESI-MS : 343.67 [M+H]⁺, 365.61 [M+Na]⁺.

Synthesis of 1-*O*-Alkyl-2-*O*-substituted-*sn*-glycerols (Scheme 2, **12a-p**)

General Method H. Protection of the 3-hydroxy group of 1-*O*-substituted-*sn*-
20 glycerols was carried out as described in the literature (e.g., Kini, G. D., Hostetler, S. E., Beadle, J. R., Aldern, K. A. Synthesis and antiviral activity of 1-*O*-octadecyl-2-*O*-alkyl-*sn*-glycero-3-foscarnet conjugates in human cytomegalovirus-infected cells, *Antiviral Research*, **1997**, 36, 115; and Huang, Z., Szoka, Z. (2008); Sterol-Modified Phospholipids: Cholesterol and Phospholipid Chimeras with Improved Biomembrane Properties. *J. Am.*
25 *Chem. Soc.*, 130, 15702-15712). Triethylamine (1.5 eq) was added to a solution of 1-*O*-alkyl-*sn*-glycerol (1 eq), N,N-dimethylaminopyridine (DMAP, 0.1 eq), and triphenyl chloride (TrCl, 1.5 eq) in anhydrous dichloromethane, and the mixture was stirred 18 hours. The reaction mixture was then quenched with water, evaporated, and adsorbed onto silica gel and purified by flash column chromatography over silica gel. An increasing
30 gradient of ethyl acetate in hexanes (0–20%) eluted the proper fractions.

11a 1-*O*-Tetradecyl-3-*O*-trityl-*sn*-glycerol – Prepared as described in the literature (Huang, Z., Szoka, Z. (2008). Sterol-Modified Phospholipids: Cholesterol and

Phospholipid Chimeras with Improved Biomembrane Properties. *J. Am. Chem. Soc.*, 130, 15702-15712).

11b 1-*O*-Hexadecyl-3-*O*-trityl-*sn*-glycerol - Prepared as described in the literature (Huang, Z., Szoka, Z. (2008). Sterol-Modified Phospholipids: Cholesterol and

5 Phospholipid Chimeras with Improved Biomembrane Properties. *J. Am. Chem. Soc.*, 130, 15702-15712).

11c 1-*O*-octadecyl-3-*O*-trityl-*sn*-glycerol. Prepared from **10c** according to General Method H. Yield 87%. ¹H NMR (CDCl₃): δ 0.9 (t, 3H), 1.3 (bs, 30H), 1.55 (m, 4H) 3.2 (m, 2H), 3.4-3.6 (m, 3H), 3.95 (m, 1H) 7.2-7.5 (m, 15H).

10 **11d** 1-*O*-oleyl-3-*O*-trityl-*sn*-glycerol. Prepared from **10d** according to General Method H. Yield 77%. ¹H NMR (CDCl₃): δ 0.88 (t, J = 7.2, 3H); 1.27 (br, 22H); 1.55 (m, 2H); 2.0 (m, 4H); 2.40 (br, 1H); 3.20 (m, 2H); 3.41-3.56 (m, 4H); 3.95 (m, 1H); 5.35 (m, 2H); 7.25 (m, 9H); 7.45 (m, 6H). ESI-MS 607.75 [M+Na]⁺

General Method I. Alkylation and deprotection of 1-*O*-alkyl-3-*O*-trityl-*sn*-glycerols was done as described previously (Kini, G. D., Hostetler, S. E., Beadle, J. R., Aldern, K. A. Synthesis and antiviral activity of 1-*O*-octadecyl-2-*O*-alkyl-*sn*-glycero-3-foscarnet conjugates in human cytomegalovirus-infected cells, *Antiviral Research*, **1997**, 36, 115). Briefly, sodium hydride (2.5 eq.) was added to a stirred solution of 1-*O*-alkyl-3-*O*-trityl-*sn*-glycerol (1 eq) in DMF at 0 °C. After 20 minutes, a bromo or
15 methanesulfonate derivative of **R**₂- (1.8 eq.) was added. The reaction mixture was then stirred at room temperature for 5 hours, or until the reaction was substantially complete by TLC. Work up and column chromatography gave 1-*O*-alkyl-2-*O*-substituted-3-*O*-trityl-*sn*-glycerols, which were detritylated with acid. Work up and column chromatography afforded 1-*O*-alkyl-2-*O*-substituted-*sn*-glycerols.
20

25 **12a** 1-*O*-Tetradecyl-2-*O*-benzyl-*sn*-glycerol – Prepared from **11a** and benzyl bromide according to General Method I. Structure was confirmed by ESI-MS: 401.51 [M+Na]⁺

12b 1-*O*-Hexadecyl-2-*O*-benzyl-*sn*-glycerol - Prepared from **11b** and benzyl bromide according to General Method I. ¹H NMR (300 MHz, CDCl₃): δ 7.36-7.27 (m, 5H), 4.65 (m, 2H), 3.79-3.59 (m, 5H), 3.55 (t, 2H), 1.57-1.51 (m, 2H), 1.29 (br s, 26H), 0.89 (t, 3H).
30

12c 1-*O*-Hexadecyl-2-*O*-(3-fluoro-4-methoxybenzyl)-*sn*-glycerol was prepared from **11b** and 3-fluoro-4-methoxybenzyl bromide according to General Method I.

12d 1-*O*-Octadecyl-2-*O*-benzyl-*sn*-glycerol,

12e 1-*O*-Octadecyl-2-*O*-benzyl-*rac*-glycerol,

5 **12f** 1-*O*-Octadecyl-2-*O*-octyl-*sn*-glycerol,

12g 1-*O*-Octadecyl-2-*O*-(cyclohexylmethyl)-*sn*-glycerol,

12h 1-*O*-Octadecyl-2-*O*-(3-fluorobenzyl)-*sn*-glycerol,

12i 1-*O*-Octadecyl-2-*O*-(4-methoxybenzyl)-*sn*-glycerol,

12j 1-*O*-Octadecyl-2-*O*-(3-fluoro-4-methoxybenzyl)-*sn*-glycerol, and

10 **12k** 1-*O*-Octadecyl-2-*O*-(pyridine-3-yl-methyl)-*sn*-glycerol were prepared from **11c** and the appropriate bromide according to General Method I.

12l 1-*O*-Olelyl-2-*O*-benzyl-*sn*-glycerol. Sodium hydride (1.3 eq.) was added to **11d** (531 mg, 0.91 mmol) in DMF (4 mL) at 0 °C. The resulting mixture was stirred at room temperature for 1 hour before benzyl bromide (1.3 eq.) was added. The reaction
15 mixture was stirred at room temperature overnight. Work up and column chromatography afforded 354 mg crude product and 200 mg of **11d** was also recovered. Deprotection afforded **12l**. ¹H NMR (300 MHz, Chloroform-*d*) δ 7.35 – 7.26 (m, 4H), 5.36 – 5.32 (m, 2H), 3.84 – 3.62 (m, 5H), 3.54 – 3.46 (m, 2H), 3.44 (t, 2H), 2.88 – 2.75 (m, 2H), 2.02 (m, 4H), 1.54 – 1.50 (pentet 2H), 1.29 (br s, 22H), 0.88 (t, 3H). 455.73 [M+Na]⁺

20 **12m** 1-*O*-Olelyl-2-*O*-(3-fluoro-4-methoxybenzyl)-*sn*-glycerol. Prepared from **11d** and 3-fluoro-4-methoxybenzyl bromide according to General Method I. ¹H NMR (300 MHz, Chloroform-*d*) δ 7.12 (m, 2H), 6.95 (t, 1H), 5.36 – 5.32 (m, 2H), 4.65 – 4.52 (dd, 2H), 3.75 – 3.70 (m, 2H), 3.67 – 3.60 (m, 2H), 3.57 – 3.55 (m, 2H), 3.45 (t, 2H), 2.01 – 1.97 (m, 2H), 1.28 (br s, 16H), 0.87 (t, 3H). ESI-MS: 503.79 [M+Na]⁺

25 **12n** 3-*O*-Octadecyl-2-*O*-benzyl-*sn*-glycerol.


12o 1-*O*-Octadecyl-2-*O*-(3-trifluoromethyl)benzyl)-*sn*-glycerol ESI-MS 525.47 [M+Na]⁺.

12p 1-*O*-Octadecyl-2-*O*-(4-trifluoromethyl)benzyl)-*sn*-glycerol. ESI-MS 503.35 [M+H]⁺.

30

Synthesis of 1-O-Alkyl-2-O-substituted-sn-glycerol esters of GS-441524 5'-monophosphate (Scheme 3. 15a-p)

The following scheme depicts embodiments of synthesis steps used to produce 1-O-alkyl-2-O-substituted-sn-glycerol esters of GS-441524 5'-monophosphate.



Scheme 3. Synthesis of 1-O-Alkyl-2-O-substituted-sn-glycerol esters of GS-441524 5'-monophosphate. *Reagents:* a) POCl₃ or bis(trichloroethyl) chlorophosphate/zinc powder; b) DCC/DMAP or DIC/NMI, pyridine; c) formic acid or con HCl/THF.

5 **General Method J.** Phosphorylation of 1-O-alkyl-2-O-substituted-sn-glycerols was accomplished as described in the literature (Kates, M., Adams, G. A., Blank, M. L., Snyder, F. M. (1991), Chemical synthesis and physiological activity of sulfonium analogues of platelet activating factor. *Lipids*, 26, 1095-1101). 1-O-alkyl-2-O-substituted-sn-glycerols (11.5 mmol) and 1-methylimidazole (14.4 mmol) were dissolved in dry
10 pyridine (100 mL) and stirred at room temperature. A solution of bis(trichloroethyl) chlorophosphate (5.5 g, 14.4 mmol) in diethyl ether (20 mL) was added dropwise over 10 minutes, then the mixture was stirred overnight. Analysis by TLC showed complete phosphorylation. Water (10 mL) was added to quench excess reagent and then the mixture was concentrated by rotary evaporation, and co-evaporated with toluene to
15 remove pyridine. The residue was adsorbed onto silica gel 60 (ca. 30 g) and purified by flash column chromatography. Gradient elution 100% hexanes to 25% EtOAc/hexanes was used to isolate protected phosphorylated product.

The products (9.65 mmol) were dissolved in a mixture of chloroform (50 mL) and glacial acetic acid (90 mL), and then vigorously stirred and cooled with an ice water bath.
20 Zinc powder (5 g) was added to the mixture, stirred for 1 hour and then the ice water bath was removed, and stirring was continued for another 2 hours. The remaining zinc was removed by vacuum filtration and the clear filtrate was concentrated by rotary evaporation. The residue was taken up in 20% MeOH/CH₂Cl₂ (250 mL) and extracted with 1 M HCl (3 x 50 mL), then the organic layer was concentrated and co-evaporated
25 with ethanol (2 x 50 mL). The waxy residue was dissolved in 1,4-dioxane, frozen, then lyophilized *in vacuo* (18 h) to provide glyceryl phosphates.

Compounds **13a**, **13b**, **13c**, **13d**, **13e**, **13f**, **13g**, **13h**, **13i**, **13j**, and **13k** were prepared according to General Method J.

13i 1-O-Oleyl-2-O-benzyl-sn-glycerol-3-phosphate. Prepared according to
30 General Method A. ¹H NMR (300 MHz, Chloroform-*d*) δ 7.36 – 7.23 (m, 5H), 5.36 – 5.32 (m, 2H), 4.69 (d, *J* = 11.9 Hz, 1H), 4.62 (d, *J* = 11.8 Hz, 1H), 4.11 – 4.09 (m, 2H), 3.80 –

3.77 (m, 2H), 3.76 – 3.69 (m, 1H), 3.53 – 3.47 (m, 1H), 3.42 (t, 2H), 2.00 (tq, $J = 7.1$, 3.7 Hz, 4H), 1.50 (m, 2H), 1.26 (br s, 22H), 0.87 (t, 3H). ESI-MS: 513.72 $[M + 1]^+$

13m 1-O-Oleyl-2-O-(3-fluoro-4-methoxybenzyl)-*sn*-glyceryl-3-phosphate.

Prepared according to General Method A.

5 **13n** 3-O-Octadecyl-2-O-benzyl-*sn*-glyceryl-1-phosphate. Prepared according to General Method A.

13o 1-O-Octadecyl-2-O-(3-trifluoromethyl)benzyl-*sn*-glyceryl-3-phosphate.

Prepared according to General Method A. ESI-MS 581.49 $[M-H]^-$.

13p 1-O-Octadecyl-2-O-(4-trifluoromethyl)benzyl-*sn*-glyceryl-3-phosphate.

10 Prepared according to General Method A. ESI-MS 581.47 $[M-H]^-$.

Coupling of Phosphates **13a-p** to GS-441524 acetamide (Remdesivir nucleoside, RVn acetamide)

15 **14a** 1-O-Tetradecyl-2-O-benzyl-*sn*-glyceryl-*phospho*-RVn acetamide – Prepared from GS-441524 acetamide and **13a** according to General Method C. Structure was confirmed by ESI-MS 770.50 $[M-H]^-$.

14b 1-O-Hexadecyl-2-O-benzyl-*sn*-glyceryl-*phospho*-RVn acetamide - Prepared from GS-441524 acetamide and **13b** according to General Method C.

20 **14c** 1-O-Hexadecyl-2-O-(3-fluoro-4-methoxy-benzyl)-*sn*-glyceryl-*phospho*-RVn acetamide - Prepared from GS-441524 acetamide and **13c** according to General Method C.

14d 1-O-Octadecyl-2-O-benzyl-*sn*-glyceryl-*phospho*-RVn acetamide - Prepared from GS-441524 acetamide and **13d** according to General Method B. N,N-Dicyclohexylcarbodiimide (DCC, 310 mg, 1.5 mmol) was added to a mixture of acetamide (300 mg, 0.91 mmol), phosphate **13d** (515 mg, 1.0 mmol), and 4-dimethylaminopyridine (DMAP, 122 mg, 1.0 mmol) in 25 mL of dry pyridine, and then the mixture was heated to 90 °C and stirred for 24h. Pyridine was then evaporated and the residue was purified by flash column chromatography on silica gel 60. Gradient elution (CH_2Cl_2 /methanol 10-20%) afforded 210 mg (28% yield) of compound **14d**. ESI MS 826.58 $[M-H]^-$.

30 **14e** 1-O-Octadecyl-2-O-benzyl-*rac*-glyceryl-*phospho*-RVn acetamide. - Prepared from GS-441524 acetamide and **13e** according to General Method C.

14f 1-O-Octadecyl-2-O-octyl-*sn*-glyceryl-*phospho*-RVn acetamide - Prepared from GS-441524 acetamide and **13f** according to General Method C.

14g 1-*O*-Octadecyl-2-*O*-(cyclohexylmethyl)-*sn*-glyceryl-*phospho*-RVn acetone – May be prepared from GS-441524 acetone and **13g** according to General Method C.

14h 1-*O*-Octadecyl-2-*O*-(3-fluoro-benzyl)-*sn*-glyceryl-*phospho*-RVn acetone - Prepared from GS-441524 acetone and **13h** according to General Method C.

5 **14i** 1-*O*-Octadecyl-2-*O*-(4-methoxy-benzyl)-*sn*-glyceryl-*phospho*-RVn acetone. – May be prepared from GS-441524 acetone and **13i** according to General Method C.

14j 1-*O*-Octadecyl-2-*O*-(3-fluoro-4-methoxy-benzyl)-*sn*-glyceryl-*phospho*-RVn acetone - Prepared from GS-441524 acetone and **13j** according to General Method C.

14k 1-*O*-Octadecyl-2-*O*-(pyridine-3-yl-methyl)-*sn*-glyceryl-*phospho*-RVn acetone – May be prepared from GS-441524 acetone and **13k** according to General Method C.

14l 1-*O*-Olelyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn acetone Prepared from GS-441524 acetone and **13l** according to General Method B.

14m 1-*O*-Olelyl-2-*O*-(3-fluoro-4-methoxy-benzyl)-*sn*-glyceryl-*phospho*-RVn acetone Prepared from GS-441524 acetone and **13m** according to General Method B.

14n 3-*O*-Octadecyl-2-*O*-benzyl-*sn*-glyceryl-1-*phospho*-RVn acetone Prepared from GS-441524 acetone and **13n** according to General Method B.

14o 1-*O*-Octadecyl-2-*O*-(3-trifluoromethyl)benzyl-*sn*-glyceryl-*phospho*-RVn acetone Prepared from GS-441524 acetone and **13o** according to General Method B.
20 ESI-MS 894.66 [M-H]⁺.

14p 1-*O*-Octadecyl-2-*O*-(4-trifluoromethyl)benzyl-*sn*-glyceryl-*phospho*-RVn acetone Prepared from GS-441524 acetone and **13p** according to General Method B.
ESI-MS 894.53 [M-H]⁺.

25 Removal of acetone protecting group

15a 1-*O*-Tetradecyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn – Prepared from Compound **14a** according to General Method E and isolated as an off-white powder. Structure was confirmed by ESI-MS [M-H]⁺ = 730.41.

15b 1-*O*-Hexadecyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn. Prepared from Compound **14b** according to General Method E. ¹H NMR (500 MHz, DMSO-d₆) δ 7.94 (s, 1H), 7.89 (s, 1H), 7.81 (s, 1H), 7.32 – 7.25 (m, 3H), 7.22 (ddd, J = 8.7, 5.4, 2.6 Hz, 1H), 6.88 (d, J = 4.5 Hz, 1H), 6.80 (d, J = 4.5 Hz, 1H), 6.24 (s, 1H), 5.95 (d, J = 4.0 Hz,

1H), 4.55 (q, J = 12.1, 12.1, 12.1 Hz, 3H), 4.09 (dt, J = 6.7, 4.3, 4.3 Hz, 1H), 3.92 (d, J = 4.5 Hz, 1H), 3.78 (dtt, J = 24.4, 7.8, 7.8, 4.4, 4.4 Hz, 2H), 3.66 – 3.55 (m, 3H), 3.43 (dd, J = 10.6, 3.5 Hz, 1H), 3.32 – 3.28 (m, 2H), 1.42 (q, J = 6.5, 6.5, 6.0 Hz, 2H), 1.20 (d, J = 7.7 Hz, 24H), 0.83 (t, J = 7.0, 7.0 Hz, 3H). LC/MS purity = 99.8%; [M+H]⁺ 760.6.

5 **15c** 1-*O*-Hexadecyl-2-*O*-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn. Prepared from Compound **14c** according to General Method E. ¹H NMR (500 MHz, DMSO-d₆) δ 7.89 (s, 1H), 7.80 (s, 1H), 7.11 (d, J = 12.3 Hz, 1H), 7.04 (d, J = 6.1 Hz, 2H), 6.87 (d, J = 4.6 Hz, 1H), 6.81 (d, J = 4.5 Hz, 1H), 6.17 (s, 1H), 4.56 (d, J = 5.0 Hz, 1H), 4.52 – 4.41 (m, 2H), 4.10 (s, 1H), 3.93 (q, J = 5.4, 5.2, 5.2 Hz, 1H), 3.78 (s, 3H), 3.63 (s, 2H), 3.57 (d, J = 4.6 Hz, 1H), 3.41 (dd, J = 10.4, 3.5 Hz, 1H), 3.29 (s, 3H), 1.41 (d, J = 6.5 Hz, 2H), 1.25 – 1.17 (m, 24H), 0.83 (t, J = 7.0, 7.0 Hz, 3H). LC/MS purity 99.8%; [M+H]⁺ 808.9.

15f 1-*O*-Octadecyl-2-*O*-octyl-*sn*-glyceryl-*phospho*-RVn – Prepared from Compound **14f** according to General Method E. ¹H NMR (500 MHz, DMSO-d₆) δ 7.89 (s, 1H), 7.79 (s, 1H), 6.88 (d, J = 4.5 Hz, 1H), 6.80 (d, J = 4.5 Hz, 1H), 6.10 (s, 1H), 5.92 (s, 1H), 4.56 (t, J = 5.2, 5.2 Hz, 1H), 4.08 (t, J = 5.7, 5.7 Hz, 1H), 3.91 (q, J = 5.0, 4.9, 4.9 Hz, 1H), 3.79 (d, J = 18.1 Hz, 3H), 3.54 (d, J = 19.3 Hz, 3H), 3.46 – 3.35 (m, 5H), 3.33 (s, 2H), 3.27 – 3.23 (m, 1H), 1.41 (dt, J = 16.0, 7.4, 7.4 Hz, 4H), 1.21 (d, J = 4.5 Hz, 36H), 0.83 (td, J = 7.1, 7.0, 5.7 Hz, 6H). LC/MS purity 99.7%; [M+H]⁺ 810.7.

20 **15g** 1-*O*-Octadecyl-2-*O*-(ethylcyclohexyl)-*sn*-glyceryl-*phospho*-RVn- May be prepared from Compound **14g** according to General Method E.

15h 1-*O*-Octadecyl-2-*O*-(3-fluoro-benzyl)-*sn*-glyceryl-*phospho*-RVn- Prepared from Compound **14h** according to General Method E. ¹H NMR (500 MHz, DMSO-d₆) δ 8.47 (s, 2H), 7.88 (s, 2H), 7.36 – 7.26 (m, 1H), 7.12 (q, J = 8.4, 6.9, 6.9 Hz, 2H), 7.03 (td, J = 8.5, 8.4, 2.9 Hz, 1H), 6.88 (d, J = 4.5 Hz, 1H), 6.80 (d, J = 4.5 Hz, 1H), 6.30 (s, 1H), 5.97 (s, 1H), 4.64 – 4.50 (m, 3H), 4.13 – 4.07 (m, 1H), 3.92 (t, J = 5.8, 5.8 Hz, 1H), 3.79 (dddd, J = 33.7, 12.0, 7.6, 4.3 Hz, 2H), 3.63 (dtt, J = 14.0, 10.1, 10.1, 5.7, 5.7 Hz, 3H), 3.43 (dd, J = 10.7, 3.4 Hz, 2H), 1.43 (p, J = 6.5, 6.5, 6.5, 6.5 Hz, 2H), 1.20 (d, J = 11.1 Hz, 30H), 0.83 (t, J = 6.9, 6.9 Hz, 3H). LC/MS purity 98.7%; [M+H]⁺ 806.8.

30 **15i** 1-*O*-Octadecyl-2-*O*-(4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn- May be prepared from Compound **14i** according to General Method E.

15j 1-*O*-Octadecyl-2-*O*-(3-fluoro-4-methoxy-benzyl)-*sn*-glyceryl-phospho-RVn-

Prepared from Compound **14j** according to General Method E. ¹H NMR (500 MHz,

DMSO-*d*₆) δ 8.43 (d, *J* = 6.7 Hz, 2H), 7.99 – 7.73 (m, 2H), 7.04 (d, *J* = 5.5 Hz, 1H), 6.87

(q, *J* = 3.7, 3.7, 3.2 Hz, 1H), 6.82 (dd, *J* = 7.3, 4.3 Hz, 1H), 6.17 – 5.74 (m, 1H), 4.59 (t, *J* = 4.9, 4.9 Hz, 1H), 4.53 – 4.41 (m, 1H), 4.11 (q, *J* = 4.9, 4.9, 4.9 Hz, 1H), 3.93 (q, *J* = 5.4, 5.4, 5.4 Hz, 1H), 3.84 (dq, *J* = 11.3, 6.5, 5.1, 5.1 Hz, 1H), 3.80 – 3.70 (m, 3H), 3.61 (ddd, *J* = 27.7, 10.9, 5.2 Hz, 3H), 3.30 (dd, *J* = 6.6, 3.0 Hz, 3H), 3.21 (dq, *J* = 9.8, 5.1, 5.1, 4.9 Hz, 1H), 1.43 (p, *J* = 6.6, 6.6, 6.5, 6.5 Hz, 2H), 1.27 – 1.16 (m, 30H), 0.83 (t, *J* = 6.8, 6.8 Hz, 3H). LC/MS purity 94.7%; [M+H]⁺ 836.8.

15k 1-*O*-Octadecyl-2-*O*-(pyridine-3-yl-methyl)-*sn*-glyceryl-phospho-RVn-

May be prepared from Compound **14k** according to General Method E.

15l 1-*O*-Oleyl-2-*O*-benzyl-*sn*-glyceryl-*phospho*-RVn Prepared from

Compound **14l** according to General Method D and isolated as an off-white solid in 84%

yield. ¹H NMR (300 MHz, CDCl₃ + CD₃OD) δ 7.79 (s, 1H), 7.38 (s, 2H), 7.27 – 7.21 (m, 5H), 6.92 (d, *J* = 6.0 Hz, 1H), 6.90 (d, *J* = 6.0 Hz, 1H), 5.30 (t, *J* = 6.0 Hz, 2H), 4.70 (d, *J* = 11 Hz, 1H), 4.62 (d, *J* = 5 Hz, 1H), 4.34 – 4.49 (m, 1H), 4.20 (m, 1H), 3.90 – 3.87 (m, 2H), 4.18 – 4.06 (m, 2H), 3.71 – 3.69 (m, 2H), 3.52 (ddd, *J* = 11.7, 3.1, 1.3 Hz, 1H), 3.35 (t, 2H), 1.96-1.93 (m, 4H), 1.49 – 1.47 (m, 2H), 1.33 – 1.23 (m, 20H), 0.83 (t, 2H). LC/MS purity 99%; [M+H]⁺ 786.78.

15m 1-*O*-Oleyl-2-*O*-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-*phospho*-RVn

Prepared from Compound **14m** according to General Method D and isolated as an off

white solid. Yield was 92%. ¹H NMR (300 MHz, CD₃OD) δ 7.74 (s, 1H), 7.32 (s, 2H), 6.99 (d, *J* = 6.0 Hz, 1H), 6.92 – 6.77 (m, 3H), 5.23 (t, *J* = 6.0 Hz, 2H), 4.67 (d, *J* = 11 Hz, 1H), 4.49 (d, *J* = 5 Hz, 2H), 4.30 (m, 1H), 4.20 (m, 1H), 3.83 – 3.81 (m, 2H), 3.75 (s, 3H), 3.63 (m, 1H), 3.31 (t, 2H), 1.91-1.89 (m, 4H), 1.44 (m, 2H), 1.18 (br s, 22H), 0.77 (t, 2H). LC/MS purity 99%; [M+H]⁺ 834.87.

15n 3-*O*-Octadecyl-2-*O*-benzyl-*sn*-glyceryl-1-*phospho*-RVn Prepared from


Compound **14n** according to General Method D and isolated as an off white solid.

15o 1-*O*-Octadecyl-2-*O*-(3-trifluoromethyl)benzyl-*sn*-glyceryl-*phospho*-RVn

Prepared from Compound **14o** according to General Method D and isolated as an off white solid. ESI-MS 854.48 [M-H]⁻.

15p 1-*O*-Octadecyl-2-*O*-(4-trifluoromethyl)benzyl-*sn*-glyceryl-*phospho*-RVn

Prepared from Compound **14p** according to General Method D and isolated as an off white solid. ESI-MS 854.46 [M-H]⁻.

5 D. Synthesis of 1-*O*-Octadecyl-2-*O*-benzyl-*sn*-glyceryl-benzyl-*phospho*-RVn

Scheme 4. Reagents: a) benzyl alcohol, PyBOP, DIEA, DMF; b) formic acid, rt


Compound **14d** (160 mg, 0.22 mmol), benzyl alcohol (48 mg, 0.45 mmol),
 10 diisopropylethylamine (DIEA, 58 mg, 0.45 mmol), and (1H-benzotriazol-1-yloxy)-
 tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 230 mg, 0.45 mmol) in dry
 DMF (5 mL) were stirred at room temperature for 3 h. DMF was then evaporated, and the
 residue was dissolved in ethyl acetate (50 mL) and washed with saturated NaHCO₃ (3 x 10
 mL). The organic layer was dried over MgSO₄ and concentrated. The residue was purified
 15 by column chromatography on silica gel, eluting with chloroform/methanol (0-15%) to
 yield **16**. ESI-MS 918.33 [M-H]⁻.

Compound **16** was added to formic acid and the deprotection was monitored by
 TLC. The mixture was concentrated under vacuum and the residue was purified by

column chromatography (silica gel, dichloromethane/methanol 10-20%) to give Compound **17**. Structure was confirmed by ESI-MS: 878.35 [M+H]⁺, 900.43 [M+Na]⁺.


E. Synthesis of GS-441524-3',5'-cyclic monophosphate, octadecyloxyethyl and 1-O-octadecyl-2-O-benzyl-sn-glycerol Esters

- In another embodiment, compounds of the disclosure are 3',5'-cyclic phosphates. The 3',5'-cyclic phosphates were prepared from known or readily prepared starting materials, following methods known to one skilled in the art of organic synthesis. As an example, 3',5'-cyclic phosphates **18-eq**, **18-ax**, **19-eq** and **19-ax** are prepared from phosphodiester **15d** and **4c** by an intramolecular esterification reaction:
- A solution of 1-O-octadecyl-2-O-benzyl-sn-glycerol-phospho-RVn (**15d**, 1 mmol) in dry DMF (5 mL) was added to a solution of diisopropylethylamine (DIEA, 58 mg, 0.45 mmol), and (1H-benzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 230 mg, 0.45 mmol) in dry DMF (5 mL) and stirred at room temperature for 3 hours until TLC indicated substantial conversion to the 3',5'-cyclic phosphate. Water (mL) was added and the solvents were evaporated under vacuum. The residue was purified by column chromatography on silica gel to yield compounds **18-eq** (ESI-MS: 770.18 [M+H]⁺), and **18-ax** (ESI-MS: 770.29 [M+H]⁺).
- Similarly, a solution of octadecyloxyethyl-phospho-RVn (**4c**, 1 mmol) in dry DMF (5 mL) was added to a solution of diisopropylethylamine (DIEA, 160 mg, 1.5 mmol), and (1H-benzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 780 mg, 1.5 mmol) in dry DMF (5 mL) and stirred at room temperature for 3 hours until TLC indicated substantial conversion to the 3',5'-cyclic phosphate. Water (1 mL) was added and the solvents were evaporated under vacuum. The residue was purified by column chromatography on silica gel to yield compounds **19-eq** (ESI-MS: 650.41 [M+H]⁺) and **19-ax** (ESI-MS: 650.42 [M+H]⁺).



5


F. Synthesis of 1-O-Octadecyl-2-O-(4-cyano)benzyl-sn-glyceryl-phospho-RVn. OD-4-CN-Bn-P-RVn was prepared using the Methods outlined above in Section C. 1-*O*-octadecyl-3-*O*-trityl-*sn*-glycerol was alkylated with 4-cyanobenzyl chloride and the trityl removed (Method I) to provide 1-*O*-octadecyl-2-*O*-(4-cyano)benzyl)-*sn*-glycerol. The alcohol was then phosphorylated (Method A), and coupled to GS-441524 acetoneide (Method C) and deprotected (Method E) to give the title compound.



1-*O*-octadecyl-2-*O*-(4-cyano)benzyl)-*sn*-glyceryl-phospho-RVn

15 *Example 2 – Assay for Anti-Coronavirus Activity in Vero E6 Cells*

In this example, reference is made to the following compounds, which include remdesivir nucleoside analogs and related intermediates:



The compounds of this example were assayed for anti-coronavirus activity in Vero E6 cells in comparison with remdesivir (RDV) and the remdesivir nucleoside (RVn). Ten thousand Vero E6 cells were seeded in 100 microliters of culture medium in 96 well plates. The following day serial two-fold dilutions of antiviral compounds or the DMSO-containing vehicle were added to each well. The USA WA-01 strain of SARS CoV-2 was added to each well at a multiplicity of infection of 0.1 thirty minutes later. Cells were incubated for 48 hours, washed twice in PBS and lysed with TRIzol. RNA was extracted using Directzol micro RNA columns. RNA was made into cDNA and assayed for the SARS CoV-2 spike protein and for a housekeeping gene (RPLPO) RNA by qPCR. Data represents the average of duplicate wells. Cellular cytotoxicity was also measured in VERO E6 cells. As below, each of the synthesized compounds exhibited enhanced anti-SARS CoV-2 activity compared to remdesivir or the remdesivir nucleoside with selectivity indices ranging from 22.8 to >227. Cytotoxicity (CC₅₀) was assessed using a commercially available MTT assay.

As shown in the following table, ODE-P-RVn (**4c**) and ODBG-P-RVn (**15d**) were 9 to 15 times more active against the USA WA-1 strain of SARS-CoV-2019 in Vero E6 cells. Likewise, HDP-P-RVn (**4b**) was 3.3-fold more active than remdesivir.

Antiviral activity in Vero E6 cells infected with SARS-CoV-2

Compound	EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)	Selectivity Index
RDV	3.16	5.50	>50	>15,8
ODE-P-RVn (4c)	0.35	0.53	>50	>143
ODBG-P-RVn (15d)	0.21	0.56	>50	>238
HDP-P-RVn (4b)	0.84	1.58	21	25.0
RVn	1.56	2.69	>50	>32.0

Abbreviations: EC₅₀, 50% effective concentration; EC₉₀, 90% effective concentration; CC₅₀, 50% cytotoxic concentration. Selective Index = CC₅₀/EC₅₀.

5 **Example 3. Additional Synthesis and Testing of RVn Monophosphate Prodrugs.**

Antiviral Activity: Also generated were concentration-response curves for ODBG-P-RVn (**15d**), ODE-P-RVn (**4c**), and HDP-P-RVn (**4b**), remdesivir (RDV) and remdesivir nucleoside (RVn) for SARS-CoV-2 infection in Vero E6 cells in two separate experiments performed in duplicate (**FIG. 1A-FIG. 1F**). Dose response curves for three
 10 remdesivir analogs (**FIG. 1A**, **FIG. 1B**, and **FIG. 1C**), remdesivir (GS-5734) (**FIG. 1D**), and remdesivir nucleoside (GS-441524) (**FIG. 1E**) against SARS-CoV-2 infection in Vero E6 cells. Vero E6 cells were pretreated with the indicated dose of the indicated drug for thirty minutes and then infected with SARS-CoV-2 isolate USA-WA1/2020 for 48 hours. The relative SARS-CoV-2 Spike RNA expression was determined by qRT-PCR. Each
 15 dose-response comparison was conducted simultaneously for all drugs on 2 separate occasions. Data from both experiments are shown at **FIG. 1A-FIG. 1F**. Data points indicate the mean relative expression from duplicate wells. Error bars represent the standard deviations (SDs). The black vertical dashed line indicates the concentrations at which there is 50% inhibition (EC₅₀). (**FIG. 1F**). Combined inhibition curves for all five
 20 compounds and DMSO on a single chart. DMSO, which was the vehicle for all compounds, had no effect on SARS-CoV-2 replication at the concentrations used. The three lipid esters of RVn-monophosphate were all substantially more active than RDV and RVn.

The following table shows the effective concentrations (EC₅₀, EC₉₀), 50%
 25 cytotoxic concentration (CC₅₀), and selectivity index of the compounds, mean ± SD. Cytotoxicity (CC₅₀) was assessed using Cell Titer Glo. The EC₅₀ values of RDV and RVn

were 4.6 and 1.7 μM , respectively. The lipid prodrugs were more active with EC_{50} s of ranging from 0.19 ± 0.023 to 0.96 ± 0.17 . ODBG-P-RVn and ODE-P-RVn were the most active and selective compounds. Based on the EC_{50} values the most active compound, ODBG-P-RVn, was 24 times more active than RDV and 8.9 times more active than RVn (p < 0.001 and 0.005) with a selectivity index of 240.

Antiviral activity, cytotoxicity and selectivity of the compounds in Vero E6 cells

Compound	EC_{50} (μM)	EC_{90} (μM)	CC_{50} (μM)	Selectivity	p value vs RDV, RVn
Remdesivir	4.6 ± 2.1	8.9 ± 4.9	>100	>21.7	-
Remdesivir nucleoside	1.7 ± 0.13	3.2 ± 0.77	>100	>58.8	-
HDP-P-RVn, 5a	0.96 ± 0.17	2.1 ± 0.78	51	52	0.02, 0.59
ODE-P-RVn, 5b	0.47 ± 0.18	1.1 ± 0.80	>100	>212	0.004, 0.047
ODBG-P-RVn, 5c	0.19 ± 0.023	0.56 ± 0.0002	46	240	<0.001, 0.005
A graph showing the CC_{50} results by Cell Titer Glo is shown in the Supplemental Materials. Abbreviations: RDV, Remdesivir (GS-5734); RVn, Remdesivir nucleoside (GS-441524); HDP-P-, hexadecyloxypropyl-P-; ODE-P-, octadecyloxyethyl-P-; ODBG-P-, 1-O-octadecyl-2-O-benzyl-glycero-3-P-; Selectivity index, $\text{CC}_{50}/\text{EC}_{50}$; statistical analysis comparing LogIC_{50} values from separate experiments by one-way ANOVA.					

Of all the perceived disadvantages of RDV, it was chosen in this example to design prodrugs of RVn which could provide oral bioavailability because an effective oral drug would allow for much earlier treatment of persons diagnosed with SARS-CoV-2 infection. As shown in this example, this was accomplished by constructing liponucleotides of RVn resembling lysophospholipids that are normally absorbed in the GI tract. The RVn liponucleotides were not metabolized rapidly in plasma and gain rapid entry to the cell often exhibiting greatly increased antiviral activity.

In contrast to the activation of RDV, which required four transformations, intracellular kinase bypass with this kind of compound generated the nucleoside monophosphate when the lipid ester moiety was cleaved in a single reaction catalyzed by acid phospholipase C or acid sphingomyelinase (sphingomyelin phosphodiesterase I).

One of the compounds, ODBG-P-RVn (**15d**) was likely to deliver relatively more drug to lung and less to liver as shown previously in lethal mousepox infection (Hostetler KY, Beadle JR, et al. Oral 1-O-octadecyl-2-O-benzyl-sn-glycero-3-cidofovir targets the lung and is effective against a lethal respiratory challenge with ectromelia virus in mice; Antiviral Res. 2007 Mar;73(3):212-8).

The synthesis of the lipid prodrugs of this example was much simpler than RDV and was readily scalable.

In this example, three lipid prodrugs of RVn were synthesized that were substantially more active than RDV or RVn in Vero E6 cells. The two most active compounds ODBG-P-RVn and ODE-P-RVn were 24 and 9.8 times more active than RDV. These compounds were expected to be orally bioavailable, stable in plasma and provide significant exposure and antiviral activity to all tissues infected with SARS-CoV-2.

Compounds: Remdesivir (GS-5734) and remdesivir nucleoside (GS-441524) were purchased from AA Blocks (San Diego, CA and Mason-Chem (Palo Alto, CA), respectively.

Cells: Vero E6 were obtained from ATCC and grown in DMEM (Corning) with 10 % FBS and Penicillin-Streptomycin (Gibco).

SARS-CoV-2 infection: SARS-CoV-2 isolate USA-WA1/2020 (BEI Resources) was propagated and infectious units quantified by plaque assay using Vero E6 (ATCC) cells. Approximately 10^4 Vero E6 cells per well were seeded in a 96 well plate and incubated overnight. Compounds or controls were added at the indicated concentrations 30 minutes prior to infection followed by the addition of SARS-CoV-2 at a multiplicity of infection equal to 0.01. After incubation for 48 hours at 37 °C and 5% CO₂, cells were washed twice with PBS and lysed in 200ul TRIzol (ThermoFisher).

RNA extraction, cDNA synthesis and qPCR: RNA was purified from TRIzol lysates using Direct-zol RNA Microprep kits (Zymo Research) according to manufacturer recommendations that included DNase treatment. RNA was converted to cDNA using the iScript cDNA synthesis kit (BioRad) and qPCR was performed using iTaq universal SYBR green supermix (BioRad) and an ABI 7300 real-time pcr system. cDNA was amplified using the following primers RPLP0 F – GTGTTCGACAATGGCAGCAT (SEQ ID NO: 1); RPLP0 R – GACACCCTCCAGGAAGCGA (SEQ ID NO: 2); SARS-

CoV-2 Spike F – CCTACTAAATTAAATGATCTCTGCTTTACT (SEQ ID NO. 3);
SARS-CoV-2 Spike R – CAAGCTATAACGCAGCCTGTA (SEQ ID NO. 4). Relative
expression of SARS-CoV-2 Spike RNA was calculated by delta-delta-Ct by first
normalizing to the housekeeping gene RPLP0 and then comparing to SARS-CoV-2
5 infected Vero E6 cells that were untreated (reference control). Curves were fit and 50
and 90% effective concentrations EC₅₀ and EC₉₀ values calculated using Prism 8.

CellTiter-glo luminescent cell viability assay: Approximately 10⁴ Vero E6 cells
per well were seeded in opaque walled 96 well cell culture plates and incubated overnight.
Compounds or controls were added at the indicated concentrations. After incubation for
10 48.5 hours at 37°C and 5% CO₂, an equal volume of CellTiter-Glo reagent (Cat. # G7570,
Promega, Madison, WI) was added, mixed and luminescence recorded on an EnSpire
Multimode Plate Reader (PerkinElmer) according to manufacturer recommendations.
Viability was calculated compared to untreated controls and CC₅₀ values were calculated
using Prism 8.

15 **Determination of Cytotoxicity:** The 50% cytotoxic concentrations (CC₅₀) were
determined with Cell Titer Glo (Cat. # G7570, Promega, Madison, WI) according to the
manufacturer's instructions. The calculated CC₅₀ values are shown in the foregoing table.

Vero E6 cells were treated with increasing concentrations of remdesivir analogs,
remdesivir (GS-5734), remdesivir nucleoside (GS441524) or DMSO vehicle (control) for
20 48.5 hrs. Relative viability was measured by CellTiter-Glo luminescent cell viability
assay, as depicted at **FIG. 2**.

Example 4 – Generation of Remdesivir triphosphate in Vero E6 cells

In this example, Vero E6 cells were plated in 6 well plates at about 3.4 x 10⁵ cells
25 per well in 2 mL of media (DMEM, 10% FBS).

Cells were then incubated at 37 °C for 24 hours. Media was then aspirated and
replaced with 2 mL of control media (fresh Dulbecco's Modified Eagle's Medium
(DMEM), 10% FBS) or 2 mL of media with drug at a concentration of 1 µM. Cell were
incubated with the various drugs for 48 hours. The media was aspirated, the cells rinsed
30 twice with phosphate-buffered saline (PBS), trypsinized for 5 minutes with 1 mL ATV,
triturated and removed to a 15 mL centrifuge tube, rinsed with 1 mL PBS, which was also
then removed to the 15 mL centrifuge tube, and the cells were trituated again. Cells were

counted in a Reichert hemocytometer using two 10 μ L samples, and the number of cells in each sample was determined.

Cells were centrifuged at 1200 rpm for 10 minutes, diluent aspirated and the pellet resuspended in 250 μ L of methanol/distilled water (70/30) and analyzed by LC/MS/MS.

- 5 The results, which are depicted at **FIG. 3**, were in picomoles/ 10^6 cells and were the average of two or three determinations. Abbreviations for **FIG. 3**: RDV, remdesivir; RVn, remdesivir nucleoside (GS-441524); ODE-P-RVn, octadecyloxyethyl-phospho-RVn (**4c**); ODBG-P-RVn, 1-O-octadecyl-2-O-benzyl-glycerol-3-phospho-RVn (**15d**)

- As depicted at **FIG. 3**, in Vero E6 cells, the synthesis of remdesivir triphosphate (RVn-TP) increased progressively to 48 hours with exposure to 1 micromolar ODE-P-RVn and ODBG-P-RVn. With RVn the levels of RVn-TP peaked at 8 hours and declined thereafter. Levels of RVn-TP with RDV were below the level of quantification at 8 and 24 hours.

Example 5 – Human Coronavirus 229E Infection

- 15 In this example, human Coronavirus 229E (ATCC) was propagated and infectious units quantified by TCID₅₀ using MRC-5 cells. For antiviral testing, approximately 104 MRC-5 cells were seeded per well in EMEM (10% FCS) at 37 °C in a 96 well plate overnight. Medium from each well was removed and cells were infected with 100 TCID₅₀ virus in 100 μ L medium for two hours.

- 20 Cells were washed one time with medium and then compounds or controls added at the indicated concentrations. After three days, CPE was observed under microscope and quantified using an MTT cell proliferation assay kit (Abcam) read on an ELx800, Universal Microplate reader (BIO-TEK Instruments, INC).

25

Effect of Compounds on HCoV-229E Replication in MRC-5 Cells

Entry	Compd	R ₁	R ₂	HCoV-229E in MRC-5 Cells		
				EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)
1	4a	eicosyl	H	±0.91 ± 1.21	±1.80 ± 1.87	±>50
2	4b	hexadecyloxypropyl	H	3.02 ± 0.36	6.60 ± 1.14	32.1 ± 17.9
3	4c	octadecyloxyethyl	H	0.41 ± 0.012	0.84 ± 0.095	>50
4	6c	octadecyloxyethyl	benzyl	0.22 ± 0.056	0.44 ± 0.085	>50
5	15a	1- <i>O</i> -tetradecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.76 ± 0.34	±3.25 ± 2.18	±>50
6	15b	1- <i>O</i> -hexadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.56 ± 0.27	0.98 ± 0.38	>50
7	15c	1- <i>O</i> -hexadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.36 ± 0.054	1.12 ± 0.10	>50
8	15d	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.10 ± 0	0.28 ± 0.013	43.1 ± 18.9
9	15e	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>rac</i> -glyceryl	H	0.21 ± 0.004	0.35 ± 0.006	30.9 ± 7.7
10	15f	1- <i>O</i> -octadecyl-2- <i>O</i> -octyl- <i>sn</i> -glyceryl	H	0.12 ± 0.057	0.29 ± 0.058	±>50
11	15g	1- <i>O</i> -octadecyl-2- <i>O</i> -(methylcyclohexyl)- <i>sn</i> -glyceryl	H	nd	nd	nd
12	15h	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F-Bn)- <i>sn</i> -glyceryl	H	1.40 ± 0.058	2.80 ± 0.04	22.9 ± 0.1
13	15i	1- <i>O</i> -octadecyl-2- <i>O</i> -(4-MeO-Bn)- <i>sn</i> -glyceryl	H	nd	nd	nd
14	15j	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.074 ± 0.01	0.17 ± 0.014	>50
15	15k	1- <i>O</i> -octadecyl-2- <i>O</i> -methylpyridinyl- <i>sn</i> -glyceryl	H	nd	nd	nd
16	15l	1- <i>O</i> -oleyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.13 ± 0.029	0.26 ± 0.0037	>50
17	15m	1- <i>O</i> -oleyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.06 ± 0.014	0.11 ± 0.02	>50
18	17	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	benzyl	1.50 ± 0.93	4.27 ± 0.88	>50
19	15o	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-CF ₃ -Bn)- <i>sn</i> -glyceryl	H	0.16±0.046	0.46±0.27	16.0±2.8
20	15p	1- <i>O</i> -octadecyl-2- <i>O</i> -(4-CF ₃ -Bn)- <i>sn</i> -glyceryl	H	0.16±0.078	0.33±0.039	13.9±8.8

nd = not determined;

The % inhibition was calculated as $(Atv - Acv)/(Acd - Acv) \times 100\%$ where Atv indicates the absorbance of the test compounds with virus infected cells and Acv and Acd indicate the absorbance of the virus control and the absorbance of the cell control, respectively. The average half-maximal effective concentration (EC₅₀) was defined as the concentration which achieved 50% inhibition of virus-induced cytopathic effects.

Example 6 – SARS-CoV-2 Infection Assay

About 12e3 TMPRSS2-Vero cells or 20e3 Huh7.5 cells were seeded per well in black with clear flat bottom 96 well plates and incubated overnight. Compounds or controls were added about 30 to about 60 minutes prior to infection at the indicated concentrations with addition of SARS-CoV-2 at a multiplicity of infection (FFU/cell) equal to 0.01 for TMPRSS2-Vero and 0.1 for Huh7.5.


After incubation for 32 hours for TMPRSS2-Vero or 48 hours for Huh7.5 at 37 °C and 5% CO₂, the medium was removed and cells were incubated in 4 % formaldehyde for 30 minutes at room temperature. Formaldehyde fixed cells were washed with PBS and permeabilized for immunofluorescence in 0.1% Triton-X 100 in PBS with 1% bovine serum albumin (BSA) fraction V (Millipore-Sigma) and stained for SARS-CoV-2 with a primary anti-Nucleocapsid antibody (GeneTex GTX135357) followed by AlexaFluor 594 secondary antibody (Thermo Fisher Scientific A-11012) with nuclear counterstain Sytox Green (Thermo Fisher Scientific).

Five images per well were obtained at 10x magnification using an Incucyte S3 (Sartorius). The percent infected cells and nuclei count were calculated using built-in image analysis tools for the Incucyte S3. Calculations for EC₅₀, EC₉₀ and CC₅₀ were carried out using the nonlinear regression analysis in GraphPad Prism 9 with the bottom and top parameters constrained to 0 and 100, respectively.

Effect of Compounds on SARS CoV-2 Replication in Huh7.5 and TMPRSS2-Vero Cells

Entry	Compd	R ₁	R ₂	Huh7.5 Cells			TMPRSS2-Vero Cells		
				EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)	EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)
1	4a	eicosyl	H	1.411 ± 0.089	4.746 ± 0.626	>20	1.171 ± 0.060	2.185 ± 0.201	>20
2	4b	hexadecyloxypropyl	H	0.19 ^a	0.40 ^a	>20	nd	nd	nd
3	4c	octadecyloxyethyl	H	0.19 ^a	0.37 ^a	>20	nd	nd	nd
4	6c	octadecyloxyethyl	benzyl	nd	nd	nd	nd	nd	nd
5	15a	1- <i>O</i> -tetradecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.156 ± 0.008	0.493 ± 0.051	>20	0.901 ± 0.022	1.694 ± 0.131	>20
6	15b	1- <i>O</i> -hexadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.130 ± 0.020	0.309 ± 0.092	>20	0.395 ± 0.021	0.810 ± 0.079	>20
7	15c	1- <i>O</i> -hexadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.049 ± 0.005	0.113 ± 0.019	>20	0.397 ± 0.026	0.869 ± 0.106	>20
8	15d	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.138 ± 0.018	0.377 ± 0.098	>20	0.205 ± 0.016	0.432 ± 0.075	>20
9	15e	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>rac</i> -glyceryl	H	0.163 ± 0.026	0.514 ± 0.172	>20	0.242 ± 0.020	0.642 ± 0.133	>20
10	15f	1- <i>O</i> -octadecyl-2- <i>O</i> -octyl- <i>sn</i> -glyceryl	H	0.710 ± 0.028	1.713 ± 0.183	>20	1.326 ± 0.052	2.349 ± 0.123	>20
11	15g	1- <i>O</i> -octadecyl-2- <i>O</i> -(methylcyclohexyl)- <i>sn</i> -glyceryl	H	nd	nd	nd	nd	nd	nd
12	15h	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F-Bn)- <i>sn</i> -glyceryl	H	0.182 ± 0.018	0.444 ± 0.083	>20	0.270 ± 0.016	0.629 ± 0.106	>20
13	15i	1- <i>O</i> -octadecyl-2- <i>O</i> -(4-MeO-Bn)- <i>sn</i> -glyceryl	H	nd	nd	nd	nd	nd	nd
14	15j	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.056 ± 0.008	0.222 ± 0.070	>20	0.174 ± 0.016	0.341 ± 0.044	>20
15	15k	1- <i>O</i> -octadecyl-2- <i>O</i> -methylpyridinyl- <i>sn</i> -glyceryl	H	nd	nd	nd	nd	nd	nd
16	15l	1- <i>O</i> -oleyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.100 ± 0.010	0.246 ± 0.062	>20	0.295 ± 0.013	0.568 ± 0.080	>20
17	15m	1- <i>O</i> -oleyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.054 ± 0.006	0.102 ± 0.013	>20	0.296 ± 0.019	0.524 ± 0.115	>20
18	17	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	benzyl	3.537 ± 0.583	16.44 ± 5.98	>20	7.443 ± 0.818	15.263 ± 5.912	>20

nd = not determined; ^aData from Example 7



5 Effect of Compounds on SARS CoV-2 Replication in Calu-3 cells

Entry	Compd	R ₁	R ₂	Calu-3 Cells		
				EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)
1	4a	eicosyl	H	0.178 ± 0.035	1.316 ± 0.566	>20
2	4b	hexadecyloxypropyl	H	nd	nd	nd
3	4c	octadecyloxyethyl	H	nd	nd	nd
4	6c	octadecyloxyethyl	benzyl	nd	nd	nd
5	15a	1- <i>O</i> -tetradecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.212 ± 0.025	0.536 ± 0.150	>20
6	15b	1- <i>O</i> -hexadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.101 ± 0.020	0.623 ± 0.272	88.23 ± 8.89
7	15c	1- <i>O</i> -hexadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.051 ± 0.006	0.221 ± 0.058	75.63 ± 3.96
8	15d	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.164 ± 0.020	0.536 ± 0.138	>20
9	15e	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>rac</i> -glyceryl	H	0.210 ± 0.062	0.763 ± 0.490	>20
10	15f	1- <i>O</i> -octadecyl-2- <i>O</i> -octyl- <i>sn</i> -glyceryl	H	0.410 ± 0.060	2.314 ± 0.738	>20
11	15g	1- <i>O</i> -octadecyl-2- <i>O</i> -(methylcyclohexyl)- <i>sn</i> -glyceryl	H	0.147 ± 0.032	1.660 ± 0.798	>20
12	15h	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F-Bn)- <i>sn</i> -glyceryl	H	0.216 ± 0.062	0.548 ± 0.368	>20
13	15i	1- <i>O</i> -octadecyl-2- <i>O</i> -(4-MeO-Bn)- <i>sn</i> -glyceryl	H	nd	nd	nd
14	15j	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.080 ± 8.741	0.101 ± 4.554	50.63 ± 5.42
15	15k	1- <i>O</i> -octadecyl-2- <i>O</i> -methylpyridinyl- <i>sn</i> -glyceryl	H	nd	nd	nd

16	15l	1- <i>O</i> -oleyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.198 ± 0.077	0.294 ± 0.093	78.96 ± 6.95
17	15m	1- <i>O</i> -oleyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.075 ± 4.752	0.090 ± 3.487	58.53 ± 6.28
18	17	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	benzyl	5.320 ± 1.393	10.92 ± 5.485	>20
19	18-eq	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	cyclic-eq	>20	>20	>20
20	18-ax	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	cyclic-ax	>20	>20	>20
21	19-eq	octadecyloxyethyl	cyclic-eq	2.758 ± 0.808	19.163 ± 12.666	>20
22	19-ax	octadecyloxyethyl	cyclic-ax	8.147 ± 8.420	>20	>20
23	15n	3- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.163 ± 0.022	0.623 ± 0.183	>20
24	15o	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-trifluoromethyl)benzyl- <i>sn</i> -glyceryl	H	0.166 ± 0.068	0.278 ± 0.109	>20

SARS-CoV-2 infection assay: Approximately 20k Calu-3 cells were seeded per well in black with clear flat bottom 96 well plates (Corning #3904) and incubated 48-72h. Compounds or controls were added 30-60 minutes prior to infection at the indicated concentrations with addition of SARS-CoV-2 at a multiplicity of infection (FFU/cell) equal to 0.01. After incubation for 44 hour at 37 °C and 5% CO₂, the medium was removed and cells were incubated in 4% formaldehyde for 30 minutes at room temperature. Formaldehyde fixed cells were washed with PBS and permeabilized for immunofluorescence in 0.1% Triton-X 100 in PBS with 1% bovine serum albumin (BSA) fraction V (Millipore-Sigma) and stained for SARS-CoV-2 with a primary anti-Nucleocapsid antibody (GeneTex GTX135357) followed by AlexaFluor 594 secondary antibody (Thermo Fisher Scientific A-11012) with nuclear counterstain Sytox Green (Thermo Fisher Scientific). Five images per well were obtained at 10x magnification using an Incucyte S3 (Sartorius).

The percent infected cells and nuclei count were calculated using built-in image analysis tools for the Incucyte S3. Calculations for EC₅₀, EC₉₀ and CC₅₀ were carried out using the nonlinear regression analysis in GraphPad Prism 9 with the bottom and top parameters constrained to 0 and 100, respectively. All work with authentic SARS-CoV-2 was conducted in Biosafety Level-3 conditions at the University of California San Diego.

Cell viability assay: For select compounds the CC₅₀ was calculated up to a maximum of 100µM using CellTiter-Glo. For these experiments, approximately 20k Calu-3 cells were seeded per well in opaque white 96-well plates (cat# 655073, Greiner Bio-One, Monroe, North Carolina) and incubated 48-72h. Compounds or controls were added and incubated for 44h at 37 °C and 5%

CO₂. After which an equal volume of CellTiter-Glo reagent (Cat. # G7570, Promega, Madison, WI) was added, mixed and luminescence recorded on a Veritas Microplate Luminometer (Turner BioSystems) according to manufacturer recommendations.

5 *Example 7 – Antiviral activity in various cell types infected with SARS-CoV-2*

Vero E6, Caco-2, and Calu-3 cell lines were obtained from ATCC. Huh7.5 cells were obtained from Apath LLC. Calu-3 and Caco-2 cells were propagated in MEM (Corning), 10% FBS, Penicillin-Streptomycin (Gibco). Vero E6 and Huh7.5 cells were propagated in DMEM (Corning) with 10% FBS and Penicillin-Streptomycin (Gibco).

10 Human PSC-lung cell generation, human lung organoids were generated as previously described (Leibel SL, McVicar RN, Winquist AM, Niles WD, Snyder EY Generation of complete multi-cell type lung organoids from human embryonic and patient-specific induced pluripotent stem cells for infectious disease modeling and therapeutics validation Curr. Protoc. Stem Cell Biol., 54 (1) (2020 Sep), Article e118). H9 embryonic stem cells
15 (WiCell) were cultured in feeder free conditions upon Matrigel (Corning #354230) coated plates in mTeSR medium (StemCellTech #85850). Media was changed daily, and stem cells were passaged using enzyme free dissociation reagent ReLeSR™ (Stem Cell Tech#05872). Cultures were maintained in an undifferentiated state, in a 5% CO₂ incubator at 37 °C.

20 For proximal lung organoid generation, human PSCs were dissociated into single cells, and then seeded on Matrigel-coated plates (BD Biosciences) at a density of 5.3×10^4 cells/cm² in Definitive Endoderm (DE) induction medium (RPMI1640, 2% B27 supplement, 1% HEPES, 1% glutamax, 50 U/mL penicillin/streptomycin), supplemented with 100 ng/mL human activin A (R&D), 5 μM CHIR99021 (Stemgent), and 10 μM
25 ROCK inhibitor, Y-27632 (R&D Systems) on day 1. On days 2 and 3 cells were cultured in DE induction media with only 100 ng/mL human activin A. Anterior Foregut Endoderm (AFE) was generated by supplementing serum free basal medium (3 parts IMDM:1 part F12, B27+N2 supplements, 50U/mL penicillin/streptomycin, 0.25% BSA, 0.05 mg/mL L-ascorbic acid, 0.4mM monothioglycerol) with 10 μM SB431542 (R&D) and 2 μM
30 Dorsomorphin (StemGent) on days 4-6. On day 7, AFE medium was changed to Lung Progenitor Cell (LPC) induction medium, containing serum free basal medium supplemented with 10 ng/mL human recombinant BMP4 (R&D), 0.1 μM all-trans retinoic acid (Sigma-Aldrich) and 3 μM CHIR99021. Media was changed every other day for 9-11

days. To generate 3D human proximal lung organoids, we modified a previously published protocol (K.B. McCauley , F. Hawkins , M. Serra , D.C. Thomas , A. Jacob , and D.N. Kotton. (2017) Efficient Derivation of Functional Human Airway Epithelium from Pluripotent Stem Cells via Temporal Regulation of Wnt Signaling. *Cell Stem Cell*; 20(6):844-857)

LPCs were dissociated in accutase for 10minutes and resuspended in Matrigel in a 12-well, 0.4µm pore size Transwell (Corning) culture insert at 5.0×10^4 cells/200ul of Matrigel. Cells were cultured in proximal lung organoid maturation media using serum free basal medium supplemented with 250ng/mL FGF2, 100ng/mL rhFGF10, 50nM dexamethasone (Dex), 100µM 8-Bromoadenosine 3',5'-cyclic monophosphate sodium salt (Br-cAMP), 100µM 3-Isobutyl-1-methylxanthine (IBMX) and 10 µM ROCK inhibitor (Y-27632). Proximal lung organoid media was changed every other day for 3 weeks. Human PSC-derived lung organoids were dissociated into single cells and seeded at 20,000 cells per well of a matrigel coated 96-well plate one day before transfection. Transwells containing the proximal organoids in matrigel were incubated in 2U/ml dispase for 30 minutes at 37 °C. Cold PBS was added to the mixture then centrifuged at 400 x g for 5 minutes.

Supernatant was carefully removed and resuspended in 2-3mls of TrypLE Express (Gibco # 12605010) for 20 minutes at 37 °C. Reaction was quenched with 2% FBS in DMEM/F12 then centrifuged at 400 x g for 5 min. The supernatant was aspirated, and the cell pellet resuspended in 1ml of quenching media supplemented with 10 µM Rock inhibitor (Y-27632). Cell count was performed and the respective volume of cells were transferred into a reagent reservoir trough and resuspended in proximal lung organoid maturation media and plated via multichannel pipette into 96 well plates at 100ul per well as monolayers.

SARS-CoV-2 infection: SARS-CoV-2 isolate USA-WA1/2020 (BEI Resources) was propagated and infectious units quantified by plaque assay using Vero E6 (ATCC) cells. Approximately 12,000 cells from each cell line were seeded per well in a 96 well plate. Vero E6 and Huh7.5 were seeded approximately 24 hours prior to treatment/infection. Calu-3 and Caco-2 were seeded approximately 48h prior to treatment/infection. Human PSC lung cell infections and cytotoxicity experiments were performed when cells reached 100% confluency. Compounds or controls were added at the indicated concentrations 30 minutes prior to infection followed by the addition of

SARS-CoV-2 at a multiplicity of infection equal to 0.01. After incubation for 48 hours at 37°C and 5% CO₂, cells were washed twice with PBS and lysed in 200ul TRIzol (ThermoFisher). All work with SARS-CoV-2 was conducted in Biosafety Level 3 conditions at the University of California San Diego with approval from the Institutional Biosafety Committee.

RNA extraction, cDNA synthesis and qPCR: RNA was purified from TRIzol lysates using Direct-zol RNA Microprep kits (Zymo Research) according to manufacturer recommendations that included Dnase treatment. RNA was converted to cDNA using the iScript cDNA synthesis kit (BioRad) and qPCR was performed using iTaq universal SYBR green supermix (BioRad) and an ABI 7300 real-time pcr system. cDNA was amplified using the following primers RPLP0 F – GTGTTCGACAATGGCAGCAT (SEQ ID NO: 1); RPLP0 R – GACACCCTCCAGGAAGCGA (SEQ ID NO: 2); SARS-CoV-2 Spike F – CCTACTAAATTAATGATCTCTGCTTTACT (SEQ ID NO: 3); SARS-CoV-2 Spike R – CAAGCTATAACGCAGCCTGTA (SEQ ID NO: 4). Relative expression of SARS-CoV-2 Spike RNA was calculated by delta-delta-Ct by first normalizing to the housekeeping gene RPLP0 and then comparing to SARS-CoV-2 infected Vero E6 cells that were untreated (reference control). Curves were fit using the nonlinear regression – log(inhibitor) vs. response (four parameter) model using Prism 9. To calculate effective concentrations EC₅₀ and EC₉₀ values, qRT-PCR values were normalized to percent inhibition and curves fit using the nonlinear regression – log(agonist) vs. response (four parameter) model with bottom and top constrained to 0 and 100 respectively using Prism 9.

Cell viability assay: Cell type were seeded as per SARS-CoV-2 infection studies in opaque walled 96-well cell culture plates or 229E infection studies in clear 96-well cell culture plates and incubated overnight. Compounds or controls were added at the indicated concentrations. For SARS-CoV-2 related studies, cells were incubated for 48.5 hours at 37°C and 5% CO₂, an equal volume of CellTiter-Glo reagent (Cat. # G7570, Promega, Madison, WI) was added, mixed and luminescence recorded on a Veritas Microplate Luminometer (Turner BioSystems) according to manufacturer recommendations. For 229E related, cells were incubated for 72 hours at 37°C and 5% CO₂, supernatants removed, 50µL of serum-free media and 50µL of MTT Reagent (Abcam ab211091) added to each well and incubated for 3hrs at 37°C. Absorbance was measured on an ELx800,

Universal Microplate reader, (BIO-TEK Instruments, INC) according to manufacturer recommendations. Percent viability was calculated compared to untreated controls and CC₅₀ values were calculated using Prism 9.

5 Antiviral Activity, Cytotoxicity and Selectivity of the Compounds in five cell types

Compound	EC ₅₀ (μ M)	EC ₉₀ (μ M)	CC ₅₀ (μ M)	Selectivity	<i>p</i> value EC ₅₀ vs RDV, RVn
Vero E6 cells					
RDV	1.13	7.05	101	89.4	-
RVn	0.38	0.77	>100	>263	-
HDP-P-RVn, 4b	0.63	0.73	>100	>158	NS
ODE-P-RVn, 4c	0.30	0.33	>100	>333	NS
ODBG-P-RVn, 15d	0.14	0.16	97.9	699	0.010, 0.311
PSC-human lung cells					
RDV	0.14	0.23	32.7 *	234	-
RVn	0.74	2.62	>100 *	>135	-
HDP-P-RVn *	0.35	0.94	ND		-
ODE-P-RVn	0.22	0.70	>100 *	>454	0.791, 0.006
ODBG-P-RVn	0.15	0.26	61.5 *	410	>0.999, 0.002
Calu-3 cells					
RDV	0.23	0.31	>100	>434	-
RVn	0.15	0.18	>100	>666	-
ODE-P-RVn	0.34	0.64	98.7	290	NS
ODBG-P-RVn	0.30	0.33	98.2	327	NS
Huh7.5 cells					
RDV	0.06	0.12	15.2	253	-
RVn	0.32	0.73	>100	>312	-
HDP-P-RVn	0.19	0.40	>100	>526	NS
ODE-P-RVn	0.19	0.37	>100	>526	NS
ODBG-P-RVn	0.14	0.15	62.9	449	NS
Caco-2 cells					
RDV	0.17	0.28	>100	>588	-
RVn	0.96	1.75	>100	>104	-
ODE-P-RVn	0.77	1.25	>100	>129	0.007, 0.971
ODBG-P-RVn	0.30	0.33	88.4	295	0.968, 0.007

Abbreviations: RDV, Remdesivir (GS-5734); RVn, Remdesivir nucleoside (GS-441524); HDP-P-, hexadecyloxypropyl-P-; ODE-P-, octadecyloxyethyl-P-; ODBG-P-, 1-O-octadecyl-2-O-benzyl-glycero-3-P-; EC₅₀: half-maximal effective concentration; CC₅₀: 50% cytotoxic concentration, Selectivity index, CC₅₀/EC₅₀; statistical analysis comparing LogEC₅₀ values from separate experiments by one-way ANOVA. CC₅₀ results by CellTiter-Glo. All experiments performed three times in duplicate except starred (*) were done twice in duplicate.

In all cell lines, there was a dose-dependent inhibition of viral RNA by ODBG-P-RVn, ODE-P-RVn, HDP-P-RVn, remdesivir (RDV) and remdesivir nucleoside (RVn). In

Vero E6 cells, the average half-maximal effective concentration (EC_{50}) and average 90% effective concentration (EC_{90}) of ODBG-P-RVn was 0.14 μ M and 0.16 μ M, respectively. The EC_{50} of ODBG-P-RVn in Vero E6 cells was significantly lower than RDV. ODE-P-RVn and HDP-P-RVn were also potently antiviral with EC_{50} values of 0.3 μ M and 0.63 μ M in Vero E6. The EC_{50} of ODBG-P-RVn and ODE-P-RVn were less than 0.35 μ M in PSC-lung and Calu-3, both models of human lung infection. The antiviral activities of ODBG-P-RVn and ODE-P-RVn were significantly better than RVn in PSC-lung cells. ODBG-P-RVn, ODE-P-RVn and HDP-P-RVn demonstrated strong antiviral activity in Huh7.5 cells with EC_{50} less than 0.2 μ M that was not significantly different from RDV or RVn. In the Caco-2 cell line, the EC_{50} of ODBG-P-RVn was 0.3 μ M which was significantly lower than RVn but similar to RDV. In the same cell line, the EC_{50} of ODE-P-RVn was 0.77 μ M, which was significantly higher than RDV.

The cytotoxicity of each compound by incubating each of these cell lines with serial dilutions of each compound from 1.23 μ M to 100 μ M for 48 hours. The average 50% cytotoxic concentrations (CC_{50}) for all compounds were greater than 60 μ M in all cell lines except for RDV which had a CC_{50} of 32.7 μ M in PSC-lung cells and 15.2 μ M in Huh7.5, a human hepatocyte cell line. The selectivity index of ODBG-P-RV ranged from 295 to 699 in the five cell types tested in this example). The range of antiviral activity and cytotoxicity of ODBG-P-RVn (EC_{50} 0.14 μ M – 0.30 μ M and CC_{50} 61.5 μ M – 98.2 μ M) was more consistent across cell types than RDV (EC_{50} 0.06 μ M – 1.13 μ M and CC_{50} 15.2 μ M – >100 μ M) (see table above). Collectively, these data demonstrate that lipid RVn monophosphate prodrugs are potent antivirals against SARS-CoV-2 *in vitro* with low toxicity and excellent selectivity indexes.

25

Example 8 - Effect of Antivirals in Human Coronavirus 229E infected cells

Human Coronavirus 229E (ATCC) was propagated and infectious units quantified by TCID₅₀ using MRC-5 cells. For antiviral testing, approximately 10⁴ MRC-5 cells were seeded per well in EMEM (10%FCS) at 37C in a 96 well plate overnight. Medium from
5 each well was removed and cells were infected with 100 TCID₅₀ virus in 100 µL medium for two hours.

Cells were washed one time with medium and then compounds or controls added at the indicated concentrations. After three days, CPE was observed under microscope and quantified using an MTT cell proliferation assay kit (Abcam) read on an ELx800,
10 Universal Microplate reader (BIO-TEK Instruments, INC). The % Inhibition was calculated as $(A_{iv} - A_{cv}) / (A_{cd} - A_{cv}) \times 100\%$ where A_{iv} indicates the absorbance of the test compounds with virus infected cells and A_{cv} and A_{cd} indicate the absorbance of the virus control and the absorbance of the cell control, respectively. The average half-maximal effective concentration (EC₅₀) was defined as the concentration which achieved 50%
15 inhibition of virus-induced cytopathic effects.

As described herein, various compounds of the disclosure inhibit the human *Alphacoronavirus* 229E. Cells were infected with 229E for 2 hours followed by treatment with the indicated dose of the indicated drug for 72 hours. The relative CPE was determined by measuring cell viability using an MTT assay. depicts the cytotoxicity in
20 MRC-5 cells incubated in the presence of the indicated drug at the indicated concentration for 72 hours, after which cell viability was measured by the CellTiter-Glo assay. Data points indicate the averages from 3 independent experiments performed in duplicate. Error bars represent the standard error mean (SEM). Both ODBG-P-RVn and RDV demonstrated a dose-dependent inhibition of cytopathic effect (CPE). The EC₅₀ values of
25 ODBG-P-RVn and RDV were 0.15µM and 0.04µM and the EC_{90s} were 0.54 mM and 0.26 mM respectively. The CC₅₀ for ODBG-P-RVn and RDV were greater than 50µM in MRC-5 cells. The most active compounds were compounds which had 3-fluoro,4-methoxy substitutions. Together with the antiviral data for SARS-CoV-2, this demonstrates that ODBG-P-RVn related analogs have antiviral activity against two
30 genetically distinct human pathogenic coronaviruses.

Example 9 - Orally administered ODBG-P-RVn (15d) achieved therapeutic plasma levels in Syrian Hamsters

ODBG-P-RVn in 0.1M sodium carbonate/bicarbonate buffer, pH 9.0, was administered to Syrian Hamsters by oral gavage every 12 hours for seven days. ODBG-P-RVn was present as the sodium salt. It was well tolerated, and no adverse clinical signs were noted. Peak plasma levels of ODBG-P-RVn were noted at 1 hour and fell by 50% in about 5 hours.

Plasma curves were generally similar at day 1 and 7 except at 16.9 mg/kg, the 7 d values were slightly higher than the levels at day 1. At 12 hours ODBG-P-RVn levels were above the EC₉₀ for ODBG-P-RVn in all cell lines studied including Vero E6 cells and PSC lung cells on both day 1 and 7. Levels of the RVn, the nucleoside metabolite of ODBG-P-RVn peaked at 3 hours after administration and declined thereafter. Plasma levels of RVn were less than the EC₉₀ for RVn in both PSC lung cells and Vero E6 cells. The observed low levels of RVn suggest that antiviral activity attributable to this metabolite will be minimal and are also consistent with finding of ODBG-P-RVn stability in human plasma. Collectively, these results suggest that ODBG-P-RVn will be effective in suppressing viral replication in a variety of tissue types *in vivo*.

FIG. 4A and FIG. 4B depict the seven day oral pharmacokinetics in Syrian hamsters. Syrian hamsters were given vehicle or ODBG-P-RVn by oral gavage every 12 hours for 7 days. Groups of 3 animals received vehicle or drug at doses of 16.9 and 13.2 mg/kg. Animals were weighed daily and monitored for clinical signs. Plasma samples were obtained at 1, 3, 6 and 12 hours on day 1 and day 7 and frozen for analysis of ODBG-P-RVn (**FIG. 4A**) and RVn (**FIG. 4B**) by LC/MS/MS.

Analytical Methods: ODBG-P-RVn: Hamster plasma samples (10 µL) containing ODBG-P-RVn and K₂EDTA as the anticoagulant were added to polypropylene tubes containing water (100 µL), internal standard solution (10 µL; 1,000 ng/mL of ODE-P-RVn in ACN:DMF (1:1, v/v)), and 10 µL of ACN:DMF (1:1, v/v). The solutions were mixed, then acidified with phosphoric acid, 85% w/v:water (1:19, v/v; 10 µL), mixed, then diluted with 200 µL of IPA, mixed, then diluted with 500 µL of water, and mixed. The samples were extracted with a Sep-Pak® tC18 96-well solid phase extraction plate (25 mg; Waters, Milford, MA). Extraction occurred under positive pressure conditions using nitrogen. Samples were washed serially with 1 mL of water:acetonitrile:formic acid (475:25:0.5, v/v/v) and 0.4 mL of water:acetonitrile:formic acid (350:150:0.5, v/v/v)

before being serially eluted with 100 μ L and 150 μ L of water: {acetonitrile:isopropyl alcohol (1:1, v/v)}:formic acid:ammonium formate:citric acid solution, 2% w/v (15:85:0.1:0.1:0.1, v/v/v/w/v). The citric acid solution was prepared as water: citric acid monohydrate (20:0.4, v/w). After elution, 100 μ L of water was added to each sample. The

5 ODBG-P-RVn extracts were analyzed using an Agilent 1200 HPLC system (Agilent, Santa Clara, CA) coupled to an API5500 mass analyzer (SCIEX, Foster City, CA). Analytes were chromatographically separated using a Dacapo DX-C18 MF column (100 x 2 mm, 2.5 μ m; ImtaktUSA, Portland, OR) using a mobile phase system consisting of Mobile Phase A (water: formic acid: [water: ammonium formate: citric acid (25:5:0.5, v/w/w)] (1,000:1:1, v/v/v) and Mobile Phase B (acetonitrile: isopropyl alcohol: formic acid: [water: ammonium formate: citric acid (25:5:0.5, v/w/w)] (800:200:1:1, v/v/v/v). The

10 total analytical run time was 4.5 minutes. The mobile phase was nebulized using heated nitrogen in a Turbo-V source/interface set to electrospray positive ionization mode. The ionized compounds were detected using multiple reaction monitoring with transitions m/z

15 788.4 > 229 (V2043) and 668.4 > 467.2 (V2041). This method is applicable for measuring ODBG-P-RVn concentrations ranging from 6.25 to 3,000 ng/mL using 10.0 μ L of plasma for extraction. The peak areas of ODBG-P-RVn and RVn were acquired using Analyst v. 1.6.2 (SCIEX, Framingham, MA). The calibration curve was obtained by fitting the peak area ratios of the analyte/I.S. and the standard concentrations to a linear equation with 1/x²

20 weighting, using Analyst. The equation of the calibration curve was then used to interpolate the concentrations of the analyte in the samples using their peak area ratios. The peak areas used for the calculations were not rounded.

Analytical Methods: RVn (GS-441524): Hamster plasma samples (20 μ L) containing GS-441524 and K₂EDTA as the anticoagulant were added to Eppendorf

25 LoBind microfuge tubes containing acetonitrile (300 μ L) and water: acetonitrile (2:8, v/v; 60 μ L). The solutions were mixed and centrifuged at 16,000 g for five minutes. The supernatant (300 μ L) was then filtered through an Ostro protein precipitation and phospholipid removal plate (25 mg; Waters, Milford, MA). Filtration occurred under positive pressure conditions using nitrogen. Collected filtered samples were capped,

30 mixed and stored at 10°C pending analysis. The GS-441524 extracts were analyzed using an Acquity UPLC system (Waters, Milford, MA) coupled to a G2-S QToF mass analyzer (Waters, Milford, MA). Analytes were chromatographically separated using a Unison-UK Amino HT column (100 x 2 mm, 3 μ m; ImtaktUSA, Portland, OR) using a mobile phase

system consisting of Mobile Phase A (0.008% ammonium hydroxide, 0.012% acetic acid in water, v/v/v) and Mobile Phase B (0.008% ammonium hydroxide, 0.012% acetic acid in acetonitrile, v/v/v). The total analytical run time was 12.5 minutes. The mobile phase was nebulized using heated nitrogen in a Z-spray source/interface set to electrospray positive ionization mode. The ionized compounds were detected using ToF MS scan monitoring in sensitivity mode scanning from 50.0 to 700 *m/z*. This method is applicable for measuring GS-441524 concentrations ranging from 1.00 to 1,000 ng/mL using 20.0 μ L of plasma for extraction. The peak areas of GS-441524 were acquired using MassLynx V4.2 (Waters, Milford, MA). The calibration curve was obtained by fitting the peak area ratios of the analyte and the standard concentrations to a linear equation with 1/*x*² weighting using MassLynx. The equation of the calibration curve was then used to interpolate the concentrations of the analyte in the samples using their peak areas. The peak areas used for the calculations were not rounded.

Example 10 - Stability of ODE-P-RVn (**4c**) and ODBG-P-RVn (**15d**) in Human Plasma

One of the disadvantages of remdesivir is instability in plasma where it has been reported to persist at virologically significant levels for less than 2 hours after intravenous infusion. (1, 2). Remdesivir also has limited stability *ex vivo* in human plasma with a reported *T*_½ of 69 minutes (Siegel D, Hui HC, Doerffler E, Clarke MO, Chun K, Zhang L, Neville S, Carra E, Lew W, Ross B, Wang Q, Wolfe L, Jordan R, Soloveva V, Knox J, Perry J, Perron M, Stray KM, Barauskas O, Feng JY, Xu Y, Lee G, Rheingold AL, Ray AS, Bannister R, Strickley R, Swaminathan S, Lee WA, Bavari S, Cihlar T, Lo MK, Warren TK, Mackman RL. Discovery and Synthesis of a Phosphoramidate Prodrug of a Pyrrolo[2,1-*f*][triazin-4-amino] Adenine C-Nucleoside (GS-5734) for the Treatment of Ebola and Emerging Viruses. *J Med Chem.* 2017 Mar 9;60(5):1648-1661).

The stability of ODE-P-RVn and ODBG-P-RVn in human plasma was evaluated with either K₂EDTA or sodium heparin as an anticoagulant.

Plasma was spiked with 2 micrograms/ml concentrations of ODE-P-RVn or ODBG-P-RVn and incubated at 37° C. Samples were taken at 0.5, 1, 2, 4, 8 and 24 hours and frozen for later analysis by LC/MS/MS by the method shown in Example C. **FIG. 5A** and **FIG. 5B** shows that both ODE-P-RVn and ODBG-P-RVn were stable for at least 24 hours in human plasma with either K₂EDTA (**FIG. 5A**) or sodium heparin (**FIG. 5B**) as

anticoagulants. (See, e.g., Warren T.K. et al. *Nature*. 2016 Mar 17;531(7594):381-5; and Tempestilli, M. et al. *J. Antimicrob Chemother*. 2020 Oct 1;75(10):2977-2980).

Example 11- Effect of Antivirals of the Replication of the Respiratory Syncytial Virus

5 Methods: The SRV A2 strain of RSV was obtained from the ATCC. Ten thousand Hep-2 cells were seeded into each well of a 96-well plate in EMEM with 10% fetal bovine serum and 1% penicillin and streptomycin. Each well was inoculated with 100 TCID₅₀ of virus. Two hours later wells were washed once with medium and serial four-fold dilutions of candidate antiviral drugs were added. Cell cultures were observed for 72 to 96 hours
10 for cytopathic effects. The concentration of drug that reduced CPE by 50 or 90% was calculated using Prism 7 software and expressed as the EC₅₀ and EC₉₀, respectively.

Cell viability assay: For select compounds the CC₅₀ was calculated up to a maximum of 100μM using CellTiter-Glo. For these experiments, approximately 20k Calu-3 cells were seeded per well in opaque white 96-well plates (cat# 655073, Greiner Bio-
15 One, Monroe, North Carolina) and incubated 48-72h. Compounds or controls were added and incubated for 44h at 37 °C and 5% CO₂. After which an equal volume of CellTiter-Glo reagent (Cat. # G7570, Promega, Madison, WI) was added, mixed and luminescence recorded on a Veritas Microplate Luminometer (Turner BioSystems) according to manufacturer recommendations.

20

Antiviral effect of compounds of the present disclosure against RSV infection in Hep-2 cells

Compd	R ₁	R ₂	Hep-2 cells		
			EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)
15d	1- <i>O</i> -octadecyl-2- <i>O</i> -benzyl- <i>sn</i> -glyceryl	H	0.010	0.019	14.7
15j	1- <i>O</i> -octadecyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.010	0.019	9.6
15m	1- <i>O</i> -oleyl-2- <i>O</i> -(3-F, 4-MeO-Bn)- <i>sn</i> -glyceryl	H	0.005	0.0094	12.5


15c	1- <i>O</i> - hexadecyl-2- <i>O</i> -(3-F, 4- MeO-Bn)- <i>sn</i> - glyceryl	H	0.034	0.068	20.6
-----	---	---	-------	-------	------

Not wishing to be bound by any particular theory, it is believed that compounds 15j, 15m, and 15c may be at least 2 to 3 times more active than compound 15d. For example, compound 15m may be more active than remdesivir.

Compounds of the disclosure had very marked antiviral activity in Hep-2 cells infected with RSV A2. Cytotoxicity was moderate and the selectivity (CC50/EC50) ranged from 605 to 2,500.

Example 12 – Preparation of Formulations

Synthesis of HIV antivirals: Synthesis of ((2*S*,5*R*)-5-(4-amino-5-fluoro-2-oxo-3,4-dihydropyrimidin-1(2*H*)-yl)-1,3-oxathiolan-2-yl)methyl benzyl (2-(octadecyloxy)ethyl) phosphate (**ODE-benzyl phospho-emtricitabine**)




Scheme 5. Example process for synthesis of emtricitabine 5'-monophosphate analogs of the disclosure. *Reagents and Conditions:* a) emtricitabine, DCC, DMAP, pyridine, 80 °C, 72 h; b) benzyl alcohol, PyBOP, DIEA, DMF, rt, 3h.

N,N-Dicyclohexylcarbodiimide (DCC, 410 mg, 2.0 mmol) was added to a solution of emtricitabine (230 mg, 0.9 mmol), octadecyloxyethyl phosphate (320 mg, 0.74 mmol), and 4-dimethylaminopyridine (DMAP, 110 mg, 0.9 mmol) in 10 mL of dry pyridine, and then the mixture was heated to 80 °C and stirred for 3 days. Pyridine was then evaporated

and the residue was purified by flash column chromatography on silica gel 60. Gradient elution (CH₂Cl₂/methanol 10-20%) afforded 220 mg (47% yield) of , octadecyloxyethyl-phospho-emtricitabine: ESI MS: 622.29 [M-H]⁻; 624.10 [M+H]⁺, 646.30 [M+Na]⁺, 662.26 [M+K]⁺.

- 5 Octadecyloxyethyl-phospho-emtricitabine (220 mg, 0.35 mmol), benzyl alcohol (76 mg, 0.70 mmol), diisopropylethylamine (DIEA, 0.12ml, 0.70 mmol), and (1H-benzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 360 mg, 0.70 mmol) in dry DMF (10 mL) were stirred at room temperature. After 3 h. DMF was evaporated under vacuum. The residue was dissolved in ethyl acetate (50 mL) washed
10 with saturated NaHCO₃ (3 x 10 mL). The organic layer was dried over MgSO₄ and ethyl acetate was evaporated. Purification of the residue by silica gel column chromatography using ethyl acetate and ethanol (0-4%) yielded 130 mg (52% yield) of octadecyloxyethyl benzyl phospho-emtricitabine. ¹H NMR (400 MHz, CDCl₃-CD₃OD) δ ppm 7.91 (d, J=5.87 Hz, 1H), 7.40-7.48 (m, 4H), 6.27-6.34 (m, 1H), 5.35-5.41 (m, 1H), 5.19 (d, J=8.43
15 Hz, 2H), 4.33-4.48 (m, 2H), 4.21-4.28 (m, 2H), 3.65-3.71 (m, 2H), 3.48-3.61 (m, 3H), 3.37-3.42 (m, 2H), 3.06 - 3.18 (m, 1H), 1.53-1.64 (m, 2H), 1.25-1.40 (m, 30H), 0.93 (t, sCJ=6.23 Hz, 3H). ESI MS: 714.32 [M+H]⁺, 736.43 [M+Na]⁺.


- Synthesis of ((2S,5R)-5-(4-amino-5-fluoro-2-oxo-3,4-dihydropyrimidin-1(2H)-yl)-1,3-oxathiolan-2-yl) methyl benzyl ((R)-2-(benzyloxy)-3-(octadecyloxy)propyl) phosphate
20 (ODBG-benzyl-phospho-emtricitabine)



ODBG-benzyl-phospho-emtricitabine

- 1-*O*-octadecyl-2-*O*-benzyl-*sn*-glyceryl phosphate (ODBG-P-) was coupled to emtricitabine in pyridine using DCC/DMAP as described in Scheme 5 to afford ODBG-phospho-emtricitabine. Esterification of the phosphate diester with benzyl alcohol was
25 accomplished using PyBOP/DIEA as in Scheme 5 to yield the phosphotriester, ODBG-benzyl-phospho-emtricitabine.

Synthesis of benzyl (2-(octadecyloxy)ethyl) (((R)-1-(2,6-diamino-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (**ODE-benzyl 9-(R)-[2-(phosphonomethoxy)-propyl]2,6-diaminopurine**)




5

Scheme 6. Example process for synthesis of 9-(R)-[2-(phosphonomethoxy)propyl]2,6-diaminopurine) analogs of the disclosure. Reagents and Conditions: a) DCC, DMAP, pyridine, 80 °C, 72 h; b) benzyl alcohol, PyBOP, DIEA, DMF, rt, 3h

N,N-Dicyclohexylcarbodiimide (3 eq) is added to a mixture of 9-(R)-[2-

- 10 (phosphonomethoxy)propyl]-2,6-diaminopurine ((R)-PMPDAP, 1 eq, see: Krečmerová, M., Jansa, P., Dračinský, M., Sázelová, P., Kašička, V., Neyts, J., Auwerx, J., Kiss, E., Goris, N., Stepan, G., Janeba, Z. 9-[2-(R)-(Phosphonomethoxy)propyl]-2,6-diaminopurine (R)-PMPDAP and its prodrugs: Optimized preparation, including identification of by-products formed, and antiviral evaluation in vitro, *Bioorganic & Medicinal Chemistry*, 2013, 21: 1199-1208.), octadecyloxyethan-1-ol (1 eq) and 4-dimethylaminopyridine (DMAP, 1 eq) in dry pyridine and the mixture is heated to 80 °C until conversion to the esterified product is complete. The monoester (ODE-(R)-PMPDAP) was isolated after column chromatography. To a mixture of ODE-(R)-PMPDAP (1 eq), benzyl alcohol (1.5 eq) and diisopropylethylamine (DIEA, 1.5 eq) in dry DMF was added (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP, 1.5 eq) and the mixture was stirred at room temperature until formation of the diester was complete. The pure product, ODE-benzyl-(R)-PMPDAP was isolated by column chromatography.
- 20

- Synthesis of benzyl (2-(octadec-9-en-1-yloxy)ethyl) (((R)-1-(2,6-diamino-9H-purin-9-yl)propan-2-yl)oxy)methyl)phosphonate (**OLE-benzyl 9-(R)-[2-(phosphonomethoxy)-propyl]2,6-diaminopurine**)L
- 25




oleyloxyethyl-benzyl 9-(R)-[2-(phosphomethoxy)propyl]2,6-diaminopurine

A mixture of (R)-PMPDAP and oleyloxyethanol in pyridine was treated with DCC/DMAP as in Scheme 6 to afford the monoester, OLE-(R)-PMPDAP.

- 5 OLE-(R)-PMPDAP was further esterified using benzyl alcohol, PyBOP and DIEA. OLE-benzyl-(R)-PMPDAP was isolated after column chromatography.

Long-acting HIV antivirals: For use in long-acting intramuscular treatment of infection with the human immunodeficiency virus, the antiviral compounds suitable for use are described in this application and in US Patent 8,835,630 and include the following:

- 10 *Compounds useful as long-acting HIV antivirals*



Antiviral activity in HIV infected human PBMCs

Compound	EC ₅₀ , nM	CC ₅₀ , μ M	Selectivity index
ODE-Bn-Tenofovir	1.7 $\pm$ 1.5	8.72	5130
ODE-PMPDAP	0.09 $\pm$ 0.05	7.0 $\pm$ 7.1	77,700
OLE-PMPDAP	0.10 $\pm$ 0.03	15.0 $\pm$ 12.3	150,000
ODE-Bn-PMPDAP	active		
OLE-Bn-PMPDAP	active		
ODE-P-Emtricitabine	9.6 $\pm$ 3.1	35.6 $\pm$ 12.9	3708
ODE-Bn-P-Emtricitabine	18.3 $\pm$ 8.7	54.7 $\pm$ 45.3	2989
Abbreviations: EC ₅₀ , 50% effective concentration; CC ₅₀ , 50% cytotoxic concentration; SI, selectivity index (CC ₅₀ /EC ₅₀). Method : inhibition of antiviral activity was measured in human PBMCs infected with HIV-1 as reported in Beadle et al. Antiviral Research vol. 171 (2019): 104614.			

Compounds of this example were active with EC₅₀ values in the 0.09 to 18
 5 nanomolar range.

Pharmaceutical Formulations: Also provided herein are pharmaceutical
 formulations. The pharmaceutical formulations included a compound as described herein,
 such as a compound of formula (I). In some embodiments, the pharmaceutical
 formulation was orally bioavailable. In some embodiments, the pharmaceutical
 10 formulation is formulated for injection, such as intramuscular injection.

The pharmaceutical formulations included one compound described herein, or
 more than one (e.g., two, three, etc.) compounds described herein. The pharmaceutical
 formulations included any one or more pharmaceutically acceptable excipients. Oil based
 formulations for compounds of the disclosure have the following composition by weight
 15 (w/w%):

Composition of intermediate- and long-acting formulations of compounds for treatment of coronavirus or HIV infections

Components	F3	F4	F7	F8
Sesame oil	92.65	73.67	-	-
Medium chain triglycerides	-	-	92.65	73.67
Benzyl alcohol	1.22	10.67	1.22	10.67
Benzyl benzoate	-	15.67	-	15.67
Ethyl alcohol	6.12	-	6.12	-

The following materials were used in the formulations of the foregoing table:

Materials and grade	Source and lot	%assay
Super refined sesame oil, NF	Croda, 0001095029	N/A
Medium chain triglycerides, USP	IOI Oleochemical, 190104	N/A
Benzyl alcohol, USP	EMD, K43136587447	98.0-100.5
Benzyl benzoate	Sigma-Aldrich, MKBQ9481V	99.5
Ethanol, USP 200 proof	Decon labs,	100

5 In some embodiments, one or more of the compounds described herein, such as those of formula (I), were soluble in an oil, such as one of formulations F3, F4, F7, and F8. When an amount of one or more compounds described herein, such as those of formula (I), were soluble in an oil, then the resulting pharmaceutical formulation may be used in any of the methods herein, including a method of treating coronavirus with a
10 single dose of the pharmaceutical formulation.

Formulations of the foregoing table were prepared using compounds of the disclosure. Antiviral prodrugs of the disclosure employing HIV antivirals or RNA virus antiviral compounds described in PCT/US2021/043094 may be utilized in the formulations. As a non limiting example the following pharmacokinetic experiments were
15 done using ODE-Bn-Tenofovir as the antiviral prodrug.

Comparison of Formulations F3, F4, F7 and F8 following a single IM administration in rats - Groups: Vehicle control, 10 mg/kg, 30 mg/kg and 100 mg/kg of formulations 3,4,7, 8.

Procedure: At time zero rats were injected intramuscularly with 100 microliters of
20 vehicle and formulations as noted. The animals were observed for general health and weights were obtained at the start and twice a week thereafter. Blood was obtained at 6, 24 and 48 hours, and at 7, 14, 28, 35 and 42 days and the plasma levels of ODE-Bn-TFV and ODE-TFV (the first metabolite, also an active anti-HIV compound) were determined by LC/MS/MS.

25 **FIG. 6A** and **FIG. 6B** depict pharmacokinetics of ODE-Bn-TFV and ODE-TFV, respectively, after intramuscular administration of 100 mg/kg in formulation F3 to rats.

Results: Rats were injected IM at day zero and weights and clinical observations were acquired twice a week for 42 days. The increases in body weight in vehicle and drug treated rats were identical and no clinical signs were noted. Comprehensive metabolic panels at study termination did not show any abnormalities in liver or kidney functions tests. CBCs were normal controls and all drug dosage groups. After administration ODE-Bn-TFV was metabolized to the more active metabolite, ODE-TFV. Clearly as seen in the ODE-Bn-TFV curves (left) and the curves for the active first metabolite, ODE-TFV (right), doses of 10 and 30 mg/kg (groups 2 & 3) were not adequate to provide antivirally significant levels of compound for 30 days and beyond. Only the 100 mg/kg dose (group 4) provided levels of ODE-Bn-TFV and ODE-TFV at a virally significant level for 28 days.

The two log₁₀ panels below show various formulations at doses of 100 mg/kg (formulation drug concentrations of 200 mg/ml). The left panel was ODE-Bn-TFV and the right panel shows plasma levels of ODE-TFV (nM/L). All four formulations given at 100 mg/kg allow for maximal plasma levels of ODE-Bn-TFV at 6 to 48 hours followed by a gradual decline to 2.87 ng/ml (F8) to 6.95 (F3). The highest values at day 28 were noted in F3, ODE-Bn-TFV 6.95 ng/ml (10.3 nM) and ODE-TFV 1.78 ng/ml (3.05 nM). These values were above the EC₅₀s for these two compounds (1.7 and 1.1 nM).

FIG. 7A and **FIG. 7B** are plots of the effects of 4 different formulations on the pharmacokinetics of ODE-Bn-TFV and ODE-TFV, respectively, after intramuscular administration of 100 mg/kg to rats.

Pharmacokinetic parameters

Pharmacokinetic Parameters				
	F3	F4	F7	F8
C _{max} , nM	130	139	76	187
T _{1/2} hr	142	171	168	278
AUC (0-inf) nM.hr	35200	33800	21400	48100
AUC (14-28 hr) nM.hr	10900	6900	3640	5420

The following figures shows the comparison between intramuscular ODE-Bn-TFV in Formulation F3 versus formulation F8 in rats. ODE-Bn-TFV in F3 provided lower early levels of drug in comparison with F8 but levels persisted longer, remaining above the HIV EC₉₀ for 28 days. The F8 formulation produced higher drug levels until day 15 and was more suitable for treatment courses of 5 to 14 days.

FIG. 8 depicts a comparison of 96 mg/kg ODE-Bn-TFV pharmacokinetics in rats using formulation F3 versus F8.

Calculated area under curve (AUC) percentages during the first and second 14 days showing the release profile are shown in the table below.

Release profile: First 14 days versus second 14 days.

Formulation	% AUC first 14 days	% AUC second 14 days
F3	69	31
F4	79	21
F7	83	17
F8	89	11

Formulation F3 provided the greatest exposure during the second 14 days. Formulation F8 and F7 showed the highest percentage exposures, 89 and 83%, during the first 14 days.

3 Month study of Formulations F3 in rats with measurements of plasma levels at 15 and 30 days following monthly IM injections of ODE-Bn-TFV - Since the initial plasma values were very high (~100 ng/ml) and declined progressively to slightly suboptimal values at day 28, it was considered important to know the PBMC TFVpp levels in a longer multi-dose study. A 3-month study was designed to test mid and trough plasma levels of ODE-Bn-TFV and ODE-TFV in male Sprague Dawley rats following monthly IM injections of F3 containing ODE-Bn-TFV.

Maximal volume IM injections of 100 microliters were given at day 0, 30 and 60 to groups of 3 rats. There were 3 groups of control animals who received monthly injections of F3 vehicle. Plasma drug levels were analyzed by LC/MS/MS. Weights were assessed over the 3-month period. Rats increased weight from 355 gm to 705 mg at study termination at 105 days. Because a constant (maximal) IM volume of 100 microliters was

given, the initial dose was 90 mg/kg and the two following injections at days 30 and 60 were lower, 62 mg/kg, due to weight increases of the rats. There were no clinical observations of significance during the 3 month study and metabolic panels and CBCs showed no abnormalities at day 105.

5 **FIG. 9** depicts rat plasma levels of ODE-Bn-TFV and ODE-TFV during a 3 month exposure to monthly intramuscular doses of ODE-Bn-TFV in formulation F3.

Plasma drug levels: 15 days following the first dose ODE-Bn-TFV levels were high and the first metabolite, ODE-TFV, were about 30% of the parent compound. However, as the 3 month study proceeded, ODE-TFV levels gradually became the
10 dominant active compound in plasma. After the last dose at 60 days, levels of ODE-TFV remained substantial until day 105 at which time they were 11 nanomolar. The EC₉₀ of ODE-TFV is 10 nM (Beadle JR, Aldern KA, Zhang XQ, Valiaeva N, Hostetler KY, Schooley RT. Octadecyloxyethyl benzyl tenofovir: A novel tenofovir diester provided sustained intracellular levels of tenofovir diphosphate. (Antiviral Res. 2019
15 Nov;171:104614). Therapeutic levels of ODE-Bn-TFV and ODE-TFV were maintained to 105 days following 3 monthly IM doses at 0, 30 and 60 days. Rats received maximal (volume) doses of 100 microliters. Since their weight doubled during treatment, the second and third doses were only 62 mg/kg. Even with the lower 2nd and 3rd doses, therapeutic levels of drug were maintained for 105 days. Therapeutic levels of the ODE-
20 TFV metabolite persisted for 105 days.

Canine Pharmacokinetics of ODE-Bn-TFV in Formulation F3 - A 30 day study in beagle dogs was done with 30 mg/kg IM doses of ODE-Bn-TFV in formulation F3.


FIG. 10 shows the levels of ODE-Bn-TFV (circles) and ODE-TFV (squares). EC₉₀ values are shown in triangles to indicate when the plasma drug levels pass below the
25 90% effective level. At 30 mg/kg intramuscular ODE-Bn-TFV in formulation F3 provided 30 to 34 days above the EC₉₀ values for both ODE-Bn-TFV and its active metabolite, ODE-TFV.

FIG. 10 depicts plasma levels of ODE-Bn-TFV and ODE-TFV in beagle dogs treated with 100 mg/kg with ODE-Bn-TFV in formulation F3.

30 There were no adverse effects except for some transient swelling at the injection sites. Pre and Post CBCs and metabolic panels before and after the study did not show any abnormalities. Of note, no abnormal liver or kidney function tests were noted. Other


compounds of the disclosure may be used in formulation F3 to provide prolonged protection from HIV infection.

Antiviral Compounds and Formulations for Intramuscular Treatment of Coronavirus Infections - In SARS-CoV-2 infection active viral replication usually lasts for 5 to 10 days. To treat with an intramuscular dose, antiviral prodrugs in formulation F8 at 200 to 400 mg/ml were expected to provide effective drug levels for at least 7 days. Non-limiting examples of effective antiviral prodrugs of the remdesivir nucleoside monophosphate may be selected from the structures shown below:




1-O-octadecyl-2-O-benzyl-*sn*-glyceryl-phospho-RVn

10



1-O-oleyl-2-O-(3-fluoro, 4-methoxybenzyl)-*sn*-glyceryl-phospho-RVn




1-O-octadecyl-2-O-benzyl-*sn*-glyceryl-benzyl-phospho-RVn

Antiviral Compounds and Formulations for Intramuscular Treatment of Human immunodeficiency infections - For use in long-acting intramuscular treatment of infection

- 5 with the human immunodeficiency virus, the antiviral compounds suitable for use are described in US Patent 8,835,630. These compounds may be used in formulations F3 or F4.

Non-limiting examples that may be used in the formulations described herein include the following compounds:

10



5 **Antiviral activity of compounds suitable for use in long-acting formulations in human PBMCs infected with HIV-1**

Compound	EC ₅₀ , nM	CC ₅₀ , μM	Selectivity index
ODE-Bn-Tenofovir	1.7±1.5	8.72	5130
ODE-PMPDAP	0.09 ± 0.05	7.0 ± 7.1	77,700
OLE-PMPDAP	0.10 ± 0.03	15.0 ± 12.3	150,000
ODE-Bn-PMPDAP	active		
OLE-Bn-PMPDAP	active		
ODE-P-Emtricitabine	9.6 ± 3.1	35.6 ± 12.9	3708
ODE-Bn-P-Emtricitabine	18.3 ± 8.7	54.7 ± 45.3	2989

Abbreviations: ODE-, octadecyloxyethyl; OLE-, oleyloxyethyl; PMP-, phosphonylpropyloxy; DAP, 2,6-diaminopurine; EC₅₀, 50% effective concentration; CC₅₀, 50% cytotoxic concentration. Method : inhibition of antiviral activity was measured in human PBMCs infected with HIV-1 as reported in Beadle et al. Antiviral Research vol. 171 (2019): 104614.

As explained herein and in this example, 22 analogs of ODE-TFV were synthesized to determine effective diester substituents in comparison with benzyl in ODE-Bn-TFV. Antiviral activities were determined in HIV infected human PBMCs. The pharmacokinetics of TFVpp were determined in HFF cells over a 28 day period.

The variation of the diester substituent on anti-HIV activity of ODE-TFV is depicted in the following table:

Substituent	EC ₅₀ nM	CC ₅₀ nM	Selectivity Index
H	1.1	1284	1170
Benzyl	1.7	8720	5130
Phenyl	2.4	3930	1640
4-methyl benzyl	2.1	3410	1620
3-methyl benzyl	3.2	3860	1210
2-methyl benzyl	2.7	1440	532
Methyl	5.2	2980	677
Ethyl	57	5365	95
Iso-propyl	183	1730	94
Cyclopropyl	413	5280	13

- 5 **FIG. 11A** and **FIG. 11B** depict the TFVpp persistence in HFF cells of ODE-TFV (**FIG. 11A**) and ODE-Et-TFV (**FIG. 11B**).

The persistence of TFVpp in HFF cells after a 3 day exposure is depicted at the following table. HFF cells were exposed to 1 micromolar drug for 3 days. At days 0, 1, 2, 4, 7, 14, 21, and 28, cells were trypsinized, counted, and processed for LC/MS measurement of TFVpp. Data were expressed at ftg/cell. nm = not meaningful; ftg – gentogram.

Compound	C _{max} , ftg/cell	T _{max} ,d	T _{1/2}	T _{1/10}	AUC
ODE-TFV	189	0	1.2	2.1	323
ODE-Bn-TFV	72.5	2	3.7	8.7	324
ODE-4MeBN-TFV	49.5	0	3.1	11.8	233
HDP-Bn-TFV	15.0	0	1.8	10.2	66.3

ODE-Me-TFV	9.3	0	2.0	11.3	37.2
ODE-Ph-TFV	8.1	1	3.0	15.1	49.2
ODE-Et-TFV	0.91	0	nm	nm	4.2
ODE-iPr-TFV	0.69	0	nm	nm	2.0

In this example, 6 oil-based formulations for ODE-Bn-TFV were prepared. The accelerated stability of 20 mg/mL formulations was determined. API loadings were assessed.

- 5 In this example, formulations of ODE-Bn-TFV for IM usage (20 mg/mL formulations) were prepared, and included the components of the following table (w/w%):

Components	F3V	F4V	F5V
Sesame Oil	92.65	73.67	89.765
Benzyl alcohol	1.22	10.67	10.20
Benzyl benzoate	0	15.67	0
Alcohol	6.12	0	0
BHT	0	0	0.0306

In this example, a two month accelerated stability study was conducted. The results and observations of this study are provided at the following tables:

2-8 °C	Appearance	Concentration (mg/g)	Recover over T ₀
F3	Colorless and Clear	20.09	99.0
F4	Light Yellow and Clear	20.45	102.1
F5	Colorless and Clear	19.85	100.7

25 °C	Appearance	Concentration (mg/g)	Recover over T ₀
F3	Colorless and Clear	20.52	101.1
F4	Yellow and Clear	19.57	97.7
F5	Colorless and Clear	19.79	100.3

40 °C	Appearance	Concentration (mg/g)	Recover over T ₀
F3	Colorless and Clear	20.56	101.3
F4	Yellow and Clear	19.08	95.3
F5	Colorless and Clear	19.15	97.10

Additional formulations were prepared for tested, as provided in the following
 5 table (w/w%):

Components	F3V	F4V	F7V	F8V
Sesame Oil	92.65	73.67	-	-
Med. Chain TGs	-	-	92.65	73.67
Benzyl alcohol	1.22	10.67	1.22	10.67
Benzyl benzoate	-	15.67	-	15.67
Alcohol	6.12	-	6.12	-

A 42 days pharmacokinetic study of formulations 3, 4, 7, and 8 of this example in
 rats was conducted. In the study, 100, 30, and 10 mg/kg of each formulation was
 10 administered by intramuscular (IM) injection to male Sprague Dawley rates (100
 microliters). Blood was taken at 0.25, 1, 2, 7, 14, 21, 28, 35, and 42 days after injection,
 and plasma levels of ODE-Bn-TFV and ODE-TFV were determined. The rats were
 subjected to twice weekly clinical observations, and weights were collected.

FIG. 12A and **FIG. 12B** depict plasma concentrations of formulation F3V
 15 containing ODE-Bn-TFV (**FIG. 12A**) and ODE-TFV (**FIG. 12B**) (ODE-Bn-TFV ED₉₀
 10.9 ng/mL; ODE-TFV ED₉₀ 6.3 nm/mL).

FIG. 12C and **FIG. 12D** depict plasma concentrations of formulation F4V containing ODE-Bn-TFV (**FIG. 12C**) and ODE-TFV (**FIG. 12D**).

FIG. 12E and **FIG. 12F** depict plasma concentrations of formulation F7V containing ODE-Bn-TFV (**FIG. 12E**) and ODE-TFV (**FIG. 12F**).

5 **FIG. 12G** and **FIG. 12H** depict plasma concentrations of formulation F8V containing ODE-Bn-TFV (**FIG. 12G**) and ODE-TFV (**FIG. 12H**).

FIG. 12I depicts mean plasma concentrations of formulations F3V, F4V, F7V, and F8V.

10 The API pharmacokinetic parameters of this example are provided at the following table:

	F3	F4	F7	F8
C _{max} , nm	130	139	76	187
T _{1/2} hour	142	171	168	278
AUC (0-inf) nM.hr	35200	33800	21400	48100
AUC (14-28 hr)nM.hr	10900	6900	3460	5420

The ODE-BN-TFV release provided observed in this example is summarized, at least in part, in the following table:

First 14 Days versus Second 14 Days		
Formulation	% AUC first 14 Days	% AUC second 14 Days
F3	69	31
F4	79	21
F7	83	17
F8	89	11

The results of this example that through chemical and formulation alterations, TFV was converted to long-acting anti-HIV status. Formulations F3 and F4 of this example released, for example, 69 and 79 % of the lead compound in the first 14 days, but this release profile may be modified as described herein. Using formulation F3, a 3 month study was conducted in rats with monthly IM injections of 100 microliters, and measuring plasma drug levels and PBMC levels of TFVpp.

105 Days Pharmacokinetic Study (Monthly IM Injections of F3 in Rats): 100 microliters of the API of formulation 3 was administered monthly by IM injection to male Sprague Dawley rats. Due to the fact that the rats gained significant weight over the 105 day period, the actual dosages based on weight were as follows: 90, 62, and 60 mg/kg. Blood was collected at 15, 30, 45, 60, 75, 90, 98, and 105 days. Plasma levels of ODE-Bn-TFV and ODE-TFV were determined. PBMCs also were obtained at the foregoing days, and $f_{mol}/10^6$ cells was determined.


FIG. 13 depicts the nanograms/mL of ODE-Bn-TFV and ODE-TFV in plasma.

FIG. 14 depicts TFV disphosphate in PBMCs.

The rats of this example gained weight during the 105 day study; about 350 grams to about 670 grams. There was no difference in growth curves between the control and treated rats. There also were not reported clinical observations in either group. CBCs in treated and control animals were normal. Complete metabolic panels were normal in both groups. Of importance was the fact that there were no changes noted in BUN or creatinine.

Example 13 – Synthesis of 2-, 3-, and 4-cyanobenzyl Analogs

The following is a scheme depicting embodiments of the synthesis of 2-, 3-, and 4-cyanobenzyl analogs.




5

Scheme 7. Synthesis of 2, 3 and 4-cyanobenzyl analogs. *Reagents:* a) sodium hydride, 2,3 or 4-(bromomethyl)benzonitrile, tetrabutyl ammonium iodide, THF; b) p-TsOH (cat.), CH₂Cl₂, MeOH; c) i) POCl₃, triethylamine, CH₂Cl₂, ii) acetone/ice water; d) RVn (GS-441524) acetonide, DIC/NMI, pyridine, 35 °C; e) formic acid, rt.

10

Example 13a. Synthesis of 1-O-Octadecyl-2-O-(4-cyanobenzyl)-sn-glyceryl-phospho-RVn (OD(4-CN-Bn)G-P-RVn)



15 Sodium hydride (440 mg, 11 mmol) was added to a cooled (0 °C) solution of (*R*)-1-((4-methoxyphenyl)diphenylmethoxy)-3-(octadecyloxy)propan-2-ol (3.39 g, 5.5 mmol) and tetrabutylammonium iodide (600 mg, 1.65 mmol) in dry THF (30 mL). After stirring vigorously for 20 min., 4-(bromomethyl)benzonitrile (1.61 g, 8.2 mmol) was added and the mixture was allowed to warm to ambient temperature and stirred overnight. The
20 reaction was then quenched with ice (25 mL), diluted with ethyl ether (75 mL), washed

with H₂O (2 x 25 mL) and dried over anhydrous MgSO₄. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 50% EtOAc- in hexanes) to yield (R)-4-(((1-((4-methoxyphenyl)diphenylmethoxy)-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile (610 mg, 15%) as a white solid. MS (ESI) *m/z* [M+Na]⁺ 754.54.


To a solution of (R)-4-(((1-((4-methoxyphenyl)diphenylmethoxy)-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile (400 mg, 0.54 mmol) in 1:1 CH₂Cl₂/MeOH (30 mL) was added p-toluenesulfonate monohydrate (5 mg, 0.03 mmol) and the mixture was stirred at room temperature until deprotection was complete (approx. 3 h) according to TLC analysis. Saturated aq NaHCO₃ (200 mg) was added, and the solvent evaporated under vacuum. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 10% EtOAc in hexanes) to afford (S)-4-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile, as clear oil (80 mg, 32%). **MS** (ESI) *m/z* [M+Na]⁺ 482.55.

A solution of (S)-4-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile (80 mg, 0.17 mmol) and triethylamine (0.08 mg, 0.8 mmol) was added to a cooled (0 °C) solution of POCl₃ (30 mg, 0.20 mmol), in CH₂Cl₂ (3 mL) and stirred for 3 h. The mixture was added to acetone/ice water, stirred 1h, then extracted with CH₂Cl₂. (R)-2-((4-cyanobenzyl)oxy)-3-(octadecyloxy)propyl dihydrogen phosphate was isolated (80 mg, 87 %) and used without further purification. **MS** (ESI) *m/z* [M-H]⁻ 538.47.

The phosphate (200 mg, 0.37 mmol) was coupled to RVn (GS-441524)-acetone (132 mg, 0.40 mmol) using diisopropylcarbodiimide (DIC, 100 mg, 0.80 mmol), N-methylimidazole (NMI, 98 mg, 1.2 mmol) in pyridine (10 mL) to yield ((3aR,4R,6R,6aR)-6-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-6-cyano-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)methyl ((R)-2-((4-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate (60 mg, 19 %, **MS** *m/z* [M-H]⁻ 851.55) which was then treated with formic acid (1 mL) and stirred 6 h. Evaporation of the solvent and purification by flash column chromatography yielded ((2R,3S,4R,5R)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-5-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methyl ((R)-2-((4-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate (20 mg, 35%) as an off white solid. **¹H NMR** (400 MHz, CD₃OD + CDCl₃) δ 7.76 (s, 1H, *H*₂), 7.60 (d, *J* = 8.0 Hz, 2H, -CH₂-aryl-*H*₃+*H*₅), 7.49 (d, *J* = 7.9 Hz, 2H, -CH₂-aryl-*H*₂+*H*₆), 6.96 (d, *J* = 4.4 Hz, 1H, *H*₉), 6.86 (d, *J*

= 4.5 Hz, 1H, *H*8), 4.80 (d, *J* = 5.3 Hz, 1H, *H*2'), 4.77 – 4.63 (m, 2H, -CH₂-aryl), 4.37 (d, *J* = 4.7 Hz, 1H, *H*4'), 4.25 (t, *J* = 5.4 Hz, 1H, *H*3'), 4.11 (d, *J* = 18.7 Hz, 2H, *H*5'), 3.90 (s, 2H, *H*3''), 3.80 – 3.68 (m, 1H, *H*2''), 3.49 (qd, *J* = 10.6, 4.9 Hz, 2H, *H*1'), 3.39 (tt, *J* = 6.0, 3.0 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.52 (p, *J* = 6.9 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.26 (br s, *J* = 5.7 Hz, 30H, -OCH₂CH₂(CH₂)₁₅-), 0.89 (t, *J* = 6.8 Hz, 3H). ¹³C NMR (101 MHz, CD₃OD + CDCl₃) δ 156.63 (*C*6), 145.57 (*C*2), 132.62, (-CH₂-aryl-*C*1), 128.47 (-CH₂-aryl-*C*3 + *C*5), 125.14 (-CH₂-aryl-*C*2 + *C*6), 119.38 (-CN), 117.29 (*C*7), 111.95 (*C*5), 111.28 (-CN'), 84.60(*C*4'), 80.18(*C*1'), 75.38 (*C*2'), 72.24 (*C*2''), 71.57 (-CH₂-aryl), 71.40 (-OCH₂CH₂(CH₂)₁₄-), 71.12 (*C*1''), 65.75 (*C*3''), 65.20 (*C*5'), 32.57 (-OCH₂CH₂(CH₂)₁₄-), 30.89 – 29.57 (m) (m, -OCH₂CH₂(CH₂)₁₄-), 26.75 (-CH₂CH₂CH₃), 23.26 (-CH₂CH₂CH₃), 14.21 (-CH₃). HRMS (ESI) *m/z* [M-H]⁻ calcd for C₄₁H₆₀N₆O₉P 811.4165, found 811.4178. HPLC purity 98.6%.

Example 13b. Synthesis of 1-*O*-Octadecyl-2-*O*-(3-cyanobenzyl)-*sn*-glyceryl-phospho-RVn (OD(3-CN-Bn)G-P-RVn)



Sodium hydride (60 mg, 2.5 mmol) was added to a cooled (0 °C) solution of (*R*)-1-(octadecyloxy)-3-(trityloxy)propan-2-ol (540 mg, 0.92 mmol) and tetrabutylammonium iodide (180 mg, 0.5 mmol) in dry THF (30 mL). The mixture was stirred vigorously for 20 min before 3-(bromomethyl)benzonitrile (480 mg, 2.6 mmol) was added, and then the mixture was allowed to warm to ambient temperature and stirred overnight. The reaction was then quenched with ice (25 mL), diluted with ethyl ether (75 mL), washed with H₂O (2 x 25 mL) and dried over anhydrous MgSO₄. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 50% EtOAc- in hexanes) to yield (*R*)-3-(((1-(octadecyloxy)-3-(trityloxy)propan-2-yl)oxy)methyl)benzonitrile (600 mg, 93%) as a white solid. MS (ESI) *m/z* [M+Na]⁺ 724.57.


To a solution of (R)-3-(((1-(octadecyloxy)-3-(trityloxy)propan-2-yl)oxy)methyl)benzonitrile (600 mg, 0.85 mmol) in 1:1 CH₂Cl₂/MeOH (30 mL) was added p-toluenesulfonate monohydrate (8 mg, 0.04 mmol) and the mixture was stirred at room temperature until deprotection was complete according to TLC analysis (approx. 3 h). Saturated aq NaHCO₃ (200 mg) was added, and the solvent evaporated under vacuum. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 10% EtOAc in hexanes) to afford (S)-3-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile as clear oil (380 mg, 97%). **MS** (ESI) *m/z* [M+H]⁺ 460.57, [M+Na]⁺ 482.52.

A solution of (S)-3-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile (360 mg, 0.78 mmol) and triethylamine (340 mg, 3.0 mmol) was added to a cooled (0 °C) solution of POCl₃ (140 mg, 2.0 mmol) in CH₂Cl₂ (5 mL) and stirred for 3 h. The mixture was added to acetone/ice water, stirred 1h, then extracted with CH₂Cl₂. Evaporation of the organic phase gave (R)-2-((3-cyanobenzyl)oxy)-3-(octadecyloxy)propyl dihydrogen phosphate (850 mg, 1.57 mmol, **MS** (ESI) *m/z* [M-H]⁻ 538.46), which was then coupled to RVn (GS-441524)-acetone (600 mg, 1.8 mmol) using DIC (450 mg, 3.6 mmol), NMI (440 mg, 5.4 mmol) in pyridine (30 mL). Reaction at 35 °C overnight followed by chromatographic purification yielded ((3aR,4R,6R,6aR)-6-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-6-cyano-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)methyl ((R)-2-((3-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate (360 mg, 26 %) **MS** (ESI) *m/z* [M+Na]⁺ 875.49.

The acetone (340 mg, 0.39 mmol) was added to formic acid (7 mL) at room temperature and stirred 3 h. Evaporation of the solvent and purification by flash column chromatography afforded ((2R,3S,4R,5R)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-5-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methyl ((R)-2-((3-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate (290 mg, 90%) as an off white solid. **¹H NMR** (400 MHz, CD₃OD + CDCl₃) δ 7.71 (s, 1H, *H*₂), 7.70 (s, 1H, -CH₂-aryl-*H*₂), 7.62 (d, *J* = 7.8 Hz, 1H, -CH₂-aryl-*H*₄), 7.56 (d, *J* = 7.8 Hz, 1H, -CH₂-aryl-*H*₅), 7.44 (t, *J* = 7.6 Hz, 1H, -CH₂-aryl-*H*₆), 6.96 (d, *J* = 4.5 Hz, 1H, *H*₉), 6.86 (d, *J* = 4.5 Hz, 1H, *H*₈), 4.72 (d, *J* = 5.3 Hz, 1H, *H*₂'), 4.72 (d, *J* = 12.4 Hz, 1H, -CH₂-aryl), 4.65 (d, *J* = 12.4 Hz, 1H, -CH₂-aryl), 4.39 (q, *J* = 4.4 Hz, 1H, *H*₄'), 4.27 (t, *J* = 5.5 Hz, 1H, *H*₃'), 4.10 (d, *J* = 11.5 Hz, 1H, *H*₅'), 3.89 (d, *J* = 11.2 Hz, 1H, *H*₅'), 3.78 (m, 2H, *H*₃'), 3.71 (q, *J* = 5.0 Hz, 1H, *H*₂'),

- 3.51-3.40 (m, 2H, *H1'*), 3.41 (tt, *J* = 5.5, 2.7 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.54 (p, *J* = 6.6 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.26 (br s, 30H, -OCH₂CH₂(CH₂)₁₅), 0.89 (t, 3H, -CH₃). ¹³C NMR (101 MHz, CD₃OD + CDCl₃) δ 156.64 (*C6*), 147.68 (*C2*), 141.50 (-CH₂-aryl-*C1*), 132.77 (-CH₂-aryl-*C2*), 131.73(-CH₂-aryl-*C4*), 131.57 (-CH₂-aryl-*C6*), 129.88 (-CH₂-aryl-*C5*), 125.25 (*C7*), 119.50 (*C5*), 117.41 (-CN), 112.62 (-CN'), 111.98 (*C8*), 102.47 (*C9*), 84.91 (*C4'*), 80.07 (*C1'*), 75.57 (*C2'*), 72.40 (-CH₂-aryl), 71.58 (-OCH₂CH₂(CH₂)₁₄-), 71.44 (*C1''*), 71.23 (*C3'*), 65.73 (*C3''*), 65.21 (*C5'*), 32.67 (-OCH₂CH₂(CH₂)₁₄-), 31.02 – 29.33 (m, -OCH₂CH₂(CH₂)₁₄-), 26.85 (-CH₂CH₂CH₃), 23.37 (-CH₂CH₂CH₃), 14.39 (-CH₃). HRMS (ESI) *m/z* [M-H]⁻ calcd for C₄₁H₆₀N₆O₉P 811.4165, found 811.4178.
- 10 HPLC purity 98.6%.

Example 13c. Synthesis of 1-*O*-Octadecyl-2-*O*-(2-cyanobenzyl)-*sn*-glyceryl-phospho-RVn (OD(2-CN-Bn)G-P-RVn)



- Sodium hydride (60 mg, 2.5 mmol) was added to a cooled (0 °C) solution of (*R*)-1-(octadecyloxy)-3-(trityloxy)propan-2-ol (587 mg, 1.0 mmol) and tetrabutylammonium iodide (180 mg, 0.5 mmol) in dry THF (30 mL). The mixture was stirred vigorously for 20 minutes, before 2-(bromomethyl)benzonitrile (455 mg, 2.5 mmol) was added, and then the solution was allowed to warm to ambient temperature overnight. The reaction was then quenched with ice (25 mL), diluted with ethyl ether (75 mL), washed with H₂O (2 x 25 mL) and dried over anhydrous MgSO₄. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 50% EtOAc- in hexanes) to yield (*R*)-2-(((1-(octadecyloxy)-3-(trityloxy)propan-2-yl)oxy)methyl)benzonitrile (580 mg, 83%) as a white solid. MS (ESI) *m/z* [M+Na]⁺ 724.62.
- 20 mL) and dried over anhydrous MgSO₄. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 50% EtOAc- in hexanes) to yield (*R*)-2-(((1-(octadecyloxy)-3-(trityloxy)propan-2-yl)oxy)methyl)benzonitrile (580 mg, 83%) as a white solid. MS (ESI) *m/z* [M+Na]⁺ 724.62.

- To a solution of (*R*)-2-(((1-(octadecyloxy)-3-(trityloxy)propan-2-yl)oxy)methyl)benzonitrile (580 mg, 0.83 mmol) in 1:1 CH₂Cl₂/MeOH (15 mL) was added
- 25

p-toluenesulfonate monohydrate (8 mg, 0.04 mmol) and the mixture was stirred at room temperature until deprotection was complete (approx. 3 h) according to TLC analysis. Saturated aq NaHCO₃ (200 mg) was added, and the solvent evaporated under vacuum. The residue was adsorbed onto silica gel and purified by flash column chromatography (gradient: 0 to 10% EtOAc in hexanes) to afford compound (*S*)-2-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile as clear oil (380 mg, 97%). **MS** (ESI) *m/z* [M+H]⁺ 460.57, [M+Na]⁺ 482.52.

A solution of (*S*)-2-(((1-hydroxy-3-(octadecyloxy)propan-2-yl)oxy)methyl)benzonitrile (360 mg, 0.78 mmol) and triethylamine (340 mg, 3.0 mmol) was added to cooled (0 °C) solution of POCl₃ (140 mg, 2.0 mmol) in CH₂Cl₂ (5 mL) and stirred for 3 h. The mixture was added to ice water, stirred 1h, then extracted with CH₂Cl₂. Evaporation of the organic phase gave phosphate (*R*)-2-((2-cyanobenzyl)oxy)-3-(octadecyloxy)propyl dihydrogen phosphate (850 mg) which was used without further purification. **MS** (ESI) *m/z* [M-H]⁻ 538.46.

The phosphate (460 mg, 0.85 mmol) was coupled to RVn (GS-441524)-acetone (280 mg, 0.85 mmol) using DIC (210 mg, 1.7 mmol), NMI (140 mg, 1.7 mmol) in dry pyridine (15 mL). Reaction at 35 °C for 23 h followed by chromatographic purification yielded ((3*aR*,4*R*,6*R*,6*aR*)-6-(4-aminopyrrolo[2,1-*f*][1,2,4]triazin-7-yl)-6-cyano-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)methyl ((*R*)-2-((2-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate (190 mg, 22 %) **MS** (ESI) *m/z* [M+H]⁺ 851.47.

The coupled acetone (190 mg, 0.22 mmol) was added to formic acid (5 mL) and stirred 3 h. Formic acid was evaporated and the residue purified by flash column chromatography to give ((2*R*,3*S*,4*R*,5*R*)-5-(4-aminopyrrolo[2,1-*f*][1,2,4]triazin-7-yl)-5-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methyl ((*R*)-2-((2-cyanobenzyl)oxy)-3-(octadecyloxy)propyl) hydrogen phosphate as an off white solid (120 mg, 67%). **¹H NMR** (400 MHz, CD₃OD + CDCl₃) δ 7.73 (s, 1H, H2), 7.65 (d, *J* = 7.6, Hz, 2H, -CH₂-aryl-H3 + H5), 7.59 (td, *J* = 7.6, 1.3 Hz, 1H, H4), 7.39 (td, *J* = 7.6, 1.3 Hz, 1H, H6), 6.97 (d, *J* = 4.6 Hz, 1H, H9), 6.86 (d, *J* = 4.5 Hz, 1H, H8), 4.81 (d, *J* = 5.7 Hz, 1H, H2'), 4.38 (q, *J* = 4.2 Hz, 1H, H4'), 4.28 (t, *J* = 5.3 Hz, 1H, H3'), 4.19 – 3.84 (m, 2H, H5'), 3.93 (tp, *J* = 10.7, 4.7 Hz, 2H, H3''), 3.85 – 3.70 (m, 1H, H2''), 3.54 (qd, *J* = 10.6, 5.0 Hz, 2H, H1''), 3.41 (td, *J* = 6.6, 2.7 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.52 (p, *J* = 6.6 Hz, 2H, -OCH₂CH₂(CH₂)₁₅-), 1.26 (m, *J* = 7.2 Hz, 30H, -OCH₂CH₂(CH₂)₁₅-), 0.89 (t, 3H, -CH₃). **¹³C NMR** (101 MHz, CD₃OD + CDCl₃) δ 156.53 (C6), 147.49 (C2), 142.96 (-CH₂-aryl-C1), 133.62, (-

- CH₂-aryl-C3), 133.23, (-CH₂-aryl-C6), 129.66, (-CH₂-aryl-C5), 128.68, (-CH₂-aryl-C4), 125.15, (C7), 117.85, (C5), 117.26, (-CN), 111.84, (-CN'), 111.58, (C8), 102.35, (C9), 84.75, (C4'), 79.89 (C1'), 78.99 (C2'), 75.45 (C2'), 72.21 (-CH₂-aryl), 71.30 (OCH₂CH₂(CH₂)₁₄-), 71.09 (C1'), 70.45 (C3'), 65.73 (C3'), 65.03 (C5'), 32.51 (-OCH₂CH₂(CH₂)₁₄-), 31.09 – 29.39 (m, -OCH₂CH₂(CH₂)₁₄-), 26.67, (-CH₂CH₂CH₃), 23.21 (-CH₂CH₂CH₃), 14.19 (-CH₃). **HRMS** (ESI) *m/z* [M+H]⁺ calcd for C₄₁H₆₂N₆O₉P 813.4310, found 8813.4302. **HPLC** purity 99.4%.

10 *Example 13d. Antiviral evaluation of cyanobenzyl analogs*

Compound	Calu 3				Huh7.5			
	EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)	SI	EC ₅₀ (μM)	EC ₉₀ (μM)	CC ₅₀ (μM)	SI
OD(4-CN-Bn)G-P-RVn	0.043 ± 0.013	0.082 ± 0.022	60.20	1400	0.024 ± 0.002	0.070 ± 0.010	47.2	1966
OD(3-CN-Bn)G-P-RVn	0.034 ± 0.007	0.065 ± 0.036	61.55	1810	0.024 ± 0.006	0.077 ± 0.017	49.6	2066
OD(2-CN-Bn)G-P-RVn	0.091 ± 0.025	0.186 ± 0.047	*82.8	909	0.0102 ± 0.017	0.281 ± 0.030	>100	>980

Claims:

1. A compound of the following formula or a pharmaceutically acceptable salt thereof:

