

CERTIFICATE OF TRANSLATION

I, SHUSAKU YAMAMOTO, a Patent Attorney, Fifteenth Floor, Crystal Tower, 1-2-27 Shiromi, Chuo-ku, Osaka 540, Japan hereby declare that I am conversant with the English and Japanese languages and that I have read the attached English translation and certify that to the best of my knowledge and belief said translation is a true and accurate translation of PCT International Application No. PCT/JP97/00161, filed on 24th January, 1997 in the name of KANEKA CORPORATION.

Dated this 15th day of July, 1998.

A handwritten signature in black ink, appearing to read 'Shusaku Yamamoto', is written over a horizontal line.

SHUSAKU YAMAMOTO

DESCRIPTION

ADSORBENT FOR IMMUNOGLOBULINS AND COMPLEXES THEREOF,
ADSORPTION METHOD, AND ADSORPTION DEVICE

5

TECHNICAL FIELD

The present invention relates to an adsorbent and
an adsorption device for adsorbing immunoglobulins and/or
10 immunoglobulin complexes, and a method for adsorbing and
removing immunoglobulins and/or immunoglobulin complexes.

BACKGROUND ART

15 In recent years, it has become clear that, in
autoimmune diseases such as rheumatoid arthritis, system-
ic lupus erythematoses, Guillain-Barré syndrome, and
idiopathic thrombocytopenic purpura, or in diseases such
as glomerular nephritis, rejection of transplanted
20 organs, tumor, and infectious disease, immunoglobulins
and/or immunoglobulin complexes present in the blood have
a close relationship with the cause or progress of
diseases and/or phenomena.

25 Under the above-mentioned circumstance, several
adsorbing and removing materials have been utilized in
the hope of preventing the progress of the above-men-
tioned diseases, relieving the conditions, and further-
more promoting healing by specifically adsorbing and
30 removing immunoglobulins and/or immunoglobulin complexes
from the body fluid such as the blood and the plasma.
For example, an immune adsorbent (Japanese Laid-open
Publication No. 62-242628) in which protein A capable of

binding to immunoglobulins is immobilized onto a silica matrix is known as an adsorbent which is highly specific to immunoglobulin G and immunoglobulin complexes thereof. Furthermore, an adsorbent (Japanese Laid-open Publication No. 57-122875) for immunoglobulins and/or immunoglobulin complexes in which a hydrophobic compound is immobilized onto an insoluble carrier has already been clinically applied in Japan.

10 However, the adsorbent which has been used (e.g., the above-mentioned adsorbent for protein A) has disadvantages. Specifically, protein A, which is a functional group, is a heterogenous protein having a molecular weight of about 42,000 daltons derived from *Staphylococcus aureus*, so that when the adsorbent is used in contact with the body fluid, protein A is eluted and its antigenicity may cause side effects. Also there is a problem of stability during sterilization of the adsorbent, therefore sterilization method is limited. Additionally, storage stability is not sufficient after protein A is immobilized onto a silica matrix; and the like. Furthermore, the adsorbent for immunoglobulins and/or immune complexes thereof described in Japanese Laid-open Publication No. 57-122875 has the disadvantages of poor adsorbing characteristics such as adsorption specificity to a substance of interest and adsorbing capacity (I. Amano et al., "Blood purification therapy, 1st part", Japanese Journal of Clinical Medicine, Vol. 49, pp. 649-654 (1991 sup.)). Accordingly, in terms of the safety, stability, adsorbing specificity, adsorbing capacity, and the like, the conventionally known adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes is not necessarily suitable for use in treating diseases

caused by the presence of the above-mentioned pathogenic immunoglobulins and/or immunoglobulin complexes in the body fluid.

5

DISCLOSURE OF THE INVENTION

The objective of the present invention is to provide an adsorbent with high stability and safety which has high adsorbing specificity to immunoglobulins and/or immunoglobulin complexes and whose adsorbing characteristics decrease less during sterilization or storage, an adsorption device using the adsorbent, and a method for adsorbing and removing immunoglobulins and/or immunoglobulin complexes.

15

The inventors of the present invention considered the factors that (1) various proteins are known to be immunoglobulin binding proteins, such as protein A, protein G, protein H, protein L, and a rheumatoid factor (N. Tsuchiya, *Clinical Immunology*, Vol. 23, pp. 896-903 (1991); P. Akesson et al., *Mol. Immunol.*, Vol. 27, pp. 523-531 (1990); and L. Bjorck, *J. Immunol.*, pp. 1194-1197 (1988)) have substantially high adsorbing specificity to immunoglobulins and/or immunoglobulin complexes, (2) a peptide having tens of amino acid residues or about ten or less amino acid residues has antigenicity to a living organism which is much smaller than that of a heterogeneous naturally-occurring macromolecule protein, and (3) such a short peptide has high stability which is represented by thermal stability, particularly high stability to sterilization and storage stability. We generated various peptides from immunoglobulin binding proteins and studied them. As a result, we succeeded in

30

obtaining a peptide which has high thermal stability, drug stability, and/or sterilization stability, when determined based on high adsorbing specificity and adsorbing capacity with respect to immunoglobulins and/or immunoglobulin complexes as indexes, and which is unlikely to have side effects caused by antigenicity, by using a partial peptide of protein G.

Protein G is one of the immunoglobulin binding proteins, which has a molecular weight of about 50,000 daltons and is present on a bacteria cell wall of Streptococcus. The cDNA of protein G has already been cloned. Based on an amino acid sequence predicted from the cDNA, protein G is assumed to consist of 448 amino acid residues. Furthermore, it has been assumed that several similar repeated amino acid sequences are present in the predicted amino acid sequence, and that the repeated amino acid sequences are binding domains to albumin and immunoglobulins. See, for example, S.R. Fahnesyook et al., J. Bacteriol., Vol. 167, pp. 870-880 (1986); U. Sjobring et al., J. Biol. Chem., Vol. 266, pp. 399-405 (1991). In B. Guss et al., EMBO J., Vol. 5, pp. 1567-1575 (1986), a repeated amino acid sequence region represented by C1, C2, or C3 suggesting an immunoglobulin (IgG) binding domain in the sequence of protein G is described. When common amino acids and amino acids which are not common in each amino acid sequence of C1, C2, and C3 are summarized, the amino acid sequence of the IgG binding domain is as represented by SEQ ID NO: 1 of Sequence Listing.

Furthermore, the inventors also studied sterilization stability of C3 peptides to γ -ray sterilization



by binding a C3 peptide, which is one of the above-mentioned peptides, to a carrier, and found sterilization stability of C3 peptides was not necessarily sufficient. Then, we generated various peptides and studied them for
5 sterilization stability by successively modifying any amino acid of a C3 peptide into another amino acid by using a genetic engineering technique. As a result, we obtained a peptide which has high affinity for immunoglobulins and/or immunoglobulin complexes, in which
10 the adsorbing capacity of a peptide ligand is hardly damaged by γ -ray sterilization, and which has excellent thermal stability and drug stability, thereby achieving the present invention.

15 In an adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes of the present invention, a peptide capable of adsorbing immunoglobulins and/or immunoglobulin complexes is immobilized onto a water-insoluble carrier, and the peptide has one or two kinds
20 of amino acid sequences selected from the amino acid sequences represented by SEQ ID NO:1 of Sequence Listing.

In an adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes of the present invention,
25 a peptide derivative is immobilized onto a water-insoluble carrier, and the peptide derivative includes at least one of deletion, substitution, insertion, and addition of an amino acid with respect to the amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing, and is
30 capable of adsorbing immunoglobulins and/or immunoglobulin complexes.

In a preferred embodiment, at least one of the

properties of thermal stability, drug stability, γ -ray sterilization stability, and high-pressure vapor sterilization stability of the above-mentioned peptide derivative is improved over that of a peptide having the amino acid sequence represented by SEQ ID NO: 1.

In a preferred embodiment, the above-mentioned peptide derivative includes the amino acid sequence represented by SEQ ID NO: 3.

In a preferred embodiment, (Lys) n or (Cys) m is added to an amino terminus and/or a carboxyl terminus of the above-mentioned peptide or peptide derivative, wherein n and m are integers of 0 to 9.

In a preferred embodiment, the above-mentioned peptide or peptide derivative consists of an amino acid sequence of 70 amino acids or less.

In a preferred embodiment, the adsorbent is subjected to high-pressure vapor sterilization.

In a preferred embodiment, the adsorbent is subjected to γ -ray sterilization.

In a preferred embodiment, the above-mentioned water-insoluble carrier is porous.

In a preferred embodiment, the above-mentioned water-insoluble carrier is hydrophilic.

In a preferred embodiment, a molecular weight of the exclusion limit of the above-mentioned water-insolu-

ble carrier is in the range of about 150,000 to about 5,000,000.

5 In a preferred embodiment, the adsorbent of the present invention is used for adsorbing and removing immunoglobulins and/or immunoglobulin complexes present in the blood, plasma, or other body fluids obtained from a living organism.

10 A method for adsorbing and removing immunoglobulins and/or immunoglobulin complexes of the present invention includes the step of bringing any of the above-mentioned adsorbent into contact with a solution containing immunoglobulins and/or immunoglobulin complexes.

15 In a device for adsorbing and removing immunoglobulins and/or immunoglobulin complexes of the present invention, any of the above-mentioned adsorbents is accommodated in a vessel having an inlet and an outlet
20 for a solution, and the device comprises effluent preventing means for preventing the adsorbent from flowing out of the vessel.

25 In a preferred embodiment, the above-mentioned solution containing immunoglobulins and/or immunoglobulin complexes is the blood, plasma, or other body fluids obtained from a living organism.

BRIEF DESCRIPTION OF THE DRAWINGS

30

Figure 1 is a schematic diagram of plasmid pUCNTMK3P47 for expressing MK3P47 peptide. In a vector for expressing other peptides described in the examples,

DNA encoding MK3P47 peptide inserted between NdeI-HindIII of pUCNTMK3P47 is replaced by DNA encoding other peptides described in each example.

5 Figure 2 shows denaturation curves of MG56, MK3G56, and MK3P47 peptides obtained by using guanidine hydrochloride.

10 Figure 3 shows denaturation curves of MK3P47, MP47K3, MP47, and MP47C peptides obtained by using guanidine hydrochloride.

15 Figure 4 shows excess heat capacity curves of MK3G56 peptide obtained by thermal denaturation with DASM-4.

20 Figure 5 shows excess heat capacity curves of MK3P47 peptide obtained by thermal denaturation with DASM-4.

Figure 6 is a schematic cross-sectional diagram of one example of a device for adsorbing immunoglobulins and/or immunoglobulin complexes of the present invention.

25 Figure 7 is a graph showing results obtained by investigating the relationship between the flow rate and the pressure loss, by using three kinds of gels.

BEST MODE FOR CARRYING OUT THE INVENTION

30

I. Definition

In the present specification, various kinds of

amino acid residues are abbreviated as follows: Ala; L-alanine residue, Asp; L-aspartic acid residue, Asn; L-asparagine residue, Cys; L-cysteine residue, Gln; L-glutamine residue, Glu; L-glutamic acid residue, Gly; L-glycine residue, Ile; L-isoleucine residue, Leu; L-leucine residue, Lys; L-lysine residue, Phe; L-phenylalanine residue, Thr; L-threonine residue, Trp; L-tryptophan residue, Tyr; L-tyrosine residue, and Val; L-valine residue. Furthermore, in the present specification, an amino acid sequence of a peptide is described in such a general manner that the amino terminus of the peptide (hereinafter, referred to as the "N-terminus") is positioned on the left side, and the carboxyl terminus of the peptide (hereinafter, referred to as the "C-terminus") is positioned on the right side.

In the present specification, a peptide derivative refers to a peptide containing at least one of deletion, substitution, insertion, or addition of an amino acid in the amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing.

In the present specification, thermal stability refers to the level of a thermal denaturation temperature of a peptide derivative. The term "having excellent thermal stability" refers to that a thermal denaturation temperature of a peptide derivative is higher than that of a peptide consisting of the amino acids represented by SEQ ID NO: 1. As an example of a method for measuring a thermal denaturation temperature, there is a method for measuring a thermal denaturation temperature in an aqueous solution such as saline by using an adiabatic type differential scanning calorimeter.

In the present specification, drug stability refers to a denaturation ratio of a peptide after drug treatment with respect to the peptide which is not subjected to drug treatment. The term "having excellent drug stability" refers to that a denaturation ratio of a peptide derivative is lower than that of a peptide consisting of the amino acids represented by SEQ ID NO: 1. As an example of a method for determining a denaturation ratio of a peptide, there is a method for comparing the fluorescent intensity of a peptide measured after the treatment in an aqueous solution containing an appropriate concentration of guanidine hydrochloride at 30°C for 10 minutes with the fluorescent intensity of a peptide before the treatment.

15 In the present specification, a γ -ray sterilization method refers to a method for killing microorganisms by irradiating them with γ -rays from a radiation source containing radioactive isotopes described in (i) Radiation Methods: An irradiation method as a "Sterilization method", described in Japanese pharmacopoeia. As an example of the γ -ray sterilization method, there is a method for irradiating each adsorbent in an aqueous solution by using a radiation source containing cobalt 60 and the like. γ -ray sterilization stability refers to a ratio of the capacity of a peptide derivative subjected to γ -ray sterilization to adsorb immunoglobulins, with respect to the capacity of the peptide not subjected to γ -ray sterilization to adsorb immunoglobulins. The term "having excellent γ -ray sterilization stability" refers to that a remaining ratio of the adsorbing capacity of a peptide derivative is higher than that of a peptide consisting of the amino acids represented by SEQ ID NO:

1.

In the present specification, a high-pressure vapor sterilization method refers to a method for killing microorganisms by heating them in saturated steam at an appropriate temperature under an appropriate pressure described in (iii) High-pressure Vapor Methods: A heating method as a "Sterilization method", described in Japanese pharmacopoeia. As an example of the high-pressure vapor sterilization method, there is a method for subjecting an adsorbent in an aqueous solution to an autoclave treatment at 121°C for 20 minutes. High-pressure vapor sterilization stability refers to a ratio of the capacity of a peptide derivative before the high-pressure vapor sterilization treatment to adsorb immunoglobulins, with respect to the capacity of the peptide derivative subjected to the treatment to adsorb immunoglobulins. The term "having excellent high-pressure vapor sterilization stability" refers to that a remaining ratio of the adsorbing capacity of a peptide derivative is higher than that of a peptide consisting of the amino acids represented by SEQ ID NO: 1

II. Preferred Embodiment

Hereinafter, the present invention will be described by way of preferred embodiments. It should be noted that the present invention is not limited by the following description.

As a peptide used for an adsorbent of the present invention, a peptide having the amino acid sequence represented by SEQ ID NO: 3 of Sequence Listing is

preferably used. The most important features in this amino acid sequence region are that the C-terminus sequence (amino acid sequence from position 47 to position 56 of the peptide represented by SEQ ID NO: 1) contained in C1, C2, and C3 of protein G is

Asp Ala Thr Lys Thr Phe Thr Val Thr Glu,

whereas corresponding sequence in the amino acid sequence represented by SEQ ID NO: 3 is

Pro Ala Thr Lys Thr Phe Thr Val Thr Glu.

More specifically, at position 47, an amino acid (Asp) is replaced by another amino acid (Pro). It was found that due to this amino acid substitutions, the sterilization stability (in particular, stability with respect to γ -ray sterilization) of the peptide represented by SEQ ID NO: 3 remarkably improves, and the drug stability based on a denaturation concentration by guanidine hydrochloride as an index and the thermal stability based on a thermal denaturation point measured by calorimetry as an index also improves. Furthermore, the pH stability of the peptide represented by SEQ ID NO: 3 was excellent. Recently, it is common to replace an amino acid by a genetic engineering technique, and thus, it is a well-known technique to study the improvement of characteristics of a peptide and a protein by amino acid substitution. However, there have been no examples showing that the sterilization stability of a peptide or a protein remarkably improved by replacing only one amino acid by another. Thus, the above-mentioned fact is a surprising finding.

An adsorbing functional group peptide used for the adsorbent of the present invention has one or two kinds of amino acid sequences selected from the amino

acid sequences represented by SEQ ID NO: 1 of Sequence Listing, and may contain the same kind of repeated amino acid sequences or a combination of different amino acid sequences. For example, a peptide selected from the amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing may contain only one kind of peptide selected from the peptides corresponding to C1, C2, and C3 in the amino acid sequence of protein G described in B. Guss et al., EMBO J., Vol. 5, pp. 1567-1575 (1986). Alternatively, the peptide may contain one selected kind of repeated peptides (i.e., a peptide containing at least two C1's, at least two C2's, or at least two C3's), or may contain two selected kinds of amino acid sequences (i.e., a peptide containing C1 and C2, C1 and C3, or C2 and C3). Furthermore, the peptide may contain two selected kinds of amino acid sequences arbitrarily repeated (e.g., a peptide containing two C1's and one C2, two C1's and one C3, one C2 and two C3's, two C1's and two C3's, or the like). The peptide may consist of substantially only the amino acid sequence represented by SEQ ID NO: 1 of Sequence Listing, or may further contain any amino acid residue or amino acid sequence at the N-terminus and/or the C-terminus of the amino acid sequence.

25

An adsorbing function group peptide derivative used for the adsorbent of the present invention is a peptide having at least one kind of amino acid sequence containing at least one of deletion, substitution, insertion, and/or addition of an amino acid with respect to the amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing, which may consist of only one kind of amino acid sequence or the combination of at least two

30

kinds of amino acid sequences. In the case of addition, any amino acid residue or amino acid sequence may be placed at the N-terminus and/or the C-terminus of the amino acid sequence. A preferred peptide derivative has
5 an amino acid sequence in which (Lys) n or (Cys) m is added to the N-terminus and/or the C-terminus with respect to at least one kind of amino acid sequence selected from the amino acid sequences represented by SEQ ID NO: 3 (herein, n and m are integers of 0 to 9). In particular,
10 a peptide derivative having (Cys) m at the C-terminus of the amino acid sequence is more preferable.

Such a peptide derivative can have the capacity to adsorb immunoglobulins and/or immunoglobulin complexes
15 in the body fluids, and more preferably has the excellent adsorbing capacity after the high-pressure vapor sterilization treatment, heat treatment, drug treatment, or γ -ray sterilization treatment, compared with the peptides represented by SEQ ID NO: 1.

20

The combination of the above-mentioned peptide and peptide derivative can be used for the adsorbent of the present invention.

25

The peptide or the peptide derivative used for the adsorbent of the present invention may be produced by a generally used peptide synthesis method, (e.g., a solid phase synthesis method, a liquid phase synthesis method
such as a stepwise elongation method, and a fragment
condensation method), or by a gene manipulation method
30 including the steps of connecting DNA encoding a peptide or a peptide derivative to a vector and introducing the vector into a host to produce a recombinant peptide or a

recombinant peptide derivative.

5 In chemical synthesis of a peptide or a peptide
derivative, it makes manipulation easier to apply a solid
phase synthesis method rather than a liquid phase synthe-
sis method. According to the solid phase synthesis
method, for example, a peptide or a peptide derivative
can be synthesized as follows. (1) An amino acid
corresponding to the C-terminus of a peptide to be
10 synthesized is bound to a support insoluble in an organic
solvent. (2) Each corresponding amino acid protecting a
functional group such as an α -amino group other than an
 α -carboxy group is bound to the amino acid binding to the
substrate successively toward the N-terminus direction by
15 condensation reaction. (3) A protective group of the α -
amino group of the bound amino acid is removed. (4) The
steps (1) to (3) are alternately repeated. As a result
of these steps of manipulation, a peptide chain is
elongated to obtain a peptide with a desired length.

20

After the peptide of interest is obtained, a
peptide chain is cleaved from the support, and protective
groups are removed. For this purpose, hydrogen fluoride
is often used. However, in terms of safety and ease of
25 handling, it is appropriate to use trifluoroacetic acid
(hereinafter, referred to as "TFA"). More specifically,
the peptide of interest is reacted in 95% TFA containing
1,2-ethanedithiol and anisole (1:3), whereby the peptide
is cleaved from the substrate and the protecting groups
30 are removed. Thus, a crude peptide is collected. The
crude peptide is precipitated with ethyl ether, whereby
a crude purified peptide is obtained.

The crude purified peptide thus obtained can be purified by various methods known to those skilled in the art. For example, the crude purified peptide is subjected to high performance liquid chromatography (hereinafter, referred to as "HPLC") using a reverse phase system column, whereby the peptide can be fractionated, collected, and purified. HPLC conditions should be optimized based on a system used generally for purifying a protein. A fraction corresponding to the peak of chromatography thus obtained, and lyophilized. The resultant purified peptide fraction is subjected to primary sequence analysis by an amino acid sequencer and amino acid composition analysis, whereby a synthesized peptide is confirmed.

In the case where a recombinant peptide or a recombinant peptide derivative is produced by a gene recombination method, based on an amino acid sequence of a peptide of interest, DNA encoding the amino acid sequence is synthesized and introduced into a phage or a plasmid vector. The resultant phage or plasmid vector is integrated into an appropriate microorganism (e.g., E. coli) to select and obtain a transformant, and the transformant is cultured by a known method. It is well within the skill of those in the art to purify and isolate a peptide of interest from this culture supernatant or cells. Furthermore, as long as an appropriate vector is selected, yeast, *Bacillus subtilis*, and the like can be easily utilized as a host.

A water-insoluble carrier used as a support of the adsorbent of the present invention is not particularly limited. Examples of the water-insoluble carrier used in the present invention include inorganic carriers such

as glass beads and silica gels; organic carriers made of synthetic macromolecules such as cross-linked polyvinyl alcohol, cross-linked polyacrylate, cross-linked polyacrylamide, and cross-linked polystyrene or polysaccharides such as crystalline cellulose, cross-linked cellulose, cross-linked agarose, and cross-linked dextran; and complexes carriers obtained by combining these organic carriers and inorganic carriers, such as a combination of two kinds of organic carriers and a combination of an organic carrier and an inorganic carrier. In the case of using a carrier for hemocatharsis, it is necessary to consider effects of a carrier on a complement system, a clotting system, and the like, when the carrier comes into contact with blood components. Therefore, hydrophilic carriers, which have relatively less possibility of non-specific adsorption and have satisfactory adsorbing selectivity with respect to immunoglobulins and/or immunoglobulin complexes, are preferably used. In the present specification, a hydrophilic carrier refers to a carrier in which, when a compound forming a carrier is formed into a flat plate, a contact angle between the flat plate and the water becomes about 60° or less. Examples of such a carrier include, but are not limited to, carriers formed of polysaccharides such as cellulose, chitosan, Sepharose and dextran, polyvinyl alcohol, a saponified ethylene-vinyl acetate copolymer, polyacrylamide, polyacrylic acid, polymethacrylic acid, poly(methyl methacrylate), polyacrylic acid-grafted polyethylene, polyacrylamide-grafted polyethylene, and glass, or carriers formed of derivatives thereof.

Among the above-mentioned water-insoluble carriers, those having an -OH group have excellent adsorbing

capacity and adsorbing selectivity. Above all, a porous cellulose gel has the following excellent points:

5 (1) The porous cellulose gel is stiff because of relatively high mechanical strength, so that it is unlikely to be damaged or generate fine powders due to stirring. Thus, when the body fluid is passed through a column packed with the gel at a high speed, the gel is unlikely to cause compaction or clogging in the column.
10 Furthermore, the porous structure of the gel is unlikely to change due to high-pressure vapor sterilization, and the like.

15 (2) Being formed of cellulose, the porous cellulose gel is hydrophilic. Since a number of hydroxyl groups available to binding of a ligand are present on the gel, there is little possibility of non-specific adsorption.

20 (3) Since the porous cellulose gel keeps its relatively high strength even in the case where a pore area is enlarged, an adsorbing capacity which is comparable to a soft gel is obtained.

25 (4) The porous cellulose gel has higher safety factor than that of a synthetic macromolecular gel.

30 Thus, the porous cellulose gel is one of those most suitable for use in the present invention. The carrier used in the present invention is not limited to those described above. The above-mentioned carriers may be used singly, or any two or more kinds thereof may be mixed.

A primary property required of the water-insoluble carrier used in the present invention is that the carrier has a number of pores with an appropriate size, i.e., the carrier is porous. Immunoglobulins and/or immunoglobulin complexes which are to be adsorbed by the adsorbent of the present invention have a molecular weight in a wide range of about 160,000 to about 1,000,000, and thus, they cannot be specified. Therefore, in order for the adsorbent to efficiently adsorb a protein, it is preferable that immunoglobulins and/or immunoglobulin complexes can enter pores with a high probability to some degree, and other proteins are prevented from entering pores as much as possible.

In the water-insoluble carrier used for the present invention, an average pore diameter of a porous carrier is preferably about 300 to about 10000 Å. The diameter of a pore is most often measured by a method of mercury porosimetry. However, in the case of a porous water-insoluble carrier used in the present invention, this method often could not be applied. As a criterion of the diameter of a pore, a molecular weight of the exclusion limit is preferably used. A molecular weight of the exclusion limit refers to a molecular weight of a molecule having the smallest molecular weight among molecules which cannot enter pores (i.e., which are excluded) in gel permeation chromatography (H. Hatano and T. Hanai, Experimental High Performance liquid Chromatography (Jikken Kosoku Ekitai Chromatography), Kagaku Dojin, 1988. A molecular weight of the exclusion limit is generally well investigated for globular proteins, dextran, polyethylene glycol, and the like. In the case of the porous water-insoluble carrier used in the present

invention, a value obtained by using a globular protein is suitably used.

As a result of studying carriers having various
5 molecular weights of the exclusion limit, the exclusion
limit of a suitable carrier, which therefore has a pore
diameter suitable for adsorbing immunoglobulins and/or
immunoglobulin complexes, was found to be about 150,000
or more in molecular weight. More specifically, in the
10 case where a carrier having a molecular weight of the
exclusion limit of less than about 150,000 is used, an
adsorbing and removing amount of immunoglobulins and/or
immunoglobulin complexes is small, resulting in decreased
practicability. Thus, a preferable molecular weight of
15 the exclusion limit of the carrier used in the present
invention is about 150,000 or more. On the other hand,
in the case where a molecular weight of the exclusion
limit exceeds about 5,000,000, no particular problems are
caused as long as the plasma or the serum is used as the
20 body fluid. However, macromolecules which have no
interaction with a ligand have a tendency to block a
binding site of a ligand, thereby substantially decreas-
ing the effective ligand amount. Furthermore, in the
case where the blood is used as the body fluid, when a
25 molecular weight of the exclusion limit exceeds about
5,000,000, there is a tendency that a ratio at which the
platelet adheres to a carrier increases. Thus, in the
case where the adsorbent of the present invention is used
in a hemocatharsis system of DHP (direct hemoperfusion)
30 type, sufficient performance cannot necessarily be
exhibited. On the other hand, in the case where a
molecular weight of the exclusion limit is about
5,000,000 or less, no particular problems are caused even

when carriers are used for any purpose. Thus, in order to use the adsorbent of the present invention for various purposes, a molecular weight of the exclusion limit is desirably about 5,000,000 or less. Accordingly, a more
5 preferable molecular weight of the exclusion limit of the carrier used in the present invention is from about 150,000 to about 5,000,000.

Next, considering the adsorbing capacity per
10 volume of an adsorbent, it is preferable that a carrier has a totally porous structure, rather than a surface porous structure, and a pore volume of 20% or more and a specific surface area of 3 m²/g or more. As a shape of a carrier, any shape such as a bead, a fiber, and/or a
15 membrane (including a hollow thread) can be selected.

In order to immobilize a ligand, it is preferable that functional groups available for an immobilization reaction of a ligand are present on the surface of a
20 carrier. Examples of the functional groups include a hydroxyl group, an amino group, an aldehyde group, a carboxyl group, a thiol group, a silanol group, an amide group, an epoxy group, a halogen group, a succinylimide group, and an acid anhydride group.

25 Next, as the carrier used in the present invention, any hard carrier and/or soft carrier can be used. In order to use a carrier as an adsorbent for an extracorporeal circulation treatment, it is important
30 that when the carrier is packed into a column and the body fluid is passed through the column, the carrier does not cause clogging. Therefore, sufficient mechanical strength is required of the carrier. Thus, the carrier

used in the present invention is preferably a hard carrier. In the present specification, a hard carrier refers to those in which, for example, in the case of a granular gel, where the gel is uniformly packed into a cylindrical column made of glass (inner diameter: 9 mm, column length: 150 mm) under the following conditions, and an aqueous fluid is passed through the column, the pressure loss ΔP and the flow rate of the aqueous fluid have a linear relationship up to a pressure of 0.3 kg/cm².

10

For example, an agarose gel (Biogel A-5m produced by Bio-Rad; particle diameter: 50 to 100 meshes), a vinyl type polymer gel (Toyopearl HW-65 produced by Toyo Soda Mfg. Co., Ltd.; particle diameter: 50 to 100 μ m), and a cellulose gel (Cellulofine GC-700m produced by Chisso Corporation; particle diameter: 45 to 105 μ m) were each uniformly packed into their own cylindrical column made of glass (inner diameter 9 mm; column length 150 mm) equipped with filters having a pore diameter of 15 μ m at both ends. Water was passed through the column by a peristaltic pump, and the relationship between the flow rate and the pressure loss ΔP was obtained (Figure 7). The flow rate (cm/min.) was plotted on an ordinate and the pressure loss (kg/cm²) was plotted on an abscissa. As can be seen in Figure 7, $\bigcirc$, $\triangle$, and $\bullet$ represent Toyopearl HW-65, Cellulofine GC700m, and Biogel-A5m, respectively. As a result, it is understood that the flow rate of the aqueous fluid through Toyopearl HW-65 and Cellulofine GC-700m increases almost in proportion to the increase in a pressure, whereas Biogel A-5m results in compaction, and the flow rate does not increase with the increase in a pressure.

30

For immobilizing a peptide or a peptide derivative onto the above-mentioned carrier, in order to improve an adsorbing efficiently by minimizing a steric hindrance of a peptide or a peptide derivative and to suppress non-specific adsorption, a peptide or a peptide derivative is preferably immobilized onto a carrier via hydrophilic spacers. As the hydrophilic spacers, for example, derivatives of polyalkylene oxide in which both ends are replaced by carboxyl groups, amino groups, aldehyde groups, epoxy groups, or the like are preferably used.

Since an organic compound used as a peptide or a peptide derivative, and spacers introduced into the above-mentioned carrier are a relatively stable substance with a low molecular weight, there is a less stringent requirement for the immobilization reaction conditions, for example, compared with the case where a protein such as an enzyme and an antibody is immobilized. Thus, an immobilization method is not particularly limited. However, considering that an adsorbent may be used for extracorporeal circulation treatment and hemocatharsis, it is more preferable to apply an immobilization method in which peptides are not easily detached from an carrier during sterilization of the adsorbent or treatment. For example, there are (1) a method for allowing a carboxyl group of a carrier to be reacted with N-hydroxysuccinimide to replace the carboxyl group into a succinimideoxycarbonyl group, and allowing a peptide or a peptide derivative to be reacted at a portion of an amino group (active ester method), (2) a method for allowing a carboxyl group or an amino group of a peptide or a peptide derivative to be condensation-reacted with

an amino group or a carboxyl group of a carrier in the presence of a condensation reagent such as dicyclohexylcarbodiimide (condensation method), and (3) a method for cross-linking a peptide or a peptide derivative to a carrier by using a compound having at least two functional groups such as glutaraldehyde (carrier cross-linking method). In order to reduce desorption and elution of peptides to a minimum, peptides are preferably bound to a carrier by a covalent bonding method.

10

There are various methods for adsorbing and removing immunoglobulins and/or immunoglobulin complexes in the body fluid by bringing a carrier on which a peptide functional group is immobilized into contact with the body fluid such as the blood and the plasma. Representative examples include (1) a method including the steps of taking out the body fluid, storing it in a bag, mixing the body fluid with an adsorbent to adsorb and remove immunoglobulins and/or immunoglobulin complexes, and filtering off the adsorbent, thereby obtaining body fluid from which immunoglobulins and/or immunoglobulin complexes has been removed, and (2) a method for charging an adsorbent into a vessel having an inlet and an outlet for the body fluid and equipped with a filter at the outlet, which passes the body fluid but does not pass the adsorbent, and flowing the body fluid through the vessel, and the like. Any method for adsorbing and removing immunoglobulins and/or immunoglobulin complexes may be used. However, the latter method is easy to operate, and furthermore, the vessel can be incorporated into an extracorporeal circulation circuit, whereby immunoglobulins and/or immunoglobulin complexes can be removed efficiently from the body fluid of a patient on line. Thus,

30

the adsorbent of the present invention is suitable for this method.

5 Next, a device for adsorbing immunoglobulins and/or immunoglobulin complexes of the present invention using an adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes will be described with reference to its schematic cross-sectional view.

10 A vessel 7 shown in Figure 6 includes an inlet or outlet 1 for a solution, an outlet or inlet 2 for a solution, an immunoglobulin and/or immunoglobulin complex adsorbent 3 of the present invention, effluent preventing means 4 and 5 for preventing the out-flow of the adsorbent, which allows a solution and a component contained
15 therein to pass therethrough but does not allow the adsorbent to pass therethrough, and a column 6. The shape and material for this vessel are not particularly limited. However, for example, a cylindrical vessel with
20 a capacity of about 20 to about 400 ml and a diameter of about 2 to about 10 cm is preferably used.

25 The present invention will be described in more detail by way of the following examples. It should be noted that the present invention is not limited to the following examples.

Examples

30

Example 1

Chemical synthesis of a peptide and
immobilization of the peptide onto Sepharose

Chemical synthesis of a peptide

Synthesis of a peptide having a sequence consisting of 57 residues of domain C3 of protein G with cysteine added to an N-terminus was conducted as follows by a solid phase synthesis method, using a Peptide Synthesizer Model 4170 (manufactured by Pharmacia LKB). Using 0.1 mmol of Fmoc-glutamine NovaSyn KA (produced by Pharmacia LKB) which is a support to which glutamine corresponding to the C-terminus is bound, a deprotection reaction and a condensation reaction are repeated in accordance with a synthesis program input to the Peptide Synthesizer, whereby a peptide chain was successively elongated by addition to the N-terminus. More specifically, the following process steps were repeated: A 9-fluorenylmethyloxycarbonyl group (hereinafter, referred to as "Fmoc") which is a protecting group of an α -amino group of the amino acid is removed by piperidine, the peptide chain was washed with N,N-dimethylformamide (hereinafter, referred to as "DMF"), a condensation reaction was conducted by using 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluroniumtetrafluorophosphate (hereinafter, referred to as "HBTU" and N,N-diisopropylethylamine (hereinafter, referred to as "DIPEA"), and the peptide chain was washed with DMF. As amino acids, Fmoc-L-Ala, Fmoc-L-Asn(Trt), Fmoc-L-Asp(OtBu), Fmoc-L-Cys(Trt), Fmoc-L-Gln(Trt), Fmoc-L-Glu(OtBu), Fmoc-L-Gly, Fmoc-L-Ile, Fmoc-L-Leu, Fmoc-L-Lys(Boc), Fmoc-L-Phe, Fmoc-L-Thr(tBu), Fmoc-L-Trp, Fmoc-L-Tyr(tBu), and Fmoc-L-Val were used. These amino acids (0.5 mmol) were used in a vial. The amount of amino acids used is about 5 times the amount of moles of a substrate which is used. (Herein, Trt, OtBu, Boc, and tBu represent a trityl group, an o-tertiary butyl ester, a tertiary butyl oxycarbonyl

group, and a tertiary butyl group, respectively.)

Deprotection reaction and cleaving off of a peptide chain

After a reaction operation was completed for all
5 the amino acids, the resultant support was washed with
tert-amyl alcohol, acetic acid, and diethyl ether,
successively, on a glass filter with a 3G-3 pore. Then,
the support was dried in a vacuum, whereby a dry support
was obtained. In a vial, 20 ml of TFA, 260 μ l of 1,2-
10 ethanedithiol, and 780 μ l of anisole were added to 1 g of
the obtained support, and the mixture was stirred at room
temperature for 1.5 hours. Thereafter, the mixture was
separated off the support by filtration through the glass
filter with a 3G-3 pore. The resultant filtrate was con-
15 densed at 35°C under reduced pressure. To the condensed
filtrate, previously cooled absolute diethyl ether was
added, and the resultant mixture was stirred until no
absolute diethyl ether is precipitated. Then, the
mixture was centrifuged, and a crude peptide precipitate
20 was collected. Furthermore, the crude peptide was washed
several times with absolute diethyl ether, and thereaf-
ter, dried under a reduced pressured, whereby a crude
purified peptide of interest was obtained.

25 Purification of a peptide

The crude purified peptide thus obtained was dis-
solved in 0.1% TFA, and the solution was filtered through
a 0.2 μ m membrane filter. The filtrate thus obtained was
subjected to high-performance liquid chromatography
30 (HPLC). For HPLC, a Model LC-10A system (manufactured by
Shimadzu Corporation) was used, and as a column, a
reverse phase type μ Bondasphere C18 (manufactured by
Nihon Millipore Waters Ltd.) was used. For a mobile

phase, using 0.1% TFA aqueous solution as A solution and 80% (v/v) acetonitrile/water containing 0.1% TFA as B solution, the column was eluted by a linear concentration gradient from A solution to B solution. Fractions corresponding to a peptide at the peak of the obtained chromatogram were collected. Collection was repeated several times, and the fractions were lyophilized, whereby a purified peptide was obtained. The peptide thus obtained was analyzed by gas-phase protein sequencer model 477 (manufactured by Applied Biosystems Co. Ltd.), and amino acid analysis using a Hitachi custom ion exchange resin. Thus, it was confirmed that a peptide having the amino acid sequence represented by ID SEQ NO: 2 was obtained.

15

Immobilization of a peptide onto Sepharose

The above-mentioned peptide was immobilized onto porous Sepharose, whereby an adsorbent was produced as follows. As Sepharose, Thiopropyl Sepharose 6B (produced by Pharmacia LKB) was used. To 50 mg of Thiopropyl Sepharose 6B, 50 ml of distilled water was added, and the mixture was allowed to stand at room temperature for 15 minutes, whereby the resin was allowed to swell. Then, the distilled water was removed and replaced by 0.1 M Tris hydrochloride (pH 7.5) coupling buffer containing 0.5 M of NaCl.

On the other hand, 4 mg of the above-mentioned purified peptide was dissolved in 400 μ l of 0.1 M Tris hydrochloride (pH 7.5) coupling buffer containing 0.5 M of NaCl, and 40 mg of the above-mentioned Thiopropyl Sepharose 6B was added to the resultant solution. The mixture was stirred at 4°C for 12 hours, whereby an

adsorbent with the purified peptide immobilized thereon was obtained. Furthermore, the resultant adsorbent with the peptide immobilized thereon was filtered by suction, and the content of the peptide in the filtrate was quantitated by an absolute calibration curve method, using HPLC. Thus, a ratio of the peptide immobilized onto the carrier was obtained. The adsorbent with the peptide immobilized thereon was thoroughly washed with 10 mM of phosphate buffer (pH 7.2) containing 150 mM of NaCl, and filtered by suction, whereby a peptide-Sepharose adsorbent of interest was obtained.

Example 2

Evaluation of adsorbing capacity of an adsorbent with a peptide immobilized thereon

Test 1

The adsorbent produced in the above-mentioned Example 1 was evaluated for adsorbing capacity of thermally coagulated complexes (model of an immune complex) in the blood, by using the serum of a healthy subject. Fifty microliters (one volume) of the above-mentioned adsorbent was collected in a vial, and 150 μ l (three volumes) of the serum of a healthy subject containing 20 μ g/ml of thermally coagulated complexes (see M. Makino, Clinical Immunology, Vol. 18, pp. 111-119 (1986)) was added to the adsorbent, and the serum and the adsorbent were incubated with shaking at 37°C for 2 hours. Furthermore, as a control, 150 μ l (3 volumes) of the serum of a healthy subject containing 20 μ g/ml of thermally coagulated complexes to 50 μ l (1 volume) of an untreated support, and the serum and the support were

incubated with shaking at 37°C for 2 hours. Thereafter, these suspensions were centrifuged at 5,000 rpm for one minute, and the amount of the complexes in the supernatants were measured by an extracorporeal diagnostic kit "FRELISA Clq-CIC" manufactured by Fujirebio Inc.

The results are shown in Table 1. The adsorption ratio of the thermally coagulated complexes was represented by an adsorption ratio (%) calculated from a value of the control serum by the following equation. Furthermore, each adsorbing capacity of immunoglobulins (IgG, IgA, IgM) was quantitated by an extracorporeal diagnostic medical product "N-Assay TIA" measurement kit commercially available from Nittobo Co., Ltd. The results are shown in Table 2. Furthermore, IgG subclasses (IgG₁, IgG₂, IgG₃, IgG₄) were quantitated by a measurement kit using an immune diffusion (SRID) method commercially available from Bindingsite. The results are shown in Table 3.

20

$$\text{Adsorption removal ratio (\%)} = \frac{\frac{\text{Complex concentration in the control solution} - \text{Complex concentration in the test solution}}{\text{Complex concentration in the control solution}} \times 100}{25}$$

Table 1

5	Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of thermally coagulated complexes (%)
10	non-immobilized thiopropyl carrier	0.0	0.0
	Thiopropyl carrier with a ligand immobilized thereon	0.8	41.7
15			

Table 2

20	Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of immuno- globulins (%)		
			IgG	IgA	IgM
25	Non-immobilized thiopropyl carrier	0.0	0.0	0.0	0.0
	Thiopropyl carrier with a ligand immobilized thereon	0.8	83.4	0.0	2.7
30					

Table 3

5	Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of IgG subclasses(%)			
			IgG ₁	IgG ₂	IgG ₃	IgG ₄
10	Non-immobilized thiopropyl carrier	0.0	0.0	0.0	0.0	0.0
	Thiopropyl carrier with a ligand immobilized thereon	0.8	93.2	78.9	63.2	65.7

15 From the above results, it was discovered that
the use of the adsorbent of the present invention enables
IgG which is a component of immune complexes and immune
complexes to be selectively adsorbed and removed, and
enables IgG₃ which is not adsorbed by protein A to be
20 adsorbed and removed.

Test 2

25 The above-mentioned adsorbent was evaluated for
adsorbing capacity of circulating immune complexes in the
blood by using the serum of a patient suffering from
rheumatism. Fifty microliters (one volume) of the above-
mentioned adsorbent was collected in a vial, and 150 μ l
(three volumes) of the serum of a patient suffering from
rheumatism was added to the adsorbent, and the serum and
30 the adsorbent were incubated with shaking at 37°C for 2
hours. Furthermore, as a control, 150 μ l (3 volumes) of
the serum of a patient suffering from rheumatism was
added to 50 μ l (1 volume) of an untreated support, and

the serum and the support were incubated with shaking at 37°C for 2 hours. Thereafter, these suspensions were centrifuged at 5,000 rpm for one minute, and the amount of the complexes in the supernatant was measured by the method described in Test 1. The results are shown in Table 4 together with the adsorption ratio (%) calculated from a value of the control serum. Furthermore, each adsorbing capacity of immunoglobulins (IgG, IgA, IgM) was quantitated by the above-mentioned method. The results are shown in Table 5.

Table 4

Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of immune complexes (%)
non-immobilized thiopropyl carrier	0.0	0.0
Thiopropyl carrier with a ligand immobilized thereon	0.8	29.3

Table 5

5	Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of immuno- globin (%)		
			IgG	IgA	IgM
10	Non-immobilized thiopropyl carrier	0.0	0.0	0.0	0.0
	Thiopropyl carrier with a ligand immobilized thereon	0.8	21.1	0.0	2.7

15

From Tables 4 and 5, it was discovered that the use of the adsorbent of the present invention enables IgG which is a component of immune complexes in the blood of a patient suffering from rheumatism and immune complexes to be selectively adsorbed and removed.

20

Example 3

High-pressure vapor sterilization stability
of an adsorbent with a peptide immobilized thereon

25

Autoclave treatment of an adsorbent

Fifty microliters of the adsorbent produced in Example 1 was suspended in 1 ml of 10 mM phosphate buffer (pH 7.2) containing 150 mM of NaCl, and the resultant suspension was heated under a high pressure at 121°C for 20 minutes in an autoclave sterilizer.

30

Evaluation of adsorbing capacity of an adsorbent subjected to an autoclave treatment

The serum was suspended in the same way as in Example 2, using the adsorbent obtained in Example 1 and the above-mentioned adsorbent treated by autoclave sterilization. The concentrations of complexes and immunoglobulins in the obtained supernatant were measured to calculate an adsorption and removal ratio. The results are shown in Tables 6 and 7.

Table 6

Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of immune complexes(%)
Non-immobilized thiopropyl carrier	0.0	0.0
Thiopropyl carrier with a ligand immobilized thereon	2.4	78.3
Carrier with a ligand immobilized thereon treated at 121°C for 20 minutes	2.4	12.5

Table 7

5	Adsorbent	Immobilized amount (mg/ml)	Adsorption ratio of immuno- globulin (%)		
			IgG	IgA	IgM
10	Non-immobilized thiopropyl carrier	0.0	0.0	0.0	0.0
	Thiopropyl carrier with a ligand immobilized thereon	2.4	58.8	0.0	1.8
15	Carrier with a ligand immobilized thereon treated at 121°C for 20 minutes	2.4	8.8		

20

From the above-mentioned test results, it was found that the adsorbent with the peptide immobilized thereon of the present invention has its adsorption removal capacity decreased by an autoclave sterilization treatment, while maintaining an adsorption removal capacity with respect to complexes and IgG.

25

Example 4

30

Production of a peptide derivative MK3P47 peptide

DNA (DNA sequence represented by SEQ ID NO: 5) encoding MK3P47 peptide represented by SEQ ID NO: 4 was

synthesized. This DNA is designed so as to be connected by utilizing each restriction enzyme site (NdeI on a 5' side; HindIII on a 3' side) of a pUCNT vector (Japanese Laid-open Publication No. 4-212692).

5

DNA having the above-mentioned sequence was connected to a pUCNT vector cleaved by digestion with restriction enzymes NdeI and HindIII (produced by Takara Shuzo Co., Ltd.) by using DNA Ligation Kit Ver. 2 produced by Takara Shuzo Co., Ltd. in accordance with a procedure, whereby a pUCNTMK3P47 vector (Figure 1) was produced.

The pUCNTMK3P47 vector DNA was introduced into E. coli HB101 strain (available from Funakoshi) by using a known method, and a transformant was selected based on resistance with respect to antibiotic ampicillin as an index. Furthermore, plasmid DNA was extracted from this transformant by a conventional method and determined for its sequence, whereby it was confirmed that the pUCNTMK3P47 vector has a DNA sequence as designed. Next, the transformant was cultured with shaking in 6 L of L-broth (5 g/L of NaCl, 10 g/L of bactotrypsin, 5 g/L of yeast extract) at 37°C for 20 hours, and cultures were centrifuged (at 4°C and 6000 rpm for 20 minutes, using a Hitachi RPR9-2 rotor) to collect the cells. The pellets thus obtained were suspended in 300 ml of a TE buffer (20 mM of Tris-HCl, 1 mM of EDTA: pH 7.5), and the resultant suspension was subjected to ultrasonication disruption (3 times (each for 6 minutes) in the ice, using BRANSON 250). Then, the suspension was centrifuged (at 4°C and 15000 rpm for 20 minutes, using a Hitachi RPR16 rotor), and a supernatant was collected. The supernatant thus

obtained was heated at 70°C for 10 minutes, centrifuged (at 4°C and 15000 rpm for 20 minutes, using a Hitachi RPR16 rotor), and 300 ml of a supernatant was collected. MK3P47 peptide was purified from the supernatant thus
5 obtained, by using HPLC chromatography (column: Waters μ BONDASPHERE 5 μ C18 300 A 19.0 $\times$ 150 mm) as follows. First, 40 ml of 0.1% TFA solution was passed through the column at a flow rate of 5 ml/min to activate the column, and 300 ml of a sample was passed through the column at
10 the same flow rate. The column was washed with 200 ml of 0.1% TFA + 14% acetonitrile solution. Then, MK3P47 peptide derivative of interest was eluted from the column by using 200 ml of 0.1% TFA + 40% acetonitrile solution, whereby fractions containing MK3P47 peptide were collect-
15 ed. The fractions were concentrated to 100 ml by an evaporator and lyophilized, whereby 1.2 g of MK3P47 peptide was collected as a high purity purified sample. This sample was used for various studies.

20

Example 5

Production of a peptide derivative MP47K3 peptide

DNA represented by SEQ ID NO: 7 encoding MP47K3
25 peptide represented by SEQ ID NO: 6 was synthesized by using the method similar to that of Example 4, and a pUCNTMP47K3 vector was produced by using the DNA and a pUCNT vector. Then, the vector was transformed into E. coli. The transformant thus obtained was cultured (6 L),
30 and the MP47K3 peptide of interest was purified, whereby 600 mg of a high purity sample was obtained.

Example 6

Production of a peptide derivative MP47 peptide

5 DNA represented by SEQ ID NO: 9 encoding MP47
peptide represented by SEQ ID NO: 8 was synthesized by
using the method similar to that of Example 4, and a
pUCNTMP47 vector was produced by using the DNA and a
pUCNT vector. Then, the vector was transformed into E.
10 coli. The transformant thus obtained was cultured (6 L),
and the MP47 peptide of interest was purified, whereby
500 mg of a high purity sample was obtained.

Example 7

15

Production of a peptide derivative MP47C peptide

DNA represented by SEQ ID NO: 11 encoding MP47C
peptide represented by SEQ ID NO: 10 was synthesized by
20 using the method similar to that of Example 4, and a
pUCNTMP47 vector was produced by using the DNA and a
pUCNT vector. Then, the vector was transformed into E.
coli. The transformant thus obtained was cultured (6 L),
and the MP47C peptide of interest was purified, whereby
25 1.3 g of a high purity sample was obtained.

Example 8

Production of MK3G56 peptide for comparison

30

DNA represented by SEQ ID NO: 13 encoding MK3G56
peptide represented by SEQ ID NO: 12 was synthesized by
using the method similar to that of Example 4, and a

pUCNTMK3G56 vector was produced by using the DNA and a pUCNT vector. Then, the vector was transformed into E. coli. The transformant thus obtained was cultured (6 L), and the MK3G56 peptide of interest was purified, whereby
5 1.0 g of a high purity sample was obtained.

Example 9

Production of a peptide derivative MK3-7M peptide

10

DNA represented by SEQ ID NO: 15 encoding MK3-7M peptide represented by SEQ ID NO: 14 was synthesized by using the method similar to that of Example 4, and a pUCNTMK3-7M vector was produced by using the DNA and a
15 pUCNT vector. Then, the vector was transformed into E. coli. The transformant thus obtained was cultured (6 L), and the MK3-7M peptide of interest was purified, whereby 800 mg of a high purity sample was obtained.

20

Example 10

Production of MG56 peptide for comparison

DNA represented by SEQ ID NO: 17 encoding MG56
25 peptide represented by SEQ ID NO: 16 was synthesized by using the method similar to that of Example 4, and a pUCNTMG56 vector was produced by using the DNA and a pUCNT vector. Then, the vector was transformed into E. coli. The transformant thus obtained was cultured (6 L),
30 and the MG56 peptide of interest was purified, whereby 400 mg of a high purity sample was obtained.

Example 11

Drug denaturation curve analysis of a peptide

5 First, 1 mg/ml of an aqueous solution of each peptide obtained in Examples 4 to 8 and Example 10 was diluted to 10 µg/ml by 0 to 8 M of a guanidine hydrochloride solution (at 30°C), and a fluorescent intensity at EM 340 nm (excited at EX 278 nm) was measured by F-2000 type spectrofluorometer manufactured by Hitachi, Ltd. The results are shown in Figures 2 and 3. The fluorescent intensity (%) was plotted on an ordinate, and the concentration of guanidine (M) was plotted on an abscissa. In Figure 2, ◇, □, and △ represent a MG56 peptide, MK3G56 peptide, and MK3P47 peptide, respectively. In Figure 3, ◇, □, △, and × represent a MK3P47 peptide, MP47K3 peptide, MP47 peptide, and MP47C peptide, respectively. Consequently, as shown in Figures 2 and 3, when the height of denaturation concentration of each peptide was compared, a relationship MP47C > MP47 > MP47K3 > MK3P47 > MG56 > MK3G56 was obtained.

Example 12

25 Measurement of a thermal
denaturation point of a peptide

30 MK3G56 peptide and MK3P47 peptide were measured for a thermal denaturation point by an adiabatic type differential scanning calorimeter DASM-4 (manufactured by Mashpriborintorg in Russia: Japanese agent, Shinkurikou Kabushikikaisha) with reference to a method described in Protein, Nucleic acid, Enzyme, Vol. 33, No. 4 (1988) pp.

337-347. The results are shown in Figures 4 and 5. The heat capacity (C_p J/g.k) was plotted on an ordinate, and the temperature ($^{\circ}\text{C}$) was plotted on an abscissa. As shown in Figures 4 (MK3G56) and Figure 5 (MK3P47), the thermal denaturation point of the MK3G56 peptide was 76.3 $^{\circ}\text{C}$, whereas that of the MK3P47 peptide was 78.1 $^{\circ}\text{C}$. This shows that an amino acid Pro at position 47 in SEQ ID NO: 3 apparently improves thermal stability.

10

Example 13

Immobilization and γ -ray
sterilization stability of K3P47 peptide

15

Water was added to 90 ml of GCL2000m (produced by Chisso Corporation, a molecular weight of the exclusion limit of a globular protein: 3,000,000) which is a cellulose type porous hard gel so that a total volume became 180 ml. Thereafter, 60 ml of 2 M sodium hydroxide was added to the mixture, and the resultant mixture was heated to 40 $^{\circ}\text{C}$. Then, 21 ml of epichlorohydrin was added to the mixture, and the mixture was allowed to react at 40 $^{\circ}\text{C}$ for one hour with stirring. After the completion of the reaction, the reaction product was thoroughly washed with water to obtain an epoxy activated gel.

25

The epoxy activated gel was suction-dried (aspirated by an aspirator on a glass filter G3 for 15 minutes). One gram of epoxy activated gel was suspended in 2.7 ml of a borate buffer (pH 10.0) (8.74 ml of 0.1 N aqueous sodium hydroxide was added to 10 ml of an aqueous solution of 0.1 M sodium tetraborate and 0.1 M potassium chloride, and water was added to the mixture so

30

that a total volume became 20 ml), and 10 mg of MK3P47 peptide (SEQ ID NO: 4) was added to the suspension. The resultant suspension was allowed to react at 40°C for 60 hours with stirring.

5

The supernatants of the reactant before and after the reaction were 2-fold diluted, respectively. Then, 100 μ l of a BCA measurement reagent (produced by Pierce) was added to each 10 μ l of the diluted supernatants, then
10 the supernatants were incubated at 37°C for 30 minutes. Thereafter, the absorbance at $\lambda = 550$ nm was measured. Calibration was conducted and the concentration of the peptide and the immobilized amount thereof before and after the reaction were calculated. As a result, the
15 reaction yield of the peptide with respect to the gel was 34%, and the immobilized amount thereof was 2.1 mg per 1 ml of gel.

The synthetic adsorbent with the peptide immobilized thereon was washed with a sufficient amount of
20 saline to obtain adsorbent GCL2000mMK3P47.

Furthermore, a part of the synthesized adsorbent GCL2000mMK3P47 was subjected to γ -ray sterilization
25 (25kGy) while being suspended in a saline, thereby obtaining adsorbent γ GCL2000mMK3P47 subjected to γ -ray sterilization. (Hereinafter, an adsorbent subjected to γ -ray sterilization will be represented with γ .)

30 Then, 300 μ l of the human serum containing 20 μ g/ml of thermally coagulated immunoglobulin complexes (hereinafter, referred to as "Model IC") was added to 100 μ l of the adsorbent GCL2000mMK3P47 or γ GCL2000mMK3P47

thus obtained, and the mixture was shaken at 37°C for 2 hours. The supernatant of the serum was collected by centrifugation, and the concentrations of immunoglobulins (hereinafter, referred to as "IgG") and IC were measured to calculate an adsorption ratio. The concentration of IgG was measured by an extracorporeal diagnostic medical product "N-Assay TIA" measurement kit available from Nittobo, and the concentration of IC was measured by an enzyme immunoassay Flerizer Clq-CIC, manufactured by Fujirebio Inc. in accordance with a manual attached thereto. The results are as shown in Table 8. Specifically, the GCL2000mMK3P47, using as a ligand, recombinant MK3P47 peptide in which Asp at position 47 of the C3 peptide was replaced by Pro, and Lys-Lys-Lys- was added to the N-terminus as a functional group for immobilization to a carrier, showed high adsorbing capacity with respect to IC and IgG, and the adsorbing capacity hardly decreased even after subjected to γ -ray sterilization.

20

Example 14

Immobilization and γ -ray sterilization stability of MK3G56 peptide

25

An adsorbent GCL2000mMK3G56 and an adsorbent γ GCL2000mMK3G56 were obtained in the same way as in Example 13, except that MK3G56 peptide obtained in Example 8 was used. The reaction yield of the peptide with respect to a gel was 40%, and the immobilized amount thereof was 2.5 mg per 1 ml of gel.

30

The IgG adsorption ratio and the IC adsorption

ratio of the GCL2000mMK3G56 or γ GCL2000mMK3G56 thus obtained are as shown in Table 8. The GCL2000mMK3G56, using as a carrier recombinant MK3G56 peptide in which Lys-Lys-Lys- was added to the N-terminus of the C3 peptide as a function group for immobilization to a carrier, exhibited a relatively high adsorption ratio with respect to IgG and IC. However, the GCL2000mMK3G56 lost most of its adsorbing capacity after being subjected to a γ -ray sterilization treatment.

Example 15

Immobilization and γ -ray sterilization stability of MP47K3 peptide

An adsorbent GCL2000mMP47K3 and an adsorbent γ GCL2000mMP47K3 were obtained in the same way as in Example 13, except that MP47K3 peptide obtained in Example 5 was used. The reaction yield of the peptide with respect to a gel was 42%, and the immobilized amount thereof was 2.6 mg per 1 ml of gel.

The IgG adsorption ratio and the IC adsorption ratio of the GCL2000mMP47K3 or γ GCL2000mMP47K3 thus obtained are as shown in Table 8. In the GCL2000mMP47K3, using as a ligand recombinant MP47K3 peptide in which Asp at position 47 of the C3 peptide was replaced by Pro, and Lys-Lys-Lys was added to the C-terminus as a functional group for immobilization to a carrier, the γ -ray sterilization stability of an adsorbing capacity with respect to IC and IgG substantially improved, compared with the case where the MK3G56 peptide (comparative example) was used as a carrier.

Example 16

Immobilization and γ -ray
sterilization stability of MP47 peptide

5

Water was added to 90 ml of GC700m (produced by Chisso Corporation, a molecular weight of the exclusion limit of a globular protein: 400,000) which is a cellulose type porous hard gel so that a total volume became 180 ml. Thereafter, 60 ml of 2 M sodium hydroxide was added to the mixture, and the resultant mixture was heated to 40°C. Then, 21 ml of epichlorohydrin was added to the mixture, and the mixture was allowed to react at 40°C for one hour with stirring. After the completion of the reaction, the reaction product was thoroughly washed with water to obtain an epoxy activated gel.

10

15

An adsorbent GC700mMP47 and an adsorbent γ GC700mMP47 were obtained in the same way as in Example 13, except that the epoxy activated gel thus obtained and MP47 peptide obtained in Example 6 were used. The reaction yield of the peptide was 42%, and the immobilized amount was 2.6 mg per 1 ml of gel.

20

25

The IgG adsorption ratio and the IC adsorption ratio of the GC700mMP47 or γ GC700mMP47 thus obtained are as shown in Table 8. In the GC700mMP47, using as a ligand recombinant peptide MP47 in which Asp at position 47 of the C3 peptide was replaced by Pro, γ -ray sterilization stability of the IC and IgG adsorbing capacity substantially improved, compared with the case where recombinant C3 peptide (MG56) described in Example 18 was used as a ligand.

30

Example 17

Immobilization and γ -ray
sterilization stability of MP47C peptide

5

An adsorbent GC700mMP47C and an adsorbent γ GC700mMP47C were obtained in the same way as in Example 13, except that the epoxy activated gel obtained in Example 16 and MP47C peptide obtained in Example 7 were
10 used. The reaction yield of the peptide with respect to a gel was 64%, and the immobilized amount was 4.0 mg per 1 ml of gel.

The IgG adsorption ratio and the IC adsorption
15 ratio of the GC700mMP47C or γ GC700mMP47C thus obtained are as shown in Table 8. In the GC700mMP47C, using as a ligand recombinant MP47C peptide in which Asp at position 47 of the C3 peptide was replaced by Pro, and -Cys was
20 added to the C-terminus as a functional group for immobilization to a carrier, the adsorption ratio with respect to IC and IgG remarkably improved, compared with the case where the MG56 corresponding to recombinant C3 peptide was used as a ligand in Example 18 and the case
25 where MP47 peptide with no -Cys added to the C-terminus was used as a ligand in Example 16. Furthermore, the adsorbing capacity hardly decreased even after γ -ray sterilization.

Example 18

Immobilization and γ -ray
sterilization stability of MG56 peptide

5

An adsorbent GC700mMG56 and an adsorbent γ GC700mMG56 were obtained in the same way as in Example 13, except that the epoxy activated gel obtained in Example 16 and MG56 peptide obtained in Example 10 were used. The reaction yield of the peptide with respect to a gel was 64%, and the immobilized amount thereof was 4.0 mg per 1 ml of gel.

The IgG adsorption ratio and the IC adsorption ratio of the GC700mMG56 or γ GC700mMG56 thus obtained are as shown in Table 8. The GC700mMG56, using as a ligand MG56 which is a recombinant C3 peptide, exhibited a relatively high adsorption ratio with respect to IC and IgG. However, the GC700mMG56 lost most of its adsorbing capacity after being subjected to a γ -ray sterilization treatment.

Example 19

25

Immobilization and γ -ray
sterilization stability of MK3-7M peptide

An adsorbent HW65MK3-7M and an adsorbent γ HW65MK3-7M were obtained in the same way as in Example 13, except that 0.3 g of AF-Torecyl Toyopearl 650 was suspended in 3 ml of 0.1 M sodium carbonate buffer containing 0.5 M sodium chloride (pH 9.0) in place of suspending the above-mentioned epoxy activated gel, 10 mg

of MK3-7M peptide obtained in Example 9 was added to the suspension, and the resultant suspension was allowed to react at 40°C for 60 hours with stirring. The reaction yield of the peptide was 57%, and the immobilized amount
5 was 3.6 mg per 1 ml of gel.

The IgG adsorption ratio and the IC adsorption ratio of the HW65MK3-7M or γ HW65MK3-7M thus obtained are as shown in Table 8. In the HW65MK3-7M which comprises
10 a recombinant peptide MK3-7M as a ligand, in which 7 mutations were introduced into the C3 peptide, and Lys-Lys-Lys- was added to the N-terminus as a functional group for immobilization to a carrier, γ -ray sterilization stability of the IC and IgG adsorbing capacity
15 substantially improved, compared with the MG56 which is the recombinant C3 peptide described in Example 18.

Table 8

Example	Adsorbent	IgG adsorption ratio (%)	IC adsorption ratio (%)
5			
	Example 13 GCL2000mMK3P47	85	35
	γGCL2000mMK3P47	75	35
	Example 14 GCL2000mMK3G56	70	30
	γGCL2000mMK3G56	21	12
10	Example 15 GCL2000mMP47K3	70	35
	γGCL2000mMP47K3	60	30
	Example 16 GC700mMP47	60	35
	γGC700mMP47	53	35
	Example 17 GC700mMP47C	90	65
15	γGC700mMP47C	82	60
	Example 18 GC700mMG56	65	30
	γGC700mMG56	20	10
	Example 19 HW65MK3-7M	75	35
	γHW65MK3-7M	65	35
20			

Example 20

High-pressure vapor sterilization stability of an
 adsorbent with a peptide derivative immobilized thereon

A part of the adsorbent GC700mMP47C obtained in
 Example 17 and a part of the adsorbent GC700mMG56 ob-
 tained in Example 18 were each subjected to autoclave
 sterilization (121°C, 20 minutes) under the condition of
 being suspended in a saline. Thus, aGC700mMP47C and
 aGC700mMG56 were obtained, respectively. The IgG
 adsorption ratio of these adsorbents was calculated by

using the method similar to that of Example 13. The results are shown in Table 9.

5 The GC700mMP47, using as a ligand recombinant peptide MP47C in which Asp at position 47 of the C3 peptide was replaced by Pro, and -Cys was added to a C-terminus as a functional group for immobilization to carriers, exhibited a very high IgG adsorption ratio (50%) after an autoclave treatment, whereas the adsorbent 10 GC700mMG56 obtained in Example 18 using recombinant C3 peptide (MG56) as a ligand completely lost an IgG adsorbing capacity after an autoclave treatment.

Table 9

15	Adsorbent	IgG adsorption ratio (%)
20	GC700mMG56	65
	aGC700mMG56	0
	GC700mMP47C	90
	aGC700MP47C	50

25 INDUSTRIAL APPLICABILITY

By using an adsorbent with a peptide immobilized thereon of the present invention, immunoglobulins and/or immunoglobulin complexes can be selectively adsorbed and removed, and IgG₃ which is not adsorbed by protein A can be adsorbed and removed. Furthermore, the adsorbent of the present invention is inexpensive, and can be subjected to an autoclave sterilization treatment due to its 30

excellent storage stability such as thermal stability. Furthermore, the adsorbent of the present invention can efficiently adsorb and remove immunoglobulins and/or immunoglobulin complexes contained in the body fluid even
5 after the adsorbent is subjected to a drug treatment and a sterilization treatment.

- 53 -

SEQUENCE LISTING

SEQ ID NO: 1

SEQUENCE LENGTH: 56 amino acids

5 SEQUENCE TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: peptide

FEATURE

10 OTHER INFORMATION: Xbb represents Val or Ile, Xcc
 represents Ile or Leu, Xdd represents Lys or Glu, Xee
 represents Glu or Ala, Xff represents Ala or Val, and Xgg
 represents Val or Glu.

SEQUENCE DESCRIPTION: SEQ ID NO: 1:

15 Thr Thr Tyr Lys Leu Xbb Xcc Asn Gly Lys Thr Leu Lys Gly Glu Thr
 1 5 10 15
 Thr Thr Xdd Ala Val Asp Ala Xee Thr Ala Glu Lys Xff Phe Lys Gln
 20 25 30
 20 Tyr Ala Asn Asp Asn Gly Val Asp Gly Xgg Trp Thr Tyr Asp Asp Ala
 35 40 45
 Thr Lys Thr Phe Thr Val Thr Glu
 25 50 55

SEQ ID NO: 2

SEQUENCE LENGTH: 57 amino acids

30 SEQUENCE TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Cys Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu Lys Gly Glu
 1 5 10 15
 5 Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys
 20 25 30
 Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr Tyr Asp Asp
 35 40 45
 10 Ala Thr Lys Thr Phe Thr Val Thr Glu
 50 55

SEQ ID NO: 3

SEQUENCE LENGTH: 56 amino acids

15 SEQUENCE TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: peptide

FEATURE

20 OTHER INFORMATION: Xbb represents Val or Ile, Xcc
 represents Ile or Leu, Xdd represents Lys or Glu, Xee
 represents Glu or Ala, Xff represents Ala or Val, Xgg
 represents Val or Glu, and Xhh represents Lys or Gly.

SEQUENCE DESCRIPTION: SEQ ID NO: 3:

25 Thr Thr Tyr Lys Leu Xbb Xcc Asn Gly Lys Thr Leu Lys Gly Glu Thr
 1 5 10 15
 Thr Thr Xdd Ala Val Asp Ala Xee Thr Ala Glu Lys Xff Phe Lys Gln
 20 25 30
 30 Tyr Ala Asn Asp Asn Gly Val Asp Gly Xgg Trp Thr Tyr Asp Pro Ala
 35 40 45
 Thr Xhh Thr Phe Thr Val Thr Glu
 50 55

SEQUENCE LENGTH: 60 amino acids

TOPOLOGY: linear

FEATURE

Met Lys Lys Lys Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu

Lys Gly Glu Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys

Ala Phe Lys Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr

Tyr Asp Pro Ala Thr Lys Thr Phe Thr Val Thr Glu

SEQ ID NO: 5

SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

CATATGAAAA AGAAGACCAC CTATAAACTG GTTATCAACG GTAAAACCTT GAAAGGTGAA 60

ACCACCACCA AGGCTGTTGA CGCTGAAACC GCTGAAAAAG CATTAAACA GTATGCTAAC 120

30 GACAACGGTG TCGACGGTGT TTGGACCTAT GACCCCGCTA CAAAAACCTT TACCGTTACC 180

GAATAAGCTT 190

- 56 -

SEQ ID NO: 6

SEQUENCE LENGTH: 60 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

5 MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu Lys Gly Glu
 1 5 10 15
 10 Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys
 20 25 30
 Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr Tyr Asp Pro
 35 40 45
 15 Ala Thr Lys Thr Phe Thr Val Thr Glu Lys Lys Lys
 50 55 60

SEQ ID NO: 7

SEQUENCE LENGTH: 190 base pairs

20 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

SEQUENCE DESCRIPTION: SEQ ID NO: 7:

25 CATATGACCA CCTATAAACT GCTTATCAAC GGTAACACCC TGAAAGGTGA AACCACCACC 60
 AAGGCTGTTG ACGCTGAAAC CGCTGAAAAA GCATTAAAC AGTATGCTAA CGACAACGGT 120
 GTCGACGGTG TTTGGACCTA TGACCCCGCT ACCAAAACCT TTACCGTTAC CGAAAAAAG 180
 30 AAGTAAGCTT 190

- 57 -

SEQ ID NO: 8

SEQUENCE LENGTH: 57 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

5 MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu Lys Gly Glu
1 5 10 15
10 Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys
20 25 30
Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr Tyr Asp Pro
35 40 45
15 Ala Thr Lys Thr Phe Thr Val Thr Glu
50 55

SEQ ID NO: 9

SEQUENCE LENGTH: 181 base pairs

20 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

SEQUENCE DESCRIPTION: SEQ ID NO: 9:

25 CATATGACCA CCTATAAACT GGTTATCAAC GGTAACCC TGAAAGGTGA AACCACCACC 60
AAGGCTGTTG ACGCTGAAAC CGCTGAAAAA GCATTAAAC AGTATGCTAA CGACAACGGT 120
GTCGACGGTG TTTGGACCTA TGACCCCGCT ACCAAAACCT TTACCGTTAC CGAATAAGCT 180
30 T 181

- 58 -

SEQ ID NO: 10

SEQUENCE LENGTH: 58 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

5 MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu Lys Gly Glu
 1 5 10 15
 10 Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys
 20 25 30
 Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr Tyr Asp Pro
 35 40 45
 15 Ala Thr Lys Thr Phe Thr Val Thr Glu Cys
 50 55

SEQ ID NO: 11

SEQUENCE LENGTH: 184 base pairs

20 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

SEQUENCE DESCRIPTION: SEQ ID NO: 11:

25 CATATGACCA CCTATAAACT GGTTATCAAC GGTAAAACCC TGAAAGGTGA AACCACCACC 60
 AAGGCTGTTG ACGCTGAAAC CGCTGAAAAA GCATTTAAAC AGTATGCTAA CGACAACGGT 120
 GTCGACGGTG TTTGGACCTA TGACCCCGCT ACCAAAACCT TTACCGTTAC CGAATGCTAA 180
 30 GCTT 184

SEQ ID NO: 12

SEQUENCE LENGTH: 60 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

5 MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 12:

	Met	Lys	Lys	Lys	Thr	Thr	Tyr	Lys	Leu	Val	Ile	Asn	Gly	Lys	Thr	Leu
	1				5				10						15	
10	Lys	Gly	Glu	Thr	Thr	Thr	Lys	Ala	Val	Asp	Ala	Glu	Thr	Ala	Glu	Lys
				20					25					30		
	Ala	Phe	Lys	Gln	Tyr	Ala	Asn	Asp	Asn	Gly	Val	Asp	Gly	Val	Trp	Thr
			35					40					45			
15	Tyr	Asp	Asp	Ala	Thr	Lys	Thr	Phe	Thr	Val	Thr	Glu				
		50					55					60				

SEQ ID NO: 13

SEQUENCE LENGTH: 190 base pairs

20 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

SEQUENCE DESCRIPTION: SEQ ID NO: 13:

25	CATATGAAAA	AGAAGACCAC	CTATAAACTG	GTTATCAACG	GTAAAACCCCT	GAAAGGTGAA..	60
	ACCACCACCA	AGGCTGTTGA	CGCTGAAACC	GCTGAAAAAG	CATTTAAACA	GTATGCTAAC	120
	GACAACGGTG	TCGACGGTGT	TTGGACCTAT	GACGACGCTA	CCAAAACCTT	TACCGTTACC	180
30	GAATAAGCTT						190

- 60 -

SEQ ID NO: 14

SEQUENCE LENGTH: 60 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

5 MOLECULE TYPE: peptide

SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Met Lys Lys Lys Thr Thr Tyr Lys Leu Ile Leu Asn Gly Lys Thr Leu
 1 5 10 15
 10 Lys Gly Glu Thr Thr Thr Lys Ala Val Asp Ala Ala Thr Ala Glu Lys
 20 25 30
 Val Phe Lys Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Glu Trp Thr
 35 40 45
 15 Tyr Asp Pro Ala Thr Lys Thr Phe Thr Val Thr Glu
 50 55 60

SEQ ID NO: 15

SEQUENCE LENGTH: 190 base pairs

20 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: other nucleic acid, synthetic DNA

SEQUENCE DESCRIPTION: SEQ ID NO: 15:

25 CATATGAAAA AGAAGACCAC CTATAAACTG ATCCTGAACG GTAAAACCAT AAAAGGTGAA 60
 ACCACCACCG GAGCTGTTGA CGCTGCTACC GCTGAAAAAG TTTTAAACA GTATGCTAAC 120
 GACAACGGTG TCGACGGTGA ATGGACCTAT GACGACGCTA CCAAAACCTT TACCGTTACC 180
 30 GAATAAGCTT 190

SEQ ID NO: 16
 SEQUENCE LENGTH: 57 amino acids
 SEQUENCE TYPE: amino acid
 TOPOLOGY: linear
 5 MOLECULE TYPE: peptide
 SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Met Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr Leu Lys Gly Glu
 1 5 10 15
 10 Thr Thr Thr Lys Ala Val Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys
 20 25 30
 Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Val Trp Thr Tyr Asp Asp
 15 35 40 45
 Ala Thr Lys Thr Phe Thr Val Thr Glu
 50 55

20 SEQ ID NO: 17
 SEQUENCE LENGTH: 181 base pairs
 SEQUENCE TYPE: nucleic acid
 STRANDEDNESS: double
 TOPOLOGY: linear
 25 MOLECULE TYPE: other nucleic acid, synthetic DNA
 SEQUENCE DESCRIPTION: SEQ ID NO: 17:

CATATGACCA CCTATAAACT GGTTATCAAC GGTAACCC TGAAAGGTGA AACCACCACC 60
 AAGGCTGTTG ACGCTGAAAC CGCTGAAAAA GCATTAAAC AGTATGCTAA CGACAACGGT 120
 30 GTCGACGGTG TTTGGACCTA TGACGACGCT ACCAAAACCT TTACCGTTAC CGAATAAGCT 180
 T 181



KA014

- 62 -

CLAIMS

1. An adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes, wherein a peptide capable of
5 adsorbing the immunoglobulins and/or the immunoglobulin complexes is immobilized onto a water-insoluble carrier, and the peptide has one or two amino acid sequence selected from amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing.

10

2. An adsorbent for adsorbing immunoglobulins and/or immunoglobulin complexes, wherein a peptide derivative is immobilized onto a water-insoluble carrier, and the peptide derivative includes at least one of deletion,
15 substitution, insertion, or addition of an amino acid with respect to the amino acid sequences represented by SEQ ID NO: 1 of Sequence Listing, and is capable of adsorbing immunoglobulins and/or immunoglobulin complexes.

20

3. An adsorbent according to claim 2, wherein at least one of properties of thermal stability, drug stability, γ -ray sterilization stability, and high-pressure vapor sterilization stability of the peptide derivative is
25 improved over that of a peptide having the amino acid sequence represented by SEQ ID NO: 1.

4. An adsorbent according to claim 2 or 3, wherein the peptide derivative includes an amino acid sequence represented by SEQ ID NO: 3.
30

5. An adsorbent according to any one of claims 1 to 4, wherein (Lys)_n or (Cys)_m is added to the amino terminus



KA014

- 63 -

and/or the carboxyl terminus of the peptide or the peptide derivative, wherein n and m are integers of 0 to 9.

5 6. An adsorbent according to any one of claim 1 to 5, wherein the peptide or the peptide derivative consists of an amino acid sequence of 70 amino acids or less.

10 7. An adsorbent according to any one of claims 1 to 6, wherein the adsorbent is subjected to high-pressure vapor sterilization.

15 8. An adsorbent according to any one of claims 1 to 6, wherein the adsorbent is subjected to γ -ray sterilization.

9. An adsorbent according to any one of claims 1 to 6, wherein the water-insoluble carrier is porous.

20 10. An adsorbent according to any one of claims 1 to 6, wherein the water-insoluble carrier is hydrophilic.

25 11. An adsorbent according to claim 9 or 10, wherein the molecular weight of the exclusion limit of the water-insoluble carrier is in the range of 150,000 to 5,000,000.

30 12. An adsorbent according to any one of claims 1 to 11, wherein the adsorbent is used for adsorbing and removing immunoglobulins and/or immunoglobulin complexes present in the blood, plasma, or other body fluids obtained from a living organism.



KA014

- 64 -

13. A method for adsorbing and removing immunoglobulins and/or immunoglobulin complexes, comprising the step of bringing the adsorbent of any one of claims 1 to 11 into contact with a solution containing immunoglobulins and/or immunoglobulin complexes.

14. A device for adsorbing and removing immunoglobulins and/or immunoglobulin complexes, the adsorbent of any one of claims 1 to 11 is accommodated in a vessel having an inlet and an outlet for a solution, and the device comprises effluent preventing means for preventing the adsorbent from flowing out of the vessel.

15. A method according to claim 13, wherein the solution containing immunoglobulins and/or immunoglobulin complexes is the blood, plasma, or other body fluids obtained from a living organism.

ABSTRACT

An adsorbent that exhibits a high specificity in adsorbing immunoglobulins and/or complexes thereof, is
5 extremely reduced in the lowering of the adsorption characteristic during sterilization or storage, is highly stable and safe, and is prepared by immobilizing on a water-insoluble support either a peptide derivative which has undergone at least one of the deletion, substitution,
10 insertion, or addition of amino acids in a peptide having a specified amino acid sequence or an amino acid sequence, or the above peptide derivative which has undergone the addition of Lys or Cys at the amino and/or carboxyl terminal thereof; a device for adsorption and
15 removal made by packing the adsorbent in a vessel equipped with effluent preventing means; and a method of adsorbing and removing immunoglobulins and/or complexes thereof contained in the blood, plasma or other body fluids with the adsorbent.

FIG. 1




FIG. 2




FIG.3




FIG.4




FIG.5




FIG. 6





FIG. 7



obtaining a peptide which has high thermal stability, drug stability, and/or sterilization stability, when determined based on high adsorbing specificity and adsorbing capacity with respect to immunoglobulins and/or immunoglobulin complexes as indexes, and which is unlikely to have side effects caused by antigenicity, by using a partial peptide of protein G.

Protein G is one of the immunoglobulin binding proteins, which has a molecular weight of about 50,000 daltons and is present on a bacteria cell wall of Streptococcus. The cDNA of protein G has already been cloned. Based on an amino acid sequence predicted from the cDNA, protein G is assumed to consist of 448 amino acid residues. Furthermore, it has been assumed that several similar repeated amino acid sequences are present in the predicted amino acid sequence, and that the repeated amino acid sequences are binding domains to albumin and immunoglobulins. See example, S.R. Fahnestock et al., J. Bacteriol., Vol. 167, pp. 870-880 (1986); U. Sjöbring et al., J. Biol. Chem., Vol. 266, pp. 399-405 (1991). In B. Guss et al., EMBO J., vol. 5, pp. 1567-1575 (1986), a repeated amino acid sequences region represented by C1, C2, or C3 suggesting an immunoglobulin (IgG) binding domain in the sequence of protein G is described. When common amino acid sequence of C1, C2, and C3 are summarized, the amino acid sequence of the IgG binding domain is as represented by SEQ ID NO: 1 of Sequence Listing.

Furthermore, the inventors also studied sterilization stability of C3 peptides to y-ray sterilization

