

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85712 A1

(51) International Patent Classification⁷: C07D 307/87

(21) International Application Number: PCT/DK01/00333

(22) International Filing Date: 10 May 2001 (10.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA200000783 12 May 2000 (12.05.2000) DK

(71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). DANCER, Robert [AU/DK]; J. M. Thielesvej 8, st th, DK-Frederiksberg C 1961 (DK).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,

CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR THE PREPARATION OF CITALOPRAM

(57) Abstract: A method for the preparation of citalopram comprising reaction of a compound of formula 5-aminomethyl-1-(3-dimethylamino-propyl)-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran with an oxidising agent to prepare citalopram.


WO 01/85712 A1

Method for the Preparation of Citalopram

The present invention relates to a method for the preparation of the well-known antidepressant drug citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile.

Background of the Invention

Citalopram is a well-known antidepressant drug that has now been on the market for some years and has the following structure:



Formula I


It is a selective, centrally acting serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities. The antidepressant activity of the compound has been reported in several publications, eg. J. Hyttel *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.* 1982, 6, 277-295 and A. Gravem, *Acta Psychiatr. Scand.* 1987, 75, 478-486. The compound has further been disclosed to show effects in the treatment of dementia and cerebrovascular disorders, EP-A 474580.

20

Citalopram was first disclosed in DE 2,657,013, corresponding to US 4,136,193. This patent publication describes the preparation of citalopram by one method and outlines a further method, which may be used for preparing citalopram.


25 According to the process described, the corresponding 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile is reacted with 3-(N,N-dimethylamino)propyl-chloride in the presence of methylsulfinylmethide as condensing agent. The starting material was prepared from the corresponding 5-bromo derivative by reaction with cuprous cyanide.

30 According to the method, which is only outlined in general terms, citalopram may be obtained by ring closure of the compound:



in the presence of a dehydrating agent and subsequent exchange of the 5-bromo group with cuprous cyanide. The starting material of Formula II is obtained from 5-bromophthalide by two successive Grignard reactions, i.e. with 4-fluorophenyl magnesium chloride and N,N-dimethylaminopropyl magnesium chloride, respectively.

A new and surprising method and an intermediate for the preparation of citalopram were described in US Patent No 4,650,884, according to which an intermediate of the formula



is subjected to a ring closure reaction by dehydration with strong sulfuric acid in order to obtain citalopram. The intermediate of Formula III was prepared from 5-cyanophthalide by two successive Grignard reactions, i.e. with 4-fluorophenyl magnesium halogenide and N,N-dimethylaminopropyl magnesium halogenide, respectively.


Further processes are disclosed in International patent application Nos. WO 98019511, WO 98019512 and WO 98019513. WO 98019512 and WO 98019513 relate to methods wherein a 5-amino-, 5-carboxy- or 5-(sec. aminocarbonyl)phthalide is subjected to two successive Grignard reactions, ring closure and conversion of the resulting 1,3-dihydroisobenzofuran derivative to the corresponding 5-cyano compound, i.e. citalopram. International patent application No. WO 98019511 discloses a process for the manufacture of citalopram wherein a (4-substituted-2-hydroxymethylphenyl-(4-fluorophenyl)methanol compound is subjected to ring closure and the resulting 5-substituted 1-(4-fluorophenyl)-1,3-dihydroisobenzofuran converted to the corresponding 5-cyano derivative which is alkylated with a (3-dimethylamino)propylhalogenide in order to obtain citalopram.

Finally, methods for preparing the individual enantiomers of citalopram are disclosed in US Patent No. 4,943,590 from which it also appears that the ring closure of the intermediate of Formula III may be carried out via a labile ester with a base.


It has now, surprisingly, been found that citalopram may be manufactured by a novel favourable and safe procedure using convenient starting materials.

Summary of the invention

A first aspect of the present invention provides a method for the preparation of citalopram comprising reaction of a compound of Formula IV



with an appropriate oxidising agent to afford citalopram




which is isolated as the base or a pharmaceutically acceptable salt thereof.

According to embodiments, there may be provided methods for preparing the intermediates of Formula IV.


3a

A second aspect of the present invention provides a compound of Formula V



Formula V

A third aspect of the present invention provides a method for the preparation of an intermediate of Formula V comprising a ring closure reaction of a compound of Formula XIII



Formula XIII


with a suitable reducing agent.

In a fourth aspect, the present invention provides an antidepressant pharmaceutical composition comprising citalopram manufactured by the method disclosed in the invention.

Furthermore, according to the invention, the compounds of Formula IV may be prepared by different methods.

One of these methods includes the following steps:


5



6-carboxy-3-(4-fluorophenyl)phthalide is reacted with an alcohol, R-OH, wherein R is preferably lower alkyl, most preferably Me, in the presence of a dehydrating agent, preferably SOCl₂.

10


The resulting compound of Formula VI is alkylated with



wherein X is a leaving group in the presence of a suitable base. X is preferably halogen or sulphonate.

Optionally, the alkylating reaction is a stepwise alkylation. In this case, the resulting compound of Formula VI is alkylated with a compound having the formula

20




wherein X' is a suitable leaving group and R' is -CH₂-O-Pg, -CH₂-NPg₁Pg₂, -CO-N(CH₃)₂, -CH(OR¹)(OR²), -C(OR⁴)(OR⁵)(OR⁶) or -COOR³, wherein Pg is a protection group for an alcohol group, Pg₁ and Pg₂ are protection groups for an amino

25

5

group, R^1 and R^2 are alkyl groups or R^1 and R^2 together form a chain of 2 to 4 carbon atoms and R^3 , R^4 , R^5 and R^6 are alkyl, alkenyl, alkynyl, aryl or aralkyl;

to form a compound of Formula XVIII



Formula XVIII

5


wherein R' is as defined above; followed by conversion of the group R' to a dimethylaminomethyl group.

The resulting compound of Formula VII is reacted with a reducing agent such as $LiAlH_4$, Red-Al, AlH_3 or activated forms of $NaBH_4$, e.g. $NaBH_4$, Me_2SO_4 ; $NaBH_4$, I_2 ; $NaBH_4$, BF_3 , Et_2O ; or B_2H_6 ; followed by treatment with acid or another dehydrating agent to perform ring closure to form the compound of Formula VIII.

The alcohol of Formula VIII is conveniently activated by tosylchloride or mesylchloride to form the corresponding substituted sulphonate; or the alcohol is converted into the corresponding benzylic halide. This conversion is preferably carried out with $SOBr_2$ or $SOCl_2$.

The corresponding sulphonate or halide is either converted directly to the compound of Formula IV by reaction with liquid ammonia; or by a reaction with a metal salt of phthalimide, preferably potassium phthalimide followed by treatment with NH_2NH_2 or by treatment with an amine in an alcohol, i.e. R^8NH_2/R^9-OH , wherein R^8 and R^9 are lower alkyl, preferably methyl or ethyl, e.g. methylamine in ethanol; or by a reaction with metal azide, MN_3 , M preferably being Na or K; followed by treatment with a reducing agent such as Pd/C and H_2 or a hydrate source such as $LiAlH_4$ or $NaBH_4$ or an activated form of it.


Another method for preparing the compound of Formula IV includes the following steps:



6-carboxy-3-(4-fluorophenyl)phthalide is conveniently reacted with a dehydrating agent such as thionylchloride, followed by aminolysis of the resulting activated acid derivative.

5

The resulting compound of Formula IX is alkylated with




wherein X is a leaving group in the presence of a suitable base. X is preferably halogen or sulphonate.

10

Optionally, the alkylating reaction is a stepwise alkylation analogous to the stepwise alkylation described above.

15 The resulting compound of Formula X is reacted with a reducing agent such as LiAlH_4 , Red-Al, AlH_3 or activated forms of NaBH_4 , e.g. NaBH_4 , Me_2SO_4 ; NaBH_4 , I_2 ; NaBH_4 , $\text{BF}_3 \cdot \text{Et}_2\text{O}$; or B_2H_6 ; followed by treatment with acid or another dehydrating agent to perform ring closure to form the compound of Formula IV.

20 According to a third method for preparing the compound of Formula IV, the corresponding 6-cyano substituted derivative of 6-carboxy-3-(4-fluorophenyl)phthalide is prepared.




The carboxy derivative is either reacted with SOCl_2 followed by treatment with ammonia and finally a dehydrating agent such as SOCl_2 to prepare the cyano derivative of Formula XI;

25

or reacted with an alcohol R-OH in the presence of acid followed by treatment with ammonia and finally reacted with SOCl_2 ; or reacted in a one-pot process such as with

$\text{SO}_2(\text{NH}_2)_2$, SOCl_2 and sulfolane, or with *tert*-butylamine, a dehydrating agent such as POCl_3 and a suitable solvent, such as toluene.

The resulting compound of Formula XI is alkylated with



5

wherein X is a leaving group in the presence of a suitable base. X is preferably halogen or sulphonate.

Optionally, the alkylating reaction is a stepwise alkylation analogous to the stepwise
10 alkylation described above.


The resulting compound of Formula XII is reacted with a reducing agent such as LiAlH_4 , Red-Al, AlH_3 or activated forms of NaBH_4 , e.g. NaBH_4 , Me_2SO ; NaBH_4 , I_2 ; NaBH_4 , $\text{BF}_3\cdot\text{Et}_2\text{O}$; or B_2H_6 ; followed by treatment with acid to perform ring closure to form the
15 compound of Formula IV.

Other reaction conditions, solvents, etc. for the reactions described above are conventional conditions for such reactions and may easily be determined by a person skilled in the art.

20 In another aspect, the present invention provides the novel intermediate of Formula V.

In a further aspect, the invention relates to methods for preparing the intermediate of Formula V.


25 One stepwise process for preparing the intermediate of Formula V is illustrated below:



m-xylene and p-fluorobenzoyl chloride, which are commercially available compounds are reacted in the presence of AlCl_3 to afford the compound of Formula XIV. This compound
30 is oxidised with permanganate, preferably KMnO_4 or NaMnO_4 , giving the resulting compound of Formula XIII, which is finally reacted conveniently with Zn in acid, preferably acetic acid.

8

Alternatively, the compound of Formula IV is prepared from the compound of Formula XIII by the following stepwise process:




The compound of Formula XIII is reacted with a reducing agent such as LiAlH_4 , Red-Al, AlH_3 or activated forms of NaBH_4 , e.g. NaBH_4 , Me_2SO , NaBH_4 , I_2 ; NaBH_4 , $\text{BF}_3 \cdot \text{Et}_2\text{O}$; or B_2H_6 ; followed by treatment with acid to perform ring closure to form the compound of Formula XV.

The alcohol of Formula XV is conveniently activated by tosylchloride or mesylchloride to form the corresponding substituted sulphonate; or the alcohol is converted into the corresponding benzylic halide. This conversion is preferably carried out with SOBr_2 or SOCl_2 .

The corresponding sulphonate or halide is either converted directly to the compound of Formula XVII by reaction with liquid ammonia; or by a reaction with a metal salt of phthalimide, preferably potassium phthalamide, followed by treatment with NH_2NH_2 or by treatment with an amine in an alcohol, i.e. $\text{R}^8\text{NH}_2/\text{R}^9\text{-OH}$, wherein R^8 and R^9 are lower alkyl, preferably methyl or ethyl, e.g. methylamine in ethanol; or by a reaction with metal azide MN_3 , M preferably being Na or K; followed by treatment with a reducing agent such as Pd/C and H_2 or a hydride source such as LiAlH_4 or NaBH_4 or an activated form thereof.

The resulting compound of Formula XVII is alkylated with



wherein X is a leaving group in the presence of a suitable base. X is preferably halogen or sulphonate.

Optionally, the alkylating reaction is a stepwise alkylation analogous to the stepwise alkylation described above.

Optionally the steps of the alkylation and the conversion to the cyano derivative are in opposite order so the conversion to the cyano derivative is performed before the alkylation.

Throughout the specification and claims, the terms lower alkyl or C₁₋₆ alkyl refer to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, 2,2-dimethyl-1-ethyl and 2-methyl-1-propyl.

Similarly, alkenyl and alkynyl, respectively, designate such groups having from two to six carbon atoms, including one double bond and triple bond respectively, such as ethenyl, propenyl, butenyl, ethynyl, propynyl, and butynyl.

The term aryl refers to a mono- or bicyclic carbocyclic aromatic group, such as phenyl and naphthyl, in particular phenyl.

The term aralkyl refers to aryl-alkyl, wherein aryl and alkyl are as defined above.

Halogen means chloro, bromo or iodo.

The compound of general Formula I may be used as the free base or as a pharmaceutically acceptable acid addition salt thereof. As acid addition salts, such salts formed with organic or inorganic acids may be used. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene sulfonic and theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids.

The acid addition salts of the compounds may be prepared by methods known in the art. The base is reacted with either the calculated amount of acid in a water miscible solvent, such as acetone or ethanol, with subsequent isolation of the salt by concentration and

cooling, or with an excess of the acid in a water immiscible solvent, such as ethylether, ethylacetate or dichloromethane, with the salt separating spontaneously.

The pharmaceutical compositions of the invention may be administered in any suitable way and in any suitable form, for example orally in the form of tablets, capsules, powders or syrups, or parenterally in the form of usual sterile solutions for injection.

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: Corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive, colouring, aroma, preservative etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilising the solution and filling it in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Examples

The invention is further illustrated by the following examples.

25

Example 1

5-Aminomethyl-1-(3-dimethylamino-propyl)-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran

1-(3-Dimethylamino-propyl)-1-(4-fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carbonitrile (5.4 g, 16.2 mmol) was dissolved in dry THF (5 mL) and diluted with dry ether (50 mL). This solution was added dropwise to a refluxing suspension of lithium aluminium hydride (2.5 g, 65 mmol) in dry ether (150 mL) over 10 – 15 minutes, after which the resulting suspension was heated at reflux for a further 4 h. The solution was allowed to cool to room temperature and was stirred at room temperature overnight. The reaction was quenched with a minimum of water, and the resulting solution/suspension was dried over anhydrous magnesium sulfate. The mixture was filtered, and the solid cake was washed with THF. The combined filtrates were evaporated to give an oil. The oil was dissolved in toluene (200 mL) and was stirred with an aqueous solution of sulfuric acid (10 mL, 70 % v/v) for 3 h. The mixture was diluted with water, and the pH was adjusted to >9

by the addition of aqueous ammonia solution (25% w/v). The toluene was separated, and the aqueous phase was extracted with further toluene. The combined toluene extracts were dried over anhydrous magnesium sulfate, filtered and evaporated to give the title compound as a yellow oil (4.4 g, 84%). ¹H NMR (CDCl₃): δ 1.25-1.40 (m, 1H), 1.40-1.55 (m, 1H), 2.11 (ddd, 1H), 2.13 (t, 3H), 2.15 (ddd, 1H), 2.21 (t, 2H), 3.85 (s, 2H), 5.11 (d, 1H), 5.14 (d, 1H), 6.96 (t, 2H), 7.15 (s, 1H), 7.21 (d, 1H), 7.22 (d, 1H), 7.45 (dd, 2H).

Example 2

Citalopram, HBr

10 A mixture of 5-aminomethyl-1-(3-dimethylamino-propyl)-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran (10 g, 30 mmol) and 5Å molecular sieves (24 g) in pyridine (150 mL) was stirred at 60 °C under an atmosphere of oxygen. Copper(I) chloride (1.8 g, 1.8 mmol) was added, and the mixture was stirred for 3 h. Further copper(I) chloride (1.8 g, 1.8 mmol) was added, and the mixture was stirred overnight. The mixture was poured onto
15 ice, and the pH of the mixture was adjusted to >9 by the addition of aqueous ammonia solution (25% w/v). The solution was diluted with toluene and filtered. The organic phase was separated, and the aqueous was washed with further toluene. The combined organic extracts were washed with water, dried over anhydrous sodium sulfate and evaporated. The residue was treated with heptane and was evaporated to give an oil (11.1 g). This oil
20 was dissolved in acetone and treated with aqueous hydrobromic acid (7 ml, 47% w/v). The solution was evaporated, and the residue was dissolved in *iso*-propanol (100 mL). The solution was stirred overnight. The resulting precipitate was filtered and dried to give the HBr salt of citalopram as a white powder (8.2 g, 66%). The filtrate was evaporated, and the oily residue was shaken with ether and allowed to stand overnight. Filtration of the
25 solution gave further HBr salt of citalopram as a brown solid (1.7 g, 14%). ¹H NMR (d⁶-DMSO): δ 1.35-1.50 (m, 1H), 1.50-1.60 (m, 1H), 2.25 (t, 2H), 2.69 (s, 3H), 3.00-3.10 (m, 2H), 5.17 (d, 1H), 5.25 (d, 1H), 7.18 (t, 2H), 7.61 (dd, 2H), 7.77 (d, 1H), 7.82 (d, 1H), 7.83 (s, 1H), 9.27 (bs, 1H).

30 Example 3

1-(4-Fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid methyl ester

A stirred suspension of 1-(4-fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid (1 g, 3.7 mmol) in thionyl chloride (25 mL) was heated at reflux for 25 min, during which time the solid dissolved. The thionyl chloride was then evaporated, and the residue
35 was dissolved in toluene, and again evaporated. The residue was stirred in methanol (25 mL) overnight, during which time a heavy precipitate formed. The solvent was evaporated, and the residue was partitioned between aqueous ammonia solution (25% w/v) and toluene. The organic phase was separated, dried over magnesium sulfate and

evaporated to give the title compound as a white solid (0.97 g, 92%). ¹H NMR (d⁶-DMSO): δ 3.92 (s, 3H), 6.85 (s, 1H), 7.26 (t, 2H), 7.42 (dd, 2H), 7.61 (d, 1H), 8.31 (dd, 1H), 8.36 (s, 1H).

5 Example 4

1-(4-Fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid amide

A stirred suspension of 1-(4-fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid (1 g, 3.7 mmol) in thionyl chloride (25 mL) was heated at reflux for 25 min, during which time the solid dissolved. The thionyl chloride was then evaporated, and the residue
10 was dissolved in toluene, and again evaporated. The residue was dissolved in toluene (15 mL) and was treated with a solution of ammonia in ether and a heavy precipitate formed. The mixture was stirred overnight, diluted with toluene and aqueous ammonia solution, and filtered. The residue was dried to give the title compound as a white solid (0.80 g, 80%). ¹H NMR (d⁶-DMSO): δ 6.81 (s, 1H), 7.25 (t, 2H), 7.40 (dd, 2H), 7.54 (d, 1H), 7.59
15 (bs, 1H), 8.24 (bs, 1H), 8.24 (dd, 1H), 8.42 (s, 1H).

Example 5

1-(4-Fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carbonitrile

A suspension of 1-(4-fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid
20 amide (13.6 g, 0.05 mole) in thionyl chloride (40 mL) and DMF (0.25 mL) was heated at reflux for 2 hours. The thionyl chloride was then evaporated, and the residue was dissolved in hot IPA (100 mL). On cooling, crystals of the title compound was formed. Yield: 7.8 g (62%). ¹H NMR (d⁶-DMSO): δ 6.87 (s, 1H), 7.26 (t, 2H), 7.42 (dd, 2H), 7.58 (d, 1H), 8.18 (dd, 1H), 8.48 (s, 1H).

25

Example 6

5-Bromomethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran

A suspension of 5-hydroxymethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran (2 g, 8.2 mmol) in toluene (20 mL) was heated until the solid dissolved. Heating was then stopped.
30 Thionyl bromide (2.2 g, 10.6 mmol) was added, and the mixture was stirred for 1 h. Silica (25 g) was added, and the mixture was filtered, and the residue was washed with a 1:1 v/v solution of ethyl acetate and heptane. The filtrate was evaporated to give the title compound as a red-orange oil (2.6 g, 90%). ¹H NMR (d⁶-DMSO): δ 4.72 (s, 2H), 5.11 (d, 1H), 5.28 (d, 1H), 6.17 (s, 1H), 7.04 (d, 1H), 7.17 (t, 2H), 7.33 (d, 1H), 7.38 (dd, 2H), 7.45
35 (s, 1H).

Example 7*5-Aminomethyl-1-(4-Fluoro-phenyl)-1,3-dihydro-isobenzofuran*

A suspension of 5-bromomethyl-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran (1.96 g, 6.4 mmol) was stirred in liquid re-distilled ammonia (200 mL) under nitrogen/ammonia at
5 -33 °C for 2½ days. The ammonia was allowed to evaporate, and the residue was stirred with a mixture of ethyl acetate and aqueous sulfuric acid (2 M). The aqueous phase was separated and was washed with ether. The aqueous phase was then basified to pH > 9 using aqueous ammonium hydroxide solution (25% w/v), and was extracted with toluene. The toluene extracts were dried over anhydrous magnesium sulfate and evaporated to give
10 the title compound as a yellow-orange oil (0.63 g, 40%). ¹H NMR (d⁶-DMSO): δ 3.72 (s, 2H), 5.09 (d, 1H), 5.25 (dd, 1H), 6.14 (s, 1H), 6.96 (d, 1H), 7.17 (t, 2H), 7.20 (d, 1H), 7.32 (s, 1H), 7.36 (dd, 2H).

Example 8*Citalopram*


To a stirred solution of 5-aminomethyl-1-(3-dimethylamino-propyl)-1-(4-fluoro-phenyl)-1,3-dihydro-isobenzofuran (0.5 g, 1.5 mmol) in dichloromethane (10 mL) was added an aqueous solution of potassium bisulfate and sodium hydroxide (19 mL; 0.2 M in K₂S₂O₈, 3.8 mmol; 0.4 M in NaOH, 7.6 mmol), followed by an aqueous solution of nickel sulfate
20 (1.5 mL, 40 mM, 61 µmol). The mixture was stirred vigorously for 4 days, and was then filtered through celite. The filtrate was partitioned between aqueous sulfuric acid (2 M) and toluene. The aqueous layer was separated, and the pH of the mixture was adjusted to >9 by the addition of aqueous ammonia solution (25% w/v). The solution was extracted with toluene, and this latter toluene extract was dried over magnesium sulfate and
25 evaporated to give the free base of citalopram as a very pale yellow oil (0.35 g, 70%).

Example 9*1-(4-Fluoro-phenyl)-3-oxo-1,3-dihydro-isobenzofuran-5-carboxylic acid*

Zink (38 g, 0.58 mol) was suspended in acetic acid (400 mL). The mixture was heated to
30 60 °C. 2,4-dicarboxy-4'-fluoro-benzophenone (21 g, 0.075 mol) was added in portions of 5 grams. After addition, the reaction mixture was heated at reflux temperature for two hours. The suspension was filtered while it was still hot. The filtrate was added to ice-water (1 kg) and the title compound was isolated by filtration. Yield 17.8 g (90%). ¹H NMR (d⁶-DMSO): δ 6.84 (s, 1H), 7.17 (t, 2H), 7.43 (dd, 2H), 7.59 (d, 1H), 8.31 (d, 1H), 8.35 (s, 1H).


Claims

1. A method for the preparation of citalopram comprising reaction of a compound of Formula IV



Formula IV


with an oxidising agent to afford citalopram



Formula I

which is isolated as the base or a pharmaceutically acceptable salt thereof.


2. The method of claim 1, characterised in that the intermediate of Formula IV is prepared by activating the alcohol of Formula VIII



Formula VIII


by a substituted sulphonate or converting the alcohol into a benzylic halide or another activated derivative followed by aminolysis to form the compound of Formula IV

15



Formula IV


3. The method of claim 2, characterised in that the intermediate of Formula VIII is
 5 prepared by reacting the compound of Formula VII



Formula VII

with a reducing agent.

- 10 4. The method of claim 3, characterised in that the intermediate of Formula VII is
 prepared by alkylating the compound of Formula VI




Formula VI

optionally by stepwise alkylation.

15


5. The method of claim 4, characterised in that the intermediate of Formula VI is
 prepared by reacting the compound of Formula V



Formula V


with an alcohol R-OH in the presence of a dehydrating agent.

- 5 6. The method of claim 1, **characterised in** that the intermediate of Formula IV is prepared by reacting the compound of Formula X



Formula X


with a reducing agent followed by ring closure to form the compound of Formula IV



Formula IV

10


7. The method of claim 6, **characterised in** that the intermediate of Formula X is prepared by alkylating the compound of Formula IX



Formula IX

optionally by stepwise alkylation.


- 5 8. The method of claim 7, characterised in that the intermediate of Formula IX is prepared by reacting the compound of Formula V



Formula V

- 10 with a dehydrating agent such as thionylchloride followed by aminolysis of the resulting activated acid derivative;


9. The method of claim 1, characterised in that the intermediate of Formula IV is prepared by reacting the compound of Formula XII



Formula XII


15 with a reducing agent followed by ring closure to form the compound of Formula IV

18



Formula IV

10. The method of claim 9, characterised in that the intermediate of Formula XII is prepared by alkylating the compound of Formula XI




Formula XI

5

optionally by stepwise alkylation.

11. The method of claim 10, characterised in that the intermediate of Formula XI is prepared by converting the compound of Formula V

10




Formula V

to the corresponding cyano substituted compound.

15


19

12. A compound of Formula V



Formula V

13. A method for the preparation of an intermediate of claim 12 comprising a ring closure
5 reaction of a compound of Formula XIII




Formula XIII

with a suitable reducing agent.

10

14. The method of claim 13, wherein the reducing agent is Zn in acid, preferably acetic acid.


15. The method of claim 1, characterised in that the intermediate of Formula IV is
15 prepared by alkylating the compound of Formula XVII



Formula XVII


optionally by stepwise alkylation to form the compound of Formula IV

20



Formula IV


16. The method of claim 15, characterised in that the intermediate of Formula XVII is prepared by aminolysis the compound of Formula XVI



Formula XVI

5

17. The method of claim 16, characterised in that the intermediate of Formula XVI is prepared by activating the alcohol of Formula XV



Formula XV


10

by a substituted sulphonate or converting the alcohol into a benzylic halide or another activated derivative.

15

21

18. The method of claim 17, characterised in that the intermediate of Formula XV is prepared by reacting the ketone of Formula XIII



Formula XIII

5 with a reducing agent followed by ring closure to form the compound of Formula XV.

19. An antidepressant pharmaceutical composition comprising citalopram manufactured by the method of any of the claims 1-11 and 13-18.