

LACTONE FORMULATIONS AND METHOD OF USE

Field of the Invention

The present inventions are generally in the fields of pharmaceutically
5 active lactones, their pharmaceutical formulations, and method of use
thereof, and methods for the synthetic preparation of chemically
functionalized lactones useful therefor as anticancer and antiinfective agents.

Background of the Invention

Despite the development of many different compounds which are
10 useful in the treatment of infection, cancer, and other disorders, there
remains a need for the development of new compounds which may be
effective at lower dosages, more selective, having fewer side effects or
capable of treating diseases or disorders where resistance to the known
compounds has developed.

15 Chemotherapeutic agents are used for the treatment of infections,
cancer, abnormal proliferation disorders (endometriosis, restenosis,
psoriasis), and other disorders. Most chemotherapeutic agents have side
effects due to lack of specificity. For example, cancer is one of the leading
causes of death. One of the primary modes of treating cancer, chemotherapy,
20 is used specifically to limit cell growth and replication. Most chemotherapy
agents also affect neoplastic and rapid proliferating cells of normal tissues
(e.g., bone marrow, hair follicles, etc.), which results in several negative side
effects including hair loss, nausea, vomiting, and suppression of bone
marrow function. Moreover, effectiveness of these agents frequently
25 diminishes over time due to the development of resistance.

It is therefore an object of this invention to provide a novel class of
compounds effective as anti-infective and/or anti-proliferative agents.

It is another object of this invention to provide an effective
antineoplastic agent with specific cytotoxicity in order to minimize side
30 effects.

It is a further object of the present invention to provide antiinfective
agents which are specific and different from many other drugs currently in

use, to provide an alternative method of treatment for drug resistant organisms.

Summary of the Invention

Compounds of Formulae Ia and Ic having a lactone structure and a methylene group at the alpha-position of the lactone structure have been discovered. The lactone compounds can be reacted with a nucleophilic agent to open the lactone ring to a compound of Formula Ib. The lactone of Formula Ia and its functional derivatives have been isolated from *Securidaca virgata*. These compounds are referred to as LMSV-6 or SecurolideTM. The purified compounds have demonstrated activity in assays for anti-bacterial and anti-fungal activities, and for treating proliferation disorders such as cancer. Based on the *in vitro* assays, the lactones are useful for treating proliferation disorders including, for example, breast cancer, colon cancer, rectal cancer, stomach cancer, pancreatic cancer, lung cancer, liver cancer, ovarian cancer, esophageal cancer, and leukemia. They are also effective for treatment of bacterial and fungal infections, including treatment of peptic ulcer disease, gingivitis and periodontitis.

The method for making a compound of Formulae Ia and Ic generally involves: a) providing a precursor having a lactone structure, and b) reacting the precursor with one or more chemical reagents to provide a product having a methylene group at the α -position of the lactone structure. The product can be treated with a nucleophilic agent such as an alcohol, alkoxide, amine, or any other neutral or anionic nucleophiles to generate a compound of Formula Ib.

Detailed Description of the Invention

I. Lactone Compositions

A. Lactones Isolated from *Securidaca virgata*.


Lactones with an exocyclic methylene group and their respective derivatives with a hydroxyl in gamma position are disclosed. The lactones and the derivatives thereof can be synthesized or isolated from natural resources. In one embodiment, the lactones and the derivatives can be isolated by means of chromatographic methods, from a plant whose

taxonomic scientific name is *Securidaca virgata*, which belongs to Polygalaceae as its botanical family. As used herein, the term "lactones" encompasses any organic chemicals having a five-member ring lactone structure in which the oxygen atom of the C=O group can be replaced by a sulfur atom or a nitrogen grouping.

5 The term "derivatives" as used herein refers to any compounds that are made from the lactones by reacting the lactones with one or more chemical reagents. The term also refers to any products obtainable by ring opening of the lactones with an organic or inorganic nucleophilic agents to form, for example, an acid, ester, amide, or

10 any other products thereof.

In one embodiment, the lactone has the following chemical structure:



Formula Ia

wherein

R₁-R₆ taken independently or R₃-R₆ taken together are a hydrogen

15 atom, a halogen atom, a hydroxyl group, or any other organic groupings containing any number of carbon atoms, preferably 1 – 8 carbon atoms, and optionally include a heteroatom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats, representative R₁-R₆ groupings being alkyl, substituted alkyl, alkenyl, substituted alkenyl,

20 alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted


25 carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl,

substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, or polypeptide group;

Z is a heteratom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats; and

X is a heteratom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats.

In another embodiment, the compound has the following chemical structure:



Formula Ib

10

wherein

R₁-R₇ taken independently or R₃-R₆ taken together may be a hydrogen atom, a halogen atom, a hydroxyl group, or any other organic groupings containing any number of carbon atoms, preferably 1 – 8 carbon atoms, and optionally include a heteroatom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats, representative R₁-R₆ groupings being alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-

25

C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, or polypeptide group;


X is a heteroatom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats;

5 Z is a heteroatom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats; and

Z' may a hydrogen atom, a halogen atom, a hydroxyl group, or any other organic composition containing from 1 – 8 carbon atoms and optionally include a heteroatom such as oxygen, sulfur, or nitrogen grouping

10 in linear, branched, or cyclic structural formats.

In still another embodiment, the lactones having an alpha-methylene group can have the structure as show below:



Formula Ic

wherein

15 R₁-R₉ taken independently or R₅ and R₆ taken together may be a hydrogen atom, a halogen atom, a hydroxyl group, or any other organic groupings containing any number of carbon atoms, preferably 1 – 8 carbon atoms, and optionally include a heteroatom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats,

20 representative R₁-R₆ groupings being alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio,

25 arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino,


substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, or
5 polypeptide group;

Y₁, Y₂, and Y₃ taken independently or Y₁ and Y₂ taken together may be a hydrogen atom, a halogen atom, a hydroxyl group, or any other organic groupings containing any number of carbon atoms, preferably 1 – 8 carbon atoms, and optionally include a heteroatom such as oxygen, sulfur, or
10 nitrogen grouping in linear, branched, or cyclic structural formats, representative R₁-R₆ groupings being alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted
15 aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted
20 phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, or polypeptide group;


Z is a heteratom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats; and

25 X is a heteratom such as oxygen, sulfur, or nitrogen grouping in linear, branched, or cyclic structural formats.

In one embodiment, the lactone is a securolide, which is a alpha-methylene-lactone (1) having the structure:




In another embodiment, the ester is methyl α -methylene- γ -hydroxybutanoate (2) as shown in the following structure:



5

In still another embodiment, the lactone is a bicyclic compound having the following structure:



(4')

10 B. Excipients

The lactone and functional derivatives can be formulated using standard techniques for enteral, parenteral, or topical administration. Effective dosages can be determined based on the *in vitro* assays known to those skilled in the art, such as the assays described in the examples.

15 Suitable pharmaceutically acceptable vehicles for parenteral delivery include sterile saline, phosphate buffered saline, and standard microparticulate formulations for injection, including polymeric microspheres, microcapsules, liposomes, and emulsions. These can include

degradable polymers such as polylactic acid and polyglycolic acid, and copolymers thereof, polyanhydrides, polyorthoesters, polyhydroxyalkanoates. For injection, the lactones will typically be formulated as solutions or suspensions in a liquid carrier.

5 For topical delivery, the lactone may be formulated in an ointment, lotion, gel, spray, or controlled or sustained release formulation (such as a transdermal patch).

For enteral delivery, the lactone may be formulated in a tablet, capsule, suspension or solution, dissolved or encapsulated in an excipient
10 such as a sugar like lactose, inert compound such as magnesium stearate, paraffin derivatives, glycols or gum arabic. The formulations may further include dyes, flavorings, preservatives, dispersing or emulsifying agents, or materials modifying release or stability properties of the formulations.

The active compound may be used in combination with a second
15 pharmaceutically acceptable antimicrobial agent, such as nitroimidazole antibiotics, e. g. tinidazole and metronidazole; tetracyclines, e. g. tetracycline, doxycycline and minocycline; penicillins, e. g. amoxicillin and mezlocillin, cephalosporins, e. g. cefaclor, cefadroxil, cephradine, cefuroxime, cefuroxime axetil, cephalexin, cefpodoxime proxetil,
20 ceftazidime and ceftriaxone; carbapenems, e. g. imipenem and meropenem; aminoglycosides, e. g. paromomycin, macrolide antibiotics, e. g. erythromycin, clarithromycin and azithromycin; lincosamide antibiotics, e. g. clindamycin; rifamycins g. rifampicin, and nitrofurantoin.

Combinations of the compounds with a pharmaceutical acid-lowering
25 agent may be used in the treatment of acid-related disorders, such as acid pump inhibitors, e. g., omeprazole and lansoprazole, or H₂ antagonists, e. g., ranitidine, cimetidine, and famotidine.

30 II. Synthesis of Lactones


The synthesis of the lactones and their respective derivatives involves the step of forming an alpha-methylene group. Generally, the method for

making a compound of Formulae Ia and Ic involves: a) providing a precursor having a lactone structure, and b) reacting the precursor with one or more chemical reagents to provide a product having a methylene group at the α -position of the lactone structure. The product can be treated with a
 5 nucleophilic agent such as an alcohol, alkoxide, amine, or any other neutral or anionic nucleophiles to generate a compound of Formula Ib.

Methods of forming the methylene group are standard techniques well documented in synthetic organic chemistry (see, for example, March, "Advanced Organic Chemistry," 4th Edition, 1992, Wiley-Interscience
 10 Publication, New York). The lactones can be synthesized, for example, by starting from a five-member lactone and then derivatizing the alpha position to add a methylene group onto the molecule.


For example, as shown by Scheme 1, reacting a compound having the structure (1) with a base, an alcohol and an ester produces an enolate, which
 15 is a precursor to **lactone** of structure (4) (See Scheme 1).

Scheme 1




The lactone (4) can be further derivatized using standard synthetic techniques available in the art of organic synthesis (see, for example, March, *Supra*). For example, carboxyl group functionalized lactones (4) can be
 20 prepared by, for example, reacting a compound having the structure (1) with an agent for alpha carboxylation of lactones (MMC, methyl-Mg-carbonate), and an aldehyde in the presence of a base such as triethylamine to yield the

lactone of structure (4), which is the isolated naturally occurring lactone (Scheme 2).




- In Scheme 2, a functionalized methylene group lactone (4), is
- 5 prepared by reacting a compound having the structure (1) with methyl-Mg-carbonate(MMC), an aldehyde in a buffer with a base NN-Dimethylformamide [DMF- $HCON(CH_3)_2$], to produce a carboxylic acid functionalized lactone (1B), and lactone (1B) is then treated with an aldehyde and Diethylamine, Sodium Acetate, Acetic Acid to produce lactone (4).
- 10 Other functionalized lactone derivatives can be readily prepared. For example, a functionalized derivative lactone (9) can be prepared by reacting a compound having the structure (5) with an inorganic acid and an alcohol by heating to reflux to produce an enol-lactone, which is then reduced with a base to produce a product of structure (7). Reaction of (7) with a base, an
- 15 alcohol and a ester produces an enolate, which is a precursor to functionalized lactones of compound (9) (Scheme 3).

Scheme 3



More functionalized lactones can be prepared by readily available synthetic method in the art (see, for example, March, "Advanced Organic Chemistry," 4th Edition, 1992, Wiley-Interscience Publication, New York).

- 5 For example, functionalized lactones (4) can be prepared by reacting a compound having the structure (1) with a base, an alcohol and a ester to produce an enolate, which is a precursor to *lactone* of structure (4) (Scheme 4).

Scheme 4

The pharmaceutically acceptable salts of the lactone compounds of the Formulae Ia-c, if in the form of an acid or a base such as an amine, can be prepared in a conventional manner by treating a solution or suspension of the compound of Formulae Ia-c with about one chemical equivalent of a pharmaceutically acceptable base or acid. Conventional concentration and recrystallization techniques are employed in isolating the salt.

III. Methods of Treatment

10 A. Disorders to be Treated

Based on the activities determined as described in the examples, the lactones are useful as anti-infectives and anti-proliferatives. In particular, the lactones can be administered in an effective amount to inhibit bacterial or fungal growth, viral disease or to treat a bacterial or fungal disease.

15 Examples of preferred bacterial disorders to be treated include peptic ulcer disease, gastritis, dyspepsia, periodontal disease and gingivitis. Fungicidal compositions are comprised of a fungicidally effective amount of a compound of formula 1 or a salt thereof and an inert pharmaceutical carrier. Examples of the pharmaceutical composition may be plain or sugar-coated

20 tablets, gelatin capsules, granules, suppositories, injectable preparations, ointments, creams or gels.

The fungicidal compositions are useful particularly on

Saccharomyces cerevisiae, *Candida albicans* and other *Candida* such as *Candida glabrata*, *krusei*, *tropicalis*, *pseudotropicalis* and *parapsilosis*, on *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus niger*, *Cryptococcus neoformans*, *Microsporum canis*, *Trichophyton rubrum*, *Trichophyton*
5 *mentagrophyte* and to combat particularly digestive, urinary, vaginal or cutaneous candidosis, cryptococcosis, for example, neuromeningeal, pulmonary or cutaneous cryptococcosis, bronchopulmonary and pulmonary aspergillosis and invasive aspergillosis of the immunodepressive system. The compositions can also be used in the prevention of mycotic affections in
10 congenital or acquired immunological suppressions.

The lactones can also be administered to treat proliferative disorders, including cancers. Representative types of cancers which have shown inhibition in cell growth or proliferation include breast cancer, lung cancer, ovarian cancer, esophageal cancer, and leukemia. Other types of abnormal
15 proliferative disorders that the lactones may be useful in the treatment of include endometriosis and restenosis, caused by abnormal overproliferation of endothelial tissue following angioplasty.

The functionalized lactone (9) above is particularly useful as an aspartic protease inhibitor, and has been demonstrated to inhibit the aspartic
20 protease of HIV-1, and thereby limit viral processing, and in particular post-translational processing of the viral gene products (gag / gag-pol).

These lactones are also useful in modulating pain response through its activity as a neurotransmitter.

B. Dosages

25 The effective amount will be determined based on the disease or disorder to be treated, the mode of administration and the formulation. Effective dosages can be routinely determined based on the effective dosages determined using *in vitro* assays such as those described in the examples.

The high activity of Securolide(LMSV-6) against *Escherichia coli*,
30 *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and its low molecular weight are advantageous. Advantages of Securolide include its facility and speediness to promote pharmacologic response; its

possibility to cross over cellular membrane barriers, where high molecular weight is the main hindrance, and its potent activity against *Pseudomonas*, which is one of the more drug resistant microorganisms.

5 The method for combating fungal infections comprises administering a fungicidally effective amount of a compound of formula I or an acid addition salt thereof by buccal, rectal, parental route, or by local route as a topical application on the skin and mucous membranes, but the preferred route is the buccal route. The usual daily dose is 1 to 5 mg/kg depending on the method of administration, the condition treated and the specific
10 compound.

The compounds may be administered alone, but will generally be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they can be administered orally or in the form of tablets
15 containing such excipients as starch or lactose, or in capsules either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. In the case of animals, they are advantageously contained in an animal feed or drinking liquid in a concentration of about 5 to 5000 ppm, preferably about 25 to about 500 ppm
20 They can be injected parenterally, for example, intramuscularly, intravenously or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which can contain other solutes, for example, enough salt or glucose to make the solution isotonic. In the case of animals, compounds can be administered intramuscularly or
25 subcutaneously at dosage levels of about 0.1 to about 50 mg/kg/day, preferably about 0.2 to about 10 mg/kg/day given in a single daily dose or up to 4 divided doses.

The compounds can be administered to humans for the treatment of *H. pylori* infections by either the oral or parenteral routes and may be
30 administered orally at dosage levels of about 0.1 to about 50 mg/kg, advantageously about 0.5 to 50 mg/kg/day given in a single dose or up to 4 divided doses. For intramuscularly or intravenous administration, dose levels

are about 0.1 to about 100 mg/kg/day, preferably about 0.5 to about 50 mg/kg/day. While intramuscularly administration may be a single dose or up to 4 divided doses, intravenous administration can include a continuous drip. Variations will necessarily occur depending on the weight and condition of the subject being treated and the particular route of administration chosen as will be known to those skilled in the art.

The second antimicrobial agent and the acid-lowering agent may be administered with the compounds in the same manner as discussed above for the compounds of the invention. Thus, depending on the particular agent, administration may be orally at about 0.1 to about 500 mg/kg, for instance at about 1 to 3 grams per day of second antimicrobial agent, and about 40 to 80 mg per day of the acid-lowering agent, or by injection at about 0.1 to about 200 mg/kg/day.

The present invention will be further understood by reference to following non-limiting examples.

Example 1: Isolation of Lactone from *Securidaca virgata*.

Securidaca virgata roots were collected at San Cristobal Dominican Republic. A sample voucher of aerial part was sent to National Botanic Garden in order to be identified. Air dried, chopped and milled roots (253 g) were extracted in a Soxhlet apparatus. The cold extract was filtered and then concentrated *in vacuo* (55°C) to afford a syrupy liquid (29.76 g) that means 11.76% of yield. This syrup (28 g, tested microbiologically) was treated with 300 ml of Sulfuric acid 0.5M and stirred for 4 hours and then extracted in a liquid-liquid extraction with chloroform successively until the syrup was separated from the acid chloroform soluble matter. Then the hydrosoluble phase was basified with Ammonium hydroxide 20% (pH =8 -9) and the hydrosoluble phase extracted again with chloroform. The chloroform extract was concentrated *in vacuo* (50°C), to yield an extract of 5.92 g (yield =21.14% with regard to crude extract and 2.33% with regard to roots extracted). The 5.92 g had a purity of 95.34 % LMSV-6 (SecurosideTM). The final chloroform extract (5.92 g) were chromatographed in basic alumina (63 -150 µm) and eluted with petroleum ether-ethylacetate mixture (30:70).

Fractions collected were concentrated to have a pure clear liquid substance (Securolide™) (1). Chromatographic fractions 72-97 (674 mg) from the liquid chromatographic column where (1) was isolated was again chromatographed to obtain an amber colored liquid (2). Compound (1) and
 5 (2) were then tested in bioassays.

Spectroscopic analysis yielded the following data for compound (1):

PHYSICAL DATA

RELATIVE DENSITY: 1.070 g/mL pH =5.853

REFRACTIVE INDEX =1.471

10 SPECTRAL DATA

Mass spectrometry: MS m/z (rel. Int.): molecular ion 98.0 [M^+], (54), 68 (100), molecular formula $C_5H_6O_2$.

Infrared Spectroscopy (IR): 1761.6cm^{-1} (lactone), 1667cm^{-1} (C=C) 810cm^{-1} (C=CH₂).

15 *Proton Nuclear Magnetic Resonance* (NMR¹H, 200 MHz, CDCl₃): δ 2.98 ppm (2 H multiplet), 4.37 ppm (1H triplet, J=7.4 Hz), 5.65 (1H triplet , J=2,6 Hz), 6.22 (1H triplet , J=2.6,)

Carbon 13 Nuclear Magnetic Resonance (NMR¹³ C 200 MHz, CDCl₃): δ 170.7 (C-1), 133.3 (C-2), 27.4 (C-3), 65.5 (C-4), 122.2 (C-5)

20 *Double Nuclear Magnetic Resonance Protonic Experiments (Irradiations)*
 double Nuclear Magnetic Resonance Proton Experiments showed coupling between each one of existing protons. Methylenic exocyclic protons δ (5.65 and 6.22), irradiation of C-3 showed coupling with C-4 and in this way were determined all protons connectivity of (1).

25 Spectroscopic analysis yielded the following data for compound (2):

Mass Spectrometry: 130.0 (M⁺, 3), 113 (M⁺-OH, 12), $C_6H_{10}O_3$.

Infrared Resonance (IR): 3614.6cm^{-1} (OH), 1629cm^{-1} (C=C) 1150.9cm^{-1} (-CO-OCH₃), 810cm^{-1} (C=CH₂).

30 *Proton Magnetic Nuclear Resonance Spectrometry* (200 MHz, CDCl₃): δ 2.54 ppm (2 H td J=6.26Hz), 3.73 (s), 3.72 (2H, t J=6.26Hz), 5.65 (1H t, J=1.2Hz), 6.21 (1H t, J=1.2Hz).

Carbon ¹³ Magnetic Nuclear Resonance Spectrometry (200 MHz, CDCl₃): δ

167.8 (C-1), 137.2 (C-2), 35.4 (C-3), 61.3 (C-4), 127.3 (C-5), 51.93 (C-6).

Ampoules consisting of a pharmaceutical composition containing the compound of the formula 1 (Securolide) had the following ingredients:

Original Quantity:		750 Amp	
5	<u>Raw material</u>	<u>Weight</u>	<u>UM</u>
	LMSV-6 (Securolide)	51.7900 mL	6.27%
	Sesame oil	760.8200 mL	92.20%
	Bencilic Alcohol	12.3750 mL	1.50%
	U. S. P.		
10	Total Volume	824.9850 mL	99.97%

Example 2: Assays for Anti-bacterial Activity.

The material obtained in Example 1 was screened for antibiotic activity against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Klebsiella pneumoniae*.

15 Materials and Methods:

Solvent: Buffer Phosphate 0.1 N, pH=8.0

Antibiotic: LMSV-6 (Pure, Securolide)

Medium coat: 2 mL/100 mL

Inoculum: 4mL/Petri Dish

20 Preparation of Culture Medium:

Culture Medium for *Staphylococcus aureus* (USP 23 <81>)

	Peptone	1.0g
	Digestive pancreatic Caseine	4.0g
	Yeast Extract	3.0g
25	Beef Extract	1.5g
	Dextrose	1.0g
	Agar	15.0g
	pH after sterilization $6.6 \pm .01$	

Culture Medium for *Pseudomonas aeruginosa* (USP 23 <81>)

30	Pancreatic digestive Casein	17.0g
	soy Papain digestive	3.0g
	Sodium Chloride	5.0g

- Dibasic Potassium Phosphate 2.5g
 Dextrose 2.5g
 Agar 20.0g
 Water to make about 1.0 Lit
- 5 pH after sterilized 7.2 ± 0.1
- Culture Medium for *Escherichia coli* and *Klebsiella pneumoniae* (USP 23
<81>)
- Peptone 5.0g
 Yeast Extract 1.5g
- 10 Beef Extract 1.5g
 Sodium Chloride 3.5g
 Dextrose 1.0g
 Dibasic Potassium Phosphate 3.68g
 Monobasic Potassium Phosphate 1.32g
- 15 Water To make about 1.0 Lit
 pH after sterilized 7.0 ± 0.05

Bacteria	LMSV-6 (puro)	Concentration ($\mu\text{L}/10\text{mL}$)							
		0	20	30	40	Ciprof	osf	is(-)	Dis(+)
<i>E. coli</i>	30(mm)	-	-	-	-	28	10	-	-
<i>Kleb</i>	40	-	-	-	-	26	22	-	-
<i>Ps. A</i>	30	-	-	-	-	12	6	-	-
<i>St. a</i>	30	-	-	-	-	20	18	-	-
<i>Str. P</i>	-	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

Unit to measure zone of inhibition is millimeters (mm)

- 20 Ciprof =Disc of sensitivity Ciprofloxacina 5 micrograms
 Fosf =Disc of sensitivity Fosfomicina 50 micrograms
 Dis(-)=Disc of sensitivity with no substance applied
 Dis(+)=Disc of sensitivity with 20 microliters of Tween 60 at 2%
 N/A =not done Test
- 25 Microorganisms used were standardized by American Type Culture

Collection (ATCC), used in first generation:

E. coli ATCC 352 18, Lot 202602, Exp. 05/2000 (19-258)

*Kleb. *Klebsiella pneumoniae*, ATCC 13882, Lot 202 174, Exp 02/2000 (19-152)

5 *Ps. A *Pseudomona aeruginosa* ATCC 27853, Lot 202992, Exp 08/2000 (19-060)

*St. A *Staphylococcus aureus* ATCC 12598, Lot 202564, Exp 05/2000 (19-137)

10 **Str. P *Streptococcus pyogenes*, ATCC 2 1547, Lote 202691, Exp 06/2000 (19-190)

*=Atmospheric oxygen, Temp = $35 \pm 2^{\circ}\text{C}$

**=Atmospheric CO₂, Temp = $35 \pm 2^{\circ}\text{C}$

Results:

15 5 microliters of LMSV-6 (Securolide) inhibits growth of *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

Dilutions :

10 microliters/10 mL, (0.001 microliters/ml)

20 microliters/10 mL, (0.002 microliters/mL)

20 30 microliters/10 mL (0.003 microliters/mL)
and 40 microliters mL(0.004 microliters/mL)

of LMSV-6 in 10 mL Tween 60 -2% did not exhibit inhibition of bacterial growth, indicating that these concentrations are below the Minimum Inhibitory concentration (MCI).

25 **Example 3: Determination of Minimum Bacterial (*Sarcina lutea* and Inhibitory Concentration, MIC, LMSV-6 (Securolide).**

Materials and Methods:

Methodology: Poured in plate, medium Mueller Hinton pH=8

Buffer pH=8 for ceftriaxone pattern dilution

30 Tween 20 at 2% for LMSV-6 (Securolide) dilution

Microorganism: *Sarcina lutea*

Sample(s): LMSV-6 (16.0 microliters/100 mL and 10 microliter applied to sensitivity disc).

Methodology: Poured in plate, medium Mueller Hinton p&8

Buffer pH=8 for ceftriaxone pattern dilution

5 Tween 20 at 2% for LMSV-6 (Securolide) dilution

Used Microorganism : *Sarcina lutea*

Sample(S): LMSV-6 (10 microliters/mL and 20 microliter applied to sensitivity disc)

Control Pattern (P): Ceftriaxone (disc with 10 mcg)

10 **Results:**

<u>APPLIED VOLUME TO DISC</u>	<u>ZONE OF INHIBITION</u>
5 microliters	42.0 mm
1.6 microliters	20.0 mm

There was no significant zone of inhibition, therefore the Minimum

15 Inhibitory Concentration is close to 0.2 microliters of LMSV-6.

The approximate MIC is 0.2 microliters of LMSV-6 3.3 mm inhibition

Example 4: *In vivo* Treatment of Bacterial Infection.

Rats were infected in surgery and then successfully treated with Securolide using doses calculated based on comparison with ceftriaxone.

20 **Example 5: Assays for Anti-neoplastic Activity**

Materials and Methods:

There are a large number of biological assays that require the measurement of surviving or proliferating mammalian cells. This can be achieved by different methods, e. g. counting cells that exclude or include a dye; measuring released ⁵¹Cr labeled protein after cell lysis; and measuring incorporation of radioactive nucleotides radioactive [³H]thymidine) or [¹²⁵I]iododeoxyuridine) during cells proliferation. Viable cells can be measured by using any of several staining methods, but a more accurate method is the multiwell scanning spectrophotometers (ELISA readers) which can measure large number of samples with a high degree of precision.

A colorimetric assay to detect living cells, utilizing a colorless substrate that is modified to a colored product by any living cell, but not by

dead cells or tissue culture medium, was used. The assay uses Tetrazolium salts MTT [3-(4,5-dimethylthiazol-2-yl) -2,5 diphenyl tetrazolium bromide] to measure the activity of various dehydrogenase enzymes (Slater et al. 1963). The Tetrazolium ring is cleaved by active mitochondria, so this
5 reaction only occurs in living cells. The cleavage of MTT into Formazan by the mitochondrial enzyme succinate dehydrogenase produces a dark blue colored compound (Formazan) whose amount is proportional to the number of cells present. The Formazan concentration in a gradient of Securoside concentrations was measured by multiwells scanning spectrophotometer
10 (ELISA reader). Data analysis was then used to establish the Inhibitory concentration fifty (IC₅₀) which is a quantitative parameter of antineoplastic activity. Different cancer cells lines were tested in order to measure indirectly the cytotoxic or antineoplastic activity (surviving cells) of LMSV-6 (Securoside). Cancer cell lines that were tested included HEP 2 (Laringe
15 carcinoma) and HELA (Cervix Carcinoma).

MATERIALS:

Culture Medium (DMEN +all)
EDTA
Trypsin-EDTA
20 DMEM + ALL and 10% TERNERO RECENTAL SERUM.
Dimethyl sulfoxide
MTT [3-(4,5-dimethylthiazol-2-yl) -2,5-diphenyl tetrazolium bromide]
Multiwell scanning spectrophotometers (ELISA readers)

METHODS:

25 Cells were cultured using standard techniques. See, for example, R. Ian Freshney "Culture of Animal Cells" A Manual of Basic Technique (Alan R Liss, Inc., New York, Second Edition) p. 245-256.

Assessment of cell growth inhibition was determined according to the methods of Skehan et al, J. Nat. Cancer Inst. 82, 1107 (1990). Cells were
30 plated between 400 and 1200 cells/well in 96 well plates and incubated at 37°C for 15-18 h prior to drug addition to allow attachment of cells.
Compounds tested were solubilized in 100% DMSO and further diluted in

RPML-1640 containing 10 mM HEPES. Each cell line was treated with 10 concentrations of compounds (5 log range). After a 72 H incubation, 100 ml of ice-cold 50%TCA was added to each well and incubated for 1 h at 4°C. Plates were then washed 5 times with tap water to remove TCA, low-
5 molecular-weight metabolites and serum proteins. Sulforhodamine B (SRB) (0.4%, 50 mL) was added to each well. Following a 5 min incubation at room temperature, plates were rinsed 5 times with 0.1% acetic acid and air dried. Bound dye was solubilized with 10 mM Tris Base (pH 10.5) for 5 min on a gyratory shaker. Optical density was measured at 570 nm.

10 See also Gerlier, et al. *J. Immunol. Meth.* 94, 57-63 (1986); Slater, et al., *Biochim. Biophys. Acta*, 77, 383-393 (1963); Kasugai, et al., *Japan J: Pharmacol.* 52,95-100 (1990); Mosmann *J. Immunol. Methods*, 65, 55-63 (1983); Denizot, et al., *J. Immunol. Methods*, 89, 271-277 (1986).

Data were fit with the Sigmoid-Emax concentration-Effect model
15 (see Holford, N. H. G.: Scheiner, L. B., "Understanding the dose-effect relationship: Clinical applications of pharmaco-kinetic-pharmacodynamic models.", *Clin. Pharmacokin.* 1981, 6,429-453) with non-linear regression, weighted by the reciprocal of the square of the predicted response. The fitting software was developed by the Roswell Park Institute with Microsoft
20 FORTRAN, and uses the Marquardt algorithm (see Marquardt, D. W., "An algorithm for least squares estimation of nonlinear parameters", *J. Soc. Ind. Appl. Math.* 1963, 11, 431-441) as adopted by Nash (see Nash J. C., "Compact numerical method for computers: Linear algebra and function minimization". John Wiley & Sons, New York, 1979) for the non-linear
25 regression. The concentration of drug which resulted in 50% growth inhibition (IC₅₀) was calculated.

Results:

LMSV-6 CYTOTOXIC ASSAY IN HEP-2 CELLS POPULATION OF 10⁵ Cells/mL (LARINGE CARCINOMA)

30 r=0.9878

IC₅₀=10.38 µ/mL

**LMSV-6 CYTOTOXIC ASSAY IN HEP-2 CELLS POPULATION OF
10⁶ Cells/mL (LARINGE CARCINOMA)**

r=0.9941

IC₅₀=37.37 µg/mL

**5 LMSV-6 CYTOTOXIC ASSAY IN HELA CELLS POPULATION OF
10⁵ Cells/mL (CERVIX CARCINOMA)**

r=0.9950

IC₅₀=7.37 µg/mL

**10 LMSV-6 CYTOTOXIC ASSAY IN HELA CELLS POPULATION OF
10⁶ Cells/mL (CERVIX CARCINOMA)**

r=0.9941

IC₅₀=27.30 µg/mL

Example 6: Testing for Inhibition of *H. pylori*

Materials and Methods:

- 15 Agar Dilution of Antimicrobial compound: 6 mg. of the compound
to be evaluated is solubilized in 0.6 ml 100% dimethylsulfoxide (DMSO)
and then brought up to 6 ml with sterile brucella broth. The final
concentration of DMSO is 10% of the total volume. Serial 2-fold dilutions (3
ml Securolide +3 ml brucella broth) are then made in sterile brucella broth. A
20 2 ml aliquot of each broth dilution within the series is placed in separate
sterile petri dishes, to which 18 ml of melted and cooled (approx. 50°C)
brucella agar supplemented with 7% horse blood is added. This yields a final
1:10 dilution of Securolide in agar, and a final concentration of DMSO of
1%. For example, if the highest concentration of drug in agar is 100 µg/ml.
25 Agar plates can be prepared one day prior to inoculating, and refrigerated
overnight.

- Inocula Preparation: *Helicobacter pylori* cultures were maintained on
trypticase soy-5% sheep blood agar plates, and transferred every 48 hours.
Helicobacter mustelae cultures were maintained on the same agar and
30 transferred every 48-60 hours, depending upon the extent of the growth of
the previous transfer. Plates are incubated at 37°C in GasPak jars with water-

activated (10 ml) CampyPak Plus (BBL Microbio. Systems) envelopes with palladium catalyst.

Helicobacter cultures can be grown in *brucella* broth supplemented with 10% fetal calf serum in 10 ml volumes in deep petri dishes. The plates
5 are incubated for 18-20 hours at 37°C in GasPak jars with water-activated (10 ml) CampyPak Plus envelopes with palladium catalyst on a shaker at 50 rpm.

Overnight cultures (approximately 10^8 CFU/ml) are diluted 10-fold in *brucella* broth (no FCS) in screwcapped tubes for use as the standard
10 inoculum. The wells of a Steer replicator are filled with 0.8 ml of the diluted organism, and approximately 2×10^4 cells in 0.002 ml are placed on the agar surface. Inoculated plates are placed in a GasPak jar to which water-activated (10 ml) Campy Pak Plus envelopes with palladium catalyst have been added, and incubated at 37°C. for 48 hours.

15 **Results:**

Following incubation, all test plates are compared to a Securolide-free growth control plate. The MIC is the concentration which inhibits growth compared to the control plate. A thin film of growth might be visible at higher concentrations but this is discounted, and not considered the true
20 MIC. Control organisms are also inoculated on each plate, and these are diluted 1000-fold for use as inocula. The control organisms include *Campylobacter jejuni*, and the screening cultures of *E. coli* [ATCC 35218, Lot 202602, Exp 05/2000 (19-258)], *Pseudomonas aeruginosa* [ATCC 27853, Lot 202992, Exp 08/2000 (19-060)], *E. cloacae*, *Providencia stuartii*
25 and *P. rettgeri*. Plates and/or inocula transfers should not be out of the microaerophilic environment longer than 2 hours. All manipulations involving *Helicobacter* cultures was performed in a laminar flow hood to decrease the chance of contaminating the cultures with mold.

The mouse model of Lee et al., Gastroenterology, 99, 1315-23 (1990)
30 is used to predict the *in vivo* activity of a compound against *H. pylori* in humans.

Helicobacter felis is grown in *brucella* broth with 10% fetal bovine

serum. A frozen culture is quickly thawed; the culture is checked for motility and 0.5 mL of the thawed frozen culture is inoculated into a deep tissue culture dish containing 9.5 mL of the brucella/serum mix. The dishes are put into a Cappy Pak jar [BBL] to insure a microaerophilic atmosphere.

5 The jar is put on a rotary shaker at 60 RPM in a 37°C incubator. After 18 hours there should be visible turbidity. The culture is checked for purity and motility under a (phase) microscope and then pooled into a flask. Swiss-Webster female mice (18-20 g) are fasted for 18 hours before infection. The mice are infected three times on alternate days during a single week. Dosing

10 begins two weeks after the last dose of organism. Treatments are given once per day for fourteen consecutive days. Sacrifice is about three weeks after completion of therapy. For each mouse, the stomach is excised and opened along the greater curvature. A plug (a 3 mm Tissue section) is taken from the antrum region of the stomach. The plug surfaced is washed, minced, and

15 dropped into a tube with 100 microliters of urease reagent. The urease reagent (pH 6.3-6.5) contains urea and phenol red. If *Helicobacter* is present, urease will break down urea producing a change of pH. Purple (alkaline) is positive for *Helicobacter*; red/yellow (no change) is negative. Any color change is recorded after 18 hours. There are usually twenty mice per

20 treatment group; the percent positive for each group is recorded.

There are several methods used clinically to determine whether *Helicobacter pylori* is present in a human subject. These are employed for initial diagnosis of infection prior to treatment, as well as for determining the success of treatment in eradicating the organism from the patient.

25 The urea breath test involves ingestion of radiolabelled urea. *H. pylori* produces an urease enzyme which degrades urea; mammalian gastric cells do not contain this enzyme. Exhalation of labeled carbon dioxide (analyzed by mass spectrometry or radioactivity, depending on the isotope employed) therefore indicates that *H. pylori* is present.

30 Serology can also be used to assess infection with *H. pylori*. Detection of serum antibodies to *H. pylori*, such as IgG and IgA, is carried out using enzyme-linked immunosorbent assay (ELISA). Numerous different

H. pylori proteins can be employed as antigens.

Endoscopy of the patient provides samples of tissue which can be cultured in a microaerophilic environment to diagnose *H. pylori* infection. Alternatively, the sample can be examined histologically by employing one of a number of stains such as Giemsa or hematoxylin-eosin. An urea test which takes advantage of the production of urease by *H. pylori*, can also be applied. This test relies on the formation of ammonia from the urea hydrolysis, which results in an observable change in pH, that is an indication of the eradication of *H. pylori*.

Results:

Comparison of patient condition by endoscopy before and after treatment with the compound of example 1 has shown a decrease or absence of *H. pylori*.

Example 7: Determination of Fungal Activity

Materials and Methods:

METHODS

Anti-fungal Activity of the Product may be determined as follow. Female mice weighing 18 to 22 g were used and a quantity of *Candida Albicans* 44858 was administered into a vein in the tail at the rate of 10^6 CFU per mouse (CFU: colony forming unit). The mice were separated into 5 batches of 5 mice and they were treated in the following manner:

One Hour After Infection

group 1: the mice were treated with product at 25 mg/kg orally

group 2: the mice were treated with product intraperitoneally at the

dose of 25 mg/kg

group 3: the mice were treated orally with Ketoconazole at 25 mg/kg

group 4: the mice were treated intraperitoneally with Ketoconazole at a dose of 25 mg/kg

group 5: the mice did not receive any anti-fungal treatment.

The dead mice were counted over a period of 22 days.

Results:

The activity of product was excellent at the dose used in the two

administration methods. The same treatments were also effective in the "topical model" with dermal fungi, for example trichophyton, and in the sublethal model.

Minimal Inhibitory Concentration (MIC)

5 *Candida albicans* cells were prepared as indicated in the J. Antimicrobial Chemotherapy 38, 579-587, and were washed 3 times with a 0.1 M phosphate solution and used immediately to determine the minimal inhibitory concentration (MIC). The MICs were determined by modification of a microtitre plate according to the standard method of the laboratory
10 clinical standards of the Comité National.

 RPMI-1640, and L-glutamine buffered at pH 7 with a 0.15 M solution of MOPS (3-[N-morpholino]propane sulfonic acid). *Candida albicans* cells (1.5×10^3 cells/ml) were added to the wells of a 96-well plate containing RPMI-1640 and dilutions of antifungal agents. The results were read 48
15 hours after incubation at 35°C and the MIC or the minimal inhibitory concentration which inhibited the growth of the *Candida albicans* cells was determined.

Minimal Fungicidal Concentration

 After reading the MIC at 48 hours, the plates were shaken and 10 µl
20 of well aliquot was removed from the wells which was placed on rectangular disks containing dextrose sugar. The plates were incubated for 48 hours at 35°C and the minimal fungicidal concentration and the concentration of the antifungal agent at which there were no number of colony forming units.

 Concomitant assays were made where LMSV-6 showed high activity
25 against *Candida albicans*.

Example 8: Treatment of Periodontal Disease


 It has been found that the Securoside is selectively effective against the specific anaerobic gram negative organisms associated with gingivitis, when topically applied to the affected gingiva.

30 The minimum inhibitory concentration (MIC) needed to kill *Bacteroides assaccharolyticus* (Forsyth strain), determined according to the procedure of Walker et al (Antimicrobial and Chemotherapy, Vol. 16, p.


452-457, (1979), is between 6 to 60 micrograms per ml. The MIC value for *B. gingivalis* is 6 micrograms per ml. The MIC value for *B. gracilis* and *Fusobacterium*, other gram negative organisms is about 25 for each; and the MIC value for the gram positive organism *Actinomyces viscosus* is 60 for
5 the aerobic strain and 60 for the anaerobic strain.

I claim:

1. An isolated compound defined by one of the following structures:



Formula Ia



Formula Ib


wherein R₁-R₇ taken independently or R₃-R₆ taken together are a hydrogen atom or a group or grouping selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;

Z is a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats;

Z' is a hydrogen atom, a halogen atom, a hydroxyl group, or an organic group having 1-8 carbon atoms which includes a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats, or

an isolated lactone defined by the following structure:



Formula Ic

wherein

R₁-R₉ taken independently or R₅ and R₆ taken together may be a hydrogen atom or any other groups or groupings selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;


Y₁, Y₂, and Y₃ taken independently or Y₁ and Y₂ taken together may be a hydrogen or any other groups or groupings selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20

cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;


Z is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats.


2. The compound of claim 1 defined by Formula Ia.
3. The compound of claim 2 wherein the X is an oxygen.
4. The compound of claim 2 wherein R₅ is an exocyclic methylene.
5. The compound of claim 1 defined by Formula Ib.
6. The compound of claim 1 which has one of the following structure.



7. The compound of claim 1 of Formula Ia wherein the lactone is isolated from a *Securidaca virgata*.
8. The compound of claim 1 of Formula Ia further comprising a pharmaceutically acceptable carrier for parenteral, enteral, or topical administration.
9. A method of treating a patient in need thereof, comprising administering an effective amount of the compound defined by one of the following structures:



Formula Ia



Formula Ib

wherein R₁-R₇ taken independently or R₃-R₆ taken together are a hydrogen atom


or a group or grouping selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;

Z is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats;

Z' is a hydrogen atom, a halogen atom, a hydroxyl group, or an organic group having 1-8 carbon atoms which includes a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats, or

an isolated lactone defined by the following structure:



Formula Ic

wherein

R₁-R₉ taken independently or R₅ and R₆ taken together may be a hydrogen atom or any other groups or groupings selected from the group consisting of


alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;

Y_1 , Y_2 , and Y_3 taken independently or Y_1 and Y_2 taken together may be a hydrogen or any other groups or groupings selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;


Z is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats.

10. The method of claim 9 wherein X is oxygen.
11. The method of claim 9 wherein R_5 is an exocyclic methylene.
12. The method of claim 9 wherein the compound has one of following structures:



13. The method of claim 8 wherein the compound of Formula Ia is administered orally, parenterally, topically, or transdermally.
14. The method of claim 9 wherein the patient has a proliferation disorder.
15. The method of claim 14 wherein the proliferation disorder is cancer.
16. The method of claim 9 wherein the patient has an infection.
17. The method of claim 16 wherein the infection is bacterial.
18. The method of claim 16 wherein the infection is fungal.
19. The method of claim 9 wherein the patient has peptic ulcer disease.
20. The method of claim 19 wherein the peptic ulcer disease is linked to the presence of *Helicobacter pylori*.
21. The method of claim 9 wherein the patient is in pain.
22. The method of claim 9 further comprising administering the compound of Formula Ia in a pharmaceutically acceptable carrier.
23. The method of claim 22 wherein the carrier for oral administration is a mouthwash, lozenge, tablet, capsule, solution, suspension, granule, or any combination thereof.
24. The method of claim 22 wherein the carrier for topical administration is a suppository, ointment, cream, gel, paste, collodion, glycerogelatin, liniment, lotion, paste, plaster, powder, tape, patch, aerosol, solution, tincture, or any combination thereof.
25. A process of isolating a lactone from a plant, the lactone being defined by the following structure:



wherein R₁-R₇ taken independently or R₃-R₆ taken together are a hydrogen atom or a group or grouping selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;


Z is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats, comprising extracting the lactone from plant material.


26. The process of claim 25 wherein the plant is *Securidaca virgata*.

27. The process of claim 25 wherein the method of isolation is extraction followed by chromatography.

28. A synthetic compound defined by one of the following structures:



Formula Ia



Formula Ib


wherein R₁-R₇ taken independently or R₃-R₆ taken together are a hydrogen atom or a group or grouping selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;

Z is a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats;

Z' is a hydrogen atom, a halogen atom, a hydroxyl group, or an organic group having 1-8 carbon atoms which includes a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteroatom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats, or

an isolated lactone defined by the following structure:



Formula Ic

wherein

R₁-R₉ taken independently or R₅ and R₆ taken together may be a hydrogen atom or any other groups or groupings selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl, sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;


Y₁, Y₂, and Y₃ taken independently or Y₁ and Y₂ taken together may be a hydrogen or any other groups or groupings selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, phenyl, substituted phenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, halo, hydroxyl, alkoxy, substituted alkoxy, phenoxy, substituted phenoxy, aroxy, substituted aroxy, alkylthio, substituted alkylthio, phenylthio, substituted phenylthio, arylthio, substituted arylthio, cyano, isocyano, substituted isocyano, carbonyl, substituted carbonyl, carboxyl, substituted carboxyl, amino, substituted amino, amido, substituted amido, sulfonyl, substituted sulfonyl,

sulfonic acid, phosphoryl, substituted phosphoryl, phosphonyl, substituted phosphonyl, polyaryl, substituted polyaryl, C1-C20 cyclic, substituted C1-C20 cyclic, heterocyclic, substituted heterocyclic, aminoacid, peptide, and polypeptide groups;

Z is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats; and

X is a heteratom selected from the group consisting of oxygen, sulfur, and nitrogen groupings in linear, branched, or cyclic structural formats.

29. The compound of claim 28 defined by any of the following structure:



30. A method of making a compound of Formulae Ia and Ic comprising:

- a) providing a precursor having a lactone structure, and
- b) reacting the precursor with one or more chemical reagents to provide a product having a methylene group at the α -position of the lactone structure.

31. A method of making a compound of Formula Ib comprising:

- a) providing a precursor having a lactone structure,
- b) reacting the precursor with one or more chemical reagents to provide a product having a methylene group at the α -position of the lactone structure, and
- c) reacting the product with a nucleophile.