(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
7 June 2012 (07.06.2012)

WIPO | PCT

(10) International Publication Number
WO 2012/075253 A2

(51) International Patent Classification:
A61K 9/16 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:
PCT/US2011/062837

(22) International Filing Date:
1 December 2011 (01.12.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/419,538 3 December 2010 (03.12.2010) US
61/436,324 26 January 2011 (26.01.2011) US

(71) Applicant *(for all designated States except US)*: **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **CAO, Yu** [CN/US]; Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936-1080 (US). **HACKL, Wolfgang** [AT/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **LI, Ping** [US/US]; Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936-1080 (US). **LI, Shoufeng** [CN/US]; Novartis Pharmaceiticals Corporation, One Health Plaza, East Hanover, New Jersey 07936-1080 (US). **MIRZA, Tahseen** [US/US]; 14232 Arbor Forest Drive, Apt. 202, Rockville, Maryland 20850 (US). **ZEHENDER, Hartmut** [DE/FR]; 16A Rue du Winkel, F-68210 Fulleren (FR). **ZHU, Jiahao** [CN/US]; Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936-1080 (US).

(74) Agent: **RUECK, Sandra**; Novartis, Patent Department, One Health Plaza, East Hanover, New Jersey 07936-1080 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

— *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

WO 2012/075253 A2

(54) **Title**: PHARMACEUTICAL COMPOSITIONS

(57) **Abstract**: The present invention relates to a pharmaceutical composition for the oral administration of a therapeutic compound of formula (I), which comprises granules that comprise at least therapeutic compound of formula (I) (see below), particularly 2-Methyl-2-[4-(3-methyl-2-oxo-8- quinolin-3-yl-2,3-dihydro-imidazo[4,5-c]quinolin-1-yl)-phenyl]-propionitrile or 8-(6-methoxy-pyridin-3-yl)-3-methyl-1-(4-piperazin-1-yl-3-trifluoromethyl-phenyl)-1,3-dihydro-imidazo[4,5-c]quinolin-2-one, or a tautomer thereof, or a pharmaceutically acceptable salt, or a hydrate or solvate thereof; at least one non-ionic surfactant that is Vitamin E-TP-GS in an amount ranging from about 15 to about 80% by weight of the composition; and at least one a dissolution enhancing agent selected from polyethylene glycol, polyethylene oxide, and any combination of the foregoing. The present invention also relates to processes for making such pharmaceutical compositions; a kit comprising such pharmaceutical composition and the instructions provide that the pharmaceutical composition may be taken immediately to about thirty minutes after the consumption of food; and related uses and methods of treatment.