


Y. et al. Liver Int 2009, 29(5), 678-85) with a relevant expression of CB2 in myofibroblasts, the cells responsible for fibrosis progression.

Activation of CB2 receptor by selective CB2 agonist has in fact been shown to exert anti-fibrotic effect in diffuse systemic sclerosis (Garcia-Gonzalez, E. et al. Rheumatology (Oxford) 2009, 48(9), 1050-6) and CB2 receptor has emerged as a critical target in experimental dermal fibrosis (Akhmetshina, A. et al. Arthritis Rheum 2009, 60(4), 1129-36) and in liver pathophysiology, including fibrogenesis associated with chronic liver diseases (Lotersztajn, S. et al. Gastroenterol Clin Biol 2007, 31(3), 255-8; Mallat, A. et al. Expert Opin Ther Targets 2007, 11(3), 403-9; Lotersztajn, S. et al. Br J Pharmacol 2008, 153(2), 286-9).

The compounds of the disclosure bind to and modulate the CB2 receptor and have lower CB1 receptor activity.

In one aspect, there is provided a compound of formula (I)



wherein


A is CH_2 or absent;

R^1 and R^2 , together with the nitrogen atom to which they are attached form substituted pyrrolidinyl or 2-oxa-6-azaspiro[3.3]heptyl, wherein substituted pyrrolidinyl is pyrrolidinyl substituted with one or two substituents independently selected from halogen, hydroxyl and alkylcarbonyloxy; and

R^3 is halophenyl, alkylsulfonylphenyl, haloalkylphenyl, halopyridinyl, alkyloxadiazolyl, alkyltriazolyl, alkyltetrazolyl, oxolanyl, cycloalkyltetrazolyl or haloalkyl-1H-pyrazolyl;

or a pharmaceutically acceptable salt or ester thereof.

In another aspect, there is provided a process for the preparation of a compound as defined herein, comprising the reaction of a compound of formula (A)



5 in the presence of X-A-R³, wherein X is a leaving group and wherein A and R¹ to R³ are as defined herein.

In another aspect, there is provided a compound as defined herein, when manufactured according to a process as defined herein.

In another aspect, there is provided a compound as defined herein for use as therapeutically active substance.

10 In another aspect, there is provided a pharmaceutical composition comprising a compound as defined herein and a therapeutically inert carrier.

In another aspect, there is provided use of a compound as defined herein for the preparation of a medicament for the treatment or prophylaxis of pain, atherosclerosis, age-related macular degeneration, diabetic retinopathy, glaucoma, retinal vein occlusion, retinopathy of prematurity, ocular ischemic syndrome, geographic atrophy, diabetes mellitus, inflammation, inflammatory bowel disease, ischemia-reperfusion injury, acute liver failure, liver fibrosis, lung fibrosis, kidney fibrosis, systemic fibrosis, acute allograft rejection, chronic allograft nephropathy, diabetic nephropathy, glomerulonephropathy, cardiomyopathy, heart failure, myocardial ischemia, myocardial infarction, systemic sclerosis, thermal injury, burning, hypertrophic scars, keloids, gingivitis pyrexia, liver cirrhosis or tumors, regulation of bone mass, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease, Parkinson's disease, stroke, transient ischemic attack or uveitis.

15
20

In another aspect, there is provided a compound as defined herein for the treatment or prophylaxis of pain, atherosclerosis, age-related macular degeneration, diabetic retinopathy, glaucoma, retinal vein occlusion, retinopathy of prematurity, ocular ischemic syndrome, geographic atrophy, diabetes mellitus, inflammation, inflammatory bowel
5 disease, ischemia-reperfusion injury, acute liver failure, liver fibrosis, lung fibrosis, kidney fibrosis, systemic fibrosis, acute allograft rejection, chronic allograft nephropathy, diabetic nephropathy, glomerulonephropathy, cardiomyopathy, heart failure, myocardial ischemia, myocardial infarction, systemic sclerosis, thermal injury, burning, hypertrophic scars, keloids, gingivitis pyrexia, liver cirrhosis or tumors, regulation of bone mass, amyotrophic
10 lateral sclerosis, multiple sclerosis, Alzheimer's disease, Parkinson's disease, stroke, transient ischemic attack or uveitis.

In the present description the term "alkyl", alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, particularly a
15 straight or branched-chain alkyl group with 1 to 6 carbon atoms and more particularly a straight or branched-chain alkyl group with 1 to 4 carbon atoms. Examples of straight-chain and branched-chain C₁-C₈ alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, particularly methyl, ethyl, propyl, butyl and pentyl. A particular example of alkyl is methyl.

20 The term "cycloalkyl", alone or in combination, signifies a cycloalkyl ring with 3 to 8 carbon atoms and particularly a cycloalkyl ring with 3 to 6 carbon atoms. Examples of cycloalkyl are cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, cycloheptyl and cyclooctyl. A particular example of "cycloalkyl" is cyclopropyl.

The terms "halogen" or "halo", alone or in combination, signifies fluorine, chlorine,
25 bromine or iodine and particularly fluorine, chlorine or bromine, more particularly fluorine and chlorine. The term "halo", in combination with another group, denotes the substitution of said group with at least one halogen, particularly substituted with one to five halogens, particularly one to four halogens, i.e. one, two, three or four halogens. Particular "halogen" are fluorine and chlorine. In the definition of R¹ and R², fluorine is a particular halogen.

30 The term "haloalkyl", alone or in combination, denotes an alkyl group substituted with at least one halogen, particularly substituted with one to five halogens, particularly one to three halogens. A particular "haloalkyl" is trifluoromethyl.

12. A compound according to any one of claims 1 to 9 for use as therapeutically active substance.
13. A pharmaceutical composition comprising a compound in accordance with any one of claims 1 to 9 and a therapeutically inert carrier.
14. The use of a compound according to any one of claims 1 to 9 for the preparation of a medicament for the treatment or prophylaxis of pain, atherosclerosis, age-related macular degeneration, diabetic retinopathy, glaucoma, retinal vein occlusion, retinopathy of prematurity, ocular ischemic syndrome, geographic atrophy, diabetes mellitus, inflammation, inflammatory bowel disease, ischemia-reperfusion injury, acute liver failure, liver fibrosis, lung fibrosis, kidney fibrosis, systemic fibrosis, acute allograft rejection, chronic allograft nephropathy, diabetic nephropathy, glomerulonephropathy, cardiomyopathy, heart failure, myocardial ischemia, myocardial infarction, systemic sclerosis, thermal injury, burning, hypertrophic scars, keloids, gingivitis pyrexia, liver cirrhosis or tumors, regulation of bone mass, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease, Parkinson's disease, stroke, transient ischemic attack or uveitis.
15. A compound according to any one of claims 1 to 9 for the treatment or prophylaxis of pain, atherosclerosis, age-related macular degeneration, diabetic retinopathy, glaucoma, retinal vein occlusion, retinopathy of prematurity, ocular ischemic syndrome, geographic atrophy, diabetes mellitus, inflammation, inflammatory bowel disease, ischemia-reperfusion injury, acute liver failure, liver fibrosis, lung fibrosis, kidney fibrosis, systemic fibrosis, acute allograft rejection, chronic allograft nephropathy, diabetic nephropathy, glomerulonephropathy, cardiomyopathy, heart failure, myocardial ischemia, myocardial infarction, systemic sclerosis, thermal

injury, burning, hypertrophic scars, keloids, gingivitis pyrexia, liver cirrhosis or tumors, regulation of bone mass, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease, Parkinson's disease, stroke, transient ischemic attack or uveitis.
