
-(3R,4S,5S,6S)-2-(acetoxymethyl)-6-(4-(3,3,6,6-tetramethyl-1,8-dioxo-2,3,4,5,6,7,8,9-octahydro-1H-xanthen-9-yl)phenoxy)tetrahydro-2H-pyran-3,4,5-triyl triacetate

-3,3,6,6-tetramethyl-9-(4-((2S,3S,4S,5S)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yloxy)phenyl)-3,4,5,6,7,9-hexahydro-1H-xanthene-1,8(2H)-dione.

8. Compound of generic formula (I'):




(I')

in which the radicals R_1, R_2, R_3, R_4, R_5, R_6 and R_7 have identical significations to those given in claims 1 to 6, but in which R_4 may in addition represent a hydrogen atom or an acetylated radical, and ~~cosmetically or~~ pharmaceutically acceptable salts, for ~~its~~ use as medicine ~~or as cosmetic active ingredient~~.

9. Compound of generic formula (I') according to claim 8, characterised in that R_4 represents a hydrogen atom for ~~its~~ use as medicine ~~or as cosmetic active ingredient~~.


10. Use of a ~~C~~compound of generic formula (I') according to one of claims 8 or 9 in the manufacture of a medicament for ~~its use in the~~ depigmentation of the skin and/or head hair and/or body hair, ~~in the treatment and/or the prevention of the ageing of the skin~~ or for ~~in~~ the treatment and/or the prevention of the inflammation of the skin.

11. Pharmaceutical or cosmetic composition, characterised in that it comprises as active ingredient at least one compound of generic formula (I') as defined according to claims 8 to 9 in association with a pharmaceutically or cosmetically acceptable excipient.



with R_1, R_2, R_3, R_5, R_6 and R_7 having the same significations as in claims 1 to 6 and R_4 represents COR_5.

16. Method for synthesising compounds of generic formula (I) according to one of claims 1 to 2 and 5 to 6, characterised in that a phenol is made to react with a glucide radical substituted by one or more acetyl radical(s) in the presence of a Lewis acid (BF_3, Et_2O):




(I)

with R_1, R_2, R_3, R_5, R_6 and R_7 having the same significations as in claims 1 to 6, and R_4 represents a glucide radical substituted by one or more acetyl radical(s).

17. The method of claim 16, ~~which may if appropriate be~~ followed by a saponification step.

18. Use of a compound of generic formula (I'):



(I')

in which the radicals R_1, R_2, R_3, R_4, R_5, R_6 and R_7 have identical significations to those given in claims 1 to 6, but in which R_4 may in addition represent a hydrogen atom or an acetylated radical, as a cosmetic active ingredient.

19. The use of claim 18, wherein R^4 represents a hydrogen atom.

20. The use of claims 18 or 19, for the treatment and/or the prevention of the ageing of the skin.