QUINAZOLINES USEFUL AS MODULATORS OF ION CHANNELS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit under 35 U.S.C. § 119 to United States Provisional application serial no. 60/607,150, filed September 2, 2004, United States Provisional application serial no. 60/607,037, filed September 2, 2004, United States Provisional application serial no. 60/607,033, filed September 2, 2004, United States Provisional application serial no. 60/607,036, filed September 2, 2004, and United States Provisional application serial no. 60/607,245, filed September 2, 2004, the entire contents of each of the above application being incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

[0002] The present invention relates to compounds useful as inhibitors of ion channels. The invention also provides pharmaceutically acceptable compositions comprising the compounds of the invention and methods of using the compositions in the treatment of various disorders.

BACKGROUND OF THE INVENTION

[0003] Na channels are central to the generation of action potentials in all excitable cells such as neurons and myocytes. They play key roles in excitable tissue including brain, smooth muscles of the gastrointestinal tract, skeletal muscle, the peripheral nervous system, spinal cord and airway. As such they play key roles in a variety of disease states such as epilepsy (See, Moulard, B. and D. Bertrand (2002) "Epilepsy and sodium channel blockers" Expert Opin. Ther. Patents **12**(1): 85-91)), pain (See, Waxman, S. G., S. Dib-Hajj, et al. (1999) "Sodium channels and pain" Proc Natl Acad Sci U S A **96**(14): 7635-9 and Waxman, S. G., T. R. Cummins, et al. (2000) "Voltage-gated sodium channels and the molecular pathogenesis of pain: a review" J Rehabil Res Dev **37**(5): 517-28), myotonia (See, Meola, G. and V. Sansone (2000) "Therapy in myotonic disorders and in muscle channelopathies" Neurol Sci **21**(5): S953-61 and Mankodi, A. and C. A. Thornton (2002) "Myotonic syndromes" Curr Opin Neurol **15**(5): 545-52), ataxia (See,

Meisler, M. H., J. A. Kearney, et al. (2002) "Mutations of voltage-gated sodium channels in movement disorders and epilepsy" Novartis Found Symp **241**: 72-81), multiple sclerosis (See, Black, J. A., S. Dib-Hajj, et al. (2000) "Sensory neuron-specific sodium channel SNS is abnormally expressed in the brains of mice with experimental allergic encephalomyelitis and humans with multiple sclerosis" Proc Natl Acad Sci U S A **97**(21): 11598-602, and Renganathan, M., M. Gelderblom, et al. (2003) "Expression of Na(v)1.8 sodium channels perturbs the firing patterns of cerebellar purkinje cells" Brain Res **959**(2): 235-42), irritable bowel (See, Su, X., R. E. Wachtel, et al. (1999) "Capsaicin sensitivity and voltage-gated sodium currents in colon sensory neurons from rat dorsal root ganglia" Am J Physiol **277**(6 Pt 1): G1180-8, and Laird, J. M., V. Souslova, et al. (2002) "Deficits in visceral pain and referred hyperalgesia in Nav1.8 (SNS/PN3)- null mice" J Neurosci **22**(19): 8352-6), urinary incontinence and visceral pain (See,Yoshimura, N., S. Seki, et al. (2001) "The involvement of the tetrodotoxin-resistant sodium channel Na(v)1.8 (PN3/SNS) in a rat model of visceral pain" J Neurosci **21**(21): 8690-6), as well as an array of psychiatry dysfunctions such as anxiety and depression (See, Hurley, S. C. (2002) "Lamotrigine update and its use in mood disorders" Ann Pharmacother **36**(5): 860-73).

[0004] Voltage gated Na channels comprise a gene family consisting of 9 different subtypes (NaV1.1-NaV1.9). These subtypes show tissue specific localization and functional differences (See, Goldin, A. L. (2001) "Resurgence of sodium channel research" Annu Rev Physiol **63**: 871-94). Three members of the gene family (NaV1.8, 1.9, 1.5) are resistant to block by the well-known Na channel blocker TTX, demonstrating subtype specificity within this gene family. Mutational analysis has identified glutamate 387 as a critical residue for TTX binding (See, Noda, M., H. Suzuki, et al. (1989) "A single point mutation confers tetrodotoxin and saxitoxin insensitivity on the sodium channel II" FEBS Lett **259**(1): 213-6).

[0005] In general, voltage-gated sodium channels (NaVs) are responsible for initiating the rapid upstroke of action potentials in excitable tissue in nervous system, which transmit the electrical signals that compose and encode normal and aberrant pain sensations. Antagonists of NaV channels can attenuate these pain signals and are useful for treating a variety of pain conditions, including but not limited to acute, chronic, inflammatory, and neuropathic pain. Known NaV antagonists, such as TTX, lidocaine (See, Mao, J. and L. L. Chen (2000) "Systemic lidocaine for neuropathic pain relief" Pain **87**(1): 7-17.) bupivacaine, phenytoin (See, Jensen, T. S. (2002) "Anticonvulsants in neuropathic pain: rationale and clinical evidence" Eur J Pain **6**

(Suppl A): 61-8), lamotrigine (See, Rozen, T. D. (2001) "Antiepileptic drugs in the management of cluster headache and trigeminal neuralgia" Headache **41 Suppl 1**: S25-32 and Jensen, T. S. (2002) "Anticonvulsants in neuropathic pain: rationale and clinical evidence" Eur J Pain **6** (Suppl A): 61-8.), and carbamazepine (See, Backonja, M. M. (2002) "Use of anticonvulsants for treatment of neuropathic pain" Neurology **59**(5 Suppl 2): S14-7), have been shown to be useful attenuating pain in humans and animal models.

[0006] Hyperalgesia (extreme sensitivity to something painful) that develops in the presence of tissue injury or inflammation reflects, at least in part, an increase in the excitability of high-threshold primary afferent neurons innervating the site of injury. Voltage sensitive sodium channels activation is critical for the generation and propagation of neuronal action potentials. There is a growing body of evidence indicating that modulation of NaV currents is an endogenous mechanism used to control neuronal excitability (See, Goldin, A. L. (2001) "Resurgence of sodium channel research" Annu Rev Physiol **63**: 871-94.). Several kinetically and pharmacologically distinct voltage-gated sodium channels are found in dorsal root ganglion (DRG) neurons. The TTX-resistant current is insensitive to micromolar concentrations of tetrodotoxin, and displays slow activation and inactivation kinetics and a more depolarized activation threshold when compared to other voltage-gated sodium channels. TTX-resistant sodium currents are primarily restricted to a subpopulation of sensory neurons likely to be involved in nociception. Specifically, TTX-resistant sodium currents are expressed almost exclusively in neurons that have a small cell-body diameter; and give rise to small-diameter slow-conducting axons and that are responsive to capsaicin. A large body of experimental evidence demonstrates that TTX-resistant sodium channels are expressed on C-fibers and are important in the transmission of nociceptive information to the spinal cord.

[0007] Intrathecal administration of antisense oligo-deoxynucleotides targeting a unique region of the TTX-resistant sodium channel (NaV1.8) resulted in a significant reduction in PGE_2-induced hyperalgesia (See, Khasar, S. G., M. S. Gold, et al. (1998) "A tetrodotoxin-resistant sodium current mediates inflammatory pain in the rat" Neurosci Lett **256**(1): 17-20). More recently, a knockout mouse line was generated by Wood and colleagues, which lacks functional NaV1.8. The mutation has an analgesic effect in tests assessing the animal's response to the inflammatory agent carrageenan (See, Akopian, A. N., V. Souslova, et al. (1999) "The tetrodotoxin-resistant sodium channel SNS has a specialized function in pain pathways" Nat

Neurosci 2(6): 541-8.). In addition, deficit in both mechano- and thermoreception were observed in these animals. The analgesia shown by the Nav1.8 knockout mutants is consistent with observations about the role of TTX-resistant currents in nociception.

[0008] Immunohistochemical, in-situ hybridization and in-vitro electrophysiology experiments have all shown that the sodium channel NaV1.8 is selectively localized to the small sensory neurons of the dorsal root ganglion and trigeminal ganglion (See, Akopian, A. N., L. Sivilotti, et al. (1996) "A tetrodotoxin-resistant voltage-gated sodium channel expressed by sensory neurons" Nature **379**(6562): 257-62.). The primary role of these neurons is the detection and transmission of nociceptive stimuli. Antisense and immunohistochemical evidence also supports a role for NaV1.8 in neuropathic pain (See, Lai, J., M. S. Gold, et al. (2002) "Inhibition of neuropathic pain by decreased expression of the tetrodotoxin-resistant sodium channel, NaV1.8" Pain **95**(1-2): 143-52, and Lai, J., J. C. Hunter, et al. (2000) "Blockade of neuropathic pain by antisense targeting of tetrodotoxin- resistant sodium channels in sensory neurons" Methods Enzymol **314**: 201-13.). NaV1.8 protein is upregulated along uninjured C-fibers adjacent to the nerve injury. Antisense treatment prevents the redistribution of NaV1.8 along the nerve and reverses neuropathic pain. Taken together the gene-knockout and antisense data support a role for NaV1.8 in the detection and transmission of inflammatory and neuropathic pain.

[0009] Several Na channel blockers are currently used or being tested in the clinic to treat epilepsy (See, Moulard, B. and D. Bertrand (2002) "Epilepsy and sodium channel blockers" Expert Opin. Ther. Patents **12**(1): 85-91.); acute (See, Wiffen, P., S. Collins, et al. (2000) "Anticonvulsant drugs for acute and chronic pain" Cochrane Database Syst Rev 3), chronic (See, Wiffen, P., S. Collins, et al. (2000) "Anticonvulsant drugs for acute and chronic pain" Cochrane Database Syst Rev **3**, and Guay, D. R. (2001) "Adjunctive agents in the management of chronic pain" Pharmacotherapy **21**(9): 1070-81), inflammatory (See, Gold, M. S. (1999) "Tetrodotoxin-resistant Na+ currents and inflammatory hyperalgesia." Proc Natl Acad Sci U S A **96**(14): 7645-9), and neuropathic pain (See, Strichartz, G. R., Z. Zhou, et al. (2002) "Therapeutic concentrations of local anaesthetics unveil the potential role of sodium channels in neuropathic pain" Novartis Found Symp **241**: 189-201, and Sandner-Kiesling, A., G. Rumpold Seitlinger, et al. (2002) "Lamotrigine monotherapy for control of neuralgia after nerve section" Acta Anaesthesiol Scand **46**(10): 1261-4); cardiac arrhythmias (See, An, R. H., R. Bangalore, et al.

(1996) "Lidocaine block of LQT-3 mutant human Na+ channels" Circ Res 79(1): 103-8, and Wang, D. W., K. Yazawa, et al. (1997) "Pharmacological targeting of long QT mutant sodium channels" J Clin Invest 99(7): 1714-20); neuroprotection (See, Taylor, C. P. and L. S. Narasimhan (1997) "Sodium channels and therapy of central nervous system diseases" Adv Pharmacol 39: 47-98) and as anesthetics (See, Strichartz, G. R., Z. Zhou, et al. (2002) "Therapeutic concentrations of local anaesthetics unveil the potential role of sodium channels in neuropathic pain." Novartis Found Symp 241: 189-201).

[0010] Various animal models with clinical significance have been developed for the study of sodium channel modulators for numerous different pain indications. E.g., malignant chronic pain, see, Kohase, H., et al., Acta Anaesthesiol Scand. 2004; 48(3):382-3; femur cancer pain (see, Kohase, H., et al., Acta Anaesthesiol Scand. 2004; 48(3):382-3); non-malignant chronic bone pain (see, Ciocon, J. O. et al., J Am Geriatr Soc. 1994; 42(6):593-6); rheumatoid arthritis (see, Calvino, B. et al., Behav Brain Res. 1987; 24(1):11-29); osteoarthritis (see, Guzman, R. E., et al., Toxicol Pathol. 2003; 31(6):619-24); spinal stenosis (see, Takenobu, Y. et al., J Neurosci Methods. 2001; 104(2):191-8); Neuropathic low back pain (see, Hines, R., et al., Pain Med. 2002; 3(4):361-5; Massie, J. B., et al., J Neurosci Methods. 2004; 137(2):283-9; neuropathic low back pain (see, Hines, R., et al., Pain Med. 2002; 3(4):361-5; Massie, J. B., et al., J Neurosci Methods. 2004; 137(2):283-9); myofascial pain syndrome (see, Dalpiaz & Dodds, J Pain Palliat Care Pharmacother. 2002; 16(1):99-104; Sluka KA et al., Muscle Nerve. 2001; 24(1):37-46); fibromyalgia (see, Bennet & Tai, Int J Clin Pharmacol Res. 1995;15(3):115-9); temporomandibular joint pain (see, Ime H, Ren K, Brain Res Mol Brain Res. 1999; 67(1):87-97); chronic visceral pain, including, abdominal (see, Al-Chaer, E. D., et al., Gastroenterology. 2000; 119(5):1276-85); pelvic/perineal pain, (see, Wesselmann et al., Neurosci Lett. 1998; 246(2):73-6); pancreatic (see, Vera-Portocarrero, L. B., et al., Anesthesiology. 2003; 98(2):474-84); IBS pain (see, Verne, G. N., et al., Pain. 2003; 105(1-2):223-30; La JH et al., World Gastroenterol. 2003; 9(12):2791-5); chronic headache pain (see, Willimas & Stark, Cephalalgia. 2003; 23(10):963-71); migraine (see, Yamamura, H., et al., J Neurophysiol. 1999; 81(2):479-93); tension headache, including, cluster headaches (see, Costa, A., et al., Cephalalgia. 2000; 20(2):85-91); chronic neuropathic pain, including, post-herpetic neuralgia (see, Attal, N., et al., Neurology. 2004; 62(2):218-25; Kim & Chung 1992, Pain 50:355); diabetic neuropathy (see, Beidoun A et al., Clin J Pain. 2004; 20(3):174-8; Courteix, C., et al., Pain. 1993; 53(1):81-8);

HIV- associated neuropathy (see, Portegies & Rosenberg, Ned Tijdschr Geneeskd. 2001; 145(15):731-5; Joseph EK et al., Pain. 2004; 107(1-2):147-58; Oh, S. B., et al., J Neurosci. 2001; 21(14):5027-35); trigeminal neuralgia (see, Sato, J., et al., Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2004; 97(1):18-22; Imamura Y et al., Exp Brain Res. 1997; 116(1):97-103); Charcot-Marie Tooth neuropathy (see, Sereda, M., et al., Neuron. 1996; 16(5):1049-60); hereditary sensory neuropathies (see, Lee, M. J., et al., Hum Mol Genet. 2003; 12(15):1917-25); peripheral nerve injury (see, Attal, N., et al., Neurology. 2004; 62(2):218-25; Kim & Chung 1992, Pain 50:355; Bennett & Xie, 1988, Pain 33:87; Decostered, I. & Woolf, C. J., 2000, Pain 87:149; Shir, Y. & Seltzer, Z. 1990; Neurosci Lett 115:62); painful neuromas (see, Nahabedian & Johnson, Ann Plast Surg. 2001; 46(1):15-22; Devor & Raber, Behav Neural Biol. 1983; 37(2):276-83); ectopic proximal and distal discharges (see, Liu, X. et al., Brain Res. 2001; 900(1):119-27); radiculopathy (see, Devers & Galer, (see, Clin J Pain. 2000; 16(3):205-8; Hayashi N et al., Spine. 1998; 23(8):877-85); chemotherapy induced neuropathic pain (see, Aley, K. O., et al., Neuroscience. 1996; 73(1):259-65); radiotherapy-induced neuropathic pain; post-mastectomy pain (see, Devers & Galer, Clin J Pain. 2000; 16(3):205-8); central pain (Cahana, A., et al., Anesth Analg. 2004; 98(6):1581-4), spinal cord injury pain (see, Hains, B. C., et al., Exp Neurol. 2000; 164(2):426-37); post-stroke pain; thalamic pain (see, LaBuda, C. J., et al., Neurosci Lett. 2000; 290(1):79-83); complex regional pain syndrome (see, Wallace, M. S., et al., Anesthesiology. 2000; 92(1):75-83; Xantos D et al., J Pain. 2004; 5(3 Suppl 2):S1); phanton pain (see, Weber, W. E., Ned Tijdschr Geneeskd. 2001; 145(17):813-7; Levitt & Heyback, Pain. 1981; 10(1):67-73); intractable pain (see, Yokoyama, M., et al., Can J Anaesth. 2002; 49(8):810-3); acute pain, acute post-operative pain (see, Koppert, W., et al., Anesth Analg. 2004; 98(4):1050-5; Brennan, T. J., et al., Pain. 1996; 64(3):493-501); acute musculoskeletal pain; joint pain (see, Gotoh, S., et al., Ann Rheum Dis. 1993; 52(11):817-22); mechanical low back pain (see, Kehl, L. J., et al., Pain. 2000; 85(3):333-43); neck pain; tendonitis; injury/exercise pain (see, Sesay, M., et al., Can J Anaesth. 2002; 49(2):137-43); acute visceral pain, including, abdominal pain; pyelonephritis; appendicitis; cholecystitis; intestinal obstruction; hernias; etc (see, Giambernardino, M. A., et al., Pain. 1995; 61(3):459-69); chest pain, including, cardiac Pain (see, Vergona, R. A., et al., Life Sci. 1984; 35(18):1877-84); pelvic pain, renal colic pain, acute obstetric pain, including, labor pain (see, Segal, S., et al., Anesth Analg. 1998; 87(4):864-

9); cesarean section pain; acute inflammatory, burn and trauma pain; acute intermittent pain, including, endometriosis (see, Cason, A. M., et al.,Horm Behav. 2003; 44(2):123-31); acute herpes zoster pain; sickle cell anemia; acute pancreatitis (see, Toma, H; Gastroenterology. 2000; 119(5):1373-81); breakthrough pain; orofacial pain, including, sinusitis pain, dental pain (see, Nusstein, J., et al., J Endod. 1998; 24(7):487-91; Chidiac, J. J., et al., Eur J Pain. 2002; 6(1):55-67); multiple sclerosis (MS) pain (see, Sakurai & Kanazawa, J Neurol Sci. 1999; 162(2):162-8); pain in depression (see, Greene B, Curr Med Res Opin. 2003; 19(4):272-7); leprosy pain; behcet's disease pain; adiposis dolorosa (see, Devillers & Oranje, Clin Exp Dermatol. 1999; 24(3):240-1); phlebitic pain; Guillain-Barre pain; painful legs and moving toes; Haglund syndrome; erythromelalgia pain (see, Legroux-Crespel, E., et al., Ann Dermatol Venereol. 2003; 130(4):429-33); Fabry's disease pain (see, Germain, D. P., J Soc Biol. 2002;196(2):183-90); Bladder and urogenital disease, including, urinary incontinence (see, Berggren, T., et al., J Urol. 1993; 150(5 Pt 1):1540-3); hyperactivity bladder (see, Chuang, Y. C., et al., Urology. 2003; 61(3):664-70); painful bladder syndrome (see, Yoshimura, N., et al., J Neurosci. 2001; 21(21):8690-6); interstitial cyctitis (IC) (see, Giannakopoulos& Campilomatos, Arch Ital Urol Nefrol Androl. 1992; 64(4):337-9; Boucher, M., et al., J Urol. 2000; 164(1):203-8); and prostatitis (see, Mayersak, J. S., Int Surg. 1998; 83(4):347-9; Keith, I. M., et al., J Urol. 2001; 166(1):323-8).

[0011] Unfortunately, as described above, the efficacy of currently used sodium channel blockers and calcium channel blockers for the disease states described above has been to a large extent limited by a number of side effects. These side effects include various CNS disturbances such as blurred vision, dizziness, nausea, and sedation as well more potentially life threatening cardiac arrhythmias and cardiac failure. Such undesirable side effects may be avoided by using a Na channel blocker that exhibit a degree of selectivity in its activity against a Na channel subtype. However, Na channel blockers currently in the market lack such selectivity. Perhaps because of this lack of molecular selectivity, drugs currently in the market exhibit use-dependent block and generally show higher affinity at depolarized potentials resulting in the preferential targeting of actively firing neurons, believed to be a key factor in the therapeutic window of existing Na channel blocking drugs. While every drug has it own unique therapeutic profile, current Na channel blockers are generally associated with central nervous system (CNS) and cardiovascular (CV) side-effects, including blood pressure changes, which are often dose-

limiting. Dizziness, sedation, nausea, ataxia, and confusion are some of the specific side-effects observed for phenytoin™, mexiletine™, and lidocaine™.

[0012] There is also a need to develop Na channel blockers that have minimal or no inhibitory activity against the hERG channel. hERG (human ether a-go-go related gene) encodes a potassium ion channel (hERG channel) that is involved in cardiac repolarization. See, e.g., Pearlstein, R., R. Vaz, et al. (2003). "Understanding the Structure-Activity Relationship of the Human Ether-a-go-go-Related Gene Cardiac K(+) Channel. A Model for Bad Behavior." J Med Chem **46**(11): 2017-22. Interaction with with the hERG channel is one indicator of potential cardaic toxicity. hERG-block increases the likelihood of cardiac QT-interval prolongation and dispersion. A subset of compounds that prolong the QT interval can cause ventricular fibrillation and cardiac failure. Belardinelli, L., C. Antzelevitch and M.A. Vos (2003). "Assessing predictors of drug-induced torsade de pointes". Trends Pharmacol Sci. **24** (12): 619-25; Al-Khatib, S. M., N. M. LaPointe, et al. (2003). "What clinicians should know about the QT interval." Jama **289**(16): 2120-7; http://www.fenichel.net/pages/site_map.htm.

[0013] There is also a need to develop Na channel blockers that have minimal or no inhibitory activity against the Cytochrome P450 enzyme family. Within this family, CYP 3A4 isoform is believed to be the major isoform present in the liver and small intestines. Other key isoforms include CYP 2D6, CYP 2C9, and CYP 1A2. See, e.g., United States patent 6,514,687, the disclosure whereof is incorporated herein by reference. A Na channel blocker that inhibits one or more of the isoforms can cause undesirable side effect or can cause undesirable drug-drug interactions when administered with another drug that interacts with that isoform. See, e.g, Davit, B., et al. (1999), "FDA Evaluations Using In Vitro Metabolism to Predict and Interpret In Vivo Metabolic Drug-Drug Interactions: Impact on Labeling," J. Clin. Pharmacol., 39: 899-910; "Drug Metabolism/Drug Interaction Studies in the Drug Development Process: Studies In Vitro, Dept. of Health and Human Services, U.S.F.D.A (http://www.fda.gov/cder/guidance.htm).

[0014] There is also a need to develop Na channel blockers that exhibit selectivity against a certain sub-type of Na channel. Particularly useful are compounds that have a desirably low activity against NaV 1.2.

[0015] There is also a need to develop Na channel blockers that have a desirably low activity against L-type calcium channel 1.2. CaV1.2 calcium channels are abundantly expressed in smooth and striated muscle, especially in the heart, brain and endocrine cells. Blocking these

channels can be therapeutically useful, but it can also result in significant side effects. The most significant concerns are impairment of cardiac contractility (that is, a negative inotropic effect) and slowing of electrical conduction in the pacemaker regions of the heart. See, e.g., Kizer, J. R., et al., "Epidemiologic Review of the Calcium Channel Blocker Drugs," Arch. Intern Med. 2001; 161: 1145-1158.


[0016] There is also a need to Na channel blockers that have a desirably low activity against potassiums channel 1.5 ("Kv1.5"; also known as KCNA5). Kv1.5 is found primarily in human atrial cells, but also in brain. See, e.g., Gutman, G. A., et al., "Compendium of Voltage-Gated Ion Channels: Potassium Channels," Pharmacol. Rev., 55: 583-585 (2003). Unwanted block of Kv1.5 could produce convulsion or ataxia.

[0017] There is also a need to develop Na channel blockers that have improved pharmacokinetic and/or pharmacodynamic properties and, therefore, are better suited for *in-vivo* administration for therapeutic purposes. Such properties include aqueous solubility, bioavailability, clearance kinetics, etc. See, e.g., Shargel, L., Yu, A., Ed's "Applied Biopharmaceutics & Pharmacokinetics", 4th Ed., McGraw-Hill, New York, 1999; Yacobi, A., Skelly, J.P., Shah, V.P., Benet, L.Z., Ed's. "Integration of Pharmacokinetics, Pharmacodynamics, and Toxicokinetics in Rational Drug Development", Plenum Press, New York, 1993; Lee, J.S., Obach, R.S., Fisher, M.B., Ed's. "Drug Metabolizing Enzymes Cytochrome P450 and Other Enzymes in Drug Discovery and Development", Marcel Dekker, New York, 2003; Birkett, D.J. "Pharmacokinetics Made Easy", McGraw-Hill Australia, Roseville, Australia, 2002; Katzung, B.G. "Basic & Clinical Pharmacology", McGraw-Hill, New York, 2001; Welling, P.G., Tse, F.L.S., Ed's. "Pharmcokinetics", Marcel Dekker, New York, 1988; Thomas, G. "Medicinal Chemistry An Introduction", Wiley & Sons, New York, 2000; and Gennaro, A. R., et al., "Remington: The Science and Practice of Pharmacy," 20th Ed., Lippincott, Williams, & Wilkins (2003).

[0018] Na channel blocker that meets one or more of the above unmet needs would be a very desirable improvement over the currently marketed Na channel blockers and would greatly benefit patients in need of a therapy therewith.

SUMMARY OF THE INVENTION

[0019] The present invention provides compounds of formula **I:**



I;


or a pharmaceutically acceptable salt or derivative thereof.

[0020] These compounds and pharmaceutically acceptable compositions are useful for treating or lessening the severity of a variety of diseases, disorders, or conditions, including, but not limited to, acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias, epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, incontinence, visceral pain, osteoarthritis pain, postherpetic neuralgia, diabetic neuropathy, radicular pain, sciatica, back pain, head or neck pain, severe or intractable pain, nociceptive pain, breakthrough pain, postsurgical pain, or cancer pain.

DETAILED DESCRIPTION OF THE INVENTION

[0021] *I. General Description of Compounds of the Invention:*


[0022] The present invention provides a compound of formula **I**:




I;

or a pharmaceutically acceptable salt or derivative thereof, wherein:


R^1 and R^2, taken together with the nitrogen atom, form a substituted ring selected from:




(A)



(B)




(C)



(D)

or



(E);

wherein, in ring (A):

each of m_1 and n_1 is independently 0-3, provided that m_1+n_1 is 2-6;

z_1 is 0-4;

Sp^1 is –O-, -S-, -NR'-, or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-, provided that Sp^1 is attached to the carbonyl group through an atom other than carbon;

ring B^1 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w_1 independent occurrences of -R^{11}, wherein w_1 is 0-4;

wherein, in ring (B):

G_2 is -N-, or CH;

each of m_2 and n_2 is independently 0-3, provided that $m_2 + n_2$ is 2-6;

p_2 is 0-2; provided that when G_2 is N, then p_2 is not 0;

q_2 is 0 or 1;

z_2 is 0-4;

Sp^2 is a bond or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, $-CO_2-$, -OCO-, $-NR'CO_2-$, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, $-SO_2-$, -NR'-, $-SO_2NR'-$, $NR'SO_2-$, or $-NR'SO_2NR'-$;

ring B^2 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4;

wherein, in ring (C) or ring (D):

G_3 is -N-, -CH-NH-, or $-CH-CH_2-NH-$;

each of m_3 and n_3 is independently 0-3, provided that m_3+n_3 is 2-6;

p_3 is 0-2;

z_3 is 0-4;

each R^{XX} is hydrogen, C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3;

provided that both R^{XX} are not simultaneously hydrogen;

R^{YY} is hydrogen, -COR', $-CO_2R'$, $-CON(R')_2$, -SOR', $-SO_2R'$, $-SO_2N(R')_2$, -COCOR', $-COCH_2COR'$, $-P(O)(OR')_2$, $-P(O)_2OR'$, or -PO(R');

wherein, in ring (E):

each of m_4 and n_4 is independently 0-3, provided that $m_4 + n_4$ is 2-6;

p_4 is 1-2;


z_4 is 0-4;

R^{YZ} is C_1-C_6 aliphatic group, optionally substituted with w_4 independent occurrences of -R^{14}, wherein w_4 is 0-3;


x and y, each, is independently 0-4;

W is OR^{XY};

R^{XY} is hydrogen or a group selected from:



or



;

wherein:

each of w_A, w_B, w_C, and w_D is independently 0 or 1;

each M is independently selected from hydrogen, Li, Na, K, Mg, Ca, Ba, -$N(R^7)_4$, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or -R^6; wherein 1 to 4 –CH_2 radicals of the alkyl or alkenyl group, other than the –CH_2 that is bound to Z, is optionally replaced by a heteroatom group selected from O, S, S(O), S(O_2), or N(R^7); and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a substituent selected from oxo, -OR^7, - R^7, $N(R^7)_2$, $N(R^7)_3$, R^7OH, -CN, -CO_2 R^7, -C(O)-$N(R^7)_2$, $S(O)_2$-$N(R^7)_2$, N(R^7)-C(O)- R^7, C(O) R^7, -$S(O)_n$- R^7, OCF_3, -$S(O)_n$-R^6, N(R^7)-$S(O)_2(R^7)$, halo, -CF_3, or -NO_2;

n is 0-2;

M' is H, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or -R^6; wherein 1 to 4 –CH_2 radicals of the alkyl or alkenyl group is optionally replaced by a heteroatom group selected from O, S, S(O), S(O_2), or N(R^7); and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a substituent selected from oxo, -O R^7, - R^7, -$N(R^7)_2$, $N(R^7)_3$, - R^7OH, -CN, -CO_2 R^7, -C(O)-$N(R^7)_2$, -$S(O)_2$-$N(R^7)_2$, -N(R^7)-C(O)- R^7, -C(O) R^7, -$S(O)_n$- R^7, -OCF_3, -$S(O)_n$-R^6, -N(R^7)-$S(O)_2(R^7)$, halo, -CF_3, or -NO_2;

Z is -CH_2-, -O-, -S-, -$N(R^7)_2$-; or,

when M is absent, then Z is hydrogen, =O, or =S;

Y is P or S, wherein when Y is S, then Z is not S;

X is O or S;

each R^7 is independently selected from hydrogen, or C_1-C_4 aliphatic, optionally substituted with up to two Q_1;

each Q_1 is independently selected from a 3-7 membered saturated, partially saturated or unsaturated carbocyclic ring system; or a 5-7 membered saturated, partially saturated or unsaturated heterocyclic ring containing one or more heteroatom or heteroatom group selected from O, N, NH, S, SO, or SO_2; wherein Q_1 is optionally substituted with up to three substituents selected from oxo, -OH, -O(C_1-C_4 aliphatic), -C_1-C_4 aliphatic, -NH_2, NH(C_1-C_4 aliphatic), -N(C_1-C_4 aliphatic)$_2$, -N(C_1-C_4 aliphatic)-C(O)- C_1-C_4 aliphatic, -(C_1-C_4 aliphatic)-OH, -CN, -CO_2H, -CO_2(C_1-C_4 aliphatic), -C(O)-NH_2, -C(O)-NH(C_1-C_4 aliphatic), -C(O)-N(C_1-C_4 aliphatic)$_2$, halo or -CF_3;

R^6 is a 5-6 membered saturated, partially saturated or unsaturated carbocyclic or heterocyclic ring system, or an 8-10 membered saturated, partially saturated or unsaturated bicyclic ring system; wherein any of said heterocyclic ring systems contains one or more heteroatoms selected from O, N, S, $S(O)_n$ or $N(R^7)$; and wherein any of said ring systems optionally contains 1 to 4 substituents independently selected from OH, C_1-C_4 alkyl, O-C_1-C_4 alkyl or O-C(O)-C_1-C_4 alkyl;

R^9 is $C(R^7)_2$, O or $N(R^7)$;

each occurrence of R^{11}, R^{12}, R^{13}, R^{14}, R^3, R^4, and R^5 is independently Q-R^X; wherein Q is a bond or is a C_1-C_6 alkylidene chain wherein up to two non-adjacent methylene units of Q are optionally and independently replaced by –NR-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -NRCO_2-, -SO_2NR-, -NRSO_2-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO_2NR-, -SO-, -SO_2-, -PO-, -PO_2-, -OP(O)(OR)-, or -POR-; and each occurrence of R^X is independently selected from -R', halogen, =O, =NR', -NO_2, -CN, -OR', -SR', -N(R')$_2$, -NR'COR', -NR'CON(R')$_2$, -NR'CO_2R', -COR', -CO_2R', -OCOR', -CON(R')$_2$, -OCON(R')$_2$, -SOR', -SO_2R', -SO_2N(R')$_2$, -NR'SO_2R', -NR'SO_2N(R')$_2$, -COCOR', -$COCH_2$COR', -OP(O)(OR')$_2$, -P(O)(OR')$_2$, -OP(O)$_2$OR', -P(O)$_2$OR', -PO(R')$_2$, or -OPO(R')$_2$; and

each occurrence of R is independently hydrogen or C_{1-6} aliphatic group having up to three substituents; and each occurrence of R' is independently hydrogen or C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' has up to four

substituents; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0023] *2. Compounds and Definitions:*

[0024] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0025] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not

substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[0026] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C_3-C_8 hydrocarbon or bicyclic C_8-C_{12} hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0027] The term "heteroaliphatic", as used herein, means aliphatic groups wherein one or two carbon atoms are independently replaced by one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon. Heteroaliphatic groups may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and include "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" groups.

[0028] The term "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[0029] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in N-substituted pyrrolidinyl)).

[0030] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[0031] The term "alkoxy", or "thioalkyl", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

[0032] The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I.

[0033] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used interchangeably with the term "aryl ring". The term "aryl" also refers to heteroaryl ring systems as defined hereinbelow.

[0034] The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

[0035] An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the unsaturated carbon atom of an aryl or heteroaryl group are generally selected from halogen; -R°; -OR°; -SR°; phenyl (Ph) optionally substituted with R°; -O(Ph) optionally substituted with R°; $-(CH_2)_{1-2}(Ph)$, optionally substituted with R°; -

CH=CH(Ph), optionally substituted with R°; $-NO_2$; -CN; $-N(R^\circ)_2$; $-NR^\circ C(O)R^\circ$; $-NR^\circ C(S)R^\circ$; $-NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ C(S)N(R^\circ)_2$; $-NR^\circ CO_2R^\circ$; $-NR^\circ NR^\circ C(O)R^\circ$; $-NR^\circ NR^\circ C(O)N(R^\circ)_2$; $-NR^\circ NR^\circ CO_2R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-CO_2R^\circ$; $-C(O)R^\circ$; $-C(S)R^\circ$; $-C(O)N(R^\circ)_2$; $-C(S)N(R^\circ)_2$; $-OC(O)N(R^\circ)_2$; $-OC(O)R^\circ$; $-C(O)N(OR^\circ)\ R^\circ$; $-C(NOR^\circ)\ R^\circ$; $-S(O)_2R^\circ$; $-S(O)_3R^\circ$; $-SO_2N(R^\circ)_2$; $-S(O)R^\circ$; $-NR^\circ SO_2N(R^\circ)_2$; $-NR^\circ SO_2R^\circ$; $-N(OR^\circ)R^\circ$; $-C(=NH)-N(R^\circ)_2$; $-P(O)_2R^\circ$; $-PO(R^\circ)_2$; $-OPO(R^\circ)_2$; $-(CH_2)_{0-2}NHC(O)R^\circ$; phenyl (Ph) optionally substituted with R°; -O(Ph) optionally substituted with R°; $-(CH_2)_{1-2}(Ph)$, optionally substituted with R°; or -CH=CH(Ph), optionally substituted with R°; wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C_{1-6} aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, -O(Ph), or $-CH_2(Ph)$, or, notwithstanding the definition above, two independent occurrences of R°, on the same substituent or different substituents, taken together with the atom(s) to which each R° group is bound, to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0036] . Optional substituents on the aliphatic group of R° are selected from NH_2, $NH(C_{1-4}$aliphatic), $N(C_{1-4}$aliphatic$)_2$, halogen, C_{1-4}aliphatic, OH, $O(C_{1-4}$aliphatic), NO_2, CN, CO_2H, $CO_2(C_{1-4}$aliphatic), $O($haloC_{1-4} aliphatic), or haloC_{1-4}aliphatic, wherein each of the foregoing C_{1-4}aliphatic groups of R° is unsubstituted.

[0037] An aliphatic or heteroaliphatic group, or a non-aromatic heterocyclic ring may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the saturated carbon of an aliphatic or heteroaliphatic group, or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and additionally include the following: =O, =S, $=NNHR^*$, $=NN(R^*)_2$, $=NNHC(O)R^*$, =NNHCO$_2$(alkyl), =NNHSO$_2$(alkyl), or $=NR^*$, where each R^* is independently selected from hydrogen or an optionally substituted C_{1-6} aliphatic group.


[0038] Unless otherwise defined above and herein, optional substituents on the nitrogen of a non-aromatic heterocyclic ring are generally selected from $-R^+$, $-N(R^+)_2$, $-C(O)R^+$, $-CO_2R^+$, $-C(O)C(O)R^+$, $-C(O)CH_2C(O)R^+$, $-SO_2R^+$, $-SO_2N(R^+)_2$, $-C(=S)N(R^{+1})_2$, $-C(=NH)-N(R^+)_2$, or $-NR^+SO_2R^+$; wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, optionally

substituted phenyl, optionally substituted -O(Ph), optionally substituted -CH_2(Ph), optionally substituted -$(CH_2)_{1-2}$(Ph); optionally substituted -CH=CH(Ph); or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R^+, on the same substituent or different substituents, taken together with the atom(s) to which each R^+ group is bound, form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0039] Optional substituents on the aliphatic group or the phenyl ring of R^+ are selected from -NH_2, -NH(C_{1-4} aliphatic), -N(C_{1-4} aliphatic)$_2$, halogen, C_{1-4} aliphatic, -OH, -O(C_{1-4} aliphatic), -NO_2, -CN, -CO_2H, -CO_2(C_{1-4} aliphatic), -O(halo C_{1-4} aliphatic), or halo(C_{1-4} aliphatic), wherein each of the foregoing C_{1-4}aliphatic groups of R^+ is unsubstituted.


[0040] The term "alkylidene chain" refers to a straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation and has two points of attachment to the rest of the molecule.

[0041] As detailed above, in some embodiments, two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0042] Exemplary rings that are formed when two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which each variable is bound include, but are not limited to the following: a) two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein) that are bound to the same atom and are taken together with that atom to form a ring, for example, N(R°)$_2$, where both occurrences of R° are taken together with the nitrogen atom to form a piperidin-1-yl, piperazin-1-yl, or morpholin-4-yl group; and b) two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein) that are bound to different atoms and are taken together with both of those atoms to form a ring, for example where a phenyl group is substituted with two occurrences of OR° 

, these two occurrences of R° are taken together with

the oxygen atoms to which they are bound to form a fused 6-membered oxygen containing ring:



. It will be appreciated that a variety of other rings can be formed when two independent occurrences of R° (or R^{+}, R, R' or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound and that the examples detailed above are not intended to be limiting.

Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C- or ^{14}C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

[0043] *3. Description of Exemplary Compounds:*

[0044] In one embodiment, R is hydrogen. Or, R is C1-C6 aliphatic. Exemplary R includes C1-C6 alkyl, e.g., methyl, ethyl, propyl, or butyl.

[0045] In one embodiment, R' is hydrogen.

[0046] In one embodiment, R' is a C1-C8 aliphatic group, optionally substituted with up to 3 substituents selected from halo, CN, CF_3, CHF_2, OCF_3, or $OCHF_2$, wherein up to two methylene units of said C1-C8 aliphatic is optionally replaced with –CO-, -CONH(C1-C4 alkyl)-, -CO_2-, -OCO-, -N(C1-C4 alkyl)CO_2-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, -SO_2N(C1-C4 alkyl)-, N(C1-C4 alkyl)SO_2-, or -N(C1-C4 alkyl)SO_2N(C1-C4 alkyl)-.

[0047] In one embodiment, R' is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen,


oxygen, or sulfur, wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF_3, CHF_2, OCF_3, $OCHF_2$, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with -CO-, -CONH(C1-C4 alkyl)-, $-CO_2-$, -OCO-, -N(C1-C4 alkyl)CO_2-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, $-SO_2N$(C1-C4 alkyl)-, N(C1-C4 alkyl)SO_2-, or -N(C1-C4 alkyl)SO_2N(C1-C4 alkyl)-.

[0048] In one embodiment, R' is an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF_3, CHF_2, OCF_3, $OCHF_2$, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with –CO-, -CONH(C1-C4 alkyl)-, $-CO_2-$, -OCO-, -N(C1-C4 alkyl)CO_2-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, $-SO_2N$(C1-C4 alkyl)-, N(C1-C4 alkyl)SO_2-, or -N(C1-C4 alkyl)SO_2N(C1-C4 alkyl)-.

[0049] In one embodiment, two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF_3, CHF_2, OCF_3, $OCHF_2$, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with –CO-, -CONH(C1-C4 alkyl)-, $-CO_2-$, -OCO-, -N(C1-C4 alkyl)CO_2-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, $-SO_2N$(C1-C4 alkyl)-, N(C1-C4 alkyl)SO_2-, or -N(C1-C4 alkyl)SO_2N(C1-C4 alkyl)-.

[0050] In another embodiment, W is OH.


[0051] In still another embodiment, R^{XY} is:




[0052] In certain embodiment, Y is P and X is O.


[0053] In another embodiment, each Z is –O-.


[0054] In yet another embodiment, R^{XY} is selected from:




[0055] In yet another embodiment, R^{XY} is selected from:












-(L)-lysine, $-PO_3Na_2$,







-(L)-tyrosine,




$-PO_3Mg$, $-PO_3(NH_4)_2$, $-CH_2-OPO_3Na_2$,





-(L)-serine, $-SO_3Na_2$,





$-SO_3Mg$, $-SO_3(NH_4)_2$, $-CH_2-OSO_3Na_2$, $-CH_2-OSO_3(NH_4)_2$,
















, acetyl,




-(L)-valine, -(L)-glutamic acid, -(L)-aspartic acid, -(L)-γ-t-butyl-aspartic acid,








-(L)-3-pyridylalanine, -(L)-histidine, -CHO,






















, PO_3K_2, PO_3Ca, PO_3-spermine, PO_3-(spermidine)$_2$ or PO_3-(meglamine)$_2$.







[0056] In yet another embodiment, R^{XY} is selected from:


R^{XY}






R^{XY}






R^{XY}






R^{XY}








R^{XY}


-SO_3H
-SO_3H





R^{XY}
PO_3K_2
PO_3Ca
PO_3Mg


[0057] In one embodiment, x is 0-2. Or, x is 1 or 2. Or, x is 1.

[0058] In one embodiment, R^3 is present at the 6- or 7-position of the quinazoline ring.

[0059] In another embodiment, R^3 is selected from halo, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, -COOR', -NRCOR', $-CON(R')_2$, $-OCON(R')_2$, COR', -NHCOOR', $-SO_2R'$, $-SO_2N(R')_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl.

[0060] In one embodiment, R^3 is independently Cl, Br, F, CF_3, $-OCF_3$, Me, Et, CN, -COOH, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-OCH_3$, $-OCH_2CH_3$, $-CH_2OH$, $-NHCOCH_3$, $-NHCOCH(CH_3)_2$, $-SO_2NH_2$, -CONH(cyclopropyl), $-CONHCH_3$, $-CONHCH_2CH_3$, or an optionally substituted group selected from -piperidinyl, piperizinyl, morpholino, phenyl, phenyloxy, benzyl, or benzyloxy.

[0061] In another embodiment, each R^3 group is independently halogen, CN, optionally substituted C_1-C_6alkyl, OR', $N(R')_2$, $CON(R')_2$, or NRCOR'.

[0062] In one embodiment, x is 1 or 2, and each R^3 group is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

[0063] In yet another embodiment, x is 1 and R^3 is at the 6-position of the quinazoline ring and is selected from -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or –CN.

[0064] In yet another embodiment, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$.

[0065] In one embodiment, R^3 is at the 6-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'.

[0066] In another embodiment, x is 1 and R^3 is at the 7-position of the quinazoline ring and is selected from -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

[0067] In yet another embodiment, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$. Or, x is 1 and R^3 is at the 7-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'.

[0068] In one embodiment, y is 0-4 and R^5 is independently halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, - -NRCOR', $-CON(R')_2$, $-S(O)_2N(R')_2$, -OCOR', -

COR', -CO_2R', -OCON(R')$_2$, -NR'SO_2R', -OP(O)(OR')$_2$, -P(O)(OR')$_2$, -OP(O)$_2$OR', -P(O)$_2$OR', -PO(R')$_2$, -OPO(R')$_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl.

[0069] In another embodiment, R^5 is independently Cl, Br, F, CF_3, Me, Et, CN, -COOH, -NH_2, –N(CH_3)$_2$, -N(Et)$_2$, -N(iPr)$_2$, -O(CH_2)$_2$OCH_3, -$CONH_2$, -$COOCH_3$, -OH, -OCH_3, -OCH_2CH_3, -CH_2OH, -NHCOCH_3, -SO_2NH_2, -SO_2NHC(CH_3)$_2$, -OCOC(CH_3)$_3$, -OCOCH_2C(CH_3)$_3$, -O(CH_2)$_2$N(CH_3)$_2$, 4-CH_3-piperazin-1-yl, OCOCH(CH_3)$_2$, OCO(cyclopentyl), -$COCH_3$, optionally substituted phenoxy, or optionally substituted benzyloxy.

[0070] In certain embodiments, each of z_1, z_2, z_3, or z_4 is independently 0-2. In other embodiments, each of z_1, z_2, z_3, or z_4 is 0 and the ring is unsubstituted. Preferred R^4 groups, when present, are each independently halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NRCOR', -CON(R')$_2$, -OCON(R')$_2$, COR', -NHCOOR', -SO_2R', -SO_2N(R')$_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl. Other exemplary R^4 groups are Cl, Br, F, CF_3, CH_3, -CH_2CH_3, CN, -COOH, –N(CH_3)$_2$, -N(Et)$_2$, -N(iPr)$_2$, -O(CH_2)$_2$OCH_3, -$CONH_2$, -$COOCH_3$, -OH, -CH_2OH, -NHCOCH_3, -SO_2NH_2, -SO_2(CH_2)$_3$$CH_3$, -$SO_2$CH($CH_3$)$_2$, -$SO_2$N($CH_3$)$_2$, -$SO_2CH_2CH_3$, -C(O)O$CH_2$CH($CH_3$)$_2$, -C(O)NH$CH_2$CH($CH_3$)$_2$, -NHCO$OCH_3$, -C(O)C($CH_3$)$_3$, -COO($CH_2$)$_2CH_3$, -C(O)NHCH($CH_3$)$_2$, -C(O)$CH_2CH_3$, or an optionally substituted group selected from piperidinyl, piperizinyl, morpholino, C_{1-4}alkoxy, phenyl, phenyloxy, benzyl, benzyloxy, -CH_2cyclohexyl, pyridyl, -CH_2pyridyl, or -CH_2thiazolyl

[0071] In certain embodiments, x is 0-2. In other embodiments, x is 1 or 2. In still other embodiments x is 1 and R^3 is substituted at the 6- or 7-position of the quinazoline ring. When the quinazoline ring is substituted (x is 1-4), R^3 groups are halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NRCOR', -CON(R')$_2$, -OCON(R')$_2$, COR', -NHCOOR', -SO_2R', -SO_2N(R')$_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl. In still other embodiments, each occurrence of R^3 is independently Cl, Br, F, CF_3, -OCF_3, Me, Et, CN, -COOH, -NH_2, –N(CH_3)$_2$, -N(Et)$_2$, -N(iPr)$_2$, -O(CH_2)$_2$OCH_3, -$CONH_2$, -$COOCH_3$, -OH, -OCH_3, -OCH_2CH_3, -

CH_2OH, $-NHCOCH_3$, $-NHCOCH(CH_3)_2$, $-SO_2NH_2$, -CONH(cyclopropyl), $-CONHCH_3$, $-CONHCH_2CH_3$, or an optionally substituted group selected from -piperidinyl, piperizinyl, morpholino, phenyl, phenyloxy, benzyl, or benzyloxy. In still other embodiments, x is 1 or 2 and each R^3 group is independently halogen, CN, optionally substituted C_1-C_6alkyl, OR', $N(R')_2$, $CON(R')_2$, or NRCOR'. In yet other embodiments, x is 1 or 2, and each R^3 group is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN. In still other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or –CN. IN yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN. In other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$. In still other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$. In other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'. In yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'.

[0072] In some embodiments, y is 0-4 and R^5 group, when present, is each independently halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, - -NRCOR', $-CON(R')_2$, $-S(O)_2N(R')_2$, -OCOR', -COR', $-CO_2R'$, $-OCON(R')_2$, $-NR'SO_2R'$, $-OP(O)(OR')_2$, $-P(O)(OR')_2$, $-OP(O)_2OR'$, $-P(O)_2OR'$, $-PO(R')_2$, $-OPO(R')_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl.

[0073] In yet other embodiments, y is 0-4 and each occurrence of R^5 is independently Cl, Br, F, CF_3, Me, Et, CN, -COOH, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-OCH_3$, $-OCH_2CH_3$, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2NHC(CH_3)_2$, $-OCOC(CH_3)_3$, $-OCOCH_2C(CH_3)_3$, $-O(CH_2)_2N(CH_3)_2$, 4-CH_3-piperazin-1-yl, $OCOCH(CH_3)_2$, OCO(cyclopentyl), $-COCH_3$, optionally substituted phenoxy, or optionally substituted benzyloxy.

[0074] In yet another embodiment, each of z_1, z_2, z_3, or z_4 is 0-4, and R^4 groups, when present, are each independently halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR',

$-CH_2SR'$, $-COOR'$, $-NRCOR'$, $-CON(R')_2$, $-OCON(R')_2$, COR', $-NHCOOR'$, $-SO_2R'$, $-SO_2N(R')_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl.

[0075] In still other embodiments, each of z_1, z_2, z_3, or z_4 is 0-4 and R^4 groups are each independently Cl, Br, F, CF_3, CH_3, $-CH_2CH_3$, CN, -COOH, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2(CH_2)_3CH_3$, $-SO_2CH(CH_3)_2$, $-SO_2N(CH_3)_2$, $-SO_2CH_2CH_3$, $-C(O)OCH_2CH(CH_3)_2$, $-C(O)NHCH_2CH(CH_3)_2$, $-NHCOOCH_3$, $-C(O)C(CH_3)_3$, $-COO(CH_2)_2CH_3$, $-C(O)NHCH(CH_3)_2$, $-C(O)CH_2CH_3$, or an optionally substituted group selected from -piperidinyl, piperizinyl, morpholino, C_{1-4}alkoxy, phenyl, phenyloxy, benzyl, benzyloxy, $-CH_2$cyclohexyl, pyridyl, $-CH_2$pyridyl, or $-CH_2$thiazolyl.

[0076] For compounds described directly above, in some embodiments, x is 0-4, and R^3 groups, when present, are each independently halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, $-OR'$, $-CH_2OR'$, $-SR'$, $-CH_2SR'$, $-COOR'$, $-NRCOR'$, $-CON(R')_2$, $-OCON(R')_2$, COR', $-NHCOOR'$, $-SO_2R'$, $-SO_2N(R')_2$, or an optionally substituted group selected from C_1-C_6aliphatic, aryl, heteroaryl, cycloaliphatic, heterocycloaliphatic, arylC_1-C_6alkyl, heteroarylC_1-C_6alkyl, cycloaliphaticC_1-C_6alkyl, or heterocycloaliphaticC_1-C_6alkyl.

[0077] In yet other embodiments, x is 1 or 2, and each occurrence of R^3 is independently Cl, Br, F, CF_3, $-OCF_3$, Me, Et, CN, -COOH, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-OCH_3$, $-OCH_2CH_3$, $-CH_2OH$, $-NHCOCH_3$, $-NHCOCH(CH_3)_2$, $-SO_2NH_2$, -CONH(cyclopropyl), $-CONHCH_3$, $-CONHCH_2CH_3$, or an optionally substituted group selected from -piperidinyl, piperizinyl, morpholino, phenyl, phenyloxy, benzyl, or benzyloxy.

[0078] In still other embodiments, x is 1 or 2 and each R^3 group is independently halogen, CN, optionally substituted C_1-C_6alkyl, OR', $N(R')_2$, $CON(R')_2$, or NRCOR'.

[0079] In yet other embodiments, x is 1 or 2, and each R^3 group is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

[0080] In still other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

[0081] In yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

[0082] In still other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$.


[0083] In yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$.

[0084] In still other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'.

[0085] In yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$.

[0086] In yet other embodiments for compounds described directly above, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN. In still other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN. In yet other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$. In still other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-OCH_3$, or $-OCH_2CH_3$. In yet other embodiments, x is 1 and R^3 is at the 6-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'. In yet other embodiments, x is 1 and R^3 is at the 7-position of the quinazoline ring and is $-CON(R')_2$, or NRCOR'.

[0087] In one embodiment, the present invention provides compounds of formula **I-A**:



I-A;

wherein x, y, n_1, m_1, z_1, R^{XY}, R^4, R^5, Sp^1, and ring B^1 are as defined above.

[0088] In one embodiment, Sp^1 is selected from -O-, -S-, or –NR'-. Or, Sp^1 is –O-. Or, Sp^1 is $-O-CH_2-$. In another embodiment, Sp^1 is –NR'-. Or, Sp^1 is –NH-. Or, Sp^1 is $-NH-CH_2-$.

[0089] In one embodiment, each of m_1 and n_1 is 1. In another embodiment, each of m_1 and n_1 is 2.

[0090] In one embodiment, ring B^1 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w independent occurrences of $-R^{11}$, wherein w_1 is 0-4.

[0091] In another embodiment, ring B^1 is a 4-8 membered, saturated, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w independent occurrences of $-R^{11}$, wherein w_1 is 0-4.

[0092] In yet another embodiment, ring B^1 is a 5-6 membered, saturated, monocyclic heterocyclic ring having having 1-2 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w independent occurrences of $-R^{11}$, wherein w_1 is 0-4.

[0093] In one embodiment, w_1 is 0.

[0094] In another embodiment, ring B^1 is tetrahydrofuranyl.

[0095] In yet another embodiment, Sp^1 is a bond, O, or $-O-CH_2-$; R is hydrogen; and n_1 and m_1 are both simultaneously 1 or 2.

[0096] In one embodiment, R is hydrogen. Or, R is C1-C6 alkyl. Preferred R include methyl, ethyl, propyl, or butyl.

[0097] In another embodiment, z_1 is 0.

[0098] According to another embodiment, ring B^1 is tetrahydrofuranyl, tetrahydro-[*2H*]-pyranyl, pyridyl, or phenyl.

[0099] According to yet another embodiment, Sp^1 is a bond, -O-, $-O-CH_2-$, or $-NH-CH_2$.

[00100] In one embodiment:

n_1 and m_1 each is 2;

R^{xy} is hydrogen;

y is 0 or 1 and R^5 is fluoro;

x is 1 and R^3 is Me at 7-position or fluoro at 6-position;


z_1 is 0;

Sp^1 is $-O-CH_2-$;

w_1 is 0; and

ring B^1 is tetrahydrofuran-3-yl, phenyl, pyridine-3-yl, pyridine-4-yl, or tetrahydro[2H]-pyran-4-yl.

[00101] According to another embodiment, the present invention provides compounds of formula **I-B**:



I-B;

wherein x, y, n_2, m_2, z_2, q_2, R, Sp^2, ring B^2, R^{XY}, R^3, R^4, and R^5 are as defined above.

[00102] In one embodiment, G_2 is N. Or, G_2 is CH.

[00103] In one embodiment, p_2 is 0. Or, p_2 is 1. Or, p_2 is 2.

[00104] In another embodiment, q_2 is 0. Or, q_2 is 1.

[00105] In one embodiment, p_2 is 1, and q_2 is 1.

[00106] In another embodiment, G_2 is CH, p_2 is 0, and q_2 is 1.

[00107] In one embodiment, m_2 and n_2 each is 1. Or, m_2 and n_2 each is 2. Or, n_2 is 1 and m_2 is 2. Or, n_2 is 1, and m_2 is 3.

[00108] In another embodiment, Sp^2 is selected from -O-, -S-, or –NR'-. In one embodiment, Sp^2 is –O-. Or, Sp^2 is –NR'-. Or, Sp^2 is –NH-.

[00109] In one embodiment, ring B^2 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4.

[00110] In another embodiment, ring B^2 is a 4-8 membered, saturated, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^2 is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4.

[00111] In yet another embodiment, ring B^2 is a 5-6 membered, saturated, monocyclic heterocyclic ring having having 1-2 heteroatoms selected from O, S, or N, wherein ring B^2 is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4.

[00112] In one embodiment, w_2 is 0.

[00113] According to yet another embodiment, Sp^2 is a bond, -O-, or $-O-CH_2-$.

[00114] In another embodiment, ring B^2 is tetrahydrofuranyl, tetrahydro[2*H*]pyranyl, or pyridyl.

[00115] In yet another embodiment,

i) Sp^2 is a bond, O, or $-O-CH_2-$;


ii) p_2 is 1;

iii) R is hydrogen; and


iv) n_2 is 1 and m_2 is 2 or 3.

[00116] In one embodiment, R is hydrogen. Or, R is C1-C6 alkyl. Preferred R include methyl, ethyl, propyl, or butyl.

[00117] In one embodiment, compounds of formula **I-B** have formula **I-B-i** or formula **I-B-ii**:



I-B-i



I-B-ii.

[00118] In one embodiment of formula **I-B-i**:

i) p_2 is 1;

ii) m_2 is 3;

iii) Sp^2 is -O-;

iv) y is 0 or 1, and R^5 is fluoro;

v) x is 1 and R^3 is 7-Me; and

vi) ring B^2 is tetrahydrofuranyl.

[00119] In another embodiment of formula **I-B-i**:

i) p_2 is 0 or 1;

ii) m_2 is 1 or 2, preferably 2;

iii) Sp^2 is -O- or -O-CH_2-;

iv) y is 0;

v) x is 1 and R^3 is 7-Me; and

vi) ring B^2 is tetrahydrofuranyl, tetrahydro[2H]pyranyl, pyridyl, or phenyl.

[00120] In one embodiment of formula **I-B-ii**:

(i) n_2 is 1, m_2 is 1 or 2, preferably 2;

(ii) y is 0 or 1, and R^5 is fluoro;

(iii) x is 1 and R^3 is 7-Me or 6-F; and

(iv) ring B^2 is cyclopropyl optionally substituted with C1-C4 alkyl, or pyridyl.

[00121] In one embodiment of formula **I-B-ii**:

(i) n_2 and m_2 both are 2;


(ii) y is 0;

(iii) x is 1 and R^3 is C1-C4 alkyl at the 7-position; and


(iv) ring B^2 is an optionally substituted tetrahydrofuranyl.

[00122] In one embodiment of formula **I-B-i** or formula **I-B-ii**, R^{XY} is hydrogen.

[00123] According to one embodiment, the present invention provides compounds of formula **I-C** or formula **I-D**:



I-C



I-D;

wherein x, y, n_3, m_3, z_3, p_3, R^{XX}, R^{YY}, R^{XY}, R^3, R^4, and R^5 are as defined above.

[00124] In one embodiment of the present invention, one R^{XX} is hydrogen and the other R^{XX} is not hydrogen.

[00125] In another embodiment of the present invention, both R^{XX} are not hydrogen.

[00126] In another embodiment, one R^{XX} is hydrogen and the other R^{XX} is C1-C6 alkyl optionally substituted with halo. Or, both R^{XX} are simultaneously C1-C6 alkyl. Exemplary alkyl include methyl, ethyl, isopropyl, *n*-propyl, *n*-butyl, *sec*-butyl, or *t*-butyl.

[00127] In one embodiment of the present invention, p_3 is 0. Or, p_3 is 1. Or, p_3 is 2.

[00128] In one embodiment of the present invention, m_3 and n_3 each is 1. Or, m_3 and n_3 each is 2. Or, m_3 and n_3 each is 3.

[00129] In one embodiment of the present invention, R^{XX} is C_{1-6} aliphatic group, wherein R^{XX} is optionally substituted with w independent occurrences of $-R^{13}$, wherein w_3 is 0-3. Or, R^{XX} is

C1-C6 alkyl group optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3.

[00130] In one embodiment of the present invention, R^{XX} is C1-C6 alkyl group.

[00131] In another embodiment of the present invention, R^{XX} is a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3.

[00132] In another embodiment, R^{XX} is a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^{XX} is optionally substituted with w independent occurrences of $-R^{13}$, wherein w_3 is 0-3.

[00133] In another embodiment, R^{XX} is an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3.

[00134] In another embodiment, R^{YY} is hydrogen, -COR', $-CO_2R'$, $-CON(R')_2$, -SOR', $-SO_2R'$, $-SO_2N(R')_2$, -COCOR', $-COCH_2COR'$, $-P(O)(OR')_2$, $-P(O)_2OR'$, or -PO(R').


[00135] Or, R^{YY} is hydrogen.

[00136] In another embodiment, R^{YY} is -COR', $-CO_2R'$, $-CON(R')_2$, -SOR', $-SO_2R'$, $-SO_2N(R')_2$, -COCOR', $-COCH_2COR'$, $-P(O)(OR')_2$, $-P(O)_2OR'$, or -PO(R').


[00137] In another embodiment, R^{YY} is R^{XY}.

[00138] In one embodiment, R is hydrogen. Or, R is C1-C6 alkyl. Preferred R include methyl, ethyl, propyl, or butyl.

[00139] In one embodiment, the present invention provides a compound of formula **I-C-i** or formula **I-D-i**:



I-C-i



I-D-i;

wherein x, y, n_3, m_3, z_3, p_3, R^{XX}, R^{YY}, R^{XY}, R^3, R^4, and R^5 are as defined above.


[00140] In one embodiment of I-C-i or I-D-i, R^{XX} is C1-C6 alkyl. In another embodiment, x is 1, and R^3 is C1-C4 alkyl at the 7-position. Or, x is 1 and R^3 is F, CN, or CF_3 at the 6-position.

[00141] In one embodiment, R^{XX} is methyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, or *t*-butyl.

[00142] In one embodiment, R^3 is C1-C6 alkyl. Or, R^3 is methyl, *n* -propyl, isopropyl, *n* -butyl, *sec*-butyl, or *t*-butyl.

[00143] In one embodiment of I-C-i or I-D-i, R^{XY} is hydrogen, and y is 0. Or, R^{XY} is hydrogen, y is 1 and R^5 is 6-F.

[00144] In another embodiment, the present invention provides a compound of formula **I-C-ii**:




I-C-ii;

wherein R^3 and R^{XX} are as defined above.

[00145] In another embodiment, R^3 is methyl at the 6- or 7-position of the quinazoline ring.

[00146] In another embodiment of formula I-C-ii, R^{XX} is $CH_2C(O)OH$ or $CH_2C(O)NH_2$.

[00147] According to one embodiment, the present invention provides a compound of formula **I-E**:



I-E;

wherein x, y, n_4, m_4, z_4, p_4, R^{YZ}, R^{XY}, R^3, R^4, and R^5 are as defined above.

[00148] In one embodiment, p_4 is 1. Or, p_4 is 2.

[00149] In one embodiment, m_4 and n_4 each is 1. Or, m_4 and n_4 each is 2. Or, m_4 and n_4 each is 3. In one embodiment, n_4 is 1 and m_4 is 3. In another embodiment, n_4 is 1 and m_4 is 2.

[00150] In one embodiment, n_4 is 1, m_4 is 3, z_4 is 0, p_4 is 1, y is 0 or 1, and x is 1.

[00151] In another embodiment, n_4 is 1, m_4 is 2, z_4 is 0, p_4 is 1, y is 0 or 1, and x is 1.

[00152] In one embodiment, n_4 is 1, m_4 is 3, z_4 is 0, p_4 is 1, y is 0 or 1, x is 1, and R and R^{XY} both are hydrogen.

[00153] In another embodiment, n_4 is 1, m_4 is 2, z_4 is 0, p_4 is 1, y is 0 or 1, x is 1, and R and R^{XY} both are hydrogen.

[00154] In one embodiment, R^{YZ} is C1-C6 alkyl, optionally substituted with w_4 independent occurrences of $-R^{14}$, wherein w_4 is 0-3. In another embodiment, R^{YZ} is C1-C4 alkyl group optionally substituted with w_4 independent occurrences of $-R^{14}$, wherein w_4 is 0-3. Or, R^Y is C1-C6 alkyl group.

[00155] In one embodiment, R is hydrogen. Or, R is C1-C6 alkyl. Preferred R include methyl, ethyl, propyl, or butyl.

[00156] In another embodiment:

(i) n_4 is 1 and m_4 is 3;

(ii) p_4 is 1;

(iii) z_4 is 0;

(iv) R^{YZ} is C1-C6 alkyl, wherein up to two -CH_2- groups therein is optionally replaced by -O-;

(v) y is 0 or 1, and R^5 is 6-fluoro; and

(vi) x is 1 and R^3 is C1-C4 alkyl.

[00157] In another embodiment:

(i) n_4 is 1 and m_4 is 2;

(ii) p_4 is 1;

(iii) z_4 is 0;

(iv) R^{YZ} is C1-C6 alkyl, wherein up to two -CH_2- groups therein is optionally replaced by -O-;

(v) y is 0 or 1, and R^5 is 6-fluoro; and

(vi) x is 1 and R^3 is C1-C4 alkyl.

[00158] In another embodiment:

(i) n_4 is 1 and m_4 is 3;

(ii) p_4 is 1;


(iii) z_4 is 0;

(iv) R^{YZ} is benzyl;

(v) y is 0 or 1, and R^5 is 6-fluoro; and

(vi) x is 1 and R^3 is C1-C4 alkyl.


[00159] In another embodiment, the present invention provides compounds of formula **I-F**:




I-F;

wherein:


R^1 and R^2, taken together with the nitrogen atom, form a substituted ring selected from:




(A)




(B)



(C)



(D) or



(E);

wherein, in ring (A):

each of m_1 and n_1 is independently 0-3, provided that m_1+n_1 is 2-6;

z_1 is 0-4;

Sp^1 is –O-, -S-, -NR'-, or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-, provided that Sp^1 is attached to the carbonyl group through an atom other than carbon;

ring B^1 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w_1 independent occurrences of -R^{11}, wherein w_1 is 0-4;

wherein, in ring (B):

G_2 is -N-, or CH;

each of m_2 and n_2 is independently 0-3, provided that $m_2 + n_2$ is 2-6;

p_2 is 0-2; provided that when G_2 is N, then p_2 is not 0;

q_2 is 0 or 1;

z_2 is 0-4;

Sp^2 is a bond or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-;

ring B^2 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4;

wherein, in ring (C) or ring (D):

G_3 is -N-, -CH-NH-, or -CH-CH_2-NH-;

each of m_3 and n_3 is independently 0-3, provided that m_3+n_3 is 2-6;

p_3 is 0-2;

z_3 is 0-4;

each R^{XX} is hydrogen, C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3;

provided that both R^{XX} are not simultaneously hydrogen;

R^{YY} is hydrogen, -COR', -CO_2R', -CON(R')$_2$, -SOR', -SO_2R', -SO_2N(R')$_2$, -COCOR', -$COCH_2$COR', -P(O)(OR')$_2$, -P(O)$_2$OR', or -PO(R');

wherein, in ring (E):

each of m_4 and n_4 is independently 0-3, provided that $m_4 + n_4$ is 2-6;

p_4 is 1-2;


z_4 is 0-4;

R^{YZ} is C_1-C_6 aliphatic group, optionally substituted with w_4 independent occurrences of -R^{14}, wherein w_4 is 0-3;

x and y, each, is independently 0-4;

W is OR^{XY};

R^{XY} is hydrogen or a group selected from:



;

wherein:

each of w_A, w_B, w_C, and w_D is independently 0 or 1;

each M is independently selected from hydrogen, Li, Na, K, Mg, Ca, Ba, -$N(R^7)_4$, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or -R^6; wherein 1 to 4 –CH_2 radicals of the alkyl or alkenyl group, other than the –CH_2 that is bound to Z, is optionally replaced by a heteroatom group selected from O, S, S(O), $S(O_2)$, or $N(R^7)$; and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a substituent selected from oxo, -OR^7, - R^7, $N(R^7)_2$, $N(R^7)_3$, R^7OH, -CN, -CO_2 R^7, -C(O)-$N(R^7)_2$, $S(O)_2$-$N(R^7)_2$, $N(R^7)$-C(O)- R^7, C(O) R^7, -$S(O)_n$- R^7, OCF_3, -$S(O)_n$-R^6, $N(R^7)$-$S(O)_2(R^7)$, halo, -CF_3, or -NO_2;

n is 0-2;

M' is H, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or -R^6; wherein 1 to 4 –CH_2 radicals of the alkyl or alkenyl group is optionally replaced by a heteroatom group selected from O, S, S(O), $S(O_2)$, or $N(R^7)$; and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a substituent selected from oxo, -O R^7, - R^7, -$N(R^7)_2$, $N(R^7)_3$, - R^7OH, -CN, -CO_2 R^7, -C(O)-$N(R^7)_2$, -$S(O)_2$-$N(R^7)_2$, -$N(R^7)$-C(O)- R^7, -C(O) R^7, -$S(O)_n$- R^7, -OCF_3, -$S(O)_n$-R^6, -$N(R^7)$-$S(O)_2(R^7)$, halo, -CF_3, or -NO_2;

Z is -CH_2-, -O-, -S-, -$N(R^7)_2$-; or,

when M is absent, then Z is hydrogen, =O, or =S;

Y is P or S, wherein when Y is S, then Z is not S;

X is O or S;

each R^7 is independently selected from hydrogen, or C_1-C_4 aliphatic, optionally substituted with up to two Q_1;

each Q_1 is independently selected from a 3-7 membered saturated, partially saturated or unsaturated carbocyclic ring system; or a 5-7 membered saturated, partially saturated or unsaturated heterocyclic ring containing one or more heteroatom or heteroatom group selected from O, N, NH, S, SO, or SO_2; wherein Q_1 is optionally substituted with up to three substituents selected from oxo, -OH, -O(C_1-C_4 aliphatic), -C_1-C_4 aliphatic, -NH_2, NH(C_1-C_4 aliphatic), -N(C_1-C_4 aliphatic)$_2$, -N(C_1-C_4 aliphatic)-C(O)- C_1-C_4 aliphatic, -(C_1-C_4 aliphatic)-OH, -CN, -CO_2H, -CO_2(C_1-C_4 aliphatic), -C(O)-NH_2, -C(O)-NH(C_1-C_4 aliphatic), -C(O)-N(C_1-C_4 aliphatic)$_2$, halo or -CF_3;

R^6 is a 5-6 membered saturated, partially saturated or unsaturated carbocyclic or heterocyclic ring system, or an 8-10 membered saturated, partially saturated or unsaturated bicyclic ring system; wherein any of said heterocyclic ring systems contains one or more heteroatoms selected from O, N, S, $S(O)_n$ or $N(R^7)$; and wherein any of said ring systems optionally contains 1 to 4 substituents independently selected from OH, C_1-C_4 alkyl, O-C_1-C_4 alkyl or O-C(O)-C_1-C_4 alkyl;

R^9 is $C(R^7)_2$, O or $N(R^7)$;

each occurrence of R^{11}, R^{12}, R^{13}, R^{14}, R^3, R^4, and R^5 is independently Q-R^X; wherein Q is a bond or is a C_1-C_6 alkylidene chain wherein up to two non-adjacent methylene units of Q are optionally and independently replaced by –NR-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -$NRCO_2$-, -SO_2NR-, -$NRSO_2$-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -$NRSO_2$NR-, -SO-, -SO_2-, -PO-, -PO_2-, -OP(O)(OR)-, or -POR-; and each occurrence of R^X is independently selected from -R', halogen, =O, =NR', -NO_2, -CN, -OR', -SR', -N(R')$_2$, -NR'COR', -NR'CON(R')$_2$, -NR'CO_2R', -COR', -CO_2R', -OCOR', -CON(R')$_2$, -OCON(R')$_2$, -SOR', -SO_2R', -SO_2N(R')$_2$, -NR'SO_2R', -NR'SO_2N(R')$_2$, -COCOR', -$COCH_2$COR', -OP(O)(OR')$_2$, -P(O)(OR')$_2$, -OP(O)$_2$OR', -P(O)$_2$OR', -PO(R')$_2$, or -OPO(R')$_2$; and













each occurrence of R is independently hydrogen or C_{1-6} aliphatic group having up to three substituents; and each occurrence of R' is independently hydrogen or C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' has up to four substituents; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s)













to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.













[00160] In compounds of formula **I-F**, exemplary embodiments of n_1, m_1, n_2, m_2, n_3, m_3, n_4, m_4, x, y, z_1, z_2, z_3, z_4, q_2, p_3, p_4, R^{XY}, R^{XX}, R^{YY}, R^{YZ}, R^3, R^4, R^5, R^{11}, R^{12}, R^{13}, R^{14}, R, Sp^1, Sp^2, ring B^1, ring B^2, etc., are as defined above for compounds of formula **I-A** through formula **I-E**.













[00161] In one embodiment, the present invention provides compounds shown below in **Table 2**.













Table 2













101 	102 	103 
104 	105 	106 
107 	108 	109 
110 	111 	112 













113	114	115
		
116	117	118
		
119	120	121
		
122	123	124
		













125	126	127
		
128	129	130
		
131	132	133
		
134	135	136
		













137	138	139
		
140	141	142
		
143	144	145
		
146	147	148
		













149	150	151
		
152	153	154
		
155	156	157
		
158	159	160
		













162	163	164
		
165	166	167
		
168	169	170
		
171	172	173
		













174	175	176
		
177	178	179
		
180	181	182
		
183	184	185
		













201	203	204
		
205	206	207
		
208	209	210
		
211	212	213
		













214	215	216
		
217	218	219
		
220	221	222
		
223	224	225
		













226	227	228
		
229	230	231
		
232	233	234
		
235	236	237
		













238	239	240
		
241	242	243
		
244	245	246
		
247	248	249
		













250	251	252
		
253	254	255
		
256	257	258
		
259	260	261
		













262	264	265
		
266	267	268
		
269	270	271
		
272	273	274
		













275	276	277
		
278	279	301
		
302	303	304
		
305	306	307
		













308	309	310
		
311	312	313
		
314	315	316
		
317	318	319
		







320	321	322
		
323	324	325
		
326	327	328
		
329	330	331
		













332	333	334
		
335	336	337
		
338	339	340
		
341	342	343
		



344	345	346
		
347	348	349
		
350	351	352
		
353	354	355
		

356	357	358
		
359	360	361
		
362	363	364
		
365	366	367
		

368	369	370
		
371	372	373
		
374	375	376
		
377	401	402
		

415	416	417
		
418	**419**	**420**
		


501	502	503
		
504	505	506
		
507	508	509
		
510	511	512
		

513	514
	

[00162] *4. General Synthetic Methodology:*


[00163] The compounds of this invention may be prepared in general by methods known to those skilled in the art for analogous compounds, as illustrated by the general schemes below, and the preparative examples that follow.

[00164] Scheme A: General Preparations via 4-Chloroquinazolines




[00165] Conditions: a) aq. NH_4OH; b) chloral hydrate, HCl, Na_2SO_4, $HONH_2$*HCl c) H_2SO_4; d) acetic acid, H_2SO_4, aq. H_2O_2; e) Et_3N, THF; f) Et_3N, DMAP, CH_2Cl_2; g) aq. NaOH; h) aq. NaOH, aq. H_2O_2 i) $POCl_3$, N,N-dimethylaniline, benzene; j) R^1R^2NH, Et_3N, CH_2Cl_2

[00166] Scheme B: General Preparations via 2,4-Dichloroquinazolines




[00167] Conditions: a) AcOH, KOCN; b) $POCl_3$; c) Et_3N, DCM, R^1R^2NH; d) $Pd(PPh_3)_4$, K_2CO_3, CH_3CN, H_2O

[00168] Scheme **I-A** for formula **I-A**:


I-A
X=Cl, Br, F, OH, OR'
PG=Boc, Bn, Bz, R3Si, etc.


[00169] Conditions: a) DCM or THF, triethylamine, 0 °C to room temperature; b) DCM or THF, triethylamine, 0 °C to room temperature; c) i. DCM or THF, triethylamine, 0 °C to room temperature, ii. Deprotect: 1:1 TFA / DCM, rt, for Boc; H_2, Pd / C for Bn; NaOH for Bz, TBAF for R3Si, etc.; d) For acid halides, DCM or THF, triethylamine; for carboxylic acids, EDC, HOBt, triethylamine, DMF; for X=OR', THF or DMF, heat.

[00170] Scheme **I-B** for compounds of formula **I-B**:




[00171] Conditions: a) DCM or THF, triethylamine, 0 °C to room temperature; b) DCM or THF, triethylamine, 0 °C to room temperature; c) i. DCM or THF, triethylamine, 0 °C to room temperature, ii. Deprotect: 1:1 TFA / DCM, rt, for Boc; H_2, Pd / C for Bn; NaOH for Bz, TBAF for R3Si, etc.; d) For acid halides, DCM or THF, triethylamine; for carboxylic acids, EDC, HOBt, triethylamine, DMF; for X=OR', THF or DMF, heat.

[00172] Scheme **I-C** for formula **I-C:**




[00173] Conditions: a) DCM or THF, triethylamine, 0 °C to room temperature; b) DCM or THF, triethylamine, 0 °C to room temperature; c) i. DCM or THF, triethylamine, 0 °C to room temperature, ii. Deprotect: 1:1 TFA / DCM, rt, for Boc; H_2, Pd / C for Bn; NaOH for Bz, TBAF for R3Si, etc.; d) For acid halides, DCM or THF, triethylamine; for carboxylic acids, EDC, HOBt, triethylamine, DMF; for X=OR', THF or DMF, heat.

[00174] Scheme **I-D** for formula **I-D**:




[00175] Conditions: a) DCM or THF, triethylamine, 0 °C to room temperature; b) DCM or THF, triethylamine, 0 °C to room temperature; c) i. DCM or THF, triethylamine, 0 °C to room temperature, ii. Deprotect: 1:1 TFA / DCM, rt, for Boc; H_2, Pd / C for Bn; NaOH for Bz, TBAF for R3Si, etc.; d) For acid halides, DCM or THF, triethylamine; for carboxylic acids, EDC, HOBt, triethylamine, DMF; for X=OR', THF or DMF, heat.

[00176] Scheme **I-E** for formula **I-E**:



[00177] Conditions: a) DCM or THF, triethylamine, 0 °C to room temperature; b) DCM or THF, triethylamine, 0 °C to room temperature; c) i. DCM or THF, triethylamine, 0 °C to room temperature, ii. Deprotect: 1:1 TFA / DCM, rt, for Boc; H_2, Pd / C for Bn; NaOH for Bz, TBAF for R3Si, etc.; d) For X=Cl, CCl_3, imidazolyl, OR', etc., DCM or THF, triethylamine, rt or heat.

[00178] Scheme **I-F** for preparing compounds of formula **I-F**:



[00179] Conditions: a) AcOH, KOCN; b) $POCl_3$; c) Et_3N, DCM, R^1R^2NH; d) $Pd(PPh_3)_4$, K_2CO_3, CH_3CN, H_2O.

[00180] *5. Uses, Formulation and Administration*

[00181] WO 2004/078733 discloses a genus of sodium channel blockers that encompasses the compounds of the present invention. However, the compounds of the present invention exhibit unexpected properties set forth below that render them therapeutically more useful.

[00182] In one embodiment, certain compounds of the present invention are useful as improved inhibitors of sodium channels.

[00183] In another embodiment, certain compounds of the present invention possess improved selectivity in inhibiting one sodium channel, e.g., NaV 1.8, over one or more of the other sodium channels. Particularly useful are compounds that have a desirably low activity against NaV 1.2 or NaV 1.5.

[00184] In another embodiment, certain compounds of the present invention are improved inhibitors of NaV 1.8.

[00185] In another embodiment, certain compounds of the present invention have improved aqueous solubility, e.g., at physiologically relevant pH.

[00186] In yet another embodiment, certain compounds of the present invention have improved pharmacokinetic and/or pharmacodynamic properties and, therefore, are better suited for *in-vivo* administration for therapeutic purposes. Such properties include oral bioavailability, clearance kinetics, efficacy, etc.

[00187] In another embodiment, certain compounds of the present invention have desirably low activity against the hERG channel.

[00188] In another embodiment, certain compounds of the present invention have desirably low activity against the key isoforms of the cytochrome P450 enzyme family, including isozymes CYP3A4, CYP2C9, CYP1A2, CYP2C19, or CYP2D6.

[00189] In another embodiment, certain compounds of the present invention have desirably low activity against the CaV 1.2 channel and/or Kv 1.5.

[00190] Thus, in one embodiment of the present invention, the compounds have one or more of the following unexpected and therapeutically beneficial features: potent inhibition of NaV 1.8 channel, selectivity for one sodium channel, e.g., NaV 1.8 over one or more of the other sodium channels, improved aqueous solubility, improved pharmacokinetic and/or pharmacodynamic properties, desirably low activity against the hERG channel, desirably low activity against the key isoforms of the cytochrome P450 enzyme family, or desirably low activity against L-type CaV 1.2 and/or Kv1.5. The presence of such features, individually or in combination, renders the compounds more suitable for administration to humans to treat various diseases as set forth below.

[00191] The phrase "desirably low activity" as used herein means a level of activity of a compound against a target/enzyme that is low enough such that said activity would be considered advantageous (e.g., mitigating a risk factor), when evaluating the suitability of said compound for administration in humans.

[00192] The present compounds are useful for the treatment of diseases, disorders, and conditions including, but not limited to acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias,

epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, and incontinence. Accordingly, in another aspect of the present invention, pharmaceutically acceptable compositions are provided, wherein these compositions comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents.

[00193] According to one embodiment, the compounds of the present invention are useful for treating a disease selected from femur cancer pain; non-malignant chronic bone pain; rheumatoid arthritis; osteoarthritis; spinal stenosis; neuropathic low back pain; neuropathic low back pain; myofascial pain syndrome; fibromyalgia; temporomandibular joint pain; chronic visceral pain, including, abdominal; pancreatic; IBS pain; chronic headache pain; migraine; tension headache, including, cluster headaches; chronic neuropathic pain, including, post-herpetic neuralgia; diabetic neuropathy; HIV- associated neuropathy; trigeminal neuralgia; Charcot-Marie Tooth neuropathy; hereditary sensory neuropathies; peripheral nerve injury; painful neuromas; ectopic proximal and distal discharges; radiculopathy; chemotherapy induced neuropathic pain; radiotherapy-induced neuropathic pain; post-mastectomy pain; central pain; spinal cord injury pain; post-stroke pain; thalamic pain; complex regional pain syndrome; phanton pain; intractable pain; acute pain, acute post-operative pain; acute musculoskeletal pain; joint pain; mechanical low back pain; neck pain; tendonitis; injury/exercise pain; acute visceral pain, including, abdominal pain; pyelonephritis; appendicitis; cholecystitis; intestinal obstruction; hernias; etc; chest pain, including, cardiac Pain; pelvic pain, renal colic pain, acute obstetric pain, including, labor pain; cesarean section pain; acute inflammatory, burn and trauma pain; acute intermittent pain, including, endometriosis; acute herpes zoster pain; sickle cell anemia; acute pancreatitis; breakthrough pain; orofacial pain, including, sinusitis pain, dental pain; multiple sclerosis (MS) pain; pain in depression; leprosy pain; behcet's disease pain; adiposis dolorosa; phlebitic pain; Guillain-Barre pain; painful legs and moving toes; Haglund syndrome; erythromelalgia pain; Fabry's disease pain; bladder and urogenital disease, including, urinary incontinence; hyperactivity bladder; painful bladder syndrome; interstitial cyctitis (IC); and prostatitis.

[00194] In another embodiment, the compounds of the present invention are useful in treating lower urinary tract disorders. See, e.g., International Patent Publication No. WO 2004/066990, the contents of which are incorporated herein by reference.

[00195] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adduct or derivative which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[00196] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of the targeted channel.

[00197] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, **1977**, *66*, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate,

heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersable products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[00198] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates,

waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[00199] *Uses of Compounds and Pharmaceutically Acceptable Compositions*

[00200] In yet another aspect, a method for the treatment or lessening the severity of acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias, epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, incontinence, visceral pain, osteoarthritis pain, postherpetic neuralgia, diabetic neuropathy, radicular pain, sciatica, back pain, head or neck pain, severe or intractable pain, nociceptive pain, breakthrough pain, postsurgical pain, or cancer pain is provided comprising administering an effective amount of a compound, or a pharmaceutically acceptable composition comprising a compound to a subject in need thereof. In certain embodiments, a method for the treatment or lessening the severity of acute, chronic, neuropathic, or inflammatory pain is provided comprising administering an effective amount of a compound or a pharmaceutically acceptable composition to a subject in need thereof. In certain other embodiments, a method for the treatment or lessening the severity of radicular pain, sciatica, back pain, head pain, or neck pain is provided comprising administering an effective amount of a compound or a pharmaceutically acceptable composition to a subject in need thereof. In still other embodiments, a method for the treatment or lessening the severity of severe or intractable pain, acute pain, postsurgical pain, back pain, or cancer pain is provided comprising administering an

effective amount of a compound or a pharmaceutically acceptable composition to a subject in need thereof.

[00201] In certain embodiments of the present invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of one or more of acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias, epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, incontinence, visceral pain, osteoarthritis pain, postherpetic neuralgia, diabetic neuropathy, radicular pain, sciatica, back pain, head or neck pain, severe or intractable pain, nociceptive pain, breakthrough pain, postsurgical pain, or cancer pain.

[00202] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of one or more of acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias, epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, incontinence, visceral pain, osteoarthritis pain, postherpetic neuralgia, diabetic neuropathy, radicular pain, sciatica, back pain, head or neck pain, severe or intractable pain, nociceptive pain, breakthrough pain, postsurgical pain, or cancer pain. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the

specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[00203] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), bucally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00204] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00205] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For

this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00206] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00207] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[00208] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00209] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain

silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[00210] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polethylene glycols and the like.

[00211] The active compounds can also be in microencapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00212] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants

or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms are prepared by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00213] It will also be appreciated that the compounds and pharmaceutically acceptable compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutically acceptable compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another agent used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated". For example, exemplary additional therapeutic agents include, but are not limited to: nonopioid analgesics (indoles such as Etodolac, Indomethacin, Sulindac, Tolmetin; naphthylalkanones such sa Nabumetone; oxicams such as Piroxicam; para-aminophenol derivatives, such as Acetaminophen; propionic acids such as Fenoprofen, Flurbiprofen, Ibuprofen, Ketoprofen, Naproxen, Naproxen sodium, Oxaprozin; salicylates such as Asprin, Choline magnesium trisalicylate, Diflunisal; fenamates such as meclofenamic acid, Mefenamic acid; and pyrazoles such as Phenylbutazone); or opioid (narcotic) agonists (such as Codeine, Fentanyl, Hydromorphone, Levorphanol, Meperidine, Methadone, Morphine, Oxycodone, Oxymorphone, Propoxyphene, Buprenorphine, Butorphanol, Dezocine, Nalbuphine, and Pentazocine). Additionally, nondrug analgesic approaches may be utilized in conjunction with administration

of one or more compounds of the invention. For example, anesthesiologic (intraspinal infusion, neural blocade), neurosurgical (neurolysis of CNS pathways), neurostimulatory (transcutaneous electrical nerve stimulation, dorsal column stimulation), physiatric (physical therapy, orthotic devices, diathermy), or psychologic (cognitive methods-hypnosis, biofeedback, or behavioral methods) approaches may also be utilized. Additional appropriate therapeutic agents or approaches are described generally in The Merck Manual, Seventeenth Edition, Ed. Mark H. Beers and Robert Berkow, Merck Research Laboratories, 1999, and the Food and Drug Administration website, www.fda.gov, the entire contents of which are hereby incorporated by reference.

[00214] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[00215] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccarides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition.

[00216] Another aspect of the invention relates to inhibiting NaV1.8 activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of the present invention or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[00217] Inhibition of NaV1.8 activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, the study of sodium ion channels in biological and pathological phenomena; and the comparative evaluation of new sodium ion channel inhibitors.

[00218] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLES


[00219] General LC/MS Methods

[00220] LC/MS data were acquired using a PESciex API-150-EX LC/MS, Shimadzu LC-8A pumps, Gilson 215 autosampler, Gilson 819 injection module, 3.0 mL/min flow rate, 10-99% CH_3CN (0.035 % TFA) / H_2O (0.05 % TFA) gradient, Phenomenex Luna 5u C18 column (50 x 4.60 mm), Shimadzu SPD-10A UV/Vis detector, Cedex 75 ELSD detector.

[00221] Example 101


[00222] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one**

[00223]



[00224] ***N*-(2-Cyano-5-methyl-phenyl)-2-methoxy-benzamide**


[00225]



[00226] To a stirred solution of 4-methyl-2-aminobenzonitrile (100 g, 0.75 mol) in 800 mL CH_2Cl_2 was added triethylamine (77.4 g, 0.76 mol) and dimethylaminopyridine (4.62 g, 0.037 mol). The solution was cooled to 0–5 °C, and *o*-anisoyl chloride (129 g, 0.75 mol) was added over 1 h while maintaining the reaction temperature at 0–5 °C. The reaction was then stirred at 30–40 °C for 3 h. Water (400 mL) was added, and the mixture was stirred for 15 minutes. The organic layer was separated, and the aqueous solution was extracted with CH_2Cl_2 (600 mL). The combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuo to yield a solid residue, to which 800 mL hexane were added. The slurry was stirred and filtered to give *N*-(2-cyano-5-methyl-phenyl)-2-methoxy-benzamide as a yellow powder (180 g, 90%). mp 147–149 °C. ^{1}H NMR ($CDCl_3$) δ 2.429 (s, 3H), 4.2 (s, 3H), 6.8–7.2 (m, 3H), 7.4–7.6 (m, 2H), 8.2–8.4 (d, 1H), 8.6 (s, 1H), 10.8 (bs, 1H); ^{13}C NMR ($CDCl_3$) δ 22.68, 55.7, 99, 111.27, 116.7, 120.3, 121.1, 124.15, 131.7, 132.25, 133.67, 141.32, 141.1, 157.2, 163. M/z (obs., $[m+H]^+$) = 268.

[00227] **2-(2-Methoxyphenyl)-7-methyl-3*H*-quinazolin-4-one**

[00228]




[00229] To a mechanically stirred suspension of *N*-(2-cyano-5-methylphenyl)-2-methoxybenzamide (180 g, 0.67 mol) in 1.8 L ethanol under an N_2 atmosphere was added 6 N sodium hydroxide solution (310 g in 1.25 L water). To the above mixture, 30% hydrogen peroxide (350 mL, 3.64 mol) was slowly added. The solution was then slowly heated to 80 °C

and maintained at this temperature for 4 h. The reaction mixture was concentrated under reduced pressure to remove ethanol, giving a suspension which was quenched with ice cold water (1.8 L) and acidified with acetic acid to pH 5–6 to give a solid residue. The solid was filtered and washed with water, then dissolved in 5.5 L CH_2Cl_2 and washed with water (2x18 L). The organic layer was dried over sodium sulfate, and the solvent was removed under reduced pressure to give a light yellow solid (100 g, 54%). mp165–170 °C. ^{1}H NMR ($CDCl_3$) δ 2.429 (s, 3H), 4.2 (s,3), 6.8–7.2 (m, 3H), 7.4–7.6 (m, 2H), 8.2–8.4 (d, 1H), 8.6 (s, 1H), 10.8 (bs, 1H); ^{13}C NMR ($CDCl_3$) δ 21.68, 55.6, 111.3, 118.2, 119.6, 121.1, 125.7, 127.14, 127.64, 130.96, 132.56, 144.9, 149.06, 150.42, 157.25, 161.52. M/z (obs., $[m+H]^+$) = 268.

[00230] 4-Chloro-2-(2-methoxy-phenyl)-7-methyl-quinazoline


[00231]



[00232] To a mechanically stirred suspension of 2-(2-methoxyphenyl)-7-methyl-3*H*-quinazolin-4-one (100 g, 0.37 mol) in 1 L toluene was added diisopropyl ethylamine (100 mL), followed by phosphorus oxychloride (69 g, 0.45 mol). The reaction was then heated to 80 °C for 4 h. The reaction mixture was distilled under reduced pressure to remove toluene, and the resulting residue was dissolved in 2.2 L CH_2Cl_2. Ice water was added, and the pH was adjusted to 8–9 with saturated aqueous sodium bicarbonate solution while maintaining the temperature below 20 °C. The resulting organic layer was separated and the aqueous solution extracted with CH_2Cl_2, then the combined the organic layers were dried over sodium sulfate and distilled under reduced pressure. The crude product was dissolved 2:1 CH_2Cl_2/hexane, and the solution was passed through silica gel (2.5 kg, 60–120 mesh), followed by washing the silica bed with 2:1 CH_2Cl_2/hexane until the product eluted. The pure fractions were collected and combined, and the solvent was removed under reduced pressure. Hexane (500 mL) was added, and the mixture was stirred and filtered to give 4-chloro-2-(2-methoxy-phenyl)-7-methyl-quinazoline as a white to off-white solid (77 g, 72%). mp 161–164 °C. ^{1}H NMR ($CDCl_3$) δ 2.6 (s, 3H), 3.9 (s, 3H), 6.9–7.2 (m, 2H), 7.4–7.6 (m, 2H), 7.7–8 (d, 2H), 8.2 (d,1H); ^{13}C NMR ($CDCl_3$) δ 22.23, 56.06, 112.2, 120.26, 120.69, 125.34, 127.94, 130.45, 131.08, 131.08. M/z (obs., $[m+H]^+$) = 285.


[00233] **2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol**

[00234]



[00235] A stirring solution of 4-chloro-2-(2-methoxyphenyl)-7-methylquinazoline (91 g, 320 mmol) and CH_2Cl_2 (2.0 L) under an N_2 atmosphere was cooled to –30 °C. Boron tribromide (957 mL, 957 mmol, 1.0 M in CH_2Cl_2) was added dropwise over a period of 30 minutes at –30 to –40 °C. The cooling bath was removed, and the mixture was allowed to warm to 25 °C. The mixture was carefully poured into a stirring solution of saturated aqueous $NaHCO_3$ (4.0 L). The organic portion was separated, dried over $MgSO_4$, and evaporated to dryness. The resulting solid was suspended in CH_2Cl_2 (400 mL) under an N_2 atmosphere, followed by the addition of triethylamine (64.8 g, 640 mmol). The solution was cooled to –10 °C. A solution of piperazine (55.0 g, 640 mmol) in CH_2Cl_2 (400 mL) was added in a single portion, and the solution was stirred at ambient temperature for 1 hour. The solution temperature rose to 23 °C upon addition of the piperazine. The solution was partitioned between CH_2Cl_2 and H_2O. The organic portion was dried over $MgSO_4$ and evaporated to dryness. The residue was purified via silica gel chromatography using 5% MeOH in CH_2Cl_2 to obtain a tan solid. The resulting solid was triturated with 1:1 Et_2O/hexanes to obtain a yellow solid which was vacuum dried to give 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol as a light yellow solid (93.0 g, 290 mmol, 91%). LC/MS: *m/z* 321.1 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00236] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one**



[00237] Method A


[00238] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol) was placed in a tube charged with a stir bar followed by (*R*)-2-hydroxy-4,4-dimethylpentanoic acid in 1 ml of DMF and triethylamine (31.57 mg, 0.312 mmol), and the reaction was cooled to 0 °C. HATU (71 mg, 0.187 mmol) was then added, and the reaction was allowed to stir at 0 °C for 10 minutes and then allowed to warm to room temperature. The reaction was complete after 40 minutes, filtered, and purified by reverse phase HPLC to give the TFA salt of (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.75 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00239] Method B

[00240] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.74 mmol) was suspended in anhydrous DMF (5 mL) and cooled to 0 °C internal temperature. Under an N_2 atmosphere, (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (125.4 mg, 0.858 mmol) was added followed by triethylamine (0.218 mL, 1.56 mmol). To this stirring solution was added HATU (356 mg, 0.936 mmol). After the complete addition of HATU, the mixture was allowed to warm to 10 °C. After 45 min the reaction was complete, and it was quenched with an equal portion of ice water. A yellow precipitate formed which was collected by vacuum filtration and dissolved in CH_2Cl_2. This solution was desiccated with Na_2SO_4, filtered, and concentrated to give a viscous yellow-orange oil. The crude material was purified via silica gel chromatography using 88% CH_2Cl_2–hexanes (1:1) and 12% EtOAc to afford (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one as a faint yellow foam (265 mg, 76%). LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.75 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.46 (dd, J = 8.2, 1.8 Hz, 1H), 8.02 (d, J = 8.5 Hz, 1H), 7.70 (s, 1H), 7.41–7.37 (m, 2H), 6.97–6.93 (m, 2H), 4.89 (d, J = 7.2 Hz, 1H), 4.49–

4.44 (m, 1H), 4.06–3.67 (m,8), 2.52 (s, 3H), 1.56 (dd, J = 14.3, 3.0 Hz, 1H), 1.42 (dd, J = 14.3, 8.8 Hz, 1H), 0.97 (s, 9H)

[00241] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one hydrochloride




[00242] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one (265 mg, 0.367 mmol) was dissolved in anhydrous CH_2Cl_2 (3 mL) followed by the addition of Et_2O (6 mL) under an N_2 atmosphere. A 2.0 M HCl solution in Et_2O (0.296 mL, 0.591 mmol) was added over a 1 minute period. The reaction solution changed from a clear yellow solution to a creamy off white slurry. After complete addition of the HCl solution, the reaction was allowed to stir for an additional 10 minutes. The product was collected by vacuum filtration, washed with 3 mL of Et_2O and dried under vacuum to obtain (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4,4-dimethylpentan-1-one hydrochloride as a white solid (261 mg, 91%). LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.33 (d, J = 7.6 Hz, 1H), 8.07 (d, J = 8.6 Hz, 1H), 7.75 (s, 1H), 7.47–7.43 (m, 2H), 7.04–6.98 (m, 2H), 4.47–4.44 (m, 1H), 4.13–4.04 (m, 4H), 3.91–3.68 (m, 4H), 2.54 (s, 3H), 1.57 (dd, J = 14.3, 3.1 Hz, 1H), 1.42 (dd, J = 14.3, 8.8 Hz, 1H), 0.97 (s, 9H).

[00243] Example 102


[00244] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one

[00245]



[00246] **2-(2-Methoxyphenyl)quinazolin-4(3*H*)-one**

[00247] Method A



[00248] To a cooled (0–5 °C) mixture of anthranilamide (350 g, 2.57 mol) and triethylamine (286 g, 2.83 mol) in THF (2.5 L) was added dropwise *o*-anisoyl chloride (437 g, 2.57 mol) while maintaining the temperature between 0–20 °C. The resulting suspension was stirred at room temperature overnight. The solvent was evaporated *in vacuo*, and the residue was washed several times with water. The wet residue was suspended in 2 M aq. NaOH (13 L), and the mixture was heated to reflux. After 20 minutes a clear solution was obtained. After 1 h of reflux the clear solution was cooled in an ice water bath and then acidified to pH 6 with conc. aq. HCl. The suspension was filtered, and the residue was washed thoroughly with water. The white solid was dried by azeotropic distillation with toluene. 2-(2-Methoxyphenyl)quinazolin-4(3*H*)-one (567 g) was obtained in 82%. ^{1}H-NMR (200 MHz, Me_2SO-d_6): δ 3.90 (s, 3 H), 7.20 (m, 2 H), 7.60 (t, 2 H), 7.85 (m, 3 H), 8.20 (d, 1 H), 12.20 (s, 1 H).


[00249] Method B

[00250] In a 2 L three-necked round-bottomed flask equipped with an overhead stirrer and reflux condenser, anthranilamide (20.0 g, 147 mmol) and potassium carbonate (28.4 g, 206 mmol) were suspended in 1 L dry ether and heated to reflux. *o*-Anisoyl chloride (32.5 g, 191 mmol) was added slowly to the refluxing mixture. After 3 hours at reflux, the reaction mixture was allowed to cool to room temperature, the ether was removed under reduced pressure, and the

resulting residue was filtered and washed with water. The resulting solid was then suspended in 600 mL of 5% aq. NaOH solution and boiled for one hour. The reaction was allowed to cool to room temperature, and it was neutralized with acetic acid, upon which 2-(2-methoxyphenyl)quinazolin-4(3*H*)-one was precipitated. The product was collected by filtration, washed with water, and dried overnight *in vacuo* to yield 27 g (73%) of pure 2-(2-methoxyphenyl)quinazolin-4(3*H*)-one. LC/MS: *m/z* 253.0 $(M+H)^+$ at 3.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (DMSO) δ 3.86 (s, 3H), δ 7.09 (t, 1H), δ 7.18 (d, 1H), δ 7.53 (m, 2H), δ 7.70 (m, 2H), δ 7.80 (m, 1H), δ 8.14 (d, 1H), δ 12.11 (s, 1H); ^{13}C NMR (DMSO) δ 55.75, δ 111.86, δ 120.89, δ 120.97, δ 122.74, δ 125.75, δ 126.45, δ 127.26, δ 130.41, δ 132.13, δ 134.32, δ 148.97, δ 152.48, δ 157.12, δ 161.35

[00251] 4-Chloro-3,4-dihydro-2-(2-methoxyphenyl)quinazoline


[00252]



[00253] A suspension of 2-(2-methoxyphenyl)quinazolin-4(3*H*)-one (567 g, 2.1 mol) in phosphoryl chloride (2 L, 21 mol) and a catalytic amount of *N,N*-dimethyl aniline was brought to reflux. The reaction started immediately with the evolution of gas (HCl) upon the addition of *N,N*-dimethyl aniline. After the production of gas had ceased the mixture was cooled to room temperature. The excess $POCl_3$ was evaporated. The resulting dark solution was cooled to room temperature and slowly poured on ice and water, while maintaining the temperature below 5 °C. The cold suspension was extracted with dichloromethane. The extract was dried over sodium sulfate and filtered, and the solvent was removed *in vacuo*. The crude material was purified by column chromatography (silica gel, CH_2Cl_2). Yield: 189 g (33%) of 4-chloro-3,4-dihydro-2-(2-methoxyphenyl)quinazoline. ^1H-NMR (300 MHz, Me_2SO-d_6): δ 3.85 (s, 3 H), 7.15 (t, 1 H), 7.25 (d, 1 H), 7.60 (t, 2 H), 7.70 (d, 1 H), 7.8 (d, 1 H), 7.9 (t, 1 H), 8.2 (d, 1 H).

[00254] 2-(4-Chloroquinazolin-2-yl)phenol


[00255]



[00256] To a solution of 4-chloro-2-(2-methoxyphenyl)quinazoline (1.0 g, 3.7 mmol) in 40 mL CH_2Cl_2 at –78 °C was added 5 equivalents of 1 M BBr_3 dropwise. After complete addition the cooling bath was removed and the reaction was quenched with $NaHCO_3$ after 90 minutes. The product was extracted twice with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 60:40 CH_2Cl_2: hexanes gave 2-(4-chloroquinazolin-2-yl)phenol (700 mg, 74%). LC/MS: *m/z* 257.1 $(M+H)^+$ at 3.75 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.44 (m, 1H), 8.24 (m, 3H), 7.89 (m, 1H), 7.49 (m, 1H), 7.05 (m, 2H).


[00257] **2-(4-(Piperazin-1-yl)quinazolin-2-yl)phenol**

[00258]



[00259] To a solution of 2-(4-chloroquinazolin-2-yl)phenol (2.0 g, 7.8 mmol) in CH_2Cl_2 at 0 °C was rapidly added a solution of piperazine (2.01 g, 23.4 mmol) and triethylamine (2.17 mL, 15.6 mmol) in 10 mL CH_2Cl_2. The reaction was warmed to room temperature and stirred for 5 hours. The reaction was quenched with 25 mL of water and extracted with (3×15) mL of CH_2Cl_2. The organic layer was dried over Na_2SO_4, filtered, and concentrated to afford 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (1.98 g, 83%). LC/MS: *m/z* 307.3 $(M+H)^+$ at 1.47 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43 (d, J = 10.1 Hz, 1H), 8.04 (d, J = 8.3 Hz, 1H), 7.85 (m, 2H), 7.53 (m, 1H), 7.38 (m, 1H), 6.95 (m, 2H), 3.85 (t, J = 4.8 Hz, 4H), 2.94 (t, J = 4.8 Hz, 4H).

[00260] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**




[00261]

[00262] To a solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.82 mmol) in CH_2Cl_2 (6 mL) was added triethylamine (227 μL, 1.63 mmol) followed by the addition of (*R*)-2-hydroxy-4-methylpentanoic acid (140 mg, 1.06 mmol) and HATU (403 mg, 1.06 mmol). The reaction mixture was stirred at room temperature for 3 h and then quenched with H_2O. The aqueous layer was extracted twice with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2:hexanes gave (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one (265 mg, 77%). LC/MS: *m/z* 421.30 $(M+H)^+$ at 2.57 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). H NMR (400 MHz, DMSO-d6) δ 8.47 (dd, *J* = 8.3, 1.7 Hz, 1H), 8.12 (d, *J* = 8.3 Hz, 1H) 7.88 (m, 2H), 7.57 (m, 1H), 7.40 (m, 1H), 6.96 (m, 2H), 4.92 (d, *J* = 7.2 Hz, 1H), 4.39 (m, 1H), 3.95 (m, 4H), 3.76 (m, 4H), 1.80 (m, 1H), 1.43 (m, 2H), 0.92 (q, *J* = 3.8 Hz, 6H).

[00263] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride

[00264]




[00265] To a solution of (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one (265 mg, 0.63 mmol) in CH_2Cl_2 (3 mL) under inert

atmosphere was added 10 mL of ether followed by the dropwise addition of 2 M HCl (0.315 mL, 0.63 mmol).The reaction was stirred for 30 min before the formed precipitate was filtered to afford (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride (261 mg, 91%). LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.60 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.28 (dd, *J* = 7.9, 1.6 Hz, 1H), 8.19 (d, *J* = 8.4 Hz, 1H), 7.96 (d, *J* = 3.9 Hz, 2H), 7.65 (m, 1H), 7.48 (m, 1H), 7.04 (m, 2H), 4.37 (m, 1H), 4.10 (m, 4H), 3.80 (m, 4H), 1.77 (m, 1H), 1.41 (m, 2H), 0.91 (dd, *J* = 6.6, 3.1 Hz, 6H).

[00266] Example 103


[00267] **(Benzo[d][1,3]dioxol-7-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00268]



[00269] **(Benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate**


[00270]



[00271] A solution of (benzo[d][1,3]dioxol-7-yl)methanol (2 g, 13.14 mmol) and di(1*H*-imidazol-1-yl)methanone (4.26 g, 26.28 mmol) in 20 mL CH_2Cl_2 was heated overnight at 50 °C. The reaction was quenched with water, extracted with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 10–70% EtOAc in CH_2Cl_2 gave (benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate (2.8 g, 86%).

[00272] **(Benzo[d][1,3]dioxol-7-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00273]




[00274] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol) , (benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate (78 mg, 0.32 mmol) and triethylamine (44.6 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (benzo[d][1,3]dioxol-7-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 499.3 $(M+H)^+$ at 2.97 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00275] Example 104


[00276] **(*R*)-3-Hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-N,N-dimethyl-4-oxobutanamide**

[00277]



[00278] **(*R*)-3-(Methoxycarbonyl)-2-hydroxypropanoic acid**


[00279]



[00280] Methyl 2-((*R*)-2,2-dimethyl-5-oxo-1,3-dioxolan-4-yl)acetate (17.1 g, 90.9 mmol) was stirred in a 1:1 mixture of THF:1 M HCl (200 mL) for 1 h at room temperature. After addition of NaCl to nearly saturate the aqueous layer, the mixture was extracted with EtOAc, and the extracts were dried over Na_2SO_4 and concentrated to obtain (*R*)-3-(methoxycarbonyl)-2-hydroxypropanoic acid as an oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 4.58–4.55 (m, 1H), 3.75 (s, 3H), 2.98–2.84 (m, 2H).


[00281] 4-((*R*)-2-Hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester

[00282]



[00283] EDCI (3.6 g, 19 mmol) was added to a solution of (*R*)-3-(methoxycarbonyl)-2-hydroxypropanoic acid (2.8 g, 19 mmol) and HOBt (2.6 g, 19 mmol) in DMF (200 mL). After stirring this mixture for 5 min, benzyl piperazine-1-carboxylate (4.2 g, 3.6 mL, 19 mmol) and triethylamine (2.6 mL, 19 mmol) were added and stirred for 3 days at room temperature. The reaction mixture was poured into water and extracted with EtOAc. After washing the organic layers with brine and water, drying over Na_2SO_4 and concentrating, purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 gave 4-((*R*)-2-hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester as an oil (2.19 g, 33%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.40–7.31 (m, 5H), 5.15 (s, 2H), 4.79–4.74 (m, 1H), 3.95 (d, J = 8.0 Hz, 1H), 3.79–3.74 (m, 1H), 3.74 (s, 3H), 3.71–3.44 (m, 7H), 2.62 (d, J = 5.8 Hz, 2H).


[00284] (*R*)-Methyl 3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoate



[00286] 4-((*R*)-2-Hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester (0.52 g, 1.5 mmol) and MeOH (15 mL) were stirred with 10% Pd/C under an H_2 atmosphere at ambient pressure overnight. After filtration and evaporation of the solvent the residue was taken up in CH_2Cl_2, and 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.40 g, 1.50 mmol) plus triethylamine (0.41 mL, 3.00 mmol) were added. The reaction mixture was stirred at room temperature overnight, washed with water, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% MeOH/ CH_2Cl_2 provided (*R*)-methyl 3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoate (0.38 g, 57%). LC/MS: *m/z* 451.1 $(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00287] **(*R*)-3-Hydroxy-4-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-butyric acid**


[00288]



[00289] $LiOH \cdot H_2O$ (19.8 mg, 0.47 mmol) was added to a solution of (*R*)-3-hydroxy-4-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-butyric acid methyl ester (71 mg, 0.16 mmol) in 2 mL THF:H_2O (1:1) and stirred at room temperature for 3 h. The reaction mixture was acidified with 1 M HCl and then extracted with EtOAc. After drying the organic layer over Na_2SO_4, it was concentrated and then purified via silica gel chromatography using 0–15% MeOH/CH_2Cl_2 to provide (*R*)-3-hydroxy-4-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-butyric acid (52 mg, 75%). LC/MS: *m/z* 437.3 $(M+H)^+$ at 2.04 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00290] **(*R*)-3-Hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-*N*,*N*-dimethyl-4-oxobutanamide**

[00291]




[00292] (*R*)-3-Hydroxy-4-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-butyric acid (17 mg, 0.039 mmol) and HATU (16 mg, 0.043 mmol) were stirred in DMF (0.5 mL). After adding dimethylamine (2 M in THF, 0.10 mL, 0.19 mmol), the reaction mixture was stirred at room temperature for 5 h. Purification via preparative reverse-phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-N,N-dimethyl-4-oxobutanamide as the TFA salt. LC/MS: *m/z* 436.3 $(M+H)^+$ at 1.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00293] Example 105


[00294] **(2-(Tetrahydro-2*H*-pyran-4-yl)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00295]



[00296] **(2-(Tetrahydro-2*H*-pyran-4-yl)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00297]




[00298] To a solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) was added 2-(tetrahydro-2*H*-pyran-4-yl)acetic acid (13.5 mg, 0.09 mmol) followed by the addition of triethylamine (25 μL), then HATU (44 mg) at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2-(tetrahydro-2*H*-pyran-4-yl)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone as the TFA salt. LC/MS: *m/z* 447.10 $(M+H)^+$ at 2.32 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00299] Example 106

[00300] **(*R*)-2-Hydroxy-1-((*R*)-2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00301]



[00302] **(*R*)-*tert*-Butyl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**




[00303]

[00304] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (450 mg, 1.66 mmol) in 10 mL DMF was added a solution of (*R*)-*tert*-butyl 2-((benzyloxy) methyl) piperazine-1-carboxylate (610 mg, 1.99 mmol) in DMF and triethylamine (0.46 mL). The reaction mixture was then refluxed at 85 °C for 30 minutes, quenched with water, extracted twice with CH_2Cl_2, dried over Na_2SO_4, and concentrated to give (*R*)-*tert*-butyl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate (760 mg, 85%). This material was used in the next step without further purification. LC/MS: *m/z* 541.5 $(M+H)^+$ at 3.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00305] **2-(4-((*R*)-3-((Benzyloxy)methyl)piperazin-1-yl)-7-methylquinazolin-2-yl)phenol**


[00306]



[00307] To a solution of (*R*)-*tert*-butyl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate (760 mg, 1.72 mmol) in 15 mL CH_2Cl_2 was added 10 mL of TFA. The reaction was stirred for 1 hour. TFA was removed under vacuum and the reaction was neutralized using a 1 M NaOH solution. The aqueous layer was extracted twice with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated to obtain 2-(4-((*R*)-3-((benzyloxy)methyl)piperazin-1-yl)-7-methylquinazolin-2-yl)phenol (570 mg, 92%). This material was used in the next step without further purification. LC/MS: *m/z* 441.5 $(M+H)^+$ at 2.44 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00308] **(*R*)-1-((*R*)-2-((Benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**


[00309]



[00310] To a solution of 2-(4-((*R*)-3-((benzyloxy)methyl)piperazin-1-yl)-7-methylquinazolin-2-yl)phenol (100 mg, 0.22 mmol) in DMF (1 mL) was added (*R*)-2-hydroxy-4-methylpentanoic acid (30 mg, 0.22 mmol) followed by the addition of triethylamine (61 μL), then HATU (109 mg) at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-1-((*R*)-2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 555.7 $(M+H)^+$ at 3.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00311] **(*R*)-2-Hydroxy-1-((*R*)-2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00312]




[00313] To a solution of (*R*)-1-((*R*)-2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one trifluoroacetate (29.6 mg, 0.053 mmol) in ethanol was added $Pd(OH)_2$ (188 mg), and the reaction was heated at 50 °C

under H_2 atmosphere at ambient pressure. The reaction was filtered, and purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-2-hydroxy-1-((*R*)-2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 465.50 $(M+H)^+$ at 2.47 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00314] Example 107


[00315] **(Pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00316]



[00317] **(Pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate**


[00318]



[00319] A solution of (pyridin-3-yl)methanol (2 g, 18.32 mmol) and di(1*H*-imidazol-1-yl)methanone (5.94 g, 36.65 mmol) in 20 mL CH_2Cl_2 was heated overnight at 50 °C. The reaction was quenched with water, extracted twice with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 10–70% EtOAc in CH_2Cl_2 gave (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (3.1 g, 84%). LC/MS: *m/z* 204.1 $(M+H)^+$ at 0.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, $CDCl_3$) δ 8.74 (d, J = 1.9 Hz, 1H), 8.68 (dd, J = 4.8, 1.4 Hz, 1H), 8.16 (s, 1H), 7.81 (m, 1H), 7.44 (s, 1H), 7.38 (m, 1H), 7.09 (s, 1H), 5.46 (s, 2H).

[00320] **(Pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00321]




[00322] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol), (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (67 mg, 0.32 mmol) and triethylamine (44.6 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 442.50 $(M+H)^+$ at 1.97 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00323] Example 108


[00324] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylpropan-1-one**

[00325]



[00326] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylpropan-1-one**

[00327]




[00328] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-hydroxy-2-methylpropanoic acid (29.6 mg, 0.284 mmol). It was followed by the addition of triethylamine (61 μL), then a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylpropan-1-one as the TFA salt. LC/MS: *m/z* 407.50 $(M+H)^+$ at 2.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00329] Example 109


[00330] **(*S*)-3-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00331]



[00332] **(*S*)-3-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00333]




[00334] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to (*S*)-3-hydroxybutanoic acid (31.0 mg, 0.297 mmol). It was followed by the addition of triethylamine (63 μL), then a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-3-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 393.1 $(M+H)^+$ at 2.04 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00335] Example 110


[00336] **2-(Trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[00337]



[00338] **2-(Trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[00339]




[00340] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-(trifluoromethyl)-2-hydroxypropanoic acid (45 mg, 0.284 mmol). It was followed by the addition of triethylamine (61 μL), then a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-(trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 461.1 $(M+H)^+$ at 2.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00341] Example 111

[00342] **(*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one**

[00343]



[00344] **(*E*)-*N*-(4-Fluorophenyl)-2-(hydroxyimino)acetamide**




[00345]

[00346] 4-Fluoroaniline (58.2 g, 0.50 mol) was added slowly to 10% aqueous HCl solution. This suspension was added to a mixture of chloral hydrate (95 g, 0.55 mol) and sodium sulfate (0.5 kg) in 750 mL water with mechanical stirring. Hydroxylamine hydrochloride (116 g, 1.63 mol) dissolved in water (250 mL) was added, and the resulting slurry was heated at 100 °C. After this temperature was reached, the heating mantle was removed immediately, and the solution was cooled to room temperature. The formed precipitate was collected by filtration, washed with water (2x300 mL), and dried in a vacuum oven at 60 °C. Yield: 78.2 g of *N*-(4-fluorophenyl)-2-hydroxyiminoacetamide as an off-white solid.


[00347] **5-Fluoroindoline-2,3-dione**

[00348]



[00349] Concentrated sulfuric acid (200 mL) was heated at 50 °C, and *N*-(4-fluorophenyl)-2-hydroxyiminoacetamide was slowly added. The black solution was carefully heated at 90 °C. At this temperature, some slight cooling was necessary to keep the temperature at 90 °C. When no more heat had developed, the reaction mixture was heated at 90 °C for an additional half hour. The dark-red solution was cooled to room temperature and poured onto 3 L ice water and 1 L ethyl acetate with vigorous stirring. The layers were separated, and the aqueous layer was extracted with ethyl acetate (1x1 L, 1x0.5 L). The combined organic extracts were dried over sodium sulfate, filtered, and evaporated to dryness. Yield: 35.3 g (52%) of a dark red solid, 5-fluoro-1*H*-indole-2,3-dione.

[00350] **2-Amino-5-fluorobenzamide**




[00351]

[00352] 5-Fluoro-1*H*-indole-2,3-dione (35.3 g, 213 mmol) was heated in acetic acid (300 mL), 1 mL concentrated sulfuric acid, and 22 mL 35% aq. hydrogen peroxide at 70 °C. The solution was kept at that temperature one and a half hours during which time a solid formed in the reaction mixture. After cooling to room temperature this solid was collected by filtration and was washed three times with water. The wet solid was suspended in 150 mL water, and 40 mL of a 25% aq. ammonia solution was added. This mixture was stirred at room temperature 3 days. The formed solid was collected by filtration and was washed twice with water. The solid was dried by azeotropic distillation with toluene (3 x 100 mL) to yield 2-amino-5-fluorobenzamide (9.5 g). The combined filtrates were extracted with ethyl acetate (2x100 mL). The combined extracts were dried over sodium sulfate, filtered, and evaporated to dryness to yield 2-amino-5-fluorobenzamide (3.5 g) as an off-white solid. Both fractions were combined for use in the next reaction step.

[00353] **6-Fluoro-2-(2-methoxyphenyl)-3*H*-quinazolin-4-one**


[00354]



[00355] *o*-Anisoyl chloride (15.7 g, 92 mmol) was added dropwise to a solution of aminobenzamide 2-amino-5-fluorobenzamide (13.0 g, 84 mmol) and triethylamine (16 mL, 110 mmol) in tetrahydrofuran (100 mL) cooled in an ice bath. Immediately a precipitate started forming. Stirring of the solution was continued for 5 hours at room temperature. The formed precipitate was collected by filtration and was washed twice with diethyl ether and dried at 50 °C *in vacuo*. The dried solid was suspended in 2 N aqueous sodium hydroxide solution (250 mL) and heated at reflux until a clear solution was obtained (3 hours). The reaction mixture was cooled to room temperature and filtered. The filtrate was acidified to pH<1 with concentrated aqueous HCl. The formed precipitate was collected by filtration and washed twice with water,

twice with methanol, and twice with diethyl ether. The solid was dried in an oven at 45 °C to yield 6-fluoro-2-(2-methoxyphenyl)-3*H*-quinazolin-4-one (18.2 g, 80%) as a white solid.


[00356] 4-Chloro-6-fluoro-2-(2-methoxyphenyl)quinazoline



[00357] A suspension of 6-fluoro-2-(2-methoxyphenyl)-3*H*-quinazolin-4-one (14.0 g, 52 mmol), *N,N*-dimethylaniline (6.6 mL, 52 mmol), and phosphorus oxychloride (4.8 mL, 52 mmol) in benzene (100 mL) was heated at reflux until a clear, dark solution was obtained (1 hour). The reaction mixture was cooled to room temperature, and the volume was reduced under reduced pressure. The black, oily residue was poured into 300 g of ice. Dichloromethane (600 mL) was added with vigorous stirring, and the temperature was kept below 5 °C at all times. The pH was monitored, and aqueous 1 N sodium hydroxide was added until the pH was 10–11. The mixture was stirred for one hour at a temperature below 5 °C, and the pH was kept between 10–11 by addition of 1 N aqueous sodium hydroxide. The layers were separated, and the organic layer was washed with ice cold 1 N aqueous sodium hydroxide (2x200 mL). Heptanes (300 mL) were added to the organic layer. This mixture was filtered through a short plug of silica gel and eluted with dichloromethane/heptanes (2:1). All fractions containing product were combined and evaporated to dryness. The residue was triturated with heptanes to yield 4-chloro-6-fluoro-2-(2-methoxyphenyl)-quinazoline (11.5 g, 76%) as a white solid.

[00358] 2-(4-Chloro-6-fluoroquinazolin-2-yl)phenol

[00359]




[00360] A solution of 4-chloro-6-fluoro-2-(2-methoxyphenyl)quinazoline (3.0 g, 10.3 mmol) in CH_2Cl_2 (15 mL) was cooled to –78 °C. Then, 1 M BBr_3 (51.95 mL, 59.95 mmol) was added dropwise. The reaction was warmed to room temperature and was quenched with $NaHCO_3$ and extracted twice with CH_2Cl_2. The organic layer was dried over $MgSO_4$, filtered,

and concentrated. Purification via silica gel chromatography using 5–20% CH_2Cl_2 in hexanes gave 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (1.61 g, 57%). LC/MS: *m/z* 275.1 $(M+H)^+$ at 3.8 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00361] 2-(6-Fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol

[00362]



[00363] To a stirring solution of 2-(6-fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (500 mg, 1.82 mmol) in CH_2Cl_2 (20 mL) at 0 °C, under an N_2 atmosphere was rapidly added a solution of piperazine (0.263 g, 7.28 mmol) and triethylamine (0.35 mL, 2.55 mmol) in CH_2Cl_2. The mixture was stirred for 1 h and then quenched with H_2O, extracted twice with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–5% MeOH in CH_2Cl_2 gave 2-(6-fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (400 mg, 68%). LC/MS: *m/z* 325.5 $(M+H)^+$ at 2.12 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.42 (m, 1H), 7.97 (m, 1H), 7.78 (m, 2H), 7.39 (m, 1H), 6.95 (m, 2H), 3.83 (t, $J = 4.9$ Hz, 4H), 2.92 (t, $J = 4.9$ Hz, 4H).

[00364] (*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one



[00365]


[00366] Method A

[00367] To a solution of 2-(6-fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.08 mmol) in DMF (1 mL) was added (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (20.28 mg, 0.14 mmol) followed by the addition of triethylamine (25 μL), then HATU (44 mg) at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))–H_2O) gave (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one as the TFA salt. LC/MS: *m/z* 453.52 $(M+H)^+$ at 3.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00368] Method B

[00369] To a solution of 2-(6-fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (200 mg, 0.61 mmol) in CH_2Cl_2 (5 mL) was added triethylamine (170 μL, 1.22 mmol) followed by the addition of (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (116 mg, 0.79 mmol), then HATU (301 mg, 0.79 mmol). An additional 3 mL of CH_2Cl_2 were added, and the reaction was stirred for 3 h. After quenching with water, the mixture was extracted twice with CH_2Cl_2. The organic phase was dried over $MgSO_4$ and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2:hexanes yielded (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one (240 mg, 86%). *m/z*: M+1 *obs* = 453.5; t_R = 3.19 minutes ^1H NMR (400 MHz, DMSO–d6) δ 8.46 (m, 1H), 8.01 (m, 1H), 7.84 (m, 2H), 7.40 (m, 1H), 6.96 (m, 2H), 4.89 (d, *J* = 7.1 Hz, 1H), 4.45 (m, 1H), 3.97 (m, 4H), 3.76 (m, 4H), 1.56 (m, 1H), 1.42 (m, 1H), 0.97 (s, 9H).

[00370] **(*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one hydrochloride**



[00371]


[00372] To a solution of (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2- hydroxy-4,4-dimethylpentan-1-one (230 mg, 0.51 mmol) in CH_2Cl_2 (3 mL)

under an inert atmosphere was added dropwise a 2 M HCl solution in ether (2.55 mL, 0.51 mmol). To it was then added ether (15 mL) which resulted in formation of a precipitate that was allowed to stir for an hour. The product was collected by vacuum filtration and dried to afford (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one hydrochloride (230 mg, 92%). LCMS: *m/z* 453.5 $(M+H)^+$ at 3.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.40 (dd, *J* = 7.8, 1.4 Hz, 1H), 8.04 (m, 1H), 7.88 (m, 2H), 7.43 (m, 1H), 6.99 (m, 2H), 4.45 (dd, *J* = 8.8, 3.0 Hz, 1H), 4.02 (m, 4H), 3.79 (m, 4H), 1.56 (m, 1H), 1.42 (m, 1H), 0.97 (s, 9H).

[00373] Example 112


[00374] **4,4,4-Trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00375]



[00376] **4,4,4-Trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**


[00377]



[00378] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (87 mg, 0.27 mmol), 4,4,4-trifluoro-2-hydroxybutanoic acid (43 mg, 0.27 mmol), HATU (0.12 g, 0.33 mmol) and triethylamine (45 μL, 0.33 mmol) were stirred in DMF (3 mL) at room temperature overnight. The mixture was diluted with water and extracted with EtOAc. The combined organic layers were washed with brine and water, dried over Na_2SO_4 and concentrated. Purification via silica gel chromatography using 0–20% EtOAc in 1:1 CH_2Cl_2: hexanes gave 4,4,4-trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one as an off-white solid (86 mg, 66%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.46 (dd, J = 8.0, 1.7 Hz, 1H), 7.77 (d, J = 8.5 Hz, 1H), 7.70 (s, 1H), 7.41–7.37 (m, 1H), 7.31 (dd, J = 8.5, 1.5 Hz, 1H), 7.05–7.03 (m, 1H), 6.97–6.93 (m, 1H), 4.80–4.75 (m, 1H), 4.07–3.68 (m, 9H), 2.56 (s, 3H), 2.50–2.39 (m, 2H); LC/MS: *m/z* 461.3 $(M+H)^+$ at 2.49 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00379] Example 113

[00380] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one**



[00381]

[00382] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one**



[00383]

[00384] Method A


[00385] Under an N_2 atmosphere, a mixture of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.08 mmol), 2-hydroxy-3-methylbutanoic acid (9 mg, 0.08 mmol), BOP (35 mg, 0.08 mmol), triethylamine (22 μL, 0.16 mmol) and DMF (0.2 mL) was stirred at room temperature for 1 hour. The reaction mixture was then purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one as the TFA salt. LC/MS: *m/z* 421.10 $(M+H)^+$ at 2.76 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00386] Method B

[00387] To a solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.78 mmol) in CH_2Cl_2 (6 mL) was added triethylamine (217 μL, 1.56 mmol) followed by the addition of 2-hydroxy-3-methylbutanoic acid (120 mg, 1.0 mmol) and HATU (380 mg, 1.00 mmol). The reaction mixture was stirred at room temperature for 3 h and then quenched with H_2O. The aqueous layer was extracted with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2:hexanes gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one (230 mg, 70%). LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.45 (m, 1H), 8.01 (d, J = 8.5 Hz, 1H), 7.70 (s, 1H), 7.39 (m, 2H), 6.95 (m, 2H), 4.79 (d, J = 7.2 Hz, 1H), 4.11 (dd, J = 7.0, 5.9 Hz, 1H), 3.87 (m, 8H), 2.52 (s, 3H), 1.91 (m, 1H), 0.88 (dd, J = 22.8, 6.7 Hz, 6H).

[00388] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one hydrochloride**

[00389]




[00390] To a solution of 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one (230 mg, 0.54 mmol) in CH_2Cl_2 (3 mL) under an inert atmosphere was added ether (12 mL) followed by the dropwise addition of 2 M HCl solution in ether (0.27 mL, 0.54 mmol) which resulted in the formation of a precipitate which was stirred for an hour and then collected by vacuum filtration and dried to afford 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylbutan-1-one hydrochloride (205 mg, 83%). LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.48 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.25 (d, J = 7.9 Hz, 1H), 8.08 (d, J = 8.5 Hz, 1H), 7.74 (s, 1H), 7.48 (t, J = 6.6 Hz, 2H), 7.04 (m, 2H), 4.10 (m, 5H), 3.79 (m, 4H), 2.53 (s, 3H), 1.90 (m, 1H), 0.87 (dd, J = 16.3, 6.7 Hz, 6H)

[00391] Example 114

[00392] **(Pyridin-4-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**


[00393]



[00394]

[00395] ***N*-(2-Cyano-5-methyl-phenyl)-2-fluoro-6-methoxy-benzamide**


[00396]



[00397] 6-Fluoro-2-anisoic acid (110 g, 0.70 mol) was added in portions over 15 minutes to a mixture of thionyl chloride (230 ml, 3.2 mol), toluene (200 mL), and DMF (1 mL). The resulting mixture was stirred overnight at room temperature. The solution was evaporated to dryness and added dropwise to an ice-bath cooled solution of anthranilonitrile (92.5 g, 0.70 mol) in pyridine (200 mL). The dropping funnel was rinsed with a minimal amount of acetonitrile. The resulting mixture was stirred overnight at room temperature under a nitrogen atmosphere and was subsequently poured into 2 L ice water. The resulting slurry was stirred vigorously for 1 hour. The formed solid was collected by filtration and was washed twice with water. The filter cake was dissolved in 2 L dichloromethane, and this solution was washed with 1 N aq. HCl (400 mL) and with saturated aq. NaCl (400 mL), dried over sodium sulfate, filtered, and evaporated to dryness to give *N*-(2-cyano-5-methylphenyl)-2-fluoro-6-methoxybenzamide (186 g, 93%) as a brownish solid. ^{1}H-NMR ($CDCl_3$, 200 MHz): δ 9.09 (s, 1H), 8.58 (s, 1H), 7.59–7.42 (m, 2H), 7.09–7.02 (m, 1H), 6.94–6.83 (m, 2H), 4.11 (s, 3H), 2.57 (s, 3H) ppm.

[00398] **2-(2-Fluoro-6-methoxy-phenyl)-7-methyl-3*H*-quinazolin-4-one**


[00399]



[00400] To a suspension of *N*-(2-cyano-5-methylphenyl)-2-fluoro-6-methoxybenzamide (31.5 g, 111 mmol) in ethanol (626 mL) was added 6 M aqueous NaOH solution (205 mL). After 10 minutes, 30% aqueous H_2O_2 (60 mL) was added, forming a slurry. The reaction was heated to reflux for 18 h and cooled to room temperature. NaOH (22.2 g, 0.56 mol) and 30% aqueous H_2O_2 (26 mL) were added, and the reaction was heated to reflux for six hours. The reaction cooled to room temperature, 30% aqueous H_2O_2 (45 mL) was added, and the reaction was heated to reflux for 18 h. The reaction was cooled to room temperature, NaOH (10 g, 0.25 mol) and 30% aqueous H_2O_2 (70 mL) were added, and the reaction was heated to reflux for six hours. The reaction was cooled to room temperature and poured over ice (800 mL). The pH was adjusted to 3–4 by addition of conc. HCl solution, and the precipitated off-white solid was filtered and washed with water (3 x 40 mL). The solid was dried under vacuum to provide 2-(2-fluoro-6-methoxy-phenyl)-7-methyl-3*H*-quinazolin-4-one (28 g, 89%).

[00401] **4-Chloro-2-(2-fluoro-6-methoxyphenyl)-7-methylquinazoline**


[00402]



[00403] Under an N_2 atmosphere, 2-(2-fluoro-6-methoxyphenyl)-7-methylquinazolin-4(3*H*)-one (20 g, 70.35 mmol) was suspended in benzene (300 mL), followed by the addition of *N*,*N*-dimethylaniline (26.8 mL, 211.05 mmol), then $POCl_3$ (13.11 mL, 140.7 mmol). The reaction was heated at reflux, and completion of product formation was observed after 1.5 h. After cooling to room temperature, the mixture was slowly poured over 1 liter of ice. The solution was then diluted with CH_2Cl_2, and the pH was adjusted to 7 using a saturated aqueous $NaHCO_3$ solution. The layers were partitioned, separated and extracted with CH_2Cl_2. All organic layers were combined, dried over Na_2SO_4, filtered, and concentrated to a dark oil. The crude material was purified by silica gel chromatography using 75% CH_2Cl_2/ 25% hexanes to obtain 4-chloro-2-(2-fluoro-6-methoxyphenyl)-7-methylquinazoline as a yellow solid (18.82 g, 88%). LC/MS: *m/z* 302.9 $(M+H)^+$ at 3.28 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, $CDCl_3$) δ 8.22 (d, J = 8.5 Hz, 1H), 7.95 (s, 1H), 7.60 (dd, J = 8.6, 1.5 Hz, 1H), 7.42–7.40 (m, 1H), 6.86–6.84 (m, 2H), 3.81 (s, 3H), 2.64 (s, 3H)

[00404] **2-(4-Chloro-7-methylquinazolin-2-yl)-3-fluorophenol**

[00405]




[00406] Under an N_2 atmosphere, 4-chloro-2-(2-fluoro-6-methoxyphenyl)-7-methylquinazolin (7.0 g, 23.12 mmol) was dissolved in CH_2Cl_2 (110 mL) and cooled to –50 °C internal temperature using a dry ice/acetone bath. A 1.0 M solution of BBr_3 in CH_2Cl_2 (115.6 mL, 115.6 mmol) was added dropwise via an addition funnel while maintaining the internal temperature at –50 °C. The reaction mixture was allowed to warm to 0 °C, and the reaction was complete after 1.5 h. It was then slowly quenched with saturated aqueous $NaHCO_3$ solution to

pH 7. After partitioning between CH_2Cl_2 and H_2O, the mixture was separated and the aqueous layer was twice extracted with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated to a brown solid. Purification via silica gel chromatography using 75% CH_2Cl_2/ 25% hexanes gave 2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol as a yellow solid (4.37 g, 66%). LC/MS: *m/z* 289.1 $(M+H)^+$ at 3.71 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, $CDCl_3$) δ 8.22 (d, J = 8.5 Hz, 1H), 7.95 (s, 1H), 7.60 (dd, J = 8.6, 1.5 Hz, 1H), 7.42–7.36 (m, 1H), 6.86–6.82 (m, 2H), 3.81 (s, 3H), 2.64 (s, 3H)


[00407] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol

[00408]



[00409] 2-(4-Chloro-7-methylquinazolin-2-yl)-3-fluorophenol (4.37 g, 15.14 mmol) was suspended in CH_2Cl_2 (65 mL) under an N_2 atmosphere and placed into an ice water bath. To this solution was added a solution of piperazine (4.00 g, 45.42 mmol) and triethylamine (4.2 mL, 30.28 mmol) in CH_2Cl_2 (15 mL) in one portion. After stirring the reaction for 30 minutes, it was partitioned between CH_2Cl_2 and H_2O and separated, and the aqueous layer was extracted twice more with CH_2Cl_2. The organic phase was dried over Na_2SO_4, filtered, and concentrated to a bright yellow solid which was purified via silica gel chromatography using a 95%/5% mixture of CH_2Cl_2/MeOH to afford 3-fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (4.32 g, 85%) as a bright yellow solid. LC/MS: *m/z* 339.3 $(M+H)^+$ at 1.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 7.92 (d, J = 8.5 Hz, 1H), 7.63 (s, 1H), 7.39–7.30 (m, 2H), 6.78 (d, J = 8.3 Hz, 1H), 6.74–6.69 (m, 1H), 3.83 (t, J = 4.8 Hz, 4H), 2.88 (t, J = 4.9 Hz, 4H).


[00410] (Pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate



[00411]

[00412] A solution of (pyridin-4-yl)methanol (2.0 g, 18.3 mmol) and di(1*H*-imidazol-1-yl)methanone (5.94 g, 36.6 mmol) in 20 mL CH_2Cl_2 was heated overnight at 50 °C The reaction was quenched with water, extracted with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 10–70% EtOAc in CH_2Cl_2 gave (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (3 g, 81%). LC/MS: *m/z* 204.3 $(M+H)^+$ at 0.38 min (v). ^1H NMR (400 MHz, $CDCl_3$) δ 8.69 (dd, J = 4.4, 1.6 Hz, 2H), 8.20 (t, J = 0.9 Hz, 1H), 7.47 (t, J = 1.5 Hz, 1H), 7.34 (dd, J = 4.4, 1.6 Hz, 2H), 7.11 (dd, J = 1.6, 0.8 Hz, 1H), 5.45 (s, 2H).

[00413] (Pyridin-4-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate




[00414]

[00415] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.15 mmol), (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (53 mg, 0.26 mmol) and triethylamine (30.4 mg, 0.3 mmol) were added into a tube, followed by the addition of DMSO (1 mL). The mixture was stirred at room temperature for 18 h and then filtered, and purified via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) giving (pyridin-4-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate. LC/MS: *m/z* 474.30 (M+H)+ at 1.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00416] Example 115


[00417] **2-Ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00418]



[00419] **2-Ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00420]




[00421] To a solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added 2-ethyl-2-hydroxybutanoic acid (39.30 mg, 0.297 mmol). It was followed by the addition of triethylamine (63 μL), then a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.51 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00422] Example 116


[00423] **(Pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00424]



[00425] **(Pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00426]




[00427] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol) , (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (67 mg, 0.32 mmol) and triethylamine (45 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 442.50 $(M+H)^+$ at 1.96 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00428] Example 117


[00429] **(*S*)-Tetrahydrofuran-3-yl 4-(6-fluoro-2-(2-hydroxyphenyl) quinazolin-4-yl)piperazine-1-carboxylate**

[00430]



[00431] **(*S*)-Tetrahydrofuran-3-yl chloroformate**


[00432]



[00433] A stirred solution of (*S*)-tetrahydrofuran-3-ol (7.9 g, 90 mmol) in anhydrous CH_2Cl_2 (50 mL) under an N_2 atmosphere was cooled in an ice bath, and a 20% solution of phosgene in toluene (134 mL, 270 mmol) was slowly added. The reaction was allowed to warm to room temperature overnight, and the solvent was removed under vacuum to afford (*S*)-tetrahydrofuran-3-yl chloroformate (12.1 g, 85%) as a clear liquid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 5.42–5.39 (m, 1H), 4.01–3.84 (m, 4H), 2.31–2.13 (m, 2H).

[00434] **(*S*)-Tetrahydrofuran-3-yl 4-(6-fluoro-2-(2-hydroxyphenyl) quinazolin-4-yl)piperazine-1-carboxylate**

[00435]




[00436] To a solution of 2-(6-fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.08 mmol) in DMF (1 mL) was added triethylamine (22 μL, 0.16 mmol) followed by the dropwise addition of (*S*)-tetrahydrofuran-3-yl chloroformate (12 mg, 0.08 mmol) at 0 °C. The reaction was complete immediately after the addition of the chloroformate. Purification using

reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-tetrahydrofuran-3-yl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00437] Example 118


[00438] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00439]



[00440] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00441]



[00442] Method A

[00443] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to (*R*)-2-hydroxybutanoic acid (31 mg, 0.297 mmol), followed by the addition of triethylamine (63 μL), then a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-2-hydroxy-1-(4-(2-(2-


hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 421.30 $(M+H)^+$ at 2.51 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00444] Method B

[00445] To a solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.82 mmol) in CH_2Cl_2 (6 mL) was added triethylamine (227 μL, 1.63 mmol) followed by the addition of (*R*)-2-hydroxybutanoic acid (110 mg, 1.06 mmol) and HATU (380 mg, 1.00 mmol). The reaction mixture was stirred at room temperature for 3 h and then quenched with H_2O. The aqueous layer was extracted twice with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2:hexanes gave (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one (280 mg, 88%). LC/MS: *m/z* 393.3 $(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.47 (m, 1H), 8.12 (d, *J* = 8.3 Hz, 1H), 7.89 (m, 2H), 7.57 (m, 1H), 7.40 (m, 1H), 6.96 (m, 2H), 4.92 (d, *J* = 7.1 Hz, 1H), 3.88 (m, 8H), 1.67 (m, 1H), 1.51 (m, 1H), 0.91 (t, *J* = 7.4 Hz, 3H).

[00446] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one hydrochloride**

[00447]




[00448] To a solution of (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one (280 mg, 0.71 mmol) in CH_2Cl_2 under an N_2 atmosphere was added a 2 M HCl solution in ether (0.355 mL, 0.71 mmol), followed by the addition of ether which resulted in the formation of precipitate which was stirred for an hour and then filtered and dried to afford (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one hydrochloride (272 mg, 90%). LC/MS: *m/z* 393.1 $(M+H)^+$ at 2.23 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.27 (dd, *J* = 7.9, 1.6 Hz,

1H), 8.19 (d, J = 8.4 Hz, 1H), 7.96 (m, 2H), 7.65 (m, 1H), 7.48 (m, 1H), 7.04 (m, 2H), 4.26 (dd, J = 7.7, 4.8 Hz, 1H), , 4.03 (m, 4H), 3.79 (m, 5H), 1.65 (m, 1H), 1.50 (m, 1H), 0.90 (t, J = 7.4 Hz, 3H).

[00449] Example 119


[00450] **(*S*)-3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid tetrahydro-furan-3-yl ester**

[00451]



[00452] **(*S*)-3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid tetrahydro-furan-3-yl ester**

[00453]




[00454] To 2-[4-(2-hydroxymethyl-piperazin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (69.2 mg, 0.19 mmol) in 650 μL of CH_2Cl_2 at 0 °C was added (*S*)-tetrahydro-furan-3-ol-chloroformate (26.8 mg, 0.17 mmol), followed by triethylamine (30 μL, 0.22 mmol). The reaction mixture was stirred for an hour and then diluted with 5 mL of CH_2Cl_2 and 5 mL of water, and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure, and the residue was subjected to purification using 30–100% EtOAc–hexanes to give the desired product. LC/MS: *m/z* 465.2 $(M+H)^+$ at 2.5 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00455] Example 120


[00456] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)hexan-1-one**

[00457]



[00458] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)hexan-1-one**

[00459]




[00460] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-hydroxyhexanoic acid (37.5 mg, 0.284 mmol). It was followed by the addition of triethylamine (61 μL), then a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)hexan-1-one as the TFA salt. LC/MS: *m/z* 435.3 $(M+H)^+$ at 2.65 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00461] Example 121


[00462] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00463]



[00464] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00465]




[00466] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to (*R*)-2-hydroxybutanoic acid (31 mg, 0.297 mmol). It was followed by the addition of triethylamine (63 μL), then a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 393.3 $(M+H)^+$ at 2.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00467] Example 122


[00468] **2-Ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00469]



[00470] **2-Ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00471]




[00472] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-ethyl-2-hydroxybutanoic acid (37.5 mg, 0.284 mmol). It was followed by the addition of triethylamine (61 μL), then a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-ethyl-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 435.3 $(M+H)^+$ at 2.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00473] Example 123


[00474] **4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-7-carboxylic acid**

[00475]



[00476] **(*R*)-α-hydroxyisocaproic acid**


[00477]



[00478] To a cooled solution (0–5 °C) of D-leucine (200 g, 1.5 mol) in sulfuric acid (3 L, 1 M) was added dropwise a solution of sodium nitrite (240 g, 3.5 mol) in water (1 L) while maintaining the temperature between 0–5 °C. The reaction mixture was stirred at room temperature overnight. The solution was saturated with sodium chloride and extracted with *tert*-butyl methyl ether (3x). The combined organic layers were dried over Na_2SO_4 and filtered, and the solvent was removed *in vacuo*. (*R*)-α-hydroxyisocaproic acid was isolated as a white solid in a yield of 67% (132 g).

[00479] **1-(4-Benzyl-piperazin-1-yl)-2-hydroxy-4-methyl-pentan-1-one**


[00480]



[00481] To a cooled solution (0–5 °C) of (*R*)-α-hydroxyisocaproic acid (64.5 g, 0.5 mol), 1-benzylpiperazine (88 g, 0.5 mol) and triethylamine (71 ml, 0.5 mol) in CH_2Cl_2 (850 ml) was added in portions HOBt (68 g, 0.5 mol) and EDCI·HCl (96 g, 0.5 mol). The reaction mixture was stirred at room temperature overnight. The organic layer was washed with water (3x) and once with brine, dried over Na_2SO_4, and filtered, and the solvent was removed *in vacuo*. Crude


1-(4-benzyl-piperazin-1-yl)-2-hydroxy-4-methyl-pentan-1-one (132 g, 91%) was used in the next step without further purification

[00481] 2-Hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one




[00482] To a solution of 1-(4-benzyl-piperazin-1-yl)-2-hydroxy-4-methyl-pentan-1-one (132 g) in methanol (1 L) was added Pd/C (20 g,10% weight Pd on carbon). The reaction mixture was stirred at room temperature overnight under an atmosphere of hydrogen. The reaction mixture was filtered through Celite, and the solvent was removed *in vacuo*. 2-Hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one was obtained as an oil (68 g, 74%).

[00483] 1,2,3,4-Tetrahydro-2,4-dioxoquinazoline-7-carboxylic acid



[00484] To a stirring mixture of dimethyl 2-aminobenzene-1,4-dioate (10.5 g, 50 mmol) and AcOH (3.4 mL, 60 mmol) in H_2O (200 mL) was added KOCN (8.1 g, 100 mmol). The reaction mixture was heated at 100 °C for 3 h. After cooling in an ice bath, NaOH (24 g, 600 mmol) was slowly added to the mixture and stirred for another 3 h at room temperature. Acidification with concentrated HCl resulted in formation of a yellow solid which was filtered, washed with water and dried under vacuum (8.3 g). NMR data showed that the solid consisted of a 3:1 mixture of the desired 1,2,3,4-tetrahydro-2,4-dioxoquinazoline-7-carboxylic acid and a side product. This mixture was used in the next step without further purification. ^{1}H NMR (400 MHz, DMSO-d6) δ 7.98 (d, J = 8.2 Hz, 1H), 7.75 (d, J = 1.2 Hz, 1H), 7.67 (dd, J = 8.2, 1.4 Hz, 1H).


[00485] 2,4-Dichloroquinazoline-7-carboxylic acid



[00490] A mixture of 1,2,3,4-tetrahydro-2,4-dioxoquinazoline-7-carboxylic acid (1.0 g, 4.9 mmol), $POCl_3$ (10 mL) and dimethylaniline (0.5 mL, 3.94 mmol) was heated at 90 °C for 3 h. After cooling and pouring the reaction mixture over ice, it was extracted with EtOAc. The combined extracts were washed with water, dried over Na_2SO_4 and concentrated. Purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 provided 2,4-dichloroquinazoline-7-carboxylic acid as a yellow solid (0.45 g, 38%). ^1H NMR (400 MHz, DMSO-d6) δ 8.46 (d, J = 1.4 Hz, 1H), 8.42 (d, J = 8.7 Hz, 1H), 8.30 (dd, J = 8.6, 1.5 Hz, 1H); LC/MS: *m/z* 243.1 $(M+H)^+$ at 2.49 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00491] **2-Chloro-4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-quinazoline-7-carboxylic acid**


[00492]



[00493] To a solution of 2,4-dichloroquinazoline-7-carboxylic acid (0.45 g, 1.9 mmol) in CH_2Cl_2 (20 mL) was added (*R*)-2-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one (0.37 g, 1.9 mmol) and triethylamine (0.52 mL, 3.7 mmol). The reaction mixture was stirred overnight at room temperature and then purified via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 to obtain 2-chloro-4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-quinazoline-7-carboxylic acid (0.58 g, 77%). LC/MS: *m/z* 407.3 $(M+H)^+$ at 2.44 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00494] **4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-7-carboxylic acid**

[00495]




[00496] A mixture of 2-chloro-4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-quinazoline-7-carboxylic acid (114 mg, 0.28 mmol), 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (88 μL, 0.42 mmol) $Pd(dppf)Cl_2$ (14 mg, 0.017 mmol) and K_2CO_3 (155 mg, 1.1 mmol) in DMF (4 mL) and H_2O (1 mL) was heated in a sealed microwave vial at 170 °C for 6 minutes. After filtration and evaporation of the solvents, the mixture was purified using preparative HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-7-carboxylic acid as the TFA salt. LC/MS: *m/z* 465.3 $(M+H)^+$ at 2.50 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00497] Example 124


[00498] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00499]



[00500] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00501]




[00502] Method A: A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to (*S*)-2-hydroxybutanoic acid (31 mg, 0.297 mmol). It was followed by the addition of triethylamine (63 μL), then a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00503] Method B: To a solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.78 mmol) in CH_2Cl_2 (6 mL) was added triethylamine (217 μL, 1.56 mmol) followed by the addition of (*S*)-2-hydroxybutanoic acid (105 mg, 1.0 mmol) and HATU (380 mg, 1.00 mmol). The reaction mixture was stirred at room temperature for 3 h and then quenched with H_2O. The aqueous layer was extracted twice with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2:hexanes gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one (300 mg, 95%). LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.45 (m, 1H), 8.01 (d, J = 8.5 Hz, 1H), 7.70 (s, 1H), 7.39 (m, 2H), 6.95 (m, 2H), 4.92 (d, J = 7.0 Hz, 1H), 4.28 (q, J = 6.5 Hz, 1H), 3.89 (m, 8H), 2.52 (s, 3H), 1.67 (m, 1H), 1.51 (m, 1H), 0.91 (t, J = 7.4 Hz, 3H)

[00504] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one hydrochloride**

[00505]




[00506] To a solution of (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one (300 mg, 0.73 mmol) in CH_2Cl_2 was added a 2 M HCl solution in ether (0.365 mL, 0.73 mmol), followed by the addition of ether until the salt precipitated out, which was stirred for an hour, collected by vacuum filtration and dried to afford (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one hydrochloride (270 mg, 84%). LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.26 (dd, J = 7.9, 1.6 Hz, 1H), 8.07 (d, J = 8.6 Hz, 1H), 7.73 (s, 1H), 7.47 (m, 2H), 7.03 (m, 2H), 4.26 (dd, J = 7.7, 4.8 Hz, 1H), 4.08 (m, 4H), 3.79 (m, 4H), 2.53 (s, 3H), 1.65 (m, 1H), 1.50 (m, 1H), 0.89 (t, J = 7.4 Hz, 3H).

[00507] Example 125


[00508] **(*R*)-4-Fluoro-2-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one**

[00509]



[00510] 4-{4-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-but-2-enoic acid ethyl ester


[00511]



[00512] To 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol (787 mg, 2.46 mmol) in 8 mL of CH_2Cl_2 at room temperature was added sequentially but-2-enedioic acid monoethyl ester (531 mg, 3.69 mmol), triethylamine(686 μL, 4.92 mmol), BOP (1.63 g, 3.69 mmol). The reaction mixture was stirred for 20 min and diluted with water and CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue was subjected to purification by normal phase LC using 10–100% EtOAc–hexanes to give 4-{4-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-but-2-enoic acid ethyl ester (1.00 g, 91% yield). LC/MS: *m/z* 447.3 $(M+H)^+$ at 2.93 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00513] 4-Hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pent-2-en-1- one

[00514]




[00515] To 4-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-oxo-but-2-enoic acid ethyl ester (655 mg, 1.47 mmol) in 2 mL of diethyl ether at –20 °C was added methyl magnesium bromide (8.4 mL, 11.7 mmol, 1.4 M THF/Toluene), and the reaction mixture

was allowed to warm to 0 °C over 15 minutes The mixture was diluted with 10 mL of water and 15 mL of CH_2Cl_2. The resulting emulsion was filtered, and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure to give the alcohol. LC/MS: *m/z* 433.5 $(M+H)^+$ at 2.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00516] 4-Fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pent-2-en-1-one

[00517]



[00518] To 4-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pent-2-en-1- one (330 mg, 0.76 mmol) at –78 °C in 3 mL of CH_2Cl_2 was added (diethylamino)sulfur trifluoride (185 mg, 1.15 mmol). The reaction mixture was stirred –78 °C for 2.5 h. The reaction mixture was diluted with 10 mL of water and 10 mL of CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue was subjected to purification by normal phase LC using 10–100% EtOAc–hexanes to give 4-fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pent-2-en-1-one (133 mg, 40% yield). LC/MS: *m/z* 435.4.3 $(M+H)^+$ at 2.92 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00519] 4-Fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one




[00520]

[00521] To 4-fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pent-2-en-1-one (228 mg) in 1.5 mL of methanol was added Pd/C (34 mg, 10% weight Pd on carbon), and the reaction mixture was hydrogenated with a hydrogen balloon for 1 h. The resulting mixture was filtered through Celite, and the solvent was removed to give 4-fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one. LC/MS: *m/z* 437.4 $(M+H)^+$ at 2.86 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00522] (*R*)-4-Fluoro-2-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one

[00523]




[00524] To 4-fluoro-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one (225 mg, 0.53 mmol) in 500 μL of THF was added a 2.7 M LDA solution (0.8 mL, 2.1 mmol) at –78 °C. The resulting solution was stirred at –78 °C for 25 min, and then (1*R*)-(-)-(10-camphorsulfonyl)oxaziridine (361 mg, 1.6 mmol) in 1.5 mL of THF was added slowly. The reaction mixture was allowed to warm to –45 °C over 30 min, and then diluted with water and CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified by preparative reverse

phase HPLC using 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*R*)-4-fluoro-2-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one as the TFA salt. LC/MS: *m/z* 453.4 $(M+H)^+$ at 2.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00525] Example 126


[00526] **3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid phenyl ester**

[00527]



[00528] **3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester**

[00529]




[00530] To 2-(4-chloro-7-methyl-quinazolin-2-yl)-phenol (245 mg, 0.91 mmol) in 3.0 mL of CH_2Cl_2 was added 3-hydroxymethyl-piperazine-1-carboxylic acid benzyl ester (226 mg, 0.58 mmol) and triethylamine (190 μL, 1.37 mmol). The reaction mixture was stirred for 12 hours at room temperature, and additional 3-hydroxymethyl-piperazine-1-carboxylic acid benzyl ester (100 mg, 0.4 mmol) and triethylamine (200 μL, 1.4 mmol) was added, and the reaction mixture was heated at 40 °C for 6 h. The reaction mixture was cooled, and diluted with 5 mL of CH_2Cl_2

and 5 mL of water, and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure, and the residue was purified by silica gel chromatography eluting with 20–85% EtOAc/hexanes to give 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester(216 mg, 65 %). LCMS: *m/z* 485 $(M+H)^+$ at 3.03 min (10%-99% CH_3CN/H_2O)

[00531] 2-[4-(2-Hydroxymethyl-piperazin-1-yl)-7-methyl-quinazolin-2-yl]-phenol


[00532]



[00533] To 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester (200 mg, 0.41 mmol) in 1.7 mL of methanol was added 39 mg of Pd/C (10% wt Pd on carbon). The reaction mixture was stirred under a hydrogen atmosphere for 3 h. The mixture was filtered through Celite, and the solvent was removed to give 2-[4-(2-hydroxymethyl-piperazin-1-yl)-7-methyl-quinazolin-2-yl]-phenol. LCMS: *m/z* 351.2 $(M+H)^+$ at 2.11 min (10%-99% CH_3CN/H_2O).

[00534] 3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid phenyl ester

[00535]




[00536] To 2-[4-(2-hydroxymethyl-piperazin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (31.9 mg, 0.091 mmol) in 300 μL of CH_2Cl_2 was added at 0 °C sequentially triethylamine (13.9 μL) and phenyl chloroformate (14.3 mg, 0.091 mmol). The reaction mixture was stirred for 30 min and was warmed to room temperature and stirred for an additional 40 minutes At the end of this period, the solvent was removed, and the residue was dissolved in DMSO and purified by preparative reverse phase HPLC using 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) as eluent to give 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid phenyl ester as the TFA salt. LC/MS: *m/z* 471.2 (M+H)+ at 2.93 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00537] Example 127


[00538] **3-Cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[00539]



[00540] **3-Cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[00541]



[00542] Method A


[00543] To 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (53 mg, 0.17 mmol) was added sequentially 3-cyclopentylpropanoyl chloride (29 mg, 0.19 mmol) in 550 μL of CH_2Cl_2 and triethylamine (28 μL, 0.2 mmol). The mixture was stirred at 0 °C for 20 minutes After adding H_2O and CH_2Cl_2, the phases were separated, and the organic layer was dried over Na_2SO_4 and concentrated under vacuum. Purification using preparative reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 3-cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 445.5 $(M+H)^+$ at 2.32 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00544] Method B

[00545] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (200 mg, 0.62 mmol) in CH_2Cl_2 (6.0 mL) was stirred under an N_2 atmosphere. Then triethylamine (170 μL, 1.24 mmol) was added, and the reaction was cooled to –10 to –20 °C. After adding 3-cyclopentylpropanoyl chloride (96 μL in 600 μL THF, 0.62 mmol), the reaction mixture was stirred for 30 minutes. CH_2Cl_2 was added, and the organic phase was washed 2x with H_2O, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 CH_2Cl_2: hexanes gave 3-cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one (240 mg, 86%). LC/MS: *m/z* 445.50 $(M+H)^+$ at 3.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.45 (dd, *J* = 8.2, 1.8 Hz, 1H), 8.00 (d, *J* = 8.5 Hz, 1H), 7.69 (s, 1H), 7.39 (m, 2H), 6.95 (m, 2H), 3.95 (dd, *J* = 19.9, 5.6 Hz, 4H), 3.74 (m, 4H), 2.52 (s, 3H), 2.38 (t, *J* = 7.8 Hz, 2H), 1.76 (m, 3H), 1.55 (m, 6H), 1.16 (m, 2H).

[00546] **3-Cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one hydrochloride**

[00547]




Under an N_2 atmosphere, a 1 M HCl solution in diethyl ether (0.54 mL, 0.54 mmol) was added dropwise to a stirring solution of 3-cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one (240 mg, 0.54 mmol) in CH_2Cl_2 (20 mL). After 10 min, ether was added until a precipitate formed. The solid was collected by filtration and dried under vacuum to give 3-cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one hydrochloride (230 mg, 88%). LC/MS: *m/z* 445.30 $(M+H)^+$ at 3.08 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.26 (d, J = 8.0 Hz, 1H), 8.09 (d, J = 8.5 Hz, 1H), 7.78 (s, 1H), 7.50–7.46 (m, 2H), 7.09 (d, J = 8.1 Hz, 1H), 7.04–7.00 (m, 1H), 4.15–4.11 (m, 4H), 3.79–3.74 (m, 4H), 2.54 (s, 3H), 2.37 (t, J = 7.7 Hz, 2H), 1.82–1.74 (m, 3H), 1.60–1.45 (m, 6H), 1.12–1.08 (m, 2H)

[00548]

[00549] Example 128


[00550] **1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-(tetrahydro-2*H*-pyran-4-yl)ethanone**

[00551]



[00552] **1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-(tetrahydro-2*H*-pyran-4-yl)ethanone**

[00553]




[00554] 2-(6-Fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.077 mmol), 2-(tetrahydro-2*H*-pyran-4-yl)acetic acid (14.3 mg, 0.10 mmol), triethylamine (22μL, 0.154 mmol), and HATU (38 mg, 0.10 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-(tetrahydro-2*H*-pyran-4-yl)ethanone. LC/MS: *m/z* 451.5 $(M+H)^+$ at 2.60 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00555] Example 129


[00556] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one**

[00557]




[00558] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one**

[00559]




[00560] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to 2-hydroxy-2-methylbutanoic acid (39.3 mg, 0.297 mmol). It was followed by the addition of triethylamine (63 μL) and a solution of HATU (113 mg) in 0.5 mL



[00557] DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one as the TFA salt. LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00558] Example 130


[00559] **(*R*)-2-Hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone**



[00560] **2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol, oxalate salt**

[00561] To 2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol (30.6 g, 95.4 mmol) in 900 mL CH_2Cl_2 was added oxalic acid (9.45 g, 105 mmol, 1.1 eq.) dissolved in 36 mL of methanol. The resulting cloudy solution was stirred for 3h and the resulting solid was filtered, washed with hexanes, and dried to give 29.3 g (75 %) the oxalate salt of 2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol. LC/MS: *m/z* 321.2 $(M+H)^+$ at 2.36 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00562] **2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol**




[00568]

[00569] To the oxalate salt of 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol (1.32 g, 3.2 mmol) in 10 mL of CH_2Cl_2 was added triethylamine (2.2 mL, 16.0 mmol) at room temperature. The resulting mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with 10 mL of water, and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure to give 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol as an oil which was used without further purification.

[00570] **(*R*)-2-Hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone**

[00571]




[00572] To 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol (64 mg, 0.19 mmol) in 600 μL of CH_2Cl_2 was added sequentially 3-(*R*)-phenyl lactic acid (35 mg, 0.21 mmol), BOP (93 mg, 0.21 mmol), and triethylamine (27.7 μL, 0.2 mmol) at room temperature. The reaction mixture was stirred for 1.5 h, diluted with 10 mL of methylene chloride, and washed with water (2 x 10 mL). The solvent was removed under reduced pressure to give an oil which was purified by normal phase LC (35%–100% EtOAc/ hexanes) to give (*R*)-2-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone. LC/MS: *m/z* 469.3 $(M+H)^+$ at 2.87 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00573] Example 131


[00574] **(*S*)-2-Hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone**

[00575]



[00576] **(*S*)-2-Hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone**

[00577]




[00578] To 2-(7-methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol (79.9 mg, 0.25 mmol) in 800 μL of CH_2Cl_2 was added sequentially 3-(*R*)-phenyl lactic acid(41.4 mg, 0.25 mmol), BOP(110 mg, 0.25 mmol), triethylamine (34.7 μL, 0.25 mmol) at room temperature. The reaction mixture was stirred for 1.5 h, diluted with 10 mL of methylene chloride, and washed with water (2 x 10 mL). The solvent was removed under reduced pressure to give an oil which was purified by normal phase LC (35%-100% EtOAc/hexanes) to give (*S*)-2-hydroxy-1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-2-phenyl-ethanone. LC/MS: *m/z* 469.4 $(M+H)^+$ at 2.88 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00579] Example 132


[00580] **(Pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00581]




[00582] **(Pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate**

[00583]



[00584] A solution of (pyridin-3-yl)methanol (2 g, 18.32 mmol) and di(1*H*-imidazol-1-yl)methanone (5.94 g, 36.65 mmol) in 20 mL CH_2Cl_2 was heated overnight at 50 °C The reaction was washed with water, dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 10–70% EtOAc in CH_2Cl_2 gave (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (3 g, 81%). LC/MS: *m/z* 204.1 $(M+H)^+$ at 0.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, $CDCl_3$) δ 8.74 (d, J = 1.9 Hz, 1H), 8.68 (dd, J = 4.8, 1.4 Hz, 1H), 8.16 (s, 1H), 7.81 (m, 1H), 7.44 (s, 1H), 7.38 (m, 1H), 7.09 (s, 1H), 5.46 (s, 2H).

[00585] **(Pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**




[00586]

[00587] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol), (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (67 mg, 0.32 mmol), and triethylamine (44.6 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (pyridin-3-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 456.5 $(M+H)^+$ at 2.04 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00588] Example 133


[00589] **1-(4-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(pyridin-2-yl)propan-1-one**

[00590]



[00591] **1-(4-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(pyridin-2-yl)propan-1-one**

[00592]




[00593] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (0.5 mL) was added to 3-(pyridin-2-yl)propanoic acid (21.23 mg, 0.14 mmol). Triethylamine (25 μL) was added, followed by a solution of HATU (45 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(pyridin-2-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 454.3 $(M+H)^+$ at 1.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00594] Example 134


[00595] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentan-1-one**

[00596]



[00597] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentan-1-one**

[00598]




[00599] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-hydroxypentanoic acid (33.6 mg, 0.28 mmol). It was followed by the addition of triethylamine (61 μL), and a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentan-1-one as the TFA salt. LC/MS: *m/z* 421.1 $(M+H)^+$ at 2.46 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00600] Example 135


[00601] **(*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one**

[00602]



[00603] **(*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one**

[00604]



[00605] Method A


[00606] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.15 mmol) was placed in a tube charged with a stir bar, (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (26 mg, 0.18 mmol) in1 ml of DMF, and triethylamine (30 mg, 41 μL, 0.29 mmol), and the tube was cooled to 0 °C. HATU (68 mg, 0.18 mmol) was added, and the reaction was stirred at 0 °C for 10 minutes and then allowed to warm to room temperature. After 40 minutes, the reaction was filtered, and purified by reverse phase HPLC to give the TFA salt of (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one. LC/MS: *m/z* 467.1 $(M+H)^+$ at 2.59 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00607] Method B

[00608] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (250 mg, 0.74 mmol) was suspended in anhydrous DMF (5 mL) and cooled to 0 °C (internal temperature). Under an N_2 atmosphere, (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (118.4 mg, 0.81 mmol) was added followed by triethylamine (0.207 mL, 1.48 mmol). To this stirring solution was added HATU (337 mg, 0.888 mmol). After the complete addition of HATU, the mixture was allowed to warm to 10 °C. After 45 min the reaction was complete, and it was quenched with an equal portion of ice water. A yellow precipitate formed which was collected by vacuum filtration, dissolved in CH_2Cl_2, and the CH_2Cl_2 solution was desiccated with Na_2SO_4, filtered, and concentrated to a viscous yellow-orange oil. The crude material was purified via silica gel chromatography using 70% CH_2Cl_2/hexanes (1:1) and 30% EtOAc to afford (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one as a yellow solid (171 mg, 50%). LC/MS: *m/z* 467.1 $(M+H)^+$ at 2.63 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.02 (d, J = 8.6 Hz, 1H), 7.69 (s, 1H), 7.43 (dd, J = 8.6, 1.5 Hz, 1H), 7.37–7.32 (m, 1H), 6.80 (d, J = 8.3 Hz, 1H), 6.76–6.71 (m, 1H), 4.87 (d, J = 7.2 Hz, 1H), 4.47–4.43 (m, 1H), 4.01–3.75 (m, 8H), 2.52 (s, 3H), 1.55 (dd, J = 14.3, 3.0 Hz, 1H), 1.40 (dd, J = 14.3, 8.8 Hz, 1H), 0.96 (s, 9H)

[00609] **(*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one hydrochloride**


[00610]



[00611] (*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one (171 mg, 0.367 mmol) was dissolved in anhydrous CH_2Cl_2 (2 mL) followed by the addition of Et_2O (6 ml) under an N_2 atmosphere. A 2.0 M HCl solution in Et_2O (0.184 mL, 0.367 mmol) was added over a 1 minute period. The reaction solution changed from a clear yellow solution to a milky white slurry. After complete addition of the HCl solution, the reaction was allowed to stir for an additional 10 minutes. The product was collected by vacuum filtration and dried under vacuum to obtain (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4,4-dimethylpentan-1-one hydrochloride as a light yellow solid (170 mg, 92%). LC/MS: *m/z* 467.3 $(M+H)^+$ at 2.60 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.17 (d, *J* = 8.3 Hz, 1H), 7.66 (s, 1H), 7.56 (d, *J* = 8.1 Hz, 1H), 7.48–7.43 (m, 1H), 6.92–6.83 (m, 2H), 4.44–4.41 (m, 1H), 3.87–3.68 (m, 8H), 2.56 (s, 3H), 1.55 (dd, *J* = 14.3, 3.0 Hz, 1H), 1.41 (dd, *J* = 14.3, 8.8 Hz, 1H), 0.96 (s, 9H).

[00612] Example 136


[00613] **(*R*)-1-(4-(2-(5-Fluoro-2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**



[00614]

[00615] **7-Methylquinazoline-2,4(1*H*,3*H*)-dione**


[00616]



[00617] To a suspension of 2-amino-4-methylbenzoic acid (58.9 g, 390 mmol) in water (1.5 L) and glacial acetic acid (50 ml) was added dropwise a solution of potassium cyanate (80.5 g, 993 mmol) in water (300 ml). After completion of the addition and stirring at room temperature for half an hour, sodium hydroxide pellets (500 g) were added at such a rate that the temperature was kept below 50 °C (with ice cooling). During the addition the mixture became a clear solution for a short period, and upon continuation of the sodium hydroxide addition a cream precipitate started forming. The suspension was cooled to 0–5 °C and the precipitate was collected by filtration and washed twice with water (150 ml). The solid was poured in water (1 L) and was acidified with concentrated aqueous HCl (30%, 150 ml). The solid was collected by filtration and washed with water (150 ml) to yield 7-methylquinazoline-2,4(1*H*,3*H*)-dione (57.0 g, 83%) after drying at 45°C under vacuum.

[00618] **2,4-Dichloro-7-methyl-quinazoline**


[00619]



[00620] 7-Methylquinazoline-2,4(1*H*,3*H*)-dione (57.0 g, 324 mmol) was refluxed overnight in phosphorus oxychloride (250 ml) in a flask equipped with a calcium chloride guard tube. The clear, dark solution was cooled in an ice bath and poured slowly into 2 L of ice water. The chocolate brown solid was collected by filtration and washed with cold water (150 ml). The solid was dissolved in dichloromethane (500 ml) and filtered. The filtrate was washed with a saturated solution of NaCl, dried over Na_2SO_4, filtered, and evaporated to dryness yielding 43.0 g of crude 2,4-dichloro-7-methyl-quinazoline. This material was dissolved in hot heptanes (0.5 L),and filtered while hot, and after cooling to room temperature, the precipitated solid was collected by filtration and washed with pentane (100 ml). The solid was purified by chromatography on silica gel with dichloromethane as the eluent to yield 2,4-dichloro-7-methyl-quinazoline (28.5 g, 41%) as an off-white solid.

[00621] **1-(4-benzyl-piperazin-1-yl)-2-(*R*)-hydroxy-4-methyl-pentan-1-one**


[00622]



[00623] (R)-α-hydroxyisocaproic acid (52.1 g, 0.394 mol) was added to a solution of 1-benzylpiperazine (69.46 g, 0.394 mol) in CH_2Cl_2 (500 mL). The mixture was cooled in ice and Et_3N (57 mL, 0.5 mol) was added, followed by HOBt (53.25 g, 0.394 mol) and EDCI·HCl (76.0 g, 0.396 mol). The reaction mixture was allowed to warm to room temperature overnight. The organic layer was washed with water (3x200 mL). The combined aqueous layers were back-extracted with CH_2Cl_2 (3x25 mL). The combined organic layers were washed with water (3x20 mL), dried over Na_2SO_4 and concentrated to give 1-(4-benzyl-piperazin-1-yl)-2-(*R*)-hydroxy-4-methyl-pentan-1-one as a brown oil (109 g, 0.375 mol, 95%). ^{1}H NMR (300 MHz, $CDCl_3$): δ 7.35-7.24 (m, 5H); 4.35 (dd, *J*= 10 Hz, 2 Hz, 1H); 3.77-3.55 (m, 4H); 3.52 (s, 2H); 3.36 (m, 2H); 4.45 (m, 4H); 1.97 (m, 1H); 1.47-1.38 (m, 1H); 1.29-1.21 (m, 1H); 0.96 (d, *J*= 6 Hz, 3H); 0.94 (d, *J*= 6 Hz, 3H).

[00624] **2-(R)-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one**


[00625]



[00626] 1-(4-benzyl-piperazin-1-yl)-2-(*R*)-hydroxy-4-methyl-pentan-1-one (109 g, 0.375 mol) was dissolved in MeOH (0.5 L). Crystals formed upon addition of 10% Pd/C (16 grams). The mixture was hydrogenated under 1-4 bar hydrogen pressure for two days. The catalyst was filtered off, the filtrate was concentrated to give 2-(R)-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one as a brownish oil (72.7 g, 0.363 mol, 97%). ^{1}H NMR (300 MHz, $CDCl_3$): δ 4.36 (dd, *J*= 10 Hz, 2 Hz, 1H); 3.76-3.66 (m, 1H); 3.62-3.52 (m, 1H); 3.37 (m, 2H); 2.99 (br. s, 2H); 2.89 (br. m, 4H); 1.96 (m, 1H); 1.48-1.38 (m, 1H); 1.29-1.21 (m, 1H); 0.96 (d, *J*= 9 Hz, 3H); 0.94 (d, *J*= 9 Hz, 3H).

[00627] **2-(R)-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one oxalate**


[00628]



[00629] The above product was dissolved in ethanol (250 mL). Oxalic acid dihydrate (45.76 g, 0.363 mol) was added. The thick slurry was diluted with ethanol (250 mL) and stirred at room temperature for 3 hours. The salt was filtered off, washed with ethanol (2x100 mL) and dried *in vacuo* over drying pearls. Yield: 89.0 grams (0.307 mol, 84%) of 2-(R)-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one oxalate as a white solid. ^{1}H NMR (300 MHz, D_2O): δ 4.68 (dd, *J*=3.3, 9.9 Hz, 1H); 3.95 (m, 4H); 3.35 (m, 4H); 1.80 (m, 1H); 1.59 (m, 1H); 1.41 (m, 1H); 0.96 (d, *J*= 6.5 Hz, 3H); 0.95 (d, *J*= 6.5 Hz, 3H).

[00630] **(*R*)-1-(4-(2-Chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**


[00631]



[00632] To a solution of 2,4-dichloro-7-methylquinazoline (2.09 g, 9.83 mmol) in CH_2Cl_2 (10 mL) at 0 °C was added triethylamine (2.74 mL, 19.66 mmol), followed by the addition of (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (1.97 g, 9.83 mmol). The reaction mixture was warmed to room temperature and stirred overnight. The mixture was then quenched with water and extracted twice with CH_2Cl_2. The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated to obtain (*R*)-1-(4-(2-chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (6.78 g, 95%). LC/MS: *m/z* 377.5 $(M+H)^+$ at 2.61 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00633] (*R*)-1-(4-(2-(5-Fluoro-2-methoxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one

[00634]




[00635] (*R*)-1-(4-(2-Chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (50 mg, 0.13 mmol), 5-fluoro-2-methoxyphenylboronic acid (27 mg, 0.16 mmol), $Pd(Ph_3P)_4$ (9.2 mg, 0.008 mmol), and K_2CO_3 (37 mg, 0.27 mmol) were placed into a microwave tube charged with a stir bar. Acetonitrile (2 mL) and H_2O (400 μL) were added, and the vessel was capped and heated at 160 °C for 12 minutes in the microwave reactor. The

reaction was partitioned between EtOAc and H_2O, the layers were separated, and the aqueous layer was extracted once more with EtOAc. The organic extracts were combined, dried over Na_2SO_4, filtered, and concentrated to an orange gel. The reaction was purified by silica gel chromatography using 10%–30% EtOAc in CH_2Cl_2/hexanes (2:1) to afford (*R*)-1-(4-(2-(5-fluoro-2-methoxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as a white foam (70%). LC/MS: *m/z* 467.30 $(M+H)^+$ at 2.38 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00636] (*R*)-1-(4-(2-(5-Fluoro-2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one

[00637]




[00638] (*R*)-1-(4-(2-(5-Fluoro-2-methoxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (30 mg, 0.064 mmol) was dissolved in 1.5 mL anhydrous CH_2Cl_2. The flask was sealed with a septum, placed under an N_2 atmosphere and cooled to –78 °C, and 0.32 mL of a 1 M solution of BBr_3 in CH_2Cl_2 was added over 2 minutes. The reaction was allowed to warm to room temperature. After 5 hours, the reaction was quenched with saturated aqueous $NaHCO_3$, and partitioned between CH_2Cl_2 and water, and the layers were separated. The organic layer was dried over Na_2SO_4, filtered, and concentrated to a red oil. The reaction was purified by reverse phase HPLC to give (*R*)-1-(4-(2-(5-fluoro-2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 453.30 $(M+H)^+$ at 3.02 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00639] Example 137


[00640] (*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one

[00641]



[00642] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00643]




[00644] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to (*S*)-2-hydroxybutanoic acid (29.6 mg, 0.284 mmol). It was followed by the addition of triethylamine (61 μL) and a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)H_2O (0.05% TFA)) gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 407.3 $(M+H)^+$ at 2.28 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00645] Example 138


[00646] **(*S*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one**

[00647]



[00648] **(*S*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one**

[00649]




[00650] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol), (*S*)-2-hydroxy-3,3-dimethylbutanoic acid (15.24 mg, 0.12 mmol), triethylamine (25μL, 0.18 mmol), and HATU (45.6 mg, 0.12 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one as the TFA salt. LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00651] Example 139


[00652] **(*S*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one**

[00653]



[00654] **(*S*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one**


[00655]



[00656] 2-(6-Fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol), (*S*)-2-hydroxy-3,3-dimethylbutanoic acid (16 mg, 0.12 mmol), triethylamine (25 mL, 0.18 mmol), and HATU (45.6 mg, 0.12 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-3,3-dimethylbutan-1-one as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.95 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00657] Example 140


(Benzo[d][1,3]dioxol-7-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate



[00658]

[00659] **(Benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate**


[00660]



[00661] A solution of (benzo[d][1,3]dioxol-7-yl)methanol (2.0 g, 13.1 mmol) and di(1*H*-imidazol-1-yl)methanone (4.26 g, 26.2 mmol) in 20 mL CH_2Cl_2 was heated overnight at 50 °C The reaction was quenched with water and extracted with CH_2Cl_2, and the combined layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 10–70% EtOAc in CH_2Cl_2 gave (benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate (2.8 g, 86%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.15 (t, J = 0.9 Hz, 1H), 7.44 (t, J = 1.4 Hz, 1H), 7.07 (dd, J = 1.6, 0.8 Hz, 1H), 6.95 (m, 2H), 6.84 (m, 1H), 6.01 (s, 2H), 5.33 (s, 2H).

[00662] **(Benzo[d][1,3]dioxol-7-yl)methyl 4-(4a,8a-dihydro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00663]




[00664] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol) , (benzo[d][1,3]dioxol-7-yl)methyl 1*H*-imidazole-1-carboxylate (78 mg, 0.32 mmol), and

triethylamine (44.6 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (benzo[d][1,3]dioxol-7-yl)methyl 4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 485.5 $(M+H)^+$ at 2.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00665] Example 141


[00666] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(1*H*-imidazol-5-yl)propan-1-one**

[00667]



[00668] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(1*H*-imidazol-5-yl)propan-1-one**

[00669]




[00670] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (87 mg, 0.27 mmol), (*S*)-2-hydroxy-3-(1*H*-imidazol-5-yl)propanoic acid (64 mg, 0.41 mmol), triethylamine (76 μL, 0.54 mmol), and BOP (180 mg, 0.41 mmol) in 1 mL of CH_2Cl_2 were stirred at room temperature for

1.5 h. The reaction mixture was diluted with H_2O and CH_2Cl_2. The organic layer was dried over Na_2SO_4 and concentrated. Purification using 1–15% MeOH in CH_2Cl_2 gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(1*H*-imidazol-5-yl)propan-1-one. LC/MS: *m/z* 459.3 $(M+H)^+$ at 2.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00671] Example 142


[00672] **4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile**

[00673]



[00674] **6-Bromo-1*H*-benzo[d][1,3]oxazine-2,4-dione**

[00675]



[00676] Bromine (35 mL, 660 mmol) was added dropwise to a suspension of isatoic anhydride (100 g, 610 mmol) in 1.6 L water at 50 °C. This temperature was maintained for an additional 2 hours. After cooling the solution to room temperature, the solid was filtered and washed twice with water and twice with acetone, yielding 125.6 g (85%) 6-bromo-1*H*-benzo[d][1,3]oxazine-2,4-dione as a pink solid.

[00677] **2-Amino-5-bromo-benzamide**




[00678]

[00679] 6-Bromo-1*H*-benzo[d][1,3]oxazine-2,4-dione (56.0 g, 230 mmol) was suspended in 1 N aq. NH_4OH (600 mL, 2.6 equivalents), and the suspension was stirred 3 d at room temperature. After filtration, the collected solid was washed with water and subsequently dissolved in tetrahydrofuran. This solution was filtered, evaporated to dryness, and dried by repeated azeotropic distillation with toluene. The solid was suspended in CH_2Cl_2, filtered, and washed once CH_2Cl_2 with yielding 35.4 g (71.2%) 2-amino-5-bromo-benzamide.

[00680] **2-(*o*-Anisoyl)-amino-5-bromo-benzamide**


[00681]



[00682] To a solution of 2-amino-5-bromo-benzamide (29.2 g, 136 mmol) and triethylamine (25.0 mL, 173 mmol) in THF (500 mL) was added dropwise *o*-anisoyl chloride (24.0 g, 140 mmol). Stirring at room temperature was continued for 3 h, after which the formed precipitate was filtered and washed once with THF and twice with dichloromethane yielding 2-(*o*-anisoyl)-amino-5-bromo-benzamide (51.4 g, 84%) with 1 equivalent of triethylamine hydrochloride.

[00683] **6-Bromo-2-(2-methoxy-phenyl)-3*H*-quinazolin-4-one**

[00684]




[00685] 2-(*o*-Anisoyl)-amino-5-bromo-benzamide (50.8 g, 105 mmol) was suspended in 2 N aq. NaOH (500 mL) and heated to reflux until a clear solution was obtained (1.5 h). The solution was cooled to room temperature and filtered. The filtrate was acidified with conc. aq. HCl, and the precipitate formed was filtered and washed twice with 1 N aq. HCl and twice with

water. The solid was dried by repeated azeotropic distillation with toluene yielding 6-bromo-2-(2-methoxy-phenyl)-3*H*-quinazolin-4-one (31.3 g, 91%).

[00686] 6-Bromo-4-chloro-2-(2-methoxyphenyl)quinazoline

[00687]



[00688] Method A

[00689] 6-Bromo-2-(2-methoxyphenyl)quinazolin-4(3*H*)-one (674 mg, 2.0 mmol), $POCl_3$ (624 mg, 4 mmol), and *N*, *N*-dimethylaniline (740 mg, 6.1 mmol) were dissolved in benzene (12 mL) and refluxed for 3 h. The reaction mixture was diluted with EtOAc, and the organic phase was washed with aqueous saturated $NaHCO_3$ (1x) and H_2O (2x), dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–5% EtOAc in CH_2Cl_2/hexanes (1:1) gave 6-bromo-4-chloro-2-(2-methoxyphenyl)quinazoline. LC/MS: *m/z* 348.9 $(M+H)^+$ at 3.66 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00690] Method B

[00691] 6-Bromo-2-(2-methoxyphenyl)quinazolin-4(3*H*)-one (0.31 g, 0.94 mmol), $POCl_3$ (86 μL, 0.94 mmol), and *N*, *N*-dimethylaniline (180 μL, 1.4 mmol) were refluxed in dry toluene for 3 h. Additional $POCl_3$ (0.94 mmol) was added, and the reaction was refluxed for one additional hour. The reaction mixture was diluted with EtOAc and water. The aqueous layer was made basic with $NaHCO_3$, and the layers were separated. After the organic phase was washed with water, dried over Na_2SO_4 and concentrated, purification via silica gel chromatography using 0–50% EtOAc in 1:1 hexanes:CH_2Cl_2 gave 6-bromo-4-chloro-2-(2-methoxyphenyl)quinazoline as a yellow solid (144 mg, 44%). ^1H NMR (400 MHz, $CDCl_3$) δ 8.45–8.45 (m, 1H), 8.04–7.99 (m, 2H), 7.82 (dd, *J* = 7.6, 1.7 Hz, 1H), 7.49–7.45 (m, 1H), 7.12–7.08 (m, 1H), 7.06 (d, *J* = 8.4 Hz, 1H), 3.90 (s, 3H); LC/MS: *m/z* 350.9 $(M+H)^+$ at 3.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00692] 2-(6-Bromo-4-chloroquinazolin-2-yl)phenol



[00693]


[00694] Method A

[00695] 6-Bromo-4-chloro-2-(2-methoxyphenyl)quinazoline (393 mg, 1.12 mmol) was dissolved in CH_2Cl_2 (3 mL), and the flask was flushed with N_2. After cooling the reaction mixture to –78 °C, 1 M BBr_3 in CH_2Cl_2 (3.37 mL, 3.37 mmol) was added dropwise, then the reaction was slowly warmed to room temperature and stirred for 2 h. After quenching the mixture with saturated aqueous $NaHCO_3$ (1x), it was transferred into a separatory funnel with CH_2Cl_2. The organic layer was washed with H_2O (2x), dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–5% EtOAc and CH_2Cl_2:hexanes (1:1) gave 2-(6-bromo-4-chloroquinazolin-2-yl)phenol (229 mg, 61%). LC/MS: *m/z* 335.30 $(M+H)^+$ at 4.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00696] Method B

[00697] A solution of 6-bromo-4-chloro-2-(2-methoxyphenyl)quinazoline (0.14 g, 0.4 mmol) in CH_2Cl_2 (mL) was cooled in a dry ice/acetone bath. A solution of 1.0 M BBr_3 in CH_2Cl_2 (1.2 mL, 1.2 mmol) was slowly added. The cooling bath was removed, and the reaction was stirred at room temperature for 2 h. The reaction was diluted with CH_2Cl_2 and made basic with a saturated $NaHCO_3$ solution. The organic layer was separated, washed with water, dried over Na_2SO_4, and evaporated to give 6-bromo-4-chloro-2-(2-hydroxyphenyl)quinazoline (0.15 g).

[00698] **(*R*)-1-(4-(6-Bromo-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**



[00699]

[00700] Method A


[00701] To a solution of 2-(6-bromo-4-chloroquinazolin-2-yl)phenol (228 mg, 0.68 mmol), triethylamine (129 μL, 0.92 mmol), and CH_2Cl_2 (6 mL) was added (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (185 mg, 0.92 mmol) dissolved in CH_2Cl_2 (3 mL). The flask was flushed with N_2 and stirred for 3 hours. The reaction mixture was then transferred into a separatory funnel with CH_2Cl_2, and the organic phase was washed with H_2O (2x), dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0 to 20% EtOAc and CH_2Cl_2:hexanes (1:1) gave (*R*)-1-(4-(6-bromo-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (285 mg, 84%). LC/MS: *m/z* 500.30 $(M+H)^+$ at 3.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00702] Method B

[00703] A solution of 6-bromo-4-chloro-2-(2-hydroxyphenyl)quinazoline (75 mg, 0.22 mmol), (*R*)-2-hydroxy-4-methyl-1-piperazin-1-yl-pentan-1-one (60 mg, 0.3 mmol), triethylamine (42 μL, 0.3 mmol), and CH_2Cl_2 (2 mL) was stirred at room temperature overnight. After diluting with CH_2Cl_2, the reaction mixture was washed with H_2O, dried over Na_2SO_4, and concentrated. The crude material was purified by silica gel chromatography using 0–10% MeOH/CH_2Cl_2 to give (*R*)-1-{4-[6-bromo-2-(2-methoxy-phenyl)-quinazolin-4-yl]-piperazin-1-yl}-2-hydroxy-4-methyl-pentan-1-one (40 mg, 36%). LC/MS: *m/z* 449.3 $(M+H)^+$ at 3.26 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00704] **4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile**

[00705]



[00706] Method A


[00707] A mixture of (*R*)-1-{4-[6-bromo-2-(2-methoxy-phenyl)-quinazolin-4-yl]-piperazin-1-yl}-2-hydroxy-4-methyl-pentan-1-one (20 mg, 0.04 mmol), $Zn(CN)_2$ (4.7 mg, 0.04 mmol), and $Pd(PPh_3)_4$ (1.4 mg, 1.4 μmol) in DMF (0.5 mL) was heated in a microwave synthesizer at 200 °C for 15 minutes Purification via preparative HPLC gave 4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile as the TFA salt. LC/MS: *m/z* 446.3 $(M+H)^+$ at 3.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00708] Method B

[00709] (*R*)-1-(4-(6-Bromo-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (188 mg, 0.38 mmol), $Zn(CN)_2$ (44 mg, 0.38 mmol), and $Pd(Ph_3)_4$ (6.5 mg, 0.0112 mmol) were dissolved in DMF (4 mL) and the reaction mixture was heated in a microwave synthesizer at 200 °C for 15 minutes EtOAc (50 mL) was added and the mixture was washed twice with H_2O. The organic layer was dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in CH_2Cl_2:hexanes (1:1) gave 4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile (127 mg, 75%). LC/MS: *m/z* 446 $(M+H)^+$ at 3.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.60 (d, J = 1.5 Hz, 1H), 8.46 (dd, J = 8.4, 1.8 Hz, 1H), 8.15 (dd, J = 8.7, 1.7 Hz, 1H), 8.00 (d, J = 8.7 Hz, 1H), 7.45–7.41 (m, 1H), 6.99–6.95 (m, 2H), 4.94 (d, J = 7.0 Hz, 1H), 4.39–4.36 (m, 1H), 4.14–4.04 (m, 4H), 3.89–3.67 (m, 4H), 1.82–1.76 (m, 1H), 1.47–1.37 (m, 2H), 0.93–0.91 (m, 6H).

[00710] **4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile hydrochloride**


[00711]



[00712] 4-[4-((*R*)-2-Hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile (137 mg, 0.31 mmol) was dissolved in the minimum amount of CH_2Cl_2. After stirring the solution under an N_2 atmosphere for 30 minutes, 1 M HCl in ether (0.31 mL, 0.31 mmol) was added dropwise to the solution and stirred for 10 minutes. Ether was added to precipitate the hydrochloride salt of 4-[4-((*R*)-2-hydroxy-4-methyl-pentanoyl)-piperazin-1-yl]-2-(2-hydroxy-phenyl)-quinazoline-6-carbonitrile, which was filtered and dried to obtain 136 mg of solid (91%). LC/MS: *m/z* 446 $(M+H)^+$ at 3.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.62 (s, 1H), 8.43–8.41 (m, 1H), 8.17 (dd, J = 8.7, 1.5 Hz, 1H), 8.02 (d, J = 8.7 Hz, 1H), 7.47–7.43 (m, 1H), 7.00–6.96 (m, 2H), 4.39–4.36 (m, 1H), 4.18–4.00 (m, 4H), 3.91–3.68 (m, 4H), 1.85–1.75 (m, 1H), 1.51–1.35 (m, 2H), 0.94–0.91 (m, 6H); ^{13}C NMR (100 MHz, DMSO–d6) δ 172.5, 161.7, 160.6, 135.1, 133.6, 132.4, 129.7, 126.9, 118.8, 118.4, 118.2, 117.4, 113.3, 107.3, 66.9, 48.7, 48.1, 43.5, 42.8, 41.1, 24.0, 23.4, 21.6.

[00713] Example 143


[00714] **(*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**



[00715]

[00716] **(*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[00717]



[00718] Method A

[00719] To a solution of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (25 mg, 0.09 mmol) in DMF (1 mL) was added triethylamine (25 μL, 0.18 mmol) followed by the addition of (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one oxalate (52 mg, 0.18 mmol) at 0 °C. The reaction was stirred for 2 hours, and purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.99 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00720] Method B

[00721] To a mixture of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (200 mg, 0.72 mmol) and CH_2Cl_2 (7 mL) was added (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one oxalate (275 mg, 0.95 mmol). The reaction was complete after two hours. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 mixture of CH_2Cl_2:hexanes gave (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one

(289 mg, 91%). LC/MS: *m/z* 439.30 $(M+H)^+$ at 3.00 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.46 (m, 1H), 8.02 (m, 1H), 7.84 (m, 2H), 7.40 (m, 1H), 6.97 (m, 2H), 4.92 (d, J = 7.2 Hz, 1H), 4.38 (m, 1H), 3.98 (m, 4H), 3.76 (m, 4H), 1.80 (m, 1H), 1.42 (m, 2H), 0.92 (q, J = 3.8 Hz, 6H).

[00722] (*R*)-1-(4-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one hydrochloride


[00723]



[00724] To a solution of (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (285 mg, 0.65 mmol) in CH_2Cl_2 (2 mL) under an N_2 atmosphere was added ether (10 mL), followed by the dropwise addition of a 2 M HCl solution in ether (0.325 mL, 0.65 mmol). A precipitate was formed which was stirred for 30 min, collected by vacuum filtration, and dried to afford (*R*)-1-(4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one hydrochloride (284 mg, 92%). LC/MS: *m/z* 439.30 $(M+H)^+$ at 3.00 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)): ^1H NMR (400 MHz, DMSO-d6) δ 8.32 (dd, J = 8.1, 1.5 Hz, 1H), 8.02 (m, 1H), 7.92 (dd, J = 9.7, 2.7 Hz, 1H), 7.85 (m, 1H), 7.45 (m, 1H), 7.01 (m, 2H), 4.37 (dd, J = 9.2, 4.1 Hz, 1H), 4.04 (m, 4H), 3.83 (m, 4H), 1.76 (m, 1H), 1.41 (m, 2H), 0.90 (dd, J = 6.6, 3.7 Hz, 6H).

[00725] Example 144


[00726] (*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one



[00727]

[00728] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**


[00729]



[00730] Under an N_2 atmosphere, BOP (137 mg, 0.31 mmol) was added in a single portion to a stirring solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (100 mg, 0.31 mmol), (*S*)-2-hydroxy-4-methylpentanoic acid (41 mg, 0.31 mmol), and triethylamine (43 μL, 0.31 mmol) in DMF (0.5 ml). After stirring the mixture for 1 h at room temperature, it was partitioned between H_2O and ether. The organic phase was washed with H_2O (3x 20 mL), dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 1:1 ethyl acetate / hexane gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one as a white solid. LC/MS: *m/z* 435.3 $(M+H)^+$ at 2.62 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.44 (dd, *J* = 8.2, 1.7 Hz, 1H), 7.99 (d, *J* = 8.5 Hz, 1H), 7.67 (s, 1H), 7.40–7.36 (m, 2H), 6.96-6.93 (m, 2H), 4.92 (d, *J* = 7.2 Hz, 1H) 4.41–4.36 (m, 1H), 4.06–3.67 (m, 8H), 2.51 (s, 3H), 1.85–1.73 (m, 1H), 1.49–1.35 (m, 2H), 0.93–0.91 (m, 6H).

[00731] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride**

[00732]




[00733] (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one (90 mg, 0.20 mmol) was dissolved in 1 mL CH_2Cl_2 and treated with 1 equivalent of 2.0 M HCl in ether (100 μL, 0.20 mmol). The formed precipitate was filtered and vacuum dried to obtain (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride. LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.62 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.27 (d, J = 6.8 Hz, 1H), 8.09 (d, J = 8.6 Hz, 1H), 7.78 (s, 1H), 7.50–7.45 (m, 2H), 7.09 (d, J = 8.2 Hz, 1H), 7.03–6.99 (m, 1H), 4.39–4.35 (m, 1H), 4.16–3.68 (m, 8H), 2.54 (s, 3H), 1.84–1.72 (m, 1H), 1.49–1.35 (m, 2H), 0.93 (d, J = 2.8 Hz, 6H).

[00734] Example 145


[00735] **(*R*)-3-Hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoic acid**

[00736]



[00737] **(*R*)-3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoic acid**

[00738]




[00739] (*R*)-Methyl 3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoate (88 mg, 0.20 mmol) and $LiOH \cdot H_2O$ (33 mg, 0.78 mmol) were stirred in THF:H_2O 1:1 at room temperature for 3 h. After acidification with 1 M HCl and extraction with EtOAc, the organic extracts were washed with water, dried over Na_2SO_4, and concentrated. The crude material was then purified via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 to obtain (*R*)-3-hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanoic acid (75 mg, 88%). LC/MS: *m/z* 437.3 $(M+H)^+$ at 2.04 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00740] Example 146

[00741] **(*R*)-1-(4-(2-(2-Chloro-6-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[00742]



[00743] **2-Amino-benzamide**


[00744]



[00745] Isatoic anhydride (40 g, 245 mmol) was suspended in 650 mL 1 N NH_4OH (2.5 equiv) and stirred at room temperature for 3 days. The precipitate was filtered and washed with water. The product was then dissolved in THF, filtered, and concentrated to dryness. The product was dried by azeotropic distillation with toluene and washed with CH_2Cl_2 to yield 10.9 g (32.7%) of 2-amino-benzamide.

[00746] **2-Chloro-*N*-(2-carboxylic acid amide phenyl)-2-methoxy-benzamide**


[00747]



[00748] 2-Amino-benzamide (6.9 g, 50.8 mmol) was dissolved in 50 mL of pyridine and cooled to 0 °C. 2-Chloro-6-methoxy-benzoyl chloride was added dropwise to the solution. After complete addition, the reaction was left to stir at room temperature for three days, which resulted in formation of a brown, cloudy solution. The reaction mixture was then poured into 150 mL of ice water. The precipitate was filtered and washed twice with water, twice with THF and finally twice with CH_2Cl_2 to obtain 2-chloro-*N*-(2-carboxylic acid amide phenyl)-2-methoxy-benzamide (13.3 g, 43.7 mmol, 86%).

[00749] **2-(2-Chloro-6-methoxy-phenyl)-3*H*-quinazolin-4-one**


[00750]



[00751] 2-Chloro-*N*-(2-carboxylic acid amide phenyl)-2-methoxy-benzamide (13 g, 42.7 mmol) was suspended in 100 mL of a 2 N NaOH solution and heated to reflux. After refluxing for 3 hours, another 25 mL of a 2 N NaOH solution was added, and the reaction was refluxed for another hour. The mixture was cooled to room temperature and acidified with acetic acid to pH 5. The formed precipitate was collected by filtration. The product was purified over alumina using EtOAc as an eluent giving 1.7 g (5.9 mmol, 14%) of 2-(2-chloro-6-methoxy-phenyl)-3*H*-quinazolin-4-one.

[00752] **4-Chloro-2-(2-chloro-6-methoxy-phenyl)-quinazoline**


[00753]



[00754] 2-(2-Chloro-6-methoxy-phenyl)-3*H*-quinazolin-4-one (1.7 g, 5.9 mmol) was dissolved in 25 mL benzene. Then, *N,N*-dimethylaniline (1.15 mL, 9 mmol)and $POCl_3$ (1.65 mL, 17.7 mmol) were added. The reaction mixture was refluxed for 3 hours during which the yellow suspension changed to a dark red color. The mixture was cooled and diluted with 50 mL toluene. The solution was poured onto ice. Saturated aq. $NaHCO_3$ was added while stirring and cooling the mixture until the pH remained constant at 7. The layers were separated, and the aqueous layer was extracted with 100 mL toluene. The toluene layers were combined and washed with 100 mL saturated aqueous NaCl solution, 150 mL 0.5 N HCl, 150 mL 5% aq. $NaHCO_3$ and saturated aqueous NaCl solution. The toluene layer was dried over Na_2SO_4, filtered, and concentrated to dryness to yield 1.87 g impure product. The product was purified over silica gel with heptane/CH_2Cl_2 (2:1) as an eluent to yield 4-chloro-2-(2-chloro-6-methoxy-phenyl)-quinazoline (1.22 g, 64%).

[00755] **3-Chloro-2-(4-chloroquinazolin-2-yl)phenol**


[00756]



[00757] To a solution of 4-chloro-2-(2-chloro-6-methoxyphenyl)quinazoline (300 mg, 0.98 mmol) in 10 mL CH_2Cl_2 was added dropwise 5 equivalents of a 1 M BBr_3 solution in CH_2Cl_2 at –78 °C. The reaction was warmed to room temperature and was complete in 30 minutes. The reaction was quenched with a saturated aqueous $NaHCO_3$ solution to pH 7. The aqueous layer was extracted with CH_2Cl_2,and the combined extracts were dried over Na_2SO_4, filtered, and concentrated to obtain 3-chloro-2-(4-chloroquinazolin-2-yl)phenol. LC/MS: *m/z* 291.3 $(M+H)^+$ at 3.16 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00758] **(*R*)-1-(4-(2-(2-Chloro-6-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[00759]




[00760] To a solution of 3-chloro-2-(4-chloroquinazolin-2-yl)phenol (42 mg, 0.14 mmol) in 2 mL CH_2Cl_2 was added triethylamine (40 μL) followed by the addition of (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (37.5 mg, 0.187 mmol). The reaction was complete after 1 hour. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-1-(4-(2-(2-chloro-6-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 455.5 $(M+H)^+$ at 2.45 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00761] Example 147


[00762] **(*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[00763]



[00764] **(*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[00765]



[00766] Method A

[00767] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol (50 mg, 0.174 mmol) in 1 mL DMF was added triethylamine (35.2 mg, 0.348 mmol), followed by the addition of (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (42.1 mg, 0.21 mmol). After stirring the reaction for 1 hour, it was filtered, and purified via preparative reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 453.3 (M+H)+ at 2.40 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00768] Method B

[00769] 2-(4-Chloro-7-methylquinazolin-2-yl)-3-fluorophenol (300 mg, 1.04 mmol) in CH_2Cl_2 (5 mL) was cooled using an ice water batch. To this stirring solution was added (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (312 mg, 1.56 mmol), followed by

triethylamine (210 mg, 291 μL, 2.08 mmol). After letting the reaction warm to room temperature, it was stirred overnight. The mixture was partitioned between water and CH_2Cl_2 and separated, and the aqueous layer was extracted with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated to a viscous yellow oil. Purification via silica gel chromatography using 0–30% EtOAc in CH_2Cl_2/hexanes (2:1) gave (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as a bright yellow foam/solid. (413 mg, 88%). LC/MS: *m/z* 453.1 $(M+H)^+$ at 2.44 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.01 (d, *J* = 8.5 Hz, 1H), 7.68 (s, 1H), 7.42 (d, *J* = 8.5 Hz, 1H), 7.37–7.32 (m, 1H), 6.80 (d, *J* = 8.3 Hz, 1H), 6.76–6.71 (m, 1H), 4.91 (d, *J* = 7.2 Hz, 1H), 4.40–4.35 (m, 1H), 4.02–3.65 (m, 8H), 2.52 (s, 3H), 1.82–1.73 (m, 1H), 1.47–1.34 (m, 2H), 0.91 (dd, *J* = 6.5, 4.1 Hz, 6H).

[00770] (*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one hydrochloride

[00771]




[00772] (*R*)-1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (406 mg, 0.898 mmol) was dissolved in anhydrous CH_2Cl_2 (3 mL) followed by the addition of Et_2O (6 mL) under an N_2 atmosphere. A 2.0 M HCl solution in Et_2O (0.449 mL, 0.898 mmol) was added over a 2 minute period. The reaction solution changed from a clear yellow solution to a turbid white slurry. After complete addition of the HCl solution, the reaction was allowed to stir for an additional 15 minutes. The product was collected by vacuum filtration and dried under vacuum to obtain (*R*)-1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one hydrochloride as a white solid (403 mg, 92%). LC/MS: *m/z* 453.5 $(M+H)^+$ at 2.44 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.20 (d, *J* = 8.7 Hz, 1H), 7.71 (s,

1H), 7.60–7.60 (m, 1H), 7.50–7.44 (m, 1H), 6.99 (d, J = 8.4 Hz, 1H), 6.87 (t, J = 9.1 Hz, 1H), 4.36–4.33 (m, 1H), 4.25–3.67 (m, 8H), 2.57 (s, 3H), 1.83–1.71 (m, 1H), 1.49–1.35 (m, 2H), 0.91 (d, J = 6.7 Hz, 6H).

[00773] Example 148


[00774] **(*R*)-Tetrahydrofuran-3-yl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00775]



[00776] **(*R*)-Tetrahydrofuran-3-yl chloroformate**


[00777]



[00778] A stirred solution of (*R*)-tetrahydrofuran-3-ol (7.9 g, 90 mmol) in anhydrous CH_2Cl_2 (50 mL) under an N_2 atmosphere was cooled in an ice bath, and a 20% solution of phosgene in toluene (134 mL, 270 mmol) was slowly added. The reaction was allowed to warm to room temperature overnight, and the solvent was removed under vacuum to afford (*R*)-tetrahydrofuran-3-yl chloroformate (12.1 g, 85%) as a clear liquid.

[00779]

[00780] **(*R*)-Tetrahydrofuran -3-yl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**




[00781]

[00782] 2-(6-Fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.08 mmol), (*R*)-tetrahydrofuran-3-yl chloroformate (12 mg, 0.08 mmol), triethylamine (22 μL, 0.154 mmol) and HATU (38 mg, 0.10 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-tetrahydrofuran-3-yl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00783] Example 149


[00784] **(2*R*)-(*R*)-Tetrahydrofuran-3-yl 2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00785]



[00786] **(2*R*)-(*R*)-Tetrahydrofuran -3-yl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**


[00787]



[00788] To a solution of 2-(4-((*R*)-3-((benzyloxy)methyl)piperazin-1-yl)-7-methylquinazolin-2-yl)phenol (75 mg, 0.17 mmol) in DMF (1 mL) was added triethylamine (47 μL) followed by the dropwise addition of (*R*)-tetrahydrofuran-3-yl chloroformate (25 mg, 0.17 mmol) at 0 °C. The reaction was warmed to room temperature and stirred for 10 minutes. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-(*R*)-tetrahydrofuran-3-yl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 455.3 $(M+H)^+$ at 2.95 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00789] **(2*R*)-(*R*)-Tetrahydrofuran -3-yl 2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00790]




[00791] To a solution of (2*R*)-(*R*)-tetrahydrofuran-3-yl 2-((benzyloxy)methyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate TFA salt (94 mg, 0.17 mmol) in ethanol was added $Pd(OH)_2$ (78 mg), and the reaction was heated at 50 °C under an H_2 atmosphere at ambient pressure. The reaction was filtered, and purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-(*R*)-tetrahydrofuran-3-yl 2-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate trifluoroacetate. LC/MS: *m/z* 465.5 $(M+H)^+$ at 2.23 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00792] Example 150


[00793] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00794]



[00795] **2-Methoxy-6-methyl-benzoic acid**


[00796]



[00797] Ethyl-2-methoxy-6-methylbenzoate (30.4 g, 0.157 mol) was added to 100 mL 3.2 M NaOH (2 equiv) and 150 mL hot EtOH. The mixture was refluxed overnight after which EtOH was removed *in vacuo*. The aqueous mixture was acidified with 5 M HCl to pH 3. CH_2Cl_2 (200 mL) was added, and the layers were separated. The water layer was extracted two times with 200 mL CH_2Cl_2, and the organic layers were combined and dried over Na_2SO_4. Filtering the Na_2SO_4 and concentrating the CH_2Cl_2 to dryness yielded 21.4 g (0.129 mol, 82.3%) of 2-methoxy-6-methyl-benzoic acid.


[00798] **2-Methoxy-6-methyl-benzoyl chloride**

[00799]



[00800] 2-Methoxy-6-methyl-benzoic acid (21.4 g, 0.129 mol) was refluxed for 3 hours in 230 mL thionyl chloride. Excess thionyl chloride was removed under reduced pressure. Co-evaporation of the residue with toluene gave 23.8 g of 2-methoxy-6-methyl-benzoyl chloride.

[00801] ***N*-(2-Cyano-5-methyl-phenyl)-2-methoxy-6-methyl-benzamide**




[00802]

[00803] 2-Amino-4-methyl-benzonitrile (15.4 g, 0.117 mol) was dissolved in 100 mL pyridine and cooled to 0 °C. To this mixture was added dropwise 2-methoxy-6-methylbenzoyl chloride (24 g, 0.13 mol, 1.1 equiv). During the addition the temperature did not exceed 2 °C. The reaction was stirred at room temperature for 48 hours. The mixture was poured into 400 mL ice water, and the precipitate was collected by filtration and washed with water. The crude product was dissolved in 600 mL CH_2Cl_2, and the solution was washed twice with 500 mL of a 1 N HCl solution and once with 400 mL of a saturated aq. NaCl solution. The CH_2Cl_2 layer was dried over Na_2SO_4, filtered, and concentrated to dryness to yield 25.44 g (0.09 mol, 77.6%) of *N*-(2-cyano-5-methyl-phenyl)-2-methoxy-6-methyl-benzamide.

[00804] **2-(2-Methoxy-6-methyl-phenyl)-7-methyl-3*H*-quinazolin-4-one**

[00805]



[00806] *N*-(2-Cyano-5-methyl-phenyl)-2-methoxy-6-methyl-benzamide (25 g, 0.09 mol) was suspended in 500 mL EtOH, and 121.3 g of 33% aq. NaOH (1 mol, 11 equiv) was added. To this was added a 35% H_2O_2 solution (50 mL, 0.58 mol), and the reaction was heated to reflux. Additional H_2O_2 was added dropwise until the reaction mixture became clear. EtOH was removed under reduced pressure, and the precipitate formed was removed by filtration. The solution was acidified with acetic acid to pH 5, and the precipitate formed was collected by filtration. The precipitate was washed twice with water and once with diethyl ether. The product was purified over alumina using EtOAc/heptane (1:1) as an eluent. Another purification was performed, over silica gel with the same eluent to yield 2-(2-methoxy-6-methyl-phenyl)-7-methyl-3*H*-quinazolin-4-one (1.61 g).

[00807] **4-Chloro-2-(2-methoxy-6-methyl-phenyl)-7-methyl-quinazoline**


[00808]



[00809] 2-(2-Methoxy-6-methyl-phenyl)-7-methyl-3*H*-quinazolin-4-one (1.61 g, 5.74 mmol) was suspended in benzene after which was added *N,N*-dimethylaniline (1.1 mL, 8.62 mmol) and $POCl_3$ (1.61 mL, 17.27 mmol). The reaction mixture was refluxed for 3 hours during which the color changed from yellow to dark red. The reaction mixture was cooled to room temperature, diluted with 40 mL toluene and poured onto ice. Saturated aq. $NaHCO_3$ was carefully added until the pH remained constant at 7 and no more gas formed. The layers were separated, and the water layer was extracted with toluene. The organic layers were combined and washed with respectively 50 mL saturated aq. NaCl, 60 mL 0.5 N HCl , 40 mL 5% $NaHCO_3$ and 50 mL saturated aq. NaCl. The solution was dried over Na_2SO_4 and evaporated to dryness to yield 1.7 g of the impure product. The product was filtered through silica gel and washed with CH_2Cl_2:heptane (2:1) to yield of 4-chloro-2-(2-methoxy-6-methyl-phenyl)-7-methyl-quinazoline (1.22 g, 71%).

[00810] 2-(4-Chloro-7-methylquinazolin-2-yl)-3-methylphenol


[00811]



[00812] To a solution of 4-chloro-2-(2-methoxy-6-methylphenyl)-7-methylquinazoline (669 mg, 2.24 mmol) in 7 mL CH_2Cl_2 was added dropwise 5 equivalents of a 1 M solution of BBr_3 in CH_2Cl_2 at –78 °C. The reaction was warmed to room temperature and was complete in 30 minutes. The reaction was quenched with a saturated aqueous $NaHCO_3$ solution until the pH was neutral. The aqueous layer was extracted with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated to obtain 2-(4-chloro-7-methylquinazolin-2-yl)-3-methylphenol. LC/MS: *m/z* 285.1 $(M+H)^+$ at 3.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00813] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00814]




[00815] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)-3-methylphenol (60 mg, 2.1 mmol) in 2 mL CH_2Cl_2 was added triethylamine followed by the addition of (*R*)-2-hydroxy-4-methyl-1-(piperazin-1-yl)pentan-1-one (54.8 mg, 2.73 mmol). The reaction was complete after 1 hour. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-2-hydroxy-1-(4-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00816] Example 151

[00817] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentane-1,4-dione**

[00818]




[00819] **Methyl 2-((*R*)-2,2-dimethyl-5-oxo-1,3-dioxolan-4-yl)acetate**

[00820] 

[00821] 2-((*R*)-2,2-Dimethyl-5-oxo-1,3-dioxolan-4-yl)acetic acid (15.8 g, 90.9 mmol) in a 3:1 mixture of THF:MeOH (100 mL) was cooled in an ice bath. After adding 2.0 M $TMSCHN_2$ (50 mL, 100 mmol) the bath was removed, and the mixture was stirred at room temperature for 3 h. The solvent was evaporated, and the crude material was purified via silica gel chromatography using 0–50% EtOAc/hexanes to give methyl 2-((*R*)-2,2-dimethyl-5-oxo-1,3-dioxolan-4-yl)acetate. ^1H NMR (400 MHz, $CDCl_3$) δ 4.73 (dd, J = 6.6, 3.8 Hz, 1H), 3.74 (s, 3H), 2.95 (dd, J = 17.0, 3.9 Hz, 1H), 2.81 (dd, J = 17.0, 6.6 Hz, 1H), 1.63 (s, 3H), 1.57 (s, 3H).


[00822] **4-((*R*)-2-Hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester**

[00823] 

[00824] Methyl 2-((*R*)-2,2-dimethyl-5-oxo-1,3-dioxolan-4-yl)acetate (17.1 g, 90.9 mmol) was stirred in a 1:1 mixture of THF:1 M HCl (200 mL) for 1 h at room temperature. After addition of NaCl to nearly saturate the aqueous layer, the mixture was extracted with EtOAc, and the extracts were dried over Na_2SO_4 and concentrated. To the resulting oil dissolved in dry DMF (500 mL) was added HOBt (13.5 g, 100 mmol) and EDCI (19.2 g, 100 mmol). After stirring for 5 min, benzyl piperazine-1-carboxylate (19.3 mL, 100 mmol) and triethylamine (13.9 mL, 100 mmol) were added to the reaction mixture, which was left stirring at room temperature overnight. Purification via silica gel chromatography using 0–10% $MeOH/CH_2Cl_2$ provided a colorless oil 4-((*R*)-2-hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester (6.71 g, 21%). LC/MS: *m/z* 351.5 $(M+H)^+$ at 2.67 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00825] 4-((*R*)-2-Hydroxy-4-oxo-pentanoyl)-piperazine-1-carboxylic acid benzyl ester


[00826] To a solution of 4-((*R*)-2-hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester (8.5 g, 24.0 mmol) in THF (240 mL) cooled in a dry ice acetone



bath, was added 1.4 M MeMgBr (61 mL, 85 mmol). The reaction was allowed to slowly warm to room temperature overnight. After quenching the mixture with saturated NH_4Cl and extracting with EtOAc, the combined organic extracts were washed with water, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 gave 4-((*R*)-2-hydroxy-4-oxo-pentanoyl)-piperazine-1-carboxylic acid benzyl ester as a colorless oil (0.9 g, 11%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.39–7.31 (m, 5H), 5.15 (s, 2H), 4.79–4.74 (m, 1H), 4.00 (d, J = 7.8 Hz, 1H), 3.75–3.45 (m, 8H), 2.85 (dd, J = 16.6, 7.3 Hz, 1H), 2.63 (dd, J = 16.6, 3.4 Hz, 1H), 2.26 (s, 3H); LC/MS: *m/z* 335.1 $(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00827] (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentane-1,4-dione


[00828]



[00829] A mixture of 4-((*R*)-2-hydroxy-4-oxo-pentanoyl)-piperazine-1-carboxylic acid benzyl ester (0.13 g, 0.39 mmol) and MeOH (4 mL) was stirred with 10 mg of Pd/C (10% wt Pd on carbon)under H_2 atmosphere at ambient pressure overnight. After filtration and evaporation of the solvent, the residue was taken up in CH_2Cl_2, and 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.11 g, 0.39 mmol) and triethylamine (0.11 mL, 0.78 mmol) were added. The reaction mixture was stirred at room temperature overnight, washed with water, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 provided (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)pentane-1,4-dione as a yellow solid (91 mg, 54%). LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00830] (Pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate

[00831]



[00832] (Pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate.

[00833]




[00834] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.16 mmol) , (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (67 mg, 0.32 mmol), and triethylamine (44.6 μL, 0.32 mmol) in DMSO (500 μL) was heated in a microwave synthesizer at 200 °C for 10 minutes. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (pyridin-4-yl)methyl 4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 456.5 $(M+H)^+$ at 2.02 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00835] Example 153


[00836] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butan-1-one**

[00837]



[00838] **1-(4-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butane-1,2-dione**

[00839]




[00840] A mixture of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (200 mg, 0.63 mmol), sodium 4-methylsulfanyl-2-oxo-butyrate (160 mg, 0.94 mmol), BOP (414 mg, 0.94 mmol), and triethylamine (348 μL, 2.5 mmol) in 2.1 mL of CH_2Cl_2 was stirred at room

temperature for 1 h. After adding saturated $NaHCO_3$ solution, the mixture was extracted with CH_2Cl_2. The organic extracts were dried over Na_2SO_4 and concentrated to give 1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butane-1,2-dione. LC/MS: *m/z* 451.2 $(M+H)^+$ at 3.10 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00841] 2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butan-1-one

[00842]



[00843] $NaBH_4$ (34 mg, 0.88 mmol) was added to 1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butane-1,2-dione (200 mg, 0.44 mmol) in 1.5 mL of MeOH, and the reaction mixture was stirred at 0 °C for 20 minutes The reaction mixture was allowed to warm to room temperature, saturated $NaHCO_3$ was added, and the aqueous layer was extracted with CH_2Cl_2. The organic layer was dried over Na_2SO_4 and concentrated. Purification using preparative reverse phase HPLC with 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-(methylthio)butan-1-one as the TFA salt. (LC/MS: *m/z* 453.4 $(M+H)^+$ at 2.73 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00844] Example 154


[00845] (*R*)-3-Hydroxy-4-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-oxobutanamide

MISSING PAGE
UPON FILING

[00852]

[00853] **2-(4-((*S*)-4-Benzyl-2-isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol**


[00854]



[00855] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (200 mg, 0.74 mmol) in 10 mL DMF was added (*S*)-1-benzyl-3-isopropylpiperazine followed by the addition of triethylamine (206 μL). The reaction was heated at 85 °C for two hours. The reaction was quenched with water after cooling it to room temperature. The aqueous layer was extracted twice with CH_2Cl_2, and the combined extracts were dried over $MgSO_4$, filtered, and concentrated. The reaction was purified via silica gel chromatography using 1:1 hexanes:CH_2Cl_2 solvent system to yield 2-(4-((*S*)-4-benzyl-2-isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol (230 mg, 64%). LC/MS: *m/z* 453.5 $(M+H)^+$ at 2.61 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00856] **2-(4-((*S*)-2-Isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol**

[00857]




[00858] 20 mg of Pd/C was added to a round-bottom flask, and the flask was flushed with nitrogen followed by evacuation of the atmosphere under vacuum. To the flask was then added a solution of 2-(4-((*S*)-4-benzyl-2-isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol (200 mg, 0.44 mmol) in methanol, followed by the addition of ammonium formate (32 mg, 0.88 mmol). The reaction was refluxed overnight. The reaction was filtered through a bed of Celite

to remove the catalyst. The solvent was evaporated to yield 2-(4-((*S*)-2-isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol (126 mg). LC/MS: *m/z* 363.5 $(M+H)^+$ at 2.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00859] (*R*)-2-Hydroxy-1-((*S*)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)-3-isopropylpiperazin-1-yl)-4-methylpentan-1-one


[00860]



[00861] To a solution of 2-(4-((*S*)-2-isopropylpiperazin-1-yl)-7-methylquinazolin-2-yl)phenol (70 mg, 0.19 mmol) in DMF (0.5 mL) was added (*R*)-2-hydroxy-4-methylpentanoic acid (32.6 mg, 0.247 mmol). It was followed by the addition of triethylamine (52 μL) and a solution of HATU (94 mg) in 0.5 mL DMF at room temperature. The reaction was complete in an hour. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-2-hydroxy-1-((*S*)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)-3-isopropylpiperazin-1-yl)-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 477.5 $(M+H)^+$ at 2.96 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00862] Example 156


[00863] (Pyridin-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate



[00864]

[00865] **(Pyridin-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00866]




[00867] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (50 mg, 0.15 mmol), (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (53 mg, 0.26 mmol), triethylamine (30.4 mg, 0.3 mmol) and DMSO (1 mL) were stirred for 18 hours at room temperature. The reaction was purified by preparative reverse phase HPLC to give (pyridin-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 474.30 $(M+H)^+$ at 1.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00868] Example 157


[00869] **3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester**

[00870]



[00871] **3-Hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester**


[00872]



[00873] To 2-(4-chloro-7-methyl-quinazolin-2-yl)-phenol (245 mg, 0.91 mmol) in 3.0 mL of CH_2Cl_2 was added sequentially 3-hydroxymethyl-piperazine-1-carboxylic acid benzyl ester (336.3 mg, 1.34 mmol) and triethylamine (190 mL, 1.37 mmol), and the reaction mixture was heated at 40 °C for 6 h. The reaction mixture was cooled and extracted with water (2 x 10 mL) and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure to give an oil. The residue was purified by normal phase LC (20–85% EtOAc–hexanes) to give 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester (216 mg, 64% yield). LC/MS: *m/z* 485.4 (M+H)+ at 3.02 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00874] Benzyl 3-(hydroxymethyl)-4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate hydrochloride

[00875]




[00876] A 2.0 M HCl solution in Et_2O (212 μL, 0.42 mmol) was slowly added at room temperature to a stirring solution of 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester (206 mg, 0.42 mmol) in 500 μL of CH_2Cl_2. The reaction was stirred for 1 hour. Solvents were removed under reduced pressure, and the residue was triturated with Et_2O and filtered to give benzyl 3-(hydroxymethyl)-4-(2-(2-

hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate hydrochloride. LC/MS: *m/z* 485.5 (M+H)+ at 3.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[00877] Example 158


[00878] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)but-3-yn-1-one**

[00879]



[00880] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)but-3-yn-1-one**

[00881]




[00882] To a mixture of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (75 mg, 0.23 mmol) in 0.78 ml CH_2Cl_2 were added successively 2-hydroxybut-3-ynoic acid (30 mg, 0.30 mmol), BOP (134 mg, 0.30 mmol), and triethylamine (36 mg, 49 μL, 0.35 mmol). The mixture was stirred at 0 °C for 30 minutes. Purification via silica gel chromatography using 0–100% ethyl acetate/hexanes gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)but-3-yn-1-one. (LC/MS: *m/z* 403.5 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00883] Example 159


[00884] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)hexan-1-one**

[00885]



[00886] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)hexan-1-one**

[00887]




[00888] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to 2-hydroxyhexanoic acid (39.3 mg, 0.297 mmol). Triethylamine (63 μL) was added at room temperature, then a solution of HATU (113 mg) in 0.5 mL DMF. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one. LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.60 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00889] Example 160


[00890] **(Tetrahydro-2*H*-pyran-2-yl)methyl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate**

[00891]



[00892] **(Tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate**


[00893]



[00894] A solution of (tetrahydro-2*H*-pyran-2-yl)methanol (1 g, 8.60 mmol) and di(1*H*-imidazol-1-yl)methanone (2.8 g, 17.2 mmol) in 17 mL CH_2Cl_2 was heated overnight at 50 °C The reaction was quenched with water, extracted with CH_2Cl_2, dried over $MgSO_4$, filtered, and concentrated. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.15 (t, J = 0.9 Hz, 1H), 7.44 (t, J = 1.4 Hz, 1H), 7.07 (dd, J = 1.6, 0.8 Hz, 1H), 6.95 (m, 2H), 6.84 (m, 1H), 6.01 (s, 2H), 5.33 (s, 2H)

[00895] **(Tetrahydro-2*H*-pyran-2-yl)methyl 4-(6-fluoro-2-(2-hydroxyphenyl) quinazolin-4-yl)piperazine-1-carboxylate**

[00896]




[00897] 2-(6-Fluoro-4-(piperazin-1-yl)quinazolin-2-yl)phenol (25 mg, 0.077 mmol), (tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate (48.5 mg, 0.23 mmol), triethylamine (22μL, 0.154 mmol), and HATU (38 mg, 0.10 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (tetrahydro-2*H*-pyran-2-yl)methyl 4-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 467.3 $(M+H)^+$ at 3.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00898] Example 162


[00899] **(Tetrahydrofuran-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00900]



[00901] **4-Benzyl-piperazine-1-carboxylic acid tetrahydro-furan-3-ylmethyl ester**

[00902]




[00903] To a cooled (0 °C) solution of triphosgene (5.0 g, 17 mmol) in 50 ml dichloromethane under a nitrogen atmosphere was added dropwise a solution of tetrahydro-3-furanmethanol (5.4 g, 53 mmol) in 10 ml dichloromethane. Pyridine (4.3 ml, 53 mmol) was added dropwise, and the solution was warmed to room temperature. After 2 hours at room temperature a mixture of triethylamine (7.5 ml, 52 mmol) and *N*-benzylpiperazine (9.5 ml, 54

mmol) was added dropwise with cooling. The resulting mixture was refluxed for 1 hour. The solution was stirred at room temperature overnight under an nitrogen atmosphere. The mixture was washed with aqueous sodium bicarbonate solution (5%, 2 x 50 ml), and with a solution of saturated aqueous NaCl (50 ml). The organic layer was dried over sodium sulfate, filtered, and evaporated to dryness. The residue was co-evaporated three times with toluene (50 ml) to yield 4-benzyl-piperazine-1-carboxylic acid tetrahydro-furan-3-ylmethyl ester (13.0 g, 81%) as a brownish oil.

[00904] Piperazine-1-carboxylic acid tetrahydro-furan-3-ylmethyl ester.


[00905]



[00906] 4-Benzyl-piperazine-1-carboxylic acid tetrahydro-furan-3-ylmethyl ester (11.0 g) was dissolved in 100 ml ethanol. Palladium on carbon (10% Pd/C, 0.5 g) was added, and a hydrogen atmosphere was applied overnight at room temperature. The solution was filtered through Celite to remove the catalyst, and the Celite cake was rinsed with 50 ml ethanol. The combined filtrates were evaporated to dryness to yield piperazine-1-carboxylic acid tetrahydro-furan-3-ylmethyl ester (8.0 g) as a colorless oil.

[00907] (Tetrahydrofuran-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate

[00908]




[00909] 2-(4-Chloro-7-methylquinazolin-2-yl)-3-fluorophenol (50 mg, 0.174 mmol) was dissolved in 1 mL DMF, followed by the addition of triethylamine (35.2 mg, 0.348 mmol). (Tetrahydrofuran-3-yl)methyl piperazine-1-carboxylate (45 mg, 0.21 mmol) was then added. After 1 hour at room temperature, the reaction was complete. It was filtered and purified by reverse phase preparative HPLC to give (tetrahydrofuran-3-yl)methyl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 467.3 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00910] Example 163


[00911] **(*S*)-Tetrahydrofuran-3-yl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00912]



[00913] **(*S*)-Tetrahydrofuran-3-yl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[00914]




[00915] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.21 mmol) was dissolved in 1 mL anhydrous DMF under an N_2 atmosphere and cooled to 0 °C. (*S*)-Tetrahydrofuran-3-yl chloroformate (34.3 mg, 0.228 mmol) was dissolved in 150 μL anhydrous DMF and added dropwise to the reaction, followed by triethylamine (42 mg, 0.41 mmol). After 1 hour the reaction was complete, and it was filtered and purified by reverse phase HPLC to give (*S*)-tetrahydrofuran-3-yl 4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazine-1-carboxylate as the TFA salt. LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.25 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00916] Example164


[00917] **(*R*)-2,6-Dihydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-6-methylheptane-1,4-dione**

[00918]



[00919] **4-((*R*)-2,6-Dihydroxy-6-methyl-4-oxo-heptanoyl)-piperazine-1-carboxylicacid benzyl ester**


[00920]



[00921] To a solution of 4-((*R*)-2-hydroxy-3-methoxycarbonyl-propionyl)-piperazine-1-carboxylic acid benzyl ester (5.26 g, 15.0 mmol) in THF (150 mL) cooled in a dry ice acetone bath was added 1.4 M MeMgBr (32 mL, 45 mmol). The reaction was allowed to slowly warm to room temperature overnight. After quenching the mixture with saturated NH_4Cl and extracting with EtOAc, the combined organic extracts were washed with water, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 gave 4-((*R*)-2-hydroxy-4-oxo-pentanoyl)-piperazine-1-carboxylic acid benzyl ester as a colorless oil (0.79 g, 15%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.40–7.31 (m, 5H), 5.15 (s, 2H), 4.71–4.68 (m, 1H), 4.05 (d, J = 6.8 Hz, 1H), 3.80–3.73 (m, 1H), 3.62–3.42 (m, 6H), 3.31–3.23 (m, 1H), 1.72–1.61 (m, 2H), 1.35 (s, 3H), 1.30 (s, 3H); LC/MS: *m/z* 351.3 $(M+H)^+$ at 2.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00922] **(*R*)-2,4-Dihydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00923]




[00924] A mixture of 4-((*R*)-2,6-dihydroxy-6-methyl-4-oxo-heptanoyl)-piperazine-1-carboxylicacid benzyl ester (0.79 g, 2.20 mmol) and MeOH (25 mL) was stirred with 40 mg of Pd/C (10% wt Pd on carbon) under H_2 atmosphere at ambient pressure overnight. After filtration and evaporation of the solvent, the residue was taken up in CH_2Cl_2, and 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.61 g, 2.20 mmol) and triethylamine (0.63 mL, 4.50 mmol) were added. The reaction mixture was stirred at room temperature overnight and then diluted with CH_2Cl_2, washed with water, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–10% MeOH/CH_2Cl_2 gave (*R*)-2,6-dihydroxy-1-(4-(2-(2-

hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-6-methylheptane-1,4-dione as a yellow solid (372 mg, 37%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.45 (dd, J = 7.9, 1.6 Hz, 1H), 7.75 (d, J = 8.5 Hz, 1H), 7.67 (s, 1H), 7.40–7.36 (m, 1H), 7.29 (dd, J = 8.5, 1.4 Hz, 1H), 7.03 (d, J = 8.2 Hz, 1H), 6.96–6.92 (m, 1H), 4.81–4.76 (m, 1H), 4.09 (d, J = 6.8 Hz, 1H), 4.03–3.77 (m, 6H), 3.75–3.67 (m, 2H), 3.28 (s, 1H), 2.55 (s, 3H), 1.78–1.69 (m, 2H), 1.40 (s, 3H), 1.33 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 173.1, 164.0, 161.3, 160.5, 150.1, 144.5, 132.7, 129.2, 127.7, 126.8, 124.4, 119.3, 118.5, 117.7, 112.7, 70.6, 66.4, 49.9, 49.0, 46.1, 44.5, 42.4, 30.6, 29.2, 21.9; LC/MS: *m/z* 451.1 $(M+H)^+$ at 2.12 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00925] Example 165


[00926] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one**

[00927]



[00928] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one**

[00929]




[00930] A solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.22 mmol) in DMF (0.5 mL) was added to 2-hydroxy-2-methylbutanoic acid (33.6 mg, 0.284 mmol). Triethylamine (61 μL) was added, followed by a solution of HATU (108 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one as the TFA salt. LC/MS: *m/z* 421.3 $(M+H)^+$ at 2.40 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00931] Example 166


(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one

[00932]



[00933] **(*S*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one**

[00934]




[00935] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol hydrochloride (30 mg, 0.09 mmol), (*S*)-2-hydroxy-3,3-dimethylbutanoic acid (16 mg, 0.12 mmol), triethylamine (37.5μL, 0.27 mmol), and HATU (45.6 mg, 0.12 mmol) were stirred in DMF (1 mL) overnight.

Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one as the TFA salt. LC/MS: *m/z* 434.53 $(M+H)^+$ at 2.61 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00936] Example 167


[00937] **1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-(4-fluorophenyl)-2-hydroxyethanone**

[00938]



[00939] **1-(4-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-(4-fluorophenyl)-2-hydroxyethanone**

[00940]




[00941] 3-Fluoro-2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol), 2-(4-fluorophenyl)-2-hydroxyacetic acid (19.62 mg, 0.12 mmol), triethylamine (25 μL, 0.18 mmol) and HATU (45.6 mg, 0.12 mmol) were stirred in DMF (1 mL) overnight. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 1-(4-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-(4-fluorophenyl)-

2-hydroxyethanone as the TFA salt. LC/MS: *m/z* 491.3 $(M+H)^+$ at 2.46 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00942] Example 168


[00943] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00944]



[00945] **(*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one**

[00946]




[00947] Under an N_2 atmosphere, BOP (138 mg, 0.31 mmol) was added in a single portion to a stirring solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (100 mg, 0.31 mmol), (*R*)-2-hydroxy-4-methylpentanoic acid (41 mg, 0.31 mmol), and triethylamine (43 µL, 0.31 mmol) in DMF (0.5 ml). After stirring the mixture for 1 h at room temperature, it was partitioned between H_2O and ether. The organic phase was washed with H_2O (3x 20 mL), dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 20%

EtOAc and 80% CH_2Cl_2 gave (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one as a light yellow solid (89 mg, 60%). LC/MS: *m/z* 435.1 (M+H)+ at 2.93 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, $CDCl_3$) δ 8.46 (dd, *J* = 7.9, 1.7 Hz, 1H), 7.76 (d, *J* = 8.5 Hz, 1H), 7.66 (s, 1H), 7.41–7.37 (m, 1H), 7.30 (dd, *J* = 8.5, 1.5 Hz, 1H), 7.06–7.04 (m, 1H), 6.98–6.94 (m, 1H), 4.51–4.45 (m, 1H), 4.04–3.80 (m, 6H), 3.72–3.62 (m, 3H), 2.56 (s, 3H), 2.12–1.99 (m, 1H), 1.57–1.49 (m, 1H), 1.38–1.32 (m, 1H), 1.05 (d, *J* = 6.6 Hz, 3H), 1.00 (d, *J* = 6.7 Hz, 3H).

[00948] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride


[00949]



[00950] (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one (90 mg, 0.20 mmol) was dissolved in 1 mL CH_2Cl_2 and treated with 1 equivalent of 2.0 M HCl in ether (100 μL, 0.20 mmol). The formed precipitate was filtered and vacuum dried to obtain (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one hydrochloride. LC/MS: *m/z* 435.1 (M+H)+ at 2.84 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, CD_3OD) δ 8.39–8.37 (m, 1H), 8.18 (d, *J* = 8.6 Hz, 1H), 7.78 (s, 1H), 7.61–7.55 (m, 2H), 7.15–7.11 (m, 2H), 4.57 (dd, *J* = 9.5, 3.5 Hz, 1H), 4.49–4.32 (m, 4H), 4.07–3.79 (m, 4H), 2.62 (s, 3H), 1.97–1.87 (m, 1H), 1.65–1.58 (m, 1H), 1.53–1.47 (m, 1H), 1.03–1.00 (m, 6H).

[00951] (*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one sulfate


[00952]



[00953] To a stirring yellow solution of (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one (200 mg, 0.46 mmol) and THF (0.9 mL) under an N_2 atmosphere was added a solution of concentrated H_2SO_4 solution (95.9%) (26 μL, 0.46 mmol) and CH_3CN (0.75 mL) in a single portion. A white precipitate formed slowly over a period of 1 h. The solid was filtered and vacuum dried to obtain (*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-4-methylpentan-1-one sulfate as a white solid (229 mg, 94%). LC/MS: *m/z* 435.3 (M+H)+ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.20 (dd, *J* = 7.9, 1.5 Hz, 1H), 8.12 (d, *J* = 8.6 Hz, 1H), 7.77 (s, 1H), 7.53–7.49 (m, 2H), 7.10–7.03 (m, 2H), 4.39–4.35 (m, 1H), 4.30–4.04 (m, 4H), 3.93–3.69 (m, 4H), 2.55 (s, 3H), 1.86–1.73 (m, 1H), 1.52–1.33 (m, 2H), 0.93–0.91 (m, 6H).

[00954] Example 169


[00955] **(*R*)-[Methyl-(2-methylamino-acetyl)-amino]-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester**



[00956]

[00957] **{[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid**

[00958]




[00959] To (*tert*-butoxycarbonyl-methyl-amino)-acetic acid (2.73 g, 14.4 mmol) in 70 mL of THF was added sequentially diisopropyl ethylamine (7.5 mL, 14.4 mmol) and HBTU (5.47 g, 14.4 mmol), and the reaction mixture was stirred at room temperature for one hour. To this reaction mixture was added methylamino-acetic acid ethyl ester (2.22 g, 14.4 mmol) in one portion, and the reaction mixture was heated at 65 °C for 12 h. The reaction mixture was cooled, diluted with a solution of saturated $NaHCO_3$, and extracted with EtOAc. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid ethyl ester.

The ester was taken up in 49 mL of methanol, and to this was added a 1 M NaOH solution (25.1 mL, 25 mmol). The reaction mixture was heated at 65 °C for 3 h. The reaction mixture was cooled, and methanol was removed under reduced pressure. The residue was diluted with 50 mL of water and acidified with glacial acetic acid until the pH was 5. This solution was extracted with EtOAc (50 mL), and the organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give {[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid as an oil.

[00960] **4-(2(*R*)-Hydroxy-4-methyl-pentanoyl)-piperazine-1-carboxylic acid benzyl ester**

[00961]




[00962] To (*R*)- 2-hydroxy-4-methyl-pentanoic acid (1.71 g, 12.9 mmol) in 48 mL of DMF was added sequentially triethylamine (4.9 mL, 35. mmol), HOBt (1.75 g, 12.9 mmol), EDCI·HCl (2.48 g, 12.9 mmol) and piperazine-1-carboxylic acid benzyl ester (2.59 g, 11.8 mmol) at room temperature. The reaction mixture was stirred for 4 h and then diluted with EtOAc (50 mL) and extracted with 50 mL of water. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue

was subjected to purification by normal phase LC using 40–85% EtOAc–hexanes to give (1.92 g, 49% yield) of 4-(2(*R*)-hydroxy-4-methyl-pentanoyl)-piperazine-1-carboxylic acid benzyl ester the desired product. LC/MS: *m/z* 335.4 $(M+H)^+$ at 2.96 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00963] 4-[2-(2-{[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetoxy)-4-methyl-pentanoyl]-piperazine-1-carboxylic acid benzyl ester

[00964]



[00965] To {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid (427 mg, 1.57 mmol) in 2 mL of THF was added sequentially HBTU (624 mg, 1.2 mmol), diisopropyl ethylamine (860 μL, 4.71 mmol) at room temperature. After 5 min, 4-(2-hydroxy-4-methyl-pentanoyl)-piperazine-1-carboxylic acid benzyl ester (403 mg 1.2 mmol) in 2 mL of THF was added to the reaction mixture, and the solution was heated at 65 °C for 12 h. The reaction mixture was cooled and diluted with 20 mL of CH_2Cl_2 and 10 mL of water. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue was subjected to purification by normal phase LC using 30–100% EtOAc–hexanes to give 4-[2-(2-{[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetoxy)-4-methyl-pentanoyl]-piperazine-1-carboxylic acid benzyl ester (283 mg, 41% yield). LC/MS: *m/z* 577 $(M+H)^+$ at 3.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[00966] {[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester.



[00967] To 4-[2-(2-{[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetoxy)-4-methyl-pentanoyl]-piperazine-1-carboxylic acid benzyl ester (208 mg, 0.49 mmol) in 1.6 mL of methanol was added 70 mg of Pd/C (10% wt Pd on carbon). The reaction mixture was hydrogenated using a balloon of H_2 for 4 h at room temperature. The mixture was then filtered through Celite, and the solvent removed to give {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid 3-methyl-1-(piperazine-1-carbonyl)-butyl ester.

The amine was taken up in 1 mL CH_2Cl_2 and treated with 2-(4-chloro-7-methyl-quinazolin-2-yl)-phenol (53 mg, 0.19 mmol) and 49 μL of triethylamine, and stirred at room temperature for 3 h. The reaction mixture was diluted with 10 mL of CH_2Cl_2 and 10 mL of water. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester. LC/MS: *m/z* 677.4 $(M+H)^+$ at 3.25 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00968] [Methyl-(2-methylamino-acetyl)-amino]-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester.




[00969]

[00970] To {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester (42 mg, 0.062 mmol) was added 300 μL of 1.25 M HCl in methanol at room temperature, and the reaction mixture was stirred for 12 h. The solvent was removed, and the residue was purified with reverse phase LC using 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) as eluent to give [methyl-(2-methylamino-acetyl)-amino]-acetic acid 1-{4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carbonyl}-3-methyl-butyl ester as the TFA salt. LC/MS: *m/z* 577.4 (M+H)+ at 2.50 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00971] Example 170


[00972] **2-(4-Fluorophenyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00973]



[00974] **2-(4-Fluorophenyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00975]




[00976] 2-(7-Methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol hydrochloride (prepared analogously to 2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol, oxalate salt, see Example 130; 30 mg, 0.09 mmol), 2-(4-fluorophenyl)-2-hydroxyacetic acid (20 mg, 0.12 mmol), triethylamine (37.5 μL, 0.27 mmol) and HATU (45 mg, 0.12 mmol) were stirred in DMF (1 mL) overnight. Purification via preparative reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-(4-fluorophenyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone as the TFA salt. LC/MS: *m/z* 473.1 $(M+H)^+$ at 2.63 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00977] Example 171


[00978] **2-(2-Isopropyl-5-methylcyclohexyloxy)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00979]



[00980] **2-(2-Isopropyl-5-methylcyclohexyloxy)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone**

[00981]




[00982] To a solution of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) was added of triethylamine (25 μL) followed by the dropwise addition of 2-(2-isopropyl-5-methylcyclohexyloxy)acetyl chloride (21 μL, 0.09 mmol) at 0 °C. The reaction was warmed to room temperature, and purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-(2-isopropyl-5-methylcyclohexyloxy)-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)ethanone as the TFA salt. LC/MS: *m/z* 517.5 $(M+H)^+$ at 3.49 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00983] Example 172


[00984] **(*R*)-2-Hydroxy-1-{3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one**

[00985]



[00986] **(*R*)-2-Hydroxy-1-{3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one**


[00987]



[00988] To 3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazine-1-carboxylic acid benzyl ester (200 mg, 0.41 mmol) in 1.7 mL of methanol was added 39 mg of Pd /C (10% weight Pd on carbon). The reaction mixture was subjected to hydrogenation using a H_2 balloon for 3 h. The reaction mixture was filtered through Celite and the solvent was removed to give 2-[4-(2-hydroxymethyl-piperazin-1-yl)-7-methyl-quinazolin-2-yl]-phenol. This amine was treated with (*R*)-2-hydroxy-4-methyl-pentanoic acid (60 mg, 0.45 mmol), BOP (200 mg, 0.45 mmol) and 115 μL of triethylamine in 1.6 mL of DMF at room temperature for 12 h. The reaction mixture was diluted with 20 mL of CH_2Cl_2 and 20 mL of water, and the organic layer was separated and dried over Na_2SO_4. The solvent was removed under reduced pressure, and the residue was subjected to purification using 60–100% EtOAc–hexanes to give (*R*)-2-hydroxy-1-{3-hydroxymethyl-4-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperazin-1-yl}-4-methyl-pentan-1-one. LC/MS: *m/z* 465 (M+H)+ at 2.77 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00989] Example 173


[00990] **(*R*)-3-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**



[00991]

[00992] **(*R*)-3-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one**

[00993]




[00994] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to (*R*)-3-hydroxybutanoic acid (31 mg, 0.30 mmol). Then, triethylamine (63 μL) was added, followed by a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-3-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)butan-1-one as the TFA salt. LC/MS: *m/z* 393.1 $(M+H)^+$ at 2.03 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[00995] Example 174


[00996] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)pentan-1-one**

[00997]



[00998] **2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)pentan-1-one**

[00999]




[001000] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to 2-hydroxypentanoic acid (35 mg, 0.30 mmol). This was followed by the addition of triethylamine (63 μL) and a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)-2-methylbutan-1-one as the TFA salt. LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.41 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001001] Example 175


[001002] **1-(4-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(piperidin-1-yl)propan-1-one**

[001003]



[001004] **1-(4-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-(piperidin-1-yl)propan-1-one**

[001005]




[001006] To 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (53 mg, 0.17 mmol) was added sequentially 3-(piperidin-1-yl)propanoyl chloride (33 mg, 0.19 mmol) in 28 μL of CH_2Cl_2 and triethylamine (28 μL, 0.2 mmol). The mixture was stirred at 0 °C for 20 minutes After adding H_2O and CH_2Cl_2, the phases were separated, and the organic layer was dried over Na_2SO_4 and concentrated under vacuum. Purification using preparative reverse phase HPLC with 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) gave 3-cyclopentyl-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 460.5 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001007] Example 176


[001008] **(2*R*,3*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylpentan-1-one**

[001009]



[001010] **(2*R*,3*R*)-2-Hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylpentan-1-one**

[001011]




[001012] To 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (80 mg, 0.25 mmol) in 800 μL of CH_2Cl_2 was added sodium of (2*R*,3*R*)-2-hydroxy-3-methyl-pentanoate (50 mg, 0.33 mmol), BOP (144 mg, 0.33 mmol), and triethylamine (52 μL, 0.38 mmol). The reaction mixture was stirred at room temperature for 2 h. After adding H_2O and CH_2Cl_2, the layers were separated, and the organic layer was dried over Na_2SO_4 and concentrated. Purification via preparative reverse phase HPLC using 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) gave (2*R*,3*R*)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-3-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.62 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001013] Example 177


[001014] **(*S*)-3,3,3-Trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001015]



[001016] **(*S*)-3,3,3-Trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001017]




[001018] A mixture of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (64 mg, 0.2 mmol), (*S*)-3,3,3-trifluoro-2-hydroxypropanoic acid (29 mg, 0.2 mmol), HATU (76 mg, 0.2 mmol) and triethylamine (28 μL, 0.2 mmol) in DMF (1 mL) was stirred at room temperature overnight. Purification via preparative HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-3,3,3-trifluoro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 447.1 $(M+H)^+$ at 2.53 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001019] Example 178


[001020] **2-(Trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl) quinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001021]



[001022] **2-(Trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl) quinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001023]




[001024] A solution of 2-(4-(piperazin-1-yl)quinazolin-2-yl)phenol (70 mg, 0.23 mmol) in DMF (0.5 mL) was added to 2-(trifluoromethyl)-2-hydroxypropanoic acid (47.0 mg, 0.297 mmol). Then, triethylamine (63 μL) was added, followed by a solution of HATU (113 mg) in 0.5 mL DMF at room temperature. The reaction was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-(trifluoromethyl)-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 447.3 $(M+H)^+$ at 2.50 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001025] Example 179


[001026] **3-Chloro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001027]



[001028] **3-Chloro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one**

[001029]




[001030] A mixture of 2-(7-methyl-4-(piperazin-1-yl)quinazolin-2-yl)phenol (121 mg, 0.38 mmol), 3-chloro-2-hydroxypropanoic acid (61 mg, 0.49 mmol), BOP (217 mg, 0.49 mmol), and triethylamine (79 μL, 0.56 mmol) in 1.2 mL of CH_2Cl_2 was stirred at room temperature for 1 h. The reaction was washed with water and the organic layer was dried over Na_2SO_4 and concentrated. Purification via preparative reverse phase HPLC using 10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) gave 3-chloro-2-hydroxy-1-(4-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)propan-1-one as the TFA salt. LC/MS: *m/z* 427.2 $(M+H)^+$ at 2.59 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001031] Example 201

[001032] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)piperidine-3-carboxamide**

[001033]



[001034] **Piperidine-3-carboxylic acid**


[001035]



[001036] To a solution of 1-(*tert*-butoxycarbonyl)piperidine-3-carboxylic acid (500 mg, 2.18 mmol) in CH_2Cl_2 (10 mL) was added TFA (5 ml). The reaction mixture was stirred for an hour. Excess TFA was removed under reduced pressure, and the piperidine-3-carboxylic acid was used without neutralization for the next step. LC/MS: *m/z* 1303 $(M+H)^+$ at 0.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001037] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxylic acid**

[001038]




[001039] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.449 mg, 1.66 mmol) in CH_2Cl_2 was added 5 equivalents of triethylamine followed by the addition of piperidine-3-carboxylic acid as a TFA salt. The reaction was stirred for 2 hours and quenched with water. The layers were separated, and the organic extracts were dried over $MgSO_4$, filtered,

and concentrated to give 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxylic acid which was used without further purification. LC/MS: *m/z* 364.3 $(M+H)^+$ at 2.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001040] 1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)piperidine-3-carboxamide

[001041]




[001042] A solution of 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxylic acid (45 mg, 0.12 mmol), (pyridin-3-yl)methanamine (14 μL, 0.136 mmol) and triethylamine (25 mg, 35 μL, 0.25 mmol) in 500 μL DMF was cooled to 0 °C, and HATU (57 mg, 0.15 mmol) was added. The reaction was warmed to room temperature, stirred overnight and purified by reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) giving 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)piperidine-3-carboxamide as the TFA salt. LC/MS: *m/z* 454.5 $(M+H)^+$ at 1.87 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001043] Example 203


[001044] (*R*)-Tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate

[001045]



[001046] Benzyl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate


[001047]



[001048] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (1.15 g, 4.26 mmol) in CH_2Cl_2 (25 mL) at 0 °C under inert atmosphere was slowly added a solution of benzyl (*S*)-piperidin-3-ylcarbamate (1.0 g, 4.26 mmol) and triethylamine (1.18 mL, 8.52 mmol) in CH_2Cl_2 (10 mL). The reaction was allowed to warm to room temperature and was then quenched with water. The mixture was extracted with CH_2Cl_2 and the organic layers were combined, dried over $MgSO_4$, and concentrated to obtain benzyl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate (2.03 g). This material was used without further purification. LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.86 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001049] 2-(4-((*S*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol

[001050]




[001051] Pd/C (175 mg, 10% weight Pd on carbon) was added to a round bottom flask and the flask was flushed with N_2. To this flask was then added MeOH (10 mL). After purging the flask again with N_2, (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate (1.75 g, 3.74 mmol) dissolved in EtOAc (60 mL) and MeOH (50 mL) was added. After flushing the flask 3 times with N_2, and evacuating it under vacuum, H_2 was passed through

the vigorously stirring mixture for 4 h until hydrogenation was complete. The mixture was filtered through a pad of Celite. The filtrate was concentrated to afford 2-(4-((*S*)-3-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.62 g, 50%). LC/MS: *m/z* 335.5$(M+H)^+$ at 1.50 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001052] (*R*)-Tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate

[001053]




[001054] 2-(4-((*S*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol), (*R*)-tetrahydrofuran-3-yl chloroformate (22.6 mg, 0.15 mmol) and triethylamine (30 mg, 0.3 mmol) were stirred in DMF (1 mL) at 0 °C. After allowing the reaction to warm to room temperature, it was purified via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (*R*)-tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 449.3.5$(M+H)^+$ at 1.52 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001055] Example 204


[001056] (*S*)-Tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate

[001057]



[001058] **(*S*)-Tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**

[001059]




[001060] 2-(4-((*S*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol), (*S*)-tetrahydrofuran-3-yl chloroformate (22.6 mg, 0.15 mmol), and triethylamine (30 mg, 0.3 mmol) were stirred in DMF (1 mL) at 0 °C. After allowing the reaction to warm to room temperature it was purified via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (*S*)-tetrahydrofuran-3-yl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001061] Example 205


[001062] **(2*R*)-Tetrahydro-N-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide**

[001063]



[001064] **(2*R*)-Tetrahydro-*N*-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide**

[001065]




[001066] 2-(4-((*S*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol), (*R*)-tetrahydrofuran-2-carboxylic acid (22.6 mg, 0.195 mmol), triethylamine (30 mg, 0.3 mmol), and HATU (74.14 mg, 0.195 mmol) were stirred at room temperature in DMF for 1 h. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-tetrahydro-*N*-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 433.3$(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001067] Example 206

[001068] **(2*R*)-Tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide**

[001069]



[001070] ***tert*-Butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**




[001071]

[001072] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.5 g, 1.84 mmol) in CH_2Cl_2 (10 mL) at 0 °C was added dropwise a solution of *tert*-butyl (*R*)-piperidin-3-ylcarbamate (0.37 g, 1.84 mmol) in CH_2Cl_2 (5 mL), then triethylamine (0.51 mL, 3.68 mmol). The mixture was allowed to warm to room temperature and was stirred for 3 h. After quenching with water, it was extracted with CH_2Cl_2. The organic layers were combined, dried over $MgSO_4$, and concentrated. Purification via silica gel chromatography using 5:1 CH_2Cl_2: hexanes gave *tert*-butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate (0.54 g, 68%). LC/MS: *m/z* 435.5$(M+H)^+$ at 2.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001073] **2-(4-((*R*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol.**


[001074]



[001075] *tert*-Butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate (0.54 g, 1.24 mmol) was dissolved in CH_2Cl_2 (15 mL) followed by the addition of TFA (8 mL). The reaction was stirred for 1.5 h, and the solvents were evaporated to an oily liquid which was diluted with CH_2Cl_2 and neutralized with a 1 M aqueous NaOH solution. The organic layer was separated, and the aqueous layer was washed two times with CH_2Cl_2. After drying the combined organic phases over $MgSO_4$, they were filtered, and concentrated to afford 2-(4-((*R*)-3-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol as a solid (0.354 g, 85%). LC/MS: *m/z* 335.7$(M+H)^+$ at 1.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001076] **(2*R*)-Tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide**

[001077]




[001078] 2-(4-((*R*)-3-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol), (*R*)-tetrahydrofuran-2-carboxylic acid (22.6 mg, 0.15 mmol), triethylamine (30 mg, 0.3 mmol), and HATU (74.14 mg, 0.195 mmol) were stirred at room temperature in DMF for 1 h. Purification via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 433.3$(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001079] Example 207


[001080] **(*R*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**

[001081]



[001082] **(*R*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**

[001083]




[001084] A solution of 2-(4-((*R*)-3-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMF (0.5 mL) was cooled to 0 °C. Then, a solution of triethylamine (30 mg, 0.3 mmol) and (*R*)-tetrahydrofuran-3-yl chloroformate (22.6 mg, 0.15 mmol) in DMF (0.5 mL) was added dropwise. The mixture was allowed to warm to room temperature over a period of 30 min before it was purified using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*R*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 449.5$(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001085] Example 208


[001086] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**

[001087]



[001088] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate**

[001089]




[001090] To a solution of 2-(4-((*R*)-3-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMF (0.5 mL) was cooled to 0 °C. A solution of (*S*)-tetrahydrofuran-3-yl chloroformate (22.6 mg, 0.15 mmol) in DMF (0.5 mL) was then added, followed by triethylamine (30 mg, 0.3 mmol). The mixture was allowed to warm to room temperature over a period of 30 min before it was purified using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 449.5$(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001091] Example 209

[001092] **(2*S*)-Tetrahydro-*N*-((1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)furan-2-carboxamide**

[001093]




[001094] **3-(Benzyloxycarbonylamino-methyl)-piperidine-1-carboxylic acid *tert*-butyl ester**

[001095] 


[001096] 3-Aminomethyl-piperidine-1-carboxylic acid *tert*-butyl ester (3.6 g, 16.8 mmol) was dissolved in 42 mL anhydrous CH_2Cl_2 under an N_2 atmosphere and cooled in an ice water bath. Triethylamine (4.7 mL, 33.6 mmol) was added followed by the dropwise addition of benzyl chloroformate (3.55 mL, 25.2 mmol). After 16 hours, the reaction mixture was partitioned between CH_2Cl_2/H_2O,and separated, and the aqueous layer was extracted twice with CH_2Cl_2. All of the organic extracts were combined, dried over Na_2SO_4, filtered, and concentrated to a light yellow oil. Purification by silica gel chromatography using 97% CH_2Cl_2/3% MeOH gave the product as a clear colorless oil (55%). LC/MS: *m/z* 349.3 $(M+H)^+$ at 3.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001097] **Benzyl (piperidin-3-yl)methylcarbamate hydrochloride**

[001098] 

[001099] Benzyl (1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate (3.22 g, 9.25 mmol) was treated with a 4.0 M HCl solution in dioxane (11.3 mL, 46.25 mmol). Formation of a white precipitate was observed. After 3 h, the reaction was complete. The solvent and excess HCl were removed under reduced pressure to obtain benzyl (piperidin-3-yl)methylcarbamate hydrochloride as a white solid. LC/MS: *m/z* 249.3 $(M+H)^+$ at 1.28 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001100] **Benzyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**




[001101]

[001102] 2-(4-Chloro-7-methylquinazolin-2-yl)phenol (1.5 g, 5.54 mmol) was suspended in anhydrous CH_2Cl_2 (15 mL) and cooled to 0 °C under an N_2 atmosphere. A mixture of benzyl (piperidin-3-yl)methylcarbamate hydrochloride (1.73 g, 6.09 mmol) in CH_2Cl_2 (20 mL), and triethylamine (23 mL, 16.67 mmol) was added dropwise. The reaction mixture was allowed to warm to room temperature, and the reaction was complete after one hour. It was then partitioned between CH_2Cl_2/H_2O, and separated, and the organic phase was dried over Na_2SO_4, filtered, and concentrated to an orange solid. Purification by silica gel chromatography using 98% CH_2Cl_2/2% EtOAc gave benzyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as a bright yellow solid (1.75 g, 66%). LC/MS: *m/z* 483.5 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001103] **2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol**

[001104]




[001105] A mixture of benzyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (1.75 g, 3.63 mmol) and EtOH/EtOAc (50 mL/20 mL) was heated to obtain a homogeneous solution. After cooling to room temperature, Pd/C (175 mg, 10% wt Pd on carbon) was added and the flask was sealed with a septum. The same flask was 3 times charged with N_2 and evacuated under vacuum. The mixture was then stirred under an H_2

atmosphere at ambient pressure for 1 h. The product was collected by filtration through a plug of Celite, eluting with MeOH. The filtrate was concentrated to a yellow solid (1.26 g) to give 2-(4-(3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol. LC/MS: *m/z* 349.3 $(M+H)^+$ at 1.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001106] (2*S*)-Tetrahydro-*N*-((1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)furan-2-carboxamide


[001107]



[001108] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled in an ice water bath. To this was added (*S*)-tetrahydrofuran-2-carboxylic acid (24.0 mg, 19.8 μL, 0.2 mmol), and triethylamine (50 μL, 0.34 mmol). After 5 minutes, HATU (78.3 mg, 0.2 mmol) was added in one portion, and the reaction was allowed to warm to room temperature overnight. Purification by reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*S*)-tetrahydro-N-((1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 447.5 $(M+H)^+$ at 2.27 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001109] Example 210


[001110] (*R*)-Tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate



[001111]

[001112] (*R*)-Tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001113]




[001114] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in DMF (0.5 mL) and placed into an ice water bath. (*R*)-tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) was dissolved in DMF (0.1 mL) and added dropwise, followed by the addition of triethylamine (50 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001115] Example 211


[001116] (*S*)-Tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001117]



[001118] (*S*)-Tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001119]




[001120] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) dissolved in DMF (0.5 mL) was cooled to 0 °C. (*S*)-Tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) was dissolved in DMF (0.1 mL) and added dropwise to the reaction mixture, followed by the addition of triethylamine (50 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. The reaction was purified by reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*S*)-tetrahydrofuran-3-yl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001121] Example 212

[001122] ***N*-((1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide**

[001123]



[001124] ***N*-((1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide**




[001125]

[001126] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in DMF (0.5 mL) and cooled to 0 °C . Cyclopropanecarbonyl chloride (15.7 mg, 0.2 mmol) was dissolved in DMF (0.1 mL) and added dropwise, followed by the addition of triethylamine (50 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave *N*-((1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 417.0 $(M+H)^+$ at 2.30 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001127] Example 213


[001128] **(2*R*)-*N*-(1-(2-(2-Chloro-6-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide**

[001129]



[001130] **3-Chloro-2-(4-chloroquinazolin-2-yl)phenol**


[001131]



[001132] To a solution of 4-chloro-2-(2-chloro-6-methoxyphenyl)quinazoline (0.3 g, 0.98 mmol) in CH_2Cl_2 at –78 °C was added 5 equivalents of 1 M BBr_3 solution (4.9 mL, 4.9 mmol) in CH_2Cl_2. The reaction was complete after 30 minutes After allowing it to warm to room temperature, the reaction mixture was quenched with aqueous $NaHCO_3$ to pH 7, the layers were separated and the aqueous layer was extracted with CH_2Cl_2. The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 2:1 CH_2Cl_2:hexanes yielded 3-chloro-2-(4-chloroquinazolin-2-yl)phenol (0.17 g, 60%). LC/MS: *m/z* 291.1$(M+H)^+$ at 3.16
min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001133] **(2*R*)-*N*-(1-(2-(2-Chloro-6-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide**

[001134]




[001135] At 0 °C, to a solution of 3-chloro-2-(4-chloroquinazolin-2-yl)phenol (42 mg, 0.144 mmol) in CH_2Cl_2 was added triethylamine (80 μL, 0.58 mmol) followed by the addition of (*R*)-tetrahydro-*N*-(piperidin-4-yl)furan-2-carboxamide a oxalate. After allowing the reaction to warm to room temperature it was purified via reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*R*)-*N*-(1-(2-(2-chloro-6-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide as the TFA salt. LC/MS: *m/z* 453.5$(M+H)^+$ at 1.98 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001136] Example 214

[001137] **(*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**


[001138]



[001139] Method 1

[001140] **Benzyl ((*S*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate**


[001141]



[001142] (*S*)-*tert*-Butyl 3-(aminomethyl)piperidine-1-carboxylate (1.00 g, 4.67 mmol) was dissolved in 14 mL anhydrous CH_2Cl_2 under an N_2 atmosphere and cooled to 0 °C. Triethylamine (1.30 mL, 945 mg, 9.34 mmol) was added followed by the dropwise addition of benzyl chloroformate (0.99 mL, 1.20 g, 7.00 mmol). After 16 h, the reaction mixture was partitioned between H_2O and CH_2Cl_2,and separated, and the aqueous layer was extracted twice with CH_2Cl_2. The organic layers were combined, dried over Na_2SO_4, filtered, and concentrated to a light yellow oil. Purification via silica gel chromatography using 97% CH_2Cl_2/3% MeOH gave benzyl ((*S*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate as a clear colorless oil (895 mg, 55%). LC/MS: *m/z* 349.5 $(M+H)^+$ at 3.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001143] **Benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**


[001144]



[001145] Benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (895 mg, 2.57 mmol) was treated with a 4.0 M HCl solution in dioxane (3.2 mL, 12.85 mmol). Formation of a white precipitate was observed. After 3 h, complete conversion of starting material was seen by TLC. The solvent and excess HCl were removed under reduced pressure to obtain benzyl ((*R*)-piperidin-3-yl)methylcarbamate hydrochloride as a white solid.

This solid was suspended in DMF/CH_2Cl_2 (3 mL/3 mL), followed by the addition of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (696 mg, 2.57 mmol), and triethylamine (1.8 mL, 1.3 g, 12.85 mmol). The mixture was stirred at room temperature under an N_2 atmosphere for 16 h. The reaction was then partitioned between H_2O/CH_2Cl_2,and separated and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated under reduced pressure. Purification via silica gel chromatography using 0–5% MeOH in CH_2Cl_2 gave benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as a thick yellow oil (610 mg, 49%). LC/MS: *m/z* 483.3 $(M+H)^+$ at 2.83 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001146] **2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol**




[001147]

[001148] To a mixture of benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (610 mg, 1.26 mmol) and EtOH (15 mL) in a round bottom flask was added Pd/C (61 mg, 10% wt Pd on carbon) and the flask was sealed with a septum. The atmosphere in the flask was evacuated, purged with N_2, and equipped with a balloon charged with H_2. The mixture was then stirred under an H_2 atmosphere at ambient pressure for 3 h. After filtration through a plug of Celite, using MeOH as the eluting solvent, the reaction mixture was concentrated to a yellow solid 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (441 mg). LC/MS: *m/z* 349.3 $(M+H)^+$ at 1.52 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43–8.45 (m, 1H), 7.95 (d, *J* = 8.5 Hz, 1H), 7.64 (s, 1H), 7.35–7.39 (m, 2H), 6.92–6.96 (m, 2H), 4.52 (d, *J* = 12.5 Hz, 1H), 4.41 (d, *J* = 13.2 Hz, 1H), 3.26–3.29 (m, 2H), 3.02–3.08 (m, 1H), 2.55–2.61 (m, 1H), 2.45–2.48 (m, 1H), 1.84–1.91 (m, 2H), 1.65–1.77 (m, 3H), 1.24–1.36 (m, 1H).

[001149] (*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001150]



[001151] Method A

[001152] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, and (*R*)-tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) dissolved in DMF (100 μL) was added


dropwise followed by triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature. The reaction was complete after two hours. Purification using reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.32 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001153] Method B

[001154] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (127 mg, 0.364 mmol) was dissolved in 5 mL anhydrous DMF and cooled to 0 °C, and a solution of (*R*)-tetrahydrofuran-3-yl chloroformate (65.4 mg, 0.436 mmol) in 200 μL DMF was added dropwise followed by the addition of triethylamine (74 mg, 0.10 mL, 0.73 mmol). The reaction was allowed to warm to room temperature. The reaction was complete after two hours. The mixture was partitioned between H_2O and CH_2Cl_2, and separated, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic phases were dried over Na_2SO_4, filtered, and concentrated to a yellow solid. Purification via silica gel chromatography using 98% CH_2Cl_2/2% MeOH gave (*R*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as an off white solid (116 mg, 69%). LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.41–8.43 (m, 1H), 7.86 (d, *J* = 8.5 Hz, 1H), 7.65 (s, 1H), 7.33–7.44 (m, 3H), 6.90–6.95 (m, 2H), 5.11 (dd, *J* = 6.2, 4.6 Hz, 1H), 4.38 (t, *J* = 14.5 Hz, 2H), 3.68–3.79 (m, 3H), 3.60–3.63 (m, 1H), 3.28–3.31 (m, 1H), 2.98–3.07 (m, 3H), 2.50 (s, 3H), 2.06–2.15 (m, 1H), 1.85–1.91 (m, 4H), 1.70 (d, *J* = 12.9 Hz, 1H), 1.24–1.37 (m, 1H).

[001155] (*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001156]




[001157] (*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (116 mg, 0.251 mmol) was suspended in 8 mL anhydrous CH_2Cl_2 and gently heated until an homogenous solution was formed. After the reaction was cooled to room temperature, a 2.0 M solution of HCl in Et_2O (0.126 mL, 0.251 mmol) was added in one portion. The reaction mixture was diluted with 25 mL Et_2O, and the product precipitated from solution. The reaction was stirred for an additional 30 minutes before the solid was filtered and dried to obtain (*R*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride as a light yellow solid (99 mg, 70%). LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.22 (d, J = 7.7 Hz, 1H), 7.95 (d, J = 8.5 Hz, 1H), 7.75 (s, 1H), 7.42–7.49 (m, 3H), 6.97–7.08 (m, 2H), 5.08 (dd, J = 6.1, 4.6 Hz, 1H), 4.52–4.54 (m, 2H), 3.66–3.78 (m, 3H), 3.58–3.60 (m, 1H), 3.48 (t, J = 10.7 Hz, 1H), 3.23 (t, J = 11.5 Hz, 1H), 3.00 (t, J = 6.3 Hz, 2H), 2.53 (s, 3H), 2.05–2.14 (m, 1H), 1.80–1.91 (m, 4H), 1.72 (d, J = 9.0 Hz, 1H), 1.34–1.43 (m, 1H).

[001158] **Method 2**

[001159] ***tert*-Butyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001160]




[001161] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.478 g, 1.76 mmol) in CH_2Cl_2 was added triethylamine (0.98 mL, 0.712 g, 7.04 mmol), and the mixture was cooled to 0 °C. To the reaction mixture was added *tert*-butyl ((*R*)-piperidin-3-yl)methylcarbamate oxalate (prepared analogously to 2-(7-Methyl-4-piperazin-1-yl-quinazolin-2-yl)-phenol, oxalate salt, see Example 130; 700 mg, 2.3 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. The reaction was quenched with water, and the aqueous layer was extracted with CH_2Cl_2. The combined organic layers were dried over

$MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–2% EtOAc in CH_2Cl_2 gave *tert*-butyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (700 mg, 88%). LC/MS: *m/z* 449.5 $(M+H)^+$ at 2.77 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001162] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol

[001163]



[001164] To ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (700 mg, 1.56 mmol) was added 20 mL CH_2Cl_2 followed by addition of 7 mL of TFA. After the reaction was stirred for 1 hour it was neutralized with a1.0 M aqueous NaOH solution. The mixture was partitioned between H_2O and CH_2Cl_2, and separated, and the aqueous layer was extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to yield 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (400 mg, 74%). LC/MS: *m/z* 349.3 $(M+H)^+$ at 1.52 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43–8.45 (m, 1H), 7.95 (d, J = 8.5 Hz, 1H), 7.64 (s, 1H), 7.35–7.39 (m, 2H), 6.92–6.96 (m, 2H), 4.52 (d, J = 12.5 Hz, 1H), 4.41 (d, J = 13.2 Hz, 1H), 3.26–3.29 (m, 2H), 3.02–3.08 (m, 1H), 2.55–2.61 (m, 1H), 2.45–2.48 (m, 1H), 1.84–1.91 (m, 2H), 1.65–1.77 (m, 3H), 1.24–1.36 (m, 1H).

[001165] (*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate




[001166]

[001167] To a solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (210 mg, 0.6 mmol) in DMF at 0 °C were added (*R*)-tetrahydrofuran-3-yl chloroformate (0.09 g, 0.6 mmol) and triethylamine (167 μL, 1.2 mmol) simultaneously. Ten to fifteen minutes after addition, the reaction was complete, and quenched with water, extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to give (*R*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (150 mg, 54%). LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.41–8.43 (m, 1H), 7.86 (d, *J* = 8.5 Hz, 1H), 7.65 (s, 1H), 7.33–7.44 (m, 3H), 6.90–6.95 (m, 2H), 5.11 (dd, *J* = 6.2, 4.6 Hz, 1H), 4.38 (t, *J* = 14.5 Hz, 2H), 3.68–3.79 (m, 3H), 3.60–3.63 (m, 1H), 3.28–3.31 (m, 1H), 2.98–3.07 (m, 3H), 2.50 (s, 3H), 2.06–2.15 (m, 1H), 1.85–1.91 (m, 4H), 1.70 (d, *J* = 12.9 Hz, 1H), 1.24–1.37 (m, 1H).

[001168] Example 215


[001169] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001170]



[001171] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001172]



[001173] Method A

[001174] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, and (*S*)-tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) in DMF (100 μL) was added dropwise followed by triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature. The reaction was complete after two hours. Purification using preparative HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001175] Method B

[001176] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (126.8 mg, 0.364 mmol) was dissolved in 5 mL anhydrous DMF and cooled to 0 °C, and a solution of (*S*)-tetrahydrofuran-3-yl chloroformate (65.4 mg, 0.436 mmol) in 200 μl of DMF was added dropwise followed by the addition of triethylamine (74 mg, 0.102 mL, 0.728 mmol). The reaction was allowed to warm to room temperature and was complete after two hours. The mixture was partitioned between H_2O and CH_2Cl_2, separated, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic phases were dried over Na_2SO_4, filtered, and concentrated to a yellow solid. Purification via silica gel chromatography using 98% CH_2Cl_2/2% MeOH gave (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as an off white solid (116 mg, 69%). LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).^{1}H NMR (400 MHz, DMSO–d6) δ 8.42 (dd, J = 8.1, 1.7 Hz, 1H), 7.86 (d, J = 8.6 Hz, 1H), 7.66 (s, 1H), 7.34–7.43 (m, 3H), 6.91–6.95 (m, 2H), 5.10–5.11 (m, 1H), 4.38 (t, J = 12.4 Hz, 2H), 3.65–3.77 (m, 4H), 2.98–3.08 (m, 3H), 2.04–2.13 (m, 1H), 1.80–1.87 (m, 4H), 1.68–1.74 (m, 1H), 1.30–1.38 (m, 1H).


[001177] Method C

[001178] To a solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (175 mg, 0.5 mmol) in DMF at 0 °C were added (*S*)-tetrahydrofuran-3-yl chloroformate (75 mg, 0.5 mmol) and triethylamine (137 μL, 1.0 mmol) simultaneously. Ten to fifteen minutes after addition, the reaction was complete, and it was quenched with water and extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and

concentrated. Purification via silica gel chromatography using 98% CH_2Cl_2/2% MeOH gave (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (150 mg, 54%). LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.42 (dd, *J* = 8.1, 1.7 Hz, 1H), 7.86 (d, *J* = 8.6 Hz, 1H), 7.66 (s, 1H), 7.34–7.43 (m, 3H), 6.91–6.95 (m, 2H), 5.10–5.11 (m, 1H), 4.38 (t, *J* = 12.4 Hz, 2H), 3.65–3.77 (m, 4H), 2.98–3.08 (m, 3H), 2.04–2.13 (m, 1H), 1.80–1.87 (m, 4H), 1.68–1.74 (m, 1H), 1.30–1.38 (m, 1H).

[001179] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001180]




[001181] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (115 mg, 0.251 mmol) was suspended in 8 mL of anhydrous CH_2Cl_2 and gently heated until an homogenous solution was formed. After cooling to room temperature, a 2.0 M solution of HCl in Et_2O (0.126 mL, 0.251 mmol) was added in one portion. The reaction mixture was diluted with 25 mL Et_2O, and the product precipitated from the solution. The reaction was stirred for an additional 30 minutes before the solid was filtered and dried to obtain (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride (108 mg, 86%) as a light yellow solid. LC/MS: *m/z* 463.5$(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).^1H NMR (400 MHz, DMSO–d6) δ 8.18 (d, *J* = 6.6 Hz, 1H), 7.94 (d, *J* = 8.6 Hz, 1H), 7.73 (s, 1H), 7.44–7.51 (m, 2H), 7.00–7.07 (m, 2H), 5.06–5.09 (m, 1H), 4.52–4.62 (m, 2H), 3.62–3.74 (m, 4H), 3.23–3.29 (m, 1H), 3.00 (d, *J* = 6.8 Hz, 2H), 2.52 (s, 3H), 2.02–2.11 (m, 1H), 1.61–2.01 (m, 4H), 1.24–1.43 (m, 2H), 0.84–0.89 (m, 1H).

[001182] Example 216


[001183] (*R*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001184]



[001185] Benzyl ((*R*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate


[001186]



[001187] (*R*)-*tert*-Butyl 3-(aminomethyl)piperidine-1-carboxylate (1.00 g, 4.67 mmol) was dissolved in 14 mL anhydrous CH_2Cl_2 under an N_2 atmosphere and cooled to 0 °C. Triethylamine (1.30 mL, 945 mg, 9.34 mmol) was added followed by the dropwise addition of benzyl chloroformate (0.99 mL, 1.20 g, 7.00 mmol). After 16 h, the reaction was complete. The mixture was partitioned between H_2O and CH_2Cl_2, and separated, and the aqueous layer was extracted twice with CH_2Cl_2. The organic layers were combined, dried over Na_2SO_4, filtered, and concentrated to a light yellow oil. Purification via silica gel chromatography using 97% CH_2Cl_2/3% MeOH gave, benzyl ((*R*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate as a clear colorless oil (1.2 g, 74%). LC/MS: *m/z* 349.5 $(M+H)^+$ at 3.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001188] Benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate


[001189]



[001190] Benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (1.2 g, 3.54 mmol) was treated with 4.0 M HCl solution in dioxane (4.3 mL, 17.2 mmol). Formation of a white precipitate was observed. The reaction was complete after three hours. The solvent and excess HCl were removed under reduced pressure to obtain benzyl ((*S*)-piperidin-3-yl)methylcarbamate hydrochloride as a white solid. This solid was suspended in DMF/CH_2Cl_2 (3 mL/3 mL), followed by the addition of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (958 mg, 3.54 mmol) and then triethylamine (1.8 mL, 1.3 g, 12.85 mmol). The mixture was stirred at room temperature under an N_2 atmosphere for 16 h. The reaction was then partitioned between H_2O and CH_2Cl_2, separated, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated under reduced pressure. Purification via silica gel chromatography using 0–5% MeOH in CH_2Cl_2 gave benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as a thick yellow oil (855 mg, 51%). LC/MS: *m/z* 483.5 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001191] **2-(4-((*R*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol**


[001192]



[001193] To a mixture of benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (855 mg, 1.77 mmol) and EtOH (15 mL) in a round bottom flask was added Pd/C (86 mg, 10% wt Pd on carbon) and the flask was sealed with a septum. The atmosphere in the flask was evacuated, purged with N_2, and equipped with a balloon charged with H_2. The mixture was stirred under an H_2 atmosphere at ambient pressure for 3 h. After filtration through a plug of Celite using MeOH as the eluting solvent, the reaction mixture was concentrated to 2-(4-((*R*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (625 mg) as a yellow solid. LC/MS: *m/z* 349.3 $(M+H)^+$ at 1.82 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001194] **(*R*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001195]



[001196] Method A

[001197] 2-(4-((*R*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which (*R*)-tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) dissolved in DMF (100 μL) was added dropwise followed by the addition of triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature, and was complete after two hours. Purification using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001198] Method B


[001199] To a stirred solution of 2-(4-((*R*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (150 mg, 0.43 mmol) under an N_2 atmosphere at 0 °C was added

triethylamine (87 mg, 0.86 mmol) followed by the dropwise addition of (*R*)-tetrahydrofuran-3-yl chloroformate (65 mg, 0.43 mmol). The reaction was allowed to warm to room temperature and was stirred for 2 h. The mixture was partitioned between H_2O and CH_2Cl_2 and separated, and the aqueous layer was extracted twice with CH_2Cl_2. The organic layers were combined, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 4:1 CH_2Cl_2:EtOAc afforded (*R*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate. LC/MS: *m/z* 463.5$(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001200] (*R*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001201]


[001202] To a solution of (*R*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-



methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (0.085 g, 0.18 mmol) in 9 mL CH_2Cl_2 was added dropwise a 2.0 M HCl solution in ether (0.09 mL, 0.18 mmol). Ether (20 mL) was then added, leading to the precipitation of (*R*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride which was filtered and dried (85 mg, 95%). LC/MS: *m/z* 463.5$(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001203] Example 217


[001204] (*S*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate



[001205]

[001206] **(*S*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001207]



[001208] Method A

[001209] 2-(4-((*R*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which (*S*)-tetrahydrofuran-3-yl chloroformate (31 mg, 0.2 mmol) dissolved in DMF (100 μL) was added dropwise followed by triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 463.5 $(M+H)^+$ at 2.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001210] Method B

[001211] To a stirred solution of 2-(4-((*R*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (200 mg, 0.57 mmol) in DMF under an N_2 atmosphere at 0 °C was added triethylamine (115 mg, 1.14 mmol) followed by the dropwise addition of (*S*)-tetrahydrofuran-3-yl chloroformate (86 mg, 0.57 mmol). The reaction was allowed to warm to room temperature and was stirred for 2 h. The mixture was partitioned between H_2O and CH_2Cl_2 and separated, and the aqueous layer was extracted twice with CH_2Cl_2. The organic layers were

combined, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 4:1 CH_2Cl_2:EtOAc afforded (*S*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate. LC/MS: *m/z* 463.5$(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001212] ((*S*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001213]




[001214] To a solution of (*S*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate in 12 mL CH_2Cl_2 was added dropwise 2.0 M HCl solution in ether (0.13 mL, 0.25 mmol). To the solution was then added 20 mL ether leading to the precipitation of ((*S*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride which was filtered and dried (116 mg, 92%). LC/MS: *m/z* 463.5$(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001215] Example 218


[001216] (2*R*)-Tetrahydro-*N*-(1-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)furan-2-carboxamide

[001217]




[001218] (*R*)-Tetrahydrofuran-2-carboxylic acid (1-benzyl-piperidin-4-yl)-amide

[001219]



[001220] A solution of (*R*)-tetrahydro-2-furoic acid (58.5 g, 504 mmol) and oxalyl chloride (86 mL, 1.0 mol) in 100 mL CH_2Cl_2 were refluxed for 2 hours in a flask equipped with a $CaCl_2$ guard tube. After the solution was cooled to room temperature the solvents and excess oxalyl chloride were removed by evaporation under reduced pressure. The resulting acid chloride was dissolved in 200 mL CH_2Cl_2 and added dropwise to a solution of 1-benzyl-4-amino piperidine dihydrochloride (142 g, 539 mmol) and triethylamine (240 mL, 1.7 mol) in 300 mL CH_2Cl_2 cooled in an ice bath. The resulting mixture was stirred for 2 hours at room temperature and subsequently washed twice with 300 mL portions of 5% aq. $NaHCO_3$ and 300 mL saturated aq. NaCl solution, dried over Na_2SO_4, filtered, and evaporated to dryness under reduced pressure. The solid residue was triturated with 500 mL heptanes, collected by filtration, and washed twice with 200 mL portions of heptanes. The solid was air-dried at 45 °C to yield (*R*)-tetrahydrofuran-2-carboxylic acid (1-benzyl-piperidin-4-yl)-amide (128 g, 88%) as an off-white solid. ^{1}H-NMR (300 MHz, $CDCl_3$): δ 7.35–7.20 (m, 5H), 6.58 (bs, 1H), 4.29 (dd, *J*= 5.9, 8.4 Hz, 1H), 3.92–3.77 (m, 2H), 3.48 (s, 2H), 2.82–2.74 (m, 2H), 2.32–2.21 (m, 1H), 2.19–1.95 (m, 3H), 1.94–1.78 (m, 4H), 1.58–1.40 (m, 2H) ppm.

[001221] (*R*)-Tetrahydro-furan-2-carboxylic acid piperidin-4-ylamide as an oxalate salt




[001222]

[001223] (*R*)-Tetrahydro-furan-2-carboxylic acid (1-benzyl-piperidin-4-yl)-amide (128 g, 444 mmol) was dissolved in 300 mL ethanol and 50 mL acetic acid. Palladium on activated carbon (5 g) was added. A hydrogen pressure of 5 bars was applied, and hydrogenation was continued until no more hydrogen was consumed (about 5 days). The suspension was filtered through Celite, and the filtrate was evaporated to dryness under reduced pressure. The residue and oxalic acid (50 g, 555 mmol) were suspended in 500 mL 2-propanol and heated to dissolve the solids. Upon cooling, the oxalate salt of *R*-tetrahydro-furan-2-carboxylic acid piperidin-4-ylamide crystallized and was collected by filtration to yield (*R*)-Tetrahydro-furan-2-carboxylic acid piperidin-4-ylamide as an oxalate salt (99.0 g, 77%) as an off-white solid. ^{1}H-NMR (300 MHz, DMSO-d_6): δ 7.84 (d, J=7.8 Hz, 1H), 5.90 (bs, 2H), 4.18–4.14 (m, 1H), 3.89–3.67 (m, 3H), 3.25–3.20 (m, 2H), 2.94–2.71 (m, 2H), 2.13–2.00 (m, 1H), 1.56–1.58 (m, 6H) ppm.

[001224] (2*R*)-Tetrahydro-*N*-(1-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)furan-2-carboxamide

[001225]




[001226] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)-3-methylphenol (60 mg, 2.1 mmol) in DMF (1 mL) was added triethylamine (1.17 mL, 8.4 mmol) and (*R*)-tetrahydro-*N*-(piperidin-4-yl)furan-2-carboxamide as an oxalate salt (80 mg, 2.73 mmol).. This mixture was

stirred for 2 h at room temperature before it was purified via reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to afford (2*R*)-tetrahydro-*N*-(1-(2-(2-hydroxy-6-methylphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 447.5$(M+H)^+$ at 2.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001227] Example 219


[001228] **{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid tetrahydro-pyran-2-ylmethyl ester**

[001229]



[001230] **(Tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate**


[001231]



[001232] A mixture of (tetrahydro-2*H*-pyran-2-yl)methanol (369 mg, 3.18 mmol) and di(1*H*-imidazol-1-yl)methanone (1.0 g, 6.36 mmol) in 0.3 M CH_3Cl (10 mL) was stirred at 50 °C for 3 h. After allowing the reaction to cool to room temperature, the solvent was evaporated under reduced pressure giving (tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate (412 mg) which was used without further purification. LC/MS: *m/z* 211.1$(M+H)^+$ at 0.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001233] **{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid tetrahydro-pyran-2-ylmethyl ester.**

[001234]




[001235] To 2-[4-(4-amino-piperidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (100 mg, 0.3 mmol) in 1 mL CH_2Cl_2 was added sequentially triethylamine (62.5 μL, 0.45 mmol) and imidazole-1-carboxylic acid tetrahydro-pyran-2-ylmethyl ester(94 mg, 0.45 mmol). The reaction mixture was stirred at room temperature for 12 h and at 45 °C for 3 h. The reaction mixture was cooled and diluted with water and CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue was purified by reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) as eluent to give the desired product {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid tetrahydro-pyran-2-ylmethyl ester as the TFA salt. LC/MS: *m/z* 477.4 (M+H)+ at 2.84 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001236] Example 220


[001237] **Tetrahydro-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2*H*-pyran-4-carboxamide**

[001238]



[001239] *tert*-Butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate


[001240]



[001241] To a cooled (0–5 °C) suspension of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (50.2 g, 186 mmol) in dichloromethane (200 mL) was slowly added a solution of *tert*-butyl piperidin-4-ylcarbamate (39.0 g, 195 mmol) and triethylamine (56 mL, 390 mmol) in dichloromethane (200 mL). The resulting mixture was stirred overnight at room temperature. Water (400 ml) was added, and the layers were separated. The aqueous layer was extracted with dichloromethane (2 x 200 mL), and the combined organic layers were dried over sodium sulfate, filtered, and evaporated to dryness under reduced pressure. The residue was purified by column chromatography (SiO_2, eluent: dichloromethane/heptanes 4:6–1:0). Two fractions were obtained: 38.7 g of *tert*-butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as a yellow solid, and a less pure fraction (26.1 g) that was purified by recrystallization from methanol to yield *tert*-butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate. Both fractions were combined (17.0 g, 69%).

[001242] 2-(4-(4-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol


[001243]



[001244] *tert*-Butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate (55.7 g, 128 mmol) was dissolved in dichloromethane (200 mL) and trifluoroacetic acid (215 ml) was slowly added (careful: immediate gas evolution!). The resulting solution was stirred overnight at room temperature under a nitrogen atmosphere and evaporated to dryness. To the residue were added equal amounts of water and dichloromethane (300 mL). The obtained emulsion was basified to pH 9 with 33% aq. NaOH. The emulsion was cleared by the addition of methanol, and the layers were separated. The aqueous layer was extracted with dichloromethane (200 mL), and the combined organic extracts were dried over sodium sulfate, filtered, and evaporated to dryness to yield crude 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol. The crude material was purified by column chromatography (SiO_2, 2% methanol in dichloromethane) to yield 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (44 g, 97%) as a yellow solid. ^{1}H-NMR (300 MHz, $CDCl_3$): δ 8.52 (dd, *J*=2.1, 8.1 Hz, 1H), 7.77 (d, *J*=8.4 Hz, 1H), 7.62 (s, 1H), 7.37 (dt, *J*=1.5, 7.2 Hz, 1H), 7.22 (dd, *J*=1.5, 9.0 Hz, 1H), 7.04 (dd, *J*=1.2, 6.9 Hz, 1H), 6.92 (dt, *J*=1.2, 7.2 Hz, 1H), 4.49–4.39 (m, 2H), 3.34 (dt, *J*=2.4, 12.2 Hz, 2H), 3.12–3.02 (m, 1H), 2.54 (s, 3H), 2.09–2.00 (m, 2H), 1.68–1.55 (m, 2H), 1.40–1.25 (m, 2H) ppm.

[001245] Tetrahydro-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2*H*-pyran-4-carboxamide

[001246]




[001247] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) was added tetrahydro-2*H*-pyran-4-carboxylic acid (17.5 mg, 0.13 mmol) followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (44 mg, 0.117 mmol). The reaction was stirred for 16 h, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford tetrahydro-*N*-(1-(2-(2-

hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2*H*-pyran-4-carboxamide as the TFA salt. LC/MS: *m/z* 447.5(M+H)$^+$ at 2.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001248] Example 221


[001249] **2-(Tetrahydro-2*H*-pyran-4-yl)-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)acetamide**

[001250]



[001251] **2-(Tetrahydro-2*H*-pyran-4-yl)-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)acetamide**


[001252]



[001253] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) was added 2-(tetrahydro-2*H*-pyran-4-yl)acetic acid (13 mg, 0.09 mmol) followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (44 mg, 0.117 mmol). The reaction was stirred for 16 h, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-(tetrahydro-2*H*-pyran-4-yl)-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)acetamide as the TFA salt. LC/MS: *m/z* 461.5(M+H)$^+$ at 2.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001254] Example 222

[001255] **2-(2-Isopropyl-5-methylcyclohexyloxy)-*N*-(1-(2-(2-hydroxyphenyl)-7-**



methylquinazolin-4-yl)piperidin-4-yl)acetamide (


[001256] **2-(2-Isopropyl-5-methylcyclohexyloxy)-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)acetamide**

[001257] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) at 0 °C was added triethylamine (25 μL, 0.18 mmol) followed by the addition of 2-(2-isopropyl-5-methylcyclohexyloxy)acetyl chloride (21 mg, 0.09 mmol). The reaction was stirred for 16 h, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-(2-isopropyl-5-methylcyclohexyloxy)-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)acetamide as the TFA salt. LC/MS: *m/z* 531.3$(M+H)^+$ at 3.08 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001258] Example 223


[001259] ***N*-(1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-3-(pyridin-2-yl)propanamide**

[001260]



[001261] *N*-(1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-3-(pyridin-2-yl)propanamide

[001262]




[001263] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (30 mg, 0.09 mmol) in DMF (1 mL) was added 3-(pyridin-2-yl)propanoic acid (20 mg, 0.13 mmol) followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (44 mg, 0.117 mmol). The reaction was stirred for 16 h, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford *N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-3-(pyridin-2-yl)propanamide as the TFA salt. LC/MS: *m/z* 468.3 $(M+H)^+$ at 1.86 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001264] Example 224


[001265] (2*R*)-*N*-(1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide

[001266]



[001267] (2*R*)-*N*-(1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide

[001268]




[001269] To a solution of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (25 mg, 0.09 mmol) in CH_2Cl_2 (1 mL) was added (*R*)-tetrahydro-*N*-(piperidin-4-yl)furan-2-carboxamide oxalate (33 mg, 0.117 mmol), followed by the addition of triethylamine (50 μL, 0.36 mmol). The reaction was stirred for 2 h before it was filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*R*)-*N*-(1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide as the TFA salt. LC/MS: *m/z* 437.1$(M+H)^+$ at 2.54 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001270] Example 225


[001271] ***N*-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-3-pyridin-3-yl-propionamide**

[001272]



[001273] ***N*-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-3-pyridin-3-yl-propionamide**

[001274]




[001275] To 2-[4-(4-amino-piperidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (238 mg, 0.71 mmol) in 2.4 mL of CH_2Cl_2 was added sequentially 3-pyridin-3-yl-propionic acid (118.3 mg, 0.78 mmol), triethylamine (129 μL, 0.92 mmol), and BOP (346 mg, 0.78 mmol) at room temperature. The reaction mixture was stirred for 40 min and diluted with water and CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give an oil. The residue was purified by reverse phase LC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) as eluent to give the desired product as the TFA salt. LC/MS: *m/z* 468.6 (M+H)+ at 2.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001276] Example 226

[001277] ***N*-(((*S*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide**

[001278]



[001279] ***N*-(((*S*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide**




[001280]

[001281] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (35 mg, 0.10 mmol) was dissolved in DMF (1 mL). Cyclopropanecarboxylic acid (9.7 mg, 0.11 mmol) was added, followed by the addition of triethylamine (28 μL, 0.2 mmol), and the mixture was cooled in an ice water bath. HATU (42 mg, 0.11 mmol) was added in one portion, and the reaction was allowed to warm to room temperature while stirring for 16 h. The reaction mixture was filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give N-(((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methyl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 417.5 $(M+H)^+$ at 2.30 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001282] Example 227


(Pyridin-3-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001283]



[001284] ***tert*-Butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate**


[001285]



[001286] To a suspension of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (2.0 g, 7.38 mmol) in CH_2Cl_2 (25 ml) at 0 °C under an N_2 atmosphere was added dropwise a solution of *tert*-butyl piperidin-4-ylcarbamate (1.92 g, 9.6 mmol) in CH_2Cl_2 (10 ml) and triethylamine (2.0 ml, 14.76 mmol). The reaction was stirred for 6 hours, then it was quenched with water (25 ml), and the layers were separated. The aqueous phase was extracted twice with CH_2Cl_2 (10 ml), and the combined organic layers were dried over $MgSO_4$, filtered, and concentrated to give *tert*-butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as a yellow solid (3.24 g, 100%). LC/MS: *m/z* 435.3 $(M+H)^+$ at 2.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001287] **2-(4-(4-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol**

[001288]




[001289] *tert*-Butyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate (3.21 g, 7.39 mmol) was dissolved in CH_2Cl_2 (55 ml). TFA (50 ml) was added and the reaction was stirred for 1 hour. After evaporating the solvents *in vacuo*, the crude material was diluted with CH_2Cl_2 and neutralized with a 1 N NaOH solution. The layers were separated, and the aqueous layer was extracted three times with CH_2Cl_2 (30 ml). The combined organic

layers were dried over $MgSO_4$, filtered, and concentrated to give 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol as a yellow solid (2.06 g, 83%). LC/MS: *m/z* 335.3 $(M+H)^+$ at 1.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001290] (Pyridin-3-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001291]




[001292] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMSO (1 mL) was added (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (53 mg, 0.263 mmol) and triethylamine (30.4 mg, 42 μL, 0.3 mmol). The reaction was stirred overnight at room temperature and then purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (pyridin-3-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as the TFA salt. LC/MS: *m/z* 470.5 $(M+H)^+$ at 1.98 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001293] Example 228


[001294] (Pyridin-4-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001295]



[001296] (Pyridin-4-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001297]




[001298] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMSO (1 mL) was added (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (53 mg, 0.263 mmol) and triethylamine (30.4 mg, 42 μL, 0.3 mmol). The reaction was stirred overnight at room temperature and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (pyridin-4-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as the TFA salt. LC/MS: *m/z* 470.5 $(M+H)^+$ at 1.98 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001299] Example 229


[001300] (Benzo[d][1,3]dioxol-7-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001301]



[001302] (Benzo[d][1,3]dioxol-7-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate

[001303]




[001304] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMSO (1 mL), was added (benzo[d][1,3]dioxol-4-yl)methyl 1*H*-imidazole-1-carboxylate (65 mg, 0.263 mmol), followed by the addition of triethylamine (30.4 mg, 42 μL, 0.3 mmol). The reaction was stirred overnight at room temperature and then purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (benzo[d][1,3]dioxol-7-yl)methyl 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as the TFA salt. LC/MS: *m/z* 513.3 $(M+H)^+$ at 2.82 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001305] Example 230


[001306] *N*-((Tetrahydrofuran-2-yl)methyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxamide

[001307]



[001308] *N*-((Tetrahydrofuran-2-yl)methyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxamide

[001309]




[001310] A solution of 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxylic acid (45 mg, 0.12 mmol) in 500 μL DMF was cooled to 0 °C, and (tetrahydrofuran-2-yl)methanamine (13.2 mg, 0.13 mmol) and triethylamine (25 mg, 35 μL, 0.25 mmol) were added. After ten minutes HATU (57 mg, 0.15 mmol) was added to the reaction mixture in one portion .The reaction was allowed to warm to room temperature, stirred overnight, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) yielding *N*-((tetrahydrofuran-2-yl)methyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidine-3-carboxamide as the TFA salt. LC/MS: *m/z* 447.3 $(M+H)^+$ at 2.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001311] Example 231


[001312] **(*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001313]



[001314] ***tert*-butyl piperidin-3-ylmethylcarbamate**

[001315]



[001316] A cold solution of HCl in MeOH (prepared by adding acetyl chloride (13.5 mL, 14.9 g, 0.19 mol) to 1 L of MeOH) was added to a cold solution of *tert*-butyl pyridin-3-ylmethylcarbamate (41.0 grams, 0.20 mol) in MeOH (100 mL). The solution was transferred to a Parr hydrogenation apparatus at 12 °C. PtO_2 (3 g) was added, and 12 bar pressure of H_2 was applied. After 16 hours ^{1}H NMR of a concentrated sample indicated the reaction to be complete. The catalyst was filtered, and conc. aq. NaOH (20 mL) was added to neutralize the HCl. The solution was concentrated to remove the bulk of the MeOH and extracted with *tert*-butyl methyl ether (4x200 mL). The combined organic layers were washed with saturated aqueous NaCl solution, dried over Na_2SO_4, and concentrated to give the product (40.16 g, 0.187 mol, 95%) as a yellow oil that crystallized upon standing. ^{1}H NMR (300 MHz, $CDCl_3$): δ 4.62 (bs. s, 1H), 3.06–2.94 (m, 4H), 2.52 (dt, *J*= 12 Hz, 3Hz, 1H), 2.28 (dd, *J*= 12 Hz, 10Hz, 1H), 1.82–1.72 (m, 1H), 1.70–1.51 (m, 3H), 1.49–1.34 (m, 1H), 1.42 (s, 9H), 1.06 (dq, *J*= 12 Hz, 4Hz, 1H).

[001317] (*R*)-*tert*-Butyl piperidin-3-ylmethylcarbamate


[001318]



[001319] To a solution of *tert*-butyl piperidin-3-ylmethylcarbamate (162 g, 0.758 mol) in EtOH was added (+)-dianisoyltartaric acid (316 g, 0.756 mol). The suspension was heated until clear and allowed to cool to room temperature overnight. The precipitated salt was recrystallized three times from EtOH. The salt was washed with EtOH (2x200 mL) and air-dried. Residual solvent was removed *in vacuo*. The salt was taken up in *tert*-butyl methyl ether and 10% aq. NaOH. The organic layer was separated, and the aqueous layer was extracted with *tert*-butyl methyl ether (3x200 mL). More product was extracted after addition of 30% aq. NaOH to the aqueous layer. The combined organic layers were washed with water and saturated aqueous NaCl solution, dried over Na_2SO_4, and concentrated to give (*R*)-*tert*-butyl piperidin-3-ylmethylcarbamate as a white crystalline solid (41.3 g, 0.192 mol, 25%). For ee determinations, samples of the salt were taken up in CH_2Cl_2 and 1 N aq. NaOH. The organic layer was washed with water, dried over Na_2SO_4, and filtered. A drop of 1-naphthyl isocyanate was added, and after 15 minutes a drop of morpholine was added to quench excess isocyanate. Volatiles were evaporated after another 15 minutes. The sample was dissolved in EtOH for chiral HPLC


(Chiralcel OD-H; heptane/EtOH/Et_2NH 90/10/0.2; 0.5 mL/min; Rt (*R*): 46 min, Rt (*S*): 57 min, Rt (*N*-(naphthalen-1-yl)morpholine-4-carboxamide): 64 min.

[001320] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol



[001321] 2-(4-Chloro-7-methylquinazolin-2-yl)-3-fluorophenol (1.0 g, 3.46 mmol) was dissolved in 15 mL anhydrous CH_2Cl_2 under an N_2 atmosphere and cooled using an ice bath and piperidin-3-ylmethyl-carbamic acid *tert*-butyl ester/oxalic acid (1.16 g, 3.81 mmol) was added in portions, followed by triethylamine (1.05 g, 1.45 mL, 10.4 mmol). The reaction was allowed to warm to room temperature and was complete after 1.5 hour. The mixture was partitioned between CH_2Cl_2 and H_2O and separated, and the aqueous layer was extracted once more with CH_2Cl_2. The organic extracts were combined, dried over Na_2SO_4, filtered, and concentrated to a yellow solid. This solid was suspended in 40 mL CH_2Cl_2 and 20 mL of TFA were added. The reaction was complete after 1 hour. The solvent and the excess TFA were removed *in vacuo*, the residue was re-dissolved in CH_2Cl_2, and the pH was adjusted to 7 using an aqueous 1 M solution of NaOH. The reaction was partitioned between CH_2Cl_2 and H_2O and separated, and the aqueous layer was extracted once more with CH_2Cl_2. The organic extracts were combined, dried over Na_2SO_4, filtered, and concentrated to yield of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (900 mg, 71% overall yield) as a yellow solid. LC/MS: *m/z* 367.3 $(M+H)^+$ at 1.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001322] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate




[001323]

[001324] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.11 mmol) was dissolved in anhydrous DMF (800 μL) and cooled to 0 °C, upon which (*S*)-tetrahydrofuran-3-yl chloroformate (16.3 mg, 0.12 mmol) dissolved in DMF (100 μL) was added dropwise followed by triethylamine (22 mg, 30.3 μL, 0.218 mmol). The reaction was allowed to warm to room temperature, and after 2 h the reaction was complete. The mixture was purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to yield (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 481.1 $(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001325] Example 232


[001326] **(2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-4,4-dimethylpentanamide**

[001327]



[001328] **(2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-4,4-dimethylpentanamide**

[001329]




[001330] 2-(4-(4-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.15 mmol) was dissolved in DMF (1 mL) and cooled to 0 °C . (2*R*)-Hydroxy-4,4-dimethyl-pentanoic acid (26.3 mg, 0.18 mmol) was added followed by the addition of triethylamine (42 μL, 0.3 mmol). After ten minutes, HATU (68 mg, 0.18 mmol) was added in one portion. The reaction was stirred at 0 °C for 10 minutes, and then allowed to warm to room temperature. The reaction was complete after 40 minutes, then filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (2*R*)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-4,4-dimethylpentanamide as the TFA salt. LC/MS: *m/z* 463.3 $(M+H)^+$ at 2.58 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001331] Example 233


[001332] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide**

[001333]



[001334] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide


[001335]



[001336] To 2-hydroxy-2-methylpropanoic acid (28 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL), followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide as the TFA salt. LC/MS: *m/z* 421.2$(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001337] 2-Ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide

[001338]



[001339] 2-Ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide

[001340]




[001341] To 2-ethyl-2-hydroxybutanoic acid (36 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 449.2$(M+H)^+$ at 2.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001342] Example 235


[001343] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide

[001344]



[001345] *tert*-Butyl 1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate


[001346]



[001347] To a suspension of 2-(4-chloroquinazolin-2-yl)phenol (2.0 g, 7.79 mmol) in CH_2Cl_2 (25 mL) at 0 °C under an N_2 atmosphere, was added dropwise *tert*-butyl piperidin-4-ylcarbamate (2.08 g, 10.13 mmol) and triethylamine (2.20 mL, 15.8 mmol) in CH_2Cl_2 (10 mL). The reaction was stirred overnight, quenched with water and extracted with CH_2Cl_2, and the combined organic layers were dried over $MgSO_4$, filtered, and concentrated to afford *tert*-butyl 1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate (3.27 g, 100% –solvent residue). LC/MS: *m/z* 421.3$(M+H)^+$ at 2.70 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001348] 2-(4-(4-Aminopiperidin-1-yl)quinazolin-2-yl)phenol

[001349]




[001350] To a solution of *tert*-butyl 1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate (3.27 g, 8.09 mmol) in CH_2Cl_2 (75 mL) was added TFA (50 mL). The reaction was complete after 45 minutes After evaporating the solvents *in vacuo*, the crude material was diluted with CH_2Cl_2 and neutralized by adding a 1 N aqueous NaOH solution. The layers were separated, and the aqueous layer was extracted with CH_2Cl_2 (2×15 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to afford 2-(4-(4-aminopiperidin-1-

yl)quinazolin-2-yl)phenol as a solid (0.09 g, 84%). LC/MS: *m/z* 321.3$(M+H)^+$ at 1.28 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001351] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide


[001352]



[001353] To 2-hydroxy-2-methylpropanoic acid (27 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylpropanamide as the TFA salt. LC/MS: *m/z* 407.5$(M+H)^+$ at 2.04 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001354] Example 236


[001355] (2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide



[001356]

[001357] **(2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide**

[001358]




[001359] To (*R*)-2-hydroxybutanoic acid (30 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) dissolved in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*R*)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 407.3$(M+H)^+$ at 2.08 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001360] Example 237


[001361] **(2*S*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide**

[001362]



[001363] **(2*S*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide**

[001364]




[001365] To (*S*)-2-hydroxybutanoic acid (30 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*S*)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 407.3$(M+H)^+$ at 2.09 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001366] Example 238


[001367] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)hexanamide**

[001368]



[001369] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)hexanamide**


[001370]



[001371] To 2-hydroxyhexanoic acid (38 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)hexanamide as the TFA salt. LC/MS: *m/z* 435.3$(M+H)^+$ at 2.4 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001372] Example 239

[001373] 2-(Trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)propanamide



[001374] 2-(Trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)propanamide

[001375]




[001376] To 2-(trifluoromethyl)-2-hydroxypropanoic acid (45 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) dissolved in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-(trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)propanamide as the TFA salt. LC/MS: *m/z* 461.1$(M+H)^+$ at 2.4 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001377] Example 240


[001378] **2-Ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide**

[001379]



[001380] **2-Ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide**

[001381]




[001382] To 2-ethyl-2-hydroxybutanoic acid (38 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-ethyl-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 435.5$(M+H)^+$ at 2.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001383] Example 241


[001384] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)pentanamide

[001385]



[001386] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)pentanamide

[001387]




[001388] To 2-hydroxypentanoic acid (34 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)pentanamide as the TFA salt. LC/MS: *m/z* 421.1 $(M+H)^+$ at 2.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001389] Example 242


[001390] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide**

[001391]



[001392] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide**

[001393]




[001394] To 2-hydroxy-2-methylbutanoic acid (34 mg, 0.28 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)quinazolin-2-yl)phenol (0.07 g, 0.22 mmol) in DMF (0.5 mL), followed by the addition of triethylamine (0.061 mL, 0.44 mmol) and a solution of HATU (0.107 g, 0.284 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide as the TFA salt. LC/MS: *m/z* 421.3$(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001395] Example 243


[001396] **(2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide**

[001397]



[001398] **(2*R*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide**

[001399]




[001400] To (*R*)-2-hydroxybutanoic acid (28 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*R*)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 421.3$(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001401] Example 244


[001402] **(2*S*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide**

[001403]



[001404] **(2*S*)-2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide**

[001405]




[001406] To (*S*)-2-hydroxybutanoic acid (28 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL), followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, then filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*S*)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)butanamide as the TFA salt. LC/MS: *m/z* 421.3$(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001407] Example 245


[001408] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)hexanamide

[001409]



[001410] 2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)hexanamide

[001411]




[001412] To 2-hydroxyhexanoic acid (36 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)hexanamide as the

TFA salt; LC/MS: m/z = 449.3 $(M+H)^+$ at 2.45 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001413] Example 246


[001414] **2-(Trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)propanamide**

[001415]



[001416] **2-(Trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)propanamide**

[001417]




[001418] To 2-(trifluoromethyl)-2-hydroxypropanoic acid (43 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

to afford 2-(trifluoromethyl)-2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)propanamide as the TFA salt. LC/MS: *m/z* 475.1$(M+H)^+$ at 2.46 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001419] Example 247


[001420] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)pentanamide**

[001421]



[001422] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)pentanamide**

[001423]




[001424] To 2-hydroxypentanoic acid (32 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL), followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and purified

using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)pentanamide as the TFA salt. LC/MS: *m/z* 435.3$(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001425] Example 248


[001426] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide**

[001427]



[001428] **2-Hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide**

[001429]




[001430] To 2-hydroxy-2-methylbutanoic acid (32 mg, 0.27 mmol) was added a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.07 g, 0.21 mmol) in DMF (0.5 mL) followed by the addition of triethylamine (0.058 mL, 0.42 mmol) and a solution of HATU (0.103 g, 0.273 mmol) in DMF (0.5 mL). The reaction was stirred overnight, filtered, and

purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-hydroxy-*N*-(1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-methylbutanamide as the TFA salt. LC/MS: *m/z* 435.3$(M+H)^+$ at 2.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001431] Example 249


[001432] **(2*R*)-*N*-(1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-hydroxy-4,4-dimethylpentanamide**

[001433]



[001434] ***tert*-Butyl 1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate**


[001435]



[001436] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol (700 mg, 2.42 mmol) in 10 ml CH_2Cl_2 at 0 °C was added triethylamine (0.67 mL, 4.8 mmol) followed by the addition of *tert*-butyl piperidin-4-ylcarbamate (630 mg, 3.14 mmol) under an N_2 atmosphere. The reaction was gradually warmed to room temperature and stirred overnight. The reaction mixture was then quenched with water, and the layers were separated. The aqueous layer was extracted twice with CH_2Cl_2, and the combined organic layers were dried over $MgSO_4$, filtered,

and concentrated to obtain *tert*-butyl 1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate (1.05 g, 96%) LC/MS: *m/z* 453.3$(M+H)^+$ at 2.60 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))


[001437] 2-(4-(4-Aminopiperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol



[001438] To a solution of *tert*-butyl 1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate (1.05 g, 2.3 mmol) in 20 ml CH_2Cl_2 was slowly added TFA (5 mL). The reaction was stirred for one hour before it was evaporated to dryness. To the residue was added CH_2Cl_2, and the reaction was neutralized using an aqueous 1 M NaOH solution. The aqueous layer was extracted twice with CH_2Cl_2, and the combined organic layers were dried over MgSO4, filtered, and concentrated to obtain 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.75 g, 92%). LC/MS: *m/z* 353.3$(M+H)^+$ at 1.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001439] (2*R*)-*N*-(1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-hydroxy-4,4-dimethylpentanamide

[001440]




[001441] To a solution of 2-(4-(4-aminopiperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (50 mg, 0.14 mmol) in 1 mL DMF at 0 °C was added (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (25 mg, 0.17 mmol), followed by the addition of triethylamine (29 mg, 0.28 mmol). HATU (65 mg, 0.17 mmol) was then added, and the reaction was stirred at 0 °C for an additional 10 minutes and then warmed to room temperature. The reaction was complete after 40 minutes, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (2*R*)-*N*-(1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-2-hydroxy-4,4-dimethylpentanamide as the TFA salt. LC/MS: *m/z* 481.3 $(M+H)^+$ at 2.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001442] Example 250


[001443] **Isobutyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001444]



[001445] **Isobutyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001446]




[001447] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which isobutyl chloroformate (27 μL, 0.2 mmol) dissolved in DMF (100 μL) was added dropwise followed by

triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature, and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.5 $(M+H)^+$ at 2.77 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001448] Example 251


[001449] **Ethyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001450]



[001451] **Ethyl (1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001452]




[001453] 2-(4-(3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which ethyl chloroformate (20μL, 0.2 mmol) dissolved in DMF (100 μL) was added dropwise followed by the addition of triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave ethyl (1-(2-(2-hydroxyphenyl)-7-

methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 421.0 $(M+H)^+$ at 2.48 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001454] Example 252


[001455] **Isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001456]



[001457] **Isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001458]




[001459] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.17 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which isobutyl chloroformate (27 μL, 0.2 mmol) dissolved in DMF (100 μL) was added dropwise followed by the addition of triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001460] Example 253


[001461] Isobutyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001462]



[001463] Isobutyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001464]




[001465] 2-(4-((*R*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (60 mg, 0.172 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which isobutyl chloroformate (27 μL, 0.21 mmol) dissolved in DMF (100 μL) was added dropwise followed by the addition of triethylamine (35 mg, 48 μL, 0.34 mmol). The reaction was allowed to warm to room temperature, and it was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.78 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001466] Example 254


[001467] Ethyl((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001468]



[001469] **Ethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001470]




[001471] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (35 mg, 0.10 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which ethyl chloroformate (10.5 μL, 0.11 mmol) dissolved in DMF (100 μL) was added dropwise followed by triethylamine (20.2 mg, 27.8 μL, 0.2 mmol). The reaction was allowed to warm to room temperature, and it was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave ethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 421.1 $(M+H)^+$ at 2.50 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001472] Example 255


[001473] **Isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001474]



[001475] **Isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**


[001476]




[001477] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (35 mg, 0.10 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which isopropyl chloroformate (1 M solution in toluene, 98.1 mg, 110 μL, 0.11 mmol) was dissolved in DMF (100 μL) and added dropwise, followed by triethylamine (20.2 mg, 27.8 μL, 0.2 mmol). The reaction was allowed to warm to room temperature, and it was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.61 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001478] Example 256

[001479] **Propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**




[001474] **Propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**



[001475] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (35 mg, 0.10 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which propyl chloroformate (12.4 μL, 0.11 mmol) dissolved in DMF (100 μL) was added dropwise followed by triethylamine (20.2 mg, 27.8 μL, 0.2 mmol). The reaction was allowed to warm to room temperature and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.61 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001476] Example 257

[001477] **2-Methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**



[001487] 2-Methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate

[001488]




[001489] 2-(4-((*S*)-3-(Aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)phenol (35 mg, 0.1 mmol) was dissolved in anhydrous DMF (1 mL) and cooled to 0 °C, upon which 2-methoxyethyl chloroformate (11.6μL, 0.1 mmol) dissolved in DMF (100 μL) was added dropwise followed by triethylamine (20.2 mg, 27.8 μL, 0.2 mmol). The reaction was allowed to warm to room temperature, and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 451.1 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001490] Example 258


[001491] 2-Methoxyethyl 1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate

[001492]



[001493] 2-(4-(4-Aminopiperidin-1-yl)-6-fluoroquinazolin-2-yl)phenol


[001494]



[001495] To a stirred solution of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (0.20 g, 0.73 mmol) in CH_2Cl_2 (5mL) was added dropwise a solution of *tert*-butyl piperidin-4-ylcarbamate (0.19 g, 0.95 mmol) and triethylamine (203 μL, 147 mg, 1.46 mmol) in CH_2Cl_2. The reaction was stirred for 3 h and then quenched with water. The aqueous layer was extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to obtain *tert*-butyl 1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate. The residue was dissolved in 10 mL CH_2Cl_2 and 3 mL TFA. The reaction was stirred for 2 hours and neutralized with a 1.0 M aqueous NaOH solution. The mixture was partitioned between H_2O/CH_2Cl_2 and separated, and the aqueous layer was extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to obtain 2-(4-(4-aminopiperidin-1-yl)-6-fluoroquinazolin-2-yl)phenol. LC/MS: *m/z* 339.3$(M+H)^+$ at 1.87 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001496] 2-Methoxyethyl 1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate

[001497]




[001498] A solution of 2-(4-(4-aminopiperidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (50 mg, 0.15 mmol) in DMF (1 mL) was cooled to –40 °C (external temp). To it was added triethylamine (41 μL, 30 mg, 0.29 mmol) and a solution of 2-methoxyethyl chloroformate (17 μL, 20 mg, 0.15 mmol) in 100 μL DMF dropwise. The reaction was allowed to warm to room temperature and was stirred for 2 h. Purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl 1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)piperidin-4-ylcarbamate as the TFA salt. LC/MS: *m/z* 441.5$(M+H)^+$ at 2.6 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA))

[001499] Example 259


[001500] **{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid *tert*-butyl ester**

[001501]



[001502] **{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid *tert*-butyl ester**

[001503]




[001504] To piperidin-4-yl-carbamic acid *tert*-butyl ester (887 mg, 4.4 mmol) in 10 mL of was CH_2Cl_2 was added sequentially triethylamine (720 μL, 5.2 mmol) and 2-(4-chloro-7-methyl-

quinazolin-2-yl)-phenol(1.0 g, 3.7 mmol). The reaction mixture was stirred at room temperature for 2 h and then diluted with water and CH_2Cl_2. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was subjected to purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-carbamic acid *tert*-butyl ester as the TFA salt. LC/MS: *m/z* 435.2 (M+H)+ at 3.03 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001505] Example 260


[001506] **Isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001507]



[001508] **Isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001509]




[001510] To a solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.11 mmol) in DMF (400 μL) at 0 °C was added dropwise isobutyl chloroformate (15.7 μL, 0.12 mmol) in DMF (400 μL), followed by the addition of triethylamine

(30 μL, 0.22 mmol). The reaction was allowed to warm to room temperature and was complete after 2h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 467.1 $(M+H)^+$ at 2.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001511] Example 261

[001512] **2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**


[001513]



[001514] Method A

[001515] **2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001516]




[001517] To a solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.11 mmol) in DMF (400 μL) at 0 °C was added dropwise 2-methoxyethyl chloroformate (12.6 μL, 0.11 mmol) in DMF (400 μL), followed by the addition of triethylamine (30 μL, 0.22 mmol). The reaction was allowed to warm to room temperature and was complete after 2h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-

methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.20 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001518] Method B

[001519] **2-Methoxyethyl ((*S*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate**


[001520]



[001521] To a solution of (*S*)-*tert*-butyl 3-(aminomethyl)piperidine-1-carboxylate (500 mg, 2.33 mmol) in CH_2Cl_2 at –10 °C was added triethylamine (650 μL, 4.66 mmol) followed by the dropwise addition of 2-methoxyethyl chloroformate (325 μL, 2.79 mmol). The reaction was warmed to room temperature and quenched with water. The aqueous layer was extracted with CH_2Cl_2, and the combined extracts were dried over $MgSO_4$, filtered, and evaporated. Purification via silica gel chromatography using 0 to 10% EtOAc in CH_2Cl_2 afforded 2-methoxyethyl ((*S*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate (496 mg, 67%). LC/MS: *m/z* 317.3 $(M+H)^+$ at 2.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001522] **2-Methoxyethyl ((*R*)-piperidin-3-yl)methylcarbamate**


[001523]



[001524] To 2-methoxyethyl ((*S*)-1-(*tert*-butoxycarbonyl)piperidin-3-yl)methylcarbamate (496 mg, 1.6 mmol) dissolved in 10 mL CH_2Cl_2 was added 5 mL TFA, and the reaction was stirred for 1 hour. After neutralizing the mixture with a 1 N NaOH solution it was extracted with CH_2Cl_2, and the combined organic layers were dried over MgSO4, filtered, and concentrated to give 2-methoxyethyl ((*R*)-piperidin-3-yl)methylcarbamate (300 mg, 87%) which was used without further purification. LC/MS: *m/z* 217.5 $(M+H)^+$ at 0.49 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001525] 2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate


[001526]



[001527] To a solution of 2-methoxyethyl ((*R*)-piperidin-3-yl)methylcarbamate (0.24 g, 1.1 mmol) and 2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol (250 mg, 0.865 mmol) in CH_2Cl_2 was added triethylamine (2.41 mL, 1.73 mmol). After stirring the reaction at room temperature for 2 h, it was quenched with water and then extracted with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0%–10% EtOAc in CH_2Cl_2 yielded 2-methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (357 mg, 88%). LC/MS: *m/z* 469.5 $(M+H)^+$ at 2.30 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 7.86 (d, J = 8.5 Hz, 1H), 7.64 (s, 1H), 7.35 (m, 3H), 6.78 (d, J = 8.1 Hz, 1H), 6.71 (m, 2H), 4.40 (d, J = 12.7 Hz, 2H), 4.05 (m, 2H), 3.48 (m, 2H), 3.30 (m, 1H), 3.25 (s, 3H), 3.05 (m, 3H), 1.82 (m, 3H), 1.63 (m, 1H), 1.32 (m, 1H).

[001528] 2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001529]




[001530] To a solution of ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (352 mg, 0.75 mmol) in CH_2Cl_2 (3 mL) was added dropwise

a 2.0 M HCl solution in ether (0.375 mL, 0.75 mmol) under an N_2 atmosphere. It was followed by the addition of 20 mL ether which lead to the precipitation of 2-methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride (350 mg, 92%) which was then filtered and dried. LC/MS: *m/z* 469.5 $(M+H)^+$ at 2.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 7.97 (d, *J* = 7.9 Hz, 1H), 7.63 (s, 1H), 7.45 (m, 3H), 6.89 (d, *J* = 8.3 Hz, 1H), 6.82 (t, *J* = 9.5 Hz, 1H), 4.54 (s, 1H), 4.03 (d, *J* = 2.7 Hz, 2H), 3.47 (t, *J* = 4.6 Hz, 4H), 3.24 (s, 3H), 2.99 (m, 2H), 2.54 (s, 3H), 1.89 (m, 3H), 1.70 (m, 1H), 1.38 (m, 1H)

[001531] Example 262


[001532] **Ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001533]



[001534] **Ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate**

[001535]



[001536] Method A

[001537] To a solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.11 mmol) in DMF (400 μL) at 0 °C was added dropwise ethyl chloroformate (10.4 μl, 0.10 mmol) in DMF (400 μL), followed by the addition of triethylamine


(30 μL, 0.22 mmol). The reaction was allowed to warm to room temperature and was complete after 2 h. Purification by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001538] Method B

[001539] To a stirred solution of 2-(4-((*S*)-3-(aminomethyl)piperidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (200 mg, 0.54 mmol) in 12 mL THF was added diisopropyl ethylamine (188 μL, 1.08 mmol) at room temperature. The mixture was cooled to –60 °C, and a solution of ethyl chloroformate (52 μL, 0.54 mmol) in 0.6 mL THF was added dropwise. After allowing the reaction to warm to room temperature, the mixture was partitioned between H_2O and CH_2Cl_2. The layers were separated, and the aqueous layer was extracted twice more with CH_2Cl_2. The organic layers were combined, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in 50:50 hexanes:CH_2Cl_2 gave ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (92 mg, 38%). LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.40 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 7.87 (d, J = 8.5 Hz, 1H), 7.64 (s, 1H), 7.34 (m, 2H), 7.21 (m, 1H), 6.78 (d, J = 8.3 Hz, 1H), 6.70 (m, 1H), 4.40 (m, 2H), 3.98 (m, 2H), 3.27 (m, 1H), 3.05 (m, 1H), 2.97 (m, 2H), 1.84 (d, J = 11.0 Hz, 3H), 1.67 (d, J = 9.4 Hz, 1H), 1.32 (dd, J = 20.8, 11.3 Hz, 1H), 1.15 (t, J = 7.1 Hz, 3H)

[001540] Ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride

[001541]




[001542] To a solution of ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate (89 mg, 0.2 mmol) in 2 mL CH_2Cl_2 under an N_2 atmosphere was added ether (10 mL) followed by the dropwise addition of a 2.0 M solution of HCl in ether (0.1 mL, 0.2 mmol) which resulted in precipitation of ethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)piperidin-3-yl)methylcarbamate hydrochloride which was then filtered and dried (85 mg, 90%). LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.37 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.26 (d, J = 8.5 Hz, 1H), 7.57 (s, 1H), 7.45 (m, 2H), 6.81 (m, 2H), 4.57 (m, 2H), 3.89 (m, 2H), 3.52 (m, 1H), 3.36 (m, 1H), 2.95 (s, 2H), 2.52 (s, 3H), 1.86 (m, 3H), 1.66 (m, 1H), 1.39 (m, 1H), 1.07 (t, J = 6.6 Hz, 3H).

[001543] Example 276


[001544] **(*R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide**

[001545]



[001546] **(*R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide**

[001547]




[001548] Method A: (*R*)-Tetrahydrofuran-2-carboxylic acid (23 mg, 0.20 mmol) and HATU (84 mg, 0.22 mmol) were dissolved in 0.75 mL DMF, then triethylamine (40 mg, 55 μL, 0.40 mmol) was added, followed immediately by 2-[4-(4-Amino-piperidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (67 mg, 0.20 mmol). The reaction was then stirred for 30 min at room temperature, diluted with 0.75 mL 1:1 methanol / DMSO, filtered, and purified by reverse phase HPLC (2–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (*R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide as the TFA salt. LC/MS: *m/z* 433.2 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001549] Method B: (*R*)-Tetrahydrofuran-2-carboxylic acid (193 mg, 1.66 mmol) was dissolved in DMF (6 mL), followed by the addition of HATU (696 mg, 1.83 mmol). The mixture was then stirred for 15 min at room temperature under an N_2 atmosphere. The 2-[4-(4-Amino-piperidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (556 mg, 1.66 mmol) was dissolved in DMF (8 mL) and added to the mixture, followed by triethylamine (336 mg, 0.463 mL, 3.32 mmol). After 30 min, theDMF was removed *in vacuo* and the crude product partitioned between water and ethyl acetate. The aqueous layer was separated, washed with ethyl acetate, and the combined organic layers dried over $MgSO_4$, filtered, and concentrated to give crude product as an orange oil. Purification via silica gel chromatography (20% ethyl acetate / 80% 1:1 CH_2Cl_2:hexane) gave pure (*R*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide (303 mg, 42%). ^1H NMR (400 MHz, DMSO-d6) δ 8.44 (dd, J = 8.2, 1.5 Hz, 1H), 7.90 (d, J = 8.5 Hz, 1H), 7.75 (d, J = 8.3 Hz, 1H), 7.67 (s, 1H), 7.40–7.36 (m, 2H), 6.97–6.93 (m, 2H), 4.48–4.45 (m, 2H), 4.20 (dd, J = 8.2, 5.3 Hz, 1H), 4.06–3.96 (m, 1H), 3.88 (dd, J = 14.3, 6.6 Hz, 1H), 3.74 (dd, J = 14.5, 6.7 Hz, 1H), 3.43–3.32 (m, 2H), 2.51 (s, 3H), 2.16–2.07 (m, 1H),

1.89–1.69 (m, 7H). ^{13}C NMR (100 MHz, DMSO-d6) δ 171.7, 162.8, 160.7, 159.5, 149.3, 144.1, 132.5, 129.0, 127.3, 125.7, 125.4, 118.9, 118.4, 117.2, 111.9, 77.6, 68.5, 48.3, 45.5, 31.3, 31.2, 30.0, 24.9, 21.2.

[001550] (*R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide hydrochloride




[001551] (*R*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide (303 mg, 0.701 mmol) was dissolved in 9 mL 2:1 dry ether / dry CH_2Cl_2 and 2.0 M HCl in ether added dropwise (0.35 mL, 0.70 mmol), producing a white precipitate which was collected by filtration to give (*R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-piperidin-4-yl}-amide hydrochloride (268 mg, 82%). LC/MS: *m/z* 433.5 $(M+H)^+$ at 2.26 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^{1}H NMR (400 MHz, DMSO-d6) δ 8.20 (dd, *J* = 7.8, 1.2 Hz, 1H), 8.00 (d, *J* = 8.6 Hz, 1H), 7.80–7.79 (m, 2H), 7.51–7.47 (m, 2H), 7.15 (d, *J* = 8.2 Hz, 1H), 7.05–7.01 (m, 1H), 4.66–4.63 (m, 2H), 4.21 (dd, *J* = 8.2, 5.2 Hz, 1H), 4.12–4.03 (m, 1H), 3.91–3.84 (m, 1H), 3.78–3.70 (m, 1H), 3.59 (t, *J* = 12.3 Hz, 2H), 2.54 (s, 3H), 2.16–2.05 (m, 1H), 1.97–1.94 (m, 2H), 1.89–1.72 (m, 5H).

[001552] Example 301


[001553] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001554]



[001555] *tert*-Butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate


[001556]



[001557] To 2-(4-chloro-7-methyl-quinazolin-2-yl)-phenol (644 mg, 2.4 mmol) in 2.9 mL of DMF at room temperature was added sequentially (*R*)- pyrrolidin-3-yl-carbamic acid *tert*-butyl ester (857 mg, 4.6 mmol) and triethylamine (662 μL, 4.8 mmol), and the reaction mixture was stirred for 12 h. The reaction mixture was diluted with water (10 mL) and CH_2Cl_2 (10 mL). The organic layer was separated and dried (Na_2SO_4), and the residue was purified by silica gel chromatography with 25–85% ethyl acetate/hexanes to give *tert*-butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (856 mg, 86%). LC/MS: *m/z* 421 $(M+H)^+$ at 2.82 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001558] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001559]



[001560] Method A

[001561] To *tert*-butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (850 mg, 2.02 mmol) was added at room temperature 4 mL of 1:1 TFA:CH_2Cl_2. The reaction mixture was stirred for 50 min, diluted with 20 mL of CH_2Cl_2, and washed with 15 mL of satd. $NaHCO_3$ solution. The organic layer was separated and dried over Na_2SO_4, and the


solvent was removed under reduced pressure to give (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol as an oil which was used without further purification.

To 45.9 mg (0.14 mmol) of the amine from above procedure was added 570 μL of CH_2Cl_2 and the solution was cooled to 0 °C. To this solution was added sequentially 24 μL (0.17 mmol) of triethylamine and 19.4 mg (0.13 mmol) of (*S*)- tetrahydro-furan-3-ol chloroformate. The reaction mixture was stirred at 0 °C for 45 min, diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 435 $(M+H)^+$ at 2.41 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001562] Method B

[001563] To *tert*-butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (907.2 mg, 2.16 mmol) was added 4 mL of 1:1 TFA:CH_2Cl_2. The mixture was stirred at room temperature for 5 hours. The reaction was diluted with a solution of saturated aqueous $NaHCO_3$ and CH_2Cl_2. The organic layer was separated, dried over Na_2SO_4, and concentrated under reduced pressure to give 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol. To this free amine (640 mg, 2 mmol) was added 8 ml of CH_2Cl_2 and triethylamine (335 μL, 2.4 mmol). After cooling the mixture to 0 °C, (*S*)-tetrahydrofuran-3-yl chloroformate (271 mg, 1.8 mmol) was added, and the reaction was allowed to stir for 30 minutes. Purification on silica gel using 30–100% ethyl acetate/hexanes gave (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate. ^1H NMR (400 MHz, $CDCl_3$) δ 14.8 (bs, 1H), 8.34 (d, J = 6.4 Hz, 1H), 7.83 (d, J = 4.0 Hz, 1H), 7.48 (bs, 1H), 7.27 (m, 1H), 7.08 (d, J = 8.1 Hz, 1H), 6.93 (d, J = 8.1 Hz, 1H), 6.82 (t, J = 7.5 Hz, 1H), 5.21 (s, 1H), 5.06 (s, 1H), 4.37 (s, 1H), 4.15 (m, 1H), 4.02 (s, 1H), 3.97 (d, J = 5.6 Hz, 1H), 3.87–3.77 (m, 5H), 2.42 (s, 3H), 2.24 (t, J = 6.0 Hz, 1H), 2.15–2.06 (m, 1H), 1.98 (q, J = 9.3 Hz, 2H).


[001564] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride



[001556] A 2.0 M HCl solution in Et_2O (192 μL, 0.38 mmol) was slowly added at –20 °C to a stirring solution of (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazoli 4-yl)pyrrolidin-3-ylcarbamate (167 mg, 0.38 mmol) in 700 μL of CH_2Cl_2. The reaction was allowed to warm to room temperature and was stirred for 25 minutes. Solvents were removed under reduced pressure, and the residue was triturated with Et_2O and filtered to give (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride. LC/MS: *m/z* 435.2 $(M+H)^+$ at 2.41 min (10%–99% CH_3CN (0.035 TFA)/H_2O (0.05% TFA)).


[001557] Example 302

[001558] **(*R,R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide**



[001559] **(*R, R*)-Tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide.**

[001560]




[001572] To 850 mg (2.02 mmol) of (*R*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester was added at room temperature 4 mL of 1:1 TFA: CH_2Cl_2. The reaction mixture was stirred for 50 min and diluted with 20 mL of CH_2Cl_2 and extracted with 15 mL of satd. $NaHCO_3$ solution. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol as an oil which was used without further purification.

To 52.6 mg (0.16 mmol) of the amine from above procedure was added 600 μL of CH_2Cl_2. To this solution was added sequentially 27.4 μL (0.19 mmol) of triethylamine and 20.9 mg (0.18 mmol) of (*R*)- tetrahydro-furan-2-carboxylic acid, 24.2 mg (0.18 mmol) of HOBt, 34.6 mg (0.18 mmol) of EDCI at room temperature. The reaction mixture was stirred for 12 h, then diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*R*, *R*)-tetrahydro-furan-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide as the TFA salt. LC/MS: *m/z* 419 $(M+H)^+$ at 2.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001573] Example 303


[001574] (2*R*)-Tetrahydro-*N*-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)furan-2-carboxamide

[001575]



[001576] (*S*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester


[001577]



[001578] To 2-(4-chloro-7-methyl-quinazolin-2-yl)-phenol(551 mg, 2.03 mmol) in 2.5 mL of DMF at room temperature was added sequentially (*S*)- pyrrolidin-3-yl-carbamic acid *tert*-butyl ester (740 mg, 3.9 mmol) and triethylamine (567 μL, 4.0 mmol), and the reaction mixture was stirred for 12 h. The reaction mixture was diluted with water (10 mL) and CH_2Cl_2 (10 mL). The organic layer was separated and dried (Na_2SO_4), and the residue was purified via silica gel chromatography with 25–85% ethyl acetate/hexanes to give of (*S*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester (694 mg, 81%). LC/MS: *m/z* 421 $(M+H)^+$ at 2.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001579] (2*R*)-Tetrahydro-*N*-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)furan-2-carboxamide

[001580]




[001581] To (*S*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester (690 mg, 1.64 mmol) was added at room temperature 3 mL of 1:1 TFA:CH_2Cl_2. The reaction mixture was stirred for 55 min, diluted with 20 mL of CH_2Cl_2 and washed with 15 mL of satd. $NaHCO_3$ solution. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give (*S*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol as an oil which was used without further purification.

To the amine from above procedure (133.3 mg, 0.42 mmol) was added 1.6 mL of CH_2Cl_2. To this solution was added sequentially triethylamine (174 μL, 1.25 mmol) and (*R*)-tetrahydro-furan-2-carboxylic acid (57.9 mg, 0.5 mmol), HOBt (67.5 mg 0.5 mmol), EDCI (95.8 mg 0.5 mmol) at room temperature. The reaction mixture was stirred for 12 h and diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (2*R*)-tetrahydro-*N*-((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 419 $(M+H)^+$ at 2.34 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001582] Example 304


[001583] **(*S*, *S*)- {1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid tetrahydro-furan-3-yl ester**

[001584]



[001585] **(*S*, *S*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid tetrahydro-furan-3-yl ester**

[001586]




[001587] To (*S*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester (690 mg, 1.64 mmol) was added at room temperature, 3 mL of 1:1 TFA:CH_2Cl_2. The reaction mixture was stirred for 55 min, diluted with 20 mL of CH_2Cl_2, and washed with 15 mL of satd. $NaHCO_3$ solution. The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give (*S*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol as an oil which was used without further purification.

To the amine from above procedure (134.7 mg, 0.42 mmol) was added 1.6 mL of CH_2Cl_2, and the solution was cooled to 0 °C. To this solution was added sequentially triethylamine (88 μL, 0.63 mmol) and (*S*)- tetrahydro-furan-3-ol chloroformate(82.3 mg, 0.55 mmol). The reaction mixture was stirred and allowed to warm from 0 °C to room temperature over 12 h and was diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*S*, *S*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid tetrahydro-furan-3-yl ester as the TFA salt. LC/MS: *m/z* 435 $(M+H)^+$ at 2.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001588] Example 305


[001589] **(*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001590]



[001591] **Benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate**


[001592]



[001593] A mixture of (*S*)-*tert*-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate (1.0 g, 5.0 mmol) and 50 mL THF was cooled in an ice bath. To this was added benzyl chloroformate (0.77 mL, 5.5 mmol), followed by triethylamine (1.39 mL, 10 mmol). After removing the ice bath, the reaction was stirred for 4 h. The mixture was poured into ice water and extracted with EtOAc. The organic extracts were washed with water, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using EtOAc in hexanes (0–40%) gave benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (1.29 g, 77%) as a colorless oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39–7.28 (m, 5H), 5.10 (s, 2H), 4.87 (s, 1H), 3.52–2.95 (m, 6H), 2.41–2.35 (m, 1H), 2.10–1.79 (m, 2H), 1.45 (s, 9H).

[001594] Benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[001595]




[001596] A solution of benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (0.20 g, 0.60 mmol) and 4 M HCl in dioxane (10 mL) was stirred for 3 h at room temperature. After evaporating the solvent under reduced pressure, the solid was triturated with Et_2O and dried under vacuum, then taken up in CH_2Cl_2 (10 mL). To this solution was added 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.16 g, 0.60 mmol) and triethylamine (0.25 mL, 1.8 mmol).

The reaction mixture was stirred at room temperature overnight, then diluted with CH_2Cl_2 and washed with water. The organic extracts were dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in hexanes gave benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as a colorless oil (0.19 g, 68%). LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.58 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001597] 2-(4-((*S*)-3-(Aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol


[001598]



[001599] A solution of benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (0.19 g, 0.41 mmol) and MeOH (5 mL) was stirred with Pd/C (20 mg, 10% weight of Pd on carbon) under an H_2 atmosphere at ambient pressure at ambient pressure overnight. Purification via silica gel chromatography using MeOH in CH_2Cl_2 (0–10%) gave 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (27 mg, 19%). LC/MS: *m/z* 335.5 $(M+H)^+$ at 1.28 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001600] (*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[001601]




[001602] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (13 mg, 0.04 mmol) and CH_2Cl_2 (0.5 mL) was cooled in an ice bath. To this mixture was added (*S*)-tetrahydrofuran-3-yl chloroformate (6 μL, 0.04 mmol), followed by triethylamine (11 μL, 0.08 mmol). After removing the ice bath, the reaction was stirred for 3 h at room temperature. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001603] Example 306


[001604] **(*R*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001605]



[001606] **Benzyl ((*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate**

[001607]




[001608] A solution of (*R*)-*tert*-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate (1.0 g, 5.0 mmol) and 50 mL THF was cooled in an ice bath. To this mixture was added benzyl

chloroformate (0.77 mL, 5.5 mmol), followed by triethylamine (1.39 mL, 10 mmol). After removing the ice bath, the reaction was stirred for 4 h. The mixture was poured into ice water and extracted with EtOAc. The organic extracts were washed with water, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using EtOAc in hexanes (0–40%) gave benzyl ((*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (1.29 g, 77%) as a colorless oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39–7.28 (m, 5H), 5.10 (s, 2H), 4.87 (s, 1H), 3.52–2.95 (m, 6H), 2.41–2.35 (m, 1H), 2.10–1.79 (m, 2H), 1.45 (s, 9H).

[001609] Benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate


[001610]



[001611] A solution of benzyl ((*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (0.20 g, 0.60 mmol) and 4 M HCl in dioxane (10 mL) was stirred for 3 h at room temperature. After evaporating the solvent under reduced pressure, the resulting solid was triturated with Et_2O and dried under vacuum, then taken up in CH_2Cl_2 (10 mL). To this solution was added 2-(4-chloro-7-methylquinazolin-2-yl)phenol (0.16 g, 0.60 mmol) and triethylamine (0.25 mL, 1.8 mmol). The reaction mixture was stirred at room temperature overnight, then diluted with CH_2Cl_2 and washed with water. The CH_2Cl_2 solution was dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in hexanes gave benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as a colorless oil (0.19 g, 68%). LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.58 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001612] 2-(4-((*R*)-3-(Aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol


[001613]



[001614] A solution of benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (0.19 g, 0.41 mmol) and MeOH (5 mL) was stirred with Pd/C (20 mg, 10% weight of Pd on carbon) under an H_2 atmosphere at ambient pressure overnight. Purification via silica gel chromatography using MeOH in CH_2Cl_2 (0–10%) gave 2-(4-((*R*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (63 mg, 45%). LC/MS: *m/z* 335.7 $(M+H)^+$ at 1.23 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001615] (*R*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[001616]




[001617] A solution of 2-(4-((*R*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (15 mg, 0.045 mmol) and CH_2Cl_2 (0.5 mL) was cooled in an ice bath. To this mixture was added (*R*)-tetrahydrofuran-3-yl chloroformate (7 μL, 0.045 mmol), followed by triethylamine (13 μL, 0.090 mmol). After removing the ice bath, the reaction was stirred at room temperature for 3 h. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*R*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-

methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001618] Example 307


[001619] **(*S*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001620]



[001621] **(*S*)-Tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001622]




[001623] A mixture of 2-(4-((*R*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (15 mg, 0.045 mmol) and CH_2Cl_2 (0.5 mL) was cooled in an ice bath. To this was added (*S*)-tetrahydrofuran-3-yl chloroformate (7 μL, 0.045 mmol), followed by triethylamine (13 μL, 0.090 mmol). After removing the ice bath, the reaction was stirred at room temperature for 3 h. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (*S*)-tetrahydrofuran-3-yl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-

yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.18 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001624] Example 308


[001625] **(2*R*)-Tetrahydro-*N*-(((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide**

[001626]



[001627] **(2*R*)-Tetrahydro-*N*-(((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide**

[001628]




[001629] To a solution of 2-(4-((*R*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (15 mg, 0.045 mmol) and DMF (0.5 mL) was added (*R*)-tetrahydrofuran-2-carboxylic acid (6 μL, 0.062 mmol), followed by the addition of HATU (26 mg, 67 mmol) and triethylamine (13 μL, 0.090 mmol). The reaction was stirred at room temperature for 3 h. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-tetrahydro-*N*-(((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 433.5 $(M+H)^+$ at 2.11 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001630] Example 309


[001631] **(*R*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester**

[001632]



[001633] **{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3*R*-yl}-carbamic acid *tert*-butyl ester**

[001634]




[001635] To a cooled (–15 °C) solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (43.8 g, 0.15 mol) in CH_2Cl_2 (125 mL) was added dropwise a solution of (3*R*)-(+)-3-(*tert*-butoxycarbonylamino)pyrrolidine (30 g, 0.16 mol) and triethylamine (38 mL, 0.27 mol) in CH_2Cl_2 (170 mL). The addition, during which the temperature stayed below 30 °C, was completed in 20 minutes. The external cooling was removed and the reaction mixture was stirred at room temperature overnight. Water (1 L) and CH_2Cl_2 (1 L) were added. The resulting precipitate was collected by filtration, washed with water and CH_2Cl_2, and air-dried to yield 33 g of pure {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3*R*-yl}-carbamic acid *tert*-butyl ester. The filtrates were evaporated to give another batch of crude {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3*R*-yl}-carbamic acid *tert*-butyl ester which was purified by trituration with CH_2Cl_2. Total yield: 43.5 g (70%). ^{1}H-NMR (300 MHz, $CDCl_3$): δ

8.45 (dd, J = 8 Hz, 1.7 Hz, 1H), 7.95 (d, J = 8.8 Hz, 1H), 7.57 (s, 1H), 7.37–7.31 (m, 1H), 7.18 (dd, J = 8.5 Hz, 1.7, 1H), 7.0 (dd, J = 8.3 Hz, 1.1 Hz, 1H), 6.9 (dt, J = 8.9 Hz, 1.4 Hz, 1H), 4.77 (bs, 1H), 4.40 (bs, 1H), 4.27–4.21 (m, 1H), 4.14–4.03 (m, 2H), 3.86 (dd, J = 11.8 Hz, 4.4 Hz, 1H), 2.50 (s, 3H), 2.34–2.28 (m, 1H), 2.08–2.02 (m, 1H), 1.46 (s, 9H) ppm.

[001636] **2-(4-(3*R* -Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol**

[001637]




[001638] {1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3*R*-yl}-carbamic acid *tert*-butyl ester (43.5 g, 100 mmol) was stirred with CF_3CO_2H (142 mL) in CH_2Cl_2 (300 mL) at room temperature overnight. The solution was concentrated to dryness, and the residue was suspended in 10% aq. Na_2CO_3 (900 mL) and stirred for 2 hours. The suspension was filtered, and the yellow solid was washed with water several times. The product was air-dried at 45 °C to yield 2-(4-(3*R*-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (34.2 g) as a light-yellow solid. ^{1}H-NMR (300 MHz, CD_3OD): δ 8.41 (dd, J = 7.4 Hz, 1.4 Hz, 1H), 8.09 (d, J = 8.5 Hz, 1H), 7.46 (s, 1H), 7.34–7.28 (m, 1H), 7.23 (dd, J = 8.8 Hz, 1.9 Hz, 1H), 6.91–6.85 (m, 2H), 4.16–4.09 (m, 2H), 4.00–3.94 (m, 1H), 3.73–3.64 (m, 2H), 2.48 (s, 3H), 2.26–2.20 (m, 1H), 1.92–1.86 (m, 1H) ppm.

[001639] **(*R*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester**

[001640] Method A

[001641]




[001642] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (50.8 mg, 0.16 mmol) was added 500 μL of CH_2Cl_2 and the solution was cooled to 0 °C. To this solution was added sequentially triethylamine (33.2 μL, 0.24 mmol) and benzyl chloroformate (29.2 mg, 0.17 mmol). The reaction mixture was stirred from 0 °C to 5 °C over 45 min, then diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give (*R*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester. LC/MS: *m/z* 455.2 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001643] Method B

[001644] **Benzyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**


[001645]



[001646] To *tert*-butyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (398 mg, 0.94 mmol) was added 3 mL of 1:1 TFA: CH_2Cl_2. The mixture was then stirred at room temperature for 30 minutes. The reaction mixture was diluted with a solution of saturated $NaHCO_3$ and CH_2Cl_2. The organic layer was separated, washed with a solution of extracted 1 N NaOH, dried over Na_2SO_4 and evaporated under reduced pressure to give the intermediate amine. To this amine (300 mg, 0.94 mmol) were added 3 ml of CH_2Cl_2 and triethylamine (145 μL, 1.04 mmol). After cooling the mixture to 0 °C, benzyl chloroformate (161.6 mg, 0.94 mmol) was added, and the reaction was stirred for 30 minutes. Purification via silica gel chromatography using 20–100% ethyl acetate/hexanes gave benzyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate. LC/MS: *m/z* 455.2 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001647] Benzyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride


[001648]



[001649] A 2.0 M HCl solution in Et_2O (318 μL, 0.636 mmol) was slowly added to a stirring solution of benzyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (289 mg, 0.636 mmol) in 2.1 mL of CH_2Cl_2. Solvents were removed under reduced pressure and the residue was triturated with Et_2O and filtered to give benzyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride. LC/MS: *m/z* 455.2 $(M+H)^+$ at 2.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001650] Example 310

[001651] (2*R*)-Tetrahydro-*N*-(((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide



[001652] (2*R*)-Tetrahydro-*N*-(((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide

[001653]




[001654] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (25 mg, 0.075 mmol) and DMF (0.5 mL) was added (*R*)-tetrahydrofuran-2-carboxylic acid (8.6 μL, 0.09 mmol), followed by the addition of HATU (34 mg, 0.09 mmol) and triethylamine (21 μL, 0.15 mmol). The reaction was stirred at room temperature for 3 h. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave (2*R*)-tetrahydro-*N*-(((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)furan-2-carboxamide as the TFA salt. LC/MS: *m/z* 433.5 $(M+H)^+$ at 2.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001655] Example 311

[001656] ***N*-(((*S*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide**

[001657]




[001658] ***N*-(((*S*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide**




[001648] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (25 mg, 0.075 mmol) and CH_2Cl_2 (0.5 mL) was cooled in an ice bath. To this mixture was added cyclopropanecarbonyl chloride (7.5 μL, 82 mmol), followed by triethylamine (21 μL, 0.15 mmol). After removing the ice bath, the reaction was stirred at room temperature for 3 h. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave *N*-(((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 403.7 $(M+H)^+$ at 2.17 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001649] Example 312

[001650] **(*R*)-Cyclopropanecarboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide**



[001651] **(*R*)-Cyclopropanecarboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide.**




[001666] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (49.3 mg, 0.15 mmol) was added 500 μL of CH_2Cl_2 and the solution was cooled to 0 °C. To this solution was added sequentially triethylamine (21.5 μL, 0.15 mmol) and cyclopropanecarbonyl chloride (16.1 mg, 0.15 mmol). The reaction mixture was stirred from 0 °C to 5 °C over 40 min and diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*R*)-cyclopropanecarboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide as the TFA salt. LC/MS: *m/z* 389 $(M+H)^+$ at 2.38 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001667] Example 313


[001668] **(*R*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester**

[001669]



[001670] **(*R*)-{1-[2-(2-Hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester.**

[001671]




[001672] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (48 mg, 0.15 mmol) was added 500 μL of CH_2Cl_2, and the solution was cooled to 0 °C. To this solution

was added sequentially triethylamine (21 μL, 0.15 mmol) and phenyl chloroformate (22.8 mg, 0.15 mmol). The reaction mixture was stirred from 0 °C to 5 °C over 40 min, diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*R*)-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester as the TFA salt. LC/MS: *m/z* 441 $(M+H)^+$ at 2.78 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001673] Example 314


[001674] **(Tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001675]



[001676] **(Tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate**


[001677]



[001678] To a mixture of (tetrahydro-2*H*-pyran-2-yl)methanol (369 mg, 3.17 mmol) and di(1*H*-imidazol-1-yl)methanone (1.03 g, 6.35 mmol) was added 10.5 mL of CH_2Cl_2. The reaction was stirred at 50 °C for 3 hours. The reaction mixture was used without further purification. LC/MS: *m/z* 211.1 $(M+H)^+$ at 0.94 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001679] **(Tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001680]




[001681] To a mixture of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (100 mg, 0.31 mmol) and (tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate (98 mg, 0.47 mmol) was added 1.04 mL CH_2Cl_2 and triethylamine (65 µL, 47 mg, 0.46 mmol). The mixture was stirred at room temperature overnight. An additional equivalent of (tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate (100 mg, 0.47 mmol) was added to the mixture, and the reaction was heated at 45 °C for 4 hours. Purification via silica gel chromatography using 10%–100% ethyl acetate/hexanes gave (tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate. LC/MS: *m/z* 463.4 $(M+H)^+$ at 2.66 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001682] Example 315


[001683] **(*R*)-5-Oxo-pyrrolidine-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide**

[001684]



[001685] **(*R*)-5-Oxo-pyrrolidine-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide**

[001686]




[001687] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (71 mg, 0.22 mmol) was added 890 μL of CH_2Cl_2. To this solution was added sequentially triethylamine (47 μL, 0.34 mmol) and 5-oxo-pyrrolidine-2-carboxylic acid (34.8 mg, 0.27 mmol) and BOP (119 mg, 0.27 mmol). The reaction mixture was stirred at room temperature for 2 h and diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure to give (*R*)-5-oxo-pyrrolidine-2-carboxylic acid {1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-amide. LC/MS: *m/z* 432.5 $(M+H)^+$ at 2.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001688] Example 316


[001689] **Tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2*H*-pyran-4-carboxamide**

[001690]



[001691] **Tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2*H*-pyran-4-carboxamide**

[001692]




[001693] To a stirred solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in 1 mL of DMF was cooled to 0 °C and tetrahydro-2*H*-pyran-4-carboxylic acid (24 mg, 0.19 mmol) was added, followed by the addition of triethylamine (32 mg, 44 μL, 0.31 mmol) and HATU (71.1 mg, 0.187 mmol). The reaction was stirred at 0 °C for 10 minutes, then gradually warmed to room temperature, then filtered. Purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave tetrahydro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2*H*-pyran-4-carboxamide as the TFA salt. LC/MS: *m/z* 433.5 $(M+H)^+$ at 2.05 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001694] Example 317


[001695] **2-(Tetrahydro-2*H*-pyran-4-yl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide**

[001696]



[001697] **2-(Tetrahydro-2*H*-pyran-4-yl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide**

[001698]




[001699] To a stirred solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in 1 mL of DMF cooled to 0 °C was added 2-(tetrahydro-2*H*-pyran-4-yl)acetic acid (27 mg, 0.19 mmol), followed by the addition of triethylamine (32 mg, 44 μL, 0.31 mmol) and HATU (71.1 mg, 0.187 mmol). The reaction was stirred at 0 °C for 10 minutes, then gradually warmed to room temperature. Filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 2-(tetrahydro-2*H*-pyran-4-yl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide as the TFA salt. LC/MS: *m/z* 477.3 $(M+H)^+$ at 2.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001700] Example 318


[001701] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-2-yl)propanamide**

[001702]



[001703] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-2-yl)propanamide**

[001704]




[001705] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.05 g, 0.15 mmol) in DMF (1 mL) was added 3-(pyridin-2-yl)propanoic acid (30 mg, 0.195 mmol), followed by the addition of triethylamine (42 μL, 0.30 mmol) and HATU (74 mg, 0.195 mmol). The reaction was stirred at room temperature for 2 h and then purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-2-yl)propanamide as the TFA salt. LC/MS: *m/z* 454.3 $(M+H)^+$ at 1.79 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001706] Example 319


[001707] **(Pyridin-3-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001708]



[001709] (Pyridin-3-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001710]




[001711] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in DMSO (0.5 mL) at room temperature was added triethylamine (43 μL, 0.31 mmol), followed by the addition of (pyridin-3-yl)methyl 1*H*-imidazole-1-carboxylate (63 mg, 0.31 mmol). The reaction was stirred at room temperature overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (pyridin-3-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 456.5 $(M+H)^+$ at 1.85 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001712] Example 320


[001713] (Pyridin-4-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001714]



[001715] (Pyridin-4-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001716]




[001717] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in DMSO (0.5 mL) at room temperature was added triethylamine (43 μL, 0.31 mmol), followed by the addition of (pyridin-4-yl)methyl 1*H*-imidazole-1-carboxylate (63 mg, 0.31 mmol). The reaction was stirred at room temperature overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (pyridin-4-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 456.5 $(M+H)^+$ at 1.84 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001718] Example 321


[001719] (Benzo[d][1,3]dioxol-7-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001720]



[001721] **(Benzo[d][1,3]dioxol-7-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001722]




[001723] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in DMSO (0.5 mL) was added triethylamine (43 µl, 0.31 mmol), followed by the addition of (benzo[d][1,3]dioxol-4-yl)methyl 1*H*-imidazole-1-carboxylate (77 mg, 0.31 mmol). The reaction was stirred at room temperature overnight, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (benzo[d][1,3]dioxol-7-yl)methyl (*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 499.3 $(M+H)^+$ at 2.57 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001724] Example 322


[001725] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-3-yl)propanamide**

[001726]



[001727] *N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-3-yl)propanamide

[001728]




[001729] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (249 mg, 0.778 mmol) was dissolved in 2.6 mL of CH_2Cl_2. 3-(Pyridin-3-yl)propanoic acid (129 mg, 0.85 mmol) was added followed by triethylamine (102 mg, 141 μL, 1.01 mmol) and BOP (378 mg, 0.85 mmol). After stirring the reaction for 35 minutes at room temperature, it was filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(pyridin-3-yl)propanamide as the TFA salt. LC/MS: *m/z* 454.4 $(M+H)^+$ at 2.08 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001730] Example 323


[001731] (*R*)-3-Cyclopentyl-N-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-propionamide

[001732]



[001733] (*R*)-3-Cyclopentyl-N-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-propionamide

[001734]




[001735] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (100 mg, 0.31 mmol) was added 1 mL of CH_2Cl_2, and the solution was cooled to 0 °C. To this solution was added sequentially triethylamine (56.6 μL, 0.41 mmol) and 3-cyclopentyl-propionyl chloride (57 mg, 0.35 mmol). The reaction mixture was stirred at room temperature for 2 h, then diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give ((*R*)-3-cyclopentyl-*N*-{1-[2-(2-hydroxy-phenyl)-7-methyl-quinazolin-4-yl]-pyrrolidin-3-yl}-propionamide as the TFA salt. LC/MS: *m/z* 445.4 $(M+H)^+$ at 2.85 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001736] Example 324


[001737] **(*R*)-N-{1-[2-(2-Hydroxy-phenyl)-quinazolin-4-yl]-pyrrolidin-3-yl}-3-piperidin-1-yl-propionamide**

[001738]



[001739] **(*R*)-N-{1-[2-(2-Hydroxy-phenyl)-quinazolin-4-yl]-pyrrolidin-3-yl}-3-piperidin-1-yl-propionamide**

[001740]




[001741] To (*R*)-2-[4-(3-amino-pyrrolidin-1-yl)-7-methyl-quinazolin-2-yl]-phenol (58.6 mg, 0.18 mmol) was added 700 μL of CH_2Cl_2. To this solution was added sequentially triethylamine (38.3 μL, 0.27 mmol), 3-piperidin-1-yl-propionic acid (37.4 mg, 0.24 mmol) and BOP (119 mg, 0.27 mmol). The reaction mixture was stirred at room temperature for 2 h and diluted with water and CH_2Cl_2 (10 mL). The organic layer was separated and dried over Na_2SO_4, and the solvent was removed under reduced pressure. The residue was purified via preparative reverse phase HPLC using 10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA) to give (*R*)-*N*-{1-[2-(2-hydroxy-phenyl)-quinazolin-4-yl]-pyrrolidin-3-yl}-3-piperidin-1-yl-propionamide as the TFA salt. LC/MS: *m/z* 460.4 $(M+H)^+$ at 2.1 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001742] Example 325


[001743] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(trifluoromethoxy)benzamide**

[001744]



[001745] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(trifluoromethoxy)benzamide**

[001746]




[001747] To a stirring solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (32 mg, 0.10 mmol) and DMF (1.0 mL) was added 3-(trifluoromethoxy)benzoyl chloride (19 μL, 0.10 mmol), followed by the addition of triethylamine (28 μL, 0.2 mmol). The reaction was stirred at room temperature overnight, filtered, and purified via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-(trifluoromethoxy)benzamide as the TFA salt. LC/MS: *m/z* 509.5 $(M+H)^+$ at 2.71 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001748] Example 326


[001749] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-methoxybenzamide**

[001750]



[001751] *N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-methoxybenzamide

[001752]




[001753] To a stirring solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (32 mg, 0.10 mmol) and DMF (1.0 mL) was added 3-methoxybenzoyl chloride (14 μL, 0.1 mmol), followed by the addition of triethylamine (28 μL, 0.2 mmol). The reaction was stirred at room temperature overnight, then filtered, and purified via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-3-methoxybenzamide as the TFA salt. LC/MS: *m/z* 455.3 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001754] Example 327


[001755] 3-Cyano-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide

[001756]



[001757] **3-Cyano-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001758]




[001759] To a stirring solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (32 mg, 0.10 mmol) and DMF (1.0 mL) was added 3-cyanobenzoyl chloride (17 mg, 0.10 mmol), followed by the addition of triethylamine (28 μL, 0.20 mmol). The reaction was stirred at room temperature overnight, then filtered, and purified via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 3-cyano-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 450.3 $(M+H)^+$ at 2.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001760] Example 328


[001761] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001762]



[001763] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001764]




[001765] To a stirring solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (32 mg, 0.1 mmol) and DMF (1.0 mL) was added nicotinoyl chloride (18 mg, 0.1 mmol), followed by the addition of triethylamine (28 μL, 0.2 mmol). The reaction was stirred at room temperature overnight, then filtered, and purified via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide as the TFA salt. LC/MS: *m/z* 426.3 $(M+H)^+$ at 1.91 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001766] Example 329

[001767] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide**


[001768]



[001769] Method A

[001770] **2-(4-Chloro-6-fluoroquinazolin-2-yl)phenol**


[001771]



[001772] A solution of 4-chloro-6-fluoro-2-(2-methoxyphenyl)quinazoline (3.0 g, 10.39 mmol) in CH_2Cl_2 (15 mL) was cooled to –78 °C. To it was added a 1.0 M BBr_3 solution in CH_2Cl_2 (52 mL, 52 mmol) dropwise. The reaction was allowed to warm to room temperature. It was neutralized with a saturated aqueous $NaHCO_3$ solution, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 5–20% CH_2Cl_2/hexanes gave 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (1.68 g, 60%). LC/MS: *m/z* 275.3 $(M+H)^+$ at 3.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001773] ***tert*-Butyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**


[001774]



[001775] At 0 °C under an N_2 atmosphere, a solution of *tert*-butyl (*R*)-pyrrolidin-3-ylcarbamate (264 mg, 1.42 mmol) and triethylamine (0.33 mL, 2.36 mmol) in CH_2Cl_2 was rapidly added to a stirring solution of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (325 mg, 1.18 mmol) in 15 mL CH_2Cl_2. The reaction mixture was stirred for 1 h before it was quenched with water, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc/CH_2Cl_2 yielded *tert*-butyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate. LC/MS: *m/z* 425.5 $(M+H)^+$ at 2.74 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001776] **2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol**


[001777]



[001778] To a solution of *tert*-butyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate (480 mg, 1.11 mmol) in CH_2Cl_2 (10 mL) was added TFA (4 mL). The reaction was stirred for an hour, diluted with 10 mL CH_2Cl_2, and neutralized with a saturated aqueous $NaHCO_3$ solution. The aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 3–20% EtOAc/CH_2Cl_2 gave 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol. LC/MS: *m/z* 325.5 $(M+H)^+$ at 1.26 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001779] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide**

[001780]




[001781] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.025 g, 0.08 mmol) in DMF (1.0 mL) was added cyclopropanecarboxylic acid (10 mg, 0.12 mmol), followed by the addition of triethylamine (22 μL, 0.16 mmol) and HATU (40 mg, 0.1 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-

fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 393.3 $(M+H)^+$ at 2.23 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001782] Method B

[001783] **Benzyl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate**


[001784]



[001785] At –10 °C, triethylamine (2.3 mL, 16.6 mmol) was added to a solution of benzyl (*R*)-pyrrolidin-3-ylcarbamate oxalate (2.0 g, 6.4 mmol) in MeOH, followed by the slow addition of Boc_2O (1.92 mL, 8.3 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. The mixture was quenched with water and extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in CH_2Cl_2 gave benzyl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate (1.85 g, 90%). LC/MS: *m/z* 321.3 $(M+H)^+$ at 3.01 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001786] **(*R*)-*tert*-Butyl 3-aminopyrrolidine-1-carboxylate**

[001787]




[001788] Under an N_2 atmosphere, a solution of benzyl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate (1.85 g, 5.75 mmol) in 10 mL MeOH was added to a

flask containing Pd/C (185 mg, 10% weight Pd on carbon). After evacuating the flask under vacuum and purging it twice with N_2, the reaction was stirred for 3 h under an H_2 atmosphere at ambient pressure. The reaction was filtered through a bed of Celite, and the solvent was evaporated under reduced pressure to obtain (*R*)-*tert*-butyl 3-aminopyrrolidine-1-carboxylate which was used without further purification. LC/MS: *m/z* 187.3 $(M+H)^+$ at 1.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001789] **(*R*)-*tert*-Butyl 3-(cyclopropanecarboxamido)pyrrolidine-1-carboxylate**


[001790]



[001791] To a mixture of (*R*)-*tert*-butyl 3-aminopyrrolidine-1-carboxylate (500 mg, 2.68 mmol) in CH_2Cl_2 (10 mL) was added cyclopropanecarboxylic acid (276 μL, 3.48 mmol), followed by the addition of HATU (1.3 g, 3.48 mmol) and triethylamine (725 μL, 5.2 mmol). The reaction was complete after stirring for 1 h. After quenching with water, the aqueous layer was extracted twice with CH_2Cl_2, and the combined organic extracts were dried over $MgSO_4$, filtered, and concentrated. The crude material was purified via silica gel chromatography 0–20% EtOAc/CH_2Cl_2 to afford (*R*)-*tert*-butyl 3-(cyclopropanecarboxamido)pyrrolidine-1-carboxylate (500 mg, 73%). LC/MS: *m/z* 255.3 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001792] ***N*-((*R*)-Pyrrolidin-3-yl)cyclopropanecarboxamide**

[001793]




[001794] TFA (1 mL) was added to a solution of (*R*)-*tert*-butyl 3-(cyclopropanecarboxamido)pyrrolidine-1-carboxylate (500 mg, 1.96 mmol) in 5 mL CH_2Cl_2. After stirring for 30 min, the reaction was quenched with 1M NaOH solution till neutral and

extracted twice with EtOAc. The organic extracts were combined, dried over $MgSO_4$, filtered, and concentrated to yield *N*-((*R*)-pyrrolidin-3-yl)cyclopropanecarboxamide (250 mg) which was used without further purification. LC/MS: *m/z* 155.3 $(M+H)^+$ at 0.6 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001795] *N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide

[001796]



[001797] A mixture of *N*-((*R*)-pyrrolidin-3-yl)cyclopropanecarboxamide (250 mg, 0.86 mmol), 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (250 mg, 0.86 mmol) and triethylamine (0.240 mL, 1.72 mmol) in CH_2Cl_2 (10 mL) was stirred at room temperature. The reaction was complete after one hour. The reaction was quenched with water, the aqueous layer was extracted twice with CH_2Cl_2, and the combined organic extracts were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc/CH_2Cl_2 gave *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide (230 mg, 68%). LC/MS: *m/z* 393.3 $(M+H)^+$ at 2.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.42–8.46 (m, 2H), 8.03 (dd, *J* = 10.5, 2.6 Hz, 1H), 7.88–7.92 (m, 1H), 7.74–7.79 (m, 1H), 7.35–7.39 (m, 1H), 6.92–6.96 (m, 2H), 4.44–4.47 (m, 1H), 4.01–4.28 (m, 3H), 3.83–3.87 (m, 1H), 2.20–2.28 (m, 1H), 1.99–2.06 (m, 1H), 1.51–1.57 (m, 1H), 0.63–0.73 (m, 4H).

[001798] *N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide hydrochloride




[001799]

[001800] To a solution of *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide (225 mg, 0.57 mmol) in 5 mL CH_2Cl_2 was added a 2 M HCl solution in ether (0.28 mL, 0.57 mmol), which resulted in the precipitation of a solid. After the addition of 20 mL ether, the reaction mixture was stirred for 1 h. The solvents were evaporated under reduced pressure to afford *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide hydrochloride (225 mg, 91%). LC/MS: *m/z* 393.3 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.23 (d, *J* = 8.3 Hz, 1H), 8.04 (d, *J* = 10.6 Hz, 1H), 7.91–7.94 (m, 1H), 7.79–7.83 (m, 1H), 7.44–7.48 (m, 1H), 7.00–7.03 (m, 2H), 4.44 (t, *J* = 4.9 Hz, 1H), 4.09–4.23 (m, 3H), 3.87–3.90 (m, 1H), 2.25–2.34 (m, 1H), 2.02–2.09 (m, 1H), 1.49–1.55 (m, 1H), 0.68–0.71 (m, 4H).

[001801] Example 330


[001802] **(2*S*)-*N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-phenylcyclopropanecarboxamide**

[001803]



[001804] **(2*S*)-*N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-phenylcyclopropanecarboxamide**

[001805]




[001806] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. (2*R*)-2-Phenylcyclopropanecarbonyl chloride (31 mg, 0.17 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (2*S*)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-phenylcyclopropanecarboxamide as the TFA salt. LC/MS (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)), LC/MS: *m/z* 465.4 $(M+H)^+$ at 2.88 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001807] Example 331


[001808] **2-Chloro-6-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001809]



[001810] **2-Chloro-6-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001811]




[001812] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 2-Chloro-6-fluorobenzoyl chloride (36 mg, 0.18 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 2-chloro-6-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 477.3 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001813] Example 332


[001814] **4-Fluoro-3-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001815]



[001816] **4-Fluoro-3-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001817]




[001818] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 4-Fluoro-3-(trifluoromethyl)benzoyl chloride (42 mg, 0.18 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 4-fluoro-3-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 511.5 $(M+H)^+$ at 3.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001819] Example 333


[001820] **3-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001821]



[001822] **3-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001823]




[001824] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 3-Fluorobenzoyl chloride (30 mg, 0.18 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 3-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 443.5 $(M+H)^+$ at 2.83 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001825] Example 334


[001826] **3-Fluoro-4-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001827]



[001828] **3-Fluoro-4-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001829]




[001830] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 3-Fluoro-4-(trifluoromethyl)benzoyl chloride (42 mg, 0.18 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 3-fluoro-4-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 511.5 $(M+H)^+$ at 3.1 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001831] Example 335


[001832] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001833]



[001834] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001835]




[001836] A solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.092 mmol) in DMF (1.0 mL) was cooled to –40 °C. To this mixture was added triethylamine (26 μL, 0.184 mmol), followed by the addition of (*S*)-tetrahydrofuran-3-yl chloroformate (0.014 g, 0.092 mmol). After allowing the reaction to warm to room temperature the mixture was filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*S*)-tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.25 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001837] Method B

[001838] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate**

[001839]




[001840] To a solution of (*R*)-*tert*-butyl 3-aminopyrrolidine-1-carboxylate (500 mg, 2.6 mmol) in 5 mL CH_2Cl_2 was added triethylamine (0.73 mL, 5.2 mmol), and the reaction was cooled to –20 °C. (*S*)-tetrahydrofuran-3-yl chloroformate (525 mg, 3.48 mmol) was added in portions over a period of 10 minutes to the above reaction mixture. After allowing the reaction to warm to room temperature, it was quenched with water and extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to afford (*S*)-

tetrahydrofuran-3-yl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate. Purification via silica gel chromatography using 0–20% EtOAc in CH_2Cl_2 gave (*S*)-tetrahydrofuran-3-yl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate (490 mg, 63%). LC/MS: *m/z* 301.3 $(M+H)^+$ at 2.35 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001841] (*S*)-Tetrahydrofuran-3-yl (*R*)-pyrrolidin-3-ylcarbamate


[001842]



[001843] TFA (1 mL) was added to a solution of (*S*)-tetrahydrofuran-3-yl (*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-ylcarbamate (490 mg, 1.63 mmol) in 5 mL CH_2Cl_2. After stirring for 30 min, the reaction was quenched with NaOH and extracted twice with EtOAc. The organic extracts were combined, dried over $MgSO_4$, filtered, and concentrated to yield (*S*)-tetrahydrofuran-3-yl (*R*)-pyrrolidin-3-ylcarbamate (230 mg) which was used without further purification. LC/MS: *m/z* 201.3 $(M+H)^+$ at 0.59 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001844] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001845]




[001846] A mixture of (*S*)-tetrahydrofuran-3-yl (*R*)-pyrrolidin-3-ylcarbamate (225 mg, 1.12 mmol), 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (250 mg, 0.86 mmol), and triethylamine (0.240 mL, 1.72 mmol) in CH_2Cl_2 (10 mL) was stirred at room temperature. The reaction was complete in one hour. The reaction was quenched with water, the aqueous layer was extracted twice with CH_2Cl_2, and the combined organic extracts were dried over $MgSO_4$, filtered, and

concentrated. Purification via silica gel chromatography using 0–10% EtOAc/CH_2Cl_2 gave (*S*)-tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate (250 mg, 66%). LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.40 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.99 (d, *J* = 2.6 Hz, 1H), 7.87–7.91 (m, 1H), 7.74–7.79 (m, 1H), 7.70 (d, *J* = 6.2 Hz, 1H), 7.35–7.39 (m, 1H), 6.91–6.96 (m, 2H), 5.15 (s, 1H), 4.21–4.25 (m, 2H), 4.12–4.14 (m, 1H), 4.01–4.06 (m, 1H), 3.87–3.89 (m, 1H), 3.65–3.78 (m, 4H), 2.02–2.26 (m, 3H), 1.82–1.89 (m, 1H).

[001847] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride

[001848]




[001849] To a mixture of (*S*)-tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate (250 mg, 0.57 mmol) and CH_2Cl_2 (25 mL) was added a 2.0 M HCl solution in ether (0.285 mL, 0.57 mmol). After the addition of ether (40 mL), the reaction was stirred for 1 h. The resulting solid was filtered and dried to afford (*S*)-tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.25 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.18 (dd, *J* = 8.1, 1.4 Hz, 1H), 8.00 (d, *J* = 9.6 Hz, 1H), 7.90–7.94 (m, 1H), 7.78–7.83 (m, 1H), 7.45–7.49 (m, 1H), 7.00–7.03 (m, 2H), 5.10 (s, 1H), 4.09–4.25 (m, 4H), 3.90–3.92 (m, 1H), 3.62–3.75 (m, 4H), 2.22–2.27 (m, 1H), 2.03–2.14 (m, 2H), 1.83–1.91 (m, 1H).

[001850] Example 336


[001851] 3-(Trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide

[001852]



[001853] **3-(Trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001854]




[001855] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (130 mg, 0.406 mmol) was dissolved in 0.7 mL anhydrous CH_2Cl_2 and cooled to 0 °C. 3-(Trifluoromethyl)benzoyl chloride (42 mg, 0.18 mmol) dissolved in 0.7 mL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (53 mg, 74 μL, 0.52 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 3-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 493.5 $(M+H)^+$ at 3.03 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001856] Example 337


[001857] **(*R*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001858]



[001859] **(*R*)-Tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001860]




To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.092 mmol) in DMF (1.0 mL) was cooled to –40 °C, triethylamine (26 μL, 0.184 mmol) was added, followed by the addition of (*R*)-tetrahydrofuran-3-yl chloroformate (0.014 g, 0.092 mmol). After warming to room temperature, the mixture was filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain (*R*)-tetrahydrofuran-3-yl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.25 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001861] Example 338


[001862] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(tetrahydro-2*H*-pyran-4-yl)acetamide**

[001863]



[001864] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(tetrahydro-2*H*-pyran-4-yl)acetamide**

[001865]




To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.092 mmol) in DMF (1.0 mL) was added 2-(tetrahydro-2*H*-pyran-4-yl)acetic acid (0.017 g, 0.12 mmol), followed by the addition of triethylamine (25.6 μL, 0.184 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(tetrahydro-2*H*-pyran-4-yl)acetamide as the TFA salt. LC/MS: *m/z* 451.5 $(M+H)^+$ at 2.15 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001866] Example 339


[001867] **(Tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001868]



[001869] **(Tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001870]




[001871] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in DMSO (0.5 mL) at room temperature was added triethylamine (43 μL, 0.31 mmol), followed by the addition of (tetrahydro-2*H*-pyran-2-yl)methyl 1*H*-imidazole-1-carboxylate (39 mg, 0.13 mmol). The reaction was stirred at room temperature overnight, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (tetrahydro-2*H*-pyran-2-yl)methyl (*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 467.3 $(M+H)^+$ at 3.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001872] Example 340


[001873] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2,2,3,3-tetramethylcyclopropanecarboxamide**

[001874]



[001875] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2,2,3,3-tetramethylcyclopropanecarboxamide**

[001876]




[001877] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.156 mmol) was dissolved in 0.52 mL of anhydrous CH_2Cl_2. 2,2,3,3-Tetramethylcyclopropanecarboxylic acid (26.63 mg, 0.18 mmol) was added, followed by triethylamine (22.14 mg, 30.49 μL, 0.22 mmol) and BOP (82.93 mg, 0.18 mmol). After stirring the mixture for 30 minutes at room temperature, it was filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2,2,3,3-tetramethylcyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 445.4 $(M+H)^+$ at 2.95 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001878] Example 341


[001879] **4-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001880]



[001881] **4-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**


[001882]



[001883] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 4-Fluorobenzoyl chloride (25 mg, 0.15 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (19 mg, 26 μL, 0.18 mmol). The reaction was stirred for 15 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 4-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 443.4 $(M+H)^+$ at 2.73 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001884] Example 342


[001885] **3-(Dimethylamino)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**



[001886]

[001887] **3-(Dimethylamino)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**


[001888]



[001889] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 3-(Dimethylamino)benzoyl chloride hydrochloride (34 mg, 0.15 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (32 mg, 44 μL, 0.31 mmol). The reaction was stirred for 1.5 hour at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 3-(dimethylamino)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 468.4 $(M+H)^+$ at 2.39 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001890] Example 343


[001891] **2-(Trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**



[001892]

[001893] **2-(Trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**


[001894]



[001895] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 2-(Trifluoromethyl)benzoyl chloride (39 mg, 0.18 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 20 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 2-(trifluoromethyl)-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 493.4 $(M+H)^+$ at 2.76 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001896] Example 344


[001897] **2-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**



[001898]


[001899] **2-Fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide**

[001900]



[001901] 2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50.6 mg, 0.158 mmol) was dissolved in 260 μL of anhydrous CH_2Cl_2 and cooled to 0 °C. 2-Fluorobenzoyl chloride (27 mg, 0.17 mmol) dissolved in 260 μL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (21 mg, 28 μL, 0.20 mmol). The reaction was stirred for 15 minutes at 0 °C and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 2-fluoro-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)benzamide as the TFA salt. LC/MS: *m/z* 443.4 $(M+H)^+$ at 2.69 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001902] Example 345




[001903] **(*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[001904]

[001905] ***tert*-Butyl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**


[001906]



[001907] A solution of *tert*-butyl (*R*)-pyrrolidin-3-ylcarbamate (368 mg, 1.97 mmol) and triethylamine (0.46 mL, 3.28 mmol) in CH_2Cl_2 was rapidly added to a stirring solution of 2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol (475 mg, 1.65 mmol) in 15 mL CH_2Cl_2 at 0 °C under an N_2 atmosphere. The reaction was stirred for 1 h before it was quenched with water, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc/CH_2Cl_2 yielded *tert*-butyl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001908] **2-(4-((*R*)-3-Aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol**


[001909]



[001910] To a solution of *tert*-butyl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (500 mg, 1.14 mmol) in CH_2Cl_2 (15 mL) was added TFA (5 mL). The mixture was stirred for 1 h and then neutralized with a 1 M NaOH solution, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 3–20% EtOAc/CH_2Cl_2 gave 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol. LC/MS: *m/z* 339.5 $(M+H)^+$ at 0.56 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001911] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate

[001912]



[001913] Method A

[001914] To a stirred solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.12 mmol) in 800 μL of anhydrous DMF cooled to 0 °C was added (*S*)-tetrahydrofuran-3-yl chloroformate (20 mg, 19 μL, 0.13 mmol) dropwise, followed by the addition of triethylamine (24 mg, 33 μL, 0.23 mmol). The reaction was warmed to room temperature and stirred overnight, and the product purified via reverse phase HPLC (10%–99%


CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate as the TFA salt. LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.05 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001915] Method B

[001916] At room temperature, *N*-ethyl-*N*-isopropylpropan-2-amine (155 mL, 0.88 mmol) was added to a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (150 mg, 0.40 mmol) in THF. The mixture was cooled in an ice bath, and (*S*)-tetrahydrofuran-3-yl chloroformate (63 mg, 0.42 mmol) was added slowly over a period of 10 minutes. After warming to room temperature, the reaction was quenched with water and extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 0–10% EtOAc in CH_2Cl_2/hexanes (1:1) gave (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (160 mg, 84%). LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.12 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.18 (d, J = 8.6 Hz, 1H), 7.70 (d, J = 6.3 Hz, 1H), 7.58 (s, 1H), 7.29–7.38 (m, 2H), 6.76 (d, J = 8.3 Hz, 1H), 6.67–6.72 (m, 1H), 5.14 (s, 1H), 4.23–4.24 (m, 1H), 3.99–4.13 (m, 3H), 3.64–3.85 (m, 5H), 2.50 (s, 3H), , 2.07–2.22 (m, 2H), 2.00–2.03 (m, 1H), 1.85–1.90 (m, 1H).

[001917] (*S*)-Tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride

[001918]




[001919] To a solution of (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate (160 mg, 0.35 mmol) in 2 mL CH_2Cl_2 was added 2 M HCL solution in ether (0.176 mL, 0.35 mmol) resulting in precipitation of a solid. After the addition of 10 mL ether, the reaction was stirred for 30 minutes, filtered and the

resulting solid was dried to obtain (*S*)-tetrahydrofuran-3-yl (*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate hydrochloride (130 mg, 76%).). LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.13 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.29 (s, 1H), 7.57 (s, 2H), 7.43–7.49 (m, 1H), 6.83–6.89 (m, 2H), 5.08 (s, 1H), 4.23–4.42 (m, 4H), 3.62–3.73 (m, 5H), 2.24–2.34 (m, 1H), 2.04–2.11 (m, 2H), 1.86–1.92 (m, 1H).

[001920] Example 346


[001921] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide**

[001922]



[001923] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide**

[001924]



[001925] Method A

[001926] To a stirred solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (40 mg, 0.12 mmol) in 800 μL of anhydrous DMF at 0 °C was added cyclopropanecarboxylic acid (11 mg, 0.13 mmol), followed by the addition of triethylamine (24


mg, 33 μL, 0.24 mmol) and HATU (60 mg, 0.16 mmol). The mixture was allowed to warm to room temperature and was stirred overnight. Filtration and purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.2 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001927] Method B

To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (100 mg, 0.295 mmol) in 10 mL DMF, at –20 °C, was added cyclopropanecarboxylic acid (23 μL, 0.30 mmol), followed by the addition of triethylamine (82 μL, 0.59 mmol) and a solution of HATU (124 mg, 0.32 mmol) in 4 mL DMF. The mixture was warmed to room temperature and stirred for 1 h. Cold water was added to the reaction mixture which resulted in the formation of a precipitate which was collected by filtration and dissolved in CH_2Cl_2. The solution was then dried over $MgSO_4$, filtered, and concentrated to afford *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide (80 mg, 66%) LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.08 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001928] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide hydrochloride**

[001929]




[001930] A 2 M HCl solution in ether (0.16 mL, 0.32 mmol) was added to a solution of *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide (133 mg, 0.32 mmol) in CH_2Cl_2 (4 mL) under an N_2 atmosphere. Additional ether was then added (15 mL) and the reaction mixture was stirred for an hour. The formed precipitate was then filtered and dried to afford *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-

7-methylquinazolin-4-yl)pyrrolidin-3-yl)cyclopropanecarboxamide hydrochloride (135 mg, 95%). LC/MS: *m/z* 407.5 $(M+H)^+$ at 2.07 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.32 (d, J = 7.6 Hz, 1H), 7.56–7.58 (m, 2H), 7.43–7.49 (m, 1H), 6.83–6.88 (m, 2H), 3.94–4.41 (m, 5H), 2.52 (s, 3H), 2.25–2.36 (m, 1H), 1.91–2.13 (m, 1H), 1.47–1.54 (m, 1H), 0.63–0.69 (m, 4H)

[001931] Example 347


[001932] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001933]



[001934] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001935]




[001936] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added isonicotinic acid (0.015 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-fluoro-2-(2-

hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide as the TFA salt. LC/MS: *m/z* 430.5 $(M+H)^+$ at 1.95 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001937] Example 348


[001938] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001939]



[001940] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001941]




[001942] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added picolinic acid (0.015 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)picolinamide as the TFA salt. LC/MS: *m/z* 430.5 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001943] Example 349


[001944] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001945]



[001946] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001947]




[001948] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added nicotinic acid (0.015 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)nicotinamide as the TFA salt. LC/MS: *m/z* 430.5 $(M+H)^+$ at 1.98 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001949] Example 350


[001950] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001951]



[001952] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001953]




[001954] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) in DMF (0.5 mL) was added isonicotinic acid (0.014 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide as the TFA salt. LC/MS: *m/z* 444.5 $(M+H)^+$ at 1.85 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001955] Example 351


[001956] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001957]



[001958] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001959]




[001960] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added picolinic acid (0.014 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide as the TFA salt. LC/MS: *m/z* 444.5 $(M+H)^+$ at 2.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001961] Example 352


[001962] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001963]



[001964] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide**

[001965]




[001966] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added nicotinic acid (0.014 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)nicotinamide as the TFA salt. LC/MS: *m/z* 444.5 $(M+H)^+$ at 1.89 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001967] Example 353


[001968] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001969]



[001970] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide**

[001971]




[001972] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added isonicotinic acid (0.015 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)isonicotinamide as the TFA salt. LC/MS: *m/z* 426.1 $(M+H)^+$ at 1.93 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001973] Example 354


[001974] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001975]



[001976] ***N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide**

[001977]




[001978] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added picolinic acid (0.015 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)picolinamide as the TFA salt. LC/MS: *m/z* 426.1 $(M+H)^+$ at 2.33 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001979] Example 355


[001980] **(*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001981]



[001982] **Benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate**


[001983]



[001984] To a stirring solution of (*S*)-*tert*-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate (1.5 g, 7.5 mmol) in 25 mL CH_2Cl_2 was added triethylamine (2.1 mL, 15 mmol) at 0 °C, followed by the dropwise addition of benzyl chloroformate (1.58 mL, 11.2 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. The mixture was quenched with water, and the aqueous layer was extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 3% MeOH in CH_2Cl_2 gave benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (1.5 g, 60%). LC/MS: *m/z*: 335.5 $(M+H)^+$ at 3.01 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001985] **Benzyl ((*R*)-pyrrolidin-3-yl)methylcarbamate**


[001986]



[001987] To a stirred solution of ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (1.5 g, 4.48 mmol) in CH_2Cl_2 (20 mL) was slowly added TFA (5 mL). The reaction was stirred for 2 h. After removing the solvents under reduced pressure, the mixture was neutralized with a 1 M NaOH solution and extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated to afford benzyl ((*R*)-pyrrolidin-3-yl)methylcarbamate (800 mg, 76%). LC/MS: *m/z* 335.3 $(M+H)^+$ at 1.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[001988] Benzyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[001989]



[001990] A solution of triethylamine (507 μL, 3.64 mmol) and benzyl ((*R*)-pyrrolidin-3-yl)methylcarbamate (0.47 g, 2 mmol) in CH_2Cl_2 was added dropwise to a solution of 2-(4-chloro-6-fluoroquinazolin-2-yl)phenol (0.5 g, 1.82 mmol) in CH_2Cl_2 (20 mL). The reaction was stirred at room temperature for 3 h. After quenching the reaction with water, the aqueous phase was extracted twice with CH_2Cl_2. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. Purification via silica gel chromatography using 5–10% EtOAc in CH_2Cl_2 yielded benzyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate. LC/MS: *m/z* 473.1 $(M+H)^+$ at 2.91 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001991] 2-(4-((*S*)-3-(Aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol




[001992]

[001993] Under an N_2 atmosphere, a mixture of benzyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (0.770 g, 1.6 mmol) and MeOH (5 mL) was added to Pd/C (77 mg, 10 % weight Pd on carbon) weighed into a 100 mL flask. After the atmosphere in the flask was evacuated and purged with N_2 three times, the reaction mixture was vigorously stirred under an H_2 atmosphere at ambient pressure overnight and then filtered through a pad of Celite, concentrated, and dried to obtain 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.45 g, 81%). LC/MS: *m/z* 459.5 $(M+H)^+$ at 2.81 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001994] (*R*)-Tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[001995]




[001996] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol) and (*R*)-tetrahydrofuran-3-yl chloroformate (13 mg, 0.09 mmol). The reaction was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (*R*)-tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the

TFA salt. LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[001997] Example 356


[001998] **(*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[001999]



[002000] **(*S*)-Tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002001]




[002002] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol) and (*S*)-tetrahydrofuran-3-yl chloroformate (13 mg, 0.09 mmol). The reaction was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (*S*)-tetrahydrofuran-3-yl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the

TFA salt. LC/MS: *m/z* 453.3 $(M+H)^+$ at 2.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002003] Example 357


[002004] ***N*-(((*S*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide**

[002005]



[002006] ***N*-(((*S*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide**

[002007]




[002008] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) and cyclopropanecarboxylic acid (10 mg, 0.12 mmol) in DMF (1.0 mL) was added triethylamine (25 μL, 0.18 mmol) followed by the addition of HATU (45 mg, 0.117 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-(((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methyl)cyclopropanecarboxamide as the TFA salt. LC/MS: *m/z* 407.3 $(M+H)^+$ at 2.26 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002009] Example 358


[002010] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide**

[002011]



[002012] ***N*-((*R*)-1-(6-Fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide**

[002013]




[002014] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added 2-(4-fluorophenyl)-2-hydroxyacetic acid (20 mg, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide as the TFA salt. LC/MS: *m/z* 477.3 $(M+H)^+$ at 2.80 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002015] Example 359


[002016] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide**

[002017]



[002018] ***N*-((*R*)-1-(2-(2-Fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide**

[002019]




[002020] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added 2-(4-fluorophenyl)-2-hydroxyacetic acid (0.020 g, 0.12 mmol), followed by the addition of triethylamine (25.6 μL, 0.184 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain *N*-((*R*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-2-(4-fluorophenyl)-2-hydroxyacetamide as the TFA salt. LC/MS: *m/z* 4491.3 $(M+H)^+$ at 2.46 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002021] Example 360


[002022] **2-(4-Fluorophenyl)-2-hydroxy-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide**

[002023]



[002024] **2-(4-Fluorophenyl)-2-hydroxy-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide**

[002025]




[002026] To a solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) was added 2-(4-fluorophenyl)-2-hydroxyacetic acid (0.02 g, 0.12 mmol), followed by the addition of triethylamine (25 μL, 0.18 mmol) and HATU (0.045 g, 0.12 mmol). The reaction was stirred at room temperature for 2 h, filtered, and purified using reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to obtain 2-(4-fluorophenyl)-2-hydroxy-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)acetamide as the TFA salt. LC/MS: *m/z* 473.1 $(M+H)^+$ at 2.32 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002027] Example 361


[002028] **(2*R*)-2-Hydroxy-*N*-((*R*)-1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-4,4-dimethylpentanamide**

[002029]



[002030] **(2*R*)-2-Hydroxy-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-4,4-dimethylpentanamide**

[002031]




[002032] To a stirred solution of 2-(4-((*R*)-3-aminopyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (50 mg, 0.16 mmol) in 1 mL of DMF cooled at 0 °C was added (*R*)-2-hydroxy-4,4-dimethylpentanoic acid (27.3 mg, 0.187 mmol), followed by the addition of triethylamine (32 mg, 44 μL, 0.31 mmol), then HATU (71.1 mg, 0.187 mmol). The reaction was stirred at 0 °C for 10 minutes, warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford (2*R*)-2-hydroxy-*N*-((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)-4,4-dimethylpentanamide as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.4 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002033] Example 362


[002034] **Benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002035]



[002036] **Benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002037]




[002038] A mixture of benzyl ((*R*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (0.20 g, 0.60 mmol) and 4 M HCl (10 mL) in dioxane was stirred at room temperature for 3 h. After evaporating the solvent under reduced pressure, the solid was triturated with Et_2O, dried under vacuum, and taken up in CH_2Cl_2 (10 mL). 2-(4-Chloro-7-methylquinazolin-2-yl)phenol (0.16 g, 0.60 mmol) was added to this solution, followed by the addition of triethylamine (0.25 mL, 1.8 mmol). The reaction mixture was stirred at room temperature overnight, then diluted with CH_2Cl_2 and washed with water. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in hexanes gave benzyl ((*R*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as a colorless oil (0.19 g, 68% yield). LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.58 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002039] Example 363


[002040] Benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002041]



[002042] Benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002043]




[002044] A mixture of benzyl ((*S*)-1-(*tert*-butoxycarbonyl)pyrrolidin-3-yl)methylcarbamate (0.20 g, 0.60 mmol) and 4 M HCl in dioxane (10 mL) was stirred for 3 h at room temperature. After evaporating the solvent under reduced pressure, the solid was triturated with Et_2O, dried under vacuum, and taken up in CH_2Cl_2 (10 mL). 2-(4-Chloro-7-methylquinazolin-2-yl)phenol (0.16 g, 0.60 mmol) was added to this solution, followed by the addition of triethylamine (0.25 mL, 1.8 mmol). The reaction mixture was stirred at room temperature overnight, then diluted with CH_2Cl_2 and washed with water. The combined organic layers were dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in hexanes gave benzyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as a colorless oil (0.19 g, 68% yield). LC/MS: *m/z* 469.1 $(M+H)^+$ at 2.58 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002045] Example 364


[002046] **Ethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002047]



[002048] **Ethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002049]




[002050] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (25 mg, 0.075 mmol) in CH_2Cl_2 (1.0 mL) was cooled in an ice bath. To this mixture was added ethyl chloroformate (7.8 μL, 0.082 mmol), followed by triethylamine (21 μL, 0.15 mmol). After removing the ice bath, the reaction was stirred for 3 h at room temperature. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave ethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 407.1 $(M+H)^+$ at 2.29 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002051] Example 365


[002052] **Isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002053]



[002054] **Isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002055]



[002056] Method A

[002057] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (25 mg, 0.075 mmol) in CH_2Cl_2 (1.0 mL) was cooled in an ice bath. To this mixture was added isobutyl chloroformate (11 μL, 0.082 mmol), followed by the addition of triethylamine (21 μL, 0.15 mmol). After removing the ice bath, the reaction was stirred for 3 h at room temperature. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 435.3 $(M+H)^+$ at 2.55 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[002058] Method B

[002059] A mixture of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (200 mg, 0.6 mmol), THF (6.0 mL), and CH_2Cl_2 was stirred under an N_2 atmosphere. Triethylamine (0.166 mL, 1.2 mmol) was added, and the reaction was cooled in an ice bath. To

this was added 1 M isobutyl chloroformate solution (78 μL in 600 μL THF, 0.6 mmol). After allowing the reaction mixture to warm to room temperature, CH_2Cl_2 was added, and the organic solution washed twice with water, then dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–20% EtOAc in CH_2Cl_2/hexanes (1:1) gave isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (183 mg, 70%). LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.63 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). LC/MS: *m/z* 435 $(M+H)^+$ at 2.63 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43 (dd, *J* = 8.2, 1.8 Hz, 1H), 8.22 (d, *J* = 8.7 Hz, 1H), 7.58 (s, 1H), 7.42–7.31 (m, 3H), 6.93–6.89 (m, 2H), 4.17–3.91 (m, 3H), 3.80–3.69 (m, 3H), 3.34–3.30 (m, 1H), 3.21–3.07 (m, 2H), 2.49 (s, 3H), 2.16–2.08 (m, 1H), 1.88–1.74 (m, 2H), 0.88 (d, *J* = 6.7 Hz, 6H).

[002060] Isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride

[002061]




[002062] A solution of isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (183 mg, 0.42 mmol) in CH_2Cl_2 (1.5 mL) was stirred under an N_2 atmosphere. A 1.0 M HCl solution in ether (0.42 ml, 0.42 mmol) was added dropwise to this solution. After 10 minutes, 5 mL ether was added, and a precipitate formed which was filtered and dried to obtain isobutyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride (169 mg, 85%). LC/MS: *m/z* 435.5 $(M+H)^+$ at 2.64 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.31 (d, *J* = 8.7 Hz, 1H), 8.24 (d, *J* = 6.8 Hz, 1H), 7.80 (s, 1H), 7.53–7.45 (m, 2H), 7.41–7.38 (m, 1H), 7.12 (d, *J* = 8.2 Hz, 1H), 7.05–7.02 (m, 1H), 4.36–3.85 (m, 5H), 3.77–3.70

(m, 2H), 3.22–3.06 (m, 2H), 2.54 (s, 3H), 2.18–2.13 (m, 1H), 1.87–1.77 (m, 2H), 0.87 (d, J = 6.7 Hz, 6H).

[002063] Example 366

[002064] **2-Methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002065]



[002066] **2-Methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002067] Method A

[002068] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (25 mg, 0.075 mmol) in anhydrous CH_2Cl_2 was cooled in an ice bath. Then 2-methoxyethyl chloroformate (8.6 μL, 0.075 mmol) was added, followed by the addition of triethylamine (16 μL, 0.11 mmol). After removing the ice bath, the reaction was stirred at room temperature for 4 h. Water and saturated aqueous $NaHCO_3$ solution were added, and the reaction mixture was stirred at room temperature overnight. After separation of the aqueous and organic layers, the organic phase was dried over Na_2SO_4, filtered, and concentrated. Purification via preparative reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)), *m/z*: M+1 *obs* = 437.3; t_R = 2.18 minutes


[002069] Method B

[002070] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (0.12 g, 0.35 mmol) and triethylamine (98 μL, 0.7 mmol) in anhydrous DMF (4.0 mL) was cooled in an ice bath. 2-Methoxyethyl chloroformate (40 mL, 0.35 mmol) was added, and

the ice bath was removed. The solution was stirred at room temperature overnight. Purification via silica gel chromatography using 0–10% MeOH in CH_2Cl_2 gave 2-methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (0.12 g, 80%). LC/MS: *m/z* 437.5 $(M+H)^+$ at 2.20 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002071] 2-Methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride

[002072]




[002073] A 1.0 M HCl solution in Et_2O (0.27 mL, 0.27 mmol) was slowly added to a stirring solution of 2-methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (0.12 g, 0.27 mmol) in CH_2Cl_2 (5 mL). The reaction was stirred for 30 minutes at room temperature, then Et_2O was slowly added to the solution until a precipitate formed. After stirring for 1 h, the solid was filtered and washed with Et_2O to obtain 2-methoxyethyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride (0.10 g, 77%). LC/MS: *m/z* 437.1 $(M+H)^+$ at 2.19 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO-d6) δ 8.31 (d, J = 8.7 Hz, 1H), 8.24 (d, J = 7.3 Hz, 1H), 7.77 (s, 1H), 7.52–7.48 (m, 3H), 7.10 (d, J = 8.1 Hz, 1H), 7.05–7.02 (m, 1H), 4.22–4.12 (m, 2H), 4.08–4.05 (m, 3H), 3.87–3.83 (m, 1H), 2H), 3.24 (s, 3H), 3.23–3.12 (m, 2H), 2.54 (s, 3H), 2.55–2.52 (m, 1H), 2.17–2.13 (m, 1H), 1.84–1.77 (m, 1H).

[002074] Example 367


[002075] Propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002076]



[002077] **Propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002078]



[002079] Method A

[002080] Propyl chloroformate (12 mg, 0.10 mmol) and triethylamine (30 mg, 42 μL, 0.30 mmol) were added to 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (33 mg, 0.10 mmol) in 1 mL anhydrous DMF. The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 421.1 $(M+H)^+$ at 2.43 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[002081] Method B

[002082] A solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (301 mg, 0.9 mmol) in CH_2Cl_2 (10 ml) was stirred under an N_2 atmosphere. Triethylamine (0.25 mL, 1.8 mmol) was added, and the solution was cooled to –30 °C. A 1 M

propyl chloroformate solution (0.1 mL in 0.9 mL CH_2Cl_2, 0.9 mmol) was added, and the reaction mixture was allowed to warm to room temperature over a period of 30 minutes. After adding CH_2Cl_2 to the reaction mixture, it was washed 2 times with water, before it was dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–20% EtOAc in CH_2Cl_2 gave propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (236 mg, 62%). LC/MS: *m/z* 421 $(M+H)^+$ at 2.54 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.43 (dd, *J* = 8.2, 1.8 Hz, 1H), 8.21 (d, *J* = 8.7 Hz, 1H), 7.58 (s, 1H), 7.39–7.31 (m, 3H), 6.93–6.89 (m, 2H), 4.09–3.86 (m, 6H), 3.78–3.73 (m, 1H), 3.21–3.09 (m, 2H), 2.49 (s, 3H), 2.15–2.09 (m, 1H), 1.83–1.74 (m, 1H), 1.62–1.48 (m, 2H), 0.89–0.86 (m, 3H).

[002083] Propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride

[002084]




[002085] A solution of propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (232 mg, 0.552 mmol) in CH_2Cl_2 (2 mL) was stirred under an N_2 atmosphere. A 2.0 M HCl solution in ether (0.276 mL, 0.552 mmol) was added dropwise to this solution. After 10 minutes, ether (8 mL) was added until a precipitate formed, which was filtered and dried to obtain propyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride (223 mg, 88%). LC/MS: *m/z* 421 $(M+H)^+$ at 2.54 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.31 (d, *J* = 8.7 Hz, 1H), 8.24 (dd, *J* = 7.9, 1.4 Hz, 1H), 7.82 (s, 1H), 7.54–7.49 (m, 2H), 7.41–7.37 (m, 1H), 7.14 (d, *J* = 8.1 Hz, 1H), 7.06–7.02 (m, 1H), 4.29–3.78 (m, 7H), 3.24–3.08

(m, 2H), 2.54 (s, 3H), 2.18–2.12 (m, 1H), 1.84–1.79 (m, 1H), 1.59–1.50 (m, 2H), 0.87 (t, J = 7.4 Hz, 3H).

[002086] Example 368


[002087] **Isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002088]



[002089] **Isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002090]




[002091] Isopropyl chloroformate (12 mg, 0.10 mmol) and triethylamine (30 mg, 42 μL, 0.30 mmol) were added to a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (33 mg, 0.10 mmol) in 1 mL anhydrous DMF. The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give isopropyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-

methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 421.1 $(M+H)^+$ at 2.42 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002092] Example 369


[002093] **Neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002094]



[002095] **Neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002096]



[002097] Method A

[002098] Neopentyl chloroformate (15 mg, 0.10 mmol) and triethylamine (30 mg, 42 μL, 0.30 mmol) were added to a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (33 mg, 0.10 mmol) in 1 mL anhydrous DMF. The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-


methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 449.3 $(M+H)^+$ at 2.67 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002099] Method B

[002100] A mixture of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)phenol (200 mg, 0.6 mmol), THF (6.0 mL), and DMF (1 mL) was stirred under an N_2 atmosphere. Triethylamine (0.166 mL, 1.2 mmol) was added, and the reaction was cooled in an ice bath. To this was added 1 M neopentyl chloroformate solution (89 µL in 600 µL THF, 0.6 mmol). After allowing the reaction mixture to warm to room temperature over a period of 30 minutes, CH_2Cl_2 was added to the reaction mixture, and it was washed once with water before it was dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–20% EtOAc in CH_2Cl_2/hexanes (1:1) gave neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (222 mg, 94%). LC/MS: *m/z* 449 $(M+H)^+$ at 2.73 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, acetic acid–d4) δ 8.42 (dd, *J* = 8.1, 1.8 Hz, 1H), 8.20 (d, *J* = 8.7 Hz, 1H), 7.57 (s, 1H), 7.40–7.29 (m, 3H), 6.92–6.89 (m, 2H), 4.11–3.92 (m, 3H), 3.78–3.61 (m, 3H), 3.33–3.28 (m, 1H), 3.23–3.10 (m, 2H), 2.49 (s, 3H), 2.16–2.08 (m, 1H), 1.83–1.75 (m, 1H), 0.89 (s, 9H).

[002101] **Neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride**

[002102]




[002103] A solution of neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate (215 mg, 0.48 mmol) in CH_2Cl_2 (2 ml) was stirred under an N_2 atmosphere. A 1.0 M HCl solution in ether (0.48 ml, 0.48 mmol) was added dropwise to this solution. After 10 minutes, ether (8 ml) was added until a precipitate formed, which was filtered

and dried to obtain neopentyl ((*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate hydrochloride (214 mg, 92%). LC/MS: *m/z* 449 $(M+H)^+$ at 2.75 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)). ^1H NMR (400 MHz, DMSO–d6) δ 8.31 (d, *J* = 8.7 Hz, 1H)1H), 7.85 (s, 1H), 7.54–7.49 (m, 2H), 7.47–7.45 (m, 1H), 7.16 (d, *J* = 8.2 Hz, 1H), 7.05–7.02 (m, 1H), 4.33–3.74 (m, 5H), 3.70–3.64 (m, 2H), 3.19–3.16 (m, 2H), 2.53 (s, 3H), 2.18–2.13 (m, 1H), 1.87–1.81 (m, 1H), 0.88 (s, 9H).

[002104] Example 370


[002105] ***tert*-Butyl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[002106]



[002107] ***tert*-Butyl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate**

[002108]




[002109] *tert*-Butyl (*S*)-pyrrolidin-3-ylcarbamate (245 mg, 1.31 mmol) was dissolved in 1.8 mL of anhydrous CH_2Cl_2 and cooled to 0 °C. 2-(4-Chloro-7-methylquinazolin-2-yl)phenol (300 mg, 1.1 mmol) dissolved in 1.8 mL of anhydrous CH_2Cl_2 was added dropwise to the mixture followed by triethylamine (134 mg, 184 μL, 1.32 mmol). The reaction was allowed to warm to room temperature and was stirred overnight. Purification via silica gel chromatography using

10–100% ethyl acetate/hexanes gave *tert*-butyl (*S*)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-ylcarbamate. LC/MS: *m/z* 421.0 $(M+H)^+$ at 2.84 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002110] Example 371


[002111] **Isopropyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002112]



[002113] **2-(4-((*S*)-3-(Aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol**

[002114]




[002115] A mixture of ((*S*)-pyrrolidin-3-yl)methanamine (0.6 g, 6.0 mmol), Na_2CO_3 (2.2 g, 21 mmol) and methyl isobutyl ketone (12 mL, 6.0 mmol) under an N_2 atmosphere was refluxed overnight using a Dean Stark apparatus. After allowing the reaction to cool to room temperature,

2-(4-chloro-7-methylquinazolin-2-yl)-3-fluorophenol (1.73 g, 5.99 mmol) was added, and the mixture was stirred overnight at room temperature under an N_2 atmosphere. The reaction was then quenched with water. The aqueous layer was extracted twice with CH_2Cl_2. The organic extracts were combined, dried over Na_2SO_4, filtered, and concentrated to obtain a crude material containing 2-(4-((*S*)-3-((*E*)-(4-methylpentan-2-ylideneamino)methyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol which was then hydrolyzed by heating at 50 °C in 80 mL of a water/isopropanol mixture (1:1) for 6 hours. The reaction was cooled to room temperature, and the aqueous layer was extracted twice with CH_2Cl_2, dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 5–20% MeOH and CH_2Cl_2 afforded 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (400 mg, 19% after 3 steps). LC/MS: *m/z* 353.1 $(M+H)^+$ at 1.22 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002116] Isopropyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002117]




[002118] Isopropyl chloroformate (10 mg, 0.09 mmol) was added to a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) and triethylamine (25 mL, 0.18 mmol) in DMF (0.6 mL) at –78 °C (external temperature). The reaction mixture was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford isopropyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002119] Example 372

[002120] **Isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002121]




[002122] **Isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002123] Isobutyl chloroformate (12 mg, 0.09 mmol) was added to a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) and triethylamine (25 mL, 0.18 mmol) in DMF (0.6 mL) at –70 °C (external temperature). The reaction mixture was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford isobutyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 453.5 $(M+H)^+$ at 2.46 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002124] Example 373


[002125] **2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002126]



[002127] **2-Methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002128]




[002129] 2-Methoxyethyl chloroformate (12 mg, 0.09 mmol) was added to a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-7-methylquinazolin-2-yl)-3-fluorophenol (0.03 g, 0.09 mmol) and triethylamine (25 mL, 0.18 mmol) in DMF (0.6 mL) at –78 °C. The reaction mixture was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford 2-methoxyethyl ((*S*)-1-(2-(2-fluoro-6-hydroxyphenyl)-7-methylquinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 455.5 $(M+H)^+$ at 2.11 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002130] Example 374


[002131] **Propyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002132]



[002133] **Propyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002134]




[002135] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol), followed by the addition of propyl chloroformate (11 mg, 0.09 mmol). The reaction was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford propyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 425.3 $(M+H)^+$ at 2.31 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002136] Example 375


[002137] **Neopentyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002138]



[002139] **Neopentyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002140]




[002141] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol), followed by the addition of neopentyl chloroformate (13 mg, 0.09 mmol). The reaction was warmed to room temperature, filtered, and purified via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to afford neopentyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 453.5 $(M+H)^+$ at 2.88 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002142] Example 376


[002143] **Isobutyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002144]



[002145] **Isobutyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate**

[002146]




[002147] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin–1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol), followed by the addition of isobutyl chloroformate (12 mg, 0.09 mmol). The reaction was warmed to room temperature, and filtered and purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave isobutyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 439.5 $(M+H)^+$ at 2.76 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002148] Example 377


[002149] 2-Methoxyethyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002150]



[002151] 2-Methoxyethyl ((*S*)-1-(6-fluoro-2-(2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate

[002152]




[002153] To a solution of 2-(4-((*S*)-3-(aminomethyl)pyrrolidin-1-yl)-6-fluoroquinazolin-2-yl)phenol (0.03 g, 0.09 mmol) in DMF (1.0 mL) at –60 °C (external temperature) was added triethylamine (25 mL, 0.18 mmol), followed by the addition of 2-methoxyethyl chloroformate (12 mg, 0.09 mmol). The reaction was warmed to room temperature, and filtered and purification via reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) gave 2-methoxyethyl ((*S*)-1-(6-fluoro-2- (2-hydroxyphenyl)quinazolin-4-yl)pyrrolidin-3-yl)methylcarbamate as the TFA salt. LC/MS: *m/z* 441.5 $(M+H)^+$ at 2.30 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002154] Example 401


[002155] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-4-yl)methyl)azetidine-3-carboxamide**

[002156]



[002157] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid**


[002158]



[002159] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (2.84 g, 10.5 mmol) and azetidine-3-carboxylic acid (1.06 g, 10.5 mmol) in DMF (100 mL) was added triethylamine (3.18 g, 4.39 mL, 31.5 mmol). The reaction was stirred at room temperature overnight then diluted with H_2O (400 mL). The pH of the solution was adjusted to 3–4 by addition of an aqueous 1 M HCl solution. The white precipitate then obtained was filtered, washed with H_2O, and purified by silica gel chromatography using 0–15% MeOH–CH_2Cl_2 giving 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid. LC/MS: *m/z* 336.3 $(M+H)^+$ at 1.97 min (10%—99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002160] 1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-4-yl)methyl)azetidine-3-carboxamide

[002161]




[002162] To a solution of 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid (34 mg, 0.1 mmol) and HATU (57 mg, 0.15 mmol) in DMF (1 mL) was added (pyridin-4-yl)methanamine (32 mg, 30μL, 0.30 mmol). The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) giving 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-

((pyridin-4-yl)methyl)azetidine-3-carboxamide as the TFA salt. LC/MS: *m/z* 426.3 $(M+H)^+$ at 1.72 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002163] Example 402


[002164] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)azetidine-3-carboxamide**

[002165]



[002166] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid**


[002167]



[002168] To a solution of 2-(4-chloro-7-methylquinazolin-2-yl)phenol (2.84 g, 10.5 mmol) and azetidine-3-carboxylic acid (1.06 g, 10.5 mmol) in DMF (100 mL) was added triethylamine (3.18 g, 4.39 mL, 31.5 mmol). The reaction was stirred at room temperature overnight then diluted with H_2O (400 mL). The pH of the solution was adjusted to 3–4 by addition of an aqueous 1 M HCl solution. The white precipitate then obtained was filtered, washed with H_2O, and purified by silica gel chromatography using 0–15% MeOH–CH_2Cl_2 giving 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid. LC/MS: *m/z* 336.3 $(M+H)^+$ at 1.97 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002169] **1-(2-(2-Hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)azetidine-3-carboxamide**

[002170]




[002171] To a solution of 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid (34 mg, 0.1 mmol) and HATU (57 mg, 0.15 mmol) in DMF (1 mL) was added (pyridin-3-yl)methanamine (32 mg, 30μL, 0.30 mmol). The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)-*N*-((pyridin-3-yl)methyl)azetidine-3-carboxamide as the TFA salt. LC/MS: *m/z* 426.3 $(M+H)^+$ at 1.72 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002172] Example 403


[002173] ***N*-(3-(Trifluoromethoxy)benzyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxamide**

[002174]



[002175] ***N*-(3-(Trifluoromethoxy)benzyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxamide**

[002176]




[002177] To a solution of 1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxylic acid (34 mg, 0.10 mmol) and HATU (57 mg, 0.15 mmol) in DMF (1 mL) was added (3-(trifluoromethoxy)phenyl)methanamine (57 μL, 0.30 mmol). The reaction was stirred at room temperature overnight, filtered, and purified by reverse phase HPLC (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give *N*-(3-(trifluoromethoxy)benzyl)-1-(2-(2-hydroxyphenyl)-7-methylquinazolin-4-yl)azetidine-3-carboxamide as the TFA salt. LC/MS: *m/z* 509.3 $(M+H)^+$ at 2.73 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).


[002178] Example 501

[002179] **1-(4-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**

[002180]




[002181] ***tert*-Butyl 4-(2-chloro-7-methylquinazolin-4-yl)piperazine-1-carboxylate**

[002182] 


[002183] A solution of 2,4-dichloro-7-methylquinazoline (0.43 g, 2.0 mmol), *tert*-butyl piperazine-1-carboxylate (0.37 g, 2.0 mmol), and triethylamine (0.56 mL, 4.0 mmol) in CH_2Cl_2 (20 mL) was stirred at room temperature overnight. The solution was washed with water, dried over Na_2SO_4, and evaporated to give *tert*-butyl 4-(2-chloro-7-methylquinazolin-4-yl)piperazine-1-carboxylate (0.72 g, 99%). LC/MS: *m/z* 363.1 $(M+H)^+$ at 3.21 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002184] **2-chloro-7-methyl-4-(piperazin-1-yl)quinazoline dihydrochloride**

[002185] 

[002186] *tert*-Butyl 4-(2-chloro-7-methylquinazolin-4-yl)piperazine-1-carboxylate (0.72 g, 2.0 mmol) in 4 M HCl in dioxane (10 mL) was stirred at room temperature for five hours. The solvent was evaporated to give 2-chloro-7-methyl-4-(piperazin-1-yl)quinazoline dihydrochloride (0.67 g, 100%). LC/MS: *m/z* 263.3 $(M+H)^+$ at 1.75 min (10%—99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002187] **1-(4-(2-Chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one**




[002188]

[002189] A solution of 2-chloro-7-methyl-4-(piperazin-1-yl)quinazoline dihydrochloride (127 mg, 0.38 mmol), 2-hydroxy-4-methylpentanoic acid (55 mg, 0.42 mmol), HATU (158 mg, 0.42 mmol), and triethylamine (163 μL, 1.2 mmol) in DMF (4 mL) was stirred at room temperature for four hours. The reaction was diluted with water and was extracted with EtOAc. The combined extracts were washed with saturated aqueous NaCl and water, dried over Na_2SO_4, and evaporated. Purification via silica gel chromatography using 0–50% EtOAc in hexanes gave 1-(4-(2-chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (23 mg, 16%). LC/MS: *m/z* 377.3 $(M+H)^+$ at 2.71 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002190] **1-Bromo-2-(difluoromethyl)benzene**


[002191]



[002192] A mixture of 2-bromo benzaldehyde (27.1 g, 0.15 mol) and (diethylamino)sulfur trifluoride (52 g, 0.32 mol) in anhydrous benzene (250 mL) was heated to reflux overnight under an N_2 atmosphere. The mixture was treated with $NaHCO_3$ (sat. aq.) and CH_2Cl_2, and the organic phase was separated. The aqueous phase was extracted with CH_2Cl_2, and the combined organic layers were washed with water and saturated aq. NaCl. Removal of the solvent gave 1-bromo-2-difluoromethylbenzene (16 g, 52%). ^1H NMR($CDCl_3$): δ 7.68 (d, 1H, J = 7.6), 7.62 (d, 1 H, J = 7.6), 7.43 (t, 1 H, J = 7.6), 7.34 (t, 1 H, J = 7.6), 7.06–6.79 (t, 1 H, J = 55.2).

[002193] **2-(Difluoromethyl)phenylboronic acid**


[002194]



[002195] To a solution of 1-bromo-2-difluoromethylbenzene (16 g, 0.077 mol) in THF (300 mL) was added *n*-BuLi (2.5 M, 46 mL, 0.12 mol) dropwise at –78 °C. After one hour, trimethyl borate (16 g, 0.15 mol) was added dropwise. The mixture was warmed to room temperature slowly overnight, and then the mixture was treated with water and ethyl acetate. The organic layer was extracted with a 10% aqueous NaOH solution. The aqueous solution was acidified with dilute aq. HCl and extracted with ethyl acetate. Removal of the solvent gave 2-(difluoromethyl)phenylboronic (0.7 g, 5.3%). ^{1}H NMR($CDCl_3$): δ 8.28 (d, 1 H, J = 7.2), 7.86 (d, 1H, J = 7.2), 7.58–7.79 (m, 1 H), 7.73 (t, 1 H, J = 7.2), 7.65 (t, 1 H).

[002196] 1-(4-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one

[002197]




[002198] A mixture of 1-(4-(2-chloro-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one (23 mg, 61 μmol), 2-(difluoromethyl)phenylboronic acid (13 mg, 73 μmol), $Pd(dppf)Cl_2$ (3 mg, 3.7 μmol), and K_2CO_3 (17 mg, 122 μmol) in DMF (1 mL) and H_2O (0.5 mL) was heated in a sealed microwave vial at 170 °C for 6 minutes. After filtration and evaporation of the solvents, the mixture was purified using preparative reverse phase HPLC (10–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)) to give 1-(4-(2-(2-(difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperazin-1-yl)-2-hydroxy-4-methylpentan-1-one as the TFA salt. LC/MS: *m/z* 469.3 $(M+H)^+$ at 2.44 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002199] Example 502


[002200] (2*R*)-*N*-(1-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide

[00220T]



[002202] *tert*-Butyl 1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate


[002203]



[002204] To a solution of 2,4-dichloro-7-methylquinazoline (1.06 g, 5.0 mmol) in CH_2Cl_2 (20 mL) cooled in an ice bath under an N_2 atmosphere was added *tert*-butyl piperidin-4-ylcarbamate (1.0 g, 5.0 mmol) dissolved in CH_2Cl_2 (30 mL). To this was added triethylamine (1.4 mL, 10.0 mmol), and the reaction was allowed to warm to room temperature. After stirring for 1 h, the solution was washed with water and dried over Na_2SO_4. The solvents were evaporated to obtain *tert*-butyl 1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate as a yellow solid (1.6 g, 85%). LC/MS: *m/z* 377.5 $(M+H)^+$ at 2.96 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002205] **1-(2-Chloro-7-methylquinazolin-4-yl)piperidin-4-amine dihydrochloride**


[002206]



[002207] Under an N_2 atmosphere, a 4 M HCl solution in dioxane (50 mL) was added to *tert*-butyl 1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-ylcarbamate (1.06 g, 2.82 mmol), and the mixture was stirred for 3 h. After evaporating the solvents, the resulting solid was taken up in CH_2Cl_2 and EtOAc and concentrated to yield 1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-amine dihydrochloride as a pale pink powder (985 mg, 100%). LC/MS: *m/z* 277.3 $(M+H)^+$ at 1.57 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002208] (*R*)-*N*-(1-(2-Chloro-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide


[002209]



[002210] To a mixture of (*R*)-tetrahydrofuran-2-carboxylic acid (0.13 mL, 1.58 mmol) in CH_2Cl_2 (2 mL) was added HATU (0.5 g, 1.58 mmol). After adding triethylamine (0.74 mL, 5.28 mmol), the solution was stirred for 30 min. 1-(2-Chloro-7-methylquinazolin-4-yl)piperidin-4-amine dihydrochloride (460 mg, 1.32 mmol) was added and the reaction was stirred overnight. The yellow, cloudy solution was washed 4 times with H_2O, and the aqueous layer was re-extracted with CH_2Cl_2. After concentrating the combined organic phases, purification via silica gel chromatography using 0–100% EtOAc in hexanes provided (*R*)-*N*-(1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide (0.17 g, 33%). LC/MS: *m/z* 475.3 $(M+H)^+$ at 2.20 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002211] 1-Bromo-2-difluoromethyl-benzene

[002212]




[002213] A solution of 2-bromobenzaldehyde (55.5 g, 300 mmol) and (diethylamino)sulfur trifluoride (75.0 g, 467 mmol) in 250 ml dichloromethane was refluxed under a nitrogen

atmosphere overnight. The cooled solution was poured into 0.5 L aqueous 15% $NaHCO_3$ and stirred until no more CO_2 was produced. The layers were separated, and the aqueous layer was extracted with 250 ml dichloromethane. The combined organic layers were washed with 250 ml 5% aq. $NaHCO_3$ and saturated aq. NaCl, dried over Na_2SO_4, filtered, and evaporated to dryness under reduced pressure. The crude material was purified by vacuum distillation, and the fraction boiling at 62–63 °C at 12 mbar was collected, yielding 1-bromo-2-difluoromethyl-benzene (42.6 g, 69%) as a light-yellow oil.


[002214] 2-(2-Difluoromethyl-phenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane

[002215]



[002216] To a solution of 1-bromo-2-difluoromethyl-benzene (19.8 g, 95.7 mmol) in dry THF (200 ml) at –78 °C under a nitrogen atmosphere was added 2.5 M *n*-BuLi in hexanes (42 ml, 105 mmol) slowly. After completion of the addition, the resulting dark solution was stirred for an additional hour at –78 °C. Subsequently, 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (25 ml, 123 mmol) was added, and the solution was slowly warmed to room temperature. After stirring overnight at room temperature under a nitrogen atmosphere the solution was poured into 400 ml water. Ethyl acetate (300 ml) was added and the layers were separated. The aqueous layer was extracted twice with ethyl acetate (150 ml and 50 ml respectively), and the combined organic layers were washed with water, dried over Na_2SO_4, filtered, and evaporated to dryness under reduced pressure. The resulting brown oil (21 g) was purified by bulb-to-bulb-distillation at $3x10^{-3}$ mbar at 90–95 °C to yield 2-(2-difluoromethyl-phenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane (14.4 g, 59%) as a slightly yellow oil.

[002217] (2*R*)-*N*-(1-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide




[002218]

[002219] To a mixture of (*R*)-*N*-(1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide (65.2 mg, 0.17 mmol), 2-(2-(difluoromethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.09 mL, 0.35 mmol), $Pd(dppf)Cl_2$ (17.1 mg, 0.02 mmol), and K_2CO_3 (48.1 mg, 0.35 mmol) was added DMF (3.3 mL), and the reaction was heated in a microwave synthesizer at 170 °C for 6 min. Since LC/MS showed that reaction was not complete, another 2 equivalents of 2-(2-(difluoromethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.09 mL, 0.38 mmol) were added, and the reaction was heated to 170 °C in a microwave synthesizer for 6 minutes The solution was diluted with EtOAc and washed with H_2O. The aqueous fraction was re-extracted with EtOAc and the organic layers were combined, dried over Na_2SO_4, and concentrated. Purification via silica gel chromatography using 0–40% EtOAc in CH_2Cl_2/hexanes (1:1) gave (2*R*)-*N*-(1-(2-(2-(difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)-tetrahydrofuran-2-carboxamide (38 mg, 47%). LC/MS: *m/z* 467.3 $(M+H)^+$ at 2.20 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002220] Example 503


[002221] **(2*S*)-*N*-((1-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide**



[002222]


[002223] ***tert*-Butyl 4-(aminomethyl)piperidine-1-carboxylate**

[002224]



[002225] A solution of 4-(aminomethyl)-piperidine (25.3 g, 222 mmol) and benzaldehyde (23.5 g, 222 mmol) in 250 mL toluene was refluxed overnight under a nitrogen atmosphere in a flask equipped with a Dean-Stark trap. The solution was cooled to 0–5 °C in an ice bath, and Boc_2O (53.2 g, 244 mmol) was added in one portion. The ice bath was removed, and stirring at room temperature was continued for 6 hours. The mixture was evaporated to dryness under reduced pressure, and the residue was stirred overnight with potassium bisulphate in water (33.2 g $KHSO_4$ in 244 mL water). The solution was extracted twice with 200 ml portions of diethyl ether and was made alkaline with 50 ml 30% aqueous sodium hydroxide solution. After saturation of the solution with sodium chloride, it was extracted with dichloromethane (1x 150 ml, 2 x 50 mL). The combined organic layers were dried over Na_2SO_4, filtered, and evaporated to dryness, yielding *tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate (41.8 g, 88%) as an orange oil.

[002226] 4-[((*S*)-2-Hydroxy-3-methyl-butyrylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester




[002227]

[002228] HOBt (23.5 g, 174 mmol) and EDCI·HCl (33.5 g, 174 mmol) were added to a mixture of *tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate (37.3 g, 174 mmol), (*S*)-(+)-2-hydroxy-3-methyl-butanoic acid (20.6 g, 174 mmol), and triethylamine (50 mL, 349 mmol) in a mixture of dichloromethane (500 mL) and DMF (250 mL) cooled in an ice bath. The reaction was stirred overnight at room temperature. An additional portion of EDCI·HCl (5 g) and triethylamine (4 mL) were added, and stirring was continued for another 4 hours. The mixture was diluted to 1 L by the addition of dichloromethane and was washed with 500 mL portions of water, 1 N HCl solution, 5% aq. $NaHCO_3$ solution, water, and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4, filtered, and evaporated to dryness under reduced pressure yielding 52 g of the crude product. This crude material was stirred with 200 mL diisopropyl ether, and the solid was collected by filtration and washed twice with 100 mL portions of diisopropyl ether to yield to 33 g (64%) of 4-[((*S*)-2-hydroxy-3-methyl-butyrylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester.

[002229] (*S*)-2-Hydroxy-3-methyl-*N*-piperidin-4-ylmethyl-butyramide


[002230]



[002231] Acetyl chloride (30 mL) was added dropwise to 500 mL methanol cooled in an ice bath, and after completion of the addition the solution was stirred at room temperature for 30 minutes. 4-[((*S*)-2-hydroxy-3-methyl-butyrylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (33 g) in 150 mL methanol was added, and stirring was continued for 2 hours at room temperature. The solution was evaporated to dryness to yield to 26.6 g of (*S*)-2-hydroxy-3-methyl-*N*-piperidin-4-ylmethyl-butyramide hydrochloride as a white solid.

[002232] (*S*)-*N*-((1-(2-Chloro-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide


[002233]



[002234] A solution of 2,4-dichloro-7-methylquinazoline (1.07 g, 5.0 mmol) in CH_2Cl_2 (20 mL) was cooled in an ice bath. (*S*)-2-Hydroxy-3-methyl-N-((piperidin-4-yl)methyl)butanamide hydrochloride (1.25 g, 5.0 mmol) suspended in 30 mL of CH_2Cl_2 was added in portions, followed by the addition of triethylamine (2.79 mL, 20.0 mmol). The reaction was allowed to warm to room temperature and was stirred for 1.5 h. After washing the mixture 4 times with H_2O, the aqueous portion was re-extracted with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4, filtered, concentrated, and dried under vacuum to obtain (*S*)-*N*-((1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide as a yellow solid (1.83 g, 94%). LC/MS: *m/z* 391.3 $(M+H)^+$ at 2.32 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002235] (2*S*)-*N*-((1-(2-(2-(Difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide

[002236]



[002237] To a solution of (*S*)-*N*-((1-(2-chloro-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide (200 mg, 0.51 mmol), 2-(2-(difluoromethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (195 μL, 0.77 mmol), and K_2CO_3 (142 mg, 1.02 mmol) in a DMF:H_2O mixture (4:1) was added Pd(dppf)Cl_2 (25.3 mg, 0.03 mmol). The reaction mixture was heated in a microwave synthesizer at 170 °C for 6 minutes The reaction was diluted with EtOAc and washed with 30 mL saturated aqueous NaCl solution. The organic portion was 3 times washed with H_2O, and the combined aqueous layers were re-extracted 3 times with EtOAc. All combined organic portions were then dried over Na_2SO_4, filtered, and concentrated. Purification via silica gel chromatography using 0–70% EtOAc in CH_2Cl_2:hexanes (1:1) yielded 177 mg (71%) of (2*S*)-*N*-((1-(2-(2-(difluoromethyl)phenyl)-7-methylquinazolin-4-yl)piperidin-4-yl)methyl)-2-hydroxy-3-methylbutanamide which crystallized as a white solid after freezing it for 1 h. LC/MS: *m/z* 483.3 $(M+H)^+$ at 2.24 min (10%–99% CH_3CN (0.035% TFA)/H_2O (0.05% TFA)).

[002238] **Table 3** below recites analytical data for exemplary compounds of the present invention.

Table 3

Cmpd No.	LC-MS M+1	LC-RT min
101	449.3	2.79
102	421	2.48
103	499.3	2.97
104	464.5	2.03
105	447.1	2.32
106	465.5	2.47
107	442.5	1.97
108	407.5	2.21
109	393.1	2.04
110	461.1	2.56
111	453.5	3.21
112	461.3	2.49
113	421.1	2.76
114	474.3	1.19
115	421.3	2.51
116	442.5	1.96
117	439.5	2.79
118	393.3	2.22
119	465.2	2.5
120	435.3	2.65
121	393.3	2.21
122	435.3	2.56
123	465.3	2.5
124	407.5	2.29
125	453.4	2.79
126	471.2	2.93
127	445.5	3.32
128	451.5	2.62
129	407.5	2.31
130	469.3	2.87
131	469.4	2.88
132	456.5	2.04
133	454.3	1.94
134	421.1	2.46
135	467.1	2.59
136	453.3	3.02
137	407.3	2.28
138	453.3	2.43
139	439.5	2.95
140	485.5	2.94
141	459.3	2.13
142	446.3	3.17
143	439.5	2.99
144	435.3	2.9
145	437.3	2.04
146	455.5	2.45
147	453.3	2.4
148	439.5	2.8
149	465.5	2.23
150	449.3	2.22
151	435.5	2.13
152	456.5	2.02
153	453.4	2.73
154	436.3	1.94
155	477.5	2.96
156	474.3	1.19
157	485.4	3.02

Cmpd No.	LC-MS M+1	LC-RT min
158	403.5	2.34
159	421.3	2.6
160	467.3	3.13
162	467.3	2.33
163	453.3	2.25
164	451.3	2.18
165	421.3	2.4
166	434.53	2.61
167	491.3	2.46
168	435.3	2.91
169	577.4	2.5
170	473.1	2.63
171	517.5	3.49
172	465	2.77
173	393.1	2.03
174	407.5	2.41
175	460.5	2.33
176	435.5	2.62
177	447.1	2.53
178	447.3	2.5
179	427.2	2.59
180	435.2	2.88
181	435.3	2.69
182	435.3	2.5
183	435.3	2.5
184	449.3	2.5
185	449.3	2.56
201	454.5	1.87
203	449.3	2.34
204	449.3	2.33
205	433.3	2.33
206	433.5	2.34
207	449.5	2.34
208	449.5	2.33
209	447.5	2.27
210	463.5	2.34
211	463.5	2.34
212	417	2.3
213	453.5	1.98
214	463.5	2.32
215	463.5	2.36
216	463.5	2.35
217	463.5	2.35
218	447.5	2.02

Cmpd No.	LC-MS M+1	LC-RT min
219	477.4	2.84
220	447.5	2.19
221	461.5	2.22
222	531.3	3.08
223	468.3	1.86
224	437.1	2.54
225	468.6	2.19
226	417.5	2.3
227	470.5	1.98
228	470.5	1.98
229	513.3	2.82
230	447.3	2.21
231	481.1	2.17
232	463.3	2.58
233	421.2	2.17
234	449.2	2.42
235	407.5	2.04
236	407.3	2.08
237	407.5	2.09
238	435.3	2.4
239	461.1	2.4
240	435.5	2.29
241	421.1	2.24
242	421.3	2.18
243	421.3	2.18
244	421.3	2.18
245	449.3	2.45
246	475.1	2.46
247	435.3	2.31
248	435.3	2.24
249	481.3	2.42
250	449.5	2.77
251	421	2.48
252	449.3	2.8
253	449.5	2.78
254	421.1	2.5
255	435.5	2.61
256	435.5	2.61
257	451.1	2.34
258	441.5	2.6
259	435.2	3.03
260	467.1	2.56
261	469.1	2.2
262	439.3	2.31

Cmpd No.	LC-MS M+1	LC-RT min
264	435.1	2.6
265	449.3	2.72
266	449.2	2.55
267	449.3	2.58
268	449.1	2.54
269	449.5	2.57
270	435.2	2.88
271	447.2	2.46
272	449.2	2.57
273	449.2	2.57
274	463.4	2.42
275	463.4	2.43
276	433.2	2.33
277	433.2	2.32
278	449.2	2.85
279	435.4	2.61
301	435	2.41
302	419	2.35
303	419	2.34
304	435	2.39
305	449.3	2.18
306	449.3	2.16
307	449.3	2.18
308	433.5	2.11
309	455.2	2.81
310	433.5	2.13
311	403.7	2.17
312	389.2	2.38
313	441	2.78
314	463.4	2.66
315	432.5	2.24
316	433.5	2.05
317	447.3	2.07
318	454.3	1.79
319	456.5	1.85
320	456.5	1.84
321	499.3	2.57
322	454.4	2.08
323	445.4	2.85
324	460.4	2.1
325	509.5	2.71
326	455.3	2.43
327	450.3	2.39
328	426.3	1.91

Cmpd No.	LC-MS M+1	LC-RT min
329	393.3	2.23
330	465.4	2.88
331	477.3	2.81
332	511.5	3.07
333	443.5	2.83
334	511.5	3.1
335	439.5	2.25
336	493.5	3.03
337	439.5	2.25
338	451.5	2.15
339	467.3	2.5
340	445.4	2.95
341	443.4	2.73
342	468.4	2.39
343	493.4	2.76
344	443.3	2.69
345	453.3	2.05
346	407.5	2.2
347	430.5	1.95
348	430.5	2.43
349	430.5	1.98
350	444.5	1.85
351	444.5	2.24
352	444.5	1.89
353	426.1	1.93
354	426.1	2.33
355	453.3	2.29
356	453.3	2.29

Cmpd No.	LC-MS M+1	LC-RT min
357	407.3	2.26
358	477.3	2.8
359	491.3	2.24
360	473.1	2.32
361	449.3	2.4
362	469.1	2.58
363	469.1	2.58
364	407.7	2.3
365	435.3	2.57
366	437.3	2.18
367	421.1	2.44
368	421.1	2.43
369	449.3	2.67
370	421	2.84
371	439.5	2.31
372	453.5	2.46
373	455.5	2.11
374	425.3	2.31
375	453.5	2.88
376	439.5	2.76
377	441.5	2.3
401	426.3	1.72
402	426.3	1.72
403	509.3	2.73
404	393.2	1.95
405	421.2	2.29
406	421.2	2.2
407	435.4	2.23

Cmpd No.	LC-MS M+1	LC-RT min
408	435.4	2.16
409	421.2	2.15
410	421.1	2.4
411	435.2	2.24
412	405.2	2.09
413	435.4	2.16
414	435.2	2.17
415	449.2	2.25
416	435.3	2.25
417	419.4	2.06
418	421.2	2.57
419	423.2	2.15
420	407.2	2.42
501	467.3	2.14
502	469.3	2.44
503	483.3	2.24
504	455.4	2.91
505	367.9	2.79
506	339.9	2.38
507	469.3	2.97
508	441.3	2.54
509	455.5	2.13
510	469.5	2.37
511	471.3	2.28
512	467.3	2.14
513	483.3	2.25
514	483.3	2.24

[002239] Methods:

[002240] (A) Micromass MUX LCT 4 channel LC/MS, Waters 60F pump, Gilson 215 4 probe autosampler, Gilson 849 injection module, 1.5 mL/min/column flow rate, 10-99% CH_3CN (0.035 % TFA) / H_2O (0.05 % TFA) gradient, Phenomenex Luna 5u C18 columns (50 x 4.60 mm), Waters MUX UV-2488 UV detector, Cedex 75 ELSD detectors.

[002241] (B) PESciex API-150-EX LC/MS, Shimadzu LC-8A pumps, Gilson 215 autosampler, Gilson 819 injection module, 3.0 mL/min flow rate, 10-99% CH_3CN (0.035 % TFA) / H_2O (0.05 % TFA) gradient, Phenomenex Luna 5u C18 column (50 x 4.60 mm), Shimadzu SPD-10A UV/Vis detector, Cedex 75 ELSD detector.

[002242] (C) PESciex API-150-EX LC/MS, Shimadzu LC-8A pumps, Gilson 215 autosampler, Gilson 819 injection module, 3.0 mL/min flow rate, 40-99% CH_3CN (0.035 % TFA) / H_2O (0.05 % TFA) gradient, Phenomenex Luna 5u C18 column (50 x 4.60 mm), Shimadzu SPD-10A UV/Vis detector, Cedex 75 ELSD detector.

[002243] ASSAYS FOR DETECTING AND MEASURING NaV INHIBITION PROPERTIES OF COMPOUNDS

[002244] A) Optical methods for assaying NaV inhibition properties of compounds:

[002245] Compounds of the invention are useful as antagonists of voltage-gated sodium ion channels. Antagonist properties of test compounds were assessed as follows. Cells expressing the NaV of interest were placed into microtiter plates. After an incubation period, the cells were stained with fluorescent dyes sensitive to the transmembrane potential. The test compounds were added to the microtiter plate. The cells were stimulated with either a chemical or electrical means to evoke a NaV dependent membrane potential change from unblocked channels, which was detected and measured with trans-membrane potential-sensitive dyes. Antagonists were detected as a decreased membrane potential response to the stimulus. The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J **69**(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol **4**(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR®) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today **4**(9): 431-439).

[002246] B) VIPR® optical membrane potential assay method with chemical stimulation

[002247] Cell Handling and Dye Loading

[002248] 24 hours before the assay on VIPR, CHO cells endogenously expressing a NaV1.2 type voltage-gated NaV are seeded in 96-well poly-lysine coated plates at 60,000 cells per well. Other subtypes are performed in an analogous mode in a cell line expressing the NaV of interest.

1) On the day of the assay, medium is aspirated and cells are washed twice with 225 µL of Bath Solution #2 (BS#2).
2) A 15 uM CC2-DMPE solution is prepared by mixing 5 mM coumarin stock solution with 10% Pluronic 127 1:1 and then dissolving the mix in the appropriate volume of BS#2.
3) After bath solution is removed from the 96-well plates, the cells are loaded with 80 µL of the CC2-DMPE solution. Plates are incubated in the dark for 30 minutes at room temperature.
4) While the cells are being stained with coumarin, a 15 µL oxonol solution in BS#2 is prepared. In addition to $DiSBAC_2(3)$, this solution should contain 0.75 mM ABSC1 and 30 µL veratridine (prepared from 10 mM EtOH stock, Sigma #V-5754).
5) After 30 minutes, CC2-DMPE is removed and the cells are washed twice with 225 µL of BS#2. As before, the residual volume should be 40 µL.
6) Upon removing the bath, the cells are loaded with 80 µL of the $DiSBAC_2(3)$ solution, after which test compound, dissolved in DMSO, is added to achieve the desired test concentration to each well from the drug addition plate and mixed thoroughly. The volume in the well should be roughly 121 µL. The cells are then incubated for 20-30 minutes.
7) Once the incubation is complete, the cells are ready to be assayed on VIPR® with a sodium addback protocol. 120 µL of Bath solution #1 is added to stimulate the NaV dependent depolarization. 200 µL tetracaine was used as an antagonist positive control for block of the NaV channel.

[002249] Analysis of VIPR® Data:

[002250] Data are analyzed and reported as normalized ratios of background-subtracted emission intensities measured in the 460 nm and 580 nm channels. Background intensities are then subtracted from each assay channel. Background intensities are obtained by measuring the emission intensities during the same time periods from identically treated assay wells in which there are no cells. The response as a function of time is then reported as the ratios obtained using the following formula:

$$(\text{intensity}_{460\,nm} - \text{background}_{460\,nm})$$

$R(t) = \frac{\text{(intensity}_{580\text{ nm}} - \text{background}_{580\text{ nm}})}{\text{(intensity}_{580\text{ nm}} - \text{background}_{580\text{ nm}})}$

[002251] The data is further reduced by calculating the initial (R_i) and final (R_f) ratios. These are the average ratio values during part or all of the pre-stimulation period, and during sample points during the stimulation period. The response to the stimulus $R = R_f/R_i$ is then calculated. For the Na^+ addback analysis time windows, baseline is 2-7 sec and final response is sampled at 15-24 sec.

[002252] Control responses are obtained by performing assays in the presence of a compound with the desired properties (positive control), such as tetracaine, and in the absence of pharmacological agents (negative control). Responses to the negative (N) and positive (P) controls are calculated as above. The compound antagonist activity A is defined as:

$A = \frac{R-P}{N-P} * 100$. where R is the ratio response of the test compound

Solutions [mM]

Bath Solution #1: NaCl 160, KCl 4.5, $CaCl_2$ 2, $MgCl_2$ 1, HEPES 10, pH 7.4 with NaOH

Bath Solution #2 TMA-Cl 160, $CaCl_2$ 0.1, $MgCl_2$ 1, HEPES 10, pH 7.4 with KOH (final K concentration ~ 5 mM)

CC2-DMPE: prepared as a 5 mM stock solution in DMSO and stored at -20°C

$DiSBAC_2(3)$: prepared as a 12 mM stock in DMSO and stored at -20°C

ABSC1: prepared as a 200 mM stock in distilled H_2O and stored at room temperature

[002253] Cell Culture

[002254] CHO cells are grown in DMEM (Dulbecco's Modified Eagle Medium; GibcoBRL #10569-010) supplemented with 10% FBS (Fetal Bovine Serum, qualified; GibcoBRL #16140-071) and 1% Pen-Strep (Penicillin-Streptomycin; GibcoBRL #15140-122). Cells are grown in

vented cap flasks, in 90% humidity and 10% CO_2, to 100% confluence. They are usually split by trypsinization 1:10 or 1:20, depending on scheduling needs, and grown for 2-3 days before the next split.

[002255] C) VIPR® optical membrane potential assay method with electrical stimulation

[002256] The following is an example of how NaV1.3 inhibition activity is measured using the optical membrane potential method#2. Other subtypes are performed in an analogous mode in a cell line expressing the NaV of interest.-

[002257] HEK293 cells stably expressing NaV1.3 are plated into 96-well microtiter plates. After an appropriate incubation period, the cells are stained with the voltage sensitive dyes CC2-DMPE/DiSBAC2(3) as follows.

[002258] ***Reagents:***

100 mg/mL Pluronic F-127 (Sigma #P2443), in dry DMSO

10 mM DiSBAC$_2$(3) (Aurora #00-100-010) in dry DMSO

10 mM CC2-DMPE (Aurora #00-100-008) in dry DMSO

200 mM ABSC1 in H_20

Hank's Balanced Salt Solution (Hyclone #SH30268.02) supplemented with 10 mM HEPES (Gibco #15630-080)

[002259] ***Loading protocol:***

[002260] **2X CC2-DMPE = 20 µM CC2-DMPE:** 10 mM CC2-DMPE is vortexed with an equivalent volume of 10% pluronic, followed by vortexing in required amount of HBSS containing 10 mM HEPES. Each cell plate will require 5 mL of 2X CC2-DMPE. 50 µL of 2X CC2-DMPE is to wells containing washed cells, resulting in a 10 µM final staining concentration. The cells are stained for 30 minutes in the dark at RT.

[002261] **2X DISBAC$_2$(3) with ABSC1 = 6µM DISBAC$_2$(3) and 1 mM ABSC1:** The required amount of 10 mM DISBAC$_2$(3) is added to a 50 ml conical tube and mixed with 1 µL 10% pluronic for each mL of solution to be made and vortexed together. Then HBSS/HEPES is added to make up 2X solution. Finally, the ABSC1 is added .

[002262] The 2X DiSBAC$_2$(3) solution can be used to solvate compound plates. Note that compound plates are made at 2X drug concentration. Wash stained plate again, leaving residual

volume of 50 µL. Add 50 uL/well of the 2X DiSBAC$_2$(3) w/ ABSC1. Stain for 30 minutes in the dark at RT.

[002263] The electrical stimulation instrument and methods of use are described in ION Channel Assay Methods PCT/US01/21652, herein incorporated by reference. The instrument comprises a microtiter plate handler, an optical system for exciting the coumarin dye while simultaneously recording the coumarin and oxonol emissions, a waveform generator, a current- or voltage-controlled amplifier, and a device for inserting electrodes in well. Under integrated computer control, this instrument passes user-programmed electrical stimulus protocols to cells within the wells of the microtiter plate.

[002264] Reagents

[002265] Assay buffer #1

140 mM NaCl, 4.5 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 10 mM HEPES, 10 mM glucose, pH 7.40, 330 mOsm

Pluronic stock (1000X): 100 mg/mL pluronic 127 in dry DMSO
Oxonol stock (3333X): 10 mM DiSBAC$_2$(3) in dry DMSO
Coumarin stock (1000X): 10 mM CC2-DMPE in dry DMSO
ABSC1 stock (400X): 200 mM ABSC1 in water

[002266] Assay Protocol

1. Insert or use electrodes into each well to be assayed.
2. Use the current-controlled amplifier to deliver stimulation wave pulses for 3 s. Two seconds of pre-stimulus recording are performed to obtain the un-stimulated intensities. Five seconds of post-stimulation recording are performed to examine the relaxation to the resting state.

[002267] Data Analysis

[002268] Data are analyzed and reported as normalized ratios of background-subtracted emission intensities measured in the 460 nm and 580 nm channels. Background intensities are then subtracted from each assay channel. Background intensities are obtained by measuring the emission intensities during the same time periods from identically treated assay wells in which

there are no cells. The response as a function of time is then reported as the ratios obtained using the following formula:

$$R(t) = \frac{(\text{intensity}_{460\ nm} - \text{background}_{460\ nm})}{(\text{intensity}_{580\ nm} - \text{background}_{580\ nm})}$$

[002269] The data is further reduced by calculating the initial (R_i) and final (R_f) ratios. These are the average ratio values during part or all of the pre-stimulation period, and during sample points during the stimulation period. The response to the stimulus $R = R_f/R_i$ is then calculated.

[002270] Control responses are obtained by performing assays in the presence of a compound with the desired properties (positive control), such as tetracaine, and in the absence of pharmacological agents (negative control). Responses to the negative (N) and positive (P) controls are calculated as above. The compound antagonist activity A is defined as:

$A = \frac{R-P}{N-P} * 100$. where R is the ratio response of the test compound.

[002271] ELECTROPHYSIOLOGY ASSAYS FOR NaV ACTIVITY AND INHBITION OF TEST COMPOUNDS

[002272] Patch clamp electrophysiology was used to assess the efficacy and selectivity of sodium channel blockers in dorsal root ganglion neurons. Rat neurons were isolated from the dorsal root ganglions and maintained in culture for 2 to 10 days in the presence of NGF (50 ng/ml) (culture media consisted of NeurobasalA supplemented with B27, glutamine and antibiotics). Small diameter neurons (nociceptors, 8-12 μm in diameter) have been visually identified and probed with fine tip glass electrodes connected to an amplifier (Axon Instruments). The "voltage clamp" mode has been used to assess the compound's IC50 holding the cells at – 60 mV. In addition, the "current clamp" mode has been employed to test the efficacy of the compounds in blocking action potential generation in response to current injections. The results of these experiments have contributed to the definition of the efficacy profile of the compounds.

[002273] VOLTAGE-CLAMP assay in DRG neurons

[002274] TTX-resistant sodium currents were recorded from DRG somata using the whole-cell variation of the patch clamp technique. Recordings were made at room temperature (~22° C) with thick walled borosilicate glass electrodes (WPI; resistance 3-4 MΩ) using an Axopatch 200B amplifier (Axon Instruments). After establishing the whole-cell configuration, approximately 15 minutes were allowed for the pipette solution to equilibrate within the cell before beginning recording. Currents were lowpass filtered between 2-5 kHz and digitally sampled at 10 kHz. Series resistance was compensated 60-70% and was monitored continuously throughout the experiment. The liquid junction potential (-7 mV) between the intracellular pipette solution and the external recording solution was not accounted for in the data analysis. Test solutions were applied to the cells with a gravity driven fast perfusion system (SF-77; Warner Instruments).

[002275] Dose-response relationships were determined in voltage clamp mode by repeatedly depolarizing the cell from the experiment specific holding potential to a test potential of +10mV once every 60 seconds. Blocking effects were allowed to plateau before proceeding to the next test concentration.

[002276] Solutions

[002277] Intracellular solution (in mM): Cs-F (130), NaCl (10), $MgCl_2$ (1), EGTA (1.5), $CaCl_2$ (0.1), HEPES (10), glucose (2), pH = 7.42, 290 mOsm.

[002278] Extracellular solution (in mM): NaCl (138), $CaCl_2$ (1.26), KCl (5.33), KH_2PO_4 (0.44), $MgCl_2$ (0.5), $MgSO_4$ (0.41), $NaHCO_3$ (4), Na_2HPO_4 (0.3), glucose (5.6), HEPES (10), CdCl2 (0.4), NiCl2 (0.1), TTX (0.25×10^{-3}).

[002279] CURRENT-CLAMP assay for NaV channel inhibition activity of compounds

[002280] Cells were current-clamped in whole-cell configuration with a Multiplamp 700A amplifier (Axon Inst). Borosilicate pipettes (4-5 MOhm) were filled with (in mM):150 K-gluconate, 10 NaCl, 0.1 EGTA, 10 Hepes, 2 $MgCl_2$, (buffered to pH 7.34 with KOH). Cells were bathed in (in mM): 140 NaCl, 3 KCl, 1 MgCl , 1 CaCl , and 10 Hepes). Pipette potential was zeroed before seal formation; liquid junction potentials were not corrected during acquisition. Recordings were made at room temperature.

[002281] Following these procedures, representative compounds of the present invention were found to possess desired voltage gated sodium channel activity and selectivity.

[002282] ASSAYS FOR DETECTING AND MEASURING L-Type CaV 1.2 INHIBITION PROPERTIES OF COMPOUNDS

[002283] A) Optical methods for assaying CaV inhibition properties of compounds:

[002284] Compounds of the invention are useful as antagonists of voltage-gated calcium ion channels. Antagonist properties of test compounds were assessed as follows. Cells expressing the CaV of interest were placed into microtiter plates. After an incubation period, the cells were stained with fluorescent dyes sensitive to the transmembrane potential. The test compounds were added to the microtiter plate. The cells were stimulated with electrical means to evoke a CaV dependent membrane potential change from unblocked channels, which was detected and measured with trans-membrane potential-sensitive dyes. Antagonists were detected as a decreased membrane potential response to the stimulus. The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J **69**(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol **4**(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR®) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today **4**(9): 431-439).

[002285] VIPR® optical membrane potential assay method with electrical stimulation

Positive Control (100% Block)

The positive control for this assay was 125uM mibefradil, achieved by adding 25uL of 250uM solution to the assay plates containing 25uL of assay buffer. Each assay plate included positive control wells.

Negative Control (No Block)

The negative (baseline) control for this assay was DMSO. This was achieved by adding 25uL of 1% DMSO (in assay buffer) to the assay plates containing 25uL of assay buffer. Each assay plate included negative control wells.

Background Subtraction

Fluorescence background from plastic in assay plates (or from the assay buffer) was assessed by running a cell-free plate through the EVIPR under the same optical configuration. The average background values for each row and each wavelength were subtracted in MOD3 prior to ratio change and activity calculations.

Reagents

Assay Buffers:

Bath Y (Prepared by Vertex Lab Support)

140mM TMA-Cl

4.5mM KCl

1mM $MgCl_2$

10mM HEPES, pH7.4

10mM glucose

Osmolarity = 295mOsm (280-310 acceptable range)

500mM $BaCl_2$ (Sigma #B0750), in H_2O

100 mg/mL Pluronic F-127 (Sigma #P2443), in dry DMSO

10mM $DiSBAC_2(3)$ (Aurora #00-100-010) in dry DMSO

10mM CC2-DMPE (Aurora #00-100-008) in dry DMSO

200mM Acid Yellow 17 (Aurora #VABSC) in H_2O

Assay Volume

50uL

DMSO Conc in Assay

0.5% (1uL of 75% DMSO/25% water, dilution factor of 160)

Incubation Time of Compounds

20-25 minutes

Instrumentation

This screen was conducted on the Allegro™ system. The system is diagrammed below: The Allegro™ is equipped with a compound plate storage unit (stacker). The stacker holds a set of trays (each tray holds 12 compound plates). The libraries were received from Compound Management, as pre-spotted (1uL/well of compound and controls) intermediate plates in 384-well format, as a 1.6mM stock solution in 75% DMSO/25% de-ionized H_2O. The plates are diluted in 80uL oxonol dye solution to create a 2X stock. Three EVIPR readers are integrated to the Allegro system by a Mitsubishi robotic arm. Only one EVIPR was used per run.

Instrumentation Settings

Optical:

Read Frequency: 10Hz

Excitation Wavelength: 400nm

Emission Wavelengths: 460nm and 560nm

Electrical Stimulation:

Pulse Width: 11.1 ms

Stimulation Current: 0.8 amps

Stimulation Frequency: 90Hz

Pre-stimulation time: 2 s

Stimulation time: 3 s

Post-stimulation time: 1 s

Waveform: Biphasic Square Wave

Plate Washer Settings:

Settings for ELx405 washer will leave a residual volume of 25uL.

Plate Type: 384

# of cycles: 3

Soak/shake: No

Dispense: dispense volume 100

	dispense flow rate	1
	dispense height	80
	horizontal disp pos	-20
	horiz y disp pos	-5
Aspirate:	aspirate height	48
	horizontal aspr pos	-18
	horiz y asp pos	-5
	aspiration rate	0
	aspiration delay	0
	final asp delay	500

Assay Procedure

Procedure run on HTS Allegro™:

1. **Carousel**: Assay plates (Cell plates) loaded into carousel module #1 (CO_2 = 5%, ambient temperature and Rh)
2. **Barrier:** Assay plates retrieved from carousel and passed through environmental barrier (The remaining steps are conducted at room temperature and ambient CO_2)
3. **Washer:** Assay plates washed with Bath Y on Biotek ELx405.
4. **MultiReagent Dispenser (MRD):** 25uL of CC2-DMPE (and equal volume Pluronic) in Bath Y added to each well to make 10uM.
5. **Barrier:** Assay plates passed through barrier.
6. **Carousel:** 30-minute incubation at room temperature.
7. **Barrier:** Assay plates passed through barrier.
8. **Washer:** Assay plates washed with Bath Y on Biotek ELx405
9. **High Density Transfer Station:**
    a. 80uL oxonol dye loading solution (4uM DiSBAC2(3), 1mM VABSC and 30mM $BaCl_2$ in BathY) added to compound plates (pre-spotted with 1uL compound) using a MultiDrop (offline)

b. Plates mixed (3 times 20uL) on CyBiWell (offline). Plates loaded onto compound tray.

c. Compound tray retrieved from compound tray stacker and compound plate barcodes read.

d. Assay plate barcode read and moved to SciClone deck

e. 25uL compound plus oxonol aspirated from compound plate on SciClone deck and transferred to assay plate.

i. Final assay volume = 50uL

ii. Final compound concentration = 10uM

f. SciClone tips washed in DMSO and 5% ethanol in water to remove external carry-over.

10. **Carousel:** Assay plates incubated for 20 minutes at RT

11. **Barrier:** Assay plates passed through final barrier

12. **Mitsubishi Robotic Arm:** Retrieves assay plate from barrier output, delivers cell plates to EVIPR 384-1, and sends command to initiate EVIPR run.

Assay Window

Assay window criteria:

Passing plates ≤ 0.5, rejected plates > 0.5

$$\textit{Assay Window} = \frac{3(SD_{FullBlock} + SD_{Baseline})}{(AVE_{Baseline} - AVE_{FullBlock})} = 1 - Z'$$

Data Reduction

The EVIPR files were reduced to decrease the amount of data pumped into the database. Two "windows" of interest were filtered out of each EVIPR file. Each window is a slice of the response measured in each well. The first window is measured before stimulation. The second

window samples the peak of the response. The ratio of the two is used to determine the response size.

Data Analysis

Once the data were collected on the VIPR, they were archived and uploaded, in reduced form, to Mod3. Once in Mod3, each individual assay plate was QC'ed (looking for acceptable window and dynamic range).

[002286] HERG Assay

[002287] hERG-inhibition was assayed in a Chinese Hamster Lung cell line (CHL) stably transduced with the structural gene for hERG. Cells express high numbers of hERG channels resulting in 500pA to 1.5nA of hERG outward K^+ currents. The method used a planar patch instrument (IonWorks HT, Molecular Devices) that allowed medium-throughput electrophysiology measurements in 384-well format. The potency of hERG inhibition was measured at 1.1 μM, 3.3μM, 10μM, and 30μM of the compound studied. The compound was added from a 3x aqueous addition buffer.

[002288] The compounds of the present invention exhibit a desirably low activity against hERG.

[002289] CYP-450 Isozyme assay

[002290] Compound preparation:

1. The desired compound was plated (2mM in 75% DMSO/25% H_2O) with a Pieso Sample Distribution Robot (PSDR™) at 8nL per well.
2. The compound was centrifuged briefly at approximately 1000rpm to shift the compound drop to the bottom of the well.
3. PVP 10K (excipient, 0.2% in 75% DMSO/25% H_2O) was plated with a PSDR™ at 100nL per well.
4. The compound and PVP 10K were centrifuged briefly at approximately 1000rpm to ensure an adequate mix of compound and excipient.
5. The dry-down of the plates was initiated using house vacuum for at least 3 hours.
6. The plates were transferred to a high vacuum (50millitorr) apparatus and the dry-down process was continued for at least 15 hours.

[002291] The following assay protocol was employed for a desired CYP-450 isozyme (CYP3A4, CYP2C9, CYP1A2, CYP2C19, or CYP2D6).

[002292] **Assay Protocol**

All reagents below were added using a Flying Reagent Dispenser (FRD™).

1. 800 nL of dH_2O was added to the 100% activity control, compound, and background control wells.
2. 800 nL of the appropriate control drug (3A4:clotrimazole, 2C9:miconazole, 1A2:ticlopidine, 2C19:lansoprazole, or 2D6:propanolol; 10uM final dissolved in dH_2O) was added to the drug control wells.
3. 200 nL of 500mM K^+ Phosphate buffer (pH 8.4)was added to the 100% activity control, drug control, and compound wells.
4. 600 nL of Control Insect Baculosomes (PanVera P2315) in 500mM K^+ Phosphate Buffer (pH 8.4) was added to the background control wells. The calculation for this reagent was based on the protein concentration of the 100% activity control wells.
5. The plate was scanned for compound fluorescence using a NanoPlate™ Fluorescence Plate Reader (NPR™).
6. 200 nL of $NADP^+$ (Sigma, 100 μM final) and substrate in 100mM K^+ Phosphate buffer (50mM K^+ Phosphate buffer for 2C9 and 2C19) was added to all wells. Fluorogenic substrate (3A4:5 μM Vivid™ 3A4 Red, 2C9:1μM Vivid™ 2C9 Green, 1A2:2μM Vivid™ 1A2 Blue, 2C19:10μM Vivid™ 2C19 Blue, and 2D6: 10μM Vivid™ 2D6 Blue) was added at a final concentration corresponding to the K_m of the substrate for its pertinent CYP450 isozyme.
7. 400 nL of the desired CYP450 isozyme and recycling buffer (3.3mM glucose-6-phosphate, 0.4units/ml glucose-6-phosphate dehydrogenase, 100mM $MgCl_2$, and 0.00025% Antifoam 289; reagents obtained from Sigma) in 100mM K^+ Phosphate Buffer (50mM K^+ Phosphate buffer for 2C9 and 2C19) were added to the 100% activity control, drug control, and compound wells. The desired isozyme was added to obtain the following final concentrations of the desired isozyme: 5nM CYP3A4, 10nM CYP2C9, 5nM CYP1A2, 5nM CYP2C19, or 20nM CYP2D6.
8. The plate was incubated for 60 minutes at room temperature.

9. The plate was scanned for solution fluorescence using a NanoPlate™ Fluorescence Plate Reader (NPR™).
10. The NPR™ data was converted into a format compatible with importation into a data visualizer and complete the analysis of data acquired.

[002293] The compounds of the present invention exhibit a desirably low activity against one or more of the CYP450 isozymes.

[00229-4] The activity of selected compounds of the present invention against NaV 1.8 channel is shown below in **Table 4**. In **Table 4**, the symbols have the following meaning:

"+++" means < 2 μM; "++" means between 2 μM and 10 μM; and "+" means > 10μM.

[00229-5] **Table 4**

Cmpd #	IC50
101	+++
102	++
103	+
104	+
105	++
106	+
107	++
108	++
109	++
110	+
111	+
112	+
113	++
114	++
115	+
116	++
117	++
118	++
119	+
120	++
121	++
122	++
123	+
124	+
125	+
126	+
127	+++
128	++
129	+
130	+
131	+
132	++
133	++
134	++
135	+++
136	+

Cmpd #	IC50
137	++
138	+
139	++
140	+
141	+
142	++
143	++
144	++
145	+
146	+
147	++
148	++
149	+
150	+
151	+
152	+
153	++
154	+
155	+
156	++
157	++
158	+
159	++
160	+
162	++
163	++
164	+
165	++
166	+
167	+
168	++
169	+
170	+
172	+
173	+
174	++

Cmpd #	IC50
175	+
176	+
177	++
178	+
179	+
180	++
181	+
182	++
183	++
184	++
185	++
201	++
203	+
204	+
205	+
206	++
207	+++
208	+++
210	++
211	+
212	+
213	+
214	+
215	+
216	+
217	+
218	++
219	+++
220	+
221	+
222	+
223	+
224	+
225	+
226	+
227	+

Cmpd #	IC50
228	+
229	+
230	++
231	++
232	++
233	+
234	+
235	++
236	+
237	+
238	+
239	+
240	+
241	+
242	++
243	+
244	+
245	+
246	+
247	+
248	++
249	++
250	+++
251	++
252	+
253	+
254	++
255	+
256	++
257	++
258	+++
259	+++
260	+++
261	++
262	++
264	+

Cmpd #	IC50
265	+
266	++
267	+
268	+
269	+
270	++
271	+
272	++
273	++
274	++
275	++
276	++
277	++
278	+
279	++
301	++
302	+
303	++
304	+
305	++
306	+
307	+
308	+
309	+++
310	++
311	+++
312	++
314	++
315	+
316	+
317	++
318	+
319	++
320	++
321	+
322	+

Cmpd #	IC50
323	+
324	+
325	+
326	+
327	+
328	+
329	++
335	++
337	++
338	+
339	++
340	+++
342	+
345	++
346	++
347	+

Cmpd #	IC50
348	+
349	+
350	+
351	++
352	+
353	+
354	++
355	+
356	+
357	+
358	+
359	+
360	+
361	++
362	+
363	++


Cmpd #	IC50
364	+++
365	+++
366	++
367	+++
368	++
369	+++
370	+++
371	+++
372	+++
373	++
374	+
375	+
376	+
377	++
401	+
402	+

Cmpd #	IC50
403	+
404	+
405	+
406	+
407	+
408	+
409	+
410	+
411	++
412	+
413	++
414	++
415	++
416	+
417	+
418	++

Cmpd #	IC50
419	++
420	+++
501	++
502	++
503	++
504	++
505	++
506	++
507	++
508	++
509	+
510	++
511	++
512	++
513	+
514	++

CLAIMS


1. A compound of formula **I**:




I;

or a pharmaceutically acceptable salt or derivative thereof, wherein:


R^1 and R^2, taken together with the nitrogen atom, form a substituted ring selected from:




(A)



(B)




(C)



(D)

or



(E);

wherein, in ring (A):

each of m_1 and n_1 is independently 0-3, provided that m_1+n_1 is 2-6;

z_1 is 0-4;

Sp^1 is –O-, -S-, -NR'-, or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR' , -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-, provided that Sp^1 is attached to the carbonyl group through an atom other than carbon;

ring B^1 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w_1 independent occurrences of -R^{11}, wherein w_1 is 0-4;

wherein, in ring (B):

G_2 is -N-, or CH;

each of m_2 and n_2 is independently 0-3, provided that $m_2 + n_2$ is 2-6;

p_2 is 0-2; provided that when G_2 is N, then p_2 is not 0;

q_2 is 0 or 1;

z_2 is 0-4;

Sp^2 is a bond or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-;

ring B^2 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B is optionally substituted with w independent occurrences of –R^{12}, wherein w_2 is 0-4;

wherein, in ring (C) or ring (D):

G_3 is -N-, -CH-NH-, or -CH-CH_2-NH-;

each of m_3 and n_3 is independently 0-3, provided that m_3+n_3 is 2-6;

p_3 is 0-2;

z_3 is 0-4;

each R^{XX} is hydrogen, C_{1-6} aliphatic group, a 3–8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently

selected from nitrogen, oxygen, or sulfur; wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3;

provided that both R^{XX} are not simultaneously hydrogen;

R^{YY} is hydrogen, -COR', $-CO_2R'$, $-CON(R')_2$, -SOR', $-SO_2R'$, $-SO_2N(R')_2$, -COCOR', $-COCH_2COR'$, $-P(O)(OR')_2$, $-P(O)_2OR'$, or -PO(R');

wherein, in ring (E):

each of m_4 and n_4 is independently 0-3, provided that $m_4 + n_4$ is 2-6;

p_4 is 1-2;


z_4 is 0-4;

R^{YZ} is C_1-C_6 aliphatic group, optionally substituted with w_4 independent occurrences of -R^{14}, wherein w_4 is 0-3;

x and y, each, is independently 0-4;

W is OR^{XY};

R^{XY} is hydrogen or a group selected from:



;

wherein:

each of w_A, w_B, w_C, and w_D is independently 0 or 1;

each M is independently selected from hydrogen, Li, Na, K, Mg, Ca, Ba, $-N(R^7)_4$, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or $-R^6$; wherein 1 to 4 $-CH_2$ radicals of the alkyl or alkenyl group, other than the $-CH_2$ that is bound to Z, is optionally replaced by a heteroatom group selected from O, S, S(O), $S(O_2)$, or $N(R^7)$; and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a substituent selected from oxo, $-OR^7$, $-R^7$, $N(R^7)_2$, $N(R^7)_3$, R^7OH, -CN, $-CO_2R^7$, $-C(O)-N(R^7)_2$, $S(O)_2-N(R^7)_2$, $N(R^7)-C(O)-R^7$, $C(O)R^7$, $-S(O)_n-R^7$, OCF_3, $-S(O)_n-R^6$, $N(R^7)-S(O)_2(R^7)$, halo, $-CF_3$, or $-NO_2$;

n is 0-2;

M' is H, C_1-C_{12}-alkyl, C_2-C_{12}-alkenyl, or $-R^6$; wherein 1 to 4 $-CH_2$ radicals of the alkyl or alkenyl group is optionally replaced by a heteroatom group selected from O, S, S(O), $S(O_2)$, or $N(R^7)$; and wherein any hydrogen in said alkyl, alkenyl or R^6 is optionally replaced with a

substituent selected from oxo, -OR^7, - R^7, -N(R^7)$_2$, N(R^7)$_3$, - R^{7}OH, -CN, -CO$_2$ R^7, –C(O)-N(R^7)$_2$, -S(O)$_2$-N(R^7)$_2$, -N(R^7)-C(O)- R^7, -C(O) R^7, -S(O)$_n$- R^7, -OCF$_3$, -S(O)$_n$-R^6, -N(R^7)-S(O)$_2$(R^7), halo, -CF$_3$, or -NO$_2$;

Z is -CH$_2$-, -O-, -S-, -N(R^7)$_2$-; or,

when M is absent, then Z is hydrogen, =O, or =S;

Y is P or S, wherein when Y is S, then Z is not S;

X is O or S;

each R^7 is independently selected from hydrogen, or C$_1$-C$_4$ aliphatic, optionally substituted with up to two Q$_1$;

each Q$_1$ is independently selected from a 3-7 membered saturated, partially saturated or unsaturated carbocyclic ring system; or a 5-7 membered saturated, partially saturated or unsaturated heterocyclic ring containing one or more hetero atom or heteroatom group selected from O, N, NH, S, SO, or SO$_2$; wherein Q$_1$ is optionally substituted with up to three substituents selected from oxo, -OH, -O(C$_1$-C$_4$ aliphatic), -C$_1$-C$_4$ aliphatic, -NH$_2$, NH(C$_1$-C$_4$ aliphatic), -N(C$_1$-C$_4$ aliphatic)$_2$, -N(C$_1$-C$_4$ aliphatic)-C(O)- C$_1$-C$_4$ aliphatic, -(C$_1$-C$_4$ aliphatic)-OH, -CN, -CO$_2$H, -CO$_2$(C$_1$-C$_4$ aliphatic), –C(O)-NH$_2$, -C(O)-NH(C$_1$-C$_4$ aliphatic), -C(O)-N(C$_1$-C$_4$ aliphatic)$_2$, halo or -CF$_3$;

R^6 is a 5-6 membered saturated, partially saturated or unsaturated carbocyclic or heterocyclic ring system, or an 8-10 membered saturated, partially saturated or unsaturated bicyclic ring system; wherein any of said heterocyclic ring systems contains one or more heteroatoms selected from O, N, S, S(O)$_n$ or N(R^7); and wherein any of said ring systems optionally contains 1 to 4 substituents independently selected from OH, C$_1$-C$_4$ alkyl, O-C$_1$-C$_4$ alkyl or O-C(O)-C$_1$-C$_4$ alkyl;

R^9 is C(R^7)$_2$, O or N(R^7);

each occurrence of R^{11}, R^{12}, R^{13}, R^{14}, R^3, R^4, and R^5 is independently Q-R^X; wherein Q is a bond or is a C$_1$-C$_6$ alkylidene chain wherein up to two non-adjacent methylene units of Q are optionally and independently replaced by –NR-, -S-, -O-, -CS-, -CO$_2$-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -NRCO$_2$-, -SO$_2$NR-, -NRSO$_2$-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO$_2$NR-, -SO-, -SO$_2$-, -PO-, -PO$_2$-, -OP(O)(OR)-, or -POR-; and each occurrence of R^X is independently selected from -R', halogen, =O, =NR', -NO$_2$, -CN, -OR', -SR', -N(R')$_2$, –NR'COR', -NR'CON(R')$_2$, -NR'CO$_2$R', -COR', -CO$_2$R', -OCOR', -CON(R')$_2$, -OCON(R')$_2$, -

SOR', $-SO_2R'$, $-SO_2N(R')_2$, $-NR'SO_2R'$, $-NR'SO_2N(R')_2$, -COCOR', $-COCH_2COR'$, $-OP(O)(OR')_2$, $-P(O)(OR')_2$, $-OP(O)_2OR'$, $-P(O)_2OR'$, $-PO(R')_2$, or $-OPO(R')_2$; and

each occurrence of R is independently hydrogen or C_{1-6} aliphatic group having up to three substituents; and each occurrence of R' is independently hydrogen or C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' has up to four substituents; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

2. The compound according to claim 1, wherein W is OH.

3. The compound according to claim 1, wherein x is 1 and R^3 is present at the 6- or 7-position of the quinazoline ring.

4. The compound according to claim 3, wherein R^3 is selected from -Cl, $-CH_3$, $-CH_2CH_3$, -F, $-CF_3$, $-OCF_3$, $-CONHCH_3$, $-CONHCH_2CH_3$, -CONH(cyclopropyl), $-OCH_3$, $-NH_2$, $-OCH_2CH_3$, or -CN.

5. The compound according to claim 4, wherein x is 1 and R^3 is at the 7-position of the quinazoline ring and is $-CH_3$.


6. The compound according to claim 1, wherein y is 0.

7. The compound according to claim 1, wherein y is 1, and R^5 is halo.

8. The compound according to claim 7, wherein R^5 is halo at the 6-position.

9. The compound according to claim 1, wherein z_1, z_2, z_3, and z_4 each is 0.

10. The compound according to claim 1, wherein said compound has formula **I-A**:



I-A.

11. The compound according to claim 10, wherein Sp^1 is -O-, -O-CH_2-, -S-, or –NH-, or -NH-CH_2-.

12. The compound according to claim 10, wherein each of m_1 and n_1 is 2.

13. The compound according to claim 10, wherein ring B^1 is a 5-6 membered, saturated, monocyclic heterocyclic ring having having 1-2 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w independent occurrences of -R^{11}, wherein w_1 is 0-4.

14. The compound according to claim 13, wherein ring B1 is tetrahydrofuranyl, tetrahydro-[*2H*]-pyranyl, pyridyl, or phenyl.

15. The compound according to claim 10, wherein Sp^1 is a bond, O, or –O-CH_2-; R is hydrogen; and n_1 and m_1 are both simultaneously 2.

16. The compound according to claim 10, wherein:

n_1 and m_1 each is 2;

R^{XY} is hydrogen;

y is 0 or 1 and R^5 is fluoro;

x is 1 and R^3 is Me at 7-position or fluoro at 6-position;


z_1 is 0;

Sp^1 is –O-CH_2-;

w_1 is 0; and

ring B^1 is tetrahydrofuran-3-yl, phenyl, pyridine-3-yl, pyridine-4-yl, or tetrahydro[2H]-pyran-4-yl.

17. The compound according to claim 1, wherein said compound has formula **I-B**:



I-B.

18. The compound according to claim 17, wherein G_2 is CH and p_2 is 1 or 2.

19. The compound according to claim 17, wherein m_2 and n_2 each is 1 or 2.

20. The compound according to claim 17, wherein m_2 is 1 and n_2 each is 2.

21. The compound according to claim 17, wherein n_2 is 1, and m_2 is 3.

22. The compound according to claim 17, wherein Sp^2 is -O- or $-O-CH_2-$.

23. The compound according to claim 17, wherein ring B^2 is a 5-6 membered, saturated, monocyclic heterocyclic ring having having 1-2 heteroatoms selected from O, S, or N, wherein ring B^2 is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4.

24. The compound according to claim 23, wherein ring B^2 is tetrahydrofuranyl, tetrahydro[2*H*]pyranyl, or pyridyl.

25. The compound according to claim 17, wherein


Sp^2 is a bond, O, or $-O-CH_2-$;

p_2 is 1;

R is hydrogen; and


n_2 is 1 and m_2 is 2 or 3.

26. The compound according to claim 17, wherein said compound has formula **I-B-i** or formula **I-B-ii**:



I-B-i

or



I-B-ii;

wherein:

m_2 is 2 or 3; and

Sp^2 is $-O-$ or $-O-CH_2$.

27. The compound of formula **I-B-i** according to claim 26, wherein:

p_2 is 1;

m_2 is 3;

Sp^2 is -O-;

y is 0 or 1, and R^5 is fluoro;

x is 1 and R^3 is 7-Me; and

ring B^2 is tetrahydrofuranyl.

28. The compound of formula **I-B-i** according to claim 26, wherein:

p_2 is 0 or 1;

m_2 is 2;

Sp^2 is -O- or -O-CH_2-;

y is 0;

x is 1 and R^3 is 7-Me; and

ring B^2 is tetrahydrofuranyl, tetrahydro[2H]pyranyl, pyridyl, or phenyl.

29. The compound of formula **I-B-ii** according to claim 26, wherein:

p_2 is 0;

m_2 is 2;

Sp^2 is a bond;


y is 0 or 1, and R^5 is fluoro;

x is 1 and R^3 is 7-Me or 6-F; and


ring B^2 is cyclopropyl optionally substituted with C1-C4 alkyl, or pyridyl.

30. The compound of formula **I-B-i** or formula **I-B-ii** according to claim 17, wherein R^{XY} is hydrogen.

31. The compound according to claim 1, wherein said compound has formula **I-C** or formula **I-D**:



I-C



I-D.

32. The compound according to claim 31, wherein one R^{XX} is hydrogen and the other R^{XX} is not hydrogen.

33. The compound according to claim 31, wherein one R^{XX} is hydrogen and the other R^{XX} is C1-C6 alkyl optionally substituted with halo.

34. The compound according to claim 31, wherein both R^{XX} are simultaneously C1-C6 alkyl.


35. The compound according to claims 33 or 34, wherein said alkyl is selected from methyl, ethyl, isopropyl, *n*-propyl, *n*-butyl, *sec*-butyl, or *t*-butyl.

36. The compound according to claim 31, wherein p_3 is 0.


37. The compound according to claim 31, wherein m_3 and n_3 each is 2.

38. The compound according to claim 31, wherein R^{YY} is hydrogen.

39. The compound according to claim 31, wherein said compound has formula **I-C-i** or formula **I-D-i**:



I-C-i



I-D-i.

40. The compound according to claim 39, wherein R^{XX} is C1-C6 alkyl.

41. The compound according to claim 39, wherein x is 1, and R^3 is C1-C4 alkyl at the 7-position.

42. The compound according to claim 39, wherein x is 1 and R^3 is F, CN, or CF_3 at the 6-position.


43. The compound according to claim 39, wherein R^{XX} is methyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, or *t*-butyl.

44. The compound according to claim 41, wherein R^3 is methyl, *n* -propyl, isopropyl, *n* -butyl, *sec*-butyl, or *t*-butyl.

45. The compound according to claim 39, wherein R^{XY} is hydrogen, and y is 0.

46. The compound according to claim 39, wherein R^{XY} is hydrogen, y is 1 and R^5 is 6-F.

47. The compound according to claim 31, wherein said compound has formula **I-C-ii**:




I-C-ii.

48. The compound according to claim 47, wherein R^3 is methyl at the 7-position of the quinazoline ring.

49. The compound according to claim 47, wherein R^{XX} is $CH_2C(O)OH$ or $CH_2C(O)NH_2$.

50. The compound according to claim 1, wherein said compound has formula **I-E**:



I-E.

51. The compound according to claim 50, wherein p_4 is 1.

52. The compound according to claim 50, wherein m_4 and n_4 each is 2.

53. The compound according to claim 50, wherein n_4 is 1, m_4 is 3, z_4 is 0, p_4 is 1, y is 0 or 1, and x is 1.

54. The compound according to claim 50, wherein n_4 is 1, m_4 is 2, z_4 is 0, p_4 is 1, y is 0 or 1, and x is 1.

55. The compound according to claim 50, wherein n_4 is 1, m_4 is 3, z_4 is 0, p_4 is 1, y is 0 or 1, x is 1, and R and R^{XY} both are hydrogen.

56. The compound according to claim 50, wherein n_4 is 1, m_4 is 2, z_4 is 0, p_4 is 1, y is 0 or 1, x is 1, and R and R^{XY} both are hydrogen.

57. The compound according to claim 50, wherein R^{YZ} is C1-C4 alkyl group optionally substituted with w_4 independent occurrences of $-R^{14}$, wherein w_4 is 0-3.

58. The compound according to claim 50, wherein:

n_4 is 1 and m_4 is 3;

p_4 is 1;

z_4 is 0;

R^{YZ} is C1-C6 alkyl, wherein up to two $-CH_2-$ groups therein is optionally replaced by -O-;

y is 0 or 1, and R^5 is 6-fluoro; and

x is 1 and R^3 is C1-C4 alkyl.

59. The compound according to claim 50, wherein:

n_4 is 1 and m_4 is 2;

p_4 is 1;

z_4 is 0;

R^{YZ} is C1-C6 alkyl, wherein up to two $-CH_2-$ groups therein is optionally replaced by -O-;

y is 0 or 1, and R^5 is 6-fluoro; and

x is 1 and R^3 is C1-C4 alkyl.

60. The compound according to claim 50, wherein:

n_4 is 1 and m_4 is 3;

p_4 is 1;


z_4 is 0;

R^{YZ} is benzyl;

y is 0 or 1, and R^5 is 6-fluoro; and

x is 1 and R^3 is C1-C4 alkyl.


61. A compound according having formula **I-F**:




I-F;

wherein:


R^1 and R^2, taken together with the nitrogen atom, form a substituted ring selected from:




(A)



(B)



(C)



(D) or (E);

wherein, in ring (A):

each of m_1 and n_1 is independently 0-3, provided that m_1+n_1 is 2-6;

z_1 is 0-4;

Sp^1 is –O-, -S-, –NR'-, or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-, provided that Sp^1 is attached to the carbonyl group through an atom other than carbon;

ring B^1 is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1-4 heteroatoms selected from O, S, or N, wherein ring B^1 is optionally substituted with w_1 independent occurrences of -R^{11}, wherein w_1 is 0-4;

wherein, in ring (B):

G_2 is -N-, or CH;

each of m_2 and n_2 is independently 0-3, provided that $m_2 + n_2$ is 2-6;

p_2 is 0-2; provided that when G_2 is N, then p_2 is not 0;

q_2 is 0 or 1;

z_2 is 0-4;

Sp^2 is a bond or a C1-C6 alkylidene linker, wherein up to two methylene units are optionally and independently replaced by -O-, -S-, –CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO_2-, -OCO-, -NR'CO_2-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -SO, -SO_2-, -NR'-, -SO_2NR'-, NR'SO_2-, or -NR'SO_2NR'-;

ring B is a 4-8 membered, saturated, partially unsaturated, or aromatic, monocyclic heterocyclic ring having having 1 -4 heteroatoms selected from O, S, or N, wherein ring B is optionally substituted with w independent occurrences of $-R^{12}$, wherein w_2 is 0-4;

wherein, in ring (C) or ring (D):

G_3 is -N-, -CH-NH-, or -CH-CH_2-NH-;

each of m_3 and n_3 is independently 0-3, provided that m_3+n_3 is 2-6;

p_3 is 0-2;

z_3 is 0-4;

each R^{XX} is hydrogen, C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^{XX} is optionally substituted with w_3 independent occurrences of $-R^{13}$, wherein w_3 is 0-3;

provided that both R^{XX} are not simultaneously hydrogen;

R^{YY} is hydrogen, -COR', -CO_2R', -CON(R')$_2$, -SOR', -SO_2R', -SO_2N(R')$_2$, -COCOR', -$COCH_2$COR', -P(O)(OR')$_2$, -P(O)$_2$OR', or -PO(R');

wherein, in ring (E):

each of m_4 and n_4 is independently 0-3, provided that m_4+n_4 is 2-6;

p_4 is 1-2;

z_4 is 0-4;

R^{YZ} is C_1-C_6 aliphatic group, optionally substituted with w_4 independent occurrences of $-R^{14}$, wherein w_4 is 0-3;

x and y, each, is independently 0-4;

each occurrence of R^{11}, R^{12}, R^{13}, R^{14}, R^3, R^4, and R^5 is independently Q-R^X; wherein Q is a bond or is a C_1-C_6 alkylidene chain wherein up to two non-adjacent methylene units of Q are optionally and independently replaced by –NR-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -NRCO_2-, -SO_2NR-, -NRSO_2-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO_2NR-, -SO-, -SO_2-, -PO-, -PO_2-, -OP(O)(OR)-, or -POR-; and each occurrence of R^X is independently selected from -R', halogen, =O, =NR', -NO_2, -CN, -OR', -SR', -N(R')$_2$, -NR'COR', -NR'CON(R')$_2$, -NR'CO_2R', -COR', -CO_2R', -OCOR', -CON(R')$_2$, -OCON(R')$_2$, -

SOR', -SO$_2$R', -SO$_2$N(R')$_2$, -NR'SO$_2$N(R')$_2$, -COCOR', -COCH$_2$COR', -OP(O)(OR')$_2$, -P(O)(OR')$_2$, -OP(O)$_2$OR', -P(O)$_2$OR', -PO(R')$_2$, or -OPO(R')$_2$; and

each occurrence of R is independently hydrogen or C_{1-6} aliphatic group having up to three substituents; and each occurrence of R' is independently hydrogen or C_{1-6} aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' has up to four substituents; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

62. The compound according to claim 1, wherein said compound is selected from Table 2.

63. A pharmaceutical composition comprising a compound according to claim 1.

64. A method for treating or lessening the severity of a disease, disorder, or condition selected from acute, chronic, neuropathic, or inflammatory pain, arthritis, migrane, cluster headaches, trigeminal neuralgia, herpetic neuralgia, general neuralgias, epilepsy or epilepsy conditions, neurodegenerative disorders, psychiatric disorders such as anxiety and depression, myotonia, arrythmia, movement disorders, neuroendocrine disorders, ataxia, multiple sclerosis, irritable bowel syndrome, incontinence, visceral pain, osteoarthritis pain, postherpetic neuralgia, diabetic neuropathy, radicular pain, sciatica, back pain, head or neck pain, severe or intractable pain, nociceptive pain, breakthrough pain, postsurgical pain, or cancer pain, said method comprising the step of administering to said patient an effective amount of a composition according to claim 62.

65. The method according to claim 63, wherein the disease, condition, or disorder is implicated in the activation or hyperactivity of voltage-gated sodium channels.

66. The method according to claim 64, wherein the disease, condition, or disorder is acute, chronic, neuropathic, or inflammatory pain.

67. The method according to claim 64, wherein the disease, condition, or disorder is radicular pain, sciatica, back pain, head pain, or neck pain.

68. The method according to claim 64, wherein the disease, condition, or disorder is severe or intractable pain, acute pain, postsurgical pain, back pain, or cancer pain.

69. The method according to claim 64, wherein said disease is selected from femur cancer pain; non-malignant chronic bone pain; rheumatoid arthritis; osteoarthritis; spinal stenosis; neuropathic low back pain; neuropathic low back pain; myofascial pain syndrome; fibromyalgia; temporomandibular joint pain; chronic visceral pain, including, abdominal; pancreatic; IBS pain; chronic headache pain; migraine; tension headache, including, cluster headaches; chronic neuropathic pain, including, post-herpetic neuralgia; diabetic neuropathy; HIV- associated neuropathy; trigeminal neuralgia; Charcot-Marie Tooth neuropathy; hereditary sensory neuropathies; peripheral nerve injury; painful neuromas; ectopic proximal and distal discharges; radiculopathy; chemotherapy induced neuropathic pain; radiotherapy-induced neuropathic pain; post-mastectomy pain; central pain; spinal cord injury pain; post-stroke pain; thalamic pain; complex regional pain syndrome; phanton pain; intractable pain; acute pain, acute post-operative pain; acute musculoskeletal pain; joint pain; mechanical low back pain; neck pain; tendonitis; injury/exercise pain; acute visceral pain, including, abdominal pain; pyelonephritis; appendicitis; cholecystitis; intestinal obstruction; hernias; etc; chest pain, including, cardiac Pain; pelvic pain, renal colic pain, acute obstetric pain, including, labor pain; cesarean section pain; acute inflammatory, burn and trauma pain; acute intermittent pain, including, endometriosis; acute herpes zoster pain; sickle cell anemia; acute pancreatitis; breakthrough pain; orofacial pain including sinusitis pain, dental pain; multiple sclerosis (MS) pain; pain in depression; leprosy pain; behcet's disease pain; adiposis dolorosa; phlebitic pain; Guillain-Barre pain; painful legs and moving toes; Haglund syndrome; erythromelalgia pain; Fabry's disease pain; bladder and urogenital disease, including, urinary incontinence; hyperactivity bladder; painful bladder syndrome; interstitial cystitis (IC); or prostatitis.