What is claimed is:

1. The compound *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification, wherein said *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered as a tablet at 600 mg twice a day in combination with trastuzumab, capecitabine, bevacizumab, paclitaxel or docetaxel.

2. The compound (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification, wherein said (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered as a tablet at 0.1 to 2000 mg a day in combination with trastuzumab, capecitabine, bevacizumab, paclitaxel or docetaxel.

3. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification.

4. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 3, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered as an oral dosage form.

5. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused

by ErbB2 over-expression or amplification according to Claims 3 or 4, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in a tablet.

6. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 3 to 5, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in combination with another therapeutic agent.

7. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 6, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

8. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 3 to 7, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used after previous treatment for brain cancer.

9. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused

by ErbB2 over-expression or amplification according to Claim 8, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

10. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 9, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

11. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 3 to 7, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used after previous treatment for breast cancer.

12. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 11, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

13. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-

dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 11, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

14. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 3 to 13, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

15. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 14, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine for use in the treatment of brain cancer is administered from about 550 to about 650 mg twice per day.

16. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 3 to 13, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

17. *N*4-(4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-

a]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to Claim 16, wherein the (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered from about 25 to about 1800 mg per day.

18. A method of treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification in a mammal comprising administering a therapeutically effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol to the mammal.

19. A method of treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification in a patient having brain cancer comprising administering a therapeutically effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol to the patient.

20. A method of treating or preventing local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification in a mammal in need of such treatment, wherein the method comprises administering to thed mammal a therapeutically effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol.

21. The method of any one of Claims 18 to 20, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered as an oral dosage form.

22. The method of any one of Claims 18 to 21, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-

yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in a tablet.

23. The method of any one of Claims 18 to 22, wherein the method includes another therapeutic agent.

24. The method of Claim 23, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

25. The method of any one of Claims 18 to 24, wherein the method is used after previous treatment for brain cancer.

26. The method of Claim 25, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

27. The method of Claim 25, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

28. The method of any one of Claims 18 to 24, wherein the method is used after previous treatment for breast cancer.

29. The method of Claim 28, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

30. The method of Claim 28, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

31. The method of any one of Claims 18 to 30, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered.

32. The method of Claim 31, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered.

33. The method of any one of Claims 18 to 30, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered.

34. The method of Claim 33, wherein about 25 to about 1800 mg per day of (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is administered.

35. The use of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in the manufacture of a medicament for the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification.

36. The use of Claim 35, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of an oral dosage form.

37. The use of Claims 35 or 36, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of a tablet.

38. The use of any one of Claims 35 to 37, wherein the use is in combination with another therapeutic agent.

39. The use of Claim 38, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

40. The use of any one of Claims 35 to 39, wherein the use is after previous treatment for brain cancer.

41. The use of Claim 40, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

42. The use of Claims 40, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal

radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

43. The use of any one of Claims 35 to 39, wherein the use is after previous treatment for breast cancer.

44. The use of Claim 43, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

45. The use of Claim 43, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

46. The use of any one of Claims 35 to 45, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

47. The use of Claim 46, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

48. The use of any one of Claims 35 to 45, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

49. The use of Claim 48, wherein about 25 to about 1800 mg per day of (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

50. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-

expression or amplification comprising administering *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine.

51. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 50, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered as an oral dosage form.

52. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claims 50 or 51, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is in a tablet.

53. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 50 to 52, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is in combination with another therapeutic agent.

54. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 54, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

55. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 50 to 54, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used after previous treatment for brain cancer.

56. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for

use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 55, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

57. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 55, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

58. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 50 to 54, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamineis used after previous treatment for breast cancer.

59. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 58, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

60. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification of Claim 58, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide,

docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

61. (2-((4-((4-([1,2,4]Triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol for use in the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification according to any one of Claims 50 to 60, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine for use in the treatment of brain cancer is administered from about 550 to about 650 mg twice per day.

62. A method of treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification in a mammal using (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol comprising administering an effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine to the mammal.

63. A method of treating local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification using (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in a patient having brain cancer comprising administering an effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine to the patient.

64. A method of treating or preventing local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification using (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in a mammal in need of such treatment, wherein the method comprises administering to the mammal an effective amount of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine.

65. The method of any one of Claims 62 to 64, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered as an oral dosage form.

66. The method of any one of Claims 62 to 65, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is in a tablet.

67. The method of any one of Claims 62 to 66, wherein the method includes another therapeutic agent.

68. The method of Claim 67, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

69. The method of any one of Claims 62 to 68, wherein the method is used after previous treatment for brain cancer.

70. The method of Claim 69, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

71. The method of Claim 69, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

72. The method of any one of Claims 62 to 68, wherein the method is used after previous treatment for breast cancer.

73. The method of Claim 72, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

74. The method of Claim 72, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

75. The method of any one of Claims 62 to 74, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is administered.

What is claimed is:

1. The use of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in the manufacture of a medicament for the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification.

2. The use of Claim 1, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of an oral dosage form.

3. The use of Claims 1 or 2, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of a tablet.

4. The use of any one of Claims 1 to 3, wherein the use is in combination with another therapeutic agent.

5. The use of Claim 4, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

6. The use of any one of Claims 1 to 5, wherein the use is after previous treatment for brain cancer.

7. The use of Claim 6, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

8. The use of Claims 6, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

9. The use of any one of Claims 1 to 5, wherein the use is after previous treatment for breast cancer.

10. The use of Claim 9, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

11. The use of Claim 9, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

12. The use of any one of Claims 1 to 11, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

13. The use of Claim 12, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

14. The use of any one of Claims 1 to 11, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

15. The use of Claim 14, wherein about 25 to about 1800 mg per day of (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.