

**ANTI-WART PHARMACEUTICAL COMPOSITION AND METHOD FOR
TREATING WART**

This application claims priority of Taiwanese application no. 100115591, filed on May 4, 2011.

5 This invention relates to an anti-wart pharmaceutical composition containing a water-soluble extract from a plant of *Solanum* genus and a method for treating wart using the water-soluble extract.

10 Wart is a kind of dermatosis that occurs on the skin or mucous of human or animal subjects, and is a benign epidermal hyperplasia caused by infection of Human papilloma virus (HPVs). In general, based on the site where the wart forms, wart can be divided into cutaneous wart and mucosal wart. The epidermal cells of skin or mucosa
15 affected by HPVs are subjected to benign proliferation and hyperkeratosis, thereby resulting in formation of papules or nodules with roughened surfaces on the skin or mucosa.

20 In clinical diagnosis, wart is divided into four types according to symptoms thereof: (1) common wart (*Verruca vulgaris*), scaly and roughened papule(s) or nodule(s) which usually occur on hands or fingers ; (2) flat wart (*Verruca plana juvenilis*), slightly raised and flat-topped papule(s) which usually occurs on faces, hands, and lower
25 legs; (3) plantar wart (*Verruca plantaris*), a thick and hyperkeratotic papule which usually occurs at pressure points of soles and toes and causes painful feeling for

a patient when the papule is pressed; and (4) genital wart (*Condyloma acuminatum*), a cauliflower-like lump formed on skin or mucosa of genitalia and anus, which is a highly infectious and sexually transmitted disease (STD).

5 Nowadays, warts are treated using the following remedies based on the type, size, number, and lesion site thereof:

 (1) Topical therapy, e.g., podophyllin (Trade name: Wartec®) and imiquimod (trade name: Aldara®) for treating
10 genital wart, 5-fluorouracil for treating genital wart and cutaneous wart, and bleomycin (Trade name: Bleocin®) for treating plantar wart and warts formed on palm;

 (2) Laser treatment, e.g., pulsed dye laser and carbon dioxide laser, especially for treating plantar wart and
15 warts formed on palm; and

 (3) Cryotherapy using liquid nitrogen and suitable for treating most warts, less effective for warts formed on palm, sole, and toe.

 Each of the aforesaid remedies is suitable only for
20 a specific type of wart, rather than all types of warts. Moreover, in clinical diagnosis, there are inferior effect problem and side effect drawback for the aforesaid remedies. For example, topical therapy used for a long period of time will cause recurrence of warts and treating effect
25 thereof is unsatisfactory. Laser treatment and cryotherapy will cause severe side effects, such as pain, ulceration, and scarring. Some patients, especially children, are

likely to refuse treatment due to the unendurable severe side effects.

To overcome the aforesaid drawbacks, researches have focused on finding active components from traditional Chinese medicines or herbs to develop a drug that is effective for treating warts and that has no undesired side effects.

It is known that *Solanum incanum* L. (also known as *Solanum incanum* Ruiz. & Pav., *Solanum undatum*, *Solanum coagulans* Forsskal in Latin, and bitter apple in English) contains steroidal glycoalkaloid which exhibits anti-cancer activity. In addition, many plants of the *Solanum* genus are reported to contain steroidal glycoalkaloid, including, for example, *Solanum indicum*, *Solanum nigrum*, also known as Long Kui in Chinese and black nightshade in English, *Solanum capsicastrum* (known as false Jerusalem cherry in English), *Solanum xanthocarpum*, *Solanum melongena*, *Solanum coagulans*, *Solanum tuberosum*, *Solanum sodomium*, *Solanum turburosum*, *Solanum aculeastrum*, *Solanum lycocarpum*, *Solanum khasianum*, *Solanum suaveolens*, *Solanum uporo*, *Solanum abutiloides*, *Solanum coccineum*, *Solanum unguiculatum*, *Solanum robustum*, *Solanum anguivi*, *Solanum platanifolium*, *Solanum mammosum*. The common steroidal glycoalkaloids contained in plants of *Solanum* genus include solamargine and solasonine.

US 7,078,063 B2 issued to the inventors of this invention discloses a water soluble extract from a plant

of *Solanum* genus, especially *Solanum incanum* L., and a process for preparing the same. The water soluble extract consists essentially of at least 60% of solamargine and solasonine.

5 In the aforesaid US patent, the inventors found that the water soluble extract exhibits inhibition of growth of tumor/cancer cells, in particular liver tumor cells, lung cancer cells, and breast cancer cells. In this invention, the inventors unexpectedly found that the water
10 soluble extract is effective in treating warts.

 Therefore, according to a first aspect, this invention provides an anti-wart pharmaceutical composition which includes a water-soluble extract from a plant of *Solanum* genus, the water-soluble extract comprising at least 60%
15 of solamargine and solasonine.

 In a second aspect, this invention provides a method for treating wart, which comprises applying to a subject in need of such treatment a water-soluble extract from a plant of *Solanum* genus, the water-soluble extract
20 comprising at least 60% of solamargine and solasonine.

 In a third aspect, this invention relates to the use of a water-soluble extract from a plant of *Solanum* genus in the manufacture of a medicament for treating wart, the water-soluble extract comprising at least 60% of
25 solamargine and solasonine.

The above and other objects, features and advantages of this invention will become apparent with reference to the following detailed description and the preferred embodiments taken in conjunction with the accompanying drawings, in which:

Fig. 1 shows photos illustrating that lesions (indicated by arrows) of test subject C are gradually ameliorated by treating the same with a *Solanum incanum* L. gel;

Fig. 2 shows photos illustrating that lesions (indicated by arrows) of test subject D are gradually ameliorated by treating the same with a *Solanum incanum* L. gel;

Fig. 3 shows photos illustrating that lesions (indicated by arrows) of test subject E are gradually ameliorated by treating the same with a *Solanum incanum* L. gel; and

Fig. 4 shows photos illustrating that lesions (indicated by arrows) of test subject F are gradually ameliorated by treating the same with a *Solanum incanum* L. gel.

For the purpose of this specification, it will be

clearly understood that the word "comprising" means "including but not limited to", and that the words "comprises," "contain" and variants thereof have a corresponding meaning.

5 Unless otherwise defined, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. One skilled in the art will recognize many methods and materials similar or equivalent to those
10 described herein, which could be used in the practice of this invention. Indeed, this invention is in no way limited to the methods and materials described. For clarity, the following definitions are used herein.

 The present invention provides an anti-wart
15 pharmaceutical composition, which includes a water-soluble extract from a plant of *Solanum* genus. The water-soluble extract comprises at least 60% of solamargine and solasonine. Preferably, the water-soluble extract comprises at least 60%-90% of solamargine and
20 solasonine.

 The process for preparing the water-soluble extract has been disclosed in US 7,078,063 B2 and includes the steps of:

 (a) subjecting a plant material of a plant of *Solanum*
25 genus to an extraction treatment using an acidic aqueous solution with a pH value of 3~5, such that an aqueous solution is obtained;

- (b) adjusting the pH value of the aqueous solution obtained in step (a) to pH 8~10 with a base, such that a precipitate is formed;
- (c) washing the precipitate formed in step (b) with water, followed by drying, such that a dried product is obtained;
- (d) admixing the dried product obtained in step (c) with chloroform, followed by addition of a suitable amount of a 100% alcohol, such that a chloroform-alcohol mixture is formed;
- (e) mixing the chloroform-alcohol mixture formed in step (d) with a water/alcohol solution having a predetermined water:alcohol ratio, such that a mixture containing a chloroform-based layer and a non-chloroform-based layer is obtained;
- (f) removing the chloroform-based layer from the mixture obtained in step (e), followed by addition of a suitable amount of water; and
- (g) obtaining a supernatant from the resultant mixture of step (f), followed by drying the supernatant, wherein the resultant dried product is able to be directly dissolved in water to form a yellowish clear and transparent aqueous solution.

Preferably, in step (a), the plant material of the plant of *Solanum* genus has been chopped in a preliminary treatment.

Preferably, in step (a), the plant material is at least

one of the fruit, root, stem, and leaf of the plant of *Solanum* genus. In a preferred embodiment of this invention, the plant material used in step (a) is the fruit of the plant of *Solanum* genus.

5 The inventors found that certain factors might affect the content and proportion of solasonine and solamargine in the water-soluble extract obtained using the aforesaid process. These factors include the species of the plant of *Solanum* genus and the part/parts of the plant used in
10 the extracting process, as well as the types of alcohol and base used. Therefore, a skilled artisan can prepare a desired water-soluble extract by selecting a suitable species of the plant of *Solanum* genus and using a suitable part or parts of the plant, in conjunction with appropriate
15 operating conditions.

 Preferably, the water-soluble extract is obtained from a plant of *Solanum* genus selected from the group consisting of *Solanum incanum* L., *Solanum indicum*, *Solanum nigrum*, *Solanum capsicastrum*, *Solanum xanthocarpum*,
20 *Solanum melongena*, *Solanum coagulans*, *Solanum tunigrum*, *Solanum sodomeum*, *Solanum turburosum*, *Solanum aculeastrum*, *Solanum lycocarpum*, *Solanum khasianum*, *Solanum suaveolens*, *Solanum uporo*, *Solanum abutiloides*, *Solanum coccineum*, *Solanum unguiculatum*, *Solanum robustum*, *Solanum anguivi*,
25 *Solanum platanifolium*, and *Solanum mammosum*. In a preferred embodiment of this invention, the water-soluble extract is obtained from *Solanum incanum* L..

In preliminary human tests done by the inventors, the results show that the water-soluble extract is effective in treating or alleviating warts including cutaneous warts and mucosal warts without undesired side effects, e.g.,
5 ulceration or erythema. Therefore, the present invention provides a method for treating wart, which comprises applying to a subject in need of such treatment the water-soluble extract.

The wart that can be treated by the anti-wart
10 pharmaceutical composition of this invention includes common wart, flat wart, plantar wart, and genital wart.

The anti-wart pharmaceutical composition of this invention can be administered via e.g., parenteral, transmucosal, transdermal, or topical route.

15 The parenteral route includes subcutaneous injection, intraepidermal injection, intradermal injection, and intralesional injection.

For the parenteral route, the anti-wart pharmaceutical composition of this invention can be
20 formulated into an injection product using techniques known to a skilled artisan, e.g., sterile aqueous solution or dispersion.

In this invention, the injection product can be prepared by mixing the water-soluble extract with a
25 pharmaceutically acceptable carrier that is widely employed in drug-manufacturing technology.

The pharmaceutically acceptable carrier comprises one

or more reagents, including, for example, water, saline, buffer solution (Phosphate Buffered Saline (PBS), Ringer's solution or Hank's solution), emulsifier, suspending agent, decomposer, pH adjusting agents, stabilizing agent, 5 chelating agent, preservative, diluents, absorption delaying agent, liposome. The choice and amount of these pharmaceutically acceptable carriers are within the expertise of those skilled in the art.

Moreover, the anti-wart pharmaceutical composition 10 of this invention can be administered by topical routes, e.g., dermal, buccal, genital, vaginal, anal, and rectal routes. In an embodiment of this invention, the anti-wart pharmaceutical composition of this invention is administered by dermal route. In another embodiment of 15 this invention, the anti-wart pharmaceutical composition of this invention is administered by genital route.

The pharmaceutical composition according to this invention can be formulated into a suitable dosage form for topical administration using technology well known 20 to those skilled in the art, which includes, external preparations, effervescent tablets, suppositories, and the like.

In a preferred embodiment of this invention, the anti-wart pharmaceutical composition is formulated into 25 an external preparation by admixing the water-soluble extract according to this invention with a base that is well known and commonly used in the art.

In this invention, the external preparation includes, emulsion, gel, ointment, cream, patch, liniment, powder, aerosol, spray, lotion, serum, paste, foam, drop, suspension, and tincture. In an embodiment of this invention, the anti-wart pharmaceutical composition is
5 in the form of gel.

For example, the base suitable for producing the external preparation according to this invention may include one or more of the following additives: water,
10 alcohols, glycol, hydrocarbons (e.g., petroleum jelly and white petrolatum), wax (e.g., paraffin and yellow wax), preserving agents, antioxidant, surfactants, absorption enhancers, stabilizing agents, gelling agents (e.g., carbopol®974P, microcrystalline cellulose, and
15 carboxymethylcellulose), active agents, humectants, odor absorbers, fragrances, pH adjusting agents, chelating agent, emulsifier, occlusive agents, emollients, thickeners, solubilizing agents, penetration enhancers, anti-irritants, colorants, and propellants. The choice
20 and amount of these additives are within the expertise of those skilled in the art.

This invention also provides a method for treating wart, which comprises applying to a subject in need of such treatment the water-soluble extract or the anti-wart
25 pharmaceutical composition.

The dosage and the frequency of administration of the anti-wart pharmaceutical composition according to this

invention may vary depending on the following factors:
the severity of the disease to be treated, the route of
administration, and the weight, age, physical condition
and response of the subject to be treated. For instance,
5 the daily dosage of the pharmaceutical composition
according to this invention may be 10 to 20 mg/cm² of the
lesion area and one to six times per day.

Example 1: Preparation of the water-soluble extract

10 The water-soluble extract was prepared based on the
procedure as disclosed in Example 1 of US 7,078,063 B2.
Specifically, 500 g of ripe fruit of *Solanum incanum* L.
was ground subsequent to addition of 1000 ml pure water.
To the resultant aqueous mixture, 99.5% of acetic acid
15 was added dropwise to adjust the pH value to 4.0, followed
by shaking at room temperature for 12 hrs. A supernatant
was obtained by centrifuging the aqueous mixture at 4500
rpm for 12 hours, and 33% NH₄OH basic solution was added
thereto dropwise to adjust the pH value of the supernatant
20 to 9.0, and a precipitate was formed. The precipitate was
obtained by conducting centrifugation at 4,500 rpm for
10 minutes(Beckman Coulter, Avanti J-25, JA-14 Rotor),

and the residual basic solution present therein was removed by washing the precipitate with water, followed by centrifugation at 4,500 rpm. The precipitate thus obtained was suspended in distilled water and subjected to
5 lyophilization (Virtis, Freezemobile 12ES) to obtain 5 g of dried powder.

2 g of the dried powder was dissolved in 50 ml chloroform in reagent grade, followed by addition of 40 ml of 100% methanol and shaking to form a uniform suspension. A
10 supernatant was obtained by centrifugation at 4,500 rpm. 70 ml of methanol:water solution (1:1) was added to the supernatant and mixed well. The mixture obtained was centrifuged at 12,000 rpm for 10 min. The resultant supernatant was taken out, and 120 ml distilled water was
15 added thereto and shaken well. The supernatant was further centrifuged at 12,000 rpm for 10 min so as to remove the precipitate. The resultant supernatant was subjected to concentration under reduced pressure at 55°C to remove methanol, followed by lyophilization to obtain dried
20 powder of the water-soluble extract.

Example 2: Therapeutic effect and safety of the water-soluble extract in treating human cutaneous wart and mucosal wart

25 In order to examine the therapeutic effect and safety of the water-soluble extract as prepared in the above Example 1 in treating human cutaneous wart and mucosal

wart, the following tests were conducted.

A. Preparation of gel containing the water-soluble extract

4 g of carbopol®974P, which was used as a gelling agent and is commercially available from Lubrizol Advanced
5 Materials, Inc., KY 40258, USA, was dissolved in 50 g pure water, followed by sequential addition of 30 g of propylene glycol and 7 g of the dried powder of the water-soluble extract obtained in Example 1 and mixing well. The mixture was heated in a heating vessel at a temperature of 50°C
10 to 60°C for 20 minutes, followed by cooling at room temperature. The cooled mixture was added with triethanolamine to adjust pH to 7.0 ± 0.5 . Subsequently, water was added until the total weight of the mixture reached 100 g, thereby obtaining a gel containing 7% (w/w) of the
15 water-soluble extract (hereinafter referred to as *Solanum incanum* L. gel).

B. Test subjects and clinical information thereof

Six test subjects (test subjects A to F) participating
20 in the following tests were enrolled from the outpatient department of National Cheng Kung University Hospital, Taiwan. The test subjects A and B suffered from mucosal wart, and the test subjects C to F suffered from cutaneous wart. The clinical information of the six test subjects,
25 including gender, age, wart type, and lesion location, are outlined in Table 1.

Table 1

Test subject	Age (y/o)	Gender	Wart type	Lesion location
A	32	Female	Mucosal wart	External genital
B	21	Female	Mucosal wart	External genital
C	28	Female	Cutaneous wart	Finger tip
D	21	Male	Cutaneous wart	Finger tip, finger
E	26	Male	Cutaneous wart	Finger
F	33	Male	Cutaneous wart	Finger tip, palm

C. Pre-clinical test

Prior to the start of the test (i.e., on Week 0),
 5 selected lesions on each of the enrolled test subjects
 were photographed. The *Solanum incanum* L. gel prepared
 in section A of Example 2 was applied to the selected lesions
 and normal skin areas that surround the selected lesions.
 Each of the normal skin areas had inner and outer peripheries
 10 and the distance between the inner and outer peripheries
 of each of the normal skin areas was not greater than 0.5
 cm. The treated dosage was 10~20 mg/cm², and twice a day.
 The selected lesions of each of the test subjects were
 photographed at predetermined evaluation times (i.e., at
 15 the ends of weeks 1, 2, 3, 4, 7, 8, 13, 20 and 29, see
 Table 2), and evaluations for therapeutic effect and safety
 of the *Solanum incanum* L. gel were conducted according
 to the procedures set forth in sections D and E below.
 The test was finished when the lesions of each of the test

subjects were completely cured and the outer appearance of the lesion skin was recovered to a normal state.

Table 2

Test subject	Evaluation time (week)									
	0 ^a	1 ^b	2	3	4	7	8	13	20	29
A	+ ^c	+	- ^d	-	-	-	-	-	-	-
B	+	+	-	-	-	-	-	-	-	-
C	+	-	-	-	+	-	+	-	-	-
D	+	-	+	+	-	-	+	-	-	-
E	+	-	+	-	-	+	-	-	-	-
F	+	-	-	-	-	+	-	⊙ ^e	※ ^f	+

^a: "0" indicates week 0, i.e., prior to application of the *Solanum incanum* L. gel

^b: "1" means week 1 after application of the *Solanum incanum* L. gel

^c: "+" means evaluation was conducted

^d: "-" means no evaluation was conducted

^e: "⊙" means evaluation was conducted only on the lesions at finger tip

^f: "※" means evaluation was conducted only on the lesion on palm

5

D. Evaluation of therapeutic effect

The photos of the selected lesions of the test subjects as obtained at Week 0 (i.e., prior to the application of the *Solanum incanum* L. gel) and as obtained at the evaluation times were compared by a dermatologist to determine the improvement of lesions treated with the *Solanum incanum* L. gel.

10

E. Evaluation of safety

Evaluation of safety was conducted by a researcher at the predetermined evaluation times. The test subjects were interrogated by the researcher in respect to the appearance of any adverse skin response (including ulceration, irritation and erythema).

Results:

A. Evaluation of the therapeutic effects of the *Solanum incanum* L. gel

Compared with the photos of the selected lesions of each of the test subjects A and B as obtained at week 0 and other predetermined evaluation times, the selected lesions of each of the test subjects A and B treated with the *Solanum incanum* L. gel were significantly improved after the end of week 1 (data not shown).

Figs. 1 to 4 respectively show photos of the selected lesions of the test subjects C to F as obtained at different evaluation times. The photos demonstrate that the lesions of each of the test subjects C to F treated with the *Solanum incanum* L. gel greatly improved with time. After the test was finished, the test subjects C to F were further subjected to the follow-up study for six months to two years to determine whether the lesion area that had already been cured by the *Solanum incanum* L. gel was recurrent. Based on the results of the follow-up study, no warts were found

to have recurred for all of the test subjects (data not shown).

The aforesaid results show that the *Solanum incanum* L. gel of this invention could be effectively used to treat cutaneous wart and mucosal wart.

B. Evaluation of the safety of the gel containing the water-soluble extract

In the safety test, only a few of the test subjects experienced temporary and mild adverse skin responses, including irritation or erythema, which, however, did not halt them from completing the test (data not shown).

In view of the aforesaid, the inventors contemplate that the water-soluble extract from a plant of *Solanum* genus according to this invention can be developed into an anti-wart drug for long-term use in treating warts without adverse side effects.

CLAIMS:

1. Use of a water-soluble extract from a plant of *Solanum* genus in the preparation of a pharmaceutical composition for treating wart, wherein the water-soluble
5 extract comprises at least 60% of solamargine and solasonine.

2. The use according to claim 1, wherein the water-soluble extract comprises at least 60%-90% of solamargine and solasonine.

10 3. The use according to claim 1, wherein the wart comprises common wart, flat wart, plantar wart, or genital wart.

4. The use according to claim 1, wherein the pharmaceutical composition further comprises a base
15 selected from the group consisting of water, alcohols, glycol, hydrocarbons, wax, preserving agents, antioxidant, surfactants, absorption enhancers, stabilizing agents, gelling agents, active agents, humectants, odor absorbers, fragrances, pH adjusting agents, chelating agent,
20 emulsifier, occlusive agents, emollients, thickeners, solubilizing agents, penetration enhancers, anti-irritants, colorants, propellants, and combinations thereof.

5. The use according to claim 4, wherein the base
25 includes a gelling agent and glycol.