

Description

The present invention relates to dithioketene derivatives which possess good therapeutic and prophylactic activity against hepatic diseases, processes for the production of said compounds and pharmaceutical compositions containing them.

Dialkyl 2-(1,3-dithietan-2-ylidene)malonates and processes for the preparation thereof are disclosed in DE-A-2,625,220, 2,316,921 and Japanese Specification No. 74 39,260.

Dialkyl 2-(1,3-dithiolan-2-ylidene)malonates are disclosed in Japanese Kokai 75 100,062, Japanese 75 39,666 and 74 39,260.

Dithiolane derivatives, having a ketone functional group in the side chain, are disclosed in Japanese Kokai Tokyo Koho JP 59 27,887.


Dithietane derivatives containing two ketone functional groups in the side chain are disclosed in Japan Kokai Tokyo Koho JP 62 158,274 and Eur. Pat. Appl. EP 234,480.

Reference is also made to USSN 206,659, which was filed by the same applicant of the present invention.

It is an object of the present invention to provide novel dithioketene derivatives which possess good therapeutic and prophylactic activity against hepatic disorders.

Another object of the present invention is to provide pharmaceutical compositions containing said compounds for example at a dose suitable for daily administration.

The present invention accordingly provides compounds of the general formula (I)



in which R¹ is a straight or branched alkyl group having 1 to 5 carbon atoms; R² is hydrogen, a straight or branched alkyl group having 1 to 5 carbon atoms, benzyl, phenyl, 4-hydroxyphenyl, halogenophenyl, 4-trifluoromethylphenyl, 2-thiazolyl, 4-methyl-2-thiazolyl, 4-phenyl-2-thiazolyl, 2-pyridyl or 1,3,4-thiadiazol-2-yl; and n is an integer of 1 or 2 but excluding compounds in which

- (i) n is 2, R¹ is ethyl and R² is benzyl;
(ii) n is 2, R¹ is isopropyl and R² is benzyl;
(iii) n is 2, R¹ is ethyl and R² is chlorophenol or bromophenol.

In the following, "dithietan-" refers to a compound of formula (I) where n=1, and "dithiolan-" refers to a compound of formula (I) where n=2.

Two particularly preferred classes of formula I are those where R¹ is methyl, ethyl, isopropyl or sec-butyl and n is 1 or 2. In one class R² is hydrogen, a straight or branched alkyl group having 1 to 5 carbon atoms, and in the other class R² is phenyl, a substituted phenyl or a heteroaromatic ring, but excluding compounds in which

- (i) n is 2, R^1 is ethyl and R^2 is benzyl;
(ii) n is 2, R^1 is isopropyl and R^2 is benzyl;
(iii) n is 2, R^1 is ethyl and R^2 is chlorophenol or bromophenol.


The compounds of the invention can also be used in the form of their pharmaceutically acceptable salts and hydrates.

The compounds according to the invention can be administered orally. They will, in general, be associated with a pharmaceutically acceptable carrier or diluent to provide a pharmaceutical dosage form.

The pharmaceutical composition can most conveniently be in the form of capsules or tablets, which may be in the form of slow releasing formulations. The composition may also take the form of a dragee or syrup.

The convenient daily dose would be of the order of 100mg to 1.0g per day and the form of dosage units contains from 50mg to 200mg.

The compounds of the present invention may be prepared from the reaction of an acid anhydride of formula (II) and an amine of formula (III):




(II)

(III)

15 in which R^1 , R^2 and n have the same meanings given previously and R^3 is tert-butyl, lower alkoxy or benzyloxy.

The reaction may be carried out by allowing the acid anhydride (II) and the amine (III) to stand in a solvent such as methylene chloride, acetonitrile or alcohol. The reaction with a weakly basic heteroaromatic amine (e.g. 2-aminopyridine, 2-aminothiazole, or 2-amino-1,3,4-thiadiazole) can be carried out by heating the reaction mixture at 50 - 100°C in a solvent such as acetonitrile or N,N-dimethylformamide.

20 In an alternative procedure, the compounds of the present invention can be prepared by reacting an active ester of formula (IV) with an amine of formula (III);




(IV)

35 wherein R^1 and n are the same as defined previously and R^4 is 1-benzotriazolyl, N-succinimidoyl or N-phthalimidoyl.

The reaction can be carried out in a solvent such as acetonitrile, methylene chloride or N,N-dimethylformamide at 10 - 100°C.

The reaction with alkyl amines or aniline derivatives can readily take place at a lower temperature (10-30°C), whereas the reaction with weakly basic heteroaromatic amines requires a high temperature (50 - 100°C).


40 In another process, the compounds of formula (I) can be prepared by reacting a monoacid of formula (V) with phosphorous pentachloride to yield an acid chloride of the monoacid which is very unstable. An amine of formula (III) is then added to the acid chloride of monoacid in the presence of pyridine or triethylamine,



(V)

55 wherein R^1 and n are the same as defined previously. This reaction can be conducted in the presence of a solvent such as methylene chloride or acetonitrile at a temperature from 0°C to -15°C.

In another alternative process, the compounds of formula (I) can be prepared by reacting the compounds of formula (VI) with the compounds of formula (VII).




(VI)



(VII)

wherein R^1 , R^2 and n are the same as defined previously; and X is a halogen.

The compound of formula (VI) can be obtained by the reaction of carbon disulfide with the compound of formula (VII) in the presence of sodium hydroxide at 15-50° C in an organic solvent, e.g. N,N-dimethylformamide or acetone,



(VIII)


wherein R^1 and R^2 are the same as defined previously.

In order to obtain the objective compounds of formula (I), the acid chloride of formula (II) and the active ester of formula (IV) are used as an intermediate in the present invention. These intermediates are novel ones, and also constitute the present invention. These intermediates may be made by the processes described below.

Acid anhydrides of formula (II), isolated in a stable form, may be prepared by reacting the monoacid of formula (V) with the acid chloride of R^3COOH , wherein R^3 is the same as defined previously, in the presence of an organic base, e.g. triethyl amine or pyridine, at 0 - 30°C.

The active ester of the formula (IV) can be prepared by reacting the monoacid of formula (V) with 1-hydroxybenzotriazole, N-hydroxysuccinimide or N-hydroxyphthalimide in the presence of a dehydrating agent such as N, N'-dicyclohexylcarbodiimide.

The monoacid of formula (V) can be prepared by hydrolyzing one ester group of formula (IX) using the corresponding alcohol as a solvent at ambient temperature;



(IX)

wherein R^1 and n are the same as defined previously.

The starting material of formula (VIII) may be prepared by the reaction of alkyl malonyl chloride with the corresponding amine of formula (III).

The compounds of formula (IX) can be prepared by similar methods described in the aforementioned literatures.

In order to make the present invention more fully understood, the following Experiments and Examples are given.

The descriptions of experiments explain the protective and therapeutic effects of the compounds.

Examples are provided to illustrate the preparation of the compounds of formula (I), the starting materials and the related intermediates (Examples 1 to 12).

Example 1 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate

Diisopropyl 2-(1,3-dithietan-2-ylidene)malonate (10g) was added to a solution of potassium hydroxide (3.0g) in isopropyl alcohol(300ml). The mixture was stirred at room temperature for 12 hrs, after which water(500ml) was added to dissolve the solid. The aqueous solution was acidified with acetic acid and the solid formed was filtered. The filter cake was purified with methylene chloride to afford the titled compound as a white solid. (6.6g, 79%)

m. p. : 123 - 124°C

¹H-NMR(CDCl₃) δ : 12.30(bs,1H), 5.11(m,1H), 4.00(s,2H), 1.27(d,6H)

IR (KBr) cm⁻¹ : 1712, 1627

Example 2 ; Ethyl 2-(1,3-dithietan-2-ylidene)-2-carboxy-acetate

Diethyl 2-(1,3-dithietan-2-ylidene)malonate was treated with a solution of potassium hydroxide in ethanol. The resultant solid was isolated and purified by the same method as in Example 1 to afford the titled compound as a white solid. (51%)

m. p. : 166 - 167°C

¹H-NMR(DMSO-d₆) δ : 4.21(q,2H), 4.14(s,2H), 1.08(t,3H)

IR (KBr) cm⁻¹ : 1701, 1637

Example 3 ; Sec-butyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate

Di-sec-butyl 2-(1,3-dithietan-2-ylidene)malonate was treated with a solution of potassium hydroxide in sec-butanol. The resultant solid was isolated and purified by the same method as in Example 1 to afford the titled compound as a white solid. (41%)

m. p. : 80 - 82° C

¹H-NMR(CDCl₃)δ : 12.28(bs,1H), 5.04(m,1H), 4.04(s,2H), 1.77(m,2H), 1.28(d,3H), 0.91(t,3H)

IR (KBr) cm⁻¹ : 1712, 1627

Example 4 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-carboxyacetate

Diisopropyl 2-(1,3-dithiolan-2-ylidene)malonate was treated with a solution of potassium hydroxide in isopropyl alcohol. The resultant solid was isolated and purified by the same method as in Example 1 to afford the titled compound as a white solid. (64%)

m. p. : 104 - 106°C

¹H-NMR(CDCl₃) δ : 13.25(bs,1H), 5.22(m,1H), 3.39(s,4H), 1.11(d,6H)

IR (KBr) cm⁻¹ : 1711, 1606

Example 5 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate

Triethylamine (5.18g) was added to a solution of isopropyl 2-(1,3-dithietan-2-ylidene) - 2 - carboxyacetate (10g) in methylene chloride (100ml). The mixture was cooled to 0°C, to which ethyl chloroformate (5.60g) was added dropwise and then stirred for an hour. The precipitate was removed by filtration. The filtrate was washed with water, dried and evaporated to afford the titled compound as a pale yellow oil. (10g, 81%).

¹H-NMR(CDCl₃) δ : 5.06(m,1H), 4.25(q,2H), 4.05(s,2H), 1.20(t,3H), 1.15(d,6H)

IR (KBr) cm⁻¹ : 1783, 1706, 1654

Example 6 ; Ethyl 2-(1,3-dithietan-2-ylidene)-2-ethoxycarboxycarbonyl)acetate

Following the procedure of Example 5, the titled compound was obtained as a white solid (99%) from ethyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate.

m. p. : 38 - 39°C

¹H-NMR(CDCl₃) δ : 4.25(2q's,4H), 4.05(s,2H), 1.32(2t's,6H)
 IR (KBr) cm⁻¹ : 1767, 1707

Example 7 ; Sec-butyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate

Following the procedure of Example 5, the titled compound was obtained as a yellow oil (97%) from sec-butyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate.

¹H-NMR(CDCl₃) δ : 5.34(m,1H), 4.32(q,2H), 4.08(s,2H), 1.72(m,2H), 1.45(t,3H), 1.34(d,3H), 0.94(t,3H)
 IR (KBr) cm⁻¹ : 1783, 1705, 1652

Example 8 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate

Following the procedure of Example 5, the titled compound was obtained as a yellow oil (97%) from isopropyl 2-(1,3-dithiolan-2-ylidene)-2-carboxyacetate.

¹H-NMR(CDCl₃) δ : 5.13(m,1H), 4.26(q,2H), 3.42(s,4H), 1.72(t,3H), 1.23(d,6H)
 IR (KBr) cm⁻¹ : 1786, 1691

Example 9 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl] acetate

N, N'-Dicyclohexylcarbodiimide (7.1g) was added to a mixture of isopropyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate (10g) and 1-hydroxybenzotriazole(4.61g) in methylene chloride(100ml). The mixture was stirred at room temperature for 4 hrs, after which the precipitate was removed by filtration. The filtrate was concentrated to afford the titled compound as a white solid. (15g, 96%)

m. p. : 146 - 148°C

¹H-NMR(CDCl₃) δ : 8.27-7.03(m,4H), 5.18(m,1H), 4.03(s,2H), 1.27(d,6H)
 IR (KBr) cm⁻¹ : 1709, 1655

Example 10 ; Ethyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl) oxycarbonyl]acetate

Following the procedure of Example 9, the titled compound was obtained as a white solid (89%) from ethyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate.

m. p. : 168 - 169°C

¹H-NMR(CDCl₃) δ : 8.27-7.06(m,4H), 4.43(q,2H), 4.11(s,2H), 1.37(t,3H)
 IR (KBr) cm⁻¹ : 1728, 1693

Example 11 ; Sec-butyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl) oxocarbonyl]acetate

Following the procedure of Example 9, the titled compound was obtained as a white solid (93%) from sec-butyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate.

m. p. : 143 - 144°C

¹H-NMR(CDCl₃) δ : 8.29-7.27(m,4H), 5.02(m,1H), 4.09(s,2H), 1.70(m,2H), 1.31(d,3H), 0.92(t,3H)
 IR (KBr) cm⁻¹ : 1712, 1663

Example 12 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl] acetate

Following the procedure of Example 9, the titled compound was obtained as a white solid (97%) from isopropyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate.

m. p. : 88 - 89°C

¹H-NMR(CDCl₃) δ : 8.28-7.15(m,4H), 5.24(m,1H), 2.31(s,4H), 1.38(d,6H)
 IR (KBr) cm⁻¹ : 1720, 1687

Example 13 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-carbamoylacetate

A solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate (10g) in methanolic ammonia (60ml) was stirred at room temperature for 4 hrs. The mixture was concentrated and purified with n-hexane and ethyl ether to give the titled compound as a white solid.(5.2g, 68%)

m. p. : 128 - 129°C

¹H-NMR(DMSO-d₆) δ : 7.68(bs,1H), 7.30(bs,1H), 5.13(m,1H), 4.03(s,2H), 1.37(d,6H)

IR (KBr) cm^{-1} : 1675, 1647

Example 14 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-(N-ethylcarbamoyl)acetate.

Following the procedure of Example 13, the titled compound was obtained as a white solid (72%) from a solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate in methanolic ethylamine.

m. p. : 65 - 67°C

$^1\text{H-NMR}(\text{DMSO-}d_6) \delta$: 8.42(bs, 1H), 5.25(m, 1H), 3.97(s, 2H), 3.37(q, 2H), 1.29(d, 6H), 1.08(t, 3H)

IR (KBr) cm^{-1} : 1683, 1613

Example 15 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-(N-benzylcarbamoyl)acetate

The mixture of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate (15g) and benzylamine (6.4ml) in methylene chloride (150ml) was stirred at room temperature for 4 hrs. The reaction mixture was then washed with dil-HCl and water to remove benzylamine. The solvent was evaporated and the residue was purified with ethyl acetate to give the titled compound as a white solid. (12.2g, 77%)

m. p. : 93 - 94°C

$^1\text{H-NMR}(\text{CDCl}_3) \delta$: 8.82(bs, 1H), 7.52(m, 5H), 5.06(m, 1H), 4.48(dd, 2H), 3.96(s, 2H), 1.28(d, 6H)

IR (KBr) cm^{-1} : 1669, 1615

Example 16 ; Methyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-fluorophenyl)carbamoyl]acetate

A mixture of methyl 2-[N-(4-fluorophenyl)carbamoyl]acetate (10g), carbon disulfide (3.2ml) and acetone (200ml) was cooled to 5 - 10°C, to which 50% aqueous sodium hydroxide (4.2ml) was added dropwise. The mixture was stirred at same temperature for 2 hrs, then dibromomethane (4ml) was added dropwise and the reaction mixture was stirred at room temperature for 1 hr and refluxed for 2 hrs. The solid was removed by filtration and the filtrate was poured onto ice-water. The solid formed was filtered, and dissolved in methylene chloride. The solution was washed with 10% aqueous sodium hydroxide solution and water, then dried. The solvent was evaporated and the residue was purified with ethyl acetate and ethyl ether to give the titled compound as a white solid. (8.5g, 60%)

m. p. : 141 - 142°C

$^1\text{H-NMR}(\text{CF}_3\text{COOD}) \delta$: 7.42-7.03(m, 4H), 4.15(s, 2H), 3.98(s, 3H)

IR (KBr) cm^{-1} : 1685, 1622

Example 17 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-chlorophenyl)carbamoyl]acetate

A solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-carboxyacetate (10g) in methylene chloride (100ml) was cooled to -10°C and thereto phosphorous pentachloride (8.9g) was added portionwise and then stirred for 1 hr. To the reaction mixture pyridine (13.8ml) and 4-chloroaniline (5.45g) were added and stirred for 2 hrs. The mixture was washed with dil-HCl, 10% aqueous sodium hydroxide solution and water and then evaporated. The residue was purified with ethyl acetate to afford the titled compound as a white solid. (8.9g, 61%).

m. p. : 164 - 165°C

$^1\text{H-NMR}(\text{CF}_3\text{COOD}) \delta$: 7.37(m, 4H), 5.21(m, 1H), 4.12(s, 2H), 1.45(d, 6H)

IR (KBr) cm^{-1} : 1678

Example 18 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-(N-phenylcarbamoyl)acetate

A solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate (10g) and aniline (3.6ml) in methylene chloride (100ml) was stirred at room temperature for 8 hrs. The reaction mixture was washed with 10% aqueous sodium hydroxide solution, dil-HCl and water and then evaporated to dryness. The residue solid was purified with ethyl acetate to give the titled compound as a white solid. (7.9g, 78%).

m. p. : 115 - 116°C

$^1\text{H-NMR}(\text{CDCl}_3) \delta$: 10.52(bs, 1H), 7.85-6.75(m, 5H), 5.16(m, 1H), 3.99(s, 2H), 1.34(s, 6H)

IR (KBr) cm^{-1} : 1671

Example 19 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-trifluoromethylphenyl)carbamoyl]acetate

Following the procedure of Example 18, the titled compound was prepared as a white solid (67%) from

isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 4-aminobenzotrifluoride.

m. p. : 151 - 152°C

¹H-NMR(CDCl₃) δ : 10.78(bs,1H), 7.68(m,4H), 5.40(m,1H), 4.02(s,2H), 1.34(d,6H)

IR (KBr) cm⁻¹ : 1686, 1623

Example 20 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-hydroxyphenyl)carbamoyl]acetate

Following the procedure of Example 18, the titled compound was prepared as a white solid (66%) from isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and p-aminophenol.

m. p. : 175 - 176°C

¹H-NMR(DMSO-d₆) δ : 10.20(s,1H), 9.15(s,1H), 7.40-6.73(dd,4H), 5.13(m,1H), 4.21(s,2H), 1.42(d,6H)

IR (KBr) cm⁻¹ : 1675, 1606

Example 21 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-[N-(4-trifluoromethylphenyl)carbamoyl]acetate

Following the procedure of Example 18, the titled compound was obtained as a white solid (80%) from isopropyl 2-(1,3-dithiolan-2-ylidene)-2-ethoxycarboxycarbonyl)acetate and 4-aminobenzotrifluoride.

m. p. : 139 - 140°C

¹H-NMR(CDCl₃) δ : 10.90(bs,1H), 7.82-7.65(dd,4H), 5.24(m,1H), 3.40(s,4H), 1.42(d,2H)

IR (KBr) cm⁻¹ : 1679, 1631

Example 22 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(2-thiazolyl)carbamoyl]acetate

The solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate (10g) and 2-aminothiazole(4.0g) in methylene chloride (100ml) was stirred at room temperature for 24 hrs. The reaction mixture was washed with 10% aqueous sodium hydroxide solution and water and then evaporated. The residue was chromatographed on silica gel column using methylene chloride as an eluent to afford the titled compound as a white solid (6.5g, 63%).

m. p. : 174 - 175°C

¹H-NMR(CDCl₃ / DMSO-d₆) δ : 11.70(bs,1H), 7.24(dd,2H), 5.29(m,1H), 4.19(s,2H), 1.29(d,6H)

IR (KBr) cm⁻¹ : 1675, 1621

Example 23 ; Ethyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-methyl-2-thiazolyl)carbamoyl]acetate

Following the procedure of Example 22, the titled compound was obtained as a white solid (39%) from ethyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 2-amino-4-methylthiazole.

m. p. : 218 - 219°C

¹H-NMR(CDCl₃) δ : 11.50(bs,1H), 6.45(s,1H), 4.39(q,2H), 4.02(s,2H), 2.34(s,3H), 1.33(t,3H)

IR (KBr) cm⁻¹ : 1685

Example 24 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-methyl-2-thiazolyl)carbamoyl]acetate

Following the procedure of Example 22, the titled compound was obtained as a white solid (46%) from isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 2-amino-4-methylthiazole.

m. p. : 197 - 198°C

¹H-NMR(CDCl₃) δ : 11.62(bs,1H), 6.44(2,1H), 5.16(m,1H), 4.06(s,2H), 2.35(s,3H), 1.43(d,6H)

IR (KBr) cm⁻¹ : 1678, 1627

Example 25 ; Sec-butyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-methyl-2-thiazolyl)carbamoyl]acetate

Following the procedure of Example 22, the titled compound was prepared as a white solid (67%) from sec-butyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 2-amino-4-methylthiazole.

m. p. : 153 - 154°C

¹H-NMR(DMSO-d₆) δ : 11.67(bs,1H), 6.47(s,1H), 5.01(m,1H), 4.07(s,2H), 2.35(s,3H), 1.49(m,2H), 1.29(d,3H), 0.93(t,3H)

IR (KBr) cm⁻¹ : 1675, 1624

Example 26 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-[N-(4-methyl-2-thiazolyl)carbamoyl]acetate

Following the procedure of Example 22, the titled compound was prepared as a white solid (41%) from isopropyl 2-(1,3-dithiolan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 2-amino-4-methylthiazole.

5 m. p. : 161 - 162°C

¹H-NMR(CDCl₃) δ : 11.85(bs,1H), 6.64(s,1H), 5.24(m,1H), 3.37(s,4H), 2.34(s,3H), 1.42(d,6H)

IR (KBr) cm⁻¹ : 1668, 1623

Example 27 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(4-phenyl-2-thiazolyl)carbamoyl]acetate

10

A solution of isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate (10g) and 2-amino-4-phenylthiazole (6.2g) in N,N-dimethylformamide (100ml) was stirred at 60°C for 24 hrs. The reaction mixture was poured onto ice-water (500ml) and the solid formed was filtered. The filter cake was dissolved in methylene chloride, and the solution was washed with 10% aqueous sodium hydroxide solution and water and then evaporated to dryness. The residue was chromatographed on silica gel column using methylene chloride as an eluent to afford the titled compound as a white solid. (6.9g, 54%).

15

m. p. : 225 - 226°C

¹H-NMR(CF₃ COOD) δ : 7.99-7.44(m,5H), 7.26(s,1H), 5.25(m,1H), 4.30(s,2H), 1.44(d,6H)

IR (KBr) cm⁻¹ : 1673

20

Example 28 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(1,3,4-thiadiazol-2-yl)carbamoyl]acetate

Following the procedure of Example 27, the titled compound (38%) was obtained from isopropyl 2-(1,3-dithietan-2-ylidene)-2-(ethoxycarboxycarbonyl)acetate and 2-amino-1,3,4-thiadiazole.

25

m. p. : 212 - 213°C

¹H-NMR(DMSO-d₆) δ : 9.16(s,1H), 5.20(m,1H), 4.26(s,2H), 1.33(d,6H)

IR (KBr) cm⁻¹ : 1676, 1625

Example 29 ; Isopropyl 2-(1,3-dithietan-2-ylidene)-2-[N-(2-pyridyl)carbamoyl]acetate

30

Following the procedure of Example 27, the titled compound (77%) was obtained from isopropyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl]acetate and 2-aminopyridine.

m. p. : 162 - 163°C

¹H-NMR(CDCl₃) δ : 11.10(s,1H), 8.72-6.93(m,4H), 5.21(m,1H), 3.99(s,2H), 1.37(d,6H)

35

IR (KBr) cm⁻¹ : 1676, 1631

Example 30 ; Ethyl 2-(1,3-dithietan-2-ylidene)-2-[N-(2-pyridyl)carbamoyl]acetate

Following the procedure of Example 27, the titled compound (51%) was prepared from ethyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl]acetate and 2-aminopyridine.

40

m. p. : 181 - 183°C

¹H-NMR(CDCl₃) δ : 10.97(bs,1H), 8.52-6.72(m,4H), 4.40(q,2H), 4.01(s,2H), 1.34(t,3H)

IR (KBr) cm⁻¹ : 1676, 1628

45

Example 31 ; Sec-butyl 2-(1,3-dithietan-2-ylidene)-2-[N-(2-pyridyl)carbamoyl]acetate

Following the procedure of Example 27, the titled compound (76%) was obtained from sec-butyl 2-(1,3-dithietan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl]acetate and 2-aminopyridine.

m. p. : 118 - 119°C

50

¹H-NMR(CDCl₃) δ : 11.03(bs,1H), 8.53-6.74(m,4H), 5.02(m,1H), 4.05(s,2H), 1.51(m,2H), 1.31(d,3H), 0.95(t,3H)

IR (KBr) cm⁻¹ : 1671, 1632

Example 32 ; Isopropyl 2-(1,3-dithiolan-2-ylidene)-2-[N-(2-pyridyl)carbamoyl]acetate

55

Following the procedure of Example 27, the titled compound was obtained as a white solid (53%) from isopropyl 2-(1,3-dithiolan-2-ylidene)-2-[(1-benzotriazolyl)oxycarbonyl]acetate and 2-aminopyridine.

m. p. : 121 - 122°C

¹H-NMR(DMSO-d₆) δ : 10.61(bs, 1H), 8.59-6.79(m,4H), 5.02(m, 1H), 3.37(bs,4H), 1.22(d,6H)
 IR (KBr) cm⁻¹ : 1666, 1629

Experiment 1 ; Protective effect against acute hepatic damage induced by carbon tetrachloride

(Principle)

Carbon tetrachloride (CCl₄) is a well-known hepatotoxic agent, thus widely used to produce experimental animal models for screening the potential drugs acting upon hepatic diseases. In this experiment test compounds were administered to mice prior to treatment with CCl₄. 24 Hours after the CCl₄ treatment, the animals were sacrificed and serum alanine aminotransferase (ALT) levels were determined. The hepatoprotective effect of a test compound was evaluated by the suppressive action against the increase of serum ALT induced by CCl₄.

(Method)

The experimental animals were divided into the normal, the CCl₄ treated and the test compound treated groups. Each group consisted of 8 mice with body weight of 20-25 grams. The test compounds were suspended in 0.2% sodium carboxymethylcellulose (CMC-Na) solution and administered orally at the dose of 50mg/125ml/kg body weight. Only the vehicle, 0.2% CMC-Na solution, was administered to the normal and to the CCl₄ treated group instead of the test compound suspension. 6 Hours after the drug administration, the CCl₄ solution in olive oil was administered orally to the CCl₄ treated-group and the test compound treated group at the dose of 50ul/25ml olive oil/kg body weight. The normal group was administered with olive oil only at the same dose. 24 Hours after the CCl₄ administration, blood samples were collected from the orbital sinus of the animals and the sera were monitored using an automatic blood analyzer (Gilford, SBA 300). The hepatoprotective effect of the test compounds was expressed by the suppressive percentage against the increase of serum ALT level induced by CCl₄ calculated by the following formula ;

$$\text{Hepatoprotective effect (\%)} = \frac{\text{ALT level (CCl}_4\text{ group)} - \text{ALT level (test group)}}{\text{ALT level (CCl}_4\text{ group)} - \text{ALT level (normal group)}} \times 100$$

(Results)

The test results are shown in Table I.

Table I (Protective effect of test compound against increase of serum ALT activity induced by CCl₄)

Compound	Inhibitory(%)	Compound	Inhibitory(%)	Compound	Inhibitory(%)
Example 13	28	Example 21	98	Example 27	11
Example 15	97	Example 22	100	Example 23	100
Example 18	100	Example 24	100	Example 29	100
Example 19	100	Example 26	98	Example 32	85

Experiment 2 ; Determination of 50% Effective Dose (ED₅₀) of test compounds**(Principle)**

5 The various doses of test compounds were administered to mice, followed by the treatment with CCl₄ after 6 hours. 24 Hours after the CCl₄ treatment, the serum ALT level of the animals were determined. The hepatoprotective effect of the test compounds and the doses were plotted to form dose-response curves, from which the ED₅₀ doses of the test compounds were estimated.

(Method)

10 The experimental animals were divided into the normal, the CCl₄ treated and the test compound treated groups. Each group consisted of 8 mice with body weight of 20 - 25 grams. The test compounds were suspended in 0.2% CMC-Na solution and administered orally at a dose of 50mg, 25mg, 12.5mg or
15 6.25mg/125ml/kg body weight. Only the vehicle, 0.2% CMC-Na solution, was administered to the normal and to the CCl₄ treated group instead of the test compound suspension. 6 Hours later, the CCl₄ solution in olive oil was administered orally to the CCl₄ treated and the test compound treated groups at the dose of 50ul/25ml olive oil/kg body weight. The normal group was administered olive oil only at the same dose. 24 Hours after
20 the CCl₄ administration, blood samples were collected from the orbital sinuses of the animals and the sera were obtained by centrifugation. The serum ALT activities were monitored and the hepatoprotective effect of the test compounds were calculated by the same method as described in experiment 1. The percentages were plotted against the corresponding doses of the test compounds to make dose-response curves, from which the 50% effective dose of test compound was estimated.

(Results)

The 50% effective doses (ED₅₀) of the test compounds are shown in Table II.

30

Table II (ED₅₀ of test compound)

35

40

Compound	ED ₅₀ (mg/kg)	Compound	ED ₅₀ (mg/kg)	Compound	ED ₅₀ (mg/kg)
Example 13	> 50	Example 21	< 50	Example 27	> 50
Example 15	10	Example 22	< 6.25	Example 28	< 6.25
Example 18	7	Example 24	< 6.25	Example 29	10
Example 19	6.25	Example 26	< 50	Example 32	< 50

45

Experiment 3 : Acute toxicity of test compounds

Each test compound was suspended in 0.2% CMC-Na solution in various concentrations to make test suspensions. The test suspensions were administered orally to male ICR mice (10 in each group) at various doses.
50 The number of dead mice was counted for 14 days and the value of median lethal dose (LD₅₀, g/kg) was calculated by the Hitchfield-Wilcoxon method.

The results are shown in Table III.


55

Table III (LD₅₀ of Test Compound)

Compound	LD ₅₀ (mg/kg)	Compound	LD ₅₀ (mg/kg)	Compound	LD ₅₀ (mg/kg)
Example 13	< 5,000	Example 22	> 5,000	Example 24	> 5,000
Example 28	> 5,000	Example 29	> 5,000		

Claims

1. Compounds of formula (I);



in which R¹ is a straight or branched alkyl group having 1 to 5 carbon atoms; R² is hydrogen, a straight or branched alkyl group having 1 to 5 carbon atoms, benzyl, phenyl, 4-hydroxyphenyl, halogenophenyl, 4-trifluoromethylphenyl, 2-thiazolyl, 4-methyl-2-thiazolyl, 4-phenyl-2-thiazolyl, 2-pyridyl or 1,3,4-thiadiazol-2-yl; and n is an integer of 1 or 2 but excluding compounds in which


- (i) n is 2, R¹ is ethyl and R² is benzyl;
 - (ii) n is 2, R¹ is isopropyl and R² is benzyl;
 - (iii) n is 2, R¹ is ethyl and R² is chlorophenyl or bromophenyl.
2. Compounds as claimed in claim 1, characterized in that R¹ represents methyl, ethyl, isopropyl, or sec-butyl.
 3. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents methyl, ethyl, isopropyl, or sec-butyl.
 4. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents benzyl.
 5. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents phenyl, 4-hydroxyphenyl, fluorophenyl, chlorophenyl or 4-trifluoromethylphenyl.
 6. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents 2-thiazolyl, 4-methyl-2-thiazolyl or 4-phenyl-2-thiazolyl.
 7. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents 1,3,4-thiadiazol-2-yl.
 8. A salt or hydrate of a compound as defined in any one of claims 1 to 7.
 9. A process for preparing thioketone derivatives of the following formula(I)

Table III (LD₅₀ of Test Compound)

Compound	LD ₅₀ (mg/kg)	Compound	LD ₅₀ (mg/kg)	Compound	LD ₅₀ (mg/kg)
Example 13	< 5,000	Example 22	> 5,000	Example 24	> 5,000
Example 28	> 5,000	Example 29	> 5,000		

Claims


1. Compounds of formula (I);



in which R¹ is a straight or branched alkyl group having 1 to 5 carbon atoms; R² is hydrogen, a straight or branched alkyl group having 1 to 5 carbon atoms, benzyl, phenyl, 4-hydroxyphenyl, halogenophenyl, 4-trifluoromethylphenyl, 2-thiazolyl, 4-methyl-2-thiazolyl, 4-phenyl-2-thiazolyl, 2-pyridyl or 1,3,4-thiadiazol-2-yl; and n is an integer of 1 or 2 but excluding compounds in which


- (i) n is 2, R¹ is ethyl and R² is benzyl;
 - (ii) n is 2, R¹ is isopropyl and R² is benzyl;
 - (iii) n is 2, R¹ is ethyl and R² is chlorophenyl or bromophenyl.
2. Compounds as claimed in claim 1, characterized in that R¹ represents methyl, ethyl, isopropyl, or sec-butyl.
 3. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents methyl, ethyl, isopropyl, or sec-butyl.
 4. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents benzyl.
 5. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents phenyl, 4-hydroxyphenyl, fluorophenyl, chlorophenyl or 4-trifluoromethylphenyl.
 6. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents 2-thiazolyl, 4-methyl-2-thiazolyl or 4-phenyl-2-thiazolyl.
 7. Compounds as claimed in claim 1 or claim 2, characterized in that R² represents 1,3,4-thiadiazol-2-yl.
 8. A salt or hydrate of a compound as defined in any one of claims 1 to 7.
 9. A process for preparing thioketone derivatives of the following formula(I)





in which R^1 is a straight or branched alkyl group having 1 to 5 carbon atoms; R^2 is hydrogen, a straight or branched alkyl group having 1 to 5 carbon atoms, benzyl, phenyl, 4-hydroxyphenyl, halogenphenyl, 4-trifluoromethylphenyl, 2-thiazolyl, 4-methyl-2-thiazolyl, 4-phenyl-2-thiazolyl, 2-pyridyl or 1,3,4-thiadiazol-2-yl; and n is an integer of 1 or 2 which comprises

(a) reacting the compound of the following formula (II)




in which R^1 and n have the same meanings as in formula (I) and R^3 is sec.- or tert.-butyl, a lower alkoxy having 1 to 4 carbon atoms or benzyloxy, with the compound of the following formula (III)




in which R^2 has the same meaning as in formula (I); or

(b) reacting the compound of the following formula (IV)



in which R^1 and n have the same meanings as in formula (I) and R^4 is 1-benzotriazolyl, N-succinimidoyl, N-phthalimidoyl or arylsulfonyl, with the compound of the above formula (III); or


(c) reacting the compound of the following formula (V)



in which R^1 and n have the same meanings as in formula (I), with phosphorus pentachloride, followed by reacting with an amine of the above formula (III) in the presence of pyridine or triethylamine; or

(d) reacting the compound of the following formula (VI)





(VI)

in which R¹ and R² have the same meanings as in formula(I), with the compound of the following formula (VII)




in which X is a halogen and n is an integer of 1 or 2.

10. Process for the preparation of compounds of the general formula I as claimed in any one of claims 1 to 7 wherein R¹, R² and n are as defined in any one of claims 1 to 7 substantially as hereinbefore described.
11. A pharmaceutical composition comprising a pharmacologically effective amount of a compound, salt or hydrate as defined in any one of claims 1 to 8, in association with a pharmaceutically acceptable carrier or diluent.
12. Compound, salt or hydrate as defined in any one of claims 1 to 8 for use as an active therapeutic or prophylactic agent against hepatic damage.

Patentansprüche

1. Verbindungen der Formel (I)



(I)


in welchen R¹ eine gerade oder verzweigte Alkylgruppe mit 1 bis 5 Kohlenstoffatomen ist; R² Wasserstoff, eine gerade oder verzweigte Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, Benzyl, Phenyl, 4-Hydroxyphenyl, Halogenphenyl, 4-Trifluormethylphenyl, 2-Thiazolyl, 4-Methyl-2-thiazolyl, 4-Phenyl-2-thiazolyl, 2-Pyridyl oder 1,3,4-Thiadiazol-2-yl ist; und n eine ganze Zahl 1 oder 2 ist, wobei Verbindungen in welchen

- (i) n 2 ist, R¹ Äthyl ist und R² Benzyl ist;
- (ii) n 2 ist, R¹ Isopropyl ist und R² Benzyl ist;
- (iii) n 2 ist, R¹ Äthyl ist und R² Chlorphenyl oder Bromphenyl ist, ausgeschlossen sind.

2. Verbindungen wie in Anspruch 1 beansprucht, dadurch gekennzeichnet, dass R¹ Methyl, Äthyl, Isopropyl oder sec-Butyl darstellt.
3. Verbindungen wie in Anspruch 1 oder Anspruch 2 beansprucht, dadurch gekennzeichnet, dass R² Methyl, Äthyl, Isopropyl oder sec-Butyl darstellt.
4. Verbindungen wie in Anspruch 1 oder Anspruch 2 beansprucht, dadurch gekennzeichnet, dass R² Benzyl darstellt.
5. Verbindungen wie in Anspruch 1 oder Anspruch 2 beansprucht, dadurch gekennzeichnet, dass R² Phenyl, 4-Hydroxyphenyl, Fluorphenyl, Chlorphenyl oder 4-Trifluormethylphenyl darstellt.




6. Verbindungen wie in Anspruch 1 oder Anspruch 2 beansprucht, dadurch gekennzeichnet, dass R² 2-Thiazolyl, 4-Methyl-2-thiazolyl oder 4-Phenyl-2-thiazolyl darstellt.
7. Verbindungen wie in Anspruch 1 oder Anspruch 2 beansprucht, dadurch gekennzeichnet, dass R² 1,3,4-Thiadiazol-2-yl darstellt.
8. Salz oder Hydrat einer Verbindung, wie sie in einem der Ansprüche 1 bis 7 definiert ist.
9. Verfahren zur Zubereitung von Thioketonderivaten der folgenden Formel (I)




in welcher R¹ eine gerade oder verzweigte Alkylgruppe mit 1 bis 5 Kohlenstoffatomen ist; R² Wasserstoff, eine gerade oder verzweigte Alkylgruppe mit 1 bis 5 Kohlenstoffatomen, Benzyl, Phenyl, 4-Hydroxyphenyl, Halogenphenyl, 4-Trifluormethylphenyl, 2-Thiazolyl, 4-Methyl-2-thiazolyl, 4-Phenyl-2-thiazolyl, 2-Pyridyl oder 1,3,4-Thiadiazol-2-yl ist; und n eine ganze Zahl 1 oder 2 ist, welches

(a) Umsetzen der Verbindung der folgenden Formel (II)




in welcher R¹ und n die gleiche Bedeutung wie in Formel (I) haben und R³ sec- oder tert-Butyl, ein Niederalkoxy mit 1 bis 4 Kohlenstoffatomen oder Benzyloxy ist, mit der Verbindung der folgenden Formel (III)



in welcher R² die gleiche Bedeutung wie in Formel (I) hat; oder


(b) Umsetzen der Verbindung der folgenden Formel (IV)




in welcher R¹ und n die gleiche Bedeutung wie in Formel (I) haben und R⁴ 1-Benzotriazolyl, N-Succinimidoyl, N-Phthalimidoyl oder Arylsulfonyl ist, mit der Verbindung der obigen Formel (III); oder

(c) Umsetzen der Verbindung der folgenden Formel (V)





in welcher R^1 und n die gleiche Bedeutung wie in Formel (I) haben, mit Phosphorpentachlorid, gefolgt von Umsetzen mit einem Amin der obigen Formel (III) im Besein vom Pyridin oder Triäthylamin; oder (d) Umsetzen der Verbindung der folgenden Formel (VI)



in welcher R^1 und R^2 die gleichen Bedeutungen wie in Formel (I) haben, mit der Verbindung der folgenden Formel (VII)




in welcher X ein Halogen und n eine ganze Zahl 1 oder 2 ist, umfasst.

10. Verfahren für die Zubereitung von Verbindungen der allgemeinen Formel 1, wie in einem der Ansprüche 1 bis 7 beansprucht, worin R^1 und R^2 und n wie in einem der Ansprüche 1 bis 7 definiert sind, im wesentlichen wie hierin zuvor beschrieben.
11. Pharmazeutische Zusammensetzung, die eine pharmakologisch wirksame Menge einer Verbindung, eines Salzes oder Hydrates wie in einem der Ansprüche 1 bis 8 definiert, zusammen mit einem pharmazeutisch annehmbaren Träger oder Diluens umfasst.
12. Verbindung, Salz oder Hydrat wie in einem der Ansprüche 1 bis 8 definiert, zur Verwendung als ein therapeutischer oder prophylaktischer Wirkstoff gegen Leberschaden.

Revendications

1. Composés de formule (I)



dans laquelle R^1 est un groupe alkyle à chaîne droite ou ramifiée ayant de 1 à 5 atomes de carbone ; R^2 est un hydrogène ou un radical alkyle à chaîne droite ou ramifiée ayant de 1 à 5 atomes de carbone, benzyle, phényle, 4-hydroxyphényle, halogénophényle, 4-trifluorométhylphényle, 2-thiazolyle, 4-méthyl-2-thiazolyle, 4-phényl-2-thiazolyle, 2-pyridyle ou 1,3,4-thiadiazolyle-2 ; et n est un entier égal à 1 ou 2, mais à l'exclusion des composés dans lesquels :


(i) n vaut 2, R^1 est le radical éthyle et R^2 est le radical benzyle ;



(ii) n vaut 2, R¹ est le radical isopropyle et R² est le radical benzyle ;


(iii) n vaut 2, R¹ est le radical éthyle et R² est le radical chlorophényle ou bromophényle.

2. Composés selon la revendication 1, caractérisés en ce que R¹ représente le radical méthyle, éthyle, isopropyle ou sec-butyle.
3. Composés selon la revendication 1 ou la revendication 2, caractérisés en ce que R² représente le radical méthyle, éthyle, isopropyle ou sec-butyle.
4. Composés selon la revendication 1 ou la revendication 2, caractérisés en ce que R² représente le radical benzyle.
5. Composés selon la revendication 1 ou la revendication 2, caractérisés en ce que R² représente le radical phényle, 4-hydroxyphényle, fluorophényle, chlorophényle ou 4-trifluorométhylphényle.
6. Composés selon la revendication 1 ou la revendication 2, caractérisés en ce que R² représente le radical 2-thiazolyle, 4-méthyl-2-thiazolyle ou 4-phényl-2-thiazolyle.
7. Composés selon la revendication 1 ou la revendication 2, caractérisés en ce que R² représente le radical 1,3,4-thiadiazolyle-2.
8. Sel ou hydrate d'un composé tel que défini dans l'une quelconque des revendications 1 à 7.
9. Procédé pour préparer des dérivés thiocétones ayant la formule (I) ci-après :




dans laquelle R¹ est un groupe alkyle à chaîne droite ou ramifiée ayant de 1 à 5 atomes de carbone ; R² est un hydrogène ou un radical alkyle à chaîne droite ou ramifiée ayant de 1 à 5 atomes de carbone, benzyle, phényle, 4-hydroxyphényle, halogénophényle, 4-trifluorométhylphényle, 2-thiazolyle, 4-méthyl-2-thiazolyle, 4-phényl-2-thiazolyle, 2-pyridyle ou 1,3,4-thiadiazolyle-2 ; et n est un entier égal à 1 ou 2, qui comprend les étapes consistant :

(a) à faire réagir le composé ayant la formule (II) ci-après :




dans laquelle R¹ et n ont les significations données pour la formule (I), et R³ est un radical sec-butyle ou tert-butyle, alcoxy inférieur ayant de 1 à 4 atomes de carbone ou benzyloxy, avec le composé ayant la formule (III) ci-après :



dans laquelle R² a les mêmes significations que dans la formule (I) ; ou bien

(b) à faire réagir le composé ayant la formule (IV) ci-après :






(IV)

dans laquelle R^1 et n ont les significations données pour la formule (I), et R^4 est le radical 1-benzotriazolyle, N-succinimidoyle, N-phthalimidoyle ou arylsulfonyl, avec le composé ayant la formule (III) ci-dessus ; ou bien


(c) à faire réagir le composé ayant la formule (V) ci-après :



(V)

dans laquelle R^1 et n ont les mêmes significations que dans la formule (I), avec du pentachlorure de phosphore, puis à procéder à une réaction avec une amine ayant la formule (III) ci-dessus en présence de pyridine ou de triéthylamine; ou bien

(d) à faire réagir le composé ayant la formule (VI) ci-après



(VI)

dans laquelle R^1 et R^2 ont les significations données pour la formule (I), avec le composé ayant la formule (VII) ci-après :



dans laquelle X est un halogène et n est un entier égal à 1 ou 2.

10. Procédé pour préparer des composés ayant la formule générale (I) selon l'une quelconque des revendications 1 à 7, dans laquelle R^1 , R^2 et n sont tels que définis dans l'une quelconque des revendications 1 à 7, essentiellement tel que décrit ci-dessus.
11. Composition pharmaceutique comprenant une quantité à effet pharmacologique d'un composé, d'un sel ou d'un hydrate selon l'une quelconque des revendications 1 à 8, en association avec un diluant ou un excipient pharmaceutiquement acceptable.
12. Composé, sel ou hydrate selon l'une quelconque des revendications 1 à 8, destiné à être utilisé comme agent thérapeutique ou prophylactique actif contre les troubles hépatiques.

