


NEW BICYCLIC DIHYDROISOQUINOLINE-1-ONE DERIVATIVES

The present invention relates to organic compounds useful for therapy or prophylaxis in a mammal, and in particular to aldosterone synthase (CYP11B2 or CYP11B1) inhibitors for the treatment or prophylaxis of chronic kidney disease, congestive heart failure,

5 hypertension, primary aldosteronism and Cushing syndrom.

The present invention provides novel compounds of formula (I)



wherein

10 R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, halocycloalkylalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl are substituted with R^{12} , R^{13} and R^{14} ;

15 or R^2 and R^4 together form a double bond, wherein in case R^2 and R^4 together form a double bond, then R^5 is H;

or R^1 and R^2 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24} ;

or R^3 and R^4 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{29} , R^{30} and R^{31} ;

5 or R^1 and R^3 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{44} , R^{45} and R^{46} ;

A^1 is CR^8 or N;

A^2 is CR^9 or N;

A^3 is CR^{10} or N;

10 A^4 is CR^{11} or N;

A^5 is CR^6 or N;

one of R^5 , R^6 , R^7 and R^8 is selected from halogen, cyano, alkoxy, hydroxyalkoxy, haloalkyl, haloalkoxy and hydroxy and the others are each independently selected from H, halogen, cyano, alkoxy, hydroxyalkoxy, haloalkoxy and hydroxy;

15 R^9 is H, halogen, hydroxy, cyano, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, alkoxycycloalkylalkyl, dialkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl or substituted heteroarylalkyl, wherein substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl and substituted heteroarylalkyl are substituted with
25 R^{32} , R^{33} and R^{34} ;

R^{10} is $-O_m-(CR^{15}R^{16})_p-(CR^{17}R^{18})_q-(CR^{19}R^{20})_r-R^{21}$;

or R⁹ and R¹⁰ together with the carbon atoms to which they are attached form a substituted cycloalkyl, a substituted heterocycloalkyl, a substituted aryl or a substituted heteroaryl, wherein substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl and substituted heteroaryl are substituted with R³⁵, R³⁶ and R³⁷;

5 R^{11} is H;

R¹⁵, R¹⁷ and R¹⁹ are each independently selected from H, alkyl, cycloalkyl, haloalkyl and halocycloalkyl;

R¹⁶, R¹⁸ and R²⁰ are each independently selected from H, hydroxy, halogen and alkyl;

or R¹⁵ and R¹⁶ together with the carbon atom to which they are attached form a cycloalkyl;

10 or R¹⁷ and R¹⁸ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁹ and R²⁰ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁵ and R¹⁷ together form -(CH₂)_v-;

or R¹⁵ and R¹⁹ together form -(CH₂)_w-;

or R¹⁷ and R¹⁹ together form -(CH₂)_x-;

15 R^{21} is H, halogen, cyano, $-OR^{25}$, $-SR^{25}$, $-S(O)R^{25}$, $-S(O)_2R^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}SO_2NR^{25}R^{27}$, $-NR^{26}C(O)R^{25}$, $-NR^{26}C(O)NR^{25}R^{27}$, $-C(O)R^{28}$, $-C(O)NR^{25}R^{26}$, cycloalkyl, substituted heterocycloalkyl, substituted heteroaryl or substituted aryl, wherein substituted heterocycloalkyl, substituted heteroaryl, substituted heteroarylalkyl and substituted aryl are substituted with R^{38} , R^{39} and R^{40} .

20 R²⁵ is H, alkyl, hydroxyalkyl, carboxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxyalkyl, alkoxyalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxyalkyl, cycloalkoxyalkyl, halocycloalkoxyalkyl, alkylcycloalkoxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, haloalkoxyalkoxyalkyl, substituted heterocycloalkyl, substituted

25 heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted heterocycloalkyl, substituted

heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R^{41} , R^{42} and R^{43} ;

R^{26} and R^{27} are each independently selected from H, alkyl, cycloalkyl, haloalkyl or halocycloalkyl;

5 or R^{15} and R^{26} together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{47} , R^{48} and R^{49} ;

or R^{17} and R^{26} together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted
10 heterocycloalkyl and substituted heteroaryl are substituted with R^{47} , R^{48} and R^{49} ;

or R^{19} and R^{26} together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{47} , R^{48} and R^{49} ;

R^{28} is H, hydroxy, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl,
15 alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxyalkylalkyl,
halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy,
cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy,
alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl,
alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl
20 substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted
heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl,
wherein substituted substituted heterocycloalkyl, substituted heterocycloalkylalkyl,
substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted
arylalkyl are substituted with R^{50} , R^{51} and R^{52} ;

25 R^{12} , R^{13} , R^{14} , R^{22} , R^{23} , R^{24} , R^{29} , R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} ,
 R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} , R^{51} and R^{52} are each independently selected
from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, alkylcarbonyl,
alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl,
halocycloalkyl, alkylcycloalkylalkyl, alkylcarbonylamino, alkylsulfonyl,

alkylsulfonylamino, alkoxyalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy,
cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy,
halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy,
alkoxycarbonyl, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy,
5 alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl,
chloropyridinylcarbonyl and heterocycloalkyl;

n is zero or 1;

m zero or 1;

p, q and r are independently selected from zero and 1;

10 v and x are independently selected from 1, 2, 3 or 4;

w is zero, 1, 2 or 3;

with the proviso that no more than two of A^2 , A^3 and A^4 are N;

and pharmaceutically acceptable salts thereof.

Herein we describe inhibitors of aldosterone synthase that have the potential to
15 protect from organ/ tissue damage caused by an absolute or relative excess of aldosterone.
Hypertension affects about 20% of the adult population in developed countries. In persons
60 years and older, this percentage increases to above 60%. Hypertensive subjects display
an increased risk of other physiological complications including stroke, myocardial
infarction, atrial fibrillation, heart failure, peripheral vascular disease and renal
20 impairment. The renin angiotensin aldosterone system is a pathway that has been linked to
hypertension, volume and salt balance and more recently to contribute directly to end
organ damage in advanced stages of heart failure or kidney disease. ACE inhibitors and
angiotensin receptor blockers (ARBs) are successfully used to improve duration and
quality of life of patients. These drugs are not yielding maximum protection. In a relatively
25 large number of patients ACE and ARB's lead to so-called aldosterone breakthrough, a
phenomenon where aldosterone levels, after a first initial decline, return to pathological
levels. It has been demonstrated that the deleterious consequences of inappropriately
increased aldosterone levels (in relation to salt intake/levels) can be minimized by

aldosterone blockade with mineralocorticoid receptor antagonists. A direct inhibition of aldosterone synthesis is expected to provide even better protection as it will also reduce non-genomic effects of aldosterone as well.

The effects of aldosterone on Na/K transport lead to increased re-absorption of sodium and water and the secretion of potassium in the kidneys. Overall this results in increased blood volume and, therefore, increased blood pressure. Beyond its role in the regulation of renal sodium re-absorption aldosterone can exert deleterious effects on the kidney, the heart and the vascular system especially in a “high sodium” context. It has been shown that under such conditions aldosterone leads to increased oxidative stress which ultimately may contribute to organ damage. Infusion of aldosterone into renally compromised rats (either by high salt treatment or by unilaterally nephrectomy) induces a wide array of injuries to the kidney including glomerular expansion, podocyte injury, interstitial inflammation, mesangial cell proliferation and fibrosis reflected by proteinuria. More specifically aldosterone was shown to increase the expression of the adhesion molecule ICAM-1 in the kidney. ICAM-1 is critically involved in glomerular inflammation. Similarly, aldosterone was shown to increase the expression of inflammatory cytokines, such as interleukin IL-1b and IL-6, MCP-1 and osteopontin. On a cellular level it was demonstrated that in vascular fibroblasts aldosterone increased the expression of type I collagen mRNA, a mediator of fibrosis. Aldosterone also stimulates type IV collagen accumulation in rat mesangial cells and induces plasminogen activator inhibitor-1 (PAI-1) expression in smooth muscle cells. In summary aldosterone has emerged as a key hormone involved in renal damage. Aldosterone plays an equally important role in mediating cardiovascular risk.

There is ample preclinical evidence that MR-antagonists (spironolactone and eplerenone) improve blood pressure, cardiac and renal function in various pre-clinical models.

More recently preclinical studies highlight the important contribution of CYP11B2 to cardiovascular and renal morbidity and mortality. The CYP11B2 inhibitor FAD286 and the MR antagonist spironolactone were evaluated in a rat model of chronic kidney disease (high angiotensin II exposure; high salt and uni-nephrectomy). Angiotensin II and high salt treatment caused albuminuria, azotemia, renovascular hypertrophy, glomerular injury,


increased PAI-1, and osteopontin mRNA expression, as well as tubulointerstitial fibrosis. Both drugs prevented these renal effects and attenuated cardiac and aortic medial hypertrophy. Following 4 weeks of treatment with FAD286, plasma aldosterone was reduced, whereas spironolactone increased aldosterone at 4 and 8 weeks of treatment.

5 Similarly only spironolactone but not FAD286 enhanced angiotensin II and salt-stimulated PAI-1 mRNA expression in the aorta and the heart. In other studies the CYP11B2 inhibitor FAD286 improved blood pressure and cardiovascular function and structure in rats with experimental heart failure. In the same studies FAD286 was shown to improve kidney function and morphology.

10 Administration of an orally active CYP11B2 inhibitor, LCI699, to patients with primary aldosteronism, lead to the conclusion that it effectively inhibits CYP11B2 in patients with primary aldosteronism resulting in significantly lower circulating aldosterone levels and that it corrected the hypokalemia and mildly decreased blood pressure. The effects on the glucocorticoid axis were consistent with a poor selectivity of the compound
15 and a latent inhibition of cortisol synthesis. Taken together these data support the concept that a CYP11B2 inhibitor can lower inappropriately high aldosterone levels. Achieving good selectivity against CYP11B1 is important to be free of undesired side effects on the HPA axis and will differentiate different CYP11B2 inhibitors.

Objects of embodiments disclosed herein are the compounds of formula (I) and their
20 aforementioned salts and esters and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts or esters, the use of the said compounds, salts or esters for the treatment or prophylaxis of illnesses, especially in the treatment or prophylaxis of chronic kidney disease, congestive
25 heart failure, hypertension, primary aldosteronism and Cushing syndrom and the use of the said compounds, salts or esters for the production of medicaments for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

In one aspect, there is provided compounds of formula (I)



wherein

R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl, cycloalkyl,
5 cycloalkylalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl,
haloalkoxyalkyl, halocycloalkylalkyl, substituted heterocycloalkyl, substituted
heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl, wherein
substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl
and substituted heteroarylalkyl are substituted with R^{12} , R^{13} and R^{14} ;

10 or R^2 and R^4 together form a double bond, wherein in case R^2 and R^4 together form a
double bond, then R^5 is H;

or R^1 and R^2 together with the carbon atom to which they are attached form a
substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted
cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24} ;

15 or R^3 and R^4 together with the carbon atom to which they are attached form a
substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted
cycloalkyl and substituted heterocycloalkyl are substituted with R^{29} , R^{30} and R^{31} ;

or R^1 and R^3 together with the carbon atom to which they are attached form a
substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted
20 cycloalkyl and substituted heterocycloalkyl are substituted with R^{44} , R^{45} and R^{46} ;

A^1 is CR^8 or N;

A² is CR⁹ or N;

A³ is CR¹⁰ or N;

A⁴ is CR¹¹ or N;

A⁵ is CR⁶ or N;

5 one of R⁵, R⁶, R⁷ and R⁸ is selected from halogen, cyano, alkoxy, hydroxyalkoxy, haloalkyl, haloalkoxy and hydroxy and the others are each independently selected from H, halogen, cyano, alkoxy, hydroxyalkoxy, haloalkoxy and hydroxy;

R⁹ is H, halogen, hydroxy, cyano, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, alkoxycycloalkylalkyl, dialkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl or substituted heteroarylalkyl, wherein substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl and substituted heteroarylalkyl are substituted with R³², R³³ and R³⁴;

R¹⁰ is -O_m-(CR¹⁵R¹⁶)_p-(CR¹⁷R¹⁸)_q-(CR¹⁹R²⁰)_r-R²¹;

20 or R⁹ and R¹⁰ together with the carbon atoms to which they are attached form a substituted cycloalkyl, a substituted heterocycloalkyl, a substituted aryl or a substituted heteroaryl, wherein substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl and substituted heteroaryl are substituted with R³⁵, R³⁶ and R³⁷;

R¹¹ is H;

25 R¹⁵, R¹⁷ and R¹⁹ are each independently selected from H, alkyl, cycloalkyl, haloalkyl and halocycloalkyl;

R¹⁶, R¹⁸ and R²⁰ are each independently selected from H, hydroxy, halogen and alkyl;

or R¹⁵ and R¹⁶ together with the carbon atom to which they are attached form a cycloalkyl;

5 or R¹⁷ and R¹⁸ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁹ and R²⁰ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁵ and R¹⁷ together form -(CH₂)_v-;

10 or R¹⁵ and R¹⁹ together form -(CH₂)_w-;

or R¹⁷ and R¹⁹ together form -(CH₂)_x-;

15 R^{21} is H, halogen, cyano, $-OR^{25}$, $-SR^{25}$, $-S(O)R^{25}$, $-S(O)_2R^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}SO_2NR^{25}R^{27}$, $-NR^{26}C(O)R^{25}$, $-NR^{26}C(O)NR^{25}R^{27}$, $-C(O)R^{28}$, $-C(O)NR^{25}R^{26}$, cycloalkyl, substituted heterocycloalkyl, substituted heteroaryl or substituted aryl, wherein substituted heterocycloalkyl, substituted heteroaryl, substituted heteroarylalkyl and substituted aryl are substituted with R^{38} , R^{39} and R^{40} .

R²⁵ is H, alkyl, hydroxyalkyl, carboxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxyalkyl, alkoxyalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxyalkyl, cycloalkoxyalkyl, halocycloalkoxyalkyl, alkylcycloalkoxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, haloalkoxyalkoxyalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R⁴¹, R⁴² and R⁴³;

R²⁶ and R²⁷ are each independently selected from H, alkyl, cycloalkyl, haloalkyl or halocycloalkyl;

or R¹⁵ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

5 or R¹⁷ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

10 or R¹⁹ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

R²⁸ is H, hydroxy, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxyalkyl, alkoxyalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R⁵⁰, R⁵¹ and R⁵²;

25 R¹², R¹³, R¹⁴, R²², R²³, R²⁴, R²⁹, R³⁰, R³¹, R³², R³³, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹, R⁴⁰, R⁴¹, R⁴², R⁴³, R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷, R⁴⁸, R⁴⁹, R⁵⁰, R⁵¹ and R⁵² are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, alkylcarbonyl, alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkylcarbonylamino, alkylsulfonyl, alkylsulfonylamino, alkoxyalkylalkyl, alkoxyalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy,

halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxy carbonyl, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, chloropyridinylcarbonyl and heterocycloalkyl;

5 n is zero or 1;

m zero or 1;

p, q and r are independently selected from zero and 1;

v and x are independently selected from 1, 2, 3 or 4;

w is zero, 1, 2 or 3;

10 with the proviso that no more than two of A^2 , A^3 and A^4 are N;

and wherein the compound is selected from

6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;

5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-nicotinonitrile;

6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-(5-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-fluoro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-methyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

20 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde;

6-Chloro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 one;
- 6-Chloro-2-[5-((S)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 2-(5-Amino-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2,2,2-trifluoro-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one
- 6-Chloro-2-[5-(cyclopropyl-hydroxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 one;
- 6-Chloro-2-[5-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-trifluoromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(1-methoxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide;

- 6-Chloro-2-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-8'-hydroxy-3,4,5',6',7',8'-hexahydro-[2,4']biisoquinoliny-1-one;
- N-(6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;
- 6-Chloro-2-{5-[hydroxy-(1-methyl-1H-imidazol-2-yl)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[(3,4-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3,5-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(4-ethyl-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(hydroxy-phenyl-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-1-phenyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[1-(3,4-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(3,5-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(6-methyl-pyrazin-2-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(morpholine-4-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N,N-dimethyl-nicotinamide;
- 6-Chloro-2-[5-(pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-methyl-nicotinamide ;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-cyclopropyl-nicotinamide;
- 10 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-(4-fluoro-phenyl)-nicotinamide;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-phenyl-nicotinamide;
- 6-Chloro-2-[5-(4,4-difluoro-piperidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-
- 20 dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyridazin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyridin-3-yl-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-2H-isoquinolin-1-one;

- 6-Chloro-2-[4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 2-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methylsulfanyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-dimethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-fluoro-4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{4-[(4-fluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(1-methyl-1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-methyl-2H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(5-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one;
- 5 3-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-isonicotinonitrile;
- 6-Chloro-2-(5-fluoro-4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-(4-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(3,5-dimethyl-3H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- (S)-1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester;

- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- (R)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 10 (S)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 8-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Methoxy-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5,6-Dichloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 2-Chloro-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 15 2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 2-Methoxy-6-pyridin-3-yl-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-5-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-7-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-7-fluoro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-4,4-dimethyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-4-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 5-Chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 10 one;
- 5-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 15 5-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- one;
- Ethanesulfonic acid [5-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;
- 5-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-2,3-dihydro-isoindol-1-one;
- 20 5-Chloro-2-(8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;
- one;
- 5-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 6-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;

- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-methyl-2-(4-methyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 5 3-Benzyl-5-chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-methoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3,3-dimethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 10 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 15 one;
- 5-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3,3-dimethyl-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 20 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5 2-Hydroxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-2-(5-imidazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2-ethyl-4-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 Propane-2-sulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-((R)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3,5-dimethyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide

N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-acetamide

10 6-Chloro-2-(5-morpholin-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1,1-dioxo-1λ6-thiomorpholin-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-propyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

25 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-1H-imidazole-2-carboxylic acid ethyl ester;

6-Chloro-2-[5-(2-hydroxymethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(oxetan-3-ylaminomethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-{5-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-isopropyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4,4-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3,3-difluoro-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.3]hept-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(3,3-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-[1,2,3]triazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

25 6-Chloro-2-(5-[1,2,3]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-chloro-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3-methyl-[1,2,4]triazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(5-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-chloro-2-[5-(3-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-(5-[1,2,4]triazol-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-methyl-benzoimidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-(5-indazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-indazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(6-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(7-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2-cyclopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(3-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-ethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 2-(5-Aminomethyl-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methyl-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[2-(1-methyl-pyrrolidin-2-yl)-ethoxymethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(5-cyclopentyloxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropylmethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-fluoro-phenoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-cyclopropylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(tetrahydro-furan-2-ylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25

- 6-Chloro-2-[5-(2,2,2-trifluoro-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclobutoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-methanesulfonyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(6-methyl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(6-morpholin-4-yl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(2,3-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(2,6-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-chloro-6-fluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3,4-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(2,5-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;

10 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-benzenesulfonamide;

N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-methanesulfonamide;

Cyclopropanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;

15 6-Chloro-2-[5-(4-fluoro-benzylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,2,2-trifluoro-ethylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-morpholin-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

20 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-propionamide;

6-Chloro-2-{5-[(2-methyl-2H-pyrazol-3-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

25 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylamino]-2-methyl-propionic acid;

6-Chloro-2-{5-[(1-methyl-1H-imidazol-4-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3,4-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(3,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3-chloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(3-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3-trifluoromethoxy-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-chloro-3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

25 6-Chloro-2-[5-(3,4-dichloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;


- 6-Chloro-2-[5-(2-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isoxazol-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-1H-imidazol-2-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5
- 6-Chloro-2-[5-(2,4-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-1-methyl-1H-pyrazole-4-carbonitrile
- 10
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-isobutyramide
- Cyclopropanecarboxylic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-4-fluorobenzamide
- 15
- 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-cyclohexylurea
- 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-(3-trifluoromethyl-phenyl)-urea
- 6-Chloro-2-(5-hydroxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 20
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetamide;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N-methylacetamide;
- [5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid methyl ester;

- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N,N-dimethyl-acetamide;
- 6-Chloro-2-(5-phenylaminomethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-{5-[(4-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(4-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[(3-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-morpholin-4-yl-2-oxo-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-N-(2-hydroxy-ethyl)-acetamide;
- 6-Chloro-2-[5-(1-methylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(1-dimethylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-1H-imidazole-2-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1-[1,2,3]triazol-2-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-imidazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(1-pyrazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(oxazol-2-ylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(1-[1,2,4]triazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(2-oxo-pyrrolidin-1-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(2-oxo-oxazolidin-3-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 N-{1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-ethyl}-methanesulfonamide;
- 6-Chloro-2-{5-[1-(3-fluoro-phenylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(1-phenylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methanesulfinyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

and pharmaceutically acceptable salts thereof.

In another aspect, there is provided a process to prepare a compound as defined herein comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



- 5 wherein $R^1, R^2, R^3, R^4, R^5, R^6, A^1, A^2, A^3, A^4, A^5$ and n are as defined herein and wherein X is halogen or triflate.

In another aspect, there is provided a compound as defined herein for use as therapeutically active substance.

- 10 In another aspect, there is provided a pharmaceutical composition comprising a compound as defined herein and a therapeutically inert carrier.

In another aspect, there is provided a compound as defined herein for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

- 15 In another aspect, there is provided use of a compound as defined herein for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

The term “alkoxy” denotes a group of the formula $-O-R'$, wherein R' is an alkyl group. Examples of alkoxy group include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Particular alkoxy group include methoxy.

The term “alkoxyalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by another alkoxy group. Examples of alkoxyalkoxy group include methoxymethoxy, ethoxymethoxy, methoxyethoxy, ethoxyethoxy, methoxypropoxy and ethoxypropoxy. Particular alkoxyalkoxy groups
5 include methoxymethoxy and methoxyethoxy.

The term “alkoxyalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an alkoxyalkoxy group. Examples of alkoxyalkoxyalkyl group include methoxymethoxymethyl, ethoxymethoxymethyl, methoxyethoxymethyl, ethoxyethoxymethyl, methoxypropoxymethyl,
10 ethoxypropoxymethyl, methoxymethoxyethyl, ethoxymethoxyethyl, methoxyethoxyethyl, ethoxyethoxyethyl, methoxypropoxyethyl and ethoxypropoxyethyl.

The term “alkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an alkoxy group. Exemplary alkoxyalkyl groups include methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl,
15 methoxypropyl, ethoxypropyl and isopropoxymethyl. Particular alkoxyalkyl group include include methoxymethyl, methoxyethyl and isopropoxymethyl.

The term “alkoxycarbonyl” denotes a group of the formula $-C(O)-R'$, wherein R' is an alkoxy group. Examples of alkoxycarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and
20 tert-butoxy. Particular alkoxycarbonyl group is a group of the formula $-C(O)-R'$, wherein R' is methoxy.

The term “alkoxycycloalkylalkyl” denotes a cycloalkylalkyl group wherein at least one of the hydrogen atoms of the cycloalkyl group is replaced by an alkoxy group. An example of alkoxycycloalkylalkyl group is cyclopropylmethoxymethyl.

25 The term “alkyl” denotes a monovalent linear or branched saturated hydrocarbon group of 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, and. Particular alkyl groups include methyl, ethyl, propyl and isopropyl. More particular alkyl groups are methyl,
30 isopropyl and ethyl.

The term “alkylcarbonyl” of the formula $-C(O)-R'$, wherein R' is an alkyl group. Examples of alkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methyl or ethyl.

The term “alkylcarbonylamino” denotes an amino group wherein one of the
5 hydrogen atoms of the $-NH_2$ group is replaced by an alkylcarbonyl group. Examples of alkylcarbonylamino groups include groups wherein R' is methyl or ethyl. Particular alkylcarbonylamino groups include groups wherein R' is ethyl.

The term “alkylcarbonylaminoalkyl” denotes an aminoalkyl group wherein one of the hydrogen atoms of the $-NH_2$ group is replaced by an alkylcarbonyl group. Examples of
10 alkylcarbonylaminoalkyl groups include groups wherein R' is methyl or ethyl.

The term “alkylcycloalkoxy” denotes a cycloalkoxy group wherein at least one of the hydrogen atoms of the cycloalkoxy group is replaced by an alkyl group. Examples of alkylcycloalkyl include methyl- cyclopropoxy, dimethyl- cyclopropoxy, methyl- cyclobutoxy, dimethyl- cyclobutoxy, methyl- cyclopentoxo, dimethyl- cyclopentoxo,
15 methyl- cyclohexyloxy and dimethyl- cyclohexyloxy.

The term “alkylcycloalkoxyalkyl” denotes a cycloalkoxyalkyl group wherein at least one of the hydrogen atoms of the cycloalkoxyalkyl group is replaced by an alkyl group. Examples of alkylcycloalkyl include methyl- cyclopropoxymethyl, dimethyl- cyclopropoxymethyl, methyl- cyclobutoxymethyl, dimethyl- cyclobutoxymethyl, methyl- cyclopentoxymethyl, dimethyl- cyclopentoxymethyl, methyl- cyclohexyloxymethyl and
20 dimethyl- cyclohexyloxymethyl.

The term “alkylcycloalkyl” denotes a cycloalkyl group wherein at least one of the hydrogen atoms of the cycloalkyl group is replaced by an alkyl group. Examples of alkylcycloalkyl include methyl- cyclopropyl, dimethyl- cyclopropyl, methyl- cyclobutyl, dimethyl- cyclobutyl, methyl- cyclopentyl, dimethyl- cyclopentyl, methyl- cyclohexyl and
25 dimethyl- cyclohexyl. Particular alkylcycloalkyl groups include methyl- cyclopropyl and dimethyl- cyclopropyl.

The term “alkylcycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by an alkylcycloalkyl group. Examples of

alkylcycloalkylalkyl include methyl-cyclopropylmethyl, dimethyl-cyclopropylmethyl, methyl-cyclopropylethyl, dimethyl-cyclopropylethyl, methyl-cyclobutylmethyl, dimethyl-cyclobutylmethyl, methyl-cyclobutylethyl, dimethyl-cyclobutylethyl, methyl-cyclopentylmethyl, dimethyl-cyclopentylmethyl, methyl-cyclopentylethyl, dimethyl-cyclopentylethyl, methyl-cyclohexylmethyl, dimethyl-cyclohexylmethyl, methyl-cyclohexylethyl, dimethyl-cyclohexylethyl, methyl-cycloheptylmethyl, dimethyl-cycloheptylmethyl, methyl-cycloheptylethyl, dimethyl-cycloheptylethyl, methyl-cyclooctylmethyl, dimethyl-cyclooctylmethyl, methyl-cyclooctylethyl and dimethyl-cyclooctylethyl.

10 The term “alkylsulfonyl” denotes a group of the formula $-S(O)_2-R'$, wherein R' is an alkyl group. Examples of alkylsulfonyl groups include groups of the formula $-S(O)_2-R'$, wherein R' is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl.

15 The term “alkylsulfonylamino” denotes a group of the formula $-NH_2-S(O)_2-R'$, wherein R' is an alkyl group. Examples of alkylsulfonyl groups include groups of the formula $-NH_2-S(O)_2-R'$, wherein R' is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl.

The term “amino” denotes a $-NH_2$ group.

20 The term “aryl” denotes a monovalent aromatic carbocyclic mono- or bicyclic ring system comprising 6 to 10 carbon ring atoms. Examples of aryl group include phenyl and naphthyl. Particular aryl group is phenyl.

The term “arylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced an aryl group. Particular arylalkyl group is phenylalkyl. More particular arylalkyl group is benzyl.

25 The term “arylhydroxyalkyl” denotes an hydroxyalkyl group wherein at least one of the hydrogen atoms of the hydroxyalkyl group has been replaced an aryl group. Particular arylalkyl group is phenylhydroxymethyl.

The term “bicyclic ring system” denotes two rings which are fused to each other via a common single or double bond (annelated bicyclic ring system), via a sequence of three or more common atoms (bridged bicyclic ring system) or via a common single atom (spiro bicyclic ring system). Bicyclic ring systems can be saturated, partially unsaturated,
5 unsaturated or aromatic. Bicyclic ring systems can comprise heteroatoms selected from N, O and S.

The term “carbonyl” denotes a -C(O)- group.

The term “carboxyl” denotes a -C(O)OH group.

The term “carboxyalkyl” denotes an alkyl group wherein at least one of the hydrogen
10 atoms of the alkyl group has been replaced by a carboxy group. Examples are carboxymethyl, carboxyethyl, carboxypropyl and 1-carboxy-2-methylpropyl. Particular example is 1-carboxy-2-methylpropyl.

The term “cyano” denotes a $\text{-C}\equiv\text{N}$ group.

The term “cycloalkoxy” denotes a group of the formula -O-R' , wherein R' is a
15 cycloalkyl group. Examples of cycloalkoxy group include cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy and cyclooctyloxy. Particular cycloalkoxy group is cyclopropoxy.

The term “cycloalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a cycloalkoxy group. Examples of
20 cycloalkoxyalkyl group include cyclopropoxymethyl, cyclopropoxyethyl, cyclobutoxymethyl, cyclobutoxyethyl, cyclopentyloxymethyl, cyclopentyloxyethyl, cyclohexyloxymethyl, cyclohexyloxyethyl, cycloheptyloxymethyl, cycloheptyloxyethyl, cyclooctyloxymethyl and cyclooctyloxyethyl.

The term “cycloalkyl” denotes a monovalent saturated monocyclic or bicyclic
25 hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments, cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means consisting of two saturated carbocycles having two carbon atoms in common. Particular cycloalkyl groups are monocyclic. Examples for monocyclic

cycloalkyl are cyclopropyl, cyclobutanyl, cyclopentyl, cyclohexyl or cycloheptyl.

Examples for bicyclic cycloalkyl are bicyclo[2.2.1]heptanyl or bicyclo[2.2.2]octanyl.

Particular monocyclic cycloalkyl groups are cyclopropyl, cyclobutanyl, cyclopentyl and

cyclohexyl. More particular monocyclic cycloalkyl group is cyclopropyl. In particular, the

- 5 cycloalkyl formed by R⁹ and R¹⁰ together with the carbon atoms to which they are attached is cyclohexyl and cyclopentyl. Further particular the cycloalkyl formed by R⁹ and R¹⁰ together with the carbon atoms to which they attached is cyclohexyl.

In particular the cycloalkyl formed by R¹ and R² together with the carbon atoms to which they attached is cyclopropyl.

- 10 The term “cycloalkylalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group is replaced by a cycloalkyl group. Examples of cycloalkylalkoxy include cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, cycloheptylmethoxy and cyclooctylmethoxy.

- The term “cycloalkylalkoxyalkyl” denotes an alkyl group wherein at least one of the
15 hydrogen atoms of the alkyl group is replaced by a cycloalkylalkoxy group. Examples of cycloalkylalkoxyalkyl include cyclopropylmethoxymethyl, cyclopropylmethoxyethyl, cyclobutylmethoxymethyl, cyclobutylmethoxyethyl, cyclopentylmethoxyethyl, cyclopentylmethoxyethyl, cyclohexylmethoxymethyl, cyclohexylmethoxyethyl, cycloheptylmethoxymethyl, cycloheptylmethoxyethyl, cyclooctylmethoxymethyl and
20 cyclooctylmethoxyethyl.

- The term “cycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a cycloalkyl group. Examples of cycloalkylalkyl include cyclopropylmethyl, cyclopropylethyl, cyclopropylbutyl, cyclobutylpropyl, 2-cyclopropylbutyl and cyclopentylbutyl. Particular examples of
25 cycloalkylalkyl groups are cyclopropylmethyl, cyclopropylbutyl and 2-cyclopropylbutyl.

The term “cycloalkylcarbonyl” of the formula -C(O)-R', wherein R' is a cycloalkyl group. Examples of cycloalkylcarbonyl groups include groups of the formula -C(O)-R', wherein R' is cyclopropyl.

The term “cycloalkylcarbonylamino” denotes an amino group wherein one of the hydrogen atoms of the -NH₂ group is replaced by an cycloalkylcarbonyl group. Examples of alkylcarbonylamino groups include groups wherein R' is cyclopropyl.

The term “cycloalkylcarbonylaminoalkyl” denotes an aminoalkyl group wherein one
5 of the hydrogen atoms of the -NH₂ group is replaced by an cycloalkylcarbonyl group. Examples of alkylcarbonylaminoalkyl groups include groups wherein R' is cyclopropyl.

The term “dialkoxyalkyl” denotes an alkyl group wherein two of the hydrogen atoms of the alkyl group have been replaced by two alkoxy group. Exemplary dialkoxyalkyl groups include dimethoxymethyl, diethoxymethyl, dimethoxyethyl, diethoxyethyl,
10 dimethoxypropyl, diethoxypropyl and diisopropoxymethyl. Particular dialkoxyalkyl group is dimethoxymethyl.

The term “haloalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by same or different halogen atoms. The term “perhaloalkoxy” denotes an alkoxy group where all hydrogen atoms of the alkoxy group
15 have been replaced by the same or different halogen atoms. Examples of haloalkoxy include fluoromethoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, trifluoromethylethoxy, trifluorodimethylethoxy and pentafluoroethoxy. Particular haloalkoxy groups are trifluoromethoxy and 2,2-difluoroethoxy.

The term “haloalkoxyalkoxy” denotes an alkoxy group wherein at least one of the
20 hydrogen atoms of the alkoxy group has been replaced by a haloalkoxy group. Examples of haloalkoxyalkyl include fluoromethoxymethoxy, difluoromethoxymethoxy, trifluoromethoxymethoxy, fluoroethoxymethoxy, difluoroethoxymethoxy, trifluoroethoxymethoxy, fluoromethoxyethoxy, difluoromethoxyethoxy, trifluoromethoxyethoxy, fluoroethoxyethoxy, difluoroethoxyethoxy,
25 trifluoroethoxyethoxy, fluoromethoxypropoxy, difluoromethoxypropoxy, trifluoromethoxypropoxy, fluoroethoxypropoxy, difluoroethoxypropoxy and trifluoroethoxypropoxy.

The term “haloalkoxyalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a haloalkoxyalkoxy group.
30 Examples of haloalkoxyalkyl include fluoromethoxymethoxymethyl,

5 difluoromethoxymethoxymethyl, trifluoromethoxymethoxymethyl,
fluoroethoxymethoxymethyl, difluoroethoxymethoxymethyl,
trifluoroethoxymethoxymethyl, fluoromethoxyethoxymethyl,
difluoromethoxyethoxymethyl, trifluoromethoxyethoxymethyl, fluoroethoxyethoxymethyl,
difluoroethoxyethoxymethyl, trifluoroethoxyethoxymethyl, fluoromethoxypropoxymethyl,
difluoromethoxypropoxymethyl, trifluoromethoxypropoxymethyl,
fluoroethoxypropoxymethyl, difluoroethoxypropoxymethyl and
trifluoroethoxypropoxymethyl.

The term “haloalkoxyalkyl” denotes an alkyl group wherein at least one of the
10 hydrogen atoms of the alkyl group has been replaced by a haloalkoxy group. Examples of
haloalkoxyalkyl include fluoromethoxymethyl, difluoromethoxymethyl,
trifluoromethoxymethyl, fluoroethoxymethyl, difluoroethoxymethyl,
trifluoroethoxymethyl, fluoromethoxyethyl, difluoromethoxyethyl, trifluoromethoxyethyl,
fluoroethoxyethyl, difluoroethoxyethyl, trifluoroethoxyethyl, fluoromethoxypropyl,
15 difluoromethoxypropyl, trifluoromethoxypropyl, fluoroethoxypropyl,
difluoroethoxypropyl and trifluoroethoxypropyl. Particular haloalkoxyalkyl is 2,2-
difluoroethoxyethyl.

The term “haloalkyl” denotes an alkyl group wherein at least one of the hydrogen
atoms of the alkyl group has been replaced by same or different halogen atoms. The term
20 “perhaloalkyl” denotes an alkyl group where all hydrogen atoms of the alkyl group have
been replaced by the same or different halogen atoms. Examples of haloalkyl include
fluoromethyl, difluoromethyl, trifluoromethyl, trifluoroethyl, trifluoromethylethyl and
pentafluoroethyl. Particular haloalkyl groups are trifluoromethyl. Also particular groups
are difluoromethyl.

25 The term “halocycloalkoxy” denotes a cycloalkoxy group wherein at least one of the
hydrogen atoms of the cycloalkoxy group has been replaced by same or different halogen
atoms, particularly fluoro atoms. Examples of halocycloalkoxy groups include
fluorocyclopropoxy, difluorocyclopropoxy, fluorocyclobutoxy and difluorocyclobutoxy.

The term “halocycloalkoxyalkyl” denotes an alkyl group wherein at least one of the
30 hydrogen atoms of the alkyl group has been replaced by a halocycloalkoxy group,

particularly fluoro atoms. Examples of halocycloalkoxyalkyl groups include fluorocyclopropoxymethyl, difluorocyclopropoxymethyl, fluorocyclobutoxymethyl and difluorocyclobutoxymethyl.

The term “halocycloalkyl” denotes a cycloalkyl group wherein at least one of the
5 hydrogen atoms of the cycloalkyl group has been replaced by same or different halogen atoms, particularly fluoro atoms. Examples of halocycloalkyl groups include fluorocyclopropyl, difluorocyclopropyl, fluorocyclobutyl and difluorocyclobutyl.

The term “halocycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a halocycloalkyl. Examples of
10 halocycloalkylalkyl groups include fluorocyclopropylmethyl, fluorocyclopropylethyl, difluorocyclopropylmethyl, difluorocyclopropylethyl, fluorocyclobutylmethyl, fluorocyclobutylethyl, difluorocyclobutylmethyl and difluorocyclobutylethyl.

The term “halogen” and “halo” are used interchangeably herein and denote fluoro, chloro, bromo, or iodo. Particular halogens are chloro and fluoro.

15 The term “heteroaryl” denotes a monovalent aromatic heterocyclic mono- or bicyclic ring system of 5 to 12 ring atoms, comprising 1, 2, 3 or 4 heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples of heteroaryl group include pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, triazinyl,
20 azepinyl, diazepinyl, isoxazolyl, benzofuranyl, isothiazolyl, benzothienyl, indolyl, isoindolyl, isobenzofuranyl, benzimidazolyl, benzoxazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzooxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl. Particular heteroaryl groups include pyrrolyl, pyrazolyl, imidazolyl, triazolyl, benzoimidazolyl, indazolyl,
25 indolyl, pyridinyl, isooxazolyl and oxazolyl.

In particular, the heteroaryl groups formed by R⁹ and R¹⁰ together with the carbon atoms to which they are attached are pyrrolyl and pyrazolyl. In particular, in the case of R²¹, the term “heteroaryl” denotes the imidazolyl, pyrazolyl, triazolyl, benzoimidazolyl, indazolyl, indolyl, pyridinyl and isooxazolyl groups. More particularly the pyrazolyl
30 group.

In particular, in the case of R^{25} , the term “heteroaryl” denotes the pyridinyl, pyrazolyl and oxazolyl groups. In particular, in the case of R^{25} , the term “heteroaryl” denotes the pyrazolyl and oxazolyl groups. Further particular heteroaryl in the case of R^{25} is pyrazolyl and pyridinyl. More particular is pyrazolyl.

5 The term “heteroarylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced a heteroaryl group. Examples are pyrazolylalkyl and imidazolylalkyl. More particular examples are pyrazolylmethyl and imidazolylmethyl.

10 In particular, in the case of R^{25} , the term “heteroarylalkyl” denotes the pyrazolylmethyl and imidazolylmethyl groups. In particular the pyrazolylmethyl group.

The term “heterocycloalkyl” denotes a monovalent saturated or partly unsaturated mono- or bicyclic ring system of 3 to 9 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. In particular embodiments, heterocycloalkyl is a monovalent saturated monocyclic ring system of 4 to 7
15 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples for monocyclic saturated heterocycloalkyl are aziridinyl, oxiranyl, azetidiny, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl,
20 thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, oxazepanyl and thiazinanyl. Examples for bicyclic saturated heterocycloalkyl are 8-aza-bicyclo[3.2.1]octyl, quinuclidinyl, 8-oxa-3-aza-bicyclo[3.2.1]octyl, 9-aza-bicyclo[3.3.1]nonyl, 3-oxa-9-aza-bicyclo[3.3.1]nonyl, 3-thia-9-aza-bicyclo[3.3.1]nonyl and 2,6-diaza-spiro[3.3]heptanyl. Examples for partly unsaturated heterocycloalkyl are
25 dihydrofuryl, imidazolinyl, dihydro-oxazolyl, tetrahydro-pyridinyl, or dihydropyranyl. More particular examples of heterocycloalkyl group are pyrrolidinyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, oxazepanyl, thiazinanyl and
30 2,6-diaza-spiro[3.3]heptanyl. More particular examples of a heterocycloalkyl are pyrrolydinyl, piperidinyl, thiomorpholinyl, thiazinanyl and 2,6-diaza-spiro[3.3]heptanyl.

In particular, in the case of R^{21} , the term “heterocycloalkyl” denotes the pyrrolidinyl, thiazinan-2-yl, isothiazolidin-2-yl, piperidinyl, morphonlinyl, thiomorphonlinyl, oxazolidinyl, imidazolidinyl, piperazinyl, 2-oxa-6-aza-spiro[3.4]heptanyl and 2-oxa-6-aza-spiro[3.4]octanyl groups. Further particularly the piperazinyl and 2-oxa-6-aza-spiro[3.4]octanyl groups. More particularly the 2-oxa-6-aza-spiro[3.4]octanyl group. Also more particularly the piperazinyl group.

In particular, in the case of R^{25} , the term “heterocycloalkyl” denotes the oxetanyl group.

In particular, in the case of R^{28} , the term “heterocycloalkyl” denotes the morpholinyl, pyrrolidinyl and piperidinyl groups.

In particular the heterocycloalkyl formed by R^{15} and R^{26} together with the nitrogen atom and carbon atom to which they are attached are azetanyl, pyrrolidinyl and piperidinyl.

The term “heterocycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced a heterocycloalkyl group. Examples are tetrahydrofuranylalkyl, pyrrolidinylalkyl and piperazin-1-ylalkyl. More particularly, tetrahydrofuranylethyl, pyrrolidinylmethyl and piperazin-1-ylmethyl.

In particular, in the case of R^9 , the term “heterocycloalkylalkyl” denotes the piperazin-1-ylalkyl group. More particularly, the piperazin-1-ylalkyl group.

In particular, in the case of R^{25} , the term “heterocycloalkylalkyl” denotes the tetrahydrofuranylalkyl and pyrrolidinylalkyl groups. More particularly, the tetrahydrofuranylethyl and pyrrolidinylmethyl groups.

The term “hydroxy” denotes a -OH group.

The term “hydroxyalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by a hydroxy group. Examples of hydroxyalkyl include hydroxyethethoxy, hydroxypropoxy and hydroxymethylpropoxy.

The term "hydroxyalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a hydroxy group. Examples of hydroxyalkyl include hydroxymethyl, hydroxyethyl, hydroxy-1-methyl-ethyl, hydroxypropyl, hydroxymethylpropyl and dihydroxypropyl. Particular examples are

5 hydroxymethyl and hydroxyethyl.

The term "oxo" denotes a divalent oxygen atom =O.

The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as

10 hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, in particular hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the

15 like. In addition these salts may be prepared by addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic

20 amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polyimine resins and the like. Particular pharmaceutically acceptable salts of compounds of formula (I) are the hydrochloride salts, methanesulfonic acid salts and citric acid salts.

25 "Pharmaceutically acceptable esters" means that compounds of general formula (I) may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compounds in vivo. Examples of such compounds include physiologically acceptable and metabolically labile ester derivatives, such as methoxymethyl esters, methylthiomethyl esters and pivaloyloxymethyl esters.

30 Additionally, any physiologically acceptable equivalents of the compounds of general

formula (I), similar to the metabolically labile esters, which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope of this invention.

The term “protecting group” (PG) denotes the group which selectively blocks a reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Protecting groups can be removed at the appropriate point. Exemplary protecting groups are amino-protecting groups, carboxy-protecting groups or hydroxy-protecting groups. Particular protecting groups are the tert-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), fluorenylmethoxycarbonyl (Fmoc) and benzyl (Bn). Further particular protecting groups are the tert-butoxycarbonyl (Boc) and the fluorenylmethoxycarbonyl (Fmoc). More particular protecting group is the tert-butoxycarbonyl (Boc).

The abbreviation uM means microMolar and is equivalent to the symbol μM .

The compounds of the present invention can also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the present invention also embraces isotopically-labeled variants of the present invention which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having the atomic mass or mass number different from the predominant atomic mass or mass number usually found in nature for the atom. All isotopes of any particular atom or element as specified are contemplated within the scope of the compounds of the invention, and their uses. Exemplary isotopes that can be incorporated in to compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, chlorine and iodine, such as ^2H (“D”), ^3H (“T”), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{32}P , ^{33}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I and ^{125}I . Certain isotopically labeled compounds of the present invention (e.g., those labeled with ^3H or ^{14}C) are useful in compound and /or substrate tissue distribution assays. Tritiated (^3H) and carbon-14 (^{14}C) isotopes are useful for their ease of preparation and detectability. Further substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g., increased in vivo half-life or reduced dosage requirements) and hence may be preferred in some circumstances. Positron emitting isotopes such as ^{15}O , ^{13}N , ^{11}C , and ^{18}F are useful for

positron emission tomography (PET) studies to examine substrate receptor occupancy. Isotopically labeled compounds of the present inventions can generally be prepared by following procedures analogous to those disclosed in the Schemes and/or in the Examples herein below, by substituting a non-isotopically labeled reagent with a isotopically labeled
5 reagent. In particular, compounds of formula (I) wherein one or more H atom have been replaced by a ^2H atom are also an embodiment of this invention.

The compounds of formula (I) can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers,
10 diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

According to the Cahn-Ingold-Prelog Convention the asymmetric carbon atom can be of the "R" or "S" configuration.

Also an embodiment of the present invention are compounds according to formula (I) as described herein and pharmaceutically acceptable salts or esters thereof, in particular
15 compounds according to formula (I) as described herein and pharmaceutically acceptable salts thereof, more particularly compounds according to formula (I) as described herein.

Also an embodiment of the present invention are compounds according to formula (I) as described herein wherein

R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl, cycloalkyl,
20 cycloalkylalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, halocycloalkylalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl are substituted with R^{12} , R^{13} and R^{14} ;
25 or R^2 and R^4 together form a double bond, wherein in case R^2 and R^4 together form a double bond, then R^5 is H;

or R¹ and R² together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R²², R²³ and R²⁴;

5 or R³ and R⁴ together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R²⁹, R³⁰ and R³¹;

or R¹ and R³ together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R⁴⁴, R⁴⁵ and R⁴⁶;

10 A¹ is CR⁸ or N;

A² is CR⁹ or N;

A³ is CR¹⁰ or N;

A⁴ is CR¹¹ or N;

A⁵ is CR⁶ or N;

15 one of R⁵, R⁶, R⁷ and R⁸ is selected from halogen, cyano, alkoxy, hydroxyalkoxy, haloalkyl, haloalkoxy and hydroxy and the others are each independently selected from H, halogen, cyano, alkoxy, hydroxyalkoxy, haloalkoxy and hydroxy;

R⁹ is H, halogen, hydroxy, cyano, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, 20 alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, alkoxycycloalkylalkyl, dialkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, substituted arylalkyl, 25 substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl or substituted heteroarylalkyl, wherein substituted arylalkyl, substituted arylhydroxyalkyl,

substituted heterocycloalkylalkyl and substituted heteroarylalkyl are substituted with R^{32} , R^{33} and R^{34} ;

R^{10} is $-O_m-(CR^{15}R^{16})_p-(CR^{17}R^{18})_q-(CR^{19}R^{20})_r-R^{21}$;

5 or R^9 and R^{10} together with the carbon atoms to which they are attached form a substituted cycloalkyl, a substituted heterocycloalkyl, a substituted aryl or a substituted heteroaryl, wherein substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl and substituted heteroaryl are substituted with R^{35} , R^{36} and R^{37} ;

R^{11} is H;

10 R^{15} , R^{17} and R^{19} are each independently selected from H, alkyl, cycloalkyl, haloalkyl and halocycloalkyl;

R^{16} , R^{18} and R^{20} are each independently selected from H, hydroxy, halogen and alkyl;

or R^{15} and R^{16} together with the carbon atom to which they are attached form a cycloalkyl;

15 or R^{17} and R^{18} together with the carbon atom to which they are attached form a cycloalkyl;

or R^{19} and R^{20} together with the carbon atom to which they are attached form a cycloalkyl;

or R^{15} and R^{17} together form $-(CH_2)_v-$;

20 or R^{15} and R^{19} together form $-(CH_2)_w-$;

or R^{17} and R^{19} together form $-(CH_2)_x-$;

25 R^{21} is H, halogen, cyano, $-OR^{25}$, $-SR^{25}$, $-S(O)R^{25}$, $-S(O)_2R^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}SO_2NR^{25}R^{27}$, $-NR^{26}C(O)R^{25}$, $-NR^{26}C(O)NR^{25}R^{27}$, $-C(O)R^{28}$, $-C(O)NR^{25}R^{26}$, cycloalkyl, substituted heterocycloalkyl, substituted heteroaryl or substituted aryl, wherein substituted heterocycloalkyl, substituted heteroaryl, substituted heteroarylalkyl and substituted aryl are substituted with R^{38} , R^{39} and R^{40} ;

- R²⁵ is H, alkyl, hydroxyalkyl, carboxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxyalkyl, cycloalkoxyalkyl, halocycloalkoxyalkyl, alkylcycloalkoxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, haloalkoxyalkoxyalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R⁴¹, R⁴² and R⁴³;
- R²⁶ and R²⁷ are each independently selected from H, alkyl, cycloalkyl, haloalkyl or halocycloalkyl;
- or R¹⁵ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;
- or R¹⁷ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;
- or R¹⁹ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;
- R²⁸ is H, hydroxy, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted

heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R^{50} , R^{51} and R^{52} ;

- 5 R^{12} , R^{13} , R^{14} , R^{22} , R^{23} , R^{24} , R^{29} , R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkylcarbonylamino, alkylsulfonyl, alkylsulfonylamino,
- 10 alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxycarbonyl, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl and heterocycloalkyl;
- 15 n is zero or 1;
- m zero or 1;
- p, q and r are independently selected from zero and 1;
- v and x are independently selected from 1, 2, 3 or 4;
- w is zero, 1, 2 or 3;
- 20 with the proviso that no more than two of A^2 , A^3 and A^4 are N;
- and pharmaceutically acceptable salts thereof.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl,

25 haloalkoxyalkyl, halocycloalkylalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl, wherein

substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl are substituted with R^{12} , R^{13} and R^{14} .

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^2 and R^4 together form a double bond, wherein
5 in case R^2 and R^4 together form a double bond, then R^5 is H.

In a further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 and R^2 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24}

10 Another further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^3 and R^4 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{29} , R^{30} and R^{31} .

15 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 and R^3 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{44} , R^{45} and R^{46} .

20 The present invention also relates to compounds according to formula (I) as described herein, wherein R^1 is H, alkyl or arylalkyl substituted with R^{12} , R^{13} and R^{14} .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is H or alkyl.

A more particular embodiment of the present invention are compounds according to
25 formula (I) as described herein, wherein R^1 is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^1 is methyl or ethyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl
5 or arylalkyl substituted with R^{12} , R^{13} and R^{14} .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 , R^2 , R^3 and R^4 are independently selected from H or alkyl.

A particular embodiment of the present invention are compounds according to
10 formula (I) as described herein, wherein R^1 is H, alkyl or arylalkyl substituted with R^{12} , R^{13} and R^{14} and wherein R^{12} , R^{13} and R^{14} are H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^2 is H or alkyl.

The present invention also relates to compounds according to formula (I) as
15 described herein, wherein R^2 is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^2 is alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^2 is methyl.

20 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 and R^2 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24} .

25 The present invention also relates to compounds according to formula (I) as described herein, wherein R^1 and R^2 together with the carbon atom to which they are attached form a cycloalkyl substituted with R^{22} , R^{23} and R^{24} .

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 and R^2 together with the carbon atom to which they are attached form a cyclopropyl.

Also an embodiment of the present invention are compounds according to formula
5 (I) as described herein, wherein R^3 is H or alkyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^3 is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^4 is H.

10 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^1 is CR^8 .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^1 is N.

A more particular embodiment of the present invention are compounds according to
15 formula (I) as described herein, wherein A^2 is CR^9 .

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^2 is N.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^3 is CR^{10} .

20 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^3 is N.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^4 is CR^{11} .

Also an embodiment of the present invention are compounds according to formula
25 (I) as described herein, wherein A^4 is N.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^5 is CR^6 .

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein A^5 is N.

- 5 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein one of R^5 , R^6 , R^7 and R^8 is selected from halogen, alkoxy and hydroxy and the others are each independently selected from H and halogen.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein one of R^5 , R^6 and R^8 are H.

- 10 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein one of R^5 , R^6 and R^8 are H and R^7 is halogen or alkoxy.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein at least one of R^5 , R^6 , R^7 and R^8 is different from H.

- 15 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^5 is H or halogen and wherein at least one of R^5 , R^6 , R^7 and R^8 is different from H.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^6 is H or halogen and wherein at least one of R^5 , R^6 , R^7 and R^8 is different from H.

- 20 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^7 is H, halogen, alkoxy or hydroxy and wherein at least one of R^5 , R^6 , R^7 and R^8 is different from H.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^7 is halogen or alkoxy.

- 25 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^7 is halogen.

Another particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁷ is chloro.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is H or halogen and wherein at least one of R^5 , R^6 , R^7 and R^8 is different from H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁹ is H, halogen, hydroxy, cyano, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, alkoxycycloalkylalkyl, dialkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl or substituted heteroarylalkyl, wherein substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl and substituted heteroarylalkyl are substituted with R³², R³³ and R³⁴.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁹ is H, halogen, cyano, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, alkoxyalkylalkyl, dialkoxyalkyl, substituted arylhydroxyalkyl or substituted heterocycloalkylalkyl, wherein substituted substituted arylhydroxyalkyl and substituted heterocycloalkylalkyl are substituted with R³², R³³ and R³⁴.

The present invention also relates to compounds according to formula (I) as described herein, wherein R⁹ is H or halogen.

A further embodiment of the present invention are compounds according to formula
 25 (I) as described herein, wherein R⁹ is H.

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁹ and R¹⁰ together with the carbon atoms to which they are attached form a substituted cycloalkyl, a substituted heterocycloalkyl, a substituted aryl or

a substituted heteroaryl, wherein substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl and substituted heteroaryl are substituted with R^{35} , R^{36} and R^{37} .

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 and R^{10} together with the carbon atoms to which they are
5 attached form a substituted cycloalkyl, a substituted aryl or a substituted heteroaryl, wherein substituted cycloalkyl, substituted aryl and substituted heteroaryl are substituted with R^{35} , R^{36} and R^{37} .

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 and R^{10} together with the carbon atoms to which they are
10 attached form a substituted cycloalkyl or a substituted aryl, wherein substituted cycloalkyl and substituted aryl are substituted with R^{35} , R^{36} and R^{37} .

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 and R^{10} together with the carbon atoms to which they are attached form a cycloalkyl substituted with R^{35} , R^{36} and R^{37} .

15 The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 and R^{10} together with the carbon atoms to which they are attached form a cycloalkyl substituted with R^{35} , R^{36} and R^{37} .

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{15} , R^{17} and R^{19} are each independently selected from H, alkyl,
20 cycloalkyl, haloalkyl or halocycloalkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{15} , R^{17} and R^{19} are each independently selected from H, alkyl, cycloalkyl and haloalkyl.

The present invention also relates to compounds according to formula (I) as
25 described herein, wherein R^{17} is H.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{19} is H, alkyl, cycloalkyl and haloalkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{19} is H or alkyl.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{19} is H.

- 5 The present invention also relates to compounds according to formula (I) as described herein, wherein R^{19} is alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{19} is methyl.

- 10 The present invention also relates to compounds according to formula (I) as described herein, wherein R^{16} , R^{18} and R^{20} are each independently selected from H, hydroxy, halogen or alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{16} , R^{18} and R^{20} are each independently selected from H, hydroxy and alkyl.

- 15 The present invention also relates to compounds according to formula (I) as described herein, wherein R^{18} is H.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{20} is H, hydroxy and alkyl.

- 20 A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{20} is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{19} and R^{20} together with the carbon atom to which they are attached form a cycloalkyl.

- 25 The present invention also relates to compounds according to formula (I) as described herein, wherein R^{19} and R^{20} together with the carbon atom to which they are attached form a cyclopropyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is H, halogen, cyano, $-OR^{25}$, $-SR^{25}$, $-S(O)R^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}C(O)R^{25}$, $-NR^{26}C(O)NR^{25}R^{27}$, $-C(O)R^{28}$, $-C(O)NR^{25}R^{26}$, cycloalkyl, substituted heterocycloalkyl, substituted heteroaryl or substituted aryl, wherein
5 substituted heterocycloalkyl, substituted heteroaryl and substituted aryl are substituted with R^{38} , R^{39} and R^{40} .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is H, halogen, $-OR^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}C(O)R^{25}$, substituted heterocycloalkyl or substituted heteroaryl,
10 wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{38} , R^{39} and R^{40} .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is H, halogen, $-OR^{25}$, $-NR^{25}R^{26}$, substituted heterocycloalkyl or substituted heteroaryl, wherein
15 substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{38} , R^{39} and R^{40} .

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is $-OR^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$ or $-NR^{26}C(O)R^{25}$.

20 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$ or $-NR^{26}C(O)R^{25}$.

An embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{21} is $-OR^{25}$.

25 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{25} is H, alkyl, hydroxyalkyl, carboxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted heterocycloalkyl, substituted

heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R^{41} , R^{42} and R^{43} .

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{25} is H, alkyl, haloalkyl, cycloalkyl, substituted
5 heteroaryl or substituted heteroarylalkyl, wherein substituted heteroaryl and substituted heteroarylalkyl are substituted with R^{41} , R^{42} and R^{43} .

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{21} is $-NR^{25}R^{26}$ and R^{25} is substituted heteroaryl or substituted heteroarylalkyl.

10 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{25} is H, alkyl, haloalkyl, cycloalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl or substituted heteroarylalkyl, wherein substituted heteroaryl and substituted heteroarylalkyl are substituted with R^{41} , R^{42} and R^{43} .

15 The present invention also relates to compounds according to formula (I) as described herein, wherein R^{26} and R^{27} are each independently selected from H, alkyl, cycloalkyl, haloalkyl or halocycloalkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{26} and R^{27} are each independently selected from H and
20 alkyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{26} is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{12} , R^{13} , R^{14} , R^{22} , R^{23} , R^{24} , R^{29} , R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} ,
25 R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, alkylcarbonyl, alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcarbonylamino, alkoxy carbonyl, alkoxyalkyl, haloalkoxy, chloropyridinylcarbonyl and heterocycloalkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{12} , R^{13} , R^{14} , R^{22} , R^{23} , R^{24} , R^{29} , R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcarbonylamino, alkoxy carbonyl, alkoxyalkyl, haloalkoxy and heterocycloalkyl.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{12} , R^{13} , R^{14} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, alkylcarbonyl, alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcarbonylamino, alkoxy carbonyl, alkoxyalkyl, haloalkoxy and heterocycloalkyl.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{12} , R^{13} , R^{14} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcarbonylamino, alkoxy carbonyl, alkoxyalkyl, haloalkoxy and heterocycloalkyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{12} , R^{13} and R^{14} are H.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{32} is alkyl or halogen, and R^{33} and R^{34} are H.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{35} is H, halogen, hydroxy, alkyl, haloalkyl or alkylcarbonylamino, and R^{36} and R^{37} are H.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{35} is H or alkyl, and R^{36} and R^{37} are H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{38} is H, halogen, hydroxy, cyano, oxo, alkyl,

alkylcarbonyl, alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkoxy carbonyl or alkoxyalkyl, R^{39} is H, halogen, oxo, alkyl or hydroxyalkyl, and R^{40} is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{38} is H, halogen, hydroxy, cyano, oxo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkoxy carbonyl or alkoxyalkyl, R^{39} is H, halogen, oxo, alkyl or hydroxyalkyl, and R^{40} is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{38} is H, alkyl alkylcarbonyl or alkylsulfonyl, and R^{39} and R^{40} is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{38} is H or alkyl, and R^{39} and R^{40} is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{41} is H, halogen, alkyl or haloalkyl, and R^{42} and R^{43} are H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{41} is alkyl, and R^{42} and R^{43} are H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{50} , R^{51} and R^{52} are each independently selected from H, halogen, hydroxy or alkyl, R^{51} is H or halogen, and R^{52} is H.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} and R^{43} are each independently selected from H and alkyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein p is zero.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein q is zero.


Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein r is zero or 1.

Another particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein m is zero.

- 5 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein n is zero.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein n is 1.

- A particular embodiment of the present invention are compounds according to
 10 formula (I) as described herein, wherein R² and R⁴ together form a double bond, R⁵ is H, n is 1 and of formula (Ia).



Particular examples of compounds of formula (I) as described herein are selected from

- 15 6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-nicotinonitrile;
 6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
 6-Chloro-2-(5-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
 6-Chloro-2-(5-fluoro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
 20 6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-
- 10 one;
- 6-Chloro-2-[5-((S)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 2-(5-Amino-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(2,2,2-trifluoro-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one
- 6-Chloro-2-[5-(cyclopropyl-hydroxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-
- 20 isoquinolin-1-one;
- 6-Chloro-2-[5-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-trifluoromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methoxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide;
- 6-Chloro-2-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-8'-hydroxy-3,4,5',6',7',8'-hexahydro-[2,4']biisoquinoliny-1-one;
- N-(6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;
- 6-Chloro-2-{5-[hydroxy-(1-methyl-1H-imidazol-2-yl)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-{5-[(3,4-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3,5-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(4-ethyl-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(hydroxy-phenyl-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-1-phenyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-{5-[1-(3,4-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(3,5-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(6-methyl-pyrazin-2-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(morpholine-4-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N,N-dimethyl-nicotinamide;
- 6-Chloro-2-[5-(pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-methyl-nicotinamide ;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-cyclopropyl-nicotinamide;
- 15 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-(4-fluoro-phenyl)-nicotinamide;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-phenyl-nicotinamide;
- 6-Chloro-2-[5-(4,4-difluoro-piperidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyridazin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-pyridin-3-yl-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 2-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 10 one;
- 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methylsulfanyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(4-dimethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{4-[(4-fluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1-methyl-1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2-methyl-2H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-fluoro-4-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(5-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one;
- 3-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-isonicotinonitrile;
- 6-Chloro-2-(5-fluoro-4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(4-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(3,5-dimethyl-3H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 (S)-1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 15 (R)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- (S)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 8-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Methoxy-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5,6-Dichloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 20 2-Chloro-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 2-Methoxy-6-pyridin-3-yl-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-5-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-7-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-7-fluoro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-4,4-dimethyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-4-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-Chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 15 one;
- 5-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 20 5-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- Ethanesulfonic acid [5-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

- 5-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5 6-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-methyl-2-(4-methyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 10 3-Benzyl-5-chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-methoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3,3-dimethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 15 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 20 one;
- 5-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- one;

- 5-Chloro-3,3-dimethyl-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 5-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 2-Hydroxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-2-(5-imidazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-ethyl-4-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 Propane-2-sulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3,5-dimethyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-acetamide
- 15 6-Chloro-2-(5-morpholin-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(1,1-dioxo-1 λ 6-thiomorpholin-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-propyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-1H-imidazole-2-carboxylic acid ethyl ester;
- 5 6-Chloro-2-[5-(2-hydroxymethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(oxetan-3-ylaminomethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-isopropyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(4,4-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,3-difluoro-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.3]hept-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(3,3-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-[1,2,3]triazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(5-[1,2,3]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-chloro-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3-methyl-[1,2,4]triazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(5-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-chloro-2-[5-(3-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(5-[1,2,4]triazol-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-methyl-benzoimidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-indazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-indazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(6-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(7-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(4-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-cyclopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2-trifluoromethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-ethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 2-(5-Aminomethyl-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methyl-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[2-(1-methyl-pyrrolidin-2-yl)-ethoxymethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopentyloxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-(5-cyclopropylmethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-fluoro-phenoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1-methyl-cyclopropylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(tetrahydro-furan-2-ylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,2,2-trifluoro-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-cyclobutoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-methanesulfonyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(6-methyl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(6-morpholin-4-yl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,3-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

25 6-Chloro-2-[5-(3,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(2,6-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-chloro-6-fluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3,4-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,5-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;
- 15 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-benzenesulfonamide;
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-methanesulfonamide;
- 20 Cyclopropanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;
- 6-Chloro-2-[5-(4-fluoro-benzylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-ethylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-(5-morpholin-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-propionamide;
- 6-Chloro-2-{5-[(2-methyl-2H-pyrazol-3-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 5 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylamino]-2-methyl-propionic acid;
- 6-Chloro-2-{5-[(1-methyl-1H-imidazol-4-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,4-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(3,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-chloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(3-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-trifluoromethoxy-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-chloro-3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(3,4-dichloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isoxazol-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(1-methyl-1H-imidazol-2-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,4-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-1-methyl-1H-pyrazole-4-carbonitrile
- 15 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-isobutyramide
- Cyclopropanecarboxylic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-4-fluorobenzamide
- 20 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-cyclohexylurea
- 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-(3-trifluoromethyl-phenyl)-urea
- 25 6-Chloro-2-(5-hydroxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetamide;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N-methyl-acetamide;
- 5 [5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid methyl ester;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N,N-dimethyl-acetamide;
- 6-Chloro-2-(5-phenylaminomethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[(4-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(4-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-{5-[(3-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-morpholin-4-yl-2-oxo-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-N-(2-hydroxy-ethyl)-acetamide;
- 6-Chloro-2-[5-(1-methylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1-dimethylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-1H-imidazole-2-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-[1,2,3]triazol-2-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(1-imidazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-pyrazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(oxazol-2-ylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(1-[1,2,4]triazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(2-oxo-pyrrolidin-1-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[1-(2-oxo-oxazolidin-3-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- N-{1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-ethyl}-methanesulfonamide;
- 6-Chloro-2-{5-[1-(3-fluoro-phenylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(1-phenylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-methanesulfinyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[4-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

and pharmaceutically acceptable salts thereof.

- 5 Also particular examples of compounds of formula (I) as described herein are selected from

5-Chloro-3,3-dimethyl-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one;

- 10 5-Chloro-3-ethyl-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(4-chloro-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(4-chloro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

6-Chloro-5'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

6-Chloro-8'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

- 15 8'-Amino-6-chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

Ethanesulfonic acid (6-chloro-1-oxo-3,4-dihydro-1H-[2,4']biisoquinoliny-8'-yl)-amide;

6'-Chloro-2'-(5-fluoropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

- 20 6'-Chloro-2'-[5-(difluoromethoxy)pyridin-3-yl]spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

2-Chloro-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

- 2-Chloro-6-(5-difluoromethoxy-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 6-(5-Fluoro-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 5 6-(5-Difluoromethoxy-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 6'-Chloro-2'-(pyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;
- 5-Chloro-3-cyclopropyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 2-Chloro-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 10 2-Ethoxy-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 2-Methoxy-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 5-Chloro-3-cyclopropyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-cyclopropyl-2-(5-difluoromethoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 15 one;
- 6-(5-Difluoromethoxy-pyridin-3-yl)-2-ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;
- 5-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 6'-Chloro-2'-(4-chloropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;
- 20 5-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- (S or R)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- (R or S)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- (R or S)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

(S or R)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one;

5 N-[(R or S)-4-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

10 Ethanesulfonic acid [4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

15 Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

20 N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinolinyl-8'-yl)-acetamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinolinyl-8'-yl)-acetamide;

25 N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide;

5 N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide;

N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

10 N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;

15 N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

20 N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

25 N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide;

Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide;

5 N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

10 5-Chloro-2-((S or R)-8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;

5-Chloro-2-((R or S)-8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

15 N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

20 N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[4-(5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

25 N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

5 N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

10 N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

15 N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

20 N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

25 N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

5 N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

10 N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

15 N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

20 N-[(S or R)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

25 N-[(R or S)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5 N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

10 5-Chloro-3,3-dimethyl-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-2,3-dihydro-isoindol-1-one;

15 2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;

2-Chloro-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

20 2-Methoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

2-Ethoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

6'-Chloro-2'-{5-[(1H-pyrazol-3-ylamino)methyl]pyridin-3-yl}spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

25 Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(6-fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(6-cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

5 Ethanesulfonic acid [5-((S or R)-5-chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

10 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-N-methyl-methanesulfonamide;

N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl} ethanesulfonamide;

N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}-N-methylethanesulfonamide;

15 Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methyl-amide;

N-{[5-(6'-Chloro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}propanamide;

20 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

N-[5-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

25 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-acetamide;

N-{[5-(5'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl)methyl}methanesulfonamide;

N-[5-(6-Cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

5 N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl)methyl}methanesulfonamide;

10 N-[5-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-(5-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

15 N-[5-(5-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

5-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 5-Chloro-3,3-dimethyl-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

25 5-Chloro-3,3-dimethyl-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 5-Chloro-2-[5-(3-isopropyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

6-Chloro-2-[5-(1,5-dimethyl-1H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 5-Chloro-3,3-dimethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

6-Chloro-2-[5-(3-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,5-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(1,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[4-chloro-5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

2-Chloro-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

25 2-Methoxy-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

5-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

(R or S)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5 (S or R)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

10 3-Chloro-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

1-Methyl-1H-imidazole-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

2-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide;

15 Pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

3-Methyl-3H-imidazole-4-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

20 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-6-methyl-nicotinamide;

3-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-isonicotinamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide;

25 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-2-methyl-nicotinamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-4-methyl-nicotinamide;

2-[5-(1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 2-[5-((R)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

10 5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-methanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 5-Chloro-2-[5-((R)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((S)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 2-[5-(1-Acetyl-piperidin-4-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(1-Acetyl-azetidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

25 5-Chloro-2-[5-(1-methanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(4-propionyl-piperazin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

10 5-Chloro-2-[5-(4-methanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(4-ethanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-{5-[4-(3-chloro-pyridine-2-carbonyl)-piperazin-1-yl]-pyridin-3-yl}-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 2-[5-(1-Acetyl-pyrrolidin-3-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one ;

20 2-[6-(1-Acetyl-piperidin-3-yl)-pyrazin-2-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[6-(1-propionyl-piperidin-3-yl)-pyrazin-2-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[6-(1-ethanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

25 5-Chloro-2-[6-(1-methanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

N-[(S or R)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

- 5 N-[4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

and pharmaceutically acceptable salts thereof.

Further particular examples of compounds of formula (I) as described herein are selected from

- 10 6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 20 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-{5-[(2-methyl-2H-pyrazol-3-ylmethyl)-amino]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

and pharmaceutically acceptable salts thereof.

10 Also further particular examples of compounds of formula (I) as described herein are selected from

(R or S)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

(S or R)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

15 N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

20 N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

25 N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5-Chloro-3,3-dimethyl-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-2,3-dihydro-isoindol-1-one;

6'-Chloro-2'-{5-[(1H-pyrazol-3-ylamino)methyl]pyridin-3-yl}spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

5 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

10 2-[5-((R)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-methanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 5-Chloro-2-[5-((R)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((S)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 2-[5-(1-Acetyl-piperidin-4-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

25 5-Chloro-2-[5-(4-methanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(4-ethanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 and pharmaceutically acceptable salts thereof.

Processes for the manufacture of compounds of formula (I) as described herein are an object of the invention.

The preparation of compounds of formula (I) of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following general schemes. The skills required for carrying out the reaction and purification of the resulting products are known to those persons skilled in the art. In case a mixture of enantiomers or diastereoisomers is produced during a reaction, these enantiomers or diastereoisomers can be separated by methods described herein or known to the man skilled in the art such as e.g. chiral chromatography or crystallization. The substituents and indices used in the following description of the processes have the significance given herein.


The following abbreviations are used in the present text:

AcOH = acetic acid, BOC = t-butyloxycarbonyl, BuLi = butyllithium, CDI= 1,1-carbonyldiimidazole, DCM = dichloromethane, DBU = 2,3,4,6,7,8,9,10-octahydro-pyrimido[1,2-a]azepine, DCE = 1,2-dichloroethane, DIBALH = di-*i*-butylaluminium hydride, DCC = N,N'-dicyclohexylcarbodiimide, DMA = *N,N*-dimethylacetamide, DMAP = 4-dimethylaminopyridine, DMF = *N,N*-dimethylformamide, EDCI = *N*-(3-dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride, EtOAc = ethylacetate, EtOH = ethanol, Et₂O = diethylether, Et₃N = triethylamine, eq = equivalents, HATU = O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, HPLC = high performance liquid chromatography, HOBT = 1-hydroxybenzo-triazole, Huenig's base = *i*Pr₂NEt = *N*-ethyl diisopropylamine, IPC= in process control, LAH = lithium aluminium hydride, LDA = lithium diisopropylamide, LiBH₄ = lithium borohydride, MeOH = methanol, NaBH₃CN, sodium cyanoborohydride, NaBH₄ = sodium borohydride, NaI =

sodium iodide, Red-Al = sodium bis(2-methoxyethoxy) aluminium hydride, RT = room temperature, TBDMSCl = t-butyldimethylsilyl chloride, TFA = trifluoroacetic acid, THF = tetrahydrofuran, quant = quantitative.

Halogen or triflate substituted heterocyclic compounds **2** react with aryl lactams **1** in
 5 solvents like 1,4-dioxane, in the presence of copper (I) iodide, potassium or cesium carbonate, a chelating 1,2-diamino compound like *N,N'*-dimethylethylenediamine or trans-1,2-diamino-hexane, at elevated temperatures, preferable with the aid of microwave heating to form lactam substituted heterocyclic compounds **3** (step a) as described in Scheme 1.

Scheme 1





X is halogen or OSO₂CF₃

10

Treatment of heteroaryl linked alcohol compounds **51** (Scheme 2b) with a base like sodium hydride in a solvent like THF or DMF and subsequently with a suitable alkylating agent such as a halide, mesylate or tosylate preferably between RT and the reflux
 15 temperature of the solvent gives compounds **52** (step a). Alternatively (Scheme 2a), heteroaryl linked alcohol compounds **51** can *e.g.* be converted into the corresponding chlorides by treatment with thionyl chloride in a solvent like dichloromethane preferably between 0 °C and room temperature (step b). Said heteroaryl linked chlorides, compounds **53**, react with nucleophilic amino-moieties **54a** or aryl, heteroaryl or heterocycloalkyl
 20 compounds **54b** per se or after anion formation *e.g.* with sodium hydride in solvents like


N,N'-dimethylformamide in a temperature range between 0 °C and about 100 °C to form adducts **55a** or **55b** (step c).

Scheme 2a




is heteroaryl or heterocyclyl substituted by R^{38} , R^{39} , R^{40} .

Scheme 2b




Alternatively (Scheme 2c), heteroaryl linked chloride compounds **53** react with boronic acids or esters **56 i**) by using *Suzuki* conditions, *e.g.* in the presence of catalysts, such as *tri*-*o*-tolylphosphine/palladium(II)acetate, *tetrakis*-(triphenylphosphine)-palladium, *bis*(triphenylphosphine)palladium(II)chloride or dichloro[1,1'-*bis*(diphenylphosphino)-ferrocene]palladium(II) optionally in the form of a dichloromethane complex (1:1), and in the presence of a base, such as aqueous or non aqueous potassium phosphate, sodium or potassium carbonate, in a solvent, such as dimethylsulfoxide, toluene, ethanol, dioxane, tetrahydrofuran or *N,N*-dimethylformamide, and in an inert atmosphere such as argon or nitrogen, in a temperature range preferably between room temperature and about 130 °C or ii) by using a nickel (0) catalyst, *e.g.* *bis*(1,5-cyclooctadiene)nickel (0) in the presence potassium phosphate, *bis*(1-methyl-1*H*-imidazol-2-yl)methane in *N,N*-dimethyl acetamide at temperatures around 100 °C giving adducts **57** (steps d).

Scheme 2c




R¹⁰¹ and R¹⁰² are H or alkyl, or R¹⁰¹ and R¹⁰² together with the boron atom to which they are attached form



R²¹ is substituted aryl or substituted heteroaryl

Aldehydes or ketones **58** (Scheme 2d) can be treated with suitable amino-moieties **59** in the presence of NaBH(OAc)₃ in a one step procedure in a solvent like methanol preferably around room temperature or in a two step procedure by first treatment with titanium (IV) isopropoxide in solvents like methanol or toluene preferably at temperatures between room temperature and the reflux temperature of the solvents followed by reaction with NaBH₄ preferably between 0 °C and room temperature which converts aldehydes or ketones **58** into amino compounds **60**; alternatively imines obtained after treatment with titanium (IV) isopropoxide can be evaporated, then be re-dissolved in a solvent like THF and being treated with a Grignard reagent R²⁰MgX, preferably between -40 °C and 0 °C leading to amino compounds **60** carrying the specific R²⁰ substituent (step e).


Scheme 2d




R^{104} is R^{25} , $-S(O)_2R^{25}$, $-C(O)R^{25}$ or $-C(O)NR^{25}R^{27}$

Heteroaryl halides **101** (Scheme 2e) react with boronic acids or esters **102** using *Suzuki* conditions as described above to give adducts **103** (step a). Phenols **104** react with alcohols **105** under *Mitsunobu* conditions *e.g.* with triphenylphosphine and di-*tert*-butyl-, diisopropyl-, diethyl-azodicarboxylate or di-(4-chlorobenzyl)azodicarboxylate as reagents in solvents like toluene, dichloromethane or tetrahydrofuran preferably at ambient temperature to give adducts **106** (step b). Compounds **106** can be transformed by additional standard modifications into synthons **107** (step c).

Scheme 2e



R^{101} and R^{102} are H or alkyl, or R^{101} and R^{102} together with the boron atom to which they are attached form



R^{21} is substituted aryl or substituted heteroaryl


X is halogen or OSO_2CF_3

R^{109} represents substituents as defined in R^{21} or substituents which can easily be transformed into R^{21} (step c).


Carbamates **201** (Scheme 3) react with polyphosphoric acid at elevated temperature (*e.g.* 100-180 °C) to form 3,4-dihydro-2*H*-isoquinolin-1-one derivatives **202** (step a). Trifluoroacetamides **203** can be cyclized to 1-(3,4-dihydro-1*H*-isoquinolin-2-yl)-2,2,2-trifluoro-ethanone compounds **204** by treatment with paraformaldehyde in a mixture of concentrated sulfuric acid and acetic acid preferably around room temperature (step b). Removal of the trifluoroacetyl group by treatment with *e.g.* potassium hydroxide in a solvent like ethanol at temperatures around room temperature gives tetrahydro-isoquinoline compounds **205** (step c). Oxidation of tetrahydro-isoquinoline compounds **205** *e.g.* with iodoso benzene and potassium bromide preferably in water gives 3,4-dihydro-2*H*-isoquinolin-1-one compounds **202** (step d). Reaction of isoindole-1,3-dione compounds **206** with a Grignard reagent R¹MgX in a solvent like THF preferably around 0 °C gives adducts **207** (step e). Subsequent treatment with triethylsilane and trifluoroboron etherate in a solvent like dichloromethane and in a temperature range preferably between -25 °C and RT gives isoindolone compounds **208** (step f). Introduction a methoxybenzyl protecting group into isoindolone compounds **209** (*e.g.* by treatment with sodium bis(trimethylsilyl) amide and 1-bromomethyl-4-methoxy-benzene in THF between 0 °C and RT) gives protected compounds **210** (step g); similarly, a methoxybenzyl protecting group can be introduced into compounds **208**. Treatment of compounds **208** carrying an additional methoxybenzyl protecting group or compounds **210** with a base like sodium hydride in a solvent like THF and then with an alkyl halide, mesylate or tosylate preferably between RT and the reflux temperature of the solvent gives compounds **211** with structurally different or structurally identical R¹ and R² groups (step h). Alternatively, treatment of compounds **208** carrying an additional methoxybenzyl protecting group or compounds **210** with a base like NaH, LDA or LiHMDS in solvents like DMF, tetrahydrofuran or 1,2-dimethoxyethane and then with one or sequentially with two different alkyl halides, mesylates or tosylates preferably between -78 °C and the reflux temperature of the solvent gives compounds **211** with structurally different or structurally identical R¹ and R² groups (step h). Removal of the protecting group, *e.g.* by treatment with trifluoroacetic acid at elevated temperature gives isoindolone compounds **212** (step i). Treatment of compounds **210** with a base like sodium hydride in a solvent like THF and then with an alpha, omega di-haloalkane or di-haloheteroalkane like *e.g.* 1,2-dibromoethane, preferably between RT and the reflux temperature of the solvent gives

spiro compounds **213** (step k) and after subsequent removal of the protecting group, spiro compounds **214** (step l).

Scheme 3



- 81a -




D is a substituted cycloalkyl or a substituted heterocycloalkyl formed by R^1 and R^2 together with the carbon atom to which they are attached, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24}

5-Halo-nicotinic acid or pyridazine carboxylic acid compounds **301** or **305** (Scheme 4a) react with acrylic acid ester compounds **302** after deprotonation with base like LDA or LiHMDS in solvents like THF preferably around -78 °C giving cyclic beta keto ester


compounds **303** and **306** (step a). Ester compounds **303** or **306** with $R^{111} = H$ can be treated with a base like NaH, LDA or LiHMDS in solvents like DMF, tetrahydrofuran or 1,2-dimethoxyethane, followed by addition of an alkyl or cycloalkyl halide, mesylate or tosylate, or *e.g.* a *N*-halobenzenesulfonamide, a reaction preferably performed between -78 °C and room temperature, to give ester compounds **303** or **306** carrying a substituent R^{111} different from H. Treatment of beta keto-ester compounds **303** or **306** with aqueous acid preferably at reflux induces ester hydrolysis and subsequent decarboxylation providing ketones **304** and **307** (step b). Ketones **304** and **307** with $R^{111} = H$ and $R^{113} = H$ can be treated with a base like NaH, LDA or LiHMDS in solvents like DMF, tetrahydrofuran or 1,2-dimethoxyethane, followed by addition of one or subsequently two different alkyl or cycloalkyl halides, mesylates or tosylates, or *e.g.* *N*-halobenzenesulfonamides, a reaction preferably performed between -78 °C and room temperature, to give ketones **304** and **307** carrying at least one of the substituents R^{111} or R^{113} different from H. Optionally, ketones **304** and **307** can be converted into the corresponding imines (*e.g.* with *N*-butylamine by using a catalyst like toluene sulfonic acid or pyridinium *p*-toluenesulfonate in a solvent like ethanol preferably at reflux); such imines with $R^{111} = H$ can be reacted with *e.g.* *N*-fluorobenzenesulfonimide using K_2CO_3 or triethylamine as base in solvents like DMF or acetonitrile or mixtures thereof, in the presence of molecular sieves preferably at room temperature to give imines carrying fluoro substituents and after imine hydrolysis (*e.g.* with hydrochloric acid in acetonitrile) ketones **304** and **307** with $R^{111} = F$. Ketones **304** and **307**, can function as compounds **2** (scheme 1) or can be converted into different compounds **2** (Scheme 1) by further structural modification using methods well known in the art as *e.g.* reduction of the keto function to a secondary hydroxy group with a reagent like sodium borohydride in methanol or reduction of the keto group to a primary or secondary amino function by reductive amination *e.g.* by reacting with an amine and $NaBH(OAc)_3$ in methanol.

Scheme 4a



Dihalopyridine or pyridazine compounds **351** (Scheme 4b) react with bromo-tetramethyl-azadisilolidine reagents **352** after deprotonation with lithium diisopropyl amide in solvents like tetrahydrofuran between -78 °C and 0 °C to give aminoalkyl substituted pyridines or pyridazine **353** (step a). After attachment of a protecting group onto compounds **353** (e.g. by introducing a BOC- or a SES-protecting group by reaction with BOC₂O or 2-trimethylsilyl-ethanesulfonyl chloride, triethylamine, DMF around 0 °C), treatment of amino protected pyridine or pyridazine compounds **354** with a base like potassium carbonate, in a solvent like toluene, and in the presence of a catalyst like *tetrakis*-(triphenylphosphine)-palladium at temperatures around 100 °C gives bicyclic compounds **355**, still carrying a protecting group (step b, c). Standard BOC removal or use of tetrabutylammonium fluoride hydrate, acetonitrile preferably between room temperature and the reflux temperature of acetonitrile then gives bicyclic compounds **356** (step d). Amino compounds **356** can function as compounds **2** (Scheme 1) either as such or after further structural modification by methods well known in the art.

Scheme 4b



prot. = protecting group


X is halogen or OSO₂CF₃R¹¹⁰ is R³⁵, R³⁶ or R³⁷.R¹¹¹ is R³⁵, R³⁶ or R³⁷.R¹¹² is R³⁵, R³⁶ or R³⁷.

5-Halo-nicotinic acid compounds **305** (Scheme 4c) react with alkene compounds **400** after deprotonation with a base like LDA or LiHMDS in solvents like THF preferably around -78 °C giving alkenes **401** (step a). Diester compounds **402** can be synthesized by methods known to persons skilled in the art such as *e.g.* by ozonolysis of alkenes **401** in the presence of methanolic NaOH to give compounds **402** which can be cyclized using Dieckmann condensation conditions to give beta keto-ester compounds **403** (step b, c). Treatment of beta keto-ester compounds **403** with aqueous acid preferably at reflux temperature induces ester

hydrolysis and subsequent decarboxylation providing ketones **404** (step e). Ester compounds **403** (R^{111} is H) can be treated with a base like NaH, LDA or LiHMDS in solvents like DMF, tetrahydrofuran or 1,2-dimethoxyethane, followed by addition of an alkyl or cycloalkyl halide, mesylate or tosylate, or *e.g.* a *N*-halobenzenesulfonamide, a reaction preferably performed between -78 °C and room temperature, to give ester compounds **403** carrying a substituent R^{111} different from H

(step d). Hydrolyses and decarboxylation as described above gives ketones **404** (step e). Ketone compounds **404** can function as compounds **2** (Scheme 1) either as such or after further structural modification by methods described herein or by methods well known in the art.

Scheme 4c



X is halogen or OSO_2CF_3

X^1 is Halogen, Mesylate or Tosylate


R^{110} is H, R^{35} , R^{36} or R^{37} .

R^{111} is H, R^{35} , R^{36} or R^{37} .

R^{112} is H, R^{35} , R^{36} or R^{37} .

R^{113} is H, R^{35} , R^{36} or R^{37} .

Also an embodiment of the present invention is a process to prepare a compound of formula (I) as defined above comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



wherein R¹, R², R³, R⁴, R⁵, R⁶, A¹, A², A³, A⁴, A⁵ and n are as defined above and wherein, X is halogen or triflate.

In particular, in the presence of copper (I) iodide, potassium or cesium carbonate, a
 5 chelating 1,2-diamino compound like *N,N'*-dimethylethylenediamine or trans-1,2-diamino-hexane, at elevated temperatures, preferable with the aid of microwave heating and in solvents like 1,4-dioxane..

Also an object of the present invention is a compound according to formula (I) as described herein for use as therapeutically active substance.

10 Likewise an object of the present invention is a pharmaceutical composition comprising a compound according to formula (I) as described herein and a therapeutically inert carrier.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease, congestive
 15 heart failure, hypertension, primary aldosteronism and Cushing syndrom.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of diabetic nephropathy.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of kidney or heart fibrosis.

20 The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of congestive heart failure.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of hypertension.

- 5 The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of primary aldosteronism.

A particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

- 10 Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of diabetic nephropathy.

Another particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of kidney or heart fibrosis.

- 15 Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease.

Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of congestive heart failure.

Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of hypertension.

- 20 Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of primary aldosteronism.

- The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and
25 Cushing syndrom.

The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of diabetic nephropathy.

5 The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of kidney or heart fibrosis.

Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease.

10 Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of congestive heart failure.

Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or
15 prophylaxis of hypertension.

Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of primary aldosteronism.

Also an object of the invention is a method for the treatment or prophylaxis of
20 chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an object of the invention is a method for the treatment or prophylaxis of diabetic nephropathy, which method comprises administering an effective amount of a
25 compound according to formula (I) as described herein.

Also an object of the invention is a method for the treatment or prophylaxis of kidney or heart fibrosis, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or prophylaxis of chronic kidney disease, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or
5 prophylaxis of congestive heart failure, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or prophylaxis of hypertension, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

10 Also an embodiment of the present invention is a method for the treatment or prophylaxis of primary aldosteronism, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a compound of formula (I) as described herein, when manufactured according to any one of the described processes.

15 Assay procedures

Herein we identified the use of the G-402 cell line as a host cell to ectopically express (transiently or stably) enzymes of the CYP11 family. Specifically we developed stable G-402 cells expressing ectopically human CYP11B1, human CYP11B2, human CYP11A1, cynmolgus CYP11B1 or cynomolgus CYP11B2 enzyme activity. Importantly
20 the identified cell line G-402 expresses co-factors (adrenodoxin and adrenodoxin reductase) important for the activity of the CYP11 family and no relevant enzyme activity of the CYP11 family (in comparison to H295R cells) was detected in these cells. Therefore the G-402 cell line is uniquely suited as a host cell for the ectopic expression of enzymes from the CYP11 family.

25 G-402 cells can be obtained from ATCC (CRL-1440) and were originally derived from a renal leiomyoblastoma.

The expression plasmids contains the ORF for either human / cyno CYP11B1 or CYP11B2 under the control of a suitable promoter (CMV-promoter) and a suitable

resistance marker (neomycin). Using standard techniques the expression plasmid is transfected into G-402 cells and these cells are then selected for expressing the given resistance markers. Individual cell-clones are then selected and assessed for displaying the desired enzymatic activity using 11-Deoxycorticosterone (Cyp11B2) or 11-Deoxycortisol (Cyp11B1) as a substrate.

G-402 cells expressing CYP11 constructs were established as described above and maintained in McCoy's 5a Medium Modified, ATCC Catalog No. 30-2007 containing 10% FCS and 400 µg/ml G418 (Geneticin) at 37 °C under an atmosphere of 5% CO₂/95% air. Cellular enzyme assays were performed in DMEM/F12 medium containing 2.5 % charcoal treated FCS and appropriate concentration of substrate (0.3-10 µM 11-Deoxycorticosterone, 11-Deoxycortisol or Corticosterone). For assaying enzymatic activity, cells were plated onto 96 well plates and incubated for 16h. An aliquot of the supernatant is then transferred and analyzed for the concentration of the expected product (Aldosterone for CYP11B2; Cortisol for CYP11B1). The concentrations of these steroids can be determined using HTRF assays from CisBio analyzing either Aldosterone or Cortisol.

Inhibition of the release of produced steroids can be used as a measure of the respective enzyme inhibition by test compounds added during the cellular enzyme assay. The dose dependent inhibition of enzymatic activity by a compound is calculated by means of plotting added inhibitor concentrations (x-axes) vs. measured steroid/product level (y-axes). The inhibition is then calculated by fitting the following 4-parameter sigmoidal function (Morgan-Mercer-Flodin (MMF) model) to the raw data points using the least squares method:

$$y = \frac{AB + Cx^D}{B + x^D}$$

wherein, A is the maximum y value, B is the EC₅₀ factor determined using XLFit, C is the minimum y value and D is the slope value.

The maximum value A corresponds to the amount of steroid produced in the absence of an inhibitor, the value C corresponds to the amount of steroid detected when the enzyme is fully inhibited.

EC50 values for compounds claimed herein were tested with the G402-based assay system described. Cyp11B2 enzyme activity was tested in presence of 1 μ M Deoxycorticosterone and variable amounts of inhibitors; Cyp11B1 enzyme activity was tested in presence of 1 μ M Deoxycortisol and variable amounts of inhibitors.

5

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
1	0.0860	4.6072
2	0.4421	7.3919
3	0.0965	3.7211
4	0.0671	1.2377
5	0.0925	4.2705
6	0.0300	1.2079
7	0.0779	0.7042

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
8	0.0449	0.8072
9	0.1170	4.7416
10	0.0260	0.8220
11	0.0127	0.2566
12	0.0357	0.1177
13	0.0543	0.8652
14	0.0550	1.0961

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
15	0.0682	0.9130
16	0.0193	0.1262
17	0.5967	14.0208
18	0.0586	0.5071
19	0.0603	0.1984
20	0.0859	1.0495
21	0.0415	0.2632
22	0.0557	0.5168
23	0.0640	0.5714

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
24	0.0056	0.0214
25	0.4010	
26	0.3214	8.1518
27	0.0753	0.7309
28	0.1035	0.7566
29	0.3184	12.9186
30	0.7512	2.0686
31	0.0502	0.3283
32	0.0468	0.5139

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
33	0.2100	0.4847
34	0.0316	0.2873
35	0.0188	0.1462
36	0.0140	0.1881
37	0.0092	0.1494
38	1.0661	8.3906
39	1.0940	10.0398
40	1.6796	10.4624
41	3.7521	6.0795

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
42	0.3269	7.8639
43	0.3724	
44	5.9322	16.9135
45	0.2953	
46	0.6978	13.8752
47	1.0252	23.2395
48	0.1413	1.1052
49	0.0496	0.1880
50	1.5035	

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
51	2.6789	
52	0.3822	2.4620
53	0.1204	10.7585
54	0.2121	
55	0.3429	9.8504
56	0.2968	13.7841
57	1.2847	7.9991
58	2.0592	
59	0.0081	0.1354

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
60	0.0707	0.7938
61	2.8309	
62	0.9355	
63	0.4479	4.4257
64	0.1268	6.3126
65	0.0973	2.3468
66	0.0069	0.0697
67	0.1021	1.2006
68	0.0067	0.1566

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
69	0.1537	7.5432
70	0.4595	0.3919
71	0.0902	1.5795
72	0.5883	16.4938
73	0.0135	0.2514
74	0.4490	
75	0.1657	2.7583
76	0.4921	13.8461
77	0.6653	12.7918

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
78	0.7612	1.4503
79	0.0051	0.0340
80	0.0047	0.0581
81	0.1781	2.3633
82	0.0789	0.5968
83	0.5896	15.7133
84	0.2251	4.8264
85	0.0596	2.8554
86	0.3767	7.7744

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
87	0.2507	9.3698
88	0.7614	
89	0.2715	16.8487
90	1.2312	
91	0.3949	
92	0.2682	11.3701
93	0.7752	14.0060
94	2.0631	
95	0.3272	10.2668

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
96	0.0458	0.5326
97	0.0169	0.2202
98	0.2274	6.8266
99	0.9008	7.2722
100	0.0172	0.4853
101	0.1042	3.5415
102	1.5376	
103	0.5639	10.9865
104	2.3394	59.7842

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
105	0.3817	4.4881
106	0.5158	8.5638
107	0.0635	0.7696
108	0.1626	9.7819
109	0.2288	1.3489
110	0.1845	1.2775
111	0.0642	0.9228
112	0.6742	11.9975
113	0.3419	1.3593

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
114	0.4240	12.7810
115	0.6825	13.5418
116	0.1322	5.1321
117	0.1267	0.8941
118	0.1229	1.2484
119	0.7219	
120	0.0247	0.2073
121	0.0071	0.0390
122	0.0423	0.1802

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
123	0.1220	7.7317
124	1.0695	0.2249
125	0.1213	3.3410
126	0.0379	3.5625
127	0.0545	0.2657
128	0.1077	4.2761
129	0.0636	3.1638
130	0.0155	0.2398
131	0.0119	0.2635

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
132	0.0378	0.5266
133	0.0986	1.3716
134	0.0186	0.4243
135	0.0931	0.5232
136	0.1022	2.6575
137	0.0101	0.1082
138	0.0162	0.1928
139	0.9532	11.1720
140	0.0118	0.0704

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
141	0.0405	1.3088
142	0.0802	0.9809
143	0.2579	4.4495
144	0.0237	0.1400
145	1.4972	1.8956
146	0.0398	0.1142
147	0.0367	0.2470
148	0.0451	0.1391
149	0.1262	1.9169

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
150	0.2266	5.3241
151	0.1005	3.9244
152	0.2946	11.5673
153	0.3021	14.9952
154	0.0747	0.1929
155	0.1315	4.0942
156	0.0439	0.3369
157	0.3520	11.1514
158	0.1823	6.0298

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
159	0.2090	4.4617
160	0.0361	0.4182
161	0.1014	0.3410
162	0.1866	2.0029
163	0.2155	3.3596
164	0.1409	0.8618
165	0.0035	0.0604
166	0.0141	0.0963
167	0.0479	0.1681

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
168	0.0034	0.0475
169	0.0052	0.0381
170	0.0035	0.0146
171	0.0172	0.0889
172	0.0223	0.1441
173	0.3340	12.2938
174	0.0258	0.2112
175	0.1051	0.6564
176	0.0098	0.0578

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
177	0.4116	6.3108
178	0.0542	0.4797
179	0.6196	7.7205
180	0.1233	0.8017
181	0.2168	1.9663
182	0.0275	0.3060
183	0.0299	0.3853
184	0.0177	0.2712
185	0.0660	0.6732

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
186	0.0232	0.2107
187	0.1368	2.2324
188	0.2275	4.1486
189	0.0553	0.7099
190	0.3279	7.3219
191	0.3944	8.5507
192	2.2023	10.3297
193	0.0201	0.2168
194	0.0203	0.2243

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
195	0.0373	0.4865
196	0.1775	1.0297
197	0.0468	0.5554
198	2.4539	0.9902
199	0.1037	0.2818
200	0.0325	0.3146
201	0.0304	0.2932
202	0.0653	0.4734
203	0.0224	0.2123

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
204	0.0408	0.9002
205	0.0005	0.0006
206	0.0017	0.0068
207	0.0028	0.0182
208	0.0021	0.0189
209	0.0009	0.0147
210	0.0186	0.1298
211	0.0044	0.0532
212	0.0084	0.0816

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
213	0.0391	0.2781
214	0.0636	0.3003
215	0.0296	0.1368
216	0.0381	0.3639
217	0.0920	0.8026
218	1.5500	14.2551
219	2.2159	17.5743
220	1.0754	
221	2.1615	10.1235

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
222	0.0803	0.3426
223	0.1155	1.4171
224	0.0372	0.2932
225	2.5506	
226	0.0534	3.4026
227	4.2089	
228	0.2634	12.3727
229	0.2038	0.7447
230	0.0070	0.1495

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
231	0.0233	0.4686
232	0.0266	1.3509
233	0.2705	21.5699
234	0.0445	1.5779
235	0.0227	0.2211
236	0.1334	4.3865
237	0.2044	3.6068
238	0.1044	5.9872
239	0.0105	0.0900

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
240	0.1148	3.2729
241	0.5820	
242	0.0256	0.2062
243	0.2678	3.3424
244	1.8627	4.6905
245	0.0011	0.0257
246	0.1023	0.5463
247	4.5758	7.4708
248		2.6463

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
249	3.2181	7.6048
250	4.4379	12.3402
251	5.8839	15.8337
252	0.2721	9.7643
253	0.5808	10.0383
254	2.1317	32.5224
255	2.8003	7.6090
256	0.7744	

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
257	0.0113	0.0954
258	0.0037	0.0418
259	0.0122	0.0738
260	0.4711	0.3450
261	0.0267	0.1052
262	0.0141	0.3002
263	0.4910	4.9268
264	0.8428	4.1868
265	0.4373	5.0201

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
266	0.0444	0.2756
267	1.6052	11.2653
268	1.6485	14.1758
269	0.0462	0.4253
270	0.0063	0.0279
271	0.0248	0.2512
272	0.0604	0.2099
273	0.0640	0.4487
274	0.1275	1.0511

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
275	0.7311	7.8164
276	0.2837	2.0297
277	0.0641	0.1634
278	0.0061	0.0344
279	0.6004	11.1051
280	1.7985	
281	0.0004	0.003
282	0.0429	0.4748

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
283	0.0667	0.6328
284	0.0205	0.2494
285	0.0076	0.3556
286	0.2655	3.0382
287	0.5546	6.4082
288	0.539	8.8668
289	0.0706	0.1856
290	0.1012	3.91
291	0.0032	0.1218

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
292	0.5152	36.2254
293	0.0164	0.2848
294	0.2042	9.8508
295	0.0488	0.8488
296	0.1316	4.4957
297	0.2072	12.784
298	0.1747	13.7042
299	0.7869	48.5358
300	0.2161	14.2793

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
301	0.1606	32.5099
302	0.0932	5.1857
303	0.0936	12.5809
304	0.0024	0.0781
305	0.0054	0.8598
306	0.0058	0.1138
307	0.3892	15.4452
308	0.3805	26.3474
309	0.016	2.4468

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
310	0.0165	3.9684
311	0.8134	6.4778
312	0.0482	4.008
313	0.1184	3.5436
314	0.0282	5.2254
315	1.0508	31.0217
316	0.0402	1.831
317	0.1088	2.6377
318	>30	

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
319	0.1089	5.1301
320	8.0313	
321	0.0556	3.995
322	5.1407	39.6231
323	0.0442	2.3486
324	2.8107	
325	0.0592	4.4874
326	1.8693	6.6127
327	1.3648	10.6987

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
328	0.1731	18.44
329	6.0154	11.9661
330	>30	
331	4.3466	35.0631
332	14.7777	
333	0.0958	5.2386
334	0.3831	34.5562
335	13.8179	
336	0.9489	29.9696

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
337	24.2997	
338	>30	
339	0.0545	7.9851
340	0.0883	14.1639
341	0.0321	1.6765
342	0.0151	0.0947
343	0.1338	4.688
344	0.0312	1.5032
345	0.0494	2.003

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
346	0.0263	2.571
347	0.162	22.9422
348	2.8265	36.9471
349	0.9822	15.1179
350	0.0594	3.6822
351	3.6617	
352	0.0965	39.4185
353	0.3963	16.1819
354	0.0122	0.3454

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
355	0.0153	8.1702
356	2.0005	0.7368
357	0.0982	3.224
358	0.005	0.4326
359	0.006	0.4113
360	0.8527	16.1218
361	0.0656	9.4139
362	0.0532	6.0869
363	0.1707	2.3291

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
364	0.1303	5.5675
365	0.3368	15.9004
366	0.0044	0.082
367	0.0621	5.2403
368	0.0121	0.7227
369	0.0403	12.4126
370	0.1955	
371	4.2585	
372	0.1	22.847

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
373	>30	
374	0.3541	
375	5.6207	
376	0.026	7.9592
377	0.0034	0.4478
378	0.2125	14.6493
379	0.0258	0.3698
380	0.0366	2.4244
381	0.0034	0.1379

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
382	0.0339	1.5202
383	0.021	0.7714
384	0.024	3.1595
385	0.8927	
386	0.006	0.4181
387	0.0922	2.8398
388	0.2717	3.9689
389	0.1154	2.9541
390	0.3159	5.4731

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
391	0.0297	1.452
392	0.0092	0.3266
393	0.2654	8.4296
394	0.0456	2.924
395	0.008	0.679
396	0.2065	5.251
397	0.2819	7.9485
398	0.2809	2.3021
399	0.3133	11.2522

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
400	0.0589	2.7442
401	0.3602	9.105
402	0.0342	2.2146
403	0.0317	1.9302
404	0.2977	5.4805
405	0.5464	35.1743
406	0.1651	3.1455
407	0.9865	20.6904
408	0.9228	14.2627

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
409	0.0069	0.4291
410	0.0165	1.7304
411	0.0196	0.772
412	0.0466	1.6678
413	0.0207	0.8128
414	0.0302	2.0382
415	0.0055	0.4838
416	0.006	0.7234
417	0.2243	9.0943

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
418	0.004	0.062
419	0.0091	0.122
420	0.0274	0.281
421	0.0007	0.0324
422	0.1544	11.0996
423	0.0637	1.0188
424	0.0062	0.1405
425	0.0562	1.2703
426	0.0121	0.5715

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
427	0.0024	0.0472
428	0.02	0.1054
429	0.0043	0.0577
430	0.0069	0.7375
431	0.0444	0.9117
432	0.0222	1.4548
433	0.0143	1.313
434	0.0066	0.2277
435	0.0088	0.256

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
436	0.0058	0.1582
437	0.0776	1.6426
438	0.0038	0.202
439	0.1189	6.9015
440	0.0176	1.2703
441	0.0166	0.9878
442	0.0056	0.4274
443	0.009	1.175
444	0.0084	0.5143

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
445	0.0074	0.9144
446	0.1852	0.2628
447	0.0019	0.2332
448	0.0212	1.9778
449	0.0028	0.2347
450	0.1219	20.1047
451	0.0389	5.3319
452	0.0268	1.7391
453	0.0051	0.8761

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
454	0.0066	0.062
455	0.0086	0.7169
456	0.0012	0.1056
457	0.0018	0.2975
458	0.0017	0.3511
459	0.0056	0.5941
460	0.0123	0.5161
461	0.004	0.3608
462	0.0326	0.7196

Example	EC₅₀ human CYP11B2 μM	EC₅₀ human CYP11B1 μM
463	0.0471	0.9322
464	0.0028	1.6504
465	0.0669	2.2689

Example	EC₅₀ human CYP11B2 μM	EC₅₀ human CYP11B1 μM
466	0.4666	21.2595
467	0.0288	2.1821
468	0.0528	2.6647

Compounds of formula (I) and their pharmaceutically acceptable salts or esters thereof as described herein have EC₅₀ (CYP11B2) values between 0.000001 uM and 1000 uM, particular compounds have EC₅₀ (CYP11B2) values between 0.00005 uM and 500 uM, further particular compounds have EC₅₀ (CYP11B2) values between 0.0005 uM and 50 uM, more particular compounds have EC₅₀ (CYP11B2) values between 0.0005 uM and 5 uM. These results have been obtained by using the described enzymatic assay.

The compounds of formula (I) and their pharmaceutically acceptable salts can be used as medicaments (e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also be effected parentally, such as intramuscularly or intravenously (e.g. in the form of injection solutions).

The compounds of formula (I) and their pharmaceutically acceptable salts can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatin capsules. Lactose, corn starch or

derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such adjuvants for tablets, dragées and hard gelatin capsules.

Suitable adjuvants for soft gelatin capsules, are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

- 5 Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose, etc.

Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

- 10 Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

- 15 The dosage can vary in wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to 4 mg per kg body weight (e.g. about 300 mg per person), divided into preferably 1-3 individual doses, which can consist, for example, of the same amounts, should be appropriate. It will,
20 however, be clear that the upper limit given herein can be exceeded when this is shown to be indicated.

In accordance with the invention, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of aldosterone mediated diseases.

- 25 The compounds of formula (I) or their pharmaceutically acceptable salts and esters herein are inhibitors of CYP11B2. The compounds of formula (I) or their pharmaceutically acceptable salts and esters herein display also variable inhibition of CYP11B1. These compounds may be used for the inhibition of CYP11B2 in combination

with variable inhibition of CYP11B1. Such compounds may be used for treatment or prophylaxis of conditions displaying excessive cortisol production/levels or both excessive cortisol and aldosterone levels (for ex. Cushing syndrome, burn trauma patients, depression, post-traumatic stress disorders, chronic stress, corticotrophic adenomas,

5 Morbus Cushing).

In accordance with the invention, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of cardiovascular conditions (including hypertension and heart failure), vascular conditions, endothelial dysfunction, baroreceptor dysfunction, renal conditions, liver
10 conditions, fibrotic diseases, inflammatory conditions, retinopathy, neuropathy (such as peripheral neuropathy), pain, insulinopathy, edema, edematous conditions, depression and the like.

Cardiovascular conditions include congestive heart failure, coronary heart disease, arrhythmia, arterial fibrillation, cardiac lesions, decreased ejection fraction, diastolic and
15 systolic heart dysfunction, fibrinoid necrosis of coronary arteries, cardiac fibrosis, hypertrophic cardiomyopathy, impaired arterial compliance, impaired diastolic filling, ischemia, left ventricular hypertrophy, myocardial and vascular fibrosis, myocardial infarction, myocardial necrotic lesions, cardiac arrhythmias, prevention of sudden cardiac death, restenosis, stroke, vascular damage.

20 Renal conditions include acute and chronic renal failure, nephropathy, end-stage renal disease, diabetic nephropathy, decreased creatinine clearance, decreased glomerular filtration rate, expansion of reticulated mesangial matrix with or without significant hypercellularity, focal thrombosis of glomerular capillaries, global fibrinoid necrosis, glomerulosclerosis, ischemic lesions, malignant nephrosclerosis (such as ischemic
25 retraction, microalbuminuria, proteinuria, reduced renal blood flow, renal arteriopathy, swelling and proliferation of intracapillary (endothelial and mesangial) and/or extracapillary cells (crescents).

Renal conditions also include glomerulonephritis (such as diffuse proliferative, focal proliferative, mesangial proliferative, membranoproliferative, minimal change
30 membranous glomerulonephritis), lupus nephritis, non-immune basement membrane

abnormalities (such as Alport syndrome), renal fibrosis and glomerulosclerosis (such as nodular or global and focal segmental glomerulosclerosis).

Liver conditions include, but are not limited to, liver steatosis, nonalcoholic steatohepatitis, liver cirrhosis, liver ascites, hepatic congestion and the like.

5 Vascular conditions include, but are not limited to, thrombotic vascular disease (such as mural fibrinoid necrosis, extravasation and fragmentation of red blood cells, and luminal and/or mural thrombosis), proliferative arteriopathy (such as swollen myointimal cells surrounded by mucinous extracellular matrix and nodular thickening), atherosclerosis, decreased vascular compliance (such as stiffness, reduced ventricular
10 compliance and reduced vascular compliance), endothelial dysfunction, and the like.

Inflammatory conditions include, but are not limited to, arthritis (for example, osteoarthritis), inflammatory airways diseases (for example, chronic obstructive pulmonary disease (COPD)), and the like.

15 Pain includes, but is not limited to, acute pain, chronic pain (for example, arthralgia), and the like.

Edema includes, but is not limited to, peripheral tissue edema, hepatic congestion, liver ascites, splenic congestion, respiratory or lung congestion, and the like.

20 Insulinopathies include, but are not limited to, insulin resistance, Type I diabetes mellitus, Type II diabetes mellitus, glucose sensitivity, pre-diabetic state, pre-diabetes, syndrome X, and the like.

Fibrotic diseases include, but are not limited to myocardial and intrarenal fibrosis, renal interstitial fibrosis and liver fibrosis.

25 Furthermore, the compounds of formula (I) or their pharmaceutically acceptable salts and esters as described herein can also be used for the treatment or prophylaxis of cardiovascular condition selected from the group consisting of hypertension, heart failure (particularly heart failure post myocardial infarction), left ventricular hypertrophy, and stroke.

In another embodiment, the cardiovascular condition is hypertension.

In particular embodiment, the cardiovascular condition is treatment-resistant hypertension.

In another embodiment, the cardiovascular condition is heart failure.

5 In another embodiment, the cardiovascular condition is left ventricular hypertrophy.

In another embodiment, the cardiovascular condition is congestive heart failure, more particularly in patients with preserved left ventricular ejection fraction.

In another embodiment, the cardiovascular condition is stroke.

10 In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis renal condition.

In another embodiment, the renal condition is nephropathy.

In another embodiment, the renal condition is auto-immune glomerulonephritis.

In another embodiment, the chronic kidney disease is diabetic nephropathy.

In another embodiment, the fibrotic disease is kidney or heart fibrosis.

15 In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis Type II diabetes mellitus.

20 In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis Type I diabetes mellitus.



In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of diabetic retinopathy.

The invention is illustrated hereinafter by Examples, which have no limiting character.


In case the preparative examples are obtained as a mixture of enantiomers, the pure enantiomers can be separated by methods described herein or by methods known to the
5 man skilled in the art, such as e.g. chiral chromatography or crystallization.

Examples

All examples and intermediates were prepared under argon atmosphere if not specified otherwise.


Intermediate A-15 **6-Chloro-5-fluoro-3,4-dihydro-2H-isoquinolin-1-one**[A] (3-Chloro-2-fluoro-phenyl)-methanol

- To a solution of 2-fluoro-3-chloro-benzaldehyde (4.74 g, 30 mmol) in THF (20 mL) was
10 added NaBH₄ (1.48 g, 40 mmol). The resulting mixture was stirred at room temperature
for 30 min before diluting with DCM. The organic layer was washed with water and brine.
The combined organic layers were dried over anhy. Na₂SO₄, filtered and concentrated in
vacuo to give desired product without further purification (4.8g, 100%). MS: 161.0
(M+H)⁺.

15 [B] 1-Bromomethyl-3-chloro-2-fluoro-benzene


To a solution of (3-chloro-2-fluoro-phenyl)-methanol (3.2 g, 20 mmol) in DCM (20 mL) was added dropwise PBr_3 (1 mL) at 0°C . The reaction mixture was stirred at room temperature for another 1 hour before quenching with satd. aq. NaHCO_3 solution. The organic layer was washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give the desired product without further purification (4.1 g, 92%). MS: 223.2 $(\text{M}+\text{H})^+$.

[C] (3-Chloro-2-fluoro-phenyl)-acetonitrile




To a solution of 1-chloro-3-bromomethyl-2-fluoro-benzene (2.22 g, 10 mmol) in CH_3CN (30 mL) was added trimethylsilyl cyanide (1.5 mL) and TBAF (1M in THF, 12 mmol, 12 mL). The resulting reaction mixture was heated at reflux temperature for 30 min. After cooling to room temperature, the volatiles were evaporated under reduced pressure. The residue was partitioned between EtOAc and water. The organic layer was then washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo. The crude product was purified by flash column chromatography to give the title compound (1.48 g, 88%). MS: 170.1 $(\text{M}+\text{H})^+$.

[D] 2-(3-Chloro-2-fluoro-phenyl)-ethylamine




To a solution of (3-chloro-2-fluoro-phenyl)-acetonitrile (1.48 g, 8.8 mmol) in anhydrous THF (20 mL) was added a solution of borane (20 mL, 1M in THF) dropwise. The resulting reaction mixture was heated at reflux temperature for 2 hours. After cooling to room temperature, MeOH was added and the mixture was stirred at room temperature for additional 30 min. After removal of volatiles under reduced pressure, desired title product (1.30 g, 90%) was obtained as oil. MS: 174.0 $(\text{M}+\text{H})^+$.

[E] [2-(3-Chloro-2-fluoro-phenyl)-ethylamine]-carbamic methyl ester



To a solution of 2-(3-chloro-2-fluoro-phenyl)-ethylamine (1.30 g, 7.9 mmol) in CH₂Cl₂ (25 mL) was added dropwise methyl chloroformate (1.04 g, 11 mmol) and Et₃N at 5 °C
5 with vigorous stirring. The solution was then stirred at room temperature for 1h. The reaction was poured into ice-water (50 mL) and extracted with CH₂Cl₂ (2x 25 mL). The organic phase was washed with H₂O (2x 25 mL), dried over Na₂SO₄, filtered and evaporated to afford the final product without purification (1.78 g, 98%). MS: 232.1 (M+H)⁺.


10 [F] 6-Chloro-5-fluoro-3,4-dihydro-2H-isoquinoline-1-one




[2-(3-Chloro-2-fluoro-phenyl)-ethylamine]-carbamic methyl ester (890 mg, 3.85 mmol) and polyphosphoric acid (15 mL) were added to a 1-L, round bottom flask equipped with a magnetic stirring bar and reflux condenser. The reaction mixture was heated in an oil bath
15 at 140-160 °C for 2 hours while keeping vigorous stirring. The reaction mixture was then allowed to cool to room temperature and poured into H₂O (100 mL). After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. MgSO₄, filtered and concentrated to give a yellowish oil which was crystallized from Et₂O to give title compound as a white solid (30 mg, 4%). MS: 200.1 (M+H)⁺.

20 Intermediate A-2

6-Chloro-3,4-dihydro-2H-isoquinolin-1-one




[A] [2-(3-Chloro-phenyl)-ethyl]-carbamic acid methyl ester



At 0 °C , methyl chloroformate (4.6 g, 48 mmol) was added dropwise to a solution of 2-(3-chloro-phenyl)-ethylamine (5.0 g, 32 mmol) and Et₃N (6.4 g, 64 mmol) in DCM (100 mL). After the addition, the mixture was stirred at room temperature for 0.5 hour. The organic layer was washed with water (3 x 30 mL), 1N HCl (20 mL) and brine (30 mL), dried over anhy. Na₂SO₄, filtered and concentrated in vacuo. After vacuum drying, the title compound was obtained (6.49 g, 95%) as a white solid. MS: 214.1 (M+H⁺).




10 [B] 6-Chloro-3,4-dihydro-2H-isoquinolin-1-one




Under N₂ protection, a mixture of [2-(3-chloro-phenyl)-ethyl]-carbamic acid methyl ester (5.0 g, 23.4 mmol) and PPA (polyphosphoric acid) (20 g) in a 250 mL round-bottom flask was vigorously stirred at 120 °C for 2 hours. After cooling to room temperature, the reaction mixture was treated with ice-water and aqueous ammonia solution to adjust the pH to 8. Then, the mixture was then extracted with EtOAc, and the organic layer was washed with brine, dried over anhy. Na₂SO₄ and filtered. After removal of solvent under reduced pressure, the crude product obtained was further washed with ethyl ether to give title compound (1.66 g, 39%) as a white solid. MS: 182.0 (M+H⁺).

20 Intermediate A-3-1

6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one


**Intermediate A-3-2****8-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one**5 **[A] 2-(3-Chloro-phenyl)-1-methyl-ethylamine**

To a solution of 3-chlorobenzylmagnesium chloride (0.1 mol) in anhydrous ether (100 mL) (prepared from 3-chlorobenzyl chloride (16.1 g, 0.1 mol) and magnesium turnings (2.4 g, 0.1 mol) in anhydrous ether (100 mL)) was added anhydrous acetonitrile (4.1 g, 0.1
10 mol) dropwise at room temperature. When the reaction mixture was stirred at 60 °C for 3 hours, it was cooled to 0 °C followed by addition of THF (50 mL). Lithium aluminum hydride (4.2 g, 0.1 mol) was then added cautiously into the above reaction mixture and it was heated to reflux temperature for 3 hours. Ice-water was added to quench the reaction. After partitioning between ether and H₂O, the separated organic layer was washed with
15 brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo. Flash column chromatography separation (silica gel, 180 g, 5% methanol in DCM) then yielded the title compound as yellowish oil (1.9 g, 11%). MS: 170.1 (M+H⁺).

[B] [2-(3-Chloro-phenyl)-1-methyl-ethyl]-carbamic acid methyl ester

At 0 °C, methyl chloroformate (0.67 g, 7.1 mmol) was added dropwise to a mixture of 2-(3-chloro-phenyl)-1-methyl-ethylamine (1.0 g, 5.9 mmol) and potassium carbonate (1.63 g, 11.8 mmol) in THF (20 mL). After the addition, the mixture was stirred at room temperature over night. After filtration and evaporation of the volatiles, the crude product
5 was purified by flash column chromatography (silica gel, 12 g, 50% hexane in dichloromethane) to give the title compound as a white solid (1.2 g, 90%). MS: 228.1 ($M+H^+$).


[C] 6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one and 8-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one




Under N_2 protection, a mixture of [2-(3-chloro-phenyl)-1-methyl-ethyl]-carbamic acid methyl ester (1.2 g, 5.3 mmol) and PPA (polyphosphoric acid) (10 g) in a 100 mL round bottom flask was stirred at 120 °C for 2 hours. After cooling to room temperature, the reaction mixture was treated with ice-water and aqueous ammonia solution to adjust the
15 pH to 8. The mixture was then extracted with ethyl acetate and the organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated in vacuo. The crude product obtained was further washed with ether to give 6-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (0.27 g, 26%) (intermediate A-3-1) as a white solid. MS: 196.1 ($M+H^+$). The ether filtrate was concentrated under reduced pressure and purified by flash
20 chromatography to afford 8-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-2) (54 mg, 5.2%) as a white solid. MS: 196.1 ($M+H^+$).



Intermediate A-3-1a

(R)-6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one




Intermediate A-3-1b**(S)-6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one**


Chiral HPLC separation of 6-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (0.4 g, 2.05 mmol, intermediate A-3-1) afforded both (R)-6-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (0.15 g, 37.5%) and (S)-6-chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (0.12 g, 30%) as white solids. MS: 196.1 (M+H⁺).

Intermediate A-4**6-Chloro-4-methyl-3,4-dihydro-2H-isoquinolin-1-one****[A] 2-(3-Chloro-phenyl)-propionitrile**


To a solution of 3-chlorobenzonitrile (15.2 g, 0.1 mol) in THF (300 mL) was added dropwise lithium *bis*(trimethylsilyl)amide (100 mL, 1M in THF, 0.1 mol) at -78 °C. After stirring at -78 °C for 1 hour, iodomethane (14.2 g, 0.1 mol) was added dropwise. After additional stirring at -78 °C for 1 hour and subsequent warming up to room temperature for 2 hours, the reaction was quenched with water and extracted with ether. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. Flash column chromatography separation (silica gel, 180 g, 1% to 4% ether in hexane) then gave the title compound as an oil (7.3 g, 44%).

[B] 2-(3-Chloro-phenyl)-propylamine hydrochloride

To a solution of 2-(3-chloro-phenyl)-propionitrile (5.0 g, 30.2 mmol) in THF (80 mL) was added borane-tetrahydrofuran complex solution (45 mL, 45.3 mmol). The reaction mixture was stirred at room temperature overnight. Ethanol (10 mL) was then added to the reaction mixture; after stirring at room temperature for 20 min, a solution of HCl in ether (2 M, 12.5 mL) was added. After stirring for another 1 hour, water was added to induce the precipitation of the product. The white solid was collected by filtration and dried in vacuo to give the title compound (3.6 g, 70%). MS (for free amine): 170.1 (M+H⁺).

10 [C] [2-(3-Chloro-phenyl)-propyl]-carbamic acid methyl ester


To a solution of 2-(3-chloro-phenyl)-ethylamine hydrochloride (6.22 g, 30.2 mmol) and Et₃N (6.1 g, 60.4 mmol) in DCM (20 mL) was added methyl chloroformate (4.28 g, 45 mmol) dropwise at 0 °C, and the resulting mixture was stirred at room temperature for 0.5 hour. The reaction mixture was then washed with water (3 x 10 mL), 1N HCl (10 mL) and brine (10mL). The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo. The residue formed was dried in vacuo to give the title compound (5.5 g, 80%) as yellow oil. MS: 228.1 (M+H⁺).

[D] [6-Chloro-4-methyl-3,4-dihydro-2H-isoquinolin-1-one

Under N₂ protection, a mixture of [2-(3-chloro-phenyl)-propyl]-carbamic acid methyl ester (3.0 g, 13.2 mmol) and PPA (polyphosphoric acid) (10 g) in a 100 mL flask was stirred at 120 °C for 2 hours. After cooling to room temperature, the reaction mixture was treated with ice-water and aqueous ammonia solution to adjust the pH to 8. The mixture was
5 extracted with EtOAc, and the organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The crude product formed was further washed with ether to give the title compound (0.52 g, 20%) as a white solid. MS: 196.1 (M+H⁺).


Intermediate A-5

6-Chloro-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one




10

[A] 2-(3-Chloro-phenyl)-2-methyl-propionitrile




To a solution of 3-chlorobenzyl nitrile (15.2 g, 0.1 mol) in DMF (100 mL) at 0 °C was added sodium hydride (6.0 g, 0.15 mol) portionwise. After stirring at 0 °C for 0.5 hours,
15 iodomethane (14.2 g, 0.1 mol) was added dropwise. The reaction mixture was warmed up to room temperature and stirred for additional 2 hours before quenching with water. The mixture was extracted with ether and H₂O, the organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The crude product was purified by flash column chromatography (silica gel, 180 g, 1% to 4% ether in hexane) to yield the title compound
20 as oil (6.4 g, 36%).

[B] 2-(3-Chloro-phenyl)-2-methyl-propylamine




To a solution of 2-(3-chloro-phenyl)-2-methyl-propionitrile (6.4 g, 35.7 mmol) in THF (80 mL) was added borane-tetrahydrofuran complex solution (71.3 mL, 71.3 mmol), and the resulting mixture was heated to reflux for 5 hours. After cooling to 0 °C, aq. HCl (2 M, 10 mL) was added dropwise to quench the reaction. The mixture was then concentrated in vacuo to afford a solid residue, which was treated with aq. ammonia solution (6M in H₂O). After extraction with ethyl acetate, the organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The crude product was purified by flash column chromatography (silica gel, 180 g, 5% methanol in dichloromethane) to yield the title compound as oil (2.8 g, 43%). MS: 184.1 (M+H⁺).

[C] [2-(3-Chloro-phenyl)-2-methyl-propyl]-carbamic acid methyl ester



To a solution of 2-(3-chloro-phenyl)-2-methyl-propylamine (1.4 g, 7.63 mmol) and Et₃N (1.54 g, 15.26 mmol) in DCM (20 mL) was added dropwise methyl chloroformate (1.08 g, 11.4 mmol) at 0 °C. After the addition, the mixture was stirred at room temperature for 0.5 hour. The reaction mixture was then washed with water (3x 10 mL), 1N HCl (10 mL) and brine (10 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The residue formed was dried in vacuo to give the title compound (1.84 g, 100%) as yellow oil. MS: 242.1 (M+H⁺).


[D] 6-Chloro-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one



Under N₂ protection, a mixture of [2-(3-chloro-phenyl)-2-methyl-propyl]-carbamic acid methyl ester (1.84 g, 7.63 mmol) and PPA (polyphosphoric acid) (8 g) in a 100 mL round bottom flask was stirred at 120 °C for 2 hours. After cooling to room temperature, the mixture was treated with ice-water and aq. ammonia solution to adjust the pH to 8. After
5 extraction with ethyl acetate, the organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo. The crude product formed was further washed with ether to give the title compound (0.32 g, 20%) as a white solid. MS: 210.1 (M+H⁺).

Intermediate A-6

6-Chloro-2*H*-isoquinolin-1-one




10

A mixture of 6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (181.5 mg, 1 mmol, intermediate A-2) and DDQ (227 mg, 1 mmol) in dioxane (3 mL) was refluxed overnight. After cooling to room temperature, the reaction mixture was treated with satd. aq. NaHCO₃ solution, and then extracted with ethyl acetate (2 x 10 mL). The organic layers were dried over anhydrous
15 Na₂SO₄, filtered, and concentrated in vacuo to afford a crude product which was then purified by silica gel flash chromatography to give title compound (108 mg, 60%) as a white solid. MS: 180.0 (M+H⁺).


Intermediate A-7

6-Chloro-7-fluoro-3,4-dihydro-2*H*-isoquinolin-1-one




20

[A] 4-Bromomethyl-2-chloro-1-fluoro-benzene




To a solution of (3-chloro-4-fluoro-phenyl)-methanol (4.3 g, 26.8 mmol) in DCM (20 mL) was added PBr_3 (1 mL) at 0 °C. The resulting mixture was stirred at room temperature for 1 hour before it was quenched with satd. aq. NaHCO_3 solution. The organic solution was washed with water, brine, dried over anhy. Na_2SO_4 and concentrated in vacuo to give the desired product (3.7 g, 61.9%). It was used directly in the next step, without further purification.

[B] (3-Chloro-4-fluoro-phenyl)-acetonitrile



To a solution of 4-bromomethyl-2-chloro-1-fluoro-benzene (3.7 g, 16.5 mmol) in CH_3CN (30 mL) was added trimethylsilyl cyanide (2.1 mL) and TBAF (4.8 g, 18.4 mmol). The resulting mixture was heated at reflux for 30 min. After cooling of the mixture to room temperature, it was extracted with EtOAc. The organic layer was washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give a crude product which was then purified by silica gel flash column chromatography to give the title compound (1.7 g, 60.7%) as an oil.


[C] 2-(3-Chloro-4-fluoro-phenyl)-ethylamine



To a solution of (3-chloro-4-fluoro-phenyl)-acetonitrile (1 g, 5.9 mmol) in THF (30 mL) was slowly added BH_3 -THF (8.26 ml). After the addition, the reaction mixture was heated to reflux for 3 hours. MeOH was added to quench the reaction and volatiles were removed under reduced pressure. The crude product was first dissolved in aq. HCl solution (30 mL) and impurities were removed by extraction with EtOAc (2x 30 mL). The pH of the aqueous


solution was adjusted to 8 using K_2CO_3 and the mixture was then extracted with DCM (3 x 30 mL). The combined organic layers were then washed with brine, dried with anhy. Na_2SO_4 , filtered and concentrated in vacuo to give the title compound (570 mg, 55.8%) as an oil.

5 [D] N-[2-(3-Chloro-4-fluoro-phenyl)-ethyl]-2,2,2-trifluoro-acetamide




To a solution of 2-(3-chloro-4-fluoro-phenyl)-ethylamine (570 mg, 3.3 mmol) and Et_3N (1 mL) in CH_2Cl_2 (25 mL) was added trifluoroacetic anhydride (761 mg, 3.6 mmol) at 5 °C dropwise and with vigorous stirring. Then, the solution was allowed to stir at room temperature for 3 hours before it was poured into ice-water (50 mL) and extracted with DCM (2x 25 mL). The organic layer was washed with H_2O (2x 25 mL), dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give the title compound (700 mg, 78.6%) as a white solid. It was used directly in the next step without further purification.


15 [E] 1-(6-Chloro-7-fluoro-3,4-dihydro-1H-isoquinolin-2-yl)-2,2,2-trifluoro-ethanone



N-[2-(3-Chloro-4-fluoro-phenyl)-ethyl]-2,2,2-trifluoro-acetamide (400 mg, 1.48 mmol) and paraformaldehyde (89 mg, 2.96 mmol) were added sequentially at 0 °C to a solution of acetic acid (3 mL) in sulfuric acid (2 mL). After stirring of the reaction mixture at room temperature for 16 hours, the clear colorless solution was poured into ice-water (20 mL). The mixture was extracted with EtOAc (2x 30 mL) and the organic layer was washed with satd. aq. $NaHCO_3$ (20 mL), H_2O (2x 25 mL), dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give the crude title product together with a regioisomer (400 mg) as white solid.

[F] 6-Chloro-7-fluoro-1,2,3,4-tetrahydro-isoquinoline


To a solution of 1-(6-chloro-7-fluoro-3,4-dihydro-1*H*-isoquinolin-2-yl)-2,2,2-trifluoroethanone (400 mg) in MeOH (15 mL) was added K₂CO₃ (572 mg, 4 mmol) in
5 H₂O (10 mL). The resulting reaction mixture was stirred at room temperature for 2 hours before it was acidified with HCl (1N) to pH 8. It was then extracted with EtOAc and the organic layer was washed with H₂O (2 x 25 mL), dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the crude title product together with a regioisomer (170 mg). Prep-TLC separation (30% EtOAc in petroleum ether) then gave the desired title
10 compound (130 mg) as a white solid.

[G] 6-Chloro-7-fluoro-3,4-dihydro-2*H*-isoquinoline-1-one


A mixture of 6-chloro-7-fluoro-1,2,3,4-tetrahydro-isoquinoline (130 mg, 0.7 mmol), potassium bromide (83.5 mg, 0.7 mmol) and iodoso benzene (0.46 g, 2.1 mmol) in water
15 (4 mL) was stirred at room temperature overnight. The mixture was then extracted with EtOAc and the organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product. Flash column chromatography separation (silica gel, 12 g, 30% EtOAc in hexane) then afforded the title compound (58 mg, 41 %) as a white solid.

Intermediate A-8

20 **2-Chloro-7,8-dihydro-6*H*-[1,6]naphthyridin-5-one**




[A] 2-Chloro-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid methyl ester



To a solution of 2-chloro-5,6,7,8-tetrahydro-[1,6]naphthyridine hydrochloride (1.54 g, 7.5 mmol) in DCM (25 mL) was added Et₃N (3.1 mL, 22 mmol). After the addition and stirring at room temperature for 5 min, the solution was cooled to 0 °C, followed by the addition of methylchloroformate (0.85 mL, 11 mmol). The resulting reaction mixture was then allowed to stir at room temperature for 2 hours. It was then washed with satd. aq. NaHCO₃ solution and brine. The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product. Flash column chromatography separation (silica gel, 40 g, 30% EtOAc in hexane) then afforded the title compound (1.53 g, 90 %) as a white solid. MS: 227.3 (M+H⁺).


[B] 2-Chloro-5-oxo-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid methyl ester



2-Chloro-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid methyl ester (226.7 mg, 1 mmol) was dissolved in CCl₄ (3.57 mL) and MeCN (0.357 mL) at room temperature before adding NaIO₄ (0.643 g) in 1 mL of H₂O, followed by RuCl₃.hydrate (62.2 mg). The reaction mixture was stirred vigorously at room temperature for 2 hours. After dilution with DCM, it was filtered through Celite and the filter cake was washed three times with DCM. The combined organic solution was concentrated in vacuo to give a crude product.

Flash column chromatography separation (silica gel, 12 g, 20% EtOAc in hexane) then afforded the title compound (143 mg, 60 %) as a white solid. MS: 241.1 ($M+H^+$).


[C] 2-Chloro-7,8-dihydro-6H-[1,6]naphthyridin-5-one




- 5 A mixture of 2-chloro-5-oxo-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid methyl ester (60 mg, 0.25 mmol) and sodium methoxide (40 mg, 0.75 mmol) in 2 mL of 1,4-dioxane was subject to microwave reaction at 110 °C for 30 min. After removal of 1,4-dioxane, the residue was dissolved in DCM and the DCM solution was washed with water and brine. The organic layer was then dried over anhy. Na_2SO_4 , filtered and concentrated
- 10 in vacuo to give a crude product. Prep-TLC separation (30% EtOAc in hexane) then afforded the title compound (28 mg, 63 %) as a white solid. MS: 183.1 ($M+H^+$).

Intermediate A-9

2-Methoxy-7,8-dihydro-6H-[1,6]naphthyridin-5-one




- 15 [A] 2-Chloro-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid tert-butyl ester



To a solution of 2-chloro-5,6,7,8-tetrahydro-[1,6]naphthyridine hydrochloride (1.54 g, 7.5 mmol) in DCM (25 mL) was added Et_3N (2 mL, 15 mmol). After the addition and stirring


at room temperature for 5 min, the solution was cooled to 0 °C, followed by the addition of (Boc)₂O (1.88 g, 8.63 mmol). The resulting reaction mixture was then allowed to stir at room temperature for 2 hours. It was then washed with H₂O and brine. The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the title
5 compound (1.87 g, 93%) as a white solid. MS: 269.1 (M+H⁺). It was used directly in the next step without further purification.

[B] 2-Chloro-5-oxo-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid tert-butyl ester



In analogy to the procedure described for the preparation of intermediates A-8 [B], 2-
10 chloro-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid tert-butyl ester was used to yield the title compound as a white solid. MS: 283.1 (M+H⁺).



[C] 2-Methoxy-7,8-dihydro-6H-[1,6]naphthyridin-5-one





In analogy to the procedure described for the preparation of intermediates A-8 [C], 2-
15 chloro-5-oxo-7,8-dihydro-5H-[1,6]naphthyridine-6-carboxylic acid tert-butyl ester was used to yield the title compound as a white solid. MS: 179.2 (M+H⁺).

Intermediate A-10

5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one



**Intermediate A-11****6-Chloro-3-methyl-2,3-dihydro-isoindol-1-one**

- 5 [A] 5-Chloro-3-hydroxy-3-methyl-2,3-dihydro-isoindol-1-one and 6-chloro-3-hydroxy-3-methyl-2,3-dihydro-isoindol-1-one



- To a solution of 5-chloro-isoindole-1,3-dione (3.63 g, 20 mmol) in DCM (150 mL) was added methylmagnesium chloride (3 M in THF, 20 mL) dropwise at 0 °C. After the addition, the mixture was stirred at 0 °C for 3 hours before it was quenched with satd. aq. NH₄Cl solution. After extraction with DCM, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product containing a mixture of two regioisomers (3.95 g, 100%). MS: 198.1 (M+H⁺).
- 10


- [B] 5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one and 6-chloro-3-methyl-2,3-dihydro-isoindol-1-one
- 15




Under N₂ protection, triethylsilane (23 g, 200 mmol) and trifluoroboron etherate (8.51 g, 60 mmol) were added successively at -15 °C to a mixture of 5-chloro-3-hydroxy-3-methyl-2,3-dihydro-isoindol-1-one and 6-chloro-3-hydroxy-3-methyl-2,3-dihydro-isoindol-1-one (3.95 g, 20 mmol) in dry DCM (100 mL). Afterwards, the reaction mixture was stirred at
5 room temperature for 2 hours and a saturated aqueous solution of NaHCO₃ (30 mL) was added. The mixture was then extracted with DCM and the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude mixture of products. The two regioisomers were separated by prep-HPLC to give the title compounds, 5-chloro-3-methyl-2,3-dihydro-isoindol-1-one (0.4 g, 11%) and 6-chloro-3-methyl-2,3-dihydro-isoindol-1-one (0.35 g, 9.6%) as white solids. MS: 182.0 (M+H⁺).
10

Intermediate A-12

5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 5-Chloro-2-(4-methoxy-benzyl)-2,3-dihydro-isoindol-1-one



15


Sodium bis(trimethylsilyl) amide (2 mL, 2 M in THF, 2 mmol) was added dropwise at 0 °C to a solution of 5-chloro-2,3-dihydro-isoindol-1-one (0.34 g, 2 mmol) in THF (10 mL). After stirring for 10 min, 1-bromomethyl-4-methoxy-benzene (0.52 g, 2.6 mmol) was added dropwise. The resulting reaction mixture was allowed to stir at room temperature for
20 48 hours before quenching with saturated aqueous ammonium chloride solution. After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the crude product (0.6 g, 100%) as a yellow oil. MS: 288.0 (M+H⁺). It was used directly in the next step, without further purification.

[B] 5-Chloro-2-(4-methoxy-benzyl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one



To a solution of 5-chloro-2-(4-methoxy-benzyl)-2,3-dihydro-isoindol-1-one (0.6 g, 2.0 mmol) in THF (10 mL) was added sodium hydride (60% in mineral oil, 0.17 g, 4.2 mmol) at room temperature. The resulting reaction mixture was stirred for 30 min before iodomethane (0.60 g, 4.2 mmol) was added. After stirring at room temperature overnight, the mixture was quenched with brine and extracted with EtOAc. The organic layer was then washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the crude product which was then purified by flash column chromatography (silica gel 14 g, 5% to 20% ethyl acetate in DCM). The title compound was obtained (0.38 g, 57%) as a white solid. MS: 316.2 (M+H⁺).


[C] 5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




A solution of 5-chloro-2-(4-methoxy-benzyl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (0.38 g, 1.2 mmol) in trifluoro acetic acid (5 mL) was heated to reflux for 20 hours. After removal of trifluoro acetic acid under reduced pressure, the crude product was purified by flash column chromatography (silica gel 14 g, 5% to 50% ethyl acetate in DCM) to give the title compound (0.20 g, 85%) as a white solid. MS: 196.1 (M+H⁺).

Intermediate A-12-1

20 **5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one**




[A] (5-Bromo-pyridin-3-yl)-methanol



Sodium borohydride (2.2 g, 59.1 mmol) was added to a suspension of 5-bromo-pyridine-3-carbaldehyde (10.0 g, 53.7 mmol) in MeOH (100 mL) at 0 °C. The mixture was stirred at 0 °C for 1 hour before it was quenched by the addition of water (5.0 mL). Evaporation of solvents afforded a light yellowish oil which was re-dissolved in EtOAc and washed with water. The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the title compound (9.6 g, 95%) as colorless oil. MS: 188.0 & 190.0 (M+H⁺).


[B] 5-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In a 25-mL sealed tube, (5-bromo-pyridin-3-yl)-methanol (900 mg, 4.8 mmol), 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12 [C], 858 mg, 4.4 mmol), CuI (200 mg, 1.1 mmol), Cs₂CO₃ (3.0 g, 9.2 mmol) and (+)-(S,S)-1,2-diaminocyclohexane (0.4 mL, 3.2 mmol) were dissolved in dioxane (8.0 mL). The resulting reaction mixture was heated at 150 °C for 3 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 125 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by silica gel

flash chromatography (30-100% EtOAc-hexane gradient) to yield the title compound (1.2 g, 90%) as a light yellow solid. MS: 303.2 (M+H⁺).


[C] 5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one





- 5 Thionyl chloride (1.4 mL, 19.0 mmol) was added slowly at 0 °C to a solution of 5-chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (1.14 g, 3.8 mmol) in DCM (50 mL). After the addition, the reaction mixture was stirred at 2-5 °C for 2 hours before it was poured into satd. aq. NaHCO₃ solution (50 mL) and extracted with EtOAc (2 x 150 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄,
10 filtered and concentrated in vacuo to give a crude product (1.42 g, 92%) as a light yellowish solid. MS: 322.1 (M+H⁺).

Intermediate A-13

5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one




- 15 [A] 5-Chloro-3-ethyl-3-hydroxy-2,3-dihydro-isoindol-1-one and 6-chloro-3-ethyl-3-hydroxy-2,3-dihydro-isoindol-1-one



To a solution of 5-chloro-isoindole-1,3-dione (5.0 g, 27.5 mmol) in DCM (200 mL) was added ethylmagnesium chloride (2 M in THF, 41.3 mL) dropwise at 0 °C. After the addition, the mixture was allowed to stir at 0 °C for 3 hours before it was quenched with satd. aq. NH₄Cl solution. After extraction with DCM, the organic layer was washed with
5 brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product containing a mixture of two regioisomers (5.82 g, 100%). MS: 212.0 (M+H⁺).


[B] 5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one



Under N₂ protection, triethylsilane (19.3 g, 166 mmol) and TFA (20 mL) were added
10 successively to a mixture of 5-chloro-3-ethyl-3-hydroxy-2,3-dihydro-isoindol-1-one and 6-chloro-3-ethyl-3-hydroxy-2,3-dihydro-isoindol-1-one (3.52 g, 16.6 mmol). After stirring at room temperature for 2 hours, the reaction mixture was concentrated under reduced pressure. The residue was treated with a satd. aq. solution of NaHCO₃ (30 mL) and extracted with DCM (2 x 30 mL). The combined organic layers were washed with brine,
15 dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude mixture of products. The desired regioisomer, 5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one was obtained by prep-HPLC separation as a white solid (0.65 g, 20%). MS: 196.1 (M+H⁺).

Intermediate A-13-1


5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one





20

Intermediate A-13-2

6-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one




[A] 5-Chloro-3-ethyl-3-hydroxy-2,3-dihydro-indol-1-one and 6-chloro-3-ethyl-3-hydroxy-2,3-dihydro-indol-1-one




- 5 To a solution of 5-chloro-indole-1,3-dione (5.0 g, 27.5 mmol) in DCM (200 mL) was added ethylmagnesium chloride (2 M in THF, 41.3 mL) dropwise at 0 °C. After the addition, the mixture was allowed to stir at 0 °C for 3 hours before it was quenched with satd. aq. NH₄Cl solution. After extraction with DCM, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product
- 10 containing a mixture of two regioisomers (5.82 g, 100%). MS: 212.0 (M+H⁺).

[B] 5-Chloro-3-ethyl-2,3-dihydro-indol-1-one and 6-Chloro-3-ethyl-2,3-dihydro-indol-1-one



and




- Under N₂ protection, triethylsilane (19.3 g, 166 mmol) and TFA (20 mL) were added
- 15 successively to a mixture of 5-chloro-3-ethyl-3-hydroxy-2,3-dihydro-indol-1-one and 6-chloro-3-ethyl-3-hydroxy-2,3-dihydro-indol-1-one (3.52 g, 16.6 mmol). After stirring at room temperature for 2 hours, the reaction mixture was concentrated under reduced pressure. The residue was treated with a satd. aq. solution of NaHCO₃ (30 mL) and extracted with DCM (2 x 30 mL). The combined organic layers were washed with brine,
- 20 dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude mixture of products. Both 5-chloro-3-ethyl-2,3-dihydro-indol-1-one (0.65 g, 20%) MS: 196.1

($M+H^+$) and 6-chloro-3-ethyl-2,3-dihydro-isoindol-1-one (0.62 g, 19%) MS: 196.1 ($M+H^+$) were obtained by prep-HPLC separation as white solids.

Intermediate A-13-1a


(R or S)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one



5

Intermediate A-13-1b


(S or R)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one




10

Intermediate A-13-2a

(R or S)-6-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one

15 **Intermediate A-13-2b**


(S or R)-6-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one



The mixture of 5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one and 6-chloro-3-ethyl-2,3-dihydro-isoindol-1-one was subject to SFC separation (IC 250 mm x 50 mm, 5 μ m, mobile phase A: supercritical CO₂, B: IPA (0.05% NH₃H₂O), A: B = 60: 40 at 140 mL/min) to give (R or S)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1a) and (S or R)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one as one pair of enantiomers and (R or S)-6-chloro-3-ethyl-2,3-dihydro-isoindol-1-one and (S or R)-6-chloro-3-ethyl-2,3-dihydro-isoindol-1-one as another pair of enantiomers.

Intermediate A-13-3


10 (S or R)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of intermediate A-12-1, (S or R)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one was used to afford the title compound as yellowish oil. MS: 322.1 (M+H⁺).

15 **Intermediate A-13-4**


(R or S)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one





In analogy to the procedure described for the preparation of intermediate A-12-1, (R or S)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one was used to afford the title compound as yellowish oil. MS: 322.1 ($M+H^+$).

5 Intermediate A-14

3-Benzyl-5-chloro-2,3-dihydro-isoindol-1-one



[A] 3-Benzyl-5-chloro-3-hydroxy-2,3-dihydro-isoindol-1-one and 3-benzyl-6-chloro-3-hydroxy-2,3-dihydro-isoindol-1-one




10

To a solution of 5-chloro-isoindole-1,3-dione (3.63 g, 20 mmol) in DCM (150 mL) was added benzylmagnesium chloride (2 M in THF, 30 mL) dropwise at 0 °C. After the addition, the reaction mixture was allowed to stir at 0 °C for 3 hours before it was quenched with satd. aq. NH_4Cl solution. After extraction with DCM, the organic layer was washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give a crude product containing a mixture of two regioisomers (5.47 g, 100%). MS: 274.1 ($M+H^+$).

15


[B] 3-Benzyl-5-chloro-2,3-dihydro-isoindol-1-one




Under N₂ protection, triethylsilane (23 g, 200 mmol) and TFA (10 mL) were added successively to a mixture of 3-benzyl-5-chloro-3-hydroxy-2,3-dihydro-isoindol-1-one and
5 3-benzyl-6-chloro-3-hydroxy-2,3-dihydro-isoindol-1-one (5.47 g, 20 mmol). After stirring at room temperature for 2 hours, the reaction mixture was concentrated under reduced pressure. The residue was treated with a satd. aq. NaHCO₃ solution (30 mL) and extracted with DCM (2 x 30 mL). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude mixture of products. The
10 desired regioisomer, 3-benzyl-5-chloro-2,3-dihydro-isoindol-1-one was obtained by prep-HPLC separation as a white solid (1.03 g, 20%). MS: 258.1 (M+H⁺).

Intermediate A-15

5-Chloro-3-cyclopropyl-2,3-dihydro-isoindol-1-one




15 [A] (2-bromo-5-chlorophenyl)(cyclopropyl)methanamine



To a solution of 2-bromo-5-chlorobenzonitrile (20 g, 93 mmol) in THF (200 mL) was added cyclopropylmagnesiumbromide (558 mL, 279 mmol) at 0 °C. The resulting reaction

- mixture was stirred at 0-5 °C for 5 hours before MeOH (100 mL) was added and stirring continued at room temperature for 15 min. NaBH₄ (7.1 g, 186 mmol) was added subsequently and the reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was extracted with AcOEt (300 mL x 3). The combined organic layers were concentrated *in vacuo* to give a crude product which was then purified by chromatography to give (2-bromo-5-chlorophenyl)(cyclopropyl)methanamine (8.2 g, yield 34%).


[B] 5-Chloro-3-cyclopropyl-2,3-dihydro-isoindol-1-one




- A mixture of (2-bromo-5-chlorophenyl)(cyclopropyl)methanamine (2 g, 7.7 mmol), Pd(dppf)Cl₂ (0.2 g), DIPEA (3 g, 23.1 mmol) in 20 mL of DMF was heated in an autoclave at 130 °C under 2 MPa of CO (g) for 16 hours. After the reaction, the mixture was diluted with AcOEt (150 mL) and washed with brine (30 mL x 3). The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to give a crude product which was purified by chromatography to give title compound (1.1 g, yield 68.7%) as a yellow solid. MS: 207.9 (M+H⁺, 1Cl).

Intermediate A-16

2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




- [A] 3-(Methoxycarbonyl)-2-methylpyridine 1-oxide




- To a stirred solution of methyl-2-methylnicotinate (95 g, 629 mmol) in DCM (1.5 L) was added m-CPBA (119 g, 692 mmol) at 0 °C. After the reaction mixture was stirred at room temperature for 16 hours, it was washed with a mixture of satd. aq. Na₂SO₃ and NaHCO₃ solution. The organic layer was then dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to give a crude product (60 g, yield 57%), which was used in the next step reaction without further purification.

[B] Methyl 2-(chloromethyl)nicotinate



- The crude 3-(methoxycarbonyl)-2-methylpyridine-1-oxide (35 g, 210 mmol) was added in small portion to POCl₃ (300 g) at room temperature. After the addition, the reaction mixture was refluxed for 3 hours before it was concentrated in *vacuo*. The residue was poured into ice-water, neutralized with aq. NaHCO₃ solution and extracted with AcOEt (125 mL x 3). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to afford a crude product which was then purified by column chromatography to give title compound (12 g, yield 30%).


[C] 2-(Chloromethyl)-3-(methoxycarbonyl)pyridine 1-oxide



- To a stirred solution of methyl-2-(chloromethyl)nicotinate (20 g, 108 mmol) in DCM (300 mL) was added m-CPBA (20.5 g, 119 mmol) at 0 °C. After it was stirred at room temperature for 16 hours, the reaction mixture was washed with a mixture of satd. aq.


Na₂SO₃ and NaHCO₃ solution. The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated in vacuo to give the crude title product (20 g, yield 92%), which was used in the next step reaction without further purification.

[D] Methyl 6-chloro-2-(chloromethyl)nicotinate




The crude of 2-(chloromethyl)-3-(methoxycarbonyl)pyridine-1-oxide (20 g, 99.5 mmol) was added in small portion to POCl₃ (200 g) at room temperature. The mixture was refluxed for 3 hours before it was concentrated *in vacuo*. The residue was poured into ice-water, neutralized with NaHCO₃ solution, and extracted with AcOEt (125 mL x 3). The combined organic layers were concentrated to give the crude title product (17 g, yield 78%), which was used in the next step reaction without further purification.

[E] 2-Chloro-6-(4-methoxybenzyl)-6,7-dihydro-5H-pyrrolo[3,4-b]pyridin-5-one




To a stirred solution of crude methyl 6-chloro-2-(chloromethyl)nicotinate (10 g, 45.4 mmol) in THF (150 mL) was added PMBNH₂ (15.5 g, 113.5 mmol) at 0 °C. The resulting reaction mixture was stirred at room temperature for 16 hours before it was concentrated under reduced pressure to give a crude product. After washing with MTBE (100 mL x 3), the title compound was obtained (8.8 g, yield 67%) as a white solid. MS: 288.8 (M+H⁺, 1Cl).

20 [F] 2-Chloro-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



To a solution of 2-chloro-6-(4-methoxy-benzyl)-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (5.8 g, 20.0 mmol) in THF (50 mL) was added sodium hydride (60% in mineral oil, 1.7 g, 42.0 mmol) at room temperature. The resulting reaction mixture was stirred for 30 min
5 before iodomethane (6.0 g, 42.0 mmol) was added. After stirring at room temperature over night, the mixture was quenched with water and extracted with EtOAc. The organic layer was then washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give the crude product which was then purified by flash column chromatography (silica gel 20 g, 5% to 20% ethyl acetate in DCM). The title compound was obtained (3.8 g, 57%)
10 as a white solid. MS: 316.2 (M+H⁺).


[G] 2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




To solution of 2-chloro-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (1.58 g, 5.0 mmol) in CH₃CN (20 mL) was added ceric ammonium nitrate
15 (8.2 g, 15.0 mmol) at room temperature. The reaction mixture was stirred at room temperature for 3 hours before water and EtOAc was added into the mixture. The organic layer was separated, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by silica gel column chromatography to give title compound (617 mg, 63%) as a solid. MS: 197.2 (M+H⁺).

20 Intermediate A-16-1

2-Chloro-6-(5-chloromethyl-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




[A] 2-Chloro-6-(5-hydroxymethyl-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



- 5 2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16 [G], 39.2 mg, 0.2 mmol), (5-bromo-pyridin-3-yl)-methanol (example 3 [A], 74.8 mg, 0.4 mmol), CuI (3.8 mg, 0.02 mmol), (1S, 2S)-cyclohexane-1,2-diamine (4.5 mg, 0.04 mmol) and Cs₂CO₃ (130 mg, 0.4 mmol) were dissolved in dioxane (5 mL). The reaction mixture was subjected to microwave reaction at 140 °C for 1 hours before it was poured into H₂O
- 10 (50 mL) and extracted with EtOAc (2x 25 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product (55.7 mg, 92%) as solid. MS: 304.1 (M+H⁺).

[B] 2-Chloro-6-(5-chloro-methyl-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




15

- Thionyl chloride (0.14 mL, 1.9 mmol) was added slowly at 0 °C to a solution of 2-chloro-6-(5-hydroxymethyl-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (55.7 mg, 0.18 mmol) in DCM (10 mL). After the addition, the reaction mixture was stirred at 2-5 °C for 0.5 hours before it was poured into satd. aq. NaHCO₃ solution (10 mL)
- 20 and extracted with DCM (2 x 15 mL). The organic layer was washed with brine, dried over

anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product (58 mg 100%) as light yellowish solid. MS: 322.1 (M+H⁺).


Intermediate A-17

2-Methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



5


[A] 2-Methoxy-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



To solution of 2-chloro-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16 [F], 3.15 g, 10 mmol) in DMF (30 mL) was added sodium methanolate (0.813 g, 15 mmol) at room temperature. The reaction mixture was stirred at room temperature for 4 hours before it was quenched with water and extracted with EtOAc. The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give title compound (2.8 g, 90%) as a solid. MS: 313.1 (M+H⁺).

15


[B] 2-Methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



To solution of 2-methoxy-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (0.31 g, 1.0 mmol) in CH₃CN (5 mL) was added ceric ammonium nitrate (1.64 g, 3.0 mmol) at room temperature. The reaction mixture was stirred at room temperature for 3 hours before water and EtOAc were added into the mixture. The organic layer was separated, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by silica gel column chromatography to give the title compound (0.12 g, 63%) as a solid. MS: 193.1 (M+H⁺).

Intermediate A-17-1


6-(5-Chloromethyl-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 318.3 (M+H⁺).

Intermediate A-18

2-Ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




[A] 2-Ethoxy-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



- To solution of 2-chloro-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16 [F], 3.15 g, 10 mmol) in DMF (30 mL) was added sodium ethoxide (1.02 g, 15 mmol) at room temperature. The reaction mixture was stirred at room temperature for 4 hours before it was quenched by water and extracted with EtOAc. The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give title product (2.9g, 89%) as a solid. MS: 327.2 (M+H⁺).


[B] 2-Ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one



- To solution of 2-methoxy-6-(4-methoxy-benzyl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (0.33 g, 1.0 mmol) in CH₃CN (5 mL) was added ceric ammonium nitrate (1.64 g, 3.0 mmol) at room temperature. The reaction mixture was stirred at room temperature for 3 hours before water and EtOAc were added into the mixture. The organic layer was separated, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by silica gel column chromatography to give the title compound (0.15 g, 72%) as a solid. MS: 207.1 (M+H⁺).

Intermediate A-18-1


6-(5-Chloromethyl-pyridin-3-yl)-2-ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one




In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 332.1 ($M+H^+$).

Intermediate A-19

5 5-Fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 4-Fluoro-2-methyl-benzoic acid methyl ester




To a stirred solution of 4-fluoro-2-methyl-benzoic acid (13 g, 84 mmol) in MeOH (500 mL) was added SOCl_2 (20 g, 168 mmol) dropwise at 0 °C. After the addition, the reaction mixture was stirred at room temperature for 16 hours. When TLC indicated that no starting material was left, the mixture was concentrated under reduced pressure to give a crude product (12.7 g, yield 90.7%), which was used in the next step reaction without further purification.

15 [B] 2-Bromomethyl-4-fluoro-benzoic acid methyl ester




A mixture of 4-fluoro-2-methyl-benzoic acid methyl ester (12.7 g, 75.5 mmol), NBS (13.4 g, 75.5 mmol), and di-benzoyl peroxide (BPO) (0.85 g, 3.5 mmol) in CCl₄ (200 mL) was heated to reflux for 3 hours until all starting material was consumed. After vacuum
 5 filtration, the filtrate concentrated *in vacuo* to give a crude product (17 g, yield 91%), which was used in the next step reaction without further purification.

[C] 5-Fluoro-2-(4-methoxy-benzyl)-2,3-dihydro-isoindol-1-one



To a stirred solution of 2-bromomethyl-4-fluorobenzoic acid methyl ester (17 g, 68.8
 10 mmol) in THF (200 mL) was added PMBNH₂ (23.6 g, 172 mmol) at 0 °C. After stirring at room temperature for 16 hours, the reaction mixture was concentrated *in vacuo* to give a crude product which was then purified by column chromatography to give the title compound (13 g, yield 69%) as a solid.

[D] 5-Fluoro-2-(4-methoxy-benzyl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one




15

To a solution of 5-fluoro-2-(4-methoxy-benzyl)-2,3-dihydro-isoindol-1-one (6 g, 22 mmol) in THF (30 mL) was added NaH (4.42 g, 110 mmol) slowly at 0 °C. After stirred at 0 °C for 30 min, MeI (18.8 g, 132 mmol) was added dropwise and the resulting reaction mixture was heated to 70 °C for 5 hours. After cooling to room temperature, the reaction mixture

was poured into aq. NH_4Cl solution and extracted with EtOAc. The organic layer was dried over anhydrous Na_2SO_4 , filtered, and then concentrated to give a crude product which was then purified by column chromatography to give the product (4.8 g, yield 72.5%) as a solid.

[E] 5-Fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




5

To a stirred solution of 5-fluoro-2-(4-methoxy-benzyl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (3.8 g, 12.7 mmol) in MeCN (80 ml) and H_2O (40 mL) was added CAN (20.9 g, 38.1 mmol) at 0 °C. After stirring for 2 hours at 0 °C, the reaction mixture was diluted with EtOAc and washed with brine. The organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated *in vacuo* to provide a crude product, which was purified by column chromatography to give the product (1.03 g, yield 45%) as a solid.

10

Intermediate A-19-1

2-(5-Chloromethyl-pyridin-3-yl)-5-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 305.1 ($\text{M}+\text{H}^+$).


15

Intermediate A-20

6-Fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 1-(2-Bromo-4-fluoro-phenyl)-1-methyl-ethylamine



To a stirred solution of 2-bromo-4-fluorobenzonitrile (5 g, 25 mmol) in ether (100 mL)
5 was added MeMgBr (25 mL, 75 mmol) at room temperature. The solution was stirred at
room temperature for 0.5 hour before Ti(Oi-Pr)₄ (41 mL, 25 mmol) was added and the
resulting mixture was heated to reflux for 6 hours. Aqueous NaOH (10%, 200 mL)
solution was then slowly added to the reaction mixture at 0 °C and the reaction mixture
was stirred at room temperature for additional 30 minutes. After dilution with aq. Na₂CO₃
10 (5%, 400 mL) solution, it was extracted with t-BuOMe (100 mL x 3). The combined
organic layers were concentrated under reduced pressure. The residue obtained was diluted
with aq. HCl solution (5%) and the aqueous layer was washed with t-BuOMe (50 mL x 2)
and basified with 20% aq. NaOH to pH~ 10. The resulting aqueous layer was further
extracted with t-BuOMe (100 mL x 3). The combined organic layers were then dried over
15 anhy. Na₂SO₄, filtered, and concentrated in vacuo to give a crude product, which was
purified by column chromatography to give the title product (3.0 g, yield 51.7%) as
yellowish oil. MS: 231.7 (M+H⁺, 1Br)


[B] 6-Fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



A mixture of 1-(2-bromo-4-fluoro-phenyl)-1-methyl-ethylamine (1.5 g, 6.52 mmol), Pd(dppf)Cl₂ (0.15 g), DIPEA (2.52 g, 19.6 mmol) in DMF (20 mL) was stirred in an autoclave under 2 MPa of CO (g) at 130 °C for 16 hours. After it was cooled to room temperature, the reaction mixture was diluted with EtOAc (300 mL). The organic layer
5 was washed with brine, filtered, and concentrated under reduced pressure to give a crude product which was purified by chromatography to give the title compound (100 mg, yield 8.6%) as a brown solid.


Intermediate A-20-1

2-(5-Chloromethyl-pyridin-3-yl)-6-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




10

[A] 6-Fluoro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one



A mixture of 6-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-20 [B], 35.8 mg, 0.2 mmol), (5-bromo-pyridin-3-yl)-methanol (74.8 mg, 0.4 mmol), CuI (3.8 mg, 0.02 mmol), (1S, 2S)-cyclohexane-1,2-diamine (4.5 mg, 0.04 mmol) and Cs₂CO₃ (130 mg, 0.4 mmol) were dissolved in dioxane (5 mL). The reaction mixture was subjected to microwave reaction at 140 °C for 1 hour before it was poured into H₂O (50 mL) and extracted with EtOAc (2x 25 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product (52 mg, 92%) as
15 a solid. MS: 287.1 (M+H⁺).
20


[B] 2-(5-Chloromethyl-pyridin-3-yl)-6-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




To a solution of 6-fluoro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (52 mg) in DCM (10 mL) was added thionyl chloride (0.14 mL, 1.9 mmol) slowly at 0 °C. After the addition, the reaction mixture was stirred at 2-5 °C for 0.5 hour
 5 before it was poured into satd. aq. NaHCO₃ solution (10 mL) and extracted with DCM (2 x 15 mL). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product (58 mg 100%) as light yellowish solid. MS: 322.1 (M+H⁺).

Intermediate A-21

10 3,3-Dimethyl-1-oxo-2,3-dihydro-1H-isoindole-5-carbonitrile




[A] 4-Bromo-2-methyl-benzoic acid methyl ester




To a solution of 4-bromo-2-methyl-benzoic acid (30.0 g, 0.14mol) in 115 mL of methanol
 15 was added thionyl chloride (20.25 mL, 0.28 mol) slowly and the reaction mixture was stirred at 70 °C for 2 hours before it was concentrated to afford a crude product which was then purified by column chromatography to give the title compound (30.03 g, 93.6%) as a solid.

[B] 4-Cyano-2-methyl-benzoic acid methyl ester




A mixture of 4-bromo-2-methyl-benzoic acid methyl ester (26.0 g, 113.5 mmol) and CuCN (12.48 g, 140.7 mmol) was heated at 180 °C for 5 hours before it was poured into ice-water. The solid precipitate was collected by vacuum filtration to give a crude product which was then purified by column chromatography to afford the title compound (12.53 g, 63%) as a solid.

[C] 2-Bromomethyl-4-cyano-benzoic acid methyl ester



A mixture of 4-cyano-2-methyl-benzoic acid methyl ester (12.5 g, 71.35 mmol), NBS (12.7 g, 71.35 mmol) and di-benzoyl peroxide (BPO) (0.8 g, 3.28 mmol) in CCl₄ (200 mL) was heated to reflux temperature for 3 hours. After it was cooled to room temperature, the reaction mixture was filtered. The filtrate was concentrated *in vacuo* to give a crude product (18.2 g) which was used in the next step reaction without further purification.


15 [D] 2-(4-Methoxy-benzyl)-1-oxo-2,3-dihydro-1H-isindole-5-carbonitrile



To a solution of 2-bromomethyl-4-cyano-benzoic acid methyl ester (18.1 g, 71.24 mmol) in THF (300 mL) was added PMBNH₂ (23.4 g, 178.1 mmol) at 0 °C and the reaction mixture was stirred at room temperature for 16 hours. After vacuum filtration, the filtrate


was concentrated *in vacuo*. The residue obtained was re-dissolved in EtOAc and washed with water and brine. The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to give a crude product which was purified by column chromatography (11.69 g, 56.0%) as a solid.

5 [E] 2-(4-Methoxy-benzyl)-3,3-dimethyl-1-oxo-2,3-dihydro-1H-isoindole-5-carbonitrile



To a solution of 2-(4-methoxy-benzyl)-1-oxo-2,3-dihydro-1H-isoindole-5-carbonitrile (11.6 g, 41.7 mmol) in THF (300 mL) was added NaH (8.34 g, 208.4 mmol, 60% in mineral oil) and the reaction mixture was stirred at room temperature for 1 hour before
10 iodomethane (35.5 g, 250.1 mmol) was added. After the addition, the reaction mixture was stirred at 70 °C for 2 hours until all the starting material was consumed. After it was cooled to room temperature, satd. aq. NH₄Cl solution was added and the mixture was extracted with EtOAc (200 mL×3). The combined organic layers were dried over anhy. MgSO₄, filtered, and concentrated under reduced pressure to give a crude product which
15 was purified by column chromatography to afford the title compound (7.22 g, 56.5%) as a solid.

[F] 3,3-Dimethyl-1-oxo-2,3-dihydro-1H-isoindole-5-carbonitrile




To a solution of 2-(4-methoxy-benzyl)-3,3-dimethyl-1-oxo-2,3-dihydro-1H-isoindole-5-carbonitrile (3.5 g, 11.42 mmol) in MeCN (70 mL) was added CAN (18.79 g, 34.27 mmol)
20 in 30 mL of water at 0 °C. The resulting reaction mixture was stirred at 0 °C for 1 hour until all the starting material was consumed. The reaction mixture was extracted between

water and EtOAc and the combined organic layers were dried over anhy. MgSO_4 , filtered, and concentrated under reduced pressure to give a crude product which was purified by column chromatography to afford the title compound (1.06 g, 49.8%) as a solid.

Intermediate A-21-1


- 5 **2-(5-Chloromethyl-pyridin-3-yl)-3,3-dimethyl-1-oxo-2,3-dihydro-1*H*-isoindole-5-carbonitrile**




In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 312.1 ($\text{M}+\text{H}^+$).

- 10 **Intermediate A-22**

6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one




[A] 1-(2-Bromo-5-chlorophenyl)cyclopropanamine



- 15 To a stirred solution of 2-bromo-5-chlorobenzonitrile (10 g, 46 mmol) and $\text{Ti}(\text{Oi-Pr})_4$ (16.64 mL, 55 mmol) in THF (200 mL) at -78°C was added EtMgBr (138 mL, 138 mmol) dropwise. The reaction mixture was allowed to warm up to room temperature and stirred

for 2 hours. $\text{BF}_3\text{-Et}_2\text{O}$ (17.2 mL) was added, and the solution was stirred for another 16 hours before it was quenched with aq. HCl solution and washed with EtOAc. The aqueous phase was adjusted to pH ~ 10 with aq. NaOH solution, and extracted with EtOAc (3 x). The combined organic layers were concentrated to give a crude product which was
5 purified by chromatography to afford title compound (2 g, yield 17.6%). MS: 246.7 ($\text{M}+\text{H}^+$, 1Cl) as oil.


[B] 6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one





A mixture of 1-(2-bromo-5-chlorophenyl)cyclopropanamine (2 g, 8.1 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$
10 (0.2 g), DIPEA (3.1 g, 24.3 mmol) in DMF (20 mL) was stirred in an autoclave under 2 MPa of CO (g) at 130 °C for 16 hours. The reaction mixture was diluted with EtOAc (300 mL), and washed with brine. The organic layer was dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to give a crude product which was purified by chromatography to afford title compound (700 mg, yield 44.6%) as a yellow solid. MS:
15 193.8 ($\text{M}+\text{H}^+$, 1Cl).

Intermediate A-22-1

6'-Chloro-2'-(5-(chloromethyl)pyridin-3-yl)spiro[cyclopropane-1,1'-isoindolin]-3'-one




In analogy to the procedure described for the preparation of intermediate A-12-1, title
20 compound was obtained as yellowish oil. MS: 320.1 ($\text{M}+\text{H}^+$).

Intermediate A-23**6'-Fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one****[A] 1-(2-Bromo-5-fluoro-phenyl)-cyclopropylamine**

5

To a stirred solution of 2-bromo-5-fluoro-benzonitrile (1 g, 5 mmol) and $\text{Ti}(\text{Oi-Pr})_4$ (1.81 mL, 5.5 mmol) in ether (20 mL) was added EtMgBr (3.67 mL, 11 mmol) at -78°C . After stirring at -78°C for 10 min, the solution was warmed up to room temperature and stirred for 1 hour. $\text{BF}_3\cdot\text{Et}_2\text{O}$ (1.25 mL) was added before it was stirred for another 1 hour. After addition of 1 N aq. HCl (15 mL) and ether (30 mL), aq. NaOH (10%, 45 mL) solution was added to give two clear phases. The mixture was extracted with EtOAc (30 mL x 5). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo to give a crude product, which was purified by column chromatography (PE:EA = 5:1) to afford the title compound (800 mg, 70%) as oil.


15 **[B] 6'-Fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one**

A solution of 1-(2-bromo-5-fluoro-phenyl)-cyclopropylamine (4.2 g, 18.3 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.42 g), DIPEA (7 g, 54.8 mmol) in 40 mL of DMF was heated at 130°C

under 2 MPa of CO for 12 hours. After it was cooled to room temperature, EtOAc was added it was washed with brine. The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo*. The crude product was purified by column chromatography to give title compound (0.3 g, yield 65.2%) as a solid, meanwhile, 3.3 g of starting material
5 was recovered.

Intermediate A-23-1


2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one




In analogy to the procedure described for the preparation of intermediate A-12-1, the title
10 compound was obtained as yellowish oil. MS: 303.1 (M+H⁺).

Intermediate A-24

5'-Fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one




[A] 1-(2-Bromo-4-fluorophenyl)cyclopropanamine



To a stirred solution of 2-bromo-4-fluorobenzonitrile (5 g, 25 mmol) and $\text{Ti}(\text{O}i\text{-Pr})_4$ (9.05 mL, 27.5 mmol) in ether (100 mL) at -78°C was added EtMgBr (18.3 mL, 55 mmol) dropwise. The solution was allowed to warm to room temperature and stirred for 1 hour before $\text{BF}_3\cdot\text{Et}_2\text{O}$ (6.25 mL) was added and stirring continued at room temperature for another 1 hour. The reaction solution was quenched with 1 N HCl solution, and washed with EtOAc . The aqueous layer was adjusted to $\text{pH}\sim 10$ with aq. NaOH (2 N) solution and then extracted with EtOAc (3 x). The combined organic layers were dried over anhyd. Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was then purified by column chromatography to give title compound (3.0 g, yield 52.2%) as yellowish oil.


10 [B] 5'-Fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one



A mixture of 1-(2-bromo-4-fluorophenyl)cyclopropanamine (3.0 g, 17.4 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.4 g), DIPEA (5.05 g, 39.1 mmol) in DMF (40 mL) was stirred in an autoclave under 2 MPa of CO (g) at 130°C for 16 hours. After it was cooled down to room temperature, the reaction mixture was diluted with EtOAc (300 mL) and washed with brine. The organic layer was dried over anhyd. Na_2SO_4 , filtered, and concentrated under reduced pressure to give a crude product which was purified by chromatography to afford title compound (600 mg, yield 26%) as a brown solid.

Intermediate A-24-1


20 **2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one**



In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 303.1 ($M+H^+$).

Intermediate A-25

6-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




5

In analogy to the procedure described for the preparation of intermediate A-20, 2-bromo-4-chloro-benzonitrile (step A) and 1-(2-bromo-4-chloro-phenyl)-1-methyl-ethylamine (step B) were used to yield the title compound as a solid (yield 40%). MS: 196.1 ($M+H^+$).

Intermediate A-25-1


10 **6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one**




In analogy to the procedure described for the preparation of intermediate A-12-1, the title compound was obtained as yellowish oil. MS: 322.1 ($M+H^+$).

Intermediate B-1

15 **3-Bromo-5-(2-isopropyl-imidazol-1-ylmethyl)-pyridine**




[A] 3-Bromo-5-chloromethyl-pyridine



To a solution of (5-bromopyridin-3-yl)methanol (3 g, 16.0 mmol) in DCM (15 mL) cooled
5 to 0 °C was added thionylchloride (7.59 g, 63.8 mmol) dropwise and the reaction mixture
was stirred at room temperature over night. The mixture was poured onto ice/water (20
mL), basified with NaOH conc. (8 mL) and extracted with EtOAc (2 x 50 mL). Combined
organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was
10 purified by silica gel flash chromatography eluting with a 0 to 40% EtOAc-heptane
gradient to give the title compound (3.08 g, 93 %) as a white solid. MS: 206.0, 207.9
(M+H⁺).

[B] 3-Bromo-5-(2-isopropyl-imidazol-1-ylmethyl)-pyridine




To a suspension of sodium hydride (60% in mineral oil, 0.073 g, 1.82 mmol) in DMF (3
15 mL) was added 2-isopropyl-1H-imidazole (0.173 g, 1.57 mmol) and the reaction mixture
was stirred at room temperature for 20 min. Then, 3-bromo-5-chloromethyl-pyridine (0.25
g, 1.21 mmol) was added and the resulting suspension was heated at 60 °C over night. The
mixture was quenched with water (2 mL) and extracted with EtOAc (2x 10 mL).
Combined organics were dried over Na₂SO₄, filtered and concentrated in vacuo. The
20 residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-

DCM gradient to give the title compound (0.275 g, 81 %) as a light yellow oil. MS: 280.0, 282.0 ($M+H^+$).

Intermediate B-2

3-Bromo-5-(2-methyl-imidazol-1-ylmethyl)-pyridine




5

In analogy to the procedure described for the preparation of intermediates B-1 [B], 2-methyl-imidazol has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a light brown solid. MS: 251.9, 254.0 ($M+H^+$).

Intermediate B-3


10 3-Bromo-5-[1,2,4]triazol-1-ylmethyl-pyridine



In analogy to the procedure described for the preparation of intermediates B-1 [B], 1H-1,2,4-triazole has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a white solid. MS: 238.9, 241.3 ($M+H^+$).

Intermediate B-4

1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidin-2-one




15

In analogy to the procedure described for the preparation of intermediates B-1 [B], pyrrolidin-2-one has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a white solid. MS: 251.1, 257.1 ($M+H^+$).

Intermediate B-5


5 1-(5-Bromo-pyridin-3-ylmethyl)-piperidin-2-one



In analogy to the procedure described for the preparation of intermediates B-1 [B], piperidin-2-one has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a colorless oil. MS: 269.2, 271.2 ($M+H^+$).

10 Intermediate B-6


3-Bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridine



In analogy to the procedure described for the preparation of intermediates B-1 [B], (S)-2-(methoxymethyl)pyrrolidine has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a yellow oil. MS: 285.0, 286.9 ($M+H^+$).

Intermediate B-7


Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide



A flask was charged with 5-bromonicotinaldehyde (2.55 g, 13.7 mmol), ethanesulfonamide (2.99 g, 27.4 mmol) and toluene (250 mL), then titanium isopropoxide (5.84 g, 20.6 mmol) was added dropwise. The reaction mixture was stirred at 115 °C over
5 night and then concentrated in vacuo. The residue was taken up in DCM (200 mL) and MeOH (200 mL) and NaBH₄ (1.04 g, 27.4 mmol) was added portionwise at 0 °C. The reaction mixture was stirred at 0 °C for 30 min and then poured into water (100 mL) and the resulting suspension was filtered through a pad of dicalite. The dicalite layer was washed with DCM (3x 100 mL). The resulting aqueous layer was separated and extracted
10 with DCM (500 mL). Combined organics were dried over Na₂SO₄, filtered and preadsorbed on silica gel. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (3.01 g, 79%) as an orange solid. MS: 279.0, 281.0 (M+H⁺).

Intermediate B-8

15 **(S)-1-(5-Bromo-pyridin-3-yl)-5-hydroxymethyl-pyrrolidin-2-one**




In a sealed tube, 3,5-dibromopyridine (0.5 g, 2.11 mmol) was combined with (S)-5-(hydroxymethyl)pyrrolidin-2-one (0.243 g, 2.11 mmol), copper (I) iodide (0.040 g, 0.021 mmol), potassium carbonate (0.583 g, 4.22 mmol) and *N,N'*-dimethylethylenediamine
20 (0.037g, 0.042 mmol) in 1,4-dioxane (20 mL). The reaction mixture was heated at 110 °C over night. The mixture was cooled to room temperature, filtered through dicalite and washed with DCM. The residue was purified by silica gel flash chromatography eluting

with a 0 to 10% MeOH-DCM gradient to give the title compound (0.140 g, 25 %) as a light yellow oil. MS: 271.1, 273.1 ($M+H^+$).

Intermediate B-9

1-(5-Bromo-pyridin-3-yl)-pyrrolidin-2-one




5

In analogy to the procedure described for the preparation of intermediate B-8, pyrrolidin-2-one has been coupled to 3,5-dibromopyridine to yield the title compound as a light yellow solid. MS: 241.0, 243.0 ($M+H^+$).

Intermediate B-10

10 3-Bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridine




In analogy to the procedure described for the preparation of example 135 [B], (S)-2-methoxymethyl-pyrrolidine was reacted with 3,5-dibromopyridine in the presence of $Pd_2(dba)_3$, *rac*-BINAP and sodium *tert*-butoxide to give the title compound as a yellow oil.


15 MS: 271.1, 273.1 ($M+H^+$).

Intermediate B-11

4-Bromo-5,6,7,8-tetrahydroisoquinolin-8-ol




[A] Ethyl 5-bromo-4-methylnicotinate



To a stirred light brown suspension of 5-bromo-4-methylnicotinic acid (10.00 g, 46.3
5 mmol) and ethanol (2.97 mL) in DCM (231 mL) at 0 °C under argon was added EDCI
(10.9 g, 55.5 mmol) and DMAP (566 mg, 4.63 mmol); stirring was continued over night
and the reaction mixture was allowed to warm up to room temperature. The reaction
mixture was then poured on aq. 10% KH₂PO₄ solution followed by extraction with AcOEt
(3x). The organic phases were washed once with aq. 10% KH₂PO₄, with sat. NaHCO₃ and
10 with aq. sat. NaCl solution. The combined organic phases were dried (Na₂SO₄), filtered
and evaporated to afford the title compound (9.49 g, 84%) as brown solid. MS: 244.0
(M+H⁺, 1Br).


[B] Methyl 4-bromo-8-oxo-5,6,7,8-tetrahydroisoquinoline-7-carboxylate



15 Ethyl 5-bromo-4-methylnicotinate (7.04 g, 28.8 mmol) in THF (28.8 mL) was added over
a period of 20 min to a solution of LDA (31.7 mmol) [generated from *N,N*-
diisopropylamine (4.52 mL, 31.7 mmol) and *n*-butyllithium (19.8 mL, 31.7 mmol, 1.6M in
hexane) in 144 mL THF] at -78 °C. The resulting dark red solution was stirred for 20 min,


then methyl acrylate (6.21 g, 72.1 mmol) in THF (28.8 mL) was added over 15 min. The reaction was stirred an additional 1.5 h, then aq. 10% AcOH was added (resulting pH = 4-5) and the reaction was allowed to warm to room temperature. After evaporation, the residue was poured on aq. sat. NaHCO₃ and extracted with EtOAc (3x). The combined organic layers were dried (Na₂SO₄) and concentrated to afford the title compound (7.80 g, 95% in 70% purity with 30% starting material) as brown solid. MS: 284.0 (M+H⁺, 1Br).

[C] 4-Bromo-6,7-dihydroisoquinolin-8(5H)-one



The crude methyl 4-bromo-8-oxo-5,6,7,8-tetrahydroisoquinoline-7-carboxylate (7.79 g, 27.4 mmol) was dissolved (small amount of not dissolved material) in aq. 6M HCl (84.1 ml, 505 mmol) and heated at reflux for 2.5 h. The acidic solution was concentrated in vacuo, suspended in water (ca. 25 mL), cooled in ice and basified with 6.0 M KOH. The aqueous solution was extracted with ether (2x) and CH₂Cl₂ (3x), the combined organic layers were dried over Na₂SO₄, filtered and concentrated to afford the title compound (4.30 g, 69%) as a brown solid. MS: 226.0 (M+H⁺, 1Br).

[D] 4-Bromo-5,6,7,8-tetrahydroisoquinolin-8-ol




4-Bromo-6,7-dihydroisoquinolin-8(5H)-one (2.135 g, 9.44 mmol) was suspended in MeOH (18.9 mL), cooled to 0 °C and treated with NaBH₄ (357 mg, 9.44 mmol) in 5 portions over 30 min; then, the reaction was stirred for 3/4 h at 0 °C (TLC after 30 min showed no more starting material). AcOH was then added dropwise until pH ~ 5-6 and the reaction mixture was evaporated. The residue was diluted with water and poured on aq.


sat. NaHCO_3 -sol. followed by extraction with EtOAc (3x). The organic layers were washed once with aq. sat. NaHCO_3 -sol. and aq. 10% NaCl-sol., dried over Na_2SO_4 and concentrated in vacuo. The residue was re-dissolved in CH_2Cl_2 , evaporated with n-pentane (3x) to around 20 ml each time, while the product precipitates; then, the solvent was
5 decanted and the residue washed with n-pentane and dried in high vacuum to afford the title compound (1.98 g, 92%) as dark brown viscous oil. MS: 227 (M^+ , 1Br).

Intermediate B-12

N-(4-Bromo-5,6,7,8-tetrahydroisoquinolin-8-yl)propionamide




10 **[A] 4-Bromo-5,6,7,8-tetrahydroisoquinolin-8-amine**



4-Bromo-6,7-dihydroisoquinolin-8(5H)-one (intermediate B-11 [C]) (4.81g, 21.3 mmol), titanium (IV) isopropoxide (12.1 g, 42.6 mmol) and ammonia, 2.0 M solution in MeOH (53.2 mL, 106 mmol) were stirred at room temperature for 5 h. The reaction was cooled to
15 0 °C and NaBH_4 (1.21 g, 31.9 mmol) was added portionwise over 10 min; the resulting mixture was stirred at room temperature for an additional 2 h. The reaction was quenched by pouring it into aq. ammonium hydroxide 25%, the precipitate was filtered and washed with EtOAc (3x, each time suspended in AcOEt and stirred for 5 min). The organic layer was separated and the remaining aqueous layer was extracted with EtOAc. The combined
20 organic extracts were extracted with 1 M HCl. The acidic aqueous extracts were washed with ethyl acetate (1x), then treated with aqueous sodium hydroxide (2 M) to give pH 10-12 and extracted with EtOAc (3x). The combined organic extracts were washed with brine,

dried (Na_2SO_4), and concentrated in vacuo to afford the title compound (4.11 g, 85%) as brown solid. MS: 225 (M^+ , 1Br).


[B] N-(4-Bromo-5,6,7,8-tetrahydroisoquinolin-8-yl)propionamide




- 5 To a stirred brown solution of 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-amine (318 mg, 1.4 mmol) and propionic acid (114 mg, 1.54 mmol) in CH_2Cl_2 (7.0 mL) at 0 °C under argon was added EDCI (63.3 mg, 0.330 μmol). Stirring was continued over night and the reaction mixture was allowed to warm up to room temperature. The reaction mixture was poured onto aq. 10% KH_2PO_4 solution followed by extraction with AcOEt (3x). The
- 10 organic phases were washed once with aq. 10% KH_2PO_4 , aq. sat. NaHCO_3 and with aq. sat. NaCl solution; the combined organic phases were dried over Na_2SO_4 , filtered and evaporated. The residue was dissolved in CH_2Cl_2 , evaporated with n-pentane (3x) to ca. 10 ml each time, while the product precipitated; then, the solvent was decanted and the residue washed twice with n-pentane to afford the title compound (0.365 g, 92%) as light
- 15 brown solid. MS: 283.0 ($\text{M}+\text{H}^+$, 1Br).

Intermediate B-13


2-(5-Bromo-pyridin-3-ylmethyl)-[1,2]thiazinane 1,1-dioxide



- In analogy to the procedure described for the preparation of intermediates B-1 [B],
- 20 [1,2]thiazinane 1,1-dioxide has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as off-white solid. MS: 305.1, 307.1 ($\text{M}+\text{H}^+$).


Intermediate B-14**3-Bromo-5-(1,1-dioxo-1⁶-isothiazolidin-2-ylmethyl)-pyridine**

In analogy to the procedure described for the preparation of intermediates B-1 [B],
5 isothiazolidine 1,1-dioxide has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as a light yellow oil. MS: 290.9, 292.8 ($M+H^+$).

Intermediate B-15**(S)-1-(5-Bromo-pyridin-3-ylmethyl)-5-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one**
10

In analogy to the procedure described for the preparation of intermediates B-1 [B], (S)-5-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one has been coupled to 3-bromo-5-chloromethyl-pyridine (intermediate B-1 [A]) to yield the title compound as colorless oil.
15 MS: 399.2, 401.2 ($M+H^+$).


Intermediate B-16**(S)-1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidine-2-carboxylic acid methyl ester**



In analogy to the procedure described for the preparation of intermediates B-1 [B], (S)-pyrrolidine-2-carboxylic acid methyl ester has been coupled to 3-bromo-5-chloromethylpyridine (intermediate B-1 [A]) to yield the title compound as light yellow oil. MS: 299.2, 301.1 ($M+H^+$).

Intermediate B-17


N-(4-Bromo-5,6,7,8-tetrahydro-isoquinolin-8-yl)-acetamide




To a stirred solution of 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-amine (intermediate B-12 [A], 318 mg, 1.4 mmol) and Et_3N (3.0 mL) in DCM (10 mL) was added acetyl chloride (0.106 mL, 1.4 mmol) at 0 °C and stirring was continued at 0 °C for 1 hour. The resulting mixture was extracted with EtOAc (2 x 100 mL). The combined organics were washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated *in vacuo* to afford a crude product which was purified by silica gel flash chromatography eluting with a 0 to 50 % EtOAc-heptane gradient to give the title compound (346 mg, 92 %) as light yellow solid. MS: 269.1 & 271.1 ($M+H^+$).

Intermediate B-18

N-(4-Bromo-6,7-dihydro-5H-[2]pyrindin-7-yl)-propionamide




[A] Ethyl 3-(3,5-dibromopyridin-4-yl)propanoate



A solution of 3,5-dibromo-4-methylpyridine (20 g, 0.08 mol) in THF (50 mL) was added
 5 over a period of 1 hour to a solution of LDA (0.088 mol) [generated from N,N-
 diisopropylamine (8.9 g, 0.088 mol) and n-butyllithium (35 mL, 0.088 mol, 2.5 M in
 hexane) in 400 mL THF] at -78°C. The resulting dark red solution was stirred at -78 °C for
 30 min before ethylbromoacetate (33.4 g, 0.2 mol) in THF (50 mL) was added over a
 period of 30 min. The reaction mixture was stirred for an additional 2.5 hour at -78 °C
 10 before 10% AcOH was added (resulting in a pH = 4-5). The reaction mixture was then
 allowed to warm up to room temperature. After evaporation of the solvents, the residue
 was poured into satd. aq. NaHCO₃ solution and extracted with EtOAc (100 mL x 3). The
 combined organic layers were dried over anhy. Na₂SO₄, filtered, and concentrated in
 vacuo. The residue was purified by column chromatography to afford the title compound
 15 (9 g, 33.5 %) as a yellow solid. MS: 337.7(M+H⁺, 2Br).

[B] 4-Bromo-5H-cyclopenta[c]pyridin-7(6H)-one




To a solution of ethyl 3-(3,5-dibromopyridin-4-yl)propanoate (4 g, 11.9 mmol) in THF (60
 mL) was slowly added n-BuLi (9.52 mL, 23.8 mmol, 2.5 M in hexane) while keeping the

inner temperature below -70 °C. After the addition, the reaction mixture was stirred below -65 °C for additional 2 hours. The reaction was quenched with water and then extracted with EtOAc (100 mL x 3). The combined organic layers were dried over anhy. Na₂SO₄, filtered, and concentrated in vacuo. The residue was then purified by column


- 5 chromatography to afford the title compound (1.2 g, 47.6%) as a solid. MS: MS: 213.7 (M+H⁺, 1Br).

[C] 4-Bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine



- CH₃COONH₄ (4.8 g, 62 mmol), NaCNBH₃ (315 mg, 5 mmol) and 4-bromo-5H-
- 10 cyclopenta[c]pyridin-7(6H)-one (1.0 g, 4.15 mmol) in EtOH (10 mL) was added to a sealed microwave vial. The mixture was subject to microwave radiation at 130 °C for 4 min. After the majority of EtOH was evaporated under reduced pressure, it was treated with aq. 2 N NaOH until pH >10 and extracted with EtOAc (50 mL x2). The combined organic layers were dried over anhy. Na₂SO₄, filtered, and concentrated under reduced
- 15 pressure to give a crude title product (800 mg) which was used in the next step reaction without further purification. MS: 212.9 (M+H⁺, 1Br).

[D] N-(4-Bromo-6,7-dihydro-5H-cyclopenta[c]pyridin-7-yl)propionamide




- To a solution of 4-bromo-6,7-dihydro-5H-cyclopenta[c]pyridin-7-amine (650 mg, 3.05
- 20 mmol, 65% purity) and Et₃N (400 mg, 3.97 mmol) in THF (20 mL) was added propionyl chloride (219 mg, 2.38 mmol) at 0 °C and stirring continued for 2 hours. Water (10 mL) and EtOAc (10 mL) were added and the aqueous layer was extracted with EtOAc (10 mL x3). The combined organic layers were dried over anhy. Na₂SO₄, filtered and concentrated

in vacuo to give a crude product which was purified by column chromatography to afford the title compound (200 mg, 37.6%) as a white solid. MS: 270.9 ($M+H^+$, 1Br).

Intermediate B-19

N-(4-Bromo-6,7-dihydro-5H-[2]pyrindin-7-yl)-acetamide




5


In analogy to the procedure describe for the preparation of intermediate B-18 (step D), the title compound was obtained as a white solid using acetyl chloride as starting material. MS: 256.9 ($M+H^+$, 1Br).

10 **Example 1**


6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one




In a sealed tube, 3-bromopyridine (0.1 g, 0.633 mmol) was combined with 6-chloro-3,4-dihydroisoquinolin-1(2H)-one (intermediate A-2) (0.115 g, 633 mmol), copper (I) iodide
15 (0.012 g, 0.063 mmol), potassium carbonate (0.175 g, 1.27 mmol) and *N,N'*-dimethylethylenediamine (0.013g, 0.127 mmol) in 1,4-dioxane (2 mL). The reaction mixture was heated at 110 °C over night. The mixture was cooled to room temperature, filtered through dicalite and washed with DCM. The residue was purified by silica gel flash chromatography eluting with a 5 to 100% EtOAc-heptane gradient to give the title
20 compound (0.107 g, 65 %) as a white solid. MS: 259.1 ($M+H^+$).

Example 2**5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-nicotinonitrile**

6-Chloro-3,4-dihydro-2*H*-isoquinolin-1-one (intermediate A-2) (36 mg, 0.2 mmol), 5-bromo-nicotinonitrile (73 mg, 0.4 mmol), CuI (3.8 mg, 0.02 mmol), (1*S*, 2*S*)-cyclohexane-1,2-diamine (4.5 mg, 0.04 mmol) and Cs₂CO₃ (130 mg, 0.4 mmol) were dissolved in dioxane (5 mL). The reaction mixture was subjected to microwave reaction at 150 °C for 2.5 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2x 25 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to yield the title compound (15 mg, 26%) as a white solid. MS: 284.1 (M+H⁺).

Example 3**6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2*H*-isoquinolin-1-one**


15 [A] (5-Bromo-pyridin-3-yl)-methanol



Sodium borohydride (2.2 g, 59.1 mmol) was added at 0 °C to a suspension of 5-bromo-pyridine-3-carbaldehyde (10.0 g, 53.7 mmol) in MeOH (100 mL). The mixture was stirred at 0 °C for 1 hour before it was quenched by addition of water (5.0 mL). Evaporation of

solvents afforded a light yellowish oil which was re-dissolved in EtOAc and washed with water. The organic layer was dried over Na₂SO₄, filtered and concentrated in vacuo to give the title compound (9.6 g, 95%) as colorless oil. MS: 188.0 & 190.0 (M+H⁺).

5 [B] 6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one







In a 25 mL sealed tube, (5-bromo-pyridin-3-yl)-methanol (900 mg, 4.8 mmol), 6-chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) (800 mg, 4.4 mmol), CuI (200 mg, 1.1 mmol), Cs₂CO₃ (3.0 g, 9.2 mmol) and (+)-(S,S)-1,2-diaminocyclohexane (0.4 mL, 3.2 mmol) were dissolved in dioxane (8.0 mL). The resulting reaction mixture was heated at 150 °C for 3 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 125 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by silica gel flash chromatography (30-100% EtOAc-hexane gradient) to yield the title compound (1.1 g, 90%) as a light yellow solid. MS: 289.2 (M+H⁺).



The following compounds listed in **Table 1** were prepared in analogy to the procedures described for the preparation of examples 1, 2 or 3[B] using appropriate starting materials.



Table 1



Ex	Compound Name	Compound Structure	Starting Materials	Prepared by analogy to example	MS (M+H) ⁺



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
41	6-Chloro-2-(5-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  293.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 1-bromo-3-chloro-benzene	2
51	6-Chloro-2-(5-fluoro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  277.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-fluoro-pyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
6	6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  293.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-chloro-pyridine	2
7	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one  337.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-iodo-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
31	6-Chloro-2-(5-methyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  273.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-methyl-pyridine	3[B]
2	5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde  287.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 5-bromopyridine-3-carbaldehyde	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
10	6-Chloro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one 	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-methoxy-pyridine	1
11	6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  317.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-isopropoxy-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>12</u>	6-Chloro-2-[5-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  317.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 2-(5-bromopyridin-3-yl)propan-2-ol	2
<u>13</u>	6-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)ethanol	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
14	6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (R)-1-1-(5-bromo-pyridin-3-yl)-ethanol	2
15	6-Chloro-2-[5-((S)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (S)-1-1-(5-bromo-pyridin-3-yl)-ethanol	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
16	6-Chloro-2-[5-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  317.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(1-methoxy-ethyl)-pyridine	2
17	2-(5-Aminopyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one  274.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 5-bromopyridin-3-ylamine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
18	6-Chloro-2-[5-(2,2,2-trifluoro-1-hydroxyethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  357.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)-2,2,2-trifluoro-ethanol	2
19	6-Chloro-2-[5-(2,2,2-trifluoro-1-methoxyethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  371.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(2,2,2-trifluoro-1-methoxy-ethyl)-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
20	6-Chloro-2-[5-(cyclopropyl-hydroxy-methyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  329.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-cyclopropyl-methanol	2
21	6-Chloro-2-[5-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  343.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(cyclopropyl-methoxy-methyl)-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
22	6-Chloro-2-(4-trifluoromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  327.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-trifluoromethyl-pyridine	3[B]
23	6-Chloro-2-[5-(2-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 2-(5-bromopyridin-3-yl)-ethanol	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
24	6-Chloro-2-[5-(1-methoxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  331.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(1-methoxy-1-methyl-ethyl)-pyridine	2
25	Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide  380.1	6-Chloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one (intermediate A-2) and ethanesulfonic acid (5-bromopyridin-3-ylmethyl)-amide (intermediate B-7)	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
26	6-Chloro-2-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  342.0	6-Chloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)-pyrrolidin-2-one (intermediate B-9)	1
27	6-Chloro-2-(1-methyl-1<i>H</i>-pyrazolo[3,4-<i>c</i>]pyridin-4-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  313.2	6-Chloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one (intermediate A-2) and 4-bromo-1-methyl-1<i>H</i>-pyrazolo[3,4-<i>c</i>]pyridine	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
28	6-Chloro-8'-hydroxy-3,4,5',6',7',8'-hexahydro-[2,4']biisoquinoliny-1-one  329.1	6-Chloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one (intermediate A-2) and 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-ol (intermediate B-11)	1
29	<i>N</i>-(6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-propionamide  384.1	6-Chloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one (intermediate A-2) and <i>N</i>-(4-bromo-5,6,7,8-tetrahydroisoquinolin-8-yl)propionamide (intermediate B-12)	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
30	6-Chloro-2-{5-[hydroxy-(1-methyl-1<i>H</i>-imidazol-2-yl)-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  369.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-(1-methyl-1<i>H</i>-imidazol-2-yl)-methanol	3[B]
31	6-Chloro-2-{5-[(3,4-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  401.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-(3,4-difluoro-phenyl)-methanol	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
32	6-Chloro-2-{5-[(3,5-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  401.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-(3,5-difluoro-phenyl)-methanol	3[B]
33	6-Chloro-2-{5-[(4-ethyl-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  393.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-(4-ethyl-phenyl)-methanol	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
34	6-Chloro-2-[5-(hydroxy-phenyl-methyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  365.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-phenyl-methanol	3[B]
35	6-Chloro-2-[5-(1-hydroxy-1-phenyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  379.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)-1-phenyl-ethanol	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
36	6-Chloro-2-{5-[1-(3,4-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  415.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)-1-(3,4-difluorophenyl)-ethanol	3[B]
37	6-Chloro-2-{5-[1-(3,5-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  415.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 1-(5-bromopyridin-3-yl)-1-(3,5-difluorophenyl)-ethanol	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
38	6-Chloro-2-(6-methyl-pyrazin-2-yl)-3,4-dihydro-2H-isoquinolin-1-one  274.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 2-bromo-6-methyl-pyrazine	3[B]
39	6-Chloro-2-[5-(morpholine-4-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  372.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (5-bromo-pyridin-3-yl)-morpholin-4-yl-methanone	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
40	6-Chloro-2-[5-(3-hydroxy-pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  372.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-(3-hydroxy-pyrrolidin-1-yl)-methanone	2
41	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-<i>N,N</i>-dimethyl-nicotinamide  330.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 5-bromo-<i>N,N</i>-dimethylnicotinamide	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
42	6-Chloro-2-[5-(pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  356.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (5-bromopyridin-3-yl)-pyrrolidin-1-yl-methanone	2
43	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-<i>N</i>-methyl-nicotinamide  316.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 5-bromo-<i>N</i>-methyl-nicotinamide	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
44	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-<i>N</i>-cyclopropyl-nicotinamide  342.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 5-bromo-<i>N</i>-cyclopropyl-nicotinamide	2
45	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-<i>N</i>-(4-fluoro-phenyl)-nicotinamide  396.0	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 5-bromo-<i>N</i>-(4-fluoro-phenyl)-nicotinamide	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
46	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-<i>N</i>-phenyl-nicotinamide  378.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 5-bromo-<i>N</i>-phenyl-nicotinamide	2
47	6-Chloro-2-[5-(4,4-difluoro-piperidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (3-bromophenyl)-(4,4-difluoropiperidin-1-yl)-methanone	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
48	6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  386.3	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridine (intermediate B-6)	1
49	6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  372.0	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridine (intermediate B-10)	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
50	6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  372.0	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (S)-1-(5-bromopyridin-3-yl)-5-hydroxymethylpyrrolidin-2-one (intermediate B-8)	1
51	6-Chloro-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one  260.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 5-bromopyrimidine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
52	6-Chloro-2-pyridazin-3-yl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  260.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromopyridazine	2
53	6-Chloro-2-pyridin-3-yl-2<i>H</i>-isoquinolin-1-one  257.1	6-Chloro-2<i>H</i>-isoquinolin-1-one (intermediate A-6) and 3-bromopyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
54	6-Chloro-2-(5-fluoro-pyridin-3-yl)-2<i>H</i>-isoquinolin-1-one  275.1	6-Chloro-2<i>H</i>-isoquinolin-1-one (intermediate A-6) and 3-bromo-5-fluoro-pyridine	3[B]
55	6-Chloro-2-[4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 1-(3-bromopyridin-4-yl)-ethanol	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
56	6-Chloro-2-(4-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one; compound with trifluoro-acetic acid  289.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (3-bromo-4-pyridin-4-yl)-methanol	2
57	2-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  314.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 1-(5-bromo-pyridin-3-yl)-cyclopropylamine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
58	6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  339.2	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(4-methyl-4H-[1,2,4]triazol-3-yl)-pyridine	2
59	6-Chloro-2-(5-methylsulfanyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  305.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-methylsulfanyl-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
60	6-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  325.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-difluoromethoxy-pyridine	2
61	6-Chloro-2-(4-dimethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  333.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-dimethoxymethyl-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
62	6-Chloro-2-[5-fluoro-4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  321.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 1-(3-bromo-5-fluoro-pyridin-4-yl)-ethanol	2
63	6-Chloro-2-{4-[(4-fluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  383.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (3-bromopyridin-4-yl)-(4-fluoro-phenyl)-methanol	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
64	6-Chloro-2-[4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  317.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-(1-methoxy-ethyl)-pyridine	2
65	6-Chloro-2-(1-methyl-1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one  312.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 7-bromo-1-methyl-1H-pyrrolo[3,2-c]-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
66	6-Chloro-2-(5-cyclopropyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  299.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-cyclopropyl-pyridine	2
67	6-Chloro-2-[5-(2-methyl-2H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  340.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(2-methyl-2H-[1,2,4]triazol-3-yl)-pyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
68	6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  315.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-cyclopropoxy-pyridine	2
69	6-Chloro-2-(4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-methoxymethyl-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
70	6-Chloro-2-[5-fluoro-4-(1-hydroxy-1-methylethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  335.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 2-(3-bromo-5-fluoro-pyridin-4-yl)-propan-2-ol	2
71	6-Chloro-2-[5-(5-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(5-methyl-pyrazol-1-ylmethyl)-pyridine	3[B]


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
72	6-Chloro-2-(1<i>H</i>-pyrrolo[3,2-<i>c</i>]pyridin-7-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  298.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 7-bromo-1<i>H</i>-pyrrolo[3,2-<i>c</i>]pyridine	2
73	6-Chloro-3,4-dihydro-[2,4']biisoquinolinyl-1-one  309.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromoisoquinoline	3[B]


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
74	3-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-isonicotinonitrile  284.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-isonicotinonitrile	2
75	6-Chloro-2-(5-fluoro-4-methoxymethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  321.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-fluoro-4-methoxymethylpyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
76	6-Chloro-2-[5-fluoro-4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  335.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-fluoro-4-(1-methoxy-ethyl)-pyridine	2
77	6-Chloro-2-(4-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  333.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-isopropoxymethyl-pyridine	2



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
78	6-Chloro-2-[4-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  343.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-4-(cyclopropyl-methoxy-methyl)-pyridine	2
79	6-Chloro-2-[5-(3,5-dimethyl-1<i>H</i>-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(3,5-dimethyl-1<i>H</i>-pyrazol-4-yl)-pyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
80	6-Chloro-2-[5-(3,5-dimethyl-3<i>H</i>-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(3,5-dimethyl-3<i>H</i>-imidazol-4-yl)-pyridine	3[B]
81	6-Chloro-2-[5-(1,1-dioxo-1λ⁶-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.3	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 2-(5-bromopyridin-3-ylmethyl)-[1,2]thiazinane 1,1-dioxide (intermediate B-13)	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
82	6-Chloro-2-[5-(1,1-dioxo-1^λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  392.1	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and 3-bromo-5-(1,1-dioxo-1^λ6-isothiazolidin-2-ylmethyl)-pyridine (intermediate B-14)	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
83	6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  386.0	6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) and (S)-1-(5-bromopyridin-3-ylmethyl)-5-(<i>tert</i>-butyl-dimethyl-silyloxymethyl)-pyrrolidin-2-one (intermediate B-15) to 2-{5-[(S)-2-(<i>tert</i>-butyl-dimethyl-silyloxymethyl)-5-oxo-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (light brown amorphous solid, MS: 500.3, 502.3 (M+H⁺)) and subsequent removal of the protecting group (4M HCl in dioxane, MeOH 2h, RT).	1



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
84	(S)-1-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester  400.0	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and (S)-1-(5-bromopyridin-3-ylmethyl)-pyrrolidine-2-carboxylic acid methyl ester (intermediate B-16)	1
85	6-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  303.1	6-Chloro-3-methyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-3-1) and 3-bromo-5-methoxy-pyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
86	6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one  303.1	6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1) and (5-bromopyridin-3-yl)-methanol	3[B]
87	6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3-methyl-3,4-dihydro-2H-isoquinolin-1-one  395.3	6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1) and 3-bromo-5-(2-isopropyl-imidazol-1-ylmethyl)-pyridine (intermediate B-1)	3[B]

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
88	6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one  273.1	6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1) and 3-bromopyridine	3[B]
89	6-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one  291.1	6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1) and 3-bromo-5-fluoropyridine	3[B]


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
90	6-Chloro-3-methyl-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one  274.1	6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1) and 5-bromopyrimidine	3[B]
91	(R)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one  273.1	(R)-6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1a) and 3-bromopyridine	3[B]

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
92	(S)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one  273.1	(S)-6-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-1b) and 3-bromo-pyridine	3[B]
93	8-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one  273.1	8-Chloro-3-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-3-2) and 3-bromo-pyridine	3[B]



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
94	6-Methoxy-2-pyridin-3-yl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  255.2	6-Methoxy-3,4-dihydroisoquinolin-1(2<i>H</i>)-one and 3-bromopyridine	1
95	5,6-Dichloro-2-pyridin-3-yl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  293.0	5,6-Dichloro-3,4-dihydroisoquinolin-1(2<i>H</i>)-one and 3-iodopyridine	1

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
96	2-Chloro-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one  290.2	2-Chloro-7,8-dihydro-6H-[1,6]naphthyridin-5-one (intermediate A-8) and 3-bromo-5-methoxy-pyridine	2
97	2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one  286.2	2-Methoxy-7,8-dihydro-6H-[1,6]naphthyridin-5-one (intermediate A-9) and 3-bromo-5-methoxy-pyridine	2



- 236a -

Ex	Compound Name/Stucture	Starting Materials	Prepare d by analogy to example
28	2-Methoxy-6-pyridin-3-yl-7,8-dihydro-6H-[1,6]naphthyridin-5-one  256.1	2-Methoxy-7,8-dihydro-6H-[1,6]naphthyridin-5-one (intermediate A-9) and 3-bromopyridine	3[B]



- 236b -

99	6-Chloro-5-fluoro-2-(5-methoxy-pyridin-3-yl)- 3,4-dihydro-2<i>H</i>-isoquinolin-1-one  307.1	6-Chloro-5-fluoro- 3,4-dihydro-2<i>H</i>- isoquinolin-1-one (intermediate A-1) and 3-bromo-5- methoxy-pyridine	2
100	6-Chloro-7-fluoro-2-(5-methoxy-pyridin-3-yl)- 3,4-dihydro-2<i>H</i>-isoquinolin-1-one  307.1	6-Chloro-7-fluoro- 3,4-dihydro-2<i>H</i>- isoquinolin-1-one (intermediate A-7) and 3-bromo-5- methoxy-pyridine	2



- 236c -

101	6-Chloro-7-fluoro-2-pyridin-3-yl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  277.1	6-Chloro-7-fluoro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-7) and 3-bromopyridine	2
102	6-Chloro-4,4-dimethyl-2-pyridin-3-yl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  287.2	6-Chloro-4,4-dimethyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-5) and 3-bromopyridine	3[B]



- 236d -

<u>103</u>	6-Chloro-2-(5-methoxy-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  317.2	6-Chloro-4,4-dimethyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-5) and 3-bromo-5-methoxy-pyridine	3[B]
<u>104</u>	6-Chloro-2-(5-fluoro-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  305.2	6-Chloro-4,4-dimethyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-5) and 3-bromo-5-fluoro-pyridine	3[B]



- 236e -

105	6-Chloro-4-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one  273.1	6-Chloro-4-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-4) and 3-bromopyridine	3[B]
106	6-Chloro-2-(5-fluoro-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one  291.1	6-Chloro-4-methyl-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-4) and 3-bromo-5-fluoro-pyridine	3[B]



- 236f -

107	6-Chloro-2-(5-methoxy-pyridin-3-yl)-4-methyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  303.2	6-Chloro-4-methyl-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-4) and 3-bromo-5-methoxy-pyridine	3[B]
108	5-Chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  245.1	5-Chloro-2,3-dihydro-isoindol-1-one and 3-iodopyridine	1



- 236g -

109	5-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  367.0	5-Chloro-2,3-dihydro-isoindol-1-one and 3-bromo-5-(2-isopropyl-imidazol-1-ylmethyl)-pyridine (intermediate B-1)	1
110	5-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  326.2	5-Chloro-2,3-dihydro-isoindol-1-one and 3-bromo-5-[1,2,4]triazol-1-ylmethyl-pyridine (intermediate B-3)	1



- 236h -

<u>111</u>	5-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  339.1	5-Chloro-2,3-dihydro-isoindol-1-one and 3-bromo-5-(2-methyl-imidazol-1-ylmethyl)-pyridine (intermediate B-2)	1
<u>112</u>	5-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  342.1	5-Chloro-2,3-dihydro-isoindol-1-one and 1-(5-bromopyridin-3-ylmethyl)-pyrrolidin-2-one (intermediate B-4)	1



- 236i -

113	5-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  372.2	5-Chloro-2,3-dihydro-isoindol-1-one and 3-bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridine (intermediate B-6)	1
114	5-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  356.3	5-Chloro-2,3-dihydro-isoindol-1-one and 1-(5-bromopyridin-3-ylmethyl)-piperidin-2-one (intermediate B-5)	1



- 236j -

115	Ethanesulfonic acid [5-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  366.0	5-Chloro-2,3-dihydro-isoindol-1-one and ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate B-7)	1
116	5-Chloro-2-(1-methyl-1<i>H</i>-pyrazolo[3,4-<i>c</i>]pyridin-4-yl)-2,3-dihydro-isoindol-1-one  299.2	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-1-methyl-1<i>H</i>-pyrazolo[3,4-<i>c</i>]pyridine	1



- 236k -

117	5-Chloro-2-(8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one  315.0	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-ol (intermediate B-11)	1
118	5-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  259.1	5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-10) and 3-bromopyridine	3[B]



- 2361 -

<u>119</u>	6-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  259.1	6-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-11) and 3-bromopyridine	3[B]
<u>120</u>	5-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one  289.1	5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-10) and 3-bromo-5-methoxy-pyridine	2



- 236m -

121	5-Chloro-3-methyl-2-(4-methyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  273.1	5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-10) and 3-bromo-5-methyl-pyridine	2
122	5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3-methyl-2,3-dihydro-isoindol-1-one  303.1	5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-10) and 1-(5-bromo-pyridin-3-yl)-ethanol	2



- 236n -

<u>123</u>	5-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one  277.1	5-Chloro-3-methyl-2,3-dihydro-isoindol-1-one (intermediate A-10) and 3-bromo-5-fluoro-pyridine	2
<u>124</u>	3-Benzyl-5-chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  335.1	3-Benzyl-5-chloro-2,3-dihydro-isoindol-1-one (intermediate A-14) and 3-bromopyridine	2



- 236o -

<u>125</u>	5-Chloro-3-ethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  273.2	5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13) and 3-bromopyridine	2
<u>126</u>	5-Chloro-3-ethyl-2-(5-fluoropyridin-3-yl)-2,3-dihydro-isoindol-1-one  291.2	5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13) and 3-bromo-5-fluoropyridine	2



- 236p -

<u>127</u>	5-Chloro-3-ethyl-2-(5-methoxy-pyridin-3-yl)- 2,3-dihydro-isoindol-1-one  303.2	5-Chloro-3-ethyl-2,3- dihydro-isoindol-1- one (intermediate A- 13) and 3-bromo-5- methoxy-pyridine	2
<u>128</u>	5-Chloro-3,3-dimethyl-2-pyridin-3-yl-2,3- dihydro-isoindol-1-one  273.1	5-Chloro-3,3- dimethyl-2,3- dihydro-isoindol-1- one (intermediate A- 12) and 3- bromopyridine	3[B]



- 236q -

<u>129</u>	5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  291.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-fluoro-pyridine	3[B]
<u>130</u>	5-Chloro-2-(5-methoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  303.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-methoxy-pyridine	3[B]

- 236r -


<u>131</u>	5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  339.2	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-difluoromethoxy-pyridine	2
<u>132</u>	5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one  317.2	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 1-(5-bromopyridin-3-yl)-ethanol	2

- 236s -


133	5-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  303.2	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and (5-bromopyridin-3-yl)-methanol	2
134	5-Chloro-3,3-dimethyl-2-[5-(2-methylimidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  367.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-(2-methylimidazol-1-ylmethyl)-pyridine (intermediate B-2)	3[B]

Example 135

6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one




[A] (S)-2-(*tert*-Butyl-dimethyl-silyloxymethyl)-pyrrolidine



To a solution of (S)-pyrrolidin-2-ylmethanol (0.69 g, 6.82 mmol) in DCM (3 mL) cooled
 5 to 0 °C was added TEA (1.38 g, 13.6 mmol) followed by TBDMS-Cl (1.03 g, 6.82 mmol)
 in DCM (3 mL). The reaction mixture was then stirred at room temperature over night and
 poured into NH₄Cl (20 mL). The aqueous layer was extracted with DCM (2 x 50 mL).
 Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to
 dryness to give the title compound (1.11g, 76%) as a yellow oil. MS: 216.2 (M+H⁺).


10 [B] 3-Bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridine



To a solution of crude (S)-2-((*tert*-butyldimethylsilyloxy)methyl)pyrrolidine (0.455 g, 2.11
 mmol) in toluene (20 mL) were added Pd₂(dba)₃ (0.039 g, 0.042 mmol) and *rac*-BINAP
 (0.066 g, 0.106 mmol). The solution was purged with argon and heated to 85 °C for 10
 15 min. After cooling to room temperature, sodium *tert*-butoxide (0.406 g, 4.22 mmol) and
 3,5-dibromopyridine (0.5 g, 2.11 mmol) were added and the reaction mixture was then


heated at 85 °C over night. The mixture was poured into sat NH₄Cl (20 mL) and the aqueous layer was extracted with DCM (2x 25mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the
 5 title compound (0.412 g, 53 %) as a yellow oil. MS: 371.0, 372.9 (M+H⁺).

[C] 2-{5-[(S)-2-(*tert*-Butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one



In analogy to the procedure described for the preparation of intermediate B-8, 6-chloro-
 10 3,4-dihydroisoquinolin-1(2*H*)-one (intermediate A-2) has been coupled to 3-bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridine to yield the title compound as a yellow oil. MS: 472.2 (M+H⁺).

[D] 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




15

To a solution of 2-{5-[(S)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (0.110 g, 0.233 mmol) in MeOH (2 mL) was added 4M HCl in dioxane (0.233 ml, 0.932 mmol) and the reaction mixture was stirred at room temperature for 2h. The mixture was evaporated to dryness, the residue
 20 was diluted with DCM (20 mL) and washed with NaHCO₃ (10 mL). The organic layer was dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica

gel flash chromatography eluting with a 0 to 5% MeOH(1% NH₄OH)-DCM gradient to give the title compound (0.06 g, 72 %) as a light yellow solid. MS: 358.0 (M+H⁺).

Example 136


5-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one




In analogy to the procedures described for the preparation of example 135, 3-bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridine (example 135[B]) has been reacted with 5-chloro-2,3-dihydro-isoindol-1-one to give 2-{5-[(S)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-5-chloro-2,3-dihydro-isoindol-1-one, which was then deprotected with 4M HCl in dioxane to give the title compound as white solid. MS: 344.0 (M+H⁺).

Example 137

6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one




[A] (R)-3-bromo-5-(3-(*tert*-butyldimethylsilyloxy)pyrrolidin-1-yl)pyridine



In analogy to the procedure described for the preparation of example 135 [A], (R)-pyrrolidin-3-ol was reacted with TBDMS-Cl in the presence of TEA to give the title compound as a yellow oil. MS: 202.2 ($M+H^+$).


5 [B] 3-Bromo-5-[(R)-3-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridine



In analogy to the procedure described for the preparation of 135 [B], (R)-3-bromo-5-(3-(*tert*-butyldimethylsilyloxy)pyrrolidin-1-yl)pyridine was reacted with 3,5-dibromopyridine in the presence of $Pd_2(dba)_3$, *rac*-BINAP and sodium *tert*-butoxide to give the title


10 compound as a yellow oil. MS: 357.1, 359.0 ($M+H^+$).

[C] 2-{5-[(R)-3-(*tert*-Butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridin-3-yl}-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one



In analogy to the procedure described for the preparation of intermediate B-8, 6-chloro-3,4-dihydroisoquinolin-1(2*H*)-one (intermediate A-2) has been coupled to 3-bromo-5-[(R)-3-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridine to give the title compound as a yellow oil. MS: 458.2 ($M+H^+$).


[D] 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one



In analogy to the procedure described in example 135 [D], 2-{5-[(R)-3-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridin-3-yl}-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one was deprotected with 4M HCl in dioxane to give the title compound as a white
5 foam. MS: 344.0 ($M+H^+$).

Example 138


5-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one



10 In analogy to the procedures described for the preparation of example 137, 3-bromo-5-[(R)-3-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridine (example 137 [B]) has been reacted with 5-chloro-2,3-dihydro-isoindol-1-one to give 2-{5-[(R)-3-(*tert*-butyl-dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridin-3-yl}-5-chloro-2,3-dihydro-isoindol-1-one, which was then deprotected with 4M HCl in dioxane to give the title compound as light
15 yellow solid. MS: 330.1 ($M+H^+$).

Example 139


2-Hydroxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6*H*-[1,6]naphthyridin-5-one




2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5one (445 mg, 1.56 mmol) (example 97) was suspended in 2 mL of aq. conc. HCl in 1,4-dioxane (2:1) and heated at 80 °C for 3 hours before the solvent was removed in vacuo. After extraction
5 between EtOAc and H₂O, the organic layer was washed with aq. satd. NaHCO₃ solution, brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to yield the title compound (4.9 mg) as a white solid. MS: 270.2 (M+H⁺).

Example 140

10 6-Chloro-2-(5-imidazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one




[A] 6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one



Thionyl chloride (1.4 mL, 19.0 mmol) was added slowly at 0 °C to a solution of 6-chloro-
15 2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 3 [B]) (1.1 g, 3.8 mmol) in DCM (50 mL). After the addition, the reaction mixture was stirred at 2-5 °C for 2 hours before it was poured into satd. aq. NaHCO₃ solution (50 mL) and extracted with EtOAc (2 x 150 mL). The organic layer was washed with brine, dried over anhy.

Na₂SO₄, filtered and concentrated in vacuo to give a crude product (1.42 g, 92%) as light yellowish solid. MS: 307.0 & 309.0 (M+H⁺).



[B] 6-Chloro-2-(5-imidazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1 one







- 5 To a solution of 1H-imidazole (50 mg, 0.73 mmol) dissolved in DMF (5.0 mL) was added NaH (60% dispersion in mineral oil, 25 mg, 0.63 mmol) at 0 °C. The reaction mixture was stirred at 2-5 °C for 0.5 h before 6-chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140 [A]) (85 mg, 0.28 mmol) was added. The mixture was then stirred at 2-5 °C for additional 2 hours before it was warmed up to room temperature
- 10 and poured into water (5.0 mL). After extraction with EtOAc (2 x 50 mL), the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to give the title compound (30 mg, 31%) as a white foam. MS: 339.2 (M+H⁺).
- 15 The following compounds listed in Table 2 were prepared in analogy to the procedure described for the preparation of example 140 using appropriate starting materials.



Table 2



Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>141</u>	6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  381.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-isopropyl-1<i>H</i>-imidazole
<u>142</u>	6-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-methyl-1<i>H</i>-imidazole



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>143</u>	6-Chloro-2-[5-(2-ethyl-4-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  381.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-ethyl-4-methyl-1<i>H</i>-imidazole
<u>144</u>	6-Chloro-2-[5-(3-hydroxy-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  372.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and piperidin-3-ol



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>145</u>	Propane-2-sulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide  394.2	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and propane-2-sulfonic acid amide
<u>146</u>	6-Chloro-2-[5-(3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  358.2	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and pyrrolidin-3-ol



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>147</u>	6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  358.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and (R)-pyrrolidin-3-ol
<u>148</u>	6-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  358.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and (S)-pyrrolidin-3-ol



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>149</u>	6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  372.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and (S)-1-pyrrolidin-2-yl-methanol
<u>150</u>	6-Chloro-2-[5-((R)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  372.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and (R)-1-pyrrolidin-2-yl-methanol



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>151</u>	6-Chloro-2-[5-(3,5-dimethyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  367.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3,5-dimethyl-1<i>H</i>-pyrazole
<u>152</u>	N-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  366.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and methanesulfonamide



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>153</u>	<i>N</i>-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-acetamide  330.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and acetamide
<u>154</u>	6-Chloro-2-(5-morpholin-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  358.3	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and morpholine



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>155</u>	6-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  356.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and pyrrolidin-2-one
<u>156</u>	6-Chloro-2-[5-(1,1-dioxo-1λ⁶-thiomorpholin-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and thiomorpholine 1,1-dioxide



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>157</u>	6-Chloro-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  358.1	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and oxazolidin-2-one
<u>158</u>	6-Chloro-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  357.1	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and imidazolidin-2-one



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>159</u>	6-Chloro-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  371.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1-methyl-imidazolidin-2-one
<u>160</u>	6-Chloro-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  339.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1<i>H</i>-pyrazole



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>161</u>	6-Chloro-2-[5-(2-propyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  381.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-propyl-1<i>H</i>-imidazole
<u>162</u>	1-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylmethyl]-1<i>H</i>-imidazole-2-carboxylic acid ethyl ester  411.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1<i>H</i>-imidazole-2-carboxylic acid ethyl ester



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>163</u>	6-Chloro-2-[5-(2-hydroxymethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  369.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and (1<i>H</i>-imidazol-2-yl)-methanol
<u>164</u>	6-Chloro-2-[5-(oxetan-3-ylaminomethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  344.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and oxetan-3-ylamine

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>165</u>	6-Chloro-2-{5-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  401.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and 2-piperazin-1-yl-ethanol
<u>166</u>	6-Chloro-2-[5-(4-isopropyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  399.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and 1-isopropyl-piperazine



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>167</u>	6-Chloro-2-[5-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  371.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1-methyl-piperazine
<u>168</u>	6-Chloro-2-[5-(4,4-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  392.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 4,4-difluoro-piperidine

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>169</u>	6-Chloro-2-[5-(3,3-difluoro-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  378.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3,3-difluoro-pyrrolidine
<u>170</u>	6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  384.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-oxa-6-aza-spiro[3.4]octane


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>171</u>	6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.3]hept-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  370.2	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-oxa-6-aza-spiro[3.3]heptane
<u>172</u>	6-Chloro-2-[5-(3,3-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  392.2	6-Chloro-2-(5-chloromethylpyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3,3-difluoropiperidine

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>173</u>	6-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  370.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and piperidin-2-one
<u>174</u>	6-Chloro-2-(5-[1,2,3]triazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  340.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2<i>H</i>-[1,2,3]triazole



- 260a -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>175</u>	6-Chloro-2-(5-[1,2,3]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  340.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1<i>H</i>-[1,2,3]triazole
<u>176</u>	6-Chloro-2-[5-(2-chloro-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  373.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-chloro-1<i>H</i>-imidazole



- 260b -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>177</u>	6-Chloro-2-[5-(3-methyl-[1,2,4]triazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  354.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3-methyl-4<i>H</i>-[1,2,4]triazole



- 260c -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
178	6-Chloro-2-[5-(5-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one; compound with 6-chloro-2-[5-(3-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one   354.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and 5-methyl-1H-[1,2,4]triazole



- 260d -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>179</u>	6-Chloro-2-(5-[1,2,4]triazol-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  340.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 4<i>H</i>-[1,2,4]triazole
<u>180</u>	6-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  340.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1<i>H</i>-[1,2,4]triazole



- 260e -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>181</u>	6-Chloro-2-[5-(2-methyl-benzoimidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  403.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-methyl-1<i>H</i>-benzoimidazole
<u>182</u>	6-Chloro-2-(5-indazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  389.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1<i>H</i>-indazole



- 260f -



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>183</u>	6-Chloro-2-(5-indazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  389.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2<i>H</i>-indazole
<u>184</u>	6-Chloro-2-[5-(6-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 6-fluoro-1<i>H</i>-indole

- 260g -



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>185</u>	6-Chloro-2-[5-(7-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 7-fluoro-1<i>H</i>-indole
<u>186</u>	6-Chloro-2-[5-(4-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  406.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 4-fluoro-1<i>H</i>-indole

- 260h -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>187</u>	6-Chloro-2-[5-(4-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 4-methyl-1<i>H</i>-pyrazole
<u>188</u>	6-Chloro-2-[5-(2-cyclopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  379.3	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-cyclopropyl-1<i>H</i>-imidazole


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>189</u>	6-Chloro-2-[5-(2-trifluoromethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  407.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-trifluoromethyl-1<i>H</i>-imidazole
<u>190</u>	6-Chloro-2-[5-(3-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3-methyl-1<i>H</i>-pyrazole

- 260j -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>191</u>	6-Chloro-2-[5-(2-ethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  367.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-ethyl-1<i>H</i>-imidazole
<u>192</u>	2-(5-Aminomethyl-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  288.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1.6 M ammonia in dioxane

Example 193


- 260k -

6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one

To a solution of 6-chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 3) (28.8 mg, 0.1 mmol) in THF (5 mL) was added NaH (8 mg, 0.2 mmol, 5 60% dispersion in mineral oil) at room temperature. After hydrogen evolution ceased, a solution of methyl iodide (0.2 mmol) in THF (5 mL) was added dropwise, and the resulting mixture was stirred at room temperature for 1 hour. The mixture was then treated with satd. aqueous NH₄Cl solution and extracted three times with EtOAc. The combined organic organic layer was washed with water and brine, dried over anhy. Na₂SO₄ and 10 concentrated in vacuo. The crude product was purified by prep-HPLC to give the title compound (4.5 mg, 14.8%) as a white solid. MS: (M+H)⁺ 303.1

Example 194**6-Chloro-2-(5-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one**

- 261 -





To solution of propan-2-ol (3.6 mg, 0.06 mmol) in dry DMF (1 mL) was added NaH (1.5 mg, 0.06 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 5 min before 6-chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2*H*-isoquinolin-1-one (15 mg, 0.05 mmol, example 140 [A]) was added. The resulting reaction mixture was then stirred at room temperature for 1 hour before it was poured into ice-water and extracted with EtOAc (2x 5 mL). The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (8 mg, 48%) as a white solid. MS: 331.1 (M+H⁺).

The following compounds listed in Table 3 were prepared in analogy to the procedure described for the preparation of example 194 using appropriate starting materials.



Table 3

Ex	Compound Name/ Structure	Starting Materials
	MS (M+H) ⁺	



- 262 -

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>195</u>	6-Chloro-2-[5-(2,2,2-trifluoro-1-methyl-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  385.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1,1,1-trifluoro-propan-2-ol
<u>196</u>	6-Chloro-2-{5-[2-(1-methyl-pyrrolidin-2-yl)-ethoxymethyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  400.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-(1-methyl-pyrrolidin-2-yl)-ethanol



- 263 -

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>197</u>	6-Chloro-2-(5-cyclopentyloxymethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  357.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and cyclopentanol
<u>198</u>	6-Chloro-2-(5-cyclopropylmethoxymethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  343.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and cyclopropyl-methanol


- 263a -


Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>199</u>	6-Chloro-2-[5-(2-fluoro-phenoxy)methyl]-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  383.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-fluoro-phenol
<u>200</u>	6-Chloro-2-[5-(1-methyl-cyclopropylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  357.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and (1-methyl-cyclopropyl)-methanol

- 263b -

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>201</u>	6-Chloro-2-[5-(tetrahydro-furan-2-ylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  373.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and (tetrahydro-furan-2-yl)-methanol
<u>202</u>	6-Chloro-2-[5-(2,2,2-trifluoro-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  371.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,2,2-trifluoro-ethanol

- 263c -

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>203</u>	6-Chloro-2-(5-cyclobutoxymethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  343.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and cyclobutanol



Example 204**6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one**



- 5 6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140 [A]) (75 mg, 0.25 mmol), Pd(PPh₃)₂Cl₂ (18 mg, 0.025 mmol), Na₂CO₃ (80 mg, 0.75 mmol) and 3,5-dimethylisoxazole-4-boronic acid (45 mg, 0.32 mmol) were dissolved in dioxane (4.0 mL). The resulting reaction mixture was heated at 120 °C for 3 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 125 mL). The combined
- 10 organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (10 mg, 10.9%) as a light yellowish solid. MS: 368.2 (M+H⁺).



- The following compounds listed in Table 4 were prepared in analogy to the procedure
- 15 described for the preparation of example 204 using appropriate starting materials.



Table 4



Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺



Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>205</u>	6-Chloro-2-[5-(4-methanesulfonyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  427.2	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 4-(methanesulfonyl)phenylboronic acid
<u>206</u>	6-Chloro-2-[5-(6-methyl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  364.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-methyl-5-pyridinylboronic acid


Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>207</u>	6-Chloro-2-[5-(6-morpholin-4-yl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  435.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A]) and 2-(morpholin-4-yl)pyridine-5-boronic acid pinacol ester
<u>208</u>	6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140[A])) and 1-methyl-1H-pyrazole-5-boronic acid pinacol ester

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>209</u>	6-Chloro-2-[5-(1-methyl-1<i>H</i>-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 1-methylpyrazole-4-boronic acid pinacol ester
<u>210</u>	6-Chloro-2-[5-(2,3-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  385.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,3-difluorophenylboronic acid

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>211</u>	6-Chloro-2-[5-(3,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  385.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3,5-difluorophenylboronic acid
<u>212</u>	6-Chloro-2-[5-(2,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  385.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,5-difluorophenylboronic acid


Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>213</u>	6-Chloro-2-[5-(2-trifluoromethyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  417.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2-trifluoromethylphenylboronic acid
<u>214</u>	6-Chloro-2-[5-(2,6-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  417.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,6-dichlorophenylboronic acid

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
215	6-Chloro-2-[5-(2-chloro-6-fluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  401.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,6-dichlorophenylboronic acid
216	6-Chloro-2-[5-(3,4-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  417.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 3,4-dichlorophenylboronic acid

Ex	Compound Name/ Structure MS (M+H) ⁺	Starting Materials
<u>217</u>	6-Chloro-2-[5-(2,5-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  417.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 140[A]) and 2,5-dichlorophenylboronic acid

Example 218

Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-amide






5



CuI (1.9 mg, 0.01 mmol), L-proline (2.3 mg, 0.02 mmol) and Cs₂CO₃ (65 mg, 0.2 mmol) were added to a dioxane (3 mL) solution of 2-(5-bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (example 7) (33.7 mg, 0.1 mmol) and ethanesulfonic acid amide (10.9 mg, 0.1 mmol). The resulting reaction mixture was subject to microwave
 10 reaction at 150 °C for 2.5 hours. The reaction mixture was filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (7.3 mg, 20%) as a white solid. MS: 366.1 (M+H⁺).



The following compounds listed in Table 5 were prepared in analogy to the procedure described for the preparation of example 218 using appropriate starting materials.



5 **Table 5**


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
219	<i>N</i>-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-yl]-benzenesulfonamide  414.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and benzenesulfonamide


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>220</u>	<i>N</i>-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-yl]-methanesulfonamide  352.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and methanesulfonamide
<u>221</u>	Cyclopropanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-yl]-amide  378.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and cyclopropanesulfonic acid amide

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>222</u>	6-Chloro-2-[5-(4-fluoro-benzylamino)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  382.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 4-fluoro-benzylamine
<u>223</u>	6-Chloro-2-[5-(2,2,2-trifluoro-ethylamino)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  356.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 2,2,2-trifluoro-ethylamine

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>224</u>	6-Chloro-2-(5-morpholin-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one  344.2	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and morpholine
<u>225</u>	N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-propionamide  330.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and propionamide

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>226</u>	6-Chloro-2-{5-[(2-methyl-2<i>H</i>-pyrazol-3-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  368.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 1-(1-methyl-1<i>H</i>-pyrazol-5-yl)methanamine
<u>227</u>	2-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-ylamino]-2-methyl-propionic acid  360.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 2-amino-2-methyl-propionic acid


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
228	6-Chloro-2-{5-[(1-methyl-1<i>H</i>-imidazol-4-ylmethyl)-amino]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  368.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 4-(aminomethyl)-1-methyl-1<i>H</i>-imidazole


Example 229**6-Chloro-2-[5-(1*H*-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one**


- 5 2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (example 7) (34 mg, 0.1 mmol), Pd(PPh₃)₄ (5.8 mg, 0.005 mmol), K₂CO₃ (27.6 mg, 0.2 mmol) and 1*H*-pyrazole-4-boronic acid pinacol ester (29.1 mg, 0.15 mmol) were dissolved in dioxane (1 mL). The resulting reaction mixture was subject to microwave reaction at 150 °C for 45 min before it was poured into H₂O (25 mL) and extracted with EtOAc (2 x 25 mL). The
- 10 combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (9 mg, 28%) as a white solid. MS: 325.1 (M+H⁺).

The following compounds listed in Table 6 were prepared in analogy to the procedure described for the preparation of example 229 using appropriate starting materials.



5 **Table 6**

Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺
230	6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  354.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 3,5-dimethylisoxazole-4-boronic acid	



Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺
<u>231</u>	6-Chloro-2-[5-(3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 3-fluorophenylboronic acid	

Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺
<u>232</u>	6-Chloro-2-[5-(3,4-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2 <i>H</i> -isoquinolin-1-one  371.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2 <i>H</i> -isoquinolin-1-one (example 7) and 3,4-difluorophenylboronic acid	



- 280a -

233	6-Chloro-2-[5-(3,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  371.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 3,5-difluorophenylboronic acid	
234	6-Chloro-2-[5-(3-chloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  369.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 3-chlorophenylboronic acid	



- 280b -

<u>235</u>	6-Chloro-2-[5-(2,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  371.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 2,5-difluorophenylboronic acid	
<u>236</u>	6-Chloro-2-[5-(3-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  403.2		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 3-trifluoromethylphenylboronic acid	



- 280c -

<u>237</u>	6-Chloro-2-[5-(3-trifluoromethoxyphenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  419.2		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 3-trifluoromethoxyphenylboronic acid	
<u>238</u>	6-Chloro-2-[5-(2-methyl-2<i>H</i>-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  339.2		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 1-methyl-1<i>H</i>-pyrazole-5-boronic acid pinacol ester	


- 280d -

<u>239</u>	6-Chloro-2-[5-(1-methyl-1<i>H</i>-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  339.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and (1-methyl-1<i>H</i>-pyrazol-4-yl)boronic acid	
<u>240</u>	6-Chloro-2-[5-(4-chloro-3-fluorophenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  387.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 4-chloro-3-fluorophenylboronic acid	

- 280e -


<u>241</u>	6-Chloro-2-[5-(3,4-dichloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  403.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 3,4-dichlorophenylboronic acid	
<u>242</u>	6-Chloro-2-[5-(2-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  403.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 2-trifluoromethylphenylboronic acid	

- 280f -

243	6-Chloro-2-(5-isoxazol-4-yl-pyridin-3-yl)-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  326.1		2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 4-isoxazoleboronic acid pinacol ester	
-------------------	---	--	--	--

Example 244

6-Chloro-2-[5-(1-methyl-1*H*-imidazol-2-yl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




5

To a stirred suspension of zinc dust (130 mg, 1.98 mmol) in THF (2.0 mL) was added 1,2-dibromoethane (0.02 mL). The mixture was heated with a hair dryer until the evolution of ethylene gas was complete. TMS-Cl (0.02 mL) and 2-bromo-1-methyl-1*H*-imidazole (0.07 mL, 0.66 mmol) were then added to the above suspension and it was allowed to stir at RT

10 for 30 minutes before the addition of 2-(5-bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (74 mg, 0.22 mmol, example 7) and Pd(PPh₃)₄ (14 mg, 0.011 mmol). The resulting reaction mixture was then heated to 90 °C for 6 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 80 mL). The combined organic layers


- 280g -

were washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (12.0 mg, 16.2%) as a white foam. MS: 339.1 ($\text{M}+\text{H}^+$).


Example 245**6-Chloro-2-[5-(2,4-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one**

- 5 To a stirred solution of 2-(5-bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) (80 mg, 0.25 mmol) and 1,4-dimethyl-1H-pyrazole (23 mg, 0.25 mmol) in NMP (2.0 mL) were added Pd(OAc)₂ (2.0 mg, 0.008 mmol), 2-dicyclohexylphosphino-2'-(N,N-dimethylamino)biphenyl (DavePhos) (6.3 mg, 0.016 mmol), Bu₄NOAc (153 mg, 0.5 mmol) and isobutyric acid (0.085 mL, 0.08 mmol). The resulting reaction mixture was
- 10 heated to 100°C for 6 hours before it was poured into H₂O (25 mL) and extracted with EtOAc (2 x 90 mL). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (8.7 mg, 9.9%) as a yellow solid. MS: 353.1 (M+H⁺).


15 **Example 246**

5-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-1-methyl-1H-pyrazole-4-carbonitrile

- The title compound was prepared in analogy to the procedure described for the preparation
- 20 of example 245 using 1-methyl-1H-pyrazole-4-carbonitrile as starting material.

Example 247***N*-[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-isobutyramide**


To a solution of 2-(5-amino-pyridin-3-yl)-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one
5 (27.3 mg, 0.1 mmol, example 15) in dry DMF (2 mL) was added at 0 °C NaH (3.0 mg,
0.12 mmol). The reaction mixture was allowed to stir at room temperature for 5 min before
isobutyryl chloride (12.7 mg, 0.12 mmol) was added into the reaction mixture. The
reaction mixture was stirred for additional 1 hour and then poured into ice-water and
extracted with EtOAc (2x 5 mL). The organic layer was dried over anhy. Na₂SO₄, filtered
10 and concentrated in vacuo to give a crude product which was then purified by prep-HPLC
to give title compound (8.1 mg, 24%) as a white solid. MS: 344.1 (M+H⁺).

Example 248**Cyclopropanecarboxylic acid [5-(6-chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-
15 pyridin-3-yl]-amide**

In analogy to the procedure described for the preparation of example 247, the title
compound was synthesized using cyclopropanecarboxylic acid chloride as corresponding
starting material. MS: 342.1 (M+H⁺).

20 Example 249


***N*-[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-4-fluoro-benzamide**



In analogy to the procedure described for the preparation of example 247, the title
5 compound was synthesized using 4-fluoro-benzoyl chloride as corresponding starting material. MS: 396.1 ($M+H^+$).

Example 250

1-[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-3-cyclohexyl-urea




10

2-(5-Amino-pyridin-3-yl)-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (27.3 mg, 0.1 mmol, example 17) was added to a solution of isocyanato-cyclohexane (12.5 mg, 0.1 mmol) in DCM (2 mL). The reaction mixture was stirred at RT for 2 hours. After evaporation of solvent under reduced pressure, a crude product was obtained which was purified by prep-
15 HPLC to give title compound as a white solid (8.0 mg, 20%). MS: 399.2 ($M+H^+$).

Example 251


1-[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-3-(3-trifluoromethyl-phenyl)-urea



In analogy to the procedure described for the preparation of example 250, the title compound was synthesized using 1-isocyanato-3-trifluoromethyl-benzene as corresponding starting material. MS: 461.3 ($M+H^+$).

5 Example 252


6-Chloro-2-(5-hydroxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one




6-Chloro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (1.44 g, 5 mmol, example 10) was dissolved in aq. HBr (48%, 20 mL). The reaction mixture was heated to reflux temperature and stirred overnight. After cooling to room temperature, it was
10 carefully neutralized with satd. aq. NaHCO_3 solution and extracted with EtOAc (2x 25 mL). The organic layer was dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give a crude product (1.02 g, 74%) as brown oil. MS: 275.1 ($M+H^+$).

Example 253

15 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetamide




[A] [5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid tert-butyl ester




NaH (38 mg, 1.5 mmol) was added to a solution of 6-chloro-2-(5-hydroxy-pyridin-3-yl)-
3,4-dihydro-2*H*-isoquinolin-1-one (288 mg, 1.0 mmol, example 252) in dry DMF (5 mL)
5 at 0 °C. After stirred at RT for 10 minutes, bromo-acetic acid tert-butyl ester (234 mg, 1.2 mmol) was added and the reaction mixture was stirred at RT for 1 hour before it was poured into ice-water and extracted with EtOAc (2x 10 mL). The combined organic layers were dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product
10 (320 mg, 82%) as a solid MS: 389.8 (M+H⁺).

[B] [5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid



Trifluoroacetic acid (3 mL) was added to a solution of 5-(6-chloro-1-oxo-3,4-dihydro-1*H*-
isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid tert-butyl ester (388 mg, 1 mmol) in DCM
15 (5 mL) and the reaction mixture was stirred at RT for 2 hours before it was poured into water and extracted with EtOAc (2x 10 mL). The organic layers were dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product (320 mg, 96%) as a solid MS: 333.1 (M+H⁺).


[C] 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yloxy]-acetamide



HATU (38 mg, 0.1 mmol), Et₃N (10.1 mg, 0.1 mmol) and ammonia (1.6 M in 1,4-dioxane, 125 μL, 0.2 mmol) were added to a solution of [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid (33.2 mg, 0.1 mmol) in DMF (2 mL). The reaction mixture was stirred at RT overnight before it was poured into water and extracted with EtOAc (2x 5 mL). The organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to give title compound (5.4 mg, 16%) as a white solid. MS: 332.1 (M+H⁺).

Example 254


10 **2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N-methylacetamide**



In analogy to the procedure described for the preparation of example 253 [C], the title compound was synthesized using methylamine as corresponding starting material. MS: 346.2 (M+H⁺).

Example 255


[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid methyl ester



In analogy to the procedure described for the preparation of example 253 [A], the title compound was synthesized using bromo-acetic acid methyl ester as corresponding starting material. MS: 347.1 ($M+H^+$).

5 **Example 256**


2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N,N-dimethyl-acetamide



10 In analogy to the procedure described for the preparation of example 253 [C], the title compound was synthesized using dimethylamine as corresponding starting material. MS: 360.1 ($M+H^+$).

Example 257

6-Chloro-2-(5-phenylaminomethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one



15 $\text{NaBH}(\text{OAc})_3$ (84 mg, 0.4 mmol) was added to a solution of 5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde (28.6 mg, 0.1 mmol, example 9) and aniline (9.3 mg 0.1 mmol) in MeOH (3mL) and the resulting reaction mixture was


- 288 -

stirred at RT overnight before it was poured into ice-water and extracted with EtOAc (2x 10 mL). The organic layers were dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to give title compound (7.2 mg, 20%) as a white solid. MS: 364.2 (M+H⁺).



5

The following compounds listed in Table 7 were prepared in analogy to the procedure described for the preparation of example 257 using appropriate starting materials.



Table 7

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
258	6-Chloro-2-{5-[(4-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one  382.2	5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde (example 9) and 4-fluorophenylamine

- 289 -


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>259</u>	6-Chloro-2-{5-[(3-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  382.2	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridine-3-carbaldehyde (example 9) and 3-fluorophenylamine
<u>260</u>	6-Chloro-2-{5-[(4-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  398.2	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridine-3-carbaldehyde (example 9) and 4-chlorophenylamine

-289a -


<u>261</u>	6-Chloro-2-{5-[(3-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  398.2	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridine-3-carbaldehyde (example 9) and 3-chloro-phenylamine
<u>262</u>	6-Chloro-2-{5-[(1<i>H</i>-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  354.2	5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridine-3-carbaldehyde (example 9) and 1<i>H</i>-pyrazol-3-ylamine

Example 263

6-Chloro-2-[5-(2-morpholin-4-yl-2-oxo-ethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




[A] (5-Bromo-pyridin-3-yl)-acetic acid methyl ester




To a solution of (5-bromo-pyridin-3-yl)-acetic acid (4.2 g, 20 mmol) in methanol (50 mL)
5 was added thionyl chloride (4.4 mL, 60 mmol) slowly at 0°C. After the addition, the
reaction mixture was warmed up and heated to reflux temperature for 2 hours. After
cooling back to room temperature, the reaction mixture was concentrated in vacuo and the
residue was re-dissolved in EtOAc (250 mL) and washed with aq. NaHCO₃ solution and
brine. The organic layer was dried over Na₂SO₄, filtered and concentrated in vacuo to give
10 a crude product (4.6 g, quant.) as light yellow solid. MS: 230.1 & 232.0 (M+H⁺).

[B] [5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid



(5-Bromo-pyridin-3-yl)-acetic acid methyl ester (230 mg, 1.0 mmol), 6-chloro-3,4-
dihydro-2H-isoquinolin-1-one (intermediate A-2) (182 mg, 1.0 mmol), CuI (40 mg, 0.21
15 mmol), Cs₂CO₃ (650 mg, 2.0 mmol) and (+)-(1S,1S)-1,2-diaminocyclohexane (0.1 mL, 0.8
mmol) were dissolved in dioxane (4.0 ml) and the resulting reaction mixture was heated at
150 °C for 2 hours before it was poured into H₂O (20 mL) and extracted with EtOAc (2 x
100 mL). The combined organic layer was washed with brine, dried over anhy. Na₂SO₄,
filtered and concentrated in vacuo to give a crude product (230 mg, 72.5%) as light brown
20 oil. MS: 317.1 (M+H⁺).


[C] 6-Chloro-2-[5-(2-morpholin-4-yl-2-oxo-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one



[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid (50 mg,
5 0.16 mmol), HATU (122 mg, 0.32 mmol), Et₃N (64 mg, 0.64 mmol) and morpholine (56 mg, 0.64 mmol) were dissolved in DCM (4.0 mL) and the resulting reaction mixture was stirred at room temperature overnight. After evaporating off the DCM, the residue was re-dissolved in EtOAc and was washed with brine. The organic layer was dried over anhy.
Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then
10 purified by prep-HPLC to give the title compound (8.5 mg, 13.9%) as light yellow oil.
MS: 386.1 (M+H⁺).

Example 264

2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-N-(2-hydroxyethyl)-acetamide




15

The title compound was synthesized in analogy to the procedure described for the preparation of example 263 [C] using [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid and 2-amino-ethanol as starting material.


MS: 360.1 (M+H⁺).

Example 265

6-Chloro-2-[5-(1-methylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one




[A] 1-(5-Bromo-pyridin-3-yl)-ethanol




Methylmagnesium bromide (1M in THF, 12 mL, 12 mmol) was added dropwise to a
5 solution of 5-bromo-pyridine-3-carbaldehyde (1.85, 10 mmol) in dry THF (20 mL) at -78 °C. The resulting mixture was slowly warmed up to 0 °C during a 2 hour period and then quenched by aq. satd. NH₄Cl solution. After extraction with DCM, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give title compound as a crude product (1.9 g, 98%).

10 [B] 3-Bromo-5-(1-chloro-ethyl)-pyridine




To a solution of 1-(5-bromo-pyridine-3-yl)-ethanol (201 mg, 1 mmol) in DCM (10 mL) was added dropwise thionyl chloride (200 uL). The resulting reaction mixture was heated to reflux for 3 hours before the solvent was removed in vacuo to give a crude product as a
15 white solid. It was used directly in the next step without further purification.

[C] [1-(5-Bromo-pyridin-3-yl)-ethyl]-methyl-amine



To a solution of 3-bromo-5-(1-chloro-ethyl)-pyridine (221 mg, 1 mmol) in DMF (10 mL) was added methylamine hydrochloride (134 mg, 2 mmol) and K_2CO_3 (572 mg, 4 mmol). The resulting reaction mixture was heated to reflux for 3 hours before it was quenched by
 5 H_2O . After extraction with DCM, the organic layer was dried over anhy. Na_2SO_4 , filtered and concentrated to give title compound as a crude product which was used directly in the next step without further purification.


[D] 6-Chloro-2-[5-(1-methylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one



10 6-Chloro-3,4-dihydro-2H-isoquinolin-1-one (intermediate A-2) (36 mg, 0.2 mmol), [1-(5-bromo-pyridin-3-yl)-ethyl]-methyl-amine (85.6 mg, 0.4 mmol), CuI (3.8 mg, 0.02 mmol), (1S, 2S)-cyclohexane-1,2-diamine (4.5 mg, 0.04 mmol) and Cs_2CO_3 (130 mg, 0.4 mmol) were dissolved in 1,4-dioxane (5 mL). The reaction mixture was subject to microwave reaction at 150 °C for 2.5 hours before it was poured into H_2O (15 mL) and extracted with
 15 $EtOAc$ (2x 10 mL). The organic layer was washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated in vacuo to give a crude product which was purified by prep-HPLC to yield the title compound (15 mg, 26%) as a white solid. MS: 316.2 ($M+H^+$).

Example 266


6-Chloro-2-[5-(1-dimethylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one
 20 one




In analogy to the procedures described for the preparation of example 265, the title compound was synthesized using dimethylamine (step C) and [1-(5-bromo-pyridin-3-yl)-ethyl]-dimethyl-amine (step D) as corresponding starting materials. MS: 330.2 (M+H⁺).

5 **Example 267**

6-Chloro-2-[5-(1-methyl-1H-imidazole-2-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one



[A] (5-Bromo-pyridin-3-yl)-(1-methyl-1H-imidazol-2-yl)-methanol




10

At -78 °C, under N₂ protection, n-BuLi (0.16 mL, 1.6 M, 1.0 mmol) was added to a solution of 1-methyl-1H-imidazole (82 mg, 1.0 mmol) in THF (10 mL) and the reaction mixture was stirred at -78 °C for 0.5 hour; then, 5-bromo-pyridine-3-carbaldehyde (186 mg, 1.0 mmol) was added to the above solution and stirring continued at -78 °C for additional 2 hours. The reaction mixture was poured into water (5.0 mL) and extracted with EtOAc (2 x 50 mL). The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product (210 mg, 78.4%) as white solid. MS: 268.1 & 270.1 (M+H⁺).

15


[B] 6-Chloro-2-[5-(1-methyl-1*H*-imidazole-2-carbonyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




(5-Bromo-pyridin-3-yl)-(1-methyl-1*H*-imidazol-2-yl)-methanol (150 mg, 0.56 mmol), 6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one (90 mg, 0.49 mmol) (intermediate A-2), CuI (30 mg, 0.16 mmol), Cs₂CO₃ (330 mg, 1.0 mmol) and (+)-(1*S*,1*S*)-1,2-diaminocyclohexane (0.08 mL, 0.64 mmol) were dissolved in dioxane (4.0 mL). The resulting mixture was heated at 150 °C for 2 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 75 mL). The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give the title compound (4.5 mg, 2.5%) as light yellow oil. MS: 367.1 (M+H⁺).

Example 268

6-Chloro-2-[5-(4-methyl-4*H*-[1,2,4]triazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




[A] [5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid methyl ester




To a solution of [5-(6-chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid (example 263 [B]) (217 mg, 0.68 mmol) in methanol (25 mL) was added thionyl chloride (0.44 mL, 6.0 mmol) slowly at 0 °C. After the addition, the reaction mixture was warmed up and heated at 80 °C for 2 hours. After cooling back to room temperature, it was concentrated in vacuo and the residue was dissolved in EtOAc (150 mL) and washed with satd. aq. NaHCO₃ solution and then brine. The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product (224 mg, quant.) as light brown oil. MS: 331.1 (M+H⁺).

10 [B] [5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid hydrazide



[5-(6-Chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid methyl ester (100 mg, 0.30 mmol) and hydrazine monohydrate (0.75 mL) were dissolved in ethanol (8.0 mL) and the reaction mixture was heated at 90 °C for 3 h before it was cooled and concentrated in vacuo to give a crude product (100 mg, quant.) as light yellow oil. MS: 331.1 (M+H⁺).


[C] 6-Chloro-2-[5-(4-methyl-4*H*-[1,2,4]triazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one




In a 5 mL sealed tube, [5-(6-chloro-1-oxo-3,4-dihydro-1*H*-isoquinolin-2-yl)-pyridin-3-yl]-acetic acid hydrazide (100 mg, 0.30 mmol) and trimethyl orthoformate (0.07 mL) were dissolved in dioxane (0.8 mL) and heated at 100 °C for 2 hours. After cooling to room temperature, methylamine (1.0 mL, 2.0 M in THF), acetic acid (0.03 mL) and dioxane (2.0 mL) were added. The vial was re-sealed and heated to 130 °C for 2 hours. The reaction mixture was concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (8.3 mg, 7.8%) as light yellowish oil. MS: 354.1 (M+H⁺).

10 **Example 269**

6-Chloro-2-[5-(1-[1,2,3]triazol-2-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one



[A] 2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2*H*-isoquinolin-1-one




15

PBr₃ (541 mg, 2 mmol) was added dropwise into a solution of 6-chloro-2-[5-(1-hydroxyethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one (example 13) (302 mg, 1 mmol) in 5 mL DCM at 0 °C. The reaction was stirred at RT for 2 hours before it was neutralized

with satd. aq. NaHCO₃ solution and extracted with DCM (10 mL, 2x). The organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to afford a crude product as an oil which was used in the next step without further purification (380 mg, 100%). MS: 365.2 (M+H)⁺.

5 [B] 6-Chloro-2-[5-(1-[1,2,3]triazol-2-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one





NaH (30 mg, 0.12 mmol) was added to a solution of 2H-[1,2,3]triazole (8.3 mg, 0.12 mmol) in DMF (5 mL) at 0 °C and the resulting reaction mixture was stirred for 10 min before 2-[5-(1-bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (36.5 mg, 0.1 mmol) was added. After additional stirring at RT for 1 hour, it was poured into ice-water (5 mL) and extracted with EtOAc (2x 10 mL). The organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (7.5 mg, 21%) as a white solid. MS: 354.2 (M+H)⁺.

The following compounds listed in Table 8 were prepared in analogy to the procedure described for the preparation of example 269 using appropriate starting materials.



20 **Table 8**

Ex	Compound Name	Compound Structure	Starting Materials	MS (M+H) ⁺



- 299 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>270</u>	6-Chloro-2-[5-(1-imidazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and 1<i>H</i>-Imidazole
<u>271</u>	6-Chloro-2-[5-(1-pyrazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  353.1	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and 1<i>H</i>-pyrazole



- 300 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>272</u>	6-Chloro-2-{5-[1-(oxazol-2-ylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  369.1	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and oxazol-2-ylamine
<u>273</u>	6-Chloro-2-[5-(1-[1,2,4]triazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  354.2	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and 1<i>H</i>-[1,2,4]triazole


- 301 -


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>274</u>	6-Chloro-2-{5-[1-(2-oxo-pyrrolidin-1-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  370.2	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and pyrrolidin-2-one
<u>275</u>	6-Chloro-2-{5-[1-(2-oxo-oxazolidin-3-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  372.1	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and oxazolidin-2-one

- 301a -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>276</u>	<i>N</i>-{1-[5-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-pyridin-3-yl]-ethyl}-methanesulfonamide  380.2	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and methanesulfonamide
<u>277</u>	6-Chloro-2-{5-[1-(3-fluoro-phenylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  396.2	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and 3-fluoro-phenylamine


- 301b -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials
278	6-Chloro-2-[5-(1-phenylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2<i>H</i>-isoquinolin-1-one  378.2	2-[5-(1-Bromo-ethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 269[A]) and phenylamine


Example 279**6-Chloro-2-(5-methanesulfinyl-pyridin-3-yl)-3,4-dihydro-2*H*-isoquinolin-1-one**

MCPBA (17.3 mg, 0.1 mmol) was added into a solution of 6-chloro-2-(5-methylsulfonyl-pyridin-3-yl)-3,4-dihydro-2*H*-isoquinolin-1-one (example 59) (30.5 mg, 0.1 mmol) in DCM (2 mL). The reaction mixture was stirred at RT for 2 hours before it was neutralized with 5% aq. NaOH and extracted with DCM (5 mL, 2x). The organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (6 mg, 19%) as a white solid. MS: 321.1 (M+H⁺).

Example 280**6-Chloro-2-[4-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2*H*-isoquinolin-1-one**




[A] 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde



CuI (190 mg, 1 mmol), (1S,2S)-cyclohexane-1,2-diamine (228 mg, 2 mmol) and Cs₂CO₃
5 (6.5 g, 20 mmol) were added to a solution of 6-chloro-3,4-dihydro-2H-isoquinolin-1-one
(1.82 g, 10 mmol, intermediate A-2) and 5-bromo-pyridine-4-carbaldehyde (3.72 g, 20
mmol) in dioxane (15 mL). The reaction mixture was heated to 150 °C using microwave
for 2.5 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 20
mL). The organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and
10 concentrated in vacuo to give a crude product (2.0 g, 70%) as brown oil. MS: 287.0
(M+H⁺).

[B] 6-Chloro-2-[4-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one




15 NaBH(OAc)₃ (84 mg, 0.4 mmol) was added into a solution of 5-(6-chloro-1-oxo-3,4-
dihydro-1H-isoquinolin-2-yl)-pyridine-4-carbaldehyde (28.6 mg, 0.1 mmol) and 1-methyl-
piperazine (10 mg 0.1 mmol) in DCM and MeOH (10 mL, 1:1). The reaction mixture was

stirred at RT overnight before it was poured into ice-water (10 mL) and extracted with EtOAc (2x 10 mL). The organic layers were washed with brine, dried over anhy. MgSO_4 , filtered and concentrated in vacuo to give a crude product which was then purified by prep-HPLC to give title compound (7.2 mg, 16%) as a white solid. MS: 371.1 ($\text{M}+\text{H}^+$).

5 **Example 281**

5-Chloro-3,3-dimethyl-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one




In analogy to the procedure described for the preparation of example 204, 5-chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and 1-methylpyrazole-4-boronic acid pinacol ester were used to yield the title compound as a light yellowish solid (yield 15%). MS: 367.2 ($\text{M}+\text{H}^+$).

The following compounds listed in Table 9 were prepared in analogy to the procedures described for the preparation of examples 1, 2 or 3[B] using appropriate starting materials, followed by separation by prep. chiral HPLC when appropriate.


Table 9

Ex	Compound Name	Compound Structure	Starting Materials	Prepared by analogy to example	MS ($\text{M}+\text{H}$) +


- 304 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
282	5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one  339.2	5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1) and 3-bromo-5-difluoromethoxy-pyridine	2



- 305 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
283	5-Chloro-3-ethyl-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  317.3	5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1) and 1-(5-bromo-pyridin-3-yl)-ethanol	2



- 306 -



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
284	5-Chloro-2-(4-chloro-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one  307.2	5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1) and 3-bromo-4-chloro-pyridine	2

- 307 -



Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
285	5-Chloro-2-(4-chloro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  307.2	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-4-chloro-pyridine	2
286	6-Chloro-5'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one  354.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-5-nitro-isoquinoline	2



- 308 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>287</u>	6-Chloro-8'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one  354.2	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-8-nitro-isoquinoline	2
<u>288</u>	8'-Amino-6-chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one  324.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-isoquinolin-8-ylamine	2


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
289	Ethanesulfonic acid (6-chloro-1-oxo-3,4-dihydro-1H-[2,4']bisoquinolinyl-8'-yl)-amide  416.0	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and ethanesulfonic acid (4-bromoisoquinolin-8-yl)-amide	2
290	6'-Chloro-2'-(5-fluoropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one  289.1	6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22) and 3-bromo-5-fluoro-pyridine	2

- 310 -


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>291</u>	6'-Chloro-2'-[5-(difluoromethoxy)pyridin-3-yl]spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one  337.2	6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22) and 3-bromo-5-difluoromethoxy-pyridine	3[B]
<u>292</u>	2-Chloro-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  292.1	2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16) and 3-bromo-5-fluoro-pyridine	2

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>293</u>	2-Chloro-6-(5-difluoromethoxy-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  340.1	2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16) and 3-bromo-5-difluoromethoxy-pyridine	2
<u>294</u>	6-(5-Fluoro-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  288.2	2-Methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-17) and 3-bromo-5-fluoro-pyridine	2


- 312 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
295	6-(5-Difluoromethoxy-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  336.2	2-Methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-17) and 3-bromo-5-difluoromethoxy-pyridine	2



- 313 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
296	6'-Chloro-2'-(pyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one  271.1	6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22) and 3-bromopyridine	2



- 314 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
297	5-Chloro-3-cyclopropyl-2-(5-fluoro-pyridin-3-yl)- 2,3-dihydro-isoindol-1-one  303.2	5-Chloro-3-cyclopropyl-2,3-dihydro-isoindol-1-one (intermediate A-15) and 3-bromopyridine	2


- 315 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>298</u>	2-Chloro-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  274.1	2-Chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16) and 3-bromopyridine	2
<u>299</u>	2-Ethoxy-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  302.2	2-Ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-18) and 3-bromo-5-fluoropyridine	2


- 316 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
300	2-Methoxy-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  N.D.	2-Methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-17) and 3-bromopyridine	2
301	5-Chloro-3-cyclopropyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one  285.1	5-Chloro-3-cyclopropyl-2,3-dihydro-isoindol-1-one (intermediate A-15) and 3-bromopyridine	2



- 317 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
302	5-Chloro-3-cyclopropyl-2-(5-difluoromethoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  351.1	5-Chloro-3-cyclopropyl-2,3-dihydro-isoindol-1-one (intermediate A-15) and 3-bromo-5-difluoromethoxy-pyridine	2


- 318 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
303	6-(5-Difluoromethoxy-pyridin-3-yl)-2-ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  350.1	2-Ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-18) and 3-bromo-5-difluoromethoxy-pyridine	2


- 319 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
304	5-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  331.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-isopropoxy-pyridine	3[B]
305	6'-Chloro-2'-(4-chloropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one  305.0	6'-Chlorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22) and 3-bromo-4-chloro-pyridine	2


- 320 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
306	5-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one  329.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) and 3-bromo-5-cyclopropoxy-pyridine	3[B]


- 321 -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
<u>307</u>	(S or R)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  291.2	(S or R)-6-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-2b) and 3-bromo-pyridine	2


- 321a -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
308	(R or S)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  291.2	(R or S)-6-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-2a) and 3-bromo-pyridine	2


- 321b -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
309	(R or S)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  291.2	(R or S)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1a) and 3-bromo-pyridine	2


- 321c -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
310	(S or R)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  291.2	(S or R)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1b) and 3-bromo-pyridine	2


- 321d -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
311	2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)- 5-chloro-2,3-dihydro-isoindol-1-one  314.2	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12 [A])	3[B]


- 321e -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
312	N-[(R or S)-4-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  398.1	(R or S)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-2a) and (+)-(R)-N-(4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-yl)-propionamide (prep. by separation of intermediate B-12 with prep. chiral HPLC)	3[B]

- 321f -


Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
313	N-[(R or S)-4-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  398.1	(S or R)-5-Chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-2b) and (+)-(R)-N-(4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-yl)-propionamide (prep. by separation of intermediate B-12 with prep. chiral HPLC)	3[B]

- 321g -

Ex	Compound Name/Structure MS (M+H) ⁺	Starting Materials	Prepared by analogy to example
314	N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  370.1	5-Chloro-2,3-dihydro-isoindol-1-one and N-(4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-yl)-propionamide (intermediate B-12)	3[B]


Example 315

Ethanesulfonic acid [4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide




5

[A] 2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one



In a 25 mL sealed tube, 5-chloro-2,3-dihydro-isoindol-1-one (801mg, 4.8 mmol), 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-ylamine (intermediate B-12 [A], 999 mg, 4.4 mmol), CuI (200 mg, 1.1 mmol), Cs₂CO₃ (3.0 g, 9.2 mmol) and (+)-(S,S)-1,2-diaminocyclohexane (0.4 mL, 3.2 mmol) were dissolved in dioxane (16 mL). The resulting reaction mixture was heated at 150 °C for 3 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 125 mL). The organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by silica gel flash chromatography (30-100% EtOAc-hexane gradient) to yield the title compound (1.2 g, 80%) as a light yellow solid. MS: 314.2 (M+H⁺).


[B] Ethanesulfonic acid [4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide



To a stirred solution of 2-(8-amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one (156 mg, 0.5 mmol) and Et₃N (210 uL) in DCM (10 mL) was added ethanesulfonyl chloride (48 uL mg, 0.5 mmol) at 0 °C and stirring continued at 0 °C for 1 h. The resulting mixture was extracted with EtOAc (2 x 50 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and concentrated to afford the title compound. The residue was separated by prep-HPLC to give a racemic mixture of title compound (170 mg, 84 %) as a light yellow solid. MS: 406.1 (M+H⁺).


Example 316

(+)-Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide



Example 317


5 (-)-Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide



The racemic mixture of ethanesulfonic acid [4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide (170 mg, example 315) was subject to chiral HPLC separation to afford (+)-ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide (47 mg, example 316), MS: 406.2 (M+H⁺) and (-)-ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide (39 mg, example 317). MS: 406.2 (M+H)⁺.


15 Example 318

(-)-N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




Example 319

(+)-N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




5 [A] 2-(7-Amino-6,7-dihydro-5H-[2]pyrindin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one



In a 25 mL sealed tube, 5-chloro-2,3-dihydro-isoindol-1-one (801mg, 4.8 mmol), 4-bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine (intermediate B-18 [C], 937 mg, 4.4 mmol), CuI (200 mg, 1.1 mmol), Cs₂CO₃ (3.0 g, 9.2 mmol) and (+)-(S,S)-1,2-diaminocyclohexane
10 (0.4 mL, 3.2 mmol) were dissolved in dioxane (16 mL). The resulting reaction mixture was heated at 150 °C for 3 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (2 x 125 mL). The organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give a crude product which was purified by silica gel flash chromatography (30-100% EtOAc-hexane gradient) to yield the title
15 compound (1.02 g, 80%) as a light yellow solid. MS: 300.2 (M+H⁺).

[B] N-[4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide



To a stirred solution of 2-(7-amino-6,7-dihydro-5*H*-[2]pyrindin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one (126 mg, 0.42 mmol) and Et₃N (1.0 mL) in DCM (10 mL) was added acetyl chloride (0.032 mL, 0.44 mmol) at 0 °C and stirred at 0 °C for 1 hour. The resulting mixture was extracted with EtOAc (2 x 100 mL) and the combined organic layers

5 were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by silica gel flash chromatography eluting with a 0 to 50% EtOAc-heptane gradient to give a racemic mixture of title compound (121 mg, 85 %) as light yellow solid. MS: 342.1 (M+H⁺). This racemic mixture was then separated by chiral HPLC to afford (-)-*N*-[(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-



10 5*H*-[2]pyrindin-7-yl]-acetamide (33 mg, example 318), MS: 342.1 (M+H⁺) and (+)-*N*-[(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5*H*-[2]pyrindin-7-yl]-acetamide (27 mg, example 319). MS: 342.1 (M+H)⁺.

The following compounds listed in Table 10 were prepared in analogy to the procedures



15 described for the preparation of examples 316 and 317 or 318 and 319 using appropriate starting materials.



Table 10

<u>Ex</u>	Compound Name	Compound Structure	Starting Materials	Sign of Optical Rotation	MS (M+S) ⁺
-----------	------------------	-----------------------	-----------------------	--------------------------------	--------------------------



<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>320</u>	N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-acetamide  370.0	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12[A])	(+)
<u>321</u>	N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-acetamide  370.0	Same as Example 320	(-)

- 327 -



<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>322</u>	N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  356.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12[A])	(-)
<u>323</u>	N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  356.1	Same as Example 322	(+)

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>324</u>	N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide  406.0	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12[A])	(-)
<u>325</u>	N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide  406.0	Same as Example 324	(+)



- 329 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>326</u>	N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  392.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12[A])	(-)
<u>327</u>	N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  392.1	Same as Example 326	(+)



- 330 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>328</u>	N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-propionamide  384.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ylamine (intermediate B-12[A])	(+)
<u>329</u>	N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1<i>H</i>-[2,4']biisoquinoliny-8'-yl)-propionamide  384.1	Same as Example 328	(-)



- 331 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>330</u>	N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-yl]-methanesulfonamide  392.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-ylamine (intermediate B-18[C])	(-)
<u>331</u>	N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-yl]-methanesulfonamide  392.1	Same as Example 330	(+)



- 332 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>332</u>	N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-yl]-acetamide  356.1	6-Chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (intermediate A-2) and 4-bromo-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-ylamine (intermediate B-18[C])	(+)
<u>333</u>	N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1<i>H</i>-isoquinolin-2-yl)-6,7-dihydro-5<i>H</i>-[2]pyrindin-7-yl]-acetamide  356.1	Same as Example 332	(-)


- 333 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>334</u>	N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  378.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine (intermediate B-18[C])	(+)
<u>335</u>	N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  378.1	Same as Example 334	(-)



- 334 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>336</u>	Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide  392.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine (intermediate B-18[C])	(+)
<u>337</u>	Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide  392.1	Same as Example 336	(-)


- 335 -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>338</u>	N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  356.2	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine (intermediate B-18[C])	(-)

- 335a -


<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>339</u>	N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  356.2	Same as Example 338	(+)
<u>340</u>	5-Chloro-2-((S)-8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one  315.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydro-isoquinolin-8-ol (intermediate B-11)	(-)

- 335b -

<u>Ex</u>	Compound Name/Structure MS (M+S)+	Starting Materials	Sign of Optical Rotation
<u>341</u>	5-Chloro-2-((R)-8-hydroxy-5,6,7,8-tetrahydro- isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one  315.1	5-Chloro-2,3-dihydro-isoindol-1-one and 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-ol (intermediate B-11)	(+)

Example 342


(+)-N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




5

Example 343

(-)-N-[(R or S)-4-((S or R)-5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide

**Example 344**


(+)-N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




5

Example 345

(-)-N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




10 **[A] 2-(7-Amino-6,7-dihydro-5H[2]pyridin-4-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one**



A mixture of 5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1, 390 mg, 2 mmol), 4-bromo-6,7-dihydro-5H-[2]pyrindin-7-ylamine (intermediate B-18[C], 318 mg, 1.5 mmol), CuI (38 mg, 0.2 mmol), (1S, 2S)-cyclohexane-1,2-diamine (45 mg, 0.4 mmol) and CsCO₃ (750 mg, 4 mmol) were dissolved in dioxane (10 mL). The resulting reaction mixture was heated at 140 °C for 12 hours before it was cooled to room temperature and filtered through a pad of silica gel (0.5 - 1 cm) and washed with diethyl ether. The combined filtrate was concentrated *in vacuo* to give a crude product which was used in the next step reaction without purification. MS: 328.1 (M+H⁺).

10 [B] N-[4-(5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide




To a solution of 2-(7-amino-6,7-dihydro-5H[2]pyridin-4-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (490 mg, 1.5 mmol) in DCM (15 mL) was added Et₃N (3 mmol) and acetyl chloride (156 mg, 2 mmol) at 0 °C. The resulting reaction mixture was stirred at room temperature for 1 hour before it was poured into 1 N HCl in dioxane (15 mL) and washed with EtOAc (15 mL x2). After the organic layer was decanted, the aqueous layer was adjusted to pH > 8 with satd. aq. sodium bicarbonate solution. It was extracted with EtOAc (20 mL x2) and the combined organic layers was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give the desired product (228 mg, 62%) as mixture of four diastereomers, which was subject to SFC separation (IC 250 mm x 50 mm, 5 um, mobile phase A: supercritical CO₂, B: IPA (0.05% NH₃H₂O), A: B = 60: 40 at 140 mL/min) to give (+)-N-[(S or R)-4-((R or S)-5-chloro-3-ethyl-1-oxo-1,3-


- dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide (31 mg, example 342), MS: 370.1 ($M+H^+$), (-)-N-[(R or S)-4-((S or R)-5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide (35 mg, example 343), MS: 370.1 ($M+H^+$), (+)-N-[(R or S)-4-((R or S)-5-chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide (39 mg, example 344), MS: 370.1 ($M+H^+$), and (-)-N-[(S or R)-4-((S or R)-5-chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide (34 mg, example 345), MS: 370.1 ($M+H^+$).

Example 346

- 10 **N-[4-(5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide**



[A] 2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one




15

- A mixture of 5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-1, 390 mg, 2 mmol), 4-bromo-5,6,7,8-tetrahydroisoquinolin-8-amine (intermediate B-12[A], 339 mg, 1.5 mmol), CuI (38 mg, 0.2 mmol), (1S, 2S)-cyclohexane-1,2-diamine (45 mg, 0.4 mmol) and CsCO₃ (750 mg, 4 mmol) were dissolved in dioxane (10 mL) and heated at 140 °C for 12 hours. After cooling to room temperature, the reaction mixture was filtered through a pad of silica gel (0.5 - 1 cm) and rinsed with diethyl ether. The combined filtrate was
- 20

concentrated *in vacuo* to afford a crude product which was used in the next step reaction without further purification. MS: 342.1 ($M+H^+$)



[B] N-[4-(5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide





5

- To a solution of 2-(8-amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one (511.5 mg, 1.5 mmol) in DCM (15 mL) was added Et_3N (3 mmol) and methanesulfonyl chloride (228 mg, 2 mmol) at 0 °C. The resulting reaction mixture was stirred at room temperature for 1 hour before it was poured into 1 N HCl in dioxane (15 mL) and washed with EtOAc (15mL x 2). After the organic layer was decanted, the aqueous layer was basified to pH > 8 with satd. aq. $NaHCO_3$ solution. After extracted with EtOAc (20 mL x2), the combined organic layers were washed with brine, dried over anhy. Na_2SO_4 , filtered and concentrated *in vacuo* to give the title compound (228 mg, 62%) as a mixture of four diastereomers. MS: 420.1 ($M+H^+$).
- 15 This mixture of diastereomers were then subject to SFC separation (IC 250 mm x 50 mm, 5 μm , mobile phase A: supercritical CO_2 , B: IPA (0.05% NH_3H_2O), A: B = 60: 40 at 140 mL/min) to give four individual diastereomers (examples 347, 348, 349 and 350). MS: 420.1 ($M+H^+$).

<u>Ex</u>	Compound	Compound	Sign of Optical	MS
	Name	Structure	Rotation	($M+S$) ⁺



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Sign of Optical Rotation
<u>347</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  420.1	(+)
<u>348</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  420.1	(-)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Sign of Optical Rotation
<u>349</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  420.1	(+)
<u>350</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide  420.1	(-)



The following compounds listed in Table 11 were prepared in analogy to the procedures described for the preparation of example 342, 343, 344, and 345 or examples 347, 348, 349, and 350 using appropriate starting materials.



5 **Table 11**



<u>Ex</u>	Compound Name/Structure MS (M+S)⁺	Optical Rotation Sign
<u>351</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide 398.1	ND


<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>352</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  398.1	ND
<u>353</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  398.1	ND



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>354</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  398.1	ND
<u>355</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  384.1	(-)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>356</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  384.1	(-)
<u>357</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  384.1	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>358</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  384.1	(+)
<u>359</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  384.3	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>360</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  384.3	(-)
<u>361</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  384.3	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>362</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide  384.3	(-)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>363</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  370.2	(-)
<u>364</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  370.2	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>365</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  370.2	(-)
<u>366</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  370.2	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>367</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  406.1	(+)
<u>368</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  406.1	(-)

<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>369</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  406.1	(+)
<u>370</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide  406.1	(-)

<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>371</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  396.1	(-)
<u>372</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  396.1	(+))

<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>373</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  396.1	(+)
<u>374</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide  396.1	(-)


<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>375</u>	N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  370.1	(-)
<u>376</u>	N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  370.1	(+)



<u>Ex</u>	Compound Name/Structure MS (M+S) ⁺	Optical Rotation Sign
<u>377</u>	N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  370.1	(+)
<u>378</u>	N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide  370.1	(-)



The following compounds listed in Table 12 were prepared in analogy to the procedure described for the preparation of example 140 using appropriate starting materials.



Table 12



5



<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>379</u>	5-Chloro-3,3-dimethyl-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one  353.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermedaite A-12-1) and 1<i>H</i>-pyrazole



<u>380</u>	2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one  368.2	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermedaite A-12-1) and 3-aminopyrazole
<u>381</u>	5-Chloro-3,3-dimethyl-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-2,3-dihydro-isoindol-1-one  368.2	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermedaite A-12-1) and 3-aminopyrazole



<u>382</u>	2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one  354.0	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one (example 140 [A]) and 3-aminopyrazole
<u>383</u>	2-Chloro-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  369.1	2-Chloro-6-(5-chloromethyl-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermediate A-16-1) and 3-aminopyrazole



<u>384</u>	2-Methoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  365.2	6-(5-Chloromethyl-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermedaite A-17-1) and 3-aminopyrazole
<u>385</u>	2-Ethoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  379.2	6-(5-Chloromethyl-pyridin-3-yl)-2-ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one (intermedaite A-18-1) and 3-aminopyrazole



<u>386</u>	6'-Chloro-2'-{5-[(1H-pyrazol-3-ylamino)methyl]pyridin-3-yl}spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one  366.2	6'-Chloro-2'-(5-(chloromethyl)pyridin-3-yl)spiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22-1) and 3-aminopyrazole
<u>387</u>	Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  394.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and ethanesulfonic acid amide



<u>388</u>	Ethanesulfonic acid [5-(6-fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  378.1	2-(5-Chloromethyl-pyridin-3-yl)-5-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-19-1) and ethanesulfonic acid amide
<u>389</u>	Ethanesulfonic acid [5-(6-cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  385.1	2-(5-Chloromethyl-pyridin-3-yl)-3,3-dimethyl-1-oxo-2,3-dihydro-1<i>H</i>-isoindole-5-carbonitrile (intermediate A- 21-1) and ethanesulfonic acid amide



<u>390</u>	Ethanesulfonic acid [5-((S or R)-5-chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  394.1	(S or R)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-3) and ethanesulfonic acid amide
<u>391</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  380.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and methanesulfonamide



<u>392</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-N-methyl-methanesulfonamide  394.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and N-methyl-methanesulfonic acid amide
<u>393</u>	N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}ethanesulfonamide  376.1	2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-23-1) and ethanesulfonamide



<u>394</u>	N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl)methyl}-N-methylethanesulfonamide  390.1	2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-23-1) and ethanesulfonic acid methylamide
<u>395</u>	Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methylamide  408.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and ethanesulfonic acid methylamide

<u>396</u>	N-{{5-(6'-Chloro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl)methyl}propanamide  356.1	6'-Chloro-2'-(5-(chloromethyl)pyridin-3-yl)spiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-22-1) and propionamide
<u>397</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydroisoindol-2-yl)-pyridin-3-ylmethyl]-propionamide  358.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and propionamide



<u>398</u>	N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide  342.1	2-(5-Chloromethyl-pyridin-3-yl)-5-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-19-1) and propionamide
<u>399</u>	N-[5-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide  358.1	(S or R)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-3) and propionamide

<u>400</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-acetamide  344.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and acetamide
<u>401</u>	N-{[5-(5'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}methanesulfonamide  362.1	2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3'-one (intermediate A-24-1) and methanesulfonamide



<u>402</u>	N-[5-(6-Cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  371.1	2-(5-Chloromethyl-pyridin-3-yl)-3,3-dimethyl-1-oxo-2,3-dihydro-1<i>H</i>-isoindole-5-carbonitrile (intermediate A-21-1) and methanesulfonamide
<u>403</u>	N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  364.4	2-(5-Chloromethyl-pyridin-3-yl)-5-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-19-1) and methanesulfonamide

404	N-{{5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl}methyl}methanesulfonamide  362.0	2'-(5-(Chloromethyl)pyridin-3-yl)-6'-fluorospiro[cyclopropane-1,1'-isoindolin]-3-one(intermediate A-23-1) and methanesulfonamide
405	N-[5-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  380.1	(S or R)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-3) and methanesulfonamide



- 370a -

406	N-[5-(5-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  364.4	2-(5-Chloromethyl-pyridin-3-yl)-6-fluoro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-20-1) and methanesulfonamide
407	N-[5-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  380.1	(R or S)-5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one (intermediate A-13-4) and methanesulfonamide



- 370b -

408	N-[5-(5-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide  380.1	6-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-25-1) and methanesulfonamide
409	5-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one  406.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and isothiazolidine 1,1-dioxide



- 370c -


<u>410</u>	5-Chloro-3,3-dimethyl-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  384.0	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and piperidin-2-one
<u>411</u>	5-Chloro-3,3-dimethyl-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  371.2	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and imidazolidin-2-one

- 370d -

<u>412</u>	5-Chloro-3,3-dimethyl-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  372.2	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and oxazolidin-2-one
<u>413</u>	5-Chloro-3,3-dimethyl-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  370.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and oxazolidin-2-one

- 370e -

<u>414</u>	5-Chloro-3,3-dimethyl-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  385.1	5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and 1-methyl-imidazolidin-2-one
<u>415</u>	5-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one  420.2	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1) and [1,2]thiazinane 1,1-dioxide

416	5-Chloro-2-[5-(3-isopropyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one  385.1	5-Chloro-2-(5-chloromethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12-1)) and 1-isopropyl-imidazolidin-2-one
-----	--	--


The following compounds listed in Table 13 were prepared in analogy to the procedure described for the preparation of example 229 using appropriate starting materials, which have been prepared as indicated or in close analogy the procedure described for the

5 preparation of example 7.



Table 13



<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
-----------	---	-----------------------

- 370g -



<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>417</u>	6-Chloro-2-[5-(1,5-dimethyl-1H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2<i>H</i>-isoquinolin-1-one (example 7) and 1,5-dimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1<i>H</i>-imidazole

- 370h -



<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>418</u>	5-Chloro-3,3-dimethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  353.1	2-(5-Bromo-pyridin-3-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole
<u>419</u>	6-Chloro-2-[5-(3-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  339.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 3-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>420</u>	6-Chloro-2-[5-(4-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one 	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 4-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole
<u>421</u>	6-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  373.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 4-chloro-1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole


- 370j -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>422</u>	6-Chloro-2-[5-(2,5-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 1,3-dimethyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole
<u>423</u>	6-Chloro-2-[5-(1,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  353.1	2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one (example 7) and 1,5-dimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole



- 370k -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>424</u>	6-Chloro-2-[4-chloro-5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one  373.1	2-(5-Bromo-4-chloro-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole
<u>425</u>	2-Chloro-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  354.1	6-(5-Bromo-pyridin-3-yl)-2-chloro-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole


- 3701 -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
426	2-Methoxy-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one  350.1	6-(5-Bromo-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole

- 370m -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>427</u>	5-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one  387.1	2-(5-Bromo-pyridin-3-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one and 4-chloro-1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole
<u>428</u>	(R or S)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  353.2	(R or S)-2-(5-Bromo-pyridin-3-yl)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole


- 370n -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
429	(S or R)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one  353.2	(S or R)-2-(5-Bromo-pyridin-3-yl)-5-chloro-3-ethyl-2,3-dihydro-isoindol-1-one and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1H-pyrazole

Example 430


3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide

5



[A] (5-Bromo-pyridin-3-yl)-methanol


- 370o -



To a suspension of 5-bromo-pyridine-3-carbaldehyde (10.0 g, 53.7 mmol) in MeOH (100 mL) was added sodium borohydride (2.2 g, 59.1 mmol) at 0 °C. The reaction mixture was


stirred at 0 °C for 1 hour before it was quenched with water (5.0 mL). Evaporation of solvents afforded light yellowish oil which was re-dissolved in EtOAc and washed with brine. The organic layer was dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give the title compound (9.6 g, 95%) as colorless oil. MS: 188.0 & 190.0 (M+H⁺).

5 [B] 3-Bromo-5-chloromethyl-pyridine




At 0 °C, to a solution of (5-bromopyridin-3-yl)methanol (3 g, 16.0 mmol) in DCM (15 mL) was added thionylchloride (7.59 g, 63.8 mmol) dropwise and the reaction mixture was stirred at room temperature over night. The mixture was poured onto ice/water (20 mL),
10 and basified with conc. NaOH solution (8 mL). The mixture was extracted with EtOAc (2 x 50 mL) and combined organic layers were dried over anhy. Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 40% EtOAc-heptane gradient to give the title compound (3.08 g, 93 %) as a white solid. MS: 206.0, 207.9 (M+H⁺).

15 [C] (5-Bromo-pyridin-3-yl)-methylamine




3-Bromo-5-chloromethyl-pyridine (10.3 g, 50 mmol) was dissolved in ammonia methanol solution (7 N, 250 ml) and heated at 60 °C overnight. After aq. NaOH (1N) solution was added to adjust the pH to >12, the mixture was extracted with EtOAc. The organic layer
20 was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by recrystallization to give title compound as a solid (2.5 g, 27%). MS: 187.1 (M+H⁺).

[D] 2-(5-Aminomethyl-pyridin-3-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isindol-1-one



- In a 25 mL sealed tube, a mixture of (5-bromo-pyridin-3-yl)-methylamine (281 mg, 1.5 mmol), 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12, 196 mg, 1.0 mmol), CuI (38 mg, 0.2 mmol), Cs₂CO₃ (652 mg, 2.0 mmol) and (+)-(S,S)-1,2-
- 5 diaminocyclohexane (68.4 mL, 0.6 mmol) were dissolved in dioxane (3 mL). The resulting reaction mixture was heated at 150 °C for 7 hours before it was poured into water (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product (320 mg, 100%). Without further purification, it was used directly in the next step. MS:
- 10 302.1 (M+H⁺).


[E] 3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide




- 15 To solution of 2-(5-aminomethyl-pyridin-3-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (30 mg, 0.1 mmol), Et₃N (15.3 mg, 0.15 mmol) and 3-methyl-pyridine-2-carboxylic acid (13.7 mg, 0.1 mmol) in DMF (3 mL) was added HATU (57.0 mg, 0.15 mmol) at room temperature. The reaction mixture was stirred at room temperature for 1 hour. Both water and EtOAc were added and the organic layer was dried over anhy.
- 20 Na₂SO₄, filtered and concentrated *in vacuo* to give a residue which was then purified by Prep HPLC to give the title compound (10 mg, 24%) as a solid. MS: 421.2 (M+H⁺).

The following compounds listed in Table 14 were prepared in analogy to the procedure described for the preparation of example 430 using appropriate starting materials.



Table 14

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>431</u>	3-Chloro-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  441.1	Example 430 (step D) and 3-chloro-pyridine-2-carboxylic acid



- 374 -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>432</u>	1-Methyl-1<i>H</i>-imidazole-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  410.2	Example 430 (step D) and 1-methyl-1<i>H</i>- imidazole-2-carboxylic acid



- 375 -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>433</u>	2-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide  441.2	Example 430 (step D) and 2-chloro-nicotinic acid
<u>434</u>	Pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  407.1	Example 430 (step D) and pyridine-2-carboxylic acid



- 376 -

<u>Ex</u>	Compound Name/Structure MS (M+H) ⁺	Starting Materials
<u>435</u>	3-Methyl-3<i>H</i>-imidazole-4-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide  410.1	Example 430 (step D) and 3-methyl-3<i>H</i>-imidazole-4-carboxylic acid
<u>436</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-6-methyl-nicotinamide  421.3	Example 430 (step D) and 6-methyl-nicotinic acid

- 376a -


<u>437</u>	3-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-isonicotinamide  441.2	Example 430 (step D) and 3-chloro-isonicotinic acid
<u>438</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide  407.2	Example 430 (step D) and nicotinic acid

- 376b -


<u>439</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-2-methyl-nicotinamide  421.3	Example 430 (step D) and 2-methyl-nicotinic acid
<u>440</u>	N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-4-methyl-nicotinamide  421.3	Example 430 (step D) and 4-methyl-nicotinic acid

Example 441

2-[5-(1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 5-Chloro-2-(5-iodo-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one



A mixture of 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12) (390
 5 mg, 2 mmol), 3,5-diiodo-pyridine (662 mg, 2 mmol), CuI (38 mg, 0.2 mmol), (1S, 2S)-
 cyclohexane-1,2-diamine (45 mg, 0.4 mmol) and K₃PO₄ (888 mg, 4 mmol) were dissolved
 in dioxane (10 mL). The reaction mixture was heated at 110 °C for 3 hours before it was
 poured into H₂O (50 mL) and extracted with EtOAc (25 mL x 2). The combined organic
 layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in
 10 vacuo to give a crude product which was then purified by silica gel flash chromatography
 (30-100% EtOAc-hexane gradient) to yield the title compound (495 mg, 62%) as a white
 solid. MS: 399.0 (M+H⁺).

[B] 3-[5-(6-Chloro-1,1-dimethyl-3-oxo-3-dihydro-isoindol-2-yl)-pyridin-3-yloxy]-
 pyrrolidine-1-carboxylic acid *tert*-butyl ester




15

A sealed microwave tube was charged with CuI (19.1 mg, 0.1 mmol), 1,10-phenanthroline
 (36 mg, 0.2 mmol), Cs₂CO₃ (750 mg, 2.0 mmol), 5-chloro-2-(5-iodo-pyridin-3-yl)-3,3-
 dimethyl-2,3-dihydro-isoindol-1-one (399 mg, 1.0 mmol), 3-hydroxy-pyrrolidine-1-
 carboxylic acid *tert*-butyl ester (374 mg, 2 mmol) and toluene (5 mL). The reaction


mixture was heated at 110 °C for 24 hours. The resulting suspension was cooled to room temperature and filtered through a pad of silica gel (0.5 - 1 cm) and rinsed with diethyl ether. The filtrate was concentrated *in vacuo* and the residue was purified by silica gel flash chromatography to give the title compound (310 mg, 68%). MS: 458 (M+H⁺).

5 [C] 5-Chloro-3,3-dimethyl-2-[5-(pyrrolidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one trifluoro acetic acid salt



To a solution of 3-[5-(6-chloro-1,1-dimethyl-3-oxo-3-dihydro-isoindol-2-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (310 mg, 0.67 mmol) in DCM was added TFA (1 mL) and the resulting mixture was stirred at room temperature for 3 hours. After the solvent was removed *in vacuo*, the crude product was obtained as yellowish oil and was used in the next step without further purification.

10 [D] 2-[5-(1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



15


To a solution of 5-chloro-3,3-dimethyl-2-[5-(pyrrolidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one trifluoro acetic acid salt (94.2 mg, 0.2 mmol) was added Et₃N (1 mmol) and acetyl chloride (23.4 mg, 0.3 mmol) at 0 °C. The resulting reaction mixture was stirred at room temperature for 1 hour before it was poured into H₂O (1 mL) and extracted with EtOAc (15 mL x 2). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product which

20

was purified by PrepHPLC to give the title compound (25.5 mg, 35%) as a white solid. MS: 400.2 ($M+H^+$).

Example 442


2-[5-((R)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, (R)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester was used (step B) to yield the title compound as a white solid. MS: 400.2 ($M+H^+$).

10 Example 443


2-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, (S)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester was used (step B) to yield the title compound as a white solid. MS: 400.2 ($M+H^+$).

Example 444


5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, propionyl
5 chloride was used (step D) to yield the title compound as a white solid. MS: 414.2
(M+H⁺).

Example 445

5-Chloro-2-[5-(1-methanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one




10

In analogy to the procedure described for the preparation of example 441, methanesulfonyl
chloride was used (step D) to yield the title compound as a white solid. MS: 436.2
(M+H⁺).

Example 446


15 **5-Chloro-2-[5-(1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one**



In analogy to the procedure described for the preparation of example 441, ethanesulfonyl chloride was used (step D) to yield the title compound as a white solid. MS: 450.2 ($M+H^+$).

Example 447


- 5 **5-Chloro-2-[5-((R)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one**



- 10 In analogy to the procedure described for the preparation of example 441, (R)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester (step B) and ethanesulfonyl chloride (step D) were used to yield the title compound as a white solid. MS: 450.2 ($M+H^+$).

Example 448

- 5-Chloro-2-[5-((S)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one**




15

In analogy to the procedure described for the preparation of example 441, (S)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester (step B) and ethanesulfonyl chloride (step D) were used to yield the title compound as a white solid. MS: 450.2 ($M+H^+$).

Example 449


2-[5-(1-Acetyl-piperidin-4-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, 4-hydroxy-
5 piperidine-1-carboxylic acid *tert*-butyl ester (step B) was used to yield the title compound
as a white solid. MS: 414.3 (M+H⁺)

Example 450

2-[5-(1-Acetyl-azetidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




10

In analogy to the procedure described for the preparation of example 441, 3-hydroxy-
cyclobutanecarboxylic acid *tert*-butyl ester (step B) was used to yield the title compound
as a white solid. MS: 386.2 (M+H⁺).

Example 451


15 **5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one**



In analogy to the procedure described for the preparation of example 441, 3-hydroxy-cyclobutanecarboxylic acid *tert*-butyl ester (step B) and propionyl chloride were used (step D) to yield the title compound as a white solid. MS: 400.1 ($M+H^+$).

5 Example 452


5-Chloro-2-[5-(1-methanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, 3-hydroxy-cyclobutanecarboxylic acid *tert*-butyl ester (step B) and methanesulfonyl chloride were used (step D) to yield the title compound as a white solid. MS: 422.1 ($M+H^+$).

Example 453


5-Chloro-2-[5-(1-ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 441, 3-hydroxy-cyclobutanecarboxylic acid *tert*-butyl ester (step B) and ethanesulfonyl chloride were used (step D) to yield the title compound as a white solid. MS: 436.1 ($M+H^+$).

Example 454


5 **5-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one**



In analogy to the procedure described for the preparation of example 137, 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12), 3-bromo-5-[(S)-3-(*tert*-butyl-
10 dimethyl-silanyloxy)-pyrrolidin-1-yl]-pyridine was used (step C) to yield the title compound as a white solid. MS: 358.1 ($M+H^+$).


Example 455

2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




15

[A] 4-(5-Bromo-pyridin-3-yl)-piperazine-1-carboxylic acid *tert*-butyl ester




A mixture of piperazine-1-carboxylic acid *tert*-butyl ester (1.86 g, 10 mmol), 3,5-dibromopyridine (2.37 g, 10 mmol), palladium(II) acetate (0.11 g, 0.5 mmol), anphos (0.35 g, 0.6 mmol), and sodium *tert*-butoxide (1.68 g, 15 mmol) in dioxane (10 mL) was heated at 130 °C in a microwave reactor for 2 hours. The mixture was then diluted with water, and extracted with EtOAc. The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash chromatography (eluent with ethyl acetate: hexane=1: 5) to afford the title compound as a white solid (1.84 g, 54%). MS: 342.2 (M⁺).

10 [B] 4-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-yl]-piperazine-1-carboxylic acid *tert*-butyl ester




5-Chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12, 690 mg, 3.51 mmol), 4-(5-bromo-pyridin-3-yl)-piperazine-1-carboxylic acid *tert*-butyl ester (1.0 g, 2.92 mmol), CuI (166.8 mg, 0.88 mmol), (1S, 2S)-cyclohexane-1,2-diamine (120 mg, 1.75 mmol) and Cs₂CO₃ (1.90 g, 5.84 mmol) were dissolved in dioxane (10 mL). The reaction mixture was subjected to microwave reaction at 150 °C for 2.5 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (50 mL x 2). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product which was purified by flash chromatography (eluent with ethyl acetate: hexane = 1: 1) to yield the title compound as yellowish oil (340 mg, 25.6%). MS: 457.3 (M+H⁺).

[C] 5-Chloro-3,3-dimethyl-2-(5-piperazin-1-yl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one



A mixture of 4-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-yl]-piperazine-1-carboxylic acid *tert*-butyl ester (0.34 g, 0.74 mmol) and trifluoroacetic acid (2 mL) in DCM (5 mL) was stirred for 1 hour at room temperature. After removal of solvent, the residue was treated with aq. ammonia to adjust the pH to ~ 9, and then extracted with DCM. The organic layer was washed with water and dried over anhy. Na₂SO₄. After removal of solvent, the crude product was obtained as yellowish oil (266 mg, 100%). MS: 357.2 (M+H⁺).


[D] 2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



To a solution of 5-chloro-3,3-dimethyl-2-(5-piperazin-1-yl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one (52 mg, 0.146 mmol) and TEA (29.5 mg, 0.292 mmol) in DCM (5 mL) was added acetyl chloride (17.2 mg, 0.219 mmol) dropwise at 0 °C. After stirring for 1 hour at room temperature, the mixture was treated with water and extracted with DCM. The organic layer was dried over anhy. Na₂SO₄. After removal of solvent, the residue was purified by preparative HPLC to afford the title compound as a white solid (23 mg, 40%). MS: 399.2 (M+H⁺).

Example 456


5-Chloro-3,3-dimethyl-2-[5-(4-propionyl-piperazin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 455, propionyl chloride was used (step D) to yield the title compound as a white solid (yield 41%). MS: 413.2 ($M+H^+$).

5 **Example 457**


5-Chloro-2-[5-(4-methanesulfonylpiperazin-1-yl)pyridin-3-yl]-3,3-dimethyl-2,3-dihydroisoindol-1-one



In analogy to the procedure described for the preparation of example 455, methanesulfonyl chloride was used (step D) to yield the title compound as a white solid (yield 35%). MS: 435.2 ($M+H^+$).

Example 458


5-Chloro-2-[5-(4-ethanesulfonylpiperazin-1-yl)pyridin-3-yl]-3,3-dimethyl-2,3-dihydroisoindol-1-one



In analogy to the procedure described for the preparation of example 455, ethanesulfonyl chloride was used (step D) to yield the title compound as a white solid (yield 40%). MS: 449.2 ($M+H^+$).

Example 459


5 **5-Chloro-2-{5-[4-(3-chloro-pyridine-2-carbonyl)-piperazin-1-yl]-pyridin-3-yl}-3,3-dimethyl-2,3-dihydro-isoindol-1-one**




A mixture of 5-chloro-3,3-dimethyl-2-(5-piperazin-1-yl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one (example 455 [C]) (52 mg, 0.146 mmol), 3-chloro-pyridine-2-carboxylic acid (25.3 mg, 0.161 mmol), HATU (83.3 mg, 0.219 mmol) and Et_3N (29.5 mg, 0.292 mmol) in DCM (5 mL) was stirred for 2 hours at room temperature. The mixture was treated with water, and extracted with DCM. The organic layer was dried over anhydrous Na_2SO_4 and after removal of solvent, the residue was purified by preparative HPLC to afford the title compound (33 mg, 45%) as a white solid. MS: 496.3 ($M+H^+$).

15 **Example 460**

2-[5-(1-Acetyl-pyrrolidin-3-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 1-(2,5-Dihydro-pyrrol-1-yl)-ethanone



To a solution of 2,5-dihydro-1H-pyrrole (1.0 g, 14.5 mmol) and triethylamine (2.93 g, 29 mmol) in DCM (20 mL) was added acetic anhydride (2.2 g, 21.7 mmol) dropwise. After the addition, the mixture was stirred for 30 minutes at room temperature before water was added. The organic layer was washed with satd. aq. sodium bicarbonate solution and brine in sequence and dried over anhy. Na₂SO₄. After removal of solvents, the crude product was obtained as a white solid (2.89 g) and was used in the next step without further purification. MS: 112.2 (M+H⁺).

[B] 1-[3-(5-Bromo-pyridin-3-yl)-pyrrolidin-1-yl]-ethanone




10

A mixture of 1-(2,5-dihydro-pyrrol-1-yl)-ethanone (70 mg, 0.63 mmol), 3-bromo-5-iodopyridine (283 mg, 1 mmol), triethyl amine (130 mg, 1.26 mmol), formic acid (41 mg, 0.88 mmol) and *tetrakis*(triphenylphosphine)palladium(0) (72.8 mg, 0.063 mmol) in DMF (3 mL) was stirred for 24 hours at 90 °C under nitrogen. After cooling to room temperature, the mixture was treated with water and extracted with ethyl acetate. The organic layer was dried over sodium sulfate. After removal of solvents, the residue was purified by flash chromatography to afford the title compound as yellowish oil (25 mg, 15%). MS: 269.1 (M+H⁺).

[C] 2-[5-(1-Acetyl-pyrrolidin-3-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one


20




A mixture of 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12, 160 mg, 0.82 mmol), 1-[3-(5-bromo-pyridin-3-yl)-pyrrolidin-1-yl]-ethanone (160 mg, 0.6 mmol), CuI (34 mg, 0.8 mmol), (1S, 2S)-cyclohexane-1,2-diamine (41 mg, 0.36 mmol) and Cs₂CO₃ (390 mg, 1.2 mmol) were dissolved in dioxane (5 mL). The reaction mixture
 5 was subjected to microwave reaction at 150 °C for 2.5 hours before it was poured into H₂O (50 mL) and extracted with EtOAc (25 mL x 2). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product which was then purified by Prep-HPLC to yield the title compound (23 mg, 10%) as a white solid. MS: 384.2 (M+H⁺).

10 **Example 461**

2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 1-(3,6-Dihydro-2H-pyridin-1-yl)-ethanone



15

In analogy to the procedure described for the preparation of example 460 [A], 1,2,3,6-tetrahydropyridine was used to yield a crude product as a white solid (yield 90%). MS: 126.0 (M+H⁺).


[B] 1-(5-Bromo-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-yl)-ethanone



20

In analogy to the procedure described for the preparation of example 460 [B], 1-(3,6-dihydro-2*H*-pyridin-1-yl)-ethanone was used to yield the tile compound as yellowish oil (yield 15%). MS: 283.2 ($M+H^+$).


[C] 2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




In analogy to the procedure described for the preparation of example 460 [C], 1-(5-bromo-3',4',5',6'-tetrahydro-2*H*-[3,4']bipyridinyl-1'-yl)-ethanone was used to yield the tile compound (28 mg, 10%) as a white solid. MS: 398.2 ($M+H^+$).

10 **Example 462**

2-[6-(1-Acetyl-piperidin-3-yl)-pyrazin-2-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one




[A] 3-(6-Chloro-pyrazin-2-yl)-piperidine-1,3-dicarboxylic acid 1-*tert*-butyl ester 3-ethyl ester




To a solution of piperidine-1,3-dicarboxylic acid 1-*tert*-butyl ester 3-ethyl ester (90 g, 350.1 mmol) in dry THF (180 mL) at 0 °C was slowly added sodium-*bis*-(trimethylsilyl)-amide (1 M in THF, 450 mL, 450 mmol). The reaction mixture was stirred at the same temperature for 0.5 hr before it was slowly added to a solution of 2,6-dichloropyrazine (60 g, 401.6 mmol) in dry THF (180 mL) under argon. The resulting reaction mixture was stirred at 0 °C for 1 hr, and then allowed to warm up to room temperature, and quenched by the addition of satd. aq. solution of NH₄Cl (360 mL). The organic layer was dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to afford a crude title product. It was then used directly in the next step without further purification.

10 [B] 3-(6-Chloro-pyrazin-2-yl)-piperidine-1,3-dicarboxylic acid 1-*tert*-butyl ester




A solution of 3-(6-chloro-pyrazin-2-yl)-piperidine-1,3-dicarboxylic acid 1-*tert*-butyl ester 3-ethyl ester (350.1 mmol) in THF (360 mL) and 1.5 N aqueous NaOH (360 mL) was heated at 80 °C for 5 hr. After it was cooled to room temperature, the aqueous layer was separated and then carefully acidified to pH = 5 by the addition of 1 N aqueous HCl solution. The resulting mixture was extracted with DCM (300 mL × 3) and the combined organic layers were dried over anhy. Na₂SO₄, filtered and concentrated under reduced pressure to give a crude title compound (46 g), which was used directly in the next step without further purification.

20 [C] 3-(6-Chloro-pyrazin-2-yl)-piperidine-1-carboxylic acid *tert*-butyl ester




- A solution of 3-(6-chloro-pyrazin-2-yl)-piperidine-1,3-dicarboxylic acid 1-*tert*-butyl ester (46 g) in *o*-xylene (300 mL) and DMA (30 mL) was heated under reflux for 5 hrs. After cooling to room temperature, it was washed with brine (200 mL $\times$ 2). The organic layer was then dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo*. The residue
- 5 obtained was then purified on a silica gel column eluting with EA:DCM:Hex = 1:1:3 to give the title compound (18 g) as a white solid. MS: 298.7 (M+H⁺).

[D] 3-[6-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyrazin-2-yl]-piperidine-1-carboxylic acid *tert*-butyl ester



- A mixture of 3-(6-chloro-pyrazin-2-yl)-piperidine-1-carboxylic acid *tert*-butyl ester (892.5 mg, 3.0 mmol), 5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (intermediate A-12, 586.5 mg, 3.0 mmol), CuI (171.4 mg, 0.9 mmol), (1*S*, 2*S*)-cyclohexane-1,2-diamine (205.2 mg, 1.8 mmol) and Cs₂CO₃ (1.956 g, 6.0 mmol) were dissolved in dioxane (15 mL). The reaction mixture was subjected to microwave reaction at 150 °C for 2.5 hours
- 15 before it was poured into H₂O (50 mL) and extracted with EtOAc (50 mL $\times$ 2). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product which was then purified by flash chromatography to afford the title compound (450 mg, 33%) as a white solid. MS: 457.2 (M+H⁺).


- 20 [E] 5-Chloro-3,3-dimethyl-2-(6-piperidin-3-yl-pyrazin-2-yl)-2,3-dihydro-isoindol-1-one



A mixture of 3-[6-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyrazin-2-yl]-piperidine-1-carboxylic acid *tert*-butyl ester (0.45 g, 0.99 mmol) and trifluoroacetic acid (2 mL) in DCM (10 mL) was stirred for 1 hour at room temperature. After removal of

solvent, the residue was treated with water, and extracted with DCM. The aqueous solution treated with satd. aq. sodium carbonate solution, and extracted with EtOAc. The combined organic layers were dried over anhy. Na_2SO_4 . After removal of solvent, the crude product was obtained as yellowish oil (352.9 mg, 100%). MS: 357.2 ($\text{M}+\text{H}^+$).


5 [F] 2-[6-(1-Acetyl-piperidin-3-yl)-pyrazin-2-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



To a mixture of 5-chloro-3,3-dimethyl-2-(6-piperidin-3-yl-pyrazin-2-yl)-2,3-dihydro-isoindol-1-one (80 mg, 0.22 mmol) and Et_3N (44.4 mg, 0.44 mmol) in DCM (2 mL) was added acetyl chloride (26.4 mg, 0.34 mmol) at 0 °C. After removal of solvent, the residue was purified by prep-HPLC to yield the title compound (30.6 mg, 35%) as a white solid. MS: 399.2 ($\text{M}+\text{H}^+$).

Example 463


15 **5-Chloro-3,3-dimethyl-2-[6-(1-propionyl-piperidin-3-yl)-pyrazin-2-yl]-2,3-dihydro-isoindol-1-one**



In analogy to the procedure described for the preparation of example 462, propionyl chloride was used (step F) to yield the title compound (36.3 mg, 40%) as a white solid. MS: 413.2 ($\text{M}+\text{H}^+$).

20 **Example 464**


5-Chloro-2-[6-(1-ethanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one



In analogy to the procedure described for the preparation of example 462, ethanesulfonyl chloride was used (step F) to yield the tile compound (29.6 mg, 30%) as a white solid. MS: 449.2 ($M+H^+$).

Example 465

5-Chloro-2-[6-(1-methanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one






10

In analogy to the procedure described for the preparation of example 462, methanesulfonyl chloride was used (step F) to yield the tile compound (19.1 mg, 20% yield) as a white solid. MS: 435.0 ($M+H^+$).


Example 466

15 N-[(S or R)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide



Example 467**N-[(R or S)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide**5 **[A] 4'-Bromo-5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinoline]**


A reaction mixture of 4-bromo-6,7-dihydro-5*H*-isoquinolin-8-one (intermediate B-11 [C], 5.0 g, 22 mmol), 2,2-dimethyl-propane-1,3-diol (2.8 g, 26.5 mmol) and toluene-4-sulfonic acid (85 mg, 0.44 mmol) in toluene (100 mL) was heated at 135 °C for 12 hours. After cooling to room temperature, it was concentrated under reduced pressure to give a residue which was extracted between EtOAc and water. The organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered, and concentrated *in vacuo* to give a crude product (6.1 g, 88.9%) as a light yellow solid. MS: 312.2 & 314.2 (M+H⁺).

15 **[B] 5-Chloro-2-(5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinolin]-4'-yl)-2,3-dihydro-1H-isoindol-1-one**

A mixture of 4'-bromo-5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinoline](5.0 g, 16 mmol), 5-chloro-2,3-dihydro-isoindol-1-one (2.7 g, 16 mmol), (+)-(1*S*,1*S*)-1,2-diaminocyclohexane (0.576 mL, 4.8 mmol), copper (I) iodide (456 mg, 2.4


mmol) and Cs_2CO_3 (10.4 g, 32 mmol) in dioxane (80 mL) was heated at 150 °C for 4 hours. After cooling to room temperature, it was diluted with water (50 mL) and extracted with EtOAc (200 mL x 2). The combined organic layers were dried over anhy. Na_2SO_4 , filtered and concentrated *in vacuo* to afford a crude product (5.5 g, 86%) as light yellow solid. MS: 399.1 ($\text{M}+\text{H}^+$).

[C] 5-Chloro-2-(5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinolin]-4'-yl)-3,3-dimethyl-2,3-dihydro-1H-isoindol-1-one



To a stirred solution of 5-chloro-2-(5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinolin]-4'-yl)-2,3-dihydro-1H-isoindol-1-one (2.38 g, 6 mmol) in THF was added LiHMDS (18 mL 1.0 M in THF, 18 mmol) at 0 °C. It was stirred at 0 °C 30 min before CH_3I (1.5 mL, 24 mmol) was added. The resulting mixture was stirred at 0 °C for another 30 min. before it was diluted with water (10 mL) and extracted with EtOAc (2 x 100 mL). The combined organic layers were dried over anhy. Na_2SO_4 , filtered and concentrated *in vacuo* to afford a crude product which was then purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (664 mg, 26 %) as a yellow foam. MS: 427.1 ($\text{M}+\text{H}^+$).


[D] 4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-isoquinolin-8-one



To a solution of 5-chloro-2-(5,5-dimethyl-6',7'-dihydro-5'H-spiro[1,3-dioxane-2,8'-isoquinolin]-4'-yl)-3,3-dimethyl-2,3-dihydro-1H-isoindol-1-one (0.66 g, 1.55 mmol) in


MeOH (2 mL) was added 4 N HCl in dioxane (1.5 mL, 6 mmol) and the reaction mixture was stirred at room temperature for 2 hours. After evaporation of the solvent, the residue was diluted with DCM (20 mL) and washed with satd. aq. NaHCO₃ (10 mL) solution. The organic layer was dried over anhy. Na₂SO₄, filtered and evaporated to dryness. The residue
 5 obtained was purified by silica gel flash chromatography eluting with a 0 to 50 % hexane – ethyl acetate gradient to give the title compound (421 mg, 80 %) as a light yellow solid. MS: 341.1 (M+H⁺).

[E] 2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one



A mixture of 4-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-isoquinolin-8-one (408 mg, 1.2 mmol), NaBH₃CN (75 mg, 1.2 mmol) and CH₃COONH₄ (1.0 g, 12 mmol) in isopropanol (10 mL) was heated to reflux for 3 hours. After cooling to room temperature, it was concentrated to afford a yellowish oil which was extracted
 15 between water with EtOAc (2 x 100 mL). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated *in vacuo* to afford the crude title compound (327 mg, 80%) as a brown solid. MS: 342.1 & 325.1 (M+H⁺).

[F] N-[4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide




To a stirred solution of 2-(8-amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one (140 mg, 0.42 mmol) and Et₃N (1.0 mL) in DCM (10

mL) was added acetyl chloride (0.032 mL, 0.44 mmol) at 0 °C and stirring was continued at 0 °C for 1 hour. After evaporation of solvent, the residue was extracted between water with EtOAc (2 x 100 mL). The combined organic layers were washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to afford a crude product which was
5 then purified by silica gel flash chromatography eluting with a 0 to 50% EtOAc-heptane gradient to give a racemic mixture of the title compound (117 mg, 73 %) as light yellow solid. MS: 384.1 (M+H⁺). This racemic mixture was then separated by chiral HPLC to afford N-[(S or R)-4-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide (35 mg, example 466), MS: 384.1 (M+H⁺) and N-
10 [(R or S)-4-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide (38 mg, example 467) MS: 384.1 (M+H)⁺.

Example 468

N-[4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide



15

In analogy to the procedure described for the preparation of Examples 466 and 467, propionyl chloride was used (step E) to afford a racemic mixture of the title compound (19.1 mg, 20%) as a white solid. MS: 398.1 (M+H⁺).

Example A

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of tablets of the following composition:

	<u>Per tablet</u>
5 Active ingredient	200 mg
Microcrystalline cellulose	155 mg
Corn starch	25 mg
Talc	25 mg
Hydroxypropylmethylcellulose	<u>20 mg</u>
10	425 mg


Example B

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of capsules of the following composition:

	<u>Per capsule</u>
15 Active ingredient	100.0 mg
Corn starch	20.0 mg
Lactose	95.0 mg
Talc	4.5 mg
20 Magnesium stearate	<u>0.5 mg</u>
	220.0 mg

CLAIMS

1. Compounds of formula (I)



wherein

- 5 R^1 , R^2 , R^3 and R^4 are independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, halocycloalkylalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted arylalkyl and substituted heteroarylalkyl are substituted with R^{12} , R^{13} and R^{14} ;
- 10

or R^2 and R^4 together form a double bond, wherein in case R^2 and R^4 together form a double bond, then R^5 is H;

- or R^1 and R^2 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{22} , R^{23} and R^{24} ;
- 15

or R^3 and R^4 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{29} , R^{30} and R^{31} ;

- or R^1 and R^3 together with the carbon atom to which they are attached form a substituted cycloalkyl or a substituted heterocycloalkyl, wherein substituted cycloalkyl and substituted heterocycloalkyl are substituted with R^{44} , R^{45} and R^{46} ;
- 20

A^1 is CR^8 or N;

A^2 is CR^9 or N;

A^3 is CR^{10} or N;

A^4 is CR^{11} or N;

A^5 is CR^6 or N;

- 5 one of R^5 , R^6 , R^7 and R^8 is selected from halogen, cyano, alkoxy, hydroxyalkoxy, haloalkyl, haloalkoxy and hydroxy and the others are each independently selected from H, halogen, cyano, alkoxy, hydroxyalkoxy, haloalkoxy and hydroxy;

- R^9 is H, halogen, hydroxy, cyano, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxycycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, alkoxycycloalkylalkyl, dialkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl or substituted heteroarylalkyl, wherein substituted arylalkyl, substituted arylhydroxyalkyl, substituted heterocycloalkylalkyl and substituted heteroarylalkyl are substituted with R^{32} , R^{33} and R^{34} ;
- 10
- 15

R^{10} is $-O_m-(CR^{15}R^{16})_p-(CR^{17}R^{18})_q-(CR^{19}R^{20})_r-R^{21}$;

- 20 or R^9 and R^{10} together with the carbon atoms to which they are attached form a substituted cycloalkyl, a substituted heterocycloalkyl, a substituted aryl or a substituted heteroaryl, wherein substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl and substituted heteroaryl are substituted with R^{35} , R^{36} and R^{37} ;

R^{11} is H;

- 25 R^{15} , R^{17} and R^{19} are each independently selected from H, alkyl, cycloalkyl, haloalkyl and halocycloalkyl;

R¹⁶, R¹⁸ and R²⁰ are each independently selected from H, hydroxy, halogen and alkyl;

or R¹⁵ and R¹⁶ together with the carbon atom to which they are attached form a cycloalkyl;

5 or R¹⁷ and R¹⁸ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁹ and R²⁰ together with the carbon atom to which they are attached form a cycloalkyl;

or R¹⁵ and R¹⁷ together form -(CH₂)_v-;

10 or R¹⁵ and R¹⁹ together form -(CH₂)_w-;

or R¹⁷ and R¹⁹ together form -(CH₂)_x-;

15 R^{21} is H, halogen, cyano, $-OR^{25}$, $-SR^{25}$, $-S(O)R^{25}$, $-S(O)_2R^{25}$, $-NR^{25}R^{26}$, $-NR^{26}SO_2R^{25}$, $-NR^{26}SO_2NR^{25}R^{27}$, $-NR^{26}C(O)R^{25}$, $-NR^{26}C(O)NR^{25}R^{27}$, $-C(O)R^{28}$, $-C(O)NR^{25}R^{26}$, cycloalkyl, substituted heterocycloalkyl, substituted heteroaryl or substituted aryl, wherein substituted heterocycloalkyl, substituted heteroaryl, substituted heteroarylalkyl and substituted aryl are substituted with R^{38} , R^{39} and R^{40} ;

R²⁵ is H, alkyl, hydroxyalkyl, carboxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxyalkyl, alkoxyalkylalkyl, halocycloalkylalkyl, cycloalkylalkoxyalkyl, cycloalkoxyalkyl, halocycloalkoxyalkyl, alkylcycloalkoxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxyalkoxyalkyl, haloalkoxyalkoxyalkyl, substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R⁴¹, R⁴² and R⁴³;

R²⁶ and R²⁷ are each independently selected from H, alkyl, cycloalkyl, haloalkyl or halocycloalkyl;

or R¹⁵ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

5 or R¹⁷ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

10 or R¹⁹ and R²⁶ together with the nitrogen atom and carbon atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R⁴⁷, R⁴⁸ and R⁴⁹;

R²⁸ is H, hydroxy, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkoxy, cycloalkylalkoxy, halocycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted
15 heteroaryl, substituted heteroarylalkyl, substituted aryl or substituted arylalkyl, wherein substituted substituted heterocycloalkyl, substituted heterocycloalkylalkyl, substituted heteroaryl, substituted heteroarylalkyl, substituted aryl and substituted arylalkyl are substituted with R⁵⁰, R⁵¹ and R⁵²;

25 R¹², R¹³, R¹⁴, R²², R²³, R²⁴, R²⁹, R³⁰, R³¹, R³², R³³, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹, R⁴⁰, R⁴¹, R⁴², R⁴³, R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷, R⁴⁸, R⁴⁹, R⁵⁰, R⁵¹ and R⁵² are each independently selected from H, halogen, hydroxy, amino, nitro, cyano, oxo, alkyl, alkylcarbonyl, alkylsulfonyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, alkylcarbonylamino, alkylsulfonyl, alkylsulfonylamino, alkoxy, cycloalkylalkoxy, cycloalkylalkoxyalkyl, cycloalkoxy, cycloalkoxyalkyl, halocycloalkoxy,
30

halocycloalkoxyalkyl, alkylcycloalkoxy, alkylcycloalkoxyalkyl, alkoxy, alkoxy carbonyl, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, haloalkoxyalkoxy, haloalkoxyalkoxyalkyl, chloropyridinylcarbonyl and heterocycloalkyl;

5 n is zero or 1;

m zero or 1;

p, q and r are independently selected from zero and 1;

v and x are independently selected from 1, 2, 3 or 4;

w is zero, 1, 2 or 3;

10 with the proviso that no more than two of A^2 , A^3 and A^4 are N;

and wherein said compound is selected from

6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;

5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-nicotinonitrile;

6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-(5-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-fluoro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 2-(5-Bromo-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridine-3-carbaldehyde;
- 5 6-Chloro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 2-(5-Amino-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one
- 20 6-Chloro-2-[5-(cyclopropyl-hydroxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-(4-trifluoromethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methoxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide;
- 6-Chloro-2-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-8'-hydroxy-3,4,5',6',7',8'-hexahydro-[2,4']biisoquinoliny-1-one;
- N-(6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;
- 15 6-Chloro-2-{5-[hydroxy-(1-methyl-1H-imidazol-2-yl)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3,4-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3,5-difluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[(4-ethyl-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(hydroxy-phenyl-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(1-hydroxy-1-phenyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-{5-[1-(3,4-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(3,5-difluoro-phenyl)-1-hydroxy-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(6-methyl-pyrazin-2-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(morpholine-4-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N,N-dimethyl-nicotinamide;
- 6-Chloro-2-[5-(pyrrolidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-methyl-nicotinamide ;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-cyclopropyl-nicotinamide;
- 15 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-(4-fluoro-phenyl)-nicotinamide;
- 5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-N-phenyl-nicotinamide;
- 6-Chloro-2-[5-(4,4-difluoro-piperidine-1-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-yl)methyl]-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-pyridazin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-pyridin-3-yl-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-hydroxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 2-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methylsulfanyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-(4-dimethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{4-[(4-fluoro-phenyl)-hydroxy-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1-methyl-1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2-methyl-2H-[1,2,4]triazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-fluoro-4-(1-hydroxy-1-methyl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(5-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(1H-pyrrolo[3,2-c]pyridin-7-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one;
- 3-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-isonicotinonitrile;
- 6-Chloro-2-(5-fluoro-4-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-fluoro-4-(1-methoxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 1-one;
- 6-Chloro-2-(4-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[4-(cyclopropyl-methoxy-methyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 1-one;
- 6-Chloro-2-[5-(3,5-dimethyl-3H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 (S)-1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-3-methyl-2-pyrimidin-5-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 15 (R)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- (S)-6-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 8-Chloro-3-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Methoxy-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5,6-Dichloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 20 2-Chloro-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 2-Methoxy-6-pyridin-3-yl-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-5-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-7-fluoro-2-(5-methoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-7-fluoro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-4,4-dimethyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4,4-dimethyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-4-methyl-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-fluoro-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxy-pyridin-3-yl)-4-methyl-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-Chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 15 one;
- 5-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 20 5-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- Ethanesulfonic acid [5-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

- 5-Chloro-2-(1-methyl-1H-pyrazolo[3,4-c]pyridin-4-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5 6-Chloro-3-methyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-methyl-2-(4-methyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3-methyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3-methyl-2,3-dihydro-isoindol-1-one;
- 10 3-Benzyl-5-chloro-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3-ethyl-2-(5-methoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;
- 5-Chloro-3,3-dimethyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;
- 15 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-methoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 5-Chloro-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- 20 one;
- 5-Chloro-2-(5-hydroxymethyl-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;
- one;

- 5-Chloro-3,3-dimethyl-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 5-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 5-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;
- 2-Hydroxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;
- 6-Chloro-2-(5-imidazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-isopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(2-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-ethyl-4-methyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-hydroxy-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 Propane-2-sulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-amide
- 6-Chloro-2-[5-(3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-((R)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-((S)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-((R)-2-hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3,5-dimethyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-acetamide
- 15 6-Chloro-2-(5-morpholin-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(1,1-dioxo-1λ6-thiomorpholin-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-propyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylmethyl]-1H-imidazole-2-carboxylic acid ethyl ester;
- 6-Chloro-2-[5-(2-hydroxymethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(oxetan-3-ylaminomethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[4-(2-hydroxy-ethyl)-piperazin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-isopropyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(4,4-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3,3-difluoro-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.3]hept-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(3,3-difluoro-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-[1,2,3]triazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-(5-[1,2,3]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-chloro-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(3-methyl-[1,2,4]triazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(5-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-chloro-2-[5-(3-methyl-[1,2,4]triazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-(5-[1,2,4]triazol-4-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-[1,2,4]triazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(2-methyl-benzoimidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-indazol-1-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-indazol-2-ylmethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(6-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(7-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-fluoro-indol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(4-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-cyclopropyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(2-trifluoromethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(3-methyl-pyrazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-ethyl-imidazol-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 15 2-(5-Aminomethyl-pyridin-3-yl)-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isopropoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-1-methyl-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[2-(1-methyl-pyrrolidin-2-yl)-ethoxymethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-cyclopentyloxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-(5-cyclopropylmethoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-fluoro-phenoxyethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1-methyl-cyclopropylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-[5-(tetrahydro-furan-2-ylmethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,2,2-trifluoro-ethoxymethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-cyclobutoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-methanesulfonyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(6-methyl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(6-morpholin-4-yl-pyridin-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,3-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

25 6-Chloro-2-[5-(3,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(2,5-difluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(2,6-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-chloro-6-fluoro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(3,4-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,5-dichloro-benzyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- Ethanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;
- 15 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-benzenesulfonamide;
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-methanesulfonamide;
- 20 Cyclopropanesulfonic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide;
- 6-Chloro-2-[5-(4-fluoro-benzylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,2,2-trifluoro-ethylamino)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-(5-morpholin-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-
propionamide;

6-Chloro-2-{5-[(2-methyl-2H-pyrazol-3-yl)methyl]-amino}-pyridin-3-yl}-3,4-
dihydro-2H-isoquinolin-1-one;

5 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-ylamino]-2-
methyl-propionic acid;

6-Chloro-2-{5-[(1-methyl-1H-imidazol-4-yl)methyl]-amino}-pyridin-3-yl}-3,4-
dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-(3,5-dimethyl-isoxazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-
isoquinolin-1-one;

6-Chloro-2-[5-(3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(3,4-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-
one;

15 6-Chloro-2-[5-(3,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-
one;

6-Chloro-2-[5-(3-chloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2,5-difluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-
one;

20 6-Chloro-2-[5-(3-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-
1-one;

6-Chloro-2-[5-(3-trifluoromethoxy-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-
isoquinolin-1-one;

25 6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-
isoquinolin-1-one;

- 6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(4-chloro-3-fluoro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(3,4-dichloro-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2-trifluoromethyl-phenyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-(5-isoxazol-4-yl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(1-methyl-1H-imidazol-2-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(2,4-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-1-methyl-1H-pyrazole-4-carbonitrile
- 15 N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-isobutyramide
- Cyclopropanecarboxylic acid [5-(6-chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-amide
- N-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-4-fluorobenzamide
- 20 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-cyclohexylurea
- 1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-3-(3-trifluoromethyl-phenyl)-urea
- 25 6-Chloro-2-(5-hydroxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetamide;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N-methyl-acetamide;
- 5 [5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-acetic acid methyl ester;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yloxy]-N,N-dimethyl-acetamide;
- 6-Chloro-2-(5-phenylaminomethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-{5-[(4-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(3-fluoro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(4-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-{5-[(3-chloro-phenylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-[5-(2-morpholin-4-yl-2-oxo-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 2-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-N-(2-hydroxy-ethyl)-acetamide;
- 6-Chloro-2-[5-(1-methylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

- 6-Chloro-2-[5-(1-dimethylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-methyl-1H-imidazole-2-carbonyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 5 6-Chloro-2-[5-(4-methyl-4H-[1,2,4]triazol-3-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-[1,2,3]triazol-2-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 10 6-Chloro-2-[5-(1-imidazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-[5-(1-pyrazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(oxazol-2-ylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 15 6-Chloro-2-[5-(1-[1,2,4]triazol-1-yl-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;
- 6-Chloro-2-{5-[1-(2-oxo-pyrrolidin-1-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 20 6-Chloro-2-{5-[1-(2-oxo-oxazolidin-3-yl)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- N-{1-[5-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-pyridin-3-yl]-ethyl}-methanesulfonamide;
- 6-Chloro-2-{5-[1-(3-fluoro-phenylamino)-ethyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;
- 25 6-Chloro-2-[5-(1-phenylamino-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-methanesulfinyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[4-(4-methyl-piperazin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

and pharmaceutically acceptable salts thereof.

5 2. A compound according to claim 1, selected from

5-Chloro-3,3-dimethyl-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3-ethyl-2-[5-(1-hydroxy-ethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

10 5-Chloro-2-(4-chloro-pyridin-3-yl)-3-ethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(4-chloro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

6-Chloro-5'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

6-Chloro-8'-nitro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

8'-Amino-6-chloro-3,4-dihydro-[2,4']biisoquinoliny-1-one;

15 Ethanesulfonic acid (6-chloro-1-oxo-3,4-dihydro-1H-[2,4']biisoquinoliny-8'-yl)-amide;

6'-Chloro-2'-(5-fluoropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

6'-Chloro-2'-[5-(difluoromethoxy)pyridin-3-yl]spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

20 2-Chloro-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

2-Chloro-6-(5-difluoromethoxy-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

6-(5-Fluoro-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

6-(5-Difluoromethoxy-pyridin-3-yl)-2-methoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

5 6'-Chloro-2'-(pyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

5-Chloro-3-cyclopropyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

2-Chloro-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

2-Ethoxy-6-(5-fluoro-pyridin-3-yl)-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

10 2-Methoxy-7,7-dimethyl-6-pyridin-3-yl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

5-Chloro-3-cyclopropyl-2-pyridin-3-yl-2,3-dihydro-isoindol-1-one;

5-Chloro-3-cyclopropyl-2-(5-difluoromethoxy-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

15 6-(5-Difluoromethoxy-pyridin-3-yl)-2-ethoxy-7,7-dimethyl-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

5-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

6'-Chloro-2'-(4-chloropyridin-3-yl)spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

5-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

(S or R)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

20 (R or S)-6-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

(R or S)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

(S or R)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

2-(8-Amino-5,6,7,8-tetrahydro-isoquinolin-4-yl)-5-chloro-2,3-dihydro-isoindol-1-one;

N-[(R or S)-4-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

5 N-[(R or S)-4-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

10 Ethanesulfonic acid [4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-amide;

15 N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

20 N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinolinyl-8'-yl)-acetamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinolinyl-8'-yl)-acetamide;

N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

25 N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide;

N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-methanesulfonamide;

5 N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

10 N-((R or S)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;

N-((S or R)-6-Chloro-1-oxo-3,4,5',6',7',8'-hexahydro-1H-[2,4']biisoquinoliny-8'-yl)-propionamide;

N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

15 N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

20 N-[(S or R)-4-(6-Chloro-1-oxo-3,4-dihydro-1H-isoquinolin-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

25 Ethanesulfonic acid [(R or S)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide;

Ethanesulfonic acid [(S or R)-4-(5-chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-amide;

N-[(S or R)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5 N-[(R or S)-4-(5-Chloro-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5-Chloro-2-((S or R)-8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;

10 5-Chloro-2-((R or S)-8-hydroxy-5,6,7,8-tetrahydro-isoquinolin-4-yl)-2,3-dihydro-isoindol-1-one;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

15 N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-acetamide;

20 N-[4-(5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

25 N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-methanesulfonamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

5 N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

10 N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

15 N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

20 N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

25 N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

5 N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

10 N-[(S or R)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

15 N-[(S or R)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

N-[(S or R)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

20 N-[(R or S)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-cyclopropyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

25 N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5 N-[(S or R)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5-Chloro-3,3-dimethyl-2-(5-pyrazol-1-ylmethyl-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

10 2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-2,3-dihydro-isoindol-1-one;

2-[5-(3-Amino-pyrazol-1-ylmethyl)-pyridin-3-yl]-6-chloro-3,4-dihydro-2H-isoquinolin-1-one;

15 2-Chloro-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

2-Methoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

20 2-Ethoxy-7,7-dimethyl-6-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

6'-Chloro-2'-{5-[(1H-pyrazol-3-ylamino)methyl]pyridin-3-yl}spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

25 Ethanesulfonic acid [5-(6-fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(6-cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-((S or R)-5-chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

5 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-N-methyl-methanesulfonamide;

10 N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}ethanesulfonamide;

N-{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}-N-methylethanesulfonamide;

Ethanesulfonic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methyl-amide;

15 N-{[5-(6'-Chloro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}propanamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

20 N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

N-[5-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-propionamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-acetamide;

25 N-{[5-(5'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}methanesulfonamide;

N-[5-(6-Cyano-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-(6-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

5 N-{{[5-(6'-Fluoro-3'-oxospiro[cyclopropane-1,1'-isoindol]-2'(3'H)-yl)pyridin-3-yl]methyl}methanesulfonamide};

N-[5-((S or R)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

10 N-[5-(5-Fluoro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-((R or S)-5-Chloro-1-ethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

N-[5-(5-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

15 5-Chloro-2-[5-(1,1-dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

20 5-Chloro-3,3-dimethyl-2-[5-(2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-oxo-oxazolidin-3-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

25 5-Chloro-3,3-dimethyl-2-[5-(3-methyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1,1-dioxo-1λ6-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(3-isopropyl-2-oxo-imidazolidin-1-ylmethyl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 6-Chloro-2-[5-(1,5-dimethyl-1H-imidazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5-Chloro-3,3-dimethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

10 6-Chloro-2-[5-(3-methyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

15 6-Chloro-2-[5-(2,5-dimethyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1,5-dimethyl-1H-pyrazol-4-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-[4-chloro-5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

2-Chloro-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

2-Methoxy-7,7-dimethyl-6-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-6,7-dihydro-pyrrolo[3,4-b]pyridin-5-one;

25 5-Chloro-2-[5-(4-chloro-2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

(R or S)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

(S or R)-5-Chloro-3-ethyl-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5 3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

10 1-Methyl-1H-imidazole-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

2-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide;

Pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

15 3-Methyl-3H-imidazole-4-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-6-methyl-nicotinamide;

20 3-Chloro-N-[5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-isonicotinamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-nicotinamide;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-2-methyl-nicotinamide;

25 N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-4-methyl-nicotinamide;

2-[5-(1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-((R)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 2-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

10 5-Chloro-2-[5-(1-methanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((R)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 5-Chloro-2-[5-((S)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(1-Acetyl-piperidin-4-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 2-[5-(1-Acetyl-azetidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-3,3-dimethyl-2-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-methanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

25 5-Chloro-2-[5-(1-ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((S)-3-hydroxy-pyrrolidin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5 5-Chloro-3,3-dimethyl-2-[5-(4-propionyl-piperazin-1-yl)-pyridin-3-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(4-methanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

10 5-Chloro-2-[5-(4-ethanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-{5-[4-(3-chloro-pyridine-2-carbonyl)-piperazin-1-yl]-pyridin-3-yl}-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(1-Acetyl-pyrrolidin-3-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one ;

2-[6-(1-Acetyl-piperidin-3-yl)-pyrazin-2-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 5-Chloro-3,3-dimethyl-2-[6-(1-propionyl-piperidin-3-yl)-pyrazin-2-yl]-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[6-(1-ethanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[6-(1-methanesulfonyl-piperidin-3-yl)-pyrazin-2-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

25 N-[(S or R)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[4-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

5 and pharmaceutically acceptable salts thereof.

3. A compound according to claim 1 or 2, selected from

6-Chloro-2-pyridin-3-yl-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(4-chloro-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-isopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

10 6-Chloro-2-[5-((R)-1-hydroxy-ethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-(5-cyclopropoxy-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

2-Methoxy-6-(5-methoxy-pyridin-3-yl)-7,8-dihydro-6H-[1,6]naphthyridin-5-one;

5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

15 5-Chloro-2-(5-fluoro-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-(5-difluoromethoxy-pyridin-3-yl)-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

6-Chloro-2-[5-(2-oxa-6-aza-spiro[3.4]oct-6-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

20 6-Chloro-2-(5-methoxymethyl-pyridin-3-yl)-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(1-methyl-1H-pyrazol-4-ylmethyl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-{5-[(2-methyl-2H-pyrazol-3-yl)methyl]-amino}-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

6-Chloro-2-[5-(2-methyl-2H-pyrazol-3-yl)-pyridin-3-yl]-3,4-dihydro-2H-isoquinolin-1-one;

5 6-Chloro-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-3,4-dihydro-2H-isoquinolin-1-one;

and pharmaceutically acceptable salts thereof.

4. A compound according to claim 1 or 2, selected from

(R or S)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

10 (S or R)-5-Chloro-3-ethyl-2-(5-fluoro-pyridin-3-yl)-2,3-dihydro-isoindol-1-one;

N-[(R or S)-4-((R or S)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-propionamide;

15 N-[(R or S)-4-((S or R)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-5,6,7,8-tetrahydro-isoquinolin-8-yl]-acetamide;

N-[(R or S)-4-((S or R)-5-Chloro-3-ethyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-methanesulfonamide;

20 N-[(R or S)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

N-[(S or R)-4-((R or S)-5-Chloro-3-methyl-1-oxo-1,3-dihydro-isoindol-2-yl)-6,7-dihydro-5H-[2]pyrindin-7-yl]-propionamide;

5-Chloro-3,3-dimethyl-2-{5-[(1H-pyrazol-3-ylamino)-methyl]-pyridin-3-yl}-2,3-dihydro-isoindol-1-one;

6'-Chloro-2'-{5-[(1H-pyrazol-3-ylamino)methyl]pyridin-3-yl}spiro[cyclopropane-1,1'-isoindol]-3'(2'H)-one;

N-[5-(6-Chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-methanesulfonamide;

5 3-Methyl-pyridine-2-carboxylic acid [5-(6-chloro-1,1-dimethyl-3-oxo-1,3-dihydro-isoindol-2-yl)-pyridin-3-ylmethyl]-amide;

2-[5-((R)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

10 2-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-(1-methanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

5-Chloro-2-[5-((R)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

15 5-Chloro-2-[5-((S)-1-ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(1-Acetyl-piperidin-4-yloxy)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

20 5-Chloro-2-[5-(1-ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-[5-(4-Acetyl-piperazin-1-yl)-pyridin-3-yl]-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;


5-Chloro-2-[5-(4-methanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

25 5-Chloro-2-[5-(4-ethanesulfonyl-piperazin-1-yl)-pyridin-3-yl]-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

2-(1'-Acetyl-1',2',3',4',5',6'-hexahydro-[3,4']bipyridinyl-5-yl)-5-chloro-3,3-dimethyl-2,3-dihydro-isoindol-1-one;

and pharmaceutically acceptable salts thereof.

5. A process to prepare a compound according to any one of claims 1 to 4 comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



wherein R¹, R², R³, R⁴, R⁵, R⁶, A¹, A², A³, A⁴, A⁵ and n are as defined in claim 1 and wherein X is halogen or triflate.

6. A compound according to any one of claims 1 to 4 for use as therapeutically active substance.
7. A pharmaceutical composition comprising a compound according to any one of claims 1 to 4 and a therapeutically inert carrier.
8. A compound according to any one of claims 1 to 4 for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

9. The use of a compound according to any one of claims 1 to 4 for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.