

C5aR ANTAGONISTS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Serial No. 61/139,919 filed
5 December 22, 2008; the entire content of which is incorporated herein by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] NOT APPLICABLE

REFERENCE TO A "SEQUENCE LISTING," A TABLE, OR A COMPUTER PROGRAM LISTING APPENDIX SUBMITTED ON A COMPACT DISK.

[0003] NOT APPLICABLE

BACKGROUND OF THE INVENTION

[0004] The complement system plays a central role in the clearance of immune complexes
and in immune responses to infectious agents, foreign antigens, virus infected cells and tumor
cells. Inappropriate or excessive activation of the complement system can lead to harmful,
and even potentially life-threatening consequences due to severe inflammation and resulting
15 tissue destruction. These consequences are clinically manifested in various disorders
including septic shock; myocardial, as well as, intestinal ischemia/reperfusion injury; graft
rejection; organ failure; nephritis; pathological inflammation; and autoimmune diseases.

[0005] The complement system is composed of a group of proteins that are normally
present in the serum in an inactive state. Activation of the complement system encompasses
25 mainly three distinct pathways, *i.e.*, the classical, the alternative, and the lectin pathway (V.
M. Holers, *In Clinical Immunology: Principles and Practice*, ed. R. R. Rich, Mosby Press;
1996, 363-391): 1) The classical pathway is a calcium/magnesium-dependent cascade, which
is normally activated by the formation of antigen-antibody complexes. It can also be
activated in an antibody-independent manner by the binding of C-reactive protein, complexed
30 with ligand, and by many pathogens including gram-negative bacteria. 2) The alternative
pathway is a magnesium-dependent cascade which is activated by deposition and activation

of C3 on certain susceptible surfaces (e.g. cell wall polysaccharides of yeast and bacteria, and certain biopolymer materials). 3) The lectin pathway involves the initial binding of mannose-binding lectin and the subsequent activation of C2 and C4, which are common to the classical pathway (Matsushita, M. *et al.*, *J. Exp. Med.* 176: 1497-1502 (1992); Suankratay, C. *et al.*, *J. Immunol.* 160: 3006-3013 (1998)).

[0006] The activation of the complement pathway generates biologically active fragments of complement proteins, e.g. C3a, C4a and C5a anaphylatoxins and C5b-9 membrane attack complexes (MAC), all which mediate inflammatory responses by affecting leukocyte chemotaxis; activating macrophages, neutrophils, platelets, mast cells and endothelial cells; and increasing vascular permeability, cytolysis and tissue injury.

[0007] Complement C5a is one of the most potent proinflammatory mediators of the complement system. (The anaphylactic C5a peptide is 100 times more potent, on a molar basis, in eliciting inflammatory responses than C3a.) C5a is the activated form of C5 (190 kD, molecular weight). C5a is present in human serum at approximately 80 µg/ml (Kohler, P. F. *et al.*, *J. Immunol.* 99: 1211-1216 (1967)). It is composed of two polypeptide chains, α and β , with approximate molecular weights of 115 kD and 75 kD, respectively (Tack, B. F. *et al.*, *Biochemistry* 18: 1490-1497 (1979)). Biosynthesized as a single-chain promolecule, C5 is enzymatically cleaved into a two-chain structure during processing and secretion. After cleavage, the two chains are held together by at least one disulphide bond as well as noncovalent interactions (Ooi, Y. M. *et al.*, *J. Immunol.* 124: 2494-2498(1980)).

[0008] C5 is cleaved into the C5a and C5b fragments during activation of the complement pathways. The convertase enzymes responsible for C5 activation are multi-subunit complexes of C4b, C2a, and C3b for the classical pathway and of (C3b)₂, Bb, and P for the alternative pathway (Goldlust, M. B. *et al.*, *J. Immunol.* 113: 998-1007 (1974); Schreiber, R. D. *et al.*, *Proc. Natl. Acad. Sci.* 75: 3948-3952 (1978)). C5 is activated by cleavage at position 74-75 (Arg-Leu) in the α -chain. After activation, the 11.2 kD, 74 amino acid peptide C5a from the amino-terminus portion of the α -chain is released. Both C5a and C3a are potent stimulators of neutrophils and monocytes (Schindler, R. *et al.*, *Blood* 76: 1631-1638 (1990); Haeffner-Cavaillon, N. *et al.*, *J. Immunol.* 138: 794-700 (1987); Cavaillon, J. M. *et al.*, *Eur. J. Immunol.* 20: 253-257 (1990)).

[0009] In addition to its anaphylatoxic properties, C5a induces chemotactic migration of neutrophils (Ward, P. A. *et al.*, *J. Immunol.* 102: 93-99 (1969)), eosinophils (Kay, A. B. *et*

al., *Immunol.* 24: 969-976 (1973)), basophils (Lett-Brown, M. A. *et al.*, *J. Immunol.* 117: 246-252 1976)), and monocytes (Snyderman, R. *et al.*, *Proc. Soc. Exp. Biol. Med.* 138: 387-390 1971)). Both C5a and C5b-9 activate endothelial cells to express adhesion molecules essential for sequestration of activated leukocytes, which mediate tissue inflammation and injury (Foreman, K. E. *et al.*, *J. Clin. Invest.* 94: 1147-1155 (1994); Foreman, K. E. *et al.*, *Inflammation* 20: 1-9 (1996); Rollins, S. A. *et al.*, *Transplantation* 69: 1959-1967 (2000)). C5a also mediates inflammatory reactions by causing smooth muscle contraction, increasing vascular permeability, inducing basophil and mast cell degranulation and inducing release of lysosomal proteases and oxidative free radicals (Gerard, C. *et al.*, *Ann. Rev. Immunol.* 12: 775-808 (1994)). Furthermore, C5a modulates the hepatic acute-phase gene expression and augments the overall immune response by increasing the production of TNF- α , IL-1- β , IL-6, IL-8, prostaglandins and leukotrienes (Lambris, J. D. *et al.*, *In: The Human Complement System in Health and Disease*, Volanakis, J. E. ed., Marcel Dekker, New York, pp. 83-118).

[0010] The anaphylactic and chemotactic effects of C5a are believed to be mediated through its interaction with the C5a receptor. The human C5a receptor (C5aR) is a 52 kD membrane bound G protein-coupled receptor, and is expressed on neutrophils, monocytes, basophils, eosinophils, hepatocytes, lung smooth muscle and endothelial cells, and renal glomerular tissues (Van-Epps, D. E. *et al.*, *J. Immunol.* 132: 2862-2867 (1984); Haviland, D. L. *et al.*, *J. Immunol.* 154:1861-1869 (1995); Wetsel, R. A., *Immunol. Leff.* 44: 183-187 (1995); Buchner, R. R. *et al.*, *J. Immunol.* 155: 308-315 (1995); Chenoweth, D. E. *et al.*, *Proc. Natl. Acad. Sci.* 75: 3943-3947 (1978); Zwirner, J. *et al.*, *Mol. Immunol.* 36:877-884 (1999)). The ligand-binding site of C5aR is complex and consists of at least two physically separable binding domains. One binds the C5a amino terminus (amino acids 1-20) and disulfide-linked core (amino acids 21-61), while the second binds the C5a carboxy-terminal end (amino acids 62-74) (Wetsel, R. A., *Curr. Opin. Immunol.* 7: 48-53 (1995)).

[0011] C5a plays important roles in inflammation and tissue injury. In cardiopulmonary bypass and hemodialysis, C5a is formed as a result of activation of the alternative complement pathway when human blood makes contact with the artificial surface of the heart-lung machine or kidney dialysis machine (Howard, R. J. *et al.*, *Arch. Surg.* 123: 1496-1501 (1988); Kirklin, J. K. *et al.*, *J. Cardiovasc. Surg.* 86: 845-857 (1983); Craddock, P. R. *et al.*, *N. Engl. J. Med.* 296: 769-774 (1977)). C5a causes increased capillary permeability and edema, bronchoconstriction, pulmonary vasoconstriction, leukocyte and platelet activation and infiltration to tissues, in particular the lung (Czermak, B. J. *et al.*, *J. Leukoc. Biol.* 64: 40-

48 (1998)). Administration of an anti-C5a monoclonal antibody was shown to reduce cardiopulmonary bypass and cardioplegia-induced coronary endothelial dysfunction (Tofukuji, M. *et al.*, *J. Thorac. Cardiovasc. Surg.* 116: 1060-1068 (1998)).

[0012] C5a is also involved in acute respiratory distress syndrome (ARDS), Chronic

5 Obstructive Pulmonary Disorder (COPD) and multiple organ failure (MOF) (Hack, C. E. *et al.*, *Am. J. Med.* 1989; 86: 20-26; Hammerschmidt DE *et al. Lancet* 1980; 1: 947-949; Heideman M. *et al. J. Trauma* 1984; 4: 1038-1043; Marc, MM, *et al.*, *Am. J. Respir. Cell and Mol. Biol.*, 2004: 31: 216-219). C5a augments monocyte production of two important pro-inflammatory cytokines, TNF- α and IL-1. C5a has also been shown to play an important role
10 in the development of tissue injury, and particularly pulmonary injury, in animal models of septic shock (Smedegard G *et al. Am. J. Pathol.* 1989; 135: 489-497; Markus, S., *et al.*, *FASEB Journal* (2001), 15: 568-570). In sepsis models using rats, pigs and non-human primates, anti-C5a antibodies administered to the animals before treatment with endotoxin or E. coli resulted in decreased tissue injury, as well as decreased production of IL-6
15 (Smedegard, G. *et al.*, *Am. J. Pathol.* 135: 489-497 (1989); Hopken, U. *et al.*, *Eur. J. Immunol.* 26: 1103-1109 (1996); Stevens, J. H. *et al.*, *J. Clin. Invest.* 77: 1812-1816 (1986)). More importantly, blockade of C5a with anti-C5a polyclonal antibodies has been shown to significantly improve survival rates in a caecal ligation/puncture model of sepsis in rats (Czermak, B.J. *et al.*, *Nat. Med.* 5: 788-792 (1999)). This model share many aspects of the
20 clinical manifestation of sepsis in humans. (Parker, S.J. *et al.*, *Br. J. Surg.* 88: 22-30 (2001)). In the same sepsis model, anti-C5a antibodies were shown to inhibit apoptosis of thymocytes (Guo, R.F. *et al.*, *J. Clin. Invest.* 106: 1271-1280 (2000)) and prevent MOF (Huber-Lang, M. *et al.*, *J. Immunol.* 166: 1193-1199 (2001)). Anti-C5a antibodies were also protective in a cobra venom factor model of lung injury in rats, and in immune complex-induced lung injury
25 (Mulligan, M. S. *et al. J. Clin. Invest.* 98: 503-512 (1996)). The importance of C5a in immune complex-mediated lung injury was later confirmed in mice (Bozic, C. R. *et al.*, *Science* 26: 1103-1109 (1996)).

[0013] C5a is found to be a major mediator in myocardial ischemia-reperfusion injury.

Complement depletion reduced myocardial infarct size in mice (Weisman, H. F. *et al.*,
30 *Science* 249: 146-151 (1990)), and treatment with anti-C5a antibodies reduced injury in a rat model of hindlimb ischemia-reperfusion (Bless, N. M. *et al.*, *Am. J. Physiol.* 276: L57-L63 (1999)). Reperfusion injury during myocardial infarction was also markedly reduced in pigs that were retreated with a monoclonal anti-C5a IgG (Amsterdam, E. A. *et al.*, *Am. J. Physiol.*

268:H448-H457 (1995)). A recombinant human C5aR antagonist reduces infarct size in a porcine model of surgical revascularization (Riley, R. D. *et al.*, *J. Thorac. Cardiovasc. Surg.* 120: 350-358 (2000)).

[0014] C5a driven neutrophils also contribute to many bullous diseases (e.g., bullous pemphigoid, pemphigus vulgaris and pemphigus foliaceus). These are chronic and recurring inflammatory disorders clinically characterized by sterile blisters that appear in the sub-epidermal space of the skin and mucosa. While autoantibodies to keratinocytes located at the cutaneous basement membranes are believed to underlie the detachment of epidermal basal keratinocytes from the underlying basement membrane, blisters are also characterized by accumulation of neutrophils in both the upper dermal layers and within the blister cavities. In experimental models a reduction of neutrophils or absence of complement (total or C5-selective) can inhibit formation of sub-epidermal blisters, even in the presence of high auto-antibody titers.

[0015] Complement levels are elevated in patients with rheumatoid arthritis (Jose, P. J. *et al.*, *Ann. Rheum. Dis.* 49: 747-752 (1990); Grant, E.P., *et al.*, *J. of Exp. Med.*, 196(11): 1461-1471, (2002)), lupus nephritis (Bao, L., *et al.*, *Eur. J. of Immunol.*, 35(8), 2496-2506, (2005)) and systemic lupus erythematosus (SLE) (Porcel, J. M. *et al.*, *Clin. Immunol. Immunopathol.* 74: 283-288 (1995)). C5a levels correlate with the severity of the disease state. Collagen-induced arthritis in mice and rats resembles the rheumatoid arthritic disease in human. Mice deficient in the C5a receptor demonstrated a complete protection from arthritis induced by injection of monoclonal anti-collagen Abs (Banda, N.K., *et al.*, *J. of Immunol.*, 2003, 171: 2109-2115). Therefore, inhibition of C5a and/or C5a receptor (C5aR) could be useful in treating these chronic diseases.

[0016] The complement system is believed to be activated in patients with inflammatory bowel disease (IBD) and is thought to play a role in the disease pathogenesis. Activated complement products were found at the luminal face of surface epithelial cells, as well as in the muscularis mucosa and submucosal blood vessels in IBD patients (Woodruff, T.M., *et al.*, *J of Immunol.*, 2003, 171: 5514-5520).

[0017] C5aR expression is upregulated on reactive astrocytes, microglia, and endothelial cells in an inflamed human central nervous system (Gasque, P. *et al.*, *Am. J. Pathol.* 150: 31-41 (1997)). C5a might be involved in neurodegenerative diseases, such as Alzheimer disease (Mukherjee, P. *et al.*, *J. Neuroimmunol.* 105: 124-130 (2000); O'Barr, S. *et al.*, *J.*

Neuroimmunol. (2000) 105: 87-94; Farkas, I., *et al. J. Immunol.* (2003) 170:5764-5771), Parkinson's disease, Pick disease and transmissible spongiform encephalopathies. Activation of neuronal C5aR may induce apoptosis (Farkas I *et al. J. Physiol.* 1998; 507: 679-687).

Therefore, inhibition of C5a and/or C5aR could also be useful in treating neurodegenerative diseases.

[0018] There is some evidence that C5a production worsens inflammation associated with atopic dermatitis (Neuber, K., *et al., Immunology* 73:83-87, (1991)), and chronic urticaria (Kaplan, A.P., *J. Allergy Clin. Immunol.* 114; 465-474, (2004).

[0019] Psoriasis is now known to be a T cell-mediated disease (Gottlieb, E. L. *et al., Nat. Med.* 1: 442-447 (1995)). However, neutrophils and mast cells may also be involved in the pathogenesis of the disease (Terui, T. *et al., Exp. Dermatol.* 9: 1-10; 2000); Werfel, T. *et al., Arch. Dermatol. Res.* 289: 83-86 (1997)). Neutrophil accumulation under the stratum corneum is observed in the highly inflamed areas of psoriatic plaques, and psoriatic lesion (scale) extracts contain highly elevated levels of C5a and exhibit potent chemotactic activity towards neutrophils, an effect that can be inhibited by addition of a C5a antibody. T cells and neutrophils are chemo-attracted by C5a (Nataf, S. *et al., J. Immunol.* 162: 4018-4023 (1999); Tsuji, R. F. *et al., J. Immunol.* 165: 1588-1598 (2000); Cavaillon, J. M. *et al., Eur. J. Immunol.* 20: 253-257 (1990)). Additionally expression of C5aR has been demonstrated in plasmacytoid dendritic cells (pDC) isolated from lesions of cutaneous lupus erythematosus and these cells were shown to display chemotactic behavior towards C5a, suggesting that blockade of C5aR on pDC might be efficacious in reducing pDC infiltration into inflamed skin in both SLE and psoriasis. Therefore C5a could be an important therapeutic target for treatment of psoriasis.

[0020] Immunoglobulin G-containing immune complexes (IC) contribute to the pathophysiology in a number of autoimmune diseases, such as systemic lupus erythematosus, rheumatoid arthritis, Sjogren's disease, Goodpasture's syndrome, and hypersensitivity pneumonitis (Madaio, M. P., *Semin. Nephrol.* 19: 48-56 (1999); Korganow, A. S. *et al., Immunity* 10: 451-459 (1999); Bolten, W. K., *Kidney Int.* 50: 1754-1760 (1996); Ando, M. *et al., Curr. Opin. Pulm. Med.* 3: 391-399 (1997)). These diseases are highly heterogeneous and generally affect one or more of the following organs: skin, blood vessels, joints, kidneys, heart, lungs, nervous system and liver (including cirrhosis and liver fibrosis). The classical animal model for the inflammatory response in these IC diseases is the Arthus reaction,

which features the infiltration of polymorphonuclear cells, hemorrhage, and plasma exudation (Arthus, M., *C.R. Soc. Biol.* 55: 817-824 (1903)). Recent studies show that C5aR deficient mice are protected from tissue injury induced by IC (Kohl, J. *et al.*, *Mol. Immunol.* 36: 893-903 (1999); Baumann, U. *et al.*, *J. Immunol.* 164: 1065-1070 (2000)). The results are

5 consistent with the observation that a small peptidic anti-C5aR antagonist inhibits the inflammatory response caused by IC deposition (Strachan, A. J. *et al.*, *J. Immunol.* 164: 6560-6565 (2000)). Together with its receptor, C5a plays an important role in the pathogenesis of IC diseases. Inhibitors of C5a and C5aR could be useful to treat these diseases.

10 Description of Related Art:

[0021] Only recently have non-peptide based C5a receptor antagonists been described in the literature (*e.g.*, Sumichika, H., *et al.*, *J. Biol. Chem.* (2002), 277, 49403-49407). Non-peptide based C5a receptor antagonist have been reported as being effective for treating endotoxic shock in rats (Stracham, A.J., *et al.*, *J. of Immunol.* (2000), 164(12): 6560-6565);


15 and for treating IBD in a rat model (Woodruff, T.M., *et al.*, *J of Immunol.*, 2003, 171: 5514-5520). Non-peptide based C5a receptor modulators also have been described in the patent literature by Neurogen Corporation, (*e.g.*, WO2004/043925, WO2004/018460, WO2005/007087, WO03/082826, WO03/08828, WO02/49993, WO03/084524); Dompe S.P.A. (WO02/029187); and The University of Queensland (WO2004/100975).

20 [0022] There is considerable experimental evidence in the literature that implicates increased levels of C5a with a number of diseases and disorders, in particular in autoimmune and inflammatory diseases and disorders. Thus, there remains a need in the art for new small organic molecule modulators, *e.g.*, agonists, preferably antagonists, partial agonists, of the C5a receptor (C5aR) that are useful for inhibiting pathogenic events, *e.g.*, chemotaxis,

25 associated with increased levels anaphylatoxin activity. The present invention fulfills this and other needs.

BRIEF SUMMARY OF THE INVENTION

[0023] In one aspect, the present invention provide compounds having the formula:



and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein

C^1 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^1 substituents;

5 C^2 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^2 substituents;

C^3 is selected from the group consisting of C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl- C_{1-4} alkyl, aryl, aryl- C_{1-4} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heterocycloalkyl or
10 heterocycloalkyl- C_{1-4} alkyl, wherein the heterocycloalkyl group or portion has from 1-3 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S, and each C^3 is optionally substituted with from 1-3 R^3 substituents;

each R^1 is independently selected from the group consisting of halogen, -CN, $-R^c$, $-CO_2R^a$,
15 $-CONR^aR^b$, $-C(O)R^a$, $-OC(O)NR^aR^b$, $-NR^bC(O)R^a$, $-NR^bC(O)_2R^c$, $-NR^a-C(O)NR^aR^b$,
 $-NR^aC(O)NR^aR^b$, $-NR^aR^b$, $-OR^a$, and $-S(O)_2NR^aR^b$; wherein each R^a and R^b is
independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to
the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
membered ring having from 0 to 2 additional heteroatoms as ring members selected from
20 N, O or S; each R^c is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic
and cyclic portions of R^a , R^b and R^c are optionally further substituted with from one to
three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and
optionally when two R^1 substituents are on adjacent atoms, are combined to form a fused
25 five or six-membered carbocyclic ring;

each R^2 is independently selected from the group consisting of halogen, -CN, $-R^f$, $-CO_2R^d$,
 $-CONR^dR^e$, $-C(O)R^d$, $-OC(O)NR^dR^e$, $-NR^eC(O)R^d$, $-NR^eC(O)_2R^f$, $-NR^dC(O)NR^dR^e$,
 $-NR^dC(O)NR^dR^e$, $-NR^dR^e$, $-OR^d$, and $-S(O)_2NR^dR^e$; wherein each R^d and R^e is
independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to
the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
membered ring having from 0 to 2 additional heteroatoms as ring members selected from
30 N, O or S; each R^f is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic

and cyclic portions of R^d , R^e and R^f are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; each R^3 is independently selected from the group consisting of halogen, $-CN$, $-R^i$, $-CO_2R^g$, $-CONR^gR^h$, $-C(O)R^g$, $-OC(O)NR^gR^h$, $-NR^hC(O)R^g$, $-NR^hC(O)_2R^i$, $-NR^gC(O)NR^gR^h$, $-NR^gR^h$, $-OR^g$, $-S(O)_2NR^gR^h$, $-X^4-R^j$, $-X^4-NR^gR^h$, $-X^4-CONR^gR^h$, $-X^4-NR^hC(O)R^g$, $-NHR^j$ and $-NHCH_2R^j$, wherein X^4 is a C_{1-4} alkylene; each R^g and R^h is independently selected from hydrogen, C_{1-8} alkyl, C_{3-6} cycloalkyl and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S and is optionally substituted with one or two oxo; each R^i is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of C_{3-6} cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g , R^h , R^i and R^j are optionally further substituted with from one to three halogen, methyl, CF_3 , hydroxy, amino, alkylamino and dialkylamino groups; and X is hydrogen or CH_3 .

[0024] In addition to the compounds provided herein, the present invention further provides pharmaceutical compositions containing one or more of these compounds, as well as methods for the use of these compounds in therapeutic methods, primarily to treat diseases associated C5a signalling activity.

[0025] In yet another aspect, the present invention provides methods of diagnosing disease in an individual. In these methods, the compounds provided herein are administered in labeled form to a subject, followed by diagnostic imaging to determine the presence or absence of C5aR7. In a related aspect, a method of diagnosing disease is carried out by contacting a tissue or blood sample with a labeled compound as provided herein and determining the presence, absence, or amount of C5aR in the sample.

BRIEF DESCRIPTION OF THE DRAWINGS

[0026] **Figure 1** provides structures and activity for representative compounds of the present invention. The compounds were prepared usually methods as described generally below, as well as methods provided in the Examples.

DETAILED DESCRIPTION OF THE INVENTION

I. Abbreviation and Definitions

[0027] The term "alkyl", by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain hydrocarbon radical, having the number of carbon atoms designated (*i.e.* C₁₋₈ means one to eight carbons). Examples of alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. The term "alkenyl" refers to an unsaturated alkyl group having one or more double bonds. Similarly, the term "alkynyl" refers to an unsaturated alkyl group having one or more triple bonds. Examples of such unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. The term "cycloalkyl" refers to hydrocarbon rings having the indicated number of ring atoms (*e.g.*, C₃₋₆cycloalkyl) and being fully saturated or having no more than one double bond between ring vertices. "Cycloalkyl" is also meant to refer to bicyclic and polycyclic hydrocarbon rings such as, for example, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, etc. The term "heterocycloalkyl" refers to a cycloalkyl group that contain from one to five heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. The heterocycloalkyl may be a monocyclic, a bicyclic or a polycyclic ring system. Non limiting examples of heterocycloalkyl groups include pyrrolidine, imidazolidine, pyrazolidine, butyrolactam, valerolactam, imidazolidinone, hydantoin, dioxolane, phthalimide, piperidine, 1,4-dioxane, morpholine, thiomorpholine, thiomorpholine-S-oxide, thiomorpholine-S,S-oxide, piperazine, pyran, pyridone, 3-pyrroline, thiopyran, pyrone, tetrahydrofuran, tetrahydrothiophene, quinuclidine, and the like. A heterocycloalkyl group can be attached to the remainder of the molecule through a ring carbon or a heteroatom.

[0028] The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having four or fewer carbon atoms. Similarly, "alkenylene" and "alkynylene" refer to the unsaturated forms of "alkylene" having double or triple bonds, respectively.

[0029] The term "heteroalkyl," by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S may be placed at any interior position of the heteroalkyl group. The heteroatom Si may be placed at any position of the heteroalkyl group, including the position at which the alkyl group is attached to the remainder of the molecule. Examples include -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Similarly, the terms "heteroalkenyl" and "heteroalkynyl" by itself or in combination with another term, means, unless otherwise stated, an alkenyl group or alkynyl group, respectively, that contains the stated number of carbons and having from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S may be placed at any interior position of the heteroalkyl group.

[0030] The term "heteroalkylene" by itself or as part of another substituent means a divalent radical, saturated or unsaturated or polyunsaturated, derived from heteroalkyl, as exemplified by -CH₂-CH₂-S-CH₂CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-, -O-CH₂-CH=CH-, -CH₂-CH=C(H)CH₂-O-CH₂- and -S-CH₂-C≡C-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (*e.g.*, alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like).

[0031] The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively. Additionally, for dialkylamino groups, the alkyl portions can be the same or different and can also be combined to form a 3-7 membered ring with the nitrogen atom to which each is attached. Accordingly, a group represented as -NR^aR^b is meant to include piperidinyl, pyrrolidinyl, morpholinyl, azetidiny and the like.

[0032] The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as "haloalkyl," are meant to include monohaloalkyl and polyhaloalkyl. For example, the term "C₁₋₄ haloalkyl" is meant to include trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0033] The term "aryl" means, unless otherwise stated, a polyunsaturated, typically aromatic, hydrocarbon group which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. The term "heteroaryl" refers to aryl groups (or rings) that contain from one to five heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl groups include phenyl, naphthyl and biphenyl, while non-limiting examples of heteroaryl groups include pyridyl, pyridazinyl, pyrazinyl, pyrimidinyl, triazinyl, quinolinyl, quinoxalinyl, quinazolinyl, cinnolinyl, phthalazinyl, benzotriazinyl, purinyl, benzimidazolyl, benzopyrazolyl, benzotriazolyl, benzisoxazolyl, isobenzofuryl, isoindolyl, indoliziny, benzotriazinyl, thienopyridinyl, thienopyrimidinyl, pyrazolopyrimidinyl, imidazopyridines, benzothiazolyl, benzofuranyl, benzothienyl, indolyl, quinolyl, isoquinolyl, isothiazolyl, pyrazolyl, indazolyl, pteridinyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiadiazolyl, pyrrolyl, thiazolyl, furyl, thienyl and the like. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0034] For brevity, the term "aryl" when used in combination with other terms (*e.g.*, aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above. Thus, the term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like).

[0035] The above terms (*e.g.*, "alkyl," "aryl" and "heteroaryl"), in some embodiments, will include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below. For brevity, the terms aryl and heteroaryl will refer to substituted or unsubstituted versions as provided below, while the term "alkyl" and related aliphatic radicals is meant to refer to unsubstituted version, unless indicated to be substituted.

[0036] Substituents for the alkyl radicals (including those groups often referred to as alkylene, alkenyl, alkynyl and cycloalkyl) can be a variety of groups selected from: -halogen, -OR', -NR'R'', -SR', -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR'S(O)₂R'', -CN and -NO₂ in a number ranging from zero to (2 m' + 1), where m' is the total number of carbon atoms in such radical. R', R'' and R''' each independently refer to hydrogen, unsubstituted C₁₋₈ alkyl, unsubstituted heteroalkyl, unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted C₁₋₈ alkyl, C₁₋₈ alkoxy or C₁₋₈ thioalkoxy groups, or unsubstituted aryl-C₁₋₄ alkyl groups. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 3-, 4-, 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-morpholinyl. The term "acyl" as used by itself or as part of another group refers to an alkyl radical wherein two substituents on the carbon that is closest to the point of attachment for the radical is replaced with the substituent =O (e.g., -C(O)CH₃, -C(O)CH₂CH₂OR' and the like).

[0037] Similarly, substituents for the aryl and heteroaryl groups are varied and are generally selected from: -halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R'', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR'S(O)₂R'', -N₃, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'' and R''' are independently selected from hydrogen, C₁₋₈ alkyl, C₃₋₆ cycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-C₁₋₄ alkyl, and unsubstituted aryloxy-C₁₋₄ alkyl. Other suitable substituents include each of the above aryl substituents attached to a ring atom by an alkylene tether of from 1-4 carbon atoms.

[0038] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U-, wherein T and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from 0 to 2. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the

aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-(CH_2)_s-X-(CH_2)_t-$, where s and t are independently integers of from 0 to 3, and X is $-O-$, $-NR'-$, $-S-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)_2NR'-$. The substituent R' in $-NR'-$ and $-S(O)_2NR'-$ is selected from hydrogen or unsubstituted C_{1-6} alkyl.

[0039] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

[0040] The term "ionic liquid" refers to any liquid that contains mostly ions. Preferably, in the present invention, "ionic liquid" refers to the salts whose melting point is relatively low (e.g., below 250 °C). Examples of ionic liquids include but are not limited to 1-butyl-3-methylimidazolium tetrafluoroborate, 1-hexyl-3-methylimidazolium tetrafluoroborate, 1-octyl-3-methylimidazolium tetrafluoroborate, 1-nonyl-3-methylimidazolium tetrafluoroborate, 1-decyl-3-methylimidazolium tetrafluoroborate, 1-hexyl-3-methylimidazolium hexafluorophosphate and 1-hexyl-3-methylimidazolium bromide, and the like.

[0041] The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of salts derived from pharmaceutically-acceptable inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, zinc and the like. Salts derived from pharmaceutically-acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted amines, cyclic amines, naturally-occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N' -dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N -ethylmorpholine, N -ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid,

either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0042] The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

[0043] In addition to salt forms, the present invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.


[0044] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0045] Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers,

regioisomers and individual isomers (*e.g.*, separate enantiomers) are all intended to be encompassed within the scope of the present invention. The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

II. Compounds

[0046] In one aspect, the present invention provides compounds having the formula **I**:



and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein

C^1 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^1 substituents;

C^2 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^2 substituents;

C^3 is selected from the group consisting of C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl- C_{1-4} alkyl, aryl, aryl- C_{1-4} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heterocycloalkyl or heterocycloalkyl- C_{1-4} alkyl, wherein the heterocycloalkyl group or portion has from 1-3 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S, and each C^3 is optionally substituted with from 1-3 R^3 substituents;

each R^1 is independently selected from the group consisting of halogen, $-\text{CN}$, $-\text{R}^c$, $-\text{CO}_2\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{C}(\text{O})\text{R}^a$, $-\text{OC}(\text{O})\text{NR}^a\text{R}^b$, $-\text{NR}^b\text{C}(\text{O})\text{R}^a$, $-\text{NR}^b\text{C}(\text{O})_2\text{R}^c$, $-\text{NR}^a-\text{C}(\text{O})\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{R}^b$, $-\text{OR}^a$, and $-\text{S}(\text{O})_2\text{NR}^a\text{R}^b$; wherein each R^a and R^b is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from

N, O or S; each R^c is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic and cyclic portions of R^a , R^b and R^c are optionally further substituted with from one to

three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and
 5 optionally when two R^1 substituents are on adjacent atoms, are combined to form a fused five or six-membered carbocyclic ring;

each R^2 is independently selected from the group consisting of halogen, $-CN$, $-R^f$, $-CO_2R^d$, $-CONR^dR^e$, $-C(O)R^d$, $-OC(O)NR^dR^e$, $-NR^eC(O)R^d$, $-NR^eC(O)_2R^f$, $-NR^dC(O)NR^dR^e$, $-NR^dC(O)NR^dR^e$, $-NR^dR^e$, $-OR^d$, and $-S(O)_2NR^dR^e$; wherein each R^d and R^e is

10 independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S; each R^f is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic and cyclic portions of R^d , R^e and R^f are optionally further substituted with from one to
 15 three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups;

each R^3 is independently selected from the group consisting of halogen, $-CN$, $-R^i$, $-CO_2R^g$, $-CONR^gR^h$, $-C(O)R^g$, $-OC(O)NR^gR^h$, $-NR^hC(O)R^g$, $-NR^hC(O)_2R^i$, $-NR^gC(O)NR^gR^h$, $-NR^gR^h$, $-OR^g$, $-S(O)_2NR^gR^h$, $-X^4-R^j$, $-X^4-NR^gR^h$, $-X^4-CONR^gR^h$, $-X^4-NR^hC(O)R^g$, $-NHR^j$

20 and $-NHCH_2R^j$, wherein X^4 is a C_{1-4} alkylene; each R^g and R^h is independently selected from hydrogen, C_{1-8} alkyl, C_{3-6} cycloalkyl and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S and is optionally substituted with one or two oxo; each R^i is independently

25 selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of C_{3-6} cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g , R^h , R^i and R^j are optionally further substituted with from one to three halogen, methyl, CF_3 , hydroxy,

30 amino, alkylamino and dialkylamino groups; and


X is hydrogen or CH_3 .

[0047] In formula I, the substituent C^1 is, in one embodiment, selected from the group consisting of phenyl, pyridyl, indolyl and thiazolyl, each of which is optionally substituted


with from 1 to 3 R^1 substituents. Preferably, each R^1 is independently selected from the group consisting of halogen, -CN, $-R^c$, $-NR^aR^b$ and $-OR^a$, and wherein each R^a and R^b is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a pyrrolidine ring; each

5 R^c is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl and C_{3-6} cycloalkyl, and wherein the aliphatic and cyclic portions of R^a , R^b and R^c are optionally further substituted with from one to three hydroxy, methyl, amino, alkylamino and dialkylamino groups; and optionally when two R^1 substituents are on adjacent atoms, are combined to form a fused five or six-membered carbocyclic ring. In selected embodiments

10 of the invention, C^1 is selected from:




[0048] Returning to formula **I**, the substituents C^2 is, in one embodiment, selected from the group consisting of phenyl, naphthyl, pyridyl and indolyl, each of which is optionally substituted with from 1 to 3 R^2 substituents. Preferably, each R^2 is independently selected from the group consisting of halogen, $-R^f$ and $-OR^d$; wherein each R^d is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl; each R^f is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl and heteroaryl, and wherein the aliphatic and cyclic portions of R^d and R^f are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups. In selected embodiments of the invention, C^2 is selected from the group consisting of:




[0049] The substituents C^3 is, in some embodiments, selected from the group consisting of C_{3-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-2} alkyl, phenyl, pyridinyl, pyrazolyl, piperidinyl, pyrrolidinyl, piperidinylmethyl and pyrrolidinylmethyl, each of which is optionally substituted with from 1 to 3 R^3 substituents. Preferably, each R^3 is independently selected from the group consisting of halogen, $-R^i$, $-CO_2R^g$, $-CONR^gR^h$, $-NR^hC(O)R^g$, $-NR^hC(O)_2R^i$, $-NR^gR^h$, $-OR^g$, $-X^4-R^j$, $-X^4-NR^gR^h$, $-X^4-CONR^gR^h$, $-X^4-NR^hC(O)R^g$, $-NHR^j$ and $-NHCH_2R^j$, wherein X^4 is a C_{1-3} alkylene; each R^g and R^h is independently selected from hydrogen, C_{1-8}

alkyl, C₃₋₆ cycloalkyl and C₁₋₈ haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 1 additional heteroatoms as ring members selected from N, O or S and is optionally substituted with one or two oxo; each Rⁱ is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of C₃₋₆ cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g, R^h, Rⁱ and R^j are optionally further substituted with from one to three halogen, methyl, CF₃, hydroxy, amino, alkylamino and dialkylamino groups. In selected embodiments of the invention, C³ is selected from the group consisting of:




In other embodiments, C^3 is selected from the group consisting of:




[0050] Returning to formula I, X is preferably H.

Subformulae of Formula I:


5 **[0051]** In one embodiment of the invention, compounds of formula **I** have subformula **1a**:



[0052] In a second embodiment of the invention, compounds of formula **I** have subformula **Ib**:




[0053] In a third embodiment of the invention, compounds of formula **I** have subformula **Ic**:




wherein X^1 is selected from the group consisting of N, CH and CR^1 ; the subscript n is an integer of from 0 to 2; X^2 is selected from the group consisting of N, CH and CR^2 ; and the subscript m is an integer of from 0 to 2.

[0054] In a fourth embodiment of the invention, compounds of formula **I** have subformula **Id**:




wherein X^1 is selected from the group consisting of N, CH and CR^1 ; the subscript n is an integer of from 0 to 2; X^2 is selected from the group consisting of N, CH and CR^2 ; and the subscript m is an integer of from 0 to 2.

[0055] In a fifth embodiment of the invention, compounds of formula **I** have subformula **Ie**:




wherein the subscript p is an integer of from 0 to 3; X^1 is selected from the group consisting of N, CH and CR^1 ; the subscript n is an integer of from 0 to 2; X^2 is selected from the group consisting of N, CH and CR^2 ; and the subscript m is an integer of from 0 to 2.

5 **[0056]** In other selected embodiments, the compounds of the invention are represented by:



wherein the substituents R^1 , R^2 and R^3 , and the subscript p all have the meanings provided with reference to formula **I**.

10 **[0057]** In still other selected embodiments, the compounds of the invention are represented by:



wherein the substituents R^1 and R^3 , and the subscript p all have the meanings provided with reference to formula **I**.

[0058] In a particularly preferred group of embodiments, the compounds of the invention are represented by formula **(Ie⁵)** wherein R^3 is a member selected from the group consisting

of $-\text{NR}^g\text{R}^h$, $-\text{NHR}^j$ and $-\text{NHCH}_2\text{R}^j$, and each R^g , R^h and R^j have the meanings provided with reference to formula I.





[0059] In another particularly preferred group of embodiments, the compounds of the invention are represented by formula (Ie⁵) wherein R^3 is a member selected from the group consisting of $-\text{X}^4-\text{NR}^g\text{R}^h$, $-\text{X}^4-\text{R}^j$ and $-\text{X}^4-\text{NR}^h\text{COR}^g$, and each of X^4 , R^g , R^h and R^j have the meanings provided with reference to formula I.

[0060] Compounds of the invention having formula I can exist in different diastereomeric forms, *e.g.*, the substituents C^1 and C^2 in subformulae Ia and Ic can be *cis* to each other or *trans* to each other. As used herein, the terms *cis* or *trans* are used in their conventional sense in the chemical arts, *i.e.*, referring to the position of the substituents to one another relative to a reference plane, *e.g.*, a double bond, or a ring system, such as a decalin-type ring system or a hydroquinolone ring system: in the *cis* isomer, the substituents are on the same side of the reference plane, in the *trans* isomer the substituents are on opposite sides. Additionally, different conformers are contemplated by the present invention, as well as distinct rotamers. Conformers are conformational isomers that can differ by rotations about one or more σ bonds. Rotamers are conformers that differ by rotation about only a single σ bond.

Preparation of Compounds


[0061] Those skilled in the art will recognize that there are a variety of methods available to synthesize molecules represented in the claims. In general, useful methods for synthesizing compounds represented in the claims consist of four parts, which may be done in any order: Formation of the piperidine ring, installation of two amide bonds, and installation and/or modification of functional groups on C^1 , C^2 , and C^3 .

[0062] Several methods for the preparation of claimed compounds are illustrated below (eq. 1-6).



[0063] Equations 1-4 demonstrate some methods of forming the piperidine ring. Coupling at the 2-position of the pyridine ring can be accomplished via transition metal mediated couplings as shown in eq. 1-2, or metal catalyzed addition of an organometallic species such as the zincate or magnesium salt (eq. 3). Subsequent to coupling at the 2-position, transition metal mediated hydrogenation of the pyridine ring yields the piperidine ring system (eq. 1-3). Another method results in elaboration of a β -amino acid to a piperidine ring as described in eq. 4. Those skilled in the art will recognize that many synthetic methodologies can yield substituted piperidines, including C-C or C-N cyclization of acyclic precursors via alkylation or ring-closing metathesis. Relative stereochemistry may be set by a variety of methods, including facial selectivity during the hydrogenation step. Absolute stereochemistry may also be set via a variety of methods, via the use of chiral ligands or a chiral auxiliary, separation of chiral diastereoisomers, use of chiral starting materials, or classical resolution. Compounds


with 2,3-*trans* stereochemistry may have the relative stereochemistry set during the piperidine formation, or may be derived via epimerization of a 2,3-*cis* piperidine as illustrated in eq. 5.



[0064] Acylation of the piperidine ring is described in equation 6. In the case of eq. 6, X may be chosen from an appropriate group such as OH, Cl and F, or from any group capable of activating a carbonyl group for addition of an amine (e.g., OSu, imidazole, etc.). Such couplings may be assisted by the use of inorganic or organic bases, activating agents such as HBTU, and also by catalysts, in particular by those catalysts known in the art which assist in the formation of amide bonds, such as DMAP, HOBT, etc. Suitable coupling partners include a carboxylic acid and a piperidine, an acyl fluoride and an amine and so forth. Those skilled in the art will recognize that there are other possible combinations which will also result in the desired product.

10

15



[0065] A variety of methods described above have been used to prepare compounds of the invention, some of which are described in the examples.

20

[0066] A family of specific compounds of particular interest having formula I consists of compounds, pharmaceutically acceptable salts, hydrates and rotomers thereof, as set forth in Figure 1.

25

III. Pharmaceutical Compositions

[0067] In addition to the compounds provided above, compositions for modulating C5a activity in humans and animals will typically contain a pharmaceutical carrier or diluent.

[0068] The term "composition" as used herein is intended to encompass a product

comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

[0069] The pharmaceutical compositions for the administration of the compounds of this invention may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy and drug delivery. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active object compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases.

[0070] The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions and self emulsifications as described in U.S. Patent Application 2002-0012680, hard or soft capsules, syrups, elixirs, solutions, buccal patch, oral gel, chewing gum, chewable tablets, effervescent powder and effervescent tablets.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents, antioxidants and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as cellulose, silicon dioxide, aluminum oxide, calcium carbonate, sodium carbonate, glucose, mannitol, sorbitol,

lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example PVP, cellulose, PEG, starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated, enterically or otherwise, by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in the U.S. Pat. Nos. 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

[0071] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil. Additionally, emulsions can be prepared with a non-water miscible ingredient such as oils and stabilized with surfactants such as mono-diglycerides, PEG esters and the like.

[0072] Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0073] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example

beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

[0074] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[0075] The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

[0076] Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. Oral solutions can be prepared in combination with, for example, cyclodextrin, PEG and surfactants.

[0077] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[0078] The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary

temperatures but liquid at the rectal temperature and will therefore melt in the rectum to

release the drug. Such materials include cocoa butter and polyethylene glycols.

Additionally, the compounds can be administered via ocular delivery by means of solutions or ointments. Still further, transdermal delivery of the subject compounds can be

accomplished by means of iontophoretic patches and the like. For topical use, creams,

ointments, jellies, solutions or suspensions, etc., containing the compounds of the present

invention are employed. As used herein, topical application is also meant to include the use of mouth washes and gargles.

[0079] The compounds of this invention may also be coupled a carrier that is a suitable

polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran

copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-

phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore,

the compounds of the invention may be coupled to a carrier that is a class of biodegradable

polymers useful in achieving controlled release of a drug, for example polylactic acid,

polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone,

polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans,

polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels.

Polymers and semipermeable polymer matrices may be formed into shaped articles, such as

valves, stents, tubing, prostheses and the like. In one embodiment of the invention, the

compound of the invention is coupled to a polymer or semipermeable polymer matrix that is formed as a stent or stent-graft device.

IV. Methods of Treating Diseases and Disorders Modulated by C5a

[0080] The compounds of the invention may be used as agonists, (preferably) antagonists,

partial agonists, inverse agonists, of C5a receptors in a variety of contexts, both *in vitro* and

in vivo. In one embodiment, the compounds of the invention are C5aR antagonist that can be

used to inhibit the binding of C5a receptor ligand (*e.g.*, C5a) to C5a receptor *in vitro* or *in*

vivo. In general, such methods comprise the step of contacting a C5a receptor with a

sufficient amount of one or more C5a receptor modulators as provided herein, in the presence of C5a receptor ligand in aqueous solution and under conditions otherwise suitable for

binding of the ligand to C5a receptor. The C5a receptor may be present in suspension (*e.g.*, in an isolated membrane or cell preparation), in a cultured or isolated cell, or in a tissue or organ.

[0081] Preferably, the amount of C5a receptor modulator contacted with the receptor should be sufficient to inhibit C5a binding to C5a receptor *in vitro* as measured, for example, using a radioligand binding assay, calcium mobilization assay, or chemotaxis assay as described herein.

[0082] In one embodiment of the invention, the C5a modulators of the invention are used to modulate, preferably inhibit, the signal-transducing activity of a C5a receptor, for example, by contacting one or more compound(s) of the invention with a C5a receptor (either *in vitro* or *in vivo*) under conditions suitable for binding of the modulator(s) to the receptor. The receptor may be present in solution or suspension, in a cultured or isolated cell preparation or within a patient. Any modulation of the signal transducing activity may be assessed by detecting an effect on calcium ion calcium mobilization or by detecting an effect on C5a receptor-mediated cellular chemotaxis. In general, an effective amount of C5a modulator(s) is an amount sufficient to modulate C5a receptor signal transducing activity *in vitro* within a calcium mobilization assay or C5a receptor-mediated cellular chemotaxis within a migration assay.

[0083] When compounds of the invention are used to inhibit C5a receptor-mediated cellular chemotaxis, preferably leukocyte (*e.g.*, neutrophil) chemotaxis, in an *in vitro* chemotaxis assay, such methods comprise contacting white blood cells (particularly primate white blood cells, especially human white blood cells) with one or more compounds of the invention. Preferably the concentration is sufficient to inhibit chemotaxis of white blood cells in an *in vitro* chemotaxis assay, so that the levels of chemotaxis observed in a control assay are significantly higher, as described above, than the levels observed in an assay to which a compound of the invention has been added.

[0084] In another embodiment, the compounds of the present invention further can be used for treating patients suffering from conditions that are responsive to C5a receptor modulation. As used herein, the term "treating" or "treatment" encompasses both disease-modifying treatment and symptomatic treatment, either of which may be prophylactic (*i.e.*, before the onset of symptoms, in order to prevent, delay or reduce the severity of symptoms) or therapeutic (*i.e.*, after the onset of symptoms, in order to reduce the severity and/or duration

of symptoms). As used herein, a condition is considered "responsive to C5a receptor modulation" if modulation of C5a receptor activity results in the reduction of inappropriate activity of a C5a receptor. As used herein, the term "patients" include primates (especially humans), domesticated companion animals (such as dogs, cats, horses, and the like) and livestock (such as cattle, pigs, sheep, and the like), with dosages as described herein.

Conditions that can be treated by C5a modulation:

[0085] Autoimmune disorders-- e.g., Rheumatoid arthritis, systemic lupus erythematosus, Guillain-Barre syndrome, pancreatitis, lupus nephritis, lupus glomerulonephritis, psoriasis, Crohn's disease, vasculitis, irritable bowel syndrome, dermatomyositis, multiple sclerosis, bronchial asthma, pemphigus, pemphigoid, scleroderma, myasthenia gravis, autoimmune hemolytic and thrombocytopenic states, Goodpasture's syndrome (and associated glomerulonephritis and pulmonary hemorrhage), immunovascularitis, tissue graft rejection, hyperacute rejection of transplanted organs; and the like.

[0086] Inflammatory disorders and related conditions-- e.g., Neutropenia, sepsis, septic shock, Alzheimer's disease, multiple sclerosis, stroke, inflammatory bowel disease (IBD), inflammation associated with severe burns, lung injury, and ischemia-reperfusion injury, osteoarthritis, as well as acute (adult) respiratory distress syndrome (ARDS), chronic pulmonary obstructive disorder (COPD), systemic inflammatory response syndrome (SIRS), atopic dermatitis, psoriasis, chronic urticaria and multiple organ dysfunction syndrome (MODS). Also included are pathologic sequelae associated with insulin-dependent diabetes mellitus (including diabetic retinopathy), lupus nephropathy, Heyman nephritis, membranous nephritis and other forms of glomerulonephritis, contact sensitivity responses, and inflammation resulting from contact of blood with artificial surfaces that can cause complement activation, as occurs, for example, during extracorporeal circulation of blood (e.g., during hemodialysis or via a heart-lung machine, for example, in association with vascular surgery such as coronary artery bypass grafting or heart valve replacement), or in association with contact with other artificial vessel or container surfaces (e.g., ventricular assist devices, artificial heart machines, transfusion tubing, blood storage bags, plasmapheresis, plateletpheresis, and the like). Also included are diseases related to ischemia/reperfusion injury, such as those resulting from transplants, including solid organ transplant, and syndromes such as ischemic reperfusion injury, ischemic colitis and cardiac

ischemia. Compounds of the instant invention may also be useful in the treatment of age-related macular degeneration (Hageman et al, *P.N.A.S.***102**: 7227-7232, 2005).

[0087] Cardiovascular and Cerebrovascular Disorders--e.g., myocardial infarction, coronary thrombosis, vascular occlusion, post-surgical vascular reocclusion, atherosclerosis, traumatic central nervous system injury, and ischemic heart disease. In one embodiment, an effective amount of a compound of the invention may be administered to a patient at risk for myocardial infarction or thrombosis (*i.e.*, a patient who has one or more recognized risk factor for myocardial infarction or thrombosis, such as, but not limited to, obesity, smoking, high blood pressure, hypercholesterolemia, previous or genetic history of myocardial infarction or thrombosis) in order to reduce the risk of myocardial infarction or thrombosis.

[0088] Diseases of Vasculitis – Vasculitic diseases are characterized by inflammation of the vessels. Infiltration of leukocytes leads to destruction of the vessel walls, and the complement pathway is believed to play a major role in initiating leukocyte migration as well as the resultant damage manifested at the site of inflammation (Vasculitis, Second Edition, Edited by Ball and Bridges, Oxford University Press, pp 47-53, 2008). The compounds provided in the present invention can be used to treat leukoclastic vasculitis, Wegener's granulomatosis, microscopic polyangiitis, Churg-Strauss syndrome, Henoch-Schonlein purpura, polyarteritis nodosa, Rapidly Progressive Glomerulonephritis (RPGN), cryoglobulinaemia, giant cell arteritis (GCA), Behcet's disease and Takayasu's arteritis (TAK).

[0089] HIV infection and AIDS -- C5a receptor modulators provided herein may be used to inhibit HIV infection, delay AIDS progression or decrease the severity of symptoms or HIV infection and AIDS.

[0090] Neurodegenerative disorders and related diseases-- Within further aspects, C5a antagonists provided herein may be used to treat Alzheimer's disease, multiple sclerosis, and cognitive function decline associated with cardiopulmonary bypass surgery and related procedures.

[0091] In one embodiment of the invention, the compounds of the invention can be used for the treatment of diseases selected from the group consisting of sepsis (and associated disorders), COPD, rheumatoid arthritis, lupus nephritis and multiple sclerosis.

[0092] Treatment methods provided herein include, in general, administration to a patient an effective amount of one or more compounds provided herein. Suitable patients include those patients suffering from or susceptible to (*i.e.*, prophylactic treatment) a disorder or disease identified herein. Typical patients for treatment as described herein include mammals, particularly primates, especially humans. Other suitable patients include domesticated companion animals such as a dog, cat, horse, and the like, or a livestock animal such as cattle, pig, sheep and the like.

[0093] In general, treatment methods provided herein comprise administering to a patient an effective amount of a compound one or more compounds provided herein. In a preferred embodiment, the compound(s) of the invention are preferably administered to a patient (*e.g.*, a human) orally or topically. The effective amount may be an amount sufficient to modulate C5a receptor activity and/or an amount sufficient to reduce or alleviate the symptoms presented by the patient. Preferably, the amount administered is sufficient to yield a plasma concentration of the compound (or its active metabolite, if the compound is a pro-drug) high enough to detectably inhibit white blood cell (*e.g.*, neutrophil) chemotaxis *in vitro*.

Treatment regimens may vary depending on the compound used and the particular condition to be treated; for treatment of most disorders, a frequency of administration of 4 times daily or less is preferred. In general, a dosage regimen of 2 times daily is more preferred, with once a day dosing particularly preferred. It will be understood, however, that the specific dose level and treatment regimen for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination (*i.e.*, other drugs being administered to the patient) and the severity of the particular disease undergoing therapy, as well as the judgment of the prescribing medical practitioner. In general, the use of the minimum dose sufficient to provide effective therapy is preferred. Patients may generally be monitored for therapeutic effectiveness using medical or veterinary criteria suitable for the condition being treated or prevented.

[0094] Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment or preventions of conditions involving pathogenic C5a activity (about 0.5 mg to about 7 g per human patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about

500 mg of an active ingredient. For compounds administered orally, transdermally, intravenously, or subcutaneously, it is preferred that sufficient amount of the compound be administered to achieve a serum concentration of 5 ng (nanograms)/mL-10 µg (micrograms)/mL serum, more preferably sufficient compound to achieve a serum

concentration of 20 ng-1 µg/ml serum should be administered, most preferably sufficient compound to achieve a serum concentration of 50 ng/ml-200 ng/ml serum should be administered. For direct injection into the synovium (for the treatment of arthritis) sufficient compounds should be administered to achieve a local concentration of approximately 1 micromolar.

[0095] Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most disorders, a dosage regimen of 4 times daily, three times daily, or less is preferred, with a dosage regimen of once daily or 2 times daily being particularly preferred. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination (*i.e.*, other drugs being administered to the patient), the severity of the particular disease undergoing therapy, and other factors, including the judgment of the prescribing medical practitioner.

[0096] In another aspect of the invention, the compounds of the invention can be used in a variety of non-pharmaceutical *in vitro* and *in vivo* application. For example, the compounds of the invention may be labeled and used as probes for the detection and localization of C5a receptor (cell preparations or tissue sections samples). The compounds of the invention may also be used as positive controls in assays for C5a receptor activity, *i.e.*, as standards for determining the ability of a candidate agent to bind to C5a receptor, or as radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT). Such methods can be used to characterize C5a receptors in living subjects. For example, a C5a receptor modulator may be labeled using any of a variety of well known techniques (*e.g.*, radiolabeled with a radionuclide such as tritium), and incubated with a sample for a suitable incubation time (*e.g.*, determined by first assaying a time course of binding). Following incubation, unbound compound is removed (*e.g.*, by washing), and bound compound detected using any method suitable for the label employed (*e.g.*, autoradiography or scintillation counting for radiolabeled compounds; spectroscopic methods may be used to detect luminescent groups and fluorescent groups). As a control, a matched

sample containing labeled compound and a greater (*e.g.*, 10-fold greater) amount of unlabeled compound may be processed in the same manner. A greater amount of detectable label remaining in the test sample than in the control indicates the presence of C5a receptor in the sample. Detection assays, including receptor autoradiography (receptor mapping) of C5a receptor in cultured cells or tissue samples may be performed as described by Kuhar in sections 8.1.1 to 8.1.9 of *Current Protocols in Pharmacology* (1998) John Wiley & Sons, New York.

[0097] The compounds provided herein may also be used within a variety of well known cell separation methods. For example, modulators may be linked to the interior surface of a tissue culture plate or other support, for use as affinity ligands for immobilizing and thereby isolating, C5a receptors (*e.g.*, isolating receptor-expressing cells) *in vitro*. In one preferred application, a modulator linked to a fluorescent marker, such as fluorescein, is contacted with the cells, which are then analyzed (or isolated) by fluorescence activated cell sorting (FACS).

[0098] In Figure 1, structures and activity are provided for representative compounds described herein. Activity is provided as follows for the binding assay as described herein: +, $500 \text{ nM} \leq \text{IC}_{50} < 2000 \text{ nM}$; ++, $50 \text{ nM} \leq \text{IC}_{50} < 500 \text{ nM}$; +++, $5 \text{ nM} \leq \text{IC}_{50} < 50 \text{ nM}$; and +++, $\text{IC}_{50} < 5 \text{ nM}$.

V. Examples

[0099] The following examples are offered to illustrate, but not to limit the claimed invention.

[0100] Reagents and solvents used below can be obtained from commercial sources such as Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). ^1H -NMR spectra were recorded on a Varian Mercury 400 MHz NMR spectrometer. Significant peaks are provided relative to TMS and are tabulated in the order: multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet) and number of protons. Mass spectrometry results are reported as the ratio of mass over charge, followed by the relative abundance of each ion (in parenthesis). In the examples, a single m/e value is reported for the $\text{M}+\text{H}$ (or, as noted, $\text{M}-\text{H}$) ion containing the most common atomic isotopes. Isotope patterns correspond to the expected formula in all

cases. Electrospray ionization (ESI) mass spectrometry analysis was conducted on a Hewlett-Packard MSD electrospray mass spectrometer using the HP1100 HPLC for sample delivery. Normally the analyte was dissolved in methanol at 0.1 mg/mL and 1 microlitre was infused with the delivery solvent into the mass spectrometer, which scanned from 100 to 1500 daltons. All compounds could be analyzed in the positive ESI mode, using acetonitrile / water with 1% formic acid as the delivery solvent. The compounds provided below could also be analyzed in the negative ESI mode, using 2 mM NH₄OAc in acetonitrile / water as delivery system.

[0101] The following abbreviations are used in the Examples and throughout the description of the invention:

EtOH: Ethanol

EtONa: Sodium ethoxide

THF: Tetrahydrofuran

TLC: Thin layer chromatography

MeOH: Methanol


[0102] Compounds within the scope of this invention can be synthesized as described below, using a variety of reactions known to the skilled artisan. One skilled in the art will also recognize that alternative methods may be employed to synthesize the target compounds of this invention, and that the approaches described within the body of this document are not exhaustive, but do provide broadly applicable and practical routes to compounds of interest.

[0103] Certain molecules claimed in this patent can exist in different enantiomeric and diastereomeric forms and all such variants of these compounds are claimed.

[0104] The detailed description of the experimental procedures used to synthesize key compounds in this text lead to molecules that are described by the physical data identifying them as well as by the structural depictions associated with them.

[0105] Those skilled in the art will also recognize that during standard work up procedures in organic chemistry, acids and bases are frequently used. Salts of the parent compounds are sometimes produced, if they possess the necessary intrinsic acidity or basicity, during the experimental procedures described within this patent.

Example 1

Synthesis of *cis*-1-(2-fluoro-6-methylbenzoyl)-2-phenylpiperidine-3-carboxylic acid (3-trifluoromethylphenyl)amide

[0106] a) $\text{Pd(PPh}_3)_4$ (3.0 g, 2.6 mmol) was added to a solution of 2-chloro-3-carboxyethylpyridine (25 g, 134.7 mmol), phenylboronic acid (21.04 g, 172.6 mmol) and K_2CO_3 (55.1 g, 399 mmol) in 1,4-dioxane (200 mL) and water (200 mL). The reaction mixture was heated at 100°C for 2 h. The solution was then cooled to room temperature and the dioxane was removed under reduced pressure. The resulting aqueous layer was extracted with ethyl acetate, and the combined organic layers were dried (Na_2SO_4), filtered through celite, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2 , 10-100% EtOAc/hexanes) to get the 2-phenylpyridine derivative in 91% yield (27.98 g). LC-MS R_t (retention time): 2.45 min, MS: (ES) m/z 228 ($\text{M}+\text{H}^+$).

[0107] b) PtO_2 (800 mg, 3.52 mmol) was added to a solution of 2-phenyl-nicotinic acid ethyl ester (20 g, 88 mmol, prepared in step a above) in EtOH (60 mL) and concentrated HCl (15 mL). The reaction mixture was hydrogenated using a Parr shaker at 40-45 psi, for 1 h. The reaction mixture was then filtered through celite, washed with EtOH, and the filtrate was

concentrated under reduced pressure. The residue was diluted with CH_2Cl_2 and washed with saturated NaHCO_3 . Purification by flash chromatography (SiO_2 , 0-20% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) gave the desired product in 85% yield (17.4 g). LC-MS R_t (retention time): 1.73 min, MS: (ES) m/z 234 ($\text{M}+\text{H}^+$).

- 5 [0108] c) Oxalyl chloride (3.2 mL, 30.75 mmol) was added to the solution of 2-fluoro-6-methylbenzoic acid (3.79 g, 24.6 mmol) in CH_2Cl_2 (20 mL) in a reaction flask at room temperature, followed by addition of a catalytic amount of DMF. The reaction was kept stirring for 2 h at room temperature. Solvent and excess oxalyl chloride were removed *in vacuo* and the residue was dried under high vacuum for 20 min. The resulting acid chloride
- 10 was dissolved in dry CH_2Cl_2 (20 mL) and cooled to 0 °C followed by the addition of the piperidine made in step b (5.56 g, 20.5 mmol) and Et_3N (8.6 mL, 61.5 mmol). The mixture was then allowed to warm to room temperature and stirred overnight. The reaction mixture was diluted with CH_2Cl_2 and water was added. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were dried (MgSO_4) and
- 15 concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2 , 10-35% $\text{EtOAc}/\text{hexanes}$) to give 7.47 g of the desired compound 99% yield). LC-MS R_t (retention time): 2.50 min and 2.58 min (two rotamers), MS: (ES) m/z 370 ($\text{M}+\text{H}^+$).

- [0109] d) Lithium aluminum hydride solution (2.0 M in THF, 8.2 mL, 16.4 mmol) was added to a solution of the ester from step c (2.98 g, 8.06 mmol) in THF (100 mL) at 0 °C. The
- 20 resulting solution was kept stirring at 0 °C for 2 h at which time the reaction was completed. 15% Aqueous NaOH (625 μL) was added drop wise to quench the reaction followed by H_2O (625 μL). To the cloudy colloidal mixture was added additional water (1.85 mL), and the mixture was kept stirring for 1 h at rt. The mixture was then filtered through a celite plug, and the filtrate was concentrated under reduced pressure. Purification by flash
- 25 chromatography (SiO_2 , 33-67% $\text{EtOAc}/\text{hexanes}$) gave 2.46 g of the desired product (93% yield). LC-MS: R_t (retention time): 1.90 min and 2.09 min (two rotamers), MS: (ES) m/z 328 ($\text{M}+\text{H}^+$).

- [0110] e) A solution of the alcohol from step d (1.42 g, 4.33 mmol) in acetic acid (65 mL) was added to a slurry of CrO_3 (2.61 g, 26.1 mmol) in H_2O (16 mL) at room temperature. The
- 30 resulting mixture was kept stirring at room temperature until the reaction was completed (90 min). The mixture was filtered through a Celite plug and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2 , 3-10% $\text{CH}_2\text{Cl}_2:\text{MeOH}$

followed by 50-67% EtOAc/hexanes) gave 1.03 g of the desired product (70% yield). LC-MS: R_t (retention time): 1.88 min and 2.12 min (two rotamers), MS: (ES) m/z 342 ($M+H^+$).


[0111] f) 3-Trifluoromethylaniline (16.2 mg, 0.1 mmol, 1.0 eq) was added to a solution of the acid prepared above (34.2 mg, 0.1 mmol) and triethylamine (6 eq) in CH_2Cl_2 (1 mL).

- 5 T3P (95.5 mg, 0.15 mmol) was then slowly added and the solution was allowed to stir at room temperature for 1.5 h. The reaction mixture was diluted with CH_2Cl_2 (1 mL), washed with 1 N aqueous HCl followed by saturated aqueous $NaHCO_3$. The organic layer was separated, dried over anhydrous $MgSO_4$, and concentrated under reduced pressure. Purification by flash chromatography (SiO_2 , 5-40% EtOAc/hexanes) gave 35 mg (73% yield) of the product as a white solid. 1H NMR (400 MHz, $CDCl_3$) δ 1.22-2.45 (m, 8 H), 2.93-3.32 (m, 3 H), 6.77-7.82 (m, 12 H), 9.10 (s, 0.38 H), 9.30 (s, 0.62 H). LC-MS: R_t (retention time) = 2.88 min, MS: (ES) m/z 485 ($M+H^+$).

Example 2

15

Synthesis of *N*-(3-*tert*-butylphenyl)-1-(5-chloro-3-methylpicolinoyl)-2-phenylpiperidine-3-carboxamide



[0112] a) 2-Chloronicotinoyl chloride (1.05 eq) dissolved in anhydrous dichloromethane (0.5 M) was added to a solution of 3-*tert*-butylaniline (1 eq) and 2 M aq K₂CO₃ (2.2 eq) in anhydrous dichloromethane (0.5 M) at 0 °C over a period of 30 min, and the reaction mixture was allowed to stir at room temperature for an additional 1.5 h. The layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layer was washed with brine, dried (MgSO₄), filtered and concentrated to give the desired amide as a foamy solid which was used as such in the next step without further purification. MS: (ES) *m/z* 289.1 (M+ H⁺).

[0113] b) Pd(PPh₃)₄ (2-5 mol%) was added to a solution of the above pyridine amide (1 eq), phenylboronic acid (1.4 eq) and 2 M aq K₂CO₃ (2.4 eq) in toluene (0.7 M) and the reaction mixture was heated at 100 °C over night (~12 h). After cooling to room temperature, the reaction mixture was filtered through celite and the celite plug was washed with EtOAc. The filtrate was diluted with water and extracted with EtOAc, dried (MgSO₄), filtered and concentrated and concentrated under reduced pressure. The residue was purified by automated flash chromatography (SiO₂, 10% to 100% gradient of EtOAc-hexanes) and dried *in vacuo* to give the 2-phenyl-3-carboxyamidopyridine in 60-75% yield, MS: (ES) *m/z* 331.2 (M+ H⁺).


[0114] c) PtO₂ (10 mol%) was added to a solution of the 2-phenylpyridine derivative prepares above (1 eq) in EtOH and concentrated HCl (excess, 4:1 ratio) and the reaction mixture was hydrogenated using a Parr shaker at 40-45 psi, for 1.5 h. It was filtered through celite, washed with EtOH, and the filtrate was concentrated. The residue was diluted with CH₂Cl₂ and washed with saturated aq NaHCO₃. The residue was then purified by automated flash chromatography (SiO₂, 1% to 30% gradient of CH₂Cl₂-MeOH) and dried *in vacuo* to give the title compound in ~85% yield as a foamy solid. MS: (ES) *m/z* 337.2 (M+ H⁺).

[0115] d) 5-Chloro-3-methylpicolinic acid (30 mg, 0.16 mmol) and *N*-(3-*tert*-butylphenyl)-2-phenylpiperidine-3-carboxamide (50 mg, 0.15mmol, prepared in step c above) were dissolved in anhydrous DMF (1 mL). *N,N*-Diisopropylethylamine (0.15 mL) was added at room temperature followed by HCTU (67 mg, 0.16 mmol). After stirring 2 h at ambient temperature, LC-MS and TLC indicated the completion of the reaction. The reaction mixture was diluted with EtOAc (50 mL) and washed with 1 N HCl (20 mL), saturated NaHCO₃ (30 mL), and brine (30 mL) and the resulting solution was concentrated under reduced pressure. The residue was purified by preparative HPLC (20 → 95% gradient of MeCN-H₂O with

0.1% TFA) and the pure fractions were lyophilized to afford the title compound (50 mg, 67% yield). HPLC retention time = 2.88 minutes. ^1H NMR (400 MHz, CDCl_3) δ 8.42 (d, 1 H, J = 0.8 Hz), 7.97 (br, 1 H), 7.59 (d, 1 H, J = 0.8 Hz), 7.56 (d, 1 H, J = 7.6 Hz), 7.34 (m, 3 H), 7.20 (m, 3 H), 7.10 (d, 1 H, J = 7.6 Hz), 6.61 (two sets of br, 1 H), 3.12 (two sets of m, 2 H), 2.94 (three sets of m, 1 H), 2.36 (s, 3 H), 2.20 (two sets of br, 2 H), 1.74 (br complex, 2 H), 1.29 (s, 9 H). MS: (ES) m/z 490.2 ($\text{M} + \text{H}^+$).

Example 3

10 Synthesis of *cis*-1-(2-methylbenzoyl)-2-(3-fluorophenyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide



15 [0116] a) To a mixture of *N*-(3-*tert*-butylphenyl)-2-chloronicotinamide (570.2 mg, 2 mmol), 3-fluorophenylboronic acid (401.2 mg, 2.8 mmol), 3 mL of toluene, and 1 mL of 2 N potassium carbonate in water was added tetrakis(triphenylphosphine)palladium(0) (234.5 mg, 0.2 mmol). The mixture was then heated at 90 °C for 3 hour under nitrogen, before it was cooled down to room temperature. The reaction mixture was then diluted with 30 mL of water and 150 mL of EtOAc. The organic layer was separated, washed with brine, and dried (Na₂SO₄). The organic solvent was removed under reduced pressure and the residue was purified by silica gel column (40% EtOAc in hexane) to give *N*-(3-*tert*-butylphenyl)-2-(3-fluorophenyl)nicotinamide (691.4 mg, 99%). MS: (ES) m/z 394.5 ($\text{M} + \text{H}^+$).


25 [0117] b) A mixture of *N*-(3-*tert*-butylphenyl)-2-(3-fluorophenyl)nicotinamide (501.2 mg, 1.4 mmol), platinum oxide (51.9 mg, 0.21 mmol), and concentrated HCl (400 μL , 5.2 mmol) in 5 mL of ethanol was stirred vigorously under hydrogen balloon overnight. The mixture

was filtered, and the solids washed with 25 mL of methanol three times. The combined solution was dried under reduced pressure. To the residue was added 30 mL of saturated sodium bicarbonate and 150 mL of EtOAc. The organic layer was separated, and dried over sodium sulfate. Evaporation of solvent gave the crude 2-(3-fluorophenyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide as a brown solid, which was taken on directly to the next step. MS: (ES) m/z 355.7 ($M+H^+$).

[0118] c) To a solution of 2-(3-fluorophenyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide (prepared above, 177.3 mg, 0.5 mmol) in 2 mL of dichloromethane was added Et₃N (100 μ L, excess), and 2-methylbenzoyl chloride (92.3 mg, 0.6 mmol) at room temperature. The resulting solution was then stirred at this temperature until completion of the reaction (10 min.). The reaction mixture was then directly loaded onto a silica gel column, and was purified by using ISCO (30% EtOAc in hexane) to give the final product 2-(3-fluorophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide (151.2mg, 64% yield). ¹H NMR (400 MHz, CDCl₃, mixture of rotomers): δ 7.91 (s, 0.6 H), 7.85 (s, 0.4 H), 7.18-7.46 (m, 9 H), 7.11 (m, 1 H), 6.95 (m, 1 H), 6.67 (d, J = 1.2 Hz, 1 H), 3.36 (d, J = 1.6 Hz, 0.4 H), 3.26 (d, J = 1.6 Hz, 1 H), 3.05 (m, 1 H), 2.89 (t, J = 1.2 Hz, 1 H), 2.45 (s, 1 H), 2.02-2.40 (m, 4 H), 1.70-1.84 (m, 3 H), 1.44-1.64 (s, 1 H), 1.32 (s, 6 H), 1.25 (s, 1 H). MS: (ES) m/z 473.2 ($M+H^+$).

Example 4

Synthesis of *cis*-1-(2-methylbenzoyl)-2-(2,2-dimethylpropyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide



[0119] a) To a stirred solution of 2-bromonicotinic acid (1.01g, 5 mmol) dissolved in anhydrous dichloromethane (8 mL) were added EDCI (1.34 g, 7 mmol) and 3-*tert*-butylaniline (0.74 g, 5 mmol) at room temperature and the reaction mixture was stirred for 12 hours. The mixture was then diluted with dichloromethane, followed by saturated sodium bicarbonate and water wash. The dichloromethane layer was dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography to obtain 2-bromo-*N*-(3-*tert*-butylphenyl)nicotinamide in 59% yield (950 mg). Rt: 2.44 min (20-100-5 method). MS: (ES) *m/z* 333, 335 (*M*+*H*⁺).

[0120] b) 2, 2-Dimethylpropylmagnesium chloride (1 M-diethylether, 4.8 mL, 4.8 mmol) was added to a suspension of copper cyanide (215 mg, 2.40 mmol) in THF (6 mL) at -78°C. After stirring at the same temperature for 1 hour, 2-bromo-*N*-(3-*tert*-butylphenyl)nicotinamide (200 mg, 0.601 mmol) was added all at once as a solid. The reaction mixture was gradually warmed to room temperature and the reaction was allowed to stir overnight. Saturated ammonium chloride solution and ethyl acetate was added, and the reaction mixture was filtered through celite and rinsed with ethyl acetate. The layers were separated and the product was extracted once more with ethyl acetate. The combined organic layers were washed with brine and dried over anhydrous sodium sulfate. After removing the


solvent under reduced pressure, the crude material was purified using silica gel column chromatography using a gradient of 20% - 50% ethyl acetate in hexanes to yield *N*-(3-*tert*-butylphenyl)-2-(2, 2-dimethylpropyl)nicotinamide (168 mg, 0.517 mmol, 86%). R_f = 0.45 (toluene: ethyl acetate = 2:1).

5 [0121] c) *N*-(3-*tert*-Butylphenyl)-2-(2, 2-dimethylpropyl)nicotinamide (168 mg, 0.517 mmol) was dissolved in ethanol (5 mL). Platinum oxide (11.6 mg, 0.0511 mmol) was added followed by concentrated hydrochloric acid (250 μ L). The reaction mixture was hydrogenated using a Parr apparatus for 1.5 hours at 45 psi. Analysis of the reaction mixture showed incomplete conversion, and the sequence was repeated one more time. Platinum
10 oxide was filtered off and the solvents were removed under reduced pressure. The crude material was neutralized using saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was then washed with brine and dried over anhydrous magnesium sulfate. Removal of solvent under reduced pressure gave the crude 2,3-*cis*-2-(2,2-dimethylpropyl)piperidine-3-carboxylic acid-(3-*tert*-butylphenyl)amide (153 mg) which was
15 used in the next step without further purification.

[0122] d) To a solution of 2,3-*cis*-2-(2,2-dimethylpropyl)piperidine-3-carboxylic acid-(3-*tert*-butylphenyl)amide (84.8 mg, 0.257 mmol) in pyridine (415 μ L, 5.13 mmol) at room temperature was added 2-methylbenzoyl chloride (81.6 mg, 0.528 mmol) in chloroform (415 μ L). A catalytic amount (not weighed) of dimethylaminopyridine was added to enhance the
20 reaction and the mixture was stirred for three days. Ethyl acetate and water was then added to the reaction mixture and the product was extracted with ethyl acetate three times. The combined organic layers were dried over anhydrous magnesium sulfate. After removal of the solvent under reduced pressure the crude material was purified via silica gel chromatography using 10% - 20% ethyl acetate in hexanes to give 2,3-*cis*-2-(2,2-dimethylpropyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid-(3-*tert*-butylphenyl)amide (47.0 mg, 0.105
25 mmol, 41%). R_f = 0.6 (hexanes: ethyl acetate = 2:1). R_t = 3.16 min., 3.26 min. (compound exists as mixtures of several conformers. 20-100-5 method.). ^1H NMR (CDCl_3) δ 9.68 (s, 1 H), 9.43 (s, 1 H), 8.33 (s, 1 H), 8.28 (s, 1 H), 6.97-7.79 (m, 8 H), 5.48 (br, 1 H), 5.39 (dd, J = 4, 10 Hz, 1 H), 5.33 (dd, J = 6, 6 Hz, 1 H), 3.38 (ddd, J = 4, 14, 14 Hz, 2 H), 3.25 (dd, J = 13, 13 Hz, 2H), 2.66 (dd, J = 4, 8.4 Hz, 1 H), 2.63 (ddd, J = 2.8, 2.8, 8 Hz, 1 H), 2.50 (s, 9 H),
30 2.40 (s, 9 H), 2.25 (s, 9 H), 2.13 (s, 9 H), 1.79-1.99 (m, 2 H), 1.23-1.56 (m, 2 H), 1.32 (s, 9 H), 1.07 (s, 9 H), 1.06 (s, 9 H), 0.97 (s, 9 H), 0.95 (s, 9 H). MS: (ES) m/z 449 ($\text{M}+\text{H}^+$).

Example 5

5 **Synthesis of *cis*-2-cyclopentyl-1-(2-methylbenzoyl)piperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide**



[0123] a) Cyclopentylzinc bromide (0.5 M, 6.5 mL, 3.26 mmol) was added to a room temperature stirred solution of the 2-chloronicotinic acid methyl ester (400 mg, 2.33 mmol), CuI (19 mg, 0.1 mmol) and Pd(dppf)Cl_2 (42 mg, 0.06 mmol) in anhydrous dimethylacetamide (1.7 mL) under nitrogen. The reaction mixture was heated to 70°C for 3.5 hours, cooled to room temperature, filtered through celite, and the cake was rinsed with ethyl acetate. The filtrate was washed with water, brine, dried (MgSO_4), filtered and concentrated under reduced. The residue was purified by flash chromatography (SiO_2 , 10-100% EtOAc/hexanes) to get the desired compound in 83% yield (400 mg). LC-MS R_t (retention time): 1.87 min; MS: (ES) m/z 206 ($\text{M}+\text{H}^+$).

[0124] b) $n\text{-BuLi}$ (1.47 mL, 3.68 mmol) was added to the 3-*tert*-butylaniline (580 mg, 3.89 mmol) at -78°C in dry THF (2 mL) under nitrogen and the solution was allowed to stir at 0°C for 10 minutes. The reaction mixture was re-cooled to -78°C and 2-cyclopentyl-nicotinic acid methyl ester (400 mg, 1.94 mmol) dissolved in dry THF (2 mL) was added to it. The reaction mixture was allowed to attain 0°C over a period of 2 hours, quenched with saturated aqueous NH_4Cl , and extracted with ethyl acetate. The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. The residue was purified by


flash chromatography (SiO₂, 10-100% EtOAc/hexanes) to give the pure compound in 91% yield (572 mg). LC-MS R_t(retention time): 2.61 min; MS: (ES) *m/z* 323 (M+H⁺).

[0125] c) To a solution of the *N*-(3-*tert*-butylphenyl)-2-cyclopentylnicotinamide (570 mg, 1.77 mmol) in ethanol (10 mL) containing concentrated HCl (1 mL) was added platinum oxide (40 mg, 0.17 mmol) and the solution was hydrogenated using a Parr shaker at 40 psi for 1.5 hour. The reaction mixture was filtered through Celite, and the cake was rinsed with ethanol. The filtrate was concentrated, and the residue was dried under high vacuum for 2 hours to get quantitative yield of the desired piperidine as a HCl salt. LC-MS R_t(retention time): 1.97 min; MS: (ES) *m/z* 329 (M+H⁺).

[0126] d) To a solution of the *cis*-2-cyclopentylpiperidine-3-carboxylic acid (3-*tert*-butylphenyl)amide prepared above (123 mg, 0.34 mmol) in dry CH₂Cl₂ (1 mL) containing Et₃N (142 μL, 1.02 mmol) was added 2-methylbenzoyl chloride (53 mg, 0.34 mmol) and the mixture was stirred at room temperature for 2 hours. The reaction mixture was then diluted with ethyl acetate (20 mL), washed with 1 N aqueous HCl, water, and brine. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure. The residue was purified by reverse phase preparative HPLC (20-95% gradient of CH₃CN-H₂O) and dried (Lyophilizer) to give the title compound in 65% yield (109 mg). ¹H NMR (400 MHz, CDCl₃): δ 1.22-1.48 (m, 11 H), 1.56-1.80 (m, 5 H), 1.84-2.06 (m, 4 H), 2.10-2.23 (m, 1 H), 2.30 (s, 1.6 H), 2.39 (s, 1.4 H), 2.41-2.50 (m, 1 H), 2.71-2.76 (m, 1 H), 3.02-3.09 (m, 1 H), 3.25-3.39 (m, 1 H), 5.11 (bs, 1 H), 7.05-7.30 (m, 6 H), 7.47-7.55 (m, 2 H), 8.32 (bs, 1 H). LC-MS R_t(retention time): 3.16 min; MS: (ES) *m/z* 447 (M+H⁺). LC-MS method: Agilent Zorbax SB-C18, 2.1× 50 mm, 5μ, 35 °C, 1 mL/min flow rate, a 2.5 min gradient of 20% to 100% B with a 1.0 min wash at 100% B; A= 0.1% formic acid/ 5% acetonitrile / 94.9 water, B = 0.1% formic acid / 5% water/ 94.9 acetonitrile.

Example 6

Synthesis of (2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid(3-chloro-4-methylphenyl)amide



[0127] b) *cis*-2-(4-tert-Butoxycarbonylamino)phenyl)piperidine-3-carboxylic acid ethyl ester was synthesized similarly as illustrated in example 1.

[0128] c:1) : *cis*-2-(4-tert-Butoxycarbonylamino)phenyl)piperidine-3-carboxylic acid ethyl ester (61 g, 174.8 mmol) and di-p-toluoyl-L-tartaric acid (62 g, 174.8 mmol) was dissolved in EtOH (500 ml). The clear solution was concentrated and pumped to dry. The obtained white salt was then dissolved into 250 ml of ethyl acetate to form a clear solution. To this solution was added 500 ml of TBME slowly. The obtained solution was left at rt undisturbed for 3 days. At this time a lot of white crystals were formed. They were then filtered and washed with 100 ml of TBME to obtain a white solid (60 g).

[0129] The above salt was re-dissolved in ethanol, concentrated and pumped to dry. The obtained salt was dissolved into 500 ml of THF, followed by adding TBME (500 ml). The obtained clear solution was left at rt undisturbed for another 2.5 days. The obtained white crystals were filtered to obtain 20.5 g (enrichment 64:1) of the salt.

[0130] c:2) To a 0 °C stirred suspension of the salt (16.7 g) in CH₂Cl₂ (150 mL) was added saturated aqueous NaHCO₃ solution (100 mL) and the reaction mixture was allowed to stir at r.t over a period of 30 minutes. The layers were separated and the aqueous layer was
5 extracted with CH₂Cl₂ (50 mL). The combined organic layer was washed with saturated aqueous NaHCO₃ (2 ×100 mL), dried and concentrated to give (2*R*,3*S*)-2-(4-tert-Butoxycarbonylaminophenyl)piperidine-3-carboxylic acid ethyl ester in 90% yield and ~ in 97% ee.

[0131] d) To a 0 °C solution of the (2*R*,3*S*)-2-(4-tert-Butoxycarbonylaminophenyl)-
10 piperidine-3-carboxylic acid ethyl ester prepared above (600 mg, 1.72 mmol) in dry CH₂Cl₂ (5 mL) containing Et₃N (480 µL, 3.44 mmol) was added 2-methylbenzoyl chloride (266 mg, 1.72 mmol) and the mixture was stirred at room temperature for over night. The reaction mixture was then diluted with CH₂Cl₂ (20 mL), washed with 1 N aqueous HCl, water, and brine. The organic layer was dried (MgSO₄), filtered and concentrated under reduced
15 pressure to give (2*R*,3*S*)-2-(4-tert-Butoxycarbonylaminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid ethyl ester in quantitative yield and the crude product was used as such in the next step.

[0132] e) 4N HCl in 1,4-dioxane (5 mL, 20 mmol) was slowly added to a 0 °C solution of the above crude product (2*R*,3*S*)-2-(4-tert-Butoxycarbonylaminophenyl)-1-(2-
20 methylbenzoyl)piperidine-3-carboxylic acid ethyl ester (840 mg, 1.72 mmol) in dry CH₂Cl₂ (4 mL). After the addition of the HCl, the reaction mixture was allowed to attain r.t and stirred for 1 h. It was diluted with CH₂Cl₂ (30 mL), cooled to 0 °C and neutralized with saturated aqueous NaHCO₃ to get the (2*R*,3*S*)-2-(4-aminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid ethyl ester (612 mg) in 97% yield over two steps.

[0133] f) Na(OAc)₃BH (495 mg, 2.33 mmol) was added to a solution of the (2*R*,3*S*)-2-(4-aminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid ethyl ester (612 mg, 1.67 mmol), cyclopentanone (140 mg, 1.67 mmol) and acetic acid (100 mg, 1.67 mmol) in dry dichloroethane at r.t and the reaction mixture was heated to 50 °C for 4 h, cooled to r.t and stirred for 48 h. It was then diluted with CH₂Cl₂ (30 mL), washed with saturated aqueous
30 NaHCO₃ solution, dried and concentrated in *vacuo*. The residue was purified by ISCO flash column using ethyl acetate and hexanes as mobile phase (40 g column, 0-40% gradient) to

afford (2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid ethyl ester (450 mg).


[0134] g) Me₃Al (290 μL, 0.57 mmol, 2M in toluene) was added to a solution of the 3-Chloro-4-methylphenylamine (65 mg, 0.46 mmol) in dry dichloroethane (1 mL) at ambient temperature. Stirred for 20 minutes, then (2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid ethyl ester (100 mg, 0.23mmol) dissolved in dry dichloroethane (1 mL) was added to it. The reaction mixture was then heated to 85 °C for 3 h, cooled to r.t, diluted with CH₂Cl₂ (20 mL), washed with saturated aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂ (20 mL) and the combined organic layer was dried (MgSO₄) and concentrated. The residue was purified by reverse phase preparative HPLC (20-95% gradient of CH₃CN-H₂O with 0.1% TFA as additive), the product containing fractions were pooled together and concentrated. The residue was diluted with CH₂Cl₂ (30 mL), washed with saturated aqueous NaHCO₃ solution. The CH₂Cl₂ layer was dried (MgSO₄) and concentrated to get the pure (2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-methylbenzoyl)piperidine-3-carboxylic acid (3-chloro-4-methylphenyl)amide in 50 % yield.

[0135] ¹H NMR (400 MHz, CDCl₃) δ 8.4(bs, 1H), 7.55 (s, 1H), 7.37-7.05 (m, 9H), 6.55-6.52 (m, 2H), 3.77-3.70(m, 1H), 3.30-3.16 (m, 1H), 3.04-2.91 (m, 2H), 2.43-1.94 (m, 8H), 1.71-1.46 (m, 11H).

Example 7

[0136] The following are representative compounds prepared and evaluated using methods similar to the examples herein. Characterization data is provided for the compounds below. Biological evaluation is shown in Figure 1 for these compounds and others prepared as described herein.


(2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (4-methyl-3-trifluoromethylphenyl)amide



[0137] ^1H NMR (400 MHz, TFA-*d*) δ 7.91 (d, J = 8.6 Hz, 1 H), 7.84 (d, J = 8.6 Hz, 1 H), 7.58-6.82 (m, 8 H), 6.75 (t, J = 8.6 Hz, 1 H), 4.10-4.00 (m, 1H), 3.60-3.47 (m, 1H), 3.45-3.41 (m, 1H), 3.33-3.25 (m, 1H), 2.44-2.22 (m, 7H), 2.04-1.92 (m, 4H), 1.82-.169 (m, 7H)

(2*R*,3*S*)-1-(2-Chlorobenzoyl)-2-(4-cyclopentylaminophenyl)piperidine-3-carboxylic acid (4-methyl-3-trifluoromethylphenyl)amide


10



[0138] ^1H NMR (400 MHz, CDCl_3) δ 9.41 (bs, 0.5H), 9.03 (bs, 0.5H), 7.55 (s, 1H), 7.49-7.39 (m, 3H), 7.31-7.27 (m, 2H), 7.18-7.04 (m, 2H), 6.83-6.74 (m, 3 H), 3.76-3.64 (m, 1H), 3.22-2.90 (m, 5H), 2.39 (s, 3H), 2.32-2.20 (m, 1H), 2.16-2.04 (m, 1H), 2.0-1.86 (m, 2H) 1.80-1.72 (m, 3H), 1.56 (bs, 5H).


(2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (3-chloro-4-methylphenyl)amide

20




[0139] ^1H NMR ($\text{DMSO}-d_6$) δ 10.22 (s, 1H), 7.67 (dd, $J = 1.8$ Hz, $J = 11.0$ Hz, 1H), 7.04-7.33 (m, 9H), 6.30 (dd, $J = 5.8$ Hz, $J = 9.4$ Hz, 1H), 5.52 (br, 1H), 3.56-3.64 (m, 1H), 3.00-3.17 (m, 2H), 2.90-2.98 (m, 1H), 2.23(2.24) (s, 3H), 1.97(2.33) (s, 3H), 1.32-2.22 (m, 12H)

5 **(2*R*,3*S*)-1-(4-Chlorobenzoyl)-2-(4-Cyclopentylaminophenyl)piperidine-3-carboxylic acid (4-methyl-3-trifluoromethylphenyl)amide**




10 [0140] ^1H NMR (400 MHz, CDCl_3) δ 8.79 (bs, 1H), 7.62 (s, 1H), 7.52-7.48 (m, 1H), 7.37-7.30 (m, 5H), 7.13 (d, $J = 8.4$ Hz, 1H), 6.52-6.50 (m, 3H), 3.75-3.69 (m, 1H), 3.44 (bs, 1H), 3.09-2.97 (m, 2H), 2.39 (s, 3H), 2.37-2.30 (m, 1H), 2.13-2.08 (m, 1H), 2.10-1.93 (m, 2H), 1.80-1.59 (m, 7H), 1.48-1.42 (m, 2H)

15 **(2*R*,3*S*)-2-(4-Cyclohexylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (3-*t*-butylphenyl)amide**




20 [0141] ^1H NMR (400 MHz, CDCl_3): δ 8.24 (m, 1H), 7.40-6.85 (m, 8H), 6.65-6.40 (m, 3H), 3.57 (s, 1H), 3.30-2.90 (m, 4H), 2.50-1.85 (m, 9H), 1.80-1.50 (m, 5H), 1.40-1.00 (m, 13H)

25 **(2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (4-methyl-3-pyrrolidin-1-yl-phenyl)amide**




[0142] ^1H NMR (400 MHz, CDCl_3): δ 7.98 (m, 1H), 7.40- 7.18 (m, 3H), 7.10- 6.80 (m, 4H), 6.64- 6.40 (m, 3H), 3.80-3.50 (m, 2H), 3.30-2.90 (m, 6H), 2.50-2.10 (m, 7H), 2.10-1.80 (m, 8H), 1.80-1.20 (m, 9H)

(2*R*,3*S*)-2-[4-(Cyclopentyloxy)phenyl]-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (3-chloro-4-methylphenyl)amide



[0143] ^1H NMR (400 MHz, CDCl_3) δ 8.68 (bs, 0.6H), 8.58 (bs, 0.4H), 7.59-7.40 (m, 3H), 7.29-6.90 (m, 4H), 6.80 (m, 2H), 6.65 (m, 1H), 4.72 (m, 1H), 3.30-2.92 (m, 3H), 2.44 (s, 1H), 2.42-2.30 (m, 1H), 2.30 (s, 1H), 2.29 (s, 2H), 2.20 (s, 2H), 2.19-2.12 (m, 1H), 2.08-1.92 (m, 2H), 1.90-1.72 (m, 7H) 1.60 (m, 2H).


($\pm$)-(2*R*,3*S*)-2-(4-Cyclopentylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (4-chloro-3-methylphenyl)amide



[0144] ^1H NMR (400 MHz, CDCl_3) δ 8.25 (bs, 0.4H), 8.16 (bs, 0.6H), 7.44-7.20 (m, 6H), 7.06-6.84 (m, 2H), 6.59-6.50 (m, 2H), 3.75 (m, 1H), 3.66 (bs, 1H), 3.26-2.92 (m, 3H), 2.43 (s, 1H), 2.42-2.30 (m, 1H), 2.30 (s, 1H), 2.29 (s, 2H), 2.20 (s, 2H), 2.19-2.12 (m, 1H), 2.08-1.92 (m, 2H), 1.80-1.58 (m, 7H) 1.45 (m, 2H).

5


(2*R*,3*S*)-2-(4-Cyclobutylaminophenyl)-1-(2-fluoro-6-methylbenzoyl)piperidine-3-carboxylic acid (3-*t*-butylphenyl)amide



10


[0145] ^1H NMR (400 MHz, CDCl_3) δ 8.20 (s, 0.6 H), 8.39 (s, 0.4 H), 7.44-6.88 (m, 10 H), 6.25 (dd, $J = 12$ Hz, $J = 6$ Hz, 1 H), 6.45 (t, $J = 8.4$ Hz, 1H), 3.87 (m, 1H), 3.26-2.95 (m, 3H), 2.46-2.05 (m, 8H), 1.86-1.61 (m, 5H), 1.34-1.11 (m, 9H)

15 **(2*R*,3*S*)-1-(2-fluoro-6-methylbenzoyl)-2-[4-(tetrahydropyran-4-ylamino)phenyl]piperidine-3-carboxylic acid (3-morpholin-4-yl-phenyl)amide**



20 [0146] ^1H NMR (400 MHz, CDCl_3) δ 7.61 (s, 1 H), 7.34-6.92 (m, 10 H), 6.78-6.65 (m, 1 H), 6.62-6.53 (m, 1H), 3.98-3.85 (m, 4H), 3.83-3.70 (m, 1H), 3.55-3.30 (m, 3H), 3.27-2.98 (M, 4H), 2.42-1.92 (m, 8H), 1.81-1.45 (m, 7H)

25 **(2*R*,3*S*)-1-(2-fluoro-6-methylbenzoyl)-2-[4-((*R*)-2-trifluoromethylpyrrolidin-1-ylmethyl)phenyl]piperidine-3-carboxylic acid (3-*t*-butylphenyl)amide**



[0147] ^1H NMR (400 MHz, CDCl_3) δ 8.01 (bs, 0.5H), 7.96 (bs, 0.5H), 7.55-7.37 (m, 3H), 7.30-7.19 (m, 6H), 7.13-7.06 (m, 1H), 7.01-6.90 (m, 1H), 6.85-6.64 (m, 1H), 4.15-4.11 (m, 1H), 3.58-3.54 (m, 1H), 3.30-3.20 (m, 2H), 3.17-2.80 (m, 2H), 2.45-2.17 (m, 4H), 2.00-1.94 (m, 2H), 1.86-1.60 (m, 8H), 1.31-1.26 (m, 7H)

Example 8

MATERIALS AND METHODS

A. Cells

1. C5a receptor expressing cells

a) *U937 Cells*

[0148] U937 cells are a monocytic cell line which express C5aR, and are available from ATCC (VA). These cells were cultured as a suspension in RPMI-1640 medium supplemented with 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, 1 mM sodium pyruvate, and 10% FBS. Cells were grown under 5% CO_2 /95% air, 100% humidity at 37°C and subcultured twice weekly at 1:6 (cells were cultured at a density range of 1×10^5 to 2×10^6 cells/mL) and harvested at 1×10^6 cells/mL. Prior to assay, cells are treated overnight with 0.5 mM of cyclic AMP (Sigma, OH) and washed once prior to use. cAMP treated U937 cells can be used in C5aR ligand binding and functional assays.

b) *Isolated human neutrophils*

[0149] Optionally, human or murine neutrophils can be used to assay for compound activity. Neutrophils may be isolated from fresh human blood using density separation and centrifugation. Briefly, whole blood is incubated with equal parts 3% dextran and allowed to separate for 45 minutes. After separation, the top layer is layered on top of 15 mls of Ficoll

(15 mls of Ficoll for every 30 mls of blood suspension) and centrifuged for 30 minutes at 400 x g with no brake. The pellet at the bottom of the tube is then isolated and resuspended into PharmLyse RBC Lysis Buffer (BD Biosciences, San Jose, CA) after which the sample is again centrifuged for 10 minutes at 400 x g with brake. The remaining cell pellet is

5 resuspended as appropriate and consists of isolated neutrophils.

B. Assays

1. Inhibition of C5aR ligand binding

[0150] cAMP treated U937 cells expressing C5aR were centrifuged and resuspended in assay buffer (20 mM HEPES pH 7.1, 140 mM NaCl, 1 mM CaCl₂, 5 mM MgCl₂, and with
10 0.1% bovine serum albumin) to a concentration of 3×10^6 cells/mL. Binding assays were set up as follows. 0.1 mL of cells was added to the assay plates containing 5 μ L of the compound, giving a final concentration of ~2-10 μ M each compound for screening (or part of a dose response for compound IC₅₀ determinations). Then 0.1 mL of ¹²⁵I labeled C5a (obtained from Perkin Elmer Life Sciences, Boston, MA) diluted in assay buffer to a final
15 concentration of ~50 pM, yielding ~30,000 cpm per well, was added, the plates sealed and incubated for approximately 3 hours at 4°C on a shaker platform. Reactions were aspirated onto GF/B glass filters pre-soaked in 0.3% polyethyleneimine (PEI) solution, on a vacuum cell harvester (Packard Instruments; Meriden, CT). Scintillation fluid (40 μ L; Microscint 20, Packard Instruments) was added to each well, the plates were sealed and radioactivity
20 measured in a Topcount scintillation counter (Packard Instruments). Control wells containing either diluent only (for total counts) or excess C5a (1 μ g/mL, for non-specific binding) were used to calculate the percent of total inhibition for compound. The computer program Prism from GraphPad, Inc. (San Diego, Ca) was used to calculate IC₅₀ values. IC₅₀ values are those concentrations required to reduce the binding of radiolabeled C5a to the receptor by
25 50%. (For further descriptions of ligand binding and other functional assays, see Dairaghi, *et al.*, *J. Biol. Chem.* **274**:21569-21574 (1999), Penfold, *et al.*, *Proc. Natl. Acad. Sci. USA.* **96**:9839-9844 (1999), and Dairaghi, *et al.*, *J. Biol. Chem.* **272**:28206-28209 (1997)).

2. Calcium mobilization

[0151] Optionally, compounds may be further assayed for their ability to inhibit calcium
30 flux in cells. To detect the release of intracellular stores of calcium, cells (e.g., cAMP

stimulated U937 or neutrophils) are incubated with 3 μ M of INDO-1AM dye (Molecular Probes; Eugene, OR) in cell media for 45 minutes at room temperature and washed with phosphate buffered saline (PBS). After INDO-1AM loading, the cells are resuspended in flux buffer (Hank's balanced salt solution (HBSS) and 1% FBS). Calcium mobilization is measured using a Photon Technology International spectrophotometer (Photon Technology International; New Jersey) with excitation at 350 nm and dual simultaneous recording of fluorescence emission at 400 nm and 490 nm. Relative intracellular calcium levels are expressed as the 400 nm/490 nm emission ratio. Experiments are performed at 37°C with constant mixing in cuvettes each containing 10^6 cells in 2 mL of flux buffer. The chemokine ligands may be used over a range from 1 to 100 nM. The emission ratio is plotted over time (typically 2-3 minutes). Candidate ligand blocking compounds (up to 10 μ M) are added at 10 seconds, followed by chemokines at 60 seconds (*i.e.*, C5a; R&D Systems; Minneapolis, MN) and control chemokine (*i.e.*, SDF-1 α ; R&D Systems; Minneapolis, MN) at 150 seconds.

3. Chemotaxis assays

[0152] Optionally, compounds may be further assayed for their ability to inhibit chemotaxis in cells. Chemotaxis assays are performed using 5 μ m pore polycarbonate, polyvinylpyrrolidone-coated filters in 96-well chemotaxis chambers (Neuroprobe; Gaithersburg, MD) using chemotaxis buffer (Hank's balanced salt solution (HBSS) and 1% FBS). C5aR ligands (*i.e.*, C5a, R&D Systems; Minneapolis, MN) are used to evaluate compound mediated inhibition of C5aR mediated migration. Other chemokines (*i.e.*, SDF-1 α ; R&D Systems; Minneapolis, MN) are used as specificity controls. The lower chamber is loaded with 29 μ l of chemokine (*i.e.*, 0.03 nM C5a) and varying amounts of compound; the top chamber contains 100,000 U937 or neutrophil cells in 20 μ l. The chambers are incubated 1.5 hours at 37°C, and the number of cells in the lower chamber quantified either by direct cell counts in five high powered fields per well or by the CyQuant assay (Molecular Probes), a fluorescent dye method that measures nucleic acid content and microscopic observation.

C. **Identification of inhibitors of C5aR**

1. Assay

[0153] To evaluate small organic molecules that prevent the C5a receptor from binding ligand, an assay was employed that detected radioactive ligand (*i.e.*, C5a) binding to cells

expressing C5aR on the cell surface (for example, cAMP stimulated U937 cells or isolated human neutrophils). For compounds that inhibited binding, whether competitive or not, fewer radioactive counts are observed when compared to uninhibited controls.

[0154] Equal numbers of cells were added to each well in the plate. The cells were then incubated with radiolabeled C5a. Unbound ligand was removed by washing the cells, and bound ligand was determined by quantifying radioactive counts. Cells that were incubated without any organic compound gave total counts; non-specific binding was determined by incubating the cells with unlabeled ligand and labeled ligand. Percent inhibition was determined by the equation:

$$\% \text{ inhibition} = (1 - [(\text{sample cpm}) - (\text{nonspecific cpm})]/[(\text{total cpm}) - (\text{nonspecific cpm})]) \times 100.$$

2. Dose Response Curves

[0155] To ascertain a candidate compound's affinity for C5aR as well as confirm its ability to inhibit ligand binding, inhibitory activity was titrated over a 1×10^{-10} to 1×10^{-4} M range of compound concentrations. In the assay, the amount of compound was varied; while cell number and ligand concentration were held constant.

D. In Vivo Efficacy Models

[0156] The compounds of interest can be evaluated for potential efficacy in treating a C5a mediated conditions by determining the efficacy of the compound in an animal model. In addition to the models described below, other suitable animal models for studying the compound of interest can be found in Mizuno, M. *et al.*, *Expert Opin. Investig. Drugs* (2005), 14(7), 807-821, which is incorporated herein by reference in its entirety.

1. Models of C5a induced Leukopenia

a) C5a induced Leukopenia in a Human C5aR knock-in Mouse Model

[0157] To study the efficacy of compounds of the instant invention in an animal model, a recombinant mouse can be created using standard techniques, wherein the genetic sequence coding for the mouse C5aR is replaced with sequence coding for the human C5aR, to create a hC5aR-KI mouse. In this mouse, administration of hC5a leads to upregulation of adhesion molecules on blood vessel walls which bind blood leukocytes, sequestering them from the blood stream. Animals are administered 20ug/kg of hC5a and 1 minute later leukocytes are quantified in peripheral blood by standard techniques. Pretreatment of mice with varying doses of the present compounds can almost completely block the hC5a induced leukopenia.

b) C5a induced Leukopenia in a Cynomolgus Model

[0158] To study the efficacy of compounds of the instant invention in a non-human primate model, C5a induced leukopenia is studied in a cynomolgus model. In this model administration of hC5a leads to upregulation of adhesion molecules on blood vessel walls which bind blood leukocytes, hence sequestering them from the blood stream. Animals are administered 10ug/kg of hC5a and 1 minute later leukocytes are quantified in peripheral blood.

Mouse model of ANCA induced Vasculitis

[0159] On day 0 hC5aR-KI mice are intravenously injected with 50mg/kg purified antibody to myeloperoxidase (Xiao et al, *J. Clin. Invest.* **110**: 955-963 (2002)). Mice are further dosed with oral daily doses of compounds of the invention or vehicle for seven days, then mice are sacrificed and kidneys collected for histological examination. Analysis of kidney sections can show significantly reduced number and severity of crescentic and necrotic lesions in the glomeruli when compared to vehicle treated animals.

2. Mouse Model of Choroidal Neovascularization

[0160] To study the efficacy of compounds of the instant invention in treatment of age related macular degeneration (AMD) the bruch membrane in the eyes of hC5aR-KI mice are ruptured by laser photocoagulation (Nozika et al, *PNAS* **103**: 2328-2333 (2006)). Mice are

treated with vehicle or a daily oral or appropriate intra-vitreous dose of a compound of the invention for one to two weeks. Repair of laser induced damage and neovascularization are assessed by histology and angiography.

5 3. Rheumatoid Arthritis Models

 a) *Rabbit model of destructive joint inflammation*

[0161] To study the effects of candidate compounds on inhibiting the inflammatory response of rabbits to an intra-articular injection of the bacterial membrane component lipopolysaccharide (LPS), a rabbit model of destructive joint inflammation is used. This study design mimics the destructive joint inflammation seen in arthritis. Intra-articular injection of LPS causes an acute inflammatory response characterized by the release of cytokines and chemokines, many of which have been identified in rheumatoid arthritic joints. Marked increases in leukocytes occur in synovial fluid and in synovium in response to elevation of these chemotactic mediators. Selective antagonists of chemokine receptors have shown efficacy in this model (see Podolin, *et al.*, *J. Immunol.* 169(11):6435-6444 (2002)).

[0162] A rabbit LPS study is conducted essentially as described in Podolin, *et al. ibid.*, female New Zealand rabbits (approximately 2 kilograms) are treated intra-articularly in one knee with LPS (10 ng) together with either vehicle only (phosphate buffered saline with 1% DMSO) or with addition of candidate compound (dose 1 = 50 μ M or dose 2 = 100 μ M) in a total volume of 1.0 mL. Sixteen hours after the LPS injection, knees are lavaged and cells counts are performed. Beneficial effects of treatment were determined by histopathologic evaluation of synovial inflammation. Inflammation scores are used for the histopathologic evaluation: 1 - minimal, 2 - mild, 3 - moderate, 4 - moderate-marked.

25 b) *Evaluation of a compound in a rat model of collagen induced arthritis*

[0163] A 17 day developing type II collagen arthritis study is conducted to evaluate the effects of a candidate compound on arthritis induced clinical ankle swelling. Rat collagen arthritis is an experimental model of polyarthritis that has been widely used for preclinical testing of numerous anti-arthritic agents (see Trentham, *et al.*, *J. Exp. Med.* **146**(3):857-868 (1977), Bendele, *et al.*, *Toxicologic Pathol.* **27**:134-142 (1999), Bendele, *et al.*, *Arthritis Rheum.* **42**:498-506 (1999)). The hallmarks of this model are reliable onset and progression

of robust, easily measurable polyarticular inflammation, marked cartilage destruction in association with pannus formation and mild to moderate bone resorption and periosteal bone proliferation.

[0164] Female Lewis rats (approximately 0.2 kilograms) are anesthetized with isoflurane and injected with Freund's Incomplete Adjuvant containing 2 mg/mL bovine type II collagen at the base of the tail and two sites on the back on days 0 and 6 of this 17 day study. A candidate compound is dosed daily in a sub-cutaneous manner from day 0 till day 17 at a efficacious dose. Caliper measurements of the ankle joint diameter were taken, and reducing joint swelling is taken as a measure of efficacy.

4. Rat model of Sepsis

[0165] To study the effect of compounds of interest on inhibiting the generalized inflammatory response that is associated with a sepsis like disease, the Cecal Ligation and Puncture (CLP) rat model of sepsis is used. A Rat CLP study is conducted essentially as described in Fujimura N, et al. (*American Journal Respiratory Critical Care Medicine* 2000; 161: 440-446). Briefly described here, Wistar Albino Rats of both sexes weighing between 200-250 g are fasted for twelve hours prior to experiments. Animals are kept on normal 12 hour light and dark cycles and fed standard rat chow up until 12 hours prior to experiment. Then animals are split into four groups; (i) two sham operation groups and (ii) two CLP groups. Each of these two groups (i.e., (i) and (ii)) is split into vehicle control group and test compound group. Sepsis is induced by the CLP method. Under brief anesthesia a midline laparotomy is made using minimal dissection and the cecum is ligated just below the ileocaecal valve with 3-0 silk, so the intestinal continuity is maintained. The antimesenteric surface of the cecum is perforated with an 18 gauge needle at two locations 1 cm apart and the cecum is gently squeezed until fecal matter is extruded. The bowel is then returned to the abdomen and the incision is closed. At the end of the operation, all rats are resuscitated with saline, 3 ml/100 g body weight, given subcutaneously. Postoperatively, the rats are deprived of food, but have free access to water for the next 16 hours until they are sacrificed. The sham operated groups are given a laparotomy and the cecum is manipulated but not ligated or perforated. Beneficial effects of treatment are measured by histopathological scoring of tissues and organs as well as measurement of several key indicators of hepatic function, renal function, and lipid peroxidation. To test for hepatic function aspartate transaminase (AST) and alanine transaminase (ALT) are measured. Blood urea nitrogen and creatinine

concentrations are studied to assess renal function. Pro-inflammatory cytokines such as TNF-alpha and IL-1beta are also assayed by ELISA for serum levels.

5. Mouse SLE model of experimental lupus nephritis.

[0166] To study the effect of compounds of interest on a Systemic Lupus Erythematosus (SLE), the MRL/lpr murine SLE model is used. The MRL/Mp-Tmfrsf6^{lpr/lpr} strain (MRL/lpr) is a commonly used mouse model of human SLE. To test compounds efficacy in this model male MRL/lpr mice are equally divided between control and C5aR antagonists groups at 13 weeks of age. Then over the next 6 weeks compound or vehicle is administered to the animals via osmotic pumps to maintain coverage and minimize stress effects on the animals. Serum and urine samples are collected bi-weekly during the six weeks of disease onset and progression. In a minority of these mice glomerulosclerosis develops leading to the death of the animal from renal failure. Following mortality as an indicator of renal failure is one of the measured criteria and successful treatment will usually result in a delay in the onset of sudden death among the test groups. In addition, the presence and magnitude of renal disease may also be monitored continuously with blood urea nitrogen (BUN) and albuminuria measurements. Tissues and organs were also harvested at 19 weeks and subjected to histopathology and immunohistochemistry and scored based on tissue damage and cellular infiltration.

6. Rat model of COPD

[0167] Smoke induced airway inflammation in rodent models may be used to assess efficacy of compounds in Chronic Obstructive Pulmonary Disease (COPD). Selective antagonists of chemokines have shown efficacy in this model (*see*, Stevenson, et al., *Am. J. Physiol Lung Cell Mol Physiol.* 288 L514-L522, (2005)). An acute rat model of COPD is conducted as described by Stevenson et al. A compound of interest is administered either systemically via oral or IV dosing; or locally with nebulized compound. Male Sprague-Dawley rats (350-400 g) are placed in Perspex chambers and exposed to cigarette smoke drawn in via a pump (50 mL every 30 seconds with fresh air in between). Rats are exposed for a total period of 32 minutes. Rats are sacrificed up to 7 days after initial exposure. Any beneficial effects of treatment are assessed by a decrease inflammatory cell infiltrate, decreases in chemokine and cytokine levels.

[0168] In a chronic model, mice or rats are exposed to daily tobacco smoke exposures for up to 12 months. Compound is administered systemically via once daily oral dosing, or

potentially locally via nebulized compound. In addition to the inflammation observed with the acute model (Stevensen et al.), animals may also exhibit other pathologies similar to that seen in human COPD such as emphysema (as indicated by increased mean linear intercept) as well as altered lung chemistry (see Martorana et al, *Am. J. Respir. Crit Care Med.* 172(7):

848-53.

7. Mouse EAE Model of Multiple Sclerosis

[0169] Experimental autoimmune encephalomyelitis (EAE) is a model of human multiple sclerosis. Variations of the model have been published, and are well known in the field. In a typical protocol, C57BL/6 (Charles River Laboratories) mice are used for the EAE model.

Mice are immunized with 200ug myelin oligodendrocyte glycoprotein (MOG) 35–55 (Peptide International) emulsified in Complete Freund's Adjuvant (CFA) containing 4 mg/ml *Mycobacterium tuberculosis* (Sigma-Aldrich) s.c. on day 0. In addition, on day 0 and day 2 animals are given 200 ng of pertussis toxin (Calbiochem) i.v. Clinical scoring is based on a scale of 0–5: 0, no signs of disease; 1, flaccid tail; 2, hind limb weakness; 3, hind limb paralysis; 4, forelimb weakness or paralysis; 5, moribund. Dosing of the compounds of interest to be assessed can be initiated on day 0 (prophylactic) or day 7 (therapeutic, when histological evidence of disease is present but few animals are presenting clinical signs) and dosed once or more per day at concentrations appropriate for their activity and pharmacokinetic properties, e.g. 100 mg/kg s.c. Efficacy of compounds can be assessed by comparisons of severity (maximum mean clinical score in presence of compound compared to vehicle), or by measuring a decrease in the number of macrophages (F4/80 positive) isolated from spinal cords. Spinal cord mononuclear cells can be isolated via discontinuous Percoll-gradient. Cells can be stained using rat anti-mouse F4/80-PE or rat IgG2b-PE (Caltag Laboratories) and quantitated by FACS analysis using 10 ul of Polybeads per sample (Polysciences).

8. Mouse Model of Kidney Transplantation

[0170] Transplantation models can be performed in mice, for instance a model of allogenic kidney transplant from C57BL/6 to BALB/c mice is described in Faikah Gueler et al, *JASN Express*, Aug 27th, 2008. Briefly, mice are anesthetized and the left donor kidney attached to a cuff of the aorta and the renal vein with a small caval cuff, and the ureters removed en block. After left nephrectomy of the recipient, the vascular cuffs are anastomosed to the recipient abdominal aorta and vena cava, respectively, below the level of the native renal

vessels. The ureter is directly anastomosed into the bladder. Cold ischemia time is 60 min, and warm ischemia time is 30 min. The right native kidney can be removed at the time of allograft transplantation or at posttransplantation day 4 for long-term survival studies.

General physical condition of the mice is monitored for evidence of rejection. Compound

- 5 treatment of animals can be started before surgery or immediately after transplantation, eg by sub cut injection once daily. Mice are studied for renal function and survival. Serum creatinine levels are measured by an automated method (Beckman Analyzer, Krefeld, Germany).

9. Mouse Model of Ischemia/Reperfusion


- 10 [0171] A mouse model of ischemia/reperfusion injury can be performed as described by Xiufen Zheng et al, *Am. J. Pathol*, Vol 173:4, Oct, 2008. Briefly, CD1 mice aged 6-8 weeks are anesthetized and placed on a heating pad to maintain warmth during surgery. Following abdominal incisions, renal pedicles are bluntly dissected and a microvascular clamp placed on the left renal pedicle for 25-30 minutes. Following ischemia the clamps are removed
15 along with the right kidney, incisions sutured, and the animals allowed to recover. Blood is collected for serum creatinine and BUN analysis as an indicator of kidney health. Alternatively animal survival is monitored over time. Compound can be administered to animals before and/or after the surgery and the effects on serum creatinine, BUN or animal survival used as indicators of compound efficacy.

20 10. Mouse Model of Tumor Growth

- [0172] C57BL/6 mice 6–16 weeks of age are injected subcutaneously with 1×10^5 TC-1 cells (ATCC, VA) in the right or left rear flank. Beginning about 2 weeks after cell injection, tumors are measured with calipers every 2–4 d until the tumor size required the mice are killed. At the time of sacrifice animals are subjected to a full necropsy and spleens
25 and tumors removed. Excised tumors are measured and weighed. Compounds may be administered before and/or after tumor injections, and a delay or inhibition of tumor growth used to assess compound efficacy.

WHAT IS CLAIMED IS:

1 1. A compound having the formula



2 and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein

3 C¹ is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group
4 has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said
5 aryl and heteroaryl groups are optionally substituted with from 1 to 3 R¹ substituents;

6 C² is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group
7 has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said
8 aryl and heteroaryl groups are optionally substituted with from 1 to 3 R² substituents;

9 C³ is selected from the group consisting of C₁₋₈ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl-C₁₋₄
10 alkyl, aryl, aryl-C₁₋₄ alkyl, heteroaryl, heteroaryl-C₁₋₄ alkyl, heterocycloalkyl or
11 heterocycloalkyl-C₁₋₄ alkyl, wherein the heterocycloalkyl group or portion has from 1-3
12 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3
13 heteroatoms as ring members selected from N, O and S, and each C³ is optionally
14 substituted with from 1-3 R³ substituents;


15 each R¹ is independently selected from the group consisting of halogen, -CN, -R^c, -CO₂R^a,
16 -CONR^aR^b, -C(O)R^a, -OC(O)NR^aR^b, -NR^bC(O)R^a, -NR^bC(O)₂R^c, -NR^a-C(O)NR^aR^b,
17 -NR^aC(O)NR^aR^b, -NR^aR^b, -OR^a, and -S(O)₂NR^aR^b; wherein each R^a and R^b is
18 independently selected from hydrogen, C₁₋₈ alkyl, and C₁₋₈ haloalkyl, or when attached to
19 the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
20 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
21 N, O or S; each R^c is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈
22 haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic
23 and cyclic portions of R^a, R^b and R^c are optionally further substituted with from one to
24 three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and
25 optionally when two R¹ substituents are on adjacent atoms, are combined to form a fused
26 five or six-membered carbocyclic ring;

27 each R² is independently selected from the group consisting of halogen, -R^f, -CO₂R^d,
28 -CONR^dR^e, -C(O)R^d, -OC(O)NR^dR^e, -NR^eC(O)R^d, -NR^eC(O)₂R^f, -NR^dC(O)NR^dR^e,
29 -NR^dC(O)NR^dR^e,


30 $-\text{NR}^{\text{d}}\text{C}(\text{O})\text{NR}^{\text{d}}\text{R}^{\text{e}}$, $-\text{NR}^{\text{d}}\text{R}^{\text{e}}$, $-\text{OR}^{\text{d}}$, and $-\text{S}(\text{O})_2\text{NR}^{\text{d}}\text{R}^{\text{e}}$; wherein each R^{d} and R^{e} is
 31 independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to
 32 the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
 33 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
 34 N, O or S; each R^{f} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
 35 haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic
 36 and cyclic portions of R^{d} , R^{e} and R^{f} are optionally further substituted with from one to
 37 three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups;
 38 each R^3 is independently selected from the group consisting of halogen, $-\text{CN}$, $-\text{R}^{\text{i}}$, $-\text{CO}_2\text{R}^{\text{g}}$,
 39 $-\text{CONR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{C}(\text{O})\text{R}^{\text{g}}$, $-\text{OC}(\text{O})\text{NR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{NR}^{\text{h}}\text{C}(\text{O})\text{R}^{\text{g}}$, $-\text{NR}^{\text{h}}\text{C}(\text{O})_2\text{R}^{\text{i}}$, $-\text{NR}^{\text{g}}\text{C}(\text{O})\text{NR}^{\text{g}}\text{R}^{\text{h}}$,
 40 $-\text{NR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{OR}^{\text{g}}$, $-\text{S}(\text{O})_2\text{NR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{X}^4-\text{R}^{\text{j}}$, $-\text{X}^4-\text{NR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{X}^4-\text{CONR}^{\text{g}}\text{R}^{\text{h}}$, $-\text{X}^4-\text{NR}^{\text{h}}\text{C}(\text{O})\text{R}^{\text{g}}$, $-\text{NHR}^{\text{j}}$
 41 and $-\text{NHCH}_2\text{R}^{\text{j}}$, wherein X^4 is a C_{1-4} alkylene; each R^{g} and R^{h} is independently selected
 42 from hydrogen, C_{1-8} alkyl, C_{3-6} cycloalkyl and C_{1-8} haloalkyl, or when attached to the
 43 same nitrogen atom can be combined with the nitrogen atom to form a five or six-
 44 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
 45 N, O or S and is optionally substituted with one or two oxo; each R^{i} is independently
 46 selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl,
 47 heterocycloalkyl, aryl and heteroaryl; and each R^{j} is selected from the group consisting of
 48 C_{3-6} cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and
 49 tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^{g} , R^{h} , R^{i} and R^{j} are
 50 optionally further substituted with from one to three halogen, methyl, CF_3 , hydroxy,
 51 amino, alkylamino and dialkylamino groups; and
 52 X is hydrogen or CH_3 .

1 2. The compound of claim 1, wherein X is hydrogen.


1 3. The compound of claim 1, having the formula:



1 4. The compound of claim 1, having the formula:



5. The compound of claim 1, having the formula:



wherein


X^1 is selected from the group consisting of N, CH and CR^1 ;

the subscript n is an integer of from 0 to 2;

X^2 is selected from the group consisting of N, CH and CR^2 ; and

the subscript m is an integer of from 0 to 2.

6. The compound of claim 1, having the formula:



wherein


X^1 is selected from the group consisting of N, CH and CR^1 ;

the subscript n is an integer of from 0 to 2;

X^2 is selected from the group consisting of N, CH and CR^2 ; and

the subscript m is an integer of from 0 to 2.

7. The compound of claim 1, having the formula:



wherein

the subscript p is an integer of from 0 to 3;


X^1 is selected from the group consisting of N, CH and CR^1 ;

the subscript n is an integer of from 0 to 2;


X^2 is selected from the group consisting of N, CH and CR^2 ; and

the subscript m is an integer of from 0 to 2.


8. The compound of claim 1, having the formula:




9. The compound of claim 1, having the formula:




10. The compound of claim 1, having the formula:




11. The compound of claim 1, having the formula:




12. The compound of claim 1, having the formula:



13. The compound of claim 1, having the formula:




14. The compound of claim 1, having the formula:



wherein R^3 is a member selected from the group consisting of $-NR^gR^h$, $-NHR^j$ and $-NHCH_2R^j$.

15. The compound of claim 1, having the formula:



wherein R^3 is a member selected from the group consisting of $-X^4-NR^gR^h$, $-X^4-R^j$ and $-X^4-NR^hCOR^g$.

16. The compound of claim 1, wherein C^1 is selected from the group consisting of phenyl, pyridyl, indolyl and thiazolyl, each of which is optionally substituted with from 1 to 3 R^1 substituents.

17. The compound of claim 1, wherein C^2 is selected from the group consisting of phenyl, naphthyl, pyridyl and indolyl, each of which is optionally substituted with from 1 to 3 R^2 substituents.

18. The compound of claim 1, wherein C^3 is selected from the group consisting of C_{3-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-2} alkyl, phenyl, pyridinyl, pyrazolyl, piperidinyl, pyrrolidinyl, piperidinylmethyl and pyrrolidinylmethyl, each of which is optionally substituted with from 1 to 3 R^3 substituents.


19. The compound of claim 16, wherein each R^1 is independently selected from the group consisting of halogen, $-CN$, $-R^c$, $-NR^aR^b$ and $-OR^a$, and wherein each R^a and R^b is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached

4 to the same nitrogen atom can be combined with the nitrogen atom to form a pyrrolidine ring;
5 each R^c is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl and
6 C₃₋₆ cycloalkyl, and wherein the aliphatic and cyclic portions of R^a, R^b and R^c are optionally
7 further substituted with from one to three hydroxy, methyl, amino, alkylamino and
8 dialkylamino groups; and optionally when two R¹ substituents are on adjacent atoms, are
9 combined to form a fused five or six-membered carbocyclic ring.

1 **20.** The compound of claim 17, wherein each R² is independently selected
2 from the group consisting of halogen, -R^f and -OR^d; wherein each R^d is independently
3 selected from hydrogen, C₁₋₈ alkyl, and C₁₋₈ haloalkyl; each R^f is independently selected from
4 the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl and
5 heteroaryl, and wherein the aliphatic and cyclic portions of R^d and R^f are optionally further
6 substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and
7 dialkylamino groups.

1 **21.** The compound of claim 18, wherein each R³ is independently selected
2 from the group consisting of halogen, -Rⁱ, -CO₂R^g, -CONR^gR^h, -NR^hC(O)R^g, -NR^hC(O)₂Rⁱ,
3 -NR^gR^h, -OR^g, -X⁴-R^j, -X⁴-NR^gR^h, -X⁴-CONR^gR^h, -X⁴-NR^hC(O)R^g, -NHR^j and -NHCH₂R^j,
4 wherein X⁴ is a C₁₋₃ alkylene; each R^g and R^h is independently selected from hydrogen, C₁₋₈
5 alkyl, C₃₋₆ cycloalkyl and C₁₋₈ haloalkyl, or when attached to the same nitrogen atom can be
6 combined with the nitrogen atom to form a five or six-membered ring having from 0 to 1
7 additional heteroatoms as ring members selected from N, O or S and is optionally substituted
8 with one or two oxo; each Rⁱ is independently selected from the group consisting of C₁₋₈
9 alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is
10 selected from the group consisting of C₃₋₆ cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl,
11 tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g,
12 R^h, Rⁱ and R^j are optionally further substituted with from one to three halogen, methyl, CF₃,
13 hydroxy, amino, alkylamino and dialkylamino groups

1 **22.** The compound of claim 20, wherein C² is selected from the group
2 consisting of:




3


1


23. The compound of claim 19, wherein C^1 is selected from the group

2

consisting of:








26. The compound of claim 1, wherein said compound is selected from the group in Table1.

27. A pharmaceutical composition comprising a compound a pharmaceutically acceptable carrier and a compound having formula



and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein

C^1 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^1 substituents;

C^2 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^2 substituents;


C^3 is selected from the group consisting of C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl- C_{1-4} alkyl, aryl, aryl- C_{1-4} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heterocycloalkyl or heterocycloalkyl- C_{1-4} alkyl, wherein the heterocycloalkyl group or portion has from 1-3 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S, and each C^3 is optionally substituted with from 1-3 R^3 substituents;

each R^1 is independently selected from the group consisting of halogen, -CN, $-R^c$, $-CO_2R^a$, $-CONR^aR^b$, $-C(O)R^a$, $-OC(O)NR^aR^b$, $-NR^bC(O)R^a$, $-NR^bC(O)_2R^c$, $-NR^a-C(O)NR^aR^b$, $-NR^aC(O)NR^aR^b$, $-NR^aR^b$, $-OR^a$, and $-S(O)_2NR^aR^b$; wherein each R^a and R^b is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S; each R^c is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic and cyclic portions of R^a , R^b and R^c are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and optionally when two R^1 substituents are on adjacent atoms, are combined to form a fused five or six-membered carbocyclic ring;

each R^2 is independently selected from the group consisting of halogen, $-R^f$, $-CO_2R^d$, $-CONR^dR^e$, $-C(O)R^d$, $-OC(O)NR^dR^e$, $-NR^eC(O)R^d$, $-NR^eC(O)_2R^f$, $-NR^dC(O)NR^dR^e$, $-NR^dC(O)NR^dR^e$, $-NR^dR^e$, $-OR^d$, and $-S(O)_2NR^dR^e$; wherein each R^d and R^e is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S; each R^f is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic and cyclic portions of R^d , R^e and R^f are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups;

each R^3 is independently selected from the group consisting of halogen, $-\text{CN}$, $-\text{R}^i$, $-\text{CO}_2\text{R}^g$, $-\text{CONR}^g\text{R}^h$, $-\text{C}(\text{O})\text{R}^g$, $-\text{OC}(\text{O})\text{NR}^g\text{R}^h$, $-\text{NR}^h\text{C}(\text{O})\text{R}^g$, $-\text{NR}^h\text{C}(\text{O})_2\text{R}^i$, $-\text{NR}^g\text{C}(\text{O})\text{NR}^g\text{R}^h$, $-\text{NR}^g\text{R}^h$, $-\text{OR}^g$, $-\text{S}(\text{O})_2\text{NR}^g\text{R}^h$, $-\text{X}^4-\text{R}^j$, $-\text{X}^4-\text{NR}^g\text{R}^h$, $-\text{X}^4-\text{CONR}^g\text{R}^h$, $-\text{X}^4-\text{NR}^h\text{C}(\text{O})\text{R}^g$, $-\text{NHR}^j$ and $-\text{NHCH}_2\text{R}^j$, wherein X^4 is a C_{1-4} alkylene; each R^g and R^h is independently selected from hydrogen, C_{1-8} alkyl, C_{3-6} cycloalkyl and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S and is optionally substituted with one or two oxo; each R^i is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of C_{3-6} cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g , R^h , R^i and R^j are optionally further substituted with from one to three halogen, methyl, CF_3 , hydroxy, amino, alkylamino and dialkylamino groups; and X is hydrogen or CH_3 .

28. A method for treating a mammal suffering from or susceptible to a disease or disorder involving pathologic activation of C5a receptors, comprising administering to the mammal an effective amount of a compound having




and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein C^1 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^1 substituents; C^2 is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R^2 substituents; C^3 is selected from the group consisting of C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl- C_{1-4} alkyl, aryl, aryl- C_{1-4} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heterocycloalkyl or heterocycloalkyl- C_{1-4} alkyl, wherein the heterocycloalkyl group or portion has from 1-3 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3

16 heteroatoms as ring members selected from N, O and S, and each C³ is optionally
 17 substituted with from 1-3 R³ substituents;
 18 each R¹ is independently selected from the group consisting of halogen, -CN, -R^c, -CO₂R^a,
 19 -CONR^aR^b, -C(O)R^a, -OC(O)NR^aR^b, -NR^bC(O)R^a, -NR^bC(O)₂R^c, -NR^a-C(O)NR^aR^b,
 20 -NR^aC(O)NR^aR^b, -NR^aR^b, -OR^a, and -S(O)₂NR^aR^b; wherein each R^a and R^b is
 21 independently selected from hydrogen, C₁₋₈ alkyl, and C₁₋₈ haloalkyl, or when attached to
 22 the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
 23 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
 24 N, O or S; each R^c is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈
 25 haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic
 26 and cyclic portions of R^a, R^b and R^c are optionally further substituted with from one to
 27 three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and
 28 optionally when two R¹ substituents are on adjacent atoms, are combined to form a fused
 29 five or six-membered carbocyclic ring;
 30 each R² is independently selected from the group consisting of halogen, -CN, -R^f, -CO₂R^d,
 31 -CONR^dR^e, -C(O)R^d, -OC(O)NR^dR^e, -NR^eC(O)R^d, -NR^eC(O)₂R^f, -NR^dC(O)NR^dR^e,
 32 -NR^dC(O)NR^dR^e, -NR^dR^e, -OR^d, and -S(O)₂NR^dR^e; wherein each R^d and R^e is
 33 independently selected from hydrogen, C₁₋₈ alkyl, and C₁₋₈ haloalkyl, or when attached to
 34 the same nitrogen atom can be combined with the nitrogen atom to form a five or six-
 35 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
 36 N, O or S; each R^f is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈
 37 haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic
 38 and cyclic portions of R^d, R^e and R^f are optionally further substituted with from one to
 39 three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups;
 40 each R³ is independently selected from the group consisting of halogen, -CN, -Rⁱ, -CO₂R^g,
 41 -CONR^gR^h, -C(O)R^g, -OC(O)NR^gR^h, -NR^hC(O)R^g, -NR^hC(O)₂Rⁱ, -NR^gC(O)NR^gR^h,
 42 -NR^gR^h, -OR^g, -S(O)₂NR^gR^h, -X⁴-R^j, -X⁴-NR^gR^h, -X⁴-CONR^gR^h, -X⁴-NR^hC(O)R^g, -NHR^j
 43 and -NHCH₂R^j, wherein X⁴ is a C₁₋₄ alkylene; each R^g and R^h is independently selected
 44 from hydrogen, C₁₋₈ alkyl, C₃₋₆ cycloalkyl and C₁₋₈ haloalkyl, or when attached to the
 45 same nitrogen atom can be combined with the nitrogen atom to form a five or six-
 46 membered ring having from 0 to 2 additional heteroatoms as ring members selected from
 47 N, O or S and is optionally substituted with one or two oxo; each Rⁱ is independently
 48 selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl,
 49 heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of

C₃₋₆ cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g, R^h, Rⁱ and R^j are optionally further substituted with from one to three halogen, methyl, CF₃, hydroxy, amino, alkylamino and dialkylamino groups; and X is hydrogen or CH₃.

29. A method of inhibiting C5a receptor-mediated cellular chemotaxis comprising contacting mammalian white blood cells with a C5a receptor modulatory amount of a compound having formula



and pharmaceutically acceptable salts, hydrates and rotomers thereof; wherein C¹ is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R¹ substituents; C² is selected from the group consisting of aryl and heteroaryl, wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S; and wherein said aryl and heteroaryl groups are optionally substituted with from 1 to 3 R² substituents; C³ is selected from the group consisting of C₁₋₈ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl-C₁₋₄ alkyl, aryl, aryl-C₁₋₄ alkyl, heteroaryl, heteroaryl-C₁₋₄ alkyl, heterocycloalkyl or heterocycloalkyl-C₁₋₄ alkyl, wherein the heterocycloalkyl group or portion has from 1-3 heteroatoms selected from N, O and S, and wherein the heteroaryl group has from 1-3 heteroatoms as ring members selected from N, O and S, and each C³ is optionally substituted with from 1-3 R³ substituents; each R¹ is independently selected from the group consisting of halogen, -CN, -R^c, -CO₂R^a, -CONR^aR^b, -C(O)R^a, -OC(O)NR^aR^b, -NR^bC(O)R^a, -NR^bC(O)₂R^c, -NR^a-C(O)NR^aR^b, -NR^aC(O)NR^aR^b, -NR^aR^b, -OR^a, and -S(O)₂NR^aR^b; wherein each R^a and R^b is independently selected from hydrogen, C₁₋₈ alkyl, and C₁₋₈ haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S; each R^c is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic

and cyclic portions of R^a , R^b and R^c are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups; and optionally when two R^1 substituents are on adjacent atoms, are combined to form a fused five or six-membered carbocyclic ring;

each R^2 is independently selected from the group consisting of halogen, $-\text{CN}$, $-\text{R}^f$, $-\text{CO}_2\text{R}^d$, $-\text{CONR}^d\text{R}^e$, $-\text{C}(\text{O})\text{R}^d$, $-\text{OC}(\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^e\text{C}(\text{O})\text{R}^d$, $-\text{NR}^e\text{C}(\text{O})_2\text{R}^f$, $-\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^d\text{R}^e$, $-\text{OR}^d$, and $-\text{S}(\text{O})_2\text{NR}^d\text{R}^e$; wherein each R^d and R^e is independently selected from hydrogen, C_{1-8} alkyl, and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S; each R^f is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl, and wherein the aliphatic and cyclic portions of R^d , R^e and R^f are optionally further substituted with from one to three halogen, hydroxy, methyl, amino, alkylamino and dialkylamino groups;

each R^3 is independently selected from the group consisting of halogen, $-\text{CN}$, $-\text{R}^i$, $-\text{CO}_2\text{R}^g$, $-\text{CONR}^g\text{R}^h$, $-\text{C}(\text{O})\text{R}^g$, $-\text{OC}(\text{O})\text{NR}^g\text{R}^h$, $-\text{NR}^h\text{C}(\text{O})\text{R}^g$, $-\text{NR}^h\text{C}(\text{O})_2\text{R}^i$, $-\text{NR}^g\text{C}(\text{O})\text{NR}^g\text{R}^h$, $-\text{NR}^g\text{R}^h$, $-\text{OR}^g$, $-\text{S}(\text{O})_2\text{NR}^g\text{R}^h$, $-\text{X}^4-\text{R}^j$, $-\text{X}^4-\text{NR}^g\text{R}^h$, $-\text{X}^4-\text{CONR}^g\text{R}^h$, $-\text{X}^4-\text{NR}^h\text{C}(\text{O})\text{R}^g$, $-\text{NHR}^j$ and $-\text{NHCH}_2\text{R}^j$, wherein X^4 is a C_{1-4} alkylene; each R^g and R^h is independently selected from hydrogen, C_{1-8} alkyl, C_{3-6} cycloalkyl and C_{1-8} haloalkyl, or when attached to the same nitrogen atom can be combined with the nitrogen atom to form a five or six-membered ring having from 0 to 2 additional heteroatoms as ring members selected from N, O or S and is optionally substituted with one or two oxo; each R^i is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, heterocycloalkyl, aryl and heteroaryl; and each R^j is selected from the group consisting of C_{3-6} cycloalkyl, pyrrolinyl, piperidinyl, morpholinyl, tetrahydrofuranyl, and tetrahydropyranyl, and wherein the aliphatic and cyclic portions of R^g , R^h , R^i and R^j are optionally further substituted with from one to three halogen, methyl, CF_3 , hydroxy, amino, alkylamino and dialkylamino groups; and

X is hydrogen or CH_3 .

30. The method of claim 28, wherein the disease or disorder is an inflammatory disease or disorder.

1 **31.** The method of claim **30**, wherein the disease or disorder is selected
2 from the group consisting of neutropenia, sepsis, septic shock, Alzheimer's disease, multiple
3 sclerosis, stroke, inflammatory bowel disease, chronic obstructive pulmonary disorder,
4 inflammation associated with burns, lung injury, osteoarthritis, atopic dermatitis, chronic
5 urticaria, ischemia-reperfusion injury, acute respiratory distress syndrome, systemic
6 inflammatory response syndrome, multiple organ dysfunction syndrome, tissue graft rejection
7 and hyperacute rejection of transplanted organs.

1 **32.** The method of claim **28**, wherein the disease or disorder is a
2 cardiovascular or cerebrovascular disorder.

1 **33.** The method of claim **32**, wherein the disease or disorder is selected
2 from the group consisting of myocardial infarction, coronary thrombosis, vascular occlusion,
3 post-surgical vascular reocclusion, arteriosclerosis, traumatic central nervous system injury
4 and ischemic heart disease.

1 **34.** The method of claim **28**, wherein the disease or disorder is an
2 autoimmune disorder.

1 **35.** The method of claim **34**, wherein the disease or disorder is selected
2 from the group consisting of rheumatoid arthritis, systemic lupus erythematosus, Guillain-
3 Barre syndrome, pancreatitis, lupus nephritis, lupus glomerulonephritis, psoriasis, Crohn's
4 disease, vasculitis, irritable bowel syndrome, dermatomyositis, multiple sclerosis, bronchial
5 asthma, pemphigus, pemphigoid, scleroderma, myasthenia gravis, autoimmune hemolytic and
6 thrombocytopenic states, Goodpasture's syndrome, immunovascularitis, tissue graft rejection
7 and hyperacute rejection of transplanted organs.

1 **36.** The method of claim **28**, wherein the disease or disorder is a
2 pathologic sequelae associated with the group consisting of insulin-dependent diabetes,
3 mellitus, lupus nephropathy, Heyman nephritis, membranous nephritis, glomerulonephritis,
4 contact sensitivity responses, and inflammation resulting from contact of blood with artificial
5 surfaces.