Figure 1



Last Injection
Active Treatment Period
Follow-Up Period
Mean (+SD) Plasma Exenatide Concentration (pg/mL)
550
500
450
400
350
300
250
200
150
100
50
0
0
4
8
12
16
20
24
28
Time (weeks)

Figure 2A




Mean Fasting Plasma Glucose (mg/dL)
225
200
175
150
125
100
0
0
3
6
9
12
15
Time (weeks)
Mean change
from baseline:
+18 ± 12 mg/dL
-43 ± 16 mg/dL *
-39 ± 9 mg/dL *

Figure 2B


Baseline (Day -3)
Mean Blood Glucose (mg/dL)
300
250
200
150
100
Pre
Post
Pre
Post
Pre
Post
3 AM
Breakfast
Lunch
Dinner



Week 15
Mean Blood Glucose (mg/dL)
300
250
200
150
100
Pre
Post
Pre
Post
Pre
Post
3 AM
Breakfast
Lunch
Dinner

Figure 2C



Mean Glycosylated Hemoglobin (%)
10
9
8
7
6
0
3
6
9
12
15
Time (weeks)
Mean change
from baseline:
+0.4±0.3%
-1.4±0.3% *
-1.7±0.3% *

Figure 2D




Subjects (%) Achieving A1C ≤7%
100
80
60
40
20
0
0%
35.7%
85.7%
N=12
Placebo LAR
N=14
0.8 mg Exenatide LAR
N=14
2.0 mg Exenatide LAR

Figure 3


Mean Change in Weight (kg)
2
1
0
-1
-2
-3
-4
-5
-6
0
3
6
9
12
15
Time (weeks)
Mean change
from baseline:
-0.04±0.7 kg
-0.03±0.7 kg
-3.8±1.4 kg *